




(22) **Date de dépôt/Filing Date:** 2008/01/18
(41) **Mise à la disp. pub./Open to Public Insp.:** 2008/07/31
(45) **Date de délivrance/Issue Date:** 2019/09/10
(62) **Demande originale/Original Application:** 2 676 417
(30) **Priorités/Priorities:** 2007/01/19 (EP07001143.2);
2007/06/20 (EP07012079.5)

(51) **Cl.Int./Int.Cl.** **C07J 31/00** (2006.01),
A61K 31/575 (2006.01), **A61P 3/00** (2006.01),
C07J 9/00 (2006.01)
(72) **Inventeur/Inventor:**
PELLICCIARI, ROBERTO, IT
(73) **Propriétaire/Owner:**
INTERCEPT PHARMACEUTICALS, INC., US
(74) **Agent:** SMART & BIGGAR

(54) **Titre : MODULATEURS DE TGR5 ET LEURS PROCEDES D'UTILISATION**
(54) **Title: TGR5 MODULATORS AND METHODS OF USE THEREOF**



(57) **Abrégé/Abstract:**

The invention relates to compounds of Formula A: (A) or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. The compounds of Formula A are TGR5 modulators useful for the treatment of various diseases, including obesity, insulin sensitivity, inflammation, cholestasis, and bile desaturation.

55929-3D1

Abstract

The invention relates to compounds of Formula A: (A) or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. The compounds of Formula A are TGR5 modulators useful for the treatment of various diseases, including obesity, insulin sensitivity, inflammation, cholestasis, and bile desaturation.

5

55929-3D1

TGR5 MODULATORS AND METHODS OF USE THEREOF

This is a divisional application of Canadian Patent Application Serial No. 2,676,417 filed on January 18, 2008.

It is to be understood that the expression "the present invention" or the like used in this specification encompasses not only the subject matter of this divisional application but that of the parent also.

Field of the Invention

The invention concerns relates to compounds that modulate TGR5 and compositions
5 useful in methods for the treatment and/or prevention of various diseases.


Background of the Invention

TGR5 receptor is a G-protein-coupled receptor that has been identified as a cell-
surface receptor that is responsive to bile acids. The primary structure of TGR5 and its
10 responsiveness to bile acids has been found to be highly conserved in TGR5 among human,
bovine, rabbit, rat, and mouse, and thus suggests that TGR5 has important physiological
functions. TGR5 has been found to be widely distributed in not only lymphoid tissues but
also in other tissues. High levels of TGR5 mRNA have been detected in placenta, spleen,
and monocytes/macrophages. Bile acids have been shown to induce internalization of the
15 TGR5 fusion protein from the cell membrane to the cytoplasm. Kawamata et al. 2003, J. Bio.
Chem., 278, 9435. TGR5 has been found to be identical to hGPCR19 reported by Takeda et
al. 2002, FEBS Lett. 520, 97-101.

Bile acids are cholesterol metabolites that are formed in the liver and secreted into the
duodenum of the intestine. Bile acids are compounds that play essential roles in the
20 absorption of dietary lipids and regulation of bile acid synthesis. For example, Farnesoid X
receptor (FXR) and pregnane X receptor (PXR) have recently been identified as specific
nuclear receptors for bile acids. Through activation of FXR, bile acids repress the expression
of the rate-limiting enzyme in bile acid synthesis, cholesterol 7 α -hydroxylase (Cyp7 α). The
activation of PXR by bile acids results in both the repression of Cyp7 α and the transcriptional
25 induction of the bile acid-metabolizing enzyme, cytochrome P450 3 α . At high
concentrations, bile acids are also known to exhibit immunosuppressive effects on cell-
mediated immunity and macrophage functions. Bile acids including deoxycholic acid (DCA)
and chenodeoxycholic acid (CDCA) have been demonstrated to have inhibitory activities on
the lipopolysaccharide (LPS)-induced promotion of cytokines in macrophages, including
30 interleukin (IL)-1, IL-6, and tumor necrosis factor alpha (TNF α).

Bile acid compounds that modulate TGR5 have been used for the treatment of various
diseases, including central nervous diseases as well as inflammatory diseases (WO 01/77325
and WO 02/84286). Specifically, bile acid compounds alkylated in position 6 of cholanic
acid have been disclosed in WO 02/072598, WO 2004/0007521, and EP 1568706 as FXR

agonists. The bile acid compound, 23-methyl-ursodeoxycholic (3- α ,7- β -dihydroxy-5- β -cholan-24-oic acid) has also been disclosed (Hepatology, 1988, 8(6), 1571-1576) for the treatment of cholestatic liver diseases.




cholanic acid

5 Modulators of TGR5 provide methods of regulating bile acid and cholesterol homeostasis, fatty acid absorption, and protein and carbohydrate digestion. There is a need for the development of TGR5 modulators for the treatment and/or prevention of various diseases. The present invention has identified compounds that modulate TGR5 as well as methods of using these compounds to treat disorders such as central nervous system diseases,
10 inflammatory diseases, and metabolic diseases such as obesity and insulin sensitivity. Methods of using the compounds of the invention also include to prevent disorders such as central nervous diseases, inflammatory diseases, and metabolic diseases e.g., metabolic syndrome, Type 2 diabetes, obesity, etc.

15 Summary of the Invention

The present invention relates to TGR5 modulators and their use to treat and/or prevent various diseases.

In one aspect, the invention relates to a compound of formula A:



(A) or a pharmaceutically acceptable salt, solvate, hydrate, or

20 prodrug thereof, wherein: R₁ is hydrogen, hydroxy, substituted or unsubstituted alkyl, or halogen; R₂ is hydrogen or α -hydroxy; R₃ is hydrogen, hydroxy, NH(CH₂)_mSO₃H, or NH(CH₂)_nCO₂H; R₄ is hydrogen, substituted or unsubstituted alkyl, or halogen; R₅ is unsubstituted or substituted alkyl, or aryl; R₆ is hydrogen, unsubstituted or substituted alkyl, or R₅ and R₆ taken together with the carbons to which they are attached form a ring of size 3,

55929-3D1

4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; R_8 is hydrogen, substituted or unsubstituted alkyl; R_9 is hydrogen, substituted or unsubstituted alkyl or taken together R_8 and R_9 form a carbonyl;

R_{10} is R_3 , SO_3H ; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. In one aspect, when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxyl or $NHCH_2CH_2SO_3H$, then R_4 is not hydrogen.

In one aspect of the invention, R_1 is hydrogen or hydroxy. R_1 is hydroxy. R_1 is hydrogen. R_2 is α -hydroxy. R_1 is hydroxy and R_2 is α -hydroxy. R_1 is hydroxy and R_2 is H. R_1 is hydroxy and R_2 is H. At least one of R_1 or R_2 is hydroxy. At least one of R_1 or R_2 is hydrogen. R_1 and R_2 are the same. R_1 and R_2 are each α -hydroxy. R_1 and R_2 are each hydrogen.

In another aspect of the invention, R_{10} is R_3 . R_3 is hydroxyl, $NH(CH_2)_mSO_3H$, or $NH(CH_2)_nCO_2H$. R_3 is hydroxyl. R_3 is not hydroxyl. R_3 is $NH(CH_2)_mSO_3H$. R_3 is $NH(CH_2)_mSO_3H$ and m is 2. R_3 is $NH(CH_2)_nCO_2H$. R_3 is $NH(CH_2)_nCO_2H$ and n is 1.

In another aspect of the invention, R_4 is hydrogen or unsubstituted alkyl. R_4 is hydrogen.

R_4 is unsubstituted alkyl. R_4 is unsubstituted alkyl. R_4 is methyl or ethyl. R_4 is methyl. R_4 is ethyl. R_3 and R_4 are the same. R_3 and R_4 are different. R_3 and R_4 are each hydrogen. R_3 is hydroxyl and R_4 is hydrogen. R_3 is $NH(CH_2)_mSO_3H$ and R_4 is hydrogen. R_3 is $NH(CH_2)_mSO_3H$, R_4 is hydrogen, and m is 2. R_3 is $NH(CH_2)_nCO_2H$ and R_4 is hydrogen. R_3 is $NH(CH_2)_nCO_2H$, R_4 is hydrogen, and n is 1. R_3 is H and R_4 is unsubstituted alkyl. R_3 is OH and R_4 is methyl. R_3 is OH and R_4 is ethyl. R_3 is OH and R_4 is methyl.

In another aspect, R_5 is unsubstituted or substituted alkyl. R_5 is in the S-configuration. R_5 is in the R-configuration. R_5 is methyl or ethyl. R_5 is S-methyl. R_5 is R-methyl. R_5 is S-ethyl. R_5 is R-ethyl. R_5 is substituted alkyl substituted with phenyl. R_5 is benzyl. R_5 is S-benzyl. R_5 is R-benzyl. R_5 is aryl. R_5 is phenyl. R_4 and R_5 are each unsubstituted alkyl. R_4 and R_5 are each unsubstituted alkyl, wherein R_5 is in the S-configuration and R_4 is in the alpha-configuration. R_4 and R_5 are each unsubstituted alkyl and R_1 is hydroxy. R_4 and R_5 are each unsubstituted alkyl and R_2 is hydrogen. R_4 and R_5 are each unsubstituted alkyl, R_1 is hydroxy, and R_2 is hydrogen.

In one aspect of the invention, R_1 , R_2 , R_3 , and R_4 are hydrogen. R_2 , R_3 , and R_4 are hydrogen. R_2 and R_3 are hydrogen. At least one of R_1 , R_2 , R_3 , or R_4 is hydrogen. At

55929-3D1

least two of R₁, R₂, R₃, or R₄ are hydrogen. At least three of R₁, R₂, R₃, or R₄ are hydrogen.
R₁, R₂, R₃, and R₄ are hydrogen.

In one aspect of the invention, R₁, R₂, and R₄ are hydrogen and R₃ is OH. R₂ and R₄
are hydrogen and R₃ is OH. R₂ is hydrogen and R₃ is OH. At least one of R₁, R₂, or R₄ is
5 hydrogen and R₃ is OH. At least two of R₁, R₂, or R₄ are hydrogen and R₃ is OH. All of R₁,
R₂ and R₄ are hydrogen and R₃ is OH. R₁, R₂ and R₄ are hydrogen and R₃ is OH.

In another aspect of the invention, at least one of R₁ or R₇ is unsubstituted alkyl. At
least one of R₁ or R₇ is methyl. At least one of R₁ or R₇ is ethyl. At least one of R₁ or R₇ is
propyl. R₁ is methyl. R₁ is ethyl. R₁ is propyl. R₇ is methyl. R₇ is ethyl. R₇ is propyl. Both
10 R₁ and R₇ are unsubstituted alkyl. Both R₁ and R₇ are methyl. Both R₁ and R₇ are ethyl. R₇
is hydrogen. R₇ is hydroxy. R₁ is hydrogen. R₁ is hydroxyl. One of R₁ or R₇ is
unsubstituted alkyl and the other R₁ or R₇ is hydrogen. One of R₁ or R₇ is unsubstituted alkyl
and the other R₁ or R₇ is hydroxy. At least one of R₁ or R₇ is unsubstituted alkyl and R₅ is
unsubstituted or substituted alkyl. At least one of R₁ or R₇ is methyl and R₅ is methyl. R₇ is
15 hydroxy and both R₁ and R₅ are unsubstituted alkyl. R₁ is hydroxyl and both R₇ and R₅ are
unsubstituted alkyl. At least one of R₁ or R₇ is unsubstituted alkyl and R₅ is unsubstituted or
substituted alkyl, further wherein R₅ is in the S-configuration. At least one of R₁ or R₇ is
unsubstituted alkyl and R₅ is unsubstituted or substituted alkyl, further wherein R₅ is in the R-
configuration.

20 In another aspect, R₁ is hydroxy and R₇ is methyl. R₁ is methyl and R₇ is hydroxy.
R₆ is unsubstituted alkyl. R₆ is methyl. R₆ is ethyl. R₆ is propyl.

In another aspect, R₈ is hydrogen. R₈ is unsubstituted alkyl. R₈ is methyl. R₈ is ethyl.
R₈ is propyl. R₂ is α -hydroxy and R₈ is unsubstituted alkyl. In another aspect, R₈ and R₉
form a carbonyl.

25 In one aspect, R₁₀ is R₃. R₃ is hydroxyl. At least one of R₈ or R₉ is hydrogen. R₈ and
R₉ are both hydrogen. At least one of R₈ or R₉ is unsubstituted alkyl. At least one of R₈ or R₉
is methyl. At least one of R₈ or R₉ is ethyl. In another aspect, R₁₀ is SO₃H.

In another aspect of the present invention, when R₂, R₄ and R₆ are each hydrogen, R₃
is hydroxyl, and one of R₁ and R₇ is hydrogen or hydroxyl, then the other R₁ or R₇ is not
30 methyl. In another aspect, when R₂ is α -OH; R₃ is hydroxyl; R₄ and R₆ are each hydrogen;
and one of R₁ and R₇ is hydrogen or hydroxyl, then the other R₁ or R₇ is not methyl. In
another aspect, the present invention does not include the following compounds:


55929-3D1

3 α ,7 α -dihydroxy-7 β -methyl-5 β -cholanoic acid, 3 α ,7 β -dihydroxy-7 α -methyl-5 β -cholanoic acid, 3 α -hydroxy-7 ϵ -methyl-5 β -cholanoic acid, 3 α ,7 β ,12 α -trihydroxy-7 α -methyl-5 β -cholan-24-oic acid; 3 α ,7 α ,12 α -trihydroxy-7 β -methyl-5 β -cholan-24-oic acid; and 3 α ,12 α -dihydroxy-7 ϵ -methyl-5 β -cholan-24-oic acid.

- 5 In another aspect of the present invention, when R₃ is hydroxyl and one of R₁ and R₇ is methyl and the other R₁ and R₇ is hydrogen or hydroxyl, then R₂, R₄ and R₆ are not all hydrogen. In another aspect, when R₂ is α -OH, R₃ is hydroxyl, and one of R₁ and R₇ is methyl and the other R₁ and R₇ is hydrogen or hydroxyl, then R₄ and R₆ are not all hydrogen.

- The invention includes a method of treating disease in a subject, comprising
 10 administering a compound of formula A to a subject in need thereof. The invention includes use of a compound of formula A or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, in the preparation of a medicament for treating and/or preventing disease involving the modulation of the TGR5 receptor comprising administering said compound to a subject in need thereof. The invention includes a composition comprising a compound of
 15 formula A or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, and at least one pharmaceutically acceptable excipient.

In one aspect the invention relates to a compound of formula I:



- or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein R₁ is
 20 hydrogen, hydroxy, or halogen; R₂ is hydrogen or α -hydroxy; R₃ is hydroxy, NH(CH₂)_mSO₃H, or NH(CH₂)_nCO₂H; R₄ is hydrogen, unsubstituted or substituted alkyl, or halogen; R₅ is unsubstituted or substituted alkyl, or aryl; R₆ is hydrogen or R₅ and R₆ taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. In one aspect, when R₅ is
 25 methyl, R₁ is hydroxyl, and R₃ is hydroxyl or NHCH₂CH₂SO₃H, then R₄ is not hydrogen.

In one aspect of the present invention, R₁ is hydrogen or hydroxy. R₁ is α -hydroxy. R₁ is β -hydroxy. R₂ is α -hydroxy. R₁ is β -hydroxy and R₂ is α -hydroxy. R₁ is β -hydroxy

55929-3D1

and R_2 is H. R_1 is α -hydroxy and R_2 is H. R_1 and R_2 are each α -hydroxy. R_1 and R_2 are each hydrogen.

In another aspect of the present invention, R_3 is hydrogen, hydroxyl, $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, or $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$. R_3 is hydroxyl. R_3 is not hydroxyl. R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$. R_3 is
 5 $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$ and m is 2. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and n is 1.

In another aspect of the present invention, R_4 is hydrogen or unsubstituted alkyl. R_4 is hydrogen. R_4 is unsubstituted alkyl. R_4 is methyl or ethyl. R_3 and R_4 are the same. R_3 and R_4 are different. R_3 and R_4 are each hydrogen. R_3 is hydroxyl and R_4 is hydrogen.

In another aspect of the present invention, R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$ and R_4 is hydrogen.
 10 R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, R_4 is hydrogen and m is 2. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and R_4 is hydrogen. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and R_4 is hydrogen and n is 1. R_3 is OH and R_4 is alkyl. R_3 is OH and R_4 is unsubstituted alkyl. R_3 is OH and R_4 is methyl.

In another aspect of the present invention, R_5 is unsubstituted or substituted alkyl. R_5 is in the S-configuration. R_5 is in the R-configuration. R_5 is methyl or ethyl. R_5 is S-methyl.
 15 R_5 is R-methyl. R_5 is S-ethyl. R_5 is R-ethyl. R_5 is alkyl substituted with aryl. R_5 is alkyl substituted with phenyl. R_5 is benzyl. R_5 is S-benzyl. R_5 is R-benzyl. R_5 is aryl. R_5 is phenyl.


In another aspect of the present invention, R_4 and R_5 are each unsubstituted alkyl. R_4 and R_5 are each unsubstituted alkyl and R_5 is in the S-configuration. R_4 is unsubstituted alkyl
 20 and R_5 is substituted alkyl. R_4 and R_5 are each unsubstituted alkyl and R_1 is α -hydroxy. R_4 and R_5 are each unsubstituted alkyl and R_2 is hydrogen. R_4 and R_5 are each unsubstituted alkyl, R_1 is α -hydroxy, and R_2 is hydrogen.

In another aspect of the present invention, R_1 , R_2 , R_3 , and R_4 are hydrogen. R_2 , R_3 , and R_4 are hydrogen. R_2 and R_3 are hydrogen. R_1 , R_2 , and R_4 are hydrogen and R_3 is OH.

25 In another aspect of the present invention, at least one of R_1 , R_2 , R_3 , or R_4 is hydrogen. At least two of R_1 , R_2 , R_3 , or R_4 are hydrogen. At least three of R_1 , R_2 , R_3 , or R_4 are hydrogen. R_1 , R_2 , R_3 , and R_4 are hydrogen.

In another aspect of the present invention, at least one of R_1 , R_2 , or R_4 is hydrogen and R_3 is OH. At least two of R_1 , R_2 , or R_4 are hydrogen and R_3 is OH. All of R_1 , R_2 , and R_4 are
 30 hydrogen and R_3 is OH.


Another aspect of the present invention includes a composition or medicament comprising a compound of formula I.



(I), or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, and at least one pharmaceutically acceptable excipient wherein R_1 is hydrogen, hydroxy, or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, $NH(CH_2)_mSO_3H$, or $NH(CH_2)_nCO_2H$; R_4 is hydrogen, unsubstituted alkyl, or halogen; R_5 is unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. In one aspect the present invention includes a composition or medicament comprising a compound of formula I with the proviso that when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxyl or $NHCH_2CH_2SO_3H$, then R_4 is not hydrogen.

Another aspect of the present invention includes a method of treating disease in a subject, comprising administering a therapeutically effective amount of a compound of formulae I, IA, II, and A to a subject in need thereof. Another aspect of the present invention includes a method of preventing disease in a subject comprising administering a prophylactically effective amount of a compound of formulae I, IA, II, and A to a subject in need thereof. The present invention includes treatment and/or prevention of diseases that involves modulation of TGR5 receptor. The present invention includes use of a compound of formulae A, I, IA, and II in preparation of a medicament for treating and/or preventing a disease that involves the modulation of TGR5 receptor. In one aspect, the disease is a metabolic disease, including obesity and insulin sensitivity. In another aspect, the disease is inflammatory disease, including rheumatoid arthritis and allergy. In another aspect, the disease is cholestasis or bile desaturation. The present invention includes where the subject is human.

The present invention also relates to compounds of formula IA and II:




R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ for the above formula IA and R₁, R₂, R₄, R₅, R₆, R₇, and R₈ for the above formula II are as described below. The present invention relates to a composition or
 5 medicament of a compound of formulae IA or formula II. The present invention includes use of a compound of formulae I, IA, A, and II in the manufacture of a composition for medicament for the treatment and/or prevention of a disease state involving modulation of TGR5.

The above description sets forth rather broadly the more important features of the present invention in order that the detailed description thereof that follows may be understood, and in
 10 order that the present contributions to the art may be better appreciated. Other objects and features of the present invention will become apparent from the following detailed description considered in conjunction with the examples.

The present invention as claimed relates to:

- a compound of formula II



15

or a salt, solvate or hydrate thereof, wherein

R₁ is hydrogen, hydroxyl, substituted or unsubstituted alkyl, or halogen;

R₂ is hydrogen or α -hydroxy;

R₄ is hydrogen, substituted or unsubstituted alkyl, or halogen;

20 R₅ is unsubstituted or substituted alkyl, or aryl;

R₆ is hydrogen, unsubstituted or substituted alkyl; or R₅ and R₆ taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms;

R₇ is hydrogen, substituted or unsubstituted alkyl, or hydroxyl; and

R₈ is hydrogen;

- 5 - a pharmaceutical composition comprising the compound as described herein or a pharmaceutically acceptable salt, solvate or hydrate thereof, and at least one pharmaceutically acceptable excipient;
- use of the compound as described herein or a pharmaceutically acceptable salt, solvate or hydrate thereof, or the composition as described herein, in the preparation of a medicament for
- 10 treating and/or preventing a disease involving modulation of the TGR5 receptor; and
- use of the compound as described herein or a pharmaceutically acceptable salt, solvate or hydrate thereof, or the composition as described herein, for treating and/or preventing a disease involving modulation of the TGR5 receptor.

15 **Description of the Invention**

The details of one or more embodiments of the invention are set forth in the accompanying description below. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the methods and materials are now described. Other features, objects, and advantages of the invention will be

20 apparent from the description. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the case of conflict, the present specification will control.

25 *Definitions*

For convenience, certain terms used in the specification, examples and appended claims are collected here.

The term “treating”, as used herein, means relieving, lessening, reducing, eliminating, modulating, or ameliorating, i.e. causing regression of the disease state or condition.

The term “preventing”, as used herein means, to completely or almost completely stop a disease state or condition, from occurring in a patient or subject, especially when the patient
 5 or subject is predisposed to such or at risk of contracting a disease state or condition. Preventing can also include inhibiting, i.e. arresting the development, of a disease state or condition, and relieving or ameliorating, i.e. causing regression of the disease state or condition, for example when the disease state or condition may already be present.

“Alkyl” includes saturated aliphatic groups, including straight-chain alkyl groups
 10 (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl), branched-chain alkyl groups (e.g., isopropyl, tert-butyl, isobutyl), cycloalkyl (e.g., alicyclic) groups (e.g., cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl), alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In certain embodiments, a straight chain or branched chain alkyl has six or fewer carbon atoms in its backbone (e.g., C₁-C₆ for straight
 15 chain, C₃-C₆ for branched chain). In some examples, a straight chain or branched chain alkyl has four or fewer carbon atoms in its backbone. Further, cycloalkyls have from three to eight carbon atoms in their ring structure.

The term “substituted alkyl” refers to an alkyl moieties having a substituent replace one or more hydrogen atoms on at least one or more carbons of the hydrocarbon backbone.
 20 Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and
 25 alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

“Aryl” includes groups with aromaticity, including 5- and 6-membered
 30 “unconjugated”, or single-ring, aromatic groups that may include from zero to four heteroatoms, as well as “conjugated”, or polycyclic, systems with at least one aromatic ring. Examples of aryl groups include benzene, phenyl, pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isooxazole, pyridine, pyrazine,

The term "ester" includes compounds and moieties which contain a carbon or a heteroatom bound to an oxygen atom which is bonded to the carbon of a carbonyl group. The term "ester" includes alkoxycarboxy groups such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, pentoxycarbonyl, etc. The alkyl, alkenyl, or alkynyl groups are as defined above.

The term "hydroxy" or "hydroxyl" includes groups with an -OH or -O[·].

The term "halogen" includes fluorine, bromine, chlorine, iodine, etc. The term "perhalogenated" generally refers to a moiety wherein all hydrogens are replaced by halogen atoms.

5 An "anionic group," as used herein, refers to a group that is negatively charged at physiological pH. Anionic groups include carboxylate, sulfate, sulfonate, sulfinic, sulfamate, tetrazolyl, phosphate, phosphonate, phosphinate, or phosphorothioate or functional equivalents thereof. "Functional equivalents" of anionic groups are intended to include bioisosteres, *e.g.*, bioisosteres of a carboxylate group. Bioisosteres encompass both classical
10 bioisosteric equivalents and non-classical bioisosteric equivalents. Classical and non-classical bioisosteres are known in the art (see, *e.g.*, Silverman, R. B. The Organic Chemistry of Drug Design and Drug Action, Academic Press, Inc.: San Diego, Calif., 1992, pp.19-23). Another anionic group is a carboxylate.

 The term "unstable functionality" refers to a substitution pattern that contains a labile
15 linkage, *e.g.*, a functionality or bond that is susceptible to hydrolysis or cleavage under physiological conditions (*e.g.*, aqueous solutions in the neutral pH range). Examples of unstable functionalities include acetals and ketals.

 The terms "crystal polymorphs" or "polymorphs" refer to the existence of more than one crystal form for a compound, salt or solvate thereof. Crystal polymorphs of the bile acid
20 analog compounds are prepared by crystallization under different conditions.

 Additionally, the compounds of the present invention, for example, the salts of the compounds, can exist in either hydrated or unhydrated (the anhydrous) form or as solvates with other solvent molecules. Nonlimiting examples of hydrates include monohydrates, dihydrates, etc. Nonlimiting examples of solvates include ethanol solvates, acetone solvates,
25 etc.

 "Solvates" means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate, when the solvent is alcohol, the solvate formed is an
30 alcoholate. Hydrates are formed by the combination of one or more molecules of water with one of the substances in which the water retains its molecular state as H₂O, such combination being able to form one or more hydrate.

It will be noted that the structure of some of the compounds of the invention include asymmetric carbon atoms. It is to be understood accordingly that the isomers arising from such asymmetry (*e.g.*, all enantiomers and diastereomers) are included within the scope of the invention, unless indicated otherwise. Such isomers can be obtained in substantially pure
5 form by classical separation techniques and by stereochemically controlled synthesis. Enantiomers (R- and S-configurations) are named according to the system developed by R.S. Cahn, C. Ingold, and V. Prelog.

Further, the structures and other compounds discussed in this application include all atropic isomers thereof. Atropic isomers are a type of stereoisomer in which the atoms of
10 two isomers are arranged differently in space. Atropic isomers owe their existence to a restricted rotation caused by hindrance of rotation of large groups about a central bond. Such atropic isomers typically exist as a mixture, however as a result of recent advances in chromatography techniques, it has been possible to separate mixtures of two atropic isomers in select cases.

15 “Stable compound” and “stable structure” are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

As used herein, the term “analog” refers to a chemical compound that is structurally similar to another but differs slightly in composition (as in the replacement of one atom by an
20 atom of a different element or in the presence of a particular functional group, or the replacement of one functional group by another functional group). Thus, an analog is a compound that is similar to or comparable in function and appearance to the reference compound.

As defined herein, the term “derivative”, *e.g.*, in the term “bile acid derivatives”,
25 refers to compounds that have a common core 4-membered ring structure, and are substituted with various groups as described herein.

The term “bioisostere” refers to a compound resulting from the exchange of an atom or of a group of atoms with another, broadly similar, atom or group of atoms. The bioisosteric replacement may be physicochemically or topologically based. Examples of
30 carboxylic acid bioisosteres include acyl sulfonimides, tetrazoles, sulfonates, and phosphonates. *See, e.g.*, Patani and LaVoie, Chem. Rev. 96, 3147-3176 (1996).

“Combination therapy” (or “co-therapy”) includes the administration of a compound of the invention and at least a second agent as part of a specific treatment regimen intended to

provide the beneficial effect from the co-action of these therapeutic agents (i.e., the compound of the invention and at least a second agent). The beneficial effect of the combination includes, but is not limited to, pharmacokinetic or pharmacodynamic co-action resulting from the combination of therapeutic agents. Administration of these therapeutic agents in combination typically is carried out over a defined time period (usually minutes, hours, days or weeks depending upon the combination selected). "Combination therapy" may, but generally is not, intended to encompass the administration of two or more of these therapeutic agents as part of separate monotherapy regimens that incidentally and arbitrarily result in the combinations of the present invention. "Combination therapy" is intended to embrace administration of these therapeutic agents in a sequential manner, that is, wherein each therapeutic agent is administered at a different time, as well as administration of these therapeutic agents, or at least two of the therapeutic agents, in a substantially simultaneous manner. Substantially simultaneous administration can be accomplished, for example, by administering to the subject a single capsule having a fixed ratio of each therapeutic agent or in multiple, single capsules for each of the therapeutic agents. Sequential or substantially simultaneous administration of each therapeutic agent can be effected by any appropriate route including, but not limited to, oral routes, intravenous routes, intramuscular routes, and direct absorption through mucous membrane tissues. The therapeutic agents can be administered by the same route or by different routes. For example, a first therapeutic agent of the combination selected may be administered by intravenous injection while the other therapeutic agents of the combination may be administered orally. Alternatively, for example, all therapeutic agents may be administered orally or all therapeutic agents may be administered by intravenous injection. The sequence in which the therapeutic agents are administered is not narrowly critical.

"Combination therapy" also embraces the administration of the therapeutic agents as described above in further combination with other biologically active ingredients and non-drug therapies (*e.g.*, surgery or mechanical treatments). Where the combination therapy further comprises a non-drug treatment, the non-drug treatment may be conducted at any suitable time so long as a beneficial effect from the co-action of the combination of the therapeutic agents and non-drug treatment is achieved. For example, in appropriate cases, the beneficial effect is still achieved when the non-drug treatment is temporally removed from the administration of the therapeutic agents, perhaps by days or even weeks.

The terms "parenteral administration" and "administered parenterally" as used herein refer to modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intra-arterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

The term "pulmonary" as used herein refers to any part, tissue or organ whose primary function is gas exchange with the external environment, *e.g.*, O₂/CO₂ exchange, within a patient. "Pulmonary" typically refers to the tissues of the respiratory tract. Thus, the phrase "pulmonary administration" refers to administering the formulations described herein to any part, tissue or organ whose primary function is gas exchange with the external environment (*e.g.*, mouth, nose, pharynx, oropharynx, laryngopharynx, larynx, trachea, carina, bronchi, bronchioles, alveoli). For purposes of the present invention, "pulmonary" also includes a tissue or cavity that is contingent to the respiratory tract, in particular, the sinuses.

The term "effective amount" means a "therapeutically effective amount" and/or a "prophylactically effective amount."

A "therapeutically effective amount" of a compound of the invention, or a combination of compounds is an amount (quantity or concentration) of compound or compounds. In one embodiment, when a therapeutically effective amount of a compound is administered to a subject in need of treatment symptoms arising from the disease are ameliorated immediately or after administration of the compound one or more times. The amount of the compound to be administered to a subject will depend on the particular disorder, the mode of administration, co-administered compounds, if any, and the characteristics of the subject, such as general health, other diseases, age, sex, genotype, body weight and tolerance to drugs. The skilled artisan will be able to determine appropriate dosages depending on these and other factors.

The term "prophylactically effective amount" means an amount (quantity or concentration) of a compound of the present invention, or a combination of compounds, that is administered to prevent or reduce the risk of a disease – in other words, an amount needed to provide a preventative or prophylactic effect. The amount of the present compound to be administered to a subject will depend on the particular disorder, the mode of administration, co-administered compounds, if any, and the characteristics of the subject, such as general

health, other diseases, age, sex, genotype, body weight and tolerance to drugs. The skilled artisan will be able to determine appropriate dosages depending on these and other factors.

The term "reducing the risk of", as used herein, means to lower the likelihood or probability of a central nervous system disease, inflammatory disease and/or metabolic disease from occurring in a patient, especially when the patient or subject is predisposed to such occurrence.

A "pharmaceutically acceptable salt" or "salt" of a compound of the invention is a product of the compound that contains an ionic bond, and is typically produced by reacting the compound with either an acid or a base, suitable for administering to a subject.

10 A "composition" is a formulation containing s compound of the invention in a form suitable for administration to a subject. In another embodiment, the pharmaceutical composition is in bulk or in unit dosage form. The unit dosage form is any of a variety of forms, including, for example, a capsule, an IV bag, a tablet, a single pump on an aerosol inhaler, or a vial. The quantity of active ingredient (*e.g.*, a formulation of a compound of the invention or salts thereof) in a unit dose of composition is an effective amount and is varied according to the particular treatment involved. One skilled in the art will appreciate that it is sometimes necessary to make routine variations to the dosage depending on the age and condition of the patient. The dosage will also depend on the route of administration. A variety of routes are contemplated, including oral, pulmonary, rectal, parenteral, transdermal, subcutaneous, intravenous, intramuscular, intraperitoneal, intranasal, and the like. Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. In another embodiment, the active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that are required.

The term "flash dose" refers to compound formulations that are rapidly dispersing dosage forms.

The term "immediate release" is defined as a release of compound from a dosage form in a relatively brief period of time, generally up to about 60 minutes. The term "modified release" is defined to include delayed release, extended release, and pulsed release. The term "pulsed release" is defined as a series of releases of drug from a dosage form. The term "sustained release" or "extended release" is defined as continuous release of a compound from a dosage form over a prolonged period.

A "subject" includes mammals, *e.g.*, humans, companion animals (*e.g.*, dogs, cats, birds, and the like), farm animals (*e.g.*, cows, sheep, pigs, horses, fowl, and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs, birds, and the like). Typically, the subject is human.

5 Compounds of the invention also include prodrugs or physiologically equivalent derivatives. A "prodrug" or "physiologically equivalent derivative" includes a precursor form of the drug which is metabolically converted *in vivo* to produce the active drug. The invention further contemplates the use of prodrugs which are converted *in vivo* to the TGR5 modulating compounds used in the methods of the invention (*see, e.g.*, R. B. Silverman,
10 1992, "The Organic Chemistry of Drug Design and Drug Action", Academic Press, Chp. 8). Such prodrugs can be used to alter the biodistribution (*e.g.*, to allow compounds which would not typically cross the blood-brain barrier to cross the blood-brain barrier) or the pharmacokinetics of the TGR5 modulating compound. For example, an anionic group, *e.g.*, a carboxylate, sulfate or sulfonate, can be esterified, *e.g.*, with an alkyl group (*e.g.*, a methyl
15 group) or a phenyl group, to yield an ester. When the ester is administered to a subject, the ester is cleaved, enzymatically or non-enzymatically, reductively or hydrolytically, to reveal the anionic group. Such an ester can be cyclic, *e.g.*, a cyclic sulfate or sulfone, or two or more anionic moieties may be esterified through a linking group. An anionic group can be esterified with moieties (*e.g.*, acyloxymethyl esters) which are cleaved to reveal an
20 intermediate TGR5 modulating compound which subsequently decomposes to yield the active TGR5 modulating compound. In one embodiment, the prodrug is a reduced form of a carboxylate, sulfate or sulfonate, *e.g.*, an alcohol or thiol, which is oxidized *in vivo* to the TGR5 modulating compound. Furthermore, an anionic moiety can be esterified to a group which is actively transported *in vivo*, or which is selectively taken up by target organs.

25 As used herein, the term "amino acid conjugates" refers to conjugates of the compounds of formulae I, IA, and A with any suitable amino acid. Taurine ($\text{NH}(\text{CH}_2)_2\text{SO}_3\text{H}$) and glycine ($\text{NHCH}_2\text{CO}_2\text{H}$) are examples of amino acid conjugates. Suitable amino acid conjugates of the compounds of formulae I, IA, and A have the added advantage of enhanced integrity in bile or intestinal fluids. Suitable amino acids are not
30 limited to taurine and glycine. The present invention encompasses amino acid conjugates of the compounds of formulae I, IA, and A.

The term "TGR5 modulator" means any compound that interacts with the TGR5 receptor. The interaction is not limited to a compound acting as an antagonist, agonist, partial

agonist, or inverse agonist of the TGR5 receptor. In one aspect, the compounds of the present invention act as an antagonist of the TGR5 receptor. In another aspect, the compounds of the present invention act as an agonist of the TGR5 receptor. In another aspect, the compounds of the present invention act as a partial agonist of the TGR5 receptor.

5 In another aspect, the compounds of the present invention as an inverse agonist of the TGR5 receptor. The profile of a ligand, traditionally, endogenous or synthetic, is characterized by its intrinsic efficacy 'e' originally described by Furchgott in 1966. It is used to express the degree to which the different ligands produce varying biological responses while occupying the same number of receptors. Generally, the term "agonist" means a compound that

10 enhances the activity of another molecule or receptor site. An agonist, by classical definition, whether a orthosteric, allosteric, inverse or a co-agonist has a property to bind to the receptor, alter its receptor state and result in a biological action. Consequently, agonism is defined as a property of an agonist or a ligand to produce a biological action. In contrast to this, an "antagonist" is essentially an agonist with high affinity to the same receptor macromolecule,

15 but with very less or negligible intrinsic efficacy, and thus sterically prevents the biological actions of an agonist. As a property, antagonism may be functional or physiological, where an agonist has a direct competition for the receptor site in former and opposing effects via a different receptor-messenger system in the later. More specifically, a TGR5 agonist is a receptor ligand or compound that binds to TGR5 and increases the concentration of cyclic

20 adenosine monophosphate (cAMP) by at least 20% in cells expressing the receptor." Conversely, a TGR5 antagonist would be a compound that antagonizes or blocks the activity of an agonist, thereby effecting a reduction in the concentration of cAMP

The term "metabolic disorders" includes but is not limited to dyslipidemia, atherosclerosis, obesity, coronary heart disease, stroke, insulin resistance/sensitivity, and


25 diabetes.

The term "inflammatory disease" means an inflammatory response which causes injury to autologous tissues. Inflammatory diseases include but are not limited to rheumatoid arthritis, osteoarthritis, cervical spondylosis, cumulative trauma disorder, allergy, endometriosis, pelvic inflammatory disease, adhesive peritonitis, appendicitis, pericarditis,

30 and pleuritis.

The present invention relates to compounds having TGR5 receptor modulating activity and their use to treat and/or prevent various diseases including, central nervous system diseases, inflammatory diseases, and metabolic diseases such as obesity and insulin

sensitivity. Further, the present invention relates to compounds of formulae A, I, IA, and II. According to one aspect, the present invention provides a compound of formula I:





(I)

or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein: R₁ is hydrogen, hydroxy, or halogen; R₂ is hydrogen or α -hydroxy; R₃ is hydroxy, NH(CH₂)_mSO₃H, or NH(CH₂)_nCO₂H; R₄ is hydrogen, unsubstituted or substituted alkyl, or halogen; R₅ is unsubstituted or substituted alkyl, or aryl; R₆ is hydrogen or R₅ and R₆ taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; m is an integer 0, 1, 2, 3, 4, or 5, and n is an integer 0, 1, 2, 3, 4, or 5. In one aspect, when R₅ is methyl, R₁ is hydroxyl, and R₃ is hydroxyl or NHCH₂CH₂SO₃H, then R₄ is not hydrogen.

In one aspect, the present invention provides compounds where R₁ is hydrogen or hydroxy. R₁ is hydroxy. R₁ is hydrogen. R₁ is α -hydroxy. R₁ is β -hydroxy.

In another aspect, the present invention provides compounds where R₁ is halogen. R₁ is fluorine. R₁ is α -fluorine. R₁ is β -fluorine. The stereochemistry of R₁ in the α - and β -configurations is shown below:

R₁ alpha (α -) configurationR₁ beta (β -) configuration

In another aspect, the present invention provides compounds where R₂ is α -hydroxy. R₂ is hydrogen. R₁ is β -hydroxy and R₂ is α -hydroxy. R₁ is β -hydroxy and R₂ is H. R₁ is α -hydroxy and R₂ is H.


In another aspect, the present invention provides compounds where at least one of R₁ or R₂ is hydroxy. In another aspect, at least one of R₁ or R₂ is hydrogen. R₁ and R₂ are the same. R₁ and R₂ are each α -hydroxy. R₁ and R₂ are each hydrogen.

In another aspect, the present invention provides compounds where R₃ is hydrogen, hydroxyl, NH(CH₂)_mSO₃H, or NH(CH₂)_nCO₂H. R₃ is hydroxyl. R₃ is not hydroxyl. R₃ is

55929-3D1

$\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$. In another aspect, R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$ and m is 2. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$. In another aspect, R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and n is 1.

In another aspect, R_4 is hydrogen or alkyl. R_4 is hydrogen. R_4 is lower alkyl. R_4 is lower alkyl and the lower alkyl group is in the alpha configuration. R_4 in the alpha configuration means that R_4 has the stereochemistry shown in the structure below.



R_4 alpha (α -) configuration

In another aspect, R_4 is halogen. R_4 is fluorine. R_4 is halogen and the halogen is in the alpha configuration. R_4 is α -fluorine.

In another aspect, R_4 is methyl or ethyl. R_4 is methyl. R_4 is ethyl. R_4 is α -methyl. R_4 is α -ethyl. R_3 and R_4 are the same. R_3 and R_4 are different. R_3 and R_4 are each hydrogen. R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$ and R_4 is hydrogen. R_3 is hydroxyl and R_4 is hydrogen. In another aspect, R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, R_4 is hydrogen and m is 2. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and R_4 is hydrogen. In another aspect, R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$, R_4 is hydrogen and n is 1.

In another aspect, R_3 is OH and R_4 is alkyl. R_3 is OH and R_4 is lower alkyl. Lower alkyl is in the alpha configuration. R_3 is OH and R_4 is methyl. R_3 is OH and R_4 is ethyl. R_3 is OH and R_4 is α -methyl. R_3 is OH and R_4 is α -ethyl.

In another aspect, R_5 is unsubstituted or substituted alkyl. R_5 is unsubstituted or substituted lower alkyl. R_5 is in the S-configuration. R_5 is in the R-configuration. R_5 is methyl or ethyl. R_5 is S-methyl. R-methyl. R_5 is S-ethyl. R-ethyl. R_5 is alkyl substituted with phenyl. R_5 is lower alkyl substituted with phenyl. R_5 is benzyl. R_5 is S-benzyl. R_5 is R-benzyl.


In another aspect, R_5 is aryl. R_5 is phenyl.

In another aspect, R_4 and R_5 are each unsubstituted alkyl. R_4 and R_5 are each lower unsubstituted alkyl. R_4 and R_5 are each lower unsubstituted alkyl and R_5 is in the S-configuration. R_4 and R_5 are each lower unsubstituted alkyl and R_4 is in the alpha configuration. In another aspect, R_4 is not hydrogen.


In another aspect, R_4 and R_5 are each lower unsubstituted alkyl and R_1 is α -hydroxy. R_4 and R_5 are each lower unsubstituted alkyl and R_2 is hydrogen. R_4 and R_5 are each lower unsubstituted alkyl, R_1 is α -hydroxy, and R_2 is hydrogen.

55929-3D1

In another aspect, R_5 and R_6 taken together with the carbons to which they are attached form a ring size of 3, 4, 5, or 6 atoms. R_5 and R_6 taken together with the carbons to which they are attached form a 3-membered ring. The 3-membered ring has the following



The 3-membered ring has the following stereochemistry:



In another aspect, R_1 , R_2 , R_3 , and R_4 are hydrogen. R_2 , R_3 , and R_4 are hydrogen. R_2 and R_3 are hydrogen. In another aspect, R_1 , R_2 , and R_4 are hydrogen and R_3 is OH. R_2 and R_4 are hydrogen and R_3 is OH. R_2 is hydrogen and R_3 is OH.

In another aspect, at least one of R_1 , R_2 , R_3 , or R_4 is hydrogen.

10 In another aspect, at least two of R_1 , R_2 , R_3 , or R_4 are hydrogen.

In another aspect, at least three of R_1 , R_2 , R_3 , or R_4 are hydrogen.

In another aspect, R_1 , R_2 , R_3 , and R_4 are hydrogen.

In another aspect, at least one of R_1 , R_2 , or R_4 is hydrogen and R_3 is OH.

In another aspect, at least two of R_1 , R_2 , or R_4 are hydrogen and R_3 is OH.

15 In another aspect, all of R_1 , R_2 , and R_4 are hydrogen and R_3 is OH.


In another aspect, the present invention does not include when R_5 is methyl, R_4 is hydrogen, and R_2 is H or OH.

In another aspect of the present invention, the compound is selected from Compounds **Ia**, **Ib**, **Ic**, **Ig**, **Ih**, **Ii**, **Io**, **Ip**, **Iq**, **Ia1**, **Ib1**, **Ic1**, **Ig1**, **Ih1**, **Ii1**, **Il1**, **Im1**, **In1**, **Io1**, **Ip1**, **Iq1**, **Ia2**, **Ib2**, **Ic2**, **Id2**, **Ie2**, **If2**, **Ig2**, **Ih2**, **Ii2**, **Il2**, **Im2**, **In2**, **Io2**, **Ip2**, **Iq2**, **Ia3**, **Ib3**, **Ic3**, **Id3**, **Ie3**, **If3**, **Ig3**, **Ih3**, **Ii3**, **Il3**, **Im3**, **In3**, **Ia4**, **Ib4**, **Ic4**, **Id4**, **Ie4**, **If4**, **Ig4**, **Ih4**, **Ii4**, **Il4**, **Im4**, **In4**, **Ia5**, **Ib5**, **Ic5**, **Id5**, **Ie5**, **If5**, **Ig5**, **Ih5**, **Ii5**, **Il5**, **Im5** and **In5**.

25 In another aspect of the present invention, the compound is not selected from Compounds **Id**, **Ie**, **If**, **Id1**, **Il**, **Im**, and **In**. In another aspect, the compound is not selected from **Iel** and **If1**.


In another aspect of the present invention, the compounds are modulate the activity of TGR5 receptor. The present invention includes compounds that are TGR5 receptor agonists. In one aspect, the present invention includes compounds that are highly selective for the TGR5 receptor over the FXR receptor.

Another aspect of the present invention includes a composition or medicament comprising a compound of formula I:



- (I), or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, and at least one pharmaceutically acceptable excipient wherein R_1 is hydrogen, hydroxy, or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, $NH(CH_2)_mSO_3H$, or $NH(CH_2)_nCO_2H$; R_4 is hydrogen, unsubstituted or substituted alkyl, or halogen; R_5 is unsubstituted or substituted lower alkyl, or aryl; R_6 is hydrogen or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. In another aspect, the present invention includes a composition or medicament comprising a compound of formula I with proviso that when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxy or $NHCH_2CH_2SO_3H$, then R_4 is not hydrogen.

- Another aspect of the invention includes a method of treating and/or preventing disease in a subject, comprising administering an effective amount of a compound of formula I to a subject in need thereof:



- (I) or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein: R_1 is hydrogen, hydroxy, or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, $NH(CH_2)_mSO_3H$, or $NH(CH_2)_nCO_2H$; R_4 is hydrogen, unsubstituted or substituted alkyl, or halogen; R_5 is unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. The present invention includes a method for treatment and/or prevention of a disease or condition that involves modulation of the TGR5 receptor.

The present invention includes a method comprising administering a compound of formula I. The present invention includes a method comprising administering a therapeutically effective amount of a compound of formula I. The present invention includes a method comprising administering a prophylactically effective amount of a compound of formula I.

In one aspect of the present invention, the compound is a regulator of a physiological function in which TGR5 is involved, or an agent for the prophylaxis or treatment of pathology or disease in which TGR5 is involved. In another aspect, the compound is a cytokine production suppressor. In another aspect, the compound is a GLP-1 secretion promoter or an insulin secretagogue. In another aspect, the compound is an anorectic agent, a pancreatic regenerator, a pancreatic β cell differentiation promoter, a pancreatic β cell growth promoter or an insulin sensitizer. In another aspect, the compound is an agent for the prophylaxis or treatment of cardiac failure, cardiac infarction, acute kidney failure, angina pectoris, arrhythmia, bronchial asthma, chronic obstructive pulmonary disease, arteriosclerosis, rheumatoid arthritis, diabetes, obesity, insulin hyposecretion, pancreatic fatigue, gastric ulcer, ulcerative colitis, allergy, osteoarthritis, erythematosus, excessive immune reaction after transplantation or infectious disease, or an immunosuppressant.

The present invention includes a method of treatment of a subject affected by a disease wherein the TGR5 receptor is involved, which method includes administration to a subject a compound of formula I.

In another aspect, the invention includes a method for treatment or prevention of a metabolic disease by administering a compound of the invention. In one aspect, the metabolic disease is obesity. In another aspect, the metabolic disease is insulin sensitivity. In another aspect, the metabolic disease is diabetes. In another aspect, the metabolic disease is insulin hyposecretion. In another aspect, the metabolic disease is pancreatic fatigue.

The present invention includes a method for treatment and/or prevention of an inflammatory disease by administering a compound of the invention. In one aspect, the inflammatory disease is rheumatoid arthritis. In another aspect, the inflammatory disease is allergy.

The present invention includes the subject is human.

The present invention includes use of the compounds of the invention for known traditional uses of the bile acids. Traditional uses of the bile acids include treatment of

cholelithiasis, bile desaturation, cholesterol metabolism, and use as an antioxidant, radical scavenger, anticholestatic, enduretic, anti-dyslipemic, and hepatocycle protector.


In another aspect, the method comprises administering a compound of formula I where if R_5 is methyl; R_1 is hydroxy; R_3 is hydroxyl or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not
5 hydrogen.

In another aspect, the method includes administering a compound selected from Compounds Ia, Ib, Ic, Id, Ie, If, Ig, Ih, Ii, Il, Im, In, Io, Ip, Iq, Ia1, Ib1, Ic1, Id1, Ie1, If1, Ig1, Ih1, Ii1, Il1, Im1, In1, Io1, Ip1, Iq1, Ia2, Ib2, Ic2, Id2, Ie2, If2, Ig2, Ih2, Ii2, Il2, Im2, In2, Io2, Ip2, Iq2, Ia3, Ib3, Ic3, Id3, Ie3, If3, Ig3, Ih3, Ii3, Il3, Im3, In3, Ia4, Ib4, Ic4, Id4,
10 Ie4, If4, Ig4, Ih4, Ii4, Il4, Im4, In4, Ia5, Ib5, Ic5, Id5, Ie5, If5, Ig5, Ih5, Ii5, Il5, Im5 and In5.

In another aspect, the method includes administering a compound selected from Compounds Ia, Ib, Ic, Ig, Ih, Ii, Io, Ip, Iq, Ia1, Ib1, Ic1, Ig1, Ih1, Ii1, Il1, Im1, In1, Io1, Ip1, Iq1, Ia2, Ib2, Ic2, Id2, Ie2, If2, Ig2, Ih2, Ii2, Il2, Im2, In2, Io2, Ip2, Iq2, Ia3, Ib3, Ic3,
15 Id3, Ie3, If3, Ig3, Ih3, Ii3, Il3, Im3, In3, Ia4, Ib4, Ic4, Id4, Ie4, If4, Ig4, Ih4, Ii4, Il4, Im4, In4, Ia5, Ib5, Ic5, Id5, Ie5, If5, Ig5, Ih5, Ii5, Il5, Im5 and In5.

In one aspect, the method of the present invention does not include administering a compound selected from Compounds Id, Ie, If, Id1, Il, Im, and In. In another aspect, the method of the present invention does not include administering Ie1 and If1.

20 Another aspect of the present invention is the use of a compound of formula I:




(I) or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, and at least one pharmaceutically acceptable excipient wherein R_1 is hydrogen, hydroxy, or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, or $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$; R_4 is hydrogen, alkyl, or halogen; R_5 is unsubstituted
25 or substituted lower alkyl, or aryl; R_6 is hydrogen or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5 for the preparation of a medicament for the treatment and/or prevention of disease wherein a modulation of TGR5 is desired. In another

aspect, the present invention includes a use for the preparation of a medicament for the treatment and/or prevention of disease wherein a modulation of TGR5 is desired comprising a compound of formula I with proviso that when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxy or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not hydrogen.

5 The present invention also provides radiolabeled compounds of formula I. Radiolabeled compounds of formula I can be prepared using conventional techniques. For example, radiolabeled compounds of formula I can be prepared by reacting the compound of formula I with tritium gas in the presence of an appropriate catalyst to produce radiolabeled compounds of formula I. In one embodiment, the compounds of formula I are tritiated.

10 Another aspect of the invention includes compounds of Formula IA:



(IA) or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof, wherein: R_1 is hydrogen, hydroxy, substituted or unsubstituted alkyl, or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, hydrogen, $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, or $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$; R_4 is hydrogen, substituted or unsubstituted alkyl, or halogen; R_5 is
15 unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen, unsubstituted or substituted alkyl, or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. In one aspect, when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxy or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not hydrogen.

20 In one aspect, R_1 is hydrogen or hydroxy. R_1 is hydroxy. R_1 is hydrogen. R_1 is hydroxy and R_2 is α -hydroxy. R_1 is hydroxy and R_2 is H. R_1 is hydroxy and R_2 is H. At least one of R_1 or R_2 is hydroxy. At least one of R_1 or R_2 is hydrogen. R_1 and R_2 are the same. R_1 is hydroxyl and R_2 is α -hydroxy. R_1 and R_2 are each hydrogen.

In one aspect, R_3 is hydrogen, hydroxy, $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, or $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$. R_3 is
25 hydroxy. R_3 is not hydroxy. R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$. R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$ and m is 2. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and n is 1.

In another aspect, R_4 is hydrogen or unsubstituted alkyl. R_4 is hydrogen. R_4 is unsubstituted alkyl. R_4 is unsubstituted alkyl. R_4 is methyl or ethyl. R_4 is methyl. R_4 is

55929-3D1

ethyl. R_3 and R_4 are the same. R_3 and R_4 are different. R_3 and R_4 are each hydrogen. R_3 is OH and R_4 is hydrogen.

In another aspect, R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$ and R_4 is hydrogen. R_3 is $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, R_4 is hydrogen, and m is 2. R_3 is $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$ and R_4 is hydrogen. R_3 is
 5 $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$, R_4 is hydrogen, and n is 1. R_3 is OH and R_4 is unsubstituted alkyl. R_3 is OH and R_4 is unsubstituted alkyl. R_3 is OH and R_4 is methyl. R_3 is OH and R_4 is ethyl. R_3 is OH and R_4 is methyl.

In one aspect, R_5 is unsubstituted or substituted alkyl. R_5 is in the S-configuration. R_5 is in the R-configuration. R_5 is methyl or ethyl. R_5 is S-methyl. R_5 is R-methyl. R_5 is S-ethyl. R_5 is R-ethyl. R_5 is substituted with phenyl. R_5 is benzyl. R_5 is S-benzyl. R_5 is R-benzyl. In another aspect, R_5 is aryl. For example, R_5 is phenyl.

R_4 and R_5 are each unsubstituted alkyl. R_4 and R_5 are each unsubstituted alkyl, further wherein R_5 is in the S-configuration. R_4 and R_5 are each unsubstituted alkyl. R_4 and R_5 are each unsubstituted alkyl and R_1 is hydroxy. R_4 and R_5 are each unsubstituted alkyl and
 15 R_2 is hydrogen. R_4 and R_5 are each unsubstituted alkyl, R_1 is hydroxy, and R_2 is hydrogen.

In one aspect, R_1 , R_2 , R_3 , and R_4 are hydrogen. R_2 , R_3 , and R_4 are hydrogen. R_2 and R_3 are hydrogen. At least one of R_1 , R_2 , R_3 , or R_4 is hydrogen. At least two of R_1 , R_2 , R_3 , or R_4 is hydrogen. At least three of R_1 , R_2 , R_3 , or R_4 is hydrogen. R_1 , R_2 , R_3 , and R_4 are hydrogen.

20 In one aspect, R_1 , R_2 , and R_4 are hydrogen and R_3 is OH. R_2 and R_4 are hydrogen and R_3 is OH. R_2 is hydrogen and R_3 is OH. At least one of R_1 , R_2 , or R_4 is hydrogen and R_3 is OH. At least two of R_1 , R_2 , or R_4 is hydrogen and R_3 is OH. At least three of R_1 , R_2 , or R_4 is hydrogen and R_3 is OH. All of R_1 , R_2 , and R_4 are hydrogen and R_3 is OH.

In another aspect, at least one of R_1 or R_7 is unsubstituted alkyl. At least one of R_1 or R_7 is methyl. At least one of R_1 or R_7 is ethyl. At least one of R_1 or R_7 is propyl. Both R_1 and R_7 are unsubstituted alkyl. Both R_1 and R_7 are methyl. Both R_1 and R_7 are ethyl. R_1 and R_7 are the same. R_1 and R_7 are different. R_7 is hydrogen. R_7 is hydroxy. One of R_1 or R_7 is unsubstituted alkyl and the remaining R_1 or R_7 is hydrogen. One of R_1 or R_7 is unsubstituted alkyl and the remaining R_1 or R_7 is hydroxy. At least one of R_1 or R_7 is unsubstituted alkyl
 25 and R_5 is unsubstituted or substituted alkyl. At least one of R_1 or R_7 is methyl and R_5 is methyl.

Both R_1 and R_5 are unsubstituted alkyl and R_7 is hydroxy. Both R_7 and R_5 are unsubstituted alkyl and R_1 is hydroxy. R_1 or R_7 is unsubstituted alkyl and R_5 is unsubstituted

55929-3D1

or substituted alkyl further wherein R_5 is in the S-configuration. R_1 or R_7 is unsubstituted alkyl and R_5 is unsubstituted or substituted alkyl, further wherein R_5 is in the R-configuration.


In another aspect, R_1 is hydroxy and R_7 is methyl. R_1 is methyl and R_7 is hydroxy. R_6 is unsubstituted alkyl. R_6 is methyl. R_6 is ethyl. R_2 and R_6 are each hydrogen. R_2 and R_6 are hydrogen and R_5 is unsubstituted alkyl. R_2 and R_6 are hydrogen, R_5 is unsubstituted alkyl, and at least one of R_1 or R_7 is unsubstituted alkyl.

In one aspect, the compound is selected from Compounds **Ia6, Ib6, Ic6, Ig6, Ih6, Ii6, Io6, Ip6, Iq6, Ia7, Ib7, Ic7, Ig7, Ih7, Ii7, Il7, Im7, In7, Io7, Ip7, Iq7, Ia8, Ib8, Ic8, Id8, Ie8, If8, Ig8, Ih8, Ii8, Il8, Im8, In8, Io8, Ip8, Iq8, Ia9, Ib9, Ic9, Id9, Ie9, If9, Ig9, Ih9, Ii9, Il9, Im9, In9, Ia10, Ib10, Ic10, Id10, Ie10, If10, Ig10, Ih10, Ii10, Il10, Im10, In10, Ia11, Ib11, Ic11, Id11, Ie11, If11, Ig11, Ih11, Ii11, Il11, Im11 and In11.**

In another aspect of the present invention, when R_2 , R_4 and R_6 are each hydrogen, R_3 is hydroxyl, and one of R_1 and R_7 is hydrogen or hydroxyl, then the other R_1 or R_7 is not methyl. In another aspect, when R_2 is α -OH; R_3 is hydroxyl; R_4 and R_6 are each hydrogen; and one of R_1 and R_7 is hydrogen or hydroxyl, then the other R_1 or R_7 is not methyl. In another aspect, the present invention does not include the following compounds: $3\alpha,7\alpha$ -dihydroxy- 7β -methyl- 5β -cholanoic acid, $3\alpha,7\beta$ -dihydroxy- 7α -methyl- 5β -cholanoic acid, 3α -hydroxy- 7ϵ -methyl- 5β -cholanoic acid, $3\alpha,7\beta,12\alpha$ -trihydroxy- 7α -methyl- 5β -cholan-24-oic acid; $3\alpha,7\alpha,12\alpha$ -trihydroxy- 7β -methyl- 5β -cholan-24-oic acid; and $3\alpha,12\alpha$ -dihydroxy- 7ϵ -methyl- 5β -cholan-24-oic acid.

In another aspect of the present invention, when R_3 is hydroxyl and one of R_1 and R_7 is methyl and the other R_1 and R_7 is hydrogen or hydroxyl, then R_2 , R_4 and R_6 are not all hydrogen. In another aspect, when R_2 is α -OH, R_3 is hydroxyl, and one of R_1 and R_7 is methyl and the other R_1 and R_7 is hydrogen or hydroxyl, then R_4 and R_6 are not all hydrogen.

Another aspect of the invention includes a composition or medicament comprising a compound of formula IA:




(IA), or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof, and at least one pharmaceutically acceptable excipient wherein:

- 5 R_1 is hydrogen, hydroxy, substituted or unsubstituted alkyl or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, or $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$; R_4 is hydrogen, substituted or unsubstituted alkyl, or halogen; R_5 is unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen, unsubstituted or substituted alkyl, or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; and m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5.

In one aspect of the invention, when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxyl or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not hydrogen.

- 15 Another aspect of the invention includes a method of treating and/or preventing disease in a subject, comprising administering a compound of formula IA to a subject in need thereof:



(IA) or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof, wherein: R_1 is hydrogen, hydroxy, substituted or unsubstituted alkyl, or halogen; R_2 is hydrogen or α -hydroxy; R_3 is hydroxy, $\text{NH}(\text{CH}_2)_m\text{SO}_3\text{H}$, or

- 20 $\text{NH}(\text{CH}_2)_n\text{CO}_2\text{H}$; R_4 is hydrogen, substituted or unsubstituted alkyl, or halogen; R_5 is unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen, unsubstituted or substituted alkyl, or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; m is an integer 0, 1, 2, 3, 4, or 5; and n is an integer 0, 1, 2, 3, 4, or 5. In one aspect, treatment and/or

prevention of the disease involves modulation of TGR5 receptor in a subject. The disease is obesity. The disease is insulin sensitivity. The disease is inflammation. The subject is human.

In one aspect, when R_5 is methyl; R_1 is hydroxy; R_3 is hydrogen or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$,
 5 then R_4 is not hydrogen. In another aspect, the method relates administering a compound selected from Compounds Ia6, Ib6, Ic6, Id6, Ie6, If6, Ig6, Ih6, Ii6, Il6, Im6, In6, Io6, Ip6, Iq6, Ia7, Ib7, Ic7, Id7, Ie7, If7, Ig7, Ih7, Ii7, Il7, Im7, In7, Io7, Ip7, Iq7, Ia8, Ib8, Ic8, Id8, Ie8, If8, Ig8, Ih8, Ii8, Il8, Im8, In8, Io8, Ip8, Iq8, Ia9, Ib9, Ic9, Id9, Ie9, If9, Ig9, Ih9, Ii9, Il9, Im9, In9, Ia10, Ib10, Ic10, Id10, Ie10, If10, Ig10, Ih10, Ii10, Im10, In10, Ia11, Ib11,
 10 Ic11, Id11, Ie11, If11, Ig11, Ih11, Ii11, Il11, Im11 and In11.

In another aspect, the method relates to administering a compound is selected from Compounds Ia6, Ib6, Ic6, Ig6, Ih6, Ii6, Io6, Ip6, Iq6, Ia7, Ib7, Ic7, Ig7, Ih7, Ii7, Il7, Im7, In7, Io7, Ip7, Iq7, Ia8, Ib8, Ic8, Id8, Ie8, If8, Ig8, Ih8, Ii8, Il8, Im8, In8, Io8, Ip8, Iq8, Ia9, Ib9, Ic9, Id9, Ie9, If9, Ig9, Ih9, Ii9, Il9, Im9, In9, Ia10, Ib10, Ic10, Id10, Ie10, If10,
 15 Ig10, Ih10, Ii10, Il10, Im10, In10, Ia11, Ib11, Ic11, Id11, Ie11, If11, Ig11, Ih11, Ii11, Il11, Im11 and In11.


In another aspect, the method relates to administering a compound of formula IA with the proviso that when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxyl or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not hydrogen.

20 The present invention includes a method of treating and/or preventing disease in a subject, comprising administering a compound of formula IA to a subject in need thereof. The disease is selected from cholestasis or bile desaturation. The present invention relates to the use of compound of formula IA in the preparation or manufacture of a medicament for treating and/or preventing disease involving the modulation of the TGR5 receptor in a
 25 subject, comprising administering said compound to a subject in need thereof.

The present invention also provides radiolabeled compounds of formula IA. Radiolabeled compounds of formula IA can be prepared using conventional techniques. For example, radiolabeled compounds of formula IA can be prepared by reacting the compound of formula IA with tritium gas in the presence of an appropriate catalyst to produce
 30 radiolabeled compounds of formula IA. In one embodiment, the compounds of formula IA are tritiated.

55929-3D1

Another aspect of the present invention includes a compound of Formula II:



(II) or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof, wherein: R_1 is hydrogen, hydroxy, substituted or unsubstituted alkyl, or halogen; R_2 is hydrogen or α -hydroxy; R_4 is hydrogen, substituted or unsubstituted alkyl, or halogen; R_5 is unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen, unsubstituted or substituted alkyl, or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; and R_8 is hydrogen, substituted or unsubstituted alkyl. In one aspect, when R_5 is methyl and R_1 is hydroxyl, then R_4 is not hydrogen.

10 In one aspect, R_1 is hydrogen or hydroxy. R_1 is hydroxy. R_1 is hydrogen. R_1 is β -hydroxy. R_2 is α -hydroxy. R_1 is hydroxy and R_2 is α -hydroxy. R_1 is hydroxy and R_2 is H. At least one of R_1 or R_2 is hydroxy. At least one of R_1 or R_2 is hydrogen. R_1 and R_2 are the same. R_1 is hydroxyl and R_2 is α -hydroxy. R_1 and R_2 are each hydrogen.

15 In another aspect, R_4 is hydrogen or unsubstituted alkyl. R_4 is hydrogen. R_4 is unsubstituted alkyl. R_4 is unsubstituted alkyl. R_4 is methyl or ethyl. R_4 is methyl. R_4 is ethyl.

In one aspect, R_5 is unsubstituted or substituted alkyl. R_5 is in the S-configuration. R_5 is in the R-configuration. R_5 is methyl or ethyl. R_5 is S-methyl. R_5 is R-methyl. R_5 is S-ethyl. R_5 is R-ethyl. R_5 is substituted with phenyl. R_5 is benzyl. R_5 is S-benzyl. R_5 is R-benzyl. R_5 is aryl. R_5 is phenyl. R_4 and R_5 are each unsubstituted alkyl. R_4 and R_5 are each unsubstituted alkyl, further wherein R_5 is in the S-configuration. R_4 and R_5 are each unsubstituted alkyl and R_1 is hydroxy. R_4 and R_5 are each unsubstituted alkyl and R_2 is hydrogen. R_4 and R_5 are each unsubstituted alkyl, R_1 is hydroxy, and R_2 is hydrogen.

25 In one aspect, R_1 , R_2 , and R_4 are hydrogen. R_2 and R_4 are hydrogen. R_2 is hydrogen. At least one of R_1 , R_2 , or R_4 is hydrogen. At least two of R_1 , R_2 , or R_4 is hydrogen. R_1 , R_2 , and R_4 are hydrogen.


In one aspect, R_1 or R_7 is unsubstituted alkyl. R_1 or R_7 is methyl. R_1 or R_7 is ethyl. R_1 or R_7 is propyl. Both R_1 and R_7 are unsubstituted alkyl. R_7 is hydrogen. R_7 is hydroxy.

One of R₁ or R₇ is unsubstituted alkyl and the remaining R₁ or R₇ is hydrogen. One of R₁ or R₇ is unsubstituted alkyl and the remaining R₁ or R₇ is hydroxy. At least one of R₁ or R₇ is unsubstituted alkyl and R₅ is unsubstituted or substituted alkyl. At least one of R₁ or R₇ is methyl and R₅ is methyl. R₇ is hydroxy and both R₁ and R₅ are unsubstituted alkyl. R₁ is hydroxy and both R₇ and R₅ are unsubstituted alkyl. At least one of R₁ or R₇ is unsubstituted alkyl and R₅ is unsubstituted or substituted alkyl, further wherein R₅ is in the S-configuration. At least one of R₁ or R₇ is unsubstituted alkyl and R₅ is unsubstituted or substituted alkyl, further wherein R₅ is in the R-configuration. R₇ is hydroxy and both R₁ and R₅ are unsubstituted alkyl, further wherein R₅ is in the S-configuration. R₇ is hydroxy and both R₁ and R₅ are unsubstituted alkyl, further wherein R₅ is in the R-configuration. R₁ is hydroxy and both R₇ and R₅ are unsubstituted alkyl, further wherein R₅ is in the S-configuration. R₁ is hydroxy and both R₇ and R₅ are unsubstituted alkyl, further wherein R₅ is in the R-configuration. R₁ is hydroxy and R₇ is methyl. R₁ is methyl and R₇ is hydroxy.

In another aspect, R₆ is unsubstituted alkyl. R₆ is methyl. R₆ is ethyl. R₈ is hydrogen. R₈ is unsubstituted alkyl. R₈ is methyl. R₈ is ethyl. R₂ is α -hydroxy and R₈ is unsubstituted alkyl.

In another aspect of the invention, the compound is selected from Compounds Ia12, Ib12, Ic12, Ig12, Ih12, Ii12, Io12, Ip12, Iq12, Ia13, Ib13, Ic13, Ig13, Ih13, Ii13, Il13, Im13, In13, Io13, Ip13, Iq13, Ia14, Ib14, Ic14, Id14, Ie14, If14, Ig14, Ih14, Ii14, Il14, Im14, In14, Io14, Ip14, Iq14, Ia15, Ib15, Ic15, Id15, Ie15, If15, Ig15, Ih15, Ii15, Il15, Im15, In15, Ia16, Ib16, Ic16, Id16, Ie16, If16, Ig16, Ih16, Ii16, Il16, Im16, In16, Ia17, Ib17, Ic17, Id17, Ie17, If17, Ig17, Ih17, Ii17, Il17, Im17 and In17.

Another aspect of the invention includes a composition or medicament comprising a compound of formula II:




25

(II), or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof, and at least one pharmaceutically acceptable excipient wherein: R₁ is hydrogen, hydroxy, substituted or unsubstituted alkyl or halogen; R₂ is hydrogen or α -hydroxy; R₄ is hydrogen, substituted or unsubstituted alkyl, or halogen; R₅ is unsubstituted

or substituted alkyl, or aryl; R_6 is hydrogen, unsubstituted or substituted alkyl, or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; and R_8 is hydrogen or substituted or unsubstituted alkyl. In one aspect, when R_5 is methyl, R_1 is hydroxyl, and R_3 is hydroxyl or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not hydrogen.

In one aspect the invention includes a method of treating and/or preventing disease in a subject, comprising administering a compound of formula II to a subject in need thereof:



or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein:

R_1 is hydrogen, hydroxy, substituted or unsubstituted alkyl, or halogen; R_2 is hydrogen or α -hydroxy; R_4 is hydrogen, substituted or unsubstituted alkyl, or halogen; R_5 is unsubstituted or substituted alkyl, or aryl; R_6 is hydrogen, unsubstituted or substituted alkyl, or R_5 and R_6 taken together with the carbons to which they are attached form a ring of size 3, 4, 5, or 6 atoms; R_7 is hydrogen, substituted or unsubstituted alkyl, or hydroxy; and R_8 is hydrogen, substituted or unsubstituted alkyl.

The invention includes a method of treatment of a disease which involves modulation of the TGR5 receptor. The invention includes a method of prevention of a disease which involves modulation of the TGR5 receptor. The disease is obesity. The disease is insulin sensitivity. The disease is inflammation. The subject is human. In one aspect, when R_5 is methyl; R_1 is hydroxy; R_3 is hydroxyl or $\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, then R_4 is not hydrogen.

One aspect of the invention relates to a method of administering a compound is selected from Compounds Ia12, Ib12, Ic12, Id12, Ie12, If12, Ig12, Ih12, Ii12, Ij12, Im12, In12, Io12, Ip12, Iq12, Ia13, Ib13, Ic13, Id13, Ie13, If13, Ig13, Ih13, Ii13, Il13, Im13, In13, Io13, Ip13, Iq13, Ia14, Ib14, Ic14, Id14, Ie14, If14, Ig14, Ih14, Ii14, Il14, Im14, In14, Io14, Ip14, Iq14, Ia15, Ib15, Ic15, Id15, Ie15, If15, Ig15, Ih15, Ii15, Il15, Im15, In15, Ia16, Ib16, Ic16, Id16, Ie16, If16, Ig16, Ih16, Ii16, Im16, In16, Ia17, Ib17, Ic17, Id17, Ie17, If17, Ig17, Ih17, Ii17, Il17, Im17 and In17.

Another aspect of the invention relates to a method of administering a compound selected from Compounds **Ia12**, **Ib12**, **Ic12**, **Ig12**, **Ih12**, **Ii12**, **Io12**, **Ip12**, **Iq12**, **Ia13**, **Ib13**, **Ic13**, **Ig13**, **Ih13**, **Ii13**, **Il13**, **Im13**, **In13**, **Io13**, **Ip13**, **Iq13**, **Ia14**, **Ib14**, **Ic14**, **Id14**, **Ie14**, **If14**, **Ig14**, **Ih14**, **Ii14**, **Il14**, **Im14**, **In14**, **Io14**, **Ip14**, **Iq14**, **Ia15**, **Ib15**, **Ic15**, **Id15**, **Ie15**, **If15**, **Ig15**, **Ih15**, **Ii15**, **Il15**, **Im15**, **In15**, **Ia16**, **Ib16**, **Ic16**, **Id16**, **Ie16**, **If16**, **Ig16**, **Ih16**, **Ii16**, **Il16**, **Im16**, **In16**, **Ia17**, **Ib17**, **Ic17**, **Id17**, **Ie17**, **If17**, **Ig17**, **Ih17**, **Ii17**, **Il17**, **Im17** and **In17**.

Another aspect of the invention includes a method related to administering a compound of formula II with the proviso that when R_5 is methyl, and R_1 is hydroxyl, then R_4 is not hydrogen.

10 The present invention includes a method of treating disease in a subject, comprising administering a therapeutically effective amount of the compound of formula II to a subject in need thereof. The present invention includes a method of preventing disease in a subject, comprising administering a prophylactically effective amount of the compound of formula II to a subject in need thereof. The disease is selected from cholestasis or bile desaturation. The
15 present invention relates to use of a compound of formula II in preparation of a medicament for treating and/ or preventing a disease involving the modulation of the TGR5 receptor in a subject, comprising administering an effective amount i.e. a therapeutically effective or prophylactically effective amount of said compound to a subject in need thereof.

The present invention also provides radiolabeled compounds of formula II.

20 Radiolabeled compounds of formula II can be prepared using conventional techniques. For example, radiolabeled compounds of formula II can be prepared by reacting the compound of formula II with tritium gas in the presence of an appropriate catalyst to produce radiolabeled compounds of formula II. In one embodiment, the compounds of formula II are tritiated.

25 Some representative compounds of the invention are shown below.

The following compounds **Ia-In5** pertain to at least formula I:

Ia: $R_1 = \alpha\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$

Ib: $R_1 = \alpha\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$

Ic: $R_1 = \alpha\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$

30 **Id:** $R_1 = \beta\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$

Ie: $R_1 = \beta\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$

If: $R_1 = \beta\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$

Ig: $R_1 = \alpha\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$

- 33 -

- 34 -

- Ia5:** $R_1 = \alpha\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$
Ib5: $R_1 = \alpha\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$
Ic5: $R_1 = \alpha\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$
Id5: $R_1 = \beta\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$
5 **Ie5:** $R_1 = \beta\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$
If5: $R_1 = \beta\text{-OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$
Ig5: $R_1 = \alpha\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$
Ih5: $R_1 = \alpha\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$
Ii5: $R_1 = \alpha\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$
10 **Ij5:** $R_1 = \beta\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$
Im5: $R_1 = \beta\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$
In5: $R_1 = \beta\text{-OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{NHCH}_2\text{CO}_2\text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$

The following compounds **In6-In11** pertain to at least formula IA:

- 15 **Ia6:** $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ib6: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ic6: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Id6: $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$
Ie6: $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$
20 **If6:** $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$
Ig6: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ih6: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ii6: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ij6: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$
25 **Im6:** $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$
In6: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$
Io6: $R_1 = \text{H}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ip6: $R_1 = \text{H}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Iq6: $R_1 = \text{H}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
30 **Ia7:** $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, $R_4 = \text{H}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ib7: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, $R_4 = \text{H}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$
Ic7: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_3 = \text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$, $R_4 = \text{H}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$

- 36 -

- 37 -

The following compounds **Ia12-In17** pertain to at least formula II:

- Ia12:** R₁= OH, R₂= H, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇= Me, R₈=H
Ib12: R₁= OH, R₂= H, R₄= H, R₅= (*S*)Me, R₆=H, R₇= Me, R₈=H
5 **Ic12:** R₁= OH, R₂= H, R₄= H, R₅= (*R*)Me, R₆=H, R₇= Me, R₈=H
Id12: R₁= Me, R₂= H, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇=OH, R₈=H
Ie12: R₁= Me, R₂= H, R₄= H, R₅= (*S*)Me, R₆=H, R₇=OH, R₈=H
If12: R₁= Me, R₂= H, R₄=H, R₅= (*R*)Me, R₆=H, R₇=OH, R₈=H
Ig12: R₁= OH, R₂= α-OH, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇= Me, R₈=H
10 **Ih12:** R₁= OH, R₂= α-OH, R₄= H, R₅= (*S*)Me, R₆=H, R₇= Me, R₈=H
Ii12: R₁= OH, R₂= α-OH, R₄= H, R₅= (*R*)Me, R₆=H, R₇= Me, R₈=H
II12: R₁= Me, R₂= α-OH, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇=OH, R₈=H
Im12: R₁= Me, R₂= α-OH, R₄= H, R₅= (*S*)Me, R₆=H, R₇=OH, R₈=H
In12: R₁= Me, R₂= α-OH, R₄= H, R₅= (*R*)Me, R₆=H, R₇=OH, R₈=H
15 **Io12:** R₁= H, R₂= H, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇= Me, R₈=H
Ip12: R₁= H, R₂= H, R₄= H, R₅= (*S*)Me, R₆=H, R₇= Me, R₈=H
Iq12: R₁= H, R₂= H, R₄= H, R₅= (*R*)Me, R₆=H, R₇= Me, R₈=H
Ia13: R₁= OH, R₂= H, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇= Me, R₈=H
Ib13: R₁= OH, R₂= H, R₄= H, R₅= (*S*)Me, R₆=H, R₇= Me, R₈=H
20 **Ic13:** R₁= OH, R₂= H, R₄= H, R₅= (*R*)Me, R₆=H, R₇= Me, R₈=H
Id13: R₁= Me, R₂= H, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇=OH, R₈=H
Ie13: R₁= Me, R₂= H, R₄= H, R₅= (*S*)Me, R₆=H, R₇=OH, R₈=H
If13: R₁= Me, R₂= H, R₄=H, R₅= (*R*)Me, R₆=H, R₇=OH, R₈=H
Ig13: R₁= OH, R₂= α-OH, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇= Me, R₈=H
25 **Ih13:** R₁= OH, R₂= α-OH, R₄= H, R₅= (*S*)Me, R₆=H, R₇= Me, R₈=H
Ii13: R₁= OH, R₂= α-OH, R₄= H, R₅= (*R*)Me, R₆=H, R₇= Me, R₈=H
II13: R₁= Me, R₂= α-OH, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇=OH, R₈=H
Im13: R₁= Me, R₂= α-OH, R₄= H, R₅= (*S*)Me, R₆=H, R₇=OH, R₈=H
In13: R₁= Me, R₂= α-OH, R₄= H, R₅= (*R*)Me, R₆=H, R₇=OH, R₈=H
30 **Io13:** R₁= H, R₂= H, R₄= H, R₅= (*S,R*)Me, R₆=H, R₇= Me, R₈=H
Ip13: R₁= H, R₂= H, R₄= H, R₅= (*S*)Me, R₆=H, R₇= Me, R₈=H
Iq13: R₁= H, R₂= H, R₄= H, R₅= (*R*)Me, R₆=H, R₇= Me, R₈=H

- 39 -

55929-3D1

- If16: $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 Ig16: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 Ih16: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 Ii16: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 5 Ij16: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 Im16: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 In16: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 Ia17: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 Ib17: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 10 Ic17: $R_1 = \text{OH}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 Id17: $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 Ie17: $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 If17: $R_1 = \text{Me}$, $R_2 = \text{H}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 Ig17: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 15 Ih17: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 Ii17: $R_1 = \text{OH}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{Me}$, $R_8 = \text{H}$
 Ij17: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S,R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 Im17: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (S)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$
 In17: $R_1 = \text{Me}$, $R_2 = \alpha\text{-OH}$, $R_4 = \alpha\text{-Me}$, $R_5 = (R)\text{Me}$, $R_6 = \text{H}$, $R_7 = \text{OH}$, $R_8 = \text{H}$

20


Citation of publications and patent documents is not intended as an admission that any is pertinent prior art, nor does it constitute any admission
 25 as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description and examples below are for purposes of illustration and not limitation of the claims that follow.

30 **EXAMPLE 1: Synthesis of TGR5 Modulators**

The compounds of the invention, and related derivatives, can be synthesized by methods known to one skilled in the art. Detailed methods for synthesizing these compounds

are described below. *See, also*, WO 02/072598, WO 2004/0007521, EP 1568706 and EP 135782. In the case of the compound where R_1 is hydrogen, R_2 and R_3 are hydroxy and R_4 is a lower alkyl group, the compound of formula (I) can be obtained in accordance with the following scheme:

5 **Scheme 1**



Scheme 1.
(i) 3,4-DHP, *p*-TSA, dioxane, r.t.; (ii) a) LDA, CH₃I, -78 °C; b) HCl, CH₃OH, r.t.; (iii) NaOH, CH₃OH, reflux.

Methyl chenodeoxycholate (1) was protected in 3- and 7-position by treatment with 3,4-dihydro-2H-pyran in dioxane in presence of catalytic amount of *p*-toluenesulfonic acid (*p*-TSA) to give the corresponding 3α,7α-tetrahydropyranyloxy analog (2). Reaction of 2 with methyl iodide (or with an appropriate alkyl halide), at -78 °C using lithium diisopropylamide as a base and tetrahydrofuran (THF) as solvent, followed by treatment with methanolic HCl afforded the corresponding methyl 23-methyl-3α,7α-dihydroxy-5β-cholan-24-oate (3). Hydrolysis with alkali of the methyl ester 3 and purification by flash chromatography yielded the desired 23(*S*)-methyl-3α,7α-dihydroxy-5β-cholan-24-oic acid (Ib) and 23(*R*)-methyl-3α,7α-dihydroxy-5β-cholan-24-oic acid (Ic).

Preparation of 23(*R*)- and 23(*S*)-methyl-3α,7α-dihydroxy-5β-cholan-24-oic acid (Ib, Ic)

a) Methyl 3α,7α-ditetrahydropyranyloxy-5β-cholan-24-oate (2)

20 *p*-Toluenesulfonic acid (78 mg, 0.41 mmol), 3,4-dihydro-2H-pyran (20.1 ml, 0.098 mol) were added to a solution of methyl 3α,7α-dihydroxy-5β-cholan-24-oate (1) (2.0 g, 4.9 mmol) in dioxane (6 mL). The reaction mixture was stirred at room temperature for 15 min. H₂O (50 mL) was then added and the mixture was partially concentrated under vacuum and extracted with EtOAc (3 x 50 mL). The combined organic fractions were washed with brine (1 x 50

mL), dried (Na_2SO_4) and evaporated under vacuum. The residue was purified by chromatography on silica gel column. Elution with light petroleum/ethyl acetate 80/20 afforded 2.5 g of the pure compound 2 (90% yield).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.64 (s, 3H, CH_3 -18), 0.89 (s, 3H, CH_3 -19), 0.92 (d, 3H, CH_3 -21), 3.31-3.67 (m, 4H, $-\text{CH}_2\text{OCH}-$), 3.65 (s, 3H, CO_2CH_3), 3.67 (m, 1H, CH-3), 3.88 (brs, 1H, CH-7), 4.67 (brs, 1H, $-\text{O-CH-O-}$), 4.73 (brs, 1H, $-\text{O-CH-O-}$).

b) Methyl 23(*R,S*)-methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oate (3)

n-Butyl lithium (4.3 mL, 2.2 M solution in hexane) were added dropwise at -78°C to a solution of diisopropylamine (1.4 mL, 10.1 mmol) in dry THF (50 mL). The system was kept to -78°C for additional 30 min and then, methyl 3 α ,7 α ,12 α -tetrahydropyranyloxy-5 β -cholan-24-oate (2) (1.8 g, 3.2 mmol) dissolved in dry THF (14 mL) was added dropwise to the mixture. After 20 min methyl iodide (1.4 mL, 22.0 mmol) dissolved in dry THF (7 mL) was slowly added and the mixture was allowed to warm to room temperature overnight. The solvents were removed under vacuum and acidified by 10% HCl and extracted with EtOAc (5 x 50 mL), washed with 5% $\text{Na}_2\text{S}_2\text{O}_3$ solution (2 x 50 mL), dried (over anhydrous Na_2SO_4), filtered, and evaporated under vacuum. The crude residue was then treated with a solution of 2N HCl in MeOH (50 mL) for 12 h. The residue was evaporated under vacuum and taken up with EtOAc (100 mL), washed with a saturated NaHCO_3 solution (2 x 50 mL), dried (Na_2SO_4) and evaporated under vacuum. The residue was purified by silica gel flash chromatography. Elution with light petroleum/ethyl acetate 70/30 afforded 1.1 g (2.7 mmol) of the pure compound 3 (84% yield).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.62 (s, 3H, CH_3 -18), 0.87 (s, 3H, CH_3 -19), 0.92 (d, 3H, CH_3 -21), 2.38 (m, 1H, CH-23), 3.27-3.40 (m, 1H, CH-3), 3.55 (brs, 1H, CH-7), 3.63 (s, 3H, CO_2CH_3).

c) 23(*R*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (Ib) and 23(*S*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (Ic)

Methyl 23-methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oate 0.97 g (2.3 mmol) was dissolved in MeOH (25 mL) and added with 10% NaOH in MeOH (5.7 mL, 14.2 mmol). The mixture was refluxed for 16 h. The mixture was acidified with 3N HCl and extracted with EtOAc (3 x 20 mL). The combined organic fractions were washed with brine (1 x 50 mL), dried (Na_2SO_4) and evaporated under vacuum. The residue was purified by silica gel flash chromatography.

Elution with $\text{CHCl}_3:\text{MeOH}$ (95/5) afforded 1.5 g (65%) of 23(*S*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid and 330 mg of 23(*R*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid.

23(*S*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (**Ib**): mp: 125-126 °C. $^1\text{H-NMR}$ (5 $\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 0.44 (s, 3H, CH_3 -18), 0.69 (s, 3H, CH_3 -19), 0.73-0.76 (d, 3H CH_3 -21), 0.93-0.97 (d, 3H, $-\text{CH}_3$), 2.36 (m, 1H, CH-23), 3.15-3.38 (m, 1H, CH-3), 3.62 (brs, 1H, CH-7). $^{13}\text{C-NMR}$ ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 11.55, 18.43, 18.87, 20.49, 22.69, 28.15, 28.57, 30.14, 32.65, 34.43, 34.61, 34.94, 35.23, 37.06, 39.17, 39.60, 40.81, 41.40, 42.57, 46.54, 50.29, 56.63, 68.24, 71.62, 179.99.

10

23(*R*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (**Ic**): mp: 163-164 °C. $^1\text{H-NMR}$ ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 0.43 (s, 3H, CH_3 -18), 0.65 (s, 3H, CH_3 -19), 0.65-0.69 (d, 3H CH_3 -21), 0.83-0.86 (d, 3H, $-\text{CH}_3$), 2.20 (m, 1H, CH-23), 3.09-3.15 (m, 1H, CH-3), 3.58 (brs, 1H, CH-7). $^{13}\text{C-NMR}$ ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 11.94, 16.40, 18.30, 20.93, 23.06, 23.89, 28.85, 30.52, 15 33.08, 34.16, 34.91, 35.38, 35.68, 37.14, 39.49, 39.64, 40.04, 40.17, 41.92, 43.05, 50.69, 57.10, 68.51, 72.01, 181.09.

EXAMPLE 2: Preparation of 23(*S*)- and 23(*R*)-methyl-6 α -methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (Ib3**, **Ic3**)**

20 The following compounds were prepared by alkylation of 6 α -methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid according to the procedure of Example 1.

23(*S*)-Methyl-6 α -methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (**Ib3**): mp: 98-100 °C. $^1\text{H-NMR}$ (CDCl_3) δ : 0.63 (s, 3H, CH_3 -18), 0.89 (s, 3H, CH_3 -19), 0.92-1.00 (m, 6H, CH_3 -21 and 25 CH_3 -6), 1.15-1.19 (d, 3H, $-\text{CH}_3$), 2.45-2.73 (m, 1H, CH-23), 3.31-3.52 (m, 1H, CH-3), 3.58 (brs, 1H, CH-7). $^{13}\text{C-NMR}$ (CDCl_3) δ : 11.76, 15.72, 18.58, 18.88, 20.63, 23.11, 23.65, 28.19, 30.21, 30.47, 32.64, 33.79, 33.97, 34.61, 35.42, 35.66, 37.03, 39.60, 40.01, 40.71, 42.71, 47.35, 50.44, 56.60, 72.34, 72.87, 182.37.

30 23(*R*)-Methyl-3 α ,7 α -dihydroxy-5 β -cholan-24-oic acid (**Ic3**): mp: 89-90 °C. $^1\text{H-NMR}$ ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 0.65 (s, 3H, CH_3 -18), 0.88 (s, 3H, CH_3 -19), 0.88-0.92 (m, 3H, CH_3 -6), 0.95-0.99 (d, 3H, CH_3 -21), 1.08-1.14 (d, 3H $-\text{CH}_3$), 2.35 (m, 1H, CH-23), 3.29-3.48 (m, 1H,

CH-3), 3.57 (brs, 1H, CH-7). ^{13}C -NMR ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 11.70, 15.66, 16.02, 18.00, 20.61, 23.09, 23.60, 28.51, 30.39, 32.61, 33.72, 33.92, 35.38, 35.65, 36.33, 39.57, 39.94, 42.77, 47.30, 50.39, 56.53, 72.22, 72.83, 180.50.

5 EXAMPLE 3: Preparation of 23(R)- and 23(S)-methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid (Ih, Ii)

The following compounds were prepared by alkylation of 3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid according to the procedure of Example 1.

10 23(S)-Methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid (Ih): mp: 237-239 °C. ^1H -NMR (CDCl_3) δ : 0.63 (s, 3H, CH_3 -18), 0.87 (s, 3H, CH_3 -19), 0.96-0.98 (m, 3H, CH_3 -21), 1.07-1.11 (d, 3H, $-\text{CH}_3$), 2.44-2.73 (m, 1H, CH-23), 3.35-3.50 (m, 1H, CH-3), 3.82 (brs, 1H, CH-7) 3.95 (brs, 1H, CH-12). ^{13}C -NMR (DMSO) δ : 12.72, 17.60, 19.24, 19.24, 23.00, 23.19, 26.59, 27.78, 28.88, 30.72, 34.77, 35.22, 35.66, 37.19, 41.84, 46.19, 47.27, 49.01, 66.69,
15 70.88, 71.45, 178.25.

23(R)-Methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid (Ii): mp: 221-223 °C. ^1H -NMR (CDCl_3) δ : 0.63 (s, 3H, CH_3 -18), 0.87 (s, 3H, CH_3 -19), 0.96-0.98 (m, 3H, CH_3 -21), 1.07-1.11 (d, 3H, $-\text{CH}_3$), 2.44-2.73 (m, 1H, CH-23), 3.35-3.50 (m, 1H, CH-3), 3.82 (brs, 1H, CH-7) 3.95 (brs, 1H, CH-12). ^{13}C -NMR (DMSO) δ : 12.76, 16.88, 17.31, 23.04, 23.24, 26.62,
20 28.12, 28.94, 30.81, 33.97, 34.80, 35.28, 35.71, 37.20, 41.85, 46.29, 47.44, 66.67, 70.86, 71.45, 178.77.

25 EXAMPLE 4: Preparation of 23(R)- and 23(S)-methyl-6 α -methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid (Ih3, Ii3)

The following compounds were prepared by alkylation of 6 α -methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid according to the procedure of Example 1.

23(S)-Methyl-6 α -methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid (Ih3): mp: 131-134
30 °C. ^1H -NMR ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ : 0.65 (s, 3H, CH_3 -18), 0.87 (s, 3H, CH_3 -19), 0.97-1.00 (m, 3H, CH_3 -21), 1.14-1.18 (d, 3H, $-\text{CH}_3$), 1.23 (m, 1H, CH-6), 2.52 (m, 1H, CH-23), 3.32-3.50 (m, 1H, CH-3), 3.55 (brs, 1H, CH-7) 3.94 (brs, 1H, CH-12). ^{13}C -NMR ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ :

12.43, 145.66, 17.62, 18.92, 22.70, 23.14, 26.21, 27.45, 28.01, 30.03, 33.44, 34.11, 34.42, 35.30, 36.71, 39.97, 40.45, 41.73, 46.45, 47.25, 72.13, 72.76, 73.01, 180.53.

23(*R*)-Methyl-6 α -methyl-3 α ,7 α ,12 α -trihydroxy-5 β -cholan-24-oic acid (**II3**): mp: 109-110
 5 °C. ¹H-NMR (CD₃OD) δ : 0.72 (s, 3H, CH₃-18), 0.91 (s, 3H, CH₃-19), 1.07-1.11 (m, 6H, -CH₃ and CH₃-21), 2.37-2.53 (m, 1H, CH-23), 3.15-3.42 (m, 1H, CH-3), 3.53 (brs, 1H, CH-7), 3.97 (brs, 1H, CH-12). ¹³C-NMR (CD₃OD) δ : 11.61, 15.04, 15.32, 16.15, 22.04, 22.75, 26.27, 27.62, 28.18, 29.61, 32.91, 33.74, 34.31, 35.06, 35.18, 36.56, 39.70, 40.25, 41.68, 46.19, 46.31, 71.76, 71.77, 72.62, 180.11.

10

Example 5: Preparation of 23(*R*)- and 23(*S*)-methyl-3 α -hydroxy-5 β -cholan-24-oic acid (Ip**, **Iq**)**

The following compounds were prepared by alkylation of 3 α -hydroxy-5 β -cholan-24-oic acid according to the procedure of Example 1.

15

23(*S*)-Methyl-3 α -hydroxy-5 β -cholan-24-oic acid (**Ip**): mp: 161-162 °C. ¹H-NMR (CDCl₃+CD₃OD) δ : 0.60 (s, 3H, CH₃-18), 0.88 (s, 3H, CH₃-19), 0.92-1.01 (m, 3H, CH₃-21), 1.13-1.16 (d, 3H, -CH₃), 2.55 (m, 1H, CH-23), 3.60 (m, 1H, CH-3). ¹³C-NMR (CDCl₃+CD₃OD) δ : 11.97, 18.52, 18.87, 20.73, 23.30, 24.14, 26.34, 27.10, 28.15, 30.18,
 20 34.48, 34.50, 35.23, 35.74, 36.06, 37.01, 40.13, 40.34, 40.74, 41.99, 42.68, 56.43, 56.75, 71.70, 181.42.




23(*R*)-Methyl-3 α -hydroxy-5 β -cholan-24-oic acid (**Iq**): mp: 152-153 °C. ¹H-NMR (CDCl₃+CD₃OD) δ : 0.63 (s, 3H, CH₃-18), 0.89 (s, 3H, CH₃-19), 0.94-1.03 (m, 3H, CH₃-21),
 25 2.45 (m, 1H, CH-23), 3.59 (m, 1H, CH-3). ¹³C-NMR (CD₃OD) δ : 11.98, 15.97, 18.00, 20.75, 23.31, 24.14, 26.34, 27.11, 28.48, 30.26, 33.68, 34.50, 35.26, 35.77, 36.15, 36.46, 39.59, 40.13, 40.36, 42.01, 42.79, 56.45, 56.76, 71.71, 181.02.

EXAMPLE 6: Biological Activity

30 The potency and efficacy of the compounds of the invention on TGR5 receptor was evaluated using in vitro assays. See, Kawamata, J. Biol. Chem 2003, Vol. 278 No. 11, p. 9435-9440). Activity on FXR was assayed by fluorescence resonance energy transfer

(FRET) for recruitment of the SRC-1 peptide to human FXR using a cell-free ELISA. See, Blanchard et al. WO 00/37077.

Table 1: EC₅₀ (μM) of Example Compounds on FXR and TGR5 Receptor

Compound	Structure	FXR Data	TGR5 Data
CDCA (<u>Cheno</u> <u>Deoxy</u> <u>Cholic</u> <u>Acid</u>)		EC ₅₀ : 8.6 μM Efficacy: 100%	EC ₅₀ : 4.0 μM Efficacy: 100%
6α-MeCDCA		EC ₅₀ : 0.21 μM Efficacy: 148%	EC ₅₀ : 0.37 μM Efficacy: 119%
23(R+S)-Me-6MeCDCA (I3a)		EC ₅₀ : 15.62 μM Efficacy: 60%	EC ₅₀ : 0.11 μM Efficacy: 123%


- 5 The compounds of the invention are potent and selective TGR5 modulators. The introduction of an alkyl group at the C-23 position of bile acid gives selectivity for the TGR5 receptor with respect to FXR. This is evident by the observation of the biological results obtained for CDCA, 6-MeCDCA and 6,23-diMe-CDCA (23-*R,S* isomers mixture) on FXR and TGR5 as shown in Table 1. 6,23-diMe-CDCA is 100-fold more potent on TGR5 with respect to the
- 10 FXR receptor.

Other Embodiments

- While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the
- 15 invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims. It will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

CLAIMS:


1. A compound of formula II



or a salt, solvate or hydrate thereof, wherein

- 5 R₁ is hydrogen, hydroxyl, substituted or unsubstituted alkyl, or halogen;
- R₂ is hydrogen or α-hydroxy;
- R₄ is hydrogen, substituted or unsubstituted alkyl, or halogen;
- R₅ is unsubstituted or substituted alkyl, or aryl;
- R₆ is hydrogen, unsubstituted or substituted alkyl; or R₅ and R₆ taken together with the carbons to
 10 which they are attached form a ring of size 3, 4, 5, or 6 atoms;
- R₇ is hydrogen, substituted or unsubstituted alkyl, or hydroxyl; and
- R₈ is hydrogen.
2. The compound of claim 1, wherein R₁ is hydrogen or hydroxy.
3. The compound of claim 1 or 2, wherein R₂ is hydrogen.
- 15 4. The compound of any one of claims 1-3, wherein R₄ is hydrogen or unsubstituted alkyl.
5. The compound of any one of claims 1-4, wherein R₅ is unsubstituted alkyl.

6. The compound of any one of claims 1-5, wherein R_6 is hydrogen.
7. The compound of any one of claims 1-6, wherein R_7 is hydrogen, unsubstituted alkyl, or hydroxy.
8. The compound of any one of claims 1-7, wherein R_5 is in the S-configuration.
- 5 9. A pharmaceutical composition comprising the compound of any one of claims 1-8 or a pharmaceutically acceptable salt, solvate or hydrate thereof, and at least one pharmaceutically acceptable excipient.
10. Use of the compound of any one of claims 1-8 or a pharmaceutically acceptable salt, solvate or hydrate thereof, or the composition of claim 9, in the preparation of a medicament
10 for treating and/or preventing a disease involving modulation of the TGR5 receptor.
11. Use of the compound of any one of claims 1-8 or a pharmaceutically acceptable salt, solvate or hydrate thereof, or the composition of claim 9, for treating and/or preventing a disease involving modulation of the TGR5 receptor.
12. The use of claim 10 or 11, wherein the disease is selected from the group
15 consisting of obesity, insulin sensitivity, inflammation, cholestasis, bile desaturation, and diabetes.
13. The use of claim 12, wherein the disease is obesity.
14. The use of claim 12, wherein the disease is insulin sensitivity.
15. The use of claim 12, wherein the disease is inflammation.
- 20 16. The use of claim 12, wherein the disease is cholestasis.
17. The use of claim 12, wherein the disease is bile desaturation.
18. The use of claim 12, wherein the disease is diabetes.



(II)