

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 December 2008 (31.12.2008)

PCT

(10) International Publication Number
WO 2009/000006 A1

(51) International Patent Classification:
C07K 16/00 (2006.01) **C40B 40/02** (2006.01)
C40B 40/10 (2006.01)

(74) Agents: **SCHWARZ, Albin** et al.; Wipplingerstrasse 30,
A-1010 Wien (AT).

(21) International Application Number:
PCT/AT2008/000232

(22) International Filing Date: 26 June 2008 (26.06.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/946,287 26 June 2007 (26.06.2007) US
61/049,826 2 May 2008 (02.05.2008) US

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (for all designated States except US): **F-STAR BIOTECHNOLOGISCHE FORSCHUNGS- UND ENTWICKLUNGSGES.M.B.H.** [AT/AT]; Gastgebasse 5-13, A-1230 Wien (AT).


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and
(75) Inventors/Applicants (for US only): **HIMMLER, Gotfried** [AT/AT]; Donau-Oder-Kanal IV, SW80, A-2301 Gross-Enzersdorf (AT). **RÜCKER, Florian** [AT/AT]; Montigasse 6, A-1170 Wien (AT). **WOZNIAK-KNOPP, Gordana** [SI/AT]; Peter-Jordanstrasse 163/2/1, A-1190 Wien (AT). **MUDDE, Geert** [NL/AT]; Hochmayrstrasse 31, A-2384 Breitenfurt (AT). **REDL, Gerda** [AT/AT]; Donau-Oder-Kanal IV, SW80, A-2301 Gross-Enzersdorf (AT).

Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
— with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: DISPLAY OF BINDING AGENTS

Figure 1.



(57) Abstract: The present invention relates to a method of preparing a genetic package displaying oligomers of modular antibody domains binding to a target and to a scaffold ligand as well as to vectors and libraries of genetic packages produced thereby. The invention further relates to methods of selecting suitable linker sequences for use in such oligomer display.

WO 2009/000006 A1

Display of binding agents

The invention relates to a method of preparing a genetic package displaying oligomers of modular antibody domains binding to a target and to a scaffold ligand and vectors and libraries of bivalent genetic packages produced by these methods. The invention further relates to methods of selecting suitable linker sequences for use in such oligomer display.

Monoclonal antibodies have been widely used as a scaffold for binding agents. The basic antibody structure will be explained here using as example an intact IgG1 immunoglobulin.

Two identical heavy (H) and two identical light (L) chains combine to form the Y-shaped antibody molecule. The heavy chains each have four domains. The amino terminal variable domains (VH) are at the tips of the Y. These are followed by three constant domains: CH1, CH2, and the carboxy-terminal CH3, at the base of the Y's stem. A short stretch, the switch, connects the heavy chain variable and constant regions. The hinge connects CH2 and CH3 (the Fc fragment) to the remainder of the antibody (the Fab fragments). One Fc and two identical Fab fragments can be produced by proteolytic cleavage of the hinge in an intact antibody molecule. The light chains are constructed of two domains, variable (VL) and constant (CL), separated by a switch.

Disulfide bonds in the hinge region connect the two heavy chains. The light chains are coupled to the heavy chains by additional disulfide bonds. Asn-linked carbohydrate moieties are attached at different positions in constant domains depending on the class of immunoglobulin. For IgG1 two disulfide bonds in the hinge region, between Cys235 and Cys238 pairs, unite the two heavy chains. The light chains are coupled to the heavy chains by two additional disulfide bonds,

between Cys229s in the CH1 domains and Cys214s in the CL domains. Carbohydrate moieties are attached to Asn306 of each CH2, generating a pronounced bulge in the stem of the Y.

These features have profound functional consequences. The variable regions of both the heavy and light chains (VH) and (VL) lie at the "tips" of the Y, where they are positioned to react with antigen. This tip of the molecule is the side on which the N-terminus of the amino acid sequence is located. The stem of the Y projects in a way to efficiently mediate effector functions such as the activation of complement and interaction with Fc receptors, or ADCC and ADCP. Its CH2 and CH3 domains bulge to facilitate interaction with effector proteins. The C-terminus of the amino acid sequence is located on the opposite side of the tip, which can be termed "bottom" of the Y.

Two types of light chain, termed lambda (λ) and kappa (κ), are found in antibodies. A given immunoglobulin either has κ chains or λ chains, never one of each. No functional difference has been found between antibodies having λ or κ light chains.

Each domain in an antibody molecule has a similar structure of two beta sheets packed tightly against each other in a compressed antiparallel beta barrel. This conserved structure is termed the immunoglobulin fold. The immunoglobulin fold of constant domains contains a 3-stranded sheet packed against a 4-stranded sheet. The fold is stabilized by hydrogen bonding between the beta strands of each sheet, by hydrophobic bonding between residues of opposite sheets in the interior, and by a disulfide bond between the sheets. The 3-stranded sheet comprises strands C, F, and G, and the 4-stranded sheet has strands A, B, E, and D. The letters A through G denote the

sequential positions of the beta strands along the amino acid sequence of the immunoglobulin fold.

The fold of variable domains has 9 beta strands arranged in two sheets of 4 and 5 strands. The 5-stranded sheet is structurally homologous to the 3-stranded sheet of constant domains, but contains the extra strands C' and C''. The remainder of the strands (A, B, C, D, E, F, G) have the same topology and similar structure as their counterparts in constant domain immunoglobulin folds. A disulfide bond links strands B and F in opposite sheets, as in constant domains.

The variable domains of both light and heavy immunoglobulin chains contain three hypervariable loops, or complementarity-determining regions (CDRs). The three CDRs of a V domain (CDR1, CDR2, CDR3) cluster at one end of the beta barrel. The CDRs are loops that connect beta strands B-C, C'-C'', and F-G of the immunoglobulin fold. The residues in the CDRs vary from one immunoglobulin molecule to the next, imparting antigen specificity to each antibody.

The VL and VH domains at the tips of antibody molecules are closely packed such that the 6 CDRs (3 on each domain) cooperate in constructing a surface (or cavity) for antigen-specific binding. The natural antigen binding site of an antibody thus is composed of the loops which connect strands B-C, C'-C'', and F-G of the light chain variable domain and strands B-C, C'-C'', and F-G of the heavy chain variable domain.

The loops which are not CDR-loops in a native immunoglobulin, or not part of the antigen-binding pocket as determined by the CDR loops and optionally adjacent loops within the CDR loop region, do not have antigen binding or epitope binding specificity, but contribute to the correct folding of the

entire immunoglobulin molecule and/or its effector or other functions and are therefore called structural loops for the purpose of this invention.

Prior art documents show that the immunoglobulin-like scaffold has been employed so far for the purpose of manipulating the existing antigen binding site, thereby introducing novel binding properties. In most cases the CDR regions have been engineered for antigen binding, in other words, in the case of the immunoglobulin fold, only the natural antigen binding site has been modified in order to change its binding affinity or specificity. A vast body of literature exists which describes different formats of such manipulated immunoglobulins, frequently expressed in the form of single-chain Fv fragments (scFv) or Fab fragments, either displayed on the surface of phage particles or solubly expressed in various prokaryotic or eukaryotic expression systems.

WO06072620A1 describes a method of engineering an immunoglobulin which comprises a modification in a structural loop region to obtain new antigen binding sites. This method is broadly applicable to immunoglobulins and may be used to produce a series of immunoglobulins targeting a variety of antigens. A CH3 library has been shown to be useful for selecting specific binders to an antigen.

Although multivalent display of proteins on genetic packages has been described (such as direct phage cloning and display, bacterial display, yeast display), prior art refers to monomeric monovalent display of binding domains, in general. WO9209690 describes phagemid particles displaying a single copy of a fusion protein on the surface of the particle. Thereby it was described to obtain high affinity binders from a library of phagemid particles, also called bacteriophages. Replicable expression vectors comprising genes encoding a

binding polypeptide and a phage coat protein are provided so to form a gene fusion encoding a fusion protein, which is a chimeric protein of a phagemid particle, the phage coat protein and the binding polypeptide.

US5223409 generally describes the method of fusing a gene encoding a protein of interest to the N-terminal domain of the gene III coat protein of the filamentous phage M13. The gene fusion is mutated to form a library of structurally related fusion proteins that are expressed in low quantity on the surface of a phagemid particle. Biological selection and screening is employed to identify novel ligands useful as drug candidates.

However, there are some limitations in using such "fusion phage" or monovalent phage display and respective single fusion proteins. Many biologicals naturally occur in oligomeric form. For the purpose of the present invention oligomeric means dimeric, trimeric or even higher polymeric forms, up to 24 monomers.

The fusion phages according to the prior art are described to display monomeric fusion proteins, mainly because it was believed that binders of highest affinity could only be selected from a library if single fusion proteins are displayed by the phagemid particles. Native proteins are however often assembled as a dimer or even at a higher degree of oligomerization. To obtain dimeric display with a single fusion protein, some techniques have been developed that involve conditional stop codons located between the coat protein and the binding polypeptide (Dall'Acqua et al *The Journal of Immunology*, 2002, 169: 5171-5180). Thereby soluble monomers of the polypeptides in addition to those fused to the phage are expressed, thus enabling the formation of a dimer. However, such stop codons requires propagation in specific

suppressor host cells that may translate a stop codon in an amino acid, to provide an appropriate amount of fusion proteins in addition to the soluble binding polypeptides. WO 03/029456 describes the use of multi-chain eukaryotic display vectors for the selection of immunoglobulin Fab fragments on the surface of yeast cells.

Prior art fusion proteins involve in some cases linker sequences to display larger binding polypeptides. Linker sequences of up to 24 amino acids are usually employed for standard purposes of displaying variable domains of an antibody. See for example, the display vector pCOMB3x (Hybrid. Hybridomics. 2003 Apr;22(2):97-108. Development of functional human monoclonal single-chain variable fragment antibody against HIV-1 from human cervical B cells. Berry JD, Rutherford J, Silverman GJ, Kaul R, Elia M, Gobuty S, Fuller R, Plummer FA, Barbas CF.)

It is an object of this invention to provide an effective method for the preparation of oligomers of modular antibody domains and to prepare such oligomers displayed on the surface of a replicable genetic package.

Brief Description of the Invention

The objects are solved by the subject matter of the present invention.

According to the invention a method of preparing a genetic package displaying oligomers of modular antibody domains binding to a target and to a scaffold ligand comprising

- providing a genetic package, and
- displaying at least two of the antibody modular domains by fusing to the outer surface of the package

is covered.

The genetic package can be displayed in a mobilized or cellular system, wherein according to the invention the mobilized system can be selected from viruses, phages, phagemids, in-vitro display systems, mRNA systems and ribosomal display systems. Alternatively, a cellular system can be selected using yeast, mammalian cells, bacterial cells, bacterial spores or insect cells.

Oligomers are possibly formed by oligomerization motifs associated with the structure of said agents, such as leucine zipper, disulfide bonds, electrostatic or hydrophobic motifs.

According to one embodiment of the invention the oligomers can be dimers, trimers or tetramers, involving the same type of monomers (homomers) or different types (heteromers). The preferred method according to the invention is specifically useful for providing homomers, in particular homodimers of oligonucleotides on a genetic package, such as a phagemid particle or a phage or yeast.

The method according to the invention can be applied to oligomers which are polypeptides with a target binding site that directs towards the surface of the genetic package and is close to said surface that are biologicals, such as polypeptides.

Alternatively, the appropriate design of such a polypeptide is employed with a binding site that is closer to the surface of the genetic package than to the surrounding environment of the genetic package. This may be advantageous for a binding polypeptide with a potential binding site that is closer to the C-terminus than to the N-terminus of the polypeptide, in particular when the binding site is engineered in a C-terminal loop position. When the potential binding site is engineered

at a position that is adjacent to the site where the genetic package particle is bound, for instance to a surface structure of a cell or a virus, it is advantageous to choose a stable construct with defined accessibility of the binding partner of said binding agent. C-terminal loop positions are, for instance, less accessible than N-terminal loop positions, when they are fused to the N-terminus of the protein 3 of a filamentous phage because they are exposed and in sterical proximity to the genetic package.

According to a preferred embodiment of the invention, however, a defined structure, such as a true oligomeric or dimeric fusion protein is provided to enable the efficient engineering of the potential binding site at an N-terminal loop position. One embodiment of the invention refers to polypeptides with at least two target binding sites, possibly engineered at the monomeric or the oligomeric target binding agent. In some cases the interaction between the monomeric structures enables additional variations of structures and thus additional potential binding sites.

According to a preferred embodiment of the invention the modular antibody is an antibody, Fc fragment, an antibody fragment with a CDR region and combinations thereof, possibly also comprising a fragment with a binding site at a structural loop position.

It can also be an antibody fragment with a CDR region like for example Fab, dAb, scFv, diabody, unibody, SMIPs, TANDABS, Fc fusion proteins and combinations thereof.

In case the genetic package is a filamentous bacteriophage, the preferred fusion structure employed with a bacteriophage involves at least part of an outer surface protein, such as

p3, p6, or p8, however, p9 or p10 may also be used for the purpose of the invention.

In case the genetic package is yeast it is preferred that the oligomer is a fusion protein comprising one of the proteins of yeast cell surface receptors selected from the group consisting of alpha-agglutinin, a-agglutinin, Aga1p, Aga2p or FLO1.

The appropriate genetic package is preferably provided in a particular form, and containing a vector encoding at least one of said fusion proteins. According to the invention there is, for example, provided a cassette vector, containing sequences encoding one or more than one fusion protein operatively linked to the genetic package. Thus, at least two of the chimeric fusion proteins are bound at the surface of the vector particle. The vector can be, for example, a phagemid.

An expression system for expressing oligomers of modular antibody domains bound at the surface of a genetic package produced according to the inventive method is also covered wherein the oligomers are encoded by a single gene and the fusion protein is displayed with at least two copies on the surface of the genetic package.

By using the method and means according to the invention it is possible to display oligomers of modular antibodies without the need of controlled expression of a soluble form of one of the oligomerization partners. The technique can be utilized for molecules such as antibody fragments, even those containing more than two immunoglobulin domains, e.g. at least four immunoglobulin domains. Thus difficult constructs involving stop codons, such as amber stop codons, can be avoided. There will be no need to get a mixture of fusion proteins and soluble monomers for the dimer display. Thus, a

preferred technique of managing mixtures of a variety of binding agents can easily be employed, while the risk of insufficient matching of the monomers is reduced. It is also possible to avoid the suppressor strains as a host cell. Conventional host cells, with non-suppressor function, can serve in a standard way to propagate the oligomeric fusion proteins.

In another embodiment according to the invention, the fusion between the binding partner and the surface protein of the genetic package is such that no conditional stop codon (i.e. no amber, ochre, opal or other similar conditional stop codon) is present in between. In such a situation, upon infection with helper phage, the binding partner is present only as a fusion protein and not in soluble form. In order for the dimer (trimer or higher) to form on the surface of the genetic package, the linker which connects the binding partner to the genetic package needs to be of sufficient length and sufficient flexibility. Linkers that fulfil this requirement can be selected using the method described above.

In another embodiment according to the invention, a helper phage can be used that has specific properties that favour the display of more than one copy of the fusion protein on the surface of the genetic package. An example for such a helper phage is the so-called hyperphage (Rondot S, Koch J, Breitling F, Dubel S. A helper phage to improve single-chain antibody presentation in phage display. Nat Biotechnol. 2001 Jan;19(1):75-78), which itself is devoid of p3, and therefore depends totally on the p3 fusion protein provided by the phagmid in order to be infective. Using such helper phage leads preferentially to phage particles that carry more than one copies of the fusion protein on their surface, and thus favors the formation of dimers of the binding protein on the surface of the genetic package.

According to a preferred embodiment of the invention each of the fusion protein monomers are prepared in the context of an oligomer, so that the portion of soluble binding agents is less than 20%, more preferably less than 10%, most preferably less than 1%.

According to the inventive method a genetic package can be prepared preferably displaying at least two fusion proteins. In a specific embodiment each monomer of the inventive oligomer is bound to the outer surface of the genetic package. The invention also provides bivalent phage displaying two fusion proteins from oligomers each containing an oligomer that dimerizes upon expression.

A library of genetic packages is also claimed, which can exemplarily comprise at least 10 variant genetic packages, wherein said variant genetic packages can display heteromers of modular antibody domains. The heteromer can be based on a scaffold comprising a target binding site. According to a specific embodiment of the invention the target binding site and the scaffold binding site can be similar or identical. A scaffold ligand according to the invention can also be a CDR target. For example, the scaffold can be a parent Fab and at least 20%, preferably at least 30%, more preferred at least 40% of the parent Fab variants are binding to the CDR-target of said parent Fab.

The inventive library can also contain variants of the oligomer produced according to the invention having differences in the amino acid sequence.

This can be provided by modifying the amino acid sequences by at least one insertion or substitution to introduce at least one foreign amino acid, and by a deletion. Foreign amino acids can be introduced by a randomization technique. The foreign

amino acids can also be selected from a specific group of amino acids to obtain a library enriched with specific amino acids at the randomized positions. When the foreign amino acid is selected from a specific group of amino acids, such as amino acids with specific polarity, or hydrophobicity, a library enriched in the specific group of amino acids at the randomized positions can be obtained according to the invention. Such libraries are also called "focused" libraries.

According to the invention there is further provided a method of selecting a linker for binding a polypeptide to the outer surface of a genetic package, comprising

- a. providing a library of genetic packages containing a variety of linkers to connect a first polypeptide to the genetic package,
- b. determining the member of the library containing a linker that does not significantly interfere with the function of said first polypeptide, and
- c. selecting said linker for connecting a second polypeptide to said genetic package.

Thereby suitable linkers to be used for fusion proteins of the same type can be obtained. By the same type of fusion proteins those are meant that link the same formats of genetic packages and binding agent to each other.

Such a method according to the invention can further be used to prepare enriched libraries containing a preselected group of linker variants that fulfil at least one selection criterion, such as flexibility and sterical accessibility of the binding site to the binding partner. This can be determined by measuring the binding properties of a well-known binding agent in the presence of a variety of linker sequences. The enriched library may thus be further selected for another criterion, such as protease resistance, which is

e.g. important for the stability in a medium containing bacterial proteases.

The selected specific linker sequence can then be used to prepare libraries of fusion proteins of the same format, however, with variants of the binding agents, thus enabling the identification, selection and preparation of those agents with the best binding properties in predetermined test systems.

In a preferred embodiment according to the invention such a linker is at least 20 amino acids long, preferably at least 25 amino acid residues, more preferably at least 30 amino acid residues, up to 50 amino acid residues. Especially when amino acids such as Gly, Ser or Ala are involved, which are responsible for the flexibility of a linker, such a linker is advantageously used for the display of a potential binding site, which is close to the surface of the genetic package and which potential binding site may not be able to bind to its partner for sterical reasons.

The linker between the protein to be displayed and the anchor protein of the genetic package (in case of filamentous phage e.g. p3, p8, pX, pIX, pVII) is especially important if the potential binding site of the displayed molecule is in spatial vicinity of the phage particle. In antibody libraries utilizing variable domains and antigen binding sites formed by CDR-loops and display of the library members as amino-terminal fusion to p3 the potential antigen binding site is directed away from the phage particle. Therefore, the linker structure between library members and the phage coat protein is not important. Engineering the bottom loops of immunoglobulin domains and performing phage display may however be an inefficient process and decrease yields of antigen binding clones or even preclude it. Varying the linker between a

library member protein and its fusion partner on the surface can solve or may at least reduce this problem.

In order to select for optimal linker sequences (in terms of length and flexibility as well as stability) a library of linkers can be prepared in which the anchor protein at the surface of the genetic replicable package is fused to a known binding protein which is for sterical reasons notoriously difficult to select for.

This library of sequences can be varied in length and amino acid content.

Selection methods of the linker library for optimal linkers depend on the application but basically it should be for selecting all properties one wishes to have in a certain methodology. Enrichment against an antigen that is difficult to select may yield linker sequences which allow library members a good access to the antigen. Incubation in protease solutions or under other harsh conditions or frequent passaging through host cells under proteolytic conditions (e.g. old microbial cultures) may be an appropriate selection for stable display linkers.

A library of linkers may be produced by any well known library technology. Synthetic linker sequence lengths may vary between 10-500 amino acids. Alternatively, linker can be complete proteins known to be of flexible nature.

The invention also provides a method of producing an oligomer of modular antibody domains binding to a target comprising the steps of:

- providing a library of oligomers of modular antibody domains produced according to the inventive method as described
- contacting said library with said target in the presence of a scaffold ligand,

- selecting a library member binding to said target in the presence of a scaffold ligand, and
- manufacturing a preparation of the functional oligomer.

The scaffold ligand can be selected from the group consisting of an effector molecule, FcRn, serum albumin, Protein A, Protein G, Protein L or a CDR target. As an example, the effector molecule can be selected from the group consisting of CD64, CD16, CD32, Fc receptors.

The oligomers can be dimers selected from the group of VH/VL, CH1/CL, CH2/CH2, CH3/CH3, Fc and Fab, or single chains thereof.

The method according to the invention can provide a library containing at least 10^2 independent clones expressing functional oligomers of modular antibody domains or variants thereof. The library member can then be selected according to the requested binding affinity, preferably it has a target binding affinity of $K_d < 10^{-8} M$. According to the invention it is also provided a pool of preselected independent clones, which is e.g. affinity matured, which pool comprises preferably at least 10, more preferably at least 100, more preferably at least 1000, more preferably at least 10000, even more than 100000 independent clones. Those libraries, which contain the preselected pools, are preferred sources to select the high affinity modular antibodies according to the invention.

Preferably the library is a yeast library and the yeast host cell exhibits at the surface of the cell the oligomers with the biological activity. The yeast host cell is preferably selected from the genera *Saccharomyces*, *Pichia*, *Hansenula*, *Schizisaccharomyces*, *Kluyveromyces*, *Yarrowia* and *Candida*. Most preferred, the host cell is *Saccharomyces cerevisiae*.

According to a specific embodiment of the invention, the target is a receptor of the erbB class. In this case, by the method of the invention an immunoglobulin can be obtained that binds to a receptor of the erbB class.

The invention further provides a high quality library containing at least 10^6 independent clones of functional dimers of modular antibody domains or variants thereof, or the pools of optimized or preselected clones, e.g. the affinity matured clones, which pools are containing at least 10 independent clones that are binding to a target and to a scaffold ligand. The target can be a ligand binding to a parent molecule subject to amino acid variation. The parent molecule can be a functional Fc or a functional Fab, or part thereof.

According to a specific embodiment of the invention the parent molecules can be varied by random or site specific mutagenesis.

The library can contain functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand, and at least 20%, preferably at least 30%, more preferred at least 40% of the functional dimers are binding to CD64. This is particularly preferred with a modular antibody that contains CH2 domains, such as an Fc scaffold.

Alternatively, the library can contain functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand, and at least 20%, preferably at least 30%, more preferred at least 40% of the functional dimers are binding to protein A. This is particularly preferred with a modular antibody that contains CH2 and CH3 domains, such as an Fc scaffold.

Alternatively, the library can contain functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand, and at least 20%, preferably at least 30%, more preferred at least 40% of the functional dimers are binding to the same CDR target. This is particularly preferred with modular antibodies containing a variable region, such as an Fab scaffold with specificity to a single CDR target.

Figures:

Figure 1:

Schematic presentation of the PCRs used for production of the fragments used for assembly of the library Fcab01. PCR primers are indicated by arrows with their respective 5'-3' orientation, and vertical lines indicate the approximate positions of the introduced restriction sites which were used for assembly of the mutated gene. The restriction sites are contained on the primers for ligations of the PCR fragments.

Figure 2:

Amino acid sequence and secondary structure of a CH3 domain (IMGT numbering). The randomization scheme is provided for the libraries Fcab01 to Fcab06.

Randomized positions in the AB and EF loop are marked with a circle. X stands for all 20 amino acids (encoded by NNB), z only for Ala, Asp, Ser, Tyr (encoded by KMT; focused library).

Detailed Description of the Invention

The oligomers of the modular antibody domains according to the present invention will be useful as stand-alone molecules, as well as fusion proteins or derivatives, most typically fused before or after modification in such a way as to be part of larger structures, e.g. of complete antibody molecules, or parts thereof. Immunoglobulins or fusion proteins as produced according to the invention thus also comprise Fc fragments,

Fab fragments, Fv fragments, single domain antibodies, single chain antibodies, in particular single-chain Fv fragments, bi- or multispecific scFv, diabodies, unibodies, multibodies, multivalent or multimers of immunoglobulin domains and others. It will be possible to use the engineered proteins to produce molecules which are monospecific, bispecific, trispecific, and may even carry more specificities. By the invention it is possible to control and preselect the valency of binding at the same time according to the requirements of the planned use of such molecules.

Specific terms as used throughout the specification have the following meaning.

The term "immunoglobulin" as used according to the present invention is defined as polypeptides or proteins that may exhibit mono- or bi- or multi-specific, or mono-, bi- or multivalent binding properties, preferably at least two, more preferred at least three specific binding sites for epitopes of e.g. antigens, effector molecules or proteins either of pathogen origin or of human structure, like self-antigens including cell-associated or serum proteins. The term immunoglobulin as used according to the invention also includes functional fragments of an antibody, such as Fc, Fab, scFv, single chain dimers of CH1/CL domains, Fv, dimers like VH/VL, CH1/CL, CH2/CH2, CH3/CH3, or other derivatives or combinations of the immunoglobulins, like single chains of pairs of immunoglobulin domains. The definition further includes domains of the heavy and light chains of the variable region (such as dAb, Fd, Vl, Vk, Vh, VHH) and the constant region or individual domains of an intact antibody such as CH1, CH2, CH3, CH4, C1 and Ck, as well as mini-domains consisting of at least two beta-strands of an immunoglobulin domain connected by a structural loop.

"Modular antibodies" as used according to the invention are defined as antigen-binding molecules, like human antibodies, composed of at least one polypeptide module or protein domain, preferably in the natural form. The term "modular antibodies" includes antigen-binding molecules that are either immunoglobulins, immunoglobulin-like proteins, or other proteins exhibiting modular formats and antigen-binding properties similar to immunoglobulins or antibodies, which can be used as antigen-binding scaffolds, preferably based on human proteins.

The term "immunoglobulin-like molecule" as used according to the invention refers to any antigen-binding protein, in particular to a human protein, which has a domain structure that can be built in a modular way. Immunoglobulin-like molecules as preferably used for the present invention are T-cell receptors (TCR), fibronectin, transferrin, CTLA-4, single-chain antigen receptors, e.g. those related to T-cell receptors and antibodies, antibody mimetics, adnectins, anticalins, phylomers, repeat proteins such as ankyrin repeats, avimers, VersabodiesTM, scorpio toxin based molecules, and other non-antibody protein scaffolds with antigen binding properties.

Ankyrin repeat (AR), armadillo repeat (ARM), leucine-rich repeat (LRR) and tetratricopeptide repeat (TPR) proteins are the most prominent members of the protein class of repeat proteins. Repeat proteins are composed of homologous structural units (repeats) that stack to form elongated domains. The binding interaction is usually mediated by several adjacent repeats, leading to large target interaction surfaces.

Avimers contain A-domains as strings of multiple domains in several cell-surface receptors. Domains of this family bind

naturally over 100 different known targets, including small molecules, proteins and viruses. Truncation analysis has shown that a target is typically contacted by multiple A-domains with each domain binding independently to a unique epitope. The avidity generated by combining multiple binding domains is a powerful approach to increase affinity and specificity, which these receptors have exploited during evolution.

Anticalins are engineered human proteins derived from the lipocalin scaffold with defined binding properties typical for humanized antibodies. Lipocalins comprise 160-180 amino acids and form conical beta-barrel proteins with a ligand-binding pocket surrounded by four loops. Small hydrophobic compounds are the natural ligands of lipocalins, and different lipocalin variants with new compound specificities (also termed 'anticalins') could be isolated after randomizing residues in this binding pocket.

Single chain or single domain antigen receptors contain a single variable domain and are 20% smaller than camelid single domain antibodies.

Phylomers are peptides derived from biodiverse natural protein fragments.

It is understood that the term "modular antibody", "immunoglobulin", "immunoglobulin-like proteins" includes a derivative thereof as well. A derivative is any combination of one or more modular antibodies of the invention and or a fusion protein in which any domain or minidomain of the modular antibody of the invention may be fused at any position of one or more other proteins (such as other modular antibodies, immunoglobulins, ligands, scaffold proteins, enzymes, toxins and the like). A derivative of the modular antibody of the invention may also be obtained by association

or binding to other substances by various chemical techniques such as covalent coupling, electrostatic interaction, disulphide bonding etc. The other substances bound to the immunoglobulins may be lipids, carbohydrates, nucleic acids, organic and inorganic molecules or any combination thereof (e.g. PEG, prodrugs or drugs). A derivative would also comprise an antibody with the same amino acid sequence but made completely or partly from non-natural or chemically modified amino acids.

A "structural loop" or "non-CDR-loop" according to the present invention is to be understood in the following manner: modular antibodies, immunoglobulins or immunoglobulin-like substances are made of domains with a so called immunoglobulin fold. In essence, strands of antiparallel beta sheets are connected by loops to form a compressed antiparallel beta barrel. In the variable region, some of the loops of the domains contribute essentially to the specificity of the antibody, i.e. the binding to an antigen by the natural binding site of an antibody. These loops are called CDR-loops. The CDR loops are located within the CDR loop region, which may in some cases include also the variable framework region (called "VFR") that is adjacent to the CDR loops. It is known that VFRs may contribute to the antigen binding pocket of an antibody, which generally is mainly determined by the CDR loops. Thus, those VFRs are considered as part of the CDR loop region, and would not be appropriately used for engineering new antigen binding sites. Contrary to those VFRs within the CDR loop region or located proximal to the CDR loops, other VFRs of variable domains would be particularly suitable for use according to the invention. Those are the structural loops of the VFRs located opposite to the CDR loop region, or at the C-terminal side of a variable immunoglobulin domain.

The term "antigen" or "target" as used according to the present invention shall in particular include all antigens and target molecules capable of being recognised by a binding site of a modular antibody. Specifically preferred antigens as targeted by the receptor molecule according to the invention are those antigens or molecules, which have already been proven to be or are capable of being immunologically or therapeutically relevant, especially those, for which a clinical efficacy has been tested.

The term "target" or "antigen" as used herein shall comprise molecules selected from the group consisting of allergens, tumor associated antigens, self antigens including cell surface receptors, enzymes, Fc-receptors, FcRn, HSA, IgG, interleukins or cytokines, proteins of the complement system, transport proteins, serum molecules, bacterial antigens, fungal antigens, protozoan antigens and viral antigens, also molecules responsible for transmissible spongiform encephalitis (TSE), such as prions, infective or not, and markers or molecules that relate to inflammatory conditions, such as pro-inflammatory factors, multiple sclerosis or alzheimer disease, or else haptens.

The term "cell surface antigens" shall include all antigens capable of being recognised by an antibody structure on the surface of a cell, and fragments of such molecules. Preferred cell surface antigens are those antigens, which have already been proven to be or which are capable of being immunologically or therapeutically relevant, especially those, for which a preclinical or clinical efficacy has been tested. Those cell surface molecules are specifically relevant for the purpose of the present invention, which mediate cell killing activity. Upon binding of the immunoglobulin according to the invention to preferably at least two of those cell surface

molecules the immune system provides for cytolysis or cell death, thus a potent means for attacking human cells may be provided.

The antigen is either recognized as a whole target molecule or as a fragment of such molecule, especially substructures of targets, generally referred to as epitopes.

Substructures of antigens are generally referred to as "epitopes" (e.g. B-cell epitopes, T-cell epitopes), as long as they are immunologically relevant, i.e. are also recognisable by natural or monoclonal antibodies. The term "epitope" as used herein according to the present invention shall mean a molecular structure which may completely make up a specific binding partner or be part of a specific binding partner to a binding site of a modular antibody or an immunoglobulin of the present invention. The term epitope may also refer to haptens. Chemically, an epitope may either be composed of a carbohydrate, a peptide, a fatty acid, an organic, biochemical or inorganic substance or derivatives thereof and any combinations thereof. If an epitope is a polypeptide, it will usually include at least 3 amino acids, preferably 8 to 50 amino acids, and more preferably between about 10-20 amino acids in the peptide. There is no critical upper limit to the length of the peptide, which could comprise nearly the full length of a polypeptide sequence of a protein. Epitopes can be either linear or conformational epitopes. A linear epitope is comprised of a single segment of a primary sequence of a polypeptide chain. Linear epitopes can be contiguous or overlapping. Conformational epitopes are comprised of amino acids brought together by folding of the polypeptide to form a tertiary structure and the amino acids are not necessarily adjacent to one another in the linear sequence. Specifically, epitopes are at least part of diagnostically relevant

molecules, i.e. the absence or presence of an epitope in a sample is qualitatively or quantitatively correlated to either a disease or to the health status of a patient or to a process status in manufacturing or to environmental and food status. Epitopes may also be at least part of therapeutically relevant molecules, i.e. molecules which can be targeted by the specific binding domain which changes the course of the disease.

As used herein, the term "specifically binds" or "specific binding" refers to a binding reaction which is determinative of the cognate ligand of interest in a heterogeneous population of molecules. Thus, under designated conditions (e.g. immunoassay conditions), the modular antibody binds to its particular target and does not bind in a significant amount to other molecules present in a sample. The specific binding means that binding is selective in terms of target identity, high, medium or low binding affinity or avidity, as selected. Selective binding is usually achieved if the binding constant or binding dynamics is at least 10 fold different, preferably the difference is at least 100 fold, and more preferred a least 1000 fold.

The term "expression system" refers to nucleic acid molecules containing a desired coding sequence and control sequences in operable linkage, so that hosts transformed or transfected with these sequences are capable of producing the encoded proteins. In order to effect transformation, the expression system may be included on a vector; however, the relevant DNA may also be integrated into the host chromosome. Alternatively, an expression system can be used for in vitro transcription/translation. The expression system preferably is employing a host cell that is either a eukaryotic or prokaryotic host cell, preferably a mammalian or yeast host cell, as well as a bacterial host cell.

The preferred expression system for the fusion proteins is a non-suppressor host cell, which would be sensitive to a stop codon, such as an amber stop codon, and would thus stop translation thereafter. In the absence of such a stop codon such non-suppressor host cells, preferably E.coli, are preferably used. In the presence of such a stop codon supressor host cells would be used.

All numbering of the amino acid sequences of the immunoglobulins is according to the IMGT numbering scheme (IMGT, the international ImMunoGeneTics, Lefranc et al., 1999, Nucleic Acids Res. 27: 209-212).

For the purposes of this invention, the term "binding agent" or "ligand" refers to a member of a binding pair, in particular binding polypeptides having the potential of serving as a binding domain for a binding partner. Examples of binding partners include pairs of binding agents with functional interactions, such as receptor binding to ligands, antibody binding to antigen or receptors, a drug binding to a target, and enzyme binding to a substrate

The term "fusion protein" or "chimeric fusion protein" as used for the purpose of the invention shall mean the molecule composed of a genetic package, at least part of an outer surface structure, such as a coat protein or part thereof, optionally a linker sequence, and a binding agent. The fusion protein is encoded by a vector with the gene of the binding agent and information to display a copy of the binding agent at the surface of the genetic package.

The term "cytotoxic activity" as used for the purpose of the invention shall mean the activity on effector cells resulting in activation of cytotoxic T-cells or on cells, which mediate

antibody-dependent cell cytotoxicity (ADCC), complement dependent cytotoxicity (CDC) and/or antibody-dependent cellular phagocytosis (ADCP). Modular antibodies according to the invention thus kill antibody-coated target cells, which they optionally bind with their Fc receptors.

"Scaffold" shall mean a temporary framework either natural or artificial used to support the molecular structure of a polypeptide in the construction of variants or a repertoire of the polypeptide. It is usually a modular system of polypeptide domains that maintains the tertiary structure or the function of the parent molecule. Exemplary scaffolds are modular antibodies, which may be mutagenized to produce variants within said scaffold, to obtain a library.

The term "scaffold ligand" as used for the purpose of the invention shall mean a ligand that binds to a scaffold or the backbone of modular antibodies, thus determining the molecular structure or primary function and specificity of said modular antibody. In preferred cases the scaffold ligand is a functional ligand, mediating a biological function upon binding, like an effector ligand. In an alternative embodiment the scaffold ligand is a functional ligand, which is a specific target bound by the CDR region, non-structural loop region or structural loop region. The same scaffold ligand can bind many variants of a modular antibody regardless of their target specificities. In general, the presence of scaffold ligand binding site indicates that the variant is expressed and folded correctly. Thus, binding of the scaffold ligand to its binding site provides a method for preselecting functional polypeptides from a repertoire of polypeptides. Designing variants of modular antibodies that keep the binding property to a scaffold ligand avoids the preparation of variants that are non-functional, for example as a result of the introduction of mutations, folding mutants or expression

mutants which would be or are incapable of binding to substantially any target or effector ligand. Such non-functional mutants sometimes are generated by the normal randomisation and variation procedures employed in the construction of polypeptide repertoires. Providing functional mutants that bind to a scaffold ligand permits the person skilled in the art to prepare a library of modular antibodies which is enriched in functional, well folded and highly expressed library members.

The term "effector ligand" as used for the purpose of the invention shall mean a ligand mediating effector functions, like an effector molecule. Exemplary effector ligands are Fc receptors or Fc receptor-like molecules interfering with immunoglobulins. An Fc receptor is a protein found on the surface of certain cells - including natural killer cells, macrophages, neutrophils, and mast cells - that contribute to the protective functions of the immune system. Its name is derived from its binding specificity for a part of an antibody known as the Fc (Fragment-crystallizable) region. Fc receptors bind to antibodies that are attached to infected cells or invading pathogens. Their activity stimulates phagocytic or cytotoxic cells to destroy microbes, or infected cells by antibody-mediated cellular phagocytosis (ADCP) or antibody-dependent cell-mediated cytotoxicity (ADCC). There are several different types of Fc receptors, which are classified based on the type of antibody that they recognize; those that bind the most common class of antibody, IgG, are called Fc-gamma receptors (Fc γ R), those that bind IgA are called Fc-alpha receptors (Fc α R) and those that bind IgE are called Fc-epsilon receptors (Fc ϵ R). Equivalent to an effector ligand and thus incorporated into the definition is any surrogate ligand that recognizes the same or similar binding site within the modular antibody, such as Protein A.

All FcγRs belong to the immunoglobulin superfamily and are the most important Fc receptors for inducing phagocytosis of opsonized (coated) microbes. This family includes several members that differ in their antibody affinities due to their different molecular structure: FcγRI (CD64), FcγRIIA (CD32a), FcγRIIB (CD32b), FcγRIIIA (CD16a), FcγRIIIB (CD16b. For instance, FcγRI binds to IgG more strongly than FcγRII and FcγRIII, and has an extracellular portion composed of three immunoglobulin (Ig)-like domains, one more domain than FcγRII and FcγRIII. These properties allow activation of FcγRI by a sole IgG molecule (or monomer), while the latter two Fcγ receptors must bind multiple IgG molecules within an immune complex to be activated.

Another FcR is expressed on multiple cell types and is similar in structure to MHC class I. This receptor also binds IgG and is involved in preservation of this antibody in order to increase its biological half-life in vivo. However, since this Fc receptor is also involved in transferring IgG from a mother either via the placenta to her fetus or in milk to her suckling infant, it is called the neonatal Fc receptor (FcRn). Recently this receptor has been implicated in being involved in homeostasis of IgG serum levels.

Antibody-Dependent Cell-Mediated Cytotoxicity (ADCC) is a mechanism of cell-mediated immunity whereby an effector cell of the immune system actively lyses a target cell that has been bound by specific antibodies. It is one of the mechanisms through which antibodies, as part of the humoral immune response, can act to limit and contain infection. Classical ADCC is mediated by natural killer (NK) cells; monocytes and eosinophils can also mediate ADCC. For example Eosinophils can kill certain parasitic worms known as helminths through ADCC. ADCC is part of the adaptive immune response due to its dependence on a prior antibody response.

The term "foreign" in the context of amino acids shall mean the newly introduced amino acids being naturally occurring, but foreign to the site of modification, or substitutes of naturally occurring amino acids. "Foreign" with reference to an antigen binding sites means that the antigen binding site is not naturally formed by the specific binding region of the agent, and a foreign binding partner, but not the natural binding partner of the agent, is bound by the newly engineered binding site.

The term "variable binding region" sometimes called "CDR region" as used herein refers to molecules with varying structures capable of binding interactions with antigens. Those molecules can be used as such or integrated within a larger protein, thus forming a specific region of such protein with binding function. The varying structures can be derived from natural repertoires of binding proteins such as immunoglobulins or phylomers or synthetic diversity, including repeat-proteins, avimers and anticalins. The varying structures can as well be produced by randomization techniques, in particular those described herein. These include mutagenized CDR or non-CDR regions, loop regions of immunoglobulin variable domains or constant domains.

Modified binding agents with different modifications at specific sites are referred to as "variants". Variants of a scaffold are preferably grouped to form libraries of binding agents, which can be used for selecting members of the library with predetermined functions. In accordance therewith, a loop region of a binding agent comprising positions within one or more loops potentially contributing to a binding site, is preferably mutated or modified to produce libraries, preferably by random, semi-random or, in particular, by site-directed random mutagenesis methods, in particular to delete,

exchange or introduce randomly generated inserts into loops, preferably into structural loops. Alternatively preferred is the use of combinatorial approaches. Any of the known mutagenesis methods may be employed, among them cassette mutagenesis. These methods may be used to make amino acid modifications at desired positions of the immunoglobulin of the present invention. In some cases positions are chosen randomly, e.g. with either any of the possible amino acids or a selection of preferred amino acids to randomize loop sequences, or amino acid changes are made using simplistic rules. For example all residues may be mutated preferably to specific amino acids, such as alanine, referred to as amino acid or alanine scanning. Such methods may be coupled with more sophisticated engineering approaches that employ selection methods to screen higher levels of sequence diversity.

The preferred cytotoxic modular antibody according to the invention with a molecular weight of less than 60 kD or up to 60 kD has a small size as compared to full length antibodies. The preferred size is up to 55kD. Modular antibody single domains usually have a molecular size of 10-15 kD, thus a molecule based on 4 modular antibody domains would have a molecular size of 40-60 kD, depending on the glycosylation or any additional conjugation of pharmacologically active substances, like toxins or peptides.

The preferred format is an oligomer, composed of modular antibody domains, preferably up to 4 domains, more preferred 3 domains, and even more preferred made up of 2 domains. Formats based on the combination of 5 modular antibody domains or more are commonly thought not to exert the specific advantages of small sized antibody fragments, which are e.g. ease of expression in various expression systems and tissue penetration.

It is feasible to provide the preferred modular antibody of the invention as a single domain antibody. However, antibody domains tend to dimerize upon expression, either as a homodimer, like an Fc, or a heterodimer, like an Fab. The dimeric structure is thus considered as a basis for the preferred stable molecule. The preferred dimers of immunoglobulin domains are selected from the group consisting of single domain dimers, like VH/VL, CH1/CL (kappa or lambda), CH2/CH2 and CH3/CH3. Dimers or oligomers of modular antibody domains can also be provided as single chain or two chain molecules, in particular those linking the C-terminus of one domain to the N-terminus of another.

Binding partners are agents that specifically bind to one another, usually through non-covalent interactions. Examples of binding partners include pairs of binding agents with functional interactions, such as receptor binding to ligands, antibody binding to antigen, a drug binding to a target, and enzyme binding to a substrate. Binding partners have found use in many therapeutic, diagnostic, analytical and industrial applications. Most prominent binding partners, also called binding pairs, are antibodies or immunoglobulins, fragments or derivatives thereof. In most cases the binding of such binding agents is required to mediate a biological effect or a function, a "functional interaction".

According to a specific embodiment of the present invention the binding agent is an immunoglobulin of human or murine origin, and may be employed for various purposes, in particular in pharmaceutical compositions. Of course, the modified immunoglobulin may also be a humanized or chimeric immunoglobulin.

The binding agent which is a human immunoglobulin is preferably selected or derived from the group consisting of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4 and IgM. The murine immunoglobulin binding agent is preferably selected or derived from the group consisting of IgA, IgD, IgE, IgG1, IgG2A, IgG2B, IgG2C, IgG3 and IgM.

Such a binding agent comprises preferably a heavy and/or light chain or a part thereof. A modified immunoglobulin according to the invention may comprise a heavy and/or light chain, at least one variable and/or constant domain, or a part thereof including a minidomain.

A constant domain is an immunoglobulin fold unit of the constant part of an immunoglobulin molecule, also referred to as a domain of the constant region (e.g. CH1, CH2, CH3, CH4, Ck, Cl).

A variable domain is an immunoglobulin fold unit of the variable part of an immunoglobulin, also referred to as a domain of the variable region (e.g. Vh, Vk, Vl, Vd)

An exemplary modular antibody according to the invention consists of a constant domain selected from the group consisting of CH1, CH2, CH3, CH4, Igk-C, Igl-C, combinations, derivatives or a part thereof including a mini-domain, with at least one loop region, and is characterised in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region binds specifically to at least one epitope of an antigen.

Another modular antibody according to the invention can consist of a variable domain of a heavy or light chain, combinations, derivatives or a part thereof including a

minidomain, with at least one loop region, and is characterised in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region binds specifically to at least one epitope of an antigen.

The modular antibody according to the present invention may comprise one or more domains (e.g. at least two, three, four, five, six, ten domains). If more than one domain is present in the modular antibody these domains may be of the same type or of varying types (e.g. CH1-CH1-CH2, CH3-CH3, (CH2)₂-(CH3)₂, with or without the hinge region). Of course also the order of the single domains may be of any kind (e.g. CH1-CH3-CH2, CH4-CH1-CH3-CH2).

The invention preferably refers to parts of antibodies, such as IgG, IgA, IgM, IgD, IgE and the like. The modular antibodies of the invention may also be a functional antibody fragment such as Fab, Fab₂, scFv, Fv, Fc, FcabTM, an antigen-binding Fc, or parts thereof, or other derivatives or combinations of the immunoglobulins such as minibodies, domains of the heavy and light chains of the variable region (such as dAb, Fd (binding site made up of one or more single domains), VL, including Vlambda (Vl) and Vkappa (Vk), VH, VHH) as well as mini-domains consisting of two beta-strands of an immunoglobulin domain connected by at least two structural loops, as isolated domains or in the context of naturally associated molecules. A particular embodiment of the present invention refers to the Fc fragment of an antibody molecule, either as antigen-binding Fc fragment (FcabTM) through modifications of the amino acid sequence or as conjugates or fusions to receptors, peptides or other antigen-binding modules, such as scFv.

The modular antibodies can be used as isolated polypeptides or as combination molecules, e.g. through recombination, fusion or conjugation techniques, with other peptides or polypeptides. The peptides are preferably homologous to immunoglobulin domain sequences, and are preferably at least 5 amino acids long, more preferably at least 10 or even at least 50 or 100 amino acids long, and constitute at least partially the loop region of the immunoglobulin domain. The preferred binding characteristics relate to predefined epitope binding, affinity and avidity.

The modular antibody according to the invention is possibly further combined with one or more modified modular antibodies or with unmodified modular antibodies, or parts thereof, to obtain a combination modular antibody. Combinations are preferably obtained by recombination techniques, but also by binding through adsorption, electrostatic interactions or the like, or else through conjugation or chemical binding with or without a linker. The preferred linker sequence is either a natural linker sequence or a functionally suitable artificial sequence.

In general the modular antibody according to the invention may be used as a building block to molecularly combine other modular antibodies or biologically active substances or molecules. It is preferred to molecularly combine at least one antibody binding to the specific partner via the variable or non-variable sequences, like structural loops, with at least one other binding molecule which can be an antibody, antibody fragment, a soluble receptor, a ligand or another antibody domain, or a binding moiety thereof. Other combinations refer to proteinaceous molecules, nucleic acids, lipids, organic molecules and carbohydrates.

The engineered molecules according to the present invention will be useful as stand-alone proteins as well as fusion proteins or derivatives, most typically fused in such a way as to be part of larger antibody structures or complete antibody molecules, or parts or fragments thereof, such as Fab fragments, Fc fragments, Fv fragments and others. It will be possible to use the engineered proteins to produce molecules which are monospecific, bispecific, trispecific, and maybe even carry more specificities at the same time, and it will be possible at the same time to control and preselect the valency of binding at the same time according to the requirements of the planned use of such molecules.

According to the present invention, the modular antibody optionally exerts one or more binding regions to antigens, including the binding site binding specifically to the cell surface target and the binding sites mediating effector function. Antigen binding sites to one or more antigens may be presented by the CDR-region or any other natural receptor binding structure, or be introduced into a structural loop region of an antibody domain, either of a variable or constant domain structure. The antigens as used for testing the binding properties of the binding sites may be naturally occurring molecules or chemically synthesized molecules or recombinant molecules, either in solution or in suspension, e.g. located on or in particles such as solid phases, on or in cells or on viral surfaces. It is preferred that the binding of an immunoglobulin to an antigen is determined when the antigen is still adhered or bound to molecules and structures in the natural context. Thereby it is possible to identify and obtain those modified immunoglobulins that are best suitable for the purpose of diagnostic or therapeutic use.

Modular antibody or immunoglobulin domains may be modified according to the present invention (as used herein the terms

immunoglobulin and antibody are interchangeable) which modifications are preferably effected in immunoglobulin domains or parts thereof that contain a loop, either a CDR-loop or a non-CDR loop, structural loops being the preferred sites of modifications or mutagenesis. In some cases it is preferable to use a defined modified structural loop or a structural loop region, or parts thereof, as isolated molecules for binding or combination purposes.

It is particularly preferred that the modular antibody according to the invention is binding to said cell surface target through at least part of a structural loop and/or CDR loop.

In an alternate embodiment it is preferred that the modular antibody according to the invention is binding to said effector ligand, or a surrogate ligand for such an effector ligand, like protein A, through at least part of a structural loop and/or CDR loop, thus mediating the effector function.

In a preferred embodiment the binding agent is binding with its native or modified binding structure or newly formed binding site, specifically to at least two such epitopes that are identical or differ from each other, either of the same antigen or of different antigens.

In a preferred domain structure of a binding agent it is preferred to modify at least one loop region resulting in a substitution, deletion and/or insertion of one or more nucleotides or amino acids, preferably a point mutation, or even the exchange of whole loops, more preferred the change of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15, up to 30 amino acids. Thereby the modified sequence comprises amino acids not included in the conserved regions of the loops, the newly introduced amino acids being naturally

occurring, but foreign to the site of modification, or substitutes of naturally occurring amino acids.

However, the maximum number of amino acids inserted into a loop region of a binding agent preferably may not exceed the number of 30, preferably 25, more preferably 20 amino acids at a maximum. The substitution and the insertion of the amino acids occurs preferably randomly or semi-randomly using all possible amino acids or a selection of preferred amino acids for randomization purposes, by methods known in the art and as disclosed in the present patent application.

The site of modification may be at a specific single loop or a loop region, in particular a structural loop or a structural loop region. A loop region usually is composed of at least one, preferably at least two, preferably at least 3 or at least 4 loops that are at the tip or the bottom of a domain, in proximity or adjacent to each other, and which may contribute to the binding of an antigen through forming an antigen binding site or antigen binding pocket. It is preferred that the one or more sites of modification are located within the area of 10 amino acids, more preferably within 20, 30, 40, 50, 60, 70, 80, 90 up to 100 amino acids, in particular within a structural region to form a surface or pocket where the antigen can sterically access the loop regions.

In this regard the preferred modifications are engineered in the loop regions of CH1, CH2, CH3 and CH4, in particular in the range of amino acids 7 to 21, amino acids 25 to 39, amino acids 41 to 81, amino acids 83 to 85, amino acids 89 to 103 and amino acids 106 to 117.

In another preferred embodiment a modification in the structural loop region comprising amino acids 92 to 98 is

combined with a modification in the structural loop region comprising amino acids 8 to 20.

The above identified amino acid regions of the respective immunoglobulins comprise loop regions to be modified. Preferably, a modification in the structural loop region comprising amino acids 92 to 98 is combined with a modification in one or more of the other structural loops.

In a preferred embodiment a modification in the structural loop region comprising amino acids 92 to 98 is combined with a modification in the structural loop region comprising amino acids 41 to 45.2.

Most preferably each of the structural loops comprising amino acids 92 to 98, amino acids 41 to 45.2 and amino acids 8 to 20 contain at least one amino acid modification.

In another preferred embodiment each of the structural loops comprising amino acids 92 to 98, amino acids 41 to 45.2, and amino acids 8 to 20 contain at least one amino acid modification.

According to another preferred embodiment the amino acid residues in the area of positions 15 to 17, 29 to 34, 41 to 45.2, 84 to 85, 92 to 100, and/or 108 to 115 of CH3 are modified.

The preferred modifications of Igk-C and Igl-C of human origin are engineered in the loop regions in the area of amino acids 8 to 20, amino acids 26 to 36, amino acids 41 to 82, amino acids 83 to 88, amino acids 92 to 100, amino acids 107 to 124 and amino acids 123 to 126.

The preferred modifications of loop regions of Igk-C and Igl-C of murine origin are engineered at sites in the area of amino acids 8 to 20, amino acids 26 to 36, amino acids 43 to 79, amino acids 83 to 85, amino acids 90 to 101, amino acids 108 to 116 and amino acids 122 to 126.

Another preferred immunoglobulin preferably used as a therapeutic according to the invention consists of a variable domain of a heavy or light chain, or a part thereof including a minidomain, with at least one loop region, preferably a structural loop region, and is characterised in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region forms a relevant binding site as described above.

According to a specific embodiment the immunoglobulin preferably used according to the invention may contain a modification within the variable domain, which is selected from the group of VH, Vkappa, Vlambda, VHH and combinations thereof. More specifically, they comprise at least one modification within amino acids 7 to 22, amino acids 39 to 55, amino acids 66 to 79, amino acids 77 to 89 or amino acids 89 to 104, where the numbering of the amino acid position of the domains is that of the IMGT.

In a specific embodiment, the immunoglobulin preferably used according to the invention is characterised in that the loop regions of VH or Vkappa or Vlambda of human origin comprise at least one modification within amino acids 7 to 22, amino acids 43 to 51, amino acids 67 to 77, amino acids 77 to 88, and amino acids 89 to 104, most preferably amino acid positions 12 to 17, amino acid positions 45 to 50, amino acid positions 68 to 77, amino acids 79 to 88, and amino acid positions 92 to

99, where the numbering of the amino acid position of the domains is that of the IMGT.

The structural loop regions of the variable domain of the immunoglobulin of human origin, as possible selected for modification purposes are preferably located in the area of amino acids 8 to 20, amino acids 44 to 50, amino acids 67 to 76, amino acids 78 to 87, and amino acids 89 to 101.

According to a preferred embodiment the structural loop regions of the variable domain of the immunoglobulin of murine origin as possible selected for modification purposes are preferably located in the area of amino acids 6 to 20, amino acids 43 to 52, amino acids 67 to 79, amino acids 79 to 87, and amino acids 91 to 100.

The immunoglobulin preferably used as a therapeutic according to the invention may also be of camelid origin. Camel antibodies comprise only one heavy chain and have the same antigen affinity as normal antibodies consisting of light and heavy chains. Consequently camel antibodies are much smaller than, e.g., human antibodies, which allows them to penetrate dense tissues to reach the antigen, where larger proteins cannot. Moreover, the comparative simplicity, high affinity and specificity and the potential to reach and interact with active sites, camel's heavy chain antibodies present advantages over common antibodies in the design, production and application of clinically valuable compounds.

According to another preferred embodiment of the present invention the structural loop regions of a modular antibody or an immunoglobulins of camelid origin are modified, e.g. within a VHH, in the region of amino acids 7 to 19, amino acids 43 to 55, amino acids 68 to 76, amino acids 80 to 87 and amino acids 91 to 101.

The preferred method of producing the modular antibody according to the invention refers to engineering a modular antibody that is binding specifically to at least one first epitope and comprising modifications in each of at least two structural loop regions, and determining the specific binding of said at least two loop regions to at least one second epitope, wherein the unmodified structural loop region (non-CDR region) does not specifically bind to said at least one second epitope. Thus, an antibody or antigen-binding structure specific for a first antigen may be improved by adding another valency or specificity against a second antigen, which specificity may be identical, either targeting different epitopes or the same epitope, to increase valency or to obtain bi-, oligo- or multispecific molecules.

On the other hand it is preferred to make use of those modular antibodies that contain native structures interacting with effector molecules or immune cells. Those native structures either remain unchanged or are modulated for an increased effector function. Binding sites for e.g. Fc receptors are described to be located in a CH2 and/or CH3 domain region, and may be mutagenized by well known techniques.

ADCC, antibody-dependent cell-mediated cytotoxicity is the killing of antibody-coated target cells by cells with Fc receptors that recognize the constant region of the bound antibody. Most ADCC is mediated by NK cells that have the Fc receptor FcγRIII or CD16 on their surface. Typical assays employ target cells, like Ramos cells, incubated with serially diluted antibody prior to the addition of freshly isolated effector cells. The ADCC assay is then further incubated for several hours and % cytotoxicity detected. Usually the Target: Effector ratio is about 1:16, but may be 1:1 up to 1 : 50.

Complement-dependent cytotoxicity (CDC) is a mechanism of killing cells in which antibody bound to the target cell surface fixes complement, which results in assembly of the membrane attack complex that punches holes in the target cell membrane resulting in subsequent cell lysis. The commonly used CDC assay follows the same procedure as for ADCC determination, however, with complement containing serum instead of effector cells.

The cytotoxic activity as determined by either of ADCC and CDC assay is proven for a modular antibody according to the invention, if there is a significant increase in the percentage of cytolysis as compared to a control. The absolute percentage increase preferably is higher than 5%, more preferably higher than 10%, even more preferred higher than 20%.

The antibody-dependent cellular phagocytosis, ADCP sometimes called ADPC, is usually investigated side by side with cytolysis of cultured human cells. Phagocytosis by phagocytes, usually human monocytes or monocyte-derived macrophages, as mediated by an antibody can be determined as follows. Purified monocytes may be cultured with cytokines to enhance expression of FcγRs or to induce differentiation into macrophages. ADCP and ADCC assays are then performed with target cells. Phagocytosis is determined as the percentage of positive cells measured by flow cytometry. The positive ADCP activity is proven with a significant uptake of the antibody-antigen complex by the phagocytes. The absolute percentage preferably is higher than 5%, more preferably higher than 10%, even more preferred higher than 20%.

In a typical assay PBMC or monocytes or monocyte derived macrophages are resuspended in RF2 medium (RPMI 1640

supplemented with 2% FCS) in 96-well plates at a concentration of 1×10^5 viable cells in 100 μ l/well. Appropriate target cells, expressing the target antigen, e.g. Her2/neu antigen and SKBR3 cells, are stained with PKH2 green fluorescence dye. Subsequently 1×10^4 PKH2- labeled target cells and an Her 2 specific (IgG1) antibody (or modular antibody) or mouse IgG1 isotype control (or modular antibody control) are added to the well of PBMC's in different concentrations (e.g. 1-100 μ g/ml) and incubated in a final volume of 200 μ l at 37°C for 24 h. Following the incubation, PBMCs or monocytes or monocyte derived macrophages and target cells are harvested with EDTA-PBS and transferred to 96-well V-bottomed plates. The plates are centrifuged and the supernatant is aspirated. Cells are counterstained with a 100- μ l mixture of RPE-conjugated anti-CD11b, anti-CD14, and human IgG, mixed and incubated for 60 min on ice. The cells are washed and fixed with 2% formaldehyde-PBS. Two-color flow cytometric analysis is performed with e.g. a FACS Calibur under optimal gating. PKH2-labeled target cells (green) are detected in the FL-1 channel (emission wavelength, 530 nm) and RPE-labeled PBMC or monocytes or monocyte derived macrophages (red) are detected in the FL-2 channel (emission wavelength, 575 nm). Residual target cells are defined as cells that are PKH2⁺/RPE⁻ Dual-labeled cells (PKH2⁺/RPE⁻) are considered to represent phagocytosis of targets by PBMC or monocytes or monocyte derived macrophages. Phagocytosis of target cells is calculated with the following equation: percent phagocytosis = $100 \times [(\text{percent dual positive}) / (\text{percent dual positive} + \text{percent residual targets})]$. All tests are usually performed in duplicate or triplicate and the results are expressed as mean \pm SD.

The effector function of the modular antibody according to the invention usually differs from any synthetic cytotoxic activity, e.g. through a toxin that may be conjugated to an

immunoglobulin structure. Toxins usually do not activate effector molecules and the biological defence mechanism. Thus, the preferred cytotoxic activity of the modular antibodies according to the invention is a biological cytotoxic activity, which usually is immunostimulatory, leading to effective cytolysis.

The modular antibody according to the invention may specifically bind to any kind of binding molecules or structures, in particular to antigens, proteinaceous molecules, proteins, peptides, polypeptides, nucleic acids, glycans, carbohydrates, lipids, organic molecules, in particular small organic molecules, anorganic molecules, or combinations or fusions thereof, including PEG, prodrugs or drugs. The preferred modular antibody according to the invention may comprise at least two loops or loop regions whereby each of the loops or loop regions may specifically bind to different molecules or epitopes.

Preferably the target antigen is selected from cell surface antigens, including receptors, in particular from the group consisting of erbB receptor tyrosine kinases (such as EGFR, HER2, HER3 and HER4, in particular those epitopes of the extracellular domains of such receptors, e.g. the 4D5 epitope), molecules of the TNF-receptor superfamily, such as Apo-1 receptor, TNFR1, TNFR2, nerve growth factor receptor NGFR, CD40, T-cell surface molecules, T-cell receptors, T-cell antigen OX40, TACI-receptor, BCMA, Apo-3, DR4, DR5, DR6, decoy receptors, such as DcR1, DcR2, CAR1, HVEM, GITR, ZTNFR-5, NTR-1, TNFL1 but not limited to these molecules, B-cell surface antigens, such as CD10, CD19, CD20, CD21, CD22, antigens or markers of solid tumors or hematologic cancer cells, cells of lymphoma or leukaemia, other blood cells including blood platelets, but not limited to these molecules.

According to a further preferred embodiment the target antigen is selected from those antigens presented by cells, like epithelial cells, cells of solid tumors, infected cells, blood cells, antigen-presenting cells and mononuclear cells. Those target antigens expressed or overexpressed by cells are preferably targeted, which are selected from the group consisting of tumor associated antigens, in particular EpCAM, tumor-associated glycoprotein-72 (TAG-72), tumor-associated antigen CA 125, Prostate specific membrane antigen (PSMA), High molecular weight melanoma-associated antigen (HMW-MAA), tumor-associated antigen expressing Lewis Y related carbohydrate, Carcinoembryonic antigen (CEA), CEACAM5, HMFG PEM, mucin MUC1, MUC18 and cytokeratin tumor-associated antigen, bacterial antigens, viral antigens, allergens, allergy related molecules IgE, cKIT and Fc-epsilon-receptorI, IRp60, IL-5 receptor, CCR3, red blood cell receptor (CR1), human serum albumin, mouse serum albumin, rat serum albumin, Fc receptors, like neonatal Fc-gamma-receptor FcRn, Fc-gamma-receptors Fc-gamma RI, Fc-gamma-RII, Fc-gamma RIII, Fc-alpha-receptors, Fc-epsilon-receptors, fluorescein, lysozyme, toll-like receptor 9, erythropoietin, CD2, CD3, CD3E, CD4, CD11, CD11a, CD14, CD16, CD18, CD19, CD20, CD22, CD23, CD25, CD28, CD29, CD30, CD32, CD33 (p67 protein), CD38, CD40, CD40L, CD52, CD54, CD56, CD64, CD80, CD147, GD3, IL-1, IL-1R, IL-2, IL-2R, IL-4, IL-5, IL-6, IL-6R, IL-8, IL-12, IL-15, IL-17, IL-18, IL-23, LIF, OSM, interferon alpha, interferon beta, interferon gamma; TNF-alpha, TNFbeta2, TNFalpha, TNFalphabeta, TNF-R1, TNF-RII, FasL, CD27L, CD30L, 4-1BBL, TRAIL, RANKL, TWEAK, APRIL, BAFF, LIGHT, VEG1, OX40L, TRAIL Receptor-1, A1 Adenosine Receptor, Lymphotoxin Beta Receptor, TACI, BAFF-R, EPO; LFA-3, ICAM-1, ICAM-3, integrin beta1, integrin beta2, integrin alpha4/beta7, integrin alpha2, integrin alpha3, integrin alpha4, integrin alpha5, integrin alpha6, integrin alphav, alphaVbeta3 integrin, FGFR-3, Keratinocyte Growth Factor, GM-CSF, M-CSF, RANKL, VLA-1, VLA-4, L-selectin, anti-

Id, E-selectin, HLA, HLA-DR, CTLA-4, T cell receptor, B7-1, B7-2, VNRintegrin, TGFbeta1, TGFbeta2, eotaxin1, BLYS (B-lymphocyte Stimulator), complement C5, IgE, IgA, IgD, IgM, IgG, factor VII, CBL, NCA 90, EGFR (ErbB-1), Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB4), Tissue Factor, VEGF, VEGFR, endothelin receptor, VLA-4, carbohydrates such as blood group antigens and related carbohydrates, Galili-Glycosylation, Gastrin, Gastrin receptors, tumor associated carbohydrates, Hapten NP-cap or NIP-cap, T cell receptor alpha/beta, E-selectin, P-glycoprotein, MRP3, MRP5, glutathione-S-transferase pi (multi drug resistance proteins), alpha-granule membrane protein(GMP) 140, digoxin, placental alkaline phosphatase (PLAP) and testicular PLAP-like alkaline phosphatase, transferrin receptor, Heparanase I, human cardiac myosin, Glycoprotein IIb/IIIa (GPIIb/IIIa), human cytomegalovirus (HCMV) gH envelope glycoprotein, HIV gp120, HCMV, respiratory syncytial virus RSV F, RSVF Fgp, VNRintegrin, Hep B gp120, CMV, gpIIbIIIa, HIV IIIB gp120 V3 loop, respiratory syncytial virus (RSV) Fgp, Herpes simplex virus (HSV) gD glycoprotein, HSV gB glycoprotein, HCMV gB envelope glycoprotein, Clostridium perfringens toxin and fragments thereof.

Preferred modular antibodies according to the invention are binding said target antigen with a high affinity, in particular with a high on and/or a low off rate, or a high avidity of binding. Usually a binder is considered a high affinity binder with a K_d of $<10^{-9}$ M. Medium affinity binders with a K_d of less than 10^{-6} up to 10^{-9} M may be provided according to the invention as well, preferably in conjunction with an affinity maturation process.

Affinity maturation is the process by which antibodies with increased affinity for antigen are produced. With structural changes of an antibody, including amino acid mutagenesis or as

a consequence of somatic mutation in immunoglobulin gene segments, variants of a binding site to an antigen are produced and selected for greater affinities. Affinity matured modular antibodies may exhibit a several logfold greater affinity than a parent antibody. Single parent antibodies may be subject to affinity maturation. Alternatively pools of modular antibodies with similar binding affinity to the target antigen may be considered as parent structures that are varied to obtain affinity matured single antibodies or affinity matured pools of such antibodies.

The preferred affinity matured variant of a modular antibody according to the invention exhibits at least a 10 fold increase in affinity of binding, preferably at least a 100 fold increase. The affinity maturation may be employed in the course of the selection campaigns employing respective libraries of parent molecules, either with modular antibodies having medium binding affinity to obtain the modular antibody of the invention having the specific target binding property of a $K_d < 10^{-8}$ M and/or a potency of $IC_{50} < 10^{-8}$ M. Alternatively, the binding potency or affinity may be even more increased by affinity maturation of the modular antibody according to the invention to obtain the high values corresponding to a K_d or IC_{50} of less than 10^{-9} M, preferably less than 10^{-10} M or even less than 10^{-11} M, most preferred in the picomolar range.

The IC_{50} , also called 50% saturation concentration, is a measure for the binding potency of a modular antibody. It is the molar concentration of a binder, which produces 50% of the maximum possible binding at equilibrium or under saturation. The potency of an antagonist is usually defined by its IC_{50} value. This can be calculated for a given antagonist by determining the concentration of antagonist needed to elicit half saturation of the maximum binding of an agonist. Elucidating an IC_{50} value is useful for comparing the potency

of antibodies or antibody variants with similar efficacies; however the dose-response curves produced by both drug antagonists must be similar. The lower the IC₅₀, the greater the potency of the antagonist, and the lower the concentration of drug that is required to inhibit the maximum biological response, like effector function or cytotoxic activity. Lower concentrations of drugs may also be associated with fewer side effects.

Usually the affinity of an antibody correlates well with the IC₅₀. The affinity of an antagonist for its binding site (K_i), is understood as its ability to bind to a receptor, which determines the duration of binding and respective agonist activity. Measures to increase the affinity by affinity maturation usually also increase the potency of binding, resulting in the respective reduction of IC₅₀ values in the same range of the K_d values.

The IC₅₀ and K_d values may be determined using the saturation binding assays well-known in the art.

The modular antibody according to the invention is preferably conjugated to a label or reporter molecule, selected from the group consisting of organic molecules, enzyme labels, radioactive labels, colored labels, fluorescent labels, chromogenic labels, luminescent labels, haptens, digoxigenin, biotin, metal complexes, metals, colloidal gold and mixtures thereof. Modified immunoglobulins conjugated to labels or reporter molecules may be used, for instance, in assay systems or diagnostic methods.

The modular antibody according to the invention may be conjugated to other molecules which allow the simple detection of said conjugate in, for instance, binding assays (e.g. ELISA) and binding studies.

In a preferred embodiment, antibody variants are screened using one or more cell-based or in vivo assays. For such assays, purified or unpurified modified immunoglobulins are typically added exogenously such that cells are exposed to individual immunoglobulins or pools of immunoglobulins belonging to a library. These assays are typically, but not always, based on the function of the immunoglobulin; that is, the ability of the antibody to bind to its target and mediate some biochemical event, for example effector function, ligand/receptor binding inhibition, apoptosis, and the like. Such assays often involve monitoring the response of cells to the antibody, for example cell survival, cell death, change in cellular morphology, or transcriptional activation such as cellular expression of a natural gene or reporter gene. For example, such assays may measure the ability of antibody variants to elicit ADCC, ADCP, or CDC. For some assays additional cells or components, that is in addition to the target cells, may need to be added, for example example serum complement, or effector cells such as peripheral blood monocytes (PBMCs), NK cells, macrophages, and the like. Such additional cells may be from any organism, preferably humans, mice, rat, rabbit, and monkey. Modular antibodies may cause apoptosis of certain cell lines expressing the target, or they may mediate attack on target cells by immune cells which have been added to the assay. Methods for monitoring cell death or viability are known in the art, and include the use of dyes, immunochemical, cytochemical, and radioactive reagents. For example, caspase staining assays may enable apoptosis to be measured, and uptake or release of radioactive substrates or fluorescent dyes such as alamar blue may enable cell growth or activation to be monitored.

In a preferred embodiment, the DELFIART EuTDA-based cytotoxicity assay (Perkin Elmer, MA) may be used.

Alternatively, dead or damaged target cells may be monitored by measuring the release of one or more natural intracellular components, for example lactate dehydrogenase.

Transcriptional activation may also serve as a method for assaying function in cell-based assays. In this case, response may be monitored by assaying for natural genes or immunoglobulins which may be upregulated, for example the release of certain interleukins may be measured, or alternatively readout may be via a reporter construct. Cell-based assays may also involve the measure of morphological changes of cells as a response to the presence of modular antibodies. Cell types for such assays may be prokaryotic or eukaryotic, and a variety of cell lines that are known in the art may be employed. Alternatively, cell-based screens are performed using cells that have been transformed or transfected with nucleic acids encoding the variants. That is, antibody variants are not added exogenously to the cells. For example, in one embodiment, the cell-based screen utilizes cell surface display. A fusion partner can be employed that enables display of modified immunoglobulins on the surface of cells (Wittrup, 2001, *Curr Opin Biotechnol*, 12:395-399).

In a preferred embodiment, the immunogenicity of the modular antibodies may be determined experimentally using one or more cell-based assays. In a preferred embodiment, ex vivo T-cell activation assays are used to experimentally quantitate immunogenicity. In this method, antigen presenting cells and naive T cells from matched donors are challenged with a peptide or whole antibody of interest one or more times. Then, T cell activation can be detected using a number of methods, for example by monitoring production of cytokines or measuring uptake of tritiated thymidine. In the most preferred embodiment, interferon gamma production is monitored using Elispot assays.

The biological properties of the modular antibody according to the invention may be characterized *ex vivo* in cell, tissue, and whole organism experiments. As is known in the art, drugs are often tested *in vivo* in animals, including but not limited to mice, rats, rabbits, dogs, cats, pigs, and monkeys, in order to measure a drug's efficacy for treatment against a disease or disease model, or to measure a drug's pharmacokinetics, pharmacodynamics, toxicity, and other properties. The animals may be referred to as disease models. Therapeutics are often tested in mice, including but not limited to nude mice, SCID mice, xenograft mice, and transgenic mice (including knockins and knockouts). Such experimentation may provide meaningful data for determination of the potential of the antibody to be used as a therapeutic with the appropriate half-life, effector function, apoptotic activity, cytotoxic or cytolytic activity. Any organism, preferably mammals, may be used for testing. For example because of their genetic similarity to humans, primates, monkeys can be suitable therapeutic models, and thus may be used to test the efficacy, toxicity, pharmacokinetics, pharmacodynamics, half-life, or other property of the modular antibody according to the invention. Tests of the substances in humans are ultimately required for approval as drugs, and thus of course these experiments are contemplated. Thus the modular antibodies of the present invention may be tested in humans to determine their therapeutic efficacy, toxicity, immunogenicity, pharmacokinetics, and/or other clinical properties. Especially those modular antibodies according to the invention that bind to single cell or a cellular complex through at least two binding motifs, preferably binding of at least three structures cross-linking target cells, would be considered effective in effector activity or preapoptotic or apoptotic activity upon cell targeting and cross-linking. Multivalent binding provides a relatively large association of

binding partners, also called cross-linking, which is a prerequisite for apoptosis and cell death.

The modular antibody of the present invention may find use in a wide range of antibody products. In one embodiment the modular antibody of the present invention is used for therapy or prophylaxis, e.g. as an active or passive immunotherapy, for preparative, industrial or analytic use, as a diagnostic, an industrial compound or a research reagent, preferably a therapeutic. The modular antibody may find use in an antibody composition that is monoclonal or polyclonal. In a preferred embodiment, the modular antibodies of the present invention are used to capture or kill target cells that bear the target antigen, for example cancer cells. In an alternate embodiment, the modular antibodies of the present invention are used to block, antagonize, or agonize the target antigen, for example by antagonizing a cytokine or cytokine receptor.

In an alternately preferred embodiment, the modular antibodies of the present invention are used to block, antagonize, or agonize growth factors or growth factor receptors and thereby mediate killing the target cells that bear or need the target antigen.

In an alternately preferred embodiment, the modular antibodies of the present invention are used to block, antagonize, or agonize enzymes and substrate of enzymes.

In a preferred embodiment, a modular antibody is administered to a patient to treat a specific disorder. A "patient" for the purposes of the present invention includes both humans and other animals, preferably mammals and most preferably humans. By "specific disorder" herein is meant a disorder that may be ameliorated by the administration of a pharmaceutical

composition comprising a modified immunoglobulin of the present invention.

In one embodiment, a modular antibody according to the present invention is the only therapeutically active agent administered to a patient. Alternatively, the modular antibody according to the present invention is administered in combination with one or more other therapeutic agents, including but not limited to cytotoxic agents, chemotherapeutic agents, cytokines, growth inhibitory agents, anti-hormonal agents, kinase inhibitors, anti-angiogenic agents, cardioprotectants, or other therapeutic agents. The modular antibody may be administered concomitantly with one or more other therapeutic regimens. For example, a modular antibody of the present invention may be administered to the patient along with chemotherapy, radiation therapy, or both chemotherapy and radiation therapy. In one embodiment, the modular antibody of the present invention may be administered in conjunction with one or more antibodies, which may or may not comprise a modular antibody of the present invention. In accordance with another embodiment of the invention, the modular antibody of the present invention and one or more other anti-cancer therapies is employed to treat cancer cells ex vivo. It is contemplated that such ex vivo treatment may be useful in bone marrow transplantation and particularly, autologous bone marrow transplantation. It is of course contemplated that the antibodies of the invention can be employed in combination with still other therapeutic techniques such as surgery.

A variety of other therapeutic agents may find use for administration with the modular antibody of the present invention. In one embodiment, the modular antibody is administered with an anti-angiogenic agent, which is a compound that blocks, or interferes to some degree, the development of blood vessels. The anti-angiogenic factor may,

for instance, be a small molecule or a protein, for example an antibody, Fc fusion molecule, or cytokine, that binds to a growth factor or growth factor receptor involved in promoting angiogenesis. The preferred anti-angiogenic factor herein is an antibody that binds to Vascular Endothelial Growth Factor (VEGF). In an alternate embodiment, the modular antibody is administered with a therapeutic agent that induces or enhances adaptive immune response, for example an antibody that targets CTLA-4. In an alternate embodiment, the modified immunoglobulin is administered with a tyrosine kinase inhibitor, which is a molecule that inhibits to some extent tyrosine kinase activity of a tyrosine kinase. In an alternate embodiment, the modular antibody of the present invention are administered with a cytokine. By "cytokine" as used herein is meant a generic term for proteins released by one cell population that act on another cell as intercellular mediators including chemokines.

Pharmaceutical compositions are contemplated wherein modular antibodies of the present invention and one or more therapeutically active agents are formulated. Stable formulations of the modular antibodies of the present invention are prepared for storage by mixing said immunoglobulin having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers, in the form of lyophilized formulations or aqueous solutions. The formulations to be used for in vivo administration are preferably sterile. This is readily accomplished by filtration through sterile filtration membranes or other methods. The modular antibody and other therapeutically active agents disclosed herein may also be formulated as immunoliposomes, and/or entrapped in microcapsules.

Administration of the pharmaceutical composition comprising a modular antibody of the present invention, preferably in the form of a sterile aqueous solution, may be done in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, intranasally, intraotically, transdermally, mucosal, topically (e.g., gels, salves, lotions, creams, etc.), intraperitoneally, intramuscularly, intrapulmonary (e.g., AERx™ inhalable technology commercially available from Aradigm, or Inhance™ pulmonary delivery system commercially available from Inhale Therapeutics), vaginally, parenterally, rectally, or intraocularly.

A preferred method according to the invention refers to a randomly modified nucleic acid molecule coding for an immunoglobulin, immunoglobulin domain or a part thereof which comprises at least one nucleotide repeating unit within a structural loop coding region having the sequence 5'-NNS-3', 5'-NNN-3', 5'-NNB-3' or 5'-NNK-3'. In some embodiments the modified nucleic acid comprises nucleotide codons selected from the group of TMT, WMT, BMT, RMC, RMG, MRT, SRC, KMT, RST, YMT, MKC, RSA, RRC, NNK, NNN, NNS or any combination thereof (the coding is according to IUPAC).

The modification of the nucleic acid molecule may be performed by introducing synthetic oligonucleotides into a larger segment of nucleic acid or by de novo synthesis of a complete nucleic acid molecule. Synthesis of nucleic acid may be performed with tri-nucleotide building blocks which would reduce the number of nonsense sequence combinations if a subset of amino acids is to be encoded (e.g. Yanez et al. Nucleic Acids Res. (2004) 32:e158; Virnekas et al. Nucleic Acids Res. (1994) 22:5600-5607).

The randomly modified nucleic acid molecule may comprise the above identified repeating units, which code for all known

naturally occurring amino acids or a subset thereof. Those libraries that contain modified sequences wherein a specific subset of amino acids are used for modification purposes are called "focused" libraries. The members of such libraries have an increased probability of an amino acid of such a subset at the modified position, which is at least two times higher than usual, preferably at least 3 times or even at least 4 times higher. Such libraries have also a limited or lower number of library members, so that the number of actual library members reaches the number of theoretical library members. In some cases the number of library members of a focused library is not less than 10^3 times the theoretical number, preferably not less than 10^2 times, most preferably not less than 10 times.

Usually libraries according to the invention comprise at least 10 fusion proteins or potential binding agents or variants of scaffold proteins, preferably at least 100, more preferred at least 1000, more preferred at least 10^4 , more preferred at least 10^5 , more preferred at least 10^6 , more preferred at least 10^7 , more preferred at least 10^8 , more preferred at least 10^9 , more preferred at least 10^{10} , more preferred at least 10^{11} , up to 10^{12} , in cases of in vitro display methods, such as ribosomal display, even higher number are feasible.

Various alternatives are available for the manufacture of the gene encoding the randomized library. It is possible to produce the DNA by a completely synthetic approach, in which the sequence is divided into overlapping fragments which are subsequently prepared as synthetic oligonucleotides. These oligonucleotides are mixed together, and annealed to each other by first heating to ca. 100°C and then slowly cooling down to ambient temperature. After this annealing step, the synthetically assembled gene can be either cloned directly, or it can be amplified by PCR prior to cloning.

Alternatively, other methods for site directed mutagenesis can be employed for generation of the library insert, such as the Kunkel method (Kunkel TA. Rapid and efficient site-specific mutagenesis without phenotypic selection. Proc Natl Acad Sci U S A. 1985 Jan;82(2):488-92) or the DpnI method (Weiner MP, Costa GL, Schoettlin W, Cline J, Mathur E, Bauer JC. Site-directed mutagenesis of double-stranded DNA by the polymerase chain reaction. Gene. 1994 Dec 30;151(1-2):119-23.).

For various purposes, it may be advantageous to introduce silent mutations into the sequence encoding the library insert. For example, restriction sites can be introduced which facilitate cloning or modular exchange of parts of the sequence. Another example for the introduction of silent mutations is the ability to "mark" libraries, that means to give them a specific codon at a selected position, allowing them (or selected clones derived from them) e.g. to be recognized during subsequent steps, in which for example different libraries with different characteristics can be mixed together and used as a mixture in the selection or panning procedure.

The invention also provides a method of producing an oligomer of modular antibody domains binding to a target comprising the steps of:

- providing a library of oligomers of modular antibody domains produced according to the inventive method as described
- contacting said library with said target in the presence of a scaffold ligand,
- selecting a library member binding to said target in the presence of a scaffold ligand, and
- manufacturing a preparation of the functional oligomer.

The scaffold ligand can be selected from the group consisting of an effector molecule, FcRn, Protein A and CDR target. As an

example, the effector molecule can be selected from the group consisting of CD64, CD32, CD16, Fc receptors.

The oligomers can be dimers selected from the group of VH/VL, CH1/CL, CH2/CH2, CH3/CH3, Fc and Fab, or single chains thereof

The method according to the invention can provide a library containing at least 10^2 independent clones expressing functional oligomers of modular antibody domains or variants thereof.

Libraries as used according to the invention preferably comprise at least 10^2 library members, more preferred at least 10^3 , more preferred at least 10^4 , more preferred at least 10^5 , more preferred at least 10^6 library members, more preferred at least 10^7 , more preferred at least 10^8 , more preferred at least 10^9 , more preferred at least 10^{10} , more preferred at least 10^{11} , up to 10^{12} members of a library, preferably derived from a parent molecule, which is a functional modular antibody as a scaffold containing at least one specific function or binding moiety, and derivatives thereof to engineer a new binding site apart from the original functional binding region of said parent moiety.

Usually the libraries according to the invention further contain variants of the modular antibody, resulting from mutagenesis or randomization techniques. These variants include inactive or non-functional antibodies. Thus, it is preferred that any such libraries be screened with the appropriate assay for determining the functional effect. Preferred libraries, according to the invention, comprise at least 10^2 variants of such modular antibodies, more preferred at least 10^3 , more preferred at least 10^4 , more preferred at least 10^5 , more preferred at least 10^6 , more preferred at least 10^7 , more preferred at least 10^8 , more preferred at least 10^9 , more preferred at least 10^{10} , more preferred at least 10^{11} , up

to 10^{12} variants or higher to provide a highly diverse repertoire of antibodies for selecting the best suitable binders. Any such synthetic libraries may be generated using mutagenesis methods as disclosed herein.

Preferably the library is a yeast library and the yeast host cell exhibits at the surface of the cell the oligomers, or monomers that form oligomers, with the biological activity. The yeast host cell is preferably selected from the genera *Saccharomyces*, *Pichia*, *Hansenula*, *Schizisaccharomyces*, *Kluyveromyces*, *Yarrowia* and *Candida*. Most preferred, the host cell is *Pichia* or *Saccharomyces cerevisiae*.

The invention further provides a high quality library containing at least 10^2 independent clones of functional dimers of modular antibody domains or variants thereof that are binding to a target and to a scaffold ligand. The target can be a ligand binding to a parent molecule subject to amino acid variation. The parent molecule can be a functional oligomer, in particular a functional Fc or a functional Fab, or part thereof.

As is well-known in the art, there is a variety of display and selection technologies that may be used for the identification and isolation of proteins with certain binding characteristics and affinities, including, for example, display technologies such as cellular and non-cellular, in particular mobilized display systems. Among the cellular systems the phage display, virus display, yeast or other eukaryotic cell display, such as mammalian or insect cell display, may be used. Mobilized systems are relating to display systems in the soluble form, such as in vitro display systems, among them ribosome display, mRNA display or nucleic acid display.

Methods for production and screening of antibody variants are well-known in the art. General methods for antibody molecular biology, expression, purification, and screening are described in *Antibody Engineering*, edited by Duebel & Kontermann, Springer-Verlag, Heidelberg, 2001; and Hayhurst & Georgiou, 2001, *Curr Opin Chem Biol* 5:683-689; Maynard & Georgiou, 2000, *Annu Rev Biomed Eng* 2:339-76.

A library according to the invention may be designed as a dedicated library that contains at least 50% specific formats, preferably at least 60%, more preferred at least 70%, more preferred at least 80%, more preferred at least 90%, or those that mainly consist of specific antibody formats. Such a preferred library mainly contains the same kind of library members having similar structural features. Specific antibody formats are preferred, such that the preferred library according to the invention is selected from the group consisting of a VH library, VHH library, Vkappa library, Vlambda library, Fab library, a CH1/CL library, an Fc library and a CH3 library. Libraries characterized by the content of composite molecules containing more than one antibody domains, such as an IgG library or Fc library are specially preferred. Other preferred libraries are those containing T-cell receptors, forming T-cell receptor libraries. Further preferred libraries are epitope or peptide libraries, wherein the fusion protein comprises a molecule with a variant of an epitope, also enabling the selection of competitive molecules having similar binding function, but different functionality. Exemplary is a TNFalpha library, wherein trimers of the TNFalpha fusion protein are displayed by a single genetic package.

Another important aspect of the invention is that each potential binding domain remains physically associated with the particular DNA or RNA molecule which encodes it, and in

addition, the fusion proteins oligomerize at the surface of a genetic package to present the binding polypeptide in the native and functional oligomeric structure. Once successful binding domains are identified, one may readily obtain the gene for expression, recombination or further engineering purposes. The form that this association takes is a "replicable genetic package", such as a virus, cell or spore which replicates and expresses the binding domain-encoding gene, and transports the binding domain to its outer surface. Another form is an in-vitro replicable genetic package such as ribosomes that link coding RNA with the translated protein. In ribosome display the genetic material is replicated by enzymatic amplification with polymerases.

Those cells or viruses or nucleic acid bearing the binding agents which recognize the target molecule are isolated and, if necessary, amplified. The genetic package preferably is M13 phage, and the protein includes the outer surface transport signal of the M13 gene III protein.

Preferably in the method of this invention the vector or plasmid of the genetic package is under tight control of the transcription regulatory element, and the culturing conditions are adjusted so that the amount or number of vector or phagemid particles displaying less than two copies of the fusion protein on the surface of the particle is less than about 20%. More preferably, the amount of vector or phagemid particles displaying less than two copies of the fusion protein is less than 10% the amount of particles displaying one or more copies of the fusion protein. Most preferably the amount is less than 1%.

The expression vector preferably used according to the invention is capable of expressing a binding polypeptide, and may be produced as follows: First a binding polypeptide gene library is synthesized by introducing a plurality of polynucleotides encoding different binding sequences. The plurality of polynucleotides may be synthesized in an appropriate amount to be joined in operable combination into a vector that can be propagated to express a fusion protein of said binding polypeptide. Alternatively the plurality of polynucleotides can also be amplified by polymerase chain reaction to obtain enough material for expression. However, this would only be advantageous if the binding polypeptide would be encoded by a large polynucleotide sequence, e.g. longer than 200 base pairs or sometimes longer than 300 base pairs. Thus, a diverse synthetic library is preferably formed, ready for selecting from said diverse library at least one expression vector capable of producing binding polypeptides having the desired preselected function and binding property, such as specificity.

The foregoing description will be more fully understood with reference to the following examples. Such examples are, however, merely representative of methods of practicing one or more embodiments of the present invention and should not be read as limiting the scope of invention.

Examples

Example 1: Construction of the non-focussed Fcab library (Fcab01) and phage surface display

The crystal structure of an IgG1 Fc fragment, which is published in the Brookhaven Database as entry 1OQO.pdb was used to aid in the design of the Fcab library.

The sequence which was used as the basis for construction of the Fcab library is given in SEQ ID No.1. In this sequence,

the first amino acid corresponds to Glu 216 of human IgG1 (EU numbering; according to the IMGT database (http://imgt.cines.fr/textes/IMGTrepertoire/Proteins/protein/human/IGH/IGHC/Hu_IGHCallgenes.html; lookup 2007 06 25), it is the first residue of the human IgG1 hinge region, which is given as: (E)PKSCDKTHTCPPCP) of the heavy constant chain hinge region of human IgG1.) The second-last residue of SEQ ID No.1 corresponds to Gly 446 of human IgG1 (EU numbering; IMGT: residue number 129 of the CH3 domain of human IgG1).

After detailed analysis of the structure of loqo.pdb and by visual inspection of the residues forming the loops which connect the beta strands, it was decided to randomize residues 144, 145 and 146, which are part of the loop connecting beta strand A-B as well as 198,199, 200, 203 and 204, which are part of the loop connecting beta strand E-F of SEQ ID No.1. In addition to the mutated residues, 5 residues were inserted at residue number 198 of SEQ ID No.1. In SEQ ID No.2, the sequence of the library insert of library Fcab01 is given in which all randomized residue positions as well as the 5 inserted residues are designated with the letter X.

The engineered gene was produced by a series of PCR reactions using degenerate primers followed by ligation of the resulting PCR products. To facilitate ligation, some of the codons of the nucleotide sequence coding for SEQ ID No.1 were modified to produce restriction sites without changing the amino acid sequences (silent mutations). For insertion into the cloning vector pHEN1 (Nucleic Acids Res. 1991 Aug 11;19(15):4133-7. Multi-subunit proteins on the surface of filamentous phage: methodologies for displaying antibody (Fab) heavy and light chains. Hoogenboom HR, Griffiths AD, Johnson KS, Chiswell DJ, Hudson P, Winter G.) in frame with the pelB secretion signal, the NcoI restriction site close to the 3' end of the pelB secretion signal was used. For the randomized residues, the

codon NNS (IUPAC code, where S means nucleotides C and G) was chosen which encodes all 20 naturally occurring amino acids, but avoids 2 out of 3 stop codons. Other codons such as for example the NNB (B meaning nucleotides T, C and G) can also be used. The engineered sequence is given as a nucleotide sequence in SEQ ID No.3. This sequence also includes the restriction sites used for cloning into the phagmid display vector pHEN1, namely an NcoI site at the 5' end and a NotI site at the 3' end.

The sequences of the PCR primers used for assembly of the mutated CH3 domain are given in SEQ ID No.4 through SEQ ID No.9.

SEQ ID No.4 (PCR primer EPKSNCO)
ccatggccgagcccaaatcttgtgacaaaactc

SEQ ID No.5 (PCR primer CH3LSAC)
agtcgagctcgtcacgggatgggggcaggg

SEQ ID No.6 (PCR primer CH3CSAC)
gtacgagctcnnnsnnsnnscaagtcagcctgacctgcctgg

SEQ ID No.7 (PCR primer CH3CHIN)
tgccaagcttgctgtagaggaagaaggagccg

SEQ ID No.8 (PCR primer CH3RHIN)
tgccaagcttaccgtgnnsnnsnnsaggtggnnnsnsgggaacgtcttctcatgctccg

SEQ ID No.9 (PCR primer CH3RNOT)
agttgcgccgctttaccggagacagggagag

Figure 1 shows a schematic presentation of the PCR fragments generated for assembly of the mutated gene, and the primers used therefore.

cDNA of the heavy chain of the human monoclonal antibody 3D6 (Felgenhauer M, Kohl J, Rüker F. Nucleotide sequences of the cDNAs encoding the V-regions of H- and L-chains of a human mono-clonal antibody specific to HIV-1-gp41. Nucleic Acids Res. 1990 Aug 25;18(16):4927.) was used as template for the PCR reactions. The 3 PCR products were digested with SacI and/or HindIII respectively and ligated together. The ligation product was further digested with NcoI and NotI and ligated into the surface display phagmid vector pHEN1, which had previously been digested with NcoI and NotI. The ligation product was then transformed into E. coli by electroporation. A number of selected clones were controlled by restriction analysis and by DNA sequencing and were found to contain the insert as planned, including the correctly inserted randomized sequences. For the following steps of phage preparation, standard protocols were followed. Briefly, the ligation mixture was transformed into E. coli TG1 cells by electroporation. Subsequently, phage particles were rescued from E. coli TG1 cells with helper phage M13-KO7. Phage particles were then precipitated from culture supernatant with PEG/NaCl in 2 steps, dissolved in water and used for selection by panning or, alternatively, they were stored at minus 80 °C.

Example 2: Construction of the focussed Fcab library (Fcab02) and phage surface display

As described in example 1, an Fcab library was prepared in which the randomized library positions are fully randomized, i.e. they are encoded by a codon such as NNS, NNB, NNK, NNN or others are used.

For clarity, the meaning of the letters such as N, B, S or K is defined by the IUPAC nucleotide ambiguity code, which is given in the following table:

Table 1. IUPAC nucleotide ambiguity code

Symbol	Meaning	Nucleic Acid
A	A	Adenine
C	C	Cytosine
G	G	Guanine
T	T	Thymine
U	U	Uracil
M	A or C	
R	A or G	
W	A or T	
S	C or G	
Y	C or T	
K	G or T	
V	A or C or G	
H	A or C or T	
D	A or G or T	
B	C or G or T	
X	G or A or T or C	
N	G or A or T or C	

Source: Nomenclature for incompletely specified bases in nucleic acid sequences: recommendations 1984. A Cornish-Bowden, *Nucleic Acids Res.* 1985 May 10; 13(9): 3021-3030.

These codons given above are designed such that all 20 amino acids are encoded by them. It may be preferable to choose subsets out of the possible amino acids. Examples can be found in the literature (Fellouse FA, Li B, Compaan DM, Peden AA, Hymowitz SG, Sidhu SS. Molecular recognition by a binary code. *J Mol Biol.* 2005 May 20;348(5):1153-62. Epub 2005 Apr 1.; Fellouse FA, Wiesmann C, Sidhu SS. Synthetic antibodies from a four-amino-acid code: a dominant role for tyrosine in antigen recognition. *Proc Natl Acad Sci U S A.* 2004 Aug 24;101(34):12467-72. Epub 2004 Aug 11.). Focused libraries which for example allow for only 4 different amino acid types can be constructed e.g. by employing the codon KMT, which codes for the amino acids Ser, Tyr, Ala and Asp.

A focused Fcab library, designated Fcab02, has been constructed in the same way as described in example 1, except that the NNS codons were replaced by KMT codons.

Therefore, the letter "X" in SEQ ID No.2 now means "S, Y, A and D" (Ser, Tyr, Ala and Asp) in order to describe the focused library Fcab02

Example 3: Construction of a phage surface display library with additional amino acid residues between the library insert (binding partner) and p3

In order to investigate accessibility of the potential binding site of the displayed protein a binding assay is performed: the phage suspension is reacted with anti-myc mAb 9E10-coated microplates (or immunotubes). After washing, the bound phages are detected with anti-M13-enzyme conjugate. As a control, helper phage - which does not display the protein fusion and the myc-tag is reacted with the plates. Other controls are reaction of phages with non-coated plates and reaction of phages with antiserum recognizing the p3 -fusion partner of the phages.

Ideally, the anti-myc-reactivity of phages displaying the p3 - fusion protein should give very clear ELISA readouts whereas helper phage reactions to anti-myc-mAb should not be above background (non-coated plates). The structure of a CH3 dimer displayed at the surface of an M13 phage through binding to protein III as an anchor is such, that each CH3 is anchored to protein III using various linker length and compositions. Thus, the CH3 dimer is preferably displayed by two anchors.

Linker optimization:

The linker between the protein to be displayed and the anchor protein of the genetic package (in case of filamentous phage e.g. p3 , p8, pX, pIX, pVII) is especially important if the potential binding site of the displayed molecule is in spatial vicinity of the phage particle. In antibody libraries

utilizing variable domains and antigen binding sites formed by CDR-loops and display of the library members as amino-terminal fusion to p3 the potential antigen binding site is directed away from the phage particle. Therefore, the linker structure between library members and the phage coat protein is less important. Engineering the bottom loops of immunoglobulin domains and performing phage display may however be an inefficient process and decrease yields of antigen binding clones or even preclude it. Varying the linker between a library member protein and its fusion partner on the surface can solve or may at least reduce this problem.

In order to select for optimal linker sequences (in terms of length and flexibility as well as stability) a library of linkers can be prepared in which the anchor protein at the surface of the genetic replicable package is fused to a known binding protein which is for sterical reasons notoriously difficult to select for.

This library of sequences can be varied in length and amino acid content.

Selection methods of the linker library for optimal linkers depend on the application but basically it should be for selecting all properties one wishes to have in a certain methodology. Enrichment against a difficult to select for antigen may yield linker sequences which allow library members a good access to the antigen. Incubation in protease solutions or under other harsh conditions or frequent passaging through host cells under proteolytic conditions (e.g. old microbial cultures) may be an appropriate selection for stable display linkers.

A library of linkers may be produced by any well known library technology. Synthetic linker sequence lengths may vary between 10-500 amino acids. Alternatively, linker can be complete proteins known to be of flexible nature.

Linker optimization Fcab01:

As an example, library Fcab01 (as described in example 1) can be used. Originally, this library is cloned in the phagmid display vector pHEN1, using NcoI and NotI restriction sites. When cloned in this manner, 18 amino acid residues are in between the C-terminal amino acid residue of the Fcab01 library insert and the N-terminal amino acid residue of phage M13 p3. The sequence of this junction region is given in SEQ ID No.10 SPGKAAAEQKLISEEDLNGAATVES - and is explained as follows: the first 4 residues, SPGK, are the 4 C-terminal residues of the Fcab01 library insert, followed by the amino acid sequence AAA, which is the amino acid residues encoded by the NotI restriction site, followed by the sequence EQKLISEEDL, which is the myc epitope, followed by NGAA, after which there is an amber stop codon, which is translated to Glutamine (Q) in amber suppressor strains of E. coli such as TG1. The C-terminal 4 residues of SEQ ID No.10, TVES, are the N-terminal 4 residues of phage M13 p3 as present in the vector pHEN1.

In order to construct a phage which displays an Fcab insert with an increased distance between the Fcab (the binding partner) and the body of the phage (the genetic package), 5 additional residues were inserted at the C-terminus of the Fcab insert FcabRGD4, directly upstream of the NotI cloning site, resulting in the clone FcabRGD4L. FcabRGD4 is an Fcab that has an integrin-binding RGD motif inserted in the EF-loop of the CH3 domain and which binds to $\alpha v \beta 3$ -integrin in ELISA. As an increased-length linker sequence, the amino acid sequence EGGGS, which appears 8 times in the phage M13 p3 sequence was used. The resulting amino acid sequence of FcabRGD4L as expressed after cloning in pHEN1 is given in SEQ ID No.11. In SEQ ID No.11, amino acid residues 198-204 represent the RGD motif, amino acid residue 237 is the C-terminal residue of the Fcab insert, residues 238-242 represent the inserted linker

sequence (which is the difference to unmodified pHEN1), which is followed by myc tag, amber stop codon and the p3 sequence.

For cloning of the construct, the FcabRGD4 sequence was amplified from pHENFcabRGD4 (SEQ ID No.12) using PCR primers EPKSNCO (SEQ ID No.4) and CH3rlink actagcggccgcagagccaccaccctccttaccgagacaggagag (SEQ ID No.13) and cloned via NcoI and NotI restriction sites into the vector pHEN1. The resulting vector, pHENFcabRGD4L (SEQ ID No.14) has the additional linker sequence at nucleotide positions 3057-3071.

The two phagemid vectors, pHENFcabRGD4 and pHENFcabRGD4L were transformed into E.coli TG1. Subsequently, phage particles were rescued from E. coli TG1 cells with helper phage M13-KO7. Phage particles were then precipitated from culture supernatant with PEG/NaCl in 2 steps, dissolved in water and used for ELISA.

Phage ELISA was performed as follows:

The phage suspension is reacted with $\alpha v\beta 3$ -integrin-coated microplates (or immunotubes). After washing, the bound phages are detected with anti-M13-enzyme conjugate. As controls, helper phage - which does not display the protein fusion and the myc-tag is reacted with the plates as well as phage particles carrying wtFcab on their surface. Other controls are reaction of phages with non-coated plates and reaction of phages with antiserum recognizing the Fcab-fusion partner of the phages. Phage particles with the increased-length linker resulting from pHENFcabRGD4L react more readily with $\alpha v\beta 3$ -integrin than phage particles with the original linker as contained in pHENFcabRGD4, and therefore give a stronger signal in ELISA.

Phage selections can be performed in which phage particles with wtFcab are mixed with small amounts of phage particles carrying either FcabRGD4 or FcabRGD4L. After several (typically 3-5) rounds of panning, preferentially phages displaying FcabRGD4L are selected.

Example 4: Fcab library design

("Fcab" is a registered trademark of f-star Biotechnologische Forschungs- und Entwicklungsges.m.b.H.)

Design of Fcab Libraries (illustrated in Figure 2): amino acid positions in non CDR-loops of CH3 constant domains of antibodies are considered for randomization. Especially loops A-B, C-D and E-F are considered as they are on one side of the domain. Some of the design criteria for randomization at a certain position are described herein.

Amino acids frequently involved in antigen antibody interactions are described herein to be included in a focused library. Here the amino acids Ala, Asp, Ser and Tyr are used to design the focused library.

Libraries with restricted amino acid utilization have been shown to be sufficient to generate binders against virtually any antigen (Sidhu & Fellhouse, NATURE CHEMICAL BIOLOGY VOLUME 2 page 682ff; Koide et al PNAS, volume 104 p6632- 6637). The advantage of such restricted (or focused) libraries is that they can be covered completely by current technologies. Ideally, the amino acid utilization reflects a natural amino acid utilization of ligand receptor binding. However, even libraries utilizing only 2 amino acids (Tyrosine and Serine) have been reported to yield good selection results (in terms of frequency of binders against different binders and in terms of affinity).

Loop flexibility:

Certain loop structures may be required by the scaffold protein in order to keep the overall natural structure. Randomizing many amino acid positions in loops and even elongation of loops may be facilitated by building certain sequences either on one or on both sides of the randomized

positions. These sequences may be flexible sequences in order to allow to compensate for any tensions with certain library sequences in such a position.

Table 2: Exemplary Fcab™ libraries, focused and non-focused

	# of randomized positions	Theoretical diversity on amino acid level	Number of independent bacterial clones
Fcab01	13	8.2×10^{16}	0.6×10^9
Fcab02	13, focused	6.7×10^7	0.6×10^9
Fcab03	13	8.2×10^{16}	1.0×10^9
Fcab04	13, focused	6.7×10^7	0.8×10^9
Fcab05	15	1.3×10^{18}	0.8×10^9
Fcab06	15, focused	1.3×10^9	1.0×10^9

Fcab01 library is described in the examples above. The sequence space of the focused library designs Fcab02, Fcab04 and Fcab06 are covered by the actual bacterial library sizes of approximately 10^9 . In contrast, the completely randomized libraries Fcab01, Fcab03 and Fcab05 are actually grossly underrepresented.

Example 5: Cloning of yeast display libraries by homologous recombination

Vector

pYD1 (Invitrogen) is used as the basic vector. The vector is modified as follows, in order to remove an XhoI site: pYD1 is cleaved with XhoI, treated with Klenow fragment of DNA polymerase and religated. The resulting sequence is given in pYD1dX (SEQ ID No.15).

pYD1dX contains a unique BamHI restriction site at position 921/925 and a unique NotI restriction site at position 963/967. It is opened with these two restriction enzymes.

An insert encoding CH1-hinge-CH2-CH3 from human IgG1 is prepared by PCR from cDNA encoding the heavy chain of a human

IgG1 monoclonal antibody. In this insert, a point mutation is introduced using standard procedures to mutate the C-terminal Cystein residue of the CH1 domain to a Serine. The insert is amplified using PCR primers that attached a BamHI and a NotI restriction site to both ends respectively. These restriction sites are then used for cloning the insert into pYD1dX to yield the display vector pYD1dXFc (SEQ Id No.16). The mutated codon at the C-terminus of the CH1 domain (Cys to Ser) is at positions 1233-1235 in the sequence pYD1dXFc. The stop codon of the insert is at position 1917/1919.

This vector is used as a positive control for the display of human CH1-hinge-CH2-CH3 on the surface of yeast and as a starting point for the construction of the vector pYD1CH12 (see below).

Cloning of libraries

Cloning of libraries in which mutations are introduced into structural loops of CH3 domains is performed in yeast by homologous recombination (gap repair). For this purpose, a recipient vector is prepared that lacks the CH3 domain: pYD1dXFc is cleaved with XhoI (position 1603/1607) and NotI (position 1921/1925), the large fragment is prepared by preparative gel electrophoresis, treated with Klenow fragment of DNA polymerase and re-ligated. This procedure reconstitutes a unique XhoI site (position 1603/1607) and yielded vector pYD1CH12 (SEQ ID No.17). pYD1CH12 is subsequently cleaved with XhoI and is used as recipient vector for gap repair in yeast.

As a source of insert, Fcab libraries Fcab01 (SEQ ID No.18), Fcab02 (SEQ ID No.19), Fcab03 (SEQ ID No.20), Fcab04 (SEQ ID No.21), Fcab05 (SEQ ID No.22) and Fcab06 (SEQ ID No.23) are used. These libraries are prepared by standard DNA synthesis, and contain randomized residues as well as inserted residues

in the AB loop (between residues 359 and 361 (EU numbering)) as well as in the EF loop (between residues 413 and 419 (EU numbering)) of the CH3 domain of human IgG1. From this synthetic DNA, the insert for gap repair in yeast is amplified by PCR using PCR primer pair gapch35 caacaaggccctgcctgccccatcgagaagaccatctccaaggccaagggccagcctcgagaaccacaggtgtacaccctgccc (SEQ ID No.24) und gapfcs3 gagaccgaggagaggggttagggataggcttaccttcgaagggccctctagactcgatcgagcggccgctcatttaccggagacagggagagctcttc (SEQ ID No.25). 100 µg of XhoI cleaved vector pYD1CH12 and 100 µg of insert are mixed and transformed in Saccharomyces strain EBY100 (Invitrogen) using the Lithium acetate procedure according to the following protocol, which is upscaled by a factor 100 to transform the required amount of cells and of DNA. Briefly, for a single transformation of 1µg vector DNA and 1 µg insert DNA, 10 ml of YPD (2% peptone, 2% dextrose (D-glucose)) are inoculated with a yeast colony and shaken overnight at 30°C. The OD600 of the overnight culture is determined and the culture diluted to an OD600 of 0.4 in 50 ml of YPD and grown for an additional 6 hours. Cells are pelleted at 2500 rpm and resuspended in 40 ml distilled water. Cells are pelleted again at 2500 rpm and resuspended in 100mM LiAc, followed by incubation at 30°C for 30 minutes. 1µg vector DNA, 1 µg insert and 50 µl denatured sheared salmon sperm DNA (2mg/ml) are mixed with 300 µl of the yeast suspension. 700 µl of a solution of 200mM Li-acetate and 40% PEG-3350 are added and mixed with the yeast/DNA suspension, followed by incubation at 30°C for 30 minutes. 88 µl DMSO are added, mixed and the mixture is incubated at 42°C for 40 minutes, followed by centrifugation in a microcentrifuge for 10 seconds. The supernatant is then removed, the cell pellet is resuspended in 1 ml distilled water. The pellet is then resuspended in 50-100 µl TE and plated on minimal dextrose plates containing leucine (10 g/l yeast nitrogen base, 20 g/l dextrose, 0.1 g/l leucine, 15 g/l agar). After incubation of the plates at 30°C for 2 to

4 days single colonies appeared that are subsequently harvested.

Cultivation - Induction

The harvested yeast libraries (yFcab libraries) are inoculated in 10ml SD-CAA medium (10 g/l yeast nitrogen base, 10 g/l casamino acids, and 20 g/l dextrose, 0.1 g/l leucine, 9.67 g/l NaH₂PO₄·2H₂O and 10.19 g/l Na₂HPO₄·7H₂O) and grown on a shaker at 250rpm at 28°C for 6-8 hours. The OD₆₀₀ of the culture is determined, and the culture is diluted to an OD₆₀₀ of 0.2, and grown under the same conditions until an OD₆₀₀ of 1-2 is reached. Cells are harvested by centrifugation (3000rpm / 5min / 4°C) and resuspended in induction medium SG/R-CAA (10 g/l yeast nitrogen base, 10 g/l casamino acids, and 20 g/l galactose, 10g/l raffinose, 0.1 g/l leucine, 9.67 g/l NaH₂PO₄·2H₂O and 10.19 g/l Na₂HPO₄·7H₂O). Cultures are induced by incubation for 2 days on a shaker at 250rpm at 20°C and subsequently analysed and sorted.

Quality control of yFcab libraries.

yFcab libraries are tested for their expression level and quality of expressed Fcab's two days after induction with SD-CAA medium. The expression level is tested using a polyclonal anti human IgG-Fc antiserum (Sigma). For this purpose 0.5x10⁶ library cells are diluted in 1 ml staining buffer (SB), which comprises PBS with 2% BSA. Cells are pelleted and stained with 100µl SB containing 1/2000 diluted anti human IgG-Fc-PE antiserum (Sigma) for 30 min on ice, washed twice with SB and subsequently analyzed in the FACS. In general 70%-80% of all cells in each library express Fcabs on their cell surface. To test correct folding of Fcabs, staining with Protein A is performed. Again 0.5x10⁶ library cells are diluted in 1 ml

staining buffer SB, cells are pelleted and stained with 100µl SB containing 1µg/ml Prot-A-FITC (Fluka) for 30 'on ice, washed twice with SB and subsequently analyzed in the FACS. In general, the yFcab libraries as described above show $\geq 40\%$ Prot A positive cells. In order to test whether the Fcabs are expressed as dimers on the surface of the cells a staining with human CD64 is performed. The affinity of CD64 is too low for efficient monomeric binding therefore CD64 complexes with dimers are used. For this purpose e.g. 1µg recombinant CD64 from R&D Systems (containing a HIS-tag) is mixed with 1 µg anti Penta HIS-alexafluor 488 (from Qiagen) in 1ml SB (total volume). yFcab libraries are tested for binding to CD64 by incubating the 5×10^5 cells with 100 µl of the complex-mixture for 30 minutes on ice, as control the cells are incubated with equivalent of the anti HIS-alexafluor 488 alone (1/200 dilution in SB). After incubation the cells are washed twice with ice cold SB and analysed in the FACS. In general $> 50\%$ of all cells in each library express dimeric Fcabs on their cell surface.

Biotinylation of antigen (Her2)

Recombinant antigen e.g. Her2 (Bendermedsystems) was done with the EZ link system of Pierce according to the manufacturers instruction. In short, the antigen is dialyzed against PBS, diluted to 1mg/ml in PBS and mixed with 10mM sulfo-LC-LC-biotin (EZ link, Pierce), which was predissolved in water. The final ratio between antigen and biotin is 1:3 and the mixture is incubated at room temperature for 30'. Afterwards the mixture is "dialyzed" against PBS using Vivaspin MWC03000 (Sartorius) columns (5x8', 4000rpm). Finally the concentration of the biotinylated antigen (Her2) is tested by HPLC and aliquots are stored at -20° C.

The quality of the biotinylated antigen is tested by ELISA. First the plates are coated with an anti-Her2 antibody (e.g. Herceptin) at 10µg/ml in PBS, 100 µl/well overnight at 4°C, after this the plate is washed 3x with washing buffer (WB) (PBS + 0.05% Tween20) and blocked by blocking buffer (BB) (PBS + 2% BSA) 1h at room temperature. After 3x washing with WB, different concentrations of Her2-biotin are added in 100 µl/well BB for 1h at room temperature, followed by 3x washing with WB. Finally the plate is incubated with 1:25000 streptavidin-HRP (GE healthcare) in BB for 1h at room temperature and washed 3x with WB. Colour is developed by adding 100 µl/well of the substrate TMB (Sigma) after ~10 minutes the reaction is stopped by adding 100 µl/well of 30% H₂SO₄. The results is analysed with an ELISA reader at 450-630nm.

Example 6: Production of antigen specific (Her2) Fcabs

Selection of antigen specific (Her2) Fcabs using FACS

First selection round:

Two days before FACS sorting a yeast library containing 2.5 x10⁸ individual Fcab clones is induced with SG/R-CAA medium to express the Fcabs on their cell surface as described above. After two days, the amount of cells covering e.g. 10 times the library (= 2.5 x10⁹) is incubated for 30 minutes on ice with 500 nM biotinylated antigen (Her2) in 2ml SB. Then the cells are washed once with cold SB and subsequently incubated for 30' on ice with streptavidin-PE (from R&D systems) diluted 1 : 100 in SB. The cells are washed twice with ice cold SB and diluted to an end concentration of 1x10⁹ cells/ml. Control stainings with 5x10⁶ cell/ml in 100 µl are made with streptavidin-PE only, in the absence of antigen. Both the complete library and the control stainings are analysed in e.g. a FACS ARIA from BD. To set the gates for sorting the control cells are used. First a FSC/SSC gate (G1) is set to

identify healthy yeast cells, from G1 a FSC-width versus FSC-area plot is made and only non-aggregating cells are selected in a new gate (G2). Cells in G2 are subsequently analysed for reactivity with streptavidin-PE using FSC versus FL-2 (PE channel). G3 is set to include 0.1% of (false) positive cells. Subsequently, at least 5×10^8 stained cells (twice the library size ideally more) are analysed with the settings as indicated above and the cells in G3 are sorted into a tube containing 2-3 ml SD-CAA medium. Roughly 5×10^5 cells (Pool1) are harvested in the first round of selection and propagated for 1 to 2 days, after which the cells can be stored at -80°C and aliquots can be induced to express the Fcabs as described above. After two more days the next selection round can take place.

Second selection round:

Pool1 selected in round 1 are induced to express the Fcab on their surface as described above. At least 5×10^6 cells (comprising multiple copies of Pool1) are incubated for 30' on ice with 500 nM biotinylated antigen (Her2) in 1 ml SB. Then the cells are washed once with cold SB and subsequently incubated for 30' on ice with streptavidin-PE (from R&D systems) diluted 1 in 100 in SB together with $2 \mu\text{g/ml}$ Protein A-FITC (Fluka). Next the cells are washed twice with ice cold SB and diluted to an end concentration of $\sim 2 \times 10^6$ cells/ml. In addition, control stainings are made in which 5×10^6 cells/ml of Pool1 in 100 μl cells are incubated with a mixture of Prot A and streptavidin-PE as indicated above, but without the incubation with the antigen (Her2). In addition, 5×10^5 cell in 100 μl of a yeast clone expressing Fcab wt non randomized Fc fragment) is stained with Prot A -FITC as described above in the absence of streptavidin-PE. Fcab-wt expressing cells are analysed in e.g. a FACS ARIA from BD to set gates for sorting. First a FSC/SSC gate (G1) is set to identify healthy yeast cells, from G1 a FSC-width versus FSC-area plot is made and only non aggregating cells are selected in new gate (G2).

Cells in G2 are subsequently analysed for Protein A expression using FSC versus FL-1 (FITC). G3 is set to cover strong Prot A positive cells (50- 60% of parent gate) and G4 is set to cover weak Prot A positive cells (20-30% of parent cells). G3+G4 will include roughly 70-80% of all cells in G2. Now the Pool cells stained for streptavidin-PE in the presence of Prot A-FITC are used to set the rest of the sorting gates. First G1 and G2 are checked with the Pool cells and if necessary adjusted. Pool cells will have lesser events in G3 and maybe also in G4 indicating that not all cells in Pool1 express Fcabs that are folded as the Fcab-wt. Using the control stained Pool cells a new gate is prepared both for G3 and G4. The new gates are set in a plot FSC and FL-2 (PE). Gate (G5) is prepared that includes 0.1% (false) streptavidin positive cells in G3 and the same is done for cells in G4 resulting in G6. In the next step at least 5×10^6 cells stained for Her2-biotin + streptavidin-PE and Prot A-FITC are sorted by the FACS-ARIA. Cells are collected from G5 (Pool2.1 and G6 (Pool2.2) in separate tubes containing 2-3 ml yeast culture medium. Between 10 and 1000 clones can be expected from both gates. Both new pools are propagated for 1 or 2 days and stored at -80° C. Cells from 2.1 and 2.2 may be either used for direct further sorting in a third round or they may be subjected, (preferably after mixing the two clone together again) to a round of additional randomization of the AB loop (affinity maturation) before they are further sorted in FACS.

Affinity maturation for selected clones/pools

For affinity maturation, diversity is introduced in selected clones or in pools of selected clones in the AB loop. For this purpose, a PCR is made with a primer that contained degenerate codons at positions 359, 360 and 361 (EU numbering) (primer Abmut, gaaccacaggtgtacaccctgccccatcccgggatgagctggnbnnbnnbca ggtcagcctgacctgcc tgggtcaaag, SEQ ID No.26) or alternatively with a primer that contained degenerate codons at positions

358, 359, 360, 361 and 362 (EU numbering) (primer Abmut2LR, gaaccacaggtgtacaccctgcccccatcccgggatgagnnbnnbnnbnnbnnbgtcagcctgacctgcctgggtcaaag, SEQ ID No.27). The second primer used in these PCRs is gapfcs3 in both cases. In order to create flanking sequences for efficient gap repair in yeast, the resulting PCR products are further amplified with the primer pair gapch35 and gapfsc3 and subsequently transformed in *Saccharomyces cerevisiae* strain EBY100 by Lithiumacetate transformation together with XhoI cleaved pYD1CH12 as described above. As alternative primers for randomization of the described residues in the AB loop, primers such as Abmut1L (gaaccacaggtgtacaccctgcccccatcccgggatgagnnbnnbnnbnnbcaggtcagcctgacctgcctgggtcaaag, SEQ ID No.28) or Abmut1R (gaaccacaggtgtacaccctgcccccatcccgggatgagctgnbnnbnnbnnbgtcagcctgacctgcctgggtcaaag, SEQ ID No.29) can also be used. In an analogous manner, residues in the EF loop can be randomized e.g. by total randomization or by randomization using spiked oligonucleotides as primers or by similar mutagenesis techniques. The Abmut primer will result in 8000 new variants (Pool2.3) of each clone and the Abmut2LR primer will lead to 3×10^6 new variants (Pool2.4). Therefore Pools 2.3. and 2.4 will both result in new libraries of approximately 10^8 individual since the starting material (Pool2.1+2.2) already contains approximately 10-1000 clones.

Third selection round

Affinity matured pools 2.3 and 2.4 and if necessary Pool2.1 (only the Prot A positive cells are preferred) are induced to express Fcabs on their cell surface as described above and subsequently sorted as described for "Second selection round", with exception that the Pools 2.3 and 2.4 are much bigger and therefore staining volumes for the pools are equal to those of the library staining described in "First selection round". In the third selection round, only Her2 positive/Prot A positive cells are sorted. Pools derived from these selections contain

typically > 20% Her2/Prot A positive cells. If not then a fourth and fifth (or even more) round(s) of selection combining Prot A with Her2 can be performed.

Clone analyses:

Individual clones from pools containing Her2/Prot A cells (>20% is preferred) are prepared either by plating the pools on agar plates with SD-CAA medium or by sorting the singles cells (=clones) directly from the FACS ARIA onto the plates without generating a pool. Clones are allowed to grow and are transferred to liquid cultures and stored in -80°C. Aliquots of the clones are subsequently induced to express Fcabs on their cell surface as described above and screened for a number of parameters in the FACS. These parameters may be: a dose response range of the antigen used for selection (Her2) with and without the presence of Prot A- FITC, CD64 staining as described above. In addition using similar staining protocols a number of irrelevant biotinylated antigen can be screened to identify non-cross reacting Fcabs.

It is to be expected that, after several rounds of selecting antigen (Her2) + Prot A positive cells, a large percentage of clones show >25% antigen (Her2) positivity when stained with 500 nM antigen (Her2) and > 70% Prot A positivity when stained with 2µg/ml Prot A-FITC. In most of the cases these clones will also show > 50% CD64 binding. Thus mimicking the Prot A and CD64 levels of non-randomized Fc fragments (Fcab wt) expressed on yeast.

Clones selected as described above with characteristics as described above are now ready to be produced as soluble molecules. This can be done by transient transfection or by stable transfection of the Fcab DNA into new host cells. For this purpose the DNA from individual yeast clones is isolated using standard procedures. The relevant DNA coding for the complete CH3 domain or only the part of the CH3 domain that is

randomized in the library is amplified by PCR and transferred into a new expression vector containing the missing part of the Fcab + a suitable promoter and one of more selection markers such as G418, that allows selection of transfected cells out of a pool of non transfected cells. The new vector is then e.g. transiently transfected into a new host cell such as HEK293 or CHO. The host cells are allowed to recover and are subsequently cultured for a number of days. The supernatant of the cultures with contain the soluble Fcab which can be used for further testing with or without purification over e.g. Prot A. Stable cell lines can also be made by standard procedures.

Table 2. Sequences of selected Her2 clones: with reference to numbering of Seq.ID No.1

Clone name	AB loop AA143ff	EF Loop AA198ff
Fcab wt	LTKNQ	-----DKSRWQQ
y-Her.C2-P3.1-1	LDNSQ (SEQ ID No.30)	IRSSVGSRRWWS (SEQ ID No.51)
y-Her.C2-P3.1-3	YEGSS (SEQ ID No.31)	ARYSPRMLRWAH (SEQ ID No.52)
y-Her.C2-P3.1-5	YMSAD (SEQ ID No.32)	SRRDSSLLRWAH (SEQ ID No.53)
y-Her.C2-P3.1-6	YRRGD (SEQ ID No.33)	APGSKGYRRWAL (SEQ ID No.54)
y-Her.C2-P3.1-8	LMSRQ (SEQ ID No.34)	DKPFWGTSRWSR (SEQ ID No.55)
y-Her.C2-P3.1-16	LHLAQ (SEQ ID No.35)	SINDLINHRWPY (SEQ ID No.56)
y-Her.C2-P3.1-18	YLSKD (SEQ ID No.36)	MWGSRDYWRWSH (SEQ ID No.57)
y-Her.C2-P3.2-3	YRSGS (SEQ ID No.37)	NSGSAMMVRWAH (SEQ ID No.58)
y-Her.C2-P3.2-9	LRDGQ (SEQ ID No.38)	QRSRLSRQRWWR (SEQ ID No.59)
y-Her.C2.P4.2-1	YSANT (SEQ ID No.39)	ARYSPRMLRWAH (SEQ ID No.60)
y-Her.C2.P4.2-3	YASNT (SEQ ID No.40)	ARYSPRMLRWAH (SEQ ID No.61)
y-Her.C2.P4.2-4	YSDGD (SEQ ID No.41)	ARYSPRMLRWAH (SEQ ID No.62)
y-Her.C2.P4.2-5	YSGGS (SEQ ID No.42)	ARYSPRMLRWAH (SEQ ID No.63)
y-Her.C2.P4.2-6	YGRDS (SEQ ID No.43)	ARYSPRMLRWAH (SEQ ID No.64)
y-Her.C2.P4.2-8	YAGGT (SEQ ID No.44)	ARYSPRMLRWAH (SEQ ID No.65)
y-Her.C2.P4.2-10	YSSDS (SEQ ID No.45)	ARYSPRMLRWAH (SEQ ID No.66)
y-Her.C2.P4.2-12	YHSGS (SEQ ID No.46)	ARYSPRMLRWAH (SEQ ID No.67)
y-Her.C2.P4.2-15	YLNS (SEQ ID No.47)	ARYSPRMLRWAH (SEQ ID No.68)
y-Her.C2.P4.2-18	YGSEE (SEQ ID No.48)	ARYSPRMLRWAH (SEQ ID No.69)
y-Her.C2.P4.2-19	YRSGE (SEQ ID No.49)	ARYSPRMLRWAH (SEQ ID No.70)
y-Her.C2.P4.2-20	YGTDD (SEQ ID No.50)	ARYSPRMLRWAH (SEQ ID No.71)

Example 7: yeast display of 4D5 Fab

For the display of a Fab fragment on yeast, the yeast display vector pYD1 (Invitrogen) (SEQ ID No.72) is modified as follows:

A NheI restriction site is introduced by site directed mutagenesis at position 581/586 to yield the modified vector pYD1Nhe (SEQ ID No.73). This vector is restricted with NheI and PmeI, to yield 3 fragments. The largest fragment is the remaining vector backbone, in which a synthetic oligonucleotide linker is inserted to yield the vector pYD1lnk (SEQ ID No. 74). A cassette which includes the MAT α transcription termination region is then amplified by PCR from the vector pYD1 and is cloned into pYD1lnk via BamHI and PstI restriction and ligation. The resulting vector is pYD1mata (SEQ ID No.75). A cassette that contains the GAL1 promotor, the gene coding for Aga2 and a synthetic linker with NotI and SfiI cloning sites is amplified by PCR from pYD1 and cloned in pYD1mata via EcoRI and PacI restriction to yield the vector pYD1gal (SEQ.ID No.76).

As an example for a Fab to be displayed on yeast, the genes coding for VH-CH1 and VL-CL respectively of the antibody 4D5 (Herceptin) are made synthetically (sequences 4D5H (SEQ ID No.77) and 4D5L (SEQ ID No.78)).

4D5H is flanked by SfiI and NotI restriction sites, and cloned into the vector pYD1gal to yield the vector pYD4D5hc (SEQ ID No.79). In this vector, the N-terminus of 4D5H is fused to the C-terminus of Aga2, and at the C-terminus of 4D5H, a hexahistidine tag is attached, followed by the stop codon. The amino acid sequence of VH-CH1 of 4D5 is given in 4D5hp (SEQ ID No.80).

4D5L is flanked by NcoI and AscI restriction sites, and cloned into the vector pYD4D5hc to yield the vector pYD4D5hl (SEQ ID No.81). 4D5L is preceded by an Aga2 secretion signal, and carries a stop codon after the C-terminal Cysteine residue of the CL domain. The amino acid sequence of VL-CL of 4D5 is given in 4D5lp (SEQ ID No.82).

For display of the 4D5 Fab, the vector pYD4D5hl is transformed into the yeast strain EBY100 (Invitrogen), transformants are selected on minimal medium without tryptophan, and expression of the recombinant protein is induced by growth on galactose containing medium according to standard protocols (Invitrogen).

Example 8: Construction of a library with randomized residues in structural loops of the CL domain of 4D5 Fab

As first step in the yeast display library construction, the wildtype CL (C kappa) domain is cut out from the display vector pYD4D5hl with restriction enzymes BsiWI and AscI. A synthetic gene encoding human C kappa domain flanked by BsiWI and AscI sites (in the context according to pYD4D5hl) is prepared in which random mutations and insertions respectively are introduced in the AB and EF loops. In this particular example, insertions of 3, 4 or 5 NNB codons are made between amino acid positions 16 and 17 of the human C kappa domain, and residue positions 92, 93, 94, 95, 97, 98 and 99 are replaced by NNB codons. (IMGT numbering, see Figure 2). An NNB codon contains all 4 nucleotides at positions 1 and 2, and C, G and T at position 3. NNB therefore encodes all 20 naturally encoded amino acids.

The library is prepared and selected following standard procedures.

As a scaffold ligand the CDR target Her2neu and 4D5 epitope is used. Those members of the library are selected for production of a cytotoxic modular antibody according to the invention, that have a binding site engineered into the CL domain, which is specifically binding to an effector molecule, such as an Fc γ receptor, or a half-life prolonging protein, such as serum albumin. The resulting Fab is tested for (i) Her2neu binding with a $K_d < 10^{-8}$ M and an $IC_{50} < 10^{-8}$ M, and (ii) effector function using a CDC and/or ADCC assay, and alternatively to determine albumin binding.

Claims:

1. Method of preparing a genetic package displaying oligomers of modular antibody domains binding to a target and to a scaffold ligand comprising
 - providing a genetic package, and
 - displaying at least two of the antibody modular domains by fusing to the outer surface of the package.
2. Method according to claim 1 wherein the genetic package is displayed in a mobilized or cellular system.
3. Method according to any one of claims 1 or 2 wherein the mobilized system is selected from the group consisting of viruses, phages, phagemids, in-vitro display systems, mRNA systems and ribosomal display systems.
4. Method according to any one of claims 1 to 3 wherein said genetic package is a filamentous bacteriophage and said oligomer is a fusion protein comprising one of the proteins p3, p6 or p8.
5. Method according to any one of claims 1 or 2 wherein the cellular system is selected from the group consisting of yeast, mammalian cells, bacterial cells, bacterial spores and insect cells.
6. Method according to any one of claims 1 to 2 and 5, wherein said genetic package is yeast and said oligomer is a fusion protein of one of the proteins of yeast cell surface receptors selected from the group consisting of comprising one of the proteins alpha-agglutinin, a-agglutinin, Aga1p, Aga2p or FLO1.
7. Method according to any one of claims 1 to 6, wherein said oligomers comprise at least one leucine zipper, disulfide bond, electrostatic or hydrophobic motif.
8. Method according to any one of claims 1 to 7, wherein said oligomers are selected from the group consisting of dimers, trimers and tetramers.

9. Method according to any one of claims 1 to 8, wherein said oligomers are homomers.
10. Method according to any one of claims 1 to 9, wherein said oligomer is a polypeptide with a target binding site that directs towards the surface of the genetic package and is close to said surface.
11. Method according to any one of claims 1 to 10, wherein said oligomer is a polypeptide with a target binding site at a C-terminal loop position.
12. Method according to any one of claims 1 to 11, wherein said oligomer is a polypeptide with a target closer to the C-terminus than to the N-terminus of the polypeptide.
13. Method according to any one of claims 1 to 12, wherein said oligomer has at least two target binding sites.
14. Method according to any one of claims 1 to 13, wherein said modular antibody is selected from the group consisting of an antibody, a Fc fragment, an antibody fragment with a CDR region and combinations thereof.
15. Method according to claim 14 wherein the modular antibody comprises a fragment with a binding site at a structural loop position.
16. Method according to any one of claims 14 or 15, wherein said antibody fragment with a CDR region is selected from the group consisting of Fab, dAb, scFv, diabody, unibody, SMIPs, TANDABS, Fc fusion proteins and combinations thereof.
17. Cassette vector for use in a method according to any of claims 1 to 16, containing sequences encoding more than one fusion proteins operatively linked to the genetic package.
18. Vector according to claim 17, which is a phagemid.
19. Expression system for expressing oligomers of modular antibody domains bound at the surface of a genetic package produced according to any one of claims 1 to 16, wherein the oligomers are encoded by a single gene and the fusion protein is displayed with at least two copies on the surface of the genetic package.

20. Method of expressing a genetic package displaying oligomers of modular antibody domains produced according to any one of claims 1 to 16 in a non-suppressor host cell.
21. Genetic package as prepared by the method according to any one of claims 1 to 16.
22. Genetic package according to claim 21 displaying at least two fusion proteins.
23. Genetic package according to any one of claims 21 and 22 wherein each monomer of said oligomer is bound to the outer surface of said genetic package.
24. Bivalent phage displaying two fusion proteins from oligomers produced according to any one of claims 1 to 16, each containing an oligomer that dimerizes upon expression.
25. Library of genetic packages according to any one of claims 21 to 24.
26. Library according to claim 25, comprising at least 10 variant genetic packages, wherein said variant genetic packages display heteromers of modular antibody domains.
27. Library according to claim 26 wherein the heteromer is based on a scaffold comprising a target binding site.
28. Library according to any one of claims 26 or 27 wherein said scaffold is a parent Fab and wherein at least 20% of the parent Fab variants are binding to the CDR-target of said parent Fab.
29. Library according to any one of claims 25 to 28 comprising variants of said oligomer with differences in the amino acid sequence.
30. Library according to any one of claim 25 to 29, wherein the amino acid sequences have been modified by at least one insertion or substitution to introduce at least one foreign amino acid, and a deletion.
31. Library according to claim 30, wherein said foreign amino acid is introduced by a randomization technique.
32. Library according to claim 30 or 31, wherein said foreign amino acid is selected from a specific group of amino acids

to obtain a library enriched with specific amino acids at the randomized positions.

33. Method of producing an oligomer of modular antibody domains binding to a target comprising the steps of:
- a. providing a library of oligomers of modular antibody domains produced according to any one of claims 1 to 16
 - b. contacting said library with said target in the presence of a scaffold ligand,
 - c. selecting a library member binding to said target in the presence of a scaffold ligand, and
 - d. manufacturing a preparation of the functional oligomer.
34. Method according to claim 33, wherein said scaffold ligand is selected from the group consisting of an effector molecule, FcRn, Protein A and CDR target.
35. Method according to any one of claims 33 or 34, wherein the effector molecule is selected from the group consisting of CD64, CD16, CD32 and Fc receptors.
36. Method according to any one of claims 33 to 35, wherein said oligomers are dimers selected from the group of VH/VL, CH1/CL, CH2/CH2, CH3/CH3, Fc and Fab, or single chains thereof
37. Method according to any one of claims 1 to 16 and 33 to 36, wherein said library contains at least 10^2 independent clones expressing functional oligomers of modular antibody domains.
38. Method according to any one of claims 1 to 16 and 33 to 37, wherein a library member is selected that has a target binding affinity of $K_d < 10^{-8} M$.
39. Method according to any one of claims 1 to 16 and 33 to 38 wherein said target is a receptor of the erbB class.
40. Immunoglobulin obtained by a method according to any one of claims 1 to 16 and 33 to 39, wherein said target is a receptor of the erbB class.

41. High quality library containing at least 10^2 independent clones of functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand.
42. High quality library according to claim 41 wherein the target is a ligand binding to a parent molecule subject to amino acid variation.
43. High quality library according to any one of claims 41 or 42 wherein the parent molecule is a functional Fc or a functional Fab or part thereof.
44. High quality library according to any one of claims 41 or 42 wherein the parent molecules are varied by random or site specific mutagenesis
45. Library according to any one of claims 41 to 44, wherein at least 20% of the functional dimers are binding to CD64.
46. Library according to any one of claims 41 to 44, wherein at least 20% of the functional dimers are binding to protein A.
47. High quality library according to any one of claims 41 to 46 wherein at least 20% of the functional dimers are binding to the same CDR target.

Figure 1.





Figure 2



INTERNATIONAL SEARCH REPORT

International application No
PCT/AT2008/000232

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K16/00 C40B40/10 C40B40/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C40B C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/71694 A (SCRIPPS RESEARCH INST [US]) 30 November 2000 (2000-11-30)	1-3, 5, 7, 8, 10, 14, 16-18, 20-27, 29-34, 36, 37, 41, 42, 44
Y	the whole document	6, 9, 11-13, 15, 28, 35, 38-40, 43, 45-47

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

13 October 2008

Date of mailing of the international search report

10/11/2008

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Pérez-Mato, Isabel

INTERNATIONAL SEARCH REPORT

International application No

PCT/AT2008/000232

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GAO CHANGSHOU ET AL: "Making artificial antibodies: A format for phage display of combinatorial heterodimeric arrays" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE, WASHINGTON, DC.; US, vol. 96, no. 11, 25 May 1999 (1999-05-25), pages 6025-6030, XP002245749 ISSN: 0027-8424 the whole document	1-3,5,7, 8,10,14, 16-18, 20-24
X	US 5 759 817 A (BARBAS CARLOS [US]) 2 June 1998 (1998-06-02) the whole document	1-5,7,8, 14, 17-27, 29-32, 41,42,44
X	US 6 057 098 A (BUECHLER JOE [US] ET AL) 2 May 2000 (2000-05-02) the whole document	1-5,7,8, 14, 17-22, 24-26, 29-32, 41,42
Y	WO 2006/072620 A (RUEKER FLORIAN [AT]; WOZNIAK-KNOPP GORDANA [AT]) 13 July 2006 (2006-07-13) the whole document	6,9, 11-13, 15,28, 35, 38-40, 43,45-47
X	WO 2006/033700 A (GENENTECH INC [US]; KAO YUNG-HSIANG [US]; VANDERLAAN MARTIN [US]) 30 March 2006 (2006-03-30) the whole document	40
Y		39
X	VAJDOS F F ET AL: "Comprehensive Functional Maps of the Antigen-binding Site of an Anti-ErbB2 Antibody Obtained with Shotgun Scanning Mutagenesis" JOURNAL OF MOLECULAR BIOLOGY, LONDON, GB, vol. 320, no. 2, 5 July 2002 (2002-07-05), pages 415-428, XP004449739 ISSN: 0022-2836	40
Y		39
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/AT2008/000232

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SPIRIDON CAMELIA I ET AL: "Targeting multiple Her-2 epitopes with monoclonal antibodies results in improved antigrowth activity of a human breast cancer cell line in vitro and in vivo" CLINICAL CANCER RESEARCH, THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, US, vol. 8, no. 6, 1 June 2002 (2002-06-01), pages 1720-1730, XP002389218 ISSN: 1078-0432	40
Y	-----	39
Y	US 2004/146976 A1 (WITTRUP K DANE [US] ET AL) 29 July 2004 (2004-07-29) the whole document	6
P,X	WO 2008/003116 A (STAR BIOTECHNOLOGISCHE FORSCHU [AT]; HIMMLER GOTTFRIED [AT]; REDL GERD) 10 January 2008 (2008-01-10) the whole document	1-47
T	RÜKER F.: "Modular Antibody Technology" F-STAR FACT SHEET, [Online] 7 February 2008 (2008-02-07), XP002499302 Retrieved from the Internet: URL: http://www.boku.ac.at/fileadmin/BOKU-Topstories/20080702_Rueker_Factsheet.pdf [retrieved on 2008-10-13] the whole document	-----

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/AT2008/000232

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0071694	A	30-11-2000	AT 317008 T 15-02-2006
			AU 782349 B2 21-07-2005
			AU 5290000 A 12-12-2000
			CA 2374505 A1 30-11-2000
			DE 60025824 T2 07-09-2006
			DK 1185636 T3 18-04-2006
			EP 1185636 A1 13-03-2002
			ES 2256015 T3 16-07-2006
			PT 1185636 T 31-05-2006
			US 2003186322 A1 02-10-2003
			US 6472147 B1 29-10-2002
US 5759817	A	02-06-1998	AT 269401 T 15-07-2004
			AU 662148 B2 24-08-1995
			AU 1785692 A 17-11-1992
			CA 2108147 A1 11-10-1992
			DE 69233367 D1 22-07-2004
			DE 69233367 T2 25-05-2005
			EP 0580737 A1 02-02-1994
			ES 2223040 T3 16-02-2005
			FI 934422 A 08-12-1993
			IE 921169 A1 21-10-1992
			JP 3672306 B2 20-07-2005
			JP 6506836 T 04-08-1994
			JP 4108430 B2 25-06-2008
			JP 2003204797 A 22-07-2003
			JP 4118896 B2 16-07-2008
			JP 2005218449 A 18-08-2005
			JP 4118938 B2 16-07-2008
			JP 2007222179 A 06-09-2007
NO 933610 A 10-12-1993			
PT 100379 A 31-08-1993			
WO 9218619 A1 29-10-1992			
US 6057098	A	02-05-2000	US 6986986 B1 17-01-2006
WO 2006072620	A	13-07-2006	AU 2006204459 A1 13-07-2006
			CA 2594356 A1 13-07-2006
			CN 101098891 A 02-01-2008
			EP 1699826 A1 13-09-2006
			JP 2008526809 T 24-07-2008
			KR 20070092242 A 12-09-2007
WO 2006033700	A	30-03-2006	AR 050353 A1 18-10-2006
			AU 2005287404 A1 30-03-2006
			BR PI0513681 A 13-05-2008
			CA 2574821 A1 30-03-2006
			CN 101023100 A 22-08-2007
			EP 1771482 A2 11-04-2007
			JP 2008507520 T 13-03-2008
			KR 20070038557 A 10-04-2007
			US 2004146976
WO 2008003116	A	10-01-2008	AT 503902 A1 15-01-2008