



(86) **Date de dépôt PCT/PCT Filing Date:** 2015/06/24
(87) **Date publication PCT/PCT Publication Date:** 2015/12/30
(85) **Entrée phase nationale/National Entry:** 2016/12/16
(86) **N° demande PCT/PCT Application No.:** US 2015/037392
(87) **N° publication PCT/PCT Publication No.:** 2015/200465
(30) **Priorité/Priority:** 2014/06/24 (US62/016,512)

(51) **Cl.Int./Int.Cl.** **C07D 403/06** (2006.01),
C07D 413/06 (2006.01)
(71) **Demandeur/Applicant:**
SHIRE HUMAN GENETIC THERAPIES, INC., US
(72) **Inventeurs/Inventors:**
DEROSA, FRANK, US;
KARVE, SHRIRANG, US;
HEARTLEIN, MICHAEL, US
(74) **Agent:** FASKEN MARTINEAU DUMOULIN LLP

(54) **Titre : COMPOSITIONS ENRICHIES STEREOCHIMIQUEMENT POUR L'ADMINISTRATION D'ACIDES NUCLEIQUES**
(54) **Title: STEREOCHEMICALLY ENRICHED COMPOSITIONS FOR DELIVERY OF NUCLEIC ACIDS**





FIG. 1



(57) Abrégé/Abstract:

Provided, in part, is a composition comprising one or more chemical entities of formula I, each of which is a compound of formula (I): a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof, the



(57) Abrégé(suite)/Abstract(continued):

composition characterized in that greater than a first threshold amount of the total amount of chemical entities of formula I in the composition: are chemical entities of formula I.a, wherein the first threshold amount is 50%; or are chemical entities of formula I.b.1, wherein the first threshold amount is 25%; or are chemical entities of formula I.b.2, wherein the first threshold amount is 25%, wherein the chemical entities of formula I.a, I.b.1, and I.b.2, are described herein, and methods of using such compositions, for example, for the delivery of a polynucleotide in vivo.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
30 December 2015 (30.12.2015)

WIPO | PCT

(10) International Publication Number
WO 2015/200465 A1

(51) International Patent Classification:

C07D 403/06 (2006.01) **C07D 413/06** (2006.01)

(21) International Application Number:

PCT/US2015/037392

(22) International Filing Date:

24 June 2015 (24.06.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/016,512 24 June 2014 (24.06.2014) US

(71) Applicant: **SHIRE HUMAN GENETIC THERAPIES, INC.** [US/US]; 300 Shire Way, Lexington, Massachusetts 02421 (US).(72) Inventors: **DEROSA, Frank**; 26 Mount Auburn St., Chelmsford, Massachusetts 01824 (US). **KARVE, Shrirang**; 4504 Stearns Hill Road, Waltham, Massachusetts 02451 (US). **HEARTLEIN, Michael**; 167 Reed Farm Road, Boxborough, Massachusetts 01719 (US).(74) Agent: **HENRY, Kevin, M.**; Two International Place, Boston, Massachusetts 02110 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

[Continued on next page]

(54) Title: STEREOCHEMICALLY ENRICHED COMPOSITIONS FOR DELIVERY OF NUCLEIC ACIDS





FIG. 1

(57) Abstract: Provided, in part, is a composition comprising one or more chemical entities of formula I, each of which is a compound of formula (I): a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof, the composition characterized in that greater than a first threshold amount of the total amount of chemical entities of formula I in the composition: are chemical entities of formula I.a, wherein the first threshold amount is 50%; or are chemical entities of formula I.b.1, wherein the first threshold amount is 25%; or are chemical entities of formula I.b.2, wherein the first threshold amount is 25%, wherein the chemical entities of formula I.a, I.b.1, and I.b.2, are described herein, and methods of using such compositions, for example, for the delivery of a polynucleotide *in vivo*.

WO 2015/200465 A1



— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))* — *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

Published:

— *with international search report (Art. 21(3))*


STEREOCHEMICALLY ENRICHED COMPOSITIONS FOR DELIVERY OF NUCLEIC ACIDS

BACKGROUND

[0001] Delivery of nucleic acids using lipids has been explored extensively for treatment of various diseases. There remains a great need for lipids and/or lipids compositions that can deliver nucleic acids, such as short interfering RNA (siRNA) and messenger RNA (mRNA) with high efficiency and low toxicity.

SUMMARY


[0002] Among other things, the present invention provides compositions comprising stereochemically enriched lipids for delivering mRNA. The invention is based, in part, on the surprising discovery that compositions comprising stereochemically enriched lipid of formula **I**, below, are highly effective and have unexpectedly low toxicity in delivering mRNA and producing encoded protein *in vivo*:



I

The present inventors found that when used for mRNA delivery, stereochemically enriched compositions of **I** have surprisingly low toxicity compared to stereochemically non-enriched, or stereochemically less enriched, compositions of the same lipid, as evidenced, for example, by the dramatically lower alanine aminotransferase (ALT) and aspartate aminotransferase (AST) expression levels. See Table 1.


[0003] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, or **I.h**, each of which is independently as defined and described below.


[0004] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**


a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition:


are chemical entities of formula **I.a**:

**I.a**

wherein the first threshold amount is 50%; or
are chemical entities of formula **I.b.1**:


**I.b.1**

wherein the first threshold amount is 25%; or
are chemical entities of formula **I.b.2**:

**I.b.2**

wherein the first threshold amount is 25%.


[0005] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,


the composition characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition:

are chemical entities of formula **I.c**:

**I.c**


wherein the first threshold amount is 6.25%; or

are chemical entities of formula **I.d**:


**I.d**

wherein the first threshold amount is 6.25%; or


are chemical entities of formula **I.e**:

**I.e**


wherein the first threshold amount is 25%; or
are chemical entities of formula **I.f**:

**I.f**

wherein the first threshold amount is 25%; or
are chemical entities of formula **I.g**:

**I.g**


wherein the first threshold amount is 12.5%; or
are chemical entities of formula **I.h**:

**I.h**

wherein the first threshold amount is 25%.

[0006] In some embodiments, the first threshold amount is 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%. In some embodiments, the first threshold amount is 50%. In some embodiments, the first threshold amount is 70%. In some embodiments, the first threshold amount is 80%. In some embodiments, the first threshold amount is 95%.

[0007] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**


a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of a first formula selected from formulae **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, or **I.h**; and

greater than or equal to a second threshold amount of the total amount of the chemical entities of the first formula in the composition are chemical entities of the same stereoisomer of the first formula.


[0008] In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**.

[0009] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition is a chemical entity of formula **I.a.i**:



I.a.i (*i.e.*, *R4-SR-cKK-E12*)


[0010] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition is a chemical entity of formula **I.a.ii**:




I.a.ii (*i.e.*, *S4-SR-cKK-E12*)

[0011] In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**.

[0012] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition is a chemical entity of formula **I.b.1.i**:


**I.b.1.i** (*i.e.*, *R4-SS-cKK-E12*)

[0013] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition is a chemical entity of formula **I.b.1.ii**:


**I.b.1.ii** (*i.e.*, *S4-SS-cKK-E12*)

[0014] In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**.

[0015] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition is a chemical entity of formula **I.b.2.i**:

**I.b.2.i** (*i.e.*, *R4-RR-cKK-E12*)


[0016] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition is a chemical entity of formula **I.b.2.ii**:



I.b.2.ii (*i.e.*, *S4-RR-cKK-E12*)

[0017] In some embodiments, the second threshold amount is 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%. In some embodiments, the second threshold amount is 50%. In some embodiments, the second threshold amount is 70%. In some embodiments, the second threshold amount is 80%. In some embodiments, the second threshold amount is 95%.

[0018] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:




I

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I**.

[0019] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of the same stereoisomer of formula **I**.

[0020] In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.i**, **I.a.ii**, **I.b.1.i**, **I.b.1.ii**, **I.b.2.i**, or **I.b.2.ii**.

[0021] In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.a.i**, **I.a.ii**, **I.b.1.i**, **I.b.1.ii**, **I.b.2.i**, or **I.b.2.ii**. In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.a.i**. In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.a.ii**. In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.b.1.i**. In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.b.1.ii**. In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.b.2.i**. In some embodiments, greater than or equal to the third threshold amount of the total amount of the composition is a chemical entity of formula **I.b.2.ii**.

[0022] In some embodiments, the third threshold amount is 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% (w/w). In some embodiments, the third threshold amount is 50% (w/w). In some embodiments, the third threshold amount is 70% (w/w). In some embodiments, the third threshold amount is 80% (w/w). In some embodiments,

the third threshold amount is 85% (w/w). In some embodiments, the third threshold amount is 95% (w/w).

[0023] In some embodiments, a provided composition further comprises one or more mRNA for mRNA delivery and expression of the encoded protein *in vivo*.

[0024] In some embodiments, the present invention provides methods for highly efficient delivery and expression of mRNA and encoded protein *in vivo*. In some embodiments, the present invention provides a method of delivery of mRNA *in vivo*, comprising administering to a subject in need of delivery a provided composition which comprises an mRNA. In some embodiments, due to their low toxicity, a provide composition permits more delivered mRNA, higher protein expression level and/or lower administration frequency, thereby providing a more potent, safer, and more patent-friendly mRNA therapy.

BRIEF DESCRIPTION OF THE DRAWING

[0025] FIG. 1 depicts the results of *in vivo* human ASS1 protein production in wild type mouse liver upon treatment with lipid nanoparticles that include compounds of formula I.

DEFINITIONS

[0026] In order for the present invention to be more readily understood, certain terms are first defined below. Additional definitions for the following terms and other terms are set forth throughout the specification. The publications and other reference materials referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference.

[0027] *Amino acid*: As used herein, term “amino acid,” in its broadest sense, refers to any compound and/or substance that can be incorporated into a polypeptide chain. In some embodiments, an amino acid has the general structure $H_E N-C(H)(R)-COHO$. In some embodiments, an amino acid is a naturally occurring amino acid. In some embodiments, an amino acid is a synthetic amino acid; in some embodiments, an amino acid is a d-amino acid; in some embodiments, an amino acid is an l-amino acid. “Standard amino acid” refers to any of the twenty standard l-amino acids commonly found in naturally occurring peptides. “Nonstandard amino acid” refers to any amino acid, other than the standard amino acids, regardless of whether it is prepared synthetically or obtained from a natural source. As used herein, “synthetic amino acid” encompasses chemically modified amino acids, including but not limited to salts, amino

acid derivatives (such as amides), and/or substitutions. Amino acids, including carboxyl- and/or amino-terminal amino acids in peptides, can be modified by methylation, amidation, acetylation, protecting groups, and/or substitution with other chemical groups that can change the peptide's circulating half-life without adversely affecting their activity. Amino acids may participate in a disulfide bond. Amino acids may comprise one or posttranslational modifications, such as association with one or more chemical entities (*e.g.*, methyl groups, acetate groups, acetyl groups, phosphate groups, formyl moieties, isoprenoid groups, sulfate groups, polyethylene glycol moieties, lipid moieties, carbohydrate moieties, biotin moieties, *etc.*). The term “amino acid” is used interchangeably with “amino acid residue,” and may refer to a free amino acid and/or to an amino acid residue of a peptide. It will be apparent from the context in which the term is used whether it refers to a free amino acid or a residue of a peptide.

[0028] *Animal*: As used herein, the term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans, at any stage of development. In some embodiments, “animal” refers to non-human animals, at any stage of development. In certain embodiments, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, and/or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, insects, and/or worms. In some embodiments, an animal may be a transgenic animal, genetically-engineered animal, and/or a clone.

[0029] *Approximately or about*: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0030] *Chemical entity*: As used herein, the term “chemical entity” includes a compound, salt, or solvate thereof, or any combination of compounds, salts, or solvates thereof.

[0031] *Delivery*: As used herein, the term “delivery” encompasses both local and systemic delivery. For example, delivery of mRNA encompasses situations in which an mRNA is delivered to a target tissue and the encoded protein is expressed and retained within the target

tissue (also referred to as “local distribution” or “local delivery”), and situations in which an mRNA is delivered to a target tissue and the encoded protein is expressed and secreted into patient’s circulation system (e.g., serum) and systematically distributed and taken up by other tissues (also referred to as “systemic distribution” or “systemic delivery”).

[0032] *Expression:* As used herein, “expression” of a nucleic acid sequence refers to translation of an mRNA into a polypeptide, assemble multiple polypeptides (e.g., heavy chain or light chain of antibody) into an intact protein (e.g., antibody) and/or post-translational modification of a polypeptide or fully assembled protein (e.g., antibody). In this application, the terms “expression” and “production,” and grammatical equivalent, are used inter-changeably.

[0033] *Functional:* As used herein, a “functional” biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized.

[0034] *Half-life:* As used herein, the term “half-life” is the time required for a quantity such as nucleic acid or protein concentration or activity to fall to half of its value as measured at the beginning of a time period.

[0035] *Improve, increase, or reduce:* As used herein, the terms “improve,” “increase” or “reduce,” or grammatical equivalents, indicate values that are relative to a baseline measurement, such as a measurement in the same individual prior to initiation of the treatment described herein, or a measurement in a control subject (or multiple control subject) in the absence of the treatment described herein. A “control subject” is a subject afflicted with the same form of disease as the subject being treated, who is about the same age as the subject being treated.

[0036] *In Vitro:* As used herein, the term “*in vitro*” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, *etc.*, rather than within a multi-cellular organism.

[0037] *In Vivo:* As used herein, the term “*in vivo*” refers to events that occur within a multi-cellular organism, such as a human and a non-human animal. In the context of cell-based systems, the term may be used to refer to events that occur within a living cell (as opposed to, for example, *in vitro* systems).

[0038] *Isolated:* As used herein, the term “isolated” refers to a substance and/or entity that has been (1) separated from at least some of the components with which it was associated when initially produced (whether in nature and/or in an experimental setting), and/or (2) produced, prepared, and/or manufactured by the hand of man. Isolated substances and/or entities may be

separated from about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% of the other components with which they were initially associated. In some embodiments, isolated agents are about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components. As used herein, calculation of percent purity of isolated substances and/or entities should not include excipients (*e.g.*, buffer, solvent, water, *etc.*).

[0039] *Local distribution or delivery:* As used herein, the terms “local distribution,” “local delivery,” or grammatical equivalent, refer to tissue specific delivery or distribution. Typically, local distribution or delivery requires a protein (*e.g.*, enzyme) encoded by mRNAs be translated and expressed intracellularly or with limited secretion that avoids entering the patient’s circulation system.

[0040] *messenger RNA (mRNA):* As used herein, the term “messenger RNA (mRNA)” refers to a polynucleotide that encodes at least one polypeptide. mRNA as used herein encompasses both modified and unmodified RNA. mRNA may contain one or more coding and non-coding regions. mRNA can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, *etc.* Where appropriate, *e.g.*, in the case of chemically synthesized molecules, mRNA can comprise nucleoside analogs such as analogs having chemically modified bases or sugars, backbone modifications, *etc.* An mRNA sequence is presented in the 5’ to 3’ direction unless otherwise indicated. In some embodiments, an mRNA is or comprises natural nucleosides (*e.g.*, adenosine, guanosine, cytidine, uridine); nucleoside analogs (*e.g.*, 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, C-5 propynyl-cytidine, C-5 propynyl-uridine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (*e.g.*, methylated bases); intercalated bases; modified sugars (*e.g.*, 2’-fluororibose, ribose, 2’-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (*e.g.*, phosphorothioates and 5’-*N*-phosphoramidite linkages).

[0041] In some embodiments, the mRNA comprises one or more nonstandard nucleotide residues. The nonstandard nucleotide residues may include, e.g., 5-methyl-cytidine (“5mC”), pseudouridine, and/or 2-thio-uridine (“2sU”). See, e.g., U.S. Patent No. 8,278,036 or WO2011012316 for a discussion of such residues and their incorporation into mRNA. The mRNA may be RNA, which is defined as RNA in which 25% of U residues are 2-thio-uridine and 25% of C residues are 5-methylcytidine. Teachings for the use of RNA are disclosed US Patent Publication US20120195936 and international publication WO2011012316, both of which are hereby incorporated by reference in their entirety. The presence of nonstandard nucleotide residues may render an mRNA more stable and/or less immunogenic than a control mRNA with the same sequence but containing only standard residues. In further embodiments, the mRNA may comprise one or more nonstandard nucleotide residues chosen from isocytosine, pseudoisocytosine, 5-bromouracil, 5-propynyluracil, 6-aminopurine, 2-aminopurine, inosine, diaminopurine and 2-chloro-6-aminopurine cytosine, as well as combinations of these modifications and other nucleobase modifications. Certain embodiments may further include additional modifications to the furanose ring or nucleobase. Additional modifications may include, for example, sugar modifications or substitutions (e.g., one or more of a 2'-O-alkyl modification, a locked nucleic acid (LNA)). In some embodiments, the RNAs may be complexed or hybridized with additional polynucleotides and/or peptide polynucleotides (PNA). In embodiments where the sugar modification is a 2'-O-alkyl modification, such modification may include, but are not limited to a 2'-deoxy-2'-fluoro modification, a 2'-O-methyl modification, a 2'-O-methoxyethyl modification and a 2'-deoxy modification. In certain embodiments, any of these modifications may be present in 0-100% of the nucleotides—for example, more than 0%, 1%, 10%, 25%, 50%, 75%, 85%, 90%, 95%, or 100% of the constituent nucleotides individually or in combination.

[0042] *Nucleic acid:* As used herein, the term “nucleic acid,” in its broadest sense, refers to any compound and/or substance that is or can be incorporated into a polynucleotide chain. In some embodiments, a nucleic acid is a compound and/or substance that is or can be incorporated into a polynucleotide chain via a phosphodiester linkage. In some embodiments, “nucleic acid” refers to individual nucleic acid residues (e.g., nucleotides and/or nucleosides). In some embodiments, “nucleic acid” refers to a polynucleotide chain comprising individual nucleic acid

residues. In some embodiments, “nucleic acid” encompasses RNA as well as single and/or double-stranded DNA and/or cDNA.

[0043] *Patient:* As used herein, the term “patient” or “subject” refers to any organism to which a provided composition may be administered, *e.g.*, for experimental, diagnostic, prophylactic, cosmetic, and/or therapeutic purposes. Typical patients include animals (*e.g.*, mammals such as mice, rats, rabbits, non-human primates, and/or humans). In some embodiments, a patient is a human. A human includes pre and post natal forms.

[0044] *Pharmaceutically acceptable:* The term “pharmaceutically acceptable” as used herein, refers to substances that, within the scope of sound medical judgment, are suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0045] *Polymer:* As used herein, a “polymer” refers to a compound comprised of at least 3 (*e.g.*, at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, etc.) repeating covalently bound structural units.

[0046] *Salt:* As used herein, the term “salt” or “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describes pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences* (1977) 66:1-19. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide,

2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2- naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C1-4alkyl)_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, sulfonate and aryl sulfonate. Further pharmaceutically acceptable salts include salts formed from the quaternization of an amine using an appropriate electrophile, e.g., an alkyl halide, to form a quaternized alkylated amino salt.

[0047] *Systemic distribution or delivery:* As used herein, the terms “systemic distribution,” “systemic delivery,” or grammatical equivalent, refer to a delivery or distribution mechanism or approach that affect the entire body or an entire organism. Typically, systemic distribution or delivery is accomplished via body’s circulation system, e.g., blood stream. Compared to the definition of “local distribution or delivery.”

[0048] *Subject:* As used herein, the term “subject” refers to a human or any non-human animal (e.g., mouse, rat, rabbit, dog, cat, cattle, swine, sheep, horse or primate). A human includes pre- and post-natal forms. In many embodiments, a subject is a human being. A subject can be a patient, which refers to a human presenting to a medical provider for diagnosis or treatment of a disease. The term “subject” is used herein interchangeably with “individual” or “patient.” A subject can be afflicted with or is susceptible to a disease or disorder but may or may not display symptoms of the disease or disorder.

[0049] *Substantially:* As used herein, the term “substantially” refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term “substantially” is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical phenomena.

[0050] *Target tissues:* As used herein, the term “target tissues” refers to any tissue that is affected by a disease to be treated. In some embodiments, target tissues include those tissues that display disease-associated pathology, symptom, or feature.

[0051] *Therapeutically effective amount:* As used herein, the term “therapeutically effective amount” of a therapeutic agent means an amount that is sufficient, when administered to a subject suffering from or susceptible to a disease, disorder, and/or condition, to treat, diagnose, prevent, and/or delay the onset of the symptom(s) of the disease, disorder, and/or condition. It will be appreciated by those of ordinary skill in the art that a therapeutically effective amount is typically administered via a dosing regimen comprising at least one unit dose.

[0052] *Treating:* As used herein, the term “treat,” “treatment,” or “treating” refers to any method used to partially or completely alleviate, ameliorate, relieve, inhibit, prevent, delay onset of, reduce severity of and/or reduce incidence of one or more symptoms or features of a particular disease, disorder, and/or condition. Treatment may be administered to a subject who does not exhibit signs of a disease and/or exhibits only early signs of the disease for the purpose of decreasing the risk of developing pathology associated with the disease.

DETAILED DESCRIPTION

[0053] The present invention provides, among other things, lipid compositions and methods for delivering mRNA *in vivo* using stereochemically enriched lipid compositions.

Lipid Compositions

[0054] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**, a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, or **I.h**.

[0055] In some embodiments, a provided composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, or **I.h**. In some

embodiments, a provided composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, or **I.h**.

[0056] As used herein, a “chemical entity” of a formula is a compound of the formula, a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof.

[0057] In some embodiments, the first threshold amount is 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%. In some embodiments, the first threshold amount is 50%. In some embodiments, the first threshold amount is 70%. In some embodiments, the first threshold amount is 80%. In some embodiments, the first threshold amount is 95%.

[0058] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0059] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**. In some embodiments, a first threshold amount is 25%. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first

threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0060] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**. In some embodiments, a first threshold amount is 25%. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0061] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.c**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.c**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.c**. In some embodiments, a first threshold amount is 6.25%. In some embodiments, a first threshold amount is 12.5%. In some embodiments, a first threshold amount is 25%. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0062] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.d**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.d**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical

entities of formula **I** in the composition are chemical entities of formula **I.d**. In some embodiments, a first threshold amount is 6.25%. In some embodiments, a first threshold amount is 12.5%. In some embodiments, a first threshold amount is 25%. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0063] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.e**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.e**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.e**. In some embodiments, a first threshold amount is 25%. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0064] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.f**. In some embodiments, the composition is characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.f**. In some embodiments, the composition is characterized in that a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.f**. In some embodiments, a first threshold amount is 25%. In some embodiments, a first threshold amount is 50%. In some embodiments, a first threshold amount is 70%. In some embodiments, a first threshold amount is 80%. In some embodiments, a first threshold amount is 90%. In some embodiments, a first threshold amount is 95%.

[0065] In some embodiments, the composition is characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.g**. In some embodiments, the composition is

[0067] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula I, each of which is a compound of formula I:




22

the composition characterized in that greater than or equal to a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of a first formula selected from formulae **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, and **I.h**; and greater than or equal to a second threshold amount of the total amount of the chemical entities of the first formula in the composition are chemical entities of the same stereoisomer of the first formula.

[0068] In some embodiments, greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of a first formula selected from formulae **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, and **I.h**. In some embodiments, a first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of a first formula selected from formulae **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, and **I.h**. In some embodiments, greater than a second threshold amount of the total amount of the chemical entities of the first formula in the composition are chemical entities of the same stereoisomer of the first formula. In some embodiments, a second threshold amount of the total amount of the chemical entities of the first formula in the composition are chemical entities of the same stereoisomer of the first formula.


[0069] In some embodiments, a first formula is formula **I.a**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.a** in the composition are chemical entities of the same stereoisomer of formula **I.a**.

[0070] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.i**:




I.a.i (i.e., *R4-SR-cKK-E12*)

[0071] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.ii**:




I.a.ii (*i.e.*, *S4-SR-cKK-E12*)

[0072] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.iii**:




I.a.iii

[0073] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.iv**:




I.a.iv


[0074] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.v**:

**I.a.v**


[0075] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.vi**:

**I.a.vi**


[0076] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.vii**:

**I.a.vii**


[0077] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.viii**:

**I.a.viii**

[0078] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.ix**:

**I.a.ix**

[0079] In some embodiments, a stereoisomer of formula **I.a** has the structure of formula **I.a.x**:

**I.a.x**

[0080] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula **I.a.i**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula **I.a.ii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula

I.a.iii. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula

I.a.iv. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula

I.a.v. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula

I.a.vi. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula

I.a.vii. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula

I.a.viii. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition are chemical entities of formula


I.a.ix.

[0081] In some embodiments, a first formula is formula **I.b.1**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.b.1** in the composition are chemical entities of the same stereoisomer of formula **I.b.1**.

[0082] In some embodiments, a stereoisomer of formula **I.b.1** has the structure of formula **I.b.1.i** (*i.e.*, *R4-SS-cKK-E12*).


[0083] In some embodiments, a stereoisomer of formula **I.b.1** has the structure of formula **I.b.1.ii** (*i.e.*, *S4-SS-cKK-E12*).

[0084] In some embodiments, a stereoisomer of formula **I.b.1** has the structure of formula **I.b.1.iii**:




I.b.1.iii

[0085] In some embodiments, a stereoisomer of formula **I.b.1** has the structure of formula **I.b.1.iv**:




I.b.1.iv

[0086] In some embodiments, a stereoisomer of formula **I.b.1** has the structure of formula **I.b.1.v**:



I.b.1.v

[0087] In some embodiments, a stereoisomer of formula **I.b.1** has the structure of formula **I.b.1.vi**:



I.b.1.vi

[0088] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition are chemical entities of formula **I.b.1.i**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition are chemical entities of formula

I.b.1.ii. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition are chemical entities of formula

I.b.1.iii. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition are chemical entities of formula

I.b.1.iv. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition are chemical entities of formula

I.b.1.v. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition are chemical entities of formula


I.b.1.vi.

[0089] In some embodiments, a first formula is formula **I.b.2**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.b.2** in the composition are chemical entities of the same stereoisomer of formula **I.b.2**.

[0090] In some embodiments, a stereoisomer of formula **I.b.2** has the structure of formula **I.b.2.i** (*i.e.*, *R4-RR-cKK-E12*).


[0091] In some embodiments, a stereoisomer of formula **I.b.2** has the structure of formula **I.b.2.ii** (*i.e.*, *S4-RR-cKK-E12*).

[0092] In some embodiments, a stereoisomer of formula **I.b.2** has the structure of formula **I.b.2.iii**:




I.b.2.iii


[0093] In some embodiments, a stereoisomer of formula **I.b.2** has the structure of formula **I.b.2.iv**:

**I.b.2.iv**

[0094] In some embodiments, a stereoisomer of formula **I.b.2** has the structure of formula **I.b.2.v**:

**I.b.2.v**

[0095] In some embodiments, a stereoisomer of formula **I.b.2** has the structure of formula **I.b.2.vi**:

**I.b.2.vi**

[0096] In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition are chemical entities of formula **I.b.2.i**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition are chemical entities of formula **I.b.2.ii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition are chemical entities of formula

I.b.2.iii. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition are chemical entities of formula

I.b.2.iv. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition are chemical entities of formula

I.b.2.v. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition are chemical entities of formula

I.b.2.vi.

[0097] In some embodiments, a first formula is formula **I.c**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.c**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.c** in the composition are chemical entities of the same stereoisomer of formula **I.c**. In some embodiments, a stereoisomer of formula **I.c** has the structure of formula **I.a.i**. In some embodiments, a stereoisomer of formula **I.c** has the structure of formula **I.b.1.i**. In some embodiments, a stereoisomer of formula **I.c** has the structure of formula **I.b.2.i**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.c** in the composition are chemical entities of formula **I.a.i**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.c** in the composition are chemical entities of formula **I.b.1.i**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.c** in the composition are chemical entities of formula **I.b.2.i**.

[0098] In some embodiments, a first formula is formula **I.d**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.d**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.d** in the composition are chemical entities of the same stereoisomer of formula **I.d**. In some embodiments, a stereoisomer of formula **I.d** has the structure of formula **I.a.ii**. In some embodiments, a stereoisomer of formula **I.d** has the structure of formula **I.b.1.ii**. In some embodiments, a stereoisomer of formula **I.d** has the structure of formula **I.b.2.ii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.d** in the composition are chemical entities of formula **I.a.ii**. In some embodiments, greater than

or equal to a second threshold amount of the total amount of chemical entities of formula **I.d** in the composition are chemical entities of formula **I.b.1.ii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.d** in the composition are chemical entities of formula **I.b.2.ii**.

[0099] In some embodiments, a first formula is formula **I.e**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.e**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.e** in the composition are chemical entities of the same stereoisomer of formula **I.e**. In some embodiments, a stereoisomer of formula **I.e** has the structure of formula **I.a.iii**. In some embodiments, a stereoisomer of formula **I.e** has the structure of formula **I.a.v**. In some embodiments, a stereoisomer of formula **I.e** has the structure of formula **I.b.1.iii**. In some embodiments, a stereoisomer of formula **I.e** has the structure of formula **I.b.2.iii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.e** in the composition are chemical entities of formula **I.a.iii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.e** in the composition are chemical entities of formula **I.a.v**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.e** in the composition are chemical entities of formula **I.b.1.iii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.e** in the composition are chemical entities of formula **I.b.2.iii**.

[00100] In some embodiments, a first formula is formula **I.f**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.f**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.f** in the composition are chemical entities of the same stereoisomer of formula **I.f**. In some embodiments, a stereoisomer of formula **I.f** has the structure of formula **I.a.vii**. In some embodiments, a stereoisomer of formula **I.f** has the structure of formula **I.a.ix**. In some embodiments, a stereoisomer of formula **I.f** has the structure of formula **I.b.1.vi**. In some embodiments, a stereoisomer of formula **I.f** has the structure of formula **I.b.2.vi**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.f**

in the composition are chemical entities of formula **I.a.iii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.f** in the composition are chemical entities of formula **I.a.ix**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.f** in the composition are chemical entities of formula **I.b.1.vi**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.f** in the composition are chemical entities of formula **I.b.2.vi**.


[00101] In some embodiments, a first formula is formula **I.g**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.g**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.g** in the composition are chemical entities of the same stereoisomer of formula **I.g**. In some embodiments, a stereoisomer of formula **I.g** has the structure of formula **I.a.iv**. In some embodiments, a stereoisomer of formula **I.g** has the structure of formula **I.a.viii**. In some embodiments, a stereoisomer of formula **I.g** has the structure of formula **I.b.1.iv**. In some embodiments, a stereoisomer of formula **I.g** has the structure of formula **I.b.2.iv**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.g** in the composition are chemical entities of formula **I.a.iv**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.g** in the composition are chemical entities of formula **I.a.viii**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.g** in the composition are chemical entities of formula **I.b.1.iv**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.g** in the composition are chemical entities of formula **I.b.2.iv**.

[00102] In some embodiments, a first formula is formula **I.h**. In some embodiments, greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.h**, and greater than or equal to a second threshold amount of the total amount of the chemical entities of formula **I.h** in the composition are chemical entities of the same stereoisomer of formula **I.h**. In some embodiments, a stereoisomer of formula **I.h** has the structure of formula **I.a.vi**. In some embodiments, a stereoisomer of formula **I.h** has the structure of formula **I.b.1.v**. In some embodiments, a

stereoisomer of formula **I.h** has the structure of formula **I.b.2.v**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.h** in the composition are chemical entities of formula **I.a.vi**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.h** in the composition are chemical entities of formula **I.b.1.v**. In some embodiments, greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.h** in the composition are chemical entities of formula **I.b.2.v**.

[00103] In some embodiments, the second threshold amount is 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%. In some embodiments, the second threshold amount is 50%. In some embodiments, the second threshold amount is 70%. In some embodiments, the second threshold amount is 80%. In some embodiments, the second threshold amount is 85%. In some embodiments, the second threshold amount is 90%. In some embodiments, the second threshold amount is 95%.

[00104] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:



I

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I**.

[00105] In some embodiments, a provided composition is characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I**. In some embodiments, a provided composition is


characterized in that a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I**.

[00106] In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**, **I.b.1**, **I.b.2**, **I.c**, **I.d**, **I.e**, **I.f**, **I.g**, or **I.h**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.c**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.d**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.e**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.f**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.g**. In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.h**.

[00107] In some embodiments, a provided composition is characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a.i**, **I.a.ii**, **I.a.iii**, **I.a.iv**, **I.a.v**, **I.a.vi**, **I.a.vii**,

I.a.viii, I.a.ix, I.b.1.i, I.b.1.ii, I.b.1.iii, I.b.1.iv, I.b.1.v, I.b.1.vi, I.b.2.i, I.b.2.ii, I.b.2.iii, I.b.2.iv, I.b.2.v, and I.b.2.vi.

[00108] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:



I

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than or equal to a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of the same stereoisomer of formula **I**.

[00109] In some embodiments, a provided composition is characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of the same stereoisomer of formula **I**. In some embodiments, a provided composition is characterized in that a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of the same stereoisomer of formula **I**.

[00110] In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.i, I.a.ii, I.a.iii, I.a.iv, I.a.v, I.a.vi, I.a.vii, I.a.viii, or I.a.ix**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.i**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.ii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.iii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.iv**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.v**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.vi**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.vii**. In some


embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.viii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.a.ix**.

[00111] In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.i**, **I.b.1.ii**, **I.b.1.iii**, **I.b.1.iv**, **I.b.1.v**, or **I.b.1.vi**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.i**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.ii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.iii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.iv**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.v**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.1.vi**.

[00112] In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.i**, **I.b.2.ii**, **I.b.2.iii**, **I.b.2.iv**, **I.b.2.v**, or **I.b.2.vi**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.i**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.ii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.iii**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.iv**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.v**. In some embodiments, a stereoisomer of formula **I** has the structure of formula **I.b.2.vi**.


[00113] In some embodiments, the third threshold amount is 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% (w/w). In some embodiments, the third threshold amount is 50% (w/w). In some embodiments, the third threshold amount is 70% (w/w). In some embodiments, the third threshold amount is 80% (w/w). In some embodiments, the third threshold amount is 85% (w/w). In some embodiments, the third threshold amount is 90% (w/w). In some embodiments, the third threshold amount is 95% (w/w). In some embodiments, the third threshold amount is 96% (w/w). In some embodiments, the third threshold amount is 97% (w/w). In some embodiments, the third threshold amount is 98% (w/w). In some embodiments, the third threshold amount is 99% (w/w).

[00114] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**


a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,

the composition characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a.i**:

**I.a.i** (i.e., *R4-SR-cKK-E12*)


wherein the third threshold amount is 50%.

[00115] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:

**I**

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,


the composition characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a.ii**:



I.a.ii (*i.e.*, *S4-SR-cKK-E12*)

wherein the third threshold amount is 50%.


[00116] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:



I

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,


the composition characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1.i**:




I.b.1.i (*i.e.*, *R4-SS-cKK-E12*)

wherein the third threshold amount is 50%.

[00117] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:




a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,
the composition characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1.ii**:




wherein the third threshold amount is 50%.

[00118] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:




a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,
the composition characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2.i**:



I.b.2.i (*i.e.*, *R4-RR-cKK-E12*)


wherein the third threshold amount is 50%.

[00119] In some embodiments, the present invention provides a composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:



I

a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,
the composition characterized in that greater than a third threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2.ii**:



I.b.2.ii (*i.e.*, *S4-RR-cKK-E12*)

wherein the third threshold amount is 50%.

[00120] In some embodiments, the second threshold amount is 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%. In some embodiments, the second threshold amount is 50%. In some embodiments, the second threshold amount is 70%. In some embodiments, the second threshold amount is 80%. In some embodiments, the second threshold amount is 95%.

Liposomes for the Delivery of Agents, Such as mRNA

[00121] Among other things, the present invention provides composition comprising a lipid compound described herein for delivery of therapeutic agents. In some embodiments, a composition provided is a lipid based nanoparticle, such as a liposome. As used herein, the term “liposome” refers to any lamellar, multilamellar, or solid lipid nanoparticle vesicle. Typically, a liposome as used herein can be formed by mixing one or more lipids or by mixing one or more lipids and polymer(s). Thus, the term “liposome” as used herein encompasses both lipid and polymer based nanoparticles. In particular, a liposome according to the present invention incorporates a lipid compound described herein as a cationic lipid component. As a non-limiting example, a liposome according to the present invention includes a composition comprising one or more chemical entities of formula **I**. A suitable liposome may also contain second or additional cationic lipids, helper lipids (e.g., non-cationic lipids and/or cholesterol-based lipids), PEG-modified lipids, and/or polymers.

[00122] In some embodiments, cationic lipid(s) (e.g., the composition comprising one or more chemical entities of formula **I**) constitute(s) about 30-50% (e.g., about 30-45%, about 30-40%, about 35-50%, about 35-45%, or about 35-40%) of the liposome by molar ratio. In some embodiments, the cationic lipid (e.g., the composition comprising one or more chemical entities of formula **I**) constitutes about 30%, about 35%, about 40 %, about 45%, or about 50% of the liposome by molar ratio. In some embodiments, the liposome comprises a second lipid or additional cationic lipids.

Second or Additional Cationic Lipids

[00123] In some embodiments, liposomes may comprise a second or additional cationic lipid. As used herein, the phrase "cationic lipid" refers to any of a number of lipid species that have a

net positive charge at a selected pH, such as physiological pH. Several cationic lipids have been described in the literature, many of which are commercially available. Particularly suitable cationic lipids for use in the compositions and methods of the invention include those described in international patent publications WO 2010/053572 (and particularly, C12-200 described at paragraph [00225]), WO 2012/170930 and WO 2013063468 each of which is incorporated herein by reference in its entirety. In certain embodiments, the compositions and methods of the invention employ a lipid nanoparticles comprising an ionizable cationic lipid described in International Patent Application No. PCT/US2013/034602, filed March 29, 2013, Publ. No. WO 2013/149140 (incorporated herein by reference), such as, e.g., (15Z, 18Z)-N,N-dimethyl-6-(9Z, 12Z)-octadeca-9, 12-dien-1-yl)tetracos-15,18-dien-1-amine (HGT5000), (15Z, 18Z)-N,N-dimethyl-6-((9Z, 12Z)-octadeca-9, 12-dien-1-yl)tetracos-4,15,18-trien-1-amine (HGT5001), and (15Z,18Z)-N,N-dimethyl-6-((9Z, 12Z)-octadeca-9, 12-dien-1-yl)tetracos-5, 15, 18-trien-1-amine (HGT5002).

[00124] In some embodiments, the second or additional cationic lipid N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride or "DOTMA" is used. (Feigner et al. (Proc. Nat'l Acad. Sci. 84, 7413 (1987); U.S. Pat. No. 4,897,355). DOTMA can be formulated alone or can be combined with the neutral lipid, dioleoylphosphatidyl-ethanolamine or "DOPE" or other cationic or non-cationic lipids into a liposomal transfer vehicle or a lipid nanoparticle, and such liposomes can be used to enhance the delivery of nucleic acids into target cells. Other suitable cationic lipids include, for example, 5-carboxyspermylglycinedioctadecylamide or "DOGS," 2,3-dioleoyloxy-N-[2(spermine-carboxamido)ethyl]-N,N-dimethyl-1-propanaminium or "DOSPA" (Behr et al. Proc. Nat'l Acad. Sci. 86, 6982 (1989); U.S. Pat. No. 5,171,678; U.S. Pat. No. 5,334,761), 1,2-Dioleoyl-3-Dimethylammonium-Propane or "DODAP", 1,2-Dioleoyl-3-Trimethylammonium-Propane or "DOTAP". Additional exemplary cationic lipids also include 1,2-distearoyloxy-N,N-dimethyl-3-aminopropane or "DSDMA", 1,2-dioleoyloxy-N,N-dimethyl-3-aminopropane or "DODMA", 1,2-dilinoleoyloxy-N,N-dimethyl-3-aminopropane or "DLinDMA", 1,2-dilinenyloxy-N,N-dimethyl-3-aminopropane or "DLenDMA", N-dioleoyl-N,N-dimethylammonium chloride or "DODAC", N,N-distearyl-N,N-dimethylammonium bromide or "DDAB", N-(1,2-dimyristyloxyprop-3-yl)-N,N-dimethyl-N-hydroxyethyl ammonium bromide or "DMRIE", 3-dimethylamino-2-(cholest-5-en-3-beta-oxybutan-4-oxy)-1-(cis, cis-9,12-octadecadienoxy)propane or "CLinDMA", 2-[5'-(cholest-5-en-3-beta-oxy)-3'-oxapentoxy]-3-

dimethyl-1-(1-(cis,cis-9', 12'-octadecadienoxy)propane or "CpLinDMA", N,N-dimethyl-3,4-dioleoyloxybenzylamine or "DMOBA", 1,2-N,N'-dioleoylcarbonyl-3-dimethylaminopropane or "DOcarbDAP", 2,3-Dilinoleoyloxy-N,N-dimethylpropylamine or "DLinDAP", 1,2-N,N'-Dilinoleoylcarbonyl-3-dimethylaminopropane or "DLincarbDAP", 1,2-Dilinoleoylcarbonyl-3-dimethylaminopropane or "DLinCDAP", 2,2-dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane or "DLin-DMA", 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane or "DLin-K-XTC2-DMA", and 2-(2,2-di((9Z,12Z)-octadeca-9,12-dien-1-yl)-1,3-dioxolan-4-yl)-N,N-dimethylethanamine (DLin-KC2-DMA)) (See, WO 2010/042877; Semple et al., Nature Biotech. 28: 172-176 (2010)), or mixtures thereof. (Heyes, J., et al., J Controlled Release 107: 276-287 (2005); Morrissey, DV., et al., Nat. Biotechnol. 23(8): 1003-1007 (2005); PCT Publication WO2005/121348A1). In some embodiments, one or more of the cationic lipids comprise at least one of an imidazole, dialkylamino, or guanidinium moiety.

[00125] In some embodiments, the second or additional cationic lipid may be chosen from XTC (2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane), MC3 (((6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate), ALNY-100 ((3aR,5s,6aS)-N,N-dimethyl-2,2-di((9Z,12Z)-octadeca-9,12-dienyl)tetrahydro-3aH-cyclopenta[d][1,3]dioxol-5-amine)), NC98-5 (4,7,13-tris(3-oxo-3-(undecylamino)propyl)-N1,N16-diundecyl-4,7,10,13-tetraazahexadecane-1,16-diamide), DODAP (1,2-dioleoyl-3-dimethylammonium propane), HGT4003 (WO 2012/170889, the teachings of which are incorporated herein by reference in their entirety), ICE (WO 2011/068810, the teachings of which are incorporated herein by reference in their entirety), HGT5000 (international patent publication WO/2013/149140, the teachings of which are incorporated herein by reference in their entirety) or HGT5001 (cis or trans) (WO/2013/149140), aminoalcohol lipidoids such as those disclosed in WO2010/053572, DOTAP (1,2-dioleoyl-3-trimethylammonium propane), DOTMA (1,2-di-O-octadecenyl-3-trimethylammonium propane), DLinDMA (Heyes, J.; Palmer, L.; Bremner, K.; MacLachlan, I. "Cationic lipid saturation influences intracellular delivery of encapsulated nucleic acids" J. Contr. Rel. 2005, 107, 276-287), DLin-KC2-DMA (Semple, S.C. et al. "Rational Design of Cationic Lipids for siRNA Delivery" Nature Biotech. 2010, 28, 172-176), C12-200 (Love, K.T. et al. "Lipid-like materials for low-dose in vivo gene silencing" PNAS 2010, 107, 1864-1869).

Non-cationic/Helper Lipids

[00126] In some embodiments, provided liposomes contain one or more non-cationic ("helper") lipids. As used herein, the phrase "non-cationic lipid" refers to any neutral, zwitterionic or anionic lipid. As used herein, the phrase "anionic lipid" refers to any of a number of lipid species that carry a net negative charge at a selected H, such as physiological pH. Non-cationic lipids include, but are not limited to, distearoylphosphatidylcholine (DSPC), dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine (DPPC), dioleoylphosphatidylglycerol (DOPG), dipalmitoylphosphatidylglycerol (DPPG), dioleoylphosphatidylethanolamine (DOPE), palmitoyloleoylphosphatidylcholine (POPC), palmitoyloleoyl-phosphatidylethanolamine (POPE), dioleoyl-phosphatidylethanolamine 4-(N-maleimidomethyl)-cyclohexane-l-carboxylate (DOPE-mal), dipalmitoyl phosphatidyl ethanolamine (DPPE), dimyristoylphosphoethanolamine (DMPE), distearoyl-phosphatidylethanolamine (DSPE), 16-O-monomethyl PE, 16-O-dimethyl PE, 18-1-trans PE, 1-stearoyl-2-oleoyl-phosphatidylethanolamine (SOPE), or a mixture thereof.

[00127] In some embodiments, such non-cationic lipids may be used alone, but are preferably used in combination with other excipients, for example, cationic lipids. In some embodiments, the non-cationic lipid may comprise a molar ratio of about 5% to about 90%, or about 10 % to about 70% of the total lipid present in a liposome. In some embodiments, a non-cationic lipid is a neutral lipid, i.e., a lipid that does not carry a net charge in the conditions under which the composition is formulated and/or administered. In some embodiments, the percentage of non-cationic lipid in a liposome may be greater than 5%, greater than 10%, greater than 20%, greater than 30%, or greater than 40%.

Cholesterol-based Lipids

[00128] In some embodiments, provided liposomes comprise one or more cholesterol-based lipids. For example, suitable cholesterol-based cationic lipids include, for example, DC-Choi (N,N-dimethyl-N-ethylcarboxamidocholesterol), 1,4-bis(3-N-oleylamino-propyl)piperazine (Gao, et al. Biochem. Biophys. Res. Comm. 179, 280 (1991); Wolf et al. BioTechniques 23, 139 (1997); U.S. Pat. No. 5,744,335), or ICE. In some embodiments, the cholesterol-based lipid may comprise a molar ration of about 2% to about 30%, or about 5% to about 20% of the total lipid present in a liposome. In some embodiments, The percentage of cholesterol-based lipid in the

lipid nanoparticle may be greater than 5, %, 10%, greater than 20%, greater than 30%, or greater than 40%.

PEGylated Lipids

[00129] In some embodiments, provided liposomes comprise one or more PEGylated lipids. For example, the use of polyethylene glycol (PEG)-modified phospholipids and derivatized lipids such as derivatized ceramides (PEG-CER), including N-Octanoyl-Sphingosine-1-[Succinyl(Methoxy Polyethylene Glycol)-2000] (C8 PEG-2000 ceramide) is also contemplated by the present invention in combination with one or more of the cationic and, in some embodiments, other lipids together which comprise the liposome. Contemplated PEG-modified lipids include, but are not limited to, a polyethylene glycol chain of up to 5 kDa in length covalently attached to a lipid with alkyl chain(s) of C₆-C₂₀ length. In some embodiments, a PEG-modified or PEGylated lipid is PEGylated cholesterol or PEG-2K. The addition of such components may prevent complex aggregation and may also provide a means for increasing circulation lifetime and increasing the delivery of the lipid-nucleic acid composition to the target cell, (Klibanov et al. (1990) FEBS Letters, 268 (1): 235-237), or they may be selected to rapidly exchange out of the formulation in vivo (see U.S. Pat. No. 5,885,613).

[00130] In some embodiments, particularly useful exchangeable lipids are PEG-ceramides having shorter acyl chains (e.g., C₁₄ or C₁₈). The PEG-modified phospholipid and derivitized lipids of the present invention may comprise a molar ratio from about 0% to about 15%, about 0.5% to about 15%, about 1% to about 15%, about 4% to about 10%, or about 2% of the total lipid present in the liposome.

[00131] According to various embodiments, the selection of second or additional cationic lipids, non-cationic lipids and/or PEG-modified lipids which comprise the lipid nanoparticle, as well as the relative molar ratio of such lipids to each other, is based upon the characteristics of the selected lipid(s), the nature of the intended target cells, the characteristics of the mRNA to be delivered. Additional considerations include, for example, the saturation of the alkyl chain, as well as the size, charge, pH, pKa, fusogenicity and toxicity of the selected lipid(s). Thus the molar ratios may be adjusted accordingly. In some embodiments, the percentage of PEG-modified lipid in a liposome may be greater than 1%, greater than 2%, greater than 5%, greater than 10%, or greater than 15%.

Polymers

[00132] In some embodiments, a suitable liposome according to the present invention further includes a polymer, in combination with one or more cationic lipids as described and, in some embodiments, other carriers including various lipids described herein. Thus, in some embodiments, liposomal delivery vehicles, as used herein, also encompass polymer containing nanoparticles. Suitable polymers may include, for example, polyacrylates, polyalkycyanoacrylates, polylactide, polylactide-polyglycolide copolymers, polycaprolactones, dextran, albumin, gelatin, alginate, collagen, chitosan, cyclodextrins, protamine, PEGylated protamine, PLL, PEGylated PLL and polyethylenimine (PEI). When PEI is present, it may be branched PEI of a molecular weight ranging from 10 to 40 kDa, e.g., 25 kDa branched PEI (Sigma #408727).

Therapeutic Agents

[00133] The present invention may be used to delivery any therapeutic agents. Specifically, any therapeutic agents to be administered to a subject may be delivered using the complexes, picoparticles, nanoparticles, microparticles, micelles, or liposomes, described herein. The agent may be an organic molecule (e.g., a therapeutic agent, a drug), inorganic molecule, nucleic acid, protein, amino acid, peptide, polypeptide, polynucleotide, targeting agent, isotopically labeled organic or inorganic molecule, vaccine, immunological agent, *etc.* In certain embodiments of the present invention, the agent to be delivered may be a mixture of agents.

[00134] In certain embodiments, the therapeutic agents are organic molecules with pharmaceutical activity, e.g., a drug. In certain embodiments, the drug is an antibiotic, anti-viral agent, anesthetic, steroidal agent, anti-inflammatory agent, anti-neoplastic agent, anti-cancer agent, antigen, vaccine, antibody, decongestant, antihypertensive, sedative, birth control agent, progestational agent, anti-cholinergic, analgesic, anti-depressant, anti-psychotic, I3-adrenergic blocking agent, diuretic, cardiovascular active agent, vasoactive agent, non-steroidal anti-inflammatory agent, nutritional agent, *etc.*

[00135] Diagnostic agents include gases; metals; commercially available imaging agents used in positron emissions tomography (PET), computer assisted tomography (CAT), single photon emission computerized tomography, x-ray, fluoroscopy, and magnetic resonance imaging (MRI); and contrast agents. Examples of suitable materials for use as contrast agents in MRI include

gadolinium chelates, as well as iron, magnesium, manganese, copper, and chromium. Examples of materials useful for CAT and x-ray imaging include iodinebased materials.

[00136] Therapeutic and prophylactic agents include, but are not limited to, antibiotics, nutritional supplements, and vaccines. Vaccines may comprise isolated proteins or peptides, inactivated organisms and viruses, dead organisms and viruses, genetically altered organisms or viruses, and cell extracts.

Polynucleotides

[00137] The present invention may be used to deliver any polynucleotide. In certain embodiments, the polynucleotide is an interfering RNA (RNAi). The phenomenon of RNAi is discussed in greater detail, for example, in the following references: Elbashir *et al.*, 2001, *Genes Dev.*, 15:188; Fire *et al.*, 1998, *Nature*, 391:806; Tabara *et al.*, 1999, *Cell*, 99:123; Hammond *et al.*, *Nature*, 2000, 404:293; Zamore *et al.*, 2000, *Cell*, 101:25; Chakraborty, 2007, *Curr. Drug Targets*, 8:469; and Morris and Rossi, 2006, *Gene Ther.*, 13:553. In certain embodiments, the polynucleotide is a dsRNA (double-stranded RNA). In certain embodiments, the polynucleotide is an siRNA (short interfering RNA). In certain embodiments, the polynucleotide is an shRNA (short hairpin RNA). In certain embodiments, the polynucleotide is an miRNA (micro RNA). Micro RNAs (miRNAs) are genomically encoded non-coding RNAs of about 21 — 23 nucleotides in length that help regulate gene expression, particularly during development. See, e.g., Bartel, 2004, *Cell*, 116:281; Novina and Sharp, 2004, *Nature*, 430:161; and U.S. Patent Publication 2005/0059005; also reviewed in Wang and Li, 2007, *Front. Biosci.*, 12:3975; and Zhao, 2007, *Trends Biochem. Sci.*, 32:189. In certain embodiments, the polynucleotide is an antisense RNA.

[00138] In certain embodiments, the polynucleotide may be provided as an antisense agent or RNA interference (RNAi). See, e.g., Fire *et al.*, *Nature* 391:806-811, 1998. Antisense therapy is meant to include, e.g., administration or *in situ* provision of single- or double-stranded oligonucleotides or their derivatives which specifically hybridize, e.g., bind, under cellular conditions, with cellular mRNA and/or genomic DNA, or mutants thereof, so as to inhibit expression of the encoded protein, e.g., by inhibiting transcription and/or translation. See, e.g., Crooke "Molecular mechanisms of action of antisense drugs" *Biochim. Biophys. Acta* 1489(1):31-44, 1999; Crooke "Evaluating the mechanism of action of antiproliferative

antisense drugs" *Antisense Nucleic Acid Drug Dev.* 10(2):123-126, discussion 127, 2000; *Methods in Enzymology* volumes 313-314, 1999. The binding may be by conventional base pair complementarity, or, for example, in the case of binding to DNA duplexes, through specific interactions in the major groove of the double helix (i.e., triple helix formation). See, e.g., Chan *et al.*, *J. Mol. Med.* 75(4):267-282, 1997.

[00139] In some embodiments, dsRNA, siRNA, shRNA, miRNA, antisense RNA, and/or RNAi can be designed and/or predicted using one or more of a large number of available algorithms. To give but a few examples, the following resources can be utilized to design and/or predict polynucleotides: algorithms found at Alnylum Online, Dharmacon Online, OligoEngine Online, Molecula Online, Ambion Online, BioPredsi Online, RNAi Web Online, Chang Bioscience Online, Invitrogen Online, LentiWeb Online GenScript Online, Protocol Online; Reynolds *et al.*, 2004, *Nat. Biotechnol.*, 22:326; Naito *et al.*, 2006, *Nucleic Acids Res.*, 34:W448; Li *et al.*, 2007, *RNA*, 13:1765; Yiu *et al.*, 2005, *Bioinformatics*, 21:144; and Jia *et al.*, 2006, *BMC Bioinformatics*, 7: 271.

[00140] The polynucleotides may be of any size or sequence, and they may be single- or double-stranded. In certain embodiments, the polynucleotide is greater than 100 base pairs long. In certain embodiments, the polynucleotide is greater than 1000 base pairs long and may be greater than 10,000 base pairs long. The polynucleotide may be provided by any means known in the art. In certain embodiments, the polynucleotide has been engineered using recombinant techniques. See, e.g., Ausubel *et al.*, *Current Protocols in Molecular Biology* (John Wiley & Sons, Inc., New York, 1999); *Molecular Cloning: A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch, and Maniatis (Cold Spring Harbor Laboratory Press: 1989). The polynucleotide may also be obtained from natural sources and purified from contaminating components found normally in nature. The polynucleotide may also be chemically synthesized in a laboratory. In certain embodiments, the polynucleotide is synthesized using standard solid phase chemistry.

[00141] The polynucleotide may be modified by chemical or biological means. In certain embodiments, these modifications lead to increased stability of the polynucleotide. Modifications include methylation, phosphorylation, end-capping, *etc.*

mRNA

[00142] The present invention can be used to deliver any mRNA. mRNA is typically thought of as the type of RNA that carries information from DNA to the ribosome. The existence of mRNA is usually very brief and includes processing and translation, followed by degradation. Typically, in eukaryotic organisms, mRNA processing comprises the addition of a “cap” on the N-terminal (5’) end, and a “tail” on the C-terminal (3’) end. A typical cap is a 7-methylguanosine cap, which is a guanosine that is linked through a 5’-5’-triphosphate bond to the first transcribed nucleotide. The presence of the cap is important in providing resistance to nucleases found in most eukaryotic cells. The tail is typically a polyadenylation event whereby a polyadenylyl moiety is added to the 3’ end of the mRNA molecule. The presence of this “tail” serves to protect the mRNA from exonuclease degradation. Messenger RNA typically is translated by the ribosomes into a series of amino acids that make up a protein.

[00143] Any mRNA capable of being translated into one or more peptides (e.g., proteins) or peptide fragments is contemplated as within the scope of the present invention. In some embodiments, an mRNA encodes one or more naturally occurring peptides. In some embodiments, an mRNA encodes one or more modified or non-natural peptides.

[00144] In some embodiments an mRNA encodes an intracellular protein. In some embodiments, an mRNA encodes a cytosolic protein. In some embodiments, an mRNA encodes a protein associated with the actin cytoskeleton. In some embodiments, an mRNA encodes a protein associated with the plasma membrane. In some specific embodiments, an mRNA encodes a transmembrane protein. In some specific embodiments an mRNA encodes an ion channel protein. In some embodiments, an mRNA encodes a perinuclear protein. In some embodiments, an mRNA encodes a nuclear protein. In some specific embodiments, an mRNA encodes a transcription factor. In some embodiments, an mRNA encodes a chaperone protein. In some embodiments, an mRNA encodes an intracellular enzyme (e.g., mRNA encoding an enzyme associated with urea cycle or lysosomal storage metabolic disorders). In some embodiments, an mRNA encodes a protein involved in cellular metabolism, DNA repair, transcription and/or translation. In some embodiments, an mRNA encodes an extracellular protein. In some embodiments, an mRNA encodes a protein associated with the extracellular matrix. In some embodiments an mRNA encodes a secreted protein. In specific embodiments,

an mRNA used in the composition and methods of the invention may be used to express functional proteins or enzymes that are excreted or secreted by one or more target cells into the surrounding extracellular fluid (e.g., mRNA encoding hormones and/or neurotransmitters).

Synthesis of mRNA

[00145] mRNAs according to the present invention may be synthesized according to any of a variety of known methods. For example, mRNAs according to the present invention may be synthesized via *in vitro* transcription (IVT). Briefly, IVT is typically performed with a linear or circular DNA template containing a promoter, a pool of ribonucleotide triphosphates, a buffer system that may include DTT and magnesium ions, and an appropriate RNA polymerase (e.g., T3, T7 or SP6 RNA polymerase), DNase I, pyrophosphatase, and/or RNase inhibitor. The exact conditions will vary according to the specific application.

[00146] In some embodiments, for the preparation of mRNA according to the invention, a DNA template is transcribed *in vitro*. A suitable DNA template typically has a promoter, for example a T3, T7 or SP6 promoter, for *in vitro* transcription, followed by desired nucleotide sequence for desired mRNA and a termination signal.

[00147] Desired mRNA sequence(s) according to the invention may be determined and incorporated into a DNA template using standard methods. For example, starting from a desired amino acid sequence (e.g., an enzyme sequence), a virtual reverse translation is carried out based on the degenerated genetic code. Optimization algorithms may then be used for selection of suitable codons. Typically, the G/C content can be optimized to achieve the highest possible G/C content on one hand, taking into the best possible account the frequency of the tRNAs according to codon usage on the other hand. The optimized RNA sequence can be established and displayed, for example, with the aid of an appropriate display device and compared with the original (wild-type) sequence. A secondary structure can also be analyzed to calculate stabilizing and destabilizing properties or, respectively, regions of the RNA.

Modified mRNA

[00148] In some embodiments, mRNA according to the present invention may be synthesized as unmodified or modified mRNA. Typically, mRNAs are modified to enhance stability. Modifications of mRNA can include, for example, modifications of the nucleotides of the RNA.

An modified mRNA according to the invention can thus include, for example, backbone modifications, sugar modifications or base modifications. In some embodiments, mRNAs may be synthesized from naturally occurring nucleotides and/or nucleotide analogues (modified nucleotides) including, but not limited to, purines (adenine (A), guanine (G)) or pyrimidines (thymine (T), cytosine (C), uracil (U)), and as modified nucleotides analogues or derivatives of purines and pyrimidines, such as e.g. 1-methyl-adenine, 2-methyl-adenine, 2-methylthio-N-6-isopentenyl-adenine, N6-methyl-adenine, N6-isopentenyl-adenine, 2-thio-cytosine, 3-methyl-cytosine, 4-acetyl-cytosine, 5-methyl-cytosine, 2,6-diaminopurine, 1-methyl-guanine, 2-methyl-guanine, 2,2-dimethyl-guanine, 7-methyl-guanine, inosine, 1-methyl-inosine, pseudouracil (5-uracil), dihydro-uracil, 2-thio-uracil, 4-thio-uracil, 5-carboxymethylaminomethyl-2-thio-uracil, 5-(carboxyhydroxymethyl)-uracil, 5-fluoro-uracil, 5-bromo-uracil, 5-carboxymethylaminomethyl-uracil, 5-methyl-2-thio-uracil, 5-methyl-uracil, N-uracil-5-oxyacetic acid methyl ester, 5-methylaminomethyl-uracil, 5-methoxyaminomethyl-2-thio-uracil, 5'-methoxycarbonylmethyl-uracil, 5-methoxy-uracil, uracil-5-oxyacetic acid methyl ester, uracil-5-oxyacetic acid (v), 1-methyl-pseudouracil, queosine, β -D-mannosyl-queosine, wybutosine, and phosphoramidates, phosphorothioates, peptide nucleotides, methylphosphonates, 7-deazaguanosine, 5-methylcytosine and inosine. The preparation of such analogues is known to a person skilled in the art e.g. from the U.S. Pat. No. 4,373,071, U.S. Pat. No. 4,401,796, U.S. Pat. No. 4,415,732, U.S. Pat. No. 4,458,066, U.S. Pat. No. 4,500,707, U.S. Pat. No. 4,668,777, U.S. Pat. No. 4,973,679, U.S. Pat. No. 5,047,524, U.S. Pat. No. 5,132,418, U.S. Pat. No. 5,153,319, U.S. Pat. Nos. 5,262,530 and 5,700,642, the disclosures of which are incorporated by reference in their entirety.

[00149] In some embodiments, mRNAs (e.g., enzyme encoding mRNAs) may contain RNA backbone modifications. Typically, a backbone modification is a modification in which the phosphates of the backbone of the nucleotides contained in the RNA are modified chemically. Exemplary backbone modifications typically include, but are not limited to, modifications from the group consisting of methylphosphonates, methylphosphoramidates, phosphoramidates, phosphorothioates (e.g. cytidine 5'-O-(1-thiophosphate)), boranophosphates, positively charged guanidinium groups etc., which means by replacing the phosphodiester linkage by other anionic, cationic or neutral groups.

[00150] In some embodiments, mRNAs (e.g., enzyme encoding mRNAs) may contain sugar modifications. A typical sugar modification is a chemical modification of the sugar of the nucleotides it contains including, but not limited to, sugar modifications chosen from the group consisting of 2'-deoxy-2'-fluoro-oligoribonucleotide (2'-fluoro-2'-deoxycytidine 5'-triphosphate, 2'-fluoro-2'-deoxyuridine 5'-triphosphate), 2'-deoxy-2'-deamine-oligoribonucleotide (2'-amino-2'-deoxycytidine 5'-triphosphate, 2'-amino-2'-deoxyuridine 5'-triphosphate), 2'-O-alkyloligoribonucleotide, 2'-deoxy-2'-C-alkyloligoribonucleotide (2'-O-methylcytidine 5'-triphosphate, 2'-methyluridine 5'-triphosphate), 2'-C-alkyloligoribonucleotide, and isomers thereof (2'-aracytidine 5'-triphosphate, 2'-arauridine 5'-triphosphate), or azidotriphosphates (2'-azido-2'-deoxycytidine 5'-triphosphate, 2'-azido-2'-deoxyuridine 5'-triphosphate).

[00151] In some embodiments, mRNAs (e.g., enzyme encoding mRNAs) may contain modifications of the bases of the nucleotides (base modifications). A modified nucleotide which contains a base modification is also called a base-modified nucleotide. Examples of such base-modified nucleotides include, but are not limited to, 2-amino-6-chloropurine riboside 5'-triphosphate, 2-aminoadenosine 5'-triphosphate, 2-thiocytidine 5'-triphosphate, 2-thiouridine 5'-triphosphate, 4-thiouridine 5'-triphosphate, 5-aminoallylcytidine 5'-triphosphate, 5-aminoallyluridine 5'-triphosphate, 5-bromocytidine 5'-triphosphate, 5-bromouridine 5'-triphosphate, 5-iodocytidine 5'-triphosphate, 5-iodouridine 5'-triphosphate, 5-methylcytidine 5'-triphosphate, 5-methyluridine 5'-triphosphate, 6-azacytidine 5'-triphosphate, 6-azauridine 5'-triphosphate, 6-chloropurine riboside 5'-triphosphate, 7-deazaadenosine 5'-triphosphate, 7-deazaguanosine 5'-triphosphate, 8-azaadenosine 5'-triphosphate, 8-azidoadenosine 5'-triphosphate, benzimidazole riboside 5'-triphosphate, N1-methyladenosine 5'-triphosphate, N1-methylguanosine 5'-triphosphate, N6-methyladenosine 5'-triphosphate, O6-methylguanosine 5'-triphosphate, pseudouridine 5'-triphosphate, puromycin 5'-triphosphate or xanthosine 5'-triphosphate.

Cap Structure

[00152] Typically, mRNA synthesis includes the addition of a “cap” on the N-terminal (5') end, and a “tail” on the C-terminal (3') end. The presence of the cap is important in providing resistance to nucleases found in most eukaryotic cells. The presence of a “tail” serves to protect the mRNA from exonuclease degradation.

[00153] Thus, in some embodiments, mRNAs (e.g., enzyme encoding mRNAs) include a 5' cap structure. A 5' cap is typically added as follows: first, an RNA terminal phosphatase removes one of the terminal phosphate groups from the 5' nucleotide, leaving two terminal phosphates; guanosine triphosphate (GTP) is then added to the terminal phosphates via a guanylyl transferase, producing a 5'5'5 triphosphate linkage; and the 7-nitrogen of guanine is then methylated by a methyltransferase. Examples of cap structures include, but are not limited to, m⁷G(5')ppp (5'(A,G(5')ppp(5')A and G(5')ppp(5')G.

[00154] In some embodiments, naturally occurring cap structures comprise a 7-methyl guanosine that is linked via a triphosphate bridge to the 5'-end of the first transcribed nucleotide, resulting in a dinucleotide cap of m⁷G(5')ppp(5')N, where N is any nucleoside. *In vivo*, the cap is added enzymatically. The cap is added in the nucleus and is catalyzed by the enzyme guanylyl transferase. The addition of the cap to the 5' terminal end of RNA occurs immediately after initiation of transcription. The terminal nucleoside is typically a guanosine, and is in the reverse orientation to all the other nucleotides, i.e., G(5')ppp(5')GpNpNp.

[00155] A common cap for mRNA produced by *in vitro* transcription is m⁷G(5')ppp(5')G, which has been used as the dinucleotide cap in transcription with T7 or SP6 RNA polymerase *in vitro* to obtain RNAs having a cap structure in their 5'-termini. The prevailing method for the *in vitro* synthesis of capped mRNA employs a pre-formed dinucleotide of the form m⁷G(5')ppp(5')G (“m⁷GpppG”) as an initiator of transcription.

[00156] To date, a usual form of a synthetic dinucleotide cap used in *in vitro* translation experiments is the Anti-Reverse Cap Analog (“ARCA”) or modified ARCA, which is generally a modified cap analog in which the 2' or 3' OH group is replaced with -OCH₃.

[00157] Additional cap analogs include, but are not limited to, chemical structures selected from the group consisting of m⁷GpppG, m⁷GpppA, m⁷GpppC; unmethylated cap analogs (e.g., GpppG); dimethylated cap analog (e.g., m^{2,7}GpppG), trimethylated cap analog (e.g., m^{2,2,7}GpppG), dimethylated symmetrical cap analogs (e.g., m⁷Gpppm⁷G), or anti reverse cap analogs (e.g., ARCA; m^{7,2'Ome}GpppG, m^{7,2'd}GpppG, m^{7,3'Ome}GpppG, m^{7,3'd}GpppG and their tetraphosphate derivatives) (see, e.g., Jemielity, J. et al., “Novel ‘anti-reverse’ cap analogs with superior translational properties”, RNA, 9: 1108-1122 (2003)).

[00158] In some embodiments, a suitable cap is a 7-methyl guanylate (“m⁷G”) linked via a triphosphate bridge to the 5'-end of the first transcribed nucleotide, resulting in m⁷G(5')ppp(5')N, where N is any nucleoside. A preferred embodiment of a m⁷G cap utilized in embodiments of the invention is m⁷G(5')ppp(5')G.

[00159] In some embodiments, the cap is a Cap0 structure. Cap0 structures lack a 2'-O-methyl residue of the ribose attached to bases 1 and 2. In some embodiments, the cap is a Cap1 structure. Cap1 structures have a 2'-O-methyl residue at base 2. In some embodiments, the cap is a Cap2 structure. Cap2 structures have a 2'-O-methyl residue attached to both bases 2 and 3.

[00160] A variety of m⁷G cap analogs are known in the art, many of which are commercially available. These include the m⁷GpppG described above, as well as the ARCA 3'-OCH₃ and 2'-OCH₃ cap analogs (Jemielity, J. et al., RNA, 9: 1108-1122 (2003)). Additional cap analogs for use in embodiments of the invention include N7-benzylated dinucleoside tetraphosphate analogs (described in Grudzien, E. et al., RNA, 10: 1479-1487 (2004)), phosphorothioate cap analogs (described in Grudzien-Nogalska, E., et al., RNA, 13: 1745-1755 (2007)), and cap analogs (including biotinylated cap analogs) described in U.S. Patent Nos. 8,093,367 and 8,304,529, incorporated by reference herein.

Tail Structure

[00161] Typically, the presence of a “tail” serves to protect the mRNA from exonuclease degradation. The poly A tail is thought to stabilize natural messengers and synthetic sense RNA. Therefore, in certain embodiments a long poly A tail can be added to an mRNA molecule thus rendering the RNA more stable. Poly A tails can be added using a variety of art-recognized techniques. For example, long poly A tails can be added to synthetic or *in vitro* transcribed RNA using poly A polymerase (Yokoe, *et al.* Nature Biotechnology. 1996; 14: 1252-1256). A transcription vector can also encode long poly A tails. In addition, poly A tails can be added by transcription directly from PCR products. Poly A may also be ligated to the 3' end of a sense RNA with RNA ligase (see, e.g., Molecular Cloning A Laboratory Manual, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1991 edition)).

[00162] In some embodiments, mRNAs (e.g., enzyme encoding mRNAs) include a 3' poly(A) tail structure. Typically, the length of the poly A tail can be at least about 10, 50, 100, 200, 300, 400 at least 500 nucleotides. In some embodiments, a poly-A tail on the 3' terminus of mRNA

typically includes about 10 to 300 adenosine nucleotides (*e.g.*, about 10 to 200 adenosine nucleotides, about 10 to 150 adenosine nucleotides, about 10 to 100 adenosine nucleotides, about 20 to 70 adenosine nucleotides, or about 20 to 60 adenosine nucleotides). In some embodiments, mRNAs include a 3' poly(C) tail structure. A suitable poly-C tail on the 3' terminus of mRNA typically include about 10 to 200 cytosine nucleotides (*e.g.*, about 10 to 150 cytosine nucleotides, about 10 to 100 cytosine nucleotides, about 20 to 70 cytosine nucleotides, about 20 to 60 cytosine nucleotides, or about 10 to 40 cytosine nucleotides). The poly-C tail may be added to the poly-A tail or may substitute the poly-A tail.

[00163] In some embodiments, the length of the poly A or poly C tail is adjusted to control the stability of a modified sense mRNA molecule of the invention and, thus, the transcription of protein. For example, since the length of the poly A tail can influence the half-life of a sense mRNA molecule, the length of the poly A tail can be adjusted to modify the level of resistance of the mRNA to nucleases and thereby control the time course of polynucleotide expression and/or polypeptide production in a target cell.

5' and 3' Untranslated Region

[00164] In some embodiments, mRNAs include a 5' and/or 3' untranslated region. In some embodiments, a 5' untranslated region includes one or more elements that affect an mRNA's stability or translation, for example, an iron responsive element. In some embodiments, a 5' untranslated region may be between about 50 and 500 nucleotides in length.

[00165] In some embodiments, a 3' untranslated region includes one or more of a polyadenylation signal, a binding site for proteins that affect an mRNA's stability of location in a cell, or one or more binding sites for miRNAs. In some embodiments, a 3' untranslated region may be between 50 and 500 nucleotides in length or longer.

[00166] Exemplary 3' and/or 5' UTR sequences can be derived from mRNA molecules which are stable (*e.g.*, globin, actin, GAPDH, tubulin, histone, or citric acid cycle enzymes) to increase the stability of the sense mRNA molecule. For example, a 5' UTR sequence may include a partial sequence of a CMV immediate-early 1 (IE1) gene, or a fragment thereof to improve the nuclease resistance and/or improve the half-life of the polynucleotide. Also contemplated is the inclusion of a sequence encoding human growth hormone (hGH), or a fragment thereof to the 3' end or untranslated region of the polynucleotide (*e.g.*, mRNA) to further stabilize the

polynucleotide. Generally, these modifications improve the stability and/or pharmacokinetic properties (e.g., half-life) of the polynucleotide relative to their unmodified counterparts, and include, for example modifications made to improve such polynucleotides' resistance to *in vivo* nuclease digestion.

[00167] According to various embodiments, any size mRNA may be encapsulated by provided liposomes. In some embodiments, the provided liposomes may encapsulate mRNA of greater than about 0.5 kb, 1 kb, 1.5 kb, 2 kb, 2.5 kb, 3 kb, 3.5 kb, 4 kb, 4.5 kb, or 5 kb in length.

Liposomes

[00168] The liposomes for use in provided compositions can be prepared by various techniques which are presently known in the art. For example, multilamellar vesicles (MLV) may be prepared according to conventional techniques, such as by depositing a selected lipid on the inside wall of a suitable container or vessel by dissolving the lipid in an appropriate solvent, and then evaporating the solvent to leave a thin film on the inside of the vessel or by spray drying. An aqueous phase may then added to the vessel with a vortexing motion which results in the formation of MLVs. Uni-lamellar vesicles (ULV) can then be formed by homogenization, sonication or extrusion of the multi-lamellar vesicles. In addition, unilamellar vesicles can be formed by detergent removal techniques.

[00169] In certain embodiments, provided compositions comprise a liposome wherein an agent, such as a nucleic acid e.g., mRNA, is associated on both the surface of the liposome and encapsulated within the same liposome. For example, during preparation of the compositions of the present invention, cationic liposomes may associate with the mRNA through electrostatic interactions. For example, during preparation of the compositions of the present invention, cationic liposomes may associate with the mRNA through electrostatic interactions.

[00170] In some embodiments, the compositions and methods of the invention comprise mRNA encapsulated in a liposome. In some embodiments, the one or more mRNA species may be encapsulated in the same liposome. In some embodiments, the one or more mRNA species may be encapsulated in different liposomes. In some embodiments, the mRNA is encapsulated in one or more liposomes, which differ in their lipid composition, molar ratio of lipid components, size, charge (Zeta potential), targeting ligands and/or combinations thereof. In some embodiments, the one or more liposome may have a different composition of cationic

lipids, neutral lipid, PEG-modified lipid and/or combinations thereof. In some embodiments the one or more liposomes may have a different molar ratio of cationic lipid, neutral lipid, cholesterol and PEG-modified lipid used to create the liposome.

[00171] The process of incorporation of a desired therapeutic agent, such as a nucleic acid (e.g., mRNA), into a liposome is often referred to as “loading”. Exemplary methods are described in Lasic, et al., FEBS Lett., 312: 255-258, 1992, which is incorporated herein by reference. The liposome-incorporated nucleic acids may be completely or partially located in the interior space of the liposome, within the bilayer membrane of the liposome, or associated with the exterior surface of the liposome membrane. The incorporation of a nucleic acid into liposomes is also referred to herein as “encapsulation” wherein the nucleic acid is entirely contained within the interior space of the liposome. The purpose of incorporating a mRNA into a transfer vehicle, such as a liposome, is often to protect the nucleic acid from an environment which may contain enzymes or chemicals that degrade nucleic acids and/or systems or receptors that cause the rapid excretion of the nucleic acids. Accordingly, in some embodiments, a suitable delivery vehicle is capable of enhancing the stability of the mRNA contained therein and/or facilitate the delivery of mRNA to the target cell or tissue.

Liposome Size

[00172] Suitable liposomes in accordance with the present invention may be made in various sizes. In some embodiments, provided liposomes may be made smaller than previously known mRNA encapsulating liposomes. In some embodiments, decreased size of liposomes is associated with more efficient delivery of mRNA. Selection of an appropriate liposome size may take into consideration the site of the target cell or tissue and to some extent the application for which the liposome is being made.

[00173] In some embodiments, an appropriate size of liposome is selected to facilitate systemic distribution of antibody encoded by the mRNA. In some embodiments, it may be desirable to limit transfection of the mRNA to certain cells or tissues. For example, to target hepatocytes a liposome may be sized such that its dimensions are smaller than the fenestrations of the endothelial layer lining hepatic sinusoids in the liver; in such cases the liposome could readily penetrate such endothelial fenestrations to reach the target hepatocytes.

[00174] Alternatively or additionally, a liposome may be sized such that the dimensions of the liposome are of a sufficient diameter to limit or expressly avoid distribution into certain cells or tissues. For example, a liposome may be sized such that its dimensions are larger than the fenestrations of the endothelial layer lining hepatic sinusoids to thereby limit distribution of the liposomes to hepatocytes.

[00175] In some embodiments, a suitable liposome has a size of or less than about 500 nm, 450 nm, 400 nm, 350 nm, 300 nm, 250 nm, 200 nm, 150 nm, 125 nm, 110 nm, 100 nm, 95 nm, 90 nm, 85 nm, 80 nm, 75 nm, 70 nm, 65 nm, 60 nm, 55 nm, or 50 nm. In some embodiments, a suitable liposome has a size no greater than about 250 nm (e.g., no greater than about 225 nm, 200 nm, 175 nm, 150 nm, 125 nm, 100 nm, 75 nm, or 50 nm). In some embodiments, a suitable liposome has a size ranging from about 10 - 250 nm (e.g., ranging from about 10 - 225 nm, 10 - 200 nm, 10 - 175 nm, 10 - 150 nm, 10 - 125 nm, 10 - 100 nm, 10 - 75 nm, or 10 - 50 nm). In some embodiments, a suitable liposome has a size ranging from about 100 - 250 nm (e.g., ranging from about 100 - 225 nm, 100 - 200 nm, 100 - 175 nm, 100 - 150 nm). In some embodiments, a suitable liposome has a size ranging from about 10 - 100 nm (e.g., ranging from about 10 - 90 nm, 10 - 80 nm, 10 - 70 nm, 10 - 60 nm, or 10 - 50 nm).

[00176] A variety of alternative methods known in the art are available for sizing of a population of liposomes. One such sizing method is described in U.S. Pat. No. 4,737,323, incorporated herein by reference. Sonicating a liposome suspension either by bath or probe sonication produces a progressive size reduction down to small ULV less than about 0.05 microns in diameter. Homogenization is another method that relies on shearing energy to fragment large liposomes into smaller ones. In a typical homogenization procedure, MLV are recirculated through a standard emulsion homogenizer until selected liposome sizes, typically between about 0.1 and 0.5 microns, are observed. The size of the liposomes may be determined by quasi-electric light scattering (QELS) as described in Bloomfield, Ann. Rev. Biophys. Bioeng., 10:421-150 (1981), incorporated herein by reference. Average liposome diameter may be reduced by sonication of formed liposomes. Intermittent sonication cycles may be alternated with QELS assessment to guide efficient liposome synthesis.

Pharmaceutical Compositions

[00177] To facilitate delivery of an agent, such as a nucleic acid e.g., mRNA, and/or expression of mRNA *in vivo*, delivery vehicles such as liposomes can be formulated in combination with one or more additional nucleic acids, carriers, targeting ligands or stabilizing reagents, or in pharmacological compositions where it is mixed with suitable excipients. Techniques for formulation and administration of drugs may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., latest edition.

[00178] Provided liposomally-encapsulated agents, such as a nucleic acid e.g., mRNA and compositions containing the same, may be administered and dosed in accordance with current medical practice, taking into account the clinical condition of the subject, the site and method of administration, the scheduling of administration, the subject's age, sex, body weight and other factors relevant to clinicians of ordinary skill in the art. The "effective amount" for the purposes herein may be determined by such relevant considerations as are known to those of ordinary skill in experimental clinical research, pharmacological, clinical and medical arts. In some embodiments, the amount administered is effective to achieve at least some stabilization, improvement or elimination of symptoms and other indicators as are selected as appropriate measures of disease progress, regression or improvement by those of skill in the art. For example, a suitable amount and dosing regimen is one that causes at least transient protein (e.g., enzyme) production.

[00179] Suitable routes of administration include, for example, oral, rectal, vaginal, transmucosal, pulmonary including intratracheal or inhaled, or intestinal administration; parenteral delivery, including intradermal, transdermal (topical), intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, and/or intranasal administration.

[00180] Alternately or additionally, liposomally encapsulated agents, such as a nucleic acid e.g., mRNA and compositions of the invention may be administered in a local rather than systemic manner, for example, via injection of the pharmaceutical composition directly into a targeted tissue, preferably in a sustained release formulation. Local delivery can be affected in various ways, depending on the tissue to be targeted. For example, aerosols containing compositions of the present invention can be inhaled (for nasal, tracheal, or bronchial delivery);

compositions of the present invention can be injected into the site of injury, disease manifestation, or pain, for example; compositions can be provided in lozenges for oral, tracheal, or esophageal application; can be supplied in liquid, tablet or capsule form for administration to the stomach or intestines, can be supplied in suppository form for rectal or vaginal application; or can even be delivered to the eye by use of creams, drops, or even injection. Formulations containing provided compositions complexed with therapeutic molecules or ligands can even be surgically administered, for example in association with a polymer or other structure or substance that can allow the compositions to diffuse from the site of implantation to surrounding cells. Alternatively, they can be applied surgically without the use of polymers or supports.

[00181] In some embodiments, provided liposomes and/or compositions are formulated such that they are suitable for extended-release of the agent, e.g., mRNA contained therein. Such extended-release compositions may be conveniently administered to a subject at extended dosing intervals. For example, in one embodiment, the compositions of the present invention are administered to a subject twice day, daily or every other day. In a preferred embodiment, the compositions of the present invention are administered to a subject twice a week, once a week, every ten days, every two weeks, every three weeks, or more preferably every four weeks, once a month, every six weeks, every eight weeks, every other month, every three months, every four months, every six months, every eight months, every nine months or annually. Also contemplated are compositions and liposomes which are formulated for depot administration (e.g., intramuscularly, subcutaneously, intravitreally) to either deliver or release a mRNA over extended periods of time. Preferably, the extended-release means employed are combined with modifications made to the mRNA to enhance stability.


[00182] Also contemplated herein are lyophilized pharmaceutical compositions comprising one or more of the liposomes disclosed herein and related methods for the use of such compositions as disclosed for example, in International Patent Application No. PCT/US2012/041663, filed June 8, 2012, Publ. No. WO 2012/170889, the teachings of which are incorporated herein by reference in their entirety. For example, lyophilized pharmaceutical compositions according to the invention may be reconstituted prior to administration or can be reconstituted *in vivo*. For example, a lyophilized pharmaceutical composition can be formulated in an appropriate dosage form (e.g., an intradermal dosage form such as a disk, rod or

membrane) and administered such that the dosage form is rehydrated over time in vivo by the individual's bodily fluids.

[00183] Provided liposomes and compositions may be administered to any desired tissue. In some embodiments, the agent, e.g., mRNA delivered by provided liposomes or compositions is expressed in the tissue in which the liposomes and/or compositions were administered. In some embodiments, the mRNA delivered is expressed in a tissue different from the tissue in which the liposomes and/or compositions were administered. Exemplary tissues in which delivered mRNA may be delivered and/or expressed include, but are not limited to the liver, kidney, heart, spleen, serum, brain, skeletal muscle, lymph nodes, skin, and/or cerebrospinal fluid.

[00184] According to various embodiments, the timing of expression of delivered agents, e.g., mRNAs, can be tuned to suit a particular medical need. In some embodiments, the expression of the protein encoded by delivered mRNA is detectable 1, 2, 3, 6, 12, 18, 24, 30, 36, 42, 48, 54, 60, 66, and/or 72 hours in serum or target tissues after a single administration of provided liposomes or compositions. In some embodiments, the expression of the protein encoded by the mRNA is detectable 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, and/or 7 days in serum or target tissues after a single administration of provided liposomes or compositions. In some embodiments, the expression of the protein encoded by the mRNA is detectable 1 week, 2 weeks, 3 weeks, and/or 4 weeks in serum or target tissues after a single administration of provided liposomes or compositions. In some embodiments, the expression of the protein encoded by the mRNA is detectable after a month or longer after a single administration of provided liposomes or compositions.

EXAMPLES

Example 1 - Synthesis of Racemic Compounds of Formula I**Scheme 1. Synthetic route to racemic compounds of Formula I**

[00185] Racemic compound **10** was prepared from protected lysine derivative **1**, Boc-Lys(Z)-OSu, by cleavage of the alpha-amino *tert*-butyl carbamate with trifluoroacetic acid and dimerization of the resulting free amine to form 3,6-dioxopiperazine **2**. Hydrogenation of **2** over catalytic palladium yields primary diamine **3**, which is treated with four equivalents of epoxide **4** and triethylamine to form the racemic 3,6-dioxopiperazine **10** of formula I.

[00186] *Dibenzyl(((2S,5S)-3,6-dioxopiperazine-2,5-diyl)bis(butane-4,1-diyl))dicarbamate* **2**: To Boc-Lys(Z)-OSu **1** (50 g) cooled with an ice bath was added TFA (60 mL) slowly. The resulting mixture was stirred for 20 min. The ice bath was removed and stirring was continued for 1 h. TFA was removed by rotary evaporation. The oily residue was dissolved in DMF (80 mL) and added slowly to stirred pyridine (anhydrous, 2.5 L). Mass spectrometry indicated completion of reaction after 1 h. Pyridine was removed by rotary evaporation and the residue

was diluted with EtOAc (2 L). After 20 min of stirring the mixture was filtered to give **2** as off-white solid (20 g after overnight vacuum drying. Yield: 73%).


[00187] *(3S,6S)-3,6-bis(4-Aminobutyl)piperazine-2,5-dione – 2HOAc 3*: To compound **2** in a mixture of AcOH (550 mL) and DCM (550 mL) was added Pd/C (10%, wet. 10 g). This mixture was stirred under a H₂ balloon for 4 h when mass spectrometry indicated completion of reaction. The reaction mixture was flushed with nitrogen for 10 min and filtered through Celite. The Celite was rinsed with MeOH (3 X 250 mL). The combined filtrate was evaporated and the residue was stirred with EtOAc (1.0 L) for 30 min. Filtration gave **3** as white solid (16.37 g after overnight vacuum drying. Yield: 114%. ¹H NMR shows clean product but with extra HOAc in sample)

[00188] *3,6-bis(4-(bis(2-Hydroxydodecyl)amino)butyl)piperazine-2,5-dione 10*: To a mixture of **3** (15.87 g, 42.2 mmol) and **4** (57 mL, 261 mmol) in EtOH (75 mL) stirred in a 500 mL pressure flask at room temperature was added Et₃N (23 mL, 165 mmol). The flask was flushed with nitrogen for 5 min and sealed. The mixture (solid and liquid slurry) was stirred for 30 min at room temperature then heated to 150-155 °C (oil bath temperature) and stirred for 5 h. A clear solution was obtained after temperature reached 150 °C. After being cooled to room temperature the reaction solution was evaporated and the residue was purified by flash column chromatography 9 times with 0-30% MeOH/DCM as eluent and 2 times with 0-30% MeOH/EtOAc as eluent. Use of DCM to 50% of 75:22:3 DCM/MeOH/NH₄OH (aq.) as eluent led to co-elution of product with a less polar side product. The side product ran closely with product on TLC with 30% MeOH/DCM as developing solvents. It was well separated from product on TLC with 30% MeOH/EtOAc as developing solvents. Pure product fractions from column purifications were pooled and evaporated. The oily residue was dissolved in Et₂O and evaporated. Drying under vacuum overnight removed all solvents and gave the racemic compound **10** as light yellow gel (9.85 g, yield: 24%). HPLC with ELSD detection showed two similar sized peaks with same product mass. Elemental Analysis: (Calc): C, 72.63; H, 12.17; N, 5.64; (Obsd): C, 72.25; H, 12.37; N, 5.68. Mass Spec: 993.8 *m/z*.

[00189] In another run with 7.15 g of **3** and 25.6 mL of **4** the crude product was purified twice with 0-30% MeOH/EtOAc as eluent to give two batches of product: the 1.55 g batch from early fractions and the 7.38 g batch from late fractions. Both batches were pure by ¹H NMR. HPLC

with ELSD detection showed two similar sized peaks with same product mass for the 7.38 g batch but only a single product peak for the 1.55 g batch.


Example 2 - Synthesis of Chiral Compounds of Formula I.b.1 (i.e., compound 10).



Scheme 2. Synthetic route to chiral compounds of Formula I.b.1 (i.e., compound 10).


[00190] Chiral compound **10** was synthesized via *N*-alkylation of protected lysine derivative **5** with two equivalents of epoxide **4** to form diol **6**. Hydrogenation of diol **6** over catalytic palladium forms α-amino acid **7**, which is divided into two portions. The first portion of α-amino acid **7** has its α-amino group protected as the *tert*-butyl carbamate, upon treatment with Boc anhydride, to form carboxylic acid **8**. The second portion of α-amino acid **7** has its free acid converted to the methyl ester to form amine **9**. Carboxylic acid **8** and amine **9** are coupled to an amide intermediate via peptide coupling reagents such as HATU and diethanolamine in aprotic solvent such as DMF, the *tert*-butyl carbamate group of the amide intermediate is cleaved with trifluoroacetic acid in dichloromethane, and the resulting amino ester product is cyclized to the piperazine-2,5-dione **10**. The stereochemistry at all chiral centers is preserved via this route.


[00191] Chiral compounds 12-15, below, are prepared via described synthetic routes using respective chiral epoxide starting materials.



Scheme 3. *Chiral compounds 12-15.*

Example 3 - *The compound of **Formula I.a.i** (i.e., R4-SR-cKK-E12)*



Scheme 4: Synthesis of the compound of Formula I.a.i (i.e., R4-SR-cKK-E12)

[00192] *Synthesis of compound 3.1:* To a suspension of Mg (60 g, 2.5 mol) in anhydrous Et₂O (1000 mL) was added one crystal of iodine, followed by addition of 1-bromononane (25 mL). The reaction was initiated and the solution began to reflux after heating the reaction flask with a water bath. The remaining 1-bromononane (360 mL, 2.0 mol total) was added through an additional funnel in 1.5 h to maintain the reflux. After addition, the reaction solution was heated at reflux with a water bath for an additional 30 min and then it was cooled to room temperature. This ether solution of **3.1** was used directly in the next reaction.

[00193] *Synthesis of compound 4.1:* To a suspension of CuI (39 g, 0.2 mol) in anhydrous THF (1000 mL) stirred with a mechanical stirrer at -78 °C in a 5 L three-necked flask was added R-epichlorohydrin (185 g, 2.0 mol) (additional funnel). After addition, the above ether solution of **3.1** was added via a cannula in 1 h. The mixture was stirred at -78 °C for 3.5 hrs, then quenched with saturated aqueous NH₄Cl (1500 mL). The organic layer was separated. The aqueous layer

was extracted with Et₂O (2000 mL). The combined organic phase was washed with brine, dried (over MgSO₄), and evaporated under vacuum. The crude product was purified by flash column chromatography (2.5 kg SiO₂, eluted with 0 – 10% EtOAc in Hexanes) to give 292 g of **4.1** (yield: 66%) as a light yellow oil.

[00194] *Synthesis of compound 5.1*: To a solution of **4.1** (292 g, 1.33 mol) in MeOH (600 mL) and THF (600 mL) was added aqueous NaOH (1.5M, 1000 mL) through an additional funnel at 0 °C. After addition the reaction mixture was stirred at room temperature for 2.5 hrs, TLC showed a major product, very minor amount of starting material (EtOAc: Hexanes = 1: 9, R_f = 0.6). THF and MeOH were removed by rotary evaporation under vacuum. The aqueous residue was extracted with Et₂O (600 mL X 3). The combined organic phase was washed with brine, dried over MgSO₄, and evaporated. The yellow oily residue was purified by column (SiO₂, 2.5 kg, eluted with 0-10% EtOAc in hexanes) to afford 205 g (84%) of pure **5.1**.

[00195] *Synthesis of compound 7.1*: Method A: To a solution of **6.1** (75 g, 0.25 mol) in a mixture of DCM (1000 mL) and MeOH (71 mL) that was being stirred at room temperature was added aqueous Na₂CO₃ (2.0 M, 135 mL). Organic layer was separated and the aqueous layer was extracted with DCM (250 mL X 2). The combined organic phase was dried over Na₂SO₄ and evaporated under vacuum. The residue was dissolved in MeOH (375 mL), then compound **5.1** (185 g, 1.0 mol) was added. The reaction mixture was stirred at room temperature for 4 days when MS and TLC showed mostly desired product. After concentration, the crude product was purified by flash column chromatography (2.0 kg SiO₂, eluted with 0 – 60% EtOAc in hexanes) to give **7.1** (131 g, 82%) as a pale viscous oil.

[00196] *Method B*: To a suspension of **8.1** (50 g, 0.2 mol) in MeOH (600 mL) was added DIPEA (45 mL), then compound **5.1** (150 g, 0.813 mol, 4.0 equiv) was added. The reaction mixture was stirred at room temperature for 7 days. Solvents were removed, the residue was purified by column (1.0 kg of SiO₂, eluted with 0 – 30% MeOH in EtOAc) to give **9.1** as a waxy solid (83 g, 67%). To a solution of **9.1** (81 g, 0.13 mol) in DMF (1000 mL) stirred at 0 °C was added HATU (50.1 g, 0.13 mol), followed by DIPEA (92 mL, 0.52 mol). The mixture was stirred at 0 °C for 40 min, and then MeOH (53.2 mL, 10.0 equiv) was added. The resulting mixture was stirred at room temperature overnight. It was then diluted with water (5000 mL) and extracted with EtOAc (500 mL X 4). The combined organic phase was washed with brine (600

mL X 3), dried over anhydrous MgSO_4 and evaporated under vacuum. The crude product was purified by column (1.0 kg of SiO_2 , eluted with 0 – 80% EtOAc in hexanes) to give **7.1** (69 g, 55% for 2 steps) as a pale viscous oil.

[00197] *Synthesis of compound 10.1:* To a solution of **7.1** (69 g, 0.11 mol) in DCM (200 mL) was added TFA (200 mL), the mixture was stirred at room temperature for 2 hrs, MS detection showed only desired product. All solvents were evaporated under vacuum to give 115 g of a brown colored oil-like product **10.1**, which was used in the next step without further purification.


[00198] *Synthesis of compound 12.1:* To a suspension of **11.1**, Boc-D-lysine (75 g, 0.305 mol) in MeOH (900 mL) were added DIPEA (68 mL) and **5.1** (196 g, 1.06 mol). The mixture was stirred at room temperature for 7 days. Volatiles were removed and the crude product was purified by column (2.5 kg of SiO_2 , eluted with 0 – 40% MeOH in EtOAc) to give 118 g (63%) of pure compound **12.1**.


[00199] *Synthesis of compound 13.1:* To a solution of **12.1** (67.5 g, 0.11 mol) in DMF (600 mL, warm up to 50 °C for 30 min to obtain a homogeneous solution) that was cooled with an ice-bath were added HATU (50 g, 0.12 mol) and DIPEA (95 mL, 0.55 mol). The resulting mixture was stirred at 0 °C for 45 min, then compound **10.1** (115 g, obtained above) in DMF (400 mL) was added using an additional funnel. The mixture was stirred at room temperature overnight. Ether (1000 mL) was added, followed by water (1000 mL). The organic phase was separated; the aqueous was extracted with ether (250 mL X 2). The combined organic phase was washed with water, dried over anhydrous MgSO_4 , filtered and concentrated. The residue was purified by column (1.0 kg of SiO_2 , eluted with 0 – 20% MeOH in EtOAc) to give 94.2 g (76%) of compound **13.1**.

[00200] *Synthesis of the compound of Formula I.a.i (i.e., R4-SR-cKK-E12):* To a solution of **13.1** (94 g, 0.084 mol) in DCM (300 mL) was added TFA (300 mL). The mixture was stirred at room temperature for 2 hrs. MS detection showed complete reaction. Solvents were evaporated under vacuum. The residue was dissolved in DCM (500 mL) and washed with aqueous Na_2CO_3 (1.0 M, 500 mL). The aqueous wash was back extracted with DCM (100 mL). The combined organic phase was dried over anhydrous Na_2SO_4 and evaporated. The residue was dissolved in MeOH (1500 mL) and cooled with an ice-bath. Aqueous NH_4OH (28%, 80 mL) was added through additional funnel. The reaction mixture was allowed to slowly warm up to room

temperature and stirred at ambient temperature for 2 days. Volatiles were evaporated under vacuum. The crude product was purified by column (1.0 kg of SiO₂, eluted with solvents: 1% NH₄OH, 4-9% MeOH, 95-90% EtOAc) to give 34 g of pure **R4-SR- cKK-E12** and 22 g of impure **R4-SR- cKK-E12**. The impure **R4-SR- cKK-E12** was re-purified by column to give 12 g of pure **R4-SR- cKK-E12**. Thus, a total 46 g (55%) of pure **R4-SR- cKK-E12** (gummy solid) was obtained.

Example 4 - The compound of *Formula I.a.ii* (i.e., *S4-SR-cKK-E12*)



Scheme 5: Synthesis of the compound of Formula I.a.ii (i.e., S4-SR-cKK-E12)

[00201] *Synthesis of compound 3.1:* To a suspension of Mg (30 g, 1.25 mol) in anhydrous Et₂O (600 mL) was added one crystal of iodine, followed by addition of 1-bromononane (30 mL). The reaction was initiated and the solution began to reflux after heating the reaction flask with a water bath. The remaining 1-bromononane (161 mL, 2.0 mol total) was added through an additional funnel in 1.5 h to maintain the reflux. After addition, the reaction solution was heated at reflux with a water bath for an additional 30 min and then it was cooled to room temperature. This ether solution of **3.1** was used directly in the next reaction.

[00202] *Synthesis of compound 4.2:* To a suspension of CuI (19 g, 0.1 mol) in anhydrous THF (1000 mL) that was being stirred with a mechanical stirrer at -78 °C in a 5 L three-necked flask was added S-epichlorohydrin (92 g, 1.0 mol) using an additional funnel. After addition, the above ether solution of **3.2** was added via a cannula in 1 h. The mixture was stirred at -78 °C for 3.5 hrs, then quenched with saturated aqueous NH₄Cl (400 mL). The organic layer was

separated. The aqueous layer was extracted with Et₂O (1000 mL). The combined organic phase was washed with brine, dried (over MgSO₄), and evaporated under vacuum. The crude product was purified by flash column chromatography (2.5 kg SiO₂, eluted with 0 – 10% EtOAc in hexanes) to give 111.6 g of **4.2** (yield: 66%) as a light yellow oil.

[00203] *Synthesis of compound 5.2:* To a solution of **4.2** (111.3 g, 0.506 mol) in MeOH (230 mL) and THF (230 mL) was added aqueous NaOH (1.5M, 395 mL) using an additional funnel at 0 °C. After addition the reaction mixture was stirred at room temperature for 2.5 hrs, TLC showed a major product and very minor amount of starting material (EtOAc: Hexanes = 1: 9, R_f = 0.6). THF and MeOH were removed by rotary evaporation under vacuum. The aqueous residue was extracted with Et₂O (200 mL X 3). The combined organic phase was washed with brine, dried over MgSO₄, and evaporated. The yellow oily residue was purified by column (SiO₂, 1.0 kg, eluted with 0-10% EtOAc in hexanes) to afford 81 g (87%) of pure **5.2**.



[00204] *Synthesis of compound 7.2:* To a solution of **6.1** (13 g, 0.044 mol) in a mixture of DCM (100 mL) and MeOH (10 mL) that was being stirred at room temperature was added aqueous Na₂CO₃ (2.0 M, 25 mL). Organic layer was separated and the aqueous layer was extracted with DCM (250 mL X 2). The combined organic phase was dried over Na₂SO₄ and evaporated under vacuum. The residue was dissolved in MeOH (60 mL), then compound **5.2** (32 g, 0.174 mol) was added. The reaction mixture was stirred at room temperature for 4 days when MS and TLC showed mostly desired product. After concentration, the crude product was purified by flash column chromatography (600 g SiO₂, eluted with 0 – 60% EtOAc in hexanes) to give **7.2** (23 g, 85%) as a pale viscous oil.

[00205] *Synthesis of compound 8.2:* To a solution of **7.2** (23 g, 0.0366 mol) in DCM (60 mL) was added TFA (60 mL), the mixture was stirred at room temperature for 2 hrs, MS detection showed only desired product. All solvents were evaporated under vacuum to give 40 g of a brown colored oil-like product **8.2**, which was used in the next step without further purification.

[00206] *Synthesis of compound 10.2:* To a suspension of **11.1**, Boc-D-lysine (14 g, 0.057 mol) in MeOH (900 mL) were added TEA (11.6 mL) and **5.2** (42 g, 0.228 mol). The mixture was stirred at room temperature for 7 days. Volatiles were removed and the crude product was purified by column (1.0 kg of SiO₂, eluted with 0 – 40% MeOH in EtOAc) to give 24 g (70%) of pure compound **10.2**.

[00207] *Synthesis of compound 11.2:* To a solution of **10.2** (9.1 g, 14.82 mmol) in DMF (120 mL) that was being cooled with an ice bath were added HATU (8.4 g, 22.23 mol) and DIPEA (25 mL, 148.2 mmol). The mixture was stirred at 0 ° C for 45 min, then compound **8.2** in DMF (80 mL) was added using an additional funnel. The resulting mixture was stirred at room temperature overnight. MS detection showed no starting material. Ether (1000 mL) was added, followed by water (1000 mL). The organic phase was separated, the aqueous was extracted with ether (200 mL X 2). The combined organic phase was washed with brine, dried with anhydrous MgSO₄, filtered and concentrated. The residue was purified by column (330 g of SiO₂, eluted with 0 – 20% MeOH in EtOAc) to give 10.6 g of desired compound **11.2**.

[00208] *Synthesis of the compound of Formula I.a.ii (i.e., S4-SR-cKK-E12):* To a solution of **11.2** (10.6 g, 0.084 mol) in DCM (30 mL) was added TFA (30 mL). The mixture was stirred at room temperature for 2 hrs. MS detection showed complete reaction. Solvents were evaporated under vacuum. The residue was dissolved in DCM (150 mL) and washed with aqueous Na₂CO₃ (1.0 M, 200 mL). The aqueous wash was back extracted with DCM (100 mL). The combined organic phase was dried over anhydrous NaSO₄ and evaporated. The residue was dissolved in MeOH (200 mL) and cooled with an ice-bath. Aqueous NH₄OH (28%, 10 mL) was added using an additional funnel. The reaction mixture was allowed to slowly warm up to room temperature and stirred at ambient temperature for 2 days. Volatiles were evaporated under vacuum. The crude product was purified by column (600 g of SiO₂, eluted with solvents: 1% NH₄OH, 4-9% MeOH, 95-90% EtOAc) give 5.1 g of pure **S4-SR-cKK-E12**.


Example 5 - The compound of **Formula I.b.1.i** (i.e., *R4-SS-cKK-E12*)**Scheme 6:** Synthesis of the compound of **Formula I.b.1.i** (i.e., *R4-SS-cKK-E12*).


[00209] Synthesis of compound **3.3** (*N*²-(*tert*-butoxycarbonyl)-*N*⁶,*N*⁶-bis((*R*)-2-hydroxydodecyl)-L-lysine): A mixture of *R*-epoxide (**5.1**, 46 g, 250 mmol), Boc-L-Lysine **8.1** (15 g, 61 mmol), and diisopropylethylamine (11 ml) in methanol (80 ml) was stirred at room temperature for 3 d. Volatiles were removed and the yellow oily residue was purified by

chromatography on silica gel (330 g) eluting with EtOAc/MeOH (100/0 to 70/30, 20 min) to give product **3.3** as a white solid (9.7 g, 26%).

[00210] *Synthesis of the compound of **Formula I.b.1.i.** (i.e., **R4-SS-cKK-E12**; ((3*S*,6*S*)-3,6-bis(4-(bis((*S*)-2-hydroxydodecyl)amino) butyl)piperazine-2,5-dione))*: To a solution of **3.3** (7.4 g, 12 mmol) and NHS (1.38 g, 12 mmol) in DCM (280 ml) was added DCC (2.97 g, 14.4 mmol). The reaction mixture was stirred at room temperature for 1.5 h. The solvent was then removed and the residue (crude **4.3**) was dissolved in TFA (30 ml). The resulting mixture was stirred at room temperature for 1 h. TFA was then removed and DCM (30 ml) was added to the residue, and co-evaporated to remove residual TFA. The crude **5.3** was dissolved in DCM (30 mL) and added to anhydrous pyridine (480 mL) at 0 ° C under N₂. The resulting mixture was stirred at room temperature overnight. Pyridine was then removed and the residue was diluted with diethyl ether (300 mL). The white solid formed was removed by filtration. The filtrate was washed with aqueous Na₂CO₃ (1M, 150 ml) and brine (150 ml), dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography five times (one 330 g column followed by four 80 g column) eluting with 3%NH₄OH/7%MeOH/90%EtOAc to give 1.05 g of pure **R4-SS-cKK-E12** as pale gum. 1.0 g of **R4-SS-cKK-E12**.

Example 6 - *The compound of **Formula I.b.1.ii** (i.e., **S4-SS-cKK-E12**)*




Scheme 7: Synthesis of compound of Formula I.b.1.ii (i.e., S4-SS-cKK-E12)


[00211] *Synthesis of compound 3 (N²-(tert-butoxycarbonyl)-N⁶,N⁶-bis((S)-2-hydroxydodecyl)-L-lysine):* A mixture of S-epoxide (**5.2**, 46 g, 250 mmol), Boc-L-Lysine **8.1** (15 g, 61 mmol) and diisopropylethylamine (11 ml) in methanol (80 ml) was stirred at room temperature for 8 d. Volatiles were removed and the yellow oily residue was purified by chromatography on silica gel (330 g) eluting with EtOAc/MeOH (100/0 to 70/30, 20 min) to give product **3.4** as a white solid (22 g, 59%).

[00212] *Synthesis of the compound of Formula I.b.1.ii (i.e., S4-SS-cKK-E12; ((3S,6S)-3,6-bis(4-(bis((S)-2-hydroxydodecyl)amino)butyl)piperazine-2,5-dione):* To a solution of **3.4** (24.5 g, 40 mmol) and NHS (4.6 g, 40 mmol) in DCM (280 ml) was added DCC (9.9 g, 48 mmol). The reaction mixture was stirred at room temperature for 3 h. The solvent was then removed and the residue (crude **4.4**) was dissolved in TFA (100 ml) at 0 °C. The resulting mixture was allowed to warm up to room temperature and stirred for 45 min. TFA was then removed and DCM (120 ml) was added to the residue, and then co-evaporated to remove residual TFA. The crude **5.4** was dissolved in anhydrous pyridine (1.6 L) at 0 °C under N₂, and the resulting mixture was stirred at

room temperature overnight. The pyridine was then removed by rotavap, and the residue was diluted with diethyl ether (1 L). The white solid formed was removed by filtration and washed with diethyl ether (200 ml). The filtrate was washed with aqueous Na_2CO_3 (1M, 500 ml) and brine (500 ml), dried over Na_2SO_4 , and concentrated. The residue was purified by column chromatography on silica gel (330 g) eluting with 0-50% (3% NH_4OH /7% MeOH /90% EtOAc)/ EtOAc to give 6.8 g TLC pure **S4-SS-cKK-E12** and 7.1 g slightly impure **S4-SS-cKK-E12**. The 6.8 g (6.8 mmol) TLC pure **S4-SS-cKK-E12** was dissolved in 120 ml of ethyl acetate. Boc_2O (0.22 g, 1.0 mmol) was added. The mixture was stirred at room temperature for 2 h. The solvent was removed and the residue was purified by column chromatography on silica gel (120 g) eluting with 0-50% (3% NH_4OH /7% MeOH /90% EtOAc)/ EtOAc to give 5.7 g (84%) of pure product **S4-SS-cKK-E12** which was free of an amine side product.

Example 7 - The compound of **Formula I.b.2.i** (i.e., **R4-RR-cKK-E12**)



Scheme 8: Synthesis of the compound of Formula I.b.2.i (i.e., R4-RR-cKK-E12)

[00213] *Synthesis of compound 3.1:* To magnesium (60 g) suspended in anhydrous Et_2O (0.9 L) was added 1-bromononane **2.1** (20 mL), followed by addition of a catalytic amount of iodine (50 mg). The resulting mixture was heated with a hot water bath until reaction of Mg with **2.1** started. The bath was removed and the remaining 1-bromononane (379.1 mL) was added to maintain a gentle reflux. After addition of **2.1** the reflux was maintained by a hot water bath for another 30 min. The resulting Grignard solution of **3.1** was cooled and used directly in next step.

[00214] *Synthesis of compound 4.1:* To CuI (38.1 g) suspended in THF (1.5 L) at -78°C was added $\text{R-(-)-epichlorohydrin}$ (156.8 mL). Then the above Grignard solution of **3.1** was added via a cannula while the reaction temperature was maintained at $<-65^\circ\text{C}$. The resulting reaction mixture was stirred at -78°C for an additional 3 hour. Then, saturated aqueous ammonium chloride solution (0.8 L) was added carefully, followed by addition of water (1.0 L). The resulting mixture was stirred and allowed to warm up to room temperature. The organic layer was separated, and the aqueous layer was extracted with Et_2O (0.5 L x 2). The organic layer was combined with the Et_2O extracts and the resulting solution was washed with brine (0.5 L x 2), and dried over anhydrous magnesium sulfate. Solvents were evaporated under vacuum, and the

resulting residue was purified on a silica gel column (hexanes to 20% EtOAc/hexanes) to provide **4.1** (243.5 g, 55%) as a yellow oil.


[00215] *Synthesis of compound 5.1:* To a solution of **4.1** (243.49 g) in 1:2.6 MeOH-THF (3.6 L) stirred at 0 °C was slowly added aqueous NaOH solution (1.5 M, 0.89 L, 1.33 mole). The resulting mixture was allowed to warm up and stirred at room temperature for 3 h. TLC analysis showed complete disappearance of **4.1**. Organic solvents were evaporated, and the aqueous layer was extracted with Et₂O (1 L + 500 mL x 2). The organic extracts were combined, washed with brine (600 mL), and dried over anhydrous magnesium sulfate. Solvents were evaporated under vacuum to give a residue which was purified on silica gel column (hexanes to 10% EtOAc/hexanes) to provide the epoxide **5.1** (193.7 g, 95%) as a light yellow oil.


[00216] *Synthesis of compound 12.1:* A mixture of *N*-Boc-D-Lysine **11.1** (49.2 g, 0.2 mole) and the epoxide **5.1** (147.2 g, 0.8 mole) in MeOH (1.04 L) was stirred at room temperature. DIPEA (51.9 g, 0.4 mole) was added. The resulting mixture was then stirred for 8 days, and then concentrated to dryness. The residue was purified on a silica gel column (2 kg, MeOH/DCM, 0 to 10%) to give 49.2 g of mostly pure **12.1** (MZ-550-180) and 58.3 g of impure **12.1**, which was purified by a second column (1.5 kg, MeOH/EtOAc, 10 to 40%) to give 41.4 g of mostly pure **12.1**. The two mostly pure batches were combined and stirred with EtOAc (0.5 L) for 3 h. The mixture was filtered to give 41.4 g pure **12.1** as a white solid. The filtrate was concentrated to dryness. The residue was stirred with EtOAc (0.1 L) for 1h and filtered to give 10.6 g of pure **12.1**. The filtrate was concentrated to dryness and the residue was purified on a silica gel column (330 g, MeOH/EtOAc, 10 to 40%) to give a third batch of 26.9 g pure **12.1**. A total of 78.9 g of pure **12.1** was obtained. Yield: 64%

[00217] *Preparation of the compound of Formula I.b.2.i (i.e., R4-RR-cKK-E12):* Batch 1: To a solution of **12.1** (6.14 g, 10 mmol) and *N*-hydroxysuccinimide (1.15 g, 10 mmol) in DCM (70 mL) was added DCC (2.47 g, 12 mmol). The resulting mixture was stirred at room temperature for 3 h. Volatiles were evaporated under vacuum to give a residue (NHS ester **6.5**), which was dissolved in TFA (25 mL) and stirred for 0.5 h. TFA was removed under vacuum, and the residue (compound **7.5**) was cooled to 0 °C. Pyridine (anhydrous, 400 mL) was added, and the reaction mixture was stirred at room temperature for 2 h. Pyridine was removed under vacuum, and the residue was suspended in Et₂O (300 mL). The solid was removed by filtration. The filtrate was

washed with 1 M Na₂CO₃ aqueous solution (150 mL) and brine (150 mL), dried over anhydrous magnesium sulfate, and concentrated to dryness. The residue was separated by column chromatograph (80 g, 7:3 MeOH-NH₃-H₂O (4 x with EtOAc)/EtOAc, 0 to 50%) to give **R4-RR-cKK-E12** as gummy solid (2.22 g). Multiple precipitations and triturations from EtOAc gave pure **R4-RR-cKK-E12** (0.46 g) as a gum.

Example 8 - The compound of *Formula I.b.2.ii* (i.e., *S4-RR-cKK-E12*)



Scheme 9: Synthesis of the compound of Formula I.b.2.ii (i.e., S4-RR-cKK-E12)

[00218] *Synthesis of compound 3.1:* To magnesium (60 g) suspended in anhydrous Et₂O (0.9 L) was added 1-bromononane **2.1** (20 mL), followed by addition of a catalytic amount of iodine (50 mg). The resulting mixture was heated with a hot water bath until reaction of Mg with **2.1** started. The bath was removed and the remaining 1-bromononane (379.1 mL) was added to maintain a gentle reflux. After addition of **2.1** the reflux was maintained by a hot water bath for another 30 min. The resulting Grignard solution of **3.1** was cooled and used directly in next step.

[00219] *Synthesis of compound 4.2:* To CuI (38.1 g) suspended in THF (1.5 L) at -78 °C was added S-(-)-epichlorohydrin (156.8 mL). Then the above Grignard solution of **3.1** was added via a cannula while the reaction temperature was maintained at <-65 °C. The resulting reaction mixture was stirred at -78 °C for an additional 3 hour. Then, saturated aqueous ammonium chloride solution (0.8 L) was added carefully, followed by addition of water (1.0 L). The resulting mixture was stirred and allowed to warm up to room temperature. The organic layer was separated, and the aqueous layer was extracted with Et₂O (0.5 L x 2). The organic layer was

combined with the Et₂O extracts and the resulting solution was washed with brine (0.5 L x 2), and dried over anhydrous magnesium sulfate. Solvents were evaporated under vacuum, and the residue was purified on a silica gel column (hexanes to 20% EtOAc/hexanes) to provide **4.2** (250.8 g, 57%) as a light yellow oil.

[00220] *Synthesis of compound 5.2:* To a solution of **4.2** (250.8 g) in 1:2.6 MeOH-THF (3.9 L) stirred at 0 °C was added slowly aqueous NaOH solution (1.5 M, 1.36 mole, 0.90 L). The resulting mixture was allowed to warm up and stirred at room temperature for 3 h. TLC analysis showed complete disappearance of **4.2**. Organic solvents were evaporated, and the aqueous layer was extracted with Et₂O (1 L + 500 mL x 2). The organic extracts were combined, washed with brine (600 mL), and dried over anhydrous magnesium sulfate. Solvents were evaporated under vacuum to give a residue which was purified on silica gel column (hexanes to 10% EtOAc/hexanes) to provide **5.2** (195.4 g, 93%) as a light yellow oil.

[00221] A mixture of *N*-Boc-D-Lysine **11.1** (49.2 g, 0.2 mole) and the epoxide **5.2** (147.2 g, 0.8 mole) in MeOH (1.04 L) was stirred at room temperature. DIPEA (51.9 g, 0.4 mole) was added. The resulting mixture was then stirred for 8 days, and then concentrated to dryness. The resulting residue was purified on a silica gel column (2 kg, MeOH/EtOAc, 10 to 30%) to give **10.2** (79.9 g, 65%) as a white solid.

[00222] *Preparation of the compound of Formula I.b.2.ii (i.e., S4-RR-cKK-E12)* To a solution of **10.2** (61.4 g, 100 mmol) and *N*-hydroxysuccinimide (11.5 g, 100 mmol) in DCM (800 mL) was added DCC (24.7 g, 120 mmol). The resulting mixture was stirred at room temperature for 4 h. Volatiles were evaporated under vacuum to give a residue (NHS ester **6.6**), which was dissolved in TFA (25 mL) and stirred for 40 min. TFA was removed under vacuum, and the residue (compound **7.6**) was cooled to 0 °C. Pyridine (anhydrous, 3.5 L) was added, and the reaction mixture was stirred at room temperature for 19 h. Pyridine was removed under vacuum, and the residue was suspended in Et₂O (3.0 L). The solid was removed by filtration. The filtrate was washed with 1 M aqueous Na₂CO₃ solution (1.0 L) and brine (1.0 L), dried over anhydrous magnesium sulfate, and concentrated to dryness. The residue was purified by column chromatograph (4 x 330 g silica gel column eluting with (7%MeOH/3%NH₃-H₂O/90%EtOAc)/EtOAc, 0 to 50%) to provide **S4-RR-cKK-E12** as a gummy solid (15.9 g, 16%)

Example 9 - Formulations

[00223] The formulations described herein consisted of a multi-component lipid mixture of varying ratios employing one or more cationic lipids, helper lipids and PEGylated lipids designed to encapsulate various nucleic acid-based materials. The cationic lipid utilized throughout is the compound of formula I (3,6-bis(4-(bis(2-hydroxydodecyl)amino)butyl)piperazine-2,5-dione). Helper lipids can include (but not exclusively) DSPC (1,2-distearoyl-sn-glycero-3-phosphocholine), DPPC (1,2-dipalmitoyl-sn-glycero-3-phosphocholine), DOPE (1,2-dioleoyl-sn-glycero-3-phosphoethanolamine), DOPC (1,2-dioleoyl-sn-glycero-3-phosphotidylcholine) DPPE (1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine), DMPE (1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine), DOPG (1,2-dioleoyl-sn-glycero-3-phospho-(1'-*rac*-glycerol)), cholesterol, etc. The PEGylated lipids can include (but not exclusively) a poly(ethylene) glycol chain of up to 5 kDa in length covalently attached to a lipid with alkyl chain(s) of C₆-C₂₀ length.

Messenger RNA Material

[00224] Human Factor IX (FIX), codon-optimized Firefly Luciferase (FFL) and codon-optimized human argininosuccinate synthetase (ASS1) messenger RNA were synthesized by *in vitro* transcription from a plasmid DNA template encoding the gene, which was followed by the addition of a 5' cap structure (Cap 1) (Fechter, P.; Brownlee, G.G. "Recognition of mRNA cap structures by viral and cellular proteins" *J. Gen. Virology* **2005**, 86, 1239-1249) and a 3' poly(A) tail of approximately 250 nucleotides in length as determined by gel electrophoresis. 5' and 3' untranslated regions present in each mRNA product are represented as X and Y, respectively and defined as stated (*vide infra*).

Codon-Optimized Human Argininosuccinate Synthetase (ASS1) mRNA:

XAUGAGCAGCAAGGGCAGCGUGGUGCUGGCCUACAGCGGCGGCCUGGACACCAGC
 UGCAUCCUGGUGUGGCUGAAGGAGCAGGGCUACGACGUGAUCGCCUACCUGGCCA
 ACAUCGGCCAGAAGGAGGACUUCGAGGAGGCCCCGCAAGAAGGCCCUAGAAGCUGGG
 CGCCAAGAAGGUGUUCUUCGAGGACGUGAGCCGCGAGUUCGUGGAGGAGUUCAU
 CUGGCCCCGCCAUCCAGAGCAGCGCCCCUGUACGAGGACCGCUACCUGCUGGGGCACC
 AGCCUGGCCCCGCCCCUGCAUCGCCCCGCAAGCAGGUGGAGAUCGCCCAGCGCGAGG
 GCGCCAAGUACGUGAGCCACGGCGCCACCGGCAAGGGCAACGACCAGGUGCGCUU
 CGAGCUGAGCUGCUACAGCCUGGCCCCCAGAUCAAGGUGAUCGCCCCCUGGCGC
 AUGCCCGAGUUCUACAACCGCUUCAAGGGCCGCAACGACCUGAUGGAGUACGCCA
 AGCAGCACGGCAUCCCCAUCCCCGUGACCCCCAAGAACCCCUGGAGCAUGGACGA

GAACCUGAUGCACAUCAGCUACGAGGCCGGCAUCCUGGAGAACCCCAAGAACCAG
 GGGGGGGGGGGCCUGUACACCAAGACCCAGGACCCCGCCAAGGGCCCCAACACCCC
 CGACAUCCUGGAGAUUCGAGUUAAGAAGGGGCGUGCCCGUGAAGGUGACCAACGU
 GAAGGACGGCACCACCCACCAGACCAGCCUGGAGCUGUUAUGUACCUGAACGAG
 GUGGCCGGCAAGCACGGCGUGGGGCCGCAUCGACAUCGUGGAGAACCGCUUCAUCG
 GCAUGAAGAGCCGCGGCAUCUACGAGACCCCGCCGGCACCAUCCUGUACCACGC
 CCACCUGGACAUCGAGGCCUUCACCAUGGACCGCGAGGUGCGCAAGAUAAGCAG
 GGCCUGGGGCCUGAAGUUCGCCGAGCUGGUGUACACCGGCUUCUGGCACAGCCCCG
 AGUGCGAGUUCGUGCGCCACUGCAUCGCCAAGAGCCAGGAGCGCGUGGAGGGCAA
 GGUGCAGGUGAGCGUGCUGAAGGGGCCAGGUGUACAUCUGGGCCGCGAGAGCCCC
 CUGAGCCUGUACAACGAGGAGCUGGUGAGCAUGAACGUGCAGGGGCGACUACGAG
 CCCACCGACGCCACCGGCUUCAUCAACAACAGCCUGCGCCUGAAGGAGUACC
 ACCGCCUGCAGAGCAAGGUGACCGCCAAGUGAY

5' and 3' UTR Sequences

X =

GGACAGAUCGCCUGGAGACGCCAUCCACGCUGUUUUGACCUCCAUAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGGCCGGGAACGGUGCAUUGGAACGCGGAUUCCTCCG
 UGCCAAGAGUGACUCACCGUCCUUGACACG

Y =

CGGGUGGCAUCCCUGUGACCCCUCCCCAGUGCCUCUCCUGGGCCUGGAAGUUGCC
 ACUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGUUGCAUC

Formulation Protocol

[00225] Aliquots of 50 mg/mL ethanolic solutions of one or more compounds of formula I, DOPE, Cholesterol and DMG-PEG2K were mixed and diluted with ethanol to 3 mL final volume. Separately, an aqueous buffered solution (10 mM citrate/150 mM NaCl, pH 4.5) of ASS1 mRNA was prepared from a 1 mg/mL stock. The lipid solution was injected rapidly into the aqueous mRNA solution and shaken to yield a final suspension in 20% ethanol. The resulting nanoparticle suspension was filtered, diafiltrated with 1x PBS (pH 7.4), concentrated and stored at 2-8°C. Final concentration = 0.64 mg/mL ASS1 mRNA (encapsulated). $Z_{ave} = 78$ nm ($Dv_{(50)} = 46$ nm; $Dv_{(90)} = 96$ nm).

Example 10 - *Analysis of ASS1 protein produced via intravenously delivered ASS1 mRNA-loaded nanoparticles:*

Injection Protocol

[00226] All studies were performed using male CD-1 mice of approximately 6-8 weeks of age at the beginning of each experiment. Samples were introduced by a single bolus tail-vein injection of an equivalent total dose of 1.0 mg/kg (or otherwise specified) of encapsulated ASS1 mRNA. Mice were sacrificed and perfused with saline at the designated time points.

Isolation of organ tissues for analysis

[00227] The liver, spleen, kidney and heart of each mouse was harvested, apportioned into separate parts, and stored in either 10% neutral buffered formalin or snap-frozen and stored at -80°C for analysis.

Isolation of plasma for analysis

[00228] All animals were euthanized by CO₂ asphyxiation at designated time points post dose administration ($\pm$ 5%) followed by thoracotomy and terminal cardiac blood collection. Whole blood (maximal obtainable volume) will be collected via cardiac puncture on euthanized animals into serum separator tubes, allowed to clot at room temperature for at least 30 minutes, centrifuged at 22°C $\pm$ 5°C at 9300 g for 10 minutes, and the serum will be extracted. For interim blood collections, approximately 40-50 μ L of whole blood will be collected via facial vein puncture or tail snip. Samples collected from non treatment animals were used as a baseline ASS1 levels for comparison to study animals.

Enzyme-Linked Immunosorbent Assay (ELISA) Analysis

[00229] *Human ASS1 ELISA:* Standard ELISA procedures were followed employing mouse anti-ASS1 2D1-2E12 IgG as the capture antibody with rabbit anti-ASS1 #3285 IgG as the secondary (detection) antibody (Shire Human Genetic Therapies). Horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG was used for activation of the 3,3',5,5'-tetramethylbenzidine (TMB) substrate solution. The reaction was quenched using 2N H₂SO₄ after 20 minutes. Detection was monitored via absorption (450 nm) on a Molecular Device SpectraMax instrument. Untreated mouse serum and organs and human ASS1 protein were used as negative and positive controls, respectively.

Example 11 - In Vivo Human ASS1 Protein Production

[00230] The production of human ASS1 protein via codon-optimized hASS1 mRNA-loaded lipid nanoparticles, comprising compounds of formula I, was tested in CD-1 mice as a single,

bolus intravenous injection, as described above. FIG. 1 depicts the amount of human ASS1 protein detected via ELISA when treating mice human ASS1 mRNA-loaded lipid nanoparticles, with various racemic and chiral compounds of formula I, at 1.0 mg/kg doses. The mice were sacrificed twenty-four hours post-injection and organs, such as livers, were harvested.

Example 12 – Toxicity Studies

[00231] Expression levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) were measured for various racemic and chiral compounds of formula I. Increased expression levels of AST and/or ALT generally caused by agents that cause liver toxicity. The chiral compounds of formula I generally yielded lower expression levels of ALT and/or AST, *i.e.*, correlating to lower liver toxicity issues, compared to stereochemically non-enriched, or stereochemically less enriched, compositions of the same lipid. See Tables 1 and 2 below.

Table 1.

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
Racemic Mixture	190 ± 43	212 ± 54	471 ± 309
	201 ± 89	403 ± 42	937 ± 337
	207 ± 84	425 ± 169	497 ± 213
	344 ± 57	555 ± 122	1387 ± 593
	426 ± 112	757 ± 158	1509 ± 598
	457 ± 274	728 ± 126	910 ± 327
	503 ± 201	653 ± 133	1010 ± 154
	618 ± 503	638 ± 273	209 ± 169

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
S ₄ with Racemic Lysine Core	170 ± 40	132 ± 44	375 ± 244
	155 ± 57	157 ± 38	674 ± 147

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
R ₄ with Racemic Lysine Core	188 ± 22	265 ± 122	823 ± 215
	236 ± 163	237 ± 139	568 ± 248

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
-----------	-----------	-----------	----------------------------

Racemic –OH with SS Lysine Core	378 ± 58	622 ± 76	117 ± 80
	618 ± 503	638 ± 273	209 ± 169

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
S ₄ -S,S-cKKE12	226 ± 71	384 ± 233	1121 ± 468

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
R ₄ -S,S-cKKE12	175 ± 102	144 ± 35	449 ± 105

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
S ₄ -S,R-cKKE12	190 ± 75	193 ± 71	2303 ± 491

Structure	ALT (U/L)	AST (U/L)	ASS1 (ng/mg Total Protein)
R ₄ -S,R-cKKE12	75 ± 13	82 ± 12	264 ± 317
	86 ± 27	119 ± 32	1369 ± 233
	94 ± 22	88 ± 16	467 ± 149
	59 ± 13	73 ± 18	401 ± 137
	139 ± 28	177 ± 73	1182 ± 150
	180 ± 19	141 ± 25	750 ± 324
	269 ± 80	424 ± 156	2790 ± 464
	123 ± 39	124 ± 22	1113 ± 35
	60 ± 4	49 ± 5	846 ± 226
	70 ± 10	78 ± 24	1082 ± 189


Table 2.

cKK-E12 of a single intravenous dose. 24 hours post-formulation used for screening was cKK-E12:DOPE:Chol:DMG-PEG2K	Structure Assignment	ALT	AST
Lot #1	‘Racemic’ Mixture	885 ± 489	982 ± 350
		207 ± 84	425 ± 169
		504 ± 317	657 ± 176
		503 ± 201	653 ± 133
Lot #2		365 ± 152	604 ± 136
		401 ± 265	586 ± 193
Lot #3		197 ± 50	309 ± 33
Lot #1	S ₄ -SS	226 ± 71	384 ± 233
Lot #1	R ₄ -SS	175 ± 102	144 ± 35
Lot #1	S ₄ -RR	152 ± 9	180 ± 42
Lot #1	R ₄ -RR	136 ± 34	194 ± 80
Lot #1	S ₄ -RS/SR	143 ± 29	240 ± 98
		189 ± 47	203 ± 87
Lot #2		190 ± 75	193 ± 71
Lot #1	R ₄ -RS/SR	86 ± 27	119 ± 32
		75 ± 13	82 ±12
		76 ± 4	79 ± 4
		94 ± 22	88 ± 16
Lot #2		269 ± 80	424 ± 156
		139 ± 28	177 ± 73
		180 ± 19	141 ± 25
		91 ±13	98 ±18
Lot #3		125 ± 47	104 ± 27
Lot #4		94 ± 24	91 ± 14
Lot #5		60 ± 4	49 ± 5
Lot #6		70 ± 10	78 ± 24
Lot #7		308 ± 115	354 ± 128
		123 ± 39	124 ± 22


[00232] While several embodiments of the present invention have been described and illustrated herein, those of ordinary skill in the art will readily envision a variety of other means and/or structures for performing the functions and/or obtaining the results and/or one or more of the advantages described herein, and each of such variations and/or modifications is deemed to be within the scope of the present invention. More generally, those skilled in the art will readily appreciate that all parameters, dimensions, materials, and configurations described herein are meant to be exemplary and that the actual parameters, dimensions, materials, and/or configurations will depend upon the specific application or applications for which the teachings of the present invention is/are used. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. It is, therefore, to be understood that the foregoing embodiments are presented by way of example only and that, within the scope of the appended claims and equivalents thereto, the invention may be practiced otherwise than as specifically described and claimed. The present invention is directed to each individual feature, system, article, material, kit, and/or method described herein. In addition, any combination of two or more such features, systems, articles, materials, kits, and/or methods, if such features, systems, articles, materials, kits, and/or methods are not mutually inconsistent, is included within the scope of the present invention.

CLAIMS

1. A composition comprising one or more chemical entities of formula **I**, each of which is a compound of formula **I**:




- a pharmaceutically acceptable salt thereof, a solvate thereof, or a solvate of a pharmaceutically acceptable salt thereof,
the composition characterized in that greater than a first threshold amount of the total amount of chemical entities of formula **I** in the composition:
are chemical entities of formula **I.a**:




wherein the first threshold amount is 50%; or

- are chemical entities of formula **I.b.1**:



wherein the first threshold amount is 25%; or


- are chemical entities of formula **I.b.2**:



I.b.2,

wherein the first threshold amount is 25%.

2. The composition of claim 1, wherein the first threshold amount is 70%.
3. The composition of claim 1, wherein the first threshold amount is 80%.
4. The composition of claim 1, wherein the first threshold amount is 95%.
5. The composition of any of claims 1-4, wherein greater than the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.a**.
6. The composition of claim 5, characterized in that greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition is a chemical entity of formula **I.a.i**:




I.a.i,

wherein the second threshold amount is 50%.

7. The composition of claim 6, wherein the second threshold amount is 70%.
8. The composition of claim 6, wherein the second threshold amount is 80%.


9. The composition of claim 6, wherein the second threshold amount is 90%.
10. The composition of claim 6, wherein the second threshold amount is 95%.
11. The composition of claim 5, characterized in that greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.a** in the composition is a chemical entity of formula **I.a.ii**:



wherein the second threshold amount is 50%.

12. The composition of claim 11, wherein the second threshold amount is 70%.
13. The composition of claim 11, wherein the second threshold amount is 80%.
14. The composition of claim 11, wherein the second threshold amount is 90%.
15. The composition of claim 11, wherein the second threshold amount is 95%.
16. The composition of any of claims 1-4, wherein greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.1**.

17. The composition of claim 16, characterized in that greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition is a chemical entity of formula **I.b.1.i**:



wherein the second threshold amount is 50%.


18. The composition of claim 17, wherein the second threshold amount is 70%.

19. The composition of claim 17, wherein the second threshold amount is 80%.

20. The composition of claim 17, wherein the second threshold amount is 90%.

21. The composition of claim 17, wherein the second threshold amount is 95%.


22. The composition of claim 16, characterized in that greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.1** in the composition is a chemical entity of formula **I.b.1.ii**:



wherein the second threshold amount is 50%.

23. The composition of claim 22, wherein the second threshold amount is 70%.


24. The composition of claim 22, wherein the second threshold amount is 80%.
25. The composition of claim 22, wherein the second threshold amount is 90%.
26. The composition of claim 22, wherein the second threshold amount is 95%.
27. The composition of any of claims 1-4, wherein greater than or equal to the first threshold amount of the total amount of chemical entities of formula **I** in the composition are chemical entities of formula **I.b.2**.
28. The composition of claim 27, characterized in that greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition is a chemical entity of formula **I.b.2.i**:



wherein the second threshold amount is 50%.

29. The composition of claim 28, wherein the second threshold amount is 70%.
30. The composition of claim 28, wherein the second threshold amount is 80%.
31. The composition of claim 28, wherein the second threshold amount is 90%.
32. The composition of claim 28, wherein the second threshold amount is 95%.

33. The composition of claim 27, characterized in that greater than or equal to a second threshold amount of the total amount of chemical entities of formula **I.b.2** in the composition is a chemical entity of formula **I.b.2.ii**:



wherein the second threshold amount is 50%.

34. The composition of claim 33, wherein the second threshold amount is 70%.

35. The composition of claim 33, wherein the second threshold amount is 80%.

36. The composition of claim 33, wherein the second threshold amount is 90%.

37. The composition of claim 33, wherein the second threshold amount is 95%.


38. The composition of any claims 1-4, characterized in that greater than or equal to a third threshold amount of the total amount of the composition are chemical entities of formula **I**, wherein the third threshold amount is 85% (w/w).

39. The composition of claim 38, wherein the third threshold amount is 90%.


40. The composition of claim 38, wherein the third threshold amount is 95%.

41. The composition of claim 38, wherein the third threshold amount is 98%.


42. The composition of claim 41, wherein greater than or equal to 90% (w/w) of the total amount of the composition is a chemical entity of formula **I.a.i**:

**I.a.i.**


43. The composition of claim 41, wherein greater than or equal to 90% (w/w) of the total amount of the composition is a chemical entity of formula **I.a.ii**:

**I.a.ii.**


44. The composition of claim 42, wherein greater than or equal to 90% (w/w) of the total amount of the composition is a chemical entity of formula **I.b.1.i**:

**I.b.1.i.**


45. The composition of claim 41, wherein greater than or equal to 90% (w/w) of the total amount of the composition is a chemical entity of formula **I.b.2.i**:

**I.b.2.i.**

46. The composition of claim 41, wherein greater than or equal to 90% (w/w) of the total amount of the composition is a chemical entity of formula **I.b.1.i**:

**I.b.1.i.**

47. The composition of claim 41, wherein greater than or equal to 90% (w/w) of the total amount of the composition is a chemical entity of formula **I.b.2.ii**:

**I.b.2.ii.**

48. A method of delivery of messenger RNA (mRNA) *in vivo*, comprising

administering to a subject in need of delivery a composition comprising an mRNA encoding a protein, encapsulated within a liposome such that the administering of the composition results in the expression of the protein encoded by the mRNA *in vivo*;

wherein the liposome comprises a cationic lipid composition of any one of claims 1-47.

49. The method of claim 48, wherein the liposome further comprises one or more non-cationic lipids, one or more cholesterol-based lipids and/or one or more PEG-modified lipids.

50. The method of claim 49, wherein the one or more non-cationic lipids are selected from DSPC (1,2-distearoyl-sn-glycero-3-phosphocholine), DPPC (1,2-dipalmitoyl-sn-glycero-3-phosphocholine), DOPE (1,2-dioleoyl-sn-glycero-3-phosphoethanolamine), DOPC (1,2-dioleoyl-sn-glycero-3-phosphotidylcholine) DPPE (1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine), DMPE (1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine), DOPG (1,2-dioleoyl-*sn*-glycero-3-phospho-(1'-*rac*-glycerol)).

51. The method of claim 49 or 50, wherein the one or more cholesterol-based lipids are cholesterol and/or PEGylated cholesterol.

52. The method of any one of claims 49-51, wherein the one or more PEG-modified lipids comprise a poly(ethylene) glycol chain of up to 5 kDa in length covalently attached to a lipid with alkyl chain(s) of C₆-C₂₀ length.

53. The method of any one of claims 48-52, wherein the liposome has a size less than about 250 nm, 200 nm, 150 nm, 100 nm, 75 nm, or 50 nm.

54. The method of any one of claims 48-53, wherein the mRNA has a length of or greater than about 0.5kb, 1 kb, 1.5 kb, 2 kb, 2.5 kb, 3 kb, 3.5 kb, 4 kb, 4.5 kb, or 5 kb.

55. The method of any one of claims 48-54, wherein the protein encoded by the mRNA is a cytosolic protein.

56. The method of any one of claims 48-54, wherein the protein encoded by the mRNA is a secreted protein.


57. The method of any one of claims 48-54, wherein the protein encoded by the mRNA is an enzyme.

58. The method of any one of claims 48-57, wherein the mRNA comprises one or more modified nucleotides.

59. The method of claim 58, wherein the one or more modified nucleotides comprise pseudouridine, N-1-methyl-pseudouridine, 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, C-5 propynyl-cytidine, C-5 propynyl-uridine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deaza-guanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and/or 2-thiocytidine.

60. The method of any one of claims 48-59, wherein the mRNA is unmodified.

61. A method of treating a disease or disorder comprising a step of delivering an mRNA encoding a therapeutic protein using a method of any of claims 48-60.

FIG. 1




FIG. 1

