

- [54] Title: INFUSION SOLUTIONS OF 1-CYCLOPROPYL-6-FLUORO-1,4-DIHYDRO-4-OXO-7-(1-PIPERAZINYL)-QUINOLINE-3-CARBOXYLIC ACID
- [75] Inventor (s): ROBERT FRANK LAMMENS, of Leverkusen; HANS FRIEDRICH MAHLER and PETER SERNO, bot of Koeln, all of Germany
- [73] Assignee (s): BAYER AKTIENGESELLSCHAFT, of Leverkusen, Germany, a corporation of Germany
- [22] Filed: October 23, 1986
- [21] Application Serial No: 34403

FOREIGN APPLICATION PRIORITY DATA

- [31] Number (s) : P3537761.5
- [32] Date (s) : October 24, 1985
- [33] Country (ies) : Federal Republic of Germany
- [52] PH Class 514/763, 514/557; 546/152, 546/255
- [51] Int. Class A61K 31/495, 31/015, 31/19; C07D215/00, C07D 215/56
- [58] Field of Search G1: 514/763, 514/557 ; 546/152, 546/255
IPC: A61K 31/495, 31/015, 31/19
- [56] Reference (s) Cited and/or Considered: C07D 215/00, 215/56

Chemical Abstract vol. 99 p. 53790 h

[57]

ABSTRACT

An aqueous infusion solution containing 0.015 to 0.5 g of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid per 100 ml of aqueous solution and an amount of at least one physiologically tolerated acid which suffices to dissolve the active compound.

PHILIPPINE PATENT OFFICE

'86 OCT 23 P3:04

Received by:

26419

SPECIFICATION

TO ALL WHOM IT MAY CONCERN:

BE IT KNOWN THAT WE

1) Robert Frank Lammens 2) Hans Friedrich Mahler 3) Peter Sern
citizens of Germany, residing at

- 1) Walter-Flex-Strasse 30e, D 5090 Leverkusen 1, Germany
- 2) Roggendorfstrasse 55, D 5000 Koeln 80, Germany
- 3) An der Ruthen 3, D 5000 Koeln 80, Germany

have invented certain new and useful Improvements in
Infusion solutions of 1-cyclopropyl-6-fluoro-1,4-dihydro-
4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid

of which the following is a full, clear and exact description.

Le A 23 890-PH

The invention relates to both infusion solutions, which are ready for use, of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid (= ciprofloxacin, to be designated active compound in the text which follows) and other presentations, which, before administration, are converted into infusion solutions of this type. The invention likewise relates to processes for the preparation of the infusion solutions and to their use for the therapeutic treatment of the human or animal body.

Solutions of lactic acid salts of piperazinyl-quinoline- and piperazinyl-azaquinolinecarboxylic acids are described in European Patent Application 84110474.8.

European Patent Application 81106511.9 relates to 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid and its pharmaceutically utilizable salts.

The present invention relates to infusion solutions of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid (= ciprofloxacin) which contain 0.015 to 0.5 g of the active compound per 100 ml of aqueous solution and an amount of a physiologically tolerated acid which suffices to dissolve the active compound and to stabilize the solution and, where appropriate, customary formulating auxiliaries.

In addition to the active compound, water and other customary formulating auxiliaries, the infusion solutions according to the invention preferably contain an amount, which suffices to dissolve the active compound and to stabilize the solution, of one or more acid(s) from the group comprising hydrochloric acid, methanesulphonic acid, propionic acid, succinic acid, glutaric acid, citric acid, fumaric acid, maleic acid,

Le A 23 890

tartaric acid, glutamic acid, gluconic acid, glucuronic acid, galacturonic acid, ascorbic acid, phosphoric acid, adipic acid, hydroxyacetic acid, sulphuric acid, nitric acid, acetic acid, malic acid, L-aspartic acid and lactic acid.

Lactic acid and hydrochloric acid or mixtures of hydrochloric acid and lactic acid are particularly preferred.

Furthermore preferred are infusion solutions which contain 0.015 to 0.5 g of the active compound per 100 ml of aqueous solution and, depending on the active compound concentration, up to 5.0 mols, in particular 0.9 to 5.0 mols, and particularly preferably 1.04 to 2.20 mols, relative to 1 mol of active compound, of one or more physiologically tolerated acids, and where several acids are present their total content does not exceed the amount of 5.0 mols, relative to 1 mol of active compound.

Moreover, the invention preferably relates to infusion solutions which contain 0.015 to 0.5 g of the active compound per 100 ml of aqueous solution and up to 5.0 mols, relative to 1 mol of active compound, of lactic acid. The amounts of lactic acid in this connection are preferably 0.99 to 1.50 mols, in particular 1.04 to 1.40 mols, relative to 1 mol of active compound, of lactic acid. Infusion solutions of the active compound which contain 1.12 to 1.24 mols, relative to 1 mol of active compound, of lactic acid are particularly advantageous.

The infusion solutions according to the invention can also be modified in such a way that they contain up to 0.5 g of the active compound per 100 ml of aqueous solution and up to 1 mol, relative to 1 mol of active compound, of lactic acid, together with another physiologically tolerated acid, with the proviso that the total amount of acid is, depending on the

Le A 23 890

active compound concentration, more than 0.9 mol but does not exceed 5.0 mol, relative to 1 mol of active compound.

5 The minimum amount of acid necessary per mol of active compound for the dissolution depends on the active compound concentration and the acid(s) used, and thus is not constant. However, within the limits according to the invention it can be determined by simple experiments. Furthermore, it should be noted
10 that the data in the amounts of acid relate only to the amounts which, according to generally known chemical laws, are not converted by the addition of bases into the corresponding salt(s). Dissociation of the acids was left out of account in the data on the amounts so
15 that they relate to the amount of dissociated and undissociated acid.

The lactic acid used in the formulations has a content of less than 25% (w/w), specifically for reasons of processing technology. The use of concentrated lactic acid - for example a 90% strength (w/w) product -
20 gives rise to difficulties when the pH of the formulations according to the invention is to be adjusted after addition of the lactic acid - for example with hydrochloric acid or sodium hydroxide solution - with
25 the objective that the adjusted pH remains constant, or changes only inconsiderably, during the remainder of the preparation process (such as, for example, a heat treatment at about 120°C for about 20 min) and/or during storage.

30 The infusion solutions according to the invention can also contain other formulating aids such as thickeners, resorbents, light-protection agents, absorption inhibitors, crystallization accelerators, absorption accelerators, crystallization retardants,
35 complexing agents, antioxidants, isotonicizing agents and/or euhydrogenating agents.

Le A 23 890

26419

The osmolality of the infusion solutions is 0.20 to 0.70 Osm/kg, preferably 0.26 to 0.39 Osm/kg and is adjusted by isotonicizing agents such as NaCl, sorbitol, mannitol, glucose, sucrose, xylitol, fructose and glycerol or mixtures of such substances. Where appropriate, it is also possible to use for this substances which are contained in conventional commercially available infusion vehicle solutions.

The customary infusion vehicle solutions include infusion solutions with the addition of electrolytes without carbohydrates, such as sodium chloride solution, Ringer lactate solution and the like, and those with carbohydrates, as well as solutions for supplying amino acids, in each case with and without a carbohydrate content. Examples of infusion vehicle solutions of these types are listed in the Rote Liste 1985, List of finished pharmaceuticals of the members of the German Association of the Pharmaceutical Industry, Editio Cantor, Aulendorf/Württ.

Preferred infusion solutions are those which, apart from water, active compound and other formulating auxiliaries, contain an amount of sodium chloride, or other auxiliaries customary for isotonicizing, such that the solution is in a form which is isotonic, or slightly hypo- or hypertonic, with the tissue fluid in the human or animal body.

The infusion solutions according to the invention have a pH of 3.0 to 5.2. pH values from 3.6 to 4.7 and 3.9 to 4.5 are preferred. pH values in the range from 4.1 to 4.3 are very particularly preferred.

A very particularly preferred embodiment of the invention comprises infusion solutions which, apart from active compound, water and other formulating auxiliaries, contain, depending on the amount of active compound, 0.99 to 1.50 mols, preferably 1.04 to 1.40 mols, of lactic acid and 0.0 to 0.80 mol of hydrochloric

Le A 23 890

acid (in each case relative to 1 mol of active compound), and, relative to 100 ml of solution, 0.6 to 2.2 g of NaCl, preferably 0.75 to 1.20 g, in particular 0.85 to 0.95 g of NaCl. The solutions thus obtained have
5 osmolalities which differ according to the amount of sodium chloride and the active compound concentration. The osmolalities relating to the amounts of sodium chloride listed above are 0.2 to 0.7, 0.26 to 0.39 and 0.28 to 0.32 Osm/kg of solution respectively. Corresponding values can also be adjusted using other isotonicizing agents or mixtures thereof, as indicated above.
10 Depending on the active compound and acid concentration, small differences from these osmolalities are perfectly possible.

15 The infusion solutions according to the invention can be in the form of dosage units, suitable for infusion, with removable contents of 40 to 600 ml, preferably 50 to 120 ml.

20 However, the invention also relates to lyophilizates which have been prepared by customary techniques and which are converted into the infusion solutions according to the invention by dissolution in solvents suitable for this purpose - such as, for example, conventional infusion vehicle solutions. Lyophilizates of this type can be obtained by freeze-drying of various starting solutions such as, for example, the infusion solutions according to the invention. It is likewise possible to freeze-dry considerably more dilute solutions as well as considerably more
25 concentrated solutions than the infusion solutions according to the invention.
30

The lyophilizates can be prepared both by freeze-drying in the final container such as, for example, in a bottle or ampule made of glass or plastic, and by bulk freeze-drying combined with dispensing the
35 lyophilizate into a container suitable for this purpose,
Le A 23 890

which takes place at a later time.

The dissolution of the lyophilizate before the administration can be brought about both by addition of a solution, which is suitable for this purpose, into the container containing the lyophilizate and by addition of the lyophilizate to a suitable solution, or by a combination of procedures of these types.

The composition of the lyophilizates can likewise vary very widely, depending on the composition of the solution which is used for the dissolution.

It can vary from pure active compound to a lyophilizate which contains all the constituents which are to be administered, apart from water.

The invention likewise relates to combinations of lyophilizates with solutions containing active compound, which are converted into the infusion solutions according to the invention before the administration.

The invention also includes concentrates and suspensions which are converted into the solutions according to the invention before the administration.

It is possible in this context for these concentrates and suspensions to have various compositions. One possibility would be that which requires merely the addition of water for dilution or dissolution in order to prepare the infusion solutions according to the invention.

This invention relates to all combinations of concentrates and/or suspensions and to solutions which are necessary for dilution or dissolution and which result in the solutions according to the invention.

The invention also relates to other presentations or combinations of presentations which finally result in the infusion solutions according to the invention - and this irrespective of the procedure.

The containers into which lyophilizates, concentrates and other presentations such as, for example,

Le A 23 890

suspensions, are dispensed can consist both of glass and of plastic. In this connection, the container materials can contain substances which confer a particular protection on the contents, such as, for example, a protection from light or a protection from oxygen.

The present invention additionally relates to a process for the preparation of infusion solutions, containing 0.015 to 0.5 % by weight of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid (= ciprofloxacin). This process comprises mixing a suitable amount of the active compound, where appropriate in the form of a salt, such as an alkali metal or alkaline earth metal salt or addition salt, of a hydrate or of a hydrate of the salt, or in the form of mixtures of these salts or hydrates, with the amount of a physiologically tolerated acid or of a mixture of several physiologically tolerated acids which, in relation to the amount which just suffices to dissolve the active compound or its salts or hydrates, represents an excess preventing separation out of the active compound, adding, where appropriate, formulating auxiliaries, and making up with water or a customary infusion vehicle solution in such a manner that the concentration of the active compound is adjusted to the range from 0.015 to 0.5 g. In this connection, it has to be remembered that when the alkali metal or alkaline earth metal salts of the active compound are used the amounts of acid which are mentioned above as being necessary for dissolution contain the amount which is needed to neutralize the active compound anion, and that when addition salts are used a part of the amounts of acid necessary is already present in the active compound salt which is to be used.

Furthermore, care has to be taken in the preparation that the solution complies with the properties relating to pH, amounts of acid and osmolalities which

Le A 23 890

have already been detailed.

In the case where the active compound is used in the salt form, it is possible and expedient to use an acid whose anion corresponds to the anion of the active compound salt or salt hydrate.

Where appropriate, the active compound is suspended in water, and up to 5 mols, relative to 1 mol of active compound, of lactic acid are added, and then, where appropriate, another physiologically tolerated acid or a mixture of such acids, in particular hydrochloric acid, is added, with the proviso that the total amount of acid does not exceed 5.0 mols, relative to 1 mol of active compound, but does exceed 0.9 mol, relative to 1 mol of active compound, and then, where appropriate, further formulating auxiliaries are added, in particular NaCl, which is also, where appropriate, produced by a neutralization reaction in the formulation mixture, and the desired active compound concentration is adjusted by making up with water.

The pH of the infusion solutions according to the invention can be adjusted with (physiologically) tolerated acids and/or bases to the abovementioned values, that is to say 3.0 to 5.2, in particular 3.6 to 4.7.

To speed up the preparation process, in particular the dissolution of solid components, it is possible gently to heat the solutions, or only a part thereof, preferably to temperatures between 20°C and 80°C.

It has been possible particularly economically to prepare the solutions according to the invention via concentrated solutions. For this purpose, the amount of active compound necessary for a batch was dissolved with the major amount of acid necessary for the complete batch (for example >95% relative to the molar basis) in a little water - where appropriate with

Le A 23 890

heating. This concentrate was then subsequently diluted. After dilution, any other auxiliaries - such as, for example, sodium chloride for isotonicizing - were added, as were the amounts of acid which were still
5 lacking, where appropriate.

The infusion solutions according to the invention are used for the therapeutic treatment of the human or animal body.

10 The infusion solutions according to the invention have a low toxicity and a broad spectrum of antibacterial activity against Gram-positive and Gram-negative microbes, in particular against Enterobacteriaceae; especially including those which are resistant to various antibiotics such as, for example, penicillins, cephalosporins, aminoglycosides, sulphonamides
15 and tetracyclines.

These valuable properties make it possible to use them in medicine.

20 The infusion solutions according to the invention are active against a very broad spectrum of microorganisms. They can be used to combat Gram-negative and Gram-positive bacteria and bacteria-like microorganisms and to prevent, ameliorate and/or heal illnesses caused by these pathogens.

25 The infusion solutions according to the invention are particularly active against bacteria and bacteria-like microorganisms. They are therefore particularly suitable for the prophylaxis and chemotherapy of local and systemic infections, caused by these
30 pathogens, in human medicine and veterinary medicine.

For example, local and/or systemic illnesses caused by the following pathogens or by mixtures of the following pathogens can be treated and/or prevented:

35 Gram-positive cocci, for example Staphylococci (Staph. aureus, Staph. epidermidis) and Streptococci (Strept. agalactiae, Strept. faecalis, Strept. pneu-

Le A 23 890

moniae, Strept. pyogenes); Gram-negative cocci (Neisseria gonorrhoeae) and Gram-negative rods such as Enterobacteriaceae, for example Escherichia coli, Haemophilus influenzae, Citrobacter (Citrob. freundii, Citrob. divernis), Salmonella and Shigella; furthermore Klebsiella (Klebs. pneumoniae, Klebs. oxytoca), Enterobacter (Ent. aerogenes, Ent. agglomerans), Hafnia, Serratia (Serr. marcescens), Proteus (Pr. mirabilis, Pr. rettgeri, Pr. vulgaris), Providencia, Yersinia, and the genus Acinetobacter. Furthermore, the antibacterial spectrum covers the genus Pseudomonas (Ps. aeruginosa, Ps. maltophilia) and strictly anaerobic bacteria such as, for example, Bacteroides fragilis, representatives of the genus Peptococcus, Peptostreptococcus and the genus Clostridium; furthermore Mycoplasmas (M. pneumoniae, M. hominis, M. urealyticum) as well as Mycobacteria, for example Mycobacterium tuberculosis.

The above list of pathogens is purely illustrative and is in no way to be interpreted as restrictive. The following may be mentioned as examples of illnesses which are caused by the said pathogens or mixed infections and can be prevented, ameliorated or healed by the compounds according to the invention:

Infectious illnesses in humans such as, for example, otitis, pharyngitis, pneumonia, peritonitis, pyelonephritis, cystitis, endocarditis, systemic infections, bronchitis (acute and chronic), septic infections, illnesses of the upper airways, diffuse panbronchiolitis, pulmonary emphysema, dysentery, enteritis, liver abscesses, urethritis, prostatitis, epididymitis, gastrointestinal infections, bone and joint infections, cystic fibrosis, skin infections, postoperative wound infections, abscesses, phlegmon, wound infections, infected burns, burns, infections in the mouth, infections after dental operations, osteomyelitis, septic

Le A 23 890

arthritis, cholecystitis, peritonitis with appendicitis, cholangitis, intraabdominal abscesses, pancreatitis, sinusitis, mastoiditis, mastitis, tonsillitis, typhoid, meningitis and infections of the nervous system, salpingitis, endometritis, genital infections, pelveoperitonitis and eye infections.

Apart from humans, it is also possible to treat bacterial infections in other species. The following may be mentioned as examples:

- 10 Pig: coli diarrhoea, enterotoxaemia, sepsis, dysentery, salmonellosis, metritis-mastitis-agalactiae syndrome, mastitis;
- ruminants (cattle, sheep, goat): diarrhoea, sepsis, bronchopneumonia, salmonellosis, pasteurellosis, mycoplasmosis, genital infections;
- 15 horse: bronchopneumonias, joint ill, puerperal and post-puerperal infections, salmonellosis;
- dog and cat: bronchopneumonia, diarrhoea, dermatitis, otitis, urinary tract infections, prostatitis;
- 20 poultry (chicken, turkey, quail, pigeon, ornamental birds and others): mycoplasmosis, E. coli infections, chronic respiratory tract illnesses, salmonellosis, pasteurellosis, psittacosis.

It is likewise possible to treat bacterial illnesses in the rearing and maintenance of productive and ornamental fishes, the antibacterial spectrum extending beyond the pathogens mentioned above to further pathogens such as, for example, Pasteurella, Brucella, Campylobacter, Listeria, Erysipelothrix, Corynebacteria, Borellia, Treponema, Nocardia, Rickettsia and Yersinia.

Examples

The molar ratio - abbreviated to R - stated in the examples which follow always relates to the substance which is mentioned first in the relevant example.

Micromole R

Examp. Formulation

No.

1	Ciprofloxacin	90 mg	272	1.00
	Lactic acid 20% (w/w)	144.3 mg	320	1.18
	Hydrochloric acid	1.5 mg	48	0.15
	Sodium chloride	5.4 g	-	-
	Water	ad 600.0 ml		
		pH: approx. 4.3		
		Osm: approx. 0.29 Osm/kg		
2	Ciprofloxacin	150 mg	453	1.00
	Lactic acid 10% (w/w)	558 mg	658	1.37
	Hydrochloric acid	7.8 mg	214	0.47
	Glucose	30.0 g	-	-
	Water	ad 600.0 ml		
		pH: approx. 3.7		
		Osm: approx. 0.29 Osm/kg		

Micromole R

Examp. No.	Formulation			
3	Ciprofloxacin	100 mg	302	1.00
	Lactic acid 25% (w/w)	629 mg	1745	5.78
	2 M NaOH solution	0.177 ml	354	1.17
	Fructose	20.0 g	-	-
	Water	400.0 ml	-	-
		ad		
	pH:	3.6 a 3.7		
	Osm:	0.29 Osm/kg		
4	Ciprofloxacin	75 mg	226	1.00
	1 M hydrochloric acid	0.203 ml	203	0.90
	Sodium chloride	4.5 g	-	-
	Water	500 ml	-	-
		ad		
		pH:	5.2	
	Osm:	0.29 Osm/kg		

Micromole R

Examp. No.	Formulation		1.00 (containing an equimolar amount of lactic acid)
5	Ciprofloxacin lactate	254 mg	604
	Lactic acid 5% (w/w)	400 mg	222
	Hydrochloric acid	10.4 mg	
	Sodium chloride	1.8 g	
	Glucose	10.0 g	
	Water	400 ml	
	ad	3.7	
	pH:	0.29 Osm/kg	
	Osm:		
6	Ciprofloxacin . 5 H ₂ O	254 mg	604
	Lactic acid 2% (w/w)	2740 mg	609
	Sodium chloride	3.6 g	-
	Water	400 ml	-
	ad	5.2	
	pH:	0.29 Osm/kg	
	Osm:		

Le A 23 890

Micromole R

Examp. Formulation

No.

Examp. No.	Formulation	Micromole R
7	Ciprofloxacin acetate	302
	Acetic acid	3.6 mg
	Lactic acid 10% (w/w)	272 mg
	Sorbitol	10.0 g
	Water	200.0 ml
	ad	4.7
	Osm:	0.30 Osm/kg
8	Ciprofloxacin	604
	Propionic acid	724
	Lactic acid 20% (w/w)	604
	Glucose	20.0 g
	Water	400.0 ml
	ad	4.7
	Osm:	0.3 Osm/kg
	1.00 (containing an equivalent amount of acetic acid)	0.20
		1.00
		-

Micromole R

Examp. Formulation
No.

Examp. No.	Formulation	Micromole R
9	Ciprofloxacin lactate . 2 H ₂ O Lactic acid 20% (w/w) 0.2 M NaOH Sorbitol Water ad pH: Osm:	755 1822 mg 5.25 ml 5.0 g 250.0 ml 3.6 0.30 Osm/kg 4045 1050 - - 1.00 (con- taining an equimolar amount of lactic acid) 5.36 1.39 - -
10	Ciprofloxacin Lactic acid 1% (w/w) 1 N hydrochloric acid Mannitol Water ad pH: Osm:	604 826 285 - 1.00 1.37 0.47 - 200 mg 7440 mg 0.285 ml 6.2 g 200 ml 3.7 0.37 Osm/kg
11	Ciprofloxacin Lactic acid 0.5% (w/w) Sodium chloride Water ad pH: Osm:	1509 1509 - 1.00 1.00 - 500 mg 27182 mg 4.5 g 500 ml 5.1 0.29 Osm/kg

Micromole R

Examp. No.	Formulation	Micromole R
12	Ciprofloxacin	1509
	Lactic acid 5% (w/w)	1509
	Glycerol	-
	1 M sodium chloride	1885
	Water	1.25
	500 mg	
	2718 mg	
	13.0 g	
	1.885 ml	
	ad	
	pH: 3.0	
	Osm: 0.29 Osm/kg	
13	Ciprofloxacin	302
	Lactic acid 0.1% (w/w)	357
	Hydrochloric acid	45
	Glucose	-
	Water	-
	100 mg	
	32.2 g	
	1.6 mg	
	4.48 g	
	ad	
	pH: 4.2	
	Osm: 0.26 Osm/kg	
14	Ciprofloxacin	151
	Lactic acid 10% (w/w)	178
	Sodium chloride	-
	Water	-
		50 mg
	160 mg	
	625 mg	
	50 ml	
	ad	
	pH: 4.4	
	Osm: 0.40 Osm/kg	

Examp. No.	Formulation	mg	Micromole R
15	Ciprofloxacin lactate . 2 H ₂ O	276 mg	604
	1 M hydrochloric acid	0.151 ml	
	Fructose	17.2 g	
	Water	200.0 ml	
	ad		
	pH:	4.2	
	Osm:	0.50 Osm/kg	
			151
			-
16	Ciprofloxacin	500 mg	1509
	Lactic acid 2% (w/w)	6727 mg	1494
	Sorbitol	25.0 g	
	Water	500 ml	
	ad		
	pH:	5.0	
	Osm:	0.29 Osm/kg	
			1.00
			0.99
17	Ciprofloxacin lactate	127 mg	302
	Maleic acid	42 mg	
	Mannitol	5.0 g	
	Water	100.0 ml	
	ad		
	pH:	3.1	
	Osm:	0.30 Osm/kg	
			364
			1.20

Examp. No.	Formulation				
18	Ciprofloxacin	200 mg	604	1.00	
	Lactic acid 20% (w/w)	297 mg	659	1.09	
	Glutaric acid	40 mg	302	0.50	
	Fructose	10.0 g			
	Water	200 ml			
		ad			
	pH:	4.3			
	Osm:	0.30 Osm/kg			
19	Ciprofloxacin	200 mg	604	1.00	
	Lactic acid 10% (w/w)	566 mg	628	1.04	
	Sodium chloride	0.9 g	-	-	
	Water	100.0 ml	-	-	
		ad			
		pH:	4.9		
	Osm:	0.29 Osm/kg			
20	Ciprofloxacin	400 mg	1207	1.00	
	Lactic acid 20% (w/w)	745 mg	1654	1.37	
	Hydrochloric acid	20.8 mg	570	0.47	
	Sodium chloride	1800 mg			
	Water	200 ml			
		ad			
	pH:	3.7			
	Osm:	0.29 Osm/kg			

Micromole R

Examp. Formulation

No.

21	Ciprofloxacin . 5 H ₂ O	509 mg	1207	1.00
	Lactic acid 5% (w/w)	2568 mg	1425	1.18
	Hydrochloric acid	6.6 mg	180	0.15
	Glucose	5.0 g	-	-
	Water	200 ml		
	ad	4.2		
	pH:	0.30 Osm/kg		
	Osm:			

22	Ciprofloxacin . 5 H ₂ O	254 mg	604	1.00
	Lactic acid 25% (w/w)	257 mg	713	1.18
	Hydrochloric acid	3.3 mg	90	0.15
	Sorbitol	5.0 g		
	Water	100.0 ml		
	ad	4.2		
	pH:	0.30 Osm/kg		
	Osm:			

Micromole R

Examp. No.	Formulation	Micromole R
23	Ciprofloxacin	604
	Lactic acid 20% (w/w)	715
	1 M hydrochloric acid	90
	Sodium chloride	
	Water	
	200 mg	1.00
	322 mg	1.18
	0.090 ml	0.15
	0.9 g	
	ad 100.0 ml	
	pH: 4.2	
	Osm: 0.30 Osm/kg	
24	Ciprofloxacin	302
	Lactic acid 0.4% (w/w)	329
	1 M hydrochloric acid	3420
	2 M sodium hydroxide solution	3420
	Sodium chloride	
	Water	
	100 mg	1.00
	7407 mg	1.09
	3.42 ml	11.32
	1.71 ml	11.32
	0.25 g	
	ad 50.0 ml	
	pH: 4.7	
	Osm: Osm/kg	

Examp. No.	Formulation	Micromole R
25	Ciprofloxacin	604
	Lactic acid 10% (w/w)	713
	Hydrochloric acid	90
	Sodium chloride	
	Water	
	200 mg	1.00
	642 mg	1.18
	3.3 mg	0.15
	0.855 g	
	ad 100.0 ml	
	pH: 4.2	
	Osm: 0.29 Osm/kg	
26	Ciprofloxacin . 5 H ₂ O	604
	Lactic acid 20% (w/w)	715
	0.1 M hydrochloric acid	90
	Sodium chloride	-
	Water	-
	254 mg	1.00
	322 mg	1.18
	0.90 ml	-
	0.945 g	-
	ad 100.0 ml	
	pH: 4.2	
	Osm: 0.32 Osm/kg	

Examp. No.	Formulation	Micromole R
27	Ciprofloxacin lactate . 2 H ₂ O Lactic acid 0.5% (w/w) Hydrochloric acid Xylitol Water ad pH: Osm:	604 111 90 - 2000 mg 3.3 mg 4.2 g 100.0 ml 4.2 0.30 Osm/kg
28	Ciprofloxacin lactate Lactic acid 20% (w/w) Hydrochloric acid Fructose Water ad pH: Osm:	604 111 90 50 mg 3.3 mg 6.0 g 100.0 ml 4.2 0.37 Osm/kg
29	Ciprofloxacin HCl . H ₂ O Lactic acid 5% (w/w) 2 M sodium hydroxide solution Sodium chloride Water ad pH: Osm:	604 715 514 1288 mg 0.257 ml 0.87g 100 ml 4.2 0.30 Osm/kg

Micromole R

Examp. Formulation

No.

30	Ciprofloxacin Na (Sodium salt of ciprofloxacin)	213 mg	604	1.00
	Lactic acid 10% (w/w)	744 mg	826	1.37
	Hydrochloric acid	32.4 mg	888	1.47
	Glucose	4.8 g		
	Water	ad 100.0 ml		
		pH: 3.7		
		Osm: 0.30 Osm/kg		

31	Calcium salt of ciprofloxacin	212 mg	604	1.00
	Lactic acid 2% (w/w)	3220 mg	715	1.18
	Hydrochloric acid	25.3 mg	693	1.15
	Glycerol	2.6 g	-	-
	Water	ad 500.0 ml		
		pH: 4.3		
		Osm: 0.30 Osm/kg		

Le A 23 890

Examp. No.	Formulation	Micromole R
32	Potassium salt of ciprofloxacin	604
	Lactic acid 20% (w/w)	615
	0.1 M hydrochloric acid	886
	Glucose	
	Water	
	233 mg	1.00
	277 mg	1.018
	8.86 ml	1.47
	5.0 g	
	ad 100.0 ml	
	pH: 4.6	
	Osm: 0.30 Osm/kg	
33	Ciprofloxacin lactate . 2 H ₂ O	604
	Lactic acid 10% (w/w)	3236
	0.1 M sodium hydroxide solution	840
	Glucose	
	Water	
	276 mg	1.00 (containing an equimolar amount of lactic acid)
	2915 mg	5.36
	8.40 ml	1.39
	5.0 g	
	ad 100.0 ml	
	pH: 3.6	
	Osm: 0.3 Osm/kg	

Examp. No.	Formulation	Micromole R
34	<p>Ciprofloxacin HCL (Hydrochloride of ciprofloxacin)</p> <p>Lactic acid 0.1% (w/w)</p> <p>2 M sodium hydroxide solution</p> <p>Hydrochloric acid</p> <p>Sodium chloride</p> <p>Water</p> <p>ad</p> <p>pH: 4.6</p> <p>Osm: 0.30 Osm/kg</p>	<p>604</p> <p>615</p> <p>604</p> <p>284</p> <p>-</p>
35	<p>Ciprofloxacin</p> <p>Tartaric acid</p> <p>Xylitol</p> <p>Water</p> <p>ad</p> <p>pH: 3.6</p> <p>Osm: 0.30 Osm/kg</p>	<p>604</p> <p>604</p> <p>1.00 (containing an equivalent molar amount of hydrochloric acid)</p> <p>1.00</p> <p>0.47</p> <p>-</p> <p>1.00</p> <p>1.00</p>

Micromole R

Examp. Formulation

No.

Examp. No.	Formulation	Micromole R
36	Ciprofloxacin gluconate	302
	159 mg	1.00 (containing an equivalent molar amount of gluconic acid)
	Gluconic acid	62
	Lactic acid 5% (w/w)	302
	Glycerol	-
	Water	-
	ad	
	pH:	
	Osm:	
	12 mg	
	544 mg	
	1.3 g	
	50.0 ml	
	4.0	
	0.30 Osm/kg	
37	Ciprofloxacin	1509
	Galacturonic acid	1811
	Sorbitol	
	Water	
	ad	
	pH:	
	Osm:	
	500 mg	
	351 mg	
	25.0 g	
	250.0 ml	
	4.6	
	0.3 Osm/kg	
38	Ciprofloxacin. 5 H ₂ O	302
	Ascorbic acid	366
	Glucose	
	Water	
	ad	
	pH:	
	Osm:	
	127 mg	
	41 mg	
	2.5 g	
	50 ml	
	4.5	
	0.30 Osm/kg	
	1.00	
	1.21	

Micromole R

Examp. No.	Formulation			
39	Ciprofloxacin	200 mg	604	1.00
	Adipic acid	106 mg	724	1.20
	Lactic acid 20% (w/w)	272 mg		
	Fructose	5.0 g		
	Water	100.0 ml		
	ad			
	pH:	4.1		
	Osm:	0.30 Osm/kg		
40	Ciprofloxacin lactate	509 mg	1207	1.00 (con- taining an equimolar amount of lactic acid)
	Hydroxyacetic acid	110 mg		
	Sodium chloride	1.8 g		
	Water	200 ml		
		ad		
	pH:	4.0		
	Osm:	0.30 Osm/kg	1448	
41	Ciprofloxacin	200 mg	604	1.00
	Malic acid	81 mg		
	Glucose	5.0 g		
	Water	100.0 ml		
		ad		
	pH:	4.2		
	Osm:	0.30 Osm/kg		

Micromole R

Examp. No.	Formulation	Micromole R
42	Ciprofloxacin	302
	L. aspartic acid	361
	Fructose	
	Water	
	100 mg	1.00
	48 mg	1.20
	2.5 g	
	50.0 ml	
	ad	
	pH:	
	Osm:	
	0.30 Osm/kg	
43	Ciprofloxacin	302
	Lactic acid 20% (w/w)	453
	0.1 M hydrochloric acid	211
	Sorbitol	
	Water	
	100 mg	1.00
	204 mg	1.50
	2.11 ml	0.70
	2.5 g	
	50.0 ml	
	ad	
	pH:	
	Osm:	
	0.30 Osm/kg	
44	Ciprofloxacin . 5 H ₂ O	754
	Lactic acid 25% (w/w)	4363
	0.2 M sodium hydroxide solution	880
	Mannitol	
	Water	
	318 mg	1.00
	1572 mg	5.78
	4.40 ml	1.17
	5.0 g	
	50.0 ml	
	ad	
	pH:	
	Osm:	
	0.35 Osm/kg	

Examp. No.	Formulation	Micromole R
45	Ciprofloxacin 200 mg Lactic acid 15% (w/w) 496 mg 0.2 M hydrochloric acid 1.42 ml Glucose 5.00 g Water 40.0 ml ad 3.7 pH: 0.33 Osm/kg Osm:	604 1.00 826 1.37 284 0.47
46	ciprofloxacin lactate . 2 H ₂ O 3858 mg Sodium chloride 4.5 g 1 M hydrochloric acid 0.754 ml Water 500ml ad 4.7 pH: 0.30 Osm/kg Osm:	7544 1.00 (con- - taining an - equimolar - amount of - lactic acid) 754 0.10
47	Ciprofloxacin 500 mg Lactic acid 1% 14814 mg Sodium chloride 0.45 g Water 100.0 ml ad 4.7 pH: 0.31 Osm/kg Osm:	1509 1.00 1645 1.09

Micromole R

Examp. No.	Formulation		Micromole R
48	Ciprofloxacin HCl . H ₂ O	244 mg	604
	Fructose	2.0 g	
	Water	ad 40.0 ml	
		pH: 4.0	
		Osm: 0.30 Osm/kg	
49	Ciprofloxacin Lactate . 2 H ₂ O	359 mg	754
	Lactic acid 20% (w/w)	2.6 mg	69
	Succinic acid	45 mg	381
	Sorbitol	2.5 g	
	Water	50.0 ml	
		ad 4.2	
		pH: 0.32 Osm/kg	
		Osm:	
50	Ciprofloxacin	500 mg	1509
	Lactic acid 10% (w/w)	1481 mg	1644
	Citric acid	159 mg	757
	Glucose	5.0 g	
	Water	100.0 ml	
		ad 3.8	
		pH: 0.33 Osm/kg	
		Osm:	

Micromole R

Examp. Formulation

No.

Examp. No.	Formulation	Micromole R
51	Ciprofloxacin lactate	302
	127 mg	1.00 (containing an equimolar amount of lactic acid)
	Ciprofloxacin fumarate	151
	68 mg	0.50 containing an equimolar amount of fumaric acid
	Ciprofloxacin	151
	Lactic acid 20%	356
	Fructose	50 mg
	Water	160 mg
	ad	2.0 g
	pH:	40.0 ml
	Osm:	3.9
		0.32 Osm/kg
52	Ciprofloxacin	754
	L-glutamic acid	754
	Mannitol	250 mg
	Water	111 mg
	ad	3.5 g
	pH:	50.0 ml
	Osm:	4.6
		0.42 Osm/kg
53	Ciprofloxacin	1509
	1 M hydrochloric acid	1510
	Glucose	500 mg
	Water	1.51 ml
	ad	12.5 g
	pH:	250 ml
	Osm:	4.0
		0.32 Osm/kg

Micromole R

Examp. No.	Formulation	1509	1780	1.00	1.18
54	ciprofloxacin Lactic acid 25% Sodium chloride Water ad pH: Osm:	500 mg 641 mg 0.0 g 100.0 ml 4.5 0.30 Osm/kg			

55	<u>Solution A</u> ciprofloxacin lactate Lactic acid 20% (w/w) Hydrochloric acid Sodium chloride Water	127 mg 25 mg 1.6 mg 0.9 g ad . 100.0 ml	302	56	45
	<u>Solution B</u> Sodium chloride Water	2.25 g 250 ml			

1.00 (containing an equimolar amount of lactic acid)

prepare the solution ready for use by mixing solution A and B

Examp. No.	Formulation	
56	<p><u>Solution A</u> Solution A from Example 55</p> <p><u>Solution B</u> Sodium chloride 4.95 g Water ad 550 ml</p> <p>Prepare the solution ready for use by mixing solution A and B</p>	
57	<p><u>Solution A</u> Ciprofloxacin 100 mg Lactic acid 6% (w/w) 537 mg 0.1 M hydrochloric acid 0.45 ml Sodium chloride ad 0.45 g Water ad 50.0 ml</p> <p><u>Solution B</u> Glucose 12.5 g Water ad 250 ml</p> <p>Prepare the solution ready for use by mixing solution A and B</p>	<p>302 1.00 358 1.19 45 0.15</p>

Examp. No.	Formulation		
58	<u>Solution A</u>		
	Ciprofloxacin	100 mg	302 1.00
	Lactic acid 2% (w/w)	1610 mg	358 1.19
	Hydrochloric acid	1.6 mg	45
	Glucose	2.5 g	
	Water	50 ml	
	<u>Solution B</u>		
	Sodium chloride	900 mg	
	Water	100 ml	
	Prepare the solution ready for use by mixing solution A and B		
59	<u>Solution A</u>		
	Potassium salt of ciprofloxacin	223 mg	604 1.00
	Lactic acid 20% (w/w)	277 mg	615 1.018
	0.1 M hydrochloric acid	8.86 ml	886 1.47
	Glucose	5.0 g	
	Water	100.0 ml	
	<u>Solution B</u>		
	Ringer lactate solution	500 ml (for example Ringer lactate DAB 7	
	Braun Melsungen		
	Rote Liste 1985 No. 51013)		
	Prepare solution ready for use by mixing solution A and B		

Examp. No.	Formulation	Micromole R
60	<p><u>Solution A</u> Ciprofloxacin lactate . 2 H₂O 276 mg</p> <p>Lactic acid 0.1% (w/w) 10.0 g Hydrochloric acid 3.3 mg Xylitol 4.2 g Water ad 100.0 ml</p> <p><u>Solution B</u> Fructose 10.0 g Water ad 100.0 ml</p> <p>Prepare the solution ready for use by mixing solution A and B</p>	<p>604 1.00 (containing an equimolar amount of lactic acid)</p> <p>111 0.18 90 0.15</p>
61	<p><u>Solution A</u> Ciprofloxacin 200 mg Lactic acid 10% (w/w) 644 mg 0.1 M hydrochloric acid 0.90 ml Sodium chloride 0.9 g Water ad 100.0 ml</p> <p><u>Solution B</u> Xylitol 12.5 g Water ad 250 ml</p> <p>Prepare the solution ready for use by mixing solution A and B</p>	<p>604 1.00 715 1.18 90 0.15</p>

Le A 23 890

Micromole R

Examp. Formulation
No.

62 Solution A
 Ciprofloxacin 200 mg 604 1.00
 Lactic acid 1.5% (w/w) 4960 mg 826 1.37
 0.1 M hydrochloric acid 2.84 ml 284 0.47
 Glucose 5.00 g
 Water ad 40.0 ml

Solution B
 Glucose 5.0 g
 Water ad 100 ml

Prepare the solution ready for use by mixing solution A and B

63 Solution A
 Ciprofloxacin 50 mg 151 1.00
 Lactic acid 20% (w/w) 393 mg 872 5.77
 0.2 M sodium hydroxide solution 0.89 ml 178 1.18
 Water ad 5.0 ml

Solution B
 Sorbitol 2.5 g
 Water ad 50.0 ml

Prepare the solution ready for use by mixing solution A and B

Micromole R

Examp. Formulation

No.

64

Solution A

Solution of Example 63

Solution B

Glucose 5.0 g
 Water ad 100.0 ml

Prepare the solution ready for use by mixing solution A and B

65

Solution A

Ciprofloxacin 100 mg
 Lactic acid 10% (w/w) 372 mg
 Hydrochloric acid 5.2 mg
 Water ad 10.0 ml

302 1.00
 413 1.37
 142 0.47

Solution B

Sodium chloride 0.45 g
 Water ad 50.0 ml

Prepare the solution ready for use by mixing solution A and B

Micromole R

Examp. Formulation
No.

69 Solution A
Ciprofloxacin lactate 254 mg 604 1.00 (con-
taining an
equimolar
amount of
lactic acid)
1.09
1.40
Lactic acid 5% (w/w) 1188 mg 659
1 M hydrochloric acid 0.120 ml 120
Water ad 20.0 ml

Solution B

Fructose 10.0 g
Water ad 100.0 ml

Prepare the solution ready for use by mixing solution A and B

70

Solution A

Solution A from Example 69

Solution B

Sodium chloride 2.25 g
Water ad 250.0 ml

Prepare the solution ready for use by mixing solution A and B

LE A 23 890

Micromole R

Examp. Formulation

No.

71

Solution A

Solution A from Example 69

Solution B

Ringer lactate solution

500 ml (for example Ringer lactate DAB 7
supplied by Braun Melsungen /
Rote Liste 1985 No. 51013)

72

Solution A

Ciprofloxacin

250mg

754

1.00

Lactic acid 20% (w/w)

465 mg

1032

1.37

0.1 M hydrochloric acid solution

1.63 ml

163

0.22

Sodium chloride

173 mg

Water

25 ml

Solution B

Glucose

5.0 g

Water

100.0 ml

Prepare the solution ready for use by mixing solution A and B

Micromole R

Examp. Formulation

No.

Le A 23 890

73

Solution A

Solution A from Example 72

Solution B

Sodium chloride

2.25 g

Water

250 ml

Prepare the solution ready for use by mixing solution A and B

74

Solution A

Ciprofloxacin lactate. 2 H₂O

276 mg

604

1.00 (containing an amount of lactic acid)

Lactic acid 10% (w/w)

20 mg

22

0.1 M hydrochloric acid solution

2.00 ml

200

Water

4.0 ml

Solution B

Mannitol

5.0 g

Water

100.0 ml

Prepare the solution ready for use by mixing solution A and B

Examp. Formulation

No.

Le A 23 890

75 Solution A
 Solution A from Example 74
Solution B
 Glucose 25.0 g
 Water ad 250.0 ml
 Prepare the solution ready for use by mixing solution A and B

76 Solution A
 Solution A from Example 74
Solution B
 Sodium chloride 4.5 g
 Water ad 500.0 ml
 Prepare the solution ready for use by mixing solution A and B

77 Solution A
 Ciprofloxacin 100 mg 302 1.00
 Lactic acid 20% (w/w) 168 mg 373 1.24
 Water ad 1.00 ml
Solution B
 Sodium chloride 0.45 g
 Water ad 50.0 ml
 Prepare the solution ready for use by mixing solution A and B

Micromole R

LE A 23 890

Examp.
No.

78

Solution A
Solution A from Example 77

Solution B
Ringer lactate solution: 100 ml (for example Ringer Lactate DAB 7
supplied by Braun Melsungen
Rote Liste 1985 No. 51013)

Prepare the solution ready for use by mixing solution A and B

79

Solution A
Ciprofloxacin 200 mg
Lactic acid 25% (w/w) 257mg
Water ad 1.00 ml

604 1.00
715 1.18

Solution B

Sodium chloride 0.9 g
Water ad 100.0 ml

Prepare the solution ready for use by mixing solution A and B

Le A	Examp. No.	Formulation	Micromole R
23	890		
	80	<u>Solution A</u>	
		Ciprofloxacin	528
		Lactic acid 20% (w/w)	667
		Water	1.00
		175 mg	1.26
		300 mg	
		ad 0.5 ml	
		<u>Solution B</u>	
		Glucose	2.5 g
		Water	50.0 ml
		ad	
		Prepare this solution ready for use by mixing solution A and B	
	81	<u>Solution A</u>	
		Ciprofloxacin	604
		Lactic acid 20% (w/w)	715
		Hydrochloric acid	90
		Sodium chloride	0.15
		Water	
		200 mg	
		322 mg	
		3.3 mg	
		0.9 g	
		ad 100.0 ml	
		<u>Solution B</u>	
		Solution for supplying amino acids:500 ml (for example Aminoplasmal LS-5	
		electrolyte-free, supplied by Braun	
		Melsungen, Rote Liste 1985 No. 51238)	
		Prepare the solution ready for use by mixing solution A and B	

Micromole R

Examp. Formulation

No.

82

Solution A

Ciprofloxacin
Lactic acid 2% (w/w)
Phosphoric acid
Water

100 mg
1481 mg
16 mg
10.0 ml
ad
pH: 3.8

302
329
165

1.00
1.09
0.50

Solution B

Sodium chloride
Water

3.6 g
400 ml
ad

Prepare the solution ready for use by mixing Solution A and B

83

Solution A

Ciprofloxacin lactate

254 mg

604

1.00 (con-
taining an
equimolar
amount of
lactic acid)

Lactic acid 0.5% (w/w)
Methanesulphonic acid
0.2 M sodium hydroxide solution
Water

979 mg
116 mg
2.4 ml
20.0 ml
ad
pH: 3.8

54
1208
480

Solution B

Glucose
Water

20.0 g
400.0 ml
ad

Prepare the solution ready for use by mixing solution A and B

Micromole R

Examp. Formulation

A No.

23				
84	<u>Lyophilizate</u>			
890	Ciprofloxacin	100 mg	302	1.00
	<u>Solution</u>			
	Lactic acid 10% (w/w)	322 mg	358	1.18
	Hydrochloric acid	1.6 mg	45	0.15
	Sodium chloride	900 mg		
	Water	ad 100.0 ml		
	Prepare solution ready for use by dissolving the lyophilizate in the solution			

85

Lyophilizate

Ciprofloxacin lactate

63.6 mg

151

1.00 (containing an equimolar amount of lactic acid)

Solution

Lactic acid 0.005% (w/w)

10.87 g

6

1.04

Sodium chloride

225 mg

Water

ad 25 ml

Prepare solution ready for use by dissolving the lyophilizate in the solution

Micromole R

Examp. No.	Formulation		
86	<u>Lyophilizate</u> Ciprofloxacin lactate	127 mg	302
	Lactic acid	1.1 mg	12
	Mannitol	5.0 g	0.04
	<u>Solution</u> Water	ad 100.0 ml	
	Prepare solution ready for use by dissolving the lyophilizate in the solution		
87	<u>Lyophilizate</u> Ciprofloxacin	100 mg	302
	<u>Solution</u> Lactic acid 20% (w/w)	186 mg	413
	0.1 M hydrochloric acid solution	0.42 ml	142
	Glucose	25.0 g	
	Water	ad 500.0 ml	
	Prepare solution ready for use by dissolving the lyophilizate in the solution		

Le A 23 890

Le A 23 890	Examp. No.	Formulation	Micromole R
	88	<u>Lyophilizate</u> Ciprofloxacin lactate	151
		63.6 mg	1.00 (containing an equimolar amount of lactic acid)
		<u>Solution</u>	
		Lactic acid 0.2% (w/w)	721
		Mannitol	4.77
		0.2 M sodium hydroxide solution	-
		Water ad 333 ml	178
		Prepare solution ready for use by dissolving the lyophilizate in the solution	
	89	<u>Lyophilizate</u> Ciprofloxacin Sodium chloride	604
		200 mg 900 mg	1.00
		<u>Solution</u>	
		Lactic acid 2% (w/w)	715
		Hydrochloric acid	90
		Water ad 100.0 ml	0.18 0.15
		Prepare solution ready for use by dissolving the lyophilizate in the the solution	

Le	Examp. No.	Formulation	Micromole R
23	92	<u>Lyophilizate</u> Ciprofloxacin lactate	604
893)		222 mg	1.00 (containing an equimolar amount of lactic acid)
		Lactic acid	111
		10 mg	
		Mannitol	
		5.0 g	
		<u>Solution</u>	
		Water	
		100.0 ml	
		Prepare solution ready for use by dissolving the lyophilizate in the solution	
	93	<u>Lyophilizate</u> Lyophilisate from Example 92	
		<u>Solution</u>	
		Water	
		200.0 ml	
		Prepare solution ready for use by dissolving the lyophilizate in the solution	

Examp. No.	Formulation	Micromole R
94	<p><u>Lyophilizate</u> Ciprofloxacin lactate</p> <p>Lactic acid <u>Solution</u> Ringer lactate solution:</p>	<p>1510</p> <p>355</p> <p>100 ml (for example Ringer lactate DAB 7 supplied by Braun Melsungen Rote Liste 1985 No. 51013)</p>
		<p>1.00 (containing an equimolar amount of lactic acid) 1.24</p>
		<p>32 mg</p> <p>636 mg</p>

Prepare solution ready for use by dissolving the lyophilizate in the solution

95	<p><u>Lyophilizate</u> Lyophilizate from Example 94 <u>Solution</u> Sodium chloride Water</p>	<p>0.9 g ad 100.0 ml</p>
		<p>Prepare solution ready for use by dissolving the lyophilizate in the solution</p>

Examp. Formulation

No.

96

Lyophilizate

Lyophilizate from Example 94

Solution

Glucose

12.5 g

Water

ad 250,0 ml

Prepare solution ready for use by dissolving the Lyophilizate in the solution

97

Lyophilizate

Lyophilizate from Example 94

Solution

Solution for supplying amino acids: 500 ml (for example aminoplasmal LS-5
electrolyte-free, supplied by Braun
Melsungen, Rote Liste 1985 No. 51238)

Prepare solution ready for use by dissolving the Lyophilizate in the solution

Le A 23 890	Examp. No.	Formulation	Micromole R
	98	<u>Lyophilizate</u> Ciprofloxacin 100 mg <u>Solution</u> Malic acid 41 mg Sorbitol 2,5 g Water ad 50,0 ml Prepare solution ready for use by dissolving the lyophilizate in the solution	302 1.00 306 1.01
	99	<u>Lyophilizate</u> Ciprofloxacin lactate 127 mg <u>Solution</u> Lactic acid 20% (w/w) 12 mg Propionic acid 45 mg Sodium chloride 4,5 g Water ad 500 ml Prepare solution ready for use by dissolving the lyophilizate in the solution	302 1.00 (containing equivalent amount of lactic acid) 27 1.09 608 2.01

26419

It will be understood that the specification and examples are illustrative but not limitative of the present invention and that other embodiments within the spirit and scope of the invention will suggest themselves to those skilled in the art.

What is claimed is:

1. An aqueous infusion solution containing 0.015 to 0.5 g of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acid per 100 ml of aqueous solution and an amount of at least one physiologically tolerated acid which suffices to dissolve the active compound.
2. An infusion solution according to claim 1, containing a mixture of physiologically tolerated acids to dissolve the active compound.
3. An infusion solution according to claim 1, wherein the acid is at least one member selected from the group consisting of hydrochloric acid, methanesulphonic acid, propionic acid, succinic acid, glutaric acid, citric acid, fumaric acid, maleic acid, tartaric acid, glutamic acid, gluconic acid, glucuronic acid, galacturonic acid, ascorbic acid, phosphoric acid, nitric acid, acetic acid, malic acid, L-aspartic acid and lactic acid.
4. An infusion solution according to claim 1, wherein the acid is selected from the group consisting of lactic acid, hydrochloric acid and mixtures of lactic acid and hydrochloric acid.

26419

5. An infusion solution according to claim 1, containing up to 5 mols per mol of active compound, of the physiologically tolerated acids.
6. An infusion solution according to claim 1, containing 0.9 to 5 mols per mol of active compound, of the physiologically tolerated acids.
7. An infusion solution according to claim 1, containing 1.04 to 2.2 mols per mol of active compound, of the physiologically tolerated acids.
8. An infusion solution according to claim 1, containing up to 5 mols per mol of active compound, of lactic acid.
9. An infusion solution according to claim 1, containing 0.99 to 1.50 mols of lactic acid per mol of active compound.
10. An infusion solution according to claim 1, containing 1.12 to 1.24 mols of lactic acid per mol of active compound.
11. An infusion solution according to claim 1, containing 0.9 to 5 mols per mol of active compound, of lactic acid plus another physiologically tolerated acid.

26419

12. An infusion solution according to claim 1, having a pH from 3 to 5.2.
13. An infusion solution according to claim 1, having a pH from 3.6 to 4.7.
14. An infusion solution according to claim 1, having a pH from 3.9 to 4.5.
15. An infusion solution according to claim 1, having a pH from 4.1 to 4.3.
16. An infusion solution according to claim 1, which is substantially isotonic.
17. An infusion solution according to claim 1, containing 0.99 to 1.5 mols of lactic acid and 0.0 to 0.80 mol of hydrochloric acid per mol of active compound, and relative to 100 ml of solution, 0.6 to 2.2 g of NaCl.
18. An infusion solution according to claim 1, containing 1.04 to 1.4 mols of lactic acid and 0.0 to 0.80 mol of hydrochloric acid per mol of active compound, and relative to 100 ml of solution, 0.75 to 1.2 g of NaCl.
19. An infusion solution according to claim 1, present in a container in from 40 to 600 ml.