

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2011342284 C1

(54) Title
Cyclosporine analogue molecules modified at amino acid 1 and 3

(51) International Patent Classification(s)
C07K 7/64 (2006.01) **A61K 38/13** (2006.01)
A61K 38/12 (2006.01) **C07K 1/113** (2006.01)

(21) Application No: **2011342284** (22) Date of Filing: **2011.12.14**

(87) WIPO No: **WO12/079172**

(30) Priority Data

(31) Number (32) Date (33) Country
61/423,576 **2010.12.15** **US**

(43) Publication Date: **2012.06.21**
(44) Accepted Journal Date: **2017.02.16**
(44) Amended Journal Date: **2017.07.13**

(71) Applicant(s)
ContraVir Pharmaceuticals, Inc.

(72) Inventor(s)
Hegmans, Alexander;Fenske, Bruce W.;Trepanier, Dan J.;Abel, Mark D.;Sugiyama, Shin;Ure, Daren R.

(74) Agent / Attorney
Davies Collison Cave Pty Ltd, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000, AU

(56) Related Art
EP 0 194 972 B1
US 7,696,166 B2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2012 (21.06.2012)

(10) International Publication Number
WO 2012/079172 A1

(51) International Patent Classification:
C07K 7/64 (2006.01) *A61K 38/13* (2006.01)
A61K 38/12 (2006.01) *C07K 1/13* (2006.01)

(21) International Application Number:
PCT/CA2011/050773

(22) International Filing Date:
14 December 2011 (14.12.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/423,576 15 December 2010 (15.12.2010) US

(71) Applicant (for all designated States except US): ISO-TECHNIKA PHARMA INC. [CA/CA]; 5120 - 75th Street, Edmonton, Alberta T6E 6W2 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HEGMANS, Alexander [CA/CA]; 5120 - 75th Street, Edmonton, Alberta T6E 6W2 (CA). FENSKE, Bruce, W. [CA/CA]; 5120 - 75th Street, Edmonton, Alberta T6E 6W2 (CA). TRE-PANIER, Dan, J. [CA/CA]; 5120 - 75th Street, Edmonton, Alberta T6E 6W2 (CA). ABEL, Mark, D. [CA/CA]; 5120 - 75th Street, Edmonton, Alberta T6E 6W2 (CA). SUGIYAMA, Shin [CA/CA]; 5120 - 75th Street, Edmonton, Alberta T6E 6W2 (CA). URE, Daren R. [CA/CA]; 3912 - 149A Street, Edmonton, Alberta T6R 1J8 (CA).

(74) Agent: RIDOUT & MAYBEE LLP; 100 Murray Street, 4th Floor, Ottawa, Ontario K1N 0A1 (CA).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,


[Continued on next page]

(54) Title: CYCLOSPORINE ANALOGUE MOLECULES MODIFIED AT AMINO ACID 1 AND 3

Table 5

Figure 1A

1-Mod Name	Modification:	1-Mod Structure		1 + 3-D-MeAla	
		Cyp A Inhibition Fold-potency vs CsA	Immunosuppression % vs CsA	Cyp A Inhibition Fold-potency vs CsA	Immunosuppression % vs CsA
404-58		0.16	<1	1.3	<1
420-171		2.3		8.7	<1
420-19		3.7	<1	9.5	3.4
416-08		3.7	<1	14.3	3
404-98		2.2	6.5	15	7.6



(57) Abstract: Analogues of cyclosporin-A are disclosed comprising modifications of the substituents as the positions of amino acids 1 and 3, according to the following Formula (I). The disclosed compounds include compounds having affinity for cyclophilin, including cyclophilin-A, and reduced immunosuppressivity in comparison with cyclosporin-A and analogs thereof modified solely at position 1.

WO 2012/079172 A1



GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD,
RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ,
DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT,
LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS,

SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

**CYCLOSPORINE ANALOGUE MOLECULES MODIFIED at AMINO ACID 1
AND 3**

FIELD OF THE INVENTION

The present invention relates to novel analogs of molecules belonging to the
5 cyclosporine family, including analogs of Cyclosporine A (CsA) and including
analogs that have reduced or no immunosuppressive activity and bind
cyclophilin (CyP).

BACKGROUND OF THE INVENTION

10

Cyclosporines are members of a class of cyclic polypeptides having potent
immunosuppressant activity. At least some of these compounds, such as
Cyclosporine A (CsA), are produced by the species *Tolyphocladium inflatum* as
secondary metabolites. CsA is a potent immunosuppressive agent that has
15 been demonstrated to suppress humoral immunity and cell-mediated immune
reactions, such as allograft rejection, delayed hypersensitivity, experimental
allergic encephalomyelitis, Freund's adjuvant arthritis and graft versus host
disease. It is used for the prophylaxis of organ rejection in organ transplants; for
the treatment of rheumatoid arthritis; and for the treatment of psoriasis.

20

Although a number of compounds in the cyclosporine family are known, CsA is
perhaps the most widely used medically. The immunosuppressive effects of
CsA are related to the inhibition of T-cell mediated activation events.
Immunosuppression is accomplished by the binding of cyclosporine to a
25 ubiquitous intracellular protein called cyclophilin (CyP). This complex, in turn,
inhibits the calcium and calmodulin-dependent serine-threonine phosphatase
activity of the enzyme calcineurin. Inhibition of calcineurin prevents the
activation of transcription factors, such as NFAT_{p/c} and NF- κ B, which are
necessary for the induction of cytokine genes (*IL-2*, *IFN- γ* , *IL-4*, and *GM-CSF*)
30 during T-cell activation.

- 2 -

Since the original discovery of cyclosporine, a wide variety of naturally occurring cyclosporines have been isolated and identified. Additionally, many cyclosporines that do not occur naturally have been prepared by partial or total synthetic means, and by the application of modified cell culture techniques.

5 Thus, the class comprising cyclosporines is substantial and includes, for example, the naturally occurring cyclosporines A through Z; various non-naturally occurring cyclosporine derivatives; artificial or synthetic cyclosporines including the dihydro- and iso-cyclosporines; derivatized cyclosporines (for example, either the 3'-O-atom of the MeBmt residue may be acylated, or a further substituent may be introduced at the sarcosyl residue at the 3-position); cyclosporines in which the MeBmt residue is present in isomeric form (e.g., in which the configuration across positions 6' and 7' of the MeBmt residue is *cis* rather than *trans*); and cyclosporines wherein variant amino acids are incorporated at specific positions within the peptide sequence.

10

15 Cyclosporine analogues containing modified amino acids in the 1-position are disclosed in WO 99/18120 and WO 03/033527, which are incorporated herein by reference in their entirety. These applications describe a cyclosporine derivative known as "ISA_{TX}247" or "ISA247" or "ISA." This analog is structurally identical to CsA, except for modification at the amino acid-1 residue. Applicants have previously discovered that certain mixtures of *cis* and *trans* isomers of ISA247, including mixtures that are predominantly comprised of *trans* ISA247, exhibited a combination of enhanced immunosuppressive potency and reduced toxicity over the naturally occurring and presently known cyclosporines.

20

25 Cyclosporine has three well established cellular targets; calcineurin, the CyP isoforms (which includes but is not limited to CyP-A, CyP-B and CyP-D), and P-glycoprotein (PgP). The binding of cyclosporine to calcineurin results in significant immunosuppression and is responsible for its traditional association with transplantation and autoimmune indications.

30

The Cyclophilin Family

CyPs (Enzyme Commission (EC) number 5.1.2.8) belong to a group of proteins that have peptidyl-prolyl cis-trans isomerase activity; such proteins are collectively known as immunophilins and also include the FK-506-binding 5 proteins and the parvulins. CyPs are found in all cells of all organisms studied, in both prokaryotes and eukaryotes and are structurally conserved throughout evolution. There are 7 major CyPs in humans, namely CyP-A, CyP-B, CyP-C, CyP-D, CyP-E, CyP-40, and CyP-NK (first identified from human natural killer cells), and a total of 16 unique proteins (Galat A. Peptidylprolyl *cis/trans* 10 isomerases (immunophilins): biological diversity - targets - functions. *Curr Top Med Chem* 2003, 3:1315-1347; Waldmeier PC et al. Cyclophilin D as a drug target. *Curr Med Chem* 2003, 10:1485-1506).

The first member of the CyPs to be identified in mammals was CyP-A. CyP-A is 15 an 18-kDa cytosolic protein and is the most abundant protein for CsA binding. It is estimated that CyP-A makes up 0.6% of the total cytosolic protein (Mikol V et al. X-ray structure of monmeric cyclophilin A-cycloporin A crystal complex at 2.1 A resolution. *J. Mol. Biol.* 1993, 234:1119-1130; Galat A, Metcalfe SM. Peptidylproline *cis/trans* isomerases. *Prog. Biophys. Mol. Biol.* 1995, 63:67-118).

20

Cellular Location of Cyclophilins

CyPs can be found in most cellular compartments of most tissues and encode unique functions. In mammals, CyP-A and CyP-40 are cytosolic whereas CyP-B and CyP-C have amino-terminal signal sequences that target them to the 25 endoplasmic reticulum protein secretory pathway (reviewed in Galat, 2003; Dornan J et al. Structures of immunophilins and their ligand complexes. *Curr Top Med Chem* 2003, 3:1392-1409). CyP-D has a signal sequence that directs it to the mitochondria (Andreeva L et al. Cyclophilins and their possible role in the stress response. *Int J Exp Pathol* 1999, 80:305-315; Hamilton GS et al. 30 Immunophilins: beyond immunosuppression. *J Med Chem* 1998, 41:5119-5143); CyP-E has an amino-terminal RNA-binding domain and is localized in the nucleus (Mi H et al. A nuclear RNA-binding cyclophilin in human T cells. *FEBS*

- 4 -

Lett 1996, 398:201-205) and CyP-40 has TPRs and is located in the cytosol (Kieffer LJ et al. Cyclophilin-40, a protein with homology to the P59 component of the steroid receptor complex. Cloning of the cDNA and further characterization. *J Biol Chem* 1993, 268:12303-12310). Human CyP-NK is the 5 largest CyP, with a large, hydrophilic and positively charged carboxyl terminus, and is located in the cytosol (Anderson SK et al. A cyclophilin-related protein involved in the function of natural killer cells. *Proc Natl Acad Sci USA* 1993, 90:542-546; Rinfret A et al. The N-terminal cyclophilin-homologous domain of a 150-kilodalton tumor recognition molecule exhibits both peptidylprolyl *cis*-10 *trans*isomerase and chaperone activities. *Biochemistry* 1994, 33:1668-1673)

Function and Activity of the Cyclophilins

CyPs are multifunctional proteins that are involved in many cellular processes. Because CyPs were highly conserved throughout evolution, this suggests an 15 essential role for CyPs. Initially, it was found that CyPs have the specific enzymatic property of catalyzing *cis-trans* isomerization of peptidyl-prolyl bonds (Galat, 1995; Fisher GA et al. A phase I study of paclitaxel (taxol) (T) in combination with SDZ valspodar, a potent modulator of multidrug resistance (MDR). *Anticancer Drugs*. 1994; 5(Suppl 1): 43). Thus, CyPs are called peptidyl-20 prolyl-*cis-trans* isomerase (PPIase), which can act as an acceleration factor in the proper folding of newly synthesized proteins. PPIases are also involved in repairing damaged proteins due to environmental stresses including thermal stress, ultraviolet irradiation, changes in the pH of the cell environment, and treatment with oxidants. This function is known as molecular chaperoning 25 activity. (Yao Q et al. Roles of Cyclophilins in Cancers and Other Organs Systems. *World J. Surg.* 2005, 29: 276-280)

In addition, the PPIase activity of CyPs has recently been shown to be involved 30 in diverse cellular processes, including intracellular protein trafficking (Andreeva, 1999; Caroni P et al. New member of the cyclophilin family associated with the secretory pathway. *J Biol Chem* 1991, 266:10739-42), mitochondrial function (Halestrap AP et al. CsA binding to mitochondrial cyclophilin inhibits the

- 5 -

permeability transition pore and protects hearts from ischaemia/reperfusion injury. *Mol Cell Biochem* 1997, 174:167–72; Connern CP, Halestrap AP. Recruitment of mitochondrial cyclophilin to the mitochondrial inner membrane under conditions of oxidative stress that enhance the opening of a calcium-
5 sensitive non-specific channel. *Biochem J* 1994, 302:321–4), pre-mRNA processing (Bourquin JP et al. A serine/arginine-rich nuclear matrix cyclophilin interacts with the C-terminal domain of RNA polymerase II. *Nucleic Acids Res* 1997, 25:2055–61), and maintenance of multiprotein complex stability (Andreeva, 1999).

10 Cyclosporine binds with nanomolar affinity to CyP-A via contacts within the hydrophobic pocket (Colgan J et al. Cyclophilin A-Deficient Mice Are Resistant to Immunosuppression by Cyclosporine. *The Journal of Immunology* 2005, 174: 6030–6038, Mikol, 1993) and inhibits PPIase activity. However, this effect is
15 thought to be irrelevant for the immunosuppression. Rather, the complex between CsA and CyP-A creates a composite surface that binds to and prevents calcineurin from regulating cytokine gene transcription (Friedman J et al. Two cytoplasmic candidates for immunophilin action are revealed by affinity for a new cyclophilin: one in the presence and one in the absence of CsA. *Cell* 1991, 66: 799–806; Liu J et al. Calcineurin is a common target of cyclophilin-CsA and FKBP-FK506 complexes. *Cell* 1991, 66: 807–815).

Homology of the Cyclophilins

CyP-A, the prototypical member of the family, is a highly conserved protein in
25 mammalian cells (Handschumacher RE et al. Cyclophilin: a specific cytosolic binding protein for CsA. *Science* 1984, 226: 544–7). Sequence homology analysis of human CyP-A shows that it is highly homologous to human CyP-B, CyP-C, and CyP-D (Harding MW, Handschumacher RE, Speicher DW. Isolation and amino acid sequence of cyclophilin. *J Biol Chem* 1986, 261:8547–55). The
30 cyclosporine binding pocket of all CyPs is formed by a highly conserved region of approximately 109 amino acids. Of the known CyPs, CyP-D has the highest homology to CyP-A. In fact, in this region the sequence identity is 100%

- 6 -

between CyP-A and CyP-D (Waldmeier 2003; Kristal BS et al. The Mitochondrial Permeability Transition as a Target for Neuroprotection. *Journal of Bioenergetics and Biomembranes* 2004, 36(4); 309-312). Therefore, CyP-A affinity is a very good predictor of CyP-D affinity, and visa versa (Hansson MJ et 5 al. The Nonimmunosuppressive Cyclosporine analogues NIM811 and UNIL025 Display Nanomolar Potencies on Permeability Transition in Brain-Derived Mitochondria. *Journal of Bioenergetics and Biomembranes*, 2004, 36(4): 407-413). This relationship has been repeatedly demonstrated empirically with Cyclosporine analogues (Hansson, 2004; Ptak RG et al. Inhibition of Human 10 Immunodeficiency Virus Type 1 Replication in Human Cells by Debio-025, a Novel Cyclophilin Binding Agent. *Antimicrobial Agents and Chemotherapy* 2008: 1302-1317; Millay DP et al. Genetic and pharmacologic inhibition of mitochondrial dependent necrosis attenuates muscular dystrophy. *Nature Medicine* 2008, 14(4): 442-447; Harris R et al. The Discovery of Novel Non- 15 Immunosuppressive Cyclosporine Ethers and Thioethers With Potent HCV Activity. Poster # 1915, *59th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD)*, 2008). The sequence homology across the CyPs suggests that all CyPs are potential targets for Cyclosporine analogues. Because of the multitude of cellular processes in which the CyPs are involved, 20 this further suggests that CsA analogues that retain significant binding to CyP can be useful in the treatment of many disease indications.

Cyclophilin Mediated Diseases

Human Immunodeficiency Virus (HIV):

25 HIV is a lentivirus of the retrovirus family and serves as an example of the involvement of CyP in the process of infection and replication of certain viruses. CyP-A was established more than a decade ago to be a valid target in anti-HIV chemotherapy (Rosenwirth BA et al. Cyclophilin A as a novel target in anti-HIV-1 chemotherapy. *Int. Antivir. News* 1995, 3:62-63). CyP-A fulfills an essential 30 function early in the HIV-1 replication cycle. It was found to bind specifically to the HIV-1 Gag polyprotein (Luban JKL et al. Human immunodeficiency virus type 1 Gag protein binds to cyclophilins A and B. *Cell* 1993, 73: 1067-1078). A

defined amino acid sequence around G89 and P90 of capsid protein p24 (CA) was identified as the binding site for CyP-A (Bukovsky AAA et al. Transfer of the HIV-1 cyclophilin-binding site to simian immunodeficiency virus from Macaca mulatta can confer both cyclosporine sensitivity and cyclosporine dependence.

5 *Proc. Natl. Acad. Sci. USA* 1997, 94: 10943–10948; Gamble TRF et al. Crystal structure of human cyclophilin A bound to the amino-terminal domain of HIV-1 capsid. *Cell* 1996, 87: 1285–1294). The affinity of CyP-A for CA promotes the incorporation of CyP-A into the virion particles during assembly (Thali MA et al. Functional association of cyclophilin A with HIV-1 virions. *Nature* 1994, 372: 363–365). Experimental evidence indicates that the CyP-A-CA interaction is essential for HIV-1 replication; inhibition of this interaction impairs HIV-1 replication in human cells (Hatzlioannou TD et al. Cyclophilin interactions with incoming human immunodeficiency virus type 1 capsids with opposing effects on infectivity in human cells. *J. Virol.* 2005, 79: 176–183; Steinkasserer AR et al.

10 15 Mode of action of SDZ NIM 811, a nonimmunosuppressive CsA analog with activity against human immunodeficiency virus type 1 (HIV-1): interference with early and late events in HIV-1 replication. *J. Virol* 1995, 69: 814–824). The step in the viral replication cycle where CyP-A is involved was demonstrated to be an event after penetration of the virus particle and before integration of the double-

20 25 stranded viral DNA into the cellular genome (Braaten DEK et al. Cyclophilin A is required for an early step in the life cycle of human immunodeficiency virus type 1 before the initiation of reverse transcription. *J. Virol* 1996 70: 3551–3560; Mlynar ED et al. The non-immunosuppressive CsA analogue SDZ NIM 811 inhibits cyclophilin A incorporation into virions and virus replication in human immunodeficiency virus type 1-infected primary and growth-arrested T cells. *J. Gen. Virol* 1996, 78: 825–835; Steinkasserer, 1995). The anti-HIV-1 activity of CsA was first reported in 1988 (Wainberg MA et al. The effect of CsA on infection of susceptible cells by human immunodeficiency virus type 1. *Blood* 1998, 72: 1904–1910). Evaluation of CsA and many derivatives for inhibition of

30 HIV-1 replication revealed that nonimmunosuppressive CsA analogs had anti-HIV-1 activities equal to or even superior to those of immunosuppressive analogs (Bartz SRE et al. Inhibition of human immunodeficiency virus replication

by nonimmunosuppressive analogs of CsA. *Proc. Natl. Acad. Sci. USA* 1995; 92: 5381–5385; Billich AF et al. Mode of action of SDZ NIM 811, a nonimmunosuppressive CsA analog with activity against human immunodeficiency virus (HIV) type 1: interference with HIV protein-cyclophilin A interactions. *J. Virol* 1995; 69: 2451–2461; Ptak, 2008).

Inflammation

Inflammation in disease involves the influx of leukocytes (white blood cells) to the area of infection. The leukocytes are drawn to the area by chemokines, a family of chemoattracting agents. *In vitro* studies have shown that extracellular CyP-A is a potent chemoattractant for human leukocytes and T cells (Kamalpreet A et al. Extracellular cyclophilins contribute to the regulation of inflammatory responses *Journal of Immunology* 2005; 175: 517–522; Yurchenko VG et al. Active-site residues of cyclophilin A are crucial for its signaling activity via CD147. *J. Biol. Chem.* 2002; 277: 22959–22965; Xu QMC et al. Leukocyte chemotactic activity of cyclophilin. *J. Biol. Chem.* 1992; 267: 11968–11971; Allain FC et al. Interaction with glycosaminoglycans is required for cyclophilin B to trigger integrin-mediated adhesion of peripheral blood T lymphocytes to extracellular matrix. *Proc. Natl. Acad. Sci. USA* 2002; 99: 2714–2719).

Furthermore, CyP-A can induce a rapid inflammatory response, characterized by leukocyte influx, when injected *in vivo* (Sherry BN et al. Identification of cyclophilin as a proinflammatory secretory product of lipopolysaccharide-activated macrophages. *Proc. Natl. Acad. Sci. USA* 1992; 89: 3511–3515). CyP-A is ubiquitously distributed intracellularly, however, during the course of inflammatory responses, CyP-A is released into extracellular tissue spaces by both live and dying cells (Sherry, 1992). Indeed, elevated levels of CyP-A have been reported in several different inflammatory diseases, including sepsis, rheumatoid arthritis, and vascular smooth muscle cell disease (Jin ZG et al. Cyclophilin A is a secreted growth factor induced by oxidative stress. *Circ. Res.* 2000; 87: 789–796; Teger, 1997; Billich, 1997). In the case of rheumatoid arthritis, a direct correlation between levels of CyP-A and the number of

- 9 -

neutrophils in the synovial fluid of rheumatoid arthritis patients was reported (Billich, 1997).

Cancer

5 CyP-A has recently been shown to be over-expressed in many cancer tissues and cell lines, including but not limited to small and non-small cell lung, bladder, hepatocellular, pancreatic and breast cancer (Li, 2006; Yang H et al. Cyclophilin A is upregulated in small cell lung cancer and activates ERK1/2 signal. *Biochemical and Biophysical Research Communications* 2007; 361: 763–767;

10 Campa, 2003). In cases where exogenous CyP-A was supplied this was shown to stimulate the cancer cell growth (Li, 2006; Yang, 2007) while CsA arrested the growth (Campa, 2003). Most recently it has been demonstrated the CyP (A and B) is intricately involved in the biochemical pathway allowing growth of human breast cancer cells and that CyP knockdown experiments decreased the cancer

15 cell growth, proliferation and motility (Fang F et al. The expression of Cyclophilin B is Associated with Malignant Progression and Regulation of Genes Implicated in the Pathogenesis of Breast Cancer. *The American Journal of Pathology* 2009; 174(1): 297-308; Zheng J et al. Prolyl Isomerase Cyclophilin A Regulation of Janus-Activated Kinase 2 and the Progression of Human Breast Cancer. *Cancer Research* 2008; 68 (19): 7769-7778). Most interestingly, CsA treatment of mice

20 xenografted with breast cancer cells induced tumor necrosis and completely inhibited metastasis (Zheng, 2008). The researchers conclude that "Cyclophilin B action may significantly contribute to the pathogenesis of human breast cancer" and that "cyclophilin inhibition may be a novel therapeutic strategy in the

25 treatment of human breast cancer" (Fang, 2009; Zheng, 2008).

Hepatitis C

Hepatitis C Virus (HCV) is the most prevalent liver disease in the world and is considered by the World Health Organization as an epidemic. Because HCV can

30 infect a patient for decades before being discovered, it is often called the "silent" epidemic. Studies suggest that over 200 million people worldwide are infected with HCV, an overall incidence of around 3.3% of the world's population. In the

- 10 -

US alone, nearly 4 million people are or have been infected with HCV and of these; 2.7 million have an ongoing chronic infection. All HCV infected individuals are at risk of developing serious life-threatening liver diseases. Current standard therapy for chronic hepatitis C consists of the combination of pegylated

5 interferon in combination with ribavirin, both generalized anti-viral agents (Craxi A et al. Clinical trial results of peginterferons in combination with ribavirin. *Semin Liver Dis* 2003; 23(Suppl 1): 35-46). Failure rate for the treatment is approximately 50 % (Molino BF. Strategic Research Institute: 3rd annual viral hepatitis in drug discovery and development world summit 2007. *AMRI Technical*

10 *Reports*; 12(1)).

It has recently been demonstrated that CyP-B is critical for the efficient replication of the HCV genome (Watashi K et al. Cyclophilin B Is a Functional Regulator of Hepatitis C Virus RNA Polymerase. *Molecular Cell* 2005, 19: 111-

15 122). Viruses depend on host-derived factors such as CyP-B for their efficient genome replication. CyP-B interacts with the HCV RNA polymerase NS5B to directly stimulate its RNA binding activity. Both the RNA interference (RNAi)-mediated reduction of endogenous CyP-B expression and the induced loss of NS5B binding to CyP-B decreases the levels of HCV replication. Thus, CyP-B

20 functions as a stimulatory regulator of NS5B in HCV replication machinery. This regulation mechanism for viral replication identifies CyP-B as a target for antiviral therapeutic strategies.

Unlike other HCV treatments, CyP inhibition does not directly target the HCV virus. It is therefore thought that resistance to CyP binding drugs will occur more slowly than current HCV treatment drugs (Manns MP, et al. The way forward in HCV treatment-finding the right path. *Nature Reviews Drug Discovery* 2007; 6: 991-1000). In addition, by interfering at the level of host-viral interaction, CyP inhibition may open the way for a novel approach to anti-HCV treatment that

25 could be complementary, not only to interferon-based treatment, but also to future treatments that directly target HCV replication enzymes such as protease and polymerase inhibitors (Flisiak R, Dumont JM, Crabbé R. Cyclophilin

- 11 -

inhibitors in hepatitis C viral infection. *Expert Opinion on Investigational Drugs* 2007, 16(9): 1345-1354). Development of new anti-HCV drugs effecting HCV viral replication has been significantly impeded by the lack of a suitable laboratory HCV model. This obstacle has only recently been overcome by the 5 development of several suitable cell culture models (Subgenomic HCV Replicon Systems) and a mouse model containing human liver cells (Goto K, et al. Evaluation of the anti-hepatitis C virus effects of cyclophilin inhibitors, CsA, and NIM811. *Biochem Biophys Res Comm* 2006; 343: 879-884; Mercer DF, et al. Hepatitis C virus replication in mice with chimeric human livers. *Nat Med* 2001; 7: 10 927-933). Cyclosporine has recently demonstrated anti-HCV activity in screening models and in small clinical trials (Watashi K, et al. CsA suppresses replication of hepatitis C virus genome in cultured hepatocytes. *Hepatology* 2003; 38:1282-1288; Inoue K, Yoshioka M. Interferon combined with cyclosporine treatment as an effective countermeasure against hepatitis C virus recurrence in liver 15 transplant patients with end-stage hepatitis C virus related disease. *Transplant Proc* 2005; 37:1233-1234).

Muscular Degenerative Disorders

CyP-D is an integral part of the mitochondrial permeability transition pore (MTP) 20 in all cells. The function of the MTP pore is to provide calcium homeostasis within the cell. Under normal conditions the opening and closing of the MTP pore is reversible. Under pathological conditions that involve an excessive calcium influx into the cell, this overloads the mitochondria and induces an irreversible opening of the MPT pore, leading to cell death or apoptosis. CsA has been 25 reported to correct mitochondrial dysfunction and muscle apoptosis in patients with Ullrich congenital muscular dystrophy and Bethlem myopathy [(Merlini L et al. CsA corrects mitochondrial dysfunction and muscle apoptosis in patients with collagen VI myopathies. *PNAS* 2008; 105(13): 5225-5229]. CsA has been demonstrated *in vitro* to dose dependently inhibit MTP opening in isolated 30 cardiac mitochondria, thereby preventing apoptosis and allowing the cell precious time for repair (Gomez L et al. Inhibition of mitochondrial permeability transition improves functional recovery and reduces mortality following acute

- 12 -

myocardial infarction in mice *Am J Physiol Heart Circ Physiol* 2007, 293: H1654–H1661). A clinical study in 58 patients who presented with acute myocardial infarction demonstrated that administration of CsA at the time of reperfusion was associated with a smaller infarct than that seen with placebo (Piot C et al. Effect 5 of Cyclosporine on Reperfusion Injury in Acute Myocardial Infarction. *New England Journal of Medicine* 2008; 395(5): 474-481)).

Chronic Neurodegenerative Diseases

CsA can act as a neuroprotective agent in cases of acute cerebral ischemia and 10 damage as a result of head trauma (Keep M, et al. Intrathecal cyclosporine prolongs survival of late-stage ALS mice. *Brain Research* 2001; 894: 327–331). Animals treated with CsA showed a dramatic 80 % survival rate relative to only a 15 10 % survival rate in the absence of treatment. It was later established that this was largely the result of the binding of CsA to mitochondrial CyP-D. It has been subsequently established that the utility of CsA extends to chronic 20 neurodegeneration, as was subsequently demonstrated in a rat model of Lou Gerhig's Disease (ALS) (US Patent No. 5,972,924), where CsA treatment more than doubled the remaining life-span. It has also recently been shown that CyP-D inactivation in CyP-D knockout mice protects axons in experimental autoimmune 25 encephalomyelitis, an animal model of multiple sclerosis (Forte M et al. Cyclophilin D inactivation protects axons in experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis. *PNAS* 2007; 104(18): 7558–7563). In an Alzheimer's disease mouse model CyP-D deficiency substantially improves learning, memory and synaptic function (Du H et al. 30 Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease *Nature Medicine* 2008, 14(10): 1097-1105). In addition, CsA has been shown to be effective in a rat model of Huntington's disease (Leventhal L et al. CsA protects striatal neurons in vitro and in vivo from 3-nitropropionic acid toxicity. *Journal of Comparative Neurology* 2000, 425(4): 471-478), and partially effective in a mouse model of 35 Parkinson's disease (Matsuura K et al. CsA attenuates degeneration of dopaminergic neurons induced by 6-hydroxydopamine in the mouse brain. *Brain*

- 13 -

Research 1996, 733(1): 101-104). Thus, mitochondrial-dependent necrosis represents a prominent disease mechanism suggesting that inhibition of CyP-D could provide a new pharmacologic treatment strategy for these diseases (Du, 2008).

5

Cellular, Tissue and Organ Injury due to a Loss of Cellular Calcium Ion (Ca²⁺) Homeostasis

Ca²⁺ is involved in a number of physiological processes at a cellular level, including healthy mitochondrial function. Under certain pathological conditions, 10 such as myocardial infarct, stroke, acute hepatotoxicity, cholestasis, and storage/reperfusion injury of transplant organs, mitochondria lose the ability to regulate calcium levels, and excessive calcium accumulation in the mitochondrial matrix results in the opening of large pores in the inner mitochondrial membrane. (Rasola A. et al. The mitochondrial permeability transition pore and its 15 involvement in cell death and in disease pathogenesis. *Apoptosis* 2007, 12: 815-833.) Nonselective conductance of ions and molecules up to 1.5 kilodaltons through the pore, a process called mitochondrial permeability transition, leads to swelling of mitochondria and other events which culminate in cell death, including the induction of apoptosis. One of the components of the MTP is CyP- 20 D. CyP-D is an immunophilin molecule whose isomerase activity regulates opening of the MPTP, and inhibition of the isomerase activity by CsA or CsA analogs inhibits creation of the MPTP, and thus prevents cell death.

Non-immunosuppressive Cyclosporine Analogue Cyclophilin Inhibitors

25 Despite the advantageous effects of CsA in the above mentioned indications, the concomitant effects of immunosuppression limit the utility of CsA as a CyP inhibitor in clinical practice. At present, there are only a few CsA analogs that have been proven to have little or reduced immunosuppressive activity (i.e., <10% of the immunosuppressive potency of CsA) and still retain their ability to 30 bind CyP (i.e., >10% CyP binding capacity as compared to CsA).

- 14 -

NIM 811 (Melle⁴-cyclosporine)

NIM 811 is a fermentation product of the fungus *Tolypocladium niveum*, which is modified at amino acid 4 and displays no immunosuppressive activity (due to lack of calcineurin binding), yet retains binding affinity for CyP-A (Rosenwirth BA

5 et al. Inhibition of human immunodeficiency virus type 1 replication by SDZ NIM 811, a nonimmunosuppressive Cyclosporine Analogue. *Antimicrob Agents Chemother* 1994, 38: 1763–1772).

DEBIO 025 (MeAla³EtVal⁴-Cyclosporin)

10 DEBIO 025 is a dual chemical modification of CsA at amino acids 3 and 4. DEBIO 025 also displays no immunosuppressive activity yet retains binding affinity for CyP-A PPIase activity (Kristal, 2004).

SCY-635 (DimethylaminoethylthioSar³-hydroxyLeu⁴-Cyclosporin)

15 SCY-635 is a dual chemical modification of CsA at amino acids 3 and 4. SCY-635 also displays no immunosuppressive activity yet retains binding affinity for CyP-A PPIase activity (PCT Publication No. WO2006/039668).

Generally, these compounds have modification on the face of CsA that is 20 responsible for binding calcineurin, and generally require the modification of amino acids 3 and 4. The modification of amino acids 3 and 4 is laborious and complex, as this approach typically involves opening up the cyclosporine ring, replacing and/or modifying those amino acids and then closing up the ring to produce the modified cyclosporine.

25 In contrast, modification of the side chain of amino acid 1 does not require opening of the cyclosporine ring. However, amino acid 1 is associated with CyP binding (as opposed to calcineurin binding) and has been modified to increase the immunosuppressive efficacy of CsA. For example US Patent No. 6,605,593, 30 discloses a single modification of amino acid 1 that results in a CsA analog with increased immunosuppressive potency.

- 15 -

Therefore, it would be desirable to have a cyclosporine analogue molecule (a "CAM") that is readily synthesized and is efficacious in the treatment of CyP mediated diseases. It is also desirable to provide a CsA analogue that provides as least some of the functionality of native CsA, but which possesses improved 5 or additional properties, effects or functions relative to native CsA.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step 10 or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an 15 acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

20 **Summary of the invention**

According to one aspect, compounds of the present invention comprise cyclosporin-A analogs that are non-immunosuppressive according to the definition herein. According to another aspect, the compounds have affinity for 25 cyclophilin, including cyclophilin-A. According to other aspects, compounds of the present invention comprise cyclosporin-A analogs that are at useful with respect to a cyclophilin-mediated disease or condition and developing therapies with respect to such diseases and conditions.


30

According to one aspect, the invention relates to a compound of Formula L:

2011342284

06 Jan 2017

- 15A -



- 16 -


Formula L

wherein

- a. R' is H or acetyl;
- 5 b. R1 is a saturated or unsaturated straight chain or branched aliphatic carbon chain from 2 to 15 carbon atoms in length;
- c. R2 is selected from the group consisting of :
 - i. H;
 - 10 ii. an unsubstituted, N–substituted, or N,N-disubstituted amide;
 - iii. a N-substituted or unsubstituted acyl protected amine;
 - iv. a carboxylic acid;
 - v. a N-substituted or unsubstituted amine;
 - 15 vi. a nitrile;
 - vii. an ester;
 - viii. a ketone;
 - ix. a hydroxy, dihydroxy, trihydroxy, or polyhydroxy alkyl; and
 - 20 x. a substituted or unsubstituted aryl;
 - xi. a saturated or unsaturated. straight or branched aliphatic chain optionally containing a substituent selected from the group consisting of hydrogen, ketones, hydroxyls, nitriles, carboxylic acids, esters, 1,3-dioxolanes, halogens, and oxo;
 - 25 xii. an aromatic group containing a substituent selected from the group consisting of halides, esters and nitro; and
 - xiii. a combination of the saturated or unsaturated, straight or branched aliphatic chain of (xi) and the aromatic group of (xii); and

d. R23 is a saturated or unsaturated straight chain or branched optionally substituted aliphatic carbon chain.

5 In accordance with one aspect, the invention relates to a compound of Formula L:



Formula L

wherein:

- a. R' is H or acetyl;
- 10 b. R1 is a saturated or unsaturated straight or branched aliphatic carbon chain from 2 to 15 carbon atoms in length;
- c. R2 is selected from the group consisting of:
 - i. an unsubstituted, N-substituted, or N,N-disubstituted amide;
 - 15 ii. a N-substituted or unsubstituted acyl protected amine;
 - iii. a carboxylic acid;
 - iv. a N-substituted or unsubstituted amine;
 - v. a nitrile;
 - vi. an ester;
 - vii. a ketone;
 - 20 viii. a hydroxy, dihydroxy, trihydroxy, or polyhydroxy alkyl; and
 - ix. a substituted or unsubstituted aryl;
 - x. a saturated or unsaturated straight or branched aliphatic carbon chain substituted with a substituent selected from the group consisting of ketones, hydroxyls, nitriles, carboxylic acids, esters, 1,3-dioxolanes,
 - 25 halogens, and oxo;
 - xi. an aromatic group substituted with a substituent selected from the group consisting of halides, esters, and nitro; and

xii. a combination of the saturated or unsaturated straight or branched aliphatic carbon chain of (x) and the aromatic group of (xi); and

d. R23 is an unsubstituted saturated or unsaturated straight or branched aliphatic carbon chain.

5


In one aspect, the substituent R1 - R2 is selected from the group consisting of:

- i.
- ii.
- iii.
- iv.
- v.
- vi.
- vii.
- viii.
- ix.
- x.
- xi.
- xii.
- xiii.
- xiv.
- xv.
- xvi.
- xvii.
- xviii.
- xix.
- xx.


2011342284

06 Jan 2017



















- 17B -




- 18 -












- 19 -

xliv. ;
 xlv. ;
 xlvi. ;
 xlvii. ;
 xlviii. ;
 xlxi. ;
 l. ;
 li. ;
 lii. ;
 liii. ;
 liv. ;
 lv. ;
 lvi. ;
 lvii. ;
 lviii. ;
 lix. ;
 lix. ;
 lxi. ;
 lxii. ;
 lxiii. ;
 lxiv. ;
 lxv. ;
 lxvi. 











- 20 -



In another aspect, R2 is selected from the group consisting of



- 21 -



wherein

5 i. R5 is a saturated or unsaturated straight chain or branched aliphatic carbon chain between 1 and 10 carbons in length; and

ii. R6 is a monohydroxylated, dihydroxylated, trihydroxylated or polyhydroxylated saturated or unsaturated straight chain or branched aliphatic carbon chain between 1 and 10 carbons in length.

10

In one aspect, R1-R2 comprises a saturated or unsaturated, straight or branched aliphatic chain of between 2 and 5 carbons optionally substituted with a substituent selected from the group consisting of

- 22 -

hydrogen, ketones, hydroxyls, nitriles, halogens, oxo, carboxylic acids, esters and 1,3-dioxolanes;

In one aspect, R23 is selected from the group consisting of:


- i. $-\text{CH}_3$
- ii. $-\text{CH}_2\text{CH}_3$
- iii. $-\text{CH}_2\text{CHCH}_2$
- iv. $-\text{CH}_2\text{CH}_2\text{CH}_2\text{I}$
- v. $-(\text{CH}_2)_3\text{CH}_2\text{I}$
- vi. $-(\text{CH}_2)_3\text{N}^+(\text{CH}_3)_3$
- vii. $-\text{CH}_2\text{CCH}$
- viii. $-\text{CH}_2\text{CO}_2(\text{t-Bu})$
- ix. $-\text{CH}_2\text{Ph}$
- x. $-\text{CH}_2\text{OH}$
- xi. $-\text{CH}(\text{OH})\text{CH}_3$
- xii. $-\text{CH}(\text{OH})(\text{t-Bu})$
- xiii. $-\text{CH}(\text{OH})\text{Ph}$
- xiv. $-\text{COOH}$
- xv. $-\text{SCH}_3$
- xvi. $-\text{S}(\text{p-Tol})$

5











10 In one aspect, R23 comprises an optionally substituted alkyl, including optionally substituted C1-C3 alkyl. Said alkyl may be substituted with amino and may comprise a C1-C3-Ala wherein said compound comprises the D-epimer. In said embodiment, R23 can be MeAla.

- 23 -


In one aspect, on the above formula L,



is selected from the group consisting of:




and



In one aspect, R23 is a straight or branched aliphatic carbon chain of 1 to 6, 1 to 5, 1 to 4, 1 to 3 or 2 carbons in length.

In another aspect, the invention relates to a compound of formula:



wherein:


R' is H or acetyl;

R1-R2 is ; and

R23 is ethyl or methyl, and

wherein the compound is a D-epimer, wherein the chiral center of the D-epimer is the carbon to which R23 is attached.

In another aspect the invention relates to a compound of formula:



wherein:

R' is H or acetyl;

R1-R2 is ; and

R23 is methyl, and

wherein the compound is a D-epimer, wherein the chiral center of the D-epimer is the carbon to which R23 is attached.

In one aspect, the invention relates to a method of treating or preventing a cyclophilin mediated disease in a mammal comprising administering the compound as described herein to the mammal under conditions to treat the cyclophilin mediated disease or injury, or use of said compound or composition to treat said disease or injury, or use of the compound to prepare a medicament for said use or treatment. Said disease or injury can be mediated by the over expression of cyclophilin or the disease is a congenital over expression of cyclophilin. Said cyclophilin mediated disease or injury can be selected from the group consisting of

- a. a viral infection;
- b. inflammatory disease;
- c. cancer;
- d. muscular disorder;
- e. neurological disorder; and
- f. injury associated with ischemia, reperfusion, loss of cellular calcium homeostasis, loss of ionic homeostasis, increase in free radical production, or toxins that induce mitochondrial dysfunction;

wherein the viral infection is optionally caused by a virus selected from the group consisting of Human Immunodeficiency virus, Hepatitis A, Hepatitis B, Hepatitis C, Hepatitis D, Hepatitis E, SARS-CoV, hCoV-NL63, hCoV-HKU-1, hCoV-OC43, hCoV-229E, coronavirus, feline infectious peritonitis virus, and transmissible gastroenteritis virus;

- 25A -

wherein the inflammatory disease is optionally selected from the group consisting of asthma, autoimmune disease, chronic inflammation, chronic prostatitis, glomerulonephritis, hypersensitivity disease, inflammatory bowel disease, sepsis, vascular smooth muscle cell disease, aneurysms, pelvic inflammatory disease, reperfusion injury, rheumatoid arthritis, transplant rejection, and vasculitis;

wherein the cancer is optionally selected from the group consisting of small and non-small cell lung cancer, bladder cancer, hepatocellular cancer, pancreatic cancer, breast cancer, glioblastoma, colorectal cancer, squamous cell carcinoma, melanoma, and prostate cancer;

wherein the muscular disorder is optionally selected from the group consisting of myocardial reperfusion injury, muscular dystrophy, collagen VI myopathies, Post-cardiac arrest syndrome (PCAS), heart failure, atherosclerosis, and abdominal aortic aneurysm;

wherein the neurological disorder is optionally selected from the group consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple systems atrophy, multiple sclerosis, cerebral palsy, epilepsy, stroke, diabetic neuropathy, amyotrophic lateral sclerosis (Lou Gehrig's Disease), bipolar disorder, excitotoxic injury, hepatic encephalopathy, hypoglycemia, manganese toxicity, neuronal target deprivation, toxic fatty acids such as arachadonic acid, mechanical nerve injury, spinal cord injury, and cerebral injury; and

- 26 -


wherein the injury associated with loss of cellular calcium homeostasis is optionally selected from the group consisting of myocardial infarct, stroke, acute hepatotoxicity, cholestasis, and storage/reperfusion injury of transplant organs.

5

In one aspect, the invention relates to a process for the preparation of a compound of the above Formula L, comprising the steps of:

10


1) reacting cyclosporin-A (CsA) with a basic lithium alkylamide, in the presence of a suitable solvent, followed by reaction with a suitable electrophile to generate a compound of Formula 1:



15


2) reacting the compound of Formula 1 with AC_2O in the presence of a suitable solvent to form a compound of Formula 2A:

- 27 -




Formula 2A

3) Reacting the compound of Formula 2A with an oxidant to form a compound of Formula 3A:



Formula 3A

10 4) Reacting the compound of Formula 3A with an electrophile to form a compound of Formula 4A:



Formula 4A

5) optionally deacylating the compound of Formula 4A.

5 In one aspect, the above preparation of Formula L comprises the addition of an excess of LiCl in said solvent to form predominantly the L-epimer of Formula L, or said preparation of Formula L is carried out in the absence of LiCl to form predominantly the D-epimer of Formula L. Said basic lithium alkylamide can comprise lithium diisopropylamide.

10

In one aspect, said electrophile is selected from the group defined in the following table, to generate the corresponding R23 set out in said table:

Electrophile	R23
methyl iodide	—CH ₃
ethyl iodide	—CH ₂ CH ₃
allyl bromide	—CH ₂ CHCH ₂
1,3-diiodopropane	—CH ₂ CH ₂ CH ₂ I
1,4-diiodobutane	—(CH ₂) ₃ CH ₂ I
trimethylammonium-3-iodopropane	—(CH ₂) ₃ N ⁺ (CH ₃) ₃
hexafluorophosphate	
propargyl bromide	—CH ₂ CCH
tert-butyl bromoacetate	—CH ₂ CO ₂ (t-Bu)
benzyl bromide	—CH ₂ Ph ²
formaldehyde	—CH ₂ OH
acetaldehyde	—CH(OH)CH ₃
pivalaldehyde	—CH(OH)(t-Bu) ³
benzaldehyde	—CH(OH)Ph
carbon dioxide	—COOH

- 29 -


dimethyl disulfide	$-\text{SCH}_3$
p-tolyl disulfide	$-\text{S}(\text{p-Tol})^4$

In one aspect, the invention relates to a method for the preparation of Formula L as defined herein comprising the steps of:


5 in the presence of a suitable solvent, to form the compound of
Formula 2:

in the presence of a suitable solvent, to form the compound of Formula I.

Formula 2.



optionally followed by formation of the aldehyde of Formula 3:




optionally followed by a Wittig reaction to generate compounds of Formula L, wherein R23 is selected from the group consisting of


- i. $-\text{CH}_3;$
- ii. $-\text{CH}_2\text{CH}_3;$
- iii. $-\text{CH}_2\text{CHCH}_2;$
- iv. $-\text{CH}_2\text{CH}_2\text{CH}_2\text{I};$
- v. $-(\text{CH}_2)_3\text{CH}_2\text{I};$

- vi. $-(CH_2)_3N^+(CH_3)_3$
- vii. $-CH_2CCH_3$
- viii. $-CH_2CO_2(t\text{-}Bu)_2$
- ix. $-CH_2Ph^2$
- x. $-CH_2OH$
- xi. $-CH(OH)CH_3$
- xii. $-CH(OH)(t\text{-}Bu)_3$
- xiii. $-CH(OH)Ph$
- xiv. $-COOH$
- xv. $-SCH_3$; and
- xvi. $-S(p\text{-}Tol)_2$

In another aspect, the invention relates to a process for the preparation of the compound of Formula L, comprising the steps of:
reacting a compound of Formula 1A with acetic anhydride in the presence of a suitable solvent, to form a compound of Formula 2:



optionally followed by formation of an aldehyde of Formula 3:



optionally followed by a Wittig reaction to generate a compound of Formula L, wherein R23 is selected from the group consisting of


i. $-\text{CH}_3$

- 30A -

- ii. $-\text{CH}_2\text{CH}_3$;
- iii. $-\text{CH}_2\text{CHCH}_2$; and
- iv. $-\text{CH}_2\text{CCH}$.

5

In one aspect, the invention relates to the preparation of Formula L as defined herein comprising the steps of: reacting a compound of Formula 5, with a basic lithium alkylamide, optionally comprising lithium diisopropylamide, in presence of a suitable electrophile in an appropriate solvent to form the compound of Formula L, wherein R23 comprises optionally substituted C1-C3 alkyl.



Formula 5

In general, for all chemical formulae disclosed in this document:

5 "Carboxylic acid" includes a group in which the carboxylic acid moiety is connected to one of the following substituents:

1. alkyl which may be substituted (for example, alkyl of 2 to 15 carbons);
2. alkenyl which may be substituted (for example, alkenyl of 2 to 15 carbons); and
- 10 3. alkynyl which may be substituted (for example, alkynyl of 2 to 15 carbons);

The substituents of the above-described may include halogen (for example, fluorine, chlorine, bromine, iodine, etc.), nitro, cyano, hydroxy, thiol which may be substituted (for example, thiol, C1-4 alkylthio, etc.), amino which may be substituted (for example, amino, mono-C1-4 alkylamino, di-C1-4 alkylamino, 5- to 6-membered cyclic amino such as tetrahydropyrrole, piperazine, piperidine, morpholine, thiomorpholine, pyrrole, imidazole, etc.), C1-4 alkoxy which may be halogenated (for example, methoxy, ethoxy, propoxy, butoxy, trifluoromethoxy, trifluoroethoxy, etc.), C1-4 alkoxy-C1-4 alkoxy which may be halogenated (for example, methoxymethoxy, methoxyethoxy, ethoxyethoxy, trifluoromethoxyethoxy, trifluoroethoxyethoxy, etc.), formyl, C2-4 alkanoyl (for example, acetyl, propionyl, etc.), C1-4 alkylsulfonyl (for example, methanesulfonyl, ethanesulfonyl, etc.), and the like, and the number of the substituents is preferably 1 to 3.

Further, the substituents of the above "amino which may be substituted" may bind each other to form a cyclic amino group (for example, a group which is formed by subtracting a hydrogen atom from the ring constituting nitrogen atom of a 5- to 6-membered ring such as tetrahydropyrrole, piperazine, piperidine, morpholine, thiomorpholine, pyrrole, imidazole, etc. so that a substituent can be attached to the nitrogen atom, or the like). The cyclic amino group may be

- 32 -

substituted and examples of the substituent include halogen (for example, fluorine, chlorine, bromine, iodine, etc.), nitro, cyano, hydroxy, thiol which may be substituted (for example, thiol, C1-4 alkylthio, etc.), amino which may be substituted (for example, amino, mono-C₁-C₄ alkylamino, di-C₁-C₄ alkylamino,

5 5- to 6-membered cyclic amino such as tetrahydropyrrole, piperazine, piperidine, morpholine, thiomorpholine, pyrrole, imidazole, etc.), carboxyl which may be esterified or amidated (for example, carboxyl, C₁-C₄ alkoxy-carbonyl, carbamoyl, mono-C₁-C₄ alkyl-carbamoyl, di-C₁-C₄ alkyl-carbamoyl, etc.), C₁-C₄ alkoxy which may be halogenated (for example, methoxy, ethoxy, propoxy, butoxy,

10 trifluoromethoxy, trifluoroethoxy, etc.), C₁-C₄ alkoxy-C₁-C₄ alkoxy which may be halogenated (for example, methoxymethoxy, methoxyethoxy, ethoxyethoxy, trifluoromethoxyethoxy, trifluoroethoxyethoxy, etc.), formyl, C₂-C₄ alkanoyl (for example, acetyl, propionyl, etc.), C₁-C₄ alkylsulfonyl (for example, methanesulfonyl, ethanesulfonyl), and the like, and the number of the

15 substituents is preferably 1 to 3.

“Amine” includes a group which may be unsubstituted or in which the amine moiety is N-substituted or N,N disubstituted having one or two substituents which may be independently selected from:

20 1. alkyl which may be substituted (for example, alkyl of 2 to 15 carbons);
2. alkenyl which may be substituted (for example, alkenyl of 2 to 15 carbons);
3. alkynyl which may be substituted (for example, alkynyl of 2 to 15 carbons);
25 4. formyl or acyl which may be substituted (for example, alkanoyl of 2 to 4 carbons (for example, acetyl, propionyl, butyryl, isobutyryl, etc.), alkylsulfonyl of 1 to 4 carbons (for example, methanesulfonyl, ethanesulfonyl, etc.) and the like);
5. aryl which may be substituted (for example, phenyl, naphthyl, etc.); and

30 the like;

and connected to a substituent independently selected from the substituents as defined for “carboxylic acid” above.

"Amide" includes a compound in which the carboxylic group of the amide moiety is

connected to a substituent independently selected from the substituents as defined for "carboxylic acid" above, connected to the amino group of the amide

5 moiety is an N-substituted or N,N disubstituted having one or two substituents, respectively, which may be independently selected from:

1. alkyl which may be substituted (for example, alkyl of 2 to 15 carbons);
2. alkenyl which may be substituted (for example, alkenyl of 2 to 15 carbons);
- 10 3. alkynyl which may be substituted (for example, alkynyl of 2 to 15 carbons);
4. formyl or acyl which may be substituted (for example, alkanoyl of 2 to 4 carbons (for example, acetyl, propionyl, butyryl, isobutyryl, etc.), alkylsulfonyl of 1 to 4 carbons (for example, methanesulfonyl, ethanesulfonyl, etc.) and the like);
- 15 5. aryl which may be substituted (for example, phenyl, naphthyl, etc.); and the like

"Aryl" may be exemplified by a monocyclic or fused polycyclic aromatic hydrocarbon group, and for example, a C6-14 aryl group such as phenyl, naphthyl, anthryl, phenanthryl or acenaphthylene, and the like are preferred, with phenyl being preferred. Said aryl may be substituted with one or more substituents, such as lower alkoxy (e.g., C1-6 alkoxy such as methoxy, ethoxy or propoxy, etc.), a halogen atom (e.g., fluorine, chlorine, bromine, iodine, etc.), lower alkyl (e.g., C1-6 alkyl such as methyl, ethyl or propyl, etc.), lower alkenyl (e.g., C2-6 alkenyl such as vinyl or allyl, etc.), lower alkynyl (e.g., C2-6 alkynyl such as ethynyl or propargyl, etc.), amino which may be substituted, hydroxyl which may be substituted, cyano, amidino which may be substituted, carboxyl, lower alkoxycarbonyl (e.g., C1-6 alkoxycarbonyl such as methoxycarbonyl or ethoxycarbonyl, etc.), carbamoyl which may be substituted (e.g., carbamoyl which may be substituted with C1-6 alkyl or acyl (e.g., formyl, C2-6 alkanoyl, benzoyl, C1-6 alkoxycarbonyl which may be halogenated, C1-6 alkylsulfonyl

- 34 -

which may be halogenated, benzenesulfonyl, etc.) which may be substituted with a 5- to 6-membered aromatic monocyclic heterocyclic group (e.g., pyridinyl, etc.), 1-azetidinylcarbonyl, 1-pyrrolidinylcarbonyl, piperidinocarbonyl, morpholinocarbonyl, thiomorpholinocarbonyl (the sulfur atom may be oxidized),

5 1-piperazinylcarbonyl, etc.), or the like. Any of these substituents may be independently substituted at 1 to 3 substitutable positions.

“Ketone” includes a compound in which the carbonyl group of the ketone moiety is connected to one or two substituents independently selected from the

10 substituents as defined above for said “carboxylic acid”.

“Ester” includes either a carboxylic or an alcohol ester wherein of the ester group is composed of one or two substituents independently selected from the substituents as defined for “carboxylic acid” or “aryl”.

15 “Alkyl” unless otherwise defined is preferably an alkyl of 1 to 15 carbon units in length.

“Aromatic group” may be exemplified by aryl as defined above, or a 5- to 6-membered aromatic monocyclic heterocyclic group such as furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, furazanyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl or the like; and a 8- to 16-membered (preferably, 10- to 12-membered) aromatic fused heterocyclic group

“Non-immunosuppressive” refers to the ability of a compound to exhibit a substantially reduced level of suppression of the immune system as compared with CsA, as measured by the compounds ability to inhibit the proliferation of

30 human lymphocytes in cell culture and preferably as measured by the method set out in Example 19 below.

- 35 -

“Analogue” means a structural analogue of CsA that differs from CsA in one or more functional groups. Preferably, such analogues preserve at least a substantial portion of the ability of CsA to bind CyP.

5 Preferred species of Formula I are those in which R' is H, R1 is a saturated or unsaturated alkyl between 2 and 15 carbons in length and R2 is selected from:

1. carboxylic acid comprising a carboxyl group;
2. N-substituted or N,N-disubstituted amide wherein the substituents are independently selected from an H, an alkyl of between 1 and 7 carbons in length, or said substituents form a heterocyclic ring of which the heterocycle is selected from O, N or S;
3. an ester of between 1 and 7 carbons in length;
4. an monohydroxylated, or dihydroxylated alkyl of between 1 and 7 carbons in length;
- 10 5. a N-substituted or unsubstituted acyl protected amine of between 1 and 7 carbons in length;
6. a nitrile;
7. a ketone wherein the carboxylic group of the ketone is connected to R1 and saturated or unsaturated alkyl chain of between 1 and 7 carbons in length;
- 15 8. phenyl, optionally substituted with one or more substituents independently selected from nitrogen dioxide, a fluorine, an amine, an ester or a carboxyl group.

20 25 The compounds of the present invention may exist in the form of optically active compounds. The present invention contemplates all enantiomers of optically active compounds within the scope of the above formulae, both individually and in mixtures of racemates. As well, the present invention includes prodrugs of the compounds defined herein.

30 According to another aspect, compounds of the present invention may be useful for treating or preventing or studying a CyP mediated disease in a mammal,

- 36 -

preferably a human. Such disease is usually mediated by the over expression of CyP, such as a congenital over expression of CyP.

CyP mediated diseases which may be treated by compounds of the present

5 invention include:

- a. a viral infection;
- b. inflammatory disease;
- c. cancer;
- 10 d. muscular degenerative disorder;
- e. neurodegenerative disorder; and
- f. injury associated with loss of cellular calcium homeostasis.

Said viral infection may be caused by a virus selected from the group consisting
15 of Human Immunodeficiency Virus, Hepatitis A, Hepatitis B, Hepatitis C,
Hepatitis D, and Hepatitis E. Said inflammatory disease is selected from the
group consisting of asthma, autoimmune disease, chronic inflammation, chronic
prostatitis, glomerulonephritis, hypersensitivity disease, inflammatory bowel
disease, sepsis, vascular smooth muscle cell disease, aneurysms, pelvic
20 inflammatory disease, reperfusion injury, rheumatoid arthritis, transplant
rejection, and vasculitis. Said cancer may be selected from the group consisting
of small and non-small cell lung, bladder, hepatocellular, pancreatic and breast
cancer. Said muscular degenerative disorder may be selected from the group
consisting of myocardial reperfusion injury, muscular dystrophy and collagen VI
25 myopathies. Said neurodegenerative disorder may be selected from the group
consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease,
Multiple Systems Atrophy, Multiple Sclerosis, cerebral palsy, stroke, diabetic
neuropathy, amyotrophic lateral sclerosis (Lou Gehrig's Disease), spinal cord
injury, and cerebral injury. Said injury associated with loss of cellular calcium
30 homeostasis may be selected from the group consisting of myocardial infarct,
stroke, acute hepatotoxicity, cholestasis, and storage/reperfusion injury of
transplant organs.

DETAILED DESCRIPTION

According to one aspect, a compound of this invention may be administered neat or with a pharmaceutical carrier to a warm-blooded animal in need thereof. The 5 pharmaceutical carrier may be solid or liquid. The compound may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral, as used herein, includes subcutaneous injections, intravenous, intramuscular, intrasternal 10 injection or infusion techniques.

The pharmaceutical compositions containing the inventive mixture may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft 15 capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to methods known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide 20 pharmaceutically elegant and palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients may also be manufactured by known methods. The excipients used may be for example, (1) inert diluents such as calcium carbonate, lactose, calcium phosphate or sodium phosphate; (2) granulating and disintegrating 25 agents such as corn starch, or alginic acid; (3) binding agents such as starch, gelatin or acacia, and (4) lubricating agents such as magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time 30 delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in the U.S.

Patent Number 4,256,108; 4,160,452; and 4,265,874 to form osmotic therapeutic tablets for controlled release.

In some cases, formulations for oral use may be in the form of hard gelatin

5 capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They may also be in the form of soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

10 Aqueous suspensions normally contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients may include: (1) suspending agents such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; or (2) dispersing or
15 wetting agents which may be a naturally-occurring phosphatide such as lecithin, a condensation product of an alkylene oxide with a fatty acid, for example, polyoxyethylene stearate, a condensation product of ethylene oxide with a long chain aliphatic alcohol, for example, heptadecaethyleneoxycetanol, a condensation product of ethylene oxide with a partial ester derived from a fatty
20 acid and a hexitol such as polyoxyethylene sorbitol monooleate, or a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride, for example polyoxyethylene sorbitan monooleate.

25 The aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate; one or more coloring agents; one or more flavoring agents; and one or more sweetening agents such as sucrose, aspartame or saccharin.

30 Oily suspension may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, a fish oil which contains omega 3 fatty acid, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax,

hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

- 5 Dispersible powders and granules are suitable for the preparation of an aqueous suspension. They provide the active ingredient in a mixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, those sweetening, 10 flavoring and coloring agents described above may also be present.

The pharmaceutical compositions containing the inventive mixture may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil such as olive oil or arachis oils, or a mineral oil such as liquid paraffin or a mixture 15 thereof. Suitable emulsifying agents may be (1) naturally-occurring gums such as gum acacia and gum tragacanth, (2) naturally-occurring phosphatides such as soy bean and lecithin, (3) esters or partial ester 30 derived from fatty acids and hexitol anhydrides, for example, sorbitan monooleate, (4) condensation products of said partial esters with ethylene oxide, for example, polyoxyethylene 20 sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example, glycerol, propylene glycol, sorbitol, aspartame or sucrose. Such formulations 25 may also contain a demulcent, a preservative, and flavoring and coloring agents.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to known methods using those suitable dispersing or wetting agents 30 and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-

- 40 -

butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including

5 synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The inventive compound may also be administered in the form of suppositories for rectal administration of the drug. Suitable compositions can be prepared by

10 mixing the compound with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

15 For topical use, suitable creams, ointments, jellies, solutions or suspensions, etc., which normally are used with cyclosporine may be employed.

In a particularly preferred embodiment, a liquid solution containing a surfactant, ethanol, a lipophilic and/or an amphiphilic solvent as non-active ingredients is

20 used. Specifically, an oral multiple emulsion formula containing the isomeric analogue mixture and the following non-medicinal ingredients: d-alpha Tocopheryl polyethylene glycol 1000 succinate (vitamin E TPGS), medium chain triglyceride (MCT) oil, Tween 40, and ethanol is used. A soft gelatin capsule (comprising gelatin, glycerin, water, and sorbitol) containing the compound and 25 the same non-medicinal ingredients as the oral solution may also preferably be used.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific


30 compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the nature and severity of the particular disease or condition undergoing therapy.

- 41 -


Methodology

The reactions set out below, are general examples of the chemical reactions able to synthesize the desired compounds modified at amino acid 1 residue (AA1) and amino acid 3 residue (AA3) of CsA. Modifications to AA1 are

5 depicted as:



and modifications to AA3 are depicted as



10

both modifications to AA1 and AA3 use reagents that have the requisite chemical properties, and it would be understood by a person skilled in the art 15 that substitutions of certain reactants may be made.

The identity and purity of the prepared compounds were generally established by methodologies including mass spectrometry, HPLC and NMR spectroscopy. Mass spectra (ESI-MS) were measured on a Hewlett Packard 1100 MSD 20 system. NMR spectra were measured on a Varian MercuryPlus 400 MHz spectrometer in deuterated solvents (DMSO for phosphonium salts, benzene for all other compounds). Analytical and preparative reversed phase HPLC was carried out on an Agilent 1100 Series system.

25 *Synthesis of Phosphonium Salt Compounds*

Phosphonium salts are prepared through reaction of triphenylphosphine or any other suitable phosphines with alkyl halides (R-X; X = Cl, Br, or I). Suitable alkyl halides are any primary or any secondary aliphatic halide of any chain length or


- 42 -

molecular weight. These alkyl halides may be branched or unbranched, saturated or unsaturated.

If the reaction is carried out in toluene (Reaction 1), the product precipitates

5 directly from the reaction solution. Unreactive substrates, however, require a more polar solvent such as dimethylformamide (DMF) (Reaction 2) to shorten reaction times and to achieve satisfactory yields.

Reaction 1:



10 Where X is a halide (including but not limited to Cl, Br, and I), and R10 is a saturated or unsaturated, straight or branched aliphatic chain, optionally containing a substituent selected from the group of ketones, hydroxyls, nitriles, carboxylic acids, esters and 1,3-dioxolanes; an aromatic group, optionally containing a substituent selected from the group of halides, esters and nitro; or a

15 combination of the aforementioned saturated or unsaturated, straight or branched aliphatic chain and the aforementioned aromatic groups.

Example 1. Synthesis of 404-15



As an illustrative example, triphenylphosphine (13 mmol) is dissolved in 50 mL





20 toluene and chloroacetone (10 mmol) is added to give a clear solution. The reaction is stirred under reflux over night. A colorless solid is filtered off, washed with toluene and hexane and dried in vacuum.

- 43 -

Using Reaction 1, the following compounds are further examples of the compounds that may be synthesized.


Compound	Reactant (R10-X)	Conditions
404-08	benzyl bromide	4 hours at reflux
404-09	methyl iodide	RT over night
404-12	4-nitrobenzyl bromide	6 hours at reflux
404-15	chloroacetone	reflux over night
404-64	4-fluorobenzyl bromide	reflux over night
404-77	methyl 3-	6 hours at reflux

- 44 -

		bromomethylbenzoate
	404-87	3-nitrobenzyl bromide
	404-161	1-bromo-2-butanone
	404-170	4-bromobutyronitrile
		reflux over night

Alternatively, suitable phosphonium salts may be synthesized through Reaction 2 as illustrated below:

5 Reaction 2:




Where X is a halide (including but not limited to Cl, Br, and I), and R10 is a saturated or unsaturated, straight or branched aliphatic chain, optionally containing a substituent selected from the group of ketones, hydroxyls, nitriles,

- 45 -

carboxylic acids, esters and 1,3-dioxolanes; an aromatic group, optionally containing a substituent selected from the group of halides, esters and nitro; or a combination of the aforementioned saturated or unsaturated, straight or branched aliphatic chain and the aforementioned aromatic groups.

5

Example 2: Synthesis of 404-51



As an illustrative example, triphenylphosphine (11 mmol) is dissolved in 10 mL DMF and 4-bromobutyric acid (10 mmol) is added. The reaction is stirred for 7 hours at 110 °C and is then allowed to cool over night. Fifty mL toluene is added 10 and a crystalline, colorless solid is collected by filtration. The product is washed with toluene and hexane and dried in vacuum over night.

If crystallization does not set in after treatment with toluene, the product is extracted with 20 mL MeOH/H₂O (1:1 mixture). The aqueous phase is washed 15 with toluene and hexane and brought to dryness. The residue is stirred with 50 mL ethyl acetate (EtOAc) at reflux temperature for 20-30 min. If a crystalline solid is obtained, the product is collected by filtration, washed with EtOAc and hexane and dried. In case the product is obtained as an oil or gum, the EtOAc is decanted and the remaining product is dried in vacuum.


20

Using Reaction 2, the following compounds are further examples of the compounds that may be synthesized.


Compound	Reactant (R11-X)	Conditions
404-14	1-bromobutane	6.5 hours at 120 °C

- 46 -


404-29 2-bromomethyl-1,3-dioxolane 120 °C over night




404-34 1-bromooctane 110 °C over night




404-51 5-bromovaleric acid 8 hours at 120 °C



404-78 6-bromohexanol 110 °C over night




404-116 4-bromobutyric acid 7 hours at 110 °C



416-01 1-bromohexane 110 °C over night


- 47 -



416-02

6-bromohexanoic acid


110 °C over night



419-132

7-bromoheptanenitrile


110 °C over night



419-134

6-chloro-2-hexanone


110 °C over night



419-136

9-bromo-1-nonanol

110 °C over night









420-32

methyl 7-bromohexanoate

110 °C over night

- 48 -

	420-78	11-bromoundecanoic acid	110 °C over night
	420-80	3-bromopropionitrile	110 °C over night
	420-82	8-bromooctanoic acid	110 °C over night
	420-90	6-bromohexanenitrile	110 °C over night
	420-94	5-chloro-2-pentanone	110 °C over night
			

- 49 -


Wittig Reaction

The Wittig reaction is broadly applicable to a wide range of substrates and reactants. The side chain, which is introduced to the substrate in the reaction, can represent any number of branched and unbranched, saturated and

5 unsaturated aliphatic compounds of variable length (R') and may contain a broad range of functional groups.

In the Wittig reaction, a base, such as potassium tert-butoxide (KOtBu) is used to generate an ylide from a phosphonium salt. The ylide reacts with the carbonyl 10 group of the substrate, CsA-aldehyde, to form an alkene. Phosphonium salts containing a carboxylic acid side chain require at least two equivalents of base to generate the ylide.

Reaction 3: Synthesis of an Acetylated Cyclosporine analogue Intermediate 15 using a Phosphonium Salt compound through a Wittig Reaction




- 50 -

Where X is a halide (including but not limited to Cl, Br, and I), and R12 is a saturated or unsaturated, straight or branched aliphatic chain, optionally containing a substituent selected from the group of ketones, hydroxyls, nitriles, carboxylic acids, esters and 1,3-dioxolanes; an aromatic group, optionally

5 containing a substituent selected from the group of halides, esters and nitro; or a combination of the aforementioned saturated or unsaturated, straight or branched aliphatic chain and the aforementioned aromatic groups.

Example 3: Synthesis of Compound 404-20 using a Phosphonium Salt Compound through a Wittig Reaction:

10



As an illustrative example, an oven dried 250 mL flask is charged under argon atmosphere with triphenylbutylphosphonium bromide (6.0 mmol) and 40 mL anhydrous tetrahydrofuran (THF). The suspension is cooled to 0 °C and potassium tert-butoxide (6.0 mmol) is added to obtain an orange color. The reaction is stirred at ambient temperature for 1-2 hours, followed by addition of CsA-aldehyde (2.0 mmol, dissolved in 20 mL anhydrous THF). Stirring is

15

- 51 -

continued for 3 hours at room temperature. The reaction is quenched with 10 mL sat. NH_4Cl and 20 mL ice-water. The layers are separated and the aqueous phase is extracted with EtOAc . The organic layers are combined, washed with brine and dried over Na_2SO_4 . The solvent is removed and the crude product is 5 purified over silica gel (hexane/acetone 3:1).

Using Reaction 3, the following compounds are further examples of the compounds that may be synthesized.

Compound	Starting Material	MS (Na^+)	Remarks
404-16	404-09	1252.9	
404-19	404-08	1328.9	
404-20	404-14	1294.9	
404-30	404-12	1373.9	stirred at 60 °C over night











- 52 -

404-31		1324.9	stirred at 60 °C for 2 days
404-33		1325.0	
404-40		1351.2	
404-43		1295.1	stirred at reflux for 10 days
404-59		1338.9	2 eq of KOtBu
404-29			
404-34			
404-15			
404-51			

- 53 -

404-65	404-64	1347.1
404-79	404-77	1386.9 stirred at RT over night
404-89	404-87	1374.1 stirred at RT for 2 days
404-134	404-116	1325.0 2 eq of KOtBu; stirred at RT for 2 days
404-163	404-161	1308.8 stirred at reflux for 15 days

- 54 -


404-187	404-170	1305.9	stirred at RT
			over night
416-04	416-02	1353.0	2 eq of KOtBu
			
416-09	416-01	1323.1	
			
420-40	420-32	1381.0	stirred at RT over night
			
420-85	420-78	1423.1	2 eq of KOtBu
			

- 55 -

420-89		420-80		1291.9
420-92		420-82		1381.1 2 eq of KOtBu
420-96		404-78		1338.9
420-101		419-132		1347.9

Deacetylation


Reaction 4: Deacetylation of Acetylated Cyclosporine Analogue



- 56 -

Where R12 is a saturated or unsaturated, straight or branched aliphatic chain, optionally containing a substituent selected from the group of ketones, hydroxyls, nitriles, carboxylic acids, esters, amides, acyl-protected amines and 1,3-dioxolanes; an aromatic group, optionally containing a substituent selected from 5 the group of halides, esters, amines and nitro; or a combination of the aforementioned saturated or unsaturated, straight or branched aliphatic chain and the aforementioned aromatic groups.

Example 4: Synthesis of Compound 404-90 though Deacetylation



10

As an illustrative example, a solution of 404-20 (0.16 mmol) in 10 mL MeOH is combined with a solution of tetramethylammoniumhydroxide pentahydrate (0.47 mmol) in 2 mL H₂O. The mixture is stirred at room temperature for 2 days. The reaction is concentrated in vacuum and 5 mL H₂O are added. The reaction is 15 extracted with EtOAc, the extract is washed with brine, dried over Na₂SO₄ and concentrated to dryness. The crude product is purified by reversed phase preparative HPLC.

Purification of deacetylated compounds is generally carried out over silica gel 20 (hexane/acetone 2:1) or by preparative HPLC. In the case of compounds 404-60, 404-137, 416-08, 420-98 and 420-100 (carboxylic acids), the reaction is acidified to pH 2-3 with 1 M HCl prior to extraction.


- 57 -

Using Reaction 4, the following compounds are further examples of the compounds that may be synthesized.


Compound	Starting Material	MS (Na ⁺)
404-22	404-16	1210.9
404-25	404-19	1287.0
404-36	404-33	1283.0
404-44	404-40	1309.1
404-58	404-57	1257.1
404-60	404-59	1297.1
404-61	404-56	1255.1

- 58 -

404-66




404-65




1305.1

404-81-1




404-79-1




1331.1

404-81-2




404-79-1




1345.1

404-85




404-83




1326.2

404-90




404-20




1253.0

404-96-1




404-94




1333.0

404-96-2




404-94




1347.0

404-97



404-89

















1331.9

404-125

404-120

1304.0


















- 59 -

		
404-130	404-128	1270.1
		
404-132	404-129	1298.0
		
404-137	404-134	1283.0
		
404-154	404-150	1338.1
		
404-157	404-155	1310.0
		
404-173	404-172	1268.9
		
404-194	404-187	1263.9
		

- 60 -

416-08	416-04	1311.0
416-13	416-09	1281.1
420-17	420-08-1	1368.0
420-30-1	420-27	1312.0
420-43	420-40	1324.9
420-47	420-46	1327.0
420-98	420-85	1381.1
420-100	420-92	1339.1
420-102	420-96	1297.0
420-108	420-101	1305.9

- 61 -

	420-117	
	420-109-1	1352.1
	420-120	
	420-110-1	1410.0
	420-122	
	420-107-2	1340.0
	420-124	
	420-109-2	1354.0
	420-125	
	420-110-2	1412.0
	420-126	
	420-107-1	1337.9
	420-131	
	420-130	1297.9
	420-132	
	420-128-1	1380.0
	420-133	
	420-134	


Hydrogenation of the Double Bond

The double bond can be hydrogenated under atmospheric pressure to obtain the saturated side chain. Functional groups such as hydroxyl, carbonyl and carboxyl

- 62 -


are stable under these conditions and do not require protection. R' represents either an acetyl group or hydrogen. In the case of α,β -unsaturated carbonyl compounds the double bond has to be reduced prior to deacetylation to avoid cyclization through a nucleophilic addition of the free hydroxy group on the 5 activated double bond.

Reaction 5:



Where R12 is a saturated or unsaturated, straight or branched aliphatic chain, 10 optionally containing a substituent selected from the group of ketones, hydroxyls, nitriles, carboxylic acids, esters, amides, acyl-protected amines and 1,3-dioxolanes; an aromatic group, optionally containing a substituent selected from the group of halides, esters, amines and nitro; or a combination of the aforementioned saturated or unsaturated, straight or branched aliphatic chain 15 and the aforementioned aromatic groups, and R' is either a H or an acetyl group.

Example 5: Synthesis of 404-56:



As an illustrative example, 404-43 (0.34 mmol) is dissolved in 40 mL anhydrous EtOH and 43 mg Pd/C (10 %) and 0.2 mL acetic acid are added. The mixture is 20 stirred under hydrogen at atmospheric pressure for 2 days. The reaction is filtered through Celite and is concentrated in vacuum. The crude product is purified by preparative HPLC.

- 63 -


Using Reaction 5, the following compounds are further examples of the compounds that may be synthesized.

Compound	Starting Material	MS (Na ⁺)
404-50	404-25	1289.1
404-56	404-43	1297.0
404-57	404-31	1327.1
404-63	404-60	1299.1
404-74	404-66	1307.1
404-92	404-90	1255.1
404-94	404-79	1388.9

- 64 -

404-168	404-134	1326.8
404-172	404-163	1310.9
420-19	416-08	1313.0
420-46	420-40	1383.1
420-68	420-134	1326.9
420-106	420-98	1383.1
420-111	420-100	1341.0
420-112	420-102	1298.9
420-130	420-123	1340.0


- 65 -



Reduction of the Nitrile Group

Reduction of the nitrile group to the corresponding primary amine can be achieved with nickel boride generated *in situ* from sodium borohydride (NaBH_4) and nickel(II)chloride (NiCl_2). Addition of a suitable trapping reagent leads to acyl-protected primary amines (carbamates or amides, respectively) and prevents the formation of secondary amines as an undesired side reaction. The double bond is partially reduced under the given conditions and a product mixture is obtained. Both, saturated and unsaturated protected amine compounds were isolated and purified. For reaction 420-123 the mixture was not separated. Instead, the mixture underwent catalytic hydrogenation to produce the fully saturated compound.


Reaction 6:



Where Acyl is any one of BOC, acetyl, or butyryl, acylating agent is any one of di-tert-butyldicarbonate, acetic anhydride, and butyric anhydride and R1 is a saturated or unsaturated straight chain or branched aliphatic group. It would be understood by one skilled in the art that the acylating agents described above may be replaced with a broad range of acylating agents to produce a similarly broad range of acyl-protected amines.

- 66 -

Example 6: Synthesis of 420-08



As an illustrative example, 404-187 (0.257 mmol) is dissolved in 15 mL methanol and cooled to 0 °C. Di-tert-butyldicarbonate (0.514 mmol) and nickel(II)chloride (0.025 mmol) are added to give a clear solution. Sodium borohydride (3.85


5 mmol) is added in portions over 1 hour. The resulting mixture is stirred at ambient temperature over night. Additional sodium borohydride (1.95 mmol) is added at 0 °C and stirring is continued for 3 hours at room temperature. HPLC shows a mixture of 420-08-1 (carbamate compound) and 420-08-2 (carbamate compound with double bond reduced). The reaction is stirred for 30 minutes
10 with diethylenetriamine (0.257 mmol) and is then concentrated in vacuum. The residue is taken up in 75 mL EtOAc, washed with 20 mL sat. NaHCO₃ solution and dried over Na₂SO₄. The solvent is removed in vacuum. The crude product is purified by preparative HPLC.

15 Using Reaction 6, the following compounds are further examples of the compounds that may be synthesized.


Compound	Starting Material	Protecting Reagent	MS (Na ⁺)
420-08-1	404-197	di-tert-butyldicarbo ne	1410.0

- 67 -

420-08-2




404-197




di-tert-butyldi-carbonate 1412.1

420-107-1




404-197




butyric anhydride 1379.9

420-107-2




404-197




butyric anhydride 1382.1

420-109-1




420-101




acetic anhydride 1394.1

420-109-2




420-101




acetic anhydride 1396.1

420-110-1




420-101




di-tert-butyldi-carbonate 1452.1

420-110-2



420-101



di-tert-butyldicarbon- te 1454.1

- 68 -

420-123 ¹	420-89	acetic anhydride	1337.9/1339.9
420-128-1	420-101	butyric anhydride	1422.1
420-128-2	420-101	butyric anhydride	1424.1

¹ mixture not separated

Amine Deprotection

The BOC protected amine (carbamate) can be converted into the free amine by

5 acidic hydrolysis using trifluoroacetic acid (TFA).


Reaction 7:



10 Where R₁ is a saturated or unsaturated, straight or branched aliphatic chain, and R' is either a H or an acetyl group.

- 69 -

Example 7: Synthesis of 420-23



As an illustrative example, 420-17 (0.026 mmol) is dissolved in 4 mL anhydrous DCM and 2 mL trifluoroacetic acid is added at 0 °C. The reaction is stirred at room temperature for 3 hours. Twenty 20 mL dichloromethane is added. The

5 reaction mixture is washed with H₂O and sat. NaHCO₃ solution and is dried over Na₂SO₄. The solvent is removed and the crude product is purified by preparative HPLC.

10 Using Reaction 7, the following compounds are further examples of the compounds that may be synthesized.

Compound	Starting Material	MS (M+1)
420-23	420-17	1246.0
420-25	420-13	1290.0
420-129	420-120	1288.0

Protection of the Amino Group


The free amino function can be protected using a wide range of protecting groups using established methods. A broader range of protecting agents is available compared to the reductive introduction starting from the nitrile.

- 70 -

Together, reactions 7 and 8 offer an alternate route to reaction 6 for the preparation of acyl-protected amino compounds.


Reaction 8:

5



Where Acyl is any one of BOC, acetyl or butyryl, acylating agent is any one of di-tert-butyldicarbonate, acetic anhydride, butyric anhydride, it would be understood by one skilled in the art that a broad range of acylating agents including, dicarbonates, anhydrides and acyl halides can be employed to produce a broad range of acyl-protected amines, and R1 is a saturated or unsaturated straight chain or branched aliphatic group.

15 Example 8: Synthesis of 420-27




As an illustrative example, 420-25 (0.039 mmol) is dissolved in 3 mL anhydrous pyridine under nitrogen. The reaction is cooled to 0 °C and acetic anhydride (0.59 mmol) is added. The mixture is stirred at ambient temperature overnight. The solvent is removed in vacuum and the residue is taken up in 25 mL EtOAc.

20 The reaction is washed with 2 x 10 mL 1 M HCl, 2 x 10 mL sat. NaHCO_3 solution and 10 mL brine and is dried over Na_2SO_4 . The solvent is removed in vacuum to give the product as a colorless solid.

Deprotection of Aldehyde

The 1,3-dioxolane moiety is converted into an aldehyde function through acidic hydrolysis.

5 Reaction 9 and Example 9: Synthesis of 404-47



As an illustrative example, a solution of 404-33 (0.246 mmol) in 20 mL formic acid is stirred at room temperature for 45 minutes. One hundred mL ice-water and 200 mL sat. NaHCO_3 solution are added slowly to the reaction (strong foaming). The reaction is extracted with 2 x 150 mL EtOAc. The combined extracts are washed with sat. NaHCO_3 solution, water and brine and are dried over Na_2SO_4 . The solvent is removed and the product is dried in vacuum.


15

Reduction of the Nitro Group

The aromatic nitro compound is reduced to the aniline through catalytic hydrogenation. The reaction leads to the reduction of the double bond.

20

Reaction 10 and Example 10: Synthesis of 404-120



As an illustrative example, 404-89 (0.13 mmol) is dissolved in 2 mL ethanol and
25 Raney-Nickel (0.18 g, 50 % in H₂O, washed 3 times with ethanol, then


- 72 -

suspended in 2 mL ethanol) and 0.1 mL acetic acid are added. The reaction is stirred at room temperature for 2 days. The reaction is filtered through Celite and the filter cake is washed with methanol. The filtrate is brought to dryness. The residue is taken up in EtOAc, washed with NaHCO₃ solution and brine and 5 is dried over Na₂SO₄. The solvent is removed in vacuum. The crude product is purified over silica gel (hexane/acetone 2:1).

Amide Synthesis

Amides are prepared from carboxylic acids by reaction of an amine with the 10 corresponding acid chloride (Reaction 11). The synthesis can also proceed directly from the acid by use of appropriate coupling reagents, such as DCC and HOBr (Reaction 12).

Reaction 11:



15


Where R1 is a saturated or unsaturated, straight or branched aliphatic chain, R15 and R16 are independently hydrogen or a saturated or unsaturated, straight

- 73 -

or branched aliphatic chain, or where NR15R16 together forms a morpholinyl moiety.

Example 11: Synthesis of 404-85

5



As an illustrative example, 365-73 (0.04 mmol) and thionylchloride (68 mmol) are combined under nitrogen atmosphere and are heated to reflux for 2 hours. The reaction is allowed to cool and is concentrated to dryness. Twenty mL toluene is added and the reaction is concentrated to dryness again (2 times). The residue 10 is taken up in 5 mL anhydrous toluene and diethylamine (0.48 mmol) is added. The reaction is stirred at room temperature over night. Five mL H_2O are added and the mixture is extracted with 20 mL EtOAc. The extract is washed with brine and dried over Na_2SO_4 . The solvent is removed in vacuum and the crude product is purified over silica gel (hexane/acetone 3:1).

15


Using Reaction 11, the following compounds are further examples of the compounds that may be synthesized.

- 74 -

Compound	Starting Material	MS (Na ⁺)	Amine
404-83	365-73	1368.2	diethylamine
404-128	404-124	1311.9	anhydrous ammonia ¹
404-129	404-124	1340.1	Dimethylamine ²

¹ passed through reaction for 10 min at 0 °C; ² 2 M solution In THF


Reaction 12:



Where R1 is a saturated or unsaturated, straight or branched aliphatic chain, R15 and R16 are independently hydrogen or a saturated or unsaturated, straight or branched aliphatic chain, or where NR15R16 together forms a morpholinyl moiety.

10

Example 12: Synthesis of 420-104



As an illustrative example, 420-98 (0.078 mmol) is dissolved in 10 mL anhydrous DCM under nitrogen atmosphere. Dicyclohexylcarbodiimide (DCC, 0.117 mmol) and 1-hydroxybenzotriazole hydrate (HOBr, 0.078 mmol) are added at 0 °C and the mixture is stirred for 15 minutes. Dimethylamine (0.78 mmol) is added to 5 give a clear, colorless solution. The cooling bath is removed after 15 minutes and stirring is continued at ambient temperature for 5 days. The reaction is transferred to a separatory funnel and 20 mL DCM and 10 mL 0.5 M HCl are added. The organic layer is taken off, dried over Na₂SO₄ and concentrated to dryness. The residue is taken up in 10 mL acetonitrile. Undissolved solid is 10 filtered off and the filtrate is concentrated in vacuum. The crude product is purified by preparative HPLC.

Using Reaction 12, the following compounds are further examples of the compounds that may be synthesized.

15

Compound	Starting Material	MS (Na ⁺)	Amine
404-150	416-04 	1380.1	Dimethyl- amine ²
404-155	404-134 	1352.1	Dimethyl- amine ²
404-156	404-60 	1324.1	Dimethyl- amine ²
404-162	416-08 	1379.9	Morpholine

- 76 -


404-164		416-08		1309.8	anhydrous ammonia ¹
404-178		404-137		1323.9	Propylamine
420-104		420-98		1408.1	Dimethyl-amine ²
420-114		420-100		1366.0	Dimethyl-amine ²
420-121		420-100		1338.0	anhydrous ammonia ¹

¹ passed through reaction for 10 min at 0 °C; ² 2 M solution in THF

Esterification

Carboxylic acid esters are prepared from the corresponding carboxylic acids and 5 an alcohol either using acidic catalysis (Reaction 13) or coupling reagents (DCC and DMAP, Reaction 14).

Reaction 13:



- 77 -

Where R1 is a saturated or unsaturated, straight or branched aliphatic chain, and R17 is a saturated or unsaturated, straight or branched aliphatic chain, optionally containing a halogen or hydroxyl substituent.

5 Example 13: Synthesis of 404-171



As an illustrative example, a mixture of 404-60 (0.059 mmol), 4 mL EtOH and 2 μ L conc. H_2SO_4 is heated to reflux for 4 hours. The solvent is evaporated and the residue is taken up in acetonitrile. The crude product is purified by 10 preparative HPLC.

Using Reaction 13, the following compounds are further examples of the compounds that may be synthesized.


Compound	Starting Material	MS (Na^+)	Reagent
404-171	404-60	1368.2	ethanol
404-182	404-60	1311.9	ethylene glycol ¹

- 78 -

420-103	420-98	1409.1	ethanol
420-113	420-100	1366.9	ethanol

[†] 3 hours at 90 °C; product extracted with EtOAc


Reaction 14:



Where R1 is a saturated or unsaturated, straight or branched aliphatic chain, and R17 is a saturated or unsaturated, straight or branched aliphatic chain, optionally containing a halogen or hydroxyl substituent.

10

Example 14: 420-24



15 As an illustrative example, 404-60 (0.053 mmol) is dissolved in 4 mL anhydrous DCM and cooled to 0 °C under nitrogen atmosphere. Dimethylaminopyridine (DMAP, 0.005 mmol), 2-fluoropropanol (0.27 mmol) and dicyclohexylcarbodiimide (DCC, 0.058 mmol) are added and the reaction is stirred for 15 min at 0 °C. The cooling bath is removed and stirring is continued

20 over night at ambient temperature. 20 mL DCM are added, the reaction is then

- 79 -

washed with H_2O and evaporated to dryness. The residue is taken up in 10 mL acetonitrile and filtered. The filtrate is concentrated in vacuum. The crude product is purified by preparative HPLC.

5 *Alcohols*

Besides direct synthesis in the Wittig reaction, alcohols are obtained through a number of reactions. Reduction of a carbonyl group with sodium borohydride leads to primary (starting from aldehyde) or secondary (starting from ketone) alcohols, respectively.


10

Oxidation of a double bond through the hydroboration method can lead to a mixture of isomers. The reaction proceeds predominantly in anti-Markovnikov orientation. In the case of a terminal olefin the primary alcohol is the main product.

15

An olefin can be converted into a diol through oxidation with hydrogen peroxide. Reaction of a carbonyl compound with a Grignard reagent leads to secondary (starting from aldehyde) and tertiary (starting from ketone) alcohols. This method allows for an extension of the carbon chain.

20 Reaction 15:



Where R' is a H or acetyl, $R1$ is a saturated or unsaturated, straight or branched

25 aliphatic chain, and $R20$ is a saturated or unsaturated, straight or branched aliphatic chain.

- 80 -

Example 15: Synthesis of 404-98



As an illustrative example, 404-61 (0.0365 mmol) is dissolved in 4.5 mL

5 anhydrous EtOH under nitrogen atmosphere. Sodium borohydride (0.15 mmol, suspended in 0.5 mL anhydrous EtOH) is added at 0 °C and the resulting mixture is stirred at ambient temperature over night. Additional sodium borohydride (0.08 mmol) is added and stirring is continued over night. The reaction is quenched with 5 mL 1 M HCl under ice-bath cooling and is extracted
 10 with EtOAc. The extract is washed with brine, dried over Na₂SO₄ and concentrated to dryness. The crude product is purified by preparative HPLC.

Using Reaction 15, the following compounds are further examples of the compounds that may be synthesized.


15

Compound	Starting Material	MS (Na ⁺)
404-98	404-61	1256.9
404-195	404-173	1271.0
404-198	404-172	1313.0

- 81 -

420-09	404-56	1298.9


Reaction 16:



5

Where R1 is a saturated or unsaturated, straight or branched aliphatic chain.

Example 16: Synthesis of 420-28-1




10 As an illustrative example, 404-16 (0.081 mmol) is dissolved under nitrogen atmosphere in 4 mL anhydrous THF. The reaction is cooled to 0 °C and $\text{BH}_3\cdot\text{THF}$ (1 M sol. in THF, 0.06 mmol) is added. The reaction is stirred at room temperature over night. HPLC shows the reaction is incomplete. Additional $\text{BH}_3\cdot\text{THF}$ (0.5 mmol) is added and stirring is continued for 4 hours at room temperature. The reaction is cooled to 0 °C and 1.0 mL 1 M NaOH and 0.30 mL 30 % hydrogen peroxide solution are added. The mixture is stirred at room temperature over night. The reaction is extracted with 25 mL EtOAc. The extract is washed with brine, dried over Na_2SO_4 and concentrated to dryness. The product is purified by preparative HPLC.

15

20


Reaction 17:



- 82 -


Where R1 is a saturated or unsaturated, straight or branched aliphatic chain, R' is either a H or an acetyl group.

5 Example 17: Synthesis of 420-49



As an illustrative example, 420-49 (0.037 mmol) is dissolved under argon atmosphere in 5 mL anhydrous THF. The reaction is cooled to -70 °C and 10 allylmagnesium chloride (1 M sol. in THF, 0.22 mmol) is added. The reaction is stirred for 15 minutes at -70 °C and is then allowed to come to room temperature. After 90 minutes the reaction is quenched with sat. NH₄Cl solution. The reaction is extracted with 25 mL EtOAc. The extract is washed with brine, dried over Na₂SO₄ and concentrated to dryness. The product is purified by 15 preparative HPLC. A mixture of acetylated and deacetylated compound is obtained.

Reaction 18:




20 Formula XLV

Formula XLVI

Where R1 is a saturated or unsaturated, straight or branched aliphatic chain, and R23 is a saturated or unsaturated, straight or branched aliphatic chain.

- 83 -

Example 18: Synthesis of 404-126




As an illustrative example, 404-16 (0.054 mmol) is dissolved in 1 mL formic acid
 5 and hydrogen peroxide (30 % aqueous solution, 0.52 mmol) is added. The
 reaction is stirred at room temperature over night and is then concentrated to
 dryness. The residue is dissolved in 25mL EtOAc, washed with sat. NaHCO_3
 solution and dried over Na_2SO_4 . The solvent is removed in vacuum. The
 reaction is taken up in 9 mL THF and 3 mL 1 M NaOH, and is stirred for 4 hours
 10 at room temperature. The solvent is removed and the residue is partitioned
 between 25 mL EtOAc and 5 mL H_2O . The organic layer is washed with brine
 and dried over Na_2SO_4 . The solvent is evaporated and the crude product is
 purified by preparative HPLC.

15 **Example 19: Modification of Amino Acid 3**

CsA undergoes substitution on AA3 as outlined below. Reaction with excess
 20 LDA (lithium diisopropylamide) leads to a hexolithio derivative containing four
 lithium azaenolate units as well as a lithium alkoxide unit on the amino acid 1
 side chain and a lithium enolate unit on AA3, respectively. Subsequent reaction
 with a suitable electrophile generates substitution products on the AA3
 (sarcosine) residue. Suitable electrophiles are e.g. alkyl halides, aldehydes,
 25 carbon dioxide and alkyl disulfides (Table 1). Both D and L epimers can be
 obtained, with the relative ratios depending on the reaction conditions. Route A
 (see below) leads predominantly to the D product, while Route B (addition of
 excess LiCl) gives mixtures of both epimers.

- 84 -



Example 19: Substitution reaction at AA3 of Cyclosporin A. D and L stereoisomers are obtained.

5

Route A: [D-MeSar]³-CsA

An oven dried flask is charged under argon atmosphere with 160 mL anhydrous THF and diisopropylamine (2.07 mL, 14.8 mmol). The solution is cooled to -78 °C and n-butyl lithium (2.5 M in hexane, 5.4 mL, 13.5 mmol) is added. After stirring for 30 minutes, CsA (2.40 g, 2.0 mmol, dissolved in 40 mL anhydrous THF) is added. The reaction is stirred for 1 hour at -78 °C. Additional n-butyl lithium (3.2 mL, 8.0 mmol) is added, followed by addition of methyl iodide (1.25 mL, 20.0 mmol). Stirring is continued at -78 °C for 1.5 hours, and then the reaction is allowed to warm to room temperature over an additional 1.5 hours. 10 20 mL H₂O are added and the THF is removed in vacuum. Additional 50 mL H₂O are added and an extraction is carried out with 150 mL EtOAc. The extract is washed with brine and dried over Na₂SO₄. The solvent is removed in vacuum and the crude product is purified over silica gel (hexane/acetone 3:1). Yield: 15 0.74 g (0.61 mmol, 30 %).

20

Route B: [MeSar]³-CsA

A dry 100 mL flask is charged under argon atmosphere with 7.5 mL anhydrous THF and diisopropylamine (0.46 mL, 3.3 mmol). The solution is cooled to 0 °C and n-butyl lithium (1.32 mL, 2.5 M solution in hexane, 3.3 mmol) is added. The

reaction is stirred for 20 minutes at 0 °C and is then cooled to -78 °C. A solution of CsA (601 mg, 0.5 mmol) and lithium chloride (636 mg, 15 mmol) in 12 mL anhydrous THF is prepared and cooled to -78 °C under argon atmosphere. The LDA solution is then transferred into this mixture through a cannula. The 5 reaction is stirred at -78 °C for 2 hours. Additional n-butyl lithium (1.20 mL, 3.0 mmol) is added, followed by methyl iodide (0.62 mL, 10 mmol). The mixture is allowed to warm to -20 °C and stirred at this temperature for 3 hours. The reaction is allowed to warm to room temperature, quenched with saturated NH₄Cl solution, extracted with EtOAc (2 x 20 mL), washed with brine and dried 10 over Na₂SO₄. The solvent is removed in vacuum and the crude product is purified over silica gel (hexane/acetone 3:1). Yield: [L-MeAla³]-CsA: 302 mg (0.25 mmol, 50 %). [D-MeAla³]-CsA: 76 mg (0.06 mmol, 12 %).

15 Table 1: Examples of possible electrophiles used for the alkylation of the 3-position of Cyclosporin.

Electrophile	R23 ¹	Route	Remarks
methyl iodide	—CH ₃	A/B	-
ethyl iodide	—CH ₂ CH ₃	A/B	-
allyl bromide	—CH ₂ CHCH ₂	B	-
1,3-diiodopropane	—CH ₂ CH ₂ CH ₂ I	B	-
1,4-diiodobutane	—(CH ₂) ₃ CH ₂ I	B	-
trimethylammonium- 3-iodopropane	—(CH ₂) ₃ N ⁺ (CH ₃) ₃	B	-
hexafluorophosphate			
propargyl bromide	—CH ₂ CCH	A	10 equiv electrophile; stirred for 1 h at room temperature after electrophile addition
tert-butyl bromoacetate	—CH ₂ CO ₂ (t-Bu)	A	5 equiv electrophile; stirred for 1 h at room temperature after electrophile addition
benzyl bromide	—CH ₂ Ph ²	A	15 equiv LDA and 30 equiv

- 86 -

			electrophile; stirred for 6 h at - 75 °C and 10 h at room temperature after electrophile addition
formaldehyde	—CH ₂ OH	A	formaldehyde prepared from paraformaldehyde at 170 °C prior to addition
acetaldehyde	—CH(OH)CH ₃	B	stirred for 2.5 h at -78 °C after electrophile addition
pivalaldehyde	—CH(OH)(t-Bu) ³	B	stirred for 70 min at -78 °C after electrophile addition
benzaldehyde	—CH(OH)Ph	B	stirred for 2.5 h at -78 °C after electrophile addition
carbon dioxide	—COOH	A	CO ₂ gas passed for 15 min through reaction mixture at - 78 °C; stirred for 1 h at -78 °C after electrophile addition
dimethyl disulfide	—SCH ₃	B	stirred for 18 h at 0 °C after electrophile addition
p-tolyl disulfide	—S(p-Tol) ⁴	B	stirred for 18 h at 0 °C after electrophile addition

¹ according to Example 19 ² Ph = phenyl; ³ t-Bu = tert-butyl; ⁴ Tol = toyl.

Examples 20 and 21, set out below, are general examples of the chemical reactions able to synthesize the desired compounds modified at amino acid 1 and 3 of CsA using reagents that have the requisite chemical properties, and it would be understood by a person skilled in the art that substitutions of certain reactants may be made.

Example 20: AA1 Modification of Alkylated CsA

10 Example 20 provides a synthetic route for the introduction of substituents at the 3-position of CsA prior to modification of the AA1 side-chain. Following the 3-

- 87 -

alkylation, a 2 step procedure leads to the acetylated aldehyde (compound **3** in the example below), which is a suitable substrate for the modification of the 1-position via Wittig reaction. This method allows introduction of residues to the AA1 side-chain that have limited stability under the reaction conditions used in 5 steps 1-3, such as strong base and oxidizing agents.


Further examples of compounds prepared using this sequence is summarized in Table 2.

Step 1: Alkylation of AA3 side-chain



Synthesis is carried out according to Route A or B, respectively, as described above.

Step 2: Acetylation of the hydroxy-group on AA1 side-chain




An oven dried flask is charged under nitrogen with $[\text{D-MeSar}]^3\text{-CsA}$ (1.84 g, 1.51 mmol), N,N-dimethylaminopyridine (19 mg, 0.15 mmol) and 20 mL anhydrous pyridine, followed by acetic anhydride (10 mL, 0.1 mol). The reaction is stirred at ambient temperature over night. The mixture is poured into 100 mL ice-water 20 and is stirred until all ice has melted. A solid is collected by filtration and dried in air. The solid is dissolved in 50 mL EtOAc and is washed with 1 M HCl (2x), sat. NaHCO_3 solution and brine. The organic phase is dried over Na_2SO_4 and evaporated. The crude product is purified over silica gel (hexane/EtOAc/MeOH 10:10:0.5).

25

- 88 -


Step 3: Aldehyde formation



To a flask containing compound **2** (800 mg, 0.636 mmol) are added 10 mL dioxane and 10 mL H_2O . NaIO_4 (544 mg, 2.54 mmol) and OsO_4 (7.9 mM solution in water/dioxane 1:1, 4.05 mL, 32 mmol) are added and the reaction is stirred at room temperature over night. 75 mL H_2O is added and the reaction is extracted with 3 x 25 mL EtOAc. The extracts are washed with water, sat. NaHCO_3 solution, water and brine (25 mL each) and are dried over MgSO_4 . The solvent is removed in vacuum and the crude product is purified over silica gel (hexane/EtOAc 3:1).

Step 4: Wittig


Reaction





An oven dried flask is charged under argon atmosphere with triphenyl-6-hexanoic acid phosphonium bromide (90 mg, 0.195 mmol) and 5 mL anhydrous THF. Potassium t-butoxide (1 M solution in THF, 0.39 mL, 0.39 mmol) is added at 0 °C and the solution is stirred for 30 minutes to give a bright orange color. Compound **3** (81 mg, 0.065 mmol, dissolved in 1 mL anhydrous THF) is added to the reaction drop-wise and stirring is continued at room temperature over night. The reaction is quenched with sat. NH_4Cl solution and is extracted with EtOAc. The extract is washed with brine and dried over Na_2SO_4 . The solvent is removed in vacuum and the crude product is purified over silica gel (toluene/acetone 3:1).

- 89 -

Step 5: Deacetylation



Compound 4 (30 mg, 0.022 mmol) is dissolved in 2 mL methanol and 0.5 mL water and tetramethylammonium hydroxide pentahydrate (12 mg, 0.066 mmol) is added. The reaction is stirred at room temperature for several days until HPLC confirms deprotection is complete. The reaction is acidified to pH 2 with 1 M HCl and concentrated in a vacuum. The residue is taken up in EtOAc, is washed with water and dried over Na_2SO_4 . The solvent is evaporated and the crude product is purified by preparative HPLC.



- 90 -



Schematic representation of 1,3-modified Cyclosporin derivatives.

Using the method of Example 20, the following compounds are further examples of the compounds that may be synthesized (X and Y in reference to the above

5 schematic representation; and reference of R in X is to indicate attachment of structure to AA1 of CsA).

Compound	X	R23	Isomer	MS (Na ⁺)
431-13		-CH ₃	L	1296.8
414-64		-CH ₃	D	1325.0
431-19		-CH ₃	L	1324.8
431-40		-CH ₂ CH ₃	D	1338.8
440-02		-CH ₃	D	1327.1
431-20		-CH ₃	L	1326.9
440-13		-CH ₂ CH ₃	D	1341.1
431-21		-CH ₃	D	1277.9
431-44		-CH ₂ CH ₃	L	1291.9
440-14		-CH ₂ CH ₃	D	1292.0
431-136		-CH ₃	L	1283.1
440-24		-CH ₃	D	1283.1
440-10-1		-CH ₃	D	1270.9

- 91 -

440-22-1		-CH ₂ CH ₃	D	1285.0
440-20		-CH ₃	D	1397.2


Alkylation of AA1 Modified Compounds

Reaction 21 introduces substituents to the AA3 residue of compounds previously modified on the AA1 side-chain. In addition to the groups available through

5 Reaction 19, this route allows the introduction of substituents at AA3 that are unstable under the reaction conditions used in Reaction 20, e.g. a thiomethyl residue could undergo oxidation during the formation of the aldehyde in step 3 of this method.

10

Example 21



A dry 25 mL flask is charged under argon atmosphere with 1.5 mL anhydrous

15 THF and diisopropylamine (87 μ L, 0.62 mmol). The solution is cooled to 0 °C and n-butyl lithium (2.5 M in hexane, 0.25 mL, 0.62 mmol) is added. The mixture is stirred for 20 minutes at 0 °C and is then cooled to -70 °C. The clear LDA solution is transferred into a solution of 404-76 (118 mg, 0.095 mmol) and lithium chloride (120 mg, 2.84 mmol) in 1.5 mL anhydrous THF at -70 °C. Stirring is

20 continued for 2 hours at -70 °C. Additional n-butyl lithium (0.23 mL, 0.58 mmol) is added, followed by methyl iodide (118 μ L, 1.89 mmol). The reaction is allowed to warm to -20 °C and is kept at this temperature over night. The reaction is quenched with sat. NH₄Cl solution and is extracted with EtOAc. The extract is washed with brine, dried over Na₂SO₄ and evaporated to dryness. The

25 crude product is purified over silica gel (hexane/acetone 3:1 \rightarrow 2:1).

- 92 -

Table 3: Examples of compounds prepared by Method 21 (X and Y according to Figure 3; and reference of R in X is to indicate attachment of structure to AA1 of CsA).

Compound	X	R23	Isomer	MS (Na ⁺)
420-176-1		-CH ₃	D	1284.9
420-176-2		-CH ₃	L	1284.9
420-177-1		-CH ₃	D	1298.8
420-177-2		-CH ₃	L	1298.8
420-180-1		-CH ₃	D	1302.8
420-182-1		-CH ₃	D	1270.8
420-182-2		-CH ₃	L	1270.8
420-186		-CH ₂ CH ₃	L	1409.0
431-42		-SCH ₃	D/L ¹	1328.8
440-03		-SCH ₃	D	1319.1
440-78		-(CH ₂) ₃ N ⁺ (CH ₃) ₃	D/L ¹	1359.9 ²
440-34-3		-CH ₂ -phenyl	D	1416.2
440-36		-CH ₃	D	1251.1


¹ isomers not separated; ² m⁺ signal.

- 93 -

Additional modifications to the functional groups at AA1 (or AA3, respectively) residue can be carried out to obtain various derivative compounds, such as esters, amides, alcohols etc. Saturated compounds can be obtained by reducing the double bond created in the Wittig reaction.


5

Example 22: Amide Formation from Carboxylic Acid – Synthesis of 440-08



440-02 (48 mg, 0.037 mmol) is dissolved under nitrogen atmosphere in 5 mL
10 anhydrous DCM and cooled to 0 °C. Dicyclohexylcarbodiimide (DCC, 11.6 mg,
0.056 mmol) and 1-hydroxybenzotriazole (HOBT, 5.0 mg, 0.037 mmol) are added
and the mixture is stirred for 15 minutes at 0 °C. Dimethylamine (2 M solution in
THF, 0.19 mL, 0.38 mmol) is added and stirring is continued at room
temperature for 3 days. The reaction is diluted with 20 mL DCM and washed
15 with 15 mL 0.5 M HCl. The organic phase is dried over Na₂SO₄ and then
brought to dryness. The crude mixture is purified by preparative HPLC.

Example 23: Ester Formation – Synthesis of 440-31




440-20 (30 mg, 0.022 mmol) is dissolved in 4 mL anhydrous EtOH and 2 μ L of conc. H_2SO_4 . The reaction is heated to reflux for 3 hours and is then allowed to cool to room temperature. The reaction is brought to dryness. The crude product is purified by preparative HPLC.

25


Example 24: Amide Formation from Nitrile Compound (Reversed Amide) – Synthesis of 440-15

- 94 -




A 50 mL flask is charged under nitrogen with 440-09 (80 mg, 0.061 mmol) and 5 mL MeOH. The reaction is cooled to 0 °C and Ni(II)Cl₂·6H₂O (1.4 mg, 0.006 mmol) and acetic anhydride (19 μ L, 0.20 mmol) are added. Sodium borohydride (104 mg, 2.75 mmol) is added in 2 batched 2 hours apart. The reaction is then allowed to warm to room temperature and is stirred over night. After the reaction is complete, 7 mL 1 M HCl is added. The solution is concentrated in vacuum to approximately half of its original volume. The resulting mixture is extracted with EtOAc, the extract is washed with sat. NaHCO₃ solution and brine and is dried over Na₂SO₄. The solvent is removed in vacuum. The product, which contains some saturated compound, is used in the following step without further purification.

15 Example 25: Synthesis of 440-25



440-15 (83 mg, 0.061 mmol) is dissolved in 10 mL anhydrous EtOH. Palladium (10 wt % on Carbon, 8 mg) and 3-4 drops of acetic acid are added. The reaction is hydrogenated under atmospheric pressure at room temperature for several days until complete by HPLC. The reaction is filtered through Celite and the filtrate is evaporated to dryness. The crude product is purified by prep. HPLC and is then subjected to the deacetylation step.

25 Example 26: Synthesis of 440-32









- 95 -

440-25 (41 mg, 0.03 mmol) is dissolved in 4 mL MeOH and tetramethylammonium hydroxide pentahydrate (16 mg, 0.09 mmol, dissolved in 1 mL H₂O) is added. The reaction is stirred at room temperature for 2 days. The reaction is concentrated in vacuum. 5 mL H₂O is added and the product is 5 extracted with EtOAc. The extract is washed with brine, dried over Na₂SO₄ and evaporated to dryness. The crude product is purified by preparative HPLC.

Table 4: Examples of derivatives of 1,3-modified Cyclosporin compounds obtained by reducing the double bond created in the Wittig reaction (X and Y 10 according to Figure 3; and reference of R in X is to indicate attachment of structure to AA1 of CsA).

Compound	X	R23	Isomer	MS (Na ⁺)
431-23		-CH ₃	L	1323.8
431-29		-CH ₃	L	1393.9
440-08		-CH ₃	D	1354.1
440-27		-CH ₂ CH ₃	D	1368.1
440-23		-CH ₃	D	1424.2

- 96 -

431-32		-CH ₃	D	1325.9
431-53		-CH ₂ CH ₃	L	1339.9
440-32		-CH ₂ CH ₃	D	1340.1
440-10-2		-CH ₃	D	1285.0
440-22-2		-CH ₂ CH ₃	D	1299.1
440-31		-CH ₃	D	1425.2

Cyclophilin A Isomerase Inhibition Assay

An enzymatic assay was used to measure the inhibition of CyP-A activity by 1,3
5 CsA analogs of the present invention, according to a protocol described in the
scientific literature with minor modifications. The assay is based on the ability of
CyP-A to catalyze a conformational change in proline-containing peptides from
cis to *trans* isomeric conformations. Briefly, a peptide substrate that includes a
nitroanilide moiety was supplied to a reaction mixture containing CyP-A, test
10 compound (CsA analog, CsA, or dimethylsulfoxide vehicle), and a second
enzyme, alpha-chymotrypsin. Each test compound was tested at 10
concentrations in triplicate or quadruplicate. The peptide was converted from the
cis conformation to the *trans* conformation both by non-catalytic and CyP
catalytic processes. The *trans* isomer of the peptide, but not the *cis* isomer, is a
15 substrate for alpha-chymotrypsin. Alpha-chymotrypsin immediately cleaved
nitroanilide from the rest of the peptide, and free nitroanilide accumulated at a
rate proportional to *cis-trans* isomerization. Since free nitroanilide is a colored
product, its accumulation was quantified by measuring its absorbance with a

- 97 -

spectrophotometer. Nitroanilide accumulation was measured for 6 minutes, and first order rate constants for each reaction were calculated using Graphpad Prism software. The CyP-A catalytic rate constant of each reaction was determined by subtracting the non-catalytic rate constant (derived from the 5 reaction without CyP-A) from the total reaction rate constant. Plots of the catalytic rate constants as a function of inhibitor concentrations demonstrated the compounds' potencies, defined by their IC₅₀ values.

DETAILED PROTOCOL

10

A. Peptide

The assay peptide was N-succinyl-alanine-alanine-proline-phenylalanine-p-nitroanilide. It was dissolved to a concentration of 3 mM in a solution of trifluoroethanolamine and lithium chloride (TFE/LiCl). TFE/LiCl was prepared 15 fresh each day by dissolving lithium chloride in trifluoroethanolamine to a concentration of 17 mg/ml. Following dissolution of LiCl, the water content of the TFE/LiCl solution was reduced by adding heat-dried molecular sieves and gentle mixing the solution for at least 30 minutes. The peptide was then dissolved in TFE/LiCl, and the solution cooled to 4°C – 8°C prior to the assays. Dissolution 20 of the peptide in dry TFE/LiCl promoted more peptide to exist in the *cis* conformation at the beginning of each assay reaction. Data analysis showed that approximately 60% of the peptide in our assays began as a *cis* isomer which is consistent with reported data in the scientific literature. In the enzyme reactions the peptide was diluted 20-fold to a final assay concentration of 150 µM.

25

B. Test Compounds

The test compounds consisted of CsA, CsA analogs, or dimethylsulfoxide (DMSO). Stock solutions of CsA and CsA analogs were made by dissolution in DMSO to a concentration of 10 mg/ml in sterile microcentrifuge tubes. Stock 30 solutions were stored at -20°C when not in use. Further dilutions of the test compounds were made on each day of the assays. DMSO and CsA were tested in every experiment to serve as the vehicle control and reference compound,

- 98 -

respectively.

The 10 mg/ml stock solutions of CsA and CsA analogs were diluted with DMSO to 50 μ M in microcentrifuge tubes, based on the molecular weights of the
5 compounds. Nine 3-fold serial dilutions of each compound in DMSO were then made in a 96-well polystyrene plate. An aliquot of DMSO-solution or DMSO vehicle alone was diluted 50-fold in reaction buffer (see below for recipe) to make final concentrations of CsA or CSA ANALOGs of 1000, 333, 111, 37, 12, 4.1, 1.4, 0.46, 0.15, and 0.05 nM. The reaction buffer solutions were stored at
10 4°C – 8°C for at least one hour prior to the assays.

C. Reaction Buffer

The starting solution (saline buffer) for the reaction buffer consisted of Hepes 50 mM, sodium chloride 100 mM, and human serum albumin 1 mg/ml, adjusted to
15 pH 8.0 with sodium hydroxide. The saline buffer was stored at 4°C when not in use. On each assay day bovine alpha-chymotrypsin was dissolved in a volume of saline buffer to a concentration of 1 mg/ml. An aliquot of the alpha-chymotrypsin solution was removed to serve as the noncatalytic control reaction buffer. Recombinant human CyP-A was added to the remainder of the
20 chymotrypsin solution to a concentration of 5 nM. The solution containing alpha-chymotrypsin and CyP-A was designated the reaction buffer and was used for preparation of the reaction solutions.

D. Reaction Protocol

25 All assay reactions were conducted in a cold room within a temperature range of 4°C – 8°C. All solutions and equipment were stored in the cold room for at least 1 hour prior to the assays. The low temperature was necessary for reactions to proceed at a sufficiently slow rate to measure with the available equipment. The measuring device was a BMG Polarstar microplate reader configured for
30 absorbance readings at OD 405 nm. Reactions were performed in 96-well, flat-bottom, polystyrene assay plates. Each assay run consisted of 12 separate reactions in one row of the plate. Peptide was aliquoted at 5 μ l per well with a

- 99 -

single-channel pipettor in one row of the plate, then the plate placed in the plate holder of the microplate reader. Reactions were begun by dispensing 95 μ l of reaction buffer into each peptide-containing well using a 12-channel pipettor and mixing each reaction thoroughly by repeat pipetting to ensure uniform dissolution
5 of the peptide. The 12 reactions in each assay run were represented by the following:

- a) 10 reactions, representing one replicate for each of the 10 concentrations of one test compound (CyP-A in reaction buffer)
- 10 b) 1 reaction with 5 μ l DMSO vehicle (CyP-A in reaction buffer)
- c) 1 reaction with 5 μ l DMSO vehicle (CyP-A A absent from reaction buffer)

Absorbance recordings were begun immediately after mixing. Approximately 15 seconds elapsed from addition of the reaction buffer to the first OD_{405} recording
15 due to mixing time and instrument setup. Subsequent readings were made at 6-second intervals for a total of 60 readings over 360 seconds. Three or four reaction runs were made for each test compound to provide data replicates.

E. Data Analysis

20 The raw data consisted of a time-dependent increase in OD_{405} . In the presence of CyP-A and the absence of inhibitor the peptide was completely converted to the *trans* isomer within approximately 150 seconds as demonstrated by a plateau in the OD_{405} . OD_{405} vs. time data were plotted with Graphpad Prism software and fitted with a one phase exponential equation to derive a first order
25 rate constant K for each reaction. In reactions without CyP-A, the rate constant entirely represented the spontaneous noncatalytic, thermal *cis*-to-*trans* isomerization of the peptide and was defined as the noncatalytic rate constant K_0 . In reactions containing CyP-A, isomerization occurred both through noncatalytic and enzyme-catalyzed processes. Thus, the rate constant K in CyP-
30 A-containing reactions represented the sum of the noncatalytic rate constant K_0 and the catalytic rate constant K_{cat} . K_{cat} was calculated by subtracting K_0 (obtained from the reaction without CyP-A) from the total rate constant K. K_{cat}

- 100 -


typically was 3-fold higher than K_0 in reactions with 5 nM CyP-A, 150 μ M peptide substrate, and no inhibitor.

Plots of K_{cat} versus inhibitor concentration were fitted with sigmoidal dose-response nonlinear regressions to demonstrate inhibitor potencies. Software-calculated EC₅₀ values represented the test compound concentrations that inhibited K_{cat} by 50%. To normalize for inter-experiment variability in assay conditions, CsA was run in every experiment as a reference compound, and CsA analog potency was expressed as a fold-potency relative to CsA based on EC₅₀ values. For example, a CsA analog EC₅₀ that was 1/2 of CsA represented a 2-fold potency compared to CsA, whereas a CSA analog IC₅₀ that was 5-fold higher than CsA represented a 0.2-fold potency compared to CsA.

Table 5, shown in the attached Figure, shows cyclophilin A inhibition and immunosuppression of CsA analogs modified at position 1 and at positions 1 and 3 according to the present invention.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound of Formula L:






















wherein:


- a. R' is H or acetyl;
- b. R1 is a saturated or unsaturated straight or branched aliphatic carbon chain from 2 to 15 carbon atoms in length;
- c. R2 is selected from the group consisting of:
 - i. an unsubstituted, N-substituted, or N,N-disubstituted amide;
 - ii. a N-substituted or unsubstituted acyl protected amine;
 - iii. a carboxylic acid;
 - iv. a N-substituted or unsubstituted amine;
 - v. a nitrile;
 - vi. an ester;
 - vii. a ketone;
 - viii. a hydroxy, dihydroxy, trihydroxy, or polyhydroxy alkyl; and
 - ix. a substituted or unsubstituted aryl;
 - x. a saturated or unsaturated straight or branched aliphatic carbon chain substituted with a substituent selected from the group consisting of ketones, hydroxyls, nitriles, carboxylic acids, esters, 1,3-dioxolanes, halogens, and oxo;
 - xi. an aromatic group substituted with a substituent selected from the group consisting of halides, esters, and nitro; and
 - xii. a combination of the saturated or unsaturated straight or branched aliphatic carbon chain of (x) and the aromatic group of (xi); and


d. R23 is an unsubstituted saturated or unsaturated straight or branched aliphatic carbon chain.


2. The compound of claim 1, wherein R1-R2 is selected from the group consisting of:


- i. ;
- ii. ;
- iii. ;
- iv. ;
- v. ;
- vi. ;
- vii. ;
- viii. ;
- ix. ;
- x. ;
- xi. ;
- xii. ;
- xiii. ;
- xiv. ;
- xv. ;
- xvi. ;
- xvii. ;
- xviii. ;
- xix. ;


xx. ;

xxi. ;


xxii. ;


xxiii. ;


xxiv. ;


xxv. ;


xxvi. ;


xxvii. ;

xxviii. ;


xxix. ;

xxx. ;


















xxxi. ;


xxxii. ;

xxxiii. ;


xxxiv. ;


xxxv. 





3. The compound of claim 1, wherein R2 is selected from the group consisting of





wherein:

- i. R5 is a saturated or unsaturated straight or branched aliphatic carbon chain between 1 and 10 carbons in length; and
- ii. R6 is a monohydroxylated, dihydroxylated, trihydroxylated, or polyhydroxylated saturated or unsaturated straight or branched aliphatic carbon chain between 1 and 10 carbons in length.

4. The compound of any one of claims 1 to 3, wherein R' is H.

5. The compound of claim 1 or 4, wherein R1-R2 comprises a saturated or unsaturated straight or branched aliphatic carbon chain between 2 and 5

carbons substituted with a substituent selected from the group consisting of ketones, hydroxyls, nitriles, halogens, oxo, carboxylic acids, esters, and 1,3-dioxolanes.

6. The compound of any one of claims 1 to 5, wherein R23 is selected from the group consisting of:


- i. $-\text{CH}_3$
- ii. $-\text{CH}_2\text{CH}_3$
- iii. $-\text{CH}_2\text{CHCH}_2$
- iv. $-\text{CH}_2\text{CCH}$

7. The compound of any one of claims 1 to 5, wherein R23 is C1-C3 alkyl.


8. The compound of claim 7, wherein R23 is ethyl and the compound is a D-epimer, wherein the chiral center of the D-epimer is the carbon to which R23 is attached.


9. The compound of claim 7, wherein R23 is methyl and the compound is a D-epimer, wherein the chiral center of the D-epimer is the carbon to which R23 is attached.

10. The compound of claim 1, wherein




is selected from the group consisting of:






11. The compound of any one of claims 1 to 5, wherein R23 is a straight or branched aliphatic carbon chain of 1 to 6, 1 to 5, 1 to 4, 1 to 3 or 2 carbons in length.

12. A compound of formula:



wherein:

R' is H or acetyl;




R23 is ethyl or methyl, and

wherein the compound is a D-epimer, wherein the chiral center of the D-epimer is the carbon to which R23 is attached.

13. The compound of claim 12, wherein R23 is methyl.

14. A compound of formula:



wherein:

R' is H or acetyl;

R1-R2 is ; and

R23 is methyl, and

wherein the compound is a D-epimer, wherein the chiral center of the D-epimer is the carbon to which R23 is attached.

15. A pharmaceutical composition comprising the compound of any one of claims 1 to 14 and one or more pharmaceutical excipients.

16. A method of treating or preventing a cyclophilin mediated disease in a mammal comprising administering the compound of any one of claims 1 to 14 or the pharmaceutical composition of claim 15 to the mammal under conditions to treat the cyclophilin mediated disease or injury, or use of said compound or composition to treat said disease or injury.

17. The method or use of claim 16, wherein the disease or injury is mediated by the over expression of cyclophilin or the disease is a congenital over expression of cyclophilin.

18. The method or use of claim 17, wherein the cyclophilin mediated disease or injury is selected from the group consisting of

a viral infection;

inflammatory disease;

cancer;

muscular disorder;

neurological disorder; and

injury associated with ischemia, reperfusion, loss of cellular calcium homeostasis, loss of ionic homeostasis, increase in free radical production, or toxins that induce mitochondrial dysfunction;

wherein the viral infection is optionally caused by a virus selected from the group consisting of Human Immunodeficiency virus, Hepatitis A, Hepatitis B, Hepatitis C, Hepatitis D, Hepatitis E, SARS-CoV, hCoV-NL63, hCoV-HKU-1, hCoV-OC43, hCoV-229E, coronavirus, feline infectious peritonitis virus, and transmissible gastroenteritis virus;

wherein the inflammatory disease is optionally selected from the group consisting of asthma, autoimmune disease, chronic inflammation, chronic prostatitis, glomerulonephritis, hypersensitivity disease, inflammatory bowel disease, sepsis, vascular smooth muscle cell disease, aneurysms, pelvic inflammatory disease, reperfusion injury, rheumatoid arthritis, transplant rejection, and vasculitis;

wherein the cancer is optionally selected from the group consisting of small and non-small cell lung cancer, bladder cancer, hepatocellular cancer, pancreatic cancer, breast cancer, glioblastoma, colorectal cancer, squamous cell carcinoma, melanoma, and prostate cancer;


wherein the muscular disorder is optionally selected from the group consisting of myocardial reperfusion injury, muscular dystrophy, collagen VI myopathies, Post-cardiac arrest syndrome (PCAS), heart failure, atherosclerosis, and abdominal aortic aneurysm;

wherein the neurological disorder is optionally selected from the group consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple systems atrophy, multiple sclerosis, cerebral palsy, epilepsy, stroke, diabetic neuropathy, amyotrophic lateral sclerosis (Lou Gehrig's Disease), bipolar disorder, excitotoxic injury, hepatic encephalopathy, hypoglycemia, manganese toxicity, neuronal target deprivation, toxic fatty acids such as arachadonic acid, mechanical nerve injury, spinal cord injury, and cerebral injury; and

wherein the injury associated with loss of cellular calcium homeostasis is optionally selected from the group consisting of myocardial infarct, stroke, acute hepatotoxicity, cholestasis, and storage/reperfusion injury of transplant organs.


19. A process for the preparation of the compound of claim 1, comprising the steps of:

1) reacting cyclosporin-A (CsA) with a basic lithium alkylamide, in the presence of a suitable solvent, followed by reaction with a suitable electrophile to generate a compound of Formula 1:




Formula 1


2) reacting the compound of Formula 1 with AC_2O in the presence of a suitable solvent to form a compound of Formula 2A:

**Formula 2A**

3) reacting the compound of Formula 2A with an oxidant to form a compound of Formula 3A:

**Formula 3A**

4) reacting the compound of Formula 3A with an electrophile to form a compound of Formula 4A:

**Formula 4A**

5) optionally deacylating the compound of Formula 4A.

20. The process of claim 19 wherein the preparation comprises the addition of an excess of LiCl in the suitable solvent in step 1) to form predominantly the L-epimer of the compound, or the preparation is carried out in the absence of LiCl in the suitable solvent in step 1) to form predominantly the D-epimer of the compound, wherein the chiral center of the D-epimer or L-epimer of the compound is the carbon atom to which R23 is attached.


21. The process of claim 19 or 20 wherein the basic lithium alkylamide is lithium diisopropylamide.

22. The process of any one of claims 19 to 21, wherein the electrophile used in step 1) is selected from the group defined in the following table, to generate the corresponding R23 in the table:


Electrophile	R23
methyl iodide	$-\text{CH}_3$
ethyl iodide	$-\text{CH}_2\text{CH}_3$
allyl bromide	$-\text{CH}_2\text{CHCH}_2$
propargyl bromide	$-\text{CH}_2\text{CCH}$

23. A process for the preparation of the compound of claim 1, comprising the steps of:

reacting a compound of Formula 1A with acetic anhydride in the presence of a suitable solvent, to form a compound of Formula 2:




optionally followed by formation of an aldehyde of Formula 3:



optionally followed by a Wittig reaction to generate a compound of Formula L, wherein R23 is selected from the group consisting of

- v. $-\text{CH}_3$;
- vi. $-\text{CH}_2\text{CH}_3$;
- vii. $-\text{CH}_2\text{CHCH}_2$; and
- viii. $-\text{CH}_2\text{CCH}$.

24. A process for the preparation of the compound of claim 1, comprising the steps of: reacting a compound of Formula 5:



Formula 5

with a basic lithium alkylamide, in the presence of a suitable electrophile in an appropriate solvent to form the compound, wherein R23 is C1-C3 alkyl.

25. The process of claim 24 wherein the basic lithium alkylamide is lithium diisopropylamide.

26. A use of the compound of any one of claims 1 to 14 or the pharmaceutical composition of claim 15 for preparing a medicament for treating or preventing a cyclophilin-mediated disease or injury in a mammal.

27. The use of claim 26, wherein the cyclophilin-mediated disease or injury is selected from the group consisting of

- a. a viral infection;
- b. inflammatory disease;
- c. cancer;
- d. muscular disorder;
- e. neurological disorder; and
- f. injury associated with ischemia, reperfusion, loss of cellular calcium homeostasis, loss of ionic homeostasis, increase in free radical production, or toxins that induce mitochondrial dysfunction;

wherein the viral infection is optionally caused by a virus selected from the group consisting of Human Immunodeficiency virus, Hepatitis A, Hepatitis B,

Hepatitis C, Hepatitis D, Hepatitis E, SARS-CoV, hCoV-NL63, hCoV-HKU-1, hCoV-OC43, hCOV-229E, coronavirus, feline infectious peritonitis virus, and transmissible gastroenteritis virus;

wherein the inflammatory disease is optionally selected from the group consisting of asthma, autoimmune disease, chronic inflammation, chronic prostatitis, glomerulonephritis, hypersensitivity disease, inflammatory bowel disease, sepsis, vascular smooth muscle cell disease, aneurysms, pelvic inflammatory disease, reperfusion injury, rheumatoid arthritis, transplant rejection, and vasculitis;

wherein the cancer is optionally selected from the group consisting of small and non-small cell lung cancer, bladder cancer, hepatocellular cancer, pancreatic cancer, breast cancer, glioblastoma, colorectal cancer, squamous cell carcinoma, melanoma, and prostate cancer;

wherein the muscular disorder is optionally selected from the group consisting of myocardial reperfusion injury, muscular dystrophy, collagen VI myopathies, Post-cardiac arrest syndrome (PCAS), heart failure, atherosclerosis, and abdominal aortic aneurysm;

wherein the neurological disorder is optionally selected from the group consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple systems atrophy, multiple sclerosis, cerebral palsy, epilepsy, stroke, diabetic neuropathy, amyotrophic lateral sclerosis (Lou Gehrig's Disease), bipolar disorder, excitotoxic injury, hepatic encephalopathy, hypoglycemia, manganese toxicity, neuronal target deprivation, toxic fatty acids such as arachadonic acid, mechanical nerve injury, spinal cord injury, and cerebral injury; and

wherein the injury associated with loss of cellular calcium homeostasis is optionally selected from the group consisting of myocardial infarct, stroke, acute hepatotoxicity, cholestasis, and storage/reperfusion injury of transplant organs.

Figure 1A

Table 5

1-Mod Name	Modification:	1-Mod Structure		1 + 3-D-MeAla	
		Cyp A Inhibition Fold-potency vs CsA	Immunosuppression % vs CsA	Cyp A Inhibition Fold-potency vs CsA	Immunosuppression % vs CsA
404-58		0.16	< 1	1.3	< 1
420-171		2.3	-	8.7	< 1
420-19		3.7	< 1	9.5	3.4
416-08		3.7	< 1	14.3	3
404-98		2.2	-	6.5	15
					7.6

				8.3 (saturated)	9.2 (saturated)
404-154		0.98	2		
420-30-1		3.3	< 1	11.7	< 1
420-103		0.19	10	1.9 (saturated)	7.7 (saturated)
404-173		1.14	2.6	3.9	4
404-76		0.65	12.5	2.8	9.7
404-194		0.61	2	4.4	1.4

Figure 1B

1-Mod Name	Modification:	1-Mod Structure		1 + 3-D-EtSar	
		Cyp A Inhibition Fold-potency vs CsA	Immunosuppression % vs CsA	Cyp A Inhibition Fold-potency vs CsA	Immunosuppression % vs CsA
404-58		0.16	< 1	0.56	< 1
420-19		3.7	< 1	4.2	2.3
404-154		0.98	2	10.6 (saturated)	4.1 (saturated)
420-30-1		3.3	< 1	8.9	1.1
404-194		0.61	2	2.1	4.9

Figure 1C