

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 October 2005 (27.10.2005)

PCT

(10) International Publication Number
WO 2005/100381 A2

- (51) International Patent Classification⁷: **C07K 7/00**
- (21) International Application Number:
PCT/US2005/011873
- (22) International Filing Date: 8 April 2005 (08.04.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/560,453 8 April 2004 (08.04.2004) US
- (71) Applicant (for all designated States except US): **MIL-LENNIUM PHARMACEUTICALS, INC.** [US/US]; 40 Landsdowne Street, Cambridge, MA 02139 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HO, Guojie** [US/US]; 33 Grindstone Lane, Sudury, MA 01776 (US). **PAONE, Antoinette, D.** [US/US]; 9 Apollo Circle, Lexington, MA 02421 (US). **FORNI, Luciano**; Chemin du Foriest, B-1420 Braine-l'Alleud (BE). **DETOL-LENAERE, Catherine**; Chemin du Foriest, B-1420 Braine-l'Alleud (BE). **BONNETT, Brice**; Chemin du Foriest, B-1420 Braine-l'Alleud (BE).
- (74) Agents: **LUCCI, Joseph** et al.; Woodcock Washburn LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 2005/100381 A2

(54) Title: PROCESSES FOR PREPARING EPTIFIBATIDE

(57) Abstract: The present invention provides, inter alia, convergent processes for preparing eptifibatide that involve coupling a 2-6 eptifibatide fragment to an activated cysteinamide residue to form a 2-7 eptifibatide fragment, attaching a mercaptopropionic acid residue to the 2-7 eptifibatide fragment through disulfide bond formation, coupling the peptide intramolecularly, and removing the protecting group, to form eptifibatide. The invention further provides products produced by the described processes, novel compounds that can be used as synthetic intermediates for the preparation of eptifibatide, and novel compounds that are structurally similar to eptifibatide.

PROCESSES FOR PREPARING EPTIFIBATIDE

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 60/560,453, filed April 8, 2004, which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] In certain embodiments, the invention relates to novel processes for preparing eptifibatide, a drug used to treat cardiovascular disease. The invention also relates to compounds that can be used as synthetic intermediates for eptifibatide, to compounds that are structurally similar to eptifibatide, and to processes for purifying eptifibatide.

BACKGROUND OF THE INVENTION

[0003] Eptifibatide is a highly specific cyclic heptapeptide antagonist of the platelet glycoprotein IIb/IIIa. It is a short-acting parenteral antithrombotic agent that is used during percutaneous coronary interventions for the treatment of unstable angina and as an adjunct to thrombolytic agents for the treatment of acute myocardial infarction. See, for example, Phillips et al., *Journal of Biological Chemistry* (1993), 268(2), 1066-73; and Scarborough, *American Heart Journal* (1999), 138(6, Pt. 1), 1093-1104. Eptifibatide is also administered to patients

undergoing balloon angioplasty, a procedure for which over 1.0 million people in the U.S. are candidates annually.

[0004] Eptifibatide is believed to work by inhibiting platelet aggregation, specifically, by blocking the platelet receptor GP IIb-IIIa. The aggregation of platelets can obstruct blood supply to the heart, causing unstable angina and, possibly, myocardial infarction (heart attack). The effects of eptifibatide are specific to platelets, avoiding interference with other normal cardiovascular processes, and the effects can be reversed when eptifibatide use is discontinued.

[0005] Eptifibatide is marketed in the U.S. under the trademark INTEGRILIN®, and is used to treat patients with acute coronary syndrome (unstable angina and non-Q-wave MI), including patients who are to be managed medically and those undergoing percutaneous coronary intervention ("PCI"). Eptifibatide is also indicated for use at the time of percutaneous coronary interventions, including procedures involving intracoronary stenting.


[0006] Many reported synthetic approaches to eptifibatide have employed known techniques of solid-phase peptide synthesis as described, for example, in U.S. Patents 5,318,899; 5,686,570 and 5,747,447. A commercial-scale, liquid phase process was also reported at the 1999 IBC Conference on Peptide Technologies, "Peptisyntha's Method of Producing GMP Peptides on an Industrial Scale". The commercial process is a convergent synthesis involving the separate preparation of two fragments: Mpa-Har-Gly and Asp-Trp-Pro. The coupling of these two fragments provides six of the seven residues needed for eptifibatide. The last residue attached is an S-trityl-protected cysteinamide as described, for example, in U.S. Patent 5,506,362. After removal of the S-trityl protecting groups (on the cysteinamide and mercaptopropionyl residues), ring closure is then achieved by disulfide bond formation. Crude eptifibatide obtained by the commercial process has a reported purity of about 80%. Two column chromatography steps improve the purity to greater than 99%.

[0007] Liquid-phase synthesis has generally been viewed as more feasible than solid-phase synthesis for the large-scale manufacture of eptifibatide. However, solubility issues and the generation of complex reaction mixtures present challenges for large-scale liquid phase processes. Complex reaction mixtures, for example, make purification of the product more difficult. Ways exist to overcome these problems, such as the use of persilylated amino acids and phase transfer reagents, as described, for example, in U.S. Patent 4,954,616, and extensive chromatographic purification, but such means add to the cost of the overall process.

[0008] A need thus exists for alternative processes for the manufacture of eptifibatide.


SUMMARY OF THE INVENTION

[0009] The invention provides, *inter alia*, processes for preparing eptifibatide. Certain processes of the invention comprise providing a compound of formula II:



(II)

wherein Har is homoarginyl; Gly is glycyl; Asp is aspartyl; Trp is tryptophanyl; Pro is prolyl; Cys-NH₂ is cysteinamide; Mpa is mercaptopropionic acid; and P₂ is a carboxyl protecting group; coupling the Har and Mpa residues to form a compound of formula III:



(III)


; and

removing P₂ from the Asp residue of the compound of formula III to form eptifibatide.

[0010] The invention also provides processes in which an amino-terminal protected homoarginine residue is coupled with a glycine residue, thereby forming a 2-3 eptifibatide fragment of the formula:




Also provided are processes in which an aspartic acid residue having a protected carboxyl side chain is coupled to a tryptophanyl-prolyl dipeptide through the tryptophanyl residue of the dipeptide, thereby forming a protected 4-6 eptifibatide fragment of the formula:




- 4 -


After deprotection, the 2-3 eptifibatide fragment and the 4-6 eptifibatide fragment, in turn, can be coupled through attachment of the Gly residue of the 2-3 eptifibatide fragment to the Asp residue of the 4-6 eptifibatide fragment, thereby forming a 2-6 eptifibatide fragment of the formula:



wherein Har is homoarginyl; Gly is glycyl; Asp is aspartyl; Trp is tryptophanyl; Pro is prolyl; P_1 is an amino protecting group; and P_2 is a carboxyl protecting group. In preferred embodiments, the 2-6 eptifibatide fragment is coupled to an activated cysteinamide residue through the Pro residue of the 2-6 eptifibatide fragment, thereby forming a 2-7 eptifibatide fragment of the formula:




wherein ACys-NH₂ is an activated cysteinamide residue. A mercaptopropionic acid residue can be attached to the 2-7 eptifibatide fragment through a disulfide linkage between the mercaptopropionic acid residue and the ACys-NH₂ residue of the 2-7 eptifibatide fragment, thereby forming a compound of formula I:



(I)

wherein Mpa is mercaptopropionic acid; Cys-NH₂ is cysteinamide. Removal of P_1 from the Har residue of the compound of formula I provides a compound of formula II:



(II)

and less than 1 % of certain process impurities. The invention also provides processes for purifying eptifibatide.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0012] In one aspect, the present invention provides convergent processes for preparing eptifibatide that involve preparation of a 2-3 eptifibatide fragment, preparation of a 4-6 eptifibatide fragment, and coupling of the 2-3 and 4-6 eptifibatide fragments to form a 2-6 eptifibatide fragment. Certain of these processes involve coupling the 2-6 eptifibatide fragment to an activated cysteinamide residue to form a 2-7 eptifibatide fragment, forming a disulfide bond between mercaptopropionic acid and the 2-7 eptifibatide fragment to form Precursor A, effecting intramolecular peptide coupling of Precursor A, and removing protecting groups from the coupling product to form eptifibatide.

[0013] In further embodiments, the invention relates to products produced by the described processes for preparing eptifibatide. In other embodiments, the invention relates to novel compounds that can be used as intermediates for the preparation of eptifibatide. In further embodiments, the invention relates to compounds that are structurally similar to eptifibatide. The invention also provides, in additional embodiments, processes for purifying eptifibatide.

[0014] As used herein, the term "carboxyl protecting group" refers to a moiety that can be selectively attached to and removed from a carboxyl group to prevent it from participating in undesired chemical reactions, without unacceptably adverse affects on desired reactions. Examples of carboxyl protecting groups include esters, such as methyl, ethyl, t-butyl, (un)substituted benzyl, and silyl esters, among others. Other carboxyl protecting groups are well known in the art and are described in detail in *Protecting Groups in Organic Synthesis*, Theodora W. Greene and Peter G. M. Wuts, 3rd Edition, 1999, published by John Wiley and Sons, Inc..

[0015] As used herein, the term "amino protecting group" refers to a moiety that can be selectively attached to and removed from a nitrogen atom to prevent it from participating in undesired chemical reactions, without unacceptably adverse affects on desired reactions. Examples of amino protecting groups include carbamates, such as Boc, Cbz, Fmoc, alloc, methyl and ethyl carbamates, among others; cyclic imide derivatives, such as phthalimide; amides, such as formyl, (un)substituted acetyl, and benzoyl; and trialkyl silyl groups, such as t-butyl dimethylsilyl and triisopropylsilyl. Other amino protecting groups are well known in the art

and are described in detail in *Protecting Groups in Organic Synthesis*, Theodora W. Greene and Peter G. M. Wuts, 3rd Edition, 1999, published by John Wiley and Son, Inc..

[0016] As used herein, the term “coupling,” and all variations thereof, refers to the formation of an amide bond, by any means, between the moieties being joined.


[0017] As used herein, the term “attaching,” and all variations thereof, refers to the formation of an amide or disulfide bond, by any means that can be used for form an amide or disulfide bond, between the moieties being joined.

[0018] As used herein, the term “activated cysteinamide residue” refers to a cysteinamide residue that is capable of forming a disulfide bond with mercaptopropionic acid.

[0019] As used herein, the term “Gly-eptifibatide” refers to a compound that is structurally related to eptifibatide but contains two adjacent glycine residues rather than a single glycine residue.


[0020] All amino acid residues referred to herein are natural amino acids that have the L-configuration.

[0021] Eptifibatide has the following chemical structure:




[0022] Eptifibatide can also be represented using amino acid designations as follows:

- 8 -



where the amino acid designations correspond to the chemical drawing shown below:



[0023] For the ease of description, the residues can also be numbered from (1) through (7). Residue (1) is mercaptopropionic acid; (2) is homoarginyl (Har); (3) is glycyl (Gly); (4) is aspartyl (Asp); (5) is tryptophanyl (Trp); (6) is prolyl (Pro); and (7) is cysteinamide (Cys-NH₂).

[0024] In certain embodiments, the present invention relates to convergent processes for preparing eptifibatide. In a first sequence of steps, a 2-3 eptifibatide fragment is prepared containing amino acids (2) and (3). In a second sequence, a 4-6 eptifibatide fragment is prepared containing amino acids (4), (5) and (6). The two fragments are then coupled to provide a single 2-6 fragment, which is a pentapeptide. The 2-6 fragment of eptifibatide is protected at the amino-terminal of the Har residue and at the aspartyl side chain carboxyl group. Scheme 1 outlines an exemplary process for the preparation of the 2-6 eptifibatide fragment:


Cbz (benzyloxycarbonyl), RCbz (benzyloxycarbonyl groups substituted on the aromatic ring), 9(2-sulfo)fluorenylmethylcarbamate, 9(2,7-dibromo)fluorenylmethylcarbamate, 2-chloro-3-indenylmethylcarbamate, Benz[f]inden-3-ylmethylcarbamate, or Alloc (Allyloxycarbonyl) groups can be used.

[0026] The t-butyl ester on the aspartyl residue can be replaced by any other protecting group that can be cleaved by acid treatment such as, for example, ODpm (diphenylmethyl ester) or protecting groups that can be cleaved by hydrogenolysis such as, for example, OBzl (benzyl ester).

[0027] The Fmoc-Har moiety in the eptifibatide fragments depicted in Sequence A of Scheme 1 can be replaced by R_{NP1} -Har, where R_{NP1} is an amino protecting group such as, for example, a Cbz (benzyloxycarbonyl) group, a RCbz (benzyloxycarbonyl substituted on the aromatic ring) group, or an Alloc (Allyloxycarbonyl) group. Similarly, the Z-Asp (or Cbz-Asp) moiety in the eptifibatide fragments depicted in Sequence B can be replaced by R_{NP2} -Asp, where R_{NP2} is an amino protecting group that can be cleaved in basic conditions such as, for example, Fmoc (fluorenylmethyloxycarbonyl), or where R_{NP2} is an amino protecting group that can be cleaved by hydrogenolysis such as, for example, a RCbz (benzyloxycarbonyl protecting group with substituted aromatic ring) group. In addition, the -OtBu group depicted in Scheme 1 can be replaced by R_{CP} , where R_{CP} is a carboxyl protecting that can be cleaved by acid treatment, such as, for example, an ODpm (diphenylmethyl ester) group. Pfp (pentafluorophenyl) can be replaced by R_{L1} , where R_{L1} is a carboxyl activating group; and Su (succinimide) can be replaced by R_{L2} , where R_{L2} is a carboxyl activating group. Independently of the nature of the fragment, both Pfp and Su can be replaced by other stable active esters such as, for example, mono and dinitrophenyl esters, tri- and pentaphenyl ester, It will be appreciated by those skilled in the art that the selection of particular protecting groups depends upon the identity of other protecting groups that exist on the same compound. In certain embodiments of the invention, a particular protecting group is chosen so that it can be removed selectively under reaction conditions that do not affect other protecting groups.

[0028] In certain embodiments of the invention, the protected 2-6 eptifibatide fragment is next coupled to an "activated" cysteinamide (7), such as, for example, 3-nitro-2-pyridinesulfonyl-cysteinamide (H-Cys(Npys)-NH₂); Nps (2-nitro-phenyl-sulfonyl); S-phenylthiocysteinamide, where the phenyl ring is substituted; S-alkylthiocysteinamide; or the S-sulfonate and S-sulfonylthiocarbonates of cysteinamide, to form the 2-7 hexapeptide fragment of


eptifibatide. The remaining eptifibatide residue is mercaptopropionic acid or Mpa-OH (1). Mpa-OH is attached to the 2-7 hexapeptide fragment under conditions that form a disulfide linkage between 1 and the 2-7 fragment. The resulting disulfide piece, designated herein as Precursor A, is a key and novel precursor for making eptifibatide. The structure of Precursor A is shown below:



where P_1 is an amino protecting group and P_2 is a carboxyl protecting group. Preferably the P_2 group is stable under conditions that are suitable for removal of the P_1 group. The selection of compatible P_1 and P_2 groups that allow for the selective removal of only one of the groups is well-known in the art. An example of a compatible pair of protecting groups is $\text{P}_1 = \text{Fmoc}$, which may be removed under basic conditions, and $\text{P}_2 = \text{t-butyl}$, which is stable under the same conditions.

[0029] Scheme 2 outlines an exemplary process for the preparation of Precursor A:


Scheme 2. Preparation of Precursor A from 2-6




- 12 -

[0030] As in Scheme 1, the Fmoc and the *t*-butyl groups shown in Scheme 2 can be replaced by other known protecting groups that are generally recognized to be useful for peptide synthesis.

[0031] In certain embodiments of the invention, removal of the P₁ protecting group from Precursor A provides another precursor, Precursor B, which is the 2-7-1 portion of eptifibatide:




[0032] Precursor B can be converted, via an intramolecular peptide coupling, to Precursor C:



The intramolecular peptide coupling can be effected, for example, in an organic solvent in the presence of a suitable coupling reagent such as, for example, a uronium-type coupling agent such as, but not limited to, *O*-[cyano(ethoxycarbonyl)methylenamino]-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TOTU), HBTU, or TBTU (2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate and tetrafluoroborate respectively); a carbodiimide type reagent such as, but not limited to, DCC (dicyclohexylcarbodiimide), DIC (diisopropylcarbodiimide), or EDC (1-Ethyl-3-(3-dimethylaminopropyl)-carbodiimide); active esters; or phosphonium type coupling reagents such as, for example, Bop (benzotriazol-1-yloxy-tri(dimethylamino)-phosphonium hexafluorophosphate) or PyBOP (benzotriazol-1-yloxy-tri(pyrrolidino)-phosphonium hexafluorophosphate). Peptide coupling is described, for example, in Humphrey and Chamberlin, *Chem. Rev.* **1997**, *97*, 2243-2266, incorporated herein by reference in its entirety.

[0033] Removal of the P₂ protecting group from Precursor C provides eptifibatide. The removal of P₂ can be effected, for example, in an organic solvent in the presence of acid, base or any other reagent or reagent system in which the protecting group is labile. Scheme 3 outlines an exemplary process for the preparation of eptifibatide from Precursor A:

Scheme 3. Preparation of Eptifibatide from Precursor A

[0034] After removal of the Fmoc protecting group, the 2-7-1 fragment undergoes an intramolecular peptide coupling to provide the *t*-butyl ester of eptifibatide. Removal of the *t*-butyl group from the aspartyl residue yields eptifibatide.

[0035] In certain embodiments, the present invention also relates to compounds that are structurally similar to eptifibatide and are prepared according to processes that are similar to the processes described above for preparing eptifibatide. Such compounds include, for example,

concentrations of 50 % of a 0.5 % aqueous solution of acetic acid and 50 % of an acetonitrile solution. In some embodiments, the stationary phase is washed with the acetic acid/acetonitrile gradient following one or more washes with a trifluoroacetic acid/acetonitrile gradient.

[0040] The invention also provides embodiments in which the ammonium acid/acetonitrile solution is a solution comprising 95 % of an aqueous 100 mM ammonium acid solution and 5 % of an acetonitrile solution. In certain aspects of the invention, the stationary phase is washed with an ammonium acid/acetonitrile solution following washing with a trifluoroacetic acid/acetonitrile gradient and/or an acetic acid/acetonitrile gradient.

[0041] The coupling of the amino acid residues that occurs in the described processes can be accomplished through known methods of peptide synthesis that are familiar to those skilled in the art. Any suitable coupling procedure for forming peptides can be employed.

[0042] The following examples are illustrative of certain embodiments of the invention and should not be considered to limit the scope of the invention.

Example 1: Preparation of Z-Asp(OtBu)-Trp-Pro-OH

[0043] Z-Asp(OtBu)-Trp-Pro-OH was obtained by known procedures starting with the commercially available Z-Asp(OtBu)-OSu and H-Trp-Pro-OH as described, for example, in Bodanszky, M. (1979), Active esters in peptide synthesis, *The peptides*, Vol. 1 (ed. E. Gross and J. Meienhofer), Chapter 3. Academic Press, London, which is incorporated herein by reference in its entirety.

Example 2: Preparation of H-Asp(O-tBu)-Trp-Pro-OH

[0044] To a 2 liter reactor charged with dimethylacetamide (DMAC, 1.2 L) at a temperature of about 20 °C was added 0.300 kg of starting material Z-Asp(OtBu)-Trp-Pro-OH, maintaining the temperature at about 20 °C. The Z-Asp(OtBu)-Trp-Pro-OH was allowed to dissolve and the reaction mixture was then purged and blanketed by a nitrogen atmosphere. Palladium on carbon (5 weight%) (0.015 kg) was added to the reaction mixture, followed by hydrogenation at a 2 bar pressure while the reaction mixture was maintained at about 20 °C.

[0045] After two hours, and every hour thereafter, a sample from the reaction was analyzed by HPLC. HPLC analysis was conducted on Purospher Star C18 55*4 mm column with a water/acetonitrile solvent mixture containing 0.1% trifluoroacetic acid (TFA) with a

gradient of 98:2 water/acetonitrile to 2:98 water/acetonitrile in 10 minutes. HPLC detection was at 215 nm, flow was 2.0 mL/min, and the temperature was 40°C.

[0046] The reaction was considered to be complete when the HPLC analysis showed the starting material to be less than or equal to 0.2% relative to the product area percent. When the reaction was shown to be complete by HPLC, an aqueous solution of p-toluenesulfonic acid (0.094 kg p-TsOH in 0.150 L of water) was added to the reaction mixture. The reaction mixture was then filtered through Celite® that was pre-washed with DMAC (Celite® washed three times with 0.6 L). After filtration of reaction mixture, the filter cake was washed three times with fresh DMAC (0.3 L). HPLC analysis after the last wash was performed to confirm that no product remained in the filter cake.

[0047] The combined filtrates were transferred to a new vessel at about 20 °C, at which temperature N-ethylmorpholine (0.066 L) was added. The temperature was maintained below about 22 °C during the addition. The mixture was then cooled to about 8-12 °C and water (3 L) was added slowly so as to maintain the temperature below about 15 °C. The mixture was then stirred at about 8-12 °C for about 30 minutes and held at that temperature for 8 hours. During this time the product crystallized out of solution. The supernatant can be analyzed by HPLC to determine the amount of product still in solution and whether further cooling is necessary. The slurry obtained was filtered and the collected solids were washed twice with a 2:1 solution of water:DMAC (0.9 L each). The solids were then washed twice with acetonitrile (0.9 L each) and dried under a vacuum below about 25 °C until at a constant weight. The water content after drying was 3.5% (Karl-Fisher analysis).

[0048] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0049] A LC/MS analysis confirmed the mass [M+H] 473.0 of H-Asp(O-tBu)-Trp-Pro-OH.

[0050] The recovery yield was 93.2 % (0.218 kg); the net yield was 93.1% (based on nitrogen content : 95.4%).

Example 3: Preparation of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-OH

[0051] To a slurry of Fmoc-Har-Gly-OH (0.163 kg net ; based on peptide content : 90.5%) in THF:DMAC (0.717 L and 0.179 L respectively) was added methanesulfonic acid (0.027 L) and pentafluorophenol (0.083 kg). The mixture was stirred at about 20 °C until the

- 17 -

solid dissolved. N-ethylmorpholine (NEM) (0.030 L) was added drop wise, followed by dicyclohexylcarbodiimide (DCC) (0.072 kg) in THF (0.179 L) and the mixture was stirred at about 20 °C. The formation of Fmoc-Har-Gly-OPfp was followed by HPLC assay. After about 17 hours, the ratio of Fmoc-Har-Gly-OH / Fmoc-Har-Gly-OPfp was 1.5/98.5.

[0052] To the reaction mixture was added H-Asp(O-tBu)-Trp-Pro-OH (0.146 kg, about 3.5 w/w% H₂O) and NEM (0.049 L). The mixture was stirred at about 20 °C for 3 hours. At that time, HPLC analysis showed the mixture to contain about 87% of product, <2% Fmoc-Har-Gly-OPfp, 6% of Fmoc-Har-Gly-OH, about 0.5% of H-Asp(O-tBu)-Trp-Pro-OH, and about 5.5% of an impurity that was identified as Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Asp(O-tBu)-Trp-Pro-OH ("d.a. impurity"). HPLC analysis was conducted on a Purospher Star C18 55*4 mm column with a water/acetonitrile solvent mixture containing 0.1% trifluoroacetic acid (TFA) with a gradient of 98:2 water/acetonitrile to 2:98 water:acetonitrile in 10 minutes. HPLC detection was at 215 nm, flow was 2.0 mL/min and temperature was 40°C.

[0053] The reaction mixture was filtered and the solid was washed twice with a 5/1 mixture of THF/DMAC (2 x 0.489 L). The filtrate was concentrated to about 0.815 L at no more than 35 °C under reduced pressure (about 20 mBar). The concentrated mixture was added slowly over 1 h to 1/4 mixture of acetonitrile/H₂O (4.1 L) containing sodium bicarbonate (0.059 kg). The addition rate was carefully controlled to avoid formation of a gummy precipitate. About 3.5% of the solution was added over about 15 min, and the addition was interrupted to stir the slurry at about 20°C for 1 h. The pasty precipitate turned to white slurry. 7.5 extra % of the solution was added over 15 to 30 min, and the addition was interrupted again to stir the slurry at about 20°C for 2 h. The remaining 89% of solution was added over about 2 h. The mixture was stirred at about 20 °C for 12 h then filtered. The solid was washed with a 1/4 mixture of acetonitrile/H₂O (3 x 0.7 L), a 2/3 mixture of acetonitrile/di-isopropylether (DIPE) (3 x 0.7 L) and DIPE (2 x 0.7 L). The solid product was dried at T ≤ 30°C until a constant weight was achieved. The water content after drying was 3.3% (Karl-Fisher analysis).

[0054] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0055] A LC/MS analysis confirmed the mass [M+H] 922.5 of Fmoc-Har-Gly-Asp(OtBu)-Trp-Pro-OH.

[0056] The recovery yield was 81.8% (0.234 kg); the net yield was 82.3% (based on nitrogen content : 96.0%).

Example 4: Preparation of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NPys)-NH₂

[0057] In a reactor charged with peptide synthesis grade DMF (0.675 L) at about 20 °C was added Fmoc-Har-Gly-Asp(OtBu)-Trp-Pro-OH (0.225 kg). The mixture was cooled to about 0°C, and H-Cys(NPys)-NH₂ was added as the solid hydrochloride salt (0.080 kg). *O*-Benzotriazol-1-yl-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TBTU) (0.082 kg) was added to the reaction mixture. The pH of the reaction mixture was adjusted to 6.5 to 7.0 by the portion-wise addition of diisopropylethylamine (DIPEA)(0.112 L), while maintaining the temperature at about 0°C. The mixture was stirred at that temperature, during which time samples were analyzed by HPLC for product formation about every 45 minutes. The pH of the reaction mixture was maintained at pH 6.5 to 7.0 with the addition of DIPEA as needed. HPLC analysis was conducted on a Platinum EPS 100-5 C18 5μ 250*4.6 mm column ; Solvent A: TFA 0.1% in water; Solvent B: TFA 0.1% in acetonitrile; gradient: 12 to 98% B in 15 minutes. HPLC detection was at 215 nm, flow 2.0 mL/min and the temperature was 40°C. The reaction was considered complete when the pentapeptide and Cys(NPys)-NH₂ starting materials were each shown to be less than 1% by HPLC. Otherwise, additional TBTU, DIPEA and the starting material shown to be deficient in the mixture were added.

[0058] The reaction mixture was added to another vessel that contained water (4.5 L) at a temperature of about 5°C. This temperature was maintained during the addition with stirring. After stirring for 10 minutes at about 5°C, the mixture was filtered and the solid was washed five times with water (0.9 L each). The solid was then washed three times with toluene (0.675 L each). The washes with toluene can be replaced by washes with diisopropyl ether. The solid was dried under vacuum at 35°C or below until a constant weight was achieved. The water content after drying was 1.2% (Karl-Fisher analysis).

[0059] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0060] A LC/MS analysis confirmed the mass [M+H] 1178.4 of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NPys)NH₂.

[0061] The recovery yield was 102.4% (0.295 kg) and the net yield 87.7% (based on peptide content : 82.0%).

- 19 -

Example 5: Preparation of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NH₂)-Mpa (Fmoc[2-7-1])

[0062] In a reactor charged with HPLC grade acetonitrile (0.570 L) and peptide synthesis grade DMF (0.285 L), under a nitrogen atmosphere at about 20°C, was added 0.285 kg of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NPys)NH₂ (Fmoc[2-7]) slowly with agitation. After dissolution of the Fmoc[2-7], the mixture was cooled to about -3°C. A solution of mercaptopropionic acid (Mpa, 0.023 kg) in acetonitrile (0.057 L), prepared at about 20°C, was added to the reaction mixture at such a rate as to maintain the reaction temperature at about -3°C. The reaction was monitored by HPLC analysis and considered completed when the analysis showed less than 1% of Fmoc[2-7] compared to Fmoc[2-7-1]. HPLC analysis was conducted on a Purospher Star C18 55*4 mm column with a water/acetonitrile solvent mixture containing 0.1% trifluoroacetic acid (TFA) with a gradient of 98:2 water/acetonitrile to 2:98 water:acetonitrile in 10 minutes. HPLC detection was at 215 nm, flow was 2.0 mL/min and the temperature was 40°C.

[0063] The reaction mixture was added to a second vessel that was charged with HPLC grade acetonitrile (5.7 L) and N-ethylmorpholine (NEM, 0.033 L) at about 20°C. After the addition was complete, the reaction was stirred at that temperature for about 30 minutes. The slurry was slowly cooled to about 0°C and stirring was continued at that temperature for about 45 minutes. The following procedure was then performed three times: (a) the agitation was stopped to allow for the precipitate and the supernatant to separate; (b) the supernatant was pumped out of the vessel (c) at about 0° C, fresh HPLC grade acetonitrile (1.425 L) was added to the reactor and agitation was re-started; and (d) the slurry was stirred for about 5 minutes at about 0° C. The remaining slurry was filtered, and the solid was washed twice with HPLC grade acetonitrile (1.140 L each) and once with toluene (1.425 L). The wash with toluene can be replaced by a wash with diisopropyl ether. The solid was dried under high vacuum at no more than 20°C.

[0064] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0065] A LC/MS analysis confirmed the mass [M+H] 1128.4 of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NH₂)-Mpa.

[0066] The recovery yield was 93.8% (0.256 kg); the net yield was quantitative (based on peptide content : 87.4%).

- 20 -

Example 6: Preparation of H-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NH₂)-Mpa ([2-7-1])

[0067] To a reactor charged with peptide synthesis grade DMF (0.750 L) at about 20°C, was added 0.250 kg of Fmoc-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NH₂)-Mpa (Fmoc[2-7-1]) slowly with agitation. The mixture was cooled to about 10°C, at which temperature diethylamine (0.034 L) was added. After addition of the diethylamine, the temperature of the mixture was allowed to rise to about 20°C. Progress of the reaction was monitored by HPLC analysis of samples taken every hour. HPLC analysis was conducted on a Platinum EPS 100-5 C18 5 μ 250*4.6 mm column ; Solvent A: TFA 0.1% in water; Solvent B: TFA 0.1% in acetonitrile; gradient: 22 to 98% B in 15 minutes. HPLC detection was at 215 nm, flow was 2.0 mL/min and the temperature was 40°C. The reaction was considered complete when the percent of Fmoc[2-7-1] was less than about 0.5% with respect to [2-7-1]. The reaction was usually complete in 3 hours.

[0068] The reaction mixture was added to a second vessel charged with ethyl acetate (5.0 L) and cooled to about 10°C. The resulting suspension was stirred for 10 minutes at about 10°C. The following procedure was then performed twice: (a) the agitation was stopped and the precipitate was allowed to separate from the supernatant for about 15 minutes; (b) the supernatant was pumped out of the vessel; (c) fresh ethyl acetate (0.750 L) was added at about 10°C; (c) the suspension was stirred at about 10°C for about 5 minutes. The slurry was filtered and the solid was washed 6 times with ethyl acetate (0.750 L each time). Additional washes with diisopropyl ether can be performed to remove the residues of ethyl acetate. The solid was dried under high vacuum at no more than 25°C for a maximum of 18 hours.

[0069] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0070] A LC/MS analysis confirmed the mass [M+H] 906.3 of H-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NH₂)-Mpa.

[0071] A GC analysis showed 3.4% residual diethylamine.

[0072] The recovery yield was 106.1% (0.213 kg) and the net yield 99.2% (based on peptide content : 81.9%).

Example 7: Preparation of [MPA-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys](NH₂) (cyclo-[1-7](OtBu)-NH₂)

[0073] To the reactor was charged peptide synthesis grade DMF (0.768 L) and the solution was cooled to about 10°C. *O*-[cyano(ethoxycarbonyl)methylenamino]-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TOTU) (0.070 kg) was added followed by dilution with peptide synthesis grade dichloromethane (1.536 L) while maintaining the temperature below 15°C. The resulting solution was cooled to about -6°C and NEM (0.024 L) was added in small portions maintaining the temperature at about -6°C. The pH was measured before and after the addition of NEM. A sample was taken for HPLC analysis.

[0074] HPLC analysis was on a Phenomenex Luna C18(2) column, 5µm, 150*4.6mm, gradient 12 to 98% B in 15 minutes. Solvent A: TFA 0.1% in water, Solvent B: TFA 0.1% in acetonitrile. Detection was at 215 nm, the flow was 2 mL/min and the temperature was 40°C.

[0075] To a separate reactor was charged peptide synthesis grade DMF (0.768 L) at about 20°C followed by 0.192 kg of H-Har-Gly-Asp(O-tBu)-Trp-Pro-Cys(NH₂)-Mpa ([2-7-1]). When the solid dissolved, the resulting solution was cooled to about 0°C. HOBt (0.026 kg) was added in small portions (weight of HOBt was corrected for purity). The resulting solution was diluted with peptide synthesis grade dichloromethane (0.384 L). A sample was taken for HPLC analysis.

[0076] The solution of [2-7-1] was added into the solution of TOTU very slowly with an addition pump over a minimum of 3 hours while the temperature was maintained at about -6°C. A sample was taken after the addition of 50% of the [2-7-1] solution for HPLC analysis. The pH was also measured at this point. HPLC and pH samples were also taken when [2-7-1] addition was complete. The pH was adjusted to 7-7.5 by the addition of NEM if needed, while the temperature was maintained at about -6°C.

[0077] The pH was checked every 15 minutes and was adjusted to 7-7.5 as necessary with NEM while the temperature was maintained at about -6°C. A sample was taken for HPLC every 45 minutes until the cyclization was complete. The reaction was considered complete when the area percent of [2-7-1] was <0.5% relative to cyclized product.

[0078] If the reaction stalled, 1.1 equivalents of TOTU per residual amount of [2-7-1] were added and the pH was adjusted to 7-7.5 as needed while the temperature was maintained at -6°C.

- 22 -

[0079] When the reaction was complete, the mixture was concentrated under a vacuum at <40°C to a final volume of about 0.8 L. The resulting viscous solution was cooled to about 5°C and added slowly to rapidly stirring ethyl acetate that was at about 0°C (7.0 L). The resulting slurry was stirred at about 0°C for 20 minutes. The following procedure was then performed three times: (a) the agitation was stopped to allow for the precipitate and supernatant to separate; (b) the supernatant was pumped out of the vessel; (c) at about 0° C, fresh ethyl acetate (1.15 L) was added to the reactor and agitation was re-started; (d) the slurry was stirred for about 20 minutes at about 0°C. The final time the above procedure was performed, a smaller volume of ethyl acetate (0.768 L) was added to the reaction and agitation was re-started for 10 minutes at 0°C. The resulting slurry was filtered and the solid was washed once with ethyl acetate (0.576 L) and three times with di-isopropyl ether (0.576 L). The solid was dried under a high vacuum at about 25°C.

[0080] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0081] A LC/MS analysis confirmed the mass [M+H] 888.2 of(cyclo-[1-7](OtBu)-NH₂).

[0082] The recovery yield was 103.7% (0.194 kg); the net yield was quantitative (based on peptide content : 79.7%).

Example 8: Preparation of [MPA-Har-Gly-Asp-Trp-Pro-Cys](NH₂) (cyclo-[1-7]-NH₂)

[0083] A mixture of dichloromethane (0.573 L), anisol (0.086 L) and TFA (0.122 L) was prepared at about 20°C. The resulting solution was cooled to 15°C and solid [1-7](OtBu)-NH₂ (0.191 kg) was added slowly. The temperature was held below 20°C during the addition of [1-7](OtBu)-NH₂. The reaction mixture was further cooled to 10°C and an extra portion of TFA (0.365 L) was added in about 10 minutes. The temperature was held below 20°C during the addition. After the addition of all the TFA, the reaction was allowed to stir at 20°C and followed by HPLC.

[0084] HPLC analysis was conducted on a Platinum EPS 100-5 C18 column, 250*4.6mm. The gradient was 22 to 98% B in 15 minutes where solvent A was 0.1% TFA in water and solvent B was 0.1% TFA in acetonitrile. Detection was at 215 nm, the flow was 1.5 mL/minute, and the temperature was 40°C. The reaction was complete when < 3.0 Area% [1-7](OtBu)-NH₂ remained relative to the product.

- 23 -

[0085] When the reaction was complete the mixture was cooled to 10°C and added to a solution of 2/1 di-isopropyl ether (DIPE)/acetonitrile (3.78 L) at 10°C. The resulting slurry was stirred for 10 minutes at 10°C and then filtered under vacuum. The solid was washed three times with a 7/3 mixture of DIPE and acetonitrile (0.573 L) and three times with DIPE (0.573 L). The product was dried at no more than 25°C.

[0086] The product was analyzed by reverse phase HPLC using an aqueous trifluoroacetic acid/acetonitrile gradient.

[0087] A LC/MS analysis confirmed the mass [M+H] 832.3 of (cyclo-[1-7]-NH₂).

[0088] The recovery yield was 0.157 kg (87.9%) and the net yield was 87.2% (based on peptide content : 79.1%).

[0089] The content in pure API was 45.8%. This value was obtained by the following HPLC method: Synergi Max-RP 4 μm 80Å 250*4.6 mm; solvent A: 52 mM H₃PO₄/CH₃CN/100mM H₇SA (86/14/0.80); Solvent B: 52 mM H₃PO₄/CH₃CN /100mM H₇SA (50/50/0.80); 220 nm; 1.3 ml/min; 50°C; gradient: 0%B over 45 minutes, then to 45% B over 13 minutes, then to 100% B over 1 minute.

Example 9: Purification of Eptifibatide

Primary-purification:

[0090] The primary purification was a trifluoroacetic acid/acetonitrile-based purification. The norms applied for the individual main fractions were ≥ 92.0 %. The stationary phase was a Kromasil C18, 10 μm, 100 A column with a 5 cm diameter. The column pressure was 50 bars, the flow rate was 50 ml/min, and the detection wavelength was 215 nm. The mobile phases were as follows:

Solvent A: TFA 0.1 % in processed water / CH₃CN (95/5); and

Solvent B: TFA 0.1 % in processed water / CH₃CN (50/50).

The column was equilibrated by elution of 100 % solvent A over 15 minutes. The purification gradient was as follows: elution of 100 % solvent A over 10 minutes; gradient: 15 % B to 45 % B over 60 minutes; and elution of 100 % solvent B over 15 minutes.

[0091] The purification was monitored using method MAD-009-SF323TG1 and the target acceptance criteria were as follows:

Main fraction (F1): ≥ 92 %; and

Side fraction (Fp): ≥ 60 % and < 92 %.

The collected fractions were stored at 2°C to 8°C.

Secondary purification:

[0092] The secondary purification was an acetic acid/acetonitrile based purification. The norms for the individual main fractions were $\geq 99.0\%$ with single impurities $< 0.3 / 0.5\%$. The stationary phase was a Kromasil C18, 10 μm , 100 A column with a diameter of 5 cm. The column pressure was 40 ± 5 bars, the flow rate was 50 ml/min, and detection was at a wavelength of 215 nm. The mobile phases were as follows:

Solvent A: AcOH 0.5 % in processed water / CH₃CN (95/5); and

Solvent B: AcOH 0.5 % in processed water / CH₃CN (50/50).

[0093] The column was equilibrated by elution of 100 % solvent A over 15 minutes. The purification gradient was as follows: elution of 100 % solvent A over 10 minutes; gradient: 0 % B to 15 % B over 5 minutes; 15 % B to 35 % B over 60 minutes; and elution of 100 % solvent B over 15 minutes.

[0094] The purification was monitored using method MAD-009-SF323TG1 and the target acceptance criteria were as follows:

Main fraction (F1): $\geq 99.0\%$ with impurities $< 0.3 / 0.5\%$; and

Side fraction (Fp): $> 80.0\%$ and $< 99.0\%$.

The collected fractions were stored at 2°C to 8°C.

Desalting/concentration:

[0095] A double desalting step in ammonium acetate solution was performed before the concentration step to remove the residual trifluoroacetate counter ion and to obtain the acetate form of the peptide. The concentration step reduced the volume of the processed solution.

[0096] The stationary phase for the NH₄OAc desalting/ AcOH concentration was a Kromasil C18, 10 μm , 100 A column with a diameter of 5 cm. The column pressure was 40 ± 5 bars, the flow rate was 50 ml/min, and the detection wavelength was 215 nm. The mobile phases were as follows:

Solvent A: AcOH 0.5 % in processed water / CH₃CN (95/5);

Solvent B: AcOH 0.5 % in processed water / CH₃CN (50/50); and

Solvent C: NH₄OAc 100 mM pH: 6.5 in processed water / CH₃CN (95/5).

- 25 -

[0097] The column was equilibrated by elution of 100 % solvent A over 15 minutes. For column loading, the main fraction from the AcOH purification was diluted with 2 volumes of processed water.

[0098] The desalting was performed as follows: 10 minutes of solvent A; 10 minutes of solvent C; 10 minutes of solvent A; and 10 minutes of solvent C.

[0099] Concentration was performed as follows: 10 minutes of solvent A and 50 % B over 15 minutes.

[0100] The desalting and concentration were monitored using method MAD-009-SF323TG1 and the target acceptance criteria were as follows:

Main fraction (F1): ≥ 99.0 % with impurities $< 0.3 / 0.5$ %; and


Side fraction (Fp): > 80.0 % and < 99.0 %.

The collected fractions were stored at 2°C to 8°C.

[0101] The entire disclosure of each patent, patent application, and publication cited or described in this document is hereby incorporated herein by reference.

What is Claimed:

1. A process comprising:
 - providing a compound of formula II:



(II)

wherein

Har is homoarginyl;

Gly is glycyl;

Asp is aspartyl;

Trp is tryptophanyl;


Pro is prolyl;

Cys-NH₂ is cysteinamide;

Mpa is mercaptopropionic acid; and

P₂ is a carboxyl protecting group; and


- coupling the Har and Mpa residues to form a compound of formula III:



(III)

2. The process of claim 1 wherein P₂ is t-butyl.
3. The process of claim 1 further comprising removing P₂ from the Asp residue of the compound of formula III to form eptifibatide.
4. The process of claim 1 further comprising removing P₁ from the Har residue of a compound of formula I:


- 27 -



(I)


wherein P₁ is an amino protecting group, thereby forming the compound of formula II.

5. The process of claim 4 wherein P₂ is stable under conditions suitable for removal of P₁.
6. The process of claim 4 wherein P₂ is t-butyl.
7. The process of claim 4 wherein P₁ is 9-fluorenylmethoxycarbonyl or benzyloxycarbonyl.
8. The process of claim 4 further comprising attaching an Mpa residue to a 2-7 eptifibatide fragment of the formula:



wherein ACys-NH₂ is an activated cysteinamide residue, through a disulfide linkage between the Mpa residue and the ACys-NH₂ residue of the 2-7 eptifibatide fragment, thereby forming the compound of formula I.

9. The process of claim 8 further comprising coupling a 2-6 eptifibatide fragment of the formula:




to an activated cysteinamide residue through the Pro residue of the 2-6 eptifibatide fragment, thereby forming the 2-7 eptifibatide fragment.

10. The process of claim 9 wherein the activated cysteinamide residue is H-Cys(Npys)-NH₂.
11. The process of claim 9 further comprising coupling a 2-3 eptifibatide fragment of the formula:



and a 4-6 eptifibatide fragment of the formula:



through attachment of the Gly residue of the 2-3 eptifibatide fragment to the Asp residue of the 4-6 eptifibatide fragment, thereby forming the 2-6 eptifibatide fragment.

12. The process of claim 11 further comprising coupling an amino-terminal protected Asp residue having a protected carboxyl side chain to a Trp-Pro dipeptide through the Trp residue of the dipeptide and removing the amino-terminal protecting group from the Asp residue to form the 4-6 eptifibatide fragment.


13. The process of claim 12 further comprising coupling an amino-terminal protected Har residue and a protected or unprotected Gly residue to form the 2-3 eptifibatide fragment.

14. A process comprising:

coupling an amino-terminal protected homoarginine residue and a protected or unprotected glycine residue, thereby forming a 2-3 eptifibatide fragment of the formula:




coupling an amino-terminal protected aspartic acid residue having a protected carboxyl side chain to a tryptophan-proline dipeptide through the tryptophan residue of the dipeptide, and removing the amino-terminal protecting group from the aspartic acid residue, thereby forming a protected 4-6 eptifibatide fragment of the formula:



coupling the 2-3 eptifibatide fragment and the 4-6 eptifibatide fragment through attachment of the Gly residue of the 2-3 eptifibatide fragment to the Asp residue of the 4-6 eptifibatide fragment, thereby forming a 2-6 eptifibatide fragment of the formula:

- 29 -



wherein

Har is homoarginyl;

Gly is glycylyl;

Asp is aspartyl;


Trp is tryptophanyl;

Pro is prolyl;

P_1 and P_3 are amino protecting groups; and


P_2 is a carboxyl protecting group;

coupling the 2-6 eptifibatide fragment to an activated cysteinamide residue through the Pro residue of the 2-6 eptifibatide fragment, thereby forming a 2-7 eptifibatide fragment of the formula:



wherein ACys-NH₂ is an activated cysteinamide;

attaching a mercaptopropionic acid residue to the 2-7 eptifibatide fragment through a disulfide linkage between the mercaptopropionic acid residue and the ACys-NH₂ residue of the 2-7 eptifibatide fragment, thereby forming a compound of formula I:



(I)


wherein

Mpa is mercaptopropionic acid;

Cys-NH₂ is cysteinamide;

removing P_1 from the Har residue of the compound of formula I, thereby forming a compound of formula II:


- 30 -



(II)

;

coupling the Har and Mpa residues of the compound of formula II, thereby forming a compound of formula III:



(III)


; and

removing P₂ from the Asp residue of the compound of formula III to form eptifibatide.

15. The process of claim 14 wherein P₂ is stable under conditions suitable for removal of P₁.
16. The process of claim 14 wherein P₂ is t-butyl.
17. The process of claim 14 wherein P₁ is 9-fluorenylmethoxycarbonyl or benzyloxycarbonyl.
18. The process of claim 14 wherein ACys-NH₂ is H-Cys(Npys)-NH₂.

- 31 -

19. A compound having formula IV:



wherein

R_1 is hydrogen or P_1 ;

P_1 is an amino protecting group; and

P_2 is a carboxyl protecting group.

20. The compound of claim 19 wherein P_2 is t-butyl.

21. The compound of claim 19 wherein R_1 is hydrogen.

22. The compound of claim 21 wherein P_2 is t-butyl.

23. The compound of claim 19 wherein R_1 is P_1 .

24. The compound of claim 23 wherein P_1 is 9-fluorenylmethoxycarbonyl or benzyloxycarbonyl.


- 32 -

25. The compound of claim 24 wherein P₂ is t-butyl.
26. A compound selected from the group consisting of:
 Fmoc-Har-Gly-OH;
 Fmoc-Har-Gly-O-P₄; and
 Fmoc-Har-Gly-Asp(O-P₅)-Trp-Pro-OH

wherein

Fmoc is 9-fluorenylmethoxycarbonyl;
 Har is homoarginine;
 Gly is glycine;
 Asp is aspartic acid;
 Trp is tryptophan;
 Pro is proline;
 P₄ and P₅ are carboxyl protecting groups.

27. The compound of claim 26 wherein P₄ is pentafluorophenol.
28. The compound of claim 26 wherein P₅ is t-butyl.
29. 3-nitro-2-pyridinesulfonyl-cysteinamide (H-Cys(Npys)-NH₂).
30. A compound having the following formula:



31. A composition comprising eptifibatide and Gly-eptifibatide.
32. The composition of claim 31 comprising at least 99 % eptifibatide and Gly-eptifibatide in the range of about 0.01 % to about 1 %.

33. The composition of claim 32 comprising at least 99 % eptifibatide and Gly-eptifibatide in the range of about 0.01 % to about 0.1 %.
34. The product produced by a process according to claim 1.
35. The product produced by a process according to claim 4.
36. The product produced by a process according to claim 8.
37. The product produced by a process according to claim 9.
38. The product produced by a process according to claim 11.
39. The product produced by a process according to claim 12.
40. The product produced by a process according to claim 13.
41. A process for purifying eptifibatide comprising:
contacting an eptifibatide solution with a stationary phase comprising octadecyl carbon chains attached to silica;
washing the stationary phase contacted with the eptifibatide solution with a trifluoroacetic acid/acetonitrile solution;
optionally washing the stationary phase contacted with the eptifibatide solution with an acetic acid/acetonitrile solution; and
washing stationary phase contacted with the eptifibatide solution with an ammonium acid/acetonitrile solution.
42. The process of claim 41 wherein the trifluoroacetic acid/acetonitrile solution is a solution comprising 95 % of a 0.1 % aqueous trifluoroacetic acid solution and 5 % of an acetonitrile solution.

- 34 -

43. The process of claim 42 further comprising washing the stationary phase contacted with the eptifibatide solution with a trifluoroacetic acid/acetonitrile solution comprising 50 % of a 0.1 % aqueous trifluoroacetic acid solution and 50 % of an acetonitrile solution.

44. The process of claim 41 wherein the acetic acid/acetonitrile solution is a solution comprising 95 % of a 0.5 % aqueous solution of acetic acid and 5 % of an acetonitrile solution.

45. The process of claim 44 further comprising washing the stationary phase contacted with the eptifibatide solution with an acetic acid/acetonitrile solution comprising 50 % of a 0.5 % aqueous solution of acetic acid and 50 % of an acetonitrile solution.

46. The process of claim 41 wherein the ammonium acid/acetonitrile solution is a solution comprising 95 % of an aqueous 100 mM ammonium acid solution and 5 % of an acetonitrile solution.