



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification³ : A61K 31/74; A01N 37/00, 37/02 A01N 37/06, 37/08, 41/02 A01N 41/06, 43/00, 43/02 A01N 43/36, 43/40, 43/48 A01N 43/64, 43/72, 43/76 A01N 43/78, 43/80, 43/84 A01N 43/86, 43/88, 43/50 A01N 43/56, 47/10, 55/02 A01N 57/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 84/ 02650</p> <p>(43) International Publication Date: 19 July 1984 (19.07.84)</p>
<p>(21) International Application Number: PCT/US84/00028</p> <p>(22) International Filing Date: 4 January 1984 (04.01.84)</p> <p>(31) Priority Application Number: 455,727</p> <p>(32) Priority Date: 5 January 1983 (05.01.83)</p> <p>(33) Priority Country: US</p> <p>(71) Applicant: UNITED STATES OF AMERICA represented by THE SECRETARY, UNITED STATES DEPARTMENT OF COMMERCE [US/US]; National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161 (US).</p> <p>(72) Inventors: VANDER MEER, Robert, Kenneth ; 3712 N.W. 16th Boulevard, Gainesville, FL 32605 (US). WILLIAMS, David, Francis ; 8427 N.W. 2nd Place, Gainesville, FL 32607 (US). LOFGREN, Clifford, Swanson ; 1321 N.W. 31st Drive, Gainesville, FL 32604 (US).</p>	<p>(74) Agents: MORGAN, Thomas, J. et al.; Irons and Sears, 1785 Massachusetts Avenue, N.W., Washington, DC 20036 (US).</p> <p>(81) Designated States: AU, BR, JP.</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: A METHOD FOR THE CONTROL OF INSECTS</p> <p>(57) Abstract</p> <p>A method has been discovered for the control of arthropod pests which comprises treating the pest with an effective amount of selected fluorocarbons.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	KR	Republic of Korea
AU	Australia	LI	Liechtenstein
BE	Belgium	LK	Sri Lanka
BG	Bulgaria	LU	Luxembourg
BR	Brazil	MC	Monaco
CF	Central African Republic	MG	Madagascar
CG	Congo	MR	Mauritania
CH	Switzerland	MW	Malawi
CM	Cameroon	NL	Netherlands
DE	Germany, Federal Republic of	NO	Norway
DK	Denmark	RO	Romania
FI	Finland	SD	Sudan
FR	France	SE	Sweden
GA	Gabon	SN	Senegal
GB	United Kingdom	SU	Soviet Union
HU	Hungary	TD	Chad
JP	Japan	TG	Togo
KP	Democratic People's Republic of Korea	US	United States of America

A METHOD FOR THE CONTROL OF INSECTS

PRIOR ART

Discovering pesticides that are effective against a broad range of insect pests and that also can be used safely on crop lands and pastures has long been a problem. One such problem area has been in the control of ants and related insects. Particularly destructive are fire ants (such as Solenopsis invicta) which sting humans and livestock, feed on crop seedlings and germinating seeds thereby reducing yields, and damaging equipment which strike their mounds. Requirements for an effective pesticide formulation for the control of fire ants has been characterized as being (1) not repellent to the ants, (2) readily transferrable from one ant to another, and (3) exhibiting delayed toxicity. Repellency can reduce or negate the effectiveness of a toxicant because the ants will avoid the treated bait. The treated bait must be transferable either by carrying it back to the nest or by trophallaxis, and the toxicity must be delayed because foraging ants constitute only a small percentage of the total colony and must survive long enough to pass the toxicant onto the main colony population, especially the queen. It is preferable that the formulations exhibit delayed toxicity over a wide range of pesticide concentration because the active ingredient becomes diluted during trophallaxis (Banks et al, ARS-S-169, Oct. 1977). Presently only one commercially available pesticide (Amdro U.S. Patent No. 4,152,436) is registered as a bait for outdoor control of fire ants such as Solenopsis invicta, S. xyloni, S. richteri, and other ants including Pheidole megacephala and Iridomyrmex humilis. Amdro is also effective against Lepidopterous larvae. However, it cannot be applied to edible crops. The insecticide

SUBSTITUTE SHEET




-2-

Mirex (U.S. Patent No. 3,220,921) is also known to be effective against fire ants, but is no longer registered for use.

SUMMARY OF THE INVENTION

We have discovered a method and a composition for the control of a population of arthropods including ants, cockroaches, flies, mosquitoes, and termites. The method comprises treating the pests with an effective amount of a toxicant substance or mixture of substances of the formula



wherein R_f is a fluoroaliphatic radical containing up to 20 carbon atoms and A is a structurally compatible residue, or agriculturally acceptable salts of the toxicant substance or substances. The compositions in accordance with the invention comprise the above toxicant substance or mixture of substances and a bait component.

The fluoroaliphatic radical, R_f , is a fluorinated, monovalent moiety which is straight chain, branched chain, and if sufficiently large, cyclic, or combinations thereof, such as alkylcycloaliphatic radicals. The skeletal chain can include catenary oxygen and/or trivalent nitrogen hetero atoms bonded only to carbon atoms, such hetero atoms providing stable linkages between fluoro-carbon groups and not interfering with the chemically inert character of the R_f radical. While R_f can have a large number of carbon atoms, compounds where R_f is not more than 20 carbon atoms will be adequate and preferred since large radicals usually represent a less efficient utilization of fluorine than is possible with smaller R_f radicals. Generally, R_f will have up to 20 carbon atoms, preferably 5 to about 12. The terminal

SUBSTITUTE SHEET



-3-

portion of the R_f group has preferably at least three fully fluorinated carbon atoms, e.g., $CF_3CF_2CF_2-$, and the preferred compounds are those in which the R_f group is fully or substantially completely fluorinated, as in the case where R_f is perfluoroalkyl, C_nF_{2n+1} .

In the above formula A is a structurally compatible residue (i.e., capable of being linked to the SO_2 radical) which includes the radicals NR_1R_2 and OR_4 wherein R_1 and R_2 are selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, aralkyl, aroyl, acyl, cycloalkyl, cycloalkenyl, cycloalkynyl, a heterocyclic ring containing atoms selected from the group consisting of C, N, S or O, hydroxyalkyl, haloalkyl, aminoalkyl, carboxyalkyl salts, esters and amides, or a group of the structure $-(C_xH_{2x}O)_n(C_yH_{2y}O)_mR_3$ wherein $n + m = 1-20$, x and y are 1-4 and R_3 is selected from the same group as R_1 and R_2 ; and wherein R_4 is H, aryl, a heterocyclic ring or an alkaline earth, alkali metal, organic amine or ammonium cation. The formula NR_1R_2 also includes radicals in which N, R_1 and R_2 are taken together to form a ring the atoms of which are selected from the group consisting of C, N, S or O.

The salts of the invention are generally metal, ammonium or organic amine and quaternary amine salts and can be prepared by treating the acid-form compound with an appropriate base under mild conditions. Among the metal salts of the invention are alkali metal (e.g., lithium, sodium and potassium), alkaline earth metal (e.g., barium, calcium and magnesium) and heavy metal (e.g., zinc and iron) salts as well as other metal salts such as aluminum. Appropriate bases for use in preparing the metal salts include metal oxides, hydroxides, carbonates, bicarbonates and alkoxides. Some salts are also prepared by cation exchange reaction (by reacting a salt of the invention

SUBSTITUTE SHEET



-4-

with an organic or inorganic salt in a cation exchange reaction). The organic amine salts include the salts of aliphatic (e.g., alkyl), aromatic and heterocyclic amines, as well as those having a mixture of these types of structures. The amines useful in preparing the salts of the invention can be primary, secondary or tertiary and preferably contain not more than 20 carbon atoms. Such amines include, for example, morpholine, methyl cyclohexylamine, glucosamine, amines derived from fatty acids, etc. The amine and ammonium salts can be prepared by reacting the acid form with the appropriate organic base or ammonium hydroxide. Any of the salts of the types set out above are agriculturally acceptable, the one chosen depending upon the particular use and upon the economics of the situation. Of particular utility are the alkali metal, alkaline earth, ammonium and amines salts.

The salts of the invention are frequently formed by reacting the precursors in aqueous solution. This solution can be evaporated to obtain the salt of the compound, usually as a dry powder. In some cases, it may be more convenient to use a non-aqueous solvent such as alcohols, acetones, etc. The resulting solution is then utilized in formulations which are treated to remove the solvent, for example, by evaporation under reduced pressure.

These pesticides which are suitable for use in the invention fulfill the above mentioned requirements for fire ant control and in addition, possess very little mammalian toxicity. They are also effective against other arthropods.

DETAILED DESCRIPTION OF THE INVENTION

Arthropod pests are suitably treated in accordance with the invention in any manner known to the prior art

SUBSTITUTE SHEET



-5-

which is compatible with the above described toxicant compounds and mixtures. Suitable treatments include applying the toxicants as sprays in solutions, emulsions and dispersions; in traps with or without pheromone attractants and the like; and in bait formulations scattered in the vicinity of nests or in crop lands.

Substances suitable for use in this invention must be effective for at least one species of arthropod. However, there are several factors which may affect the effectiveness of specific substances with specific insects and with specific treatment techniques. These factors include:

1. Odor repellency.
2. Taste repellency.
3. Solubility in carriers such as solvents and baits.
4. Enzymatic effects.
5. Degradation by atmospheric oxygen, UV radiation and the like.

Using the invention for control of ants, in particular imported fire ants, it is preferable to prepare bait formulations into which the toxicants can be incorporated. The term "bait" is understood by those skilled in the art to be any substance that will entice the insect to ingest the toxicant. Suitable baits include edible oils and fats, vegetable seed meals, meat by-products such as blood, fish meal, syrups, honey, sucrose and other sugars, peanut butter, cereals and the like (see U.S. Patent No. 3,220,921). Preferred baits for fire ants are mixtures of edible oils (as solvents for the toxicant compounds) with granular carriers such as corncob grits, pregel defatted corn grits and the like. These impregnated granular bait formulations readily fall to the ground when dispersed by areal or ground applicator where the ants forage. When

SUBSTITUTE SHEET

-6-

found by the ants they are carried into the nest where the toxicants are ingested and distributed to workers and queen.

These compounds of the invention are known; but they are not known to be useful in pesticide formulations. Disclosures relevant to their preparation are found in the following U.S. patents:

2,003,615
2,346,612
2,732,378
2,759,019
2,803,656
2,915,554
3,398,182
3,787,351 and
4,424,178

The tests in the following examples indicate the effectiveness of the invention but are not intended to limit the invention's scope which is defined by the claims. All percentages and parts are by weight unless otherwise specified.

For fire ants any compound showing % mortality which is significantly greater than the bait without the toxicant is considered to be effective for the purposes of this invention. The use of preferred toxicants should result in less than 15% mortality at 24 hours and more than 50% and most preferably more than 85% mortality by the end of the test. The preferred compounds also should have at least a 10-fold difference between maximum and minimum dosages exhibiting delayed toxicity.

SUBSTITUTE SHEET

-7-

EXAMPLE 1

Each compound to be screened for fire ant toxicity was tested in 3 replications of 20 worker ants (Solenopsis invicta) which were placed in 30 ml cups for 14 days.

Cotton swabs saturated with soybean oil containing 1.0% of a test compound were offered to the ants in the cups for a 24 hour period. The swabs were removed and the ants remained without food for 24 hours. Cotton swabs saturated with SBO only were then placed in the cups and left there for the remainder of the testing period. Mortality counts were recorded for the test compounds and for a standard fire ant toxicant, Table 1.

EXAMPLE 2

Some of the compounds tested in Example 1 were retested as in Example 1 at concentrations of 0.01%, 0.10% and 1.0%, Table 2.

EXAMPLE 3

Some of the compounds listed in Table 1, which were not readily soluble in SBO were retested in the same manner as in Example 1 with the exception that SBO was replaced with a 1:1 v/v mixture of honey and water, Table 3.

EXAMPLE 4

Some of the preferred compounds from Example 2 were tested against duplicate laboratory colonies of fire ants.

SUBSTITUTE SHEET

-8-

The colonies consisted of a queen, eggs, larvae, pupae and greater than 40,000 workers. The test compounds were dissolved in SBO at 1.0% concentration and impregnated on pregel defatted corn grits so that the corn grits contained 30% SBO mixture. The test compound, therefore, was 0.30% of the total bait weight.

Five grams of the bait was made available to each colony for 4 days. The bait was removed and the colonies fed a standard diet for the remainder of the test which consisted of 1:1 honey-water mixture, boiled eggs and frozen fly pupae and cockroaches. Two colonies were treated with bait without toxicant as a control. Observation on the status of the queen and workers are recorded in Table 4. The ultimate fate of the colony is indicated as QD (queen dead) or CN (colony normal), Table 4.

EXAMPLE 5

Mixed sexes of adult house flies (Musca domestica) from an insecticide-resistant laboratory strain were fed after emergence exclusively on a fly food bait (6 parts sugar, 6 parts powdered nonfat dry milk, and 1 part powdered egg yolk) containing 1% of the test compounds. The bait was prepared by adding 10 ml of a solution or suspension of the test compound in a volatile solvent to 10 g of fly food in a small container. The solvent was allowed to evaporate for 4 to 6 hours, then the treated fly food was repulverized. The container of treated fly food and a container of water was placed in a cage with 100 newly emerged adult flies. Percent mortality for two replicate tests was determined after 3 days and compared to the results of feeding untreated fly food, Table 5.

SUBSTITUTE SHEET



-9-

Some of the above compounds may be more effective than the data indicates. If solubilities were poor in the SBO or honey-water formulation, concentrations of toxicants may not have reached the desired level. Many of these compounds showing poor effectiveness may be highly effective in other formulations.

EXAMPLE 6

Compounds 29757, 29758 and 29759 (see Table 1 for structures) were each dissolved in SBO at 1% concentration. Bait was prepared by impregnating the SBO mixture on pregel defatted corn grits so that the corn grit mixture contained 30% SBO mixture and 0.3% test compound. Compound 29759 was also used at 2.5% concentration so that it was 0.75% of the total bait weight.

Treated baits were scattered by tractor on field plots containing a number of active fire ant mounds at a concentration of one pound per acre. Other plots containing fire ant mounds were treated in the same manner with untreated pregel defatted corn grits and with grits treated with a standard fire ant toxicant as control. Results were evaluated as described by D. P. Harlan, W. H. Banks and C. E. Stringer, *Southwest Entomologist*, Vol. 6, pp. 150-157, 1981, Table 6.

EXAMPLE 7

Compounds 29756, 29757, 29759 and 29778 were used to treat Orlando normal colonies of American cockroaches (Periplaneta americana) each containing 10 adults of mixed sexes as described by G. S. Burden, *Pest Control* Vol. 48, pp. 22-24, 1980. A bait was formulated which consisted of a 3:1 mixture of cornmeal and powdered sugar containing

SUBSTITUTE SHEET



-10-

2.0% of the above compounds or a standard cockroach toxicant (trichlorfon). Untreated 3:1 mixture of cornmeal and powdered sugar was used as a control. A container with 2 grams of candidate or standard bait and a container with 2 grams of normal diet (Purina Lab Chow 5001) were placed in each arena with the cockroaches and left until the test was ended. Percent mortality is shown in Table 7.

EXAMPLE 8

Orlando normal colonies of German cockroaches (Blattella germanica) containing 25 adults were treated in the same manner as in Example 7. Results are shown in Table 8.

EXAMPLE 9

Twenty-five late 3rd- and early 4th-stage larvae of Anopheles quadrimaculatus were placed in a 500 ml glass jar (9 x 8.5 cm diam) containing 100 ml of well water, 0.05 g ground hog supplement for larval food, and a known amount of the candidate compound in not more than 1.0 ml acetone. The jars were covered with cloth netting and held in constant temperature incubators at 26.7-28.9°C (80-84°F) and 65-75% RH; a low level of illumination (ca. 0.5 footcandles inside the incubators) was maintained during nonworking hours. Seven days after set up, the jars were examined for the number of dead pupae, the number of adults that were dead or unable to complete eclosion, and the number of exuviae. The live adults were observed for gross abnormalities. Tests were replicated for at

SUBSTITUTE SHEET



-11-

at least 1 concn. and a standard larvicide, methoprene, was used as a control with each test series.

Several concns. of each compound were tested so that a dose-response relationship could be established. Using the Statistical Analysis System supported by NERDC, the resulting data were corrected for control mortality (Abbott's formula) and a probit analysis was made using log transformed mortality data. This analysis provided estimates for the LC-50 and LC-90, in ppm, for the compound.

EXAMPLE 10

Compounds were screened as mosquito larvicides (Insect Growth Regulators, IGR) by exposing early 4th-instar larvae of Anopheles quadrimaculatus to solutions or suspensions of the compounds in water (duplicate tests). The compounds were dissolved in acetone and added to water; water-soluble compounds remained in solution and the others became finely divided suspensions. Mosquito larvae were added to the treated water and mortality was determined after 24 hours of exposure. The compounds were initially tested at concns. of 10 and 1 parts per million. If 50% mortality occurred at 1 ppm, additional tests were conducted with lower concns. A standard larvicide, temephos (Abate), was tested as a control concurrently.

Results are shown in Table 10.

The following indicates the criteria for compounds tested against mosquitoes:

SUBSTITUTE SHEET



Classification of IGR's Against Mosquitoes

<u>Class</u>	<u>Criteria</u>	<u>LC-90 (ppm)</u>
1	Ineffective at screening dose	> 1.000
2	Partially effective but not promising	0.101-1.000
3	Effective enough to justify full investigation	0.021-0.100
4	Exceptional	≤ 0.020


Results are shown in Table 9.

SUBSTITUTE SHEET



TABLE I

Number	Structure	Mortality at Specified Days						
		%						
		1	2	3	6	8	10	14
29752	$\begin{array}{c} \text{C}_2\text{H}_5 \quad \text{O} \\ \quad \parallel \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OP(OH)}_2 \end{array}$	2	3	3	3	3	3	10
29753	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{O})_3\text{H} \end{array}$	0	0	0	8	65	87	93
29754	$\begin{array}{c} \text{C}_4\text{H}_9 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OH} \end{array}$	0	0	0	3	32	72	93
29755	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_6\text{F}_{13}\text{SO}_2\text{NC}_2\text{H}_4\text{OH} \end{array}$	0	0	0	17	38	87	97
29756	$\begin{array}{c} \text{CH}_3 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_4\text{H}_8\text{OH} \end{array}$	3	3	13	73	90	97	98
29757	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_5 \end{array}$	33	88	100				
29758	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_3 \end{array}$	18	88	100				
29759	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NH}_2$	18	63	85	88	97	100	
29760	$\text{C}_8\text{F}_{17}\text{SO}_3\text{-} \begin{array}{c} \text{C}_6\text{H}_4 \\ \\ \text{NH}_2 \end{array}$	0	0	0	0	0	3	70
29761	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{NH}_2 \end{array}$	2	2	2	2	2	2	10

29762	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{NH})_2\text{C}_2\text{H}_4\text{NH}_2 \end{array}$	2	2	2	2	2	2	10
29763	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_6\text{F}_{13}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{O})_{12}\text{H} \end{array}$	2	3	3	7	7	13	40
29764	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_{10}\text{F}_{21}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{O})_{14}\text{H} \end{array}$	0	0	0	0	0	0	17
29765	$\begin{array}{c} \text{C}_{12}\text{H}_{25} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OH} \end{array}$	0	0	0	2	17	47	83
29766	Adduct from							
	<p>1 mole</p> 	0	0	0	3	3	3	23
	1 mole $\text{C}_{18}\text{H}_{37}\text{OH}$ and							
	<p>2 moles</p> $\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OH} \end{array}$							
29767	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2-\text{C}_6\text{H}_5 \end{array}$	0	0	0	2	25	80	97
29768	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{O})_{12.5}\text{CH}_3 \end{array}$	0	0	0	7	42	87	97

29769	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{O})_{17}\text{CH}_3 \end{array}$	0	0	13	70	90	95	98
29770	$\begin{array}{c} \text{H} \\ \\ \text{C}_6\text{F}_{13}\text{SO}_2\text{NC}_2\text{H}_5 \end{array}$	68	100					
29771	$\begin{array}{c} \text{C}_2\text{H}_5 \quad \text{O} \\ \quad \parallel \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OCC}_{17}\text{H}_{35} \end{array}$	0	2	2	48	90	100	
29772	$\begin{array}{c} \text{C}_4\text{H}_9 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{O}(\text{C}_3\text{H}_6\text{O})_8\text{H} \end{array}$	0	0	0	2	5	17	62
29773	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{O}(\text{C}_3\text{H}_6\text{O})_8\text{H} \end{array}$	0	2	2	2	2	12	43
29774	$\begin{array}{c} \text{H} \\ \\ \text{C}_4\text{F}_9\text{SO}_2\text{NC}_{12}\text{H}_{25} \end{array}$	0	0	0	0	2	3	7
29775	$\begin{array}{c} \text{H} \\ \\ \text{C}_4\text{F}_9\text{SO}_2\text{NC}_3\text{H}_6\text{N}(\text{CH}_3)_2 \end{array}$	0	0	3	3	3	5	13
29776	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_3\text{H}_6\text{N}(\text{CH}_3)_2 \end{array}$	0	0	0	0	0	3	23
29777	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_{12}\text{H}_{25} \end{array}$	0	0	5	47	82	98	100
29778	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N} \begin{array}{c} \diagup \text{N} \\ \diagdown \end{array}$	0	0	70	93	98	98	100
29779	$\text{C}_2\text{F}_5\text{C}_6\text{F}_{10}\text{SO}_3\text{K}$	0	2	2	7	7	10	15








29780	$C_7F_{15}CO_2NH_4$	0	0	2	2	3	3	5
29781	$ \begin{array}{c} H \\ \\ C_8F_{17}SO_2NCH_2- \text{C}_6\text{H}_4-SO_3Na \end{array} $	0	0	0	18	40	80	97
29782	$ \begin{array}{c} C_2H_5 \\ \\ C_8F_{17}SO_2NC_2H_4OH \end{array} $	0	0	5	15	53	83	100
50950	$C_8F_{17}SO_3K$	0	0	0	0	0	0	0
10700	$ \begin{array}{c} H \\ \\ C_8F_{17}SO_2NNa \end{array} $	0	3	7	58	70	80	87
10701	$ \begin{array}{c} CH_3 \\ \\ C_8F_{17}SO_2NNa \end{array} $	0	15	80	97	98	98	100
10702	$C_6F_{13}SO_2NH_2$	3	8	40	80	83	85	92
10703	$CF_3SO_2NH_2$	3	13	25	58	70	72	72
10704	$ \begin{array}{c} H \\ \\ C_4F_9SO_2NCH_3 \end{array} $	68	97	100				
10705	$ \begin{array}{c} CH_3 \quad O \\ \quad / \quad \backslash \\ C_8F_{17}SO_2NCH_2CH-CH_2 \end{array} $	2	2	2	3	8	15	30
10706	$ \begin{array}{c} CH_3 \quad O \\ \quad \\ C_8F_{17}SO_2N-C_2H_4CNH_2 \end{array} $	0	0	0	0	3	5	10







-17-




10707	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_5)_2$	17	100						
10708	$\begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ \quad \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{CH}_2)_{10}\text{NSO}_2\text{C}_8\text{F}_{17} \end{array}$	0	2	2	7	7	7	7	
10709	$\begin{array}{c} \text{C}_2\text{H}_5 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{C}\equiv\text{CH} \end{array}$	0	2	2	48	53	72	85	
10710	$\begin{array}{c} \text{H} \\ \\ n\text{-C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{CH}=\text{CH}_2 \end{array}$	2	25	55	90	95	98	98	
10711	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{CH}=\text{CH}_2 \end{array}$	0	12	40	93	97	98	98	
10712	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}(\text{CH}_3)_2 \end{array}$	8	88	100					
10713	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}(\text{CH}_3)_3 \end{array}$	0	0	0	0	0	0	5	
10714	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_6\text{H}_5 \end{array}$ (mixed isomers)	75	82	82	90	90	92	93	
10715	$\begin{array}{c} \text{H} \\ \\ n\text{-C}_8\text{F}_{17}\text{SO}_2\text{NC}_6\text{H}_5 \end{array}$ (recrystallized linear isomers)	68	82	85	92	92	92	100	
10716	$\begin{array}{c} \text{H} \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_6\text{H}_5 \end{array}$ (branched isomers)	52	63	63	77	83	88	90	
10717	$\begin{array}{c} \text{CH}_3 \\ \\ \text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}=\text{CH}_2 \end{array}$	23	87	98	100				

SUBSTITUTE SHEET



10718		0	2	3	3	3	5	12
10719		2	5	7	7	7	7	7
10720		0	0	0	3	7	12	32
10721	C ₈ H ₁₇ SO ₂ NH ₂	5	7	8	12	12	12	12
10722		0	0	0	2	3	3	3
10723		2	2	2	5	5	7	8
10724		0	0	0	2	2	2	3
10725		0	0	0	0	7	18	52

10726		0	0	0	0	0	0	0
10727	C ₈ F ₁₇ SO ₃ H	0	3	3	12	12	22	22
10728	C ₆ F ₁₃ SO ₃ H	0	0	0	3	7	7	8
10729		0	0	0	7	13	25	50
10730		0	2	2	3	20	30	62
10731	C ₈ F ₁₇ SO ₂ N(C ₂ H ₄ OH) ₂	0	0	0	2	5	10	35
10732		0	0	0	2	2	2	5
10733		5	50	70	93	97	98	100
10734		2	3	5	5	8	20	57
10735	C ₈ F ₁₇ C ₂ H ₄ SH	3	3	3	3	3	5	5

10736	$[\text{CF}_2\text{CF}_2\text{SO}_2\text{NH}_2]_2$	0	2	2	3	3	3	5
10737	$\text{n-C}_3\text{F}_7\text{CNH}_2$ $\begin{array}{c} \text{O} \\ \\ \text{C} \end{array}$	0	3	3	10	10	10	25
10738	$\text{C}_5\text{F}_{11}\text{CNH}_2$ $\begin{array}{c} \text{O} \\ \\ \text{C} \end{array}$	3	3	3	5	5	5	5
10739	$\text{C}_7\text{F}_{15}\text{CNH}_2$ $\begin{array}{c} \text{O} \\ \\ \text{C} \end{array}$	0	0	0	2	2	2	2
10740	$\text{C}_7\text{F}_{15}\text{CNCH}_2\text{C}_6\text{H}_5$ $\begin{array}{c} \text{H} \\ \\ \text{C} \\ \\ \text{O} \end{array}$	0	0	0	0	0	3	3
10741	$\text{CF}_3\text{SO}_2\text{N}$ $\begin{array}{l} \text{H} \\ \diagup \\ \text{N} \\ \diagdown \\ \text{CH}_3 \end{array}$	3	3	3	32	35	38	48
10742	$\text{CF}_3\text{SO}_2\text{N}(\text{CH}_3)_2$	82	82	82	82	82	82	90
10743	$\text{CF}_3\text{SO}_2\text{NH}_2$ 	0	0	0	12	13	15	20
10744	$\text{C}_2\text{F}_5\text{SO}_2\text{NH}_2$	3	17	22	35	40	45	50
10745	$\text{C}_4\text{F}_9\text{SO}_2\text{NH}_2$	0	0	0	3	3	3	5
10746	$\text{CF}_3\text{SO}_2\text{N}$ -C(=O)- 	0	5	8	13	15	17	23
10747	$\text{C}_4\text{F}_9\text{SO}_2\text{N}$ -C(=O)- 	0	0	0	32	48	53	73


10748	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}-\overset{\text{H O}}{\underset{\text{ }}{\text{C}}}-\text{C}_6\text{H}_5$	3	13	17	37	42	47	85
10749	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}-\overset{\text{C}_2\text{H}_5}{\text{C}}-(\text{C}_2\text{H}_4\text{O})_7\text{CH}_3$	2	5	5	38	57	80	88
10750	$\text{C}_8\text{F}_{17}\text{SO}_3^{\ominus} \text{N}^{\oplus}(\text{C}_2\text{H}_5)_4$	0	0	0	3	3	3	7
10751	$\text{C}_6\text{F}_{10}(\text{CH}_2\text{OH})_2$ (Cyclic)	2	5	5	23	28	32	58
10752	$\text{FCH}_2\text{SO}_2\text{NH}_2$	0	0	2	30	33	43	53
10753	$\text{HCF}_2\text{SO}_2\text{NH}_2$	0	2	2	13	18	20	30
10754	$\text{CF}_3\text{CH}_2\text{SO}_2\text{NH}_2$	0	0	0	2	3	5	13
	Standard	0	12	97	100			
	SBO	0	0	0	5	8	9	13

TABIE 2

Number	Structure	Conc. &	Percent Mortality at Specified Days													
			1	2	3	6	8	10	14	17	21					
29753	$\overset{\text{C}_2\text{H}_5}{\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{O})_3\text{H}}$	0.01	0	0	0	0	3	7	13	20						
		0.1	0	0	0	0	0	3	7	27						
		1.0	0	0	2	52	87	98	99	100						
29754	$\overset{\text{C}_4\text{H}_9}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OH}}$	0.01	0	0	0	0	0	0	0	2						
		0.1	0	2	2	3	3	25	48	78						
		1.0	0	0	0	0	0	40	92	98	100					
29755	$\overset{\text{C}_2\text{H}_5}{\text{C}_6\text{F}_{13}\text{SO}_2\text{NC}_2\text{H}_4\text{OH}}$	0.01	0	0	0	5	7	12	15	22	28					
		0.1	0	0	0	2	2	7	10	23	40					
		1.0	0	0	0	12	45	80	95	98	98					
29756	$\overset{\text{CH}_3}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_4\text{H}_8\text{OH}}$	0.01	0	0	2	5	8	8	8	10	13					
		0.1	0	0	0	2	5	30	75	85	92					
		1.0	0	2	10	83	85	95	100							
29757	$\overset{\text{H}}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_5}$	0.01	0	0	0	0	2	2	10	22	50					
		0.1	0	0	2	80	97	97	98	98	100					
		1.0	25	100												
29758	$\overset{\text{H}}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_3}$	0.01	0	0	2	3	7	7	7	23	40					
		0.1	0	0	7	88	97	98	100							
		1.0	17	93	100											



SUBSTITUTE SHEET



29759	C ₈ F ₁₇ SO ₂ NH ₂	0.01 0.1 1.0	0 0 43	0 0 85	0 0 98	3 2 100	7 33 77	7 77	10 92	20 95	23 98
29765	^{C₁₂H₂₅} C ₈ F ₁₇ SO ₂ NC ₂ H ₄ OH	0.01 0.1 1.0	0 0 0	0 0 0	0 0 0	0 0 2	0 0 32	0 0 77	0 2 88	0 3 88	0 15 100
29767	^{C₂H₅} C ₈ F ₁₇ SO ₂ NCH ₂ - 	0.01 0.1 1.0	0 0 0	0 0 0	0 2 2	0 2 42	3 3 83	3 8 100	3 18	3 42	3 42
29769	^{C₂H₅} C ₈ F ₁₇ SO ₂ N(C ₂ H ₄ O) ₁₇ CH ₃	0.01 0.1 1.0	0 0 0	0 0 0	3 2 32	7 3 100	7 5	10 10	13 25	17 48	17 48
29770	^H C ₆ F ₁₃ SO ₂ NC ₂ H ₅	0.01 0.1 1.0	2 0 47	2 2 98	3 2 98	5 5 100	7 7	8 17	10 48	13 58	15 70
29771	^{C₂H₅} C ₈ F ₁₇ SO ₂ NC ₂ H ₄ OCC ₁₇ H ₃₅	0.01 0.1 1.0	0 2 0	0 2 0	0 7 87	0 10 98	0 17 98	8 20 98	10 32 98	13 58 98	18 58 98
29772	^{C₄H₉} C ₈ F ₁₇ SO ₂ NC ₂ H ₄ O(C ₃ H ₆ O) ₈ H	0.01 0.1 1.0	0 0 0	0 0 2	2 0 2	2 0 2	2 0 40	2 3 87	2 15 97	2 15 97	2 45 100

SUBSTITUTE SHEET



29773	^{C₂H₅} C ₈ F ₁₇ SO ₂ NC ₂ H ₄ O(C ₃ H ₆ O) ₈ H	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3
		1.0	0	0	0	0	3	5	23	37	45	60								
29776	^H C ₈ F ₁₇ SO ₂ NC ₃ H ₆ N(CH ₃) ₂	0.01	0	0	0	0	2	2	2	2	2	2	2	2	2	2	2	2	2	2
		0.1	0	0	0	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3
		1.0	0	0	0	0	0	0	0	0	7	35								
29777	^H C ₈ F ₁₇ SO ₂ NC ₁₂ H ₂₅	0.01	0	0	0	5	5	5	5	7	13	17								
		0.1	0	0	0	0	2	2	2	20	50	80								
		1.0	0	2	3	78	97	100												
29778	^N C ₈ F ₁₇ SO ₂ N 	0.01	2	3	3	8	8	8	8	13	15	18								
		0.1	0	2	3	10	10	10	52	67	80	88								
		1.0	0	2	17	92	100													
29779	C ₂ F ₅ C ₆ F ₁₀ SO ₃ K	0.01	0	0	0	2	2	2	2	2	2	2								
		0.1	0	0	0	0	2	3	3	3	3	3								
		1.0	0	0	0	0	2	2	2	5	5	7								
29781	^H C ₈ F ₁₇ SO ₂ NCH ₂ - 	0.01	0	0	0	2	2	5	5	5	7	7								
		0.1	0	0	0	3	3	5	8	8	12	23								
		1.0	0	0	0	77	93	95	100											
29782	^{C₂H₅} C ₈ F ₁₇ SO ₂ NC ₂ H ₄ OH	0.01	0	0	2	2	2	2	2	2	2	3								
		0.1	0	0	0	0	2	2	2	8	40	60								
		1.0	0	0	0	45	67	88	98	98	100									

SUBSTITUTE SHEET



10700	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NNA}^{\text{H}}$	0.01	2	3	3	7	7	7	8	12	15
		0.1	2	3	3	8	12	40	70	82	95
		1.0	0	2	10	75	88	93	98	100	
10701	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NNA}^{\text{CH}_3}$	0.01	0	0	0	0	0	0	0	2	7
		0.1	0	0	0	18	65	83	95	97	98
		1.0	23	87	100						
10702	$\text{C}_6\text{F}_{13}\text{SO}_2\text{NH}_2$	0.01	0	0	0	2	2	2	2	5	12
		0.1	3	7	7	7	7	17	63	77	92
		1.0	0	3	30	67	75	87	95	98	100
10703	$\text{CF}_3\text{SO}_2\text{NH}_2$	0.01	0	0	0	3	3	3	7	8	17
		0.1	3	3	3	5	5	5	17	27	58
		1.0	2	8	18	33	42	50	67	73	82
10704	$\text{C}_4\text{F}_9\text{SO}_2\text{NCH}_3^{\text{H}}$	0.01	0	2	3	7	7	8	8	15	27
		0.1	2	2	2	3	3	3	3	3	10
		1.0	87	98	98	98	98	100			
10707	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_5)_2$	0.01	0	0	0	2	5	10	20	50	60
		0.1	0	7	13	78	92	98	100		
		1.0	30	100							
10709	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_5)_2\text{C}\equiv\text{CH}$	0.01	5	5	5	5	5	5	15	15	30
		0.1	2	2	2	2	3	5	43	73	87
		1.0	0	0	0	2	45	60	90	93	100

SUBSTITUTE SHEET



10710	$\overset{\text{H}}{\text{n-C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{CH}=\text{CH}_2$	0.01 0.1 1.0	2 3 13	2 3 53	2 3 80	2 3 48	2 60 100	2 78	2 93	12 37	37 98	75 100
10711	$\overset{\text{H}}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{CH}=\text{CH}_2$	0.01 0.1 1.0	2 3 23	2 3 57	2 3 72	2 3 88	2 42 100	5 85	5 97	5 97	7 98	28 100
10712	$\overset{\text{H}}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}(\text{CH}_3)_2$	0.01 0.1 1.0	2 0 83	2 0 97	2 10 100	2 75	2 93	2 98	3 100	5 98	27	65
10714	$\overset{\text{H}}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_6\text{H}_5}$ (mixed isomers)	0.01 0.1 1.0	0 0 83	0 2 87	0 2 88	2 8 95	2 53 97	3 70 97	3 93 100	7 93 100	17 95	27 98
10715	$\overset{\text{H}}{\text{n-C}_8\text{F}_{17}\text{SO}_2\text{NC}_6\text{H}_5}$ (recrystallized linear isomer)	0.01 0.1 1.0	2 0 88	2 0 92	2 0 93	2 18 97	2 63 98	2 87 98	2 90 98	3 90 98	12 95 98	27 93 100
10716	$\overset{\text{H}}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_6\text{H}_5}$ (branched isomer)	0.01 0.1 1.0	2 0 78	5 3 83	5 3 87	8 12 100	8 40	8 70	8 93	8 93	13 97	27 97
10717	$\overset{\text{CH}_3}{\text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}=\text{CH}_2$	0.01 0.1 1.0	0 0 100	0 7	0 33	8 77	8 90	13 92	25 100	37	57	

SUBSTITUTE SHEET



10733	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\overset{\text{C}_2\text{H}_5}{\text{C}_2\text{H}_4})\text{Cl}$	0.01	0	0	0	0	3	3	3	5	23	47
		0.1	0	0	2	22	83	95	97	100		
		1.0	57	87	98	100						
10742	$\text{CF}_3\text{SO}_2\text{N}(\text{CH}_3)_2$	0.01	0	0	0	0	0	3	12	13	30	
		0.1	0	0	3	3	5	8	17	35	53	
		1.0	88	90	90	97	98	98	98	98	98	100
10749	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\overset{\text{C}_2\text{H}_5}{\text{C}_2\text{H}_4})_7\text{CH}_3$	0.01	2	3	3	3	3	5	8			
		0.1	5	5	8	15	20	23	40			
		1.0	2	5	5	38	57	80	88			
----	-----	Copolymer of										
	30%	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\overset{\text{C}_4\text{H}_9}{\text{C}_2\text{H}_4})_2$										
	70%	$\text{CH}_2=\overset{\text{O}}{\overset{\text{H}}{\text{C}}}-\text{CH}_2$	0.01	-	-	-	-	-	-	-	-	-
		$\text{CH}_2=\overset{\text{O}}{\overset{\text{O}}{\text{C}}}-\text{CH}(\text{O}(\text{C}_2\text{H}_4\text{O}))_10-$	0.1	0	8	10	18	18	52	-	87	
		$(\text{C}_3\text{H}_6\text{O})_{22}(\text{C}_2\text{H}_4\text{O})_{10}-\text{CH}=\text{CH}_2$	1.0	3	3	7	13	15	47	-	73	

SUBSTITUTE SHEET











-----	50% Copolymer of	0.01	2	3	5	7	10	12	13	15	22	25	30	58	80	78
	C_2H_5															
	30% $\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4$ -	0.1	5	7	7	15	22	25	30	58	80	78				
	$\text{O} \begin{matrix} \text{CH}_3 \\ \\ \text{OC}-\text{C}=\text{CH}_2 \end{matrix}$	1.0	2	3	5	7	10	12	13	15	22	25	30	58	80	78
	70% $\text{CH}_3\text{O}(\text{C}_2\text{H}_4\text{O})_{16}$															
	$\text{O} \begin{matrix} \text{H} \\ \\ \text{R}-\text{C}=\text{CH}_2 \end{matrix}$															
	50% $\text{C}_2\text{H}_5\text{OOCCH}_3$															
Standard		0.01	0	0	0	0	0	12	37	48	62	83				
		0.1	3	5	20	88	97	100								
		1.0	2	75	100											
SBO			0	1	1	3	6	11	13	19	22					

SUBSTITUTE SHEET







TABLE 3

Percent Mortality at Specified Days



Numbers	Structure	Conc. %	Percent Mortality at Specified Days										
			1	2	3	6	8	10	14				
10950	C ₈ F ₁₇ SO ₃ K	1.0	2	2	23	87	100						
10700	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NNa}$ 	1.0	2	37	73	98	100						
10701	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NNa}$ 	1.0	62	97	100								
10705	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{CH}_2)_2\text{CH}(\text{O})\text{CH}_2$ 	1.0	0	0	0	0	0	0	0	0	8	42	
10706	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4)_2\text{CNH}_2$ 	1.0	0	0	2	2	2	2	2	7	62		
10708	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{CH}_2)_{10}\text{SO}_2\text{C}_8\text{F}_{17}$ 	1.0	0	0	0	0	0	0	0	0	0	35	
10710	$\bar{n}\text{-C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{CH}=\text{CH}_2$ 	1.0	0	12	55	97	97	97	98	100			
10712	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}(\text{CH}_3)_2$ 	1.0	23	100									
10719	$[\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_5)_2\text{H}_4\text{OCNH}-$ 	1.0	0	0	3	10	33	55	85				



10720	$(C_8F_{17}SO_2NC_2H_4O-$ $(CH_2CHO)5C$  CH_2Cl	1.0	10	22	33	33	33	33	35	43
10722	$C_8H_{17}SO_2NC_{12}H_{25}$	1.0	0	0	0	0	2	3	5	
10723	$C_{17}H_{35}C$ 	1.0	0	0	0	0	0	0	0	0
10724	$C_8F_{17}SO_2NC_{12}H_{25}CO_2H$	1.0	0	0	2	7	7	10	12	
10725	$C_8F_{17}SO_3-$ 	1.0	0	0	2	2	3	5	8	
10726	$C_8F_{17}SO_3-$ 	1.0	0	0	0	0	0	2	5	
10727	$C_8F_{17}SO_3H$	1.0	0	37	62	95	100			
10728	$C_6F_{13}SO_3H$	1.0	15	77	95	100				
10729	$C_8F_{17}SO_2NC_2H_5CO_2CH_3$	1.0	0	0	0	22	40	58	90	

SUBSTITUTE SHEET



10730	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_2\text{H}_4\text{OH}$ C_3H_7	1.0	0	0	0	0	0	0	2	0	2	12	77
10731	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_4\text{OH})_2$	1.0	0	0	0	0	0	0	0	0	25	57	
10732	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NCH}_2\text{CH}-\text{CH}_2$ CH_3 OH OH	1.0	0	0	0	0	0	0	0	0	0	2	
10734	$\text{C}_8\text{F}_{17}\text{SO}_2\text{NC}_4\text{H}_9\text{SH}$ CH_3	1.0	0	0	0	0	0	2	2	2	2	5	
10738	$\text{C}_5\text{F}_{11}\text{CNH}_2$	1.0	0	0	0	2	2	2	2	2	7	10	
10739	$\text{C}_7\text{F}_{15}\text{CNH}_2$	1.0	0	0	0	0	2	2	3	17	35		
10740	$\text{C}_7\text{F}_{15}\text{CNCH}_2\text{C}_6\text{H}_5$ OH	1.0	0	0	0	0	0	0	2	2	2	2	
10743	$\text{CF}_3\text{SO}_2\text{NH}_2 \cdot \text{HN}$ 	1.0	0	2	2	15	27	42	57				
10748	$\text{C}_8\text{F}_{17}\text{SO}_2\text{N}-\text{C}$ 	1.0	0	0	0	2	3	13	63				
10750	$\text{C}_8\text{F}_{17}\text{SO}_3^- \oplus \text{N}(\text{C}_2\text{H}_5)_4$	1.0	10	47	95	100							
10752	$\text{FCH}_2\text{SO}_2\text{NH}_2$	1.0	0	0	0	2	5	17	45				

SUBSTITUTE SHEET



10754	CF ₃ CH ₂ SO ₂ NH ₂	1.0	0	0	0	0	0	0	0	2	18	2	25	52	62
-----	Copolymer of														
	30% C ₈ F ₁₇ SO ₂ NC ₂ H ₄ -	1.0	0	0	0	7	12	20	20	20	13	18	40	73	
	70% CH ₂ =CHCO(C ₂ H ₄ O) ₁₀ -														
	(C ₃ H ₆ O) ₂₂ (C ₂ H ₄ O) ₁₀ -														
	50% of Copolymer														
	30% C ₈ F ₁₇ SO ₂ NC ₂ H ₄ -	1.0	0	0	0	2	2	10	10	13	18	40	73		
	70% CH ₃ O(C ₂ H ₄ O) ₁₆ C(C ₂ H ₄ O) ₁₆ -														
	50% C ₂ H ₄ OCCH ₃														

SUBSTITUTE SHEET




Standard	1.0	0	0	100	0	0	0	0	0	1	1
Honey:Water											

SUBSTITUTE SHEET



TABLE 4


Number	Structure	Percent worker mortality after following days of initial exposure to bait					
		2	10	14	16	23	26
29757	$C_8F_{17}SO_2\overset{H}{N}C_2H_5$	40	87	95	QD		
29757		10	50	75	92 QD		
29758	$C_8F_{17}SO_2\overset{H}{N}CH_3$	10	30	50	75	90	92 QD
29758		5	28	35 QD			
29759	$C_8F_{17}SO_2NH_2$	15	50	65 QD			
29759		8	25	30	30	50 QD	
29778	$C_8F_{17}SO_2N$ 	0	10	35	50 QD		
29778		0	15 QD				
Control		0	1	2	2	2	2 CN
Control		0	2	3	3	5	5 CN

SUBSTITUTE SHEET



-35-

TABLE 5

Number	Structure	% Mortality After 3 Days
29756	$C_8F_{17}SO_2N(CH_3)C_4H_8OH$	0
29757	$C_8F_{17}SO_2N(H)C_2H_5$	0
29758	$C_8F_{17}SO_2N(H)CH_3$	0
29759	$C_8F_{17}SO_2NH_2$	80
29778	$C_8F_{17}SO_2N$ 	60
Untreated Fly Food		0

SUBSTITUTE SHEET



TABLE 6
Population Index

Treatment	Conc. %	Population Index		
		Pre-treatment	6 weeks	% Reduction
Untreated		305	305	0
		130	155	0
		105	185	0
Standard	1	120	19	84
		140	0	100
		255	15	94
29758	1	170	0	100
		190	7	96
		170	0	100
29757	1	150	0	100
		160	48	70
		160	32	80
29759	2.5	195	2	99
		180	0	100
		175	25	86
29759	1	165	0	100
		195	25	87
		180	81	55

SUBSTITUTE SHEET



-37-

TABLE 7

Treatment	Percent Mortality After Indicated Days				
	1	2	3	7	10
29756	0	0	0	60	100
	0	0	20	100	
29757	0	10	30	90	100
	0	40	40	100	
29758	0	30	40	80	100
	0	20	30	90	100
29759	20	50	90	100	
	30	80	90	100	
29778	0	20	40	100	
	0	50	80	100	
Standard	100				
	100				
Cornmeal-Powdered Sugar Mixture	0	0	0	0	0
	0	0	0	0	0

SUBSTITUTE SHEET



TABLE 8

Treatment	Percent Mortality After Indicated Days				
	1	2	3	7	10
29756	44	88	100		
	56	100			
	68	88	96	100	
	68	88	100		
29757	88	100			
	92	96	100		
	92	92	100		
	88	96	96	100	
29758	84	84	92	100	
	80	92	100		
	76	88	96	100	
	88	92	96	100	
29759	84	92	100		
	96	96	100		
	100				
	96	100			
29778	100				
	92	96	100		
	96	96	100		
	96	100			
Standard	100				
	92	96	96	96	100
	100				
	100				
Cornmeal-powdered Sugar Mixture	0	0	0	0	0
	0	0	0	0	0
	0	0	0	0	0
	0	0	0	0	0

SUBSTITUTE SHEET



-39-

TABLE 9

Chemical Code No.	Lethal Concentration (ppm)	
	LC-50	LC-90
29756	0.0167	0.0486
29757	.0054	.0200
29758	.0067	.0274
29759	.0029	.0081
29778	.0209	.0374
Control	.0016	.0069

SUBSTITUTE SHEET



-40-

TABLE 10

Chemical Code No.	% Mortality After 24 hrs., Conc. (ppm)				
	10	1.0	0.1	0.01	0.001
29756	100	96	0		
29757	100	96	34		
29758	100	82	78		
29759	100	16	0		
29778	44	22	0		
Control				98	38


SUBSTITUTE SHEET



CLAIMS

1. A composition for the control of a population of arthropods which comprises

(1) An effective amount of a toxicant substance or mixture of substances of the formula



wherein R_f is a fluoroaliphatic radical containing up to 20 carbon atoms, A is a structurally compatible residue, or agriculturally acceptable salts of said substance or substances, and

(2) A bait component or an agriculturally acceptable carrier.

2. A composition according to Claim 1 wherein R_f contains from 5-12 carbon atoms.

3. A composition according to Claim 1 wherein A is NR_1R_2 and wherein R_1 and R_2 are selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, aralkyl, aroyl, acyl, cycloalkyl, cycloalkenyl, cycloalkynyl, a heterocyclic ring containing atoms selected from the group consisting of C, O, N and S, hydroxyalkyl, haloalkyl, aminoalkyl, carboxyalkyl, carboxyalkyl salts, esters and amides or a group of the structure $-(C_xH_{2x}O)_n-$ $(C_yH_{2y}O)_m$ R_3 in which $n + m$ are 1-20, x and y are 1-4 and R_3 is selected from the same group as R_1 and R_2 .

4. A composition according to Claim 1 wherein A is NR_1R_2 and N, R_1 and R_2 taken together form a ring from the atoms of which are selected from the group consisting of C, O, N and S.

5. A composition according to Claim 1 wherein A is OR_4 and R_4 is H, aryl, a heterocyclic ring, and alkaline earth, alkali metal, organic amine or ammonium cation.

SUBSTITUTE SHEET




-42-

6. A composition according to Claim 3 wherein A is $-\text{NH}_2$.


7. A composition according to Claim 3 wherein R_1 is H or alkyl and R_2 is H, alkyl, alkenyl, alkynyl or aryl.

8. A composition according to Claim 3 wherein R_1 is H or alkyl, R_2 is $-(\text{C}_x\text{H}_{2x}\text{O})_n(\text{C}_y\text{H}_{2y}\text{O})_m$, R_3 is H or alkyl, $n + m = 1-20$, and x and $y = 1-4$.


9. A composition according to Claim 1 of the formula



10. A composition according to Claim 3 of the formula



11. A composition according to Claim 4 of the formula




SUBSTITUTE SHEET




- 43 -


12. A composition according to Claim 5 of the formula




13. A composition according to Claim 6 of the formula




14. A composition according to Claim 7 of the composition




-44-



15. A composition according to Claim 8 of the formula



16. A method for the control of a population of arthropods which comprises treating said arthropods with an effective amount of a toxicant substance or mixture of substances of the formula



SUBSTITUTE SHEET



- 45 -

wherein R_f is a fluoroaliphatic radical containing up to 20 carbon atoms, and A is a structurally compatible residue, agriculturally acceptable salt thereof.

17. A method according to Claim 16 wherein R_f contains 5-12 carbon atoms.

18. A method according to Claim 16 wherein A is NR_1R_2 and wherein R_1 and R_2 are selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, aralkyl, aroyl, acyl, cycloalkyl, cycloalkenyl, cycloalkynyl, a heterocyclic ring containing atoms from the group consisting of C, O, N and S, hydroxyalkyl, haloalkyl, aminoalkyl, carboxyalkyl, carboxyalkyl salts, esters, and amides, or a group of the structure $(C_xH_{2x}O)_n(C_yH_{2y}O)_mR_3$ in which $n + m = 1-20$, x and y = 1-4, and R_3 is selected from the same group as R_1 and R_2 .

19. A method according to Claim 16 wherein A is NR_1R_2 and N, R_1 and R_2 taken together form a ring the atoms of which are selected from the group consisting of C, O, N and S.


20. A method according to Claim 16 wherein A is OR_4 wherein R_4 is H, aryl, a heterocyclic ring, an alkaline earth, alkali metal, organic amine or ammonium cation.

21. A method according to Claim 18 wherein A is NH_2 .

22. A method according to Claim 18 wherein R_1 is H, or alkyl and R_2 is H, alkyl, alkenyl, alkynyl, or aryl.

23. A method according to Claim 18 wherein R_1 is H, or alkyl, R_2 is $-(C_xH_{2x}O)_n(C_yH_{2y}O)_m-R_3$, R_3 is H, or alkyl, $n + m = 1-20$, and x and y = 1-4.







24. A method according to Claim 16 wherein the toxicant substance has the formula




SUBSTITUTE SHEET





25. A method according to Claim 18 wherein the toxicant substance or mixture of substances is selected from the group consisting of






26. A method according to Claim 19 wherein the toxicant substance has a formula







27. A method according to Claim 20 wherein the toxicant substance or mixture of substances is selected from the group consisting of










-47-



28. A method according to Claim 21 wherein the toxicant substance or mixture of substances is selected from the group consisting of





29. A method according to Claim 22 wherein the toxicant substance or mixture of substances is selected from the group consisting of












SUBSTITUTE SHEET





30. A method according to Claim 23 wherein the toxicant substance or mixture of substances is selected from the group consisting of



31. The method of Claim 16 wherein said pests are fire ants.

32. The method of Claim 31 wherein said compound is formulated with a suitable bait.

33. The method of Claim 32 wherein said bait is selected from the group consisting of soybean oil, honey-water or other suitable food attractant.

-49 -

34. The method of Claim 33 wherein said bait is impregnated onto a carrier.


35. The method of Claim 34 wherein said bait carrier is selected from the group consisting of corncob grits, pregel defatted corn grits or other granular materials that are compatible with the food attractant and readily fall to the ground surface where fire ants forage.

36. The method of Claim 16 wherein said compound or mixture of compounds is in a sprayable solution, emulsion, or dispersion.

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. **PCT/US 84/00028**

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ³		
According to ⁹ International Patent Classification (IPC) or to both National Classification and IPC INT. CL. ⁹ A61K 31/74; A01N 37/00; 37/02; <u>CONT.</u>		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
US	424/78; 211; 244; 245; 246; 248.5; 249; 263; 267; 269; 270; 272; 273R; 274; 278; 289; 295; 298; 300; 303; 305; 307; 311; 312; 317; 318; and 321	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁵		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
Y	US, A, 4,176,176, Published, 27 November, 1979, Cella et al	1-3,5-8, 12-15
Y	US, A, 2,915,554, Published, 01 December, 1959, Ahlbrecht et al.	1-3,8,15
Y	US, A, 2,803,656, Published, 20 August, 1957 Ahlbercht et al.	1-3,8,15
X	US, A, 3,991,209, Published, 09 November, 1976 Forsyth et al.	1,3,6,7,13
Y	US, A, 3,734,962, Published, 22 May, 1973 Niederdrum et al.	1-3,8,15
Y	GB, A, 738,758, Published 19 October, 1955 Farbenfabriken Bayer, Aktien-Gesellschaft, LeverKusen-Bayerwerk, Germany	1-3,7,16-18, 22, 31-36
<p>¹⁵ * Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ³ 17 April 1984	Date of Mailing of this International Search Report ³ 30 APR 1984	
International Searching Authority ¹ ISA/US	 Allen Jay Robinson	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No ¹⁸
Y	JP, B, 48-32650, Published, 10 August 1973 Kumiai Chemical Ind. Co. Ltd.	1,3,6,7, 13,16,18, 21,22,28, 31-36
Y	US, A, 3,398,182, Published, 20 August 1968 Guenthner et al.	1-3,15
Y	US, A, 2,759,019, Published, 14 August 1956, Brown et al	1,2
A	US, A, 2,732,378, Published, 24 January 1956, Cavallito	1-36
A	US, A, 2,346,612, Published, 11 April 1944, Rothrock	1-36
Y	US, A, 3,220,921, Published, 30 November 1965, Greenbaum et al	1-36
A	US, A, 4,068,000, Published, 10 January 1978, Edwards	1-36
A	US, A, 4,107,332, Published, 15 August 1978, Chan et al	1-36
A	US, A, 3,248,419, Published, 26 April 1966, Hauptschein et al	1-36
Y	US, A, 3,795,743, Published, 05 March 1974, Okuda et al	1,3,6,7, 16,18,21, 22,36
A	US, A, 4,101,468, Published, 18 July 1978, Perrey et al	1-36
Y	JP, B, 48-34208, Published, 19 October 1973, Kumiai Chemical Ind. Co. Ltd.	1,3,7,10 14,15,16, 18,22,25, 29,30,36

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category*	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No ¹⁸
A	US, A, 4,281,011, Published, 28 July 1981, Siegler et al	1-36
Y	JP, A, 57-156407, Published, 27 September 1982, Sumitomo Chemical KK	1,3,7,16, 18,22,31-36
X	JP, A, 57-156407, Published, 27 September 1982, Sumitomo Chemical KK	1,3,7
Y	US, A, 3,803,199, Published, 09 April 1974, Voss et al	1-3,5, 16-18,20, 31-36
Y	US, A, 3,818,102, Published, 18 June 1974, Partos	1,2,5,16, 18,20, 31-36
Y	JP, B, 49-35415, Published, 21 September 1974, Kumiai Chem. Ind. Co. Ltd.	1,5,12, 16,20,27, 31-36
Y	JP, B, 48-18453, Published, 06 June 1973, Kumiai Chemical Ind. Co. Ltd.	1,3,10, 16,18,25, 31-36
Y	GB, A, 873,590, Published, 26 July 1961, Farben Fabriken Bayer Aktien Gesellschaft Lever Kusen Bayerwerk, Germany	1-3,7, 16-18,22 31-36
Y	JP, B, 48-34213, Published, 19 October 1973, Kumiai Chemi- cal Ind. Co., Ltd.	1, 3, 10 16, 18, 25, 31-36

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹⁰

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. Claim numbers because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. Claim numbers, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

VI. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ¹¹

This International Searching Authority found multiple inventions in this international application as follows:

SEE PCT/ISA/206 mailed 06 March 1984

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

The additional search fees were accompanied by applicant's protest.

No protest accompanied the payment of additional search fees.

PCT/ISA/210 SUPPLEMENTAL SHEET FOR

1. Classification of Subject matter

PCT/US84/00028

1. CLASSIFICATION OF SUBJECT MATTER

37/06; 37/08; 41/02; 41/06; 43/00; 43/02; 43/36;
43/40; 43/48; 43/64; 43/72; 43/76; 43/78; 43/80;
43/84; 43/86; 43/88; 43/50; 43/56; 47/10; 55/02;
and 57/00

US CL. 424/78; 211; 244; 245; 246; 248.5; 249; 263; 267;
269; 270; 272; 273R; 274; 278; 289; 295; 298; 300;
303; 305; 307; 311; 312; 317; 318; and 321