



US009752134B2

(12) **United States Patent**
Landvik et al.

(10) **Patent No.:** **US 9,752,134 B2**
(45) **Date of Patent:** ***Sep. 5, 2017**

(54) **POLYPEPTIDES HAVING GLUCOAMYLASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME**

(71) Applicants: **NOVOZYMES A/S**, Bagsvaerd (DK); **NOVOZYMES NORTH AMERICA, INC.**, Franklinton, NC (US)

(72) Inventors: **Sara Landvik**, Bagsvaerd (DK); **Marc Dominique Morant**, Bagsvaerd (DK); **Keiichi Ayabe**, Chiba (JP); **Guillermo Coward-Kelly**, Franklinton, NC (US)

(73) Assignees: **NOVOZYMES NORTH AMERICA, INC.**, Franklinton, NC (US); **NOVOZYMES A/S**, Bagsvaerd (DK)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **14/934,911**

(22) Filed: **Nov. 6, 2015**

(65) **Prior Publication Data**

US 2016/0053245 A1 Feb. 25, 2016

Related U.S. Application Data

(60) Division of application No. 14/790,411, filed on Jul. 2, 2015, which is a division of application No. 13/916,042, filed on Jun. 12, 2013, now Pat. No. 9,102,928, which is a continuation of application No. 13/500,325, filed as application No. PCT/US2010/058424 on Nov. 30, 2010, now Pat. No. 8,557,541.

(60) Provisional application No. 61/406,741, filed on Oct. 26, 2010, provisional application No. 61/265,526, filed on Dec. 1, 2009.

(51) **Int. Cl.**
C12N 9/24 (2006.01)
C12N 9/34 (2006.01)
C12P 7/10 (2006.01)
C12P 19/02 (2006.01)

C12P 19/14 (2006.01)
C12N 15/81 (2006.01)

(52) **U.S. Cl.**
CPC **C12N 9/2428** (2013.01); **C12N 15/81** (2013.01); **C12P 7/10** (2013.01); **C12P 19/02** (2013.01); **C12P 19/14** (2013.01); **C12Y 302/01003** (2013.01); **Y02E 50/16** (2013.01); **Y02E 50/17** (2013.01)

(58) **Field of Classification Search**
CPC C12N 9/24
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,326,548 B2 2/2008 Udagawa
7,871,800 B2 1/2011 Landvik
8,148,127 B2 4/2012 Udagawa

FOREIGN PATENT DOCUMENTS

WO 2004056981 A2 7/2004
WO 2006/069289 A2 6/2006
WO 2006/069290 A2 6/2006
WO 2007/134207 A2 11/2007
WO 2007/144424 A2 12/2007
WO 2008074884 A2 6/2008

OTHER PUBLICATIONS

Horst et al, 1960, Archiv Fur Mikrobiol 35 (3), 258-278—English Translation filed.

Wang et al, 2002, Acta Microbial Sinica 42(2), 220-225 (English Abstract Only).

Primary Examiner — Maryam Monshipouri

(74) *Attorney, Agent, or Firm* — David Fazzolare

(57) **ABSTRACT**

The present invention relates to polypeptides having glucoamylase activity and polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

8 Claims, No Drawings

**POLYPEPTIDES HAVING GLUCOAMYLASE
ACTIVITY AND POLYNUCLEOTIDES
ENCODING SAME**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This application is a division of U.S. application Ser. No. 14/790,411 filed Jul. 2, 2015, now pending, which is a division of U.S. application Ser. No. 13/916,042 filed Jun. 12, 2013, now U.S. Pat. No. 9,102,928, which is continuation of U.S. application Ser. No. 13/500,325 filed May 3, 2012, now U.S. Pat. No. 8,557,541, which is a 35 U.S.C. 371 national application of PCT/US2010/058424 filed Nov. 30, 2010, which claims priority or the benefit under 35 U.S.C. 119 of U.S. provisional application Nos. 61/265,526 and 61/406,741 filed Dec. 1, 2009 and Oct. 26, 2010, respectively, the contents of which are fully incorporated herein by reference.

REFERENCE TO A SEQUENCE LISTING

This application contains a Sequence Listing in computer readable form, which is incorporated herein by reference.

REFERENCE TO A DEPOSIT OF BIOLOGICAL
MATERIAL

This application contains a reference to a deposit of biological material, which deposit is incorporated herein by reference. For complete information see last paragraph of the description.

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to polypeptides having glucoamylase activity and polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods for producing and using the polypeptides, and to the use of glucoamylases of the invention for starch conversion to producing fermentation products, such as ethanol, and syrups, such as glucose. The invention also relates to a composition comprising a glucoamylase of the invention.

Description of the Related Art

Glucoamylase (1,4-alpha-D-glucan glucohydrolase, EC 3.2.1.3) is an enzyme, which catalyzes the release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules. Glucoamylases are produced by several filamentous fungi and yeast, with those from *Aspergillus* being commercially most important.

Commercially, glucoamylases are used to convert starchy material, which is already partially hydrolyzed by an alpha-amylase, to glucose. The glucose may then be converted directly or indirectly into a fermentation product using a fermenting organism. Examples of commercial fermentation products include alcohols (e.g., ethanol, methanol, butanol, 1,3-propanediol); organic acids (e.g., citric acid, acetic acid, itaconic acid, lactic acid, gluconic acid, gluconate, lactic acid, succinic acid, 2,5-diketo-D-gluconic acid); ketones (e.g., acetone); amino acids (e.g., glutamic acid); gases (e.g., H₂ and CO₂), and more complex compounds, including, for example, antibiotics (e.g., penicillin and tetracycline); enzymes; vitamins (e.g., riboflavin, B₁₂, beta-carotene); hormones, and other compounds which are difficult to

produce synthetically. Fermentation processes are also commonly used in the consumable alcohol (e.g., beer and wine), dairy (e.g., in the production of yogurt and cheese) industries.

The end product may also be a syrup. For instance, the end product may be glucose, but may also be converted, e.g., by a glucose isomerase to fructose or a mixture composed almost equally of glucose and fructose. This mixture, or a mixture further enriched with fructose, is the most commonly used high fructose corn syrup (HFCS) commercialized throughout the world.

It is an object of the present invention to provide polypeptides having glucoamylase activity and polynucleotides encoding the polypeptides and which provide a high yield in fermentation product production processes, such as ethanol production processes, including one-step ethanol fermentation processes from un-gelatinized raw (or uncooked) starch.

Uniprot: B0CVJ1 discloses a polypeptide from *Laccaria bicolor* and WO 2006/069289 describes a glucoamylase from *Trametes cingulata*.

SUMMARY OF THE INVENTION

Polypeptides produced by the fungus *Gloeophyllum* and having glucoamylase activity have been identified and characterized. More particularly the *Gloeophyllum* sp. is selected from the group consisting of *G. abietinum*, *G. sepiarium*, and *G. trabeum*.

Accordingly, the present invention relates in a first aspect to an isolated polypeptide having glucoamylase activity, selected from the group consisting of: (a) a polypeptide comprising an amino acid sequence having preferably at least 82%, more preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18; (b) a polypeptide comprising an amino acid sequence having preferably at least 82%, more preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity to the catalytic domain shown as amino acids 19 to 474 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14 or amino acids 19 to 471 of SEQ ID NO: 16 or of SEQ ID NO: 18; c) a polypeptide encoded by a polynucleotide that hybridizes under preferably at least medium-high stringency conditions, and most preferably at least high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ

ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, or (iii) a full-length complementary strand of (i) or (ii); (d) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having preferably at least 82%, more preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17; and (e) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18.

The present invention relates in a second aspect to an isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of the first aspect.

In further aspects the invention relates to a nucleic acid construct, a recombinant expression vector, a recombinant host cell, a transgenic plant, a plant part or plant cell comprising the polynucleotide of the second aspect.

In still further aspects the invention relates to a method of producing the polypeptide, uses of the polypeptide and a composition comprising an alpha-amylase and the polypeptide.

DEFINITIONS

Glucoamylase: The term glucoamylase (1,4-alpha-D-glucan glucohydrolase, EC 3.2.1.3) is defined as an enzyme, which catalyzes the release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules. For purposes of the present invention, glucoamylase activity is determined according to the procedure described in the 'Materials & Methods'-section below.

The polypeptides of the present invention have at least 20%, preferably at least 40%, preferably at least 45%, more preferably at least 50%, more preferably at least 55%, more preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 100% of the glucoamylase activity of the mature polypeptide of SEQ ID NO: 2 or a homologous sequence thereof, SEQ ID NO: 4 or a homologous sequence thereof, SEQ ID NO: 6 or a homologous sequence thereof, SEQ ID NO: 8 or a homologous sequence thereof, SEQ ID NO: 10 or a homologous sequence thereof, SEQ ID NO: 12 or a homologous sequence thereof, SEQ ID NO: 14 or a homologous sequence thereof, or SEQ ID NO: 18 or a homologous sequence thereof.

Isolated polypeptide: The term "isolated polypeptide" as used herein refers to a polypeptide that is isolated from a source. Preferably, the polypeptide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by SDS-PAGE.

Substantially pure polypeptide: The term "substantially pure polypeptide" denotes herein a polypeptide preparation that contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polypeptide material with which it is natively or recombinantly associated. It is, therefore, preferred that the substantially pure polypeptide is at least 92% pure, preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, more preferably at least 98% pure, even more preferably at least 99%, most preferably at least 99.5% pure, and even most preferably 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides of the present invention are preferably in a substantially pure form, i.e., that the polypeptide preparation is essentially free of other polypeptide material with which it is natively or recombinantly associated. This can be accomplished, for example, by preparing the polypeptide by well-known recombinant methods or by classical purification methods.

Mature polypeptide: The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. In one aspect, the mature polypeptide is amino acids 18 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14 or amino acids 18 to 576 of SEQ ID NO: 16, or of SEQ ID NO: 18 based on the SignalP program (Nielsen et al., 1997, *Protein Engineering* 10: 1-6) that predicts amino acids 1 to 17 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16 and SEQ ID NO: 18 are a signal peptide. The sequence defined by amino acids 19 to 474 (particularly 19 to 471) of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14 or amino acids 19 to 471 of SEQ ID NO: 16 or of SEQ ID NO: 18 is the catalytic domain. The sequence defined by amino acids 480 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14 or amino acids 483 to 576 of SEQ ID NO: 16 or SEQ ID NO: 18 is a starch binding domain.

Mature polypeptide coding sequence: The term "mature polypeptide coding sequence" is defined herein as a nucleotide sequence that encodes a mature polypeptide having glucoamylase activity. Preferably, the mature polypeptide coding sequence is nucleotides 52 to 1719 of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, or nucleotides 52 to 1728 of SEQ ID NO: 15 or of SEQ ID NO: 17.

Identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity".

For purposes of the present invention, the degree of identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *Trends in Genetics* 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substi-

tution matrix. The output of Needle labeled “longest identity” (obtained using the `-nobrief` option) is used as the percent identity and is calculated as follows:

$$\frac{(\text{Identical Residues} \times 100)}{(\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})}$$

For purposes of the present invention, the degree of identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *supra*), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled “longest identity” (obtained using the `-nobrief` option) is used as the percent identity and is calculated as follows:

$$\frac{(\text{Identical Deoxyribonucleotides} \times 100)}{(\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})}$$

Homologous sequence: The term “homologous sequence” is defined herein as a nucleotide sequence/polypeptide sequence having a % identity to the mature polypeptide encoding part of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 or to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18, respectively, of at least 82%, preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, or even at least 99%.

Polypeptide fragment: The term “polypeptide fragment” is defined herein as a polypeptide having one or more (several) amino acids deleted from the amino and/or carboxyl terminus of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18; or a homologous sequence thereof; wherein the fragment has glucoamylase activity. Preferably, a fragment contains at least 500 amino acid residues, more preferably at least 450 amino acid, and most preferably at least 400 amino acid residues, of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14 or amino acids 19 to 471 of SEQ ID NO: 16 or of SEQ ID NO: 18 which comprises the catalytic domain of the polypeptide of the invention.

Subsequence: The term “subsequence” is defined herein as a nucleotide sequence having one or more (several) nucleotides deleted from the 5' and/or 3' end of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17; or a homologous sequence thereof; wherein the

subsequence encodes a polypeptide fragment having glucoamylase activity. Preferably, a subsequence contains at least 1500 nucleotides, more preferably at least 1400 nucleotides, and most preferably at least 1200 nucleotides of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 or a homologous sequence thereof.

Allelic variant: The term “allelic variant” denotes herein any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

Isolated polynucleotide: The term “isolated polynucleotide” as used herein refers to a polynucleotide that is isolated from a source. Preferably, the polynucleotide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by agarose electrophoresis.

Substantially pure polynucleotide: The term “substantially pure polynucleotide” as used herein refers to a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered protein production systems. Thus, a substantially pure polynucleotide contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polynucleotide material with which it is natively or recombinantly associated. A substantially pure polynucleotide may, however, include naturally occurring 5' and 3' untranslated regions, such as promoters and terminators. It is preferred that the substantially pure polynucleotide is at least 90% pure, preferably at least 92% pure, more preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, even more preferably at least 98% pure, most preferably at least 99%, and even most preferably at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form, i.e., that the polynucleotide preparation is essentially free of other polynucleotide material with which it is natively or recombinantly associated. The polynucleotides may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

Coding sequence: When used herein the term “coding sequence” means a nucleotide sequence, which directly specifies the amino acid sequence of its protein product. The boundaries of the coding sequence are generally determined by an open reading frame, which usually begins with the ATG start codon or alternative start codons such as GTG and TTG and ends with a stop codon such as TAA, TAG, and TGA. The coding sequence may be a DNA, cDNA, synthetic, or recombinant nucleotide sequence.

cDNA: The term “cDNA” is defined herein as a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic cell. cDNA lacks intron sequences that are usually present in the corresponding genomic DNA. The initial,

primary RNA transcript is a precursor to mRNA that is processed through a series of steps before appearing as mature spliced mRNA. These steps include the removal of intron sequences by a process called splicing. cDNA derived from mRNA lacks, therefore, any intron sequences.

Nucleic acid construct: The term “nucleic acid construct” as used herein refers to a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or which is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic. The term nucleic acid construct is synonymous with the term “expression cassette” when the nucleic acid construct contains the control sequences required for expression of a coding sequence of the present invention.

Control sequences: The term “control sequences” is defined herein to include all components necessary for the expression of a polynucleotide encoding a polypeptide of the present invention. Each control sequence may be native or foreign to the nucleotide sequence encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleotide sequence encoding a polypeptide.

Operably linked: The term “operably linked” denotes herein a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of the polynucleotide sequence such that the control sequence directs the expression of the coding sequence of a polypeptide.

Expression: The term “expression” includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

Expression vector: The term “expression vector” is defined herein as a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide of the present invention and is operably linked to additional nucleotides that provide for its expression.

Host cell: The term “host cell”, as used herein, includes any cell type that is susceptible to transformation, transfection, transduction, and the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention.

Modification: The term “modification” means herein any chemical modification of the polypeptide consisting of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18; or a homologous sequence thereof; as well as genetic manipulation of the DNA encoding such a polypeptide. The modification can be a substitution, a deletion and/or an insertion of one or more (several) amino acids as well as replacements of one or more (several) amino acid side chains.

Variant: When used herein, the term “variant” means a polypeptide having glucoamylase activity comprising an alteration, i.e., a substitution, insertion, and/or deletion of one or more (several) amino acid residues at one or more (several) positions. A substitution means a replacement of an amino acid occupying a position with a different amino acid;

a deletion means removal of an amino acid occupying a position; and an insertion means adding 1-3 amino acids adjacent to an amino acid occupying a position.

DETAILED DESCRIPTION OF THE INVENTION

Polypeptides Having Glucoamylase Activity

In a first aspect, the present invention relates to isolated polypeptides comprising an amino acid sequence having a degree of identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18 of preferably at least 82%, more preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity, which have glucoamylase activity (hereinafter “homologous polypeptides”). Preferably, the homologous polypeptides have an amino acid sequence that differs by ten amino acids, preferably by five amino acids, more preferably by four amino acids, even more preferably by three amino acids, most preferably by two amino acids, and even most preferably by one amino acid from the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18.

A polypeptide of the present invention preferably comprises the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18 or an allelic variant thereof; or a fragment thereof having glucoamylase activity. Preferably, the polypeptide comprises the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18. In another preferred aspect, the polypeptide comprises the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18 or an allelic variant thereof; or a fragment thereof having glucoamylase activity. In another preferred aspect, the polypeptide consists of the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18 or an allelic variant thereof; or a fragment thereof having glucoamylase activity. In another preferred aspect, the polypeptide consists of the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18.

In a second aspect, the present invention relates to isolated polypeptides having glucoamylase activity that are encoded by polynucleotides that hybridize under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID

NO: 17, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, (iii) a subsequence of (i) or (ii), or (iv) a full-length complementary strand of (i), (ii), or (iii) (J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, N.Y.). A subsequence of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 contains at least 100 contiguous nucleotides or preferably at least 200 contiguous nucleotides. Moreover, the subsequence may encode a polypeptide fragment having glucoamylase activity. Preferably, the complementary strand is the full-length complementary strand of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17.

The nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17; or a subsequence thereof; as well as the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18; or a fragment thereof; may be used to design nucleic acid probes to identify and clone DNA encoding polypeptides having glucoamylase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 14, preferably at least 25, more preferably at least 35, and most preferably at least 70 nucleotides in length. It is, however, preferred that the nucleic acid probe is at least 100 nucleotides in length. For example, the nucleic acid probe may be at least 200 nucleotides, preferably at least 300 nucleotides, more preferably at least 400 nucleotides, or most preferably at least 500 nucleotides in length. Even longer probes may be used, e.g., nucleic acid probes that are preferably at least 600 nucleotides, more preferably at least 700 nucleotides, even more preferably at least 800 nucleotides, or most preferably at least 900 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ³³P, ³²P, ³H, ³⁵S, biotin, or avidin). Such probes are encompassed by the present invention.

A genomic DNA or cDNA library prepared from such other strains may, therefore, be screened for DNA that hybridizes with the probes described above and encodes a polypeptide having glucoamylase activity. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that is homologous with SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17; or a subsequence thereof; the carrier material is preferably used in a Southern blot.

For purposes of the present invention, hybridization indicates that the nucleotide sequence hybridizes to a labeled

nucleic acid probe corresponding to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17; the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17; its full-length complementary strand; or a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film.

Preferably, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 2, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is the polynucleotide sequence contained in plasmid in *E. coli* strain DSM 23222, wherein the polynucleotide sequence thereof encodes a polypeptide having glucoamylase activity. In another preferred aspect, the nucleic acid probe is the mature polypeptide coding region contained in plasmid in *E. coli* strain DSM 23222.

Preferably, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17. In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17.

For long probes of at least 100 nucleotides in length, very low to very high stringency conditions are defined as pre-hybridization and hybridization at 42° C. in 5×SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures for 12 to 24 hours optimally.

For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2×SSC, 0.2% SDS preferably at 45° C. (very low stringency), more preferably at 50° C. (low stringency), more preferably at 55° C. (medium stringency), more preferably at 60° C. (medium-high stringency), even more preferably at 65° C. (high stringency), and most preferably at 70° C. (very high stringency).

For short probes that are about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at about 5° C. to about 10° C. below the calculated T_m using the calculation according to Bolton and McCarthy (1962, Proceedings of the National Academy of Sciences USA 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1×Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures for 12 to 24 hours optimally.

For short probes that are about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in

6xSSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6xSSC at 5° C. to 10° C. below the calculated T_m .

In a third aspect, the present invention relates to isolated polypeptides having glucoamylase activity encoded by polynucleotides comprising or consisting of nucleotide sequences that have a degree of identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 of at least 82%, more preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99%, which encode an active polypeptide. See polynucleotide section herein.

The present invention also relates to artificial variants comprising a substitution, deletion, and/or insertion of one or more (or several) amino acids of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18; or a homologous sequence thereof. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R. L. Hill, 1979, *In, The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline, and alpha-methyl serine) may be substituted for amino acid residues of a wild-type polypeptide. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, and preferably, are commercially available, and include pipercolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the

thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

Essential amino acids in the parent polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for glucoamylase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, *Science* 255: 306-312; Smith et al., 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver et al., 1992, *FEBS Lett.* 309: 59-64. The identities of essential amino acids can also be inferred from analysis of identities with polypeptides that are related to a polypeptide according to the invention.

Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, *Biochem.* 30: 10832-10837; U.S. Pat. No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire et al., 1986, *Gene* 46: 145; Ner et al., 1988, *DNA* 7: 127).

Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

The total number of amino acid substitutions, deletions and/or insertions of the mature polypeptide, such as amino acids 18 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14 or amino acids 18 to 576 of SEQ ID NO: 16 or SEQ ID NO: 18, or the catalytic domain, such as amino acids 19 to 474 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14 or amino acids 19 to 471 of SEQ ID NO: 16 or of SEQ ID NO: 18, is 10, preferably 9, more preferably 8, more preferably 7, more preferably at most 6, more preferably 5, more preferably 4, even more preferably 3, most preferably 2, and even most preferably 1.

Sources of Polypeptides Having Glucoamylase Activity

A polypeptide of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a nucleotide sequence is produced by the source or by a strain in which the nucleotide sequence from the source has been inserted. Preferably, the polypeptide obtained from a given source is secreted extracellularly.

A polypeptide having glucoamylase activity of the present invention may also be bacterial polypeptide, or a yeast polypeptide, or more preferably a filamentous fungal polypeptide such as an *Acremonium*, *Agaricus*, *Alternaria*, *Artomyces*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Gloeophyllum*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocalimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotriconympha*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolyopocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* polypeptide having glucoamylase activity.

In a more preferred aspect, the polypeptide is a *Gloeophyllum* sp. polypeptide, such as a *Gloeophyllum sepiarium*, *Gloeophyllum trabeum*, or *Gloeophyllum abietinum* polypeptide having glucoamylase activity. It will be understood that for the aforementioned species the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen and Zellkulturen GmbH (DSMZ), Centraalbureau voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

Furthermore, such polypeptides may be identified and obtained from other sources including microorganisms isolated from nature (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms from natural habitats are well known in the art. The polynucleotide may then be obtained by similarly screening a genomic or cDNA library of such a microorganism. Once a polynucleotide sequence encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are well known to those of ordinary skill in the art (see, e.g., Sambrook et al., 1989, supra).

Polypeptides of the present invention also include fused polypeptides or cleavable fusion polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by fusing a nucleotide sequence (or a portion thereof) encoding another polypeptide to a nucleotide sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fused polypeptide is under control of the same promoter(s) and terminator.

A fusion polypeptide can further comprise a cleavage site. Upon secretion of the fusion protein, the site is cleaved releasing the polypeptide having glucoamylase activity from the fusion protein. Examples of cleavage sites include, but are not limited to, a Kex2 site that encodes the dipeptide Lys-Arg (Martin et al., 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-76; Svetina et al., 2000, *J. Biotechnol.* 76: 245-251; Rasmussen-Wilson et al., 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward et al., 1995, *Biotechnology* 13: 498-

503; and Contreras et al., 1991, *Biotechnology* 9: 378-381), an Ile-(Glu or Asp)-Gly-Arg site, which is cleaved by a Factor Xa protease after the arginine residue (Eaton et al., 1986, *Biochem.* 25: 505-512); an Asp-Asp-Asp-Asp-Lys site, which is cleaved by an enterokinase after the lysine (Collins-Racie et al., 1995, *Biotechnology* 13: 982-987); a His-Tyr-Glu site or His-Tyr-Asp site, which is cleaved by Genenase I (Carter et al., 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248); a Leu-Val-Pro-Arg-Gly-Ser site, which is cleaved by thrombin after the Arg (Stevens, 2003, *Drug Discovery World* 4: 35-48); a Glu-Asn-Leu-Tyr-Phe-Gln-Gly site, which is cleaved by TEV protease after the Gln (Stevens, 2003, supra); and a Leu-Glu-Val-Leu-Phe-Gln-Gly-Pro site, which is cleaved by a genetically engineered form of human rhinovirus 3C protease after the Gln (Stevens, 2003, supra).

Polynucleotides

The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that encode polypeptides having glucoamylase activity of the present invention.

Preferably, the nucleotide sequence comprises or consists of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17. In another preferred aspect, the nucleotide sequence comprises or consists of the sequence contained in plasmid which is contained in *E. coli* DSM 23222. In another preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence of SEQ ID NO: 1. In another more preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence contained in plasmid contained in *E. coli* DSM 23222. The present invention also encompasses nucleotide sequences that encode polypeptides comprising or consisting of the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18 or the mature polypeptides thereof, which differ from SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 or the mature polypeptide coding sequence thereof by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 that encodes fragments of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18 that have glucoamylase activity.

The present invention also relates to mutant polynucleotides comprising or consisting of at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, in which the mutant nucleotide sequence encodes the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18.

The techniques used to isolate or clone a polynucleotide encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the polynucleotides of the present invention from such genomic DNA can be effected, e.g., by using the well known polymerase chain

reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, e.g., Innis et al., 1990, PCR: A Guide to Methods and Application, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligated activated transcription (LAT) and nucleotide sequence-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Penicillium*, or another or related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the nucleotide sequence.

The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that have a degree of identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17 of preferably at least 82%, more preferably at least 83%, more preferably at least 84%, more preferably at least 85%, more preferably at least 86%, more preferably at least 87%, more preferably at least 88%, more preferably at least 89%, more preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity, which encode an active polypeptide.

Modification of a nucleotide sequence encoding a polypeptide of the present invention may be necessary for the synthesis of polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, e.g., artificial variants that differ in specific activity, thermostability, pH optimum, or the like. The variant sequence may be constructed on the basis of the nucleotide sequence presented as the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, e.g., a subsequence thereof, and/or by introduction of nucleotide substitutions that do not give rise to another amino acid sequence of the polypeptide encoded by the nucleotide sequence, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford et al., 1991, *Protein Expression and Purification* 2: 95-107.

It will be apparent to those skilled in the art that such substitutions can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by an isolated polynucleotide of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, e.g., Cunningham and Wells, 1989, supra). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for glucoamylase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional

labeling (see, e.g., de Vos et al., 1992, supra; Smith et al., 1992, supra; Wlodaver et al., 1992, supra).

The present invention also relates to isolated polynucleotides encoding polypeptides of the present invention, which hybridize under very low stringency conditions, preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, or (iii) a full-length complementary strand of (i) or (ii); or allelic variants and subsequences thereof (Sambrook et al., 1989, supra), as defined herein. Preferably, the complementary strand is the full-length complementary strand of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17.

The present invention also relates to isolated polynucleotides obtained by (a) hybridizing a population of DNA under very low, low, medium, medium-high, high, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having glucoamylase activity. Preferably, the complementary strand is the full-length complementary strand of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17.

Hybrid Enzymes

The present invention also relates to hybrid enzymes comprising a catalytic domain having enzyme activity (e.g., starch degrading enzyme activity, such as alpha-amylase, amylopullulanase, beta-amylase, CGTase, glucoamylase, isoamylase, maltogenic amylase, or pullulanase activity), and a carbohydrate-binding module (CBM). The hybrid enzyme may further comprise a linker.

The hybrid may be produced by fusing a first DNA sequence encoding a catalytic domain and a second DNA sequence encoding a carbohydrate-binding module, or the hybrid may be produced as a completely synthetic gene based on knowledge of the amino acid sequences of suitable CBMs, linkers and catalytic domains.

The term "hybrid enzyme" (also referred to as "fusion protein", "hybrid", hybrid polypeptide" or "hybrid protein") is used herein to characterize the hybrid polypeptides of the invention comprising a catalytic module having enzyme activity (e.g., starch degrading enzyme activity, such as alpha-amylase, amylopullulanase, beta-amylase, CGTase, glucoamylase, isoamylase, maltogenic amylase, or pullulanase activity) and a carbohydrate-binding module wherein the catalytic domain and the carbohydrate-binding module are derived from different sources. The term "source"

includes, but is not limited to, a parent enzyme or a variant thereof, e.g., an amylase or glucoamylase, or other catalytic activity comprising a suitable catalytic module and/or a suitable CBM and/or a suitable linker. However the CBM may also be derived from a polypeptide having no catalytic activity. The catalytic domain and the carbohydrate binding module may be derived from the same microbial strain, from strains within the same species, from closely related species or less related organisms. Preferably the catalytic domain and the carbohydrate binding module of the hybrids are derived from different sources, e.g., from different enzymes from the same strain and/or species, or e.g., from strains within different species.

In one aspect the hybrid enzyme comprises the CBM (also known as a carbohydrate binding domain or CBD) according to the invention and a catalytic domain. The catalytic domain is in a particular embodiment a glucoamylase catalytic domain.

Nucleic Acid Constructs

The present invention also relates to nucleic acid constructs comprising an isolated polynucleotide of the present invention operably linked to one or more (several) control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

An isolated polynucleotide encoding a polypeptide of the present invention may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide's sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotide sequences utilizing recombinant DNA methods are well known in the art.

The control sequence may be an appropriate promoter sequence, a nucleotide sequence that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter sequence contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any nucleotide sequence that shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus oryzae* TAKA amylase, *Aspergillus awamori* or *Aspergillus niger* glucoamylase (glaA), *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Rhizomucor miehei* aspartic proteinase, *Rhizomucor miehei* lipase, *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger*

neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase); and mutant, truncated, and hybrid promoters thereof.

In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos et al., 1992, *Yeast* 8: 423-488.

The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleotide sequence encoding the polypeptide. Any terminator that is functional in the host cell of choice may be used in the present invention.

Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos et al., 1992, supra.

The control sequence may also be a suitable leader sequence, a nontranslated region of an mRNA that is important for translation by the host cell. The leader sequence is operably linked to the 5' terminus of the nucleotide sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used in the present invention.

Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3' terminus of the nucleotide sequence and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell of choice may be used in the present invention.

Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, *Aspergillus niger* glucoamylase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Molecular Cellular Biology* 15: 5983-5990.

The control sequence may also be a signal peptide coding sequence that codes for an amino acid sequence linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of

the coding sequence of the nucleotide sequence may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding sequence that is foreign to the coding sequence. The foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, the foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell of choice, i.e., secreted into a culture medium, may be used in the present invention.

Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Aspergillus oryzae* TAKA amylase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, *Humicola lanuginosa* lipase, and *Rhizomucor miehei* aspartic proteinase.

It may also be desirable to add regulatory sequences that allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the lac, tac, xyl and trp operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the nucleotide sequence encoding the polypeptide would be operably linked with the regulatory sequence.

Expression Vectors

The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleic acids and control sequences described herein may be joined together to produce a recombinant expression vector that may include one or more (several) convenient restriction sites to allow for insertion or substitution of the nucleotide sequence encoding the polypeptide at such sites. Alternatively, a polynucleotide sequence of the present invention may be expressed by inserting the nucleotide sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about expression of the nucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids.

The vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

The vectors of the present invention preferably contain one or more (several) selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

Selectable markers for use in a filamentous fungal host cell include, but are not limited to, amdS (acetamidase), argB (ornithine carbamoyltransferase), bar (phosphinothricin acetyltransferase), hph (hygromycin phosphotransferase), niaD (nitrate reductase), pyrG (orotidine-5'-phosphate decarboxylase), sC (sulfate adenylyltransferase), and trpC (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are the amdS and pyrG genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the bar gene of *Streptomyces hygroscopicus*.

The vectors of the present invention preferably contain an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleotide sequences for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which have a high degree of identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleotide sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" is defined herein as a nucleotide sequence that enables a plasmid or vector to replicate in vivo.

Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems et al., 1991, *Gene* 98: 61-67; Cullen et al., 1987, *Nucleic Acids Research* 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and

construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of the gene product. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, e.g., Sambrook et al., 1989, supra).

Host Cells

The present invention also relates to recombinant host cells, comprising an isolated polynucleotide of the present invention, which are advantageously used in the recombinant production of the polypeptides. A vector comprising a polynucleotide of the present invention is introduced into a host cell so that the vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, e.g., a prokaryote or a eukaryote.

The prokaryotic host cell may be any gram-positive bacterium or gram-negative bacterium. Gram-positive bacteria include, but are not limited to, *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*, *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, and *Streptomyces*. Gram-negative bacteria include, but are not limited to, *E. coli*, *Campylobacter*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, and *Ureaplasma*.

The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

Preferably, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by Hawksworth et al., *In, Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth et al., 1995, supra, page 171) and all mitospore fungi (Hawksworth et al., 1995, supra).

In a more preferred aspect, the fungal host cell is a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner, F. A., Passmore, S. M., and Davenport, R. R., eds, *Soc. App. Bacteriol. Symposium Series No. 9*, 1980).

In an even more preferred aspect, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

In a most preferred aspect, the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*,

Saccharomyces diastaticus, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* cell. In another most preferred aspect, the yeast host cell is a *Kluyveromyces lactis* cell. In another most preferred aspect, the yeast host cell is a *Yarrowia lipolytica* cell.

In another more preferred aspect, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth et al., 1995, supra). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

In an even more preferred aspect, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filobasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolyocladium*, *Trametes*, or *Trichoderma* cell.

In a most preferred aspect, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger* or *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcocroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis cargeia*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermisporea*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pamicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes versicolor*, *Trametes villosa*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known per se. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 0238023 and Yelton et al., 1984, *Proceedings of the National Academy of Sciences USA* 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Malardier et al., 1989, *Gene* 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J. N. and Simon, M. I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzy-*

mology 194: 182-187, Academic Press, Inc., New York; Ito et al., 1983, *Journal of Bacteriology* 153: 163; and Hinnen et al., 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920.

Methods of Production

The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. Preferably, the cell is of the genus *Gloeophyllum*. In a more preferred aspect, the cell is of the species *Gloeophyllum sepiarium*, *Gloeophyllum trabeum*, or *Gloeophyllum abietinum*.

The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell, as described herein, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell under conditions conducive for production of the polypeptide, wherein the host cell comprises a mutant nucleotide sequence having at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, or SEQ ID NO: 17, wherein the mutant nucleotide sequence encodes a polypeptide that comprises or consists of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18, and (b) recovering the polypeptide.

In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods well known in the art. For example, the cell may be cultivated by shake flask cultivation, and small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted into the medium, it can be recovered from cell lysates.

The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide as described herein.

The resulting polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

The polypeptides of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), elec-

trophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure polypeptides.

Plants

The present invention also relates to plants, e.g., a transgenic plant, plant part, or plant cell, comprising an isolated polynucleotide encoding a polypeptide having glucoamylase activity of the present invention so as to express and produce the polypeptide in recoverable quantities. The polypeptide may be recovered from the plant or plant part. Alternatively, the plant or plant part containing the recombinant polypeptide may be used as such for improving the quality of a food or feed, e.g., improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as *Festuca*, *Lolium*, temperate grass, such as *Agrostis*, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn).

Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, e.g., epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilisation of the invention are also considered plant parts, e.g., embryos, endosperms, aleurone and seeds coats.

Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

The transgenic plant or plant cell expressing a polypeptide of the present invention may be constructed in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more (several) expression constructs encoding a polypeptide of the present invention into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

The expression construct is conveniently a nucleic acid construct that comprises a polynucleotide encoding a polypeptide of the present invention operably linked with appropriate regulatory sequences required for expression of the nucleotide sequence in the plant or plant part of choice. Furthermore, the expression construct may comprise a selectable marker useful for identifying host cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences is determined, for example, on the basis of when, where, and how the polypeptide is desired to be expressed. For instance, the expression of the gene encoding a polypeptide of the present invention may be constitutive or inducible, or may be developmental, stage or tissue specific,

and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are, for example, described by Tague et al., 1988, *Plant Physiology* 86: 506.

For constitutive expression, the 35S-CaMV, the maize ubiquitin 1, and the rice actin 1 promoter may be used (Franck et al., 1980, *Cell* 21: 285-294, Christensen et al., 1992, *Plant Mol. Biol.* 18: 675-689; Zhang et al., 1991, *Plant Cell* 3: 1155-1165). organ-specific promoters may be, for example, a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards & Coruzzi, 1990, *Ann. Rev. Genet.* 24: 275-303), or from metabolic sink tissues such as meristems (Ito et al., 1994, *Plant Mol. Biol.* 24: 863-878), a seed specific promoter such as the glutelin, prolamin, globulin, or albumin promoter from rice (Wu et al., 1998, *Plant and Cell Physiology* 39: 885-889), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* (Conrad et al., 1998, *Journal of Plant Physiology* 152: 708-711), a promoter from a seed oil body protein (Chen et al., 1998, *Plant and Cell Physiology* 39: 935-941), the storage protein napA promoter from *Brassica napus*, or any other seed specific promoter known in the art, e.g., as described in WO 91/14772. Furthermore, the promoter may be a leaf specific promoter such as the rbcS promoter from rice or tomato (Kyozuka et al., 1993, *Plant Physiology* 102: 991-1000), the chlorella virus adenine methyltransferase gene promoter (Mitra and Higgins, 1994, *Plant Molecular Biology* 26: 85-93), or the aldP gene promoter from rice (Kagaya et al., 1995, *Molecular and General Genetics* 248: 668-674), or a wound inducible promoter such as the potato pin2 promoter (Xu et al., 1993, *Plant Molecular Biology* 22: 573-588). Likewise, the promoter may be inducible by abiotic treatments such as temperature, drought, or alterations in salinity or induced by exogenously applied substances that activate the promoter, e.g., ethanol, oestrogens, plant hormones such as ethylene, abscisic acid, and gibberellic acid, and heavy metals.

A promoter enhancer element may also be used to achieve higher expression of a polypeptide of the present invention in the plant. For instance, the promoter enhancer element may be an intron that is placed between the promoter and the nucleotide sequence encoding a polypeptide of the present invention. For instance, Xu et al., 1993, supra, disclose the use of the first intron of the rice actin 1 gene to enhance expression.

The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

The nucleic acid construct is incorporated into the plant genome according to conventional techniques known in the art, including *Agrobacterium*-mediated transformation, virus-mediated transformation, microinjection, particle bombardment, biolistic transformation, and electroporation (Gasser et al., 1990, *Science* 244: 1293; Potrykus, 1990, *Bio/Technology* 8: 535; Shimamoto et al., 1989, *Nature* 338: 274).

Presently, *Agrobacterium tumefaciens*-mediated gene transfer is the method of choice for generating transgenic dicots (for a review, see Hooykas and Schilperoort, 1992, *Plant Molecular Biology* 19: 15-38) and can also be used for transforming monocots, although other transformation methods are often used for these plants. Presently, the method of choice for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the transforming DNA) of embryonic calli or developing embryos (Christou, 1992, *Plant Journal* 2: 275-281; Shimamoto, 1994, *Current Opinion Biotechnology*

5: 158-162; Vasil et al., 1992, *Bio/Technology* 10: 667-674). An alternative method for transformation of monocots is based on protoplast transformation as described by Omiruleh et al., 1993, *Plant Molecular Biology* 21: 415-428.

Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well-known in the art. Often the transformation procedure is designed for the selective elimination of selection genes either during regeneration or in the following generations by using, for example, co-transformation with two separate T-DNA constructs or site specific excision of the selection gene by a specific recombinase.

The present invention also relates to methods of producing a polypeptide of the present invention comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide having glucoamylase activity of the present invention under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

Compositions

The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the glucoamylase activity of the composition has been increased, e.g., with an enrichment factor of at least 1.1.

The composition may comprise a polypeptide of the present invention as the major enzymatic component, e.g., a mono-component composition. Alternatively, the composition may comprise multiple enzymatic activities, such as an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, haloperoxidase, invertase, laccase, lipase, mannosidase, oxidase, pectinolytic enzyme, peptidoglutaminase, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase. The additional enzyme(s) may be produced, for example, by a microorganism belonging to the genus *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*; *Fusarium*, preferably *Fusarium bactridoides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium gramineum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcocroum*, *Fusarium sulphureum*, *Fusarium toruloseum*, *Fusarium trichothecioides*, or *Fusarium venenatum*; *Humicola*, preferably *Humicola insolens* or *Humicola lanuginosa*; or *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride*.

The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art.

Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other condi-

tions under which the composition is used may be determined on the basis of methods known in the art.

Combination of Glucoamylase and Acid Alpha-Amylase

According to this aspect of the invention a glucoamylase of the invention may be combined with an alpha-amylase, preferably acid alpha-amylase in a ratio of between 0.3 and 5.0 AFAU/AGU. More preferably the ratio between acid alpha-amylase activity and glucoamylase activity is at least 0.35, at least 0.40, at least 0.50, at least 0.60, at least 0.7, at least 0.8, at least 0.9, at least 1.0, at least 1.1, at least 1.2, at least 1.3, at least 1.4, at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.85, or even at least 1.9 AFAU/AGU. However, the ratio between acid alpha-amylase activity and glucoamylase activity should preferably be less than 4.5, less than 4.0, less than 3.5, less than 3.0, less than 2.5, or even less than 2.25 AFAU/AGU. In AUU/AGI the activities of acid alpha-amylase and glucoamylase are preferably present in a ratio of between 0.4 and 6.5 AUU/AGI. More preferably the ratio between acid alpha-amylase activity and glucoamylase activity is at least 0.45, at least 0.50, at least 0.60, at least 0.7, at least 0.8, at least 0.9, at least 1.0, at least 1.1, at least 1.2, at least 1.3, at least 1.4, at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.9, at least 2.0, at least 2.1, at least 2.2, at least 2.3, at least 2.4, or even at least 2.5 AUU/AGI. However, the ratio between acid alpha-amylase activity and glucoamylase activity is preferably less than 6.0, less than 5.5, less than 4.5, less than 4.0, less than 3.5, or even less than 3.0 AUU/AGI.

Above composition is suitable for use in a starch conversion process mentioned below for producing syrup and fermentation products, such as ethanol.

Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

Uses

The present invention is also directed to processes/methods for using the polypeptides having glucoamylase activity of the invention.

Uses according to the invention include starch conversion of starch to, e.g., syrup and fermentation products, including ethanol and beverages. Examples of processes where a glucoamylase of the invention may be used include the ones described in WO 92/20777, WO 03/066816, WO 03/066826, WO 2004/080923, and WO 2004/081193, which are hereby all incorporated by reference.

Production of Fermentation Products

Process for Producing Fermentation Products from Gelatinized Starch Containing Material

In this aspect the present invention relates to a process for producing a fermentation product, especially ethanol, from starch-containing material, which process includes a liquefaction step and sequentially or simultaneously performed saccharification and fermentation steps.

The invention relates to a process for producing a fermentation product from starch-containing material comprising the steps of:

(a) liquefying starch-containing material; preferably using an alpha-amylase

(b) saccharifying the liquefied material obtained in step (a) using a glucoamylase of the invention; and

(c) fermenting the saccharified material using a fermenting organism.

The fermentation product, such as especially ethanol, may optionally be recovered after fermentation, e.g., by distillation. Suitable starch-containing starting materials are listed

in the section "Starch-containing materials"-section below. Contemplated enzymes are listed in the "Enzymes"-section below. The liquefaction is preferably carried out in the presence of an alpha-amylase. The fermentation is preferably carried out in the presence of yeast, preferably a strain of *Saccharomyces*. Suitable fermenting organisms are listed in the "Fermenting Organisms"-section below. In preferred embodiments step (b) and (c) are carried out sequentially or simultaneously (i.e., as SSF process).

In a particular embodiment, the process of the invention further comprises, prior to the step (a), the steps of:

x) reducing the particle size of the starch-containing material, preferably by milling; and

y) forming a slurry comprising the starch-containing material and water.

The aqueous slurry may contain from 10-40 wt. %, preferably 25-35 wt. % starch-containing material. The slurry is heated to above the gelatinization temperature and alpha-amylase, preferably bacterial and/or acid fungal alpha-amylase, may be added to initiate liquefaction (thinning). The slurry may in an embodiment be jet-cooked to further gelatinize the slurry before being subjected to an alpha-amylase in step (a) of the invention.

More specifically liquefaction may be carried out as a three-step hot slurry process. The slurry is heated to between 60-95° C., preferably 80-85° C., and alpha-amylase is added to initiate liquefaction (thinning). Then the slurry may be jet-cooked at a temperature between 95-140° C., preferably 105-125° C., for 1-15 minutes, preferably for 3-10 minute, especially around 5 minutes. The slurry is cooled to 60-95° C. and more alpha-amylase is added to finalize hydrolysis (secondary liquefaction). The liquefaction process is usually carried out at pH 4.5-6.5, in particular at a pH between 5 and 6. Milled and liquefied whole grains are known as mash.

The saccharification in step (b) may be carried out using conditions well known in the art. For instance, a full saccharification process may last up to from about 24 to about 72 hours, however, it is common only to do a pre-saccharification of typically 40-90 minutes at a temperature between 30-65° C., typically about 60° C., followed by complete saccharification during fermentation in a simultaneous saccharification and fermentation process (SSF process). Saccharification is typically carried out at temperatures from 30-65° C., typically around 60° C., and at a pH between 4 and 5, normally at about pH 4.5.

The most widely used process in fermentation product, especially ethanol, production is the simultaneous saccharification and fermentation (SSF) process, in which there is no holding stage for the saccharification, meaning that fermenting organism, such as yeast, and enzyme(s) may be added together. SSF may typically be carried out at a temperature between 25° C. and 40° C., such as between 29° C. and 35° C., such as between 30° C. and 34° C., such as around 32° C. According to the invention the temperature may be adjusted up or down during fermentation.

In accordance with the present invention the fermentation step (c) includes, without limitation, fermentation processes used to produce alcohols (e.g., ethanol, methanol, butanol); organic acids (e.g., citric acid, acetic acid, itaconic acid, lactic acid, gluconic acid); ketones (e.g., acetone); amino acids (e.g., glutamic acid); gases (e.g., H₂ and CO₂); antibiotics (e.g., penicillin and tetracycline); enzymes; vitamins (e.g., riboflavin, B12, beta-carotene); and hormones. Preferred fermentation processes include alcohol fermentation processes, as are well known in the art. Preferred fermentation processes are anaerobic fermentation processes, as are well known in the art.

Processes for Producing Fermentation Products from Un-Gelatinized Starch-Containing

In this aspect the invention relates to processes for producing a fermentation product from starch-containing material without gelatinization of the starch-containing material (i.e., uncooked starch-containing material). In one embodiment only a glucoamylase of the invention is used during saccharification and fermentation. According to the invention the desired fermentation product, such as ethanol, can be produced without liquefying the aqueous slurry containing the starch-containing material. In one embodiment a process of the invention includes saccharifying (milled) starch-containing material, e.g., granular starch, below the gelatinization temperature in the presence of a glucoamylase of the invention to produce sugars that can be fermented into the desired fermentation product by a suitable fermenting organism.

Accordingly, in this aspect the invention relates to a process for producing a fermentation product from starch-containing material comprising:

(a) saccharifying starch-containing material with a mature glucoamylase according to the invention, preferably having the sequence shown as amino acids 18 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14 or amino acids 18 to 576 of SEQ ID NO: 16 or SEQ ID NO: 18, or a glucoamylase having at least 82% identity thereto, at a temperature below the initial gelatinization temperature of said starch-containing material,

(b) fermenting using a fermenting organism.

Steps (a) and (b) of the process of the invention may be carried out sequentially or simultaneously. In an embodiment a slurry comprising water and starch-containing material is prepared before step (a).

The fermentation process may be carried out for a period of 1 to 250 hours, preferably is from 25 to 190 hours, more preferably from 30 to 180 hours, more preferably from 40 to 170 hours, even more preferably from 50 to 160 hours, yet more preferably from 60 to 150 hours, even yet more preferably from 70 to 140 hours, and most preferably from 80 to 130 hours.

The term "initial gelatinization temperature" means the lowest temperature at which gelatinization of the starch commences. Starch heated in water begins to gelatinize between 50° C. and 75° C.; the exact temperature of gelatinization depends on the specific starch, and can readily be determined by the skilled artisan. Thus, the initial gelatinization temperature may vary according to the plant species, to the particular variety of the plant species as well as with the growth conditions. In the context of this invention the initial gelatinization temperature of a given starch-containing material is the temperature at which birefringence is lost in 5% of the starch granules using the method described by Gorinstein and Lii, 1992, *Starch/Stärke* 44(12): 461-466 (1992).

Before step (a) a slurry of starch-containing material, such as granular starch, having 10-55 wt. % dry solids, preferably 25-40 wt. % dry solids, more preferably 30-35 wt. % dry solids of starch-containing material may be prepared. The slurry may include water and/or process waters, such as stillage (backset), scrubber water, evaporator condensate or distillate, side stripper water from distillation, or other fermentation product plant process water. Because the process of the invention is carried out below the gelatinization temperature and thus no significant viscosity increase takes place, high levels of stillage may be used if desired. In an embodiment the aqueous slurry contains from about 1 to

about 70 vol. % stillage, preferably 15-60% vol. % stillage, especially from about 30 to 50 vol. % stillage.

The starch-containing material may be prepared by reducing the particle size, preferably by dry or wet milling, to 0.05 to 3.0 mm, preferably 0.1-0.5 mm. After being subjected to a process of the invention at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or preferably at least 99% of the dry solids of the starch-containing material is converted into a soluble starch hydrolysate.

The process of the invention is conducted at a temperature below the initial gelatinization temperature. Preferably the temperature at which step (a) is carried out is between 30-75° C., preferably between 45-60° C.

In a preferred embodiment step (a) and step (b) are carried out as a sequential or simultaneous saccharification and fermentation process. In such preferred embodiment the process is typically carried at a temperature between 25° C. and 40° C., such as between 29° C. and 35° C., such as between 30° C. and 34° C., such as around 32° C. According to the invention the temperature may be adjusted up or down during fermentation.

In an embodiment simultaneous saccharification and fermentation is carried out so that the sugar level, such as glucose level, is kept at a low level such as below 6 wt. %, preferably below about 3 wt. %, preferably below about 2 wt. %, more preferred below about 1 wt. %, even more preferred below about 0.5 wt. %, or even more preferred 0.25% wt. %, such as below about 0.1 wt. %. Such low levels of sugar can be accomplished by simply employing adjusted quantities of enzyme and fermenting organism. A skilled person in the art can easily determine which quantities of enzyme and fermenting organism to use. The employed quantities of enzyme and fermenting organism may also be selected to maintain low concentrations of maltose in the fermentation broth. For instance, the maltose level may be kept below about 0.5 wt. % or below about 0.2 wt. %.

The process of the invention may be carried out at a pH in the range between 3 and 7, preferably from pH 3.5 to 6, or more preferably from pH 4 to 5.

Starch-Containing Materials

Any suitable starch-containing starting material, including granular starch, may be used according to the present invention. The starting material is generally selected based on the desired fermentation product. Examples of starch-containing starting materials, suitable for use in a process of present invention, include tubers, roots, stems, whole grains, corns, cobs, wheat, barley, rye, milo, sago, cassava, tapioca, sorghum, rice, peas, beans, or sweet potatoes, or mixtures thereof, or cereals, sugar-containing raw materials, such as molasses, fruit materials, sugar cane or sugar beet, potatoes, and cellulose-containing materials, such as wood or plant residues, or mixtures thereof. Contemplated are both waxy and non-waxy types of corn and barley.

The term "granular starch" means raw uncooked starch, i.e., starch in its natural form found in cereal, tubers or grains. Starch is formed within plant cells as tiny granules insoluble in water. When put in cold water, the starch granules may absorb a small amount of the liquid and swell. At temperatures up to 50° C. to 75° C. the swelling may be reversible. However, with higher temperatures an irreversible swelling called "gelatinization" begins. Granular starch to be processed may be a highly refined starch quality, preferably at least 90%, at least 95%, at least 97% or at least 99.5% pure or it may be a more crude starch containing

material comprising milled whole grain including non-starch fractions such as germ residues and fibers. The raw material, such as whole grain, is milled in order to open up the structure and allowing for further processing. Two milling processes are preferred according to the invention: wet and dry milling. In dry milling whole kernels are milled and used. Wet milling gives a good separation of germ and meal (starch granules and protein) and is often applied at locations where the starch hydrolysate is used in production of syrups. Both dry and wet milling is well known in the art of starch processing and is equally contemplated for the process of the invention.

The starch-containing material is reduced in particle size, preferably by dry or wet milling, in order to expose more surface area. In an embodiment the particle size is between 0.05 to 3.0 mm, preferably 0.1-0.5 mm, or so that at least 30%, preferably at least 50%, more preferably at least 70%, even more preferably at least 90% of the starch-containing material fit through a sieve with a 0.05 to 3.0 mm screen, preferably 0.1-0.5 mm screen.

Fermentation Products

The term "fermentation product" means a product produced by a process including a fermentation step using a fermenting organism. Fermentation products contemplated according to the invention include alcohols (e.g., ethanol, methanol, butanol); organic acids (e.g., citric acid, acetic acid, itaconic acid, lactic acid, gluconic acid); ketones (e.g., acetone); amino acids (e.g., glutamic acid); gases (e.g., H₂ and CO₂); antibiotics (e.g., penicillin and tetracycline); enzymes; vitamins (e.g., riboflavin, B₁₂, beta-carotene); and hormones. In a preferred embodiment the fermentation product is ethanol, e.g., fuel ethanol; drinking ethanol, i.e., potable neutral spirits; or industrial ethanol or products used in the consumable alcohol industry (e.g., beer and wine), dairy industry (e.g., fermented dairy products), leather industry and tobacco industry. Preferred beer types comprise ales, stouts, porters, lagers, bitters, malt liquors, happoushu, high-alcohol beer, low-alcohol beer, low-calorie beer or light beer. Preferred fermentation processes used include alcohol fermentation processes, as are well known in the art. Preferred fermentation processes are anaerobic fermentation processes, as are well known in the art.

Fermenting Organisms

"Fermenting organism" refers to any organism, including bacterial and fungal organisms, suitable for use in a fermentation process and capable of producing desired a fermentation product. Especially suitable fermenting organisms are able to ferment, i.e., convert, sugars, such as glucose or maltose, directly or indirectly into the desired fermentation product. Examples of fermenting organisms include fungal organisms, such as yeast. Preferred yeast includes strains of *Saccharomyces* spp., in particular, *Saccharomyces cerevisiae*. Commercially available yeast include, e.g., Red Star™/Lesaffre Ethanol Red (available from Red Star/Lesaffre, USA) FALI (available from Fleischmann's Yeast, a division of Burns Philp Food Inc., USA), SUPERSTART (available from Alltech), GERT STRAND (available from Gert Strand AB, Sweden) and FERMIOL (available from DSM Specialties).

Enzymes

Glucoamylase

The glucoamylase is preferably a glucoamylase of the invention. However, as mentioned above a glucoamylase of the invention may also be combined with other glucoamylases. The term "glucoamylase" (1,4-alpha-D-glucan glucohydrolase, EC 3.2.1.3) is an enzyme, which catalyzes the

release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules.

The glucoamylase may added in an amount of 0.001 to 10 AGU/g DS, preferably from 0.01 to 5 AGU/g DS, such as around 0.1, 0.3, 0.5, 1 or 2 AGU/g DS, especially 0.1 to 0.5 AGU/g DS or 0.02-20 AGU/g DS, preferably 0.1-10 AGU/g DS.

Alpha-Amylase

The alpha-amylase may according to the invention be of any origin. Preferred are alpha-amylases of fungal or bacterial origin.

In a preferred embodiment the alpha-amylase is an acid alpha-amylase, e.g., fungal acid alpha-amylase or bacterial acid alpha-amylase. The term "acid alpha-amylase" means an alpha-amylase (EC 3.2.1.1) which added in an effective amount has activity optimum at a pH in the range of 3 to 7, preferably from 3.5 to 6, or more preferably from 4-5.

Bacterial Alpha-Amylases

According to the invention a bacterial alpha-amylase may preferably be derived from the genus *Bacillus*.

In a preferred embodiment the *Bacillus* alpha-amylase is derived from a strain of *B. licheniformis*, *B. amyloliquefaciens*, *B. subtilis* or *B. stearothermophilus*, but may also be derived from other *Bacillus* sp. Specific examples of contemplated alpha-amylases include the *Bacillus licheniformis* alpha-amylase (BLA) shown in SEQ ID NO: 4 in WO 99/19467, the *Bacillus amyloliquefaciens* alpha-amylase (BAN) shown in SEQ ID NO: 5 in WO 99/19467, and the *Bacillus stearothermophilus* alpha-amylase (BSG) shown in SEQ ID NO: 3 in WO 99/19467. In an embodiment of the invention the alpha-amylase is an enzyme having a degree of identity of at least 60%, preferably at least 70%, more preferred at least 80%, even more preferred at least 90%, such as at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to any of the sequences shown as SEQ ID NO: 1, 2, 3, 4, or 5, respectively, in WO 99/19467.

The *Bacillus* alpha-amylase may also be a variant and/or hybrid, especially one described in any of WO 96/23873, WO 96/23874, WO 97/41213, WO 99/19467, WO 00/60059, and WO 02/10355 (all documents hereby incorporated by reference). Specifically contemplated alpha-amylase variants are disclosed in U.S. Pat. Nos. 6,093,562, 6,187,576, and 6,297,038 (hereby incorporated by reference) and include *Bacillus stearothermophilus* alpha-amylase (BSG alpha-amylase) variants having a deletion of one or two amino acid in position 179 to 182, preferably a double deletion disclosed in WO 96/23873—see, e.g., page 20, lines 1-10 (hereby incorporated by reference), preferably corresponding to delta (181-182) compared to the wild-type BSG alpha-amylase amino acid sequence set forth in SEQ ID NO: 3 disclosed in WO 99/19467 or deletion of amino acids 179 and 180 using SEQ ID NO: 3 in WO 99/19467 for numbering (which reference is hereby incorporated by reference). Even more preferred are *Bacillus* alpha-amylases, especially *Bacillus stearothermophilus* alpha-amylase, which have a double deletion corresponding to delta (181-182) and further comprise a N193F substitution (also denoted 1181*+G182*+N193F) compared to the wild-type BSG alpha-amylase amino acid sequence set forth in SEQ ID NO: 3 disclosed in WO 99/19467.

The alpha-amylase may also be a maltogenic alpha-amylase. A "maltogenic alpha-amylase" (glucan 1,4-alpha-maltohydrolase, EC 3.2.1.133) is able to hydrolyze amylose and amylopectin to maltose in the alpha-configuration. A maltogenic alpha-amylase from *Bacillus stearothermophilus* strain NCIB 11837 is commercially available from Novozymes NS, Denmark. The maltogenic alpha-amylase is

described in U.S. Pat. Nos. 4,598,048, 4,604,355 and 6,162,628, which are hereby incorporated by reference.

Bacterial Hybrid Alpha-Amylases

A hybrid alpha-amylase specifically contemplated comprises 445 C-terminal amino acid residues of the *Bacillus licheniformis* alpha-amylase (shown as SEQ ID NO: 4 in WO 99/19467) and the 37 N-terminal amino acid residues of the alpha-amylase derived from *Bacillus amyloliquefaciens* (shown as SEQ ID NO: 3 in WO 99/19467), with one or more, especially all, of the following substitutions:

G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S (using the *Bacillus licheniformis* numbering). Also preferred are variants having one or more of the following mutations (or corresponding mutations in other *Bacillus* alpha-amylase backbones): H154Y, A181T, N190F, A209V and Q264S and/or deletion of two residues between positions 176 and 179, preferably deletion of E178 and G179 (using the SEQ ID NO: 5 numbering of WO 99/19467).

The bacterial alpha-amylase may be added in amounts as are well-known in the art. When measured in KNU units (described below in the "Materials & Methods"-section) the alpha-amylase activity is preferably present in an amount of 0.5-5,000 NU/g of DS, in an amount of 1-500 NU/g of DS, or more preferably in an amount of 5-1,000 NU/g of DS, such as 10-100 NU/g DS.

Fungal Alpha-Amylases

Fungal acid alpha-amylases include acid alpha-amylases derived from a strain of the genus *Aspergillus*, such as *Aspergillus kawachii*, *Aspergillus niger*, or *Aspergillus oryzae* alpha-amylases.

A preferred acid fungal alpha-amylase is a Fungamyl-like alpha-amylase which is preferably derived from a strain of *Aspergillus oryzae*. In the present disclosure, the term "Fungamyl-like alpha-amylase" indicates an alpha-amylase which exhibits a high identity, i.e., more than 70%, more than 75%, more than 80%, more than 85%, more than 90%, more than 95%, more than 96%, more than 97%, more than 98%, more than 99% or even 100% identity to the mature part of the amino acid sequence shown in SEQ ID NO: 10 in WO 96/23874.

Another preferred acid alpha-amylase is derived from a strain *Aspergillus niger*. In a preferred embodiment the acid fungal alpha-amylase is the one from *A. niger* disclosed as "AMYA_ASPNG" in the Swiss-prot/TrEMBL database under the primary accession no. P56271 and described in more detail in WO 89/01969 (Example 3). The acid *Aspergillus niger* acid alpha-amylase is also shown as SEQ ID NO: 1 in WO 2004/080923 (Novozymes) which is hereby incorporated by reference. Also variants of said acid fungal amylase having at least 70% identity, such as at least 80% or even at least 90% identity, such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to SEQ ID NO: 1 in WO 2004/080923 are contemplated. A suitable commercially available acid fungal alpha-amylase derived from *Aspergillus niger* is SP288 (available from Novozymes NS, Denmark).

In a preferred embodiment the alpha-amylase is derived from *Aspergillus kawachii* and disclosed by Kaneko et al., 1996, *J. Ferment. Bioeng.* 81: 292-298, "Molecular-cloning and determination of the nucleotide-sequence of a gene encoding an acid-stable alpha-amylase from *Aspergillus kawachii*"; and further as EMBL:#AB008370.

The fungal acid alpha-amylase may also be a wild-type enzyme comprising a carbohydrate-binding module (CBM) and an alpha-amylase catalytic domain (i.e., a none-hybrid),

or a variant thereof. In an embodiment the wild-type acid alpha-amylase is derived from a strain of *Aspergillus kawachii*.

Fungal Hybrid Alpha-Amylases

In a preferred embodiment the fungal acid alpha-amylase is a hybrid alpha-amylase. Preferred examples of fungal hybrid alpha-amylases include the ones disclosed in WO 2005/003311 or U.S. Application Publication no. 2005/0054071 (Novozymes) or U.S. application No. 60/638,614 (Novozymes) which is hereby incorporated by reference. A hybrid alpha-amylase may comprise an alpha-amylase catalytic domain (CD) and a carbohydrate-binding domain/module (CBM) and optional a linker.

Specific examples of contemplated hybrid alpha-amylases include those disclosed in U.S. application No. 60/638,614 including Fungamyl variant with catalytic domain JA118 and *Athelia rolfsii* SBD (SEQ ID NO: 100 in U.S. application No. 60/638,614), *Rhizomucor pusillus* alpha-amylase with *Athelia rolfsii* AMG linker and SBD (SEQ ID NO: 101 in U.S. application No. 60/638,614) and *Meripilus giganteus* alpha-amylase with *Athelia rolfsii* glucoamylase linker and SBD (SEQ ID NO: 102 in U.S. application No. 60/638,614).

Other specific examples of contemplated hybrid alpha-amylases include those disclosed in U.S. Application Publication no. 2005/0054071, including those disclosed in Table 3 on page 15, such as *Aspergillus niger* alpha-amylase with *Aspergillus kawachii* linker and starch binding domain. Commercial Alpha-Amylase Products

Preferred commercial compositions comprising alpha-amylase include MYCOLASE from DSM (Gist Brocades), BANTM, TERMAMYLTM SC, FUNGAMYLTM, LIQUOZYMETM X and SANTM EXTRA L (Novozymes NS) and CLARASETM L-40,000, DEX-LOTM, SPEZYMETM FRED, SPEZYMETM AA, SPEZYMETM Ethyl, GC358, GC980, SPEZYMETM RSL, and SPEZYMETM DELTA AA (Genencor Int.), and the acid fungal alpha-amylase sold under the trade name SP288 (available from Novozymes NS, Denmark).

An acid alpha-amylase may according to the invention be added in an amount of 0.1 to 10 AFAU/g DS, preferably 0.10 to 5 AFAU/g DS, especially 0.3 to 2 AFAU/g DS.

Production of Syrup

The present invention also provides a process of using a glucoamylase of the invention for producing syrup, such as glucose and the like, from starch-containing material. Suitable starting materials are exemplified in the "Starch-containing materials"-section above. Generally, the process comprises the steps of partially hydrolyzing starch-containing material (liquefaction) in the presence of alpha-amylase and then further saccharifying the release of glucose from the non-reducing ends of the starch or related oligo- and polysaccharide molecules in the presence of glucoamylase of the invention.

Liquefaction and saccharification may be carried out as described above for fermentation product production.

The glucoamylase of the invention may also be used in immobilized form. This is suitable and often used for producing specialty syrups, such as maltose syrups, and further for the raffinate stream of oligosaccharides in connection with the production of fructose syrups, e.g., high fructose syrup (HFS).

Consequently, this aspect of the invention relates to a process of producing syrup from starch-containing material, comprising

(a) liquefying starch-containing material in the presence of an alpha-amylase, and

(b) saccharifying the material obtained in step (a) using a glucoamylase of the invention.

A syrup may be recovered from the saccharified material obtained in step (b).

Details on suitable conditions can be found above.

Brewing

A glucoamylase of the invention can also be used in a brewing process. The glucoamylase of the invention is added in effective amounts which can be easily determined by the skilled person in the art.

The invention described and claimed herein is not to be limited in scope by the specific embodiments herein disclosed, since these embodiments are intended as illustrations of several aspects of the invention. Any equivalent embodiments are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. In the case of conflict, the present disclosure including definitions will control.

Various references are cited herein, the disclosures of which are incorporated by reference in their entireties. The present invention is further described by the following examples which should not be construed as limiting the scope of the invention.

Materials & Methods

Yeast:

RED STAR™ available from Red Star/Lesaffre, USA

Media and Reagents:

Chemicals used as buffers and substrates were commercial products of at least reagent grade.

PDA: 39 g/L Potato Dextrose Agar, 20 g/L agar, 50 ml/L glycerol

Methods

Unless otherwise stated, DNA manipulations and transformations were performed using standard methods of molecular biology as described in Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, N.Y.; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology", John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for *Bacillus*". John Wiley and Sons, 1990.

Glucoamylase Activity

Glucoamylase activity may be measured in AGI units or in Glucoamylase Units (AGU).

Glucoamylase Activity (AGI)

Glucoamylase (equivalent to amyloglucosidase) converts starch into glucose. The amount of glucose is determined here by the glucose oxidase method for the activity determination. The method described in the section 76-11 Starch—Glucoamylase Method with Subsequent Measurement of Glucose with Glucose Oxidase in "Approved methods of the American Association of Cereal Chemists". Vol. 1-2 AACC, from American Association of Cereal Chemists, (2000); ISBN: 1-891127-12-8.

One glucoamylase unit (AGI) is the quantity of enzyme which will form 1 micro mole of glucose per minute under the standard conditions of the method.

Standard Conditions/Reaction Conditions:

Substrate: Soluble starch, concentration approx. 16 g dry matter/L.

Buffer: Acetate, approx. 0.04 M, pH=4.3

pH: 4.3

Incubation temperature: 60° C.

Reaction time: 15 minutes

Termination of the reaction: NaOH to a concentration of approximately 0.2 g/L (pH-9)

Enzyme concentration: 0.15-0.55 AAU/mL.

The starch should be Lintner starch, which is a thin-boiling starch used in the laboratory as colorimetric indicator. Lintner starch is obtained by dilute hydrochloric acid treatment of native starch so that it retains the ability to color blue with iodine.

Glucoamylase Activity (AGU)

The Novo Glucoamylase Unit (AGU) is defined as the amount of enzyme, which hydrolyzes 1 micromole maltose per minute under the standard conditions 37° C., pH 4.3, substrate: maltose 23.2 mM, buffer: acetate 0.1 M, reaction time 5 minutes.

An autoanalyzer system may be used. Mutarotase is added to the glucose dehydrogenase reagent so that any alpha-D-glucose present is turned into beta-D-glucose. Glucose dehydrogenase reacts specifically with beta-D-glucose in the reaction mentioned above, forming NADH which is determined using a photometer at 340 nm as a measure of the original glucose concentration.

AMG incubation:

| | |
|-------------------------|-----------------|
| Substrate: | maltose 23.2 mM |
| Buffer: | acetate 0.1M |
| pH: | 4.30 ± 0.05 |
| Incubation temperature: | 37° C. ± 1 |
| Reaction time: | 5 minutes |
| Enzyme working range: | 0.5-4.0 AGU/mL |

Color reaction:

| | |
|-------------------------|-----------------------------|
| GlucDH: | 430 U/L |
| Mutarotase: | 9 U/L |
| NAD: | 0.21 mM |
| Buffer: | phosphate 0.12M; 0.15M NaCl |
| pH: | 7.60 ± 0.05 |
| Incubation temperature: | 37° C. ± 1 |
| Reaction time: | 5 minutes |
| Wavelength: | 340 nm |

A folder (EB-SM-0131.02/01) describing this analytical method in more detail is available on request from Novozymes NS, Denmark, which folder is hereby included by reference.

Alpha-Amylase Activity (KNU)

Alpha-amylase activity may be determined using potato starch as substrate. This method is based on the break-down of modified potato starch by the enzyme, and the reaction is followed by mixing samples of the starch/enzyme solution with an iodine solution. Initially, a blackish-blue color is formed, but during the break-down of the starch the blue color gets weaker and gradually turns into a reddish-brown, which is compared to a colored glass standard.

One Kilo Novo alpha amylase Unit (KNU) is defined as the amount of enzyme which, under standard conditions (i.e., at 37° C. ± 0.05; 0.0003 M Ca²⁺; and pH 5.6) dextrinizes 5260 mg starch dry substance Merck Amylum soluble.

A folder EB-SM-0009.02/01 describing this analytical method in more detail is available upon request to Novozymes NS, Denmark, which folder is hereby included by reference.

Acid Alpha-Amylase Activity

When used according to the present invention the activity of any acid alpha-amylase may be measured in AFAU (Acid Fungal Alpha-amylase Units). Alternatively activity of acid alpha-amylase may be measured in AAU (Acid Alpha-amylase Units).

Acid Alpha-Amylase Units (AAU)

Acid alpha-amylase activity can be measured in AAU (Acid Alpha-amylase Units), which is an absolute method. One Acid Amylase Unit (AAU) is the quantity of enzyme converting 1 g of starch (100% of dry matter) per hour under standardized conditions into a product having a transmission at 620 nm after reaction with an iodine solution of known strength equal to the one of a color reference.

Standard Conditions/Reaction Conditions:

Substrate: Soluble starch. Concentration approx. 20 g DS/L.

Buffer: Citrate, approx. 0.13 M, pH=4.2

Iodine solution: 40.176 g potassium iodide+0.088 g iodine/L

City water: 15°-20° dH (German degree hardness)

pH: 4.2

Incubation temperature: 30° C.

Reaction time: 11 minutes

Wavelength: 620 nm

Enzyme concentration: 0.13-0.19 AAU/mL


Enzyme working range: 0.13-0.19 AAU/mL

The starch should be Lintner starch, which is a thin-boiling starch used in the laboratory as colorimetric indicator. Lintner starch is obtained by dilute hydrochloric acid treatment of native starch so that it retains the ability to color blue with iodine. Further details can be found in EP 0140410 B2, which disclosure is hereby included by reference.

Acid Alpha-Amylase Activity (AFAU)

Acid alpha-amylase activity may be measured in AFAU (Acid Fungal Alpha-amylase Units), which are determined relative to an enzyme standard. 1 AFAU is defined as the amount of enzyme which degrades 5.260 mg starch dry matter per hour under the below mentioned standard conditions.

Acid alpha-amylase, an endo-alpha-amylase (1,4-alpha-D-glucan-glucanohydrolase, EC 3.2.1.1) hydrolyzes alpha-1,4-glucosidic bonds in the inner regions of the starch molecule to form dextrans and oligosaccharides with different chain lengths. The intensity of color formed with iodine is directly proportional to the concentration of starch. Amylase activity is determined using reverse colorimetry as a reduction in the concentration of starch under the specified analytical conditions.



Standard Conditions/Reaction Conditions

Substrate: Soluble starch, approx. 0.17 g/L

Buffer: Citrate, approx. 0.03 M

Iodine (I₂): 0.03 g/L

CaCl₂: 1.85 mM

pH: 2.50±0.05

Incubation temperature: 40° C.

Reaction time: 23 seconds

Wavelength: 590 nm

Enzyme concentration: 0.025 AFAU/mL

Enzyme working range: 0.01-0.04 AFAU/mL

A folder EB-SM-0259.02/01 describing this analytical method in more detail is available upon request to Novozymes NS, Denmark, which folder is hereby included by reference.

Example 1

Simultaneous Saccharification and Fermentation (SSF) with *Gloeophyllum sepiarium* AMG

The SSF performance of *Gloeophyllum* sp. glucoamylases was tested at different enzyme doses. Fermentation was run under the following conditions:

Substrate: Ground corn was slurried with backset and adjusted its dry substance to approximately 32% (w/w). It was then liquefied at 85° C. and pH 5.8. The liquefied mash had a DE of 13.4.

Temperature: 32° C.

Initial pH: 5.0

Enzyme dose: *Gloeophyllum* sp. AMG produced in *Aspergillus niger* at 30, 40, 55 and 70 micrograms enzyme protein/g DS. Enzymes were compared to a purified sample of the commercial *Talaromyces emersonii* AMG dosed at same dosages. The highest dose of *Talaromyces emersonii* AMG is equivalent to an industry relevant amount of 0.56 AGU/g DS. A control for maximum obtainable saccharification was prepared using excess amounts of commercial AMG and alpha-amylase.

Fermentation

To the substrate for SSF, 1000 ppm urea as nitrogen source and 3 ppm penicillin for bacterial control were added; the pH was adjusted to 5.0 with H₂SO₄. Aliquots of 5 g mash were transferred to 15 ml centrifuge tubes with a hole drilled at the top for CO₂ release. Enzymes and yeast were added and the tubes were placed in a water bath without stirring at 32° C. for 54 hrs. Samples were analyzed in HPLC for determination of ethanol produced during fermentation. The results are shown in the tables below.

TABLE 1

Ethanol g/L produced during SSF with *Gloeophyllum sepiarium* AMG at 30, 40, 55 and 70 micrograms enzyme protein/g DS as compared to *Talaromyces emersonii* AMG.
Control resulted in 133.08 g/L ethanol

| | Enzyme dose (micrograms enzyme protein/g DS) | | | |
|----------------------------------------------|----------------------------------------------|-------|-------|-------|
| | 30 | 40 | 55 | 70 |
| <i>Talaromyces emersonii</i> (SEQ ID NO: 20) | 110.3 | 119.8 | 124.9 | 126.9 |
| <i>Gloeophyllum sepiarium</i> (SEQ ID NO: 2) | 115.5 | 122.6 | 130.8 | 130.7 |

TABLE 2

Ethanol g/L produced during SSF with *Gloeophyllum sepiarium* AMG at 30, 40, 55 and 70 micrograms enzyme protein/g DS as compared to *Talaromyces emersonii* AMG.
Control resulted in 131.9 g/L ethanol

| | Enzyme dose (micrograms enzyme protein/g DS) | | | |
|----------------------------------------------|----------------------------------------------|-------|-------|-------|
| | 30 | 40 | 55 | 70 |
| <i>Talaromyces emersonii</i> (SEQ ID NO: 20) | 110.4 | 118.9 | 123.4 | 125.6 |
| <i>Gloeophyllum sepiarium</i> (SEQ ID NO: 4) | 113.0 | 121.1 | 129.0 | 129.5 |

TABLE 3

| Ethanol yield produced during SSF with <i>Gloeophyllum</i> sp. AMG at 30, 40, 55 and 70 micrograms enzyme protein/g DS as compared to <i>Talaromyces emersonii</i> AMG and <i>Trametes cingulata</i> AMG. Control resulted in 100% yield. | | | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|-----|-----|-----|
| | Enzyme dose (micrograms enzyme protein/g DS) | | | |
| | 30 | 40 | 55 | 70 |
| <i>Gloeophyllum sepiarium</i> (SEQ ID NO: 2) | 87% | 92% | 98% | 98% |
| <i>Gloeophyllum trabeum</i> (SEQ ID NO: 6) | 93% | 97% | 97% | 98% |
| <i>Gloeophyllum trabeum</i> (SEQ ID NO: 14) | 88% | 97% | 97% | 98% |
| <i>Trametes cingulata</i> AMG G1 (SEQ ID NO: 22) | 87% | 90% | 97% | 98% |
| <i>Talaromyces emersonii</i> , T-AMG (SEQ ID NO: 20) | 83% | 90% | 94% | 95% |

Example 2

Specific Activity of *Gloeophyllum* sp. AMGs Compared to *Trametes cingulata* AMG

Specific activity was calculated by dividing the glucoamylase AGU-activity (according to the procedure described above) and enzyme protein from each preparation (AA analysis). The table below shows the results for the different *Gloeophyllum* sp. AMGs compared to the *Trametes cingulata* AMG.

| Specific activity | [AGU/mg] |
|-------------------------------------------------|----------|
| <i>Trametes cingulata</i> AMG G1, SEQ ID NO: 22 | 5.6 |
| <i>Gloeophyllum sepiarium</i> AMG, SEQ ID NO: 4 | 6.3 |
| <i>Gloeophyllum sepiarium</i> AMG, SEQ ID NO: 2 | 7.2 |
| <i>Gloeophyllum trabeum</i> , SEQ ID NO: 16 | 6.1 |
| <i>Gloeophyllum trabeum</i> , SEQ ID NO: 18 | 6.0 |
| <i>Gloeophyllum trabeum</i> , SEQ ID NO: 14 | 7.2 |
| <i>Gloeophyllum trabeum</i> , SEQ ID NO: 6 | 6.5 |

10 Deposit of Biological Material

The following biological material has been deposited under the terms of the Budapest Treaty with Deutsche Sammlung von Mikroorganismen and Zellkulturen GmbH (DSMZ), Mascheroder Weg 1 B, D-38124 Braunschweig, Germany, and given the following accession number: Deposit: *E. coli* strain NN059223 with plasmid comprising sequence D7279 (SEQ ID NO: 1) Accession Number: DSM 23222 Date of Deposit: Jan. 13, 2010

20 The strain has been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by foreign patent laws to be entitled thereto. The deposit represents a substantially pure culture of the deposited strain. The deposit is available as required by foreign patent laws in countries wherein counterparts of the subject application or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 1722

<212> TYPE: DNA

<213> ORGANISM: *Gloeophyllum sepiarium*

<400> SEQUENCE: 1

```

atgtaccgct tecttgtctg tgcgctgggg cttgcggcat cagttctcgc ccagtcggtc      60
gacagctatg ttagcagcga aggtcccata gccaaaggcg gcgtccttgc taacattggg      120
ccgaacggct ccaaggcctc tggcgcatcc gctggtgttg tggtcgcgag ccctagcacg      180
tcggaccccg actattggta cacttggacg cgtgactcgt cctcgtatt caagtcaact      240
attgaccagt acaccaccgg catcgacagc acgagctctc tgaggactct catcgacgat      300
ttcgtaactg ccgaggctaa tctccagcaa gtctctaacc ctagtggtag cctcaccacc      360
ggtggcttgg gagagcccaa gttcaacgtc gacgaaactg catttactgg tgcattgggg      420
cgaccccaac gcgacggacc tgccctccgc tcgactgcat tgatcacgta cggtaaactg      480
ctgttgcca acgaaatac gagctatggt acgagcaatc tgtggccgat catccagaac      540
gaccttggtt atgtcgtgtc atactggaac cagtctacct acgacctctg ggaggaagta      600
gactcgtcat cgttcttca c tactgcagta cagcaccgtg ctctccgtga aggtcgggcc      660
ttcgtaccg ccateggtea gacttcgacg gtcagcagct atacgactca ggccgacaat      720
cttctgtgct tcttgcagtc ttaactggaac ccgagcgggt gttacatcac tgctaactact      780
ggcggcggcc gttccggcaa ggatgccaac acacttctgg catccattca cacgtacgac      840
cccagcgggg gctgcgacgc tgcgacttcc cagccctgct ctgacaaggc actgtcgaac      900
ctgaaggtct acgtcgactc tttccgctcg gctactcca tcaacagtgg tgcgcctct      960

```

-continued

```

aacgctgcgc tcgccacggg tcgttatccc gaggatagct accaggggtgg aaacccttgg 1020
tacctcacca catttgcggt cgccgagcaa ctctatgatg ctctcaatgt ctgggagtcg 1080
cagggttccc tcgaggtcac ctccacctcc cttgccttct tccagcagtt ctcacccggc 1140
gtcactgctg gcacctactc ttctagctcc agcacatata gcacctcac gtctgccatc 1200
aagaactttg ccgatggatt tgcgcctatc aatgctaagt acacgccatc caacgggtggc 1260
ctggcggaac aatacagcaa gagcgacggt tctcccctta gcgcgggtgga cttgacgtgg 1320
agctacgctt cggctttgac ggcgtttgaa gcaaggaaca atactcagtt cgccggctgg 1380
ggcgctgcag gcctgactgt gccttcctct tgctccggca actctggtgg gccgaccgtt 1440
gctgtcacat tcaacgtgaa cgccgagact gtgtggggag agaacatcta tcttactggt 1500
tccgtcgatg ctctggagaa ctggtcggcc gacaatgccc tctgctctc atcgctaata 1560
taccgcacct ggagtatcac cgtcaacttg ccggcgagca ctgctattga gtacaagtac 1620
atccgcaaaa ataatggggc cgttacctgg gagtcagacc ccaacaatag catcactact 1680
ccggccagcg gctcgacgac cgagaatgac acttggcggt ga 1722

```

<210> SEQ ID NO 2

<211> LENGTH: 573

<212> TYPE: PRT

<213> ORGANISM: *Gloeophyllum sepiarium*

<400> SEQUENCE: 2

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Ser Val Leu
1           5           10          15
Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Ile Ala Lys
20          25          30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
35          40          45
Ala Ser Ala Gly Val Val Val Ala Ser Pro Ser Thr Ser Asp Pro Asp
50          55          60
Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
65          70          75          80
Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Thr
85          90          95
Leu Ile Asp Asp Phe Val Thr Ala Glu Ala Asn Leu Gln Gln Val Ser
100         105         110
Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
115         120         125
Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
130         135         140
Asp Gly Pro Ala Leu Arg Ser Thr Ala Leu Ile Thr Tyr Gly Asn Trp
145         150         155         160
Leu Leu Ser Asn Gly Asn Thr Ser Tyr Val Thr Ser Asn Leu Trp Pro
165         170         175
Ile Ile Gln Asn Asp Leu Gly Tyr Val Val Ser Tyr Trp Asn Gln Ser
180         185         190
Thr Tyr Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr Thr
195         200         205
Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Ala
210         215         220
Ile Gly Gln Thr Ser Gln Val Ser Ser Tyr Thr Thr Gln Ala Asp Asn
225         230         235         240

```


-continued

```

attgaccagt acaccaccgg catcgacagc acgagctctc tgaggactct catcgacgat 300
ttcgtaactg ccgagggtaa tctocagcaa gtctctaacc ctagtggtag cctcaccacc 360
ggtggcttgg gagagccaaa gttcaacgtc gacgaaactg catttactgg tgcattgggt 420
cgaccccaac gcgacggatc tgccctccgc tcgactgcat tgatcacata cggtactgg 480
ctggtgtcca acgccaatac gagctatgtt acgagcaaac tgtggccgat catccagaac 540
gatcttggt atgtcgtgtc atactggaac cagtctacct acgatctctg ggaagaagta 600
gactcgtcat cgttcttcac tactgcagtg cagcatcgtg ctctcccgca aggtgcggcg 660
ttcgctacag caatcgggtca gacttcgcag gttagcagct ataccacgca ggcggacaat 720
ctcttggtct tcttcgagtc ttactggaac ccgagcggtg gttacatcac tgtaaacact 780
ggtggcggcc gttccggcaa ggatgccaac actcttctgg catccattca cacgtaacgac 840
cccagcgcgc gctcgcagc tgcgacttcc caaccctgct ctgacaaggc actgtcaaac 900
ctgaaagtct atgtcgactc gttccgctcg gtctactcta ttaacagtgg tatcgctct 960
aacgctgctg tcgccacggg tcggtacccc gaggacagct accaggggtg aaacccttgg 1020
tacctacca catttgccgt cgcagcaaa ctctatgatg ctctcaatgt ctgggagtcg 1080
cagggttccc tcgaggtcac ttccacctcc cttgccttct tccagcagtt ttcattctggc 1140
gtcactgctg gcacctactc ttctagctcc agcacgtaca gcacctcac gtctgccatc 1200
aaaagccttg ccgacggatt tgtcactatc aacgctaagt acacgccatc caacggtggc 1260
ctggcggaac aatacagcaa gagcgacggc tctcccctta gcgcggtgga cttgacgtgg 1320
agctacgctt cggctttgac ggcggttgaa gcaaggaacg atactcagtt cgcggctgg 1380
ggcgctgcag gcctgactgt ccttctctct tgctccgcca actcgggtgg gccgaccgtc 1440
gctgttatc tcaacgtgaa cgcagagact gtgtggggag agaacatcta tcttactggt 1500
tcagtcgatg ctctggagaa ctggtcggcc gacaatgccc tctgctctc atcggctaata 1560
taccgcacct ggagtatcac cgtcaacttg ccggcgagca ctgctattga gtacaagtac 1620
atccgcaaaa ataatggggc cgttacctgg gagtcagacc ccaacaatag taccactact 1680
ccggccagcg gctcgcagc cgagaatgac acttggcgtt ga 1722

```

<210> SEQ ID NO 4

<211> LENGTH: 573

<212> TYPE: PRT

<213> ORGANISM: Gloeophyllum sepiarium

<400> SEQUENCE: 4

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Ser Val Leu
 1           5           10           15
Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Ile Ala Lys
 20          25          30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
 35          40          45
Ala Ser Ala Gly Val Val Val Ala Ser Pro Ser Thr Ser Asp Pro Asp
 50          55          60
Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
 65          70          75          80
Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Thr
 85          90          95
Leu Ile Asp Asp Phe Val Thr Ala Glu Ala Asn Leu Gln Gln Val Ser
100         105         110

```


-continued

| 530 | 535 | 540 | |
|-----------------------------|---------------------|---------------------|-----|
| Asn Gly Ala Val Thr Trp | Glu Ser Asp Pro Asn | Asn Ser Ile Thr Thr | |
| 545 | 550 | 555 | 560 |
| Pro Ala Ser Gly Ser Thr Thr | Glu Asn Asp Thr | Trp Arg | |
| | 565 | 570 | |

<210> SEQ ID NO 5
 <211> LENGTH: 1722
 <212> TYPE: DNA
 <213> ORGANISM: Gloeophyllum trabeum

<400> SEQUENCE: 5

| | |
|--------------------------------------------------------------------|------|
| atgtaccgct tccttgtctg tgcgctgggg ettgeggcaa cagttctcgc ccagtcggtc | 60 |
| gacagctatg ttagcagcga aggtcccata gccaaaggcgg gcgtccttgc taacattggg | 120 |
| ccgaacggct ccaaggcctc tggcgcaccc gctggtgttg tggtcgcgag ccctagcacg | 180 |
| tcggaccctg actattggta cacttggacg cgtgactcat ccctcgtttt caagtcactc | 240 |
| attgaccagt acaccactgg catcgacagc acgagctctc tgaggactct catcgacgat | 300 |
| ttcgtaactg ccgaggctaa tctccagcaa gtctctaacc ctagtgttac cctcaccacc | 360 |
| ggtggccttg gagagcccaa gttcaacgtc gacgaaactg catttactgg tgcattgggt | 420 |
| cgaccccaac gcgacggacc tgccctccgc tcgactgcat tgatcacgta cggttaactgg | 480 |
| ctggtgtcca acgaaaatac gagctatgtt acgagcattc tgtggccgat catccagaac | 540 |
| gaccttggtt atgtcgtgtc atattggaac cagtctacct acgacctctg ggaggaagta | 600 |
| gactcgtcat cgttcttcac taccgcagta cagcaccgtg ctctccgtga aggtgcggcc | 660 |
| ttcgctaccg ccctcgggtc gacttcgcag gtcagcagct atacgactca ggcggacaat | 720 |
| cttctgtgct tcttgcagtc ttactggaac ccgagcggtg gttacatcac tgetaacact | 780 |
| ggcggcggcc gttcgggcaa ggatgccaac acacttctgg catccattca cacgtaacgac | 840 |
| cccagcgcgg gctcgcagc tcgcacttcc cagccctgct ctgacaaggc actgtcgaac | 900 |
| ctgaaggctt acttcgactc tttccgctcg gtctactcca tcaacagtgg tgtcgcctct | 960 |
| aacgctgccc tcgccacggg tcgttatccc gaggatagct accaggggtg aaacccttgg | 1020 |
| tacctcacca catttgcggg cgcgcagcaa ctctatgat ctctcaatgt ctgggagtcg | 1080 |
| cagggttccc tcgaggctac ctccacctcc cttgccttct tccagcagtt ctcatccggc | 1140 |
| gtcactgctg gcacctactc ttctagctcc agcacataca gcacctcac gtctgacctc | 1200 |
| aagaactttg ccgatggatt tgcctctatc aatgctaagt acacgccatc caacggtggc | 1260 |
| ctggcggaac aatacagcaa gagcgaaggc tctcccctta gcgcgggtgga cttgacgtgg | 1320 |
| agctacgctt cggctttgac ggcggttgaa gcaaggaaca atactcagtt cgcggctggg | 1380 |
| ggcgtcgcag gcctgactgt gccttctctc tgctccggca actctggtgg gccgaccgtt | 1440 |
| gctgtcacat tcaacgtgaa cgcgcagact gtgtggggag agaacatcta tcttactggt | 1500 |
| tccgtcgatg ctctggagaa ctggtcggcc gacaatgccc tctgctctc atcggtaat | 1560 |
| taccgcacct ggagtatcac cgtcaacttg ccggcgagca ctgctattga gtacaagtac | 1620 |
| atccgcaaaa ataatggggc cgttacctgg gagtcagacc ccaacaatag catcaactact | 1680 |
| ccggccagcg gctcgcagac cgagaatgac acttggcgtt aa | 1722 |

<210> SEQ ID NO 6
 <211> LENGTH: 573
 <212> TYPE: PRT
 <213> ORGANISM: Gloeophyllum trabeum

-continued

<400> SEQUENCE: 6

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Thr Val Leu
 1 5 10 15
 Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Ile Ala Lys
 20 25 30
 Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
 35 40 45
 Ala Ser Ala Gly Val Val Val Ala Ser Pro Ser Thr Ser Asp Pro Asp
 50 55 60
 Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
 65 70 75 80
 Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Thr
 85 90 95
 Leu Ile Asp Asp Phe Val Thr Ala Glu Ala Asn Leu Gln Gln Val Ser
 100 105 110
 Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
 115 120 125
 Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
 130 135 140
 Asp Gly Pro Ala Leu Arg Ser Thr Ala Leu Ile Thr Tyr Gly Asn Trp
 145 150 155 160
 Leu Leu Ser Asn Gly Asn Thr Ser Tyr Val Thr Ser Ile Leu Trp Pro
 165 170 175
 Ile Ile Gln Asn Asp Leu Gly Tyr Val Val Ser Tyr Trp Asn Gln Ser
 180 185 190
 Thr Tyr Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr Thr
 195 200 205
 Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Ala
 210 215 220
 Ile Gly Gln Thr Ser Gln Val Ser Ser Tyr Thr Thr Gln Ala Asp Asn
 225 230 235 240
 Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr Ile
 245 250 255
 Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr Leu
 260 265 270
 Leu Ala Ser Ile His Thr Tyr Asp Pro Ser Ala Gly Cys Asp Ala Ala
 275 280 285
 Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val Tyr
 290 295 300
 Phe Asp Ser Phe Arg Ser Val Tyr Ser Ile Asn Ser Gly Val Ala Ser
 305 310 315 320
 Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Gln Gly
 325 330 335
 Gly Asn Pro Trp Tyr Leu Thr Thr Phe Ala Val Ala Glu Gln Leu Tyr
 340 345 350
 Asp Ala Leu Asn Val Trp Glu Ser Gln Gly Ser Leu Glu Val Thr Ser
 355 360 365
 Thr Ser Leu Ala Phe Phe Gln Gln Phe Ser Ser Gly Val Thr Ala Gly
 370 375 380
 Thr Tyr Ser Ser Ser Ser Ser Thr Tyr Ser Thr Leu Thr Ser Ala Ile
 385 390 395 400
 Lys Asn Phe Ala Asp Gly Phe Val Ala Ile Asn Ala Lys Tyr Thr Pro

-continued

```

aagagctttg ccgatggatt tgcgctatc aatgctaagt acacgccatc caacgggtggc 1260
ctggcggaac aatatagcaa gagcgacggc tctcccctta gcgcggtgga cttgacgtgg 1320
agctacgctt cggctttgac ggcgtttgaa gcaaggaacg aactcagtt cgccggtg 1380
ggcgctgcaa gcctgactgt gccttcctct tgctccggca actctggtgg gccgaccgtc 1440
gctgtcacat tcaacgtgaa cgccgagact gtgtggggag agaacatcta tcttactggt 1500
tccgtogatg ctctggagaa ctggtcggcc gacaatgccc tctgtctctc atcggctaac 1560
taccgacct ggagatcac cgtcaacttg ccggcgagca ctgccattga gtacaagtac 1620
atccgcaaaa ataatggggc agttacctgg gagtcagacc ccaacaatag catcactact 1680
cggccagcgc gctcgacgac cgagaatgac acttggcgtt aa 1722

```

```

<210> SEQ ID NO 8
<211> LENGTH: 573
<212> TYPE: PRT
<213> ORGANISM: Gloeophyllum sepiarium

```

```

<400> SEQUENCE: 8

```

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Ser Val Leu
1           5           10          15
Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Ile Ala Lys
20          25          30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
35          40          45
Ala Ser Ala Gly Val Val Val Ala Ser Pro Ser Thr Ser Asp Pro Asp
50          55          60
Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
65          70          75          80
Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Thr
85          90          95
Leu Ile Asp Asp Phe Val Thr Ala Glu Ala Asn Leu Gln Gln Val Ser
100         105         110
Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
115         120         125
Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
130         135         140
Asp Gly Pro Ala Leu Arg Ser Thr Ala Leu Ile Thr Tyr Gly Asn Trp
145         150         155         160
Leu Leu Ser Asn Gly Asn Thr Ser Tyr Val Thr Ser Asn Leu Trp Pro
165         170         175
Ile Ile Gln Asn Asp Leu Gly Tyr Val Val Ser Tyr Trp Asn Gln Ser
180         185         190
Thr Tyr Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr Thr
195         200         205
Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Ala
210         215         220
Ile Gly Gln Thr Ser Gln Val Ser Ser Tyr Thr Thr Gln Ala Asp Asn
225         230         235         240
Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr Ile
245         250         255
Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr Leu
260         265         270
Leu Ala Ser Ile His Thr Tyr Asp Pro Ser Ala Gly Cys Asp Ala Ala

```

-continued

| 275 | | | | | 280 | | | | | 285 | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Phe | Gln | Pro | Cys | Ser | Asp | Lys | Ala | Leu | Ser | Asn | Leu | Lys | Val | Tyr |
| 290 | | | | | | 295 | | | | | 300 | | | | |
| Val | Asp | Ser | Phe | Arg | Ser | Val | Tyr | Ser | Ile | Asn | Ser | Gly | Ile | Ala | Ser |
| 305 | | | | | 310 | | | | | 315 | | | | | 320 |
| Asn | Ala | Ala | Val | Ala | Thr | Gly | Arg | Tyr | Pro | Glu | Asp | Ser | Tyr | Gln | Gly |
| | | | | 325 | | | | | 330 | | | | | 335 | |
| Gly | Asn | Pro | Trp | Tyr | Leu | Thr | Thr | Phe | Ala | Val | Ala | Glu | Gln | Leu | Tyr |
| | | | 340 | | | | | 345 | | | | | 350 | | |
| Asp | Ala | Leu | Asn | Val | Trp | Glu | Leu | Gln | Gly | Ser | Leu | Glu | Val | Thr | Ser |
| | | 355 | | | | 360 | | | | | | 365 | | | |
| Thr | Ser | Leu | Ala | Phe | Phe | Gln | Gln | Phe | Ser | Ser | Gly | Val | Thr | Ala | Gly |
| 370 | | | | | | 375 | | | | | 380 | | | | |
| Thr | Tyr | Ser | Ser | Ser | Ser | Ser | Thr | Tyr | Ser | Thr | Leu | Thr | Ser | Ala | Ile |
| 385 | | | | | 390 | | | | | 395 | | | | | 400 |
| Lys | Ser | Phe | Ala | Asp | Gly | Phe | Val | Ala | Ile | Asn | Ala | Lys | Tyr | Thr | Pro |
| | | | | 405 | | | | | 410 | | | | | | 415 |
| Ser | Asn | Gly | Gly | Leu | Ala | Glu | Gln | Tyr | Ser | Lys | Ser | Asp | Gly | Ser | Pro |
| | | | 420 | | | | | 425 | | | | | 430 | | |
| Leu | Ser | Ala | Val | Asp | Leu | Thr | Trp | Ser | Tyr | Ala | Ser | Ala | Leu | Thr | Ala |
| | | 435 | | | | | 440 | | | | | | 445 | | |
| Phe | Glu | Ala | Arg | Asn | Asp | Thr | Gln | Phe | Ala | Gly | Trp | Gly | Ala | Ala | Ser |
| 450 | | | | | | 455 | | | | | 460 | | | | |
| Leu | Thr | Val | Pro | Ser | Ser | Cys | Ser | Gly | Asn | Ser | Gly | Gly | Pro | Thr | Val |
| 465 | | | | | 470 | | | | | 475 | | | | | 480 |
| Ala | Val | Thr | Phe | Asn | Val | Asn | Ala | Glu | Thr | Val | Trp | Gly | Glu | Asn | Ile |
| | | | | 485 | | | | | 490 | | | | | 495 | |
| Tyr | Leu | Thr | Gly | Ser | Val | Asp | Ala | Leu | Glu | Asn | Trp | Ser | Ala | Asp | Asn |
| | | | 500 | | | | | 505 | | | | | | 510 | |
| Ala | Leu | Leu | Leu | Ser | Ser | Ala | Asn | Tyr | Pro | Thr | Trp | Ser | Ile | Thr | Val |
| | | 515 | | | | | 520 | | | | | | 525 | | |
| Asn | Leu | Pro | Ala | Ser | Thr | Ala | Ile | Glu | Tyr | Lys | Tyr | Ile | Arg | Lys | Asn |
| 530 | | | | | | 535 | | | | | 540 | | | | |
| Asn | Gly | Ala | Val | Thr | Trp | Glu | Ser | Asp | Pro | Asn | Asn | Ser | Ile | Thr | Thr |
| 545 | | | | | 550 | | | | | 555 | | | | | 560 |
| Pro | Ala | Ser | Gly | Ser | Thr | Thr | Glu | Asn | Asp | Thr | Trp | Arg | | | |
| | | | | 565 | | | | | | 570 | | | | | |

<210> SEQ ID NO 9

<211> LENGTH: 1722

<212> TYPE: DNA

<213> ORGANISM: Gloeophyllum sepiarium

<400> SEQUENCE: 9

```

atgtaccgct tccttgtctg tgcgctcggg cttgctgcaa cagttctcgc ccagtcggtc    60
gacagctatg ttagcagcga agggcccgta gccaaaggcgg gcgtccttgc taacattgga    120
ccgaacggct ccaaggcctc tggcgcgtcc gccggtggtg tggtagcgag cccgagcagc    180
tccgaccccg actattggta cacttggacg cgtgactcgt ccctcgtttt caagtcactc    240
attgaccagt acaccaccgg catcgacagc acgagctctc tgaggactct catcgacgat    300
ttcgtaaactg ccgaggctaa tetccagcaa gtctetaacc ctagtggtag cctcactacc    360
ggtagccttg gagagccaaa gttcaacgtc gacgaaactg catttactgg tgcattgggt    420
cgaccccaac gcgacggacc tgcctcctg tgcactgcat tgatcacata cggtaatgg    480

```

-continued

```

ctgttgcca acgaaatac gagctatgtt acgagcaatc tgtggccgat catccagaac 540
gaccttggtt atgtcgtgtc atactggaac cagtctacct acgacctctg ggaggaagta 600
gactcgtcat cgttcttcac tactgcagta cagcaccgtg ctctccgtga aggtgcggcg 660
ttcgtaccg ccacgggtca gacttcgcag gtcagcagct atacgactca ggcggacaat 720
cttctgtgct tcttgcaagc atactggaac ccgagcgggtg gttatatcac tgctaact 780
ggtggcggcc gttccggcaa ggatgccaac actcttctgg catccattca cacatacgat 840
cccagcgcg gctcgcagc tgccacttcc cagccctgct ctgacaagge gctgtcgaat 900
ctgaaggtct acgtcgaatc cttccgctca atctactcca tcaacagtgg tatcgcctct 960
aacgccgcg tcgccacggg tcgttaccac gaggacagct accaggggtg aaacccttgg 1020
tacctcacca catttgccgt cgcggagcaa ctctatgatg ctctcaatgt ctgggagtcg 1080
cagggttccc tcgaggtcac ctccacctcc cttgccttct tccagcagtt ctcacccggc 1140
gtcactgctg gcacctactc ttctagctcc agcacataca gcacctcac gtctgccatc 1200
aagagctttg ccgatggatt tgcctctatc aatgctaagt acacgccatc caacgggtgg 1260
ctggcggaac aatatagcaa gagcgacggc tctcccctta gcgcgggtgga cttgacgtgg 1320
agctacgctt cggctttgac ggcgtttgaa gcaaggaacg aactcagtt cgcggctgg 1380
ggcgtgcag gcctgactgt gccttctct tgcctcggca actctggtgg gccgaccgtc 1440
gcggtccat tcaacgtgaa cgcggagact gtgtggggag agaacatcta tcttactggt 1500
tccgtcagtg ctctggagaa ctggtcggcc gacaatgccc tccgtctctc atcggetaac 1560
taccgcacct ggagtatcac cgtcaacttg ccggcgagca ctgccattga gtacaagtac 1620
atccgcaaaa ataatggggc cgtcacctgg gagtcagacc ccaacaatag catcactact 1680
cggccagcgc gctcgcagac cgagaatgac acttggcgtt aa 1722

```

```

<210> SEQ ID NO 10
<211> LENGTH: 573
<212> TYPE: PRT
<213> ORGANISM: Gloeophyllum sepiarium

```

```

<400> SEQUENCE: 10

```

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Thr Val Leu
1           5           10          15
Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Val Ala Lys
20          25          30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
35          40          45
Ala Ser Ala Gly Val Val Val Ala Ser Pro Ser Thr Ser Asp Pro Asp
50          55          60
Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
65          70          75          80
Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Thr
85          90          95
Leu Ile Asp Asp Phe Val Thr Ala Glu Ala Asn Leu Gln Gln Val Ser
100         105         110
Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
115         120         125
Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
130         135         140
Asp Gly Pro Ala Leu Arg Ser Thr Ala Leu Ile Thr Tyr Gly Asn Trp

```

-continued

| | | | |
|-----------------------------------------------------------------|-----|-----|-----|
| 145 | 150 | 155 | 160 |
| Leu Leu Ser Asn Gly Asn Thr Ser Tyr Val Thr Ser Asn Leu Trp Pro | 165 | 170 | 175 |
| Ile Ile Gln Asn Asp Leu Gly Tyr Val Val Ser Tyr Trp Asn Gln Ser | 180 | 185 | 190 |
| Thr Tyr Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr Thr | 195 | 200 | 205 |
| Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Ala | 210 | 215 | 220 |
| Ile Gly Gln Thr Ser Gln Val Ser Ser Tyr Thr Thr Gln Ala Asp Asn | 225 | 230 | 235 |
| Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr Ile | 245 | 250 | 255 |
| Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr Leu | 260 | 265 | 270 |
| Leu Ala Ser Ile His Thr Tyr Asp Pro Ser Ala Gly Cys Asp Ala Ala | 275 | 280 | 285 |
| Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val Tyr | 290 | 295 | 300 |
| Val Asp Ser Phe Arg Ser Ile Tyr Ser Ile Asn Ser Gly Ile Ala Ser | 305 | 310 | 315 |
| Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Gln Gly | 325 | 330 | 335 |
| Gly Asn Pro Trp Tyr Leu Thr Thr Phe Ala Val Ala Glu Gln Leu Tyr | 340 | 345 | 350 |
| Asp Ala Leu Asn Val Trp Glu Ser Gln Gly Ser Leu Glu Val Thr Ser | 355 | 360 | 365 |
| Thr Ser Leu Ala Phe Phe Gln Gln Phe Ser Ser Gly Val Thr Ala Gly | 370 | 375 | 380 |
| Thr Tyr Ser Ser Ser Ser Ser Thr Tyr Ser Thr Leu Thr Ser Ala Ile | 385 | 390 | 395 |
| Lys Ser Phe Ala Asp Gly Phe Val Ala Ile Asn Ala Lys Tyr Thr Pro | 405 | 410 | 415 |
| Ser Asn Gly Gly Leu Ala Glu Gln Tyr Ser Lys Ser Asp Gly Ser Pro | 420 | 425 | 430 |
| Leu Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ser Ala Leu Thr Ala | 435 | 440 | 445 |
| Phe Glu Ala Arg Asn Asp Thr Gln Phe Ala Gly Trp Gly Ala Ala Gly | 450 | 455 | 460 |
| Leu Thr Val Pro Ser Ser Cys Ser Gly Asn Ser Gly Gly Pro Thr Val | 465 | 470 | 475 |
| Ala Val Thr Phe Asn Val Asn Ala Glu Thr Val Trp Gly Glu Asn Ile | 485 | 490 | 495 |
| Tyr Leu Thr Gly Ser Val Asp Ala Leu Glu Asn Trp Ser Ala Asp Asn | 500 | 505 | 510 |
| Ala Leu Leu Leu Ser Ser Ala Asn Tyr Pro Thr Trp Ser Ile Thr Val | 515 | 520 | 525 |
| Asn Leu Pro Ala Ser Thr Ala Ile Glu Tyr Lys Tyr Ile Arg Lys Asn | 530 | 535 | 540 |
| Asn Gly Ala Val Thr Trp Glu Ser Asp Pro Asn Asn Ser Ile Thr Thr | 545 | 550 | 555 |
| Pro Ala Ser Gly Ser Thr Thr Glu Asn Asp Thr Trp Arg | 565 | 570 | |

-continued

```

<210> SEQ ID NO 11
<211> LENGTH: 1722
<212> TYPE: DNA
<213> ORGANISM: Gloeophyllum sp

<400> SEQUENCE: 11
atgtaccgct tccttgtctg tgccctcggg cttgctgcaa cagttctcgc ccagtcggtc   60
gacagctatg ttagcagcga agggcccgta gccaaaggcgg gcgtccttgc taacattgga   120
ccgaacggct ccaaggcctc tggcgcgtcc gccgggtgtg tggtagcgag cccgagcacg   180
tcggaccccg actattggta cacttggacg cgtgactcgt ccctcgtttt caagtcactc   240
attgaccagt acaccaccgg catcgacagc acgagctctc tgaggactct catcgacgat   300
ttcgtaaactg ccgaggctaa tctccagcaa gtctetaacc ctagtggtag cctcactacc   360
ggtggccttg gagagccaaa gttcaacgtc gacgaaactg catttactgg tgcattgggt   420
cgaccccaac gcgacggacc tgccctccgt tcgactgcat tgatcacata cggttaactgg   480
ctggtgtcca acgaaaatac gagctatgtt acgagcaatc tgtggccgat catccagaac   540
gaccttggtt atgtcgtgtc atactggaac cagtctacct acgacctctg ggaggaagta   600
gactcgtcat cgttcttccac tactgcagta cagcaccgtg ctctccgtga aggtgcggcg   660
ttcgtaccg ccacgtgta gacttcgcag gtcagcagct atacgactca ggcggacaat   720
cttctgtgct tcttgcagtc atactggaac ccgagcgggt gttatatcac tgctaact   780
ggtggcggcc gttccggcaa ggatgccaac actcttctgg catccattca cacatacgat   840
cccagcgcg gctgcgacgc tgccacttcc cagccctgct ctgacaagge gctgtcgaat   900
ctgaaggtct acgtcgactc cttccgctcg atctactcca tcaacagtgg tatcgcctct   960
aacgccgcg tcgccacggg tcggtacccc gaggacagct accaggggtg aaacccttgg   1020
tacctcacca catttgccgt cgcgagcaa ctctatgatg ctctcaatgt ctgggagtcg   1080
cagggttccc tcgaggtcac ctccacctcc cttgccttct tccagcagtt ctcatccggc   1140
gtcactgctg gcacctactc ttctagctcc agcacataca gcacctcac gtctgccatc   1200
aagagccttg ccgatggatt tgtcgtatc aatgctaagt acacgccatc caacgggtggc   1260
ctggcggaac aatatagcaa gagcgacggc tctcccctta gcgcgggtga cttgacgtgg   1320
agctacgctt cggcttggac ggcggttgaa gcaaggaacg aactcagtt cgcggctgg   1380
ggcgtgcag gcctgactgt gccttctct tgcctcggca actctggtgg gccgaccgtc   1440
gctgtccat tcaacgtgaa cgcgagact gtgtggggag agaacatcta tcttactggt   1500
tccgtcgatg ctctggagaa ctggtcggcc gacaatgcc tctgctctc atcggttaac   1560
taccgcacct ggagtatcac cgtcaacttg ccggcgagca ctgocattga gtacaagtac   1620
atccgcaaaa ataatggggc cgtcacctgg gagtcagatc ccaacaatag catcactact   1680
cggccagcgc gctcgacgac cgagaatgac acttggcgtt aa                               1722

```

```

<210> SEQ ID NO 12
<211> LENGTH: 573
<212> TYPE: PRT
<213> ORGANISM: Gloeophyllum sp

```

```

<400> SEQUENCE: 12

```

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Thr Val Leu
1           5           10          15
Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Val Ala Lys

```

-continued

| 20 | | | | 25 | | | | 30 | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Gly | Val | Leu | Ala | Asn | Ile | Gly | Pro | Asn | Gly | Ser | Lys | Ala | Ser | Gly |
| | 35 | | | | | | 40 | | | | | 45 | | | |
| Ala | Ser | Ala | Gly | Val | Val | Val | Ala | Ser | Pro | Ser | Thr | Ser | Asp | Pro | Asp |
| | 50 | | | | | | 55 | | | | 60 | | | | |
| Tyr | Trp | Tyr | Thr | Trp | Thr | Arg | Asp | Ser | Ser | Leu | Val | Phe | Lys | Ser | Leu |
| | 65 | | | | 70 | | | | | 75 | | | | | 80 |
| Ile | Asp | Gln | Tyr | Thr | Thr | Gly | Ile | Asp | Ser | Thr | Ser | Ser | Leu | Arg | Thr |
| | | | 85 | | | | | | 90 | | | | | 95 | |
| Leu | Ile | Asp | Asp | Phe | Val | Thr | Ala | Glu | Ala | Asn | Leu | Gln | Gln | Val | Ser |
| | | 100 | | | | | | 105 | | | | | 110 | | |
| Asn | Pro | Ser | Gly | Thr | Leu | Thr | Thr | Gly | Gly | Leu | Gly | Glu | Pro | Lys | Phe |
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Asn | Val | Asp | Glu | Thr | Ala | Phe | Thr | Gly | Ala | Trp | Gly | Arg | Pro | Gln | Arg |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Asp | Gly | Pro | Ala | Leu | Arg | Ser | Thr | Ala | Leu | Ile | Thr | Tyr | Gly | Asn | Trp |
| | 145 | | | | 150 | | | | | 155 | | | | | 160 |
| Leu | Leu | Ser | Asn | Gly | Asn | Thr | Ser | Tyr | Val | Thr | Ser | Asn | Leu | Trp | Pro |
| | | | 165 | | | | | | 170 | | | | | 175 | |
| Ile | Ile | Gln | Asn | Asp | Leu | Gly | Tyr | Val | Val | Ser | Tyr | Trp | Asn | Gln | Ser |
| | | 180 | | | | | | | 185 | | | | 190 | | |
| Thr | Tyr | Asp | Leu | Trp | Glu | Glu | Val | Asp | Ser | Ser | Ser | Phe | Phe | Thr | Thr |
| | | 195 | | | | | 200 | | | | | 205 | | | |
| Ala | Val | Gln | His | Arg | Ala | Leu | Arg | Glu | Gly | Ala | Ala | Phe | Ala | Thr | Ala |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Ile | Gly | Gln | Thr | Ser | Gln | Val | Ser | Ser | Tyr | Thr | Thr | Gln | Ala | Asp | Asn |
| | 225 | | | | 230 | | | | | 235 | | | | | 240 |
| Leu | Leu | Cys | Phe | Leu | Gln | Ser | Tyr | Trp | Asn | Pro | Ser | Gly | Gly | Tyr | Ile |
| | | | 245 | | | | | | 250 | | | | | 255 | |
| Thr | Ala | Asn | Thr | Gly | Gly | Gly | Arg | Ser | Gly | Lys | Asp | Ala | Asn | Thr | Leu |
| | | 260 | | | | | | | 265 | | | | 270 | | |
| Leu | Ala | Ser | Ile | His | Thr | Tyr | Asp | Pro | Ser | Ala | Gly | Cys | Asp | Ala | Ala |
| | | 275 | | | | | 280 | | | | | 285 | | | |
| Thr | Phe | Gln | Pro | Cys | Ser | Asp | Lys | Ala | Leu | Ser | Asn | Leu | Lys | Val | Tyr |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Val | Asp | Ser | Phe | Arg | Ser | Ile | Tyr | Ser | Ile | Asn | Ser | Gly | Ile | Ala | Ser |
| | 305 | | | | 310 | | | | | 315 | | | | | 320 |
| Asn | Ala | Ala | Val | Ala | Thr | Gly | Arg | Tyr | Pro | Glu | Asp | Ser | Tyr | Gln | Gly |
| | | | 325 | | | | | | 330 | | | | | 335 | |
| Gly | Asn | Pro | Trp | Tyr | Leu | Thr | Thr | Phe | Ala | Val | Ala | Glu | Gln | Leu | Tyr |
| | | 340 | | | | | | 345 | | | | | 350 | | |
| Asp | Ala | Leu | Asn | Val | Trp | Glu | Ser | Gln | Gly | Ser | Leu | Glu | Val | Thr | Ser |
| | | 355 | | | | | 360 | | | | | 365 | | | |
| Thr | Ser | Leu | Ala | Phe | Phe | Gln | Gln | Phe | Ser | Ser | Gly | Val | Thr | Ala | Gly |
| | 370 | | | | | 375 | | | | | 380 | | | | |
| Thr | Tyr | Ser | Ser | Ser | Ser | Thr | Tyr | Ser | Thr | Leu | Thr | Ser | Ala | Ile | |
| | 385 | | | | 390 | | | | 395 | | | | | 400 | |
| Lys | Ser | Phe | Ala | Asp | Gly | Phe | Val | Ala | Ile | Asn | Ala | Lys | Tyr | Thr | Pro |
| | | | 405 | | | | | | 410 | | | | | 415 | |
| Ser | Asn | Gly | Gly | Leu | Ala | Glu | Gln | Tyr | Ser | Lys | Ser | Asp | Gly | Ser | Pro |
| | | | 420 | | | | | | 425 | | | | 430 | | |
| Leu | Ser | Ala | Val | Asp | Leu | Thr | Trp | Ser | Tyr | Ala | Ser | Ala | Leu | Thr | Ala |
| | | 435 | | | | | 440 | | | | | 445 | | | |

-continued

Phe Glu Ala Arg Asn Asp Thr Gln Phe Ala Gly Trp Gly Ala Ala Gly
 450 455 460
 Leu Thr Val Pro Ser Ser Cys Ser Gly Asn Ser Gly Gly Pro Thr Val
 465 470 475 480
 Ala Val Thr Phe Asn Val Asn Ala Glu Thr Val Trp Gly Glu Asn Ile
 485 490 495
 Tyr Leu Thr Gly Ser Val Asp Ala Leu Glu Asn Trp Ser Ala Asp Asn
 500 505 510
 Ala Leu Leu Leu Ser Ser Val Asn Tyr Pro Thr Trp Ser Ile Thr Val
 515 520 525
 Asn Leu Pro Ala Ser Thr Ala Ile Glu Tyr Lys Tyr Ile Arg Lys Asn
 530 535 540
 Asn Gly Ala Val Thr Trp Glu Ser Asp Pro Asn Asn Ser Ile Thr Thr
 545 550 555 560
 Pro Ala Ser Gly Ser Thr Thr Glu Asn Asp Thr Trp Arg
 565 570

<210> SEQ ID NO 13

<211> LENGTH: 1722

<212> TYPE: DNA

<213> ORGANISM: Gloeophyllum trabeum

<400> SEQUENCE: 13

```

atgtaccgct tccttgtctg tgcgctgggg cttgcggcgt cagttctcgc ccagtcggtc   60
gacagctatg ttagcagcga aggtcccata gccaaaggcgg gcgtocttgc taacattgga   120
ccgaacggct ccaaggcctc tggcgcaccc gctggcgctg tggtcgcaag cccaagcacg   180
tcggaccccg actattggta cacttggacg cgtgactcgt ccctcgtttt caagtcactc   240
attgaccagt acaccaccgg catcgacggc acgagctctc tgaggactct catcgacgat   300
ttcgtaactg ccgaggctaa tcttcagcaa gtctctaacc ctagtggtag cctcaccacc   360
ggtggccttg gagagccaaa gttcaacgtc gacgaaactg catttactgg tgcattgggt   420
cgacccaac gtgacggacc tgccctcgc tcgactcgc tgatcacata cggtaaactg   480
ctgttgcca acgggaatac gagctatgtt acgagcaatc tgtggccgat catccagaac   540
gaccttgggt atgtcgtgtc atactggaac cagtccactg acgatctctg ggaggaagta   600
gactcgtcgt cattctcacc tactgcagtg cagcaccctg ctctcctgga aggagcggcg   660
ttcgtaccg ctatcggtca gacttcgcag gttagcagct atacgactca ggcggacaat   720
cttctgtgct tcttgcagtc ttactggaac ccgagtgggt gttacatcac tgctaactc   780
ggcggcggcc gttccggcaa ggatgccaac actcttctgg catccattca tacgtacgac   840
cccagcgcgg gctcgcgacg tcgcactttc cagccctgct ctgacaaggc attgtcgaac   900
ctgaaagtct acgttgactc tttccgctcg gtctattcca tcaacagtgg tatcgcctct   960
aacgccgcgg tcgccacggg tcggttaccac gaggacagct accaggggtg aaatccttgg  1020
tacctcacca catttgcggt cgcgagcaa ctctatgat ctctcaatgt ctgggagtcg  1080
cagggtccc tcgaggtcac ctccacctcc cttgccttct tccagcagtt ctcatccggt  1140
gtcactgctg gcacctactc ttctagctcc agcacgtaca gcacctcac gtctgctatc  1200
aagagctttg ccgatggatt tgcgctgctc aacgctaagt acacgccatc caacgggtggc  1260
ctggcggaac aatacagcaa gagcgcgggt tctcccctta gcgcgggtgga cttgacgtgg  1320
agctacgctt cggctttgac ggcgcttgaa gcaaggaaca aactcagtt cgcggctgg  1380

```

-continued

```

ggcgctgcag gcctgactgt gccttcctct tgctccggca actctggtgg gccgaccgtc 1440
getgttacat tcaacgtgaa cgccgagact gtgtggggag agaacatcta tcttactggt 1500
tcagtcgatg ctctggagaa ctggtcggcc gacaatgccc tctgctctc atcggctaata 1560
taccgcacct ggagtatcac cgtaacttg ccggcgagca ctgctattga gtacaagtac 1620
atccgcaaaa ataatggggc cgttacctgg gagtcagacc ccaacaatag catcactact 1680
ccggctagcg gctcgacgac cgagaatgac acttgccggtt aa 1722

```

<210> SEQ ID NO 14

<211> LENGTH: 573

<212> TYPE: PRT

<213> ORGANISM: Gloeophyllum trabeum

<400> SEQUENCE: 14

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Ala Ala Ser Val Leu
1           5           10           15
Ala Gln Ser Val Asp Ser Tyr Val Ser Ser Glu Gly Pro Ile Ala Lys
20           25           30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
35           40           45
Ala Ser Ala Gly Val Val Val Ala Ser Pro Ser Thr Ser Asp Pro Asp
50           55           60
Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
65           70           75           80
Ile Asp Gln Tyr Thr Thr Gly Ile Asp Gly Thr Ser Ser Leu Arg Thr
85           90           95
Leu Ile Asp Asp Phe Val Thr Ala Glu Ala Asn Leu Gln Gln Val Ser
100          105          110
Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
115          120          125
Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
130          135          140
Asp Gly Pro Ala Leu Arg Ser Thr Ala Leu Ile Thr Tyr Gly Asn Trp
145          150          155          160
Leu Leu Ser Asn Gly Asn Thr Ser Tyr Val Thr Ser Asn Leu Trp Pro
165          170          175
Ile Ile Gln Asn Asp Leu Gly Tyr Val Val Ser Tyr Trp Asn Gln Ser
180          185          190
Thr Tyr Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr Thr
195          200          205
Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Ala
210          215          220
Ile Gly Gln Thr Ser Gln Val Ser Ser Tyr Thr Thr Gln Ala Asp Asn
225          230          235          240
Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr Ile
245          250          255
Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr Leu
260          265          270
Leu Ala Ser Ile His Thr Tyr Asp Pro Ser Ala Gly Cys Asp Ala Ala
275          280          285
Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val Tyr
290          295          300
Val Asp Ser Phe Arg Ser Val Tyr Ser Ile Asn Ser Gly Ile Ala Ser
305          310          315          320

```


-continued

```

ttcgctacca agatcgggtca gacctcctcg gtcagcagct acacaaccca agcggcgaat 720
ctactttgct ttttgagtc ttactggaac cccacttccg gatatatcac cgtaaacact 780
ggcgggtggtc ggtccggcaa ggacgccaac accctcttgg catccatcca cacttaacgac 840
cccagcggcg gctgcgatgc cagcaccctc cagccctgct cggacaaaagc cctctcgaat 900
ctgaaggttt acgctgactc cttccgttct gtctactcca tcaacagcgg tattgcctct 960
aacgcccgtg tagccactgg tcgctaccgg gaagacagct accagggcgg gaacctatgg 1020
tacctcacta cgttcgcogt cgccgagcag ctctatgacg ccctcaatgt ctgggctgct 1080
cagggctccc tcaatgtcac ctccatctcc ctcccccttct tccagcagtt ctctctagt 1140
gtcactgccc gcacttatgc ttgagctcc accaacttaca cgactctgac ctccgccatt 1200
aagagcttcc cggatggatt cgtcgtatc aacgcccagt acacgcccgc caacgggtggc 1260
ctcgtgagc agttcagcag gagcaacggc tctcccgtca gcgctgttga cttgacatgg 1320
agctatgcat ccgcaattgac cgcgtttgaa gcgaggaata ataactcagtt cgcgggtgg 1380
ggcgcggtag gtttactgt gccgacctcg tgctccagca acagtgggtgg aggcggagga 1440
tcaactgtcc cgtgacgtt caacgtgaac gcccaaacgg tttggggcga aaactctac 1500
atactggct cggttgacgc tctgagtaac tggctctctg acaacgcccct cttgctctcg 1560
tctgccaact acccgacctg gagcattacc gtgaatttac ccgagcagc tgccattcag 1620
tataagtata tcccaagaa caacggagct gtcacctggg aatccgatcc caacaacagc 1680
ataactactc cagccagcgg ctccgtgacc gagaatgaca cttggcggtta a 1731

```

<210> SEQ ID NO 16

<211> LENGTH: 576

<212> TYPE: PRT

<213> ORGANISM: Gloeophyllum trabeum

<400> SEQUENCE: 16

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Leu Gly Thr Val Leu
1           5           10           15
Ala Gln Ser Val Asp Ser Tyr Val Gly Ser Glu Gly Pro Ile Ala Lys
20          25          30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
35          40          45
Ala Ala Ala Gly Val Val Val Ala Ser Pro Ser Lys Ser Asp Pro Asp
50          55          60
Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
65          70          75          80
Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Ser
85          90          95
Leu Ile Asp Ser Phe Val Ile Ala Glu Ala Asn Ile Gln Gln Val Ser
100         105         110
Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
115         120         125
Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
130         135         140
Asp Gly Pro Ala Leu Arg Ala Thr Ala Leu Ile Thr Tyr Gly Asn Trp
145         150         155         160
Leu Leu Ser Asn Gly Asn Thr Thr Trp Val Thr Ser Thr Leu Trp Pro
165         170         175
Ile Ile Gln Asn Asp Leu Asn Tyr Val Val Gln Tyr Trp Asn Gln Thr
180         185         190

```

-continued

Thr Phe Asp Leu Trp Glu Glu Val Asn Ser Ser Ser Phe Phe Thr Thr
 195 200 205
 Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Lys
 210 215 220
 Ile Gly Gln Thr Ser Ser Val Ser Ser Tyr Thr Thr Gln Ala Ala Asn
 225 230 235 240
 Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Thr Ser Gly Tyr Ile
 245 250 255
 Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr Leu
 260 265 270
 Leu Ala Ser Ile His Thr Tyr Asp Pro Ser Ala Gly Cys Asp Ala Thr
 275 280 285
 Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val Tyr
 290 295 300
 Val Asp Ser Phe Arg Ser Val Tyr Ser Ile Asn Ser Gly Ile Ala Ser
 305 310 315 320
 Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Gln Gly
 325 330 335
 Gly Asn Pro Trp Tyr Leu Thr Thr Phe Ala Val Ala Glu Gln Leu Tyr
 340 345 350
 Asp Ala Leu Asn Val Trp Ala Ala Gln Gly Ser Leu Asn Val Thr Ser
 355 360 365
 Ile Ser Leu Pro Phe Phe Gln Gln Phe Ser Ser Ser Val Thr Ala Gly
 370 375 380
 Thr Tyr Ala Ser Ser Ser Thr Thr Tyr Thr Thr Leu Thr Ser Ala Ile
 385 390 395 400
 Lys Ser Phe Ala Asp Gly Phe Val Ala Ile Asn Ala Gln Tyr Thr Pro
 405 410 415
 Ser Asn Gly Gly Leu Ala Glu Gln Phe Ser Arg Ser Asn Gly Ser Pro
 420 425 430
 Val Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ser Ala Leu Thr Ala
 435 440 445
 Phe Glu Ala Arg Asn Asn Thr Gln Phe Ala Gly Trp Gly Ala Val Gly
 450 455 460
 Leu Thr Val Pro Thr Ser Cys Ser Ser Asn Ser Gly Gly Gly Gly Gly
 465 470 475 480
 Ser Thr Val Ala Val Thr Phe Asn Val Asn Ala Gln Thr Val Trp Gly
 485 490 495
 Glu Asn Ile Tyr Ile Thr Gly Ser Val Asp Ala Leu Ser Asn Trp Ser
 500 505 510
 Pro Asp Asn Ala Leu Leu Leu Ser Ser Ala Asn Tyr Pro Thr Trp Ser
 515 520 525
 Ile Thr Val Asn Leu Pro Ala Ser Thr Ala Ile Gln Tyr Lys Tyr Ile
 530 535 540
 Arg Lys Asn Asn Gly Ala Val Thr Trp Glu Ser Asp Pro Asn Asn Ser
 545 550 555 560
 Ile Thr Thr Pro Ala Ser Gly Ser Val Thr Glu Asn Asp Thr Trp Arg
 565 570 575

<210> SEQ ID NO 17

<211> LENGTH: 1731

<212> TYPE: DNA

<213> ORGANISM: Gloeophyllum trabeum

-continued

<400> SEQUENCE: 17

```

atgtaccgct tccttgtctg tgctctcggg cttctgggga cagtccctgc tcagtcagtc      60
gacagttatg tcggcagcga aggccccata gcaaaggccg gcgtccttgc caacattggg      120
ccgaacggct caaaggcctc tggtgacgcc gccggcgtgg tgggtggtag ccccagcaag      180
tcggatcccg actattggta cacttggacg cgtgactcgt cactcgtttt caagtctctc      240
attgatcagt acaccactgg tatcgacagc acgagttcgt tgaggtctct gatagacagt      300
ttcgttatg  cggaggccaa cattcagcag gtctctaadc ccagcggcac tcttactacc      360
ggcggccttg gagagccaaa attcaatgtc gatgaaactg cattcacccg tgcattgggt      420
cgaccccagc gcgacggacc tgcgtccgt  gcgactgctt tgatcaccta cggttaactgg      480
ctcttgtcaa acgggaacac gacctgggtt accagtagc  tgtggccgat catccagaac      540
gatctcaact acgtcgttca gtactggaac cagaccact  tcgacctctg ggaagaagtg      600
aactcttct  cgttcttcc  cactgcagtg cagcaccgtg ccttgcgcga agggcagca      660
ttcgtacca  agatcggta  gacctcctc  gtcagcagct acacaacca agcggcgaat      720
ctactttgct ttttgcagtc ttactggaac cccacttccg gatatatcac cgtaaacact      780
ggcggtggtc ggtccggcaa ggacgccaac accctcttgg catccatcca cacttacgac      840
cccagcgcgg gctcgcgatg cagcaccctc cagccctgct ccgacaaagc cctctcgaat      900
ctgaaggttt acgtcgcactc cttccgttct gtctactcca tcaacagcgg tattgcctct      960
aacgccctg  tcgccactgg tcgctaccg  gaagacagct accagggcgg gaacctatgg      1020
tacctcacta cgttcgcctg cgcgagcag ctctatgacg ccctcaatgt ctgggctgct      1080
cagggtctcc tcaatgtcac ctccatctcc ctccccctt  tccagcagtt ctctctagt      1140
gtcaactgcg gcacttaagc ttcagctcc  accacttaca cgactctgac ctccgccatt      1200
aagagcttcc cggatggatt cgtcgtatc aacgcccagt acacgccgtc caacgggtggc      1260
ctcgtgagc  agttcagcag gagcaacggc gctcccgtca gcgctgtga tttgacatgg      1320
agctatgcat ctgcattgac cgcggttgaa gcgaggaata ataactcagtt cgcggctgg      1380
ggcgcggtag gtttactgt  gccacctc  tgctccagca acagtgggtg aggcggagga      1440
tcgactgtcg ccgtgacgtt caactggaac gcccaaacgg tttggggcga aaacatctac      1500
atcaactggc cggttgacgc tctgagtaac tggctctccg acaacgccct cttgctctcg      1560
tctgccaact acccgacctg gagcattacc gtgaatttac ccgcgagcac tgccattcag      1620
tataagtata tccgcaagaa caacggagct gtcacctggg aatccgatcc caacaacagc      1680
ataactactc cagccagcgg ctccgtgacc gagaatgaca cttggcggtta a      1731

```

<210> SEQ ID NO 18

<211> LENGTH: 576

<212> TYPE: PRT

<213> ORGANISM: Gloeophyllum trabeum

<400> SEQUENCE: 18

```

Met Tyr Arg Phe Leu Val Cys Ala Leu Gly Leu Leu Gly Thr Val Leu
1           5           10          15
Ala Gln Ser Val Asp Ser Tyr Val Gly Ser Glu Gly Pro Ile Ala Lys
20          25          30
Ala Gly Val Leu Ala Asn Ile Gly Pro Asn Gly Ser Lys Ala Ser Gly
35          40          45
Ala Ala Ala Gly Val Val Val Ala Ser Pro Ser Lys Ser Asp Pro Asp
50          55          60

```

-continued

Tyr Trp Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe Lys Ser Leu
 65 70 75 80
 Ile Asp Gln Tyr Thr Thr Gly Ile Asp Ser Thr Ser Ser Leu Arg Ser
 85 90 95
 Leu Ile Asp Ser Phe Val Ile Ala Glu Ala Asn Ile Gln Gln Val Ser
 100 105 110
 Asn Pro Ser Gly Thr Leu Thr Thr Gly Gly Leu Gly Glu Pro Lys Phe
 115 120 125
 Asn Val Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln Arg
 130 135 140
 Asp Gly Pro Ala Leu Arg Ala Thr Ala Leu Ile Thr Tyr Gly Asn Trp
 145 150 155 160
 Leu Leu Ser Asn Gly Asn Thr Thr Trp Val Thr Ser Thr Leu Trp Pro
 165 170 175
 Ile Ile Gln Asn Asp Leu Asn Tyr Val Val Gln Tyr Trp Asn Gln Thr
 180 185 190
 Thr Phe Asp Leu Trp Glu Glu Val Asn Ser Ser Ser Phe Phe Thr Thr
 195 200 205
 Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Ala Phe Ala Thr Lys
 210 215 220
 Ile Gly Gln Thr Ser Ser Val Ser Ser Tyr Thr Thr Gln Ala Ala Asn
 225 230 235 240
 Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Thr Ser Gly Tyr Ile
 245 250 255
 Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr Leu
 260 265 270
 Leu Ala Ser Ile His Thr Tyr Asp Pro Ser Ala Gly Cys Asp Ala Thr
 275 280 285
 Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val Tyr
 290 295 300
 Val Asp Ser Phe Arg Ser Val Tyr Ser Ile Asn Ser Gly Ile Ala Ser
 305 310 315 320
 Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Gln Gly
 325 330 335
 Gly Asn Pro Trp Tyr Leu Thr Thr Phe Ala Val Ala Glu Gln Leu Tyr
 340 345 350
 Asp Ala Leu Asn Val Trp Ala Ala Gln Gly Ser Leu Asn Val Thr Ser
 355 360 365
 Ile Ser Leu Pro Phe Phe Gln Gln Phe Ser Ser Ser Val Thr Ala Gly
 370 375 380
 Thr Tyr Ala Ser Ser Ser Thr Thr Tyr Thr Thr Leu Thr Ser Ala Ile
 385 390 395 400
 Lys Ser Phe Ala Asp Gly Phe Val Ala Ile Asn Ala Gln Tyr Thr Pro
 405 410 415
 Ser Asn Gly Gly Leu Ala Glu Gln Phe Ser Arg Ser Asn Gly Ala Pro
 420 425 430
 Val Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ser Ala Leu Thr Ala
 435 440 445
 Phe Glu Ala Arg Asn Asn Thr Gln Phe Ala Gly Trp Gly Ala Val Gly
 450 455 460
 Leu Thr Val Pro Thr Ser Cys Ser Ser Asn Ser Gly Gly Gly Gly Gly
 465 470 475 480

-continued

```

gagctgggca actggtccac ggccagcgcg atccccctcc gcgcggatgc ttacaccaac 1680
agcaaccgcg tctggtactg gaccgtcaat ctgccccctg gcaccagctt cgagtacaag 1740
ttcttcaaga accagacgga cgggaccatc gtctgggaag acgacccgaa cgggtcgtac 1800
acggtcccag cgtactgtgg gcagactacc gccattcttg acgatagttg gcagtga 1857

```

```

<210> SEQ ID NO 20
<211> LENGTH: 618
<212> TYPE: PRT
<213> ORGANISM: Talaromyces emersonii

```

```

<400> SEQUENCE: 20

```

```

Met Ala Ser Leu Val Ala Gly Ala Leu Cys Ile Leu Gly Leu Thr Pro
1           5           10           15
Ala Ala Phe Ala Arg Ala Pro Val Ala Ala Arg Ala Thr Gly Ser Leu
20          25          30
Asp Ser Phe Leu Ala Thr Glu Thr Pro Ile Ala Leu Gln Gly Val Leu
35          40          45
Asn Asn Ile Gly Pro Asn Gly Ala Asp Val Ala Gly Ala Ser Ala Gly
50          55          60
Ile Val Val Ala Ser Pro Ser Arg Ser Asp Pro Asn Tyr Phe Tyr Ser
65          70          75          80
Trp Thr Arg Asp Ala Ala Leu Thr Ala Lys Tyr Leu Val Asp Ala Phe
85          90          95
Ile Ala Gly Asn Lys Asp Leu Glu Gln Thr Ile Gln Gln Tyr Ile Ser
100         105        110
Ala Gln Ala Lys Val Gln Thr Ile Ser Asn Pro Ser Gly Asp Leu Ser
115         120        125
Thr Gly Gly Leu Gly Glu Pro Lys Phe Asn Val Asn Glu Thr Ala Phe
130         135        140
Thr Gly Pro Trp Gly Arg Pro Gln Arg Asp Gly Pro Ala Leu Arg Ala
145         150        155        160
Thr Ala Leu Ile Ala Tyr Ala Asn Tyr Leu Ile Asp Asn Gly Glu Ala
165         170        175
Ser Thr Ala Asp Glu Ile Ile Trp Pro Ile Val Gln Asn Asp Leu Ser
180         185        190
Tyr Ile Thr Gln Tyr Trp Asn Ser Ser Thr Phe Asp Leu Trp Glu Glu
195         200        205
Val Glu Gly Ser Ser Phe Phe Thr Thr Ala Val Gln His Arg Ala Leu
210         215        220
Val Glu Gly Asn Ala Leu Ala Thr Arg Leu Asn His Thr Cys Ser Asn
225         230        235        240
Cys Val Ser Gln Ala Pro Gln Val Leu Cys Phe Leu Gln Ser Tyr Trp
245         250        255
Thr Gly Ser Tyr Val Leu Ala Asn Phe Gly Gly Ser Gly Arg Ser Gly
260         265        270
Lys Asp Val Asn Ser Ile Leu Gly Ser Ile His Thr Phe Asp Pro Ala
275         280        285
Gly Gly Cys Asp Asp Ser Thr Phe Gln Pro Cys Ser Ala Arg Ala Leu
290         295        300
Ala Asn His Lys Val Val Thr Asp Ser Phe Arg Ser Ile Tyr Ala Ile
305         310        315        320
Asn Ser Gly Ile Ala Glu Gly Ser Ala Val Ala Val Gly Arg Tyr Pro
325         330        335

```

-continued

Glu Asp Val Tyr Gln Gly Gly Asn Pro Trp Tyr Leu Ala Thr Ala Ala
 340 345 350
 Ala Ala Glu Gln Leu Tyr Asp Ala Ile Tyr Gln Trp Lys Lys Ile Gly
 355 360 365
 Ser Ile Ser Ile Thr Asp Val Ser Leu Pro Phe Phe Gln Asp Ile Tyr
 370 375 380
 Pro Ser Ala Ala Val Gly Thr Tyr Asn Ser Gly Ser Thr Thr Phe Asn
 385 390 395 400
 Asp Ile Ile Ser Ala Val Gln Thr Tyr Gly Asp Gly Tyr Leu Ser Ile
 405 410 415
 Val Glu Lys Tyr Thr Pro Ser Asp Gly Ser Leu Thr Glu Gln Phe Ser
 420 425 430
 Arg Thr Asp Gly Thr Pro Leu Ser Ala Ser Ala Leu Thr Trp Ser Tyr
 435 440 445
 Ala Ser Leu Leu Thr Ala Ser Ala Arg Arg Gln Ser Val Val Pro Ala
 450 455 460
 Ser Trp Gly Glu Ser Ser Ala Ser Ser Val Pro Ala Val Cys Ser Ala
 465 470 475 480
 Thr Ser Ala Thr Gly Pro Tyr Ser Thr Ala Thr Asn Thr Val Trp Pro
 485 490 495
 Ser Ser Gly Ser Gly Ser Ser Thr Thr Thr Ser Ser Ala Pro Cys Thr
 500 505 510
 Thr Pro Thr Ser Val Ala Val Thr Phe Asp Glu Ile Val Ser Thr Ser
 515 520 525
 Tyr Gly Glu Thr Ile Tyr Leu Ala Gly Ser Ile Pro Glu Leu Gly Asn
 530 535 540
 Trp Ser Thr Ala Ser Ala Ile Pro Leu Arg Ala Asp Ala Tyr Thr Asn
 545 550 555 560
 Ser Asn Pro Leu Trp Tyr Val Thr Val Asn Leu Pro Pro Gly Thr Ser
 565 570 575
 Phe Glu Tyr Lys Phe Phe Lys Asn Gln Thr Asp Gly Thr Ile Val Trp
 580 585 590
 Glu Asp Asp Pro Asn Arg Ser Tyr Thr Val Pro Ala Tyr Cys Gly Gln
 595 600 605
 Thr Thr Ala Ile Leu Asp Asp Ser Trp Gln
 610 615

<210> SEQ ID NO 21

<211> LENGTH: 1725

<212> TYPE: DNA

<213> ORGANISM: Trametes cingulata

<400> SEQUENCE: 21

```

atgcgtttca cgctcctcac ctccctcctg gccctcgccc tcggcgcggt cgcgcagtcg      60
agtgcggcgc acgcgtacgt cgcgtccgaa tcgcccacgc ccaaggcggg tgtgctcgcc      120
aacatcgggc ccagcggctc caagtccaac ggagcaaagg caggcatcgt gattgcaagt      180
ccgagcacat ccaaccggaa ctacctgtac acatggagcc gcgactcgtc cctcgtgttc      240
aaggcgctca tcgaccagtt caccactggc gaagatacct cgctccgaac tctgattgac      300
gagttcacct cggcggaggg catactccag caggtgccga acccgagcgg gacagtcagc      360
actggaggcc tcggcgagcc caagttcaac atcgacgaga ccgcggtcac ggatgcctgg      420
ggtcgtcctc agcgcgatgg tcccgtcttc cgggcgactg ccatcatcac ctacgccaac      480

```

-continued

```

tggctcctcg acaacaagaa cacgacctac gtgaccaaca ctctctggcc tatcatcaag 540
ctcgacctcg actacgtcgc cagcaactgg aaccagtcca cgtttgatct ctgggaggag 600
attaactcct cgtcgttctt cactacggcc gtccagcacc gtgctctgcg cgagggcgcg 660
actttcgcta atcgcatcgg acaaacctcg gtggtcagcg ggtacaccac ccaagcaaac 720
aaccttctct gcttctcgca gtctactggt aacccccacc gcggctatat caccgcaaac 780
acggggcgcg gccgctctgg caaggacgcg aacaccgttc tcacgtcgat ccacaccttc 840
gaccggcgcg ctggatgoga cgtgtttacg ttccagccgt gctcggacaa ggcgctgctg 900
aacttgaagg tgtacgtoga tgcgttcocg tcgatctact ccatcaacag cgggatcgcc 960
tcgaatgcgg ccggttgctac cggccgctac cccgaggaca gctacatggg cggaaacca 1020
tggtaacctca ccacctcgcg cgtcgtgag cagctctacg atgcgctcat tgtgtggaac 1080
aaacttggcg ccctgaaagt cagcagcacc tccctcccct tcttcagca gttctcgta 1140
ggcgtcacgg tcggcaccta tgcctcatcc tcgtccacct tcaagacgct cacttccgcc 1200
atcaagacct tcgccagcgg cttcctcgcg gtcaacgcca agtacacgcc ctogaacggc 1260
ggccttgctg aacagtacag ccggagcaac ggctcgcggc tcagcgtgtg ggacctgacg 1320
tggagctatg ctgctgcctc cagctcgttt gctgcgcgct caggcaagac gtatgcgagc 1380
tggggcgcgg cgggtttgac tgtcccagc acttgctcgg ggagtggcgg tgctgggact 1440
gtggccgta ccttcaacgt gcaggcgacc accgtgttcg gcgagaacat ttacatcaca 1500
ggctcgttcc ccgctctcca gaactggctg cccgacaacg cgctcatcct ctcagcggcc 1560
aactaccca cttggagcat caccgtgaac ctgccggcga gcacgacgat cgagtacaag 1620
tacattcgca agttcaacgg cgcggtcacc tgggagtcgg acccgaacaa ctgatcacg 1680
acgcccgcga gcggcacggt caccacagaac gacacctggc ggtag 1725

```

<210> SEQ ID NO 22

<211> LENGTH: 574

<212> TYPE: PRT

<213> ORGANISM: *Trametes cingulata*

<400> SEQUENCE: 22

```

Met Arg Phe Thr Leu Leu Thr Ser Leu Leu Gly Leu Ala Leu Gly Ala
1           5           10           15
Phe Ala Gln Ser Ser Ala Ala Asp Ala Tyr Val Ala Ser Glu Ser Pro
20           25           30
Ile Ala Lys Ala Gly Val Leu Ala Asn Ile Gly Pro Ser Gly Ser Lys
35           40           45
Ser Asn Gly Ala Lys Ala Gly Ile Val Ile Ala Ser Pro Ser Thr Ser
50           55           60
Asn Pro Asn Tyr Leu Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe
65           70           75           80
Lys Ala Leu Ile Asp Gln Phe Thr Thr Gly Glu Asp Thr Ser Leu Arg
85           90           95
Thr Leu Ile Asp Glu Phe Thr Ser Ala Glu Ala Ile Leu Gln Gln Val
100          105          110
Pro Asn Pro Ser Gly Thr Val Ser Thr Gly Gly Leu Gly Glu Pro Lys
115          120          125
Phe Asn Ile Asp Glu Thr Ala Phe Thr Asp Ala Trp Gly Arg Pro Gln
130          135          140

```

-continued

Arg Asp Gly Pro Ala Leu Arg Ala Thr Ala Ile Ile Thr Tyr Ala Asn
 145 150 155 160

Trp Leu Leu Asp Asn Lys Asn Thr Thr Tyr Val Thr Asn Thr Leu Trp
 165 170 175

Pro Ile Ile Lys Leu Asp Leu Asp Tyr Val Ala Ser Asn Trp Asn Gln
 180 185 190

Ser Thr Phe Asp Leu Trp Glu Glu Ile Asn Ser Ser Ser Phe Phe Thr
 195 200 205

Thr Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Thr Phe Ala Asn
 210 215 220

Arg Ile Gly Gln Thr Ser Val Val Ser Gly Tyr Thr Thr Gln Ala Asn
 225 230 235 240

Asn Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Thr Gly Gly Tyr
 245 250 255

Ile Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr
 260 265 270

Val Leu Thr Ser Ile His Thr Phe Asp Pro Ala Ala Gly Cys Asp Ala
 275 280 285

Val Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val
 290 295 300

Tyr Val Asp Ala Phe Arg Ser Ile Tyr Ser Ile Asn Ser Gly Ile Ala
 305 310 315 320

Ser Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Met
 325 330 335

Gly Gly Asn Pro Trp Tyr Leu Thr Thr Ser Ala Val Ala Glu Gln Leu
 340 345 350

Tyr Asp Ala Leu Ile Val Trp Asn Lys Leu Gly Ala Leu Asn Val Thr
 355 360 365

Ser Thr Ser Leu Pro Phe Phe Gln Gln Phe Ser Ser Gly Val Thr Val
 370 375 380

Gly Thr Tyr Ala Ser Ser Ser Ser Thr Phe Lys Thr Leu Thr Ser Ala
 385 390 395 400

Ile Lys Thr Phe Ala Asp Gly Phe Leu Ala Val Asn Ala Lys Tyr Thr
 405 410 415

Pro Ser Asn Gly Gly Leu Ala Glu Gln Tyr Ser Arg Ser Asn Gly Ser
 420 425 430

Pro Val Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ala Ala Leu Thr
 435 440 445

Ser Phe Ala Ala Arg Ser Gly Lys Thr Tyr Ala Ser Trp Gly Ala Ala
 450 455 460

Gly Leu Thr Val Pro Thr Thr Cys Ser Gly Ser Gly Gly Ala Gly Thr
 465 470 475 480

Val Ala Val Thr Phe Asn Val Gln Ala Thr Thr Val Phe Gly Glu Asn
 485 490 495

Ile Tyr Ile Thr Gly Ser Val Pro Ala Leu Gln Asn Trp Ser Pro Asp
 500 505 510

Asn Ala Leu Ile Leu Ser Ala Ala Asn Tyr Pro Thr Trp Ser Ile Thr
 515 520 525

Val Asn Leu Pro Ala Ser Thr Thr Ile Glu Tyr Lys Tyr Ile Arg Lys
 530 535 540

Phe Asn Gly Ala Val Thr Trp Glu Ser Asp Pro Asn Asn Ser Ile Thr
 545 550 555 560

-continued

Thr Pro Ala Ser Gly Thr Phe Thr Gln Asn Asp Thr Trp Arg
 565 570

The invention claimed is:

1. A recombinant host cell comprising a nucleic acid construct comprising a polynucleotide encoding a glucoamylase according to the invention operably linked to one or more control sequences that direct the production of the polypeptide in the expression host, wherein the host cell is a yeast cell and wherein the glucoamylase is selected from the group consisting of:

a polypeptide having glucoamylase activity, having at least 90% identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, or SEQ ID NO: 18.

2. A recombinant host cell comprising a nucleic acid construct comprising a polynucleotide encoding a glucoamylase comprising a catalytic domain according to the invention operably linked to one or more control sequences that direct the production of the glucoamylase comprising a catalytic domain polypeptide in the expression host, wherein the host cell is a yeast cell and wherein the glucoamylase catalytic domain is selected from the group consisting of:

a polypeptide having at least 90% identity, to the sequence of amino acids 19 to 474 of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14 or amino acids 19 to 471 of SEQ ID NO: 16 or SEQ ID NO: 18.

3. The recombinant host cell according to claim 1, wherein the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

4. The recombinant host cell according to claim 3, wherein the yeast host cell is a *Saccharomyces* cell.

5. The recombinant host cell according to claim 4, wherein the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* cell.

6. The recombinant host cell according to claim 5, wherein the yeast host cell is a *Saccharomyces cerevisiae* cell.

7. The recombinant host cell according to claim 6, wherein the polypeptide having glucoamylase activity is selected from SEQ ID NO: 14 or a polypeptide having at least 90% identity to the mature polypeptide of SEQ ID NO: 14.

8. The recombinant host cell according to claim 6, wherein the glucoamylase catalytic domain is selected from amino acids 19 to 474 SEQ ID NO: 14 or a polypeptide having at least 90% identity to amino acids 19 to 474 of SEQ ID NO: 14.

* * * * *