



(51) International Patent Classification:

A61P 35/04 (2006.01) G01N 33/574 (2006.01)
C12Q 1/25 (2006.01)

(21) International Application Number:

PCT/US2013/040917

(22) International Filing Date:

14 May 2013 (14.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/646,586 14 May 2012 (14.05.2012) US
61/646,593 14 May 2012 (14.05.2012) US

(72) Inventor; and

(71) Applicant : PESTELL, Richard, G. [US/US]; 901 North
Penn Street, Apt. R1902, Philadelphia, PA 19123 (US).

(74) Agents: UDAL, Robert, P. et al.; Morgan, Lewis &
Bockius LLP, 1701 Market Street, Philadelphia, PA 19103
(US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: USING MODULATORS OF CCR5 FOR TREATING CANCER

(57) Abstract: This disclosure is directed, in part, to a method of determining whether a subject having cancer is at risk for developing metastasis of the cancer. In one embodiment, the method comprises (a) obtaining a biological sample from the subject having cancer; (b) determining CCR5 expression level and/or expression level of at least one of CCR5 ligands in the biological sample; and (c) if the expression level of CCR5 and/or of at least one of CCR5 ligands determined in step (b) is increased compared to CCR5 expression level and/or expression level of at least one of CCR5 ligands in a control sample, then the subject is identified as likely at risk for developing metastasis of the cancer.



USING MODULATORS OF CCR5 FOR TREATING CANCER

[0001] Use Of Modulators Of CCR5 In The Treatment Of Cancer And Cancer Metastasis

CROSS-REFERENCE TO RELATED APPLICATIONS

[0002] This application claims the benefit of U.S. Provisional Patent Application Nos. 61/646,586, filed May 14, 2012, and entitled "Use Of Modulators Of CCR5 In The Treatment Of Cancer And Cancer Metastasis"; and 61/646,593, filed May 14, 2012, and entitled "Use Of Modulators Of CCR5 In The Treatment Of Cancer And Cancer Metastasis", both of which are incorporated by reference herein in their entirety.

FIELD OF THE INVENTION

[0003] This disclosure is directed, in part, to a method of determining whether a subject having cancer is at risk for developing metastasis of the cancer and a method of blocking cancer methastasis.

STATEMENT REGARDING FEDERALLY SPONSORED R&D

[0004] This invention was supported in part by PASPA-UNAM (M.A.V-V.), NIH grants R01CA070896, R01CA075503, R01CA132115, R01CA107382, R01CA086072 (R.G.P.), R01CA120876 (M.P.L), the Kimmel Cancer Center NIH Cancer Center Core grant P30CA056036 (R.G.P.), generous grants from the Dr. Ralph and Marian C. Falk Medical Research Trust and the Margaret Q. Landenberger Research Foundation, and a grant from Pennsylvania Department of Health (R.G.P.). Accordingly, the United States government has certain rights in the invention.

BACKGROUND OF THE INVENTION

[0005] Breast cancer causes the death of 40,000 women in the USA and 410,000 women in the world annually.¹ Despite advances in the treatment of the disease, 20% to 30% of patients with early breast cancers will experience relapse with distant metastatic disease.² In those patients, metastasis is the main cause of death. Patients with basal tumors have increased risk of metastasis and lower survival rate.^{3,4} Kennecke *et al.* studied 3,726 breast cancer patients and reported that the basal tumors have higher frequencies of metastases and reduced time from identification of metastases to death compared to that of patients with luminal A or B tumors.⁴ The absence of AR,

ER, and HER-2 commonly found in basal breast tumors⁵ means that they are unlikely to respond to hormone therapies or HER-2 targeted therapies. Currently, chemotherapy, radiation, and surgery are the only choices for patients with basal breast cancers, but all demonstrate poor outcomes.⁶ The need for a specific targeted therapy for basal breast cancer remains urgent.

5

BRIEF SUMMARY OF THE INVENTION

[0006] Certain aspects of embodiments disclosed herein by way of example are summarized below. It should be understood that these aspects are presented merely to provide the reader with a brief summary of certain forms an invention disclosed and/or claimed herein might take and that these aspects are not intended to limit the scope of any invention disclosed and/or claimed herein.

10 Indeed, any invention disclosed and/or claimed herein may encompass a variety of aspects that may not be set forth below.

[0007] The present disclosure generally relates to various methods, including methods of determining whether a subject has cancer or is at risk for developing cancer and/or is at risk for developing cancer metastasis. In some embodiments, the methods of the present invention include methods of treating, preventing, or managing a neoplasm or a cancer metastasis in a patient. In some embodiments, the methods of the present invention include in vivo methods for down regulating CCR5 expression in a set of one or more prostate cancer cell lines derived from transduction of murine epithelial or prostate epithelial cells and transformed by at least one oncogene selected from the group consisting of NeuT, Ha-Ras, and c-Src. In some exemplary
15
20
25
embodiments drugs that target the HIV receptor CCR5, which the virus uses to enter and infect host cells, are used to prevent migration and spread of cancer cells from their primary tissue to secondary sites in the body of the patient. In one embodiment, a drug that targets the HIV receptor CCR5 may be used as an adjuvant therapy, or adjuvant care, wherein the drug is administered to the patient in addition to a primary, main or initial treatment for cancer. In some embodiments, such adjuvant therapy may be administered concomitantly or concurrently as other therapies for cancer. In one embodiment, such adjuvant therapy may be given concurrently as other adjuvant therapies or following other adjuvant therapies.

[0008] In some embodiments, when CCR5 receptor antagonists are used as adjuvant therapy they improve prognosis in cancer patients. In some embodiments, where a plurality of anticancer
30
drugs are used in combination for cancer therapy, a CCR5 receptor antagonist may be included as an adjuvant therapy in order to improve the therapeutic effect by blocking metastasis of the cancer being treated, thereby contributing to improving clinical outcomes of the cancer therapy.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0009] The foregoing summary, as well as other features, aspects, and advantages of the present invention and the following detailed description of embodiments of the invention, will be better understood when read in conjunction with the claims and the appended drawings of an exemplary
 5 embodiments, wherein:

[0010] Figure 1A illustrates a heatmap of the expression of CCL5 and its receptor CCR5 in samples from patients with breast cancer divided by molecular subtype (luminal A, luminal B, basal, normal-like, and Her-2) based on their gene expression pattern;

[0011] Figures 1B1 to 1B5 illustrate scatter plots and correlation analysis (Student *t* test) of the
 10 expression of CCL5 and CCR5 among the breast cancer molecular subtypes of luminal A, luminal B, basal, normal-like, and Her-2;

[0012] Figure 1C illustrates quantification of the proportions of the breast cancer samples overexpressing CCL5 and CCR5, fraction of the bars representing upper right quadrants of the scatter plots shown in Figure B;

15 [0013] Figure 1D illustrates metastasis-free Kaplan–Meier plots and log-rank analysis for the different molecular subtypes of breast cancer in the analyzed database, which is described in Materials and Methods section of this of this application;

[0014] Figure 2A illustrates flow cytometric histograms of the CCR5 expression in MDA-MB-231 breast cancer cells identified a subpopulation of CCR5⁺ cells;

20 [0015] Figures 2B1 and 2B2 illustrate induction of calcium signaling in cells loaded with Fluo-4-AM before the sequential addition of CCL5 (60 µg/mL) and FBS (5%);

[0016] Figure 2C illustrates 3D invasion into collagen gels by breast cancer cell lines, using CCL5 (15 µg/mL) as chemoattractant;

[0017] Figure 2D illustrates mean distances of invasion ± SEM from 3 independent experiments;

25 [0018] Figure 2E, 3D invasion assays for MCF-10A cells and MCF-10A-NeuT, -Ras, and -Src derivatives showing that CCL5-induced invasion is activated by oncogenic transformation;

[0019] Figure 2F illustrates quantification (F, mean ± SEM, n = 3) 3D invasion assays for MCF-10A cells and MCF-10A-NeuT, -Ras, and -Src derivatives showing that CCL5-induced
 30 invasion is activated by oncogenic transformation shown in Figure 2E;

[0020] Figure 2G illustrates CCR5⁺ and CCR5[−] subpopulations from SUM-159 cell line that

were isolated by FACS and their respected invasion into collagen gels evaluated using fetal bovine serum (FBS) as chemoattractant;

[0021] Figure 2H illustrates quantification of the invasion of CCR5+ and CCR5- subpopulations into collagen as mean $\pm$ SEM of the two independent experiments shown in Figure 2G;

[0022] Figures 3A1 to 3A4 illustrate(s) intensity versus time analysis of Fluo-4 AM-loaded MDA-MB-231 cells treated with the CCR5 antagonists maraviroc or vicriviroc (100 nmol/L) for 30 minutes before the addition of CCL5 (60 μ g/mL);

[0023] Figure 3B illustrates a comparison of the fraction of cells with increased fluorescence intensity upon addition of CCL5;

[0024] Figure 3C1 to 3C4 show CCL5-induced calcium signaling blocked by CCR5 antagonists in Hs578T cells;

[0025] Figure 3D illustrates quantification (mean $\pm$ SEM) of 3 to 4 independent experiments shown in Figures 3A-3C;

[0026] Figure 4A illustrates 3D reconstruction of FBS-induced invasion into collagen gels by Hs578T breast cancer cells in presence of CCR5 antagonists (100 nmol/L);

[0027] Figure 4B illustrates quantifications (mean $\pm$ SEM, $n = 3$) and analysis (Bonferroni t test) of the FBS-induced invasion into collagen gels by Hs578T breast cancer cells shown in Figure 4A;

[0028] Figure 4C illustrates 3D reconstruction of FBS-induced invasion into collagen gels by SUM-159 breast cancer cells in presence of CCR5 antagonists (100 nmol/L);

[0029] Figure 4D illustrates quantifications (mean $\pm$ SEM, $n = 3$) and analysis (Bonferroni t test) of the FBS-induced invasion into collagen gels by SUM-159 breast cancer cells shown in Figure 4A;

[0030] Figure 5A shows exemplary in vivo bioluminescent images (BLIs) of vehicle- or maraviroc-treated (8 mg/kg every 12 hours) of nonobese diabetic (NOD)/severe combined immunodeficiency (SCID) mice (hereinafter abbreviated as NOD/SCID mice);

[0031] Figure 5B shows quantification (mean $\pm$ SEM, $n = 6$) of in vivo bioluminescent images (BLIs) in the control (vehicle-treated NOD/SCID mice, red line/squares) and treated groups (maraviroc-treated NOD/SCID mice, blue line/triangles);

[0032] Figures 5C1 and 5C2 show the presence of pulmonary tumors and the differences between treatments corroborated by ex vivo imaging (left) and India ink staining (right);

[0033] Figure 5D showing the fraction of mice with metastatic tumors being significantly larger

in the control group ($P < 0.0001$, Fisher exact test);

[0034] Figure 5E showing histological analysis (hematoxylin and eosin staining, $\times 100$) of the area covered by metastatic tumors in lung slides;

[0035] Figure 5F shows quantification of the area covered by metastatic tumors in lung slides shown in Figure 5E;

[0036] Figure 6A, effect of CCR5 antagonist on breast cancer cell viability. MDA-MB-231 cells were exposed to increasing concentrations of maraviroc (inverted triangles) or vicriviroc (squares) for 48 hours and the cell viability was evaluated by MTT assay.

[0037] Figure 6B illustrates flow cytometric analysis showing CCR5 expression in MDA-MB-231 cells stably transfected with pcDNA3.1⁺/Zeo⁺ (MDA.Vector) or human CCR5 cloned into pCDNA3⁺Zeo⁺ (MDA.CCR5).

[0038] Figure 6C illustrates a comparison of *in vitro* proliferation rates of MDA-MB-231 in MDA.Vector and MDA.CCR5;

[0039] Figure 6D illustrates evaluation of the *in vivo* effect of maraviroc on growth of established metastasis in mice, wherein treatment of mice was initiated 10 days after injection of MDA.pFULG cells as illustrated, and *in vivo* bioluminescence imaging (BLI) of the mice for evaluating the efficacy of the treatment was carried out in days: 0, 10, 17, 24, 31 and 38;

[0040] Figure 6E illustrates quantification (mean $\pm$ SEM, $n = 5$) of *in vivo* BLI in the control (red/squares) and treated groups (blue/triangles) showed no differences in the growth rate;

[0041] Figure 6F illustrates an schema of the experimental design used to evaluate CCR5 role in lung colonization by MDA-MB-231 cells;

[0042] Figure 6G illustrates exemplary confocal images of eGFP⁺ cells in lungs of mice 24 hours after injection of the mice with MDA.pFULG cells, wherein cells expressing eGFP were counted in 3 random fields of 2 different histologic sections (separated 700 μ m from each other) per mouse ($n = 5$ mice per group);

[0043] Figure 6H illustrates quantification (H) of the number of eGFP⁺ cells in lungs of mice 24 hours after injection of the mice with MDA.pFULG cells, wherein cells expressing eGFP were counted in 3 random fields of 2 different histologic sections (separated 700 μ m from each other) per mouse ($n = 5$ mice per group);

[0044] Figure 7A illustrates GSEA analysis using KEGG and GO of tumor samples to determine the gene expression signaling pathway associated with enrichment of CCR5 and CCL5,

wherein genes in the examined breast cancer dataset are ranked by a signal-to-noise metric representing their differential expression in highest 2.5% CCL5/CCR5-expressing samples versus lowest (2.5%) CCL5/CCR5 expressing samples (N=54), depicted with a color gradient, where red indicates positive correlation with CCL5/CCR5 expression and blue represents negative correlation with CCL5/CCR5 expression;

[0045] Figures 7B1 to 7B5 illustrate scatter plots and correlation analysis using students' *t*-tests for the expression of CCR1 and CCR5 among the breast cancer molecular subtypes from a set of 2250 human breast cancer data shown in Figure 1A;

[0046] Figures 7C1 to 7C5 and show scatter plots and correlation analysis for the expression of CCR3 and CCL5 among the breast cancer molecular subtypes from a set of 2250 human breast cancer data shown in Figure 1A;

[0047] Figures 7D1 and 7D2 show Kaplan Meier curves for patients with breast cancer samples enriched for the highest level of CCR5 (up of 50%) versus the lower level CCR5 expression (lower 50%);

[0048] Figure 8A illustrates heat map of CCL5 expression and its receptors CCR5, CCR1 and CCR3 in healthy population.

[0049] Figure 8B illustrates heat map representing cross correlation of CCL5 expression and its receptors CCR5, CCR1 and CCR3 amongst healthy population;

[0050] Figures 8C and 8D illustrate heat map representing cross correlation of CCL5 expression and its receptors CCR5, CCR1 and CCR3 amongst breast cancer patients derived from a collection of 2550 breast cancers;

[0051] Figure 9A illustrates flow cytometry plot of the CCR5 expression in HS578T cells stably transfected with allophycocyanin (APC)-labeled antibody to CCR5;

[0052] Figures 9B1 and 9B2 illustrate induction of calcium signaling in HS578T cells loaded with Fluo-4-AM before the sequential addition of CCL5 (60 µg/mL) and fetal bovine serum (FBS) (5%), wherein RFI represents relative fluorescence intensities. The red traces are of cells that did not respond to the addition of CCL5 ligand-but do respond to FBS (fetal bovine serum), the green line is the trace for the cells that do respond to the addition of CCL5, and also respond to FBS. This data tells us that the MDA-MB-231 cells are a population of cells in which some cells have the receptor and respond to CCL5- and some cells that do not have the CCR5 receptor and do not respond to its ligand CCL5;

[0053] Figure 9C illustrates flow cytometry plot of the CCR5 expression in SUM159 cells stably

transfected with allophycocyanin (APC)-labeled antibody to CCR5.

[0054] Figures 9D1 and 9D2 illustrate induction of calcium signaling in SUM159 cells loaded with Fluo-4-AM before the sequential addition of CCL5 (60 $\mu\text{g/mL}$) and fetal bovine serum (FBS) (5%), wherein RFI represents relative fluorescence intensities. The red traces are of cells that did not respond to the addition of CCL5 ligand-but do respond to FBS (fetal bovine serum), the green line is the trace for the cells that do respond to the addition of CCL5, and also respond to FBS. This data tells us that the SUM-159 cells are a population of cells in which some cells have the receptor and respond to CCL5- and some cells that do not have the CCR5 receptor and do not respond to its ligand CCL5;

[0055] Figures 10A1 and 10A2 and illustrates fluorescence-activated cell sorting (FACS) analysis for the abundance of CCR5 receptor in either the SUM159-vector control cells or SUM159 cells stably overexpressing the CCR5 receptor. APC-labeled antibody to CCR5 was used to follow the CCR5 positive cells;

[0056] Figure 10B illustrates schematic representation of the timeline for induction of calcium signaling in which CCL5 is added to the SUM159-vector control cells or the SUM159 cells stably overexpressing the CCR5 receptor at 60 seconds and FBS is added at 320 seconds;

[0057] Figure 10C illustrates induction of calcium signaling (the Ca^{+2} response to CCL5 versus FBS) in the SUM159-vector control cells loaded with Fluo-4-AM before and upon the addition of CCL5 or FBS;

[0058] Figure 10D illustrates the average fluorescence for the population of SUM159-vector control cells;

[0059] Figure 10E illustrates induction of calcium signaling (the Ca^{+2} response to CCL5 versus FBS) in the SUM159 cells stably overexpressing the CCR5 receptor at 60 seconds and FBS is added at 320 second;

[0060] Figure 10F illustrates the average fluorescence for the population of SUM159-CCR5 cells;

[0061] Figure 11A illustrates that fact that maraviroc reduces MDA-MB-231 breast cancer lung metastasis burden as evidenced by the quantification results of weekly BLI, conducted for 5 weeks on maraviroc-treated NOD/SCID mice, wherein the lung metastatic tumor radiance antemortem was used as a surrogate measurement of tumor burden to the lungs (per Figure 5B).

[0062] Figure 11B illustrates the fact that maraviroc reduces MDA-MB-231 breast cancer lung

metastasis burden as evidenced by the quantification illustrates results of weekly BLI, conducted for 5 weeks on vehicle-treated NOD/SCID mice, wherein the lung metastatic tumor radiance antemortem was used as a surrogate measurement of tumor burden.

[0063] Figures 12A to 12F illustrate that fact that pulmonary tumors are enriched in CCR5⁺ cells as evidenced by the FACS analysis of the proportion of CCR5⁺ cells into metastatic tumors caused by tail vein injection of MDA.pFULG cells. The proportion of CCR5⁺ cells within eGFP⁺ MDA.pFULG cells in culture (Figure 12A) was compared with that of cells isolated from metastatic tumors (Figure 12B). Analysis of data (Figure 12C) showed an eight-fold increase in CCR5⁺ fraction in tumors (mean $\pm$ SEM, n=6; Student's *t* test);

[0064] Figures 13A1 to 13A5 and illustrates cell surface expression of CCR1 protein in cell lines MDA-MB-23, HS578T, SUM159, MCF-7, Jurkat cells by FACS analysis;

[0065] Figures 13B1 to 13B5 and illustrate[[s]] cell surface expression of CCR3 in cell lines MDA-MB-23, HS578T, SUM159, MCF-7, Jurkat cells by FACS analysis;

[0066] Figures 13C1 to 13C5 illustrate[[s]] cell surface expression of CCR5 protein in cell lines MDA-MB-23, HS578T, SUM159, MCF-7, Jurkat cells by FACS analysis;

[0067] Figure 13D shows relative abundance of cell surface expression of CCR1, CCR3 and CCR5 of cell lines MDA-MB-23, HS578T, SUM159, MCF-7, Jurkat cells as determined by FACS analysis;

[0068] Figure 14A illustrates immunohistochemical staining of CCR5 in breast cancer tissue, showing staining was localized primarily to the breast cancer epithelial cell compared with normal breast tissue;

[0069] Figure 14B illustrates immunohistochemical staining of CCR5 in normal breast tissue, showing staining of CCR5 in normal breast tissue is very low, demonstrating lack of CCR5 in normal breast compared with breast tumor;

[0070] Figure 14C Immunohistochemical staining of CCR5 in breast cancer tissue, demonstrating CCR5 immunohistochemical staining being localized primarily to the breast cancer epithelial cell compared with normal breast tissue; and

[0071] Figure 14D shows immunohistochemical staining of CCR5 in normal breast tissue, demonstrating lack of CCR5 in normal breast tissue (of different patients than shown in Figure 14 B) compared with breast tumor.

DETAILED DESCRIPTION OF THE INVENTION

[0072] Chemokine (C-C motif) ligand 5 (“CCL5”), also known as RANTES (an acronym for Regulated on Activation, Normal T cell Expressed and Secreted) is a protein which in humans is encoded by the CCL5 gene. Its receptor, C-C chemokine receptor type 5 (“CCR5” or “CD195”) is a protein found on the white blood cells. CCR5 is the main coreceptor used by macrophage (M)-tropic strains of human immunodeficiency virus type 1 (HIV-1) and HIV-2, which are responsible for viral transmission. CCR5 therefore plays an essential role in HIV pathogenesis. A number of inflammatory CC-chemokines, including MIP-1 alpha, MIP-1 beta, RANTES, MCP-2, and HCC-1[9-74] act as CCR5 agonists, while MCP-3 is a natural antagonist of the receptor. CCR5 is mainly expressed in memory T-cells, macrophages, and immature dendritic cells, and is upregulated by proinflammatory cytokines.

[0073] Classes of antiretroviral medications with activity against HIV include nucleoside analogs, nonnucleoside reverse transcriptase inhibitors, protease inhibitors, fusion inhibitors, integrase inhibitors, and CCR5 receptor antagonists. CCR5 antagonists exert their antiviral activity against HIV by blocking entry of CCR5-tropic viruses into the CD4 T cell. As a result, CCR5 antagonists have historically only been associated with expression in inflammatory cells in the immune system.

[0074] Prior to the present disclosure, the roles of the chemokine CCL5 and its receptor CCR5 in cancer progression were unclear. As disclosed herein, patients with cancers expressing CCL5 and its receptor CCR5, present new drug targets. As disclosed herein, treating cancers that expressed CCL5 and its receptor CCR5 with drugs that blocking their activities respecting CCL5 and its receptor CCR5 selectively affect these cancer cells that express mutant chemokine CCL5 and its receptor CCR5. In one embodiment, a subpopulation of human breast cancer cell lines were found to express CCR5 displayed a functional response to CCL5.

[0075] A microarray analysis conducted on 2,254 human breast cancer specimens found increased expression of CCL5 and its receptor CCR5, but not CCR3, in the basal and HER-2 genetic subtypes. The subpopulation of human breast cancer cell lines found to express CCR5 displayed a functional response to CCL5. Also, oncogene transformation induced CCR5 expression, and the subpopulation of cancer cells that expressed functional CCR5 also displayed increased invasiveness.

[0076] In one embodiment, the CCR5 antagonists developed initially to block CCR5 HIV co-receptor function, reduced *in vitro* invasion of cancer cells without affecting cell proliferation or viability. In one embodiment, the CCR5 antagonists include maraviroc and vicriviroc. In one embodiment, the subpopulation of cancer cells that expressed functional CCR5 include basal breast cancer cells. In one embodiment, the CCR5 antagonists include maraviroc and vicriviroc reduced *in*

vitro invasion of basal cancer cells without affecting the basal cancer cell proliferation or viability. In one embodiment, the CCR5 antagonists include maraviroc and vicriviroc reduced *in vivo* invasion of basal cancer cells without affecting the basal cancer cell proliferation or viability.

[0077] However, as disclosed herein, CCL5 and its receptor CCR5 were found to be expressed in cancer cells, including breast cancer cells, were also found to regulate cancer metastasis, spread of cancer from its primary site to other sites in the body (e.g., brain, liver, lung). Moreover, it was found that blocking the CCR5 receptor with CCR5 antagonists, such as Maraviroc and Vicriviroc, prevented migration and spread of the cancer from its primary site to other sites in the body.

[0078] As described herein, CCR5 and CCL5 were found to play a key role in cancer invasiveness. For example, it was shown that CCR5 antagonists slowed down and/or prevented the invasion of secondary sites in the body by cancers that express CCL5 and/or its receptor CCR5, demonstrating usefulness of CCR5 antagonists as viable adjuvant therapy for reducing the risk of metastasis in cancer patients, including patients having basal breast cancer molecular subtype that express CCL5 and/or its receptor CCR5.

[0079] Cell receptor status can determine whether or not a cancer patient is susceptible to a particular anti-cancer treatment. For example, patients diagnosed with the basal breast cancer subtype, where the basal breast cancer subtype does not express the androgen or estrogen receptors or HER-2, current treatment choices, including chemotherapy, radiation, or surgery, are not only the only choices for these patients, but all show poor outcomes for these patients. Notwithstanding the fact that currently there are no other effective therapies available to them, the patients with this variant of basal breast cancer are further disadvantaged by the fact that this variant of basal breast cancer is also typically associated with metastasis. New cancer treatments present the only hope of cancer free survival for these patients. Accordingly, an urgent need exists for a specific targeted therapy for the basal breast cancer subtype.

[0080] In one aspect the present invention relates to the use of CCR5 modulators to treat, prevent, or manage a neoplasm or metastasis of the neoplasm. In one embodiment the neoplasm is cancer. Exemplary cancers and related disorders that can be treated, prevented, or managed in accordance with the exemplary embodiments of the methods of the present invention include, but are not limited to, leukemias, such as but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias, such as, myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia leukemias and myelodysplastic syndrome; chronic leukemias, such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease,

non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenstrom's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and
5 connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors such as but not limited to, glioma, astrocytoma, brain stem glioma,
10 ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including but not limited to ductal carcinoma, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer such
15 as but not limited to pheochromocytom and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer such as but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers such as but limited to Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipius; eye cancers such
20 as but not limited to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers such as squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer such as squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers such as but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers such as but not limited to endometrial
25 carcinoma and uterine sarcoma; ovarian cancers such as but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers such as but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers such as but not limited to, adenocarcinoma,
30 fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers such as but not limited to hepatocellular carcinoma and hepatoblastoma; gallbladder cancers such as adenocarcinoma; cholangiocarcinomas such as but not limited to pappillary, nodular, and diffuse;

lung cancers such as non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers such as but not limited to germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers such as but not limited to, prostatic intraepithelial neoplasia, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penal cancers; oral cancers such as but not limited to squamous cell carcinoma; basal cancers; salivary gland cancers such as but not limited to adenocarcinoma, mucoepidermoid carcinoma, and adenoicystic carcinoma; pharynx cancers such as but not limited to squamous cell cancer, and verrucous; skin cancers such as but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers such as but not limited to renal cell carcinoma, adenocarcinoma, hypemephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/or uterer); Wilms' tumor; bladder cancers such as but not limited to transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas.

[0081] In one embodiment, antagonists of CCR5 are used to treat CCR5-expressing neoplasm or metastasis of the CCR5-expressing neoplasm. In one embodiment, antagonists of CCR5 are used to prevent the neoplasm or metastasis of said neoplasm. In one embodiment, antagonists of CCR5 are used to manage the neoplasm metastasis of said neoplasm. In one embodiment, antagonists of CCR5 are used slow the progression of the CCR5-expressing neoplasm or metastasis of the CCR5-expressing neoplasm. In one embodiment, antagonists of CCR5 are used to delay metastasis of the CCR5-expressing neoplasm.

[0082] In one embodiment, antagonists of CCR5 that are suitable for use in accordance with the exemplary methods of the present invention include, but are not limited to, the chemical compounds that are described in U.S. Patent No. 6,667,314 by Perros *et al.* The chemical compounds of Perros *et al.* and all formulations and dosage forms including them are incorporated by reference into the present application. Preferred examples of the compounds by Perros *et al.* include:

4,4-difluoro-N-[(1S)-3-[(1R,5S)-3-(3-methyl-5-propan-2-yl-1,2,4-triazol-4-yl)-8-azabicyclo[3.2.1]octan-8-yl]-1-phenylpropyl]cyclohexane-1-carboxamide ("Maraviroc");
N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropylcyclobutanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropylcyclopentanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropyl-4,4,4-trifluorobutanamide;

5 N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropyl-4,4-difluorocyclohexanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-(3-fluorophenyl)propyl-4,4-difluorocyclohexanecarboxamide; and pharmaceutically acceptable salts or solvates thereof.

10 [0083] In one embodiment, the modulators of CCR5 receptor that are suitable for treating the neoplasm or metastasis of said neoplasm; or preventing the neoplasm or metastasis of said neoplasm; or managing the neoplasm metastasis of said neoplasm; or slowing the progression of the neoplasm or metastasis of said neoplasm; or delaying the neoplasm or metastasis of said neoplasm; include, but are not limited to the chemical compounds that are described in U.S. Patent No.
15 6,586,430 by Armour *et al.* The chemical compounds of Armour *et al.* and all formulations or dosage forms including them are incorporated by reference into the present application. Preferred examples of the compounds by Armour *et al.* include:

N-{3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

20 N-[(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

25 N-[(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-2H-pyran-4-carboxamide;

1-Acetyl-N-[(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]3-azetidine carboxamide;

1-Hydroxy-N-[(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

30 2-Methyl-N-[(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopropanecarboxamide;

2-Cyclopropyl-N-[(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo [3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

- N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) tetrahydro-3-furancarboxamide;
 3,3,3-Trifluoro-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) propanamide;
- 5 N-((1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) tetrahydro-2-furancarboxamide;
 1-(Acetylamino)-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) cyclopentanecarboxamide;
 N-((1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) acetamide;
- 10 1-Methoxy-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) cyclopentanecarboxamide;
 1-Amino-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) cyclopentanecarboxamide;
- 15 1-Methyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-oxo-4-pyrrolidinecarboxamide;
 1-Acetyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)3-azetidinecarboxamide;
- 20 N-((1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) acetamide;
 N-((1S)-3-[6-(2-Methyl-1H-benzimidazol-1-yl)-3-azabicyclo[3.1.0]hex-3-yl]-1-phenylpropyl) cyclobutanecarboxamide;
- 25 2-Cyclopropyl-N-((1S)-3-[3-exo-(3-{4-[(methylsulfonyl)amino]benzyl}-1,2,4-oxadiazol-5-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl) acetamide;
 N-((1S)-3-[7-exo-(2-Methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl) cyclobutanecarboxamide;
- 30 2-Cyclopropyl-N-((1S)-3-[7-exo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl) acetamide;
 3,3,3-Trifluoro-N-((1S)-3-[7-exo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl) propanamide;
- N-((1S)-3-[7-endo-(2-Methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl) cyclobutanecarboxamide;
 2-Cyclopropyl-N-((1S)-3-[7-endo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-

- azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}acetamide;
- N-((1S)-3-[7-exo-(2-Methyl-1H-benzimidazol-1-yl)-3-thia-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)cyclobutanecarboxamide;
- 2-Cyclopropyl-N-[(1S)-3-(3-endo-[[2-(4-fluorophenyl)acetyl]amino]-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]acetamide;
- N-[(1S)-3-(3-[[3-endo-(4-Fluorophenyl)propanoyl]amino]-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]cyclobutanecarboxamide;
- N-[(1S)-3-(3-[[3-exo-(4-Fluorophenyl)propanoyl]amino]-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]cyclobutanecarboxamide;
- 2-Cyclopropyl-N-[(1S)-3-(3-exo-[[2-(4-fluorophenyl)acetyl]amino]-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]acetamide;
- N-((1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- N-((1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)tetrahydro-3-furancarboxamide;
- N-((1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)tetrahydro-2H-pyran-4-carboxamide;
- N-((1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)tetrahydro-2-furancarboxamide;
- 1-Acetyl-N-((1S)-3-[3-endo-(1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- N-((1S)-3-[3-endo-(1H-Benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- Methyl 3-(((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)amino)carbonyl]-1-azetidinecarboxylate;
- N-((1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide 1-Acetyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-azetidinecarboxamide;
- 2-[Acetyl(methyl)amino]-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
- 3-[Acetyl(methyl)amino]-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;

- 2-Methoxy-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
- 3-Methoxy-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;
- 5 1-Acetyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-pyrrolidinecarboxamide;
- 1-Methyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-oxo-4-pyrrolidinecarboxamide;
- 1-Acetyl-N-((1S)-3-[3-exo-(2-ethyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- 10 N-((1S)-3-[3-exo-(2-Ethyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- 1-Acetyl-N-((1S)-1-phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)-3-azetidinecarboxamide;
- 15 N-((1S)-1-Phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)-1-propionyl-3-azetidinecarboxamide;
- N-((1S)-1-Phenyl-3-{3-exo[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)acetamide;
- 2-[Acetyl(methyl)amino]-N-((1S)-1-phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)acetamide;
- 20 1-Acetyl-N-((1S)-3-[3-exo-(1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- N(1S)-3-[3-exo-(1H-Benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- 25 1-acetyl-N(1S)-3-[3-exo-(5-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- N-((1S)-3-[3-exo-(5-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- 1-Acetyl-N-((1S)-3-[3-exo-(5-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- 30 N-((1S)-3-[3-exo-(5-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-

- phenylpropyl}-3-azetidinecarboxamide;
1-methyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
(2S)-1-acetyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-azetidinecarboxamide;
5 (2R)-1-acetyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-azetidinecarboxamide;
2-[acetyl(methyl)amino]-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
10 3-[acetyl(methyl)amino]-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;
1-acetyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-pyrrolidinecarboxamide;
N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-(trifluoromethyl)cyclopropanecarboxamide;
15 2-methoxy-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
3-methoxy-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;
20 1-Acetyl-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
1-Methyl-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
25 N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
2-Methoxy-N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
30 N-((1S)-3-[3-exo-(4-Fluoro-2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
3-Methoxy-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;

- 2-[Acetyl(methyl)amino]-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
- 3-[Acetyl(methyl)amino]-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;
- 5 N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-methyl-3-oxetanecarboxamide;
- 3-Ethyl-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-oxetanecarboxamide;
- N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-oxetanecarboxamide;
- 10 3-Ethyl-N-((1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-oxetanecarboxamide;
- N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-methyl-3-oxetanecarboxamide;
- 15 N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-oxetanecarboxamide;
- N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-methyl-3-azetidinecarboxamide;
- 20 1-Acetyl-N-((1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;
- N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;
- 25 N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-methoxyacetamide;
- N-((1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
- N-((1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-methoxypropanamide;
- 30 2-[Acetyl(methyl)amino]-N-((1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;
- 3-[Acetyl(methyl)amino]-N-((1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-

azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide; and pharmaceutically acceptable salts thereof.

[0084] In yet another embodiment, modulators of CCR5 receptor that are suitable for treating neoplasm or metastasis of the neoplasm; or preventing the neoplasm or metastasis of the neoplasm; or managing the neoplasm metastasis of the neoplasm; or slowing the progression of the neoplasm or metastasis of the neoplasm; or delaying the neoplasm or metastasis of said neoplasm; include, but are not limited to the chemical compounds that are described in U.S. Patent Nos. 6,689,765 and 7,384,944 both by Baroudy *et al.* The chemical compounds of Baroudy *et al.* and all formulations or dosage forms including them are incorporated by reference into the present application. A preferred example of the compounds by Baroudy *et al.* includes: (4,6-dimethylpyrimidin-5-yl)-[4-[(3S)-4-[(1R)-2-methoxy-1-[4-(trifluoromethyl)phenyl]ethyl]-3-methylpiperazin-1-yl]-4-methylpiperidin-1-yl]methanone (Vieriviroc, also previously named SCH 417690 and SCH-D), which has an alternative IUPAC name of 5-({4-[(3S)-4-{2-methoxy-1-[4-(trifluoromethyl)phenyl]ethyl}-3-methylpiperazin-1-yl]-4-methylpiperidin-1-yl}carbonyl)-4,6-dimethylpyrimidine; and pharmaceutically acceptable salts or solvates thereof.

[0085] In one aspect, the present invention provides a method of determining whether a subject has cancer or is at risk for developing cancer and/or is at risk for developing metastasis of. In one embodiment, the method includes obtaining a biological sample from a subject having or suspected of having cancer and assessing the level of expression of CCR5 and/or of at least one of CCR5 ligands in the biological sample. In one embodiment, the expression level of CCR5 and/or of at least one of CCR5 ligands in the biological sample is compared to an expression level of CCR5 and/or of at least one of CCR5 ligands in a control sample. In one embodiment, if the expression level of CCR5 and/or of at least one of CCR5 ligands in the biological sample is higher than the level of expression of CCR5 and/or of at least one of CCR5 ligands in the control sample, then the subject is diagnosed as likely to have cancer. In one embodiment, if the expression level of CCR5 and/or of at least one of CCR5 ligands in the biological sample is higher than the level of expression of CCR5 and/or of at least one of CCR5 ligands in the control sample, then the subject is diagnosed as at increased risk for developing cancer. In one embodiment, if the expression level of CCR5 and/or of at least one of CCR5 ligands in the biological sample is higher than the level of expression of CCR5 and/or of at least one of CCR5 ligands in the control sample, then the subject is diagnosed as at increased risk for developing cancer metastasis.

[0086] The term "subject" as used herein is intended to include animals. In particular embodiments, the subject is a mammal, a human or nonhuman primate, a dog, a cat, a horse, a cow

or a rodent.

[0087] In another aspect, the present invention provides a method for molecular classification of cancer based on a level of expression of CCR5 and/or of at least one of CCR5 ligands in the biological sample of the cancer. In one embodiment, the method for molecular classification of cancer comprises (a) obtaining a biological sample of cancer from subject; (b) determining level of expression of CCR5 and/or level of expression of at least one of CCR5 ligands in the biological sample; and (c) if the level of expression of CCR5 and/or level of expression of at least one of CCR5 ligands determined in step (b) is higher than the level of expression of CCR5 and/or of at least one of CCR5 ligands in a control sample, then the cancer is classified as CCR5-expressing cancer. In one embodiment, a cancer subject whose cancer has been classified as CCR5-expressing cancer is diagnosed as likely at risk for developing metastasis of the cancer. In one embodiment, the cancer is breast cancer. In one embodiment, the cancer is prostate cancer. In one embodiment, the cancer is selected from the group consisting of leukemias, such as but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias, such as, myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia leukemias and myelodysplastic syndrome; chronic leukemias, such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease, non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenstrom's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors such as but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including but not limited to ductal carcinoma, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer such as but not limited to pheochromocytom and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid

cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer such as but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers such as but limited to Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; eye cancers such as but not limited to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers such as squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer such as squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers such as but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers such as but not limited to endometrial carcinoma and uterine sarcoma; ovarian cancers such as but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers such as but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers such as but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers such as but not limited to hepatocellular carcinoma and hepatoblastoma; gallbladder cancers such as adenocarcinoma; cholangiocarcinomas such as but not limited to pappillary, nodular, and diffuse; lung cancers such as non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers such as but not limited to germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers such as but not limited to, prostatic intraepithelial neoplasia, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penial cancers; oral cancers such as but not limited to squamous cell carcinoma; basal cancers; salivary gland cancers such as but not limited to adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; pharynx cancers such as but not limited to squamous cell cancer, and verrucous; skin cancers such as but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers such as but not limited to renal cell carcinoma, adenocarcinoma, hypemephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/or uterer); Wilms' tumor; bladder cancers such as but not limited to transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma,

hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas.

[0088] In another aspect, the present invention provides for a method of treating, preventing, or managing a CCR-5 expressing neoplasm or a metastasis of the CCR5-expressing neoplasm in a subject. In one embodiment, the a method of treating or managing a CCR-5 expressing neoplasm or a metastasis of the CCR5-expressing neoplasm in subject having the CCR5-expressing neoplasm or at risk for developing metastasis of the CCR5-expressing neoplasm, comprises administering to the subject a CCR5 modulator. In one embodiment, the CCR5 modulator comprises a CCR5 antagonist. In one embodiment, the CCR5 antagonist comprises 4,4-difluoro-N-[(1S)-3-[(1R,5S)-3-(3-methyl-5-propan-2-yl-1,2,4-triazol-4-yl)-8-azabicyclo[3.2.1]octan-8-yl]-1-phenylpropyl]cyclohexane-1-carboxamide ("Maraviroc"). In one embodiment, the CCR5 modulator comprises a CCR5 antagonist. In one embodiment, the CCR5 antagonist comprises (4,6-dimethylpyrimidin-5-yl)-[4-[(3S)-4-[(1R)-2-methoxy-1-[4-(trifluoromethyl)phenyl]ethyl]-3-methylpiperazin-1-yl]-4-methylpiperidin-1-yl]methanone ("Vicriviroc"). In one embodiment, the CCR5 antagonist is selected from the group consisting of 4,4-difluoro-N-[(1S)-3-[(1R,5S)-3-(3-methyl-5-propan-2-yl-1,2,4-triazol-4-yl)-8-azabicyclo[3.2.1]octan-8-yl]-1-phenylpropyl]cyclohexane-1-carboxamide ("Maraviroc") and (4,6-dimethylpyrimidin-5-yl)-[4-[(3S)-4-[(1R)-2-methoxy-1-[4-(trifluoromethyl)phenyl]ethyl]-3-methylpiperazin-1-yl]-4-methylpiperidin-1-yl]methanone ("Vicriviroc").

[0089] In one embodiment, suitable CCR5 antagonist is selected from the group consisting of: 4,4-difluoro-N-[(1S)-3-[(1R,5S)-3-(3-methyl-5-propan-2-yl-1,2,4-triazol-4-yl)-8-azabicyclo[3.2.1]octan-8-yl]-1-phenylpropyl]cyclohexane-1-carboxamide ("Maraviroc");

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropylcyclobutanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropylcyclopentanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropyl-4,4,4-trifluorobutanamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropyl-4,4-difluorocyclohexanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-

(3-fluorophenyl)propyl-4,4-difluorocyclohexanecarboxamide; and pharmaceutically acceptable salts or solvates thereof.

[0090] In one embodiment, suitable CCR5 antagonist is selected from the group consisting of:

N-{3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

N-{(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-2H-pyran-4-carboxamide;

1-Acetyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}3-azetidine carboxamide;

1-Hydroxy-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

2-Methyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopropanecarboxamide;

2-Cyclopropyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-3-furancarboxamide;

3,3,3-Trifluoro-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-2-furancarboxamide;

1-(Acetylamino)-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

1-Methoxy-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)cyclopentanecarboxamide;

1-Amino-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)cyclopentanecarboxamide;

5 1-Methyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-oxo-4-pyrrolidinecarboxamide;

1-Acetyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)3-azetidinecarboxamide;

10 N-((1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;

N-((1S)-3-[6-(2-Methyl-1H-benzimidazol-1-yl)-3-azabicyclo[3.1.0]hex-3-yl]-1-phenylpropyl)cyclobutanecarboxamide;

2-Cyclopropyl-N-((1S)-3-[3-exo-(3-{4-[(methylsulfonyl)amino]benzyl}-1,2,4-oxadiazol-5-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;

15 N-((1S)-3-[7-exo-(2-Methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)cyclobutanecarboxamide;

2-Cyclopropyl-N-((1S)-3-[7-exo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)acetamide;

20 3,3,3-Trifluoro-N-((1S)-3-[7-exo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)propanamide;

N-((1S)-3-[7-endo-(2-Methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)cyclobutanecarboxamide;

2-Cyclopropyl-N-((1S)-3-[7-endo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)acetamide;

25 N-((1S)-3-[7-exo-(2-Methyl-1H-benzimidazol-1-yl)-3-thia-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl)cyclobutanecarboxamide;

2-Cyclopropyl-N-[(1S)-3-(3-endo-{[2-(4-fluorophenyl)acetyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]acetamide;

N-[(1S)-3-(3-{[3-endo-(4-Fluorophenyl)propanoyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-

phenylpropyl]cyclobutanecarboxamide;

N-[(1S)-3-(3-{[3-exo-(4-Fluorophenyl)propanoyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]cyclobutanecarboxamide;

2-Cyclopropyl-N-[(1S)-3-(3-exo-{[2-(4-fluorophenyl)acetyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]acetamide;

N-[(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-1-propionyl-3-azetidinecarboxamide;

N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]tetrahydro-3-furancarboxamide;

10 N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]tetrahydro-2H-pyran-4-carboxamide;

N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]tetrahydro-2-furancarboxamide;

15 1-Acetyl-N-[(1S)-3-[3-endo-(1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-3-azetidinecarboxamide;

N-[(1S)-3-[3-endo-(1H-Benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-1-propionyl-3-azetidinecarboxamide;

Methyl 3-[(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]amino)carbonyl]-1-azetidinecarboxylate;

20 N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-1-propionyl-3-azetidinecarboxamide 1-Acetyl-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-2-azetidinecarboxamide;

2-[Acetyl(methyl)amino]-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]acetamide;

25 3-[Acetyl(methyl)amino]-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]propanamide;

2-Methoxy-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]acetamide;

3-Methoxy-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-

yl]-1-phenylpropyl}propanamide;

1-Acetyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-pyrrolidinecarboxamide;

1-Methyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-oxo-4-pyrrolidinecarboxamide;

1-Acetyl-N-((1S)-3-[3-exo-(2-ethyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinedicarboxamide;

N-((1S)-3-[3-exo-(2-Ethyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinedicarboxamide;

1-Acetyl-N-((1S)-1-phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)-3-azetidinedicarboxamide;

N-((1S)-1-Phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)-1-propionyl-3-azetidinedicarboxamide;

N-((1S)-1-Phenyl-3-{3-exo[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)acetamide;

2-[Acetyl(methyl)amino]-N-((1S)-1-phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)acetamide;

1-Acetyl-N-((1S)-3-[3-exo-(1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinedicarboxamide;

N(1S)-3-[3-exo-(1H-Benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinedicarboxamide;

1-acetyl-N(1S)-3-[3-exo-(5-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinedicarboxamide;

N-((1S)-3-[3-exo-(5-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinedicarboxamide;

1-Acetyl-N-((1S)-3-[3-exo-(5-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinedicarboxamide;

N-((1S)-3-[3-exo-(5-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinedicarboxamide;

N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

1-methyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

5 (2S)-1-acetyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-2-azetidinecarboxamide;

(2R)-1-acetyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-2-azetidinecarboxamide;

10 2-[acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

3-[acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

1-acetyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-pyrrolidinecarboxamide;

15 N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-1-(trifluoromethyl)cyclopropanecarboxamide;

2-methoxy-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

20 3-methoxy-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

1-Acetyl-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

25 1-Methyl-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-1-propionyl-3-azetidinecarboxamide;

2-Methoxy-N-{(1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-

azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

N-{(1S)-3-[3-exo-(4-Fluoro-2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

3-Methoxy-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

2-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

3-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

10 N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-methyl-3-oxetanecarboxamide;

3-Ethyl-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

15 N-{(1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

3-Ethyl-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-methyl-3-oxetanecarboxamide;

20 N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

25 N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-1-methyl-3-azetidinecarboxamide;

1-Acetyl-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-1-propionyl-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-2-methoxyacetamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

5 N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-methoxypropanamide;

2-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

10 3-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide; and pharmaceutically acceptable salts thereof.

[0091] As disclosed herein, one aspect of the present invention provides for a method of treating, preventing, or managing a CCR-5 expressing neoplasm or a metastasis of the CCR5-expressing neoplasm in a subject by administering to the subject a CCR5 receptor antagonist. In one embodiment, CCR5 receptor antagonists are administered in the form of pharmaceutical formulations or dosage forms that include the CCR5 receptor antagonists. In one embodiment, the CCR5 receptor antagonists are used in the form of acids, esters, or other suitable chemical derivatives. In one embodiment, the CCR5 receptor antagonists are in the form of pharmaceutically acceptable salts derived from various organic and inorganic acids and bases in accordance with procedures well known in the art. In one embodiment, the expression "pharmaceutically acceptable salt" as used herein is intended to mean an active ingredient comprising a CCR5 receptor antagonist utilized in the form of a salt thereof, especially where the salt form confers on the CCR5 receptor antagonist improved pharmacokinetic properties as compared to the free form of the CCR5 receptor antagonist or other previously disclosed salt form. In one embodiment, a pharmaceutically acceptable salt form of the CCR5 receptor antagonist may also initially confer a desirable pharmacokinetic property on the CCR5 receptor antagonist which it did not previously possess, and may even positively affect the pharmacodynamics of the CCR5 receptor antagonist with respect to its therapeutic activity in the body. In one embodiment, the pharmacokinetic properties of the CCR5 receptor antagonist which may be favorably affected include, e.g., the manner in which the CCR5 receptor antagonist is transported across cell membranes, which in turn may directly and positively affect the absorption, distribution, biotransformation or excretion of the CCR5 receptor antagonist.

While the route of administration of the pharmaceutical composition is important and various anatomical, physiological and pathological factors can critically affect bioavailability, the solubility of the CCR5 receptor antagonist is usually dependent upon the character of the particular salt form thereof which it utilized. Further, an aqueous solution may provide the most rapid absorption of an active ingredient into the body of a patient being treated, while lipid solutions and suspensions, as well as solid dosage forms, may result in less rapid absorption. Oral ingestion of the CCR5 receptor antagonist is the most preferred route of administration for reasons of safety, convenience, and economy, but absorption of such an oral dosage form can be adversely affected by physical characteristics such as polarity, emesis caused by irritation of the gastrointestinal mucosa, destruction by digestive enzymes and low pH, irregular absorption or propulsion in the presence of food or other drugs, and metabolism by enzymes of the mucosa, the intestinal flora, or the liver. Formulation of the CCR5 receptor antagonist into different pharmaceutically acceptable salt forms may be effective in overcoming or alleviating one or more of the above-recited problems encountered with absorption of oral dosage forms. Well-known pharmaceutically acceptable salts include, but are not limited to acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, besylate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecysulfate, ethanesulfonate, fumarate, glucoheptanoate, gluconate, glycerophosphate, hemisuccinate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, isethionate, lactate, lactobionate, maleate, mandelate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, oleate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphonate, picrate, pivalate, propionate, salicylate, sodium phosphate, stearate, succinate, sulfate, sulfosalicylate, tartrate, thiocyanate, thiomalate, tosylate, and undecanoate.

[0092] Base salts of the compounds of suitable CCR5 receptor antagonists include, but are not limited to ammonium salts; alkali metal salts such as sodium and potassium; alkaline earth metal salts such as calcium and magnesium; salts with organic bases such as dicyclohexylamine, meglumine, N-methyl-D-glucamine, trishydroxymethylmethylethylamine (tromethamine), and salts with amino acids such as arginine, lysine, etc. Compounds of the present invention which comprise basic nitrogen-containing groups may be quaternized with such agents as (C₁-C₄) alkyl halides, e.g., methyl, ethyl, iso-propyl and tert-butyl chlorides, bromides and iodides; di(C₁-C₄) alkyl sulfate, e.g., dimethyl, diethyl and diamyl sulfates; (C₁₀-C₁₈) alkyl halides, e.g., decyl, dodecyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; and aryl-(C₁-C₄) alkyl halides, e.g., benzyl chloride and

phenethyl bromide. Such salts permit the preparation of both water-soluble and oil-soluble compounds of the present invention.

[0093] Among the above-recited pharmaceutical salts those which are preferred include, but are not limited to acetate, besylate, citrate, fumarate, gluconate, hemisuccinate, hippurate, hydrochloride, hydrobromide, isethionate, mandelate, meglumine, nitrate, oleate, phosphonate, pivalate, sodium phosphate, stearate, sulfate, sulfosalicylate, tartrate, thiomalate, tosylate, and tromethamine.

[0094] Multiple salts forms are included within the scope of the present invention where a CCR5 receptor antagonist contains more than one group capable of forming such pharmaceutically acceptable salts. Examples of typical multiple salt forms include, but are not limited to bitartrate, diacetate, difumarate, dimeglumine, diphosphate, disodium, and trihydrochloride.

Suitable CCR5 receptor antagonists can be administered alone but will generally be administered in admixture with one or more suitable pharmaceutical excipients, diluents or carriers selected with regard to the intended route of administration and standard pharmaceutical practice.

[0095] For example, a CCR5 receptor antagonist can be administered orally or sublingually in the form of tablets, capsules, ovules, elixirs, solutions or suspensions, which may contain flavouring or colouring agents, for immediate or controlled release applications. Such tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine, disintegrants such as starch (preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc may be included. Solid compositions of a similar type may also be employed as fillers in gelatin capsules. Preferred excipients in this regard include lactose or milk sugar as well as high molecular weight polyethylene glycols. For aqueous suspensions and/or elixirs, the CCR5 receptor antagonist may be combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

[0096] Suitable CCR5 receptor antagonists can also be injected parenterally, for example, intravenously, intraperitoneally, intrathecally, intraventricularly, intrastemally, intracranially, intramuscularly or subcutaneously, or they may be administered by infusion techniques. They are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions should be

suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

[0097] For oral and parenteral administration to human subjects, the daily dosage level of the CCR5 receptor antagonist will usually be from 1 microgram/kg to 25 mg/kg (in single or divided doses). Thus tablets or capsules of the CCR5 receptor antagonist may contain from 0.05 mg to 1.0 g of active compound for administration singly or two or more at a time, as appropriate. The physician in any event will determine the actual dosage which will be most suitable for any individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention.

Suitable CCR5 receptor antagonists can also be administered intranasally or by inhalation and are conveniently delivered in the form of a dry powder inhaler or an aerosol spray presentation from a pressurised container or a nebuliser with the use of a suitable propellant, eg

dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134a), carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound, eg using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a powder mix of a CCR5 receptor antagonist and a suitable powder base such as lactose or starch.

[0098] Aerosol or dry powder formulations are preferably arranged so that each metered dose or "puff" contains from 20 µg to 20 mg of a CCR5 receptor antagonist for delivery to the subject. The overall daily dose with an aerosol will be in the range of from 20 µg to 20 mg which may be administered in a single dose or, more usually, in divided doses throughout the day.

Alternatively, Suitable CCR5 receptor antagonists can be administered in the form of a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. The CCR5 receptor antagonist may also be transdermally administered by the use of a skin patch. They may also be administered by the ocular route, particularly for treating neurological disorders of the eye.

[0099] For ophthalmic use, the compounds can be formulated as micronised suspensions in

isotonic, pH adjusted, sterile saline, or, preferably, as solutions in isotonic, pH adjusted, sterile saline, optionally in combination with a preservative such as benzylalkonium chloride. Alternatively, they may be formulated in an ointment such as petrolatum.

For application topically to the skin, the compounds of the formula (I) can be formulated as a

5 suitable ointment containing the active compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, they can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol,
10 liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benyl alcohol and water.

[00100] In some embodiments, the compounds described herein can modulate CCR5 chemokine receptor activity and consequent or associated pathogenic processes subsequently mediated by the CCR5 receptor and its ligands. The expression "modulate CCR5 chemokine receptor activity" as
15 used herein is intended to refer to manipulation of the basic physiological processes and agencies which involve CCR5 chemokine receptors and their ligands. Included within the scope of this intended meaning are all types and subtypes of CCR5 receptors, in whatever tissues of a particular patient they are found, and in or on whatever components of the cells comprising those tissues they may be located. Most commonly, CCR5 receptors are situated on the cell membranes of particular
20 cell types such as monocytes. CCR5 receptors participate in and define, along with various endogenous ligands to which they are naturally bound, signaling pathways which control important cellular and tissue functions by means of the influence which they exert on the movement of agents such as the chemokines, into and out of those cells and tissues.

[00101] The dosage and dose rate of the compounds of Formula (I) effective for treating or
25 preventing diseases and conditions in a patient which are mediated by or associated with modulation of CCR5 chemokine receptor activity as described herein, as well as for favorably affecting the outcome thereof in the patient, in accordance with the methods of treatment of the present invention comprising administering to the patient a therapeutically effective amount of a CCR5 receptor antagonist, will depend on a variety of factors such as the nature of the CCR5 receptor antagonist,
30 the size of the patient, the goal of the treatment, the nature of the pathology being treated, the specific pharmaceutical composition used, the concurrent treatments that the patient may be subject to, and the observations and conclusions of the treating physician.

[00102] Generally, however, the effective therapeutic dose of a suitable CCR5 receptor antagonist which will be administered to a subject will be between about 10 µg (0.01 mg)/kg and about 60.0 mg/kg of body weight per day, preferably between about 100 µg (0.1 mg)/kg and about 10 mg/kg of body weight per day, more preferably between about 1.0 mg/kg and about 6.0 mg/kg of body weight per day, and most preferably between about 2.0 mg/kg and about 4.0 mg/kg of body weight per day of the CCR5 receptor antagonist.

[00103] Included within the scope of the present invention are embodiments comprising coadministration of, and compositions which contain, in addition to a CCR5 receptor antagonist as active ingredient, additional therapeutic agents and active ingredients. Such multiple drug regimens, often referred to as combination therapy, may be used in the treatment and prevention of any of the diseases or conditions mediated by or associated with CCR5 chemokine receptor modulation, particularly cancer metastasis. The use of such combinations of therapeutic agents is especially pertinent with respect to the treatment, prevention, or management of cancer metastasis within a subject in need of treatment cancer, prevention cancer, or management of risk of cancer metastasis.

[00104] Exemplary CCR5 receptor antagonists may be administered in accordance with a regimen of 1 to 4 times per day, preferably once or twice per day. The specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing.

[00105] As disclosed in U.S. Patent No. 6,667,314 by Perros et al. CCR5 antagonists can be administered orally, buccally or sublingually in the form of tablets, capsules, multi-particulates, gels, films, ovules, elixirs, solutions or suspensions, which may contain flavouring or colouring agents, for immediate-, delayed-, modified-, sustained-, pulsed- or controlled-release applications. The CCR5 receptor antagonists may also be administered as fast-dispersing or fast-dissolving dosage forms or in the form of a high energy dispersion or as coated particles. Suitable formulations of the compounds of the CCR5 receptor antagonist may be in coated or uncoated form, as desired. Such solid pharmaceutical compositions, for example, tablets, may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate, glycine and starch (preferably corn, potato or tapioca starch), disintegrants such as sodium starch glycollate, croscarmellose sodium and certain complex silicates, and granulation binders such as

polyvinylpyrrolidone, hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

[00106] "Metastasis", as used herein, is defined as the transfer of malignant tumor cells, or neoplasm, via the circulatory or lymphatic systems or via natural body cavities, usually from the primary site of neoplasia to a distant site in the body, and subsequent development of secondary tumors or colonies in the new location. In some exemplary embodiments of the methods of the present invention, metastasis comprises a tumor metastasis in or more organs selected from the group consisting of liver, brain, bladder, lung, adrenal gland, kidney, bone, skin or pancreas or control kidney and combinations thereof.

[00107] In another aspect, the present invention provides a method of identifying a compound that reduces or prevents or treats cancer metastasis. In one embodiment, the method identifies a candidate compound that selectively interferes with proliferation or viability of neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand. In one embodiment, the method identifies a candidate compound that selectively blocks activity of CCR5 and/or at least one of CCR5 receptor ligand in neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand. In one exemplary embodiment, the method for identifying a candidate compound that reduces or prevents or treats cancer metastasis in neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand, comprises (a) contacting a one or more neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand with one or more candidate compounds; and (b) detecting activity of and/or proliferation or viability of the one or more neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand, wherein decreased activity and/or decreased proliferation and/or decreased viability of the one or more neoplastic cells (relative to as compared to a control sample) identifies the candidate compound as a compound that that selectively reduces or prevents or treats cancer metastasis in neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand. In one embodiment, if proliferation of the one or more neoplastic cells is decreased/depressed compared to untreated (control) neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand, the candidate compound is identified as a compound that selectively reduces or prevents or treats metastasis of a neoplasm cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand. In one embodiment, if viability of the one or more neoplastic cells is decreased/depressed compared to untreated (control) neoplastic cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand,

the candidate compound is identified as a compound that that selectively reduces or prevents or treats metastasis of a neoplasm cells that over express CCR5 and/or over express at least one of CCR5 receptor ligand. In one embodiment, the one or more CCR5 receptor ligands comprise CCL5. In one embodiment, the one or more CCR5 receptor ligands comprise CCL8. In one
5 embodiment, the one or more CCR5 receptor ligands comprise CCL7.

[00108] For the study disclosed herein, CCL5 and CCR5 expression in human breast cancer cell lines were investigated as well as the effect of CCR5 antagonists in vitro and in vivo. An interrogation was conducted using a microarray dataset to evaluate CCR5 and CCL5 expression in the context of 2,254 patient breast cancer samples. Samples in the dataset were assigned to five
10 breast cancer subtypes, including luminal A, luminal B, normal-like, basal and HER-2 overexpressing disease. The analysis revealed an increased expression of CCL5 and CCR5 in patients with basal and HER-2 subtypes. 58% the cancer samples indicated a positive CCR5 and CCL5 signature. It was found that oncogenes turn on the CCR5 receptor in normal breast cells as they became transformed into cancer cells. Metastasis those cells was also found to be regulated by
15 CCR5.

[00109] To evaluate the functional relevance of CCR5 in cellular migration and invasion in vitro, the drugs were tested in 3-D invasion assays with two different cell lines. It was found that both antagonists inhibited breast cancer cell invasiveness.

[00110] To evaluate the functional relevance of CCR5 in cellular migration and invasion in vivo, mice were injected with the antagonists and invasiveness of the basal breast cancer cells to other
20 tissue, i.e. lung, was tracked with bioluminescence imaging. It was found that mice treated with the drug showed a more than 90% reduction in both the number and size of pulmonary metastases compared to untreated mice. This and the other preclinical studies provide the rational basis for studying the use of CCR5 antagonists as new treatments to block the dissemination of basal breast
25 cancers. These findings may also have implications for other cancers where CCR5 promotes metastasis, such as prostate and gastric.

[00111] Referring to the drawings in detail, wherein like reference numerals indicate like elements throughout, there are exemplary embodiments of the present invention shown in Figures 1A-14D.

[00112] Referring to Figure 1A, there is shown a heatmap of the expression of CCL5 and its receptor CCR5 in samples from patients with breast cancer divided by molecular subtypes of breast cancer, namely luminal A, luminal B, basal, normal-like, and Her-2, based on their gene expression pattern. The heatmap shows that relative abundances of CCL5 and CCR5 are increased (over
30

expressed) in patients with the basal and HER-2 breast cancer subtypes. The heatmap also shows that CCL5 and CCR5 are over expressed in the Her-2 breast cancer subtype.

[00113] Referring to Figures 1B1 to 1B5, there is shown fluorescence-activated cell sorting (FACS) scatter plots and correlation analysis (Student *t* test) of the expression of CCL5 and CCR5 among the breast cancer molecular subtypes whose expression of CCL5 and its receptor CCR5 is shown Figure 1A. Consistent with the expression of CCL5 and its receptor CCR5 observed in A, the scatter plots show CCL5 and CCR5 are over expressed in patients with the basal and HER-2 breast cancer subtypes.

[00114] Referring to Figure 1C, there is shown quantification of the proportions of the breast cancer samples overexpressing CCL5 and CCR5 (fraction of the bars representing upper right quadrants of the scatter plots shown in Figures 1B1 to 1B5) displayed in Figures 1B1 to 1B5. The number of samples in each subtype is indicated at the top of the bar.

[00115] Referring to Figure 1D, there is shown metastasis-free Kaplan–Meier plots and log-rank analysis for the different genetic subtypes in the analyzed database described in Materials and Methods section of the present disclosure. The metastasis-free Kaplan–Meier plots show that patients with the basal or HER-2 subtypes of breast cancer display increased probability to develop metastasis.

[00116] Figures 2A-2H, show that human breast cancer cell lines that express CCR5 respond to CCL5. In Figure 1A, flow cytometric histograms of the CCR5 expression in MDA-MB-231 breast cancer cells identified a subpopulation of CCR5⁺ cells. In Figures 2B and 2B2, induction of calcium signaling in cells loaded with Fluo-4-AM before the sequential addition of CCL5 (60 µg/mL) and FBS (5%) is shown. A fraction of cells responded to CCL5 (closed arrowheads in the middle of micrographs) whereas the rest did not (open arrowheads). The average changes in fluorescence on 5 responsive (green line) and 5 nonresponsive (red line) cells are represented in the far right graphs. Data shown are representative of 3 to 5 independent experiments for each cell line (Bar, 100 µm). In Figure 2C, 3D invasion into collagen gels by breast cancer cell lines, using CCL5 (15 µg/mL) as chemoattractant is shown. Figure 2D shows mean distances of invasion ± SEM from 3 independent experiments whose 3D invasion are shown in Figure 2C. Figure 2E shows 3D invasion assays, and their corresponding quantification shown in Figure 2F (mean ± SEM, *n* = 3), for MCF-10A cells and MCF-10A-NeuT, -Ras, and -Src derivatives showing that CCL5-induced invasion is activated by oncogenic transformation. Figure 2G shows CCR5⁺ cells display increased invasiveness. CCR5⁺ and CCR5[−] subpopulations from SUM-159 cell line were isolated by FACS and invasion into collagen gels was evaluated using FBS as chemoattractant. Quantification of

FACS and invasion into collagen gels experiments of the samples in Figure 2G is shown in Figure 2H as mean $\pm$ SEM of 2 independent experiments. Statistical analysis was conducted using the Student *t* test.

[00117] Figures 3A1-3D show that CCR5 antagonists block CCL5-induced calcium signaling. In Figures 3A1 to 3A4, there is shown intensity versus time analysis of Fluo-4 AM-loaded MDA-MB-231 cells treated with the CCR5 antagonists maraviroc or vicriviroc (100 nmol/L) for 30 minutes before the addition of CCL5 (60 μ g/mL). Micrographs illustrate the axis (x-x') of the pseudoline scan plot. Those axes were used to construct the adjacent intensity versus time plots. In Figure 3B, there is shown comparison of the fraction of cells with increased fluorescence intensity upon addition of CCL5. In Figures 3C1 to 3C4, CCL5-induced calcium signaling was also blocked by CCR5 antagonists in Hs578T cells. The corresponding quantification is shown in Figure 3D. The data in Figures 3B and 3D are mean $\pm$ SEM of 3 to 4 independent experiments. Statistical analysis was conducted using the Student *t* test.

[00118] Figures 4A-4D show that CCR5 antagonists block FBS-induced breast cancer cell invasion. In Figure 4A, there is shown 3D reconstruction of FBS-induced invasion into collagen gels by Hs578T breast cancer cells in presence of CCR5 antagonists (100 nmol/L). In Figure 3C there is shown 3D reconstruction of FBS-induced invasion into collagen gels by SUM-159 breast cancer cells in presence of CCR5 antagonists (100 nmol/L). The corresponding quantifications (mean $\pm$ SEM, *n* = 3) and analysis (Bonferroni *t* test) are displayed in Figures 4B and 4D.

[00119] Referring to Figures 5A-5F, the data presented show that the CCR5 antagonist maraviroc inhibits lung metastases *in vivo*. In Figure 5A, MDA-MB-231 cells transduced with Luc2-eGFP fusion protein were injected into the tail vein of NOD/SCID mice and the *in vivo* bioluminescent signal was quantified weekly. Representative *in vivo* images of vehicle- or maraviroc-treated (8 mg/kg every 12 hours) mice are shown in Figure 5A. In Figure 5B, there is shown quantification (mean $\pm$ SEM, *n* = 6) of BLI in the control (red line) and treated groups (blue line). Statistical comparison (*, *P* = 0.048) was carried out using Student *t* test with Welch correction for heterogeneous variances. As shown in Figures 5C1 and 5C2, the presence of pulmonary tumors and the differences between treatments were corroborated by *ex vivo* imaging (left) and India ink staining (right). As shown in Figure 5D, the fraction of mice with metastatic tumors was significantly larger in the control group (*P* < 0.0001, Fisher exact test). Figure 5E shows histologic analysis (hematoxylin and eosin staining, $\times$ 100) and the corresponding quantification (shown in Figure 5F) of the area covered by metastatic tumors in lung slides. Tumor area was quantified with the Nikon Elements BR 3.0 software analyzing at low magnification ($\times$ 40) 2 random fields of 2

different histologic sections (separated 600 μ m from each other) per mouse. Statistical analysis was conducted using the Student *t* test with Welch correction for heterogeneous variances (*n* = 33 and 12 for control and treated groups, respectively).

[00120] Referring to Figures 6A-6H, it is shown that maraviroc reduces lung colonization but does not modify cell proliferation. In Figure 6A, there is shown the effect of CCR5 antagonist on breast cancer cell viability. MDA-MB-231 cells were exposed to increasing concentrations of maraviroc (inverted triangles) or vicriviroc (squares) for 48 hours and the cell viability was evaluated by MTT assay. Graph is from a representative experiment carried out by sextuplicate. No statistical differences were found (ANOVA) in 3 independent experiments. In Figure 6B, there is shown CCR5 expression in MDA-MB-231 cells stably transfected with pcDNA3.1+/Zeo+ (MDA.Vector) or human CCR5 cloned into pCDNA3+Zeo+ (MDA.CCR5). In Figure 6C, comparison of *in vitro* proliferation rates of MDA.Vector versus MDA.CCR5 showed no differences (ANOVA). Representative experiment from 2 carried out by sextuplicate. In Figure 6D, to evaluate the *in vivo* effect of maraviroc on growth of established metastasis, treatment of mice was initiated 10 days after injection of MDA.pFULG cells as illustrated. In Figure 6E, quantification (mean $\pm$ SEM, *n* = 5) of *in vivo* BLI in the control (red/squares) and treated groups (blue/triangles) showed no differences in the growth rate. There is no effect of maraviroc of the growth rate of the tumors once established in the lungs. Thus, the reduction in tumor mass in the lungs is due to the inhibition of "homing" or spread of tumors to the lung. Figure 6F shows schema of the experimental design used to evaluate CCR5 role in lung colonization. Figure 6G shows representative confocal images and quantification (Figure 6H) of the number of eGFP+ cells in lungs 24 hours after injection of MDA.pFULG cells. Cells expressing eGFP were counted in 3 random fields of 2 different histologic sections (separated 700 μ m from each other) per mouse (*n* = 5 mice per group). Statistical analysis was conducted using Student *t* test. Bar in micrographs, 100 μ m.

[00121] Reference is now made to Figures 7A1-7D2. To determine the gene expression signaling pathway associated with enrichment of CCR5 and CCL5, GSEA analysis using KEGG and GO was conducted of the tumor samples discussed herein. Figure 7A shows the GSEA analysis. These studies showed enrichment for gene expression of pathways including lymphocyte activation, Janus-activated kinase (JAK)-STAT signaling, and Toll-like receptor activation. The receptors for CCL5 include CCR1 and CCR3. Increased expression of CCL5 associated with increased CCR1, but not CCR3, in the basal and HER-2 genetic tumor type (Figures 7B1-7B5 and 7C1 to 7C5). In ER-negative patients receiving chemotherapy, there was an insignificant trend toward reduced metastasis-free survival and relapse-free survival in the increased CCR5 population, compared with

the population with reduced CCR5 expression (Figure 7D1 and 7D2).

[00122] In Figures 8A-8D, a comparison of expression levels for CCL5 versus CCR5, CCR1, and CCR3, comparing normal breast with breast cancer showed increased correlation between receptor and ligand expression levels in tumors compared with healthy breast tissue. Figure 8A illustrates heatmap of CCL5 expression and its receptors CCR5, CCR1 and CCR3 in healthy population. The healthy population was previously described as part of a collection of 2550 breast cancers (Ertel, A., Dean, J. L., Rui, H., Liu, C., Witkiewicz, A. K., Knudsen, K. E., and Knudsen, E. S. RB-pathway disruption in breast cancer: differential association with disease subtypes, disease-specific prognosis and therapeutic response. *Cell Cycle*, 9: 4153-4163, 2010.).

[00123] Referring to Figures 9A-9D, three human breast cancer cell lines with a basal phenotype and molecular signature: MDA-MB-231, Hs578T, and SUM-159 (34-37) were used as models in the studies. Analysis of CCR5 expression by FACS showed that a small subpopulation of cells were positive for the receptor in all 3 cell lines (Figure 2A for MDA-MB-231 and Figure 9A and Figure 9C for Hs578T and SUM-159). Because CCR5 activation induces calcium flux (38, 39), the activation of calcium signaling was assessed by CCL5. Addition of CCL5 to the cultures induced immediate calcium fluxes in a subpopulation of cells (Figure 2B for MDA-MB-231 and Figures 9B and 9D for Hs578T and SUM-159), providing evidence that CCR5 is functional in basal breast cancer cells. As a positive control, the same cultures were exposed to 5% FBS (40). Calcium flux, assessed by relative fluorescence intensity, increased in more than 95% of the cells after FBS addition (Figure 2B and Figures 9B, 9D1, and 9D2). To further distinguish CCL5-dependent signaling, SUM159 cells were stably transduced with a CCR5 expression vector and the Ca^{+2} response to CCL5 versus FBS was conducted (Figures 10C and 10D vs. Figures 10E and 10F). CCR5 induced Ca^{+2} signaling in the CCR5-overexpressing cells, whereas both lines responded similarly to FBS induced Ca^{+2} activation (Figures 10A-10F).

[00124] In view of the finding that CCR5 inhibition by CCR5 antagonists reduced calcium signaling and cell invasion, the in vivo effect of maraviroc on lung metastasis was determined. MDA-MB-231 cells were transduced within the Luc2-eGFP lentiviral vector (MDA.pFULG cells) in an experimental metastasis model. The Luc2 gene is a codon-optimized version of Luc and cells expressing this reporter were 10 to 100 times brighter than the unmodified Luc gene (30). After injection of MDA.pFULG cells into the tail vein of mice, noninvasive BLI enabled the early detection of breast cancer metastasis (41). Weekly BLI was conducted for 5 weeks and the radiance antemortem was used as a surrogate measurement of tumor burden. Mice treated with maraviroc (8 mg/kg twice daily) showed a significant reduction in both the number and the size of pulmonary

metastases compared with vehicle-treated mice (Figures 5A and 5B, Figures 11A and 11B). To avoid the possibility that metastases were missed because of inappropriate imaging, ex vivo imaging, India ink staining (Figures 5C1 and 5C2), and histology (Figure 5E) of the lungs were conducted. Histologic analysis corroborated that tumor burden corresponds to bioluminescence, as previously shown (30). Metastatic tumors were still detectable in 50% of the maraviroc-treated mice, but their mean size was reduced by 65% (Figures 5D and 5F). Interestingly, analysis of CCR5 expression in lungs from control mice showed an 8-fold enrichment of the CCR5+ fraction (Figures 12A to 12F). Collectively, these results provide evidence that CCR5 antagonists reduce breast cancer metastasis in vivo.

[00125] Given the aggressive clinical behavior of basal breast cancer and the lack of targeted therapies for it, the importance of the CCL5/CCR5 was evaluated in invasion and metastasis in the human breast cancer cell lines MDA-MB-231, Hs578T and SUM-159. These cell lines reflect the clinicopathologic features of the basal subtype of breast cancer (including the lack of HER-2, ER, and progesterone receptor), a basal-like molecular signature, the activation of specific signaling pathways (e.g., hypoxic or EGF receptor responses) and over expression of epithelial-mesenchymal transition proteins (FN, VIM, ad matrix metalloproteinase 2; refs. 34-37). Only a small fraction of cells within the cell lines used in this study expressed CCR5 as evaluated by FACS analysis. The studies confirmed the expression of CCR5 in MDA-MB-23 cells by reverse transcriptase PCR and showed the presence of the CCR5 protein by FACS analysis (Figures 13A1-13D1), and showed that CCR5 immunohistochemical staining was localized primarily to the breast cancer epithelial cell, compared with normal breast tissue (Figures 14A-14D).

[00126] The results disclosed herein show that CCL5 activates calcium flux in basal-like human breast cancer cells. By using the selective CCR5-antagonists maraviroc and vicriviroc (both with IC50 below 30 nmol/L; refs. 44, 45), it was shown that CCL5-activated signaling is mediated by CCR5. However, the fraction of CCL5-responsive cells (10% and 12% for MDA-MB-231 and Hs578T cells, respectively) is higher than the percentage of CCR5-expressing cells determined by FACS. This may be due to the greater sensitivity of the Ca²⁺ activation assays compared with the sensitivity of analysis by FACS. In addition, CCL5-induced calcium redistribution is not completely blocked by CCR5 antagonists. This may be caused by the expression of other receptors to CCL5, namely CCR1 and CCR3. CCR5 has been identified as the main CCL5 receptor in MDA-MB-231 cells (13) and CCR1 and CCR3 transcripts are absent in both MDA-MB-231 or Hs578T cell lines (8) and breast tumor samples (11). CCR1 and CCR3 were able to be detected by FACS (Figures 13A1-13D), suggesting a possible mechanism for the incomplete response to the CCR5

antagonist.

[00127] In one aspect, the present invention provides an in vivo method for identifying a candidate compound that down regulate expression of CCR5 and/or one or more of its ligand in tumor cells overexpressing endogenous or virally transduced oncogene selected from the group consisting of NeuT, Ha-Ras, c-Src and combinations thereof. In one embodiment, the method comprises (a) administering a candidate compound to an animal model of the tumor cells; (b) and measuring the level of CCR5 RNA or protein expression in the animal model, wherein if the level of CCR5 RNA or protein expression in the animal model is decreased compared to the level of CCR5 RNA or protein expression in untreated animal model, then the candidate compound is identified as a compound that down regulates expression of CCR5 and/or one or more of its ligand in tumor cells overexpressing endogenous or virally transduced oncogene selected from the group consisting of NeuT, Ha-Ras, c-Src and combinations thereof. In one embodiment, the tumor cells comprise mammalian prostate cancer cells. In one embodiment, tumor cells comprise mammalian prostate cancer cell line having at least one or more of a set of primary mammalian epithelial cells which have been infected with a retroviral vector carrying an oncogene selected from the group consisting of NeuT, Ha-Ras, c-Src and combinations thereof.

[00128] An exemplary mammalian prostate cancer cell lines suitable for use in the present invention include, but are not limited to, mammalian prostate cancer cell lines disclosed in US Provisional Patent Application No. 61/646,586, filed May 14, 2012, whose contents are incorporated by reference herein in their entirety. In one embodiment, the mammalian prostate cancer cell line comprises at least one or more of a set of primary mammalian epithelial cells which have been infected with a retroviral vector carrying an oncogene. In one embodiment, the oncogene is selected from the group consisting of c-Myc, Ha-Ras, NeuT, c-Src and combinations thereof. In one embodiment of the mammalian prostate cancer cell line, an oncogene selected from the group consisting of c-Myc, Ha-Ras, NeuT, c-Src and combinations thereof is expressed. The mammalian prostate cancer cell line can include any suitable mammalian cell, including primary murine epithelial cells. In some embodiments, the primary mammalian epithelial cells are derived from any immune competent mammal. In one embodiment, the primary mammalian epithelial cells are derived from an immune competent mammal selected from the group consisting of rodents, including rats and mice.

[00129] In some embodiments, the suitable animal model of cancer comprises an immune

competent mammal implanted with a cancer cell line transformed with one or more of a set of oncogenes selected from the group consisting of c-Myc, Ha-Ras, NeuT, c-Src and combinations thereof. In one embodiment, the animal model of cancer is an immunocompetent transgenic mouse created using the mammalian prostate cancer cell line of the present invention develops a prostate tumor capable of producing a detectable molecular genetic signature based on an expression level of one or more of a set of oncogenes selected from the group consisting of c-Myc, Ha-Ras, NeuT, c-Src and combinations thereof.

[00130] In some embodiments, the suitable animal model of cancer is produced by an in vitro method. In one embodiment, in vitro method for producing the suitable animal model includes production of immortalized primary mammalian epithelial cells. In one embodiment, the method for the in vitro production of immortalized primary mammalian epithelial cells, comprises infecting primary mammalian epithelial cells with a retroviral vector carrying an oncogene selected from the group consisting of c-Myc, Ha-Ras, NeuT, c-Src and combinations thereof to provide infected cells, wherein the primary mammalian epithelial cells are capable of being infected by said retroviral vector and under conditions whereby the c-Myc, Ha-Ras, NeuT, c-Src and combinations thereof are expressed in said infected cells.

I. EXAMPLES

A. MATERIALS AND METHODS

A.1. Breast cancer patients data set and statistical analysis

[00131] A microarray data set that was previously compiled (21) from the public repositories Gene Expression Omnibus (23) and ArrayExpress (24) was used to evaluate CCR5 and CCL5 expression in the context of clinical samples. Samples in this data set were assigned to 5 canonical breast cancer subtypes, including luminal A, luminal B, normal-like, basal, and HER-2-overexpressing disease. The classification of microarray samples among these 5 subtypes was achieved by computing their correlation against an expression profile centroid representative of each subtype and assigning samples to the subtype with the highest corresponding correlation coefficient (25). Samples with a maximum correlation coefficient below 0.3 were considered unclassified. Analysis of CCL5 and CCR5 transcript was then conducted specifically among the luminal A, luminal B, basal, normal-like, and HER-2 subtypes. Differential expression of the averaged gene signature magnitude among these sample subsets was evaluated using 2-tailed Student *t* test.

Kaplan–Meier analysis was used to evaluate survival trends within the sample subsets. Scatter plots of CCL5 versus CCR5 samples were also generated to observe coregulation patterns specific to each subtype. For these scatter plots, gene profiles were median-centered and scaled to unitary SD.

A.2. Cell Lines And Cell Culture

5 [00132] MDA-MB-231, MCF-7, and Hs578T cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FBS. SUM-159 cells (kindly provided by Dr. Stephen Ethier, Wayne State University, Detroit, MI) were maintained in Ham's F-12 supplemented with 4 µg/mL of insulin, 1 µg/mL of hydrocortisone, and 5% FBS. Oncogene-transformed derivatives of MCF-10A cells (MCF10A-NeuT, MCF10A-Src, and MCF10A-Ras; ref. 26) were
10 maintained in DMEM:Ham's F-12 (50 of 50) supplemented with 4 mg/mL of insulin, 10 ng/mL of EGF, and 1 mg/mL of hydrocortisone. A total of 100 µg/mL of each penicillin and streptomycin were included in all media. Cells were cultured in 5% CO₂ at 37°C. For *in vitro* treatments, maraviroc was dissolved in dimethyl sulfoxide (DMSO) and diluted in culture medium. The final concentration of DMSO in treated and control cultures was 0.5%. Vicriviroc was dissolved in
15 culture medium.

A.3. Fluorescence-Activated Cell-Sorting Analysis

[00133] Cell labeling and fluorescence-activated cell-sorting (FACS) analysis for CCR5 were based on prior publications (27) with minor modifications. Before labeling, the cells were blocked with normal mouse IgG (1 of 100) and purified rat anti-mouse Fcγ III/II receptor antibody (1 of 100;
20 Pharmingen) for 30 minutes and then incubated with allophycocyanin (APC)-labeled CCR5 antibody (R&D Systems). All experiments were conducted at 4°C. Sample analysis was conducted on FACSCalibur flow cytometer (BD Biosciences). These data were analyzed with FlowJo software (Tree Star, Inc.).

A.4. Invasion Assay

25 [00134] The 3-dimensional invasion assay was conducted as previously reported (12). Briefly, 100 µL of 1.67 mg/mL Rat Tail collagen type I (BD Biosciences) was pipetted into the top chamber of a 24-well 8-µm pore Transwell (Corning). The Transwell was incubated at 37°C overnight to allow the collagen to solidify. A total of 30,000 cells were then seeded on the bottom of the Transwell membrane and allowed to attach. Serum-free growth medium was placed into the bottom

chamber, whereas 15 ng/mL CCL5 or 5% FBS was used as a chemoattractant in the medium of the upper chamber. The cells were then chemoattracted across the filter through the collagen above for 3 days. Cells were fixed in 4% formaldehyde, permeabilized with 0.2% Triton-X in PBS, and then stained with 40 µg/mL propidium iodide (PI) for 2 hours. Fluorescence was analyzed by confocal z-sections (one section every 20 µm) at ×10 magnification from the bottom of the filter using a Zeiss LSM 510 Meta inverted confocal microscope at the Kimmel Cancer Center Bioimaging Facility.

A.5. Intracellular Calcium Assay

[00135] Calcium responses induced either by CCL5 or FBS in human cancer cell lines were monitored under fluorescence confocal microscope as previously reported (28). Briefly, breast cancer cells were seeded in 4-well labtek chambers (Nunc) at 10^4 cells/cm² and incubated for 1 day. After 12-hour starvation, cells were labeled by incubating them with 2 mmol/L Fluo-4-AM (Molecular Probes) in HBSS for 30 minutes, washed twice, and incubated for additional 30 minutes before imaging under the microscope. Time-lapse images were collected using a Zeiss LSM 510 Meta inverted confocal microscope with the incubator at 37°C. Relative intracellular Ca²⁺ concentration was determined by the changes in fluorescent intensity (FI) of Fluo-4-AM upon the addition of CCL5 (60 ng/mL) or FBS (5%) and was calculated as $(FI_t - FI_0)/FI_0$.

A.6. MTT Assay

[00136] The MTT assay is a colorimetric assay for measuring the activity of cellular enzymes that reduce the tetrazolium dye, 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, a yellow tetrazole (MTT), to its insoluble formazan, giving a purple color. These assay measures cellular metabolic activity via NAD(P)H-dependent cellular oxidoreductase enzymes and may, under defined conditions, reflect the number of viable cells (cell proliferation). Tetrazolium dye assay can also be used to measure cytotoxicity (loss of viable cells) or cytostatic activity (shift from proliferative to resting status) of potential medicinal agents and toxic materials. MTT Assay usually done in the dark because MTT reagent is sensitive to light.

[00137] The effects of CCR5 antagonists on cell viability and proliferation rate were estimated using the soluble tetrazolium salt MTT assay (29). MTT is reduced by the mitochondria of viable cells, and the amount of reduced formazan is proportional to the number of viable cells. After 72 hours of exposure to the drugs, cells were incubated with 1 mg/mL of MTT for 90 minutes. Then, the reduced (insoluble and colored) formazan was dissolved in DMSO and measured

spectrophotometrically at 570 nm. The effect of CCR5 over expression in breast cancer cell proliferation was studied in MDA-MB-231 cells transfected with full-length human CCR5 subcloned into pcDNA3.1⁺/Zeo⁺ vector (kindly provided by Dr. Eleanor Fish, University of Toronto, Toronto, ON, Canada) and selected with Zeocin (200 µg/mL) as previously described (18).

5 MTT assays were conducted in sextuplicate using 96-well microplates.

A.7. Viral Cell Transduction

[00138] A lentiviral vector encoding firefly luciferase 2 (Luc2)-eGFP fusion protein was a generous gift from Dr. Sanjiv S. Gambhir (School of Medicine, Stanford University, Stanford, CA; ref. 30). Lentivirus propagation was conducted following the protocol described by Zahler and
10 colleagues (31). Breast cancer cell lines were transduced at a multiplicity of infection of 20 in the presence of 8 mg/mL polybrene (Sigma) for 24 hours (30, 31).

A.8. Experimental Metastasis Assay And Bioluminescence Imaging

[00139] MB-MDA-231 cells expressing Luc2-eGFP (called MDA.pFLUG for the rest of the article) were detached with a nonenzymatic cell dissociation buffer (4 mmol/L EDTA in Ca²⁺ and
15 Mg²⁺-free PBS), resuspended in Dulbecco's PBS without Ca²⁺ and Mg²⁺ and immediately injected into the tail vein of 8-week-old, female nonobese diabetic/severe combined immunodeficient (NOD/SCID) mice (NCI, Bethesda MD). Each mouse received 10⁶ cells. Mice were treated by oral gavage with maraviroc (8 mg/kg every 12 hours) or vehicle (5% DMSO in acidified water; ref. 32). Treatment was started immediately after injection or 10 days later for the experiments analyzing the
20 proliferation of established metastasis. For *in vivo* bioluminescence imaging (BLI), mice were given an intraperitoneal (i.p.) injection with 200 µL of d-luciferin (30 mg/mL). Mice were anesthetized with isoflurane (2% in 1 L/min oxygen), and bioluminescence images were acquired 10 to 15 minutes after d-luciferin injection using the IVIS XR system (Caliper Life Sciences). Acquisition times ranged from 10 seconds (for later time points) to 5 minutes (for early time points). Data are
25 expressed as total photon flux and were analyzed using Living Image 3.0 software (Caliper Life Sciences). For *ex vivo* BLI, d-luciferin was diluted in PBS to a final concentration of 300 µg/mL and used to soak freshly isolated lungs for 2 to 3 minutes before imaging. Some lungs were stained with India ink, as previously reported (33), or processed and stained with hematoxylin and eosin to corroborate the presence of pulmonary tumors. For homing assays, mice were euthanized 24 hours
30 after the intravenous injection of MDA.pFULG cells. Lungs were perfused with PBS, fixed with

freshly prepared formaldehyde (4% in PBS), and frozen in optimum cutting temperature (Sakura Finetek). Cryosections (10 µm) were counterstained with 4',6-diamidino-2-phenylindole analyzed by confocal microscopy. Animal experiments were approved by the Thomas Jefferson University's Institutional Animal Care and Use Committee.

5 A.9. Reagents And Antibodies

[00140] CCL5 (catalog no. 278-RN) and anti-CCR5 APC antibody (catalog no. FAB1802A) were purchased from R&D Systems. A rabbit anti-human CCR5 polyclonal antibody (GenScript; catalog no. A00979) was used for immunohistochemical staining. Rat tail collagen type I was purchased from BD Biosciences. Vicriviroc and maraviroc were obtained from Selleck Chemicals. Luciferin
10 was obtained from Gold Biotechnology.

B. RESULTS

B.1. Active CCL5/CCR5 Signaling In Basal Breast Cancer

[00141] To examine the relative abundance of CCL5 and its receptor CCR5 by genetic subtype, interrogation was conducted within a combined microarray database comprising 2,254 human breast
15 cancer samples from 27 independent studies (21). The relative abundances of CCL5 and CCR5 were significantly increased in the basal and HER-2 subtypes compared with the normal-like, luminal A and luminal B subtypes (Figure 1A). The increased expression of CCL5 and CCR5 correlated positively in individual breast cancer samples and the correlation was highly significant in the basal and HER-2 subtypes (Figures 1B1 to 1B5). The proportion of patients with a
20 CCL5/CCR5-positive signature was more than 58% in the basal and HER-2 subtypes (Figure 1C). In agreement with previous reports, clinical information of the cases in this database showed that the probability to develop metastasis is increased in the basal, luminal B, and HER-2 subtypes (Figure 1D).

[00142] To determine the gene expression signaling pathway associated with enrichment of
25 CCR5 and CCL5, GSEA analysis using KEGG and GO was conducted of these tumor samples (Figure 7A). These studies showed enrichment for gene expression of pathways including lymphocyte activation, Janus-activated kinase (JAK)-STAT signaling, and Toll-like receptor activation (Figure 7A). The receptors for CCL5 include CCR1 and CCR3. Increased expression of CCL5 associated with increased CCR1, but not CCR3, in the basal and HER-2 genetic tumor type
30 (Figures 7B1 to 7B5 and 7C1 to 7C5). In ER-negative patients receiving chemotherapy, there was

an insignificant trend toward reduced metastasis-free survival and relapse-free survival in the increased CCR5 population, compared with the population with reduced CCR5 expression (Figures 7D1 and 7D2). A comparison of expression levels for CCL5 versus CCR5, CCR1, and CCR3, comparing normal breast with breast cancer showed increased correlation between receptor and ligand expression levels in tumors compared with healthy breast tissue (Figure 8A).

B.2. Ccl5 Promotes Breast Cancer Ca^{2+} Signaling And Cellular Invasion

[00143] Three human breast cancer cell lines with a basal phenotype and molecular signature: MDA-MB-231, Hs578T, and SUM-159 (34–37) were used as models in the studies described herein. Analysis of CCR5 expression by FACS showed that a small subpopulation of cells were positive for the receptor in all 3 cell lines (Figure 2A for MDA-MB-231 and Figures 9A and 9C for Hs578T and SUM-159). Because CCR5 activation induces calcium flux (38, 39), the activation of calcium signaling was assessed by CCL5. Addition of CCL5 to the cultures induced immediate calcium fluxes in a subpopulation of cells (Figure 2B for MDA-MB-231 and Figures 9B1/9B2 and 9D1/9D2 for Hs578T and SUM-159), providing evidence that CCR5 is functional in basal breast cancer cells. As a positive control, the same cultures were exposed to 5% FBS (40). Calcium flux, assessed by relative fluorescence intensity, increased in more than 95% of the cells after FBS addition (Figure 2B1/2B2 and Figures 9B1/9B2 and 9D1/9D2). To further distinguish CCL5-dependent signaling, SUM159 cells were stably transduced with a CCR5 expression vector and the Ca^{+2} response to CCL5 versus FBS was conducted (Figures 10C and 10D vs. 10E and 10F). CCR5 induced Ca^{+2} signaling in the CCR5-overexpressing cells, whereas both lines responded similarly to FBS induced Ca^{+2} activation (Figures 10A1–10F).

[00144] Next, the effect of CCR5 activation on breast cancer cell invasion was assessed using 3D migration assays. CCL5 induced invasion of the basal MDA-MB-231, Hs578T, SUM-159 but not the luminal MCF-7 cells (Figures 2C and 2D). CCL5 promoted invasion of MCF-10A cells engineered to express either NeuT, H-Ras, or c-Src oncogenes, compared with MCF10A vector-transduced cells (Figure 2E and 2F), suggesting that CCL5 responsiveness may be acquired during transformation and requires specific cooperative oncogenic signals. The finding that CCL5 induced cellular invasion led us to examine the migratory capacity of CCR5^{+} cells versus that of CCR5^{-} cells. Within the same SUM-159 breast cancer cell line, CCR5^{+} cells showed an approximately 40-fold greater cellular invasiveness (Figure 2G and 2H), indicating that the expression of CCR5 correlates with a proinvasive phenotype.

B.3. CCR5 Antagonists Block Breast Cancer Calcium Signaling And Cell Invasion

[00145] The importance of CCR5 in HIV infection led to the development of different drugs that target this receptor. Therefore, examination of whether the CCR5 antagonists maraviroc and vicriviroc were capable of blocking the CCL5/CCR5 signaling in basal breast cancer cells were conducted. Both CCR5 antagonists blocked CCL5-induced calcium mobilization. In MDA-MB-231 cells, maraviroc and vicriviroc inhibited calcium responses by 65% and 90%, respectively (Figure 3A and 3B). Similar observations were made with both drugs in Hs578T cells (Figures 3C1 to 3C4 and 3D), indicating that CCR5 expressed in different basal breast cancer cells is sensitive to pharmacologic inhibition.

[00146] To evaluate the functional relevance of CCR5 in cellular migration and invasion, the effects of maraviroc and vicriviroc were tested in 3D invasion assays. Using 2 different cell lines, it was found that both CCR5 antagonists inhibited FBS-induced breast cancer cell invasion at the clinically relevant concentration of 100 nmol/L (Figures 4A-4D). Thus, the proinvasive effect of CCR5 can be abrogated by using specific antagonists.

B.4. CCR5 Inhibition Blocks Breast Cancer Metastasis *In Vivo*

[00147] In view of the finding that CCR5 inhibition by CCR5 antagonists reduced calcium signaling and cell invasion, the *in vivo* effect of maraviroc on lung metastasis was determined. MDA-MB-231 cells transduced within the Luc2-eGFP lentiviral vector (MDA.pFULG cells) were used in an experimental metastasis model. The *Luc2* gene is a codon-optimized version of *Luc* and cells expressing this reporter were 10 to 100 times brighter than the unmodified *Luc* gene (30). After injection of MDA.pFULG cells into the tail vein of mice, noninvasive BLI enabled the early detection of breast cancer metastasis (41). Weekly BLI was conducted for 5 weeks and the radiance antemortem was used as a surrogate measurement of tumor burden. Mice treated with maraviroc (8 mg/kg twice daily) showed a significant reduction in both the number and the size of pulmonary metastases compared with vehicle-treated mice (Figures 5A and B, Figures 11A and 11B). To avoid the possibility that metastases were missed because of inappropriate imaging, *ex vivo* imaging, India ink staining (Figures 5C1 and 5C2), and histology (Figure 5E) of the lungs were conducted. Histologic analysis corroborated that tumor burden corresponds to bioluminescence, as previously shown (30). Metastatic tumors were still detectable in 50% of the maraviroc-treated mice, but their mean size was reduced by 65% (Figures 5D and 5F). Interestingly, analysis of CCR5 expression in lungs from control mice showed an 8-fold enrichment of the CCR5⁺ fraction (Figures 12A to 12F).

Collectively, these results provide evidence that CCR5 antagonists reduce breast cancer metastasis *in vivo*.

B.5. CCR5 Antagonist Impairs Lung Colonization But Not Cell Proliferation Or Tumor Growth

- 5 [00148] It was determined whether the reduction in metastatic tumors by maraviroc involved changes in cellular proliferation and/or target organ colonization. The effect of CCR5 inhibition on cell viability and proliferation both *in vitro* and *in vivo* was analyzed. Maraviroc or vicriviroc treatment of MDA-MB-231 cells for 48 hours did not affect the MTT reduction, which was used as a surrogate measurement of cancer cell number (Figure 6A). In agreement, over expression of
- 10 CCR5 in MDA-MB-231 cells did not modify their proliferation rate compared with cells transfected with the empty vector (Figures 6B and 6C). Finally, maraviroc treatment of mice with established pulmonary metastasis did not modify tumor growth (Figures 6D and 6E), indicating that CCR5 activation does not promote the proliferation of basal breast cancer cells *in vitro* nor in the pulmonary microenvironment of immunocompromised mice.
- 15 [00149] On a different *in vivo* experiment, the effect of maraviroc on breast cancer cell homing to lungs was examined. To reach a steady-state concentration in plasma and tissues, mice were given 10 administrations of maraviroc (twice a day for 5 days) before the intravenous injection of MDA.pFULG cells (Figure 6F). Inoculation of equal numbers of MDA.pFULG cells in control and treated groups was corroborated by BLI immediately after injection. Maraviroc reduced the number
- 20 of eGFP⁺ cells in the lungs by 40% (Figures 6G and 6H), suggesting that the *in vivo* antimetastatic effect of maraviroc is caused by a reduction in the number of cancer cells that colonize the target organ from the circulation.

C. Discussion

- [00150] The current studies show for the first time that: (i) enrichment of CCL5/CCR5 expression
- 25 occurs in patients with basal and Her2 positive genetic subtypes of breast cancer; (ii) oncogenic transformation of immortalized human breast cells by distinct oncogenes induces CCL5 responsiveness; and (iii) maraviroc, an FDA-approved drug for the treatment of CCR5-trophic HIV infection, reduce metastatic tumor burden *in vivo*.

- [00151] Previous studies showed that CCL5 levels are elevated in breast primary and metastatic
- 30 tumors (9–11), suggesting a role of CCL5 in the acquisition of malignancy. The present disclosure show that increased expression of CCL5 and CCR5 are associated and that CCL5/CCR5 expression

levels are different among the different genetic subtypes of breast cancer. Increased expression of CCL5 and CCR5 is found in the basal and HER-2 subtypes. In agreement, increased CCL5 expression has been found predominantly in ER-negative patients (42). Increased CCL5 also correlated with increased CCR1 in basal and Her2 genetic subtypes of breast cancer. A trend toward reduced metastasis-free survival and relapse-free survival was observed among the CCR5-overexpressing tumors in patients who received chemotherapy.

[00152] Given the aggressive clinical behavior of basal breast cancer and the lack of targeted therapies for it, the importance of the CCL5/CCR5 axis in invasion and metastasis was evaluated in the human breast cancer cell lines MDA-MB-231, Hs578T and SUM-159. These cell lines reflect the clinicopathologic features of the basal subtype of breast cancer (including the lack of HER-2, ER, and progesterone receptor), a basal-like molecular signature, the activation of specific signaling pathways (e.g., hypoxic or EGF receptor responses) and over expression of epithelial-mesenchymal transition proteins (FN, VIM, and matrix metalloproteinase 2; refs. 34-37). Only a small fraction of cells within the cell lines used in this study expressed CCR5 as evaluated by FACS analysis. Our findings are consistent with studies by Müller and colleagues who showed CCR5 expression in MDA-MB-231 by quantitative real-time PCR (8). These studies confirmed the expression of CCR5 in MDA-MB-23 cells by reverse transcriptase PCR and showed the presence of the CCR5 protein by FACS analysis (Figures 13A1-13D), and showed that CCR5 immunohistochemical staining was localized primarily to the breast cancer epithelial cell, compared with normal breast tissue (Figures 14A-14D).

[00153] The results herein show that CCL5 activates calcium flux in basal-like human breast cancer cells, as previously described in cells of the immune system (39, 43) and CCR5-transfected cells (27, 44, 45). By using the selective CCR5-antagonists maraviroc and vicriviroc (both with IC₅₀ below 30 nmol/L; refs. 44, 45), it was shown that CCL5-activated signaling is mediated by CCR5. However, the fraction of CCL5-responsive cells (10% and 12% for MDA-MB-231 and Hs578T cells, respectively) is higher than the percentage of CCR5-expressing cells determined by FACS. This may be due to the greater sensitivity of the Ca²⁺ activation assays compared with the sensitivity of analysis by FACS. In addition, CCL5-induced calcium redistribution is not completely blocked by CCR5 antagonists. This may be caused by the expression of other receptors to CCL5, namely CCR1 and CCR3. CCR5 has been identified as the main CCL5 receptor in MDA-MB-231 cells (13) and CCR1 and CCR3 transcripts are absent in both MDA-MB-231 or Hs578T cell lines (8) and breast tumor samples (11). CCR1 and CCR3 were able to be detected by FACS (Figures 13A1-13D), suggesting a possible mechanism for the incomplete response to the CCR5 antagonist.

[00154] It was observed that the subpopulation of CCR5⁺ cells displayed increased invasiveness, indicating that CCR5 favors cell migration and invasion in basal-like breast cancer cells. The failure of luminal-like MCF-7 cells to respond to CCL5 is in agreement with previous publications (12). These studies also showed that CCR5 inhibition with either maraviroc or vicriviroc reduced *in vitro* FBS-induced breast cancer cellular invasion without affecting cellular viability. The finding that CCR5 antagonists block FBS-induced invasion is novel and suggested that CCR5 activation contribute to the production of metastasis *in vivo* where different chemotactic and growth signals are present. The mechanisms involved in CCR5 regulation of FBS-activated invasiveness are uncharacterized but they may include heterodimerization and ligand affinity regulation of other GPCRs (46), or the transactivation of growth factor receptor (47) or integrin-mediated signaling (48), as described in noncancerous cells.

[00155] The *in vivo* antimetastatic effect of maraviroc was shown by injecting MDA.pFULG cells into the circulation of immunodeficient mice and treating them with clinically relevant doses of the drug. In humans, oral doses of 300 mg produce an average $C_{\max}$ of 1,200 nmol/L (49), whereas in mice 16 mg/kg produce an average $C_{\max}$ of 1,045 nmol/L (32). Because the drug is taken twice a day in the clinical setting, 16 mg/kg/d divided into 2 doses and administered during the experiments described herein. Maraviroc significantly reduced the pulmonary tumor burden. Although it has been proposed that pharmacologic CCR5 inhibition may be beneficial for patients with breast cancer, to our knowledge this is the first study showing that systemic administration of a CCR5 antagonist reduces metastatic colonization of basal breast cancer cells.

[00156] The antimetastatic effect of maraviroc is not caused by alterations in growth of established metastasis. CCR5 activation by CCL5 drives proliferation in CCR5-transfected MCF-7 breast cancer cells (18) and prostate cancer cells (50), but this study and others (13) showed that the CCL5/CCR5 axis does not play a role in cell proliferation or survival in the basal-like MDA-MB-231 cells. Furthermore, inhibition of CCR5 surface expression through a dominant-negative form of CCR5 (CCR5 Δ 32) in MDA-MB-231 cells does not change *in vivo* proliferation or apoptotic response (17). On the other hand, it was found that maraviroc reduced lung colonization by MDA.pFULG cancer cells. This result is consistent with previous studies in which inhibition of CCR5 expression within breast cancer cells or administration of anti-CCL5 neutralizing antibody to tumor-bearing mice reduced the enhanced metastatic capability induced by coinjection of mesenchymal stem cells (MSC; ref. 13). The authors identified cancer cell extravasation as the crucial metastatic step affected by CCL5/CCR5 inhibition (13). Together, these data support a role for CCR5 antagonists in blocking the ability of basal breast cancer cells to reach the metastatic sites

instead of inhibiting their proliferation or survival after arrival. Blocking the homing of cancer cells to metastatic sites is a desirable characteristic in a true antimetastatic drug (51). Therefore, CCR5 antagonists may be useful as adjuvant therapy for breast basal tumors with CCR5 over expression or other tumor types where CCR5 promotes metastasis, such as prostate cancer (50) or gastric cancer (52).

[00157] It will be appreciated by those skilled in the art that changes could be made to the exemplary embodiments shown and described above without departing from the broad inventive concept thereof. It is understood, therefore, that this invention is not limited to the exemplary embodiments shown and described, but it is intended to cover modifications within the spirit and scope of the present invention as defined by the claims. For example, specific features of the exemplary embodiments may or may not be part of the claimed invention and features of the disclosed embodiments may be combined. Unless specifically set forth herein, the terms “a”, “an” and “the” are not limited to one element but instead should be read as meaning “at least one”.

[00158] It is to be understood that at least some of the figures and descriptions of the invention have been simplified to focus on elements that are relevant for a clear understanding of the invention, while eliminating, for purposes of clarity, other elements that those of ordinary skill in the art will appreciate may also comprise a portion of the invention. However, because such elements are well known in the art, and because they do not necessarily facilitate a better understanding of the invention, a description of such elements is not provided herein.

[00159] Further, to the extent that the method does not rely on the particular order of steps set forth herein, the particular order of the steps should not be construed as limitation on the claims. The claims directed to the method of the present invention should not be limited to the performance of their steps in the order written, and one skilled in the art can readily appreciate that the steps may be varied and still remain within the spirit and scope of the present invention.

[00160] All references, including publications, patent applications, and patents, cited herein are hereby incorporated by reference to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein.

II. REFERENCES

1. Parkin DM, Fernandez LM. Use of statistics to assess the global burden of breast cancer. *Breast J* 2006; 12 Suppl 1: S70–80.
2. Group EBCTC. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet* 2005; 365: 1687–717.
3. Meyers MO, Klauber-Demore N, Ollila DW, Amos KD, Moore DT, Drobish AA, *et al.* Impact of breast cancer molecular subtypes on locoregional recurrence in patients treated with neoadjuvant chemotherapy for locally advanced breast cancer. *Ann Surg Oncol* 2011; 18: 2851–7.
4. Kennecke H, Yerushalmi R, Woods R, Cheang MC, Voduc D, Speers CH, *et al.* Metastatic behavior of breast cancer subtypes. *J Clin Oncol* 2010; 28: 3271–7.
5. Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, *et al.* Molecular portraits of human breast tumours. *Nature* 2000; 406: 747–52.
- 15 6. Reis-Filho JS, Lakhani SR . Breast cancer special types: why bother? *J Pathol* 2008; 216: 394–8.
7. Kakinuma T, Hwang ST . Chemokines, chemokine receptors, and cancer metastasis. *J Leukoc Biol* 2006; 79: 639–51.
8. Muller A, Homey B, Soto H, Ge N, Catron D, Buchanan ME, *et al.* Involvement of chemokine receptors in breast cancer metastasis. *Nature* 2001; 410: 50–6.
- 20 9. Luboshits G, Shina S, Kaplan O, Engelberg S, Nass D, Lifshitz-Mercer B, *et al.* Elevated expression of the CC chemokine regulated on activation, normal T cell expressed and secreted (RANTES) in advanced breast carcinoma. *Cancer Res* 1999; 59: 4681–7.
10. Niwa Y, Akamatsu H, Niwa H, Sumi H, Ozaki Y, Abe A . Correlation of tissue and plasma RANTES levels with disease course in patients with breast or cervical cancer. *Clin Cancer Res* 2001; 7: 285–9.
- 25 11. Zhang Y, Yao F, Yao X, Yi C, Tan C, Wei L, *et al.* Role of CCL5 in invasion, proliferation and proportion of CD44+/CD24- phenotype of MCF-7 cells and correlation of CCL5 and CCR5 expression with breast cancer progression. *Oncol Rep* 2009; 21: 1113–21. Medline
- 30 12. Jiao X, Katiyar S, Willmarth NE, Liu M, Ma X, Flomenberg N, *et al.* c-Jun induces

mammary epithelial cellular invasion and breast cancer stem cell expansion. *J Biol Chem* 2010; 285: 8218–26.

13. Karnoub AE, Dash AB, Vo AP, Sullivan A, Brooks MW, Bell GW, *et al.* Mesenchymal stem cells within tumour stroma promote breast cancer metastasis. *Nature* 2007; 449: 557–63.
14. Locati M, Deuschle U, Massardi ML, Martinez FO, Sironi M, Sozzani S, *et al.* Analysis of the gene expression profile activated by the CC chemokine ligand 5/RANTES and by lipopolysaccharide in human monocytes. *J Immunol* 2002; 168: 3557–62.
15. Robinson SC, Scott KA, Balkwill FR . Chemokine stimulation of monocyte matrix metalloproteinase-9 requires endogenous TNF-alpha. *Eur J Immunol* 2002; 32: 404–12.
16. Robinson SC, Scott KA, Wilson JL, Thompson RG, Proudfoot AE, Balkwill FR . A chemokine receptor antagonist inhibits experimental breast tumor growth. *Cancer Res* 2003; 63: 8360–5.
17. Manes S, Mira E, Colomer R, Montero S, Real LM, Gomez-Mouton C, *et al.* CCR5 expression influences the progression of human breast cancer in a p53-dependent manner. *J Exp Med* 2003; 198: 1381–9.
18. Murooka TT, Rahbar R, Fish EN . CCL5 promotes proliferation of MCF-7 cells through mTOR-dependent mRNA translation. *Biochem Biophys Res Commun* 2009; 387: 381–6.
19. Stormes KA, Lemken CA, Lepre JV, Marinucci MN, Kurt RA . Inhibition of metastasis by inhibition of tumor-derived CCL5. *Breast Cancer Res Treat* 2005; 89: 209–12.
20. Jayasinghe MM, Golden JM, Nair P, O'Donnell CM, Werner MT, Kurt RA . Tumor-derived CCL5 does not contribute to breast cancer progression. *Breast Cancer Res Treat* 2008; 111: 511–21.
21. Ertel A, Dean JL, Rui H, Liu C, Witkiewicz AK, Knudsen KE, *et al.* RB-pathway disruption in breast cancer: differential association with disease subtypes, disease-specific prognosis and therapeutic response. *Cell Cycle* 2010; 9: 4153–63.
22. Wilkin TJ, Su Z, Krambrink A, Long J, Greaves W, Gross R, *et al.* Three-year safety and efficacy of vicriviroc, a CCR5 antagonist, in HIV-1-infected treatment-experienced patients. *J Acquir Immune Defic Syndr* 2010; 54: 470–6.
23. Barrett T, Troup DB, Wilhite SE, Ledoux P, Rudnev D, Evangelista C, *et al.* NCBI GEO:

mining tens of millions of expression profiles—database and tools update. *Nucleic Acids Res* 2007; 35: D760–5.

24. Brazma A, Parkinson H, Sarkans U, Shojatalab M, Vilo J, Abeygunawardena N, *et al.* ArrayExpress—a public repository for microarray gene expression data at the EBI. *Nucleic Acids Res* 2003; 31: 68–71.
25. Hu Z, Fan C, Oh DS, Marron JS, He X, Qaqish BF, *et al.* The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics* 2006; 7: 96.
26. Liu M, Casimiro MC, Wang C, Shirley LA, Jiao X, Katiyar S, *et al.* p21CIP1 attenuates Ras- and c-Myc-dependent breast tumor epithelial mesenchymal transition and cancer stem cell-like gene expression in vivo . *Proc Natl Acad Sci U S A* 2009; 106: 19035–9.
27. Nguyen DH, Taub D . Cholesterol is essential for macrophage inflammatory protein 1 beta binding and conformational integrity of CC chemokine receptor 5. *Blood* 2002; 99: 4298–306.
28. Janowski E, Jiao X, Katiyar S, Lisanti MP, Liu M, Pestell RG, *et al.* c-Jun is required for TGF-beta-mediated cellular migration via nuclear Ca(2) signaling. *Int J Biochem Cell Biol* 2011; 43: 1104–13.
29. Velasco-Velazquez MA, Agramonte-Hevia J, Barrera D, Jimenez-Orozco A, Garcia-Mondragon MJ, Mendoza-Patino N, *et al.* 4-Hydroxycoumarin disorganizes the actin cytoskeleton in B16-F10 melanoma cells but not in B82 fibroblasts, decreasing their adhesion to extracellular matrix proteins and motility. *Cancer Lett* 2003; 198: 179–86.
30. Liu H, Patel MR, Prescher JA, Patsialou A, Qian D, Lin J, *et al.* Cancer stem cells from human breast tumors are involved in spontaneous metastases in orthotopic mouse models. *Proc Natl Acad Sci U S A* 2010; 107: 18115–20.
31. Zahler MH, Irani A, Malhi H, Reutens AT, Albanese C, Bouzahzah B, *et al.* The application of a lentiviral vector for gene transfer in fetal human hepatocytes. *J Gene Med* 2000; 2: 186–93.
32. Walker DK, Abel S, Comby P, Muirhead GJ, Nedderman AN, Smith DA . Species differences in the disposition of the CCR5 antagonist, UK-427,857, a new potential treatment for HIV. *Drug Metab Dispos* 2005; 33: 587–95.
33. Wu K, Katiyar S, Li A, Liu M, Ju X, Popov VM, *et al.* Dachshund inhibits oncogene-induced breast cancer cellular migration and invasion through suppression of interleukin-8.

Proc Natl Acad Sci U S A 2008; 105: 6924–9.

34. Charafe-Jauffret E, Ginestier C, Monville F, Finetti P, Adelaide J, Cervera N, *et al.* Gene expression profiling of breast cell lines identifies potential new basal markers. *Oncogene* 2006; 25: 2273–84.
- 5 35. Riaz M, Elstrodt F, Hollestelle A, Dehghan A, Klijn JG, Schutte M . Low-risk susceptibility alleles in 40 human breast cancer cell lines. *BMC Cancer* 2009; 9: 236.
36. Hollestelle A, Nagel JH, Smid M, Lam S, Elstrodt F, Wasielewski M, *et al.* Distinct gene mutation profiles among luminal-type and basal-type breast cancer cell lines. *Breast Cancer Res Treat* 2010; 121: 53–64.
- 10 37. Kao J, Salari K, Bocanegra M, Choi YL, Girard L, Gandhi J, *et al.* Molecular profiling of breast cancer cell lines defines relevant tumor models and provides a resource for cancer gene discovery. *PLoS ONE* 2009; 4: e6146.
38. Mueller A, Mahmoud NG, Goedecke MC, McKeating JA, Strange PG . Pharmacological characterization of the chemokine receptor, CCR5. *Br J Pharmacol* 2002; 135: 1033–43.
- 15 39. Petkovic V, Moghini C, Paoletti S, Uguccioni M, Gerber B. I-TAC/CXCL11 is a natural antagonist for CCR5. *J Leukoc Biol* 2004; 76: 701–8.
40. Miyashita M, Smith MW, Willey JC, Lechner JF, Trump BF, Harris CC . Effects of serum, transforming growth factor type beta, or 12-O-tetradecanoyl-phorbol-13-acetate on ionized cytosolic calcium concentration in normal and transformed human bronchial epithelial cells. *Cancer Res* 1989; 49: 63–7.
- 20
41. Prescher JA, Contag CH . Guided by the light: visualizing biomolecular processes in living animals with bioluminescence. *Curr Opin Chem Biol* 2010; 14: 80–9.
42. Yaal-Hahoshen N, Shina S, Leider-Trejo L, Barnea I, Shabtai EL, Azenshtein E, *et al.* The chemokine CCL5 as a potential prognostic factor predicting disease progression in stage II breast cancer patients. *Clin Cancer Res* 2006; 12: 4474–80.
- 25
43. Shideman CR, Hu S, Peterson PK, Thayer SA . CCL5 evokes calcium signals in microglia through a kinase-, phosphoinositide-, and nucleotide-dependent mechanism. *J Neurosci Res* 2006; 83: 1471–84.
44. Dorr P, Westby M, Dobbs S, Griffin P, Irvine B, Macartney M, *et al.* Maraviroc (UK-427,857), a potent, orally bioavailable, and selective small-molecule inhibitor of chemokine
- 30

receptor CCR5 with broad-spectrum anti-human immunodeficiency virus type 1 activity. *Antimicrob Agents Chemother* 2005; 49: 4721–32.

- 5 45. Strizki JM, Tremblay C, Xu S, Wojcik L, Wagner N, Gonsiorek W, *et al.* Discovery and characterization of vicriviroc (SCH 417690), a CCR5 antagonist with potent activity against human immunodeficiency virus type 1. *Antimicrob Agents Chemother* 2005; 49: 4911–9.
46. Isik N, Hereld D, Jin T . Fluorescence resonance energy transfer imaging reveals that chemokine-binding modulates heterodimers of CXCR4 and CCR5 receptors. *PLoS ONE* 2008; 3: e3424.
- 10 47. Mira E, Lacalle RA, Gonzalez MA, Gomez-Mouton C, Abad JL, Bernad A, *et al.* A role for chemokine receptor transactivation in growth factor signaling. *EMBO Rep* 2001; 2: 151–6.
48. Thirkill TL, Lowe K, Vedagiri H, Blankenship TN, Barakat AI, Douglas GC . Macaque trophoblast migration is regulated by RANTES. *Exp Cell Res* 2005; 305: 355–64.
- 15 49. Abel S, van der Ryst E, Rosario MC, Ridgway CE, Medhurst CG, Taylor-Worth RJ, *et al.* Assessment of the pharmacokinetics, safety and tolerability of maraviroc, a novel CCR5 antagonist, in healthy volunteers. *Br J Clin Pharmacol* 2008; 65 Suppl 1: 5–18.
50. Vaday GG, Peehl DM, Kadam PA, Lawrence DM. Expression of CCL5 (RANTES) and CCR5 in prostate cancer. *Prostate* 2006; 66: 124–34.
51. Perret GY, Crepin M . New pharmacological strategies against metastatic spread. *Fundam Clin Pharmacol* 2008; 22: 465–92.
- 20 52. Sugasawa H, Ichikura T, Tsujimoto H, Kinoshita M, Morita D, Ono S, *et al.* Prognostic significance of expression of CCL5/RANTES receptors in patients with gastric cancer. *J Surg Oncol* 2008; 97: 445–50.

CLAIMS

1. A method of determining whether a subject having cancer is at risk for developing metastasis of the cancer, the method comprising:

5 (a) obtaining a biological sample from the subject having cancer;

(b) determining CCR5 level of expression and/or level of expression of at least one of CCR5 ligands in the biological sample; and

(c) if the CCR5 level of expression and/or the level of expression of at least one of CCR5 ligands determined in step (b) is increased compared to CCR5 level of expression and/or
10 level of expression of at least one of CCR5 ligands in a control sample, then the subject is identified as likely at risk for developing metastasis of the cancer.

2. The method of claim 1, further comprising treating, preventing, or managing the cancer metastasis if the subject is identified as likely at risk for developing metastasis of the cancer by administering to the subject a CCR5 antagonist.

15 3. The method of claim 2, wherein the CCR5 antagonist blocks metastasis of the cancer.

4. The method of claim 2, wherein the CCR5 antagonist is administered to the subject in addition to a primary, main or initial treatment for the cancer.

5. The method of claim 1, wherein said cancer is selected from the group consisting of acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic leukemia, promyelocytic leukemia, myelomonocytic leukemia, monocytic leukemia, erythroleukemia
20 leukemia, myelodysplastic syndrome; chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas, Hodgkin's disease, non-Hodgkin's disease; multiple myelomas, smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary
25 plasmacytoma; Waldenstrom's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas, bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma,
30 lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors, glioma,

astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including but not limited to ductal carcinoma, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer, pheochromocytom and adrenocortical carcinoma; thyroid cancer, papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers, Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipius; eye cancers, ocular melanoma, iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers, squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers, squamous cell carcinoma, and adenocarcinoma; uterine cancers, endometrial carcinoma, uterine sarcoma; ovarian cancers, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers, hepatocellular carcinoma and hepatoblastoma; gallbladder cancers, adenocarcinoma; cholangiocarcinomas, pappillary cancer, nodular cancer, diffuse cancer; lung cancers, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers, germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers, prostatic intraepithelial neoplasia, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penal cancers; oral cancers, squamous cell carcinoma; basal cancers; salivary gland cancers, adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; pharynx cancers, squamous cell cancer, and verrucous; skin cancers, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers, renal cell carcinoma, adenocarcinoma, hypemephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/or uterer); Wilms' tumor; bladder cancers, transitional cell carcinoma, squamous cell cancer,

adenocarcinoma, carcinosarcoma, myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas.

5 6. The method of claim 1, wherein said cancer is selected from the group consisting of basal breast cancer and Her2 positive breast cancer.

7. The method of claim 2, wherein said CCR5 antagonist is selected from the group consisting of 4,4-difluoro-N-[(1S)-3-[(1R,5S)-3-(3-methyl-5-propan-2-yl-1,2,4-triazol-4-yl)-8-azabicyclo[3.2.1]octan-8-yl]-1-phenylpropyl]cyclohexane-1-carboxamide ("Maraviroc") and (4,6-
10 dimethylpyrimidin-5-yl)-[4-[(3S)-4-[(1R)-2-methoxy-1-[4-(trifluoromethyl)phenyl]ethyl]-3-methylpiperazin-1-yl]-4-methylpiperidin-1-yl]methanone ("Vicriviroc").

8. The method of claim 2, wherein said CCR5 antagonist is selected from the group consisting of: 4,4-difluoro-N-[(1S)-3-[(1R,5S)-3-(3-methyl-5-propan-2-yl-1,2,4-triazol-4-yl)-8-azabicyclo[3.2.1]octan-8-yl]-1-phenylpropyl]cyclohexane-1-carboxamide ("Maraviroc");

15 N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropylcyclobutanecarboxamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropylcyclopentanecarboxamide;

20 N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropyl-4,4,4-trifluorobutanamide;

N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-phenylpropyl-4,4-difluorocyclohexanecarboxamide;

25 N-(1S)-3-3-(3-Isopropyl-5-methyl-4H-1,2,4-triazol-4-yl)-exo-8-azabicyclo[3.2.1]oct-8-yl-1-(3-fluorophenyl)propyl-4,4-difluorocyclohexanecarboxamide; and pharmaceutically acceptable salts or solvates thereof.

9. The method of claim 2, wherein said CCR5 antagonist is selected from the group consisting of:

N-{3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

N-{(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclobutanecarboxamide;

5 N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-2H-pyran-4-carboxamide;

1-Acetyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}3-azetidine carboxamide;

10 1-Hydroxy-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

2-Methyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopropanecarboxamide;

2-Cyclopropyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo [3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

15 N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-3-furancarboxamide;

3,3,3-Trifluoro-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

20 N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}tetrahydro-2-furancarboxamide;

1-(Acetylamino)-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

N-{(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

25 1-Methoxy-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

1-Amino-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}cyclopentanecarboxamide;

1-Methyl-N-{(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-2-oxo-4-pyrrolidinecarboxamide;

1-Acetyl-N-{(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}3-azetidinecarboxamide;

5 N-{(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

N-{(1S)-3-[6-(2-Methyl-1H-benzimidazol-1-yl)-3-azabicyclo[3.1.0]hex-3-yl]-1-phenylpropyl}cyclobutanecarboxamide;

10 2-Cyclopropyl-N-{(1S)-3-[3-exo-(3-{4-[(methylsulfonyl)amino]benzyl}-1,2,4-oxadiazol-5-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

N-{(1S)-3-[7-exo-(2-Methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}cyclobutanecarboxamide;

2-Cyclopropyl-N-{(1S)-3-[7-exo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}acetamide;

15 3,3,3-Trifluoro-N-{(1S)-3-[7-exo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}propanamide;

N-{(1S)-3-[7-endo-(2-Methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}cyclobutanecarboxamide;

20 2-Cyclopropyl-N-{(1S)-3-[7-endo-(2-methyl-1H-benzimidazol-1-yl)-3-oxa-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}acetamide;

N-{(1S)-3-[7-exo-(2-Methyl-1H-benzimidazol-1-yl)-3-thia-9-azabicyclo[3.3.1]non-9-yl]-1-phenylpropyl}cyclobutanecarboxamide;

2-Cyclopropyl-N-[(1S)-3-(3-endo-{[2-(4-fluorophenyl)acetyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]acetamide;

25 N-[(1S)-3-(3-{[3-endo-(4-Fluorophenyl)propanoyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]cyclobutanecarboxamide;

N-[(1S)-3-(3-{[3-exo-(4-Fluorophenyl)propanoyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]cyclobutanecarboxamide;

2-Cyclopropyl-N-[(1S)-3-(3-exo-{[2-(4-fluorophenyl)acetyl]amino}-8-azabicyclo[3.2.1]oct-8-yl)-1-phenylpropyl]acetamide;

N-[(1S)-3-[3-exo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-1-propionyl-3-azetidinecarboxamide;

5 N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]tetrahydro-3-furancarboxamide;

N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]tetrahydro-2H-pyran-4-carboxamide;

10 N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]tetrahydro-2-furancarboxamide;

1-Acetyl-N-[(1S)-3-[3-endo-(1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-3-azetidinecarboxamide;

N-[(1S)-3-[3-endo-(1H-Benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-1-propionyl-3-azetidinecarboxamide;

15 Methyl 3-[(1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]amino)carbonyl]-1-azetidinecarboxylate;

N-[(1S)-3-[3-endo-(2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-1-propionyl-3-azetidinecarboxamide 1-Acetyl-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-2-azetidinecarboxamide;

20 2-[Acetyl(methyl)amino]-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]acetamide;

3-[Acetyl(methyl)amino]-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]propanamide;

25 2-Methoxy-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]acetamide;

3-Methoxy-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]propanamide;

1-Acetyl-N-[(1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl]-3-pyrrolidinecarboxamide;

1-Methyl-N-((1S)-3-[3-endo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-oxo-4-pyrrolidinecarboxamide;

1-Acetyl-N-((1S)-3-[3-exo-(2-ethyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

5 N-((1S)-3-[3-exo-(2-Ethyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;

1-Acetyl-N-((1S)-1-phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)-3-azetidinecarboxamide;

10 N-((1S)-1-Phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)-1-propionyl-3-azetidinecarboxamide;

N-((1S)-1-Phenyl-3-{3-exo[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)acetamide;

2-[Acetyl(methyl)amino]-N-((1S)-1-phenyl-3-{3-exo-[2-(trifluoromethyl)-1H-benzimidazol-1-yl]-8-azabicyclo[3.2.1]oct-8-yl}propyl)acetamide;

15 1-Acetyl-N-((1S)-3-[3-exo-(1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

N(1S)-3-[3-exo-(1H-Benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;

20 1-acetyl-N(1S)-3-[3-exo-(5-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)3-azetidinecarboxamide;

N-((1S)-3-[3-exo-(5-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;

1-Acetyl-N-((1S)-3-[3-exo-(5-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)3-azetidinecarboxamide;

25 N-((1S)-3-[3-exo-(5-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;

N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

1-methyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

(2S)-1-acetyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-azetidinecarboxamide;

5 (2R)-1-acetyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-2-azetidinecarboxamide;

2-[acetyl(methyl)amino]-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;

10 3-[acetyl(methyl)amino]-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;

1-acetyl-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-pyrrolidinecarboxamide;

N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-(trifluoromethyl)cyclopropanecarboxamide;

15 2-methoxy-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;

3-methoxy-N-((1S)-3-[3-exo-(2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)propanamide;

20 1-Acetyl-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

1-Methyl-N-((1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-3-azetidinecarboxamide;

25 N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)-1-propionyl-3-azetidinecarboxamide;

2-Methoxy-N-((1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl)acetamide;

N-{(1S)-3-[3-exo-(4-Fluoro-2-Methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

3-Methoxy-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

5 2-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

3-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide;

10 N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-methyl-3-oxetanecarboxamide;

3-Ethyl-N-{(1S)-3-[3-exo-(4-fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-2-methyl-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

15 3-Ethyl-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-methyl-3-oxetanecarboxamide;

20 N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-oxetanecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-1-methyl-3-azetidinecarboxamide;

25 1-Acetyl-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-1-propionyl-3-azetidinecarboxamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-2-methoxyacetamide;

N-{(1S)-3-[3-exo-(4-Fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

5 N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}-3-methoxypropanamide;

2-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}acetamide;

10 3-[Acetyl(methyl)amino]-N-{(1S)-3-[3-exo-(4-fluoro-1H-benzimidazol-1-yl)-8-azabicyclo[3.2.1]oct-8-yl]-1-phenylpropyl}propanamide; and pharmaceutically acceptable salts thereof.

10. The method of claim 4, wherein the metastasis is a tumor metastasis in or more organs selected from the group consisting of liver, brain, bladder, lung, adrenal gland, kidney, bone and combinations thereof.

15 11. The method of claim 2, wherein the CCR5 antagonist is administered concomitantly or concurrently as other therapies for cancer.

12. The method of claim 2, wherein the CCR5 antagonist is administered concurrently as other adjuvant therapies or following other adjuvant therapies.

20 13. The method of claim 2, wherein the CCR5 receptor antagonist improves prognosis in the subject having cancer.

14. The method of claim 2, wherein the CCR5 receptor antagonist improves therapeutic effect of concomitantly or concurrently administered cancer therapy by blocking metastasis of the cancer being treated, thereby contributing to improving clinical outcome of the cancer therapy.

Fig. 1A









Fig. 1B1**Fig. 1B2****Fig. 1B3****Fig. 1B4****Fig. 1B5**

Fig. 1C








Fig. 1D









Fig. 2E




Fig. 2F





Fig. 2G






Fig. 3D




Fig. 3C2




Fig. 3C4






Fig. 3C1 Hs578T



Control

Fig. 3C3



Maraviroc
(100 nM)

Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D






Fig. 5A









Fig. 5E




Fig. 5D





Fig. 6F




Fig. 6G




Fig. 6H






Fig. 7A









Fig. 7D1




Fig. 7D2




Fig. 8A




Fig. 8B




Fig. 8C




Fig. 8D




Fig. 9A




Fig. 9B1





Fig. 9B2






Fig. 10A1

SUM159-vector




Fig. 10A2

SUM159-CCR5




Fig. 10B



Fig. 10C




Fig. 10D




Fig. 10E






Fig. 10F








Fig. 13A1




Fig. 13A2





Fig. 13A3



CCR1-APC



Fig. 13A4

— No stain

— IgG

Fig. 13A5

— aCCR1



CCR1-APC





Fig. 13C






Fig. 14A

Breast Tumor




Fig. 14B

Normal Breast




Fig. 14C

Breast Tumor





Fig. 14D

Normal Breast



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US13/40917

A. CLASSIFICATION OF SUBJECT MATTER IPC(8) - A61P 35/04; C12Q 1/25; G01N 33/574 (2013.01) USPC - 435/7.23; 514/235.2, 256 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC(8): A61P 35/04; C12Q 1/25; G01N 33/574 (2013.01) USPC: 435/7.23; 514/235.2, 256 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) MicroPatent (US-G, US-A, EP-A, EP-B, WO, JP-bib, DE-C,B, DE-A, DE-T, DE-U, GB-A, FR-A); DialogPro; Google Scholar; IP.com; assess*, evaluat*, determin*, identifi*, patient*, subject*, cancer*, metastas*, neoplas*, tumor*, risk*, susceptib*, measur*, detect*, CCR*2, cytokin*, express*, level*, amount*, quantiti*, azabicyclooctan*, treat*, antagoni*		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 2002/0048786 A1 (ROSEN, CA, et al.) 25 April 2002; paragraphs [0117]-[0118], [0420], [0476], [0502]-[0503], [0506]-[0507], [0510], [0531], [0584], [0588], [0590], [0669], [0673], [0707], [0709], [0753]	1-5, 11-15 --- 7-10
Y	POTTER, SM et al., 'Systemic chemokine Levels in Breast Cancer Patients and their Relationship with Circulating Menstrual Hormones' Breast Cancer Res Treat 115, 2009, pages 279-287; Retrieved from the Internet <URL: <http://peer.ccsd.cnrs.fr/docs/00/47/82/37/PDF/PEER_stage2_10.1007%252Fs10549-008-0078-2.pdf>; abstract; page 280, column 1, paragraph 1; page 280, column 2, paragraph 2; page 281, column 1, paragraph 1, table 1; page 281, column 2, paragraph 3; page 282, column 1, paragraph 2; page 282, column 2, paragraph 3; page 283, column 2, paragraphs 1-2; page 284, column 1, paragraph 1 - column 2, paragraph 2; page 285, column 1, paragraphs 1-2	7
Y	WO 2006/067584 A1 (PRYDE, DC et al.) 29 June 2006; page 1, lines 4-6, 25-30; page 20, lines 32-37; page 21, lines 1-23; page 22, lines 3-6; page 23, lines 9-10; page 29, lines 4-6, 19-22; page 30, lines 13-14	8-9
Y	US 6,586,430 B1 (ARMOUR, DR et al.) 01 July 2003; column 1, lines 17-24; column 110, lines 5-25	10
A	US 2006/0251651 A1 (SHIBAYAMA, S et al.) 09 November 2006; entire document	1-5, 7-15
P, Y	WO 2012/122499 A2 (PESTELL, RG) 13 September 2012; abstract; paragraphs [0017], [0034], [0190]-[0195]	1-5, 7-15
☐ Further documents are listed in the continuation of Box C. ☐		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 14 October 2013 (14.10.2013)		Date of mailing of the international search report 21 OCT 2013
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201		Authorized officer: Shane Thomas PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774