



US 20100254941A1

(19) **United States**(12) **Patent Application Publication**
Novick et al.(10) **Pub. No.: US 2010/0254941 A1**(43) **Pub. Date: Oct. 7, 2010**(54) **INTERLEUKIN-18 BINDING PROTEINS,
THEIR PREPARATION AND USE FOR THE
TREATMENT OF SEPSIS**(75) Inventors: **Daniela Novick**, Rehovot (IL);
Charles Dinarello, Boulder, CO
(US); **Menachem Rubinstein**,
Rehovot (IL); **Soo Hyun Kim**,
Aurora, CO (US); **Giamila**
Fantuzzi, Denver, CO (US); **Leonid**
L. Reznikov, Aurora, CO (US);
Boris Schwartzburd, Rehovot (IL)

Correspondence Address:

Mintz Levin Cohn Ferris Glovsky and Popeo, P.C.
One Financial Center
Boston, MA 02111 (US)(73) Assignee: **Yeda Research Institute and
Development Ltd.**, Rehovot (IL)(21) Appl. No.: **12/715,089**(22) Filed: **Mar. 1, 2010****Related U.S. Application Data**(63) Continuation of application No. 11/505,554, filed on
Aug. 16, 2006, now Pat. No. 7,704,944, which is a
continuation-in-part of application No. 09/790,338,
filed on Feb. 21, 2001, now Pat. No. 7,220,717, which
is a continuation-in-part of application No. 09/485,
632, filed on Oct. 12, 2000, now Pat. No. 6,605,280.(30) **Foreign Application Priority Data**


Aug. 14, 1997	(IL)	121554
Aug. 27, 1997	(IL)	121639
Sep. 29, 1997	(IL)	121860
Nov. 6, 1997	(IL)	122134
Jul. 22, 1998	(IL)	125463
Aug. 13, 1998	(IL)	PCT/IL98/00379
Feb. 21, 2000	(EP)	00103590.6
Feb. 21, 2000	(EP)	00103597.1
Nov. 23, 2000	(EP)	00125633.8

Publication Classification

(51) Int. Cl.	
A61K 38/21	(2006.01)
A61K 39/395	(2006.01)
A61K 38/19	(2006.01)
A61K 35/00	(2006.01)
A61K 31/713	(2006.01)
A61P 31/02	(2006.01)
A61P 9/00	(2006.01)
(52) U.S. Cl.	424/85.4 ; 424/158.1; 514/1.4; 424/85.7; 424/85.6; 424/93.2; 514/44 R; 424/134.1

(57) **ABSTRACT**

The present invention provides a method of treatment or prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS), including severe sepsis, septic shock, and sepsis related to cardiac dysfunction.






FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D








FIG. 3E






FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B




FIG. 13




FIG. 14



FIG. 15A



FIG. 15B




FIG. 15C




FIG. 16




FIG. 17




FIG. 18




FIG. 19A




FIG. 19B




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27

INTERLEUKIN-18 BINDING PROTEINS, THEIR PREPARATION AND USE FOR THE TREATMENT OF SEPSIS

RELATED APPLICATIONS

[0001] This application is a continuation of U.S. Ser. No. 11/505,554, filed Aug. 16, 2006; which is a continuation-in-part application of U.S. Ser. No. 09/790,338, filed Feb. 21, 2001 (now U.S. Pat. No. 7,220,217); which is a continuation-in-part application of U.S. Ser. No. 09/485,632, filed Oct. 12, 2000 (now U.S. Pat. No. 6,605,280); and claims the benefit of EP application 00103590.6, filed Feb. 21, 2000; EP application 00103597.1, filed Feb. 21, 2000; EP application 00125633.8, filed Nov. 23, 2000; PCT Application IL98/00379, filed Aug. 13, 1998; Israel application 125463, filed Jul. 22, 1998; Israel application 122134, filed Nov. 6, 1997; Israel application 121860, filed Sep. 29, 1997; Israel application 121639, filed Aug. 27, 1997; and Israel application 121554, filed Aug. 14, 1997. Each of these applications is incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0002] The present invention relates to interleukin-18 (IL-18) binding proteins, hereinafter IL-18BP, capable of blocking the activity of IL-18. This invention relates to soluble IL-18BPs obtainable from body fluids, to soluble IL-18BPs obtainable by expression of suitable DNA vectors in host cells, to virus-encoded homologues of IL-18BP obtainable by expression of suitable DNA vectors in host cells, to vectors expressing the various IL-18BPs, to vectors useful for expression of IL-18BPs in humans and other mammals, to antibodies against IL-18BPs, to therapeutic use of said IL-18BPs in blocking IL-18 activity, to therapeutic use of said expression vectors in blocking IL-18 activity and to use of the antibodies.

[0003] More specifically, the present invention provides a method of treatment and prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS), including severe sepsis, septic shock, and sepsis related to cardiac dysfunction.

BACKGROUND OF THE INVENTION

[0004] In 1989, an endotoxin-induced serum activity that induced interferon- γ (IFN- γ) obtained from mouse spleen cells was described (Micallef et al., 1996). This serum activity functioned not as a direct inducer of IFN- γ but rather as a co-stimulant together with IL-2 or mitogens. An attempt to purify the activity from post-endotoxin mouse serum revealed an apparently homogeneous 50-55 kDa protein. Since other cytokines can act as co-stimulants for IFN- γ production, the failure of neutralizing antibodies to IL-1, IL-4, IL-5, IL-6, or TNF to neutralize the serum activity suggested it was a distinct factor. In 1995, the same scientists demonstrated that the endotoxin-induced co-stimulant for IFN- γ production was present in extracts of livers from mice preconditioned with *P. acnes* (Novick et al., 1992). In this model, the hepatic macrophage population (Kupffer cells) expand and in these mice, a low dose of bacterial lipopolysaccharide (LPS), which in non-preconditioned mice is not lethal, becomes lethal. The factor, named IFN- γ -inducing factor (IGIF) and later designated interleukin-18 (IL-18), was purified to homogeneity from 1,200 grams of *P. acnes*-treated mouse livers. Degenerate oligonucleotides derived from amino acid sequences of purified IL-18 were used to clone a

murine IL-18 cDNA (Novick et al., 1992). IL-18 is an 18-19 kDa protein of 157 amino acids, which has no obvious similarities to any peptide in the databases. Messenger RNAs for IL-18 and interleukin-12 (IL-12) are readily detected in Kupffer cells and activated macrophages. Recombinant IL-18 induces IFN- γ more potently than does IL-12, apparently through a separate pathway (Novick et al., 1992). Similar to the endotoxin-induced serum activity, IL-18 does not induce IFN- γ by itself, but functions primarily as a co-stimulant with mitogens or IL-2. IL-18 enhances T cell proliferation, apparently through an IL-2-dependent pathway, and enhances Th1 cytokine production in vitro and exhibits synergism when combined with IL-12 in terms of enhanced IFN- γ production (Maliszewski et al., 1990).

[0005] Neutralizing antibodies to mouse IL-18 were shown to prevent the lethality of low-dose LPS in *P. acnes* preconditioned mice. Others had reported the importance of IFN- γ as a mediator of LPS lethality in pre-conditioned mice. For example, neutralizing anti-IFN- γ antibodies protected mice against Shwartzman-like shock (Fantuzzi et al., 1998), and galactosamine-treated mice deficient in the IFN- γ receptor were resistant to LPS-induced death (Byrn, 1990). Hence, it was not unexpected that neutralizing antibodies to murine IL-18 protected *P. acnes*-preconditioned mice against lethal LPS (Novick et al., 1992). Anti-murine IL-18 treatment also protected surviving mice against severe hepatic cytotoxicity.

[0006] After the murine form was cloned, the human cDNA sequence for IL-18 was reported in 1996 (Okamura et al., 1995). Recombinant human IL-18 exhibits natural IL-18 activity (Okamura et al., 1995). Human recombinant IL-18 is without direct IFN- γ -inducing activity on human T-cells, but acts as a co-stimulant for production of IFN- γ and other T-helper cell-1 (Th1) cytokines (Okamura et al., 1995). To date, IL-18 is thought of primarily as a co-stimulant for Th1 cytokine production (IFN- γ , IL-2 and granulocyte-macrophage colony stimulating factor) (Izaki, 1978) and also as a co-stimulant for FAS ligand-mediated cytotoxicity of murine natural killer cell clones (Novick et al., 1989).

[0007] By cloning IL-18 from affected tissues and studying IL-18 gene expression, a close association of this cytokine with an autoimmune disease was found. The non-obese diabetic (NOD) mouse spontaneously develops autoimmune insulinitis and diabetes, which can be accelerated and synchronized by a single injection of cyclophosphamide. IL-18 mRNA was demonstrated by reverse transcriptase PCR in NOD mouse pancreas during early stages of insulinitis. Levels of IL-18 mRNA increased rapidly after cyclophosphamide treatment and preceded a rise in IFN- γ mRNA, and subsequently diabetes. Interestingly, these kinetics mimic that of IL-12-p40 mRNA, resulting in a close correlation of individual mRNA levels. Cloning of the IL-18 cDNA from pancreas RNA followed by sequencing revealed identity with the IL-18 sequence cloned from Kupffer cells and in vivo pre-activated macrophages. Also NOD mouse macrophages responded to cyclophosphamide with IL-18 gene expression while macrophages from Balb/c mice treated in parallel did not. Therefore, IL-18 expression is abnormally regulated in autoimmune NOD mice and closely associated with diabetes development (Novick et al., 1992).

[0008] IL-18 plays a potential role in immunoregulation or in inflammation by augmenting the functional activity of Fas ligand on Th1 cells (Conti et al., 1997). IL-18 is also expressed in the adrenal cortex and therefore might be a

secreted neuro-immunomodulator, playing an important role in orchestrating the immune system following a stressful experience (Chater, 1986).

[0009] In vivo, IL-18 is formed by cleavage of pro-IL-18, and its endogenous activity appears to account for IFN- γ production in *P. acnes* and LPS-mediated lethality. Because of its activity, blocking the biological activity of IL-18 in human disease is a therapeutic strategy in many diseases. This can be accomplished using soluble receptors or blocking antibodies to the cell-bound IL-18 receptor.

[0010] Cytokine binding proteins (soluble cytokine receptors) correspond to the extracellular ligand binding domains of their respective cell surface cytokine receptors. They are derived either by alternative splicing of a pre-mRNA, common to the cell surface receptor, or by proteolytic cleavage of the cell surface receptor. Such soluble receptors have been described in the past, including among others, the soluble receptors of IL-6 and IFN- γ (Nakamura et al., 1989), TNF (Dao et al., 1996; Engelmann et al., 1989), IL-1 and IL-4 (John, 1986), IFN- α/β (Mizushima and Nagata, 1990) and others. One cytokine-binding protein, named osteoprotegerin (OPG, also known as osteoclast inhibitory factor—OCIF), a member of the TNFR/Fas family, appears to be the first example of a soluble receptor that exists only as a secreted protein (Anderson, 1997; Bolton, 1980). The present invention deals with soluble IL-18 binding proteins.

[0011] Recently, it has been suggested that the interleukin IL-18 is involved in the progression of pathogenicity in chronic inflammatory diseases, including endotoxin shock, hepatitis, and autoimmune-diabetes (Kahiwamura and Okamura, 1998). A further indication of a possible role of IL-18 in the development of liver injury resulted from experiments published by Tsuij et al. (Tsuij et al., 1999), showing an elevated level of IL-18 in lipopolysaccharide-induced acute liver injury in a mouse model. However, the mechanism of the multi-functional factor IL-18 in the development of liver injury has not been elucidated so far.

[0012] Liver damage or injury may have diverse causes. It may be due to viral or bacterial infections, alcohol abuse, immunological disorders, or cancer, for example.

[0013] Viral hepatitis, due to Hepatitis B virus and Hepatitis C virus, for example, are poorly managed diseases that afflict large number of people worldwide. The number of known hepatitis viruses known is constantly increasing. Apart from Hepatitis B and C virus, at least four other viruses causing virus-associated hepatitis have been discovered so far, called Hepatitis A, D, E and G-Virus.

[0014] Alcoholic liver disease is another widespread disease associated with chronic consumption of alcohol. Immune hepatitis is a rare autoimmune disease that is poorly managed. Liver injury also includes damages of the bile ducts. Primary biliary cirrhosis (PBC) is an autoimmune liver disease characterized by destruction of the intrahepatic bile ducts.

[0015] Several studies have demonstrated that damage to the liver in diseases such as alcoholic hepatitis, liver cirrhosis, viral hepatitis and primary biliary cirrhosis is associated with T-helper cell-1 (Th1) responses. In one study, a novel liver injury model was established in mice by targeting of ovalbumin-containing liposomes into the liver, followed by adoptive transfer of ovalbumin-specific Th1 cells. Combined treatment of mice with ovalbumin-containing liposomes and Th1 cell transfer caused an increase in serum transaminase activity that was paralleled with an elevation of serum IFN- γ

levels. In sharp contrast, ovalbumin-specific Th2 cell transfer resulted in an increase of serum IL-4 levels but did not induce liver injury. The liver injury was blocked by anti-IFN- γ antibodies and anti-tumor necrosis factor (TNF)- α antibodies. These findings indicate that Th1 cells are the major effector cells in acute liver injury (Nishimura and Ohta, 1999). In another set of studies it was shown that mice over-expressing IFN- γ exhibit spontaneous hepatitis without any pathogen or any other stimulant (Okamoto et al., 1998).

[0016] Another study implicated Th1 responses in primary biliary cirrhosis (PBC). PBC is an autoimmune liver disease characterized by destruction of the intrahepatic bile ducts. It is generally believed that cellular immune mechanisms, particularly involving T cells, result in this bile duct damage. The relative strength of Th1 and Th2 responses has recently been proposed to be an important factor in the pathophysiology of various autoimmune diseases. In this study, the Th subset balance in PBC was evaluated by detection of cytokines specific to the two T-cell subsets, i.e., IFN- γ for Th1 cells and IL-4 for Th2 cells. IFN- γ and IL-4 messenger RNA (mRNA) positive cells were counted in liver sections from 18 patients with PBC and 35 disease controls including chronic active hepatitis C, extrahepatic biliary obstruction, and normal liver, using nonisotopic in situ hybridization and immunohistochemistry. Mononuclear cells expressing IFN- γ and IL-4 mRNA were aggregated in inflamed portal tracts in PBC livers, but were rarely present in extrahepatic biliary obstruction, alcoholic fibrosis, or normal liver sections. The IFN- γ and IL-4 mRNA positive cells in PBC livers were detected in significantly higher numbers than in control livers ($P < 0.01$). Moreover, IFN- γ mRNA expression was more commonly detected than IL-4 expression in PBC livers, and the levels of IFN- γ mRNA expression were highly correlated with the degree of portal inflammatory activity. IFN- γ mRNA-positive cells were detected primarily around damaged bile ducts that were surrounded by lymphoid aggregates. The data indicate that Th1 cells are the more prominent T-cell subset in the lymphoid infiltrates in PBC (Harada et al., 1997).

[0017] The cytokine pattern on viral antigen recognition is also believed to exert a profound influence on the resolution of viral infections and viral clearance. One study investigated whether a cytokine imbalance oriented toward Th2 type response plays a role in chronic hepatitis B. Cytokine profiles of peripheral blood mononuclear cells associated with chronic hepatitis B were analyzed by RT-PCR. Upon hepatitis B surface antigen (HbsAg) stimulation, expression of IFN- γ , IL-2, IL-4, and IL-10 was detected in 41%, 8%, 41%, and 50% of the patients, respectively. Among these cytokines, the expression of the Th1 cytokine IFN- γ was associated with high levels of serum AST/ALT (Aspartate aminotransferase/Alanine aminotransferase), representing typical markers of liver damage. Th2 type cytokines were not shown to exert a protective effect on hepatocytes. In conclusion, production of a Th1 cytokine, IFN- γ , by HBsAg-reactive cells was associated with hepatocyte damage in chronic hepatitis B (Lee et al., 1999). High levels of the FAS ligand and its receptor (CD95) were reported in liver of hepatitis B patients (Luo et al., 1997). FAS ligand is considered to be one of the major cytotoxic agents leading to hepatocyte apoptosis.

[0018] Another study identified factors associated with the progression of liver injury in 30 hepatitis C virus/RNA (HCV/RNA)-positive untreated patients with chronic hepatitis. Necroinflammatory and architectural damage were evaluated using Ishak's score. Activated hepatic stellate cells (HSC)

were visualized by immunohistochemistry for α -smooth muscle actin (α SMA) and quantitated by morphometry. Plasma HCV/RNA was evaluated using a competitive RT-PCR method. To study the type of immune response involved in the progression of liver injury, IFN- γ -positive cells (as expression of a Th1-like response) were evaluated by immunohistochemistry and quantitated by morphometry. It was found that HSC were mostly detected close to areas of lobular necroinflammation or lining fibrotic septa. The α SMA- and Sirius Red-positive parenchyma correlated significantly with necroinflammatory and architectural scores. IFN γ -positive cells were detected in periportal areas associated with the inflammatory infiltrates and significantly correlated with architectural damage. It was therefore concluded that HSC activation and progression of liver injury are associated with a Th1-like response (Baroni et al., 1999). Similarly to the case of Hepatitis B, FAS ligand and its receptor were found in liver and sera of hepatitis C patients (Hiramatsu et al., 1994; Okazaki et al., 1996; Lio et al., 1998).

[0019] Th1 cytokines and other Th1 markers were found to be associated with alcoholic hepatitis and liver cirrhosis. Inflammatory stimuli and lipid peroxidation activate nuclear factor κ B (NF- κ B) and upregulate proinflammatory cytokines and chemokines. In one study, the relationship between pathological liver injury, endotoxemia, lipid peroxidation, and NF- κ B activation and imbalance between pro- and anti-inflammatory cytokines was evaluated. Rats (5 per group) were fed ethanol and a diet containing saturated fat, palm oil, corn oil, or fish oil by intragastric infusion. Dextrose isocalorically replaced ethanol in control rats. Pathological analysis was performed and measurements of endotoxin were taken, lipid peroxidation, NF- κ B, and messenger RNA (mRNA) levels of proinflammatory cytokines (TNF α , IL-1 beta, IFN- γ , and IL-12), C—C chemokines (regulated upon activation, normal T cell expressed and secreted [RANTES], monocyte chemoattractant protein [MCP]-1, macrophage inflammatory protein [MIP]-1- α), C—X—C chemokines (cytokine induced neutrophil chemoattractant [CINC], MIP-2, IP-10, and epithelial neutrophil activating protein [ENA]-78), and anti-inflammatory cytokines (IL-10, IL-4, and IL-13). Activation of NF- κ B and increased expression of proinflammatory cytokines C—C and C—X—C chemokines was seen in the rats exhibiting necroinflammatory injury (fish oil-ethanol and corn oil-ethanol). These groups also had the highest levels of endotoxin and lipid peroxidation. Levels of IL-10 and IL-4 mRNA were lower in the group exhibiting inflammatory liver injury. Thus, activation of NF- κ B occurs in the presence of proinflammatory stimuli and results in increased expression of Th1 proinflammatory cytokines and chemokines (Naji et al., 1999). FAS ligand and its receptor are also elevated in alcoholic liver diseases, suggesting once again that Th1 cytokines are involved in the autoimmune processes induced in alcoholic hepatitis (Galle et al., 1995; Taieb et al., 1998; Fiore et al., 1999).

[0020] TNF- α has also emerged as a common pathway in the pathogenesis of alcohol-related hepatic necro-inflammation. Increased levels of hepatic and serum TNF have been documented in animal models of alcoholic liver disease and in human alcoholic liver disease. This dysregulated TNF metabolism has been postulated to play a role in many of the metabolic complications and the liver injury of alcoholic liver disease (Grove et al., 1997; McClain and Cohen, 1989). For instance it was found in one study that patients with alcoholic hepatitis had higher TNF- α levels (mean, 26.3 ng/L; 95% CI,

21.7 to 30.9) than normal subjects (6.4 ng/L; CI, 5.4 to 7.4). Patients who subsequently died had a higher TNF- α level (34.7 ng/L; CI, 27.8 to 41.6) than survivors (16.6 ng/L; CI, 14.0 to 19.2). In patients with alcoholic hepatitis, TNF- α levels correlated positively with serum bilirubin ($r=0.74$; $P=0.0009$) and serum creatinine ($r=0.81$; $P=0.0003$). Patients with alcoholic hepatitis had higher TNF- α levels than patients with inactive alcoholic cirrhosis (11.1 ng/L; CI, 8.9 to 13.3) and severely alcoholic persons without liver disease (6.4 ng/L; CI, 5.0 to 7.8). Patients with abnormal renal function had lower TNF- α levels (14.1 ng/L; CI, 5.4 to 22.8) than patients with alcoholic hepatitis. It was therefore concluded that elevations in TNF- α in alcoholic hepatitis are most marked in severe cases, suggesting that TNF- α plays a role in the pathogenesis (Bird et al., 1990).

[0021] TNF mediates many of the biologic actions of endotoxin. Recent studies have shown that TNF administration may cause liver injury and that TNF may mediate the lethality of the hepatotoxin galactosamine. One of the most potent TNF inducers is endotoxin. Because patients with alcoholic liver disease frequently have endotoxemia and because many of the clinical manifestations of alcoholic hepatitis are known biologic actions of TNF, its activity was evaluated in patients with alcoholic hepatitis. Basal and lipopolysaccharide-stimulated TNF release from peripheral blood monocytes, a major source of TNF production, was determined in 16 patients with alcoholic hepatitis and 16 healthy volunteers. Eight of 16 alcoholic hepatitis patients and only two of 16 healthy volunteers had detectable spontaneous TNF activity (p less than 0.05). After lipopolysaccharide stimulation, mean monocyte TNF release from alcoholic hepatitis patients was significantly increased to over twice that of healthy controls (25.3 ± 3.7 vs. 10.9 ± 2.4 units per ml, p less than 0.005). It was therefore concluded that monocytes from alcoholic hepatitis patients have significantly increased spontaneous and lipopolysaccharide-stimulated TNF release compared to monocytes from healthy volunteers (McClain and Cohen, 1989).

[0022] Lipopolysaccharide (LPS)-binding protein (LBP) and CD14 play key intermediary roles in the activation of cells by endotoxin. Gut-derived LPS has been postulated to participate in promoting pathological liver injury in alcoholic liver disease. It was demonstrated that rats fed intragastrically with ethanol in oil for 4 weeks had elevated levels of CD14 and LBP in their Kupffer cells and hepatocytes, respectively. Expression of CD14 mRNA was also elevated in nonmyeloid cells. Enhanced LBP and CD14 expression rapidly increases the LPS-induced expression of various pro-inflammatory cytokines and correlates with the presence of pathological liver injury in alcoholic liver injury (Su et al., 1998; Lukkari et al., 1999).

[0023] IL-6 is a 26 kd cytokine that plays a major role in the acute phase response, especially the hepatic aspects of the acute phase response. Patients with alcoholic hepatitis manifest many aspects of the acute phase response. Serial plasma IL-6 levels in 30 consecutive patients with moderate to severe alcoholic hepatitis was measured during 6 month. Mean admission plasma IL-6 activity was markedly increased (49.8 ± 8.5 U/ml, normal less than 5 U/ml) in patients with alcoholic hepatitis, and levels decreased with clinical improvement to 15.6 ± 6.1 U/ml at 6 months. Admission IL-6 activity correlated significantly ($r=0.82$) with the severity of liver disease as assessed by the discriminant function of Maddrey. IL-6 activity fell over time in a pattern similar to

that of bilirubin and C-reactive protein. This and additional studies suggest that plasma IL-6 is probably a marker of inflammation and severity of disease in alcoholic hepatitis (Sheron et al., 1991; Hill et al., 1992).

[0024] IL-8, a cytokine produced by a number of cells, including monocytes, macrophages, Kupffer cells and hepatocytes, can activate neutrophils. Peripheral neutrophilia and liver neutrophil infiltration are frequently noted in patients with alcoholic liver disease. It was found that serum IL-8 levels were markedly elevated in patients with alcoholic hepatitis (437+/-51 pg/ml) when compared to all other groups ($p<0.05$). Levels of IL-8 in patients with alcoholic fatty liver, alcoholic cirrhosis and viral hepatitis were higher than those in controls and in patients with non-alcoholic fatty liver. In addition, IL-8 levels were higher in patients who died ($p=0.007$), and correlated with biochemical and histological parameters, and severity of liver injury: serum aspartate aminotransferase, alanine aminotransferase, total bilirubin, prothrombin time, indocyanine green retention ratio, TNF- α and peripheral neutrophil count in patients with alcoholic hepatitis. After a 2-year follow up, patients with IL-8 above 479 pg/ml had a higher mortality rate in the alcoholic hepatitis group ($p=0.033$). These findings suggest that IL-8 as well as some other chemokines are activated in alcoholic liver disease, especially in alcoholic hepatitis, and is closely correlated with liver injury (Martinez et al., 1992; Hill et al., 1992).

[0025] Induction of adhesion molecules such as ICAM-1 is associated with the activation and attraction of a special population of inflammatory cells characteristic for alcoholic hepatitis. Frozen liver samples from patients who died with signs of acute alcoholic hepatitis show elevated ICAM-1 expression in the membranes of hepatocytes, as well as the occurrence of CD11b positive polymorphonuclear leukocytes (neutrophils) suggesting a possible major role of the beta 2-integrin Mac-1 as a ligand for ICAM-1. It was concluded that in alcoholic hepatitis cytokines may be responsible for the induction of the adhesion molecule ICAM-1 on hepatocytic membranes and activate a defined population of inflammatory cells, thus contributing to the characteristic histological picture of acute alcoholic hepatitis with its concentration of neutrophils especially in areas with ballooned Mallory body-containing hepatocytes (Afford et al., 1998).

[0026] A significant increase in both NK cells (CD3-/CD56+) and the cytotoxic T cells coexpressing the CD3 and the CD56 molecules together with an increased NK cytotoxic activity were observed in patients having alcoholic hepatitis. Interestingly these abnormalities persisted during the withdrawal period (Ohlinger et al., 1993).

[0027] Arthritis is a disease involving joint inflammation. The joints show swelling, stiffness, tenderness, redness or warmth. The symptoms may be accompanied by weight loss, fever or weakness. When these symptoms last for more than two weeks, inflammatory arthritis e.g. rheumatoid arthritis may be the cause. Joint inflammation may also be caused by infection, which can lead to septic arthritis. A very common type of arthritis is degenerative joint disease (osteoarthritis). Joint inflammation is not a prominent feature of osteoarthritis.

[0028] The medicaments commonly prescribed for arthritis and related conditions are non-steroidal anti-inflammatory drugs (NSAIDs). NSAIDs include aspirin and aspirin-like drugs. They reduce inflammation, which is the cause for joint pain, stiffness and swelling of the joints. However, NSAIDs are unspecific drugs having a number of side effects, involv-

ing bleeding of the stomach (Homepage of the Department of Orthopaedics of the University of Washington on Arthritis, Frederick Matsen (Chairman)). In addition to NSAIDs, Celebrex™, a cyclooxygenase (COX-2) inhibitor, is used to relieve the signs and symptoms of osteoarthritis and rheumatoid arthritis in adults. It is also indicated for the treatment of patients with familial adenomatous polyposis.

[0029] Further, TNF antagonists are used for the treatment of arthritis. TNF antagonists are described, for example, in WO 9103553.

[0030] Recent studies indicate that the interleukin IL-18 plays a proinflammatory role in joint metabolism. Olee et al. (1999) showed that IL-18 is produced by articular chondrocytes and induces proinflammatory and catabolic responses. The IL-18 mRNA was induced by IL-18 in chondrocytes. Chondrocytes produced the IL-18 precursor and in response to IL-1 stimulation secreted the mature form of IL-18. Studies on IL-18 effects on chondrocytes further showed that it inhibits TGF- β -induced proliferation and enhances nitric oxide production. IL-18 stimulated the expression of several genes in normal human articular chondrocytes including inducible nitric oxide synthase, inducible cyclooxygenase, IL-6, and stromelysin. Gene expression was associated with the synthesis of the corresponding proteins. Treatment of normal human articular cartilage with IL-18 increased the release of glycosaminoglycans. These findings identified IL-18 as a cytokine that regulates chondrocyte responses and contributes to cartilage degradation.

[0031] The localisation of Interleukin-1 β -converting enzyme (ICE)/caspase-1 in human osteoarthritic tissues and its role in the maturation of interleukin-1 beta and interleukin-18 have been shown by Saha et al. (1999). Saha et al. studied the expression and production of caspase-1 in human normal and osteoarthritic (OA) cartilage and synovium, quantitated the level of ICE in OA chondrocytes, and examined the relationship between the topographic distribution of ICE, interleukin-1 β (IL-1 β), and IL-18, as well as apoptosis of chondrocytes. The experiments performed in this study indicated that ICE was expressed and synthesised in both human synovial membrane and cartilage, with a significantly greater number of cells staining positive in OA tissue than in normal tissue. ICE production was preferentially located in the superficial and upper intermediate layers of articular cartilage. The production of mature IL-1 β in OA cartilage explants and chondrocytes was completely blocked by treatment with a specific ICE inhibitor, which also markedly diminished the number of IL-18-positive cells. The relationship between active IL-1 β and ICE suggests that ICE may promote OA progression by activating this proinflammatory cytokine, and that IL-18 may play a role in cartilage pathology.

[0032] Gracie et al. (1999) suggested a proinflammatory role for IL-18 in rheumatoid arthritis. Gracie et al. detected the IL-18 mRNA and protein within rheumatoid arthritis synovial tissues in significantly higher levels than in osteoarthritis controls. It was also shown that a combination of IL-12 or IL-15 with IL-18 induced the IFN- γ production by synovial tissues in vitro. Furthermore, IL-18 administration of collagen/incomplete Freund's adjuvant-immunized mice facilitated the development of an erosive, inflammatory arthritis, suggesting that IL-18 may be proinflammatory in vivo.

[0033] However, so far, apart from chemical compounds, only the blockade of TNF α and IL-18 by using soluble receptors or monoclonal antibodies have been shown to decrease murine collagen-induced arthritis (CIA, which is a mouse

model for rheumatoid arthritis) (Williams et al., 1994), and were therefore suggested as a therapeutic for rheumatoid arthritis.

[0034] The term “systemic inflammatory response syndrome (SIRS)” describes the familiar clinical syndrome of sepsis, independent of its cause. SIRS can result from trauma, pancreatitis, drug reactions, autoimmune disease, and other disorders; when it arises in response to infection, sepsis is said to be present (Nathens and Marshall 1996).

[0035] Septic shock is the most common cause of death in medical and surgical intensive-care units (Aestiz and Rackow 1998). The terms sepsis, severe sepsis and septic shock are used to identify the continuum of the clinical response to infection. Patients with sepsis present evidences of infection and clinical manifestations of inflammation. Patients with severe sepsis develop hypoperfusion with organ dysfunction. Septic shock is manifested by hypoperfusion and persistent hypotension. Mortality ranges from 16% in patients with sepsis to 40-60% in patients with septic shock. Bacterial infection is the most common cause of septic shock. The most frequent sites of infection are the lungs, abdomen, and urinary tract. General anti-inflammatory therapies such as the use of corticosteroids, failed to show improvement in survival from sepsis and septic shock. Three monoclonal antibodies specific to endotoxin have been tested in clinical trials and have failed to improve survival rates. Therapy with antagonists to tumor necrosis factor, interleukin 1, bradykinin, ibuprofen, and platelet activating-factor, did not show improvement on survival from septic shock (Aestiz and Rackow 1998).

[0036] Sepsis is caused inter alia by Gram-positive bacteria e.g., *Staphylococcus epidermidis*. Lipoteichoic acids and peptidoglycans, which are the main cell wall components of the *Staphylococcus* species, are thought to be the inducers of cytokine release in this condition (Grupta et al. 1996 and Cleveland et al. 1996). However, other Gram-positive components are considered as stimulators of cytokine synthesis as well (Henderson et al. 1996). Production of IL-1, TNF- α and IFN- γ are thought to be the major contributors in the pathogenesis of septic shock (Dinarello 1996 and Okusawa et al. 1988). In addition, the chemokine IL-8 was shown to be induced by neutrophils in response to *S. epidermidis* (Hachicha et al. 1998). Important factors in the regulation of IL-1, TNF- α and IFN- γ induction by *S. epidermidis* are IL-18, IL-12, IL-1 β and TNF- α . IL-1 β and TNF- α can be considered as a co-stimuli for IFN- γ production by T lymphocytes in a manner similar to IL-18. These two cytokines have a co-stimulatory activity on IFN- γ production in the context of IL-12 or bacterial stimulation (Skeen et al. 1995 and Tripp et al. 1993).

[0037] Recently it was reported by Nakamura et al. (2000) that concomitant administration of IL-18 and IL-12 to mice results in high toxicity similar to that found in endotoxin-induced septic shock.

[0038] The levels of IL-18 have been shown to be elevated in sera from patients with sepsis (Endo et al. 2000) but this increase correlated with creatinine levels suggesting that the elevated levels of IL-18 may result from renal failure. In another study (Grobmyer et al. 2000), the levels of IL-18 in 9 subjects with sepsis during the first 96 hours following hospital admission were high and did not correlate with creatinine levels.

[0039] Dayer (1999) summarized the different and partially contradicting functions of IL-18. IL-18 is a pleiotropic interleukin having both inflammatory enhancing and attenuating

functions. On the one hand, it enhances production of the proinflammatory cytokines like TNF α , therefore promoting inflammation. On the other hand, it induces the production of NO, an inhibitor of caspase-1, thus blocking the maturation of IL-1 β and IL-18, and possibly attenuating inflammation. This ambiguous role of IL-18 seriously questioned the efficacy of IL-18 inhibitors in inflammatory diseases. Furthermore, because of the interaction of a huge variety of different cytokines and chemokines in the regulation of inflammation, it could not have been expected to obtain a beneficial effect by blocking only one of the players in such a complicated scenario.

[0040] Netea et al. (2000) reported that administration of anti-IL-18 polyclonal antibody protected mice against the deleterious effects of both LPS derived from *E. coli* or *S. typhimurium* species tested, supporting the concept that IL-18 has an important pathogenic role in lethal endotoxemia. However, the uncertainty regarding the efficiency of a potential treatment formulation based on administration of IL-18 inhibitors, is manifested in that IL-18 knock out mice are not less susceptible to sepsis as compared to wild type animals (Sakao et al. 1999) and in that recent reports suggest that the proinflammatory activity of IL-18 is essential to host defences against severe infections (Nakanishi et al. 2001 and Foss et al. 2001).

[0041] Thus, there exists a need to provide means to treat and/or prevent sepsis despite the above-mentioned considerations regarding the use of IL-18 inhibitors.

SUMMARY OF THE INVENTION

[0042] The present invention provides IL-18 binding proteins (IL-18BPs) and virally encoded IL-18BP homologues (hereinafter, viral IL-18BPs), and fused proteins, muteins, functional derivatives and active fragments thereof, capable of binding to IL-18. The invention also provides a process for obtaining IL-18BPs by isolating them from human fluids to obtain IL-18BP, or by recombinant means, which also allows for attainment of the various IL-18BPs. The invention also provides expression vectors of IL-18BPs, suitable for expression of IL-18BP in humans and other mammals. The IL-18BPs and the expression vectors of the present invention are useful for blocking the biological activities of IL-18.

[0043] Replicable expression vehicles containing DNAs suitable for expression of the various IL-18BPs in host cells, host cells transformed herewith and proteins and polypeptides produced by expression of such hosts are also provided.

[0044] The invention further provides antibodies to the IL-18BPs and the viral IL-18BPs, suitable for affinity purification and immunoassays of same.

[0045] The invention further provides pharmaceutical compositions consisting of suitable vehicles and IL-18BPs, or viral IL-18BPs, or vectors for expressing same in humans and other mammals, for the treatment of diseases or conditions which require modulation of IL-18 activity.

[0046] These diseases or conditions include autoimmune diseases, type I diabetes, rheumatoid arthritis, graft rejections, inflammatory bowel disease, sepsis, multiple sclerosis, ischemic heart diseases (including heart attacks), ischemic brain injury, chronic hepatitis, psoriasis, chronic pancreatitis, acute pancreatitis and the like.

[0047] It is a specific object of the present invention to provide for a novel means for treating and/or preventing liver injury. It has been found that an IL-18 inhibitor is effective in the prevention and treatment of liver damages. The invention

therefore also relates to the use of an IL-18 inhibitor for the manufacture of a medicament for treatment and/or prevention of liver injury. More specifically, the invention relates to the treatment and/or prevention of liver injuries caused by alcoholic hepatitis, viral hepatitis, immune hepatitis, fulminant hepatitis, liver cirrhosis, and primary biliary cirrhosis.

[0048] It has also been found in accordance with the present invention that an inhibitor of IL-18 is effective in the therapy of arthritis. The therapeutic effect includes decreasing the severity of the disease, as well as preventing the spreading of the disease. The invention therefore relates to the use of an inhibitor of IL-18 for treatment and/or prevention of arthritis. This finding is unexpected, since from the state of the art outlined above, it could not have been concluded that a blockade of one specific factor involved in arthritis, namely interleukin IL-18, would lead to the alleviation of arthritis or even the curing of a diseased arthritic joint.

[0049] It has also been found that the administration of an IL-18 inhibitor significantly diminishes cartilage erosion in a murine model of arthritis. The present invention thus also relates to the use of an inhibitor of IL-18 in the manufacture of a medicament for treatment and/or prevention of cartilage destruction.

[0050] In order to apply a gene therapeutic approach to deliver the IL-18 inhibitor to the diseased tissue or cell, it is a further object of the invention to use an expression vector comprising the coding sequence of an IL-18 inhibitor for the treatment and/or prevention of arthritis.

[0051] The invention relates to the use of an inhibitor of IL-18 in the manufacture of a medicament for the treatment and/or prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS) selected from severe sepsis and septic shock, and also for sepsis related cardiac dysfunction.

[0052] More specifically the invention relates to the use of an inhibitor of IL-18, selected from caspase-1 (ICE) inhibitors, antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signaling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, inhibitors of IL-18 production and IL-18 binding proteins, isoforms, muteins, fused proteins, functional derivatives, active fractions or circularly permuted derivatives thereof having at least essentially the same activity as an IL-18 binding protein.

[0053] The IL-18 binding protein used according to the invention is an isoform, a mutein, fused protein (e.g., Ig fused), functional derivative (e.g., PEG-conjugated), active fraction or circularly permuted derivative thereof.

[0054] The antibodies used according to the invention may be anti IL-18 specific antibodies selected from chimeric, humanized and human antibodies.

[0055] In addition, the invention relates to the use of an inhibitor in the manufacture of a medicament further comprising an IL-12 inhibitor, preferably an IL-12 neutralizing antibody, an interferon, preferably interferon .alpha. or .beta., a tumor necrosis factor inhibitor, preferably soluble TNFRI or TNFRII, and/or an IL-1 inhibitor, preferably IL-1 receptor antagonist for simultaneous, sequential or separate administration.

[0056] In one aspect, the invention provides the use of an expression vector comprising the coding sequence of an inhibitor of IL-18 selected from caspase-1 (ICE) inhibitors, antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signaling pathway,

antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, inhibitors of IL-18 production and IL-18 binding proteins, isoforms, muteins, fused proteins, or circularly permuted derivatives thereof having at least essentially the same activity as an IL-18 binding protein, in the manufacture of a medicament for the treatment and/or prevention of sepsis, for example by gene therapy.

[0057] In another aspect, the invention provides the use of a vector for inducing and/or enhancing the endogenous production of an inhibitor of IL-18 in a cell, in the manufacture of a medicament for the treatment and/or prevention of sepsis.

[0058] In addition, the invention provides the use of a cell that has been genetically modified to produce an inhibitor of IL-18 in the manufacture of a medicament for the treatment and/or prevention of sepsis.

[0059] In another embodiment, the invention relates to a method for the treatment and/or prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS), including sepsis related cardiac dysfunction, comprising administering to a subject in need thereof a pharmaceutically effective amount of inhibitor of IL-18 selected from caspase-1 (ICE) inhibitors, antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signaling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, inhibitors of IL-18 production, and IL-18 binding proteins, isoforms, muteins, fused proteins, functional derivatives, active fractions or circularly permuted derivatives thereof having essentially the same activity as an IL-18 binding protein. The method may comprise co-administration of a therapeutically effective amount an inhibitor of cytokines selected from IL-12 inhibitor, preferably neutralizing antibodies, Tumor Necrosis Factor inhibitors, preferably soluble portion of TNFRI or TNFRII, IL-1 inhibitors, preferably the IL-1 receptor antagonists and IL-8 inhibitors, and/or an interferon, preferably interferon-.alpha. or -.beta.

[0060] In addition, the invention provides a method for the treatment and/or prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS) comprising administering to a subject in need thereof a pharmaceutically effective amount of vector coding the sequence of an inhibitor of IL-18 selected from caspase-1 (ICE) inhibitors, antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signaling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, inhibitors of IL-18 production, and IL-18 binding proteins, isoforms, muteins, fused proteins or circularly permuted derivatives thereof.

[0061] In a further embodiment, the invention relates to a method for the treatment and/or prevention of sepsis and other diseases characteristic to the SIRS comprising administering to a subject in need thereof a pharmaceutically effective amount of a vector for inducing and/or enhancing the endogenous production of an inhibitor of IL-18 in a cell.

[0062] In addition, the invention provides a method for the treatment and/or prevention of sepsis and other diseases characteristic to the SIRS comprising administration to a subject in need thereof a cell that has been genetically modified to produce an inhibitor of IL-18.

[0063] The invention also relates to the use of an inhibitor of IL-18 in the manufacture of a medicament for the treatment and/or prevention of sepsis and other diseases characteristic

to the Systemic Inflammatory Response Syndrome (SIRS) selected from severe sepsis, septic shock, and also for sepsis related to cardiac dysfunction.

[0064] The invention includes a method of treating or preventing sepsis associated with an excess of IL-18, where the method comprises administering to a subject in which such treatment or prevention is desired, a composition comprising (a) a polypeptide comprising an amino acid sequence selected from the group consisting of AA1-AA192 of SEQ ID NO:2, or AA29-AA192 of SEQ ID NO:2, (b) a nucleic acid molecule comprising a nucleic acid sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of AA1-AA192 of SEQ ID NO:2, or AA29-AA192 of SEQ ID NO:2, or (c) a mutein of any one of the sequences in (a) or (b), characterized in that the mutein i) has at least 90% identity to at least one of the sequences in (a); ii) comprises the amino acid sequence of SEQ ID NO:10; and iii) binds IL-18 in an amount sufficient to treat or prevent the disease in the subject.

[0065] In one embodiment, the method of treating or preventing sepsis includes administration to a human subject. In another embodiment, the polypeptide is glycosylated at one or more sites. Alternatively, the polypeptide is not glycosylated. In another embodiment, the polypeptide further comprises at least one moiety attached to one or more functional groups which occur as one or more side chains on the amino acid residues or the N- or C-terminal groups.

[0066] In another aspect of the method of treating sepsis, the polypeptide is circularly permuted. In another aspect, the polypeptide is a non-viral protein. In another the polypeptide is a human protein. In another aspect, the polypeptide is a fused protein. In the polypeptide comprises an Ig fusion. In another aspect, the polypeptide is soluble. In another aspect, the polypeptide is pegylated. In another aspect, at least one moiety is a polyethylene glycol moiety.

BRIEF DESCRIPTION OF THE DRAWINGS

[0067] FIG. 1 shows SDS-PAGE (sodium dodecyl sulfate polyacrylamide gel electrophoresis) of ligand affinity purified IL-18 binding protein. Crude urinary proteins (concentrated by ultrafiltration of 500 L normal human urine) were loaded on an IL-18-agarose column. The column was washed and bound proteins eluted at pH 2.2. Eluted fractions were neutralized and aliquots were analyzed by SDS-PAGE (10% acrylamide) under non-reducing conditions and silver staining. The lanes are: 1: crude urinary proteins (1.5 µg, loaded on the gel); 2-9: elutions 1-8, respectively, from the IL-18-agarose column; 10: molecular weight markers, in kD, as indicated on the right side. An arrow indicates the band corresponding to IL-18BP.

[0068] FIG. 2 shows an autoradiogram of SDS-PAGE (7.5% acrylamide) of complexes consisting of ¹²⁵I-IL-18 (apparent molecular weight 19 kD), cross-linked to the following preparations of soluble IL-18 binding protein: Lane 1: Wash of the IL-18 affinity column. Lane 2: Elution 2 of the IL-18 affinity column. Lane 3: Elution 3 of the IL-18 affinity column. Molecular weight markers are indicated on the right side (in kD). An arrow indicates the cross-linked product (58 kD).

[0069] FIG. 3 shows inhibition of IL-18-induced production of IFN-γ by IL-18BP

[0070] (A) Mouse splenocytes were stimulated (24 hr, 37° C.) with the indicated combinations of LPS (1 µg/ml) and human IL-18 (5 ng/ml), added either directly,

or after pre-mixing (1 h, 37° C.) with urinary IL-18BP. The level of mulFN-γ in the culture was determined after 24 hr.

[0071] (B) Mouse splenocytes were incubated (24 h) with LPS (1 µg/ml) together with murine IL-18 (10 ng/ml) pre-mixed (1 h, 37° C.) with increasing concentrations of human IL-18BP.

[0072] (C) Mouse splenocytes were incubated (24 h) with LPS (10 µg/ml) together with increasing concentrations of human IL-18BP.

[0073] (D) Mouse splenocytes were incubated (24 h) with Con A (1 µg/ml), together with increasing concentrations of human IL-18BP.

[0074] (E) Human peripheral blood mononuclear cells (PBMC) of three donors were stimulated with IL-12 (10 ng/ml) and huIL-18 (25 ng/ml), added either alone, or after pre-mixing (1 h, 37° C.) with urinary IL-18BP. Human KG-1 cells were stimulated with TNF-α (20 ng/ml) and huIL-18 (25 ng/ml), added either alone, or after pre-mixing (1 h, 37° C.) with urinary IL-18BP.

[0075] FIG. 4 shows the effect of recombinant IL-18BP on human and mouse IL-18 activity. His₆-tagged IL-18BP was transiently expressed in COS7 cells and purified.

[0076] (A) Human IL-18 (5 ng/ml) was pre-mixed with either His₆-tagged-IL-18BP or RPMI and added to mouse spleen cells together with LPS (1 µg/ml). IFN-γ production was measured after 24 h.

[0077] (B) Mouse IL-18 (10 ng/ml) was pre-mixed with either His₆-tagged-IL-18BP or RPMI and added to mouse spleen cells together with LPS (1 µg/ml). IFN-γ production was measured after 24 h.

[0078] (C) Human IL-18 (25 ng/ml) was pre-mixed with either COS7-IL-18BP or RPMI and added to Human PBMC in the presence of IL-12 (10 ng/ml).

[0079] (D) Human IL-18 (25 ng/ml) was pre-mixed with either COS7-IL-18BP or RPMI and added to Human KG-1 cells in the presence of TNF-α (20 ng/ml).

[0080] FIG. 5 shows a histogram depicting the serum levels of IFN-γ (pg/ml) after injection of various amounts of recombinant IL18BP (0; 0.04; 0.4; 4 mg/kg) into mice 1 h before the injection of LPS. Blood samples were taken 5 h after LPS injection and analyzed by ELISA for circulating IFN-γ.

[0081] FIG. 6 shows a histogram depicting the serum levels of Alanine aminotransferase. Mice were injected with increasing doses of recombinant human IL18BP (0; 0.04; 0.4; 4 mg/kg) before injection of LPS into *P. acnes* sensitized mice. Blood samples were taken 5 h after LPS injection and serum levels of Alanine aminotransferase (ALT) were measured. SF=Sigma-Frankel: 1 SF Unit of AST/ALT will form 4.82×10^{-4} µmol glutamate/minute at pH 7.5 at 25° C.

[0082] FIG. 7 shows the survival time of the mice after LPS injection. Mice were injected with different doses of recombinant human IL18BP (0; 0.04; 0.004; 4 mg/kg) 20 min before injection of LPS into *P. acnes* sensitized mice. Triangles: 4 mg/kg; small diamond: 0.4; big diamond: 0.04; circles: no IL18BP (only LPS).

[0083] FIG. 8 shows a histogram depicting serum levels of IFN-γ, measured 5 h after injection of different amounts of IL18BP (0; 0.4; 4 mg/kg), which was administered 20 min before LPS injection into *P. acnes* sensitized mice.

[0084] FIG. 9 shows the survival of mice injected either with polyclonal IL-18 antiserum or normal rabbit serum (NDS=control) 30 min before injection with 40 mg/ml (lethal dosis) of LPS derived from *E. coli* (FIG. 9 A) or *S. typhimur-*

rium (FIG. 9 B). Triangles: mice were injected with IL-18 antiserum; circles: mice were injected with NDS. On the x-axis, the days after LPS challenge are depicted. * $p < 0.05$.

[0085] FIG. 10 shows a histogram, in which the mean $\pm$ SEM of five mice per group treated in the following way are depicted. Mice were injected intraperitoneally (i.p.) either anti-IL-18 antiserum, soluble TNF- α receptors (TNF-sRp55) or vehicle (saline), immediately followed by the intravenous (i.v.) administration of Concanavalin A (Con A; FIG. 10 A) or PEA (*Pseudomonas aeruginosa*, FIG. 10 B). ** $p < 0.01$; *** $p < 0.001$ vs. ConA or PEA alone; # $p < 0.01$ vs. either TNFsRp55 or anti-IL-18 factorial ANOVA.

[0086] FIG. 11 shows the effect of IL-18BP on clinical scores in a murine model of arthritis

[0087] (A) shows a diagram depicting the clinical scores measured after daily administration of different amounts of IL-18BP or IFN- β or vehicle (NaCl) i.p. (intraperitoneally) to mice. Symbols: Filled triangles: 10 000 IU IFN- β ; open triangles: 10 mg/kg IL-18BP, reversed triangles: 3 mg/kg IL-18BP, diamonds: 1 mg/kg IL-18BP; circles: 0.5 mg/kg IL-18BP; open squares: 0.25 mg/kg IL-18BP, and filled squares: NaCl. The days of treatment are depicted on the x-axis, the clinical scores (mean values) are depicted on the y-axis. Statistics were calculated by the Mann Whitney test.

[0088] (B) shows a histogram depicting the AUC (area under the curve) derived from the graph of FIG. 11A. n=number of animals.

[0089] FIG. 12 shows the effect of IL-18BP on paw swelling

[0090] (A) shows a diagram depicting the results obtained by measuring the paw thickness (swelling) of diseased hind paws of individual animals treated with different amounts of IL-18BP. The y-axis shows the change of paw thickness in millimeters from the beginning of treatment. The symbols are as in FIG. 11.

[0091] (B) shows a histogram depicting the AUC derived from FIG. 11A. n=number of animals.

[0092] FIG. 13 shows the analysis of the number of diseased hind paws at the time of acute arthritis, i.e. spreading of the disease to additional joints. Symbols: Filled squares: NaCl (control), triangles: 10 mg/kg IL-18BP, reversed triangles: 3 mg/kg IL-18BP, diamonds: 1 mg/kg IL-18BP, circles: 0.5 mg/kg IL-18BP and open squares: 0.25 mg/kg IL-18BP.

[0093] FIG. 14 shows a histogram depicting the erosion scores of the cartilage of diseased joints.

[0094] FIG. 15 shows the histopathology of mouse joints. At the end of the experiment, the paw that first developed arthritis was dissected away, fixed and processed as described in Example 2 below. A) normal mouse joint; B) joint from an arthritic mouse; C) joint from a mouse treated with rhIL-18BP.

[0095] FIG. 16 shows a histogram depicting the levels of anti-collagen type II antibodies of the isotype IgG1 (open columns) or IgG2a (hatched columns) of mice treated with 3 mg/kg of IL-18BP or saline (vehicle), respectively. Measurements were taken on day 4 (D4) or day 8 (D8) of the disease.

[0096] FIG. 17 shows a histogram depicting IL-6 levels in pg/ml of animals treated with 1, 3 or 10 mg/kg of IL-18BP, 10 000 IU of Interferon β (IFN- β), normal mouse serum (NMS) or saline (NaCl), respectively.

[0097] FIG. 18 shows the distribution of serum IL-18BP in healthy individuals. Sera of male and female healthy indi-

viduals ages 20-60, as assayed for the presence of IL-18BP by a specific ELISA test (n=107).

[0098] FIG. 19A shows the average levels of IL-18 and IL-18BP in sepsis patients and healthy subjects. Serum IL-18 and IL-18BP were determined in the healthy individuals (see FIG. 18) and in 198 samples from 42 sepsis patients immediately upon hospital admission and during hospitalisation.

[0099] FIG. 19 B shows the distribution of the individual levels of IL-18 and IL-18BP of the healthy subjects and patients described in FIG. 19A. The mean serum levels of IL-18 and IL-18BP in healthy subjects are indicated by a dashed vertical and horizontal line, respectively.

[0100] FIG. 20 shows a comparison between the total and free IL-18 in individual sepsis patients upon hospital admission. The level of free IL-18 (closed circles) in sera was calculated based on the concentration of total IL-18 (open circles) and IL-18BP, taking into account a stoichiometry of 1:1 of IL-18 to IL-18BP in a complex and a calculated K_d of 400 μ M. Each vertical line links total and free IL-18 in an individual serum sample.

[0101] FIG. 21 shows the effect of IL-18BP on *S. epidermidis*-induced IFN- γ production in whole blood. Whole blood was stimulated with *S. epidermidis* alone or *S. epidermidis* and recombinant IL-18BP at indicated concentrations. After 48 hours of incubation, the blood culture was lysed and IFN- γ was measured. The results are expressed as percentage of *S. epidermidis* IFN- γ induction. $P < 0.01$ vs. *S. epidermidis* alone. The data represents the means $\pm$ standard error of the mean (SEM) of six experiments. SEM represents the spread of the mean of a sample. SEM gives an idea of the accuracy of the mean. $SEM = SD / (\text{square root of sample size})$. The data was analysed with paired Student's test.

[0102] FIG. 22 shows the inhibitory effect exerted by the double activity of IL-18BP and anti IL-12 monoclonal antibodies on the production of IFN- γ by whole blood treated with *S. epidermidis*. Whole blood was stimulated with *S. epidermidis* in the presence or in the absence of IL-18BP (125 ng/ml), IL-12 Mab (2.5 μ g/ml) and both. After 48 hours of incubation, the blood culture was lysed and IFN- γ was measured. The data is expressed as percent of IFN- γ induction. $P < 0.01$ vs. *S. epidermidis* alone. The data represents the mean $\pm$ standard error of the mean (SEM) of six experiments. SEM represents the spread of the mean of a sample. SEM gives an idea of the accuracy of the mean. $SEM = SD / (\text{square root of sample size})$. The data was analysed with paired Student's test.

[0103] FIG. 23 shows the quantitation of IL-18BP as reflected in an ELISA standard curve. Recombinant human IL-18BP was serially diluted and subjected to ELISA as described in Example 30. The data shows the mean absorbance $\pm$ SE (standard error) of 10 experiments.

[0104] FIG. 24 shows the inhibitory effect of IL-18 on the quantitation of IL-18BP by ELISA assay. The percent of inhibition of the signal is shown.

[0105] FIG. 25 shows the immunological cross reactivity of human IL-18BP isoforms. The IL-18PELISA was performed with all isoforms (a-d) of IL-18BP. Stocks (3.2 μ g/ml) of the IL-18BP isoforms were serially diluted and tested in IL-18BP ELISA.

[0106] FIG. 26 shows myocardial tissue content of IL-18 following LPS administration. Mice were injected with *E. coli* LPS (0.5 mg/Kg, intraperitoneally). Hearts were homog-

enized at the indicated times at the x axis. ELISA was carried out to determine myocardial IL-18 content. (n=4-5 per group).

[0107] FIG. 27 shows the effect of anti IL-18 antibody on LPS-induced myocardial dysfunction. Mice were injected with either vehicle (intraperitoneally injected saline n=8) or *E. coli* LPS (0.5 mg/Kg injected intraperitoneally, n=8) and the left ventricular developed pressure (LVDP) was determined at 6 hours by isolated heart perfusion. In separate experiments, mice were pretreated with either normal rabbit serum (NRS, n=5) or anti-IL-18 antibody (Anti IL-18, n=8) 30 minutes prior to LPS.

DETAILED DESCRIPTION OF THE INVENTION

[0108] The present invention relates to various IL-18BPs and viral IL-18BPs which bind to IL-18 and thus, are capable of blocking the biological activities of IL-18. The term, "IL-18BPs and viral IL-18BPs," includes muteins of IL-18BPs and viral IL-18BPs, derivatives of IL-18BPs and viral IL-18BPs and truncated forms of IL-18BPs and viral IL-18BPs and salts thereof. The invention further relates to replicable expression vehicles, suitable for expression of various IL-18BPs or viral IL-18BPs in host cells and host bacteria. The invention further relates to expression vectors, suitable for expression of various IL-18BPs or viral IL-18BPs in humans and in other mammals.

[0109] The replicable expression vehicles containing DNAs suitable for expression of various IL-18BPs and viral IL-18BPs in host cells, according to the invention, may be PCR products, cDNA, synthetic DNA and combinations thereof. DNA molecules hybridizing to the above DNAs under stringent conditions and encoding proteins or polypeptides having the same activity as IL-18BP are also included in the present invention.

[0110] The expression vectors, suitable for expression of various IL-18BPs or viral IL-18BPs in humans and in other mammals, according to the present invention, may be viral vectors or other types of vectors to which an IL-18BP gene or an IL-18BP cDNA or a DNA encoding a viral IL-18BP was inserted in a way that enables efficient expression of an IL-18BP or a viral IL-18BP in humans and other mammals. DNA molecules hybridizing to the above DNAs under stringent conditions and encoding proteins or polypeptides having the same activity as IL-18BP, are also included in the present invention.

[0111] Isolation of IL-18BP may be accomplished in accordance with the invention, e.g. by passing a human fluid, such as urine or serum, through a chromatographic column to which IL-18 is coupled, and thereafter, eluting the bound IL-18BP.

[0112] The various IL-18BPs and viral IL-18BPs can also be prepared by recombinant means, i.e. by expressing IL-18BP in a suitable host, after adding promoters, expression enhancers, etc., suitable for the particular host employed.

[0113] The various IL-18BPs and viral IL-18BPs and vectors for expressing IL-18 Bp in humans and other mammals may be employed in the treatment and alleviation of conditions in which IL-18 is involved or caused by an excess of exogenously administered or endogenously produced IL-18. Such conditions are, e.g., autoimmune diseases, type I diabetes, rheumatoid arthritis, graft rejections, inflammatory bowel disease, sepsis, multiple sclerosis, ischemic heart dis-

eases (including heart attacks), ischemic brain injury, chronic hepatitis, psoriasis, chronic pancreatitis, acute pancreatitis and the like.

[0114] According to the present invention, IL-18BP was isolated from normal human urine by one chromatographic step. A preparation of crude human urinary proteins was loaded on a column consisting of human IL-18 bound to agarose. The column was washed and bound proteins were eluted at low pH. Eluted fractions were neutralized and aliquots were analyzed by SDS-PAGE (10% acrylamide) under non-reducing conditions and silver staining. A protein band of ~40 kD was specifically obtained in the eluted fractions (FIG. 1).

[0115] The ~40 kD protein obtained in the first step was identified as an IL-18 binding protein by its ability to specifically cross-link with ¹²⁵I-IL-18 (FIG. 2). The ~40 kD protein was further characterized by N-terminal protein sequence analysis. Aliquots from the eluted protein were subjected to SDS-PAGE, electroblotted to a PVDF membrane and subjected to protein microsequence analysis. Similarly, aliquots from the eluted protein were subjected to direct protein microsequence analysis. In both cases, two polypeptide sequences were obtained. A major sequence and a minor sequence, the latter corresponding to a fragment of human defensin (accession number p11398), starting at amino acid 65. Subtraction of the known defensin sequence provided the following new sequence:

T-P-V-S-Q-Q-x-x-x-A-A-A- (SEQ ID NO: 11)
1 . . . 5 . . . 10 . .

[0116] wherein x represents a yet undetermined amino acid.

[0117] In order to obtain a longer and more accurate sequence and in order to identify potential cysteine residues, an aliquot of the eluted fraction was reduced with DTT under denaturing conditions, reacted with 4-vinylpyridine, desalted by a micro-ultrafiltration device (Ultrafree, cutoff 10,000 Da, Millipore) and subjected to protein microsequence analysis. After sequencing cycle No. 1 the residual protein was reacted with o-phthalaldehyde to block all N-terminal polypeptides other than Pro and sequencing was then resumed. In this way the following single protein sequence was obtained:

(SEQ ID NO: 10)
TPVSQXXXAA XASVRSTKDP CPSQPPVFPA AKQCPALEVT
1 10 20 30 40
(T = Thr; P = Pro; V = Val; S = Ser; Q = Gln; X =
Unknown; A = Ala; R = Arg; K = Lys; D = Asp; C =
Cys; F = Phe; L = Leu; E = Glu)

[0118] The resulting sequence is significantly different from that of any other known protein, as determined by searching protein databases. However, searching the database of The Institute of Genomic Research (TIGR) by the tblastn search program provided a cDNA file, denoted THC123801, whose open reading frame (218 codons) contains a sequence highly homologous to that of the N-terminal sequence of IL-18BP. The homology is hereby shown:

```

1  . . . . . TPVSQXXXAAXASVRSTKDPSPVFPAAKQCPALEVT . . . 40 (SEQ ID NO: 12)
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
51 VTLVLRATXVXQTTTAATASVRSTKDPSPVFPAAKQCPALEVTWPE 100 (SEQ ID NO: 13)

```

[0119] (The upper sequence (1-40) (SEQ ID NO: 12) is that of the isolated IL-18BP; the lower sequence (51-100) (SEQ ID NO: 13) is derived by translation of TIGR file THC123801).

[0120] The affinity-purified urinary IL-18BP retained the ability to bind its labeled ligand (^{125} I-IL-18), and following covalent cross-linking, a complex of molecular weight 58 kD was formed. The molecular weight of this complex corresponded to a 1:1 ratio of the ~40 kD IL-18BP and the 19 kD IL-18 (FIG. 2).

[0121] The affinity-purified urinary IL-18BP blocked the biological activity of human as well as mouse IL-18. Thus when IL-18BP was added to either human or mouse IL-18 it blocked the ability of IL-18 to induce the production of interferon- γ when added together with lipopolysaccharide to cultures of mouse spleen cells (FIG. 3).

[0122] Using the partial sequence of the IL-18BP cDNA as provided by the TIGR database, a DNA probe was prepared by reverse-transcription PCR with specific sense and anti-sense primers and RNA from the human Jurkat T cells. The resulting PCR product was confirmed by DNA sequence analysis. This PCR product was labeled with ^{32} [P] and used as a probe for screening of four human cDNA libraries, derived from peripheral blood monocytes, from the Jurkat T-cell line, from PBMC and from human spleen. The various independent cDNA clones corresponded to four IL-18BP splice variants. All splice variants coded for putative soluble secreted proteins. The most abundant one (IL-18BP_a) had an open reading frame of 192 codons, coding for a signal peptide of 28 amino acid residues followed by a mature putative IL-18BP_a, whose first 40 residues matched perfectly with the N-terminal protein sequence of the urinary IL-18BP (SEQ ID NO:1 and SEQ ID NO:2). The position of the cysteine residues suggested that this polypeptide belongs to the immunoglobulin (Ig) super-family. Interestingly, each of the four Gln residues within mature IL-18BP_a was a potential N-glycosylation site. The three other variants of IL-18BP were less abundant than IL-18BP_a. They included a shorter 1 kb IL-18BP_b cDNA, coding for a signal peptide of 28 amino acid residues followed by a mature protein of 85 amino acid residues (SEQ ID NO:3 and SEQ ID NO:4). A third variant, IL-18BP_c was represented by a 2.3 kb cDNA, coding for a signal peptide of 28 amino acid residues followed by a mature IL-18BP of 169 amino acid residues (SEQ ID NO:5 and SEQ ID NO:6). The fourth variant, IL-18BP_d, coded for a signal peptide of 28 amino acid residues followed by a mature IL-18BP of 133 amino acid residues (SEQ ID NO:7 and SEQ ID NO:8).

[0123] To further study the possible existence of additional IL-18BP splice variants, a human genomic library was screened with a probe corresponding to full length IL-18BP cDNA. Five genomic clones, differing in length, were identified in this library. These clones were subjected to DNA sequence analysis with external and internal primers. Altogether, a 7.1 kb contig was assembled from these clones (SEQ ID NO:9). No exon coding for a trans-membrane (TM) receptor was identified within the 7.1 kb contig. All variants shared a common translation start site, coded for the same signal

peptide of 28 amino acid residues and soluble mature proteins of varying sizes and C-terminal sequences. The IL-18BP locus contains an additional gene, coding for the nuclear mitotic apparatus protein 1 (NUMA1), positioned at the minus strand. This finding localizes the IL-18BP gene to human chromosome 11q13 (Broach et al., 1981).

[0124] An homology search was done with the complete protein sequence of IL-18BP_a and the GenPept database, using the Smith Watermann algorithm. It was found that homologues of IL-18BP are expressed in several Poxviruses as secreted proteins of a previously unknown function. It was previously reported that viruses code for various cytokine receptors and that such virally encoded molecules serve as decoy receptors that inhibit immune responses by neutralizing their corresponding cytokine (reviewed by Spriggs, M K, 1994, Curr. Opin. Immunol., 6, 526-529). Therefore the invention further relates to virally encoded homologues of IL-18BP that may also serve as blockers of the biological activity of IL-18. Examples of virus-encoded homologues of IL-18BP are provided in Table 1.

TABLE 1

Virus-encoded proteins, showing high homology to human IL-18BP	
GenPept sequence	Virus type
MCU60315_54	U60315 Molluscum contagiosum virus subtype 1
MCU60315_53	U60315 Molluscum contagiosum virus subtype 1
SWPHLSB_12	L22013 Swinepox virus
CV41KBPL_14	Cowpox virus
VVCGAA_5	Variola virus
U01161_3 174	Ectromelia virus (mouse Poxvirus)
VVU18340_6	Variola virus
VVU18338_7	Variola virus
VVU18337_7	Variola virus
VARCG_7 173	Variola major virus
MCU60315_51	Molluscum contagiosum virus
HNABV_1	New Hepatitis non-A, non-B associated virus

[0125] IL-18BP_a was expressed in monkey COS7 cells. For this purpose, the cDNA of IL-18BP_a was inserted into the mammalian expression vector pEF-BOS. A cassette coding for an (His)₆ sequence was added to the 3'-end of the IL-18BP ORFs in frame, in order to facilitate purification of the recombinant protein. COS7 cells were transiently transfected with the expression vector and serum-free medium of these cells (150 ml) was concentrated and purified by metal chelate chromatography. IL-18BP_a ran as a single band upon SDS-PAGE with silver staining under reducing and non-reducing conditions and had the same apparent molecular mass as that of the urinary IL-18BP. Protein sequence analysis of this preparation revealed the same N-terminal sequence as that of the urinary IL-18BP. Immunoblot analysis of IL-18BP_a with antibodies raised against the urinary IL-18BP revealed the same molecular mass band as that of the urinary protein. Furthermore, using immunoprecipitation followed by SDS-PAGE and autoradiography, IL-18BP_a was able to displace

urinary ^{125}I -IL-18BP from binding to the antibody. Therefore, IL-18BP_a corresponds structurally to the IL-18BP isolated from urine.

[0126] Crude and purified IL-18BP_a were tested for their ability to inhibit the biological activity of IL-18. IL-18BP_a inhibited the activity of human and mouse IL-18 in murine splenocytes, PBMC and the human KG-1 cell line (FIG. 4). In contrast, IL-18BP_b did not significantly inhibit the activity of IL-18. These results confirm the identity of IL-18BP_a cDNA as the one coding for a biologically active IL-18BP.

[0127] The invention further relates to active muteins and fragments of IL-18BPs and viral IL-18BPs and to fused proteins consisting of wild type IL-18BPs and viral IL-18BPs or their active muteins or their active fragments, fused to another polypeptide or protein and exhibiting a similar ability to block the biological activities of IL-18 or its homologues.

[0128] As used herein the term “muteins” refers to analogs of an IL-18BP, or analogs of a viral IL-18BP, in which one or more of the amino acid residues of a natural IL-18BP or viral IL-18BP are replaced by different amino acid residues, or are deleted, or one or more amino acid residues are added to the natural sequence of an IL-18BP, or a viral IL-18BP, without changing considerably the activity of the resulting products as compared with the wild type IL-18BP or viral IL-18BP. These muteins are prepared by known synthesis and/or by site-directed mutagenesis techniques, or any other known technique suitable therefor.

[0129] Any such mutein preferably has a sequence of amino acids sufficiently duplicative of that of an IL-18BP, or sufficiently duplicative of a viral IL-18BP, such as to have substantially similar activity to IL-18BP. One activity of IL-18BP is its capability of binding IL-18. As long as the mutein has substantial binding activity to IL-18, it can be used in the purification of IL-18, such as by means of affinity chromatography, and thus can be considered to have substantially similar activity to IL-18BP. Thus, it can be determined whether any given mutein has substantially the same activity as IL-18BP by means of routine experimentation comprising subjecting such a mutein, e.g., to a simple sandwich competition assay to determine whether or not it binds to an appropriately labeled IL-18, such as radioimmunoassay or ELISA assay.

[0130] In a preferred embodiment, any such mutein has at least 40% identity or homology with the sequence of either an IL-18BP or a virally-encoded IL-18BP homologue. More preferably, it has at least 50%, at least 60%, at least 70%, at least 80% or, most preferably, at least 90% identity or homology thereto.

[0131] Muteins of IL-18BP polypeptides or muteins of viral IL-18BPs, which can be used in accordance with the present invention, or nucleic acid coding therefor, include a finite set of substantially corresponding sequences as substitution peptides or polynucleotides which can be routinely obtained by one of ordinary skill in the art, without undue experimentation, based on the teachings and guidance presented herein. For a detailed description of protein chemistry and structure, see Schulz, G. E. et al., *Principles of Protein Structure*, Springer-Verlag, New York, 1978; and Creighton, T. E., *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, 1983, which are hereby incorporated by reference. For a presentation of nucleotide sequence substitutions, such as codon preferences, see Ausubel et al, supra, at §§ A.1.1-A.1.24, and Sambrook et al, supra, at Appendices C and D.

[0132] Preferred changes for muteins in accordance with the present invention are what are known as “conservative” substitutions. Conservative amino acid substitutions of IL-18BP polypeptides or proteins or viral IL-18BPs, may include synonymous amino acids within a group which have sufficiently similar physicochemical properties that substitution between members of the group will preserve the biological function of the molecule, Grantham, *Science*, Vol. 185, pp. 862-864 (1974). It is clear that insertions and deletions of amino acids may also be made in the above-defined sequences without altering their function, particularly if the insertions or deletions only involve a few amino acids, e.g., under thirty, and preferably under ten, and do not remove or displace amino acids which are critical to a functional conformation, e.g., cysteine residues, Anfinsen, “*Principles That Govern The Folding of Protein Chains*”, *Science*, Vol. 181, pp. 223-230 (1973). Proteins and muteins produced by such deletions and/or insertions come within the purview of the present invention.

[0133] Preferably, the synonymous amino acid groups are those defined in Table I. More preferably, the synonymous amino acid groups are those defined in Table 2; and most preferably the synonymous amino acid groups are those defined in Table 31.

TABLE 1

Preferred Groups of Synonymous Amino Acids	
Amino Acid	Synonymous Group
Ser	Ser, Thr, Gly, Asn
Arg	Arg, Gln, Lys, Glu, His
Leu	Ile, Phe, Tyr, Met, Val, Leu
Pro	Gly, Ala, Thr, Pro
Thr	Pro, Ser, Ala, Gly, His, Gln, Thr
Ala	Gly, Thr, Pro, Ala
Val	Met, Tyr, Phe, Ile, Leu, Val
Gly	Ala, Thr, Pro, Ser, Gly
Ile	Met, Tyr, Phe, Val, Leu, Ile
Phe	Trp, Met, Tyr, Ile, Val, Leu, Phe
Tyr	Trp, Met, Phe, Ile, Val, Leu, Tyr
Cys	Ser, Thr, Cys
His	Glu, Lys, Gln, Thr, Arg, His
Gln	Glu, Lys, Asn, His, Thr, Arg, Gln
Asn	Gln, Asp, Ser, Asn
Lys	Glu, Gln, His, Arg, Lys
Asp	Glu, Asn, Asp
Glu	Asp, Lys, Asn, Gln, His, Arg, Glu
Met	Phe, Ile, Val, Leu, Met
Trp	Trp

TABLE 2

More Preferred Groups of Synonymous Amino Acids	
Amino Acid	Synonymous Group
Ser	Ser
Arg	His, Lys, Arg
Leu	Leu, Ile, Phe, Met
Pro	Ala, Pro
Thr	Thr
Ala	Pro, Ala
Val	Val, Met, Ile
Gly	Gly
Ile	Ile, Met, Phe, Val, Leu
Phe	Met, Tyr, Ile, Leu, Phe
Tyr	Phe, Tyr
Cys	Cys, Ser

TABLE 2-continued

More Preferred Groups of Synonymous Amino Acids	
Amino Acid	Synonymous Group
His	His, Gln, Arg
Gln	Glu, Gln, His
Asn	Asp, Asn
Lys	Lys, Arg
Asp	Asp, Asn
Glu	Glu, Gln
Met	Met, Phe, Ile, Val, Leu
Trp	Trp

TABLE 3

Most Preferred Groups of Synonymous Amino Acids	
Amino Acid	Synonymous Group
Ser	Ser
Arg	Arg
Leu	Leu, Ile, Met
Pro	Pro
Thr	Thr
Ala	Ala
Val	Val
Gly	Gly
Ile	Ile, Met, Leu
Phe	Phe
Tyr	Tyr
Cys	Cys, Ser
His	His
Gln	Gln
Asn	Asn
Lys	Lys
Asp	Asp
Glu	Glu
Met	Met, Ile, Leu
Trp	Met

[0134] Examples of production of amino acid substitutions in proteins which can be used for obtaining muteins of IL-18BP polypeptides or proteins, or muteins of viral IL-18BPs, for use in the present invention include any known method steps, such as presented in U.S. Pat. Nos. RE 33,653, 4,959,314, 4,588,585 and 4,737,462, to Mark et al; 5,116,943 to Kothe et al., 4,965,195 to Namen et al; 4,879,111 to Chong et al; and 5,017,691 to Lee et al; and lysine substituted proteins presented in U.S. Pat. No. 4,904,584 (Shaw et al).

[0135] In another preferred embodiment of the present invention, any mutein of an IL-18BP or a viral IL-18BP, has an amino acid sequence essentially corresponding to that of an IL-18BP, or to a viral IL-18BP. The term “essentially corresponding to” is intended to comprehend proteins with minor changes to the sequence of the natural protein which do not affect the basic characteristics of the natural proteins, particularly insofar as their ability to bind IL-18 and to thereby inhibit its biological activity. The type of changes which are generally considered to fall within the “essentially corresponding to” language are those which would result from conventional mutagenesis techniques of the DNA encoding these proteins, resulting in a few minor modifications, and screening for the desired activity in the manner discussed above. Muteins in accordance with the present invention include proteins encoded by a nucleic acid, such as DNA or RNA, which hybridizes to DNA or RNA, which encodes an IL-18BP or encodes a viral IL-18BP, in accor-

dance with the present invention, under stringent conditions. The invention also includes such nucleic acid, which is also useful as a probe in identification and purification of the desired nucleic acid. Furthermore, such nucleic acid would be a prime candidate to determine whether it encodes a polypeptide, which retains the functional activity of an IL-18BP of the present invention. The term “stringent conditions” refers to hybridization and subsequent washing conditions, which those of ordinary skill in the art conventionally refer to as “stringent”. See Ausubel et al., Current Protocols in Molecular Biology, supra, Interscience, N.Y., §§6.3 and 6.4 (1987, 1992), and Sambrook et al., supra. For example high stringent conditions include washing conditions 12-20° C. below the calculated T_m of the hybrid under study in, e.g., 2×SSC and 0.5% SDS for 5 minutes, 2×SSC and 0.1% SDS for 15 minutes; 0.1×SSC and 0.5% SDS at 37° C. for 30-60 minutes and then, a 0.1×SSC and 0.5% SDS at 68° C. for 30-60 minutes. Those of ordinary skill in this art understand that stringency conditions also depend on the length of the DNA sequences, oligonucleotide probes (such as 10-40 bases) or mixed oligonucleotide probes. If mixed probes are used, it is preferable to use tetramethyl ammonium chloride (TMAC) instead of SSC. See Ausubel, supra.

[0136] The term “fused protein” refers to a polypeptide comprising an IL-18BP, or a viral IL-18BP, or a mutein thereof, fused with another protein, which, e.g., has an extended residence time in body fluids. An IL-18BP or a viral IL-18BP, may thus be fused to another protein, polypeptide or the like, e.g., an immunoglobulin or a fragment thereof.

[0137] The term “salts” herein refers to both salts of carboxyl groups and to acid addition salts of amino groups of an IL-18BP, a viral IL-18BP, muteins, or fused proteins thereof. Salts of a carboxyl group may be formed by means known in the art and include inorganic salts, for example, sodium, calcium, ammonium, ferric or zinc salts, and the like, and salts with organic bases as those formed, for example, with amines, such as triethanolamine, arginine or lysine, piperidine, procaine and the like. Acid addition salts include, for example, salts with mineral acids such as, for example, hydrochloric acid or sulfuric acid, and salts with organic acids such as, for example, acetic acid or oxalic acid. Of course, any such salts must have substantially similar activity to IL-18BP.

[0138] “Functional derivatives” as used herein cover derivatives of IL-18BPs or a viral IL-18BP, and their muteins and fused proteins, which may be prepared from the functional groups which occur as side chains on the residues or the N- or C-terminal groups, by means known in the art, and are included in the invention as long as they remain pharmaceutically acceptable, i.e. they do not destroy the activity of the protein which is substantially similar to the activity of IL-18BP, or viral IL-18BPs, and do not confer toxic properties on compositions containing it. These derivatives may, for example, include polyethylene glycol side-chains, which may mask antigenic sites and extend the residence of an IL-18BP or a viral IL-18BP in body fluids. Other derivatives include aliphatic esters of the carboxyl groups, amides of the carboxyl groups by reaction with ammonia or with primary or secondary amines, N-acyl derivatives of free amino groups of the amino acid residues formed with acyl moieties (e.g. alkanoyl or carbocyclic aroyl groups) or O-acyl derivatives of free hydroxyl groups (for example that of seryl or threonyl residues) formed with acyl moieties.

[0139] As “active fractions” of an IL-18BP, or a viral IL-18BP, muteins and fused proteins, the present invention

covers any fragment or precursors of the polypeptide chain of the protein molecule alone or together with associated molecules or residues linked thereto, e.g., sugar or phosphate residues, or aggregates of the protein molecule or the sugar residues by themselves, provided said fraction has substantially similar activity to IL-18BP.

[0140] Various recombinant cells such as prokaryotic cells, e.g., *E. coli*, or other eukaryotic cells, such as yeast or insect cells can produce IL-18BPs or viral IL-18BPs. Methods for constructing appropriate vectors, carrying DNA that codes for an IL-18BP and suitable for transforming (e.g., *E. coli*, mammalian cells and yeast cells), or infecting insect cells in order to produce a recombinant IL-18BP or a viral IL-18BP are well known in the art. See, for example, Ausubel et al., eds. "Current Protocols in Molecular Biology" Current Protocols, 1993; and Sambrook et al., eds. "Molecular Cloning: A Laboratory Manual", 2nd ed., Cold Spring Harbor Press, 1989.

[0141] Cells that produce IL-18 BPs or viral IL-18BPs are made according to known procedures in the art, including for example, by gene activation (See, PCT publication WO90/11354, U.S. Pat. Nos. 5,272,071, and 5,641,670, which are incorporated herein by reference in their entirety), and transduction.

[0142] For the purposes of expression of IL-18BP proteins, or viral IL-18BPs, DNA encoding an IL-18BP or a viral IL-18BP, their fragments, muteins or fused proteins, and the operably linked transcriptional and translational regulatory signals, are inserted into eukaryotic vectors which are capable of integrating the desired gene sequences into the host cell chromosome. In order to be able to select the cells which have stably integrated the introduced DNA into their chromosomes, one or more markers which allow for selection of host cells which contain the expression vector is used. The marker may provide for prototrophy to an auxotrophic host, biocide resistance, e.g., antibiotics, or resistance to heavy metals, such as copper, or the like. The selectable marker gene can either be directly linked to the DNA gene sequences to be expressed, or introduced into the same cell by cotransfection. Additional elements may also be needed for optimal synthesis of single chain binding protein mRNA. These elements may include splice signals, as well as transcription promoters, enhancers, and termination signals.

[0143] Said DNA molecule to be introduced into the cells of choice will preferably be incorporated into a plasmid or viral vector capable of autonomous replication in the recipient host. Preferred prokaryotic plasmids are derivatives of pBr322. Preferred eukaryotic vectors include BPV, vaccinia, SV40, 2-micron circle, etc., or their derivatives. Such plasmids and vectors are well known in the art (Anderson, 1997; Bolton, 1980; Botstein, 1982; Broach, 1981; Kendall et al., 1987). Once the vector or DNA sequence containing the construct(s) has been prepared for expression, the expression vector may be introduced into an appropriate host cell by any of a variety of suitable means, such as transformation, transfection, lipofection, conjugation, protoplast fusion, electroporation, calcium phosphate precipitation, direct microinjection, etc.

[0144] Host cells to be used in this invention may be either prokaryotic or eukaryotic. Preferred prokaryotic hosts include bacteria such as *E. coli*, *Bacillus*, *Streptomyces*, *Pseudomonas*, *Salmonella*, *Serratia*, etc. The most preferred prokaryotic host is *E. coli*. Bacterial hosts of particular interest include *E. coli* K12 strain 294 (ATCC 31446), *E. coli*

X1776 (ATCC 31537), *E. coli* W3110 (F⁻, lambda⁻, phototropic (ATCC 27325). Under such conditions, the protein will not be glycosylated. The prokaryotic host must be compatible with the replicon and control sequences in the expression plasmid.

[0145] However, since IL-18BPs are glycosylated proteins, eukaryotic hosts are preferred over prokaryotic hosts. Preferred eukaryotic hosts are mammalian cells, e.g., human, monkey, mouse and Chinese hamster ovary (CHO) cells, because they provide post-translational modifications to protein molecules including correct folding, correct disulfide bond formation, as well as glycosylation at correct sites. Also yeast cells and insect cells can carry out post-translational peptide modifications including high mannose glycosylation.

[0146] A number of recombinant DNA strategies exist which utilize strong promoter sequences and high copy number of plasmids, which can be utilized for production of the desired proteins in yeast and in insect cells. Yeast and insect cells recognize leader sequences on cloned mammalian gene products and secrete mature IL-18BP. After the introduction of the vector, the host cells are grown in a selective medium, which selects for the growth of vector-containing cells. Expression of the cloned gene sequence(s) results in the production of an IL-18BP, a viral IL-18BP, fusion proteins, or muteins or fragments thereof. The above-mentioned cloning, clone isolation, identification, characterization and sequencing procedures are described in more detail hereinafter in the Examples.

[0147] The expressed proteins are then isolated and purified by any conventional procedure involving extraction, precipitation, chromatography, electrophoresis, or the like, or by affinity chromatography, using, e.g., an anti-IL-18BP monoclonal antibodies immobilized on a gel matrix contained within a column. Crude preparations containing said recombinant IL-18BP are passed through the column whereby IL-18BP will be bound to the column by the specific antibody, while the impurities will pass through. After washing, the protein is eluted from the gel under conditions usually employed for this purpose, i.e. at a high or a low pH, e.g. pH 11 or pH 2.

[0148] The invention further relates to vectors useful for expression of an IL-18BP or a viral IL-18BP or their derivatives in mammals and more specifically in humans. Vectors for short and long-term expression of genes in mammals are well known in the literature. Studies have shown that gene delivery to e.g., skeletal muscle, vascular smooth muscle and liver result in systemic levels of therapeutic proteins. Skeletal muscle is a useful target because of its large mass, vascularity and accessibility. However, other targets and particularly bone marrow precursors of immune cells have been used successfully. Currently available vectors for expression of proteins in e.g., muscle include plasmid DNA, liposomes, protein-DNA conjugates and vectors based on adenovirus, adeno-associated virus and herpes virus. Of these, vectors based on adeno-associated virus (AAV) have been most successful with respect to duration and levels of gene expression and with respect to safety considerations (Kessler, P. D. 1996, Proc. Natl. Acad. Sci. USA 93, 14082-14087).

[0149] Procedures for construction of an AAV-based vector have been described in detail (Snyder et al, 1996, Current Protocols in Human Genetics, Chapters 12.1.1-12.1.17, John Wiley & Sons) and are incorporated into this patent. Briefly plasmid psub201, containing the wild-type AAV genome is cut with the restriction enzyme Xba I and ligated with a

construct consisting of an efficient eukaryotic promoter, e.g., the cytomegalovirus promoter, a Kozak consensus sequence, a DNA sequence coding for an IL-18BP or a viral IL-18BP, or their muteins or fusion proteins or fragments thereof, a suitable 3' untranslated region and a polyadenylation signal, e.g., the polyadenylation signal of simian virus 40. The resulting recombinant plasmid is cotransfected with an helper AAV plasmid e.g., pAAV/Ad into mammalian cells e.g., human T293 cells. The cultures are then infected with adenovirus as a helper virus and culture supernatants are collected after 48-60 hours. The supernatants are fractionated by ammonium sulfate precipitation, purified on a CsCl density gradient, dialyzed and then heated at 56° C. to destroy any adenovirus, whereas the resulting recombinant AAV, capable of expressing IL-18BP or a viral IL-18BP, or their muteins or fusion proteins remains stable at this step.

[0150] So far, the physiological role of the soluble cytokine receptors has not been established. The soluble receptors bind their specific ligands and in most cases inhibit their biological activity, as was shown, e.g., in the TNF system (Dao et al., 1996; Engelmann et al., 1989). In very few cases, e.g., IL-6, the soluble receptor enhances the biological activity. The recombinant soluble TNF receptor, also known as TBP (TNF binding protein) was found to prevent septic shock in animal models, while soluble forms of IL-1 receptor were found to have profound inhibitory effects on the development of in vivo alloreactivity in mouse allograft recipients.

[0151] Similarly, the IL-18BPs and viral IL-18BPs of the present invention may find use as modulators of IL-18 activity, e.g. in type I diabetes, in sepsis, in autoimmune diseases, in graft rejections, rheumatoid arthritis, inflammatory bowel disease, sepsis, multiple sclerosis, ischemic heart disease including acute heart attacks, ischemic brain injury, chronic hepatitis, psoriasis, chronic hepatitis and acute hepatitis. It may thus be used, e.g. in any disease in which endogenous production or exogenous administration of IL-18 induces the disease or aggravates the situation of the patient.

[0152] The present invention further relates to pharmaceutical compositions comprising a pharmaceutically acceptable carrier and an IL-18BP or a viral IL-18BP of the invention or their active muteins, fused proteins and their salts, functional derivatives or active fractions thereof.

[0153] The present invention further relates to pharmaceutical compositions comprising a pharmaceutically acceptable carrier and e.g., a viral vector such as any one of said AAV-based viral vectors or another vector expressing an IL-18BP or viral IL-18BP or their muteins, fragments or fusion proteins thereof and suitable for administration to humans and other mammals for the purpose of attaining expression in vivo of IL-18BP or a viral IL-18BP or their muteins or fragments or fusion protein of the invention.

[0154] The pharmaceutical compositions of the invention are prepared for administration by mixing an IL-18BP or a viral IL-18BP, or their derivatives, or vectors for expressing same with physiologically acceptable carriers, and/or stabilizers and/or excipients, and prepared in dosage form, e.g., by lyophilization in dosage vials. The method of administration can be via any of the accepted modes of administration for similar agents and will depend on the condition to be treated, e.g., intravenously, intramuscularly, subcutaneously, by local injection or topical application, or continuously by infusion, etc. The amount of active compound to be administered will depend on the route of administration, the disease to be treated and the condition of the patient. Local injection, for

instance, will require a lower amount of the protein on a body weight basis than will intravenous infusion.

[0155] Accordingly, IL-18BPs, or viral IL-18BPs, or vectors expressing same in vivo are indicated for the treatment of autoimmune diseases, Type I diabetes, rheumatoid arthritis, graft rejections, inflammatory bowel disease, sepsis, multiple sclerosis, ischemic heart disease including acute heart attacks, ischemic brain injury, chronic hepatitis, psoriasis, chronic pancreatitis and acute pancreatitis and similar diseases, in which there is an aberrant expression of IL-18, leading to an excess of IL-18 or in cases of complications due to exogenously administered IL-18.

[0156] The invention also includes antibodies against an IL-18BP or a viral IL-18BP, as well as against their muteins, fused proteins, salts, functional derivatives and active fractions. The term "antibody" is meant to include polyclonal antibodies, monoclonal antibodies (MAbs), chimeric antibodies, anti-idiotypic (anti-Id) antibodies to antibodies that can be labeled in soluble or bound form, as well as fragments thereof provided by any known technique, such as, but not limited to enzymatic cleavage, peptide synthesis or recombinant techniques.

[0157] Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen. A monoclonal antibody contains a substantially homogeneous population of antibodies specific to antigens, which population contains substantially similar epitope binding sites. MAbs may be obtained by methods known to those skilled in the art. See, for example Kohler and Milstein, *Nature* 256:495-497 (1975); U.S. Pat. No. 4,376,110; Ausubel et al, eds., supra, Harlow and Lane, *ANTIBODIES: A LABORATORY MANUAL*, Cold Spring Harbor Laboratory (1988); and Colligan et al., eds., *Current Protocols in Immunology*, Greene Publishing Assoc. and Wiley Interscience, N.Y., (1992, 1993), the contents of which references are incorporated entirely herein by reference. Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, GILD and any subclass thereof. A hybridoma producing a MAb of the present invention may be cultivated in vitro, in situ or in vivo. Production of high titers of MAbs in vivo or in situ makes this the presently preferred method of production.

[0158] Chimeric antibodies are molecules, different portions of which are derived from different animal species, such as those having the variable region derived from a murine MAb and a human immunoglobulin constant region. Chimeric antibodies are primarily used to reduce immunogenicity in application and to increase yields in production, for example, where murine MAbs have higher yields from hybridomas but higher immunogenicity in humans, such that human/murine chimeric MAbs are used. Chimeric antibodies and methods for their production are known in the art (Cabilly et al, *Proc. Natl. Acad. Sci. USA* 81:3273-3277 (1984); Morrison et al., *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984); Boulianne et al., *Nature* 312:643-646 (1984); Cabilly et al., *European Patent Application* 125023 (published Nov. 14, 1984); Neuberger et al., *Nature* 314:268-270 (1985); Taniguchi et al., *European Patent Application* 171496 (published Feb. 19, 1985); Morrison et al., *European Patent Application* 173494 (published Mar. 5, 1986); Neuberger et al., *PCT Application WO 8601533*, (published Mar. 13, 1986); Kudo et al., *European Patent Application* 184187 (published Jun. 11, 1986); Morrison et al., *European Patent Application* 173494 (published Mar. 5, 1986); Sahagan et al., *J. Immunol.*

137:1066-1074 (1986); Robinson et al., International Patent Publication, WO 9702671 (published 7 May 1987); Liu et al., Proc. Natl. Acad. Sci. USA 84:3439-3443 (1987); Sun et al., Proc. Natl. Acad. Sci. USA 84:214-218 (1987); Better et al., Science 240:1041-1043 (1988); and Harlow and Lane, ANTI-BODIES: A LABORATORY MANUAL, supra. These references are entirely incorporated herein by reference.

[0159] An anti-idiotypic (anti-Id) antibody is an antibody, which recognizes unique determinants generally, associated with the antigen-binding site of an antibody. An Id antibody can be prepared by immunizing an animal of the same species and genetic type (e.g., mouse strain) as the source of the MAb with the MAb to which an anti-Id is being prepared. The immunized animal will recognize and respond to the idiotypic determinants of the immunizing antibody by producing an antibody to these idiotypic determinants (the anti-Id antibody). See, for example, U.S. Pat. No. 4,699,880, which is herein entirely incorporated by reference.

[0160] The anti-Id antibody may also be used as an "immunogen" to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody. The anti-anti-Id may be epitopically identical to the original MAb, which induced the anti-Id. Thus, by using antibodies to the idiotypic determinants of a MAb, it is possible to identify other clones expressing antibodies of identical specificity.

[0161] Accordingly, MAbs generated against IL-18BP and related proteins of the present invention may be used to induce anti-Id antibodies in suitable animals, such as BALB/c mice. Spleen cells from such immunized mice are used to produce anti-Id hybridomas secreting anti-Id MAbs. Further, the anti-Id MAbs can be coupled to a carrier such as keyhole limpet hemocyanin (KLH) and used to immunize additional BALB/c mice. Sera from these mice will contain anti-anti-Id antibodies that have the binding properties of the original MAb specific for an IL-18BP epitope or epitopes of a viral IL-18BP.

[0162] The anti-Id MAbs thus have their own idiotypic epitopes, or "idiotopes" structurally similar to the epitope being evaluated, such as an IL-18BP or a viral IL-18BP.

[0163] The term "antibody" is also meant to include both intact molecules as well as fragments thereof, such as, for example, Fab and F(ab')₂, which are capable of binding antigen. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl et al., J. Nucl. Med. 24:316-325 (1983)). It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies useful in the present invention may be used for the detection and quantitation of an IL-18BP or a viral IL-18BP, according to the methods disclosed herein for intact antibody molecules. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments).

[0164] An antibody is said to be "capable of binding" a molecule if it is capable of specifically reacting with the molecule to thereby bind the molecule to the antibody. The term "epitope" is meant to refer to that portion of any molecule capable of being bound by an antibody which can also be recognized by that antibody. Epitopes or "antigenic determinants" usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and have specific three dimensional structural characteristics as well as specific charge characteristics.

[0165] An "antigen" is a molecule or a portion of a molecule capable of being bound by an antibody which is additionally capable of inducing an animal to produce antibody capable of binding to an epitope of that antigen. An antigen may have one or more than one epitope. The specific reaction referred to above is meant to indicate that the antigen will react, in a highly selective manner, with its corresponding antibody and not with the multitude of other antibodies which may be evoked by other antigens.

[0166] The antibodies, including fragments of antibodies, useful in the present invention may be used to detect quantitatively or qualitatively an IL-18BP or a viral IL-18BP, or related proteins in a sample or to detect presence of cells, which express such proteins of the present invention. This can be accomplished by immunofluorescence techniques employing a fluorescently labeled antibody (see below) coupled with light microscopic, flow cytometric, or fluorometric detection.

[0167] The antibodies (or fragments thereof) useful in the present invention may be employed histologically, as in immunofluorescence or immunoelectron microscopy, for in situ detection of an IL-18BP or a viral IL-18BP, and related proteins of the present invention. In situ detection may be accomplished by removing a histological specimen from a patient, and providing the a labeled antibody of the present invention to such a specimen. The antibody (or fragment) is preferably provided by applying or by overlaying the labeled antibody (or fragment) to a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of an IL-18BP or a viral IL-18BP, or related proteins but also its distribution on the examined tissue. Using the present invention, those of ordinary skill will readily perceive that any of wide variety of histological methods (such as staining procedures) can be modified in order to achieve such in situ detection.

[0168] Such assays for an IL-18BP or a viral IL-18BP, or related proteins of the present invention typically comprises incubating a biological sample, such as a biological fluid, a tissue extract, freshly harvested cells such as lymphocytes or leukocytes, or cells which have been incubated in tissue culture, in the presence of a detectably labeled antibody capable of identifying IL-18BP or related proteins, and detecting the antibody by any of a number of techniques well-known in the art.

[0169] The biological sample may be treated with a solid phase support or carrier such as nitrocellulose, or other solid support or carrier which is capable of immobilizing cells, cell particles or soluble proteins. The support or carrier may then be washed with suitable buffers followed by treatment with a detectably labeled antibody in accordance with the present invention. The solid phase support or carrier may then be washed with the buffer a second time to remove unbound antibody. The amount of bound label on said solid support or carrier may then be detected by conventional means.

[0170] By "solid phase support", "solid phase carrier", "solid support", "solid carrier", "support" or "carrier" is intended any support or carrier capable of binding antigen or antibodies. Well-known supports or carriers, include glass, polystyrene, polypropylene, polyethylene, dextran, nylon amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled

molecule is capable of binding to an antigen or antibody. Thus, the support or carrier configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports or carriers include polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

[0171] The binding activity of a given lot of antibody in accordance with the present invention may be determined according to well known methods. Those skilled in the art will be able to determine operative and optimal assay conditions for each determination by employing routine experimentation.

[0172] Other such steps as washing, stirring, shaking, filtering and the like may be added to the assays as is customary or necessary for the particular situation.

[0173] One of the ways in which an antibody in accordance with the present invention can be detectably labeled is by linking the same to an enzyme and use in an enzyme immunoassay (EIA). This enzyme, in turn, when later exposed to an appropriate substrate, will react with the substrate in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorometric or by visual means. Enzymes which can be used detectably label the antibody include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colorimetric methods, which employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

[0174] Detection may be accomplished using any of a variety of other immunoassays. For example, by radioactivity labeling the antibodies or antibody fragments, it is possible to detect an IL-18BP or a viral IL-18BP, through the use of a radioimmunoassay (RIA). A good description of RIA may be found in *Laboratory Techniques and Biochemistry in Molecular Biology*, by Work, T. S. et al., North Holland Publishing Company, NY (1978) with particular reference to the chapter entitled "An Introduction to Radioimmuno Assay and Related Techniques" by Chard, T., incorporated by reference herein. The radioactive isotope can be detected by such means as the use of a gamma counter or a scintillation counter or by autoradiography.

[0175] It is also possible to label an antibody in accordance with the present invention with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wavelength, its presence can be then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

[0176] The antibody can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriamine pentaacetic acid (ETPA). The antibody can also be detectably

labeled by coupling it to biotin. Biotinylated antibody can then be detected by avidin or streptavidin coupled to a fluorescent compound or to an enzyme such as peroxidase or to a radioactive isotope and the like.

[0177] The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, thermotropic acridinium ester, imidazole, acridinium salt and oxalate ester.

[0178] Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

[0179] An antibody molecule of the present invention may be adapted for utilization in an immunometric assay, also known as a "two-site" or "sandwich" assay. In a typical immunometric assay, a quantity of unlabeled antibody (or fragment of antibody) is bound to a solid support or carrier and a quantity of detectably labeled soluble antibody is added to permit detection and/or quantitation of the ternary complex formed between solid-phase antibody, antigen, and labeled antibody.

[0180] Typical, and preferred, immunometric assays include "forward" assays in which the antibody bound to the solid phase is first contacted with the sample being tested to extract the antigen from the sample by formation of a binary solid phase antibody-antigen complex. After a suitable incubation period, the solid support or carrier is washed to remove the residue of the fluid sample, including unreacted antigen, if any, and then contacted with the solution containing an unknown quantity of labeled antibody (which functions as a "reporter molecule"). After a second incubation period to permit the labeled antibody to complex with the antigen bound to the solid support or carrier through the unlabeled antibody, the solid support or carrier is washed a second time to remove the unreacted labeled antibody.

[0181] In another type of "sandwich" assay, which may also be useful with the antigens of the present invention, the so-called "simultaneous" and "reverse" assays are used. A "simultaneous" assay involves a single incubation step as the antibody bound to the solid support or carrier and labeled antibody are both added to the sample being tested at the same time. After the incubation is completed, the solid support or carrier is washed to remove the residue of fluid sample and uncomplexed labeled antibody. The presence of labeled antibody associated with the solid support or carrier is then determined as it would be in a conventional "forward" sandwich assay.

[0182] In the "reverse" assay, stepwise addition first of a solution of labeled antibody to the fluid sample followed by the addition of unlabeled antibody bound to a solid support or carrier after a suitable incubation period is utilized. After a second incubation, the solid phase is washed in conventional fashion to free it of the residue of the sample being tested and the solution of unreacted labeled antibody. The determination

of labeled antibody associated with a solid support or carrier is then determined as in the “simultaneous” and “forward” assays.

[0183] The present invention also provides DNA molecules encoding any of the proteins of the present invention as defined above, replicable expression vehicles comprising any such DNA molecules, host cells transformed with any such expression vehicles including prokaryotic and eukaryotic and host cells, preferably CHO cells. The invention also includes a process for the production of expression vectors coding for any of the proteins of the present invention for the purpose of their expression in humans and other mammals.

[0184] The invention also includes a process for the production of any of the proteins of the present invention by culturing a transformed cell in accordance with the present invention and recovering the protein encoded by the DNA molecule and the expression vehicle within such transformed host cell.

[0185] In addition to the use of an IL-18BP or a viral IL-18BP, in modulating the activity of IL-18, they can, of course, also be employed for the purification of IL-18 itself. For this purpose, IL-18BP or a viral IL-18BP is coupled to an affinity column and crude IL-18 is passed through. The IL-18 can then be recovered from the column by, e.g., elution at low pH.

[0186] The present invention further relates to the treatment and/or prevention of liver injury using inhibitors of IL-18. An inhibitor according to the invention can be an inhibitor of IL-18 production and/or action.

[0187] The term “inhibitor of IL-18” within the context of this invention refers to any molecule modulating IL-18 production and/or action in such a way that IL-18 production and/or action is attenuated, reduced, or partially, substantially or completely prevented or blocked.

[0188] An inhibitor of production can be any molecule negatively affecting the synthesis, processing or maturation of IL-18 on the level of the gene, the mRNA or the protein. The inhibitors considered according to the invention can be, for example, suppressors of gene expression of the interleukin IL-18, antisense mRNAs reducing or preventing the transcription of the IL-18 mRNA or leading to degradation of the mRNA, proteins impairing correct folding, or partially or substantively preventing secretion of IL-18, proteases degrading the IL-18, once it has been synthesized, and the like.

[0189] An inhibitor of IL-18 action can be an IL-18 antagonist, for example. Antagonists can either bind to or sequester the IL-18 molecule itself with sufficient affinity and specificity to partially or substantially neutralize the IL-18 or IL-18 binding site(s) responsible for IL-18 binding to its ligands (like, e.g. to its receptors). An antagonist may also inhibit the IL-18 signaling pathway, activated within the cells upon IL-18/receptor binding.

[0190] Inhibitors of IL-18 action may be also soluble IL-18 receptors or molecules mimicking the receptors, or agents blocking the IL-18 receptors, IL-18 antibodies, like monoclonal antibodies, for example, or any other agent or molecule preventing the binding of IL-18 to its targets, thus diminishing or preventing triggering of the intra- or extracellular reactions mediated by IL-18.

[0191] The invention particularly relates to for the treatment or prevention of both acute and chronic liver diseases,

such as alcoholic hepatitis, viral hepatitis, immune hepatitis, fulminant hepatitis, liver cirrhosis, and primary biliary cirrhosis, for example.

[0192] The IL-18 inhibitors contemplated herein can further be used for the treatment of acute hepatic poisoning caused by an high amount of paracetamol. Such an acute hepatic poisoning may be due to an overdose, be it accidental or on purpose.

[0193] As shown in the examples below, it has been surprisingly found that IL-18 inhibitors are particularly effective in the prevention and treatment of fulminant hepatitis (acute hepatitis). Therefore, the invention preferably relates to the prevention and/or treatment of fulminant hepatitis.

[0194] It has been surprisingly found that a combination of an inhibitor of IL-18 with an inhibitor of tumor necrosis factor (TNF) leads to a complete blockade of liver injury in a murine model of disease. The invention therefore further relates to the use of an IL-18 inhibitor in combination with a TNF-inhibitor for the treatment and/or prevention of liver injury.

[0195] An inhibitor of TNF can be a TNF antagonist, for example. TNF antagonists exert their activity in several ways. First, antagonists can bind to or sequester the TNF molecule itself with sufficient affinity and specificity to partially or substantially neutralize the TNF epitope or epitopes responsible for TNF receptor binding (hereinafter termed “sequestering antagonists”). A sequestering antagonist may be, for example, an antibody directed against TNF.

[0196] Alternatively, TNF antagonists can inhibit the TNF signaling pathway activated by the cell surface receptor after TNF binding (hereinafter termed “signaling antagonists”). Both groups of antagonists are useful, either alone or together, in combination with an IL-18 inhibitor, in the prevention of therapy of liver injury.

[0197] TNF antagonists are easily identified and rated by routine screening of candidates for their effect on the activity of native TNF on susceptible cell lines in vitro, for example human B cells, in which TNF causes proliferation and immunoglobulin secretion. The assay contains TNF formulation at varying dilutions of candidate antagonist, e.g. from 0.1 to 100 times the molar amount of TNF used in the assay, and controls with no TNF or only antagonist (Tucci et al., 1992).

[0198] Sequestering antagonists are the preferred TNF antagonists to be used according to the present invention. Amongst sequestering antagonists, those polypeptides that bind TNF with high affinity and possess low immunogenicity are preferred. Soluble TNF receptor molecules and neutralizing antibodies to TNF are particularly preferred. For example, soluble TNF receptors TNF-RI and TNF-RII are useful in the present invention.

[0199] Truncated forms of these receptors, comprising the extracellular domains of the receptors or functional portions thereof, are highly preferred TNF inhibitors, according to the present invention. Truncated soluble TNF type-I and type-II receptors are described in EP914431, for example.

[0200] The IL-18 inhibitor can be used simultaneously, sequentially or separately with the TNF inhibitor.

[0201] Preferably, a combination of an IL-18 antibody or antiserum and a soluble receptor of TNF, having TNF inhibiting activity, is used.

[0202] The present invention is further based on the finding of a beneficial effect of an IL-18 inhibitor in arthritis. The invention therefore further relates to the use of an IL-18 inhibitor for the manufacture of a medicament for the treatment and/or prevention of arthritis.

[0203] According to the present invention, the term “arthritis” includes all different types of arthritis and arthritic conditions, both acute and chronic arthritis, as defined for example in the Homepage of the Department of Orthopaedics of the University of Washington on Arthritis. Examples for arthritic conditions are ankylosing spondylitis, back pain, carpal deposition syndrome, Ehlers-Danlos-Syndrome, gout, juvenile arthritis, lupus erythematosus, myositis, osteogenesis imperfecta, osteoporosis, polyarthritis, polymyositis, psoriatic arthritis, Reiter’s syndrome, scleroderma, arthritis with bowel disease, Behcets’s disease, children’s arthritis, degenerative joint disease, fibromyalgia, infectious arthritis, Lyme disease, Marfan syndrome, osteoarthritis, osteonecrosis, Pagets Disease, Polymyalgia rheumatica, pseudogout, reflex sympathetic dystrophy, rheumatoid arthritis, rheumatism, Sjogren’s syndrome, familial adenomatous polyposis and the like.

[0204] Preferably, according to the invention, inhibitors of IL-18 are provided for treatment and/or prevention of inflammatory arthritis. Inflammatory arthritis is classified as a chronic arthritis, according to the persistent, continuous or recurring course of the disease.

[0205] In a preferred embodiment of the invention, the inflammatory arthritis is rheumatoid arthritis (RA). RA causes inflammation in the lining of the joints (the synovial membrane, a one cell layer epithelium) and/or internal organs. The disease tends to persist for many years, typically affects many different joints throughout the body and ultimately can cause damage to cartilage, bone, tendons, and ligaments. The joints that may be affected by RA are the joints located in the neck, shoulders, elbows, hips, wrists, hands, knees, ankles and feet, for example. In many cases, the joints are inflamed in a symmetrical pattern in RA.

[0206] RA is prevalent in about 1% of the population in the United States, being distributed within all ethnic groups and ages. It occurs all over the world, and women outnumber men by 3 to 1 among those having RA.

[0207] As shown in the examples below, an inhibitor of IL-18 has been proven to exhibit a highly efficacious beneficial effect on cartilage erosion. The invention therefore further relates to the use of an inhibitor of IL-18 in the manufacture of a medicament for treatment and/or prevention of cartilage destruction, i.e. to the use of an IL-18 inhibitor as a chondroprotective agent. The IL-18 inhibitor may be used in any condition in which cartilage destruction or erosion occurs. Cartilage destruction is the progressive decline in the structural integrity of joint articular cartilage. It occurs for example in conditions affecting articular cartilage such as rheumatoid arthritis, juvenile rheumatoid arthritis, or osteoarthritis, but also in infectious synovitis, for instance.

[0208] In a preferred embodiment, the inhibitor of IL-18 is selected from inhibitors of caspase-1 (ICE), antibodies directed against IL-18, antibodies directed against any of the IL-18 receptor subunits, inhibitors of the IL-18 signalling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, and IL-18 binding proteins, isoforms, muteins, fused proteins, functional derivatives, active fractions or circularly permuted derivatives thereof having the same activity.

[0209] The term “IL-18 binding proteins” is used herein synonymously with “IL18BP”. It comprises IL-18 binding proteins as defined in detail above, including splice variants and/or isoforms of IL-18 binding proteins. In particular, human isoforms a and c of IL-18BP are useful in accordance

with the present invention. The proteins useful according to the present invention may be glycosylated or non-glycosylated, they may be derived from natural sources, like urine, or produced recombinantly. Recombinant expression may be carried out in prokaryotic expression systems like *E. coli*, or in eukaryotic, preferably in mammalian, expression systems.

[0210] The terms “muteins”, “functional derivatives”, “active fractions” and “circularly permuted derivatives” have been defined above.

[0211] Functional derivatives of IL-18BP may be conjugated to polymers in order to improve the properties of the protein, such as the stability, half-life, bioavailability, tolerance by the human body, or immunogenicity. To achieve this goal, IL18-BP may be linked e.g. to Polyethylenglycol (PEG). PEGylation may be carried out by known methods, described in WO 92/13095, for example.

[0212] The term “fused protein” refers to a polypeptide comprising an IL-18BP, or a viral IL-18BP, or a mutein thereof, fused with another protein, which, e.g., has an extended residence time in body fluids. An IL-18BP or a viral IL-18BP, may thus be fused to another protein, polypeptide or the like, e.g., an immunoglobulin or a fragment thereof. The fusion may be direct, or via a short linker peptide which can be as short as 1 to 3 amino acid residues in length or longer, for example, 13 amino acid residues in length. Said linker may be a tripeptide of the sequence E-F-M (Glu-Phe-Met), for example, or a 13-amino acid linker sequence comprising Glu-Phe-Gly-Ala-Gly-Leu-Val-Leu-Gly-Gly-Gln-Phe-Met (SEQ ID NO: 14) introduced between the IL-18BP sequence and the immunoglobulin sequence. Fusion proteins comprising IL-18BP fused to all or part of an immunoglobulin are highly preferred. For example, IL-18BP may be fused to the constant regions of Ig molecules, preferably to heavy chain regions, like the CH2 and CH3 domains of human IgG1, for example. The generation of specific fusion proteins comprising IL-18BP and a portion of an immunoglobulin are described in example 11 of WP99/09063. Other isoforms of Ig molecules are also suitable for the generation of fusion proteins according to the present invention, such as isoforms IgG₂ or IgG₄ or other Ig classes, like IgM or IgA, for example. Fusion proteins may be monomeric or multimeric, hetero- or homomultimeric.

[0213] In a further preferred embodiment of the invention, the inhibitor of IL-18 is an IL-18 antibody. Anti-IL-18 antibodies may be polyclonal or monoclonal, chimeric, fully humanised, or even fully human. Recombinant antibodies and fragments thereof are characterised by high affinity binding to IL-18 in vivo and low toxicity. The antibodies which can be used in the invention are characterised by their ability to treat patients for a period sufficient to have good to excellent regression or alleviation of the pathogenic condition or any symptom or group of symptoms related to a pathogenic condition, and a low toxicity.

[0214] Neutralising antibodies are readily raised in animals such as rabbits, goat or mice by immunisation with IL-18. Immunised mice are particularly useful for providing sources of B cells for the manufacture of hybridomas, which in turn are cultured to produce large quantities of anti-IL-18 monoclonal antibodies.

[0215] Chimeric antibodies are immunoglobulin molecules characterised by two or more segments or portions derived from different animal species. Generally, the variable region of the chimeric antibody is derived from a non-human mammalian antibody, such as murine monoclonal antibody,

and the immunoglobulin constant region is derived from a human immunoglobulin molecule. Preferably, both regions and the combination have low immunogenicity as routinely determined (Elliott, M. J., Maini, R. N., Feldmann, M., Long-Fox, A., Charles, P., Bijl, H., and Woody, J. N., 1994). Humanised antibodies are immunoglobulin molecules created by genetic engineering techniques in which the murine constant regions are replaced with human counterparts while retaining the murine antigen binding regions. The resulting mouse-human chimeric antibody preferably have reduced immunogenicity and improved pharmacokinetics in humans (Knight, D. M., Trinh, H., Le, J., Siegel, S., Shealy, D., McDonough, M., Scallon, B., Moore, M. A., Vilcek, J., and Daddona, P., 1993).

[0216] Thus, in a further preferred embodiment, IL-18 antibody is a humanised IL-18 antibody. Preferred examples of humanized anti-IL-18 antibodies are described in the European Patent Application EP 0 974 600, for example.

[0217] In yet a further preferred embodiment, the IL-18 antibody is fully human. The technology for producing human antibodies is described in detail e.g. in WO00/76310, WO99/53049, U.S. Pat. No. 6,162,963 or AU5336100. Fully human antibodies are preferably recombinant antibodies, produced in transgenic animals, e.g. xenomice, comprising all or parts of functional human Ig loci.

[0218] In a highly preferred embodiment of the present invention, the inhibitor of IL-18 is a IL-18BP, or an isoform, a mutein, fused protein, functional derivative, active fraction or circularly permuted derivative thereof essentially having at least an activity similar to IL-18BP. Preferred active fractions have an activity which is better than the activity of IL-18BP or have further advantages, like a better stability or a lower toxicity or immunogenicity, or they are easier to produce in large quantities.

[0219] Interferons are predominantly known for inhibitory effects on viral replication and cellular proliferation. Interferon γ , for example, plays an important role in promoting immune and inflammatory responses. Interferon β (IFN- β , an interferon type I), is said to play an anti-inflammatory role. Studies published by Triantaphyllopoulos et al. (1999) indicated that IFN- β has a beneficial effect in the therapy of rheumatoid arthritis, as shown in a mouse model of the disease, the collagen-induced arthritis (CIA) model. This beneficial effect of IFN- β was confirmed in the examples below.

[0220] The invention also relates to the use of a combination of an inhibitor of IL-18 production and/or action and an interferon in the manufacture of a medicament for the treatment of arthritis, in particular rheumatoid arthritis.

[0221] Interferons may be conjugated to polymers in order to improve the stability of the proteins. A conjugate between Interferon β and the polyol Polyethyleneglycol (PEG) has been described in WO99/55377, for instance.

[0222] The inhibitor of IL-18 production and/or action is preferably used simultaneously, sequentially, or separately with the interferon.

[0223] In another preferred embodiment of the invention, the interferon is Interferon- β (IFN- β), more preferably IFN- β 1a.

[0224] The invention further relates to the use of a combination of IL-18 inhibitors, interferons and TNF antagonists. The combination is suitable for the treatment and/or prevention of arthritis, in particular rheumatoid arthritis, and

for the treatment and/or prevention of liver injury. The active components may be used simultaneously, sequentially, or separately.

[0225] The invention further relates to the use of an expression vector comprising a coding sequence of an inhibitor of IL-18 in the preparation of a medicament for the prevention and/or treatment of arthritic conditions or arthritis, in particular rheumatoid arthritis, or for the treatment of liver injury. A gene therapeutical approach is thus used for treating and/or preventing the disease. Advantageously, the expression of the IL-18 inhibitor will then be in situ, thus efficiently blocking IL-18 directly in the tissue(s) or cells affected by the disease.

[0226] In order to treat and/or prevent arthritis, the gene therapy vector comprising the sequence of an inhibitor of IL-18 production and/or action may be injected directly into the diseased joint, for example, thus avoiding problems usually involved in systemic administration of gene therapy vectors, like dilution of the vectors, reaching of the target cells and side effects.

[0227] The use of a vector for inducing and/or enhancing the endogenous production of an inhibitor of IL-18 in a cell normally silent for expression of an IL-18 inhibitor, or which expresses amounts of the inhibitor which are not sufficient are also contemplated according to the invention. The vector may comprise regulatory sequences functional in the cells desired to express the inhibitor or IL-18. Such regulatory sequences may be promoters or enhancers, for example. The regulatory sequence may then be introduced into the right locus of the genome by homologous recombination, thus operably linking the regulatory sequence with the gene, the expression of which is required to be induced or enhanced. The technology is usually referred to as endogenous gene activation (EGA), and it is described e.g. in WO 91/09955. It will be understood by the person skilled in the art that it is also possible to shut down IL-18 expression using this technique, by introducing a negative regulation element, like e.g. a silencing element, into the gene locus of IL-18, thus leading to down-regulation or prevention of IL-18 expression. The person skilled in the art will understand that such down-regulation or silencing of IL-18 expression has the same effect as using an IL-18 inhibitor in order to prevent and/or treat disease.

[0228] The invention further relates to pharmaceutical compositions for prevention and/or treatment of inflammatory arthritis or liver injury, comprising an inhibitor of IL-18. The composition may comprise antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signalling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, and IL-18 binding proteins, isoforms, muteins, fused proteins, functional derivatives, active fractions or circularly permuted derivatives thereof having the same activity.

[0229] IL-18BP and its isoforms, muteins, fused proteins, functional derivatives, active fractions or circularly permuted derivatives as described above are the preferred active ingredients of the pharmaceutical compositions.

[0230] In a further preferred embodiment, the pharmaceutical composition further comprises an interferon, preferably IFN- β .

[0231] In yet another preferred embodiment, the pharmaceutical composition comprises a TNF antagonist. Advantageously, anti-TNF antibodies neutralising TNF action are used. Highly preferred are TBPI and TPBII as TNF antagonists.

[0232] The active ingredients of the pharmaceutical composition according to the invention, i.e. the IL-18 inhibitor and/or interferon and/or TNF antagonist can be administered to an individual in a variety of ways. The routes of administration include intradermal, transdermal (e.g. in slow release formulations), intramuscular, intraperitoneal, intravenous, subcutaneous, oral, epidural, topical, and intranasal routes. Any other therapeutically efficacious route of administration can be used, for example absorption through epithelial or endothelial tissues or by gene therapy wherein a DNA molecule encoding the active agent is administered to the patient (e.g. via a vector) which causes the active agent to be expressed and secreted in vivo. In addition, the protein(s) according to the invention can be administered together with other components of biologically active agents such as pharmaceutically acceptable surfactants, excipients, carriers, diluents and vehicles.

[0233] In another aspect, the present invention relates to the administration of IL-18 inhibitors to prevent or treat sepsis and is based on the findings described in the examples. It was found in according with the present invention that the levels of effective (free) circulating IL-18 are significantly elevated in serum of sepsis patients in comparison to healthy individuals, and that the inhibition of IL-18 causes a decrease in IFN- γ induction by the Gram-positive bacterium *Staphylococcus epidermidis*.

[0234] Sepsis according to the present invention, comprises SIRS, severe sepsis, septic shock, endotoxic shock etc., and may be caused by Gram-positive or Gram-negative bacteria.

[0235] In addition the prevention/treatment of sepsis can be approached by administration of an IL-18 inhibitor in combination of other cytokine inhibitors potentially able to increase its effect.

[0236] Other cytokine inhibitors can be used together with inhibitors of IL-18 in the treatment of sepsis. An inhibitor of IL-18 may be used in combination with an IL-12 Inhibitor.

[0237] The term "inhibition of IL-12" within the context of this invention refers to any molecule modulating IL-12 production and/or action in such a way that IL-12 production and/or action is attenuated, reduced, or partially, substantially or completely prevented or blocked.

[0238] An inhibitor of IL-12 production can be any molecule negatively affecting its synthesis e.g. interferon α/β (Karp et al. 2000), or molecules negatively affecting processing or maturation of IL-12. The inhibitors, considered according to the invention can be, for example, suppressors of gene expression of the interleukin IL-12, antisense mRNAs reducing or preventing the transcription of the IL-12 mRNA or ribozymes leading to degradation of the IL-12 mRNA, proteins impairing correct folding, or partially or substantially preventing secretion of IL-12, proteases degrading IL-12 and the like.

[0239] IL-12 antagonists exert their activity in several ways. Antagonists can bind to or sequester the IL-12 molecule itself with sufficient affinity and specificity to partially or substantially neutralize the IL-12 epitope or epitopes responsible for IL-12 receptor binding (hereinafter termed "sequestering antagonists"). A sequestering antagonist may be, for example, an antibody directed against IL-12, a truncated form of IL-12 receptor, comprising the extracellular domains of the receptor or functional portions thereof etc. An antagonist may also inhibit the IL-12 signaling pathway, which is activated within the cells upon IL-12/receptor binding (hereinafter termed "signalling antagonists"). All groups

of antagonists are useful, either alone or together, in combination with an IL-18 inhibitor, in the therapy of sepsis. IL-12 antagonists are easily identified and evaluated by routine screening of candidates for their effect on the activity of native IL-12 on susceptible cell lines in vitro. For example mouse splenocytes in which phorbol ester and IL-12 causes proliferation. The assay contains IL-12 formulation at varying dilutions of candidate antagonist, e.g., from 0.1 to 100 times the molar amount of IL-12 used in the assay, and controls with no IL-12, antagonist only or phorbol ester and IL-2 (Tucci et al., 1992).

[0240] Sequestering antagonists are the preferred IL-12 antagonists to be used according to the present invention. Amongst sequestering antagonists, antibodies that neutralize IL-12 activity are preferred. The simultaneous, sequential, or separate use of the IL-18 inhibitor with the IL-12 antagonist is preferred, according to the invention. Anti IL-12 is the preferred IL-12 antagonist to be used in combination with an IL-18 inhibitor as well as antibody derivatives, fragments, regions and biologically active portions of the antibody. The IL-18 inhibitor can be used simultaneously, sequentially or separately with the IL-12 inhibitor.

[0241] In addition, an inhibitor of IL-18 may be used in combination with inhibitors of other cytokines, known to play an important role in septic shock e.g. IL-1, TNF, IL-8 etc Dinarello 1996 and Okusawa et al. 1988. An example for an inhibitor of IL-1 is IL1-receptor antagonist, and for TNF the soluble portion of receptors TNFR1 and TNFR2.

[0242] The invention further relates to the use of an expression vector comprising the coding sequence of an inhibitor of IL-18 in the preparation of a medicament for the prevention and/or treatment of sepsis. A gene therapeutical approach is thus used for treating and/or preventing the disease. Advantageously, the expression of the IL-18 inhibitor will then be in situ, thus efficiently blocking IL-18 directly in the tissue(s) or cells affected by the disease.

[0243] The use of a vector for inducing and/or enhancing the endogenous production of an inhibitor of IL-18 in a cell normally silent for expression of an IL-18 inhibitor, or which expresses amounts of the inhibitor which are not sufficient, are also contemplated according to the invention. The vector may comprise regulatory sequences functional in the cells desired to express the inhibitor or IL-18. Such regulatory sequences may be promoters or enhancers, for example. The regulatory sequence may then be introduced into the right locus of the genome by homologous recombination, thus operably linking the regulatory sequence with the gene, the expression of which is required to be induced or enhanced. The technology is usually referred to as "endogenous gene activation" (EGA), and it is described, e.g., in WO 91/09955.

[0244] The invention comprise also the administration of genetically modified cells, able to produce an inhibitor of IL-18 for the treatment or prevention of sepsis.

[0245] It will be understood by the person skilled in the art that it is also possible to shut down IL-18 expression using the same technique, i.e., by introducing a negative regulation element, e.g., a silencing element, into the gene locus of IL-18, thus leading to down-regulation or prevention of IL-18 expression. The person skilled in the art will understand that such down-regulation or silencing of IL-18 expression has the same effect as the use of an IL-18 inhibitor in order to prevent and/or treat disease.

[0246] The definition of "pharmaceutically acceptable" is meant to encompass any carrier, which does not interfere with

effectiveness of the biological activity of the active ingredient and that is not toxic to the host to which it is administered. For example, for parenteral administration, the active protein(s) may be formulated in a unit dosage form for injection in vehicles such as saline, dextrose solution, serum albumin and Ringer's solution.

[0247] For parenteral (e.g. intravenous, subcutaneous, intramuscular) administration, the active protein(s) can be formulated as a solution, suspension, emulsion or lyophilised powder in association with a pharmaceutically acceptable parenteral vehicle (e.g. water, saline, dextrose solution) and additives that maintain isotonicity (e.g. mannitol) or chemical stability (e.g. preservatives and buffers). The formulation is sterilized by commonly used techniques.

[0248] The bioavailability of the active protein(s) according to the invention can also be ameliorated by using conjugation procedures which increase the half-life of the molecule in the human body, for example linking the molecule to polyethylenglycol, as described in the PCT Patent Application WO 92/13095.

[0249] The therapeutically effective amounts of the active protein(s) will be a function of many variables, including the type of antagonist, the affinity of the antagonist for IL-18, any residual cytotoxic activity exhibited by the antagonists, the route of administration, the clinical condition of the patient (including the desirability of maintaining a non-toxic level of endogenous IL-18 activity

[0250] A "therapeutically effective amount" is such that when administered, the IL-18 inhibitor results in inhibition of the biological activity of IL-18. The dosage administered, as single or multiple doses, to an individual will vary depending upon a variety of factors, including IL-18 inhibitor pharmacokinetic properties, the route of administration, patient conditions and characteristics (sex, age, body weight, health, size), extent of symptoms, concurrent treatments, frequency of treatment and the effect desired. Adjustment and manipulation of established dosage ranges are well within the ability of those skilled in the art, as well as in vitro and in vivo methods of determining the inhibition of IL-18 in an individual.

[0251] According to the invention, the inhibitor of IL-18 is used in an amount ranging between about 0.1 to 5000 µg/kg of body weight or about 1 to 1000 µg/kg of body weight. Amounts of about 10 to 500 µg/kg of body weight are preferred, and amounts of about 50 to 250 µg/kg of body weight are further preferred.

[0252] Inhibitors of IL-18 may also be used in an amount of about 1 to 50 µg/kg of body weight.

[0253] The route of administration which is preferred according to the invention is administration by subcutaneous route. Intramuscular administration is further preferred according to the invention.

[0254] In further preferred embodiments, the inhibitor of IL-18 is administered daily or every other day.

[0255] The daily doses are usually given in divided doses or in sustained release form effective to obtain the desired results. Second or subsequent administrations can be performed at a dosage which is the same, less than or greater than the initial or previous dose administered to the individual. A second or subsequent administration can be administered during or prior to onset of the disease.

[0256] According to the invention, the IL-18 inhibitor can be administered prophylactically or therapeutically to an individual prior to, simultaneously or sequentially with other

therapeutic regimens or agents (e.g. multiple drug regimens), in a therapeutically effective amount, in particular with an interferon and/or a TNF antagonist. Active agents that are administered simultaneously with other therapeutic agents can be administered in the same or different compositions. Other agents which may be used in combination with IL-18 inhibitors, in particular with IL-18BP, are COX-2 inhibitors. COX-2 inhibitors are known in the art. Specific COX-2 inhibitors are disclosed in WO 01/00229, for example.

[0257] The invention further relates to a method for the preparation of a pharmaceutical composition comprising admixing an effective amount of an IL-18 inhibitor with a pharmaceutically acceptable carrier.

[0258] Having now described the invention, it will be more readily understood by reference to the following examples that are provided by way of illustration and are not intended to be limiting of the present invention.

EXAMPLES

PART I: Examples 1 to 17

Isolation, Production and Biological Activities of IL-18BP

Example 1: Isolation of an IL-18 Binding Protein

[0259] *E. coli* IL-18 (2.5 mg, Peprotech, N.J.) was coupled to Affigel-10 (0.5 ml, BioRad), according to the manufacturer's instructions and packed into a column. Crude urinary proteins (1000-fold concentrated, 500 ml) were loaded onto the column at a flow rate of 0.25 ml/min. The column was washed with 250 ml 0.5M NaCl in phosphate buffered saline (PBS). Bound proteins were then eluted with 25 mM citric acid, pH 2.2 and benzamidine (1 mM), immediately neutralized by 1M Na₂CO₃. Fractions of 1 ml were collected. The fractions were analyzed by SDS-PAGE and silver staining. The IL-18 binding protein eluted in fractions 2-8 as a ~40,000 Dalton protein (FIG. 1). The ~40 kD band, corresponding to the IL-18BP exhibited a distinct yellow color upon silver staining. The various fractions were analyzed by cross-linking with ¹²⁵I-IL-18, SDS-PAGE and autoradiography as described in Example 2. An IL-18 binding protein was found in fractions 2-8, eluted from the IL-18-agarose column (FIG. 2).

Example 2: Cross-Linking of Affinity-Purified IL-18BP to Labeled IL-18

[0260] Samples (40 µl) of IL-18BP from the affinity purification step were incubated (70 min. at 4° C.) with ¹²⁵I-IL-18 (5,000,000 cpm). Disuccinimidyl suberate (DSS), dissolved in dimethyl sulfoxide (Me₂SO, 20 mM), was then added to a final concentration of 2 mM and the mixture was left for 20 min. at 4° C. The reaction was stopped by the addition of 1M Tris-HCl pH 7.5, and 1M NaCl to a final concentration of 100 mM. A sample buffer containing Dithiothreitol (DTT, 25 mM final) was added and the mixtures were analyzed by SDS-PAGE (7.5% acrylamide) followed by autoradiography (FIG. 2).

[0261] A specific band of molecular weight 58 kD, probably consisting of a ~40 kD protein cross-linked to the ~20 kD ¹²⁵I-IL-18, was observed in fractions eluted from the IL-18

affinity column (lanes 2 and 3) but not in the column wash (lane 1), containing all other crude urinary proteins.

Example 3: Protein Sequence Analysis

[0262] Eluted fractions from the affinity column of Example 1 were resolved by SDS-PAGE (10% acrylamide) under non-reducing conditions, electroblotted on a PVDF membrane (Pro-Blot, Applied Biosystems, USA). The membrane was stained with coomassie blue, the ~40 kD band was excised and subjected to protein sequence analysis by a Pro-

prudent not to assign a specific amino acid residue at said cycles.

[0268] The resulting sequence is significantly different from that of any other known protein, as determined by searching protein databases. However, searching the TIGR database by the tblastn search program provided a cDNA file, denoted THC123801, whose open reading frame (218 codons), when translated contains a sequence highly homologous to that of the N-terminal sequence of IL-18BP. The homology is hereby shown:

```

1 . . . . . TPVSQXXXAAXASVRSTKDCPSQPPVFPAKQCPALEVT. 40 (SEQ ID NO: 12)
      | | | | |
51 VTLLVRATXVXQTTAATASVRSTKDCPSQPPVFPAKQCPALEVTWPE 100 (SEQ ID NO: 13)

```

cise microsequencer (Applied Biosystems, USA). The following major sequence was obtained:

T-P-V-S-Q-Q-x-x-x-A-A-A (SEQ ID NO: 11)

1 . . . 5 10 . .

[0263] wherein x represents a yet undetermined amino acid.

[0264] In addition, a minor sequence was obtained:

A-x-Y-x-R-I-P-A-x-A-I-A (SEQ ID NO: 15)

1 . . . 5 10 . .

[0265] Because of this double sequence it was not possible to obtain a longer sequence data. The minor sequence was identified as a fragment of human defensin, (accession No. p11398) starting at amino acid 65 of defensin. The major sequence could not be associated with any other known protein, as determined by searching all available databases in NCBI and TIGR by the blastp and tblastn search programs.

[0266] In order to obtain a longer and more accurate sequence and in order to identify potential cysteine residues, another aliquot of the fraction eluted from the IL-18-agarose column was reduced with DTT in 6 M guanidine HCl, reacted with 4-vinyl pyridine, desalted by a micro-ultrafiltration device (Ultrafree, cutoff 10,000 Daltons, Millipore) and subjected to protein microsequence analysis. After cycle No. 1 of sequencing, the filter was reacted with o-phthalaldehyde to block all N-terminal polypeptides other than Pro. In this way only the major sequence was obtained as follows:

```

                                (SEQ ID NO: 10)
TPVSQXXXAA XASVRSTKDP CPSQPPVFPA AKQCPALEVT
1          10          20          30          40
(T = Thr; P = Pro; V = Val; S = Ser; Q = Gln; X =
Unknown; A = Ala; R = Arg; K = Lys; D = Asp; C =
Cys; F = Phe; L = Leu; E = Glu)

```

[0267] In cycles 6,7,8 and 11 a low level of a Thr signal was obtained. Because of this low level we considered it more

[0269] (The upper sequence (1-40) (SEQ ID NO: 12) is that of the IL-18BP isolated according to the invention; the lower sequence (51-100) (SEQ ID NO: 13) is deduced by translation of the cDNA of TIGR file THC123801).

[0270] The putative protein sequence, obtained by translating file THC123801, was ambiguous at residues 2 and 4 of the IL-18BP. It confirms the identity of amino acid residues 6, 7 and 8 of IL-18BP as Thr and seems to do so also for residue 11.

Example 4: The IL-18BP is a Glycoprotein

[0271] Aliquot (0.3 ml) of eluted fractions of Example 1 were further purified by size exclusion chromatography on a Superose 12 column (1×30 cm, Pharmacia, Sweden). The column was pre-equilibrated and eluted with phosphate buffered saline and sodium azide (0.02%) at a flow rate of 0.5 ml/min. and fractions of 1 min. were collected. The IL-18 binding protein eluted in fractions 20-25 as a ~40,000 Dalton protein, as determined by SDS-PAGE and silver staining. A sample containing the ~40,000 Dalton protein (fraction 23, 50 µl, ~50 ng protein) was reacted with N-glycosidase F (PNGase F, Biolab) according to the manufacturers instructions. Briefly, the aliquot was denatured by boiling in the presence of 5% SDS for 10 min., 10×G7 buffer (2.5 ?l), 10% NP-40 (2.5 µl) and PNGase F (1 µl), 1 h at 37° C. The sample was analyzed by SDS-PAGE (10% acrylamide) under non-reducing conditions and compared with undigested IL-18BP from the same Superose 12 fraction. It was found that the ~40 kD band of IL-18BP disappeared in the PNGase-treated fraction. New bands, corresponding to 30 kD (just above the PNGase band) and 20 kD were obtained. The elimination of the ~40 kD band indicates that this band is an N-glycosylated protein.

Example 5: Blocking of the Biological Activity of IL-18 by IL-18BP

[0272] The ability of the IL-18BP isolated from urine to block IL-18 activity was determined by measuring the IL-18-induced production of IFN-γ in mononuclear cells. IL-18 induces IFN-γ when added together with either low concentration of LPS, IL-12, IL-2, or other stimulants. The activity of IL-18 was tested in murine splenocytes, in human peripheral blood mononuclear cells (PBMC) and in the human KG-1 cell line. Spleen cells were prepared from a healthy mouse, washed and suspended in RPMI 1640 medium supplemented with 10% fetal bovine serum at 5×10⁶ cells/ml.

1.0 ml cultures were stimulated with LPS (either 0.5 or 1 µg/ml) together with recombinant human or murine IL-18 (either 0.5 or 5 ng/ml). Human IL-18 binding protein (0.5 or 50 ng/ml) was added to the recombinant IL-18 before adding to spleen cells. After culturing for 24 h, the spleen cells were subjected to three freeze (−70° C.) and thaw (room temperature) cycles, the cellular debris was removed by centrifugation and the supernatants were assayed for IFN-γ using the ELISA kits for mouse IFN-γ (Endogen). As shown in FIG. 3A, IL-18BP blocked the activity of huIL-18 in murine splenocytes in a dose-dependent manner. In contrast, as a control, soluble interferon-α/6 receptor had no effect. The activity of recombinant murine IL-18 was similarly inhibited by the human IL-18BP, suggesting that human IL-18BP recognizes murine IL-18 (FIG. 3B). Endogenous IL-18 is induced in murine splenocytes by high concentrations of LPS, leading to production of IFN-γ. Indeed, IFN-γ induction by LPS (10 µg/ml) was also inhibited by the urinary IL-18BP (FIG. 3C). Concanavalin A (con A) activates T-cells to produce IFN-γ in the absence of IL-18 (Fantuzzi et al., 1998). Indeed, induction of IFN-γ by Con A was not inhibited by IL-18BP even at high concentrations (FIG. 3D). This observation demonstrated that IL-18BP was a specific inhibitor of IL-18 bioactivity rather than a non-specific inhibitor of IFN-γ production. IL-18BP also inhibited the activity of human IL-18 in human KG-1 cells induced by a combination of IL-18 and TNF-α (FIG. 3E).

[0273] The above data demonstrate that urinary IL-18BP inhibits human as well as murine IL-18 activity as measured by co-induction of IFN-γ in human and murine mononuclear cells. The concentration of IL-18BP which reduced IL-18 activity by >90% was comparable to that of IL-18 itself, suggesting a high affinity interaction between these two proteins.

Example 6: Isolation of cDNA Clones Coding for IL-18BP

[0274] Total RNA from Jurkat T-cells (CRL 8163, American Type Culture Collection) was reverse-transcribed with SuperScript RNase H[−] reverse transcriptase (Gibco-BRL) and random primers (Promega, Madison Wis.). The resulting cDNA fragments were then amplified by PCR, using Taq DNA polymerase (Sigma) and primers corresponding to TIGR clone THC123801 nucleotides 24-44 (sense) and 500-481 (reverse). The amplification was done in 30 cycles of annealing (55° C., 2 min) and extension (70° C., 1 min). The resulting PCR products were resolved by agarose (1%) gel electrophoresis, eluted and cloned into pGEM-Teasy TA cloning vector (Promega). DNA from individual clones was sequenced with T7 and SP6 primers.

[0275] The resulting 477 bp fragment was ³²P-labeled by random priming. This probe was used for screening various human cDNA and genomic libraries. Duplicate nitrocellulose filters were lifted and hybridized with the probe at 60° C. in a buffer consisting of 6×SSC, 10×Denhardt's solution, 0.1% SDS and 100 µg/ml Salmon sperm DNA. The filters were washed and exposed overnight at −80° C. to Kodak XAR film. Double positive clones were plaque-purified. Plasmids were excised from the XpCEV9 clones and self-ligated. cDNA clones from other libraries were isolated according to the manufacturer's instructions. Automated DNA sequence analysis of the isolated clones was performed with Models 373A and 377 sequencers (Applied Biosystems) using sense

and antisense primers. Standard protocols were used for these cloning procedures (Sambrook et al., 1989).

[0276] The following libraries were screened: a human monocyte cDNA library, constructed in λpCEV9 cloning vector (Gutkind, 1991), kindly provided by T. Miki; a human Jurkat leukemic T-cell cDNA library, a human peripheral blood leukocyte cDNA library and a human spleen cDNA library, all from Clontech (Palo Alto, Calif.). A human placenta genomic library in lambda FIX II vector was from Stratagene (La Jolla, Calif.).

[0277] All cDNA clones corresponded to four different IL-18BP splice variants were obtained and characterized. All splice variants coded for putative soluble secreted proteins. The most abundant one (IL-18BP_a) had an open reading frame of 192 codons, coding for a signal peptide of 28 amino acid residues followed by a mature putative IL-18BP_a, whose first 40 residues (SEQ ID NO:10) matched perfectly with the N-terminal protein sequence of the urinary IL-18BP (SEQ ID NO:2). The position of the cysteine residues suggested that this polypeptide belongs to the immunoglobulin (Ig) superfamily. Each of the four Gln residues within mature IL-18BP_a was a potential N-glycosylation site. The other three splice variants of IL-18BP were significantly less abundant.

[0278] Another 1 kb IL-18BP_b cDNA coded for a mature protein of 85 amino acid residues (SEQ ID NO:4). A third variant, IL-18BP_c, was represented by a 2.3 kb cDNA, coding for a mature IL-18BP of 169 amino acid residues (SEQ ID NO:6). The fourth variant, IL-18BP_d coded for a mature IL-18BP of 133 amino acid residues (SEQ ID NO:8). In-exon splicing occurred at two sites along the pro-mRNA. These events and an additional 5' exon in IL-18BP_d gave rise to 3 different 5' UTRs in the various cDNA clones. It is therefore quite possible that different IL-18BP variants may be generated in response to distinct transcription regulation signals.

[0279] No cDNA coding for a receptor with a transmembrane domain was found so far.

Example 7: Construction of a Mammalian Expression Vector, Production of Recombinant IL-18BP, and Evaluation of the Biological Activities of Recombinant IL-18BP

[0280] The coding region of the IL-18BP_a cDNA was amplified by PCR with the sense primer

(SEQ ID NO: 16)
5' TATATCTAGAGCCACCATGAGACACAACCTGGACACCA

[0281] and the reverse primer:

(SEQ ID NO: 17)
5' ATATCTAGATTAATGATGATGATGATGATGACCCCTGCTGTGGAC
TGC

[0282] The PCR products were cut with Xba I and cloned into the Xba I site of the pEF-BOS expression vector (25), to yield pEF-BOS-IL-18BP_a. The constructs were confirmed by DNA sequencing.

[0283] Batches of 6×10⁷ COS7 cells in 1.4 ml TD buffer, containing pEF-BOS-IL-18BP_a plasmid DNA (10 µg) and DEAE-dextran (120 µg) were incubated for 30 min at room temperature, as described (Sampayrat and Dana, 1981). The cells were then washed with DMEM-10% FBS, plated for 4 hr in DMEM-10, washed and incubated for 3-5 days in

serum-free DMEM. Culture medium was collected, concentrated 6-fold by ultrafiltration (10 kD cutoff) and the IL-18BP-His₆ was isolated on a Talon column (Clontech) with imidazole as eluant according to the manufacturer's instructions.

[0284] Immunological cross-reactivity of the urinary and the COS7-expressed IL-18BP was assessed as follows: Urinary IL-18BP (5 µg) was labeled with ¹²⁵I by the chloramine T procedure. Supernatants of COS7 cells (250 µl) were mixed (1 h, room temperature final volume 500 µl) with the antibody to urinary IL-18BP, diluted 1:1000 in phosphate-buffered saline (PBS), 0.05% Tween 20 and 0.5% bovine serum albumin (Wash Buffer). ¹²⁵I-labeled urinary IL-18BP (10⁶ cpm) was then added and after 1 h protein G-sepharose (20 µl) was added. The mixture was suspended (1.5 h, 4° C.), the beads were then isolated and washed wash 3× Wash Buffer and once in PBS. The beads were then eluted with a Sample buffer, resolved by SDS-PAGE (10% acrylamide under reducing conditions followed by Autoradiography).

[0285] IL-18BP_a ran as a single band upon SDS-PAGE with silver staining under reducing and non-reducing conditions and had the same apparent molecular mass as that of the urinary IL-18BP (data not shown). Protein sequence analysis of this preparation revealed the same N-terminal sequence as that of the urinary IL-18BP, indicating that the latter was not degraded at its N-terminus.

[0286] Immunoblot analysis of IL-18BP_a with antibodies raised against the urinary IL-18BP revealed the same molecular mass band as that of the urinary protein. Furthermore, using immunoprecipitation followed by SDS-PAGE and autoradiography, IL-18BP_a was able to displace urinary ¹²⁵I-IL-18BP from binding to the antibody. Therefore, IL-18BP_a corresponds structurally to the urinary IL-18BP.

[0287] Crude and purified IL-18BP_a was tested for its ability to inhibit the biological activity of IL-18. IL-18BP_a inhibited in a dose dependent manner the IFN-γ inducing activity of human and mouse IL-18 in murine splenocytes, PBMC and the human KG-1 cell line (FIG. 4).

[0288] The results of the various bioassays as well as the mobility shift assay (Example 8) demonstrated that inhibition of IL-18 activity is an intrinsic property of the cloned IL-18BP and not that of any accompanying impurities in urinary IL-18BP, such as the co-eluting fragment of defensin.

Example 8: Electrophoretic Mobility Shift Assays

[0289] The effect of the urinary and the recombinant IL-18BP on IL-18-induced activation of NF-κB in human KG-1 cells was also studied. Human KG-1 cells (4×10⁶ in 1 ml RMPI) were stimulated with either huIL-18 (10 ng/ml) or huIL-18 pre-mixed with an IL-18BP (20 min, room temperature). After 20 min at 37° C., cells were washed three times with ice-cold PBS and immediately frozen in liquid nitrogen. Cell pellets were resuspended in three times the packed cell volume in buffer A (20 mM Tris pH 7.6, 0.4M NaCl, 0.2 mM EDTA, glycerol (20% by volume), 1.5 mM MgCl₂, 2 mM dithiothreitol (DDT), 0.4 mM PMSF, 1 mM Na₃VO₄, 2 µg/ml each of leupeptin, pepstatin and aprotinin). Cell debris was removed by centrifugation (15,000×g, 15 min.), aliquots of the supernatant were frozen in liquid nitrogen and stored at -80° C. Protein concentration was determined by a Bradford assay (Bio-Rad) using bovine serum albumin as standard. A double-stranded oligonucleotide corresponding to NF-κB binding element (10 pmol, Promega) was labeled with [³²P] dCTP (300 Ci/mmol) and T4 polynucleotide kinase (New

England Biolabs). Free nucleotides were removed by a spin column. Extracts (10 µg protein) of cells treated with IL-18 or IL-18 plus IL-18BP were incubated (15 min., room temperature) with the labeled probe (3×10⁴ cpm) together with poly dI.dC (500 ng, Pharmacia) and denatured salmon sperm DNA (100 ng, Sigma) in 20 µl buffer consisting of HEPES (pH 7.5, 10 mM), 60 mM KCl, 1 mM MgCl₂, 2 mM EDTA, 1 mM DTT and glycerol (5% by volume). The mixtures were then loaded onto 5% non-denaturing polyacrylamide gels. Electrophoresis was performed at 185 V in 0.5×TBE (40 mM Tris HCl, 45 mM boric acid and 2.5 mM EDTA). Gels were vacuum dried and autoradiographed overnight at -80° C. IL-18 was found to induce the formation of the p50 NF-κB homodimer and the p65/p50 NF-κB heterodimer. Urinary as well as recombinant IL-18BP inhibited the activation of NF-κB by IL-18, as determined by an electrophoretic mobility shift assay with KG-1 cell extracts binding a radiolabeled oligonucleotide corresponding to the NF-κB consensus sequence.

Example 9: Expression of IL-18BP in *E. coli*, Yeast and Insect Cells

[0290] IL-18BP may also be produced by other recombinant cells such as prokaryotic cells, e.g., *E. coli*, or other eukaryotic cells, such as yeast and insect cells. Well known methods are available for constructing appropriate vectors, carrying DNA that codes for either IL-18BP and suitable for transforming *E. coli* and yeast cells, or infecting insect cells in order to produce recombinant IL-18BP. For expression in yeast cells, the DNA coding for IL-18BP (Example 6) is cut out and inserted into expression vectors suitable for transfection of yeast cells. For expression in insect cells, a DNA coding for IL-18BP is inserted into baculovirus and the insect cells are infected with said recombinant baculovirus. For expression in *E. coli*, the DNA coding for IL-18BP is subjected to site directed mutagenesis with appropriate oligonucleotides, so that an initiation ATG codon is inserted just prior to the first codon of mature IL-18BP. Alternatively, such DNA can be prepared by PCR with suitable sense and antisense primers. The resulting cDNA constructs are then inserted into appropriately constructed prokaryotic expression vectors by techniques well known in the art (Manatis, 1982).

Example 10: Construction of Adeno-Associated Expression Vector for In Vivo Expression of IL-18BP_a

[0291] A functional gene coding for IL-18BP_a is constructed based on plasmid pcDNA3 (Invitrogen, San Diego Calif.). The IL-18BP cDNA with a Kozak consensus sequence at the 5' end is ligated into the Xba I site of pcDNA3 in a way that destroys the restriction site. New Xba I sites are inserted by site-directed mutagenesis before the neomycin cassette (base 2151 of the original pcDNA3 sequence) and after the SV40 polyadenylation signal (base 3372 of the original pcDNA3 sequence). This construct is then cut with Xba I and the resulting 4.7 kb minigen is inserted at the Xba I site of plasmid psub201 as described (Snyder et al, 1996, Current Protocols in Human Genetics, Chapters 12.1.1-12.1.17, John Wiley & Sons). The resulting recombinant plasmid is cotransfected with the helper AAV plasmid pAAV/Ad into human T293 cells. The cultures are then infected with adenovirus as a helper virus and the cells are collected after 48-60 hours of

incubation. The cells are subjected to 3 freeze-thaw cycles, cell debris is removed by centrifugation, and the supernatant is brought to 33% saturation with ammonium sulfate. The mixture is then centrifuged and rAAV is precipitated from the supernatant by bringing the ammonium sulfate to 50% saturation. The virus is further purified by CsCl, dialyzed and finally heated for 15 min at 56° C. to destroy any adenovirus.

Example 11: Construction of Recombinant Fusion Proteins of IL-18BP

[0292] The production of proteins comprising IL-18BP fused to the constant region of IgG2 heavy chain may be carried out as follows: the DNA of IL-18BP is subjected to site-directed mutagenesis with appropriate oligonucleotides so that a unique restriction site is introduced immediately before and after the coding sequences. A plasmid bearing the constant region of IgG2 heavy chain, e.g. pRKCO42Fc1 (Byrn, 1990) is subjected to similar site-directed mutagenesis to introduce the same unique restriction site as close as possible to Asp 216 of IgG1 heavy chain in a way that allows translation in phase of the fused protein. A dsDNA fragment, consisting of 5' non-translated sequences and encoding for IL-18BP is prepared by digestion at the unique restriction sites or alternatively by PCR with appropriately designed primers. The mutated pRKCD42Fc1 is similarly digested to generate a large fragment containing the plasmid and the IgG1 sequences. The two fragments are then ligated to generate a new plasmid, encoding a polypeptide precursor consisting of IL-18BP and about 227 C-terminal amino acids of IgG1 heavy chain (hinge region and CH2 and CH3 domains). The DNA encoding the fused proteins may be isolated from the plasmid by digestion with appropriate restriction enzymes and then inserted into efficient prokaryotic or eukaryotic expression vectors.

Example 12: Production of Chemically Modified IL-18BPs

[0293] In order to increase the half-life of the IL-18BPs in plasma, IL-18BPs which are chemically modified with polyethylene glycol (PEG) may be made. The modification may be done by cross linking PEG to a cysteine residue of the IL-18BP molecules. Mutant IL-18BPs may be constructed which contain an extra cysteine residue at the amino terminus, glycosylation sites, and the carboxyl terminus of each IL-18BP. The mutagenesis may be carried out by PCR using oligonucleotides containing the desired mutation. These mutant proteins are expressed in the usual manner as well known in the art. Pegylation of these proteins will be carried out and the activity will be assessed.

Example 13: Preparation of Polyclonal Antibodies to IL-18BP

[0294] Rabbits were initially injected subcutaneously with 5 µg of a pure preparation of the urinary IL-18BP, emulsified in complete Freund's adjuvant. Three weeks later, they were injected again subcutaneously with 5 µg of the IL-18BP preparation in incomplete Freund's adjuvant. Two additional injections of IL-18BP as solution in PBS were given at 10 day intervals. The rabbits were bled 10 days after the last immunization. The development of antibody level was followed by a radioimmunoassay. ¹²⁵I-labeled IL-18BP (166,000 cpm) was mixed with various dilutions (1:50, 1:500, 1:5,000 and 1:50,000) of the rabbit serum. A suspension of protein-G

agarose beads (20 µl, Pharmacia) was added in a total volume of 200 µl. The mixture was left for 1 hour at room temperature, the beads were then washed 3 times and bound radioactivity was counted. Rabbit antiserum to human leptin was used as a negative control. The titer of the IL-18R antiserum was between 1:500 and 1:5000, while that of the negative control was less than 1:50.

Example 14: Preparation of Monoclonal Antibodies to IL-18BP

[0295] Female Balb/C mice (3 months old) are first injected with 2 µg purified IL-18BP in an emulsion of complete Freund's adjuvant, and three weeks later, subcutaneously in incomplete Freund's adjuvant. Three additional injections are given at 10 day intervals, subcutaneously in PBS. Final boosts are given intraperitoneally 4 and 3 days before the fusion to the mouse showing the highest binding titer as determined by IRIA (see below). Fusion is performed using NSO/1 myeloma cell line and lymphocytes prepared from both the spleen and lymph nodes of the animal as fusion partners. The fused cells are distributed into microculture plates and the hybridomas are selected in DMEM supplemented with HAT and 15% horse serum. Hybridomas that are found to produce antibodies to IL-18BP are subcloned by the limiting dilution method and injected into Balb/C mice that had been primed with pristane for the production of ascites. The isotypes of the antibodies are defined with the use of a commercially available ELISA kit (Amersham, UK).

[0296] The screening of hybridomas producing anti-IL-18BP monoclonal antibodies is performed as follows: Hybridoma supernatants are tested for the presence of anti-IL-18BP antibodies by an inverted solid phase radioimmunoassay (IRIA). ELISA plates (Dynatech Laboratories, Alexandria, Va.) are coated with Talon-purified IL-18BPα-His₆ (10 µg/ml, 100 µl/well). Following overnight incubation at 4° C., the plates are washed twice with PBS containing BSA (0.5%) and Tween 20 (0.05%) and blocked in washing solution for at least 2 hrs at 37° C. Hybridoma culture supernatants (100 Owen) are added and the plates are incubated for 4 hrs at 37° C. The plates are washed 3 times and a conjugate of goat-anti-mouse horseradish peroxidase (HRP, Jackson Labs, 1:10,000, 100 Owen) is added for 2 hrs at room temperature. The plates are washed 4 times and the color is developed by ABTS (2,2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid, Sigma) with H₂O₂ as a substrate. The plates are read by an automatic ELISA reader. Samples giving OD that are at least 5 times higher than the negative control value are considered positive.

Example 15: Affinity Chromatography of IL-18BP with Monoclonal Antibodies

[0297] Antibodies against IL-18BP are utilized for the purification of IL-18BP by affinity chromatography. Ascitic fluid containing the monoclonal antibody secreted by the hybridoma is purified by ammonium sulfate precipitation at 50% saturation followed by extensive dialysis against PBS. About 10 mg of immunoglobulins are bound to 1 ml Affigel 10 (BioRad USA), as specified by the manufacturer.

[0298] 250 ml of human urinary proteins (equivalent to 250 l of crude urine) are loaded on 0.5 ml of the anti IL-18BP antibody column at 4° C. at a flow rate of 0.25 ml/min. The column is washed with PBS until no protein is detected in the washings. IL-18BP is eluted by 25 mM citric acid buffer, pH

2.2 (8×1 column volume fractions) and immediately neutralized by 1 M Na₂CO₃. Further purification of this preparation is obtained by size exclusion chromatography.

Example 16: ELISA Test

[0299] Microtiter plates (Dynatech or Maxisorb, by Nunc) are coated with anti-IL-18BP monoclonal antibody (serum free hybridoma supernatant or ascitic fluid immunoglobulins) overnight at 4° C. The plates are washed with PBS containing BSA (0.5%) and Tween 20 (0.05%) and blocked in the same solution for at least 2 hrs at 37° C. The tested samples are diluted in the blocking solution and added to the wells (100 µl/well) for 4 hrs at 37° C. The plates are then washed 3 times with PBS containing Tween 20 (0.05%) followed by the addition of rabbit anti-IL-18BP serum (1:1000, 100 µl/well) for further incubation overnight at 4° C. The plates are washed 3 times and a conjugate of goat-anti-rabbit horseradish peroxidase (HRP, Jackson Labs, 1:10,000, 100 Owen) was added for 2 hrs at room temperature. The plates were washed 4 times and the color is developed by ABTS (2,2'-azino-bis(3-ethyl-benzthiazoline-6-sulfonic acid, Sigma) with H₂O₂ as a substrate. The plates are read by an automatic ELISA reader.

Example 17: Non-Glycosylated Human IL-18BP is Biologically Active

[0300] Purified recombinant IL-18BP_a was tested for its ability to inhibit the biological activity of IL-18. IL-18BP_a inhibited in a dose dependent manner the IFN-γ inducing activity of human and mouse IL-18 in murine splenocytes, PBMC and the human KG-1 cell line (FIG. 4).

[0301] Purified IL-18BP_a having an His₆ tag at the C-terminus (1.5 µg, 50 µl) was adjusted to pH 7.5 and mixed with N-glycosidase F (3 µl, 500,000 U/ml, PNGase F, New England Biolabs). The mixture was incubated for 24 h at 37° C. under non-denaturing conditions. Aliquots from the sample and from undigested IL-18BP-His₆ were analyzed by SDS-PAGE under non-reducing conditions followed by immunoblotting with antibodies to IL-18PB. It was found that the ~40 kD band of IL-18BP-His₆ disappeared in the PNGase-treated fraction and a new ~20 kD band was obtained. The molecular mass of the product and the specificity of PNGase F indicated that IL-18BP-His₆ was fully deglycosylated.

[0302] The PNGase-treated fraction, undigested IL-18BP-His₆ and control sample containing PNGase in buffer were absorbed separately on Talon beads, washed with phosphate buffer and eluted with imidazole (100 mM). The eluted fractions were subjected to bioassay using human IL-18 (20 ng/ml), LPS (2 µg/ml) and murine splenocytes. The results are shown in the following Table 4:

TABLE 4

Sample	IFN-γ (ng/ml)
Control	7.5
Non-digested IL-18BP-His ₆	0
PNGase-treated IL-18BP-His ₆	0

[0303] Therefore, it is concluded that deglycosylated IL-18BP is biologically active as a modulator of IL-18 activity.

PART II: Examples 18 to 25

Relating to the Use of IL-18 Inhibitors in Liver Injury

Example 18: Production of IL-18BP-His Tag

[0304] Purified recombinant human IL18BP containing a his-tag (r-hIL-18BP-His tag) was produced in CHO cells. The production of recombinant proteins in eukaryotic cells is known by the person skilled in the art. Well known methods are available for constructing appropriate vectors, carrying DNA that codes for IL-18BP and suitable for transfection of eukaryotic cells in order to produce recombinant IL-18BP. For expression in cells, the DNA coding for IL-18BP (see, e.g. (Novick et al., 1999) is cut out and inserted into expression vectors suitable for transfection of cells. Alternatively, such DNA can be prepared by PCR with suitable sense and antisense primers. The resulting cDNA constructs are then inserted into appropriately constructed eukaryotic expression vectors by techniques well known in the art (Maniatis, 1982). The recombinant protein was purified to over 95% purity and found to be biologically active in-vitro and in-vivo with a high affinity to its ligand.

Example 19: Protective Effect of IL18BP Against Endotoxin-Induced Death in the Murine Model

[0305] A murine model was used to test whether IL18BP, an inhibitor of IL-18, would protect mice against a high dose of lipopolysaccharides (LPS). LPS elicits acute liver injury, followed by rapid death of the mice.

[0306] 4 mg/kg of recombinant, human IL-18BP (rhIL18BP_{his}) containing a his-tag (resulting from recombinant production of the protein) was injected intraperitoneally (i.p.) into C57BL/6 mice. 1 h later, 60 mg/kg LPS were injected (lethal doses). The survival of mice was compared to a group of animals who received LPS alone (no IL18BP).

[0307] Five out of 7 mice injected with rhIL-18BP-his survived the LPS injection in contrast to the control mice, in which all animals died within 3 days.

[0308] Blood samples were taken 5 h after the LPS injection in the absence or presence of increasing doses of rhIL-18BP-his and analyzed by ELISA for circulating IFN-γ (FIG. 5). 0.4 and 4 mg/kg rhIL-18BP induced a 2 fold reduction in serum IFN-γ. This inhibition was lost at lower doses of rhIL-18BP (0.004 and 0.4 mg/kg).

Example 20: IL18BP has a Protective Effect Against Liver Injury in a Murine Model

[0309] A mouse model of fulminant hepatitis was used to test the effect of IL18BP. Mice develop acute liver injury when subjected to a sequential administration of *Propionibacterium acnes* (*P. acnes*) and lipopolysaccharide (LPS).

[0310] Mice were injected with increasing doses of rhIL-18BP-his (4; 0.4; 0.04; 0 mg/kg) at various times (1 h, 20 min, simultaneously) before the injection of LPS in C57BL/6 *P. acnes* sensitized mice. When rhIL-18BP-his was given i.p. at the same time as LPS, none of the mice survived and levels of circulating IFN-γ and TNF-α were unaffected. Surprisingly, rhIL-18BP (4 and 0.4 mg/kg) induced a 70% reduction of circulating Alanine aminotransferase (a marker of liver injury), as shown in FIG. 6.

[0311] In addition to this, the survival of mice was monitored (FIG. 7): When rhIL-18BP was given i.p. 20 minutes before LPS, the two highest doses of IL-18BP (4 and 0.4

mg/kg) delayed the death of the mice by 10 h as compared to the control mice who received NaCl instead of IL-18BP

[0312] The results of the measurement of serum IFN- γ levels are shown in FIG. 8. rhIL-18BP (4 mg/kg) inhibited 90% of circulating IFN- γ levels and 80% of circulating Alanine aminotransferase (not shown).

[0313] When rhIL-18BP-his was given 1 h before LPS, survival curves and levels of circulating IFN- γ were similar to what was observed when rhIL-18BP-his was given 20 min before LPS, but levels of circulating alanine aminotransferase were unaffected (not shown).

[0314] In addition to this, murine liver tissue was analyzed by hematoxylin-eosine staining, as well as by tunnel microscopy. The livers of mice, in which severe hepatitis had been induced before, showed severe necrosis as compared to normal liver tissue. In contrast to this, liver tissue of mice treated with IL-18BP showed significantly less necrotic foci than untreated mice.

Example 21: Anti-IL-18 Antibodies Protect Against Lethal Endotoxemia

[0315] In order to evaluate, whether blockade of IL-18 with IL-18 antibodies would protect mice against lethal doses of bacterial lipopolysaccharides, C57BL/6J mice were first injected with a neutralizing rabbit anti-mouse IL-18 antibody (polyclonal) or normal rabbit serum (NDS) as a control. 30 min after antibody treatment, a lethal dosis of LPS derived either from *E. coli* (FIG. 9 A) or *S. typhimurium* (FIG. 9 B) was injected. Experiments involved 10-12 mice/group, and were performed twice on two different occasions.

[0316] As shown in FIG. 9A, treatment of the mice with the anti-IL-18 antiserum prevented the mortality induced by 40 mg/kg *E. coli* LPS. 100% of the mice survived after anti-IL-18 treatment vs. 10% survival in mice treated with normal rabbit serum ($p < 0.005$).

[0317] FIG. 9B shows that the antibody treated mice were also protected against *S. typhimurium* lethal effects (50% vs. 0% survival; $p < 0.05$).

Example 22: Blockade of IL-18 and TNF- α Protects Mice from ConA- and PEA-Induced Hepatotoxicity

[0318] Two experimental models of hepatotoxicity were used to evaluate the role of IL-18 and TNF- α in liver injury. Injection of Concanavalin A (Con A) and *Pseudomonas aeruginosa* (PEA) into mice both induce liver injury, and are models of T cell mediated hepatitis.

[0319] C57BL/6J mice were pretreated with an anti-IL-18 antiserum or a soluble TNF- α receptor, TNFRp55. Serum Alanine aminotransferase (ALT) levels were measured as indicators of hepatic injury (FIG. 10).

[0320] As shown in FIG. 10 A, both IL-18 antiserum and soluble TNF-receptors significantly reduced ConA-induced serum ALT levels, as compared to a control injection of the vehicle alone (pyrogen free saline). A co-administration of soluble TNF receptor and IL-18 antiserum led to a complete inhibition of Con-A induced liver injury.

[0321] As shown in FIG. 10 B, in PEA-injected mice, neutralization of either TNF- α inhibitors or anti-IL-18 antibodies

resulted in 93% and 83% inhibition of serum ALT levels, respectively. A combined blockade of both resulted in 99% protection.

Example 23: Plasma Levels of IL-18-Binding Protein are Elevated in Patients with Chronic Liver Disease

[0322] IL-18 BP plasma levels were measured in 133 patients with chronic liver disease (CLD) of varying etiologies and 31 healthy controls by a specific ELISA, using an IL-18BP monoclonal antibody.

[0323] Plasma levels of IL-18 BP were significantly higher in CLD patients (12.91 ± 0.89 ng/ml; average $\pm$ SEM) than in healthy subjects (4.96 ± 0.43 ng/ml, $p < 0.001$). Cirrhotic patients had significantly higher levels than patients with non-cirrhotic CLD (19.23 ± 1.28 ng/ml, $n = 67$, vs. 6.49 ± 0.51 ng/ml, $n = 66$, $p < 0.001$). Patients with stage B of the Child-Pugh classification had higher levels of IL-18 BP than those with stage A (22.48 ± 2.44 ng/ml vs. 9.57 ± 1.25 ng/ml, $p < 0.001$). However, there was no significant difference between Child B and C (22.48 ± 2.44 ng/ml vs. 20.62 ± 4.75 ng/ml, $p = 0.7$). Plasma levels of IL-18 BP correlated positively with GOT, bilirubin and erythrocyte sedimentation rate. Negative correlation was found with prothrombin time.

[0324] In conclusion, the results show that IL-18 BP plasma levels are elevated in CLD and correlate with the severity of disease independent of the etiology of disease. Although an endogenous antagonist of the proinflammatory IL-18, increased levels of IL-18 BP seem not to be sufficient to counteract the overwhelming proinflammatory mediators in CLD.

Example 24: Inhibition of Alcoholic Hepatitis by IL-18BP

[0325] Four groups of rats (5 per group) are fed ethanol and a diet containing corn oil by intragastric infusion for 4 weeks. Dextrose isocalorically replaces ethanol in control rats. The rats are injected daily with mouse IL-18BP (1 mg/kg), or saline. Pathological analysis is performed on liver sections and measurements of liver enzymes in serum, TNF- α , Fas ligand and IFN- γ are taken. Necroinflammatory injury and expression of liver enzymes, TNF- α , Fas ligand, and IFN- γ are seen in the ethanol-fed rats that were injected with saline.

[0326] Rats injected with mouse IL-18BP are protected from necroinflammatory injury and the levels of liver enzymes, TNF- α , Fas ligand and IFN- γ are significantly reduced ($> 90\%$).

Example 25: Inhibition of Concanavalin A-induced Hepatitis by IL-18BP

[0327] Balb/c mice are injected with 12 mg/kg Concanavalin A (Con A) with or without injection of murine IL-18BP (1 mg/kg), 2 h prior to Con A administration and then daily. Liver damage is evaluated by determining serum levels of liver enzymes, TNF- α , Fas ligand and IFN- γ . Hepatic histopathology is compared to mice treated with Concanavalin A only.

[0328] Pretreatment with IL-18BP significantly reduces serum levels of liver enzymes and TNF- α with no evidence of

inflammation in histopathologic examination compared to control mice treated with Con A.

PART III: Examples 26 and 27

Relating to the Use of IL-18 Inhibitors in Arthritis

Example 26: Production of IL-18BP-His Tag

[0329] For the experiment described in detail in Example 27 below, recombinant human IL-18BP containing a his-tag of 6 residues (r-hIL-18BP-His tag) was produced in CHO cells and purified as described by Kim et al., 2000. The recombinant protein was purified to over 95% purity and found to be biologically active in-vitro and in-vivo with a high affinity to its ligand.

[0330] The production of recombinant proteins in other eukaryotic systems, with or without tags facilitating purification of the recombinant proteins, is known by the person skilled in the art. Well known methods are available for constructing appropriate vectors, carrying DNA that codes for IL-18BP and suitable for transfection of eukaryotic cells in order to produce recombinant IL-18BP. For expression in cells, the DNA coding for IL-18BP (see, e.g. Novick et al., 1999) is cut out from the cloning vector and inserted into expression vectors suitable for transfection of cells. Alternatively, such DNA can be prepared by PCR with suitable sense and antisense primers. The resulting cDNA constructs are then inserted into appropriately constructed eukaryotic expression vectors by techniques known in the art (Maniatis et al., 1982).

Example 27: Blockade of Endogenous IL-18 in a Murine Model of Arthritis

[0331] Methods

[0332] Induction of Collagen-Induced Arthritis (CIA)

[0333] CIA was induced in male DBA/1 mice (8-12 weeks old) by immunisation with native type II bovine collagen (CII) as previously described (Plater-Zyberk et al., 1995). From day 25 post-CII immunisation, mice were examined daily for onset of disease.

[0334] Treatment with rhIL-18BP-6his

[0335] Treatment of CII-immunised DBA/1 mice was started on the first appearance of clinical sign of disease. Recombinant, human IL-18BP containing a tag of 6 histidines (rh-IL-18BP-6his) was used to neutralise endogenous IL18 in the collagen treated mice. rh-IL-18BP-6his was injected daily for 7 seven days at 5 different concentrations 10, 3, 1, 0, 5, 0, 25 mg/kg/injection intraperitoneally (i.p.). The placebo control mice received the vehicle only (0.9% NaCl).

[0336] Assessment of Disease Development

[0337] Clinical Evaluation (Clinical Scores)

[0338] From first appearance of clinical symptoms, mice were examined every day by an investigator blinded to the treatment. Each limb was graded for disease severity (scores 0-3.5, max score=14/mouse). The progression of edema (inflammation) was measured on the first paws that showed signs of disease using precision calipers (Protest 2T, Kroepelin Langenmesstechnik)

[0339] Disease progression was assessed daily for 8 days post-onset at which time all mice were sacrificed and paws collected for histopathological examination.

[0340] Histological Assessment of Cartilage Erosions and Microscopical Inflammation

[0341] At termination of the experiments, i.e. at day 8 post-onset, mice were killed and the paw that first developed sign of disease was dissected away. Joints were fixed, decalcified and embedded in paraffin. Standard sections (5 to 7 μ m) of the joints were made and stained with hematoxylin/eosin/Safranin O. Each joint was scored by 2 investigators unaware of the treatment protocol (no destruction of cartilage or bone=0; localised cartilage erosions=1-2; more extended erosions=3; general cartilage destruction and presence of bone erosions=4). The final scores of each mouse was the mean of the result on all the scored joints. Microscopical inflammation or synovitis was scored from 0 to 4, as follows: no inflammation=0; slight thickening of lining layer and/or some infiltrating cells in sublining layer=1-2; thickening of lining layer and/or more pronounced influx of cells of sublining layer=3; presence cells in the synovial space and synovium highly infiltrated with many inflammatory cells=4.

[0342] Determination of Anti-Collagen Antibodies.

[0343] Antibodies against bovine type II collagen were examined by using an enzyme-linked immunosorbent assay (ELISA). Titers of IgG1 and IgG2a were measured. Briefly, plates were coated with 10 μ g of bovine collagen and thereafter-nonspecific binding sites were blocked with 0.1 M ethanolamine (Sigma). Serial 1:2 dilutions of the sera were added followed by incubation with isotype specific goat anti-mouse peroxidase (Southern Biotechnology Associates, Birmingham, Ala., USA) and substrate (5-aminosalicylic acid, Sigma). Plates were read at 492 nm. Titers were expressed as mean $\pm$ SD dilution, which gives the half-maximal value.

[0344] IL-6 Assays

[0345] Levels of IL-6 were determined by commercial ELISA (R&D systems, Minneapolis, Minn., USA). IL-6 bioactivity was determined by a proliferative assay using B9 cells. Briefly, 5×10^3 B9-cells in 200 μ l 5% FCS-RPMI 1640 medium per well were plated in a round-bottom microtitre plate and incubated for 3 days using human recombinant IL-6 (R&D systems, Minneapolis, Minn., USA) as standards. At the end of the incubation, 0.5 μ Ci of ^{3} [H]thymidine (NEN-Dupont, Boston, Mass., USA) was added per well. Three hours later, cells were harvested and thymidine incorporation was determined. Detection limit for the IL-6 bioassay was 1 pg/ml.

[0346] Statistical Analysis

[0347] Significance of differences was assessed by the Mann Whitney test using SigmaStat statistical analysis program and the GraphPad Prism program.

[0348] Results

[0349] A mouse experimental model, CIA (collagen induced arthritis), was used for assessing the effectiveness of IL-18BP as an agent for the treatment of arthritis. Administration of collagen and incomplete Freund's adjuvant in DBA/1 mice induces the development of an erosive, inflammatory arthritis and represents an ideal opportunity to explore the therapeutic potential of IL-18BP. To this end, endogenous IL-18 was neutralised using IL-18BP and the effect on various pathogenic parameters was evaluated.

[0350] A dose-related study was performed. Three groups of collagen-induced arthritic DBA/1 mice were treated therapeutically (i.e. after onset of disease) with 5 doses of IL-18BP i.p. (intraperitoneal). IL-18BP at concentrations of 10, 3, 1, 0.5 or 0.25 mg/kg was administered at the first clinical sign of disease. Injection of physiological saline (sodium chloride, NaCl) was used as a control. In addition to this, 10 000 IU of interferon β (IFN- β) were administered i.p. to assess the

effect of IFN in this experimental model of arthritis. The effect on disease severity was monitored by daily visual scoring of each individual paw as described above. The mice were sacrificed at day 8 post-onset.

[0351] The following values were measured:

[0352] visual clinical scores (0-3.5 per paw) (FIGS. 11A and B)

[0353] joint swelling/edema (in mm, measured with calipers) of first diseased paw, provided it was a hind paw (FIG. 12)

[0354] number of paws recruited into the disease (FIG. 13)

[0355] erosion scores of first diseased paw (0-4 cartilage destruction, FIG. 14).

[0356] Histopathological analysis of the paw that first developed arthritis (FIG. 15)

[0357] Levels of anti-collagen type II antibodies (FIG. 16)

[0358] Levels of IL-6 (FIG. 17).

[0359] Clinical Severity of Disease

[0360] As shown in FIGS. 11A and B, the clinical severity of disease was significantly diminished in the groups treated with 1 mg/ml ($P<0.01$) and 0.5 mg/ml ($0.01<P<0.05$) of rhIL-18BP. Mice receiving the low dose of rhIL-18BP (0.25 mg/kg) or the high dose of 10 mg/kg had a clinical score similar to the placebo group. The dose of 1 mg/kg of IL-18BP was approximately as effective as Interferon β (designated IFNb in FIG. 11A).

[0361] Joint Inflammation and Paw Swelling (Edema)

[0362] Macroscopical inflammation (swelling) was studied by measuring paw edema from day 1 after onset of disease until day 8, the end of the experiment. The results are shown in FIGS. 12A and B. The effective doses of IL-18BP were 1, 3 and 10 mg/kg. Administration of lower doses did not result in a beneficial effect on the swelling of paws. As shown in FIGS. 12A and 12B, Interferon- β (IFNb) at a concentration of 10 000 IU showed a beneficial effect on paw swelling.

[0363] Microscopic synovitis was examined at the end of the experiment on histopathological sections and was expressed as scores ("synovitis score"). The results of inflammation (swelling) and synovitis score are summarized in Table 1. Whilst treatment with rhIL-18BP at dosages 1 and 3 mg/kg resulted in a trend towards reduction of swelling, treatment with any dosage of IL-18BP had only a limited effect on the inflammatory synovitis (Table 5).

TABLE 5

EFFECT OF IL-18BP TREATMENT ON JOINT INFLAMMATION			
Treatment		Swelling (AUC, mean $\pm$ sem)	Synovitis score (mean $\pm$ sem)
RhIL-18BP-6his	3 mg/kg	1.55 $\pm$ 0.64	2.17 $\pm$ 0.5
	1 mg/kg	1.53 $\pm$ 0.60	1.98 $\pm$ 0.4
	0.5 mg/kg	3.38 $\pm$ 0.70	1.92 $\pm$ 0.5
	0.25 mg/kg	3.06 $\pm$ 0.90	2.08 $\pm$ 0.5
Placebo		3.11 $\pm$ 0.77	2.23 $\pm$ 0.3

[0364] FIG. 13 shows that the number of paws affected by the disease was diminished after administration of IL-18BP. In particular, therapeutic injections of IL-18BP at doses of 1 and 0.5 mg/kg reduced the number of paws recruited into the disease, demonstrating that blockade of IL-18 in vivo halts the spreading of arthritis to additional joints. Treatment with

1 and 0.5 mg/kg of IL-18BP even appears capable of reverting some of the arthritic joints to normality.

[0365] Protection from Joint Destruction

[0366] Treatment of mice with rhIL-18BP resulted in protection of joints from destruction (FIG. 14). A semi-quantitative scoring system demonstrated that bone erosion showed a dose-related protective effect that was significant at 10 and 3 mg/kg ($P<0.05$, FIG. 14). Mice receiving 1 mg/kg of rhIL-18BP presented less erosion than mice receiving vehicle only. No protection was observed at doses of 0.5 mg/kg and 0.25 mg/kg. Interestingly, the effect on joint protection at doses of 3 and 10 mg/kg IL-18BP were comparable to or even more pronounced than the beneficial effect of 10 000 IU of Interferon β (IFN- β).

[0367] FIG. 15 shows the histology of a healthy (A) and a diseased (B) joint in comparison to a joint derived from an animal treated with IL-18BP (C). Sections were taken at the end of the experiment from those paws which first developed arthritis

[0368] The joint from an arthritic mouse showed severe destructive arthritis with cartilage depletion and erosions and numerous infiltrating cells in the inflamed synovium. In the joint from a mouse treated with rhIL-18BP, the cartilage appeared almost normal despite the presence of inflammatory cells in the synovial space. There was not only a higher amount of cartilage, but the cartilage has also a smoother appearance.

[0369] Anti-IL-18 Treatment Modulates Levels of Anti-Type II Collagen Antibodies

[0370] CIA mice have elevated levels of IgG1 and IgG2a anti-type II collagen antibodies in the circulation. Antibodies of the isotype IgG1 are associated to TH2 mediated diseases, whereas antibodies of the isotype IgG2a and IgG2b are associated to TH1 mediated diseases. Arthritis is usually classified as a TH1 mediated disease.

[0371] Anti-type II collagen (CII) IgG1 and IgG2a antibody isotypes were determined in the sera of animals that were treated with IL-18BP (FIG. 16). Levels of anti-CII of the IgG isotypes IgG1 and IgG2a were not significantly modified by IL-18BP treatment at day 4 or 8 (D4, D8) of clinical disease. However, a 2.6 and 3.4 fold decrease in collagen-specific IgG1/IgG2a ratios was observed after 8 days of rhIL-18BP-treatment, at 1 and 3 mg/kg respectively. FIG. 11 shows the experiment in which 3 mg/kg were used. Essentially the same results were obtained using an amount of 1 mg/kg of IL-18BP. The reduced IgG1/IgG2a ratio of anti-CII antibodies indicate a diminished concentration of anti-type II collagen antibodies of the isotype IgG2a and an elevated concentration of anti-type II collagen antibodies of the isotype IgG1, suggesting that there is a shift towards TH2-mediated disease in this model of arthritis.

[0372] Reduction of IL-6 Levels after IL-18 Neutralisation

[0373] To gain some insight into the effects of IL-18 blockade, IL-6 was measured in the sera of IL-18BP treated animals. FIG. 17 shows that the levels of bioactive IL-6 was significantly reduced in the animals having received IL-18BP treatment at all doses measured, i.e. at 1, 3 and 10 mg/kg as well as with Interferon- β (IFNb). Immunoactive levels of IL-6 measured in the sera of the animals treated with 3 mg/kg rhIL-18BP were significantly reduced ($P<0.0023$) as compared with saline-treated animals. IL-6 serum levels of diseased animals treated with 1, 3 or 10 mg of IL-18BP or 10 000 IU of IFN- β were similar to normal mouse serum (NMS)

derived from healthy animals, i.e. from those animals not having an inflammatory disease.

[0374] These findings demonstrate that IL-18 controls IL-6 levels during the onset of the disease. Since IL-6 is a marker of inflammation, these findings show that treatment of diseased mice with IL-18BP reduces inflammation in the animal.

[0375] From the experiments outlined above, the following conclusions can be drawn:

[0376] Administration of IL-18BP decreases the clinical severity of arthritis

[0377] IL-18BP inhibits further progression or spreading of the disease

[0378] Administration of IL-18BP decreases oedema

[0379] Administration of IL-18BP decreases cartilage destruction

[0380] Serum IL-6 levels are diminished and IgG1/IgG2a anti-CII ratios decreased after IL-18BP therapy.

[0381] The data presented above indicate that neutralisation of IL-18 bioactivity after disease onset represents a disease-modifying anti-rheumatic therapy. The results clearly demonstrate that IL18 blockade reduces the clinical progression of arthritis and more importantly stops progression of cartilage and bone destruction. IL18 blockade by IL-18BP, anti-IL-18 antibodies or by any other IL-18 blocking agent therefore represents a new disease-modifying anti-rheumatic therapy.

PART IV: Examples 28-34

Relating to the Use of IL-18 Inhibitors in Sepsis

Example 28: Levels of Serum IL-18BP and IL-18 in Healthy Individuals and Sepsis Patients

[0382] Measurement of specific circulating cytokines, and their natural inhibitors in health and disease provides information about their involvement in progression and severity of a disease. As mentioned in the background section, one of the key mediators of sepsis is IFN- γ . Since IL-18 is a coinducer of IFN- γ , the level of IL-18 and its natural inhibitor, IL-18BP splice variant a, were monitored in sepsis patients and compared to the levels found in healthy subjects by using specific ELISAs (Example 30).

[0383] Levels of IL-18 and IL-18BP in healthy subjects.

[0384] The mean level of IL-18 in 107 healthy donors was 64 ± 17 pg/ml as measured by the ECL assay (Pomerantz et al. 2001). The ECL assay was tested for interference by related proteins. It was found that it was not affected by the presence of mature IL-1 β or proIL1 β , whereas pro-IL-18 was cross-reactive. Therefore, as much as 20% of the detected mature IL-18 in human serum samples in the present study may be pro-IL-18. The ECL assay of IL-18 was not affected by IL18BP at a concentration ≤ 160 ng/ml.

[0385] The levels of IL-18BP were tested in the serum from the 107 healthy individuals by ELISA (Example 30). The levels of IL-18BP ranged from 0.5 ng/ml to as high as 7 ng/ml, with an average of 2.15 ± 0.15 ng/ml (FIG. 18).

[0386] Because both IL-18 and IL-18BP are concomitantly present in the serum, some of the IL-18 may be present in a complex with IL-18BP. The level of free IL-18 was calculated based on the average level of total IL-18 (2.15 ng/ml). Free IL-18 was determined according to the law of mass action. The calculation was based on the following parameters: the concentrations of total IL-18 as determined by the ECL assay; the concentration of total IL-18BP as determined by the ELISA; a 1:1 stoichiometry in the complex

of IL-18BP and IL-18 and a dissociation constant (Kd) of 0.4 nM (Novick et al. 1999 and Kim et al. 2000). In an equilibrium system $L + R \rightleftharpoons LR$ where L represents IL-18 and R represents IL-18BP the following equations are applicable:

$$Kd = [LR] / [L_{free}] [R_{free}]$$

$$L_{free} = L_{total} - LR$$

$$R_{free} = R_{total} - LR$$

[0387] By substituting: $L_{total} = 64 \pm 17$ pg/ml (mean level). $R_{total} = 2.15 \pm 0.15$ ng/ml (average) and $Kd = 0.4$ nM, it was found that in healthy subjects about 51.2 pg/ml IL-18 (about 80% from total) is in its free form.

[0388] Levels of IL-18 and IL-18BP in Sepsis Patients.

[0389] The levels of IL-18 and of IL-18BP were tested in 192 sera samples from 42 septic patients immediately upon admission and during hospitalisation. The levels of both IL-18 and IL-18BP were significantly more elevated in sepsis patients in comparison with the healthy subjects (FIG. 19A), and a broad distribution of the values was observed (FIG. 19B). Moreover, these levels were even higher in these patients at the day of admission, showing a 22 fold increase in the level of IL-18 compared with healthy individuals (1.5 ± 0.4 ng/ml versus 0.064 ± 0.17 ng/ml) and a 13 fold increase in the level of IL-18BP (28.6 $\pm$ 4.5 versus 2.15 ± 0.15 ng/ml, FIG. 19A). No statistically significant correlation between creatinine levels and either IL-18 or IL-18BP concentrations in these sera could be observed (assessed by APACHE II score Knaus et al. 1993), suggesting that elevated IL-18 and IL-18BP levels in these patients were not due to renal failure.

[0390] Because serum IL-18 and IL-18BP levels in sepsis patients varied considerably (FIG. 19B) the levels of free serum IL-18 in individual samples were calculated. The calculations were done as previously described, using the same three equations and substitution of the Kd, L_{total} , R_{total} values found experimentally. The calculations show that IL-18BP reduced the level of free IL-18 in most patients (FIG. 20). Notably, this effect was particularly strong when total IL-18 was very high. For example, in two patients, serum IL-18 reached 0.5 nM (10 ng/ml), a 150-fold increase over healthy individuals. However, considering binding to the circulating IL-18BP (27 ng/ml and 41.2 ng/ml), the level of free IL-18 in these two patients dropped by 78% and 84%, respectively (FIG. 20). Therefore, most of the serum IL-18 is blocked by the circulating IL-18BP. Yet, according to these calculations, the remaining free IL-18 is still higher than that found in healthy individuals. Therefore, administration of exogenous IL-18BP to septic patients is expected to further lower the free circulating IL-18 level, causing alleviation of the disease outcome.

Example 29: Effect of IL-18BP on *S. epidermidis*-induced IFN- γ Production in Cells Present in Whole Blood Samples from Healthy Subjects

[0391] The Gram-positive bacteria *Staphylococcus epidermidis* is known to cause sepsis. Induction of cytokines, e.g., IL-1, IFN- γ and TNF- α are key mediators for this pathology. Since IL-18 is a co-inducer of IFN- γ , the effect of its inhibition on *Staphylococcus epidermidis* IFN- γ induction, was tested. The IL-18BP, used as the IL-18 inhibitor, was a recombinant histidine tagged version produced in CHO cells.

[0392] 0.5 ml of blood was mixed in 5 ml 12x75 mm round-bottom polypropylene tubes (Falcon, Becton Dickinson Labware, Franklin Lakes, NJ) with 0.5 ml of RPMI growth medium (Cellgro Mediatech, Hendon, Va. supplemented with 10 mM L-glutamine, 100 U/ml penicillin 100 µg/ml streptomycin and 10% FBS [Gibco BRL, Grand Island, N.Y.]) containing *S. epidermidis* (from ATCC) with or without IL-18BP. The samples were incubated at 37° C. (5% CO₂) for 48 hours followed by treatment with triton X-100 (Bio-Rad Laboratories, Richmond, Calif.) at 0.5% final concentration to lyse the cells present in the whole blood sample. The tubes containing the samples were inverted several times until the blood sample showed a clear appearance. IFN-γ concentration in the lysed samples of blood, which represents both intracellular and secreted cytokine, was measured by electrochemiluminescence (ECL) as described by Pomerantz et al. (2001).

[0393] For this study whole blood from healthy, non-smoking volunteers was used. The blood was taken by venipuncture and kept in heparinized tubes. *S. epidermidis* was propagated in brain-heart infusion broth for 24 hours, washed in pyrogen free saline and boiled as described by Aiura et al. (1993). The concentration of heat-killed bacteria was adjusted to 10 bacterial cells per white blood cell (WBC).

[0394] The results in FIG. 21 show that IL-18BP_a is able to inhibit the *S. epidermidis* induction of IFN-γ.

[0395] Since IFN-γ is known to be co-stimulated by IL-18 and IL-12, the effect of the combination of IL-18BP_a and anti IL-12 specific antibodies (anti human IL-12 Mab 11.5.14, Preprotech, Rocky Hill, N.J.) on IFN-γ induction directed by *S. epidermidis*, was tested. This induction was tested in the presence of 125 ng/ml IL-18BP_a (histidine tagged IL-18BP_a produced in COS cells and purified over a talon column as described by Novick et al. 1999) and 2.5 µg/ml anti human IL-12 specific antibody. The results shown in FIG. 22 suggest that anti IL-12 specific antibodies potentiate the inhibitory effect of IL-18BP_a on the IFN-γ induction by *S. epidermidis*.

Example 30: Development of an IL-18BP Specific ELISA Test

[0396] ELISA comprised two anti IL-18BP_a antibodies: murine Mab 582.10, a subclone of Mab 582 described in Example 4, as the capture antibody and rabbit polyclonal antibody for detection (Example 4). Microtiter 96-well ELISA plates (Maxisorb; Nunc A/S, Roskilde, Denmark) were coated with anti IL-18BP_a MAb No. 582.10 (a subclone of Mab 582, 4 µg/ml in PBS) overnight at 4° C. The plates were washed with PBS containing 0.05% Tween 20 (washing solution) and blocked (2 h, 37° C.) with BSA stock solution (KPL, Geithersburg, Md.) diluted 1:10 in water. BSA stock solution was diluted 1:15 in water (diluent) for the dilution of all tested samples and the detecting antibody. Sera samples were diluted at least 1:5 in the diluent and 100 µl aliquots were added to the wells. Highly pure rIL-18BP_a (prepared in CHO and purified by immunoaffinity using protein G-purified Mab N 430 specific for IL-18BP shown in Table 6, example 31) was diluted by 7 serial two-fold dilutions (4 to 0.062 ng/ml) and added to each ELISA plate for the generation of a standard curve. The plates were incubated for 2 hrs at 37° C. and washed 3 times with the washing solution. Rabbit anti IL-18BP_a serum (1:5000 in diluent, 100 µl/well) was added and the plates were further incubated for 2 hrs at 37° C. The plates were washed 3 times, a conjugate of goat-anti-rabbit horseradish peroxidase (HRP, Jackson ImmunoResearch

Labs, 1:10,000 in PBS, 100 µl/well) was added and the plates were incubated for 1 h at 37° C. The plates were washed 3 times and developed by the addition of OPD Peroxidase substrate (o-phenylenediamine dihydrochloride tablets, Sigma) for 30 min at room temperature. The reaction was stopped by 3N HCl (100 µl) and the absorbance at 492 nm was determined by an ELISA reader. The OD was plotted as a function of IL-18BP_a concentration and linearity was observed in a range of 0.12-2.00 ng/ml IL-18BP_a (FIG. 11).

[0397] The IL-18BP_a standard curve in the absence of serum or in the presence of up to 20% human serum remained the same. Since IL-18BP_a binds IL-18 with a very high affinity, it was necessary to determine whether ELISA distinguishes between free and bound IL-18BP_a. It was found that IL-18 interferes with the IL-18 ELISA however, the interference is insignificant in serum samples. In 90% of sepsis patients, serum IL-18 levels were under 4 ng/ml (see below). Such sera samples have elevated IL-18BP_a, requiring a 5 to 20-fold dilution prior to measuring IL-18BP_a. At such dilutions, IL-18 interference with the IL-18BP_a ELISA is less than 10% (FIG. 24). Other related cytokines, including proIL-18, used at a 10-fold molar excess over IL-18BP_a, as well as a 200-fold excess of mature IL-1β do not interfere with the assay.

[0398] The most abundant isoform of human IL-18BP is IL-18BP_a, whereas isoforms b, c and d are minor splice variants (Novick et al. 1999 and Kim et al. 2000). IL-18BP_a exhibits the highest affinity for IL-18 (Kim et al. 2000). The affinity of IL-18BP_c for IL-18 is 10 fold lower, whereas isoforms b and d do not bind or neutralize IL-18. It was therefore important to determine the cross-reactivity of IL-18BP_a in ELISA with the other isoforms of IL-18BP. As shown in FIG. 25, human IL-18BP_c generated a 10-fold lower signal on a weight basis compared with huIL-18BP_a, whereas human isoforms b and d generated an insignificant signal in the ELISA.

Example 31: Generation of Anti IL-18BP Specific Antibodies

[0399] A rabbit was injected with rIL-18BP_a-His6 for the generation of polyclonal antibodies.

[0400] For the production of monoclonal antibodies, female Balb/C mice were injected 5 times with 10 µg of recombinant histidine tagged IL-18BP_a (rIL-18BP_a-His6). The mouse exhibiting the highest titer as determined by an inverted radioimmunoassay (IRIA example 32) or solid phase RIA (sRIA, example 32) was given a final boost intraperitoneally 4 and 3 days before fusion. Lymphocytes were prepared from spleen and fusion to NSW myeloma cells was performed. Hybridomas that were found to produce antibodies to IL-18BP_a were subcloned by limiting dilution. The clones generated were assayed by IRIA (Example 32). Representative hybridomas are shown in Table 6.

TABLE 6

Representative hybridomas producing anti IL-18BP _a		
Mab No.	IRIA ₁ (cpm)	sRIA ₂ (cpm)
148	23912	7941
297	1652	6483
369	316	12762
430	6887	3254
433	1009	15300

TABLE 6-continued

Representative hybridomas producing anti IL-18BP _a		
MAB No.	IRIA ₁ (cpm)	sRIA ₂ (cpm)
460	3199	4326
485	400	13010
582	15000	17897
601	1046	1928

^aPlates coated with goat anti mouse antibodies and hybridomas detected with ¹²⁵I IL-18BP_a.

^bPlates coated with urinary IL-18BP and hybridomas detected with ¹²⁵I goat anti mouse antibodies.

[0401] Hybridomas secreting antibodies directed against the histidine tag were discarded. Antibodies were further characterized by sRIA for their ability to recognize the naturally occurring IL-18BP (purified from urine). Binding characteristics of several positive hybridomas are shown in Table 3. Hybridomas No. 148, 430, 460 and 582 were positive with both, recombinant and urinary IL-18BP. Antibodies suitable for Western blotting, immunoprecipitation, immunoaffinity purification and for development of a specific ELISA were obtained. Positive clones producing anti IL-18BP specific antibodies were injected into Balb/C mice pre-primed with pristane for the production of ascites. The isotypes of the antibodies were defined with the use of an anti mouse IgG ELISA (Amersham-Pharmacia Biotech). Mab 582, which was highly reactive with the recombinant and the native IL-18BP (Table 6) was used for the assembly of the IL-18BP specific ELISA (Example 30).

[0402] Antibodies were tested by sRIA in the presence of IL-18 (Example 32) for their ability to recognize a complex of IL-18BP_a with IL-18. Most antibodies were unable to recognize IL-18BP when it was complexed to IL-18. Therefore, these antibodies appear to be directed against the ligand-binding domain of IL-18BP_a.

Example 32: Radioimmunoassays for the Detection of Anti IL-18BP Antibody Producing Cell Clones

[0403] Inverted Radioimmunoassay (IRIA).

[0404] PVC microtiter plates (Dynatech Laboratories, Alexandria, Va.) were coated overnight at 4° C. with affinity-purified goat anti-mouse F (ab) 2 antibodies (10 µg/ml, 100 µl/well; Jackson ImmunoResearch Labs). The plates were then washed twice with PBS containing 0.05% Tween 20 (washing solution) and blocked with BSA (0.5% in washing solution) for 2 hrs at 37° C. Hybridoma culture supernatants (100 µl/well) were added and the plates were incubated for 2 hrs at room temperature. The plates were washed three times, ¹²⁵I-rIL-18BP_a-His6 (105 cpm in 100 µl) was added to each well and the plates were incubated for 5 hrs at 22° C. The plates were then washed three times and individual wells were counted in a gamma counter. Hybridomas generating supernatants, which exhibited bound radioactivity at levels 5 folds higher than the negative control, were considered positive.

[0405] Solid Phase RIA (sRIA).

[0406] rIL-18BP_a (5 µg/ml) was used as the capture antigen and ¹²⁵I-goat anti mouse antibodies (100 µl, 10⁵ cpm) were used for detection. Blocking and washings were done as above. Positive clones were further screened by sRIA for their ability to recognize the IL-18BP isolated from concentrated human urine (Novick et al. 1999). Microtiter plates were coated with ligand-affinity purified urinary IL-18BP (1

µg/ml), blocked and washed as above. Hybridoma supernatants (100 µl) were added and detection was done with ¹²⁵I-goat anti mouse antibodies (10⁵ cpm in 100 µl).

[0407] sRIA for Antibodies Specific for the Ligand-binding Site of IL-18BP.

[0408] Microtiter plates were coated with either urinary IL-18BP (0.5 µg/ml) or rIL-18BP_a (5 µg/ml), blocked and washed as in sRIA. Recombinant human IL-18 (50 µl) was added to a final concentration of 1.5 µg/ml (15 min at room temperature), followed by the addition of hybridoma supernatants (50 µl). Detection was done with ¹²⁵I-goat anti mouse antibodies (10⁵ cpm in 100 µl, 2 h at room temperature).

Example 33: Myocardial IL-18 Content Following LPS

[0409] TNF α and IL-1 β have been implicated in cardiac dysfunction during sepsis, since IL-18 is a pro-inflammatory cytokine known to mediate the production of TNF α and IL-1 β , the concentration of IL-18 on LPS-induced cardiac dysfunction was tested.

[0410] Mice were treated with either vehicle (saline) or LPS. Hearts were harvested at 2, 4 and 6 hours following LPS administration and homogenized to determine myocardial IL-18 content by ELISA (Kit purchased from R&D Systems (Minneapolis Minn.)). The results in FIG. 26 show a two-fold increase in myocardial IL-18 content at 4 hours following LPS administration, which indicates that IL-18 is involved in cardiac dysfunction during sepsis.

Example 34: Effect of Neutralization of IL-18 on LPS-Induced Myocardial Dysfunction

[0411] In the previous example an increase in the IL-18 content has been found in cardiac dysfunction induced by LPS. Therefore, the effect of neutralization of IL-18 on LPS-induced myocardial dysfunction was evaluated.

[0412] Myocardial function was determined by an isovolumetric, nonrecirculating Langendorff technique as described previously (Meng et al. 1998). Isolated hearts were perfused with normothermic Krebs-Henseleit solution containing 11.0 mmol/l glucose, 1.2 mmol/l CaCl₂, 4.7 mmol/l KCl, 25 mmol/l NaHCO₃, 119 mmol/l NaCl, 1.17 mmol/l MgSO₄ and 1.18 mmol/l KH₂PO₄. A latex balloon was inserted in the left ventricle via the left atrium and inflated with water to achieve a left ventricular and diastolic pressure (LVEDP) of 10 mmHg. Pacing wires were attached to the right atrium and hearts were paced at 300 beats per minute. Coronary flow was quantified by collecting the effluent from the pulmonary arteries. Myocardial temperature was maintained at 37° C. Left ventricular developed pressure (LVDP), its first derivatives (+dP/dt, -dP/dt) and LVEDP were continuously recorded by a computerized pressure amplifier-digitizer (MacLab 8, AD Instrument, Cupertino, Calif.). After a 20 minutes equilibration period LVDP and +/-dP/dt were determined at varied LVEDP levels (10, 15, and 20 mmHg).

[0413] Following LPS administration, the left ventricular developed pressure (LVDP) was reduced by 38% compared to saline controls (36.3+/-1.9 mmHg vs. 59.1+/- mmHg, P<0.001, FIG. 27). Pre treatment of mice 30 minutes prior to LPS administration with normal rabbit serum (NRS) had minimal influence on LPS-induced myocardial dysfunction; however pretreatment with IL-18 neutralizing antibody abrogated dysfunction (FIG. 27). Coronary flow was not different between the groups (not shown).

[0414] The results indicate that neutralization of IL-18 protects against LPS-induced myocardial dysfunction.

[0415] The foregoing description of the specific embodiments reveal the general nature of the invention so that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without departing from the generic concept, and, therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology employed herein is for the purpose of description and not of limitation.

REFERENCES

- [0416] 1. Aestiz, M. E. and Rackow, E. C. Septic shock. *The Lancet*, 1998, 351: p. 1501-5.
- [0417] 2. Afford, S. C., et al., Distinct patterns of chemokine expression are associated with leukocyte recruitment in alcoholic hepatitis and alcoholic cirrhosis. *J Pathol*, 1998, 186(1): p. 82-9.
- [0418] 3. Anderson, D. M., et al., A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature*, 1997, 390(6656): p. 175-179.
- [0419] 4. Baroni, G. S., et al., Hepatic stellate cell activation and liver fibrosis are associated with necroinflammatory injury and Th1-like response in chronic hepatitis C. *Liver*, 1999, 19(3): p. 212-9.
- [0420] 5. Bird, G. L., et al., Increased plasma tumor necrosis factor in severe alcoholic hepatitis. *Ann Intern Med*, 1990, 112(12): p. 917-20.
- [0421] 6. Bolton, D. P., et al. *J. Clin. Hematol. Oncol.* 1980, 10: p. 39-48.
- [0422] 7. Botstein, D., et al. *Miami Wint. Symp.* 1982, 19: p. 265-274.
- [0423] 8. Broach, J. R., in "The Molecular Biology of the Yeast *Saccharomyces*: Life Cycle and Inheritance", Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., p. 445-470 (1981).
- [0424] 9. Broach, J. R., *Cell* 1982, 28: p. 203-204.
- [0425] 10. Byrn R. A. et al., *Nature (London)* 1990, 344: p. 667-670.
- [0426] 11. Car, B. D., V. M. Eng, B. Schnyder, L. Ozmen, S. Huang, P. Gallay, D. Heumann, M. Aguet, and B. Ryffel. Interferon gamma receptor deficient mice are resistant to endotoxic shock. *J. Exp. Med.* 1994, 179: p. 1437-44 issn: 0022-1007.
- [0427] 12. Chater, K. F. et al., in "Sixth International Symposium on Actinomycetales Biology", Akademiai Kaido, Budapest, Hungary (1986), p. 45-54.
- [0428] 13. Conti, B., J. W. Jahng, C. Tinti, J. H. Son, and T. H. Joh. Induction of interferon-gamma inducing factor in the adrenal cortex. *J. Biol. Chem.* 1997, 272: p. 2035-2037.
- [0429] 14. Dao, T., K. Ohashi, T. Kayano, M. Kurimoto, and H. Okamura. Interferon-gamma-inducing factor, a novel cytokine, enhances Fas ligand-mediated cytotoxicity of murine T helper 1 cells. *Cell-Immunol.* 1996, 173: p. 230-5 issn: 0008-8749.
- [0430] 15. Dayer, J-M. *J. Clin. Inv.* 1999, 104: p. 1337-1339
- [0431] 16. DiDonato, J A, Hayakawa, M, Rothwarf, D M, Zandi, E. and Karin, M. *Nature* 1997, 388: p. 16514-16517.
- [0432] 17. Dinarello C A. Biologic basis for interleukin-1 in disease. *Blood* 1996, 87: p. 2095-147.
- [0433] 18. Elliott, M. J., Maini, R. N., Feldmann, M., Long-Fox, A., Charles, P., Bijl, H., and Woody, J. N. *Lancet* 1994, 344: p. 1125-1127.
- [0434] 19. Endo, S., Inada, K., Yamada, Y., Wakabayashi, G., Ishikura, H., Tanaka, T. and Sato, S. Interleukin 18 (IL-18) levels in patients with sepsis. *J Med* 2000, 31: p. 15-20.
- [0435] 20. Engelmann, H., D. Aderka, M. Rubinstein, D. Rotman, and D. Wallach. A tumor necrosis factor-binding protein purified to homogeneity from human urine protects cells from tumor necrosis factor toxicity. *J. Biol. Chem.* 1989, 264: p. 11974-11980.
- [0436] 21. Engelmann, H., D. Novick, and D. Wallach. Two tumor necrosis factor-binding proteins purified from human urine. Evidence for immunological cross-reactivity with cell surface tumor necrosis factor receptors. *J. Biol. Chem.* 1990, 265: p. 1531-1536.
- [0437] 22. Fantuzzi, G., et al., IL-18 regulation of IFN-g production and cell proliferation as revealed in interleukin-1b converting enzyme-deficient mice. *Blood*, 1998, 91: p. 2118-2125.
- [0438] 23. Fiore, G., et al., Liver tissue expression of CD80 and CD95 antigens in chronic hepatitis C: relationship with biological and histological disease activities. *Microbios*, 1999, 97(386): p. 29-38.
- [0439] 24. Foss D. L., Zilliox M. J., Murtaugh M. P. Bacterially induced activation of interleukin-18 in porcine intestinal mucosa. *Vet Immunol Immunopathol.* 2001, 78: p. 263-77.
- [0440] 25. Galle, P. R., et al., Involvement of the CD95 (APO-1/Fas) receptor and ligand in liver damage. *J Exp Med*, 1995, 182(5): p. 1223-30.
- [0441] 26. Gracie, A J, Forsey, R J, Chan, W L, Gilmour, A, Leung, B P, Greer, A R, Kennedy, K, Carter, R, Wei, X-Q, Xu, D., Field, M, Foulis, A, Liew, F Y, and McInnes, I B. *J. Clin. Inv.* 1999, 104: p. 1393-1401
- [0442] 27. Grantham Science, 1974, 185: p. 862-864.
- [0443] 28. Grobmyer, S. R., Lin, E., Lowry, S. F., Rivadeneira, D. E., Potter, S., Bane, P. S., and Nathan, C. F. Elevation of IL-18 in Human Sepsis. *J Clin Immunol* 2000, 20: p. 212-5.
- [0444] 29. Grove, J., et al., Association of a tumor necrosis factor promoter polymorphism with susceptibility to alcoholic steatohepatitis [see comments]. *Hepatology*, 1997, 26(1): p. 143-6.
- [0445] 30. Gryczan, T., "The Molecular Biology of the Bacilli", Academic Press, NY (1982), p. 307-329).
- [0446] 31. Gutkind, J. S., et al., A novel c-fgr exon utilized in Epstein-Barr virus-infected B lymphocytes but not in normal monocytes. *Molec. Cell. Biol.*, 1991, 11: p. 1500-1507.
- [0447] 32. Gupta, D., Kirkland, T. N., Viriyakosol, S., and Dziarski, R. CD14 is a cell-activating receptor for bacterial peptidoglycan. *J Biol Chem* 1996, 271: p. 23310-6.
- [0448] 33. Hachicha, M., Rathanaswami, P., Naccache, P. H. and McColl, S. R. Regulation of chemokine gene expression in human peripheral blood neutrophils phagocytosing microbial pathogens. *J Immunol* 1998, 160: p. 449-54.
- [0449] 34. Harada, K., et al., In situ nucleic acid hybridization of cytokines in primary biliary cirrhosis: predominance of the Th1 subset. *Hepatology*, 1997, 25(4): p. 791-6.
- [0450] 35. Henderson, B., Poole, S., and Wilson, M. Bacterial modulins: a novel class of virulence factors which

- cause host tissue pathology by inducing cytokine synthesis. *Microbiol. Rev* 1996, 60: p. 316-41.
- [0451] 36. Heremans, H., J. Van Damme, C. Dillen, R. Dijkmans, and A. Billiau. Interferon gamma, a mediator of lethal lipopolysaccharide-induced Shwartzman-like shock reactions in mice. *J. Exp. Med.* 1990, 171: p. 1853-69 issn: 0022-1007.
- [0452] 37. Hill, D. B., et al., Increased plasma interleukin-6 concentrations in alcoholic hepatitis. *J Lab Clin Med*, 1992, 119(5): p. 547-52.
- [0453] 38. Hill, D. B., L. S. Marsano, and C. J. McClain, Increased plasma interleukin-8 concentrations in alcoholic hepatitis. *Hepatology*, 1993, 18: p. 576-580.
- [0454] 39. Hiramatsu, N., et al., Immunohistochemical detection of Fas antigen in liver tissue of patients with chronic hepatitis C. *Hepatology*, 1994, 19(6): p. 1354-9.
- [0455] 40. Huang, Y. S., et al., Serum levels of interleukin-8 in alcoholic liver disease: relationship with disease stage, biochemical parameters and survival. *J Hepatol*, 1996, 24(4): p. 377-84.
- [0456] 41. Iio, S., et al., Serum levels of soluble Fas antigen in chronic hepatitis C patients. *J Hepatol*, 1998, 29(4): p. 517-23.
- [0457] 42. Izaki, K. *Jpn. J. Bacteriol.* 1978, 33: p. 729-742.
- [0458] 43. John, J. F., et al. *Rev. Infect. Dis.* 1986, 8: p. 693-704.
- [0459] 44. Kahiwamura, S., Okamura, H. *Nippon. Rinsho.* 1998, 56, p. 1798-1806.
- [0460] 45. Karp, C. L., Biron, C. A., and Irani, D. N. Interferon b in multiple sclerosis: is IL-12 suppression the key? *Immunology Today* 2000, 21: p. 24-28.
- [0461] 46. Kendall, K. J. et al. *J. Bacteriol.* 1987, 169: p. 4177-4183.
- [0462] 47. Kim S. H., Eisenstein M., Reznikov L., Fantuzzi G., Novick D., Rubinstein M., Dinarello C. A. Structural requirements of six naturally occurring isoforms of the IL-18 binding protein to inhibit IL-18. *Proc Natl Acad Sci USA* 2000, 97: p. 1190-1195.
- [0463] 48. Knight D. M., Trinh H., Le J., Siegel S., Shealy D., McDonough M., Scallan B., Moore M. A., Vilcek J., Daddona P., et al. Construction and initial characterization of a mouse-human chimeric anti-TNF antibody. *Mol Immunol* 1993 Nov. 30(16): p. 1443-53
- [0464] 49. Kohno, K., J. Kataoka, T. Ohtsuki, Y. Suemoto, I. Okamoto, M. Usui, M. Ikeda, and M. Kurimoto. IFN-gamma-inducing factor (IGIF) is a costimulatory factor on the activation of Th1 but not Th2 cells and exerts its effect independently of IL-12. *J. Immunol.* 1997, 158: p. 1541-1550.
- [0465] 50. Lee, M., et al., Expression of Th1 and Th2 type cytokines responding to HBsAg and HBxAg in chronic hepatitis B patients. *J Korean Med Sci*, 1999, 14(2): p. 175-81.
- [0466] 51. Lukkari, T. A., et al., Short-term ethanol exposure increases the expression of Kupffer cell CD14 receptor and lipopolysaccharide binding protein in rat liver. *Alcohol Alcohol*, 1999, 34(3): p. 311-9.
- [0467] 52. Luo, K. X., et al., In situ investigation of Fas/FasL expression in chronic hepatitis B infection and related liver diseases. *J Viral Hepat*, 1997, 4(5): p. 303-7.
- [0468] 53. Maliszewski, C. R., T. A. Sato, T. Vanden Bos, S. Waugh, S. K. Dower, J. Slack, M. P. Beckmann, and K. H. Grabstein. Cytokine receptors and B cell functions. I. Recombinant soluble receptors specifically inhibit IL-1- and IL-4-induced B cell activities in vitro. *J. Immunol.* 1990, 144: p. 3028-3033.
- [0469] 54. Maniatis, T., in "Cell Biology: A Comprehensive Treatise, Vol. 3: Gene Expression", Academic Press, NY, pp. 563-608 (1980).
- [0470] 55. Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, 1982.
- [0471] 56. Martinez, F., et al., Ethanol and cytokine secretion. *Alcohol*, 1992, 9(6): p. 455-8.
- [0472] 57. McClain, C. J., et al., Tumor necrosis factor and alcoholic liver disease. *Alcohol Clin Exp Res*, 1998, 22(5 Suppl): p. 248S-252S.
- [0473] 58. McClain, C. J. and D. A. Cohen, Increased tumor necrosis factor production by monocytes in alcoholic hepatitis. *Hepatology*, 1989, 9(3): p. 349-51.
- [0474] 59. Micallef, M. J., T. Ohtsuki, K. Kohno, F. Tanabe, S. Ushio, M. Namba, T. Tanimoto, K. Torigoe, M. Fujii, M. Ikeda, S. Fukuda, and M. Kurimoto. Interferon-gamma-inducing factor enhances T helper 1 cytokine production by stimulated human T cells: synergism with interleukin-12 for interferon-gamma production. *Eur J Immunol* 1996, 26: p. 1647-51 issn: 0014-2980.
- [0475] 60. Mizushima, S., and Nagata, S. pEF-BOS, a powerful mammalian expression vector. *Nucleic Acid Res.* 1990, 18: p. 5322-5328.
- [0476] 61. Nanji, A. A., et al., Activation of nuclear factor κ B and cytokine imbalance in experimental alcoholic liver disease in the rat. *Hepatology*, 1999, 30(4): p. 934-43.
- [0477] 62. Nakamura, K., H. Okamura, K. Nagata, T. Komatsu, and T. Tamura. Purification of a factor which provides a costimulatory signal for gamma interferon production. *Infect-Immun* 1993, 61: p. 64-70 issn: 0019-9567.
- [0478] 63. Nakamura, K., H. Okamura, M. Wada, K. Nagata, and T. Tamura. Endotoxin-induced serum factor that stimulates gamma interferon production. *Infect-Immun* 1989, 57: p. 590-5 issn: 0019-9567.
- [0479] 64. Nakamura, S., Otani, T., Ijiri, Y., Otoda, R., Kurimoto, M. and Orita, K. IFN- γ -dependent and -independent mechanisms in adverse effects caused by concomitant administration of IL-18 and IL-12. *The Journal of Immunology* 2000, 164: p. 3330-6.
- [0480] 65. Nakanishi K., Yoshimoto T., Tsutsui H., Okamura H. Interleukin-18 is a unique cytokine that stimulates both Th1 and Th2 responses depending on its cytokine milieu. *Cytokine Growth Factor Rev* 2001, 12: p. 53-72.
- [0481] 66. Nathans A. B., Marshall J C. Sepsis, SIRS, and MODS: what's in a name? *World J Surg* 1996, 20: p. 386-91.
- [0482] 67. Netea, M. G., Fantuzzi, G., Kullberg, B. J., Stuyt, R. J. L., Pulido, E. J., McIntyre, Jr. R. C., Joosten, L. A. B., Van der Meer, J. W. M. and Dinarello, C. A. Neutralization of IL-18 reduces neutrophil tissue accumulation and protects mice against lethal *Escherichia coli* and *Salmonella typhimurium* endotoxemia. *The Journal of Immunology*, 2000, p. 2644-2649.
- [0483] 68. Nishimura, T. and A. Ohta, A critical role for antigen-specific Th1 cells in acute liver injury in mice. *J. Immunol*, 1999, 162: p. 6503-6509.
- [0484] 69. Novick, D., B. Cohen, and M. Rubinstein. The Human Interferon alpha/beta Receptor-Characterization and Molecular Cloning. *Cell* 1994, 77: p. 391-400.

- [0485] 70. Novick, D., B. Cohen, and M. Rubinstein. Soluble Interferon- α Receptor Molecules Are Present in Body Fluids. *FEBS Lett* 1992, 314: p. 445-448.
- [0486] 71. Novick, D., H. Engelmann, D. Wallach, and M. Rubinstein. Soluble cytokine receptors are present in normal human urine. *J. Exp. Med.* 1989, 170: p. 1409-1414.
- [0487] 72. Novick, D., Kim, S.-H., Fantuzzi, G., Reznikov, L., Dinarello, C., and Rubinstein, M. *Immunity* 1999, 10: p. 127-136.
- [0488] 73. Ohlinger, W., et al., Immunohistochemical detection of tumor necrosis factor- α , other cytokines and adhesion molecules in human livers with alcoholic hepatitis. *Virchows Arch A Pathol Anat Histopathol*, 1993, 423 (3): p. 169-76.
- [0489] 74. Okamura, H., H. Tsutsui, T. Komatsu, M. Yutsudo, A. Hakura, T. Tanimoto, K. Torigoe, T. Okura, Y. Nukada, K. Hattori, K. Akita, M. Namba, F. Tanabe, K. Konishi, S. Fukuda, and M. Kurimoto. Cloning of a new cytokine that induces IFN- γ production by T cells. *Nature* 1995, 378: p. 88-91.
- [0490] 75. Okamoto, T., et al., Induction of Fas ligand and Fas antigen mRNA expressions in interferon- γ transgenic mouse liver. *Jpn J Pharmacol*, 1998, 78(2): p. 233-5.
- [0491] 76. Okazaki, M., et al., Hepatic Fas antigen expression before and after interferon therapy in patients with chronic hepatitis C. *Dig Dis Sci*, 1996, 41(12): p. 2453-8.
- [0492] 77. Okamoto, T., K. Yamamura, and O. Hino, The mouse interferon- γ transgene chronic hepatitis model (Review). *Int J Mol Med*, 1999, 3(5): p. 517-20.
- [0493] 78. Okusawa, S., Gelfand, J. A., Ikejima, T., Connolly, R. J. and Dinarello, C. A. Interleukin 1 induces a shock-like state in rabbits. Synergism with tumor necrosis factor and the effect of cyclooxygenase inhibition *J Clin Invest* 1988, 81: p. 1162-72.
- [0494] 79. Olee T., Hashimoto S., Quach J., Lotz M. *J Immunol* 1999, 162(2): p. 1096-100
- [0495] 80. Parnet, P., Garka, K. E., Bonnert, T. P., Dower, S. K., and Sims, J. E. *J. Biol. Chem.* 196, 271: p. 3967-3970.
- [0496] 81. Plater-Zyberk C., Bonnefoy J. Y. Marked amelioration of established collagen-induced arthritis by treatment with antibodies to CD23 in vivo. *Nat Med* 1995, 1: p. 781-785.
- [0497] 82. Saha N., Moldovan F., Tardif G., Pelletier J. P., Cloutier J. M., Martel-Pelletier J. *Arthritis Rheum* 1999, 42(8): p. 1577-87.
- [0498] 83. Sakao, Y., Takeda, K., Tsutsui, H., Kaisho, T., Nomura, F., Okamura, H., Nakanishi, K. and Akira S. IL-18-deficient mice are resistant to endotoxin-induced liver injury but highly susceptible to endotoxin shock. *Int Immunol* 1999, 3: p. 471-80.
- [0499] 84. Skeen, M. J. and Ziegler, H. K. Activation of gamma delta T cells for production of IFN- γ is mediated by bacteria via macrophage-derived cytokines IL-1 and IL-12. *J Immunol*. 1995, 154: p. 5832-41.
- [0500] 85. Rothe, H., N. A. Jenkins, N. G. Copeland, and H. Kolb. Active stage of autoimmune diabetes is associated with the expression of a novel cytokine, IGIF, which is located near Idd2. *J. Clin. Invest.* 1997, 99: p. 469-74 issn: 0021-9738.
- [0501] 86. Sambrook, J., E. F. Fritsch, and M. T., *Molecular Cloning: A laboratory manual*. 2nd ed. ed. 1989, Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory.
- [0502] 87. Simonet, W. S., et al., Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell*, 1997, 89(2): p. 309-319.
- [0503] 88. Sheron, N., et al., Elevated plasma interleukin-6 and increased severity and mortality in alcoholic hepatitis. *Clin Exp Immunol*, 1991, 84(3): p. 449-53.
- [0504] 89. Sompayrac, L. H. and K. L. Danna, Efficient infection of monkey cells with DNA of simian virus 40. *Proc. Nat'l. Acad. Sci. USA*, 1981, 78: p. 7575-7578.
- [0505] 90. Sparks, C. A., et al., Assignment of the nuclear mitotic apparatus protein NuMA gene to human chromosome 11q13. *Genomics*, 1993, 17: p. 222-224.
- [0506] 91. Su, G. L., et al., CD14 and lipopolysaccharide binding protein expression in a rat model of alcoholic liver disease. *Am J Pathol*, 1998, 152(3): p. 841-9.
- [0507] 92. Taieb, J., et al., Raised plasma soluble Fas and Fas-ligand in alcoholic liver disease [letter]. *Lancet*, 1998, 351(9120): p. 1930-1.
- [0508] 93. Triantaphyllopoulos, K. A., Williams, R., Taylor, H., and Chemakovsky, Y. *Arthritis and Rheumatism* 1999, 42: p. 90-99.
- [0509] 94. Tsutsui, H., K. Nakanishi, K. Matsui, K. Higashino, H. Okamura, Y. Miyazawa, and K. Kaneda. IFN- γ -inducing factor up-regulates Fas ligand-mediated cytotoxic activity of murine natural killer cell clones. *J. Immunol.* 1996, 157: p. 3967-73 issn: 0022-1767.
- [0510] 95. Tsuij, H., Mukaida, N., Harada A., Kaneko, S., Matsushita, E., Nakanuma, Y., Tsutsui, H., Okamura, H., Nakanishi, K., Tagawa, Y., Iwakura, Y., Kobayashi, K., and Matsushima, K. *J. Immunol.* 1989, 162: p. 1049-1055.
- [0511] 96. Tucci, A., James, H., Chicheportiche, R., Bonnefoy, J. Y., Dayer, J. M., and Zubler, R. H., *J. Immunol.* 1992, 148: p. 2778-2784.
- [0512] 97. Ushio, S., M. Namba, T. Okura, K. Hattori, Y. Nukada, K. Akita, F. Tanabe, K. Konishi, M. Micallef, M. Fujii, K. Torigoe, T. Tanimoto, S. Fukuda, M. Ikeda, H. Okamura, and M. Kurimoto. 1996. Cloning of the cDNA for human IFN- γ -inducing factor, expression in *Escherichia coli*, and studies on the biologic activities of the protein. *J. Immunol.* 156:4274-4279. 34. Okayama, H. and Berg, P. A cDNA cloning vector that permits expression of cDNA inserts in mammalian cells. *Mol. Cell. Biol.* 1983, 3: p. 280-289.
- [0513] 98. Williams R. O., Mason L. J., Feldmann M., Maini R. N. Synergy between anti-CD4 and anti-tumor necrosis factor in the amelioration of established collagen-induced arthritis. *Proc Natl Acad Sci USA* 1994, Mar. 29 91(7), p. 2762-6.
- [0514] 99. Yasuda, H., et al., Identity of osteoclastogenesis inhibitory factor (OCIF) and osteoprotegerin (OPG): a mechanism by which OPG/OCIF inhibits osteoclastogenesis in vitro. *Endocrinology*, 1998, 139: p. 1329-1337.
- [0515] 100. Yoshimoto T., Takeda, K., Tanaka, T., Ohkusu, K., Kashiwamura, S., Okamura, H., Akira, S. and Nakanishi, K. *J. Immunol.* 1998, 161: p. 3400-3407.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 1348

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

gagaagagga cgttgtcaca gataaagagc caggctcacc agctcctgac gcatgcatca    60
tgaccatgag acacaactgg acaccagacc tcagcccttt gtgggtcctg ctctgtgtg    120
ccccagtcgt cactctcctg gtcagagcca cacctgtctc gcagaccacc acagctgccca    180
ctgctcagtg tagaagcaca aaggaccctt gccctcccca gcccacagtg tccccagcag    240
ctaagcagtg tccagcattg gaagtgaact ggccagaggt ggaagtgcc aagaatggaa    300
cgctgagctt atcctgtgtg gcctgcagcc gcttcccaaa cttcagcacc ctctactggc    360
tgggcaatgg ttcttctcatt gagcacctcc caggccgact gtgggagggg agcaccagcc    420
gggaacgtgg gagcacaggt acgcagctgt gcaaggcctt ggtgctggag cagctgaccc    480
ctgcctgca cagcaccaac ttctcctgtg tgctcgtgga ccctgaacag gttgtccagc    540
gtcagctcgt cctggcccag ctctgggctg ggctgagggc aaccttgccc cccacccaag    600
aagccctgcc ctccagccac agcagtcacc agcagcaggg ttaagactca gcacagggcc    660
agcagcagca caaccttgac cagagcttgg gtcctacctg tctacctgga gtgaacagtc    720
cctgactgcc tgtaggtcgc gtggatgcgc aacacacccc ctcttctctc gctttgggtc    780
ccttctctca ccaaattcaa actccattcc cacctaccta gaaaatcaca gctccttat    840
aatgcctcct cctcctgcca ttctctctcc acctatccat tagccttctt aacgtcctac    900
tcctcacact gctctactgc tcagaaacca ccaagactgt tgatgcctta gccttgcaact    960
ccagggccct acctgcattt cccacatgac tttctggaag cctcccaact attcttgctt   1020
ttcccagaca gtcaccaact ccatgtctct gctcatttag tcccgcttc ctcaccgccc   1080
cagcagggga acgtcaagc ctggttgaaa tgctgcctct tcagtgaagt cactctcttt   1140
cagctctggc cgcattctgc agacttecta tottcgtgct gtatgttttt tttttcccc   1200
ttcactctaa tggactgttc caggaagggt atgggggcac cagctgcttc ggatccacac   1260
tgtatctgtg tcatccccac atgggtctct ataaaggatt attcaatgga aaaaaaaaaa   1320
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa                                     1348

```

<210> SEQ ID NO 2

<211> LENGTH: 192

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp Val Leu Leu
1           5           10           15

Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro Val Ser
20           25           30

Gln Thr Thr Thr Ala Ala Thr Ala Ser Val Arg Ser Thr Lys Asp Pro
35           40           45

Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys Gln Cys Pro Ala

```

-continued

50	55	60	
Leu Glu Val Thr Trp	Pro Glu Val Glu Val	Pro Leu Asn Gly Thr Leu	
65	70	75	80
Ser Leu Ser Cys Val	Ala Cys Ser Arg Phe	Pro Asn Phe Ser Ile Leu	
	85	90	95
Tyr Trp Leu Gly Asn Gly	Ser Phe Ile Glu His Leu	Pro Gly Arg Leu	
	100	105	110
Trp Glu Gly Ser Thr	Ser Arg Glu Arg Gly	Ser Thr Gly Thr Gln Leu	
	115	120	125
Cys Lys Ala Leu Val	Leu Glu Gln Leu Thr	Pro Ala Leu His Ser Thr	
	130	135	140
Asn Phe Ser Cys Val	Leu Val Asp Pro Glu	Gln Val Val Gln Arg His	
	145	150	155
Val Val Leu Ala Gln	Leu Trp Ala Gly Leu	Arg Ala Thr Leu Pro Pro	
	165	170	175
Thr Gln Glu Ala Leu	Pro Ser Ser His Ser	Pro Gln Gln Gln Gly	
	180	185	190
 <210> SEQ ID NO 3			
<211> LENGTH: 1038			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
 <400> SEQUENCE: 3			
gagaagagga cggtgtcaca gataaagagc caggctcacc agctcctgac gcatgcatca			60
tgaccatgag acacaactgg acaccagacc tcagcccttt gtgggtcctg ctctgtgtg			120
cccacgtcgt cactctcctg gtcagagcca cacctgtctc gcagaccacc acagctgcca			180
ctgcctcagt tagaagcaca aaggaccctt gccctcccca gccccagtg ttcccagcag			240
ctaagcagtg tccagcattg gaagtacact ggccagaggt ggaagtgcc a ctgagctggg			300
ctgaggggcaa ccttgcccc caccgaagaa gccctgccct ccagccacag cagtccacag			360
cagcaggggtt aagactcagc acagggccag cagcagcaca accttgacca gagcttgggt			420
cctacctgtc tacctggagt gaacagtccc tgactgctg taggtgcgt ggatgcgcaa			480
cacacccct ccttctctgc ttgggtccc ttctctcacc aaattcaaac tccattccca			540
cctacctaga aaatcacagc ctcttataa tgctctctcc tcctgccatt ctctctccac			600
ctatccatta gccttcctaa cgtctactc ctcaactgc tctactgtc agaaaccacc			660
aagactgttg atgccttagc cttgcactcc agggccctac ctgcatttcc cacatgactt			720
tctggaagcc tcccaactat tcttgctttt ccagacagc tcccactccc atgtctctgc			780
tcatttagtc ccgtcttctt caccgcccc gaagggaac gctcaagcct ggttgaaatg			840
ctgcctcttc agtgaagtca tctcttttca gctctggcgg cattctgcag acttctctac			900
ttcgtgctgt atgttttttt ttccccctt cactetaatg gactgttcca gggaagggat			960
gggggcacca gctgcttcgg atccacactg tatctgtgtc atccccacat gggctctcat			1020
aaaggattat tcaatgga			1038

<210> SEQ ID NO 4
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 4

```

Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp Val Leu Leu
1           5           10           15

Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro Val Ser
20           25           30

Gln Thr Thr Thr Ala Ala Thr Ala Ser Val Arg Ser Thr Lys Asp Pro
35           40           45

Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys Gln Cys Pro Ala
50           55           60

Leu Glu Val Thr Trp Pro Glu Val Glu Val Pro Leu Ser Trp Ala Glu
65           70           75           80

Gly Asn Leu Ala Pro His Pro Arg Ser Pro Ala Leu Gln Pro Gln Gln
85           90           95

Ser Thr Ala Ala Gly Leu Arg Leu Ser Thr Gly Pro Ala Ala Ala Gln
100          105          110

```

Pro

<210> SEQ ID NO 5

<211> LENGTH: 7063

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

gaattcgcg ccgctgcac gccagagggg ctaggatgag agacagaggg tgtgatggtg      60
gggtgctggga aatgtacccg accttggggc tgggtggctgg gggagtgggt agcctgggaa    120
aggccaggat gtggacggac tggataggca ttgagcctga agtgggtcaa cttgggggttc    180
cccagtgcct aggaaagttg tccccctgaa tgtcagtgtg aaggtgaagg aggaagcaga    240
tgccctgttca tatggaacaa aagacctggc tgtgaagagg ggaggcggac accaaagtcc    300
tgacacttgg gcgggacaga attgatctgt gagagactca tctagtccat accctagggtg    360
accctggggg tggctggggg gtagattaga gatcccagtc tgggtatcctc tggagagtag    420
gagtgccagg agctgaagggt ttctggccac tgaactttgg ctaaagcaga ggtgtcacag    480
ctgctcaaga ttccctggtt aaaaagtga aagtgaatag agggtcgggg cagtgccttc    540
ccagaaggat tgctcggcat cctgcccttc ccagaagcag ctctggtgct gaagagagca    600
ctgcctccct gtgtgactgg gtgagtcct attctctctt tgggtctcaa ttttgccttc    660
cctaataaag gggtaaagatt ggactaggta agcatcttac aaccatttgt ggtcatgaga    720
gtcgggggtg ggaaggattg tcaactgacc ccccagctc tgtttctaag tgctgaaaga    780
gctccaggct atgctacggg aggagaagcc agctactgag gaaaagccag ctactgagaa    840
aaagcgggag tggtttacca ttctcctccc ccacctttca ccagagaaga ggacgttgtc    900
acacataaag agccaggctc accagctcct gacgcatgca tcatgaccat gagacacaa    960
tggacaccag acctcagccc tttgtgggtc ctgctcctgt gtgccacgt cgctcacttc   1020
ctggtcagag ccacacctgt ctgcagacc accacagctg ccactgcctc agttagaagc   1080
acaaaggacc cctgcccctc ccagccccc gtgttcccag cagctaagca gtgtccagca   1140
ttggaagtga cctggccaga ggtggaagt ccaactgaatg gaacgctgag cttatcctgt   1200
gtggcctgca gccgtctccc caacttcagc atcctctact ggctgggcaa tggttccttc   1260
attgagcacc tcccaggccg actgtgggag gggagcacca gccgggaacg tgggagcaca   1320

```

-continued

ggtagcgagc	tgtgcaaggc	cttgggtgctg	gagcagctga	ccctgcccct	gcacagcacc	1380
aacttctcct	gtgtgctcgt	ggacctgaa	caggttgctc	agcgtcacgt	cgctcctggc	1440
cagctctggg	tgaggagccc	aaggagaggc	ctccaggaac	aggaggagct	ctgcttccat	1500
atgtggggag	gaaaggggtg	gctctgccag	agcagcctgt	gaactaatgc	ccagcattcc	1560
tcaaggtcag	ccagacaaaa	aggaacttag	gtcttgggca	gaggaggtgt	agcctggggc	1620
aaagtgatga	gatgtccctc	ctttccttgg	cctgatcctt	gtctgccttc	acttccctag	1680
gctgggctga	gggcaacctt	gccccccacc	caagaagccc	tgccctccag	ccacagcagt	1740
ccacagcagc	agggtaaga	ctcagcacag	ggccagcagc	agcacaacct	tgaccagagc	1800
ttgggtccta	cctgtctacc	tggagtgaac	agtcctgac	tgctgtagg	ctgcgtggat	1860
gcgcaacaca	ccccctcctt	ctctgctttg	ggtcccttct	ctcaccaaat	tcaaactcca	1920
ttccaccta	cctagaaaat	cacagcctcc	ttataatgcc	tcctcctcct	gccattctct	1980
ctccacctat	ccattagcct	tcctaacgtc	ctactcctca	cactgctcta	ctgctcagaa	2040
accaccaaga	ctgttgatgc	cttagccttg	cactccaggg	ccctacctgc	atttcccaca	2100
tgactttctg	gaagcctccc	aactattctt	gcttttccca	gacagctccc	actcccatgt	2160
ctctgctcat	ttagtcccgt	cttctcacc	gccccagcag	gggaacgctc	aagcctgggt	2220
gaaatgctgc	ctcttcagtg	aagtcatcct	ctttcagctc	tggccgcatt	ctgcagactt	2280
cctatcttgc	tgctgtatgt	tttttttttc	ccccttcact	ctaattggact	gttccagggg	2340
agggatgggg	gcagcagctg	cttcggatcc	acactgtatc	tgtgtcatcc	ccacatgggt	2400
cctcataaag	gattattcaa	tggaggcatc	ctgacatctg	ttcatttagg	cttcagttcc	2460
actcccagga	actttgcctg	tcccacgagg	gagtatggga	gagatggact	gccacacaga	2520
agctgaagac	aacacctgct	tcagggggaa	acaggcgctt	gaaaaagaaa	agagagaaca	2580
gcccataaat	ctccccggga	gcagaggcca	ctaattggaga	gtgggaagag	cctgggaaaga	2640
tgtggcctca	ggaaaaggga	tgagagaaag	gaggtggtat	ggaagactca	gcaggaacaa	2700
ggtaggcttc	aaagagccta	tattcctctt	tttccacac	cgatcaagtc	aactcagtac	2760
tcacgggaga	aaaatagact	ttatttaca	gtaataacat	ttagaaaaga	tccatccccg	2820
gcccttaaaa	accttcccat	cactccaaat	cccacccag	tgcaagtctg	gggaaggtag	2880
ggtgtgagct	gctgtgaag	gctgtccccc	aacccactc	ctgagacaca	gggcccattc	2940
gtcctgggaa	agagcctcct	ctggcaggtg	ctcccaccag	gtcagacca	gtcctggact	3000
tcaagagtga	gggcccctgc	tgggcccagc	caccaggaca	gcaggaacca	gggcctactc	3060
ctcttatggt	cccttctaga	tccagaggct	aagagggaaga	ctggccaggc	ccaaggaccc	3120
agccatcaaa	accagcctca	aatctggttg	tgatggagaa	gtgactttgc	tttaagaaaa	3180
aaggaggcaa	ggtagggaga	gcgcccacac	tgtccatgct	ccaggccccc	tgggcccagct	3240
ccgagaaggc	gccagtgaag	gaccagggac	caggccaggg	tgcgggcagg	catcactgtc	3300
tctagggggt	tggctactgt	tggcctggga	gctgagagaa	ggcactgaga	gggacagtag	3360
gcggagggac	aggtgacggc	agcatcgggg	acacagggtg	ggccactcac	tggtactggc	3420
ccttttagtg	tttgcttgaa	agagacacag	tcacatggcc	agatgagaac	ttgcgatact	3480
agcctgcacc	cactggctgg	gaagatctct	tcctgctccc	acgcccctgt	ctggatcccc	3540
tcccttggtg	gccccagggt	tatcagttgc	tggctgtgcc	tgagcagctc	tgggtgctct	3600

-continued

ccatgagaat	ggggccatct	gtcttctctc	cttgagagagg	agctaccagg	acagggacac	3660
ctcttacccc	acaccctcca	gcagcctggc	gtggcccat	cttgatgct	acttggtggg	3720
gcggtctggg	gggtgccc	gctctcatcg	ggtttccctc	ccccatcctg	ccagtgcctc	3780
taccttgccc	ttggctcgag	gggtggcacc	aatggcggca	gcagtggcgg	cgctggctgt	3840
gggtgtggca	atgcgcggag	aacggcgggt	tccactgcga	gtgtggggg	aagccttggg	3900
cagggccttc	tttgaggctc	cccgcgcgag	aaggtgttc	cctagcttct	tgggtgtgtt	3960
gaggatgctg	aaggccatcg	actggcgcg	gtcagcctgc	aaggaagggc	tgtcagaccg	4020
ggagacccaa	tgtctgcctc	ccaggccagc	gtgtgtgtcc	acgtgtacc	agcaaggctc	4080
cggcaggcgg	tgccttcac	ccccttcagc	cccagcctca	cctgtttagt	agaagctgga	4140
gctgctttct	tctgggcctc	agtagtgctc	tgtttgcgcc	cttcatgtcg	gtctcgggga	4200
gtcatggggc	gtgggaaaca	gctgtgtggc	ttcttagact	atggagaaga	ggacagttag	4260
gcagacagta	gcaagaggag	tcacatctga	agccagggtg	cttgtcctct	cagagctgag	4320
tggaccttgt	aagtcaacgt	gcaacctgct	ccccttccca	actctgggcc	agatccttcc	4380
cttcccaaca	gttcccatcc	atgggtcagg	cccttgagga	gagggaaaga	gagggggaag	4440
tgagggaaag	agagagaagg	ctccctttag	tccttggtga	gctgggcctg	acctgagcac	4500
agtgtgggag	taacaccacg	gagccaccgc	gcctacctca	ggagtccag	ggccctggtg	4560
gggtcttagg	gagaccctgt	tgcgtgctg	cgggtggtg	atgccagtgc	cctcggttat	4620
ctggattggc	tgcattgctg	ctcggcgcag	ggctctcttg	gggtctccag	tttcatctc	4680
ctcatctgtg	atggtgcccc	ggctcaggga	aggctgcatt	gggtgaagag	gtggtcagtg	4740
gaccatagct	gtatggagat	ggaggaggac	ctggggctgt	tccagaactc	tacactcgcc	4800
cgacacttat	ggtcgggacc	cttctgcct	acgaggtaga	aagacacaag	cctcctttcc	4860
tgttctgctt	tctacctaa	ccctgggcaa	atggcacaag	cagtgcagtc	ctgaccagat	4920
tcctctctga	gctctgcct	accccccagg	acttcacccc	tgagtgcct	ccagctgtct	4980
gttccacctg	gaacatgaga	aggtcacccc	ttccctctt	cggccagtca	gtgatccagg	5040
gccctagtgc	tcaggctaga	tcagcagggt	ggattccaag	gaagggcagg	gatgggaggc	5100
cctgcacagt	gacccagcgc	ctcaccctgg	actccaggga	tagcaggtct	tcagatgtgg	5160
ggggcacact	cgattgcgct	gctgcagctc	tgcaatgcgg	ttccagtcac	ccagctgctc	5220
aggctcatcc	tggcaagtgc	ccatgtagaa	gctgttctt	cctgtggaag	gcagggaagt	5280
gggaacaaat	gagcctggag	tcggcaggtc	acctcctggc	cctggcatct	tgccagcctt	5340
tgtgtccacc	taccccataa	acttgaagcc	cggcacacca	gtctgattca	gtgccgcagg	5400
tgcaggagta	cggcacacag	actatttcta	tcctaggggc	ttgctcacca	ccttctccct	5460
ggagagggca	gaagagggtc	cacgcagaga	ctgctactac	atcttattca	cctgccaagg	5520
cttggtggcc	aacaccacga	ggaacaaatt	aaggaccggg	aattaattcc	caggggctcc	5580
ctggtgcccc	aaggacaaga	gcttccaaga	agagtctggc	cagcctggcc	tttccagcag	5640
cccatcaccg	cctgagaagg	gcattggagg	ctcccccacg	ctaagtgtca	caattgtgct	5700
gggaatcccc	ggcccttaac	tctggctaag	agtgccccc	acacagccag	cccctagatg	5760
ggcaggtaag	gaaggccctg	aggctgcagg	aaggaggggc	agggtggagct	ggatggtagc	5820
aaggaggcca	gccttgatt	tttaaaaagc	tttctcttt	tcctgtgccc	acgatccacc	5880

-continued

```

ttccagtcta attttggggt atagtaagtc cctgtagtcc cctcacctgg aggggccccca 5940
ctggacacccc cggcctggga acgacgagca gaactgcgag tgggtggggcg gtagccaggc 6000
aagctgagca gggctgagtt gccataatcg ggagaaccca ggcgagctag agactgagta 6060
gaggaggtgg ctgcgaggct agcctgggaa gcaggagcag accgcgtgct gtagaacgat 6120
gagttggcgc tgtctggctc ttccacatct agcttctgga agacagagtg aatctgttgc 6180
agtgtacagt ccctggcact gtacagaagc ttccatttcc cttccgaagc cctcagatcc 6240
cacggcacat ccatgtattc ccaactgctt tgcaaaggtc cttaaagtgt gtgtctgcaa 6300
gaaatggggc ttgtcgacag aagccctcac aaggtggtgc tgatgttgc aagactcttc 6360
tacgcatttt tttcatggag tctattcata atgctttgag gtagggaatg cagagtgttt 6420
atcgggcccat tttggagatg aagtgc aaataaagtg actagcccca aatcacactg 6480
ctaggaagta tcagagctgg ggctaggccc catgtctcct gactagtcag gctcatccca 6540
cagcctctgc tgtccctcag tccaaacttc cagggccctt accatgttcc agaacttccc 6600
ccaacttctt ggtagcaggg ggcaccctaa acacacaggt cccccctgct gtaccagggg 6660
ccccctctcc cctcctccca aacctcccct tcaagatgtg gaaacaaagg caagggcctg 6720
cagcctgtca ggcagtcac tgggcagcaa caatgcctct cagctgcatg gggcatgctg 6780
ggaggcacag gatgggctgc agcttcgcca cgttctctcc cttcacctg cacaggctca 6840
gtgtacgca tggagagaat gctagcctta gtcaggaggc agggatctaa tcctagccct 6900
gcctttttct tcagaagtgc ccttaaccaa gtcactgccc tttttaagac ctctcagctt 6960
tcccactgta acatggactg gctgctcatc cctccctgct cctgactgag tgcccagtgc 7020
aaagatgccc ttgagaggaa gtgggaattg ctgacctgac gac 7063

```

```

<210> SEQ ID NO 6
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 6

```

```

Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp Val Leu Leu
1           5           10           15
Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro Val Ser
20           25           30
Gln Thr Thr Thr Ala Ala Thr Ala Ser Val Arg Ser Thr Lys Asp Pro
35           40           45
Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys Gln Cys Pro Ala
50           55           60
Leu Glu Val Thr Trp Pro Glu Val Glu Val Pro Leu Asn Gly Thr Leu
65           70           75           80
Ser Leu Ser Cys Val Ala Cys Ser Arg Phe Pro Asn Phe Ser Ile Leu
85           90           95
Tyr Trp Leu Gly Asn Gly Ser Phe Ile Glu His Leu Pro Gly Arg Leu
100          105          110
Trp Glu Gly Ser Thr Ser Arg Glu Arg Gly Ser Thr Gly Thr Gln Leu
115          120          125
Cys Lys Ala Leu Val Leu Glu Gln Leu Thr Pro Ala Leu His Ser Thr
130          135          140

```

-continued

Asn	Phe	Ser	Cys	Val	Leu	Val	Asp	Pro	Glu	Gln	Val	Val	Gln	Arg	His
145					150					155					160
Val	Val	Leu	Ala	Gln	Leu	Trp	Val	Arg	Ser	Pro	Arg	Arg	Gly	Leu	Gln
				165					170					175	
Glu	Gln	Glu	Glu	Leu	Cys	Phe	His	Met	Trp	Gly	Gly	Lys	Gly	Gly	Leu
				180				185					190		
Cys	Gln	Ser	Ser	Leu											
				195											

<210> SEQ ID NO 7
 <211> LENGTH: 1360
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

gcggccgcgt cgaccacgca gctaaacaca gctaacttga gtcttgagc tcctaaaggg      60
aagcttcttg aaaggaaggc tcttcaggac ctcttaggag ccaagaaga ggacgttgtc      120
acagataaag agccaggctc accagctcct gacgcatgca tcattgacct gagacacaac      180
tggaacaccag acctcagccc tttgtgggtc ctgctcctgt gtgccacgt cgtcactctc      240
ctggtcagag ccacacctgt ctgcagacc accacagctg cactgcctc agttagaagc      300
acaaaggacc cctgcccctc ccagccccc gtgttcccag cagctaagca gtgtccagca      360
ttggaagtga cctggccaga ggtggaagtg cactgaatg gaacgctgag cttatcctgt      420
gtggcctgca gccgcttccc caacttcagc atcctctact ggctgggcaa tggttccttc      480
attgagcacc tccagggccg actgtgggag gggagcacca gccgggaacg tgggagcaca      540
ggctgggctg agggcaacct tgccccccac ccaagaagcc ctgccctcca gccacagcag      600
tccacagcag caggggtaag actcagcaca gggccagcag cagcacaacc ttgaccagag      660
cttggttctt acctgtctac ctggagtga cagtccctga ctgcctgtag gctgcgtgga      720
tgcgcaacac accccctcct tctctgcttt gggtcccttc tctcacaaa ttcaaactcc      780
attcccacct acctagaaaa tcacagcctc cttataatgc ctctcctcc tgcattctc      840
tctccacctc tccattagcc ttcctaactg cctactctc acactgctct actgctcaga      900
aaccaccaag actgttgatg ccttagcctt gcaactccag gccctacctg catttccac      960
atgactttct ggaagcctcc caactattct tgcttttccc agacagctcc cactcccatg     1020
tctctgctca tttagtcctc tcttcctcac cgcctcagca ggggaacgct caagcctggt     1080
tgaaatgctg cctcttcagt gaagtcaccc tctttcagct ctggccgcat tctgcagact     1140
tctatctctc gtgctgatg tttttttttt ccccttcac tctaattggac tgttccaggg     1200
aagggatggg ggcagcagct gcttcggatc cacactgtat ctgtgtcacc cccacatggg     1260
tcctcataaa ggattattca atggaggcat cctgacatct gtccatttag gcttcagttc     1320
cactcccagg aactttgcct gtcccacgag ggagtatggg                               1360

```

<210> SEQ ID NO 8
 <211> LENGTH: 161
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Met	Arg	His	Asn	Trp	Thr	Pro	Asp	Leu	Ser	Pro	Leu	Trp	Val	Leu	Leu
1				5					10					15	

-continued

Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro Val Ser
 20 25 30
 Gln Thr Thr Thr Ala Ala Thr Ala Ser Val Arg Ser Thr Lys Asp Pro
 35 40 45
 Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys Gln Cys Pro Ala
 50 55 60
 Leu Glu Val Thr Trp Pro Glu Val Glu Val Pro Leu Asn Gly Thr Leu
 65 70 75 80
 Ser Leu Ser Cys Val Ala Cys Ser Arg Phe Pro Asn Phe Ser Ile Leu
 85 90 95
 Tyr Trp Leu Gly Asn Gly Ser Phe Ile Glu His Leu Pro Gly Arg Leu
 100 105 110
 Trp Glu Gly Ser Thr Ser Arg Glu Arg Gly Ser Thr Gly Trp Ala Glu
 115 120 125
 Gly Asn Leu Ala Pro His Pro Arg Ser Pro Ala Leu Gln Pro Gln Gln
 130 135 140
 Ser Thr Ala Ala Gly Leu Arg Leu Ser Thr Gly Pro Ala Ala Ala Gln
 145 150 155 160
 Pro

<210> SEQ ID NO 9

<211> LENGTH: 7812

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```

gtcgacggta cccccgggaa agatttaata cgactcacta tagggcgggg cagaattgat      60
ctgtgagaga ctcatctagt tcatacccta ggtgacctg ggggtggcat ggggtagat      120
tagagatccc agtctggtat cctctggaga gtaggagtc caggagctga aggtttctgg      180
ccactgaact ttggctaaag cagaggtgtc acagctgctc aagattccct gggtaaaaag      240
tgaaagttaa atagagggtc ggggcagtgc tttcccagaa ggattgctcg gcatcctgcc      300
cttcccagaa gcagctctgg tgctgaagag agcactgcct cctgtgtga ctgggtgagt      360
ccatattctc tctttgggtc tcaattttgc cttccctaat gaaggggtaa gattggacta      420
ggtaagcatc ttacaacctt ttgtggtcat gagagctggg gtggggaagg attgtcactt      480
gaccccccca gctctgtttc taagtgtga aagagctcca ggctatgcta cgggaggaga      540
agccagctac tgaggaaaag ccagctactg agaaaaagcg ggagtgggtt accattctcc      600
tccccacctt ttcaccagag aagaggacgt tgtcacacat aaagagccag gctcaccagc      660
tcctgacgca tgcacatga ccatgagaca caactggaca ccaggtaggc cttggggcta      720
cgcatgggca ggccgggtag ggtgaggtct atgaacagaa tggagcaatg ggctaaccgc      780
gagccttcac tccaaggcaa accaccagc gcacctggtg ctgttgcttt aagaacctgg      840
gcagatattg tagctctggc tccagtctaa agcttctctg tactctgttc aataaagggc      900
taaggggtgg gtgctgaggg gtccctcttc ccgctctgat tcctggcta gaaccagac      960
atctctgggc tggagttaca tccttaccgc ggcagccac tctgtctcca gagccgtga     1020
cctgtaactg tcctttcttc agacctcagc cctttgtggg tcctgtctct gtgtgcccac     1080
gtcgtcactc tcctggtcag agccacacct gtctcgaga ccaccacagc tgccactgcc     1140

```

-continued

tcagttagaa	gcacaaagga	cccctgcccc	tcccagcccc	cagtgttccc	agcagctaag	1200
cagtgtccag	cattggaagt	gacctggcca	gaggtggaag	tgccactgag	taagaagcac	1260
agtggtagg	ggtaggctat	gggcacagag	gttcccaggg	tggggtgac	tccagagcgc	1320
cagtccccct	ctgcccattgt	accaccagct	gagccagctg	ggctgagcac	gcaccattct	1380
ccctccccc	cccagtgta	tgggtgcagg	cttggcgag	ctcccaagat	gctccctatc	1440
aaataggaca	gagaactcaa	gacataagta	atggtcacag	gacctcccag	agccttggtt	1500
gcagtggacc	ccaagggcag	cccctccacc	cagagcctgc	tggcctctgg	ccatctcaga	1560
ggagcagcag	ccatccagca	ctgcctctgt	cacctgggct	ccaagtac	cgaggctggg	1620
cactagaaaa	ggtcatccctg	aggagacagg	ttcagaagag	gattcatcac	gtgaaccaag	1680
gaccattcct	cacattcccc	gtgtttagg	ctagggcctc	tggagacaa	ctgcaattct	1740
gtaacggacg	ttccaccta	ggtggtgtgc	agagcagttc	tctaggttcc	agatgcatgg	1800
ggactggggg	gagctggcag	agagggcaca	gcagagcagg	gtaggggaag	ggcctgctct	1860
tctgaagagc	taactgctgc	ctgtgtccct	agatggaacg	ctgagcttat	cctgtgtggc	1920
ctgcagccgc	ttcccact	tcagcatcct	ctactggctg	ggcaatggtt	ccttcattga	1980
gcacctccca	ggccgactgt	gggaggggag	caccaggtga	gggtcgcagc	agccaggtgg	2040
gtgggaagga	agccttctgc	ggccttctca	tgacctttcc	ttcccttcgg	ctccagccgg	2100
gaacgtggga	gcacaggtac	gcagctgtgc	aaggccttgg	tgtgagca	gctgacctct	2160
gccctgcaca	gcaccaactt	ctcctgtgtg	ctcgtggacc	ctgaacaggt	tgtccagcgt	2220
cacgtcgtcc	tggccagct	ctgggtgagg	agcccaagga	gaggcctcca	ggaacaggag	2280
gagctctgct	tccatatgtg	gggaggaaag	ggtgggctct	gccagagcag	cctgtgaact	2340
aatgccacgc	attcctcaag	gtcagccaga	caaaaaggaa	cttaggtctt	gggcagagga	2400
ggtgtagcct	ggggcaaagt	gatgagatgt	ccctccttcc	cttggcctga	tcttgtctg	2460
ccttcacttc	cctaggtctg	gctgagggca	accttgcccc	ccaccaaga	agcctgcccc	2520
tccagccaca	gcagtcaca	gcagcagggt	taagactcag	cacagggccca	gcagcagcac	2580
aaccttgacc	agagcttggg	tcctacctgt	ctacctggag	tgaacagtc	ctgactgcct	2640
gtaggctgcg	tggatgcgca	acacaccccc	tccttctctg	ctttgggtcc	cttctctcac	2700
caaatcaaaa	ctccattccc	acctacctag	aaaatcacag	cctccttata	atgcctctct	2760
ctcctgccc	tctctctcca	cctatccatt	agccttctca	acgtctact	cctcacactg	2820
ctctactgct	cagaaaccac	caagactgtt	gatgccttag	ccttgcactc	cagggcccta	2880
cctgcatttc	ccacatgact	ttctggaagc	ctcccaacta	ttcttgcttt	tcccagacag	2940
ctccactccc	catgtctctg	ctcatttagt	ccgtcttccc	tcaccgcccc	agcaggggaa	3000
cgctcaagcc	tggttgaaat	gctgcctctt	cagtgaagtc	atcctcttcc	agctctggcc	3060
gcattctgca	gacttctctat	cttcgtgctg	tatgtttttt	tttcccccct	tcactcta	3120
ggactgttcc	aggggaaggga	tgggggcagc	agctgcttcc	gatccacact	gtatctgtgt	3180
catcccccaca	tgggtctcca	taaaggatta	ttcaatggag	gcacccctgac	atctgttcat	3240
ttaggcttca	gttccactcc	caggaaacttt	gcctgtccca	cgaggagagta	tgggagagat	3300
ggactgccac	acagaagctg	aagacaacac	ctgcttcagg	ggaacacagg	cgcttgaaaa	3360
agaaaagaga	gaacagccca	taatgctccc	cgggagcaga	ggccactaat	ggagagtggg	3420

-continued

aagagcctgg	aaagatgtgg	cctcaggaaa	agggatgaga	gaaaggaggt	ggtatggaag	3480
actcagcagg	aacaaggtag	gcttcaaaga	gcctatattc	ctctttttcc	cacaccgatc	3540
aagtcaactc	agtactcacg	ggagaaaaat	agactttatt	tacaagtaat	aacatttaga	3600
aaagatccat	ccccggccct	taaaaaacct	cccatcactc	caaatcccac	cccagtgcga	3660
gtctggggaa	ggtagggtgt	gagctgctgc	tgaaggctgt	cccccaacc	cactcctgag	3720
acacaggggc	catccgtcct	gggaaagagc	atcctctggc	aggtgctccc	accaggtcag	3780
accagtcct	ggacttcaag	agtgaggggc	cctgctgggc	ccagccacca	ggacagcagg	3840
aaccaggggc	tactcctctt	atggtccctt	ctagatccag	aggctaagag	gaagactggc	3900
caggcccaag	gacccagcca	tcaaaaccag	cctcaaactc	ggttgtgatg	gagaagtga	3960
tttgtcttaa	gaaaaaagga	ggcaaggtag	ggagagcgcc	cacactgtcc	atgctccagg	4020
ccccctgggc	cagctccgag	aaggcgccag	tgaaggacca	gggaccaggc	cagggtgcgg	4080
gcaggcatca	ctgtctctag	gggtttggct	actgttgccc	tgggagctga	gagaaggcac	4140
tgagagggac	agtaggcgga	ggaccagggt	acggcagcat	cggggacaca	ggtggggcca	4200
ctcactggta	ctggcccttt	agtgtcttgc	ctgaaagaga	cacagtcaca	tggccagatg	4260
agaacttgcg	atactagcct	gcacccactg	gctgggaaga	tctcttctcg	ctcccacgcc	4320
cctgtctgga	tccccctcct	tgtgagcccc	agggttatca	gttctgtggc	gtgectgagc	4380
agctctgggt	gctctccatg	agaatggggc	catctgtctt	ctctccttgg	agaggagcta	4440
ccaggacagg	gacacctctt	acccacaccc	ctccagcagc	ctggcggtgg	cccatcttgg	4500
atgtactctg	gtggggcggt	ctgggggggt	cccatgctct	catcggtttt	ccctcccca	4560
tcttgccagt	gcctctacct	tggccttggc	tcgaggggtg	gcaccaatgg	cggcagcagt	4620
ggcgcgctg	gctgtggtgg	tggcaatcg	cggagaacgg	cgggttccac	tgcgagtgtt	4680
gggggaagcc	ttggacaggg	ccttctttga	ggctccccgc	cgcagaaggc	tgttccctag	4740
cttcttgggt	gtgttaggga	tgtctgaagg	catcgactgg	cgcgggtcag	cctgcaagga	4800
agggctgtca	gaccgggaga	cccaatgctg	ccttcccagg	ccagcgtgct	gtgccacgct	4860
gtaccagcaa	ggtcccgcga	ggcgctgct	tcacccccct	tcagccccag	cctcacctgt	4920
ttagtagaag	ctggagctgc	tttcttctgg	gcctcagtag	tgtctgtttt	gcgcccttca	4980
tgtcgtctc	ggggagtcac	ggggcggtgg	aaacagctgg	tggccttctt	agactatgga	5040
gaagaggaca	gttaggcaga	cagttagcaag	aggagtcaca	tctgaagcca	ggtgtcttgt	5100
cctctcagag	ctgagtgga	cttgtaagtc	aacgtgcaac	ctgctcccc	tcccaactct	5160
gggocagatc	cttcccttcc	caacagttcc	catccatggg	tcaggccctt	ggagagaggg	5220
aaagagaggg	ggaagtgagg	gaaggagaga	gaaggctccc	tttagtcctt	ggtgagctgg	5280
gcctgacctg	agcacagtgc	tggagtaaca	cccaggagcc	accgcgccta	cctcaggagt	5340
tccagggccc	tgggtgggct	ctaggggagac	cgttttgccg	tgtgcccggg	tggatgatgc	5400
agtgcctcgc	gctatctgga	ttggctgcat	gtggctcgg	cgcagggtct	cttgggggtc	5460
tccagttttc	atctcctcat	ctgtgatggt	gcccaggctc	agggaaggct	gcatgggtgg	5520
aagaggtggt	cagtggacca	tagctgtatg	gagatggagg	aggacctggg	gctgttccag	5580
aactctacac	tcgcccagca	cttatggtcg	ggacccttcc	tgcctacgag	gtagaaagac	5640
acaagcctcc	tttctgttcc	tgctttctac	ctaagccctg	ggcaaatggc	acaagcagtg	5700

-continued

cagtcctgac cagattcctc tctgagctcc tgcctacccc cagggacttc acccctgagt	5760
gccctccagc tgtctgttcc acctggaaca tgagaaggtc accccttccc ctcttcggcc	5820
agtcagtgat ccaggccctc agtgctcagg ctatgacagc aggtgggatt ccaaggaagg	5880
gcagggatgg gaggcctcgc acagtgaccc caggcctcac cctggactcc agggatagca	5940
ggtcttcaga tgtggggggc aactcagatt gcgctgctgc agctctgcaa tgcggttcca	6000
gtcatccagc tgctcaggct catcctggca agtgcccatg tagaagctgt tccttcctgt	6060
ggaaggcagg gaagtgggaa caaatgagcc tggagtgggc aggtcacctc ctggccctgg	6120
catcttgcca gcctttgctg ccacctaccc cataaacttg aagcccggca caccagtctg	6180
attcagtgcc gcaggtcgag gagtacggca cacagactat ttctatccta ggggcttgct	6240
caccaccttc tccctggaga gggcagaaga ggtcacacgc agagactgct actacatctt	6300
attcacctgc caaggtctgg tggccaacac ccagaggaac aaattaagga ccgggaatta	6360
attcccaggg gtcctctggt gccc aaagga caagagcttc caagaagagt ctggccagcc	6420
tggcctttcc agcagcccat caccgctga gaaggcatg gaggactccc cacagctaag	6480
tgtcacaatt gtgctgggaa tcccggggcc ttaactctgg ctaagagtgc ccccaacaca	6540
gccagccctc agatgggcag gtaagggaagg ccctgaggct gcaggaagga ggggcagggtg	6600
gagctggatg gtagcaagga ggcacgctt ggatttttaa aaagctttcc tcttttccct	6660
gtgccacgat ccaccttcca gtctaatttt ggggtatagt aagtcctgt agtccctca	6720
cctggagggg cccactgga cccccggc tgggaacgac gagcagaact gcgagtgggtg	6780
ggcggttagc caggcaagct gagcagggtc gagggtccat aatcgggaga acccaggcga	6840
gctagagact gagtagagga ggtggctcgc aggttagcct gggaaagcagg agcagaccgc	6900
gtgctgtaga acgatgagtt ggcgctgtct ggctcttcca catctagctt ctggaagaca	6960
gagtgaatct gttgagtgat acagtccctg gcaactgtaca gaagcttccc attcccttcc	7020
gaagccctca gatccacagg cacatccatg tattcccaac tgctttgcaa aggtccttaa	7080
agtgtgtgtc tgcaaaaaat gggccttgct gacagaagcc ctcacaaggt ggtgctgatg	7140
ttgtcaagac tcttctacgc atttttttca tggagtctat tcataatgct ttgaggtagg	7200
gaatgcagag tgtttatcgg cccattttgg agatgaagtg caaagaaata aagtgactag	7260
ccccaaatca cactgctagg aagtatcaga gctggggcta ggcctcatgt ctctgacta	7320
gtcaggctca tcccacagcc tctgtgtcc ctcagtccaa acttcagggt cccttaccat	7380
gttcagaac tcccccaac ttcttggtag cagggggcac cctaaacaca cagggtcccc	7440
ctgctgtacc agggggcccc tctccctcc tcccaaacct ccccttcaag atgtggaaac	7500
aaaggcaagg gcctgcagcc tgtcaggcag tccactgggc agcaacaatg cctctcagct	7560
gcatggggca tgctgggagg cacaggatgg gctgcagctt cgccacgttc tctccctca	7620
ccctgcacag gctcagtgct acgcatggag agaagctag ccttagtcag gaggcaggga	7680
tctaactcta gccctgcctt tttcttcaga agtgccctta accaagtcac tgcctttttt	7740
aagacctctc agctttccca ctgtaacatg gactggctgc tcacccctcc ctgctcctga	7800
ctgagtgcgc ag	7812

<210> SEQ ID NO 10

<211> LENGTH: 40

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(8)
<223> OTHER INFORMATION: X is any amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: X is any amino acid

<400> SEQUENCE: 10

Thr Pro Val Ser Gln Xaa Xaa Xaa Ala Ala Xaa Ala Ser Val Arg Ser
1 5 10 15

Thr Lys Asp Pro Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys
20 25 30

Gln Cys Pro Ala Leu Glu Val Thr
35 40

<210> SEQ ID NO 11
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(9)
<223> OTHER INFORMATION: X is any amino acid

<400> SEQUENCE: 11

Thr Pro Val Ser Gln Gln Xaa Xaa Xaa Ala Ala Ala
1 5 10

<210> SEQ ID NO 12
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(8)
<223> OTHER INFORMATION: X is any amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: X is any amino acid

<400> SEQUENCE: 12

Thr Pro Val Ser Gln Xaa Xaa Xaa Ala Ala Xaa Ala Ser Val Arg Ser
1 5 10 15

Thr Lys Asp Pro Cys Pro Ser Gln Pro Pro Val Phe Pro Ala Ala Lys
20 25 30

Gln Cys Pro Ala Leu Glu Val Thr
35 40

<210> SEQ ID NO 13
<211> LENGTH: 50
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: X is any amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE

-continued

<222> LOCATION: (11)..(11)

<223> OTHER INFORMATION: X is any amino acid

<400> SEQUENCE: 13

Val Thr Leu Leu Val Arg Ala Thr Xaa Val Xaa Gln Thr Thr Thr Ala
1 5 10 15

Ala Thr Ala Ser Val Arg Ser Thr Lys Asp Pro Cys Pro Ser Gln Pro
 20 25 30

Pro Val Phe Pro Ala Ala Lys Gln Cys Pro Ala Leu Glu Val Thr Trp
 35 40 45

Pro Glu
 50

<210> SEQ ID NO 14

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically Synthesized

<400> SEQUENCE: 14

Glu Phe Gly Ala Gly Leu Val Leu Gly Gly Gln Phe Met
1 5 10

<210> SEQ ID NO 15

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (2)..(2)

<223> OTHER INFORMATION: X is any amino acid

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (4)..(4)

<223> OTHER INFORMATION: X is any amino acid

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: X is any amino acid

<400> SEQUENCE: 15

Ala Xaa Tyr Xaa Arg Ile Pro Ala Xaa Ala Ile Ala
1 5 10

<210> SEQ ID NO 16

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically Synthesized

<400> SEQUENCE: 16

tatatctaga gccaccatga gacacaactg gacacca

-continued

```

<210> SEQ ID NO 17
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically Synthesized

<400> SEQUENCE: 17
atatctagat taatgatgat gatgatgatg accctgctgc tgtggactgc

```

50

We claim:

1. A method for treatment and/or prevention of a sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS) comprising administering to a subject in need thereof a pharmaceutically effective amount of an inhibitor of IL-18 selected from a group consisting of caspase-1 (ICE) inhibitors, antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signaling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, and inhibitors of IL-18 production, and a pharmaceutically acceptable carrier.

2. The method according to claim 1, further comprising administering a therapeutically effective amount of an IL-12 inhibitor.

3. The method according to claim 1, further comprising administering a cytokine inhibitor selected from a group consisting of a Tumor Necrosis Factor (TNF) inhibitor, an IL-1 inhibitor and an IL-8 inhibitor.

4. The method according to claim 2, further comprising administering a cytokine inhibitor selected from a group consisting of a Tumor Necrosis Factor (TNF) inhibitor, an IL-1 inhibitor and an IL-8 inhibitor.

5. The method according to claim 3, wherein the TNF inhibitor is a soluble portion of TNFRI or TNFRII.

6. The method according to claim 1, wherein the method further comprises administering an interferon.

7. The method according to claim 2, wherein the method further comprises administering an interferon.

8. The method according to claim 3, wherein the method further comprises administering an interferon.

9. A method according to claim 6, wherein the interferon is interferon- α or interferon- β .

10. A method for the treatment and/or prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS) comprising administering to a subject in need thereof a pharmaceutically effective amount of a vector coding the sequence of an inhibitor of IL-18 selected from a group consisting of caspase-1 (ICE) inhibitors, antibodies against IL-18, antibodies against any of the IL-18 receptor subunits, inhibitors of the IL-18 signaling pathway, antagonists of IL-18 which compete with IL-18 and block the IL-18 receptor, inhibitors of IL-18 production, and IL-18 binding proteins, isoforms, muteins, fused proteins or circularly permuted derivatives thereof and a pharmaceutically acceptable carrier.

11. A method for the treatment and/or prevention of sepsis and other diseases characteristic to the Systemic Inflammatory Response Syndrome (SIRS) comprising administering to a subject in need thereof a pharmaceutically effective amount of

- i) a vector for inducing and/or enhancing the endogenous production of an inhibitor of IL-18 in a cell or
- ii) a cell that has been genetically modified to produce an inhibitor of IL-18.

12. The method according to claim 1, wherein the sepsis is related to cardiac dysfunction.

13. The method according to claim 10, wherein the fused protein comprises an Ig fusion.

14. The method according to claim 1, wherein the inhibitor of IL-18 is an anti IL-18 specific antibody selected from chimeric, humanized and human antibodies.

15. The method according to claim 2, wherein the IL-12 inhibitor is a neutralizing antibody.

* * * * *