



(86) Date de dépôt PCT/PCT Filing Date: 2004/09/10

(87) Date publication PCT/PCT Publication Date: 2005/11/17

(45) Date de délivrance/Issue Date: 2016/02/09

(85) Entrée phase nationale/National Entry: 2006/03/10

(86) N° demande PCT/PCT Application No.: US 2004/029575

(87) N° publication PCT/PCT Publication No.: 2005/108565

(30) Priorité/Priority: 2003/09/15 (US60/503,251)

(51) Cl.Int./Int.Cl. *C12N 9/24* (2006.01)


(72) Inventeurs/Inventors:
CLARKSON, KATHLEEN A., US;
FENEL, FRED, FI

(73) Propriétaire/Owner:
DANISCO US INC., US

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : XYLANASES MODIFIEES AYANT UNE MEILLEURE STABILITE THERMIQUE ET UNE MEILLEURE
STABILITE EN MILIEU ALCALIN

(54) Title: MODIFIED XYLANASES HAVING INCREASED THERMOSTABILITY AND ALKALINE STABILITY



(57) Abrégé/Abstract:

The invention is directed to modified xylanases having increased stability in harsh industrial environments, such as increased pH and/or temperature.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
17 November 2005 (17.11.2005)

PCT

(10) International Publication Number
WO 2005/108565 A2

(51) International Patent Classification⁷: **C12N 9/24**

(21) International Application Number:
PCT/US2004/029575

(22) International Filing Date:
10 September 2004 (10.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/503,251 15 September 2003 (15.09.2003) US

(71) Applicant (for all designated States except US): **GENENCOR INTERNATIONAL, INC.** [US/US]; 925 Page Mill Road, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CLARKSON, Kathleen, A.** [US/US]; 53 28th Street, San Francisco, CA 94110 (US). **FENEL, Fred** [FI/FI]; Vellamonkatu 29 D 41, FIN-00550 Helsinki 7 (FI).

(74) Agent: **ANDERTON, H., Thomas, Jr.**; Genencor International, Inc., 925 Page Mill Road, Palo Alto, CA 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: MODIFIED ENZYMES, METHODS TO PRODUCE MODIFIED ENZYMES AND USES THEREOF



(57) Abstract: The invention is directed to modified xylanases having increased stability in harsh industrial environments, such as increased pH and/or temperature.

WO 2005/108565 A2

WO 2005/108565 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

MODIFIED XYLANASES HAVING INCREASED THERMOSTABILITY AND ALKALINE STABILITY

5

FIELD OF THE INVENTION

The invention is directed to modified enzymes having increased stability in harsh industrial environments, such as increased pH and/or temperature.

10

BACKGROUND OF THE INVENTION

Xylanases have been found in at least a hundred different organisms. Xylanases are glycosyl hydrolases which hydrolyse β -1,4-linked xylopyranoside chains. Within the sequence-based classification of glycosyl hydrolase families established by Henrissat and Bairoch (1993), most xylanases are found in families 10 and 11. Common features for family 11 members include high genetic homology, a size of about 20 kDa and a double displacement catalytic mechanism (Tenkanen *et al.*, 1992; Wakarchuk *et al.*, 1994). The families have now been grouped, based on structure similarities, into Clans (Henrissat and Davies, 1995). Family 11 glycosyl hydrolases, which are primarily xylanases, reside in Clan C along with family 12 enzymes, all of which are known to be cellulases.

20

Xylanases can be often used for important applications such as the bleaching of pulp, modification of textile fibers and in animal feed (e.g., xylanases can aid animal digestion, Prade, 1996). Xylanases are useful for production of human foods as well. For example, xylanase improves the properties of bread dough and the quality of bread. Xylanases can also aid the brewing process by improving filterability of xylan containing beers. Xylanases can be employed in the decomposition of vegetative matter including disposal/use of agricultural waste and waste resulting from processing of agricultural products, including production of fuels or other biobased products/materials from biomass.

25

Often, however, extreme conditions in these applications, such as high temperature and/or pH, etc, render the xylanases less effective than under normal conditions. During pulp bleaching, for example, material that comes from an alkaline wash stage can have a high temperature, sometimes greater than 80 °C, and a high pH, such as a pH greater than 10. Since most xylanases do not function well under those conditions, pulp must be cooled and the alkaline pH neutralized before the normal xylanase can function. Taking some of

30

these steps into account, the process can become more expensive since it must be altered to suit the xylanase.

In another example, xylanases are also useful in animal feed applications. There, the enzymes can face high temperature conditions for a short time (e.g. – 0.5 - 5 min at 95 °C or higher) during feed preparation. Inactivation of the enzyme can occur under these temperature conditions, and the enzymes are rendered useless when needed at a lower temperature such as, for example, ~37 °C.

Xylanases with improved qualities have been found. Several thermostable, alkalophilic and acidophilic xylanases have been found and cloned from thermophilic organisms (Bodie *et al.*, 1995; Fukunaga *et al.*, 1998). However, it is often difficult to produce the enzymes in economically efficient quantities. *T. reesei*, on the other hand, produces xylanases, which are not as thermostable as xylanases from thermophilic organisms. *T. reesei* is known to produce different xylanases of which xylanases I and II (XynI and XynII, respectively) are the best characterized (Tenkanen *et al.*, 1992). XynI has a size of 19 kDa, a pI of 5.5 and a pH of between 3 and 4. XynII has a size of 20 kDa, a pI of 9.0 and a pH optimum of 5.0-5.5 (Törrönen and Rouvinen, 1995). These xylanases exhibit a favorable pH profile, specificity and specific activity in a number of applications, and can be produced economically in large-scale production processes.

Efforts have been made to engineer a xylanase with favorable qualities. For example, some have tried to improve the stability of the *Bacillus circulans* xylanase by adding disulphide bridges which bind the N-terminus of the protein to the C-terminus and the N-terminal part of the α -helix to the neighbouring β -strand (Wakarchuk *et al.*, 1994). Also, Campbell *et al.* (1995) modified *Bacillus circulans* xylanase by inter- and intramolecular disulphide bonds in order to increase thermostability. Similarly, the stability of *T. reesei* xylanase II has been improved by changing the N-terminal region to a respective part of a thermophilic xylanase (Sung *et al.*, 1998). In addition to the improved thermostability, the activity range of the enzyme was broadened to include an alkaline pH. Single point mutations have also been used to increase the stability of *Bacillus pumilus* xylanase (Arase *et al.*, 1993).

By comparing the structures of thermophilic and mesophilic enzymes much information has been obtained (Vogt *et al.*, 1997). Structural analysis of thermophilic

xylanases has also given information about factors influencing the thermostability of xylanases (Gruber *et al.*, 1998; Harris *et al.*, 1997).

Currently, however, there is a need for enzymes, especially xylanases, with improved properties in industrial conditions.

5

SUMMARY OF THE INVENTION

The current invention relates to modified enzymes. Specifically, the invention relates to modified enzymes with improved performance at extreme conditions of pH and temperature.

10 In a first aspect, the invention is drawn to a modified xylanase comprising a polypeptide having an amino acid sequence as set forth in SEQ ID NO:1, wherein the sequence has at least one substituted amino acid residue at a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 15 165, 169, 180, 184, 186, 188, 190 and +191. Preferably, the substitution is selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. Preferably, the modified xylanase has at least one substitution selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Also, preferably, the modified xylanase exhibits improved thermophilicity, 20 alkalophilicity or a combination thereof, in comparison to a wild-type xylanase.

In a second aspect, the invention is drawn to a modified enzyme, the modified enzyme comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted amino acid residue at a position equivalent to a position selected from the group 25 consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 30 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C.

In a preferred embodiment of the invention, the modified enzyme is a glycosyl hydrolase of Clan C comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Preferred modified enzymes are as disclosed herein.

In a preferred embodiment, the modified enzyme is a family 11 xylanase comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Preferred modified family 11 enzymes are as disclosed herein.

In another preferred embodiment, the modified enzyme is a family 12 cellulase comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180,

184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C, wherein the position is an equivalent position, as defined herein. Preferred family 12 modified enzymes are as disclosed herein.

In a preferred embodiment, the family 12 cellulase is *Trichoderma* EGIII cellulase as set forth in SEQ ID NO:3, the modification comprises at least one amino acid selected from the group consisting of: 2, 13, 28, 34, 77, 80, 86, 122, 123, 134, 137, 140, 164, 174, 183, 209, 215 and 218, the position numbering being with respect to SEQ ID NO:3. In a preferred embodiment, the substitution is at least one mutation selected from the group consisting of T2C, N13H, S28K, T34C, S77C, P80R, S86C, G122C, K123W, Q134H, Q134K, Q134R, V137H, G140C, N164C, N164K, N174C, K183H, N209C, A215D and N218C, position numbering being with respect to SEQ ID NO:3.

Embodiments of the first and second aspects of the invention, as disclosed above, also provide for nucleic acids encoding any of the modified enzymes, as set forth above, as well as complements. In another preferred embodiment, the invention provides for compositions comprising at least one modified enzyme, as disclosed herein, and another ingredient. In another preferred embodiment, the invention provides vectors comprising a modified enzyme, as disclosed herein, cells comprising the modified enzyme and methods of expressing the modified enzyme.

In a third aspect, the invention is drawn to a method of modifying an enzyme comprising modifying a first site in the enzyme so that the first site can bind to a second site in the enzyme. In a preferred embodiment, the first site is in a loop or sequence adjacent to a β -sheet. In a preferred embodiment, the second site is located in a β -sheet.

In a preferred embodiment, the modified enzyme is a xylanase. For example, in a preferred embodiment, the invention is drawn to a modified xylanase, wherein the xylanase is modified by at least one of the following methods: (i) by modifying an N-terminal sequence so that the N-terminal sequence is bound by a disulphide bridge to an adjacent β -strand; (ii) by modifying a C-terminal sequence so that the C-terminal sequence is bound to an adjacent β -strand; (iii) by modifying an α -helix or sequence adjacent to an α -helix, so

that the α -helix, or sequence adjacent to the α -helix, is bound more tightly to the body of the protein; (iv) by modifying a sequence adjacent to the β -strand so that the sequence adjacent to the β -strand can be bound more tightly to an adjacent sequence. For example, in a preferred embodiment, modification can occur in a β -strand next to the cord.

5

BRIEF DESCRIPTION OF FIGURES

Figure 1 shows an amino acid alignment among family 11 xylanases. The amino acid numbering is compared with *T.Reesei* Xylanase II, as indicated at the top of the sequences. The residues common to at least 75% of family 11 xylanases are underlined.

10 The following are aligned (by abbreviation) in the figure: XYN2_TRIRE Endo-1,4-beta-xylanase 2 precursor (EC 3.2.1.8) (Xylanase 2) (1,4-beta-D-xylan xylanohydrolase 2) - *Trichoderma reesei* (*Hypocrea jecorina*) >sp|P36217|; XYN1_TRIRE Endo-1,4-beta-xylanase 1 precursor (EC 3.2.1.8) (Xylanase 1) (1,4-beta-D-xylan xylanohydrolase 1) - *Trichoderma reesei* (*Hypocrea jecorina*) >sp|P36218|; XYN2_BACST Endo-1,4-beta-

15 xylanase precursor (EC 3.2.1.8) (Xylanase) (1,4-beta-D-xylan xylanohydrolase) - *Bacillus stearothermophilus* >sp|P45703|; XYN1_HUMIN Endo-1,4-beta-xylanase 1 precursor (EC 3.2.1.8) (Xylanase 1) (1,4-beta-D-xylan xylanohydrolase 1) - *Humicola insolens* >sp|P55334|; XYN1_ASPAW Endo-1,4-beta-xylanase I precursor (EC 3.2.1.8) (Xylanase I) (1,4-beta-D-xylan xylanohydrolase I) - *Aspergillus awamori* >sp|P55328|; XYNA_BACST

20 Endo-1,4-beta-xylanase A precursor (EC 3.2.1.8) (Xylanase A) (1,4-beta-D-xylan >sp|P45705|.

Figure 2 shows an amino acid alignment of family 12 Cellulases with XynII. The following are aligned (by abbreviation) in the figure: 1ENX *XylanaseII Trichoderma reesei*, and cell12 family members Q8NJY2 *Aspergillus awamori*, Q8NJY3 *Humicola*

25 *grisea*, Q8NJY4 *Trichoderma viride*, Q8NJY5 *Hypocrea koningii*, Q8NJY6 *Hypocrea schweinitzii*, Q8NJY7 *Stachybotrys echinata*, Q8NJY8 *Bionectria ochroleuca*, Q8NJY9 *Bionectria ochroleuca*, Q8NJZ0 *Bionectria ochroleuca*, Q8NJZ1 *Bionectria ochroleuca*, Q8NJZ2 *Fusarium solani* (subsp. *Cucurbitae*), Q8NJZ3 *Fusarium solani* (subsp. *cucurbitae*), Q8NJZ4 *Fusarium equiseti* (*Fusarium scirpi*), Q8NJZ5 *Emericella*

30 *desertorum*, Q8NJZ6 *Chaetomium brasiliense*, Q9KIH1 *Streptomyces sp. 11AG8*. In the Figure, the two arrows indicates the position of the disulphide bridges (signal sequence not removed).

Figure 3 shows the nucleotide sequence of the *Trichoderma reesei* oligonucleotides used in mutagenesis of the xylanase, with the codon changes underlined.

Figure 4 shows a graph comparing activity with respect to temperature of the wild-type XynII with the Y2 and Y5 mutated xylanases. Mutated xylanases have the following mutations: K58R and an aspartic acid added to the C-terminal serine at position 190 (+191D) (=Y2); T2C, T28C, K58R +191D, (=Y5). The figure exemplifies that a salt bridge, alone, does not increase thermophilicity and thermal stability. Rather, introduction of a disulphide bridge increases stability and temperature dependent activity. Activity is measured as per *Bailey et al.*, 1992.

Figure 5 shows a graph comparing the activity with respect to pH of the XynII wild-type with the Y5 mutated xylanase with the following mutations: T2C, T28C, K58R with an added aspartic acid added to the C-terminal serine position 190 (+191D). Activity is measured as per *Bailey et al.*, 1992

Figure 6 shows a graph comparing the activity with respect to temperature of the XynII wild-type with the Y5 mutated xylanase with the following mutations: T2C, T28C, K58R with an added aspartic acid added to the C-terminal serine position 190 (+191D). Activity is measured as per *Bailey et al.*, 1992.

Figure 7 shows a graph comparing the residual activity at pH 5.0, with inactivation at pH 8 with respect to temperature of the wild type XynII xylanase with the Y5 mutated xylanase having the following mutations: T2C, T28C, K58R with an added aspartic acid added to the C-terminal serine position 190 (+191D). Activity is measured as per *Bailey et al.*, 1992.

Figure 8 shows a graph comparing the residual activity at pH 5.3, with inactivation at pH 8 with respect to temperature of the Y5 mutated xylanase with a XynII xylanase (SS105/162) having the following additional mutations Q162C and L105C. Activity is measured as per *Bailey et al.*, 1992.

Figure 9 shows a graph comparing the residual activity at pH 5, with inactivation at pH 9 with respect to temperature of the Y5 mutated xylanase with a XynII xylanase (P9) having the following additional mutations: F93W, N97R and H144K. Activity is measured as per *Bailey et al.*, 1992.

Figure 10 shows a graph comparing the residual activity at pH 5, with inactivation at pH 5 with respect to temperature of the Y5 mutated xylanase with a XynII xylanase

(P12) having the following additional mutations H144C and N92C. Activity is measured as per *Bailey et al.*, 1992.

Figure 11 shows a graph comparing the residual activity at pH 5, with inactivation at pH 9 with respect to temperature of the Y5 mutated xylanase with a XynII xylanase
5 (P12) having the following additional mutations H144C and N92C. Activity is measured as per *Bailey et al.*, 1992.

Figure 12 shows a graph comparing the residual activity at pH 5.2, with inactivation at pH 8 with respect to temperature of the Y5 mutated xylanase with a XynII (P15) xylanase having the following additional mutations: F180Q, H144C and N92C. Activity is
10 measured as per *Bailey et al.*, 1992.

Figure 13 shows a graph comparing the residual activity at pH 5, with inactivation at pH 8 with respect to temperature of the Y5 mutated xylanase with a XynII xylanase (P21) having the following additional mutations: H22K, F180Q, H144C and N92C. Activity is measured as per *Bailey et al.*, 1992.

15 Figure 14 shows a graph comparing the residual activity at pH 5.17 with inactivation at pH 7.8, with respect to temperature of the Y5 mutated xylanase with a XynII xylanase (P20) having the following additional mutations: H22K and F180Q. Activity is measured as per *Bailey et al.*, 1992.

Figure 15 shows a graph comparing the activity at pH 8 with respect to temperature
20 of the Y5 mutated xylanase with a XynII xylanase (J17) having the following additional mutation: V108H. Activity is measured as per *Bailey et al.*, 1992.

Figure 16 shows a graph comparing the activity at pH 8 with respect to temperature of the Y5 mutated xylanase with a XynII xylanase (J21) having the following additional mutations: S65C and S186C (J21 in the graph). Activity is measured as per *Bailey et al.*,
25 1992.

Figure 17 shows a structural alignment of *Trichoderma reesei* xylanaseII (XynII, PDB 1 ENX, in blue;) and *Trichoderma reesei* endoglucanaseIII (Cal12A, PDB 1H8V, in red).

Figure 18 sets forth the nucleotide amino acid of sequence of XynII.

30 Figure 19 sets forth the nucleotide amino acid of sequence of EGIII.

Figure 20 sets forth the nucleotide amino acid of sequence of Xyn1I.

Detailed Description of the Preferred Embodiments

The invention will now be described in detail by way of reference only using the following definitions and examples. Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of
5 ordinary skill in the art to which this invention belongs. Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 2D ED., John Wiley and Sons, New York (1994), and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY, Harper Perennial, NY (1991) provide one of skill with a general dictionary of many of the terms used in this invention. Although any methods and materials similar or equivalent to those
10 described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. Practitioners are particularly directed to Sambrook *et al.*, 1989, and Ausubel
15 FM *et al.*, 1993, for definitions and terms of the art. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary.

The headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the specification as a
20 whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole.

As used herein, the term "polypeptide" refers to a compound made up of a single chain of amino acid residues linked by peptide bonds. The term "protein" herein may be synonymous with the term "polypeptide" or may refer, in addition, to a complex of two or more polypeptides.

25 As used herein, the term "expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of the gene. The process includes both transcription and translation.

As used herein, the term “gene” means the segment of DNA involved in producing a polypeptide chain, that may or may not include regions preceding or following the coding region.

As used herein, when referring to position numbering, the term “equivalent” refers to positions as determined by sequence and structural alignments with *Trichoderma reesei* xylanase II (xynII) as a reference sequence or reference structure, as provided herein (see, for example, Figure 2 for a multiple sequence alignment and *Trichoderma reesei* xylanaseII with other sequences, and Figure 17 for a structural alignment of *Trichoderma reesei* Xyn II with *Trichoderma reesei* endoglucanaseIII). Position numbering shall be with respect to *Trichoderma reesei* xynII, as set forth in SEQ ID NO:1. The numbering system, even though it may use a specific sequence as a base reference point, is also applicable to all relevant homologous sequences. Sequence homology between proteins may be ascertained using well-known alignment programs and as described herein and by using hybridisation techniques described herein.

As used herein, the term “adjacent” refers to close linear and/or close spatial proximity between amino acid residues or regions or areas of a protein. For example, a first residue or first region or first area which is adjacent to a second residue or second region or second area (in a linear sense), respectively, shall have preferably about 7, preferably about 5, preferably about 2 intervening amino acid residues between them. Alternatively, for example, when a first set of residues or a first region or first area is adjacent to a second set of residues or a second region or second area, then the first set of residues or first region or first area shall be proximal (in space, as shown, for example, by the tertiary structure of a protein) to the second set of residues or second region or second area. One skilled in the art, when possible, would know how to solve the tertiary structure of a protein.

As used herein, when referring to sequence positions, the designation “+” followed by an integer shall mean that a polypeptide has been modified to include additional amino acid(s) at the putative position, as specified by the integer. For example, the designation +191 shall mean that a polypeptide which normally has 190 amino acids in the amino acid sequence has an added amino acid.

As used herein, the term “nucleic acid molecule” includes RNA, DNA and cDNA molecules. It will be understood that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding a given protein, such as the mutant proteins of the invention, may be produced.

As used herein, the term “disulphide bridge” or “disulphide bond” refers to the bond formed between the sulphur atoms of cysteine residues in a polypeptide or a protein. In this invention, a disulphide bridge or disulphide bond may be non-naturally occurring and introduced by way of point mutation.

5 As used herein, the term “salt bridge” refers to the bond formed between oppositely charged residues, amino acids in a polypeptide or protein. In this invention, a salt bridge may be non-naturally occurring and introduced by way of point mutation.

As used herein, an “enzyme” refers to a protein or polypeptide that catalyzes a chemical reaction.

10 As used herein, the term “activity” refers to a biological activity associated with a particular protein, such as enzymatic activity associated with a protease. Biological activity refers to any activity that would normally be attributed to that protein by one skilled in the art.

As used herein, the term “xylanase” refers to glycosyl hydrolases that hydrolyse β -1,4-linked xylopyranoside chains.

As used herein, “XynI” refers to the *Trichoderma reesei* xylanase, xylanase I. XynI has a size of 19 kDa, a pI of 5.5 and a pH optimum of between 3 and 4.

As used herein, “XynII” refers to the *Trichoderma reesei* xylanase, xylanase II. XynII has a size of 20 kDa, a pI of 9.0 and a pH optimum of between 5 and 5.5.

20 As used herein, “xylopyranoside” refers to a β -1,4-linked polymer of xylose, including substituted polymers of xylose, i.e. branched β -D-1,4-linked xylopyranose polymers, highly substituted with acetyl, arabinosyl and uronyl groups (see, for example, Biely, P. (1985) Microbial Xylanolytic Systems. Trends Biotechnol., 3, 286-290.).

As used herein, the term “glycosyl hydrolase” refers to an enzyme which hydrolyzes the glycosidic bond between two or more carbohydrates or between a carbohydrate and a non-carbohydrate moiety. Enzymatic hydrolysis of the glycosidic bond takes place via general acid catalysis and requires two critical residues: a proton donor and a nucleophile/base. The IUB-MB Enzyme nomenclature of glycosyl hydrolases is based on substrate specificity and occasionally on molecular mechanism.

30 As used herein, the term “hydrolase” refers to an enzyme that catalyzes a reaction whereby a chemical bond is enzymatically cleaved with the addition of a water molecule.

As used herein, “hydrolysis” refers to the process of the reaction whereby a chemical bond is cleaved with the addition of a water molecule.

As used herein, “Clan C” refers to groupings of families which share a common three-dimensional fold and identical catalytic machinery (*see*, for example, Henrissat, B. and Bairoch, A., (1996) *Biochem. J.*, 316, 695-696).

As used herein, “family 11” refers to a family of enzymes as established by Henrissat and Bairoch (1993) *Biochem J.*, 293, 781-788 (*see*, also, Henrissat and Davies (1997) *Current Opinion in Structural Biol.* 1997, &:637-644). Common features for family 11 members include high genetic homology, a size of about 20 kDa and a double displacement catalytic mechanism (*see* Tenkanen *et al.*, 1992; Wakarchuk *et al.*, 1994). The structure of the family 11 xylanases includes two large β -sheets made of β -strands and α -helices. Family 11 xylanases include the following: *Aspergillus niger* XynA, *Aspergillus kawachii* XynC, *Aspergillus tubigensis* XynA, *Bacillus circulans* XynA, *Bacillus pumilus* XynA, *Bacillus subtilis* XynA, *Neocallimastix patriciarum* XynA, *Streptomyces lividans* XynB, *Streptomyces lividans* XynC, *Streptomyces thermoviolaceus* XynII, *Thermomonospora fusca* XynA, *Trichoderma harzianum* Xyn, *Trichoderma reesei* XynI, *Trichoderma reesei* XynII, *Trichoderma viride* Xyn.

As used herein, “family 12” refers to a family of enzymes established by Henrissat and Bairoch (1993) in which known glycosyl hydrolases were classified into families based on amino acid sequence similarities. To date all family 12 enzymes are cellulases. Family 12 enzymes hydrolyze the β -1,4-glycosidic bond in cellulose via a double displacement reaction and a glucosyl-enzyme intermediate that results in retention of the anomeric configuration of the product. Structural studies of family 12 members reveal a compact β -sandwich structure that is curved to create an extensive substrate binding site on the concave face of the β -sheet.

As used herein, the term “protease” refers to an enzyme that degrades by hydrolyzing at least some of their peptide bonds.

As used herein, “peptide bond” refers to the chemical bond between the carbonyl group of one amino acid and the amino group of another amino acid.

As used herein, “wild-type” refers to a sequence or a protein that is native or naturally occurring.

As used herein, "point mutations" refers to a change in a single nucleotide of DNA, especially where that change shall result in a change in a protein.

As used herein, "mutant" refers to a version of an organism or protein where the version is other than wild-type. The change may be affected by methods well known to one skilled in the art, for example, by point mutation in which the resulting protein may be referred to as a mutant.

As used herein, "mutagenesis" refers to the process of affecting a change from a wild-type into a mutant.

As used herein, "substituted" and "modified" are used interchangeably and refer to a sequence, such as an amino acid sequence comprising a polypeptide, that includes a deletion, insertion, replacement or interruption of a naturally occurring sequence. Often in the context of the invention, a substituted sequence shall refer, for example, to the replacement of a naturally occurring residue.

As used herein, "modified enzyme" refers to an enzyme that includes a deletion, insertion, replacement or interruption of a naturally occurring sequence.

As used herein, " β -strands" refers to that portion of an amino acid sequence that forms a linear sequence that occurs in a β -sheets.

As used herein, " β -sheets" refers to the sheet-type structure that results when amino acids hydrogen-bond to each other to form a sheet like structure.

As used herein, " α -helix" refers to the structure that results when a single polypeptide chain turns regularly about itself to make a rigid cylinder in which each peptide bond is regular hydrogen-bonded to other peptide bonds in the nearby chain.

As used herein, "thumb" refers to a loop between β -strands B7 and B8 in XynI and in XynII (*see*, for example, in Torronen, A. and Rouvinen, J.; Biochemistry 1995, 34, 847-856).

As used herein, "cord" refers to a loop between β -strands B7 and B8 which make a thumb and a part of the loop between β -strands B6a and B9 which crosses the cleft on one side (*see*, for example, Torronen, A. and Rouvinen, J.; Biochemistry 1995, 34, 847-856).

As used herein, "alkaline" refers to the state or quality of being basic.

As used herein, "alkalophilic" refers to the quality of being more robust in an alkaline atmosphere than a non-alkalophilic member. For example, an alkalophilic organism refers to an organism that survives or thrives under alkaline conditions where a

normal organism may not, and an alkalophilic protein is one whose activity is active or more robust under alkaline conditions where a normal protein would be less active.

As used herein, “acidic” refers to the state or quality of being acidic.

As used herein, “acidophilic” refers to the quality of being more robust in an acidic
5 atmosphere than a non-acidophilic member. For example, an acidophilic organism refers to an organism that survives or thrives under acidic conditions where a normal organism may not, and an acidophilic protein is one whose activity is active or more robust under acidic conditions where a normal protein would be less active.

As used herein, “thermostable” refers to the quality of being stable in an atmosphere
10 involving temperature. For example, a thermostable organism is one that is more stable under specified temperature conditions than a non-thermostable organism.

As used herein, “thermostability,” refers to the quality of being thermostable.

As used herein, “thermophilic” refers to the quality of being more robust in an hot
atmosphere than a non-thermophilic member. For example, a thermophilic organism refers
15 to an organism that survives or thrives under hot conditions where a normal organism may not, and a thermophilic protein is one whose activity is active or more robust under hot conditions where a normal protein would be less active.

As used herein, “mesophilic” refers to the quality of being more robust in an normal
atmosphere than a non-mesophilic member. For example, a mesophilic organism refers to
20 an organism that survives or thrives under normal conditions where another organism may not, and a mesophilic protein is one whose activity is active or more robust under normal conditions where another protein would be less active.

As used herein, “oligonucleotides” refers to a short nucleotide sequence which may
be used, for example, as a primer in a reaction used to create mutant proteins.

As used herein, “codon” refers to a sequence of three nucleotides in a DNA or
25 mRNA molecule that represents the instruction for incorporation of a specific amino acid into a polypeptide chain.

As used herein, “Y5” refers to a mutant xylanase as disclosed, for example, in
publication number WO 01/27252.

30 As used herein, the following designations shall refer to the following mutants:

“P2” = N97R + H144K / Y5

“P3” = F93W + H144K in Y5

“P8” = F180Q in Y5
 “P9” = N97R in F93W + H144K in Y5
 “P12” = H144C + N92C in Y5
 “P15” = F180Q in H144C + N92C in Y5
 5 “P16” = N97R in H144C + N92C in Y5
 “P18” = H22K in Y5
 “P20” = H22K + F180Q in Y5
 “P21” = H22K + F180Q + H144C + N92C in Y5
 “J17” = V108H in Y5
 10 “J21” = S65C + S186C in Y5

wherein position numbering shall be with respect to XynII.

The present invention relates to modified enzymes with improved performance in extreme conditions, such as temperature and pH.

15 In a first aspect, the invention is drawn to a modified xylanase comprising a polypeptide having an amino acid sequence as set forth in SEQ ID NO:1, wherein the sequence has at least one substituted amino acid residue at a position selected from the group consisting of: 2, 5, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 20 165, 169, 180, 184, 186, 188, 190 and +191, where position numbering is with respect to SEQ ID NO:1. Preferably, the substitution is selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. Preferably, the modified xylanase has at least one substitution selected from the group consisting of H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Also, preferably, the 25 modified xylanase exhibits improved thermophilicity, alkalophilicity or a combination thereof, in comparison to a wild-type xylanase.

In a second aspect, the invention is drawn to a modified enzyme, the modified enzyme comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one 30 substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 44, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191, wherein position numbering is with respect to SEQ ID NO:1. In a preferred embodiment, the amino acid sequence has at least one 35 substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a

preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C.

In a preferred embodiment of the invention, the modified enzyme is a glycosyl
5 hydrolase of Clan C comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, , 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 110, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157,
10 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K,
15 S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Preferred modified enzymes are as disclosed herein.

In a preferred embodiment, the modified enzyme is a family 11 xylanase comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted amino acid
20 residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10 , 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38 , 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group
25 consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Preferred modified family 11 enzymes are as disclosed herein.

30 In another preferred embodiment, the modified enzyme is a family 12 cellulase comprising an amino acid sequence, the amino acid sequence being homologous to the sequence set forth in SEQ ID NO:1, the amino acid sequence having at least one substituted

amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 144, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, 190 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 144, 162, 180, 186 and +191. In a preferred embodiment, the amino acid sequence has at least one substituted amino acid residue selected from the group consisting of: H22K, S65C, N92C, F93W, N97R, V108H, H144C, H144K, F180Q and S186C. Preferred family 12 modified enzymes are as disclosed herein.

In a preferred embodiment, the family 12 cellulase is *Trichoderma* EGIII cellulase as set forth in SEQ ID NO:3, the modification comprises at least one amino acid selected from the group consisting of: 2, 13, 28, 34, 77, 80, 86, 122, 123, 134, 137, 140, 164, 174, 183, 209, 215 and 218, position numbering being with respect to SEQ ID NO:3. In a preferred embodiment, the substitution is at least one mutation selected from the group consisting of T2C, N13H, S28K, T34C, S77C, P80R, S86C, G122C, K123W, Q134H, Q134K, Q134R, V137H, G140C, N164C, N164K, N174C, K183H, N209C, A215D and N218C, position numbering being with respect to SEQ ID NO:3.

XynII exhibits a significant amino acid homology with other members of family 11, approximately 20-90%, as well as overall structural similarity. Homology, as used herein, may be determined by one skilled in the art; specifically, homologies of at least 20%, preferably 30% or more, preferably 40% or more, preferably 50% or more, preferably 60% or more, preferably 70% or more, preferably 80% or more, preferably 90% or more, preferably 95% or more and preferably 97% or more are contemplated (as calculated at the amino acid level and the nucleotide level and as used herein). There are structural similarities between family 11 and family 12 enzymes. Beta proteins have two stacked beta sheets, and one alpha helix packed against one of the beta sheets forming a so-called beta-jelly roll structure. (see Stirk, H.J., Woolfson, D.N., Hutchison, E.G. and Thornton, J.M. (1992) Depicting topology and handedness in jellyroll structures. *FEBS Letters* 308 p1-3).

Based on this structural similarity, both enzyme families have been assigned to a "super family" referred to as Clan C (see Sandgren, M. et. al., J. Mol. Bio. (2001) 308, 295-310.)).

Although the sequence homology between families 11 and 12 is low, the overall structural similarity of the two families is remarkable as seen by comparing figures 2 and 16. The length of the loops connecting the two beta-sheets comprises the major structural differences between the families (Sandgren et. al., J. Mol., Biol., 2001). Presently, no
 5 family 11 enzymes are known to contain N terminal disulphide bridges while many family 12 cellulases, in general appear to contain a disulphide bridge near the N-terminus (e.g, between residues 4 and 32 in *T. reesei* Cel 12A). That disulphide bridge in family 12 enzymes is located near the position where a disulphide was introduced into the *Trichoderma* (Y5) variant, although further away from the N-terminus (see, for example,
 10 publication WO 01/27252). The importance of a restriction stabilizing the N-terminal region of family 11 enzymes was examined in *Trichoderma reesei* xylanase II (XynII). By inserting a non-natural disulphide bridge between residues (T2C and T28C), an increase in T_m of 11 °C was achieved. In these two structurally similar families, family 11 and family 12, the N-terminal disulphide bridges play a similar roles regarding stability. This has
 15 been demonstrated by replacing the cysteine at position 32 with an alanine in Cel12A resulting in a significant decrease in T_m of 18.5 °C. Interestingly, the magnitude of the change in stability for adding a non-natural N-terminal disulphide into XynII is comparable to that of removing a natural one from Cel 12A (see table A).

Table A

20

Enzyme	Delta T_m	T_m (degrees C)
WT Cel12A		54.4
C32A	-18.5	35.9
WT xynII		58.6
Y5	+10.7	69.3

Table A shows the melting temperatures, T_m of the wild type Cel12A compared to the variant with the substitution at position 32, and the wild type XynII compared to the Y5 variant of this enzyme.

25

The three dimensional structures of the N-terminal disulphide bridges of the three publicly known structures for family 12 glycosyl hydrolases (*Trichoderma reesei*- PDB 1H8V, *Aspergillus niger*- PDB 1KS5, *Streptomyces lividans*- PDB 2NLR), show a shift in

the position of the disulphide bridge as compared to the non-natural disulphide bridge at sites 2 and 28 in Y5 xylanase. Table B shows the position of the disulphide bridge in a Y5 xylanase ("PDB 1ENX" being wild type XynII xylanase) and in the three known family 12 structures. The structural positions of the mutations at 2 and 28 of Y5 xylanase can be translated to the corresponding residues in the Cel 12 structures. In each case, the non-native disulphide from Y5 is closer to the N-terminus; and for the *A. niger* structure (PDB 1KS5) a disulphide could be designed that would utilize the N-terminal residue itself (at residues Q1C, V35C, according to *A. niger* numbering). Instead of being limited by the natural sequence, X-ray data could be used to design extensions and truncations of the N-terminus to facilitate non-native disulphides that specifically attach to the new N-terminal residues.

Table B

Code	WT N-terminal S-S position	Corresponding site to 2-28 of xynII	Where (according to structure) could a S-S be inserted at the N-terminal
PDB 1ENX	No	-	
Y5	C2-C28	T2-T28	T2C-T28C
PDB 1H8V	C4-C32	T2-T34	T2C-T34C
PDB 1KS5	C4-C32	T2-Y34	Q1C-V35C
PDB 2NLR	C5-C31	T3-T33	T3C-T33C

A large number of family 12 sequences (Table C) are known which could potentially be stabilized through an N-terminal disulphide bridge, particularly those molecules where a non-native disulphide bridge could be introduced or a native disulphide could be moved closer to an N-terminus. Table C lists a number of sequences where a predicted removal of the signal sequence produces mature protein sequences very similar to the ones of the known family 12 structures. Table C also lists the distance between the two N-terminal cysteines (26-28 amino acids) similar to the disulphide bond of Y5. In the cleavage site predictions, a signal sequences is theoretically removed by the means of known, acknowledged parameters (*see, for example, "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites". Henrik Nielsen, Jacob Engelbrecht, Søren Brunak and Gunnar von Heijne, Protein Engineering 10, 1-6 (1997).*

A large group of sequences of unknown three dimensional structures in Table C fall within the structurally similar group of family 12 enzymes, which have in a similar manner a cysteine residue at the N-terminal at site 5 +/- 2 residues, forming a disulphide bridge

with residue 32 +/- 7, such that the first beta strand or strands of the beta sheet can be bound to the adjacent beta sheet. All of these sequences could be treated in the manner described in the discussion around table B to improve stability.

5

Table C

ID	Sequence	Eucaryote/ Gram-/ Gram+	Predicted cleavage site	Number of adequate cysteine (1 st in ss bond)	aa's to 2 nd cysteine in ss bond
Q8NJY2	Endoglucanase {GENE:CEL12B} <i>Aspergillus awamori</i> (var. kawachi)	Eu	16-17	6	28
Q8NJY4	Endoglucanase {GENE:CEL12A} - <i>Trichoderma viride</i>	Eu	16-17	4	28
Q8NJY5	Endoglucanase {GENE:CEL12A} - <i>Hypocrea koningii</i>	Eu	16-17	4	28
Q8NJY6	Endoglucanase {GENE:CEL12A} - <i>Hypocrea schweinitzii</i>	Eu	16-17	4	28
Q8NJY7	Endoglucanase {GENE:CEL12A} - <i>Stachybotrys echinata</i>	Eu	16-17	4	28
Q8NJY8	Endoglucanase {GENE:CEL12D} - <i>Bionectria ochroleuca</i>	Eu	17-18	4	28
Q8NJY9	Endoglucanase {GENE:CEL12C} - <i>Bionectria ochroleuca</i>	Eu	17-18	3	28
Q8NJZ1	Endoglucanase {GENE:CEL12A} - <i>Bionectria ochroleuca</i>	Eu	18-19	4	28
Q8NJZ4	Endoglucanase {GENE:CEL12A} - <i>Fusarium equiseti</i> (<i>Fusarium scirpi</i>)	Eu	17-18	4	28
Q9KIH1	Cellulase 12A {GENE:CEL12A} - <i>Streptomyces</i> sp. 11AG8	Gram+	31-32	5	26

10

Table D lists further a number of sequences of family 12 enzymes with uncleaved signal sequence. They all have cysteines 30-39 amino acids apart, and after a removal of the signal sequence (removal can be performed as in table C) are structurally capable of forming a disulphide bridge at the N-terminal (as seen in the publicly known structures, see table B). The proposed mutation site correlates to the corresponding site of the disulphide

bridge between sites 2-28 of the Y5 mutant. The glycosyl hydrolase sequences were aligned using the program MOE (Chemical Computing Corp) using standard sequence matching methods.

5	Table D			
	Sequence code	enzyme	Species	Mutations
	Tr O94218	Cel12	<i>Aspergillus aculeatus</i>	D22C/ G52C
10	Sp P22669	Cel12	<i>Aspergillus aculeatus</i>	Q20C/ T52C
	Sp Q12679	Cel12	<i>Aspergillus awamori</i>	T18C/ Y50C
	Tr O13454	Cel12	<i>Aspergillus oryzae</i>	E18C/ Y50C
	Sp P16630	Cel12	<i>Erwina carotovora</i>	A32C/ I68C
	Tr O31030	Cel12	<i>Pectobacterium carotovora</i>	A32C/ V68C
15	Tr Q9V2TO	Cel12	<i>Pyrococcus furiosus</i>	P57C/ T96C
	Tr O33897	Cel12	<i>Rhodothermus marinus</i>	E40C / E70C
	Tr Q9RJY3	Cel12	<i>Streptomyces coelicolor</i>	T43C/ T73C
	Tr O08468	Cel12	<i>Streptomyces halstedii</i>	L40C/ T70C
	Tr Q59963	Cel12	<i>Streptomyces rochei</i>	T40C / T70C
20	Tr Q9KIH1	Cel12	<i>Streptomyces</i> sp. 11AG8	Q34C / N64C
	Tr Q60032	Cel12	<i>Thermotoga maritima</i>	V2C / K38C
	Tr Q60033	Cel12	<i>Thermotoga maritime</i>	V20C/ K56C
	Tr O08428	Cel12	<i>Thermotoga neopolitana</i>	V2C/ R38C
	Tr P96492	Cel12	<i>Thermotoga neopolitana</i>	V20C / K56C
25	AF435072	Cel12A	<i>Aspergillus Kawachi</i>	Q20C / T52C
	AF434180	Cel12A	<i>Chaetium brasiliense</i>	S28C / Y61C
	AF434181	Cel12A	<i>Emericella desertorum</i>	D30C / G63C
	AF434182	Cel12A	<i>Fusarium equiseti</i>	D19C / H51C
	AF434183	Cel12A	<i>Nectria ipomoeae</i>	Q25C / T58C
30	AF434184	Cel12B	<i>Nectria ipomoeae</i>	T32C / T65C
	AF435063	Cel12A	<i>Bionectria ochroleuca</i>	T20C / Y52C
	AF435064	Cel12B	<i>Bionectria ochroleuca</i>	T34C / T66C
	AF435065	Cel12C	<i>Bionectria ochroleuca</i>	A18C / T50C
	AF435066	Cel12D	<i>Bionectria ochroleuca</i>	S19C / Y51C
35	AF435071	Cel12A	<i>Humicola grisea</i>	S34C / Y67C
	AF435068	Cel12A	<i>Hypochrea schweinitzii</i>	T18C / T50C
	AF435067	Cel12A	<i>Stachybotrys echinata</i>	S18C / Y50C

Not only does the N-terminal region show high structural similarity between families 11 and 12; both families show a hand like structure, the one of a “partly closed right hand” as described in Törrönen et al. 1997. The two β -sheets form “fingers”, and a twisted pair from one β -sheet and the α -helix forms a “palm”. The long loop between β -strands B7 and B8 makes the “thumb” and a part of the loop between β -strands B6b (residues 95-102 in xynII and 125-131 in Cel12A) and B9 forms a “cord”, which crosses the cleft on one side (Törrönen A. and Rouvinen, J. Biochem. 1995, 34, 847-0856). The stabilizing effect of

inserting rigidifying substitutions between beta strand B6b and the adjacent loop and/or the “cord” is seen in the mutation at sites 92, 93, 144 (N92C-H144C, at least one of the following mutations N97R, F93W + H144K (XynII), and can in a similar way be introduced into corresponding sites in family 12.

5

Table E shows the numbering of a selection of structurally equivalent sites between xynII and Cel 12A. The high structural similarity between the two families enables a large number of similar substitutions (see Sandgren et. al., J. Mol., Biol., 2001 for structural comparison).

10

Table E

Examples of equivalent sites	
<u>XynII</u>	<u>Cel12A</u>
T2C	T2C
T28C	T34C
N92C	G122C
H144C, K	N164C, K
F93W	K123W
Q162H	K183H

The modified enzymes of the invention may comprise one or more mutations in addition to those set out above. Other mutations, such as deletions, insertions, substitutions, transversions, transitions and inversions, at one or more other locations, may also be included. Likewise, the modified enzyme may be missing at least one of the substitutions set forth above.

The modified enzyme may also comprise a conservative substitution that may occur as a like-for-like substitution (e.g., basic for basic, acidic for acidic, polar for polar etc.) Non-conservative substitutions may also occur, i.e. from one class of residue to another or alternatively involving the inclusion of unnatural amino acids such as ornithine, diaminobutyric acid ornithine, norleucine ornithine, pyriylalanine, thienylalanine, naphthylalanine and phenylglycine.

The sequences may also have deletions, insertions or substitutions of amino acid residues that produce a silent change and result in a functionally equivalent substance. Deliberate amino acid substitutions may be made on the basis of similarity in amino acid properties (such as polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues) and it is therefore useful to group amino acids together in functional groups. Amino acids can be grouped together based on the properties of their side chain alone. However it is more useful to include mutation data as well. The sets of amino acids thus derived are likely to be conserved for structural reasons. These sets can be described in the form of a Venn diagram (Livingstone C.D. and Barton G.J. (1993) "Protein sequence alignments: a strategy for the hierarchical analysis of residue conservation" *Comput. Appl Biosci.* 9: 745-756)(Taylor W.R. (1986) "The classification of amino acid conservation" *J.Theor.Biol.* 119; 205-218). Conservative substitutions may be made, for example according to the table below which describes a generally accepted Venn diagram grouping of amino acids.

15

Set		Sub-set	
Hydrophobic	F W Y H K M I L V A G C	Aromatic	F W Y H
		Aliphatic	I L V
Polar	W Y H K R E D C S T N Q	Charged	H K R E D
		Positively charged	H K R
		Negatively charged	E D
Small	V C A G S P T N D	Tiny	A G S

Variant amino acid sequences may also include suitable spacer groups inserted between any two amino acid residues of the sequence including alkyl groups such as methyl, ethyl or propyl groups in addition to amino acid spacers such as glycine or β -

alanine residues. A further form of variation involves the presence of one or more amino acid residues in peptoid form.

Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences. % homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence is directly compared with the corresponding amino acid in the other sequence one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues.

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (Devereux et al

1984 Nuc. Acids Research 12 p387). Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel et al., 1999 Short Protocols in Molecular Biology, 4th Ed – Chapter 18), FASTA (Altschul et al., 1990 J. Mol. Biol. 403-410) and the GENEWORKS suite of comparison tools. Both
5 BLAST and FASTA are available for offline and online searching (see Ausubel et al., 1999, Short Protocols in Molecular Biology, pages 7-58 to 7-60). However, for some applications, it is preferred to use the GCG Bestfit program. BLAST 2 Sequences is also available for comparing protein and nucleotide sequence (see FEMS Microbiol Lett 1999 174(2): 247-50; FEMS Microbiol Lett 1999 177(1): 187-8 and tatiana@ncbi.nlm.nih.gov).

10 Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of
15 programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). For some applications, it is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

Alternatively, percentage homologies may be calculated using the multiple
20 alignment feature in DNASISTM (Hitachi Software), based on an algorithm, analogous to CLUSTAL (Higgins DG & Sharp PM (1988), Gene 73(1), 237-244).

Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

25 Embodiments of the first and second aspects of the invention, as disclosed above, provide a nucleic acid encoding any of the modified enzymes, as set forth above, as well as complements thereof. In another preferred embodiment, the invention provides for compositions comprising at least one modified enzyme, as disclosed herein, and another ingredient. In another preferred embodiment, the invention provides vectors comprising a
30 modified enzyme, as disclosed herein, cells comprising the modified enzyme and methods of expressing the modified enzyme.

One skilled in the art will be aware of the relationship between nucleic acid sequence and polypeptide sequence, in particular, the genetic code and the degeneracy of this code, and will be able to construct such modified enzymes without difficulty. For example, one skilled in the art will be aware that for each amino acid substitution in the modified enzyme sequence there may be one or more codons which encode the substitute amino acid. Accordingly, it will be evident that, depending on the degeneracy of the genetic code with respect to that particular amino acid residue, one or more modified enzyme nucleic acid sequences may be generated corresponding to that modified enzyme polypeptide sequence.

Mutations in amino acid sequence and nucleic acid sequence may be made by any of a number of techniques, as known in the art. In particularly preferred embodiments, the mutations are introduced into parent sequences by means of PCR (polymerase chain reaction) using appropriate primers, as illustrated in the Examples. The parent enzymes may be modified at the amino acid level or the nucleic acid level to generate the modified enzyme sequences described herein. Therefore, a preferred embodiment provides for the generation of modified enzymes by introducing one or more corresponding codon changes in the nucleotide sequence encoding a modified enzyme.

It will be appreciated that the above codon changes can be made in any modified enzyme nucleic acid sequence. For example, sequence changes can be made to any of the homologous sequences described herein.

The modified enzyme may comprise the "complete" enzyme, i.e., in its entire length as it occurs in nature (or as mutated), or it may comprise a truncated form thereof. The modified enzyme derived from such may accordingly be so truncated, or be "full-length". The truncation may be at the N-terminal end or the C-terminal end. The modified enzyme may lack one or more portions, such as sub-sequences, signal sequences, domains or moieties, whether active or not.

A nucleotide sequence encoding either an enzyme which has the specific properties as defined herein or an enzyme which is suitable for modification, such as a modified enzyme, may be identified and/or isolated and/or purified from any cell or organism producing said enzyme. Various methods are well known within the art for the identification and/or isolation and/or purification of nucleotide sequences. By way of

example, PCR amplification techniques to prepare more of a sequence may be used once a suitable sequence has been identified and/or isolated and/or purified.

By way of further example, a genomic DNA and/or cDNA library may be constructed using chromosomal DNA or messenger RNA from the organism producing the enzyme. If the amino acid sequence of the enzyme or a part of the amino acid sequence of the enzyme is known, labelled oligonucleotide probes may be synthesised and used to identify enzyme-encoding clones from the genomic library prepared from the organism. Alternatively, a labelled oligonucleotide probe containing sequences homologous to another known enzyme gene could be used to identify enzyme-encoding clones. In the latter case, hybridisation and washing conditions of lower stringency are used.

Alternatively, enzyme-encoding clones could be identified by inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming enzyme-negative bacteria with the resulting genomic DNA library and then plating the transformed bacteria onto agar plates containing a substrate for enzyme thereby allowing clones expressing the enzyme to be identified.

In a yet further alternative, the nucleotide sequence encoding the modified enzyme may be prepared synthetically by established standard methods, e.g. the phosphoroamidite method described by Beaucage S.L. *et al.*, (1981) *Tetrahedron Letters* 22, p 1859-1869 or the method described by Matthes *et al.*, (1984) *EMBO J.* 3, p 801-805. In the phosphoroamidite method, oligonucleotides are synthesised, e.g. in an automatic DNA synthesiser, purified, annealed, ligated and cloned in appropriate vectors.

The nucleotide sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin in accordance with standard techniques. Each ligated fragment corresponds to various parts of the entire nucleotide sequence. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202 or in Saiki R K *et al.*, (*Science* (1988) 239, pp 487-491).

The nucleotide sequences described here, and suitable for use in the methods and compositions described here may include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones and/or the addition of acridine

or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of this document, it is to be understood that the nucleotide sequences described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the *in vivo* activity or life span of nucleotide sequences.

5 A preferred embodiment of the invention provides for nucleotide sequences and the use of nucleotide sequences that are complementary to the sequences presented herein, or any derivative, fragment or derivative thereof. If the sequence is complementary to a fragment thereof then that sequence can be used as a probe to identify similar coding sequences in other organisms etc.

10 Polynucleotides which are not 100% homologous to the modified enzyme sequences may be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing DNA libraries made from a range of individuals, for example individuals from different populations. In addition, other homologues may be obtained and such homologues and fragments thereof in general will
15 be capable of selectively hybridising to the sequences shown in the sequence listing herein. Such sequences may be obtained by probing cDNA libraries made from or genomic DNA libraries from other species and probing such libraries with probes comprising all or part of any one of the sequences in the attached sequence listings under conditions of medium to high stringency. Similar considerations apply to obtaining species homologues and allelic
20 variants of the polypeptide or nucleotide sequences described here.

Variants and strain/species homologues may also be obtained using degenerate PCR which will use primers designed to target sequences within the variants and homologues encoding conserved amino acid sequences. The primers used in degenerate PCR will contain one or more degenerate positions and will be used at stringency conditions lower
25 than those used for cloning sequences with single sequence primers against known sequences. Conserved sequences can be predicted, for example, by aligning the amino acid sequences from several variants/homologues. Sequence alignments can be performed using computer software known in the art as described herein.

Alternatively, such polynucleotides may be obtained by site directed mutagenesis of
30 characterised sequences, as provided herein. This may be useful where, for example, silent codon sequence changes are required to optimise codon preferences for a particular host cell in which the polynucleotide sequences are being expressed. Other sequence changes

may be desired in order to introduce restriction enzyme recognition sites, or to alter the property or function of the polypeptides encoded by the polynucleotides.

The polynucleotides may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a revealing label by
5 conventional means using radioactive or non-radioactive labels or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 15, preferably at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term polynucleotides.

Polynucleotides such as DNA polynucleotides and probes may be produced
10 recombinantly, synthetically or by any means available to those of skill in the art. They may also be cloned by standard techniques. In general, primers will be produced by synthetic means, involving a stepwise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

15 Longer polynucleotides will generally be produced using recombinant means, for example using a PCR (polymerase chain reaction) cloning techniques. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector. Preferably, the variant sequences are at least as biologically active as the sequences presented herein.

20 A preferred embodiment of the invention includes sequences that are complementary to the modified enzyme or sequences that are capable of hybridising either to the nucleotide sequences of the modified enzymes (including complementary sequences of those presented herein), as well as nucleotide sequences that are complementary to sequences that can hybridise to the nucleotide sequences of the modified enzymes
25 (including complementary sequences of those presented herein). A preferred embodiment provides polynucleotide sequences that are capable of hybridising to the nucleotide sequences presented herein under conditions of intermediate to maximal stringency.

A preferred embodiment includes nucleotide sequences that can hybridise to the nucleotide sequence of the modified enzyme nucleic acid, or the complement thereof, under
30 stringent conditions (e.g. 50°C and 0.2xSSC). More preferably, the nucleotide sequences can hybridise to the nucleotide sequence of the modified enzyme, or the complement thereof, under high stringent conditions (e.g. 65°C and 0.1xSSC).

It may be desirable to mutate the sequence in order to prepare a modified enzyme. Accordingly, a mutant may be prepared from the modified enzymes provided herein. Mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites. A suitable method is disclosed in Morinaga *et al.*, (*Biotechnology* (1984) 2, p646-649). Another method of
5 introducing mutations into enzyme-encoding nucleotide sequences is described in Nelson and Long (*Analytical Biochemistry* (1989), 180, p 147-151). A further method is described in Sarkar and Sommer (*Biotechniques* (1990), 8, p404-407 – “The megaprimer method of site directed mutagenesis”). Other methods to mutate the sequence are employed and
10 disclosed herein.

In a preferred embodiment, the sequence for use in the methods and compositions described here is a recombinant sequence – i.e. a sequence that has been prepared using recombinant DNA techniques. Such techniques are explained, for example, in the literature, for example, J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular*
15 *Cloning: A Laboratory Manual*, Second Edition, Books 1-3, Cold Spring Harbor Laboratory Press.

Another embodiment provides for compositions and formulations comprising modified enzymes. The compositions include the modified enzyme together with another component.

20 Another embodiment provides vectors comprising the modified enzyme, cells comprising the modified enzyme and methods of expressing the modified enzyme. The nucleotide sequence for use in the methods and compositions described herein may be incorporated into a recombinant replicable vector. The vector may be used to replicate and express the nucleotide sequence, in enzyme form, in and/or from a compatible host cell. Expression may be controlled using control sequences, e.g., regulatory sequences. The
25 enzyme produced by a host recombinant cell by expression of the nucleotide sequence may be secreted or may be contained intracellularly depending on the sequence and/or the vector used. The coding sequences may be designed with signal sequences which direct secretion of the substance coding sequences through a particular prokaryotic or eukaryotic cell
30 membrane. Polynucleotides can be incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. The vector

comprising the polynucleotide sequence may be transformed into a suitable host cell.

Suitable hosts may include bacterial, yeast, insect and fungal cells.

Modified enzymes and their polynucleotides may be expressed by introducing a polynucleotide into a replicable vector, introducing the vector into a compatible host cell
5 and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell.

The modified enzyme nucleic acid may be operatively linked to transcriptional and translational regulatory elements active in a host cell of interest. The modified enzyme nucleic acid may also encode a fusion protein comprising signal sequences such as, for
10 example, those derived from the glucoamylase gene from *Schwanniomyces occidentalis*, α -factor mating type gene from *Saccharomyces cerevisiae* and the TAKA-amylase from *Aspergillus oryzae*. Alternatively, the modified enzyme nucleic acid may encode a fusion protein comprising a membrane binding domain.

The modified enzyme may be expressed at the desired levels in a host organism
15 using an expression vector. An expression vector comprising a modified enzyme nucleic acid can be any vector capable of expressing the gene encoding the modified enzyme nucleic acid in the selected host organism, and the choice of vector will depend on the host cell into which it is to be introduced. Thus, the vector can be an autonomously replicating vector, i.e. a vector that exists as an episomal entity, the replication of which is independent
20 of chromosomal replication, such as, for example, a plasmid, a bacteriophage or an episomal element, a minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome.

The expression vector typically includes the components of a cloning vector, such
25 as, for example, an element that permits autonomous replication of the vector in the selected host organism and one or more phenotypically detectable markers for selection purposes. The expression vector normally comprises control nucleotide sequences encoding a promoter, operator, ribosome binding site, translation initiation signal and optionally, a repressor gene or one or more activator genes. Additionally, the expression vector may
30 comprise a sequence coding for an amino acid sequence capable of targeting the modified enzyme to a host cell organelle such as a peroxisome or to a particular host cell compartment. Such a targeting sequence includes but is not limited to the sequence SKL.

For expression under the direction of control sequences, the nucleic acid sequence the modified enzyme is operably linked to the control sequences in proper manner with respect to expression.

Preferably, a polynucleotide in a vector is operably linked to a control sequence that
5 is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The control sequences may be modified, for example, by the addition of further transcriptional regulatory elements to make the level of transcription directed by the control sequences more responsive to transcriptional modulators. The control sequences may in particular comprise promoters.

10 In the vector, the nucleic acid sequence encoding for the modified enzyme is operably combined with a suitable promoter sequence. The promoter can be any DNA sequence having transcription activity in the host organism of choice and can be derived from genes that are homologous or heterologous to the host organism. Examples of
15 suitable promoters for directing the transcription of the modified nucleotide sequence, such as modified enzyme nucleic acids, in a bacterial host include the promoter of the *lac* operon of *E. coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis* α -amylase gene (*amyL*), the *aprE* promoter of *Bacillus subtilis*, the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus amyloliquefaciens* α -amylase gene (*amyQ*), the promoters of the
20 *Bacillus subtilis* *xylA* and *xylB* genes and a promoter derived from a *Lactococcus* sp.-derived promoter including the P170 promoter. When the gene encoding the modified enzyme is expressed in a bacterial species such as *E. coli*, a suitable promoter can be selected, for example, from a bacteriophage promoter including a T7 promoter and a phage lambda promoter. For transcription in a fungal species, examples of useful promoters are
25 those derived from the genes encoding the, *Aspergillus oryzae* TKA amyase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral α -amylase, *A. niger* acid stable α -amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase or *Aspergillus nidulans* acetamidase. Examples of suitable promoters for the expression in a yeast species include
30 but are not limited to the Gal 1 and Gal 10 promoters of *Saccharomyces cerevisiae* and the *Pichia pastoris* AOX1 or AOX2 promoters.

Examples of suitable bacterial host organisms are gram positive bacterial species such as *Bacillaceae* including *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus megaterium* and *Bacillus thuringiensis*, *Streptomyces* species such as *Streptomyces murinus*, lactic acid bacterial species including *Lactococcus* spp. such as *Lactococcus lactis*, *Lactobacillus* spp. including *Lactobacillus reuteri*, *Leuconostoc* spp., *Pediococcus* spp. and *Streptococcus* spp. Alternatively, strains of a gram-negative bacterial species belonging to *Enterobacteriaceae* including *E. coli*, or to *Pseudomonadaceae* can be selected as the host organism. A suitable yeast host organism can be selected from the biotechnologically relevant yeasts species such as but not limited to yeast species such as *Pichia* sp., *Hansenula* sp or *Kluyveromyces*, *Yarrowinia* species or a species of *Saccharomyces* including *Saccharomyces cerevisiae* or a species belonging to *Schizosaccharomyce* such as, for example, *S. Pombe* species. Preferably a strain of the methylotrophic yeast species *Pichia pastoris* is used as the host organism. Preferably the host organism is a *Hansenula* species. Suitable host organisms among filamentous fungi include species of *Aspergillus*, e.g. *Aspergillus niger*, *Aspergillus oryzae*, *Aspergillus tubigensis*, *Aspergillus awamori* or *Aspergillus nidulans*. Alternatively, strains of a *Fusarium* species, e.g. *Fusarium oxysporum* or of a *Rhizomucor* species such as *Rhizomucor miehei* can be used as the host organism. Other suitable strains include *Thermomyces* and *Mucor* species.

Host cells comprising polynucleotides may be used to express polypeptides, such as the modified enzymes disclosed herein, fragments, homologues, variants or derivatives thereof. Host cells may be cultured under suitable conditions which allow expression of the proteins. Expression of the polypeptides may be constitutive such that they are continually produced, or inducible, requiring a stimulus to initiate expression. In the case of inducible expression, protein production can be initiated when required by, for example, addition of an inducer substance to the culture medium, for example dexamethasone or IPTG. Polypeptides can be extracted from host cells by a variety of techniques known in the art, including enzymatic, chemical and/or osmotic lysis and physical disruption. Polypeptides may also be produced recombinantly in an *in vitro* cell-free system, such as the TnTTM (Promega) rabbit reticulocyte system.

In a third aspect, the invention is drawn to a method of modifying an enzyme comprising modifying a first site in the enzyme part of a structurally defined region so that the first site can bind to a second site. In a preferred embodiment, the first site is in a loop or sequence adjacent to a β -sheet. In a preferred embodiment, the second site is located in a β -sheet. In a preferred embodiment, the modified enzyme is a xylanase or Clan C.

In a preferred embodiment, the invention is drawn to a modified xylanase or a method of modifying a xylanase (or modified enzyme), according to at least one of the following: (i) modifying the N-terminal sequence so that the N-terminal region is bound by a disulphide bridge to an adjacent β -strand (*see* Gruber, *et al.*, 1998 in *T. reesei* XynII the amino acids 1-4 and 24-30 respectively); (ii) modifying the C-terminal (in *T. reesei* XynII amino acids 183-190, *see* Gruber, *et al.*, 1998) so that it is bound to an adjacent β -strand; (iii) modifying an α -helix of the enzyme so that it can be bound more tightly to the body of the protein; (iv) modifying at least one adjacent loop so that it binds adjacent beta strand B6a (in *T. reesei* XynII amino acids 91-94, Gruber, *et al.*, 1998) or (v) modifying residue equivalent to XynII, as provided above.

As another embodiment, (per the examples) mutagenesis may be used to create disulphide bridges, salt bridges and separate point mutations at different regions. For example, the enzyme may be modified to create at least one disulphide bridge, so that at least one disulphide bridge may: 1) stabilize the N-terminal region or bind the N-terminal beta strand to the adjacent beta sheet (positions 2-28, 5-19, 7-16, 10-29 in XynII, or an equivalent position, as disclosed herein); 2) stabilize the alpha helix region (positions 105-162, 57-153, 110-151, 111-151, in XynII, or an equivalent position as disclosed herein); 3) stabilize the C-terminal region (positions 63-188, 61-190, 36-186 or 34-188 in XynII, or an equivalent position as disclosed herein); or 4) stabilize the loop by binding to the beta strand such as B6b (92-144, 113-143 in XynII or an equivalent position as disclosed herein) and/or 5) stabilize the beta sheet (positions 26-38, 61-149, 63-147, 65-186, 67-184 in XynII, or an equivalent position, as provided herein).

Salt bridges may be created at different sites of the enzyme: (e.g., positions 22, 180, 58 or +191D in XynII, or an equivalent position, as provided herein) and single point mutations may be introduced at different sites of the molecule (e.g., positions 108, 26, 30, 67, 93, 97, 132, 157, 160, 165, 169 or 186 in XynII, or an equivalent position, as provided herein) thereby increasing the thermostability and/or thermophilicity and or alkalophilicity

the protein. As with the Y5 mutant, the C-terminus may be bound more tightly to the body of the enzyme by adding as a recombinant change one amino acid (e.g. aspartic acid or glutamic acid) which then can form a salt bridge from the C-terminus to the body of the enzyme. If appropriate, a suitable amino acid replacement can be made in the body of the protein, so as to enable the formation of a salt bridge or to stabilize the enzyme in the C-terminal part via the α -helix or a region near the α -helix.

Additional mutants can be created according to this aspect of the invention. The structure of the N-terminal beta strand A1 or N-terminal loop in family 11 and 12 enzymes is described as the beta strand, a part of the beta sheet A prior to/up to a beta bend structure leading to beta strand B1 or the N-terminal loop prior to the first beta strand of the beta sheet. (see, Törrönen et al., Biochemistry 1995, 34, 847-856; Sandgren, et. al., J. Mol. Bio. (2001) 308, 295-310; Gruber, *et al.*, 1998). The B1 beta strand of the N-terminal region is described as the beta strand part of the beta sheet B prior to/up to a beta bend structure leading to beta strand B2 or the loop prior to the first beta strand of the beta sheet. The beta strand A1 region is bound preferably to beta strand A2 or to any other adjacent region (XynII or an equivalent thereof). The beta strand B1 region is bound preferably to beta strand B2 or to any other adjacent region (XynII or an equivalent thereof). In XynII A1 comprises residues 1-4, A2 residues 25-30, B1 residues 6-10 and B2 residues 13-19.

The structure of the C-terminal beta strand A4 or C-terminal loop in family 11 and 12 enzymes is the beta strand part of the beta sheet A between beta strands A3 and A5 or the loop as following beta sheet A4 (see Törrönen et al., Biochemistry 1995, 34, 847-856; Sandgren, et. al., J. Mol. Bio. (2001) 308, 295-310; Gruber, *et al.*, 1998). The beta strand A4 region is bound preferably to beta strand A3 or A5, or to any other adjacent region. In XynII A4 is residues 183-190, A3 is residues 33-39 and A5 is residues 61-69. The cord of family 11 and 12 is described as the loop connecting beta strands B6b and B9. The beta strand of family 11 and 12 B6b is described as the beta strand prior to the cord (Törrönen et al., Biochemistry 1995, 34, 847-856; Sandgren, et. al., J. Mol. Bio. (2001) 308, 295-310; Gruber, *et al.*, 1998). The beta strand B6b region may be bound to the cord or to the loop between beta strands A6 and B7, or to any other adjacent region. In XynII, B6b is residues 90-94 and B9 is residues 103-110, the cord is 95-102, beta strand A6 is residues 148-152, beta strand B7 is residues 134-142 and the loop between beta stands A6 and B7 is residues 143-147.

The helix of family 11 and 12 enzymes is described as region following beta strand A6 and forming a helical structure parallel to beta strand B9 (Törrönen et al., Biochemistry 1995, 34, 847-856; Sandgren, et. al., J. Mol.). The helix of family 11 and 12 enzymes is bound preferably to beta strand B9 or any other adjacent region. In XynII the helix is
5 residues 153-162, beta strand A6 is residues 148-152 and beta strand B9 is residues 103-110.

EXAMPLES

EXAMPLE 1.

10 Plasmids used for xylanase II expression and mutagenesis template

The open reading frame encoding *Trichoderma reesei* XYNII gene product was amplified by polymerase chain reaction (PCR) from the *T. reesei* cDNA library. XYNII cDNA was cloned into pKKtac (VTT, Espoo, Finland) or alternatively into pALK143 (ROAL, Rajamäki, Finland).

15 EXAMPLE 2.

Site-directed mutagenesis for generation of mutant of xylanase II

Expression vectors containing cDNA-encoding xylanase II as described in Example 1 were used as template in the stepwise site-directed mutagenesis in consecutive PCR amplifications. Synthetic oligonucleotide primers containing the altered codons for the
20 mutations X-Y were used for insertion of the desired alteration into the native xylanase II primary amino acid sequence. By this approach the residues of sites 92, 93 and 144 of the wild-type enzyme mutants were generated to bind the loop N143- S146 of xynII to the neighbouring β -strand. Additionally, mutagenesis was performed to generate the mutations at sites 22, 65, 97 and 108 into the xylanase primary sequence. The oligonucleotide
25 sequences used in the mutagenesis are shown Figure 3. PCR was carried out as described in the Quick Change Site-directed mutagenesis (Stratagene, La Jolla, Ca, USA) according to standard PCR procedures. *Pfu*Turbo (Stratagene) was used as DNA polymerase to amplify plasmid DNA. Plasmid DNA from the site-directed mutagenesis PCR amplification was transformed to *E. coli* XL-1 blue and the transformed bacterial cells were
30 then propagated on LB, with ampicillin 100 ug/ml for plasmid DNA selection and amplification of the mutated DNA. Plasmids were isolated and sequenced to confirm that they contain the desired mutations. The mutated plasmid DNA encoding the mutant

variants was over- expressed in *E. coli* to examine the influence of the mutagenesis on the *T. reesei* xylanase Y5 mutants enzymatic properties.

EXAMPLE 3.

5 **Production of the modified *XYNII* gene products in *E. coli* strain and assay for xylanase activity**

E. coli strains over-expressing the mutated variants of the xylanase II were cultivated on plates supplemented with 1% birchwood xylan (Sigma, Steinheim, Germany) coupled with Rhemazol Brilliant Blue. Rhemazol Brilliant Blue coupled to xylan was
10 utilized to detect xylanase activity that was readily visualized by a characteristic halo formation due to the blue colour disappearance around the bacterial colonies expressing xylanase activity (Biely *et al.*, 1985).

The mutated xylanase genes (see above; Example 2) were expressed in *E. coli* at +37°C in shake flasks in LB culture medium. Cell cultures expressing the enzyme variants
15 were centrifuged and the cell pellet separated from the supernatant harbouring the enzyme that was secreted from the cells into the culture medium. The xylanase enzyme activity assay was performed according to standard methods. The growth medium containing the secreted xylanase mutants were incubated for 10 min in 1% birchwood xylan (Sigma) at 50°C in 50mM citrate-phosphate buffer (ph 5.0 -t) and 50 mM Tris-HCl at pH 7-9. (Bailey
20 *et al.*, 1992). If needed, heat inactivated growth medium was used to dilute the samples. The enzymatic activity of the mutant variants was examined in comparison to the wild type and the Y5 mutation enzyme at varying conditions (*see*, for Bailey *et al.*, 1992).

EXAMPLE 4.

25 **Determination of the temperature dependent stability and pH dependent activity of the xylanase II mutants**

Activity as a function of temperature;

The xylanase activity of the mutant variants was determined at varying temperatures and selected pH values (see Figures herein). The mutants were incubated for 10 min with
30 1% birchwood xylan (Sigma) in 50mM citrate-phosphate buffer (ph 4.5-7) or 50 mM Tris-HCl at pH 7-9. The relative amount of released reducing sugars was detected with the DNS method assay as described in example 3.

Residual activity

The mutant variants were incubated for 10 minutes at varying temperatures without substrate. After the inactivation, the samples were cooled on ice and the residual activity
5 was determined by DNS-method as described in example 3.

pH dependent activity

The pH-dependent xylanase activity was determined by detecting the enzyme activity at varying pH ranging from XX – YY for 10 min in 1% birchwood xylan at selected
10 temperatures (see pictures) in 50mM citrate-phosphate buffer (ph 4.5-7) and 50 mM Tris-HCl at pH 7,5-9. This was followed by the DNS assay as described in example 3.

EXAMPLE 5.**Preparation and Testing of mutant xylanases for improved properties**

15 Mutant xylanases were prepared having substitutions at one or more substitutions at different regions of the molecule. The substitutions were either separate point mutations in contact with other separate point mutations or they were prepared to act on a structural element found commonly in both family 11 and family 12 enzymes. The enzyme assays were performed as outlined in the examples. Examples of “structural” substitutions are
20 disclosed herein and shown in the examples.

The disulphide bridge can be placed between sites 2 and 28 (T2C, T28C). Figure 4 shows the importance of the N-terminal region in substituting residues of the wt for a more thermophilic variant. In a similar way removal of the native disulphide bridge (residues C4 and C32, Cell12A numbering) of T.reesei EGIII affects greatly the stability of the enzyme,
25 as shown in the figures provided and tables herein (*see*, especially, Table A).

The region of the beta sheet common to both family 11 and 12 named beta strand B6b (as in Gruber et al), is shown to be of importance for stability, especially at alkali conditions. This effect is seen in the substitutions (as compared to the Y5 variant) as improved stability at pH 9 vs pH5 for P12, as shown in the figures (*see*, for example, Figure
30 9, Figure 10 and Figure 11).

The importance of the region is clearly demonstrated by a different set of mutations (although in the same region) affecting the same beta strand. When sites 93, 97 and 144 are

substituted (F93W, N97R, H144K, *P9* in the graph), a similar effect in stabilization of the enzyme as when substituting the sites 92 and 144 (N92C, H144C= *P12* in the graph) can be seen in the Figure 9.

An example of the improved characteristics of separate substitutions at sites 22 and
5 180 is seen below. The variant containing the substitutions H22K and F180Q (*P20* in Figure 14) shows enhanced thermal stability over Y5 at pH 7.8.

Also the C-terminal region is of important for stability. In the substitution S65C, S186C (*J21* in the graph) the enzyme shows improved activity with respect to temperature at pH 8.

10

15

All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains.

20

25

30

The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether
5 or not the excised material is specifically recited herein.

SEQUENCE LISTING

<110> Genencor International, Inc.

<120> Modified Enzymes, Methods to Produce
Modified Enzymes and Uses Thereof

<130> 11816-119

<140> CA 2,538,837

<141> 2004-09-10

<150> US 60/503,251

<151> 2003-09-15

<160> 51

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 223

<212> PRT

<213> Trichoderma reesei

<400> 1

Met	Val	Ser	Phe	Thr	Ser	Leu	Leu	Ala	Gly	Val	Ala	Ala	Ile	Ser	Gly	1	5	10	15
Val	Leu	Ala	Ala	Pro	Ala	Ala	Glu	Val	Glu	Ser	Val	Ala	Val	Glu	Lys	20	25	30	
Arg	Gln	Thr	Ile	Gln	Pro	Gly	Thr	Gly	Tyr	Asn	Asn	Gly	Tyr	Phe	Tyr	35	40	45	
Ser	Tyr	Trp	Asn	Asp	Gly	His	Gly	Gly	Val	Thr	Tyr	Thr	Asn	Gly	Pro	50	55	60	
Gly	Gly	Gln	Phe	Ser	Val	Asn	Trp	Ser	Asn	Ser	Gly	Asn	Phe	Val	Gly	65	70	75	80
Gly	Lys	Gly	Trp	Gln	Pro	Gly	Thr	Lys	Asn	Lys	Val	Ile	Asn	Phe	Ser	85	90	95	
Gly	Ser	Tyr	Asn	Pro	Asn	Gly	Asn	Ser	Tyr	Leu	Ser	Val	Tyr	Gly	Trp	100	105	110	
Ser	Arg	Asn	Pro	Leu	Ile	Glu	Tyr	Tyr	Ile	Val	Glu	Asn	Phe	Gly	Thr	115	120	125	
Tyr	Asn	Pro	Ser	Thr	Gly	Ala	Thr	Lys	Leu	Gly	Glu	Val	Thr	Ser	Asp	130	135	140	
Gly	Ser	Val	Tyr	Asp	Ile	Tyr	Arg	Thr	Gln	Arg	Val	Asn	Gln	Pro	Ser	145	150	155	160
Ile	Ile	Gly	Thr	Ala	Thr	Phe	Tyr	Gln	Tyr	Trp	Ser	Val	Arg	Arg	Asn	165	170	175	
His	Arg	Ser	Ser	Gly	Ser	Val	Asn	Thr	Ala	Asn	His	Phe	Asn	Ala	Trp	180	185	190	
Ala	Gln	Gln	Gly	Leu	Thr	Leu	Gly	Thr	Met	Asp	Tyr	Gln	Ile	Val	Ala	195	200	205	
Val	Glu	Gly	Tyr	Phe	Ser	Ser	Gly	Ser	Ala	Ser	Ile	Thr	Val	Ser		210	215	220	

<210> 2

<211> 781

<212> DNA

<213> Trichoderma reesei

42

<400> 2

```

atggtctcct tcacctccct cctcgccggc gtcgccgcca tctcggggcgt cttggccgct      60
cccgccgccc aggtcgaatc cgtggctgtg gagaagcgcc agacgattca gcccggcacg      120
ggctacaaca acggctactt ctactcgta tggaaacgat gccacggcgg cgtgacgtac      180
accaatggtc ccggcgggca gttctccgtc aactgggtcca actcgggcaa ctttgtcggc      240
ggcaagggat ggcagcccgg caccaagaac aagtaagact acctactctt accccctttg      300
accaacacag cacaacacaa tacaacacat gtgactacca atcatggaat cggatctaac      360
agctgtgttt tcaaaaaaaaa gggatcatcaa cttctcgggc agctacaacc ccaacggcaa      420
cagctacctc tccgtgtacg gctgggtccc caaccccttg atcgagtact acatcgtcga      480
gaactttggc acctacaacc cgtccacggg cgccaccaag ctgggcgagg tcacctccga      540
cggcagcgtc tacgacattt accgcacgca gcgcgtcaac cagccgtcca tcatcggcac      600
cgccaccttt taccagtact ggtccgtccg ccgcaaccac cgctcgagcg gctccgtcaa      660
cacggcgaac cacttcaacg cgtggggtca gcaaggcctg acgctcggga cgatggatta      720
ccagattggt gccgtggagg gttactttag ctctggctct gcttccatca ccgtcagcta      780
a                                                                                   781

```

<210> 3

<211> 234

<212> PRT

<213> Trichoderma reesei

<400> 3

```

Met Lys Phe Leu Gln Val Leu Pro Ala Leu Ile Pro Ala Ala Leu Ala
 1              5              10              15
Gln Thr Ser Cys Asp Gln Trp Ala Thr Phe Thr Gly Asn Gly Tyr Thr
      20              25              30
Val Ser Asn Asn Leu Trp Gly Ala Ser Ala Gly Ser Gly Phe Gly Cys
      35              40              45
Val Thr Ala Val Ser Leu Ser Gly Gly Ala Ser Trp His Ala Asp Trp
      50              55              60
Gln Trp Ser Gly Gly Gln Asn Asn Val Lys Ser Tyr Gln Asn Ser Gln
      65              70              75              80
Ile Ala Ile Pro Gln Lys Arg Thr Val Asn Ser Ile Ser Ser Met Pro
      85              90              95
Thr Thr Ala Ser Trp Ser Tyr Ser Gly Ser Asn Ile Arg Ala Asn Val
      100             105             110
Ala Tyr Asp Leu Phe Thr Ala Ala Asn Pro Asn His Val Thr Tyr Ser
      115             120             125
Gly Asp Tyr Glu Leu Met Ile Trp Leu Gly Lys Tyr Gly Asp Ile Gly
      130             135             140
Pro Ile Gly Ser Ser Gln Gly Thr Val Asn Val Gly Gly Gln Ser Trp
      145             150             155             160
Thr Leu Tyr Tyr Gly Tyr Asn Gly Ala Met Gln Val Tyr Ser Phe Val
      165             170             175
Ala Gln Thr Asn Thr Thr Asn Tyr Ser Gly Asp Val Lys Asn Phe Phe
      180             185             190
Asn Tyr Leu Arg Asp Asn Lys Gly Tyr Asn Ala Ala Gly Gln Tyr Val
      195             200             205
Leu Ser Tyr Gln Phe Gly Thr Glu Pro Phe Thr Gly Ser Gly Thr Leu
      210             215             220
Asn Val Ala Ser Trp Thr Ala Ser Ile Asn
225              230

```

<210> 4

<211> 826

<212> DNA

<213> Trichoderma reesei

<400> 4

43

```

atgaagttcc ttcaagtcct ccctgccctc ataccggccg ccctggccca aaccagctgt      60
gaccagtggg caaccttcac tggcaacggc tacacagtca gcaacaacct ttggggagca      120
tcagccggct ctggatttgg ctgcgtgacg gcggtatcgc tcagcggcgg ggcctcctgg      180
cacgcagact ggcagtggtc cggcggccag aacaacgtca agtcgtacca gaactctcag      240
attgccattc cccagaagag gaccgtcaac agcatcagca gcatgcccac cactgccagc      300
tggagctaca gcgggagcaa catccgcgct aatgttgcgt atgacttggt caccgcagcc      360
aaccggaatc atgtcacgta ctcgggagac tacgaactca tgatctggta agccataaga      420
agtgaccctc cttgatagtt tcgactaaca acatgtcttg aggcttggca aatacggcga      480
tattggggccg attgggtcct cacaggggaa agtcaacgtc ggtggccaga gctggacgct      540
ctactatggc tacaacggag ccatgcaagt ctattccttt gtggcccaga ccaacactac      600
caactacagc ggagatgtca agaacttctt caattatctc cgagacaata aaggatacaa      660
cgctgcaggc caatatgttc ttagtaagtc accctcactg tgactgggct gagtttggtg      720
caacgtttgc taacaaaacc ttcgtatagg ctaccaattt ggtaccgagc ccttcacggg      780
cagtgggaact ctgaacgtcg catcctggac cgcactatc aactaa      826

```

<210> 5

<211> 222

<212> PRT

<213> *Trichoderma reesei*

<400> 5

```

Met Val Ser Phe Thr Ser Leu Leu Ala Ala Ser Pro Pro Ser Arg Ala
 1              5              10              15
Ser Cys Arg Pro Ala Ala Glu Val Glu Ser Val Ala Val Glu Lys Arg
              20              25              30
Gln Thr Ile Gln Pro Gly Thr Gly Tyr Asn Asn Gly Tyr Phe Tyr Ser
              35              40              45
Tyr Trp Asn Asp Gly His Gly Gly Val Thr Tyr Thr Asn Gly Pro Gly
              50              55              60
Gly Gln Phe Ser Val Asn Trp Ser Asn Ser Gly Asn Phe Val Gly Gly
65              70              75              80
Lys Gly Trp Gln Pro Gly Thr Lys Asn Lys Val Ile Asn Phe Ser Gly
              85              90              95
Ser Tyr Asn Pro Asn Gly Asn Ser Tyr Leu Ser Val Tyr Gly Trp Ser
              100              105              110
Arg Asn Pro Leu Ile Glu Tyr Tyr Ile Val Glu Asn Phe Gly Thr Tyr
              115              120              125
Asn Pro Ser Thr Gly Ala Thr Lys Leu Gly Glu Val Thr Ser Asp Gly
              130              135              140
Ser Val Tyr Asp Ile Tyr Arg Thr Gln Arg Val Asn Gln Pro Ser Ile
145              150              155              160
Ile Gly Thr Ala Thr Phe Tyr Gln Tyr Trp Ser Val Arg Arg Asn His
              165              170              175
Arg Ser Ser Gly Ser Val Asn Thr Ala Asn His Phe Asn Ala Trp Ala
              180              185              190
Gln Gln Gly Leu Thr Leu Gly Thr Met Asp Tyr Gln Ile Val Ala Val
              195              200              205
Glu Gly Tyr Phe Ser Ser Gly Ser Ala Ser Ile Thr Val Ser
              210              215              220

```

<210> 6

<211> 227

<212> PRT

<213> *Humicola insolens*

<400> 6

```

Met Val Ser Leu Lys Ser Val Leu Ala Ala Ala Thr Ala Val Ser Ser
 1              5              10              15
Ala Ile Ala Ala Pro Phe Asp Phe Val Pro Arg Asp Asn Ser Thr Ala
              20              25              30

```

44

```

Leu Gln Ala Arg Gln Val Thr Pro Asn Ala Glu Gly Trp His Asn Gly
   35          40          45
Tyr Phe Tyr Ser Trp Trp Ser Asp Gly Gly Gly Gln Val Gln Tyr Thr
   50          55          60
Asn Leu Glu Gly Ser Arg Tyr Gln Val Arg Trp Arg Asn Thr Gly Asn
  65          70          75          80
Phe Val Gly Gly Lys Gly Trp Asn Pro Gly Thr Gly Arg Thr Ile Asn
          85          90          95
Tyr Gly Gly Tyr Phe Asn Pro Gln Gly Asn Gly Tyr Leu Ala Val Tyr
          100          105          110
Gly Trp Thr Arg Asn Pro Leu Val Glu Tyr Tyr Val Ile Glu Ser Tyr
          115          120          125
Gly Thr Tyr Asn Pro Gly Ser Gln Ala Gln Tyr Lys Gly Thr Phe Tyr
          130          135          140
Thr Asp Gly Asp Gln Tyr Asp Ile Phe Val Ser Thr Arg Tyr Asn Gln
 145          150          155          160
Pro Ser Ile Asp Gly Thr Arg Thr Phe Gln Gln Tyr Trp Ser Ile Arg
          165          170          175
Lys Asn Lys Arg Val Gly Gly Ser Val Asn Met Gln Asn His Phe Asn
          180          185          190
Ala Trp Gln Gln His Gly Met Pro Leu Gly Gln His Tyr Tyr Gln Val
          195          200          205
Val Ala Thr Glu Gly Tyr Gln Ser Ser Gly Glu Ser Asp Ile Tyr Val
          210          215          220
Gln Thr His
225

```

<210> 7

<211> 210

<212> PRT

<213> *Bacillus stearothermophilus*

<400> 7

```

Met Lys Leu Lys Lys Lys Met Leu Thr Leu Leu Leu Thr Ala Ser Met
  1          5          10          15
Ser Phe Gly Leu Phe Gly Ala Thr Ser Ser Ala Ala Thr Asp Tyr Trp
          20          25          30
Gln Tyr Trp Thr Asp Gly Gly Gly Met Val Asn Ala Val Asn Gly Pro
          35          40          45
Gly Gly Asn Tyr Ser Val Thr Trp Gln Asn Thr Gly Asn Phe Val Val
          50          55          60
Gly Lys Gly Trp Thr Val Gly Ser Pro Asn Arg Val Ile Asn Tyr Asn
  65          70          75          80
Ala Gly Ile Trp Glu Pro Ser Gly Asn Gly Tyr Leu Thr Leu Tyr Gly
          85          90          95
Trp Thr Arg Asn Ala Leu Ile Glu Tyr Tyr Val Val Asp Ser Trp Gly
          100          105          110
Thr Tyr Arg Pro Thr Gly Asn Tyr Lys Gly Thr Val Asn Ser Asp Gly
          115          120          125
Gly Thr Tyr Asp Ile Tyr Thr Thr Met Arg Tyr Asn Ala Pro Ser Ile
          130          135          140
Asp Gly Thr Gln Thr Phe Gln Gln Phe Trp Ser Val Arg Gln Ser Lys
 145          150          155          160
Arg Pro Thr Gly Ser Asn Val Ser Ile Thr Phe Ser Asn His Val Asn
          165          170          175
Ala Trp Arg Ser Lys Gly Met Asn Leu Gly Ser Ser Trp Ala Tyr Gln
          180          185          190
Val Leu Ala Thr Glu Gly Tyr Gln Ser Ser Gly Arg Ser Asn Val Thr
          195          200          205
Val Trp

```

45

210

<210> 8
 <211> 229
 <212> PRT
 <213> *Trichoderma reesei*

<400> 8
 Met Val Ala Phe Ser Ser Leu Ile Cys Ala Leu Thr Ser Ile Ala Ser
 1 5 10 15
 Thr Leu Ala Met Pro Thr Gly Leu Glu Pro Glu Ser Ser Val Asn Val
 20 25 30
 Thr Glu Arg Gly Met Tyr Asp Phe Val Leu Gly Ala His Asn Asp His
 35 40 45
 Arg Arg Arg Ala Ser Ile Asn Tyr Asp Gln Asn Tyr Gln Thr Gly Gly
 50 55 60
 Gln Val Ser Tyr Ser Pro Ser Asn Thr Gly Phe Ser Val Asn Trp Asn
 65 70 75 80
 Thr Gln Asp Asp Phe Val Val Gly Val Gly Trp Thr Thr Gly Ser Ser
 85 90 95
 Ala Pro Ile Asn Phe Gly Gly Ser Phe Ser Val Asn Ser Gly Thr Gly
 100 105 110
 Leu Leu Ser Val Tyr Gly Trp Ser Thr Asn Pro Leu Val Glu Tyr Tyr
 115 120 125
 Ile Met Glu Asp Asn His Asn Tyr Pro Ala Gln Gly Thr Val Lys Gly
 130 135 140
 Thr Val Thr Ser Asp Gly Ala Thr Tyr Thr Ile Trp Glu Asn Thr Arg
 145 150 155 160
 Val Asn Glu Pro Ser Ile Gln Gly Thr Ala Thr Phe Asn Gln Tyr Ile
 165 170 175
 Ser Val Arg Asn Ser Pro Arg Thr Ser Gly Thr Val Thr Val Gln Asn
 180 185 190
 His Phe Asn Ala Trp Ala Ser Leu Gly Leu His Leu Gly Gln Met Asn
 195 200 205
 Tyr Gln Val Val Ala Val Glu Gly Trp Gly Gly Ser Gly Ser Ala Ser
 210 215 220
 Gln Ser Val Ser Asn
 225

<210> 9
 <211> 211
 <212> PRT
 <213> *Aspergillus awamori*

<400> 9
 Met Lys Val Thr Ala Ala Phe Ala Gly Leu Leu Val Thr Ala Phe Ala
 1 5 10 15
 Ala Pro Val Pro Glu Pro Val Leu Val Ser Arg Ser Ala Gly Ile Asn
 20 25 30
 Tyr Val Gln Asn Tyr Asn Gly Asn Leu Gly Asp Phe Thr Tyr Asp Glu
 35 40 45
 Ser Ala Gly Thr Phe Ser Met Tyr Trp Glu Asp Gly Val Ser Ser Asp
 50 55 60
 Phe Val Val Gly Leu Gly Trp Thr Thr Gly Ser Ser Asn Ala Ile Thr
 65 70 75 80
 Tyr Ser Ala Glu Tyr Ser Ala Ser Gly Ser Ser Ser Tyr Leu Ala Val
 85 90 95
 Tyr Gly Trp Val Asn Tyr Pro Gln Ala Glu Tyr Tyr Ile Val Glu Asp
 100 105 110
 Tyr Gly Asp Tyr Asn Pro Cys Ser Ser Ala Thr Ser Leu Gly Thr Val

46

		115					120				125					
Tyr	Ser	Asp	Gly	Ser	Thr	Tyr	Gln	Val	Cys	Thr	Asp	Thr	Arg	Thr	Asn	
	130					135					140					
Glu	Pro	Ser	Ile	Thr	Gly	Thr	Ser	Thr	Phe	Thr	Gln	Tyr	Phe	Ser	Val	
145					150					155					160	
Arg	Glu	Ser	Thr	Arg	Thr	Ser	Gly	Thr	Val	Thr	Val	Ala	Asn	His	Phe	
				165					170					175		
Asn	Phe	Trp	Ala	Gln	His	Gly	Phe	Gly	Asn	Ser	Asp	Phe	Asn	Tyr	Gln	
			180				185						190			
Val	Met	Ala	Val	Glu	Ala	Trp	Ser	Gly	Ala	Gly	Ser	Ala	Ser	Val	Thr	
		195					200					205				
Ile	Ser	Ser														
	210															

<210> 10

<211> 330

<212> PRT

<213> Bacillus stearothermophilus

<400> 10

Met	Cys	Ser	Ser	Ile	Pro	Ser	Leu	Arg	Glu	Val	Phe	Ala	Asn	Asp	Phe	
1				5					10					15		
Arg	Ile	Gly	Ala	Ala	Val	Asn	Pro	Val	Thr	Leu	Glu	Ala	Gln	Gln	Ser	
			20					25					30			
Leu	Leu	Ile	Arg	His	Val	Asn	Ser	Leu	Thr	Ala	Glu	Asn	His	Met	Lys	
		35				40						45				
Phe	Glu	His	Leu	Gln	Pro	Glu	Glu	Gly	Arg	Phe	Thr	Phe	Asp	Ile	Ala	
50						55					60					
Ile	Lys	Ser	Ser	Thr	Ser	Pro	Phe	Ser	Ser	His	Gly	Val	Arg	Gly	His	
65					70					75					80	
Thr	Leu	Val	Trp	His	Asn	Gln	Thr	Pro	Ser	Trp	Val	Phe	Gln	Asp	Ser	
				85					90					95		
Gln	Gly	His	Phe	Val	Gly	Arg	Asp	Val	Leu	Leu	Glu	Arg	Met	Lys	Ser	
			100					105					110			
His	Ile	Ser	Thr	Val	Val	Gln	Arg	Tyr	Lys	Gly	Lys	Val	Tyr	Cys	Trp	
		115					120					125				
Asp	Val	Ile	Asn	Glu	Ala	Val	Ala	Asp	Glu	Gly	Ser	Glu	Trp	Leu	Arg	
	130					135					140					
Ser	Ser	Thr	Trp	Arg	Gln	Ile	Ile	Gly	Asp	Asp	Phe	Ile	Gln	Gln	Ala	
145					150					155					160	
Phe	Leu	Tyr	Ala	His	Glu	Ala	Asp	Pro	Glu	Ala	Leu	Leu	Phe	Tyr	Asn	
				165					170					175		
Asp	Tyr	Asn	Glu	Cys	Phe	Pro	Glu	Lys	Arg	Glu	Lys	Ile	Tyr	Thr	Leu	
			180					185					190			
Val	Lys	Ser	Leu	Arg	Asp	Lys	Gly	Ile	Pro	Ile	His	Gly	Ile	Gly	Met	
		195					200					205				
Gln	Ala	His	Trp	Ser	Leu	Asn	Arg	Pro	Thr	Leu	Asp	Glu	Ile	Arg	Ala	
	210					215					220					
Ala	Ile	Glu	Arg	Tyr	Ala	Ser	Leu	Gly	Val	Ile	Leu	His	Ile	Thr	Glu	
225					230					235					240	
Leu	Asp	Ile	Ser	Met	Phe	Glu	Phe	Asp	Asp	His	Arg	Lys	Asp	Leu	Ala	
				245					250					255		
Ala	Pro	Thr	Asn	Glu	Met	Val	Glu	Arg	Gln	Ala	Glu	Arg	Tyr	Glu	Gln	
			260					265					270			
Ile	Phe	Ser	Leu	Phe	Lys	Glu	Tyr	Arg	Asp	Val	Ile	Gln	Asn	Val	Thr	
		275					280					285				
Phe	Trp	Gly	Ile	Ala	Asp	Asp	His	Thr	Trp	Leu	Asp	His	Phe	Pro	Val	
	290					295					300					
Gln	Gly	Arg	Lys	Asn	Trp	Pro	Leu	Leu	Phe	Asp	Glu	Gln	His	Asn	Pro	
305					310					315					320	

47

Lys Pro Ala Phe Trp Arg Val Val Asn Ile
 325 330

<210> 11

<211> 190

<212> PRT

<213> *Trichoderma reesei*

<400> 11

Gln	Thr	Ile	Gln	Pro	Gly	Thr	Gly	Tyr	Asn	Asn	Gly	Tyr	Phe	Tyr	Ser
1				5					10					15	
Tyr	Trp	Asn	Asp	Gly	His	Gly	Gly	Val	Thr	Tyr	Thr	Asn	Gly	Pro	Gly
			20					25					30		
Gly	Gln	Phe	Ser	Val	Asn	Trp	Ser	Asn	Ser	Gly	Asn	Phe	Val	Gly	Gly
		35					40					45			
Lys	Gly	Trp	Gln	Pro	Gly	Thr	Lys	Asn	Lys	Val	Ile	Asn	Phe	Ser	Gly
	50					55					60				
Ser	Tyr	Asn	Pro	Asn	Gly	Asn	Ser	Tyr	Leu	Ser	Val	Tyr	Gly	Trp	Ser
65					70					75				80	
Arg	Asn	Pro	Leu	Ile	Glu	Tyr	Tyr	Ile	Val	Glu	Asn	Phe	Gly	Thr	Tyr
				85					90					95	
Asn	Pro	Ser	Thr	Gly	Ala	Thr	Lys	Leu	Gly	Glu	Val	Thr	Ser	Asp	Gly
			100					105						110	
Ser	Val	Tyr	Asp	Ile	Tyr	Arg	Thr	Gln	Arg	Val	Asn	Gln	Pro	Ser	Ile
		115					120					125			
Ile	Gly	Thr	Ala	Thr	Phe	Tyr	Gln	Tyr	Trp	Ser	Val	Arg	Arg	Asn	His
	130					135					140				
Arg	Ser	Ser	Gly	Ser	Val	Asn	Thr	Ala	Asn	His	Phe	Asn	Ala	Trp	Ala
145					150					155					160
Gln	Gln	Gly	Leu	Thr	Leu	Gly	Thr	Met	Asp	Tyr	Gln	Ile	Val	Ala	Val
				165					170					175	
Glu	Gly	Tyr	Phe	Ser	Ser	Gly	Ser	Ala	Ser	Ile	Thr	Val	Ser		
			180					185					190		

<210> 12

<211> 237

<212> PRT

<213> *Aspergillus awamori*

<400> 12

Met	Lys	Ala	Phe	His	Leu	Leu	Ala	Ala	Leu	Ser	Gly	Ala	Ala	Val	Ala
1				5					10					15	
Gln	Gln	Ala	Gln	Leu	Cys	Asp	Gln	Tyr	Ala	Thr	Tyr	Thr	Gly	Gly	Val
			20					25					30		
Tyr	Thr	Ile	Asn	Asn	Asn	Leu	Trp	Gly	Lys	Asp	Ala	Gly	Ser	Gly	Ser
		35					40					45			
Gln	Cys	Thr	Thr	Val	Asn	Ser	Ala	Ser	Ser	Ala	Gly	Thr	Ser	Trp	Ser
	50					55					60				
Thr	Lys	Trp	Asn	Trp	Ser	Gly	Gly	Glu	Asn	Ser	Val	Lys	Ser	Tyr	Ala
65					70					75				80	
Asn	Ser	Gly	Leu	Ser	Phe	Asn	Lys	Lys	Leu	Val	Ser	Gln	Ile	Ser	His
				85					90					95	
Ile	Pro	Thr	Ala	Ala	Arg	Trp	Ser	Tyr	Asp	Asn	Thr	Cys	Ile	Arg	Arg
			100					105					110		
Gly	Arg	Ala	Tyr	Asp	Leu	Phe	Thr	Ala	Ala	Asp	Ile	Asn	His	Val	Thr
		115					120					125			
Trp	Ser	Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Ala	Arg	Tyr	Gly	Gly
	130					135					140				
Val	Gln	Pro	Leu	Gly	Ser	Gln	Ile	Ala	Thr	Ala	Thr	Val	Glu	Gly	Gln
145					150					155					160

48

Thr	Trp	Glu	Leu	Trp	Tyr	Gly	Val	Asn	Gly	Ala	Gln	Lys	Thr	Tyr	Ser
				165					170					175	
Phe	Val	Ala	Ala	Asn	Pro	Ile	Thr	Ser	Phe	Gln	Gly	Asp	Ile	Asn	Asp
			180					185					190		
Phe	Phe	Lys	Tyr	Leu	Thr	Gln	Asn	His	Gly	Phe	Pro	Ala	Ser	Ser	Gln
		195					200					205			
Tyr	Leu	Ile	Thr	Leu	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Gly	Pro
	210					215					220				
Ala	Thr	Leu	Asn	Val	Ala	Asp	Trp	Ser	Ala	Ser	Val	Gln			
225					230					235					

<210> 13

<211> 233

<212> PRT

<213> Trichoderma viride

<400> 13

Met	Lys	Phe	Leu	Gln	Ile	Ala	Pro	Thr	Leu	Leu	Pro	Val	Ala	Leu	Ala
1				5					10					15	
Gln	Ser	Ser	Cys	Ser	Gln	Tyr	Ala	Thr	Phe	Ser	Gly	Gly	Asn	Tyr	Ala
			20					25					30		
Leu	Ser	Asn	Asn	Leu	Trp	Gly	Gln	Ser	Ala	Gly	Ser	Gly	Ser	Gly	Cys
		35					40					45			
Ile	Thr	Asp	Val	Ser	Leu	Gly	Gly	Ser	Ala	Val	Trp	Ser	Thr	Thr	Trp
	50					55					60				
Asp	Trp	Ser	Gly	Gly	Gln	Ser	Asn	Val	Lys	Gly	Tyr	Pro	Asn	Ile	Ala
65					70					75					80
Leu	Asn	Ile	Pro	Asn	Lys	Arg	Leu	Val	Ser	Ser	Ile	Ser	Ser	Met	Pro
				85					90					95	
Thr	Thr	Ala	Gln	Trp	Ser	Tyr	Ser	Gly	Ser	Ser	Ile	Arg	Ala	Asp	Val
			100					105					110		
Ala	Tyr	Asp	Leu	Phe	Thr	Ala	Ser	Asn	Pro	Asn	His	Val	Thr	Tyr	Ser
		115					120					125			
Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Gly	Lys	Tyr	Gly	Asp	Ile	Gln
	130					135					140				
Pro	Ile	Gly	Ser	Ser	Gln	Gly	Thr	Val	Asn	Val	Gly	Gly	Thr	Ser	Trp
145					150					155					160
Asn	Leu	Trp	Tyr	Gly	Pro	Asn	Gly	Ser	Met	Gln	Val	Tyr	Ser	Phe	Val
				165					170					175	
Ala	Pro	Gly	Asn	Leu	Thr	Asn	Trp	Ser	Gly	Asp	Val	Lys	Asn	Phe	Tyr
			180					185					190		
Thr	Tyr	Leu	Gln	Asn	Asn	Lys	Gly	Tyr	Pro	Ala	Ser	Ser	Gln	Tyr	Val
		195					200					205			
Leu	Ser	Tyr	Gln	Phe	Gly	Thr	Glu	Ala	Phe	Thr	Gly	Ser	Gly	Thr	Leu
	210					215					220				
Asn	Asn	Thr	Trp	Thr	Ala	Ser	Ile	Asn							
225					230										

<210> 14

<211> 234

<212> PRT

<213> Hypocrea koningii

<400> 14

Met	Lys	Leu	Ile	His	Val	Leu	Pro	Ala	Leu	Ile	Pro	Ala	Ala	Leu	Ala
1				5					10					15	
Gln	Thr	Ser	Cys	Asp	Gln	Tyr	Ala	Val	Phe	Thr	Gly	Ser	Asp	Tyr	Thr
			20					25					30		
Val	Ser	Asn	Asn	Leu	Trp	Gly	Gln	Ser	Ala	Gly	Ser	Gly	Phe	Gly	Cys
		35					40					45			

49

Val	Thr	Ala	Glu	Ser	Leu	Ser	Gly	Ser	Ala	Ser	Trp	His	Ala	Asp	Trp
50						55					60				
Gln	Trp	Ser	Gly	Gly	Gln	Asn	Asn	Val	Lys	Ser	Tyr	Gln	Asn	Ser	Gln
65					70				75						80
Ile	Pro	Ile	Pro	Gln	Lys	Arg	Thr	Val	Asn	Ser	Ile	Ser	Ser	Met	Pro
				85					90					95	
Thr	Thr	Ala	Ser	Trp	Ser	Tyr	Thr	Gly	Ser	Asp	Ile	Arg	Ala	Asn	Val
			100					105					110		
Ala	Tyr	Asp	Leu	Phe	Thr	Ala	Ala	Asn	Pro	Asn	His	Val	Thr	Tyr	Ser
		115					120					125			
Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Gly	Arg	Tyr	Gly	Asp	Ile	Gly
130						135					140				
Pro	Ile	Gly	Ser	Ser	Gln	Gly	Thr	Val	Asn	Val	Gly	Gly	Gln	Ser	Trp
145					150					155					160
Thr	Leu	Tyr	Tyr	Gly	Tyr	Asn	Gly	Ala	Met	Gln	Val	Tyr	Ser	Phe	Val
				165					170					175	
Ala	Gln	Thr	Asn	Thr	Thr	Ser	Tyr	Ser	Gly	Asp	Val	Lys	Asn	Phe	Phe
			180					185					190		
Asn	Tyr	Leu	Arg	Asp	Asn	Lys	Gly	Tyr	Asn	Ala	Ala	Gly	Gln	Tyr	Val
		195					200					205			
Leu	Ser	Tyr	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Ser	Gly	Thr	Leu
210						215					220				
Asn	Val	Ala	Ser	Trp	Thr	Ala	Ser	Ile	Asn						
225					230										

<210> 15

<211> 234

<212> PRT

<213> Hypocrea schweinitzii

<400> 15

Met	Lys	Phe	Leu	Gln	Val	Leu	Pro	Ala	Ile	Leu	Pro	Ala	Ala	Leu	Ala
1				5					10					15	
Gln	Thr	Ser	Cys	Asp	Gln	Tyr	Ala	Thr	Phe	Ser	Gly	Asn	Gly	Tyr	Ile
			20					25					30		
Val	Ser	Asn	Asn	Leu	Trp	Gly	Ala	Ser	Ala	Gly	Ser	Gly	Phe	Gly	Cys
		35				40						45			
Val	Thr	Ser	Val	Ser	Leu	Asn	Gly	Ala	Ala	Ser	Trp	His	Ala	Asp	Trp
50						55					60				
Gln	Trp	Ser	Gly	Gly	Gln	Asn	Asn	Val	Lys	Ser	Tyr	Gln	Asn	Val	Gln
65					70					75					80
Ile	Asn	Ile	Pro	Gln	Lys	Arg	Thr	Val	Asn	Ser	Ile	Gly	Ser	Met	Pro
				85					90					95	
Thr	Thr	Ala	Ser	Trp	Ser	Tyr	Ser	Gly	Ser	Asp	Ile	Arg	Ala	Asn	Val
			100					105					110		
Ala	Tyr	Asp	Leu	Phe	Thr	Ala	Ala	Asn	Pro	Asn	His	Val	Thr	Tyr	Ser
		115					120					125			
Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Gly	Lys	Tyr	Gly	Asp	Ile	Gly
130						135					140				
Pro	Ile	Gly	Ser	Ser	Gln	Gly	Thr	Val	Asn	Val	Gly	Gly	Gln	Thr	Trp
145					150					155					160
Thr	Leu	Tyr	Tyr	Gly	Tyr	Asn	Gly	Ala	Met	Gln	Val	Tyr	Ser	Phe	Val
				165					170					175	
Ala	Gln	Ser	Asn	Thr	Thr	Ser	Tyr	Ser	Gly	Asp	Val	Lys	Asn	Phe	Phe
			180					185					190		
Asn	Tyr	Leu	Arg	Asp	Asn	Lys	Gly	Tyr	Asn	Ala	Gly	Gly	Gln	Tyr	Val
		195					200					205			
Leu	Ser	Tyr	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Ser	Gly	Thr	Leu
210						215					220				
Asn	Val	Ala	Ser	Trp	Thr	Ala	Ser	Ile	Asn						

50

225

230

<210> 16

<211> 237

<212> PRT

<213> *Stachybotrys echinata*

<400> 16

Met	Lys	Val	Ala	Ala	Leu	Leu	Val	Ala	Leu	Ser	Pro	Leu	Ala	Phe	Ala		
1				5					10					15			
Gln	Ser	Leu	Cys	Asp	Gln	Tyr	Ser	Tyr	Tyr	Ser	Ser	Asn	Gly	Tyr	Glu		
			20					25					30				
Phe	Asn	Asn	Asn	Met	Trp	Gly	Arg	Asn	Ser	Gly	Gln	Gly	Asn	Gln	Cys		
		35				40						45					
Thr	Tyr	Val	Asp	Tyr	Ser	Ser	Pro	Asn	Gly	Val	Gly	Trp	Arg	Val	Asn		
	50				55						60						
Trp	Asn	Trp	Ser	Gly	Gly	Asp	Asn	Asn	Val	Lys	Ser	Tyr	Pro	Tyr	Ser		
65				70						75					80		
Gly	Arg	Gln	Leu	Pro	Thr	Lys	Arg	Ile	Val	Ser	Trp	Ile	Gly	Ser	Leu		
			85						90					95			
Pro	Thr	Thr	Val	Ser	Trp	Asn	Tyr	Gln	Gly	Asn	Asn	Leu	Arg	Ala	Asn		
			100					105					110				
Val	Ala	Tyr	Asp	Leu	Phe	Thr	Ala	Ala	Asn	Pro	Asn	His	Pro	Asn	Ser		
		115					120					125					
Ser	Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Gly	Arg	Leu	Gly	Asn	Val		
	130					135					140						
Tyr	Pro	Ile	Gly	Asn	Gln	Val	Ala	Thr	Val	Asn	Ile	Ala	Gly	Gln	Gln		
145					150					155					160		
Trp	Asn	Leu	Tyr	Tyr	Gly	Tyr	Asn	Gly	Ala	Met	Gln	Val	Tyr	Ser	Phe		
			165					170						175			
Val	Ser	Pro	Asn	Gln	Leu	Asn	Tyr	Phe	Ser	Gly	Asn	Val	Lys	Asp	Phe		
			180					185					190				
Phe	Thr	Tyr	Leu	Gln	Tyr	Asn	Arg	Ala	Tyr	Pro	Ala	Asp	Ser	Gln	Tyr		
		195				200					205						
Leu	Ile	Thr	Tyr	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Gln	Asn	Ala		
	210					215					220						
Val	Phe	Thr	Val	Ser	Asn	Trp	Ser	Ala	Gln	Gln	Asn	Asn					
225					230						235						

<210> 17

<211> 238

<212> PRT

<213> *Fusarium equiseti*

<400> 17

Met	Lys	Ser	Thr	Leu	Leu	Leu	Ala	Gly	Ala	Phe	Ala	Pro	Leu	Ala	Phe		
1				5					10					15			
Ala	Lys	Asp	Leu	Cys	Glu	Gln	Tyr	Gly	Tyr	Leu	Ser	Ser	Asp	Gly	Tyr		
			20					25					30				
Ser	Leu	Asn	Asn	Asn	Val	Trp	Gly	Lys	Asp	Ser	Gly	Thr	Gly	Asp	Gln		
		35				40						45					
Cys	Thr	His	Val	Asn	Trp	Asn	Asn	Ala	Asn	Gly	Ala	Gly	Trp	Asp	Val		
	50				55						60						
Glu	Trp	Asn	Trp	Ser	Gly	Gly	Lys	Asp	Asn	Val	Lys	Ser	Tyr	Pro	Asn		
65				70						75					80		
Ser	Ala	Leu	Leu	Ile	Gly	Glu	Asp	Lys	Lys	Thr	Ile	Ser	Ser	Ile	Thr		
			85					90						95			
Asn	Met	Gln	Ser	Thr	Ala	Glu	Trp	Lys	Tyr	Ser	Gly	Asp	Asn	Leu	Arg		
		100						105					110				
Ala	Asp	Val	Ala	Tyr	Asp	Leu	Phe	Thr	Ala	Ala	Asp	Pro	Asn	His	Glu		

51

		115					120					125					
Thr	Ser	Ser	Gly	Glu	Tyr	Glu	Leu	Met	Val	Trp	Leu	Ala	Arg	Ile	Gly		
	130					135					140						
Gly	Val	Gln	Pro	Ile	Gly	Ser	Leu	Gln	Thr	Ser	Val	Thr	Ile	Glu	Gly		
145					150					155					160		
His	Thr	Trp	Glu	Leu	Trp	Val	Gly	Met	Asn	Gly	Ser	Met	Lys	Val	Phe		
				165					170					175			
Ser	Phe	Val	Ala	Pro	Thr	Pro	Val	Asn	Asn	Phe	Asn	Ala	Asp	Ile	Lys		
			180					185					190				
Gln	Phe	Trp	Asp	Tyr	Leu	Thr	Lys	Ser	Gln	Asn	Phe	Pro	Ala	Asp	Asn		
	195						200					205					
Gln	Tyr	Leu	Leu	Thr	Phe	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Asp		
	210					215					220						
Asn	Ala	Lys	Phe	Thr	Val	Thr	Asn	Phe	Asn	Ala	His	Leu	Lys				
225					230					235							

<210> 18

<211> 237

<212> PRT

<213> Bionectria ochroleuca

<400> 18

Met	Lys	Thr	Gly	Ile	Ala	Tyr	Leu	Ala	Ala	Val	Leu	Pro	Leu	Ala	Met		
1				5					10					15			
Ala	Glu	Ser	Leu	Cys	Asp	Gln	Tyr	Ala	Tyr	Leu	Ser	Arg	Asp	Gly	Tyr		
			20					25					30				
Asn	Phe	Asn	Asn	Asn	Glu	Trp	Gly	Ala	Ala	Thr	Gly	Thr	Gly	Asp	Gln		
		35					40					45					
Cys	Thr	Tyr	Val	Asp	Ser	Thr	Ser	Ser	Gly	Gly	Val	Ser	Trp	His	Ser		
	50					55					60						
Asp	Trp	Thr	Asn	Ser	Gly	Ser	Glu	Ser	Glu	Ile	Lys	Ser	Tyr	Pro	Tyr		
65					70					75					80		
Ser	Gly	Leu	Asp	Leu	Pro	Glu	Lys	Lys	Ile	Val	Thr	Ser	Ile	Gly	Ser		
				85					90					95			
Ile	Ser	Thr	Gly	Ala	Glu	Trp	Ser	Tyr	Ser	Gly	Ser	Asn	Ile	Arg	Ala		
			100					105					110				
Asp	Val	Ala	Tyr	Asp	Ile	Phe	Thr	Ala	Ala	Asp	Pro	Asn	His	Ala	Thr		
		115					120					125					
Ser	Ser	Gly	Asp	Tyr	Glu	Val	Met	Ile	Trp	Leu	Ala	Asn	Leu	Gly	Gly		
	130					135					140						
Leu	Thr	Pro	Ile	Gly	Ser	Pro	Ile	Gly	Thr	Val	Lys	Ala	Ala	Gly	Arg		
145					150						155				160		
Asp	Trp	Glu	Leu	Trp	Asp	Gly	Tyr	Asn	Gly	Ala	Met	Arg	Val	Tyr	Ser		
				165					170					175			
Phe	Val	Ala	Pro	Ser	Gln	Leu	Asn	Ser	Phe	Asp	Gly	Glu	Ile	Met	Asp		
			180					185					190				
Phe	Phe	Tyr	Val	Val	Lys	Asp	Met	Arg	Gly	Phe	Pro	Ala	Asp	Ser	Gln		
	195						200					205					
His	Leu	Leu	Thr	Val	Gln	Phe	Gly	Thr	Glu	Pro	Ile	Ser	Gly	Ser	Gly		
	210					215					220						
Ala	Lys	Phe	Ser	Val	Ser	His	Trp	Ser	Ala	Lys	Leu	Gly					
225					230					235							

<210> 19

<211> 236

<212> PRT

<213> Bionectria ochroleuca

<400> 19

Met Lys Phe Gln Leu Leu Ser Leu Thr Ala Phe Ala Pro Leu Ser Leu

52

1				5					10					15		
Ala	Ala	Leu	Cys	Gly	Gln	Tyr	Gln	Ser	Gln	Ser	Gln	Gly	Gly	Tyr	Ile	
			20					25					30			
Phe	Asn	Asn	Asn	Lys	Trp	Gly	Gln	Gly	Ser	Gly	Ser	Gly	Ser	Gln	Cys	
		35				40						45				
Leu	Thr	Ile	Asp	Lys	Thr	Trp	Asp	Ser	Asn	Val	Ala	Phe	His	Ala	Asp	
	50					55					60					
Trp	Ser	Trp	Ser	Gly	Gly	Thr	Asn	Asn	Val	Lys	Ser	Tyr	Pro	Asn	Ala	
65				70					75					80		
Gly	Leu	Glu	Phe	Ser	Arg	Gly	Lys	Lys	Val	Ser	Ser	Ile	Gly	Thr	Ile	
				85					90				95			
Asn	Gly	Gly	Ala	Asp	Trp	Asp	Tyr	Ser	Gly	Ser	Asn	Ile	Arg	Ala	Asn	
			100					105					110			
Val	Ala	Tyr	Asp	Ile	Phe	Thr	Ser	Ala	Asp	Pro	Asn	His	Val	Thr	Ser	
		115					120					125				
Ser	Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Gly	Lys	Leu	Gly	Asp	Ile	
	130					135					140					
Tyr	Pro	Ile	Gly	Asn	Ser	Ile	Gly	Arg	Val	Lys	Ala	Ala	Asn	Arg	Glu	
145				150					155					160		
Trp	Asp	Leu	His	Val	Gly	Tyr	Asn	Gly	Ala	Met	Lys	Val	Phe	Ser	Phe	
				165					170					175		
Val	Ala	Pro	Ser	Pro	Val	Thr	Arg	Phe	Asp	Gly	Asn	Ile	Met	Asp	Phe	
			180					185					190			
Phe	Tyr	Val	Met	Arg	Asp	Met	Gln	Gly	Tyr	Pro	Met	Asp	Lys	Gln	Tyr	
		195					200					205				
Leu	Leu	Thr	Leu	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Ser	Asn	Ala	
	210					215					220					
Lys	Phe	Ser	Cys	Trp	Tyr	Phe	Gly	Ala	Lys	Ile	Lys					
225					230					235						

<210> 20

<211> 240

<212> PRT

<213> Bionectria ochroleuca

<400> 20

Met	Lys	Ala	Asn	Ile	Val	Ile	Leu	Ser	Leu	Phe	Ala	Pro	Leu	Ala	Ala
1				5				10					15		
Val	Ala	Gln	Thr	Leu	Cys	Gly	Gln	Tyr	Ser	Ser	Asn	Thr	Gln	Gly	Gly
			20					25					30		
Tyr	Ile	Phe	Asn	Asn	Asn	Met	Trp	Gly	Met	Gly	Ser	Gly	Ser	Gly	Ser
		35				40					45				
Gln	Cys	Thr	Tyr	Val	Asp	Lys	Val	Trp	Ala	Glu	Gly	Val	Ala	Trp	His
	50				55					60					
Thr	Asp	Trp	Ser	Trp	Ser	Gly	Gly	Asp	Asn	Asn	Val	Lys	Ser	Tyr	Pro
65				70					75					80	
Tyr	Ser	Gly	Arg	Glu	Leu	Gly	Thr	Lys	Arg	Ile	Val	Ser	Ser	Ile	Lys
			85					90						95	
Ser	Ile	Ser	Ser	Gly	Ala	Asp	Trp	Asp	Tyr	Thr	Gly	Ser	Asn	Leu	Arg
			100				105						110		
Ala	Asn	Ala	Ala	Tyr	Asp	Ile	Phe	Thr	Ser	Ala	Asn	Pro	Asn	His	Ala
	115					120						125			
Thr	Ser	Ser	Gly	Asp	Tyr	Glu	Val	Met	Ile	Trp	Leu	Gly	Arg	Tyr	Gly
	130					135					140				
Gly	Val	Tyr	Pro	Ile	Gly	Asn	Ser	Ile	Gly	Thr	Val	Arg	Ala	Ala	Gly
145				150					155					160	
Arg	Asp	Trp	Ala	Leu	His	Ile	Gly	Tyr	Asn	Gly	Ala	Met	Lys	Val	Phe
			165					170						175	
Ser	Phe	Val	Ala	Ala	Asn	Pro	Val	Thr	Arg	Phe	Asp	Gly	Glu	Ile	Met
			180					185					190		

53

Asp	Phe	Phe	Tyr	Leu	Leu	Arg	Asp	Met	Gln	Gly	Tyr	Pro	Met	Thr	Ser
		195					200					205			
Gln	Tyr	Leu	Leu	Thr	Leu	Gln	Phe	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Ser
	210					215					220				
Gly	Ala	Lys	Phe	Asn	Cys	Trp	Tyr	Phe	Gly	Ala	Thr	Leu	Ser	Tyr	Trp
225					230					235					240

<210> 21
 <211> 254
 <212> PRT
 <213> Humicola grisea

<400> 21

Met	Leu	Lys	Ser	Ala	Leu	Leu	Leu	Gly	Ala	Ala	Ala	Val	Ser	Val	Gln
1				5					10					15	
Ser	Ala	Ser	Ile	Pro	Thr	Ile	Pro	Ala	Asn	Leu	Glu	Pro	Arg	Gln	Ile
			20				25						30		
Arg	Ser	Leu	Cys	Glu	Leu	Tyr	Gly	Tyr	Trp	Ser	Gly	Asn	Gly	Tyr	Glu
		35					40					45			
Leu	Leu	Asn	Asn	Leu	Trp	Gly	Lys	Asp	Thr	Ala	Thr	Ser	Gly	Trp	Gln
	50					55					60				
Cys	Thr	Tyr	Leu	Asp	Gly	Thr	Asn	Asn	Gly	Gly	Ile	Gln	Trp	Asn	Thr
65					70					75					80
Ala	Trp	Glu	Trp	Gln	Gly	Ala	Pro	Asp	Asn	Val	Lys	Asn	Tyr	Pro	Tyr
				85				90						95	
Val	Gly	Lys	Gln	Ile	Gln	Arg	Gly	Arg	Lys	Ile	Ser	Asp	Ile	Asn	Ser
			100				105						110		
Met	Arg	Thr	Ser	Val	Ser	Trp	Thr	Tyr	Asp	Arg	Thr	Asp	Leu	Arg	Ala
		115					120					125			
Asn	Val	Ala	Tyr	Asp	Val	Phe	Thr	Ala	Arg	Asp	Pro	Asp	His	Pro	Asn
	130					135					140				
Trp	Gly	Gly	Asp	Tyr	Glu	Leu	Met	Ile	Trp	Leu	Ala	Arg	Tyr	Gly	Gly
145					150					155					160
Ile	Tyr	Pro	Ile	Gly	Thr	Phe	His	Ser	Gln	Val	Asn	Leu	Ala	Gly	Arg
				165					170					175	
Thr	Trp	Asp	Leu	Trp	Thr	Gly	Tyr	Asn	Gly	Asn	Met	Arg	Val	Tyr	Ser
			180				185						190		
Phe	Leu	Pro	Pro	Ser	Gly	Asp	Ile	Arg	Asp	Phe	Ser	Cys	Asp	Ile	Lys
		195				200						205			
Asp	Phe	Phe	Asn	Tyr	Leu	Glu	Arg	Asn	His	Gly	Tyr	Pro	Ala	Arg	Glu
	210					215					220				
Gln	Asn	Leu	Ile	Val	Tyr	Gln	Val	Gly	Thr	Glu	Cys	Phe	Thr	Gly	Gly
225					230					235					240
Pro	Ala	Arg	Phe	Thr	Cys	Arg	Asp	Phe	Arg	Ala	Asp	Leu	Trp		
				245					250						

<210> 22
 <211> 247
 <212> PRT
 <213> Chaetomium brasiliense

<400> 22

Met	Lys	Leu	Thr	Leu	Val	Leu	Phe	Val	Ser	Ser	Leu	Ala	Ala	Ala	Thr
1				5					10					15	
Pro	Leu	Gly	Trp	Arg	Glu	Arg	Arg	Gln	Gln	Val	Ser	Leu	Cys	Gly	Gln
			20					25					30		
Ser	Ser	Ser	Trp	Ser	Gly	Asn	Gly	Tyr	Gln	Leu	Asn	Asn	Asn	Leu	Trp
		35					40					45			
Gly	Gln	Ser	Arg	Ala	Thr	Ser	Gly	Ser	Gln	Cys	Thr	Tyr	Leu	Asp	Ser
	50					55					60				

54

```

Ser Ser Asn Ser Gly Ile His Trp His Thr Thr Trp Thr Trp Glu Gly
65      70      75      80
Gly Glu Gly Glu Val Lys Ser Tyr Ala Tyr Ser Gly Arg Gln Val Ser
      85      90      95
Thr Gly Leu Thr Ile Ala Ser Ile Asp Ser Met Gln Thr Ser Val Ser
      100      105      110
Trp Glu Tyr Asn Thr Thr Asp Ile Gln Ala Asn Val Ala Tyr Asp Ile
      115      120      125
Phe Thr Ala Glu Asp Pro Asp His Glu His Ser Ser Gly Asp Tyr Glu
      130      135      140
Val Met Ile Trp Leu Ala Arg Tyr Asn Asn Val Ser Pro Ile Gly Ser
145      150      155      160
Ser Val Ala Thr Ala Thr Val Gly Gly Asp Thr Trp Asp Leu Phe Ala
      165      170      175
Gly Ala Asn Gly Asp Met Glu Val Tyr Ser Phe Val Ala Glu Asn Thr
      180      185      190
Met Asn Ser Phe Ser Gly Asp Val Lys Asp Phe Phe Asp Tyr Leu Glu
      195      200      205
Gln Asn Val Gly Phe Pro Val Asp Asp Gln Tyr Leu Leu Val Phe Glu
      210      215      220
Leu Gly Ser Glu Ala Phe Thr Gly Gly Pro Ala Thr Leu Ser Val Ser
225      230      235      240
Gln Phe Ser Ala Asn Ile Ala
      245

```

<210> 23

<211> 357

<212> PRT

<213> Bionectria ochroleuca

<400> 23

```

Met Lys Ser Ile Ile Ser Phe Phe Gly Leu Ala Thr Leu Val Ala Ala
1      5      10      15
Ala Pro Ser Gln Asn Pro Thr Arg Thr Gln Pro Leu Glu Lys Arg Ala
      20      25      30
Thr Thr Leu Cys Gly Gln Trp Asp Ser Val Glu Thr Gly Gly Tyr Thr
      35      40      45
Ile Tyr Asn Asn Leu Trp Gly Gln Asp Asn Gly Ser Gly Ser Gln Cys
      50      55      60
Leu Thr Val Glu Gly Val Thr Asp Gly Leu Ala Ala Trp Ser Ser Thr
65      70      75      80
Trp Ser Trp Ser Gly Gly Ser Ser Ser Val Lys Ser Tyr Ser Asn Ala
      85      90      95
Val Leu Ser Ala Glu Ala Ala Arg Ile Ser Ala Ile Ser Ser Ile Pro
      100      105      110
Ser Lys Trp Glu Trp Ser Tyr Thr Gly Thr Asp Ile Val Ala Asn Val
      115      120      125
Ala Tyr Asp Leu Phe Ser Asn Thr Asp Cys Gly Asp Thr Pro Glu Tyr
      130      135      140
Glu Ile Met Ile Trp Leu Ser Ala Leu Gly Gly Ala Gly Pro Ile Ser
145      150      155      160
Ser Thr Gly Ser Ser Ile Ala Thr Val Thr Ile Ala Gly Ala Ser Trp
      165      170      175
Asn Leu Trp Gln Gly Gln Asn Asn Gln Met Thr Val Phe Ser Phe Val
      180      185      190
Ala Glu Ser Asp Gln Lys Ser Phe Ser Gly Asp Leu Asn Asp Phe Ile
      195      200      205
Gln Tyr Leu Val Asp Ser Gln Gly Tyr Ser Gly Ser Gln Cys Leu Tyr
      210      215      220
Ser Ile Gly Ala Gly Thr Glu Pro Phe Thr Gly Thr Asp Ala Glu Phe

```

55

225					230					235				240
Ile	Thr	Thr	Gly	Tyr	Ser	Val	Ser	Val	Ser	Ala	Gly	Asp	Ser	Gly
				245					250					255
Asp	Glu	Thr	Thr	Thr	Ser	Ser	Gln	Ala	Gln	Ser	Ser	Thr	Val	Glu
			260					265					270	Thr
Ser	Thr	Ala	Thr	Gln	Pro	Gln	Ser	Ser	Ser	Thr	Val	Val	Pro	Thr
		275					280					285		Val
Thr	Leu	Ser	Gln	Pro	Ser	Asn	Glu	Ser	Thr	Thr	Thr	Pro	Val	Gln
	290					295					300			Ser
Gln	Pro	Ser	Ser	Val	Glu	Thr	Thr	Pro	Thr	Ala	Gln	Pro	Gln	Ser
305					310					315				320
Ser	Val	Gln	Thr	Thr	Thr	Thr	Ala	Gln	Ala	Gln	Pro	Thr	Pro	Glu
			325				330						335	Arg
Ala	Ala	Pro	Asp	Ala	Gly	Ser	Ala	Glu	Leu	Leu	Ser	Ser	Ala	Thr
			340				345						350	Met
His	Leu	Asp	Arg	Arg										
		355												

<210> 24

<211> 247

<212> PRT

<213> Emericella desertorum

<400> 24

Met	Lys	Leu	Leu	Ala	Leu	Ser	Leu	Val	Ser	Leu	Ala	Ser	Ala	Ala	Ser
1				5					10					15	
Ala	Ala	Ser	Ile	Leu	Ser	Asn	Thr	Phe	Thr	Arg	Arg	Ser	Asp	Phe	Cys
			20					25					30		
Gly	Gln	Trp	Asp	Thr	Ala	Thr	Val	Gly	Asn	Phe	Ile	Val	Tyr	Asn	Asn
		35					40					45			
Leu	Trp	Gly	Gln	Asp	Asn	Ala	Asp	Ser	Gly	Ser	Gln	Cys	Thr	Gly	Val
	50					55					60				
Asp	Ser	Ala	Asn	Gly	Asn	Ser	Ile	Ser	Trp	His	Thr	Thr	Trp	Ser	Trp
65					70				75						80
Ser	Gly	Gly	Ser	Ser	Ser	Val	Lys	Ser	Tyr	Ala	Asn	Ala	Ala	Tyr	Gln
			85					90						95	
Phe	Thr	Ser	Thr	Lys	Leu	Asn	Ser	Leu	Ser	Ser	Ile	Pro	Thr	Ser	Trp
			100					105					110		
Lys	Trp	Gln	Tyr	Ser	Thr	Thr	Asp	Ile	Val	Ala	Asn	Val	Ala	Tyr	Asp
		115					120					125			
Leu	Phe	Thr	Ser	Ser	Ser	Ala	Gly	Gly	Asp	Ser	Glu	Tyr	Glu	Ile	Met
	130					135					140				
Ile	Trp	Leu	Ala	Ala	Leu	Gly	Gly	Ala	Gly	Pro	Ile	Ser	Ser	Thr	Gly
145					150					155					160
Ser	Ser	Ile	Ala	Thr	Val	Thr	Leu	Gly	Gly	Val	Thr	Trp	Ser	Leu	Tyr
			165					170						175	
Ser	Gly	Pro	Asn	Gly	Ser	Met	Gln	Val	Tyr	Ser	Phe	Val	Ala	Ser	Ser
			180					185					190		
Thr	Thr	Glu	Ser	Phe	Ser	Ala	Asp	Leu	Met	Asp	Phe	Ile	Asn	Tyr	Leu
		195					200					205			
Ala	Glu	Asn	Gln	Gly	Leu	Ser	Ser	Ser	Gln	Tyr	Leu	Thr	His	Val	Gln
	210					215					220				
Ala	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Thr	Asp	Ala	Thr	Leu	Thr	Val	Ser
225					230					235					240
Ser	Tyr	Ser	Val	Ser	Val	Ser									
				245											

<210> 25

<211> 244

<212> PRT

56

<213> Fusarium solani

<400> 25

Met	Lys	Ser	Ala	Ile	Val	Ala	Ala	Leu	Ala	Gly	Leu	Ala	Ala	Ala	Ser	1	5	10	15
Pro	Thr	Arg	Leu	Ile	Pro	Arg	Gly	Gln	Phe	Cys	Gly	Gln	Trp	Asp	Ser	20	25	30	
Glu	Thr	Ala	Gly	Ala	Tyr	Thr	Ile	Tyr	Asn	Asn	Leu	Trp	Gly	Lys	Asp	35	40	45	
Asn	Ala	Glu	Ser	Gly	Glu	Gln	Cys	Thr	Thr	Asn	Ser	Gly	Glu	Gln	Ser	50	55	60	
Asp	Gly	Ser	Ile	Ala	Trp	Ser	Val	Glu	Trp	Ser	Trp	Thr	Gly	Gly	Gln	65	70	75	80
Gly	Gln	Val	Lys	Ser	Tyr	Pro	Asn	Ala	Val	Val	Glu	Ile	Glu	Lys	Lys	85	90	95	
Thr	Leu	Gly	Glu	Val	Ser	Ser	Ile	Pro	Ser	Ala	Trp	Asp	Trp	Thr	Tyr	100	105	110	
Thr	Gly	Asn	Gly	Ile	Ile	Ala	Asn	Val	Ala	Tyr	Asp	Leu	Phe	Thr	Ser	115	120	125	
Ser	Thr	Glu	Ser	Gly	Asp	Ala	Glu	Tyr	Glu	Phe	Met	Ile	Trp	Leu	Ser	130	135	140	
Ala	Leu	Gly	Gly	Ala	Gly	Pro	Ile	Ser	Asn	Asp	Gly	Ser	Pro	Val	Ala	145	150	155	160
Thr	Val	Glu	Leu	Ala	Gly	Thr	Ser	Trp	Lys	Leu	Tyr	Gln	Gly	Lys	Asn	165	170	175	
Asn	Gln	Met	Thr	Val	Phe	Ser	Phe	Val	Ala	Glu	Ser	Asp	Val	Asn	Asn	180	185	190	
Phe	Cys	Gly	Asp	Leu	Ala	Asp	Phe	Thr	Asp	Tyr	Leu	Val	Asp	Asn	His	195	200	205	
Gly	Val	Ser	Ser	Ser	Gln	Ile	Leu	Gln	Ser	Val	Gly	Ala	Gly	Thr	Glu	210	215	220	
Pro	Phe	Glu	Gly	Thr	Asn	Ala	Val	Phe	Thr	Thr	Asn	Asn	Tyr	His	Ala	225	230	235	240
Asp	Val	Glu	Tyr																

<210> 26

<211> 250

<212> PRT

<213> Fusarium solani

<400> 26

Met	Lys	Phe	Phe	Gly	Val	Val	Ser	Ala	Phe	Leu	Ala	Ala	Thr	Ala	Val	1	5	10	15
Ala	Thr	Pro	Thr	Thr	Pro	Thr	Glu	Thr	Ile	Glu	Lys	Arg	Asp	Thr	Thr	20	25	30	
Trp	Cys	Asp	Ala	Phe	Gly	Ser	Leu	Ala	Thr	Ser	Gly	Tyr	Thr	Val	Tyr	35	40	45	
His	Asn	Asn	Trp	Gly	Lys	Gly	Asp	Ala	Thr	Ser	Gly	Ser	Gln	Cys	Thr	50	55	60	
Thr	Phe	Thr	Ser	Val	Ser	Asn	Asn	Asn	Phe	Val	Trp	Ser	Thr	Ser	Trp	65	70	75	80
Thr	Trp	Ala	Gly	Gly	Ala	Gly	Lys	Val	Lys	Ser	Tyr	Ser	Asn	Val	Ala	85	90	95	
Leu	Glu	Lys	Ile	Asn	Lys	Lys	Ile	Ser	Asp	Ile	Lys	Ser	Val	Ser	Thr	100	105	110	
Arg	Trp	Ile	Trp	Arg	Tyr	Thr	Gly	Thr	Lys	Met	Ile	Ala	Asn	Val	Ser	115	120	125	
Tyr	Asp	Leu	Trp	Phe	Ala	Pro	Thr	Ala	Ser	Ser	Asn	Asn	Ala	Tyr	Glu	130	135	140	

57

Ile	Met	Ile	Trp	Val	Gly	Ala	Tyr	Gly	Gly	Ala	Leu	Pro	Ile	Ser	Thr
145					150					155					160
Pro	Gly	Lys	Gly	Val	Ile	Asp	Arg	Pro	Thr	Leu	Ala	Gly	Ile	Pro	Trp
				165					170						175
Asp	Val	Tyr	Lys	Gly	Pro	Asn	Gly	Asp	Val	Thr	Val	Ile	Ser	Phe	Val
			180					185					190		
Ala	Ser	Ser	Asn	Gln	Gly	Asn	Phe	Gln	Ala	Asp	Leu	Lys	Glu	Phe	Leu
		195				200					205				
Asn	Tyr	Leu	Thr	Ser	Lys	Gln	Gly	Leu	Pro	Ser	Asn	Tyr	Val	Ala	Thr
	210					215					220				
Ser	Phe	Gln	Ala	Gly	Thr	Glu	Pro	Phe	Glu	Gly	Thr	Asn	Ala	Val	Leu
225					230					235					240
Lys	Thr	Ser	Ala	Tyr	Thr	Ile	Ser	Val	Asn						
				245					250						

<210> 27

<211> 371

<212> PRT

<213> Streptomyces sp. 11AG8

<400> 27

Met	Arg	Ser	His	Pro	Arg	Ser	Ala	Thr	Met	Thr	Val	Leu	Val	Val	Leu
1				5					10					15	
Ala	Ser	Leu	Gly	Ala	Leu	Leu	Thr	Ala	Ala	Ala	Pro	Ala	Gln	Ala	Asn
			20					25					30		
Gln	Gln	Ile	Cys	Asp	Arg	Tyr	Gly	Thr	Thr	Thr	Ile	Gln	Asp	Arg	Tyr
		35					40					45			
Val	Val	Gln	Asn	Asn	Arg	Trp	Gly	Thr	Ser	Ala	Thr	Gln	Cys	Ile	Asn
	50					55				60					
Val	Thr	Gly	Asn	Gly	Phe	Glu	Ile	Thr	Gln	Ala	Asp	Gly	Ser	Val	Pro
65					70					75					80
Thr	Asn	Gly	Ala	Pro	Lys	Ser	Tyr	Pro	Ser	Val	Tyr	Asp	Gly	Cys	His
				85					90					95	
Tyr	Gly	Asn	Cys	Ala	Pro	Arg	Thr	Thr	Leu	Pro	Met	Arg	Ile	Ser	Ser
		100						105					110		
Ile	Gly	Ser	Ala	Pro	Ser	Ser	Val	Ser	Tyr	Arg	Tyr	Thr	Gly	Asn	Gly
		115					120					125			
Val	Tyr	Asn	Ala	Ala	Tyr	Asp	Ile	Trp	Leu	Asp	Pro	Thr	Pro	Arg	Thr
	130					135					140				
Asn	Gly	Val	Asn	Arg	Thr	Glu	Ile	Met	Ile	Trp	Phe	Asn	Arg	Val	Gly
145					150					155					160
Pro	Val	Gln	Pro	Ile	Gly	Ser	Pro	Val	Gly	Thr	Ala	His	Val	Gly	Gly
				165					170					175	
Arg	Ser	Trp	Glu	Val	Trp	Thr	Gly	Ser	Asn	Gly	Ser	Asn	Asp	Val	Ile
			180					185					190		
Ser	Phe	Leu	Ala	Pro	Ser	Ala	Ile	Ser	Ser	Trp	Ser	Phe	Asp	Val	Lys
		195				200					205				
Asp	Phe	Val	Asp	Gln	Ala	Val	Ser	His	Gly	Leu	Ala	Thr	Pro	Asp	Trp
	210					215					220				
Tyr	Leu	Thr	Ser	Ile	Gln	Ala	Gly	Phe	Glu	Pro	Trp	Glu	Gly	Gly	Thr
225					230					235					240
Gly	Leu	Ala	Val	Asn	Ser	Phe	Ser	Ser	Ala	Val	Asn	Ala	Gly	Gly	Gly
				245					250					255	
Asn	Gly	Gly	Thr	Pro	Gly	Thr	Pro	Ala	Ala	Cys	Gln	Val	Ser	Tyr	Ser
			260					265					270		
Thr	His	Thr	Trp	Pro	Gly	Gly	Phe	Thr	Val	Asp	Thr	Thr	Ile	Thr	Asn
		275					280					285			
Thr	Gly	Ser	Thr	Pro	Val	Asp	Gly	Trp	Glu	Leu	Asp	Phe	Thr	Leu	Pro
	290					295					300				
Ala	Gly	His	Thr	Val	Thr	Ser	Val	Trp	Asn	Ala	Leu	Ile	Ser	Pro	Ala

58

305					310					315					320
Ser	Gly	Ala	Val	Thr	Ala	Arg	Ser	Thr	Gly	Ser	Asn	Gly	Arg	Ile	Ala
				325					330					335	
Ala	Asn	Gly	Gly	Thr	Gln	Ser	Phe	Gly	Phe	Gln	Gly	Thr	Ser	Ser	Gly
			340					345					350		
Ala	Gly	Phe	Thr	Ala	Pro	Ala	Gly	Ala	Arg	Leu	Asn	Gly	Thr	Ser	Cys
		355					360					365			
Thr	Val	Arg													
	370														

<210> 28
 <211> 221
 <212> PRT
 <213> conArtificial Sequence

<220>
 <223> consensus sequence

<221> VARIANT
 <222> (1)...(221)
 <223> Xaa = Any Amino Acid

<400> 28															
Cys	Xaa	Gln	Tyr	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Gly	Tyr	Xaa	Xaa	Xaa	Asn
1				5					10					15	
Asn	Xaa	Trp	Gly	Xaa	Xaa	Xaa	Xaa	Xaa	Ser	Gly	Xaa	Gln	Cys	Thr	Xaa
			20					25					30		
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Trp	Xaa	Xaa	Xaa	Trp
		35				40					45				
Xaa	Trp	Ser	Gly	Gly	Xaa	Xaa	Xaa	Val	Lys	Ser	Tyr	Xaa	Xaa	Xaa	Xaa
	50				55						60				
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Ser	Xaa	Ile	Xaa	Ser	Xaa
65				70						75					80
Xaa	Xaa	Xaa	Xaa	Xaa	Trp	Xaa	Tyr	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Ala	Asn
				85				90						95	
Val	Ala	Tyr	Asp	Leu	Phe	Thr	Xaa	Xaa	Xaa	Pro	Xaa	His	Xaa	Xaa	Xaa
			100					105					110		
Xaa	Gly	Xaa	Tyr	Glu	Xaa	Met	Ile	Trp	Leu	Xaa	Xaa	Xaa	Gly	Gly	Xaa
		115					120					125			
Xaa	Pro	Ile	Gly	Ser	Xaa	Xaa	Xaa	Xaa	Val	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
	130				135					140					
Gly	Xaa	Xaa	Trp	Xaa	Leu	Xaa	Xaa	Gly	Xaa	Asn	Gly	Xaa	Met	Xaa	Val
145				150						155					160
Xaa	Ser	Phe	Val	Ala	Xaa	Ser	Ser	Ser	Ser	Ser	Ser	Phe	Xaa	Gly	Asp
			165					170						175	
Xaa	Xaa	Xaa	Phe	Xaa	Xaa	Tyr	Leu	Xaa	Xaa	Xaa	Xaa	Gly	Xaa	Pro	Xaa
			180					185					190		
Xaa	Xaa	Gln	Tyr	Leu	Xaa	Xaa	Xaa	Gln	Xaa	Gly	Thr	Glu	Pro	Phe	Thr
		195					200					205			
Gly	Xaa	Xaa	Ala	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Ala		
	210					215						220			

<210> 29
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 29
gaacgatggc aagggcggcg tgacg 25

<210> 30
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 30
cttctcgggc tgctacaacc caaacgg 27

<210> 31
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 31
acatcgtcga gtgttttggc acctac 26

<210> 32
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 32
catcgtcgag aactggggca cctacaacc 29

<210> 33
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 33
ggcacctacc gaccgtccac g 21

<210> 34
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 34
caagctgggc gagcacacct ccgac 25

<210> 35
<211> 21

<212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 35
 cgccgcaact gtcgctcgag c

21

<210> 36
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 36
 gtggagggtt accaaagctc tggctctgc

29

<210> 37
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 37
 tctggctctg cttgcatcac cgtcagc

27

<210> 38
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 38
 gagaagcgcc agtgcattca gcccggc

27

<210> 39
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 39
 gtgacgtact gcaatggtcc cggcggg

27

<210> 40
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

61

<400> 40
ggcaccaaga acaggggtcat caacttctcg ggc 33

<210> 41
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 41
tccatcacccg tcagcgatta aagggggctc ttc 33

<210> 42
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 42
cccagacgat tcagtgcggc acgggctaca ac 32

<210> 43
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 43
cttctactcg tactggtgcg atggccacgg cg 32

<210> 44
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 44
cgattcagcc cggctgcggc tacaacaacg gc 32

<210> 45
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> synthetic oligonucleotide

<400> 45
caacggctac ttctactgct actggaacga tggcc 35

<210> 46

<211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 46
 ccggcacggg ctactgcaac ggctacttct actc

34

<210> 47
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 47
 ggcgtagcgt acacctgcgg tccggcgagg c

31

<210> 48
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 48
 ggcgccacca agtgcggcga ggtcacc

27

<210> 49
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic oligonucleotide

<400> 49
 gcgtgggctc agtgcggcct gacgctcg

28

<210> 50
 <211> 752
 <212> DNA
 <213> Trichoderma reesei

<400> 50
 atggttgcc tttccagcct catctgcgct ctcaccagca tcgccagtac tctggcgatg 60
 cccacaggcc tcgagcctga gagcagtgtc aaçgtcacag agcgtggcat gtacgacttt 120
 gttcttgag ctcaaatga tcatcgccgt cgtgctagca tcaactacga ccaaaactac 180
 caaactggcg gacaagtcag ctattcgcc tccaacactg gcttctcagt gaactggaac 240
 actcaagatg actttgttgt gggcgttggt tggacgactg gatcttctgc gtaggaggac 300
 tcctcatcat tctgcacttt gaaagcatct tctgaccaa agcttctctt agtcccatca 360
 actttggcgg ctcttttagt gtcaacagcg gaactggcct gctttccgtc tatggctgga 420
 gcaccaacc actggttgag tactacatca tggaggacaa ccacaactac ccagcacagg 480
 gtaccgtcaa gggaaccgtc accagcgacg gagccactta caccatctgg gagaataccc 540
 gtgtcaacga gccttccatc cagggcacag cgacottcaa ccagtacatt tccgtgcgga 600
 actcgcccag gaccagcgga actgttactg tgcagaacca cttcaatgct tgggcctcgc 660

63

ttggcctgca ccttgggcag atgaactacc aggttgctgc tgtcgaaggc tgggggtggtgta 720
 gtgggttctgc ctcacagagt gtcagcaact ag 752

<210> 51

<211> 248

<212> PRT

<213> Trichoderma reesei

<400> 51

Met	Val	Ala	Phe	Ser	Ser	Leu	Ile	Cys	Ala	Leu	Thr	Ser	Ile	Ala	Ser	1	5	10	15
Thr	Leu	Ala	Met	Pro	Thr	Gly	Leu	Glu	Pro	Glu	Ser	Ser	Val	Asn	Val	20	25	30	
Thr	Glu	Arg	Gly	Met	Tyr	Asp	Phe	Val	Leu	Gly	Ala	His	Asn	Asp	His	35	40	45	
Arg	Arg	Arg	Ala	Ser	Ile	Asn	Tyr	Asp	Gln	Asn	Tyr	Gln	Thr	Gly	Gly	50	55	60	
Gln	Val	Ser	Tyr	Ser	Pro	Ser	Asn	Thr	Gly	Phe	Ser	Val	Asn	Trp	Asn	65	70	75	80
Thr	Gln	Asp	Asp	Phe	Val	Val	Gly	Val	Gly	Trp	Thr	Thr	Gly	Ser	Ser	85	90	95	
Ala	Glu	Asp	Ser	Ser	Ser	Phe	Cys	Thr	Leu	Lys	Ala	Ser	Ser	Asp	Gln	100	105	110	
Lys	Leu	Leu	Leu	Val	Pro	Ser	Thr	Leu	Ala	Ala	Leu	Leu	Val	Ser	Thr	115	120	125	
Ala	Glu	Leu	Ala	Cys	Phe	Pro	Ser	Met	Ala	Gly	Ala	Pro	Thr	His	Trp	130	135	140	
Leu	Ser	Thr	Thr	Ser	Trp	Arg	Thr	Thr	Thr	Thr	Gln	His	Arg	Val		145	150	155	160
Pro	Ser	Arg	Glu	Pro	Ser	Pro	Ala	Thr	Glu	Pro	Leu	Thr	Pro	Ser	Gly	165	170	175	
Arg	Ile	Pro	Val	Ser	Thr	Ser	Leu	Pro	Ser	Arg	Ala	Gln	Arg	Pro	Ser	180	185	190	
Thr	Ser	Thr	Phe	Pro	Cys	Gly	Thr	Arg	Pro	Gly	Pro	Ala	Glu	Leu	Leu	195	200	205	
Leu	Cys	Arg	Thr	Thr	Ser	Met	Leu	Gly	Pro	Arg	Leu	Ala	Cys	Thr	Leu	210	215	220	
Gly	Arg	Thr	Thr	Arg	Leu	Ser	Leu	Ser	Lys	Ala	Gly	Val	Val	Val	Val	225	230	235	240
Leu	Pro	His	Arg	Val	Ser	Ala	Thr									245			

CLAIMS

1. A modified xylanase comprising a polypeptide having an amino acid sequence with at least 80% sequence identity to SEQ ID NO: 11 wherein the polypeptide has a substituted amino acid residue at position 144 corresponding to a histidine to cysteine (H144C) substitution wherein the position of the substituted amino acid is numbered from the amino acid after the signal and pro signal sequence as set forth in SEQ ID NO: 1, and wherein the modified xylanase has xylanase activity.
2. The xylanase according to claim 1, further comprising at least one substituted amino acid residue at a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188, and 190.
3. The xylanase according to claim 1 or 2, further comprising an addition at position +191.
4. The xylanase according to any one of claims 1 to 3, comprising at least one substitution at a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 162, 180, and 186.
5. The xylanase according to claim 4, wherein the modified xylanase has at least one substitution selected from the group consisting of H22K, S65C, N92C, F93W, N97R, V108H, F180Q and S186C.
6. The xylanase according to claim 5, wherein the xylanase has the following mutations: F93W and N97R.
7. The xylanase according to any one of claims 1 to 3, wherein the xylanase has the following mutation N92K.

8. The xylanase according to claim 5, wherein the xylanase has the following mutations: F180Q and N92C.
9. The xylanase according to claim 5, wherein the xylanase has the following mutations: H22K and F180Q.
10. The xylanase according to claim 5, wherein the xylanase has the following mutation: V108H.
11. The xylanase according to claim 5, wherein the xylanase has the following mutations: S65C and S186C.
12. The xylanase according to claim 5, wherein the xylanase has the following mutations: H22K, F180Q and N92C.
13. The xylanase according to any one of claims 1 to 12, comprising a polypeptide having an amino acid sequence with at least 90% sequence identity to SEQ ID NO: 11.
14. The xylanase according to any one of claims 1 to 12, comprising a polypeptide having an amino acid sequence with at least 97% sequence identity to SEQ ID NO: 11.
15. The xylanase according to any one of claims 1 to 12, comprising a polypeptide having an amino acid sequence with at least 90% sequence identity to SEQ ID NO: 1.
16. The xylanase according to any one of claims 1 to 12, comprising a polypeptide having an amino acid sequence with at least 97% sequence identity to SEQ ID NO: 1.
17. The xylanase according to any one of claims 1 to 16, wherein the modified xylanase has increased thermostability or pH-stability under basic conditions with respect to the corresponding wild-type xylanase.
18. A nucleic acid encoding a xylanase according to any one of claims 1 to 17.

19. A modified family 11 xylanase comprising an amino acid sequence, the amino acid sequence having a substituted amino acid residue at a position equivalent to 144 in SEQ ID NO: 1, wherein the amino acid at the position equivalent to 144 is substituted to cysteine and the position of the substituted amino acid is numbered from the amino acid after the signal and pro signal sequence, wherein said modified xylanase has at least 90% sequence identity to SEQ ID NO: 1 and has xylanase activity.

20. The xylanase according to claim 19, wherein the amino acid sequence has at least one further substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 22, 28, 58, 65, 92, 93, 97, 105, 108, 162, 180 and 186.

21. The xylanase according to claim 19 or 20, wherein the amino acid sequence has an addition at a position equivalent to +191.

22. The xylanase according to any one of claims 19 to 21, further comprising at least one substituted amino acid residue at a position equivalent to a position selected from the group consisting of: 2, 5, 7, 10, 11, 16, 19, 22, 26, 28, 29, 30, 34, 36, 38, 57, 58, 61, 63, 65, 67, 92, 93, 97, 105, 108, 110, 111, 113, 132, 143, 147, 149, 151, 153, 157, 160, 162, 165, 169, 180, 184, 186, 188 and 190.

23. The xylanase according to claim any one of claims 19 to 22, wherein the xylanase has a H144C substitution.

Figure 1

	10	20	30	40	50
XYN2_TRIRE	---MVSFTSLLAASPP-SRASCRPAAEV---ESVAVEKRQTIQ-----P				
XYN1_HUMIN	---MVSLKSVLAAATAVSSAIAAPFDFVPRDNSTALQARQVTP-----N				
XYNA_BACST	-----MKLKKKMLTLLLTASMSFGLF-----G				
XYN1_TRIRE	---MVAFSSLICALTSIASTLAMPTGLEPESSVNVTERGMYDFVLGAHND				
XYN1_ASPAW	-----MKVTAAAFAGLLVTAFAPVPEPVLVS-----				
XYN2_BACST	MCSSIPSLREVFANDFRIGA AVNPVTLEAQQLIRHVNSLTAENHMKFE				

	60	70	80	90	100
XYN2_TRIRE	GTGYNNGYFYSYWNDGHGGV TYTNGPGGQFSVNWS--NSG-NFVGGKGWQ				
XYN1_HUMIN	AEGWHNGYFYSWWSDGGGQVQYTNLEGSRYQVRWR--NTG-NFVGGKGW N				
XYNA_BACST	ATSSAATDYWQYWT DGGGMVNAVNGPGGNYSVTWQ--NTG-NFVVGKGW T				
XYN1_TRIRE	HRRRASINYDQNYQTG-GQVSYSPTSNTG-FSVNWN--TQD-DFVVGVGW T				
XYN1_ASPAW	--RSAGINYVQNYNGNLGDFTYDESAGT-FSMYWEDGVSS-DFVVG LGW T				
XYN2_BACST	HLQPEEGRFTFDIAIKSSTSPFSSHGVRGHTLVWHNQTPSWVFQDSQGHF				

	110	120	130	140	150
XYN2_TRIRE	PGTKNKVINFS-GSYNPNGNSYLSVYGWSRNPLIEYYIVENF---GTYNP				
XYN1_HUMIN	PGT-GRTINYG-GYFNPQGNGLAVYGWTRNPLVEYYVIESY---GTYNP				
XYNA_BACST	VGSPNRVINYNAGIWEPSGNGYLTLYGWTRNALIEYYVVD SW---GTYRP				
XYN1_TRIRE	TGS-SAPINFGGSFSVNSGTGLLSVYGWSTNPLVEYYIMED-----NHNY				
XYN1_ASPAW	TGS-SNAITYSAEYSASGSSSYLAVYGWVNYPQAEYYIVEDY---GDYNP				
XYN2_BACST	VGRDVLLERMKSHISTVVQRYKGKVCWDVINEA VADEGSEWLRSSTWRQ				

	160	170	180	190	200
XYN2_TRIRE	STGATKLGEVTSDGSVYDIYRTQRVNQPSIIGTATFYQYWSVRRNHRSSG				
XYN1_HUMIN	GSQAQYKGT FYTDGDQYDIFVSTRYNQPSIDGTRTFQQYWSIRKNKRVGG				
XYNA_BACST	T--GNYKGT VNSDGGTYDIYTTMRYNAPSIDGTQTFQQFWSVRQSKRPTG				
XYN1_TRIRE	PAQGT VKGT VTS DGATYTIWENTRVNEPSIQGTATFNQYISVRNSPRTSG				
XYN1_ASPAW	CSSATSLGTVYSDGSTYQVCTDTRTNEPSITGTSTFTQYFSVRESTRTSG				
XYN2_BACST	IIGDDFIQQAFLYAHEADPEALLFYNDYNECFPEKREKIYTLVKSLRDKG				

Figure 1

	210	220	230	240	250
XYN2_TRIRE	S---VNTANHFNAWA-QQGLTLGTMD-YQIVAVEGYFSSGSASITVS--				
XYN1_HUMIN	S---VNMQNHFNAWQ-QHGMP LGQHY-YQVVATEGYQSSGESDIYVQTH				
XYNA_BACST	SNV-SITFSNHVNAWR-SKGMNLGSSWAYQVLATEGYQSSGRSNVTW--				
XYN1_TRIRE	T---VTVQNHFNAWA-SLGLHLGQMN-YQVVAVEGWGGSGSASQSVSN-				
XYN1_ASPAW	T---VTVANHFNFWA-QHGFGNSDFN-YQVMAVEAWSGAGSASVTISS-				
XYN2_BACST	IPIHGIGMQAHWSLNRPTLDEIRAAIERYASLG VILHITELDISMFEFDD				

	260	270	280	290	300
XYN2_TRIRE	-----				

Figure 1

XYN1_HUMIN	-----
XYNA_BACST	-----
XYN1_TRIRE	-----
XYN1_ASPAW	-----
XYN2_BACST	HRKDLAAPTNE MVERQAERYEQIFSLFKEYRDV IQNVTFWGIADDHTWLD

	310	320	330
XYN2_TRIRE	-----		
XYN1_HUMIN	-----		
XYNA_BACST	-----		
XYN1_TRIRE	-----		
XYN1_ASPAW	-----		
XYN2_BACST	HFPVQGRKNWPLLFDEQHNP KP AFWRVVNI		

FIGURE 2




Figure 3

H22K	5'- GAACGATGGCA <u>AAGGG</u> CGGCGTGACG -3'
S65C	5'- CTTCTCGGGCTGCTACAACCCAAACGG -3'
N92C	5'- ACATCGTCGAGT <u>GTTT</u> TGGCACCTAC -3'
F93W	5'- CATCGTCGAGAACTGGGGGCACCTACAACC -3'
N97R	5'- GGCACCTACCGA <u>CCGT</u> CCACG -3'
V108H	5'- CAAGCTGGGCGAGCACACCTCCGAC -3'
H144C	5'- CGCCGCAACTGTCGCTCGAGC -3'
F180Q	5'- GTGGAGGGTTACCAAAGCTCTGGCTCTGC -3'
S186C	5'- TCTGGCTCTGCTTGCATCACCGTCAGC -3'
T2C	5'-GAGAAGCGCCAGTGCATTAGCCCGGC-3'
T28C	5'-GTGACGTACTGCAATGGTCCCGGGCGGG-3'
K58R	5'-GGCACCAAGAACAGGGTTCATCAACTTCTCGGGC-3'
191D	5'-TCCATCACCGTCAGCGATTAAAGGGGGCTCTTC-3'
P5C	5'-CCCAGACGATTCAGTGCGGCACGGGCTACAAC-3'
N19C	5'-CTTCTACTCGTACTGGTGCGATGGCCACGGCG-3'
T7C	5'-CGATTCAGCCCGGCTGCGGCTACAACAACGGC-3'
S16C	5'-CAACGGCTACTTCTACTGCTACTGGAACGATGGCC-3'
N10C	5'-CCGGCACGGGCTACTGCAACGGCTACTTCTACTC-3'
N29C	5'-GGCGTGACGTACACCTGCGGTCCCGGGCGGGC-3'
L105C	5'-GGCGCCACCAAGTGCGGGCGAGGTCACC-3'
Q162C	5'-GCGTGGGCTCAGTGCGGCCTGACGCTCG-3'

Figure 4.




Figure 5




Figure 6




Figure 7




Figure 8





Figure 9.


Figure 10.

Figure 11.





Figure 12.

Figure 13.




Figure 14.





Figure 15.

Figure 16.




Figure 17




Figure 18

Trichoderma reesei Xyl II protein (high pl xylanase)
the full sequence, including signal and pro sequence

MVSFTSLLAGVAISGVLAAPAAEVESVAVEKRQTQPQTGYNNGYFYSWNDGHGGVITYTNGPGGQFVNSNSGNFVG
GKGWQPGTKNKVINFSGSYNPNNGNSYLSVYGWSRNPLEYIYVENFGTYNPNSTGATKLGEVTSDSGSVDIYRTQRVNQPS
IIGTATFYQYWSVRRNHRSSGSVNTANHFNAWAQQGLTLGTM DYQIVAVEGYFSSGSASITVS (SEQ. ID NO: 1)

Trichoderma reesei XynII gene (high pl xylanase)
DNA from start codon to stop codon (includes a single intron)

ATGGTCTCCTTCACCTCCCTCCCTCGCCGGCGTGCGCCGCGCATCTCGGGCGTCTTGGCCGCTCCCGCCGAGGTCGAATC
CGTGGCTGTGGAGAAGCGCCAGACGATTACGCCGGCACGGCTACAACAACGGCTACTTCTACTCGTACTGGAACGATG
GCCACGGCGCGTGACGTACACCAATGTCCCGGGCGGAGTTCTCCGTCAACTGGTCCAACTCGGGCAACTTTGTCGGC
GGCAAGGGATGGCAGCCCGCACCAAGTAAGACTACTCTTACCCCTTTGACCAACACAGCACACACAA
TACAACACATGTGACTACCAATCATGGAATCGGATCTAACAGCTGTGTTTCAAAAAGGGTCATCAACTTCTCGGGC
AGCTACAACCCCAACGGCAACAGCTACCTCTCCGTGTACGGCTGGTCCCGCAACCCCTGATCGAGTACTACATCGTCGA
GAACTTTGGCACCTACAACCCGTCCACGGCGCCACCAAGCTGGGCGAGGTCACCTCCGACGGCAGCGTCTACGACATTT
ACCGCACGACGCGTCAACCAAGCCGTCCATCATCGGCACCGCACCTTTTACCAGTACTGTCGTCGCGCGCAACCAC
CGCTCGAGCGGCTCCGTCAACACACGGCGAACCACCTTCAACGCGTGGGCTCAGCAAGGCCTGACGCTCGGGACGATGGATTA
CCAGATTGTTGCCGTGGAGGGTTACTTTAGCTCTGGCTCTGCTCCATCACCGTCAGCTAA (SEQ. ID NO: 2)

Figure 19

Trichoderma reesei EGL III protein (endoglucanase III)
the full sequence, including signal sequence

MKFLQVLPALIPAALAQTSQDQWATFTGNGYTVSNNLWGASAGSGFGCVTAVSLSGASWHADWQWSGGQNNVKSQNSQ
IAIPQKRTVNSISSMPTTASWSYSGSNIRANVAYDLFTAANPNHVTYSGDYELMIWLKYGDIGPIGSSQGTNVVGGQSW
TLYYGYNGAMQVYSFVAQTNTTNYSGDVKNFFNYLRDNKGYNAAGQYVLSYQFQTEPFTGSGTLNVASWTASIN
(SEQ. ID NO: 3)

Trichoderma reesei EG III gene (endoglucanase III)
DNA from start codon to stop codon (includes two introns)

ATGAAGTTCCTTCAAGTCCTCCCTGCCCTCATACCGCGCCCTGGCCCAACCAAGCTGTGACCAGTGGGCAACCTTCAC
TGGCAACGGGTACACAGTCAGCAACAACCTTTGGGAGCATCAGCCGGCTCTGGATTTGGCTGCGTGACGGCGGTATCGC
TCAGCGCGGGCCTCCTGGCACGCAGACTGGCAGTGGTCCGGCGGCCAGAACACGTCAAGTCGTACCAGAACTCTCAG
ATTGCCATTCCCCAGAAAGAGGACCGTCAACAGCATCAGCAGCATGCCACCACTGCCAGCTGGAGCTACAGCGGAGCAA
CATCCGCGCTAATGTTGCGTATGACTTGTACCCGACGCCAACCGAATCATGTCACTCGGGAGACTACGAACTCA
TGATCTGGTAAGCCATAAGAAGTGACCCCTCCTTGATAGTTTCGACTAACAAACATGTCTTGAGGCTTGCAAAATACGGCGA
TATTGGCCGATTGGGTCCTCACAGGGAACAGTCAACGTCGGTGGCCAGAGCTGGACGCTCTACTATGGCTACAACGGAG
CCATGCAAGTCTATTCCCTTTGTGGCCAGACCAACTACCAACTACAGCGGAGATGTCAAGAACTTCTTCAATTATCTC
CGAGACAATAAAGGATACAAACGCTGCAGGCCAATATGTTCTTAGTAAGTCACCCCTCACTGTGACTGGGCTGAGTTTGTG
CAACGTTTGCTAACAAACCTTCGTATAGGCTACCAATTTGGTACCGAGCCCTTCACGGGCAGTGGAACCTCTGAACGTCG
CATCCTGGACCGCATCTATCAACTAA (SEQ. ID NO: 4)

Figure 20

atggtgcctttccagccatctcgcgctctaccagcatcgccagtactctggcgatgccacaggccctgagcctgagagcagtgtaacgtcacagagcgtggcatgtacgactttgttctt
ggagctcacaaatgatcatcgccgtcgtgctagcatcaactacgaccaaactaccgacgacaaactggcggacaagtcagctattcgccttccaacactggccttctcagtgaactgggaacactcaagatg
actttgttgaggcggttggtggacgactggatcttctcgtaggaggactcctcatcttgcactttgaaagcattctcttagtcccatcaactttggcggctcttttagtg
tcaacagcggaaactggcctgctttccgtctatggctggagcaccacccactgggtgagtactacatcatggaggacaaccacaactaccagcacagggtaccgtcaagggaaccgtcacc
agcgacggaggccacttacaccatctgggagatacccggtgtcaacgagccttccatccaggcacagcgacctcaacttccgtcggaaactcggccaggaccagcgggaactg
ttactgtgcagaaccacttcaatgcttgggcctcgcttggcctgcaccttgggcagatgaactaccagggtgtcgtgctggaaggctggggtgtagtggtctgcctcacagagtgcagcaac
tag

Xylanase I Amino Acid sequence

MVAFSSLICALTSIASTLAMPTGLEPESSVNVTERGMYDFVLGAHNDHRRRASINYDQNYQTGGQVSYSPSNTGFSVNW
N
TQDDFVVGWTTGSSA.EDSSSFCTLKASSDQKLLVPSTLAALLVSTAELACFPSMAGAPTHWLSTTSWRTTTTQHR
VPSREPSATEPLTPSGRIPVSTSLPSRAQRPSTSTFP CGTRPGPAELLLCRTTSM LGPRLACTLGR.TTRL SLSKAGVV
VVLPHRVSAT

