



- (51) **International Patent Classification:**
A61K 8/49 (2006.01) *A61Q 5/10* (2006.01)
- (21) **International Application Number:**
PCT/US20 12/025371
- (22) **International Filing Date:**
16 February 2012 (16.02.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/445,275 22 February 2011 (22.02.2011) US
- (71) **Applicant (for all designated States except US):** **THE PROCTER & GAMBLE COMPANY** [US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio 45202 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **LIM, Muil** [KR/KR]; 1008-802 Beong-Gil 47, Seungigol-Gil Chuncheon-Si, Kangwon-DO, 200-775 (KR). **MURPHY, Bryan, Patrick** [US/US]; 974 Arnold Palmer Drive, Loveland, Ohio 45140 (US). **POPP, Margaret, Ann** [US/US]; P&G, MVIC 11810 East Miami River Road, Cincinnati, Ohio 45252 (US). **SUTTON, Richard, Matthew Charles** [GB/US]; 12141 Paulmeadows Drive, Cincinnati, Ohio 45249 (US). **ZHANG, Guiru** [US/US]; 3657 Avalon Trail, Lebanon, Ohio 45036 (US).
- (74) **Common Representative:** **THE PROCTER & GAMBLE COMPANY**; Eileen L. Hughett, Global Patent



Services, 299 East Sixth Street, Sycamore Building, 4th Floor, Cincinnati, Ohio 45202 (US).

- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) **Title:** OXIDATIVE DYEING COMPOSITIONS COMPRISING AN 1-HEXYL/HEPTYL-4,5-DIAMINOPYRAZOLE AND A BENZO[1,3]DIOXOL-5-YLAMINE AND DERIVATIVES THEREOF



(57) **Abstract:** A composition for the oxidative dyeing of keratin fibers, in particular human keratin fibers, comprising (A) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I), its physiologically compatible water-soluble salt, or mixtures thereof; and (B) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II), its physiologically compatible water-soluble salt, or mixtures thereof; and (C) an oxidizing agent, wherein **R₁**, **R₂**, **R₃**, **R₄**, **R₅**, **R₆** and **R₇** are as defined herein and **a** = 1 or 2.

Oxidative Dyeing Compositions Comprising an 1-Hexyl/Heptyl-4,5-diaminopyrazole and a
Benzo[1,3]dioxol-5-ylamine and Derivatives Thereof

FIELD OF THE INVENTION

The present invention relates to a composition for the oxidative dyeing of keratin fibers,
5 in particular human keratin fibers, comprising (A) a 1-hexyl/heptyl-4,5-diaminopyrazole
compound as defined in formula (I) hereafter, its physiologically compatible water-soluble salt or
mixtures thereof, and (B) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II) as
defined hereafter, its physiologically compatible water-soluble salt, or mixtures thereof, and (C)
an oxidizing agent.

10

BACKGROUND OF THE INVENTION

The oxidative dyeing of hair is one of the most extensively used methods to color hair. In
this process, oxidative hair coloring precursors are used in combination with an oxidizing agent,
commonly a peroxy oxidizing agent. The precursors are generally small molecules capable of
diffusing into hair, generally comprising primary intermediates and couplers. A wide variety of
15 primary intermediates and couplers have been employed in such oxidative hair coloring
compositions. Selections of specific combinations of couplers and primary intermediates allow
reaching the desired nuance of hair color.

Besides providing the desired color, the dye precursors used for oxidative hair dyeing
have to also meet some additional requirements. Indeed, the combination of couplers and primary
20 intermediates used should deliver the desired intensity together with good wash and bleeding
fastness and be acceptable from a toxicological and dermatological point of view.

Pyrazole compounds have been disclosed as oxidative dye precursors. A colorant for
keratin fibers, which is not mixed with an oxidant before use and free from manganese salts and
consists of a combination suitable for coloring keratin fibers in the presence of atmospheric
25 oxygen in a suitable cosmetic medium is known. The combination comprises: (a) 4,5-
diaminopyrazole compound(s); and (b) compound(s) selected from 5-amino-2-methylphenol(s)
and/or m-phenylenediamine compound(s) or their physiologically-compatible salts. See for
example EP 1,166,748A2.

Oxidation colorant compositions containing a combination of at least one
30 diaminopyrazole derivative; and at least one heterocyclic coupler (A) selected from indole

derivatives, benzimidazole derivatives, benzomorpholine derivatives, pyridine derivatives, indoline derivatives, quinoline derivatives, sesamol derivatives and their acid-addition salts are known.

Furthermore, these disclosures do not disclose an example with a selection of a C6/C7
5 straight chain alkyl substituent on the N-1 position of a 4,5-diaminopyrazole. See for example EP 0728464.


The compositions, known in the prior art, comprising pyrazole compounds are usually not satisfactory, especially in terms of intensity, wash fastness and bleeding.

Thus, there is a need to develop new compositions for the oxidative dyeing of hair
10 providing satisfactory color intensity along with good wash fastness and bleeding properties.

The invention provides novel dye compositions that overcome known drawbacks of pyrazole compounds. In particular, the invention provides good hair color intensity together with good wash and bleeding fastness. Indeed, it has surprisingly been found that the modification of the pyrazole ring system on the N-1 position by a C6/C7 straight alkyl chain combined with
15 benzo[1,3]dioxol-5-ylamine couplers provides excellent properties of resistance to the various treatments which keratinous fibers may undergo.

SUMMARY OF THE INVENTION

The present invention relates to a composition for the oxidative dyeing of keratin fibers, in particular human keratin fibers, comprising (A) a 1-hexyl/heptyl-4,5-diaminopyrazole
20 compound of the general formula (I), its physiologically compatible water-soluble salt, or mixtures thereof; and (B) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II), its physiologically compatible water-soluble salt, or mixtures thereof; and (C) an oxidizing agent.



wherein R₁, R₂, R₃, R₄, R₅, R₆ and R₇ are as defined in the claims and a = 1 or 2.

The present invention also relates to a method of dyeing hair comprising the steps of
25 applying this composition to hair, and to a kit comprising: (i) a tint component comprising at least (a) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I), its

physiologically compatible water-soluble salt, or mixtures thereof; and, (b) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II), its physiologically compatible water-soluble salt, or mixtures thereof, and (ii) a developer component comprising (c) an oxidizing agent.

DETAILED DESCRIPTION OF THE INVENTION

5 Definitions

As used herein, the term "composition for the oxidative dyeing of keratin fibers" means a ready-to-use composition in a suitable carrier medium for dyeing keratin fibers, in particular human hair, comprising oxidative dye precursors (primary intermediates and couplers) and an oxidizing agent. These compositions can typically be the result of a mixture of two compositions
10 namely a tint component comprising the dye precursors and usually an alkalizing agent such as ammonia and/or monoethanolamine and a developer component comprising the oxidizing agent.


As used herein, the term "keratin" refers to a scleroprotein found in epidermal tissues and modified into hard structures such as horns, hair, and nails. As used herein, the term "hair" refers to keratinous fibers on a living, e.g. a person, or non-living body, e.g. in a wig, hairpiece, or other
15 aggregation of non-living keratinous fibers. Mammalian, preferably human, hair is preferred. Notably, hair, wool, fur, and other keratinous fibers are suitable substrates for coloring by the compounds and compositions described herein.

As used herein, the term "dye precursors" refers to compounds that may be used in the composition to act as primary intermediates, couplers, or both, in order to provide color to
20 ketatinous fibers.

It is to be understood that when this development refers to a particular structure, all of the reasonable additional tautomeric structures are included. In the art, tautomeric structures are frequently represented by one single structure and the invention follows this general practice.

Pyrazole compound

25 The 1-hexyl/heptyl-4,5-diaminopyrazole compound of the invention is of the general formula (I), its physiologically compatible water-soluble salt, or mixtures thereof:



wherein a is equal to one or two;

wherein **R_i** is selected from the group consisting of:

(a) C-linked substituents selected from the group consisting of:

5 (i) substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic substituents, in particular alkyl or hydroxy substituted alkyl,

(ii) substituted or unsubstituted, mono- or poly- unsaturated aromatic or heteroaromatic substituents, in particular aryl or heteroaryl,

10 wherein said C-linked substituents comprise from 1 to 6 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, F, N, P and Si;

(b) S-linked substituents selected from the group consisting of SA¹, SO₂A¹, SO₃A¹, SSA¹, SOA¹, SOzNA², SNA*A², and SONA²;

(c) O-linked substituents selected from the group consisting of OA¹, ONA²;

15 (d) N-linked substituents selected from the group consisting of NA¹A²; (NA¹A²A³)⁺, NA², NO₂; NA*A²;

(e) halogens selected from the group consisting of F, Cl, Br, and I; and

(f) hydrogen;


20 wherein A¹, A², and A³ are independently selected from the group consisting of hydrogen; substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic or aromatic or heteroaromatic substituents, preferably substituted or unsubstituted alkyl, or A¹ and A² together with nitrogen atoms to which they are bound form a ring; wherein said substituents or ring comprise from 1 to 6 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, S, N, P, and Si;

25 R₂ and R₃ are selected independently of each other from the group consisting of a hydrogen atom; a C1-C6 alkyl substituent; a trifluoromethyl substituent; a C1-C6 aminoalkyl substituent; a C1-C6 hydroxyalkyl substituent; C1-C6 an alkoxyalkyl substituent.

Suitable salts of formula **(I)** are for example chlorides, bromides, sulfates, malates, tartrates, lactates and acetates. Of particular interest is the hemisulfate salt, which can provide a better stability during storage than other salts such as chloride or sulfate.

30 In a preferred embodiment the 1-hexyl/heptyl-4,5-diaminopyrazole compound **(I)** is a compound with formula **(1.1)**:

5




wherein a is equal to one or two.

The 1-hexyl/heptyl-4,5-diaminopyrazole compounds of formula (I) are combined with a benzo[1,3]dioxol-5-ylamine compound of formula (II) in order to provide a resulting hair color with excellent washfastness, bleeding fastness, and intensity.

Benzo[1,3]dioxol-5-ylamine compound

The benzo[1,3]dioxol-5-ylamine compound of the invention is according to the general formula (II), its physiologically compatible water-soluble salt, or mixtures thereof;



wherein R_4 , R_5 , R_6 , R_7 are substituents selected independently of each other from the group consisting of:

(a) C-linked substituents selected from the group consisting of:

(i) substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic substituents, in particular alkyl or hydroxy substituted alkyl

(ii) substituted or unsubstituted, mono- or poly- unsaturated aromatic or heteroaromatic substituents, in particular aryl or heteroaryl,

wherein said C-linked substituents comprise from 1 to 10 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, F, N, P and Si;

(b) S-linked substituents selected from the group consisting of SA^1 , SO_2A^1 , SO_3A^1 , SSA^1 , SOA^1 , $SOzNA^2$, SNA^1A^2 , and $SONA^2$;

(c) O-linked substituents selected from the group consisting of OA^1 , ONA^2 ;

(d) N-linked substituents selected from the group consisting of NA^1A^2 ; $(NA^1A^2A^3)^+$, NA^2 , NO_2 ; NA^2 ;

(e) halogens selected from the group consisting of F, Cl, Br, and I; and

(f) hydrogen;


5 wherein A^1 , A^2 , and A^3 are independently selected from the group consisting of: hydrogen; substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic or aromatic or heteroaromatic substituents, preferably substituted or unsubstituted alkyl, or A^1 and A^2 together with nitrogen atoms to which they are bound form a ring; wherein
10 said substituents or ring comprise from 1 to 10 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, S, N, P, and Si;

Suitable salts of formula (II) are for example chlorides, bromides, sulfates, malates, tartrates, lactates and acetates.

In preferred embodiments, R_6 are selected from hydrogen, C1-C6 alkyl, C2-C6 hydroxyalkyl and mixtures thereof. Preferably R_4 , R_5 and R_7 are selected from hydrogen, CH_3
15 and mixtures thereof.

Preferred benzo[1,3]dioxol-5-ylamine derivatives of formula (II) are benzo[d][1,3]dioxol-5-amine, 2-(benzo[d][1,3]dioxol-5-ylamino)ethanol, 3-(benzo[d][1,3]dioxol-5-ylamino)propan-1-ol, 4-(benzo[d][1,3]dioxol-5-ylamino)butan-1-ol, 5-(benzo[d][1,3]dioxol-5-ylamino)pentan-1-ol, 6-(benzo[d][1,3]dioxol-5-ylamino)hexan-1-ol, N-
20 pentylbenzo[d][1,3]dioxol-5-amine, N-ethylbenzo[d][1,3]dioxol-5-amine, N-propylbenzo[d][1,3]dioxol-5-amine, N-butylbenzo[d][1,3]dioxol-5-amine, N-pentylbenzo[d][1,3]dioxol-5-amine and N-hexylbenzo[d][1,3]dioxol-5-amine.


In a more preferred embodiment the benzo[1,3]dioxol-5-ylamine derivative (II) is a compound with formula (II.1), also referred to as aminol:



(II.1)

25

The hair dyeing composition of the present invention may further comprise an auxiliary coupler. In a preferred embodiment the auxiliary coupler is any of the compounds with formula (III.1) to (III.8) below and combinations thereof:



It has surprisingly been found that the modification of the pyrazole ring system in N-l-
 position by a C-6/C-7 straight alkyl chain provides a dye precursor which couples with a
 5 benzo[1,3]dioxol-5-ylamine compound to give excellent properties of resistance to the various
 treatments which keratinous fibers may undergo, in particular washfastness and resistance to
 bleeding.

Oxidizing Agent

The compositions of the invention comprise an oxidizing agent. Typical suitable
 10 oxidizing agents for the oxidative dyeing of keratin fibers may be selected from hydrogen
 peroxide, sodium periodate, urea peroxide, melamine peroxide, perborates, percarbonates,
 perphosphates, persulfates, oxidizing enzymes such as uricases, oxidases, and
 peroxidases, and mixtures thereof. Hydrogen peroxide, perborates, or percarbonates may be
 preferred.

15 Another potential oxidizing agent for use herein is a source of peroxymonocarbonate
 ions. Preferably, such a source is formed in situ from a source of hydrogen peroxide and a
 hydrogen carbonate ion source. Such an oxidizing agent has been found to be particularly
 effective at a pH of up to and including 9.5, preferably from about 7.5 to about 9.5 more
 preferably about pH 9. Moreover, this system is also particularly effective in combination with a
 20 source of ammonia or ammonium ions.

Accordingly, any source of these peroxymonocarbonate ions may be utilized. Suitable
 sources for use herein include sodium, potassium, guanidine, arginine, lithium, calcium,
 magnesium, barium or ammonium salts of carbonate, carbamate and hydrocarbonate ions and
 mixtures thereof. In particular, sodium carbonate, sodium hydrogen carbonate, potassium
 25 carbonate, potassium hydrogen carbonate, guanidine carbonate, guanidine hydrogen carbonate,

lithium carbonate, calcium carbonate, magnesium carbonate, barium carbonate, ammonium carbonate, ammonium hydrogen carbonate and mixtures thereof may be preferred. Percarbonate salts may also be utilized to provide both the source of carbonate ions and as an oxidizing agent. Preferred sources of carbonate ions, carbamate and hydrocarbonate ions are sodium hydrogen carbonate, potassium hydrogen carbonate, ammonium carbonate, ammonium carbamate, and mixtures thereof.

The hair dyeing composition may usually comprise from about 1% to about 15% by total weight of the composition, typically from about 1.5% to about 10% by weight, and more typically from about 2% to about 8% by weight of the oxidizing agent relative to the total weight of the composition.

The oxidizing agent may be provided in a developer component which is mixed to a tint component to obtain the composition of the invention. The developer component may be based on any desired formulation chassis, including any commercial product, for example an oil-in-water emulsion. Typical developer components comprise about 6% or about 9% of $\frac{3}{4}$ O₂ by total weight of the composition. A commercial example is the Welloxon® Emulsion with respectively about 6% and about 9% H₂O₂, marketed by Wella and comprising as INCI ingredients: Water, Hydrogen Peroxide, Cetearyl Alcohol, Ceteareth-25, Salicylic Acid, Phosphoric Acid, Disodium Phosphate, Etidronic Acid.

The hair dyeing composition of the invention may be formed as thick liquid, cream, gel, emulsion, foam, aerosol mousse or as a solid form to which water is added to generate the oxidant and form a thickened vehicle suitable for hair coloring. They may comprise in addition to the ingredients indicated above further ingredients in order to further enhance the properties of the composition, including but not limited to: solvents; oxidative dyes, direct dyes; oxidizing agents; radical scavengers; thickeners and or rheology modifiers; chelants; pH modifiers and buffering agents; carbonate ion sources; peroxy monocarbonate ion sources; anionic, cationic, nonionic, amphoteric or zwitterionic surfactants, or mixtures thereof; anionic, cationic, nonionic, amphoteric or zwitterionic polymers, or mixtures thereof; fragrances; enzymes; dispersing agents; peroxide stabilizing agents; antioxidants; natural ingredients, e.g. proteins and protein compounds, and plant extracts; conditioning agents including silicones and cationic polymers, ceramides, preserving agents; and opacifiers and pearling agents (such as titanium dioxide and mica). Some adjuvants referred to above, but not specifically described below, which are suitable are listed in the International Cosmetics Ingredient Dictionary and Handbook, (8th ed.; The

Cosmetics, Toiletry, and Fragrance Association). Particularly, vol. 2, sections 3 (Chemical Classes) and 4 (Functions) are useful in identifying specific adjuvants to achieve a particular purpose or multipurpose. A few of these ingredients are discussed hereinbelow, whose disclosure is of course non-exhaustive.

5 **Alkalizing agent**

The composition for the oxidative dyeing of keratin fibers may further comprise, generally in the tint component, an alkalizing agent as known in the art. Any alkalizing agent known in the art may be used such as ammonia, alkanolamines for example monoethanolamine, diethanolamine, triethanolamine, monopropanolamine, dipropanolamine, tripropanolamine, 2-
10 amino-2-methyl-1,3-propanediol, 2-amino-2-methyl-1-propanol, and 2-amino-2-hydroxymethyl-1,3-propanediol, guanidium salts, alkali metal and ammonium hydroxides such as sodium hydroxide, alkali metal and ammonium carbonates, and mixtures thereof. Typical alkalizing agents are ammonia and/or monoethanolamine.

Typically, the compositions for the oxidative dyeing of keratin fibers comprise from
15 about 0.1% to about 10%, preferably from about 0.5% to about 6%, more preferably from about 1% to about 4% by weight of the alkalizing agent relative to the total weight of the composition.

Primary intermediates

In addition to the pyrazole compounds of the invention, the hair dyeing compositions of the invention may comprise further primary intermediates. Suitable primary intermediates for use
20 in the compositions described herein include, but are not limited to: toluene-2,5-diamine, p-phenylenediamine, N-phenyl-p-phenylenediamine, N,N-bis(2-hydroxyethyl)-p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, hydroxypropyl-bis-(N-hydroxyethyl)-p-phenylenediamine), 2-methoxymethyl-p-phenylenediamine, 2-(1,2-dihydroxyethyl)-p-phenylenediamine, 2,2'-(2-(4-aminophenylamino)ethylazanediy)diethanol, 2-(2,5-diamino-4-
25 methoxyphenyl)propane-1,3-diol, 2-(7-amino-2H-benzo[b][1,4]oxazin-4(3H)-yl)ethanol, 2-chloro-p-phenylenediamine, p-aminophenol, p-(methylamino)phenol, 4-amino-m-cresol, 2-methoxy-p-phenylenediamine, 2,2'-methylenebis-4-aminophenol, 2,4,5,6-tetraminopyrimidine, 2,5,6-triamino-4-pyrimidinol, 1-hydroxyethyl-4,5-diaminopyrazole sulfate, 4,5-diamino-1-methylpyrazole, 4,5-diamino-1-ethylpyrazole, 4,5-diamino-1-isopropylpyrazole, 4,5-diamino-1-
30 butylpyrazole, 4,5-diamino-1-pentylpyrazole, 4,5-diamino-1-benzylpyrazole, 2,3-diamino-6,7-dihydropyrazolo[1,2-a]pyrazol-1(5H)-one dimethosulfonate, salts thereof (for example chlorides, bromides, sulfates, malates, tartrates, lactates and acetates) and mixtures thereof.

Of particular interest are toluene-2,5-diamine, p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, 2-methoxymethyl-p-phenylenediamine, 2-(2,5-diamino-4-methoxyphenyl)propane-1,3-diol, 2-chloro-p-phenylenediamine, 2-methoxy-p-phenylenediamine, 2-((2-(4-amino-phenylamino)-ethyl)-(2-hydroxyethyl)-amino)-ethanol, their salts and combinations.

Typically, the compositions for the oxidative dyeing of keratin fibers comprise from about 0.1% to about 10%, preferably from about 0.3% to about 6%, more preferably from about 0.5% to about 4% by weight of the primary intermediates relative to the total weight of the composition.

10 Couplers

In addition to the benzo[1,3]dioxol-5-ylamine compounds of the invention, the hair dyeing compositions of the invention may comprise couplers to obtain various shades. Suitable couplers for use in the compositions described herein include, but are not limited to: resorcinol, 4-chlororesorcinol, 2-chlororesorcinol, 2-methylresorcinol, 4,6-dichlorobenzene-1,3-diol, 2,4-dimethylbenzene-1,3-diol, m-aminophenol, 4-amino-2-hydroxytoluene, 2-methyl-5-hydroxyethylaminophenol, 3-amino-2,6-dimethylphenol, 3-amino-2,4-dichlorophenol, 5-amino-6-chloro-o-cresol, 5-amino-4-chloro-o-cresol, 6-hydroxybenzomorpholine, 2-amino-5-ethylphenol, 2-amino-5-phenylphenol, 2-amino-5-methylphenol, 2-amino-6-methylphenol, 2-amino-5-ethoxyphenol, 5-methyl-2-(methylamino)phenol, 2,4-diaminophenoxyethanol, 2-amino-4-hydroxyethylaminoanisole, 1,3-bis-(2,4-diaminophenoxy)-propane, 2,2'-(2-methyl-1,3-phenylene)bis(azanediyl)diethanol, benzene-1,3-diamine, 2,2'-(4,6-diamino-1,3-phenylene)bis(oxy)diethanol, 3-(pyrrolidin-1-yl)aniline, 1-(3-(dimethylamino)phenyl)urea, 1-(3-aminophenyl)urea, 1-naphthol, 2-methyl-1-naphthol, 1,5-naphthalenediol, 2,7-naphthalenediol or 1-acetoxy-2-methylnaphthalene, 4-chloro-2-methylnaphthalen-1-ol, 4-methoxy-2-methylnaphthalen-1-ol, 2,6-dihydroxy-3,4-dimethylpyridine, 2,6-dimethoxy-3,5-pyridinediamine, 3-amino-2-methylamino-6-methoxypyridine, 2-amino-3-hydroxypyridine, 2,6-diaminopyridine, pyridine-2,6-diol, 5,6-dihydroxyindole, 6-hydroxyindole, 5,6-dihydroxyindoline, 3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one, 1,2,4-trihydroxybenzene, 2-(benzo[d][1,3]dioxol-5-ylamino)ethanol (also known as hydroxyethyl-3,4-methylenedioxy aniline), and mixtures thereof.

When the composition is obtained by mixing a tint component and a developer component, additional primary intermediates and couplers may be preferably incorporated in the tint component.

Typically, the compositions for the oxidative dyeing of keratin fibers comprise from about 0.1% to about 10%, preferably from about 0.3% to about 6%, more preferably from about 0.5% to about 4% by weight of the couplers relative to the total weight of the composition.

Direct Dyes

The compositions of the present invention may also comprise compatible direct dyes, in an amount sufficient to provide additional coloring, particularly with regard to intensity. Typically, such an amount will range from about 0.05% to about 4%, by weight of the direct dyes relative to the total weight of the composition. When the composition is obtained by mixing a tint component and a developer component, the direct dyes are usually incorporated in the tint component.

The following direct dyes are commonly used: Acid dyes such as Acid Yellow 1, Acid Orange 3, Acid Black 1, Acid Black 52, Acid Orange 7, Acid Red 33, Acid Yellow 23, Acid Blue 9, Acid Violet 43, HC Blue 16, Acid Blue 62, Acid Blue 25, Acid Red 4, Basic Dyes such as Basic Brown 17, Basic Red 118, Basic Orange 69, Basic Red 76, Basic Brown 16, Basic Yellow 57, Basic Violet 14, Basic Blue 7, Basic Blue 26, Basic Red 2, Basic Blue 99, Basic Yellow 29, Basic Red 51, Basic Orange 31, Basic Yellow 87, 4-(3-(4-amino-9,10-dioxo-9,10-dihydroanthracen-1-ylamino)propyl)-4-methylmorpholin-4-ium-methylsulfate, (E)-1-(2-(4-(4,5-dimethylthiazol-2-yl)diazenyl)phenyl)(ethylamino)ethyl)-3-methyl-1H-imidazol-3-ium chloride, (E)-4-(2-(4-(dimethylamino)phenyl)diazenyl)-1-methyl-1H-imidazol-3-ium-3-yl)butane-1-sulfonate, (E)-4-(4-(2-methyl-2-phenylhydrazono)methyl)pyridinium-1-yl)butane-1-sulfonate, N,N-dimethyl-3-(4-(methylamino)-9,10-dioxo-4a,9,9a,10-tetrahydroanthracen-1-ylamino)-N-propylpropan-1-aminium bromide, Disperse Dyes such as Disperse Red 17, Disperse Violet 1, Disperse Red 15, Disperse Violet 1, Disperse Black 9, Disperse Blue 3, Disperse Blue 23, Disperse Blue 377, Nitro Dyes such as 1-(2-(4-nitrophenylamino)ethyl)urea, 2-(4-methyl-2-nitrophenylamino)ethanol, 4-nitrobenzene-1,2-diamine, 2-nitrobenzene-1,4-diamine, Picramic acid, HC Red No. 13, 2,2'-(2-nitro-1,4-phenylene)bis(azanediyl)diethanol, HC Yellow No. 5, HC Red No. 7, HC Blue No.2, HC Yellow No. 4, HC Yellow No. 2, HC Orange No. 1, HC Red No. 1, 2-(4-amino-2-chloro-5-nitrophenylamino)ethanol, HC Red No. 3, 4-amino-3-nitrophenol, 4-(2-hydroxyethylamino)-3-nitrophenol, 2-amino-3-nitrophenol, 2-(3-(methylamino)-4-

nitrophenoxy)ethanol, 3-(3-amino-4-nitrophenyl)propane-1,2-diol, HC Yellow No. 11, HC Violet No. 1, HC Orange No. 2, HC Orange No. 3, HC Yellow No. 9, HC Red No. 10, HC Red No. 11, 2-(2-hydroxyethylamino)-4,6-dinitrophenol, HC Blue No. 12, HC Yellow No. 6, HC Yellow No. 12, HC Blue No. 10, HC Yellow No. 7, HC Yellow No. 10, HC Blue No. 9, 2-chloro-6-
5 (ethylamino)-4-nitrophenol, 6-nitropyridine-2,5-diamine, HC Violet No. 2, 2-amino-6-chloro-4-nitrophenol, 4-(3-hydroxypropylamino)-3-nitrophenol, HC Yellow No. 13, 6-nitro-1,2,3,4-tetrahydroquinoxaline, HC Red No. 14, HC Yellow No. 15, HC Yellow No. 14, N2-methyl-6-nitropyridine-2,5-diamine, N1-allyl-2-nitrobenzene-1,4-diamine, HC Red No. 8, HC Green No.1, HC Blue No. 14, and Natural dyes such as Annato, Anthocyanin, Beetroot, Carotene, Capsanthin,
10 Lycopene, Chlorophyll, Henna, Indigo, Cochineal.

Thickeners

The hair dyeing compositions of the present invention may comprise a thickener in an amount sufficient to provide the composition with a viscosity so that it can be readily applied to the hair without unduly dripping off the hair and causing mess. Typically, such an amount will be
15 at least 0.05%, preferably at least 0.5%, more preferably at least 1%, by weight of thickener relative to the total weight of the composition. When the composition is obtained by mixing several components, the thickener may be present in any of the components.

Preferred for use herein are salt tolerant thickeners, including but not limited to: xanthan, guar, hydroxypropyl guar, scleroglucan, methyl cellulose, ethyl cellulose (available as
20 AQUACOTE(TM)), hydroxyethyl cellulose (NATROSOL(TM)), carboxymethyl cellulose, hydroxypropylmethyl cellulose, microcrystalline cellulose, hydroxybutylmethyl cellulose, hydroxypropyl cellulose (available as KLUCCEL(TM)), hydroxyethyl ethyl cellulose, cetyl hydroxyethyl cellulose (available as NATROSOL(TM)) Plus 330), N-vinylpyrrolidone (available as POVIDONE(TM)), Acrylates/Ceteth-20 Itaconate Copolymer (available as
25 STRUCTURE(TM) 3001), hydroxypropyl starch phosphate (available as STRUCTURE(TM) ZEA), polyethoxylated urethanes or polycarbamyl polyglycol ester (e.g. PEG-150/Decyl/SMDI copolymer (available as ACULYN(TM) 44), PEG-150/Stearyl/SMDI copolymer available as ACULYN(TM) 46), Acrylates/Beheneth-25 Methacrylate Copolymer (available as ACULYN(TM) 28), Acrylates/Vinyl Neodecanoate Crosspolymer (available as ACULYN(TM)
30 38), Acrylates/Steareth-20 Methacrylate Crosspolymer (available as ACULYN(TM) 88), PEG-150 Distearate (available as ACULYN(TM) 60), trihydroxystearin (available as THIXCIN(TM)), acrylates copolymer (e.g. available as ACULYN(TM) 33) or hydrophobically

modified acrylate copolymers (e.g. Acrylates/Stearth-20 Methacrylate Copolymer (available as ACULYN(TM) 22), non-ionic amphiphilic polymers comprising at least one fatty chain and at least one hydrophilic unit selected from polyether urethanes comprising at least one fatty chain.

Also preferred for use herein are thickeners based on lamellar gel network systems, comprising at least one surfactant or amphiphile having an HLB of 6 or less and a melting point of at least 30 °C, preferably selected from fatty alcohols comprising from 14 to 30 carbon atoms, or oxyethylenated fatty alcohols comprising from 16 to 30 carbon atoms and 2 units or less of ethylene oxide, and further comprising at least one ionic or nonionic surfactant, preferably selected from:

- 10 - anionic surfactants selected from C8-C30 alkyl sulfates, preferably C12-C18 alkyl sulfates,
- anionic surfactants according to the formula R_nX_mYM , wherein R is independently selected from alkyl, alkenyl or alkylaryl groups having from 8 to 30 carbon atoms, X is independently selected from polar groups comprising at least one carbon atom and at least
15 one oxygen or nitrogen atom, Y is an anionic group selected from carboxylates, sulfates, sulfonates or phosphates, n and m are independently 1 or 2, and M is hydrogen or a salt forming cation and mixture thereof, most preferably selected from C8-C30 alkyl ether phosphates having from 1 to 20, preferable 2 to 10 ethylene oxide units (e.g. available as CRODAFOS(TM) CES);
- non-ionic surfactant comprising one or more polyethyleneoxide chains,
20 preferably each polyethyleneoxide chain has on average at least 50 ethylene oxide units and most preferably 100 to 200 ethylene oxide units (e.g. available as VOLPO(TM) S200),
- cationic surfactants selected from quaternary ammonium salts or amido-amines having at least one fatty chain, preferably comprising at least 16 carbon atoms and most preferably at least 20 carbon atoms,
- 25 - and mixture thereof.

Examples of such lamellar gel network systems are disclosed in EP1,832,273 and EP2,103,299.

The composition preferably comprises a mixture of ceteryl alcohol and dicetyl phosphate and ceteth-10 phosphate (e.g. available as CRODAFOS(TM) CES).

Chelants

The compositions of the present invention may comprise chelants in an amount sufficient to reduce the amount of metals available to interact with formulation components, particularly oxidizing agents, more particularly peroxides and percarbonates. Typically, such an amount range from at least 0.15%, preferably at least 0.25%, by weight of the chelants relative to the total weight of the composition. Suitable chelants for use herein include but are not limited to: diethylenetriamine-N,N',N''-polyacids, diethylenetriaminepentaacetic acid (DTPA), diethylenetriaminepenta(methylene phosphonic acid) (DTPMP), diamine-N,N'-dipolyacid, monoamine monoamide-N,N'-dipolyacid, and N,N'-bis(2-hydroxybenzyl)ethylenediarnine-N,N'-diacetic acid chelants (preferably EDDS (ethylenediaminedisuccinic acid)), carboxylic acids (preferably aminocarboxylic acids), phosphonic acids (preferably aminophosphonic acids) and polyphosphoric acids (in particular straight polyphosphoric acids), their salts and derivatives. When the composition is obtained by mixing a tint component and a developer component, the chelants may be incorporated in the tint component or in the developer component or in both. A chelant is usually present in developer components for stability reason.

pH Modifiers

The compositions of the present invention may comprise in addition to the alkalinizing agent discussed above a pH modifier and/or buffering agent in an amount that is sufficiently effective to adjust the pH of the composition to fall within a range from about 3 to about 13, preferably from about 8 to about 12, more preferably from about 9 to about 11.

Radical Scavengers

According to the present invention, the compositions may comprise a radical scavenger. As used herein the term radical scavenger refers to a species that can react with a radical, to convert the radical species by a series of fast reactions to an unreactive or less reactive species. The radical scavenger is also preferably selected such that it is not an identical species as the alkalinizing agent and is present in an amount sufficient to reduce the damage to the hair during the coloring /bleaching process. The compositions of the present invention comprise a radical scavenger from about 0.1% to about 10%, preferably from about 1% to about 7% by weight of the radical scavenger relative to the total weight of the composition. Suitable radical scavengers for use herein may be selected from the classes of alkanolamines, amino sugars, amino acids, esters of amino acids and mixtures thereof. Suitable compounds include 3-substituted-pyrazol-5-ones, 3-carboxy-1H-pyrazol-5-one, 3-methyl-1-phenyl-pyrazol-

5-one, 3-methyl-1-p-tolyl-pyrazol-5-one, 3-methyl-1-(4-sulfophenyl)-pyrazol-5-one, 3-methyl-1-(4-sulfoamidophenyl)-pyrazol-5-one, 3-methyl-1-(3-sulfophenyl)-pyrazol-5-one, 3-methyl-1-(3-sulfoamidophenyl)-pyrazol-5-one, 3-methyl-1-(2-chloro-5-sulfophenyl)-pyrazol-5-one, 3-methyl-1-(2,5-dichloro-4-sulfophenyl)-pyrazol-5-one, 3-methyl-1-(4-chlorophenyl)-pyrazol-5-one, 3-methyl-1-(4-carboxyphenyl)-pyrazol-5-one, 3-carboxy-1-phenyl-pyrazol-5-one, 3-carboxy-1-(4-sulfophenyl)-pyrazol-5-one, 1,3-diphenyl-pyrazol-5-one, methyl pyrazol-5-one-3-carboxylate, 3-amino-1-propanol, 4-amino-1-butanol, 5-amino-1-pentanol, 1-amino-2-propanol, 1-amino-2-butanol, 1-amino-2-pentanol, 1-amino-3-pentanol, 1-amino-4-pentanol, 3-amino-2-methylpropan-1-ol, 1-amino-2-methylpropan-2-ol, 3-aminopropane-1,2-diol, glucosamine, N-acetylglucosamine, glycine, arginine, lysine, proline, glutamine, histidine, sarcosine, serine, glutamic acid, tryptophan, or mixtures thereof, or the salts, such as the potassium, sodium, or ammonium salts thereof, or mixtures thereof. In some embodiments, the inventive compositions may comprise glycine, sarcosine, lysine, serine, 2-methoxyethylamine, glucosamine, glutamic acid, morpholine, piperidine, ethylamine, 3-amino-1-propanol, or mixtures thereof.

15 **Method of hair dyeing**

In order to use the dyeing composition, the tint component and developer components are usually mixed immediately prior to use and a sufficient amount of the mixture is applied to the hair, according to the hair abundance, generally from about 60 to about 250 grams. Upon such preparation the composition is applied to the hair to be dyed and remains in contact with the hair for an amount of time effective to dye the hair. Typically, the hair dye composition is allowed to act on the hair from about 2 to about 60, preferably about 15 to about 45, more preferably about 30 minutes, at a temperature ranging from 15 °C to about 50 °C. Thereafter, the hair is rinsed with water to remove the composition and dried. If necessary, the hair is washed with a shampoo and rinsed, e.g., with water or a weakly acidic solution, such as a citric acid or tartaric acid solution, and dried. Optionally, a separate conditioning product may also be provided.

The method of dyeing hair with the composition may therefore comprise the steps of:

(i) providing a tint component comprising (a) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I) as defined in claim 1, its physiologically compatible, water-soluble salt, or mixtures thereof and, (b) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II) as defined in claim 1, its physiologically compatible water-soluble salt, or mixtures thereof;

(ii) providing a developer component comprising (c) an oxidizing agent;

(iii) mixing the tint component and the developer component to obtain a composition for the oxidative dyeing of keratin fibers according to the composition of the invention;

(iv) applying said composition for the oxidative dyeing of keratin fibers onto the hair.

The method may further comprise waiting a period of time, typically between 2 minutes
5 and 60 minutes, and then rinsing the composition from the hair.

The compositions can be applied on hair via applicator bottle or brush. It can be used on full head or partly on single strands (highlight application) as common highlight applicator foils, caps and special applicators can be used, but also freehand techniques such as balayage, with brush and/or combs can be possible. The composition can also be applied as a mousse via a
10 manual spray, a pressurized container or an aerosol mousse.

The composition may be dispensed as a solid form to which water is added to generate the oxidant and form a thickened vehicle suitable for hair coloring.

The dye combination of the invention may also be used in three components system. See for example disclosed US2010/0223739A2 assigned to L'Oreal. Such a process and kit for
15 lightening or dyeing keratin fibers may comprise the following composition applied to the hair fibers: an aqueous cosmetic composition (A) comprising at least one fatty substance and at least one surfactant; a cosmetic composition (B) comprising at least one alkaline agent and the oxidative dyes of the invention and if present direct dyes and other oxidative dyes, a cosmetic composition (C) comprising at least one oxidizing agent, wherein the amount of the at least one
20 fatty substance in composition (A) is greater than 20% by weight relative to the total weight of composition (A).

Methods of making - Kit

The composition, and its tint component and developer component, may be manufactured by conventional processes known in the art for manufacturing oxidative dyeing products, and ad-
25 mixing the ingredients of each component composition in suitable vessels, followed by packaging in appropriate individual containers. The components may be for example packaged in plastic or aluminium bottles.

In particular, the present invention may be provided as a kit comprising different components to be mixed by the consumer or salon stylist to obtain a hair dyeing composition
30 according to the invention. Such a kit may comprise:

(i) a tint component comprising (a) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I) as defined above, its physiologically compatible water-soluble salt, or mixtures thereof, and (b) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II) as defined above, its physiologically compatible water-soluble salt, or mixtures thereof, and

5 ii) a developer component comprising an oxidizing agent.

The kit may be presented in a single package comprising separate containers for the tint component, the developer component, and optionally a conditioner, a color refresher or other hair treatment product, instructions for use, gloves. The instructions for use include the steps of the method described above and optionally provide visual cues or pictures for the desired steps of the method. Kits are usually sold in retail products with enough material in each component for preparing a hair dyeing composition for one use.

The composition may be dispensed as a foam using for example manually-actuable, non-aerosol dispenser such as a pump or squeeze foamer, aerosol mousse. See for example EP 613,728 B1, WO 97/013585 A1, EP 1,716,933A1, US 3,709,437, US 3,937,364, US 4,022,351, 15 US 4,147,306, US 4,184,615, US 4,615,467 and FR 2,604,622. One particular example of a squeeze foamer useful herein is able to dispense from an upright or inverted position such as the one discussed in US 6,604,693 assigned to Taplast, and more specifically, at column 2, line 65, through column 4, line 67 of that patent.

The composition may be dispensed as a solid form to which water is added to generate the oxidant and form a thickened vehicle suitable for hair coloring.

EXAMPLES

The following are non-limiting examples of the compositions of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention, which would be recognized by one of ordinary skill in the art. All concentrations are listed as weight percent, unless otherwise specified.

The compositions tested were formulated by mixing a tint component according to the formulation below with a developer component in a 1:1 ratio. The developer component is a commercially available Welloxon developer at 6% (20 vol) peroxide concentration. In all the tests, white hair was used (1.5 g/ tress). 3 g tint component + 3 g developer component were

applied for each tress for 30 minutes at 30 °C. The tresses were rinsed for 2 minutes including 30 second shampooing.


Table 1: Tint Component Formulation

Ingredients	Weight%
Lanolin alcohol	2.0
Glyceryl Stearate	5.5
Glycol Distearate	2.0
Cetearyl alcohol and SLS	17.5
Sodium cocoyl isethionate	0.46
Sodium laureth sulfate	4.0
Ascorbic acid	0.3
Sodium sulfate	0.001
Sodium sulfite	0.4
Disodium EDTA	0.1
Alkyl Pyrazole Salt	X
Benzo[1,3]dioxol-5-ylamine	Y
Ammonia, 28.3% in water	6.43
Fragrance	0.25
Reverse osmosis water	QS to 100% Total Wt

X and Y were calculated to arrive at concentration for each compound of 0.050 M in the tint component formulation, except for the Towel Bleeding test where the molar concentration was raised to 0.070M.

Color intensity

The tint component indicated above was formulated with alkyldiaminopyrazoles having varying alkyl chain length and benzo[1,3]dioxol-5-ylamine of formula (II.1):



10

(II.1)

After mixing with the developer component, the resulting mixture was applied to white hair and rinsed. The resulting color was measured using a colorimeter and characterized by the L* value. An L* of 100 is considered white and L* of 0 is considered black, therefore the higher the L* value the lower the color intensity. The measured L* values for the different primary dye combinations are summarized in Table 2 below.

15

Table 2: Comparison of L* values




	L*
C5 	23.27
C6 	24.99
C8 	33.68

Table 2 demonstrates that the L* values of hair tresses dyed with the C8 alkyldiaminopyrazole combined with benzo[1,3]dioxol-5-ylamine are significantly higher compared to the other alkyldiaminopyrazoles with the benzo[1,3]dioxol-5-ylamine, correlating with weaker color. As the C8 alkyldiaminopyrazole with the selected couplers clearly does not give the desired color intensity, it will not be further analyzed or compared to the present invention.


Towel Bleeding

A consumer relevant measure of the ability to see differences in washfastness is to assess how much color from wet dyed hair tresses transfers to a white cloth, such as a white wash cloth. The amounts of towel bleeding resulting from washed tresses dyed with a C5 alkyldiaminopyrazole and C6 alkyldiaminopyrazole with compound of formula (II.1) were measured using conventional ΔE^* measurements. The testing method is summarized below:

After washing a dyed hair switch by the method described above, place the switch on a white cloth (50% cotton, 50% polyester) with an average color reading of $L^*=95.8$, $a^*=-1.40$ and $b^*=0.89$. Fold the cloth in half over the hair switches. Place a 2.27 kg weight on the folded

cloth for 5 min. Upon removal of the hair, visual inspection shows a larger and more intense stain for the C5 alkyldiaminopyrazole compared to the C6 alkyldiaminopyrazole compound. The noticeability of the stain on the fabric was calculated using the ΔE measurements for C5 and C6 relative to the fabric. (see ΔE^* values below). A higher ΔE^* value shows more color transfer to the cloth (bleeding).

Table 3: Total change in cloth color (ΔE^*) due to dye transfer from wet tresses dyed with C5 and C6 alkyldiaminopyrazoles.

+	C5	C6
 (II.1)	$\Delta E = 16.63$	$\Delta E = 6.76$

It was found that the C6 alkyldiaminopyrazole dye had a stain intensity 40% less noticeable than the C5 alkyldiaminopyrazole dye after one wash cycle.

10 Additional example

The color intensity and the towel bleeding test as indicated above were conducted under the same conditions with a tint component comprising 2,5-toluenediamine sulfate (DTS) as a further primary intermediate as indicated in Table 4 below.

Table 4: Tint Component Formulation





Ingredients	Weight %
Lanolin alcohol	2.0
Glyceryl stearate	5.5
Glycol Distearate	2.0
Cetearyl alcohol and SLS (9:1)	17.5
Sodium cocoyl isethionate	0.46
Sodium laureth sulfate	4.0
Ascorbic acid	0.3
Sodium sulfate	0.001
Sodium sulfite	0.4
Disodium EDTA	0.1
C5 pyrazole salt	0.869
Or	Or
C6 pyrazole salt	0.925
Or	Or
C8 Pyrazole salt	1.042
Benzo[1,3]dioxol-5-ylamine	1.088

2,5-toluenediamine sulfate	0.22
Ammonia, 28.3% in water	6.43
Fragrance	0.25
Reverse osmosis water	QS to 100% Total Wt

The amount of pyrazole varies to achieve the same molar concentration of about 0.050M.

The Color intensity results are as indicated in Table 5, the Towel Bleeding results as in Table 6.

Table 5: Comparison of L* values with DTS


	L* values
	 (II.1)
C5 	21.3
C6 	22.69
C8 	28.17

5

Table 5 demonstrates that the L* values of hair tresses dyed with the C8 alkyldiaminopyrazole combined with 2,5-toluenediamine and benzo[1,3]dioxol-5-ylamine are significantly higher compared to the other alkyldiaminopyrazoles with 2,5-toluenediamine and, correlating with weaker color. As the C8 alkyldiaminopyrazole with 2,5-toluenediamine and benzo[1,3]dioxol-5-ylamine clearly does not give the desired color intensity, it will not be further analyzed or compared to the present invention.

10

Table 6: Total change in cloth color (ΔE).

	C5	C6
 <p>(II.1)</p> <p>$\Delta E = 15.62$ $\Delta E = 5.50$</p>		


Concluding remark

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm."

CLAIMS

What is claimed is:

1. A composition for the oxidative dyeing of keratin fibers, in particular human keratin fibers, comprising:
- 5 - (A) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I), a physiologically compatible water-soluble salt thereof, or mixtures thereof;



wherein a is equal to one or two;

10 wherein R_i is selected from the group consisting of:

(a) C-linked substituents selected from the group consisting of:

(i) substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic substituents, in particular alkyl or hydroxy substituted alkyl,

15 (ii) substituted or unsubstituted, mono- or poly- unsaturated aromatic or heteroaromatic substituents, in particular aryl or heteroaryl,

wherein said C-linked substituents comprise from 1 to 6 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, F, N, P and Si;

20 (b) S-linked substituents selected from the group consisting of SA^1 , SO_2A^1 , SO_3A^1 , SSA^1 , SOA^1 , $SOzNA^2$, SNA^1A^2 , and $SONA^2$;

(c) O-linked substituents selected from the group consisting of OA^1 , ONA^2 ;

(d) N-linked substituents selected from the group consisting of NA^2 ; $(NA^1A^2A^3)^+$, NA^2A^2 , NO_2 ; NA^*A^2 ;

(e) halogens selected from the group consisting of F, Cl, Br, and I; and


25 (f) hydrogen;

wherein A^1 , A^2 , and A^3 are selected independently of each other from the group consisting of: hydrogen; substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated,

aliphatic or heteroaliphatic or aromatic or heteroaromatic substituents, preferably substituted or unsubstituted alkyl, or A¹ and A² together with nitrogen atoms to which they are bound form a ring; wherein said substituents or ring comprise from 1 to 6 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, S, N, P, and Si;

5 R₂ and R₃ are selected independently of each other from the group consisting of a hydrogen atom; a C1-C6 alkyl substituent; a trifluoromethyl substituent; a C1-C6 aminoalkyl substituent; a C1-C6 hydroxyalkyl substituent; C1-C6 an alkoxyalkyl substituent;

- (B) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II), a physiologically compatible water-soluble salt thereof, or mixtures thereof;



10

wherein R₄, R₅, R₆, R₇ are substituents selected independently of each other from the group consisting of:

(a) C-linked substituents selected from the group consisting of:

15 (i) substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic substituents, in particular alkyl or hydroxy substituted alkyl

(ii) substituted or unsubstituted, mono- or poly- unsaturated aromatic or heteroaromatic substituents, in particular aryl or heteroaryl,

20 wherein said C-linked substituents comprise from 1 to 10 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, F, N, P and Si;

(b) S-linked substituents selected from the group consisting of SA¹, SO₂A¹, SO₃A¹, SSA¹, SOA¹, SOzNA², SNA², and SONA²;

(c) O-linked substituents selected from the group consisting of OA¹, ONA²;

25 (d) N-linked substituents selected from the group consisting of NA¹A²; (NA¹A²A³)⁺, NA², NO₂; NA²;

(e) halogens selected from the group consisting of F, Cl, Br, and I; and


(f) hydrogen;

wherein A¹, A², and A³ are selected independently of each other from the group consisting of: hydrogen; substituted or unsubstituted, straight or branched or cyclic, saturated or unsaturated, aliphatic or heteroaliphatic or aromatic or heteroaromatic substituents, preferably substituted or unsubstituted alkyl, or A¹ and A² together with nitrogen atoms to which they are bound form a ring; wherein said substituents or ring comprise from 1 to 10 carbon atoms and from 0 to 5 heteroatoms selected from the group consisting of O, S, N, P, and Si;


and

- (C) an oxidizing agent.

2. A composition according to claim 1, wherein the 1-hexyl/heptyl-4,5-diaminopyrazole compound (I) is a compound with formula (1.1):




3. A composition according to claim 1 or 2 wherein the benzo[1,3]dioxol-5-ylamine derivative of formula (II) is selected from benzo[d][1,3]dioxol-5-amine, 2-(benzo[d][1,3]dioxol-5-ylamino)ethanol, 3-(benzo[d][1,3]dioxol-5-ylamino)propan-1-ol, 4-(benzo[d][1,3]dioxol-5-ylamino)butan-1-ol, 5-(benzo[d][1,3]dioxol-5-ylamino)pentan-1-ol, 6-(benzo[d][1,3]dioxol-5-ylamino)hexan-1-ol, N-pentylbenzo[d][1,3]dioxol-5-amine, N-ethylbenzo[d][1,3]dioxol-5-amine, N-propylbenzo[d][1,3]dioxol-5-amine, N-butylbenzo[d][1,3]dioxol-5-amine, N-pentylbenzo[d][1,3]dioxol-5-amine, N-hexylbenzo[d][1,3]dioxol-5-amine, aminol as represented in formula (II.1), salts thereof and mixtures thereof, preferably aminol:



4. A composition according to any of the preceding claims, further comprising a primary intermediate selected from toluene-2,5-diamine, p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, 2-methoxymethyl-p-phenylenediamine, 2-(2,5-diamino-4-

methoxyphenyl)propane-1,3-diol, 2-chloro-p-phenylenediamine, 2-methoxy-p-phenylenediamine, 2-((2-(4-amino-phenylamino)-ethyl)-(2-hydroxyethyl)-amino)-ethanol, salts thereof and combination thereof.

5. A composition according to any of the preceding claims, further comprising an auxiliary coupler selected from the compounds having formula (III.1) to (III.8) below and combinations thereof:



6. A composition according to any one of the preceding claims further comprising at least one additional component selected from the group consisting of alkalizing agents, auxiliary primary intermediates, auxiliary couplers, direct dyes, thickeners, chelants, pH modifiers and/or buffering agents, radical scavenger systems and mixtures thereof.
7. A composition according to any of the preceding claims, further comprising fatty alcohols comprising from 14 to 30 carbon atoms, or oxyethylenated fatty alcohols comprising from 16 to 30 carbon atoms and 2 units or less of ethylene oxide.
8. A composition according to claim 7, further comprising at least one ionic or nonionic surfactant selected from:
- anionic surfactants selected from C8-C30 alkyl sulfates, preferably C12-C18 alkyl sulfates,
 - anionic surfactants according to the formula R_nX_mYM , wherein R is independently selected from alkyl, alkenyl or alkylaryl groups having from 8 to 30 carbon atoms, X is independently selected from polar groups comprising at least one carbon atom and at least one oxygen or nitrogen atom, Y is an anionic group selected from carboxylates, sulfates,

sulfonates or phosphates, n and m are independently 1 or 2, and M is hydrogen or a salt forming cation and mixture thereof, most preferably selected from C8-C30 alkyl ether phosphates having from 1 to 20, preferably 2 to 10 ethylene oxide units;

5 - non-ionic surfactant comprising one or more polyethyleneoxide chains, preferably each polyethyleneoxide chain has on average at least 50 ethylene oxide units and most preferably 100 to 200 ethylene oxide units;

- cationic surfactants selected from quaternary ammonium salts or amido-amines having at least one fatty chain, preferably comprising at least 16 carbon atoms and most preferably at least 20 carbon atoms; and

10 - mixtures thereof.

9. A composition according to claim 8 comprising a mixture of cetearyl alcohol and dicetyl phosphate and ceteth-10 phosphate.

10. A composition according to any of the preceding claims further comprising a chelant selected from the group consisting of: diethylenetriamine-N,N',N''-polyacids, diethylenetriaminepentaacetic acid (DTPA), diethylenetriaminepenta(methylene phosphonic acid) (DTPMP), diamine-N,N'-dipolyacid, monoamine monoamide-N,N'-dipolyacid, and N,N'-bis(2-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid chelants (preferably EDDS (ethylenediaminedisuccinic acid)), carboxylic acids (preferably aminocarboxylic acids), phosphonic acids (preferably aminophosphonic acids) and polyphosphoric acids (in particular straight polyphosphoric acids), their salts and derivatives .

15

20

11. A composition according to any of the preceding claims wherein said oxidizing agent is selected from the group consisting of hydrogen peroxide, sodium periodate, urea peroxide, melamine peroxide, perborates, percarbonates, perphosphates, persilicates, persulfates, oxidizing enzymes such as uricases, oxidases, and peroxidases, a source of peroxymonocarbonate ions and mixtures thereof.

25

12. A composition according to any of the preceding claims, wherein said composition is dispensed as a foam.

13. A method of dyeing hair comprising the steps of:

- (i) providing a tint component comprising (a) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I) as defined in claim 1, a physiologically compatible water-soluble salt or mixtures thereof; and (b) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II) as defined in claim 1, a physiologically compatible water-soluble salt or mixtures thereof;
- 5 (ii) providing a developer component comprising (c) an oxidizing agent;
- (iii) mixing the tint component and the developer component to obtain a composition for the oxidative dyeing of keratin fibers according to any of the preceding claims;
- (iv) applying said composition for the oxidative dyeing of keratin fibers onto the hair.
- 10 14. An oxidative hair dyeing kit comprising: (i) a tint component comprising (a) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I) as defined in claim 1, a physiologically compatible water-soluble salt thereof or mixtures thereof; and, (b) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II) as defined in claim 1, a physiologically compatible water-soluble salt thereof or mixtures thereof and
- 15 ii) a developer component comprising an oxidizing agent.
15. A tint component comprising (a) a 1-hexyl/heptyl-4,5-diaminopyrazole compound of the general formula (I) as defined in claim 1, a physiologically compatible, a physiologically compatible water-soluble salt thereof or mixtures thereof and; (b) a benzo[1,3]dioxol-5-ylamine compound of the general formula (II) as defined in claim 1, a physiologically
- 20 compatible water-soluble salt thereof or mixtures thereof.