



- (51) **International Patent Classification:** Not classified
- (74) **Agents:** SUBRAMANIAM, Hariharan et al.; Subramaniam, Nataraj & Associates, E-556, Greater Kailash-II, New Delhi - 110 048 (IN).
- (21) **International Application Number:** PCT/IN2012/000148
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (22) **International Filing Date:** 2 March 2012 (02.03.2012)
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
590/MUM/2011 3 March 2011 (03.03.2011) IN
1948/MUM/2011 6 July 2011 (06.07.2011) IN
2777/MUM/2011 30 September 2011 (30.09.2011) IN
- (71) **Applicant** (for all designated States except US): CADILA HEALTHCARE LIMITED [IN/IN]; Zydus Tower, Satellite Cross Roads, Ahmedabad - 380 015, Gujarat (IN).
- (72) **Inventors; and**
- (75) **Inventors/Applicants** (for US only): PANDEY, Bipin [IN/IN]; CADILA HEALTHCARE LIMITED, Zydus Tower, Satellite Cross Roads, Ahmedabad - 380 015, Gujarat (IN). DAVE, Mayank Ghanshyambhai [IN/IN]; CADILA HEALTHCARE LIMITED, Zydus Tower, Satellite Cross Roads, Ahmedabad - 380 015, Gujarat (IN). KOTHARI, Himanshu M. [IN/IN]; CADILA HEALTHCARE LIMITED, Zydus Tower, Satellite Cross Roads, Ahmedabad - 380 015, Gujarat (IN). SHUKLA, Bhavin Shriprasad [IN/IN]; CADILA HEALTHCARE LIMITED, Zydus Tower, Satellite Cross Roads, Ahmedabad - 380 015, Gujarat (IN).
- Declarations under Rule 4.17:**
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- Published:**
- without international search report and to be republished upon receipt of that report (Rule 48.2(g))



(54) **Title:** NOVEL SALTS OF DPP-IV INHIBITOR

(57) **Abstract:** The present invention discloses new salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine i.e. gentsiate, adipate and trifluoro acetic acid salts. The invention also describes the new, crystalline and amorphous forms of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentsiate. The present invention also discloses novel crystalline salt form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride and novel crystalline and amorphous salt forms of besylate and process for their preparation and isolation.

NOVEL SALTS OF DPP-IV INHIBITOR

FIELD OF INVENTION


The present invention discloses certain new salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as well as their processes of preparation.

The present invention also discloses certain novel crystalline forms of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride and novel crystalline and amorphous forms of the besylate salt and processes for their preparation and isolation.

BACKGROUND OF THE ART

The invention is related to novel salts and certain novel crystalline and amorphous forms of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine salts.

WO 03/004498 and US patent No. 6,699,871 both assigned to Merck and Co., describes a class of beta-amino tetrahydrotriazolo[4,3-a]pyrazines, which are inhibitors of DPP-IV. Disclosed therein are compounds, whose general formula is,



Specifically disclosed in WO 03/004498 is (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro [1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine. WO 03/004498 is silent as to the preparation of and the nature of specific crystal forms of the salts. WO 2005/003135 describes dihydrogenphosphate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and crystalline hydrates thereof, in particular a crystalline monohydrate.

In WO 2005/003135 it is said that (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine dihydrogenphosphate salt and crystalline hydrates have advantages in the preparation of pharmaceutical compositions, such as ease of processing, handling, and dosing. In particular, they exhibit improved physical and chemical stability, such as stability to stress, high temperatures and humidity, as well as improved physicochemical

properties, such as solubility and rate of dissolution. WO2005/020920 describes the crystalline anhydrate Form I, Form II and Form III as well as solvates of the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine dihydrogenphosphate salt.

5 WO2005/030127 describes novel crystalline anhydrate Form IV of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine dihydrogenphosphate salt.

WO2006/033848 describes the amorphous form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine dihydrogenphosphate salt.

WO2005/072530 describes crystalline hydrochloric acid, benzenesulfonic acid, p-toluenesulfonic acid, 10-camphorsulfonic acid, and tartaric acid salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and hydrates thereof.

15 WO2007/035198 describes (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine dodecylsulfate salt, in particular, a crystalline anhydrate form thereof.

In addition, the hemifumarate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine has been described by D. Kim et al. in J. Med. Chem. 2005, 48, 141-151.

Furthermore, International Patent Publication No. WO 2008/000418 discloses the preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride in amorphous form.

25 In addition, International Patent Publication No. WO 2009/085990 describes other acid addition salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, including salts of di-p-tolyl-L-tartaric acid, phosphoric acid, sulfuric acid, hydrobromic acid, methanesulfonic acid, acetic acid, benzoic acid, oxalic acid, succinic acid, mandelic acid, fumaric acid, and lactic acid.

30 Patent Publication No. WO 2010/00469 describes different crystalline form of salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine such as hydrochloride, succinate, lactate, maleate, citrate and mesylate.

International Patent Publication No. WO 2010/092090 describes other acid addition salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, i.e. salts of D-glucuronic acid, L-glucuronic acid, glutaric acid, lactic acid, L-mandelic acid, D-mandelic acid and sulfuric acid.

International Patent Publication No. WO 2010/012781 describes other acid addition salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, i.e., salts of ethanedisulfonic acid, galactaric acid, thiocynic acid, and glutaric acid.

International Patent Publication No. WO 2010/117738 disclosed different crystalline forms of acetate, oxalate and fumarate of compound of formula I.

International Patent Publication No. WO 2012/007455 disclosed orotate salt of compound of formula I.

Different salts and new crystalline forms of the same pharmaceutically active moiety differ in their physical properties such as melting point, solubility, etc. These properties may appreciably influence pharmaceutical properties such as dissolution rate and bioavailability which in turn may have an impact on their efficacy. Thus there is a continuing need to obtain new salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine having improved physical and/or chemical properties. The present invention satisfies this need by providing new salts of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

In view of the foregoing, it would be desirable to provide new crystalline forms of salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine. We herein disclose new salts i.e. (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate, (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine adipate and novel trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

We herein also disclose new crystalline salt form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-

trifluorophenyl)butan-2-amine hydrochloride and new crystalline and amorphous salt forms of besylate which have not been disclosed in prior art.

These salts and new crystalline salt forms can show certain superior pharmaceutical properties compared to (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

All of these salts may be present either in substantially crystalline or amorphous forms or may be present as partially crystalline forms.

In an embodiment, the present invention provide new salts and new crystalline salt forms of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine which can be used as an intermediate to prepare pure (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and/or pure (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine phosphate. Further, new salts and new crystalline salt forms of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine have desired physicochemical properties such as improved stability and/or solubility. This compound also shows superior efficacy over the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine from which it is derived.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a powder X-ray diffraction (XRPD) pattern of crystalline gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention.

FIG. 2 is a powder X-ray diffraction (XRPD) pattern of amorphous gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention

FIG. 3 is a powder X-ray diffraction (XRPD) pattern of crystalline adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention.

FIG. 4 is a powder X-ray diffraction (XRPD) pattern of crystalline hydrochloride salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention.

FIG. 5 is a powder X-ray diffraction (XRPD) pattern of crystalline besylate salt (I) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention

FIG. 6 is a powder X-ray diffraction (XRPD) pattern of crystalline trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention.

FIG. 7 is a powder X-ray diffraction (XRPD) pattern of crystalline besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention.

FIG. 8 is a powder X-ray diffraction (XRPD) pattern of amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine according to the present invention.

FIG. 9- Effect of single dose oral administration of ZY compounds at 10 mg/kg, PO on OGTT in male C57 mice.

FIG. 10- Effect of single dose oral administration of ZY compounds at 10mg/kg, PO on OGTT in male C57 mice.

OBJECT OF THE INVENTION

In an embodiment the present invention provides gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and processes for their preparation.

In a preferred embodiment the present invention provides a crystalline form of gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which can be characterized by its powder X-ray diffraction (PXRD) pattern and other characteristic properties as described hereinafter and process of preparation thereof.

In another preferred embodiment the present invention provides the amorphous form of the gentisate salt of 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which can be characterized by its powder X-ray diffraction (PXRD) pattern and other characteristic properties as described hereinafter and process for preparation thereof.

In an embodiment the present invention provides an adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and processes for their preparation.

In an embodiment the present invention provides a novel trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and processes for its preparation.

In an embodiment the present invention provides a new crystalline form of the hydrochloride salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which can be characterized by its powder X-ray diffraction (PXRD) pattern and other characteristic properties as described hereinafter and process for preparation thereof.

In an embodiment the present invention provides a new crystalline besylate salt(I) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which can be characterized by its powder X-ray diffraction (PXRD) pattern and other characteristic properties as described hereinafter and process for preparation thereof.

In an embodiment the present invention provides a new crystalline besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which can be characterized by its powder X-ray diffraction (PXRD) pattern and other characteristic properties as described hereinafter and process for preparation thereof.

In an embodiment the present invention provides an amorphous form of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which can be characterized by its powder X-ray diffraction (PXRD) pattern and other characteristic properties as described hereinafter and process for preparation thereof.

In a still further embodiment is provided a process for preparation of chemically and chirally pure (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine base through the intermediate formation of above disclosed novel salts and new crystalline salt forms.


DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "reflux temperature" refers to the boiling point of the solvent used.

As used herein, the term "PXRD" refers to powder X-ray diffraction.

As used herein, the term "(2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine" refers to Sitagliptin.

As used herein, the term "THF" refers to tetrahydrofuran, the term "DCM" refers to dichloro methane, the term "DIPE" refers to di-isopropyl ether, the term "MTBE" refers to methyl t-butyl ether, the term "IPA" refers to isopropyl alcohol, the term "IPAc" refers to isopropyl acetate, the term "phosphate" refers to dihydrogen phosphate salt, the term "besylate" refers to benzenesulphonic acid salt
 5 (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine is having structural formula (I).



In one of the embodiment the invention discloses novel gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.
 10

In an embodiment this salt may be present either in crystalline or amorphous form or suitable mixtures of crystalline and amorphous forms. In a further embodiment, the crystalline and/or amorphous forms may exist either in hydrated, solvated as well as non-solvated forms.
 15

Hydrates of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of crystalline (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate can be synthesized by process as mentioned in Example 17.
 20

The terms 'dissolving', 'contacting', 'slurring', 'stirring', 'mixing' are interchangeable terms and doesn't affect the scope of the present invention.

In another embodiment, the invention provides a process for the synthesis of gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine which comprises
 25

- (a) dissolving of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in a suitable solvent;
 - (b) thereafter addition of gentisic acid and reacting under suitable conditions and;
 - (c) removing the solvent under suitable conditions to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-
- 30

trifluorophenyl)butan-2-amine gentisate.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like, water and suitable mixtures of one or more of the solvents described above.

In another embodiment, the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisic acid salt of the present invention can be prepared by reacting a (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine salt of a first acid with a gentisic acid, provided that the gentisic acid is stronger and sets free the first, weaker acid.

Another aspect of the present invention provides for the new gentisate salt of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in a crystalline form.

The crystalline form of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt have a characteristic XRD peaks at about 8.0, 13.73, 16.00, 18.66, 22.61, 25.62 and $26.48^{\circ} \pm 0.2$ degrees 2θ .

The crystalline form is further characterized by an additional XPRD peaks at about 6.53, 18.02, 19.68, 20.32, 23.40, 24.73, 27.05, 27.68, $29.38^{\circ} \pm 0.2^{\circ}$ degrees 2θ .

Fig.1 illustrate the XRD of the crystalline form of 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt.

In another embodiment, the invention provides a process for the synthesis of crystalline (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt, said process comprising crystallization of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt in the presence of suitable solvent and optionally further adding suitable anti solvents and subsequently removing the solvent, when necessary.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like; hydrocarbons such as hexane, heptanes,

cyclohexane, toluene, xylene, chlorobenzene and the like and suitable mixtures of one or more of the solvents described above. Preferably a mixture of alcohol and hydrocarbon.

Another aspect of the present invention includes the new gentisate salt of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in amorphous form.

Anhydrous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of amorphous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate, can be prepared by processes as mentioned in Example 18 and 19.

Fig.2 illustrate the XRD of an amorphous form of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt.

In another embodiment, the invention provides a process for the synthesis of the amorphous form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt, said process comprising reaction of the free base with the gentisic acid in suitable solvent under suitable conditions and optionally adding suitable anti solvents and removing the solvent when necessary.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like and water and suitable mixtures of one or more of the solvents described above to prepare gentisate salt.

In another embodiment, the invention provides a process for the synthesis of amorphous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt, said process comprising dissolving or slurring of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt in the presence of suitable solvent and optionally further adding suitable anti solvents and subsequently removing the solvent, when necessary.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like; esters such as ethyl acetate, isopropyl acetate and the like;

chlorinated solvents such as chloroform, dichloromethane and the like; nitriles such as acetonitrile and the like; ethers such as diethyl ether, 1,4-dioxane, DIPE, MTBE, THF and the like and suitable mixtures of one or more of the solvents described above.

In one of the embodiment the invention discloses novel adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

In an embodiment this salt may be present either in crystalline or amorphous form or suitable mixtures of crystalline and amorphous forms. In a further embodiment, the crystalline and/or amorphous forms may exist either in hydrated, solvated as well as non-solvated forms.

Hydrates of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine adipate, can be synthesized by process as mentioned in Example 17.

In another embodiment, the invention provides a process for the synthesis of the adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

Adipate salts may be prepared by reaction of the free base (formula I) under suitable reaction conditions with the adipic acid in water or one or more suitable organic solvents or in a mixture of water and one or more suitable organic solvents and removing the solvent when necessary.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like and suitable mixtures thereof.

Furthermore, the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine adipate salts of the present invention can be prepared by reacting a (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine salt of a first acid with adipic acid, provided that the adipic acid is stronger and sets free the first, weaker acid.

The crystalline form of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

adipate salt is having a characteristic XRD peaks at about 8.05, 17.64, 18.29, 21.65, 23.38, 23.85 and 24.38 ± 0.2 degrees 2θ .

The crystalline form is further characterized by an additional XPRD peaks at about 5.98, 10.03, 13.78, 15.72, 16.15, 22.34, 22.85, 25.09, 25.45 and $29.43 \pm 0.2^\circ$ degrees 2θ .

Fig.3 illustrate the XRD of a crystalline form of 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine adipate salt.

In one of the embodiments, the invention discloses novel trifluoro acetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

In an embodiment this salt may be present either in crystalline or amorphous form or suitable mixtures of crystalline and amorphous forms. In a further embodiment, the crystalline and/or amorphous forms may exist either in hydrated, solvated as well as non-solvated forms.

Anhydrous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoro acetic acid salt, can be synthesized by processes as mentioned in Examples 18 and 19 of this specification.

In another embodiment, the invention provides a process for the synthesis of trifluoro acetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

Trifluoro acetic acid salts can be prepared by reaction of the free base (formula I) under suitable reaction conditions with the trifluoro acetic acid in water or one or more suitable organic solvents or in a mixture of water and one or more suitable organic solvents or mixture of suitable solvents and subsequently removing water or the solvent used, under suitable conditions when necessary.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like and suitable mixtures of one or more of the solvents described above. In another embodiment, the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-

trifluorophenyl)butan-2-amine trifluoro acetic acid salt of the present invention can be prepared by reacting a (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine salt of a first acid with a trifluoro acetic acid, provided that the trifluoro acetic acid is stronger and sets free the first, weaker acid.

Suitable weaker acid salt of first acid is selected from HCl, acetic acid.

Another aspect of the present invention provides for the new trifluoroacetic acid salt of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in a crystalline form.

The crystalline form of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoroacetic acid salt have a characteristic XRD peaks at about 6.64, 9.39, 10.99, 13.40, 15.32, 17.15, 19.20, 19.66, 21.48, 22.46, 24.68, 26.89, 27.86, 29.34, 31.24, 32.77 and $34.82^{\circ} \pm 0.2$ degrees 2θ .

Fig.6 illustrate the XRD of crystalline form of 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoroacetic acid salt.

Another aspect of the invention relates to a process for the preparation of crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoroacetic acid salt, said process comprising crystallization of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoro acetic acid salt in the presence of suitable solvent and optionally further adding suitable anti solvents and subsequently removing the solvent, when necessary.

The suitable solvents used may be selected from alcohols such as methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol and the like; esters such as ethyl acetate, isopropyl acetate and the like; ethers such as diethyl ether, 1,4-dioxane, DIPE, MTBE, THF and the like and suitable mixtures of one or more of the solvents described above.

In an embodiment is provided a novel crystalline hydrochloride salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

Another aspect of the invention relates to a process for the preparation of a new crystalline hydrochloride salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

5 Anhydrous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride salt, can be synthesized by processes as mentioned in Example 18 and 19.

10 A new crystalline hydrochloride salt can be prepared by reaction of the free base (formula I) under suitable reaction conditions with the hydrochloric acid in water or one or more suitable organic solvents or in a mixture of water and one or more suitable organic solvents and removing the solvent under suitable conditions when necessary.

The suitable solvents used may be selected from alcohols such as methanol, 15 ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol, hexanol, heptanol, octanol, decanol and the like; ethers such as diethyl ether, 1,4-dioxane, DIPE, MTBE, THF and the like and suitable mixtures of one or more of the solvents described above.

20 The crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride salt have a characteristic XRD peaks at about 6.50, 8.05, 13.79, , 18.05, 19.52, 22.65, 23.40, 25.39 and $27.09^{\circ} \pm 0.2$ degrees 2θ .

The crystalline form is further characterized by an additional XPRD peaks at about 16.03, 18.71, 22.36, 24.79, $26.49 \pm 0.2^{\circ}$ degrees 2θ .

25 Fig.4 illustrate the XRD of crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride salt.

In one of the embodiment the invention discloses a novel crystalline form of besylate salt(I) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3- 30 a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

Anhydrous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-

a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt(I), can be synthesized by processes as mentioned in Example 18 and 19.

Another aspect of the invention relates to a process for the preparation of a novel crystalline form of besylate salt(I) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

A novel crystalline form of besylate salt(I) can be prepared by reaction of the free base (formula I) under suitable reaction conditions with the benzenesulfonic acid in one or more suitable organic solvents and removing the solvent when necessary.

The suitable solvents used may be selected from esters such as ethyl acetate, isopropyl acetate and the like; hydrocarbons such as hexane, heptanes, cyclohexane, toluene, xylene, chlorobenzene and the like and suitable mixtures of one or more of the solvents described above.

The crystalline form of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt is having a characteristic XRD peaks at about 6.81, 18.31, 22.99, 23.70 and $25.75^{\circ} \pm 0.2$ degrees 2θ .

The crystalline form is further characterized by an additional XPRD peaks at about 12.62, 17.05, 19.35, 22.13, $27.56^{\circ} \pm 0.2^{\circ}$ degrees 2θ .

Fig.5 illustrate the XRD of crystalline form of 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate(I) salt.

In an embodiment is provided a new crystalline form II of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and its hydrate.

Anhydrous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt(II), can be synthesized by processes as mentioned in Example 18 and 19.

Another aspect of the invention relates to a process for the preparation of a new crystalline form II of besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-

dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

A new crystalline besylate salt(II) can be prepared by reaction of the free base under suitable reaction conditions with the benzenesulphonic acid in water or one or more suitable organic solvents or in a mixture of water and one or more suitable organic solvents or a suitable mixture of one or more organic solvents and subsequently removing the solvent under suitable conditions as are well known, when necessary.

The suitable solvents used may be selected from a mixture of esters and water, mixture of esters, ethers and water, mixture of esters, alcohols and water, mixture of ester, hydrocarbons, water and mixture of esters, ketones and water.

The suitable alcohols used is selected from methanol, ethanol, isopropanol, butanol, 1,2-dimethoxy ethanol, 2-methoxy ethanol, 2-ethoxy ethanol, ethylene glycol, hexanol, heptanol, octanol, decanol and the like; esters used is selected from ethyl acetate, isopropyl acetate and the like; chlorinated solvents used is selected from chloroform, dichloromethane and the like; hydrocarbons used is selected from hexane, heptanes, cyclohexane, toluene, xylene, chlorobenzene and the like; ketones used is selected from acetone and the like; ethers used is selected from diethyl ether, 1,4-dioxane, DIPE, MTBE, THF and the like.

The crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt have a characteristic XRD peaks at about 5.69, 7.45, 10.08, 11.41, 13.09, 18.34, 22.52 and $22.97^{\circ} \pm 0.2$ degrees 2θ .

The crystalline form II is further characterized by an additional XPRD peaks at about 16.63, 20.82, 21.95, 23.75, 24.49, 25.02, 26.43, 27.66 and $30.19 \pm 0.2^{\circ}$ degrees 2θ .

Fig.7 illustrate the XRD of crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt.

In other embodiments the present invention further includes substantially an amorphous form of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and processes to prepare the same.

Anhydrous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine required for the

preparation amorphous form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt, can be synthesized by processes as mentioned in example 18 and 19.

Substantially amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in the present invention includes more than about 80%, more than about 90%, more than about 95%, or more than about 99%, by weight of amorphous content, the corresponding balance amount comprising of crystalline substances.

In an aspect, the present invention provides processes for the preparation of amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, which comprises:

- a) providing a solution of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in a suitable organic solvent; and
- b) recovering an amorphous form of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

The step a) may involve dissolving the active substance in a solvent that is suitable for easy, commercially viable solvent removal via distillation, etc.

A solution of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine may be obtained by dissolving besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in suitable solvent, or by reacting (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine with benzene sulphonic acid, in a suitable solvent under suitable reaction conditions. The solvent that can be used for preparing amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine may be from the various classes of solvents such as for example alcohols, halogenated hydrocarbons, water or mixtures of any two or more thereof. Alcohol solvents include for example methanol, ethanol, denatured spirits, n-propanol, isopropanol, n-butanol, isobutanol, and t-butanol and the like. Halogenated

hydrocarbons include for example dichloromethane, 1,2-dichloroethane, chloroform, carbon tetrachloride and the like.

This list is not intended to be exhaustive, and combinations of solvents that are useful can include more than one member of a class, and/or can be from different
5 classes.

These and other classes of solvents known to a person skilled in the art are all contemplated without limitation. The organic solvent acceptable for the practice of the process described herein preferably provide sufficient solubility for the active substance, and do not cause any undesirable chemical reactions with the besylate salt of
10 (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, or (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, such as degradation or side reaction, under the conditions of processing.

The recovering step b) may involve removing the solvent by distillation or by
15 filtration.

It is generally preferred that rapid drying is utilized to provide the amorphous form of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine with desired stability, moisture content and residual solvent characteristics. The resultant product
20 may be dried using any methods of drying including spray drying, rotational evaporation (such as using a Buchi Rotavapor), agitated thin film drying, spin-flash drying, fluid-bed drying, lyophilization, or other techniques known in the art.

The process may also include further drying of the product obtained from the solution by vacuum drying over a desiccant, such as phosphorous pentoxide (P₂O₅).
25 The product can also be obtained with other suitable drying agents such as potassium carbonate (K₂CO₃), sodium carbonate (Na₂CO₃), silica gel and the like, as will be apparent to the skilled artisan.

Fig. 8 illustrates the XRD of an amorphous form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-
30 trifluorophenyl)butan-2-amine besylate salt.

Analytical methods:

The complete x-ray powder spectrum, which was recorded with a Rigaku multiflex 2.0 Kilowatt X-ray powder diffractometer model using copper radiation. The X-ray diffraction pattern was recorded by keeping the instrument parameters listed below:

- i) X-ray: Cu/40kv/30mA, Diverging slit: 1°, Scattering slit: 1°, Receiving slit: 0.15 mm, Monochromator RS: 0.8 mm, Counter: Scintillation counters;
Scan mode: Continuous, Scan speed : 4.000 deg./min., Sampling width : 0.010°,
Scan axes : 2 theta vs CPS; Scan range : 4° to 40.0°, Theta offset: 0.000.
- 5 ii) Differential scanning calorimetric analysis was carried out in a DSC-60 model from Shimadzu (S/W: TA-60WS Aquisition version 2.1.0.0) by keeping following parameters,
Sample Size: Approx. 1-2mg, Sample Pans: Hermetic/Crimping Pans,
Start Temperature: 50°C, End Temperature: 300°C, Rate of Heating: 10 °C/min., Purge
10 Gas: Nitrogen, Flow rate: 20 ml/min
- iii) The infrared (IR) spectrum has been recorded on a Shimadzu FTIR-8400 model spectrophotometer, between 450 cm⁻¹ and 4000 cm⁻¹, with a resolution of 4 cm⁻¹ in a KBr pellet.
- iv) (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-
15 yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and its salts were analyzed for purity by analytical HPLC at λ_{max} 210nm using column YMC-C8 , 250 mm x 4.6mm x 4 mm or its equivalent on AGILENT 1100 series under the following conditions,
Detector: UV absorption photometer Wave length: 210 nm Column temp. : 25 °C
Flow rate: 1.0 mL/min. Injection Vol.: 10 μL
20 Mobile Phase: 10mM KH₂PO₄ (pH-6.8): Acetonitrile (55:45)
(2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]-triazolo[4,3-a]pyrazin-7(8H)-yl]-
1-(2,4,5-trifluorophenyl)butan-2-amine and its salts were analyzed for chiral purity by HPLC at λ_{max} 268 nm using column Chiral-Cel OJ-H, 250mm x 4.6mm x 5μ or its equivalent on Shimadzu LCVP model under the following conditions,
25 Detector: UV absorption photometer Wave length: 268 nm Column temp. : 35 °C
Flow rate: 0.8 mL/min. Injection Vol.: 10 μL
Mobile Phase: 0.1 % diethyl amine in [n-Hexane: Ethanol (90:10)]
- v) Melting points were taken on VEEGO make model VMP-D melting point apparatus and are uncorrected.
- 30 (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoro acetic acid salt, crystalline form II and amorphous forms of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

and crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride salt may also be useful as a therapeutic agent for treatment of certain disorders.

5 (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoro acetic acid salt, crystalline form II and amorphous forms of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride salt prepared according to the present invention has chemical and polymorphic stability on storage at $25^{\circ}\text{C} \pm 2^{\circ}\text{C} / 60\% \pm 5\% \text{RH}$ for 3 months. Results of the stability of, crystalline form II and amorphous forms of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride salt and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoroacetate are shown in table 1.

20 **Table-1: STABILITY STUDY ON STORAGE ($25^{\circ}\text{C} \pm 2^{\circ}\text{C} / 60\% \pm 5\% \text{RH}$)**

Salt	Test	Initial	After 3 Month
besylate amorphous	Description	white colored powder	white colored pow
	Purity (By HPLC)	99.41 %	99.52 %
	Water (by KFR)	1.01 %	3.14 %
	Polymorphism	XRD	amorphous pattern
besylate crystalline(II)	Description	white colored powder	Off white colore powder
	Purity (By HPLC)	99.87 %	99.97 %
	Water (by KFR)	2.77 %	2.90 %
	Polymorphism	XRD	Crystalline pattern
Hydrochloride crystalline	Description	white colored powder	white colored pow
	Purity (By HPLC)	99.82 %	99.91 %
	Water (by KFR)	3.54 %	4.01 %
	Polymorphism	XRD	Crystalline pattern

TFA salt	Description		white colored powder	white colored pov
	Purity (By HPLC)		97.89 %	97.5%
	Water (by KFR)		2.77 %	3.45 %
	Polymorphism	XRD	Crystalline pattern	No Change

The novel gentisate and adipate salt of the present application have some pharmaceutical advantages over the free base in the preparation of pharmaceutical drug product because both the novel salts are very stable.

5 It is to be noted that WO03004498 does not provide any biological activity data of the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine. (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate, the crystalline form II and amorphous forms of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine of the present invention is derived from (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine, and therefore, we are providing comparison of the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate, crystalline form II and amorphous forms of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine with crystalline monohydrate form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine phosphate salt and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

15 Comparative efficacy study of (i) (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate (ii) crystalline form of the besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (iii) amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and vs. crystalline monohydrate form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

phosphate salt and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SB):

Test material particulars:

- 5 (i) (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate (SG).
- (ii) crystalline besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SB_{II}).
- 10 (iii) amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SB_a).
- (iv) Crystalline monohydrate form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine phosphate salt (SP).
- 15 (v) (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SB):

Study Protocol:

Eight to ten weeks old male C57BLKS/J mice obtained from Animal Research Facility of Zydus Research Centre were used. The mice were allowed ad libitum access to chow and water. Animals were on a 12-hour light, 12-hour dark cycle. Protocol for use of animals for conducting this study has been reviewed and approved by Institutional Animal Ethics Committee (IAEC).

20

Oral glucose tolerance test (OGTT) was performed on these animals. The mice were deprived of food for 18 h and then drug or vehicles were administered with a glucose load (3 g/kg) by oral route. Blood (200 μ L) samples were collected at 0, 30, 60, 120 and 240 min. Serum glucose concentrations were determined using commercially available kits purchased from RFCL (New Delhi, India) using spectrophotometric assay. The glucose values were then analyzed using GraphPad Prism version 5.04 for Windows; GraphPad Software, San Diego, CA, USA. The area under the curve for 120 min was calculated, based on which the percentage inhibition was obtained for each treatment.

30

Results:

Group	% change in AUC glucose
Vehicle control	-
(SB _{II}) (10mg/kg)	-45.7±1.5
(SB) (10mg/kg)	-36.5±1.6
(SG) (10mg/kg)	-39.4±1.8
(SBa) (10mg/kg)	-43.3±1.6
(SP) (10mg/kg)	-31.9±1.8

The results indicate that crystalline besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SB_{II}) and amorphous besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SBa) are better than crystalline monohydrate form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine phosphate salt (SP) when tested in the acute oral glucose tolerance test in C57 mice.

Therefore, the present (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate, the crystalline form II and amorphous forms of besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine have potentially significant therapeutic advantages compared to the (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (SB) from which it is derived and also better than marketed crystalline monohydrate form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine phosphate salt (SP).

The invention is further exemplified by the following non-limiting examples, which are illustrative representing the preferred modes of carrying out the invention. The invention's scope is not limited to these specific embodiments only but should be read in conjunction with what is disclosed anywhere else in the specification together with those information and knowledge which are within the general understanding of a person skilled in the art.

Example 1

Preparation of adipic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

5 In a 25 mL three neck flask ethanol (10.0 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (2.0 g) were taken. It was heated to 68-70 °C and adipic acid (0.717 g) was added. The reaction mixture was stirred for 2 hrs at 78-80 °C, gradually cooled to 25-30 °C and again stirred for 2 to 3 hrs. Solid salt precipitated out.
10 The salt was filtered and washed with ethanol which provided the adipic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained (Wt.: 1.54 g, % Yield: 56.8 %, Purity by HPLC: 99.29 %, % chiral purity by HPLC of R-isomer 100.0 %), m.p.: 138-142 °C, XRD: crystalline-Fig (3)

15

Example 2

Preparation of gentisic acid salt (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine and its amorphous form.

20 In a 25 mL round bottom flask water (10 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (2.0 g) were taken. The reaction mixture was heated up to 98-100 °C to obtain clear solution. To this clear solution gentisic acid (0.756 g) was added and stirred at 98-100 °C for 1 to 2 hrs. The solvent was distilled out at reduced
25 pressure to obtain a solid mass. (Wt.: 2.7 g).

The solid was taken into DIPE (20 mL) and stirred for 1 hr then solvent decanted. Solid was then dissolved into IPA. The solvent was distilled out to obtain the gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as an amorphous solid.
30 (Wt.: 1.91g, % purity: 93.47 %, % water: 3.99% w/w, XRD=amorphous Fig (2).

Example 3

Preparation of an amorphous form of gentisic acid salt (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

In a dry, 50 mL round bottom flask methanol (20 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (4.0 g) were taken. To this clear solution gentisic acid (1.51 g) was added. The reaction mixture was heated up to 65-67 °C and stirred for 2 hrs. The solvent was distilled out at reduced pressure to obtain a solid mass. (Wt.: 5.5 g).

% water: 0.82, XRD: Amorphous Fig (2). % Purity-99.34 %, TGA-4.4 %, DSC-87.3 and 213.2

Example 4

Preparation of an amorphous form of gentisic acid salt (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

The gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was formed by mixing gentisic acid (0.189 g) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.500 g) in methanol (2.5 mL), followed by stirring at reflux temperature for 2 h. The solvent was distilled out at reduced pressure to obtain solid gentisate salt (Wt.: 0.680 g)

The salt (0.600 g) was dissolved into IPA (1.2 mL). To the solution DIPE (2.4 mL) was added. It was stirred for 1 h at 25-30 °C, solid was precipitated then solvent decanted. The residual solvent was distilled out under reduced pressure to obtain the gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as an amorphous solid. (Wt.: 0.260 g, % purity: 99 %, % water: 1.99% w/w, XRD=amorphous Fig (2)).

Example 5

Preparation of an amorphous form of gentisic acid salt (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

The gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was formed by mixing gentisic acid (0.113 g), (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.300 g) in a mixture of isopropylalcohol (0.6 mL) and chloroform (1.5 mL), followed by stirring at reflux temperature for 2 h. The solvent was distilled out at reduced pressure to obtain solid gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as an amorphous solid. [Wt.: 0.410 g, % purity: 98.9 %, XRD=amorphous Fig (2)].

10

Example 6

Preparation of an amorphous form of gentisic acid salt (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

The gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was formed by mixing gentisic acid (0.113 g), (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.300 g) in a mixture of isopropylalcohol (0.6 mL) and acetonitrile (1.5 mL), followed by stirring at reflux temperature for 2 h. The solvent was distilled out at reduced pressure to obtain solid gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as an amorphous solid. [Wt.: 0.410 g, % purity: 98.9 %, XRD=amorphous Fig (2)].

20

Example 7

Preparation of crystalline form of gentisic acid salt (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

25

In a dry, 25 mL round bottom flask methanol (10 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (2.0 g) were taken. To this clear solution gentisic acid (0.756 g) was added. The reaction mixture was heated up to 65-67 °C and stirred for 2 hrs. The solvent was distilled out at reduced pressure to obtain the gentisic acid salt.

30

The gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (450 mg) was dissolved in ethanol (4.5 mL) and toluene (4.5 mL) was added

subsequently, stirred for 2 to 3 hrs. Solid salt was precipitated out, filtered and washed with toluene. Subsequently, solid was dried to obtain the gentisic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine. (Wt.: 257 mg, XRD: crystalline-Fig (1))

5

Example 8

Preparation of hydrochloride salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

In a dry, 50 mL three neck flask methanol (2.5 mL), IPA (21.3mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (2.0 g) were taken. The reaction mixture was heated up to 40-45 °C to obtain clear solution. To this clear solution IPA-HCl was added upto acidic pH and then (2.5mL) diethyl ether was added into the reaction mixture. It was stirred for 1 hr at 25-30 °C. Solid salt was precipitated out. It was then heated upto 55 °C and then cooled to 25-30 °C. The solid salt was filtered and washed with IPA. The hydrochloride salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained (Wt. 1.1g, % Purity by HPLC: 99.98 %, % chiral purity by HPLC: >99 %, m.p. = 168 -170 °C, XRD=crystalline (Fig-4).

20

Example 9

Preparation of besylate salt(I) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

In a dry, 50 mL round bottom flask isopropyl acetate (25 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.5 g) were taken. The reaction mixture was heated up to 45-50 °C and a solution of benzenesulfonic acid (0.194 g) in (2.3mL) IPAc was added. Upon complete addition, solid salt was precipitated out. It was then cooled to 25-30 °C. The salt was filtered and washed with hexane. The besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained (Wt. 0.63g, % Purity by HPLC: 99.76 %, % chiral purity by HPLC: >99 %), m.p. = 171-174 °C, XRD=crystalline (Fig.5).

30

Example 10

Preparation of hydrated besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

5 In a dry, 50 mL round bottom flask isopropyl acetate (250 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (5.0 g) were taken. The reaction mixture was heated up to 45-50 °C and a solution of benzenesulfonic acid (1.93 g) in a mixture of water (0.5 mL) and isopropyl acetate (23mL) was slowly added. It was stirred for 2 h at 50 °C. It
10 was then cooled to 25-30 °C. The salt was filtered and washed with IPA and Hexane. The besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained. It was dried. (Wt. 6.7 g, % Water- 2.96 %, % Purity by HPLC: > 99.0 %, % chiral purity by HPLC: > 99 %), m.p. = 172-174 °C, XRD=crystalline (Fig. 7).

Example 11

Preparation of hydrated besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.

15 In a dry, 50 mL round bottom flask chloroform (5 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.5 g) and water (0.5 mL) were taken. The mixture was
20 stirred at 25-30 °C for 5-10 min. and solid benzenesulfonic acid (0.193 g) was added. It was heated to reflux temperature and stirred for 2 h. It was then cooled to 25-30 °C and stirred for 68 h at 25-30 °C. The salt was filtered and washed with chloroform. The
25 besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained. It was dried. (Wt. 0.660 g, % Water- 1.93 %, % Purity by HPLC: > 99.0 %, % chiral purity by HPLC: > 99 %).

Example 12

30 **Preparation of hydrated besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.**

To the 25 mL round bottom flask isopropyl acetate (5 mL), tetrahydrofuran (0.25 mL), water (0.5 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-

dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.5 g) were added. The mixture was stirred at 25-30 °C for 5-10 min. and solid benzenesulfonic acid (0.194 g) was added. It was heated to reflux temperature and stirred for 2 h. It was then cooled to 25-30 °C and stirred for 15-30 min. at 25-30 °C. The salt was filtered and washed with isopropyl acetate. The besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained. It was dried. (Wt. 0.680 g, % Water- 2.2 %, % Purity by HPLC: > 99.0 %, % chiral purity by HPLC: > 99 %).

Example 13

10 **Preparation of hydrated besylate salt(II) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.**

To the 25 mL round bottom flask isopropyl acetate (5 mL), toluene (0.25 mL), water (0.5 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (0.5 g) were added. The mixture was stirred at 25-30 °C for 5-10 min. and solid benzenesulfonic acid (0.194 g) was added. It was heated to reflux temperature and stirred for 2 h. It was then cooled to 25-30 °C and stirred for 15-30 min. at 25-30 °C. The salt was filtered and washed with isopropyl acetate. The besylate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was obtained. It was dried. (Wt. 0.660 g, % Water- 1.61 %, % Purity by HPLC: > 99.0 %, % chiral purity by HPLC: > 99 %).

Example 14

25 **Preparation of Trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.**

In a dry, 25 mL round bottom flask isopropanol (15 mL) and (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (1.0 g.) were taken. The reaction mixture was heated up to 78-80 °C and trifluoroacetic acid (0.28 g) was added. It was stirred for 1.5 h at 80-82 °C. It was then cooled to 25-30 °C. Then the solvent was evaporated. The salt was dissolved into IPA and precipitated with DIPE. The trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine was filtered and washed with DIPE. It was dried. (Wt.

0.89g, % Purity by HPLC: > 98 %, % chiral purity by HPLC: > 99 %),
XRD=crystalline (Fig. 6).

Example 15

Preparation of Trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-
5 **dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-**
amine.

The trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-
dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine
(6.5 g) was stirred with isopropylacetate (90 mL) for 2 hrs at 25-30 °C. The salt was
10 filtered and washed with isopropylacetate. It was dried (Wt. 5.4 g, % Purity by HPLC:
> 98 %, % chiral purity by HPLC: > 99 %, % water-3.14 %).

XRD=crystalline (Fig. 6).

Example 16

Preparation of amorphous benzenesulfonic acid salt of (2R)-4-oxo-4-[3-
15 **(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-**
trifluorophenyl)butan-2-amine.

In a dry, 25 mL round bottom flask methanol (2 mL) and (2R)-4-oxo-4-[3-
(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-
trifluorophenyl)butan-2-amine (1.0 g), were taken. Then benzenesulfonic acid (0.38 g)
20 was added. It was stirred for 3h at 28 °C. Further, dichloromethane (10 mL) was added
and the reaction mixture was stirred for 1h at 28 °C. Then the solvent was distilled out
at 25-30 °C under reduced pressure. The amorphous besylate salt of (2R)-4-oxo-4-[3-
(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-
trifluorophenyl)butan-2-amine was obtained (Wt. 1.2g, % Purity by HPLC: >98 %, %
25 chiral purity by HPLC: > 99 %), XRD=amorphous (Fig. 8).

Example 17

Preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-
a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine from Monohydrate
phosphate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-
30 **a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine .**

Monohydrate phosphate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-
dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine
(2 g) and water (10 mL) were taken in a 50 ml three neck flask. To the solution sodium
chloride (3.4 g) , sodium bicarbonate (0.33 g) in small lots and ethyl acetate (20 mL)

were added at 25-30 °C. It was stirred for 15-30 min. at 40-42°C. It was transferred into a separating funnel; organic layer was collected. The aqueous layer was extracted twice with ethylacetate (20 mL); organic layer was collected. All the organic layers were combined. The solvent was distilled out at reduced pressure to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (Wt. 1.8 g, % Water- 1.18 %, % Purity by HPLC: > 99.0 %).

Example 18

Preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine from Monohydrate phosphate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine .

Monohydrate phosphate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (2 g) and water (10 mL) were taken in a 50 ml three neck flask. To the solution sodium chloride (3.4 g) , 10 % aq. sodium hydroxide (3 mL) and ethyl acetate (20 mL) were added at 25-30 °C. It was stirred for 15-30 min. at 25-30 °C. It was transferred into a separating funnel; organic layer was collected. The aqueous layer was saturated with sodium chloride (3.4 g) and extracted twice with ethylacetate (20 mL). All the organic layers were combined and dried over anhydrous sodium sulfate. The solvent was distilled out at reduced pressure to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (Wt. 1.6 g, % Water- 0.39 %, % Purity by HPLC: > 99.0 %).

Example 19

Preparation of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine

The (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine (21.0 g, Chiral Purity- 96.4 %, prepared as per method disclosed in patent WO 2010/032264) was dissolved in isopropylalcohol (42.0 mL) at 68-70 °C. Then it was cooled to 45-48 °C and n-heptane (168 mL) was added drop by drop over a period of 30-45 min. The solid was precipitated. It was cooled to 25-30 °C and stirred for 1 h. The solid was filtered and washed with a mixture of isopropylalcohol and n-heptane [63 mL (1:4)]. It was dried at reduced pressure to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-

dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine
(17.1 g, Purity- 99.5 %, Chiral Purity- 99.5 %, % Water – 0.16 %.

5

10

15

20

25

30

Claims:

1. The gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine including their polymorphs and hydrates.
- 5 2. Gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 1, which is in crystalline form.
3. Crystalline gentisate salt as claimed in claim 2 which is characterized by PXRD pattern with peaks at about 8.0, 13.73, 16.00, 18.66, 22.61, 25.62 and 26.48 ° \pm 0.2°
10 two-theta (2 θ).
4. Gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 3, further characterized by a PXRD pattern substantially as depicted in Figure 1.
5. A process for preparing the gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-
15 dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 1 comprising
 - (a) dissolving (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in a suitable solvent;
 - 20 (b) thereafter addition of gentisic acid and reacting under suitable conditions;
 - (c) removing the solvent under suitable conditions to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate.
6. The process as claimed in claim 5, wherein the suitable solvent used is selected
25 from alcohols, water or suitable mixture thereof.
7. A process for preparing the crystalline form of gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claims 2-4 comprising
 - (a) crystallization of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-
30 dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt in the presence of suitable solvent and optionally further adding suitable anti solvents and

- (b) removing the solvent under suitable conditions to obtain the crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate.
8. The process as claimed in claim 7, wherein the suitable solvent used is selected
5 from hydrocarbons and mixture of alcohols and hydrocarbons.
9. Amorphous gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine.
10. The amorphous form of gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 9, characterized by a PXRD pattern as depicted in Figure
10 2.
11. A process for preparing the amorphous form of gentisate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claims 9-10 comprising
15 (a) dissolving or slurring of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate salt in the presence of suitable solvent and optionally further adding suitable anti solvents and
20 (b) removing the solvent under suitable conditions to obtain amorphous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine gentisate
12. The process as claimed in claim 11, wherein the suitable solvent used is selected from suitable alcohol, water, mixture of alcohol and water; mixture of alcohol and
25 ether, mixture of alcohol and halogenated hydrocarbon and mixture of alcohol and nitrile.
13. The adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine including their polymorphs and hydrates.
- 30 14. Adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 13, which is characterized by PXRD pattern with peaks at about 8.05, 17.64, 18.29, 21.65, 23.38, 23.85 and $24.38^\circ \pm 0.2^\circ$ two-theta. (2 θ).

15. Adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claims 13-14, further characterized by a PXRD pattern substantially as depicted in Figure 3.
- 5 16. A process for preparing the adipate salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claims 13-15 comprising
- (a) dissolving of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine in a suitable solvent;
- 10 (b) thereafter addition of adipic acid and reacting under suitable conditions and;
- (c) removing the solvent under suitable conditions to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine adipate.
- 15 17. The process as claimed in claim 16, wherein the suitable solvent used is selected from alcohols, water or suitable mixture thereof.
18. The trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine including their polymorphs and hydrates.
- 20 19. Trifluoroacetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 18, which is characterized by PXRD pattern with peaks at about 6.64, 9.39, 10.99, 13.40, 15.32, 17.15, 19.20, 19.66, 21.48, 22.46, 24.68, 26.89, 27.86, 29.34, 31.24, 32.77 and $34.82 \pm 0.2^\circ$ two-theta (2θ).
- 25 20. Trifluoro acetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 18, further characterized by a PXRD pattern substantially as depicted in Figure 6.
21. A process for preparing the trifluoro acetic acid salt of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claim 18 comprising
- 30 (a) dissolving of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine or its salts with first acid in a suitable solvent;

- (b) thereafter addition of trifluoroacetic acid and reacting under suitable conditions;
- (c) removing the solvent under suitable conditions to obtain (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoroacetic acid salt.
- 5 22. The process as claimed in claim 21, wherein the suitable solvent used is selected from alcohols or suitable mixture thereof.
23. The process as claimed in claim 21, wherein the suitable salt of first acid is selected from HCl, acetic acid.
24. A process for preparing the crystalline form trifluoro acetic acid salt of (2R)-4-oxo-
10 4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine as claimed in claims 19-20 comprising
- (a) crystallization of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoro acetic acid salt in the presence of suitable solvent and optionally
15 further adding suitable anti solvents and
- (b) removing the solvent under suitable conditions to obtain crystalline (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine trifluoroacetic acid salt.
25. The process as claimed in claim 24, wherein the suitable solvent used is selected
20 from alcohols, esters and ethers or suitable mixture thereof.
26. The crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride.
27. The crystalline (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-
25 a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride as claimed in claim 26 having characteristic powder X-ray diffraction pattern having peaks expressed as 2θ at about 6.50, 8.05, 13.79, , 18.05, 19.52, 22.65, 23.40, 25.39 and 27.09 ± 0.2 degree two theta.
28. The crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-
30 dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride as claimed in claim 26, further characterized by a PXRD pattern as depicted in Figure 4.

29. A process for preparing the crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride as claimed in claim 26 comprising
- 5 (a) reaction of the free base (formula I) under suitable reaction conditions with the hydrochloric acid in suitable solvent;
- (b) suitably removing the solvent to obtain the crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine hydrochloride.
30. The process as claimed in claim 29, wherein the suitable solvent used is selected
- 10 from alcohols, ethers and water or suitable mixtures thereof.
31. The crystalline form (I) of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate.
32. The crystalline (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-
- 15 a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate (I) as claimed in claim 31 having characteristic powder X-ray diffraction pattern having peaks expressed as 2θ at about 6.81, 18.31, 22.99, 23.70 and 25.75 ± 0.2 degree two theta.
33. The crystalline form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-
- 20 dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate as claimed in claim 31, characterized by a PXRD pattern as depicted in Figure 5.
34. A process for preparing the crystalline form (I) of (2R)-4-oxo-4-[3-
- (trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-
- 25 trifluorophenyl)butan-2-amine besylate as claimed in claim 31 comprising
- (a) reaction of the free base (formula I) under suitable reaction conditions with the benzenesulfonic acid in suitable solvent;
- (b) suitably removing the solvent to obtain the crystalline form of (2R)-4-oxo-4-
- 30 [3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate.
35. The process as claimed in claim 34, wherein the suitable solvent used is selected from esters and hydrocarbons or suitable mixture of thereof.

36. The crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate.
37. The crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate as claimed in claim 36 having characteristic powder X-ray diffraction pattern having peaks expressed as 2θ at about 5.69, 7.45, 10.08, 11.41, 13.09, 18.34, 22.52 and 22.97 ± 0.2 degree two theta.
38. The crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate as claimed in claim 36, characterized by a PXRD pattern as depicted in Figure 7.
39. A process for preparing the crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate as claimed in claim 36 comprising
- (a) reaction of the free base (formula I) under suitable reaction conditions with the benzenesulfonic acid in suitable solvent;
- (b) suitably removing the solvent to obtain the crystalline form II of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate.
40. The process as claimed in claim 39, wherein the suitable solvent used is selected from a mixture of esters and water, mixture of esters, ethers and water, mixture of esters, alcohols and water, mixture of ester, hydrocarbons and water, mixture of esters, ketones and water.
41. Amorphous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate.
42. The amorphous form of (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate as claimed in claim 41, further characterized by a PXRD pattern as depicted in Figure 8.
43. A process for preparing amorphous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate as claimed in claim 41 comprising

- 5 (a) dissolving or slurring of (2R) 4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate salt in the presence of suitable solvent and optionally further adding suitable anti solvents
- 5 (b) removing the solvent under suitable conditions to obtain amorphous (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine besylate
- 10 44. The process as claimed in claim 43, wherein the suitable solvent used is selected from alcohols, halogenated hydrocarbons and water or suitable mixtures thereof.
- 15

Fig. 1




Fig.2




Fig. 3




Fig-4




Fig-5




Fig-6




Fig-7




Fig-8




Figure: 9- Effect of single dose oral administration of ZY compounds at 10mg/kg, PO on OGTT in male C57 mice.

Effect of single dose oral administration of ZY compounds at 10mg/kg, PO on OGTT in male C57 mice




Figure: 10- Effect of single dose oral administration of ZY compounds at 10mg/kg, PO on OGTT in male C57 mice.

