



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2637726 C 2014/04/15

(11)(21) **2 637 726**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2007/01/25
(87) Date publication PCT/PCT Publication Date: 2007/08/02
(45) Date de délivrance/Issue Date: 2014/04/15
(85) Entrée phase nationale/National Entry: 2008/07/18
(86) N° demande PCT/PCT Application No.: US 2007/001875
(87) N° publication PCT/PCT Publication No.: 2007/087344
(30) Priorité/Priority: 2006/01/25 (US60/761,832)

(51) Cl.Int./Int.Cl. *A61K 31/495* (2006.01),
A61P 37/08 (2006.01)

(72) **Inventeurs/Inventors:**
ASOTRA, SATISH, CA;
ZADYKOWICZ, JERZY, CA;
VANAM, KALPANA R., US;
GAO, SHEN, CA

(73) **Propriétaire/Owner:**
TARO PHARMACEUTICALS NORTH AMERICA, INC.,
KY


(74) **Agent:** METHOD LAW PROFESSIONAL
CORPORATION

(54) Titre : COMPOSITIONS ANTIHISTAMINIQUES ET LEUR UTILISATION

(54) Title: ANTI-HISTAMINE COMPOSITIONS AND USE THEREOF

(57) **Abrégé/Abstract:**

The present invention provides for a storage stable pharmaceutical liquid solution for oral administration having a pharmaceutically effective amount of an antihistamine and having a purity equal to or greater than about 99% by weight-based HPLC assay, residual solvents of less than about 0.5%, and a total impurity of less than about 0.2%. The storage stable solution preferably contains cetirizine. The present invention further provides a process of preparing the storage stable pharmaceutical liquid solution as well as a method of treating a mammal with a therapeutically effective amount of cetirizine in the stable pharmaceutical liquid solution.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
2 August 2007 (02.08.2007)

PCT

(10) International Publication Number
WO 2007/087344 A3(51) International Patent Classification:
A61K 31/19 (2006.01) A61K 31/135 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2007/001875

(22) International Filing Date: 25 January 2007 (25.01.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/761,832 25 January 2006 (25.01.2006) US

(71) Applicant (for all designated States except US): TARO PHARMACEUTICALS U.S.A., INC. [US/US]; 3 Skyline Drive, Hawthorne, New York 10532 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ASOTRA, Satish [CA/CA]; 16 Langford Court, Brampton, Ontario L6W 4K4 (CA). ZADYKOWICZ, Jerzy [CA/CA]; 3867 Arvona Place, Mississauga, Ontario L5M 6L4 (CA). RAO, Kalpana [—/US]; Albertson, New York (US). GAO, Shen [CA/CA]; 23 Meadow Brook Court, Botonin, Ontario L7E 2Y6 (CA).

(74) Agents: GOLLIN, Michael A. et al.; Venable LLP, P.O. Box 34385, Washington, DC 20043-9998 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
13 December 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTI-HISTAMINE COMPOSITIONS AND USE THEREOF

(57) Abstract: The present invention provides for a storage stable pharmaceutical liquid solution for oral administration having a pharmaceutically effective amount of an antihistamine and having a purity equal to or greater than about 99% by weight-based HPLC assay, residual solvents of less than about 0.5%, and a total impurity of less than about 0.2%. The storage stable solution preferably contains cetirizine. The present invention further provides a process of preparing the storage stable pharmaceutical liquid solution as well as a method of treating a mammal with a therapeutically effective amount of cetirizine in the stable pharmaceutical liquid solution.

ANTI-HISTAMINE COMPOSITIONS AND USE THEREOF

BACKGROUND OF THE INVENTION

5

[001] Cetirizine is a metabolite of the anti-histamine, hydroxyzine. Similar to other second-generation anti-H₁ receptor antagonists, cetirizine does not cross the blood brain barrier and is considered a peripherally active anti-histamine with few central nervous system effects. Studies have shown that cetirizine provides safe and effective, symptomatic relief of seasonal allergies.

10 Advantages to using cetirizine as an anti-histamine include minimal sedative effect, little anticholinergic activity and a longer acting duration.

[002] Cetirizine is water soluble, and rapidly absorbed. Tablets and syrup formulations of cetirizine have been commercially manufactured and marketed. However, cetirizine pharmaceutical formulations are subject to degradation that occurs during manufacture and storage. U.S. Patent No.

15 6,171,618 teaches that an esterification process of the cetirizine occurs when low molecular weight esters or alcohols are present in a reactive mixture with cetirizine. In particular, high boiling point alcohols, especially glycerin, should not be present during the manufacturing and in formulations of cetirizine. The resultant ester impurities can cause the formulations to deviate from regulatory purity requirements and decrease the shelf-life of the formulations.

20 [003] Spill resistant pharmaceutical formulations have been described in U.S. Patent Nos. 6,102,254, 6,071,523 and 6,399,079. The spill resistant formulations have been described by the physical properties of (a) extrudability under light manual pressure from a squeezable container or a proxy (e.g. a syringe with a 5 mm orifice), and (b) spreadability in a spoon bowl measured by extruding the formulation into a spoon bowl and determining whether the material levels or spreads 25 to the edges of the spoon bowl. Spreadability also contributes to accuracy of measurement.

[004] Surprisingly, we have discovered a spill resistant cetirizine formulation, containing high levels of low molecular weight polyhydric alcohol that does not form the most common cetirizine formulation degradative products, has a high degree of purity and increased storage time.

SUMMARY OF THE INVENTION

[005] The invention is to a stabilized pharmaceutical liquid formulation comprising an antihistamine and having a purity equal to or greater than about 99% by weight-based HPLC assay, residual solvents of less than about 0.5%, and a total impurity of less than about 0.2%.

5 Surprisingly, the impurities are often less than 0.1% of the active. The stability of the active ingredient during the shelf storage of the product allows for an extended shelf life of the product compared to other antihistamine formulations. Consequently, one embodiment of the invention allows for antihistamine solutions that have a storage stability of up to 24 months. Another 10 embodiment would allow for a storage stability of up to 36 months.

[006] One embodiment of the invention is directed to a stabilized liquid formulation of cetirizine, or a salt thereof. The particular formulations described herein will contain cetirizine, a carrier component and a thickener. In one embodiment the carrier may be glycerin and the thickener is a carbomer. The formulation may contain at least one organoleptic agent, and other pharmaceutically 15 acceptable excipients may be present to provide a palatable, stable, and pleasing composition.

[007] In addition to pharmaceutically acceptable excipients, the formulation may comprise other active pharmaceutical ingredients in combination with the antihistamine. Some embodiments of the inventive formulation may include a decongestant, an analgesic, an antitussive, an expectorant, or any combination of two or more thereof.

20 [008] The storage stable solution of the present invention has a pH of between about 5.4 and 8.0, more preferably between 6.2 and 7.3.

[009] Another embodiment of the invention is directed towards a method of treating a human in need of cetirizine by administering a therapeutically effective amount of the storage stable pharmaceutical liquid solution of the invention.

25 [010] An additional embodiment of the invention is a method of preventing the formation of impurities in a pharmaceutical liquid formulation comprising mixing an antihistamine in a storage stabilizing vehicle.

[011] Additionally, there is disclosed the method of providing an extended product shelf life, that method comprising mixing the antihistamine with a storage stable vehicle.

DETAILED DESCRIPTION

[012] In describing embodiments of the present invention, specific terminology is employed for the sake of clarity. However, the invention is not intended to be limited to the specific terminology so selected. It is to be understood that each specific element includes all technical equivalents, which operate in a similar manner to accomplish a similar purpose.

10

[013] The inventive solution has, as the active pharmaceutical agent, an antihistamine, evenly dissolved in a semi-solid aqueous vehicle. A preferred antihistamine is cetirizine. Cetirizine, a piperazine derivative and carboxylated metabolite of hydroxyzine, is a potent histamine H₁-receptor antagonist with antiallergic properties. Cetirizine is used herein to describe not only the free compound of cetirizine, but also any pharmaceutically acceptable salt thereto. Preferred are the acid addition salts, in particular, cetirizine dihydrochloride. The term cetirizine also includes racemic mixtures and the enantiomers of cetirizine.

[014] The solution provides a method of treating allergic conditions, such as allergic rhinitis and chronic uticaria in mammals. The methods include administering to a subject suffering from an allergic condition a storage stable pharmaceutical composition according to the invention. In one embodiment, the subject is a human. In another embodiment the allergic condition is allergic rhinitis or chronic idiopathic uticaria. The storage stable pharmaceutical composition can be administered to a patient in a dosage range of, for example, 0.5 mg to about 15 mg per day, preferably about 1 mg to about 12 mg per day, and more preferably 5 to 10 mg per day.

[015] The inventive solution comprises an aqueous and solution-stabilizing vehicle that is pharmaceutically acceptable. By a stabilizing solution, it is meant that not more than about 0.2%, preferably not more than about 0.1%, of the cetirizine is degraded into known or unknown impurities under varying temperature and humidity conditions. Impurities are the unwanted chemicals that develop during formulation or upon aging of the formulated active pharmaceutical

ingredient. An impurity can be any component present in the intermediate or active ingredient that is not the desired entity. An impurity may be a characterized or unknown chemical entity. The presence of these impurities, even in small amounts, may have an effect on the safety and efficacy of the pharmaceutical product. Organic impurities may arise during the manufacturing process and/or storage of the drug substance. They may be identified or unidentified, volatile or non-volatile, and include starting materials, by-products of the synthesis procedure or degradation products. Although the end products are washed with solvents, there is always the chance of having residual unreacted starting materials. Impurities can also be formed by degradation of the end product during manufacturing of bulk drugs. In general, a pharmaceutical formulation may contain all of the above-mentioned types of organic impurities at levels varying from negligible to significant.

[016] There are several known impurities of cetirizine that are monitored during production of pharmaceutical formulations. 4-Chloro Benzhydryl Piperazine is a starting material in the manufacture of cetirizine active pharmaceutical ingredient. This compound is not a degradation product, and its impurity level in the formulation would depend on the amount that is in starting amount of cetirizine. Cetamide (2-[4-(4-chloro-benzhydryl]-1-piperazinyl)] ethoxy)acetamide is an intermediate in the process of preparing the active pharmaceutical ingredient. Thus, cetamide is a byproduct of the production steps and its impurity level in the formulation would also depend on the amount that is in the originally used cetirizine. Glycerin-cetirizine esters are formed between the cetirizine carboxylic group and the hydroxyl groups of polyhydric alcohols over time. These esters are degradation products and have been previously described for the manufactured cetirizine oral solutions, and in U.S. Patent No. 6,171,618.

[017] A HPLC method was developed and validated for the determination of cetirizine dihydrochloride as well as its related impurities in the commercial oral solution and the inventive formulation. Test samples of the cetirizine solutions were monitored for active and impurities. The testing was done at room temperature and at accelerated conditions of 30°C and 40° C. Cetirizine and its impurities were measured by HPLC employing either Agilent™ (Hewlett-Packard) 1100 HPLC and / or Waters Alliance™ HPLC System 2695. Chromatograms from the HPLC show the peaks of detectable chemicals in the solution. The method was validated with respect to linearity and range, method accuracy, method precision, method specificity, limit of quantitation and method

robustness. The validation method proved the HPLC measurements to be accurate, precise, specific and robust. The measurement of impurities was made by measuring the Relative Retention Time which is a ration of the retention time of the analyte to the retention time of the active, cetirizine.

[018] The guidelines for the International Conference on Harmonization (ICH), require that, for 5 a maximum daily dosage of 10mg/day for children, any impurity greater than 0.5% in a solution be qualified, any impurity in an amount of greater than 0.2% be identified and any impurity greater than 0.1% be reported. In general, liquid dosage forms are much more susceptible to both degradation and microbiological contamination. In this regard, water content, pH of the solution/suspension and the mutual interactions of ingredients are critical factors. The limit of 10 detection is the minimum concentration (% w/w) at which the analyte can reliably be detected. The limit of quantitation is the minimum concentration (% w/w) at which the analyte can reliably be quantified. Limits of detection and quantitation were determined by comparing measured signals from samples with known low concentrations of analyte to measured signals from blank samples.

[019] The storage stabilizing vehicle of the present invention comprises a thickener component 15 and a carrier component, and may include organoleptic components. Carbomers (Merck Index 12th ed., no. 1878) can be used as thickeners in semi-solid pharmaceutical formulations (see Mchita et al., US patent 6,071,523). Carbomer 934P (Carbopol[®] 974P) is a suitable thickener or gelling agent. Suitable concentrations from about 0.2 to about 1.0 %, and more specifically from about 0.40 to 20 about 0.45 %, w/w. Its rheology supports a high yield value (Handbook of Pharmaceutical Excipients Third Ed., A.H. Kibbe (Ed.), Pharmaceutical Press, London, UK, 2000, Pg. 442, 79, 53 ("Handbook of Pharm. Excipients"). Carbomers are slightly acidic and must be neutralized e.g. with sodium hydroxide (as needed to neutralize the carborner up to about 0.08 % in particular formulations) with a preferred pH range being about 6.2 to about 7.3, providing the maximum viscosity plateau. The vehicle carrier component comprises propylene glycol up to about 20%, or 25 from about 3 to about 10%. Glycerin up to about 50% may be present. Additionally, sorbitol, up to 10%, may be added as a vehicle and stabilizer. Purified water comprises the bulk of the carrier component comprising from about 29% to about 64% of the formulation.

[020] The vehicle of the present invention has been described in U.S. Publication No. 2005-0175642 A1, as a stabilizing composition for a suspension of the 30 piperidine antihistamine, Loratadine.

[021] The solutions of the present invention may also contain Eddate Disodium (EDTA). EDTA is a chelating agent that forms a stable water-soluble complex with alkaline earth and heavy metal ions. It is useful as an antioxidant synergist, sequestering metal ions that might otherwise catalyze autoxidation reactions. EDTA may also have synergistic effects as an antimicrobial when used in combination with other preservatives (Handbook of Pharmaceutical Excipients 4th Ed.).

[022] Microbiological growth resulting from the growth of bacteria, fungi, and yeast in a humid and warm environment may result in oral liquid products that are unusable for human consumption. Microbial contamination may occur during the shelf life and subsequent consumer-use of a multiple-dose product due to inappropriate use of certain preservatives in the preparations, or because of the semi-permeable nature of primary containers. The solution has antimicrobial activity. Propylparaben (up to about 0.04%) and butylparaben (0.018% to about 0.18%) are suitable. Other antimicrobial excipients may also be used.

[023] Water Activity is a measure of the free or unbound water in a product. Water activity is a critical factor that determines shelf life. Most bacteria will not grow at water activities below 0.80, and most molds cease to grow at water activities below 0.70. By measuring water activity, it is possible to predict which microorganisms will and will not be potential sources of spoilage. Water activity determines the lower limit of available water for microbial growth. In addition to influencing microbial spoilage, water activity can have a major impact their color, taste, and aroma. Water activity was measured again using an AwQuick™ unit (Rotronic Instrument Corporation, Huntington, NY). Initially, the water activity was found to be too high and so sorbitol was introduced to decrease the water level. Water activity measurements are shown in table V.

The organoleptic ingredients improve the taste and appearance and do not negatively affect the solution, stability. The organoleptic agents in the following examples include coloring and flavoring agents, sweeteners and masking agents. The pharmaceutical formulation of the present invention may also contain additional active pharmaceutical ingredients, such as for example, decongestants, analgesics, antitussives and expectorants. Any specific drug within these therapeutic classes is suitable for inclusion in the present invention. Illustrative examples include analgesics such as aspirin, acetaminophen, naproxen, ketoprofen and ibuprofen; decongestants such as pseudoephedrine or phenylpropanolamine; antitussives such as codeine, hydrocodone, or dextromethorphan; and expectorants such as guaifenesin, including salts thereof.

EXAMPLES

[024] **Example 1:** The storage stable solution was formulated to contain the following ingredients:

5 **Table I: Cetirizine Storage Stable Formulation**

INGREDIENTS	% (w/v)
Cetirizine	0.087
Purified Water	38.69
Eddate Disodium USP	0.04
Glycerin	50
Sorbitol (Crystalline)	5.0
Propylene Glycol	5.0
Carbomer 934P (Carbopol® 974P)	0.43
Sodium Hydroxide	0.0116
Sucralose Liquid Concentrate	0.20
Butylparaben	0.04
Masking Agent	0.20
Artificial Grape Flavor	0.20

[025] **Example 2:** As a comparative example, a cetirizine hydrochloride syrup (5mg/5mL) was prepared. The composition of the syrup is as follows:

10 **Table II: Cetirizine Syrup Formulation**

INGREDIENTS	% (w/v)
Cetirizine	0.087
Purified Water	q.s.
Glycerin	24.7
Sucrose	45.1
Propylene Glycol	12.2

Sodium Acetate (anhydrous)	0.12
Glacial Acetic Acid	0.22
Butylparaben	0.02
Methylparaben	0.18
Flavoring	0.15 to 0.25

[026] **Example 3: Impurities**

Table III: Three exhibit batches of cetirizine spill resistant solution were manufactured with the formulation as given in Example 1. Cetirizine solution sample 1, sample 2, and sample 3 were 5 placed in amber bottles, and placed in an inverted position for a three-month time period for regulatory stability testing. The environmental conditions for the exhibit batches were either room temperature with average relative humidity (RH) ($25\pm2^{\circ}\text{C}$; $60\pm5\%\text{RH}$), or high temperature with high relative humidity ($40\pm2^{\circ}\text{C}$; $75\pm5\%\text{RH}$). Total unidentified degradation products of the cetirizine stabilized solution were never above 0.1%. In contrast, after three months of stability 10 testing with the cetirizine syrup as prepared in example 2, glycerin-cetirizine ester was the main degradation product. The total impurities seen in the cetirizine syrup formulations were ten times more than the total impurities seen in the stabilized solution. Additionally, 0.6% of the total impurities measured in the syrup were identified as glycerin-cetirizine ester. Surprisingly, there was 15 no detection of any glycerin-cetirizine ester in the stabilized solution.

15

Table III: Total Impurities Cetirizine Oral Formulations (HPLC % total active)

Sample	Temperature and Relative Humidity Conditions	
	$25\pm2^{\circ}\text{C}$; $60\pm5\%\text{RH}$	$40\pm2^{\circ}\text{C}$; $75\pm5\%\text{RH}$
Cetirizine Solution Sample 1	0.07%	0.06%
Cetirizine Solution Sample 2	0.09%	0.08%
Cetirizine Solution Sample 3	0.09%	0.07%

Cetirizine Syrup	0.2%	0.7% (0.6% identified as glycerin-cetirizine ester)
------------------	------	---

[027] **Table IV:** The impurities of the spill resistant stabilized cetirizine, as described in example 1, was compared to a sample of the commercially available cetirizine syrup. Both samples were placed in amber bottles, and placed in an inverted position for a three-month time period for 5 regulatory stability testing. The environmental conditions for the exhibit batches were either room temperature with average relative humidity (RH) (25±2°C; 60±5%RH), or high temperature with high relative humidity (40±2°C; 75±5%RH). Total unidentified degradation products of the cetirizine stabilized solution were never above 0.2% for both room temperature and accelerated conditions. In contrast, after six months of stability testing with the commercial syrup impurity 10 levels of 0.29% at room temperature and 3.51% at high temperature were recorded.

Table IV: Comparison of Impurities in Cetirizine Spill Resistant Solution Vs Cetirizine Commercial Syrup

	Time	Cetirizine Spill Resistant 5mg/5ml		Commercial Syrup 5mg/5ml	
		RT	40° C	RT	40° C
Impurities, individual unidentified	Initial	LT ² 0.05%	LT 0.05%	RRT 1.3: 0.07%	RRT 1.3: 0.07%
Impurities, total	Initial	LT 0.05%	LT 0.05%	0.07%	0.07%
	3 months	0.08%	0.15%	0.13%	1.83%
	6 months	0.06%	0.05%	0.29%	3.51%

1 – Results shown are the average values from analysis of three lots.

2 – LT = Less Than

Example 5: Water Activity

Most bacteria will not grow at water activities below 0.91, and most molds cease to grow at water activities below 0.80. Water activity was measured using an AwQuick unit (Rotronic Instrument Corporation, Huntington, NY). Water activity measurements are shown in table VII.

5

Table V: Water Activities Cetirizine Hydrochloride Spill Resistant

Lot #	Sorbitol (%)	Water (%)	Butylparaben (%)	Water Activity A_w
1	0	~44%	0.018	0.766
2	0	~44%	0.0144	0.771
3	0	~44%	0.009	0.766
4	5	~39%	0.018	0.723
5	5	~39%	0.0144	0.724
6	5	~39%	0.009	0.722

WO 2007/087344

PCT/US2007/001875

CLAIMS

We claim:

1. A storage-stabilized pharmaceutical liquid formulation comprising cetirizine or a salt thereof in a semi-solid spill resistant aqueous vehicle comprising glycerin in an amount of up to 50% by weight, propylene glycol in an amount of up to 20% by weight, and carbomer in an amount of 0.2 to 1.0% by weight, wherein the cetirizine has a purity equal to or greater than 99% by weight-based HPLC assay, residual solvents of less than 0.5% by weight, and a total impurity of less than 0.2% by weight.
2. The formulation of claim 1 wherein the total impurity is less than 0.1%
3. The formulation of claim 1 having a storage stability of up to 24 months.
4. The formulation of claim 1 having a storage stability of up to 36 months.
5. The formulation of claim 1, further comprising at least one organoleptic agent.
6. The formulation of claim 1 further comprising a decongestant, an analgesic, an antitussive, an expectorant, or any combination of two or more thereof.
7. The formulation of claim 1 having a pH of between 5.4 and 8.0.
8. The formulation of claim 1 having a pH of between 6.2 and 7.3.
9. Use of a therapeutically effective amount of the formulation of claim 1 in the preparation of a medicament for treating allergic conditions.

WO 2007/087344

PCT/US2007/001875

10. A method of preventing formation of impurities in a pharmaceutical liquid formulation comprising mixing cetirizine or a salt thereof with a storage stabilizing vehicle comprising glycerin in an amount of up to 50% by weight, propylene glycol in an amount of up to 20% by weight, and carbomer in an amount of 0.2 to 1.0% by weight.
11. A method of providing cetirizine or a salt thereof in a liquid formulation having an extended product shelf life, that method comprising mixing cetirizine or a salt thereof with a storage stabilizing vehicle comprising glycerin in an amount of up to 50% by weight, propylene glycol in an amount of up to 20% by weight, and carbomer in an amount of 0.2 to 1.0% by weight.
12. The formulation of claim 1, wherein the propylene glycol is in an amount of from 3% by weight to 10% by weight.
13. The formulation of claim 1, wherein the formulation further comprises sorbitol in an amount of up to 10% by weight.