



(51) International Patent Classification:

B01D 53/52 (2006.01) **C08L 95/00** (2006.01)
C08K 5/3492 (2006.01)

(21) International Application Number:

PCT/US2018/024272

(22) International Filing Date:

26 March 2018 (26.03.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/478,427 29 March 2017 (29.03.2017) US

(71) Applicant: **ECOLAB USA, INC.** [US/US]; 1 Ecolab Place, St. Paul, Minnesota 55102 (US).

(72) Inventor: **SOLOMON, Kim R.**; 337 Devin Lane, River Falls, Wisconsin 54022 (US).

(74) Agent: **BABYCH, Eric D.**; Brinks Gilson & Lione, 405 S. Main Street, Suite 1000, Salt Lake City, Utah 84111 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: DISSOLUTION OF HEXAMINE IN NON-AQUEOUS SOLVENT

(57) Abstract: The present disclosure generally relates to scavenging hydrogen sulfide. The disclosure pertains to non-aqueous and non-volatile compositions that include a monolignol alcohol and hydrogen sulfide scavenging compounds which are hexamine or hexamine derivatives. The hydrogen sulfide scavenging compound may be hexamine in some aspects. The compositions may also include a C₂-glycol. The compositions disclosed are stable and can be used, for example, in removing hydrogen sulfide from hot asphalt.



DISSOLUTION OF HEXAMINE IN NON-AQUEOUS SOLVENT

BACKGROUND

1. Field of the Invention


[0001] The present disclosure generally relates to scavenging hydrogen sulfide. More particularly, the disclosure pertains to the use of non-aqueous hexamine compositions to scavenge hydrogen sulfide present in asphalt.

2. Description of the Related Art

[0002] Petroleum asphalt is produced as a residue of a thermal separation refinery process. The thermal separation process causes thermal cracking to occur which frequently causes hydrogen sulfide to be present in the asphalt stream. In fact, thermal cracking continues in the asphalt even after the asphalt has left the vacuum distillation section of the operation, particularly at high temperature. In order to permit the safe loading, handling, and storage of the asphalt, it is necessary to reduce the hydrogen sulfide to safe levels in the asphalt. This has been done in the past by weathering of the hot asphalt for sufficient time for the hydrogen sulfide to be reduced to safe levels. This not only takes a considerable amount of time (several days), but it releases hydrogen sulfide to the vapor space in the storage, which could create hazardous conditions. Moreover, recent emphasis on environmental regulations in Europe stresses the limits on the hydrogen sulfide content of vent gas.

BRIEF SUMMARY

[0003] In some embodiments, a composition is provided that includes a monolignol alcohol and a compound of formula (I) or formula (II)



where R_1 may be a substituted or unsubstituted C_{1-32} alkyl, a substituted or unsubstituted C_{2-32} alkenyl, or a substituted or unsubstituted C_{2-32} alkynyl. The substituted C_{1-32} alkyl, the substituted C_{2-32} alkenyl, and the substituted C_{2-32} alkynyl may be substituted with hydroxyl or C_{1-5} alkyl.

[0004] In some embodiments, the monolignol alcohol is selected from the group consisting of p-coumaryl alcohol, coniferyl alcohol, sinapyl alcohol, cinnamyl alcohol, and any combination thereof. The monolignol alcohol may be cinnamyl alcohol. The monolignol alcohol may be coniferyl alcohol. The compound may be hexamine.

[0005] In some embodiments, the composition comprises a C_{2-8} polyol. The C_{2-8} polyol may be glycerin.

[0006] In some embodiments, the composition comprises the compound of formula (I), cinnamyl alcohol, and glycerin.


[0007] In some embodiments, the composition comprises about 2 to about 20 weight percent of the compound of formula (I) or formula (II). The composition may comprise about 10 to about 98 weight percent of the monolignol alcohol. The composition may comprise about 2 to about 80 weight percent of the C_{2-8} polyol.

[0008] In some embodiments, a weight ratio of the monolignol alcohol to the C_{2-8} polyol is about 10:1 to about 20:1. In some embodiments, a weight ratio of the compound of formula (I) or formula (II) to monolignol alcohol is about 1:15 to about 1:1.

[0009] In some embodiments, the composition comprises an amount of water that is less than about 1% by weight. In some embodiments, the composition excludes water.

[0010] The composition may comprise a pH greater than about 7.

[0011] In some embodiments, a method is provided for scavenging hydrogen sulfide in an industrial process. The method may include adding an effective amount of a composition to a medium in the industrial process. The composition may include a monolignol alcohol and a compound of formula (I) or formula (II)




where R_1 is a substituted or unsubstituted C_{1-32} alkyl, a substituted or unsubstituted C_{2-32} alkenyl, or a substituted or unsubstituted C_{2-32} alkynyl, wherein the substituted C_{1-32} alkyl, the substituted C_{2-32} alkenyl, and the substituted C_{2-32} alkynyl are substituted with hydroxyl or C_{1-5} alkyl.

[0012] In some embodiments, the composition of the method further comprises a C_{2-8} polyol and the monolignol alcohol is cinnamyl alcohol.

[0013] In some embodiments, the effective amount is in a range from about 0.00001 ppm to about 100,000 ppm.

[0014] In some embodiments, the medium is selected from the group consisting of crude oil, diesel fuel, asphalt and any combination thereof.

[0015] The present disclosure also provides the use of a composition in a method of scavenging hydrogen sulfide. The composition comprises a monolignol alcohol and a compound of formula (I) or formula (II)



, where R_1 is a substituted or unsubstituted C_{1-8} alkyl, a substituted or unsubstituted C_{2-8} alkenyl, or a substituted or unsubstituted C_{2-8} alkynyl, wherein the substituted C_{1-8} alkyl, the substituted C_{2-8} alkenyl, and the substituted C_{2-8} alkynyl are substituted with hydroxyl or C_{1-5} alkyl.

[0016] The foregoing has outlined rather broadly the features and technical advantages of the present disclosure in order that the detailed description that follows may be better understood. Additional features and advantages of the disclosure will be described hereinafter that form the subject of the claims of

this application. It should be appreciated by those skilled in the art that the conception and the specific embodiments disclosed may be readily utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present disclosure. It should also be realized by those skilled in the art that such equivalent embodiments do not depart from the spirit and scope of the disclosure as set forth in the appended claims.

DETAILED DESCRIPTION

[0017] Various embodiments are described below. The relationship and functioning of the various elements of the embodiments may better be understood by reference to the following detailed description. However, embodiments are not limited to those illustrated below. In certain instances details may have been omitted that are not necessary for an understanding of embodiments disclosed herein.

[0018] Environmental standards and safety standards require the removal of hydrogen sulfide from oil, asphalt, or other materials. Various strategies and chemicals are used to remove hydrogen sulfide.

[0019] Hexamine can be used as a hydrogen sulfide scavenger. Hydrogen sulfide scavenger refers to compounds that may eliminate hydrogen sulfide or reduce its concentration in a medium such as oil, asphalt, or diesel.

Hexamine decomposes in the presence of acid to ammonia and formaldehyde. Formaldehyde or ammonia can then react with hydrogen sulfide converting the volatile and toxic gas into a nonvolatile material.

Despite the usefulness of hexamine, it has limited solubility in common solvents. Some solvents that can effectively dissolve hexamine are volatile and incompatible with compositions that need to be added to hot asphalt.

Additionally, hexamine is available in at least two different forms. Hexamine can be in a hexahydrate form that melts below room temperature at 13 °C, which is generally described as an “aqueous solution” of hexamine.


Hexamine can be in a pure solid form with a melting point of about 280 °C.

[0020] Hexamine has limited solubility in most common organic solvents at 20 °C: 13.4 g in chloroform, 7.25 g in methanol, 2.89 g in ethanol, 0.65 g in acetone, 0.23 g in benzene, 0.14 g in xylene, 0.06 g in ether, and near zero in

petroleum ether. Chloroform is not suitable due to toxicity, and ethanol and acetone are too volatile for use in use in hot asphalt, which can be up to 300-400 °F.

[0021] The present disclosure seeks to solve the problems associated with the solubility of hexamine. The compositions may be non-aqueous and non-volatile, which is ideal for applications involving hot material such as asphalt. The compositions may be halogen-free, which is desirable for improved worker safety.

[0022] In some embodiments, a composition is disclosed for scavenging hydrogen sulfide. The composition may include a hydrogen sulfide scavenger. The hydrogen sulfide scavenger may be a compound of formula (I) or formula (II) as shown below,



where R_1 may be a substituted or unsubstituted C_{1-32} alkyl, a substituted or unsubstituted C_{2-32} alkenyl, or a substituted or unsubstituted C_{2-32} alkynyl. The substituted C_{1-32} alkyl, substituted C_{2-32} alkenyl, and substituted C_{2-32} alkynyl may be substituted with a hydroxyl group or a C_{1-5} alkyl.

[0023] In some embodiments, R_1 may be a substituted or unsubstituted C_{1-8} alkyl, a substituted or unsubstituted C_{2-8} alkenyl, or a substituted or unsubstituted C_{2-8} alkynyl. The substituted C_{1-8} alkyl, substituted C_{2-8} alkenyl, and substituted C_{2-8} alkynyl may be substituted with a hydroxyl group or a C_{1-5} alkyl.

[0024] The term “alkyl,” as used herein, refers to a linear or branched hydrocarbon radical, preferably having 1 to 32 carbon atoms (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, or 32 carbons). Examples of unsubstituted alkyl groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, s-pentyl, t-pentyl, and the like.

[0025] The term “alkenyl,” as used herein, refers to a straight or branched hydrocarbon radical, preferably having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, or 32 carbons, and having one or more carbon-carbon double bonds. Alkenyl groups include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl (allyl), iso-propenyl, 2-methyl-1-propenyl, 1-butenyl, and 2-butenyl.

[0026] The term “alkynyl,” as used herein, refers to a straight or branched hydrocarbon radical, preferably having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, or 32 carbons, and having one or more carbon-carbon triple bonds. Alkynyl groups include, but are not limited to, ethynyl, propynyl, and butynyl.

[0027] In some embodiments, the compound of formula (I) may be hexamine, which is also known as hexamethylenetetramine, methenamine, urotropine, or 1,3,5,7-tetraazatricyclo[3.3.1.1^{3,7}]decane. Hexamine is readily available from common chemical suppliers.

[0028] The composition may also include a monolignol alcohol. Monolignol alcohols may be isolated from various species of plants and trees. The monolignol alcohol may be p-coumaryl alcohol, coniferyl alcohol, sinapyl alcohol, or cinnamyl alcohol.

[0029] In certain embodiments, the composition may have about 2 to about 20 weight percent of the compound of formula (I) or formula (II). The amount of the compound of formula (I) or formula (II) in the composition may range from about 5 to about 20 weight percent, about 5 to about 15 weight percent, or about 7 to about 12 weight percent. In some embodiments, the amount of the compound of formula (I) or formula (II) may be about 7 weight percent, 8 weight percent, about 9 weight percent, about 10 weight percent, about 11 weight percent, about 12 weight percent, about 13 weight percent, about 14 weight percent, or about 15 weight percent.

[0030] Monolignol alcohols originates from certain plants, such as grass, and can be purchased from chemical suppliers at varying degrees of purity. The monolignol alcohol in the composition may include a mixture of one or more of p-coumaryl alcohol, coniferyl alcohol, sinapyl alcohol, or cinnamyl alcohol, including any other monolignol alcohols.

[0031] In other embodiments, the monolignol alcohol may be p-coumaryl alcohol.

[0032] In some embodiments, the monolignol alcohol may be coniferyl alcohol.

[0033] In some embodiments, the monolignol alcohol may be sinapyl alcohol.

[0034] In some embodiments, the monolignol alcohol may be cinnamyl alcohol.

[0035] In certain embodiments, the monolignol alcohol may be a mixture of one or more of p-coumaryl alcohol, coniferyl alcohol, sinapyl alcohol, or cinnamyl alcohol.

[0036] In some embodiments, the composition may consist of the compound of formula (I) or formula (II) and a monolignol alcohol. The composition may consist of the compound of formula (I) and cinnamyl alcohol. The composition may consist of hexamine and cinnamyl alcohol.

[0037] In some embodiments, the composition may have about 10 to about 98 weight percent of the monolignol alcohol. In other embodiments, the monolignol alcohol present in the composition may range from about 20 to about 98 weight percent, about 30 to about 98 weight percent, about 40 to about 98 weight percent, about 50 to about 98 weight percent, about 60 to about 98 weight percent, about 70 to about 98 weight percent, and about 80 to about 98 weight percent.

[0038] In other embodiments, the composition may include a C₂₋₈ polyol. A polyol may be defined as an alcohol having multiple hydroxyl groups. In some embodiments, the composition may include a C₃₋₈ polyol, a C₄₋₈ polyol, or a C₅₋₈ polyol. The composition may include a C₃ polyol.

[0039] In some embodiments, the C₂₋₈ polyol may be glycerin. Glycerin is hydroscopic and may contain water. Water content in commercially available glycerin can be as low as about 1% by weight or less. In some embodiments, the composition may include an amount of water that is less than about 1% by weight. The glycerin may be prepared such that all the water may be removed by using molecular sieves, for example. In certain embodiments, the composition may exclude water.

[0040] In some embodiments, the composition may have about 2 to about 80 weight percent of the C₂₋₈ polyol. The amount of C₂₋₈ polyol in the composition may range from about 2 to about 70 weight percent, about 2 to about 60 weight percent, about 2 to about 50 weight percent, about 2 to about 40 weight percent, about 2 to about 30 weight percent, or about 2 to about 25 weight percent.


[0041] In some embodiments, the composition may include the compound of formula (I) or formula (II), cinnamyl alcohol, and a C₂₋₈ polyol. The composition may include the compound of formula (I) or formula (II), cinnamyl alcohol, and glycerin. In other embodiments, the composition may consist of the compound of formula (I) or formula (II), cinnamyl alcohol, and a C₂₋₈ polyol. The composition may consist of hexamine, cinnamyl alcohol, and glycerin.

[0042] In some embodiments, the weight ratio of the monolignol alcohol to the C₂₋₈ polyol may be about 10:1 to about 20:1, about 1:2 to about 20:1, about 1:1 to about 20:1, or about 1:10 to about 10:1. In certain embodiments, the weight ratio of the monolignol alcohol to the C₂₋₈ polyol may be about 15:1.

[0043] In some embodiments, the weight ratio of the compound of formula (I) or formula (II) to monolignol alcohol may range from about 1:20 to about 1:1. In other embodiments the weight ratio may range from about 1:15 to about 1:1, about 1:15 to about 1:2, or about 1:15 to about 1:5. In some embodiments, the weight ratio of the compound of formula (I) or formula (II) to monolignol alcohol may be about 1:8.

[0044] In some embodiments, the composition may have a pH greater than about 7. The composition may have a pH of about 7 to about 9. Generally, it may be desirable to maintain the pH at neutral or alkaline conditions before adding to the asphalt to avoid acid hydrolysis of hexamine.

[0045] In some embodiments, a method is provided for scavenging hydrogen sulfide in an industrial process. The method may include adding an effective amount of a composition to a medium in the industrial process. The composition may include a monolignol alcohol and a compound of formula (I) or formula (II)



where R_1 is a substituted or unsubstituted C_{1-32} alkyl, a substituted or unsubstituted C_{2-32} alkenyl, or a substituted or unsubstituted C_{2-32} alkynyl, wherein the substituted C_{1-32} alkyl, the substituted C_{2-32} alkenyl, and the substituted C_{2-32} alkynyl are substituted with hydroxyl or C_{1-5} alkyl

[0046] In some embodiments, R_1 may be a substituted or unsubstituted C_{1-8} alkyl, a substituted or unsubstituted C_{2-8} alkenyl, or a substituted or unsubstituted C_{2-8} alkynyl, wherein the substituted C_{1-8} alkyl, the substituted C_{2-8} alkenyl, and the substituted C_{2-8} alkynyl may be substituted with hydroxyl or C_{1-5} alkyl.

[0047] The effective amount added to the medium in an industrial process can be readily determined by one of ordinary skill in the art. An effective amount would depend at least partially on the medium's hydrogen sulfide content. Generally, if a medium contains high levels of hydrogen sulfide then more of the composition may be required. In some embodiments, the effective amount may be in a range from about 0.00001 ppm to about 100,000 ppm.

[0048] The industrial process may utilize or process crude oil, diesel fuel, asphalt or the like. The composition may be added to a medium in the industrial process, such as crude oil, diesel fuel, asphalt, or combinations thereof in order to remove unwanted hydrogen sulfide.

[0049] The composition may be added to the process continuously or dosed intermittently. A programmed logic controller may control the addition of the composition to the process. The method may include sensors that detect levels of hydrogen sulfide. The sensors may be in communication with the PLC. Once levels of hydrogen sulfide approach or reach a predetermined threshold the composition may be added to process.

[0050] The compositions and methods disclosed herein may be especially useful for removing hydrogen sulfide from hot asphalt where the use of aqueous or volatile solvents presents safety or toxicity concerns.

Examples

[0051] Example 1 Dissolution of Hexamine in Cinnamyl Alcohol and Glycerin

[0052] Slurries were prepared by mixing hexamine in cinnamyl alcohol or a mixture of cinnamyl alcohol and glycerin at about room temperature. The compositions were then allowed to sit undisturbed for two days and the clear supernatant decanted off and analyzed for composition. As the results below illustrate, clear stable solutions of hexamine in various ratios of glycerin and cinnamyl alcohol were obtained.

Table 1. Composition of Stable Solutions by Weight%

Hexamine	Cinnamyl Alcohol	Glycerin (<1% water)
7	93	0
10	84	6
11	78	11
10	68	22
7	59	34
7	50	43
8	37	55
5	32	63
8	16	76

[0053] All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While this invention may be embodied in many different forms, there are described in detail herein specific preferred embodiments of the invention. The present disclosure is an exemplification of the principles of the invention and is not intended to limit the invention to the particular

embodiments illustrated. In addition, unless expressly stated to the contrary, use of the term “a” is intended to include “at least one” or “one or more.” For example, “a compound” is intended to include “at least one compound” or “one or more compounds.”

[0054] Any composition disclosed herein may comprise, consist of, or consist essentially of any of the compounds / components disclosed herein. In accordance with the present disclosure, the phrases “consist essentially of,” “consists essentially of,” “consisting essentially of,” and the like, limit the scope of a claim to the specified materials or steps and those materials or steps that do not materially affect the basic and novel characteristic(s) of the claimed invention.


[0055] Any ranges given either in absolute terms or in approximate terms are intended to encompass both, and any definitions used herein are intended to be clarifying and not limiting. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Moreover, all ranges disclosed herein are to be understood to encompass any and all subranges (including all fractional and whole values) subsumed therein.

[0056] Furthermore, the invention encompasses any and all possible combinations of some or all of the various embodiments described herein. It should also be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the invention and without diminishing its intended advantages. It is therefore intended that such changes and modifications be covered by the appended claims.

CLAIMS

What is claimed is:

1. A composition, comprising: a monolignol alcohol and a compound of formula (I) or formula (II)




where R₁ is a substituted or unsubstituted C₁₋₈ alkyl, a substituted or unsubstituted C₂₋₈ alkenyl, or a substituted or unsubstituted C₂₋₈ alkynyl, wherein the substituted C₁₋₈ alkyl, the substituted C₂₋₈ alkenyl, and the substituted C₂₋₈ alkynyl are substituted with hydroxyl or C₁₋₅ alkyl.

2. The composition of claim 1, wherein the monolignol alcohol is selected from the group consisting of p-coumaryl alcohol, coniferyl alcohol, sinapyl alcohol, cinnamyl alcohol, and any combination thereof.
3. The composition of claim 1 or claim 2, wherein the monolignol alcohol is cinnamyl alcohol.
4. The composition of claim 1 or claim 2, wherein the monolignol alcohol is coniferyl alcohol.
5. The composition of any one of claims 1 to 4, wherein the compound is hexamine.
6. The composition of any one of claims 1 to 5, further comprising a C₂₋₈ polyol.

7. The composition of claim 6, wherein the C₂₋₈ polyol is glycerin.
8. The composition of claim 1, comprising the compound of formula (I), cinnamyl alcohol, and glycerin.
9. The composition of any one of claims 1 to 8, wherein the composition comprises about 2 to about 20 weight percent of the compound of formula (I) or formula (II).
10. The composition of any one of claims 1 to 9, wherein the composition comprises about 10 to about 98 weight percent of the monolignol alcohol.
11. The composition of any one of claims 6 to 10, wherein the composition comprises about 2 to about 80 weight percent of the C₂₋₈ polyol.
12. The composition of any one of claims 6 to 11, wherein a weight ratio of the monolignol alcohol to the C₂₋₈ polyol is about 10:1 to about 20:1.
13. The composition of any one of claims 1 to 12, wherein a weight ratio of the compound of formula (I) or formula (II) to monolignol alcohol is about 1:15 to about 1:1.
14. The composition of any one of claims 1 to 11, wherein the composition comprises an amount of water that is less than about 1% by weight.
15. The composition of any one of claims 1 to 14, wherein the composition excludes water.
16. The composition of any one of claims 1 to 15, wherein the composition comprises a pH greater than about 7.
17. A method for scavenging hydrogen sulfide in an industrial process, comprising:

adding an effective amount of a composition to a medium in the industrial process, wherein the composition comprises a monolignol alcohol and a compound of formula (I) or formula (II)




where R_1 is a substituted or unsubstituted C_{1-8} alkyl, a substituted or unsubstituted C_{2-8} alkenyl, or a substituted or unsubstituted C_{2-8} alkynyl, wherein the substituted C_{1-8} alkyl, the substituted C_{2-8} alkenyl, and the substituted C_{2-8} alkynyl are substituted with hydroxyl or C_{1-5} alkyl.

18. The method of claim 17, further comprising a C_{2-8} polyol, and wherein the monolignol alcohol is cinnamyl alcohol.

19. The method of any one of claims 17 to 18, wherein the effective amount is in a range from about 0.00001 ppm to about 100,000 ppm.

20. The method of any one of claims 17 to 19, wherein the medium is selected from the group consisting of crude oil, diesel fuel, asphalt and any combination thereof.

21. Use of a composition in a method of scavenging hydrogen sulfide, the composition comprising: a monolignol alcohol and a compound of formula (I) or formula (II)



where R_1 is a substituted or unsubstituted C_{1-8} alkyl, a substituted or unsubstituted C_{2-8} alkenyl, or a substituted or unsubstituted C_{2-8} alkynyl, wherein the substituted C_{1-8} alkyl, the substituted C_{2-8} alkenyl, and the substituted C_{2-8} alkynyl are substituted with hydroxyl or C_{1-5} alkyl.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/024272

A. CLASSIFICATION OF SUBJECT MATTER

INV. B01D53/52 C08K5/3492 C08L95/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

B01D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	RU 2 269 567 C1 (YAKUT CONS SCI RES INST) 10 February 2006 (2006-02-10) the whole document & DATABASE WPI Week 200617 Thomson Scientific, London, GB; AN 2006-162079 & RU 2 269 567 C1 (YAKUT CONS SCI RES INST) 10 February 2006 (2006-02-10) abstract	1-21
Y	----- CN 102 140 364 A (XUN YANSEN; JIANMIN XIANG) 3 August 2011 (2011-08-03) the whole document -/-	1-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 May 2018

Date of mailing of the international search report

17/05/2018

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, Marc

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/024272

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	& DATABASE WPI Week 201214 Thomson Scientific, London, GB; AN 2011-K85600 & CN 102 140 364 A (SUN G) 3 August 2011 (2011-08-03) abstract ----- RU 2 479 615 C2 (FAKHRIEV A M) 20 April 2013 (2013-04-20) the whole document & DATABASE WPI Week 201364 Thomson Scientific, London, GB; AN 2013-G06409 & RU 2 479 615 C2 (FAKHRIEV A M) 20 April 2013 (2013-04-20) abstract	1-21
Y	----- WO 93/01126 A1 (GATLIN LARRY W [US]) 21 January 1993 (1993-01-21) claim 13	1-21
Y	----- RU 2 302 523 C1 (FAKHRIEV AKHMATFAIL MAGSUMOVIC [RU]; FAKHRIEV RUSTEM AKHMATFAILOVIC [R]) 10 July 2007 (2007-07-10) the whole document & DATABASE WPI Week 200750 Thomson Scientific, London, GB; AN 2007-511912 & RU 2 302 523 C1 (FAKHRIEV A M) 10 July 2007 (2007-07-10) abstract -----	1-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2018/024272

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
RU 2269567	C1	10-02-2006	NONE

CN 102140364	A	03-08-2011	NONE

RU 2479615	C2	20-04-2013	NONE

WO 9301126	A1	21-01-1993	AU 2319592 A 11-02-1993
			GB 2274651 A 03-08-1994
			US 5167479 A 01-12-1992
			WO 9301126 A1 21-01-1993

RU 2302523	C1	10-07-2007	NONE
