



(22) **Date de dépôt/Filing Date:** 2005/06/03

(41) **Mise à la disp. pub./Open to Public Insp.:** 2006/02/23

(62) **Demande originale/Original Application:** 2 577 447

(30) **Priorités/Priorities:** 2004/08/16 (US60/602,208);
2004/12/01 (US60/632,209)

(51) **Cl.Int./Int.Cl.** **C12N 5/0775** (2010.01),
A61K 35/28 (2015.01), **C12N 5/073** (2010.01),
C12N 5/077 (2010.01), **C12Q 1/68** (2006.01)

(71) **Demandeur/Applicant:**
CELLRESEARCH CORPORATION PTE LTD, SG

(72) **Inventeurs/Inventors:**
PHAN, TOAN-THANG, SG;
LIM, IVOR JIUN, SG

(74) **Agent:** BLAKE, CASSELS & GRAYDON LLP

(54) **Titre : ISOLEMENT DE CELLULES SOUCHES/PROGENITRICES ISSUES DE LA MEMBRANE AMNIOTIQUE DU CORDON
OMBILICAL**

(54) **Title: ISOLATION OF STEM/PROGENITOR CELLS FROM AMNIOTIC MEMBRANE OF UMBILICAL CORD**

(57) **Abrégé/Abstract:**

The present invention relates to a method for isolating stem/progenitor cells from the amniotic membrane of umbilical cord, wherein the method comprises separating the amniotic membrane from the other components of the umbilical cord in vitro, culturing the amniotic membrane tissue under conditions allowing cell proliferation, and isolating the stem/progenitor cells from the tissue cultures. The isolated stem cell cells can have embryonic stem cell- like properties and can be used for various therapeutic purposes. In one embodiment, the invention relates to the isolation and cultivation of stem cells such as epithelial and/or mesenchymal stem/progenitor cells under conditions allowing the cells to undergo mitotic expansion. Furthermore, the invention is directed to a method for the differentiation of the isolated stem/progenitor cells into epithelial and/or mesenchymal cells.

ABSTRACT

The present invention relates to a method for isolating stem/progenitor cells from the amniotic membrane of umbilical cord, wherein the method comprises separating the amniotic membrane from the other components of the umbilical cord in vitro, culturing the amniotic membrane tissue under conditions allowing cell proliferation, and isolating the stem/progenitor cells from the tissue cultures. The isolated stem cell cells can have embryonic stem cell-like properties and can be used for various therapeutic purposes. In one embodiment, the invention relates to the isolation and cultivation of stem cells such as epithelial and/or mesenchymal stem/progenitor cells under conditions allowing the cells to undergo mitotic expansion. Furthermore, the invention is directed to a method for the differentiation of the isolated stem/progenitor cells into epithelial and/or mesenchymal cells.

ISOLATION OF STEM/PROGENITOR CELLS FROM AMNIOTIC MEMBRANE OF UMBILICAL CORD

5 **[0001]** The present invention relates to a method for isolating
stem/progenitor cells from the amniotic membrane of umbilical cord, wherein
the method comprises separating the amniotic membrane from the other
components of the umbilical cord *in vitro*, culturing the amniotic membrane
tissue under conditions allowing cell proliferation, and isolating the
stem/progenitor cells from the tissue cultures. In particular, the invention relates
10 to the isolation and cultivation of stem cells having embryonic properties such
as epithelial and/or mesenchymal stem/progenitor cells under conditions
allowing the cells to undergo mitotic expansion. Furthermore, the invention is
directed to a method for the differentiation of the isolated stem/progenitor cells
into epithelial and/or mesenchymal cells and therapeutic uses of these
15 stem/progenitor cells:

BACKGROUND OF THE INVENTION

20 **[0002]** Stem cells are a cell population possessing the capacities to self-
renew indefinitely and to differentiate in multiple cell or tissue types. Embryonic
stem cells (from approximately days 3 to 5 after fertilisation) proliferate
indefinitely and can differentiate spontaneously into all tissue types: they are
thus termed pluripotent stem cells (reviewed, for example, in Smith, A.G. (2001)
Annu. Rev. Cell. Dev. Biol. **17**, 435-462). Adult stem cells, however, are more
25 tissue-specific and may have less replicative capacity: they are thus termed
multipotent stem cells (reviewed, for example, in Paul, G. et al. (2002) *Drug
Discov. Today* **7**, 295-302). The "plasticity" of embryonic and adult stem cells
relies on their ability to trans-differentiate into tissues different from their origin
and, perhaps, across embryonic germ layers.

30 **[0003]** The ability of stem cells to self-renew is critical to their function as
reservoir of primitive undifferentiated cells. In contrast, most somatic cells have
a limited capacity for self-renewal due to telomere shortening (reviewed, for
example, in Dice, J.F. (1993) *Physiol. Rev.* **73**, 149-159). Stem cell-based
therapies thus have the potential to be useful for the treatment of a multitude of

human and animal diseases.

[0004] Stem cells as well as stem/progenitor cells can be derived from different sources. The "multi-lineage" potential of embryonic and adult stem cells has been extensively characterized. Even though the potential of embryonic stem cells is enormous, their use implies many ethical problems. Therefore, non-embryonic stem cells derived from the bone marrow stroma, fat tissue, dermis and umbilical cord blood have been proposed as alternative sources. These cells can differentiate *inter alia* into chondrocytes, adipocytes, osteoblasts, myoblasts, cardiomyocytes, astrocytes, and tenocytes *in vitro* and undergo differentiation *in vivo*, making these stem cells - in general referred to as mesenchymal stem cells - promising candidates for mesodermal defect repair and disease management.

[0005] In clinical use, however, harvesting of such mesenchymal stem cells causes several problems. The collection of the cells is a mental and physical burden to the patient as a surgical procedure is required to obtain the cells (for example, the collection of bone marrow is an invasive technique performed with a biopsy needle that requires local or even general anesthesia). Furthermore, in many cases the number of stem cells extracted is rather low. More importantly, no epithelial cells are derived or differentiated from these cells. This prompted the search for other possible sources of stem cells.

[0006] Umbilical cord blood has been identified as a rich source of haematopoietic stem/progenitor cells. However, the existence of mesenchymal stem/progenitor cells is discussed controversially. On the one hand, such cells could not be isolated or successfully cultured from term umbilical cord blood (Mareschi, K. et al. (2001) *Haematologica* 86, 1099-1100). At the same time, results obtained by Campagnoli, C. et al. (*Blood* (2001) 98, 2396-2402) as well as Erices, A. et al. (*Br. J. Haematol.* (2000) 109, 235-242) suggest that mesenchymal stem cells are present in several fetal organs and circulate in the blood of pre-term fetuses simultaneously with hematopoietic precursors. Accordingly, International Patent Application WO 03/070922 discloses isolation and culture-expansion methods of mesenchymal stem/progenitor cells from umbilical cord blood and a differentiation method of such cells into various mesenchymal tissues. Isolation efficiencies of about 60% have been reported (Bieback, K. et al. (2004) *Stem Cells* 22, 625-634). In the same study, both the

time period from collection of the umbilical cord blood to isolation of the cells and the volume of the blood sample used have been determined as crucial parameters for achieving such a yield. However, it is still a matter of debate whether these stem/progenitor cells are indeed derived of umbilical cord tissue.

5 **[0007]** Recently, mesenchymal stem/progenitor cells have been successfully isolated from umbilical cord tissue, namely from Wharton's jelly, the matrix of umbilical cord, (Mitchell, K.E. et al. (2003) *Stem Cells* **21**, 50-60; U.S. Patent 5,919,702; US Patent Application 2004/0136967). These cells have been shown to have the capacity to differentiate, for example, into a neuronal
10 phenotype and into cartilage tissue, respectively. Furthermore, mesenchymal stem/progenitor cells have also been isolated from the endothelium and the subendothelial layer of the umbilical cord vein, one of the three vessels (two arteries, one vein) found within the umbilical cord (Romanov, Y.A. et al. (2003) *Stem Cells* **21**, 105-110; Covas, D.T. et al. (2003) *Braz. J. Med. Biol. Res.* **36**,
15 1179-1183).

[0008] However, none of these approaches employed thus far has resulted in the isolation or cultivation of epithelial stem/progenitor cells as a source for epithelial cell-based therapies such as skin resurfacing, liver repair, bladder tissue engineering and other engineered surface tissues. Thus, there is
20 still a need for methods and reliable sources useful for the isolation and cultivation of epithelial stem/progenitor cells. Furthermore, rapid and efficient methods which are ethically acceptable and do not pose a biomedical burden on the patient for the isolation of epithelial and mesenchymal stem/progenitor cells are still required in order to provide such cells in a sufficient amount for
25 various applications in regenerative medicine and tissue engineering.

SUMMARY OF THE INVENTION

[0009] The invention provides a method for isolating stem/progenitor
30 cells from the amniotic membrane of umbilical cord, the method comprising:

- (a) separating the amniotic membrane from the other components of the umbilical cord *in vitro*;
- (b) culturing the amniotic membrane tissue obtained in step (a) under conditions allowing cell proliferation; and

(c) isolating the stem/progenitor cells.

[0010] In one embodiment, the invention provides a method, further comprising:

5 (a") separating the cells from the amniotic membrane tissue before cultivation by a technique selected from the group consisting of enzymatic digestion and direct tissue explant.

[0011] In one preferred embodiment, the invention provides a method for isolating stem/progenitor cells that have embryonic stem cell-like properties.

10 **[0012]** In another preferred embodiment, the invention provides a method for isolating epithelial and/or mesenchymal stem/progenitor cells.

[0013] In another embodiment, the invention provides a method further comprising:

15 (d) culturing the stem/progenitor cells under conditions allowing the cells to undergo clonal expansion.

[0014] In yet another embodiment, the invention provides a method further comprising:

20 (e) culturing the stem/progenitor cells under conditions allowing the differentiation of said cells into epithelial cells and/or mesenchymal cells; and
(f) isolating the differentiated cells.

[0015] In yet another embodiment, the invention provides a method, further comprising:

25 (g) preserving the isolated stem/progenitor cells for further use.

[0016] In yet a further embodiment, the invention comprising a method of cultivating stem/progenitors cells of the invention, comprising:

Obtaining a tissue explant from the amniotic membrane of umbilical cord;

Cultivating the tissue explant in suitable cultivation media and cultivation conditions over a suitable period of time.

30 In yet other embodiments, the invention is directed to therapeutic uses of the stem/progenitor cells or cellular extracts thereof. One of these embodiments provide a method of treating a subject having a disorder comprising administering to the subject an effective amount of a stem/progenitor cell isolated by the inventive method of explained above. Another embodiment

provides a corresponding pharmaceutical composition.

BRIEF DESCRIPTION OF THE DRAWINGS

5 **[0017]** The invention will be better understood with reference to the detailed description when considered in conjunction with the non-limiting examples and the drawings, in which:

10 **[0018]** **Figure 1** depicts epithelial cell outgrowth from umbilical cord amniotic membrane by the method of direct tissue explant (40 x magnification) at day 2 (**Fig. 1A**) and day 5 (**Fig. 1B, C**) of tissue culture. Cell culture plastic surfaces were coated with collagen 1/collagen 4 mixtures (1:2; Becton Dickinson) before placing the amniotic membrane on the surface. The amniotic membrane specimens were submerged in 5 ml EpiLife medium or Medium 171 (both from Cascade Biologics). Medium was changed every 2 or 3 days and cell
15 outgrowth by explant was monitored under light microscopy. Microphotographs were taken at different time intervals as stated above. The observed polyhedral cell morphology is typical of epithelial cells.

20 **[0019]** **Figure 2** depicts enzymatic digestion of the umbilical cord segments yielding similar epithelial (40 x magnification) cells at day 2 (**Fig. A, C**) and day 5 (**Fig. B, D**). Umbilical cord amniotic membrane was divided into small pieces of 0.5 cm x 0.5 cm and digested in 0.1% (w/v) collagenase type 1 solution (Roche Diagnostics) at 37°C for 8 hours. The samples were vortexed every 30 min for 3 min. Cells were harvested by centrifugation at 4000 rpm for 30 min. Cell pellets were resuspended in EpiLife medium or Medium 171 (both
25 from Cascade Biologics) supplemented with 50 µg/ml insulin-like growth factor-1 (IGF-1), 50 µg/ml platelet-derived growth factor-BB (PDGF-BB), 5 µg/ml transforming growth factor-β1 (TGF-β1) and 5 µg/ml insulin (all obtained from R&D Systems), counted and seeded on 10 cm tissue culture dishes pre-coated with collagen 1/collagen 4 mixtures (1:2; Becton Dickinson) at density of 1×10^6
30 cells/dish. After 24 hours, attached cells were washed with warm phosphate buffered saline (PBS) and the culture medium was replaced with EpiLife medium or Medium 171 (both from Cascade Biologics). The medium was changed every 2 or 3 days, and cell outgrowth was monitored under light microscopy. Microphotographs were taken at different time intervals as stated

above. Once again the cells demonstrated typical epithelial cell polyhedral morphology.

[0020] **Figure 3** depicts outgrowing mesenchymal cells explanted from umbilical cord amniotic membrane. Cellular outgrowth was observed as early as 48 hours after placement in tissue culture dishes using DMEM supplemented with 10% fetal calf serum (FCS) as culture medium (40 x magnification) (**Fig. 3A, C**). The explants were submerged in 5 ml DMEM (Invitrogen) supplemented with 10% fetal bovine serum (Hyclone) (DMEM/10% FBS). Medium was changed every 2 or 3 days. Cell outgrowth was monitored under light microscopy. Microphotographs were taken at different time intervals. The cells were characterized by their spindle shaped morphology, and migrated and expanded both easily and quickly *in vitro*, closely resembling fibroblasts (**Fig. 3B, D**).

[0021] **Figure 4** (40 x magnification) depicts mesenchymal cells from umbilical cord amniotic membrane cells isolated by collagenase enzymatic digestion. **Fig. 4A** shows mesenchymal cells isolated from umbilical cord amniotic membrane at day 2. Cell proliferation was observed at day 5 (**Fig. 4B**). Umbilical cord amniotic membrane was divided into small pieces of 0.5cm x 0.5cm and digested in 0.1% (w/v) collagenase type1 solution (Roche Diagnostics) at 37°C for 6 hours. The samples were vortexed every 15 min for 2 min. Cells were harvested by centrifugation at 4000 rpm for 30 min. Cell pellets were resuspended in DMEM/10% FBS, counted and seeded on 10cm tissue culture dish at density of 1×10^6 cells/dish. Medium was changed every 2 or 3 days. Cell outgrowing was monitored under light microscopy. Microphotographs were taken at different time intervals. Once again, cells demonstrated spindle shaped morphology typical of mesenchymal cells as fibroblasts.

[0022] **Figure 5** (40 x magnification) depicts the morphology in serum-free culture condition (DMEM) and serum culture condition (DMEM/10%FCS) of umbilical cord amniotic membrane mesenchymal cells (UCMC, **Fig. 5E, F, G, H**) isolated according to the method of the invention, normal dermal fibroblasts (NF109 cells, **Fig. 5A, B**) and adipose-derived mesenchymal cells (ADMC, **Fig. 5C, D**). **Fig. 5** shows changes in cell morphology of NF and ADCMC cultured in serum starvation conditions (DMEM only) reflected by flatter cells and less dense cytoplasm as compared with serum rich conditions (DMEM/10%FCS)

where cells are more rounded with a dense cytoplasm (Fig. 5A, B, C, D). No change in morphology was observed in both UCMC groups cultured under identical conditions of serum-free vs. serum rich media (Fig. 5E, F, G, H), indicating a difference in behavior and physiology of these latter mesenchymal cells.

[0023] Figure 6 (40 x magnification) depicts UCMC isolated according to the invention cultured in DMEM/10%FCS at days 3 and 7 without a 3T3 feeder layer. The cells are seen to be growing well, and are forming a colony (vertical growth) instead of exhibiting radial spread. Once again, this indicates a difference in behavior of these mesenchymal cells as compared to their more differentiated counterparts.

[0024] Figure 7 (40 x magnification) depicts colony formation of umbilical cord epithelial cells (UCEC) cultured on a 3T3 feeder layer at days 3 and 7. This appearance is similar to that of normal skin derived epithelial keratinocyte stem cells. In the latter, the 3T3 feeder layer maintains stemness of the cells.

[0025] Figure 8 (40 x magnification) depicts obvious colony formation of umbilical cord mesenchymal cells (UCMC) isolated according to the invention cultured on a 3T3 feeder layer at days 3 and 7. The 3T3 feeder layer normally suppresses the growth of differentiated mesenchymal cells as human dermal fibroblasts. Once again, this indicates a difference in behavior of these mesenchymal cells as compared to their more differentiated counterparts.

[0026] Figure 9-1 to Figure 9-26 show Western blot analysis by which the expression of several embryonic stem cell markers in UCEC and UCMC isolated according to the invention, was compared to the expression of these markers in human dermal fibroblasts (NF), in bone marrow mesenchymal cells (BMSC) and adipose-derived mesenchymal cells (ADMC).

[0027] Figure 10 shows indirect immunofluorescent analysis of markers of epithelial cells expressed in umbilical cord epithelial stem cells such as cytokeratins (CK)-general, CK17, CK6, CK10, CK19, CK18, CK16, CK15

(**Fig.10-1**); Hemidesmosome components-integrin alpha6, integrin beta4; Desmosome components (**Fig.10-2**); Basement membrane components-laminin1, laminin5, collagen IV, collagen VII (**Fig.10-3**) and other important extracellular matrix components as integrin-beta1 and fibronectin (**Fig.10-4**).

5 **[0028] Figure 11** shows cytokine array analysis of secreted cytokines and growth factors by umbilical cord mesenchymal stem cells (UCMC) in comparison with human bone-marrow mesenchymal stem cells.

10 **[0029] Figure 12** shows cytokine array analysis of secreted cytokines and growth factors by umbilical cord epithelial stem cells (UCEC) in comparison with human epidermal keratinocytes.

15 **[0030] Figure 13** shows UCMC cells cultured in DMEM supplemented with 10 % fetal calf serum (FCS) (**Fig.13-1**), serum-free media PTT-1 (**Fig.13-2**), in serum-free media PTT-2 (**Fig.13-3, Fig.13-4**) and in serum-free media PTT-3 (**Fig.13-5**). **Fig. 13** also shows the growth of adipose derived stromal cells (**Fig.13-6**) and bone marrow derived stromal cells (**Fig.13-7**) in serum free medium PTT-3.

20 **[0031] Figure 14** shows global gene expression in umbilical cord epithelial and mesenchymal stem cells analyzed by DNA microarray. UCEC expressed a total of 28055 genes and UCMC expressed a total of 34407 genes. There are 27308 overlapping genes expressing in both cell types. 747 genes expressed were unique to UCEC and 7099 genes expressed were unique to UCMC. The selected genes of interest are presented in this Figure. Both stem cell types expressed 140 genes related to embryonic stem cells and embryonic development.

25 **[0032] Figure 15** shows a schematic illustration of expansion of umbilical cord epithelial and mesenchymal stem cells using repetitive explants of umbilical cord lining membrane tissues.

30 **[0033] Figure 16** depicts a cross section of an umbilical cord demonstrating the umbilical cord amniotic lining membrane (LM), the contained Wharton's jelly (WJ), as well as two umbilical arteries (UA) and one umbilical vein (UV) supported within this jelly.

DETAILED DESCRIPTION

[0034] The invention is based on the surprising finding that the amniotic membrane of umbilical cord represents a source, from which stem/progenitor cells such as mesenchymal and epithelial stem/progenitor cells can be successfully isolated and expanded under *in vitro* conditions. Even more surprising is the finding that these cells show embryonic stem cell-like characteristics. The amniotic membrane (also called amniotic lining membrane), i.e. thin innermost membranous sac enclosing the placenta and developing embryo of mammals, has recently been used as a natural substrate in ocular surface reconstruction and as a biological substrate for expanding limbal epithelial stem cells (cf., e.g., Anderson, D.F. et al. (2001) *Br. J. Ophthalmol.* **85**, 567-575; Grüterich, M. et al. (2003) *Surv. Ophthalmol.* **48**, 631-646). However, no methods have been described thus far for the isolation of stem/progenitor cells from the amniotic membrane, at least for humans, nor has the amniotic membrane covering the umbilical cord been reported as a source for stem cells.

[0035] The invention provides a method for isolating stem/progenitor cells from the amniotic membrane of umbilical cord, the method comprising:

- (a) separating the amniotic membrane from the other components of the umbilical cord *in vitro*;
- (b) culturing the amniotic membrane tissue obtained in step (a) under conditions allowing cell proliferation; and
- (c) isolating the stem/progenitor cells.

[0036] The term "stem/progenitor cell" as used herein refers to any cell derived of umbilical cord having the capacities to self-renew indefinitely and to differentiate in multiple cell or tissue types such as endothelial cells, epithelial cells, fibroblasts, myocytes or neurons. Furthermore, the cells may be derived of any mammalian species, such as mouse, rat, guinea pig, rabbit, goat, dog, cat, sheep, monkey or human, with cells of human origin being preferred in one embodiment.

[0037] The term "embryonic stem cell-like properties" refers to the ability of the cells derived of umbilical cord that they can - almost like or exactly like embryonic stem cells - differentiate spontaneously into all tissue types, meaning that they are pluripotent stem cells.

[0038] The term "amniotic membrane" as used herein refers to the thin innermost membranous sac enclosing the developing embryo of mammals.

During pregnancy, the fetus is surrounded and cushioned by a liquid called amniotic fluid. This fluid, along with the fetus and the placenta, is enclosed within a sac called the amniotic membrane, which also covers the umbilical cord. The amniotic fluid is important for several reasons. It cushions and protects the fetus, allowing the fetus to move freely. The amniotic fluid also allows the umbilical cord to float, preventing it from being compressed and cutting off the fetus' supply of oxygen and nutrients derived from the circulating blood within the placental blood vessels. The amniotic sac contains the amniotic fluid which maintains a homeostatic environment protecting the fetal environment from the outside world. This barrier additionally protects the fetus from organisms (like bacteria or viruses) that could travel up the vagina and potentially cause infection.

[0039] Media and reagents for tissue culture are well known in the art (cf., for example, Pollard, J.W. and Walker, J.M. (1997) *Basic Cell Culture Protocols, Second Edition*, Humana Press, Totowa, NJ; Freshney, R.I. (2000) *Culture of Animal Cells, Fourth Edition*, Wiley-Liss, Hoboken, NJ). Examples of suitable media for incubating/transporting umbilical cord tissue samples include, but are not limited to, Dulbecco's Modified Eagle Medium (DMEM), RPMI media, Hanks' Balanced Salt Solution (HBSS) phosphate buffered saline (PBS), and L-15 medium, with the latter one being preferred in some embodiments. Examples of appropriate media for culturing stem/progenitor cells according to the invention include, but are not limited to, Dulbecco's Modified Eagle Medium (DMEM), DMEM-F12, RPMI media, EpiLife medium, and Medium 171, with the latter being preferred in some embodiments. The media may be supplemented with fetal calf serum (FCS) or fetal bovine serum (FBS) as well as antibiotics, growth factors, amino acids, inhibitors or the like, which is well within the general knowledge of the skilled artisan.

[0040] In one embodiment, the invention provides a method, further comprising:

(a") separating these stem/progenitor cells from the amniotic membrane tissue by a enzymatic digestion and/or direct tissue explant technique before cultivation. The term "enzymatic digestion technique" as used herein means that enzymes are added to cleave the cells from the main tissue mass (here the amniotic membrane of the umbilical cord). The separated cells

are subsequently collected. The term "direct tissue explant technique" as used herein means that the tissue is first placed in media without enzymes. Then under careful conditions the cells separate from the main tissue mass by itself- and the cells are then harvested for collection.

5 **[0041]** Methods for separating cells of a particular tissue or organ by treatment with enzymes or by direct tissue explant are well known in the art (cf., for example, Pollard, J.W. and Walker, J.M. (1997) *Basic Cell Culture Protocols, Second Edition*, Humana Press, Totowa, NJ; Freshney, R.I. (2000) *Culture of Animal Cells, Fourth Edition*, Wiley-Liss, Hoboken, NJ). Any enzyme catalyzing
10 tissue dissociation may be used for performing the methods of the present invention. In preferred embodiments, collagenase is used for that purpose. The enzyme may be used as a crude preparation or in purified form. It may be purified from any prokaryotic or eukaryotic organism (with *Clostridium histolyticum* being most preferred) or produced recombinantly by means of gene
15 technology. Any type of collagenase may be employed, i.e. type 1, type 2, type 3, type 4, or any combination thereof. In some embodiments the use of collagenase type 1 is being preferred.

[0042] In one embodiment, the invention provides a method for isolating stem/progenitor cells that have embryonic stem cell-like properties. These cells
20 can ultimately be differentiated into, but not limited to, by morphology, epithelial or mesenchymal cells.

[0043] Accordingly, in another embodiment, the invention provides a method for isolating epithelial and/or mesenchymal stem/progenitor cells, wherein in accordance with the above disclosure these cells may have
25 embryonic stem cell-like properties.

[0044] Epithelial stem/progenitor cells include any cells exhibiting a epithelial cell like morphology (i.e. a polyhedral shape) that can be differentiated into any type of epithelial cell such as, but not limited to, skin epithelial cells, hair follicular cells, cornea epithelial cells, conjunctival epithelial cells, retinal
30 epithelial cells, liver epithelial cells, kidney epithelial cells, pancreatic epithelial cells, oesophageal epithelial cells, small intestinal epithelial cells, large intestinal epithelial cells, lung and airway epithelial cells, bladder epithelial cells or uterine epithelial cells.

[0045] Mesenchymal stem/progenitor cells include any cells exhibiting a

mesenchymal cell like morphology (i.e. a spindle-like shape) that can be differentiated into any type of mesenchymal cell such as, but not limited to, skin fibroblasts, chondrocytes, osteoblasts, tenocytes, ligament fibroblasts, cardiomyocytes, smooth muscle cells, skeletal muscle cells, adipocytes, cells
 5 derived from endocrine glands, and all varieties and derivatives of neurectodermal cells.

[0046] In another embodiment, the invention provides a method further comprising:

(d) culturing the stem/progenitor cells under conditions allowing the
 10 cells to undergo clonal expansion.

[0047] The term "clonal expansion" (sometimes also referred to as "mitotic clonal expansion") relates to a process that occurs early in the differentiation program of a cell, by which stem/progenitor cells become committed to a particular lineage and then undergo terminal differentiation. It is
 15 well known in the art that the conditions to induce clonal expansion of progenitor cells may vary significantly between different cell types. Without being limited to a particular method, the induction of clonal expansion is generally achieved by cultivating the stem/progenitor cells in a medium that has been optimized for cell proliferation. Such media are commercially available from many providers.
 20 Non-limiting examples of such media are KGM[®]-Keratinocyte Medium (Cambrex), MEGM-Mammary Epithelial Cell Medium (Cambrex), EpiLife medium (Cascade Biologics) or Medium 171 (Cascade Biologics). Alternatively, a culture medium may be supplemented with reagents inducing cell proliferation such as growth factors. Such reagents may be admixed in a single solution
 25 such as the Human Keratinocyte Growth Supplement Kit (Cascade Biologics), to name one example, or may be supplemented individually. Such reagents include, but are not limited to, growth factors (such as epidermal growth factor, insulin-like growth factor-1, platelet-derived growth factor-BB, transforming growth factor- β 1, insulin, for example), hormones (such as a bovine pituitary
 30 extract), hydrocortisone, transferrin and the like in any suitable combination to induce clonal expansion of a given cell type. The term "clonal expansion" also includes cultivation of the cell *in vivo*, for example, by injection of the cells into mammals such as humans, mice, rats, monkeys, apes to name only a few.

[0048] In yet another embodiment, the invention provides a method

further comprising:

- (e) culturing the stem/progenitor cells under conditions allowing the differentiation of said cells into epithelial cells and/or mesenchymal cells; and
- (f) isolating the differentiated cells.

5 **[0049]** In yet another embodiment, the invention provides a method, further comprising:

- (g) preserving the isolated stem/progenitor cells for further use.

10 **[0050]** Methods and protocols for preserving and storing of eukaryotic cells, and in particular mammalian cells, are well known in the art (cf., for example, Pollard, J.W. and Walker, J.M. (1997) *Basic Cell Culture Protocols, Second Edition*, Humana Press, Totowa, NJ; Freshney, R.I. (2000) *Culture of Animal Cells, Fourth Edition*, Wiley-Liss, Hoboken, NJ). Any method maintaining the biological activity of the isolated epithelial or mesenchymal stem/progenitor cells may be utilized in connection with the present invention. In one preferred
15 embodiment, the stem/progenitor cells are maintained and stored by using cryo-preservation.

20 **[0051]** Accordingly, the invention is also directed to a progenitor/stem cell derived from the amniotic membrane of umbilical cord by means of the above methods. In addition, the invention is also directed to a cell bank comprising or consisting of one or more progenitor/stem cells that have been isolated as described here. This cell bank of progenitor/ stem cells may be autologous to an individual or pooled, and subsequently can be employed by further differentiation for regenerative medicine, tissue repair and regeneration, for example.

25 **[0052]** In accordance with the above, the invention is also directed to a pharmaceutical composition comprising a stem/progenitor cell isolated from the amniotic membrane of umbilical cord by the above inventive method. The pharmaceutical composition can be of any kind, and usually comprises the stem/progenitor cells or a cellular extract thereof together with a suitable
30 therapeutically acceptable carrier/excipient. In some embodiments, the pharmaceutical composition is adapted for systemic or topical application.

[0053] A pharmaceutical composition adapted for topical application may be in liquid or viscous form. Examples thereof include an ointment, a cream, and a lotion and the like. Examples for pharmaceutical compositions that are

suitable for systemic use are liquid compositions, wherein the stem/progenitor cells or the cellular extract are dissolved in a buffer that is acceptable for injection or infusion, for example.

5 **[0054]** Accordingly, the invention also relates to a method of treating a subject having a disorder. This method comprises administering to the subject an effective amount either of a stem/progenitor cell isolated as explained herein or of a cellular extract derived from such a cell.

10 **[0055]** In principle, any condition which is suitable for being treated by means of stem cells/progenitor cells can be treated with a cell or a cellular extract of present invention. In some embodiments, the disorder is selected from the group consisting of neoplastic disease, accelerated skin aging and skin disorders, tissue disorders, visceral endocrine deficiencies, and neural disorders.

15 **[0056]** The tissue disorder to be treated can be a congenital or an acquired tissue deficiency. Examples of visceral endocrine deficiency that can be treated with a cell of the invention include, but are not limited to, Diabetes mellitus associated with insulin deficiency, testosterone deficiency, anemia, hypoglycemia, hyperglycemia, pancreatic deficiency, adrenal deficiency, and thyroid deficiencies.

20 **[0057]** Examples of neural disorders that can be treated include, but are not limited to, Alzheimer's disease, Parkinson's disease, Jacob Kreutzfeld's disease, Lou Gehrig's disease, Huntington's disease and neural neoplastic conditions.

25 **[0058]** An example of a skin disease is a wound or a damaged part of the skin, for example, sun burned skin. Also aging of the skin is considered to be a skin disease herein. Topical or similar delivery of stem/progenitor cells of the invention or cellular extracts thereof, for example, as a constituent in lotions or creams or any other suitable vehicle may thus be used for repair of sun damaged skin and in addition may slow also down the aging process of skin
30 (anti-aging properties) by replenishing, and thus fortifying, deficient growth factors and related peptide elements, without which skin aging would be accelerated. The stem/progenitor cells may also migrate to injured regions of the body such as surface wounds to form the necessary required cellular elements necessary for the local reparative processes (cf. *The Journal of*

Immunology, 2001, 166: 7556–7562; or *International Journal of Biochemical and Cell Biology* 2004; 36: 598-606.

[0059] The neoplastic disease may be cancer, in particular as recent studies have demonstrated that stem cells may selectively target neoplastic tumor tissue (*Journal of the National Cancer Institute* 2004; 96 (21): 1593-1603) allowing for directed delivery of antineoplastic agents such as interferon to neoplastic foci. The cancer can be any kind of cancer, including those cancers that are able to form solid tumors, ranging from skin cancer to cancer of the internal organs. Examples of cancers to be treated include, squamous cell carcinoma, breast ductal and lobular carcinoma, hepatocellular carcinoma, nasopharyngeal carcinoma, lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular malignant melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, testicular cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, non-Hodgkin's lymphoma, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemias, solid tumors of childhood, lymphocytic lymphoma, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer or any combination of such cancers, including disseminated (metastasising) forms thereof. In case of treatment of a neoplastic disease the umbilical cord amnion derived stem cells and/or their cellular extracts disclosed herein can be administered systemically both as a direct treatment and/or as a carrier vehicle. In the latter case of anti-neoplastic tumor therapy, the cells comprise an anti-neoplastic agent.

[0060] In another pharmaceutical use, stem/progenitor cells of the present invention can be used for gene therapy. For this purpose, the cells can be transformed with a nucleic acid encoding the protein that is to be produced in the cells. The nucleic acid can be introduced into a cells of the invention using

any of the various methods that are well known to the skilled person, for example, using a viral vector and/or a lipid containing transfection composition such as as IBAfect (IBA GmbH, Göttingen, Germany), Eugene (Roche), GenePorter (Gene Therapy Systems), Lipofectamine (Invitrogen), Superfect
 5 (Qiagen), Metafecten (Biontex) or those ones described in the PCT application WO 01/015755). In a related embodiment, the cells of the invention, after being transformed with a nucleic acid encoding a polypeptide of choice, can be used of recombinantly producing this polypeptide.

[0061] As mentioned above, stem cell extracts are rich in a variety of
 10 growth factors and peptides that are relevant for normal tissue physiology. Such growth factors and/or peptides may be deficient in exposed parts of the body, such as the skin, which is the surface layer of all human beings protecting the body from external elements for the maintenance of internal homeostasis. Therefore in a further embodiment, stem/progenitor cells of the invention or
 15 cellular extracts thereof are suitable for the treatment and/or maintenance of internal homeostasis.

[0062] In a further embodiment and in line with the above disclosure, the stem/progenitor cells of the invention can be used for the production of any biological molecule. The biological molecule can be, for instance, any molecule
 20 that is naturally produced in the cells or a molecule the coding nucleic acid of which has been introduced into the cells via recombinant DNA technology. Examples of molecules that can be produced by the cells of the invention include, but are not limited to, a protein such as a cytokine, a growth factor such as insulin-like growth factor (IGF), epidermal growth factor (EGF), transforming
 25 growth factor beta (TGF-beta), Activin A, a bone morphogenetic protein (BMP), PDGF or a hormone as insulin or erythropoietin or a transporter protein such as transferrin, a peptide such a growth factor or hormone (e.g. luteinizing hormone (LH), follicle stimulating hormone (FSH)), a small organic molecule such as a steroid hormone, an oligo- or polysaccharide, for example, heparin or heparan
 30 sulfate (cf., example WO 96/23003, or WO 96/02259 in this regard), a proteoglycan, a glycoprotein such as collagen or laminin, or a lipid, to name only a few.

[0063] In a further aspect and in accordance with recent approaches (see, for example, Amit, M et al., Human feeder layers for human embryonic

stell cells, Biol Reprod 2003; 68: 2150-2156), the stem/progenitor cells described here can be used as feeder layer for the cultivation of other embryonic stem cells, in particular human embryonic stem cells. In one of these embodiments the cells of the present invention are preferably of human origin, since using human cells as feeder layer minimizes the risk of contaminating the cell culture with animal-derived components such as animal pathogens or immunogens. In this respect, it is to be noted that the cells of the invention can be cultivated under serum free conditions. Accordingly, employing the cells as feeder layer and cultivating the cell culture under with serum free media as the one described herein later, or in Draper et al. (Culture and characterization of human embryonic stem cell lines, Stem Cells Dev 2004, 13:325-336) or in the International patent application WO 98/30679, for example.

[0064] In this connection, it is noted that in transplantation surgery and cell-based therapy high quantities of low passage cells with a minimal proportion of senescent cells (i.e., large proportion of high quality cells) are crucial and are required to be derived within the shortest possible time during cell expansion. For example, mesenchymal stem cells from bone marrow and cord blood are low in quantity and therefore require expansion over many passages for a long period of time in order to achieve the sufficient number of cells required for cell transplant. The high passage cells however tend to deteriorate in quality and may lead to cell senescence or cancerous transformation.. It has been found here that high quantities of cells of the present invention can be obtained by low passage numbers using a repetitive explantation technique. The present invention thus also relates to a method of cultivating stem/progenitors cells of the invention, wherein this method comprises:

Obtaining a tissue explant from the amniotic membrane of umbilical cord;
Cultivating the tissue explant in suitable cultivation media and cultivation conditions over a suitable period of time,
Optionally exposing the tissue explant to fresh cultivation media and continuing the cultivation under suitable conditions over a suitable period of time (cf., Fig. 15).

[0065] The cultivation can be carried out in for as many cycles (passages) as wanted and be stopped once the desired number of cells has

been obtained. Exposing the tissue explant to fresh cultivation can be carried out by removing the used cell cultivation medium from the vessel used for growing the cells and adding fresh media to that vessel. Instead of replacing the media in the used vessel, exposing to fresh cultivation media can be achieved
 5 by transferring the tissue explant to a new vessel which is filled with cultivation media. The tissue explant used for cultivation/propagation of the cells can be obtained by any suitable method, for example by the "direct tissue explant technique" as explained above (in which the tissue is first placed in media without enzymes, and then under careful conditions the cells separate from the
 10 main tissue mass by itself- and the cells are then harvested for collection)..

[0066] The cultivation of the tissue explants can be carried out in any media that is suitable for cultivation of mammal cells. Examples include the conventional and commercially available media that are given above with respect to the cultivation or the clonal expansion of the cells of the invention
 15 such as, but not limited to, KGM®-Keratinocyte Medium (Cambrex), MEGM-Mammary Epithelial Cell Medium (Cambrex) EpiLife medium (Cascade Biologics), Medium 171 (Cascade Biologics), DMEM, DMEM-F12 or RPMI media. The cultivation is typically carried out at conditions (temperature, atmosphere) that are normally used for cultivation of cells of the species of
 20 which the cells are derived, for example, at 37°C in air atmosphere with 5%CO₂. In one embodiment, the cultivation is carried out using serum free, in particular bovine serum free media. The cultivation (in one passage) is performed for any suitable time the cells need for growth, typically, but by no means limited to, for about 1 to several days, for example to about 7 or about 8 days.

[0067] The inventions illustratively described herein may suitably be
 25 practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "comprising", "including", "containing", etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as
 30 terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed

by preferred embodiments and optional features, modification and variation of the inventions embodied therein herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention.

5 **[0068]** The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material
10 is specifically recited herein.

[0069] Other embodiments are within the following claims and non-limiting examples. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or
15 subgroup of members of the Markush group.

EXAMPLES

Example 1: Collection of umbilical cord tissue

20 **[0070]** Umbilical cord tissue is collected immediately after delivery of the child. The specimen is rinsed clean and immediately transferred into a 500ml sterile glass bottle containing culture transport medium (L-15 medium supplemented with 50 IU/ml penicillin, 50 µg/ml streptomycin, 250 µg/ml fungizone, 50 µg/ml gentamicin; all reagents purchased from Invitrogen) prior to
25 transport to the laboratory. In the laboratory, stem cell extraction is conducted in a laminar flow hood under sterile conditions. The specimen is first transferred to a sterile stainless steel tray. All remaining blood in the cord vessels is removed by multiple syringing washes using warm phosphate-buffered saline (PBS) supplemented with 5 IU/ml heparin (from Sigma). Plain PBS without heparin is
30 used in the final washes. The umbilical cord tissue specimen is then cut into pieces 2cm in length and transferred into 10 cm diameter cell culture dishes, where further washing and disinfection is performed with 70% ethanol followed by multiple washes using PBS containing an antibiotic mixture (50 IU/ml penicillin, 50 µg/ml streptomycin, 250 µg/ml fungizone, 50 µg/ml gentamicin; all

purchased from Invitrogen) until the solution becomes clear.

Example 2: Cell separation/cultivation

5 **[0071]** Dissection of umbilical cord tissue is first performed to separate the umbilical cord amniotic membrane from Wharton's jelly (i.e. the matrix of umbilical cord) and other internal components. The isolated amniotic membrane is then cut into small pieces (0.5 cm x 0.5 cm) for cell isolation. Explant is performed by placing the pieces of umbilical cord amniotic membrane on tissue culture dishes at different cell culture conditions for isolation of either epithelial or mesenchymal stem cells.

10 **[0072]** For mesenchymal cell separation/cultivation, the explants were submerged in 5 ml DMEM (Invitrogen) supplemented with 10% fetal bovine serum (Hyclone) (DMEM/10% FBS) and maintained in a CO₂ cell culture incubator at 37°C. The medium was changed every 2 or 3 days. Cell outgrowth was monitored under light microscopy. Outgrowing cells were harvested by trypsinization (0.125% trypsin/0.05% EDTA) for further expansion and cryo-preservation using DMEM/10% FBS.

15 **[0073]** For epithelial cell separation/cultivation, cell culture plastic surfaces were coated with collagen 1/ collagen 4 mixtures (1:2) before placing the tissue samples on the surface. The tissue samples were submerged in 5 ml EpiLife medium or Medium 171 (both from Cascade Biologics). The medium was changed every 2 or 3 days. Cell outgrowth from tissue culture explants was monitored under light microscopy. Outgrowing cells were harvested by trypsinization (0.125% trypsin/0.05% EDTA) using EpiLife medium or Medium 20 171.

25 **[0074]** For the enzymatic extraction method of cells, umbilical cord amniotic membrane was divided into small pieces of 0.5 cm x 0.5 cm and digested in 0.1% (w/v) collagenase type1 solution (Roche Diagnostics) at 37°C for 6 hours. The samples were vortexed every 15 min for 2 min. Cells were harvested by centrifugation at 4000 rpm for 30 min. Two different approaches were employed to isolate either epithelial or mesenchymal stem cells.

30 **[0075]** For isolation of epithelial stem cells, cell pellets were resuspended in EpiLife medium or Medium 171 (both from Cascade Biologics) supplemented with 50 µg/ml insulin-like growth factor-1 (IGF-1), 50 µg/ml

platelet-derived growth factor-BB (PDGF-BB), 5 µg/ml transforming growth factor-β1 (TGF-β1), and 5 µg/ml insulin (all obtained from R&D Systems), counted and seeded on 10 cm tissue culture dishes pre-coated with collagen 1/collagen 4 mixtures (1:2; Becton Dickinson) at density of 1×10^6 cells/dish.

5 After 24 hours, attached cells were washed with warm PBS and medium was replaced with supplement-added EpiLife medium or Medium 171. The medium was changed every 2 or 3 days. Cell growth and expanding clonal formation was monitored under light microscopy. At a confluence of about 70%, cells were sub-cultured by trypsinization (0.125% trypsin/0.05% EDTA) for further

10 expansion and cryo-preservation.

[0076] For isolation of mesenchymal stem cells, cell pellets were resuspended in DMEM/10% FBS, counted and seeded on 10 cm tissue culture dishes at density of 1×10^6 cells/dish. The culture medium was changed every 2 or 3 days. Cell growth and expansion was monitored under light microscopy. At

15 a confluence of about 90%, cells were sub-cultured as outlined above.

[0077] For cultivation of epithelial and mesenchymal stem cells on feeder layer, umbilical cord lining membrane was digested by collagenase treatment, counted and seeded on 10cm tissue culture dishes coated with lethally irradiated or Mitomycin C treated 3T3 fibroblasts (feeder layer) in

20 Green's medium. The culture medium was changed every 2 or 3 days. Colony formation was monitored under light microscopy and photographed.

Example 3: Identification of stem/progenitor cells

[0078] *Epithelial cells:* Fig. 1 shows pictures of outgrowing epithelial cells

25 from umbilical cord amniotic membrane prepared by the method using tissue explant (40 x magnification). Pictures were taken at day 2 (Fig. 1A) and day 5 (Fig. 1B, C) of tissue culture. Cell morphology analysis demonstrated polyhedral shaped epithelial-like cells. Enzymatic digestion of the umbilical cord segments yielded similar (Fig. 2), epithelial cells at day 2 (Fig. A, C) and day 5

30 (Fig. B, D) (40 x magnification). Fig.7 shows pictures of colony formation of epithelial stem cells from umbilical cord amniotic membrane cultured on feeder layer using Green's method (40 x magnification). A colony of polyhedral shaped epithelial-like cells expanded rapidly from day 3 to day 7.

[0079] *Mesenchymal cells:* Outgrowth of mesenchymal cells explanted

from umbilical cord amniotic membrane was observed as early as 48 hours after placement in tissue culture dishes using DMEM supplemented with 10% fetal calf serum (FCS) as culture medium (**Fig. 3A, C**) (40 x magnification). The cells were characterized by their spindle shaped morphology, and migrated and expanded both easily and quickly *in vitro*, closely resembling fibroblasts (**Fig. 3B, D**) (40 x magnification). Similar observations were noted in the cell group isolated by collagenase enzymatic digestion (**Fig. 4**). **Fig. 4A** shows mesenchymal cells isolated from umbilical cord amniotic membrane at day 2. Cell proliferation was observed at day 5 (**Fig. 4B**) (40 x magnification). Figures 6 and 8 show pictures of colony formation of mesenchymal stem cells from umbilical cord amniotic membrane cultured on non-feeder layer and feeder layer condition in DMEM/10%FCS (40 x magnification). The colonies of elongated shaped fibroblastic-like cells expanded rapidly from day 3 to day 7.

[0080] Western blot analysis (**Fig.9**) shows that mesenchymal stem cells from umbilical cord amniotic membrane (UCMC) and umbilical cord epithelial cells (UCEC) isolated in accordance with the invention expressed the POU5f1 gene which encodes the transcription factor Octamer-4 (Oct-4) a specific marker of embryonic stem cells (cf. Niwa, H., Miyazaki, J., and Smith, A.G. (2000). Nat. Genet. 24, 372–376). Thus, this analysis indicates the embryonic-like properties of these stem cells. These cells also highly expressed the other growth factors such as connective tissue growth factor (CTGF), vascular endothelial growth factor (VEGF), placenta-like growth factor PLGF, STAT3, stem cell factor (SCF), Hepatoma-derived Growth Factor (HDGF), Fibroblast Growth Factor-2 (FGF-2), Platelet-derived Growth Factor (PDGF), alpha-Smooth Muscle Actin (α -SMA), Fibronectin, Decorin, Syndecan-1,2,3,4. In **Fig. 9**, the expression of these genes is compared to human dermal fibroblasts, bone marrow mesenchymal cells (BMSC) and adipose-derived mesenchymal cells (ADMC). Highly secreted Activin A and Follistatin (both of which proteins are well known to promote tissue repair and regeneration, enhanced angiogenesis, and maintain embryonic stem cell culture, so that expression of the respective genes is a sign for the embryonic properties and ability of the cells to differentiate) were detected by ELISA assay (data not shown) in supernatants of umbilical cord mesenchymal and epithelial stem cell culture in comparison with bone marrow, adipose derived stem cells, human dermal fibroblasts and

epidermal keratinocytes. Also these results indicate that the cells of the invention are promising candidates in therapeutic application of these cells areas such as regenerative medicine, aging medicine, tissue repair and tissue engineering.

5 **[0081]** Mesenchymal cells were further characterized by analysis of secreted cytokines and growth factors in comparison with human bone-marrow mesenchymal stem cells. The umbilical cord epithelial stem cells (UCEC) were analysed in comparison with human epidermal keratinocytes. This analysis was carried out as follows: Briefly, UMMC, UCEC, dermal fibroblasts, bone-marrow
10 mesenchymal cells, epidermal keratinocytes were cultured in growth media until 100% confluence (37°C, 5% CO₂) and then synchronized in starvation medium (serum-free DMEM) for 48 hours. The next day, the medium was replaced the next against fresh serum-free DMEM and the cells then were cultivated for another 48 hours. Conditioned media were collected, concentrated and
15 analyzed using a Cytokine Array (RayBiotech, Inc, GA, USA).

[0082] The results of this analysis show that UCMC secrete Interleukin-6 (IL-6); (MCP1); hepatocyte growth factor (HGF); Interleukin-8 (IL8); sTNFR1; GRO; TIMP1; TIMP2; TRAILR3; uPAR; ICAM1; IGFBP3; IGFBP6 (**Fig.11**), whereas UCEC secrete IGFBP-4; PARC; EGF; IGFBP-2; IL-6; Angiogenin;
20 GCP-2; IL1R α ; MCP-1; RANTES; SCF; TNF β ; HGF; IL8; sTNFR; GRO; GRO- α ; Amphiregulin; IL-1R4/ST2; TIMP1; TIMP2; uPAR; VEGF (**Fig.12**).

[0083] Accordingly, this shows that both cells types secrete large amounts of cytokines and growth factors that play important roles in developmental biology, tissue homeostasis, tissue repair and regeneration and
25 angiogenesis. This further demonstrates the versatility of the cells of the invention for use in the respective therapeutic applications.

[0084] In addition, the cells of the invention were further examined with respect to their safety profile using mouse teratoma formation assay as an indicator. Six SCID mice were used in these experiments. A suspension of more
30 than 2 million UCMC was injected with a sterile 25G needle into the thigh muscle of each SCID mouse. Animals were kept up to 6 months and tumor formation was assessed. No tumor formation was observed in these mice (data not shown). This indicates that the cells of the invention are safe and do not have any capability to form tumors, benign or otherwise.

Example 4: Cultivation of stem/progenitor cells in serum free media

[0085] UCMC cells were cultured in DMEM containing 10 FCS and in serum-free media, PTT-1, PTT-2 and PTT-3. The three media PTT-1, PTT-2
 5 and PTT-3 were prepared by one of the present inventors, Dr Phan. In brief, these 3 media do not contain fetal bovine or human serum, but contain different cytokines and growth factors such as IGF, EGF, TGF-beta, Activin A, BMPs, PDGF, transferrin, and insulin. The growth factor components vary between media to assess differential growth characteristics. The cultivation was carried
 10 out as follows: Different proportions of growth factors and cytokines were added in basal media. UCMC were thawed and maintained in these media for 10 days. Cell proliferation was monitored under light microscopy.

[0086] Fig. 13 shows good UCMC growth in the 4 different media groups (Fig. 13-1 to Fig.13-5), wherein the morphology of UCMC cells is different
 15 depending on the ratio or proportion of cytokines or growth factors present in the respective media. In contrast, bone marrow and adipose-derived mesenchymal cells did not grow well in these serum-free media (Fig.13-6 and Fig.13-7). Accordingly, the good growth of the UCMC demonstrates the robustness of the cells of the invention and their high viability, indicating that
 20 their growth characteristics are superior to conventional sources of mesenchymal stem cells as bone marrow derived and adipose-derived mesenchymal cells. In this respect, it is worth to note that (bovine) serum free medium was used in these experiments and that the majority of human mesenchymal cells do not grow well in serum-free medium systems. Thus,
 25 using the cells of the invention in connection with defined serum-free media technologies is a big advantage in cell therapy as the risks of using fetal bovine serum for cell culture and expansion are removed. (Although use of bovine serum has been practiced for a long time and typically optimizes cell growth, concerns of its used have been raised as to the transmission of zoonoses as
 30 Bovine Spongiform Encephalopathy (Mad Cow Disease)).

Example 5: Characterization of the gene expression profile of umbilical cord epithelial and mesenchymal stem cells

[0087] The gene expression profile of umbilical cord epithelial and

mesenchymal stem cells was analyzed using a DNA microarray. For this purpose, UCMC and UCEC were cultured in growth media at 37°C, 5% CO₂ until 100% confluence. Cells were synchronized in basal media for 48 hours then replaced with fresh basal media for another 48 hours. Total RNA was harvested and sent to Silicon Genetics Microarray Service. Data analysis was performed using GeneSpring 7.2). **Fig. 14** summarizes the global gene expression. UCEC expressed a total of 28055 genes and UCMC expressed a total of 34407 genes. There are 27308 overlapping genes expressing in both cell types. 747 genes expressed were unique to UCEC and 7099 genes expressed were unique to UCMC. The selected genes of interest are presented in **Fig. 14**.

[0088] Both stem cell types expressed 140 genes related to embryonic stem cells and embryonic development, further supporting that the cells of the invention have embryonic stem cell-like properties: Nanog; Alpha-fetal protein; Pre-B-cell leukemia transcription factor 3; Laminin alpha 5; Carcinoembryonic antigen-like 1; abhydrolase domain containing 2; Delta-like 3 (Drosophila); Muscleblind-like (Drosophila); GNAS complex locus; Carcinoembryonic antigen-related cell adhesion molecule 3; Palmitoyl-protein thioesterase 2; Pregnancy specific beta-1-glycoprotein 2; Carcinoembryonic antigen-like 1; Embryonic ectoderm development; Maternal embryonic leucine zipper kinase; Chorionic somatomammotropin hormone 2; Forkhead box D3; radical fringe homolog (Drosophila); Kinesin family member 1B; Myosin, heavy polypeptide 3, skeletal muscle, embryonic; Split hand/foot malformation (ectrodactyly) type 3; TEA domain family member 3; Laminin, alpha 1; Chorionic somatomammotropin hormone 1; placental lactogen; Corticotropin releasing hormone receptor 1; thyrotrophic embryonic factor; Aryl-hydrocarbon receptor nuclear translocator 2; Membrane frizzled-related protein; Neuregulin 1; Collagen, type XVI, alpha 1; Neuregulin 1; Chorionic somatomammotropin hormone 1 (placental lactogen); CUG triplet repeat, RNA binding protein 1; Chorionic somatomammotropin hormone 1 (placental lactogen) Bystin-like; MyoD family inhibitor; Retinoic acid induced 2; GNAS complex locus; Pre-B-cell leukemia transcription factor 4; Laminin, alpha 2 (merosin, congenital muscular dystrophy); SMAD, mothers against DPP homolog 1 (Drosophila); Homo sapiens transcribed sequence with moderate similarity to protein pir:D28928 (H.sapiens) D28928 pregnancy-

specific beta-1 glycoprotein IB, abortive - human (fragment); Kinesin family member 1B; Bruno-like 4, RNA binding protein (Drosophila); Embryo brain specific protein; Pregnancy-induced growth inhibitor; SMAD, mothers against DPP homolog 5 (Drosophila); Chorionic somatomammotropin hormone

5 2; Adenylate cyclase activating polypeptide 1 (pituitary); Carcinoembryonic antigen-related cell adhesion molecule; Laminin, alpha 3; Protein O-fucosyltransferase 1; Jagged 1 (Alagille syndrome); Twisted gastrulation homolog 1 (Drosophila); ELAV (embryonic lethal, abnormal vision, Drosophila)-like 3 (Hu antigen C); Thyrotrophic embryonic factor; Solute carrier family 43,

10 member 3; Inversin; nephronophthisis 2 (infantile); inversion of embryonic turning; Homo sapiens inversin (INVS), transcript variant 2, mRNA; Homo sapiens transcribed sequences; Homeo box D8; Embryonal Fyn-associated substrate; ELAV (embryonic lethal, abnormal vision, Drosophila)-like 1 (Hu antigen R); Basic helix-loop-helix domain containing, class B, 2; Oxytocin

15 receptor; Teratocarcinoma-derived growth factor 1; Fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor); Adrenomedullin; Nuclear receptor coactivator 6-CUG triplet repeat, RNA binding protein 1; Twisted gastrulation homolog 1 (Drosophila); Carcinoembryonic antigen-related cell adhesion molecule 4; Protein tyrosine phosphatase, receptor

20 type, R; Acrg embryonic lethality (mouse) minimal region ortholog; EPH receptor A3; Delta-like 1 (Drosophila); Nasal embryonic LHRH factor; Transcription factor CP2-like 1; Split hand/foot malformation (ectrodactyly) type 3; Jagged 2; Homo sapiens transcribed sequence; Neuregulin 1; Split hand/foot malformation (ectrodactyly) type 1; Solute carrier family 43, member 3;

25 Hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl-Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), alpha subunit; Fucosyltransferase 10 (alpha (1,3) fucosyltransferase); Acrg embryonic lethality (mouse) minimal region ortholog; Carcinoembryonic antigen-related cell adhesion molecule 7; Nucleophosmin/nucleoplasmin, 2; Fc fragment of IgG,

30 receptor, transporter, alpha; Twisted gastrulation homolog 1 (Drosophila); Homo sapiens similar to vacuolar protein sorting 35; maternal-embryonic 3 (LOC146485), mRNA; abhydrolase domain containing 2; T, brachyury homolog (mouse); A disintegrin and metalloproteinase domain 10; Ribosomal protein L29; Endothelin converting enzyme 2; ELAV (embryonic lethal, abnormal vision,

Drosophila)-like 1 (Hu antigen R); Trophinin; Homeo box B6; Laminin, alpha 4; Homeo box B6; hypothetical protein FLJ13456; NACHT, leucine rich repeat and PYD containing 5; ELAV (embryonic lethal, abnormal vision, Drosophila)-like 1 (Hu antigen R); Undifferentiated embryonic cell transcription factor 1;
 5 Pregnancy-associated plasma protein A, pappalysin 1; Secretoglobin, family 1A, member 1 (uteroglobin); Parathyroid hormone-like hormone; Carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein); Laminin, alpha 1.

[0089] Both stem cell types also expressed thousands of genes related to
 10 developmental biology, cell growth and differentiation, cell homeostasis, cell and tissue repair and regeneration. Examples of such growth factors and their receptors is as follows: (G-CSF, FGFs, IGFs, KGF, NGF, VEGFs, PIGF, Angiopoietin, CTGF, PDGFs, HGF, EGF, HDGF, TGF-beta, Activins and Inhibins, Follistatin, BMPs, SCF/c-Kit, LIF, WNTs, SDFs, OncostatinM,
 15 Interleukins, Chemokines and many others); MMPs, TIMPs extracellular matrices (collagens, laminins, fibronectins, vitronectins, tenascins, integrins, syndecans, decorin, fibromodulin, proteoglycans, sparc/osteonectin, mucin, netrin, glypican, cartilage associated protein, matrilin, hyaluronan, fibulin, ADAMTS, biglycan, discoidin, desmosome components, ICAMs, cadherins,
 20 catenins and many others); cytokeratin.

[0090] There are groups of genes present only in UCMC. These genes are related to the following: Normal Physiological Processes (Insulin-like growth factor 1 (somatomedin C); Insulin-like 4 (placenta); Relaxin 1; Plasminogen; Insulin-like growth factor 1 (somatomedin C); Insulin-like 5; Insulin-like growth
 25 factor 1 (somatomedin C); Insulin-like growth factor 2 (somatomedin A), Homeostasis (Radial spokehead-like 1; Hemochromatosis; Chemokine (C-C motif) ligand 5; Interleukin 31 receptor A; Chemokine (C-X-C motif) ligand 12 (stromal cell-derived factor 1); Nuclear receptor subfamily 3, group C, member 2; Hemochromatosis; Chemokine (C-C motif) ligand 23; Chemokine (C-C motif)
 30 ligand 23; Ferritin mitochondrial; Peroxisome proliferative activated receptor, gamma, coactivator 1, alpha; Surfactant, pulmonary-associated protein D; Chemokine (C-C motif) ligand 11; Chemokine (C-C motif) ligand 3; Egl nine homolog 2 (C. elegans); Peroxisome proliferative activated receptor, gamma, coactivator 1, beta; Chemokine (C-C motif) ligand 1; Chemokine (C-X-C motif)

ligand 12 (stromal cell-derived factor 1); ATPase, Na⁺/K⁺ transporting, alpha 2
 (+) polypeptide; Chemokine (C motif) ligand 2; Hemopexin; Ryanodine receptor
 3), Morphogenesis (Spectrin, alpha, erythrocytic 1 (elliptocytosis 2); Homeo box
 D3; Eyes absent homolog 1 (Drosophila); Ras homolog gene family, member J;
 5 Leukocyte specific transcript 1; Ectodysplasin A2 receptor; Glypican 3; Paired
 box gene 7; Corin, serine protease; Dishevelled, dsh homolog 1 (Drosophila);
 Ras homolog gene family, member J; T-box 3 (ulnar mammary syndrome);
 Chondroitin beta1,4 N-acetylgalactosaminyltransferase; Chondroitin beta1,4 N-
 acetylgalactosaminyltransferase; SRY (sex determining region Y)-box 10;
 10 Myosin, heavy polypeptide 9, non-muscle; Luteinizing
 hormone/choriogonadotropin receptor; radical fringe homolog (Drosophila);
 Secreted frizzled-related protein 5; Wingless-type MMTV integration site family,
 member 11; Eyes absent homolog 2 (Drosophila); Muscleblind-like
 (Drosophila); T-box 5; Mab-21-like 1 (C. elegans); Growth arrest-specific 2; Sex
 15 comb on midleg homolog 1 (Drosophila); T-box 6; Filamin-binding LIM protein-
 1; Melanoma cell adhesion molecule; Twist homolog 1 (acrocephalosyndactyly
 3; Saethre-Chotzen syndrome) (Drosophila); Homeo box A11; Keratocan;
 Fibroblast growth factor 1 (acidic); Carboxypeptidase M; CDC42 effector protein
 (Rho GTPase binding) 4; LIM homeobox transcription factor 1, beta; Engrailed
 20 homolog 1; Carboxypeptidase M; Fibroblast growth factor 8 (androgen-
 induced); Fibroblast growth factor 18; Leukocyte specific transcript 1;
 Endothelin 3; Paired-like homeodomain transcription factor 1), Embryonic
 Development (Pregnancy specific beta-1-glycoprotein 3; ELAV (embryonic
 lethal, abnormal vision, Drosophila)-like 4 (Hu antigen D); G protein-coupled
 25 receptor 10; Ectodysplasin A2 receptor; ATP-binding cassette, sub-family B
 (MDR/TAP), member 4; Pregnancy specific beta-1-glycoprotein 11; Nasal
 embryonic LHRH factor; Relaxin 1; Notch homolog 4 (Drosophila); Pregnancy
 specific beta-1-glycoprotein 6; pih-2P; Homo sapiens pregnancy-induced
 hypertension syndrome-related protein (PIH2); Oviductal glycoprotein 1,
 30 120kDa (mucin 9, oviductin); Progesterone-associated endometrial protein;
 Myosin, light polypeptide 4, alkali; atrial, embryonic; Prolactin; Notch homolog 4
 (Drosophila); Pre-B-cell leukemia transcription factor 1; radical fringe homolog
 (Drosophila); Corticotropin releasing hormone; Nuclear receptor subfamily 3,
 group C, member 2; Neuregulin 2; Muscleblind-like (Drosophila); Myosin, light

polypeptide 4, alkali; atrial, embryonic; Homo sapiens cDNA FLJ27401 fis,
 clone WMC03071; Extraembryonic, spermatogenesis, homeobox 1-like; Insulin-
 like 4 (placenta); Human processed pseudo-pregnancy-specific glycoprotein
 (PSG12) gene, exon B2C containing 3' untranslated regions of 2 alternative
 5 splice sites C1 and C2; Fms-related tyrosine kinase 1 (vascular endothelial
 growth factor/vascular permeability factor receptor); Pre-B-cell leukemia
 transcription factor 1; Pregnancy specific beta-1-glycoprotein 3;
 carcinoembryonic antigen-related cell adhesion molecule 1 (biliary
 glycoprotein); Steroid sulfatase (microsomal), arylsulfatase C, isozyme S;
 10 Homeo box B6; Protein O-fucosyltransferase 1; LIM homeobox transcription
 factor 1, beta; Carcinoembryonic antigen-related cell adhesion molecule 1
 (biliary glycoprotein); Follicle stimulating hormone, beta polypeptide;
 Angiotensinogen (serine (or cysteine) proteinase inhibitor, clade A (alpha-1
 antiproteinase, antitrypsin), member 8); Carcinoembryonic antigen-related cell
 15 adhesion molecule 6 (non-specific cross reacting antigen); Protein kinase C,
 alpha binding protein; Collectin sub-family member 10 (C-type lectin); Laminin,
 alpha 1), the Extracellular Space (Carboxylesterase 1 (monocyte/macrophage
 serine esterase 1); Fibroblast growth factor 5; Progastricsin (pepsinogen C);
 Sperm associated antigen 11; Proprotein convertase subtilisin/kexin type 2 ;
 20 Hyaluronan binding protein 2; Sema domain, immunoglobulin domain (Ig), short
 basic domain, secreted, (semaphorin) 3F; Interleukin 2; Chymotrypsin-like;
 Norrie disease (pseudoglioma); mucin 5, subtypes A and C,
 tracheobronchial/gastric; Carboxypeptidase B2 (plasma, carboxypeptidase U);
 radical fringe homolog (Drosophila); Pregnancy specific beta-1-glycoprotein 11;
 25 Meprin A, alpha (PABA peptide hydrolase) ; Tachykinin, precursor 1 (substance
 K, substance P, neurokinin 1, neurokinin 2, neuromedin L, neurokinin alpha,
 neuropeptide K, neuropeptide gamma) ; Fibroblast growth factor 8 (androgen-
 induced); Fibroblast growth factor 13; Hemopexin; Breast cancer 2, early onset;
 Fibroblast growth factor 14; Retinoschisis (X-linked, juvenile) 1; Chitinase 3-like
 30 1 (cartilage glycoprotein-39); Dystonin; Secretoglobin, family 1D, member 2;
 Noggin; WAP four-disulfide core domain 2; CD5 antigen-like (scavenger
 receptor cysteine rich family); Scrapie responsive protein 1; Gremlin 1 homolog,
 cysteine knot superfamily (Xenopus laevis); Interleukin 16 (lymphocyte
 chemoattractant factor); Chemokine (C-C motif) ligand 26; Nucleobindin 1;

Fibroblast growth factor 18; Insulin-like growth factor binding protein 1;
 Surfactant, pulmonary-associated protein A1; Delta-like 1 homolog (*Drosophila*);
 Cocaine- and amphetamine-regulated transcript; Meprin A, beta; Interleukin
 17F; Complement factor H; Cysteine-rich secretory protein 2; Dystonin; WAP
 5 four-disulfide core domain 1; Prolactin; Surfactant, pulmonary-associated
 protein B; Fibroblast growth factor 5; Dickkopf homolog 2 (*Xenopus laevis*);
 Sperm associated antigen 11; Chemokine (C-C motif) ligand 11; Meprin A,
 alpha (PABA peptide hydrolase); Chitinase 3-like 2; C-fos induced growth factor
 (vascular endothelial growth factor D); Chemokine (C-C motif) ligand 4;
 10 Poliovirus receptor; Hyaluronoglucosaminidase 1; Oviductal glycoprotein 1,
 120kDa (mucin 9, oviductin); Chemokine (C-X-C motif) ligand 9; Secreted
 frizzled-related protein 5; Amelogenin (amelogenesis imperfecta 1, X-linked);
 Relaxin 1; Sparc/osteonectin, cwcw and kazal-like domains proteoglycan
 (testican); Chemokine (C-C motif) ligand 26; Fibroblast growth factor 1 (acidic);
 15 Angiopoietin-like 2; Fms-related tyrosine kinase 1 (vascular endothelial growth
 factor/vascular permeability factor receptor); Dystonin; Insulin-like 4 (placenta);
 Transcobalamin II; macrocytic anemia; Chemokine (C-C motif) ligand 1; Insulin-
 like growth factor binding protein, acid labile subunit; Complement factor H;
 Pregnancy specific beta-1-glycoprotein 6; Silver homolog (mouse);
 20 Proteoglycan 4; Fibroblast growth factor 16; Cytokine-like protein C17;
 Granulysin; Angiopoietin 2; Chromogranin B (secretogranin 1); Sema domain,
 immunoglobulin domain (Ig), and GPI membrane anchor, (semaphorin) 7A;
 Pleiotrophin (heparin binding growth factor 8, neurite growth-promoting factor
 1); Chloride channel, calcium activated, family member 3; Secretoglobin, family
 25 1D, member 1; Fibulin 1; Phospholipase A2 receptor 1, 180kDa), and the
 Extracellular Matrix (ADAMTS-like 1; Periostin, osteoblast specific factor;
 Glypican 5; Leucine rich repeat neuronal 3; Transglutaminase 2 (C polypeptide,
 protein-glutamine-gamma-glutamyltransferase); A disintegrin-like and
 metalloprotease (repolysin type) with thrombospondin type 1 motif, 2;
 30 Microfibrillar-associated protein 4; Glypican 3; Collagen, type V, alpha 3; Tissue
 inhibitor of metalloproteinase 2; Keratocan; Cartilage oligomeric matrix protein;
 Lumican; Hyaluronan and proteoglycan link protein 3; Statherin; A disintegrin-
 like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 3;
 Spondin 1, extracellular matrix protein; Chitinase 3-like 1 (cartilage glycoprotein-

39); Collagen, type IV, alpha 3 (Goodpasture antigen); Wingless-type MMTV integration site family, member 7B; Collagen, type VI, alpha 2; Lipocalin 7; Hyaluronan and proteoglycan link protein 4; A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2); Fibronectin 1; Matrilin 1, cartilage matrix protein; Hypothetical protein FLJ13710; Chondroitin beta1,4 N-acetylgalactosaminyltransferase; Matrix metalloproteinase 16 (membrane-inserted); Von Willebrand factor; Collagen, type VI, alpha 2; Transmembrane protease, serine 6 ; Matrix metalloproteinase 23B ; Matrix metalloproteinase 14 (membrane-inserted);

10 Leucine rich repeat neuronal 3; SPARC-like 1 (mast9, hevin); Sparc/osteonectin, cwcvc and kazal-like domains proteoglycan (testican) 3; Dermatotontin; collagen, type XIV, alpha 1 (undulin); Amelogenin, Y-linked; Nidogen (enactin); ADAMTS-like 2; Hyaluronan and proteoglycan link protein 2; Collagen, type XV, alpha 1; Glypican 6; Matrix metalloproteinase 12

15 (macrophage elastase); Amelogenin (amelogenesis imperfecta 1, X-linked); A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 15; Transmembrane protease, serine 6; A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 16; Sparc/osteonectin, cwcvc and kazal-like domains proteoglycan (testican); A

20 disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 20; Collagen, type XI, alpha 1; Hyaluronan and proteoglycan link protein 1; Chondroitin beta1,4 N-acetylgalactosaminyltransferase ; Asporin (LRR class 1) ; Collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant) ; Secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early

25 T-lymphocyte activation 1); Matrix Gla protein; Fibulin 5; collagen, type XIV, alpha 1 (undulin); Tissue inhibitor of metalloproteinase 3 (Sorsby fundus dystrophy, pseudoinflammatory); Collagen, type XXV, alpha 1; Cartilage oligomeric matrix protein ; Collagen, type VI, alpha 1 ; Chondroadherin ; Collagen, type XV, alpha 1 ; A disintegrin-like and metalloprotease (reprolysin

30 type) with thrombospondin type 1 motif, 16; Collagen, type IV, alpha 4 ; Dentin matrix acidic phosphoprotein ; Collagen, type IV, alpha 1; Thrombospondin repeat containing 1; Matrix metalloproteinase 16 (membrane-inserted); Collagen, type I, alpha 2; Fibulin 1; Tectorin beta; Glycosylphosphatidylinositol specific phospholipase D1; Upregulated in colorectal cancer gene 1).

Cytoskeleton: (Filamin B, beta (actin binding protein 278); Centrin, EF-hand protein, 1; FERM domain containing 3; Bridging integrator 3; Parvin, gamma; Rho guanine nucleotide exchange factor (GEF) 11; Tyrosine kinase 2; Kelch-like 4 (Drosophila); Spectrin, beta, erythrocytic (includes spherocytosis, clinical
 5 type I); Arg/Abl-interacting protein ArgBP2; Advillin; Spectrin repeat containing, nuclear envelope 1; Catenin (cadherin-associated protein), delta 1; Erythrocyte membrane protein band 4.1 like 5; Catenin (cadherin-associated protein), alpha 2; Chemokine (C-C motif) ligand 3; Sarcoglycan, gamma (35kDa dystrophin-associated glycoprotein); Nebulin; Thymosin, beta, identified in neuroblastoma
 10 cells; 3-phosphoinositide dependent protein kinase-1; Wiskott-Aldrich syndrome protein interacting protein; Dystonin; Huntingtin interacting protein 1; KIAA0316 gene product; Tropomodulin 4 (muscle); Deleted in liver cancer 1; Villin-like; Syntrophin, beta 1 (dystrophin-associated protein A1, 59kDa, basic component 1); Protein kinase, cGMP-dependent, type I; Homo sapiens similar to keratin 8;
 15 cytokeratin 8; keratin, type II cytoskeletal 8 (LOC345751), mRNA; Adducin 1 (alpha); Protein kinase C and casein kinase substrate in neurons 3; Dystonin; Kell blood group; Filamin A interacting protein 1; Growth arrest-specific 2; ; Chromosome 1 open reading frame 1; Stathmin-like 2; Spectrin, alpha, erythrocytic 1 (elliptocytosis 2); FKSG44 gene; Kinesin family member 1C;
 20 Tensin; Kaptin (actin binding protein); Neurofibromin 2 (bilateral acoustic neuroma); Pleckstrin homology, Sec7 and coiled-coil domains 2 (cytohesin-2); Actin-related protein T1; Wiskott-Aldrich syndrome-like; Kelch-like 4 (Drosophila); Fascin homolog 1, actin-bundling protein (Strongylocentrotus purpuratus); Amphiphysin (Stiff-Man syndrome with breast cancer 128kDa
 25 autoantigen); Polycystic kidney disease 2-like 1; Ankyrin 2, neuronal; CDC42 binding protein kinase alpha (DMPK-like); Hypothetical protein FLJ36144; Arg/Abl-interacting protein ArgBP2; Formin-like 3; Catenin (cadherin-associated protein), beta 1, 88kDa; Profilin 2; Synaptopodin 2-like; Syntrophin, gamma 2; Phospholipase D2; Engulfment and cell motility 2 (ced-12 homolog, C. elegans);
 30 Neurofilament, light polypeptide 68kDa; Dystonin; Actin-like 7B; Kinesin family member 1C; PDZ and LIM domain 3; Adducin 2 (beta); obscurin, cytoskeletal calmodulin and titin-interacting RhoGEF; Tubulin, beta polypeptide paralog; Filamin A interacting protein 1; Talin 1; Homo sapiens similar to [Segment 1 of 2] Piccolo protein (Aczonin) (LOC375597); CDC42 effector protein (Rho

GTPase binding) 4; Syndecan 1; Filamin A, alpha (actin binding protein 280); Profilin 2; Tensin like C1 domain containing phosphatase; Hypothetical protein MGC33407; Rho family GTPase 1; Flavoprotein oxidoreductase MICAL2; Ca²⁺-dependent secretion activator; Rabphilin 3A-like (without C2 domains);

5 Myosin XVA; Protein kinase, cGMP-dependent, type I; Myosin regulatory light chain interacting protein; Kinesin family member 13B; Muscle RAS oncogene homolog; Spectrin, beta, non-erythrocytic 1; TAO kinase 2; Filamin B, beta (actin binding protein 278); Neurofibromin 2 (bilateral acoustic neuroma); Catenin (cadherin-associated protein), alpha 3; obscurin, cytoskeletal

10 calmodulin and titin-interacting RhoGEF; Coronin, actin binding protein, 1A; Erythrocyte membrane protein band 4.1-like 1; Spectrin, beta, non-erythrocytic 4; Thymosin, beta 4, Y-linked; Tektin 2 (testicular); Ras homolog gene family, member J; Serine/threonine kinase with Dbp- and pleckstrin homology domains; Dystrobrevin, beta; Actin, gamma 2, smooth muscle, enteric; Tara-like protein;

15 Caspase 8, apoptosis-related cysteine protease; Kelch repeat and BTB (POZ) domain containing 10; Mucin 1, transmembrane; Microtubule-associated protein tau; Tensin; Ras homolog gene family, member F (in filopodia); Adducin 1 (alpha); Actinin, alpha 4; Erythrocyte membrane protein band 4.1 (elliptocytosis 1, RH-linked); Bicaudal D homolog 2 (Drosophila); Ankyrin 3, node of Ranvier

20 (ankyrin G); Myosin VIIA (Usher syndrome 1B (autosomal recessive, severe)); Catenin (cadherin-associated protein), alpha 2; Homo sapiens similar to keratin 8, type II cytoskeletal - human (LOC285233); Fascin homolog 3, actin-bundling protein, testicular; Ras homolog gene family, member J; Beaded filament structural protein 2, phakinin; Desmin; Myosin X; Signal-induced proliferation-associated gene 1; Scinderin; Coactosin-like 1 (Dictyostelium); Engulfment and cell motility 2 (ced-12 homolog, *C. elegans*); Tubulin, beta 4; Ca²⁺-dependent secretion activator; FERM domain containing 4A; Actin, alpha 1, skeletal muscle; Talin 1; Caldesmon 1; Filamin-binding LIM protein-1; Microtubule-associated protein tau; Syntrophin, alpha 1 (dystrophin-associated protein A1,

25 59kDa, acidic component); Adducin 2 (beta); Filamin A interacting protein 1; PDZ and LIM domain 3; Erythrocyte membrane protein band 4.1 like 4B; FYN binding protein (FYB-120/130); Bridging integrator 3). Extracellular: (A disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 20; SPARC-like 1 (mast9, hevin); Serine (or cysteine) proteinase

30

inhibitor, clade G (C1 inhibitor), member 1, (angioedema, hereditary); Urocortin; Chymotrypsin-like; Platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog); BMP-binding endothelial regulator precursor protein; Complement factor H; Chorionic somatomammotropin

5 hormone-like 1; Chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated); Fibronectin 1; Pregnancy specific beta-1-glycoprotein 3; A disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 3; CocoaCrisp; Insulin-like 4 (placenta); Wingless-type MMTV integration site family, member 11; Cartilage oligomeric matrix protein;

10 Transmembrane protease, serine 6; C-fos induced growth factor (vascular endothelial growth factor D); Family with sequence similarity 12, member B (epididymal); Protein phosphatase 1, regulatory subunit 9B, spinophilin; Transcobalamin II; macrocytic anemia; Coagulation factor V (proaccelerin, labile factor); Phospholipase A2, group IID; Tumor necrosis factor, alpha-induced

15 protein 6; Collagen, type XV, alpha 1; Hyaluronan and proteoglycan link protein 3; collagen, type XIV, alpha 1 (undulin) ; Interleukin 19; Protease inhibitor 15; Cholinergic receptor, nicotinic, beta polypeptide 1 (muscle); Lysyl oxidase-like 3; Insulin-like growth factor binding protein 5; Growth hormone 1; Casein beta; NEL-like 2 (chicken); I factor (complement); Chemokine (C-C motif) ligand 23 ;

20 Interferon, alpha 2; Matrix metalloproteinase 16 (membrane-inserted); Matrix metalloproteinase 12 (macrophage elastase); Glypican 5; Pregnancy specific beta-1-glycoprotein 3; Fibroblast growth factor 6; Gremlin 1 homolog, cysteine knot superfamily (*Xenopus laevis*); Protein S (alpha); Chondroitin beta1,4 N-acetylgalactosaminyltransferase; Glycosylphosphatidylinositol specific

25 phospholipase D1; Fibroblast growth factor 1 (acidic); Spondin 1, extracellular matrix protein; Bone morphogenetic protein 1; Surfactant, pulmonary-associated protein B; Dentin matrix acidic phosphoprotein ; Lipoprotein, Lp(a) ; Mucin 1, transmembrane; Mannan-binding lectin serine protease 1 (C4/C2 activating component of Ra-reactive factor); Meprin A, beta; Secretoglobin,

30 family 1D, member 1; Asporin (LRR class 1); Chemokine (C-C motif) ligand 25; Cytokine-like protein C17; Insulin-like 5; Meprin A, alpha (PABA peptide hydrolase) ; Scrapie responsive protein 1; Fibroblast growth factor 18; Chemokine (C-X-C motif) ligand 9; Inhibin, beta B (activin AB beta polypeptide); Fibroblast growth factor 8 (androgen-induced); Granulysin; Cocaine- and

amphetamine-regulated transcript; Collagen, type I, alpha 2; Chemokine (C-C motif) ligand 17; Chemokine (C-C motif) ligand 23 ; Sparc/osteonectin, cwcw and kazal-like domains proteoglycan (testican) 3; Gamma-aminobutyric acid (GABA) A receptor, beta 3; Defensin, alpha 4, corticostatin; Leucine rich repeat
 5 neuronal 3; Glypican 6; Mitogen-activated protein kinase kinase 2; Coagulation factor XI (plasma thromboplastin antecedent); Chemokine (C-C motif) ligand 5; Dystonin; Frizzled-related protein; Coagulation factor XIII, A1 polypeptide; Insulin-like growth factor 1 (somatomedin C); Hypothetical protein MGC45438; Sperm associated antigen 11; Insulin-like growth factor 1 (somatomedin C);
 10 Periostin, osteoblast specific factor; Alpha-2-macroglobulin; Gamma-aminobutyric acid (GABA) A receptor, alpha 5; Serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 3; Silver homolog (mouse); Frizzled-related protein; Chondroadherin; Chondroitin beta1,4 N-acetylgalactosaminyltransferase; 5-hydroxytryptamine (serotonin)
 15 receptor 3, family member C; Collagen, type VI, alpha 2; Toll-like receptor 9; Amelogenin, Y-linked; Vascular endothelial growth factor B; Radial spokehead-like 1; Fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor); Protease inhibitor 16; Interleukin 2; Clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J); Follicle stimulating hormone, beta polypeptide; A disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 16; Lysozyme (renal amyloidosis); radical fringe homolog (Drosophila); Insulin-like growth factor binding protein 5; Taxilin; Apolipoprotein A-V; Platelet derived growth factor C;
 25 Chemokine (C-C motif) ligand 3-like 1; Fibroblast growth factor 16; Collagen, type VI, alpha 2; Serine (or cysteine) proteinase inhibitor, clade C (antithrombin), member 1; Chemokine (C-C motif) ligand 11; Collagen, type IV, alpha 4; Bruton agammaglobulinemia tyrosine kinase; Insulin-like growth factor 2 (somatomedin A); Kazal-type serine protease inhibitor domain 1; Fibrinogen,
 30 A alpha polypeptide; Chemokine (C-C motif) ligand 1; Inhibin, beta E; Sex hormone-binding globulin; Collagen, type IV, alpha 1; Lecithin-cholesterol acyltransferase; Cysteine-rich secretory protein 2; Hyaluronan and proteoglycan link protein 1; Natriuretic peptide precursor C; Ribonuclease, RNase A family, k6; Fibroblast growth factor 14; ADAMTS-like 2; Collagen, type IV, alpha 3

(Goodpasture antigen); Angiopoietin 2; Apolipoprotein L, 3; Chemokine (C-X-C motif) ligand 12 (stromal cell-derived factor 1); Hyaluronan binding protein 2; Coagulation factor VII (serum prothrombin conversion accelerator); collagen, type XIV, alpha 1 (undulin) ; Oviductal glycoprotein 1, 120kDa (mucin 9, oviductin); Matrilin 1, cartilage matrix protein; mucin 5, subtypes A and C, tracheobronchial/gastric; Tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin); Transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase); Keratocan; Collagen, type V, alpha 3; WAP four-disulfide core domain 2; Chemokine (C-X3-C motif) ligand 1; Serine (or cysteine) proteinase inhibitor, clade D (heparin cofactor), member 1; Secretory protein LOC348174; Coagulation factor X; Interleukin 16 (lymphocyte chemoattractant factor); Pancreatic lipase-related protein 2; HtrA serine peptidase 3; Glycine receptor, alpha 3; CD5 antigen-like (scavenger receptor cysteine rich family); Hypothetical protein MGC39497; Coagulation factor VIII, procoagulant component (hemophilia A); Dermatopontin; Noggin; Secreted LY6/PLAUR domain containing 1; ADAMTS-like 1; Alpha-1-B glycoprotein; Chromosome 20 open reading frame 175; Wingless-type MMTV integration site family, member 8B; Fibulin 1; Fibulin 5; Cathepsin S; Nidogen (enactin); Chemokine (C-C motif) ligand 26; Endothelial cell-specific molecule 1; Chitinase 3-like 1 (cartilage glycoprotein-39); Gamma-aminobutyric acid (GABA) A receptor, beta 1; Secretoglobulin, family 1D, member 2; Mannan-binding lectin serine protease 1 (C4/C2 activating component of Ra-reactive factor); ADAMTS-like 1; Sema domain, immunoglobulin domain (Ig), and GPI membrane anchor, (semaphorin) 7A; A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 15; Proprotein convertase subtilisin/kexin type 2 ; Insulin-like growth factor 1 (somatomedin C); Retinoschisis (X-linked, juvenile) 1; A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 16; Chemokine (C motif) ligand 2; Fibroblast growth factor 5; Sperm associated antigen 11; Microfibrillar-associated protein 4; Poliovirus receptor; Extracellular signal-regulated kinase 8; Transmembrane protease, serine 6; Protein kinase C, alpha; Chitinase 3-like 2; Interleukin 9; Apolipoprotein L, 6; Surfactant, pulmonary-associated protein A1; Collagen, type VI, alpha 1; Apolipoprotein L, 6; Hypothetical protein FLJ13710; Carboxypeptidase B2 (plasma, carboxypeptidase U) ;

Bactericidal/permeability-increasing protein-like 2; Fibroblast growth factor 5;
 Secreted phosphoprotein 1 (osteopontin, bone sialoprotein 1, early T-
 lymphocyte activation 1); HtrA serine peptidase 3; Deleted in liver cancer 1;
 Endothelial cell-specific molecule 1; Von Willebrand factor; A disintegrin-like
 5 and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 5
 (aggrecanase-2); Sema domain, immunoglobulin domain (Ig), short basic
 domain, secreted, (semaphorin) 3A; Chemokine (C-X-C motif) ligand 12
 (stromal cell-derived factor 1); Statherin; Extracellular signal-regulated kinase 8;
 Tissue inhibitor of metalloproteinase 3 (Sorsby fundus dystrophy,
 10 pseudoinflammatory); Platelet factor 4 (chemokine (C-X-C motif) ligand 4);
 Surfactant, pulmonary-associated protein D; Complement factor H; Delta-like 1
 homolog (Drosophila); WAP four-disulfide core domain 1; Insulin-like growth
 factor binding protein, acid labile subunit; Breast cancer 2, early onset; Pre-B
 lymphocyte gene 1; Corticotropin releasing hormone; Hypothetical protein
 15 DKFZp434B044; Prolactin-induced protein; RAS guanyl releasing protein 4;
 Progastricsin (pepsinogen C); Sema domain, immunoglobulin domain (Ig), short
 basic domain, secreted, (semaphorin) 3F; Upregulated in colorectal cancer
 gene 1; Proteoglycan 4; Cholinergic receptor, nicotinic, delta polypeptide;
 Cartilage oligomeric matrix protein; ABO blood group (transferase A, alpha 1-3-
 20 N-acetylgalactosaminyltransferase; transferase B, alpha 1-3-
 galactosyltransferase); Interleukin 12A (natural killer cell stimulatory factor 1,
 cytotoxic lymphocyte maturation factor 1, p35); Fibroblast growth factor 7
 (keratinocyte growth factor); Kin of IRRE like 3 (Drosophila); Cholinergic
 receptor, nicotinic, alpha polypeptide 2 (neuronal); Palate, lung and nasal
 25 epithelium carcinoma associated; Collagen, type XV, alpha 1; Pleiotrophin
 (heparin binding growth factor 8, neurite growth-promoting factor 1);
 Angiopoietin-like 2; Norrie disease (pseudoglioma); Chemokine (C-C motif)
 ligand 3; Chitinase 3-like 1 (cartilage glycoprotein-39); Inter-alpha (globulin)
 inhibitor H3; Amelogenin (amelogenesis imperfecta 1, X-linked); Epidermal
 30 growth factor (beta-urogastrone); Fibroblast growth factor 13; Wingless-type
 MMTV integration site family, member 7B; Cholinergic receptor, nicotinic,
 gamma polypeptide; Pregnancy specific beta-1-glycoprotein 6; Matrix
 metalloproteinase 14 (membrane-inserted); Chemokine (C-C motif) ligand 26;
 Interferon, alpha 6; Tachykinin, precursor 1 (substance K, substance P,

neurokinin 1, neurokinin 2, neuromedin L, neurokinin alpha, neuropeptide K,
 neuropeptide gamma); Secreted frizzled-related protein 5; Hyaluronan and
 proteoglycan link protein 4; Complement component 4B; Matrix
 metalloproteinase 16 (membrane-inserted); Fibroblast growth factor 7
 5 (keratinocyte growth factor); Apolipoprotein C-II; Chloride channel, calcium
 activated, family member 3; Tetranectin (plasminogen binding protein);
 Collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal
 dominant); KIAA0556 protein; Chemokine (C-C motif) ligand 4; Hemopexin;
 Inter-alpha (globulin) inhibitor H1; Relaxin 1; Matrix Gla protein; A disintegrin-
 10 like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 2;
 Interferon (alpha, beta and omega) receptor 2; Acid phosphatase, prostate;
 Guanine nucleotide binding protein (G protein), gamma 8; Matrix
 metalloproteinase 23B; Meprin A, alpha (PABA peptide hydrolase);
 Hyaluronoglucosaminidase 1; Angiotensinogen (serine (or cysteine) proteinase
 15 inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 8); Cartilage
 intermediate layer protein, nucleotide pyrophosphohydrolase; Purinergic
 receptor P2X, ligand-gated ion channel, 7; Glypican 3; Tectorin beta; Interferon,
 alpha 5; Lipocalin 7; Platelet factor 4 variant 1; Nucleobindin 1; Collagen, type
 XI, alpha 1; Gastric inhibitory polypeptide; Thrombospondin repeat containing 1;
 20 5-hydroxytryptamine (serotonin) receptor 3 family member D; Collagen, type
 XXV, alpha 1; Growth differentiation factor 9; Hypothetical protein
 DKFZp434B044; Endothelin 3; Chemokine (C motif) ligand 2; Prokineticin 2;
 Tumor necrosis factor receptor superfamily, member 11b (osteoprotegerin);
 Tissue inhibitor of metalloproteinase 2; Dystonin; Chromogranin B
 25 (secretogranin 1); Hyaluronan and proteoglycan link protein 2; Leucine rich
 repeat neuronal 3 ; Lumican ; Matrilin 1, cartilage matrix protein ; Phospholipase
 A2, group IIA (platelets, synovial fluid); Carboxylesterase 1
 (monocyte/macrophage serine esterase 1); Sparc/osteonectin, cwcw and kazal-
 like domains proteoglycan (testican); Dickkopf homolog 2 (*Xenopus laevis*);
 30 Gamma-aminobutyric acid (GABA) A receptor, alpha 3; Pregnancy specific
 beta-1-glycoprotein 11; Insulin-like growth factor binding protein 1; Defensin,
 beta 106; Interleukin 17F; Ligand-gated ion channel subunit ; Phospholipase A2
 receptor 1, 180kDa; I factor (complement); Dystonin; LAG1 longevity assurance
 homolog 1 (*S. cerevisiae*); Prolactin; Testis expressed sequence 264; Sema

domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3D; secreted frizzled-related protein 2; secreted frizzled-related protein 4).

[0091] There are groups of genes present only in UCEC. These genes are related to the following: Homeostasis (Albumin; Calcium-sensing receptor; Aquaporin 9; Lactotransferrin. Morphogenesis: Homeo box HB9; Epithelial V-like antigen 1). Embryonic Development (Relaxin 2; Carcinoembryonic antigen-related cell adhesion molecule 8; Indoleamine-pyrrole 2,3 dioxygenase; EPH receptor A3; Thyrotrophic embryonic factor; Pregnancy specific beta-1-glycoprotein 1; Laminin, alpha 3), the Extracellular Space (Surfactant, pulmonary-associated protein A1; Pregnancy specific beta-1-glycoprotein 1; Lactotransferrin; TGF-alpha; Albumin; FGF-23; S100 calcium binding protein A9 (calgranulin B)), the Extracellular Matrix (Laminin, beta 4; Laminin, alpha 3; Zona pellucida glycoprotein 4. Structural Molecule Activity: Chromosome 21 open reading frame 29; Laminin, alpha 3; Microtubule-associated protein 2; Laminin, beta 4; Keratin 6B; Ladinin 1; Keratin 6A; Occludin; Loricrin; Erythrocyte membrane protein band 4.1 (elliptocytosis 1, RH-linked); Crystallin, beta A2; eye lens structural protein; Contactin associated protein-like 4; Claudin 19; Hypothetical protein LOC144501; Keratin 6E; Keratin 6L; Lens intrinsic membrane protein 2, 19kDa), the Cytoskeleton (Microtubule-associated protein 2; Erythrocyte membrane protein band 4.1 like 5; Homo sapiens trichohyalin (THH); Keratin 6B; Keratin 6A; Epithelial V-like antigen 1; Hook homolog 1 (Drosophila); Loricrin; Erythrocyte membrane protein band 4.1 (elliptocytosis 1, RH-linked); Tropomodulin 1; MAP/microtubule affinity-regulating kinase 1; Keratin 6E; Actin binding LIM protein family, member 2), Cell Adhesion Molecules (Cadherin 19, type 2; Myeloid/lymphoid or mixed-lineage leukemia; Chromosome 21 open reading frame 29; Kin of IRRE like 2; Laminin, alpha 3; Sialoadhesin; CD84 antigen (leukocyte antigen); Lectin, galactoside-binding, soluble, 2 (galectin 2); Epithelial V-like antigen 1; CD96 antigen; Tubulointerstitial nephritis antigen; Carcinoembryonic antigen-related cell adhesion molecule 8; IL-18; Immunoglobulin superfamily, member 1; Integrin, beta 8; Ornithine carbamoyltransferase; Integrin, beta 6; Contactin associated protein-like 4; Collagen, type XVII, alpha 1; Cadherin-like 26; Mucin and cadherin-like), Cell Differentiation proteins (Protein tyrosine phosphatase,

receptor-type, Z polypeptide 1; Laminin, alpha 3; CD84 antigen (leukocyte antigen); EDRF2; Homo sapiens erythroid differentiation-related factor 2; Tumor protein p73-like; NB4 apoptosis/differentiation related protein; Homo sapiens PNAS-133; Similar to seven in absentia 2; Interleukin 24; Keratin 6B; Keratin 6A; Dehydrogenase/reductase (SDR family) member 9; Gap junction protein, beta 5 (connexin 31.1); Iroquois homeobox protein 4; Ventral anterior homeobox 2; Chemokine (C-X-C motif) ligand 10; Tumor necrosis factor receptor superfamily, member 17; Calcium channel, voltage-dependent, beta 2 subunit; Parkinson disease (autosomal recessive, juvenile) 2, parkin; Kallikrein 7 (chymotryptic, stratum corneum); Glial cells missing homolog 2; AP-2 alpha; Protein tyrosine phosphatase, receptor-type, Z polypeptide 1; Troponin T1; Sciellin; Glucosaminyl (N-acetyl) transferase 2, I-branching enzyme; Collagen, type XVII, alpha 1; Suppressor of cytokine signaling 2; Distal-less homeo box 1; Zygote arrest 1; Interleukin 20; Growth differentiation factor 3; FGF-23; Wingless-type MMTV integration site family, member 8A. Extracellular: Chromosome 21 open reading frame 29; Laminin, alpha 3; Laminin, beta 4; Interleukin 24; Pregnancy specific beta-1-glycoprotein 1; Chemokine (C-X-C motif) ligand 11; Surfactant, pulmonary-associated protein A1; Prepronociceptin; 5-hydroxytryptamine (serotonin) receptor 3B; Carcinoembryonic antigen-related cell adhesion molecule 8; Chemokine (C-X-C motif) ligand 10; IL-18 (interferon-gamma-inducing factor); Lactotransferrin; Albumin; Fas ligand (TNF superfamily, member 6); Cholinergic receptor, nicotinic, beta polypeptide 4; Cathelicidin antimicrobial peptide; Airway trypsin-like protease; S100 calcium binding protein A9 (calgranulin B); TGF-alpha; Kallikrein 10; Serine protease inhibitor, Kunitz type 1; WNT1 inducible signaling pathway protein 3; Relaxin 2; Interferon, kappa; Defensin, beta 103A; IL-20; Zona pellucida glycoprotein 4; Growth differentiation factor 3; FGF-23; Wingless-type MMTV integration site family, member 8A; Complement factor H-related 5; Developmental proteins (EPH receptor A3; NIMA (never in mitosis gene a)-related kinase 2; Zinc finger protein 282; TANK-binding kinase 1; MRE11 meiotic recombination 11 homolog A; E2F transcription factor 2; Protein tyrosine phosphatase, receptor-type, Z polypeptide 1; Homo sapiens clone 161455 breast expressed mRNA from chromosome X; Laminin, alpha 3; v-myb myeloblastosis viral oncogene homolog (avian)-like 1; Regulator of G-protein

signalling 11; Microtubule-associated protein 2; Transmembrane protein 16A;
 Adenomatosis polyposis coli 2; Homeo box HB9; Centromere protein F,
 350/400ka (mitosin); CD84 antigen (leukocyte antigen); EDRF2; Homo sapiens
 erythroid differentiation-related factor 2; Tumor protein p73-like; NB4
 5 apoptosis/differentiation related protein; Homo sapiens PNAS-133; Forkhead
 box P2; Homo sapiens gastric-associated differentially-expressed protein
 YA61P (YA61); Tenascin N; Chromosome 6 open reading frame 49; Zinc finger
 protein 462; Zinc finger protein 71 (Cos26); SRY (sex determining region Y)-box
 7; Triggering receptor expressed on myeloid cells-like 4; Interleukin 24;
 10 Pregnancy specific beta-1-glycoprotein 1; Chondroitin sulfate proteoglycan 5
 (neuroglycan C); Keratin 6B; Keratin 6A; Dehydrogenase/reductase (SDR
 family) member 9; Epithelial V-like antigen 1; Gap junction protein, beta 5
 (connexin 31.1); G protein-coupled receptor 51; Interferon regulatory factor 6;
 Neurotrophin 5 (neurotrophin 4/5); CD96 antigen; Iroquois homeobox protein 4;
 15 Interleukin 1 receptor-like 1; G-2 and S-phase expressed 1; Nuclear receptor
 subfamily 2, group E, member 3; Ventral anterior homeobox 2; Zinc finger
 protein 215; DNA segment on chromosome 4 (unique) 234 expressed
 sequence; Carcinoembryonic antigen-related cell adhesion molecule 8;
 Chemokine (C-X-C motif) ligand 10; IL-18; Indoleamine-pyrrole 2,3
 20 dioxygenase; Albumin; Calcium-sensing receptor (hypocalciuric hypercalcemia
 1, severe neonatal hyperparathyroidism); Fas ligand (TNF superfamily, member
 6); TNFR superfamily, member 17; Calcium channel, voltage-dependent, beta 2
 subunit; Parkinson disease (autosomal recessive, juvenile) 2, parkin; Kallikrein
 7 (chymotryptic, stratum corneum); Glial cells missing homolog 2; TGF- α ;
 25 Thyrotrophic embryonic factor; AP-2 α (activating enhancer binding protein
 2 α); Kallikrein 10; Regulator of G-protein signalling 7; Protein tyrosine
 phosphatase, receptor-type, Z polypeptide 1; Serine protease inhibitor, Kunitz
 type 1; WNT1 inducible signaling pathway protein 3; Zic family member 3
 heterotaxy 1 (odd-paired homolog, Drosophila); TTK protein kinase; Troponin
 30 T1, skeletal, slow; Sciellin; TGFB-induced factor 2-like, X-linked; Kallikrein 8
 (neuropsin/ovasin); Glucosaminyl (N-acetyl) transferase 2, I-branching enzyme;
 Ankyrin repeat domain 30A; Relaxin 2; Collagen, type XVII, α 1; Gene
 differentially expressed in prostate; Phosphatase and actin regulator 3;
 Suppressor of cytokine signaling 2; Nuclear receptor subfamily 4, group A,

member 3; Angiotensin I converting enzyme (peptidyl-dipeptidase A) 1; Hypothetical protein MGC17986; Distal-less homeo box 1; LAG1 longevity assurance homolog 3 (*S. cerevisiae*); Zygote arrest 1; Interferon, kappa; IL-20; ICEBERG caspase-1 inhibitor; Growth differentiation factor 3; FGF-23; Testis expressed sequence 15; Wingless-type MMTV integration site family, member 8A; SRY (sex determining region Y)-box 7; Carnitine deficiency-associated, expressed in ventricle 1; Prokineticin 1; CAMP responsive element binding protein 3-like 3; Caspase recruitment domain family, member 15; FLJ23311 protein).

10

Example 6: Direct differentiation of umbilical cord epithelial stem cells (UCEC) into skin epidermal keratinocytes

[0092] For differentiation into skin epidermal keratinocytes, umbilical cord epithelial stem cells, UCEC cells, were cultured according to a standard protocol for the cultivation of keratinocytes. Cell isolation techniques were as described above. UCEC were then cultured in serum-free keratinocyte growth media, KGM, KGM-2 (Cambrex), EpiLife (Cascade Biologics) or in Green's medium in the presence of irradiated or Mytomycin-C treated 3T3 mouse embryonic feeder layer at 37°C, 5% CO₂). UCEC cell morphology thus differentiated resembled human epidermal keratinocytes. Epithelial cells have similar morphology under light microscope and can be easily turned into fibroblasts using conventional and commercially available media (cf., Figure 2).

[0093] Immunofluorescent analysis shows that the cultivated UCEC also express epidermal keratinocyte molecular markers such as keratins, desmosome, hemidesmosome and basement membrane components (see also Fig. 10 that shows that UCEC are qualified to be epithelial cells in general by expressing a variety of these epithelial cell markers). Accordingly, these results show that umbilical cord epithelial progenitor/stem cells of the present invention can be differentiated into skin cells such as epidermal keratinocytes which can be used for wound healing and have great potential for the development of cultured skin equivalents.

30

Example 7: Expansion of umbilical cord epithelial and mesenchymal stem cells using repetitive tissue explants of umbilical cord lining membrane

tissues

[0094] Umbilical cord epithelial and mesenchymal stem cells of the invention were expanded using repetitive explants of umbilical cord amniotic membrane tissue as follows. Briefly, at day 1 of process, tissue explants were
 5 plated onto tissue culture dishes in growth media (DMEM/10%FCS, EpiLife, KGM, KGM-2 or M171) at 37°C, 5%CO₂; media was changed every 2 or 3 days. Cell outgrowths started and continued migrating from the explants for 7 days. After that, tissue explants were transferred to other dishes to allow further cell outgrowth. This process was continued until the explants had diminished in
 10 size, preventing further explantation. In this connection it is noted that the explants progressively shrink in size until they are too small for further tissue explant since during the process of cells outgrowing and migrating from tissue explants, the cells produce proteases to digest and break down tissue. **Fig. 15** schematically illustrates the rapid and robust expansion process of umbilical
 15 cord epithelial and mesenchymal stem cells achieved using this protocol. Thus, this study demonstrates the high yield of UCMC and UMEC cells can be obtained from this source, further reflecting the high viability and pro-growth characteristics of these cells in comparison with other sources of cells as bone-marrow or adipose-derived stem cells. In addition, being a solid tissue, the
 20 successful repetitive explant technique used herein demonstrates that the cells of the invention can be uniformly extracted from the entire tissue instead of only certain portions. This allows the maximum number of cells that can be derived at a low passage instead of passing the cells through many generations causing deterioration of cells.

25 **Example 8: Direct differentiation of umbilical cord mesenchymal cells (UCMC) into skin dermal fibroblasts**

[0095] For differentiation into skin dermal fibroblasts, umbilical cord mesenchymal stem cells, UCMC cells were cultured according to a standard protocol for the cultivation of fibroblasts. Cell isolation techniques were as
 30 described above in Example 6. UCMC were then cultured in DMEM or commercially available fibroblast growth media (FGM). UCMC cell morphology thus differentiated resembled human dermal fibroblasts. Mesenchymal cells have similar morphology under light microscope and can be easily turned into fibroblasts using conventional and commercially available media (cf., Figure 3).

35

CLAIMS

What is claimed is:

1. A method for isolating mesenchymal stem/progenitor cells from the amniotic membrane of umbilical cord, the method comprising:
 - (a) separating the amniotic membrane from the other components of the umbilical cord *in vitro*;
 - (b) culturing the amniotic membrane tissue obtained in step (a) as tissue explant or culturing amniotic membrane cells separated from the amniotic membrane tissue obtained in step (a) under conditions allowing cell proliferation of mesenchymal stem/progenitor cells without differentiation of the mesenchymal stem/progenitor cells; and
 - (c) isolating the mesenchymal stem/progenitor cells, wherein the mesenchymal stem/progenitor cells have a spindle shape, express the gene POU5f1 which encodes the transcription factor Octamer-4 (Oct-4), and secrete Activin A and Follistatin.
2. The method of claim 1,
wherein the amniotic membrane cells are separated from the amniotic membrane tissue obtained in (a) by enzymatic separation.
3. The method of any one of claims 1 to 2, further comprising:
 - (d) culturing the mesenchymal stem/progenitor cells under conditions allowing the cells to undergo clonal expansion.
4. The method of claim 3, further comprising obtaining a mesenchymal cell differentiated from the epithelial stem/progenitor cells by:
 - (e) culturing the mesenchymal stem/progenitor cells under conditions allowing the differentiation of said cells into mesenchymal cells; and
 - (f) isolating the differentiated mesenchymal cells.
5. The method of claim 4, wherein the differentiated mesenchymal cells are selected from the group consisting of: skin fibroblasts, chondrocytes, osteoblasts, tenocytes, ligament

fibroblasts, cardiomyocytes, smooth muscle cells, skeletal muscle cells, adipocytes, cells derived from endocrine glands, and neurectodermal cells.

6. The method of any one of claims 1 to 3, further comprising
(g) preserving the isolated stem/progenitor cells for further use.
7. The method of claim 6, wherein preserving is carried out by using cryo-preservation.
8. The mesenchymal stem/progenitor cell isolated from the amniotic membrane of umbilical cord by means of the method of any one of claims 1 to 3 or 6 to 7.
9. A cell bank comprising a mesenchymal stem/progenitor cell as defined in claim 8.
10. A pharmaceutical composition comprising a mesenchymal stem/progenitor cell as defined in claim 8, the composition further comprising a pharmaceutically acceptable carrier.
11. The pharmaceutical composition of claim 10, wherein the pharmaceutical composition is adapted for systemic or topical application.
12. The pharmaceutical composition of claim 11, wherein the pharmaceutical composition adapted for topical application is selected from the group consisting of an ointment, a cream, and a lotion.
13. The use of a mesenchymal stem/progenitor cell as defined in claim 8 for the manufacture of a pharmaceutical composition for treating a subject having a disorder, wherein the disorder is selected from the group consisting of tissue disorder, neoplastic disease, visceral endocrine deficiency, skin disease, wherein the skin disease is accelerated aging or a wound.
14. The use of claim 13, wherein the tissue disorder is a congenital or an acquired tissue deficiency.
15. The use of claim 13, wherein the visceral endocrine deficiency is selected from the group consisting of insulin deficiency, Diabetes mellitus associated with insulin deficiency,

testosterone deficiency, anemia, hypoglycemia, hyperglycemia, pancreatic deficiency, adrenal deficiency, and thyroid abnormality.

16. The use of claim 13, wherein the neoplastic disease is cancer.

17. The use of claim 16, wherein the cancer is a selected from the group consisting of squamous cell carcinoma, breast ductal and lobular carcinoma, hepatocellular carcinoma, nasopharyngeal carcinoma, lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular malignant melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, testicular cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, non-Hodgkin's lymphoma, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemias, solid tumors of childhood, lymphocytic lymphoma, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer and any combination of such cancers.

18. The use of a mesenchymal stem/progenitor cell as defined in claim 8 for the production of a biological molecule, wherein the biological molecule is selected from the group consisting of a protein, a peptide, a small organic molecule, an oligosaccharide, a polysaccharide, a proteoglycan and a lipid.

19. The use of a mesenchymal stem/progenitor cell as defined in claim 8 as a feeder layer in the cultivation of mammalian cells.

20. A method of isolating mesenchymal stem/progenitors cells from the amniotic membrane of umbilical cord, comprising:


Obtaining a tissue explant from the amniotic membrane of umbilical cord;

Cultivating the tissue explant in suitable cultivation media and cultivation conditions allowing cell proliferation of mesenchymal stem/progenitor cells without differentiation of the mesenchymal stem/progenitor cells over a suitable period of time, wherein the mesenchymal stem/progenitor cells have a spindle shape, express the gene POU5f1 which encodes the transcription factor Octamer-4 (Oct-4) and secrete Activin A and Follistatin.

21. The method of claim 20, further comprising exposing the tissue explant to fresh cultivation media and continuing the cultivation under suitable conditions over a suitable period of time.



A




C




B

FIGURE 1


FIGURE 2




A



B



C



D





FIGURE 3



B




A

FIGURE 4


Figure 5: Morphology of normal dermal fibroblasts (NF) and adipose-derived mesenchymal cells (ADMC) or umbilical cord mesenchymal stem cells (UCMC) in serum-free (DMEM) and serum medium (DMEM/10%FCS)

A




NF109 in 10% FCS

B




NF109 in DMEM

C




ADMC3 in 10% FCS

D




ADMC3 in DMEM

Figure 5: continued...



E

UCMC3 in DMEM/10%FCS




G

UCMC4 in DMEM/10%FCS



F


UCMC3 in DMEM




H

UCMC4 in DMEM

Figure 6: Colony formation of umbilical cord mesenchymal stem cells cultured in non-feeder layer condition in DMEM/10%FCS




Day 3




Day 7

Figure 7: Colony formation of umbilical cord epithelial stem cells cultured in feeder layer condition in Green's medium





Day 3





Day 7

Figure 8: Colony formation of umbilical cord mesenchymal stem cells cultured on feeder layer condition in Green's medium



Day 7



Day 3

FIGURE 9-1: Expression of Oct-4 in Umbilical Cord Epithelial Cells and Mesenchymal Cells




FIGURE 9-2: Expression of STAT3 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-3: Expression of STAT3 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-4: Expression of Placental Growth Factor (PLGF) in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-5: Expression of Placental Growth Factor (PLGF) in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-6: Expression of Connective Tissue Growth Factor (CTGF) in Umbilical Cord Mesenchymal Stem Cells (UCMC)






**FIGURE 9-7: Expression of Connective Tissue Growth Factor (CTGF) in Umbilical Cord
Mesenchymal Stem Cells (UCMC)**



1 2 3 4 5 6 7 8 9 10 11 12 13 14

- | | |
|-----------------|-------------------|
| 1 UCMC-3 | 9 hBMSC-1 |
| 2 UCMC-3 | 10 hBMSC-1 |
| 3 UCMC-3 | 11 ADMC |
| 4 UCMC-3 | 12 ADMC |
| 5 UCMC-2 | 13 hBMSC-3 |
| 6 UCMC-2 | 14 hBMSC-3 |
| 7 UCMC-3 | |
| 8 UCMC-3 | |

FIGURE 9-8: Expression of Platelet Derived Growth Factor (PDGF) in Umbilical Cord Mesenchymal Stem Cells (UCMC)



A vertical strip of a textured, woven fabric, possibly a book cover or endpaper. The fabric features a repeating pattern of light and dark squares, creating a checkered or plaid appearance. The texture is highly detailed, showing the individual threads and the interlacing of the weave. The lighting is even, highlighting the tactile quality of the material.

1 UCMC-3
2 UCMC-3
3 UCMC-3
4 UCMC-3
5 UCMC-2
6 UCMC-2
7 UCMC-3
8 UCMC-3

9 NF (ear)
10 NF (ear)
11 NF 106
12 NF 106
13 NF 112
14 NF 112

A vertical strip of a woven fabric, likely silk, showing a complex pattern of dark and light threads forming a series of vertical bands and horizontal lines. The pattern is dense and intricate, with varying shades of grey and black on a lighter background.

1	2	3	4	5	6	7	8	9	10	11	12
1	UCMC-3						7	NF (ear)			
2	UCMC-3						8	NF (ear)			
3	UCMC-2						9	NF 106			
4	UCMC-2						10	NF 106			
5	UCMC-3						11	NF 112			
6	UCMC-3						12	NF 112			

**FIGURE 9-11: Expression of Vascular Endothelial Growth Factor (VEGF) in Umbilical Cord
Mesenchymal Stem Cells (UCMC)**




FIGURE 9-12: Expression of Fibroblast Growth Factor-2 (FGF-2) in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-13: Expression of Fibroblast Growth Factor-2 (FGF-2) in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-14: Expression of Hepatoma Derived Growth Factor (HDGF) in Umbilical Cord Mesenchymal Stem Cells (UCMC)




**FIGURE 9-15: Expression of Hepatoma Derived Growth Factor (HDGF) in Umbilical Cord
Mesenchymal Stem Cells (UCMC)**




FIGURE 9-16: Expression of Stem Cell Growth Factor (SCF) in Umbilical Cord Mesenchymal Stem Cells (UCMC)




**FIGURE 9-17: Expression of alpha-Smooth Muscle Actin (α -SMA) in Umbilical Cord
Mesenchymal Stem Cells (UCMC)**





FIGURE 9-18: Expression of Fibronectin in Umbilical Cord Mesenchymal Stem Cells (UCMC)



1 UCMC-3
2 UCMC-3
3 UCMC-3
4 UCMC-3
5 UCMC-2
6 UCMC-2
7 UCMC-3
8 UCMC-3

9 hBMSC-1
10 hBMSC-1
11 ADMC
12 ADMC
13 hBMSC-3
14 hBMSC-3

FIGURE 9-19: Expression of Fibronectin in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-20: Expression of Decorin in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-21: Expression of Syndecan-1 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-22: Expression of Syndecan-2 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-23: Expression of Syndecan-2 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-24: Expression of Syndecan-3 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




FIGURE 9-25: Expression of Syndecan-3 in Umbilical Cord Mesenchymal Stem Cells (UCMC)





FIGURE 9-26: Expression of Syndecan-4 in Umbilical Cord Mesenchymal Stem Cells (UCMC)




1 2 3 4 5 6 7 8 9 10 11 12 13 14

**1 UCMC-1
2 UCMC-1
3 UCMC-3
4 UCMC-3
5 UCMC-2
6 UCMC-2
7 UCMC-4
8 UCMC-4**


**9 UCMC-5
10 UCMC-5
11 NF 106
12 NF 106
13 NF 112
14 NF 112**

FIGURE 10-1: Expression of Cytokeratins in Umbilical Cord Epithelial Cells


CK general




CK17




CK6




CK10



CK19



CK18



CK16



CK15






**FIGURE 10-2: Expression of Hemidesmosome and Desmosome
in Umbilical Cord Epithelial Cells**

Integrin $\alpha 6$



Integrin $\beta 4$



Desmosome





**FIGURE 10-3: Expression of Basement Membrane Proteins
in Umbilical Cord Epithelial Cells**


Laminin 1



Laminin 5



Collagen IV




Collagen VII



**FIGURE 10-4: Expression of Integrin- β 1 and Fibronectin
in Umbilical Cord Epithelial Cells**

Integrin β 1



Fibronectin




FIGURE 11: Expression Profile of Secreted Cytokines and Growth Factors by Umbilical Cord Mesenchymal Cells




FIGURE 11: Expression Profile of Secreted Cytokines and Growth Factors by Umbilical Cord Mesenchymal Cells





FIGURE 11: Expression Profile of Secreted Cytokines and Growth Factors by Bone Marrow Mesenchymal Cells



42/65

1,2-IL6; 3,4-MCP1; 5,6-RANTES

FIGURE 11: Expression Profile of Secreted Cytokines and Growth Factors by Bone Marrow Mesenchymal Cells






FIGURE 12: Expression Profile of Secreted Cytokines and Growth Factors by Umbilical Cord Epithelial Cells




1,2-IGFBP-2; 3,4-IL6; 5,6-IL1R α ; 7,8-MCP-1

FIGURE 12: Expression Profile of Secreted Cytokines and Growth Factors by Umbilical Cord Epithelial Cells



**1,2-IL8; 3,4-sTNFR; 5,6-GRO; 7,8-Amphiregulin; 9,10-TIMP1;
11,12-TIMP2; 13,14-uPAR;15,16-VEGF**

FIGURE 12: Expression Profile of Secreted Cytokines and Growth Factors by Human Epidermal Keratinocytes



1,2-IL6; 3,4-IL1R α ; 5,6-MCP-1; 7,8-RANTES; 9,10-TNF β

FIGURE 12: Expression Profile of Secreted Cytokines and Growth Factors by Human Epidermal Keratinocytes




FIGURE 12: Expression Profile of Secreted Cytokines and Growth Factors by Umbilical Cord Epithelial Cells




Fig.12
Spotting of Chip used for Cytokine Array (6.1)

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
1	POS	POS	PO S	PO S	Blank	Angiog enin	BDN F	BLC	BMP- 4	BMP -6	CK b 8-1	CNT F	EGF	Eota xin
2	NEG	NEG	NE G	NE G	Blank	Angiog enin	BDN F	BLC	BMP- 4	BMP -6	CK b 8-1	CNT F	EGF	Eota xin
3	Eotaxi n-2	Eotax in-3	FGF -6	FGF -7	Fit-3 Ligand	Fractal kine	GCP -2	GD NF	GM- CSF	I-309	IFN-g	IGFB P-1	IGFB P-2	IGFB P-4
4	Eotaxi n-2	Eotax in-3	FGF -6	FGF -7	Fit-3 Ligand	Fractal kine	GCP -2	GD NF	GM- CSF	I-309	IFN-g	IGFB P-1	IGFB P-2	IGFB P-4
5	IGF-I	IL-10	IL- 13	IL- 15	IL-16	IL-1a	IL-1b	IL- 1ra	IL-2	IL-3	IL-4	IL-5	IL-6	IL-7
6	IGF-I	IL-10	IL- 13	IL- 15	IL-16	IL-1a	IL-1b	IL- 1ra	IL-2	IL-3	IL-4	IL-5	IL-6	IL-7
7	Leptin	LIGH T	MC P-1	MC P-2	MCP-3	MCP-4	CSF	MD C	MIG	MIP- 1 d	MIP- 3a	NAP- 2	NT-3	PAR C
8	Leptin	LIGH T	MC P-1	MC P-2	MCP-3	MCP-4	CSF	MD C	MIG	MIP- 1 d	MIP- 3a	NAP- 2	NT-3	PAR C
9	PDGF -BB	RANT ES	SCF	SDF -1	TARC	TGF- b1	TGF- b3	TNF -a	TNF- b	Blan k	Blank	Blank	Blank	Blank
10	PDGF -BB	RANT ES	SCF -1	SDF -1	TARC	TGF- b1	TGF- b3	TNF -a	TNF- b	Blan k	Blank	Blank	POS	POS

Fig.12
Spotting of Chip used for Cytokine Array (7.1)

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
1	POS	POS	POS	POS	Blank	Acrrp30	AgR P	Angiopoietin-2	Amphiregulin	Axl	bFGF	b-NGF	BTC	CCL-28
2	NEG	NEG	NEG	NEG	Blank	Acrrp30	AgR P	Angiopoietin-2	Amphiregulin	Axl	bFGF	b-NGF	BTC	CCL-28
3	CTA CK	Dtk	EGF-R	ENA-78	Fas	FGF-4	FGF-9	GCSF	GITR-Ligand	GITR	GR O	GR O-a	HCC-4	HGF
4	CTA CK	Dtk	EGF-R	ENA-78	Fas	FGF-4	FGF-9	GCSF	GITR-Ligand	GITR	GR O	GR O-a	HCC-4	HGF
5	ICA M-1	ICAM-3	IGFBP-3	IGFBP-6	IGF-I SR	IL-1 R4/ST2	IL-1 RI	IL-11	IL-12 p40	IL-12 p70	IL-17	IL-2 Ra	IL-6 R	IL-8
6	ICA M-1	ICAM-3	IGFBP-3	IGFBP-6	IGF-I SR	IL-1 R4/ST2	IL-1 RI	IL-11	IL-12 p40	IL-12 p70	IL-17	IL-2 Ra	IL-6 R	IL-8
7	I-TAC	Lymphotactin	MIF	MIP-1a	MIP-1b	MIP-3b	MS P-a	NT-4	Osteoprotegerin	Oncostatin M	PIGF	sgp1 30	sTNF RII	sTNF -RI
8	I-TAC	Lymphotactin	MIF	MIP-1a	MIP-1b	MIP-3b	MS P-a	NT-4	Osteoprotegerin	Oncostatin M	PIGF	sgp1 30	sTNF RII	sTNF -RI
9	TEC K	TIMP-1	TIMP-2	Thrombospondin	TRAIL R3	TRAIL R4	uPAR	VEGF	VEGF-D	Blank	Blank	Blank	Blank	Blank
10	TEC K	TIMP-1	TIMP-2	Thrombospondin	TRAIL R3	TRAIL R4	uPAR	VEGF	VEGF-D	Blank	Blank	Blank	POS	POS

FIGURE 13-1: UCMC-16 cultured in DMEM/10%FCS, at day 10



FIGURE 13-2: UCMC-15 cultured in PTT-1 medium, day 4






FIGURE 13-3: UCMC-16 cultured in PTT-1 at day 10






FIGURE 13-4: UCMC-16 cultured in PTT-2 at day 10







FIGURE 13-5: UCMC-16 cultured in PTT-3 at day 10




**FIGURE 13-6: Adipose-derived stromal cells cultured in PTT-3 at day 10.
Cells did not grow well in serum free medium**






FIGURE 13-7: Bone marrow-derived stromal cells cultured in PTT-3 (MeE) at day 10. Cells did not grow well in serum free medium




FIGURE 14: Global gene expression in Umbilical Cord Epithelial and Mesenchymal Stem Cells




FIGURE 14: Gene expression present in BOTH, Umbilical Cord Epithelial and Mesenchymal Stem Cells






FIGURE 14: Gene expression present in BOTH, Umbilical Cord Epithelial and Mesenchymal Stem Cells

663	genes in both-protein binding
666	genes in both-cell death
747	genes in both-structural molecule activity
774	genes in both-extracellular
808	genes in both-cell cycle
817	genes in both-protein biosynthesis
825	genes in both-DNA metabolism
899	genes in both-energy pathway
926	genes in both-cytoskeleton organization and biogenesis
975	genes in both-receptor binding
987	genes in both-protein metabolism
1048	genes in both-cell-cell signaling
1138	genes in both-cell differentiation
1317	genes in both-cytosol
1364	genes in both-kinase
1364	genes in both-protein kinase activity
1473	genes in both-host-pathogen interaction
1483	genes in both-protein transport
1637	genes in both-ion channel activity
1667	genes in both-ion transport
1774	genes in both-cell organization, biogenesis
1774	genes in both-organelle organization biogenesis
1968	genes in both-nucleotide binding
2067	genes in both-receptor activity
2067	genes in both-signal transducer activity
2108	genes in both-signal transduction
2125	genes in both-cell growth
2503	genes in both-transcription regulator activity
2906	genes in both-transferase
2947	genes in both-transcription
3435	genes in both-plasma membrane
3436	genes in both-transcription factor activity
3509	genes in both-regulation of gene expression
3509	genes in both-regulation of gene expression, epigenesis
3634	genes in both-DNA binding

**FIGURE 14: Gene
expression present
in BOTH, Umbilical
Cord Epithelial and
Mesenchymal
Stem Cells**





Only in UCMC





FIGURE 14: Gene expression present UNIQUELY in Umbilical Cord Mesenchymal Stem Cells (UCMC)

Figure 15




FIGURE 16: Histology of normal umbilical cord

