

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/86, 15/11, 15/40, 15/51, C07K 14/18 // C12Q 1/68		A1	(11) International Publication Number: WO 97/07224 (43) International Publication Date: 27 February 1997 (27.02.97)
(21) International Application Number: PCT/US96/13198		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 14 August 1996 (14.08.96)		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 60/002,265 14 August 1995 (14.08.95) US 08/580,038 21 December 1995 (21.12.95) US 08/639,857 19 April 1996 (19.04.96) US			
(71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).			
(72) Inventors: SIMONS, John, N.; 738 N. Allegheny Road, Grayslake, IL 60030 (US). DESAI, Suresh, M.; 1408 Amy Lane, Libertyville, IL 60048 (US). MUSHAHWAR, Isa, K.; 18790 Arbor Boulevard, Grayslake, IL 60030 (US).			
(74) Agents: POREMBSKI, Priscilla, E. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).			

(54) Title: REAGENTS AND METHODS USEFUL FOR CONTROLLING THE TRANSLATION OF HEPATITIS GBV PROTEINS

(57) Abstract

Reagents and composition for controlling the translation of hepatitis GB virus (HGBV)-A, -B or -C peptides from viral nucleic acid. These reagents and methods comprise control elements of the 5'NTR region of the HGBV-A, -B, or -C viral genome.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

REAGENTS AND METHODS USEFUL FOR CONTROLLING THE TRANSLATION OF HEPATITIS GBV PROTEINS

Related Applications

5 This application is a continuation-in-part of U.S. Serial No. 08/580,038, filed December 21, 1995, which is a continuation-in-part application of, and claimed the benefit of, U.S. provisional application Serial No. 60/002,265 filed August 14, 1995, which is related to patent application U.S. Serial No. 60/002,255 filed August 14, 1995, which is related to patent applications U.S. 10 Serial No. 08/480,995 filed June 7, 1995, U.S. Serial No. 08/473,475 filed June 7, 1995 and U.S. Serial No. 08/417,629, filed April 6, 1995, which are continuation-in-part applications of U.S. Serial No. 08/424,550 filed June 5, 1995, which is a continuation-in-part application of U.S. Serial No. 08/377,557 filed January 30, 1995, which is a continuation-in-part of U.S. Serial No. 15 08/344,185 filed November 23, 1994 and U.S. Serial No. 08/344,190 filed November 23, 1994, which are each continuation-in-part applications of 08/283,314 filed July 29, 1994, which is a continuation-in-part application of U.S. Serial No. 08/242,654, filed May 13, 1994, which is a continuation-in-part application of U.S. Serial No. 08/196,030 filed February 14, 1994, all of which 20 enjoy common ownership and each of which is incorporated herein by reference.

Background of the Invention

This invention relates generally to the family of hepatitis GB viruses (HGBV) and more particularly, relates to reagents such as antisense nucleic acid sequences and methods utilizing these nucleic acid sequences which are useful for controlling translation of HGBV-A, -B, or -C, both *in vivo* and *in vitro*, by either increasing or decreasing the expressions of HGBV-A, -B or -C proteins.

Recently, a new family of flaviviruses detected in patients with clinically diagnosed hepatitis was reported. This new family of viruses has been named the "GB" viruses, after the initials of the patient first infected with the virus. These viruses have been reported by J. N. Simons et al., Proc. Natl. Acad. Sci. USA 92:3401-3405 (1995); and J. N. Simons et al., Nature Medicine 1(6):564-569 (1995). Three members of the family have been identified to date: GBV-A, GBV-B and GBV-C. T. P. Leary, et al., J. Med. Virol. 48:60-67 (1995) While 35 HGBV-A appears at this time to be of non-human primate source, HGBV-C is

clearly of human source. Currently, the source of HGBV-B is unknown. These viruses are thought to play a role in transmittable hepatitis disease of viral origin.

The GB viruses appear to be members of the *Flaviviridae* family. They possess RNA genomes approximately 9.5 kb in size which contain a single long open reading frame (ORF). Structural and nonstructural proteins are encoded in the N-terminal one-third and C-terminal two-thirds of the putative viral polyproteins, respectively. Phylogenetic analyses of the nonstructural helicase and replicase genes demonstrate that these viruses are related to, but distinct from, the HCV genus of the *Flaviviridae*. See, for example, T. P. Leary, et al., *supra* and 10 A. S. Muerhoff et al., *J. Virol.* 69:5621-5630 (1995). Specifically, GBV-A and GBV-C appear most closely related as they share a common ancestor, while the GBV-A/C ancestor, GBV-B and HCV all appear to be equally divergent from other members of the *Flaviviridae*.

However, when the 5' nontranslated regions (NTRs) and structural genes 15 are examined, a more striking division between the GB viruses and the other members of the *Flaviviridae* becomes apparent. GBV-B appears similar to the HCV and pestivirus genera of the *Flaviviridae*. Conserved sequences present in the 5' NTRs of HCV and pestiviruses are found in the 5' NTR of GBV-B, and GBV-B and HCV share closely related RNA secondary structures within the 5' 20 NTR. (M. Honda, E. A. Brown and S. M. Lemon, manuscript submitted). Moreover, a basic (pI = 11.1) core protein is present at the N-terminus of the GBV-B putative polyprotein precursor, and two putative envelope glycoproteins with several potential N-linked glycosylation sites are located downstream of core in GBV-B. A. S. Muerhoff et al., *supra*. These structural proteins appear in all 25 members of the *Flaviviridae* examined to date. See, for example, M. S. Collett et al., *J. Gen. Virol.* 69:2637-2643 (1989) and R. H. Miller and R. H. Purcell, *Proc. Natl. Acad. Sci. USA* 87:2057-2061 (1990).

In contrast to GBV-B, examination of GBV-A and GBV-C reveals marked 30 differences between these viruses and other genera of the *Flaviviridae*. GBV-A and GBV-C contain long 5' NTRs that have limited sequence identity at the 5'NTR to each other but no identity to the 5' NTRs GBV-B, HCV or pestiviruses. GBV-A and GBV-C also encode putative envelope proteins that contain relatively few potential N-linked glycosylation sites. Most strikingly, clearly discernible basic core proteins are not found in the cDNA sequences cloned thus far from these 35 viruses.

The absence of core proteins would distinguish GBV-A and GBV-C from other genera of the *Flaviviridae*. However, several important aspects of the structure of the GBV-A and GBV-C genomes remain undefined. Primary among these is the identification of the AUG codons at which translation of the viral polyproteins initiate. The sequence of GBV-A contains two potential in-frame initiator AUG codons 27 nucleotides (9 amino acids) and immediately upstream of the putative E1 signal sequence. Similarly, multiple GBV-C sequences possess two to three potential in-frame initiator AUG codons. *See*, T. P. Leary et al., *supra*; and J. Linnen et al., *Science* 271:505-508 (1996). However, none of these AUGs have been demonstrated to serve as the initiator codon, and initiation at any of these sites would result in a severely truncated core protein at best. It is conceivable that deletions during the cloning of these virus RNAs could have resulted in the elimination of core sequences or a disruption of the true ORF in this region of the genome, as suggested by Leary et al., *supra*. However, multiple RT-PCR products generated from the 5' ends of GBV-A and GBV-C using a variety of primers, polymerases and conditions (unpublished data), in addition to determining the 5' end sequences of over 35 separate GBV-C isolates (U.S. Serial No. 08/580,038, filed December 21, 1995, previously incorporated herein by reference) provide no support for the existence of additional sequence missing from the previously described cDNA clones. Thus, it is possible that the 5' ends of these viruses are complete (or nearly complete), and that GBV-A and GBV-C do not encode core proteins.

Of the genera that comprise the *Flaviviridae*, the viruses classified in the flaviviruses genus (e.g., yellow fever virus, dengue virus) contain relatively short 5' NTRs of 97 to 119 nucleotides. In these viruses, translational initiation is thought to utilize a conventional eukaryotic ribosome scanning mechanism in which ribosomes bind the RNA at a 5' cap structure and scan in a 3' direction until encountering an AUG codon in a favorable context for initiation. *See*, M. Kozak, *Cell* 44:283-292 (1989) and M. Kozak, *J. Cell. Biol.* 108:229-241 (1989).

In contrast to the flavivirus genus, genomic RNAs from members of the pestiviruses and HCV genera contain relatively long 5' NTRs of 341 to 385 nucleotides which in some ways are similar to those of picornaviruses. Extensive studies of the picornavirus 5' NTRs reveal that translation initiation occurs through a mechanism of internal ribosome entry. R.J. Jackson et al., *Mol. Biol. Reports* 19:147-159 (1994); K. Meerovitch and N. Sonnenberg, *Semin. Virol.* 67:3798-3807 (1993). This internal entry requires a defined segment of the viral 5' NTR

known as an "internal ribosome entry site" (IRES) or "ribosome landing pad." The RNA comprising the cis-acting IRES forms highly ordered structures which interact with trans-acting cellular translation factors to bind the 40S ribosome subunit at an internal site on the viral message, often many hundreds of nucleotides downstream of the 5' end of the molecule. Such translation initiation functions in a 5' cap-independent fashion, and is generally not influenced by structure or sequence upstream of the IRES.

Practically, the ability of a sequence to function as an IRES is assessed by insertion of the sequence between two cistrons of a bicistronic RNA. If the 10 intercistronic sequence contains an IRES, there is significant translation of the downstream cistron which is generally independent of the translational activity of the upstream cistron. Studies of the 5' NTRs of HCV and pestiviruses using bicistronic mRNAs demonstrate the presence of IRESs in these sequences. *See*, for example, T. L. Poole et al., Virology 206:750-754 (1995); R. Rijnbrand et al., 15 FEBS Letters 365:115-119 (1995); K. Tsukiyama-Kohara et al., J. Virol. 66:1476-1483 (1992); and C. Wang et al., J. Virol. 67:3338-3344 (1993).

Structural changes in the IRES influence the rate of translation initiation. Thus, by modifying a virus's IRES, one can control the amount of viral protein being made. Control of the translation process of the nucleic acids of GB viruses 20 could provide an effective means of treating viral disease. The ability to control translation could result in a decrease of the expression of viral proteins. Also, the ability to increase expression may prove useful by producing greater amounts of GB viral proteins which could be utilized in a variety of ways, both diagnostically and therapeutically. Further, the ability to increase translation of the GB viruses *in* 25 *vivo* may provide a means for increasing immune stimulation in an individual.

It therefore would be advantageous to provide reagents and methods for controlling the translation of HGBV proteins from HGBV nucleic acids. Such reagents would comprise antisense nucleic acid sequences or other compound which may specifically destabilize (or stabilize) the IRES structure. Such nucleic 30 acid sequences or compounds could greatly enhance the ability of the medical community to provide a means for treating an individual infected with GB virus(es). In addition, IRESs are among the most highly conserved nucleotide sequences. Identification of such a sequence immediately suggests a target for probe-based detection reagents. Diagnostic or screening tests developed from 35 these reagents could provide a safer blood and organ supply by helping to eliminate GBV in these blood and organ donations, and could provide a better

understanding of the prevalence of HGBV in the population, epidemiology of the disease caused by HGBV and the prognosis of infected individuals. Additionally, these consereved structures may provide a means for purifying GBV proteins for use in diagnostic assays.

5

Summary of the Invention

The present invention provides unique reagents comprising nucleic acid sequences for HGBV-A, -B or -C that are useful for controlling the translation of HGBV nucleic acids to proteins. These nucleic acid sequences may be DNA or RNA, derivatized DNA or RNA, PNA in both the antisense or sense orientations.

The present invention also provides a method for controlling the translation of HGBV nucleic acids to HGBV proteins, comprising contacting a first nucleic acid sequence with HGBV nucleic acid sequence under conditions which permit hybridization of the first nucleic acid sequence and the HGBV nucleic acid sequence, and altering the level of translation of the HGBV nucleic acid. The first nucleic acid sequence is an antisense nucleic acid sequence which is substantially complementary to a sequence of the sense strand within the 5' NTR region of the HGBV nucleic acid sequence. The sense strand is of genomic or messenger RNA that is subjected to the translation process. The method described herein is performed in an individual infected with HGBV.

The present invention also provides a method of enhancing the translation of a nucleic acid comprising operably linking a nucleic acid with a nucleic acid having a seequence corresponding to the sequence of GBV-A, -B or -C 5' region, to form a combined nucleic acid capable of being translated.

Further, the invention herein provides a composition for enhancing the translation of a nucleic acid, which composition comprises a nucleic acid having a sequence corresponding to the squence of GBV-A, -B, or -C 5' region, for operable linkage to nucleic acid to be translated. Further, a composition for contrlling translation of hepatitis GB virus -A, -B, or -C from GBvirus -A, -B or -C nucleic acids is provided, which comprises a first non-naturally occurring nucleic acid having a sequence complementary to, or capable of being transcribed to form, a nucleic acid having a sequence complementary to, a sequence of the sense strand within the 5'-NTR region of HGBV-A, -B, or -C, wherein said first nucleic acid comprises a sequence selected from the 5' NTR region of GBV-A, -B, or -C, and a cleavage are at which the full length GBV-A, -B, or -C RNA is cleaved to form a subgenomic HGBV-A, -B, or -C RNA. The first nucleic acid

can be a nucleic acid analog, and it can be linked to a cholesteryl moiety at the 3' end.

Brief Description of the Drawings

5 FIGURE 1 presents the alignment of GBV-A and GBV-C 5' sequences and amino acid alignment of their respective ORF's. The putative E1 signal sequence in GBV-C and the Asn-Cys-Cys motif are underlined.

10 FIGURE 2A presents a schematic representation of monocistronic T7 templates, wherein viral RNA sequence is represented as a bold line, the positions of the AUG codons (AUG) and ORFs (box) are indicated;

15 FIGURE 2B shows a PhosphorImager scan of products generated by IVTT reactions programmed with pA15-707/CAT (A15-707, lane 1), pC1-631/CAT (C1-631, lane 2), pHAV-CAT1 (HAV, lane 3), pC631-1/CAT (C631-1, lane 4), SspI-linearized pA15-707/CAT (A15-707-SspI, lane 5) and pC1-631/CAT (C1-631-SspI, lane 6).

FIGURE 3A presents the organization of site-specific mutants of GBV-CAT monocistronic templates;

20 FIGURE 3B shows a PhosphorImager scan of IVTT products generated from GBV-CAT mutant templates, wherein Lanes 1, 4 and 7 are control reactions programmed with pA15-707/CAT, pC1-631/CAT and pHAV-CAT1, respectively; products generated from reactions programmed with the mutant templates are found in lanes 2 (pAmut1/CAT), 3 (pAmut2/CAT), 5 (pCmut1/CAT) and 6 (pCmut2/CAT).

25 FIGURES 4A AND 4B show an Edman degradation of ³H-Leu-labeled GBV-CAT fusion products, wherein IVTT reactions programmed with pA15-707/CAT are presented in 4A and those programmed with pC1-631/CAT are presented in 4B.

30 FIGURES 5A and 5B show the translation of monocistronic RNAs containing 3' GBV deletions. FIGURE 5A presents a schematic of the monocistronic templates and FIGURE 5B presents a PhosphorImager scan of IVTT products generated with pA15-665/CAT (lane 2), pA15-629/CAT (lane 3), pA15-596/CAT (lane 4), pC1-592/CAT (lane 6), pC1-553/CAT (lane 7) and pC1-526/CAT (lane 8). Control reactions are shown in lanes 1 (pA15-707/CAT), 5 (pC1-631/CAT) and 9 (pHAV-CAT1).

35 FIGURES 6A and 6B show the translation of bicistronic GBV and HCV vectors, wherein FIGURE 6A presents a schematic of the bicistronic T7 templates,

and FIGURE 6B presents the luciferase activity (Luc-A, light units X 10⁻³), luciferase protein (Luc-P, band volume X 10⁻³) and protein production of IVTTs programmed with the bicistronic vectors.

FIGURE 7A, B, C and D presents a schematic that depicts a preliminary 5 model of the secondary RNA structures which are present near the 5' end of the GBV-C genome (GenBank accession no. U36380) (SEQUENCE ID NO 3), wherein major putative structural domains are labeled I - V with roman numerals; base pairs which are sites of covariant nucleotide substitutions in different strains of GBV-C are shown in boxes; the putative initiator AUG codon (first in-frame 10 AUG codon which is conserved in all GBV-C sequences) is located between domains IV and V (highlighted bases); (Inset) presents the preliminary model of GBV-A domain V; and covariance between GBV-A and sequences from GBV-A-like viruses found indigenous to tamarins are boxed.

15 **Detailed Description of the Invention**

The present invention provides reagents and methods useful for controlling the translation of HGBV-A, HGBV-B or HGBV-C nucleic acid to protein.

The term "Hepatitis GB Virus" or "HGBV", as used herein, collectively denotes a viral species which causes non-A, non-B, non-C, non-D, non-E 20 hepatitis in man, and attenuated strains or defective interfering particles derived therefrom. This may include acute viral hepatitis transmitted by contaminated foodstuffs, drinking water, and the like; hepatitis due to HGBV transmitted via person to person contact (including sexual transmission, respiratory and parenteral routes) or via intravenous drug use. The methods as described herein will allow 25 the treatment of individuals who have acquired HGBV. Individually, the HGBV isolates are specifically referred to as "HGBV-A", "HGBV-B" and "HGBV-C." As described herein, the HGBV genome is comprised of RNA. Analysis of the nucleotide sequence and deduced amino acid sequence of the HGBV reveals that 30 viruses of this group have a genome organization similar to that of the *Flaviridae* family. Based primarily, but not exclusively, upon similarities in genome organization, the International Committee on the Taxonomy of Viruses has recommended that this family be composed of three genera: Flavivirus, Pestivirus, and the hepatitis C group. Similarity searches at the amino acid level reveal that the hepatitis GB virus subclones have some, albeit low, sequence resemblance to 35 hepatitis C virus. It now has been demonstrated that HGBV-C is not a genotype

of HCV. See, for example, U.S. Serial No. 08/417,629, filed April 6, 1995, previously incorporated herein by reference.

The term "similarity" and/or "identity" are used herein to describe the degree of relatedness between two polynucleotides or polypeptide sequences. The 5 techniques for determining amino acid sequence "similarity" and/or "identity" are well-known in the art and include, for example, directly determining the amino acid sequence and comparing it to the sequences provided herein; determining the nucleotide sequence of the genomic material of the putative HGBV (usually via a cDNA intermediate), and determining the amino acid sequence encoded therein, 10 and comparing the corresponding regions. In general, by "identity" is meant the exact match-up of either the nucleotide sequence of HGBV and that of another strain(s) or the amino acid sequence of HGBV and that of another strain(s) at the appropriate place on each genome. Also, in general, by "similarity" is meant the exact match-up of amino acid sequence of HGBV and that of another strain(s) at 15 the appropriate place, where the amino acids are identical or possess similar chemical and/or physical properties such as charge or hydrophobicity. The programs available in the Wisconsin Sequence Analysis Package, Version 8 (available from the Genetics Computer Group, Madison, Wisconsin, 53711), for example, the GAP program, are capable of calculating both the identity and 20 similarity between two polynucleotide or two polypeptide sequences. Other programs for calculating identity and similarity between two sequences are known in the art.

Additionally, the following parameters are applicable, either alone or in combination, in identifying a strain of HGBV-A, HGBV-B or HGBV-C. It is 25 expected that the overall nucleotide sequence identity of the genomes between HGBV-A, HGBV-B or HGBV-C and a strain of one of these hepatitis GB viruses will be about 45% or greater, since it is now believed that the HGBV strains may be genetically related, preferably about 60% or greater, and more preferably, about 80% or greater.

30 Also, it is expected that the overall sequence identity of the genomes between HGBV-A and a strain of HGBV-A at the amino acid level will be about 35% or greater since it is now believed that the HGBV strains may be genetically related, preferably about 40% or greater, more preferably, about 60% or greater, and even more preferably, about 80% or greater. In addition, there will be 35 corresponding contiguous sequences of at least about 13 nucleotides, which may be provided in combination of more than one contiguous sequence. Also, it is

expected that the overall sequence identity of the genomes between HGBV-B and a strain of HGBV-B at the amino acid level will be about 35% or greater since it is now believed that the HGBV strains may be genetically related, preferably about 40% or greater, more preferably, about 60% or greater, and even more preferably, 5 about 80% or greater. In addition, there will be corresponding contiguous sequences of at least about 13 nucleotides, which may be provided in combination of more than one contiguous sequence. Also, it is expected that the overall sequence identity of the genomes between HGBV-C and a strain of HGBV-C at the amino acid level will be about 35% or greater since it is now believed that the 10 HGBV strains may be genetically related, preferably about 40% or greater, more preferably, about 60% or greater, and even more preferably, about 80% or greater. In addition, there will be corresponding contiguous sequences of at least about 13 nucleotides, which may be provided in combination of more than one contiguous sequence.

15 A polynucleotide "derived from" a designated sequence for example, the HGBV cDNA, or from the HGBV genome, refers to a polynucleotide sequence which is comprised of a sequence of approximately at least about 6 nucleotides, is preferably at least about 8 nucleotides, is more preferably at least about 10-12 nucleotides, and even more preferably is at least about 15-20 nucleotides 20 corresponding, i.e., similar to or complementary to, a region of the designated nucleotide sequence. Preferably, the sequence of the region from which the polynucleotide is derived is similar to or complementary to a sequence which is unique to the HGBV genome. Whether or not a sequence is complementary to or similar to a sequence which is unique to an HGBV genome can be determined by 25 techniques known to those skilled in the art. Comparisons to sequences in databanks, for example, can be used as a method to determine the uniqueness of a designated sequence. Regions from which sequences may be derived include but are not limited to regions encoding specific epitopes, as well as non-translated and/or non-transcribed regions.

30 The derived polynucleotide will not necessarily be derived physically from the nucleotide sequence of HGBV, but may be generated in any manner, including but not limited to chemical synthesis, replication or reverse transcription or transcription, which are based on the information provided by the sequence of bases in the region(s) from which the polynucleotide is derived. In addition, 35 combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be consistent with an intended use.

The term "polynucleotide" as used herein means a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, the term includes double- and single-stranded DNA, as well as double- and single-stranded RNA. It 5 also includes modifications, either by methylation and/or by capping, and unmodified forms of the polynucleotide.

The terms "polynucleotide," "oligomer," "oligonucleotide," "oligo" and "primer" are used interchangeably herein.

"HGBV containing a sequence corresponding to a cDNA" means that the 10 HGBV contains a polynucleotide sequence which is similar to or complementary to a sequence in the designated DNA. The degree of similarity or complementarity to the cDNA will be approximately 50% or greater, will preferably be at least about 70%, and even more preferably will be at least about 90%. The sequence which corresponds will be at least about 70 nucleotides, preferably at least about 80 15 nucleotides, and even more preferably at least about 90 nucleotides in length. The correspondence between the HGBV and the cDNA can be determined by methods known in the art, and include, for example, a direct comparison of the sequenced material with the cDNAs described, or hybridization and digestion with single strand nucleases, followed by size determination of the digested fragments.

"Purified viral polynucleotide" refers to an HGBV genome or fragment thereof which is essentially free, i.e., contains less than about 50%, preferably less than about 70%, and even more preferably, less than about 90% of polypeptides with which the viral polynucleotide is naturally associated. Techniques for purifying viral polynucleotides are well known in the art and include, for example, 20 disruption of the particle with a chaotropic agent, and separation of the polynucleotide(s) and polypeptides by ion-exchange chromatography, affinity chromatography, and sedimentation according to density. Thus, "purified viral 25 polypeptide" means an HGBV polypeptide or fragment thereof which is essentially free, that is, contains less than about 50%, preferably less than about 70%, and even more preferably, less than about 90% of cellular components with which the 30 viral polypeptide is naturally associated. Methods for purifying are known to the routineer.

"Polypeptide" as used herein indicates a molecular chain of amino acids and does not refer to a specific length of the product. Thus, peptides, 35 oligopeptides, and proteins are included within the definition of polypeptide. This term, however, is not intended to refer to post-expression modifications of the

polypeptide, for example, glycosylations, acetylations, phosphorylations and the like.

A "polypeptide" or "amino acid sequence" derived from a designated nucleic acid sequence or from the HGBV genome refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence or a portion thereof wherein the portion consists of at least 3 to 5 amino acids, and more preferably at least 8 to 10 amino acids, and even more preferably 15 to 20 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence.

A "recombinant polypeptide" as used herein means at least a polypeptide of genomic, semisynthetic or synthetic origin which by virtue of its origin or manipulation is not associated with all or a portion of the polypeptide with which it is associated in nature or in the form of a library and/or is linked to a polynucleotide other than that to which it is linked in nature. A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence of HGBV or from an HGBV genome. It also may be generated in any manner, including chemical synthesis or expression of a recombinant expression system, or isolation from mutated HGBV.

The term "synthetic peptide" as used herein means a polymeric form of amino acids of any length, which may be chemically synthesized by methods well-known to the routineer. These synthetic peptides are useful in various applications.

"Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting microorganisms or higher eucaryotic cell lines cultured as unicellular entities refer to cells which can be, or have been, used as recipients for recombinant vector or other transfer DNA, and include the original progeny of the original cell which has been transfected.

As used herein "replicon" means any genetic element, such as a plasmid, a chromosome or a virus, that behaves as an autonomous unit of polynucleotide replication within a cell. That is, it is capable of replication under its own control.

A "vector" is a replicon in which another polynucleotide segment is attached, such as to bring about the replication and/or expression of the attached segment.

The term "control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism.

In prokaryotes, such control sequences generally include promoter, ribosomal binding site and terminators; in eukaryotes, such control sequences generally include promoters, terminators and, in some instances, enhancers. The term "control sequence" thus is intended to include at a minimum all components whose presence is necessary for expression, and also may include additional components whose presence is advantageous, for example, leader sequences.

5 presence is necessary for expression, and also may include additional components whose presence is advantageous, for example, leader sequences.

"Operably linked" refers to a situation wherein the components described are in a relationship permitting them to function in their intended manner. Thus, for example, a control sequence "operably linked" to a coding sequence is ligated 10 in such a manner that expression of the coding sequence is achieved under conditions compatible with the control sequences.

The term "open reading frame" or "ORF" refers to a region of a polynucleotide sequence which encodes a polypeptide; this region may represent a portion of a coding sequence or a total coding sequence.

15 A "coding sequence" is a polynucleotide sequence which is transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5' -terminus and a translation stop codon at the 3' -terminus. A coding sequence can include, but is not limited to, 20 mRNA, cDNA, and recombinant polynucleotide sequences.

The term "immunologically identifiable with/as" refers to the presence of epitope(s) and polypeptide(s) which also are present in and are unique to the designated polypeptide(s), usually HGBV proteins. Immunological identity may be determined by antibody binding and/or competition in binding. These 25 techniques are known to the routineer and also are described herein. The uniqueness of an epitope also can be determined by computer searches of known data banks, such as GenBank, for the polynucleotide sequences which encode the epitope, and by amino acid sequence comparisons with other known proteins.

As used herein, "epitope" means an antigenic determinant of a polypeptide. 30 Conceivably, an epitope can comprise three amino acids in a spatial conformation which is unique to the epitope. Generally, an epitope consists of at least five such amino acids, and more usually, it consists of at least eight to ten amino acids. Methods of examining spatial conformation are known in the art and include, for example, x-ray crystallography and two-dimensional nuclear magnetic resonance.

35 The term "individual" as used herein refers to vertebrates, particularly members of the mammalian species and includes but is not limited to domestic

animals, sports animals, primates and humans; more particularly the term refers to tamarins and humans.

A polypeptide is "immunologically reactive" with an antibody when it binds to an antibody due to antibody recognition of a specific epitope contained within the polypeptide. Immunological reactivity may be determined by antibody binding, more particularly by the kinetics of antibody binding, and/or by competition in binding using as competitor(s) a known polypeptide(s) containing an epitope against which the antibody is directed. The methods for determining whether a polypeptide is immunologically reactive with an antibody are known in the art.

As used herein, the term "immunogenic polypeptide containing an HGBV epitope" means naturally occurring HGBV polypeptides or fragments thereof, as well as polypeptides prepared by other means, for example, chemical synthesis or the expression of the polypeptide in a recombinant organism.

The term "transformation" refers to the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion. For example, direct uptake, transduction, or f-mating are included. The exogenous polynucleotide may be maintained as a non-integrated vector, for example, a plasmid, or alternatively, may be integrated into the host genome.

"Treatment" refers to prophylaxis and/or therapy.

The term "plus strand" (or "+") as used herein denotes a nucleic acid that contains the sequence that encodes the polypeptide. The term "minus strand" (or "-") denotes a nucleic acid that contains a sequence that is complementary to that of the "plus" strand.

"Positive stranded genome" of a virus denotes that the genome, whether RNA or DNA, is single-stranded and encodes a viral polypeptide(s).

The term "test sample" refers to a component of an individual's body which is the source of the analyte (such as, antibodies of interest or antigens of interest). These components are well known in the art. These test samples include biological samples which can be tested by the methods of the present invention described herein and include human and animal body fluids such as whole blood, serum, plasma, cerebrospinal fluid, urine, lymph fluids, and various external secretions of the respiratory, intestinal and genitorurinary tracts, tears, saliva, milk, white blood cells, myelomas and the like; biological fluids such as cell culture supernatants; fixed tissue specimens; and fixed cell specimens.

"Purified HGBV" refers to a preparation of HGBV which has been isolated from the cellular constituents with which the virus is normally associated, and from other types of viruses which may be present in the infected tissue. The techniques for isolating viruses are known to those skilled in the art and include, 5 for example, centrifugation and affinity chromatography.

"PNA" denotes a "peptide nucleic analog" which may be utilized in various diagnostic, molecular or therapeutic methods. PNAs are neutrally charged moieties which can be directed against RNA or DNA targets. PNA probes used in assays in place of, for example, DNA probes, offer advantages not achievable 10 when DNA probes are used. These advantages include manufacturability, large scale labeling, reproducibility, stability, insensitivity to changes in ionic strength and resistance to enzymatic degradation which is present in methods utilizing DNA or RNA. These PNAs can be labeled with such signal generating compounds as fluorescein, radionucleotides, chemiluminescent compounds, and the like. PNAs 15 or other nucleic acid analogs such as morpholino compounds thus can be used in various methods in place of DNA or RNA. It is within the scope of the routineer that PNAs or morpholino compounds can be substituted for RNA or DNA with appropriate changes if and as needed in reagents and conditions utilized in these methods.

20 The detection of HGBV in test samples can be enhanced by the use of DNA hybridization assays which utilize DNA oligomers as hybridization probes. Since the amount of DNA target nucleotides present in a test sample may be in minute amounts, target DNA usually is amplified and then detected. Methods for amplifying and detecting a target nucleic acid sequence that may be present in a test 25 sample are well-known in the art. Such methods include the polymerase chain reaction (PCR) described in U.S. Patents 4,683,195 and 4,683,202 which are incorporated herein by reference, the ligase chain reaction (LCR) described in EP-A-320 308, gap LCR (GLCR) described in European Patent Application EP-A-439 182 and U.S. Patent No. 5,427,930 which are incorporated herein by reference, 30 multiplex LCR described in International Patent Application No. WO 93/20227, NASBA and the like. These methods have found widespread application in the medical diagnostic field as well as in the fields of genetics, molecular biology and biochemistry.

35 The reagents and methods of the present invention are made possible by the provision of a family of closely related nucleotide sequences present in the plasma, serum or liver homogenate of an HGBV infected individual, either tamarin or

human. This family of nucleotide sequences is not of human or tamarin origin, since it hybridizes to neither human nor tamarin genomic DNA from uninfected individuals, since nucleotides of this family of sequences are present only in liver (or liver homogenates), plasma or serum of individuals infected with HGBV. In 5 addition, the family of sequences has shown no significant identity at the nucleic acid level to sequences contained within the HAV, HBV, HCV, HDV and HEV genome, and low level identity, considered not significant, as translation products. Infectious sera, plasma or liver homogenates from HGBV infected humans contain these polynucleotide sequences, whereas sera, plasma or liver homogenates from 10 non-infected humans has not contained these sequences. Northern blot analysis of infected liver with some of these polynucleotide sequences has demonstrated that they are derived from a large RNA transcript similar in size to a viral genome. Sera, plasma or liver homogenates from HGBV-infected humans contain 15 antibodies which bind to this polypeptide, whereas sera, plasma or liver homogenates from non-infected humans do not contain antibodies to this polypeptide; these antibodies are induced in individuals following acute non-A, non-B, non-C, non-D and non-E hepatitis infection. By these criteria, it is believed that the sequence is a viral sequence, wherein the virus causes or is associated with non-A, non-B, non-C, non-D and non-E hepatitis.

20 Using determined portions of the isolated HGBV nucleic acid sequences as a basis, oligomers of approximately eight nucleotides or more can be prepared, either by excision or synthetically, which hybridize with the HGBV genome and are useful in identification of the viral agent(s), further characterization of the viral genome, as well as in detection of the virus(es) in diseased individuals. The 25 natural or derived probes for HGBV polynucleotides are a length which allows the detection of unique viral sequences by hybridization. While six to eight nucleotides may be a workable length, sequences of ten to twelve nucleotides are preferred, and those of about 20 nucleotides may be most preferred. These sequences preferably will derive from regions which lack heterogeneity. These 30 probes can be prepared using routine, standard methods including automated oligonucleotide synthetic methods. A complement of any unique portion of the HGBV genome will be satisfactory. Complete complementarity is desirable for use as probes, although it may be unnecessary as the length of the fragment is increased.

35 Synthetic oligonucleotides may be prepared using an automated oligonucleotide synthesizer such as that described by Warner, DNA 3:401 (1984).

If desired, the synthetic strands may be labeled with ^{32}P by treatment with polynucleotide kinase in the presence of ^{32}P -ATP, using standard conditions for the reaction. DNA sequences including those isolated from genomic or cDNA libraries, may be modified by known methods which include site directed

5 mutagenesis as described by Zoller, *Nucleic Acids Res.* 10:6487 (1982). Briefly, the DNA to be modified is packaged into phage as a single stranded sequence, and converted to a double stranded DNA with DNA polymerase using, as a primer, a synthetic oligonucleotide complementary to the portion of the DNA to be modified, and having the desired modification included in its own sequence. Culture of the

10 transformed bacteria, which contain replications of each strand of the phage, are plated in agar to obtain plaques. Theoretically, 50% of the new plaques contain phage having the mutated sequence, and the remaining 50% have the original sequence. Replicates of the plaques are hybridized to labeled synthetic probe at temperatures and conditions suitable for hybridization with the correct strand, but

15 not with the unmodified sequence. The sequences which have been identified by hybridization are recovered and cloned.

Polymerase chain reaction (PCR) and ligase chain reaction (LCR) are techniques for amplifying any desired nucleic acid sequence (target) contained in a nucleic acid or mixture thereof. In PCR, a pair of primers are employed in excess

20 to hybridize at the outside ends of complementary strands of the target nucleic acid. The primers are each extended by a polymerase using the target nucleic acid as a template. The extension products become target sequences themselves, following dissociation from the original target strand. New primers are then hybridized and extended by a polymerase, and the cycle is repeated to geometrically increase the

25 number of target sequence molecules. PCR is disclosed in U.S. patents 4,683,195 and 4,683,20, previously incorporated herein by reference.

LCR is an alternate mechanism for target amplification. In LCR, two sense (first and second) probes and two antisense (third and fourth) probes are employed in excess over the target. The first probe hybridizes to a first segment of the target

30 strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being positioned so that the primary probes can be ligated into a fused product. Further, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar ligatable fashion. If the target is initially double

35 stranded, the secondary probes will also hybridize to the target complement in the first instance. Once the fused strand of sense and antisense probes are separated

from the target strand, it will hybridize with the third and fourth probes which can be ligated to form a complementary, secondary fused product. The fused products are functionally equivalent to either the target or its complement. By repeated cycles of hybridization and ligation, amplification of the target sequence is 5 achieved. This technique is described in EP-A-320,308, hereby incorporated by reference. Other aspects of LCR technique are disclosed in EP-A-439,182, which is incorporated herein by reference.

The 5'-NTR region of HGBV-A is approximately 592 nucleotides long (SEQUENCE ID NOs 23). This region in HGBV-B is approximately 445 10 nucleotides long (SEQUENCE ID NO 32), and the 5'NTR region of HGBV-C is approximately 533 nucleotides in length (SEQUENCE ID NO 4). To functionally characterize the 5' ends of GBV-A and GBV-C RNAs, the sites and mechanism of translation initiation of both monocistronic and bicistronic RNAs were examined in a cell-free *in vitro* translation system. Weak IRES elements were found to be 15 present in the 5' RNAs of GBV-A and GBV-C suggesting that these sequences are complete or nearly complete. In addition, the position of the initiating AUG codons in the monocistronic RNAs, and presumably in the viral genomic RNA as well, demonstrated that GBV-A and GBV-C do not contain core proteins at the N-termini of their polyproteins. Thus, GBV-A and GBV-C appear unique from other 20 members of the *Flaviviridae* and may constitute a separate group within this family. Consistent with this hypothesis, we also discovered that the secondary structures of the 5' ends of these viruses are different from the conserved structures present in the 5' NTRs of the pestiviruses, HCV and GBV-B.

The present invention provides nucleic acids that are capable of interacting 25 with distinct cis-acting control elements of HGBV and thus are capable of blocking, enhancing or suppressing the translation of HGBV nucleic acids.

In a first embodiment, a method for controlling the translation of HGBV nucleic acids to proteins is provided. This method comprises the steps of contacting a first *non-naturally occurring* nucleic acid with HGBV nucleic acid. 30 This first nucleic acid has a sequence that is complementary to a sequence of the sense strand within the 5'NTR region of HGBV-A, -B or -C. This first nucleic acid is contacted with an HGBV nucleic acid for times and under conditions suitable for hybridization to occur, and thus form a hybridization product. The hybridization results in the alteration of the level of translation of the HGBV 35 nucleic acid.

The antisense nucleic acid of the present invention is RNA, DNA or a modified nucleic acid such as a PNA or morpholino compound, degradation-resistant sulfurized and thiophosphate derivatives of nucleic acids, and the like. Modified nucleic acids preferably will be able to increase the intracellular stability 5 and/or permeability of the nucleic acid, increase the affinity of the nucleic acid for the sense strand or decrease the toxicity of the nucleic acid. Such advantages are well known in the art, and are described in, for example, S. T. Crooke et al., eds., Antisense Research and Applications, CRC Press (1993).

Antisense nucleic acids thus can be modified or altered to contain modified 10 bases, sugars or linkages, be delivered in specialized systems such as liposomes or by gene therapy, or may have attached moieties. Such attached moieties, such as hydrophobic moieties such as lipids and in particular, cholesterols, can enhance the interaction of the nucleic acid with cell membranes. In addition, such attached moieties can act as charge neutralizers of the phosphate backbone (for example, 15 polycationic moieties such as polylysine). These moieties can be attached at either the 5' or the 3' end of the nucleic acids, and also can be attached through a base, sugar or internucleotide linkage. Other moieties can act as capping groups which are specifically placed at the 3' or the 5' ends of the nucleic acids to prevent exonuclease degradation. These capping groups include, for example, hydroxyl 20 protecting groups including glycols such as polyethylene glycols (PEG), tetraethylene glycol (TEG) and the like.

The first nucleic acid will have at least 10 nucleotides in a sequence substantially complementary to a sequence of the sense strand within the 5'NTR 25 region of HGBV-A, -B, or -C. Preferably, the first nucleic acid has about 12 nucleotides in such a complementary sequence; more preferably, the first nucleic acid has about 15 nucleotides; and still more preferably, the first nucleic acid has about 20 nucleotides. It is preferred that such a first nucleic acid have less than 100 nucleotides in such a complementary sequence, and more preferably, a first nucleic acid will have less than 50 nucleotides. Most preferably, the first nucleic 30 acid will have between 20 to 30 nucleotides that are capable of forming a stable hybridization product with a sense sequence of the 5'NTR region of HGBV-A, -B or -C.

The 5'NTR region of HGBV-A is set forth in SEQUENCE ID NO 23; the 35 5'NTR region of HGBV-B is set forth in SEQUENCE ID NO 32; and the 5'NTR region of HGBV-C is set forth in SEQUENCE ID NO 4. The nucleic acid can be placed in the cell through several ways known to those in the art. For example,

cells can be transfected with a second nucleic acid capable of generating the first nucleic acid as a transcription product (for example, by including the second nucleic acid in a viral carrier as detailed by U.S. Patent 4,493,002, incorporated herein by reference, or by gene therapy methods such as including the second nucleic acid in a retroviral vector). Gene therapy methods are known to those of skill in the art.

The present invention further encompasses means for placing the first nucleic acid or the second nucleic acid into cells infected with HGBV-A, -B or -C or into cells which are to be protected from HGBV infection. Examples of such means include but are not limited to vectors, liposomes and lipid suspensions, such as N-(1-(2,3-dioleyloxy)propyl)-N, N, N-trimethylammonium methylsulfate (DOTAP), N-[1-(2,3-dioleyloxy)propyl]-N, N, N-trimethylammonium chloride (DOTMA), and the like. The lipid may be covalently linked directly to the first nucleic acid in an alternative embodiment.

The antisense nucleic acid also may be linked to moieties that increase cellular uptake of the nucleic acid. Such moieties may be hydrophobic (such as, phospholipids or lipids such as steroids [for example, cholesterol]) or may be polycationic moieties that are attached at any point to the antisense nucleic acid, including at the 5' or 3' ends, base, sugar hydroxyls and internucleoside linkages. A moiety known to increase uptake is a cholestryl group, which may be attached through an activated cholestryl chloroformate or cholic acid, by means known in the art.

Further, enhancement of translation may allow for stronger immune responses. Blocking or decreasing translation of viral nucleic acid may decrease the pathology of the viral infection.

Nucleic acid or nucleic acid analogs can be provided as compositions for pharmaceutical administration. Injection preparations and suppositories may usually contain 1-10 mg of the nucleic acid or nucleic acid analog per dose (ampule or capsule). For humans the daily dose of about 0.1 to 1000 mg, preferably 1-100 mg (from about 10-20 mg/kg to 1000 to 2000 mg/kg body weight) is the daily dosage. As is known to those in the art, however, a particular dose for a particular individual depends on a variety of factors, including but not limited to, effectiveness of the particular nucleic acid or nucleic acid analog used, the age, weight and general state of health of the individual, the diet and sex of the individual, the mode of administration of the dosage, the rate of elimination and half life of the composition, whether this composition is used in combination with

other medications and the clinical severity of the individual's disease. Such compositions which are pharmaceutical articles of manufacture include articles whose active ingredients are contained in an effective amount of attain the intended purpose. A preferred range has been described hereinabove, and determination of 5 the most effective amounts for treatment of each HGBV infection is well within the skill of the rountineer.

In addition to the nucleic acid and nucleic acid analogs of the present invention, contemplated pharmaceutical preparations may contain suitable excipients and auxiliaries which facilitate processing of the active compounds. 10 These preparations can be administered orally, rectally, parenterally, buccally or sublingually. All may contain from 0.1 to 99% by weight of active ingredients, together with an excipient. A preferred method of administration is parenteral, especially intravenous administration.

15 Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble or water-dispersible form. Additionally, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils (for example, sesame oil or synthetic fatty acid esters such as ethyoleate or triglycerides). Aqueous injection suspensions may contain substances which 20 increase the viscosity of the suspension, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. The suspension also may contain stabilizers.

25 It is within the scope of the present invention that the compositions described herein may be administered encapsulated in liposomes, pharmaceutical compositions wherein the active ingredient is contained either dispersed or variously present in corpuscles consisting of aqueous concentric layers adherent to lipidic layers. Methods of utilizing this technology are known in the art.

The present invention will now be described by way of examples, which are meant to illustrate, but not to limit, the spirit and scope of the invention.

30

EXAMPLES

Example 1. Internal ribosome entry site in 5' NTR of GBV-B

Several positive strand RNA viruses, such as picornaviruses and 35 pestiviruses, possess large 5' nontranslated regions (NTRs). These large NTRs control the initiation of cap-independent translation by functioning as internal ribosome entry sites (IRESs) (Pelletier and Sonenberg, Nature (London) 334:320-

325). The IRES is thought to form a specific RNA structure which allows ribosomes to enter and begin translation of an RNA without using the cellular machinery required for cap-dependent translation initiation. The large 5' NTR of HCV has been shown to possess an IRES (Tsukiyama-Kohara et al. *J. Virol.* 66:1476-1483, 1992; Wang *et al.* *J. Virol.* 67:3338-3344, 1993; Rijnbrand et al. *FEBS Letters* 365:115-119, 1995). Due to the high level of sequence conservation between the 5' NTRs of GBV-B and HCV, it was reasoned that GBV-B may also contain an IRES.

5 To test for IRES function in GBV-B (SEQUENCE ID NO 32), the 5' NTR of this virus was used to replace the 5' NTR of hepatitis A virus (HAV) in the pLUC-HAV-CAT plasmid described by Whetter *et al.* (*J. Virol.* 68:5253-5263, 1994). The 5' NTR of GBV-B was amplified from a plasmid clone using 10 SEQUENCE ID NO. 58 (UTR-B.1) and SEQUENCE ID. NO. 59 (NTR-B-a1) as primers. Briefly, a 50 μ l PCR was set up using a Perkin-Elmer PCR kit as 15 described by the manufacturer with 1 μ M primers, 2 mM MgCl₂ and approximately 10 ng of plasmid. This reaction was amplified for 20 cycles (94°C, 20 sec; 55°C, 30 sec; 72°C, 30 sec) followed by a final extension at 72°C for 10 min. The completed reaction then was held at 4°C. This product was extracted with phenol:chloroform and precipitated as described in the art. The 3' terminal 20 adenosine residues added by the AmpliTaq[®] polymerase were removed from this product by incubation with T4 DNA polymerase and deoxynucleotide triphosphates as described (Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 1989). After heat inactivation, the product 25 was digested with Xba I and gel purified as described in the art. The purified product was ligated to pHAV-CAT1 (Whetter *et al.* *J. Virol.* 68:5253-5263, 1994) that had been cut with HindIII, end-filled with Klenow polymerase and deoxynucleotide triphosphates, heat-inactivated, digested with Xba I, treated with bacterial alkaline phosphatase, extracted with phenol:chloroform, and precipitated as described in the art. The constructed plasmid, pGBB-CAT1, was digested with 30 Sac I, blunt-ended with T4 DNA polymerase and deoxynucleotide triphosphates, heat-inactivated, and digested with Not I as described in the art. The 1.3 kbp product from these reactions was gel purified and cloned into pLUC-HAV-CAT (Whetter *et al.* *J. Virol.* 68:5253-5263, 1994) that had been digested with HindIII, end-filled with Klenow polymerase and deoxynucleotide triphosphates, heat- 35 inactivated, digested with Not I, treated with bacterial alkaline phosphatase, extracted with phenol:chloroform, and precipitated as described in the art. The

resultant plasmid, pLUC-GBB-CAT was used in *in vitro* transcription-translation experiments to test for an IRES function.

An *in vitro* transcription-translation assay was performed using the TNTTM T7 coupled reticulocyte lysate system from Promega (Madison, WI) as described by the manufacturer. The plasmids tested were pLUC-GBB-CAT (described above), pLUC-HAV-CAT (positive control from Whetter *et al.* *J. Virol.* 68:5253-5263, 1994), and pLUC-Δ355-532 (negative control from Whetter *et al.* *J. Virol.* 68:5253-5263, 1994). The products (labeled with ³⁵S-methionine) were run on a 10% Laemmli gel as described in the art. The gel was fixed in 10% methanol, 20% acetic acid for 10 minutes, dried down and exposed to a PhosphoImager[®] screen (Molecular Dynamics, Sunnyvale, CA). The products were visualized with the PhosphoImager[®]. In addition, the reactions were examined for Luc and CAT activity using commercially available kits (Promega, Madison, WI)(data not shown).

All three reactions contained luciferase activity and a band consistent with the size expected for luciferase (transcribed from the LUC gene in the plasmid). LUC expression, which is a measure of the level of translation that initiates from the 5' end of the mRNA, appeared to be equivalent in the three reactions. Thus, equivalent amounts of RNA templates were present in a translatable form in these three reactions. The pLUC-HAV-CAT and the pLUC-GBB-CAT reactions also possessed chloramphenicol acetyltransferase (CAT) activity and contained a band consistent with the size expected for CAT (from the the CAT gene in the plasmid). This band is not seen in the pLUC-Δ355-532 negative control. CAT expression measures the level of internal translation initiation. Because translation of the CAT gene requires the existence of an IRES in this plasmid construct, the 5' NTR of GBV-B must be providing this function. Therefore, similar to HCV, GBV-B's 5'NTR contains an IRES. Further studies of these plasmids, both *in vitro* and *in vivo* are ongoing to better characterize the IRES in GBV-B.

30 Example 2. Internal ribosome entry site in 5' NTR of GBV-A and -C

A. Plasmids. Various monocistronic and bicistronic plasmids were constructed with PCR-amplified sequences of GBV-A and GBV-C. PCRs utilized components of the GeneAmp PCR Kit with AmpliTaq (Perkin-Elmer) as directed by the manufacturer with final reaction concentrations of 1 μM for oligonucleotide primers and 2 mM MgCl₂. PCR products were digested with restriction

endonucleases, gel purified and cloned using standard procedures as described by J. Sambrook et al., Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor (1989). Monocistronic fusions between GBV sequences and bacterial chloramphenicol acetyltransferase (CAT) were generated by replacing the hepatitis A virus (HAV) HindIII/XbaI fragment of pHAV-CAT1 (described by L. E. Whetter et al., J. Virology 68:5253-5263 (1994) with PCR-amplified cDNA from the 5' ends of GBV-A and GBV-C. The bicistronic constructs were generated in pT7/CAT/ICS/Luc, described by D. Macejak et al., in M. A. Brinton et al., eds., New Aspects of Positive-Strand RNA Viruses, American Society for Microbiology, Washington, D. C., 1990, p. 152-157, and provided as a gift by P. Sarnow, in a two step procedure. First, monocistronic fusions between GBV and luciferase (Luc) were constructed by inserting GBV sequences into the HindIII/NcoI-cut pT7/CAT/ICS/Luc.

Bicistronic vectors were constructed by cloning the HindIII/blunt/SacI GBV fragment from these monocistronic vectors into pT7/CAT/ICS/Luc which had been digested with SalI (blunt) and SacI. The sequence of the cloned inserts and ligation junctions were confirmed by dsDNA sequencing (Sequenase 2.0, USB, Cleveland). Nomenclature (e.g. A15-707) describes the source (GBV-A) and range (nts 15 to 707) of sequence incorporated into the various vectors.

GBV-A sequences (GenBank accession no. U22303) were amplified from a plasmid clone. PCRs for the GBV-A monocistronic and bicistronic constructs utilized the sense primer 5'-TATAATAAAGCTTGCCCCGGACCTCCCACCGAG-3' (HindIII site underlined) (SEQUENCE ID NO 5) coupled with 5'-GCTCTAGATCGGGAACAACAATTGGAAAG (SEQUENCE ID NO 6), 5'-GCTCTAGAGCACTGGTGCCGCGAGT (SEQUENCE ID NO 11), 5'-GCTCTAGAGAGGGGAAGCAAACCA (SEQUENCE ID NO 12) and 5'-GCTCTAGACATGGTGAATGTGTCGACCA (Xba I sites underlined) (SEQUENCE ID NO 13) for the monocistronic vectors pA15-707/CAT, pA15-665/CAT, pA15-629/CAT and pA15-596/CAT, respectively; and 5' CCATAATCATGAGGGAAACAACAATTGGAAAG (SEQUENCE ID NO 17), 5'-CCATAATCATGAGCCGCGAGTTGAAGAGCAC (SEQUENCE ID NO 24), and 5' GCCAAGCCATGGTGAATGTG 3' (BspHI or NcoI sites underlined) (SEQUENCE ID NO 25) for the bicistronic vectors pCAT/A15-705/Luc, pCAT/A15-657/Luc and pCAT/A15-596/Luc, respectively. In addition, a GBV-A sequence amplified with 5'-TATAATAAAGCTTGCCGCGAGTTGAAGAGCAC (SEQUENCE ID NO 21) and 5'-

CCATAATCATGAGCCCCGGACCTCCCACCGAG (SEQUENCE ID NO 22)

were used to construct pCAT/A657-15/Luc which contain GBV-A sequences in the antisense orientation.

GBV-C sequences were amplified from a plasmid generated during the 5 cloning of GBV-C 5' sequences, as described in U.S. Serial No. U.S. Serial No. 08/580,038, previously incorporated herein by reference. The sequence of this GBV-C cDNA (nts 1 to 631, SEQUENCE ID NO 4) corresponds to nts 30 to 659 of GenBank accession no. U44402, the longest GBV-C isolate reported to date and nts 13 to 643 of SEQUENCE ID NO. 3. PCRs for the GBV-C monocistronic 10 and bicistronic plasmids utilized the sense primer 5'-

TATAATAAAGCTTCACTGGGTGCAAGCCCCA (HindIII site underlined)

(SEQUENCE ID NO 7) coupled with 5'-

GCTCTAGAGCGCAACAGTTGTGAGGAA (SEQUENCE ID NO 8), 5'-

GCTCTAGAACAAGCGTGGGTGGCCGGGG (SEQUENCE ID NO 14), 5'-

15 GCTCTAGAGACCGACGAGAAGGAGCAGAAG (SEQUENCE ID NO 15) and 5'-GCTCTAGACATGATGGTATAGAAAAGAG (Xba I site underlined) (SEQUENCE ID NO 16) for the monocistronic vectors pC1-631/CAT, pC1-592/CAT, pC1-553/CAT and pC1-526/CAT, respectively; and 5'-

CATGCCATGGCGCAACAGTTGTGAGGAA (SEQUENCE ID NO 18), 5'-

20 GTATTGCGCCATGGCTCGACAAGCGTGGGTGGCCGGGG (SEQUENCE ID NO 26), and 5'-GGACTGCCATGGTGGTATAGAAAAGAG (NcoI sites underlined) (SEQUENCE ID NO 27) for the bicistronic vectors pCAT/C1-629/Luc, pCAT/C1-596/Luc and pCAT/C1-526/Luc, respectively. Additional GBV-C sequences were amplified with 5'-

25 GCTCTAGACACTGGGTGCAAGCCCCA (XbaI site underlined) (SEQUENCE ID NO 9) and 5'-TATAATAAGCTTGGCGCAACAGTTGTGAG (HindIII site underlined) (SEQUENCE ID NO 10) for the monocistronic pC631-1/CAT plasmid, and 5'-TATAATAAGCTTCGACAAGCGTGGGTGGCCGGGG 3' (HindIII site underlined) (SEQUENCE ID NO 28) and 5'-

30 GTATTGCGCCATGGCACTGGGTGCAAGCCCCAGAA (NcoI site underlined) (SEQUENCE ID NO 29) for the bicistronic pCAT/C596-1/Luc plasmid. Both of these plasmids contain GBV-C sequences in the antisense orientation.

HCV sequences were amplified from a plasmid clone of a genotype 1a isolate using the sense primer 5'-

35 TATAATAAGCTTCACTCCCCTGTGAGGAACTAC (HindIII site underlined) (SEQUENCE ID NO 19) coupled with 5'-

GTATTGCGTCATGATGGTTTTCTTGGGTTAG (SEQUENCE ID NO 20) or 5'-CCATAATCATGATGCACGGTCTACGAGACCT (BspHI sites underlined) (SEQUENCE ID NO 30) to generate the bicistronic vectors pCAT/HCV39-377/Luc and pCAT/HCV39-345/Luc, respectively.

5 Site-specific nucleotide changes were generated in pA15-707/CAT and pC1-631/CAT using the MORPH™ site-specific plasmid DNA mutagenesis kit (5 Prime --> 3 Prime, Inc., Boulder, CO) as directed by the manufacturer.

Nucleotide changes were confirmed by dsDNA sequencing as described above.

B. In vitro transcription/translation. *In vitro* transcription/translation (IVTT)

10 reactions were performed with the TNT™ T7 Coupled Reticulocyte Lysate System (Promega) according to manufacturer's instructions. Reactions (25 µl) contained 20 units rRNasin (Promega), 20 µCi ³⁵S-cysteine (1000 Ci/mmol, Amersham), and 0.5 µg of plasmid template. After incubation at 30°C for 60 minutes, 5 µl aliquots were denatured (5 minutes, 99°C) in an equal volume of 2X SDS/PAGE 15 loading buffer (125 mM Tris, pH 6.8, 4% SDS, 20% glycerol, 10% 2-mercaptoethanol and 0.2 mg/ml bromophenol blue) and electrophoretically separated on 10 to 20% SDS-polyacrylamide gels (Bio-Rad). The gels were fixed in 10% methanol, 20% acetic acid, dried and analyzed with a PhosphorImager SI™ using ImageQuaNT™ software (Molecular Dynamics, Inc.). Image exposure 20 time, white-black range and product quantitations are presented hereinbelow corresponding figure descriptions.

C. Reporter gene enzymatic assays. Luciferase assays were performed by mixing 50 µl of 1X Luciferase Assay Reagent (Promega) with 1 µl of a 10-fold dilution of a rabbit reticulocyte lysate reaction. Activity was assayed immediately by a 5 25 second count in a Clinilumat LB9502 Luminometer (Berthold Systems Inc., Pittsburgh). CAT assays were completed with a commercially available kit (Promega) according to manufacturer's instructions. Briefly, 5 µl of lysate was incubated with [³H]chloramphenicol and n-butyryl CoA in a 125 µl reaction for one hour at 37 °C. Butyrylated [³H]chloramphenicol products were isolated by 30 xylene extraction and quantitated by liquid scintillation counting.

D. Secondary RNA structure. A model of the secondary structure of the 5' nontranslated RNA of the GBV-C genome was constructed using a combination of phylogenetic and thermodynamic approaches. A first level phylogenetic analysis considered nucleotide sequences representing the 5' RNA of GBV-C strains 35 present in 35 different patient sera, as presented in U.S. Serial No. 08/580,038, filed December 21, 1995, previously incorporated herein by reference. These

were aligned with the program PILEUP (Wisconsin Sequence Analysis Package, version 8, September 1994; Genetics Computer Group, Madison, Wisconsin) and subjected to a manual search for covariant nucleotide substitutions indicative of conserved helical structures. In addition to canonical Watson-Crick base pairs, G-
5 U base pairs were considered acceptable for this analysis. Conserved helical structures identified by the presence of one or more covariant nucleotide substitutions were forced to base pair in the subsequent computer-based folding of the prototype GBV-C sequence (GenBank accession no. U36380) (SEQUENCE ID NO 3) which used the program MFOLD. Separate MFOLD analyses were
10 carried out with sequences representing nts 1-611, 43-522 (both closed at 273-418), 273-418, and 43-180 of SEQUENCE ID NO 3. MFOLD predicts a series of alternative structures with different predicted folding energies. These were reviewed to determine which predicted structures were most permissive for covariant and noncovariant nucleotide substitutions present in the other GBV-C
15 sequences. Where no predicted structure could accommodate most nucleotide substitutions, the sequence was left single stranded in the final model. A second level phylogenetic analysis involved the alignment of GBV-C sequences with the 5' RNA sequences of 5 separate GBV-A strains (as described in G. G. Schlauder et al., *Lancet* 346:447 [1995] and J. N. Simons et al., *Proc Natl. Acad. Sci. USA*
20 92:3401-3405 [1995]), followed by a manual search for covariant substitutions indicative of similar structures in the 5' sequences of these related viruses.

E. Results.

1. Translation of monocistronic transcripts containing 5' GBV RNA. A common Asn-Cys-Cys motif homologous to the HCV E1 Asn-Ser-Cys motif is
25 found near the N-termini of the putative E1 proteins of GBV-A, GBV-B and GBV-C (T. P. Leary et al., *supra* and FIGURE 1). Located near the N-termini of the GBV-A and GBV-C large ORFs, this tripeptide sequence appears to be the 5' most conserved motif between HCV and the GB viruses. Because it is within the coding regions of GBV-B and HCV and in-frame with the long ORF, this
30 sequence was believed likely to be translated in GBV-A and GBV-C as well. To determine whether the 5' ends of GBV-A and -C could direct translation, nts 15 to 707 of GBV-A (SEQUENCE ID NO 23) and nts 1 to 631 of GBV-C (SEQUENCE ID NO 4) were cloned into plasmid vectors to create pA15-707/CAT and pC1-631/CAT, respectively. These vectors contained a T7 promoter driving
35 transcription of the 5' GBV sequences, which were ligated in-frame (relative to the Asn-Cys-Cys motif) with the bacterial chloramphenicol acetyltransferase (CAT)

gene, as shown in FIGURE 2A. For GBV-C, only AUGs conserved in all isolated examined are depicted.

In vitro transcription-translation (IVTT) reactions containing rabbit reticulocyte lysates were programmed with pA15-707/CAT, pC1-631/CAT and a positive control plasmid, pHAV-CAT1, which contained the 5' NTR of hepatitis A virus (HAV) inserted upstream of CAT. All three plasmid DNAs directed the translation of discreet products migrating with somewhat different molecular masses in SDS-PAGE, as shown in FIGURE 2B. Referring to the FIGURE 2B, the image was generated from a 16 h exposure with a linear range of 7 to 200.

10 GBV-CAT product in lanes 1 and 2 are present at 26 to 27% of the level of the CAT product made from pHAV-CAT1 (lane 3) when the number of Cys residues have been normalized for each product. The products derived from pA15-707/CAT and pC1-631/CAT were slightly larger than that derived from pHAV-CAT1, indicating that translation was initiating upstream of the site of GBV-CAT fusion. In contrast, no product was detected in IVTT reactions programmed with pC631-1/CAT which contained the GBV-C sequences inserted in the antisense orientation relative to CAT. Only the pHAV-CAT1-programmed reaction possessed detectable CAT activity (data not shown). The absence of activity in the products of reactions programmed with pA15-707/CAT and pC1-631/CAT was

15 likely to be due to the misfolding of the CAT protein as a result of its fusion with the N-terminal segment of the GBV polyprotein.

20

To confirm that the products of the reactions programmed with pA15-707/CAT and pC1-631/CAT were in fact GBV-CAT fusion proteins, the pA15-707/CAT and pC1-631/CAT plasmids were digested with *Ssp*I prior to being used to program reactions. *Ssp*I linearized these plasmids within the CAT coding region so that run-off transcripts produced from these plasmids would lack sequences encoding the C-terminal 45 amino acids of CAT. As expected, reactions programmed with the *Ssp*I-digested pA15-707/CAT and pC1-631/CAT DNAs (FIGURE 2B, lanes 5 and 6, respectively) contained products that were approximately 5 kDa smaller than those found in reactions programmed with undigested pA15-707/CAT and pC1-631/CAT plasmids (lanes 1 and 2 of FIGURE 2B, respectively).

2. Site of translation initiation in GBV-A and GBV-C. The apparent molecular masses of the GBV-CAT fusion proteins shown in FIGURE 2B suggested possible sites of translation initiation. As indicated in FIGURE 1, the GBV-A and GBV-C ORFs that were ligated to CAT in pA15-707/CAT and pC1-

631/CAT each contained two in-frame AUG codons that might serve as potential sites of translation initiation within the sequence immediately upstream of CAT. These were the fourth and fifth AUG codons in each of the GBV-A and GBV-C sequences (see FIGURE 2A). If initiation occurred at the fourth AUG, the 5 resultant fusion proteins would contain 46 amino acids of GBV-A (adding 5.1 kDa to the 24 kDa of CAT) (SEQUENCE ID NO 30) or 67 amino acids of GBV-C (adding 7.5 kDa to CAT) (SEQUENCE ID NO 31), respectively. In contrast, initiation at the fifth AUG in these transcripts would produce CAT fusion proteins containing 38 and 36 amino acids of GBV-A and GBV-C encoded protein, 10 respectively, adding 4.1 kDa to CAT. The apparent molecular mass of the ~28 kDa fusion proteins detected in the reactions programmed with pA15-707/CAT and pC1-631/CAT suggested that translation initiates at the fifth AUG in each transcript (i.e., the second in-frame Met codons in the long ORF, which are located at nt 594 15 of the GBV-A sequence [SEQUENCE ID NO 23] and nt 524 of the GBV-C sequence [SEQUENCE ID NO 4]). To identify the sites of translation initiation, the first and second in-frame AUG codons in GBV-A (SEQUENCE ID NO 23) and GBV-C (SEQUENCE ID NO 4) were changed to UAG stop codons producing pAmut1/CAT, pAmut2/CAT, pCmut1/CAT and pCmut2/CAT, as shown in FIGURE 3A. These plasmids were used to program IVTT reactions. 20 GBV-CAT fusion proteins were detected in reactions programmed with pAmut1/CAT and pCmut1/CAT, as shown in FIGURE 3B, lanes 2 and 5, respectively). Referring to FIGURE 3B, image characteristics are identical to those of FIGURE 2B. The GBV-CAT proteins in Lanes 1 and 4 are present at 35 to 41% of the level of CAT produced from pHAV-CAT1 template (lane 7). Amut 25 1 (lane 2) is 94% of A15-707 (lane 1); Cmut1 (lane 5) is 42% of C1-631 (lane 4). Reactions programmed with pAmut2/CAT and pCmut2/CAT (FIGURE 3B, lanes 3 and 6, respectively) did not produce detectable quantities of fusion protein. Thus, because the 28 kDa GBV-CAT protein was detected when the first in-frame 30 AUG codon (nt. 570 in GBV-A [SEQUENCE ID NO 23] and nt. 431 in GBV-C [SEQUENCE ID NO 4]) was replaced with a stop codon, initiation did not occur at this position. However, mutation of the second in-frame AUG codon (nt. 594 in GBV-A [SEQUENCE ID NO 23] and nt. 524 in GBV-C [SEQUENCE ID NO 4]) completely abrogated protein production directed by these constructs, 35 consistent with the second in-frame AUG being the site of translation initiation in both GBV-A (SEQUENCE ID NO 23) and GBV-C (SEQUENCE ID NO 4). In a related experiment, IVTT reactions programmed with a plasmid containing GBV-C

sequence with an AUG to ACG change at the position of the second in-frame AUG (nt 524) produced protein of identical size to pC1-631/CAT, although at a diminished level (data not shown). Because initiation has been found to occur with lower efficiency at ACG codons in other mRNAs (R. Böck et al., EMBO J 5 13:3608-3617 [1994]), these data are consistent with translation of the GBV-C/CAT fusion protein initiating at the ACG codon.

The number and position of Leu residues immediately downstream of the initiator Met in both GBV-A (SEQUENCE ID NO 23) and GBV-C (SEQUENCE ID NO 4) provided a biochemical method to confirm the position of the initiation site in the GBV-CAT fusion proteins. IVTT reactions containing ³H-Leu were programmed with pA15-707/CAT and pC1-631/CAT. Reaction products were separated by SDS-PAGE, transferred onto a solid support, and the 28 kDa protein bands were excised. The N-terminal amino acids of the resultant GBV-CAT fusion proteins were sequentially removed by Edman degradation and each fraction was analyzed by scintillation counting. These results are shown in FIGURE 4A and 4B. The ³H-Leu profile obtained from the pA15-707/CAT product was consistent with the expected sequence of GBV-A downstream of the second in-frame AUG, as shown in FIGURE 4A) assuming that the N-terminal Met residue is removed (see, F. Sherman et al., Bioessays 3:27-31 [1985]). Some trailing of the ³H signal was noted which may be attributed to incomplete removal of the N-terminal Met. However, for the pC1-631/CAT product, the ³H-Leu profile exactly matched the expected amino acid sequence downstream of the second in-frame AUG for GBV-C, as shown in FIGURE 4B). Referring to FIGURE 4B, CPM following each degradation cycle is plotted above the predicted N-terminal sequences (minus initiator Met) of HGBV-A (SEQUENCE ID NO 30) and GBV-C (SEQUENCE ID NO 31). These experiments thus confirm that translation is initiated at nt 594 of the GBV-A sequence (SEQUENCE ID NO 23) and nt 524 of the GBV-C sequence (SEQUENCE ID NO 4). The relative length of the 5' nontranslated RNA segments and the multiple AUG codons (some of which are in good context for translation initiation) upstream of the authentic initiator AUG in these transcripts both suggest that translation is initiated on these RNAs by internal ribosomal entry, rather than by a conventional 5' scanning mechanism. Thus, we concluded that it is likely that the GBV-A and GBV-C 5' sequences contain an IRES.

35 3. GBV coding sequence is required for efficient translation of monocistronic RNAs. The results of the *in vitro* translation reactions described

above demonstrated that initiation begins at the Met residue positioned immediately upstream of the putative E1 signal sequence in both pA15-707/CAT and pC1-631/CAT. To determine the 3' limits of the apparent IRES in GBV-A and GBV-C, and whether any amount of GBV sequence is necessary for protein production in the IVTT assays, several 3' deletions were made which reduced the amount of GBV sequence in the GBV-CAT fusion proteins. A schematic of these constructs is shown in FIGURE 5A. Protein production was observed in reactions programmed with the deletion constructs pA15-665/CAT and pC1-592/CAT, which encode 72 and 69 nucleotides of the GBV-A (SEQUENCE ID NO 23) and GBV-C (SEQUENCE ID NO 4) coding sequence fused to CAT, respectively, and as shown in FIGURE 5B, lanes 2 and 6). Referring to FIGURE 5B, image characteristics are identical to those of FIGURE 2B. GBV-CAT protein (lanes 1, 2, 5 and 6) is present at 20 to 36% of the level of CAT produced from the pHAV-CAT1 template (lane 9). In contrast, no protein was detected in reactions programmed with the deletion constructs pA15-596/CAT, pC1-526/CAT, pA15-629/CAT or pC1-553/CAT which contain three (pA15-596/CAT, pC1-526/CAT), 36 (pA15-629/CAT) or 30 (pC1-553/CAT) nts of the GBV coding sequence ligated in-frame with CAT. These results demonstrate, rather surprisingly, that sequences downstream of the predicted initiator AUG are necessary for efficient translation initiation *in vitro*. Given that the authentic initiator codons are in good context in both GBV-A (SEQUENCE ID NO 23) and GBV-C (SEQUENCE ID NO 4), these data provide further evidence that translation is not initiated by a conventional 5' scanning mechanism.

The quantity of CAT produced from the control plasmid, pHAV-CAT1 (seen in FIGURE 5B, lane 9), was considerably greater than that produced from either the GBV-A (SEQUENCE ID NO 23) or GBV-C (SEQUENCE ID NO 4) monocistronic constructs. This is of interest, because the HAV IRES has been known to direct the internal initiation of translation with very low efficiency relative to other picornaviral IRES elements (L. E. Whetter et al., J. Virol. 68:5253-5263 [1994]). The low production of GBV-CAT proteins was believed not likely to be due to differences in T7 transcriptional efficiency in these IVTT assays, as similar results were obtained with reactions programmed with equal amounts of RNA (data not shown). Thus, it appears that the level of GBV-CAT protein reflects the extremely low efficiency with which the GBV IRESs direct internal initiation *in vitro*.

4. Translation of bicistronic GBV RNAs. In an effort to formally demonstrate that the 5' RNA sequences of GBV-A and GBV-C contain IRESs, these sequences were inserted between CAT and luciferase (Luc) genes to create bicistronic T7 transcriptional units. These results are graphically shown in FIGURE 6A. IVTT reactions programmed with the bicistronic constructs produced equivalent amounts of CAT activity and CAT protein, as shown in FIGURE 6B). Referring to FIGURE 6B, CAT activity was equivalent in the reactions shown ($157,000 \pm 3,550$ cpm). The PhosphorImager scan was generated from a 72 h exposure with a linear range of 25 to 600. Band volumes are reported in FIGURE 6B without background subtraction. This confirmed that essentially equivalent amounts of RNA were being transcribed in each reaction. In contrast, the level of Luc activity and amount of Luc protein produced was dependent on the sequence cloned into the intercistronic space upstream of Luc. Although much less than the level of Luc produced from two positive control plasmids containing the IRES of HCV in the intercistronic space (270,000 to 540,000 light units, FIGURE 6B), detectable levels of Luc activity were produced only in reactions programmed with GBV bicistronic constructs containing GBV-A (SEQUENCE ID NO 23) and GBV-C sequences (SEQUENCE ID NO 4) in the sense orientation (10,300 to 13,300 light units, FIGURE 6B). Although the quantities of Luc produced were barely detectable by SDS-PAGE, PhosphorImager analysis of these gels indicated that Luc enzymatic activity did not correlate with the protein detected in the IVTT assays (FIGURE 6B, Luc-A versus Luc-P). This was most likely due to altered activity as a result of the GBV fusion. Of greater importance, however, was the fact that no detectable protein and only minimal Luc activities (130 and 2020 light units) were produced in reactions programmed with bicistronic constructs containing GBV-A (SEQUENCE ID NO 23) and GBV-C sequences (SEQUENCE ID NO 4) in the antisense orientation. These results suggest that these viruses utilize internal ribosome entry for initiation of translation, but the extraordinarily low activities of the putative GBV IRES elements when placed in a bicistronic context raises a number of issues which are discussed hereinbelow.

5. Secondary structure of the 5' NTR of GBV-C. The results presented above suggested that translation of the GBV-A and GBV-C polyproteins is initiated by an unusual mechanism of internal ribosomal entry, which is likely to be controlled by RNA structures within the 5' nontranslated RNA, and which is also dependent upon sequence downstream of the initiator AUG (see FIGURE 5).

Thus, we attempted to characterize the secondary structure near the 5' end of GBV-C RNA using a combination of phylogenetic analysis and thermodynamic predictions. Covariant nucleotide substitutions indicative of conserved base-pair interactions were identified by manual search of an alignment of 41 different GBV-C sequences. These were used to constrain the folding of the RNA by the computer program, MFOLD. Alternative structures were reviewed to determine which were most permissive for observed variations in the nucleotide sequence, resulting in the model for secondary structure shown in FIGURE 7A-D. Referring to FIGURE 7A-D, the model structure resulted from a combination of 5 phylogenetic analysis and computational thermodynamic prediction. With minor variation, the structure shown can be assumed by all available known GBV-C sequences. The predicted secondary structure of the 5' NTR of GBV-C is very 10 different from that of HCV (E. A. Brown et al., Nuc. Acid Res. 20:5041-5045 [1992] and M. Honda et al., manuscript submitted) suggesting that the 5' NTRs of 15 these viruses have distinctly different evolutionary histories.

The model suggests that the 5' RNA of GBV-C contains 4 major secondary structure domains upstream of the authentic initiator AUG at nt 524 which is conserved in all GBV-C sequences (domains I - IV in FIGURE 7A-D). Domain I consists of an extended stem-loop structure, which is highly conserved 20 in nucleotide sequence between nts 68-152, but which contains several covariant nucleotide substitutions within the flanking RNA segments near its base (FIGURE 7A-D, boxed base pairs). The predicted structure of the conserved sequence between nts 68-152 is confirmed by the presence of covariant nucleotide substitutions in alignments of GBV-C with GBV-A, which shares a very similar 25 overall 5' NTR secondary structure (not shown). Domain II contains two small stem-loops (IIa and IIb), both of which are supported by the presence of covariant substitutions in different GBV-C strains. The larger, complex stem-loops which comprise domains III and IV of the model structure are also well supported by covariant substitutions among different GBV-C strains (FIGURE 7A-D). Of 30 particular interest, given the requirement for the inclusion of coding sequence for efficient translation of monocistronic GBV transcripts (FIGURE 5), is evidence suggesting the existence of a very stable, conserved stem-loop containing 9-10 G-C base-pairs within the ORF, downstream of the putative 5' NTR (see below) (FIGURE 7A-D). The existence of this stable helical structure is supported by the 35 presence of a single covariant substitution among different GBV-C strains. This stem-loop appears to be an extension of a larger, well conserved structure (domain

V, FIGURE 7A-D), located 20 nts downstream of the putative initiator AUG. Importantly, a very similar structure is present near the 5' end of the ORF of GBV-A (FIGURE 7A-D, inset).

F. Discussion.

5 Monocistronic mRNAs containing the 5' ends of the GBV-A and GBV-C genomic RNAs fused to CAT directed the production of GBV-CAT fusion proteins in IVTT reactions. Site-specific mutagenesis and Edman degradation of the translation products indicated that translation of these transcripts, and presumably GBV-A and GBV-C genomic RNAs as well, initiates immediately 10 upstream of the putative E1 envelope signal sequence, at the AUG located at nt 594 in GBV-A (SEQUENCE ID NO 23) and nt 524 in the GBV-C sequence (SEQUENCE ID NO 4). The site of initiation identified in GBV-C is corroborated by analysis of the 5' RNA sequences obtained from 35 different GBV-C positive individuals. When these sequences are aligned, the only conserved AUG codon 15 which is in-frame with the GBV-C polyprotein is the AUG at nt 524. Downstream of this AUG codon, nucleotide substitutions in the different GBV-C strains generally result in either silent or conservative amino acid changes. In contrast, upstream of this AUG codon nucleotide substitutions, deletions and insertions drastically change the encoded amino acid sequence in different strains. These data 20 suggest that there is a selective pressure acting downstream of the AUG at nt 524 to maintain a protein coding sequence while no selective pressure exists to maintain such a sequence upstream of this codon.

The fact that translation initiates at the fifth AUG codon in both viral 25 RNAs, many hundreds of nucleotides from the 5' end, is strongly reminiscent of translation in the picornaviruses and HCV, and suggests that translation may be initiated by binding of the 40S ribosomal subunit at an internal site on the RNA. Thus, it seems likely that the 5' NTRs of these viruses may contain an IRES. 30 Because the functional activities of the IRES elements of HCV and the picornaviruses are known to be highly dependent on RNA secondary structure within the 5' NTR, we sought evidence for conserved secondary RNA structures 35 within the 5' NTRs of these viruses. Although the 5' nucleotide sequences of the GBV-C and GBV-A virus genomes have only ~50% nucleotide identity within the 500 nts preceding the initiator AUG of GBV-C, we found the secondary structures of these RNAs to be remarkably similar. Each of the major secondary structural domains shown for GBV-C in FIGURE 7A-D is conserved in the structure of GBV-A with only minimal changes (data not shown). However, both the GBV-A

and GBV-C 5' NTR structures are very different from those of the pestiviruses, HCV, and GBV-B, despite the fact that these viruses share a common genome organization as well as multiple sequence motifs within their nonstructural proteins (T. P. Leary et al., *supra* and A. S. Muerhoff et al., *supra*). While the 5' NTRs of 5 GBV-B, HCV and the pestiviruses are particularly closely related to each other at the structural level (E. A. Brown et al., *supra* and M. Honda et al., *supra*), the prominent domain III pseudoknot and complex stem-loop III structures of these viruses are completely lacking in GBV-C and GBV-A. In addition there is no clear-cut structural relatedness to HCV or the pestiviruses in any of the upstream 10 secondary structures of GBV-A and GBV-C. Thus, similar to the existence of two distinct types of 5' NTR structures among the picornaviruses (one in the cardioviruses, aphthoviruses, and hepatoviruses, and another in the enteroviruses and rhinoviruses [R. J. Jackson et al., *Mol. Biol. Reports* 19:147-159 {1994}]), there are two distinct types of 5' NTR structures present in the flaviviruses. This 15 has interesting implications for the evolution of these agents.

A prominent feature of the 5' NTR sequences of GBV-C and GBV-A is the presence of a short oligopyrimidine tract located just upstream of the initiator AUG. While this tract is somewhat variable in sequence, it is present in all of the GBV-C sequences and is positioned approximately 21 nts upstream of the initiator 20 AUG. Thus, this region of the 5' NTR bears remarkable similarity to the "box A" / "box B" motif identified at the 3' end of picornaviral 5' NTRs by Pilipenko et al. (E. V. Pilipenki et al., *Cell* 68:119-131 [1992]), including the distance (20 to 25 nts) between the start of the pyrimidine tract and the first downstream AUG in GBV-C (the initiator AUG), which Pilipenko et al. found to be critical to 25 poliovirus IRES-directed translation. It is interesting that the segment intervening between the oligopyrimidine tract and the first downstream AUG is somewhat shorter in the GBV-A viruses (approximately 17 nts). By analogy with the picornaviruses (Pilipenko et al., *supra*), this might be expected to result in a preference for initiation of translation at the second in-frame AUG codon in GBV- 30 A (nt +25 with respect to the first AUG). We confirmed this experimentally (see FIGURE 4A). The striking differences between the 5' NTR structures of these viruses and that of HCV, coupled with these similarities between the translation of GBV-A and GBV-C and picornaviral 5' NTRs, suggests that the mechanism of translation might be closer to that of picornaviruses than HCV. In HCV, relatively 35 strong evidence supports the concept that the 40S ribosomal subunit binds RNA directly at the site of translation initiation (Honda et al., *supra*). In contrast, the

40S subunit appears to scan for a variable distance from an upstream primary binding site to the initiator AUG in some picornaviruses (R. J. Jackson et al, supra). Given the variable distances between the authentic initiator codons and the upstream oligopyrimidine tracts in GBV-A and GBV-C, this appears likely to be 5 the case with GBV-A (and possibly also GBV-C).

Both GBV-A and GBV-C contain a very stable stem-loop structure within the translated open reading frame (domain V, FIGURE 7D.). This conserved structure is located about 20 nts downstream of the initiator AUG in GBV-C, although it is possible that additional, less well conserved base-pair interactions 10 may bring the base of this structure closer to the AUG. It is tempting to speculate that this stem-loop may function to enhance initiation by a scanning 40S ribosomal subunit, much as M. Kozak, Proc. Natl. Acad. Sci USA 87:8301-8305 (1990) has shown that stable stem-loops placed downstream of an AUG can result in a "pausing" of the ribosome over the AUG, enhancing the likelihood of initiation at 15 that codon. This phenomenon may explain why the efficient translation of reporter proteins fused to the 5' NTR requires inclusion of the most 5' sequence of the GBV-C open reading frame. If so, this would provide a novel mechanism by which sequence within the open-reading frame can contribute to regulation of translation in flaviviruses. Both HCV and the GBV-B viruses differ from GBV-A 20 and GBV-C in that their initiator AUG is located within the loop segment of a stem-loop which straddles the 5' end of the open reading frame (M. Honda, supra). Initiation of translation of these viral RNAs is thus dependent upon melting of this stem-loop while, in the case of GBV-A and GBV-C, initiation of 25 translation is likely to be dependent on maintenance of the integrity of the domain V stem-loop.

The domain V stem-loop for which is required for efficient translation of the monocistronic transcripts does not appear to be required for efficient translation in the bicistronic transcripts (compare FIGURES 5 and 6). This apparent discrepancy may be a result of the different reporter genes being utilized in these 30 transcripts. Similar findings have been reported for HCV. Specifically, Reynolds et al., supra, using bicistronic vectors with the IRES-dependent reporter genes secreted alkaline phosphatase or a truncated influenza virus nonstructural protein, show efficient translation directed by the 5' end of HCV requires the inclusion of coding sequences. In contrast, Wang et al., supra, using monocistronic and 35 bicistronic vectors with luciferase as the IRES-dependent reporter gene, find the inclusion of HCV coding sequences is not necessary for efficient translation.

Addressing these conflicting results, Reynolds et al., supra, hypothesize that the 5' end of the luciferase gene may complement the function provided by the HCV coding sequences. A similar argument may explain the discordance between the results obtained with the monocistronic GBV-CAT constructs and the bicistronic 5 GBV-Luc constructs.

Although all of these observations suggest the strong likelihood that GBV-A and GBV-C translation is initiated by internal ribosomal entry, only minimal translation of the downstream cistron was noted from bicistronic transcripts containing the 5' NTRs of these viruses in the intercistronic space. Translation 10 directed by the GBV-A and GBV-C 5' NTRs within a bicistronic context was only 2 to 5% that of the HCV IRES in rabbit reticulocyte lysates *in vitro* (FIGURE 6). The very low activities of the GBV-A and GBV-C IRESs suggest several 15 possibilities. First, it is possible that these viruses may in fact have IRES elements with extraordinarily low activity. This is supported by a very low level of translation directed by monocistronic transcripts containing the 5' ends of GBV-A and GBV-C in the *in vitro* system. Specifically, after adjustment for the number of Cys residues in each construct, GBV-CAT fusion proteins were translated from pA15-707/CAT and pC1-631/CAT transcripts at only 20 to 41% of the level 20 produced by the IRES of HAV. The HAV IRES is known to have very low activity, in the range of 2% of the Sabin poliovirus type I IRES within HAV permissive cells (see, D. E. Schultz et al., *J. Virol.* 70:1041-1049 [1996] and L. E. Whetter et al., supra). Thus, the low GBV IRES activity noted *in vitro* may be 25 a true reflection of the strength of these translation elements. Limiting production of viral proteins within an infected host might act to reduce recognition of the infection by the immune system and thus promote viral persistence. Alternatively, it is possible that the low IRES activity detected in reticulocyte lysates reflects a requirement for a specific host cell translation factor which is absent in reticulocyte lysates. The nuclear autoantigen, La, is an example of such a specific cellular factor. It is required for efficient translation directed by the poliovirus IRES, but is 30 not present in sufficient amounts in reticulocyte lysates. K. Meerovitch et al., *J. Virol.* 67:3798-3807 (1993). It is difficult to comment more specifically on this possibility, since the cellular tropisms of GBV-A and GBV-C are unknown. Yet a third possibility is that the low translational activity of the GBV-A and GBV-C 5' NTRs may reflect a requirement for additional, yet to be identified 5' viral 35 sequences that may be present in these viral genomes. It is also conceivable that translation is initiated by a mechanism distinct from both the classic 5' scanning

and IRES-directed translation initiation mechanism. For example, relatively efficient translation initiation at an internal site in monocistronic transcripts but low translational activity in the bicistronic context could be explained by a mechanism involving "ribosome shunting" (J. Fütterer et al., Cell 73:789-802 [1993])

5 following recognition of the 5' end of the RNA by the 40S ribosome subunit. Further studies will be required to distinguish between these different possibilities.

The proteins located at or near the amino termini of the polyproteins of yellow fever virus (protein C), a flavivirus, bovine viral diarrhea virus, a

pestivirus, and HCV (core) are small and highly basic (Q.-L. Choo et al., Proc.

10 Natl. Acad. Sci. USA 88:2451-2455 [1991]; M.S. Collett et al., supra; R. H. Miller et al., Proc. Natl. Acad. Sci. USA 87:2057-2061 [1990]). Because GBV-A and GBV-C are phylogenetically related to these viruses (12, 18) it was expected that such a protein would be encoded in these viruses. However, the position of the initiation codons in GBV-A and GBV-C eliminates the possibility of a basic

15 core protein being located at the N-termini of the viral polyproteins. The possibility that the core coding sequences may have been deleted during RT-PCR amplification or cloning of the 5' ends of GBV-A and GBV-C is unlikely for several reasons. First, identical deletions would have had to occur consistently in each of the several clones generated during the sequencing of GBV-A and GBV-C,

20 in addition to the 42 separate GBV-C isolates described by U.S. Serial No. 08/580,038, filed December 21, 1995 and previously incorporated herein by reference, and the 2 HGV isolates described by Linnen et al., supra. This consistency, in addition to the correspondence between PCR and infective titers for

25 GBV-A (G. G. Schlauder et al., J. Med. Virol. 46:81-90 [1995] and J. N. Simons, Proc. Natl. Acad. Sci USA, supra), argues against GBV-A and GBV-C

sequences being derived from defective interfering particles in the cloning sources. Second, the deletion of core sequences would have had to occur without disturbing the translational activity of the 5' ends of these viruses. But because proper initiation requires sequences located in the coding regions of GBV-A and GBV-C,

30 the coupling between the translational activity and the coding regions appear to make this an impossibility. Finally, several RT-PCR experiments using different virus isolates, different primer combinations, and different RT-PCR conditions and polymerases provide no evidence for additional virus sequence (data not shown).

35 The lack of a core-like protein at the N-terminus of the viral polyprotein distinguishes GBV-A and GBV-C from all other members of the *Flaviviridae*. In

fact, searches of all six potential reading frames of the three full length GBV-C sequences (T. P. Leary et al., *supra* and L. Linnen et al., *supra*) or the GBV-A sequence (SEQUENCE ID NO. 23) present in GenBank does not reveal a conserved open reading frame encoding a core-like protein. Thus, these viruses 5 appear distinct from enveloped viruses in general as they do not appear to encode a basic protein which mediates the packaging of the viral nucleic acid into the virion envelope. Core-less infectious particles have been generated artificially using the vesicular somatitis virus glycoprotein. M. M. Rolls et al., *Cell* 79:497-506 (1994). Thus, it is possible that GBV-A and GBV-C may be truly "core-less" 10 enveloped viruses. However, it is possible that a cellular RNA-binding protein has been appropriated by these viruses to facilitates the specific and efficient packaging of the virion RNA into the envelope. Whether GBV-A and GBV-C contain core proteins and the source of these cores awaits the biochemical characterization of these viruses.

15

The present invention is intended to be limited only by the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Simons, J. N.
Desai, S. M
Mushahwar, I. K.

(ii) TITLE OF INVENTION: REAGENTS AND METHODS USEFUL FOR CONTROLLING THE TRANSLATION OF HEPATITIS GB PROTEINS

(iii) NUMBER OF SEQUENCES: 32

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Abbott Laboratories
(B) STREET: 100 Abbott Park Rd
(C) CITY: Abbott Park
(D) STATE: IL
(E) COUNTRY: USA
(F) ZIP: 60064

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Porembski, Priscilla E.
(B) REGISTRATION NUMBER: 33,207
(C) REFERENCE/DOCKET NUMBER: 5793.US.P1

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 708-937-0378
(B) TELEFAX: 708-938-2623

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCACAAACAC TCCAGTTGT TAC

23

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCTCTAGACA TGTGCTACGG TCTACGAG

28

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9126 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCCCCCCCCCC GGCACTGGGT GCAAGCCCCA GAAACCGACG CCTACTGAAG TAGACGTAAT	60
GGCCCCCGCGC CGAACCGGCG ACCGGCCAAA AGGTGGTGGA TGGGTGATGA CAGGGTTGGT	120
AGGTCTGAAA TCCCCGTCAT CCTGGTAGCC ACTATAGGTG GGTCTTAAGG GGAGGGCTACG	180
GTCCCTCTTG CGCATATGGA GGAAAAGCGC ACGGTCCACA GGTGTTGGTC CTACCGGTGT	240
AATAAGGACC CGGCGCTAGG CACGCCGTTA AACCGAGCCC GTTACTCCCC TGGGCAAACG	300
ACGCCACGT ACGGTCCACG TCGCCCTTCA ATGTCTCTCT TGACCAATAG GCGTAGCCGG	360
CGAGTTGACA AGGACCAGTG GGGGCCGGGC GGGAGGGGGA AGGACCCCCA CCGCTGCCCT	420
TCCCCGGGAG GCGGGAAATG CATGGGGCCA CCCAGCTCCG CGGCAGCCTA CAGCCGGGGT	480
AGCCCAAGAA CCTTCGGGTG AGGGCGGGTG GCATTTCTTT TCCTATACCG ATCATGGCAG	540
TCCTTCTGCT CCTACTCGTG GTGGAGGCCG GGGCTATTTT AGCCCCGGCC ACCCATGCTT	600
GTAGCGCGAA AGGGCAATAT TTBCACCAA ACTGTTGCGC CCTGGAGGAC ATAGGCTTCT	660

GCCTGGAGGG CGGATGCCTG GTGGCTCTGG GGTGCACCAT TTGCACCGAC CGCTGCTGGC	720
CACTGTATCA GGCGGGTTTG GCCGTGCGC CCGCAAGTC CGCCGCCAG TTGGTGGGG	780
AACTCGGTAG TCTCTACGGG CCCTTGTCTGG TCTCGGCTTA TGTGGCCGGG ATCCTGGGC	840
TTGGGGAGGT CTACTCGGGG GTCCTCACCG TCAGGGTGGC GTTGACGCGC AGGGTCTACC	900
CGGTCCCGAA CCTGACGTGT GCAGTAGAGT GTGAGTTGAA GTGGGAAAGT GAGTTTGGA	960
GATGGACTGA ACAGCTGGCC TCAAACACT GGATTCTGGA ATACCTCTGG AAGGTGCCTT	1020
TCGACTTTG GCGGGGAGTG ATGAGCCTTT CTCCTCTCTT GGTGTGCGTG GCGGCCCTCC	1080
TCCTGCTGGA GCAGCGTATT GTCATGGTCT TCCTCCTGGT CACTATGGCG GGCATGTCAC	1140
AAGGCGCGCC CGCCTCAGTG TTGGGGTCAC GGCCCTTCGA GGCCGGCTG ACTTGGCAGT	1200
CTTGTTCCTTG CAGGTCGAAC GGGTCCCGCG CGCCGACAGG GGAGAGGGTT TGGAACGTG	1260
GGAACGTCAC ACTTTTGTGT GACTGCCCA ACGGTCTTGT GGTGTGGTC CGGCCCTTT	1320
GCCAGGCAAT CGGATGGGC GACCCTATCA CTCATTGGAG CCACGGACGA AATCAGTGGC	1380
CCCTTCTTG TCCCCAATT GTCTACGGCG CCGTTTCAGT GACCTGCGTG TGGGGTTCTG	1440
TGTCTTGGTT TGCTTCCACT GGGGGTCGCG ACTCCAAGGT TGATGTGTGG AGTTTGGTTC	1500
CAGTTGGCTC TGCCAGCTGT ACCATAGCCG CACTGGGATC TTCGGATCGC GACACAGTGG	1560
TTGAGCTCTC CGAATGGGA ATCCCCTGCG CCACTTGTAT CCTGGACAGG CGGCCTGCCT	1620
CGTGTGGCAC CTGTGTGAGG GACTGCTGGC CCGAGACCGG GTCGGTACGT TTCCCATTC	1680
ACAGGTGTGG CGCGGGACCG AGGCTGACCA GAGACCTTGA GGCTGTGCC CTCGTCAATA	1740
GGACAACCTCC CTTCACCAATA AGGGGGCCCC TGGGCAACCA GGGGCGAGGC GACCCGGTGC	1800
GGTGCCTT GGGTTTGGG TCCTACACCA TGACCAAGAT CCGAGACTCC TTACACTTGG	1860
TGAAATGTCC CACCCCAGCC ATTGAGCCTC CCACCGGAAC GTTTGGGATC TTCCAGGAG	1920
TCCCCCCCCT TAACAACCTGC ATGCTTCTCG GCACTGAGGT GTCAGAGGTA TTGGGTGGGG	1980
CGGGCCTCAC TGGGGGGTTT TACGAACCTC TGGTGCAGCG GTGTTCAGAG CTGATGGTC	2040
GGCGGAATCC GGTCTGCCCG GGGTTTCAT GGCTCTCTTC GGGACGGCCT GATGGGTCA	2100
TACATGTACA GGGCCACTTG CAGGAGGTGG ATGCGGGCAA CTTCATCCG CCCCCACGCT	2160
GGTTGCTCTT GGACTTTGTA TTTGTCCGT CATACTGAT GAAGCTGGCA GAGGCACGGT	2220
TGGTCCCGCT GATCCTCCTC CTGCTATGGT GGTGGGTGAA CCAGTTGGCG GTCCTTGKAC	2280
TGSCGGCTGC KCRCGCCGCC GTGGCTGGAG AGGTGTTGC GGGCCCTGCC TTGTCTGGT	2340
GTCTGGGCCT ACCCTTCGTG AGTATGATCC TGGGGCTAGC AACACCTGGTG TTGTACTTCC	2400

GCTGGATGGG TCCTCAACGC CTGATGTTCC TCGTGTGTG GAAGCTCGCT CGGGGGGCTT	2460
TCCCGCTGGC ATTACTGATG GGGATTTCCG CCACTCGCGG CGGCACCTCT GTGCTTGGCG	2520
CCGAATTCTG CTTTGATGTC ACCTTTGAAG TGGACACGTC AGTCTTGGGT TGGGTGGTTG	2580
CTAGTGTGGT GGCTTGGGCC ATAGCGCTCC TGAGCTCTAT GAGCGCGGGG GGGTGGAAGC	2640
ACAAAGCCAT AATCTATAGG ACGTGGTGT AAGGGTACCA GGCYCTTCGC CAGCGCGTGG	2700
TGCGTAGCCC CCTCGGGGAG GGGCGGCCA CCAAGCCGCT GACGATAGCC TGGCGTCTGG	2760
CCTCTTACAT CTGGCCGGAC GCTGTGATGT TGGTGGTTGT GGCCATGGTC CTCCTCTTCG	2820
GCCTTTTCA CGCGCTCGAT TGGCCTTGG AGGAGCTCCT TGTGTCGCGG CCTTCGTTGC	2880
GTCGTTGGC AAGGGTGGTG GAGTGTGTG TGATGGCGGG CGAGAAGGCC ACTACCGTCC	2940
GGCTTGTGTC CAAGATGTGC GCGAGAGGGG CCTACCTGTT TGACCACATG GGGTCGTTCT	3000
CGCGCGCGGT CAAGGAGCGC TTGCTGGAGT GGGACGCGGC TTTGGAGMCC CTGTCATTCA	3060
CTAGGACGGA CTGCCGCATC ATACGAGACG CCGCCAGGAC TCTGAGCTGC GGCCAATGCG	3120
TCATGGGCTT GCCCGTGGTG GCTAGGCGCG GCGATGAGGT CCTGGTTGGG GTCTTCAGG	3180
ATGTGAACCA CTTGCCTCCG GGGTTGYTC CTACAGCGCC TGTGTCATC CGTCGGTGCG	3240
GAAAGGGCTT CCTCGGGGTC ACTAAGGCTG CCTTGACTGG TCGGGATCCT GACTTACACC	3300
CAGGAAACGT CATGGTTTG GGGACGGCTA CCTCGCGCAG CATGGAACG TGCTTAAACG	3360
GGTTGCTGTT CACGACATTC CATGGGGCTT CTTCCGAAC CATTGCGACA CCTGTGGGGG	3420
CCCTTAACCC AAGGTGGTGG TCGGCCAGTG ATGACGTAC GGTCTATCCC CTCCCCGATG	3480
GAGCTAACTC GTTGGTCCCC TGCTCGTGTG AGGCTGAGTC CTGTTGGTC ATYCGATCCG	3540
ATGGGGCTCT TTGCCATGGC TTGAGCAAGG GGGACAAGGT AGAACTGGAC GTGGCCATGG	3600
AGGTTGCTGA CTTTCGTGGG TCGTCTGGGT CTCCTGTCCCT ATGCGACGAG GGGCACGCTG	3660
TAGGAATGCT CGTGTCCGTC CTTCATTGGG GGGGGAGGGT GACCGCGGCT CGATTCACTC	3720
GGCCGTGGAC CCAAGTCCCA ACAGACGCCA AGACTACCAC TGAGCCACCC CCGGTGCCAG	3780
CTAAAGGGT TTTCAAAGAG GCTCCTCTTT TCATGCCAAC AGGGGCGGGG AAAAGCACAC	3840
GCGTCCCTTT GGAATATGGA AACATGGGGC ACAAGGTCT GCTTCTCAAC CCGTCGGTTG	3900
CCACTGTGAG GGCCATGGC CCTTACATGG AGAAGCTGGC GGGGAAACAT CCTAGCATT	3960
TCTGTGGACA CGACACAAACA GCTTTCACAC GGATCACGGA CTCTCCATTG ACGTACTCTA	4020
CCTATGGGAG GTTTCTGGCC AACCCGAGGC AGATGCTGAG GGGAGTTCC GTGGTCATCT	4080

GTGATGAGTG CCACAGTCAT GACTCAACTG TGTTGCTGGG TATAGGCAGG GGCAGGGAGC	4140
TGGCGCGGGG GTGTGGAGTG CAATTAGTGC TCTACGCTAC TGCGACTCCC CCGGGCTCGC	4200
CTATGACTCA GCATCCATCC ATAATTGAGA CAAAGCTGGA CGTCGGTGAG ATCCCCTTT	4260
ATGGGCATGG TATCCCCCTC GAGCGTATGA GGACTGGTCG CCACCTTGTA TTCTGCCATT	4320
CCAAGGCGGA GTGCGAGAGA TTGGCCGGCC AGTTCTCCGC GCGGGGGGTT AATGCCATCG	4380
CCTATTATAG GGGTAAGGAC AGTTCCATCA TCAAAGACGG AGACCTGGTG GTTTGTGCGA	4440
CAGACGCGCT CTCTACCGGG TACACAGGAA ACTTCGATTG TGTCACCGAC TGTCGGTTAG	4500
TGGTGGAGGA GGTGTTGAG GTGACCCCTG ATCCCACCAT TACCAATTCC TTGCGGACTG	4560
TCCCTGCTTC GGCTGAATTG TCGATGCAGC GGCGCGGACG CACGGGAGA GGTGCGTCGG	4620
GCCGCTACTA CTACGCTGGG GTCGGTAAGG CTCCCGCGGG GGTGGTGCAG TCTGGTCCGG	4680
TCTGGTCGGC AGTGGAAAGCT GGAGTGACCT GGTATGGAAT GGAACCTGAC TTGACAGCAA	4740
ACCTTCTGAG ACTTTACGAC GACTGCCCTT ACACCGCAGC CGTCGCAGCT GACATTGGTG	4800
AAGCCGCGGT GTTCTTGCG GGCCTCGCGC CCCTCAGGAT GCATCCCGAT GTTAGCTGGG	4860
CAAAAGTTCG CGGCGTCAAT TGGCCCCCTCC TGGTGGGTGT TCAGCGGACG ATGTGTCGGG	4920
AAACACTGTC TCCCAGCCCG TCGGACGACC CTCAGTGGGC AGGTCTGAAA GGCCCGAATC	4980
CTGCCCCACT ACTGCTGAGG TGGGGCAATG ATTTGCCATC AAAAGTGGCC GGCCACCACCA	5040
TAGTTGACGA TCTGGTCCGT CGGCTCGGTG TGGCGGAGGG ATACGTGCGC TGTGATGCTG	5100
GRCCCCATCCT CATGGTGGGC TTGGCCATAG CGGGCGGCAT GATCTACGCC TCTTACACTG	5160
GGTCGCTAGT GGTGGTAACA GACTGGAATG TGAAGGGAGG TGGCAATCCC CTTTATAGGA	5220
GTGGTGACCA GGCCACCCCT CAACCCGTGG TGCAGGTCCC CCCGGTAGAC CATCGGCCGG	5280
GGGGGGAGTC TGCGCCAGCG GATGCCAAGA CAGTGACAGA TGCGGTGGCA GCCATCCAGG	5340
TGAACGTGCA TTGGTCTGTG ATGACCCCTGT CGATCGGGGA AGTCCTCACC TTGGCTCAGG	5400
CTAAGACAGC CGAGGCCTAC GCAGCTACTT CCAGGTGGCT CGCTGGCTGC TACACGGGA	5460
CGCGGGCCGT CCCCCACTGTA TCAATTGTTG ACAAGCTCTT CGCCGGGGGT TGGGCCGCCG	5520
TGGTGGGTCA CTGTCACAGC GTCATTGCTG CGGTGGTGGC TGCCTATGGG GTTTCTCGAA	5580
GTCCTCCACT GGCCCGCGCG GCATCCTACC TCATGGGGTT GGGCGTCGGA GGCAACGCAC	5640
AGGCGCGCTT GGCTTCAGCT CTTCTACTGG GGGCTGCTGG TACGGCTCTG GGGACCCCTG	5700
TCGTGGGACT CACCATGGCG GGGGCCTTCA TGGGCGGTGC CAGCGTGTCC CCCTCCCTCG	5760
TCACTGTCCCT ACTTGGGACT GTGGGAGGTT GGGAGGGCGT TGTCAACGCT GCCAGTCTCG	5820

TCTTCGACTT CATGGCTGGG AAACTTCAA CAGAAGACCT TTGGTATGCC ATCCCGGTAC	5880
TCACTAGTCC TGGRGCGGGC CTCGCAGGGGA TTGCCCTTGG TCTGGTTTG TACTCAGCAA	5940
ACAACTCTGG CACTACCACA TGGCTGAACC GTCTGCTGAC GACGGTGCCA CGGTCACTTT	6000
GCATACCCGA CAGCTACTTC CAACAGGCTG ACTACTGCGA CAAGGTCTCG GCAATGCTGC	6060
GCCGCCTGAG CCTTACTCGC ACCGTGGTGG CCCTGGTCAA CAGGGAGCCT AAGGTGGATG	6120
AGGTCCAGGT GGGGTACGTC TGGGATCTGT GGGAGTGGGT AATGCGCCAG GTGCGCATGG	6180
TGATGTCTAG ACTCCGGGC CTCTGCCCTG TGGTGTCACT CCCCTGTGG CACCGCGGGG	6240
AGGGGTGGTC CGGTGAATGG CTTCTCGATG GGCACGTGGA GAGTCGTTGT CTGTGCGGGT	6300
GTGTAATCAC CGGCGACGTC CTCAATGGGC AACTCAAAGA TCCAGTTAC TCTACCAAGC	6360
TGTGCAGGCA CTACTGGATG GGAACGTGTC CGGTCAACAT GCTGGCTAC GGGGAAACCT	6420
CACCTCTTCT CGCCTCTGAC ACCCGAAGG TGGTACCCCTT CGGGACGTG GGGTGGGCTG	6480
AGGTGGTGGT GACCCCTACC CACGTGGTGA TCAGGCGCAC GTCCCTGTTAC AACTGCTTC	6540
GCCAGCAAAT TCTTCAGCA GCTGTAGCTG AGCCCTACTA CGTTGATGGC ATTCCGGTCT	6600
CTTGGGAGGC TGACGCGAGA GCGCCGGCCA TGGTCTACGG TCCGGCCAA AGTGTACCA	6660
TTGATGGGGA GCGCTACACC CTTCCGCACC AGTTGCGGAT GCGGAATGTG GCGCCCTCTG	6720
AGGTTTCATC CGAGGTCAGC ATCGAGATCG GGACGGAGAC TGAAGACTCA GAACTGACTG	6780
AGGCCGATTG GCCACCAGCG GCTGCTGCC TCCAAGCGAT AGAGAATGCT GCGAGAATTG	6840
TCGAACCGCA CATCGATGTC AYCATGGAGG ATTGCAGTAC ACCCTCTCTC TGTGGTAGTA	6900
GCCGAGAGAT GCCTGTGTGG GGAGAAGACA TACCCCGCAC TCCATCGCCT GCACTTATCT	6960
CGGTTACGGA GAGCAGCTCA GATGAGAAGA CCCTGTCGGT GACCTCCTCG CAGGAGGACA	7020
CCCCGTCTC AGACTCATTT GAAGTCATCC AAGAGTCTGA TACTGCTGAA TCAGAGGAAA	7080
GCGTCTCAA CGTGGCTCTT TCCGTACTAA AAGCATTATT TCCACAGAGC GTGCCACAC	7140
GAAAGCTAAC GGTAAAGATG TCTTGCTGTG TTGAGAAGAG CGTAACACGC TTCTTTCTT	7200
TAGGGTTGAC CGTGGCTGAC GTGGCTAGCC TGTGTGAGAT GGAGATCCAG AACCATACAG	7260
CCTATTGTGA CAAGGTGCGC ACTCCGCTCG AATTGCAAGT TGGGTGCTTG GTGGCAATG	7320
AACTTACCTT TGAATGTGAC AAGTGTGAGG CACGCCAAGA GACCCTGCC TCCTTCTCCT	7380
ACATATGGTC CGGGGTCCCA CTTACTCGGG CCACTCCGGC CAAACCAACCA GTGGTGAGGC	7440
CGGTGGGTC CTTGTTGGTG GCAGACACCA CCAAGGTCTA CGTGACCAAT CCGGACAATG	7500

TTGGGAGGAG GGTTGACAAG GTGACTTCT GGCGCGCTCC TCGGGTACAC GACAAGTTCC	7560
TCGTGGACTC GATCGAGCGC GCTCGGAGAG CTGCTCAAGG CTGCCTAACG ATGGGTTACA	7620
CTTATGAGGA GGCAATAAAGG ACTGTTAGGC CGCATGCTGC CATGGGCTGG GGATCTAAGG	7680
TGTCGGTCAG GGACTTGGCC ACCCCTGCGG GGAAGATGGC TGTCATGAC CGGCTTCAGG	7740
AGATACTTGA AGGGACTCCA GTCCCTTTA CCCTGACTGT CAAAAAGGAG GTGTTCTCA	7800
AAGATCGTAA GGAGGGAGAAG GCCCCCGCC TCATTGTGTT CCCCCCCCTG GACTTCCGGA	7860
TAGCTGAAAA GCTCATTCTG GGAGACCCGG GGCGGGTTGC AAAGGCGGTG TGGGGGGGGG	7920
CTTACGCCTT CCAGTACACC CCCAACCAGC GGGTTAAGGA GATGCTAAAG CTGTGGGAAT	7980
CAAAGAAGAC CCCGTGCCGCC ATCTGTGTGG ATGCCACTTG CTTCGACAGT AGCATTACTG	8040
ARGAGGACGT GGCACTAGAG ACAGAGCTTT ACGCCCTGGC CTCGGACCAT CCAGAATGGG	8100
TGCGCGCCCT GGGGAAATAC TRTGCCTCTG GCACAATGGT GACCCCGGAA GGGGTGCCAG	8160
TGGGCGAGAG GTATTGTAGG TCCTCGGGTG TGTAAACCAC AAGTGCTAGC AACTGTTGA	8220
CCTGCTACAT CAAAGTGAGA GCCGCCTGTG AGAGGATCGG ACTGAAAAAT GTCTCGCTTC	8280
TCATCGCGGG CGATGACTGC TTAATTGTGT GCGAGAGGCC TGTATGCGAC CCTTGCGAGG	8340
CCCTGGGCCG AGCCCTGGCT TCGTACGGGT ACGCGTGTGA GCCCTCGTAT CACGCTTCAC	8400
TGGACACAGC CCCCTCTGC TCCACTTGGC TTGCTGAGTG CAATGCGGAT GGGRAAAGGC	8460
ATTTCTTCCT GACCACGGAC TTTCGGAGAC CACTCGCTCG CATGTCGAGC GAGTACAGTG	8520
ACCCATGGC TTCGGCCATT GGTTACATTC TCCTCTATCC CTGGCRTCCC ATCACACGGT	8580
GGGTCATCAT CCCGCATGTG CTAACATGCG CTTCTTCCG GGGTGGTGGC ACACSGTCTG	8640
ATCCGGTTTG GTGTCAGGTT CATGGTAACT ACTACAAGTT TCCCCTGGAC AACTGCTTA	8700
ACATCATCGT GGCCCTCCAC GGACCAGCAG CGTTGAGGGT TACCGCAGAC ACAACCAAAA	8760
CAAAGATGGA GGCTGGGAAG GTTCTGAGCG ACCTCAAGCT CCCTGGTCTA GCGTCCACC	8820
GCAAGAAGGC CGGGCATTG CGAACACGCA TGCTCCGGTC GCGCGGGTGG GCGGAGTTGG	8880
CTAGGGGCCT GTTGTGGCAT CCAGGACTCC GGCTTCCTCC CCCTGAGATT GCTGGTATCC	8940
CAGGGGGTTT CCCTCTGTCC CCCCCCTACA TGGGGGTGGT TCATCAATTG GATTTCACAG	9000
CSCAGCGGAG TCGCTGGCGG TGGTTGGGT TCTTAGCCCT GCTCATCGTA GCGCTTTG	9060
GGTGAACTAA ATTCACTCTGT TGCAGGCCGGA GTCAGACCTG AGCCCCGTTTC AAAAGGGAT	9120
TGAGAC	9126

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 635 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CACTGGGTGC AAGCCCCAGA AACCGACGCC TATCTAAGTA GACGCAATGA CTCGGCGCCA	60
ACTCGGCGAC CGGCCAAAAG GTGGTGGATG GGTGATGACA GGGTTGGTAG GTCGTAAATC	120
CCGGTCACCT TGGTAGGCCAC TATAAGTGGG TCTTAAGAGA AGGTTAAGAT TCCTCTTGTG	180
CCTGCGGCGA GACCGCGCAC GGTCCACAGG TGTTGGCCCT ACCGGTGTGA ATAAGGGCCC	240
GACGTCAGGC TCGTCGTTAG ACCGAGCCCG TCACCCACCT GGGCAAACGT CGCCCACGTA	300
CGGTCCACGT CGCCCTTCAA TGTCTCTCTT GACCAATAGG CTTAGCCGGC CGAGTTGACA	360
AGGACCAGTG GGGGTGGGG GCTTGGGGAG GGACCCCAAG TCCTGCCCTT CCCGGTGGGC	420
CGGGAAATGC ATGGGGCCAC CCAGCTCCGC GGCGGCCTGC AGCCGGGGTA GCCCAAGAAT	480
CCTTCGGGTG AGGGCGGGTG GCATTTCTCT TTTCTATAACC ATCATGGCAG TCCTTCTGCT	540
CCTTCTCGTG GTCGAGGCCG GGGCCATTCT GGCCCCGGCC ACCCACGCTT GTCGAGCGAA	600
TGGGGCAATA CTTCCCTACAA AACTGTTGCG CCCTG	635

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TATAATAAGC TTGCCCCGGA CCTCCACCG AG	32
------------------------------------	----

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCTCTAGATC GGGAACAAACA ATTGGAAAG

29

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TATAATAAGC TTCACTGGGT GCAAGCCCCA GAA

33

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCTCTAGAGG CGCAACAGTT TGTGAGGAA

29

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GCTCTAGACA CTGGGTGCAA GCCCCA

26

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TATAATAAGC TTGGCGCAAC AGTTTGTGAG

30

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GCTCTAGAGC ACTGGTGCCG CGAGT

25

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

49

GCTCTAGAGA GGGGGAAGCA AACCA

25

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCTCTAGACA TGGTGAATGT GTCGACCAC

29

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GCTCTAGAAC AAGCGTGGGT GGCGGGGG

28

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GCTCTAGAGA CCACGAGAAG GAGCAGAAG

29

(2) INFORMATION FOR SEQ ID NO:16:

50

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GCTCTAGACA TGATGGTATA GAAAAGAG

28

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CCATAATCAT GAGGGAACAA CAATTGGAAA G

31

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CATGCCATGG CGCAACAGTT TGTGAGGAA

29

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid

51

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TATAATAAAG CTTCACTCCC CTGTGAGGAA CTAC

34

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GTATTGCGTC ATGATGGTTT TTCTTTGGGG TTTAG

35

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TATAATAAGC TTGCCGCGAG TTGAAGAGCA C

31

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CCATAATCAT GAGCCCCGG ACCTCCCACC GAG

33

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9493 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CGTGGGAGTC CGGGGCCCG GACCTCCCAC CGAGGTGGGG GGAAAGGGGC CCTGGACCGG	60
CCGGGTGGAA GGCCCGGAAC CGGTCCATCT TCCTCAAGGT TGAGGAAGGG GTACGTCTAT	120
CGGTCCGGTC GGTCCGAAAG GCGTCTGGAT GCCTAGTGTGTT AGGGTTCGTA GGTGGTAAAT	180
CCCAGCTAGG CGTGAAAGCG CTATAGGATA GGCTTATCCC GGTGACCGCT GCCCCGGAAC	240
CAGCCCCGCG GKTCTTTGGA CACGGTCCAC AGGTTGGGGG TACCGGTGTG AATAACCCCC	300
CGACTGAAGC GTCAGTCGTT AAACGGAGAC GGTCTCCTGA GATCGCAACG ACGCCCCACG	360
TACGGGAACG CCGCCAAAAC CTTCGGGACA GCTATGCGGG TTGACAATCC CAGTGGGGGG	420
CCGGGGACCA GCTGATTACT TGTCTCGCA GTTCCTCTTG AGACTGCCG AAAGGCAGCC	480
ACGGGGCCAC CAAGGCGGCG CAGCGCTGCA TGCAGCAAGG GGAAAAATCC TTGGGGTGAC	540
CCCTGGTGGC AATCCCTTCC CTTAGGAGCA TGAGTGTGGT CGACACATTC ACCATGGCTT	600
GGCTGTGGTT GCTGGTTTGC TTCCCCCTCG CGGGGGGGGT GCTCTTCAAC TCGGGCACC	660
AGTGCTCAA TGGGGACCAT TATGTGCTTT CCAATTGTTG TTCCCGAGAC GAGGTTTACT	720
TCTGTTTCGG GGACGGATGT CTGGTGGCTT ATGGCTGTAC TGTTGCACA CACTCTTGCT	780
GGAAAGCTCTA CCGGCCTGGG GTGGCTACTC GGCCCGGGTC CGAACCAAGT GAGCTGCTGG	840
GGAGATTGG GAGTGTAAATT GGTCCGGTGT CGGCTTCGGC TTACACCGCT GGAGTCCTCG	900
GGTTGGGTGA ACCTTACAGT TTGGCCTTCT TGGGGACGTT CCTCACCAGT CGCCTCTCAC	960
GGATTCCCAA CGTCACCTGC GTGAAGGCTT GTGACCTTGA GTTTACCTAC CCAGGCTTGT	1020

CCATCGATTT	TGACTGGCG	TTTACCAAGA	TCTTGCAGTT	GCCGGCCAAG	CTGTGGCGAG	1080	
GCCTAACGGC	RGCWCCGGTC	TTGAGCCTCC	TCGTGATCCT	CATGCTGGTC	CTCGAGCAGC	1140	
GCCTCCTGAT	AGCCTTCCTA	CTGCTTTGG	TAGTGGCGA	GGCTCAGAGG	GGGATGTTCG	1200	
ACAACTGCGT	GTGTGGTTAC	TGGGGGGGCA	AGAGGCCCCC	GTCGGTGACC	CCGCTGTACC	1260	
GTGGCAACGG	TACTGTGGTG	TGTGACTGTG	ATTTTGGAAA	AATGCATTGG	GCCCCCCCCC	1320	
TGTGTTCCGG	YCTGGTGTGG	CGGGACGGTC	ATAGGAGGGG	CACCGTGC	GACCTCCCCC	1380	
CGGTTTGC	CCGGGAGGTT	CTCGGCACGG	TGACAGTCAT	GTGTCAGTGG	GGTTCTGCCT	1440	
ACTGGATTTG	GAGATTGGG	GA	CTGGGTTG	CATTGTACGA	CGAGCTACCA	CGATCAGCTC	1500
TCTGTACTTT	CTTCTCAGGT	CATGGTCCAC	AACCTAAAGA	TCTCTCAGTC	TTGAATCCAT	1560	
CCGGGGCACC	TTGTGCTTCT	TGCGTCGTTG	ACCAGAGGCC	GCTGAAATGT	GGTTCTGCG	1620	
TCCCGCAGTG	CTGGGAGACG	GGGGGT	GGTCGATGA	GTGCGGTGTC	GGTACTCGGA	1680	
TGACGAAGCA	CCTCGAGGCC	GTCCTGGTTG	ATGGAGGTGT	GGAGTCCAAG	GTGACAACGC	1740	
CCAAGGGTGA	GCGCCCCAAA	TACATAGGTC	AGCACGGTGT	GGGAACCTAC	TACGGCGCTG	1800	
TCCGTAGCCT	CAACATCAGT	TACCTAGTGA	CTGAGGTGGG	GGGCTATTGG	CATGCGCTGA	1860	
AGTGCCCGTG	CGACTTTGTG	CCCCGAGTGC	TCCCAGAAAG	AATTCCAGGT	AGGCCTGTGA	1920	
ATGCATGTCT	AGCTGGGAAG	TCTCCGCACC	CGTCGCAAG	TTGGGCTCCC	GGTGGGTTTT	1980	
ACGCCCCCGT	GTTCACCAAG	TGCAACTGGC	CGAAGACCTC	CGGAGTGGAT	GTGTGTCCTG	2040	
GGTTGCTTT	CGATTTCCCT	GGTGATCACA	ACGGCTTCAT	CCATGTTAAA	GGCAACAGAC	2100	
AGCAGGTTA	CAGTGGTCAG	CGAAGGTCTT	CGCCGGCTTG	GTTGCTTACT	GACATGGTCC	2160	
TGGCCCTGTT	GGTGGTGATG	AAGTTGGCTG	AGGCTAGAGT	TGTCCCCCTG	TTTATGCTGG	2220	
CAATGTGGTG	GTGGTTGAAT	GGAGCATCTG	CTGCCACTAT	TGTCA	TACATGGTCC	2280	
TCACGAAGTC	CACTGAAAGT	GTTCCATTGT	GGACTCCGCC	CACTGTTCCA	ACTCCATCTT	2340	
GCCCCAATT	TACCACCGGA	GTCGGGACT	CTACCTACAA	TGCTGGTTGC	TACATGGTGG	2400	
CAGGCCTGGC	GGCCGGGGCT	CAGGCGGTCT	GGGGTGCTGC	CAATGATGGT	GCTCAGGCCG	2460	
TCGTTGGTGG	CATCTGGCCC	GCGTGGCTCA	AGCTGCGAAG	CTTCGCTGCC	GGTCTGGCCT	2520	
GGTTGTAAA	TGTTGGGGCT	TACTTGCCGG	TCGTCGAGGC	CGCVCTGGCT	CCCGAGCTGG	2580	
TGTGCACCCC	GGTGGTCGGC	TGGGCAGCCC	AGGAGTGGTG	GTTCA	TGTCTGGGTG	2640	
TGATGTGTGT	CGTGGCGTAC	CTGAATGTCC	TGGGCTCTGT	RAGGGCTGCC	GTGCTTGTGG	2700	

CGATGCACCT	CGCAAGGGT	GCTCTGCCGC	TGGTATTGGT	GGTAGCTGCC	GGGGTRACCC	2760
GGGAGCGGCA	CAGCGCTTA	GGGCTTGAGG	TGTGCTTCGA	TCTGGATGGT	GGAGACTGGC	2820
CRGACGCCAG	TTGGTCTTGG	GGTTTAGCAG	GCGTGGTGAG	CTGGGCCCTC	CTGGTGGGGG	2880
GTCTGATGAC	CCACGGTGGC	CGATCAGCCA	GAYTGACTTG	GTAYGCCAGG	TGGGCCGTCA	2940
ATTAYCAGAG	GGTCGYCGG	TGGGTGAACA	ACTCACCGGT	TGGAGCYTTT	GGYCGTTGGM	3000
GGCGYGCCTG	GAAAGCYTGG	TTRGKGTGG	CTTGGTTCTT	CCCCCAGACA	GTTGCCACAG	3060
TYTCCGTCAT	CTTCATACTC	TGTTGAGCA	GTTTAGATGT	CATTGATTTC	ATCTTGGARG	3120
TACTCTTGGT	TAACTCACCA	AATCTCGCGC	GCTTGGCGCG	RGTGCTGGAC	TCCTTAGCTC	3180
THGCTGAGGA	GCGGCTGGCC	TGCTCTTGGC	TGGTGGCGT	CCTGCGCAAG	CGGGGGCGTCC	3240
TCCTCTACGA	GCACGCYGGT	CACACTAGCA	GGCGCGGTGC	TGCCCGCTTG	CGAGAGTGGG	3300
GYTTTGGCCT	YGAGCCKGTT	AGYATAACCA	AGGAAGATTG	YCYATTGTT	CGGGACTCTG	3360
CTCGTGTGTT	GGGCTGTGGA	CAATTGGTCC	ATGGGAAACC	AGTGGTCGCG	AGGCGAGGCG	3420
ACGAGGTGTT	GATCGGCTGT	GTGAACAGTC	GGTCGACCT	TCCGCCTGGC	TTTGTTCCTA	3480
CTGCTCCCGT	GGTSCTTCAT	CARGCWGGCA	ARGGRTTYTT	YGGGGTTGTG	AAGACMTCCA	3540
TGACAGGCAA	GGACCCGTCC	GAACACCACG	GRAACGTGGT	GGTCCTWGGG	ACTTCAACAA	3600
CKCGTTCCAT	GGGCTGCTGC	GTGAACGGAG	TAGTGTACAC	RACATACCAT	GGYACCAACG	3660
CCCGRCKAT	GGCGGGGCCK	TTTGGKCCYG	TCAAYGCTCG	GTGGTGGTCW	GCGAGYGACG	3720
ACGTCACGGT	YTACCCGCTC	CCWAATGGYG	CTTCTTGCCT	YCARGCWTGY	AAGTGCCAAC	3780
CAACTGGGGT	GTGGGTGATC	CGGAATGACG	GAGCTTTTG	CCATGGAAC	CTCGGCAAGG	3840
TGGTGGATT	AGATATGCC	GCTGAGTTGT	CAGACTTTCG	CGGGTCTTCT	GGATCACCAA	3900
TCTTGTGCGA	TGAGGGTCAT	GCTGTTGGCA	TGCTGATTTC	GGTGTTCAT	AGGGGGAGTA	3960
GGGTTTCCTC	GGTGCAGGTAT	ACCAAACCTT	GGGAAACTCT	CCCTCGGGAG	ATTGAGGCTC	4020
GATCGGAGGC	CCCCCCTGTG	CCAGGAACCA	CTGGATACAG	GGAGGCGCCA	CTGTTCCCTGC	4080
CCACCGGAGC	TGGCAAGTCG	ACGCGCGTGC	CGAATGAGTA	CGTCAAGGCT	GGACACAARG	4140
TGCTTGTACT	AAACCCATCC	ATTGCCACAG	TGAGGGCCAT	GGGCCCTTAC	ATGGAAAAGT	4200
TAACCGGCAA	ACATCCGTG	GTTACTGTG	GCCATGACAC	TACTGCATAT	TCCAGGACTA	4260
CTGACTCATC	TTTGACCTAC	TGTACATACG	GCAGGTTTAT	GGCCAATCCC	AGGAAATACT	4320
TGCGGGGGAA	CGACGTCGTA	ATTTGCGACG	AGTTGCACGT	CACCGACCCG	ACCTCAATT	4380
TGGGGATGGG	TCGGGCGAGG	TTACTCGCTC	GCGAGTGCAGG	CGTACGCCTC	CTGCTTTCG	4440

CTACGGCGAC CCCACCGGTC TCTCCGATGG CGAAGCATGA ATCTATTCA	4500
TGGGCAGTGA GGGGGAGGTC CCCTCTATT GCCAATTCCCT CCCACTGAGT AGGTATGCTA	4560
CTGGGAGACA CCTGCTGTTT TGTCAATTCCA AGGTAGARTG CACTAGGTTA TCCTCAGCTT	4620
TGGCCAGCTT TGGTGTCAAC ACCGTTGTGT ACTTCAGAGG CAAAGAAACT GACATTCCAA	4680
CTGGTGACGT GTGCCTTGC GCCACAGACG CACTTCCAC TGGTTACACT GGCAATTTG	4740
ACACCGTAAC AGACTGTGGT TTAATGGTTG AGGAGGTA	4800
ACACTACAT CGGTGTGAAG ACCGTCCCGG CCCCTGCCGA ACTGAGGGCT CAGAGGCGTG	4860
GTAGGTGTGG CCGTGGGAAA GCGGGCAC	4920
CGGGAACSGT TCGGTCTGGG GCTCTCTGGG CAGCTGTTGA GGCTGGHGTC TCGTGGTATG	4980
GCCTAGAGCC CGATGCTATT GGAGACCTGC TTAGGGCCTA CGACTCGTGT CCTTATACTG	5040
CTGCCATCAG TGCGTCCATC GGAGAGGCCA TTGCCTTTT TACTGGYCTA GTGCCAATGA	5100
GGAAATTATCC TCAGGTGGTT TGGGCAAGC AGAAGGGRCA CAACTGGCCA CTCTGGTGG	5160
GTGTGCAGAG GCACATGTGT GAGGACGCGG GCTGTGGTCC KCCCGCTAAT GGTCCCGAAT	5220
GGAGCGGCAT CAGGGGAAA GGGCCTGTTCC CCGATGGGT GGTGACTTGC	5280
CTGAGTCGGT GGCTCCGCAT CACTGGTTG ATGACCTACA GGCCCGGCTC GGTGTGGCCG	5340
AGGGTTACAC TCCCTGCATT GCTGGACCGG TGCTTTGGT CCGTTGGCG ATGGCGGGGG	5400
GGGCTATCCT GGCACACTGG ACGGGGTCTC TGGTTGTAGT GACCAGTTGG GTTGTCAATG	5460
GGAACGGTAA CCCGCTGATA CAAAGCGCCT CTAGGGCGT GGCKACYAGC GGTCCATACC	5520
CAGTACCCCC AGATGGTGGT GAACGGTACC CATCAGACAT CAAGCCAATY ACTGAGGCTG	5580
TGACCACCCCT TGAGACTGCG TGCGGYTGGG GCCCAGCCGC GGCBAGTCTG GCTTATGTGA	5640
AGGCCTGTGA AACTGGAACC ATGTTGGCTG ACAARGCGAG TGCTGCGTGG CAGGCTTGGG	5700
CTGCAAACAA CTTTGTGCCT CCACCAGCAT CACACTCAAC TTCCCTGTTR CAGAGCTTGG	5760
AYGCTGCGTT CACTTCAGCT TGGGATAGCG TGTTCACTCA CGGCCGTTCC TTGCTTGTG	5820
GGTTCACAGC TGCTTACGGC GCTCGGCAGA ACCCACCGCT GGGCGTCGGA GCCTCTTCT	5880
TGCTGGGCAT GTCATCGAGC CACYTRACTC ACGTCAGACT TGCTGCTGCG TTGCTCCTCG	5940
GCGTCGGGGG TACCGTCCCTA GGCACGCCCTG CTACTGGGCT TGCTATGGCG GGTGCCTACT	6000
TCGCKGGGGG CAGCGTTACC GCTAACTGGC TGAGTATCAT TGTGGCTCTA ATCGGAGGCT	6060
GGGAGGGGGC RGTKAACGCA GCCTCACTCA CCTTCGAYCT CCTGGCKGGG AAGTTACAAG	6120

CKAGYGAYGC TTGGTGCCR GTCA GYTGCY TGGCCTCTCC GGGGCTTCG GTGGCYGGTG	6180
TGGCDCTVGG YCTDYTGCTV TGGTCTGTCA ARAAGGGTGT GGGWCARGAY TGGGTTAACAA	6240
GAYTGTGAC GATGATGCCA CGCAGTTCGG TGATGCCTGA CGATTCTTC CTCAAAGATG	6300
AGTTCGTCAC CAAGGTGTCT ACTGTCCTGC GAAAGTTGTC ATTGTCAAGA TGGATCATGA	6360
CTCTTGTGGA CAAGCGGGAG ATGGAGATGG AGACMCCCCTC TTCTCAGATT GTTTGGGACT	6420
TGCTTGACTG GTGCATCCGG CTRGGTCGGT TCCTGTACAA TAAACTYATG TTTGCTCTCC	6480
CTAGGTTGCG CCTGCCGCTT ATCGGTTGCA GTACCGGTG GGGTGGCCCG TGGGAGGGCA	6540
ATGGTCATTT GGAAACAAGG TGTACTTGTG GCTGTGTGAT TACCGGTGAT ATTACAGATG	6600
GTATATTGCA CGACCTACAT TATA CCTCCC TACTGTGCAG ACATTACTAC AAGAGGACAG	6660
TGCCTGTTGG CGTCATGGGC AATGCTGAGG GAGCAGTCCC CCTTGTCGCT ACTGGCGGTG	6720
GAATCAGGAC TTACCAAATT GGGACTTCTG ACTGGTTGA GGCTGTGGTC GTGCATGGGA	6780
CAATCACGGT GCACGCCACC AGTTGCTATG AGTTGAAAGC TGCTGACGTT CGGAGGGCGG	6840
TGCGAGCCGG CCCGACTTAC GTTGGTGGCG TACCTGCAG CTGGAGCGCG CCGTGTACTG	6900
CGCCTGCGCT CGTTTACAGG CTAGGCCAGG GCATCAAAAT CGATGGAGCG CGCCGACTGT	6960
TGCCCTGTGA CTTAGCACAG GGAGCGCGCC ACCCCCCGGT ATCTGGCAGT GTGCCGGTA	7020
GTGGTTGGAC AGATGAGGAC GAGAGGGACT TGGTGGAAAC CAAGGCTGCC GCCATCGAGG	7080
CCATTGGGGC GGCTTGAC CTCCCTTCAC CGGAGGCTGC TCAGGCCGCT CTAGAGGCTT	7140
TGGAGGAGGC TGCCGTGTCC CTGTTGCCCGT CATTATGGGT GATGACTGTT	7200
CATGCCGGGA TGAGGCCTTC CAAGGCCACT TCATCCCAGA ACCCAATGTG ACAGAGGTAC	7260
CCATTGAGCC CACGGTCGGA GACGTGGAGG CACTCAAGCT GCGGGCTGCA GACCTGACCG	7320
CCAGGTTGCA AGACTTGGAG GCCATGGCTC TCGCCCGCGC TGAGTCAATC GAGGATGCTC	7380
GCGCAGCTTC GATGCCTTCG CTCACCGAGG TGGACTCAAT GCCATCATTG GAGTCGAGCC	7440
CTTGCTCCTC CTTTGAACAA ATCTCTTAA CTGAAAGTGA CCCTGAGACT GTCGTCGAGG	7500
CTGGCTTACC CTTGGAGTTC GTGAACTCCA ACACCGGGCC GTCTCCGGCT CGGAGGATTG	7560
TCAGAATCCG ACAGGCTTGC TGTTGTGACA GATCCACAAT GAAGGCCATG CCGTTGTCGT	7620
TCACTGTCGG GGAGTGCCCTC TTCGTTACTC GCTATGACCC GGACGGTCAC CAACTGTTG	7680
ACGAGCGAGG TCCGATAGAG GTATCTACTC CTATATGTGA AGTGATTGGG GACATCAGGC	7740
TTCAGTGTGA CCAAATTGAG GAAACTCCAA CATCTTACTC TTACATCTGG TCAGGGGCGC	7800
CCTTGGGTAC TGGGAGAAGT GTCCCCAAC CCATGACGCG CCCTATAGGG ACCCATCTGA	7860

CTTGTGACAC TACCAAAGTT TATGTTACTG ACCCTGATCG GGCGCTGAG CGGGCCGAGA	7920
AGGTTACAAT CTGGAGGGGT GATAGGAAGT ATGACAAGCA TTATGAGGCT GTCGTTGAGG	7980
CTGCTCTGAA AAAGGCAGCC GCGACGAAGT CTCATGGCTG GACCTATTCC CAGGCTATAG	8040
CTAAAGTTAG GCGCCGAGCA GCCGCTGGAT ACGGCAGCAA GGTGACCGCC TCCACATTGG	8100
CCACTGGTTG GCCTCACGTG GAGGAGATGC TGGACAAAAT AGCCAGGGGA CAGGAAGTTC	8160
CTTCACTTT TGTGACCAAG CGAGAGGTTT TCTTCTCCAA AACTACCCGT AAGCCCCAA	8220
GATTCAAGT TTTCCCACCT TTGGACTTCA GGATAGCTGA AAAGATGATT CTGGGTGACC	8280
CCGGCCTCGT TGCAAAGTCA ATTCTGGGTG ACGCTTATCT GTTCCAGTAC ACGCCCAATC	8340
AGAGGGTCAA AGCTCTGGTT AAGGCGTGGG AGGGGAAGTT GCATCCCGCT GCGATCACTG	8400
TGGACGCCAC TTGTTTCGAC TCATCGATTG ATGAGCACGA CATGCAGGTG GAGGCTTCGG	8460
TGTTTGCAGGC GGCTAGTGAC AACCCCTCAA TGGTACATGC TTTGTGCAAG TACTACTCTG	8520
GTGGCCCTAT GGTTTCCCCA GATGGGGTTC CCTTGGGGTA CCGCCAGTGT AGGTCGTCGG	8580
GCGTGTAAAC AACTAGCTCG GCGAACAGCA TCACCTGTTA CATTAAAGTC AGCGCGGCCT	8640
GCAGGGCGGGT GGGGATTAAG GCACCATCAT TCTTTATAGC TGGAGATGAT TGCTTGATCA	8700
TCTATGAAAAA TGATGGAACT GATCCCTGCC CTGCTCTTAA GGCTGCCCTG GCCAACTATG	8760
GATACAGGTG TGAACCAACA AAGCATGCTT CACTGGACAC AGCTGAGTGT TGCTCGGCCT	8820
ACTTGGCTGA GTGCGTAGCT GGGGGTGCCA AGCGCTGGTG GTTGAGCACG GACATGAGGA	8880
AGCCGCTCGC AAGGGCGTCT TCCGAATATT CGGACCCAAT CGGCAGTGCT TTAGGGACCA	8940
TCTTGATGTA TCCCCGGCAT CCAATCGTGC GGTATGTTCT AATACCACAC GTACTAATAA	9000
TGGCTTACAG GAGTGGCAGC ACACCGGATG AGTTGGTTAT GTGTCAGGTT CAGGGAAATC	9060
ATTACTCTTT CCCGCTGCAG CTGCTGCCCTC GCGTCTTGGT CTCTCTACAT GGTCCGTGGT	9120
GCCTACAAGT CACCACGGAC AGTACGAAGA CTAGGATGGA GGCAGGCTCA GCSTTGCAGG	9180
ATTTAGGAAT GAAATCCCTA GCCTGGCACC GCCGACGTGC CGGAAATGTG CGCACTCGCC	9240
TCCTGAGGGG AGGCAAGGAG TGGGGGCACC TGGCCAGAGC CCTCCTCTGG CAYCCAGGKT	9300
TGAAGGAGCA YCCCCCRCCC ATAAATTACAC TTCCAGGTTT TCAGCTGGCG ACGCCTTACG	9360
AACACCATGA AGAGGTCTTG ATCTCGATCA AGAGTCGACC ACCTTGGATA AGGTGGATTG	9420
TTGGTGCTTG TCTCTCGTTG CTGGCCGCCT TGCTGTGAAT TCGCTCCAGG CAGTAGGACC	9480
TTCCGGGTAG GGG	9493

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCATAATCAT GAGCCGCGAG TTGAAGAGCA C

31

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GCCAAGCCAT GGTGAATGTG

20

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 38 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GTATTGCGCC ATGGCTCGAC AAGCGTGGGT GGCCGGGG

38

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GGACTGCCAT GGTGGTATAG AAAAGAG

27

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

TATAATAAGC TTCTCGACAA GCGTGGGTGG CCGGGG

36

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GTATTGCGCC ATGGCACTGG GTGCAAGCCC AGAA

34

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 46 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Met Ser Val Val Asp Thr Phe Thr Met Ala Trp Leu Trp Leu Leu Val
5 10 15

Cys Phe Pro Leu Ala Gly Gly Val Leu Phe Asn Ser Arg His Gln Cys
20 25 30

Phe Asn Gly Asp His Tyr Val Leu Ser Asn Cys Cys Ser Arg
35 40 45

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 67 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Met Gly Pro Pro Ser Ser Ala Ala Ala Cys Ser Arg Gly Ser Pro Arg
5 10 15

Ile Leu Arg Val Arg Ala Gly Gly Ile Ser Leu Phe Tyr Thr Ile Met
20 25 30

Ala Val Leu Leu Leu Leu Val Val Glu Ala Gly Ala Ile Leu Ala
35 40 45

Pro Ala Thr His Ala Cys Arg Ala Asn Gly Gln Tyr Phe Leu Thr Asn
50 55 60

Cys Cys Ala
65

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9143 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

ACCACAAACA	CTCCAGTTG	TTACACTCCG	CTAGGAATGC	TCCTGGAGCA	CCCCCCCCTAG	60
CAGGGCGTGG	GGGATTTCCC	CTGCCCGTCT	GCAGAAGGGT	GGAGCCAACC	ACCTTAGTAT	120
GTAGGCGGCG	GGACTCATGA	CGCTCGCGT	ATGACAAGCG	CCAAGCTGA	CTTGGATGGC	180
CCTGATGGGC	GTTCATGGGT	TCGGTGGTGG	TGGCGTTTA	GGCAGCCTCC	ACGCCCACCA	240
CCTCCCAGAT	AGAGCGGCGG	CACTGTAGGG	AAGACCGGGG	ACCGGTCACT	ACCAAGGACG	300
CAGACCTCTT	TTTGAGTATC	ACGCCTCCGG	AAGTAGTTGG	GCAAGCCCAC	CTATATGTGT	360
TGGGATGGTT	GGGGTTAGCC	ATCCATACCG	TACTGCCTGA	TAGGGTCCTT	GCGAGGGAT	420
CTGGGAGTCT	CGTAGACCCT	AGCACATGCC	TGTTATTTCT	ACTCAAACAA	GTCCTGTACC	480
TGCGCCCAGA	ACGCGCAAGA	ACAAGCAGAC	GCAGGCTTCA	TATCCTGTGT	CCATTAAAAC	540
ATCTGTTGAA	AGGGGACAAC	GAGCAAAGCG	CAAAGTCCAG	CGCGATGCTC	GGCCTCGTAA	600
TTACAAAATT	GCTGGTATCC	ATGATGGCTT	GCAGACATTG	GCTCAGGCTG	CTTGCCAGC	660
TCATGGTTGG	GGACGCCAAG	ACCCTGCCA	TAAGTCTCGC	AATCTGGAA	TCCTTCTGGA	720
TTACCCTTG	GGGTGGATTG	GTGATGTTAC	AACTCACACA	CCTCTAGTAG	GCCCGCTGGT	780
GGCAGGAGCG	GTCGTTCGAC	CAGTCTGCCA	GATAGTACGC	TTGCTGGAGG	ATGGAGTCAA	840
CTGGGCTACT	GGTTGGTCG	GTGTCCACCT	TTTGTGGTA	TGTCTGCTAT	CTTGCCCTG	900
TCCCTGTAGT	GGGGCGCGGG	TCACTGACCC	AGACACAAAT	ACCACAATCC	TGACCAATTG	960
CTGCCAGCGT	AATCAGGTTA	TCTATTGTT	TCCTTCCACT	TGCCTACACG	AGCCTGGTTG	1020
TGTGATCTGC	GCGGACGAGT	GCTGGTTC	CGCCAATCCG	TACATCTCAC	ACCCTTCCAA	1080
TTGGACTGGC	ACGGACTCCT	TCTTGGCTGA	CCACATTGAT	TTTGTATGG	GCGCTTTGT	1140
GACCTGTGAC	GCCCTTGACA	TTGGTGAGTT	GTGTGGTGCG	TGTGTATTAG	TCGGTGACTG	1200
GCTTGTCAAG	CACTGGCTTA	TTCACATAGA	CCTCAATGAA	ACTGGTACTT	GTTACCTGGA	1260
AGTGCCCACT	GGAATAGATC	CTGGGTTCC	AGGGTTTATC	GGGTGGATGG	CCGGCAAGGT	1320
CGAGGCTGTC	ATCTTCTTGA	CCAAACTGGC	TTCACAAAGTA	CCATACGCTA	TTGCGACTAT	1380
GTTTAGCAGT	GTACACTACC	TGGCGGTGG	CGCTCTGATC	TACTATGCCT	CTCGGGGCAA	1440
GTGGTATCAG	TTGCTCCTAG	CGCTTATGCT	TTACATAGAA	GCGACCTCTG	GAAACCCTAT	1500
CAGGGTGCC	ACTGGATGCT	CAATAGCTGA	GTGTTGCTCG	CCTTGATGA	TACCATGTCC	1560
TTGCCACTCT	TATTTGAGTG	AGAATGTGTC	AGAAGTCATT	TGTTACAGTC	CAAAGTGGAC	1620

CAGGCCTGTC ACTCTAGAGT ATAACAACTC CATATCTGG TACCCCTATA CAATCCCTGG	1680
TGCGAGGGGA TGTATGGTTA AATTCAAAAA TAACACATGG GGTTGCTGCC GTATTCGCAA	1740
TGTGCCATCG TACTGCACTA TGGCACTGA TGCAGTGTGG AACGACACTC GCAACACTTA	1800
CGAAGCATGC GGTGTAACAC CATGGCTAAC AACCGCATGG CACAACGGCT CAGCCCTGAA	1860
ATTGGCTATA TTACAATACC CTGGGTCTAA AGAAATGTTT AAACCTCATA ATTGGATGTC	1920
AGGCCATTTG TATTTGAGG GATCAGATAC CCCTATAGTT TACTTTATG ACCCTGTGAA	1980
TTCCACTCTC CTACCACCGG AGAGGTGGC TAGGTTGCC GGTACCCCAC CTGTGGTACG	2040
TGGTTCTTGG TTACAGGTTT CGCAAGGGTT TTACAGTGAT GTGAAAGACC TAGCCACAGG	2100
ATTGATCACC AAAGACAAAG CCTGGAAAAA TTATCAGGTC TTATATTCCG CCACGGGTGC	2160
TTTGTCTCTT ACGGGAGTTA CCACCAAGGC CGTGGTGCTA ATTCTGTTGG GGTTGTGTGG	2220
CAGCAAGTAT CTTATTTAG CCTACCTCTG TTACTTGTC CTTTGTTTG GGCGCGCTTC	2280
TGGTTACCTT TTGCGTCTG TGCTCCCATC CCAGTCGTAT CTCCAAGCTG GCTGGGATGT	2340
TTTGTCTAAA GCTCAAGTAG CTCCTTTGC TTTGATTTTC TTCATCTGTT GCTATCTCCG	2400
CTGCAGGCTA CGTTATGCTG CCCTTTAGG TTTGTGCC ATGGCTGCGG GCTTGCCCC	2460
AACTTTCTTT GTGCAGCAG CTGCTGCCA ACCAGATTAT GACTGGTGGG TGCGACTGCT	2520
AGTGGCAGGG TTAGTTTGT GGGCCGGCCG TGACCGTGGT CCACGTATAG CTCTGCTTGT	2580
AGGTCTTGG CCTCTGGTAG CGCTTTAAC CCTCTTGCT TTGGCTACGC CTGCTTCAGC	2640
TTTGACACC GAGATAATTG GAGGGCTGAC AATACCACCT GTAGTAGCAT TAGTTGTCAT	2700
GTCTCGTTTT GGCTTCTTG CTCACTGTT ACCTCGCTGT GCTTAGTTA ACTCCTATCT	2760
TTGGCAACGT TGGGAGAATT GGTTTGGAA CGTTACACTA AGACCGGAGA GGTTTCTCCT	2820
TGTGCTGGTT TGTTTCCCCG GTGCGACATA TGACACGCTG GTGACTTTCT GTGTGTGTCA	2880
CGTAGCTCTT CTATGTTAA CATCAGTGC AGCATCGTTC TTTGGACTG ACTCTAGGGT	2940
TAGGGCCCAT AGAATGTTGG TGCGTCTCGG AAAGTGTCA GCTTGGTATT CTCATTATGT	3000
TCTTAAGTTT TTCCTCTTAG TGTTTGGTGA GAATGGTGTG TTTTCTATA AGCACTTGCA	3060
TGGTGATGTC TTGCCTAATG ATTTGCCTC GAAACTACCA TTGCAAGAGC CATTTCCTCC	3120
TTTTGAAGGC AAGGCAAGGG TCTATAGGAA TGAAGGAAGA CGCTTGGCGT GTGGGGACAC	3180
GGTTGATGGT TTGCCCGTTG TTGCGCGTCT CGGCGACCTT GTTTGCAG GGTTAGCTAT	3240
GCCGCCAGAT GGGTGGGCCA TTACCGCACC TTTTACGCTG CAGTGTCTCT CTGAACGTGG	3300
CACGCTGTCA GCGATGGCAG TGGTCATGAC TGGTATAGAC CCCCGAAGTT GGACTGGAAC	3360

TATCTTCAGA TTAGGATCTC TGGCCACTAG CTACATGGGA TTTGTTTGTG ACAACGTGTT	3420
GTATACTGCT CACCATGGCA GCAAGGGGCG CCGGTTGGCT CATCCACAG GCTCCATACA	3480
CCCAATAACC GTTGACGCGG CTAATGACCA GGACATCTAT CAACCACCAT GTGGAGCTGG	3540
GTCCCTTACT CGGTGCTCTT GCGGGGAGAC CAAGGGGTAT CTGGTAACAC GACTGGGGTC	3600
ATTGGTTGAG GTCAACAAAT CCGATGACCC TTATTGGTGT GTGTGCGGGG CCCTTCCCCT	3660
GGCTGTTGCC AAGGGTTCTT CAGGTGCCCG GATTCTGTGC TCCTCCGGGC ATGTTATTGG	3720
GATGTTCAC C GCTGCTAGAA ATTCTGGCGG TTCAGTCAGC CAGATTAGGG TTAGGCCGTT	3780
GGTGTGTGCT GGATACCATC CCCAGTACAC AGCACATGCC ACTCTTGATA CAAAACCTAC	3840
TGTGCCTAAC GAGTATTCAG TGCAAATTTT AATTGCCCGG ACTGGCAGCG GCAAGTCAAC	3900
CAAATTACCA CTTTCTTACA TGCAAGGAGAA GTATGAGGTC TTGGTCCTAA ATCCCAGTGT	3960
GGCTACAACA GCATCAATGC CAAAGTACAT GCACGCGACG TACGGCGTGA ATCCAAATTG	4020
CTATTTTAAT GGCAAATGTA CCAACACAGG GGCTTCACTT ACGTACAGCA CATATGGCAT	4080
GTACCTGACC GGAGCATGTT CCCGGAACTA TGACGTCATC ATTTGTGACG AATGCCATGC	4140
TACCGATGCA ACCACCGTGT TGGGCATTGG AAAGGTTCTA ACCGAAGCTC CATCCAAAAA	4200
TGTTAGGCTA GTGGTTCTTG CCACGGCTAC CCCCCCTGGGA GTAATCCCTA CACCACATGC	4260
CAACATAACT GAGATTCAAT TAACCGATGA AGGCACATAC CCCTTCATG GAAAAAAGAT	4320
TAAGGAGGAA AATCTGAAGA AAGGGAGACA CCTTATCTTT GAGGCTACCA AAAAACACTG	4380
TGATGAGCTT GCTAACGAGT TAGCTCGAAA GGGATAACA GCTGTCCTT ACTATAGGGG	4440
ATGTGACATC TCAAAAATCC CTGAGGGCGA CTGTGTAGTA GTGCCACTG ATGCCCTTGTG	4500
TACAGGGTAC ACTGGTGACT TTGATTCCGT GTATGACTGC AGCCTCATGG TAGAAGGCAC	4560
ATGCCATGTT GACCTTGACC CTACTTTCAC CATGGGTGTT CGTGTGTGCG GGGCTCAGC	4620
AATAGTTAAA GGCCAGCGTA GGGCCCGCAC AGGCCGTGGG AGAGCTGGCA TATACTACTA	4680
TGTAGACGGG AGTTGTACCC CTTGGGTAT GGTTCTGAA TGCAACATTG TTGAAGCCTT	4740
CGACGCAGCC AAGGCATGGT ATGGTTGTC ATCAACAGAA GCTCAAACCA TTCTGGACAC	4800
CTATCGCACC CAACCTGGGT TACCTGCGAT AGGAGCAAAT TTGGACGAGT GGGCTGATCT	4860
CTTTTCTATG GTCAACCCCG AACCTTCATT TGTCAATACT GCAAAAAGAA CTGCTGACAA	4920
TTATGTTTG TTGACTGCAG CCCAACTACA ACTGTGTGTCAGTATGGCT ATGCTGCTCC	4980
CAATGACGCA CCACGGTGGC AGGGAGCCCG GCTTGGAAA AAACCTTGTG GGGTTCTGTG	5040

GCGCTTGGAC GGCGCTGACG CCTGTCCTGG CCCAGAGCCC AGCGAGGTGA CCAGATACCA	5100
AATGTGCTTC ACTGAAGTCA ATACTTCTGG GACAGCCGCA CTCGCTGTTG GCGTTGGAGT	5160
GGCTATGGCT TATCTAGCCA TTGACACTTT TGGCGCCACT TGTGTGCGGC GTTGTGGTC	5220
TATTACATCA GTCCCTACCG GTGCTACTGT CGCCCCAGTG GTTGACGAAG AAGAAATCGT	5280
GGAGGAGTGT GCATCATTCA TTCCCTTGGGA GGCCATGGTT GCTGCAATCG ATAAGCTGAA	5340
GAGTACAATA ACCACAAACTA GTCCCTTCAC ATTGGAAACC GCCCTTGAAA AACTTAACAC	5400
CTTTCTTGGG CCTCATGCAG CTACAATCCT TGCTATCATA GAGTATTGCT GTGGCTTAGT	5460
CACTTTACCT GACAATCCCT TTGCATCATG CGTGTGTTGCT TTCATTGCGG GTATTACTAC	5520
CCCACTACCT CACAAGATCA AAATGTTCCCT GTCATTATTGTT GGAGGGCAA TTGCGTCAA	5580
GCTTACAGAC GCTAGAGGCG CACTGGCGTT CATGATGGCC GGGGCTGCGG GAACAGCTCT	5640
TGGTACATGG ACATCGGTGG GTTTTGTCTT TGACATGCTA GGCGGCTATG CTGCCGCCTC	5700
ATCCACTGCT TGCTTGACAT TTAAATGCTT GATGGGTGAG TGGCCCACTA TGGATCAGCT	5760
TGCTGGTTA GTCTACTCCG CGTTCAATCC GGCGCAGGA GTTGTGGCG TCTTGTCA	5820
TTGTGCAATG TTTGCTTGA CAACAGCAGG GCCAGATCAC TGGCCCAACA GACTTCTTAC	5880
TATGCTTGCT AGGAGCAACA CTGTATGTAA TGAGTACTTT ATTGCCACTC GTGACATCCG	5940
CAGGAAGATA CTGGGCATTC TGGAGGCATC TACCCCCCTGG AGTGTCAATAT CAGCTTGCAT	6000
CCGTTGGCTC CACACCCCGA CGGAGGATGA TTGCGGCCTC ATTGCTTGGG GTCTAGAGAT	6060
TTGGCAGTAT GTGTGCAATT TCTTTGTGAT TTGCTTAAAT GTCCCTAAAG CTGGAGTTCA	6120
GAGCATGGTT AACATTCTG GTTGTCTTT CTACAGCTGC CAGAAGGGGT ACAAGGGCCC	6180
CTGGATTGGA TCAGGTATGC TCCAAGCAGC CTGTCCATGC GGTGCTGAAC TCATCTTTTC	6240
TGTTGAGAAT GTTTTGCAA AACTTTACAA AGGACCCAGA ACTTGTCAA ATTACTGGAG	6300
AGGGGCTGTT CCAGTCAACG CTAGGCTGTG TGGGTCGGCT AGACCGGACC CAACTGATTG	6360
GACTAGTCTT GTCGTCAATT ATGGCGTTAG GGACTACTGT AAATATGAGA AATTGGGAGA	6420
TCACATTTTT GTTACAGCAG TATCCTCTCC AAATGTCTGT TTCACCCAGG TGCCCCAAC	6480
CTTGAGAGCT GCAGTGGCCG TGGACGGCGT ACAGGTTCAAG TGTTATCTAG GTGAGCCAA	6540
AACTCCTTGG ACGACATCTG CTTGCTGTTA CGGTCCGGAC GGTAAGGGTA AAACGTAA	6600
GCTTCCCTTC CGCGTTGACG GTCACACACC TGGTGTGCGC ATGCAACTTA ATTTGCGTGA	6660
TGCACTTGAG ACAAAATGACT GTAATTCCAT AAACAACACT CCTAGTGATG AAGCCGAGT	6720
GTCCGCTCTT GTTTCAAAC AGGAGTTGCG GCGTACAAAC CAATTGCTTG AGGCAATTTC	6780

AGCTGGCGTT GACACCACCA AACTGCCAGC CCCCTCCATC GAAGAGGTAG TGGTAAGAAA	6840
GCGCCAGTTC CGGGCAAGAA CTGGTTCGCT TACCTTGCT CCCCTCCGA GATCCGTCCC	6900
AGGAGTGTCA TGTCTGAAA GCCTGCAACG AAGTGACCCG TTAGAAGGTC CTTCAAACCT	6960
CCCTTCTTCA CCACCTGTT TACAGTTGGC CATGCCGATG CCCCTGTTGG GAGCAGGTGA	7020
GTGTAACCTT TTCACTGCAA TTGGATGTGC AATGACCGAA ACAGGCGGAG GCCCTGATGA	7080
TTTACCCAGT TACCCCTCCA AAAAGGAGGT CTCTGAATGG TCAGACGGAA GTTGGTCAAC	7140
GAATACAACC GCTTCCAGCT ACCTTACTGG CCCCCCGTAC CCTAAGATAC GGGGAAAGGA	7200
TTCCACTCAG TCAGCCCCCG CCAAACGGCC TACAAAAAAG AAGTTGGAA AGAGTGAGTT	7260
TTCGTGCAGC ATGAGCTACA CTTGGACCGA CGTGATTAGC TTCAAAACTG CTTCTAAAGT	7320
TCTGTCTGCA ACTCGGGCCA TCACTAGTGG TTTCTCAAA CAAAGATCAT TGGTGTATGT	7380
GAATGAGCCG CGGGATGCGG AGCTTAGAAA ACAAAAGTC ACTATTAATA GACAACCTCT	7440
GTTCCCCCA TCATACCACA AGCAAGTGAG ATTGGCTAAG GAAAAAGCTT CAAAGTTGT	7500
CGGTGTCACTG TGGGACTATG ATGAAGTAGC AGCTCACACG CCCTCTAAGT CTGCTAAGTC	7560
CCACATCACT GGCCTTCGGG GCACTGATGT TCGTTCTGGA GCAGCCCGCA AGGCTGTTCT	7620
GGACTTGCAG AAGTGTGTCG AGGCAGGTGA GATACCGAGT CATTATCGGC AAACTGTGAT	7680
AGTTCCAAAG GAGGAGGTCT TCGTGAAGAC CCCCCAGAAA CCAACAAAGA AACCCCCAAG	7740
GCTTATCTCG TACCCCCACC TTGAAATGAG ATGTGTTGAG AAGATGTACT ACGGTCAGGT	7800
TGCTCCTGAC GTAGTTAAAG CTGTCATGGG AGATGCGTAC GGGTTGTCG ACCCACGTAC	7860
CCGTGTCAAG CGTCTGTTGT CGATGTGGTC ACCCGATGCA GTGGAGGCCA CATGCGATAC	7920
AGTGTGTTT GACAGTACCA TCACACCGA GGATATCATG GTGGAGACAG ACATCTACTC	7980
AGCAGCTAAA CTCAGTGACC AACACCGAGC TGGCATTAC ACCATTGCGA GGCAGTTATA	8040
CGCTGGAGGA CCGATGATCG CTTATGATGG CCGAGAGATC GGATATCGTA GGTGTAGGTC	8100
TTCCGGCGTC TATACTACCT CAAGTTCCAA CAGTTGACC TGCTGGCTGA AGGTAAATGC	8160
TGCAGCCGAA CAGGCTGGCA TGAAGAACCC TCGCTTCCTT ATTTGCGGCG ATGATTGCAC	8220
CGTAATTGGG AAGAGCGCCG GAGCAGATGC AGACAAACAA GCAATGCGTG TCTTGCTAG	8280
CTGGATGAAG GTGATGGGTG CACCACAAGA TTGTGTGCCT CAACCCAAAT ACAGTTGGAA	8340
AGAATTAACA TCATGCTCAT CAAATGTTAC CTCTGGAATT ACCAAAAGTG GCAAGCCTTA	8400
CTACTTTCTT ACAAGAGATC CTCGTATCCC CCTTGGCAGG TGCTCTGCCG AGGGTCTGGG	8460

ATACAACCCC AGTGCTGCGT GGATTGGGT A TCTAATACAT CACTACCCAT GTTTGTGGGT	8520
TAGCCGTGTG TTGGCTGTCC ATTTCATGGA GCAGATGCTC TTTGAGGACA AACTTCCCGA	8580
GA C TGTGACC TTTGACTGGT ATGGGAAAAA TTATACGGTG CCTGTAGAAG ATCTGCCAG	8640
CATCATTGCT GGTGTGCACG GTATTGAGGC TTTCTCGGTG GTGCCTACA CCAACGCTGA	8700
GATCCTCAGA GTTTCCAAT CACTAACAGA CATGACCATG CCCCCCTGC GAGCCTGGCG	8760
AAAGAAAGCC AGGGCGGTCC TCGCCAGCGC CAAGAGGC GT GGCGGAGCAC ACGAAAATT	8820
GGCTCGCTTC CTTCTCTGGC ATGCTACATC TAGACCTCTA CCAGATTGG ATAAGACGAG	8880
CGTGGCTCGG TACACCAC TT TCAATTATTG TGATGTTAC TCCCCGGAGG GGGATGTGTT	8940
TGTTACACCA CAGAGAAGAT TGCAGAAGTT TCTTGTGAAG TATTGGCTG TCATTGTTTT	9000
TGCCCTAGGG CTCATTGCTG TTGGACTAGC CATCAGCTGA ACCCCCAAAT TCAAAATTAA	9060
TTAACAGTTT TTTTTTTTTT TTTTTTTTTT TTTTAGGGCA GCGGCAACAG GGGAGACCCC	9120
GGGCTTAACG ACCCCGCGAT GTG	9143

WHAT IS CLAIMED IS:


1. A method for controlling the translation of HGBV nucleic acids to HGBV proteins, comprising
 - a. contacting a first non-naturally occurring nucleic acid sequence with HGBV nucleic acid sequence under conditions which permit hybridization of the first nucleic acid sequence and the HGBV nucleic acid sequence, and
 - b. altering the level of translation of the HGBV nucleic acid.
2. The method of claim 1 wherein said first nucleic acid sequence is an antisense nucleic acid sequence which is substantially complementary to a sequence of the sense strand within the 5' NTR region of the HGBV nucleic acid sequence.
3. The method of claim 1 wherein said first nucleic acid is a nucleic acid analog.
4. The method of claim 3 wherein said nucleic acid analog is selected from the group consisting of a morpholino compound, a peptide nucleic analog and a phosphorothioate nucleic acid analog.
5. A method of enhancing the translation of a nucleic acid comprising operably linking a nucleic acid with a nucleic acid having a sequence corresponding to the sequence of GBV-A, -B or -C 5' region, to form a combined nucleic acid capable of being translated.
6. A composition for enhancing the translation of a nucleic acid, which composition comprises a nucleic acid having a sequence corresponding to the sequence of GBV-A, -B, or -C 5' region, for operable linkage to nucleic acid to be translated.
7. A composition for controlling translation of hepatitis GB virus -A, -B, or -C from GBvirus -A, -B or -C nucleic acid, which comprises a first non-naturally occurring nucleic acid having a sequence complementary to, or capable of being transcribed to form, a nucleic acid having a sequence complementary to, a sequence of the sense strand within the 5'-NTR region of HGBV-A, -B, or -C, wherein said first nucleic acid comprises a sequence selected from the 5' NTR

region of GBV-A, -B, or -C, and a cleavage area at which the full length GBV-A, -B, or -C RNA is cleaved to form a subgenomic HGBV-A, -B, or -C RNA.

8. The composition of claim 7 wherein said first nucleic acid is be a nucleic acid analog.

9. The composition of claim 8 wherein said nucleic acid analog is selected from the group consisting of morpholino compounds, peptide nucleic analogs and a phosphorothioate nucleic acid analog

10. The composition of claim 7 wherein said first nucleic acid is linked to a cholesteryl moiety at the 3' end.



SUBSTITUTE SHEET (RULE 26)

2 / 13




FIG. 2A

3 / 13




FIG. 2B

4 / 13




FIG. 3A

5 / 13




FIG.3B

6/13




FIG. 4B




FIG. 4A

SUBSTITUTE SHEET (RULE 26)

7/13




FIG. 5A

8/13




FIG.5B

9 / 13




FIG. 6A

10 / 13






FIG.6B



11 / 13

FIG.7A



CONTINUED ON FIG.7B

CONTINUED ON FIG. 7B

SUBSTITUTE SHEET (RULE 26)





FIG.7D

FIG.7C
SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/13198

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/86 C12N15/11 C12N15/40 C12N15/51 C07K14/18
//C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,94 08002 (CHIRON CORP) 14 April 1994 see page 3, line 4 - page 4 see page 6, line 12 - page 12 see examples 8-10 see claims ---	1-10
Y	EMBL DATABASE ENTRY HG22303; ACCESSION NUMBER U22303 , 19 May 1995, XP002023170 SIMONS, J. ET AL.: "Identification of two flavivirus-like genomes in the GB hepatitis agent" see abstract & PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 92, 11 April 1995, WASHINGTON US, pages 3401-3405, cited in the application --- -/-	1-10

 Further documents are listed in the continuation of box C. Patent family members are listed in annex.

° Special categories of cited documents :

- °A° document defining the general state of the art which is not considered to be of particular relevance
- °E° earlier document but published on or after the international filing date
- °L° document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- °O° document referring to an oral disclosure, use, exhibition or other means
- °P° document published prior to the international filing date but later than the priority date claimed

- °T° later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- °X° document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- °Y° document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- °&° document member of the same patent family

Date of the actual completion of the international search 21 January 1997	Date of mailing of the international search report 24.01.97
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016	Authorized officer Andres, S

INTERNATIONAL SEARCH REPORT

International Application No PCT/US 96/13198

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EMBL DATABASE ENTRY HG22304; ACCESSION NUMBER U22304, 19 May 1995, XP002023171 SIMONS, J. ET AL.: "Identification of two flavivirus-like genomes in the GB hepatitis agent" see abstract & PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 92, 11 April 1995, WASHINGTON US, pages 3401-3405, cited in the application ---	1-10
A	WO,A,94 18217 (ABBOTT LAB ;TASSOPOULOS NIC C (GR); HATZAKIS ANGELOS E (GR); TROON) 18 August 1994 see the whole document ---	1-10
P,X	WO,A,95 32291 (GENELABS TECH INC) 30 November 1995 see page 71, line 17 - page 72, line 16 ---	1-3,7,8
T	JOURNAL OF VIROLOGICAL METHODS, (OCT 1996) VOL. 62, NO. 1, PP. 55-62., XP000616255 MUERHOFF, A. ET AL.: "IDENTIFICATION OF CONSERVED NUCLEOTIDE-SEQUENCES WITHIN THE GB VIRUS-C 5'-UNTRANSLATED REGION - DESIGN OF PCR PRIMERS FOR DETECTION OF VIRAL-RNA" see the whole document -----	1-5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/13198

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: 1-5
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-5 (as far as in vivo methods are concerned) are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/13198

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9408002	14-04-94	AU-A-	5141993	26-04-94
		CA-A-	2145290	14-04-94
		CA-A-	2158427	29-03-94
		EP-A-	0662128	12-07-95
		EP-A-	0718400	26-06-96
		JP-T-	8502167	12-03-96
-----	-----	-----	-----	-----
WO-A-9418217	18-08-94	AU-A-	3606193	29-08-94
-----	-----	-----	-----	-----
WO-A-9532291	30-11-95	AU-A-	2594195	18-12-95
		AU-A-	2689595	18-12-95
		WO-A-	9532292	30-11-95
-----	-----	-----	-----	-----