



(86) Date de dépôt PCT/PCT Filing Date: 2004/10/18
(87) Date publication PCT/PCT Publication Date: 2005/05/12
(45) Date de délivrance/Issue Date: 2015/04/14
(85) Entrée phase nationale/National Entry: 2006/04/21
(86) N° demande PCT/PCT Application No.: US 2004/034367
(87) N° publication PCT/PCT Publication No.: 2005/041937
(30) Priorité/Priority: 2003/10/23 (US60/513,618)

(51) Cl.Int./Int.Cl. *A61K 9/19* (2006.01),
A61K 31/496 (2006.01), *A61K 47/26* (2006.01),
A61K 47/38 (2006.01), *A61P 25/18* (2006.01)

(72) Inventeurs/Inventors:
KOSTANSKI, JANUSZ W., US;
MATSUDA, TAKAKUNI, JP;
NERURKAR, MANOJ, US;
NARINGREKAR, VIJAY H., US


(73) Propriétaire/Owner:
OTSUKA PHARMACEUTICAL CO., LTD., JP

(74) Agent: SMART & BIGGAR

(54) Titre : FORMULATION D'ARIPIRAZOLE LYOPHILISEE STERILE A LIBERATION CONTROLEE ET FORMULATION
INJECTABLE DE CELLE-CI
(54) Title: CONTROLLED RELEASE STERILE FREEZE-DRIED ARIPIRAZOLE FORMULATION AND INJECTABLE
FORMULATION THEREOF

(57) **Abrégé/Abstract:**

A controlled release sterile freeze-dried aripiprazole formulation is provided which is formed of aripiprazole of a desired mean particle size and a vehicle therefore, which upon constitution with water and intramuscular injection releases aripiprazole over a period of at least about one week and up to about eight weeks. A method for preparing the controlled release freeze-dried aripiprazole formulation is also provided.



75535-6

Abstract:

A controlled release sterile freeze-dried aripiprazole formulation is provided which is formed of aripiprazole of a desired mean particle size and a vehicle therefore, which upon constitution with water and intramuscular injection releases aripiprazole over a period of at least about one week and up to about eight weeks. A method for preparing the controlled release freeze-dried aripiprazole formulation is also provided.

75535-6

CONTROLLED RELEASE STERILE FREEZE-DRIED ARIPIPRAZOLE FORMULATION AND INJECTABLE FORMULATION THEREOF


FIELD OF THE INVENTION

The present invention relates to a controlled release sterile freeze-dried aripiprazole formulation, an injectable formulation which contains the sterile freeze-dried aripiprazole and which releases aripiprazole over at least a one week period, and a method for preparing the above formulation.

BACKGROUND OF THE INVENTION

U.S. Patent No. 5,006,528 to Oshiro et al. discloses 7-[(4-phenylpiperazino)-butoxy] carbostyrils, which include aripiprazole, as dopaminergic neurotransmitter antagonists.

Aripiprazole which has the structure



is an atypical antipsychotic agent useful in treating schizophrenia. It has poor aqueous solubility ($<1\mu\text{g/mL}$ at room temperature).

U.S. Patent No. 6,267,989 to Liversidge, et al. discloses a method for preventing crystal growth and particle aggregation in nanoparticulate compositions wherein a nanoparticulate composition is reduced to an optimal effective average particle size employing aqueous milling techniques including ball milling.

U.S. Patent No. 5,314,506 to Midler, et al. relates to a process for the direct crystallization of a pharmaceutical having high surface area particles of high purity and stability wherein impinging jet streams are employed to achieve high intensity

75535-6

micromixing of particles of the pharmaceutical followed by nucleation and direct production of small crystals.

A long-acting aripiprazole sterile injectable formulation has merit as a drug dosage form in that it may increase the compliance of patients and thereby lower the rate of relapse in the treatment of schizophrenia. Examples of known long acting drug products for the treatment of schizophrenia include haloperidol decanoate and fluphenazine decanoate both of which have an ester compound of low water solubility dissolved in sesame oil. Microcapsules containing Risperidone (WO95/13814) and Olanzapine (WO99/12549) are also known.

10

BRIEF DESCRIPTION OF THE INVENTION

In accordance with the present invention, there is provided a sterile freeze-dried aripiprazole formulation which upon constitution with water for injection releases aripiprazole over a period of at least about one week, and preferably over a period of two, three or four weeks and up to six weeks or more.

15

In an embodiment, the invention relates to a lyophilized formulation for release of aripiprazole over a period of about two to about four weeks, which comprises: (a) aripiprazole monohydrate having a mean particle size of 1 to 10 microns; (b) carboxymethylcellulose or its sodium-salt, hydroxypropyl cellulose, hydroxypropylethyl cellulose, hydroxypropylmethyl cellulose, or polyvinylpyrrolidone; (c) mannitol, sucrose, maltose, lactose, xylitol or sorbitol; and (d) sodium phosphate, potassium phosphate or TRIS buffer.

20

The freeze-dried aripiprazole formulation of the invention includes:

(a) aripiprazole, and

(b) a vehicle for the aripiprazole,

75535-6

which formulation upon reconstitution with water forms an injectable suspension which, upon injection, preferably intramuscularly, releases aripiprazole over a period of at least one week, preferably two, three or four weeks, and up to six weeks or more.

In one embodiment, the invention relates to a sterile freeze-dried controlled
5 release aripiprazole formulation which comprises:

(a) aripiprazole having a mean particle size of 1 to 10 microns, and

(b) a vehicle therefore,

which formulation upon reconstitution with water forms a sterile injectable formulation in the
form of a sterile homogeneous suspension which upon injection releases aripiprazole over a
10 period of at least two weeks.

The freeze-dried aripiprazole formulation of the invention will preferably
include:

(a) aripiprazole,

(b) one or more suspending agents,

15 (c) optionally one or more bulking agents,

(d) optionally one or more buffers, and

(e) optionally one or more pH adjusting agents.

In another embodiment, the invention relates to a sterile freeze-dried controlled
release aripiprazole formulation, which comprises:

20 (a) aripiprazole having a mean particle size of 1 to 10 microns, and

(b) one or more suspending agents,

75535-6

wherein said sterile freeze-dried controlled release aripiprazole formulation is obtained by freeze-drying an aqueous suspension comprising (a) and (b), and

wherein upon reconstitution, which comprises adding water to the freeze-dried formulation, a sterile injectable formulation is obtained which is a homogeneous suspension and which contains from 0.1 to 600 mg of aripiprazole per mL, and wherein upon injection into a subject, said sterile injectable formulation releases aripiprazole over a period of at least two weeks from the date of administration.

A mean particle size of the freeze-dried aripiprazole formulation within the range from about 1 to about 30 microns is essential in formulating an injectable which releases aripiprazole over a period of at least about one week and up to six weeks or more, for example up to 8 weeks.

It has been found that the smaller the mean particle size of the freeze-dried aripiprazole, the shorter the period of extended release. Thus, in accordance with the present invention, when the mean particle size is about 1 micron, the aripiprazole will be released over a period of less than three weeks, preferably about two weeks. When the mean particle size is more than about 1 micron, the aripiprazole will be released over a period of at least two weeks, preferably about three to four weeks, and up to six weeks or more. Thus, in accordance with the present invention, the aripiprazole release duration can be modified by changing the particle size of the aripiprazole in the freeze-dried formation.

The term "mean particle size" refers to volume mean diameter as measured by laser-light scattering (LLS) methods. Particle size distribution is measured by LLS methods and mean particle size is calculated from the particle size distribution.

In addition, in accordance with the present invention, a controlled release sterile injectable aripiprazole formulation in the form of a sterile suspension, that is, the freeze-dried formulation of the invention suspended in water for injection, is provided which, upon injection, preferably intramuscularly, releases aripiprazole over a period of at least one week, which includes:

- (a) aripiprazole,
- (b) a vehicle therefor, and
- (c) water for injection.

The controlled release sterile injectable formulation of the invention in the form of a sterile suspension allows for high drug loadings per unit volume of the formulation and therefore permits delivery of relatively high doses of aripiprazole in a small injection volume (0.1 – 600 mg of drug per 1 mL of suspension).

75535-6

In another embodiment, the invention relates to a controlled release sterile aripiprazole injectable formulation in the form of a sterile homogeneous suspension which upon injection releases aripiprazole over a period of at least two weeks, which comprises: (a) aripiprazole having a mean particle size of 1 to 10 microns in an amount of 100 to 400 mg per mL, (b) a vehicle therefore, and (c) water for injection.

5 Further, in accordance with the present invention, a method is provided for preparing the sterile freeze-dried aripiprazole formulation described above which includes the steps of:

(a) preparing sterile bulk aripiprazole, preferably having a desired particle size distribution and mean particle size within the range from about 5 to about 100 microns,

(b) preparing a sterile vehicle for the sterile bulk aripiprazole,

(c) combining the sterile bulk aripiprazole and the sterile vehicle to form a sterile primary suspension,

5 (d) reducing the mean particle size of aripiprazole in the sterile primary suspension to within the range from about 1 to about 30 microns, to form a final sterile suspension, and

(e) freeze drying the final sterile suspension to form a sterile freeze-dried suspension of the aripiprazole of desired polymorphic form (anhydrous, monohydrate, or a mixture of both).

10 In carrying out the above method, the reduction of the mean particle size of the sterile primary suspension to the desired mean particle size is preferably carried out employing an aseptic wet milling procedure, which more preferably is aseptic wet ball milling. Aseptic wet milling is preferred in forming a homogeneous and sterile aripiprazole formulation of desired mean particle size distribution.

15 In addition, in accordance with the present invention, a method for freeze drying the final sterile suspension of aripiprazole is provided which produces a sterile freeze-dried aripiprazole of desired polymorphic form, that is anhydrous, monohydrate, or a mixture of both.

20 As an unexpected observation, it has been discovered that a suspension of aripiprazole suspended in an aqueous solvent system will maintain a substantially
25 constant aripiprazole drug plasma concentration when administered by injection; preferably as an intra-muscular injection. No large "burst phenomenon" is observed and it is considerably surprising that a constant aripiprazole drug plasma concentration can be maintained from one (1) to more than eight (8) weeks employing the aripiprazole suspension of the invention. The daily starting dose for an orally
30 administered aripiprazole formulation is fifteen (15) milligrams. In order to administer a drug dose equivalent to one (1) to more than eight (8) weeks of the oral dosage quantity requires the administration of a very large amount of the drug as a

75535-6

single dose. The aqueous aripiprazole injectable formulation of the invention may be administered to deliver large amounts of the drug without creating patient compliance problems.

The aripiprazole injectable formulation of the invention may include
5 anhydrous or monohydrate crystalline forms of aripiprazole or an admixture containing both. If the monohydrate is used, the maintenance of an extended drug plasma concentration is possible.

The aripiprazole injectable formulation of the invention can be administered as an aqueous ready-to-use suspension; however, by freeze-drying this suspension a
10 more useful drug product can be supplied.

BRIEF DESCRIPTION OF FIGURES

Figure 1 is a graph depicting mean plasma concentrations versus time profiles of the Example 1 formulation of the invention in rats;

15 Figure 2 is a graph depicting mean plasma concentrations versus time profiles of the Example 1 formulation of the invention in dogs; and

Figure 3 is a graph depicting mean plasma concentrations vs. time profiles of the Example 1 formulation of the invention in humans.

20 DETAILED DESCRIPTION OF THE INVENTION

The controlled release sterile injectable aripiprazole formulation of the invention may include aripiprazole in an amount within the range from about 1 to about 40%, preferably from about 5 to about 20%, and more preferably from about 8 to about 15% by weight based on the weight of the sterile injectable formulation.

25 As indicated, desired mean particle size of the aripiprazole is essential in producing an injectable formulation having the desired controlled release properties of the aripiprazole. Thus, to produce desired controlled release, the aripiprazole should have a mean particle size within the range from about 1 to about 30 microns, preferably from about 1 to about 20 microns, and more preferably from about 1 to about
30 10 to 15 microns.

Where the desired controlled release period is at least about two weeks, up to six weeks or more, preferably about three to about four weeks, the aripiprazole will

75535-6

have a mean particle size within the range from about 1 to about 20, preferably from about 1 to about 10 microns, more preferably from about 2 to about 4 microns, and most preferably about 2.5 microns. The aripiprazole having a mean particle size of about 2.5 microns will have a particle size distribution as follows:

5	<u>Preferred</u>	<u>More Preferred</u>
	95% < 20 microns	95% < 8 microns
	90% < 15 microns	90% < 6 microns
	50% < 10 microns	75% < 3 microns
	10% < 2 microns	50% < 1.5 microns
10		10% < 0.5 microns

The aripiprazole formulation of the invention will preferably be formed of:

- A. aripiprazole,
- B. a vehicle therefor, which includes:
 - (a) one or more suspending agents,
 - 15 (b) one or more bulking agents,
 - (c) one or more buffers, and
 - (d) optionally one or more pH adjusting agents.

The suspending agent may be present in an amount within the range from about 0.2 to about 10% by weight, preferably for about 0.5 to about 5% by weight
 20 based on the total weight of the sterile injectable formulation. Examples of suspending agents suitable for use include, but are not limited to, one, two or more of the following: sodium carboxymethyl cellulose, hydroxypropyl cellulose, carboxymethyl cellulose, hydroxypropylethyl cellulose, hydroxypropylmethyl cellulose, and polyvinylpyrrolidone, with sodium carboxymethyl cellulose and
 25 polyvinylpyrrolidone being preferred. Other suspending agents suitable for use in the vehicle for the aripiprazole include various polymers, low molecular weight oligomers, natural products, and surfactants, including nonionic and ionic surfactants, such as cetyl pyridinium chloride, gelatin, casein, lecithin (phosphatides), dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride,
 30 calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers (e.g., macrogol ethers such as cetomacrogol 1000), polyoxyethylene castor oil derivatives, polyoxyethylene

sorbitan fatty acid esters (e.g., the commercially available Tweens® such as e.g., Tween 20® and Tween 80® (ICI Specialty Chemicals)); polyethylene glycols (e.g., Carbowaxs 3350® and 1450®, and Carbopol 934® (Union Carbide)), dodecyl trimethyl ammonium bromide, polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl celluloses (e.g., HPC, HPC-SL, and HPC-L), methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethyl-cellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol (PVA), 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde (also known as tyloxapol, superione, and triton), poloxamers (e.g., Pluronic F68® and F108®, which are block copolymers of ethylene oxide and propylene oxide); poloxamines (e.g., Tetronic 908®, also known as Poloxamine 908®, which is a tetrafunctional block copolymer derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine (BASF Wyandotte Corporation, Parsippany, N.J.)); a charged phospholipid such as dimyristoyl phosphatidyl glycerol, dioctylsulfosuccinate (DOSS); Tetronic 1508® (T-1508) (BASF Wyandotte Corporation), dialkylesters of sodium sulfosuccinic acid (e.g., Aerosol OT®, which is a dioctyl ester of sodium sulfosuccinic acid (American Cyanamid)); Duponol P®, which is a sodium lauryl sulfate (DuPont); Tritons X-200®, which is an alkyl aryl polyether sulfonate (Rohm and Haas); Crodestas F-110®, which is a mixture of sucrose stearate and sucrose distearate (Croda Inc.); p-isononylphenoxypoly-(glycidol), also known as Olin-10G® or Surfactant 10-G® (Olin Chemicals, Stamford, Conn.); Crodestas SL-40® (Croda, Inc.); and SA9OHCO, which is $C_{18}H_{37}CH_2(CON(CH_3)) - CH_2(CHOH)_4(CH_2OH)_2$ (Eastman Kodak Co.); decanoyl-N-methylglucamide; n-decyl β -D-glucopyranoside; n-decyl β -D-maltopyranoside; n-dodecyl β -D-glucopyranoside; n-dodecyl β -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl- β -D-glucopyranoside; n-heptyl β -D-thioglucoside; n-hexyl β -D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl β -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- β -D-glucopyranoside; octyl β -D-thioglucopyranoside; and the like.

75535-6

Most of these suspending agents are known pharmaceutical excipients and are described in detail in the *Handbook of Pharmaceutical Excipients*, published jointly by the American Pharmaceutical Association and The Pharmaceutical Society of Great Britain (The Pharmaceutical Society of Great Britain (The Pharmaceutical Press, 1986). The suspending agents are commercially available and/or can be prepared by techniques known in the art.

Carboxymethyl cellulose or the sodium salt thereof is particularly preferred where the desired mean particle size is about 1 micron or above.

The bulking agent (also referred to as a cryogenic/lyophilize protecting agent) may be present in an amount within the range from about 1 to about 10% by weight, preferably from about 3 to about 8% by weight, more preferably from about 4 to about 5% by weight based on the total weight of the sterile injectable formulation. Examples of bulking agents suitable for use herein include, but are not limited to, one, two or more of the following: mannitol, sucrose, maltose, xylitol, glucose, starches, sorbital, and the like, with mannitol being preferred for formulations where the mean particle size is about 1 micron or above. It has been found that xylitol and/or sorbitol enhances stability of the aripiprazole formulation by inhibiting crystal growth and agglomeration of drug particles so that desired particle size may be achieved and maintained.

The buffer may be employed in an amount to adjust pH of an aqueous suspension of the freeze-dried aripiprazole formulation to from about 6 to about 8, preferably about 7. To achieve such pH, usually the buffer, depending on type, will be employed in an amount within the range from about 0.02 to about 2% by weight, preferably from about 0.03 to about 1% by weight, and more preferably about 0.1% by weight based on the total weight of the sterile injectable formulation. Examples of buffers suitable for use herein include, but are not limited to, one, two or more of the following: sodium phosphate, potassium phosphate, or TRIS buffer, with sodium phosphate being preferred.

The freeze-dried formulation of the invention may optionally include a pH adjusting agent which may then be employed in an amount to adjust pH of the aqueous suspension of the freeze-dried aripiprazole within the range from about 6 to about 7.5, preferably about 7 and may be an acid or base depending upon whether the pH of the

75535-6

aqueous suspension of the freeze-dried aripiprazole needs to be raised or lowered to reach the desired neutral pH of about 7. Thus, where the pH needs to be lowered, an acidic pH adjusting agent such as hydrochloric acid or acetic acid, preferably hydrochloric acid, may be employed. When the pH needs to be raised, a basic pH
5 adjusting agent will be employed such as sodium hydroxide, potassium hydroxide, calcium carbonate, magnesium oxide or magnesium hydroxide, preferably sodium hydroxide.

The freeze-dried aripiprazole formulations may be constituted with an amount of water for injection to provide from about 10 to about 400 mg of aripiprazole
10 delivered in a volume of 2.5 mL or less, preferably 2 mL for a two to six week dosage.

In carrying out the method for preparing the freeze-dried aripiprazole formulation of the invention, it is required that everything be sterile so that sterile aripiprazole and sterile vehicle are combined aseptically to form a sterile suspension and that the sterile suspension be freeze-dried in a manner to form sterile freeze-dried
15 powder or cake. Thus, an aseptic procedure is employed to produce sterile bulk aripiprazole of desired particle size distribution. The sterile bulk aripiprazole will have a mean particle size within the range from about 5 to about 100 microns, preferably from about 10 to about 90 microns.

Preferably, an impinging jet crystallization method is employed to produce
20 sterile aripiprazole of desired small particle size and a narrow size distribution, high surface area, high chemical purity, high stability due to improved crystal structure, while employing continuous processing.

The impinging jet crystallization utilizes two jet streams that strike each other head-on. One of the streams carries a solution rich in the aripiprazole and the other
25 carries an anti-solvent, such as water. The two streams strike each other which allows rapid homogeneous mixing and supersaturation due to high turbulence and high intensity of micromixing upon impact. This immediate achievement of supersaturation initiates rapid nucleation. In general, the average crystal size decreases with increasing supersaturation and decreasing temperature of the anti-
30 solvent. Therefore, in order to obtain the smallest particle size, it is advantageous to have the highest possible concentration of the rich solution and the lowest temperature of the anti-solvent.

75535-6

The vehicle for the sterile bulk aripiprazole including suspending agent, bulking agent, buffer, optional pH adjusting agent and water is prepared and subjected to sterilization. Thereafter, the sterile bulk aripiprazole and sterile vehicle are aseptically combined to form a sterile primary suspension and the particle size of the aripiprazole is reduced to a desired level. This is preferably carried out employing an aseptic wet milling procedure wherein sterile particles of aripiprazole dispersed in the sterile vehicle are subjected to grinding means in the presence of grinding media to reduce the particle size of the aripiprazole to within the range of about 1 to about 10 microns, depending upon desired controlled release period.

10 The aseptic wet milling procedure is preferably wet ball milling. When the desired mean particle size of the aripiprazole is above about 1 micron, the primary suspension (combined aripiprazole-vehicle) is passed through a wet ball mill a single time (single pass) at from about 5 to about 15 L/hr, preferably from about 8 to about 12 L/hr, and more preferably about 10 L/hr, to reduce mean particle size of the aripiprazole to within the desired range, for example, from about 1 to about 5 microns.

In addition to ball mills, such as Dyno mills, other low and high energy mills may be employed such as a roller mill, and high energy mills may be employed such as Netzsch mills, DC mills and Planetary mills. However, it is essential that the milling procedure and equipment employed be capable of producing sterile aripiprazole formulation of desired mean particle size.

Other techniques for particle size reduction which may be employed include aseptic controlled crystallization, high shear homogenization, high pressure homogenization and microfluidization to produce particles having a mean particle size ranging from about 1 to about 100 microns.

The resulting final suspension is aseptically filled into sterile vials and aseptically loaded into a sterilized freeze drier. It is essential that a carefully designed freeze drying cycle be applied in order to form and/or maintain the desired crystalline form of the aripiprazole, which is known to exist in monohydrate form (Aripiprazole Hydrate A) as well as in a number of anhydrous forms, namely Anhydrous Crystals B, Anhydrous Crystals C, Anhydrous Crystals D, Anhydrous Crystals E, Anhydrous

75535-6

Crystals F, and Anhydrous Crystals G, all of which may be employed in the formulation of the invention.

The aripiprazole monohydrate (grains) or hydrate as referred to below employed in the present invention has the physicochemical properties given in (1)-(5) below. This aripiprazole hydrate is described hereinafter as "Aripiprazole Hydrate A".

(1) It has an endothermic curve characterized by the appearance of a small peak at about 71°C and a gradual endothermic peak around 60°C to 120°C.

(2) It has an ¹H-NMR spectrum which has characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, J = 7.4 Hz, 2H), 2.97 ppm (brt, J = 4.6 Hz, 4H), 3.92 ppm (t, J = 6.3 Hz, 2H), 6.43 ppm (d, J = 2.4 Hz, 1H), 6.49 ppm (dd, J = 8.4 Hz, J = 2.4 Hz, 1H), 7.04 ppm (d, J = 8.1 Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

(3) It has a powder x-ray diffraction spectrum which has characteristic peaks at 2θ = 12.6°, 15.4°, 17.3°, 18.0°, 18.6°, 22.5° and 24.8°.

(4) It has clear infrared absorption bands at 2951, 2822, 1692, 1577, 1447, 1378, 1187, 963 and 784 cm⁻¹ on the IR (KBr) spectrum.

(5) It has a mean grain size of 50 μm or less.

Aripiprazole Hydrate A is manufactured by milling conventional aripiprazole hydrate.

Conventional milling methods can be used to mill the aripiprazole hydrate. For example, the aripiprazole hydrate can be milled in a milling machine. A widely used milling machine can be used, such as an atomizer, pin mill, jet mill or ball mill. Of these, the atomizer is preferred.

Regarding the specific milling conditions when using an atomizer, a rotational speed of 5000-15000 rpm could be used for the main axis, for example, with a feed rotation of 10-30 rpm and a screen hole size of 1-5 mm.

The mean grain size of the Aripiprazole Hydrate A obtained by milling should normally be 50 μm or less, preferably 30 μm or less. Mean grain size can be ascertained by the grain size measurement method described hereinafter.

75535-6

Grain Size Measurement: 0.1 g of the grains to be measured were suspended in a 20 ml n-hexane solution of 0.5 g soy lecithin, and grain size was measured using a size distribution meter (Microtrack HRA, Microtrack Co.).

5 The aripiprazole anhydrous crystals employed in the present invention have the physicochemical properties given in (6)-(10) below. These aripiprazole anhydrous crystals are referred to hereinafter as "Aripiprazole Anhydrous Crystals B".

(6) They have an ^1H -NMR spectrum, which has (DMSO- d_6 , TMS). ^1H -NMR spectrum, which has characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, $J = 7.4$ Hz, 2H), 2.97 ppm (brt, $J = 4.6$ Hz, 4H), 3.92 ppm (t, $J = 6.3$ Hz, 2H), 6.43 ppm (d, $J = 2.4$ Hz, 1H), 6.49 ppm (dd, $J = 8.4$ Hz, $J = 2.4$ Hz, 1H), 7.04 ppm (d, $J = 8.1$ Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

15 (7) They have a powder x-ray diffraction spectrum which has characteristic peaks at $2\theta = 11.0^\circ$, 16.6° , 19.3° , 20.3° and 22.1° .

(8) They have clear infrared absorption bands at 2945, 2812, 1678, 1627, 1448, 1377, 1173, 960 and 779 cm^{-1} on the IR (KBr) spectrum.

20 (9) They exhibit an endothermic peak near about 141.5°C in thermogravimetric/differential thermal analysis (heating rate $5^\circ\text{C}/\text{min}$).

(10) They exhibit an endothermic peak near about 140.7°C in differential scanning calorimetry (heating rate $5^\circ\text{C}/\text{min}$).

Aripiprazole Anhydrous Crystals B employed in the present invention have low hygroscopicity. For example, Aripiprazole Anhydrous Crystals B employed in the present invention maintain a water content of 0.4% or less after 24 hours inside a desiccator set at a temperature of 60°C and a humidity of 100%.

Well-known methods of measuring water content can be used as long as they are methods commonly used for measuring the water content of crystals. For example, a method such as the Karl Fischer method can be used.

30 The Aripiprazole Anhydrous Crystals B employed in the present invention are prepared for example by heating the aforementioned Aripiprazole Hydrate A at 90-

75535-6

125°C. The heating time is generally about 3-50 hours, depending on heating temperature. The heating time and heating temperature are inversely related, so that for example the heating time will be longer the lower the heating temperature, and shorter the higher the heating temperature. Specifically, if the heating temperature of Aripiprazole Hydrate A is 100°C, the heating time should normally be 18 hours or more or preferably about 24 hours. If the heating temperature of Aripiprazole Hydrate A is 120°C, on the other hand, the heating time can be about 3 hours. The Aripiprazole Anhydrous Crystals B of the present invention can be prepared with certainty by heating Aripiprazole Hydrate A for about 18 hours at 100°C, and then heating it for about 3 hours at 120°C.

Furthermore, the Aripiprazole Anhydrous Crystals B employed in the present invention are prepared for example by heating conventional aripiprazole anhydrous crystals at 90-125°C. The heating time is generally about 3-50 hours, depending on heating temperature. The heating time and heating temperature are inversely related, as described above. Specifically, if the heating temperature of the aripiprazole anhydrous crystals is 100°C, the heating time can be about 4 hours, and if the heating-temperature is 120°C the heating time can be about 3 hours.

The aripiprazole anhydrous crystals which are the raw material for preparing the Aripiprazole Anhydrous Crystals B employed in the present invention are prepared for example by Method a or b below.

20 **Method a:**

Aripiprazole Anhydrous Crystals are prepared by well-known methods, as by reacting 7-(4-bromobutoxy)-3,4-dihydrocarbostyryl with 1-(2,3-dichlorophenyl) piperidine and recrystallizing the resulting raw aripiprazole crystals with ethanol as described in Example 1 of Japanese Unexamined Patent Publication No. 191256/1990.

25 **Method b:**

Aripiprazole Anhydrous Crystals are prepared by heating conventional aripiprazole hydrate at a temperature of at least 60°C and less than 90°C. The heating

75535-6

time is generally about 1-30 hours, depending on heating temperature. The heating time and heating temperature are inversely related, as described above. Specifically, if the heating temperature of the aripiprazole hydrate is about 60°C, the heating time can be about 8 hours, while if the heating temperature is 80°C, the heating time can be about 4 hours.

Method b is described in the Proceedings of the 4th Japanese-Korean Symposium on Separation Technology (October 6-8, 1996).

Furthermore, the Aripiprazole Anhydrous Crystals B employed in the present invention are prepared for example by heating conventional aripiprazole hydrate at 90-125°C. The heating time is generally about 3-50 hours, depending on heating temperature. The heating time and heating temperature are inversely related. Specifically, if the heating temperature of the aripiprazole hydrate is 100°C, the heating time can be about 24 hours, while if the heating temperature is 120°C, the heating time can be about 3 hours.

The aripiprazole hydrate which is the raw material for preparing the Aripiprazole Anhydrous Crystals B employed in the present invention is prepared for example by Method c below.

Method c:

Aripiprazole hydrate is obtained by dissolving the aripiprazole anhydrous crystals obtained by Method a above in a hydrous solvent, and heating and then cooling the resulting solution. Using this method, aripiprazole hydrate is precipitated as crystals in the hydrous solvent.

An organic solvent containing water is usually used as the hydrous solvent. The organic solvent should be one which is miscible with water, such as for example an alcohol such as methanol, ethanol, propanol or isopropanol, a ketone such as acetone, an ether such as tetrahydrofuran, dimethylformamide, or a mixture thereof, with ethanol being particularly desirable. The amount of water in the hydrous solvent can be 10-25% by weight of the solvent, or preferably close to 20% by weight.

As mentioned above, the Aripiprazole Anhydrous Crystals B employed in the present invention are prepared by heating at 90-125°C of the Aripiprazole Hydrate A,

75535-6

conventional aripiprazole anhydrous crystals or conventional aripiprazole hydrate, and the Aripiprazole Hydrate A, conventional aripiprazole anhydrous crystals or conventional aripiprazole hydrate may be used either individually or in combination.

5 The above crystal forms and other crystal forms of aripiprazole and methods for making such crystal forms which may be employed herein include Hydrate A and Anhydrous Crystals B as well as Anhydrous Crystals C, Anhydrous Crystals D, Anhydrous Crystals E, Anhydrous Crystals F, and Anhydrous Crystals G, as disclosed in PCT WO 03/26659, published April 4, 2003.

10 If the monohydrate form of the aripiprazole in the freeze-dried formulation is desired then the freeze drying cycle should include cooling of the formulation to about -40°C at a suitable cooling rate. The primary drying should be performed at a temperature below around 0°C and a suitable vacuum and duration.

15 If the anhydrous form of the aripiprazole in the freeze-dried formulation is desired then the freeze drying cycle should include three phases: freezing, primary drying, and secondary drying. The freezing phase should include cooling of the formulation to about -40°C at a suitable cooling rate. The primary drying should be performed at a temperature below about 0°C and a suitable vacuum and duration. The secondary drying should be performed at a temperature above about 0°C and a suitable vacuum and duration.

20 The vials with the resulting freeze-dried aripiprazole suspension are aseptically stoppered under atmospheric pressure or partial vacuum and sealed.

Preferred injectable formulations in the form of aqueous suspensions are set out below:

75535-6

Preferred Particle Size of About 1-10 Microns (Preferably About 2.5 Microns)

	Range % w/v	Preferred Range % w/v
(1) Sterile Aripiprazole – (mean particle size from about 1 to about 5 microns)	1 to 40	8 to 15
(2) Suspending Agent (preferably carboxymethyl cellulose Na salt)	0.2 to 10	0.5 to 5
(3) Bulking Agent (preferably mannitol)	1 to 10	4 to 5
(4) Buffer (preferably sodium phosphate) (adjust pH to about 6 to about 7.5)	0.02 to 2	0.03 to 1
(5) pH Adjusting Agent (preferably sodium hydroxide) (adjust pH to about 6 to about 7.5)	As needed	As needed
(6) Water for injection	As needed	As needed

5 The aripiprazole will be present in the aqueous injectable formulation in an amount within the range from about 1 to about 40% (w/v), preferably from about 5 to about 20 % (w/v), and more preferably from about 8 to about 15% (w/v), based on the total injectable formulation.

In preferred embodiments, the aripiprazole will be present in the aqueous injectable formulation to provide from about 50 to about 400 mg/2 mL of formulation, preferably from about 100 to about 200 mg/mL of formulation.

10 Preferred individual dose injectable formulations in accordance with the invention are as follows:

aripiprazole	100 mg	200 mg	400 mg
carboxymethyl cellulose	9 mg	9 mg	9 mg
mannitol	45 mg	45 mg	45 mg
Na phosphate	0.8 mg	0.8 mg	0.8 mg
sodium hydroxide	qs to adjust pH to 7	qs to adjust pH to 7	qs to adjust pH to 7
water for injection	qs to 1 mL	qs to 1 mL	qs to 1 mL

15 The aripiprazole formulations of the invention may be used in the treatment of schizophrenia and related disorders such as bipolar disorder and dementia in human patients. The preferred dosage employed for the injectable formulations of the invention will be a single injection or multiple injections containing from about 100 to about 400 mg

aripiprazole/mL given one to two times monthly. The injectable formulation is preferably administered intramuscularly, although subcutaneous injections are acceptable as well.

The following examples represent preferred embodiments of the invention.

5

EXAMPLES

EXAMPLE 1

An aripiprazole injectable (IM Depot) aqueous suspension (200 mg
5 aripiprazole/2 mL, 200 mg/vial) was prepared as follows.

ARIPIRAZOLE MICROSUSPENSION PREPARED BY MEDIA MILLING

A microparticulate dispersion of aripiprazole was prepared using a DYNO®-
MILL (Type KDL A, manufactured by Willy A. Bachoffen AG Maschinenfabrik,
10 Basel, Switzerland).

The following ingredients were added to a 3L glass jacketed vessel maintained
at 15°C ($\pm 5^\circ\text{C}$) to form a sterile primary suspension:

	Aripiprazole	100 g
	Carboxymethylcellulose, Sodium Salt 7L2P	9.0 g
15	Mannitol	45 g
	Sodium Phosphate, Monobasic	0.8 g
	Sodium Hydroxide Solution, 1N	q.s. to adjust pH to 7.0
	Water for Injection, USP	q.s. to 1040 g

The primary suspension was mixed at 500-1000 rpm for about 0.5 hour and
20 then at 300-500 rpm for an additional 1 hour under 20 "Hg (± 5 "Hg) vacuum.

The media mill was prepared accordingly for the media milling process. The
grinding container was partially filled with zirconium oxide beads and the dispersion
was passed through the mill operating at the following conditions:

	Grinding container:	water jacketed 0.6 L stainless steel vessel
25	Coolant temperature:	15°C ($\pm 5^\circ\text{C}$)
	Agitation speed:	2500 rpm
	Grinding medium:	500 mL of very-high-density (VHD) zirconium oxide beads
	Suspension flow rate:	10 L/h
30	Milling time:	6 minutes

After a single pass milling, a sample of the processed suspension was removed and evaluated for particle size distribution using Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer. The particles were determined to have a mean particle size of 2.5 microns (μ) and the following particle size distribution: 10% < 0.4 μ , 50% < 1.6 μ , 75% < 3.3 μ , 90% < 5.9 μ , and 95% < 7.6 μ .

2.5 mL of the above suspension were aseptically filled into sterilized vials which were then aseptically partially stoppered with sterilized stoppers. The vials were aseptically transferred to a freeze dryer and lyophilized according to the following cycle:

- (a) thermal treatment: freeze product at -40°C over 0.1-1 h and keep at -40°C for at least 3 h,
- (b) cool the condenser to -50°C or below,
- (c) primary drying: lower chamber pressure to approximately 100 microns Hg and increase product temperature to -5°C over approximately 2 h; continue primary drying at -5°C and 100 microns Hg for at least 48 h,
- (d) stopper the vials under atmospheric pressure or partial vacuum using sterile nitrogen or air and remove from the freeze dryer,
- (e) seal the vials with the appropriate seals and label.

20

EXAMPLE 2

An aripiprazole injectable (IM Depot) aqueous suspension (200 mg aripiprazole/2 mL, 200 mg/vial) was prepared as follows.

25

ARIPIRAZOLE MICROSUSPENSION PREPARED BY IMPINGING JET CRYSTALLIZATION

A microparticulate dispersion of aripiprazole was prepared using impinging jet crystallization.

The following procedure was employed to form a sterile bulk aripiprazole:

1. Suspend 100 g of aripiprazole in 2000 mL of 95% ethanol. Heat the suspension to 80°C until it becomes a clear solution.

2. Polish filter the aripiprazole solution into a holding vessel and maintain at 80° C.
3. Polish filter 2000 mL water to another holding vessel and heat to 80° C.
4. Pump the aripiprazole solution through a 0.02 inch diameter nozzle at 0.25 kg/min and impinge it with the 30° C water pumped at 0.25 kg/min through a 0.02 inch diameter nozzle to form a crystal slurry which is collected in an impingement vessel.
5. Agitate the newly formed crystal slurry in the impingement vessel while continuously transferring it to a receiver to maintain a constant volume in the impingement vessel.
6. At the end of impingement, cool the slurry in the receiver to room temperature.
7. Filter the slurry.
8. Dry the wet cake at 35° C under vacuum to yielding 100 g (96% recovery) of aripiprazole with reduced particle size (90% < 100 µm).

The following ingredients were added to a 3L glass jacketed vessel maintained at 15°C (±5°C) to form a sterile primary suspension:

	Aripiprazole (prepared by impinging jet crystallization):	100 g
20	Carboxymethylcellulose, Sodium Salt 7L2P	9.0 g
	Mannitol	45 g
	Sodium Phosphate, Monobasic	0.8 g
	Sodium Hydroxide Solution, 1N	q.s. to adjust pH to 7.0
	Water, USP	q.s. to 1040 g

- 25 The sterile suspension was mixed at 500-1000 rpm for about 0.5 hour and then at 300-500 rpm for an additional 1 hour under 20 "Hg (± 5"Hg) vacuum.

The sterile suspension was found to contain particles having mean particle size of 2.5 microns and the following particle size distribution:

	10%	<	0.4 μ
	50%	<	1.6 μ
5	75%	<	3.3 μ
	90%	<	5.9 μ
	95%	<	7.5 μ

2.5 mL of the above suspension were aseptically filled into sterilized vials which were then aseptically partially stoppered with sterilized stoppers. The vials were aseptically transferred to a freeze dryer and lyophilized according to the following cycle:

- (a) thermal treatment: freeze product at -40°C over 0.1-1 h and keep at -40°C for at least 6 h,
- (b) cool the condenser to -50°C or below,
- 15 (c) primary drying: lower chamber pressure to approximately 100 microns Hg and increase product temperature to -5°C over approximately 2 h; continue primary drying at -5°C and 100 microns Hg for at least 48 h,
- (d) stopper the vials under atmospheric pressure or partial vacuum using sterile nitrogen or air and remove from the freeze dryer,
- 20 (e) seal the vials with the appropriate seals and label.

EXAMPLE 3 (Animal PK Data)

A. Single-dose I.M. depot study in rats

25 Aripiprazole I.M. depot formulation prepared in Example 1 was injected into the thigh muscle of fifteen rats (M-males, F-females) at doses of 12.5, 25, and 50 mg/kg. Blood samples for the evaluation of systemic exposure after aripiprazole I.M. depot administration were collected on days 1 (6 h postdose), 2, 4, 7, 10, 15, 22, 28, 36, and 43 and analyzed for aripiprazole. Figure 1 shows mean plasma concentrations
30 vs. time profiles of aripiprazole in rats.

B. Single-dose I.M. depot study in dogs

The aripiprazole intramuscular (I.M). depot formulation prepared in Example 1 was injected into the thigh muscle of five dogs (M-males, F-females) at doses of 100, 200, and 400 mg. Blood samples for the evaluation of systemic exposure after aripiprazole I.M. depot administration were collected on days 1 (10 and 30 minutes, and 1, 3, and 8 h postdose), 2, 4, 7, 10, 15, 22, 28, 36, and 42 and analyzed for aripiprazole. Figure 2 shows mean plasma concentrations vs. time profiles of aripiprazole in dogs.

PK profiles

Mean aripiprazole rats' serum concentration-time profiles are shown graphically in Fig.1. Aripiprazole aqueous suspensions showed steady serum concentration for at least 4 weeks in the rats' model.

Mean aripiprazole dogs' serum concentration-time profiles are shown graphically in Fig.2.

Aripiprazole aqueous suspensions showed steady serum concentration for 3-4 weeks in the dogs' model.

EXAMPLE 4 (Human PK Data)**Single-dose I.M. depot study in patients**

Aripiprazole I.M. depot formulation prepared in Example 1 was administered intramuscularly to patients diagnosed with chronic, stable schizophrenia or schizoaffective disorder at. The study design included administration of a 5-mg dose of aripiprazole solution to all subjects followed by a single dose of IM depot at 15, 50, and 100 mg per patient. Samples for PK analysis were collected until plasma concentrations of aripiprazole were less than the lower limit of quantification (LLQ) for 2 consecutive visits.

Figure 3 shows mean plasma concentrations vs. time profiles of aripiprazole in subjects 2 and 3 dosed with 15 mg of IM Depot, and subjects 4 and 5 who received 50 mg of IM Depot. In all cases aripiprazole plasma levels showed a fast onset of release and sustained release for at least 30 days.

75535-6

CLAIMS:

1. A sterile freeze-dried controlled release aripiprazole formulation which comprises:

(a) aripiprazole having a mean particle size of 1 to 10 microns, and

5 (b) a vehicle therefore,

which formulation upon reconstitution with water forms a sterile injectable formulation in the form of a sterile homogeneous suspension which upon injection releases aripiprazole over a period of at least two weeks.

10 2. A sterile freeze-dried controlled release aripiprazole formulation, which comprises:

(a) aripiprazole having a mean particle size of 1 to 10 microns, and

(b) one or more suspending agents,

wherein said sterile freeze-dried controlled release aripiprazole formulation is obtained by freeze-drying an aqueous suspension comprising (a) and (b), and

15 wherein upon reconstitution, which comprises adding water to the freeze-dried formulation, a sterile injectable formulation is obtained which is a homogeneous suspension and which contains from 0.1 to 600 mg of aripiprazole per mL, and wherein upon injection into a subject, said sterile injectable formulation releases aripiprazole over a period of at least two weeks from the date of administration.

20 3. The freeze-dried formulation as defined in claim 1, wherein upon reconstitution with water the sterile injectable formulation is obtained which contains from 0.1 to 600 mg of aripiprazole per mL.

4. The freeze-dried formulation as defined in claim 1 or 3, wherein said vehicle comprises:

75535-6

(b) one or more suspending agents,

(c) one or more bulking agents, and

(d) one or more buffering agents.

5. The freeze-dried formulation as defined in claim 2, which further comprises:

5 (c) one or more bulking agents, and

(d) one or more buffering agents.

6. The freeze-dried formulation as defined in claim 4 or 5, wherein the sterile injectable formulation comprises:

10 (a) aripiprazole having a mean particle size of 1 to 10 microns in an amount within the range from 1 to 40%,

(b) suspending agent in an amount within the range from 0.2 to 10%,

(c) bulking agent in an amount within the range from 1 to 10%, and

(d) buffering agent in an amount within the range from 0.02 to 2% to adjust the pH of the sterile injectable formulation within the range from 6 to 7.5,

15 wherein all of the above % are % by weight based on the volume of the sterile injectable formulation.

7. The freeze-dried formulation as defined in claim 6, wherein the sterile injectable formulation comprises:

20 (a) aripiprazole having a mean particle size of 1 to 10 microns in an amount within the range from about 5 to about 20%,

(b) suspending agent in an amount within the range from about 0.5 to about 5%,

75535-6

(c) bulking agent in an amount within the range from about 4 to about 5%, and

(d) buffering agent in an amount within the range from about 0.03 to about 1% to adjust the pH of the sterile injectable formulation within the range from 6 to 7.5,

wherein all of the above % are % by weight based on the volume of the sterile injectable formulation.

8. The freeze-dried formulation as defined in any one of claims 4 to 7, wherein the suspending agent is carboxymethylcellulose or its sodium-salt, hydroxypropyl cellulose, hydroxypropylethyl cellulose, hydroxypropylmethyl cellulose, or polyvinylpyrrolidone; the bulking agent is mannitol, sucrose, maltose, lactose, xylitol or sorbitol; and the buffering agent is sodium phosphate, potassium phosphate or TRIS buffer.

9. The freeze-dried formulation as defined in any one of claims 4 to 8, which further comprises:

(e) a pH adjusting agent.

10. The freeze-dried formulation as defined in any one of claims 4 to 8, comprising:

(a) aripiprazole having a mean particle size of 1 to 10 microns,

(b) carboxymethyl cellulose or its sodium salt,

(c) mannitol, and

(d) sodium phosphate to adjust pH to about 7.

11. The freeze-dried formulation as defined in claim 9 or 10, which further comprises:

(e) sodium hydroxide to adjust pH to 7.

75535-6

12. The freeze-dried formulation as defined in any one of claims 1 to 11, which upon reconstitution with water and injection releases aripiprazole over a period of at least three weeks.

13. The freeze-dried formulation as defined in any one of claims 1 to 12, which
5 upon reconstitution with water and injection releases aripiprazole over a four week period.

14. The freeze-dried formulation as defined in any one of claims 1 to 13, wherein said freeze-dried formulation is in the form of a powder or cake.

15. The freeze-dried formulation as defined in any one of claims 1 to 14, wherein the aripiprazole is in the form of anhydrous crystals or is in the form of a monohydrate.

10 16. The freeze-dried formulation as defined in any one of claims 1 to 15, wherein the aripiprazole is in the form of a monohydrate.

17. The freeze-dried formulation as defined in any one of claims 1 to 15, wherein the aripiprazole is Aripiprazole Hydrate A.

18. The freeze-dried formulation as defined in any one of claims 1 to 17, wherein
15 the aripiprazole has a mean particle size of 1 to 5 microns.

19. The freeze-dried formulation as defined in any one of claims 1 to 18, wherein the aripiprazole has a mean particle size within the range from 2 to 4 microns.

20. The freeze-dried formulation as defined in any one of claims 1 to 19, wherein the aripiprazole has a mean particle size of 2.5 microns.

20 21. The freeze-dried formulation as defined in any one of claims 1 to 19, wherein the aripiprazole has a mean particle size of about 2.5 microns and a particle size distribution as follows:

95% < 20 microns

90% < 15 microns

75535-6

50% < 10 microns

10% < 2 microns.

22. The freeze-dried formulation as defined in any one of claims 1 to 21, wherein the sterile injectable formulation contains from 100 to 400 mg of aripiprazole per mL.

5 23. The freeze-dried formulation as defined in any one of claims 1 to 21, wherein the sterile injectable formulation contains from 200 to 400 mg of aripiprazole per mL.

24. The freeze-dried formulation as defined in any one of claims 1 to 21, wherein the sterile injectable formulation contains about 200 mg of aripiprazole per mL.

10 25. The freeze-dried formulation as defined in any one of claims 1 to 21, which contains about 300 mg of aripiprazole.

26. The freeze-dried formulation as defined in any one of claims 1 to 21, which contains about 400 mg of aripiprazole.

15 27. The freeze-dried formulation as defined in any one of claims 1 to 26, wherein the sterile injectable formulation upon injection maintains a substantially constant aripiprazole plasma concentration over the period of release.

28. The freeze-dried formulation as defined in any one of claims 1 to 27, wherein the sterile injectable formulation is an intramuscular formulation.

29. The freeze-dried formulation as defined in claim 1, which is obtained by a process which comprises the steps of:

20 (a) preparing sterile bulk aripiprazole having a desired particle size distribution,

(b) preparing a sterile vehicle for the sterile bulk aripiprazole,

75535-6

(c) combining said sterile aripiprazole and said sterile vehicle to form a sterile primary suspension,

(d) reducing the mean particle size of the aripiprazole in said sterile primary suspension to within the range from 1 to 10 microns to form a sterile final suspension, and

5 (e) freeze-drying said sterile final suspension to form the freeze-dried formulation.

30. The freeze-dried formulation as defined in claim 2, which is obtained by a process which comprises the steps of:

(a) preparing sterile bulk aripiprazole having a desired particle size
10 distribution,

(b) preparing a sterile vehicle comprising one or more sterile suspending agents for the sterile bulk aripiprazole,

(c) combining said sterile aripiprazole and said sterile vehicle comprising one or more sterile suspending agents to form a sterile primary suspension,

15 (d) reducing the mean particle size of the aripiprazole in said sterile primary suspension to within the range from 1 to 10 microns to form a sterile final suspension containing from 0.1 to 600 mg of aripiprazole monohydrate per mL, and

(e) freeze-drying said sterile final suspension to form the freeze-dried formulation.

20 31. The freeze-dried formulation as defined in claim 29 or 30, wherein the step of reducing the mean particle size of the aripiprazole in said sterile primary suspension is carried out employing wet milling.

32. The freeze-dried formulation as defined in claim 31, wherein the wet milling comprises wet ball milling.

75535-6

33. The freeze-dried formulation as defined in any one of claims 29 to 32, wherein the aripiprazole is in the form of a monohydrate.

34. The freeze-dried formulation as defined in any one of claims 29 to 33, wherein the freeze drying step of the sterile final suspension is carried out in three phases:

5 (1) a freezing phase which includes cooling of the sterile final suspension at -40°C,

(2) a primary drying phase which is performed at below 0°C, and

(3) a secondary drying phase which is performed at above 0°C, to form aripiprazole in anhydrous form.

10 35. A method for preparing the sterile freeze-dried formulation as defined in claim 1, which comprises the steps of:

(a) preparing sterile bulk aripiprazole having a desired particle size distribution,

(b) preparing a sterile vehicle for the sterile bulk aripiprazole,

15 (c) combining said sterile aripiprazole and said sterile vehicle to form a sterile primary suspension,

(d) reducing the mean particle size of the aripiprazole in said sterile primary suspension to within the range from 1 to 10 microns to form a sterile final suspension, and

20 (e) freeze-drying said sterile final suspension to form the freeze-dried formulation.

36. A method for preparing the freeze-dried formulation as defined in claim 2, which comprises the steps of:

75535-6

(a) preparing sterile bulk aripiprazole having a desired particle size distribution,

(b) preparing a sterile vehicle comprising one or more suspending agents for the sterile bulk aripiprazole,

5 (c) combining said sterile aripiprazole and said sterile vehicle comprising one or more suspending agents to form a sterile primary suspension,

(d) reducing the mean particle size of the aripiprazole in said sterile primary suspension to within the range from 1 to 10 microns to form a sterile final suspension containing from 0.1 to 600 mg of aripiprazole per mL, and

10 (e) freeze-drying said sterile final suspension to form the freeze-dried formulation.

37. The method as defined in claim 35 or 36, wherein the step of reducing the mean particle size of the aripiprazole in said sterile primary suspension is carried out employing wet milling.

15 38. The method as defined in claim 37, wherein the wet milling comprises wet ball milling.

39. The method as defined in any one of claims 35 to 38, wherein said freeze-drying step is carried out by cooling the sterile final suspension to -40°C and drying said cooled sterile final suspension at below 0°C, to form freeze-dried aripiprazole in the form of
20 its monohydrate.

40. The method as defined in any one of claims 35 to 39, wherein the freeze drying step of the sterile final suspension is carried out in three phases:

(1) a freezing phase which includes cooling of the sterile final suspension at -40°C,

75535-6

(2) a primary drying phase which is performed at below 0°C, and

(3) a secondary drying phase which is performed at above 0°C, to form aripiprazole in anhydrous form.

41. A controlled release injectable formulation prepared by mixing the freeze-dried

5 formulation as defined in any one of claims 1 to 34 with sterile water for injection.

42. The formulation as defined in claim 41, comprising one of the following

aripiprazole	100 mg	200 mg	400 mg
carboxymethyl cellulose	9 mg	9 mg	9 mg
mannitol	45 mg	45 mg	45 mg
Na phosphate	0.8 mg	0.8 mg	0.8 mg
sodium hydroxide	qs to adjust pH to 7	qs to adjust pH to 7	qs to adjust pH to 7
water for injection	qs to 1 mL	qs to 1 mL	qs to 1 mL

43. A product comprising the freeze-dried formulation as defined in any one of

10 claims 1 to 34 and sterile water for use on reconstitution of the freeze-dried formulation.

44. A lyophilized formulation for release of aripiprazole over a period of about two to about four weeks, which comprises:

(a) aripiprazole having a mean particle size of 1 to 10 microns;

(b) carboxymethylcellulose or its sodium-salt, hydroxypropyl cellulose,
15 hydroxypropylethyl cellulose, hydroxypropylmethyl cellulose, or polyvinylpyrrolidone;

(c) mannitol, sucrose, maltose, lactose, xylitol or sorbitol; and

(d) sodium phosphate, potassium phosphate or TRIS buffer.

75535-6

45. The lyophilized formulation as defined in claim 44, which comprises:

(a) aripiprazole monohydrate having a mean particle size of 2 to 4 microns;

(b) carboxymethyl cellulose or its sodium salt;

(c) mannitol;

5 (d) sodium phosphate;

and which further comprises:

(e) sodium hydroxide.

46. The lyophilized formulation as defined in claim 44 or 45, wherein the mean particle size is 2.5 microns.

10 47. The lyophilized formulation as defined in claim 44 or 45, wherein the mean particle size is about 2.5 microns and with a particle size distribution as follows:

95% < 20 microns

90% < 15 microns

50% < 10 microns

15 10% < 2 microns.

48. The lyophilized formulation as defined in any one of claims 44 to 47, which is for release of aripiprazole over a period of about two weeks.

49. The lyophilized formulation as defined in any one of claims 44 to 47, which is for release of aripiprazole over a period of about three weeks.

20 50. The lyophilized formulation as defined in any one of claims 44 to 47, which is for release of aripiprazole over a period of about four weeks.

75535-6

51. The lyophilized formulation as defined in any one of claims 44 to 50, wherein (a) is aripiprazole hydrate A.

52. The lyophilized formulation as defined in any one of claims 44 to 51 for maintaining a substantially constant aripiprazole plasma concentration over the period of
5 release.

53. An injectable formulation comprising the lyophilized formulation as defined in any one of claims 44 to 52 and sterile water for injection.

54. The injectable formulation as defined in claim 53, which is an intramuscular formulation.

10 55. A controlled release sterile aripiprazole injectable formulation in the form of a sterile homogeneous suspension which upon injection releases aripiprazole over a period of at least two weeks, which comprises:

(a) aripiprazole having a mean particle size of 1 to 10 microns in an amount of 100 to 400 mg per mL,

15 (b) a vehicle therefore, and

(c) water for injection.

56. The formulation as defined in claim 55, wherein said vehicle comprises one or more suspending agents.

57. The formulation as defined in claim 55, wherein the vehicle comprises:

20 (1) one or more suspending agents,

(2) optionally one or more bulking agents, and

(3) optionally one or more buffering agents.

58. The formulation as defined in claim 57, which comprises:

• 75535-6

(a) the aripiprazole is present in an amount within the range from 1 to 40%,

(b) the suspending agent is present in an amount within the range from 0.2 to 10%,

(c) the bulking agent is present in an amount within the range from 2 to 10%,

5 (d) the buffer is present in an amount within the range from 0.02 to 2% to adjust the pH of the suspension within the range from 6 to 7.5,

wherein all of the above % are % by weight based on the volume of suspension.

59. The formulation as defined in claim 58, which comprises:

10 (a) aripiprazole having a mean particle size of 1 to 10 microns in an amount within the range from about 5 to about 20%,

(b) suspending agent in an amount within the range from about 0.5 to about 5%,

(c) bulking agent in an amount within the range from about 4 to about 5%, and

15 (d) buffering agent in an amount within the range from about 0.03 to about 1% to adjust the pH of the sterile injectable formulation within the range from 6 to 7.5,

wherein all of the above % are % by weight based on the volume of the sterile injectable formulation.

60. The formulation as defined in any one of claims 55 to 59, further comprising a
20 pH adjusting agent.

61. The formulation as defined in any one of claims 57 to 59, wherein the suspending agent is carboxymethylcellulose or its sodium-salt, hydroxypropyl cellulose, carboxymethyl cellulose, hydroxypropylethyl cellulose, hydroxypropylmethyl cellulose, or

75535-6

polyvinylpyrrolidone; the bulking agent is mannitol, sucrose, maltose, lactose, xylitol or sorbitol; and the buffer is sodium phosphate, potassium phosphate or TRIS buffer.

62. The formulation as defined in any one of claims 57 to 61, which comprises:

(a) aripiprazole,

5 (b) carboxymethyl cellulose or its sodium salt,

(c) mannitol,

(d) sodium phosphate to adjust pH to 7,

(e) optionally sodium hydroxide to adjust pH to 7, and

(f) water for injection.

10 63. The formulation as defined in any one of claims 55 to 62, wherein the aripiprazole has a mean particle size within the range from 2 to 4 microns.

64. The formulation as defined in any one of claims 55 to 63, wherein the aripiprazole has a mean particle size of 2.5 microns.

15 65. The formulation as defined in any one of claims 55 to 64, which contains from 200 to 400 mg of aripiprazole per mL.

66. The formulation as defined in any one of claims 55 to 65, wherein the aripiprazole is in anhydrous form or in the form of a monohydrate.

67. The formulation as defined in claim 66, wherein the aripiprazole is in the form of Aripiprazole Anhydride Crystals B or Aripiprazole Hydrate A.

20 68. The formulation as defined in claim 66, wherein the aripiprazole is in the form of a monohydrate.

75535-6

69. The formulation as defined in claim 66, wherein the aripiprazole is Aripiprazole Hydrate A.

70. The formulation as defined in any one of claims 55 to 69, which upon injection releases aripiprazole over a period of about two weeks.

5 71. The formulation as defined in any one of claims 55 to 69, which upon injection releases aripiprazole over a period of about three weeks.


72. The formulation as defined in any one of claims 55 to 69, which upon injection releases aripiprazole over a period of about four weeks.

73. The formulation as defined in any one of claims 55 to 72, which upon injection
10 maintains a substantially constant aripiprazole plasma concentration over the period of release.

74. The formulation as defined in any one of claims 55 to 73, which is an intramuscular formulation.


1/3

FIG. 1



2/3

FIG. 2



3/3

FIG. 3

Preliminary Human PK Results for Aripiprazole IM DEPOT

