

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2012307006 B2**

(54) Title
Plants having enhanced nitrogen efficiency

(51) International Patent Classification(s)
C12N 15/82 (2006.01) **C12N 5/10** (2006.01)
A01H 5/00 (2006.01) **C12N 15/113** (2010.01)
A01H 5/10 (2006.01) **C12N 15/54** (2006.01)
C12N 5/04 (2006.01)

(21) Application No: **2012307006** (22) Date of Filing: **2012.09.07**

(87) WIPO No: **WO13/033846**

(30) Priority Data

(31)	Number	(32)	Date	(33)	Country
	61/532,016		2011.09.07		US

(43) Publication Date: **2013.03.14**

(44) Accepted Journal Date: **2017.08.03**

(71) Applicant(s)
Bayer Cropscience AG

(72) Inventor(s)
Good, Allen;Lock, Yee Ying;Beatty, Perrin H.

(74) Agent / Attorney
Davies Collison Cave Pty Ltd, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000, AU

(56) Related Art
US 7390937 B2
WO 2007076115 A2
GAO C ET AL, "Evolutionary and expression study of the aldehyde dehydrogenase (ALDH) gene superfamily in rice (Oryza sativa)", GENE, vol. 431, no. 1-2, pages 86 - 94.



(51) International Patent Classification:

C12N 15/82 (2006.01) C12N 15/54 (2006.01)
A01H 5/00 (2006.01) C12N 5/10 (2006.01)
A01H 5/10 (2006.01) C12N 5/04 (2006.01)
C12N 15/113 (2010.01)

(21) International Application Number:

PCT/CA2012/050622

(22) International Filing Date:

7 September 2012 (07.09.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/532,016 7 September 2011 (07.09.2011) US

(72) Inventors; and

(71) Applicants : **GOOD, Allen** [CA/CA]; 5727 - 107 Street,
Edmonton, Alberta T6H 2X6 (CA). **LOCK, Yee Ying**
[MY/CA]; Suite 3B, 8918-112 ST NW, Edmonton, Alberta
T6G2C5 (CA). **BEATTY, Perrin H.** [CA/CA]; 11615 25
Avenue, Edmonton, Alberta T6J 3R8 (CA).

(74) Agents: **SECHLEY, Konrad** et al.; Gowling Lafleur
Henderson LLP, 550 Burrard Street, Suite 2300, Van-
couver, British Columbia V6C 2B5 (CA).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

[Continued on next page]

(54) Title: PLANTS HAVING ENHANCED NITROGEN EFFICIENCY




FIG. 3

(57) Abstract: A transgenic plant comprising a polynucleotide encoding a nitrogen utilization protein operably linked to a PBpr1 promoter is provided. The transgenic plant exhibits increased nitrogen use efficiency, increased biomass and/or increased seed yield. Seeds from such transgenic plants, genetic constructs to prepare such plants, methods of generating and growing transgenic plants are also provided.



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, — *with sequence listing part of description (Rule 5.2(a))*
ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

PLANTS HAVING ENHANCED NITROGEN EFFICIENCY

FIELD OF INVENTION

[0001] The present invention relates to monocot plants having enhanced yield and/or nitrogen utilization efficiency (NUE), to methods for enhancing yield and NUE in monocot plants, and to methods of increasing biomass and seed yield in monocot plants, for example, when grown under nitrogen limiting conditions.

BACKGROUND OF THE INVENTION

[0002] The productivity of plants is limited by the three primary nutrients: nitrogen, phosphorous and potassium, in most natural and agricultural ecosystems. Generally nitrogen is the most important of the three limiting nutrients and the major component in fertilizers. Since nitrogen is usually the rate-limiting element in plant growth, most field crops have a fundamental dependence on inorganic nitrogenous fertilizer. The nitrogen source in fertilizer is usually ammonium nitrate, potassium nitrate, or urea.

[0003] Increased nitrogen use efficiency by plants has a number of beneficial effects, for example, increased growth and yield when compared to conventional plants grown in nitrogen poor soils, and reduced requirement for the addition of nitrogenous fertilizers to crops. Fertilizers account for a significant percentage of the costs associated with crop production, therefore using less fertilizer would reduce the producers' costs. A reduction in fertilizer application would also lessen the environmental damage resulting from extensive nitrogenous fertilizer use. Excess fertilizer application causes increased eutrophication, acid rain, soil acidification and the greenhouse effect. These environmental disasters cause further problems such as fish kills, loss of biodiversity, increased algal blooms, loss of arable land and accelerated global climate change, affecting the world population on both social and economic scales.

[0004] Monocots represent a large percentage of the crops grown in the world with approximately 217 million hectares of wheat and 158 million hectares of both maize and rice planted in 2007. Approximately half of the global calorie and protein requirement is derived from wheat, rice and maize. Rice is routinely used as a model

crop for genetic and physiological studies in other monocot crops including maize, wheat, sugarcane, barley, sorghum, rye and grass. Rice has a small, diploid genome that is well conserved and syntenic across monocots.

[0005] Promoters are nucleic acid sequences that allow for regulation of transcription of a gene or nucleotide sequence. Promoters can allow for constitutive expression, such as the well known cauliflower mosaic virus promoter CaMV 35S, inducible expression, such as the stress inducible promoter *rd29A* (Pino et al., 2007), tissue specific expression, such as root specific *OsANT1* (U.S. Patent Application Publication No. 2009/0288224) and developmentally specific expression, such as the senescence induced *IPT* promoters (Ma et al., 2009). Promoters can also be weak or strong, suggesting that whenever or wherever they are induced, they will allow for expression of the attached gene or nucleotide sequence at varying levels.

[0006] When a promoter is fused to the 5' end of a gene or nucleotide sequence, it can regulate the expression of that gene or nucleotide sequence. However not all promoters will successfully express all genes or nucleotide sequences in all plant types. For example, a dicot promoter may not perform in the same manner when placed in a monocot system, and vice versa. Similarly, a monocot promoter may not function in the same manner when placed in a monocot of a different Genus, such as a rice promoter placed into wheat. For transgenic studies, there are different types of promoters that can be used, depending on the goal of the experiment. Promoters are often classified as constitutive, tissue specific and/or inducible. Many transgenic studies to date have used generic constitutive promoters such as the cauliflower mosaic virus (*CAMV35S*) and maize ubiquitin 1 promoter (*ubi-1*) to drive target gene over-expression in plants. This can be a disadvantage as it could be energetically unfavorable for plants to express the gene at all times and it could produce abnormal development, since expression levels of the transgene is not regulated (Shelton *et al.*, 2002). For example, constitutive over-expression of the cellulose synthase like gene *CsLF6* by the oat globulin promoter *ProASGL* frequently resulted in reduced germination of seeds or seedlings with necrosis on the leaf tips leading to death in severe cases (Burton *et al.*, 2011).

[0007] Using inducible or tissue specific promoters may be a better option to drive transgene expression. Inducible promoters will only drive gene expression when a specific physical, environmental, biological or chemical stimulus is applied. The heat inducible promoter of the *Hvhsp17* gene from wheat can be used for high target gene expression when plants are exposed to 38 to 40°C for 1 to 2 hours (Freeman *et al.*, 2011). This allows for the short term gene or nucleotide sequence expression and control of developmental expression but is limited to tissues that are not severely affected by temperature changes. Over-expression of the *Triticum aestivum* NAC protein (*TaNAC69*), encoding a transcription factor involved in drought stress by the drought inducible promoter *HvDhn4s*, produced wheat plants with significantly higher shoot biomass at the early vegetative stage under mild salt stress and water limitation, compared to the wild-type. Conversely, the *HvDhn8s* constitutive promoter driving *TaNAC69* showed no significant difference from untransformed controls (Xue *et. al.*, 2011). When the drought and cold stress inducible promoter *rdA29* was used to over-express the *DREB1A* gene, which encodes a transcription factor involved in stress tolerance in *Arabidopsis*, normal plants were generated, while constitutive expression driven by the *CAMV35S* promoter resulted in growth retardation under normal growing conditions (Kasuga *et al.*, 1999).

[0008] Tissue specific promoters are involved in organ specific and developmental stage specific expression of transgenes. For example, in potatoes, *StRCap* is engineered into the leaves as a defense mechanism against predatory insects, but is not expressed in the eating parts of the plant that produces this toxin (Weber, 2003; Park and Jones, 2008). The use of the 35S promoter has also raised concerns of food safety where the toxin produced in non-target organs might cause the potatoes to be unsafe for consumption. In addition, it may be metabolically taxing for the plant to be constantly producing a secondary metabolite, regardless of the developmental stage or organ, therefore causing plants to be less healthy and potentially compromising yields.

[0009] In the case of NUE plant engineering, tissue specific expression of genes might increase the efficacy of N uptake, utilization or remobilization in the plant. In contrast, the use of a constitutive promoter might prove to be a waste of energy

because over-expression of non-rate limiting enzymes in certain organs may not produce any phenotype or may even decrease yield. Development of an NUE plant may also involve the transformation of multiple genes or gene stacking to achieve a satisfactory NUE phenotype, since N metabolism and transport are very complex processes.

[0010] In addition, the use of the *CAMV35S* promoter may not produce any phenotype because the gene expression or protein expression was not sufficient in a specific organ or developmental phase. Also, plants could turn off the expression of the transgene when it proves to be energetically unfavourable. When *HvAlaAT* was driven by the *CAMV35S* promoter, it did not exhibit any NUE phenotype. However, when the root specific *btg26* promoter was used, it produced plants that had higher NUE (Good *et al.*, 2007).

[0011] Similar to over-expression of target genes, promoters when used in a different species may not mimic the expression patterns of its native species. Seed specific promoters from barley (*B-hor* and *D-hor*) and wheat (*HMW-Glu*) did not direct seed specific expression in rice, instead the promoters drove high expression levels in leaf, shoot and maternal seed tissues of rice plants (Wu *et al.*, 1998; Qu and Takaiwa, 2004; Furtado *et al.*, 2008).

[0012] PBpr1 promoters are from genes encoding methyl-melonate semialdehyde dehydrogenase (MMSDH). The *PBpr1* promoter is upstream of the *OsALDH6* gene, which encodes a gene for methylmalonate semialdehyde dehydrogenase in rice (Accession number: gene: AK 121280.1 and mRNA: AF045770.1). AK121280.1 and AF045770.1 are splice variants of each other where AF045770 is shorter than the AK121280.1. AK121280.1 appears to be a hypothetical protein designated by GenBank, while AF045770.1 has been characterized (Oguchi *et al* (2004). *OsALDH6* is homologous to the *ALDH6B2* gene in *Arabidopsis*, which also encodes for a methylmalonate semialdehyde dehydrogenase. The *OsALDH6* gene is highly expressed in young roots and stems (Gao and Han, 2009).

[0013] Methylmalonate semialdehyde dehydrogenase (MMSDH) catalyzes the irreversible oxidative decarboxylation of malonate semialdehydes to acetyl-CoA and

methylmalonate-semialdehyde to propionyl-CoA in the distal portions of the valine and pyrimidine catabolic pathways. Since MMSDH generates acetyl-CoA, it is an important factor in the glyoxylate pathway, TCA cycle and fatty acid production.

[0014] MMSDH is down-regulated during oxidative stress due to the restriction in the TCA cycle and the production of ATP (Sweetlove *et al.*, 2002). In two week old rice plants, MMSDH mRNA was found at high levels in roots and leaf sheaths while protein accumulation was found highest in roots followed by leaf blades (Oguchi *et al.*, 2004). With the addition of auxin, MMSDH levels in roots increased along with an increase in rooting. Tanaka *et al.* (2005) has suggested that MMSDH is involved in root growth, tissue differentiation and thickening growth due to its expression in crown roots, lateral roots and root hairs. MMSDH was found to be localized in the mitochondrial matrix of Arabidopsis, rice, human, bovine and rats which suggests a similarity of function of MMSDH among all these organisms.

[0015] This background information is provided for the purpose of making known information believed by the applicant to be of possible relevance to the present invention. No admission is necessarily intended, nor should be construed, that any of the preceding information constitutes prior art against the present invention.

SUMMARY OF THE INVENTION

[0016] An object of the invention to provide plants having enhanced nitrogen efficiency. In accordance with one aspect of the present invention, there is provided a transgenic plant or plant part comprising a polynucleotide encoding a nitrogen utilization protein operably linked to a PBpr1 promoter.

[0017] In accordance with another aspect, there is provided a seed obtained from a transgenic plant of the invention.

[0018] In accordance with another aspect of the invention, there is provided a plant cell transformed with a polynucleotide encoding a nitrogen utilization protein operably linked to a PBpr1 promoter.

[0019] In accordance with another aspect of the invention, there is provided a genetic construct comprising a polynucleotide encoding a nitrogen utilization protein operably linked to a PBpr1 promoter.

[0020] In accordance with another aspect, there is provided a vector comprising a genetic construct of the invention.

[0021] In accordance with another aspect, there is provided a host cell comprising a genetic construct or a vector according of the invention.

[0022] In accordance with another aspect of the invention, there is provided a method of generating a plant having increased nitrogen use efficiency comprising: transforming a plant cell with a genetic construct or vector of the invention, and growing the transformed plant cell to produce a plant that expresses the nitrogen utilization protein from the PBpr1 promoter, thereby generating a plant having increased nitrogen use efficiency, wherein the increased nitrogen use efficiency is relative to the nitrogen use efficiency of a wild-type plant grown under identical conditions.

[0023] In accordance with another aspect of the invention, there is provided a method of generating a plant having increased biomass comprising:
transforming a plant cell with a genetic construct or vector of the invention, and growing the transformed plant cell to produce a plant that expresses the nitrogen utilization protein from the PBpr1 promoter, thereby generating a plant having increased biomass, wherein the increased biomass is relative to the biomass of a wild-type plant grown under identical conditions.

[0024] In accordance with another aspect of the invention, there is provided a method of generating a plant having increased seed yield comprising: transforming a plant cell with a genetic construct or vector of the invention, and growing the transformed plant cell to produce a plant that expresses the nitrogen utilization protein from the PBpr1 promoter, thereby generating a plant having increased seed yield, wherein the increased seed yield is relative to the seed yield of a wild-type plant grown under identical conditions.

[0025] In accordance with another aspect of the invention, there is provided a method for producing a plant having increased nitrogen use efficiency, increased biomass, increased seed yield, or a combination thereof, comprising: (a) providing a plant, a plant part or a seed comprising a genetic construct or vector of the invention, and (b) growing a plant from the plant, plant part or seed, thereby producing a plant having increased nitrogen uptake, increased biomass, increased seed yield, or a combination thereof, wherein the increased nitrogen uptake, increased biomass and increased seed yield is relative to the nitrogen uptake, biomass and seed yield of a wild-type plant grown under identical conditions.

[0026] In accordance with another aspect, there is provided a use of a transgenic plant of the invention for producing progeny.

[0027] In accordance with another aspect of the invention, there is provided a use of a genetic construct or vector of the invention to increase nitrogen use efficiency, biomass or seed yield in a plant.

[0028] In accordance with another aspect of the invention, there is provided a process for increasing nitrogen use efficiency, biomass, seed yield, or a combination thereof, in a plant comprising: providing a seed comprising a genetic construct or the vector of the invention, applying a herbicide, insecticide, fertilizer, or combination thereof, to the seed, and growing a plant from the seed, wherein the plant has increased nitrogen use efficiency, increased biomass, increased seed yield, or a combination thereof, and wherein the increased nitrogen use efficiency, increased biomass or increased seed yield is relative to the nitrogen use efficiency, biomass or seed yield of a wild-type plant grown under identical conditions.

[0029] In certain embodiments, the present invention provides a method of generating the plant with increased nitrogen use efficiency comprising, introducing a nucleic acid encoding the one or more nitrogen utilization protein operably linked to a root-epidermis-specific promoter, and producing the plant, the plant comprising elevated levels of one or more nitrogen utilization protein specifically localized to the root epidermis.

[0030] In certain embodiments, the present invention provides a method for directing tissue-specific expression of a target gene or nucleotide sequence in a plant, including producing a plant from a transformed plant cell such that tissue-specific expression of a target gene or nucleotide sequence occurs within a selected tissue of the plant, wherein the transformed plant cell contains a target gene or nucleotide sequence in operative linkage with a *PBpr1* promoter element. In one embodiment, the tissue-specific expression takes place in the root of the plant.

[0031] In certain embodiments, the present invention provides a method for directing tissue-specific and environmental or developmentally-regulated expression of a target gene or nucleotide sequence in a plant, including producing a plant from a transformed plant cell such that tissue-specific and environmentally or developmentally-regulated expression of a target gene or nucleotide sequence occurs within a selected tissue of the plant, wherein the transformed plant cell contains a target gene or nucleotide sequence in operative linkage with a *PBpr1* promoter element. In one embodiment, the tissue-specific expression takes place in the root or leaves of the plant and the environmentally or developmentally-regulated expression takes place under conditions of nutrient stress.

[0032] In certain embodiments, the present invention provides a method for increasing biomass of a plant growing under one or more environmentally adverse conditions comprising; transforming a plant with a target gene or nucleotide sequence in operative linkage with a *PBpr1* promoter element to produce a transformed plant, the target gene or nucleotide sequence encoding an enzyme involved in nitrogen assimilation or metabolism; and growing the transformed plant. In one embodiment, the enzyme is alanine aminotransferase (AlaAT).

[0033] In certain embodiments, the present invention provides a method for increasing biomass of a plant growing under conditions of low nitrogen comprising; transforming a plant with a target gene or nucleotide sequence in operative linkage with a *PBpr1* promoter element to produce a transformed plant, the target gene or nucleotide sequence encoding an enzyme involved in nitrogen assimilation; and growing the transformed plant. In one embodiment, the enzyme is AlaAT.

[0034] In certain embodiments, the present invention provides a method for increasing the biomass of a plant growing under one or more environmentally adverse conditions comprising; transforming a plant with a gene or nucleotide sequence encoding AlaAT in operative linkage with a promoter element to produce a transformed plant; and growing the transformed plant. In one embodiment, the promoter is *PBpr1*.

[0035] In certain embodiments, the present invention provides a method for increasing seed yield of a plant comprising; transforming the plant with a target gene or nucleotide sequence or nucleotide sequence in operative linkage with a *PBpr1* promoter element, to produce a transformed plant, the target gene or nucleotide sequence or nucleotide sequence encoding an enzyme involved in nitrogen assimilation or metabolism; and growing the transformed plant. In one embodiment, the enzyme is AlaAT.

[0036] In certain embodiments, the present invention provides a method for increasing seed yield of a plant growing under conditions of low nitrogen comprising; transforming a plant with a target gene or nucleotide sequence or nucleotide sequence in operative linkage with a *PBpr1* promoter element to produce a transformed plant, the target gene or nucleotide sequence or nucleotide sequence encoding an enzyme involved in nitrogen assimilation; and growing the transformed plant. Preferably, the enzyme is AlaAT.

[0037] In certain embodiments, the present invention provides a method for increasing seed yield of a plant growing under conditions of high nitrogen comprising; transforming a plant with a target gene or nucleotide sequence or nucleotide sequence in operative linkage with a *PBpr1* promoter element to produce a transformed plant, the target gene or nucleotide sequence or nucleotide sequence encoding an enzyme involved in nitrogen assimilation; and growing the transformed plant. In one embodiment, the enzyme is AlaAT.

[0038] This summary of the invention does not necessarily describe all features of the invention and is not intended to be limiting.

BRIEF DESCRIPTION OF THE DRAWINGS

[0039] These and other features of the invention will become more apparent from the following description in which reference is made to the appended drawings.

[0040] FIGURE 1 shows a construct map of *PBpr1::HvAlaAT* in pCAMBIA1300. The *HvAlaAT* is driven by *PBpr1* promoter and terminated by a nos terminator. The selection marker is kanamycin for plasmid selection and hygromycin.

[0041] FIGURE 2 shows a comparison of total above ground biomass (g) and seed yield (g) of T₁ PBpr1 lines, NB and AGR1/7 at maturation. Each measurement is from 5 to 7 replications and error bars are standard deviations. Black bars: Total above ground biomass (g); Grey bars: Total seed yield (g).

[0042] FIGURE 3 shows the number of tillers and productive tillers produced by T₁ PBpr1 plants compared to NB and AGR1/7 at maturation. Each measurement refers to an average of at least 4 replications and error bars show SD. Black bars: Total number of tillers; Grey bars: Number of seed producing tillers.

[0043] FIGURE 4 shows a comparison between T₂ PBpr1 plants, NB and AGR1/7 above ground biomass (g) and seed yield (g) at maturity. Values from each line are from 7 replications and error bars represents SD. Black bars: Total above ground biomass (g); Grey bars: Total seed yield (g).

[0044] FIGURE 5 shows (A) T₃ PBpr1 plant dry above ground biomass (g) at 52 DAG, each line was represented by 5 replication plants and error bars are SD, and (B) the fresh weight (g) of three *PBpr1:AlaAT* over-expressing rice lines versus WT (NB; Nipponbare) at maturity during soil growth conditions.

[0045] FIGURE 6 shows shoot specific AlaAT activity in $\mu\text{M NADH/min.mg}$ of protein of T₁ PBpr1 lines grown in soil at 40 days after germination (DAG) compared to NB and AGR1/7. All lines labeled with N are null segregants. Each data point for the transgenic plants was taken from an average of 3 replications and error bars refer to SD of the 3 replicates.

[0046] FIGURE 7 shows alanine aminotransferase activity ($\mu\text{M NADH}/\text{min} \cdot \text{mg}$ of protein) of WT (NB), AGR 1/7 (*OsANT1:AlaAT*) and three independent PBpr1:AlaAT over-expressing T₃ rice lines (PBpr1-11, PBpr1-12, PBpr1-21) at five (5) different growth points during soil growth under high N conditions. Error bars refer to SD.

[0047] FIGURE 8 shows T₂ PBpr1 lines shoot AlaAT activity in $\mu\text{M NADH}/\text{min} \cdot \text{mg}$ of protein at 28 and 52 days old plant that were grown hydroponically. PBpr1-21-2 and PBpr1-21-3 are siblings of the same line. Each data point referred to 5 replications and error bars refer to their SD.

[0048] FIGURE 9 shows T₂ PBpr1 lines root AlaAT activity at 28 and 52 days old plant grown hydroponically. Each data point referred to at least 5 replications and error bars refer to their SD. PBpr1-21-2 and PBpr1-21-3 are siblings of the same line. Black bars: 28 day old plants; grey bars: 52 day old plants.

[0049] FIGURE 10 shows an immunoblot analysis of protein extracts of 52 Day old T₃ PBpr1 lines detected with *HvAlaAT*-2 specific 1^o antibody serum. Lane 1: AGR1/7, lane 2: wild-type Nipponbare, lane 3: PBpr1-10, lane 4: PBpr1-11, lane 5: PBpr1-11-2N, lane 6: PBpr1-12 and lane 7: PBpr1-21. Protein concentration was standardized across all lanes by using the Bradford protein quantifaction assay.

[0050] FIGURE 11 shows an immunoblot analysis of protein extract from T₂ hydroponically grown roots (A) and shoots (B) of 28 DAG rice lines with *HvAlaAT* 1^o antibody detection to show amounts of *HvAlaAT* protein in each sample. Lane 1: PBpr1-11, lane 2: PBpr1-11-2N, lane 3: PBpr1-21-2, lane 4: PBpr1-21-3, lane 5: PBpr1-21-3 replicate, lane 6: PBpr1-21-N, lane 7: AGR1/7 and lane 8: wild-type Nipponbare. Protein concentration was standardized across all lanes by using the Bradford protein quantifaction assay.

[0051] FIGURE 12 shows a Log₂ of the relative quantification (ddCT) PBpr1 shoots, T₂ PBpr1 shoots and T₂ PBpr1 roots, AGR1/7 and null segregants of the PBpr1 lines relative to NB plants. 7 transcripts (*HvAlaAT*: Barley AlaAT, *GRP*: Glycine rich protein, *LRR*: Leucine rich repeat) were measured by qRT-PCR. Plants were raised in hydroponic conditions and harvested at 52 days after germination.

Tissue samples used were the same with those of the AlaAT activity assays in Figure 9. (2 fold change represents 1 ddCT. $\text{Log}_2 2 = 1$).

[0052] FIGURE 13 shows the nucleotide sequence of the PBpr1 promoter including the ATG start and initial nucleotides of the coding sequence (SEQ ID NO:6) and showing the *EcoRI* and *SacI* restriction enzyme sites used for cloning.

[0053] FIGURE 14 shows (A) the nucleotide sequence of the barley (*Hordeum vulgare*) AlaAT gene (SEQ ID NO:2) and (B) the amino acid sequence of the AlaAT protein (SEQ ID NO:3).

[0054] FIGURE 15 shows GUS staining of (A) 3 day old *PBpr1::GUSplus* (PBpr1G line) seedling, (B) 3 day old *OsANT1::GUSplus* (*OsANT1::GUS* lines) seedling as a positive control. (C) to (M) are GUS stained tissues from 7 to 10 day old seedlings of PBpr1G and *OsANT1::GUS* lines. (C) stem of PBpr1G, (D), (G), and (H) basal region of the shoot/shoot root interface containing remnant seed PBpr1G, (E) basal region of the shoot/shoot root interface containing remnant seed *OsANT1GUS*, (F) stem of *OsANT1GUS*, (I) leaf sheath and leaf blade interface *OsANT1GUS*, (J) leaf blades and veins PBpr1G, (K) leaf blades and veins *OsANT1GUS*, (L) roots and root hairs PBpr1G, and (M) roots and root hairs *OsANT1GUS*.

[0055] FIGURE 16 presents a table showing the foliar amino acid concentrations of wildtype (cv. Nipponbare) rice versus barley *AlaAT* over-expressing rice transgenic rice lines; AGR 1/7 (*OsANT1:HvAlaAT*) and PBpr1-11, PBpr1-12, PBpr1-21 (*OsPBpr1::HvAlaAT*). Amino acid concentration (nM/g FW) are averages of 4 replicates.

[0056] FIGURE 17 shows (A) PBpr2 promoter driving expression of AlaAT in rice, T1 generation, grown in soil growth chamber; comparison of plant height, total biomass and seed weight to wild-type rice cv. Nipponbare (NB), and (B) PBpr2 promoter driving expression of AlaAT in rice, T2 generation, grown in soil growth chamber; comparison of total biomass and seed weight to wild-type rice cv. Nipponbare (NB).

[0057] FIGURE 18 shows (A) the nucleotide sequence of the PBpr1 promoter (SEQ ID NO:1), and (B) the nucleotide sequence of the PBpr2 promoter (SEQ ID NO:4).

[0058] FIGURE 19 shows an alignment of the nucleotide sequences of the OsANT1 promoter (SEQ ID NO:5), the PBpr1 promoter (SEQ ID NO:1) and the PBpr2 promoter (SEQ ID NO:4).

DETAILED DESCRIPTION OF THE INVENTION

[0059] The present invention relates to monocot plants having enhanced yield and/or nitrogen utilization efficiency (NUE), to methods for enhancing yield and NUE in monocot plants, and to methods of increasing biomass and seed yield in monocot plants. This invention also relates to PBpr1 promoters, which are promoters from genes encoding methyl-melonate semialdehyde dehydrogenase (MMSDH).

[0060] Given the worldwide requirements for monocots and the diminishing fertility of existing fields, it is desirable to generate monocot plants that are able to grow under suboptimal nutrient conditions. One means for accomplishing this goal is to generate monocot plants that can use nitrogen more efficiently. Such monocot plants have the advantage of being able to grow in soils that are poorer in nitrogen, as a result of being able to more efficiently use the nitrogen that is available, with no loss in yield. Additionally, such monocot plants may also demonstrate enhanced yield in soils that have normal nitrogen levels as well. Nitrogen use efficiency in plants is a result of two main subcomponents; N uptake efficiency and N utilization efficiency. A plant exhibiting a nitrogen use efficiency phenotype may have improvements in its' ability to take up nitrogen from the soil, which is a desired trait in plants growing in lower nutrient-available soils. Or the NUE plant may have improvements in its' ability to utilize the N that it has taken up so that the available N (whether it be low or high) is efficiently incorporated into the subcellular components (such as nucleic acids, proteins, storage etc.), translocated to the necessary tissues and remobilized at the correct developmental stage into seed. Or, another possibility is that the NUE plant has improvements in both N uptake and utilization. Any one of these possibilities would allow for an increased yield from a NUE crop grown in normal nitrogen conditions because those plants would be able to take up a non-

limiting amount of nitrogen and be able to use the nitrogen to increase biomass and seed yield, either by increased number of seeds or an increase in seed weight, or both. Certain embodiments of the invention provide for transgenic plants having improved nitrogen utilization efficiency compared to wild-type plants.

5 [0061] Plants face a wide variety of non-optimal environmental conditions during growth and development. Such conditions may include water limitation, excess salinity, alkaline or acidic soil, infestation by pests, disease, or temperature stress, any of which individually may significantly hinder crop growth and/or yields. Certain
10 embodiments of the present invention provide methods by which plants, and seeds thereof, may be engineered to grow and thrive under changing environmental conditions usually not conducive to the development of the plant. In some embodiments, methods are provided to develop plants that can maintain or increase their biomass and yield while being grown under conditions that are not optimal, in terms of nutrient applications are described.

15 [0062] In certain embodiments, the present invention advantageously allows the user to develop plants that have an environmental benefit in that they can maintain yield, while reducing the need for high levels of nutrient application. In certain embodiments, the present invention allows the user to develop plants that, under high nutrient levels, have improved nutrient up-take allowing the plants to extract more
20 nutrients from their environment during times of nutrient abundance. Using the methods and compositions of certain embodiments of the invention, plants may be improved for growth and development under environmental conditions usually unsuitable for growth of the plant. In some embodiments, the methods and compositions of the invention permit the genetic engineering of a plant to alter one or
25 more plant characteristics in only selected tissues of the plant.

[0063] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

30 [0064] As used herein, the terms “comprising,” “having,” “including” and “containing,” and grammatical variations thereof, are inclusive or open-ended and do

not exclude additional, unrecited elements and/or method steps. The term “consisting essentially of” when used herein in connection with a composition, use or method, denotes that additional elements and/or method steps may be present, but that these additions do not materially affect the manner in which the recited composition, method or use functions. The term “consisting of” when used herein in connection with a composition, use or method, excludes the presence of additional elements and/or method steps. A composition, use or method described herein as comprising certain elements and/or steps may also, in certain embodiments consist essentially of those elements and/or steps, and in other embodiments consist of those elements and/or steps, whether or not these embodiments are specifically referred to.

[0065] In addition, the use of the singular includes the plural, and “or” means “and/or” unless otherwise stated.

[0066] As used herein, the term “about” refers to an approximately +/-10% variation from a given value. It is to be understood that such a variation is always included in any given value provided herein, whether or not it is specifically referred to.

[0067] The term “substantially identical” as used herein in relation to a nucleotide or amino acid sequence indicates that, when optimally aligned, for example using the methods described below, the nucleotide or amino acid sequence shares at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity with a defined second nucleotide or amino acid sequence (or “reference sequence”). “Substantial identity” may be used to refer to various types and lengths of sequence, such as full-length sequence, functional domains, coding and/or regulatory sequences, promoters, and genomic sequences. Percent identity between two amino acid or nucleotide sequences can be determined in various ways that are within the skill of a worker in the art, for example, using publicly available computer software such as Smith Waterman Alignment (Smith, T. F. and M. S. Waterman (1981) *J Mol Biol* 147:195-7); “BestFit” (Smith and Waterman, *Advances in Applied Mathematics*, 482-489 10 (1981)) as incorporated into GeneMatcher Plus™, Schwarz and Dayhof (1979) *Atlas of Protein Sequence and Structure*, Dayhof, M. O., Ed pp 353-358; BLAST program (Basic

Local Alignment Search Tool (Altschul, S. F., W. Gish, *et al.* (1990) *J Mol Biol* 215: 403-10), and variations thereof including BLAST-2, BLAST-P, BLAST-N, BLAST-X, WU-BLAST-2, ALIGN, ALIGN-2, CLUSTAL, and Megalign (DNASTAR) software. In addition, those skilled in the art can determine appropriate parameters for measuring alignment, including algorithms needed to achieve maximal alignment over the length of the sequences being compared. In general, for amino acid sequences, the length of comparison sequences will be at least 10 amino acids. One skilled in the art will understand that the actual length will depend on the overall length of the sequences being compared and may be at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100, at least 110, at least 120, at least 130, at least 140, at least 150, or at least 200 amino acids, or it may be the full-length of the amino acid sequence. For nucleotide sequences, the length of comparison sequences will generally be at least 25 nucleotides, but may be at least 50, at least 100, at least 125, at least 150, at least 200, at least 250, at least 300, at least 350, at least 400, at least 450, or at least 500, or it may be the full-length of the nucleotide sequence.

[0068] The term “tissue-specific expression” of a target nucleotide sequence is known in the art and includes the expression of a target nucleotide sequences predominantly in selected tissues. Tissue-specific expression may result in expression of the target nucleotide sequence only in the selected tissue, in which case the target nucleotide sequence may be present in multiple tissues, but is expressed in only a subset of those tissues. Alternatively, tissue-specific expression may result in higher levels of expression of the target nucleotide sequence in selected tissues, in which case expression may also be observed in other tissues, but at a lower level than in the target tissue(s), for example, 50% lower, 40% lower, 30% lower, 25% lower or 20% lower. Such selective expression may be due to the influence of one or more regulatory genetic elements, for example but not limited to promoter elements, repressor elements, enhancer elements, or other regulatory factors that may interact with DNA, RNA.

[0069] The terms “target gene” or “target nucleotide sequence” are art-recognized, and include nucleotide sequences which are desired to be expressed in one or more

selected plant tissues. Non-limiting examples of target genes or nucleotide sequences which may be utilized in conjunction with the methods of the invention include genes or nucleotide sequences involved in nitrogen assimilation and/or utilization, genes or nucleotide sequences involved stress resistance, genes or nucleotide sequences
5 disease and pest resistance, and genes or nucleotide sequences involved in nutrient uptake and utilization. Such genes or nucleotide sequences are well known to one of skill in the art.

[0070] The terms “operative linkage,” “operatively linked” and “operably linked” are used interchangeably herein to refer to the placement of a target nucleotide
10 sequence relative to a nucleic acid regulatory sequence such that the expression of the target nucleotide sequence is controlled by the regulatory sequence. This regulatory sequence can have a positive effect (increase) on the expression of the target gene or nucleotide sequence (*e.g.*, the regulatory sequence is a promoter or an enhancer element), or the regulatory sequence can reduce the expression of the target gene or
15 nucleotide sequence (*e.g.*, the regulatory sequence is a repressor element). The regulatory sequence may be physically located 5’ or 3’ of the target gene or nucleotide sequence, may be within the coding sequence of the target gene or nucleotide sequence, or may be contained on an intron within the target gene or nucleotide sequence.

[0071] The term “nulls” is art-recognized and includes a plant that has undergone
20 tissue culturing but does not carry a transgene or selectable marker.

[0072] Production of plants which express one or more target genes or nucleotide sequences in a tissue-specific fashion are described herein. In certain embodiments, invention provides seeds containing one or more target genes or nucleotide sequences
25 under the control of a promoter element which specifically directs the tissue-specific expression of the gene or nucleotide sequence. In certain embodiments, the methods of the invention allow production of plants having one or more desired traits or properties in selected tissues; *e.g.*, to alter specifically the genetic and/or physiological properties of the fruit or the roots of the plant. Certain embodiments of
30 the invention further provide methods of producing plants having root and

leaf-specific expression of one or more desired nucleotide sequences using the *PBpr1* promoter element.

[0073] The methods of the invention for the production of plants having tissue-specific expression of one or more target genes or nucleotide sequences are accomplished through the use of a genetic regulatory element which directs the tissue-specific expression of the target gene(s). Regulatory elements may be either negative or positive in activity: a plant tissue-specific promoter or enhancer element permits the expression of the target gene or nucleotide sequence(s) in one or more specific tissues, whereas a plant tissue-specific repressor suppresses the expression of the target genes or nucleotide sequences in one or more specific tissues, while expression in the other tissue(s) continues unabated. For the purposes of the invention, it will be understood that promoter sequences constitute the preferred genetic regulatory elements of the invention.

[0074] The tissue-specific promoters useful in the present invention can be either homologous or heterologous to the plant to which they are to be used. Promoters which direct the expression of an operatively linked gene or nucleotide sequence in one or more plant tissues while excluding expression of the linked gene or nucleotide sequence in one or more other plant tissues may be used in the methods and constructs of the invention. Appropriate plant tissues include but are not limited to *e.g.*, root, leaf, petal, sepal, stamen, anther, stigma, ovary, style, pistil, epidermis, phloem, xylem, cortex, pith, cambium, stem or trunk, root hair, petiole, fruit and tuber.

[0075] Preferably, the promoter is a *PBpr1* promoter. In accordance with certain embodiments of the invention, the promoter is a *PBpr1* promoter having a nucleotide sequence substantially identical to the sequence as set forth in SEQ ID NO:1 (Figure 13).

[0076] It will be understood by one skilled in the art that modifications may be made to the promoters useful in the methods and constructs of the invention to improve or modulate the activity of the promoter. Multiple copies of a selected promoter may be operatively linked to a single target gene or nucleotide sequence to

thereby alter the expression level of the linked gene or nucleotide sequence, or a selected promoter may be operatively linked to one or more target genes or nucleotide sequences such that the expression of each target gene or nucleotide sequence is coordinately regulated. A promoter may be of any size appropriate to permit the tissue-specific functioning of the promoter. A promoter may be modified (*e.g.*, by mutagenesis, deletion, insertion, or truncation) to alter the degree to which the operatively linked gene or nucleotide sequence is expressed in the selected tissue, or to alter the specificity of tissue expression directed by the promoter. Further, the placement of the promoter relative to the operatively linked target gene or nucleotide sequence may be modulated (*e.g.*, moved further away or closer together) to attain a desired level of promoter-directed expression.

[0077] A promoter may direct the expression of an operatively linked gene or nucleotide sequence in one or more selected tissues of the plant, and also direct expression of the gene or nucleotide sequence in response to a specific environmental or physiological condition. For example, promoters may be activated under conditions of stress (*e.g.*, drought stress, saline stress, temperature stress, oxygen stress, pH stress, or heavy metal stress), under conditions of nutrient deprivation, under conditions of attack by disease or pests. In another example, promoters may be activated under specific developmental conditions (*e.g.*, upon sprouting, fruiting, or seed production) or in response to a change in environmental conditions (*e.g.*, from nutrient deprivation to provision or from hypoxia to normoxia).

[0078] In certain embodiments, the present invention provides for constructs comprising a target gene under control of a PBpr1 promoter that, when transformed into an appropriate plant host, result in tissue-specific expression of the gene product. For example, expression may preferentially be in the roots and/or shoots of the plant. In certain embodiments, the present invention provides for constructs comprising a target gene under control of a PBpr1 promoter that, when transformed into an appropriate plant host, result in developmentally regulated expression of the gene product.

[0079] A target gene or nucleotide sequence of the invention is a gene or nucleotide sequence which it is desired to be expressed in a plant. General classes of target genes

or nucleotide sequences which may be advantageously employed in the methods and constructs of the invention include genes or nucleotide sequences encoding plant structural proteins, genes or nucleotide sequences encoding proteins involved in the transport and/or uptake of nutrients, genes or nucleotide sequences encoding enzymes and proteins involved in nutrient utilization, genes or nucleotide sequences encoding proteins involved in plant resistance to herbicides, genes or nucleotide sequences encoding proteins involved in plant resistance to nematodes, viruses, insects, or bacteria, genes or nucleotide sequences encoding proteins involved in plant resistance to stress (*e.g.*, osmotic, temperature, pH, or oxygen stress), genes or nucleotide sequences encoding proteins involved in stimulation or continuation of plant growth, or genes or nucleotide sequences encoding proteins involved in phytoremediation. Further, the target gene or nucleotide sequence may be a nucleotide sequence which, when transcribed, is antisense to a native sequence, the transcription and translation of which is desired to be suppressed.

[0080] In certain embodiments of the invention, the target genes or nucleotide sequences are those encoding enzymes in the assimilation and/or metabolism of nitrogen. The genes or nucleotide sequences of interest may include genes or nucleotide sequences which encode proteins involved in assimilating ammonia into amino acids or use the formed amino acids in biosynthetic reactions, that is, "nitrogen utilization proteins". Examples of nitrogen utilization proteins include, but are not limited to, nitrate ammonium and amino acid transporters, glutamine synthetase (GS), asparagine synthetase (AS), glutamate synthase (also known as glutamate 2:oxoglutarate amino transferase and GOGAT), asparaginase (ANS), glutamate dehydrogenase (GDH), aspartate aminotransferase (AspAT) and alanine aminotransferase (AlaAT) and those genes or nucleotide sequences which may be involved in Nutrient Use Efficiency (NUE) as described by Beatty et al. (2009), and in U.S. Patent No. 7,589,257. Sequences of appropriate target genes and nucleotide sequences are readily available from public sources, for example GenBank, or can be determined by standard techniques well-known to those of skill in the art.

[0081] The target gene or nucleotide sequence may be a gene or nucleotide sequence naturally expressed in the selected plant, or it may be heterologous to the

selected plant. The gene or nucleotide sequence may originate from a variety of sources, including viral, bacterial, plant or animal sources.

[0082] Certain embodiment of the invention provide for genetic constructs comprising the target gene or nucleotide sequence operably linked to a *PBrp1* promoter. In certain embodiments, the gene or nucleotide sequence is heterologous to the *PBrp1* promoter to which it is linked.

[0083] The target gene or nucleotide sequence can be modified as required. For example, the gene or nucleotide sequence is modified to be transcribable and translatable in a plant system, for example by codon optimization. In certain embodiments, the gene or nucleotide sequence can be modified such that it contains all of the necessary poly-adenylation sequences, start sites and termination sites which allow the coding sequence to be transcribed into messenger ribonucleic acid (mRNA) and the mRNA to be translated into a functional protein in the selected plant system.

[0084] Certain embodiments of the invention thus provide for genetic constructs comprising the target gene or nucleotide sequence operably linked to a *PBrp1* promoter and further comprising one or more poly-adenylation sequences, transcription start sites, and/or transcription termination sequences. Such target gene or nucleotide sequence modifications and the methods by which they may be made are well known in the art. In certain embodiments, the target gene or nucleotide sequence encodes a nitrogen utilization protein., for example, a nitrate transporter, an ammonium transporter, an amino acid transporter, glutamine synthetase (GS), asparagine synthetase (AS), glutamate synthase (also known as glutamate 2:oxoglutarate amino transferase and GOGAT), asparaginase (ANS), glutamate dehydrogenase (GDH), aspartate aminotransferase (AspAT), or alanine aminotransferase (AlaAT). Other examples include those genes or nucleotide sequences which may be involved in Nutrient Use Efficiency (NUE) as described by Beatty et al. (2009), and in U.S. Patent No. 7,589,257.

[0085] In certain embodiments, the present invention provides for vectors comprising the genetic constructs. One skilled in the art will appreciate that the precise vector used is not critical to the instant invention and suitable vectors can be

readily selected by the skilled person. Examples of expression vectors and cloning vehicles include, but are not limited to, viral particles, baculovirus, phage, plasmids, phagemids, cosmids, fosmids, bacterial artificial chromosomes, retrovirus vectors, viral DNA (for example, vaccinia, adenovirus, fowl pox virus, pseudorabies and derivatives of SV40), P1-based artificial chromosomes, yeast plasmids, yeast artificial chromosomes, and other known vectors specific for specific host cells of interest.

[0086] Certain embodiments provide for host cells comprising the genetic constructs or vectors according to the invention. The genetic construct or vector can be introduced into a suitable host cell by one of a variety of standard methods. Such methods are well-known in the art and include, for example, stable or transient transfection, lipofection, electroporation, and infection with recombinant viral vectors. One skilled in the art will understand that selection of the appropriate host cell will be dependent upon the vector chosen. Examples of host cells include, but are not limited to, bacterial, yeast, insect, plant and mammalian cells. In certain embodiments, the host cell is a plant cell.

[0087] The methods and genetic constructs disclosed herein may be used to produce a plant or a plant part of any species capable of utilizing the promoter such that the transgenic (non-natural) plant, has tissue-specific expression of one or more desired genes or nucleotide sequences. The invention is intended to be particularly applicable to, for example, crop plants (especially those of the genus *Oryza*), ornamental plants, and trees (particularly conifers and the genus *Populus*). Particularly suitable plants for the practice of the present invention include, but are not limited to, canola, barley, sugar cane, corn, canola, tobacco, soybean, cotton, alfalfa, tomato, wheat, potato, aspen, cottonwood, conifers and poplar, or parts of any of these plants, for example, roots, root tips, leaves, stems, flowers, apical buds, meristematic tissues, and the like. In certain embodiments, plants for use in the invention include canola, barley, maize, rice, tobacco, soybean, cotton, alfalfa, tomato, wheat, potato, and certain tree genera, including conifers and *Populus* species.

[0088] In certain embodiments, the methods, uses and constructs of the invention are applied to plants of the genus *Oryza* and other genera closely related to *Oryza*, such as *Porteresia* and *Leersia*. Examples of species in these genera include, but are

not limited to, *Oryza sativa*, *Oryza punctata*, *Oryza officinalis*, *Oryza malapuzhanensis*, *Oryza latifolia*, *Oryza australiensis*, *Oryza brachyantha*, *Oryza granulate*, *Oryza longiglumis*, *Oryza schlechteri*, *Porteresia coarctata*, *Leersia perrieri*, *Leersia hexandra* and *Leeria tisserantii*.

[0089] The transgenic (non-natural) plants, plant parts, and seeds produced according to the present invention may be further useful in breeding programs for the production of plant species having more than one desired trait. For example two transgenic plants of the invention each having expression of a desired transgene in differing plant tissues may be crossed to result in progeny transgenic plants having tissue-specific expression of both transgenes; or two transgenic plants of the invention each having expression of a different desired transgene in the same plant tissue may be crossed to result in progeny transgenic plants having tissue-specific expression of both transgenes. In this fashion it is possible to produce transgenic plants having a combination of desirable traits in selected tissue(s) of the plant.

[0090] Furthermore, it will be understood by one skilled in the art that different species of plants may be more or less amenable to genetic manipulation in general, and that, therefore, it may be advantageous to first transform a related species of the desired plant by the methods and with the constructs of the invention and to subsequently introduce the tissue-specific expression of the target gene or nucleotide sequence into the desired plant species by cross-breeding techniques. Such techniques and appropriately related plant species are well known to one skilled in the art.

[0091] Plant cells or protoplasts that have been transformed with the gene construct of the present invention can be regenerated into differentiated plants using standard nutrient media supplemented with shoot-inducing or root-inducing hormone, using methods known to those skilled in the art (see, for example, U.S. Patent No. 4,634,674 and references therein). Seeds may additionally be harvested from such transgenic plants using methods well known in the art and further used to re-grow the transgenic plants and hybrids of the invention.

Uses

[0092] The methods and constructs of the invention allow the production of plants and seeds having expression of one or more desired genes or nucleotide sequences, for example, in one or more selected tissues of the plant. Thus, in certain
5 embodiments, the methods and constructs of the invention permit the production of plants having one or more desired traits limited to selected plant tissues, thereby enabling the targeting of a trait to the tissue to which it is best suited, or avoiding the expression of a desirable gene or nucleotide sequence in a tissue where its effects are unwanted. There are a wide variety of specific applications of the invention,
10 including, but not limited to, the production of plants having increased yield, stress tolerance, having improved nutrient uptake and/or utilization, having improved nutrient content and/or yields of desired compounds, and having phytoremediative properties. Specific applications of the invention are further described below.

[0093] One application of the invention is in the production of plants better able to
15 thrive on nutrient-poor soils. It is well known in the art that certain plant species, particularly crop plants, deplete the soil of nutrients necessary to sustain growth, such as nitrogen, phosphate, and potassium. In order to replenish the lacking nutrients, it is necessary either to fertilize the soil (an expensive and environmentally damaging procedure) or to cultivate plants known to deposit the depleted nutrient into the soil
20 (*e.g.*, clover or soybean in the case of nitrogen depletion), which may be crops that are less profitable or nutritive and therefore less desirable to grow. Frequent fertilizing to maintain optimal nutrients in the soil is costly in terms of agriculture outputs such as labour and fuel therefore crop plants are usually in a state of either nutrient abundance or depletion. Another application of the invention is in the production of plants better
25 able to capture and utilize nitrogen when it is present in either an adequate or abundant supply in the soil before nitrogen loss due to leaching, volatilization or microbial degradation occurs.

[0094] In certain embodiments, the methods of the invention permit the targeted expression of genes or nucleotide sequences involved in nutrient uptake (*e.g.*,
30 transport molecules) to those tissues in which the uptake occurs (*e.g.*, the root or root hairs) to thereby improve the ability of the plant to absorb the nutrient from the

environment. In some embodiments, the invention may be used to produce plants which express heterologous nutrient utilization nucleotide sequences, or optimized (for example, optimized for plant expression) native nutrient utilization nucleotide sequences, that permit more efficient use of the nutrient, such that less of the nutrient is required for the normal growth and functioning of the plant. In certain embodiments, the expression of such sequences is targeted to selected tissues (*e.g.*, the root or leaves). In certain embodiments, the methods of the invention allow for expression of genes or nucleotide sequences involved in the use and uptake of nutrients not normally used by the plant in those plant tissues which are directly exposed to the different nutrient (*e.g.*, root and leaf). In this fashion, plants which are able to grow and thrive on different nutrient sources (*e.g.*, different nitrogen sources) may be produced. Particularly useful target genes or nucleotide sequences for the optimization of nitrogen efficiency of the plant include: nitrate transporters, ammonia transporters and amino acid transporters, glutamine synthetase (GS), asparagine synthetase (AS), glutamate synthase (also known as glutamate 2:oxoglutarate amino transferase and GOGAT), asparaginase (ANS), glutamate dehydrogenase (GDH), alanine dehydrogenase, aspartate aminotransferase (AspAT) and alanine aminotransferase (AlaAT) and those genes or nucleotide sequences which may be involved in Nutrient Use Efficiency described by Beatty et al. (2009), and U.S. Patent No. 7,589,257.

[0095] In certain embodiments, plants produced by the methods of the invention can more efficiently utilize fertilizer input by rapidly taking up the nitrogen in the fertilizer and storing it at the time of application, to thereby reduce the amounts of nitrogenous fertilizer which are lost to leaching, etc. This may permit a reduction in the amount of nitrogenous fertilizer required to be applied to a crop, to obtain crop yields comparable to those obtained using normal cultivation techniques and plants which have not been modified according to the present invention. Additional agronomic advantages can include faster growth and crop yield, where nitrogenous fertilizer input is maintained at levels used in common crop cultivation techniques.

Development of Cereal Crops

[0096] As described below, transformed *Oryza sativa* plants ectopically expressing a nucleotide sequence encoding an enzyme involved in nitrogen assimilation or metabolism, for example but not limited to alanine dehydrogenase, glutamine synthetase, asparagine synthetase, glutamate synthase, asparaginase, glutamate dehydrogenase, aspartate aminotransferase, alanine aminotransferase, and those nucleotide sequences which may be involved in Nutrient Use Efficiency described by Beatty et al. (2009), and U.S. Patent No. 7,589,257 were developed using a *PBpr1* promoter and grown under laboratory conditions to determine if the beneficial effects on plant growth and yield under controlled growth conditions could be obtained. For example, it was examined whether the amount of available nitrogen would have an effect on plant biomass and seed yield on plants expressing *PBpr1/AlaAT* grown under laboratory conditions when compared to similar plants that did not ectopically express the target nucleotide sequence.

[0097] Seed yield in control and transgenic plants grown under laboratory conditions was examined. Transgenic plants expressing a gene or nucleotide sequence encoding an enzyme involved in nitrogen assimilation or metabolism exhibited higher biomass and seed yields than control plants in the presence of an adequate supply of nitrogen. Collectively these results indicate that transgenic plants ectopically expressing an enzyme involved in nitrogen assimilation or metabolism are capable of optimising the utilization of available nitrogen under a range of environmental conditions thereby resulting in an increase in plant biomass, seed yield or a combination thereof.

[0098] Therefore, in one aspect, the present invention provides a method for increasing seed yield of a plant comprising; transforming the plant with a target gene or nucleotide sequence in operative linkage with a *PBpr1* promoter element, to produce a transformed plant, the target gene or nucleotide sequence encoding an enzyme involved in nitrogen assimilation or metabolism; and growing the transformed plant.

[0099] The above description is not intended to limit the claimed invention in any manner, furthermore, the discussed combination of features might not be absolutely necessary for the inventive solution.

[00100] To gain a better understanding of the invention described herein, the following examples are set forth. It will be understood that these examples are intended to describe illustrative embodiments of the invention and are not intended to limit the scope of the invention in any way.

EXAMPLES

EXAMPLE 1: PBpr1 Promoter Fused to the *alaAT* Gene

Construction of binary vectors and Agrobacterium mediated transformation

[00101] The *PBpr1* promoter was selected and designed to be cloned using GeneArt (Invitrogen, Life Technologies, Carlsbad, CA, USA). The *HvAlaAT* cDNA (SEQ ID NO: 2) was introduced into pCAMBIA1300 using a PstI/HindIII ligation while the *PBpr1* promoter was introduced upstream of *HvAlaAT* to drive expression using EcoRI/SacI sites (Figures 1,13,14). The *PBpr1* promoter was introduced into pCAMBIA1305.1 to drive *GUSplus* for promoter pattern analysis using the EcoRI/NcoI sites. The constructs were transformed separately into *Agrobacterium tumefaciens* strain EHA105 by the freeze thaw method (Weigel and Glazebrook, 2002). Rice callus (*Oryza sativa* c.v. Nipponbare (NB)) was transformed with both constructs, using an *Agrobacterium* transformation system developed in our laboratory (Shrawat and Good, 2011). Lines transformed with *PBpr1::HvAlaAT* are referred to herein as *PBpr1* lines while lines transformed with *PBpr1::GUSplus* are referred to herein as *PBpr1G* lines.

[00102] Lines transformed with *PBpr1::HvAlaAT* were previously named *Ap* lines. Correspondence between the previous nomenclature and the nomenclature used herein is shown in Table 1. Likewise, lines transformed with *PBpr1::GUSplus* were previously named *ApG* lines.

[00103] Based on seed yield, total above soil biomass and tiller number at maturation, two T₀ lines containing *PBpr1::HvAlaAT* were selected to be continued through future generations. Leaf tissue was collected from two week old plants to determine if they were transgenic by means of a genomic DNA polymerase chain reaction (PCR) analysis using primers specific to the hygromycin resistant gene.

Transgenic plants and null segregants to be continued to further generations were selected based on uniform size and height. Primary transgenics are designated as T₀ lines.

Soil growth chamber experiments

[00104] T₀, T₁, and T₂ rice plant generations were planted in soilless-potting mixture Sunshine Mix #4 (Sun Gro Horticulture) in growth chambers at 28°C, 70% relative humidity, 14h/10h light/dark photoperiod, photon flux density 750 µE at bench height, as described in Shrawat et al., 2008).

[00105] In the T₃ generation, three seeds were germinated in a 7 inch round pot and allowed to grow for two weeks. Two of the three seedlings were culled based on uniformed height and size of plants remaining in each pot. A fertilization regime similar to that of the T₂ generation was started on the three week old plants. All statistical comparisons carried out for these growth chamber experiments were done using a Student's t-test.

Hydroponic growth chamber experiment

[00106] The hydroponic experiment was carried out in a growth chamber at 28°C, 70% humidity, 14h/10h light/dark photoperiod, photon flux density 750 µE at bench height, as described by Shrawat et al., 2008. Plant material was harvested on 28 and 52 day old plants for AlaAT assays and RNA experiments.

Selection of homozygous lines and maintenance of null lines

[00107] T₂ seed obtained from T₁ plants were germinated on MS and hygromycin and the ratio of resistance to the antibiotic was measured to determine homozygosity to heterozygosity as described in Shrawat et al., 2008. Null lines were identified by PCR at the T₁ plant stage using hygromycin specific primers. Subsequently, null lines were selected and maintained for a comparison between NB, AGR1/7 and *PBpr1* lines.

AlaAT Extraction and enzyme assays

[00108] Both shoot and root tissue was harvested for AlaAT enzyme assays in the hydroponic experiments, while only shoot tissue was harvested in potted experiments as per Muench and Good, 1994.

Protein SDS-PAGE electrophoresis and immunoblot analysis

5 [00109] In order to validate the results of the AlaAT enzyme assays, SDS-PAGE electrophoresis and immunoblot analyses was carried out using the protocol of Muench and Good (1994).

RNA extraction, cDNA synthesis and quantitative real time PCR

10 [00110] RNA extraction, cDNA synthesis and quantitative real time PCR was carried out using the protocol of Beatty et. al., 2009. Gene specific primer and probe pairs that were previously used by Beatty *et al.* (2009) were used to detect differences in gene expression due to over expression of *HvAlaAT*. NB plants were used as a negative control and 18srRNA was used as an endogenous control.

Bioinformatics analysis of promoter

15 [00111] The *PBpr1* nucleotide sequence was selected based on homology to the *OsANT1* promoter using NCBI's BLASTn program, and it was further investigated using the gene bioinformatic tools of NCBI Genbank and Refseq.

GUS histochemical staining

20 [00112] A total of 20 T₁ seeds of *PBpr1::GUSplus* transgenic lines, named PBpr1G lines, were sterilized and germinated in sterile liquid MS (4.4 g L⁻¹, pH 5.8) for staining. From each line, five seeds were selected at each sampling time for staining. Seedlings were sampled at 3 DAG and 7 to 10 DAG and stained for GUS expression.

RESULTS:

Heterozygous T₀ seed yield and biomass preliminary screens

25 [00113] At the T₀ stage, the independent PBpr1 plants had higher above ground biomass and seed yield compared to NB (Table 1). Two independent *PBpr1* lines

were selected based on high seed yield and biomass to continue on to the T₁ generation. When comparing the six lines selected with NB, they had significantly higher seed yield ($p<0.001$) and above ground biomass ($p<0.001$) compared to NB.

Table 1: Plant height, biomass and seed yield of T₀ PBpr1 plants and untransformed Nipponbare plants. SD refers to standard deviations.

Lines	Previous name of line	Plant height (cm)	Total above ground biomass (g)	% Change from NB total above ground biomass	Seed yield (g)	% change from NB seed yield	Total above ground biomass without seeds (g)
<i>PBpr1</i> -3	Ap10	86.5	11.6	6	3.13	-20	8.47
<i>PBpr1</i> -5	Ap12	69.2	12.5	15	3.34	-15	9.16
<i>PBpr1</i> -7	Ap14	80.8	13.2	21	3.37	-14	9.83
<i>PBpr1</i> -8	Ap15	79.2	14.2	30	4.27	9	9.93
<i>PBpr1</i> -9	Ap16	74.0	12.4	14	4.04	3	8.36
<i>PBpr1</i> -10	Ap17	73.8	17.9	64	7.33	87	10.57
<i>PBpr1</i> -11	Ap18	84.3	17.7	62	10.01	156	7.69
<i>PBpr1</i> -12	Ap19	87.2	20.3	86	8.31	113	11.99
<i>PBpr1</i> -13	Ap20	85.0	17.8	63	7.01	79	10.79
<i>PBpr1</i> -14	Ap21	78.5	12.2	12	4.62	18	7.58
<i>PBpr1</i> -15	Ap22	78.0	12.1	11	3.49	-11	8.61
<i>PBpr1</i> -16	Ap23	79.1	13.8	27	4.00	2	9.80
<i>PBpr1</i> -17	Ap24	86.5	17.2	58	6.54	67	10.66
<i>PBpr1</i> -18	Ap25	67.3	13.5	24	4.25	9	9.25
<i>PBpr1</i> -19	Ap26	80.2	15.9	46	5.51	41	10.39
<i>PBpr1</i> -20	Ap27	85.4	11.7	7	4.16	6	7.54
<i>PBpr1</i> -21	Ap28	86.0	17.4	60	7.33	87	10.07
<i>PBpr1</i> -22	Ap29	76.0	21.4	96	7.50	92	13.9
average		78.0	14.1	39	5.45	40	9.70
SD		6.8	3.63		2.05		1.62
Nipponbare		75.2	10.9		3.91		7.03
SD		6.3	2.17		1.63		1.25

Heterozygous T₁ PBpr1 seed yield and biomass preliminary screens

[00114] For further phenotypic analysis and to obtain homozygous seed, T₁ plants of *PBpr1* -11 and *PBpr1*-21 were grown to maturity and biomass and seed yield data collected. *PBpr1* -11 and *PBpr1*-21 had significantly higher above ground biomass

than NB ($p < 0.05$) and AGR1/7 ($p < 0.001$) (Figure 2). In terms of seed production, *PBpr1-11* and *PBpr1-21* had significantly higher seed yield than NB ($p < 0.05$).

Growth chamber biomass and seed yield analysis of homozygous lines

[00115] T_2 homozygous *PBpr1* lines were grown in soil conditions to maturity and total above ground biomass and seed yield data was collected (Figure 4). Two of the five *PBpr1* lines had higher above ground biomass than NB. Only one of these lines, *PBpr1-21*, was significantly larger than NB and AGR1/7 in terms of biomass ($p < 0.001$). *PBpr1-10* and *PBpr1-11* were shown to have higher seed yield than NB and AGR 1/7, but these differences were not significant ($p > 0.05$).

*T_3 above ground biomass of homozygous *PBpr1* lines*

[00116] Homozygous *PBpr1* T_3 plants were harvested at 52 DAG in order to assess the changes of biomass at a different developmental stage (Figure 5). *PBpr1-11* and *PBpr1-21*, had higher dry above ground biomass than NB and the nulls at 52 DAG. *PBpr1-21* had significantly higher above ground biomass than both NB and nulls ($p < 0.03$). *PBpr1-11-2N* plants had similar above ground biomass compared to NB, which was consistent with previous experiments in the T_1 and T_2 generations. *PBpr1-11*, on the other hand, had higher biomass than NB and nulls.

**PBpr1* biomass, seed yield and NUE*

[00117] *PBpr1-21*, over all 3 generations, has significantly higher above ground biomass than NB. In the T_1 and T_2 generation, *PBpr1-21* is higher in above ground biomass than AGR1/7 but not significantly (Figure 2 and 4). *PBpr1-11* also exhibits a consistent trend of higher above ground biomass than NB in the T_1 and T_3 generation but not in the T_2 generation (Figures 2, 4 and 5). Across all generations, *PBpr1-21* had higher seed yield than NB and AGR1/7 while *PBpr1-11* had higher seed yield than NB in the T_1 but not in the T_2 generation.

*T_1 *PBpr1* lines tiller productivity*

[00118] In addition to measuring biomass and seed yield at maturity, the number of productive and non-productive tillers was measured. Tiller productivity and plant

height was only recorded in soil grown plants. Tiller productivity of the PBpr1 lines, in comparison with AGR1/7 and NB was determined and all PBpr1 lines had more tillers and more productive tillers than NB (Figure 3). PBpr1-11 and PBpr1-21 had significantly ($p<0.01$) more tillers and more productive tillers than AGR1/7. PBpr1-11 had the highest total number of tillers and productive tillers among all PBpr1 lines, AGR1/7 and NB. This was consistent with other parameters assessing overall plant size as PBpr1-11 produced the largest plants among all other PBpr1 lines. In comparison to AGR1/7, PBpr1-11 and PBpr1-21 had significantly more productive tillers ($p<0.05$).

T₁ screens for AlaAT activity for HvAlaAT over-expression in soil grown plants

[00119] Shoots of T₁ soil grown PBpr1 plants were harvested between active tillering and maximum tillering stage (40 DAG) to determine their specific AlaAT activity. The different PBpr1 lines had 2.5 to 8 times higher shoot AlaAT activity compared to the null segregants and NB (Figure 6). AlaAT specific activity was highest in PBpr1-12 and PBpr1-21 at between 3.3 to 4.3 $\mu\text{moles NADH min}^{-1} \text{mg of protein}$. AGR1/7 containing *OsANT1::HvAlaAT* had higher specific AlaAT activity than nulls and NB in agreement with Shrawat *et al.* (2008). Out of all the PBpr1 lines, only PBpr1-12 had higher average specific AlaAT activity than AGR1/7, but PBpr1-11 and PBpr1-21 had comparable levels of AlaAT activity as AGR1/7. The AGR1/7 used was a T₅ homozygous line while the T₁ PBpr1 lines tested were still segregating. Null segregants of the different PBpr1 lines (PBpr1-10-N, PBpr1-11-N, PBpr1-12-N and PBpr1-21-N) had similar AlaAT activity as NB. Once these lines were confirmed as true nulls, they were used as negative controls.

T₃ homozygous shoot AlaAT activity assay from soil grown plants

[00120] Independent plants of the T₃ generation from the lines PBpr1-10, PBpr1-11, PBpr1-12 and PBpr1-21 were identified as homozygous plants and were raised in the growth chamber until 52 DAG when tissue samples were harvested for AlaAT activity. Trends observed in the T₃ generation were consistent with those of the T₁ plants (Figure 7). PBpr1 lines had up to 5.3 times higher shoot AlaAT activity than NB and the nulls. In agreement with T₁ studies in soil, PBpr1-12 had the highest

AlaAT activity followed by PBpr1-11 and PBpr1-21 while PBpr1-10 had the lowest AlaAT activity.

T₂ homozygous AlaAT activity from hydroponic grown plants

[00121] To assess the AlaAT activity of roots, homozygous PBpr1 lines were selected to be grown hydroponically. PBpr1 roots and shoot tissue were harvested at the beginning of active tillering (28 DAG) and at maximum tillering (52 DAG).

Shoot AlaAT activity

[00122] The PBpr1 lines exhibit up to 8 times higher shoot AlaAT activity compared to nulls and NB (Figure 8). At both 28 DAG and 52 DAG, PBpr1-11 and PBpr1-21 had up to 1.2 and 1.7 times higher AlaAT activity than AGR1/7 respectively. Shoot AlaAT activity of PBpr1-11 and PBpr1-21 was higher at 52 DAG than 28 DAG, whereas native shoot AlaAT activity (the PBpr1 null lines and NB) was observed to drop slightly between 28 and 52 DAG.

[00123] Between 28 DAG and 52 DAG, AlaAT activities in some of the transgenic, nulls and NB shoots were observed to have decreased with the exception of PBpr1-21. For all PBpr1 lines, the changes in shoot AlaAT activity was only between 1.1 to 1.2 times between 28 and 52 DAG.

Root AlaAT activity

[00124] Although all PBpr1 lines had higher root AlaAT activity than NB, similar levels of root AlaAT activity were observed between PBpr1 lines and their corresponding null siblings (Figure 9). AGR1/7 was the only transgenic line that had clearly higher level of root AlaAT activity than all other lines. However, root AlaAT activity was observed to have dropped in all cases at 52 DAG compared to 28 DAG. The decrease was evident in null segregants and NB where levels of root AlaAT activity were observed to have dropped 2 to 3 times. A smaller decrease of 1.2 to 1.4 times in root AlaAT activity of 28 to 52 DAG was observed in the PBpr1 lines and AGR1/7. With the decrease of root AlaAT activity at 52 DAG, PBpr1 lines and AGR1/7 had up to 3 times higher AlaAT activity than NB and nulls. Shoot AlaAT

activities in the non-transgenics are always low at both 28 and 52 DAG (Figure 8). Root AlaAT activity, on the other hand, is high at 28 DAG and decreases drastically at 52 DAG. Overall root AlaAT activity in the non-transgenics was always higher than shoot AlaAT activity.

HvAlaAT immunoblotting to validate AlaAT assays

[00125] Fifty two day old soil grown T₃ PBpr1 lines were harvested to determine the stability of the *PBpr1* promoter in over-expressing *HvAlaAT*, by AlaAT activity assays and immunoblots. T₃ shoots of all the PBpr1 lines tested had high levels of AlaAT protein according to the immunoblot (Figure 10'). In addition, *HvAlaAT* protein levels in AGR1/7 were found to be comparable but slightly higher than that of the PBpr1 lines. This was consistent with the results obtained in the AlaAT assays (Figure 7). The 52kDa *HvAlaAT* band is absent in the nulls and NB indicating the absence of *HvAlaAT* protein.

[00126] Roots of hydroponically grown T₂ plants were harvested at 28 days (Figure 11). In lines that did not carry the *HvAlaAT* transgene insertion, the *HvAlaAT* antibody bound to native AlaAT. PBpr1 and AGR1/7 roots produced a thicker and more intense band that masked the two nonspecific thinner bands representing native AlaAT at the 52kDa region (Figure 11A). Therefore roots of 28 day old PBpr1 plants had higher levels of *HvAlaAT* protein than the nulls and NB.

[00127] Shoot *HvAlaAT* protein levels of hydroponically grown T₂ PBpr1 plants were also assessed at 28 (Figure 11B). *HvAlaAT* protein was observed to be high in the shoots of all PBpr1 lines. Again, AGR1/7 had the same level of banding intensity as all the PBpr1 lines indicating high levels of *HvAlaAT* protein compared to nulls and NB. At 28 days, protein levels of *HvAlaAT* appeared to be present and high in PBpr1 lines and absent in the non-transgenics.

qRT-PCR analysis to determine gene expression changes from over-expression of HvAlaAT

[00128] The mRNA expression levels of three target genes or nucleotide sequences (barley AlaAT (*HvAlaAT*), leucine rich repeat (*LRR*) and a glycine rich protein (GRP)

were selected from a list of genes or nucleotide sequences that had been found to be differentially expressed in a microarray experiment comparing AGR1/7 to NB (Beatty *et al.*, 2009).

[00129] *HvAlaAT* was approximately 400 to 1400 fold higher in mRNA expression in the T₃, soil grown, PBpr1 lines and AGR1/7 shoots compared to NB, except for the nulls which did not differentially express *HvAlaAT* compared to NB (Figure 12A). The qRT-PCR analysis, AlaAT assays and immunoblotting confirmed that the PBpr1 lines were over-expressing *HvAlaAT* compared to the null segregant. Among the PBpr1 lines tested, PBpr1-21 had the highest *HvAlaAT* expression at 1400 fold higher followed by PBpr1-12 with a 1000 fold higher expression relative to NB (Figure 2.18). *GRP* was up-regulated approximately 10 to 30 fold in all PBpr1 lines, null lines and AGR1/7. *LRR* was 9 to 20 fold up-regulated in the PBpr1 lines and AGR1/7 compared to NB and the nulls, which were 3.4 fold up-regulated compared to NB.

[00130] Hydronically grown T₂ PBpr1 shoots were analyzed using qRT-PCR to determine mRNA expression levels compared to NB. In agreement with previous AlaAT assays and immunoblots, *HvAlaAT* mRNA was highly expressed in all PBpr1 lines with 400 to 800 fold higher expression relative to NB with PBpr1-21 shoots having the highest up-regulation, while the null line did not differentially express *HvAlaAT* relative to NB (Figure 12B). All the PBpr1 lines, null line and AGR1/7 exhibited 20 to 40 fold up-regulation of *GRP* relative to NB. *LRR* was most highly up-regulated in AGR1/7 at a 29 fold change relative to NB, while the transgenic PBpr1 lines had a 7 to 11 fold up-regulation and the null line had 16 fold up-regulation.

[00131] Hydroponically grown T₂ PBpr1 roots RNA was analyzed for changes in transcript levels in comparison to AGR1/7 and NB. *HvAlaAT* mRNA was highly expressed in PBpr1 lines at an approximately 700 to 2000 fold increase compared to NB. AGR1/7 roots had 2500 fold higher expression of *HvAlaAT* relative to NB (Figure 12C). As expected, the null line did not show differential expression of *HvAlaAT* relative to NB. *GRP* was highly up-regulated at 200 to 500 fold in the PBpr1 lines, null line and AGR1/7 relative to NB.

[00132] *GRP* was more highly upregulated in roots than in shoots relative to NB in both the *PBpr1* lines and *AGR1/7*. Similarly *HvAlaAT* mRNA was also more highly expressed in roots than in shoots of all the transgenic lines relative to NB (Figure 12B and C).

5 *GUS histochemical staining of PBpr1G plants*

[00133] Three and 7 to 10 day old *PBpr1G* plants were stained for GUS expression to determine the expression pattern of the *PBpr1* promoter at the seedling stage (Figure 15). GUS stains appeared at the tip of the newly emerged shoots in 3 day old plants. In 7 to 10 day old *PBpr1G* plants, staining was observed in the interface
10 between the shoots and the roots, within the remnant seed coat. Faint GUS staining was also found in the veins of the leaf blades and tillers as well as exterior of the tiller of *PBpr1G* plants, which may be endogenous GUS stain. With the *OsANT1::GUSplus* lines used as a positive control, GUS staining appeared within one hour of incubation in GUS staining buffer and GUS stains appeared in the root hairs, root tips and veins
15 of the leaves.

PBpr1 over-expresses *HvAlaAT*.

[00134] This conclusion is supported by the following observations: high levels of AlaAT activity were observed in all the *PBpr1* lines consistently throughout all generations analyzed; and immunodetection showed that there is a high level of
20 *HvAlaAT* protein in transgenic plants and finally, qPCR analysis demonstrated an increase in *HvAlaAT* mRNA in the *PBpr1* lines.

Expression from PBpr1 promoter

[00135] At the transcript level, the *OsANT1* and *PBpr1* promoters both drive high level of *HvAlaAT* gene expression in roots and shoots compared to Nipponbare (NB)
25 but *OsANT1* drives higher levels of *HvAlaAT* gene expression in the roots than *PBpr1* while *PBpr1* drives higher levels of *HvAlaAT* gene expression in shoots than *OsANT1* (Figure 12).

[00136] Pattern analysis of *PBpr1::GUSplus* lines (PBpr1G lines) showed endogenous GUS staining in 3 day old and 7 to 10 day old seedlings while intense GUS staining was observed in the *OsANT1::GUSplus* line. PBpr1 may not show GUS expression but yet it drives high levels of *HvAlaAT* expression. This agrees with the fact that PBpr1 may be developmentally regulated in roots until after active tillering while *OsANT1* drives high levels of expression at all times in both roots and shoots.

Transgenic HvAlaAT

[00137] At the protein level, *PBpr1::HvAlaAT* lines (PBpr1 lines) and *OsANT1::HvAlaAT* lines (AGR1/7) demonstrated higher levels of *HvAlaAT* protein in both roots and shoots from active to maximum tillering compared to NB (Figures 10 and 11).

[00138] At the enzymatic level, *AlaAT* activity of AGR1/7 was higher than NB at both active and maximum tillering in both roots and shoots. The PBpr1 lines exhibit similarly high levels of shoot *AlaAT* activity from active to maximum tillering compared to NB while root *AlaAT* activity of PBpr1 lines in roots was only observed to be higher than NB after active tillering and hence its activity seemed to be developmentally regulated in roots instead of constitutively regulated as seen in roots of AGR 1/7. In shoots, *AlaAT* activity appears to be high at all times using the *PBpr1* promoter.

HvAlaAT over-expressing plant lines

[00139] Morphologically, the *PBpr1* lines produced higher seed yield and biomass compared to NB and AGR1/7 in the T₁, T₂ and T₃ generations. In addition, the *PBpr1* lines exhibit increased tillering compared to NB and AGR1/7.

[00140] With the *OsANT1* promoter, an increase in gene expression resulted in higher levels of *HvAlaAT* protein and consequently higher levels of *AlaAT* activity in both the roots and shoots of AGR1/7. In contrast, although the PBpr1 promoter drove high levels of *HvAlaAT* expression and *HvAlaAT* protein levels, this did not result in an increase in *AlaAT* activity until after active tillering in roots. In shoots, an increase

in *HvAlaAT* gene expression driven by PBpr1 promoter resulted in both increased protein levels and AlaAT activity. This suggests that PBpr1 may be developmentally regulated in rice roots.

[00141] The nitrogen use efficiency (NUE) of three independent PBpr1 lines, AGR 1/7 and NB were calculated from the seed yield per gram of nitrogen applied per plant (Table 2). The three *PBpr1* lines were all higher in NUE than AGR 1/7 by 5-12 % and NB by 11-18%. The highest NUE line was *PBpr1*-21 and the lowest was NB.

[00142] Table 3 shows yield tests and agronomic traits from three *PBpr1:AlaAT* over-expressing rice lines (PBpr1-11, PBpr1-12, PBpr1-21 (*OsPBpr1::HvAlaAT*)) versus WT (NB; Nipponbare) at maturity during soil growth under high N conditions.

[00143] Figure 16 shows the foliar amino acid concentrations of wildtype (cv. Nipponbare) rice versus barley *AlaAT* over-expressing rice transgenic rice lines; AGR 1/7 (*OsANT1:HvAlaAT*) and PBpr1-11, PBpr1-12, PBpr1-21 (*OsPBpr1::HvAlaAT*). Measurements were taken at 28, 40, 52 and 95 days after germination (DAG) corresponding to active tillering, mid-tillering, maximum tillering and flag leaf emergence, respectively. Amino acid concentration (nM/g FW) are averages of 4 replicates.

Table 2: NUE of T₂ homozygous PBpr1 lines, AGR1/7 and Nipponbare (NB) plants (based on seed yield per gram of nitrogen applied per plant and the above ground biomass without seed. Each data point is taken from at least 4 replicates. (NUE= Grain yield/ N applied))

	Above ground biomass without seed (g)	Seed yield (g)	NUE (yield .unit applied ⁻¹)
PBpr1-10	18.26	8.55	43.63
PBpr1-11	18.03	8.29	42.27
PBpr1-21	24.20	8.83	45.07
NB	19.75	7.48	38.16
AGR1/7	19.94	7.89	40.23

Table 3. Yield tests and agronomic traits from three PBpr1:AlaAT over-expressing rice lines versus WT.

Trait	NB	PBpr1-11	PBpr1-12	PBpr1-21
Average days to flowering	92-105	88-99	93-100	94-110
Producing tillers	6.5	6.6	6.4	7.2
Nonproducing tillers	0.25	0.6	0.2	0.4
Seed weight	10.3±0.5	9.9±0.5	8.5±0.2	9.7±0.5
Fresh weight biomass per plant (g)	24.9±1.4	26.9±2.0	25.5±1.9	25.7±2.7
Dry weight per plant (g)	14.6±0.9	17.4±1.5	17.5±1.3	16.7±2.0
% dry weight of NB		108%	102%	103%

EXAMPLE 2: PBpr2 Promoter Fused to the *alaAT* Gene

[00144] The choice of promoter to use to drive expression of a gene, such as the alanine aminotransferase gene, in crop plants is important for optimal expression of the gene at the right time (developmental stage) and the right place in the plant (tissue specificity). The promoter chosen to drive the expression will affect the phenotypic outcome for the plant. There are examples in the literature where the promoter chosen to drive a gene of interest affects the phenotypic outcome. For example, Myashita *et al.*, (2007) used the CaMV35S promoter to drive expression of AlaAT in Arabidopsis and observed no phenotype. Furtado *et al.*, (2008) studied the ability of various promoters to drive expression of green fluorescent protein (GFP) in transgenic rice and showed that three out of five putative endosperm specific promoters from rice, barley and wheat did not allow GFP expression in glumes/lemma/palea of the flower tissues, while the other two did allow GFP expression in those plant parts. The choice of promoter to use therefore requires both careful design and experiments to determine its efficacy to improve NUE.

[00145] Prior to the studies outlined in Example 1, a promoter named "PBpr2" was selected because its native gene is homologous to rice aldehyde dehydrogenases, such as the OsANT1 gene. The sequence of the PBpr2 promoter is provided in Figure 18 (SEQ ID NO: 4). Based on sequence similarity the PBpr2 gene is a putative aldehyde dehydrogenase, however, its function is unknown.

[00146] An alignment of the sequences of the PBpr1, PBpr2 and OsANT1 promoters (SEQ ID NOs: 1, 4 and 5, respectively) is shown in Figure 19. Although neither the Softberry nor the PlantpromoterDB software predicted a TATA box in PBpr1 at the same location as is present for OsANT1 or PBpr2, there is a conserved TATGAG sequence at that location. Also, PBpr1 includes a regulatory element about 30 nt downstream of the TATA boxes, and although the promoter prediction software does not predict a regulatory element at the corresponding position for the other two promoters, it is in a section of DNA with high conservation between the three sequences. The sequence identity between the PBpr1 and PBpr2 sequences, based on the alignment shown in Figure 19, is approximately 50% over the entire length (733 nucleotides) of the PBpr1 sequence.

[00147] The PBpr2 promoter was fused to the alaAT gene and a Nos terminator sequence and used to drive expression of AlaAT in rice following protocols similar to those outlined in Example 1. The results of these experiments are shown in Figure 17 and show that PBpr2 does not increase plant height, above ground biomass or seed yield compared to control wild-type (Nipponbare) rice, and so does not promote an NUE phenotype.

[00148] Collectively, the results from Examples 1 and 2 show that despite their similarities, including an approximately 50% sequence identity, the PBpr1 and PBpr2 promoters when fused to the AlaAT gene have different effects on phenotypic outcome.

[00149] The disclosures of all patents, patent applications, publications and database entries referenced in this specification are hereby specifically incorporated by reference in their entirety to the same extent as if each such individual patent, patent application, publication and database entry were specifically and individually indicated to be incorporated by reference.

[00150] Although the invention has been described with reference to certain specific embodiments, various modifications thereof will be apparent to those skilled in the art without departing from the spirit and scope of the invention. All such modifications as

2012307006 06 Jul 2017

would be apparent to one skilled in the art are intended to be included within the scope of the following claims.

Throughout this specification, unless the context requires otherwise, the word "comprise" or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or method step or group of elements or integers or method steps but not the exclusion of any element or integer or method step or group of elements or integers or method steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgement or admission or any form of suggestion that the prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

2012307006 06 Jul 2017

REFERENCES

- Beatty, P.H., Shrawat, A.K., Carroll, R.T., Zhu, T., and Good, A.G. (2009) *Plant Biotechnology Journal* **7**, 562–576
- Burton, R.A., *et al.* (2011). *Plant Biotechnology Journal* **9**, 117-135.
- Freeman, J., Sparks, C.A, West, J., Shewry, P.R., and Jones, H.D. (2011). *Plant biotechnology journal* **9** (7): 788-796.
- Furtado, A., Henry, R.J., and Takaiwa, F. (2008) *Plant Biotechnology Journal* **6**, 679–693.
- Gao, C., and Han, B. (2009). *Gene* **431**, 86-94.
- Good A.G. and Beatty P.H. (2011b) Biotechnological Approaches to Improving Nitrogen Use Efficiency in Plants: Alanine Aminotransferase as a Case Study. In, *The Molecular and Physiological Basis of Nutrient Use Efficiency in Crops*, First Edition. Edited by Malcolm J. Hawkesford, Peter Barraclough. John Wiley & Sons, Inc. Published 2011 by John Wiley & Sons, Inc.
- Good, A.G., Johnson, S.J., DePauw, M.D., Carroll, R.T., Savidov, N., Vidamir, J., Lu, Z., Taylor, G. and Stroehler, V. (2007) *Canadian Journal of Botany* **85**, 252-262.
- Kasuga, M., Liu, Q., Miura, S., Yamaguchi-Shinozaki, K., and Shinozaki, K. (1999). *Nature Biotechnology* **17**, 287–291
- Ma et al., 2009, *Plant Cell Rep.* 28(11):1759-65. Epub 2009 Oct 10.
- Muench, D.G., and Good, A.G. (1994) *Plant Molecular Biology* **24**, 417–427.
- Oguchi, K., Tanaka, N., Komatsu, S., and Akao, S. (2004) *Plant cell reports* **22**, 848-858.
- Park, T.H., and Jones, J.D.G. (2008) *Euphytica* **166**, 331-339.

2012307006 06 Jul 2017

- Pino et al., 2007 *Plant Biotechnol J.* 5(5):591-604. Epub 2007 Jun 8.
- Qu, L.Q., and Takaiwa, F. (2004) *Plant Biotechnology Journal* **2**, 113–125.
- Shelton, A.M., Zhao, J., and Roush, R.T. (2002). *Annual Reviews in Entomology* **47**, 845–881
- Shrawat, A.K., and Good, A.G. (2011). *Methods in molecular biology* **710**, 355-372.
- Shrawat, A.K., Carroll, R.T., DePauw, M., Taylor, G.J. and Good, A.G. (2008) *Plant Biotechnology Journal* **6**, 722–732.
- Sweetlove, L.J., Heazlewood, J.L., Herald, V., Holtzapffel, R., Day, D.A., Millar, A.H. (2002) *Plant Journal* **32**, 891–904
- Tanaka, N., Takahashi, H., Kitano, H., Matsuoka, M., Akao, S., Uchimiya, H., and Komatsu, S. (2005). *Journal of proteome research* **4** (5), 1575-1582.
- Weber, D. (2003). *Pesticide Outlook* **14**, 256-259.
- Weigel, D., and Glazebrook, J. (2002) *Arabidopsis: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Wu, C.Y., Adachi, T., Hatano, T., Washida, H., Suzuki, A., and Takaiwa, F. (1998) *Plant Cell Physiology* **39**, 885–889.
- Xue, G-P., Way, H.M., Richardson, T., Drenth, J., Joyce, P.A., and McIntyre, C.L. (2011). *Molecular Plant* 1-16.

2012307006 11 Jul 2017

CLAIMS:

1. A transgenic monocot plant or monocot plant part comprising a polynucleotide encoding a nitrogen utilization protein selected from glutamine synthetase, glutamate synthetase-1, glutamate dehydrogenase, an aminotransferase, alanine aminotransferase, aspartate aminotransferase, asparagine synthetase and asparaginase operably linked to a PBpr1 promoter, wherein the PBpr1 promoter comprises a nucleotide sequence substantially identical to the sequence as set forth in SEQ ID NO:1.
2. The transgenic monocot plant or monocot plant part according to Claim 1, wherein the plant is rice, barley, wheat or maize.
3. The transgenic monocot plant or monocot plant part according to Claim 1, wherein the plant is rice.
4. The transgenic monocot plant or monocot plant part according to any one of Claims 1 to 3, wherein the nitrogen utilization protein is an alanine aminotransferase.
5. The transgenic plant or plant part according to any one of Claims 1 to 3, wherein the nitrogen utilization protein is alanine aminotransferase comprising the amino acid sequence of SEQ ID NO:3.
6. A monocot seed obtained from the transgenic monocot plant according to any one of Claims 1 to 5 comprising said polynucleotide encoding said nitrogen utilization protein operably linked to said PBpr1 promoter as described in any one of Claims 1, 4 or 5.
7. A monocot plant cell transformed with a polynucleotide encoding a nitrogen utilization protein selected from glutamine synthetase, glutamate synthetase-1, glutamate dehydrogenase, an aminotransferase, alanine aminotransferase, aspartate aminotransferase, asparagine synthetase and asparaginase operably linked to a PBpr1 promoter, wherein the

2012307006 11 Jul 2017

PBpr1 promoter comprises a nucleotide sequence substantially identical to the sequence as set forth in SEQ ID NO:1.

8. The monocot plant cell according to Claim 7, wherein the plant cell is a rice plant cell, a barley plant cell, a wheat plant cell or a maize plant cell.

9. The monocot plant cell according to Claim 7, wherein the plant cell is a rice plant cell.

10. The monocot plant cell according to any one of Claims 7 to 9, wherein the nitrogen utilization protein is an alanine aminotransferase.

11. The monocot plant cell according to any one of Claims 7 to 9, wherein the nitrogen utilization protein is alanine aminotransferase comprising the amino acid sequence of SEQ ID NO:3.

12. A genetic construct comprising a polynucleotide encoding a nitrogen utilization protein selected from glutamine synthetase, glutamate synthetase-1, glutamate dehydrogenase, an aminotransferase, alanine aminotransferase, aspartate aminotransferase, asparagine synthetase and asparaginase operably linked to a PBpr1 promoter, wherein the PBpr1 promoter comprises a nucleotide sequence substantially identical to the sequence as set forth in SEQ ID NO:1.

13. The genetic construct according to Claim 12, wherein the nitrogen utilization protein is an alanine aminotransferase.

14. The genetic construct according to Claim 12, wherein the nitrogen utilization protein is alanine aminotransferase comprising the amino acid sequence of SEQ ID NO:3.

2012307006 11 Jul 2017

15. The genetic construct according to any one of Claims 12 to 14, wherein the construct further comprises a transcription termination sequence, a polyadenylation region, or both.

16. A vector comprising the genetic construct according to any one of Claims 12 to 14.

17. A method of generating a monocot plant having increased nitrogen use efficiency comprising:

transforming a monocot plant cell with the genetic construct according to any one of Claims 12 to 14 or the vector according to Claim 16; and

growing the transformed monocot plant cell to produce a monocot plant that expresses the nitrogen utilization protein from the PBpr1 promoter,

thereby generating a monocot plant having increased nitrogen use efficiency, wherein the increased nitrogen use efficiency is relative to the nitrogen use efficiency of a wild-type monocot plant grown under identical conditions.

18. A method of generating a monocot plant having increased biomass comprising:

transforming a monocot plant cell with the genetic construct according to any one of Claims 12 to 14 or the vector according to Claim 16; and

growing the transformed monocot plant cell to produce a monocot plant that expresses the nitrogen utilization protein from the PBpr1 promoter,

thereby generating a monocot plant having increased biomass, wherein the increased biomass is relative to the biomass of a wild-type monocot plant grown under identical conditions.

19. A method of generating a monocot plant having increased seed yield comprising:

transforming a monocot plant cell with the genetic construct according to any one of Claims 12 to 14 or the vector according to Claim 16; and

growing the transformed monocot plant cell to produce a monocot plant that expresses the nitrogen utilization protein from the PBpr1 promoter,

2012307006 11 Jul 2017

thereby generating a monocot plant having increased seed yield, wherein the increased seed yield is relative to the seed yield of a wild-type monocot plant grown under identical conditions.

20. A method for producing a monocot plant having increased nitrogen use efficiency, increased biomass, increased seed yield, or a combination thereof, comprising:

(a) providing a monocot plant, a monocot plant part or a monocot seed comprising the genetic construct according to any one of Claims 12 to 14 or the vector according to Claim 16; and

(b) growing a monocot plant from the monocot plant, monocot plant part or monocot seed,

thereby producing a monocot plant having increased nitrogen uptake, increased biomass, increased seed yield, or a combination thereof, wherein the increased nitrogen uptake, increased biomass and increased seed yield is relative to the nitrogen uptake, biomass and seed yield of a wild-type monocot plant grown under identical conditions.

21. The method of Claim 20, further comprising providing a fertilizer to the plant or the habitat of the plant in step (b).

22. The method of Claim 21, wherein the fertilizer is a nitrogen containing fertilizer.

23. The method according to any one of Claims 17 to 22, wherein the monocot plant is rice, barley, wheat or maize.

24. The method according to any one of Claims 17 to 22, wherein the monocot plant is rice.

25. The method of any one of Claims 17 to 22, wherein the increased nitrogen use efficiency, increased biomass or increased seed yield is observed in the mature monocot plant.

2012307006 11 Jul 2017

26. Use of a transgenic monocot plant according to any one of Claims 1 to 5 for producing progeny.

27. Use of the genetic construct according to any one of Claims 12 to 14 or the vector according to Claim 16 to increase nitrogen use efficiency, biomass or seed yield in a monocot plant.

28. A process for increasing nitrogen use efficiency, biomass, seed yield, or a combination thereof, in a monocot plant comprising:

providing a monocot seed comprising the genetic construct according to any one of Claims 12 to 14 or the vector according to Claim 16;

applying a herbicide, insecticide, fertilizer, or combination thereof, to the monocot seed; and

growing a monocot plant from the monocot seed, wherein the plant has increased nitrogen use efficiency, increased biomass, increased seed yield, or a combination thereof, and wherein the increased nitrogen use efficiency, increased biomass or increased seed yield is relative to the nitrogen use efficiency, biomass or seed yield of a wild-type monocot plant grown under identical conditions.










FIG. 1


**FIG. 2**


**FIG. 3**


**FIG. 4**

A.**B.****FIG. 5**

**FIG. 6**

**FIG. 7**

**FIG. 8**

**FIG. 9**





FIG. 10

**FIG. 11**




FIG. 12

EcoRI
 + GAATTCGTGAA AGTTCCGTC CAAATCGCAC CTTTTAACCG TTGAAAAAC ATACAAACGA AAAATAATCT ATATCTTAAT CAGGAAGAAA GAGTACGAAA
 - CTTAAGACTT TCAAAGGCAG GTTTAGCGTG GAAAAATTGGC AAACCTTTTG TAIGTTTGCT TTTTATTAGA TATAGAATTA GTCCTTCTTT CTCATGCTTT

101 TGGTGAACCG TCGAAACTAT TCATATACGT CGTCTGTCTC ATGAAAAAAA AAATCAATCC AGAAGGATAC GAGACACTTT TACTTCAACA AATATAGACA
 ACCACTTGGC AGCTTTGATA AGTATATGCA GCAGACAGAG TACTTTT TTTAGTTAGG TCTTCCTATG CTCTGTGAAA ATGAAGTTGT TTATATCTGT

201 TGAGCTTATT CTA TAGGTT TGGTTGTTTA ATAAGACGAA AGAATACAT TGGTTAGTTT TTCAATAAAA AATAATCGTT TGACTGACAT AAACCTAGGA
 ACTCGAATAA GATGATCCAA ACCAACAAAT TATTCTGCTT TCATTATGTA ACCAATCAAA AAGTAATTTT TTATTAGCAA ACTGACTGTA TTTGGATCCT

301 AATACTGGAT TAAGATAGAT CAGTAGGATT AAGATCCACT GATGTAATTT CCCACTGATT TGGTGGCTGA CATGTGGACC TGAGAGTTGT GTGGGCTCAC
 TTATGACCTA ATTCTATCTA GTCATCCTAA TTCTAGGTGA CTACATTAAA GGGTGACTAA ACCACCGACT GTACACCTGG ACTCTCAACA CACCCGAGTG

401 ATGTCAAATC ACGGTGAACA GTACGTCACG ATATGTTAGA GGTTCCTCTT CCGGAGATAC TTATACGAAT TTGCGGAAA CCTGCAAACT TTGATGGACG
 TACAGTTTAG TGCCACTTGT CATGCAGTGC TATACAATCT CCAAGGAGAA GGCTCTATG AATATGCTTA AAACGCCTTT GGACGTTTGA AACTACCTGC

501 ATTGAGCGA GTT TAGTTCT AAATTTTTC TTCAAACCTC TAACTTTTTC ATCACATCGT TTCAATTTCA ATCAAACCTC CAATGTTGAC GTGAACATAA
 TAACTCCGCT CAAATCAAGA TTAAAAAAG AAGTTTGAAG ATTGAAAAAG TAGTGTAGCA AAGTTAAAGT TAGTTTGAAG GTTACAACTG CACTTGATT

601 CACACCTATG AGATATGAGA ACGGGGTTGA CACTTGACAA GTCCTGACAT GCTGTGTTGG CGTGGGCCCC ACCTGCCACG TCAGGTCCAG CTCGGGTGG
 GTGTGGATAC TCTATACTCT TCGCCCAACT GTGAACTGTT CAGGACTGTA CGACACAACC GCACCCGGGG TGGACGGTGC AGTCCAGGTC GAGGCCACC

SacI
 701 TTGGGTTTGG TGCTTTCCGA TAGGCA**GAG CTC** ATGGCTG CCACC
 AACCCAAACC ACGAAAGGCT ATCCGT**GCTC GAG** TACCGAC GGTGG

FIG. 13


A.

atggctgccaccgtcgccgtggacaacctgaaccccaagggttttaaaatgtgag
tatgctgtgctggagagattgtcatccatgctcagcgcttgaggaacagcta
aagactcaaccaggggtctctaccttttgatgagatcctctattgtaacattggg
aaccacaaatctcttggtcagcaaccagttacattcttcagggagggttcttgcc
ctttgtgatcatccagacctgttgcaaagagaggaaatcaaacattgttcagt
gctgattctatctctcgagcaaagcagattcttgccatgatacctggaagagca
acaggagcatacagccatagccaggggtattaaaggacttcgtgatgcaattgct
tctgggatcgcttcacgagatggattccctgctaattgctgatgacattttctc
acagatggagcaagtccctggggtgcacctgatgatgcaattactgataaggaat
gagaaagatggcattcttggtcccagattcctcagtagcccttgtagctcggttcc
atagctcttcatggcggagctcttggtccatactatctcaatgaatcgacgggc
tgggggtttggaacctctgatgttaagaagcaacttgaagatgctcggtcaaga
ggcatcaacgttagggctttggtggttatcaatccaggaaatccaactggacag
gtacttgctgaagaaaaccaatatgacatagtgaagttctgcaaaaatgaggggt
cttggtcttctagctgatgaggtataccaagagaacatctatgttgacaacaag
aaattccactctttcaagaagatagttagatccttgggatacggcgaggaggat
ctccctctagtagtatcatatcaatctgtttctaaaggatattatgggtgagtggtg
aaaagagggtggttactttgagattactggcttcagtgtccagtaagagagcag
atctacaaaatagcatcagtgaacctatgctccaatatcactggccagatcctt
gctagtcttgctcatgaacccaccaaaggctagtgtgatgaatcatacgttccatac
aaggcagaaaaagatggaatcctcgcatcttttagctcgtcgtgcaaggcattg
gagcatgcattcaataaacttgagggaattacttgcaacgaggctgaaggagca
atgtacgtgttccctcaaactctgtctgccacagaaggcaattgaggctgctaaa
gctgctaacaaagcacctgatgcattctatgctcttcgtctcctcgagtcgact
ggaatcgctgttggtccctggatcaggatttgccaggttcttgccacatggcac
ttcaggtgcacgatccttccgcaggaggataagatcccggcagtcacatctccgc
ttcacgggtgttccatgaggcggttcatgtcagagtatcgtgactaaactggt

B.

MAATVAVDNLPKVLKCEYAVRGEIVIHAQRLQEQLKTQPGSLPFDEILYCNIG
NPQSLGQQPVTFREVLALCDHPDLLQREEIKTLFSADSI SRAKQILAMIPGRA
TGAYSHSQGIKGLRDAIASGIASRDGFANADDIFLTDGASPGVHLMQLLIRN
EKDGILVPI PQYPLYSASIALHGGALVPYYLNESTGWGLETS DVKKQLEDARSR
GINVRALVVINPGNPTGQVLA EENQYDIVKFCKNEGLVLLADEVYQENIYVDNK
KFHSFKKIVRSLGYGEEDLPLVSYQSVSKGYYGECGRGGYFEITGFSAPVREQ
IYKIASVNLCSNITGQILASLVMNPPKASDES YASYKAEKD GILASLARRAKAL
EHAFNKLEGITCNEAEGAMYVFPQICLPQKAIEA AKAANKAPDAFYALRLLEST
GIVVVP GSGFGQVPGTWH
FRCTILPQEDKIPAVISRFTVFHEAFMSEYRD



FIG. 14

**FIG. 15**

16/20

Amino acids	28 DAG					40 DAG				
	NB	AGR 1/7	PBpr1-11	PBpr1-12	PBpr1-21	NB	AGR 1/7	PBpr1-11	PBpr1-12	PBpr1-21
Glu	5313.8	4344.2	4321.1	4421.3	4426.4	4322.8	5299.0	5771.8	5042.7	5018.9
Ala	3613.0	3143.7	3478.5	4549.0	3578.7	3442.4	4460.8	3801.1	4232.9	3692.7
Asp	1578.7	1103.0	1258.7	1155.1	1034.8	1200.4	1264.9	1554.6	1168.4	1228.7
Ser	1568.5	930.4	1566.1	1006.9	590.8	982.8	583.2	1392.0	635.4	492.0
Thr	664.0	489.2	736.5	545.0	438.4	453.7	535.2	686.9	482.7	457.2
Asn	644.2	338.9	1129.6	620.8	389.2	79.5	80.1	157.3	56.3	58.4
Gln	597.5	1342.8	2805.3	1180.6	952.2	199.6	429.1	629.7	253.1	309.1
Gly	304.9	276.7	426.3	530.2	354.8	197.9	276.2	330.6	315.2	254.7
GABA	293.4	173.5	394.8	258.9	280.3	229.3	201.2	196.9	163.2	144.7
Val	204.0	246.5	414.6	279.6	239.7	198.5	213.4	246.6	231.0	225.8
His	167.6	189.3	202.2	164.8	133.1	181.4	156.2	173.0	182.2	140.6
Hyp	166.2	92.9	144.9	140.6	108.3	152.4	134.3	152.9	126.1	107.3
Pro	113.4	90.0	107.8	86.9	99.1	83.0	65.8	114.0	86.4	74.7
Lys	56.2	56.0	75.8	70.2	48.5	52.5	42.6	63.1	50.3	45.4
Leu	39.0	58.7	57.6	53.6	61.8	37.6	49.0	53.1	47.5	53.3
Tyr	32.3	50.4	86.7	34.4	49.2	nd	nd	109.9	nd	nd
Arg	nd	nd	294.2	81.1	nd	41.5	41.5	nd	44.1	52.3
Cys	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Met	nd	nd	24.8	nd	nd	nd	nd	nd	13.9	nd
Trp	nd	16.4	58.0	16.7	nd	nd	nd	24.3	nd	nd
Phe	nd	nd	53.2	17.2	14.3	nd	19.1	42.5	38.3	35.4
Ile	nd	27.1	124.9	38.9	35.5	42.4	nd	55.3	33.1	41.2
TOTAL	15356.8	12969.6	17761.6	15251.8	12835.0	11897.7	13851.7	15555.5	13202.8	12432.4
Amino acids	52 DAG					95 DAG				
	NB	AGR 1/7	PBpr1-11	PBpr1-12	PBpr1-21	NB	AGR 1/7	PBpr1-11	PBpr1-12	PBpr1-21
Glu	5210.8	5186.8	4880.6	5340.4	5340.4	4393.2	5500.2	4431.5	4512.8	3750.9
Ala	3080.7	3011.1	3536.2	2836.1	2836.1	2388.6	1832.1	2142.7	2721.5	1801.4
Asp	1392.8	1423.2	1255.4	1673.0	1673.0	1023.3	1667.9	1431.4	1266.8	1129.9
Ser	801.9	1199.7	1018.8	971.0	971.0	622.8	791.5	533.6	512.5	520.8
Thr	414.3	524.8	491.9	454.0	454.0	252.7	347.8	261.3	280.4	185.1
Asn	169.1	254.6	139.0	165.3	165.3	296.5	161.9	39.1	103.3	25.9
Gln	428.1	412.0	240.6	369.6	369.6	405.1	282.3	202.4	216.0	261.9
Gly	371.0	397.2	377.2	378.4	378.4	351.3	333.2	374.1	490.4	327.8
GABA	212.9	198.3	148.3	116.8	116.8	197.8	111.2	139.6	128.4	122.6
Val	289.0	247.6	272.1	224.4	224.4	396.4	367.8	253.2	226.0	226.7
His	333.7	307.3	304.7	291.4	291.4	430.3	451.4	377.8	386.2	425.3
Hyp	111.1	152.4	115.6	97.9	97.9	122.8	93.5	85.5	132.5	52.8
Pro	nd	45.9	57.2	45.8	45.8	237.1	128.7	114.7	52.6	54.4
Lys	nd	nd	nd	nd	nd	220.8	nd	nd	nd	nd
Leu	nd	nd	24.1	nd	nd	76.8	112.9	29.0	nd	nd
Tyr	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Arg	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Cys	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Met	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Trp	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Phe	nd	33.1	31.9	41.3	41.3	103.4	120.7	80.8	61.5	58.0
Ile	40.3	nd	32.1	31.7	31.7	85.7	137.2	27.1	nd	nd
TOTAL	12855.	13393.	12925.	13037.	13037.	11604.	12440.	10523.	11091.	8943.5
	7	9	8	2	2	4	2	9	0	

FIG. 16
SUBSTITUTE SHEET (RULE 26)

A.**B.****FIG. 17**

A.

GAATTCTGAAAGTTTCCGTCCAAATCGCACCTTTTAACCGTTTGAAAA
ACATACAAACGAAAAATAATCTATATCTTAATCAGGAAGAAAGAGTA
CGAAATGGTGAACCGTCGAAACTATTCATATACGTCGTCTGTCTCATG
AAAAAAAAAATCAATCCAGAAGGATACGAGACACTTTTACTTCAACA
AATATAGACATGAGCTTATTCTACTAGGTTTGGTTGTTTAATAAGACG
AAAGAAATACATTGGTTAGTTTTTCATTAAAAAATAATCGTTTGACTG
ACATAAACCTAGGAAATACTGGATTAAAGATAGATCAGTAGGATTAAG
ATCCACTGATGTAATTTCCCACTGATTTGGTGGCTGACATGTGGACCTG
AGAGTTGTGTGGGCTCACATGTCAAATCACGGTGAACAGTACGTCACG
ATATGTTAGAGGTTCCCTCTTCCGGAGATACTTATACGAATTTTGCGGA
AACCTGCAAACCTTTGATGGACGATTGAGGCGAGTTTAGTTCTAAATTT
TTTCTTCAAACCTTCTAACTTTTTTCATCACATCGTTTCAATTTCAATCAA
CTTCCAATGTTGACGTGAACTAAACACACCTATGAGATATGAGAAGCG
GGTTGACACTTGACAAGTCCTGACATGCTGTGTTGGCGTGGGCCCCAC
CTGCCACGTCAGGTCCAGCTCCGGGTGGTTGGGTTTGGTGCTTTCCGAT
AGGCACGAGCTC

B.

GAATTCATATAGTATCTTTGTATGTTGAGCATTAGTATTCAGTGAGTTA
AAACTTTTATAGTTGTTAGTGGATGAGTTTGAGTTTTAGGATTTAGTGGG
GTATTATAATTTTCATGGTAAGATAGGATGAGTATATATAATTAGGGAT
TAAATTACAATATTTTAAGGATCATATCGATATTCAAAACAATATGTTT
TTTTAAGATGGATGGAATCAATTTTCTGAGATATGTCCCGCTGGGGTG
TCCACAATCTGCACGCCATATCTGACTACACATATAACAACGCCCATCC
TCTGTCGATCTCCTTGTGTAAACACAACCGTCCGTGACACGTACGTAC
ATGGGGGATAGCTTGCCTCTCCGTTCAAGTTGCGTGACAACATAAAAA
TACTAACACAATGTCTTAATCCTCCAAAAAAAACCTCCTAAATAATTA
ATTAAAGATAAGGCAAGAAATTAATCCAAATAGTCCAAAACAATAGG
ATAAAGAGGAGAGCAAAACCTCCAAATGGTCACACGTCTCTCGGTGTC
ATTTTCTCGTCGGCTTGCTTTGTTTGATTCCAGAGTTTCCCAAACCTCTCT
AGAGAGCAAGTCAAACAGATCTCATCGATCGCCCAAGAAGAAGACAC
ATCCACAAATAATAATCTCATCTTATCGTACTAATTTATCTTAATACAG
TGTAATAAACTATATCAGTTTGAGATATTTTTTTCTGTCTAAATTATC
TCCAAACCGTAGGAATTTTCGCACCCGACGATCTCTTTCTTGATTTT
CAAACCAAACGGACCAATCTCTCTCGCGTGTATATATAGCGAGGTGGA
GCTCGAGCTCCATGCCTGCTTCCTCCTGACTCCTCTCCTGCATTCTCTC
GCTTCTGACGGCGGCAACTAACCTTTTTGGGATTTCGATTTCGAGGAGGA
GGTGATGGGATCC

FIG. 18

		1	50
O. sativa	(1)	AGGAAGTGAATTTTAGCGTAGCTGGTTGGTAGCGTAATTGCGTAAAGTTC	
PBPr1 promoter	(1)	-----	
PBPr2 promoter	(1)	-----GAATTCATATAGTATCTTTGTATGTTGAGCATTAGTATCAGTGT	
Consensus	(1)	ATT TA GTA CT TGT TGT G G A T G T AGT	
		51	100
O. sativa	(51)	CTTCAATTTTGGCTATATCTCAGTCGAA-AGATTTTCTTATCTCTCAC	
PBPr1 promoter	(1)	-----	
PBPr2 promoter	(45)	AGTTAAACCTTATAGTTGTTAGTGGATGAGTTTGGATTTTAGGATTTAG	
Consensus	(51)	TT AA TT TA T T A T GA AG TT T TTA T A	
		101	150
O. sativa	(100)	TCGATTTCTCACCAAATTTACAGTGTATTCTCTGTA---AGTTACAG	
PBPr1 promoter	(1)	-----	
PBPr2 promoter	(95)	TGGGTAATTATAAATTCATGGTAAGATAGGATGAGTATATATAATTAGGG	
Consensus	(101)	T G T T A T AT AG T T TA A TTA G	
		151	200
O. sativa	(147)	TGTAATTTATGAACTTACACCTGTAACCTTTGTAAGTTACAGTGTAATTT	
PBPr1 promoter	(1)	---GAATTCGAAAGTTCCGTC-CAATTCGCACCTTTTAACC---GTT	
PBPr2 promoter	(145)	ATTAATTTACAATATTTAAGGATCATATCGATATTCAAAACAATATGTT	
Consensus	(151)	TAAATTATGAAA TTTCAGT TCA ATCG TA TTTAAAC TA GTT	
		201	250
O. sativa	(197)	TTGAATCTTTCACATGTAAATTTTAAATTTTGTATTGGATTGTTCTTTT	
PBPr1 promoter	(43)	TGAAAA---ACATACAAACGAAAAAATAATCTAT-----ATCTTAAT	
PBPr2 promoter	(195)	TTTTTA---AGATGGATGGAAATCAATTTCTGA-----GATATGTC	
Consensus	(201)	TT AAA ACATG AAA ATAAATTTCTAT GTCTT TT	
		251	300
O. sativa	(247)	CTTGAGGATATGGTAATTTAATGTTCTATTAGGTTTCTTAAT-TGCTT	
PBPr1 promoter	(81)	CAGGAAGAAAGAGTACGAAATGGTGAACCGTCGAAACTATTCATATACGT	
PBPr2 promoter	(233)	CCGCTGGGTGTCCAAATCTGCACGCCATATCTGACTACACATATACAA	
Consensus	(251)	C GGAGGA AG GTAC ATATGGT AC T GTGACTATTTCATATAC T	
		301	350
O. sativa	(296)	TTGCTTTTATTATATCTATCGGAATTTAATACAAAGATTAAAAATCTG	
PBPr1 promoter	(131)	CGTCTGTCTCATGAAAAAATAATCAATC--CAGAAGGATACGAGACACT	
PBPr2 promoter	(283)	CGCCCATCCTCTGTCTGATCTCTTGTGTAACACAACCGTCGGTGACACG	
Consensus	(301)	CGTCC TCTTATGA AA A C T T T AACACAA GAT CGAGACACG	
		351	400
O. sativa	(346)	TGTAAT---ACGATTTATAAAAACTTTTCGAAAGATGTATAGGTACTCCCA	
PBPr1 promoter	(179)	TTTACTTCAACAAATATAGACATGAGCTTAT-TCTAC-TAGGTTTGGTTG	
PBPr2 promoter	(333)	T-----ACGTACATGGGGGATAGCTTGCCCTCTCCGTTCACTTGCCTG	
Consensus	(351)	T T T ACGAATATAGA AT AGCTTA TCT C TAGGTTTGC TG	
		401	450
O. sativa	(393)	AGCCCTTTTAAAGAAAGTTTTCAGACAAAAGTTTTTGGATGAAAGGTAG	
PBPr1 promoter	(227)	TTTAATAAGACGAAAGAAATACATGG-TTAGTTTTTCATTAATAAAAA---	
PBPr2 promoter	(375)	ACAACATAATAAATACTAACACAATGTCCTTAATCCTCCAAAAAATACTC	
Consensus	(401)	A ACTAATAAGAAAGTAATACAATG TTAGTTTTTCAATAAAAAA	
		451	500
O. sativa	(443)	TTATAGGGAAAAAGGAATGTGCGTTTATGTTTATTGGCATTGCTTATTAG	
PBPr1 promoter	(273)	-TAATCGTTTGACTGAC-ATAAACCTAGGAAATACTGGAT----TAAGAT	
PBPr2 promoter	(425)	CTAAATAAATAATTAAGATAAGGCAAGAAATTAATCCAAA---TAGTCC	
Consensus	(451)	TAAA G TTAA TGAA ATAAG CTAGGAATTA TGCAT TA TA	
		501	550
O. sativa	(493)	CAACCAAAAACTAATCTATAAGTAAATCTTTATATACGTGCGCTTAA-T	
PBPr1 promoter	(317)	AGATCAGTAGG--AT-TAAGATC----CACTGATGTAATTT-CCCAC-T	
PBPr2 promoter	(472)	AAAAACAATAGG--AT-AAAGAGGAGAGCAAAACTCCAAATGGTCAACAGT	
Consensus	(501)	AAA CAATAGG AT TAAGAG A A CA T AT TAA TG TC CAC T	
		551	600
O. sativa	(542)	AATTCAAAAGCAAAATTCATGTAAAAATAAATGCGATGAAG--AAACTTTA	

FIG. 19

PBPr1 promoter	(357)	GATTGGTGGC--TGACATGTGGACCTGAGACTTGTGTGGGCTCACATGT	
PBPr2 promoter	(519)	CTCTCGGIGTCATTTCTCGTCCGCTTGCTTTGTTTGAATTCAGAGTTTC	
Consensus	(551)	ATTCTGGTGGCA TTTCATGT GACTTGA TG T TGA G CA ACTTT	601 650
O. sativa	(590)	AAAAGTTATCAAATTTAGATTTTATTAAATTTAGTTTACAAGAGCGCTA	
PBPr1 promoter	(405)	CAAATCAGCG---TGAACAGTACGTACAG-ATATGTTAGAGGTCCTCTT	
PBPr2 promoter	(569)	CAAACCTCTC---TAGAGAGCAAGTCAAAAGATCTCATCGATCGCCCAA	
Consensus	(601)	CAAA CTCTC T AGAGTA GTCAAA ATATGTTA CGAT GC CTA	651 700
O. sativa	(640)	CGATGAAGGCTTTAAAGAGTGGGAAAAATA---AAACCTTTGACCTTTTC	
PBPr1 promoter	(451)	CC-GGAGATACTT-----A---TACGAATTTTGCGGAAACCTGCAAACTT--	
PBPr2 promoter	(616)	GAAGAAGACACATCCCAAATAATAATCTCATCTTATCGTAATAATTTAT	
Consensus	(651)	C AGGAGA ACTT A A ATA GAAT T AAACCT C AACTT	701 750
O. sativa	(686)	TGGACTTCAC-CAAACAGCTCAGCTTTGGCTTCGTGCCGCTCTGTCCTC	
PBPr1 promoter	(492)	--TGATGGAC-GATTGAG---GCGAGTTAGTTCTAAATTTTCTTTCAA	
PBPr2 promoter	(666)	CTTAATACAGTGTAAATA---ACTATATCAGTTTGAGATAATTTTCTCT	
Consensus	(701)	TAAT CAC GAAA AG ACGATTTCAGTTT A AT TTTTCTTCC	751 800
O. sativa	(735)	G--TGCTACTGCTA-CCCGCTCCCTGACCCCAACCCGC-CACTCCACGCTCCCT	
PBPr1 promoter	(536)	ACTTCTAACTTTTTCATCACATCGTTTCAATTT---CAATCAA--ACTTC	
PBPr2 promoter	(713)	G--TCTAAATAT-CTCCAAACCGTAGGAATTTTGCACCCGACGATCTC	
Consensus	(751)	G TCTAACTT T C CCA ACCGT CAATTT C CACTC ACGATCTC	801 850
O. sativa	(782)	TTCTCTCTCCCTTTCCGCTGACACACAGTCCCACTCCGACCGCTCCGTAT	
PBPr1 promoter	(581)	CAATGTTGACGT-----GAACTAAACACACCTAT-----GAGA-TAT	
PBPr2 promoter	(760)	TTTCTCTTGATTTTTC---AAACCAAACGGACCAATCTCTCTGCG-CTGTAT	
Consensus	(801)	TT TCTTGAC TT C GAACCAAAC CACCAATC C CG G GTAT	851 900
O. sativa	(832)	AAGTATCCCTTCCTTACCGCCGGCCAGCCACAGCCACCGCTCCCCCCACC	
PBPr1 promoter	(617)	GAGAAGCGGGT---TGACACTTGA---CAAGTCCTGA--CATGCTG-TGT	
PBPr2 promoter	(807)	ATATAGCGAGG---TGGAGCTCGAGCTCCATGCCCTGCTTCCCTCTTG-ACT	
Consensus	(851)	AAGTAGCG GT TG CGCT GA CCA CCTGC CCTCTG ACT	901 950
O. sativa	(882)	CCACCCCGATCCCCCTCCCGGCCGTACGGGCGCAGAAGGAACCCGCTCTTCT	
PBPr1 promoter	(658)	TGGCGTGGGCCCCACCTGCCACGTCAGGTC---CAGCTCGGGTGGTT-	
PBPr2 promoter	(853)	CCCTCTCTGCAATCTCTGCTTCTGACGCG--GCAACTAACCTTTTGTG-	
Consensus	(901)	CC C CCGGCCCCCTCTCCC CGT AGGGCG GCAGTACCCGT TT	951 1000
O. sativa	(932)	AGAAGGAGGAGGAGGCTA-CTCTCTCTCTCTCTCTTCTGCCATG	
PBPr1 promoter	(703)	----GGGTTTGGTGCTTTCGATAGGCACGAGCTC-----ATG	
PBPr2 promoter	(900)	----GGATTGATTTCAGGAGGAGGTGATGGGATCC-----ATG	
Consensus	(951)	GGATT GGTGCG T CG GTCACG GCTC ATG	

Potential TATA box
Regulatory element group
Pyrimidine (Y) patch

FIG. 19 (con.)

pctca2012050622-seq1.txt
SEQUENCE LISTING

<110> GOOD, Allen
LOCK, Yee Ying
BEATTY, Perrin H.

<120> PLANTS HAVING ENHANCED NITROGEN EFFICIENCY

<130> V84334W0

<150> 61/532,016

<151> 2011-09-07

<160> 6

<170> PatentIn version 3.5

<210> 1

<211> 733

<212> DNA

<213> artificial sequence

<220>

<223> PBpr1 promoter

<400> 1

gaattctgaa agtttccgtc caaatcgcac cttttaaccg tttgaaaaac atacaaacga	60
aaaataatct atatcttaat caggaagaaa gagtacgaaa tgggtgaaccg tcgaaactat	120
tcatatacgt cgtctgtctc atgaaaaaaaa aaatcaatcc agaaggatac gagacacttt	180
tacttcaaca aatatagaca tgagcttatt ctactagggtt tgggtgttta ataagacgaa	240
agaaatacat tgggttagttt ttcattaaaa aataatcgtt tgactgacat aaacctagga	300
aatactggat taagatagat cagtaggatt aagatccact gatgtaattt cccactgatt	360
tgggtggctga catgtggacc tgagagttgt gtgggctcac atgtcaaatac acggtgaaca	420
gtacgtcacg atatgttaga gggtcctctt ccggagatac ttatacgaat tttgcggaaa	480
cctgcaaact ttgatggacg attgaggcga gtttagttct aaattttttc ttcaaacttc	540
taactttttc atcacatcgt ttcaatttca atcaaacttc caatgttgac gtgaactaaa	600
cacacctatg agatatgaga agcgggttga cacttgacaa gtcctgacat gctgtgttgg	660
cgtgggccc acctgccacg tcaggtccag ctccgggtgg ttgggtttgg tgctttccga	720
taggcacgag ctc	733

<210> 2

<211> 1455

<212> DNA

<213> Hordeum vulgare

<400> 2

atggctgcca ccgtcgccgt ggacaacctg aacccaagg ttttaaaatg tgagtatgct	60
gtgcgtggag agattgtcat ccatgctcag cgcttgacag aacagctaaa gactcaacca	120
gggtctctac cttttgatga gatcctctat tgtaacattg ggaaccaca atctcttgggt	180
cagcaaccag ttacattctt caggagggtt cttgcccttt gtgatcatcc agacctgttg	240
caaagagagg aaatcaaac attgttcagt gctgattcta tttctcgagc aaagcagatt	300

pctca2012050622-seq1.txt

```

cttgccatga tacctggaag agcaacagga gcatacagcc atagccaggg tattaaagga      360
cttcgtgatg caattgcttc tgggatcgct tcacgagatg gattccctgc taatgctgat      420
gacatttttc tcacagatgg agcaagtcct ggggtgcacc tgatgatgca attactgata      480
aggaatgaga aagatggcat tcttgtcccg attcctcagt accccttgta ctcggtttcc      540
atagctcttc atggcggagc tcttgtccca tactatctca atgaatcgac gggctggggg      600
ttggaaacct ctgatgtaa gaagcaactt gaagatgctc ggtcaagagg catcaacggt      660
agggctttgg tggttatcaa tccaggaaat ccaactggac aggtacttgc tgaagaaaac      720
caatatgaca tagtgaagtt ctgcaaaaat gagggctcttg ttcttctagc tgatgaggta      780
taccaagaga acatctatgt tgacaacaag aaattccact ctttcaagaa gatagtgaga      840
tccttgggat acggcgagga ggatctccct ctagtatcat atcaatctgt ttctaaggga      900
tattatgggtg agtgtggtaa aagaggtggg tactttgaga ttactggctt cagtgtctca      960
gtaagagagc agatctacaa aatagcatca gtgaacctat gctccaatat cactggccag     1020
atccttgcta gtcttgcat gaaccacca aaggctagtg atgaatcata cgcttcatac     1080
aaggcagaaa aagatggaat cctcgcatct ttagctcgtc gtgcgaaggc attggagcat     1140
gcattcaata aacttgaggg aattacttgc aacgaggctg aaggagcaat gtacgtgttc     1200
cctcaaactc gtctgccaca gaaggcaatt gaggtgcta aagctgctaa caaagcacct     1260
gatgcattct atgctcttcg tctcctcgag tcgactggaa tcgtcgttgt ccctggatca     1320
ggatttggcc aggttcctgg cacatggcac ttcaggtgca cgatccttcc gcaggaggat     1380
aagatcccg cagtcatctc ccgcttcacg gtgttccatg aggcgttcat gtcagagtat     1440
cgtgactaaa ctggt                                     1455

```

```

<210> 3
<211> 482
<212> PRT
<213> Hordeum vulgare

```

```
<400> 3
```

```
Met Ala Ala Thr Val Ala Val Asp Asn Leu Asn Pro Lys Val Leu Lys
1          5          10          15
```

```
Cys Glu Tyr Ala Val Arg Gly Glu Ile Val Ile His Ala Gln Arg Leu
          20          25          30
```

```
Gln Glu Gln Leu Lys Thr Gln Pro Gly Ser Leu Pro Phe Asp Glu Ile
          35          40          45
```

```
Leu Tyr Cys Asn Ile Gly Asn Pro Gln Ser Leu Gly Gln Gln Pro Val
          50          55          60
```

```
Thr Phe Phe Arg Glu Val Leu Ala Leu Cys Asp His Pro Asp Leu Leu
65          70          75          80
```

pctca2012050622-seq1.txt

Gln Arg Glu Glu Ile Lys Thr Leu Phe Ser Ala Asp Ser Ile Ser Arg
85 90 95

Ala Lys Gln Ile Leu Ala Met Ile Pro Gly Arg Ala Thr Gly Ala Tyr
100 105 110

Ser His Ser Gln Gly Ile Lys Gly Leu Arg Asp Ala Ile Ala Ser Gly
115 120 125

Ile Ala Ser Arg Asp Gly Phe Pro Ala Asn Ala Asp Asp Ile Phe Leu
130 135 140

Thr Asp Gly Ala Ser Pro Gly Val His Leu Met Met Gln Leu Leu Ile
145 150 155 160

Arg Asn Glu Lys Asp Gly Ile Leu Val Pro Ile Pro Gln Tyr Pro Leu
165 170 175

Tyr Ser Ala Ser Ile Ala Leu His Gly Gly Ala Leu Val Pro Tyr Tyr
180 185 190

Leu Asn Glu Ser Thr Gly Trp Gly Leu Glu Thr Ser Asp Val Lys Lys
195 200 205

Gln Leu Glu Asp Ala Arg Ser Arg Gly Ile Asn Val Arg Ala Leu Val
210 215 220

Val Ile Asn Pro Gly Asn Pro Thr Gly Gln Val Leu Ala Glu Glu Asn
225 230 235 240

Gln Tyr Asp Ile Val Lys Phe Cys Lys Asn Glu Gly Leu Val Leu Leu
245 250 255

Ala Asp Glu Val Tyr Gln Glu Asn Ile Tyr Val Asp Asn Lys Lys Phe
260 265 270

His Ser Phe Lys Lys Ile Val Arg Ser Leu Gly Tyr Gly Glu Glu Asp
275 280 285

Leu Pro Leu Val Ser Tyr Gln Ser Val Ser Lys Gly Tyr Tyr Gly Glu
290 295 300

Cys Gly Lys Arg Gly Gly Tyr Phe Glu Ile Thr Gly Phe Ser Ala Pro
305 310 315 320

Val Arg Glu Gln Ile Tyr Lys Ile Ala Ser Val Asn Leu Cys Ser Asn
325 330 335

Ile Thr Gly Gln Ile Leu Ala Ser Leu Val Met Asn Pro Pro Lys Ala
340 345 350

pctca2012050622-seq1.txt

Ser Asp Glu Ser Tyr Ala Ser Tyr Lys Ala Glu Lys Asp Gly Ile Leu
355 360 365

Ala Ser Leu Ala Arg Arg Ala Lys Ala Leu Glu His Ala Phe Asn Lys
370 375 380

Leu Glu Gly Ile Thr Cys Asn Glu Ala Glu Gly Ala Met Tyr Val Phe
385 390 395 400

Pro Gln Ile Cys Leu Pro Gln Lys Ala Ile Glu Ala Ala Lys Ala Ala
405 410 415

Asn Lys Ala Pro Asp Ala Phe Tyr Ala Leu Arg Leu Leu Glu Ser Thr
420 425 430

Gly Ile Val Val Val Pro Gly Ser Gly Phe Gly Gln Val Pro Gly Thr
435 440 445

Trp His Phe Arg Cys Thr Ile Leu Pro Gln Glu Asp Lys Ile Pro Ala
450 455 460

Val Ile Ser Arg Phe Thr Val Phe His Glu Ala Phe Met Ser Glu Tyr
465 470 475 480

Arg Asp

<210> 4
<211> 931
<212> DNA
<213> artificial sequence

<220>
<223> PBpr2 promoter

<400> 4
gaattcatat agtatctttg tatgttgagc attagtattc agtgagttaa aactttatag 60
ttgttagtgg atgagtttga gttttaggat ttagtggggg attataattt catggtaaga 120
taggatgagt atatataatt agggattaaa ttacaatatt ttaaggatca tatcgatatt 180
caaaacaata tgttttttta agatggatgg aatcaatttt ctgagatatg tcccgctggg 240
gtgtccacaa tctgcacgcc atatctgact acacatatac aacgcccata ctctgtcgat 300
ctccttgtgt aaacacaacc gtccgtgaca cgtacgtaca tgggggatag cttgcctctc 360
cgttcagttg cgtgacaact aataaaatac taacacaatg tcttaatcct ccaaaaaaaaa 420
actcctaaat aattaattaa agataaggca agaaattaat ccaaatagtc caaaacaata 480
ggataaagag gagagcaaaa ctccaaatgg tcacacgtct ctcggtgtca ttttctcgtc 540
ggcttgcttt gtttgattcc agagtttccc aaactctcta gagagcaagt caaacagatc 600
tcatcgatcg cccaagaaga agacacatcc acaaataata atctcatctt atcgactaa 660

pctca2012050622-seq1.txt

tttatcttaa tacagtgtaa taaactatat cagtttgaga ttttttttc ctgtctaaat	720
tatctccaaa ccgtaggaat tttcgcaccc gacgatctct tttcttgatt tttcaaacca	780
aacggaccaa tctctctcgc gtgtatatat agcgagggtg agctcgagct ccatgcctgc	840
ttcctcctga ctctctctct gcattctctc gcttctgacg gcggcaacta acctttttgg	900
gattcgattc gaggaggagg tgatgggatc c	931

<210> 5
 <211> 973
 <212> DNA
 <213> artificial sequence

<220>
 <223> OsANT1 promoter

<400> 5	
aggaagtgat ttttagcgta gctgtgtttg tagcgtaatt gcgtaaagtc ctttcaattt	60
tgctatatct cactcgaaag attttttctt atctctcact cgattttctc actcaaattt	120
acagtgtatt ttcttgtaag ttacagtgtg atttatgaaa cttacactgt aacttttgta	180
agttacactg taatttttga atcttcacat gtaaatttta aattttgtat tggatttggg	240
ctttttcttg aggatatggg aatttaatgt tcattatggg gtttcttaat tgctttttgc	300
tttttattat atctatcgga ttttaataca aagattaaaa atctgtgtga tacgattata	360
aaaatctttc gaaagatgta taggtactcc caagcccttt taagaaagtt tttcaagaca	420
aaagtttttg gatgaaagggt agttataggg aaaaaggaat gtgcgtttat gtttatttgc	480
attgcttatt agcaaccaa aactaatcta taagtaaadc ttttatatac gtgcgcttaa	540
taattcaaaa gcaaatcat gtaaaataaa atgcgatgaa gaaactttta aaagttatca	600
aatttagatt ttattaaatt ttagtttaca agagcgctac gatgaaggct ttaaaaagat	660
gggaaaataa aacctttgac ctttctggac ttcaccaaac agctcacgct ttcggcttcg	720
tgccgtctcg tcccggtgcta ctgctacccc ctctgaccc caccgcccac tccacgtcc	780
cttctcctcc ctttcccgtg acacacagtc cccactccac cgctccgta taagtatccc	840
ttccttaccg ccggccagcc acagccaccg cctccccac cccacccga tcccctcccc	900
gccgtacggg cgcagaagga acccgtcttc tagaaggagg aggagggcta cctctctctc	960
tctctcttct gcc	973

<210> 6
 <211> 745
 <212> DNA
 <213> artificial sequence

<220>
 <223> PBpr1 promoter sequence including start codon

<400> 6	
gaattctgaa agtttccgtc caaatcgcac cttttaaccg ttgaaaaac atacaaacga	60
aaaataatct atatcttaat caggaagaaa gagtacgaaa tggatgaaccg tcgaaactat	120

pctca2012050622-seq1.txt

tcatatacgt cgtctgtctc atgaaaaaaaa aaatcaatcc agaaggatac gagacacttt	180
tacttcaaca aatatagaca tgagcttatt ctactagggtt tggttgttta ataagacgaa	240
agaaatacat tggttagttt ttcattaaaa aataatcggtt tgactgacat aaacctagga	300
aatactggat taagatagat cagtaggatt aagatccact gatgtaattt cccactgatt	360
tggtaggctga catgtggacc tgagagttgt gtgggctcac atgtcaaadc acggtgaaca	420
gtacgtcacg atatgttaga gggtcctctt ccggagatac ttatacgaat ttgtaggaaa	480
cctgcaaact ttgatggacg attgaggcga gtttagttct aaatTTTTTc ttcaaacttc	540
taactTTTTc atcacatcgt ttcaatttca atcaaacttc caatgttgac gtgaactaaa	600
cacacctatg agatatgaga agcgggttga cacttgacaa gtcctgacat gctgtgttgg	660
cgtgggcccc acctgccacg tcaggtccag ctccgggtgg ttgggtttgg tgctttccga	720
taggcacgag ctcatggctg ccacc	745