



US 20130210007A1

(19) **United States**(12) **Patent Application Publication**
Godler(10) **Pub. No.: US 2013/0210007 A1**(43) **Pub. Date: Aug. 15, 2013**(54) **TREATMENT AND DIAGNOSIS OF
EPIGENETIC DISORDERS AND
CONDITIONS**(75) Inventor: **David Eugeny Godler, Ormond (AU)**(73) Assignee: **MURDOCH CHILDRENS
RESEARCH INSTITUTE,
PARKVILLE, VICTORIA (AU)**(21) Appl. No.: **13/814,911**(22) PCT Filed: **Aug. 11, 2011**(86) PCT No.: **PCT/AU11/01024**§ 371 (c)(1),
(2), (4) Date: **Feb. 7, 2013**(30) **Foreign Application Priority Data**

Aug. 11, 2010 (AU) 2010903595

Publication Classification(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G06F 19/24 (2006.01)(52) **U.S. Cl.**CPC . **C12Q 1/68** (2013.01); **G06F 19/24** (2013.01)
USPC **435/6.11; 702/20**(57) **ABSTRACT**

The present disclosure relates generally to the field of epigenetics and in particular epigenetic profiles associated with a pathological condition. The present specification teaches screening of individuals and populations for epigenetic profiles associated with a pathological condition. The epigenetic profiles can be from an intron, an intron/exon boundary or a splicing region. Epigenetic profiles are disclosed from the following sites in the FMR locus: FREES, intron 2 of FMR1, the genomic FREE2 region as a whole or specific FREE2 fragments including FREE2 (D) or FREE2 (E). Kits and diagnostic assays are also taught herein as are computer programs to monitor changes in epigenetic patterns and profiles. Further enabled herein is a method for screening for agents which can reduce or mask the adverse effects of epigenetic modification and the use of these agents in therapy and prophylaxis.




Figure 1A




Figure 2




Figure 3




Figure 4A




Figure 4B




Figure 4B continued




Figure 4B continued




Figure 4B continued




Figure 4B continued




Figure 4B continued






Figure 4C continued




Figure 4D




Figure 4D continued




Figure 5




Figure 6




Figure 7

TREATMENT AND DIAGNOSIS OF EPIGENETIC DISORDERS AND CONDITIONS

[0001] This application is associated with and claims priority from Australian Provisional Patent Application No. 2010903595, filed on 11 Aug. 2010, entitled "Treatment and diagnosis of epigenetic disorders and conditions", the entire contents of which, are incorporated herein by reference.

FIELD

[0002] The present disclosure relates generally to the field of epigenetics and in particular epigenetic profiles associated with a pathological condition. The present specification teaches screening of individuals and populations for epigenetic profiles associated with a pathological condition. Kits and diagnostic assays are also taught herein as are computer programs to monitor changes in epigenetic patterns and profiles. Further enabled herein is a method for screening for agents which can reduce or mask the adverse effects of epigenetic modification and the use of these agents in therapy and prophylaxis.

BACKGROUND

[0003] Bibliographic details of the publications referred to in this specification are also collected at the end of the description.

[0004] Reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common general knowledge in any country.

[0005] It is apparent that DNA methylation and other epigenetic modifications play a role in the regulation of gene expression in higher organisms. The importance of epigenetic modification has been highlighted by its involvement in several human diseases. Methylation, for example, of cytosine at the 5' position is the only known methylation modification of genomic DNA. In particular, methylation of CpG islands within regulatory regions of the genome appears to be highly tissue specific. Methylation of cytosines distal to the islands is also important. These regions are called "shores" or "island shores" (Irizarry et al., *Nature Genetics* 41(2):178-186, 2009). Epigenetic modifications include histone modification, changes in acetylation, methylation, ubiquitylation, phosphorylation, sumoylation, activation or deactivation, chromatin altered transcription factor levels and the like.

[0006] Another genetic condition which can affect gene expression arises from expansion or increase in the number of repeats in a specific tandem repeat array. Such nucleotide expansion can result in repeat expansion disease conditions. A critical threshold of repeat expansion determines the level of pathogenicity (Orr and Zoghbi, *Ann Rev Neurosci* 30:575-621, 2007). Many diseases arise from expansion of a repeat located in an open reading frame resulting in a protein with a long polyQ² tract that is toxic to neurons (On and Zoghbi, 2007 supra). Other expansion disease conditions such as Fragile X syndrome (FXS), Fragile XE mental retardation (FRAXE), Fragile X-associated primary ovarian insufficiency (FXPOI), Fragile type, folic acid type, rare 12 (FRA12A), mental retardation (MR), Friedrich's ataxia (FRDA) and myotonic dystrophy (DM), arise from altered transcription of the repeats which are not translated.

[0007] A particular type of expansion disorder is referred to as a trinucleotide repeat disorder (also known as trinucleotide

repeat expansion disorder, triplet repeat expansion disorder and codon reiteration disorder) and results from trinucleotide repeats in certain genetic loci. An example occurs in the Fragile X Mental Retardation genetic locus ("FMR genetic locus").

[0008] The FMR genetic locus includes the FMR1 gene which is composed of 17 exons, spanning 38 Kb, and encodes Fragile X Mental Retardation Protein (FMRP), essential for normal neurodevelopment (Verkerk et al., *Cell* 65(5):905-914, 1991; Terracciano et al., *Am J Med Genet C Semin Med Genet* 137C(1):32-37, 2005). A CGG repeat segment is located within the 5' untranslated region (UTR) of the gene. Its normal range is <40 repeats. When expanded, these repeats have been implicated in a number of pathologies, including the Fragile X syndrome (FXS), Fragile X-associated Tremor Ataxia Syndrome (FXTAS) and Fragile X-associated primary ovarian insufficiency (FXPOI; formerly referred to as Premature Ovarian Failure [POF]). FXS is neurodevelopmental in nature with a frequency of 1/1400 males and 1/8000 females, associated with a Fragile site at the Xq27.3 locus (Jin and Warren, *Hum. Mol. Genet* 9(6):901-908, 2000).

[0009] This syndrome is caused by a CGG expansion to "full mutation" (FM) which comprises >200 repeats, leading to a gross deficit of FMRP and subsequent synaptic abnormalities (Pieretti et al., *Cell* 66(4):817-822, 1991; Irwin et al., *Cereb Cortex* 10(10):1038-1044, 2000). The FXS clinical phenotype ranges from learning disabilities to severe mental retardation and can be accompanied by a variety of physical and behavioral characteristics. FXTAS is prevalent in ~30% of premutation individuals (PM), comprising 55 to 199 repeats (Nolin et al., *Am J Hum Genet* 72(2):454-464, 2003) and is a progressive neurodegenerative late-onset disorder with a frequency of 1/3000 males in the general population (Jacquemont et al., *Am J Ment Retard* 109(2):154-164, 2004), manifesting as tremor, imbalance and distinct MRI and histological changes (Hagerman et al., *Neurology* 57(1):127-130, 2001; Jacquemont et al., *J Med Genet* 42(2):e14, 2005; Loesch et al., *Clin Genet* 67(5):412-417, 2005). It is often associated with 'toxicity' of elevated FMR1 mRNA, which has been linked to the intranuclear inclusions and cell death observed during neurodegeneration (Jin et al., *Neuron* 39(5):739-747, 2003).

[0010] FXTAS can occur in females carrying PM, but with much lower frequency as can be expected from X-linked inheritance. The intermediate or Gray Zone (GZ) alleles comprising 41 to 54 repeats (Bodega et al., *Hum Reprod* 21(4):952-957, 2006) are the most common form of the expansion, 1 in 30 males and 1 in 15 females. As with PM alleles, increased levels of FMR1 mRNA have been reported in the GZ individuals, proportional to the size of CGG expansion (Kenneson et al., *Hum Mol Genet* 10(14):1449-1454, 2001; Mitchell et al., *Clin Genet* 67(1):38-46, 2005; Loesch et al., *J Med Genet* 44(3):200-204, 2007). Female carriers of both PM and GZ allelic types have an increased risk of developing POF (Allingham-Hawkins et al., *Am J Med Genet* 83(4):322-325, 1999; Sullivan et al., *Hum Reprod* 20(2):402-412, 2005) which has incidence of approximately 1% in the general population, and often unknown etiology (Coulam, *Fertil Steril* 38(6):645-655, 1982).

[0011] Expansion related abnormalities in FMR1 are involved in pathologies with a wide spectrum of pathomechanisms all pointing to involvement of multiple factors at the Xq27.3 locus in addition to FMR1. A number of antisense

transcripts have been described embedded within the FMR1 sequence, ASFMR1 (Ladd et al., *Hum Mol Genet* 16(24): 3174-3187, 2007) and FMR4 (Khalil et al., *PLoS ONE* 3(1): e1486, 2008).

[0012] The ASFMR1 and FMR4 transcripts have been suggested to share the bi-directional promoter with FMR1, which is heavily regulated by the state of the surrounding chromatin environment (Pietrobono et al., *Nucleic Acids Res* 30(14): 3278-3285, 2002; Chiurazzi et al., *Hum Mol Genet* 7(1): 109113, 1998).

[0013] Transcription of ASFMR1 is also regulated by another promoter located in the exon 2 of FMR1, with the resulting transcript spanning the CGG repeat in the antisense direction (Ladd et al., 2007, supra), and an open reading frame (ORF) with the CGG encoding a polyproline peptide (Ladd et al., 2007, supra). FMR4, however, is a long non-coding RNA, involved in regulation of apoptosis (Khalil et al., 2008, supra).

[0014] The length of the CGG repeat has been reported to effect transcription of all three genes FMR1, FMR4 and ASFMR1 (Ladd et al., 2007, supra; Khalil et al., 2008, supra). However, although it is well documented that FMR1 transcription is promoter methylation dependent, linked to the CGG expansion size, the relationship between FMR4 and ASFMR1 transcription and methylation remains elusive.

[0015] One of the current problems is in the diagnosis of subjects with FM in the FMR genetic locus. Diagnostic assays targeting only the CGG expansion have hitherto been inconclusive. Therefore, currently Southern DNA analysis, which is expensive and time consuming, is used as a gold standard assay for diagnosis in many laboratories.

[0016] Despite the availability of a range of methylation and nucleotide expansion assays (see, for example, Rein et al., *Nucleic Acids Res.* 26:2255, 1998 in relation to methylation assays), selection of regions to amplify and screen is an important aspect of determining an epigenetic profile characteristic of a disease condition. There is a need to identify crucial regions which are associated with epigenetic change linked to a pathological condition to assay and/or therapeutically target.

SUMMARY

[0017] Throughout this specification, unless the context requires otherwise, the word "comprise" or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or method step or group of elements or integers or method steps but not the exclusion of any element or integer or method step or group of elements or integers or method steps.

[0018] Nucleotide and amino acid sequences are referred to by a sequence identifier number (SEQ IN NO:). The SEQ ID NOs: correspond numerically to the sequence identifiers <400>1 (SEQ ID NO:1), <400>2 (SEQ ID NO:2), etc. A summary of sequence identifiers is given in Table 1.

[0019] Aspects enabled herein are predicated in part on the determination of an association between epigenetic modification of intronic regions including intron/exon boundaries and splicing regions within a genetic locus and a pathological condition including a trinucleotide expansion disorder. In an embodiment, the epigenetic modification occurs in:

[0020] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or

[0021] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

within a genetic locus.

[0022] In an embodiment, the epigenetic modification occurs within a genetic locus which leads to a condition including a pathoneurological condition such as a pathoneurodevelopmental and pathoneurodegenerative condition as well as a non-neurological condition. Conditions and disorders including trinucleotide expansion disorders associated with a change in the epigenetic profile from that observed in healthy controls associated with a change in epigenetic profile in intronic regions including intron/exon boundary regions, include Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidol-luysianatrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCAT), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCAB) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome. Reference to a "control" means relative to a healthy subject which means a subject with a normal size of expansion repeats and/or who is phenotypically normal meaning that the subject does not have symptoms of, for example, a trinucleotide expansion disorder and/or is the epigenetic (e.g. methylation) profile observed in healthy control subjects.

[0023] It is proposed herein that epigenetic changes in an intron, intron/exon boundary and/or splicing region within a particular genetic locus are associated with the development, progression and severity of a range of pathological conditions including trinucleotide expansions disorders such as but not limited to those listed above. In relation to the FMR genetic locus, the epigenetic modification may occur in:

[0024] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or

[0025] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

within the genetic locus. These regions also include promoter regions. Epigenetic changes may occur on either strand of double stranded genomic DNA including either strand of a promoter or other regulatory region and either strand may be targeted for epigenetic analysis. The ASFMR1 promoter is an example. The location of this promoter can be seen in FIGS. 6A and 6C and has a transcription start site in FREE3 of intron 2. Hence, another aspect of the present disclosure provides a method for identifying FXS or a related condition in a human subject, the method comprising screening for a change relative to a control in the extent of epigenetic modification in the FMR genetic locus at a location selected from:

[0026] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ

ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions; and

[0027] (ii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

wherein a change in extent of epigenetic modification relative to a control is indicative of the presence or severity of the pathological condition or a propensity to develop same.

[0028] In an embodiment, a method is provided for identifying a trinucleotide expansion disorder in a mammalian subject including a human, the method comprising screening for a change relative to a healthy control in the extent of epigenetic modification within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus; wherein a change in extent of epigenetic modification relative to the control is indicative of the presence or severity of the trinucleotide expansion disorder or a propensity to develop same wherein the intron, intron/exon boundary and/or splicing region is selected from the list consisting of:

[0029] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0030] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions; and

[0031] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions.

[0032] By “epigenetic modification” is meant changes in extent, level and/or profile or epigenetic modification including changes in methylation including hypermethylation and hypomethylation, histone modification, acetylation, ubiquitylation, phosphorylation and/or sumoylation, as well as changes in chromatin altered transcription factor levels and the like leading to activation or deactivation of genetic locus expression. The epigenetic modification extends to an increase or decrease in epigenetic change relative to a normal control. It also extends to either strand of these target regions which includes either strand of a promoter region. In an embodiment, epigenetic modification includes the methylation state of CpG and CpNpG sites within an intron of a genetic locus. In an embodiment, the genetic locus is the FMR genetic locus which includes FMR1, FMR4 and ASFMR1 genes. In an embodiment, the epigenetic modification occurs in the FMR genetic locus within a region selected from:

[0033] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0034] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0035] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0036] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0037] (v) the FREE3 region; and

[0038] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 region alone or in combination with the FREE1 region (D) region or FREE2 (D)/(E) boundary.

[0039] These regions include either strand of a double stranded genomic sequence and includes either strand of the promoter region such as the ASFMR1 promoter with a transcription initiation site in FREE3 of intron 2.

[0040] Furthermore, the epigenetic profile of the FMR genetic locus is also informative as to the spectrum of disease conditions associated with the genetic locus, such as whether the subject is normal or has a PM, GZ or FM pathology and/or whether the epigenetic change and/or CGG expansion is heterozygous or homozygous at the FMR allele. Reference to “FREE2” include FREE2 (A), FREE2 (B), FREE2 (C), FREE2 (D) and FREE2 (E) including any exon/intron boundaries therein such as the FREE2(D)/FREE2 (E) boundary. The boundary regions include a promoter region. A “promoter region” includes either or both nucleic acid strands within double stranded genomic DNA.

[0041] Accordingly, an aspect enabled herein is a method for identifying an epigenetic profile in the genome of a cell indicative of a pathological condition, the method comprising screening for a change relative to a control in the extent of epigenetic modification within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of a genetic locus wherein the extent of epigenetic change relative to a control is indicative of the presence or severity of the pathological condition or a propensity to develop same.

[0042] In an embodiment, the epigenetic profile is determined within a genetic locus, the epigenetic profile of which, is associated with a pathological condition including a trinucleotide disorder associated with a change in epigenetic profile from that observed in healthy controls selected from Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidol-luysianatrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCAT), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome.

[0043] In an embodiment, the epigenetic change is within the FMR genetic locus and is associated with one or more of FXS, FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and/or cognitive impairment.

[0044] In relation to the latter aspect, the method comprises screening for a change relative to the control in the extent of epigenetic modification within the FMR genetic locus within a region selected from:

[0045] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0046] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0047] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0048] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0049] (v) the FREE3 region; and

[0050] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including FREE2 (D) or FREE2(D)/(E) boundary. These regions may be assessed on either strand of double stranded genomic DNA and include either strand of a promoter region. An example of a promoter region is the ASFMR1 promoter with an initiation site in FREE3 of intron 2 (see FIG. 6).

[0051] The extent of epigenetic change is indicative of the presence of severity of the pathological condition or a propensity to developing same. As indicated above, a "pathological condition" includes a trinucleotide expansion disorder.

[0052] A further aspect taught herein is a method for identifying an epigenetic profile in the genome of a cell indicative of a pathological condition associated with the FMR genetic locus, the method comprising extracting genomic DNA from the cell and subjecting the DNA to an amplification reaction using primers selective for (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region, within the FMR genetic locus including the FMR1 gene, subjecting the amplified DNA to an epigenetic assay to determine the extent of epigenetic modification of the DNA wherein a change in the extent of epigenetic modification is indicative of the presence or severity of the pathological condition or propensity to develop same. The region within the FMR genetic locus is selected from:

[0053] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0054] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0055] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0056] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0057] (v) the FREE3 region; and

[0058] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0059] A similar method applies to epigenetic changes in other genetic loci. Reference to the "FMR genetic locus"

includes the FMR1, FMR4 and ASFMR1 genes and corresponds to Xq27.3 as well as promoter regions associated with these sites. The term "FMR locus" means the "FMR genetic locus". In an embodiment, an aspect taught herein determines that the intronic region downstream of intron 1 comprises Fragile X-related Epigenetic Element 3 as defined by SEQ ID NO:1 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions; or is intron 2 as defined by SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions. The nucleotide sequence of intron 1 of the FMR1 gene is set forth in SEQ ID NO:3. The genomic nucleotide sequence of FREE2 region alone or in combination with the FREE1 region (D), FREE2 (E) and FREE3 are set forth in SEQ ID NOs:48, 49 and 47, respectively, and the present disclosure extends to homologs thereof having at least 80% identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or a complement thereof under medium stringency conditions. The present disclosure further contemplates amplifying all or part of an expansion mutation and/or detecting extent of epigenetic change therein in combination with an epigenetic change (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus including the FMR1 gene. The extent of epigenetic change in two or more of an intron, an intron/exon boundary and/or a splicing region or in one seventh or greater of an intron within the FMR genetic locus may be determined alone or in combination with extent of (CGG)_n expansion and/or any other epigenetic change therein. The determination of epigenetic change may also be conducted in combination with an assay as contemplated by International Patent Application No. PCT/AU2010/000169 filed on 17 Feb. 2010, the contents of which are incorporated herein by reference in their entirety. In an embodiment, the epigenetic modification is a change in extent of methylation which includes hypermethylation and hypomethylation and profile of methylation,

[0060] Another aspect of the present disclosure contemplates a method for identifying a pathological condition in a mammalian subject including a human, the method comprising screening for a change relative to a control in the extent of change in methylation or other epigenetic modification within a region selected from:

[0061] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO: 1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0062] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0063] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ

ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0064] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0065] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0066] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein a change in epigenetic modification relative to a control is indicative of the presence or severity of the pathological condition or a propensity to develop same. In an embodiment, the pathological condition is a trinucleotide expansion disorder.

[0067] As indicated above, the region within the FMR genetic locus assayed for epigenetic change is selected from:

[0068] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0069] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0070] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0071] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0072] (v) the FREE3 region; and

[0073] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0074] Hence, in relation to detecting epigenetic changes in (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region or FREE3 within intron 2, the present disclosure enables the diagnosis monitoring or analyzing of a spectrum of neurodegenerative or neurodevelopmental pathologies such as Fragile X-related conditions including FXS, FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment. Certain tri-nucleotide disorders are also included.

[0075] A "modified" X-chromosome includes an inactivated X-chromosome or an X-chromosome having a skewed X-inactivation, or inversion, insertion, deletion, duplication or is a hybrid.

[0076] The epigenetic profile is determined in the genome of a cell of a subject. Any cell may be tested such as a cell from a post-natal or pre-natal human or embryo. More particularly, the cell is a cultured or uncultured chorionic villi sample (CVS) cell, a lymphoblast cell, a blood cell, a buccal cell, epithelial cells, fibroblast cells, an amniocyte or an EBV transformed lymphoblast cell line.

[0077] In a particular embodiment of the present disclosure, the epigenetic modification is methylation of CpG and/or CpNpG sites. Methylation is determined by a range of assays including bisulfite MALDI-TOF methylation assay. In an alternative embodiment, methylation is determined by use of methylation sensitive PCR, methylation specific melting

curve analysis (MS-MCA) or high resolution melting (MS-FIRM); quantification of methylation by MALDI-TOF MS; methylation specific MLPA; methylated-DNA precipitation and methylation-sensitive restriction enzymes (COMPARE-MS); single molecule (SMRT) sequencing; or methylation sensitive oligonucleotide microarray; or antibodies. Other methods include NEXT generation (GEN) and DEEP sequencing or pyrosequencing. However, any assay of methylation status may be employed. Regardless of the method, either strand of genomic double stranded DNA may be assessed for its epigenetic profile.

[0078] Further taught herein is a method for screening for an agent which modulates epigenetic change of (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within a genetic locus, the method comprising screening for a change relative to a control in the extent of epigenetic modification within the intron 1n the presence or absence of an agent to be tested, wherein an agent is selected if it induces a change in the epigenetic modification.

[0079] In an embodiment, the epigenetic modification is associated with a polyglutamine (polyQ) disease such as Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidololysianatrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCAT), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome. The "association" is based on a change in epigenetic profile from that of healthy controls.

[0080] In an embodiment, the agent modulates genetic change in the FMR genetic locus within a region selected from:

[0081] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0082] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0083] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0084] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0085] (v) the FREE3 region; and

[0086] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0087] In an embodiment, a method is also provided for screening for an agent which modulates epigenetic modification of an FMR genetic locus in a mammalian cell including a human cell, the method comprising screening for a change relative to a healthy control in the extent of epigenetic change in (i) two or more of (a) an intron; (b) an intron/exon bound-

ary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus; wherein the intron, intron/exon boundary and/or splicing region is selected from the list consisting of:

[0088] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0089] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions; and

[0090] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

in the presence or absence of an agent to be tested wherein the agent is selected if it induces a change in extent of epigenetic modification. Such an agent is useful in treating a trinucleotide expansion disorder or other pathological conditions.

[0091] A method is also provided for screening for an agent which modulates epigenetic change of an FMR genetic locus in a mammalian cell including a human cell, the method comprising screening for a change relative to a control in the extent of epigenetic modification within a region selected from:

[0092] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0093] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0094] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0095] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0096] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0097] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

in the presence or absence of an agent to be tested wherein the agent is selected if (i) it induces a change in extent of epigenetic modification and/or (ii) causes an improvement in disease phenotype based on the type and degree of epigenetic modification. A change in extent of methylation includes hypermethylation, hypomethylation and a change in the profile of methylation.

[0098] As indicated above, in an embodiment, the epigenetic modification is methylation. De-methylation as well as pro-methylation agents are contemplated herein.

[0099] Further enabled herein is a method for monitoring the treatment of a disease condition including trinucleotide expansion disorders associated with a change in epigenetic profile from that observed in healthy controls such as Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallid-oligodystrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCA7), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome, the method comprising screening for a change relative to the control in the extent of epigenetic modification within a genetic locus wherein the epigenetic profile of the genetic locus is associated with the disease or condition, wherein the extent of epigenetic change is indicative of the presence or severity of the pathological condition, wherein the treatment modulates the extent of epigenetic change of (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the genetic locus, the method comprising monitoring for a change relative to a control in a pre- and post-treatment sample in the extent of epigenetic modification within the intron, wherein a change in extent of epigenetic modification after or during treatment is indicative of effective treatment.

[0100] In an embodiment, the genetic locus is the FMR genetic locus and the disease or condition is FXS or a related condition such as FXTAS, FXPOI, autisms, mental retardation, a modified X-chromosome or cognitive impairment.

[0101] By "monitoring" in this context includes diagnosis of disease, monitoring progress of the disease before or after treatment, prognosis of the disease development or remission as well as the pharmacoresponsiveness or pharmacosensitivity of a subject or agent.

[0102] The present disclosure also teaches the use of an epigenetic profile within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of a genetic locus in a cell in the manufacture of an assay to identify an epigenetic profile of gene associated with a pathological condition. In an embodiment, the genetic locus is the FMR genetic locus and epigenetic change is monitored within a region selected from:

[0103] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0104] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0105] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0106] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0107] (v) the FREE3 region; and

[0108] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0109] An embodiment herein is directed to the use of an epigenetic profile within the FMR genetic locus in a mammalian cell including a human cell, the epigenetic profile including methylation of CpG and/or CpNpG sites located in a region selected from:

[0110] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0111] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0112] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0113] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0114] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0115] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

in the manufacture of an assay to identify an epigenetic profile of an FMR locus-associated pathological condition.

[0116] The assay taught herein may also be used alone or in combination with assays to detect extent of a nucleotide expansion such as a (CGG)_n expansion, such as using PCR and Southern blot assays. This is useful in determining homozygosity, heterozygosity and mosaicism of a disease or condition. The assay of the present disclosure is also useful in population studies such as epidemiological studies as well as studies based on ethnic populations. Accordingly, another aspect enabled herein provides a method of identifying epigenetic profile in populations of subjects indicative of a pathological condition associated with epigenetic modifications or changes in an intron, intron/exon boundary and/or splicing region, the method comprising screening for a

change, relative to a control in a statistically significant number of subjects, in the extent of epigenetic change within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of a genetic locus, the epigenetic change including extent of methylation of CpG and/or CpNpG sites located within the intron, intron/exon boundary and/or splicing region wherein a change in extent of epigenetic modification is indicative of the presence or severity of the pathological condition or a propensity to develop same.

[0117] In an embodiment, the epigenetic modification is determined in the FMR genetic locus within a region selected from:

[0118] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0119] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0120] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0121] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0122] (v) the FREE3 region; and

[0123] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0124] Contemplated herein is a method of identifying a methylation or other epigenetic profile in a population of subjects indicative of a pathological condition associated with the FMR locus, the method comprising screening for a change, relative to a control, in a statistically significant number of subjects in the extent of epigenetic modification including extent of change in methylation of CpG and/or CpNpG sites within a region selected from:

[0125] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0126] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0127] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0128] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0129] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0130] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein a change in extent of epigenetic modification is indicative of the presence of the pathological condition or a propensity to develop same in the population.

[0131] In accordance with this method the assay may comprise the further step of determining the extent of a nucleotide expansion such as a (CGG)_n expansion such as by PCR and/or Southern blot analysis. The regions investigated for epigenetic change within the FMR genetic locus include:

[0132] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0133] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0134] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0135] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0136] (v) the FREE3 region; and

[0137] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0138] Aspects herein extend to the use of the epigenetic profile of an intron within a genetic locus to determine the status, prognosis or disease development or recovery and/or treatment options including responsiveness of the subject to pharmacological agents and/or behavioral intervention strategies.

[0139] Computer programs to monitor changes in epigenetic modification or profile over time that may assist in making decisions regarding treatment options including responsiveness of the subject to pharmacological agents and/or behavioral intervention strategies, are also enabled herein.

[0140] Accordingly, another aspect provides a method of allowing a user to determine the status, prognosis and/or treatment response of a subject with respect to an FMR locus-associated pathology, the method including:

[0141] (a) receiving data in the form of extent of methylation or other epigenetic modification at a site within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of a genetic locus associated with the pathology, wherein the extent of methylation or other epigenetic modification provides a correlation to the presence, state, classification or progression of the pathology;

[0142] (b) transferring the data from the user via a communications network;

[0143] (c) processing the subject data via multivariate or univariate analysis to provide a disease value;

[0144] (d) determining the status of the subject in accordance with the results of the disease value in comparison with predetermined values; and

[0145] (e) transferring an indication of the status of the subject to the user via the communications network.

[0146] In an embodiment, the genetic locus is the FMR genetic locus. The epigenetic profile is determined within the FMR genetic locus from a region selected from:

[0147] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0148] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0149] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0150] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0151] (v) the FREE3 region; and

[0152] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0153] In an embodiment, a method is provided of allowing a user to determine the status, prognosis and/or treatment response of a subject with respect to an FMR locus-associated pathology, the method including:

[0154] (a) receiving data in the form of extent of methylation or other epigenetic modification at a site selected from:

[0155] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0156] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0157] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0158] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0159] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0160] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein the extent of methylation or other epigenetic modification provides a correlation to the presence, state, classification or progression of the pathology;

[0161] (b) transferring the data from the user via a communications network;

[0162] (c) processing the subject data via multivariate or univariate analysis to provide a disease index value;

[0163] (d) determining the status of the subject in accordance with the results of the disease index value in comparison with predetermined values; and

[0164] (e) transferring an indication of the status of the subject to the user via the communications network.

[0165] A further embodiment enabled herein is a kit comprising primers which amplify regions of the FMR genetic locus, comprising CpG and/or CpNpG sites located within a region selected from:

[0166] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0167] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0168] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0169] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0170] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0171] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

in the manufacture of a diagnostic kit or device to detect epigenetic modification of the FMR locus-associated with a pathological condition.

[0172] In an embodiment, the epigenetic modification relates to extent of, or change in, methylation at CpG and/or CpNpG sits within the selected regions of the FMR genetic locus, defined as FREE3, intron 2 and an intron, intron/exon boundary and/or splicing region downstream of intron 2 of the FMR gene. In an embodiment, the epigenetic modification occurs in the FMR genetic locus within a region selected from:

[0173] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0174] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0175] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0176] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0177] (v) the FREE3 region; and

[0178] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0179] In an embodiment, the primers useful in practicing the subject assay are selected from the list consisting of SEQ ID NOs:6 through 11. Those sequences include tag sequences. The present disclosure extends to the primer only portions of SEQ ID NOs:6 through 11.

TABLE 1

Summary of sequence identifiers	
SEQUENCE ID NO:	DESCRIPTION
1	Nucleotide sequence of FREE3 within the FMR1 gene
2	Nucleotide sequence of intron 2 of the FMR1 gene
3	Nucleotide sequence of intron 1 of FMR1 gene
4	Nucleotide sequence of FREE2 (B)
5	Nucleotide sequence of FREE2 (C)
6	Forward primer and tag sequence for FREE2 (B)
7	Reverse primer and tag sequence for FREE2 (B)
8	FREE 2 (C) forward primer and tag
9	FREE2 (C) reverse primer and tag
10	FREE3 forward primer and tag
11	FREE3 reverse primer and tag
12	Nucleotide sequence of regulatory motif GATA-1
13	Nucleotide sequence of regulatory motif HSF2
14	Nucleotide sequence of regulatory motif C/EBP
15	Nucleotide sequence of regulatory motif CdxA
16	Nucleotide sequence of regulatory motif AML-1a
17	Nucleotide sequence of regulatory motif AML-1a
18	Nucleotide sequence of regulatory motif CdxA
19	Nucleotide sequence of regulatory motif CdxA
20	Nucleotide sequence of regulatory motif CdxA
21	Nucleotide sequence of regulatory motif HFH-1/HFH-2
22	Nucleotide sequence of regulatory motif Cdx2
23	Nucleotide sequence of regulatory motif SRY
24	Nucleotide sequence of regulatory motif SRY
25	Nucleotide sequence of regulatory motif SRY
26	Nucleotide sequence of regulatory motif S8
27	Nucleotide sequence of regulatory motif SRY
28	Nucleotide sequence of regulatory motif CdxA
29	Nucleotide sequence of regulatory motif Oct-1
30	Nucleotide sequence of intron1 downstream of FREE2 (C)
31	Nucleotide sequence of exon2 upstream of FREE3
32	Nucleotide sequence of CGG amplification primer (r)
33	Nucleotide sequence of CGG amplification primer (f)
34	ASFMR1 (-1) forward primer
35	ASFMR1 (-1) reverse primer
36	ASFMR1 (-1) probe
37	ASFMR1 (-2) forward primer
38	ASFMR1 (-2) reverse primer
39	ASFMR1 (-2) probe
40	ASFMR1 (-3) forward primer
41	ASFMR1 (-3) reverse primer
42	ASFMR1 (-3) probe
43	Forward primer FREE2 (D)
44	Reverse primer FREE2 (D)
45	Forward primer FREE2 (E)
46	Reverse primer FREE2 (E)
47	Genomic target sequence FREE3
48	Genomic target sequence FREE2 (D)
49	Genomic target sequence FREE2 (E)
50	Forward primer FREE3
51	Forward primer FREE3
52	Tag forward primer FREE2 (D)
53	Tag reverse primer FREE2 (D)
54	Tag forward primer FREE2 (E)
55	Tag reverse primer FREE2 (E)
56	Tag forward primer FREE3
57	Tag reverse primer FREE3

[0180] A list of abbreviations used herein is provided in Table 2.

TABLE 2

Abbreviations	
ABBREVIATION	DESCRIPTION
Ab	Antibody
ASFMR1	Antisense Fragile X mental retardation 1 gene

TABLE 2-continued

Abbreviations	
ABBREVIATION	DESCRIPTION
(CGG) _n	CGG repeat element located within 5' untranslated region of the FMR1 gene
CpG	Cytosine and guanine separated by a phosphate (C-phosphate-G), which links the two nucleosides together in DNA
CpNpG	Cytosine and guanine separated by a nucleotide (N) where N is any nucleotide but guanine. The cytosine and N nucleotide are phosphorylated.
CVS	Cultured or uncultured Chorionic Villi Sample
DM	Myotonic dystrophy
DNA	Deoxyribonucleic acid
DRPLA	dentatonubropallid oluysiantrophy
FIQ	Full scale IQ
FM	Full Mutation
FMR	Fragile X mental retardation genetic locus comprising of FMR1 and FMR4 genes
FMR1	Fragile X mental retardation 1 gene
FMRP	Fragile X mental retardation protein
FRA12A	Fragile type, folic acid type, rare 12
FRAXE	Fragile X E mental retardation
FRDA	Friedrich's ataxia
FREE	Fragile X related Epigenetic Element (e.g. FREE2 and FREE3)
FREE2 (D)/(E) boundary	Boundary of FREE2 (D) and FREE2 (E)
FXPOI	Fragile X-associated primary ovarian insufficiency
FXS	Fragile X Syndrome
FXTAS	Fragile X-associated Tremor Ataxia Syndrome
GZ	Gray Zone
HD	Huntington's disease
HRM	Heat Resolution Melt
MR	Mental retardation
ORF	Open Reading Frame
PCR	Polymerase Chain Reaction
PM	Premutation
POF	Premature Ovarian Failure
PolyQ	Polyglutamine
SBMA	spinobullar muscular atrophy (Kennedy disease)
SCA1	spinocerebellar ataxia Type 1
SCA17	spinocerebellar ataxia Type 17
SCA2	spinocerebellar ataxia Type 2
SCA3	spinocerebellar ataxia Type 3
SCA6	spinocerebellar ataxia Type 6
SCA7	spinocerebellar ataxia Type 7
SCA8	spinocerebellar ataxia Type 8

BRIEF DESCRIPTION OF THE FIGURES

[0181] Some figures contain color representations or entities. Color photographs are available from the Patentee upon request or from an appropriate Patent Office. A fee may be imposed if obtained from a Patent Office.

[0182] FIG. 1A is a representation of the intron and exon regions 5' of the FMR1 CGG expansion (sequence numbering from GenBank L29074 L38501). (A) is a diagrammatic representation of the organization of the FMR genetic locus in relation to FMR1 and ASFMR1 transcription start sites, FMR1 promoter, the Fragile X-related epigenetic elements (FREE), FMR1 gene has 17 exons, and encodes FMRP. A CGG repeat is located within the 5' (UTR) of the FMR1 gene. ASFMR1 spans the CGG expansion in the antisense direction and is also regulated by another promoter located in the exon 2 of FMR1. The FREE2 located downstream of the CGG expansion. The FREE3 region is located within intron 2 of FMR1 downstream of the second ASFMR1 promoter. (B) Primers utilized for MALDI-TOF methylation analysis targeted 4 regions at the Xq27.3 locus designated as FREE2(A)

[described as amplicon 5 in Godler et al., *Hum Mol Genet* 10(8):1618-1632. [Epub 2010]; Godler et al., *J. Mol Diagn.* 2011 [Epub ahead of print] PMID:21723415; HMG); FREE2 (B); FREE2(C) and FREE3 (color coded). Individual CPG sites within each region are numbered accordingly. Prominent transcription factor binding sites and methylation sensitive restriction enzyme recognition sites are indicated in capital font, and are listed/identified in Table 1. << Indicates ASFMR1 transcription start site.

[0183] FIG. 2 is a graphical representation of the methylation pattern variation between healthy controls and FXS individuals within the body of the FMR1 gene, 5' of the CGG expansion. DNA from lymphoblasts of (A) healthy controls (n=4) and (B) Fragile X syndrome affected patients (n=3). Methylation of individual CpGs, were analyzed within the 9.762 kb region 5' CGG expansion, inclusive of intron 1, exon 2 and intron 2 (sequence numbering from GenBank L29074 L38501) using 3 SEQUENOM mass spectrometry assays (see Table 2). *—represent missing values. OL—represent CpG unit/s with overlaying fragment peak; DUP—represent CpG unit's with peak/s of the same size.

[0184] FIGS. 3A through C are graphical representations of a spiking experiment indicating the quantification limits of the MALDI-TOF methylation analysis of the Fragile X Syndrome (FXS) DNA within the 9.762 kb region 5' CGG expansion, inclusive of intron 1, exon 2 and intron 2 (sequence numbering from GenBank L29074 L38501) using 3 SEQUENOM mass spectrometry assays (A: FREE2(B); B: FREE2 (C); C: FREE3). Healthy control DNA was spiked with FXS DNA at 1:0; 2:1; 1:1; 1:2; 0:1 ratios corresponding to 0, 33.3, 50, 66.6, 100% FXS DNA in the sample. The spiked DNA samples were analyzed using MALDI-TOF methylation analysis at three sequential regions at the Xq27.3 locus (see FIG. 1 for locations). The methylated vs unmethylated ratios at each analysable CpG unit were expressed as output methylation ratios on Y axis, with FXS DNA input % expressed on the X axis (each point represents mean of duplicate PCRs from a single bisulfate converted DNA mixture). Methylation output ratios for CpG sites within FREE2B and FREE2C amplicons (A and B) were positively correlated with increasing FXS DNA input %; while FREE3 Methylation output ratios were negatively correlated with increasing FXS DNA input %.

[0185] FIG. 4A is a diagrammatic representation of the intron and exon regions at the Xq27.3 locus (sequence numbering from GenBank L29074 L38501), locations of FMR1 and ASFMR1 transcription start sites and alternative splicing events. The locations of target sequences for FMR1 and ASFMR1 real-time PCR assays used are also indicated: ASFMR1 (−1) real-time assay: detects unspliced and splice variant C (positioned −282 to −343 from FMR1 transcription start site), ASFMR1 (−2) real-time assay: detects unspliced only (positioned −588 to −663 from FMR1 transcription start site), ASFMR1 real-time assay: detects all (positioned −1299 to −1360 from FMR1 transcription start site).

[0186] FIG. 4B is a graphical representation of standard curve and amplification real-time PCR plots showing that in the FXS cell lines with fully methylated FMR1 promoter and silenced FMR1 and FMRP, ASFMR1 is expressed. RNA was extracted from 3 FXS cell lines whose methylation profiles are presented in FIG. 2; Sample 849 was taken from the male 490 CGG repeat line; Sample 862 was taken from the male 530 CGG repeat line; Sample 865 was taken from the female 563 and 47 CGG repeat line. Each RNA sample was split in

two, with one half subjected to RNase A treatment prior to ASFMR1 (−3) relative standard curve analysis. The ASFMR1 (−3) real-time PCR analysis was performed in quadruplicate reactions. The difference in Ct values between RNase A treated and untreated samples represents the level of ASFMR1 expression.

[0187] FIGS. 4C and D are graphical representations of standard curve and amplification real-time PCR plots indicating that in the FXS cell lines, ASFMR1RNA forms RNA:DNA complexes. FXS RNA samples were treated with TURBO DNase (C) and RQ1 DNase (D) respectively. These DNase treatments caused complete loss of real-time-PCR signal for the ASFMR1(−3) assay. Because DNase can only degrade RNA molecules if they form complexes with DNA, loss of ASFMR1 after DNase treatment suggests that ASFMR1RNA forms RNA:DNA complexes in FXS samples with fully methylated FMR1 promoter and silenced FMR1 expression.

[0188] FIGS. 5A and B are graphical representations of different FMR1 and ASFMR1 transcripts in RNA samples from lymphoblast lines of 6 male controls, two FXS males (samples 849 and 862) and one FXS female (865). The control and FXS RNA samples were either treated with TURBO DNase (A), RQ1 DNase (B), RNase A (C), or were untreated. Addition of TURBO DNase or RQ1 DNase buffers to RNA samples without DNase were included as additional controls in A and B. The FMR1 and ASFMR1 transcripts were quantified using real-time RT-PCR relative standard curve method, normalized to mRNA levels of three internal control genes, GUS, GAPDH and B2M. FMR15' and 3' assays showed no signal for the FXS RNA samples, while similar levels were detected in all control samples (upper two panels in A, B and C). TURBO and RQ1 DNase treatment caused ~50% decrease in the FMR1 levels in most of the control samples; while RNase A treatment caused complete loss of FMR1 and ASFMR1 signals. While decrease of ASFMR1 (−1)(−2) and (−3) levels was also observed in all control samples caused by TURBO and RQ1 DNase treatment, in FXS samples (with analogous to control ASFMR1 levels in the untreated samples) TURBO and RQ1 DNase treatment resulted in complete loss of signal for all three ASFMR1 assays. Because DNase can only degrade RNA molecules if they form complexes with DNA, this suggests that ASFMR1 RNA forms RNA:DNA complexes more readily in FXS samples than in controls. Increase in RNA:DNA interaction of ASFMR1 in FXS may lead to methylated FMR1 promoter and adjacent regions (FIG. 1) and silencing FMR1 expression leading to loss of FMRP and the resulting FXS clinical phenotype.

[0189] FIG. 6(A) is a representation of the intron and exon regions 5' of the FMR1 CGG expansion (sequence numbering from GenBank L29074 L38501) in relation to FMR1 and ASFMR1 transcription start sites, FMR1 promoter, the Fragile X-related epigenetic elements 2 and 3 (FREE). A CGG repeat is located within the 5' (UTR) of the FMR1 gene. ASFMR1 spans the CGG expansion in the antisense direction and is also regulated by another promoter located in the exon 2 of FMR1. The FREE2 located downstream of the CGG expansion. The FREE3 region is located within intron 2 of FMR1 and spans the potential second ASFMR1 promoter of transcription in the antisense direction. (B) Primers utilized for MALDI-TOF methylation analysis targeted 5 regions at the Xq27.3 locus designated as FREE2(A) (described as amplicon 5 in Godler et al., *Hum Mol Genet*, 2010; [Epub

ahead of print] doi:10.1093/hmg/ddq 1037); FREE2(D); FREE2(E), and FREE3 (color coded). Individual CPG sites within each region are numbered accordingly. Prominent transcription factor binding sites and methylation sensitive restriction enzyme recognition sites are indicated in capital font, and are listed/identified in Tables 3 and 4. << Indicates ASFMR1 transcription start site. The red arrow indicates the FREE2 3' Boundary located at CpG1 of FREE2(E) which is underlined in the sequence. (C) is a representation of the CG dinucleotide density in the regions proximal to the FREE2 3' Boundary. The CpG sites that have been analysed for methylation status are represented in Blue on the X axis, while sites not covered by our assays are represented in red. The coordinates of the FREE2(D) and FREE2(E) assays on the GenBank L29074 L38501 are also indicated.

[0190] FIG. 7 is a graphical representation of the methylation pattern variation in lymphoblasts and blood between (A) healthy controls, (B) unmethylated FM 'high functioning' males with full scale IQ (FIQ) greater than 70, (C) FXS affected individuals full scale IQ (FIQ) less than 70. Methylation output ratio (Y axis) of individual CpG units (X axis), were analyzed within the 27 kb region 3' CGG expansion, inclusive of intron 1, exon 2 and intron 2 (sequence numbering from GenBank L29074 L38501). The SEQUENOM mass spectrometry assays FREE2(D) and FREE2(E) (see Tables 3 and 4) were the most proximal to the FREE2 3' Boundary highlighted with the RED arrow as the CpG 1 of the FREE2 (E) assay. It is evident from the representation that this CpG unit is the last one within the FREE2 region which is unmethylated in healthy controls and high functioning FM individuals while being hypermethylated in FXS affected individuals. It is evident from the representation that the FREE2 region methylation status is an important biomarker of the FXS phenotype as it is hypermethylated in high functioning FM individuals as well as in healthy controls, while being hypomethylated in FXS affected individuals. It is also evident from the representation that the methylation patterns are consistent between lymphoblasts and blood. LB-represents lymphoblasts.

DETAILED DESCRIPTION

[0191] Taught herein is to a method for identifying an epigenetic profile of an intron, intron/exon boundary and/or splicing region within a genetic locus associated with or indicative, instructive or informative of a pathological condition. The epigenetic modification occurs in:

[0192] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or

[0193] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

within a genetic locus. In an embodiment, the pathological condition is a trinucleotide expansion disorder such as in association with a change in epigenetic profile from that of a healthy control subject.

[0194] By "epigenetic profile" includes epigenetic modifications such as methylation including hypermethylation and hypomethylation, RNA/DNA interactions, expression profiles of non-coding RNA, histone modification, changes in acetylation, ubiquitylation, phosphorylation and sumoylation, as well as chromatin altered transcription factor levels and the like leading to activation or deactivation of genetic locus expression. Particularly, the extent of methylation, RNA/DNA interaction and non-coding RNA expression are

determined as well as any changes therein. In an aspect, the epigenetic modification is an elevation in methylation, a decrease in methylation or an alteration in distribution of methylation sites. The epigenetic profile may be determined on either strand of genomic double stranded DNA or an amplified fragment thereof. Hence, primers may be generated to amplify either strand of a genomic DNA target.

[0195] The pathological condition may be a neurological or non-neurological condition. Insofar as the condition is neurological, it may be described as a neuropathological condition or a pathoneurological condition which encompasses neurodegenerative and neurodevelopmental disorders. Non-neurological pathologies are also contemplated herein as well as any nucleotide expansion disease or condition. Reference to a pathological conditions includes a trinucleotide expansion disorder. Reference to a “control” means relative to a healthy subject which means a subject with a normal size of expansion repeats and/or who is phenotypically normal meaning that the subject does not have symptoms of, for example, a trinucleotide expansion disorder and/or is the epigenetic profile observed in healthy control subjects.

[0196] In an embodiment, the pathological disease or condition including trinucleotides expansion disorders associated with a change in the epigenetic profile from that observed in healthy controls associated with the intronic epigenetic change is Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidol-oluyisiantrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCAT), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCAT), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome. The present disclosure also identifies nucleotide expansion diseases and conditions. In an embodiment, the genetic locus is the FMR genetic locus and the pathology is FXS or related condition such as FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome or cognitive impairment.

[0197] A method is enabled for identifying an epigenetic profile in a genome of a cell indicative of a pathological condition selected from Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidol-oluyisiantrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCA7), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome, the method comprising screening for a change relative to the control in the extent of epigenetic modification

in a genetic locus within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of a genetic locus associated with the pathological condition wherein the extent of epigenetic change is indicative of the presence or severity of the pathological condition or a propensity to develop same. In an embodiment, the genetic locus is the FMR locus and the region assayed for epigenetic change is selected from:

[0198] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0199] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0200] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0201] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0202] (v) the FREE3 region; and

[0203] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0204] In an embodiment, a method is provided for identifying an epigenetic profile in a genome of a cell indicative of FXS or related condition the method comprising screening for a change relative to the control in the extent of epigenetic modification in the FMR genetic locus within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of a genetic locus associated with the pathological condition wherein the extent of epigenetic change is indicative of the presence or severity of the pathological condition or a propensity to develop same. A related condition includes FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment.

[0205] In relation to this aspect, an “intron”, “intron/exon boundary” and “splicing region”, are regarded as an intron, intron/exon boundary and splicing region within a genetic locus or a gene within a genome. The intron, intron/exon boundary and splicing region may also encode a regulatory RNA species. Either strand of a double stranded genomic DNA or an amplified fragment or region thereof may be assayed for its epigenetic profile.

[0206] In an embodiment, the pathological condition is associated with an epigenetic profile of the FMR genetic locus. For the purposes of the present disclosure, the “FMR genetic locus” includes the FMR1, FMR4 and ASFMR1 genes as well as promoter and regulatory regions and introns and exons and intron/exon boundaries. In particular, the FMR genetic locus comprises a promoter region, a (CGG)_n region proximal to the promoter and exonic and intronic regions of the FMR1, FMR4 and ASFMR1 genes as depicted in FIGS. 1A and 4A and 6A. The promoter is generally referred to as the “FMR1 promoter” or “ASFMR1 promoter” for the promoter with an initiation site in FREE3 of intron 2 (see FIG. 6). The FMR locus includes introns, intron/exon boundaries and splicing regions wherein it is proposed herein that epigenetic changes occur within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions; and/or (ii) approximately one seventh or greater of an intron includ-

ing an intron/exon boundary and/or a splicing region of the FMR1 genetic locus including the FMR1 gene or a part thereof such as FREE3 or FREE2 region alone or in combination with the FREE1 region (D) or FREE2 (E) which are indicative or diagnostic of a pathological condition or its severity involving the FMR1, FMR4 and/or ASFMR1 genes. FREE3, FREE2 (D) and FREE2 (E) are further defined below. As indicated above, when determining an epigenetic profile either or both strands of the double stranded genomic DNA or an amplified product therefrom may be assayed. This also applies to a promoter region or other regulatory region.

[0207] In an embodiment, the epigenetic profile is determined within a genetic locus which enables the determination of an epigenetic profile in a genome of a cell indicative of a pathological condition and in particular a trinucleotide expansion disorder selected from Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidoluysianatrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCAT), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCA7), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome. Epigenetic changes in intronic, intronic/exonic boundaries and promoter or other regulatory regions in certain genetic loci are instructive to the development of these conditions.

[0208] In an embodiment, the genetic locus is the FMR1 genetic locus and the method comprises screening for a change relative to the control in the extent of epigenetic modification within the FMR1 genetic locus within a region selected from:

[0209] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0210] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0211] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0212] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0213] (v) the FREE3 region; and

[0214] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0215] The extent of epigenetic change is indicative of the presence or severity of the pathological condition or a propensity to developing same. Disease conditions contemplated herein associated with the FMR1 genetic locus include the trinucleotide expansion disorder FXS and related conditions such as FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment. In an embodiment, the present disclosure teaches a method for identifying FXS or a related condition in a human subject, the method

comprising screening for a change relative to a control in the extent of epigenetic modification in the FMR1 genetic locus at a location selected from:

[0216] Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions; and

[0217] (ii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

wherein a change in extent of epigenetic modification relative to a control is indicative of the presence or severity of the pathological condition or a propensity to develop same.

[0218] In an embodiment a method is provided for identifying a trinucleotide expansion disorder in a mammalian subject including a human, the method comprising screening for a change relative to a healthy control in the extent of epigenetic modification within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR1 genetic locus; wherein a change in extent of epigenetic modification relative to the control is indicative of the presence or severity of the trinucleotide expansion disorder or a propensity to develop same wherein the intron, intron/exon boundary and/or splicing region is selected from the list consisting of:

[0219] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0220] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions; and

[0221] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions.

[0222] The present disclosure teaches the manufacture of an assay to identify an epigenetic profile of an FMR1 genetic locus-associated pathological condition. Reference to an FMR1 genetic locus-associated pathological condition includes a trinucleotide expansion disorder associated with a change in epigenetic profile from that observed in a healthy subject.

[0223] The FMR1 genetic locus is depicted in part in FIGS. 1A, 4A and 6A. Reference to the "FMR1 genetic locus" includes the FMR1, FMR4 and ASFMR1 genes and corre-

sponds to Xq27.3. The term “FMR locus” means the “FMR genetic locus”. In an embodiment, an aspect taught herein determines that the intronic region downstream of intron 1 comprises Fragile X-related Epigenetic Element 3 as defined by SEQ ID NO:1 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions; or is intron 2 as defined by SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions. The nucleotide sequence of intron 1 of the FMR1 gene is set forth in SEQ ID NO:3. The nucleotide sequence of genomic FREE2 (D), FREE2 (E) and FREE3 are set forth in SEQ ID NOs:48, 49 and 47, respectively and the present disclosure extends to homology thereof having at least 80% identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or a complement thereof under medium stringency conditions. The present disclosure extends to both strands and hence, for example, reference to a particular SEQ ID NO: includes the corresponding complementary sequence.

[0224] The present disclosure further contemplates amplifying all or part of an expansion mutation and/or and detecting extent of epigenetic change therein in combination with an epigenetic change (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus including the FMR1 gene. The extent of epigenetic change in two or more of an intron, an intron/exon boundary and/or a splicing region or in one seventh or greater of an intron within the FMR genetic locus may be determined alone or in combination with extent of (CGG)_n expansion and/or any other epigenetic change therein. The determination of epigenetic change may also be conducted in combination with an assay as contemplated by International Patent Application No. PCT/AU2010/000169 filed on 17 Feb. 2010, the contents of which are incorporated herein by reference in their entirety. In an embodiment, the epigenetic modification is a change in extent of methylation which includes hypermethylation and hypomethylation and profile of methylation. Without limiting the present disclosure to any one theory or mode of action, epigenetic changes in these introns may affect the ability of the introns, intron/exon boundaries and/or splicing regions to transcribe regulatory RNAs which in turn have an effect on bidirectional transcription capability.

[0225] Hence, an aspect taught herein is a method for identifying a pathological condition in a mammalian subject including a human, the method comprising screening for a change relative to a control in the extent of epigenetic modification within a region selected from:

[0226] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0227] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which

hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0228] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0229] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0230] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0231] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein a change in extent of genetic modification relative to a control is indicative of the presence or severity of the pathological condition or a propensity to develop same.

[0232] The present disclosure teaches a method for identifying in a genome of a mammalian cell including a human cell, a pathological condition associated with methylation and other epigenetic change within the FMR locus, the method comprising extracting genomic DNA from the cell and subjecting the DNA to an amplification reaction using primers selective of a region of the FMR genetic locus comprising CpG and/or CpNpG sites, the CpG and CpNpG sites located in a region selected from:

[0233] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0234] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0235] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0236] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0237] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0238] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

and subjecting the amplified DNA to a methylation or other epigenetic assay to determine the extent of epigenetic modification of the DNA wherein a change in extent of epigenetic modification relative to a control is indicative of the presence

or severity of the pathological condition or propensity to develop same. In an embodiment, the above assay is useful in detecting FXS or a related condition. Examples of related conditions include FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment.

[0239] In an embodiment, the epigenetic modification is methylation of CpG and/or CpNpG sites and the assay identifies the extent of methylation change in either strand of double stranded genomic DNA or an amplified fragment thereof including either strand of a promoter region. This change may be an elevation or increase in methylation or a decrease in methylation relative to a control. Alternatively, the epigenetic modification is extent of change in RNA/DNA interaction and/or change in profile of expression of expression of non-coding RNA. Yet in another embodiment, the epigenetic profile is a change in histone modification, changes in acetylation, ubiquitylation, phosphorylation, sumoylation, activation or deactivation, chromatin altered transcription factor levels and the like.

[0240] In accordance with the present disclosure, a method is provided wherein the extent of methylation or other epigenetic modification provides a quantitative or semi-quantitative or qualitative indication of extent of change in epigenetic profile in (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus including the FMR1 gene, ASFMR1 gene and promoter gene (including an FMR1 promoter and an ASFMR1 promoter) and as such the level of epigenetic modification defines the severity of the pathological condition alone or in combination with the extent of (CGG)_n expansion. The number of repeats indicate whether a subject is a healthy control or has a Gray Zone (GZ) pathology, premutation (PM) pathology or full mutation (FM) pathology. The method disclosed herein may also be used in conjunction with other assays such as Southern blot or PCR to measure (CGG)_n expansion. Examples of pathology conditions associated with inronic epigenetic changes include the polyQ and non-polyQ conditions listed above.

[0241] The present disclosure, however, is not limited to the FMR genetic locus and pathological conditions only associated therewith. Rather, the present disclosure extends to any epigenetic modification in any genetic locus selected from (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh greater of an intron including an intron/exon boundary and/or splicing region and which epigenetic change is associated with a pathological condition.

[0242] By “approximately one seventh or greater” means from about 14% or greater or nucleotides capable of epigenetic change or modification have undergone a change. This includes 14.5, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 67, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and 100% of the nucleotides.

[0243] The regions within the FMR genetic locus identified above include a region selected from:

[0244] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0245] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0246] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0247] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0248] (v) the FREE3 region; and

[0249] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0250] As taught herein a “pathological condition” or “disease condition” includes an abnormal condition including a neurodevelopmental condition or a neurodegenerative condition or a non-neurological condition as defined by objective or subjective manifestations of disease. In an embodiment, it is a trinucleotide expansion disorder. A particular condition is FXS or a related condition such as FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment. The assay herein described is useful for diagnosing nucleotide expansion diseases or conditions. The assay enabled herein includes a genetic determination to be made to complement other symptom-based diagnoses such as based on behavioral studies or may be made in its own right. The assay may be part of a suit of diagnostic or prognostic genetic assays of embryos, pre- and post-natal subjects. The terms “method”, “assay”, “system”, “test”, “determination”, “prognostic”, “diagnostic”, “report” and the like may all be used to describe the methylation assay of selected regions of the FMR genetic locus or other genetic locus. The epigenetic assay such as a methylation assay determines the epigenetic profile or extent of epigenetic change compared to a control which suggests or indicates or is instructive of a disease or condition associated with epigenetic modification of an intron within a genetic locus. The present assay is also useful in population studies such as epidemiological studies including studies of ethnic populations.

[0251] Accordingly, the present disclosure further provides a method of identifying a methylation or other epigenetic profile in populations of subjects indicative of a pathological condition, the method comprising screening for a change relative to a control in a statistically significant number of subjects the extent of epigenetic modification in (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within a genetic locus wherein a change in extent of epigenetic modification is indicative of the presence or severity of the pathological condition or a propensity to develop same.

[0252] In an embodiment, the genetic locus is the FMR genetic locus and the region screened for epigenetic change is selected from:

[0253] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0254] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0255] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0256] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0257] (v) the FREE3 region; and

[0258] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0259] The present disclosure teaches a method of identifying a methylation or other epigenetic profile in a population of subjects indicative of a pathological condition associated with the FMR locus, the method comprising screening for a change, relative to a control, in a statistically significant number of subjects in the extent of epigenetic modification including extent of change in methylation of CpG and/or CpNpG sites within a region selected from:

[0260] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0261] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0262] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0263] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0264] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0265] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein a change in extent of epigenetic modification is indicative of the presence of the pathological condition or a propensity to develop same in the population. In an embodiment, the pathological condition is FXS or a related condition such as FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment.

[0266] In accordance with this method, a further step may be conducted of determining the extent of (CGG)_n expansion such as by PCR and/or Southern blot analysis of bisulfite converted and/or non converted DNA. Furthermore, this assay may be conducted with one or more assays contemplated and described in International Patent Application NO. PCT/AU2010/000169 filed on 17 Feb. 2010, the contents of which are incorporated by reference in their entirety.

[0267] In an embodiment, the extent of methylation or change in extent of methylation is detected and associated with the pathology condition such as but not limited to an expansion disease or condition.

[0268] An epigenetic map and in particular a methylation map of introns, intron/exon boundaries and/or splicing

regions within the FMR locus has thus been constructed in accordance with the present disclosure using standard techniques such as high throughput mass spectrometry in the genome of various cells. Any cell type cell may be assayed. These cells include cultured or uncultured Chorionic Villi Sample (CVS) cells, lymphoblasts, blood cells, buccal cells, epithelial cells, fibroblast cells, an amniocyte and EBV transformed lymphoblast cell lines from male and female subjects with either no symptoms or from a spectrum of a pathological condition such as Fragile X mental retardation symptoms. In an embodiment a Fragile X-related Epigenetic Element 3 (FREE3) has been identified within intron 2 of the FMR1 gene. It is proposed that this region [FREE3] or other regions of intron 2 or other introns or parts thereof including intron/exon boundaries and splicing regions downstream of intron 2 of FMR1 or elsewhere in the FMR genetic locus are responsible for the regulation of transcription of FMR4 and ASFMR1 and FMR1 and expression of FMRP. Another region is an ASFMR1 promoter having an initiation site in FREE3 of intron 2 (see FIG. 6A).

[0269] In an embodiment, the present disclosure teaches that the extent of methylation in CpG and/or CpNpG sites located within the region downstream of intron 1 or part thereof such as FREE3 closely corresponds to a healthy condition or a level or severity of disease within the spectrum of PM to FM including GZ subjects such a correspondence may be in a further association with other epigenetic modifications within the FMR genetic locus and/or CGG expansion. Furthermore, using the methylation assay, methylation levels of the FREE3 region provide fully quantitative results, which also reflect the degree of X-chromosome modification in females. This can be more informative than methylation patterns of FMR1 CpG islands only which may be biased due to its proximity to a nucleotide expansion, and hence can only provide a qualitative assessment of methylation. Other regions of interest include FREE2 (A), (B), (C) and (D) and the FREE2 (D)/FREE2 (E) boundary defined herein.

[0270] Hence, in an embodiment, the present disclosure contemplates a change in extent of methylation which includes an increase or decrease in extent of methylation. There may also be no change in the extent of methylation within an intron of a genetic locus. However, the present disclosure extends to the detection of the change in extent of any epigenetic modification. Such a change or level of methylation in an intron is proposed to be associated with a pathological condition or its severity. In this context, an "intron" includes an intron/exon boundary and/or a splicing region.

[0271] A "normal" or "control" in the assay of the present disclosure may be a control genome from a healthy individual performed at the same time or the epigenetic pattern may be compared to a statistically validated standard. In relation to a nucleotide expansion disease condition, a healthy individual includes a subject with a nucleotide repeat within the normal range with no clinically apparent pathological phenotype. For example, in relation to (CGG)_n expansion conditions within the FMR genetic locus, this includes when n is <40.

[0272] The present disclosure also explores the relationship between transcription and epigenetic profile of introns or parts thereof and pathological conditions. A "part" includes an intron/exon boundary and splicing region. In an embodiment, methylated CpG sites are identified within FREE3 or intron 2 of the FMR1 gene in subjects with Fragile X mental retardation conditions. In another embodiment, the methylated CpG sites are identified in (i) two or more of (a) an

intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus. Fragile X mental retardation conditions include FXS as well as FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment.

[0273] As used herein, the terms “subject”, “patient”, “individual”, “target” and the like refer to any organism or cell of the organism on which an assay of the present disclosure is performed whether for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include both male and female humans but the present disclosure extends to experimental animals such as non-human primates, (e.g., mammals, mice, rats, rabbits, pigs and guinea pigs/hamsters). The “subject” may also be referred to as a population since the present disclosure teaches an assay useful in populations studies including epidemiological studies or assays of ethnic population. In a particular embodiment, the subject is a human. The test may be tailored to human females or human males or pre-natal humans. A control subject has epigenetic (e.g. methylation) profile of a healthy subject.

[0274] The terms “Fragile X mental retardation-like condition” and FMR condition” refer to a neurological disease, disorder and/or condition characterized by one or more of the following symptoms: (1) behavioral symptoms, including but not limited to hyperactivity, stereotypy, anxiety, seizure, impaired social behavior, and/or cognitive delay; (2) defective synaptic morphology, such as an abnormal number, length, and/or width of dendritic spines; and/or (3) defective synaptic function, such as enhanced long-term depression (LTD); and/or reduced long-term potentiation (LTP); and/or impaired cognitive ability. The pathological condition is a disease, disorder, and/or condition caused by and/or associated with epigenetic changes within an intron or part thereof within the FMR genetic locus such as downstream of intron 1 of the FMR1 gene. Such epigenetic changes may be alone or in combination with one or more of the following: (1) a mutation in FMR1 or FMR4 or ASFMR1; (2) defective FMR1/FMR4/ASFMR1 expression; (3) increased and/or decreased levels of FMRP; (4) defective FMRP function; (5) increased and/or decreased expression of genes or genetic functions regulated by FMR1, FMRP, FMR4 transcript or ASFMR1 transcript; (6) the increased methylation of FMR locus at CpG or CpNpG sites in the region upstream of FMR1 promoter and/or the region downstream of the (CGG)_n portion of the FMR1 promoter but not including the (CGG)_n portion; (7) an increased and/or decreased function of the FMR locus via miRNAs and/or members of the miRNA pathway; (8) an increased and/or decreased ability of FMRP to interact with its known target RNAs, such as RNAs encoding Rac1, microtubule-associated protein 1B, activity-regulated cytoskeleton-associated protein, and/or alpha-calcium/calmodulin-dependent protein kinase II; (9) symptoms of FXS, FXTAS, FXPOI, mental retardation, a modified X-chromosome, autism and/or autism spectrum disorders; and/or (10) cognitive impairment. Generally, the FMR condition is a trinucleotide expansion disorder, particularly associated with a changed epigenetic profile from that of a healthy subject.

[0275] Those of ordinary skill in the art will appreciate that the teachings of the present disclosure are applicable to any neurodevelopmental or neurodegenerative disorders linked, associated or otherwise influenced by the function of the

FMR genetic locus or genes therein such as FMR1, FMR4 and ASFMR1 as well as their promoters or other regulatory regions. Non-neurological disorders are also contemplated herein including FXPOI and other tri-nucleotide expansion disorders.

[0276] Furthermore, the present disclosure teaches a range of nucleotide expansion disorders. Conditions and disorders contemplated herein include diseases such as Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington’s disease (HD), dentatorubropallidol-luysianatrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCAT), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich’s ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCAB) and spinocerebella ataxias Type 12 (SCA12), Klinefelter’s syndrome and Turner’s syndrome.

[0277] The term “genomic DNA” includes all DNA in a cell, group of cells, or in an organelle of a cell and includes exogenous DNA such a transgenes introduced into a cell. Either strand or both strands of double stranded DNA may be assayed.

[0278] In a particular aspect, the present disclosure enables the determination of the presence of an FMR genetic locus-associated pathology based on extent of methylation of CpG/CpNpG sites located within (i) an intron downstream of intron 1 of the FMR1 gene or part of an intron; (ii) two or more or (a) an intron; (b) an intron/exon boundary; (c) a splicing region; and/or (iii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus. The downstream FMR1 introns may extend beyond the FMR1 gene. In an embodiment, the extent of methylation in part of intron 2 [FREE3] is identified in the FMR1 gene which includes all or part of an ASFMR1 promoter with a transcription start site in FREE3 of intron 2 (see FIG. 6A).

[0279] Hence, the present disclosure teaches a method for identifying a methylation or other epigenetic profile in the genome of a cell indicative of a pathological condition associated with the FMR genetic locus, the method comprising screening for a change relative to the control in the extent of epigenetic modification of CpG and/or CpNpG sites located within:

[0280] (i) (a) FREE3; (b) intron 2; and (c) an intron downstream of intron 2 or a homolog thereof or a portion or fragment thereof within the FMR1 gene;

[0281] (ii) two or more of (a) an intron; (b) an intron/exon boundary; (c) a splicing region within the FMR genetic locus; and/or

[0282] (iii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein a change in the extent of epigenetic modification is indicative of the presence of the pathological condition or a propensity to develop same. The nucleotide sequences of FREE3 and intron 2 are set forth in SEQ ID NOs: 1 and 2, respectively and the present disclosure extends to their homologs and portions and parts thereof having at least 80%

identity thereto or a nucleotide sequence capable of hybridizing to these sequences or their complementary forms under medium stringency conditions. Reference to FREE3 and an intron such as intron 2 includes portions, fragments, parts, regions and domains thereof. The nucleotide genomic sequences of genome FREE2 (D), FREE2 (E) and FREE3 are set forth in SEQ ID NOs:48, 49 and 47, respectively and the present disclosure extends to their homologs and portions and parts thereof having at least 80% identity thereto or a nucleotide sequence capable of hybridizing to these sequences or their complementary forms under medium stringency conditions. Reference to FREE3 and an intron such as intron 2 includes portions, fragments, parts, regions and domains thereof as well as one or both strands of double stranded genomic DNA.

[0283] In a particular embodiment, the epigenetic modification is methylation and RNA/DNA interactions.

[0284] The present disclosure further contemplates a method for identifying a pathological condition in a subject associated with methylation within the FMR locus, the method comprising extracting genomic DNA from a cell of the subject and subjecting the DNA to an amplification reaction using primers selective of a region of the FMR genetic locus comprising CpG and/or CpNpG sites, the CpG and CpNpG sites located in the FMR genetic locus within a region selected from:

[0285] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0286] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0287] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0288] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0289] (v) the FREE3 region; and

[0290] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

and subjecting the DNA to a methylation assay to determine the extent of methylation of the DNA wherein a change in extent of methylation relative to a control is indicative of the presence or severity of the pathological condition or propensity to develop same.

[0291] In an embodiment, the present disclosure teaches a method for identifying FXS or a related condition associated with methylation of the FMR genetic locus, the method comprising extracting genomic DNA from a cell of the subject and subjecting the DNA to an amplification reaction using primers selective of a region of the FMR genetic locus comprising CpG and/or CpNpG sites, the CpG and CpNpG sites located in the FMR genetic locus within a region selected from:

[0292] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0293] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0294] (iii) the Y boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0295] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0296] (v) the FREE3 region; and

[0297] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

and subjecting the DNA to a methylation assay to determine the extent of methylation of the DNA wherein a change in extent of methylation relative to a control is indicative of the presence or severity of the pathological condition or propensity to develop same.

[0298] Any methylation assay may be employed such as methylation sensitive PCR, methylation specific melting curve analysis (MS-MCA) or high resolution melting (MS-HRM) [Dahl et al., *Clin Chem* 53(4):790-793, 2007; Wojdacz et al., *Nucleic Acids Res.* 35(6):e41, 2007]; quantification of CpG methylation by MALDI-TOF MS (Tost et al., *Nucleic Acids Res* 31(9):e50, 2003); methylation specific MLPA (Nygren et al., *Nucleic Acids Res.* 33(14):e128, 2005); methylated-DNA precipitation and methylation-sensitive restriction enzymes (COMPARE-MS) [Yegnashubramanian et al., *Nucleic Acids Res.* 34(3):e19, 2006] or methylation sensitive oligonucleotide microarray (Gitan et al., *Genome Res.* 12(1):158-164, 2002), as well as via antibodies. Other assays include NEXT generation (GEN) and DEEP sequencing or pyrosequencing. Another assay is single molecule (SMRT) sequencing.

[0299] Insofar as the methylation assay may involve an amplification, an amplification methodology may be employed. Amplification methodologies contemplated herein include the polymerase chain reaction (PCR) such as disclosed in U.S. Pat. Nos. 4,683,202 and 4,683,195; the ligase chain reaction (LCR) such as disclosed in European Patent Application No. EP-A-320 308 and gap filling LCR (GLCR) or variations thereof such as disclosed in International Patent Publication No. WO 90/01069, European Patent Application EP-A-439 182, British Patent No. GB 2,225,112A and International Patent Publication No. WO 93/00447. Other amplification techniques include Q β replicase such as described in the literature; Strand Displacement Amplification (SDA) such as described in European Patent Application Nos. EP-A-497 272 and EP-A-500 224; Self-Sustained Sequence Replication (3SR) such as described in Fahy et al., *PCR Methods Appl.* 1(1) 25-33, 1991 and Nucleic Acid Sequence-Based Amplification (NASBA) such as described in the literature.

[0300] A PCR amplification process is particularly useful in the practice of the present disclosure.

[0301] In an embodiment, prior to the PCR, either essentially all cytosines in the DNA sample are selectively delaminated, but 5-methylcytosines remain essentially unchanged or essentially all 5-methylcytosines in the DNA sample are selectively delaminated, but cytosines remain essentially unchanged. Cytosine-guanine (CpG) dinucleotides and CpNpG trinucleotides are detected, allowing conclusions about the methylation state of cytosines in the CpG dinucleotides and CpNpG trinucleotide in the DNA sample. This delamination is generally performed using a bisulfite reagent. After bisulfite treatment, the 5-methylcytosines residues are converted to thymine (T).

[0302] The sample DNA is only amplified by chosen PCR primers if a certain methylation state is present at a specific site in the sample DNA the sequence context of which is essentially complementary to one or more of the chosen PCR primers. This can be done using primers annealing selectively

to bisulfite treated DNA which contains in a certain position either a TG or a CG or CNG, depending on the methylation status in the genomic DNA. Primers are designed based on particular regions around CpG and/or CpNpG sites or other FMR1 intronic regions. Introns or parts thereof including intron/exon boundaries and splicing regions downstream of intron 2 of FMR1 or downstream of the FMR1 gene itself are also contemplated herein as are (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region, of the FMR genetic locus.

[0303] A technology which can alternatively be employed for methylation analysis utilizes base-specific cleavage followed by MALDI-TOF mass spectrometry on DNA after bisulfite treatment, where all the 5-methylcytosines residues are converted to thymine (T) or where all unmethylated cytosines residues are not converted to thymine (T). Primers are designed based on particular regions around CpG and/or CpNpG sites or other FMR1 intronic regions or downstream thereof. Primer sequences are designed to amplify without bias both converted and unconverted sequences using the PCR amplification process under the medium to high stringency conditions. The PCR products are in vitro transcribed and subjected to base specific cleavage and fragmentation analysis using MALDI-TOF MS. The size ratio of the cleaved products provides quantitative methylation estimates for CpG sites within a target region. The shift in mass for non-methylated (NM) from methylated (M) fragments for a single CpG site is -16 daltons due to the presence of an adenosine residue in the place of a guanosine. A software is then used to calculate methylation for each fragment based on this difference in mass, where the output methylation ratios are the intensities of methylated signal/[methylated+unmethylated signal]. If the fragment size overlaps for different CpGs, their methylation output ratio is calculated based on the sum of intensities for methylated/[methylated+unmethylated signal]. To distinguish how well the methylation output ratio for multiple fragments of a similar size represented methylation of separate CpG sites, for some amplicons both cytosine and thymidine cleave reactions can be performed (that produced fragments of different size) prior to fragment analysis. Silent peaks (S)—fragments of unknown origin, should not be taken into consideration if their size does not overlap with the fragments of interest. Methylation of CpG sites that have silent peaks (S) that overlap with the fragments of interest should be included in the analysis.

[0304] Hence, a method is provided for determining the methylation profile of one or more CpG or CpNpG sites located within the genome of a eukaryotic cell or group of cells, the method comprising obtaining a sample of genomic DNA from the cell or group of cells and subjecting the genomic DNA to primer-specific amplification within an intron of a genetic locus and assaying for extent of methylation relative to a control, including a change in the extent of methylation and associating this change with a pathological condition.

[0305] A “nucleic acid” as used herein, is a covalently linked sequence of nucleotides in which the 3' position of the phosphorylated pentose of one nucleotide is joined by a phosphodiester group to the 5' position of the pentose of the next nucleotide and in which the nucleotide residues are linked in specific sequence; i.e. a linear order of nucleotides. A “polynucleotide” as used herein, is a nucleic acid containing a

sequence that is greater than about 100 nucleotides in length. An “oligonucleotide” as used herein, is a short polynucleotide or a portion of a polynucleotide. An oligonucleotide typically contains a sequence of about two to about one hundred bases. The word “oligo” may be used in place of the word “oligonucleotide”. The term “oligo” also includes a particularly useful primer length in the practice of the present disclosure of up to about 10 nucleotides.

[0306] As used herein, the term “primer” refers to an oligonucleotide or polynucleotide that is capable of hybridizing to another nucleic acid of interest under particular stringency conditions. A primer may occur naturally as in a purified restriction digest or be produced synthetically, by recombinant means or by PCR amplification. The primer may be selected to amplify either or both strands of double stranded genomic DNA. This includes a promoter or other regulatory region. The terms “probe” and “primers” may be used interchangeably, although to the extent that an oligonucleotide is used in a PCR or other amplification reaction, the term is generally “primer”. The ability to hybridize is dependent in part on the degree of complementarity between the nucleotide sequence of the primer and complementary sequence on the target DNA.

[0307] The terms “complementary” or “complementarily” are used in reference to nucleic acids (i.e. a sequence of nucleotides) related by the well-known base-pairing rules that A pairs with T or U and C pairs with G. For example, the sequence 5'-A-G-T-3' is complementary to the sequence 3'-T-C-A-5' in DNA and 3'-U-C-A-5' in RNA. Complementarity can be “partial” in which only some of the nucleotide bases are matched according to the base pairing rules. On the other hand, there may be “complete” or “total” complementarity between the nucleic acid strands when all of the bases are matched according to base-pairing rules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands as known well in the art. This is of particular importance in detection methods that depend upon binding between nucleic acids, such as those of the present disclosure. The term “substantially complementary” is used to describe any primer that can hybridize to either or both strands of the target nucleic acid sequence under conditions of low stringency as described below or, preferably, in polymerase reaction buffer heated to 95° C. and then cooled to room temperature. As used herein, when the primer is referred to as partially or totally complementary to the target nucleic acid, that refers to the 3'-terminal region of the probe (i.e. within about 10 nucleotides of the 3'-terminal nucleotide position).

[0308] As discussed above, when reference is made to a genomic nucleotide sequence, the present disclosure extends to its complementary strand sequence. Either or both strands may be assayed from epigenetic change.

[0309] Reference herein to a stringency in relation to hybridization includes and encompasses from at least about 0 to at least about 15% v/v formamide and from at least about 1 M to at least about 2 M salt for hybridization, and at least about 1 M to at least about 2 M salt for washing conditions. Generally, low stringency is at from about 25-30° C. to about 42° C. The temperature may be altered and higher temperatures used to replace formamide and/or to give alternative stringency conditions. Alternative stringency conditions may be applied where necessary, such as medium stringency, which includes and encompasses from at least about 16% v/v

to at least about 30% v/v formamide and from at least about 0.5 M to at least about 0.9 M salt for hybridization, and at least about 0.5 M to at least about 0.9 M salt for washing conditions, or high stringency, which includes and encompasses from at least about 31% v/v to at least about 50% v/v formamide and from at least about 0.01 M to at least about 0.15 M salt for hybridization, and at least about 0.01 M to at least about 0.15 M salt for washing conditions. In general, washing is carried out $T_m = 69.3 + 0.41 (G+C) \%$ (Marmur and Doty, *J. Mol. Biol.* 5: 109, 1962). However, the T_m of a duplex DNA decreases by 1°C . with every increase of 1% in the number of mismatch base pairs (Bonner and Laskey, *Eur. J. Biochem.* 46: 83, 1974). Formamide is optional in these hybridization conditions. Accordingly, particularly preferred levels of stringency are defined as follows: low stringency is $6\times\text{SSC}$ buffer, 0.1% w/v SDS at $25\text{--}42^\circ \text{C}$.; a moderate stringency is $2\times\text{SSC}$ buffer, 0.1% w/v SDS at a temperature in the range 20°C . to 65°C .; high stringency is $0.1\times\text{SSC}$ buffer, 0.1% w/v SDS at a temperature of at least 65°C . Reference to at least "80% identity" includes 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and 100%.

[0310] The present disclosure enables the determination of a methylation or other epigenetic profile of sites within an intron, intron/exon boundary and/or splicing region of a genetic locus in a genome of a eukaryotic cell or group of cells, the method comprising obtaining a sample of genomic DNA from the cell or group of cells, subjecting the digested DNA to an amplification reaction using primers selected to amplify a region of the genetic locus selected from:

[0311] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) splicing region; and/or

[0312] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the genetic locus;

and then subjecting the amplified DNA to methylation or other epigenetic detection means to determine relative to control the extent of methylation or other epigenetic modification wherein a change in epigenetic modification or other epigenetic modification relative to the control is indicative of a pathological condition associated with the genetic locus.

[0313] Examples of pathological conditions include Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chromosome, Huntington's disease (HD), dentatorubropallidolucyosian trophy (DMA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCA7), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome.

[0314] In an embodiment, the present disclosure enables determination of a methylation profile of the sites within the FMR locus in a genome of a eukaryotic cell or group of cells, the methylation profile comprising the extent or level of methylation within the FMR locus, the method comprising obtaining a sample of genomic DNA from the cell or group of

cells, subjecting the digested DNA to an amplification reaction using primers selected to amplify a region of the FMR genetic locus selected from:

[0315] (i) (a) FREE3; (b) intron 2; and (c) an intron, intron/exon boundary and/or splicing region downstream of intron 2 or part thereof;

[0316] (ii) two or more of (a) an intron; (b) an intron/exon boundary; (c) a splicing region; and

[0317] (iii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

and then subjecting the amplified DNA to methylation detection means to determine relative to control the extent of methylation wherein a change in methylation relative to the control is indicative of a pathological condition associated with the FMR genetic locus.

[0318] In an embodiment, the region amplified within the FMR genetic locus is selected from:

[0319] (i) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0320] (ii) the FREE2 region alone or in combination with the FREE1 region;

[0321] (iii) the FREE3 region; and

[0322] (iv) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene including the FREE2 (D) region or FREE2 (D)/(E) boundary.

[0323] The present disclosure further provides a methylation profile of the sites within the FMR locus in a genome of a eukaryotic cell or group of cells, the methylation profile comprising the extent or level of methylation within the FMR locus, the method comprising obtaining a sample of genomic DNA from the cell or group of cells, subjecting the digested DNA to an amplification reaction using primers selected to amplify FREE3 within the FMR1 gene and then subjecting the amplified DNA to methylation detection means to determine relative to control the extent of methylation wherein a change in methylation relative to the control is indicative of a pathological condition associated with the FMR genetic locus.

[0324] In an embodiment, the present disclosure enables determination of a methylation profile of the sites within the FMR locus in a genome of a eukaryotic cell or group of cells, the methylation profile comprising the extent or level of methylation within the FMR locus, the method comprising obtaining a sample of genomic DNA from the cell or group of cells, subjecting the digested DNA to an amplification reaction using primers selected to amplify all or part of FREE3 and then subjecting the amplified DNA to methylation detection means to determine relative to control the extent of methylation wherein a change in methylation relative to the control is indicative of a pathological condition associated with the FMR genetic locus. An adverse change in methylation or other epigenetic profile is associated with FXS or related condition such as FXTAS, FXPOI, autism, mental retardation and cognitive impairment.

[0325] As indicated above, the cells may be a lymphoblast, a CVS cell, a blood cell, a buccal cell, epithelial cell, fibroblast cell, an amniocyte or an EBV transformed lymphoblast cell line. In addition, the methylation profile may be determined or one or both alleles a genetic locus and in selected cells where mosaicism has occurred. In particular, the extent of methylation can determine homozygosity or heterozygosity

or mosaicism. Reference to “mosaicism” includes the situation wherein two or more populations of cells have different genotypes or epigenetic profiles at the genetic locus.

[0326] The diagnostic assay herein can also detect heterozygosity or mosaicism where the methylation pattern is indicative of, for example, in relation to an FMR genetic locus-associated pathology, an FM. The latter may also be conducted in combination with an assay to detect (CGG)_n expansion.

[0327] The present disclosure also teaches kits for determining the methylation or other epigenetic profile of one or more nucleotides at one or more sites within the genome of a eukaryotic cell or group of cells. The kits may comprise many different forms but in one embodiment, the kits comprise reagents for the bisulfate methylation assay.

[0328] A further embodiment of the present disclosure is a kit for the use in the above methods comprising primers to amplify an intron within a genetic locus.

[0329] In an embodiment, the present disclosure provides a use of primers which amplify regions of the FMR genetic locus, comprising CpG and/or CpNpG sites located within:

[0330] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region;

[0331] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

[0332] (iii) the 3' boundary of the FREE2 region and the FMR1 promoter located between CpG unit 1 of the FREE2 (E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon;

[0333] (iv) the FREE2 region alone or in combination with the FREE1 region;

[0334] (v) the FREE3 region; and

[0335] (vi) an intron including an intron/exon boundary and/or splicing region downstream of intron 1 within the FMR1 gene.

in the manufacture of a diagnostic kit or device to detect methylation of the FMR locus-associated with a pathological condition.

[0336] In relation to one embodiment, a kit is provided for the use in the above methods comprising primers identified by SEQ ID NOs:6 through 11 to amplify an intronic site within the FMR1 genetic locus. The nucleotide sequences in SEQ ID NOs:6 through 11 comprise primer and tag sequences. The present disclosure extends to SEQ ID NO:6 through 11 as well as primer only portions therein. The primers may also include primers disclosed in PCT/AU2010/000169.

[0337] The kit may also comprise instructions for use.

[0338] Conveniently, the kits are adapted to contain compartments for two or more of the above-listed components. Furthermore, buffers, nucleotides and/or enzymes may be combined into a single compartment.

[0339] As stated above, instructions optionally present in such kits instruct the user on how to use the components of the kit to perform the various methods of the present disclosure. It is contemplated that these instructions include a description of the detection methods of the subject disclosure, including detection by gel electrophoresis.

[0340] The present disclosure further enables kits which contain a primer for a nucleic acid target of interest with the primer being complementary to a predetermined nucleic acid target. In another embodiment, the kit contains multiple primers or probes, each of which contains a different base at an interrogation position or which is designed to interrogate different target DNA sequences. In a contemplated embodi-

ment, multiple probes are provided for a set of nucleic acid target sequences that give rise to analytical results which are distinguishable for the various probes. The multiple probes may be in microarray format for ease of use.

[0341] The kit may comprise a vessel containing a purified and isolated enzyme whose activity is to release one or more nucleotides from the 3' terminus of a hybridized nucleic acid probe and a vessel containing pyrophosphate. In one embodiment, these items are combined in a single vessel. It is contemplated that the enzyme is either in solution or provided as a solid (e.g. as a lyophilized powder); the same is true for the pyrophosphate. Preferably, the enzyme is provided in solution. Some contemplated kits contain labeled nucleic acid probes. Other contemplated kits further comprise vessels containing labels and vessels containing reagents for attaching the labels. Microtiter trays are particularly useful and these may comprise from two to 100,000 wells or from about six to about 10,000 wells or from about six to about 1,000 wells.

[0342] Another important application is in the high throughput screening of agents which are capable of demethylation genomes and in particular intronic regions within genomes. This may be important, for example, in de-differentiating cells and/or treating pathological conditions.

[0343] The present disclosure further enables a method for screening for an agent which modulates methylation or other epigenetic modification of a genetic locus, the method comprising screening for a change relative to a control in the extent of methylation or other epigenetic modification in an intron, intron/exon boundary and/or splicing region within the genetic locus which is associated with a pathological condition in the presences or absence of an agent to be tested, wherein an agent is selected if it induces a change in the extent of methylation or other epigenetic change. Agents include de-methylation agents and hyper-methylation agents, global and site specific.

[0344] In an embodiment, a method is also provided for screening for an agent which modulates epigenetic modification of an FMR genetic locus in a mammalian cell including a human cell, the method comprising screening for a change relative to a healthy control in the extent of epigenetic change in (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus; wherein the intron, intron/exon boundary and/or splicing region is selected from the list consisting of:

[0345] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0346] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0347] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide

sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0348] in the presence or absence of an agent to be tested wherein the agent is selected if it induces a change in extent of epigenetic modification. Such an agent is useful in the treatment of a trinucleotide expansion disorder such as associated with a change in epigenetic profile from that of a healthy subject.

[0349] In an embodiment, a method is provided for screening for an agent which modulates methylation of an FMR genetic locus in a mammalian cell including a human cell, the method comprising screening for a change relative to a control in the extent of methylation in a region selected from:

[0350] (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0351] (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0352] (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0353] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0354] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and

[0355] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

in the presence or absence of an agent to be tested wherein the agent is selected if it induces a change in extent of methylation.

[0356] The present disclosure further enables a method for monitoring the treatment of a genetic locus-associated disease including a nucleotide expansion disease in which the treatment modulates the methylation of the genetic locus, the method comprising monitoring for a change relative to a control or a pre and post-treatment sample in the extent of methylation within an intron, intron/exon boundary and/or splicing region of the genetic locus.

[0357] By "monitoring" includes diagnosis, prognosis, pharmacoresponsiveness, pharmacosensitivity, level of disease progression or remission, improving or declining health of a subject and the like.

[0358] As indicated above, conditions and disorders contemplated herein include a range of nucleotide expansion diseases such as but not limited to Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, a modified X-chro-

mosome, Huntington's disease (HD), dentatorubropallidoluysianatrophy (DRPLA), spinobulbar muscular atrophy or Kennedy disease (SBMA), spinocerebella ataxia Type 1 (SCA1), spinocerebella ataxia Type 2 (SCA2), spinocerebella ataxia Type 3 or Machado-Joseph disease (SCA3), spinocerebella ataxia Type 6 (SCA6), spinocerebella ataxia Type 7 (SCAT), spinocerebella ataxia Type 17 (SCA17), Fragile XE mental retardation (FRAXE), Friedrich's ataxia (FRDA), Fragile type, folic acid type, rare 12 (FRA12A), myotonic dystrophy (DM), spinocerebella ataxia (SCA8) and spinocerebella ataxias Type 12 (SCA12), Klinefelter's syndrome and Turner's syndrome. Reference to a "modified" X-chromosome includes skewed X-inactivation, inversions, deletions, duplications, hybrids and any modification leading to X-chromosome inactivation. A particular condition associated with epigenetic changes to the FMR genetic locus include FXS and related disorders such as FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment.

[0359] The present disclosure further teaches the identification of genes having introns with CpG or CpNpG sites or other methylation-sensitive restriction sites. The identification of these sites permits identification of potential regulatory regions which can be targeted by agonists or antagonists of abnormal gene expression.

[0360] In cases where the gene is methylated and silenced in affected individuals or tissues, compounds are screened in high throughput fashion in stable cell lines or individuals to identify drugs that result in demethylation and reactivation of the affected gene. Alternatively, a normal active copy of the affected gene is transfected as a transgene into cells to correct the defect. Such transgenes are introduced with modulating sequences that protect the transgene from methylation and keep it unmethylated and transcriptionally active.

[0361] In cases where the gene is unmethylated and transcriptionally active or transcriptionally over-active in affected individuals or tissues, compounds are screened in high throughput fashion in stable cell lines to identify drugs that result in methylation and silencing of the affected gene. Alternatively, a transgene encoding a double stranded RNA homologous to the affected sequences or homologs thereof, are transfected as a transgene into cells to methylate the gene, silence it and thereby correct the defect. Such double stranded RNA-encoding transgenes are introduced with modulating sequences which protect it from methylation, keep it transcriptionally active and producing double stranded RNA.

[0362] The present disclosure further provides a computer program and hardware which monitors the changing state, if any, of extent of methylation over time or in response to therapeutic and/or behavioral modification. Such a computer program has important utility in monitoring disease progression, response to intervention and may guide modification of therapy or treatment. The computer program is also useful in understanding the association between increasing methylation and disease progression.

[0363] The computer program monitors in a quantitative or semi-quantitative manner one or more features including extent of methylation or other epigenetic modification in an intron of a genetic locus. In addition, the length of a nucleotide expansion may be determined or any epigenetic changes therein. In relation to a neuropathological condition, a behavioral assessment may be made using criteria associated with normal subjects or subjects considered to be suffering with a disease condition. For example, cognitive ability can be mea-

sured as well as the general phenotype or clinical manifestations in subjects with a neurodevelopmental or neurodegenerative condition or other condition associated with nucleotide expansion.

[0364] Thus, in accordance with the present disclosure, values are assigned to the listed features which are stored in a machine-readable storage medium, which is capable of processing the data to provide an extent of disease progression or change in methylation or other epigenetic modification for a subject.

[0365] In an aspect, the disclosure teaches a computer program product for assessing progression of a pathological condition associated with the FMR locus in a subject, the product comprising:

[0366] (1) assigning a value to one or more of:

[0367] (a) change in of methylation or other epigenetic modification relative to a control in FREE3 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0368] (b) change of methylation or other epigenetic modification relative to a control in intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0369] (c) change in methylation or other epigenetic modification relative to a control in genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0370] (d) change of methylation in an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0371] (e) two or more of (i) an intron; (ii) an intron/exon boundary; (iii) a splicing region within the FMR genetic locus;

[0372] (f) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

[0373] (g) length of (CGG)_n expansion within the FMR genetic locus when considered in combination with (a) and/or (b);

[0374] (h) general phenotype or clinical manifestations in subjects with a neurodevelopmental or neurodegenerative condition;

[0375] (i) behavioral assessment criteria associated with normal subjects, PM subjects, GZ subjects and FM subjects;

[0376] (j) cognitive ability;

[0377] (k) extent of transcription of genes within the FMR locus with the proviso that if any one of (d) through (k) is determined then one or more of (a) through (c) is also determined;

[0378] (2) means to converting the value to a code; and

[0379] (3) means to store the code in a computer readable medium and compare code to a knowledge database to determine whether the code corresponds to a pathological condition.

[0380] In a related aspect, the disclosure teaches a computer for assessing an association between extent of methylation or other epigenetic modification within the FMR locus, the FMR locus and progression of a disease condition wherein the computer comprises:

[0381] (1) a machine-readable data storage medium comprising a data storage material encoded with machine-readable data, wherein the machine-readable data comprise values associated with the features of one or more of:

[0382] (a) change in of methylation or other epigenetic modification relative to a control in FREE3 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions upstream;

[0383] (b) change of methylation or other epigenetic modification relative to a control in CpG and/or CpNpG islands and island shores in intron 2 of the FMR1 gene comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0384] (c) change in methylation or other epigenetic modification relative to a control in genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0385] (d) change in methylation of an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;

[0386] (e) two or more of (i) an intron; (ii) an intron/exon boundary; (iii) splicing region within the FMR genetic locus;

[0387] (f) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

[0388] (g) length of (CGG)_n expansion within the FMR genetic locus when considered in combination with (a) and/or (b);

[0389] (h) general phenotype or clinical manifestations in subjects with a neurodevelopmental or neurodegenerative condition;

[0390] (i) behavioral assessment criteria associated with normal subjects, PM subjects, GZ subjects and FM subjects;

[0391] (j) cognitive ability;

[0392] (k) extent of transcription of genes within the FMR locus with the proviso that if one or more of (d) through (k) is determined, then one or more of (a) through (c) or (k) is also determined;

[0393] (2) means to converting the value to a code; and

[0394] (3) means to store the code in a computer readable medium and compare code to a knowledge database to determine whether the code corresponds to a pathological condition.

[0395] The computer system of the present disclosure may also be linked to detection systems such as MALDI-TOF mass spectrometry machines.)

[0396] The present disclosure further provides a web-based system where data on extent of methylation within a genetic locus (optionally together with clinical phenotype) are provided by a client server to a central processor which analyzes and compares to a control and optionally considers other information such as patient age, sex, weight and other medical conditions and then provides a report, such as, for example, a risk factor for disease severity or progression or status or response to treatment or an index of probability of a genetic locus-associated pathology in a subject.

[0397] Hence, knowledge-based computer software and hardware also form part of the present disclosure.

[0398] In an embodiment, the assays herein may be used in existing or newly developed knowledge-based architecture or platforms associated with pathology services. For example, results from the assays are transmitted via a communications network (e.g. the internet) to a processing system in which an algorithm is stored and used to generate a predicted posterior probability value which translates to the index of disease probability which is then forwarded to an end user in the form of a diagnostic or predictive report.

[0399] The assay may, therefore, be in the form of a kit or computer-based system which comprises the reagents necessary to detect the extent of methylation or other epigenetic modification within the genetic locus and includes computer hardware and/or software to facilitate determination and transmission of reports to a clinician.

[0400] The assay of the present disclosure permits integration into existing or newly developed pathology architecture or platform systems. For example, the present disclosure contemplates a method of allowing a user to determine the status of a subject with respect to an FMR locus-associated pathology, the method including:

[0401] (a) receiving data in the form of extent of methylation or other epigenetic modification at a site within:

[0402] (A) the FMR1 gene selected from:

[0403] (i) Fragile X-related Epigenetic Element 3 [FREE3] comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0404] (ii) intron 2 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions; and

[0405] (iii) change in methylation or other epigenetic modification relative to a control in genomic FREE2

region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0406] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 or a homolog thereof or a fragment or portion thereof;

[0407] (B) the FMR genetic locus selected from:

[0408] (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; or

[0409] (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region;

wherein the extent of methylation or other epigenetic modification provides a correlation to the presence, state, classification or progression of the pathology; by transferring the data from the user via a communications network;

[0410] (c) processing the subject data via multivariate or univariate analysis to provide a disease value;

[0411] (d) determining the status of the subject in accordance with the results of the disease value in comparison with predetermined values; and

[0412] (e) transferring an indication of the status of the subject to the user via the communications network. Reference to the multivariate or univariate analysis includes an algorithm which performs the multivariate or univariate analysis function.

[0413] Conveniently, the method generally further includes:

[0414] (a) having the user determine the data using a remote end station; and

[0415] (b) transferring the data from the end station to the base station via the communications network.

[0416] The base station can include first and second processing systems, in which case the method can include:

[0417] (a) transferring the data to the first processing system;

[0418] (b) transferring the data to the second processing system; and

[0419] (c) causing the first processing system to perform the multivariate analysis function to generate the disease value.

[0420] The method may also include:

[0421] (a) transferring the results of the multivariate or univariate analysis function to the first processing system; and

[0422] (b) causing the first processing system to determine the status of the subject.

[0423] In this case, the method also includes at least one of:

[0424] (a) transferring the data between the communications network and the first processing system through a first firewall; and

[0425] (b) transferring the data between the first and the second processing systems through a second firewall.

[0426] The second processing system may be coupled to a database adapted to store predetermined data and/or the multivariate analysis and/or univariate analysis function, the method including:

[0427] (a) querying the database to obtain at least selected predetermined data or access to the multivariate or univariate analysis function from the database; and

[0428] (b) comparing the selected predetermined data to the subject data or generating a predicted probability.

[0429] The second processing system can be coupled to a database, the method including storing the data in the database.

[0430] The method can also include having the user determine the data using a secure array, the secure array of elements capable of determining the extent of methylation in an intron with a genetic locus and having a number of features each located at respective position(s) on the respective code. In this case, the method typically includes causing the base station to:

[0431] (a) determine the code from the data;

[0432] (b) determine a layout indicating the position of each feature on the array; and

[0433] (c) determine the parameter values in accordance with the determined layout, and the data.

[0434] The method can also include causing the base station to:

[0435] (a) determine payment information, the payment information representing the provision of payment by the user; and

[0436] (b) perform the comparison in response to the determination of the payment information.

[0437] The present disclosure also teaches a base station for determining the status of a subject with respect to a pathology associated with a genetic locus such as the FMR locus, the base station including:

[0438] (a) a store method;

[0439] (b) a processing system, the processing system being adapted to;

[0440] (c) receive subject data from the user via a communications network, the data; including extent of methylation within the genetic locus wherein the level or methylation or epigenetic modification relative to a control provides a correlation to the presence, state, classification or progression of the pathology;

[0441] (d) performing an algorithmic function including comparing the data to predetermined data;

[0442] (e) determining the status of the subject in accordance with the results of the algorithmic function including the comparison; and

[0443] (f) output an indication of the status of the subject to the user via the communications network.

[0444] The processing system can be adapted to receive data from a remote end station adapted to determine the data.

[0445] The processing system may include:

[0446] (a) a first processing system adapted to:

[0447] (i) receive the data; and

[0448] (ii) determine the status of the subject in accordance with the results of the multivariate or univariate analysis function including comparing the data; and

[0449] (b) a second processing system adapted to:

[0450] (i) receive the data from the processing system;

[0451] (ii) perform the multivariate or univariate analysis function including the comparison; and

[0452] (iii) transfer the results to the first processing system.

[0453] The determination of the extent of methylation or other epigenetic modification within the FMR locus at a site within the FMR1 gene selected from:

[0454] (i) Fragile X-related Epigenetic Element 3 [FREE3] comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0455] (ii) intron 2 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0456] (iii) change in methylation or other epigenetic modification relative to a control in genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0457] (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 or a homolog thereof or a portion or fragment thereof;

[0458] (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; or

[0459] (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

enables establishment of a diagnostic or prognostic rule based on the extent of methylation relative to controls. Alternatively, the diagnostic or prognostic rule is based on the application of a statistical and machine learning algorithm. Such an algorithm uses relationships between methylation profiles and disease status observed in training data (with known disease status) to infer relationships which are then used to predict the status of patients with unknown status. An algorithm is employed which provides an index of probability that a patient has an FMR locus-associated pathology. The algorithm performs a multivariate or univariate analysis function.

[0460] Hence, the present disclosure teaches a diagnostic rule based on the application of statistical and machine learning algorithms. Such an algorithm uses the relationships between epigenetic profile and disease status observed in training data (with known disease status) to infer relationships which are then used to predict the status of patients with unknown status. Practitioners skilled in the art of data analysis recognize that many different forms of inferring relationships in the training data may be used without materially changing the present disclosure.

[0461] The present disclosure teaches a knowledge base of training data comprising extent of methylation within a genetic locus such as the FMR genetic locus from a subject with locus-associated pathology to generate an algorithm which, upon input of a second knowledge base of data comprising levels of the same biomarkers from a patient with an unknown pathology, provides a probability that predicts the nature of unknown pathology or response to treatment.

[0462] The term "training data" includes knowledge of the extent of methylation relative to a control. A "control" includes a comparison to levels in a healthy subject devoid of a pathology or is cured of the condition or may be a statistically determined level based on trials.

[0463] The present disclosure contemplates, therefore, the use of the methylation, including epigenetic profile of intronic sites within the FMR genetic locus and in particular the FMR1 gene to assess or determine the status of a subject with respect to disease, to stratify a subject relative to normal controls or unhealthy subjects, to provide a prognosis of recovery or deterioration and/or to determine the pharmacoresponsiveness or pharmacosensitivity of a subject to treatment or an agent for use in treatment and/or determine applicability for other treatment options including behavioural intervention, and the like. By "intronic sites" includes intron/exon boundaries and splicing regions.

[0464] Hence, another aspect enabled herein is a method of allowing a user to determine the status, prognosis and/or treatment response of a subject with respect to an FMR locus-associated pathology, the method including:

[0465] (a) receiving data in the form of extent of methylation or other epigenetic modification at a site in:

[0466] (i) FMR1 gene selected from:

[0467] (ii) Fragile X-related Epigenetic Element 3 [FREE3] comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

[0468] (iii) intron 2 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;

[0469] (iv) change in methylation or other epigenetic modification relative to a control in genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

[0470] (v) an intron, intron/exon boundary and/or splicing region downstream of intron 2 or a homolog or a portion or fragment thereof;

[0471] (vi) two or more of (a) an intron; (b) an intron/exon boundary; (c) a splicing region within the FMR locus; or

[0472] (vii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;

wherein the extent of methylation or epigenetic modification provides a correlation to the presence, state, classification or progression of the pathology;

[0473] (b) transferring the data from the user via a communications network;

[0474] (c) processing the subject data via multivariate or univariate analysis to provide a disease value;

[0475] (d) determining the status of the subject in accordance with the results of the disease value in comparison with predetermined values; and

[0476] (e) transferring an indication of the status of the subject to the user via the communications network.

[0477] Aspects disclosed herein are further described by the following non-limiting Example. In these Examples, materials and methods as outlined below were employed.

Patient Samples

[0478] FXS and premutation carrier EBV transformed lymphoblast cell lines were obtained from the tissue culture storage repository of the Murdoch Childrens Research Institute, Melbourne, Victoria, Australia or purchased from Coriell.

DNA Extraction

[0479] DNA for CGG repeat size PCR and methylation analysis was obtained either from 200 μ l venous blood samples anti-coagulated with EDTA or from EBV transformed lymphoblasts 1 to 5×10^6 cells per sample and extracted using a BIO ROBOT M48 DNA Extractor, as per manufacturer's instructions (Qiagen Inc., Hilden, Germany). DNA for Southern blot or methylation analysis was extracted from 3 ml blood samples anti-coagulated with EDTA or from EBV transformed lymphoblasts 5 to 10×10^6 cells per sample.

CGG Repeat Size PCR Amplification

[0480] CGG repeat size for all samples was initially assessed using a fully validated PCR assay with precision of $\pm$ one triplet repeat across the normal and GZ ranges, performed using a fragment analyzer (MegaBace, GE Healthcare), with the higher detection limit of 170 repeats, as previously described (Khaniani et al., *Mol Cytogenet* 1(1):5, 2008). Briefly, PCR amplifications were performed using primers:

```
(i) r (5'-GCTCAGCTCCGTTTCGGTTTCACTTCCGGT-3'
    [SEQ ID NO: 32]);
    and
(ii) f (5'-AGCCCGCAGCTTCCACCAACAGCTCCTCCA-3'
    [SEQ ID NO: 33]),
```

(Fu et al., *Cell* 67(6):1047-1058, 1991) in a total volume of 25 μ l containing 50 ng of genomic DNA, 0.75 pmol of each primer, 8 μ l of 5 $\times$ Q-Solution (Qiagen Inc., Hilden, Germany), 2.5 μ l of 10 $\times$ PCR Buffer and 1 unit of HotStarTaq Plus DNA polymerase (Qiagen Inc., Hilden, Germany) in a Gene Amp ® PCR System 9700. The PCR cycling profile was as follows: initial denaturation at 98 $^{\circ}$ C. for 5 minutes; 35 cycles at 98 $^{\circ}$ C. for 45 seconds, 70 $^{\circ}$ C. for 45 seconds, and 72 $^{\circ}$ C. for 2 minutes, and a final extension at 72 $^{\circ}$ C. for 10 minutes. Alleles were sized by capillary electrophoresis using an automatic sequencer (MegaBACE $^{\text{TM}}$ 1000—GE HealthCare Amersham) with size standards (HealthCare) and controls of lengths 10, 23, 29, 30, 52 and 74 repeats determined by sequencing in-house or obtained from Coriell Cell Repositories web site (<http://www.phppo.cde.gov/dls/genetics/qcmaterials/>).

CGG Repeat Size by Southern Blot

[0481] CGG sizes were assessed using a fully validated Southern Blot procedure with appropriate normal and abnormal controls for samples where the products could not be amplified using PCR (Fu et al 1991, supra; Francis et al., *Mol Diagn* 5(3):221-225, 2000). Briefly, 5 mg of DNA was digested with Pst1 (Boehringer Mannheim, Castle Hill, Australia), separated on 1% w/v agarose gels, and analyzed by

Southern blot hybridization. The FMR-1 gene was detected using Southern blot analysis with probe Fxa3 and an X-chromosome control probe, pS8 (Yu et al., *Science* 252(5010): 1179-1181, 1991). Probes were labeled using random oligonucleotide priming (Boehringer, Mannheim) with [α^{32} -P] CTP (NEN Dupont, Boston, Mass.). Autoradiography was performed at -80° C., with intensifying screens and Kodak XAR films (Sigma-Aldrich).

Methylation Sensitive Southern Blot Analysis

[0482] Methylation of the classical FMR1 CpG island was assessed using a fully validated methyl sensitive Southern Blot procedure with appropriate normal and abnormal controls, as previously described (Tassone et al., *J. Mol. Diagn* 10:43-49, 2008). Briefly, EcoRI and NruI digestion was performed on 7 to 9 μ g of DNA, and separated on a 0.8% w/v agarose/Tris acetate EDTA (TAE) gel. The DNA was denatured with HCL and NaOH, transferred to a charged nylon membrane and analyzed by Southern blot hybridization. The FMR1 alleles were detected using Southern blot analysis with probe StB12.3, labeled with Dig-11-dUTP by PCR (PCR Dig Synthesis kit; Roche Diagnostics). Autoradiography was performed with intensifying screens and Fuji Medical X-Ray film (Bedford, UK) and FMR1 methylation values for the expanded alleles were calculated as previously described (Tassone et al., 2008 supra). The FMR1 activation ratios for female samples were calculated based on the following formula: optically scanned density of the 2.8 kb band/combined densities of the 2.8 kb and 5.2 kb bands (where the 2.8 kb band represents the proportion of normal active X and the 5.2 kb band represents the proportion of normal inactive X), as previously described (de Vries et al., *Am J Hum Genet.* 58:1025-1032, 1996).

MALDI-TOF Methylation Analysis

Bisulfite Treatment

[0483] Bisulfite treatment of genomic DNA at 0.5 μ g per sample was performed using XCEED kit from MethylEasy (Human Genetic Signatures, Sydney, Australia) for sample sets of $n < 40$. For sample sets $n > 40$ 96 well Methylamp kit from Epigentek (Brooklyn, N.Y., USA) was used. Duplicate bisulfite reactions were made from each sample, and six of the same control DNA samples spiked with DNA from an FXS patient cell line at 0, 33.3, 50, 66.6 or 100% were included as standards within each run, as an indicator of the inter-run variation in the degree of bisulfite related bias. Protocols were performed according to the manufacturer's instructions. Briefly, for the MethylEasy conversion, 20 μ l of genomic DNA (0.5 μ g total) was mixed with 2.2 μ l of 3 μ l NaOH, and incubated at 37° C. for 15 minutes, then denatured by 45 minute incubation at 80° C. 240 μ l of the reagent #3 (XCEED kit, Human Genetic Signatures, Sydney, Australia) were then added to the mixture, which was transferred into the purification column and spun down at 10,000 g for 1 minute. The captured DNA was then washed in Reagent #4 (XCEED kit, Human Genetic Signatures, Sydney, Australia), and DNA eluted twice by placing 50 μ l of the pre-warmed solution #5 (XCEED kit, Human Genetic Signatures, Sydney, Australia) onto column membrane, which was incubated for 1 minute at room temperature, and spun down at 10,000 g for 1 minute. The eluted DNA was then incubated at 95° C. for 20 minutes, with resulting final concentration at ~ 20 ng/ μ l per sample.

[0484] For the Methylamp conversion, 7 μ l of genomic DNA (0.5 μ g total) was mixed with 5 μ l of the CF3 (Methylamp kit, Epigentek, Brooklyn, N.Y., USA) solution diluted 1:10 in distilled water, in each well of the 96 well plate. The DNA was denatured by placing the plate at 65° C. for 90 minutes. It was then captured in the filter plate and washed in 150 μ l of the CF5 solution (Methylamp kit, Epigentek, Brooklyn, N.Y., USA), then twice in 250 μ l of 80% v/v ethanol. The filter plate was then incubated in the CF3/90% ethanol solution, and washed twice in 90% v/v ethanol, as per manufacturer's instructions. The modified and cleaned DNA was then eluted with 40 μ l of the CF6 solution (Methylamp kit, Epigentek, Brooklyn, N.Y., USA), with resulting converted DNA final concentration at ~ 20 ng/ μ l per sample. For the short term storage the converted DNA was kept at -20° C., and for storage of more than 3 months it was kept at -80° C.

PCR and In Vitro Transcription

[0485] The primers used to amplify the target regions and the annealing temperatures are listed in Tables 3 and 4. Each bisulfite converted sample was analyzed in duplicate PCR reactions, carried out in a total volume of 5 μ l using 1 pmol of each primer, 40 μ M dNTP, 0.2 U Hot Star Taq DNA polymerase (Qiagen Inc., Hilden, Germany), 1.5 mM $MgCl_2$ and buffer supplied with the enzyme (final concentration $1\times$). The reaction mix was pre-activated for 15 min at 95° C., followed by 45 cycles of amplification at 94° C. for 20 s, primer specific annealing for 30 s and 72° C. for 1 min followed by 72° C. for 3 min. The PCR products were run on 1.5% w/v agarose gel to confirm successful PCR amplification and efficiency. The DNA was then cleaned up and the T or C-cleavage reactions were carried out (1-cleave for Amplicons 1 to 5, C-cleave for Amplicon 5 only) as per manufacturer's instructions (SEQUENOM, San Diego, Calif.). Briefly, unincorporated dNTPs were dephosphorylated by adding 1.7 μ l H_2O and 0.3 U Shrimp Alkaline Phosphatase (SAP) [SEQUENOM, San Diego] to PCR products, which were incubated at 37° C. for 20 min, and 10 min at 85° C. to heat-inactivate the SAP. The transcription was performed on 2 μ l of template DNA in the 6.5 μ l reaction consisting of 20 U of the T7 R&DNA polymerase (Epicentre, Madison, Wis.) to incorporate either dCTP or dTTP; Ribonucleotides at 1 mM and the dNTP substrate at 2.5 mM, with other components used as recommended (SEQUENOM, San Diego). RNase A (SEQUENOM, San Diego) was then added to the mix to cleave the in vitro transcript. The mix was diluted to 27 μ l in H_2O , and 6 mg CLEAN Resin (SEQUENOM, San Diego, Calif.) was added for conditioning of the phosphate backbone prior to MALDI-TOF MS. The SEQUENOM Nanodispenser was then used to spot the samples onto a SpectroCHIP for subsequent analysis. MassARRAY mass spectrometer (Bruker-SEQUENOM) was then used to collect mass spectra, which were analysed using the EpiTYPER software (Bruker-SEQUENOM). The calculation of the output methylation ratios for each CpG unit were based on the ratio of the signal intensities for the fragment from a methylated CpG unit/[methylated+unmethylated CpG units]. Further details are described in (Godler et al., 2010 supra).

RNA Extractions and Quality Assessments

[0486] Total RNA was extracted and purified using the Rneasy extraction kit, as per manufacturer's instructions (Qiagen Inc., Hilden, Germany). RNA concentrations were

measured in triplicate using a NanoDrop ND-1000 Spectrophotometer, with purity being determined by the A260/A280 ratio using the expected values between 1.8 and 2. Total RNA quality and the degree of DNA contamination was also assessed using capillary electrophoresis Standard Sens Kit (Bio-rad), which involved descriptive comparison of chromatographic features based on previous publications using this system (Fleige and Pfaffl, *Mol Aspects Med* 27(2-3):126-139, 2006). Each RNA sample was then diluted to 30 ng/ul, to be used in for reverse transcription real-time PCR analysis, where mRNA quality at the Xq27.3 region was initially assessed by examining the relationship between 5' and 3' levels of FMR1 mRNA.

Standard Reverse Transcription Real-Time PCR

[0487] Reverse transcription was performed one reaction per sample using the Multiscribe Reverse Transcription System, 50 units/ul (Applied Biosystems). The 7900HT Fast Real Time PCR (Applied Biosystems) was used to quantify FMR1-5', FMR1-3', ASFMR1 (-1), (-2), (-3), GAPDH, B2M, and GUS, using the relative standard curve method. The target gene and the internal control gene dynamic linear ranges were performed on a series of doubling dilutions of an RNA standard (160-4 ng/ul). Since, both ASFMR1 assays do not target an exon/exon boundary, to minimize the impact of potential DNA contamination on the expression results, a no reverse transcription enzyme control was included for every sample. The difference between the plus and minus no reverse transcriptase control was considered as the ASFMR1 expression value for each sample. Previously published sequences were be used for primers and probe for: FMR1-5' and GUS (32); FMR1-3' (41). The following ASFMR1 primers and probes were designed using Primer Express 3.0 (Applied Biosystems):

ASFMR1 (-1)
Fw Primer [SEQ ID NO: 34]
(CCGCGGAATCCCAGAGA);
Rv Primer: [SEQ ID NO: 35]
(CAGTGGCGTGGGAAATCAA);
Probe: [SEQ ID NO: 36]
(FAM-TGGGATAACCGGATGCA-MGB).

-continued

ASFMR1 (-2)
Fw Primer: [SEQ ID NO: 37]
(ACACCCCTGTGCCCTTTAAGG);
Rv Primer: [SEQ ID NO: 38]
(TCAAAGCTGGGTCTGAGGAAAG);
Probe: [SEQ ID NO: 39]
(VIC-TCGGGATCTCAAATGT-TAMRA).
ASFMR1 (-3)
Fw Primer: [SEQ ID NO: 40]
(CCCCAGAATGAGAGGATGTTG);
Rv Primer: [SEQ ID NO: 41]
(GCCCTAGATCCACCGCTTTAA);
Probe: [SEQ ID NO: 42]
(FAM-TGCTGGTGGAATC-MGB).

[0488] FMR1-5', FMR1-3', ASFMR1 primers and probes were be used at concentrations of 18 μ M and 2 μ M, respectively. GAPDH and B₂M primer/probe mixes will be obtained from PrimerDesign (PerfectProbe ge-PP-12-hu kit) and used at concentration of 2 μ M. All of the above assays were singleplexed, with each sample assayed in duplicate 10 μ l PCR reactions. The reactions consisted of 5.8 mM MgCl₂, 1 μ l Buffer A (Applied Biosystems), 3.35 μ l Rnase-free water, 1.2 mM dNTPs, 0.01 units/ μ l of AmpliTaq Gold, 0.5 μ l of Taq-Man probe and 0.5 μ l forward and 0.5 μ l reverse primers, and 1 μ l of the reverse transcription (cDNA) reaction. The annealing temperature for thermal cycling protocol was 60° C. for 40 cycles. The samples were quantified in arbitrary units (au) in relation to the standard curves performed on each plate, standardized to the mean of the 3 internal control genes (GUS, GAPDH and B₂M).

Amplicons

[0489] Amplicons were amplified using the primers and conditions shown in Tables 3 and 4. Table 5 shows prominent regulatory motif locations inclusive and proximal to FREE2 in FREE3. Amplicon 5 is as described by Godler et al., *Hum Mol Genet* 10(8):1618-1632. [Epub 2010]; Godler et al., *J. Mol Diagn.* 2011 [Epub ahead of print] doi: 10.1093/hmg/ddq1037, the contents of which are incorporated by reference.

TABLE 3

Amplicon details used for MALDI-TOF methylation analysis of the regions greater than 0.2 kb 3' of the CGG expansion at the Xq27.3 locus					
Amplicon No.	Annealing Temperature	Size (kb)	Distance 3' of CGG end (kb):	Primer sequence (in capitals) and tag (in lower case)	
FREE2(B)	(I) 94° C. 4 min; 25 cycles of: touchdown PCR -0.5° C. per cycle-94° C. 20 s; 64° C. 30 s; 72° C. 1 min. (II) 20 cycles of: 94° C. for 20s, 59° C. for 30s,	500	0.207	Fw: 5'-aggaagagagGGTTTTTTGAAATTTTGGATTTA-3' Rv: 5'-cagtaatacgactcactataggagaaggctTAAACCTATTAAAA ACCCCTCTCC-3'	

TABLE 3-continued

Amplicon details used for MALDI-TOF methylation analysis of the regions greater than 0.2 kb 3' of the CGG expansion at the Xq27.3 locus				
Amplicon No.	Annealing Temperature	Size (kb)	Distance 3' of CGG end (kb):	Primer sequence (in capitals) and tag (in lower case)
	72° C. for 1 min. (III) 72° C. for 3 min; 4° C. forever			
FREE2 (C)	(I) 94° C. 4 min; 25 cycles of: touchdown PCR -0.5° C. per cycle-94° C. 20 s; 64° C. 30 s; 72° C. 1 min. (II) 20 cycles of: 94° C. for 20 s, 59° C. for 30 s, 72° C. for 1 min. (III) 72° C. for 3 min; 4° C. forever	302	0.504	Fw: 5'-aggaagagagTAAGAGGGTTTTAGGTTTTTTTGG-3' Rv: 5'-cagtaatacgactcactataggagaaggctAAAACATATACATTCCTAAATTACCCC-3'
FREE3 (ASFMR1)	(I) 94° C. 4 min; 25 cycles of: touchdown PCR -0.5° C. per cycle-94° C. 20 s; 64° C. 30 s; 72° C. 1 min. (II) 20 cycles of: 94° C. for 20 s, 59° C. for 30 s, 72° C. for 1 min. (III) 72° C. for 3 min; 4° C. forever	327	9.739	Fw: 5'-aggaagagagTTTTTTTATATAGGTATTGTAAAGGATG-3' Rv: 5'-cagtaatacgactcactataggagaaggctTCTCTAATTCTTTCTTCACATTCAAAA-3'

TABLE 4

Amplicon details used to define FREE2 3' Border region using MALDI-TOF methylation analysis within the FMR1 intro 1 sequence at the Xq27.3 locus				
Amplicon No.	Annealing Temperature	Size (kb)	Distance 3' of CGG end (kb):	Primer sequence (in capitals) and tag (in lower case)
FREE2 (D) (pp4)	(I) 95° C. 15 min (II) 10 cycles of: 94° C. 20 s; 56° C. 30s; 72° C. 1 min. 35 cycles of: 94° C. for 20 s, 62° C. for 30 s, 72° C. for 1 min. (III) 72° C. for 3 min; 4° C. forever	374	0.790	Fw: 5'-aggaagagagAAAAGTTTTAGGAAGATTTTAATATGG-3' Rv: 5'-cagtaatacgactcactataggagaaggctAAAAACA CAATAAACCCATAAATACC-3'
FREE2 (E) (pp6)	(I) 95° C. 15 min (II) 10 cycles of: 94° C. 20 s; 56° C. 30 s; 72° C. 1 min. 35 cycles of: 94° C. for 20 s, 62° C. for 30 s, 72° C. for 1 min. (III) 72° C. for 3 min; 4° C. forever	360	1.424	Fw: 5'-aggaagagagGAATGGTTTGAATGTTTATAGATAGGAT-3' Rv: 5'-cagtaatacgactcactataggagaaggctACCAAAAA TCTAATAACCAAAACCAC-3'
FREE3 (ASFMR1)	(I) 95° C. 15 min (II) 10 cycles of: 94° C. 20 s; 56° C. 30 s; 72° C. 1 min. 35 cycles of: 94° C. for 20 s, 62° C. for 30 s, 72° C. for 1 min.	327	9.908	Fw: 5'-aggaagagagTTTTTTTATATAGGTATTGTAAAGGATG-3' Rv: 5'-cagtaatacgactcactataggagaaggctTCTCTAATTTCTTCTTCACATTCAAAA-3'

TABLE 4-continued

Amplicon details used to define FREE2 3' Border region using MALDI-TOF methylation analysis within the FMR1 intro 1 sequence at the Xq27.3 locus				
Amplicon No.	Annealing Temperature	Distance		Primer sequence (in capitals) and tag (in lower case)
		Size (kb)	3' of CGG 3' end (kb):	
	30s, 72° C. for 1 min. (III) 72° C. for 3 min; 4° C. forever			

TABLE 5

Prominent regulatory motif locations inclusive and proximal to FREE2 and FREE3 regions					
TRANSCRIPTION FACTOR SITES/POTENTIAL REGULATORY MOTIFS:	SEQUENCE ON THE SENSE STRAND	STRAND	SEQUENCE HOMOLOGY (%)	AMPLICON	CpG UNIT LOCATION:
GATA-1	GGCGATGGCT	LEADING	95	FREE2 (A)	CpG15 and CpG16
HSF2	TGAATATTCG	LEADING	96	FREE2 (B)	CpG7 and CpG8/9
C/EBP	AAGTTTCCAAAGA	LAGGING	95	FREE2 (D)	CpG6 and CpG7
CdxA	TATTATTATT	LAGGING	99	FREE2 (D)	CpG6 and CpG7
AML-1a	ACCACA	LAGGING	100	FREE2 (D)	3' of CpG7
AML-1a	TGTGGTGT	LEADING	100	FREE2 (D)	3' of CpG7
CdxA	TATAAAT	LAGGING	100	N/A	Between FREE2 (D) and FREE2 (E)
CdxA	TATAAAT	LAGGING	100	N/A	Between FREE2 (D) and FREE2 (E)
CdxA	AATAATAT	LEADING	99	N/A	Between FREE2 (D) and FREE2 (E)
HFH-1/HFH-2	AAATAACAAT	LAGGING	97	FREE2 (E)	CpG1 and CpG2/3
CdxA	CATAAAT	LAGGING	100	FREE2 (E)	CpG1 and CpG2/3
SRV	TTTGTTT	LAGGING	100	N/A	3' of FREE2 (E) CpG6
SRV	TTTGTTT	LAGGING	100	N/A	3' of FREE2 (E) CpG6
SRV	TTGTTTA	LAGGING	99	N/A	3' of FREE2 (E) CpG6
S8	TTTATTTAATTAAGTT	LEADING	96	N/A	3' of FREE2 (E) CpG6
SRV	AAACAAA	LEADING	100	5' of FREE3	5' of FREE3 CpG1
CdxA	TATAATT	LEADING	99	FREE3	CpG1
Oct-1	TTTATGCTAATT	LEADING	99	FREE3	Between CpG1 and CpG2

Example 1

Mapping Methylation of the FMR Genetic Locus
Using High Throughput Mass Spectrometry

[0490] The structure of the FMR genetic locus is shown in FIG. 1A and FIG. 6A and comprises the FMR1 promoter, and FMR1 and ASFMR1 genes. A CGG repeat is located within the 5' (UTR) of the FMR1 gene. ASFMR1 spans the CGG expansion in the antisense direction and is also regulated by another promoter located in the exon 2 of FMR1. The FREE2 located downstream of the CGG expansion. The FREE3 region is located within intron 2 of FMR1 downstream of the second ASFMR1 promoter.

[0491] The primers utilized for MALDI-TOF methylation analysis targeted 4 regions at the Xq27.3 locus designated as FREE2(A) [described as amplicon 5 in Godler et al., 2010 supra]; FREE2(B); FREE2(C) and FREE3 (color coded). Individual CPG sites within each region are numbered accordingly. Prominent transcription factor binding sites and methylation sensitive restriction enzyme recognition sites are indicated in capital font, and are listed/identified in Tables 3 and 4. Numerous HpaII/MspI sites (CCGG) are located throughout the FREE2 A, B and C region.

[0492] Regions identified as biologically significant showed consistent differences in methylation between healthy controls and FXS samples (FIG. 1B and FIGS. 2A and B). These include HpaII/MspI sites throughout FREE2 A, B and C regions including but not restricted to the FREE2B CCGG sites located at CpGs 6, 9, 13 and between CpGs 25 and 26; as well as FREE2 (C) CCGG site located at CpG1. These would be sensitive to HpaII methylation specific digestion, which can be followed by PCR or other restriction enzyme based methods to assay differential methylation between healthy controls and FXS samples, and potentially carriers of smaller expansion alleles.

[0493] Other regions identified as biologically significant that showed consistent differences in methylation between healthy controls and FXS samples (FIG. 1B and FIGS. 2A and B and Table 5) included: (I) GATA-1 site (FREE2B between CpG 15 and 16); (II) HSF2 site (FREE2C between CpG 7, 8 and 9); (III) an SKY site located upstream of FREE3; (IV) a CdxA/TATA box site located at CpG1 of FREE3; (V) an Oct-1 site located between CpG sites 1 and 2 within FREE3. Differential methylation of any of these sites in diseased individuals compared to controls may have an affect of relevant transcription factor binding and/or further epigenetic modification; which would inturn affect transcription of FMR1, ASFMR1 and/or FMR4. Or may result or reflect aberrant non coding RNA expression and/or RNA:DNA interactions or stability of RNA:DNA hybrids (FIGS. 4B and C and FIG. 5).

Example 2

Determining the Impact of Technical Variation on
Quantitative Analysis of Methylation and Evidence
for Disease Specific Methylation within Intron 1 and
Intron 2 of FMR1

[0494] DNA from lymphoblasts of healthy controls with 30 CGG repeats, normal levels of FMR1 mRNA and FMRP, and DNA from lymphoblasts of FXS patient with 530 CGG, silenced FMR1 transcription and absence of FMRP were mixed at ratios of 1:0; 2:1; 1:1; 1:2; 0:1 corresponding to 0,

33.3, 50, 66.6, 100% FXS DNA in the sample (FIG. 3). The spiked DNA samples were bisulfite converted in duplicate reactions. Each reaction was amplified with primer sets (forward and reverse primers) as listed by SEQ ID NOs: which corresponded to 3 SEQUENOM mass spectrometry assays (A: FREE2(B); B: FREE2(C); C: FREE3). The spiked DNA samples were analysed using MALDI-TOF methylation analysis at three sequential regions at the Xq27.3 locus (see FIG. 1 for locations). The methylated vs unmethylated ratios at each analysable CpG unit were expressed as output methylation ratios on Y axis, with FXS DNA input % expressed on the X axis (each point represents mean of duplicate PCRs from a single bisulfite converted DNA mixture). Methylation output ratios for CpG sites within FREE2B and FREE2C amplicons (A and B) were positively correlated with increasing FXS DNA input %; while FREE3 Methylation output ratios were negatively correlated with increasing FXS DNA input % with high Pierson's correlation. This clearly demonstrates that the FREE2 region comprising a large portion of FMR1 intron 1 is hypermethylated in FXS sample while FREE3 region within intron 2 of FMR1 is hypomethylated in the FXS sample. This methylation pattern is reversed in the healthy control sample, and supports the differential methylation patterns within FREE2 and FREE3 related to the disease state as shown in FIGS. 2A and B.

Example 3

Evidence for Expression of ASFMR1 in FXS and
Disease Specific RNA:DNA Interactions

[0495] Standard curve and amplification real-time PCR plots (of assays described in FIG. 4A) show that in the FXS cell lines, ASFMR1 RNA with fully methylated FMR1 promoter and silenced FMR1 and FMRP, ASFMR1 is expressed (FIG. 4B). RNA was extracted from 3 FXS cell lines whose methylation profiles are presented in FIG. 2; Sample 849 was taken from the male 490 CGG repeat line; Sample 862 was taken from the male 530 CGG repeat line; Sample 865 was taken from the female 563 and 47 CGG repeat line. Each RNA sample was split in two, with one half subjected to RNase A treatment prior to ASFMR1 (-3) relative standard curve analysis. The ASFMR1 (-3) real-time PCR analysis was performed in quadruplicate reactions. The difference in Ct values between RNase A treated and untreated samples represents the level of ASFMR1 expression.

[0496] Standard curve and amplification real-time PCR plots (FIGS. 4C and D) also indicate that in the FXS cell lines, ASFMR1RNA forms RNA:DNA complexes. FXS RNA samples were treated with TURBO DNase (C) and RQ1 DNase (D) respectively. These DNase treatments caused complete loss of real-time-PCR signal for the ASFMR1(-3) assay. Because DNase can only degrade RNA molecules if they form complexes with DNA, loss of ASFMR1 after DNase treatment suggests that ASFMR1RNA forms RNA:DNA complexes in FXS samples with fully methylated FMR1 promoter and silenced FMR1 expression.

[0497] Expression of different FMR1 and ASFMR1 transcripts (detailed in FIG. 4A) was detected in RNA samples from lymphoblast lines of 6 male controls, two FXS males (samples 849 and 862) and one FXS female (865) [FIGS. 5A and B]. The control and FXS RNA samples were either treated with TURBO DNase (A), RQ1 DNase (B), RNase A (C), or were untreated. Addition of TURBO DNase or RQ1 DNase buffers to RNA samples without DNase were

included as additional controls in (FIGS. 5A and B). The FMR1 and ASFMR1 transcripts were quantified using real-time RT-PCR relative standard curve method, normalized to mRNA levels of three internal control genes, GUS, GAPDH and B2M. FMR1 5' and 3' assays showed no signal for the FXS RNA samples, while similar levels were detected in all control samples (upper two panels in FIGS. 5A, B and C). TURBO and RQ1 DNase treatment caused ~50% decrease in the FMR1 levels in most of the control samples; while RNase A treatment caused complete loss of FMR1 and ASFMR1 signals. While decrease of ASFMR1 (-1), (-2) and (-3) levels was also observed in all control samples caused by TURBO and RQ1 DNase treatment, in FXS samples (with analogous to control ASFMR1 levels in the untreated samples) TURBO and RQ1 DNase treatment resulted in complete loss of signal for all three ASFMR1 assays. Because DNase can only degrade RNA molecules if they form complexes with DNA, this suggests that ASFMR1 RNA forms RNA:DNA complexes more readily in FXS samples than in controls. Increase in RNA:DNA interaction of ASFMR1 in FXS may lead to methylated FMR1 promoter and adjacent regions (FIG. 1) and silencing FMR1 expression leading to loss of FMRP and the resulting FXS clinical phenotype.

Example 4

Characterizing the 3' Boundary of the FMR1 Promoter Located within the FREE2 Region—1.310 from the 5' End of the FMR1 Intron 1—Relative to Transcription

[0498] There is significant inhibition of transcription of the FMR4, ASFMR1 and FMR1 genes in FXS patients which is likely caused by the methylation of the CpG units in Amplicon 1 (FREE1 region), and/or Amplicon 5 (FREE2 region) in association with the FMR1 promoter. It is demonstrated here that the FREE2 region and FMR1 promoter expands 1302 base pair into the FMR1 intron 1 from 5' end of the intron (FIGS. 6A through C). In subjects with FM affected with FXS with cognitive impairment (IQ<70) with no FMR1 mRNA or FMRP and significantly decreased from normal ASFMR1/FMR4 mRNA levels, FREE2 CpG units from FREE2(A) amplicon expanding to CpG unit 1 on the FREE2(E) amplicon have methylation status approaching 100% (FIG. 7C). In these same cell lines the FREE1 and FREE2 regions and the FMR1 promoter were fully methylated. In healthy controls and the 'high functioning' FM males with IQ>70 the same region (FREE2 CpG units from FREE2(A) amplicon expanding to CpG unit 1 on the FREE2(E)) have methylation status approaching 0% (FIGS. 7A and B). The DNA regions 5' of the CpG unit 1 (eg CpG2/3 of FREE2E on the FREE2(E) do not show any difference in methylation between affected FM, healthy controls and 'high functioning full mutation males, as these regions have methylation status approaching 100% in all groups examined (FIG. 7). Therefore, the assay enabled herein clearly identifies the 3' boundary of the FREE2 region and the FMR1 promoter, which is located between CpG unit 1 of the FREE2(E) amplicon and CpG unit 2/3 of the FREE2 (E) amplicon.

[0499] These data also indicate that methylation of the FREE1 region is closely related to inhibition of bi-directional transcription and translation of the FMR locus required for normal neuronal development. As a consequence, this can lead to pathological conditions such as FXS, mental retardation and autism. Hence, the assay examining one or more

biomarker sites herein can be used to diagnose, make a prognosis and detect the presence or predisposition to FXS, and potentially any other neuropathological condition/s associated with elevated methylation and/or altered distribution of methylated sites in the FMR locus described herein.

Example 5

Methylation Status of FREE3 in "High Functioning" Males and FM Carrier Females

[0500] To determine if FREE3 region is informative in biological settings, methylation of FREE3 was examined in blood and lymphoblast DNA samples of 'high functioning' FM males with IQ>70, FMR1 mRNA expression in blood which was 2.6 fold elevated above the normal levels and FMRP expression which was moderately reduced. The methylation results were compared to healthy control males and FXS affected FM males with no FMR1 expression and no FMRP expression and hypermethylated FMR1 promoter encompassing the FMR1 CpG island, FREE1 and FREE2 regions. It is evident from the representations in FIG. 6 that the FREE3 region methylation status is an important biomarker of the FXS phenotype as it is hypermethylated in high functioning FM individuals as well as in healthy controls, while being hypomethylated in FXS affected individuals. It is also evident from the representation that the methylation patterns are consistent between lymphoblasts and blood.

[0501] These data indicate that decreased methylation of the FREE3 region is closely related to inhibition of bi-directional transcription and translation of the FMR locus required from normal neuronal development as ASFMR1 transcription start site is located between CpG units 1 and 2 of the FREE3 region. As a consequence, this can lead to pathological conditions such as FXS, mental retardation and autism. Hence, the assay examining one or more biomarker sites herein can be used to diagnose, make a prognosis and detect the presence or predisposition to FXS, and potentially any other neuropathological condition/s associated with elevated methylation and/or altered distribution of methylated sites in the FMR locus described herein.

[0502] Those skilled in the art will appreciate that aspects of the disclosure described herein are susceptible to variations and modifications other than those specifically described. It is to be understood that these aspects include all such variations and modifications. These aspects also include all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of the steps or features.

BIBLIOGRAPHY

- [0503]** Allingham-Hawkins et al., *Am J Med Genet* 83(4): 322-325, 1999
- [0504]** Bodega et al., *Hum Reprod* 21(4):952-957, 2006
- [0505]** Bonner and Laskey, *Eur. J. Biochem.* 46: 83, 1974
- [0506]** Chiurazzi et al., *Hum Mol Genet* 7(1):109-113, 1998
- [0507]** Coulam, *Fertil Steril* 38(6):645-655, 1982
- [0508]** Dahl et al., *Clin Chem* 53(4):790-793, 2007
- [0509]** de Vries et al., *Am J Hum Genet.* 58:1025-1032, 1996
- [0510]** Fahy et al., *PCR Methods Appl.* 1(1):25-33, 1991
- [0511]** Fleige and Pfaffl, *Mol Aspects Med* 27(2-3):126-139, 2006

- [0512] Francis et al., *Mol Diagn* 5(3):221-225, 2000
 [0513] Fu et al., *Cell* 67(6):1047-1058, 1991
 [0514] Gitan et al., *Genome Res.* 12(1):158-164, 2002
 [0515] Godler et al., *Hum Mol Genet* 10(8):1618-1632, [Epub 2010]
 [0516] Godler et al., *J Mol Diagn*, 2011 [Epub ahead of print]
 [0517] Hagerman et al., *Neurology* 57(1):127-130, 2001
 [0518] Irizarry et al., *Nature Genetics* 41(2):178-186, 2009
 [0519] Irwin et al., *Cereb Cortex* 10(10):1038-1044, 2000
 [0520] Jacquemont et al., *Am Ment Retard* 109(2):154-164, 2004
 [0521] Jacquemont et al., *J Med Genet* 42(2):e14, 2005
 [0522] Jin and Warren, *Hum. Mol. Genet* 9(6):901-908, 2000
 [0523] Jin et al., *Neuron* 39(5):739-747, 2003
 [0524] Kenneson et al., *Hum Mol Genet* 10(14):1449-1454, 2001
 [0525] Khalil et al., *PLoS ONE* 3(1):e1486, 2008
 [0526] Khaniani et al., *Mol Cytogenet* 1(1):5, 2008
 [0527] Ladd et al., *Hum Mol Genet* 16(24):3174-3187, 2007
 [0528] Loesch et al., *Clin Genet* 67(5):412-417, 2005
 [0529] Loesch et al., *J Med Genet* 44(3):200-204, 2007
 [0530] Marmur and Doty, *J. Mol. Biol.* 5: 109, 1962
 [0531] Mitchell et al., *Clin Genet* 67(1):38-46, 2005
 [0532] Nolin et al., *Am J Hum Genet* 72(2):454-464, 2003
 [0533] Nygren et al., *Nucleic Acids Res.* 33(14):e128, 2005
 [0534] Orr and Zoghbi, *Ann Rev Neurosci* 30:575-621, 2007
 [0535] Pieretti et al., *Cell* 66(4):817-822, 1991
 [0536] Pietrobono et al., *Nucleic Acids Res* 30(14):3278-3285, 2002
 [0537] Rein et al., *Nucleic Acids Res.* 26:2255, 1998
 [0538] Sullivan et al., *Hum Reprod* 20(2):402-412, 2005
 [0539] Tassone et al., *J. Mol. Diagn* 10:43-49, 2008
 [0540] Terracciano et al., *Am J fed Genet C Semin Med Genet* 137C(1):32-37, 2005
 [0541] Tost et al., *Nucleic Acids Res* 31(9):e50, 2003
 [0542] Verkerk et al., *Cell* 65(5):905-914, 1991
 [0543] Wojdacz et al., *Nucleic Acids Res.* 35(6):e41, 2007
 [0544] Yegnasubramanian et al., *Nucleic Acids Res.* 34(3):e19, 2006
 [0545] Yu et al., *Science* 252(5010):1179-1181, 1991

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 57

<210> SEQ ID NO 1

<211> LENGTH: 326

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

tttttcttac acaggcattg taaaggatgt tcatgaagat tcaataacag ttgcatttga      60
aaacaagtaa gtgtctcggt atattatattt aatgatgagg ttctttaata ttttatgcta      120
attctattct tcatttttta aaaattcaag tccagtttga gtgcttttca ggaatggatc      180
ttcatgttac tgactgagaa gtttctgaac aactcagtat taaactaatg gaatgactgt      240
ttctgctaata gtcttgaggg tcccttattg tatggtattg atccttacgt cttaattccc      300
ttgaatgtga agaaagaaac cagaga                                           326

```

<210> SEQ ID NO 2

<211> LENGTH: 3556

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

agtaagtgtc tcgttatata attttaatga tgaggttctt taatatTTTA tgctaattct      60
attcttcatt ttttaaaaat tcaagtcag tttgagtgtc tttcaggaat ggatcttcat      120
gttactgact gagaagtttc tgaacaactc agtattaaac taatggaatg actgtttctg      180
ctaagtccct ggaggtcctt tattgtatgg tattgatcct taogtcttaa ttcccttgaa      240
tgtgaagaaa gaaaccagag agtcttgtgt attagtaact ggaagtgcgc ataggagtat      300
ttgtaaaatt taaaaagata gtaatgaaaa gattcatggg tattgctaaa aagttgtatg      360
tgagttcttt aacattaaat ctaaaactga tggttttagt tgtctagtat attttatgga      420
aacagtcatg tttacctatt atctgatttt ttttaatgtt tcatatgttc agcaatagca      480
gtagctgac tttctgttct tcttttgcta gatgaagaaa taaaaacagt cataggccta      540

```

-continued

ggaatattaa	ctgtatgaaa	gcatacatag	tatagatggt	aacattttat	tggagaacac	600
aagtctcttg	actaaatggt	ttgaatgcta	ataaaggcta	ttttcagggt	agctgttggt	660
aagattgtaa	agtacatata	aactccttag	tcaaagtgta	gatgtggcta	tgatcttagg	720
atcttactaa	actctgatgg	atgggttaaca	gttatcattt	ttttggctct	tatataccaa	780
gaaaattaat	aatatatcaa	aagcaggctg	caaactctata	gagacagaaa	gtagattagt	840
gattgcttgt	gctggggctg	gtggggagaa	catagctaaa	gagtacagga	tttttgtgtg	900
tgtggcaatg	aaaatgttgt	aaaatgact	ggtaattatt	gcacataact	gtaaaccctc	960
tataaaccat	tgaattgtac	atttaaattg	atgaattata	atgatatgtg	gattatgtct	1020
gctaaaaatg	agtatgtgac	ccaggcctac	tgagcaataa	atacacacat	cttgcttcta	1080
aagtttaaaa	ccaggcctgt	gtttcagtgt	ctagaaaata	tatttcataa	ttgaacttag	1140
cagattcttt	tggctagcaa	caggacaaa	tagtatcctt	agatgcctga	aacacgtgta	1200
gggggacaga	gataaaagga	ttaaagcat	ctaaccctat	ttctggctct	ggactaataa	1260
tcgcagacca	aaatacaagt	ttgaatagta	gaataccttt	ccctgaatac	ataaagccaa	1320
aagttttagt	atttgtctct	tcgctgttct	atatatccat	ggctatgcaa	tagacaacag	1380
catcacacca	acagaggaat	ctttgataac	ttggagggtg	tttttttctt	acaattatac	1440
actttatcta	cttacttaca	tcactaggta	aatgctgttt	cccataaagg	gagagtgtgc	1500
tttatttgct	actcttttag	tttttctgt	agtcctataat	ctaaagtgtt	aagcaaacac	1560
tcatttctct	tatgcattta	agaatttgtc	tagtttatac	attgatttat	tagatcactt	1620
ctttttaatt	ttcaatcaga	tttctaccat	aaacttaaaa	ttatttcata	tttggaacct	1680
ccaattatct	atctcttgta	ccatatctgc	atgaaagtgt	ctcttcaagc	tctttgtcat	1740
cctgggagtt	tgaaatagat	cattgatctt	cccttggtgt	catggagtat	gtttgcaata	1800
caacactggg	tttgagcaca	tcaataggat	ttaataggtc	tccaccaaag	taatctcttt	1860
gcccttctct	gtgtgtttcc	tttttggttt	tctgttagaa	acaaaacgaa	aacagtgttt	1920
tgtaaaacac	cagattgtat	ttacccttct	tttcctttca	atcaggaaga	agctcctgtt	1980
actgatagca	gattgtttta	ggtgctttgt	ttttagggtc	ttcctaaacc	ttccataaag	2040
cataaagggt	ttttcatatt	tcacatcttt	taataaggaa	ttgaattttt	tatctgtata	2100
cacttaaaat	cattatcttt	tccttggaac	cattgagtta	tatattatta	tatgctttta	2160
aatcaaatct	agcaatgaat	gtataacatc	tattagggtg	gtataataac	tcagaggagg	2220
gttcagttaa	tttaggccct	aatcagattt	ccacaaattc	tgacttaata	tttgcccgct	2280
tatataacag	ctcttcttta	acaaaaacaa	gtacttttct	caatagaatt	ttactaagaa	2340
agctctttag	taaaacatcg	acattataca	tacaacatat	ctcagtatct	gctgatgaag	2400
aacacaaaaa	agaaccacga	tgtgactgct	ccggaagtgt	aatcctcagt	atttttgcaa	2460
agtttgtctt	tcagtatttt	atttgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtgtgtctat	2520
atatatatat	attttttttt	ttttaaagac	aggatctcac	tctgtcacct	aggtcggagt	2580
gcagtggcat	gatcatggct	cactgtaacc	ttgaactcct	gagcttgagc	tatcctccca	2640
cctcagcctc	ccgagttagct	gggactatag	gcacatacca	ctgcacctaa	tttttttttt	2700
tttttaataa	tttgtgttaa	agatcagggtc	ttaccttggt	gcccaggtcg	ctcttgaagt	2760
cctggcctga	agcagtgtct	ccacctcagc	ctcccaaagc	tctgggatta	taggcttgag	2820

-continued

ccaccgcac	ctaataat	atattttt	at	ggatataaaa	aataattt	gg	tatctttcag	2880
agttgtttaa	tatcatttta	aatttaaaaa	cataggcaac	ttaaactcct	ataggctgtc			2940
tccatcggt	ttctgtggt	taggagaccc	caccatccca	gtgcatgtg	ataacgtcat			3000
actgacgag	atccagctac	ccacagcaag	aattgaccac	ctcgtgggat	ctaaaattta			3060
aagggggaaa	agtgagttgt	gaattgctaa	tgtgctgata	gccccatttt	gcttggaat			3120
tagagggcag	ttttgtggt	ccttggaatg	tggttaaaat	tcttctgcaa	gtggaagcat			3180
atttatatta	ctaacaatta	ctggtactaa	tattcaaata	tgaaggaaa	tttctgtgt			3240
ggacttatgt	ttaaagctct	tagaagtga	aattattgga	aagaagactt	gttttgaaaa			3300
tcataatgtt	gctgtattgt	gtttagagaa	atattccaaa	cgggagtagg	ctgctgtgct			3360
gcatgcagac	tttctgaaa	tgttactata	ttgccgttat	gtccactca	gcaaaaactg			3420
atgattttaa	agcttttgtc	tcttgaaact	agaaaaagag	gggtcagcct	taacccaaaag			3480
ttgatggcag	agtggtcatt	atttcagtta	aacatgaaaa	gcatgttaaa	taattgtatg			3540
tttgcttatt	tacagc							3556

<210> SEQ ID NO 3

<211> LENGTH: 9703

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

gtacttggt	ctagggcag	ccccatcttc	gcccttcctt	ccctcccttt	tcttcttgg	60
gtcggcgga	ggcaggccc	gggcccctct	cccagaccc	gcgcctgggt	gccagggcac	120
gctcggcgg	atgttgttg	gaggaagga	ctggacttgg	ggcctgttgg	aagcccctct	180
ccgactccga	gaggccctag	cgcctatcga	aatgagagac	cagcaggag	agggttctct	240
ttcggcgcc	agccccgcc	gggtgagctg	gggatggcg	aggccggcg	gcaggtaacta	300
gagccggcg	ggaaggccg	aaatcggcgc	taagtgcagg	cgatggctta	ttcccccttt	360
cctaacaac	atctcccag	gggatccgg	cctgtcgtgt	gggtagtgt	ggaggagcgg	420
ggggcgctt	agccggccg	cctcctgcag	cgccaagagg	gcttcaggtc	tcctttggct	480
tctcttttc	ggtctagcat	tgggacttcg	gagagctcca	ctgttctggg	cgagggtctgt	540
gaagaaagag	tagtaagaag	cggtagtcgg	caccaaata	caatggcaac	tgatttttag	600
tggcttctct	ttgtggattt	cggaggagat	tttagatcca	aaagtttcag	gaagacccta	660
acatggccca	gcagtgcatt	gaagaagttg	atcatcgtga	atattcgcgt	cccccttttt	720
gttaaaccgg	gtaaattcag	gaatgcacat	gcttcagcgt	ctaaaaccat	tagcagcgt	780
gctacttaaa	aattgtgtgt	gtgtgtttaa	gtttccaaag	acctaaatat	atgccatgaa	840
acttcaggta	attaactgag	agtatattat	tactagggca	tttttttttt	aactgagcga	900
aaatattttt	gtgcccttaa	gaacttgacc	acatttcctt	tgaatttggt	gtgttgcagt	960
ggactgaatt	gttgaggctt	tatataggca	ttcatgggtt	tactgtgctt	tttaaagtta	1020
caccattgca	gatcaactaa	cacctttcag	ttttaaaagg	aagatttaca	aatttgatgt	1080
agcagtagtg	cgtttgttgg	tatgtagggt	ctgtataaat	tcactataaa	attctcattt	1140
ccttttgaat	gtctataacc	tctttcaata	atatcccacc	ttactacagt	attttggcaa	1200
tagaagggtc	gtgtggaagg	aaggctggaa	aatagctatt	agcagtgctc	aacacaattc	1260
ttaaattgtat	tgtagaatgg	cttgaatgtt	tcagacagga	cacgtttggc	tataggaaaa	1320

-continued

taaacaattg actttattct gtgtttacca attttatgaa gacatttga gatcagtata	1380
tttcataaat gagtaaagta tgtaactgt tccatacttt gagcaciaag ataaagcctt	1440
ttgctgtaaa aggaggcaaa aggtaacccc gcgtttatgt tcttaacagt ctcataaata	1500
tgaaattggt tcagttgact ctgcagtcaa aattttaatt tcattgattt tattgatcca	1560
taatttcttc tggtagtgtt gcgtagaatc gttcacggtc ctatagtagt ggttttggtc	1620
actagatttc tggcactaat aactataata catatacata tatatgtgtg agtaacggct	1680
aatggtagg caagattttg attgacctgt gatataaact tagattggat gccactaaag	1740
tttgcttctc acagagggca agtagcacat tatggccttg aagtacttat tgttctcttc	1800
cagcaactta tgatttgctc cagtgatttt gcttgcacac tgactggaat ataagaaatg	1860
ccttctattt ttgctattaa ttccctcctt ttttgttttg ttttgtaacg aagttgttta	1920
acttgaagg gaataagaa taggttggtt gcccttagt tccctgagga gaaatgttaa	1980
tacttgaaca agtgtgtgtc agacaaattg ctgttatgtt tatttaatta agtttgattt	2040
ctaagaaaat ctcaaatggt ctgcactgat ggaagaacag tttctgtaac aaaaaagctt	2100
gaaattttta tatgacttat aatactgctg tgagttttaa aagtaaagca aaagtaaact	2160
gagttgcttg tccagtggga tggacaggaa agatgtgaaa taaaaaccaa tgaataatga	2220
actgctgtgg agaagtgtta catatatgga aaaagaaata ggaaccttgt tcatcaaatt	2280
gatagaaaag cttttaaaac taaacaaatc aaacaacttg agtataatgg aattcagact	2340
ttgatttgcc taacataacc accatatttg caaggacagc tctctatctt ctggtgttta	2400
ttcttaaaaa cttaaaagt agatttagcg atcaccagag ccactacttt tatgcttagg	2460
tatttgtttg acttagaaaa aattgggtcac gtgtaccact ttatagtgcc ctgcagggtg	2520
taagatatga aggcactttg acttacacct cataaaatct ttacaaagta ttttctaaat	2580
gaataatgat gaaataaagt ctttattcta ggtgcactcg cccacataa tttgttttct	2640
ttggactaga agttttgatg tgttgagaat ggtaaatgaat taactccatt ttaaatgtag	2700
aatgctatc actccaatat gaatgcccta atgaatccta agattttagt gttttgtgta	2760
ctagtatgaa aattactaaa gatggaaaaa tcacatgttg gagacataag atacaaacct	2820
ttttgttttc tgaataatac acctctgatt tctgattcct tgttgtaata tgggtgaatt	2880
atactagatt gtaattttgt tgtagatta tactttttta agttcagtggt ttgaggacag	2940
actttcattt ggttagtagt attatggcag ctacagccta aatatgataa agtgatcaat	3000
caaaaggata tttttaatga agatattagt ggtctaacat gtcatttcag atacatagct	3060
gaaatgtagt aaaatcagtt ttactacaaa taaacttgca taagggttat aaatttataa	3120
gtttataaat caacttgggt aaagtgtaaa taaacttgca ctctgggttt ctctgaagtc	3180
tcctgagcta actttgcata aagggtgtat tctgtacttc gaggaagtga attattgggg	3240
tcaaccacat ttttttctct tcctacagtc tgattgcctt ttttagtttt taggatcttt	3300
gtggctgcat catTTTTTccc cttttgaact gtgcattttc taaccccata cttaaatatt	3360
ctcataacct ccaaattatt aattagatgc aacattcagt ggtatattac tggagtttct	3420
gatttctgcc cactatagga atgtgcttcc tgagaagatt gggatcgtga ttataataat	3480
agttaacagg gtagtagtac tttctagggt ccaggcactg ttctctctga tactttattt	3540
gatgtattgt tgttattccc attctttaaa tgatgcacag agaggtagg taagtgactt	3600

-continued

actaccaagt gtcagggcca ttaagggcca ggattctgaa ttctgaaat gatgaaattt	3660
agcttgaaga aattggtttg atttcctgct tagttttcaa tttcatgggtg gtctttgatt	3720
gtattttgtg ctataacact gccttagcat cctataacta tagttacagt gttatattac	3780
cattttttat tgttaataca aagccatcat gaaataattc agtttatgtg ccagcttttt	3840
ttgttactaa ttcttgaacc ttggcgctgt acttcttcat gtggatgcct gtttaaggaaa	3900
gataaagtta gaaatctttg accctgctag gaaatttgc tttgttatat tgggagctca	3960
taaaactgaa gtattcaaaa gttagaatac atacacacaa gaaaaattag taactaattt	4020
aataatgttt tctttgcaca tgtctctgtt gtcttttggc cagagtgaag ctaaatgtgt	4080
ttttcacata attttagacc tatatgaagt cctggacatg tggatgtgtt ggaaggactg	4140
ttgatgaggt ttattgtctc tctttattct tttatgttgt tagtgtcccc atacaacggg	4200
gcggggggag tggggacaaa atgataaact gctttatata tgaagccttg ggtttgaatc	4260
atactgttat cacatttctc atgtccctat cctgtctcct gcagggtttc tttcttgggt	4320
ttcttttttg gttttcttcc ttcctttctt cctttctttt cctttcttcc ctttcttctc	4380
tttttcttcc cttttcttct ttcctttctt tcttctattt ctctcctttg ctctttagt	4440
ttgtttcttt gtttcttttg tttgtttctt tgtttctttg gttgtttctt ttgtttcttt	4500
agtttgtttc tttgtttctt tgtttcttcc aacagggtctc actgtcggcc aggttggagt	4560
gcagtgggtg gatctcagct cactgcaacc tccacctctc gggttcaagc cagtctcatg	4620
cttcagcctt ccgggtagct gagattacag gcattgtgcca ccattcctgg ctattttttg	4680
catttttagt agagatgagg ttctgccatg ttggccaggc tgatctcaag ttatctgcct	4740
gcctcagccc tccaaagtgc tgggaattaca ggcatgagac accgggtcca gcttccagca	4800
ggttttcttt aggagggata ttttacaatg ctgtaagttt ttcctaacga gaattatcat	4860
agcactacat gttctgtctt cagtaagtga tacaagctta ctgatgatgt ttagttatg	4920
ttcattgggtg gtcgggtgta cattgaaact ttaacacata atagcctctc ctgtgagcag	4980
tggttctctg ttgtaagata ctttactaag ggaaggaatg tgagggtctg ctggggagag	5040
tttaccctaa taaggatgga ctttctgtct ttgtttctac agtcttggtg atagaatgtt	5100
tgaatagata gctctaggca ttacatactt tcataaatat gattattgta attacctctt	5160
tggcccagtt gctagtaaat tagggacccc ttaatgattt atttctgtt tattcacctt	5220
gatgaagaac ttgtatctct tttaaactgt actttatcgc ctttctcaaa ttccaagatt	5280
ctcatcacat ttttttctt cccaaactct aaataacctt ttaatattaa gtatctttgt	5340
ggaaacattg ttttcttttt ctatcccaat ttttaaagct tttttaaaaa aaagagtgt	5400
ttgttggga tgtacatttt ccaaatgcaa aaacatttat gattctgtgt ctcttataaa	5460
atatgacact ctctactttt ctctcattta tttagtgcc cctatgtgtg taatttcatt	5520
accacagca gtcttaggag gctggtcag ttccttattt gcagatgagg aatctgaggt	5580
ccagagatca cttcttgggt agagtctcac agctattaag tattagagcc aagattttga	5640
acgtaggctt gattcacagc aaaaccgtta accactaagt aactgactc cagtaagagc	5700
cctagtcctc acccaatata ctttaattcc cctgtgcatt cattcaaatt cattgaattt	5760
gctgcctttg gaaacctctc aggaacctcc tcaacctctc ttctctacag acatcagctt	5820
tgcttgatag gtagggatca tagcaaaaca cagttttcca aggtggtgat aggtggagt	5880
atagtgtctt ggagatggcc aaagaaggaa ggtatgagt tatctgtggg tgggtgagt	5940

-continued

gtggataagg ggaaggacag agccaaaagc gacggctatt ggaaaaacta tgatgagaaa	6000
caggaagatg gaaccttggt ggaattagtg aaagacctta gaattcacag gaggtatttg	6060
tccttcacgc gagtgtgac caaacgtaac ctatgagttt cttttattcc acttattaaa	6120
gcagcaacca aaggtattat atacctctg tattcactta aatgactga tttgaaaaa	6180
gtcatgcaaa catccattta cagataagcc tcattaactc aaaggcagtg gccctgttg	6240
gctctgatga ttactcaaac cattctgac actctgacac tacatccgga aatcatccg	6300
aatgcattc agagcctgca aaagctttt ttttttttt ttttttttt gaatagctac	6360
agcacttgca gatcttcctc ctttgagaga atatttgatt ttaggaaata agcaaatgta	6420
gataagtatg attcaactgg gcaataactt ttaggcaaag agaaaaaac aaaatatagc	6480
aatgtagtaa taaggctgat attatagtgc ttttatgatg agtctgaaaa cagtctccaa	6540
catttgtaaa tgttttaaat cgggtctgtc tactacagta gcccttaagc catatgtagc	6600
tattgagcac ttgacatggt gcaagagtga attggagact gattttaatt ttatttaatt	6660
ttaatgtaag tagtcgcaca tgaccogtg ctactatggt agatggcata gttttagaca	6720
agggcagtag tcttggttag atacttgatt tacttagaac atttctttac ctactgttaa	6780
caaggttcta atagctgatt aaaggacaac atttttagca tgtaatatag agtaaggaa	6840
taatgttaat tactgccaa atgtataaac attatgaaac cttaaacaag gatgaacaca	6900
gaagcagtg cccttctttg taaataagg gtcagttact tcctaatagg tgtcttagtt	6960
ttagttaata atctaatagc acccccaaaa agcaaactgt taatttgta ttagccatcc	7020
tgtaagaca agggagaaat cggggccagg agagtcctct tcccattct ccataattt	7080
tacgacctat agaaataaga gcctaagagt gaccagttcc tgcaacctaa ctcaaaatta	7140
ggcctctctg ttgaagtaat agtgaggcaa acaatccctt gaagcactgg ggcactgcac	7200
cttatgaaat ctaataatc taaaatatca ttggcagtat taagctagag ttctcaaat	7260
atgctgtaga aaatattggt catttaata tagctgagta aaaataagca agatcaaat	7320
gataagaaaa tgttgatttc tccccttttg aaccagtaac taactataag ggtatatacc	7380
catgcttaac ttaaaaaata ttatttagcc accttgggta tagcacaaca tatggatgcc	7440
attatagtcc accttgatct tacaaggaag ctttctttta gcgtagcttt acttttattt	7500
aagcattatt gaagaagctt ggtatctctg ttttaagttgc ttctgtatca gtgttttcca	7560
gggccctgtc tcagcttcca tagttttcct taagagcagc taattaatgc ttcactccat	7620
atgctttaat ttgatattt gggaagtctt atttttttaa ataactttt tatatcatag	7680
gcctttgaag acatatattgt ttacaaagac atctgcattg aaccatttgt ttttaaaaac	7740
atgtatcggg catgtcctat ataccagaaa ctatgccatt tactgaggac accgaagtta	7800
gagaatggtc ttaatccgaa ctagtattg tctcactatc caatatggat atcaaacatt	7860
cagcgtcatg aggatttact actgtgtttt ctcccaagag ttttataatt ttagctctta	7920
tatttaggtc tttgatgcat attcattgat tttttttgt atgcagtatg aggtaaaggt	7980
tgaaacgtat tctttttgca tgtgggtgtc cacttgctcc agcttcattt tttgaaaaa	8040
ctatactttc ccattgaat tgtcttggct tctttgttga caaaggattt attatttctg	8100
ccacaacca gctgtttctc actgtctaaa ccttatgcc tgetgccccg taggcttagt	8160
ggaatgcctc tccatcctcc actctctttt gtttagctag tacttcctta tcccttaagg	8220

-continued

ggtcagtagg aacctcacct tcttcagtat gttttccttt gtcttatcaa cctaccacca	8280
tcacccacac acataccttt tccctaaggc tgggatagga acctcttagg caatgatcta	8340
tttgataagt actcactgag ttcatactag acactctgca gtatatccgt aatgaagta	8400
aaacctgaca ccacctcag aaaacttact gttagtgaga aaatgggcta taataataga	8460
gtgggtaag tttattgtag ggataagcat aggatgctat gagaatgcag aggacaggaa	8520
tttaacctag acttctcagt tctccggtg aagttgacat ctgaactgaa atctggaaga	8580
ccagtaagag atagctatgt aaaaagaggg gaaggacaat aggaagaag gaatggagag	8640
aggcccgaa actacagagt atggcacaag tagtttagca ttgttgggtc acaaattcta	8700
aggatatgaa agataaagt gaagaagtgg tggggctggg gaggacagct cctgtagagc	8760
cttttacagc gggagaagga gtttggtat ggtcctgcag gcaggagaa accactgaag	8820
gggagaaaca tacagaataa atttgagtaa gaagcttaa tctcccaag cctttttaa	8880
ataaattaag atataaagc tttgggttc tttatgctt gtctattct tctaattgtc	8940
caaaacaaaa caaaacctt cctttctgt acctattaa aggttaattt tataaagta	9000
cagacagcat gctgattaaa gaattcttga taattagcta ttttgtctgt ctttgtgtag	9060
attactaaca attgtgtcat cagatttaa agatctaaac ctctggactt tatatatatt	9120
ttctacaagt actgtgagat tgagaactta attcaactct agtaactgga ctttttagtg	9180
ttgtttgcag catatgattg ttaaagaaag tttagtatat ttgtgtgtgc gtatatatat	9240
atatatagat agatagatag atttgaggtt atatttaac attgactatt ggtattctaa	9300
gattttaagg agagaaaggt aggttacaat atatggctga tatatggtc tggacaatt	9360
ttcttagttt tcagtaattc tacctaaat gtacgtccag ctttgactc actaccacat	9420
tgtaatttcc agtatacttg tctatttttc gagatgttga aatctgtgga gaattttttc	9480
tcatgtttag tgttttagct atgtttatct ctagggtata gcacatacaa ggtggcaata	9540
tagatgtgct taatgaataa aatgatcttc aaaaactgac cttcatttga agttctattt	9600
tattcataaa cacataaaac gtttggtatc actgtaaaat ttaactaaaa acaaaaacta	9660
tctttaagct cacaagttaa tttaacgttt tttcttacac agg	9703

<210> SEQ ID NO 4

<211> LENGTH: 497

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

tgggcctggt ggaagccct ctccgactcc gagaggccct agcgccctac gaaatgagag	60
accagcgagg agaggttctc ttccggcgcc gagtccccgc cggggtgagc tgggatgggc	120
gagggccggc ggcaggtact agagccgggc ggggaaggcc gaaatcggcg ctaagtgacg	180
gcgatggctt attccccctt tcctaaacat catctcccag cgggatccgg gcctgtcgtg	240
tgggtagttg tggaggagcg gggggcgctt cagccgggcc gcctcctgca gcgccaagag	300
ggcttcaggt ctcccttggc ttctcttttc cggcttagca ttgggacttc ggagagctcc	360
actgttctgg gcgaggcgct tgaagaaaga gtagtaagaa gcggtagtcg gcaccaaatc	420
acaatggcaa ctgattttta gtggcttctc tttgtggatt tcggaggaga ttttagatcc	480
aaagtttcag gaagacc	497

-continued

<210> SEQ ID NO 5
<211> LENGTH: 303
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```
caagagggct tcaggctctc tttggcttct ctttctctgg tctagcattg ggacttcgga      60
gagctccact gttctgggcg agggctgtga agaaagagta gtaagaagcg gtagtcggca      120
ccaaatcaca atggcaactg atttttagta gcttctcttt gtggatttca gagaagattt      180
tagatccaaa agtttcagaa agaccctaac atggcccagc agtgcattga agaagttgat      240
catcgtgaat attcgcgtcc ccctttttgt taaacgggga aattcaggaa tgcacattgc      300
ttc                                                                    303
```

<210> SEQ ID NO 6
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```
aggaagagag ggtttttttg aaatttttgg attta                                35
```

<210> SEQ ID NO 7
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```
cagtaatacg actcactata gggagaaggc ttaaaccta ttaaaaaccc ctctcc          56
```

<210> SEQ ID NO 8
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```
aggaagagag taagagggtt ttaggttttt tttgg                                35
```

<210> SEQ ID NO 9
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```
cagtaatacg actcactata gggagaaggc taaaacatat acattcctaa atttaccac      59
```

<210> SEQ ID NO 10
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

```
aggaagagag ttttttttat ataggtattt gtaaaggatg                          40
```

<210> SEQ ID NO 11
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

-continued

cagtaatacgc actcactata gggagaaggc ttctctaatt tctttcttca cattcaaaa 59

<210> SEQ ID NO 12
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

ggcgcgtggct 10

<210> SEQ ID NO 13
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

tgaatattcg 10

<210> SEQ ID NO 14
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

aagtttccaa aga 13

<210> SEQ ID NO 15
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

tattattatt 10

<210> SEQ ID NO 16
<211> LENGTH: 6
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

accaca 6

<210> SEQ ID NO 17
<211> LENGTH: 7
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

tgtggtg 7

<210> SEQ ID NO 18
<211> LENGTH: 7
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

tataaat 7

<210> SEQ ID NO 19
<211> LENGTH: 7
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 19

tataaat

7

<210> SEQ ID NO 20

<211> LENGTH: 8

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

aataatat

8

<210> SEQ ID NO 21

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

aaataaaca t

11

<210> SEQ ID NO 22

<211> LENGTH: 7

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

cataaat

7

<210> SEQ ID NO 23

<211> LENGTH: 7

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

tttgttt

7

<210> SEQ ID NO 24

<211> LENGTH: 7

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

tttgttt

7

<210> SEQ ID NO 25

<211> LENGTH: 7

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

ttgttta

7

<210> SEQ ID NO 26

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

tttattta taaagt

16

<210> SEQ ID NO 27

-continued

<211> LENGTH: 7
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

aaacaaa

7

<210> SEQ ID NO 28
 <211> LENGTH: 7
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

tataatt

7

<210> SEQ ID NO 29
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

tttatgctaa tt

12

<210> SEQ ID NO 30
 <211> LENGTH: 1520
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

taacatggcc cagcagtgca ttgaagaagt tgatcatcgt gaattatcgc gtcccccttt	60
ttgtaaacgg ggtaaattca ggaatgcaca tgcttcagcg tctaaaacca ttagcagcgc	120
tgctacttaa aaattgtgtg tgtgtgttta agtttccaaa gacctaaata tatgccatga	180
aacttcaggt aattaactga gagtatatta ttactagggc attttttttt taactgagcg	240
aaaaatattt tgtgccccta agaacttgac cacatttcct tgaattgtgg tgttgacgtg	300
gactgaattg ttgaggcttt atataggcat tcatgggttt actgtgcttt ttaaagttac	360
accattgcag atcaactaac acctttcagt tttaaaagga agatttaca atttgatgta	420
gcagtagtgc gtttgttgtt atgtaggtgc tgtataaatc atctataaat tctcatttcc	480
ttttgaatgt ctataacctc ttcaataata tcccacctta ctacagtatt ttggcaatag	540
aagggtgctg tggaaggaag gctggaaaat agctattagc agtgtccaac acaattctta	600
aatgtattgt agaatggcct gaatgtttca gacaggacac gtttggctat aggaaaataa	660
acaattgact tattctgtgt ttaccaattt tatgaagaca tttggagatc agtatatttc	720
ataaatgagt aaagtatgta aactgttcca tactttgagc acaaagataa agccttttgc	780
tgtaaaaagga ggcaaaagggt aaccccgctt ttatgttctt aacagtctca tgaatatgaa	840
attgtttcag ttgactctca gtcaaaaatt taatttcatt gattttattg atccataatt	900
tcttctgggt agtttgcgta gaatcggtca cggtcctaga ttagtggttt tggtcactag	960
atttctggca ctaataacta taatacatat acatatatat gtgtgagtaa cggctaattg	1020
ttaggcaaga tttgattgac ctgtgatata aacttagatt ggatgccact aaagtttgct	1080
tatcacagag ggcagtagca cattatggcc ttgaagtact tattgttctc ttccagcaac	1140
ttatgatttg ctccagtgat tttgcttgca cactgactgg aatataagaa atgccttcta	1200
tttttgctat taattccctc cttttttgtt ttgttttgta acgaagtgtg ttaacttgaa	1260

-continued

```

ggatgaatgaa gaataggttg gttgcccctt agttcccctga ggagaatggt aatacttgaa 1320
caagtgtgtg tcagacaaat tgctgttatg tttatttaaat taagtgtgat ttctaagaaa 1380
atctcaaagt gtctgcactg atggaagaac agtttctgta acaaaaaagc ttgaaatttt 1440
tatatgactt ataatactgc tgtgagtttt aaaagtaaag caaaagtaaa ctgagttgct 1500
tgtccagtgg gatggacagg 1520

```

```

<210> SEQ ID NO 31
<211> LENGTH: 237
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 31
tcgagatggt gaaatctgtg gagaattttt ctcattgtta gtgttttagc tatgtttatc 60
tctagggtat agcacatata aggtggcaat atagatgtgc ttaatgaata aaatgatctt 120
caaaactgac cttcatttga gttctatttt attcataaac acataaaacg tttggtatca 180
ctgtaaaatt taactaaaaa caaaaactat ctttaagctc acaagttaat ttaacgt 237

```

```

<210> SEQ ID NO 32
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 32
gctcagctcc gtttcggttt cacttcgggt 30

```

```

<210> SEQ ID NO 33
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 33
agccccgcac ttccaccacc agctcctcca 30

```

```

<210> SEQ ID NO 34
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 34
ccgcggaatc ccagaga 17

```

```

<210> SEQ ID NO 35
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 35
cagtggcgtg ggaaatcaa 19

```

```

<210> SEQ ID NO 36
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 36
tgggataacc ggatgca 17

```

-continued

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37

acaccctgtg ccctttaagg

20

<210> SEQ ID NO 38
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

tcaaagctgg gtctgaggaa ag

22

<210> SEQ ID NO 39
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

tcgggatctc aaaatgt

17

<210> SEQ ID NO 40
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

ccccagaatg agaggatgtt g

21

<210> SEQ ID NO 41
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

gccctagatc caccgcttta a

21

<210> SEQ ID NO 42
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

tgctggtgga actc

14

<210> SEQ ID NO 43
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer FREE2 (D)

<400> SEQUENCE: 43

aaaagtttta ggaagatttt aatatgg

27

<210> SEQ ID NO 44
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: artificial sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Reverse primer FREE2 (D)

<400> SEQUENCE: 44

aaaaaacaca ataaacccat aaatacc                27

<210> SEQ ID NO 45
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer FREE2 (E)

<400> SEQUENCE: 45

gaatggtttg aatgttttag ataggat                27

<210> SEQ ID NO 46
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse primer FREE2 (E)

<400> SEQUENCE: 46

accaaaaaatc taataaccaa aaccac                26

<210> SEQ ID NO 47
<211> LENGTH: 327
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: FREE3

<400> SEQUENCE: 47

tttttcttac acaggcattt gtaaaggatg ttcataaga ttcaataaca gttgcatttg    60
aaaacaagta agtgtctcgt tatataattt taatgatgag gttctttaat attttatgct    120
aattctattc ttcatttttt aaaaattcaa gtccagtttg agtgcttttc aggaatggat    180
cttcagtgtt ctgactgaga agtttctgaa caactcagta ttaaaactaat ggaatgactg    240
tttctgctaa tgtctcggag gtcccttatt gtatggtatt gatccttacg tcttaattcc    300
cttgaatgtg aagaaagaaa ccagaga                327

<210> SEQ ID NO 48
<211> LENGTH: 374
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: FREE2 (D)

<400> SEQUENCE: 48

aaaagtttca ggaagaccct aacatggccc agcagtgcac tgaagaagtt gatcatcgtg    60
aatattcgcg tccccctttt tgtaaaccgg ggtaaattca ggaatgcaca tgcttcagcg    120
tctaaaaacca ttagcagcgc tgctaactaa aaattgtgtg tgtgtgttta agtttccaaa    180
gacctaaata tatgccatga aacttcaggt aattaactga gagtatatta ttactagggc    240
attttttttt taactgagcg aaaatatttt tgtgccccta agaacttgac cacatttcct    300
ttgaatttgt ggtgttcgag tggactgaat tgttgaggct ttatataggc attcatgggt    360
ttactgtgct tttt                374

```

-continued

<210> SEQ ID NO 49
<211> LENGTH: 360
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: FREE2 (E)

<400> SEQUENCE: 49

```
gaatggcttg aatgtttcag acaggacacg tttggctata ggaaaataaa caattgactt    60
tattctgtgt ttaccaattt tatgaagaca tttggagatc agtatatttc ataaatgagt    120
aaagtatgta aactgttcca tactttgagc acaaagataa agccttttgc tgtaaaagga    180
ggcaaaagggt aaccccgcggt ttatgttctt aacagtctca tgaatatgaa attgtttcag    240
ttgactctgc agtcaaaatt ttaatttcat tgattttatt gatccataat ttcttctggt    300
gagtttgcgt agaatcggtc acgggtcctag attagtgggt ttggtcacta gattttctggc    360
```

<210> SEQ ID NO 50
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer FREE3

<400> SEQUENCE: 50

```
ttttttttat ataggtattt gtaaaggatg    30
```

<210> SEQ ID NO 51
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer FREE3

<400> SEQUENCE: 51

```
tctctaattt ctttcttcac attcaaaa    28
```

<210> SEQ ID NO 52
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Tag forward primer FREE2 (D)

<400> SEQUENCE: 52

```
aggaagagag    10
```

<210> SEQ ID NO 53
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Tag reverse primer FREE2 (D)

<400> SEQUENCE: 53

```
cagtaatacg actcactata gggagaaggc t    31
```

<210> SEQ ID NO 54
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Tag forward primer FREE2 (E)

-continued

<400> SEQUENCE: 54

aggaagagag

10

<210> SEQ ID NO 55

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Tag reverse primer FREE2 (E)

<400> SEQUENCE: 55

cagtaatacg actcactata gggagaaggc t

31

<210> SEQ ID NO 56

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Tag forward primer FREE3

<400> SEQUENCE: 56

aggaagagag

10

<210> SEQ ID NO 57

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Tag reverse primer FREE3

<400> SEQUENCE: 57

cagtaatacg actcactata gggagaaggc t

31

1. A method for identifying a trinucleotide expansion disorder in a mammalian subject including a human, said method comprising screening for a change relative to a healthy control in the extent of epigenetic modification within (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus; wherein a change in extent of epigenetic modification relative to the control is indicative of the presence or severity of the trinucleotide expansion disorder or a propensity to develop same wherein the intron, intron/exon boundary and/or splicing region is selected from the group consisting of:

(i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

(ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions; and

(iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions.

2. The method of claim 1 wherein the trinucleotide expansion disorder is selected from the group consisting of Fragile X syndrome (FXS), Fragile X-associated tremor or ataxia (FXTAS), Fragile X-associated primary ovarian insufficiency (FXPOI), autism, mental retardation, cognitive impairment, and a modified X-chromosome.

3. The method of claim 1 wherein the epigenetic modification is methylation.

4. The method of claim 1 wherein a cell from the subject is a cultured or uncultured Chorionic Villi Sample (CVS) cell, a lymphoblast cell, a blood cell, buccal cell, epithelial cell, fibroblast cell, an amniocyte or an EBV transformed lymphoblast cell line.

5. The method according to claim 1 further comprising determining the length of (CGG)_n expansion within the FMR genetic locus leading to a (CGG)_n expansion pathology selected from the group consisting of a Gray Zone (GZ) pathology, a premutation (PM) pathology and a full mutation (FM) pathology.

6. The method of claim 1 wherein the trinucleotide expansion disorder is FXS or a related condition in a human subject, and wherein the screening is of an epigenetic modification in the FMR genetic locus at a location selected from the group consisting of (i) and (iii).

7. A method for screening for an agent which modulates epigenetic modification of an FMR genetic locus in a mammalian cell including a human cell, said method comprising screening for a change relative to a healthy control in the extent of epigenetic change in (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region of the FMR genetic locus; wherein the intron, intron/exon boundary and/or splicing region is selected from the group consisting of:

- (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;
- (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions; and
- (iii) genomic FREE2 region as a whole or specific fragments of FREE2 including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

in the presence or absence of an agent to be tested wherein the agent is selected if it induces a change in extent of epigenetic modification.

8. The method of claim 7 wherein the epigenetic modification is methylation.

9.-15. (canceled)

16. The method according to claim 1, said method comprising extracting genomic DNA from a cell from said subject and subjecting the DNA to an amplification reaction using primers selective of (i) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing region; and/or (ii) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the genetic locus.

17. The method of claim 16 wherein the trinucleotide expansion disorder is FXS or a related condition selected from the group consisting of FXTAS, FXPOI, autism, mental retardation, a modified X-chromosome and cognitive impairment.

18. (canceled)

19. A kit for the use in a method of claim 1 comprising primers which amplify a region with the FMR genetic locus, said region in:

Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined

by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;

- (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;
- (iii) genomic FREE2 region as a whole or specific FREE2 fragments including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;
- (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;
- (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and
- (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus.

20. The kit of claim 19 wherein the primers are selected from the group consisting of SEQ ID NOs:6 through 11.

21. A computer program product for assessing progression of a pathological condition associated with the FMR locus in a subject, the product comprising:

- (1) assigning a value to one or more of:
 - (a) change in of methylation or other epigenetic modification relative to a control in FREE3 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;
 - (b) change of methylation or other epigenetic modification relative to a control in intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or portion or part thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;
 - (c) change in methylation or other epigenetic modification relative to a control in genomic FREE2 region as a whole or specific FREE2 fragments including FREE2 (D), FREE2 (E) or FREE3 comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;

- (d) change of methylation in an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;
 - (e) two or more of (i) an intron; (ii) an intron/exon boundary; (iii) a splicing region within the FMR genetic locus;
 - (f) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;
 - (g) length of (CGG)_n expansion within the FMR genetic locus when considered in combination with (a) and/or (b);
 - (h) general phenotype or clinical manifestations in subjects with a neurodevelopmental or neurodegenerative condition;
 - (i) behavioral assessment criteria associated with normal subjects, PM subjects, GZ subjects and FM subjects;
 - (j) cognitive ability;
 - (k) extent of transcription of genes within the FMR locus with the proviso that if any one of (d) through (k) is determined then one or more of (a) through (c) is also determined;
- (2) means to converting the value to a code; and
- (3) means to store the code in a computer readable medium and compare code to a knowledge database to determine whether the code corresponds to a pathological condition.
- 22.** A computer for assessing an association between extent of methylation or other epigenetic modification within the FMR locus, the FMR locus and progression of a disease condition wherein the computer is programmed by the program of claim **21**.
- 23.** A method of identifying epigenetic profile in a population of human subjects indicative of a pathological condition associated with the FMR locus, said method comprising identifying a trinucleotide expansion disorder according to claim **1** in a plurality of members of said population.
- 24.** A method of allowing a user to determine the status, prognosis and/or treatment response of a subject with respect to an FMR locus-associated pathology, the method including:
- (1) receiving data in the form of extent of methylation or other epigenetic modification at a site in:
 - (i) Fragile X-related Epigenetic Element 3 in FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:1 or a homolog thereof or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:1 or which hybridizes to SEQ ID NO:1 or its complementary form under medium stringency conditions;
 - (ii) intron 2 of FMR1 comprising the nucleotide sequence set forth in SEQ ID NO:2 or a homolog thereof or a portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:2 or which hybridizes to SEQ ID NO:2 or its complementary form under medium stringency conditions;
 - (iii) genomic FREE2 region as a whole or specific FREE2 fragments including FREE2 (D), FREE2 (E) or FREES comprising the nucleotide sequence set forth in SEQ ID NO:48 or 49 or 47, respectively or a homolog or portion or fragment thereof defined by having at least 80% nucleotide sequence identity to SEQ ID NO:48 or 49 or 47 or which hybridizes to SEQ ID NO:48 or 49 or 47 or its complementary form under medium stringency conditions;
 - (iv) an intron, intron/exon boundary and/or splicing region downstream of intron 2 of FMR1 or a homolog thereof or a portion or fragment thereof;
 - (v) two or more of (a) an intron; (b) an intron/exon boundary; and/or (c) a splicing regions within the FMR genetic locus; and
 - (vi) approximately one seventh or greater of an intron including an intron/exon boundary and/or a splicing region within the FMR genetic locus;
 wherein the extent of methylation or epigenetic modification provides a correlation to the presence, state, classification or progression of the pathology;
 - (2) transferring the data from the user via a communications network;
 - (3) processing the subject data via multivariate or univariate analysis to provide a disease index value;
 - (4) determining the status of the subject in accordance with the results of the disease index value in comparison with predetermined values; and
 - (5) transferring an indication of the status of the subject to the user via the communications network.

* * * * *