

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 July 2002 (25.07.2002)

PCT

(10) International Publication Number  
**WO 02/057409 A2**

(51) International Patent Classification<sup>7</sup>: C12N

(21) International Application Number: PCT/US01/44216

(22) International Filing Date:  
26 November 2001 (26.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/723,535 27 November 2000 (27.11.2000) US

(71) Applicant (*for all designated States except US*): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BENNETT, C., Frank** [US/US]; 1347 Cassins Street, Carlsbad, CA 92008 (US). **COWSERT, Lex, M.** [US/US]; 1299 Parrott Drive, San Mateo, CA 94402 (US).

(74) Agents: **LICATA, Jane, Massey** et al.; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: ANTISENSE MODULATION OF SRC-2 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of SRC-2. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding SRC-2. Methods of using these compounds for modulation of SRC-2 expression and for treatment of diseases associated with expression of SRC-2 are provided.



WO 02/057409 A2

**ANTISENSE MODULATION OF SRC-2 EXPRESSION****FIELD OF THE INVENTION**

5           The present invention provides compositions and methods for modulating the expression of SRC-2. In particular, this invention relates to antisense compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding human SRC-2. Such  
10 oligonucleotides have been shown to modulate the expression of SRC-2.

**BACKGROUND OF THE INVENTION**

          Steroid, thyroid and retinoid hormones produce a  
15 diverse array of physiologic effects through the regulation of gene expression. Upon entering the cell, these hormones bind to a unique group of intracellular nuclear receptors which have been characterized as ligand-dependent transcription factors. This complex then moves  
20 into the nucleus where the receptor and its cognate ligand interact with the transcription preinitiation complex affecting its stability and ultimately the rate of transcription of the target genes.

          The interactions of the liganded receptor with the  
25 specific elements in the promoter region are mediated by two classes of molecules; corepressors which inhibit transactivation and coactivators which enhance transactivation.

          SRC-2 (also known as steroid receptor coactivator-2, TIF2, GRIP-1 and NcoA-2) is a member of the growing family  
30 of transcription coactivators.

          Disclosed and claimed in PCT Publication WO 98/56806 is a substantially purified nucleic acid molecule, comprising a nucleotide sequence encoding a NCoA-2 (SRC-2)  
35 polypeptide. Generally disclosed are nucleotide

-2-

sequences that bind to a nucleic acid molecule of the invention, such nucleotide sequences being useful, for example, as probes or as antisense molecules, which can inhibit the expression of a nucleic acid molecule encoding a p/CIP. Disclosed and claimed in PCT Publication WO98/02455 is an isolated nucleic acid molecule comprising a polynucleotide encoding the TIF2 (SRC-2) polypeptide or fragments thereof, a polynucleotide having a nucleotide sequence at least 90% identical to the TIF2 (SRC-2) polypeptide or fragments thereof, and a polynucleotide having a nucleotide sequence complementary to said polynucleotides.

Characterization studies of SRC-2 demonstrated that the protein could mediate transcriptional transactivation by steroid receptors including the androgen, estrogen and progesterone receptors (Voegel et al., *Embo J.*, 1996, 15, 3667-3675) as well as non-steroid receptors which include thyroid, retinoid, glucocorticoid and Vitamin-D receptors (Voegel et al., *Embo J.*, 1998, 17, 507-519). Furthermore, expression of the transcript was found in most human tissues including the pancreas, skeletal muscle, liver, lung, placenta, brain and heart, with lower levels in the kidney (Voegel et al., *Embo J.*, 1996, 15, 3667-3675).

SRC-2 has recently been implicated in the altered modulation of gene transcription in a distinct subtype of acute myeloid leukemia (Carapeti et al., *Blood*, 1998, 91, 3127-3133). These studies showed that the chromosomal translocation inv(8)(p11q13) resulted in a fusion protein of MOZ (monocytic leukemia zinc finger protein) to SRC-2. The resulting protein retains the acetyltransferase domain of both proteins and therefore is thought to contribute to the disease process through a mechanism involving abnormal histone acetylation.

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of SRC-2.

-3-

Consequently, there remains a long felt need for additional agents capable of effectively inhibiting SRC-2 function.

It is anticipated that antisense oligonucleotides  
5 against SRC-2 may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the effective and specific modulation of SRC-2 expression.

## 10 SUMMARY OF THE INVENTION

The present invention is directed to antisense compounds, particularly oligonucleotides, which are targeted to a nucleic acid encoding SRC-2, and which modulate the expression of SRC-2. Pharmaceutical and  
15 other compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating the expression of SRC-2 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of  
20 the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of SRC-2 by administering a therapeutically or prophylactically effective amount of one or more of the  
25 antisense compounds or compositions of the invention.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric antisense compounds, particularly oligonucleotides, for use in  
30 modulating the function of nucleic acid molecules encoding SRC-2, ultimately modulating the amount of SRC-2 produced.

This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding SRC-2. As used herein, the terms "target  
35 nucleic acid" and "nucleic acid encoding SRC-2" encompass

-4-

DNA encoding SRC-2, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal  
5 function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be  
10 interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated  
15 by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of SRC-2. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression  
20 of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a  
25 particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose  
30 expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding SRC-2. The targeting process also includes determination of a site or sites  
35 within this gene for the antisense interaction to occur

-5-

such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding SRC-2, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to

-6-

about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice

-7-

sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

10        Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

15        In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds.

20        "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and

25

30

35

-8-

the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

-9-

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides  
5 composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms  
10 because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form  
15 of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 30  
20 nucleobases (i.e. from about 8 to about 30 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 25 nucleobases. As is known in the art, a nucleoside is a base-sugar  
25 combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar  
30 portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to  
35 form a linear polymeric compound. In turn the respective

-10-

ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are  
5 commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds  
10 useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the  
15 backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to  
20 be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates  
25 including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having  
30 normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

-11-

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

-12-

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, **1991**, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular  $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$  [known as a methylene (methyylimino) or MMI backbone],  $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$  and  $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$  [wherein the native phosphodiester backbone is represented as  $-\text{O}-\text{P}-\text{O}-\text{CH}_2-$ ] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl;

-13-

O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>,  
5 O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>,  
10 SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the  
15 pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*,  
20 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>,  
25 also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other  
30 positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar  
35 mimetics such as cyclobutyl moieties in place of the

-14-

pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.:

4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878;  
5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811;  
5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053;  
5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920,

certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azouracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al.,

-15-

*Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B. , ed., CRC Press, 1993.

5 Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-  
10 propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are  
15 presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified  
20 nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121,  
25 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by  
30 reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular

-16-

uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1994**, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, **1992**, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1993**, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, **1992**, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, **1991**, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, **1990**, 259, 327-330; Svinarchuk et al., *Biochimie*, **1993**, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, **1995**, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, **1990**, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, **1995**, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, **1995**, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, **1995**, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, **1996**, 277, 923-937).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469;

-17-

5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098;  
5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475;  
5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142;  
5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923;  
5 5,599,928 and 5,688,941, certain of which are commonly  
owned with the instant application, and each of which is  
herein incorporated by reference.

It is not necessary for all positions in a given  
compound to be uniformly modified, and in fact more than  
10 one of the aforementioned modifications may be  
incorporated in a single compound or even at a single  
nucleoside within an oligonucleotide. The present  
invention also includes antisense compounds which are  
chimeric compounds. "Chimeric" antisense compounds or  
15 "chimeras," in the context of this invention, are  
antisense compounds, particularly oligonucleotides, which  
contain two or more chemically distinct regions, each made  
up of at least one monomer unit, i.e., a nucleotide in the  
case of an oligonucleotide compound. These  
20 oligonucleotides typically contain at least one region  
wherein the oligonucleotide is modified so as to confer  
upon the oligonucleotide increased resistance to nuclease  
degradation, increased cellular uptake, and/or increased  
binding affinity for the target nucleic acid. An  
25 additional region of the oligonucleotide may serve as a  
substrate for enzymes capable of cleaving RNA:DNA or  
RNA:RNA hybrids. By way of example, RNase H is a cellular  
endonuclease which cleaves the RNA strand of an RNA:DNA  
duplex. Activation of RNase H, therefore, results in  
30 cleavage of the RNA target, thereby greatly enhancing the  
efficiency of oligonucleotide inhibition of gene  
expression. Consequently, comparable results can often be  
obtained with shorter oligonucleotides when chimeric  
oligonucleotides are used, compared to phosphorothioate  
35 deoxyoligonucleotides hybridizing to the same target

-18-

region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

5 Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred  
10 to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355;  
15 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through  
20 the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well  
25 known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized in vitro and do not include antisense  
30 compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with  
35 other molecules, molecule structures or mixtures of

-19-

compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the

5 preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804;

10 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass

15 any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for

20 example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that

25 is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE

30 [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 to Imbach et al.

The term "pharmaceutically acceptable salts" refers

35 to physiologically and pharmaceutically acceptable salts

-20-

of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are  
5 formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are  
10 N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form  
15 with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective  
20 salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt  
25 of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically  
30 acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic,  
35 sulfonic, sulfo or phospho acids or N-substituted sulfamic

-21-

acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid,

-22-

gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of SRC-2 is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding SRC-2, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding SRC-2 can be detected by means known in the art.

Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of SRC-2 in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense

-23-

compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

Compositions and formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets.

Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

-24-

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those

-25-

skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### Emulsions

5           The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*,  
10   Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*,  
15   Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases  
20   intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a  
25   water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to  
30   the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed.

-26-

Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex  
5 formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water  
10 stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into  
15 the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of  
20 stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids  
25 (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation  
30 of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker,  
35 Inc., New York, N.Y., 1988, volume 1, p. 199).

-27-

Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations.

Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgate, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

-28-

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth),  
5 cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming  
10 strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides  
15 that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-  
20 hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated  
25 hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for  
30 their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of  
35 ease of formulation, efficacy from an absorption and

-29-

bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

10        In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and  
15        then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are  
20        stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215).  
25        Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the  
30        microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric

-30-

packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

5       The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of  
10       solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-  
20       ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate  
25       (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into  
30       the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art.  
35       The aqueous phase may typically be, but is not limited to,

-31-

water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, **1994**, *11*, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, **1993**, *13*, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, **1994**, *11*, 1385; Ho et al., *J. Pharm. Sci.*, **1996**, *85*, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the

-32-

gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

5           Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and  
10   nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee  
15   *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

          There are many organized surfactant structures  
20   besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the  
25   standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

          Liposomes are unilamellar or multilamellar vesicles  
30   which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as

-33-

efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each  
5 with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes  
10 obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical*  
15 *Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

20 Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As  
25 the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many  
30 drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the

-34-

administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, **1987**, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, **1992**, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions

-35-

generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

-36-

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside  $G_{M1}$ , or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, **1987**, 223, 42; Wu et al., *Cancer Research*, **1993**, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, **1987**, 507, 64) reported the ability of monosialoganglioside  $G_{M1}$ , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, **1988**, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside  $G_{M1}$  or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-*sn*-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

-37-

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et

-38-

al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry *et al.* discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa *et al.* discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman *et al.* describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love *et al.* discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants,

-39-

both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different  
5 surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants  
10 find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters,  
15 propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included  
20 in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include  
25 carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most  
30 important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include

-40-

quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry  
5 either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations  
10 and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

In one embodiment, the present invention employs  
15 various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily  
20 cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration  
25 enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating  
30 non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present  
35 invention, surfactants (or "surface-active agents") are

-41-

chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of  
5 oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic*  
10 *Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include,  
15 for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-  
20 monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and diglycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et  
25 al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

Bile salts: The physiological role of bile includes  
30 the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic

-42-

derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-

-43-

339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), *N*-acyl derivatives of collagen, laureth-9 and *N*-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including

-44-

glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

#### Carriers

5           Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (*i.e.*, does not possess biological activity *per se*) but is  
10       recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid  
15       and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic  
20       acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid  
25       (Miyao *et al.*, *Antisense Res. Dev.*, **1995**, 5, 115-121; Takakura *et al.*, *Antisense & Nucl. Acid Drug Dev.*, **1996**, 6, 177-183).

#### Excipients

          In contrast to a carrier compound, a "pharmaceutical  
30       carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in  
35       mind, so as to provide for the desired bulk, consistency,

-45-

*etc.*, when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (*e.g.*, pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, *etc.*); fillers (*e.g.*, lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, *etc.*); lubricants (*e.g.*, magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, *etc.*); disintegrants (*e.g.*, starch, sodium starch glycolate, *etc.*); and wetting agents (*e.g.*, sodium lauryl sulphate, *etc.*).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

-46-

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense

-47-

mechanism. Examples of such chemotherapeutic agents include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine (CA), 5-fluorouracil (5-FU), floxuridine (5-FUdR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck*  
10 *Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 1206-1228). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to  
15 ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense  
20 chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds,  
25 particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or  
30 sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be  
35 treated, with the course of treatment lasting from several

-48-

days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on  $EC_{50}$ s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

#### EXAMPLES

##### Example 1

##### 30 Nucleoside Phosphoramidites for Oligonucleotide Synthesis Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyl diisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside

-49-

amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was  
5 utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids*  
10 *Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

#### **2'-Fluoro amidites**

##### **2'-Fluorodeoxyadenosine amidites**

2'-fluoro oligonucleotides were synthesized as  
15 described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was  
20 synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-  
25 arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-  
30 dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

##### **2'-Fluorodeoxyguanosine**

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS)

-50-

protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP  
5 to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and  
10 5'-DMT-3'-phosphoramidites.

#### **2'-Fluorouridine**

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was  
15 treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by  
20 selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **25 2'-O-(2-Methoxyethyl) modified amidites**

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

#### **30 2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL).

-51-

The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

**2'-O-Methoxyethyl-5-methyluridine**

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with

-52-

the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-**

5 **methyluridine**

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and  
10 the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70%  
15 product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL). The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was  
20 obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give  
25 a total yield of 183 g (57%).

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a  
30 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of  
35 the reaction, as judged by TLC, MeOH (50 mL) was added and

-53-

the mixture evaporated at 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product).

The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-

-54-

dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and  $\text{NH}_4\text{OH}$  (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel.

MeOH (400 mL) saturated with  $\text{NH}_3$  gas was added and the vessel heated to  $100^\circ\text{C}$  for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in  $\text{CHCl}_3$  (700 mL) and extracted with saturated  $\text{NaHCO}_3$  (2x300 mL) and saturated NaCl (2x300 mL), dried over  $\text{MgSO}_4$  and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5%  $\text{Et}_3\text{NH}$  as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in  $\text{CH}_2\text{Cl}_2$  (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxytetra(isopropyl)phosphite (40.5 mL, 0.123 M) were added

-55-

with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

**2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites**

**2'-(Dimethylaminooxyethoxy) nucleoside amidites**

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

**5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine**

O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22, ethyl acetate) indicated a complete reaction. The

-56-

solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (R<sub>f</sub> 0.67 for desired product and R<sub>f</sub> 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl

-57-

acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g.

5 The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-**

**butyldiphenylsilyl-5-methyluridine**

10 5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The  
15 reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting  
20 deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained  
25 was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

**5'-O-tert-butyldiphenylsilyl-2'-O-[(2-**

30 **formadoximinooxy)ethyl]-5-methyluridine**

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-**

butyldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h

-58-

the mixture was filtered, the filtrate was washed with ice cold  $\text{CH}_2\text{Cl}_2$  and the combined organic phase was washed with water, brine and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy) ethyl]-5-methyluridine as white foam (1.95 g, 78%).

**5'-O-*tert*-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine**

5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium *p*-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in  $\text{CH}_2\text{Cl}_2$ ). Aqueous  $\text{NaHCO}_3$  solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction

-59-

mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

**2'-O-(dimethylaminoxyethyl)-5-methyluridine**

10 Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-*tert*-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

20 **5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine**

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

-60-

5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine  
5 (1.08g, 1.67mmol) was co-evaporated with toluene (20mL).  
To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-  
10 N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated,  
15 then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites]  
25 are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

N<sup>2</sup>-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-  
30 [(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-

-61-

O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine  
5 deaminase. (McGee, D. P. C., Cook, P. D., Guinasso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-  
10 dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may  
15 phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside  
20 amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside  
25 amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL  
30 bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-,2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The

-62-

crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent.

As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour.

The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified

-63-

by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

## Example 2

### Oligonucleotide synthesis

5           Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

10           Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the  
15           capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution.           Phosphinate oligonucleotides are prepared  
20           as described in U.S. Patent 5,508,270, herein incorporated by reference.

          Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

25           3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

          Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent  
30           5,366,878, herein incorporated by reference.

          Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

35           3'-Deoxy-3'-amino phosphoramidate oligonucleotides

- 64 -

are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by  
5 reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### 10 Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino  
15 linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are  
20 prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and  
25 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

### Example 4

#### 30 PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*,  
35 1996, 4, 5-23. They may also be prepared in accordance

-65-

with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

**Example 5****Synthesis of Chimeric Oligonucleotides**

5 Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked  
10 nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are  
15 also known in the art as "hemimers" or "wingmers".

**[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric****Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligo-  
20 nucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The  
25 standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1  
30 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness.  
35 The pellet is resuspended in 1M TBAF in THF for 24 hrs at

-66-

room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-

(Methoxyethyl)] Chimeric Phosphorothioate

10 Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

15 [2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy

Phosphorothioate]--[2'-O-(2-Methoxyethyl)

Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,4,5-benzodithiole-3-one 1,1-dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

30 Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

-67-

**Example 6****Oligonucleotide Isolation**

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* **1991**, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

**Example 7****20 Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or

- 68 -

patented methods. They are utilized as base protected beta-cyanoethyldiisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

10 **Example 8**

**Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

25 **Example 9**

**Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following four cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type

-69-

chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

T-24 cells:

5       The transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were  
10       seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.  
15

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.  
20

A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in  
25       DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely  
30       passaged by trypsinization and dilution when they reached 90% confluence.

NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs  
35       were routinely maintained in Fibroblast Growth Medium

-70-

(Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

HEK cells:

5 Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely  
10 maintained for up to 10 passages as recommended by the supplier.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates,  
15 wells were washed once with 200  $\mu$ L OPTI-MEM<sup>TM</sup>-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM<sup>TM</sup>-1 containing 3.75  $\mu$ g/mL LIPOFECTIN<sup>TM</sup> (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh  
20 medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line,  
25 the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a  
30 phosphorothioate backbone which is targeted to human c-Ha-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCCCAAGGA**, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is

-71-

targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of c-Ha-ras or c-raf mRNA is then utilized in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

**Example 10****Analysis of oligonucleotide inhibition of SRC-2 expression**

Antisense modulation of SRC-2 expression can be assayed in a variety of ways known in the art. For example, SRC-2 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996.

Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be

-72-

"multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated  
5 cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of  
10 dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed  
15 samples, the primer-probe set specific for that target is deemed as multiplexable. Other methods of PCR are also known in the art.

SRC-2 protein levels can be quantitated in a variety of ways well known in the art, such as  
20 immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to SRC-2 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation,  
25 Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc.,  
30 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art  
35 and can be found at, for example, Ausubel, F.M. et al.,

-73-

*Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current*  
5 *Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp.  
10 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

**Example 11****Poly(A)+ mRNA isolation**

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for  
15 poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was  
20 washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was  
25 transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper  
30 towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate

-74-

was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

## 5 Example 12

### Total RNA Isolation

Total mRNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for  
10 cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 µL cold PBS. 100 µL Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100 µL of 70% ethanol was then added to each well and the contents mixed  
15 by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the  
20 RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10  
25 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 µL water into each  
30 well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 µL water.

-75-

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

**Example 13****Real-time Quantitative PCR Analysis of SRC-2 mRNA Levels**

Quantitation of SRC-2 mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye

-76-

from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25 µL PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl<sub>2</sub>, 300 µM each of dATP, dCTP and dGTP, 600 µM of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 µL poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension). SRC-2 probes and primers were designed to hybridize to the human SRC-2 sequence, using published sequence information (GenBank accession number X97674, incorporated herein as SEQ ID NO:3).

For SRC-2 the PCR primers were:  
forward primer: CTCCAGGGCAGAGACAAGAAAG (SEQ ID NO: 4)  
reverse primer: CACGATTACGTTTTTCAGTGTTCTT (SEQ ID NO: 5)  
and the PCR probe was: FAM-TCCTGACCAACTTGGACCCAGCCC-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster

-77-

City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

For GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 7)

5 reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 8) and  
the PCR probe was: 5' JOE-CAAGCTTCCCGTTCTCAGCC- TAMRA 3'  
(SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster  
City, CA) is the fluorescent reporter dye) and TAMRA (PE-  
Applied Biosystems, Foster City, CA) is the quencher dye.

#### 10 **Example 14**

##### **Northern blot analysis of SRC-2 mRNA levels**

Eighteen hours after antisense treatment, cell  
monolayers were washed twice with cold PBS and lysed in 1  
mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total  
15 RNA was prepared following manufacturer's recommended  
protocols. Twenty micrograms of total RNA was  
fractionated by electrophoresis through 1.2% agarose gels  
containing 1.1% formaldehyde using a MOPS buffer system  
(AMRESCO, Inc. Solon, OH). RNA was transferred from the  
20 gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia  
Biotech, Piscataway, NJ) by overnight capillary transfer  
using a Northern/Southern Transfer buffer system (TEL-TEST  
"B" Inc., Friendswood, TX). RNA transfer was confirmed by  
UV visualization. Membranes were fixed by UV cross-  
25 linking using a STRATALINKER™ UV Crosslinker 2400  
(Stratagene, Inc, La Jolla, CA).

Membranes were probed using QUICKHYB™ hybridization  
solution (Stratagene, La Jolla, CA) using manufacturer's  
recommendations for stringent conditions with a SRC-2  
30 specific probe prepared by PCR using the forward primer  
CTCCAGGGCAGAGACAAGAAAG (SEQ ID NO: 4) and the reverse  
primer CACGATTACGTTTTTCAGTGTTCT (SEQ ID NO: 5). To  
normalize for variations in loading and transfer  
efficiency membranes were stripped and probed for

-78-

glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA). Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

**Example 15****Antisense inhibition of SRC-2 expression- phosphorothioate oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human SRC-2 RNA, using published sequences (GenBank accession number X97674, incorporated herein as SEQ ID NO: 3). The oligonucleotides are shown in Table 1. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. X97674), to which the oligonucleotide binds. All compounds in Table 1 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. All cytidine residues are 5-methylcytidines. The compounds were analyzed for effect on SRC-2 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1

Inhibition of SRC-2 mRNA levels by phosphorothioate oligodeoxynucleotides

| ISIS# | REGION      | TARGET | SEQUENCE            | %          | SEQ ID |
|-------|-------------|--------|---------------------|------------|--------|
|       |             | SITE   |                     | Inhibition | NO.    |
| 29929 | 5' UTR      | 71     | atcggctcgggtcgggtca | 62         | 10     |
| 29930 | Start Codon | 151    | actcatcttgaacacata  | 0          | 11     |

-79-

|       |                |      |                      |    |    |
|-------|----------------|------|----------------------|----|----|
| 29931 | Start<br>Codon | 157  | catcccactcatcttgaa   | 0  | 12 |
| 29932 | Start<br>Codon | 164  | tttctcccatcccactca   | 13 | 13 |
| 29933 | Coding         | 225  | ccaagttggtcaggacat   | 0  | 14 |
| 29934 | Coding         | 358  | tgcacatttgtcaggttt   | 82 | 15 |
| 29935 | Coding         | 438  | tgcacttcacatctatgttg | 50 | 16 |
| 29936 | Coding         | 495  | ggccccagcgcaccccttg  | 55 | 17 |
| 29937 | Coding         | 548  | tgccttccaggttcacta   | 1  | 18 |
| 29938 | Coding         | 595  | gttataaccttagatactg  | 16 | 19 |
| 29939 | Coding         | 642  | ccaacatgcaagatgcta   | 17 | 20 |
| 29940 | Coding         | 692  | catttactatagactttg   | 21 | 21 |
| 29941 | Coding         | 778  | ttctgaatcaggtaaagg   | 0  | 22 |
| 29942 | Coding         | 845  | gagagacagcgaagcact   | 71 | 23 |
| 29943 | Coding         | 936  | agaactgggtctttccttc  | 8  | 24 |
| 29944 | Coding         | 1075 | ctgcgcatggaacttctg   | 55 | 25 |
| 29945 | Coding         | 1265 | gaagcatatgtaaagata   | 0  | 26 |
| 29946 | Coding         | 1475 | aaccaccaaacctgcca    | 58 | 27 |
| 29947 | Coding         | 1667 | aaaactgactgggtggga   | 0  | 28 |
| 29948 | Coding         | 1805 | aagccaacgatgacccta   | 80 | 29 |
| 29949 | Coding         | 2060 | tctgccctttgctgtcat   | 41 | 30 |
| 29950 | Coding         | 2162 | gcaagctacctgtggagt   | 50 | 31 |
| 29951 | Coding         | 2371 | attctctttcttcttggg   | 29 | 32 |
| 29952 | Coding         | 2492 | ctattaattttgtgttac   | 0  | 33 |
| 29953 | Coding         | 2613 | gggaaaagctgtggtaat   | 21 | 34 |
| 29954 | Coding         | 2748 | ctctgtgaaattcgcagt   | 0  | 35 |
| 29955 | Coding         | 2944 | ccctaaatttccttggtt   | 18 | 36 |
| 29956 | Coding         | 3063 | ttcatggcactggtggta   | 15 | 37 |
| 29957 | Coding         | 3342 | ggacatagcaagtcacat   | 38 | 38 |
| 29958 | Coding         | 3552 | tgctgtgggaaaacgggc   | 57 | 39 |
| 29959 | Coding         | 3746 | gctgtgcttgaggcgat    | 57 | 40 |
| 29960 | Coding         | 3872 | ggatttccctctgtctct   | 68 | 41 |
| 29961 | Coding         | 4012 | agggttgcacatagttgc   | 39 | 42 |
| 29962 | Coding         | 4182 | gctgggttggcctgagac   | 0  | 43 |
| 29963 | Coding         | 4407 | gtaggcacggaggtcact   | 0  | 44 |
| 29964 | Stop<br>Codon  | 4539 | cagcaatatattccgtgtt  | 12 | 45 |
| 29965 | Stop<br>Codon  | 4549 | tcagcagtgtcagcaata   | 0  | 46 |
| 29966 | Stop<br>Codon  | 4555 | ctggcttcagcagtgta    | 28 | 47 |
| 29967 | 3' UTR         | 4695 | gacgccaccctgggaacc   | 26 | 48 |
| 29968 | 3' UTR         | 4828 | ctgtcaacatttacttgg   | 0  | 49 |

As shown in Table 1, SEQ ID NOS 10, 15, 16, 17, 23, 25, 27, 29, 30, 31, 32, 38, 39, 40, 41, 42, 47 and 48

-80-

demonstrated at least 25% inhibition of SRC-2 expression in this assay and are therefore preferred.

#### Example 16:

#### Antisense inhibition of SRC-2 expression- phosphorothioate

#### 5 2'-MOE gapmer oligonucleotides

In accordance with the present invention, a second series of oligonucleotides targeted to human SRC-2 were synthesized. The oligonucleotide sequences are shown in Table 2. Target sites are indicated by nucleotide  
10 numbers, as given in the sequence source reference (Genbank accession no. X97674, incorporated herein as SEQ ID NO: 3), to which the oligonucleotide binds.

All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length,  
15 composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate  
20 (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from two experiments. If present, "N.D." indicates "no data".

25

Table 2

Inhibition of SRC-2 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and  
a deoxy gap

30

| ISIS# | REGION         | TARGET | SEQUENCE            | %          | SEQ<br>ID<br>NO. |
|-------|----------------|--------|---------------------|------------|------------------|
|       |                | SITE   |                     | Inhibition |                  |
| 29969 | 5' UTR         | 71     | atcggctcgggtcgggtca | 44         | 10               |
| 29970 | Start<br>Codon | 151    | actcatcttgaacacata  | 15         | 11               |

- 81 -

|       |                |      |                     |    |    |
|-------|----------------|------|---------------------|----|----|
| 29971 | Start<br>Codon | 157  | catcccactcatcttgaa  | 35 | 12 |
| 29972 | Start<br>Codon | 164  | tttctcccatcccactca  | 51 | 13 |
| 29973 | Coding         | 225  | ccaagttggtcaggacat  | 26 | 14 |
| 29974 | Coding         | 358  | tgcacatttgtcaggttt  | 85 | 15 |
| 29975 | Coding         | 438  | tgcacttcatctatgttg  | 85 | 16 |
| 29976 | Coding         | 495  | ggccccagcgcacccctg  | 66 | 17 |
| 29977 | Coding         | 548  | tgccttccagggttacta  | 86 | 18 |
| 29978 | Coding         | 595  | gttataccttagatactg  | 2  | 19 |
| 29979 | Coding         | 642  | ccaacatgcaagatgcta  | 77 | 20 |
| 29980 | Coding         | 692  | catttactatagactttg  | 6  | 21 |
| 29981 | Coding         | 778  | ttctgaatcaggtaaagg  | 58 | 22 |
| 29982 | Coding         | 845  | gagagacagcgaagcact  | 57 | 23 |
| 29983 | Coding         | 936  | agaactgggtctttccttc | 0  | 24 |
| 29984 | Coding         | 1075 | ctgcgcatggaacttctg  | 69 | 25 |
| 29985 | Coding         | 1265 | gaagcatatgtaaagata  | 0  | 26 |
| 29986 | Coding         | 1475 | aaccaccaaacctgccca  | 21 | 27 |
| 29987 | Coding         | 1667 | aaaactgactgggtggga  | 27 | 28 |
| 29988 | Coding         | 1805 | aagccaacgatgacccta  | 6  | 29 |
| 29989 | Coding         | 2060 | tctgccctttgctgtcat  | 77 | 30 |
| 29990 | Coding         | 2162 | gcaagctacctgtggagt  | 81 | 31 |
| 29991 | Coding         | 2371 | attctctttcttcttggg  | 16 | 32 |
| 29992 | Coding         | 2492 | ctattaattttgtgttac  | 16 | 33 |
| 29993 | Coding         | 2613 | gggaaaagctgtggtaat  | 0  | 34 |
| 29994 | Coding         | 2748 | ctctgtgaaattcgcagt  | 5  | 35 |
| 29995 | Coding         | 2944 | ccctaaatttccttggtt  | 34 | 36 |
| 29996 | Coding         | 3063 | ttcatggcactggtggta  | 77 | 37 |
| 29997 | Coding         | 3342 | ggacatagcaagtcacat  | 4  | 38 |
| 29998 | Coding         | 3552 | tgctgtgggaaaacgggc  | 41 | 39 |
| 29999 | Coding         | 3746 | gctgtgcttgaggcgat   | 69 | 40 |
| 30000 | Coding         | 3872 | ggatttccctctgtctct  | 58 | 41 |
| 30001 | Coding         | 4012 | agggttgctcatagttgc  | 29 | 42 |
| 30002 | Coding         | 4182 | gctgggttggcctgagac  | 58 | 43 |
| 30003 | Coding         | 4407 | gtaggcacggaggtcact  | 71 | 44 |
| 30004 | Stop<br>Codon  | 4539 | cagcaatatttccgtgtt  | 52 | 45 |
| 30005 | Stop<br>Codon  | 4549 | tcagcagtgtcagcaata  | 41 | 46 |
| 30006 | Stop<br>Codon  | 4555 | ctggcttcagcagtgtca  | 65 | 47 |
| 30007 | 3' UTR         | 4695 | gacgccaccctgggaacc  | 5  | 48 |
| 30008 | 3' UTR         | 4828 | ctgtcaacatttacttgg  | 0  | 49 |

As shown in Table 2, SEQ ID NOS 10, 12, 13, 15, 16, 17, 18, 20, 22, 23, 25, 30, 31, 36, 37, 39, 40, 41, 43,

- 82 -

44, 45, 46 and 47 demonstrated at least 30% inhibition of SRC-2 expression in this experiment and are therefore preferred.

**Example 17****5 Western blot analysis of SRC-2 protein levels**

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for  
10 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to SRC-2 is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary  
15 antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

-83-

**What is claimed is:**

1. An antisense compound 8 to 30 nucleobases in length targeted to the 5'-untranslated region, start  
5 codon, coding region, stop codon or 3'-untranslated region of a nucleic acid molecule encoding human SRC-2, wherein said antisense compound specifically hybridizes with and inhibits the expression of human SRC-2.

2. The antisense compound of claim 1 which is an  
10 antisense oligonucleotide.

3. The antisense compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 10, 15, 16, 17, 23, 25, 27, 29, 30, 31, 32, 38, 39, 40, 41, 42, 47, 48, 12, 13, 18, 20, 22, 36, 37, 43, 44, 45  
15 or 46.

4. The antisense compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 10, 15, 16, 17, 23, 25, 30, 31, 39, 40, 41 or 47.

5. The antisense compound of claim 2 wherein the  
20 antisense oligonucleotide comprises at least one modified internucleoside linkage.

6. The antisense compound of claim 5 wherein the modified internucleoside linkage is a phosphorothioate linkage.

25 7. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

8. The antisense compound of claim 7 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

30 9. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

10. The antisense compound of claim 9 wherein the modified nucleobase is a 5-methylcytosine.

11. The antisense compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

12. A pharmaceutical composition comprising the  
5 antisense compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The pharmaceutical composition of claim 12 further comprising a colloidal dispersion system.

14. The pharmaceutical composition of claim 12  
10 wherein the antisense compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of SRC-2 in human cells or tissues comprising contacting said cells or tissues with the antisense compound of claim 1 so that  
15 expression of SRC-2 is inhibited.

16. A method of treating a human having a disease or condition associated with SRC-2 comprising administering to said animal a therapeutically or prophylactically effective amount of the antisense  
20 compound of claim 1 so that expression of SRC-2 is inhibited.

17. The method of claim 16 wherein the disease or condition is a hormonal disorder.

18. The method of claim 16 wherein the disease or  
25 condition is an inflammatory disorder.

19. The method of claim 16 wherein the disease or condition is a hyperproliferative disorder.

20. The method of claim 19 wherein the hyperproliferative disorder is cancer.

## SEQUENCE LISTING

<110> Isis Pharmaceuticals, Inc.  
C. Frank Bennett  
Lex M. Cowser

<120> ANTISENSE MODULATION OF SRC-2 EXPRESSION

<130> RTS-0226

<150> 09/723,535

<151> 2000-11/27

<160> 49

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctcctcaggg

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2

atgcattctg cccccaagga

20

<210> 3

<211> 6156

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (163)..(4557)

<400> 3

ggcggccgca gcctcggcta cagcttcggc ggcgaaggtc agcgccgacg gcagccggca 60

cctgacggcg tgaccgaccc gagccgattt ctcttggtt tggctacaca cttatagatc 120

ttctgcactg ttacaggca cagttgctga tatgtgttca ag atg agt ggg atg 174  
Met Ser Gly Met

1

gga gaa aat acc tct gac ccc tcc agg gca gag aca aga aag cgc aag 222

|          |     |     |     |     |           |     |     |     |     |           |     |     |     |     |           |     |
|----------|-----|-----|-----|-----|-----------|-----|-----|-----|-----|-----------|-----|-----|-----|-----|-----------|-----|
| Gly<br>5 | Glu | Asn | Thr | Ser | Asp<br>10 | Pro | Ser | Arg | Ala | Glu<br>15 | Thr | Arg | Lys | Arg | Lys<br>20 |     |
| gaa      | tgt | cct | gac | caa | ctt       | gga | ccc | agc | ccc | aaa       | agg | aac | act | gaa | aaa       | 270 |
| Glu      | Cys | Pro | Asp | Gln | Leu       | Gly | Pro | Ser | Pro | Lys       | Arg | Asn | Thr | Glu | Lys       |     |
|          |     |     |     | 25  |           |     |     |     | 30  |           |     |     |     | 35  |           |     |
| cgt      | aat | cgt | gaa | cag | gaa       | aat | aaa | tat | ata | gaa       | gaa | ctt | gca | gag | ttg       | 318 |
| Arg      | Asn | Arg | Glu | Gln | Glu       | Asn | Lys | Tyr | Ile | Glu       | Glu | Leu | Ala | Glu | Leu       |     |
|          |     |     | 40  |     |           |     |     | 45  |     |           |     |     | 50  |     |           |     |
| att      | ttt | gca | aat | ttt | aat       | gat | ata | gac | aac | ttt       | aac | ttc | aaa | cct | gac       | 366 |
| Ile      | Phe | Ala | Asn | Phe | Asn       | Asp | Ile | Asp | Asn | Phe       | Asn | Phe | Lys | Pro | Asp       |     |
|          |     | 55  |     |     |           |     | 60  |     |     |           |     | 65  |     |     |           |     |
| aaa      | tgt | gca | atc | tta | aaa       | gaa | act | gtg | aag | caa       | att | cgt | cag | atc | aaa       | 414 |
| Lys      | Cys | Ala | Ile | Leu | Lys       | Glu | Thr | Val | Lys | Gln       | Ile | Arg | Gln | Ile | Lys       |     |
|          | 70  |     |     |     |           | 75  |     |     |     |           | 80  |     |     |     |           |     |
| gaa      | caa | gag | aaa | gca | gca       | gct | gcc | aac | ata | gat       | gaa | gtg | cag | aag | tca       | 462 |
| Glu      | Gln | Glu | Lys | Ala | Ala       | Ala | Ala | Asn | Ile | Asp       | Glu | Val | Gln | Lys | Ser       |     |
|          | 85  |     |     |     | 90        |     |     |     |     | 95        |     |     |     |     | 100       |     |
| gat      | gta | tcc | tct | aca | ggg       | cag | ggg | gtc | atc | gac       | aag | gat | gcg | ctg | ggg       | 510 |
| Asp      | Val | Ser | Ser | Thr | Gly       | Gln | Gly | Val | Ile | Asp       | Lys | Asp | Ala | Leu | Gly       |     |
|          |     |     |     | 105 |           |     |     |     | 110 |           |     |     |     | 115 |           |     |
| cct      | atg | atg | ctt | gag | gcc       | ctt | gat | ggg | ttc | ttc       | ttt | gta | gtg | aac | ctg       | 558 |
| Pro      | Met | Met | Leu | Glu | Ala       | Leu | Asp | Gly | Phe | Phe       | Phe | Val | Val | Asn | Leu       |     |
|          |     |     | 120 |     |           |     |     | 125 |     |           |     |     | 130 |     |           |     |
| gaa      | ggc | aac | gtt | gtg | ttt       | gtg | tca | gag | aat | gtg       | aca | cag | tat | cta | agg       | 606 |
| Glu      | Gly | Asn | Val | Val | Phe       | Val | Ser | Glu | Asn | Val       | Thr | Gln | Tyr | Leu | Arg       |     |
|          |     | 135 |     |     |           |     | 140 |     |     |           |     | 145 |     |     |           |     |
| tat      | aac | caa | gaa | gag | ctg       | atg | aac | aaa | agt | gta       | tat | agc | atc | ttg | cat       | 654 |
| Tyr      | Asn | Gln | Glu | Glu | Leu       | Met | Asn | Lys | Ser | Val       | Tyr | Ser | Ile | Leu | His       |     |
|          | 150 |     |     |     |           | 155 |     |     |     |           | 160 |     |     |     |           |     |
| gtt      | ggg | gac | cac | acg | gaa       | ttt | gtc | aaa | aac | ctg       | ctg | cca | aag | tct | ata       | 702 |
| Val      | Gly | Asp | His | Thr | Glu       | Phe | Val | Lys | Asn | Leu       | Leu | Pro | Lys | Ser | Ile       |     |
|          | 165 |     |     |     | 170       |     |     |     |     | 175       |     |     |     |     | 180       |     |
| gta      | aat | ggg | gga | tct | tgg       | tct | ggc | gaa | cct | ccg       | agg | cgg | aac | agc | cat       | 750 |
| Val      | Asn | Gly | Gly | Ser | Trp       | Ser | Gly | Glu | Pro | Pro       | Arg | Arg | Asn | Ser | His       |     |
|          |     |     |     | 185 |           |     |     |     | 190 |           |     |     |     | 195 |           |     |
| acc      | ttc | aat | tgt | cgg | atg       | ctg | gta | aaa | cct | tta       | cct | gat | tca | gaa | gag       | 798 |
| Thr      | Phe | Asn | Cys | Arg | Met       | Leu | Val | Lys | Pro | Leu       | Pro | Asp | Ser | Glu | Glu       |     |
|          |     | 200 |     |     |           |     |     | 205 |     |           |     |     | 210 |     |           |     |
| gag      | ggg | cat | gat | aac | cag       | gaa | gct | cat | cag | aaa       | tat | gaa | act | atg | cag       | 846 |
| Glu      | Gly | His | Asp | Asn | Gln       | Glu | Ala | His | Gln | Lys       | Tyr | Glu | Thr | Met | Gln       |     |
|          |     | 215 |     |     |           |     | 220 |     |     |           |     | 225 |     |     |           |     |
| tgc      | ttc | gct | gtc | tct | caa       | cca | aag | tcc | atc | aaa       | gaa | gaa | gga | gaa | gat       | 894 |
| Cys      | Phe | Ala | Val | Ser | Gln       | Pro | Lys | Ser | Ile | Lys       | Glu | Glu | Gly | Glu | Asp       |     |
|          | 230 |     |     |     |           | 235 |     |     |     |           |     | 240 |     |     |           |     |

3

|                                                                 |      |
|-----------------------------------------------------------------|------|
| ttg cag tcc tgc ttg att tgc gtg gca aga aga gtt ccc atg aag gaa | 942  |
| Leu Gln Ser Cys Leu Ile Cys Val Ala Arg Arg Val Pro Met Lys Glu |      |
| 245 250 255 260                                                 |      |
| aga cca gtt ctt ccc tca tca gaa agt ttt act act cgc cag gat ctc | 990  |
| Arg Pro Val Leu Pro Ser Ser Glu Ser Phe Thr Thr Arg Gln Asp Leu |      |
| 265 270 275                                                     |      |
| caa ggc aag atc acg tct ctg gat acc agc acc atg aga gca gcc atg | 1038 |
| Gln Gly Lys Ile Thr Ser Leu Asp Thr Ser Thr Met Arg Ala Ala Met |      |
| 280 285 290                                                     |      |
| aaa cca ggc tgg gag gac ctg gta aga agg tgt att cag aag ttc cat | 1086 |
| Lys Pro Gly Trp Glu Asp Leu Val Arg Arg Cys Ile Gln Lys Phe His |      |
| 295 300 305                                                     |      |
| gcg cag cat gaa gga gaa tct gtg tcc tat gct aag agg cat cat cat | 1134 |
| Ala Gln His Glu Gly Glu Ser Val Ser Tyr Ala Lys Arg His His His |      |
| 310 315 320                                                     |      |
| gaa gta ctg aga caa gga ttg gca ttc agt caa atc tat cgt ttt tcc | 1182 |
| Glu Val Leu Arg Gln Gly Leu Ala Phe Ser Gln Ile Tyr Arg Phe Ser |      |
| 325 330 335 340                                                 |      |
| ttg tct gat ggc act ctt gtt gct gca caa acg aag agc aaa ctc atc | 1230 |
| Leu Ser Asp Gly Thr Leu Val Ala Ala Gln Thr Lys Ser Lys Leu Ile |      |
| 345 350 355                                                     |      |
| cgt tct cag act act aat gaa cct caa ctt gta ata tct tta cat atg | 1278 |
| Arg Ser Gln Thr Thr Asn Glu Pro Gln Leu Val Ile Ser Leu His Met |      |
| 360 365 370                                                     |      |
| ctt cac aga gag cag aat gtg tgt gtg atg aat ccg gat ctg act gga | 1326 |
| Leu His Arg Glu Gln Asn Val Cys Val Met Asn Pro Asp Leu Thr Gly |      |
| 375 380 385                                                     |      |
| caa acg atg ggg aag cca ctg aat cca att agc tct aac agc cct gcc | 1374 |
| Gln Thr Met Gly Lys Pro Leu Asn Pro Ile Ser Ser Asn Ser Pro Ala |      |
| 390 395 400                                                     |      |
| cat cag gcc ctg tgc agt ggg aac cca ggt cag gac atg acc ctc agt | 1422 |
| His Gln Ala Leu Cys Ser Gly Asn Pro Gly Gln Asp Met Thr Leu Ser |      |
| 405 410 415 420                                                 |      |
| agc aat ata aat ttt ccc ata aat ggc cca aag gaa caa atg ggc atg | 1470 |
| Ser Asn Ile Asn Phe Pro Ile Asn Gly Pro Lys Glu Gln Met Gly Met |      |
| 425 430 435                                                     |      |
| ccc atg ggc agg ttt ggt ggt tct ggg gga atg aac cat gtg tca ggc | 1518 |
| Pro Met Gly Arg Phe Gly Gly Ser Gly Gly Met Asn His Val Ser Gly |      |
| 440 445 450                                                     |      |
| atg caa gca acc act cct cag ggt agt aac tat gca ctc aaa atg aac | 1566 |
| Met Gln Ala Thr Thr Pro Gln Gly Ser Asn Tyr Ala Leu Lys Met Asn |      |
| 455 460 465                                                     |      |
| agc ccc tca caa agc agc cct ggc atg aat cca gga cag ccc acc tcc | 1614 |
| Ser Pro Ser Gln Ser Ser Pro Gly Met Asn Pro Gly Gln Pro Thr Ser |      |
| 470 475 480                                                     |      |

|                                                                 |      |
|-----------------------------------------------------------------|------|
| atg ctt tca cca agg cat cgc atg agc cct gga gtg gct ggc agc cct | 1662 |
| Met Leu Ser Pro Arg His Arg Met Ser Pro Gly Val Ala Gly Ser Pro |      |
| 485 490 495 500                                                 |      |
| cga atc cca ccc agt cag ttt tcc cct gca gga agc ttg cat tcc cct | 1710 |
| Arg Ile Pro Pro Ser Gln Phe Ser Pro Ala Gly Ser Leu His Ser Pro |      |
| 505 510 515                                                     |      |
| gtg gga gtt tgc agc agc aca gga aat agc cat agt tat acc aac agc | 1758 |
| Val Gly Val Cys Ser Ser Thr Gly Asn Ser His Ser Tyr Thr Asn Ser |      |
| 520 525 530                                                     |      |
| tcc ctc aat gca ctt cag gcc ctc agc gag ggg cac ggg gtc tca tta | 1806 |
| Ser Leu Asn Ala Leu Gln Ala Leu Ser Glu Gly His Gly Val Ser Leu |      |
| 535 540 545                                                     |      |
| ggg tca tcg ttg gct tca cca gac cta aaa atg ggc aat ttg caa aac | 1854 |
| Gly Ser Ser Leu Ala Ser Pro Asp Leu Lys Met Gly Asn Leu Gln Asn |      |
| 550 555 560                                                     |      |
| tcc cca gtt aat atg aat cct ccc cca ctc agc aag atg gga agc ttg | 1902 |
| Ser Pro Val Asn Met Asn Pro Pro Pro Leu Ser Lys Met Gly Ser Leu |      |
| 565 570 575 580                                                 |      |
| gac tca aaa gac tgt ttt gga cta tat ggg gag ccc tct gaa ggt aca | 1950 |
| Asp Ser Lys Asp Cys Phe Gly Leu Tyr Gly Glu Pro Ser Glu Gly Thr |      |
| 585 590 595                                                     |      |
| act gga caa gca gag agc agc tgc cat cct gga gag caa aag gaa aca | 1998 |
| Thr Gly Gln Ala Glu Ser Ser Cys His Pro Gly Glu Gln Lys Glu Thr |      |
| 600 605 610                                                     |      |
| aat gac ccc aac ctg ccc ccg gcc gtg agc agt gag aga gct gac ggg | 2046 |
| Asn Asp Pro Asn Leu Pro Pro Ala Val Ser Ser Glu Arg Ala Asp Gly |      |
| 615 620 625                                                     |      |
| cag agc aga ctg cat gac agc aaa ggg cag acc aaa ctc ctg cag ctg | 2094 |
| Gln Ser Arg Leu His Asp Ser Lys Gly Gln Thr Lys Leu Leu Gln Leu |      |
| 630 635 640                                                     |      |
| ctg acc acc aaa tct gat cag atg gag ccc tcg ccc tta gcc agc tct | 2142 |
| Leu Thr Thr Lys Ser Asp Gln Met Glu Pro Ser Pro Leu Ala Ser Ser |      |
| 645 650 655 660                                                 |      |
| ttg tcg gat aca aac aaa gac tcc aca ggt agc ttg cct ggt tct ggg | 2190 |
| Leu Ser Asp Thr Asn Lys Asp Ser Thr Gly Ser Leu Pro Gly Ser Gly |      |
| 665 670 675                                                     |      |
| tct aca cat gga acc tcg ctc aag gag aag cat aaa att ttg cac aga | 2238 |
| Ser Thr His Gly Thr Ser Leu Lys Glu Lys His Lys Ile Leu His Arg |      |
| 680 685 690                                                     |      |
| ctc ttg cag gac agc agt tcc cct gtg gac ttg gcc aag tta aca gca | 2286 |
| Leu Leu Gln Asp Ser Ser Ser Pro Val Asp Leu Ala Lys Leu Thr Ala |      |
| 695 700 705                                                     |      |
| gaa gcc aca ggc aaa gac ctg agc cag gag tcc agc agc aca gct cct | 2334 |
| Glu Ala Thr Gly Lys Asp Leu Ser Gln Glu Ser Ser Ser Thr Ala Pro |      |

| 710                                                                                                                                                   | 715                                                       | 720                                                           |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------|---------------------------------------------------------------|------|
| gga tca gaa gtg act att<br>Gly Ser Glu Val Thr<br>725                                                                                                 | aaa caa gag ccg<br>Lys Gln Glu Pro<br>730                 | gtg agc ccc aag aag aaa<br>Val Ser Pro Lys Lys Lys<br>735 740 | 2382 |
| gag aat gca cta ctt cgc tat ttg cta gat<br>Glu Asn Ala Leu Leu Arg Tyr Leu Leu Asp<br>745 750                                                         | aaa gat gat act aaa gat<br>Lys Asp Asp Thr Lys Asp<br>755 |                                                               | 2430 |
| att ggt tta cca gaa ata acc ccc aaa ctt gag aga ctg gac agt aag<br>Ile Gly Leu Pro Glu Ile Thr Pro Lys Leu Glu Arg Leu Asp Ser Lys<br>760 765 770     |                                                           |                                                               | 2478 |
| aca gat cct gcc agt aac aca aaa tta ata gca atg aaa act gag aag<br>Thr Asp Pro Ala Ser Asn Thr Lys Leu Ile Ala Met Lys Thr Glu Lys<br>775 780 785     |                                                           |                                                               | 2526 |
| gag gag atg agc ttt gag cct ggt gac cag cct ggc agt gag ctg gac<br>Glu Glu Met Ser Phe Glu Pro Gly Asp Gln Pro Gly Ser Glu Leu Asp<br>790 795 800     |                                                           |                                                               | 2574 |
| aac ttg gag gag att ttg gat gat ttg cag aat agt caa tta cca cag<br>Asn Leu Glu Glu Ile Leu Asp Asp Leu Gln Asn Ser Gln Leu Pro Gln<br>805 810 815 820 |                                                           |                                                               | 2622 |
| ctt ttc cca gac acg agg cca ggc gcc cct gct gga tca gtt gac aag<br>Leu Phe Pro Asp Thr Arg Pro Gly Ala Pro Ala Gly Ser Val Asp Lys<br>825 830 835     |                                                           |                                                               | 2670 |
| caa gcc atc atc aat gac ctc atg caa ctc aca gct gaa aac agc cct<br>Gln Ala Ile Ile Asn Asp Leu Met Gln Leu Thr Ala Glu Asn Ser Pro<br>840 845 850     |                                                           |                                                               | 2718 |
| gtc aca cct gtt gga gcc cag aaa aca gca ctg cga att tca cag agc<br>Val Thr Pro Val Gly Ala Gln Lys Thr Ala Leu Arg Ile Ser Gln Ser<br>855 860 865     |                                                           |                                                               | 2766 |
| act ttt aat aac cca cga cca ggg caa ctg ggc agg tta ttg cca aac<br>Thr Phe Asn Asn Pro Arg Pro Gly Gln Leu Gly Arg Leu Leu Pro Asn<br>870 875 880     |                                                           |                                                               | 2814 |
| cag aat tta cca ctt gac atc aca ttg caa agc cca act ggt gct gga<br>Gln Asn Leu Pro Leu Asp Ile Thr Leu Gln Ser Pro Thr Gly Ala Gly<br>885 890 895 900 |                                                           |                                                               | 2862 |
| cct ttc cca cca atc aga aac agt agt ccc tac tca gtg ata cct cag<br>Pro Phe Pro Pro Ile Arg Asn Ser Ser Pro Tyr Ser Val Ile Pro Gln<br>905 910 915     |                                                           |                                                               | 2910 |
| cca gga atg atg ggt aat caa ggg atg ata gga aac caa gga aat tta<br>Pro Gly Met Met Gly Asn Gln Gly Met Ile Gly Asn Gln Gly Asn Leu<br>920 925 930     |                                                           |                                                               | 2958 |
| ggg aac agt agc aca gga atg att ggt aac agt gct tct cgg cct act<br>Gly Asn Ser Ser Thr Gly Met Ile Gly Asn Ser Ala Ser Arg Pro Thr<br>935 940 945     |                                                           |                                                               | 3006 |
| atg cca tct gga gaa tgg gca ccg cag agt tcg gct gtg aga gtc acc                                                                                       |                                                           |                                                               | 3054 |

|      |     |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|------|-----|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Met  | Pro | Ser  | Gly  | Glu  | Trp  | Ala  | Pro  | Gln  | Ser  | Ser  | Ala  | Val  | Arg  | Val  | Thr  |      |
| 950  |     |      |      |      |      | 955  |      |      |      |      | 960  |      |      |      |      |      |
| tgt  | gct | gct  | acc  | acc  | agt  | gcc  | atg  | aac  | cgg  | cca  | gtc  | caa  | gga  | ggg  | atg  | 3102 |
| Cys  | Ala | Ala  | Thr  | Thr  | Ser  | Ala  | Met  | Asn  | Arg  | Pro  | Val  | Gln  | Gly  | Gly  | Met  |      |
| 965  |     |      |      |      | 970  |      |      |      |      | 975  |      |      |      |      | 980  |      |
| att  | cgg | aac  | cca  | gca  | gcc  | agc  | atc  | ccc  | atg  | agg  | ccc  | agc  | agc  | cag  | cct  | 3150 |
| Ile  | Arg | Asn  | Pro  | Ala  | Ala  | Ser  | Ile  | Pro  | Met  | Arg  | Pro  | Ser  | Ser  | Gln  | Pro  |      |
|      |     |      |      | 985  |      |      |      |      | 990  |      |      |      |      | 995  |      |      |
| ggc  | caa | aga  | cag  | acg  | ctt  | cag  | tct  | cag  | gtc  | atg  | aat  | ata  | ggg  | cca  | tct  | 3198 |
| Gly  | Gln | Arg  | Gln  | Thr  | Leu  | Gln  | Ser  | Gln  | Val  | Met  | Asn  | Ile  | Gly  | Pro  | Ser  |      |
|      |     |      | 1000 |      |      |      |      | 1005 |      |      |      |      | 1010 |      |      |      |
| gaa  | tta | gag  | atg  | aac  | atg  | ggg  | gga  | cct  | cag  | tat  | agc  | caa  | caa  | caa  | gct  | 3246 |
| Glu  | Leu | Glu  | Met  | Asn  | Met  | Gly  | Gly  | Pro  | Gln  | Tyr  | Ser  | Gln  | Gln  | Gln  | Ala  |      |
|      |     | 1015 |      |      |      |      | 1020 |      |      |      |      | 1025 |      |      |      |      |
| cct  | cca | aat  | cag  | act  | gcc  | cca  | tgg  | cct  | gaa  | agc  | atc  | ctg  | cct  | ata  | gac  | 3294 |
| Pro  | Pro | Asn  | Gln  | Thr  | Ala  | Pro  | Trp  | Pro  | Glu  | Ser  | Ile  | Leu  | Pro  | Ile  | Asp  |      |
|      |     | 1030 |      |      |      | 1035 |      |      |      |      | 1040 |      |      |      |      |      |
| cag  | gcg | tct  | ttt  | gcc  | agc  | caa  | aac  | agg  | cag  | cca  | ttt  | ggc  | agt  | tct  | cca  | 3342 |
| Gln  | Ala | Ser  | Phe  | Ala  | Ser  | Gln  | Asn  | Arg  | Gln  | Pro  | Phe  | Gly  | Ser  | Ser  | Pro  |      |
| 1045 |     |      |      | 1050 |      |      |      |      |      | 1055 |      |      |      |      | 1060 |      |
| gat  | gac | ttg  | cta  | tgt  | cca  | cat  | cct  | gca  | gct  | gag  | tct  | ccg  | agt  | gat  | gag  | 3390 |
| Asp  | Asp | Leu  | Leu  | Cys  | Pro  | His  | Pro  | Ala  | Ala  | Glu  | Ser  | Pro  | Ser  | Asp  | Glu  |      |
|      |     |      |      | 1065 |      |      |      |      | 1070 |      |      |      |      | 1075 |      |      |
| gga  | gct | ctc  | ctg  | gac  | cag  | ctg  | tat  | ctg  | gcc  | ttg  | cgg  | aat  | ttt  | gat  | ggc  | 3438 |
| Gly  | Ala | Leu  | Leu  | Asp  | Gln  | Leu  | Tyr  | Leu  | Ala  | Leu  | Arg  | Asn  | Phe  | Asp  | Gly  |      |
|      |     |      | 1080 |      |      |      |      | 1085 |      |      |      |      | 1090 |      |      |      |
| ctg  | gag | gag  | att  | gat  | aga  | gcc  | tta  | gga  | ata  | ccc  | gaa  | ctg  | gtc  | agc  | cag  | 3486 |
| Leu  | Glu | Glu  | Ile  | Asp  | Arg  | Ala  | Leu  | Gly  | Ile  | Pro  | Glu  | Leu  | Val  | Ser  | Gln  |      |
|      |     | 1095 |      |      |      |      | 1100 |      |      |      |      | 1105 |      |      |      |      |
| agc  | caa | gca  | gta  | gat  | cca  | gaa  | cag  | ttc  | tca  | agt  | cag  | gat  | tcc  | aac  | atc  | 3534 |
| Ser  | Gln | Ala  | Val  | Asp  | Pro  | Glu  | Gln  | Phe  | Ser  | Ser  | Gln  | Asp  | Ser  | Asn  | Ile  |      |
|      |     | 1110 |      |      |      | 1115 |      |      |      |      | 1120 |      |      |      |      |      |
| atg  | ctg | gag  | cag  | aag  | gcg  | ccc  | gtt  | ttc  | cca  | cag  | cag  | tat  | gca  | tct  | cag  | 3582 |
| Met  | Leu | Glu  | Gln  | Lys  | Ala  | Pro  | Val  | Phe  | Pro  | Gln  | Gln  | Tyr  | Ala  | Ser  | Gln  |      |
| 1125 |     |      |      |      | 1130 |      |      |      |      | 1135 |      |      |      |      | 1140 |      |
| gca  | caa | atg  | gcc  | cag  | ggg  | agc  | tat  | tct  | ccc  | atg  | caa  | gat  | cca  | aac  | ttt  | 3630 |
| Ala  | Gln | Met  | Ala  | Gln  | Gly  | Ser  | Tyr  | Ser  | Pro  | Met  | Gln  | Asp  | Pro  | Asn  | Phe  |      |
|      |     |      |      | 1145 |      |      |      |      | 1150 |      |      |      |      | 1155 |      |      |
| cac  | acc | atg  | gga  | cag  | cgg  | cct  | agt  | tat  | gcc  | aca  | ctc  | cgt  | atg  | cag  | ccc  | 3678 |
| His  | Thr | Met  | Gly  | Gln  | Arg  | Pro  | Ser  | Tyr  | Ala  | Thr  | Leu  | Arg  | Met  | Gln  | Pro  |      |
|      |     |      | 1160 |      |      |      |      | 1165 |      |      |      |      | 1170 |      |      |      |
| aga  | ccg | ggc  | ctc  | agg  | ccc  | acg  | ggc  | cta  | gtg  | cag  | aac  | cag  | cca  | aat  | caa  | 3726 |
| Arg  | Pro | Gly  | Leu  | Arg  | Pro  | Thr  | Gly  | Leu  | Val  | Gln  | Asn  | Gln  | Pro  | Asn  | Gln  |      |
|      |     | 1175 |      |      |      |      | 1180 |      |      |      |      | 1185 |      |      |      |      |

|                                                                                                                                                           |      |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| cta aga ctt caa ctt cag cat cgc ctc caa gca cag cag aat cgc cag<br>Leu Arg Leu Gln Leu Gln His Arg Leu Gln Ala Gln Gln Asn Arg Gln<br>1190 1195 1200      | 3774 |
| cca ctt atg aat caa atc agc aat gtt tcc aat gtg aac ttg act ctg<br>Pro Leu Met Asn Gln Ile Ser Asn Val Ser Asn Val Asn Leu Thr Leu<br>1205 1210 1215 1220 | 3822 |
| agg cct gga gta cca aca cag gca cct att aat gca cag atg ctg gcc<br>Arg Pro Gly Val Pro Thr Gln Ala Pro Ile Asn Ala Gln Met Leu Ala<br>1225 1230 1235      | 3870 |
| cag aga cag agg gaa atc ctg aac cag cat ctt cga cag aga caa atg<br>Gln Arg Gln Arg Glu Ile Leu Asn Gln His Leu Arg Gln Arg Gln Met<br>1240 1245 1250      | 3918 |
| cat cag caa cag caa gtt cag caa cga act ttg atg atg aga gga caa<br>His Gln Gln Gln Gln Val Gln Gln Arg Thr Leu Met Met Arg Gly Gln<br>1255 1260 1265      | 3966 |
| ggg ttg aat atg aca cca agc atg gtg gct cct agt ggt atg cca gca<br>Gly Leu Asn Met Thr Pro Ser Met Val Ala Pro Ser Gly Met Pro Ala<br>1270 1275 1280      | 4014 |
| act atg agc aac cct cgg att ccc cag gca aat gca cag cag ttt cca<br>Thr Met Ser Asn Pro Arg Ile Pro Gln Ala Asn Ala Gln Gln Phe Pro<br>1285 1290 1295 1300 | 4062 |
| ttt cct cca aac tac gga ata agt cag caa cct gat cca ggc ttt act<br>Phe Pro Pro Asn Tyr Gly Ile Ser Gln Gln Pro Asp Pro Gly Phe Thr<br>1305 1310 1315      | 4110 |
| ggg gct acg act ccc cag agc cca ctt atg tca ccc cga atg gca cat<br>Gly Ala Thr Thr Pro Gln Ser Pro Leu Met Ser Pro Arg Met Ala His<br>1320 1325 1330      | 4158 |
| aca cag agt ccc atg atg caa cag tct cag gcc aac cca gcc tat cag<br>Thr Gln Ser Pro Met Met Gln Gln Ser Gln Ala Asn Pro Ala Tyr Gln<br>1335 1340 1345      | 4206 |
| gcc ccc tcc gac ata aat gga tgg gcg cag ggg aac atg ggc gga aac<br>Ala Pro Ser Asp Ile Asn Gly Trp Ala Gln Gly Asn Met Gly Gly Asn<br>1350 1355 1360      | 4254 |
| agc atg ttt tcc cag cag tcc cca cca cac ttt ggg cag caa gca aac<br>Ser Met Phe Ser Gln Gln Ser Pro Pro His Phe Gly Gln Gln Ala Asn<br>1365 1370 1375 1380 | 4302 |
| acc agc atg tac agt aac aac atg aac atc aat gtg tcc atg gcg acc<br>Thr Ser Met Tyr Ser Asn Asn Met Asn Ile Asn Val Ser Met Ala Thr<br>1385 1390 1395      | 4350 |
| aac aca ggt ggc atg agc agc atg aac cag atg aca gga cag atc agc<br>Asn Thr Gly Gly Met Ser Ser Met Asn Gln Met Thr Gly Gln Ile Ser<br>1400 1405 1410      | 4398 |
| atg acc tca gtg acc tcc gtg cct acg tca ggg ctg tcc tcc atg ggt<br>Met Thr Ser Val Thr Ser Val Pro Thr Ser Gly Leu Ser Ser Met Gly<br>1415 1420 1425      | 4446 |

```

ccc gag cag gtt aat gat cct gct ctg agg gga ggc aac ctg ttc cca      4494
Pro Glu Gln Val Asn Asp Pro Ala Leu Arg Gly Gly Asn Leu Phe Pro
      1430                      1435                      1440

aac cag ctg cct gga atg gat atg att aag cag gag gga gac aca aca      4542
Asn Gln Leu Pro Gly Met Asp Met Ile Lys Gln Glu Gly Asp Thr Thr
      1445                      1450                      1455                      1460

cgg aaa tat tgc tga cactgctgaa gccagttgct tcttcagctg accggggtca      4597
Arg Lys Tyr Cys

cttgc tcaaaa acacttccag tctggagagc tgtgtctatt tgtttcaacc caactgacct      4657

gccagccggt tctgctagag cagacaggcc tggccctggt tcccagggtg gcgtccactc      4717

ggctgtggca ggaggagctg cctcttctct tgacagctctg aagctcgcat ccagacagtc      4777

gctcagtctg ttccctgcat tcaccttagt gcaacttaga tctctctctc ccaagtaaat      4837

gttgacaggc caatttcata cccatgtcag attgaatgta tttaaatgta tgtatttaag      4897

gagaaccatg ctcttggttct gttcctgttc gggtccagac actgggttct tgctttgttt      4957

tccctggcta acagtctagt gccaaagatt aagattttat ctgggggaaa gaaaagaatt      5017

ttttaaaaaa ttaaaactaaa gatgttttaa gctaaagcct gaatttgga tggaagcagg      5077

acagacaccg tggacagcgc tgtatttaca gacacacca gtgcgtgaag accaacaag      5137

tcacagtcgt atctctagaa agctctaaag accatgttgg aaagagtctc cagttactga      5197

acagatgaaa aggagcctgt gagagggctg ttaacattag caaatatatt ttcttggttt      5257

tttctttggt aaaaccaaac tggttcacct gaatcatgaa ttgagaagaa ataattttca      5317

tttctaaatt aagtcctttt tagtttgatc agacagcttg aatcagcatc tcttcttccc      5377

tgtcagcctg actcttccct tccctctctc cattcccat actccctatt ttcatctctt      5437

ttttaaaaaa taatataagc tacagaaacc aggtaagccc tttatttcct taaatgtttt      5497

gccagccact taccaattgc taagtattga atttcagaaa aaaaaaatgc atttactggc      5557

aaggagaaga gcaaagttaa ggcttgatac caatcgagct aaggatacct gctttggaag      5617

catgtttatt ctgttcccca gcaactctgg cctccaaaat gggagaaacg ccagtgtgtt      5677

taaattgata gcagatatca cgacagattt aacctctgcc atgtgttttt tattttgttt      5737

tttagcagtg ctgactaagc cgaagttttg taagggtacat aaaatccaat ttatatgtaa      5797

acaagcaata atttaagttg agaacttatg tgttttaatt gtataatttt tgtgaggtat      5857

acatattgtg gaattgactc aaaaatgagg tacttcagta ttaaattaga tatcttcata      5917

gcaatgtctc ctaaagggtgt tttgtaaagg atatcaatgc cttgattaga cctaatttgt      5977

```

9

agacttaaga ctttttatatt tctaaacctt gtgattctgc ttataagtca tttatctaata 6037

ctatatgata tgcagccgct gtaggaacca attcttgatt tttatatggt tatattcttt 6097

cttaatgaac cttagaaaga ctacatgtta ctaagcaggc cacttttatg gttgttttt 6156

<210> 4

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 4

ctccagggca gagacaagaa ag 22

<210> 5

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 5

cacgattacg tttttcagtg ttctt 25

<210> 6

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 6

tcctgaccaa cttggaccca gccc 24

<210> 7

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 7

gaagggtgaag gtcggagtc 19

10

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 8  
gaagatggtg atgggatttc

20

<210> 9  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 9  
caagcttccc gttctcagcc

20

<210> 10  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 10  
atcggtcgg gtcggtca

18

<210> 11  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 11  
actcatcttg aacacata

18

<210> 12  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

11

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 12

catccactc atcttgaa

18

&lt;210&gt; 13

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 13

tttctcccat ccactca

18

&lt;210&gt; 14

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 14

ccaagttggt caggacat

18

&lt;210&gt; 15

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 15

tgcacatttg tcaggttt

18

&lt;210&gt; 16

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 16

tgcacttcac ctatgttg

18

&lt;210&gt; 17

12

<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 17  
ggccccagcg catccttg

18

<210> 18  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 18  
tgccttccag gttcacta

18

<210> 19  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 19  
gttatacctt agatactg

18

<210> 20  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 20  
ccaacatgca agatgcta

18

<210> 21  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

13

<400> 21  
catttactat agactttg 18

<210> 22  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 22  
ttctgaatca ggtaaagg 18

<210> 23  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 23  
gagagacagc gaagcact 18

<210> 24  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 24  
agaactggtc tttccttc 18

<210> 25  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 25  
ctgcgcatgg aacttctg 18

<210> 26  
<211> 18

14

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 26

gaagcatatg taaagata

18

&lt;210&gt; 27

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 27

aaccaccaaa cctgccca

18

&lt;210&gt; 28

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 28

aaaactgact gggtggga

18

&lt;210&gt; 29

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 29

aagccaacga tgacccta

18

&lt;210&gt; 30

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

15

<400> 30  
tctgcccttt gctgtcat 18

<210> 31  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 31  
gcaagctacc tgtggagt 18

<210> 32  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 32  
attctctttc ttcttggg 18

<210> 33  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 33  
ctattaattt tgtgttac 18

<210> 34  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 34  
gggaaaagct gtggtaat 18

<210> 35  
<211> 18  
<212> DNA

16

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 35

ctctgtgaaa ttcgcagt

18

&lt;210&gt; 36

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 36

ccctaaattt ccttggtt

18

&lt;210&gt; 37

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 37

ttcatggcac tggtggtta

18

&lt;210&gt; 38

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 38

ggacatagca agtcatct

18

&lt;210&gt; 39

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 39

17

tgctgtggga aaacgggc 18

<210> 40

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 40

gctgtgcttg gaggcgat 18

<210> 41

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 41

ggatttccct ctgtctct 18

<210> 42

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 42

agggttgctc atagttgc 18

<210> 43

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 43

gctgggttgg cctgagac 18

<210> 44

<211> 18

<212> DNA

<213> Artificial Sequence

18

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 44

gtaggcacgg aggtcact

18

&lt;210&gt; 45

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 45

cagcaatatt tccgtgtt

18

&lt;210&gt; 46

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 46

tcagcagtgt cagcaata

18

&lt;210&gt; 47

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 47

ctggcttcag cagtgtca

18

&lt;210&gt; 48

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 48

gacgccaccc tgggaacc

18

19

&lt;210&gt; 49

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 49

ctgtcaacat ttacttgg