



(12)

Rettet oversættelse af
europæisk patentskriftPatent- og
Varemærkestyrelsen

(51) Int.Cl.: **A 61 K 39/00 (2006.01)**
C 07 K 16/18 (2006.01)
C 12 P 21/08 (2006.01)

(45) Oversættelsen bekendtgjort den: **2024-08-19**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2023-04-19**

(86) Europæisk ansøgning nr.: **20174805.0**

(86) Europæisk indleveringsdag: **2016-08-18**

(87) Den europæiske ansøgnings publiceringsdag: **2021-01-27**

(30) Prioritet: **2015-08-19 US 201562207164 P**

(62) Stamansøgningsnr: **16837818.0**

(84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(73) Patenthaver: **Astrazeneca AB, 151 85 Södertälje, Sverige**

(72) Opfinder: **DEPAZ, Roberto, c/o Medimmune LLC, One Medimmune Way, Gaithersburg, Maryland 20878, USA**
DEJESUS, Natalie, c/o Medimmune LLC, One Medimmune Way, Gaithersburg, Maryland 20878, USA
BEE, Jared, c/o Medimmune LLC, One Medimmune Way, Gaithersburg, Maryland 20878, USA

(74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**

(54) Benævnelse: **STABIL ANTI-IFNAR1-FORMULERING**

(56) Fremdragne publikationer:
WO-A2-2009/100309
US-A1- 2014 127 227
SZYM CZYK KATARZYNA ET AL: "Aggregation behavior of Triton X-114 and Tween 80 at various temperatures and concentrations studied by density and viscosity measurements", JOURNAL OF THERMAL ANALYSIS AND CALORIMETRY, KLUWER, DORDRECHT, NL, vol. 126, no. 1, 5 July 2016 (2016-07-05), pages 315-326, XP036052597, ISSN: 1388-6150, DOI: 10.1007/S10973-016-5631-3 [retrieved on 2016-07-05]
BRUCE A. KERWIN: "Polysorbates 20 and 80 used in the formulation of protein biotherapeutics: Structure and degradation pathways", DISSOLUTION PROFILE OF NOVEL COMPOSITE PELLET CORES BASED ON DIFFERENT RATIOS OF MICROCRYSTALLINE CELLULOSE AND ISOMALT, vol. 97, no. 8, 1 August 2008 (2008-08-01) , pages 2924-2935, XP055015864, ISSN: 0022-3549, DOI: 10.1002/jps.21190
WANG W ET AL: "ANTIBODY STRUCTURE, INSTABILITY, AND FORMULATION", JOURNAL OF PHARMACEUTICAL SCIENCES, AMERICAN CHEMICAL SOCIETY AND AMERICAN PHARMACEUTICAL ASSOCIATION, US, vol. 96, no. 1, 1 January 2007 (2007-01-01), pages 1-26, XP009084505, ISSN: 0022-3549, DOI: 10.1002/jps.20727
SHARMA V K ET AL: "The formulation and delivery of monoclonal antibodies", 1 January 2009 (2009-01-01), THERAPEUTIC MONOCLONAL ANTIBODIES : FROM BENCH TO CLINIC, WILEY, HOBOKEN, NJ, PAGE(S) 675 -

711, XP009143115, ISBN: 978-0-470-11791-0 * page 14 *

Meera Agarkhed ET AL: "Effect of Polysorbate 80 Concentration on Thermal and Photostability of a Monoclonal Antibody", AAPS PharmSciTech, vol. 14, no. 1, 14 November 2012 (2012-11-14), pages 1-9, XP055350741, DOI: 10.1208/s12249-012-9878-0

HA E ET AL: "PEROXIDE FORMATION IN POLYSORBATE 80 AND PROTEIN STABILITY", JOURNAL OF PHARMACEUTICAL SCIENCES, AMERICAN CHEMICAL SOCIETY AND AMERICAN PHARMACEUTICAL ASSOCIATION, US, vol. 91, no. 10, 1 October 2002 (2002-10-01), pages 2252-2264, XP001148131, ISSN: 0022-3549, DOI: 10.1002/JPS.10216

Naoto Inoue ET AL: "Arginine and lysine reduce the high viscosity of serum albumin solutions for pharmaceutical injection", Journal of Bioscience and Bioengineering, vol. 117, no. 5, 1 May 2014 (2014-05-01), pages 539-543, XP055135662, ISSN: 1389-1723, DOI: 10.1016/j.jbiosc.2013.10.016

Naoto Inoue ET AL: "Specific Decrease in Solution Viscosity of Antibodies by Arginine for Therapeutic Formulations", Molecular Pharmaceutics, vol. 11, no. 6, 2 June 2014 (2014-06-02), pages 1889-1896, XP055134481, ISSN: 1543-8384, DOI: 10.1021/mp5000218

DESCRIPTION

[0001] This application incorporates by reference a Sequence Listing submitted with the application via EFS-Web as a text file entitled "IFNAR-350WO1_SL.txt" created on August 18, 2016 and having a size of 2,490.

BACKGROUND OF THE INVENTION

Field of the Invention

[0002] The present invention relates to a stable, low viscosity antibody formulation, wherein the formulation comprises a high concentration of an antibody that specifically binds interferon alpha receptor 1 (IFNAR1) or an antigen-binding fragment thereof. In embodiments, the antibody formulation comprises anifrolumab or an antigen-binding fragment thereof.

[0003] In some embodiments, the invention relates to a stable antibody formulation comprising about 100 mg/mL to about 200 mg/mL of an antibody or fragment thereof that specifically binds IFNAR1, about 25 mM to about 130 mM of lysine or a lysine salt; an uncharged excipient; a surfactant; and a formulation buffer. In some embodiments, the invention is directed to a container, dosage form and kit. In some embodiments, the invention is directed to a method of making and using the stable antibody formulation.

Background

[0004] Antibodies have been used in the treatment of various diseases and conditions due to their specificity of target recognition, thereby generating highly selective outcomes following systemic administration. In order for antibodies to remain effective, they must maintain their biological activity during their production, purification, transport and storage. New production and purification techniques have been developed to provide for large amounts of highly purified monoclonal antibodies to be produced. However, challenges still exist to stabilize these antibodies for transport and storage, and yet even more challenges exist to provide the antibodies in a dosage form suitable for administration.

[0005] Denaturation, aggregation, contamination, and particle formation can be significant obstacles in the formulation and storage of antibodies. Due to the wide variety of antibodies, there are no universal formulations or conditions suitable for storage of all antibodies. Optimal formulations of one antibody are often specific to that antibody. Additionally, antibody formulations may need to be further tailored to a specific antibody depending on the concentration of the antibody, and/or a desired physical property, e.g., viscosity, of the antibody formulation. Antibody storage formulations are often a significant part of the research and development process for a commercial antibody.

[0006] Various methods have been proposed to overcome the challenges associated with antibody stability. For example, in some instances, the antibody is often lyophilized, and then reconstituted shortly before administration. However, reconstitution adds an additional step to the administration process, and could introduce contaminants to the formulation. Additionally, reconstituted antibodies could suffer from aggregation and particle formation. Thus, a need exists to provide stable antibody formulations that can overcome the challenges associated with transport and storage.

SUMMARY OF THE INVENTION

[0007] The present invention relates to a stable, low viscosity antibody formulation, wherein the formulation comprises a high concentration of anti- interferon alpha receptor 1 antibody or an antigen binding fragment thereof. The invention is defined by the claims. The invention is directed to an antibody formulation comprising:

1. a. 125 to 175 mg/mL anifrolumab;
2. b. 0.02% to 0.08% of a surfactant, wherein the surfactant is selected from the group consisting of polysorbate 20,

polysorbate 80 and poloxamer;

3. c. 100 mM to 150mM of an uncharged excipient, wherein the uncharged excipient is selected from the group consisting of glucose, sucrose, trehalose and glycerol;
4. d. a formulation buffer; and e. 25 to 130 mM of a lysine salt;

wherein the formulation is at a pH of 5.5 to 6.5, and wherein the formulation has a viscosity of equal to or less than 20 mPas at 25°C

[0008] Described is also an antibody formulation comprising about 100 mg/mL to about 200 mg/mL anifrolumab or an antigen binding fragment thereof; about 40 mM to about 60 mM lysine HCl; about 100 mM to about 160 mM trehalose dihydrate; about 0.02% to about 0.1% polysorbate 80; about 15 mM to about 35 mM histidine/histidine HCl, wherein the formulation is at a pH of from about 5.5 to 6.5.

[0009] In additional embodiments, the invention is directed to an antibody formulation comprising about 145 mg/mL to about 155 mg/mL anifrolumab ; about 45 mM to about 55 mM lysine HCl; about 120 mM to about 140 mM trehalose dihydrate; about 0.04% to about 0.08% polysorbate 80; and about 20 mM to about 30 mM histidine/histidine HCl, wherein the formulation is at a pH of from about 5.8 to about 6.1.

[0010] Described is also In some embodiments, the invention is directed to an antibody formulation comprising:
about 150 mg/mL anifrolumab ; about 50 mM lysine HCl;
about 130 mM trehalose dihydrate; about 0.05% polysorbate 80; and about 25 mM histidine/histidine HCl, wherein the formulation is at a pH of about 5.9.

[0011] Described is also In some embodiments, the invention is directed to an antibody formulation comprising:
150 mg/mL anifrolumab 50 mM lysine HCl; 130 mM trehalose dihydrate; 0.05% polysorbate 80; 25 mM histidine/histidine HCl, wherein the formulation is at a pH of about 5.9.

[0012] Described is also In additional embodiments, the invention is directed to an antibody formulation comprising: 150 mg/mL anifrolumab; 50 mM lysine HCl; 130 mM trehalose dihydrate; 0.05% polysorbate 80; 25 mM histidine/histidine HCl, wherein the formulation is at a pH of 5.9.

[0013] Described is also a pharmaceutical unit dosage form suitable for parenteral administration to a human which comprises any one of the antibody formulations described herein in a suitable container.

[0014] Described is also a kit comprising any antibody formulation described herein, a container as described herein, a unit dosage form as described herein, or a pre-filled syringe as described herein.

[0015] Described is also a method of producing a stable antibody formulation, the method comprising: purifying an antibody to about 100 mg/mL to about 200 mg/mL of an anti-IFNAR antibody or antigen-binding fragment thereof, placing the isolated antibody in a stabilizing formulation to form the stable antibody formulation, wherein the resulting stable antibody formulation comprises 100mg/mL to 200mg/mL of the antibody;
25 mM to 130 mM of lysine or a lysine salt; 100 mM to 150 mM uncharged excipient; 0.02% to 0.1% of a surfactant; and a formulation buffer.

[0016] Described is also a method of treating a type I IFN-mediated disease or disorder in a subject in need thereof, the method comprising administering a therapeutically effective amount of the antibody formulation of any one of the antibody formulations described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] For the purpose of illustrating the invention, there are depicted in the drawings certain embodiments of the invention. However, the invention is not limited to the precise arrangements and instrumentalities of the embodiments

depicted in the drawings.

Figure 1 shows viscosity as a function of pH in a high throughput screening bead-based method.

Figure 2 shows viscosity as a function of lysine HCl concentration using a high throughput screening bead-based method.

Figure 3 shows viscosity as a function of concentration of antibody (anifrolumab) concentration for formulations containing 25 mM histidine/histidine-HCl, 25 mM lysine, 130 mM trehalose at pH 6.0 (circles); and 25 mM histidine/histidine-HCl, 50 mM lysine, 130 mM trehalose at pH 6.0 (diamonds).

Figure 4 shows viscosity as a function of antibody (anifrolumab) concentration for solutions containing: 0 mM lysine (diamonds); 5 mM lysine (squares); 12.5 mM lysine (triangles); 25 mM lysine (x's); and 50 mM lysine (asterisks).

Figure 5 shows viscosity as a function of lysine HCl concentration for solutions containing: 135 mg/mL anifrolumab (diamonds); 150 mg/mL anifrolumab (squares); and 180 mg/mL anifrolumab (triangles).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0018] Before describing the present invention in detail, it is to be understood that this invention is not limited to specific compositions or process steps, as such can vary. It must be noted that, as used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise. The terms "a" (or "an"), as well as the terms "one or more," and "at least one" can be used interchangeably herein.

[0019] Furthermore, "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term "and/or" as used in a phrase such as "A and/or B" herein is intended to include "A and B," "A or B," "A," (alone) and "B" (alone). Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

[0020] Throughout the present disclosure, all expressions of percentage, ratio, and the like are "by weight" unless otherwise indicated. As used herein, "by weight" is synonymous with the term "by mass," and indicates that a ratio or percentage defined herein is done according to weight rather than volume, thickness, or some other measure.

[0021] The term "about" is used herein to mean approximately, in the region of, roughly, or around. When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" is used herein to modify a numerical value above and below the stated value by a variance of 10%.

[0022] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention is related. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular Biology, 3rd ed., 1999, Academic Press; and the Oxford Dictionary Of Biochemistry And Molecular Biology, Revised, 2000, Oxford University Press, provide one of skill with a general dictionary of many of the terms used in this invention.

[0023] Units, prefixes, and symbols are denoted in their Système International de Unites (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, amino acid sequences are written left to right in amino to carboxy orientation. The headings provided herein are not limitations of the various aspects or embodiments of the invention, which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification in its entirety.

[0024] It is understood that wherever embodiments are described herein with the language "comprising," otherwise

analogous embodiments described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0025] Amino acids are referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, are referred to by their commonly accepted single-letter codes.

[0026] As used herein, the term "injection force" is the amount of pressure (in Newtons) required to pass the antibody formulation through a needle.

[0027] As used herein, the term "autoimmune disease" refers to a disorder, disease state or condition associated with the formation of autoantibodies reactive with the patient's own cells to form antigen-antibody complexes. The term "autoimmune disease" includes conditions such as, e.g., systemic lupus erythematosus, as well as those disorders which are triggered by a specific external agent, e.g., acute rheumatic fever. Examples of autoimmune disorders include, but are not limited to, autoimmune hemolytic anemia, autoimmune hepatitis, Berger's disease, chronic fatigue syndrome, Crohn's disease, dermatomyositis, fibromyalgia, Graves' disease, Hashimoto's thyroiditis, idiopathic thrombocytopenia purpura, lichen planus, multiple sclerosis, myasthenia gravis, psoriasis, rheumatic fever, rheumatoid arthritis, scleroderma, Sjogren's syndrome, systemic lupus erythematosus, type 1 diabetes, ulcerative colitis, and vitiligo. In specific aspects, the autoimmune disease is systemic lupus erythematosus (SLE), scleroderma (SSe), myositis, or lupus nephritis.

[0028] The terms "Interferon alpha receptor-1," "IFNARI," and "IFNAR" are used interchangeably, and include variants, isoforms, species homologs of human IFNAR1, and analogs having at least one common epitope with IFNARI. See, e.g., de Weerd et al., *J. Bioi. Chern.* 282:20053-20057 (2007). Accordingly, human antibodies specific for human IFNARI, in certain cases, cross-react with IFNARI from species other than human, or other proteins which are structurally related to human IFNAR1 (e.g., human IFNAR1 homologs). In other cases, the antibodies can be completely specific for human IFNARI and not exhibit species or other types of cross-reactivity. The complete cDNA sequence of human IFNAR1 has the Genbank accession number NM 000629.

[0029] The terms "type I interferon" or "type I IFN" as used herein refer to members of the type I interferon family of molecules that are ligands for IFNAR1 (i.e., members of the type I interferon family of molecules that are capable of binding IFNAR1). Examples of type I interferon ligands are interferon alpha 1, 2a, 2b, 4, 5, 6, 7, S, 10, 14, 16, 17, 21, interferon beta and interferon omega.

[0030] The term "type I IFN-mediated disease or disorder" refers to any type I IFN or IFN a inducible disease, disorder, or condition that exhibits a type I IFN pharmacodynamic ("PD") marker expression profile or gene signature (type I IFN GS). A PD marker expression profile and a gene signature will be understood to be equivalent. These diseases, disorders, or conditions include those with an autoimmune component such as systemic lupus erythematosus (SLE), scleroderma, lupus nephritis, o myositis. A type I IFN-mediated disease or disorder can be treated by administering a small molecule or a biological agent, e.g., an antibody or an antigen binding fragment thereof. If the therapeutic agent is a biological agent, it may be an antibody specific for any subtype(s) of type I IFN or IFNa. For instance, the antibody may be specific for any one of IFN α 1, IFN α 2, IFN α 4, IFN α 5, IFN α 6, IFN α 7, IFN α 8, IFN α 10, IFN α 14, IFN α 17, IFN α 21, IFN β , or IFN ω . Alternatively, the antibody may be specific for any two, any three, any four, any five, any six, any seven, any eight, any nine, any ten, any eleven, any twelve type I IFN or IFNa subtypes. If the antibody is specific for more than one type I IFN subtype, the antibody may be specific for IFN α 1, IFN α 2, IFN α 4, IFN α 5, IFN α 8, IFN α 10, and IFN α 21; or it may be specific for IFN α 1, IFN α 2, IFN α 4, IFN α 5, IFN α 8, and IFN α 10; or it may be specific for IFN α 1, IFN α 2, IFN α 4, IFN α 5, IFN α 8, and IFN α 21; or it may be specific for IFN α 1, IFN α 2, IFN α 4, IFN α 5, IFN α 10, and IFN α 21. A therapeutic agent that modulates IFNa activity may neutralize IFNa activity. A type I IFN-mediated disease or disorder can also be treated with antibodies specific for a type I IFN receptor, e.g., IFNAR1. In some aspects, anti-IFNAR1 antibodies can cross-react with IFNAR1 from species other than human. In other aspects, the anti-IFNAR1 antibodies can be specific for IFNAR1 only and not exhibit species or other types of cross-reactivity. In some aspects, the anti-IFNAR1 antibodies exhibit reduced binding affinities for FC ligands and have reduced or ablated effector function (ADCC and/or CDC), reduced or ablated binding to Fc ligands, or reduced or ablated toxicities as compared to an unmodified antibody.

[0031] The term "MEDI-546" refers to an Fc-modified version of the anti-IFNAR 9D4 antibody described in U.S. Patent No. 7,662,381. The terms "MEDI-546" and "anifrolumab" are used interchangeably herein. The sequence of MEDI-546 is described in U.S. 2011-0059078. MEDI-546 comprises a combination of three mutations: L234F, L235E, and P331S,

wherein the numbering is according to the EU index as set forth in Kabat, introduced into the lower hinge and CH2 domain of human IgG 1, which cause a decrease in their binding to human Fc_YRI (CD64), Fc_YRIIA (CD32A), Fc_YRIII (CD16) and C1q. See, e.g., U.S. 2011/0059078 and Oganesyan et al. *Acta Crystallographica D* 64:700-704 (2008), which are hereby incorporated by reference in their entireties. The VH and V_k sequences of MEDI-546 are shown in TABLE 1.

TABLE 1

MEDI-546 VH (SEQ ID NO:1)	EVQLVQSGAEVKPGESLKISCKGSGYIFTNYWIAWVRQMPGKC LESMGIYPGDSDIRYSPSFQGQVTISADKSITTAYLQWSSLKAS DTAMYYCARHDIEGFDYWGRGTLVTVSS
MEDI-546 V _k (SEQ ID NO:2)	EIVLTQSPGTLSLSPGERATLSCRASQSVSSFFAWYQQKPGQAPR LLIYGASSRATGIPDRLSGSGSGTDFTLTIRLEPEDFAVYYCQ QYDSSAITFGQGTRLEIK

[0032] The term "antibody or antigen-binding fragment thereof that modulates type I IFN activity" refers to an antibody (see *infra*) in its broadest sense capable of modulating type I IFN activity in a patient. The term "modulating" as used herein includes the inhibition or suppression of a type I IFN activity as well as the induction or enhancement of a type I IFN activity. In specific aspects, the type I IFN activity is IFN_α activity. In some aspects, the suppression of a type IFN GS is a suppression of a type I IFN activity. In some aspects, the antibody or antigen-binding fragment thereof is monoclonal. In specific aspects, the antibody or antigen-binding fragment thereof that modulates type I IFN activity specifically binds to a type I IFN receptor such as IFNAR1. In some specific aspects, the antibody or antigen binding fragment thereof specifically binds to subunit 1 of IFNAR1.

[0033] The term "antibody" is used herein in its broadest sense and includes, e.g., monoclonal antibodies, polyclonal antibodies, multivalent antibodies, multispecific antibodies, chimeric antibodies, and humanized antibodies. The term "antibody" includes whole antibodies. The term "antibody" also refers to a protein comprising at least two immunoglobulin heavy (H) chains and two immunoglobulin light (L) chains inter-connected by disulfide bonds, or an antigen binding portion thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FRI, CDRI, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies can mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

[0034] The term "antigen-binding fragment" refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., IFNAR). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen binding fragment" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird et al. (1988) *Science* 242:423-426; and Huston et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding fragment" of an antibody. These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact

antibodies.

[0035] An "isolated antibody," as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds IFNAR is substantially free of antibodies that specifically bind antigens other than IFNAR). An isolated antibody that specifically binds IFNAR can, however, have cross-reactivity to other antigens, such as IFNAR molecules from other species. Moreover, an isolated antibody can be substantially free of other cellular material and/or chemicals.

[0036] The term "monoclonal antibody" as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody displays a single binding specificity and affinity for a particular epitope.

[0037] The term "human antibody," as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from human germline immunoglobulin sequences. Furthermore, if the antibody contains a constant region, the constant region also is derived from human germline immunoglobulin sequences. The human antibodies of the disclosure can include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or sitespecific mutagenesis in vitro or by somatic mutation in vivo). However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[0038] The term "human monoclonal antibody" refers to antibodies displaying a single binding specificity which have variable regions in which both the framework and CDR regions are derived from human germline immunoglobulin sequences. In one embodiment, the human monoclonal antibodies are produced by a hybridoma which includes a B cell obtained from a transgenic nonhuman animal, e.g., a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

[0039] The term "recombinant human antibody", as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom (described further below), (b) antibodies isolated from a host cell transformed to express the human antibody, e.g., from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial human antibody library, and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable regions in which the framework and CDR regions are derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies can be subjected to in vitro mutagenesis (or, when an animal transgenic for human Ig sequences is used, in vivo somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire in vivo.

[0040] The term "antibody" as used herein also includes "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison et al, Proc. Natl. Acad. Sci. USA 81:6851-6855 (1984)).

[0041] Basic antibody structures in vertebrate systems are relatively well understood. See, e.g., Harlow et al. (1988) *Antibodies: A Laboratory Manual* (2nd ed.; Cold Spring Harbor Laboratory Press).

[0042] In the case where there are two or more definitions of a term that is used and/or accepted within the art, the definition of the term as used herein is intended to include all such meanings unless explicitly stated to the contrary. A specific example is the use of the term "complementarity determining region" ("CDR") to describe the non-contiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. This particular region has been described by Kabat et al. (1983) U.S. Dept. of Health and Human Services, "Sequences of Proteins of Immunological Interest" and by Chothia and Lesk, J. Mol. Biol. 196:901-917 (1987), which are incorporated herein by reference, where the definitions include overlapping or subsets of amino acid residues when compared against each

other. Nevertheless, application of either definition to refer to a CDR of an antibody or variants thereof is intended to be within the scope of the term as defined and used herein. The appropriate amino acid residues that encompass the CDRs as defined by each of the above cited references are set forth below in TABLE 2 as a comparison. The exact residue numbers that encompass a particular CDR will vary depending on the sequence and size of the CDR. Those skilled in the art can routinely determine which residues comprise a particular CDR given the variable region amino acid sequence of the antibody.

TABLE 2 CDR Definitions¹

	Kabat	Chothia
VH CDR1	31-35	26-32
VH CDR2	50-65	52-58
VH CDR3	95-102	95-102
VL CDR1	24-34	26-32
VL CDR2	50-56	50-52
VL CDR3	89-97	91-96

¹Numbering of all CDR definitions in TABLE 2 is according to the numbering conventions set forth by Kabat et al. (see below).

[0043] Kabat et al. also defined a numbering system for variable domain sequences that is applicable to any antibody. One of ordinary skill in the art can unambiguously assign this system of "Kabat numbering" to any variable domain sequence, without reliance on any experimental data beyond the sequence itself. As used herein, "Kabat numbering" refers to the numbering system set forth by Kabat et al. (1983) U.S. Dept. of Health and Human Services, "Sequence of Proteins of Immunological Interest." Unless otherwise specified, references to the numbering of specific amino acid residue positions in an anti-IFNAR antibody or antigen-binding fragment, variant, or derivative thereof of the present disclosure are according to the Kabat numbering system.

[0044] The terms "treat" or "treatment" as used herein refer to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) an undesired physiological change or disorder in a subject, such as the progression of an inflammatory disease or condition. Beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. The term "treatment" also means prolonging survival as compared to expected survival if not receiving treatment. Those in need of treatment include those already with the condition or disorder as well as those prone to have the condition or disorder or those in which the condition or disorder is to be prevented.

[0045] Terms such as "treating" or "treatment" or "to treat" refer to both (1) therapeutic measures that cure, slow down, lessen symptoms of, and/or halt progression of a diagnosed pathologic condition or disorder and (2) prophylactic or preventative measures that prevent and/or slow the development of a targeted pathologic condition or disorder. Thus, those in need of treatment include those already with the disorder; those prone to have the disorder; and those in whom the disorder is to be prevented.

[0046] The terms "effective amount" or "amount effective to" or "therapeutically effective amount" includes reference to a dosage of a therapeutic agent sufficient to produce a desired result.

[0047] By "subject" or "patient" is meant any subject, particularly a mammalian subject, for whom diagnosis, prognosis, or therapy is desired. As used herein, the terms "subject" or "patient" include any human or nonhuman animal. The term "nonhuman animal" includes all vertebrates, e.g., mammals and non-mammals, such as nonhuman primates, sheep, dogs, cats, horses, cows, bears, chickens, amphibians, reptiles, etc. As used herein, phrases such as "a patient having a type I IFN-mediated disease or disorder" includes subjects, such as mammalian subjects, that would benefit from the administration of an antibody or antigenbinding fragment thereof that modulates type I IFN activity, e.g., for detection, imaging, or other diagnostic procedure, and/or from treatment, i.e., palliation or prevention of a disease, with such antibody or antigen-binding thereof.

Antibody Formulations

[0048] In embodiments, the present invention provides an antibody formulation comprising:

1. a. about 125 to about 175 mg/mL anifrolumab;
2. b. about 0.02% to about 0.08% of a surfactant, wherein the surfactant is selected from the group consisting of polysorbate 20, polysorbate 80 and poloxamer;
3. c. about 100 mM to about 150mM of an uncharged excipient, wherein the uncharged excipient is selected from the group consisting of glucose, sucrose, trehalose and glycerol;
4. d. a formulation buffer; and
5. e. about 25 to 130 mM of a lysine salt;

wherein the formulation is at a pH of 5.5 to 6.5, and wherein the formulation has a viscosity of equal to or less than 20 mPas at 25°C.

[0049] Described formulations can comprise an antibody or antigen-binding fragment thereof that comprises a VH domain sequence comprising from 0 to 5 amino acid substitutions from a reference VH having the amino acid sequence SEQ ID NO: 1

[0050] Described formulations can comprise comprise an antibody or antigen-binding fragment comprises a Vk domain comprising from 0-5 amino acid substitutions from a reference Vk having the amino acid sequence SEQ ID NO:2.

[0051] In some embodiments, the antibody in the antibody formulation is purified prior to being added to the antibody formulation. The terms "isolate," and "purify" refer to separating the antibody from an impurity or other contaminants in the composition which the antibody resides, e.g., a composition comprising host cell proteins. In some embodiments, at least 50%, 70%, 80%, 90%, 95%, 98%, 99%, 99.5%, or 99.9% (w/w) of an impurity is purified from the antibody. For example, in some embodiments, purification of an antibody, e.g. anti-IFNAR1 antibody, would comprise separating the antibody from 99% (w/w) of the host cell proteins present originally in the composition.

[0052] In some embodiments, the terms "isolate," and "purify" refer to separating an antibody, e.g. anti- IFNAR1 antibody, from an impurity or other contaminants in the composition to an extent consistent with guidelines of a governmental organization, e.g., the World Health Organization or the United States Food and Drug Administration.

[0053] Methods of purifying an antibody are known to those of skill in the art. Suitable techniques for carrying out purification include various types of chromatography, such as affinity chromatography, hydrophobic interaction, ion exchange (such as cation exchange chromatography or mixed-mode chromatography), and filtration.

[0054] Affinity chromatography refers to a separation method whereby an antibody, by virtue of its specific binding properties, is bound to an affinity ligand for the antibody. The functional affinity ligand can be immobilized on a solid or semi-solid support so that when a composition comprising the antibody is passed over the ligand and the solid support, the antibody having a specific binding affinity to the ligand adsorbs to the ligand, and one or more other impurities are not adsorbed (or are bound at a lower affinity) and are separated from the antibody. Examples of impurities that do not typically bind (or do not bind well) include process-related impurities (e.g., host cell proteins, DNA, medium components) and some product-related impurities (e.g., antibody fragments). In some embodiments, the solid support comprising the ligand is washed one or more times with a buffer to remove additional impurities before the adsorbed antibody is removed from the ligand and the support. After one or more impurities have been removed, the adsorbed antibody can be removed (eluted) from the ligand and the support, resulting in isolation of the antibody from the original composition. Methods of removing the antibody from the ligand and support are dependent on the ligand and are known to those of skill in the art and can include, e.g., changes in environment, e.g., pH, addition or chaotropic agents or denaturants, or addition of commercially available elution buffers. In some embodiments, more than one affinity purification process can be employed on an antibody composition. Various affinity ligands are known in the art, including Protein A and Protein G (and combinations thereof). Immobilized ligands are commercially available. For example, Protein A affinity systems include MabSelect, MabSelect SuRe, MabSelect Xtra, MabSelect SuRe LX, Sepaharose CL-4B, ProSep vA, ProSep vA Ultra, and Ceramic HyperD.


[0055] Ion exchange chromatography includes cation exchange chromatography and mixed chromatography. Cation exchange chromatography refers to any method by which an antibody and some impurity or impurities can be separated based on charge differences using a cation exchange matrix. A cation exchange matrix generally comprises covalently bound, negatively charged groups. Weak or strong cation exchange resins may be employed. Commonly, strong cation exchange resins comprise supported organic groups comprising sulphonate acid or sulphonate groups, depending upon the pH. Weak cation exchanger resins commonly comprise supported organic groups comprising carboxylic acid or carboxylate groups, depending upon the pH. In certain embodiments, multimodal cation exchange resins can be used, which incorporate additional binding mechanisms as well as the ionic interactions, for example one or more of hydrogen bonding interactions and hydrophobic interactions. Examples of suitable cation exchange resins are well known in the art, and can include, but are not limited to Fractogel, carboxymethyl (CM), sulfoethyl(SE), sulfopropyl(SP), phosphate(P) and sulfonate(S), PROPAC WCX-10™ (Dionex), Capto S, S-Sepharose FF, Fractogel EMD SO₃M, Toyopearl Megacap II SP 550C, Poros 50 HS, and SP-sepharose matrix. In some embodiments, more than one cation exchange chromatography process can be employed on the composition.

[0056] Mixed mode chromatography refers to a method that utilizes more than one form of interaction between the stationary phase and analytes in order to achieve their separation from impurities (e.g., process-related impurities, such as host-cell proteins, DNA, and/or endogenous or adventitious viruses). Examples of suitable anion exchange matrices are known in the art, and can include, but are not limited to, Capto Adhere, Sartobind Q, Natrix Q, Chromasorb Q, and Mustang Q.

[0057] In some embodiments, additional filtration steps can be used to remove impurities. For example, in some embodiments nanofiltration or ultrafiltration is used. Nanofiltration comprises passing the composition through a matrix having a pore size of, e.g., less than 75 nm, less than 50 nm, and even less than 15 nm, to separate impurities, e.g., viruses, from the antibody. Commercially available nanofilters and ultrafilters that can be employed are manufactured by various vendors such as Millipore Corporation (Billerica, Mass., e.g., Viresolve Pro and Viresolve Pro+), Pall Corporation (East Hills, N.Y.), GE Healthcare Sciences (Piscataway, N.J.), and Sartorius Corporation (Goettingen, Germany).

[0058] In some embodiments, the antibody used in the described formulations e.g., an anti-IFNAR, or antigen binding fragment thereof comprises the Kabat-defined VH sequence in SEQ ID NO: 1 and the Kabat-defined Vk of SEQ ID NO:2, wherein the antibody is in the formulation at a concentration of 10 mg/ml to 300 mg/ml, 30 mg/ml to 250 mg/ml, 50 mg/ml to 200 mg/ml, 100 mg/ml to 200 mg/ml, 125 mg/ml to 175 mg/ml, 130 mg/ml to 170 mg/mL, 135 mg/ml to 165 mg/mL, 140 mg/ml to 160 mg/mL, 145 mg/ml to 155 mg/mL, 130 mg/ml, 135 mg/ml, 140 mg/ml, 145 mg/ml, 146 mg/ml, 147 mg/ml, 148 mg/ml, 149 mg/ml, 150 mg/ml, 151 mg/ml, 152 mg/ml, 153 mg/ml, 154 mg/ml, 155 mg/ml, 156 mg/ml, 157 mg/ml, 158 mg/ml, 159 mg/ml or 160 mg/ml. In some embodiments, the antibody, is in concentration of about 50 mg/ml, 55 mg/ml, 60 mg/ml, 65 mg/ml, 70 mg/ml, 75 mg/ml, 80 mg/ml, 85 mg/ml, 90 mg/ml, 95 mg/ml, 100 mg/ml, 125 mg/ml, 130 mg/ml, 140 mg/ml, 150 mg/ml, 160 mg/ml, 170 mg/ml, 180 mg/ml, 190 mg/ml, 200 mg/ml.

[0059] The antibody formulations of the present invention comprise lysine. Lysine is an essential amino acid having the following structure:



Lysine, as used herein, can include the free base form of lysine, as well as any and all salts thereof. In embodiments, the salt form of lysine is lysine acetate, lysine monochloride, lysine dichloride, lysine L-aspartate, and lysine L-glutamate. In some embodiments, lysine includes a pharmaceutically acceptable salt thereof. For example, lysine would include lysine hydrochloride. Lysine, as used herein, also includes all enantiomers (e.g., L-lysine and S-lysine), and any combination of enantiomers (e.g., 50% L-lysine and 50% S-lysine; 90%-100% L-lysine and 10%-0% S-lysine, etc.). In some embodiments, the term "lysine" includes greater than 99% L-lysine and less than 1% S-lysine. In some embodiments, the term "lysine" includes a enantomerically pure L-lysine. In some embodiments, lysine is a pharmaceutical grade lysine.


[0060] The described antibody formulations can comprise about 10 to about 100 mM lysine, about 20 to about 90 mM lysine, about 30 mM lysine to about 80 mM lysine, about 40 to about 70 mM lysine, about 45 to about 65 mM lysine, about 45 to about 60 mM lysine, about 50 to about 55 mM lysine in the antibody formulation, e.g., an antibody formulation comprising 100 to 200 mg/mL antibody, or about 150 mg/mL antibody. The described antibody formulations can comprise

about 50 mM lysine HCl in an antibody formulation comprising 100 to 200 mg/mL antibody, or about 150 mg/mL antibody, an uncharged excipient, a surfactant, and a formulation buffer.

[0061] The described antibody formulations can comprise an uncharged excipient. The term excipient refers to a pharmacologically inactive substance formulated with the antibody as described herein. In some embodiments, the excipient can assist in the prevention of denaturation or otherwise assist in stabilizing the antibody. Suitable excipients that may be used in the pharmaceutical compositions are known in the art. Examples can be taken e.g. from the handbook: Gennaro, Alfonso R.: "Remington's Pharmaceutical Sciences", Mack Publishing Company, Easton, Pa., 1990. In some embodiments, the excipient is an "uncharged" excipient, i.e., the excipient does not carry either a positive "+" or negative "-" charge. The excipient is selected from the group consisting of fructose, glucose, mannose, sorbose, xylose, lactose, maltose, sucrose, dextran, pullulan, dextrin, cyclodextrins, soluble starch, trehalose, sorbitol, erythritol, isomalt, lactitol, maltitol, xylitol, glycerol, lactitol, hydroxyethyl starch, watersoluble glucans.

[0062] The uncharged excipient is about 1mM to about 1 M, about 2 mM to about 500 mM, about 5 mM to about 400 mM, about 10 mM to about 300 mM or about 20 mM to about 250 mM in the antibody formulation. The uncharged excipient is about 30 mM to about 230 mM, about 40 mM to about 220 mM, about 50 mM to about 210 mM, about 60 mM to about 210 mM, about 70 mM to about 200 mM, about 80 mM to about 190 mM, about 90 mM to about 180 mM, about 100 mM to about 170 mM, about 110 mM to about 160 mM, about 120 mM to about 150 mM, about 125 mM to about 145 mM, about 125 mM to about 140 mM, about 120 mM, about 125 mM, about 130 mM, about 135 mM, about 140 mM, about 150 mM, about 160 mM, or about 170 mM in the antibody formulation, e.g., an antibody formulation comprising 100 to 200 mg/mL antibody. In one embodiment, the uncharged excipient is about 130 mM in the antibody formulation. The uncharged excipient is about 50 mM to about 500 mM, about 100 mM to about 450 mM, about 110 mM to about 350 mM, about 120 mM, about 125 mM, about 130 mM, about 140 mM, or about 145 mM in the antibody formulation, e.g., an antibody formulation comprising 100 to 200 mg/mL antibody or about 150 mg/mL antibody. The uncharged excipient is about 130 mM in the antibody formulation.

[0063] In some embodiments, the uncharged excipient is trehalose, as represented by the formula:




[0064] The trehalose can be about 1mM to about 1 M, about 2 mM to about 500 mM, about 5 mM to about 400 mM, about 10 mM to about 300 mM or about 20 mM to about 250 mM in the antibody formulation. The trehalose can be about 30 mM to about 230 mM, about 40 mM to about 220 mM, about 50 mM to about 210 mM, about 60 mM to about 210 mM, about 70 mM to about 200 mM, about 80 mM to about 190 mM, about 90 mM to about 180 mM, about 100 mM to about 170 mM, about 110 mM to about 160 mM, about 120 mM to about 150 mM, about 125 mM to about 145 mM, about 125 mM to about 140 mM, about 120 mM, about 125 mM, about 130 mM, about 135 mM, about 140 mM, about 150 mM, about 160 mM, or about 170 mM in the antibody formulation, e.g., an antibody formulation comprising 100 to 200 mg/mL antibody. The trehalose can be about 130 mM in the antibody formulation. The trehalose can be about 50 mM to about 500 mM, about 100 mM to about 450 mM, about 110 mM to about 350 mM, about 120 mM, about 125 mM, about 130 mM, about 140 mM, or about 145 mM in the antibody formulation, e.g., an antibody formulation comprising 100 to 200 mg/mL antibody or about 150 mg/mL antibody. The trehalose can be about 130 mM in the antibody formulation.

[0065] Various other components can be included in the antibody formulation. The antibody formulation can comprise a buffer (e.g., histidine, acetate, phosphate or citrate buffer), and/or a stabilizer agent (e.g. human albumin), etc. The antibody formulation can comprise pharmaceutically acceptable carriers, including, e.g., ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, sucrose, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, polyethylene-polyoxypropylene-block polymers, and polyethylene glycol.

[0066] The disclosed antibody formulations can further comprise a surfactant. In some embodiments, the surfactant is selected from the group consisting of Triton X-100, Tween 80, polysorbate 20, polysorbate 80, nonoxynol-9, polyoxamer, stearyl alcohol, sodium dodecyl sulfate, and sorbitan monostearate.

[0067] In some embodiments, the surfactant is polysorbate 80, i.e., polyoxyethylene (20) sorbitan monooleate, as represented by the formula:



Polysorbate 80 (PS-80) is available commercially from several commercial vendors, e.g., Alkest® TW 80 (Univar®), and Tween® 80 (Sigma-Aldrich®). Applicants have found that in some instances, controlling the concentration of PS-80 in the antibody formulation adds stability and reduces the amount of particle formation when stored for extended periods of time.

[0068] PS-80 can be about 0.01% to about 0.1%, about 0.02% to about 0.09%, about 0.02% to about 0.08%, about 0.03% to about 0.08%, about 0.04% to about 0.07%, about 0.05% to about 0.06%, about 0.02%, about 0.03%, about 0.04%, about 0.05%, about 0.06% about 0.07% of the antibody formulation, e.g., an antibody formulation comprising about 100 mg/mL to about 200 mg/mL antibody or about 150 mg/mL antibody. In some embodiments, PS-80 is about 0.05% in the antibody formulation.

[0069] In some embodiments, the antibody formulation further comprises histidine/histidine HCl buffer. The described antibody formulation comprises about 1 mM to about 100 mM, about 5 mM to about 80 mM histidine/histidine HCl buffer, about 10 mM to about 60 mM histidine/histidine HCl buffer, about 15 mM to about 50 mM histidine/histidine HCl buffer, about 15 mM to about 30 mM histidine/histidine HCl buffer, or about 25 mM histidine/histidine HCl buffer in the antibody formulations, e.g., an antibody formulation comprising 100 to 200 mg/mL antibody or about 150 mg/mL antibody. In one embodiment, histidine/histidine HCl buffer is about 25 mM in the antibody formulation.

[0070] Disclosed is an antibody formulation comprising: 150 mg/mL anifrolumab or an antigen binding fragment thereof; 50 mM lysine HCl; 130 mM uncharged excipient; 0.05% surfactant; 25 mM formulation buffer, wherein the formulation is at a pH of about 5.9.

[0071] Disclosed is an antibody formulation comprising: 150 mg/mL anifrolumab; 50 mM lysine HCl; 130 mM uncharged excipient; 0.05% surfactant; 25 mM formulation buffer, wherein the formulation is at a pH of about 5.9.

[0072] In additional embodiments, the invention is directed to an antibody formulation comprising: 150 mg/mL anifrolumab; 50 mM lysine HCl; 130 mM trehalose dihydrate; 0.05% polysorbate 80; 25 mM histidine/histidine HCl, wherein the formulation is at a pH of 5.9.

[0073] Disclosed is a stable antibody formulation comprising: (a) about 100 mg/mL to about 200 mg/mL of an antibody wherein the antibody is anifrolumab, (b) about 0.02% to about 0.1% polysorbate-80, (c) about 100 mM to about 160 mM trehalose, (d) about 40 mM to about 60 mM L-lysine HCl, (e) and 15-35 mM histidine/histidine HCl. In embodiments, the formulation pH is about 5.5 to about 6.5.

[0074] In further embodiments, the invention is directed to a stable antibody formulation comprising: (a) about 145 mg/mL to about 155 mg/mL of an antibody thereof, wherein the antibody is anifrolumab, (b) about 0.04% to about 0.08 polysorbate-80, (c) about 120-140 mM trehalose dihydrate, (d) about 45-55 mM L-lysine HCl, and (e) about 20-30 mM histidine/histidine HCl. In embodiments, the formulation pH is about 5.8 to about 6.1.

[0075] In further embodiments, the invention is directed to a stable antibody formulation comprising: (a) about 150 mg/mL of an antibody wherein the antibody is anifrolumab, (b) about 0.05% polysorbate-80, (c) about 130 mM trehalose dihydrate, (d) about 50 mM L-lysine HCl, and (e) about 25 mM histidine/histidine HCl. In embodiments, the formulation pH is about 5.9.

[0076] In some embodiments, the invention is directed to an antibody formulation comprising: 150 mg/mL anifrolumab 50 mM lysine HCl; 130 mM trehalose dihydrate; 0.05% polysorbate 80; 25 mM histidine/histidine HCl, wherein the

formulation is at a pH of about 5.9.

[0077] In additional embodiments, the invention is directed to an antibody formulation comprising: 150 mg/mL anifrolumab; 50 mM lysine HCl; 130 mM trehalose dihydrate; 0.05% polysorbate 80; 25 mM histidine/histidine HCl, wherein the formulation is at a pH of 5.9.

[0078] In some embodiments, various components can be omitted from the antibody formulation, or can be "substantially free" of that component. The term "substantially free" as used herein refers to an antibody formulation, said formulation containing less than 0.01%, less than 0.001%, less than 0.0005%, less than 0.0003%, or less than 0.0001% of the designated component.

[0079] The antibody formulations can have different osmolarity concentrations. Methods of measuring osmolarity of antibody formulations are known to those in the art, and can include, e.g., an osmometer (e.g., an Advanced Instrument Inc 2020 freezing point depression osmometer). In some embodiments, the formulation has an osmolarity of between 200 and 600 mosm/kg, between 260 and 500 mosm/kg, or between 300 and 450 mosm/kg.

[0080] The described antibody formulations can have various pH levels. In some embodiments, the pH of the antibody formulation is between 4 and 7, between 4.5 and 6.5, or between 5 and 6. The pH of the described antibody formulation is 5.0. In some embodiments, the pH of the antibody formulation is 6.0. Various means may be utilized in achieving the desired pH level, including, but not limited to the addition of the appropriate buffer.

[0081] The antibody formulations described herein have various viscosities. Methods of measuring viscosity of antibody formulations are known to those in the art, and can include, e.g., a rheometer (e.g., Anton Paar MCR301 Rheometer with either a 50 mm, 40 mm or 20 mm plate accessory). Viscosities were reported at a high shear limit of 1000 per second shear rate. The described antibody formulation can have a viscosity of less than 20 centipoise (cP), less than 18 cP, less than 15 cP, less than 13 cP, or less than 11 cP. The described antibody formulation can have a viscosity of less than 13 cP. One of skill in the art will appreciate that viscosity is dependent on temperature, thus, unless otherwise specified, the viscosities provided herein are measured at 25°C unless otherwise specified.

[0082] The injection force is correlated with the amount of resistance provided by the antibody formulation when administering the antibody formulation to a subject. The injection force will be dependent on the gauge of the administering needle, as well as temperature. The described antibody formulation can have an injection force of less than 15 N, 12 N, 10 N, or 8 N when passed through a 27 Ga spinal thin wall (STW) needle. In some embodiments, the antibody formulation has an injection force of less than 15 N, 12 N, 10 N, or 8 N when passed through a 29 Ga STW needle.

[0083] The antibody formulations of the present invention are an aqueous solution. The antibody formulation has not been subjected to freezing temperatures, and/or have not been frozen, i.e., they have remained in a liquid state. In some embodiments, the antibody in the antibody formulation has not been subjected to lyophilization.

[0084] As used herein, the term stability generally is related to maintaining the integrity or to minimizing the degradation, denaturation, aggregation or unfolding of a biologically active agent such as a protein, peptide or another bioactive macromolecule. As used herein, "improved stability" generally means that, under conditions known to result in degradation, denaturation, aggregation or unfolding, the protein (e.g., antibody such as anifrolumab), peptide or another bioactive macromolecule of interest maintains greater stability compared to a control protein, peptide or another bioactive macromolecule.

[0085] In some embodiments, stability refers to an antibody formulation having low to undetectable levels of particle formation. The phrase "low to undetectable levels of particle formation" as used herein refers to samples containing less than 1000 particles/mL, less than 700 particles/ml, less than 650 particles/ml, less than 500 particles/ml, less than 400 particles/ml, less than 200 particles/ml, less than 100 particles/ml or less than 1 particle/ml as determined by HIAC analysis or visual analysis, wherein the particles detected are greater than 10 micron in size, after storage at about 40°C for about 18 months. In some embodiments, no particles in the antibody formulation are detected, either by HIAC analysis or visual analysis.

[0086] In some embodiments, stability refers to reduced fragmentation of the antibody. In embodiments, the

fragmentation rate of the antibody, e.g., anifrolumab, in the formulations of the invention is about 2.0 to 4.0 percent per month for 12 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In embodiments, the fragmentation rate of the antibody, e.g., anifrolumab, in the formulations of the invention is about 2.0 to 4.0 percent per month for 6 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In embodiments, the fragmentation rate of the antibody, e.g., anifrolumab, in the formulations of the invention is about 2.0 to 4.0 percent per month for 2 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In embodiments, the fragmentation rate of the antibody, e.g., anifrolumab, in the formulations of the invention is about 3.0 to 4.0 percent per month for 2 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column.

[0087] In further embodiments, stability refers to reduced aggregation of the antibody. In embodiments, the aggregation rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 0.5 to 2.5% per month for 12 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In embodiments, the aggregation rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 0.5 to 2.5% per month for 6 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In embodiments, the aggregation rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 0.5 to 2.5% per month for 2 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In additional embodiments, the aggregation rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 1 to 2% per month for 2 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column.

[0088] In further embodiments, stability refers to reduced purity losses. In embodiments, the purity loss rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 3 to 5% per month for 12 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In embodiments, the purity loss rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 3 to 5% per month for 6 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column. In additional embodiments, the purity loss rate of the antibody formulations of the present invention containing, e.g., anifrolumab, is about 3.5 to 4.5% per month for 2 months as determined by HP-SEC analysis performed on an Agilent HPLC system with a TSK-Gel G3000 column.

[0089] One of skill in the art will appreciate that stability of a protein is dependent on other features in addition to the composition of the formulation. For example, stability can be affected by temperature, pressure, humidity, pH, and external forms of radiation. Thus, unless otherwise specified, stability referred to herein is considered to be measured at 40°C, one atmosphere pressure, 50% relative humidity, pH of 6.0, and normal background levels of radiation. Stability of the antibody in the antibody formulation can be determined by various means. In some embodiments, the antibody stability is determined by size exclusion chromatography (SEC). SEC separates analytes (e.g., macromolecules such as proteins and antibodies) on the basis of a combination of their hydrodynamic size, diffusion coefficient, and surface properties. Thus, for example, SEC can separate antibodies in their natural three-dimensional conformation from antibodies in various states of denaturation, and/or antibodies that have been degraded. In SEC, the stationary phase is generally composed of inert particles packed into a dense three-dimensional matrix within a glass or steel column. The mobile phase can be pure water, an aqueous buffer, an organic solvent, mixtures of these, or other solvents. The stationary-phase particles have small pores and/or channels which will only allow species below a certain size to enter. Large particles are therefore excluded from these pores and channels, but the smaller particles are removed from the flowing mobile phase. The time particles spend immobilized in the stationary-phase pores depends, in part, on how far into the pores they can penetrate. Their removal from the mobile phase flow causes them to take longer to elute from the column and results in a separation between the particles based on differences in their size.

[0090] In some embodiments, SEC is combined with an identification technique to identify or characterize proteins, or fragments thereof. Protein identification and characterization can be accomplished by various techniques, including but not limited chromatographic techniques, e.g., high-performance liquid chromatography (HPLC), immunoassays, electrophoresis, ultraviolet/visible/infrared spectroscopy, raman spectroscopy, surface enhanced raman spectroscopy, mass spectroscopy, gas chromatography, static light scattering (SLS), Fourier Transform Infrared Spectroscopy (FTIR), circular dichroism (CD), urea-induced protein unfolding techniques, intrinsic tryptophan fluorescence, differential scanning calorimetry, and/or ANS protein binding.

[0091] In some embodiments, protein identification is achieved by high-pressure liquid chromatography. Various

instruments, and apparatuses are known to those of skill in the art to perform HPLC. Generally HPLC involves loading a liquid solvent containing the protein of interest onto a separation column, in which the separation occurs. The HPLC separation column is filled with solid particles (e.g. silica, polymers, or sorbents), and the sample mixture is separated into compounds as it interacts with the column particles. HPLC separation is influenced by the liquid solvent's condition (e.g. pressure, temperature), chemical interactions between the sample mixture and the liquid solvent (e.g. hydrophobicity, protonation, etc.), and chemical interactions between the sample mixture and the solid particles packed inside of the separation column (e.g. ligand affinity, ion exchange, etc.).

[0092] In some embodiments, the SEC and protein identification occurs within the same apparatus, or simultaneously. For example, SEC and HPLC can be combined, often referred to as HP-SEC.

[0093] In some embodiments, the antibody formulation comprises about 100 mg/ml to about 200 mg/ml antibody wherein the antibody is anifrolumab, wherein said formulation is stable upon storage at about 40° C for 1 to 24 months. In some embodiments, the formulation is stable upon storage at about 25° C for 1 to 18 months. In some embodiments, the formulation is stable upon storage at about 5° C for 1 to 6 months. In some embodiments, the formulation is stable upon storage at about 5° C for 1 to 3 months. In some embodiments, the formulation is stable upon storage at about 5° C for 1 to 12 months. In some embodiments, the formulation is stable upon storage at about 5° C for at least 18 months. In some embodiments, the formulation is stable upon storage at about 5° C for at least 24 months, or 36 months.

[0094] The term "stable" can be relative and not absolute. Thus, in some embodiments the antibody is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by HP-SEC when the antibody is stored at 2°C to 8°C for 6 months. In some embodiments, the antibody is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by SEC HPLC when the antibody is stored at 2°C to 8°C for 12 months. In some embodiments, the antibody in the antibody formulation is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by HP-SEC when the antibody is stored at 2°C to 8°C for 18 months. In some embodiments, the antibody in the antibody formulation is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by SEC HPLC when the antibody is stored at 2°C to 8°C for 24 months.

[0095] In some embodiments, the antibody is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by HP-SEC when the antibody is stored at 23°C to 27°C for 3 months. In some embodiments, the antibody is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by HP-SEC when the antibody is stored at 23°C to 27°C for 6 months. In some embodiments, the antibody is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by HP-SEC when the antibody is stored at 23°C to 27°C for 12 months. In some embodiments, the antibody is stable if less than 20%, less than 15%, less than 10%, less than 5% or less than 2% of the antibody is degraded, denatured, aggregated or unfolded as determined by HP-SEC when the antibody is stored at 23°C to 27°C for 24 months.

[0096] In some embodiments the antibody is stable if less than 6%, less than 4%, less than 3%, less than 2% or less than 1% of the antibody is degraded, denatured, aggregated or unfolded per month as determined by HP-SEC when the antibody is stored at 40°C. In some embodiments the antibody is stable if less than 6%, less than 4%, less than 3%, less than 2% or less than 1% of the antibody is degraded, denatured, aggregated or unfolded per month as determined by HP-SEC when the antibody is stored at 5°C.

[0097] In embodiments the antibody is stable if 1%, 2%, 3%, 4%, 5% or 6% (or about 1% to 6%) of the antibody is degraded, denatured, aggregated or unfolded per month for 1-3 months, 1 to 6 months, 1 to 12 months, 1 to 18 months, or 1 to 24 months as determined by HP-SEC when the antibody is stored at 5°C.

[0098] In some embodiments, the antibody formulations of the present invention can be considered stable if the antibody exhibits very little to no loss of the binding activity of the antibody (including antibody fragments thereof) of the formulation compared to a reference antibody as measured by antibody binding assays known to those in the art, such as, e.g., ELISAs, etc., over a period of 8 weeks, 4 months, 6 months, 9 months, 12 months or 24 months. In some embodiments,

the antibody stored at about 40° C for at least 1 month retains at least 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, or at least about 99% of binding ability to an INFAR1 receptor polypeptide compared to a reference antibody which has not been stored. In some embodiments, the antibody stored at about 5° C for at least 6 months retains at least 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, or at least about 99% of binding ability to an INFAR1 receptor polypeptide compared to a reference antibody which has not been stored. In some embodiments, the antibody stored at about 40° C for at least 1 month retains at least 95% of binding ability to an INFAR1 receptor polypeptide compared to a reference antibody which has not been stored. In some embodiments, the antibody stored at about 5° C for at least 6 months retains at least 95% of binding ability to an INFAR1 receptor polypeptide compared to a reference antibody which has not been stored.

[0099] Applicants have found the antibody formulations provided herein result in greatly reduced particle formation as determined by visual inspection, micro-flow imaging (MFI), or size-exclusion chromatography (SEC).

[0100] In some embodiments, the formulation is substantially free of particles upon storage at about 40° C for at least 1 month as determined by visual inspection. In some embodiments, the formulation is substantially free from particles upon storage at about 5° C for at least 6 months, at least 9 months, at least 12 months, at least 15 months, at least 18 months, at least 24 months, or at least 36 months as determined by visual inspection.

[0101] In some embodiments, the antibody formulation of the present invention can be used for pharmaceutical purposes. Antibodies used in pharmaceutical applications generally must have a high level of purity, especially in regard to contaminants from the cell culture, including cellular protein contaminants, cellular DNA contaminants, viruses and other transmissible agents. See "WHO Requirements for the use of animal cells as in vitro substrates for the production of biologicals: Requirements for Biological Substances No. 50. " No. 878. Annex 1, 1998. In response to concerns about contaminants, The World Health Organization (WHO) established limits on the levels of various contaminants. For example, the WHO recommended a DNA limit of less than 10 ng per dose for protein products. Likewise, the United States Food and Drug Administration (FDA) set a DNA limit of less than or equal to 0.5 pg/mg protein. The described antibody formulations meeting or exceeding contaminant limits as defined by one or more governmental organizations, e.g., the United States Food and Drug Administration and/or the World Health Organization.

[0102] In some embodiments, the antibody formulation described herein is pharmaceutically acceptable. "Pharmaceutically acceptable" refers to an antibody formulation that is, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity or other complications commensurate with a reasonable benefit/risk ratio.

[0103] Purity of the antibody formulations can vary. In some embodiments, the therapeutic antibody of interest, e.g., anti-IFNAR1 antibody, is greater than 90% (wt/wt) of the total polypeptides present in the antibody formulation. In some embodiments, the therapeutic antibody of interest, e.g., anti- IFNAR1 is greater than 95% (wt/wt), 98% (wt/wt), 99% (wt/wt), 99.5% (wt/wt) or 99.9% (wt/wt) of the total polypeptide present in the antibody formulation.

[0104] Disclosed are methods of treating a type I IFN-mediated disease or disorder in a subject in need thereof, by administering a therapeutically effective amount of the antibody formulation described herein. In embodiments, the disease or disorder is selected from the group consisting of systemic lupus erythematosus (SLE), insulin dependent diabetes mellitus, inflammatory bowel disease, multiple sclerosis, psoriasis, autoimmune thyroiditis, rheumatoid arthritis, glomerulonephritis, scleroderma, myositis and lupus nephritis. In further embodiments, the disease or disorder is an inflammatory bowel disease such as, Crohn's disease, ulcerative colitis, and Celiac's disease. In further embodiments, the disease or disorder is a pulmonary disease or disorder, such as systemic lupus erythematosus.

[0105] The antibody formulation disclosed herein can be administered to a subject through various means. In some embodiments, the antibody formulation is suitable for parenteral administration, e.g., via inhalation (e.g., powder or aerosol spray), transmucosal, intravenous, subcutaneous, or intramuscular administration. In some embodiments, the formulation is an injectable formulation. Also disclosed is a sealed container comprising any of the antibody formulations as described herein.

[0106] Disclosed are various pharmaceutical dosage forms. Various dosage forms could be applicable to the formulations provided herein. See, e.g., Pharmaceutical Dosage Form: Parenteral Medications, Volume 1, 2nd Edition. A pharmaceutical unit dosage can comprise the antibody formulation in a suitable container, e.g. a vial or syringe A

pharmaceutical unit dosage form can comprise an intravenously, subcutaneously, or intramuscularly delivered antibody formulation. A pharmaceutical unit dosage form can comprise aerosol delivered antibody formulation. A pharmaceutical unit dosage form can comprise a subcutaneously delivered antibody formulation. A pharmaceutical unit dosage form can comprise an aerosol delivered antibody formulation. A pharmaceutical unit dosage form can comprise an intranasally administered antibody formulation.

[0107] The antibody formulations can be prepared as unit dosage forms by preparing a vial containing an aliquot of the aqueous antibody formulation for a one-time use. For example, a unit dosage per vial may contain 1 ml, 2 ml, 3 ml, 4 ml, 5 ml, 6 ml, 7 ml, 8 ml, 9 ml, 10 ml, 15 ml, or 20 ml of different concentrations of an antibody that specifically binds to IFNAR1 receptor ranging from about 0.1 mg/ml to about 300 mg/ml. If necessary, these preparations can be adjusted to a desired concentration by adding a sterile diluent to each vial. In a specific embodiment, the aqueous antibody formulations are formulated into single dose vials as a sterile liquid that contains about 2 mg/mL to about 20 mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody is anifrolumab. In another specific embodiment, the aqueous antibody formulations of the present invention are formulated into single dose vials as a sterile liquid that contains about 100 mg/mL to about 200 mg/mL of an antibody wherein the antibody is anifrolumab, about 45 to 55 mM lysine HCl, about 0.04% to about 0.08% polysorbate-80, about 120 mM to about 140 mM trehalose, and about 20-30 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 6. In one embodiment, the antibody of the invention is supplied at 140 to 160 mg/ml in 3 cc USP Type I borosilicate amber vials (West Pharmaceutical Services-Part No. 6800-0675). In another embodiment, the antibody of the invention is supplied at 150 mg/ml in 3 cc USP Type I borosilicate amber vials.

[0108] The antibody formulations of the present invention can be prepared as unit dosage forms by preparing a pre-filled syringe containing an aliquot of the aqueous antibody formulation for a one-time use. For example, a unit dosage per pre-filled syringe may contain 0.1 ml, 0.2 ml, 0.3 ml, 0.4 ml, 0.5 ml, 0.6 ml, 0.7 ml, 0.8 ml, 0.9 ml, 1 ml, 2 ml, 3 ml, 4 ml, 5 ml, 6 ml, 7 ml, 8 ml, 9 ml, 10 ml, 15 ml, or 20 ml of different concentrations of an antibody or antigen binding fragment thereof that specifically binds to an IFNAR1 polypeptide ranging from about 140 mg/ml to about 160 mg/ml. In a specific embodiment, the aqueous antibody formulations of the present invention are formulated into single dose pre-filled syringes as a sterile liquid that contains about 145 mg/mL to about 155 mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody is anifrolumab, about 45 to 55 mM lysine HCl, about 0.04% to about 0.08% polysorbate-80, about 120 mM to about 140 mM trehalose, and about 20-30 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 6. In a specific embodiment, the aqueous antibody formulations of the present invention are formulated into single dose pre-filled syringes as a sterile liquid that contains about 145 mg/mL to about 155 mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody wherein the antibody is anifrolumab, about 45 to 55 mM lysine HCl, about 0.04% to about 0.08% polysorbate-80, about 120 mM to about 140 mM trehalose, and about 20-30 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 6.

[0109] Various dosage amounts can be administered in a single use. For example, in some embodiments 0.1 mg, 0.2 mg, 0.3 mg, 0.4 mg, 0.5 mg, 0.6 mg, 0.7 mg, 0.8 mg, 0.9 mg, 1.0 mg, 1.1 mg, 1.2 mg, 1.3 mg, 1.4 mg, 1.5 mg, 1.6 mg, 1.7 mg, 1.8 mg, 1.9 mg, 2 mg, 3 mg, 4 mg, 5 mg, 6 mg, 7 mg, 8 mg, 9 mg, 10 mg, 12 mg, 14 mg, 16 mg, 18 mg, 20 mg, 30 mg, 40 mg, 50 mg, 70 mg, or 100 mg of antibody can be administered in a single dose.

[0110] Various types of syringes can be used. The syringe can be filled with the antibody formulation immediately prior to administration to a subject, e.g., less than 1 week, 1 day, 6 hours, 3 hours, 2 hours, 1 hour, 30 minutes, 20 minutes, or 10 minutes prior to administration to a subject. In some embodiments, the syringe is filled with the antibody formulation at the point of retail sale, or by the facility for which treatment of the subject occurs. In some embodiments, the syringe is pre-filled, e.g., the syringe is filled with the antibody formulation greater than 1 day, 2 days, 4 days, 1 week, 2 weeks, 1 month, 2 months, 3 months, 6 months, 12 months, 18 months, 24 months, 3 years, or 4 years prior to administration to a subject. In some embodiments, the pre-filled syringe comprises a needle, e.g., a 27G regular wall needle, a 27G thin wall needle, a 29G regular wall needle, or a 29G thin wall needle. In some embodiments, the pre-filled syringe comprises a 27 G spinal thin wall needle.

[0111] In some embodiments, any syringe suitable for administration to the desired subject can be used. In some embodiments, the syringe is a plastic syringe or a glass syringe. In some embodiments, the syringe is made of materials that are substantially free from tungsten. In some embodiments, the syringe is coated with silicone. In some embodiments, the pre-filled syringe comprises a plunger having a fluoropolymer resin disk. Examples of syringes can include, but are not limited to BD Hypak™ SCF 1 MLL 27G1/2-5B BD260L WL, 0.4 mg silicon oil MDN with 27 G STW

needle; Hypak™ for Biotech 1 ml Long (Becton Dickinson), with a Becton Dickinson Hypak 1 mL long plunger stopper 4023 Flurotec Daikyo Si1000 (Catalog #47271919); C3Pin; Hypak™ for Biotech 0.8 mg silicone oil (Becton Dickinson); and CZ syringes (West, Catalog # 19550807).

[0112] The aqueous antibody formulations of the present invention may be sterilized by various sterilization methods, including sterile filtration, radiation, etc. In a specific embodiment, the difiltered antibody formulation is filter-sterilized with a presterilized 0.2 micron filter. Sterilized aqueous antibody formulations of the present invention may be administered to a subject to prevent, treat and/or manage an immune response, e.g. an inflammatory response.

[0113] The invention further provides pre-filled syringes comprising the antibody formulations of the captioned invention.

[0114] The invention further provides pre-filled syringes comprising the antibody formulations of the captioned invention. In some embodiments, the pre-filled syringe comprises (a) about 100 mg/mL to about 200 mg/mL anifrolumab about 40 to 60 mM lysine HCl, about 100 mM to about 160 mM trehalose dihydrate, 0.02% to about 0.1% polysorbate 80, and about 15 mM to about 35 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 5.5 to about 6.5.

[0115] The invention further provides pre-filled syringes comprising the antibody formulations of the captioned invention. In some embodiments, the pre-filled syringe comprises (a) about 145 mg/mL to about 155 mg/mL anifrolumab about 45 to 55 mM lysine HCl, 120 mM to about 140 mM trehalose dihydrate, 0.04% to about 0.08% polysorbate 80, and 20 mM to about 30 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 5.8 to 6.1.

[0116] The invention further provides pre-filled syringes comprising the antibody formulations of the captioned invention. In some embodiments, the pre-filled syringe comprises (a) about 150mg/mL of an antibody wherein the antibody is anifrolumab, about 50 mM lysine HCl, about 0.05% polysorbate-80, about 130 mM trehalose dihydrate, and about 25 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 5.9.

[0117] The invention further provides pre-filled syringes comprising the antibody formulations of the captioned invention. In some embodiments, the pre-filled syringe comprises (a) about 150mg/mL of anifrolumab, about 50 mM lysine HCl, about 0.05% polysorbate-80, about 130 mM trehalose dihydrate, and about 25 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 5.9.

[0118] In a specific embodiment, the antibody formulations of the present invention are formulated into single dose pre-filled syringes as a sterile liquid that contains about 145 mg/mL to about 155 mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody wherein the antibody is anifrolumab, about 45 to 55 mM lysine HCl, about 0.04% to about 0.08% polysorbate-80, about 120 mM to about 140 mM trehalose, and about 20-30 mM histidine/histidine HCl. In embodiments, the pH of the formulation is about 6.

[0119] In embodiments, the invention is directed to a prefilled syringe comprising an antibody formulation of the present invention, wherein the pre-filled syringe has an average glide force of between 1 and 20 N when equipped with a 27 Gauge spinal thin-wall (STW) needle. In embodiments, the invention is directed to a prefilled syringe comprising an antibody formulation of the present invention, wherein the pre-filled syringe has an average glide force of between 5 and 15 N when equipped with a 27 Gauge spinal thin-wall (STW) needle. In embodiments, the invention is directed to a prefilled syringe comprising an antibody formulation of the present invention, wherein the pre-filled syringe has an average glide force of about 8 N when equipped with a 27 Gauge spinal thin-wall (STW) needle.

[0120] Described is a kit comprising any of the antibody formulations described herein, the containers described herein, the unit dosage forms described herein, or the pre-filled syringe described herein.

[0121] Described is a method of producing a stable antibody formulation comprising an antibody, the method comprising: (a) purifying an antibody or antigen binding fragment thereof to about 100 mg/mL to about 200 mg/mL, wherein the antibody is anifrolumab; and (b) placing the isolated antibody in a stabilizing formulation to form the stable antibody formulation, wherein the resulting stable antibody formulation comprises: (i) about 100 mg/mL to about 200 mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody is anifrolumab, and (ii) about 25 mM to about 130 mM lysine or a lysine salt; (iii) about 100 mM to about 150 mM trehalose (iv) about 0.02% to about 0.1% polysorbate-80.

[0122] Also described is a method of producing a stable antibody formulation comprising an antibody, the method

comprising: (a) purifying an antibody or antigen binding fragment thereof to about 100 mg/mL to about 200 mg/mL, wherein the antibody is anifrolumab; and (b) placing the isolated antibody in a stabilizing formulation to form the stable antibody formulation, wherein the resulting stable antibody formulation comprises: about 100 mg/mL to about 200 mg/mL of the antibody; about 45 mM to about 55 mM lysine HCl or a lysine salt; about 100 mM to about 150 mM uncharged excipient; about 0.02% to about 0.1% of a surfactant; and a formulation buffer.

[0123] Also described is a method of producing a stable antibody formulation comprising an antibody, the method comprising: (a) purifying an antibody or antigen binding fragment thereof to about 145 mg/mL to about 155 mg/mL, wherein the antibody is anifrolumab; and (b) placing the isolated antibody in a stabilizing formulation to form the stable antibody formulation, wherein the resulting stable antibody formulation comprises about 145 mg/mL to about 155 mg/mL of the antibody, about 25 mM to about 130 mM of lysine or a lysine salt; about 120 mM to about 140 mM trehalose dihydrate; about 0.04% to about 0.08% polysorbate 80; about 20 mM to about 30 mM histidine/histidine HCl, wherein the formulation is at a pH of about 5.8 to 6.1.

[0124] Also described is a method of making a stable antibody formulation, the method comprising (a) purifying an antibody or antigen binding fragment thereof to about 150 mg/mL, wherein the antibody is anifrolumab; and (b) placing the isolated antibody in a stabilizing formulation to form the stable antibody formulation, wherein the resulting stable antibody formulation comprises: (i) about 150 mg/mL mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody is anifrolumab, and (ii) about 50 mM lysine HCl; (iii) about 130 mM trehalose, (iv) about 0.05% % polysorbate-80.

[0125] Described is a method of producing a reconstituted antibody or antigen-binding fragment thereof formulation comprising anifrolumab, the method comprising: (a) purifying the antibody from a cell culture; (b) lyophilizing the isolated antibody; (c) adding the lyophilized antibody to a aqueous solution to form a reconstituted antibody formulation, wherein the reconstituted antibody formulation comprises: (i) about 100 mg/mL to about 200 mg/mL of an antibody or antigen binding fragment thereof, wherein the antibody is anifrolumab, and (ii) about 45 to about 55 mM lysine HCl.

[0126] Described is an antibody formulation comprising an antibody or an antigen binding fragment thereof wherein the antibody is anifrolumab, wherein the antibody formulation is essentially free of particles. In some embodiments, the term "essentially free of particles" refer to the absence of visible particles when viewed under a light box. In some embodiments, the term "essentially free of particles" is synonymous with the phrase "low to undetectable levels of particle formation" as described previously. In some embodiments, essentially free of particles refers to samples containing less than 30 particles/mL, less than 20 particles/ml, less than 20 particles/ml, less than 15 particles/ml, less than 10 particles/ml, less than 5 particles/ml, less than 2 particles/ml or less than 1 particle/ml wherein the particles are greater than 25 μ m and the particle count is determined by HIAC analysis or visual analysis. In some embodiments, essentially free of particles refers to samples containing 1 to 50 particles/mL, 2 to 40 particles/ml, 3-30 particles/ml, 4 to 25 particles/ml, or 5 to 20 particles/ml wherein the particles are greater than 25 μ m and the particle count is determined by HIAC analysis or visual analysis. In some embodiments, the term "visible particles" refers to particles greater than 25 μ m.

[0127] In some embodiments, essentially free of particles refers to samples containing 1 to 200 particles/mL, 10 to 150 particles/ml, about 30 particles/ml to about 100 particles/ml, or 40 to 80 particles/ml, wherein the particles are greater than 5 μ m and the particle count is determined by HIAC analysis or visual analysis. In some embodiments, the term "visible particles" refers to particles greater than 5 μ m. In some embodiments, no particles in the antibody formulation are detected, either by HIAC analysis or visual analysis.

[0128] Described is an antibody formulation comprising an antibody or antigen binding fragment thereof wherein the antibody is anifrolumab, wherein the antibody formulation is essentially free of particles for at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, at least 8 months, at least 10 months, at least 12 months, or at least 18 months when stored at 38 ° to 42° C. Described is an antibody formulation comprising an antibody or antigen binding fragment thereof, wherein the antibody is anifrolumab, wherein the antibody formulation is essentially free of particles for at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, at least 8 months, at least 10 months, at least 12 months, at least 18 months, at least 24 months, at least 30 months, at least 36 months, or at least 48 months when stored at 2-6° C.

[0129] Also described is a method of purifying an antibody or antigen-binding fragment thereof wherein the antibody is anifrolumab, the method comprising (i) obtaining a cell culture comprising the antibody, (ii) performing affinity

chromatography on the antibody, (iv) performing cation exchange on the antibody, (v) performing mixed mode chromatography on the antibody.

[0130] Described is a method of purifying an antibody or antigen binding fragment thereof wherein the antibody is anifrolumab, the method comprising (i) obtaining a cell culture comprising the antibody, (ii) binding the antibody to a Protein A column, (iii) eluting the antibody from the Protein A column, (iv) performing cation exchange on the antibody, (v) performing mixed mode chromatography on the antibody. In some embodiments, the method of purifying an antibody further comprises a viral inactivation process. In some embodiments, the viral inactivation step is performed by lowering the pH to less than 4.0. In some embodiments, the method further comprises a diafiltration process. In some embodiments, the method further comprises a filtration process. In some embodiments, the filtration process is sufficient to remove active virus particles.

[0131] Also described is a method of treating a patient. In some embodiments, the method comprises administering the antibody formulations described herein, the containers described herein, the unit dosage forms described herein, or the pre-filled syringe described herein to a subject in need thereof.

[0132] Antibody formulations can be suitable for treatment of pulmonary disease or disorder by administering the antibody formulation described herein. In some embodiments, the invention is directed to a method of treating a patient with an eosinophilic disease or disorder by administering the antibody formulation described herein. Also described is a method of treating a pulmonary disease or disorder in a subject, the method comprising administering the antibody formulations described herein. In some embodiments, the invention is directed to a method of treating an eosinophilic disease or disorder in a subject, the method comprising administering the antibody formulations described herein. Described is the treatment of pulmonary diseases or disorders e.g., asthma, COPD, eosinophilic asthma, combined eosinophilic and neutrophilic asthma, aspirin sensitive asthma, allergic bronchopulmonary aspergillosis, acute and chronic eosinophilic bronchitis, acute and chronic eosinophilic pneumonia, Churg-Strauss syndrome, hypereosinophilic syndrome, drug, irritant and radiation-induced pulmonary eosinophilia, infection-induced pulmonary eosinophilia (fungi, tuberculosis, parasites), autoimmune-related pulmonary eosinophilia, eosinophilic esophagitis, or Crohn's disease or combination thereof in a subject, the method comprising administering the antibody formulations described herein. Described is the treatment of asthma in a subject, the method comprising administering the antibody formulations described herein. Also described is the treatment of COPD in a subject, the method comprising administering the antibody formulations described herein.

[0133] In some embodiments, a therapeutically effective amount of the antibody formulations described herein is administered to treat a condition. As used herein, the term "therapeutically effective amount" refers to the amount of a therapy (e.g., an antibody that immunospecifically binds to an IFNAR1 receptor polypeptide), that is sufficient to reduce the severity of a disease or disorder (e.g., a disease or disorder characterized by aberrant expression and/or activity of an IFNAR1 polypeptide, a disease or disorder characterized by aberrant expression and/or activity of an IFNAR1 receptor or one or more subunits thereof, an autoimmune disease, an inflammatory disease, a proliferative disease, or an infection or one or more symptoms thereof), reduce the duration of a condition, ameliorate one or more symptoms of such a disease or disorder, prevent the advancement of such a disease or disorder, cause regression of such a disease or disorder, or enhance or improve the therapeutic effect(s) of another therapy. In some embodiments, the therapeutically effective amount cannot be specified in advance and can be determined by a caregiver, for example, by a physician or other healthcare provider, using various means, for example, dose titration. Appropriate therapeutically effective amounts can also be determined by routine experimentation using, for example, animal models.

[0134] The terms "therapies" and "therapy" can refer to any protocol(s), method(s), and/or agent(s) that can be used in the prevention, treatment, management, or amelioration of a disease or disorder (e.g., a disease or disorder characterized by aberrant expression and/or activity of an IFNAR1 polypeptide, a disease or disorder characterized by aberrant expression and/or activity of an IFNAR1 receptor or one or more subunits thereof, an autoimmune disease, an inflammatory disease, a proliferative disease, or an infection (preferably, a respiratory infection) or one or more symptoms thereof). In certain embodiments, the terms "therapy" and "therapy" refer to biological therapy, supportive therapy, and/or other therapies useful in treatment, management, prevention, or amelioration of such a disease or disorder or one or more symptoms known to skilled medical personnel.

[0135] As used herein, the term "therapeutic protocol" refers to a regimen for dosing and timing the administration of one or more therapies (e.g., therapeutic agents) that has a therapeutic effective.

[0136] The route of administration of the antibody formulation can be via, for example, oral, parenteral, inhalation or topical modes of administration. The term parenteral as used herein includes, e.g., intravenous, intraarterial, intraperitoneal, intramuscular, subcutaneous, rectal or vaginal administration. The antibody can be an anti-IFNAR1 antibody and the route of administration is intramuscular injection. While all these forms of administration are clearly contemplated in some embodiments, the antibody formulation is suitable for administration via injection, in particular for intravenous or intraarterial injection or drip.

[0137] In some embodiments, the compositions and methods enable a manufacturer to produce an antibody formulation suitable for administration to a human in a more efficient manner, either by reducing costs, reducing method steps, reducing opportunities for error, reducing opportunities for introduction of unsafe or improper additives, reducing waste, increasing storage time, etc.

EXAMPLES

[0138] The invention is now described with reference to the following examples. These examples are illustrative only and the invention should in no way be construed as being limited to these examples

Example 1

Materials and Methods

Materials

[0139] All the materials used were of USP or Multicompodial grade. All the solutions and buffers were prepared using USP or HPLC water and were filtered before further use. Samples for stability studies were prepared under aseptic conditions in the Biosafety Cabinet Hood (BSC). Bulk material was stored at 2-8°C. Stability studies were carried out using the supplies listed in TABLE 3.

TABLE 3

Component	Purification process
MEDI-546	From cNS0 19B4 clone cell line
Vials	3 cc Fiolax vials (Schott, p/n: 1230392)
Stoppers	West 13 mm HBR Teflon2 442/50 grey rubber stopper (item#: 10124671)
Overseals	13 mm aluminium TruEdge® Flip-Off®

Protein Concentration Determination

[0140] Protein concentrations were determined by measuring absorbance at 280 nm with an Agilent UV-Vis spectrophotometer using a procedure adapted from SOP DV-6233. A measured extinction coefficient of 1.39 (mg/mL)-~cm ~ was used. For the samples, density correction factors were applied as per TD-0025 for sugar or non-sugar containing formulations.

Cone and Plate Viscosity Measurements

[0141] Viscosities were measured using an Anton Paar MCR301 Rheometer with cone and plate accessory. To minimize volume required a 20 mm cone was used with single replicate measurements for screening purposes. Results were reported at the high-shear limit of 1000 per second shear rate.

Purity Determination by Size Exclusion Chromatography (HPSEC)

[0142] SEC analysis was performed on an Agilent HPLC system with a TSK-Gel G3000 as per the current Formulation Sciences guidelines.

Visual Appearance

[0143] Visual inspection of the samples were performed by examining the samples in their respective container for particles, color, and clarity using particle standards and guidance adapted from the standard operating procedure: Visual Appearance Evaluation of Protein Drug Substance and Drug Product.

Studies to Assess Impact of Excipient Levels on Stability and Viscosity

[0144] In order to study the impacts of concentration, trehalose level and lysine HCl level (factors) on stability and viscosity (responses) a design of experiments approach was utilized. A Box Behnken design was prepared using JMP9.1 software (SAS, Inc., Cary, NC). The design included factors: concentration (from 100 to 200 mg/mL); trehalose level (from 0 to 211 mM); and lysine HCl level (from 25 to 130 mM). The pH was set to 6.0 based on the previous formulation stability data for the cycle 1 formulations. The polysorbate level was set at 0.02%. The low lysine level was defined at 25 mM based on preliminary high-throughput viscosity screening studies where a large drop in viscosity over the range 0 to 25 mM lysine HCl was observed at high concentrations (see Example 2). Stability was evaluated by performing an accelerated stability study at 40°C with assessment of visual appearance and purity by HP-SEC (monomer loss, aggregation, fragmentation) as readouts. Subvisible particle counts at 5°C after 1.8 months were also measured but not further analyzed because no meaningful differentiation was observed between samples.

Robustness and Impact of Lysine HCl Level on Viscosity Profile

[0145] A stock of MEDI-546 at ~ 200 mg/mL in 25 mM histidine/histidine HCl, 25 mM lysineHCl (note: this is the low lysine HCl bracket), 130 mM trehalose dihydrate, pH 6.0 was prepared and used to prepare a dilution series formulated with 0.02% polysorbate 80. The concentration and viscosities were measured and plotted. The nominal formulation condition (50 mM lysine HCl) was also prepared and measured. A 0.5 M lysineHCl stock was prepared as used to spike the MEDI-546 stock to the nominal 50 mM lysine level. The same dilution and formulation was performed and the viscosities and concentrations were measured.

Assessment of Functionality of the Lead Formulation in 27Ga STWPrefilled Syringes

[0146] Prefilled syringes ("PFS") (BD Hypak SCF 1MLL 27G1/2-SB); 0.4 mg silicone oil MDN with 27 G STW thin-wall needle) were filled with MEDI-546 at 150 mg/mL in 25 mM histidine/histidineHCl, 50 mM lysine HCl, 130 mM trehalose dihydrate, 0.05% polysorbate 80, pH 5.9. The viscosity of this lot was measured as 9.4 mPas and the concentration as 150.7 mg/mL. BD Hypak SCF 1MLL 4023 FLUR Daikyo SI1000 stoppers were applied with vacuum stoppering. An Instron 5542 (Norwood, MA 02062) was used to measure the gliding performance at 260 mm/min. Additionally, three analysts assessed the time to inject with and without arthritis simulation gloves (Arthritis gloves from Georgia Tech Research Institute and Injection trainer from Limbsnthings, UK).

Example 2**High-Throughput Viscosity Screening Results Summaries**

[0147] High-throughput viscosity screening was performed with nanoparticles. In brief, nanoparticles of known size were measured in water (known viscosity) and the samples (unknown viscosity) and the ratio used to determine the unknown viscosity of the samples. These data were collected for screening and trending purposes and not for absolute viscosity determination.

[0148] The HTS screening results for viscosity vs. pH is shown in Figure 1. The results suggested that viscosity increased with pH. Based on this potential impact of pH on the viscosity and possible Dorman-type effects during TFF we made the judgment that the formulation pH should be selected at 5.9. For PFS development additional robustness around the impact of pH will need to be performed. The HTS screening results for viscosity vs. lysine HCl (Figure 2) showed a large drop in viscosity from zero to 25 mM lysine HCl. Hands-on laboratory observations of sample preparation also indicated that without lysine samples were more viscous and harder to handle at higher concentrations. For this reason the experiments started with 25 mM lysine HCl as the low level.

Example 3

Overall Assessment of Stability and Viscosity Results for the Formulations

[0149] TABLE 4 gives a summary of viscosity and accelerated stability results for the antibody formulations. TABLE 5 gives a summary of the subvisible particles results (by HIAC) for Formulations after 1.8 months at 5°C. An overall assessment of the results show that the purity loss rates at 40°C were all acceptable, ranging from 3.8 to 4.8 percent per month. The fragmentation rate ranged from 2.8 to 3.4 percent per month and the aggregation rate ranged from 0.7 to 2.0 percent per month. The visual inspections after 1 month at 40°C indicate similar performance for all formulations (all were standard 1 for particles, <III or II for clarity and Y6 for color). The subvisible particle (HIAC) data showed that all samples had less than 670 particles per mL greater than 10 microns in size. The viscosity ranged from 2.8 to 39.7 mPas for all of the formulations. Several of the formulations at 150 mg/mL had acceptable viscosity values.

[0150] Additional analysis of the data in the Examples below shows how the data was used and interpreted to select an appropriate and robust formulation that maximizes stability while providing an acceptable viscosity.

TABLE 4 - Summary of Viscosity and Accelerated Stability Results for Formulations

DOE ID	LysineHCl (mM)	Trehalose (mM)	Target Cone. (mg/mL)	Viscosity (mPa.s)	Aggregation Rate at 40°C (%/month)	Fragmentation Rate at 40°C (%/month)	Purity Loss Rate at 40°C (%/month)	Visual Appearance After & month at 40°C (particles, color, clarity)
1	130	211	150	13.8	1.1	2.8	3.8	1, Y6, <III
2	77.5	105.5	150	11.5	1.4	2.9	4.4	1, Y6, <III
5	130	105.5	200	25.9	15	2.9	4.4	1, Y6, <III
6	77.5	0	100	2.8	1.3	3.1	4.5	1, Y6, <III
7	25	105.5	200	29.6	1.9	2.8	4.7	1, Y6, <III
8	130	0	150	9.8	1.5	3.0	4.5	1, Y6, <III
9	77.5	105.5	150	12.6	1.4	2.9	4.3	1, Y6, <III
10	77.5	105.5	150	12.8	1.5	2.9	4.4	1, Y6, II
11	130	105.5	100	4.0	1.1	3.0	4.1	1, Y6, <III
12	77.5	0	200	39.7	2.0	2.8	4.8	1, Y6, <III
13	25	0	150	12.1	1.7	2.8	4.6	1, Y6, <III
14	25	211	150	16.7	1.3	2.9	4.2	1, Y6, II
15	77.5	211	200	33.9	1.3	2.9	4.2	1, Y6, <III

DOE ID	LysineHCl (mM)	Trehalose (mM)	Target Cone. (mg/mL)	Viscosity (mPa.s)	Aggregation Rate at 40°C (%/month)	Fragmentation Rate at 40°C (%/month)	Purity Loss Rate at 40°C (%/month)	Visual Appearance After & month at 40°C (particles, color, clarity)
16	25	105.5	100	4.1	0.9	3.4	4.3	1, Y6, <III
17	77.5	211	100	5.1	0.7	3.3	4.0	1, Y6, <III

TABLE 5 - Summary of Subvisible Particles Results for Formulations after 1.8 months at 5°C

DOE ID	LysineHCl (mM)	Trehalose (mM)	Target Conc. (mg/mL)	SVP $\geq 2 \mu\text{m}$ (HIAC) per mL	SVP $\geq 10\mu\text{m}$ (HIAC) per mL	SVP $\geq 25\mu\text{m}$ (HIAC) per mL
1	130	211	150	880	20	0
2	77.5	105.5	150	2640	80	20
5	130	105.5	200	3493	0	0
6	77.5	0	100	4120	80	0
7	25	105.5	200	3627	53	0
8	130	0	150	4540	60	20
9	77.5	105.5	150	5940	220	40
10	77.5	105.5	150	5640	20	0
11	130	105.5	100	1387	67	0
12	77.5	0	200	1680	53	0
13	25	0	150	1691	40	0
14	25	211	150	3980	20	0
15	77.5	211	200	44853	667	53
16	25	105.5	100	5800	93	13
17	77.5	211	100	5440	120	13

Example 4

Analysis of Results

[0151] TABLE 6 summarizes the interpretation and conclusions drawn from the data:

TABLE 6 - Output for Significance of Factors on Measured Responses

	Concentration	Lysine HCl Level	Trehalose Level
Viscosity Results	Significant Impact: Viscosity increased with concentration.	Not significant (for range tested above 25 mM)	Not significant. Minor trend for increase in viscosity with higher trehalose.
Viscosity Conclusions	Data suggests 150 mg/mL is feasible (<20 mPas) but that 200 mg/mL will have too high a viscosity (>30 mPas). Lysine was shown to decrease viscosity in prior work. A bracket with 25 mM as the low end and 50 mM as the nominal level would provide appropriate robustness.		
Aggregation Rate	Significant: aggregation increased with concentration	Not significant	Significant: trehalose decreased aggregation by a small amount.
Aggregation Rate Conclusions	Strategy to maximize trehalose with isotonic constraint (little incremental benefit of much higher trehalose was observed to justify hypertonicity) will minimize aggregation.		

	Concentration	Lysine HCl Level	Trehalose Level
Fragmentation Rate	Significant: fragmentation slightly decreased with concentration. Rate ranged from 2.8 to 3.4 percent per month at 40°C.	Not significant	Not significant
Fragmentation Rate Conclusions	Fragmentation is not likely an issue over entire knowledge space at 5°C. Differences in rate were very minor and could be artifacts of method variability due to SEC shoulder 'pop' on stability.		
Overall Conclusions.	Overall, a lead formulation at 150 mg/mL with 50 mM lysineHCl to minimize viscosity and 130 mM trehalose to minimize aggregation will achieve isotonic condition when 25 mM histidine buffer is present. Polysorbate level will require additional optimization.		

[0152] Based on the analysis results, a formulation having MEDI-546 at 150 mg/mL in 25 mM histidine/histidine HCl, 50 mM lysine HCl, 130 mM trehalose dehydrate, 0.05% polysorbate 80 and a pH of 5.9 was further assessed.

Example 6

Confirmation of Viscosity and Syringe Functionality for Antibody Formulation

[0153] Ideally the glide force for a prefilled syringe should be as low as possible to ensure functionality. In this work we are targeting a viscosity of less than 20 mPas and a glide force of less than 15 N. Prior experience has been for formulations with nominal viscosities less than 15 mPas to have acceptable gliding performance when equipped with 27 Gauge thin-walled needles. Up to 20 mPas may be feasible in 27 Ga STW PFS. The formulation and PFS should include robustness to account for variability in the syringes and the formulation.

[0154] In this section we document additional data demonstrating the robustness of a formulation viscosity vs. concentration curve when compared with a lower bracket (25 mM) lysineHCl formulation. Gliding performance and injection time feasibility results with the nominal formulation in 27 Ga STW PFS are also assessed in this section.

Robustness and Impact of Lysine HCl Level on Viscosity Profile

[0155] A dilution series of MEDI-546 over 100 to 200 mg/mL was prepared in two formulations. They contained MEDI-546 in 25 mM histidine/histidine-HCl, 130 mM trehalose dehydrate with 0.02% polysorbate 80 at pH 6.0 with either 25 mM or 50 mM lysine HCl. The viscosity curves are shown in Figure 3. The results show that the viscosity of the formulations with both low (25 mM) and nominal (50 mM) lysine levels is acceptable at 150 mg/mL (<15 mPas) and that they are both below about 20 mPas at 165 mg/mL. Therefore the results that the nominal level of 50 mM lysine HCl is appropriately bracketed by the 25 mM lower level in terms of viscosity performance were confirmed. Based on the impact of pH on the viscosity and possible Dorman-type effects during TFF the antibody formulation pH was selected to be 5.9.

Assessment of Functionality of the Antibody Formulation in Prefilled Syringes

[0156] Table 7 gives the results of the functionality assessment of a formulation having MEDI-546 at 150 mg/mL in 25 mM histidine/histidine HCl, 50 mM lysine HCl, 130 mM trehalose dehydrate, 0.05% polysorbate 80 and a pH of 5.9.

[0157] The forces required to inject MEDI-546 Drug Product were within acceptable ranges. Table 8 gives the results of the user injection assessment (laboratory analysts). The users reported that 'Injection Force was easy'; 'arthritis gloves made handling of the APFS difficult'; 'SSI device worked.' Some variability in injection speed was noted based on users being new to using injection trainer and gloves (note: force to inject was not considered an issue). These results indicate that this formulation is suitable for use in a PFS with a 27 Ga STW needle. These studies did not identify any issues with the viscosity of the antibody formulations.

Table 7 - Functionality Assessment of Antibody Formulation

Replicate	Break Loose Force (N)	Average Glide Force (N)	Max. Glide Force (N)	Abs. Max. Force (N)
1	8.0	7.3	8.0	8.5
2	7.0	8.7	9.1	9.1
3	7.8	7.7	8.2	8.5

Table 8 - User Injection Time Assessment

Analyst	Injection time (s) - using arthritis gloves and APFS (flange, SSI device)	Injection time (s) - bare PFS
1	9.8	15.6
2	17	10.4
3	8.1	5.9

Example 7**Lysine and Protein Effects on Viscosity**

[0158] The viscosity of solutions of lysine at different concentrations were measured as a function of protein (anifrolumab) concentration, as shown in Table 9 and Figure 4.

Table 9

0 mM lysine	
	cone (mg/mL)
	130.5
	145.8
	180.5
5 mM lysine	
	cone (mg/mL)
	133.5
	147.8
	174.4
12.5 mM lysine	
	cone (mg/mL)
	134.6
	147.5
	176.4
25 mM lysine	

cone (mg/mL)	viscosity (cP)
135.9	6.1
148.2	8.6
181.6	17.5
50 mM lysine	
cone (mg/mL)	viscosity (cP)
135.8	5.6
148.3	6.8
178.8	11.1

[0159] The viscosity of solutions of antibody (anifrolumab) at different concentrations were measured as a function of protein concentration, as shown Table 10 and Figure 5.

Table 10

Target 135 mg/mL	
mM lysine	viscosity (cP)
0	8.3
5	8.6
12.5	6.6
25	6.1
50	5.6

Target 150 mg/mL	
mM lysine	viscosity (cP)
0	12.3
5	13.5
12.5	11.9
25	8.6
50	6.8

Target 180 mg/mL	
mM lysine	viscosity (cP)
0	38.4
5	33.0
12.5	20.1
25	17.5
50	11.1

[0160] The examples shown above illustrate various aspects of the invention and practice of the methods of the invention.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US7662381B \[0031\]](#)
- [US20110059078A \[0031\] \[0031\]](#)
- [US4816567A \[0040\]](#)

Non-patent literature cited in the description

- **JUO, PEI-SHOW**Concise Dictionary of Biomedicine and Molecular BiologyCRC Press20020000 [\[0022\]](#)
- The Dictionary of Cell and Molecular BiologyAcademic Press19990000 [\[0022\]](#)
- Oxford Dictionary Of Biochemistry And Molecular BiologyOxford University Press20000000 [\[0022\]](#)
- **WEERD et al.**J. Bioi. Chern., 2007, vol. 282, 20053-20057 [\[0028\]](#)
- **OGANESYAN et al.**Acta Crystallographica, 2008, vol. D 64, 700-704 [\[0031\]](#)
- **WARD et al.**Nature, 1989, vol. 341, 544-546 [\[0034\]](#)
- **BIRD et al.**Science, vol. 242, 423-426 [\[0034\]](#)
- **HUSTON et al.**Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 5879-5883 [\[0034\]](#)
- **MORRISON et al.**Proc. Natl. Acad. Sci. USA, 1984, vol. 81, 6851-6855 [\[0040\]](#)
- **HARLOW et al.**Antibodies: A Laboratory ManualCold Spring Harbor Laboratory Press19880000 [\[0041\]](#)
- **KABAT et al.**Sequences of Proteins of Immunological InterestU.S. Dept. of Health and Human Services19830000 [\[0042\]](#)
- **CHOTHIALESKJ.** Mol. Bioi., 1987, vol. 196, 901-917 [\[0042\]](#)
- **KABAT et al.**Sequence of Proteins of Immunological InterestU.S. Dept. of Health and Human Services19830000 [\[0043\]](#)
- **GENNARO, ALFONSO R.**Remington's Pharmaceutical SciencesMack Publishing Company19900000 [\[0061\]](#)
- WHO Requirements for the use of animal cells as in vitro substrates for the production of biologicals: Requirements for Biological Substances No. 5019980000 [\[0101\]](#)
- Pharmaceutical Dosage Form: Parenteral Medicationsvol. 1, [\[0106\]](#)

Patentkrav

1. Antistofformulering, som omfatter:

a. 100 til 200 mg/ml anifrolumab;

5 b. 0,02 % til 0,08 % af et overfladeaktivt stof, hvor det overfladeaktive stof er valgt fra gruppen, der består af polysorbat 20, polysorbat 80 og poloxamer;

c. 100 mM til 150 mM af et uladet hjælpestof, hvor det uladede hjælpestof er valgt fra gruppen, der består af fructose,

10 glucose, mannose, sorbose, xylose, lactose, maltose, saccharose, dextran, pullulan, dextrin, cyclodextriner, opløselig stivelse, trehalose, sorbitol, erythritol, isomalt, lactitol, maltitol, xylitol, glycerol, lactitol, hydroxyethylstivelse og vandopløselige glucaner;

15 d. en formuleringsbuffer; og

e. 25 til 130 mM af et lysinsalt:

hvor formuleringen er ved et pH på 5,5 til 6,5,

og hvor formuleringen har en viskositet lig med eller mindre end 20 mPas ved 25 °C.

20

2. Antistofformulering ifølge krav 1, hvor formuleringen er ved et pH på 5,9.

25 3. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor lysinsaltet er valgt fra gruppen, der består af lysinacetat, lysinmonochlorid, lysindichlorid, lysin-L-aspartat, lysin-L-glutamat og lysin-HCl.

30 4. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor det overfladeaktive stof er polysorbat 80.

35 5. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor formuleringen omfatter 0,05 % af et overfladeaktivt stof.

6. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor formuleringsbufferen er en acetatbuffer, TRIS-buffer, HEPES-buffer, hydrochloridbuffer, argininbuffer,

glycinbuffer, citratbuffer, histidinbuffer eller TES-buffer.

7. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor formuleringsbufferen er en histidinbuffer,
5 eventuelt hvor histidinbufferen omfatter histidinhydrochlorid.

8. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor formuleringsbufferen er histidin/histidinhydrochlorid.

10

9. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor formuleringsbufferen omfatter 10 mM til 40 mM histidin/histidinhydrochlorid, eventuelt hvor bufferen omfatter 25 mM histidin/histidinhydrochlorid.

15

10. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor det uladede hjælpestof er valgt fra gruppen, der består af glucose, saccharose, trehalose og glycerol, eventuelt hvor det uladede hjælpestof er trehalose,
20 eventuelt hvor det uladede hjælpestof er trehalosedihydrat.

11. Antistofformulering ifølge et hvilket som helst af ovennævnte krav, hvor formuleringen er egnet til intravenøs, subkutan eller intramuskulær administration.

25

12. Fyldt sprøjte, som indeholder antistofformuleringen ifølge et hvilket som helst af kravene 1 til 11.

DRAWINGS

FIG. 1




FIG. 2




FIG. 3
MEDI-546 Viscosity Screening




FIG. 4

Viscosity vs Protein Conc





FIG. 5

Viscosity vs Lysine Conc



SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

