



(86) Date de dépôt PCT/PCT Filing Date: 2012/02/10
(87) Date publication PCT/PCT Publication Date: 2012/08/16
(85) Entrée phase nationale/National Entry: 2013/08/08
(86) N° demande PCT/PCT Application No.: CN 2012/071017
(87) N° publication PCT/PCT Publication No.: 2012/106995
(30) Priorité/Priority: 2011/02/11 (EP11154188.4)


(51) Cl.Int./Int.Cl. *C07D 231/56* (2006.01),
A61K 31/41 (2006.01), *A61K 31/416* (2006.01),
A61K 31/422 (2006.01), *A61K 31/437* (2006.01),
A61P 29/00 (2006.01), *A61P 37/00* (2006.01),
A61P 43/00 (2006.01), *C07D 403/10* (2006.01),
C07D 403/12 (2006.01), *C07D 471/04* (2006.01)

(71) Demandeurs/Applicants:
MERCK SHARP & DOHME CORP., US;
MERCK SHARP & DOHME B.V., NL

(72) Inventeurs/Inventors:
KARSTENS, WILLEM FREDERIK JOHAN, NL;
STELT, MARIO VAN DER, NL;
CALS, JOS, NL; ...

(54) Titre : INHIBITEURS DE RORGAMMAT

(54) Title: RORGAMMAT INHIBITORS



(I)

(57) Abrégé/Abstract:

The present invention relates to compounds according to Formula (I) or a pharmaceutically acceptable salt or solvate thereof. Such compounds can be used in the treatment of RORgammaT-mediated diseases or condition.



(72) **Inventeurs(suite)/Inventors(continued)**: AZEVEDO, RITA CORTE REAL GONCALVES, NL; BARR, KENNETH JAY, US; ZHANG, HONGJUN, US; BERESIS, RICHARD THOMAS, CN; ZHANG, DONGSHAN, CN; DUAN, XIAOBANG, CN

(74) **Agent**: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
16 August 2012 (16.08.2012)



(10) International Publication Number
WO 2012/106995 A1

(51) International Patent Classification:

C07D 231/56 (2006.01) *A61K 31/422* (2006.01)
C07D 403/10 (2006.01) *A61K 31/437* (2006.01)
C07D 403/12 (2006.01) *A61P 43/00* (2006.01)
C07D 471/04 (2006.01) *A61P 37/00* (2006.01)
A61K 31/416 (2006.01) *A61P 29/00* (2006.01)
A61K 31/41 (2006.01)

(21) International Application Number:

PCT/CN2012/071017

(22) International Filing Date:

10 February 2012 (10.02.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11154188.4 11 February 2011 (11.02.2011) EP

(71) Applicants (for all designated States except US): **MERCK SHARP & DOHME CORP.** [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **N.V. ORGANOON** [NL/NL]; Kloosterstraat 6 NL-5349 AB Oss (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KARSTENS, Willem Frederik Johan** [NL/NL]; Dotterbloem 28, 5351 MB, Berghem (NL). **STELT, Mario van der** [NL/NL]; Sterrenbos 41, 3511 ET, Utrecht (NL). **CALS, Jos** [NL/NL]; Irenelaan 30, 5342 EL, Oss (NL). **AZEVEDO, Rita Corte Real Goncalves** [PT/NL]; Oberon 14, 5221 LV 's-Hertogenbosch (NL). **BARR, Kenneth Jay** [US/US]; 261 Commonwealth Avenue, No. 7, Boston, MA 02116 (US). **ZHANG, Hongjun** [CN/US]; 8 Town House Drive, Newton, MA 02460 (US). **BERESIS, Richard Thomas** [US/CN]; Apartment 2504, Building 1, 2504 Shimmen #2 Road, Shanghai 200040 (CN). **ZHANG, Dong-**

shan [CN/CN]; Shanghai ChemPartner Co., Ltd., No.10 Building, 965 Halei Road, Zhangjiang Hi-tech Park, Pudong New Area, Shanghai 201203 (CN). **DUAN, Xiaobang** [CN/CN]; Shanghai ChemPartner Co., Ltd., No.10 Building, 965 Halei Road, Zhangjiang Hi-tech Park, Pudong New Area, Shanghai 201203 (CN).

(74) Agent: **LIU, SHEN & ASSOCIATES**; A0601, Huibin Building, No.8 Beichen Dong Street, Chaoyang District, Beijing 100101 (CN).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: RORGAMMAT INHIBITORS



(I)

(57) Abstract: The present invention relates to compounds according to Formula (I) or a pharmaceutically acceptable salt or solvate thereof. Such compounds can be used in the treatment of RORgammaT-mediated diseases or condition.

RORgammaT INHIBITORS

BACKGROUND OF THE INVENTION

5 Upon activation by antigen-presenting cells naïve T helper cells undergo clonal expansion and will ultimately differentiate in cytokine secreting effector T cells, such as Th1 and Th2 subtypes. A third and distinct effector subset has been identified, which plays a key role in providing immunity to bacteria and fungi at mucosal surfaces (Kastelein et al., *Annu. Rev. Immunol.* 25: 221-242, 2007). This effector T helper cell subset can be distinguished
10 based on its ability to produce large quantities of IL-17/F, IL-21 and IL-22, and is named Th17 (Miossec et al., *New Eng. J. Med.* 2361: 888-898, 2009).

 Different T helper subsets are characterized by the expression of lineage specific master transcription factors. Th1 and Th2 effector cells express Tbet and GATA3, respectively. A Thymocyte/T cell specific variant of Retinoic Acid Receptor-related Orphan Receptor (ROR),
15 RORgammaT, is highly expressed in Th17 cells (He et al., *Immunity* 9: 797-806, 1998). RORgammaT belongs to the nuclear hormone receptor superfamily (Hirose et al., *Biochem. Biophys. Res. Comm.* 205: 1976-1983, 1994). RORgammaT is a truncated form of RORgamma, lacking the first N-terminal 21 amino acids and is, in contrast to RORgamma which is expressed in multiple tissues (heart, brain, kidney, lung, liver and muscle),
20 exclusively expressed in cells of the lymphoid lineage and embryonic lymphoid tissue inducers (Sun et al., *Science* 288: 2369-2372, 2000; Eberl et al., *Nat Immunol.* 5: 64-73, 2004).

 Studies using heterozygous knock-in mice replacing the RORgammaT open reading frame with GFP, revealed a constitutive expression of GFP in approximately 10% of the
25 CD4⁺ T cells in the small intestinal lamina propria (LP), co-expressing the Th17 cytokines IL-17/F and IL-22 (Ivanov et al., *Cell* 126: 1121-1133, 2006). In mice deficient for RORgammaT, the number of Th17 cells was markedly decreased in the LP and in vitro stimulation of CD4⁺ T cells, under Th17 polarizing conditions resulted in a drastic decrease of IL-17 expression. These results were further substantiated via forced expression of
30 RORgammaT in naïve CD4⁺ T cells, which resulted in an induction of IL-17/F and IL-22 (Ivanov et al., *Cell* 126: 1121-1133, 2006). Taken together demonstrating the importance of RORgammaT in differentiation and stabilization of the Th17 lineage. In addition, a ROR

family member, RORalpha has been demonstrated to be involved in Th17 differentiation and stabilization (Yang et al., *Immunity* 28: 29-39, 2008).

Recently, RORgammaT was shown to play a crucial role in non-Th17 lymphoid cells. In these studies, RORgammaT was critically important in innate lymphoid cells expressing
5 Thy1, SCA-1 and IL-23R proteins. Genetic disruption of RORgamma in a mouse colitis model dependent on these innate lymphoid cells, prevented colitis development (Buonocore et al., *Nature* 464: 1371-1375, 2010). In addition, RORgammaT was shown to play a crucial role in other non-Th17 cells, such as mast cells (Hueber et al., *J. Immunol.* 184: 3336-3340, 2010). Finally, RORgammaT expression and secretion of Th17-type of cytokines was
10 reported for Lymphoid Tissue Inducer cells, NK T-cells, NK cells (Eberl et al., *Nat. Immunol.* 5: 64-73, 2004) and gamma-delta T-cells (Sutton et al., *Nat. Immunol.* 31: 331-341, 2009; Louten et al., *J. Allergy Clin. Immunol.* 123: 1004-1011, 2009), suggesting an important function for RORgammaT in these subtypes of cells.

Based on the role of IL-17 producing cells (either Th17 or non-Th17 cells)
15 RORgammaT has been identified as a key mediator in the pathogenesis of several diseases (Louten et al., *J. Allergy Clin. Immunol.* 123: 1004-1011, 2009; Annunziato et al., *Nat. Rev. Rheumatol.* 5: 325-331, 2009). This was confirmed using several disease models representative of autoimmune diseases. Genetic ablation of the RORgamma gene in mice prevented the development of experimental autoimmune diseases, such as experimental
20 autoimmune encephalomyelitis (EAE) and colitis (Ivanov et al., *Cell* 126:1121-33, 2006; Buonocore et al., *Nature* 464: 1371-1375, 2010).

Being a critical mediator in Th17-cells and other non-Th17 cells, inhibition of RORgammaT is expected to have a beneficial effect on autoimmune diseases, such as, but not limited to rheumatoid arthritis, psoriasis, multiple sclerosis, inflammatory bowel disease,
25 Crohn's disease, and asthma (Annunziato et al., *Nat. Rev. Immunol.* 5: 325-331, 2009; Louten et al., *J. Allergy Clin. Immunol.* 123: 1004-1011, 2009). Inhibition of RORgammaT may also be beneficial in other diseases, which are characterized by increased levels of Th17 cells and/or elevated levels of Th17 hallmark cytokines such as IL-17, IL-22 and IL-23. Examples of such diseases are Kawasaki Disease (Jia et al., *Clin. Exp. Immunol.* 162: 131-137, 2010)
30 and Hashimoto's thyroiditis (Figueroa-Vega et al., *J. Clin. Endocrinol. Metab.* 95: 953-62, 2010). Another example includes infectious diseases, such as, but not limited to, mucosal

leishmaniasis (Boaventura et al., *Eur. J. Immunol.* 40: 2830-2836, 2010). In each of the above examples the inhibition may be enhanced by simultaneous inhibition of RORalpha.

Compounds modulating RORgammaT have been reported. Examples of agonists include T0901317 and SR1078 (Wang et al., *ACS Chem. Biol.* 5:1029-1034, 2010). In addition, antagonist have been reported such as 7-oxygenated sterols (Wang et al., *J. Biol. Chem.* 285: 5013-5025, 2009) and compounds described in EP2181710 A1.

Numerous immune and inflammatory disorders continue to afflict millions of patients worldwide. Although significant advances have been made in treating these disorders, current therapies do not provide satisfactory results for all patients due to, for example, detrimental side effects or insufficient efficacy. One exemplary immune disorder in need of better therapy is psoriasis. Various therapeutics have been developed in an attempt to treat psoriasis. However, the traditional therapies for psoriasis often have toxic adverse effects. An exemplary inflammatory disorder in need of better treatment is rheumatoid arthritis. Numerous therapeutics have been developed in an attempt to treat this disorder. However, some patients develop resistance to current therapies.


Accordingly, a need exists for improved treatments for immune disorders and inflammatory disorders. The present invention addresses this need and provides other related advantages.

SUMMARY OF THE INVENTION

The present invention provides compounds which inhibit RORgammaT, their use for the treatment of RORgammaT-mediated diseases or conditions, in particular autoimmune diseases and inflammatory diseases, as well as pharmaceutical compositions comprising such compounds and pharmaceutical carriers.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a compound according to Formula I



I

5 or a pharmaceutically acceptable salt or solvate thereof wherein,

X is C(O), SO or SO₂,

Y is CH or N;

n = 0, 1, 2, 3 or 4;

A⁴-A⁷ are independently N or CR⁴, CR⁵, CR⁶ or CR⁷, respectively, with the proviso that no
10 more than two of the four positions A⁴-A⁷ can be simultaneously N;

R¹ is (i) (C₃₋₇)cycloalkyl or (C₃₋₅)heterocycloalkyl, both optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally
15 substituted with one or more halogens; (ii) (C₂₋₉)heteroaryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens; or (iii) (C₆₋₁₄)aryl, optionally substituted with one or
20 more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy are optionally substituted with one or more halogens;


R² is C(O)OH, 5-tetrazoyl, HOC(CF₃)₂, C(O)OH(C₁₋₁₀)alkyl, (C₁₋₁₀)alkylsulfoxyaminocarbonyl; or carbamoyl;

R³ is hydrogen, halogen, cyano, nitro, hydroxy, (C₁₋₃)alkylC(O)O-, (C₁₋₄)alkyl, or (C₁₋

₄)alkoxy, wherein (C₁₋₄)alkyl and (C₁₋₄)alkoxy are optionally substituted with one or more halogens;


R⁴-R⁷ independently are H, halogen, amino, cyano, hydroxy, (C₁₋₃)alkoxy, (C₁₋₄)alkyl, (C₀₋₁₀)alkyl)aminocarbonyl, (di)(C₁₋₆)alkylaminocarbonyl or amino(C₁₋₄)alkyl, wherein (C₁₋

5 ₃)alkoxy, (C₁₋₄)alkyl, (C₀₋₁₀)alkyl)aminocarbonyl, (di)(C₁₋₆)alkylaminocarbonyl and amino(C₁₋₄)alkyl are optionally substituted with one or more halogen, hydroxyl or (C₁₋₃)alkoxy; or a

group having the formula , optionally substituted with one or more of the following: (C₁₋₁₀)alkyl, halogen, amino, cyano, hydroxy, (C₁₋₃)alkoxy, and wherein m is 1, 2, 3, or 4.

10

This invention also provides a compound according to Formula Ia



15

Ia

or a pharmaceutically acceptable salt or solvate thereof

wherein X represents C(O), SO or SO₂,

Y is CH or N;

20 A⁴-A⁷ are independently N or CR⁴, CR⁵, CR⁶ or CR⁷, respectively, with the proviso that no more than two of the four positions A⁴-A⁷ can be simultaneously N;

R¹ is (C₃₋₇)cycloalkyl or (C₃₋₅)heterocycloalkyl, both optionally substituted with one or more groups selected from halogen, amino, cyano, hydroxy, H₂NC(O)-, or (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all optionally substituted with one

25 or more halogens or;

R¹ is (C₂₋₉)heteroaryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, or (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all optionally substituted with one or more halogens or;

5 R¹ is (C₆₋₁₄)aryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, or (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all optionally substituted with one or more halogens;

R² is C(O)OH, 5-tetrazoyl, or HOC(CF₃)₂;

10 R³ is independently selected from hydrogen, halogen, cyano, nitro or (C₁₋₄)alkyl, optionally substituted with one or more halogens; and

R⁴-R⁷ independently are H, halogen, amino, cyano, hydroxy, (C₁₋₃)alkoxy, (C₁₋₄)alkyl, optionally substituted with one or more halogens.

In the first embodiment of the compounds of Formula I and Ia are compounds wherein
15 A⁴, A⁵, A⁶, A⁷ are selected from the group consisting of: (i) CR⁴, CR⁵, CR⁶, CR⁷; (ii) N, CR⁵, CR⁶, CR⁷; (iii) CR⁴, N, CR⁶, CR⁷; (iv) CR⁴, CR⁵, N, CR⁷; (v) CR⁴, CR⁵, CR⁶, N; (vi) N, N, CR⁶, CR⁷; (vii) CR⁴, N, N, CR⁷; (viii) CR⁴, CR⁵, N, N; (ix) N, CR⁵, N, CR⁷; (x) CR⁴, N, CR⁶, N; and (xi) N, CR⁵, CR⁶, N.

In a subset of the first embodiment are compounds wherein A⁴, A⁵, A⁶, A⁷ are selected
20 from the group consisting of: (i) CR⁴, CR⁵, CR⁶, CR⁷; (ii) N, CR⁵, CR⁶, CR⁷; and (iii) CR⁴, N, CR⁶, CR⁷.

In a second subset of the first embodiment are compounds wherein A⁴, A⁵, A⁶, A⁷ is (i) CR⁴, CR⁵, CR⁶, CR⁷, or (ii) N, CR⁵, CR⁶, CR⁷; and Y is N.

In a third subset of the first embodiment are compounds wherein
25 X is C(O) or SO₂;

R¹ is (i) (C₃₋₇)cycloalkyl or (C₃₋₅)heterocycloalkyl, both optionally substituted with one or more groups selected from (C₁₋₄)alkyl or halogen; (ii) (C₂₋₉)heteroaryl, optionally substituted with one or more groups selected from halogen, amino or (C₁₋₄)alkyl; or (iii) (C₆₋₁₄)aryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro,
30 hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl, (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens.

In a fourth subset of the first embodiment are compounds wherein
R¹ is (C₂₋₉)heteroaryl or (C₆₋₁₄)aryl, both optionally substituted with one or more groups
selected from halogen, amino, cyano, nitro, hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally
5 substituted with one or more halogen.

In a fifth subset of the first embodiment are compounds according to Formula I wherein
R¹ is (C₆₋₁₄)aryl, optionally substituted with one or more groups selected from halogen,
amino, cyano, nitro, hydroxy, or (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all
optionally substituted with one or more halogen.

10 In a sixth subset of the first embodiment are compounds wherein R¹ is phenyl, naphthyl,
pyridinyl, quinolinyl, benzooxadiazolyl, thiophenyl, or isoxazolyl, each optionally substituted
with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and
(C₁₋₃)alkoxy are optionally substituted with one or more halogens.


15 In a seventh subset of the first embodiment are compounds according to Formula I and
Ia wherein (C₆₋₁₄)aryl in R¹ is phenyl.

In an eighth subset of the first embodiment are compounds wherein R¹ is phenyl,
optionally substituted with one or more groups selected from halogen, amino, cyano, nitro,
hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl,
20 (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens.

In a ninth subset of the first embodiment, are compounds wherein R² is C(O)OH.

A tenth subset of the invention relates to compounds according to Formula I and Ia
wherein Y is N.


25 In a second embodiment of the compound of Formula I is a compound having Formula
Ib



Ib

- 5 and a pharmaceutically acceptable salt or solvate thereof, wherein A^4 - A^7 , R^1 - R^3 , X, Y and n are defined as set forth hereinabove for Formula I.

In a third embodiment of the compound of Formula I is a compound having Formula Ic



Ic

or a pharmaceutically acceptable salt or solvate thereof wherein,

X is C(O) or SO₂

- 15 R^8 is selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy are optionally substituted with one or more halogens;


x is 0, 1, 2, 3, 4 or 5; and

A^4 - A^7 , R^1 - R^3 , Y and n are defined as set forth hereinabove for Formula I.

.

In a fourth embodiment of the compound of Formula I is a compound having Formula Id


5



Id

10 and a pharmaceutically acceptable salt or solvate thereof, wherein A^4 - A^7 , R^2 , R^3 , X, Y and n are defined as set forth hereinabove for Formula I.

In a fifth embodiment of the compound of Formula I is a compound having Formula Ie



15

Ie

and a pharmaceutically acceptable salt or solvate thereof, wherein A^4 - A^7 , R^3 , X and n are defined as set forth hereinabove for Formula I.


5

In a further embodiment of the compounds of Formula I, Ia, Ib, Ic, Id, and Ie, are compounds wherein

R^4 is selected from H, halogen, amino, and (C_{1-4}) alkyl optionally substituted with one or more halogen;

10 R^5 is H;

R^6 is selected from H, dimethylaminocarbonyl, $-CH_2NH_2$, cyano,

, and each optionally substituted with one or more halogen, hydroxyl, $-CH_2OH$, $-CH_2OCH_3$ or methoxy; and

R^7 is selected from H and halogen.

15

A still further embodiment of the compounds of Formula I, Ia, Ib, Ic, Id, and Ie, are compounds wherein one of R^4 , R^5 and R^7 is other than hydrogen.

The invention also relates to those compounds wherein all specific definitions for A^1 through A^4 , R^1 through R^8 , X, Y, n and x and all substituent groups in the various aspects of the inventions defined here above occur in any combination within the definition of the compound of Formula I.

Non-limiting examples of the compounds of the present invention include:

25 4-(1-(2,6-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2,6-dichlorobenzoyl)-5-fluoro-1H-indazol-3-yl)benzoic acid;

4-(1-(2,6-dichlorobenzoyl)-7-fluoro-1H-indazol-3-yl)benzoic acid;

4-(1-(2,6-dichlorobenzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;

4-(1-(2,6-dichlorobenzoyl)-6-fluoro-1H-indazol-3-yl)benzoic acid;

30 4-(1-(2,6-dichlorobenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;

4-(1-(2-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;

- 4-(1-(2,6-dichlorobenzoyl)-1H-pyrrolo[3,2-b]pyridin-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acid;
1,1,1,3,3,3-hexafluoro-2-(4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)phenyl)propan-2-ol;
- 5 4-(1-(2-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(3,5-dichloroisonicotinoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-bromobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-(methoxycarbonyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(cyclohexanecarbonyl)-1H-indazol-3-yl)benzoic acid;
- 10 4-(1-(2-bromo-6-chlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-fluoro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,4,6-trichlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,6-dimethylbenzoyl)-1H-indazol-3-yl)benzoic acid;
- 15 4-(1-(2-ethoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,3-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-chlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-methyl-6-nitrobenzoyl)-1H-indazol-3-yl)benzoic acid;
- 20 4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(quinolin-8-ylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-bromophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,5-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(3-chloro-2-methylphenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
- 25 4-(1-(2,3-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,3,4-trichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-cyanophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-(trifluoromethoxy)phenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(naphthalen-1-ylsulfonyl)-1H-indazol-3-yl)benzoic acid;
- 30 4-(1-(2,6-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(benzo[c][1,2,5]oxadiazol-4-ylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-cyano-5-methylphenylsulfonyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;
 4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-2-(trifluoromethyl)benzoic acid;
 4-(1-(2-chlorophenylsulfonyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acid;
 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(3-chlorophenylsulfonyl)-1H-indazole;
 5 methyl 3-(3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-ylsulfonyl)thiophene-2-carboxylate;
 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(2-bromo-4-fluorophenylsulfonyl)-1H-indazole;
 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(3-bromo-5-chlorothiophen-2-ylsulfonyl)-1H-indazole;
 4-(3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-ylsulfonyl)-3,5-dimethylisoxazole;
 (3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-yl)(2,6-dichlorophenyl)methanone;
 10 (3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-yl)(2,3-dichlorophenyl)methanone;
 methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoate;
 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoic acid;
 Sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoate;
 15 methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoate;
 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
 sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoate;
 methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-fluorobenzoate;
 20 sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-fluorobenzoate;
 (2-chloro-6-(trifluoromethyl)phenyl)(4-fluoro-3-(4-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)-1H-indazol-1-yl)methanone;
 25 3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;
 2-(3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)phenyl)acetic acid;
 2-(4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)phenyl)acetic acid;
 2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;
 Sodium 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoate;
 30 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzamide;

- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluoro-5-methylbenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-methylbenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-methoxybenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluorobenzoic acid;
- sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluorobenzoate;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-methylbenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-methoxybenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,6-difluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-isopropylbenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3,5-dimethoxybenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-cyano benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,3-difluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,5-difluorobenzoic acid;
- 2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluoro-2-methoxybenzoic acid;
- 5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoic acid;
- 3-fluoro-4-(4-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
- 4-(4-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
- 3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl) benzoic acid;
- 2-(4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl) phenyl)acetic acid;

- (2-chloro-6-(trifluoromethyl)phenyl)(3-(4-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)-1H-pyrazolo[4,3-b]pyridin-1-yl)methanone;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-methylbenzoic acid;
- 5 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-fluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-methoxybenzoic acid;
- 10 2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-5-fluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2,5-difluorobenzoic acid;
- 15 5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-hydroxybenzoic acid;
- 4-(1-(2-fluoro-6-methoxybenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;
- 3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-(trifluoromethyl)-1H-indazol-3-yl)benzoic acid;
- 20 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-isopropylbenzoic acid;
- sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-fluorobenzoate;
- sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-3-fluorobenzoate;
- 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
- 3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
- 25 sodium 3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoate;
- 4-(7-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
- 3-fluoro-4-(7-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
- 4-(6-bromo-1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;
- 4-(6-bromo-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
- 30 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-methoxy-1H-indazol-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl)benzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl) benzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl)-3-fluorobenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-hydroxy-1H-indazol-3-yl)benzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-hydroxy-1H-indazol-3-yl)-3-
5 fluorobenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(hydroxymethyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(hydroxymethyl)-1H-indazol-3-yl)-3-
fluorobenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(methoxymethyl)-1H-indazol-3-yl)benzoic acid;
10 1-(2-chloro-6-(trifluoromethyl)benzoyl)-3-(4-(methoxy carbonyl)phenyl)-1H-indazole-6-
carboxylic acid;
methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-
indazol-3-yl)benzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxy ethylcarbamoyl)-1H-indazol-3-
15 yl)benzoic acid;
sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-
indazol-3-yl)benzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)benzoic
acid;
20 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-
yl)benzoic acid;
sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-
indazol-3-yl)benzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-3-
25 fluorobenzoic acid;
sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-
3-fluorobenzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)-2-
fluorobenzoic acid;
30 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)-3-
fluorobenzoic acid;

- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-2-fluorobenzoic acid;
- 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;
- 5 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-hydroxypropylcarbamoyl)-1H-indazol-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-hydroxypropylcarbamoyl)-1H-indazol-3-yl)-10 3-fluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-methoxypropylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxypropylcarbamoyl)-1H-indazol-3-yl)benzoic acid;
- 15 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(4-hydroxybutylcarbamoyl)-1H-indazol-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-hydroxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-hydroxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;
- 20 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-methoxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-fluoroazetidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid;
- 25 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3,3-difluoroazetidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid;
- 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(pyrrolidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid;
- 4-(6-(azetidine-1-carbonyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
- 30 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoic acid;

2-acetoxy-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoate;

5 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluoro-6-hydroxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-hydroxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluoro-2-hydroxybenzoic acid;

10 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3,6-difluoro-2-hydroxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-hydroxybenzoic acid;

2-acetoxy-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-

15 yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-2-hydroxybenzoic acid;

4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;

4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-3-

20 fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-(difluoromethyl)benzoic acid;
and

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-cyano-1H-indazol-3-yl)benzoic acid.

25 The terms used herein have their ordinary meaning and the meaning of such terms is independent at each occurrence thereof. That notwithstanding, and except where stated otherwise, the following definitions apply throughout the specification and claims. Chemical names, common names, and chemical structures may be used interchangeably to describe the same structure. If a chemical compound is referred to using both a chemical structure and a
30 chemical name, and an ambiguity exists between the structure and the name, the structure predominates. These definitions apply regardless of whether a term is used by itself or in combination with other terms, unless otherwise indicated. Hence, the definition of "alkyl"

applies to "alkyl" as well as the "alkyl" portions of "hydroxyalkyl," "fluoroalkyl," "alkoxy", etc.

As used herein, and throughout this disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

5 The term "alkyl," as used herein, refers to an aliphatic hydrocarbon group having one of its hydrogen atoms replaced with a bond having the specified number of carbon atoms. In different embodiments, an alkyl group contains, for example, from 1 to 6 carbon atoms (C₁-C₆ alkyl) or from 1 to 3 carbon atoms (C₁-C₃ alkyl). Non-limiting examples of alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl,
10 neopentyl, isopentyl, n-hexyl, isohexyl and neohexyl. In one embodiment, an alkyl group is linear. In another embodiment, an alkyl group is branched.

Unless specified otherwise, "alkyl" includes both branched- and straight-chain saturated aliphatic hydrocarbon groups, including all isomers, having the specified number of carbon atoms; for example, "C₁₋₆ alkyl" (or "C₁-C₆ alkyl") includes all of the hexyl alkyl and pentyl
15 alkyl isomers as well as n-, iso-, sec- and t-butyl, n- and isopropyl, ethyl and methyl. "Alkylene" refers to both branched- and straight-chain saturated aliphatic hydrocarbon groups, including all isomers, having the specified number of carbons, and having two terminal end chain attachments; for example, the term "A-C₄alkylene-B" represents, for example, A-CH₂-CH₂-CH₂-B, A-CH₂-CH₂-CH(CH₃)-CH₂-B, A-CH₂-CH(CH₂CH₃)-B, A-CH₂-
20 C(CH₃)(CH₃)-B, and the like. "Alkoxy" represents a linear or branched alkyl group of indicated number of carbon atoms attached through an oxygen bridge; for example "C₁-C₆ alkoxy" includes -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -O(CH₂)₅CH₃, and the like.

Unless otherwise specifically noted as only "unsubstituted" or only "substituted", alkyl groups are unsubstituted or substituted with 1 to 3 substituents on each carbon atom, with
25 halo, C₁-C₂₀ alkyl, CF₃, NH₂, N(C₁-C₆ alkyl)₂, NO₂, oxo, CN, N₃, -OH, -O(C₁-C₆ alkyl), C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₀-C₆ alkyl) S(O)₀₋₂-, (C₀-C₆ alkyl)S(O)₀₋₂(C₀-C₆ alkyl)-, (C₀-C₆ alkyl)C(O)NH-, H₂N-C(NH)-, H₂N-C(O)(NH)-, -O(C₁-C₆ alkyl)CF₃, (C₀-C₆ alkyl)C(O)-, (C₀-C₆ alkyl)OC(O)-, (C₀-C₆ alkyl)O(C₁-C₆ alkyl)-, (C₀-C₆ alkyl)C(O)₁₋₂(C₀-C₆ alkyl)-, (C₀-C₆ alkyl)OC(O)NH-, -NH(C₁-C₆ alkyl)NHC(O)NH(C₁-C₆ alkyl),
30 NHC(O)OC₁-C₆ alkyl, -NH(C₁-C₆ alkyl)NHSO₂(C₁-C₆ alkyl), -(C₀-C₆ alkyl)NHSO₂(C₁-C₆ alkyl), aryl, aralkyl, heterocycle, heterocyclylalkyl, halo-aryl, halo-aralkyl, halo-heterocycle,

halo-heterocyclalkyl, cyano-aryl, cyano-alkyl, cyano-heterocycle and cyano-heterocyclalkyl.


The term "alkenyl" means a straight or branched carbon chain having the specified number of carbon atoms with at least one carbon-carbon double bond. Examples of alkenyl include, but are not limited to, vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, 2,4-hexadienyl, and the like.


The term "alkynyl" means a straight or branched carbon chain having the specified number of carbon atoms with at least one carbon-carbon triple bond. Examples of alkynyl include, but are not limited to ethynyl, propargyl, 1-propynyl, 2-butyne, and the like.

The term "carbocycle" (and variations thereof such as "carbocyclic" or "carbocycl") as used herein, unless otherwise indicated, refers to (i) a C₃ to C₈ monocyclic, saturated or unsaturated ring or (ii) a C₇ to C₁₂ bicyclic saturated or unsaturated ring system. Each ring in (ii) is either attached via a bond to, or fused (including spirofused) to, the other ring, and each ring is saturated or unsaturated. The carbocycle may be attached to the rest of the molecule at any carbon atom which results in a stable compound.

Saturated carbocyclics form a subset of carbocycles in which the entire ring system (mono- or polycyclic) is saturated. Saturated monocyclic carbocyclic rings are also referred to as cycloalkyl rings, e.g., cyclopropyl, cyclobutyl, etc. The fused bicyclic carbocycles are a further subset of the carbocycles in which a C₇ to C₁₀ bicyclic ring system in which each ring is saturated or unsaturated and two adjacent carbon atoms (or in the case of spirofused, one carbon atom) are shared by each of the rings in the ring system. A saturated bicyclic carbocycle is one in which both rings are saturated. An unsaturated bicyclic carbocycle is one in which one ring is unsaturated and the other is unsaturated or saturated. Unless otherwise noted, carbocycle is unsubstituted or substituted with C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, aryl, halogen, NH₂ or OH. A subset of the fused bicyclic unsaturated carbocycles are those bicyclic carbocycles in which one ring is a benzene ring and the other ring is saturated or unsaturated, with attachment via any carbon atom that results in a stable compound.

Representative examples of this subset include the following:






Aromatic carbocycles form another subset of the carbocycles. The term "aryl" refers to aromatic mono- and poly-carbocyclic ring systems in which the individual carbocyclic rings in the polyring systems are fused or attached to each other via a single bond. Suitable aryl groups include phenyl, naphthyl, and biphenyl.

The term "cycloalkyl" means a cyclic ring of an alkane having the specified total ring carbon atoms; for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl.

The term "heterocycle" (and variations thereof such as "heterocyclic" or "heterocyclyl") broadly refers to (i) a stable 4- to 8-membered, saturated or unsaturated monocyclic ring, or (ii) a stable 7- to 12-membered bicyclic ring system, wherein each ring in (ii) is either attached via a bond to, or fused (including spirofused) to, the other ring, and each ring is saturated or unsaturated, and the monocyclic ring or bicyclic ring system contains one or more heteroatoms (e.g., from 1 to 6 heteroatoms, or from 1 to 4 heteroatoms) selected from N, O and S and a balance of carbon atoms (the monocyclic ring typically contains at least one carbon atom and the ring systems typically contain at least two carbon atoms); and wherein any one or more of the nitrogen and sulfur heteroatoms is optionally oxidized, and any one or more of the nitrogen heteroatoms is optionally quaternized. Unless otherwise specified, the heterocyclic ring may be attached at any heteroatom or carbon atom, provided that attachment results in the creation of a stable structure. Unless otherwise specified, when the heterocyclic ring has substituents, it is understood that the substituents may be attached to any atom in the ring, whether a heteroatom or a carbon atom, provided that a stable chemical structure results.


Saturated heterocyclics form a subset of the heterocycles; i.e., the term "saturated heterocyclic" generally refers to a heterocycle as defined above in which the entire ring system (whether mono- or poly-cyclic) is saturated. The term "saturated heterocyclic ring" refers to a 4- to 8-membered saturated monocyclic ring or a stable 7- to 12-membered bicyclic ring system which consists of carbon atoms and one or more heteroatoms selected from N, O and S. Representative examples include piperidinyl, piperazinyl, azepanyl, pyrrolidinyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, isothiazolidinyl, 1,4-dioxanyl, 1,4-thioxanyl, tetrahydropyranyl, tetrahydrofuryl (or tetrahydrofuranyl), tetrahydrothienyl, and tetrahydrothiopyranyl.



Heteroaromatics form another subset of the heterocycles; i.e., the term "heteroaromatic" (alternatively "heteroaryl") generally refers to a heterocycle as defined above in which the entire ring system (whether mono- or poly-cyclic) is an aromatic ring system. The term "heteroaromatic ring" refers a 5- or 6-membered monocyclic aromatic ring or a 7- to 12-membered bicyclic aromatic ring, and which consists of carbon atoms and one or more heteroatoms selected from N, O and S. In the case of substituted heteroaryl rings containing at least one nitrogen atom (e.g., pyridine), such substitutions can be those resulting in N-oxide formation. Representative examples of monocyclic heteroaromatic rings include pyridyl, pyrrolyl, pyrazinyl, pyrimidinyl, pyridazinyl, thienyl (or thiophenyl), thiazolyl, furanyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, and thiadiazolyl. Examples of bicyclic heteroaromatic rings include benzotriazolyl, indolyl, benzoxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, isoindolyl, indazolyl, quinoxalyl, quinazolinyl, cinnolyl, quinolyl, isoquinolyl, naphthyridinyl, pyrazolo[3,4-b]pyridine, imidazo[2,1-b](1,3)thiazole, (i.e.,



), 6-(1-pyrrolyl)-3-pyridyl, 4-(1-pyrrolyl)phenyl, 4-(pyrid-3-yl)phenyl, and 4-(pyrid-4-yl)phenyl.

Another subset of heterocycles are unsaturated heterocycles in which one or both rings are unsaturated (provided the entire ring system is not aromatic). Representative examples of unsaturated heterocycles include dihydrofuranyl, dihydrothienyl, dihydropyranyl, dihydroimidazolyl, indolyl, isoindolyl, chromanyl, isochromanyl, tetrahydroquinolyl, tetrahydroisoquinolyl, tetrahydronaphthyridinyl, 2,3-dihydrobenzofuranyl, 1,4-

benzoxazinyl, 1,3-benzoxazolyl, 2,3-dihydrobenzo-1,4-dioxinyl (i.e., ) , and benzo-

1,3-dioxolyl (i.e., ). In certain contexts herein,  is alternatively referred to as phenyl having as a substituent methylenedioxy attached to two adjacent carbon atoms. Also included are groups such as chromone and coumarin.

Unless otherwise specifically noted as only unsubstituted or only substituted, cycloalkyl, cycloalkenyl, cycloalkyl, aryl (including phenyl) and heteroaryl groups are unsubstituted or substituted (also referred to as "optionally substituted"). Unless the substituents are specifically provided, substituents for substituted or optionally substituted cycloalkyl,


cycloalkenyl, aryl (including phenyl, and as an isolated substituent or as part of a substituent such as in aryloxy and aralkyl), heteroaryl (as an isolated substituent or as part of a substituent such as in heteroaryloxy and heteroaralkyl) are one to three groups independently selected from halogen (or halo), C₁-C₆ alkyl optionally substituted with one to five fluorine, NH₂, N(C₁-C₆ alkyl)₂, NO₂, oxo, CN, N₃, -OH, -O(C₁-C₆ alkyl) optionally substituted with one to five fluorine, C₃-C₁₀ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₀-C₆ alkyl)S(O)₀₋₂-, aryl-S(O)₀₋₂-, (C₀-C₆ alkyl)S(O)₀₋₂(C₀-C₆ alkylene)-, (C₀-C₆ alkyl)C(O)NH-, H₂N-C(NH)-, (C₀-C₆ alkyl)C(O)-, (C₀-C₆ alkyl)OC(O)-, (C₀-C₆alkyl)O(C₁-C₆ alkylene)-, (C₀-C₆ alkyl)-C(O)₁₋₂(C₀-C₆ alkylene)-, (C₀-C₆ alkyl)₂NC(O)-, (C₀-C₆ alkyl)OC(O)NH-, aryl, aralkyl, heteroaryl, heteroaralkyl, halo-aryl, halo-aralkyl, halo-heteroaryl, halo-heteroaralkyl, cyano-aryl, cyano-aralkyl, cyano-heteroaryl and cyano-heteroaralkyl.


The term "halogen" (or "halo") refers to fluorine, chlorine, bromine and iodine (alternatively referred to as fluoro (F), chloro (Cl), bromo (Br), and iodo (I)).

The term "haloalkyl" means alkyl having the specified number of carbon atoms in which from one to all of the hydrogen atoms have been replaced by a halogen atom.

The terms "aralkyl" and "heteroaralkyl" refer to an aryl/heteroaryl linked to the rest of the molecule via a C₁ to C₄ alkylene.

The term "C₀" as employed in expressions such as "C₀₋₆ alkylene" means a direct covalent bond; or when employed in expressions such as "C₀₋₆ alkyl" means hydrogen. Similarly, when an integer defining the presence of a certain number of atoms in a group is equal to zero, it means that the atoms adjacent thereto are connected directly by a bond; for


example, in the structure , wherein s is an integer equal to zero, 1 or 2, the

structure is  when s is zero; or it means that the indicated atom is absent; for example -S(O)₀- means -S-.

Unless expressly stated to the contrary, an "unsaturated" ring is a partially or fully unsaturated ring. For example, an "unsaturated monocyclic C₆ carbocycle" refers to cyclohexene, cyclohexadiene, and benzene.

Unless expressly stated to the contrary, all ranges cited herein are inclusive. For example, a heterocycle described as containing from "1 to 4 heteroatoms" means the heterocycle can contain 1, 2, 3 or 4 heteroatoms.

When any variable occurs more than one time in any constituent or in any formula depicting and describing compounds of the invention, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. For variable definitions containing terms having repeated terms, e.g., (CR_iR_j)_r, where r is the integer 2, R_i is a defined variable, and R_j is a defined variable, the value of R_i may differ in each instance in which it occurs, and the value of R_j may differ in each instance in which it occurs. For example, if R_i and R_j are independently selected from the group consisting of methyl, ethyl, propyl and butyl, then (CR_iR_j)₂ can be



The term (C₁₋₆)alkyl as used hereinabove means a branched or unbranched alkyl group having 1-6 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, n-pentyl and n-hexyl. Preferred is (C₁₋₄)alkyl.

The term (C₁₋₅)alkyl means a branched or unbranched alkyl group having 1-5 carbon atoms, for example methyl, ethyl, propyl, isopropyl, butyl, tert-butyl and n-pentyl.

The term (C₁₋₄)alkyl as used herein means a branched or unbranched alkyl group having 1-4 carbon atoms, being methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl.

The term (C₁₋₃)alkoxy means an alkoxy group having 1-3 carbon atoms, the alkyl moiety being branched or unbranched.

The term (C₁₋₃)alkoxycarbonyl means an alkoxycarbonyl group having 1-3 carbon atoms in the alkoxy moiety, the alkoxy moiety having the same meaning as previously defined.

The term (di)(C₁₋₆)alkylaminocarbonyl means an alkylaminocarbonyl group, the amino group of which is monosubstituted or disubstituted independently with an alkyl group which contains 1-6 carbon atoms and which has the same meaning as previously defined. Preferred alkyl group is (C₁₋₄)alkyl.

The term (C₃₋₇)cycloalkyl means a cycloalkyl group having 3-7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. 5-6 Carbon atoms are preferred.

The term (C₃₋₅)heterocycloalkyl means a heterocycloalkyl group having 3-5 carbon atoms, including 1-3 heteroatoms selected from N, O and/or S, which may be attached via a nitrogen if feasible, or a carbon atom. Preferred number of heteroatoms is one or two. Most preferred number is one. Preferred heteroatoms are N or O. Most preferred are piperazinyl, tetrahydropyranyl, morpholinyl and pyrrolidinyl.

The term (C₂₋₉)heteroaryl means an aromatic group having 2-9 carbon atoms and 1-3 heteroatoms selected from N, O and S, like imidazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, thiophenyl or furyl, pyrazolyl, isoxazolyl or quinolyl. Preferred number of heteroatoms is one or two. Preferred heteroaryl groups are pyrazolyl, thiophenyl, isoxazolyl, pyridyl and quinolyl. The (C₂₋₅)heteroaryl group may be attached via a carbon atom or a nitrogen, if feasible.

The term (C₆₋₁₄)aryl means an aromatic hydrocarbon group having 6-14 carbon atoms, such as phenyl, naphthyl, tetrahydronaphthyl, indenyl, anthracyl, More preferred are (C₆₋₁₀)aryl groups. The most preferred aromatic hydrocarbon group is phenyl.

As used herein, the term "X_a-X_b", shall have the same meaning as the term "X_{a-b}", wherein X is any atom and a and b are any integers. For example, "C₁-C₄" shall have the same meaning as "C₁₋₄". Additionally, when referring to a functional group generically, "A^x" shall have the same meaning, and be interchangeable with, "AX", wherein "A" is any atom and "x" or "X" are any integer. For example, "R¹" shall have the same meaning, and be interchangeable with, "R1".

In the above definitions with multifunctional groups, the attachment point is at the last group.

The term "substituted" means that one or more hydrogens on the designated atom/atoms is/are replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. "Stable compound" or "stable structure" is defined as a compound or structure that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

When, in the definition of a substituent, is indicated that “all of the alkyl groups” of said substituent are optionally substituted, this also includes the alkyl moiety of an alkoxy group.

The term “optionally substituted” means optional substitution with the specified groups, radicals or moieties.

5 The use of the terms "salt", "solvate", “ester”, "prodrug" and the like, is intended to equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the inventive compounds.

The term "effective amount" as used herein, refers to an amount of the compound of Formula (I) and/or an additional therapeutic agent, or a composition thereof, that is effective
10 in producing the desired therapeutic, ameliorative, inhibitory or preventative effect when administered to a subject suffering from an RORgammaT-mediated disease or disorder. In the combination therapies of the present invention, as effective amount can refer to each individual agent or to the combination as a whole, wherein the amounts of all agents administered are together effective, but wherein the component agent of the combination may
15 not be present individually in an effective amount.

A “subject” is a human or non-human mammal. In one embodiment, a subject is a human. In another embodiment, a subject is a chimpanzee.

It should be noted that any carbon as well as heteroatom with unsatisfied valences in the text, schemes, examples and tables herein is assumed to have the sufficient number of
20 hydrogen atom(s) to satisfy the valences.

The compounds of this invention include the prodrugs, hydrates or solvates of the compounds.

Optical Isomers - Diastereomers - Geometric Isomers – Tautomers

25 The compounds of Formula I may contain asymmetric or chiral centers, and, therefore, exist in different stereoisomeric forms. It is intended that all stereoisomeric forms of the compounds of Formula (I) as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, the present invention embraces all geometric and positional isomers. For example, if a compound of Formula (I) incorporates a double bond or
30 a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within the scope of the invention.

Compounds described herein may contain an asymmetric center and may thus exist as enantiomers. Where the compounds according to the invention possess two or more asymmetric centers, they may additionally exist as diastereomers. The present invention includes all such possible stereoisomers as substantially pure resolved enantiomers, racemic mixtures thereof, as well as mixtures of diastereomers. The above Formula I is shown without a definitive stereochemistry at certain positions. The present invention includes all stereoisomers of Formula I and pharmaceutically acceptable salts thereof. Diastereoisomeric pairs of enantiomers may be separated by, for example, fractional crystallization from a suitable solvent, and the pair of enantiomers thus obtained may be separated into individual stereoisomers by conventional means, for example by the use of an optically active acid or base as a resolving agent or on a chiral HPLC column. Further, any enantiomer or diastereomer of a compound of the general Formula I may be obtained by stereospecific synthesis using optically pure starting materials or reagents of known configuration.

When compounds described herein contain olefinic double bonds, unless specified otherwise, such double bonds are meant to include both E and Z geometric isomers.

Some of the compounds described herein may exist with different points of attachment of hydrogen, referred to as tautomers. For example, compounds including carbonyl $-\text{CH}_2\text{C}(\text{O})-$ groups (keto forms) may undergo tautomerism to form hydroxyl $-\text{CH}=\text{C}(\text{OH})-$ groups (enol forms). Both keto and enol forms, individually as well as mixtures thereof, are included within the scope of the present invention.

Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g. chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (e.g. hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of Formula (I) may be atropisomers (e.g. substituted biaryls) and are considered as part of this invention. Enantiomers can also be separated by use of chiral HPLC column.

It is also possible that the compounds of Formula I may exist in different tautomeric forms, and all such forms are embraced within the scope of the invention. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the invention.

All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this invention, as are positional isomers. Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the IUPAC 1974 Recommendations.

15 Salts

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids. When the compound of the present invention is acidic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic bases, including inorganic bases and organic bases. Salts derived from such inorganic bases include aluminum, ammonium, calcium, copper (ic and ous), ferric, ferrous, lithium, magnesium, manganese (ic and ous), potassium, sodium, zinc and the like salts. Preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts prepared from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines derived from both naturally occurring and synthetic sources. Pharmaceutically acceptable organic non-toxic bases from which salts can be formed include, for example, arginine, betaine, caffeine, choline, N,N'-dibenzyl-ethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, dicyclohexylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

When the compound of the present invention is basic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic inorganic and organic acids. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, 5 lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The compounds of Formula I can form salts which are also within the scope of this invention. Reference to a compound of Formula I herein is understood to include reference to 10 salts thereof, unless otherwise indicated.

The term pharmaceutically acceptable salt represents those salts which are, within the scope of medical judgement, suitable for use in contact for the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well 15 known in the art. They may be obtained during the final isolation and purification of the compounds of the invention, or separately by reacting the free base function with a suitable mineral acid such as hydrochloric acid, phosphoric acid, or sulfuric acid, or with an organic acid such as for example ascorbic acid, citric acid, tartaric acid, lactic acid, maleic acid, malonic acid, fumaric acid, glycolic acid, succinic acid, propionic acid, acetic acid, 20 methanesulfonic acid, and the like. The acid function can be reacted with an organic or a mineral base, like sodium hydroxide, potassium hydroxide, calcium hydroxide, calcium carbonate, ammonium (e.g. diethylamine) or lithium hydroxide.

Solvates

25 The present invention includes within its scope solvates of compounds of Formula I. As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (i.e., a compound of Formula I) or a pharmaceutically acceptable salt thereof and a solvent that does not interfere with the biological activity of the solute. Examples of solvents include, but are not limited to water, ethanol, and acetic acid. When the solvent is water, the 30 solvate is known as hydrate; hydrate includes, but is not limited to, hemi-, mono, sesqui-, di- and trihydrates.

The compounds of the invention may form hydrates or solvates. It is known to those of skill in the art that charged compounds form hydrated species when lyophilized with water, or form solvated species when concentrated in a solution with an appropriate organic solvent. One or more compounds of the invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. "Solvate" may also mean a physical association of a compound of this invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanولات, methanولات, and the like. "Hydrate" is a solvate wherein the solvent molecule is H₂O.

Prodrugs

The present invention includes within its scope the use prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible in vivo into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with a compound of formula I or with a compound which may not be a compound of formula I, but which converts to a compound of formula I in vivo after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," ed. H. Bundgaard, Elsevier, 1985.

A discussion of prodrugs is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems (1987) 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, (1987) Edward B. Roche, ed., American Pharmaceutical Association and Pergamon Press. The term "prodrug" means a compound (e.g, a drug precursor) that is transformed in vivo to yield a compound of Formula I or a pharmaceutically acceptable salt, hydrate or solvate of the compound. The transformation may occur by various mechanisms (e.g. by metabolic or chemical processes), such as, for example, through hydrolysis in blood. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as

Novel Delivery Systems,” Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

5 Isotopes

In the compounds of generic Formula I, the atoms may exhibit their natural isotopic abundances, or one or more of the atoms may be artificially enriched in a particular isotope having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number predominantly found in nature. The present invention is meant
10 to include all suitable isotopic variations of the compounds of generic Formula I. For example, different isotopic forms of hydrogen (H) include protium (^1H) and deuterium (^2H). Protium is the predominant hydrogen isotope found in nature. Enriching for deuterium may afford certain therapeutic advantages, such as increasing in vivo half-life or reducing dosage requirements, or may provide a compound useful as a standard for characterization of
15 biological samples. Isotopically-enriched compounds within generic Formula I can be prepared without undue experimentation by conventional techniques well known to those skilled in the art or by processes analogous to those described in the Schemes and Examples herein using appropriate isotopically-enriched reagents and/or intermediates.

20 Utilities

Compounds of the present invention are inhibitors of Retinoic Acid Receptor-related Orphan Receptor gamma t (RORgammaT), and as such are useful in the treatment of diseases and conditions in which inhibition of RORgammaT is desirable, such as autoimmune and inflammatory diseases and disorders.

25 Accordingly, another embodiment of the present invention provides a method for treating a disease or condition mediated by RORgammaT in a subject comprising administering to the subject an amount of a compound having Formula I, Ia, Ib, Ic, Id or Ie or a pharmaceutically acceptable salt or solvate thereof, that is effective for treating the disease or condition mediated by RORgammaT in the subject.

30 The compounds according to the invention can be used in therapy.

A further aspect of the invention resides in the use of compounds according to the invention or a pharmaceutically acceptable salt thereof for the treatment of RORgammaT-mediated diseases or RORgammaT mediated conditions.

Another aspect of the invention resides in the use of compounds or a pharmaceutically acceptable salt thereof having the general formula I for the treatment of autoimmune diseases, in particular those diseases in which Th17 cells and non-Th17 cells, which express Th17 hallmark cytokines play a prominent role. These include, but are not limited to, the treatment of rheumatoid arthritis, psoriasis, inflammatory bowel disease, Crohn's disease and multiple sclerosis.

In another aspect, compounds or a pharmaceutically acceptable salt thereof having the general formula I can be used for treatment of inflammatory diseases in which Th17 cells and/or non-Th17 cells, which express Th17 hallmark cytokines play a prominent role such as, but not limited to respiratory diseases, osteoarthritis and asthma. Also, compounds or a pharmaceutically acceptable salt thereof having the general formula I can be used for treatment of infectious diseases in which Th17 cells and/or non-Th17 cells, which express Th17 hallmark cytokines play a prominent role such as, but not limited to mucosal leishmaniasis.

Compounds or a pharmaceutically acceptable salt thereof having the general formula I can also be used for treatment of other diseases in which Th17 cells and/or non-Th17 cells, which express Th17 hallmark cytokines play a prominent role such as, but not limited to Kawasaki disease and Hashimoto's thyroiditis.

In one aspect is the disease or condition is an autoimmune disease or inflammatory disease. The disease or condition includes, but is not limited to, multiple sclerosis, inflammatory bowel disease, Crohn's disease, psoriasis, rheumatoid arthritis, asthma, osteoarthritis, Kawasaki disease, Hashimoto's thyroiditis or mucosal leishmaniasis.

In another aspect, the compounds according to the invention can be used in therapies to treat or prevent multiple sclerosis, inflammatory bowel disease, Crohn's disease, psoriasis, rheumatoid arthritis, asthma, osteoarthritis, Kawasaki disease, Hashimoto's thyroiditis and mucosal leishmaniasis.

In another aspect the compounds according to the invention can be used to treat or prevent psoriasis.

In yet another aspect the compounds according to the invention can be used to treat

inflammatory bowel disease.

This aspect of the present invention further includes the use of a compound of Formula I, Ia, Ib, Ic, Id or Ie, or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for the treatment of a disease or condition mediated by
5 RORgammaT.

Route of Administration/Dosage

The compounds of this invention can be administered for the treatment or prevention of afflictions, diseases and illnesses according to the invention by any means that effects contact
10 of the active ingredient compound with the site of action in the body of a warm-blooded animal. For example, administration can be oral, topical, including transdermal, ocular, buccal, intranasal, inhalation, intravaginal, rectal, intracisternal and parenteral. The term "parenteral" as used herein refers to modes of administration which include subcutaneous, intravenous, intramuscular, intraarticular injection or infusion, intrasternal and intraperitoneal.
15 For the purpose of this disclosure, a warm-blooded animal is a member of the animal kingdom possessed of a homeostatic mechanism and includes mammals and birds.

The compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but are generally administered with a
20 pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage administered will be dependent on the age, health and weight of the recipient, the extent of disease, kind of concurrent treatment, if any, frequency of treatment and the nature of the effect desired. Usually, a daily dosage of active ingredient compound
25 will be from about 1.0-2000 milligrams per day. Ordinarily, from 10 to 500 milligrams per day in one or more applications is effective to obtain desired results. These dosages are the effective amounts for the treatment and prevention of afflictions, diseases and illnesses described above, e.g., autoimmune and inflammatory diseases and disorders.

Compositions include e.g. those suitable for oral, sublingual, subcutaneous, intravenous,
30 intramuscular, nasal, local, or rectal administration, and the like, all in unit dosage forms for administration.

For oral administration, the active ingredient may be presented as discrete units, such as tablets, capsules, powders, granulates, solutions, suspensions, and the like.

For parenteral administration, the pharmaceutical composition of the invention may be presented in unit-dose or multi-dose containers, e.g. injection liquids in predetermined
5 amounts, for example in sealed vials and ampoules, and may also be stored in a freeze dried (lyophilized) condition requiring only the addition of sterile liquid carrier, e.g. water, prior to use.

Mixed with such pharmaceutically acceptable auxiliaries, e.g. as described in the standard reference, Gennaro, A.R. et al., Remington: *The Science and Practice of Pharmacy*
10 (20th Edition., Lippincott Williams & Wilkins, 2000, see especially Part 5: Pharmaceutical Manufacturing), the active agent may be compressed into solid dosage units, such as pills, tablets, or be processed into capsules or suppositories. By means of pharmaceutically acceptable liquids the active agent can be applied as a fluid composition, e.g. as an injection preparation, in the form of a solution, suspension, emulsion, or as a spray, e.g. a nasal spray.

15 For making solid dosage units, the use of conventional additives such as fillers, colorants, polymeric binders and the like is contemplated. In general any pharmaceutically acceptable additive which does not interfere with the function of the active compounds can be used. Suitable carriers with which the active agent of the invention can be administered as solid compositions include lactose, starch, cellulose derivatives and the like, or mixtures
20 thereof, used in suitable amounts. For parenteral administration, aqueous suspensions, isotonic saline solutions and sterile injectable solutions may be used, containing pharmaceutically acceptable dispersing agents and/or wetting agents, such as propylene glycol or butylene glycol.

25 Pharmaceutical Compositions

Another aspect of the present invention provides pharmaceutical compositions comprising a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof and one or more pharmaceutically acceptable excipients. The term "excipient" and "carrier" may be used interchangeably. The term "composition", as in pharmaceutical composition, is
30 intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation

of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients.

Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of Formula I, additional active ingredient(s), and
5 pharmaceutically acceptable excipients.

The pharmaceutical compositions of the present invention comprise a compound represented by Formula I (or pharmaceutically acceptable salts thereof) as an active ingredient, a pharmaceutically acceptable carrier and optionally other therapeutic ingredients or adjuvants. The compositions include compositions suitable for oral, rectal, topical, and
10 parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

15 The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, troches, dragées, granules and powders, or in liquid dosage forms, such as elixirs, syrups, emulsions, dispersions, and suspensions. The active ingredient can also be administered parenterally, in sterile liquid dosage forms, such as dispersions, suspensions or solutions. Other dosage forms that can also be used to administer the active ingredient as an
20 ointment, cream, drops, transdermal patch or powder for topical administration, as an ophthalmic solution or suspension formation, i.e., eye drops, for ocular administration, as an aerosol spray or powder composition for inhalation or intranasal administration, or as a cream, ointment, spray or suppository for rectal or vaginal administration.

Gelatin capsules contain the active ingredient and powdered carriers, such as lactose,
25 starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the
30 gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propylparaben, and chlorobutanol.

Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field.

For administration by inhalation, the compounds of the present invention may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery system for inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a compound of Formula I in suitable propellants, such as fluorocarbons or hydrocarbons.

For ocular administration, an ophthalmic preparation may be formulated with an appropriate weight percent solution or suspension of the compounds of Formula I in an appropriate ophthalmic vehicle, such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention include, but are not limited to, hard and soft gelatin capsules, tablets, parenteral injectables, and oral suspensions.

A large number of unit capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to
5 increase palatability or delay absorption.

A parenteral composition suitable for administration by injection is prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol. The solution is made to volume with water for injection and sterilized.

An aqueous suspension is prepared for oral administration so that each 5 milliliters
10 contain 100 milligrams of finely divided active ingredient, 100 milligrams of sodium carboxymethyl cellulose, 5 milligrams of sodium benzoate, 1.0 grams of sorbitol solution, U.S.P., and 0.025 milliliters of vanillin.

The same dosage forms can generally be used when the compounds of this invention are administered stepwise or in conjunction with another therapeutic agent. When drugs are
15 administered in physical combination, the dosage form and administration route should be selected depending on the compatibility of the combined drugs. Thus the term coadministration is understood to include the administration of the two agents concomitantly or sequentially, or alternatively as a fixed dose combination of the two active components.

The present invention also relates to a pharmaceutical composition comprising
20 compounds or pharmaceutically acceptable salts thereof having the general formula I in admixture with pharmaceutically acceptable auxiliaries and optionally other therapeutic agents. The auxiliaries must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof.

The invention further includes a pharmaceutical composition, as hereinbefore described,
25 in combination with packaging material suitable for said composition, said packaging material including instructions for the use of the composition for the use as hereinbefore described.

The exact dose and regimen of administration of the active ingredient, or a pharmaceutical composition thereof, may vary with the particular compound, the route of
30 administration, and the age and condition of the individual subject to whom the medicament is to be administered.

In general parenteral administration requires lower dosages than other methods of administration which are more dependent upon absorption. However, a dosage for humans preferably contains 0.0001-100 mg per kg body weight. The desired dose may be presented as one dose or as multiple subdoses administered at appropriate intervals throughout the day.

5 The dosage as well as the regimen of administration may differ between a female and a male recipient.

Combination Therapy

Compounds of the present invention, and their salts and solvates, and physiologically functional derivatives thereof, may be employed alone or in combination with other therapeutic agents for the treatment of diseases and conditions associated with inappropriate IL-17 pathway activity. Combination therapies according to the present invention thus comprise the administration of at least one compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, or a physiologically functional derivative thereof, and the use of at least one other pharmaceutically active agent. The compound(s) of formula (I) and the other pharmaceutically active agent(s) may be administered together or separately and, when administered separately this may occur simultaneously or sequentially in any order. The amounts of the compound(s) of formula (I) and the other pharmaceutically active agent(s) and the relative timings of administration will be selected in order to achieve the desired combined therapeutic effect. For the treatment of the inflammatory and autoimmune diseases, rheumatoid arthritis, psoriasis, inflammatory bowel disease, SLE, uveitis, atopic dermatitis, COPD, asthma and allergic rhinitis a compound of formula (I) may be combined with one or more other active agents such as: (1) TNF- α inhibitors; (2) non-selective COX-I/COX-2 inhibitors; (3) COX-2 inhibitors; (4) other agents for treatment of inflammatory and autoimmune diseases including glucocorticoids, methotrexate, leflunomide, sulfasalazine, azathioprine, cyclosporin, tacrolimus, penicillamine, bucillamine, actarit, mizoribine, lobenzarit, ciclesonide, hydroxychloroquine, d-penicillamine, aurothiomalate, auranofin or parenteral or oral gold, cyclophosphamide, Lymphostat-B, BAFF/APRIL inhibitors and CTLA-4-Ig or mimetics thereof; (5) leukotriene biosynthesis inhibitor, 5-lipoxygenase (5-LO) inhibitor or 5-lipoxygenase activating protein (FLAP) antagonist; (6) LTD4 receptor antagonist; (7) PDE4 inhibitor; (8) antihistamine H1 receptor antagonists; (9) α_1 - and α_2 -adrenoceptor agonist; (10) anticholinergic agents; (11) β -adrenoceptor agonists; (12) insulin-

10
15
20
25
30

like growth factor type I (IGF-1) mimetic; (13) glucocorticosteroids; (14) kinase inhibitors such as inhibitors of the Janus Kinases (JAK 1 and/or JAK2 and/or JAK 3 and/or TYK2), p38 MAPK and IKK2; (15) B-cell targeting biologics such as rituximab; (16) selective costimulation modulators such as abatacept; (17) interleukin inhibitors, such as IL-1 inhibitor anakinra, IL-6 inhibitor tocilizumab, and IL12/IL-23 inhibitor ustekinumab. It could also be combined with anti-IL17 antibodies to obtain additive/synergistic responses for the treatment of inflammatory and autoimmune diseases.

It will be clear to a person skilled in the art that, where appropriate, the other therapeutic ingredient(s) may be used in the form of salts, for example as alkali metal or amine salts or as acid addition salts, or prodrugs, or as esters, for example lower alkyl esters, or as solvates, for example hydrates, to optimise the activity and/or stability and/or physical characteristics, such as solubility, of the therapeutic ingredient. It will be clear also that, where appropriate, the therapeutic ingredients may be used in optically pure form.

The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical composition and thus pharmaceutical compositions comprising a combination as defined above together with a pharmaceutically acceptable diluent or carrier represent a further aspect of the invention. These combinations are of particular interest in respiratory diseases and are conveniently adapted for inhaled or intranasal delivery.

The individual compounds of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical compositions. Preferably, the individual compounds will be administered simultaneously in a combined pharmaceutical composition. Appropriate doses of known therapeutic agents will be readily appreciated by those skilled in the art.

Accordingly, the pharmaceutical compositions of the present invention include those that also comprise at least one additional therapeutically active agent, in addition to the compound of Formula I, Ia, Ib, Ic, Id or Ie.









The invention further includes a compound of Formula I in combination with one or more other drug(s).











Synthesis








Table 1 below lists commercial sources, and previously disclosed synthetic routes, for chemical materials employed in the synthesis of intermediates and examples of the instant invention. This list is not intended to be exhaustive, exclusive, or limiting in any way.








5


Table 1

Structure	Source
	Oakwood
	Aldrich
	Frontier
	Oakwood
	Sinova
	Sinova
	Aldrich
	Sinova

	Sinova
	Aldrich
	Sinova
	Alfa
	Alfa
	<i>Bioorganic & Medicinal Chemistry Letters</i> , 2006, 695-700
	Acros
	Combi-blocks
	Combi-blocks
	Alfa

 <chem>COC(=O)c1ccc(Cl)c(c1)B(O)O</chem>	Combi-blocks
 <chem>NC(=O)c1ccc(B(O)O)cc1</chem>	Alfa
 <chem>COC(=O)c1cc(C)cc(B(O)O)c1</chem>	Combi-blocks
 <chem>COC(=O)c1cc(OC)cc(B(O)O)c1</chem>	Combi-blocks
 <chem>COC(=O)c1cc(F)cc(B(O)O)c1</chem>	Alfa
 <chem>CC1=CC=C(C(=C1)C(=O)O)B(O)O</chem>	Combi-blocks
 <chem>COC(=O)c1cc(OC)cc(B(O)O)c1</chem>	Anichem

 <chem>COC(=O)c1cc(F)c(B(O)O)cc1F</chem>	Aobchem
 <chem>COC(=O)c1ccc(C#N)cc1B(OC1(C)C(C)(C)C1(C)C)OC2(C)C(C)(C)C2(C)C</chem>	Chemmaker
 <chem>OC(=O)c1cc(F)c(B(O)O)cc1F</chem>	Combi-blocks
 <chem>CC(C)(C)OC(=O)c1cc(Cl)c(B(OC2(C)C(C)(C)C2(C)C)OC3(C)C(C)(C)C3(C)C)c(F)c1</chem>	Combi-blocks
 <chem>OC(=O)c1cc(OC)c(B(O)O)cc1F</chem>	Anisyn
 <chem>COC(=O)c1cc(O)cc(B(O)O)c1</chem>	Anisyn
 <chem>COC(=O)c1cc(B(O)O)c(F)cc1</chem>	Alfa




Abbreviations used herein are as follows: Boc: *tert*-butoxycarbonyl; CDCl₃: chloroform-d; DiPEA: *N,N*-diisopropylethylamine; DMAP: 4-dimethylaminopyridine; DMF: *N,N*-dimethylformamide; Et₃N or TEA: triethyl amine; HPLC: high performance liquid chromatography; TFA: trifluoro acetic acid; MS: mass spectrum; (PPh₃)₄Pd: tetrakis-(triphenylphosphine)palladium(0); THF: tetrahydrofuran; TLC: thin layer chromatography; SiO₂: silica-gel; DME: dimethoxy ethane; m: molar; t-BuOK; potassium *tert*-butoxide; APCI-MS: atmospheric pressure chemical ionization mass spectrometry; ESI-MS; electrospray ionization-mass spectroscopy; EtOAc: Ethyl acetate; PE: Petroleum ether; EA: Ethyl acetate; DCM: Dichloro methane; Dppf: 1,1'-Bis(diphenylphosphino)ferrocene; AcOH: Acetic acid; DMAC: *N,N*-Dimethylacetamide; PYAOP: (7-Azabenzotriazol-1-yloxy)tripyrrolid in ophosphonium hexafluorophosphate; Pd(dppf)Cl₂: [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium (II); BnBr: Benzyl bromide; DAST: (Diethylamino)sulfur trifluoride; Ac₂O: Acetic anhydride; LiHMDS: Lithium bis(trimethylsilyl)amide; PhNTf₂: *N*-Phenyl-bis(trifluoromethanesulfonimide); S-Phos: 2-Dicyclohexylphosphino-2',6'-dimethoxybiphenyl; X-Phos: 2-Dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl; CPME: Cyclopentyl methyl ether.

20

Intermediates

Example i-1: Preparation of methyl 2-isopropyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-1)

Scheme i-1.



i) Preparation of methyl 4-(benzyloxy)-2-hydroxybenzoate (i-1b). To a solution of methyl 2,4-dihydroxybenzoate (**i-1a**) (1.68 g, 10 mmol) in acetone (30 mL) was added K_2CO_3 (2.76 g, 20 mmol). The mixture was stirred at rt for 5 h, followed by the addition of BnBr (1.88 g, 11 mmol). The mixture was heated at 60 °C for 24 h. The reaction mixture was cooled down, filtered. The filtrate was concentrated to afford the title compound. LCMS (ESI) calc'd for $\text{C}_{15}\text{H}_{14}\text{O}_4$ $[\text{M}+\text{H}]^+$: 259.1, found: 259.1.

ii) Preparation of methyl 4-(benzyloxy)-2-(trifluoromethylsulfonyloxy)benzoate (i-1c). To a solution of methyl 4-(benzyloxy)-2-hydroxybenzoate (**i-1b**) (1.1 g, 4.26 mmol) in THF (20 ml) at -78 °C was added LiHMDS (1M in THF, 5.1 ml, 5.1 mmol) dropwise. Then the mixture was warmed up to -40 °C and stirred at this temperature for 3 h. PhNTf_2 (1.52 g, 4.26

mmol) was added, and the mixture was stirred at rt for 24 h. The mixture was then diluted with H₂O, and extracted with EtOAc. The organic layer was dried, concentrated, and purified by flash chromatography (PE/EA=10:1) to afford the title compound. LCMS (ESI) calc'd for C₁₆H₁₃F₃O₆S [M+H]⁺: 391, found: 391.

5

iii) Preparation of methyl 4-(benzyloxy)-2-(prop-1-en-2-yl)benzoate (i-1d). Methyl 4-(benzyloxy)-2-(trifluoromethylsulfonyloxy)benzoate (**i-1c**) (6.8 g, 17.4 mmol), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (2.9 g, 17.4 mmol), Pd(OAc)₂ (195 mg, 0.87 mmol), S-Phos (71 mg, 0.174 mmol) and K₃PO₄ (11 g, 52 mmol) were mixed in THF
10 (70 ml). The mixture was stirred at 75 °C for 24 h. The mixture was diluted with H₂O, and extracted with EtOAc. The combined organics were dried and concentrated. The residue was purified by flash chromatography (PE/EA=10:1) to afford the title compound. LCMS (ESI) calc'd for C₁₈H₁₈O₃ [M+H]⁺: 391, found: 391.

iv) Preparation of methyl 4-hydroxy-2-isopropylbenzoate (i-1e). To a solution of methyl 4-(benzyloxy)-2-(prop-1-en-2-yl)benzoate (**i-1d**) (4.6 g, 16.3 mmol) in methanol (100ml) was added Pd/C (0.46 g). The mixture was stirred at rt under H₂ atmosphere for 24 h. Then the mixture was filtered, and the filtrate was concentrated to afford the title compound. LCMS (ESI) calc'd for C₁₁H₁₄O₃ [M+H]⁺: 195.1, found: 195.1.

20

v) Preparation of methyl 2-isopropyl-4-(trifluoromethylsulfonyloxy)benzoate (i-1f). To a solution of methyl 4-hydroxy-2-isopropylbenzoate (**i-1e**) (2.45 g, 12.6 mmol) in THF (100 ml) at -78 °C was added LiHMDS (1M in THF, 15 ml, 15 mmol) dropwise. Then the mixture was warmed up to -40 °C and stirred for 3 h. PhNTf₂ (4.5 g, 12.6 mmol) was added and the
25 mixture was stirred at rt for 24 h. The reaction mixture was diluted with H₂O and extracted with EtOAc. The organic layer was dried and concentrated. The residue was purified by flash chromatography (PE/EA=8:1) to afford the title compound. LCMS (ESI) calc'd for C₁₂H₁₃F₃O₅S [M+H]⁺: 327, found: 327.


vi) Preparation of methyl 2-isopropyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan- 2-yl)benzoate (i-1). Methyl 2-isopropyl-4-(trifluoromethylsulfonyloxy)benzoate (**i-1f**) (1.5 g, 4.6 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (1.75 g, 6.9 mmol),

30

Pd(dppf)Cl₂.CH₂Cl₂ (188 mg, 0.23 mmol), dppf (102 mg, 0.184 mmol) and KOAc (1.13 g, 11.5 mmol) were mixed in 1,4-dioxane (35 ml). The mixture was stirred at 80 °C for 24 h. The mixture was then cooled down and concentrated under reduced pressure. The residue was diluted with H₂O, extracted with EtOAc. The combined organics were dried and concentrated. The residue was purified by flash chromatography (PE/EA=10:1) to afford the title compound. LCMS (ESI) calc'd for C₁₇H₂₅BO₄ [M+H]⁺: 305.2, found: 305.2.

Example i-2: Preparation of methyl 2-fluoro-5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-2)

Scheme i-2.




i) Preparation of methyl 4-bromo-2-fluoro-5-methylbenzoate (i-2b). To a solution of 4-bromo-2-fluoro-5-methylbenzoic acid (**i-2a**) (1.87 g, 8 mmol) in CH₃OH (50 mL) at 0 °C was added SOCl₂ (6 mL, 80 mmol) dropwise. Then the mixture was heated to 80 °C for 2 h. The solvent was removed *in vacuo* and the residue was diluted with EtOAc and washed with H₂O, brine, dried over anhydrous Na₂SO₄, and concentrated. The resulting crude solid was washed with PE to give the title compound as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 7.80 (1H, d), 7.37 (1H, d), 3.92 (3H, s), 2.39 (3H, s).

ii) Preparation of methyl 2-fluoro-5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-2). To a flask was added compound (**i-2b**) (830 mg, 3.3 mmol), compound (**i-2c**) (915 mg, 3.6 mmol), dppf (25 mg, 0.132 mmol), Pd(dppf)Cl₂.CH₂Cl₂ (134 mg, 0.165 mmol) and 1,4-dioxane (20 mL). The reaction mixture was heated at 80 °C for 2 h. The mixture was diluted with H₂O, and the organic layer was separated. The aqueous layer was

extracted with CH_2Cl_2 . The combined organics were washed with H_2O and brine, dried over Na_2SO_4 , and concentrated. The residue was purified by flash chromatography (Pentane/EtOAc) to afford the title compound. LCMS (ESI) calc'd for $\text{C}_{15}\text{H}_{20}\text{BFO}_4$ $[\text{M}+\text{H}]^+$: 295.1, found: 295.1.

5

Example i-3: Preparation of methyl 3,5-dimethoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-3)



10

i) Preparation of methyl 4-bromo-3,5-dimethoxybenzoate (i-3b).

Prepared following a similar procedure of compound **i-2b**

LCMS (ESI) calc'd for $\text{C}_{10}\text{H}_{11}\text{BrO}_4$ $[\text{M}+\text{H}]^+$: 275.1, found: 275.1.


15 ii) Preparation of methyl 3,5-dimethoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-3)

Prepared following a similar procedure of compound **i-2**

LCMS (ESI) calc'd for $\text{C}_{16}\text{H}_{23}\text{BO}_6$ $[\text{M}+\text{H}]^+$: 323.2, found: 323.2.

20 Example i-4: Preparation of methyl 2,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-4)


Scheme i- 4.



i) Preparation of methyl 2-fluoro-5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-4). To a flask was added compound (**i-4a**) (412 mg, 2.0 mmol), compound (**i-2c**) (1.52 g, 6 mmol), X-Phos (95 mg, 0.2 mmol), $\text{Pd}(\text{OAc})_2$ (44 mg, 0.2 mmol) and CPME (20 mL). The reaction mixture was heated at 85 °C for 2 h. The mixture was then cooled down, diluted with H_2O , and the organic layer was separated. The aqueous layer was extracted with Et_2O . The combined organics were washed with H_2O and brine, dried over Na_2SO_4 , and concentrated. The residue was purified by flash chromatography (DCM) to afford the title compound. LCMS (ESI) calc'd for $\text{C}_{14}\text{H}_{17}\text{BF}_2\text{O}_4$ $[\text{M}+\text{H}]^+$: 299.1, found: 299.1.

Example i-5: Preparation of 3-fluoro-5-methoxy-4-(methoxycarbonyl)phenyl boronic acid (i-5)

Scheme i-5.



i) Preparation of methyl 4-bromo-2,6-difluorobenzoate (i-5b)

Prepared following a similar procedure of compound **i-2b**

LCMS (ESI) calc'd for $\text{C}_8\text{H}_5\text{BrF}_2\text{O}_2$ $[\text{M}+\text{H}]^+$: 250.9, Found: 250.9.

ii) Preparation of methyl 4-bromo-2-fluoro-6-methoxybenzoate (i-5c).


To a solution of compound **(i-5b)** (827 mg, 3.3 mmol) in DMF (20 mL) at 0 °C was added a solution of sodium methoxide (25% in MeOH, 0.75 mL, 3.3 mmol). The reaction mixture was stirred at 0 °C for 10 min and then at rt for 30 min. The mixture was then brought back to 0 °C and quenched with EtOAc and 1M HCl. The organic layer was separated and the aqueous
 5 phase was extracted with EtOAc. The combined organics were dried over Na₂SO₄ and concentrated. The crude oil was purified by flash chromatography (Pentane/EtOAc) to afford the title compound. LCMS (ESI) calc'd for C₉H₈BrFO₃ [M+H]⁺: 263.0, found: 263.0; ¹H NMR (400 MHz, CDCl₃) δ 6.98 (1H, t), 6.91 (1H, q), 3.76 (3H, s), 3.74 (3H, s).

10 **ii) Preparation of 3-fluoro-5-methoxy-4-(methoxycarbonyl)phenylboronic acid (i-5)**

Prepared following a similar procedure of compound **i-2**

LCMS (ESI) calc'd for C₉H₁₀BFO₅ [M+H]⁺: 229.1, found: 229.1.

Example i-6: Preparation of 4-borono-3,6-difluoro-2-methoxybenzoic acid (i-6)



i) Preparation of 3,6-difluoro-2-methoxybenzoic acid (i-6b).


A solution of 1,4-difluoro-2-methoxybenzene (**i-6a**) (2.88 g, 20 mmol) in THF (5 mL) was
 20 added to a solution of LDA freshly prepared from n-BuLi (10M solution in hexane, 2.2 mL, 22 mmol) and diisopropylamine (2.28 g, 22.8 mmol) in THF (20 mL) at -60 °C. After 30min, a freshly prepared solution of CO₂ in diethyl ether (20 mL) was quickly added. Then aq. H₂SO₄ was added, and the resulting precipitate was collected and washed with H₂O, and dried to give the title compound (**i-6b**) as white crystals. LCMS (ESI) calc'd for C₈H₆F₂O₃ [M+H]⁺:
 25 189, found: 189.

ii) Preparation of 4-borono-3,6-difluoro-2-methoxybenzoic acid (i-6).

To a solution of 3,6-difluoro-2-methoxybenzoic acid (**i-6b**) (0.94 g, 5 mmol) in THF (5 mL) at -70 °C was added a solution of LDA [freshly prepared from diisopropylamine (1.51 g, 15 mmol) and n-BuLi (10M solution in hexane, 1.5 mL, 15mol) in THF (15 mL). The resultant solution was stirred for 15 mins, followed by the addition of B(OEt)₃ (1.46 g, 10 mmol). The mixture was stirred for 15mins and then hydrolyzed with dilute aq. H₂SO₄. The organic phase was separated and the aqueous layer was extracted with Et₂O. The combined organics were concentrated and the residue solid was washed with H₂O and hexane, dried to give the title compound as white crystals. LCMS (ESI) calc'd for C₈H₇BF₂O₅ [M+H]⁺: 233, found: 233.

Example i-7: methyl 5-fluoro-2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-7)

Scheme i-7.



i) Preparation of 4-bromo-2,5-difluorobenzoic acid (i-7b)

To a solution of 1,4-dibromo-2,5-difluorobenzene (**i-7a**) (1.1 g, 4 mmol) in Et₂O (10 mL) at -78 °C was added n-BuLi (2.5M in hexane, 1.6 mL, 4 mmol) dropwise. The mixture was stirred for 30 min at -78 °C and then quenched with an excess of freshly crushed dry ice. After 15 min, the mixture was brought to rt and diluted with H₂O. The aqueous layer was separated, and the organic layer was extracted with 10% aq. Na₂CO₃. The combined aqueous layers were acidified with 1M HCl and extracted with EtOAc. The combined organic layers

were dried with Na₂SO₄ and concentrated to give the title compound. LCMS (ESI) calc'd for C₇H₃BrF₂O₂ [M+H]⁺: 237, found: 237.

ii) Preparation of methyl 4-bromo-2,5-difluorobenzoate (i-7c)

5 A mixture of 4-bromo-2,5-difluorobenzoic acid (**i-7b**) (0.48 g, 2 mmol) and methanol (10 mL, saturated with gaseous HCl) was heated at 65 °C for 2 h. The mixture was concentrated. Then MeOH was added and the mixture was concentrated again. The residue was purified by flash chromatography (Pentane/EtOAc=4:1) to give the title compound. LCMS (ESI) calc'd for C₈H₅BrF₂O₂ [M+H]⁺: 251, found: 251.

10

iii) Preparation of methyl 4-bromo-5-fluoro-2-methoxybenzoate (i-7d)


To a solution of methyl 4-bromo-2,5-difluorobenzoate (**i-7c**) (0.5 g, 2 mmol) in DMF (5 mL) at 0 °C was added NaOMe (25% in MeOH, 0.45 mL, 2 mmol). The reaction mixture was stirred at 0 °C for 10 min and then at rt for 30min. The mixture was then brought back to 0 °C
15 and quenched with EtOAc and 1M HCl. The organic layer was separated, and the aqueous phase was extracted with EtOAc, dried over Na₂SO₄, concentrated and purified by flash chromatography (Pentane/EtOAc=9:1) to give the title compound. LCMS (ESI) calc'd for C₉H₈BrFO₃ [M+H]⁺: 263, found: 263

20 **iv) Preparation of methyl 5-fluoro-2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (i-7)**

A mixture of Bis(pinacolato)diboron (0.72 g, 2.86 mmol), Pd(OAc)₂ (50mg), X-Phos (100 mg), methyl 4-bromo-5-fluoro-2-methoxybenzoate (**i-7d**) (0.5 g, 1.9 mmol), and KOAc (100 mg) in DMAc (20 mL) were heated at 80 °C for 24 h. The mixture was then cooled down,
25 diluted with EtOAc and H₂O. The organic layer was separated and the aqueous layer was extracted with EtOAc. The organics were combined, dried, concentrated and purified by flash chromatography (Pentane/EtOAc=10:1) to give the title compound. LCMS (ESI) calc'd for C₁₅H₂₀BFO₅ [M+H]⁺: 311, found: 311

30 **Example i-8: Preparation of (2-chloro-6-(trifluoromethyl)phenyl)(3-iodo-6-(methoxymethyl)-1H-indazol-1-yl)methanone (i-8)**

Scheme i-8.




i) Preparation of (2-chloro-6-(trifluoromethyl)phenyl)(6-(hydroxymethyl)-3-iodo-1H-indazol-1-yl)methanone (i-8b). To a solution of compound (i-8a) (prepared following the procedure of **Scheme D**) (175 mg, 0.35 mmol) in THF (20 mL) was added $\text{BH}_3 \cdot \text{THF}$ (1.75 mL, 1.75 mmol) and the mixture was refluxed for 12 h. The mixture was cooled down and quenched with excess MeOH. The mixture was concentrated to give the title compound. LCMS (ESI) calc'd for $\text{C}_{16}\text{H}_9\text{ClF}_3\text{IN}_2\text{O}_2$ $[\text{M}+\text{H}]^+$: 480.9, found: 480.9.

ii) Preparation of (2-chloro-6-(trifluoromethyl)phenyl)(3-iodo-6-(methoxy methyl)-1H-indazol-1-yl)methanone (i-8). To a mixture of (i-8b) (0.2 g, 0.416 mmol) in THF (5 mL) was added NaH (25 mg, 0.62 mmol) and the mixture was stirred at room temperature for 30 min. Then iodomethane (0.1 mL, 0.83 mmol) was added dropwise. The reaction mixture was stirred at rt for 24 h. Ammonium Hydroxide (5 mL) was added to quench the reaction. The solvent was removed under reduced pressure and the residue was diluted with aq. NH_4Cl . The mixture was extracted with EtOAc, and the combined organics were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep-TLC (PE:EA 4:1) to provide the title compound as a pale yellow solid. LCMS (ESI) calc'd for $\text{C}_{17}\text{H}_{11}\text{ClF}_3\text{IN}_2\text{O}_2$ $[\text{M}+\text{H}]^+$: 495, found: 495.


Method for Preparing Compounds of the Invention

Methods for preparing the compounds of this invention are illustrated in the following schemes. Other synthetic protocols will be readily apparent to those skilled in the art. The examples illustrate the preparation of the compounds of Formula I and as such are not to be considered as limiting the invention set forth in the claims appended hereto. Unless otherwise indicated, all variables are as previously defined.


SCHEME 1




The heterocyclic derivatives of Formula I, wherein A⁴ to A⁷, R¹, R², R³, X and Y are as previously defined, can be prepared from compounds II. Palladium catalyzed Suzuki–Miyaura arylation of compounds II using appropriately substituted phenylboronic acids III-a (corresponding boronic esters III-b may also be used) directly affords the desired compounds I. In a typical procedure, a mixture of compound II, a palladium catalyst (*e.g.* Pd(PPh₃)₄), base (*e.g.* aqueous NaOH, K₂CO₃ or the like) and an aryl boronic acid of general Formula III-a or aryl boronic ester of general formula III-b in a suitable solvent such as dioxane or toluene/EtOH, is heated under an atmosphere of nitrogen under microwave irradiation or using conventional heating.




Alternatively, compound I may be obtained from compounds of Formula IV, in which FG is a functional group (*e.g.* ester, cyano, protected tetrazole) that can be easily converted to groups defined for R². Non limiting examples of suitable functional groups in compounds IV are a *tert*-butyl ester or methyl ester or a tetrazole, containing an acid labile group such as a *para*-methoxy benzyl protecting group, which can be treated with acid or base to afford compounds of Formula I-a or I-b in which R² is carboxyl or 5-tetrazolyl, respectively.



Compounds II are prepared by straightforward acylation or sulfonylation of compounds V. For example, treatment of compounds V with a strong base (such as potassium *tert*-butoxide or NaH) and the appropriately substituted sulfonyl chlorides in a suitable solvent such as DMF or THF will give compounds II-a in which X = SO₂. Alternatively, treatment of V with sulfonyl chlorides and a tertiary amine base (*e.g.* DiPEA or triethylamine) in a solvent such as dichloromethane can also give compounds II-a. Treatment of V with acyl chlorides in pyridine or Et₃N will yield compounds II-b, in which X = CO. In case the acid chlorides are not available, they can be easily prepared from the corresponding carboxylic acids using known methods (*e.g.* treatment with SOCl₂ or similar reagents). Compounds V are commercially available or prepared by methods known in the art, such as halogenation of compounds VI. Compounds VI are also commercially available or easily prepared by standard organic synthesis techniques.



Compounds IV can be prepared in the same way as compounds I, by Suzuki-Miyaura reaction between compounds of general Formula II and an appropriately substituted
 5 arylboronic acid or ester.



Alternatively, a Suzuki reaction on compounds VII (obtained by Boc-protection of
 compounds V) will give compounds VIII (in most cases the Boc-group is cleaved under the
 10 reaction conditions, although a separate deprotection may be necessary). Now, acylation or
 sulfonation as described for the conversion of V to II, yields compounds IV. The functional
 group (FG in formula IV) may be a *tert*-butyl ester, or a *para*-methoxy benzyl protected
 tetrazole, which after treatment with trifluoro acetic acid will give a carboxylate or 5-
 tetrazolyl, respectively, as required for compounds I. Finally, a Suzuki reaction on
 15 unprotected compounds V, may also directly lead to compounds VIII.
 The invention is illustrated by the following examples.

General

The following procedure was used for Examples 1-23. ^1H NMR spectra were recorded on a Bruker spectrometer (400 MHz) with deuteriochloroform as the solvent unless stated otherwise. Chemical shifts are reported as δ values (parts per million) relative to tetramethylsilane as an internal standard.

5 MS: Electro Spray spectra were recorded on the Applied Biosystems API-165 single quad MS in alternating positive and negative ion mode using Flow Injection. The mass range was 120-2000 Da and scanned with a step rate of 0.2 Da. and the capillary voltage was set to 5000 V. N₂-gas was used for nebulasation.

LC-MS spectrometer (Waters) Detector: PDA (200-320 nm), Mass detector: ZQ

10 Eluens : A: acetonitrile with 0.05% trifluoroacetic acid , B: acetonitrile / water = 1/9 (v/v) with 0.05% trifluoroacetic acid

Column 1: Chromolith Performance, RP-18e, 4.6x100 mm,

Gradient method: Flow: 4 mL/min

	Time (min)	A (%)	B (%)
15	0,0	100	0
	3.60	0	100
	4.00	0	100
	4.05	100	0
	6.00	100	0

20 Column 2: XBridge C18, 3.5 μm , 4.6x20mm

Gradient method: Flow: 4 ml/min

	Time (min.)	A (%)	B (%)
	0.0	100	0
	1.60	0	100
25	3.10	0	100
	3.20	100	0
	5.00	100	0

UPLC : Water acquity UPLC system; Column : BEH C18 1.7 μm , 2.1 x 100 mm, Detector: PDA (200-320 nm), Mass detector: SQD


30 Eluens : A: acetonitrile with 0.035% trifluoroacetic acid , B: acetonitrile / water = 1/9 (v/v) with 0.035% trifluoroacetic acid

Method 60_100			Method 40_80		Method 0_60	
Flow: 0.75 mL/min			Flow: 0.65 mL/min		Flow: 0.60 mL/min	
Time (min)	A (%)	B (%)	A (%)	B (%)	A (%)	B (%)
0.0	40	60	60	40	100	0
3.00	0	100	20	80	40	60
3.20	0	100	0	100	0	100
3.69	0	100	0	100	0	100
3.70	40	60	60	40	100	0

All target compounds were characterized and determined as at least >95% pure by ¹H NMR, MS and analytical HPLC.

Examples

Example 1



4-(1-(2,6-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid

i) A solution of 3-bromo-4-azaindole (100 mg, 0.508 mmol) and 2,6-dichlorobenzoyl chloride (159 mg, 0.761 mmol) in 4 ml of pyridine was stirred for 1 h at 150 °C in a microwave reactor. After cooling to room temperature the reaction mixture was diluted with water and the product was extracted into CH₂Cl₂. The organic layer was washed with water, dried over a phase separation filter and concentrated under reduced pressure. The residue was purified on SiO₂, using 10% ethylacetate in heptane as the eluent, to give (3-bromo-1H-indazol-1-yl)(2,6-dichlorophenyl)methanone (140 mg) as a yellow solid.

ii) To a microwave reaction vial were added the product obtained in the previous step (53 mg, 0.143 mmol) in 2 ml dioxane, 4-(*tert*-butoxycarbonyl)phenylboronic acid (47.7 mg, 0.215 mmol) and a 2M aqueous solution of sodium carbonate (0.286 ml, 0.573 mmol). After

purging the vial with nitrogen for about 5 minutes, Pd(PPh₃)₄ (8.28 mg, 7.16 μmol) was added and the reaction mixture was stirred for 30 minutes at 100 °C in a microwave reactor. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate and washed with water, brine and dried over magnesium sulfate.

5 After filtration the solvent was evaporated under reduced pressure and the desired product, tert-butyl 4-(1-(2,6-dichlorobenzoyl)-1H-indazol-3-yl)benzoate, was obtained as a yellow solid (90 mg). The product was used in the next step without further purification.

iii) A solution of the product obtained in the previous step (90 mg, 0.22 mmol) and trifluoroacetic acid (5 ml, 67.3 mmol) in 20 ml CH₂Cl₂ was stirred for 1h at room temperature


10 After completion, the reaction mixture was concentrated under reduced pressure and the residue was purified on SiO₂, using 10% ethylacetate in heptane as the eluent to give the title compound 4-(1-(2,6-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid as a white solid (50 mg). ESI MS = m/z 411 [M+H]⁺

15 **Example 2**

Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-5-fluoro-1H-indazole, the following compound was prepared.

4-(1-(2,6-dichlorobenzoyl)-5-fluoro-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 429 [M+H]⁺




20

Example 3

25 Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-7-fluoro-1H-indazole, the following compound was prepared.

4-(1-(2,6-dichlorobenzoyl)-7-fluoro-1H-indazol-3-yl)benzoic acid




ESI MS = m/z 429 [M+H]⁺

Example 4

- 5 Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-4-fluoro-1*H*-indazole, the following compound was prepared.

4-(1-(2,6-dichlorobenzoyl)-4-fluoro-1*H*-indazol-3-yl)benzoic acid



ESI MS = m/z 429 [M+H]⁺

10

Example 5

Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-6-fluoro-1*H*-indazole, the following compound was prepared.

4-(1-(2,6-dichlorobenzoyl)-6-fluoro-1*H*-indazol-3-yl)benzoic acid


15 ESI MS = m/z 429 [M+H]⁺

Example 6

Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-1*H*-pyrazolo[4,3-*b*]pyridine, the following compounds were prepared.


20 **6A:** 4-(1-(2,6-dichlorobenzoyl)-1*H*-pyrazolo[4,3-*b*]pyridin-3-yl)benzoic acid

ESI MS = m/z 412 [M+H]⁺



6B: 4-(1-(2-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid

ESI MS = m/z 412 [M+H]⁺




5

Example 7

Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-1H-pyrrolo[3,2-b]pyridine, the following compound was prepared.

4-(1-(2,6-dichlorobenzoyl)-1H-pyrrolo[3,2-b]pyridin-3-yl)benzoic acid



10 ESI MS = m/z 412 [M+H]⁺




Example 8

Following a procedure analogous to that described in **Example 1**, starting from 3-bromo-1H-pyrazolo[4,3-c]pyridine, the following compound was prepared.

15

4-(1-(2,6-dichlorobenzoyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acidESI MS = m/z 412 [M+H]⁺5 **Example 9**1,1,1,3,3,3-hexafluoro-2-(4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)phenyl)propan-2-ol

- i) To a suspension of 3-bromoindazole (500 mg, 2.54 mmol) in 8 ml CH₂Cl₂ was added triethyl amine (1.06 ml, 7.61 mmol) at room temperature. To this yellow solution was added 2-(trifluoromethyl)benzenesulfonyl chloride (0.392 ml, 2.54 mmol) and the reaction mixture was stirred for 3 h at room temperature. The reaction mixture was concentrated under reduced pressure and the residue was purified on SiO₂ using 5% to 20% ethyl acetate in heptane to give 3-bromo-1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazole (894 mg)
- ii) Following a procedure analogous to that described in **Example 1, step ii**, using 1,1,1,3,3,3-hexafluoro-2-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propan-2-ol as the boronic ester, the title compound 1,1,1,3,3,3-hexafluoro-2-(4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)phenyl)propan-2-ol was prepared (9 mg).
- ESI MS = m/z 569 [M+H]⁺

Example 104-(1-(2-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid

- 5 **i)** To a mixture of *tert*-butyl 3-bromo-1H-indazole-1-carboxylate (5 g, 16.83 mmol,) and 4-(*tert*-butoxycarbonyl)phenylboronic acid (4.52 g, 20.36 mmol) in 30 ml of dioxane and 30 ml water was added sodium carbonate (50.5 mmol, 5.35 g). The mixture was purged with N₂ and subsequently, Pd(PPh₃)₄ (0.486 g, 0.421 mmol) was added. The reaction mixture was heated to 100°C overnight under a nitrogen atmosphere.
- 10 After cooling to room temperature, the reaction mixture was diluted with water and the product was extracted into ethyl acetate. The combined organic layers were washed with water, brine and dried over magnesium sulfate. After filtration, the solvent was removed under reduced pressure and the residue was purified on SiO₂, using 0% to 25 % ethylacetate in heptane as the eluent, to give *tert*-butyl 4-(1H-indazol-3-yl)benzoate as a yellow solid.
- 15 **ii)** To a solution of *tert*-butyl 4-(1H-indazol-3-yl)benzoate (40 mg, 0.136 mmol) in 1 ml of pyridine was added 2-(trifluoromethyl)benzoyl chloride (356 mg, 1.699 mmol). The reaction mixture was stirred at 150 °C in a microwave reactor for 1h. After cooling, water was added and the product was extracted into ethylacetate. The organic layer was washed with water, brine and dried over magnesium sulfate. After filtration, the
- 20 solvent was removed under reduced pressure and the residue was purified on SiO₂, using 3% to 20% ethylacetate in heptane as the eluent, to give *tert*-butyl 4-(1-(2-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoate as a white solid.
- iii)** A solution of the compound obtained in the previous step (19.4 mg, 0.042 mmol) and trifluoroacetic acid (0.5 ml, 6.73 mmol,) in 2 ml CH₂Cl₂ was stirred for 1h at room
- 25 temperature.

After the reaction was complete the reaction mixture was concentrated under reduced pressure to give the title compound 4-(1-(2-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoic acid as a light yellow solid.

ESI MS = m/z 411 $[M+H]^+$


5

Example 11

Following a procedure analogous to that described in **Example 10**, the following compounds were prepared.


11A: 4-(1-(3,5-dichloroisonicotinoyl)-1*H*-indazol-3-yl)benzoic acid

10 ESI MS = m/z 412 $[M+H]^+$



11B: 4-(1-(2-bromobenzoyl)-1*H*-indazol-3-yl)benzoic acid




ESI MS = m/z 421/423 $[M+H]^+$







15


11C: 4-(1-(2-(methoxycarbonyl)benzoyl)-1*H*-indazol-3-yl)benzoic acid

ESI MS = m/z 401 $[M+H]^+$

**11D:** 4-(1-(cyclohexanecarbonyl)-1*H*-indazol-3-yl)benzoic acidESI MS = m/z 349 $[M+H]^+$ 5 **11E:** 4-(1-(2-bromo-6-chlorobenzoyl)-1*H*-indazol-3-yl)benzoic acidESI MS = m/z 457 $[M+H]^+$ 10 **11F:** 4-(1-(2-fluoro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoic acidESI MS = m/z 429 $[M+H]^+$


**11G:** 4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acidESI MS = m/z 391 [M+H]⁺5 **11H:** 4-(1-(2,4,6-trichlorobenzoyl)-1H-indazol-3-yl)benzoic acidESI MS = m/z 445 / 447 [M+H]⁺**11I:** 4-(1-(2,6-dimethylbenzoyl)-1H-indazol-3-yl)benzoic acidESI MS = m/z 371 [M+H]⁺10 **11J:** 4-(1-(2-ethoxybenzoyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 387 [M+H]⁺



11K: 4-(1-(2,3-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid


ESI MS = m/z 411 [M+H]⁺



5


11L: 4-(1-(2-chlorobenzoyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 377 [M+H]⁺




11M: 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid

10 ESI MS = m/z 445 [M+H]⁺



11N: 4-(1-(2-methyl-6-nitrobenzoyl)-1H-indazol-3-yl)benzoic acid


ESI MS = m/z 402 [M+H]⁺



5

Example 12

4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)benzoic acid



- i)** To a solution of *tert*-butyl 4-(1H-indazol-3-yl)benzoate, **Example 10, step i**, (58 mg, 0.197 mmol) in 2 ml CH₂Cl₂ were added triethyl amine (60 mg, 0.592 mmol) and 2-(trifluoromethyl)benzenesulfonyl chloride (48 mg, 0.197 mmol). The reaction mixture was stirred overnight at room temperature. The reaction mixture was directly purified on SiO₂ using 0% to 50% ethyl acetate in heptane as the eluent to give *tert*-butyl 4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)benzoate (102 mg) as a clear oil.

ii) Following a procedure analogous to that described in **Example 10, step iii**, the title compound, 4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1*H*-indazol-3-yl)benzoic acid, was prepared (28 mg)

ESI MS = m/z 447 $[M+H]^+$


5

Example 13

Following a procedure analogous to that described in **Example 11**, the following compounds were prepared.


13A: 4-(1-(quinolin-8-ylsulfonyl)-1*H*-indazol-3-yl)benzoic acid

10 ESI MS = m/z 429 $[M+H]^+$



13B: 4-(1-(2-bromophenylsulfonyl)-1*H*-indazol-3-yl)benzoic acid


ESI MS = m/z 457/459 $[M+H]^+$



15


13C: 4-(1-(2,5-dichlorophenylsulfonyl)-1*H*-indazol-3-yl)benzoic acid

ESI MS = m/z 447 $[M+H]^+$



13D: 4-(1-(3-chloro-2-methylphenylsulfonyl)-1H-indazol-3-yl)benzoic acid


ESI MS = m/z 427 [M+H]⁺



5


13E: 4-(1-(2,3-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 447 [M+H]⁺




13F: 4-(1-(2,3,4-trichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 481/483 [M+H]⁺



Example 14



5 4-(1-(2-cyanophenylsulfonyl)-1H-indazol-3-yl)benzoic acid

i) To a solution of 3-bromo-1H-indazole (200 mg, 1.02 mmol) in 2 ml CH₂Cl₂ was added triethylamine (308 mg, 3.05 mmol) and 2-cyanobenzene-1-sulfonyl chloride (205 mg, 1.02 mmol). The reaction mixture was stirred overnight at room temperature. After completion the reaction mixture was concentrated under reduced pressure and the product was purified on
10 SiO₂ using 0% to 50% ethylacetate in heptane as the eluent to give 4-(1-(2-
cyanophenylsulfonyl)-1H-indazol-3-yl)benzoic acid (360 mg) as a yellow solid.

ii) To a microwave reaction vial were added the product obtained in the previous step (50 mg, 0.138 mmol) in 2 ml toluene and 0.5 ml ethanol, 4-(*tert*-butoxycarbonyl)phenylboronic acid (50 mg, 0.166 mmol) and a 2M aqueous solution of potassium carbonate (0.345 ml, 0.690
15 mmol). After purging the vial with nitrogen for about 5 minutes, Pd(PPh₃)₄ (3.99 mg) was added and the reaction mixture was stirred for 30 minutes at 100 °C in a microwave reactor. After cooling to room temperature, the reaction mixture was diluted with water and product was extracted into ethyl acetate. The organic layer was washed with water, brine and dried over magnesium sulfate.

After filtration the solvent was evaporated under reduced pressure and the product was purified on SiO₂ using 0% to 50% ethylacetate in heptane as the eluent to give the desired product, tert-butyl 4-(1-(2-cyanophenylsulfonyl)-1H-indazol-3-yl)benzoate (12 mg).

iii) Following a procedure analogous to that described in **Example 10, step iii**, the title compound, 4-(1-(2-cyanophenylsulfonyl)-1H-indazol-3-yl)benzoic acid, was prepared (13 mg)


ESI MS = m/z 404 [M+H]⁺

Example 15

Following a procedure analogous to that described in **Example 14**, the following compounds were prepared.


15A: 4-(1-(2-(trifluoromethoxy)phenylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 463 [M+H]⁺




15B: 4-(1-(naphthalen-1-ylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 429 [M+H]⁺




15C: 4-(1-(2,6-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 447 [M+H]⁺




15D: 4-(1-(benzo[c][1,2,5]oxadiazol-4-ylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 421 [M+H]⁺




5 **15E:** 4-(1-(2-cyano-5-methylphenylsulfonyl)-1H-indazol-3-yl)benzoic acid

ESI MS = m/z 418 [M+H]⁺



10 **Example 16**



4-(1-(2-chlorophenylsulfonyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acid

i) To a solution of 3-bromo-1H-pyrazolo[4,3-c]pyridine (198 mg, 1.0 mmol) in 10 ml THF was added portion wise at 0 °C sodium hydride (101 mg, 4 mmol, 95%). After addition was complete, 2-chlorobenzenesulfonyl chloride (495 mg, 2.347 mmol) was added and the reaction mixture was stirred overnight at room temperature. The reaction mixture was quenched by the addition of water. The product was extracted into ethyl acetate and the organic layer was washed with brine. The organic solvent was evaporated under reduced pressure to give 3-bromo-1-(2-chlorophenylsulfonyl)-1H-pyrazolo[4,3-c]pyridine as a yellow oil. The product was used in the next step without further purification.


ii) To a microwave reaction vial were added the product obtained in the previous step (67 mg, 0.135 mmol) in 3 ml DME, *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (49.2 mg, 0.162 mmol) and 0.472 ml of a 2M aqueous solution of sodium carbonate (0.944 mmol). After purging the vial with nitrogen for about 5 minutes, Pd(PPh₃)₄ (31.2 mg, 0.027 mmol) was added and the reaction mixture was stirred for 40 minutes at 100 °C in a microwave reactor. After cooling to room temperature, the reaction mixture was diluted with water and washed with ethyl acetate. The organic layer was washed with water, brine and dried over magnesium sulfate.

After filtration the solvent was evaporated under reduced pressure and the product was purified preparative HPLC using 20% to 100% CH₃CN in water as the eluent to give *tert*-butyl 4-(1-(2-chlorophenylsulfonyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoate as a yellow solid (16 mg).

iii) Following a procedure analogous to that described in **Example 10, step iii**, the title compound, 4-(1-(2-chlorophenylsulfonyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acid, was prepared (7 mg)

ESI MS = m/z not measured $[M+H]^+$

Example 17



5

4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-3-fluorobenzoic acid

- i) 3-bromo-1H-indazole (3 gram; 15 mmol) was dissolved in 30 ml DMF, followed by the addition of sodiumhydride (850 mg, 21 mmol). After stirring for 20 min at room temperature, 3,5-dimethylbenzene-1-sulfonyl chloride (4.4 gram, 21 mmol) was added and the reaction mixture was stirred for another 2 h at room temperature. The reaction was quenched by the addition of water and the formed solids were filtered off and washed with little methanol. The solids were dried to obtain 3-bromo-1-(3,5-dimethylphenylsulfonyl)-1H-indazole (2.1 gram) as a white solid.
- ii) The product obtained in the previous step (100 mg, 0.27 mmol), $Pd(PPh_3)_4$ (63 mg, 0.05 mmol) and 4-borono-3-fluorobenzoic acid (101 mg, 0.55 mmol) were stirred in 6.5 ml DME and 3.5 ml of an aqueous solution of $NaHCO_3$ (69 mg, 0.82 mmol) was added. The reaction mixture was heated in a microwave reactor for 30 min at $100^\circ C$. After cooling, the reaction mixture was filtered over cotton and the volatiles were evaporated under reduced pressure. The crude material was purified by prep. HPLC (Gemini NX column (100 * 30 mm, 5 μ m); 40 ml/min; Eluens A: 20 mmol/L NH_4HCO_3 (aq) and Eluens B: acetonitrile; Gradient: 0-7 min 70%A and 30% B; 7-7.1 min 50%A and 50% B; 7.1-8.5 min 100% B) to give the title compound 4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-3-fluorobenzoic acid (25 mg) as an off white solid.


ESI MS m/z 425 $[M+H]^+$

Example 18

Following a procedure analogous to that described in **Example 17**, the following compounds was prepared.


5 4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-2-(trifluoromethyl)benzoic acid

ESI MS m/z 475 [M+H]⁺



10 **Example 19**

5-(4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1H-tetrazole



- i) 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)benzonitrile (8.0 g, 34.8 mmol), trimethylsilyl azide, (8.04 g, 69.8 mmol) and dibutyl-stannanone (868 mg, 3.5 mmol, 0.1 equiv) were dissolved in 1,2-dimethoxy ethane (110 ml). The mixture was divided in two portions and transferred in two microwave vessels then irradiated in a microwave reactor simultaneously at 150 °C for 10 min. After cooling to room temperature, trimethylsilyl azide (4.02 g, 35 mmol) and dibutyl-stannanone (434 mg, 1.75 mmol) were added into each microwave vessel and the mixtures were irradiated again at 150 °C for 10 min. After cooling
- 15

to room temperature, the mixtures were combined and purified by column chromatography on florisil (200 g). It was first eluted with 20% dichloromethane in hexane to remove the impurities, then with 10% methanol in dichloromethane to elute the product. 1-(4-

Methoxybenzyl)-5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1*H*-tetrazole 2

was isolated as a white powder (8.01 g).

APCI MS m/z 273 $[M + H]^+$.

ii) The product obtained in the previous step (8.40 g, 30.9 mmol) was dissolved in acetonitrile (150 ml). First potassium carbonate powder (5.12 g, 37.0 mmol, 1.2 equiv), then 4-

methoxybenzyl chloride (6.3 mL, 46.3 mmol) were added. After stirring at 40 °C for 12 hours

the reaction mixture was diluted with ethyl-acetate (100 ml), washed with water (2×100 ml) and dried over magnesium sulfate. After filtration the solvent was evaporated under reduced


pressure. The residue was suspended in hexane and the formed crystals were filtered. Title

compound 1-(4-Methoxybenzyl)-5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1*H*-tetrazole was obtained as a white powder as a mixture of regio-isomers (9.67 g).

APCI MS m/z 393 $[M + H]^+$.

Example 20

3-(4-(1*H*-tetrazol-5-yl)phenyl)-1-(3-chlorophenylsulfonyl)-1*H*-indazole



i) *Tert*-butyl 3-bromo-1*H*-indazole-1-carboxylate (5.0 g, 16.8 mmol) and a mixture of the regio-isomers of 1-(4-methoxybenzyl)-5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1*H*-tetrazole (**Example 19, step ii**) (7.92 g, 20.2 mmol) were dissolved in 100 ml of a 1 to 1 mixture of 1,4-dioxane and water under a nitrogen atmosphere. Sodium carbonate (5.35 g, 50.5 mmol) was added and the system was purged with nitrogen.. Then $\text{Pd}(\text{PPh}_3)_4$

(486 mg, 0.42 mmol) was added and the reaction mixture was heated overnight at reflux temperature under a nitrogen atmosphere. After cooling to room temperature the mixture was diluted with water and extracted with ethyl acetate. The combined organic phases were washed with brine, dried over magnesium sulfate. After filtration the solvent was removed under reduced pressure. The residue was purified on SiO₂ using 20 to 30% ethyl acetate in hexane as the eluent to give 3-(4-(1-(4-methoxybenzyl)-1*H*-tetrazol-5-yl)phenyl)-1*H*-indazole as a mixture of two regioisomers in a 2 to 1 ratio (2.8 g) as a white solid.

Major regio-isomer (1.91 g, 30%) as a white powder.

APCI MS m/z 383 [M + H]⁺,

ii) To a solution of the product obtained in the previous step (100 mg, 0.26 mmol) in 2ml anhydrous THF was added *t*-BuOK (32 mg, 0.29 mmol). After stirring the reaction mixture at room temperature for 20 min, 3-Chlorobenzene-1-sulfonyl chloride (41 mg, 0.29 mmol) was added and the mixture was stirred at room temperature for an additional 2 h. After completion, the solvent was evaporated under reduced pressure, the residue was dissolved in dichloromethane and washed with water. The organic phase was evaporated and the residue was dried under reduced pressure to afford 1-(3-chlorophenylsulfonyl)-3-(4-(1-(4-methoxybenzyl)-1*H*-tetrazol-5-yl)phenyl)-1*H*-indazole (146 mg, crude) as a brown oil which was used in the next step without any further purification.

ESI MS m/z 557 [M + H]⁺

iii) The product obtained in the previous step (81 mg) was dissolved in 1 ml trifluoroacetic acid and stirred for 2 days at room temperature. The solvent was evaporated under reduced pressure and the residue was purified by preparative HPLC using eluent system 1 then eluent system 2 to afford 3-(4-(1*H*-tetrazol-5-yl)phenyl)-1-(3-chlorophenylsulfonyl)-1*H*-indazole (26 mg) as a white powder:


ESI MS m/z 437 [M + H]⁺

Example 21


Following a procedure analogous to that described in **Example 20**, the following compounds were prepared.

21A: methyl 3-(3-(4-(1*H*-tetrazol-5-yl)phenyl)-1*H*-indazol-1-ylsulfonyl)thiophene-2-carboxylate

ESI MS = m/z 467 [M+H]⁺




21B: 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(2-bromo-4-fluorophenylsulfonyl)-1H-indazole




ESI MS = m/z 499/501[M+H]⁺

5 **21C:** 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(3-bromo-5-chlorothiophen-2-ylsulfonyl)-1H-indazole



ESI MS = m/z 523 [M+H]⁺


21D: 4-(3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-ylsulfonyl)-3,5-dimethylisoxazole



ESI MS = m/z 422 $[M+H]^+$

Example 22

(3-(4-(1*H*-tetrazol-5-yl)phenyl)-1*H*-indazol-1-yl)(2,6-dichlorophenyl)methanone



5

i) To a solution of a mixture of regio-isomers of 3-(4-(1-(4-Methoxybenzyl)-1*H*-tetrazol-5-yl)phenyl)-1*H*-indazole (**Example 20, step i**) (50 mg, 0.13 mmol) was dissolved in anhydrous THF (1 ml). After addition of *t*-BuOK (16 mg, 0.14 mmol,) the reaction mixture was stirred at room temperature for 15 min. Subsequently, 2,6-dichlorobenzoyl chloride (21 μ l, 0.14 mmol) was added and the reaction mixture was stirred for another 2h at room temperature. The solvent was evaporated under reduced pressure, the residue was dissolved in dichloromethane and washed with water, an aqueous solution of 5% K_2CO_3 and water. The organic solvent was evaporated under reduced pressure and the residue was dried in vacuo. The product was purified by preparative HPLC to afford (2,6-dichlorophenyl)(3-(4-(1-(4-methoxybenzyl)-1*H*-tetrazol-5-yl)phenyl)-1*H*-indazol-1-yl)methanone as a mixture of 2 regioisomers in a 98 to 2 ratio (59 mg, 74%).

ESI MS m/z 555 $[M + H]^+$.

ii) Following a procedure analogous to that described in **Example 20, step iii**, the product obtained in the previous step was converted to the title compound (3-(4-(1*H*-tetrazol-5-yl)phenyl)-1*H*-indazol-1-yl)(2,6-dichlorophenyl)methanone (16 mg).

20


ESI MS m/z 435 $[M + H]^+$.

Example 23

Following a procedure analogous to that described in **Example 22**, the following compound

5 was prepared.

Org 356723-0



(3-(4-(*1H*-tetrazol-5-yl)phenyl)-*1H*-indazol-1-yl)(2,3-dichlorophenyl)methanone

ESI MS = m/z $[M+H]^+$

10


15

20

25

Example 24A: Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl) benzoyl)-4-fluoro-1*H*-indazol-3-yl)-3-fluorobenzoate (24A)

SCHEME A.



5

i) Preparation of 4-fluoro-3-iodo-1*H*-indazole (A-2). To a solution of 4-fluoroindazole **A-1** (5.00 g, 36.7 mmol) in DMF (80 mL) was added I_2 (18.6 g, 73.5 mmol) and KOH (7.73 g, 134 mmol) successively at rt. After 2 h, the reaction mixture was poured into aq. 10% NaHSO₃ (200 mL) and extracted with EtOAc (200 mL*3). The combined organic layers were washed with H₂O and brine, dried over Na₂SO₄, and concentrated. The crude solid was washed with PE to give the title compound as a yellow solid. LCMS (ESI) calc'd for C₇H₅FIN₂ [M+H]⁺: 262.9, found: 262.9.

10

ii) Preparation of (2-chloro-6-(trifluoromethyl)phenyl)(4-fluoro-3-iodo-1*H*-indazol-1-yl)methanone (A-4). To a flask was added compound **A-2** (5.24 g, 20 mmol), compound **A-3** (4.86 g, 20 mmol), DMAP (2.44 g, 20 mmol) and DCM (30 mL), followed by the addition of TEA (5.8 mL, 40 mmol) slowly. The reaction mixture was stirred at rt for 24 h. The mixture was diluted with H₂O, and the organic layer was separated. The aqueous layer was extracted with CH₂Cl₂. The combined organics were washed with H₂O, brine, dried over Na₂SO₄, and

15


concentrated. The residue was purified by flash chromatography (Pentane/EtOAc) to afford the title compound. LCMS (ESI) calc'd for $C_{15}H_7ClF_4IN_2O$ $[M+H]^+$: 468.9, found: 468.9.

iii) **Preparation of methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoate (A-5).** To a mixture of **A-4** (300 mg, 0.64 mmol), 2-fluoro-4-(methoxycarbonyl)phenylboronic acid (190 mg, 0.96 mmol), Pd(dppf)Cl₂ (52 mg, 0.064 mmol) and KOAc (190 mg, 1.92 mmol) was added dioxane (10 ml) and H₂O (2 ml), and the mixture was heated at 90 °C under microwave for 2 h. The mixture was cooled down, diluted with CH₂Cl₂ (50 ml). The organic layer was separated, washed with brine, dried over Na₂SO₄, concentrated. The residue was purified by flash chromatography (Pentane/EtOAc) to give the title compound. LCMS (ESI) calc'd for $C_{23}H_{13}ClF_5N_2O_3$ $[M+H]^+$, 495, found: 495.

iv) **Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoic acid (A-6).** To a solution of **A-5** (180 mg, 0.36 mmol) in THF (5mL) and H₂O (5mL) was added LiOH (350 mg, 1.44 mmol), and the mixture was stirred at rt for 24 h. The mixture was neutralized with 2N HCl to PH = 3~4. The mixture was then concentrated under reduced pressure. The residue was filtered and washed with H₂O to afford the title compound as white solid. LCMS (ESI): calc'd for $C_{22}H_{11}ClF_5N_2O_3$ $[M+H]^+$: 481, found: 481; ¹H NMR (400 MHz, CDCl₃) δ 8.45-8.47 (1H, d), 7.88-7.95 (2H, m), 7.66-7.71 (3H, m), 7.58-7.64 (2H, m), 7.13-7.17 (1H, m).

v) **Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoate (24A).** To a suspension of compound **A-6** (160 mg, 0.33 mmol) in H₂O (10 mL) was added 0.1M NaOH (3.3 ml, 0.33 mmol) and the mixture was kept stirring at 0 °C for 30 mins. The reaction mixture was dried under lyophilization to afford the title compound. LCMS (ESI) calc'd for $C_{22}H_{11}ClF_5N_2O_3$ $[M-Na+2H]^+$: 481, found: 481; ¹HNMR (400 MHz, DMSO) δ 8.36-8.38 (1H, d), 7.96-8.03 (2H, m), 7.66-7.71 (1H, m), 7.85-7.89 (1H, m), 7.72-7.74 (1H, m), 7.65(1H, s), 7.40-7.45(2H, m).

Example 24B: Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoate (24B)



5

i) Preparation of (3-bromo-1*H*-indazol-1-yl)(2-chloro-6-(trifluoromethyl)phenyl)methanone (B-2). To a solution of 3-bromo-1*H*-indazole (2 g, 0.75 mmol), DMAP (1 g, 8.5 mmol) and Et₃N (1.85 ml, 12.7 mmol) in CH₂Cl₂ (20 mL) at 0°C was added a solution of compound A-3 (4 g, 20.4 mmol) in CH₂Cl₂ (5 mL). The mixture was stirred at rt for 24 h, diluted with CH₂Cl₂, washed with 1M HCl. The organic layer was washed with sat. NaHCO₃ and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by flash chromatography (PE/EA=10/1) to give the title compound. LCMS (ESI) calc'd for C₁₅H₈BrClF₃N₂O [M+H]: 404.9, Found: 404.9.

10

15

ii) Preparation of methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoate (B-3). A mixture of compound B-2 (3.0 g, 7.4 mmol), 4-(methoxycarbonyl)phenylboronic acid (1.6 g, 8.9 mmol), Pd(PPh₃)₄ (0.43g, 0.37 mmol) and K₂CO₃ (2.04 g, 10.8 mmol) in dioxane (10 mL) and H₂O (2 mL) was heated at 140°C for 2 h under microwave. The mixture was cooled down, diluted with CH₂Cl₂ and H₂O. The organic

layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (PE/EA = 10/1) to give the title compound as a yellow solid. LCMS (ESI) calc'd for C₂₃H₁₅ClF₃N₂O₃ [M+H]⁺: 459, found: 459.


5 **iii) Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoic acid (B-4).** A mixture of compound **B-3** (1.3 g, 2.8 mmol) and LiOH (0.35 g, 8.4 mmol) in THF (5 mL) and H₂O (5 mL) was stirred at rt for 24 h. Then the mixture was neutralized with AcOH to PH < 7, diluted with CH₂Cl₂ and H₂O. The organic layer was separated and washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by prep HPLC to
10 give the title compound as a white solid. LCMS (ESI) calc'd for C₂₂H₁₃ClF₃N₂O₃ [M+H]⁺: 445, found: 445; ¹H NMR (400 MHz, CDCl₃) δ 8.68-8.71 (1H, d), 8.19-8.21 (2H, m), 8.03-8.05 (1H, d), 7.94-7.96 (2H, m), 7.70-7.75 (3H, m), 7.26-7.63 (2H, m).

iv) Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoate (24B). To a suspension of compound **B-4** (0.87g, 2.0 mmol) in H₂O (10 mL) at 0
15 °C was added 0.1 M NaOH (2.0 ml, 2.0 mmol) and the mixture was kept stirring at at 0°C for 30 mins. The mixture was dried under lyophilization to give the title compound. ¹H NMR (400 MHz, DMSO) δ 8.54-8.56 (1H, d), 8.22-8.24 (1H, d), 8.03-8.05 (1H, d), 7.98-8.00 (3H, m), 7.82-7.88 (2H, m), 7.71-7.73 (2H, m), 7.63-7.67 (1H, t).

20

Example 24C: Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-pyrazolo [4,3-*b*]pyridin-3-yl)-3-fluorobenzoate (24C)

SCHEME C.



i) Preparation of (3-bromo-1H-pyrazolo[4,3-b]pyridin-1-yl)(2-chloro-6-(trifluoromethyl)phenyl)methanone (C-2). To a mixture of compound **A-3** (1.2 g, 5.0 mmol), compound **C-1** (1.0 g, 5.0 mmol), and DMAP (0.61 g, 5.0 mmol) in CH₂Cl₂ (20 mL) was added TEA (0.55 g, 5.5 mmol) dropwise. The mixture was stirred at rt for 24 h., diluted with H₂O, and the organic layer was separated. The aqueous layer was extracted with CH₂Cl₂. The combined organics were washed with H₂O, brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (PE/EA from 50/1 to 10/1) to give the title solid. LCMS (ESI) calc'd for C₁₄H₇BrClF₃N₃O [M+H]⁺: 405.9, found: 405.9.


ii) Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-fluorobenzoic acid (C-3). A mixture of **C-2** (120 mg, 0.3 mmol), 2-fluoro-4-(methoxycarbonyl)phenylboronic acid (89 mg, 0.45 mmol), Pd(PPh₃)₄ (18 mg, 0.015 mmol) and K₂CO₃ (128 mg, 0.75 mmol) in dioxane (3 mL) and H₂O (0.6 mL) was heated at 140°C for 3 h under microwave. The mixture was cooled down and diluted with CH₂Cl₂ and H₂O. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (PE/EA=1/1) to give the title compound as a yellow solid. LCMS (ESI) calc'd for C₂₂H₁₃ClF₄N₃O₃ [M+H]⁺: 464, found: 464; ¹H NMR (400 MHz,

DMSO) δ 13.51 (1H, s), 8.98-8.91 (2H, m), 8.42-8.38 (2H, t), 8.06-7.95 (3H, m), 7.90-7.87 (2H, m), 7.87-7.80 (1H, m).













iii) Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-fluorobenzoate (24C). To a suspension of compound C-3 (135 mg, 0.29 mmol) in H₂O (20 mL) was added 0.1 M NaOH (2.9 ml, 0.29 mmol) at 0 °C, and the mixture was kept stirring for 30 min at 0 °C. The mixture was then dried under lyophilization to give the title compound. LCMS (ESI) calc'd for C₂₁H₁₁ClF₄N₃O₃ [M-Na+2H]⁺: 464, found: 464, ¹H NMR (400 MHz, CD₃OD) δ 8.95-8.93 (1H, m), 8.89-8.87 (1H, m), 8.13-8.09 (1H, t), 7.90-7.87 (3H, m), 7.81-7.75 (3H, m).
















The following examples shown in TABLE 2 were prepared following similar procedures described for Examples #24A, B, C in Schemes A-C, which can be achieved by those of ordinary skill in the art.
















TABLE 2

































Ex.	IUPAC name	A ring	P	Q	NMR or LCMS [M+H] ⁺ (found)













24D	(2-chloro-6-(trifluoromethyl)phenyl)(4-fluoro-3-(4-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)-1H-indazol-1-yl)methanone				585
24E	3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid				463
24F	2-(3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)phenyl)acetic acid				477
24G	2-(4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)phenyl)acetic acid				477













24H	2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid				497
24I	Sodium 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoate				497
24J	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzamide				462
24K	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluoro-5-methylbenzoic acid				495
24L	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-methylbenzoic acid				477
















24M	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-methoxybenzoic acid				494
24N	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluorobenzoic acid				481
24O	sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluorobenzoate				481 [M-Na+2H] ⁺
24P	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-methylbenzoic acid				483
24Q	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-methoxybenzoic acid				493
















	acid				
24R	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,6-difluorobenzoic acid				499
24S	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-isopropylbenzoic acid				505
24T	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3,5-dimethoxybenzoic acid				523
24U	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-cyano benzoic acid				488
24V	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,3-difluorobenzoic acid				499
















24W	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,5-difluorobenzoic acid				499
24X	2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluorobenzoic acid				515
24Y	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluoro-2-methoxybenzoic acid				511
24Z	5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoic acid				479
24AA	3-fluoro-4-(4-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)-				427













	yl)benzoic acid				
24AB	4-(4-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid				409
24AC	3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid				446
24AD	2-(4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)phenyl)acetic acid				460
24AE	(2-chloro-6-(trifluoromethyl)phenyl)(3-(4-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)-1H-pyrazolo[4,3-b]pyridin-1-yl)methanone				568
















24AF	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-methylbenzoic acid				460
24AG	3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid				480
24AH	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-fluorobenzoic acid				464
24AI	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-methoxybenzoic acid				476

24AJ	2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-5-fluorobenzoic acid				498
24AK	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2,5-difluorobenzoic acid				482
24AL	5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-hydroxybenzoic acid				462
24AM	4-(1-(2-fluoro-6-methoxybenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid				392
24AN	3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid				410

24AO	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-(trifluoromethyl)-1H-indazol-3-yl)benzoic acid				513
24AP	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-isopropylbenzoic acid				487
24AQ	sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-fluorobenzoate				463 [M-Na+2H] ⁺
24AR	sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-3-fluorobenzoate				463 [M-Na+2H] ⁺
24AS	3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-				480


	3-yl)benzoic acid				
24AT	3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid				409
24AU	sodium 3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoate				409 [M-Na+2H] ⁺
24AV	4-(7-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid				409
24AW	3-fluoro-4-(7-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid				427
24AX	4-(6-bromo-1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid				543

24AY	4-(6-bromo-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid				¹ H NMR (400 MHz, DMSO) δ 13.25 (1H, s), 8.72 (1H, s), 8.23 (1H, d, <i>J</i> =8.4 Hz), 8.01-8.11 (4H, m), 7.86-7.96 (4H, m)
24AZ	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-methoxy-1H-indazol-3-yl)benzoic acid				493
24BA	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl)benzoic acid				461
24BB	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl)-3-fluorobenzoic acid				479

24BC	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-hydroxy-1H-indazol-3-yl)benzoic acid				479
24BD	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-hydroxy-1H-indazol-3-yl)-3-fluorobenzoic acid				497
24BE	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(hydroxymethyl)-1H-indazol-3-yl)benzoic acid				475
24BF	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(hydroxymethyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				493
24BG	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(methoxymethyl)-1H-indazol-3-yl)benzoic acid				489

Example 25A: Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxy ethylcarbamoyl)-1*H*-indazol-3-yl)benzoate (25A)

Scheme D.



i) Preparation of methyl 1*H*-indazole-6-carboxylate (D-2).

To a solution of methyl 3-amino-4-methylbenzoate **D-1** (5.0 g, 30 mmol) in AcOH (140 mL) at 0 °C was added a solution of sodium nitrite (2.1 g, 30 mmol) in H₂O (3.5 mL). After addition, the mixture was stirred at rt for 24 h. Then half of the solvent was removed under reduced pressure and the resulting mixture was diluted with H₂O, and extracted with EtOAc. The combined organics were washed H₂O and brine, and concentrated to give the title compound. LCMS (ESI) calc'd for C₉H₉N₂O₂ [M+H]⁺: 177, found 177.

ii) Preparation of methyl 3-iodo-1*H*-indazole-6-carboxylate (D-3).

To a solution of methyl *1H*-indazole-6-carboxylate **D-2** (5.0 g, 28 mmol) in DMAC (50 mL) at 0 °C was added iodine (14.4 g, 56.7 mmol) and potassium hydroxide (6.3 g, 114 mmol) portionwise. After addition, the mixture was kept stirring for 1 h, then quenched with aq. Na₂S₂O₃ and extracted with EtOAc. The combined organics were concentrated and triturated
5 with hexane. The precipitate was collected by filtration to afford the title compound as a brown solid. LCMS (ESI) calc'd for C₉H₈IN₂O₂ [M+H]⁺: 302.9, found: 302.9.

iii): Preparation of methyl 1-(2-chloro-6-(trifluoromethyl)benzoyl)-3-iodo- *1H*-indazole-6-carboxylate (D-4).

10 To a solution of methyl 3-iodo-*1H*-indazole-6-carboxylate **D-3** (11.7 g, 38.7 mmol), DMAP (4.72 g, 38.7 mmol) in CH₂Cl₂ (30 mL) was added Et₃N (11.2 mL, 77 mmol) dropwise. The reaction mixture was stirred at rt for 24 h, diluted with H₂O, and the organic layer was separated. The aqueous layer was extracted with CH₂Cl₂ twice. The combined organics were washed with H₂O, brine, dried over Na₂SO₄, concentrated. The residue was purified by
15 column chromatography (PE /EtOAc from 50/1 to 10/1) to give the title compound. LCMS (ESI) calc'd for C₁₇H₁₀ClF₃IN₂O₃ [M+H]⁺: 508.9, found: 508.9.

iv): Preparation of 1-(2-chloro-6-(trifluoromethyl)benzoyl)-3-iodo-*1H*- indazole -6-carboxylic acid (D-5).

20 To a solution of **D-4** (16.5 g, 32.5 mmol) in THF (10 mL) and H₂O (50 mL) was added LiOH (3.40 g, 162 mmol), and the mixture was stirred at rt for 24 h. The solvent was removed under reduced pressure and the residue was taken up in H₂O, and neutralized with 5% aqueous HCl to PH = 4~5. The white precipitate was collected by filtration, washed with H₂O and hexane to give the title compound as an off-white solid. LCMS (ESI) calc. C₁₆H₈ClF₃IN₂O₃ [M+H]⁺:
25 494.9, found: 494.9.

v) Preparation of 1-(2-chloro-6-(trifluoromethyl)benzoyl)-3-(4-(methoxycarbonyl)phenyl)-*1H*-indazole-6-carboxylic acid (D-6).

A mixture of **D-5** (300 mg, 0.61 mmol), 4-(methoxycarbonyl)phenylboronic acid (165 mg, 0.92 mmol), Pd(dppf)Cl₂ (50 mg, 0.061 mmol) and KOAc (181 mg, 1.83 mmol) in dioxane
30 (10 mL) and H₂O (2 mL) was heated at 95 °C for 2 h under microwave. The mixture was then cooled to rt, diluted with EtOAc and H₂O. The organic layer was washed with brine, dried

over Na₂SO₄ and concentrated. The residue was purified by flash chromatography (PE/EtOAc =20/1) to give the title compound as a white solid. LCMS (ESI) calc'd for C₂₄H₁₅ClF₃N₂O₅ [M+H]⁺: 503.1, found: 503.1.

5 **vi) Preparation of methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1*H*-indazol-3-yl)benzoate (D-7).**

To a solution of compound **D-6** (180 mg, 0.36mmol) in CH₂Cl₂ (15 mL) was added PYAOP (374 mg, 0.72mmol), followed by the addition of Et₃N (0.16mL, 1.1 mmol). The mixture was stirred at rt for 2 h, diluted with EtOAc. The organic layer was separated, washed with brine,
10 dried over Na₂SO₄, concentrated to give the title compound as a white solid. LCMS (ESI) calc'd for C₂₆H₂₀ClF₃N₃O₅ [M+H]⁺: 546.1, Found: 546.1.

vii) Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1*H*-indazol-3-yl)benzoic acid (D-8).


15 A mixture of **D-7** (191mg, 0.35 mmol) and LiOH (42mg, 1.75 mmol) in THF (10 mL) and H₂O (5 mL) was stirred at rt for 2 h. The solvent was removed under reduced pressure, and the residue was dissolved in H₂O and neutralized with aqueous 5% HCl to PH = 4~5. The precipitated was collected by filtration, washed with H₂O and hexane to the title compound as an off-white solid. LCMS (ESI) calc'd for C₂₅H₁₈ClF₃N₃O₅ [M+H]⁺: 532.1, found: 532.1; ¹H
20 NMR (400 MHz, MeOD) δ 9.109 (1H, s), 8.29-8.31(1H, d, J = 8.4 Hz), 8.17-8.19 (2H, d, J = 8.4 Hz), 8.07-8.10 (1H, m), 7.97-8.00 (2H, d, J = 8.8 Hz), 7.88-7.99 (2H, m), 7.78-7.82 (1H, m), 3.79-3.82 (2H, m), 3.60-3.63 (2H, m).

25 **viii) Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1*H*-indazol-3-yl)benzoate (25A)**










To a suspension of compound **D-8** (185 mg, 0.35 mmol) in H₂O (10 mL) was added 0.1M NaOH (3.5 ml, 0.35 mmol) at 0 °C, and the mixture was kept stirring for 30 mins. The mixture was then dried under lyophilizaiton to give the title compound. LCMS (ESI) calc. C₂₅H₁₈ClF₃N₃O₅ [M-Na+2H]⁺: 532, found: 532; ¹H NMR (400 MHz, MeOD) δ 9.09 (1H, s),
30 8.28-8.30 (1H, d, J = 8.8 Hz), 8.05-8.09 (3H, m), 7.86-7.91 (4H, m), 7.77-7.86 (1H, m), 3.79-3.82 (2H, m), 3.60-3.63 (2H, m).










The following examples shown in **TABLE 3** were prepared following similar procedures described for **Example #25A** in **Scheme D**, which can be achieved by those of ordinary skill in the art.













5 **TABLE 3**
























Ex.	IUPAC name		P	Q	LCMS [M+H] ⁺ (found)
25B	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)benzoic acid				516
25C	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethyl)carbamoyl)-1H-indazol-3-yl)benzoic acid				532










	hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoic acid				
25D	sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoate				554
25E	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				534
25F	sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoate				556

25G	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)-2-fluorobenzoic acid				550
25H	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				550
25I	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-2-fluorobenzoic acid				534

25J	3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)benzoic acid				550
25K	3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoic acid				566
25L	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-hydroxypropylcarbamoyl)-1H-indazol-3-yl)benzoic acid				546
25M	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-hydroxypropylcarbamoyl)-1H-indazol-3-yl)-2-fluorobenzoic acid				564


	indazol-3-yl)-3-fluorobenzoic acid				
25N	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-methoxypropyl carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				580
25O	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxypropyl carbamoyl)-1H-indazol-3-yl)benzoic acid				546
25P	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(4-hydroxybutyl carbamoyl)-1H-indazol-3-yl)benzoic acid				560
25Q	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-methyl-2-hydroxyethyl carbamoyl)-1H-indazol-3-yl)benzoic acid				546

	((2-hydroxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)benzoic acid				
25R	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-hydroxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				564
25S	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-methoxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				578
25T	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-fluorobenzoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid				546

	fluoroazetidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid				
25U	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3,3-difluoroazetidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid				564
25V	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(pyrrolidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid				542
25W	4-(6-(azetidine-1-carbonyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid				528

Example 26: Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro- *1H*-indazol-3-yl)-2-hydroxybenzoic acid (26A), 2-acetoxy-4-(1-(2-chloro-6-(tri fluoromethyl) benzoyl)-4-fluoro-*1H*-indazol-3-yl) benzoic acid (26B) and sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-*1H*-indazol-3-yl)-2-hydroxy benzoate (26C)

Scheme E.



i) Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-*1H*-indazol-3-yl)-2-hydroxybenzoic acid (26A)

To a solution of compound E-1 (prepared following a similar procedure in Scheme A) (100 mg, 0.19 mmol) in DCM (10 mL) was added BBr_3 (1M, 0.37 mL, 0.37 mmol) at rt, and the mixture was kept stirring for 48 h. Then MeOH (~10 mL) was added to quench excess BBr_3 and the mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC to give the title compound as a yellow solid. LCMS (ESI) calc'd for $\text{C}_{22}\text{H}_{12}\text{ClF}_4\text{N}_2\text{O}_4$ $[\text{M}+\text{H}]^+$: 479, found: 479; ^1H NMR (400 MHz, MeOD) δ 8.46 (1H, d, $J=8.4$ Hz), 7.95-7.76 (5H, m), 7.35-7.29 (3H, m).

ii) Preparation of 2-acetoxy-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid (26B)



A suspension of compound **26A** (48 mg, 0.1 mmol), Ac₂O (0.8 mL) and MgI₂ (28 mg, 0.1 mmol) in anhydrous ethyl ether (3 mL) was refluxed at 40 °C for 0.5 h. The reaction was cooled to rt, quenched with H₂O, extracted with ether, and concentrated. The crude product was dissolved in THF (10 mL) /H₂O (10 mL) and refluxed at 80 °C for 1 h. The mixture was cooled to rt, diluted with H₂O, and extracted with EtOAc. The combined organic extracts were concentrated and purified by flash chromatography PE/EA (5/1) to give the title compound as a white solid. LCMS (ESI): calc'd for C₂₂H₁₂ClF₄N₂O₄ [M+H]⁺: 521, found: 521, ¹H NMR (400 MHz, MeOD) δ 8.47 (1H, d, J=8.0 Hz), 8.11 (1H, d, J=8.0 Hz), 7.90-7.76 (5H, m), 7.53 (1H, t, J=1.6 Hz), 3.34 (1H, q, J=8.0 Hz), 2.31 (3H, s).

iii) Preparation of sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoate (26C)










To a suspension of compound **26A** (48 mg, 0.1 mmol) in H₂O (20 mL) was added 0.1 M NaOH (1 mL), and the mixture was sonicated for ~10 mins. The mixture was then dried under lyophilization to give the title compound. LCMS (ESI) calc'd for C₂₂H₁₁ClF₄N₂NaO₄ [M+H]⁺: 501, found: 501; ¹H NMR (400 MHz, MeOD) δ 8.45 (1H, d, J=8.0), 7.91-7.76 (5H, m), 7.31 (1H, q, J=8.0 Hz), 7.20-7.17 (2H, m).







The following examples shown in **TABLE 4** were prepared following similar procedures described for **Examples #26A, B, C** in **Scheme E**, which can be achieved by those of ordinary skill in the art.

TABLE 4




	IUPAC name	A ring	P	Q	LCMS [M+H] ⁺ (found)
26D	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluoro-6-hydroxybenzoic acid				497
26E	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-hydroxybenzoic acid				479

26F	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluoro-2-hydroxybenzoic acid				497
26G	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3,6-difluoro-2-hydroxybenzoic acid				515
26H	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-hydroxybenzoic acid				462

26I	2-acetoxy-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid				504
26J	4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-2-hydroxybenzoic acid				532

Example 27A: Preparation of 4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid (27A)

Scheme F.



i) Preparation of methyl 4-(6-carbamoyl-1-(2-chloro-6-(trifluoromethyl) benzoyl)-1H-indazol-3-yl)benzoate (F-2)

To a solution of compound **D-6** (180 mg, 0.36 mmol) in DCM (15 mL) was added PYAOP (374 mg, 0.72 mmol), followed by the addition of TEA (0.16 mL, 1.1 mmol). The mixture was stirred at rt for 2 h, diluted with EtOAc, washed with brine, dried over Na₂SO₄ and concentrated to give the title compound. LCMS (ESI) calc'd for C₂₄H₁₆ClF₃N₃O₄ [M+H]⁺: 502, found: 502.1.

ii) Preparation of (6-(aminomethyl)-3-(4-(hydroxymethyl)phenyl)-1H-indazol-1-yl)(2-chloro-6-(trifluoromethyl)phenyl)methanone (F-3)

To a solution of compound **F-2** (175 mg, 0.35 mmol) in THF (20 mL) was added BH₃.THF (1.75 mL, 1.75 mmol) and the mixture was refluxed for 12 h. Then MeOH was added to quench the excess BH₃, and the mixture was concentrated under reduced pressure to give the title compound as a white solid. LCMS (ESI): calc'd for C₂₃H₁₈ClF₃N₃O₂ [M+H]⁺: 460.1, found: 460.1.

iii) Preparation of tert-butyl (1-(2-chloro-6-(trifluoromethyl)benzoyl)-3-(4-(hydroxymethyl)phenyl)-1H-indazol-6-yl)methylcarbamate (F-4)

To a solution of compound **F-3** (138 mg, 0.30 mmol) in CH₂Cl₂ (20 mL) was added (Boc)₂O (138 mg, 0.30 mmol), followed by the addition of excess. The mixture was stirred at rt for 2 h. The mixture was diluted with H₂O, extracted with EtOAc. The combined organics were washed with H₂O, brine, dried over Na₂SO₄, concentrated to give the title compound. LCMS (ESI) calc. C₂₈H₂₆ClF₃N₃O₄ [M+H]⁺: 560.1, found: 560.1.

iv) Preparation of 4-(6-(((tert-butoxycarbonylamino)methyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)benzoic acid (F-5)


To a solution of compound **F-4** (162 mg, 0.29 mmol) in acetone (10 mL) was added Jones reagent (2 mL), and the mixture was kept stirring at rt for 12 h. MeOH (20 mL) was added to quench excess Jones reagent, and the mixture was filtered and rinsed with EtOAc. The organic layer was separated and concentrated to give the title compound. LCMS (ESI): calc'd for C₂₈H₂₄ClF₃N₃O₅ [M+H]⁺: 574.1, Found: 574.1.

v) Preparation of 4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl) benzoyl) -1*H*-indazol-3-yl)benzoic acid (27A)

To a solution of compound **F-5** (100 mg, 0.17 mmol) in CH₂Cl₂ (10 mL) was added excess amount of TFA. The mixture was stirred at rt for 2 h. The mixture was concentrated, diluted with H₂O, and extracted with EtOAc. The combined organics were washed with H₂O, brine, dried over Na₂SO₄ and concentrated to give the title compound. LCMS (ESI): calc'd for C₂₃H₁₆ClF₃N₃O₃ [M+H]⁺: 474.1, found: 474.1; ¹H NMR (400 MHz, MeOD) δ 8.76 (1H, s), 8.23-8.25 (1H, m), 8.04-8.06 (2H, d, J = 6.8Hz), 7.88-7.91 (2H, m), 7.77-7.85 (3H, m), 7.66-7.68 (1H, d, J = 8.8 Hz), 4.35 (2H, s).


The following examples shown in **TABLE 5** were prepared following similar procedures described for **Example #27A** in **Scheme F**, which can be achieved by those of ordinary skill in the art.

TABLE 5



Ex.	IUPAC name	$\begin{matrix} R^9 \\ \\ R^{10}N \end{matrix}$	P	Q	LCMS [M+H] ⁺ (found)
27B	4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid	H_2N			492

Example 28: Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)-2-(difluoromethyl)benzoic acid



5

i) Preparation of 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isobenzofuran-1(3H)-one (G-1)

A mixture of 5-bromoisobenzofuran-1(3H)-one (1.5 g, 7.0 mmol), 4,4,4',4', 5,5,5',5'-
 10 octamethyl-2,2'-bi(1,3,2-dioxaborolane) (2.1 g, 8.5 mmol), KOAc (2.1 g, 21 mmol), dppf (0.16 g, 0.28 mmol) and Pd(dppf)Cl₂ (0.26 g, 0.35 mmol) in dioxane (30ml) was heated at 100 °C for 4 h. The mixture was cooled down and concentrated under reduced pressure. The residue was diluted with H₂O, and extracted with EtOAc. The combined organic layers were

washed with brine, dried over Na_2SO_4 , and concentrated. The residue was purified by flash chromatography (PE: EA = 5:1) to give the title compound as a white solid. LCMS (ESI):calc. $\text{C}_{14}\text{H}_{18}\text{BO}_4$ $[\text{M}+\text{H}]^+$: 261, found: 261.

5 **ii) 5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)isobenzofuran-1 (3*H*)-one (G-2)**

A mixture of 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isobenzofuran-1(3*H*)-one (**G-1**) (0.97 g, 2.48 mmol), (3-bromo-1*H*-indazol-1-yl)(2-chloro-6-(trifluoromethyl)phenyl)methanone (1.0 g, 2.48 mmol), K_2CO_3 (1.03 g, 7.44 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (0.143 g, 0.12 mmol) were suspended in 1,4-dioxane (15 mL) and H_2O (3 mL). The reaction mixture was heated at 100 °C in a microwave reactor for 2 h. The reaction mixture was cooled to rt and diluted with H_2O . The aqueous layer was extracted with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by flash chromatography (PE: EA = 4:1) to afford the title compound as
10 a pale yellow oil. LCMS (ESI) calc'd for $\text{C}_{23}\text{H}_{13}\text{ClF}_3\text{N}_2\text{O}_3$ $[\text{M}+\text{H}]^+$: 457, found: 457.

iii) Preparation of benzyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)-2-(hydroxymethyl)benzoate (G-3)

A mixture of 5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)isobenzofuran-1(3*H*)-one (**G-2**) (0.6 g, 1.32 mmol) and LiOH (0.16 g, 6.58 mmol) in THF (6 mL) and H_2O (3mL) was stirred at rt for 3 h. The solvent was removed under reduced pressure, and the residue was dissolved in DMF (10 mL), followed by the addition of BnBr (0.78 mL, 6.6 mmol). The resulting mixture was stirred at rt for 3 h. The mixture was diluted with H_2O , and EtOAc. The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by flash chromatography (PE: EA = 4:1) to give the title compound as a colorless oil. LCMS (ESI) calc'd for $\text{C}_{30}\text{H}_{19}\text{ClF}_3\text{N}_2\text{O}_3$ for $[\text{M}+\text{H}]^+$: 585, found: 585.

30 **iv) Preparation of benzyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)-2-formylbenzoate (G-4)**

To a mixture of benzyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)-2-(hydroxymethyl)benzoate (**G-3**) (0.4 g, 0.71 mmol) in DCM (10 mL) was added Dess-Martin

periodinane (0.45 g, 1.07 mmol), and the mixture was stirred at rt for 24 h. The mixture was diluted with H₂O and extracted with DCM. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated. The resulting crude oil was used in the next step without further purification. LCMS (ESI) calc'd for C₃₀H₁₉ClF₃N₂O₄ [M+H]⁺: 563, found: 563.

v) Preparation of benzyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)-2-(difluoromethyl)benzoate (G-5)


To a solution of compound (G-4) (0.35 g, 0.62 mmol) in DCM (6 mL) at 0 °C was added a solution of DAST (0.42 mL, 3.1 mmol) in DCM (4 mL) dropwise. The mixture was warmed to rt and stirred for 2 h. The reaction mixture was then diluted with H₂O, and extracted with DCM. The combined organics were washed with brine, dried over Na₂SO₄, and concentrated. The residue was purified by flash chromatography (PE: EA = 10:1) to give the title compound as a white solid. LCMS (ESI) calc'd for C₃₀H₁₉ClF₅N₂O₃ [M+H]⁺: 585, found: 585.

vi) 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1*H*-indazol-3-yl)-2-(difluoromethyl)benzoic acid (28)

A mixture of compound (G-5) (0.17 mg, 0.30 mmol) and LiOH (63 mg, 1.5 mmol) in THF (6 mL) and H₂O (3 mL) was stirred at rt for 24 h. The mixture was acidified with 2M HCl to pH = ~3, and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified with prep-HPLC to give the title compound as a white solid. LCMS (ESI) calc'd for C₂₃H₁₃ClF₅N₂O₃ [M+H]⁺: 495, found: 495; ¹H NMR (400 MHz, CDCl₃) δ 8.70 (1H, d, *J* = 8 Hz), 8.35 (1H, s), 8.25 (1H, d, *J* = 8 Hz), 8.02-8.05 (2H, m), 7.71-7.77 (3H, m), 7.49-7.64 (3H, m).

Example 29: Preparation of 4-(1-(2-chloro-6-(trifluoromethyl) benzoyl)-6-cyano -1*H*-indazol- 3-yl)benzoic acid

Scheme H.



i) Preparation of methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-cyano-1H-indazol-3-yl)benzoate (H-2)

To a solution of compound **H-1** (100 mg, 0.186 mmol) (prepared according to a similar procedure in Scheme A) in DMF (2 mL) was added Zn(CN)_2 (44 mg, 0.37 mmol) and $\text{Pd(PPh}_3)_4$ (22 mg, 0.019 mmol). The mixture was purged with N_2 and then stirred at 100 °C for 1 h under microwave. The reaction mixture was cooled down to rt and diluted with H_2O , and extracted with EtOAc. The combined organics were dried over anhydrous Na_2SO_4 , filtered and concentrated. The crude solid was purified by Prep-TLC (20*20cm)(PE/EA=3/1) to obtain the title compound as a brown solid. LCMS (ESI) calc'd for $\text{C}_{24}\text{H}_{14}\text{F}_3\text{N}_3\text{O}_3\text{Cl}$ $[\text{M}+\text{H}]^+$: 484, Found: 484.

ii) Preparation of 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-cyano-1H-indazol-3-yl)benzoic acid (29)

A mixture of compound **H-2** (49 mg, 0.10 mmol), and LiOH (37 mg, 0.88 mmol) in dioxane (10 mL) and H_2O (2 mL) was stirred at rt for 24 h. The mixture was acidified with AcOH to $\text{PH} = \sim 4$, and extracted with DCM. The combined organics were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep HPLC to give the title compound as a white solid. LCMS (ESI) calc'd for $\text{C}_{23}\text{H}_{12}\text{F}_4\text{N}_3\text{O}_3\text{Cl}$ $[\text{M}+\text{H}]^+$: 470, found: 470; ^1H NMR (400 MHz, DMSO) δ 13.23 (1H, s), 8.99 (1H, s), 8.49 (1H, d, $J=8.4$ Hz), 8.03-8.12 (5H, m), 7.89-7.98 (3H, m).

Biological Assays

The compounds of the invention inhibit RORgammaT activity. Activation of RORgammaT activity can be measured using e.g. biochemical TR-FRET assay. In such an assay, interaction of cofactor-derived peptides with human RORgammaT-Ligand Binding Domain (LBD) can be measured. The TR-FRET technique is a sensitive biochemical proximity assay that will give information concerning the interaction of a ligand with the LBD, in the presence of cofactor derived peptides (Zhou et al., Methods 25:54-61, 2001).

To identify novel antagonists of RORgammaT, an assay was developed which employs the interaction of RORgammaT with its co-activator peptide SRC1_2. This peptide mimics the recruitment of co-activators to RORgammaT through its interaction with the LXXLL (eg NR box) motifs (Xie et al., J. Immunol. 175: 3800-09, 2005; Kurebayashi et al., Biochem. Biophys. Res. Commun. 315: 919-27, 2004; Jin et al., Mol. Endocrinology 24:923-29, 2010). The assay was run following three slightly different protocols, i.e. Protocol A, Protocol B, and Protocol C, which are described below. The TR-FRET assay following Protocol A is described below and the results thereof shown in Table 1 below.

TR-FRET assay (Protocol A)

In the current setting, a peptide from Steroid Receptor Co-factor 1 box 2 (SRC1_2) is aminoterminal biotinylated (biotinyl-CPSSHSSLTERHKILHRLQLQEGSPS, NEOMPS, France) and the applied RORgamma-LBD contains a 6-His and GST tag. The 6His(GST)RORgamma-LBD was expressed and purified from baculovirus infected Sf-21 insect cells (Bac-to-BAC, Invitrogen) and purified using glutathione sepharose chromatography (GE healthcare). Biotin and His function as anchoring points for streptavidin-allophycocyanin (Strep-APC, Perkin Elmer, C130-100) and anti-His-Europium Cryptate (Perkin Elmer, AD0111), respectively. Since energy transfer between the streptavidin and His-coupled labels APC and europium cryptate can only take place when these labels are in close proximity, measurement of energy transfer in a mixture of biotinylated peptide, His-GR-LBD, streptavidin-APC and anti-His-Europium Cryptate will be indicative for RORgamma-LBD/peptide binding. Final assay conditions in 50 µl are: TR-FRET buffer [50 mM KCl, 50 mM TRIS, 1 mM Na-EDTA, 0.1 mM DTT, (pH 7.0)]; 6His(GST)RORgamma-LBD 10 nM, Anti-6His-Eu 1.25 nM, SRC1_2 peptide 0.1 µM, Strep-APC 8 nM. Changes in peptide binding caused by dose-curves of compounds added to the

reaction mixture (DMSO concentration 0.1 %), can likewise be monitored. Fluorescence emission at 665 nm (excitation wavelength 340 nm) was read in each well of 384 well plates (Perkin Elmer, 60007299) using an Envision fluorescence reader in Time Resolved mode (Perkin Elmer) after overnight incubation at 4°C. Dose-response data are curve-fitted (IBDS Software, ActivityBase) and pEC50 values calculated. pEC50 values for Examples 1-23 using Protocol A are provided in Table 6 below. All compounds of the invention have a pEC50 of 4 or higher. Preferred are compounds with a pEC50 of more than 5.

Table 6

Example	pEC50
1	7.97
2	6.61
3	7.73
4	8.08
5	6.76
6A	7.87
6B	6.65
7	7.85
8	5.31
9	5.03
10	7.64
11A	6.47
11B	7.35
11C	6.29
11D	6.28
11E	7.57

11F	7.4
11G	7.22
11H	7.15
11I	7.14
11J	5.72
11K	7.67
11L	7.58
11M	7.98
11N	7.59
12	7.17
13A	6.81
13B	7.1
13C	7.12
13D	7.28
13E	7.58
13F	6.78
14	5.71
15A	6.56
15B	6.74
15C	7.85
15D	5.35
15E	6.29
17	5.91
18	5.48

16	6.02
20	5.45
21A	6.41
21B	6.24
21C	6.21
21D	6.13
22	5.3
23	5.99

TR-FRET assay (Protocol B)

HIS-tagged ROR γ -LBD was recombinantly expressed in SF9 cells using a baculovirus expression system. The protein was not purified. Cells were lysed and the lysate was used as a source for ROR γ -LBD for the assay. A 1:80 dilution of ROR γ -LBD lysate in assay buffer (25 mM HEPES pH 7.0, 100 mM NaCl, 0.01% Tween, 0.1% BSA) was prepared and 5 μ L was added to each well (ROR γ -LBD final concentration ~10 nM). Control wells received lysate from SF9 cells not expressing ROR γ -LBD.

Compounds to be tested were diluted to 100x final test concentration in DMSO and further diluted to 4x final test concentration using assay buffer to provide the test compound mixture. An aliquot (5 μ L) of the test compound mixture was added to each well.

A 4x stock of biotinylated-LXXLL peptide from SRC1-2 (Biotin-CPSSHSSLTERHKILHRLQEGSPS) was prepared in assay buffer and a 5 μ L aliquot added to each well (450 nM final concentration). A 4x solution of europium tagged anti-HIS antibody (2 nM final concentration) and APC conjugated streptavidin (60 nM final concentration) were prepared and a 5 μ L aliquot added to each well.

The final assay mixture was incubated for 4 hours to overnight, and the fluorescence signal was measured on an Envision plate reader: (Excitation filter = 340 nm; APC emission = 665 nm; Europium emission = 615 nm; dichroic mirror = D400/D630; delay time = 100 μ s, integration time = 200 μ s).

IC50 values for test compounds were calculated from the quotient of the fluorescence signal at 665 nm divided by the fluorescence signal at 615 nm using GraphPad Prism software.

5 TR-FRET assay (Protocol C)

HIS-tagged ROR γ -LBD protein was expressed in SF9 cells using a baculovirus expression system. The ROR γ -LBD protein was purified by glutathione sepharose chromatography. Separately, SF9 cells not expressing any recombinant protein were lysed and the lysate was added to the purified ROR γ -LBD at 0.25 μ l lysate (from 10,000 SF9
10 cells)/nM purified protein. The mixture was then diluted in assay buffer (50 mM Tris pH 7.0, 50 mM KCl, 1 mM EDTA, 0.1 mM DTT) to obtain ROR γ -LBD final concentration of 3 nM in 384-well assay plate.

Compounds to be tested were injected to the assay plate using Acoustic Droplet Ejection technology by Echo 550 liquid handler (Labcyte, CA).

15 A stock of biotinylated-LXXLL peptide from coactivator SRC1 (Biotin-CPSSHSSLTERHKILHRLQEGSPS) was prepared in assay buffer and added to each well (100 nM final concentration). A solution of Europium tagged anti-HIS antibody (1.25 nM final concentration) and APC conjugated streptavidin (8 nM final concentration) were also added to each well.

20 The final assay mixture was incubated for overnight at 4°C, and the fluorescence signal was measured on an Envision plate reader: (Excitation filter = 340 nm; APC emission = 665 nm; Europium emission = 615 nm; dichroic mirror = D400/D630; delay time = 100 μ s, integration time = 200 μ s). IC50 values for test compounds were calculated from the quotient of the fluorescence signal at 665 nm divided by the fluorescence signal at 615 nm.

25 Table 7 below tabulates the biological data disclosed for Examples 24A-29 as a range of FRET IC50 values, which was reported as ranges according to the annotation delineated below:

- * IC50 <50 nM
- ** IC50 >50 and <500 nM
- 30 *** IC50 >500 and <10000 nM
- **** IC50 >10000 nM

Table 7

Examples	Fret IC ₅₀	Assay protocol
24A	*	B
24B	*	B
24C	*	B
24D	***	B
24E	***	B
24F	***	B
24G	***	B
24H	*	B
24I	*	B
24J	**	B
24K	*	B
24L	*	B
24M	*	B
24N	*	B
24O	*	B
24P	*	B
24Q	**	B
24R	*	B
24S	***	B
24T	****	B
24U	*	B
24V	*	B

24W	*	B
24X	**	B
24Y	*	B
24Z	****	B
24AA	*	B
24AB	*	B
24AC	***	B
24AD	***	B
24AE	***	B
24AF	*	B
24AG	*	B
24AH	*	B
24AI	**	B
24AJ	**	B
24AK	*	B
24AL	****	C
24AM	*	B
24AN	*	B
24AO	*	B
24AP	***	B
24AQ	*	B
24AR	*	B
24AS	*	C
24AT	*	B


24AU	*	C
24AV	**	C
24AW	**	C
24AX	*	B
24AY	**	B
24AZ	**	B
24BA	*	B
24BB	*	B
24BC	***	B
24BD	***	B
24BE	*	B
24BF	*	B
24BG	**	C
25A	*	B
25B	*	B
25C	*	B
25D	*	B
25E	*	B
25F	*	B
25G	**	B
25H	*	B
25I	**	B
25J	*	B
25K	*	B

25L	*	B
25M	*	B
25N	*	C
25O	*	B
25P	*	B
25Q	*	B
25R	*	B
25S	*	C
25T	*	C
25U	*	C
25V	*	C
25W	*	C
26A	*	B
26B	*	B
26C	*	B
26D	*	B
26E	*	B
26F	*	C
26G	*	C
26H	*	B
26I	*	B
26J	*	B
27A	*	B
27B	*	B

28	*	B
29	**	B

CLAIMS

1. A compound according to Formula I



I

or a pharmaceutically acceptable salt or solvate thereof wherein,

X is C(O), SO or SO₂,

Y is CH or N;

n = 0, 1, 2, 3 or 4;

A⁴-A⁷ are independently N or CR⁴, CR⁵, CR⁶ or CR⁷, respectively, with the proviso that no more than two of the four positions A⁴-A⁷ can be simultaneously N;


R¹ is (i) (C₃₋₇)cycloalkyl or (C₃₋₅)heterocycloalkyl, both optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens; (ii) (C₂₋₉)heteroaryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens; or (iii) (C₆₋₁₄)aryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl,

(C₁₋₄)alkyl or (C₁₋₃)alkoxy are optionally substituted with one or more halogens;


R² is C(O)OH, 5-tetrazoyl, HOC(CF₃)₂, C(O)OH(C₁₋₁₀)alkyl, (C₁₋₁₀)alkylsulfoxyaminocarbonyl; or carbamoyl;

R³ is hydrogen, halogen, cyano, nitro, hydroxy, (C₁₋₃)alkylC(O)O-, (C₁₋₄)alkyl, or (C₁₋₄)alkoxy, wherein (C₁₋₄)alkyl and (C₁₋₄)alkoxy are optionally substituted with one or more halogen;

R⁴-R⁷ independently are H, halogen, amino, cyano, hydroxy, (C₁₋₃)alkoxy, (C₁₋₄)alkyl, (C₀₋₁₀)alkylaminocarbonyl, (di)(C₁₋₆)alkylaminocarbonyl or amino(C₁₋₄)alkyl, wherein (C₁₋₃)alkoxy, (C₁₋₄)alkyl, (C₀₋₁₀)alkylaminocarbonyl, (di)(C₁₋₆)alkylaminocarbonyl and amino(C₁₋₄)alkyl are optionally substituted with one or more halogen, hydroxyl or (C₁₋

₃)alkoxy; or a group having the formula , optionally substituted with one or more of the following: (C₁₋₁₀)alkyl, halogen, amino, cyano, hydroxy, (C₁₋₃)alkoxy, and wherein m is 1, 2, 3, or 4.

2. A compound according to Formula Ia



Ia

or a pharmaceutically acceptable salt or solvate thereof

wherein X represents C(O), SO or SO₂,

Y is CH or N;

A⁴-A⁷ are independently N or CR⁴, CR⁵, CR⁶ or CR⁷, respectively, with the proviso that

no more than two of the four positions A⁴-A⁷ can be simultaneously N;

R¹ is (C₃₋₇)cycloalkyl or (C₃₋₅)heterocycloalkyl, both optionally substituted with one or more groups selected from halogen, amino, cyano, hydroxy, H₂NC(O)-, or (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all optionally substituted with one or more halogens or;

R¹ is (C₂₋₉)heteroaryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, or (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all optionally substituted with one or more halogens or;

R¹ is (C₆₋₁₄)aryl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, or (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, all optionally substituted with one or more halogens;

R² is C(O)OH, 5-tetrazoyl, or HOC(CF₃)₂;

R³ is independently selected from hydrogen, halogen, cyano, nitro or (C₁₋₄)alkyl, optionally substituted with one or more halogens; and

R⁴-R⁷ independently are H, halogen, amino, cyano, hydroxy, (C₁₋₃)alkoxy, (C₁₋₄)alkyl, optionally substituted with one or more halogens.

3. The compound of claim 1, wherein A⁴, A⁵, A⁶, A⁷ are selected from the group consisting of: (i) CR⁴, CR⁵, CR⁶, CR⁷; (ii) N, CR⁵, CR⁶, CR⁷; and (iii) CR⁴, N, CR⁶, CR⁷.

4. The compound of claim 3, wherein A⁴, A⁵, A⁶, A⁷ is (i) CR⁴, CR⁵, CR⁶, CR⁷, or (ii) N, CR⁵, CR⁶, CR⁷; and Y is N.

5. The compound of claim 4, wherein

X is C(O) or SO₂;

R¹ is (i) (C₃₋₇)cycloalkyl or (C₃₋₅)heterocycloalkyl, both optionally substituted with one or more groups selected from (C₁₋₄)alkyl or halogen; (ii) (C₂₋₉)heteroaryl, optionally substituted with one or more groups selected from halogen, amino or (C₁₋₄)alkyl; or (iii) (C₆₋₁₄)aryl, optionally substituted with one or more groups selected from halogen,

amino, cyano, nitro, hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl, (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens.

6. The compound of claim 5, wherein


R¹ is (C₂₋₉)heteroaryl or (C₆₋₁₄)aryl, both optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens.

7. The compound of claim 6, wherein R¹ is phenyl, naphthyl, pyridinyl, quinolinyl, benzooxadiazolyl, thiophenyl, or isoxazolyl, each optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens.

8. The compound of claim 7, wherein R¹ is phenyl, optionally substituted with one or more groups selected from halogen, amino, cyano, nitro, hydroxy, (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (C₁₋₄)alkyl and (C₁₋₃)alkoxy are optionally substituted with one or more halogens.

9. The compound of claim 8, wherein R² is C(O)OH.

10. The compound of claim 3 having Formula Ib




Ib

and a pharmaceutically acceptable salt or solvate thereof.

5

11. The compound of claim 10 having Formula Ic



Ic

10

or a pharmaceutically acceptable salt or solvate thereof wherein,


X is C(O) or SO₂

R⁸ is selected from halogen, amino, cyano, nitro, hydroxy, H₂NC(O)-, (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy, wherein (C₁₋₃)alkoxycarbonyl, (di)(C₁₋₆)alkylaminocarbonyl, (C₁₋₄)alkyl or (C₁₋₃)alkoxy are optionally substituted with one or more halogens; and x is 0, 1, 2, 3, 4 or 5.

15

12. The compound of claim 11 having Formula Id


20



Id

5 and a pharmaceutically acceptable salt or solvate thereof.

13. The compound of claim 12 having Formula Ie



Ie

and a pharmaceutically acceptable salt or solvate thereof.

15 14. A compound according to claim 1 selected from:

- 4-(1-(2,6-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-5-fluoro-1H-indazol-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-7-fluoro-1H-indazol-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;
5 4-(1-(2,6-dichlorobenzoyl)-6-fluoro-1H-indazol-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;
4-(1-(2-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-1H-pyrrolo[3,2-b]pyridin-3-yl)benzoic acid;
4-(1-(2,6-dichlorobenzoyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acid;
10 1,1,1,3,3,3-hexafluoro-2-(4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)phenyl)propan-2-ol;
4-(1-(2-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(3,5-dichloroisonicotinoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-bromobenzoyl)-1H-indazol-3-yl)benzoic acid;
15 4-(1-(2-(methoxycarbonyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(cyclohexanecarbonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-bromo-6-chlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-fluoro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
20 4-(1-(2,4,6-trichlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,6-dimethylbenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-ethoxybenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,3-dichlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-chlorobenzoyl)-1H-indazol-3-yl)benzoic acid;
25 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-methyl-6-nitrobenzoyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-(trifluoromethyl)phenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(quinolin-8-ylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2-bromophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
30 4-(1-(2,5-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(3-chloro-2-methylphenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
4-(1-(2,3-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;

- 4-(1-(2,3,4-trichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 4-(1-(2-cyanophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 4-(1-(2-(trifluoromethoxy)phenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 4-(1-(naphthalen-1-ylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 5 4-(1-(2,6-dichlorophenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 4-(1-(benzo[c][1,2,5]oxadiazol-4-ylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 4-(1-(2-cyano-5-methylphenylsulfonyl)-1H-indazol-3-yl)benzoic acid;
 4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;
 4-(1-(3,5-dimethylphenylsulfonyl)-1H-indazol-3-yl)-2-(trifluoromethyl)benzoic acid;
 10 4-(1-(2-chlorophenylsulfonyl)-1H-pyrazolo[4,3-c]pyridin-3-yl)benzoic acid;
 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(3-chlorophenylsulfonyl)-1H-indazole;
 methyl 3-(3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-ylsulfonyl)thiophene-2-
 carboxylate;
 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(2-bromo-4-fluorophenylsulfonyl)-1H-indazole;
 15 3-(4-(1H-tetrazol-5-yl)phenyl)-1-(3-bromo-5-chlorothiophen-2-ylsulfonyl)-1H-
 indazole;
 4-(3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-ylsulfonyl)-3,5-dimethylisoxazole;
 (3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-yl)(2,6-dichlorophenyl)methanone;
 (3-(4-(1H-tetrazol-5-yl)phenyl)-1H-indazol-1-yl)(2,3-dichlorophenyl)methanone;
 20 methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-
 fluorobenzoate;
 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-fluorobenzoic
 acid;
 Sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-
 25 fluorobenzoate;
 methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoate;
 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;
 sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoate;
 methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-
 30 fluorobenzoate;
 sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-
 fluorobenzoate;

(2-chloro-6-(trifluoromethyl)phenyl)(4-fluoro-3-(4-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)-1H-indazol-1-yl)methanone;
3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;
2-(3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)phenyl)acetic
5 acid;
2-(4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)phenyl)acetic
acid;
2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic
acid;
10 Sodium 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzamide;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluoro-5-methylbenzoic acid;
15 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-methylbenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-methoxybenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluorobenzoic
20 acid;
sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluorobenzoate;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-methylbenzoic
acid;
25 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-methoxybenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,6-difluorobenzoic acid;
4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-isopropylbenzoic acid;
30 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3,5-dimethoxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-cyano benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,3-difluorobenzoic acid;

5 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2,5-difluorobenzoic acid;

2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluorobenzoic acid;

10 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluoro-2-methoxybenzoic acid;

5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoic acid;

3-fluoro-4-(4-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;

4-(4-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;

15 3-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl) benzoic acid;

2-(4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)phenyl)acetic acid;

20 (2-chloro-6-(trifluoromethyl)phenyl)(3-(4-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)-1H-pyrazolo[4,3-b]pyridin-1-yl)methanone;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-3-methylbenzoic acid;

3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;

25 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-methoxybenzoic acid;

30 2-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-5-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2,5-difluorobenzoic acid;

5-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-hydroxybenzoic acid;

4-(1-(2-fluoro-6-methoxybenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;

3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-(trifluoromethyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-isopropylbenzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-fluoro benzoate;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-3-fluoro benzoate;

3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;

3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;

sodium 3-fluoro-4-(1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoate;

4-(7-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;

3-fluoro-4-(7-fluoro-1-(2-fluoro-6-methoxybenzoyl)-1H-indazol-3-yl)benzoic acid;

4-(6-bromo-1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl) benzoic acid;

4-(6-bromo-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-methoxy-1H-indazol-3-yl)

benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl)benzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl) benzoate;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-hydroxy-1H-indazol-3-yl)-3-

fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-hydroxy-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-6-hydroxy-1H-indazol-3-yl)-3-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(hydroxymethyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(hydroxymethyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(methoxymethyl)-1H-indazol-3-yl)benzoic acid;

5 1-(2-chloro-6-(trifluoromethyl)benzoyl)-3-(4-(methoxy carbonyl)phenyl)-1H-indazole-6-carboxylic acid;

methyl 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoate;

10 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxy ethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoate;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

15 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoate;

20 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoate;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)-2-fluorobenzoic acid;

25 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-2-fluorobenzoic acid;

30 3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

3-chloro-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxyethylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-hydroxypropylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-hydroxypropylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

5 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-methoxypropylcarbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(2-hydroxypropylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

10 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(4-hydroxybutylcarbamoyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-hydroxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-hydroxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

15 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-((2-methoxyethyl)(methyl)carbamoyl)-1H-indazol-3-yl)-3-fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3-fluoroazetidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid;

20 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(3,3-difluoroazetidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(pyrrolidine-1-carbonyl)-1H-indazol-3-yl)benzoic acid;

4-(6-(azetidine-1-carbonyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)benzoic acid;

25 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoic acid;

2-acetoxy-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)benzoic acid;

30 sodium 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-hydroxybenzoate;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-2-fluoro-6-hydroxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3-

hydroxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-5-fluoro-2-

hydroxybenzoic acid;

5 4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-4-fluoro-1H-indazol-3-yl)-3,6-difluoro-2-

hydroxybenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-yl)-2-

hydroxybenzoic acid;

2-acetoxy-4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-pyrazolo[4,3-b]pyridin-3-

10 yl)benzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-(dimethylcarbamoyl)-1H-indazol-3-yl)-2-

hydroxybenzoic acid;

4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl) benzoyl)-1H-indazol-3-yl)benzoic
acid;

15 4-(6-(aminomethyl)-1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-3-

fluorobenzoic acid;

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-1H-indazol-3-yl)-2-(difluoro

methyl)benzoic acid; and

4-(1-(2-chloro-6-(trifluoromethyl)benzoyl)-6-cyano-1H- indazol-3-yl)benzoic acid.

20

15. A pharmaceutical composition comprising a compound of claim 1 or a pharmaceutically acceptable salt or solvate thereof and one or more pharmaceutically acceptable excipients.

25 16. The pharmaceutical composition of claim 15, which further comprises at least one additional therapeutically active agent.

17. Use of a compound of claim 1, or a pharmaceutically acceptable salt or solvate thereof, in the manufacture of a medicament for the treatment of a disease or condition mediated
30 by Retinoic acid receptor-related Orphan Receptor gamma t (RORgammaT).


18. A method for treating a disease or condition mediated by RORgammaT in a subject comprising administering to the subject an amount of a compound of claim 1, or a pharmaceutically acceptable salt or solvate thereof, that is effective for treating the disease or condition mediated by RORgammaT in the subject.

5

19. The method of claim 18, wherein the disease or condition is an autoimmune disease or inflammatory disease.

20. The method of claim 19, wherein the disease or condition is multiple sclerosis,
inflammatory bowel disease, Crohn's disease, psoriasis, rheumatoid arthritis, asthma,
osteoarthritis, Kawasaki disease, Hashimoto's thyroiditis or mucosal leishmaniasis.

10



(I)