

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number
WO 01/36595 A2

(51) International Patent Classification⁷: **C12N 5/00**

(US). **FALCO, S., Carl** [US/US]; 1902 Millers Road, Wilmington, DE 19810 (US).

(21) International Application Number: PCT/US00/31600

(22) International Filing Date:
16 November 2000 (16.11.2000)

(74) Agent: **FELTHAM, S., Neil**; E.I. Du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).

(25) Filing Language: English

(81) Designated States (*national*): AU, BR, CA, HU, IL, JP, KR, MX, NZ, PL, US.

(26) Publication Language: English

(30) Priority Data:
09/442,021 17 November 1999 (17.11.1999) US

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(71) Applicant (*for all designated States except US*): **E.I. DU PONT DE NEMOURS AND COMPANY** [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).


Published:
— *Without international search report and to be republished upon receipt of that report.*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **YADAV, Nerendra, S.** [IN/US]; 3704 Knole Lane, Chadds Ford, PA 19317

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR CONDITIONAL TRANSGENE EXPRESSION AND TRAIT REMOVAL IN PLANTS



(57) **Abstract:** This invention relates to constructs for the conditional or regulated expression or excision of transgenes in plants using site-specific recombinase systems. The constructs comprise a variety of constitutive, inducible, tissue specific or developmental stage-specific promoters operably linked to either a transgene or the elements of one or more site-specific recombinase system. By matching promoters, responsive to various inducers, plant tissues or plant developmental states with the recombinase systems, stop fragments and transgenes, virtually any trait may be expressed or excised at any plant development stage or in any plant generation.

WO 01/36595 A2

TITLE

METHODS FOR CONDITIONAL TRANSGENE EXPRESSION AND TRAIT REMOVAL IN PLANTS

This application is a continuation in part of USSN 09/442,021, filed
5 November 17, 1999.

FIELD OF INVENTION

The present invention relates to the field of molecular biology and the
genetic transformation of plants with foreign gene fragments. More particularly,
the invention relates to methods for the conditional or regulated expression or
10 excision of genetic traits from plant progeny using site specific recombinase
constructs.

BACKGROUND OF THE INVENTION

The advent of genetically modified crops holds the promise of improving
crop yield and productivity as well as reducing the number of pesticides and other
15 toxic compounds needed for crop success. These benefits are conferred via the
transformation of crop plants with new genes beneficial to growth or conveying
disease or pathogen resistance. However, transformation events often result in the
presence of ancillary genetic material that is not useful for the expression of the
desired trait. These ancillary sequences are necessary for the transformation
20 processes, but they do not positively contribute to the final cultivar and in fact
lessen its desirability to the consumer. The presence of these undesirable
sequences may also complicate the regulatory procedures necessary to bring the
cultivar to the market place. A reliable method for eliminating the unwanted
ancillary sequences would thus improve commercial viability by increasing public
25 acceptance and simplify the regulatory process. Additionally, it will be useful to
have certain traits only present during intermediate generations and then removed
in harvested generations. The prior art has not recognized the importance of these
problems, nor has it worked to provide a solution.

Plants are increasingly being looked to as platforms for the production of
30 materials, foreign to plant systems. As the art of genetic engineering advances it
will be possible to engineer plants for the production of a multiplicity of
monomers and polymers, currently only available by chemical synthetic means.
The accumulation of these materials in various plant tissues will be toxic at some
level and it will be useful to tightly regulate the relevant genes to prevent
35 expression in inappropriate plant tissues.

Currently few methods exist that provide for tightly regulated transgene
expression. Non-specific expression of transgenes in non-target cells, tissues, or
generation hinders plant transgenic work. This is important where the goal is to

produce such high levels of materials in transgenic plants that may be phytotoxic or adversely affects normal plant development . Conditional transgene expression would enable economic production of desired chemicals, monomers, and polymers at levels likely to be phytotoxic to growing plants by restricting
5 their production to production tissue of transgenic plants either just prior to or after harvest of the crop biomass used for extracting the desired product. Therefore, lack of a commercially usable conditional expression system and the difficulty in attaining a reliable, high-level expression both limit development of transgene expression in plants.

10 Conditional or regulated expression has been reported in plants (see De Veylder, L. et al., *Plant Cell Physiol.* 38:568-577 (1997); Gatz, C., *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48:89-108 (1997); Hansen, G. et al., *Mol. Gen. Genet.* 254:337-343 (1997); Jepson, I., PCT Int. Appl. WO 9706269 A1 (1997); Jepson, I, et al. PCT Int. Appl. WO 9711189 A2 (1997), and other references
15 within this application). However, when tested stringently for basal non-specific expression, very few have been strictly specific (Odell J. T. et al., *Plant Physiol.* 106:447-458 (1994); van der Geest et al., *Plant Physiol.*, 109(4), 1151-58 (1995); Ma et al., *Aust. J. Plant Physiol.*, 25(1), 53-59 (1998); Czako et al., *Mol. Gen. Genet.*, 235(1), 33-40 (1992)). Such promoters are not suitable for some
20 applications, such as the use of transgenes for expressing novel phytotoxic proteins, enzymes that lead to the biosynthesis of phytotoxic products, and/or gene silencing. Site-specific recombination in plants (Odell et al., *Plant Physiol.* 106:447-458 (1994); Odell et al., PCT Int. Appl. WO 9109957 (1991); Surin et al., PCT Int. Appl WO 9737012 (1997)) and the reduction in the proficiency of
25 Cre-mediated recombination by mutant lox P sites and their use in increasing the frequency of Cre-lox based integration have been reported (Albert et al., *Plant J.* 7:649-59 (1995); Araki et al., *Nucleic Acids Res.* 25:868-872 (1997)). However, the use of the mutant sites to enhance the specificity Cre-mediated recombination in conjunction with chimeric Cre genes under the control of available regulated
30 promoters has not been demonstrated. Thus, there is a need for an appropriately stringent, site-specific recombination system for a commercially-attractive, conditional site-specific recombination.

Directed excision of a transgene from the plant genome has been reported using recombinase specific-sites and a recombinase. In Russel et al. (*Mol. Gen. Genet.* 234:49-59 (1992)), utility of these techniques is evaluated to remove a
35 selectable marker for transformation from the plant, while leaving a preferred non-selectable trait. Variation in efficiency of excision of transgenes in different plants was also examined and comparison was made between introduction of the

cre gene by re-transformation or by cross pollination. However, incorporation of promoters for conditional or regulated expression was not attempted.

Ow et al. (PCT Int. Appl. WO 9301283 A1 (1992)) also examine directed excision of selectable markers for transformation from plants, while leaving a preferred non-selectable trait that could be operably linked to control sequences capable of effecting the timing of said expression in higher plants. Their disclosed technique is limited by the requirement that the recombinase for excision must be introduced via a second round of transformation either directly to the initially transformed plants or by cross pollination of the independently regenerated plants from first and second round transformants.

In one disclosure, Oliver et al. (US 5,977,441) demonstrate limited expression of a desired introduced gene in a transgenic plant, according to a particular stage of plant development, a particular plant tissue, particular environmental conditions, or a particular time or location. This is achieved via: 1) the insertion of a transiently-active promoter in functional relation with a structural gene whose expression results in an altered plant phenotype, but separated by a blocking sequence that is flanked by specific excision sequences; and 2) a second DNA sequence that comprises an inducible promoter operably linked to a gene encoding a site-specific recombinase. These methods ultimately permit the expression of certain plant traits to be under external control by application of an external stimulus, through hybridization, or by direct introduction of a recombinase into a plant. However, Oliver et al. did not consider the need for eliminating unwanted ancillary sequences, such as selection markers.

Surin et al. (PCT Int. Appl. WO 9737012 A1 (1997)) disclose a method whereby single-step excision of transgenes is achieved, such as selectable marker genes or reporter genes, from genetically-transformed organisms. This is possible by the incorporation of both a recombinase genetic unit (comprising a promoter, recombinase gene, and terminator) and a transgene unit (comprising a promoter, transgene, and terminator) within a single expression cassette, all flanked by recombinase sites. This genetic unit facilitates multiple sequential genetic transformation events using the same selectable marker gene and provides means for tightly regulating transgene expression in genetically-manipulated plants. Additionally, the need for multiple transformation events or sexual crossing to produce a single cell comprising both the loci for DNA recombination and the site-specific recombinase is alleviated, since their method facilitates precise excision of genetic material in a single generation using promoters that are differentially activated. Although Surin et al. allow for the incorporation of a second separate expression cassette, their techniques are limited in that a

maximum of two transgenic units may be expressed within the plant at different time periods, the first of which must be removed prior to the expression of the second. At no point may the second transgene be removed from the genome if desired.

5 Hodges et al. (US 6,110,736) describe a method based on homologous recombination which enables the targeting of a length of DNA to a specific non-lethal site in the host's genome, and provides for the removal of any randomly inserted DNA sequences, using site-specific recombinase sites and the corresponding recombinase protein. The need met through the application of the
10 site-specific recombinase system for gene excision is specifically to maintain control over the copy number and the location of the inserted DNA. A further embodiment of the invention combines use of cre/lox and FLP/FRT to ultimately leave only the desired DNA sequences integrated into the chromosome while the selectable marker is removed from the chromosome. However, this work relies is
15 confined to instances where precise homologous recombination can be executed, which necessarily requires precise engineering of the vector sequences to match sequences within the chromosome.

 Finally, Perez and Flament (PCT Int. Appl. WO 9838323 (1998)) discuss the insertion of a of a male sterility gene (AMS) to create a cytoplasmic or nuclear male
20 sterile plant. This plant avoids the dissemination of pollen, and therefore necessarily prevents dissemination of any transgenes that are linked to the male sterility genes. In further embodiments of their invention, the AMS gene can serve as a positive marker for the screening of plants having integrated a transgene of interest; or, the gene, when genetically linked to a transgene, can be excised using a system of
25 transposition (e.g. systems of recombination such as Cre/lox or FLP/ FRT). Limitations of these techniques focus on the requirement that a transgene must be directly linked to a transgene, thereby requiring linked expression. No allowance is made for conditional expression, whereby expression of one gene may activate, or excise, another later in the life cycle.

30 The methods described above are useful for removal of transgenes and ancillary genetic material from plants but are limited in their ability to regulate transgene expression at various times in a plant life cycle or in hybrid progeny. The problem to be solved therefore is to develop a method for the tightly regulated conditional expression and removal of genetic traits in plants, both in initial
35 transformants and in hybrid.

 Applicant has solved the stated problem by providing developmentally regulated germ-like promoters that allow using at least two different recombinases for the selective expression and/or removal of genetic traits.

SUMMARY OF THE INVENTION

The present invention provides methods for the conditional or regulated expression of a transgene in a plant, or the excision of a trait or marker in a plant using a suite of constitutive, inducible, tissue specific or developmental stage-specific promoters in combination with at least one recombinase system. Transgenes may encode specific traits, including transformation markers. By pairing the promoters with the appropriate recombinase and/or transgenes virtually any trait may be expressed at any time during a plant life cycle. Similarly, traits whose usefulness is limited to one portion of the life cycle or to only one generation may be selectively excised from subsequent generations.

For example, it is contemplated that different site specific recombinase systems under the control of different promoters may be combined or linked to enable a series of genetic switches for the regulated expression as well as removal of transgenes staggered over time during a single life cycle of the organism. Removal of transgene(s) from transgenic crops is useful for safety and protection of the environment, enhanced breeding, or for conditional transgene expression in only one generation, such as male sterility. For example, combinations of recombinase elements comprising two or more site-specific recombination (SSR) systems may be designed to effect regulated expression and/or removal of trait gene. This coupling of conditional and tissue-specific promoters with two or more site-specific recombinations, such that the conditional expression of one recombinase activates another later in the life cycle, allows their use as a series of genetic switches.

Although SSRs have been used singly as genetic switches, two (or more) SSRs under the control of different constitutive or regulated promoters can be used as a series of genetic switches within a plant's life cycle, such that conditional expression of one recombinase element at one stage activates another recombinase element at a later stage in the same or the subsequent generation. Thus, one can conditionally trigger the process at a convenient developmental stage, such as germination, but get delayed effects at later stages. Trait gene removal may be unlinked or linked to trait gene expression. The latter provides a more stringent control of trait gene expression. One of the key aspects of the invention is that expression of the various recombinase enzymes does not have to occur immediately upon enablement or priming of the recombinase elements (i.e., removal of the stop fragment) but are rather controlled solely by the choices of the respective promoters.

Accordingly the invention provides a trait removal construct comprising:

a) A first recombinase element having the general structure:

P1-R and;

b) A second recombinase element selected from the group of general structures consisting of K-TG, and RS-K-TG-RS;

wherein,

- 5 (i) P₁ is a first promoter;
- (ii) R is a recombinase coding sequence and 3' region
- (iii) TG is a transgene;
- (iv) RS is a recombinase site responsive to the recombinase
- (v) K is selected from the group consisting of;

10 1) P₂-RS-STP-RS and

2) P₂

wherein P₂ is a second promoter, RS is a recombinase site responsive to the recombinase and STP is a stop fragment;

wherein P₁ and P₂ are operably linked to their down stream elements and

15 wherein P₁ is activated prior to or at the same time as P₂ in the plant life cycle and wherein expression of the recombinase results in excision of any element contained between the recombinase site responsive to the recombinase.

20 In an alternate embodiment the invention provides a trait removal construct comprising:

a) a first recombinase element selected from the group of general structures consisting of :

P₁-R₁ and P₁-R₂, and;

25 b) a second recombinase element having the general structure RS₂-K-TG-RS₂;

wherein,

- (i) P₁ is a first promoter;
- (ii) R₁ is a first recombinase coding sequence and 3' region;
- (iii) R₂ is a second recombinase coding sequence and 3' region;
- 30 (iv) RS₂ is a second recombinase site responsive to the second recombinase;
- (v) TG is a transgene sequence and 3' region ;
- (vi) K is selected from the group consisting of;

1) P₂-RS₁-STP-RS₁ and

35 2) P₂-RS₁-TG

wherein P₂ is a second promoter, RS₁ is a first recombinase site responsive to the first recombinase, STP is a stop fragment and TG is a transgene sequence and 3' region;

wherein P_1 and P_2 are operably linked to their down stream elements, and wherein P_1 is activated earlier than or at the same time as P_2 in the plant life cycle, and wherein expression of the first recombinase results in excision of any element contained
 5 between the first recombinase sites responsive to the first recombinase and expression of the second recombinase results in excision of any element contained between the second recombinase sites responsive to the second recombinase.

In another embodiment the invention provides a trait removal construct
 10 comprising:

- a) a first recombinase element selected from the group of general structures consisting of P_1 - R_1 and RS_2 - P_1 - R_1 - RS_2 ;
 - b) a second recombinase element selected from the group of general structures Z - Y and RS_2 - Z - Y - RS_2 ;
 - 15 c) a third recombinase element selected from the group of general structures Q - X and RS_2 - Q - X - RS_2 ;
- Wherein:
- (i) P_1 is a first promoter;
 - (ii) R_1 is a first recombinase coding sequence and 3' region;
 - 20 (iii) RS_2 is a second recombinase site responsive to a second recombinase;
 - (iv) Z has the general formula, P_2 - RS_1 -STP- RS_1 , wherein P_2 is a second promoter, RS_1 is a first recombinase site responsive to a first recombinase, and STP is a stop fragment;
 - 25 (v) Y is selected from the group consisting of R_2 and TG wherein R_2 is a second recombinase coding sequence and 3' region and TG is a transgene sequence and 3' region;
 - (vi) Q has the general formula P_3 -RS-STP-RS, wherein P_3 is a third promoter, RS is a recombinase site selected
 30 from the group consisting of RS_1 and RS_2 and STP is a stop fragment;
 - (vii) X is selected from the group consisting of TG and R
 Wherein TG is a transgene sequence and 3' region and R is a recombinase coding sequence and 3' region selected from the group
 35 consisting of R_1 and R_2 ;
- wherein P_1 , P_2 and P_3 are operably linked to their down stream elements and wherein P_1 is activated earlier than P_2 in the plant life cycle, and wherein P_2 is activated earlier than P_3 in the plant

life cycle, and wherein expression of the first recombinase results in excision of any element contained between the first recombinase sites responsive to the first recombinase and expression of the second recombinase results in excision of any element contained between the second recombinase sites responsive to the second recombinase.

The constructs of the present invention will comprise a first promoter selected from the group consisting of:

- (a) constitutive plant promoters;
- (b) plant tissue-specific promoters;
- (c) plant development-specific promoters;
- (d) inducible plant promoters;
- (e) viral promoters;
- (f) male germline-specific promoters;
- (g) female germline-specific promoters;
- (h) male/female germline-specific promoters;
- (i) flower-specific promoters; and
- (j) vegetative shoot apical meristem-specific promoters.

Where the constructs of the present invention contain a second promoter that promoter is selected from the group consisting of: constitutive plant promoters; plant tissue-specific promoters; plant development stage-specific promoters; inducible plant promoters; and viral promoters.

Where the constructs of the present invention comprise a transgene for expression the transgene is selected from the group consisting of: genes encoding a transformation marker; genes encoding a morphological trait; genes conveying sterility; genes conveying specific phenotype on a plant or plant cell; and hormone biosynthetic genes.

Additionally the different components of the invention are heritable independently and may be introduced together into a transgenic plant or brought together by crossing transgenic plants carrying the separate components, such as by the method to produce TopCross[®] high oil corn seed (U.S. Patent No. 5,704,160). Also provided are methods of making the expression cassettes and methods of using them to produce transformed plant cells having an altered genotype and/or phenotype.

BRIEF DESCRIPTION OF FIGURES AND SEQUENCE DESCRIPTIONS

Figure 1 illustrates a scheme for the use of a single site-specific recombinase system for the conditional expression of a transgene gene in a first generation plant.

Figure 2 illustrates a scheme for the use of a single site-specific recombinase system for the regulated excision of a transgene in a first generation plant.

5 Figure 3 illustrates a scheme for the use of a dual site-specific recombinase system for the conditional expression of a transgene in a second generation plant.

Figure 4 illustrates a scheme for the use of a dual site-specific recombinase system for the conditional expression of a transgene in a first generation plant and its subsequent removal.

10 Figure 5 illustrates a scheme for the use of a dual site-specific recombinase system to effect conditional male sterility in a first generation plant.

Figure 6 illustrates a scheme for the use of a dual site-specific recombinase system to effect conditional gametophytic male sterility in a second generation plant.

15 Figure 7 illustrates a scheme for the use of a dual site-specific recombinase system to effect conditional male sterility in a Top Cross ® second generation.

Figure 8 illustrates a scheme for the use of a dual site-specific recombinase system to effect conditional male sterility for hybrid seeds.

The following sequence descriptions and sequences listings attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825. The Sequence Descriptions contain the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IYUB standards described in *Nucleic Acids Research* 13:3021-3030 (1985) and in the *Biochemical Journal* 219 (No. 2):345-373 (1984) which are
25 herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

SEQ ID NO:1 is the primer used as P298.

SEQ ID NO:2 is the primer used as P299.

SEQ ID NO:3 is the primer used as P192.

30 SEQ ID NO:4 is the primer used as P194.

SEQ ID NO:5 is the primer used as P198.

SEQ ID NO:6 is the primer used as P227.

SEQ ID NO:7 is the primer used as P228.

35 SEQ ID NO:8 is the primer used as PH785 for PCR amplification of the Arabidopsis Apetala 3 (AP3) promoter.

SEQ ID NO:9 is the primer used as PH786 for PCR amplification of the Arabidopsis Apetala 3 (AP3) promoter.

SEQ ID NO:10 is the primer used as PH783 for PCR amplification of the BCP1 promoter.

SEQ ID NO:11 is the primer used as PH784 for PCR amplification of the BCP1 promoter.

5 SEQ ID NO:12 is the primer used as PH788 for PCR amplification of the Arabidopsis Erecta (ER) promoter.

SEQ ID NO:13 is the primer used as PH790 for PCR amplification of the Arabidopsis Erecta (ER) promoter.

10 SEQ ID NO:14 is the primer used as PH795 for PCR amplification of the TA29 promoter from plasmid pTZALG.

SEQ ID NO:15 is the primer used as PH815 for PCR amplification of the TA29 promoter from plasmid pTZALG.

SEQ ID NO:16 is the primer used as P321 for PCR amplification of the Arabidopsis Pistilata (PI) promoter.

15 SEQ ID NO:17 is the primer used as P322 for PCR amplification of the Arabidopsis Pistilata (PI) promoter.

SEQ ID NO:18 is the primer used as PH806 for PCR amplification of the Arabidopsis Heat Shock (HSP) promoter 18.2.

20 SEQ ID NO:19 is the primer used as PH807 for PCR amplification of the Arabidopsis Heat Shock (HSP) promoter 18.2.

SEQ ID NO:20 is the primer used as P355 for PCR amplification of the Arabidopsis Apetala 1 (AP1) promoter.

SEQ ID NO:21 is the primer used as P356 for PCR amplification of the Arabidopsis Apetala 1 (AP1) promoter.

25 SEQ ID NO:22 is the primer used as P353 for PCR amplification of the Arabidopsis Agamous (AG) promoter.

SEQ ID NO:23 is the primer used as P354 for PCR amplification of the Arabidopsis Agamous (AG) promoter.

DETAILED DESCRIPTION OF THE INVENTION

30 The present invention provides constructs and methods for the conditional or regulated expression or excision of transgenes in plants by employing one or more site-specific recombinase systems and transgenes under the control of a variety of constitutive, inducible, tissue specific or development-specific promoters. Regulated expression of genetic traits is useful in plant breeding and
35 agronomic applications. Additionally transgene removal or excision from germline (pollen and/or seed) can be used not only for its containment in the production field but also for obtaining marker-free transgenics. Persistence of transformation marker, which is required to identify the rare transformed cells, in

transgenic plants may be undesirable because of regulatory concerns/requirements, its detrimental effect on transgenic plants, and/or it prevents recurrent use of the marker for stacking trait transgenes.

5 The following terms and definitions shall be used to fully understand the specification and claims.

“Gene” refers to a nucleic acid fragment that expresses mRNA, functional RNA, or specific protein, including regulatory sequences. The term “native gene” refers to gene as found in nature. The term “chimeric gene” refers to any gene that contains 1) DNA sequences, including regulatory and coding sequences, that are
10 not found together in nature, or 2) sequences encoding parts of proteins not naturally adjoined, or 3) parts of promoters that are not naturally adjoined. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or comprise regulatory sequences and coding sequences derived from the same source, but arranged in a
15 manner different from that found in nature. A “transgene” refers to a gene that has been introduced into the genome by transformation and is stably maintained. Transgenes may include, for example, genes that are either heterologous or homologous to the genes of a particular plant to be transformed. Additionally, transgenes may comprise native genes inserted into a non-native organism, or
20 chimeric genes. The term “endogenous gene” refers to a native gene in its natural location in the genome of an organism.

“Coding sequence” refers to a DNA or RNA sequence that codes for a specific amino acid sequence and excludes the non-coding sequences. The terms “open reading frame” and “ORF” refer to the amino acid sequence encoded
25 between translation initiation and termination codons of a coding sequence. The terms “initiation codon” and “termination codon” refer to a unit of three adjacent nucleotides (‘codon’) in a coding sequence that specifies initiation and chain termination, respectively, of protein synthesis (mRNA translation).

“Regulatory sequences” and “suitable regulatory sequences” each refer to
30 nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences include enhancers, promoters, translation leader sequences, introns, and polyadenylation signal sequences. They include
35 natural and synthetic sequences as well as sequences which may be a combination of synthetic and natural sequences. As is noted above, the term “suitable regulatory sequences” is not limited to promoters, however, some suitable regulatory sequences useful in the present invention will include, but are not

limited to constitutive plant promoters, plant tissue-specific promoters, plant developmental stage-specific promoters, inducible plant promoters and viral promoters.

5 The “3’ region” means the 3’ non-coding regulatory sequences located downstream of a coding sequence. This DNA can influence the transcription, RNA processing or stability, or translation of the associated coding sequence (e.g. for a recombinase, a transgene, etc.).

“Promoter” refers to a nucleotide sequence, usually upstream (5’) to its coding sequence, which controls the expression of the coding sequence by providing the recognition for RNA polymerase and other factors required for proper transcription. “Promoter” includes a minimal promoter that is a short DNA sequence comprised of a TATA- box and other sequences that serve to specify the site of transcription initiation, to which regulatory elements are added for control of expression. “Promoter” also refers to a nucleotide sequence that includes a
10 minimal promoter plus regulatory elements that is capable of controlling the expression of a coding sequence or functional RNA. This type of promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a DNA sequence which can stimulate promoter activity and may be an innate element of
15 the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. It is capable of operating in both orientations (normal or flipped), and is capable of functioning even when moved either upstream or downstream from the promoter. Both enhancers and other upstream promoter elements bind sequence-specific DNA-binding proteins that mediate their effects.
20 Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even be comprised of synthetic DNA segments. A promoter may also contain DNA sequences that are involved in the binding of protein factors which control the effectiveness of transcription initiation in response to physiological or
25 developmental conditions.
30

“Conditionally activating” refers to activating a transgene that is normally not expressed. In context of this invention it refers to expression of recombinase R1 either by a cross or, if it is inducible, also by an inducer.

“Constitutive expression” refers to expression using a constitutive or
35 regulated promoter. “Conditional” and “regulated expression” refer to expression controlled by regulated promoter. “Transient” expression in context of this invention refers to expression only in specific developmental stages or tissue in one or two generation.

- “Constitutive promoter” refers to promoters that direct gene expression in all tissues and at all times. “Regulated promoter” refers to promoters that direct gene expression not constitutively but in a temporally- and/or spatially-regulated manner and include tissue-specific, developmental stage-specific, and inducible promoters. It includes natural and synthetic sequences as well as sequences which may be a combination of synthetic and natural sequences. Different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro et al., *Biochemistry of Plants* 15:1-82, 1989. Since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity. Typical regulated promoters useful in plants include but are not limited to safener-inducible promoters, promoters derived from the tetracycline-inducible system, promoters derived from salicylate-inducible systems, promoters derived from alcohol-inducible systems, promoters derived from glucocorticoid-inducible system, promoters derived from pathogen-inducible systems, and promoters derived from ecdysome-inducible systems.
- “Tissue-specific promoter” refers to regulated promoters that are not expressed in all plant cells but only in one or more cell types in specific organs (such as leaves, shoot apical meristem, flower, or seeds), specific tissues (such as embryo or cotyledon), or specific cell types (such as leaf parenchyma, pollen, egg cell, microspore- or megaspore mother cells, or seed storage cells). These also include “developmental-state specific promoters” that are temporally regulated, such as in early or late embryogenesis, during fruit ripening in developing seeds or fruit, in fully differentiated leaf, or at the onset of senescence. It is understood that the developmental specificity of the activation of a promoter and, hence, of the expression of the coding sequence under its control, in a transgene may be altered with respect to its endogenous expression. For example, when a transgene under the control of a floral promoter is transformed into a plant, even when it is the same species from which the promoter was isolated, the expression specificity of the transgene will vary in different transgenic lines due to its insertion in different locations of the chromosomes.
- “Non-specific expression” refers to constitutive expression or low level, basal (‘leaky’) expression in nondesired cells, tissues, or generation.

“Inducible promoter” refers to those regulated promoters that can be turned on in one or more cell types by a stimulus external to the plant, such as a chemical, light, hormone, stress, or a pathogen.

5 “Promoter activation” means that the promoter has become activated (or turned “on”) so that it functions to drive the expression of a downstream genetic element. Constitutive promoters are continually activated. A regulated promoter may be activated by virtue of its responsiveness to various external stimuli (inducible promoter), or developmental signals during plant growth and differentiation, such as tissue specificity (floral specific, anther specific, pollen
10 specific seed specific etc) and development-stage specificity (vegetative or floral shoot apical meristem-specific, male germline specific, female germline specific etc).

“Operably-linked” refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other.
15 For example, a promoter is operably-linked with a coding sequence or functional RNA when it is capable of affecting the expression of that coding sequence or functional RNA (i.e., that the coding sequence or functional RNA is under the transcriptional control of the promoter). Coding sequences can be operably-linked to regulatory sequences in sense or antisense orientation. “Unlinked” means that
20 the associated genetic elements are not closely associated with one another and function of one does not affect the other.

“Expression” refers to the transcription and stable accumulation of sense (mRNA) or functional RNA. Expression may also refer to the production of protein. “Overexpression” refers to the level of expression in transgenic
25 organisms that exceeds levels of expression in normal or untransformed organisms.

“Altered levels” refers to the level of expression in transgenic organisms that differs from that of normal or untransformed organisms.

30 The term “altered plant trait” means any phenotypic or genotypic change in a transgenic plant relative to the wildtype or non-transgenic plant host.

“Transcription Stop Fragment” refers to nucleotide sequences that contain one or more regulatory signals, such as polyadenylation signal sequences, capable of terminating transcription. Examples include the 3' non-regulatory regions of genes encoding nopaline synthase and the small subunit of ribulose biphosphate
35 carboxylase.

“Translation Stop Fragment” refers to nucleotide sequences that contain one or more regulatory signals, such as one or more termination codons in all three frames, capable of terminating translation. Insertion of translation stop

fragment adjacent to or near the initiation codon at the 5' end of the coding sequence will result in no translation or improper translation. Excision of the translation stop fragment by site-specific recombination will leave a site-specific sequence in the coding sequence that does not interfere with proper translation using the initiation codon.

“Stop fragment” or “Blocking fragment” refers to a DNA fragment that is flanked by site-specific sequences that can block the transcription and/or the proper translation of a coding sequence resulting in an inactive transgene. When the blocking fragment contains polyadenylation signal sequences and other sequences encoding regulatory signals capable of terminating transcription it can block the transcription of a coding sequence when placed in the 5' non-translated region, i.e., between the transcription start site and the ORF. When inserted in the coding sequence a blocking fragment can block proper translation by disrupting its open reading frame. DNA rearrangement by site-specific recombination can restore transcription and/or proper translatability. For example, excision of the blocking fragment by site-specific recombination leaves behind a site-specific sequence that allows transcription and/or proper translatability. A Transcription or Translational Stop Fragment will be considered a blocking fragment. A “stop fragment” can also block transcription by disrupting the gene in the non-transcribed region, for example by its presence and/or orientation in promoter sequences either between the upstream promoter elements and the “TATA” box or between the TATA box and the transcription start site.

This process of excision of the stop fragment or blocking fragment will be referred to herein as “unblocking”. When the blocking fragment is removed from the DNA by site-specific recombination, it will be appreciated by one skilled in the art that a site-specific sequence remains which can be transcribed and/or translated properly.

“Priming” or “enabling” refers to the removal of blocking sequences upstream of a promoter and/or gene, such that the gene can become activated in response to the appropriate environmental cue, stage of development, or presence in a specific tissue/ cell type. When a genetic element is enabled or primed by the removal of a blocking fragment, the promoter element may or may not be free to drive the expression of the downstream element. For example, a genetic construct comprising an inducible promoter separated by a stop fragment from a downstream gene to which it is operably linked, will be primed by the removal of the stop fragment, however will not express the downstream element until it is activated or induced. Thus, activation of a blocked gene will require enabling it and activation of the promoter driving the gene.

“Recombinase element” refers to a DNA element comprising a promoter operably linked to a gene encoding a site-specific recombinase, or to other genetic elements flanked by site-specific recombinase sequences. Recombinase elements of the present invention may optionally contain blocking or stop fragments to allow for more highly regulated gene expression.

The term “floxed” will refer to the flanking of a genetic element with tandemly (i.e., directly, repeated) site-specific sequences. The floxed element may be recombinase element or any other genetic element.

The term “recombinase” refers to enzyme(s) that carry out site-specific recombination that alters the DNA structure and includes transposases, lambda integration/excision enzymes, as well as site-specific recombinases. Well-known examples of recombinases can be found in *Cre-lox*, *FLP/FRT*, *R/RS*, *Gin/gix*, a pSR1 system, a *cer* system, and a *fim* system (for example, N. L. Craig, *Annu Rev. Genet.*, vol. 22, p.17, 1988; Odell et al., Use of site-specific recombination systems in plants. *Homologous Recomb. Gene Silencing Plants* (1994), 219-70. Editor(s): Paszkowski, Jerzy. Publisher: Kluwer, Dordrecht, Germany). Additionally, site-specific recombination systems have been identified in microorganisms such as phage, bacterium (e.g., *E. coli*), yeast and the like. This includes the *E. coli* lambda att P system (Zubko et al. (2000) *Nature Biotechnology* 18:442) integration and excision and the *Streptomyces* phage C31 integrase (Groth et al. (2000) *Proc. Natl Acad. Sci. USA* 97:5995). When the site-specific recombination system separated from these microorganisms with the use of a *Cre/lox* system derived from P1 phage (WO 93/01283) is introduced into organisms (including plants) different from the organism from which this system had been derived, it behaves in the same way as in the original organism. The site-specific recombination system of yeast (*Zygosaccharomyces rouxii*) (pSR1 system (H. Matsuzaki et al., *J. Bacteriology*, vol.172, p.610, 1990)) can also be used in accordance with the present invention. This pSR1 system also maintains its inherent function in higher plants (H. Onouchi et al., *Nucleic Acid Res.*, vol.19, p.6373, 1991).

“Recombinase site” or “site-specific recombinase sequence” means a DNA sequence that a recombinase will recognize and bind to. It will be appreciated that this may be a wild type or mutant recombinase site, as long as functionality is maintained and the recombinase enzyme may still recognize the site, bind to the DNA sequence, and catalyze the recombination between two adjacent recombinase sites.

“Trait removal construct” is defined herein as any assembly of DNA constructs comprising (at least): a recombinase element, site specific recombinase

sites responsive to the recombinase enzyme, and a transgene. The trait removal construct becomes functional upon the introduction of each element into a single organism, either by co-transformation, sequential transformations, or genetic crosses after single transformation in a plant.

5 “Production tissue” refers to mature, harvestable tissue consisting of non-dividing, terminally-differentiated cells. It excludes young, growing tissue consisting of germline, meristematic, and not-fully-differentiated cells.

 “Germline” refers to cells that are destined to be gametes. Thus, the genetic material of germline cells is heritable.

10 “Common germline” refers to all germline cells prior to their differentiation into the male and female germline cells and, thus, includes the germline cells of developing embryo, vegetative SAM, floral SAM, and flower. Thus, site-specific excision in common germline results in excision from both male and female gametes.

15 “Male germline” refers to cells of the sporophyte (anther primordia, anther, microspore mother cells) or gametophyte (microspore, pollen) that are destined to be male gametes (sperms) and the male gametes themselves.

 “Female germline” refers to cells of the sporophyte (pistil primordia, pistil, ovule, macrospore mother cells) or gametophyte (macrospore, egg cell) that are
20 destined to be female gametes or the female gametes themselves.

 “Somatic” cells are all other cells in the organism that are not germline cells.

 “Common germline” promoter refers to a promoter that is activated in germline cells prior to their differentiation into the male and female germlines. It
25 also refers to a promoter that is activated in both male and female germline cells and to a set of promoters, one specific to the male germline and the other to the female germline. Thus, site-specific excision in common germline results in excision from both male and female gametes.

 “Floral common germline” promoter refers to a promoter of flower or
30 flower primordia genes whose expression occurs in “common germlines”. It does not include male germline or female germline promoters, which are also expressed in the flower.

 “Male germline” promoter refers to a promoter whose expression occurs in male not female germline in the flower.

35 “Female germline” promoter refers to a promoter whose expression occurs in female not male germline in the flower.

" Flower" or "floral" -specific promoter refers to a promoter whose expression occurs in the flower or flower primordia. They include floral common germline, male germline, and female germline promoters.

5 "Genetically linked" refers to physical linkage of transgene such that they co-segregate in progeny.

"Genetically unlinked" refers to the lack of physical linkage of transgene such that they do not co-segregate in progeny.

"Seed-specific promoter refers to promoter that is expressed only in the seed.

10 "Plant developmental stage-specific promoter" refers to a promoter that is expressed not constitutively but at specific plant developmental stage or stages. Plant development goes through different stages and in context of this invention the germline goes through different developmental stages starting, say, from fertilization through development of embryo, vegetative shoot apical meristem, floral shoot
15 apical meristem, anther and pistil primordia, anther and pistil, micro- and macrospore mother cells, and macrospore (egg) and microspore (pollen).

"Vegetative shoot apical meristem" refers to the cells found in the shoot apex of vegetative shoots that give rise to leaves and shoots.

20 "Floral shoot apical meristem" refers to the cells found in the shoot apex of floral meristem shoots that give rise to flowers and inflorescences.

"Morphological trait" refers to traits of morphology, such as shoots, roots, calli, tumors, flowers, or leaves

"Tumorigenic" genes refers to genes that cause plant tumors, such as the T-DNA genes of *Agrobacterium tumefaciens*.

25 "Root inducing" genes refers to genes, such as rol A,B, and C genes of *Agrobacterium rhizogenes*, that cause root formation.

The term "Lethal gene that block development" refers to a gene that express a toxin, such as alpha chain of diphtheria toxin, barnase, or interferes with normal plant development, such as rolB (Roder et al. (1994) *Mol. Gen. Genet.* 30 243:32). It also includes transgenes that silence or co-suppress plant genes required for normal development (see Chuang CF, Meyerowitz EM (2000) Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A*.97:4985-90)

35 "Conditional and transient expression" refers to expression of a trait gene only in the selected generation or two. In context of this invention, expression is triggered in first generation and upon useful trait expression, the trait gene is removed from the germline.

“Restoring male fertility” refers to removing a male sterility gene by excision or counteracting it by the expression of a suppressor that can be either a transgene for silencing the male sterility gene or a transgene that encode a protein that cointeracts the male sterility factor, such as barstar against barnase ichiels, F., and M. Williams, 1998, Improved barstar gene and its expression in male-sterile barnase-producing plants to restore fertility, PCT Int. Appl., WO, (Plant Genetic Systems, N.V., Belg.; Michiels, Frank; Williams, Mark)., p. 54 pp.

“Synthetic anther promoter” refers to G9/SGB6 hybrid promoter (U.S. Pat No. 5470359; 5837850).

“Pollen-specific: promoters refers to promoters that are only expressed in pollen, such as LAT52 Twell et al. (1998) *Trends in Plant Sciences* 3:305.

“Sterility” means the inability of a plant to reproduce sexually or to set seeds. In cross pollinating plants it will include inability to form a functional pollen as well as inability to set seeds. Sterility can also result from abnormal plant development that prevents flower formation. A “sterility gene” refers to any gene that conveys sterility to the plant and includes genes that prevent both pollen formation and seed set such as by preventing flower formation. “Male sterility” means the inability of a plant to produce functional pollen as a consequence of mechanical or hand detasseling, incorporation of genetic male sterility, or by other means. “Female sterility” means the inability of a plant to set seeds.

Male sterility result from the expression of the male sterility gene in diploid cells (sporophytic), haploid (gametophytic) cells, or both types of cells. Gametophytic male sterility refers to a gene whose expression occurs only in the pollen, for example expression of barnase or the alpha chain of diphtheria toxin under pollen-specific promoter.

“Gametophytic male sterility” refers to sterility that is expressed only in pollen.

“Gametophytic male fertility restorer” a transgene that restores fertility by removing or suppressing the male sterility gene.

“Barnase” and “Barstar” refer to a RNase toxin and its antidote as described by Michiels, F., and M. Williams, 1998, Improved barstar gene and its expression in male-sterile barnase-producing plants to restore fertility, PCT Int. Appl., WO, (Plant Genetic Systems, N.V., Belg.; Michiels, Frank; Williams, Mark)., p. 54 pp.

“Activating transgene” refers to expression of a transgene. In context of this invention, it refers to both enabling a blocked gene or enabling a blocked gene followed by activation of its promoter.

“Transformation” refers to the transfer of a foreign gene into the genome of a host organism. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al.(1987) *Meth. Enzymol.* 143:277) and particle-accelerated or “gene gun” transformation technology (Klein et al.(1987) *Nature* (London) 327:70-73; U.S. Patent No. 4,945,050). The terms “transformed”, “transformant” and “transgenic” refer to plants or calli that have been through the transformation process and contain a foreign gene integrated into their chromosome. The term “untransformed” refers to normal plants that have not been through the transformation process.

10 “Stably transformed” refers to cells that have been selected and regenerated on a selection media following transformation.

“Genetically stable” and “heritable” refer to chromosomally-integrated genetic elements that are stably maintained in the plant and stably inherited by progeny through successive generations.

15 “Wild-type” refers to the normal gene, virus, or organism found in nature without any known mutation.

“Genome” refers to the complete genetic material of an organism.

“Genetic trait” means a genetically determined characteristic or condition, which is transmitted from one generation to another. “Homozygous” state means a genetic condition existing when identical alleles reside at corresponding loci on homologous chromosomes. In contrast, “heterozygous” state means a genetic condition existing when different alleles reside at corresponding loci on homologous chromosomes. A “hybrid” refers to any offspring of a cross between two genetically unlike individuals. “Inbred” or “inbred lines” or “inbred plants” means a substantially homozygous individual or variety. This results by the continued mating of closely related individuals, especially to preserve desirable traits in a stock.

The term “ortholog” or “orthologous genes” refer to genes related by common phylogenetic descent. Orthologous genes are those genes from one species which corresponds to a gene in another species that is related via a common ancestral species (a homologous gene), but which has evolved to become different from the gene of the other species.

30 “Selfing” or “self fertilization” refers to the transfer of pollen from an anther of one plant to the stigma (a flower) of that same said plant. Selfing of a hybrid (F1) results in a second generation of plants (F2)

35 “TopCross® high oil corn seed method” refers to a commercial method of making hybrid corn seeds in the field, as described, for example, in U.S. Patent

No. 5,704,160. A "TopCross® pollinator refers to the parent line of the cross that provides the pollen.

The term "sporophyte" means the diploid phase or cells of a plant.

5 The term "gametophyte" means haploid phase or cells of a plant. This is the stage in a plant's life cycle between meiosis and fertilization. The male gametophyte includes the haploid phase or cells of the pollen and the female gametophyte includes the haploid phase or cells of the egg cell.

10 The term "plant life cycle" means a complete sequence of developmental events in the life of a plant, such as from fertilization to the next fertilization or from flowering in one generation to the next.

"Primary transformant" and "T₀ generation" refer to transgenic plants that are of the same genetic generation as the tissue which was initially transformed (i.e., not having gone through meiosis and fertilization since transformation).

15 "Secondary transformants" and the "T₁, T₂, T₃, etc. generations" refer to transgenic plants derived from primary transformants through one or more meiotic and fertilization cycles. They may be derived by self-fertilization of primary or secondary transformants or crosses of primary or secondary transformants with other transformed or untransformed plants.

20 The term "generation" means a plant life cycle starting from fertilization to fertilization. In the context of this invention, a "first generation plant" is defined as the plant in which the first recombination event occurs, while a "second generation plant" is the progeny seed and plant of the first generation plant.

25 A "restorer gene" is defined herein as a gene whose expression restores fertility. A gametophyte fertility restorer gene refers to genes whose expression overcome gametophytic sterility, such as a barstar gene (Michiels, F., and M. Williams, 1998, Improved barstar gene and its expression in male-sterile barnase-producing plants to restore fertility, PCT Int. Appl., WO, (Plant Genetic Systems, N.V., Belg.; Michiels, Frank; Williams, Mark), P. 54 pp.).

The following abbreviations will be used herein:

30 "STP" is the abbreviation for stop fragment or blocking fragment.

"AP1" is the abbreviation for *Arabidopsis Apetala 1* gene (Liljegren et al, (1999) *The Plant Cell* 11:1007.)

"AG" is the abbreviation for *Arabidopsis agamous* gene (Yanofsky et al. 91990) *Nature* 346:35).

35 "PI" is the abbreviation for *Arabidopsis Pistillata* gene (Goto and Meyerowitz (1994) *Genes and development* 8:1548-1560).

"LFY" is the abbreviation for *Arabidopsis Leafy* gene (Nilsson et al. (1998) *The Plant Journal* 15:799-804).

“ANT” is the abbreviation for *Arabidopsis Aintegumenta* gene (Mizukami and Fischer (2000) *Proc. Natl. Acad. Sci. USA* 97:942-947).

“CLV3” is the abbreviation for *Arabidopsis Clavata 3* gene (Fletcher et al. (1999) *Science* 283:1911-1914).

5 “WUS” is the abbreviation for *Arabidopsis Wushel* gene (Mayer et al. (1998) *Cell* 95(6), 805-815).

“STM” is the abbreviation for *Arabidopsis Shoot Meristemless* gene (Long et al. (1996) *Nature* 379:66-69).

10 “rol C” is the abbreviation for the root locus C gene that causes root formation (see Constantino et al. (1994) *Genetics* 94:203).

“IPT” is the abbreviation for the *isopentyl transferase* gene (Ebumina et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2117-2121).

“KNAT” is the abbreviation for the a Knox class of genes (see Reiser et al. (2000) *Plant Mol. Biol.* 42:151-166).

15 “Lec1” is the abbreviation for *Arabidopsis Leafy Cotyledon 1* (Lotan et al, 1998. *Cell* 93: 1195-1205) gene.

“OSHI ” is the abbreviation for a rice homeobox gene (Sentoku et al. (2000) *Developmental Biology* 220:358-364).

20 “Kn1” is the abbreviation for corn Knotted 1 gene (Vollbrecht, E. et al. (1991) *Nature* 350:241-243).

“Gmf” means gametophytic male fertile.

“Gms” means gametophytic male sterile.

“TG” is the abbreviation for transgene.

25 “SAM” is the abbreviation for shoot apical meristem. SAM can be vegetative or floral.

“SSR” is the abbreviation for site-specific recombination.

“SAP is the abbreviation for Synthetic anther promoter as described in U.S. Pat No. 5470359; 5837850.

“HSP” is the abbreviation for heat shock protein.

30 The present invention provides constructs and methods for the conditional or regulated expression or excision of transgenes in plants by employing one or more site-specific recombinase systems and transgenes under the control of a variety of constitutive, inducible, tissue specific or development-specific promoters. The invention makes use of a variety of constructs referred to herein
35 as recombinase elements. Each recombinase element comprises at least one promoter functional in a plant cell. Additionally the recombinase element may comprise a number of other components including sequence encoding a site specific recombinase or a transgene or may comprise a stop fragment or site-

specific sequences responsive to a recombinase. The recombinase elements are introduced into plants in a variety of combinations so as to provide for the conditional expression or excision of specific genetic traits encoded by the transgenes. By matching promoters, responsive to various inducers, plant tissues or plant developmental states with the recombinase systems, stop fragments and transgenes, virtually any trait may be expressed or excised at any plant development stage or in any plant generation.

The present system of gene expression has several advantages over current methods. For example, regulated expression under some current promoters may result in non-specific or "leaky" expression.

Promoters

The present invention makes use of a variety of plant promoters to drive the expression of either a recombinase or a transgene as part of the recombinase elements of the invention.

Regulated expression of transgene expression is possible by placing the transgene or recombinase system under the control of promoters that may be conditionally regulated. Any promoter functional in a plant will be suitable including but not limited to constitutive plant promoters, plant tissue-specific promoters, plant development-specific promoters, inducible plant promoters, viral promoters, male germline-specific promoters, female germline-specific promoters, flower-specific promoters, and vegetative shoot apical meristem-specific promoters.

Several tissue-specific regulated genes and/or promoters have been reported in plants. These include genes encoding the seed storage proteins (such as napin, cruciferin, beta-conglycinin, and phaseolin), zein or oil body proteins (such as oleosin), or genes involved in fatty acid biosynthesis (including acyl carrier protein, stearyl-ACP desaturase, and fatty acid desaturases (fad 2-1)), and other genes expressed during embryo development (such as Bce4, see, for example, EP 255378 and Kridl et al., *Seed Science Research* (1991) 1:209-219). Particularly useful for seed-specific expression is the pea vicilin promoter (Czako et al., *Mol. Gen. Genet.* (1992), 235(1), 33-40). Other useful promoters for expression in mature leaves are those that are switched on at the onset of senescence, such as the SAG promoter from *Arabidopsis* (Gan et al., "Inhibition of leaf senescence by autoregulated production of cytokinin", *Science* (Washington, D.C.) (1995), 270 (5244), 1986-8).

A class of fruit-specific promoters expressed at or during anthesis through fruit development, at least until the beginning of ripening, is discussed in U.S. 4,943,674, the disclosure of which is hereby incorporated by reference.

cDNA clones that are preferentially expressed in cotton fiber have been isolated (John et al., Gene expression in cotton (*Gossypium hirsutum* L.) fiber: cloning of the mRNAs, *Proc. Natl. Acad. Sci. U.S.A.* (1992), 89 (13), 5769-73). cDNA clones from tomato displaying differential expression during fruit development
5 have been isolated and characterized (Mansson et al., *Mol. Gen. Genet.* (1985) 200:356-361; Slater et al., *Plant Mol. Biol.* (1985) 5:137-147). The promoter for polygalacturonase gene is active in fruit ripening. The polygalacturonase gene is described in U.S. Patent No. 4,535,060 (issued August 13, 1985), U.S. Patent No. 4,769,061 (issued September 6, 1988), U.S. Patent No. 4,801,590 (issued
10 January 31, 1989) and U.S. Patent No. 5,107,065 (issued April 21, 1992), which disclosures are incorporated herein by reference.

Mature plastid mRNA for psbA (one of the components of photosystem II) reaches its highest level late in fruit development, in contrast to plastid mRNAs for other components of photosystem I and II which decline to nondetectable
15 levels in chromoplasts after the onset of ripening (Piechulla et al., *Plant Mol. Biol.* (1986) 7:367-376). Recently, cDNA clones representing genes apparently involved in tomato pollen (McCormick et al., *Tomato Biotechnology* (1987) Alan R. Liss, Inc., New York) and pistil (Gasser et al., *Plant Cell* (1989), 1:15-24) interactions have also been isolated and characterized.

20 Other examples of tissue-specific promoters include those that direct expression in leaf cells following damage to the leaf (for example, from chewing insects), in tubers (for example, patatin gene promoter), and in fiber cells (an example of a developmentally-regulated fiber cell protein is E6 (John et al., Gene expression in cotton (*Gossypium hirsutum* L.) fiber: cloning of the mRNAs, *Proc.*
25 *Natl. Acad. Sci. U.S.A.* (1992), 89(13), 5769-73)). The E6 gene is most active in fiber, although low levels of transcripts are found in leaf, ovule and flower.

The tissue-specificity of some "tissue-specific" promoters may not be absolute and may be tested by one skilled in the art using the *diphtheria* toxin sequence. One can also achieve tissue-specific expression with "leaky"
30 expression by a combination of different tissue-specific promoters (Beals et al., (1997) *Plant Cell*, vol 9, 1527-1545). Other tissue-specific promoters can be isolated by one skilled in the art (see U.S. 5,589,379).

Germline specific promoters, responsive to male, female, or both male-female specific cell lineages are also useful in the present invention. For instance
35 transgenes can be expressed or removed from pollen by site-specific recombinase expression under the control of male germline-specific genes in anther primordia genes, such as *Arabidopsis* *Apetalla* 3 and *Pistilata* (PI) or their orthologs from other plant species, in sporophytic anther tissue (eg. Bcp I and TA29 promoters, or

gametophytic pollen. Similarly, transgenes can be expressed or removed from ovules by site-specific recombinase expression under the control of female germline-specific genes in ovule primordia. Transgenes can be expressed or removed from both male- and female-specific germlines by expression of site-specific recombinase gene under the control of promoter for genes common to both male and female lineages in flower, such as *Arabidopsis agamous* gene or its orthologs in other species, in floral meristem, such as *Arabidopsis Apetala 1*, *Leafy*, and *Erecta* or their orthologs from other species, and in vegetative shoot apical meristem, such as *Arabidopsis WUSCHEL* (WUS) and *SHOOT*
5 *MERISTEMLESS* (STM) or their orthologs from other species. Promoters of shoot apical meristem are especially useful for removing or expressing transformation marker genes early in tissue-culture following selection or in planta following a transformation phenotype.

Similarly, several inducible promoters ("gene switches") have been reported. Many are described in the review by Gatz (*Current Opinion in Biotechnology*, 1996, vol. 7, 168-172; Gatz, C. Chemical control of gene expression, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* (1997), 48, 89-108). These include tetracycline repressor system, *Lac* repressor system, copper-inducible systems, salicylate-inducible systems (such as the PR1a system), glucocorticoid-
15 (Aoyama T. et al., *N-H Plant Journal* (1997) vol 11:605-612) and ecdysone-inducible systems. Also, included are the benzene sulphonamide- (U.S. 5,364,780) and alcohol- (WO 97/06269 and WO 97/06268)-inducible systems and glutathione S-transferase promoters. Other studies have focused on genes inducibly regulated in response to environmental stress or stimuli such as increased salinity, drought,
20 pathogen, and wounding. (Graham et al., *J. Biol. Chem.* (1985) 260:6555-6560; Graham et al., *J. Biol. Chem.* (1985) 260:6561-6554) (Smith et al., *Planta* (1986) 168:94-100). Accumulation of a metallocarboxypeptidase-inhibitor protein has been reported in leaves of wounded potato plants (Graham et al., *Biochem Biophys Res Comm* (1981) 101:1164-1170). Other plant genes have been reported to be
25 induced methyl jasmonate, elicitors, heat-shock, anerobic stress, or herbicide safeners.

Site-Specific Recombinase Systems

The present invention provides site-specific recombinase systems for use in the regulated expression or excision of transgenes. A site-specific
35 recombination system consists of two elements, (1) a recombination site (corresponding to the removable DNA element of the present invention) having a characteristic DNA sequence, and (2) an enzyme that binds to the DNA sequence specifically and catalyzes the recombination between DNA sequences if two or

more of the sequences exist (recombinase). When the two DNA sequences are oriented in the same direction at a given interval on the same DNA molecule, the region held by these DNA sequences is excised from the DNA molecule, such as a plasmid, chromosome or the like. When the two DNA sequences are oriented in opposite directions on the same DNA molecule, the region held by these DNA sequences is inverted.

Use of developmentally-regulated or chemically-induced promoters for conditional transgene expression is usually limited either by their insufficient strength in the "fully-on" stage or, more often, by their basal non-specific (i.e., "leaky" expression) in the 'off' stage, depending on the application.

One can increase both the level and specificity of conditional expression by putting the coding sequence of the gene of interest under the control of a strong constitutive or regulated promoter for expression in the production tissue in such a manner that the gene is transcriptionally inactive unless it undergoes a site-specific recombination through the conditional expression of the cognate site-specific recombinase. Thus, conditional expression of the gene of interest is now dependent on the conditional expression of the recombinase. In this manner, determinants for high-level expression and for specificity are separated and one can then focus on the basal non-specific (i.e., "leaky") expression of recombinase. The site-specific sequences and their cognate recombinase enzymes can be from any natural site-specific recombination systems. Well-known examples include *Cre-lox*, *FLP/FRT*, *R/RS*, *Gin/gix*, a *pSR1* system, a *cer* system, and a *flm* system (for example, N. L. Craig, *Annu Rev. Genet.*, vol.22, p.17, 1988; Odell et al., Use of site-specific recombination systems in plants. *Homologous Recomb. Gene Silencing Plants* (1994), 219-70. Editor(s): Paszkowski, Jerzy. Publisher: Kluwer, Dordrecht, Germany). Additionally, site-specific recombination systems have been identified in microorganisms such as phage, bacterium (e.g., *E. coli*), yeast and the like. When the site-specific recombination system separated from these microorganisms with the use of a *Cre/lox* system derived from P1 phage (WO 93/01283) is introduced into organisms (including plants) different from the organism from which this system had been derived, it behaves in the same way as in the original organism. The site-specific recombination system of yeast (*Zygosaccharomyces rouxii*) (*pSR1* system (H. Matsuzaki et al., *J. Bacteriology*, vol.172, p.610, 1990)) can also be used in accordance with the present invention. This *pSR1* system also maintains its inherent function in higher plants (H. Onouchi et al., *Nucleic Acid Res.*, vol.19, p.6373, 1991).

Since the levels of the recombinase enzyme required are not expected to be high, several "specific" promoters can be used that may otherwise be too weak to

express the gene of interest. Furthermore, since site-specific recombination depends on a threshold level of the recombinase, there may be a tolerance for leaky transcription that results in sub-threshold levels of recombinase.

Furthermore, increased “tissue-selectivity” to available regulated promoters is provided by decreasing the efficiency of wild-type Cre-mediated recombination, raising the threshold of recombinase required by using either a mutant site for site-specific recombination and/or a mutant recombinase that are not proficient in recombination. Such mutants are well known, at least for the Cre-lox system. The applicants have shown that when using safener-inducible Cre expression to activate the expression of a transgene (35S:luciferase), the use of a mutant lox site (lox72) and a wild type lox P site in Cre-mediated activation of the transgene reduces the basal activity of the promoter compared to using both wild type lox P sites.

The non-specificity of recombinase expression can be further reduced (i.e., its expression specificity further increased) by other post-transcriptional approaches including: 1) using a chimeric recombinase gene that is poorly translated (such as having a non-ideal context sequence around the initiation codon following Kozak’s rule or having additional short ORFs in the 5' untranslated region as in yeast GCN4 mRNA, or having 3' UTR sequences that makes mRNA unstable as described by Pamela Green (Department of Biochemistry, Michigan State University, East Lansing, MI 48824-1312, U.S.A.); 2) using a mutant recombinase that has less cellular stability (i.e., shorter half-life). Such mutants could be made by adding PEST sequences (Sekhar et al., *Jrl. Receptor Signal Transduction Res.* 18 (2-3), 113-132 (1998)).

Once a system is developed in a given crop, it can be easily adapted for conditional expression of a variety of target trait genes.

Transgenes

Transgenes of the present invention will be those that convey a desirable phenotype on the transformed plant, or those that encode markers useful in breeding. Particularly useful transgenes will include, but not be limited to genes conveying specific phenotype on a plant or plant cell, genes encoding a transformation marker, genes encoding a morphological trait, genes conveying sterility, and hormone biosynthetic genes.

Transgenes can encode functional RNAs or foreign proteins. Foreign proteins will typically encode proteins that may be foreign to plant hosts. Such foreign proteins will include, for example, enzymes for primary or secondary metabolism in plants, proteins that confer disease or herbicide resistance, commercially useful non-plant enzymes, and proteins with desired properties

useful in animal feed or human food. Additionally, foreign proteins encoded by the transgenes will include seed storage proteins with improved nutritional properties, such as the high-sulfur 10 kD corn seed protein or high-sulfur zein proteins. Additional examples of a transgene suitable for use in the present invention include genes for disease resistance (e.g., gene for endotoxin of *Bacillus thuringiensis*, WO 92/20802)), herbicide resistance (mutant acetolactate synthase gene, WO 92/08794)), seed storage protein (e.g., glutelin gene, WO 93/18643)), fatty acid synthesis (e.g., acyl-ACP thioesterase gene, WO 92/20236)), cell wall hydrolysis (e.g., polygalacturonase gene (D. Grierson et al., *Nucl. Acids Res.*, vol.14, p.8595, 1986)), anthocyanin biosynthesis (e.g., chalcone synthase gene (H. J. Reif et al., *Mol. Gen. Genet.*, vol.199, p.208, 1985)), ethylene biosynthesis (e.g., ACC oxidase gene (A. Slater et al., *Plant Mol. Biol.*, vol.5, p.137, 1985)), active oxygen-scavenging system (e.g., glutathione reductase gene (S. Greer & R. N. Perham, *Biochemistry*, vol.25, p.2736, 1986)), and lignin biosynthesis (e.g., phenylalanine ammonia-lyase gene, cinnamyl alcohol dehydrogenase gene, o-methyltransferase gene, cinnamate 4-hydroxylase gene, 4-coumarate-CoA ligase gene, cinnamoyl CoA reductase gene (A. M. Boudet et al., *New Phytol.*, vol.129, p.203, 1995)).

Transgenes may function as transformation markers. Transformation markers include selectable genes, such as antibiotic or herbicide resistance genes, which are used to select transformed cells in tissue culture, non-destructive screenable reporters, such as green fluorescent and luciferase genes, or a morphological marker, such as "shooty", "rooty", or "tumorous" phenotype.

Additionally transgenes may encode proteins that affect plant morphology and thus may also be used as markers. Morphological transformation marker genes include cytokinin biosynthetic genes, such as the bacterial gene encoding isopentenyl transferase (IPT). IPT gene was proposed as a marker for transformation by Ebumina et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2117-2121), and Kunkel et al. *Nat. Biotechnol.* (1999), 17(9), 916-919. In the former case, the IPT gene was inserted inside a transposable element, whose excision following transformation resulted in the loss of the transposable element and the IPT gene. However, this method is inefficient (see Kunkel, *supra*), especially because of its low frequency of loss (1% or less). Kunkel, *supra*) proposed the use of an inducible IPT gene. However, this is also undesirable, since the bacterial IPT gene is not lost following transformation and that could be of concern from a regulatory point of view. Furthermore, it does not allow its use for retransformation for trait stacking. Thus, there is a need for an efficient regulated removal of morphological markers. Other morphological markers

include developmental genes that can induce ectopic shoots, such as *Arabidopsis* *STM*, *KNAT 1*, or *AINTEGUMANTA*, *Lec 1*, *Brassica* "Babyboom" gene, rice *OSH1* gene, or maize *Knotted (Kn1)* genes. Yet other morphological markers are the wild type T-DNA of *Ti* and *Ri* plasmids of *Agrobacterium* that induce tumors or hairy roots, respectively, or their constituent T-DNA genes for distinct morphological phenotypes, such as shooty (e.g., cytokinin biosynthesis gene) or rooty phenotype (e.g. *rol C* gene). Use of a morphological transformation marker to identify transformed tissue/organ and its subsequent removal (leaving behind the transgene of interest) restores normal morphology and development to transgenic tissues. This is especially useful for *in planta* transformation, where the morphological marker is used to obtain abnormal transgenic organs that are then corrected by site-specific recombination to form morphologically and developmentally normal transgenic plants without going through the time and labor intensive tissue culture methods for transformation.

15 Plant Hosts

The present invention additionally provides plant hosts for transformation with the present constructs. Moreover, the host plant for use in the present invention is not particularly limited. Examples of herbaceous plant used as the host plant include tobacco (*Tabacum*), tomato (*Lycopersicom*), sweet potato (*Ipomoea*), potato (*Solanum*), carrot (*Dacus*), lettuce (*Lactuca*), cauliflower (*Brassica*), cabbage (*Brassica*), oilseed rape (*Brassica*), sunflower (*Helianthus*), sugar beet (*Beta*), asparagus (*Asparagus*), banana (*Musa*), cotton (*Gossypium*), *arabidopsis* (*Arabidopsis*), alfalfa (*Medicago*), peas (*Pisum*), soybean (*Glycine*), rice (*Oryza*), corn (*Zea*), and rye (*Secale*). Examples of arboreous plant used as the host plant include poplar (*Populus*), eucalypti (*Eucalyptus*), acacia (*Acacia*), pear (*Pyrus*), apple (*Malus*), grape (*Vitis*), walnut (*Juglans*), plum (*Prunus*), rose (*Rosa*), and spruce (*Picea*). However, the host plants for use in the present invention are not limited thereto.

Plant Transformation

30 One skilled in the art recognizes that the expression level and regulation of a transgene in a plant can vary significantly from line to line. Thus, one has to test several lines to find one with the desired expression level and regulation. Once a line is identified with the desired regulation specificity of a chimeric Cre transgene, it can be crossed with lines carrying different inactive replicons or
35 inactive transgene for activation.

A variety of techniques are available and known to those skilled in the art for introduction of constructs into a plant cell host. These techniques include transformation with DNA employing *A. tumefaciens* or *A. rhizogenes* as the

transforming agent, electroporation, particle acceleration, etc. (See for example, EP 295959 and EP 138341). It is particularly preferred to use the binary type vectors of Ti and Ri plasmids of *Agrobacterium spp.* Ti-derived vectors transform a wide variety of higher plants, including monocotyledonous and dicotyledonous plants, such as soybean, cotton, rape, tobacco, and rice (Pacciotti et al. (1985) *Bio/Technology* 3:241; Byrne et al. (1987) *Plant Cell, Tissue and Organ Culture* 8:3; Sukhapinda et al. (1987) *Plant Mol. Biol.* 8:209-216; Lorz et al. (1985) *Mol. Gen. Genet.* 199:178; Potrykus (1985) *Mol. Gen. Genet.* 199:183; Park et al., *J. Plant Biol.* (1995), 38(4), 365-71; Hiei et al., *Plant J.* (1994), 6:271-282). The use of T-DNA to transform plant cells has received extensive study and is amply described (EP 120516; Hoekema, In: The Binary Plant Vector System, Offset-drukkerij Kanters B.V.; Alblaserdam (1985), Chapter V, Knauf, et al., *Genetic Analysis of Host Range Expression by Agrobacterium* In: Molecular Genetics of the Bacteria-Plant Interaction, Puhler, A. ed., Springer-Verlag, New York, 1983, p. 245; and An, et al., *EMBO J.* (1985) 4:277-284). For introduction into plants, the chimeric genes of the invention can be inserted into binary vectors as described in the examples.

Other transformation methods are available to those skilled in the art, such as direct uptake of foreign DNA constructs (see EP 295959), techniques of electroporation (see Fromm et al. (1986) *Nature* (London) 319:791) or high-velocity ballistic bombardment with metal particles coated with the nucleic acid constructs (see Kline et al. (1987) *Nature* (London) 327:70, and see U.S. Patent No. 4,945,050). Once transformed, the cells can be regenerated by those skilled in the art. Of particular relevance are the recently described methods to transform foreign genes into commercially important crops, such as rapeseed (see De Block et al. (1989) *Plant Physiol.* 91:694-701), sunflower (Everett et al. (1987) *Bio/Technology* 5:1201), soybean (McCabe et al. (1988) *Bio/Technology* 6:923; Hinchey et al. (1988) *Bio/Technology* 6:915; Chee et al. (1989) *Plant Physiol.* 91:1212-1218; Christou et al. (1989) *Proc. Natl. Acad. Sci USA* 86:7500-7504; EP 301749), rice (Hiei et al., *Plant J.* (1994), 6:271-282), and corn (Gordon-Kamm et al. (1990) *Plant Cell* 2:603-618; Fromm et al. (1990) *Biotechnology* 8:833-839).

Transgenic plant cells are then placed in an appropriate selective medium for selection of transgenic cells which are then grown to callus. Shoots are grown from callus and plantlets generated from the shoot by growing in rooting medium. The various constructs normally will be joined to a marker for selection in plant cells. Conveniently, the marker may be resistance to a biocide (particularly an antibiotic such as kanamycin, G418, bleomycin, hygromycin, chloramphenicol,

herbicide, or the like). The particular marker used will allow for selection of transformed cells as compared to cells lacking the DNA which has been introduced. Components of DNA constructs including transcription cassettes of this invention may be prepared from sequences which are native (endogenous) or foreign (exogenous) to the host. By "foreign" it is meant that the sequence is not found in the wild-type host into which the construct is introduced. Heterologous constructs will contain at least one region which is not native to the gene from which the transcription-initiation-region is derived.

To confirm the presence of the transgenes in transgenic cells and plants, a Southern blot analysis can be performed using methods known to those skilled in the art. Expression products of the transgenes can be detected in any of a variety of ways, depending upon the nature of the product, and include Western blot and enzyme assay. One particularly useful way to quantitate protein expression and to detect replication in different plant tissues is to use a reporter gene, such as GUS. Once transgenic plants have been obtained, they may be grown to produce plant tissues or parts having the desired phenotype. The plant tissue or plant parts, may be harvested, and/or the seed collected. The seed may serve as a source for growing additional plants with tissues or parts having the desired characteristics.

Description of the Preferred Embodiments

The present invention solves the problem of the conditional expression of various genetic traits in plants during specific times in a plant life cycle, in specific plant tissues, or in a specific generation by tying the expression of these traits to regulated promoters will additionally timing expression through the judicious use of site-specific recombinase systems. The constructs of the invention are referred to as recombinase elements. Each recombinase element will comprise at least one plant promoter. The promoters may be constitutive, inducible, tissue specific or developmental stage-specific promoters. Combinations of developmentally-regulated germline promoters are particularly useful. One recombinase element will typically consist of a single promoter operably linked to a recombinase. Second, third and fourth recombinase elements may consist of different promoters that will either drive the expression of a second or third recombinase or the expression of a transgene. Transgenes of the present invention will encode genetic traits, or various transformation or morphological markers. By configuring the recombinase elements and placing them in different parental plants it is possible to have the transgene expressed in specific tissues or during specific times in a plant life cycle or in a specific generation and then, if desirable, have the transgene selectively excised when no longer convenient. It will be appreciated that any number of recombinase elements may be combined

with these essential components to effect the regulate expression and expression of transgenes.

The invention provides constructs and methods for the conditional or regulated expression or excision of a transgene in a plant system. In its most basic form this may be accomplished through the use of a single site-specific recombina-
5 recombina- system and a pair of regulated promoters. Such as scheme is illustrated in Figure 1. Referring to Figure 1, a first recombina- element comprising a promoter (P1) operably linked to a recombina- (R1) coding sequence, is provided. Similarly a second recombina- element comprising a
10 second, different promoter (P2) placed upstream of a stop fragment (STP) which in turn is upstream of a transgene (TG) encoding a trait. The stop fragment (STP) is bounded by site specific sequences responsive to the recombina-, or recombina- sequences (RS1). Typically the first and second recombina- elements are provided in different plants, although it will be appreciated that, in
15 the case of an inducible P1 promoter, a single plant may be co-transformed with both elements. After crossing, or co-transformation, where the first and second recombina- elements are present in the same tissue, activation of the first promoter (P1) results in expression of the recombina- (R1) which in turn excises the stop fragment (STP) from the second recombina- element. Activation of the
20 second promoter effects expression of the transgene (TG) encoding a trait. One utility of such a scheme is to activate a transgene, whose expression is detrimental to normal plant development, only in the first generation. Such transgenes include those that result in too high levels of a desired product to be phytotoxic are not expressed during breeding but only in the harvestable generation.

25 In an alternate embodiment the invention provides a constructs and methods for removal or excision of the transgene encoding a trait at some specified developmental stage in a plant life cycle of the first or the second generation. A simple example of this method is illustrated in Figure 2. Referring to Figure 2, again two recombina- elements are provided. The first recombina-
30 element comprises a germline promoter (P1) operably linked to a recombina- coding sequence (R1). The second recombina- element comprises a promoter (P2) operably linked to a transgene (TG) encoding a trait where the entire recombina- element is bounded by recombina- sequences (RS1) responsive to the recombina-. Typically the first recombina- element is provided in a first
35 plant whereas the second recombina- element is provided in a second plant although it will be appreciated that, in the case of an inducible P1 promoter, a single plant may be co-transformed with both elements. Where P1 is a developmentally regulated germline promoter, a cross of the first and second

plants results in activation of P1 and the expression of recombinase (R1) in the first generation. When TG is expressed in vegetative tissues and where P1 is a common or male germline promoter that is activated late in the plant life cycle to express the recombinase (R1) and cause the excision of the P2-TG chimeric element in the first generation from the seed or pollen, respectively. One utility of this scheme is to remove a transgene from pollen to prevent its environmental impact, escape to wild type relatives, and escape to neighbor crops. Another utility is to remove the trait gene from progeny seeds to prevent residual or volunteer plants in the next growing season, reduce regulatory concerns, and provide a tighter control on unwanted gene flow.

Yet another utility is for marker excision. i.e., removal of transformation marker for easier regulatory approval and for reusing the marker for retransformation. Antibiotic or herbicide resistance genes are used as selectable markers in plant transformation to select for the rare transgenic cells from nontransgenic ones. However, the presence of such markers in transgenic plants is undesirable because of regulatory concerns/requirements as well as because it prevents recurrent use of the selectable marker for stacking trait transgenes. The use of a bacterial IPT gene as a marker for transformation was proposed by Ebumina et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2117-2121) and Kunkel et al. *Nat. Biotechnol.* (1999), 17(9), 916-919. In the former case, the IPT gene was inserted inside a transposable element, whose random excision following transformation resulted in the loss of the transposable element and the IPT gene. However, this method is inefficient (see Kunkel, *supra*), especially because of its low frequency of loss (1% or less). Kunkel, *supra*) proposed the use of a chemically-inducible IPT gene. However, this is also undesirable, since the bacterial IPT gene is not lost following transformation and that could be of concern from a regulatory point of view. Furthermore, it does not allow its use for retransformation for trait stacking. Therefore, there is a need for an efficient, regulatable excision of a constitutively expressed transformation marker gene. We propose inserting a transformation marker gene within site-specific sequences such that it is efficiently excised upon the expression of a chimeric recombinase gene under the control of a regulated or an inducible promoter following transformation. SAM-specific genes could be particularly useful in that their promoters could express Cre and cause SSR in shoot apices following transformation of undifferentiated cells/calli. This would be useful to remove the selectable genes early in tissue-culture but after the selection. This can be tested with a non-destructive excision reporter such as GFP or luciferase either in agrobacterium-mediated transformation of tobacco (leafdisc method) or

Arabidopsis, or by biolistic bombardment in soybean or corn embryogenic cultures. The said invention provides for such promoters (AP3, BCP1, HSP18.2)

In a particularly useful embodiment, the invention may be used for *in planta* transformation to circumvent the need to go through time-consuming and laborious plant tissue culture method of transformation. This relies on the use of a scorable (non-selectable) marker transgene that induces morphological or visible changes in transgenic tissue (such as shoots and/or roots. Examples of such genes are STM, KNAT 1 homoeobox genes, Lec 1, or ANT gene, or even agrobacterium T-DNA that can induce proliferation and redifferentiation *in planta*. Subsequent loss of these markers from the germline under the control of a regulated site-specific recombinase (as above) will allow recovery of morphologically normal transformants on the plant. The regulated recombinase gene can either be within the site-specific sequences or outside. The promoter for the recombinase gene can be from a developmentally regulated germline specific gene, such as those involved in meristem identity, organ primordia, or anther/pistil. It can also be inducible by a chemical, such as a safener (e.g., IN2 promoter) or by heat shock, such as *Arabidopsis* HSP18.2 gene. The said invention provides for such promoters (AP3, BCP1, HSP).

In other preferred embodiments the invention will make use of two or more developmentally staggered site-specific recombinations systems. Although SSRs have been used singly as genetic switches, two (or more) SSRs under the control of different constitutive or regulated promoters can be used as a series of genetic switches within a plant's life cycle, such that conditional expression of one recombinase (R1) at one stage activates another recombinase (R2) at a later stage. Thus, one can conditionally trigger the process at a convenient developmental stage, such as germination, but get delayed effects at later stages.

Conditionality to the first SSR is provided by either chemical application or a genetic cross that combines its recombinase gene with its cognate target gene/s. The latter is more amenable for hybrid crops. Chemical application on seeds or during germination is likely to overcome the chemical's cost and problem with its biokinetics into target cells. Chemical application can also be done in the prior generation by using a relay of three, rather than two, site-specific recombination systems. Thus, the chemical can be applied to germinating seeds in the last generation of seed production to induce one type of SSR that results in another type, say in late seed development of progeny seeds, that, in turn, results in a third type of SSR to express in early seeds to remove the trait gene. In another embodiment R1 can be chemically repressible, such that the application of the chemical represses SSR I (R1) to allow production of seeds with the transgenic

trait. Here, in the absence of the chemical, such as in the farmers' field, the crop is genetically triggered to enable trait gene expression and/or its subsequent removal on cue.

Such combinations of two or more different site-specific recombinations, whether linked or unlinked, provide novel and useful tools to control transgene expression and/or removal in the first, second, or third generations that are not currently available in agricultural biotechnology. Thus, a pair of developmentally staggered SSRs may be used as ON-ON or ON-OFF (transgene removal) switches. The salient feature in both schemes, is that expression of R2 and/or trait genes does not have to occur immediately upon enablement (i.e., removal of the stop fragment) by R1 but are rather controlled solely by the choices of P2 and P3 promoters. While the following embodiments involve two site-specific recombinations, it is recognized that the strategy can be extended to include addition site-specific recombinations for controlling transgene expression even beyond the second generation or to include more elaborate transgene activation or removal.

In one embodiment, two SSRs may be staggered as ON-ON switches to get trait expression later than possible with only one SSR. One such utility is for allowing trait expression only in the second generation. This is important when trait expression is detrimental to normal plant development, such as seed germination or seedling growth, and it is undesirable for trait expression during breeding or in the farmer's field but desirable in the generation that is harvested and processed. Thus, in an alternate preferred embodiment the invention will make use of at least three separate recombinase elements as shown in Figure 3. Referring to Figure 3, a first recombinase element is provided consisting of a first promoter (P1) operably linked to a first recombinase coding sequence (R1). A second recombinase element is provided consisting of a second different promoter (P2) upstream of a stop fragment (STP) which is in turn upstream of a second recombinase coding sequence (R2), where the stop fragment (STP) is bounded by recombinase sequences responsive to the first recombinase (RS2). A third recombinase element is provided consisting of a third promoter (P3) upstream of a stop fragment (STP) which is in turn upstream of a trait transgene (TG), where the stop fragment (STP) is bounded by recombinase sequences (RS2) that are responsive to the second recombinase. It will be appreciated that, when P1 is an inducible promoter, the first, second and third recombinase elements may be combined into a plant through any means including crossing, or co-transformation and when P1 is not an inducible promoter, the first and the second elements are brought together in the first generation by a cross. In one particularly useful

embodiment a first plant is provided having the first and third recombinase elements while a second plant is provided having the second and third recombinase elements. Crossing the first and second plant will give rise to the first generation plant in which the first recombinase is expressed excising the stop
5 fragment (STP) from the second recombinase element and allowing the expression of recombinase R2 under the control of P2 promoter, which recombinase, in turn, excises the stop fragment from the third recombinase element, allowing expression of the trait gene under the control of P3 promoter in the second and subsequent generations. Here P1, P2, and P3 promoters are not all activated
10 simultaneously. They may each be activated at a different stage: first P1 in first generation, then P2 in first or second generation, and last P3 in the second generation; alternatively, P1 and P2 are activated together in the first generation and P3 in the second generation; alternatively, P1 is activated in the first generation and P2 and P3 are activated together in second generation. Also
15 promoters for expression in first generation express in common germline, while promoters P3 and P2, when it occurs in second generation, may be in second generation germline or somatic cells and may be developmental stage-specific or chemically inducible. One utility of this scheme is to express traits, such as developmental traits, only in second and subsequent generation. Such
20 developmental traits include apomixes, parthenocarpy, flowering (Nilsson et al. (1998) *The Plant Journal* 15:799-804), self-incompatibility (Stone et al. (1999) *Science* 286:1729), altered flowering time (Mandel and Yanofsky *Nature* (1995) 377:522; Weigel and Nilsson (1995) *Nature* 377:495), type of pollination (selfing vs. crossing), and barriers to cross pollination, pollination control, self-
25 incompatibility, sterility, etc. Some of these traits will require expression of a trait transgene encoded polypeptide, such as a toxin or a transcriptional factor that regulates endogenous genes, others will not, such as when a trait transgene co-suppresses or silences an endogenous gene (see Chuang CF, Meyerowitz EM (2000) Specific and heritable genetic interference by double-stranded RNA in
30 *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A*.97:4985-4990).

Another utility of this application is to allow trait gene expression in germinating seeds or seedlings of the second generation after harvest in contained bioreactors. This is important when trait expression of toxic material is undesirable in the field. In an extension of this concept, another transgene is
35 activated that results in trait gene removal in the second generation or the second generation has a lethal gene that block development or a sterility gene that prevents flowering and seed set to further genetically contain the trait gene.

In another embodiment of the invention two or more site-specific recombination may allow conditional and transient transgene expression by linking trait activation and trait removal.

Here two SSRs may be developmentally staggered, such that R2
5 recombinase removes a transgene. This will result in an ON-OFF (by transgene removal) switch to get transient trait expression. Trait gene removal may be unlinked (Figures 2, 8) or linked (Figures 4,5,6) to trait gene expression. The latter provides a more stringent control of trait gene expression. These Figures show that only the trait gene is removed, although all transgenes could be removed
10 by having them flanked by appropriate recombinase sites.

For example, in some situations it may be desirable to express a trait at one point in a plant life cycle, but then have that trait removed in a later generation. Such a scheme is illustrated in Figure 4. Referring to Figure 4, two plants are provided. The first plant will contain a first recombinase element comprising a
15 first promoter (P1) operably linked to a first recombinase coding sequence (R1). The second plant will contain both second and third recombinase elements. The second element will consist of a second promoter (P2) upstream of a stop fragment (STP) which in turn is upstream of a transgene encoding a trait (TG). The entire second recombinase element is flanked by recombinase sequences
20 responsive to a second recombinase (RS2). The third recombinase element is comprised of a third promoter (P3) upstream of a stop fragment (STP) which is in turn upstream of a second (different from R1) recombinase coding sequence (R2). In both the second and third recombinase elements the stop fragments (STP) are flanked by recombinase sequences responsive to the first recombinase (RS1).

It will be appreciated that, when P1 is an inducible promoter, the first,
25 second and third recombinase elements may be combined into a plant through any means including crossing, or co-transformation and when P1 is not an inducible promoter, the first and the second elements are brought together in the first generation by a cross. Thus, where the first promoter (P1) is a developmentally
30 regulated germline promoter, crossing the first and second plants will result in the activation of P1 and the expression of the first recombinase (R1). Expression of the first recombinase (R1) primes both the second and third recombinase elements by excising the stop fragment (STP) in each. Where the second promoter (P2) is activated prior to the third promoter (P3), the transgene is expressed. The third
35 promoter (P3) is a common or male germline promoter, whose activation will cause the expression of R2 and the removal of the trait gene from the progeny seed of the second generation or pollen of the first generation. Thus, the invention

provides for both conditional expression of the trait and its subsequent excision when no longer useful.

Here P1, P2, and P3 promoters are not all activated simultaneously and P3 promoter is always activated after useful expression of the trait gene under P2 promoter. Promoter P1 is activated in the germline of first generation, P2 promoter is activated in germline or somatic cells of the first or second generation, and P3 promoter is activated in the common or male germline of the first or second generation following useful trait expression under P2. The three promoters may each be activated at different developmental stages; for example, promoter P1 is activated first in the common germline of first generation, then P2 in somatic or somatic plus germline cells of first or second generation, and last P3 in the common or male germline cells of first or second generation following useful trait expression; alternatively, P1 and P2 may be activated together in the first generation and P3 later in the first generation or in the second generation; or P1 may be activated first in the common germline of the first generation and P2 in somatic cells and P3 in germline cells of first or second generation. This scheme combines two utilities: first, trait genes can be activated and expressed only in the desired generation, especially useful when the target product is phytotoxic; and second, after trait genes have served their usefulness, they are removed from pollen or seeds for better genetic containment and prevention of unwanted gene flow.

Other utilities include molecular approaches for controlling developmental traits, such as flowering (Nilsson et al. (1998) *The Plant Journal* 15:799-804), self-incompatibility (Stone et al. (1999) *Science* 286:1729), altered flowering time (Mandel and Yanofsky *Nature* (1995) 377:522; Weigel and Nilsson (1995) *Nature* 377:495), apomixes, parthenocarpy, type of pollination (selfing vs. crossing), and barriers to cross pollination, etc.

One particularly useful trait is conditional male sterility, which is important for hybrid seed production. The present invention is particularly useful for controlling male sterility by providing a simple molecular biology approach for conditional male sterility. Male sterility can be sporophytic or gametophytic. For example, Referring to Figure 5, two male fertile plants are provided. The first plant contains a first recombinase element consisting of a first promoter (P1) operably linked to a first recombinase coding sequence (R1). The second plant contains both a second and third recombinase element. The second element will consist of a second promoter (P2) upstream of a stop fragment (STP) which in turn is upstream of a transgene encoding the male sterile trait (MS). The entire second recombinase element is flanked by recombinase sequences responsive to a

second recombinase (RS2). The third recombinase element is comprised of a third promoter (P3) upstream of a stop fragment (STP) which is in turn upstream of a second recombinase coding sequence (R2). In both the second and third recombinase elements the stop fragments (STP) are flanked by recombinase sequences responsive to the first recombinase (RS1).

It will be appreciated that, when P1 is an inducible promoter, the first, second and third recombinase elements may be combined into a plant through any means including crossing, or co-transformation and when P1 is not an inducible promoter, the first and the second elements are brought together in the first generation by a cross. Thus, where the first promoter (P1) is a constitutive or developmentally regulated common germline promoter, crossing the first and second plants results in expression of the first recombinase (R1) and the priming of both the second and third recombinase elements by removal of the stop fragments (STP). Where the second promoter (P2) is anther specific the male sterile gene is expressed in the sporophyte cells of the anther to render the first generation male sterile. Where the third promoter (P3) is a seed specific promoter, the male sterility transgene is excised in the F1 hybrid seed to render it male fertile.

The dominant male sterility gene can encode a toxin (e.g., barnase, avidin, RIP) gene or a co-suppressor of a fertility gene (e.g., corn MS45 gene). This element can also be a constitutive promoter expressing a co-suppressor of an anther-specific fertility gene (e.g., corn MS45 gene).

The present invention may be specifically applied for the conditional control of male sterility in corn according to the TopCross® method of breeding. When male sterility is required even in the next generation, such as in corn TopCross® the above scheme can be modified to omit fertility restoration and incorporate a second conditional male sterility system. For example, referring to Figure 7, two male fertile plants are provided. The first plant comprises a first and third recombinase element while the second plant comprises a second and third element. The first recombinase element comprises a first promoter (P1) operably linked to a first recombinase coding sequence (R1). The second recombinase element comprises a second promoter (P2) upstream of a stop fragment (STP) which is in turn upstream of a transgene encoding male sterility (MS), wherein the stop fragment (STP) is bounded by site-specific recombinase sequences responsive to the first recombinase (RS1). The third recombinase element comprises a second promoter (P2) upstream of stop fragment (STP), which is in turn upstream of a transgene encoding male sterility (MS), wherein the stop fragment (STP) is bounded by site-specific recombinase sequences responsive to

the second (and different from R1) recombinase (RS2). A third male fertile plant is also provided having a fourth recombinase element comprising a first promoter (P1) operably linked to a second recombinase coding sequence (R2)

A cross of the first and second plants produces a first generation plant.

- 5 Where the first promoter is constitutive or germline specific P1 is activated and R1 is expressed. The expression of R1 excises the stop fragment (STP) from the second recombinase element. Where P2 is activated in the first generation the MS transgene is expressed resulting in a male sterile plant. Crossing the male sterile plant with the third plant produces an F1 hybrid sterile plant. The plant is sterile
10 since, where P1 is activated in the hybrid, R2 is expressed causing the excision of stop fragment (STP) from the third recombinase element, allowing the expression of the second MS gene, where P2 is activated. Crossing this male sterile F1 hybrid with a TopCross® pollinator results then in a TopCross® hybrid.

- The methods of the present invention may also be applied to provide
15 conditional male sterility in seeds. For example, referring to Figure 8, three plants are provided. The first plant comprises a first recombinase element, the second comprises a second recombinase element and the third comprises a third element. The first element comprises a first promoter (P1) operably linked to a first recombinase coding sequence (R1). The second element comprises an anther
20 promoter upstream of a stop fragment (STP) which is in turn upstream of a male sterility encoding gene (MS). The stop fragment (STP) is bounded by site-specific recombinase sequences responsive to the first recombinase (RS1) and the entire element is bounded by site-specific recombinase sequences responsive to a second recombinase (RS2). The third element comprises the first promoter (P1) operably
25 linked to a second recombinase coding sequence (R2).

- It will be appreciated that, when P1 is an inducible promoter, the first and second recombinase elements may be combined into a plant through any means including crossing, or co-transformation. Where P1 is constitutive or developmentally regulated common germline promoter, crossing the first and
30 second plants results in a first generation plant where R1 is expressed and the stop fragment is excised from the second recombinase element. Where the anther promoter is activated the male sterility transgene (MS) is expressed and the plant is male sterile. Crossing this male sterile plant with the third plant results in the removal of the MS gene and restoration of fertility in subsequent generations.

- 35 The present invention will also allow for the conditional control of gametophytic male sterility to ensure that a trait transgene that is present in a heterozygous state in one of two parent lines of a cross is expressed in all second generation (F2) grain. For example, referring to Figure 6, two male fertile plants

are provided. The first plant comprises second and third recombinase elements. The second plant comprises first and third recombinase elements and a transgene physically linked to a gametophytic male fertility restorer gene. The first recombinase element comprises a first promoter (P1) operably linked to a first
5 recombinase coding sequence (R1). The second recombinase element comprises a first promoter (P1) upstream of a stop fragment (STP) which is in turn upstream of second recombinase coding sequence (R2), wherein the stop fragment (STP) is bounded by site-specific recombinase sequences responsive to the first recombinase (RS1). The third recombinase element comprises a pollen specific
10 promoter upstream of a stop fragment (STP) which is in turn upstream of a transgene encoding gametophytic male sterility (gms) coding sequence, wherein the stop fragment (STP) is bounded by site-specific recombinase sequences responsive to the second recombinase (RS2). The male sterility gene can encode barnase. The fourth element is linked trait gene (TG) and a gametophytic male
15 fertility (gmf) restorer gene. The gmf restorer gene can be barstar, when the sterility gene is barnase.

When the first and second plants are crossed a first generation plant is produced. Where the first promoter (P1) is a constitutive or developmentally regulated common germline promoter; the first promoter is activated causing the
20 expression of the first recombinase (R1). Expression of the R1 resulted in removal of the stop fragment (STP) from the second recombinase element and expression of the second recombinase (R2). Expression of R2 results in the removal of the stop fragment (STP) from the third recombinase element and the priming of that element for expression of the gametophytic male sterility (gms)
25 gene under the control of a pollen promoter. This system allows for the expression of the gametophytic male sterility gene in pollen without the trait gene-gmf gene but not in pollen with the trait gene-gmf gene.

In order to accomplish the above mentioned embodiments, this invention discloses the use of promoters for either male germline, floral common germline,
30 and vegetative SAM common germline SSR. It also provides for an excision reporter construct that one skilled in the art can screen promoters with the desired activation specificities.

Site-specific recombinations in germline may be accomplished by the regulated expression of Cre using promoters from germline cell-specific genes
35 promoters. These SSRs may be specific for male, female, or common germlines. Examples of likely genes for germline promoters are:

- a) AP3 and PI homeotic genes, and anther genes (SAP, Bcp1, etc) for male germline, or male gametophyte promoters, such as LAT52 (also see Twell et al. (1998) *Trends in Plant Sciences* 3:305)
- b) Petunia's FBP7 and FBP11 involved in the formation of ovule
5 primordium, Arabidopsis ANT and HLL genes involved in initiation of ovule protrusion for female germline; and
- (i) shoot apical meristem (SAM) genes, organ primordia-specific genes, and floral homeotic genes, such as AG, LFY and ER for common germline. This category is especially useful for removing
10 selectable genes early in tissue-culture following selection in tissue culture. Examples of such homeotic genes are: shoot apical meristem (SAM) genes include WUS and STM, maize KN1, rice OSH1, and UNUSUAL FLORAL ORGANS (UFO) gene.

It is also disclosed that the specificity of germline promoters can be the
15 same as expected for the endogenous gene from which the promoter is derived or be changed depending on the transgenic line carrying the construct. Thus, AP3 was shown to confer vegetative SAM common germline, floral SAM common germline, and floral male germline excisions depending on the line. Such variation in transgene expression is not uncommon and one skilled in the art can
20 screen and identify lines with the desired germline specificity.

EXAMPLES

The present invention is further defined in the following Examples. These Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one
25 skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

GENERAL METHODS

Standard recombinant DNA and molecular cloning techniques used in the
30 Examples are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, (1989) (Maniatis) and by T. J. Silhavy, M. L. Bannan, and Enquist, L. W. *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1984) and by Ausubel, F.
35 M. et al., *Current Protocols in Molecular Biology*, pub. by Greene Publishing Assoc. and Wiley-Interscience (1987).

Restriction enzyme digestions, phosphorylations, ligations and transformations were done as described in Sambrook, J. et al., *supra*. Restriction

enzymes were obtained from New England Biolabs (Boston, MA), GIBCO/BRL (Gaithersburg, MD), or Promega (Madison, WI). Taq polymerase was obtained from Perkin Elmer (Branchburg, NJ). Growth media was obtained from GIBCO/BRL (Gaithersburg, MD).

- 5 The *Agrobacterium tumefaciens* strain LBA4404 was obtained from Dr. R. Schilperoot, Leiden (Hoekema et al. *Nature* 303:179-180, (1983)).

Transformation Protocols

- Biolistic transformations were done essentially as described in U.S. Patent No. 4,945,050, hereby incorporated by reference. Briefly, gold particles (1 mm in diameter) are coated with DNA using the following technique. Ten ug of plasmid DNAs are added to 50 uL of a suspension of gold particles (60 ug per uL). Calcium chloride (50 uL of a 2.5 M solution) and spermidine free base (20 uL of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 min, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 uL of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 uL of ethanol. An aliquot (5 mL) of the DNA-coated gold particles can be placed in the center of a flying disc (Bio-Rad Labs, 861 Ridgeview Dr, Medina, OH). The particles are then accelerated into the corn tissue with a PDS-1000/He (Bio-Rad Labs, 861 Ridgeview Dr., Medina, OH), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.
- 10
15
20

- Where *Agrobacterium* transformations were done the procedure was accomplished essentially as described Park et al., *J. Plant Biol.* (1995), 38(4), 365-71.
- 25

EXAMPLE 1

Making An Excision Reporter in Binary Vector (pBE894) To Test Germline Specificity of Promoters

- In order to identify male germline and common germline promoters, an excision reporter was made. In this reporter, the plant kanamycin resistance gene flanked by lox sites is inserted as a blocking fragment between a constitutive promoter and the luciferase coding region. The blocking fragment blocks the translation of luciferase by interrupting the luciferase coding sequence and such that upon cre lox excision, there is a single copy of lox site is left behind as a translational fusion with the Luciferase ORF that allows luciferase expression. The Cre gene under the control of regulated germline promoters is maintained outside the lox sites, i.e., not flanked by lox sites.
- 30
35

The excision reporter in binary plasmid, named pBE894, was made through the following intermediate plasmids:

5 pGV853 is 35S:Lox:yeast 2u:Lox:Luc (translational fusion). The coding sequence of luciferase was isolated from a 35S:luciferase gene (Promega) by PCR using primers P298 and P299 and cloned by *in vivo* recombination in yeast (as described in PCT WO 99/22003) to replace the GUS coding sequence in pGV796, previously digested with Eco RI and Sma I, to result in plasmid pGV853 carrying 35S:Lox:yeast 2u:Lox:Luc. The N terminus of the luciferase was translationally fused to the Lox sequence.

10 pGV789 is a floxed Lox 66':Avr II:Lox 71' as a translation stop in GUS gene. It was made as follows: PCR products A and B were made on pML63 carrying a 35S promoter:GUS gene using primer pairs P192, P194 and P192, P198, respectively, both were cut with Avr II, ligated, and used as template for second PCR using primers P192 and P198 to result in PCR product C, which was
15 cut with Bgl II and Bcl I and cloned into Bgl II - Bcl I cut pML63. One skilled in the art can deduce the translation fusion from the sequences of the PCR primers used.

pGV796 is a floxed WT Lox P:Avr II:WT Lox P as a translation stop in GUS gene. It was made by PCRing the yeast trp 2 μ m fragment by using PCR
20 primers P227 P228 (that introduced wild type lox P sites at both ends) on a plasmid containing the yeast 2 μ sequence and cloning it by yeast homologous recombination into Avr II cut pGV789, which has a unique Avr II sites between mutant Lox sites: 35S promoter: Lox 66'-Avr II-Lox 71' as a translation stop in GUS gene.

25 pGV890 is 35S:Luc blocked by a yeast 2 μ fragment flanked by Lox sites. It was made by converting the Bam HI site of pGV853 to an Eco RI site. For this, pGV853 was digested with Bam HI, filled in, ligated to Eco RI linkers (New England Biolab. #1020), digested with Eco RI, and self religated.

pGV891 is 35S:Luc blocked by NPT II gene yeast (as described in PCT
30 WO 99/22003) flanked by Lox sites. It was made by inserting the 1781 bp Avr II fragment from pGV801 containing the kanamycin resistance gene (nos:NPT II;3'nos gene) for plant transformation into the Avr II site of pGV890 in the desired orientation, i.e., same as the Luc coding sequence.

pBE892 is bar binary with floxed NPT II gene and "outside" SCP:Cre.
35 The SCP promoter is described in Bowen, Benjamin A.; Bruce, Wesley B.; Lu, Guihua; Sims, Lynne E.; Tagliani, Laura A. Synthetic constitutive promoters for high-level expression of foreign genes in plants. U.S. (2000), 31 pp., Cont.-in-part of U.S. Ser. No. 661,601, abandoned. CODEN: USXXAM US 6072050 A

20000606. It was made by cloning the Eco RI/Xba I fragment of pGV891 into Eco RI/Xba I of the bar binary pBE673 (as described in PCT WO 99/22003).

pGV895 is SCP:mCre. The Bam HI/Eco RI fragment of pHP15254 carrying bacterial Cre sequence was replaced by the Bam HI/Eco RI fragment of pHP16072 carrying the maize codon optimized Cre w/ the potato ST-LS1 intron (described in WO 9925840).

pGV897 is SCP:mCre. It was made by introducing a Hind III site at the Eco RI site in pGV895 by ligating Hind III linkers (NEB #1050), digesting with Hind III, and religating.

pBE894 is bar binary with floxed NPT II gene and "outside" SCP:Cre. It was made by digesting pBE892' with Xba I/Hind III and cloning in the 2159 bp Xba I/Hind III fragment from pGV897 carrying the SCP:Cre gene.

To test the vector *Agrobacterium tumefaciens* strain C58 was transformed and the resultant strain CA894 was used to transform 100 tobacco leaf discs as well known in the art. Table 1 shows that there were only 3 out of 90 discs that were alive on Kan after 4 weeks. Of these three only one showed luciferase expression by imaging. On the other hand 114/132 (86%) discs were resistant on Bar and almost all showed luciferase expression. Thus, the excision reporter works well.

Table 1

Resistance in Tobacco Leaf Discs after 4 weeks

Agrobacterium	Bar selection	Kan selection
None	0/6 resistant	0/6 resistant
CA894	114/132 resistant	3/90 resistant

EXAMPLE 2

Identifying Promoters for Male and Common Germline Expression in *Arabidopsis thaliana*

The Xba I-Bam HI fragment of pBE894 carrying the SCP1 promoter was replaced by an Xba I-Bam HI fragment carrying one of several different developmentally-regulated promoters. These promoter regions, the PCR primers used to isolate them from genomic DNA of *Arabidopsis*, and the resultant binary plasmids are:

Promoter for *Apetala 3* (AP3) gene from *Arabidopsis thaliana* Col. (pBE913).

Promoter for *Bcp 1* (BCP 1) gene from *Arabidopsis thaliana* Col. (pBE914).

Promoter for *Erecta* (ER) gene from *Arabidopsis thaliana* Col. (pBE915).

Synthetic anther promoter (SAP) (pBE928) G9/SGB6 hybrid promoter
5 (U.S. Pat No. 5470359; 5837850)

Promoter for *Pistilata* (PI) gene from *Arabidopsis thaliana* Col. (pBE929).

Promoter for TA29 from tobacco (Hsu, Francis C.; Odell, Joan Tellefsen; Shen, Jennie Bih Jien. Induction of male sterility in crop plants with heterologous genes expressed from tissue-specific promoters. PCT Int. Appl. (1992), 92 pp.
10 CODEN: PIXXD2 WO 9204454 A1, 19920319 CAN 117:209112) in pTZALG pBE855).

	<u>Promoter</u>	<u>Length</u>	<u>Nucleotide positions</u> (Genbank Accession #)	<u>PCR Primers</u>	
				<u>UP</u>	<u>LP</u>
15	Apetala 3 (AP3)	605 bp	1151-1755 (U30729)	PH785	PH786
	BCP 1 (BCP 1)	579 bp	48943-49528 (U30729)	PH783	PH784
	Erecta (ER)	1187 bp	616-1802 (D83257)	PH788	PH790
	TA29	1525 bp	(from plasmid pTZALG)	PH795	PH815
	Pistilata (PI)	299 bp	2997 3296 (AB035137)	P321	P322

20

Oligos used for PCR of promoters from *Arabidopsis* genomic DNA.

AP3 PCR primers

PH785:

5'-AGT CTA GAC CCG GGA TGG AAG TGA CGA TTA-3' (SEQ ID NO:8)

25 PH786:

5'-GAG GAT CCC GGG TCT TCT CTC TTT GTT T-3' (SEQ ID NO:9)

BCP1 PCR primers

PH783:

30 5'-TAT CTA GAC CCG GGT CTC GAT CCG ATC GAA-3' (SEQ ID NO:10)

PH784:

5'-TTG GAT CCC GGG TTC TCT CTC TCC TTC TTA-3' (SEQ ID NO:11)

ER PCR primers

35 PH788:

5'-GGT CTA GAC CCG GGA CTT TTT GAG AAA AG-3' (SEQ ID NO:12)

PH790:

5'-ATG GAT CCC GGG TTC TCA CAC ACA GTC TTA-3' (SEQ ID NO:13)

PI PCR primersP321:

5'-CGT CTA GAC CCG GGA TGT TGT CTT CAA GGC-3' (SEQ ID NO:16)

5 P322:

5'-ATG GAT CCC GGG TTC TCA CAC ACA GTC TTA-3' (SEQ ID NO:17)

For PCR, genomic DNA (50 – 200 ng for each 50 µl reaction) isolated from *Arabidopsis thaliana* var. Columbia, was used as the PCR reaction template unless otherwise indicated. Reaction mixtures included final concentrations of the following: 1 µM each of a 5' and 3' primer designed to amplify the desired promoter, 200 µM of each of 4 nucleotides dATP, dCTP, dGTP, dTTP; 1.25 µM Amplitaq (PE Applied Biosystems, Foster City, CA). PCR reactions were carried out in a Perkin Elmer 9600 thermocycler (PE Applied Biosystems, Foster City, CA). The thermocycler was programmed as follows: a 2 min 96°C initial denaturation step was followed by 25 cycles of 94°C, 45 sec; 55°C, 45 sec; 72°C, 90 sec, and ended with an 8 minute 72°C final extension cycle. The desired PCR products were cloned into a PCR cloning vector pGEM-T (Promega Corp., Madison, WI) according to the manufacturer's protocols. After the presence of a cloned insert was established, the ends of the insert were sequenced to confirm that the desired promoter had been amplified and cloned. The cloned promoter fragments were isolated as Xba I- Bam HI fragments and cloned into binary plasmid pBE894.

TA29 promoter was isolated as a Xba I- Nco I fragment and used to replace a 35S promoter in front of the bacterial (not maize optimized) Cre ORF without intron. The chimeric gene was then cloned into a bar binary vector pBE673, described in PCT application WO 99/22003.

Binary plasmids pBE913, pBE914, pBE915, pBE928, pBE929, and pBE855 were transformed into *Agrobacterium tumefaciens* strain C58 as described in PCT application WO 99/22003.

Transformants selected on LB medium containing 100 mg/L kanamycin. Transformed bacteria were then used to transform *Arabidopsis thaliana* var. Col by the floral dip method of Clough SJ, Bent AF (1998) Floral Dip: A Simplified Method for *Agrobacterium*-Mediated Transformation of *Arabidopsis thaliana*. *Plant J* 16:735-743. Specifically, several pots of healthy *Arabidopsis* plants were grown under long day conditions. After 4 to 6 weeks the first bolts were clipped. This encourages the proliferation of secondary bolts in 4 to 6 days. A starter culture (~200 mL of LB with antibiotics) of C58 *Agrobacterium* strain carrying the

gene of interest on a binary vector was grown for 2 days at 20 °C with agitation. The starter culture was used to inoculate a 2 L culture which was incubated overnight. The culture was spun down at 5,000 rpm for 10 min at RT. The cells were resuspended in 5% sucrose with 0.05% Silwet L-77 (Lehle Seeds) to an

5 OD600 of 0.8 or higher. The above ground parts of the *Arabidopsis* plants were dipped in the Agro/sucrose solution for 2-3 seconds with agitation. The pots were laid on their side, covered with a plastic dome and placed in low light conditions for a couple of days, then uprighted and placed back into full light. After ~6 weeks the plants were allowed to dry down and the seed was harvested. Seeds

10 were sterilized using 80% ethanol with 0.01% Triton X-100 for 10 min with agitation, 33% bleach with 0.01% Triton X-100 for 10 min with agitation followed by 5 sterile water rinses. The seeds were then suspended in 8 mL of 0.1% sterile agar and spread onto plates. The plates consist of 1x MS, 1% sucrose, 0.8% agar, 100 mg/L Timentin (Smith Kline Beecham), 10 mg/L

15 Benomyl (DuPont) and antibiotic selection. Kanamycin sulfate (Kan) was used at 50 mg/L and glufosinate ammonium (Bar) at 20 mg/L. Resistant seedlings were removed from the plates, imaged for luciferase expression non-destructively (Tables 2-7). Lines that had no or very slight Luciferase expression (Luc +) were potted and selfed. Progeny T2 seeds of were analyzed for resistance to Kan or

20 Bar.

The results with AP3, Bcp 1, PI, ER, and SAP show that:

- (i) all these promoters were not activated in developing embryos/seeds, since there was no significant difference in transformation efficiency on Kan and Bar plates (except possibly

25 for ER:Cre transformants, Table 5), indicating no significant excision in seeds,

- (ii) all these promoters have non-specific expression in some lines, since in all seedlings resistant to Kan or Bar there was varying amounts of luciferase expression, including none, very slight

30 (Luc +), medium (Luc ++), and high (Luc ++++). AP3 and PI constructs showed a similar distribution in Luc expression suggesting similar profile in expression specificity. AP3 and PI appear to have the most lines that show leaky expression, while Bcp 1 and SAP promoters show the least. For example, 40% of the

35 Kan resistant seedlings showed high luc expression. In these lines AP3 is apparently serving as vegetative SAM promoter.

- (iii) There was no toxicity from Cre expression in these lines,

- (iv) ER gave the least transformants (Table 4) and were replated (Table 5). The second plating suggest that ER:Cre may be toxic. It is not known if this toxicity, if real, is from Cre or the ER promoter.

- 5 Lines that had no or very slight Luciferase expression (Luc +) were potted and selfed. Progeny T2 seeds of were analyzed for resistance to Kan or Bar (Tables 8 and 9). These results show that AP3:Cre is a good promoter not only for vegetative SAM excision but also for floral common germline and male germline excisions. Thus, of the 17 AP3 lines analyzed at T2 stage, 9 showed 100%
 10 excision from the common (male and female) germ lines, 4 showed 98% excision, 1 showed 92 % excision, and one (AB07) showed male germline excision. With Bcp 1, some lines showed neither Bar or Kan resistance and may represent escapes during T1 selection. But there was one line (BK21) that showed male germline excision since Kan segregated 3:1 and Bar segregated 15:1 (two loci).
 15 When lines with common or male germline excision had more than one locus for the reporter construct, excision occurred in all loci.

Table 2

Analysis of transformed Arabidopsis seedlings (T1 progeny of plants transformed with CA913 carrying AP3: Cre) for Luciferase expression following selection on

20

Kan or Bar.

Selection	Luc OFF	Luc ON +	Luc ON++	Luc ON++++	Total	TE (%)
Kan	3	5	6	10	24	0.26
Bar	9	5	10	8	32	0.35
Total	12	10	16	18	56	
%	21	18	29	32		

Table 3

Analysis of transformed Arabidopsis seedlings (T1 progeny of plants transformed with CA914 carrying Bcp 1: Cre) for Luciferase expression following selection on

Kan or Bar.

Selection	Luc OFF	Luc ON +	Luc ON++	Luc ON++++	Total	TE (%)
Kan	18	9	0	3	30	0.33
Bar	2	0	0	0	2	0.09
Total	20	9	0	3	32	
%	63	28	0	9		

25

Table 4

Analysis of transformed Arabidopsis seedlings (T1 progeny of plants transformed with CA915 carrying ER: Cre) for Luciferase expression following selection on

Kan or Bar.

Selection	Luc OFF	Luc ON +	Luc ON++	Luc ON++++	Total	TE (%)
Kan	4	2	0	2	8	0.07
Bar	0	1	0	1	2	0.09
Total	4	3	0	3	10	
%	40	30	0	30		

5

Table 5

Analysis of transformed Arabidopsis seedlings (T1 progeny of plants transformed with CA915 carrying ER: Cre) for Luciferase expression following selection on

Kan or Bar.

Selection	Luc OFF	Luc ON +	Luc ON++	Luc ON++++	Total	TE (%)
Kan	0	3	0	1	4	0.02
Bar	2	1	1	6	10	0.24
Total	2	4	1	7	14	
%	14	29	7	50		

10

Table 6

Analysis of transformed Arabidopsis seedlings (T1 progeny of plants transformed with CA928 carrying SAP: Cre) for Luciferase expression following selection on

Kan or Bar.

Selection	Luc OFF	Luc ON +	Luc ON++	Luc ON++++	Total	TE (%)
Kan	20	6	2	1	29	0.22
Bar	5	5	0	0	10	0.39
Total	25	11	2	1	39	
%	64	28	5	3		

15

Table 7

Analysis of transformed Arabidopsis seedlings (T1 progeny of plants transformed with CA929 carrying PI: Cre) for Luciferase expression following selection on Kan or Bar.

Selection	Luc OFF	Luc ON +	Luc ON++	Luc ON++++	Total	TE (%)
Kan	22	23	34	17	96	1.00
Bar	ND	ND	ND	ND	ND	0.84
Total	22	23	34	17	96	
%	23	24	35	18		

5

Table 8

Analysis of T2 progeny seeds from Arabidopsis transformants with CA913 (AP3:Cre) for Resistance to Bar and Kan.

T1		T2 Kanamycin Sensitivity				T2 Bar Sensitivity			
I.D.	Luc	R	S	Ratio R:S	X ² P-Value (fits ratio)	R	S	Ratio R:S	X ² P-Value (fits ratio)
AB01	Off	0	130	0.00		156	0	NA	
AB02	Off	0	137	0.00		137	0	NA	
AB03	Off	0	130	0.00		168	0	NA	
AB04	Off	0	142	0.00		175	0	NA	
AB05	On+	2	136	0.01		97	40	2.43	0.26 (for 3:1)
AB06	On+	0	104	0.00		116	29	4.00	0.16 (for 3:1)
AB07	On+	77	59	1.31	0.12 (for 1:1)	102	39	2.62	0.47 (for 3:1)
AB08	Off	0	137	0.00		96	44	2.18	0.08 (for 3:1)
AB09	On+	8	122	0.07		72	49	1.47	
AB12	Off	0	133	0.00		133	0	NA	
AB14	Off	0	131	0.00		139	0	NA	
AK02	On+++	2	152	0.01		115	49	2.35	0.15 (for 3:1)
AK03	On+	3	158	0.02		153	10	NA	0.95

									(for 15:1)
AK05	Off	163	0	NA		0	179	NA	
AK06	On+++	2	53	0.04		54	9	6.00	
AK08	On+	2	181	0.01		142	38	3.74	0.23 (for 3:1)
AK10	On+	0	150	0.00		114	31	3.68	0.31 (for 3:1)

Table 9

Analysis of T2 progeny seeds from Arabidopsis transformants with CA914 (Bcp 1:Cre) for Resistance to Bar and Kan.

	T1	T2 Kanamycin Sensitivity				T2 Bar Sensitivity			
ID	Luc	R	S	Ratio R:S	X ² P-Value (fits ratio)	R	S	Ratio R:S	X ² P-Value (fits ratio)
BB03	On+	105	39	2.69	0.56 (for 3:1)	84	13	6.46	
BB04	On+	56	98	0.57		39	84	0.46	
BK01	Off	0	157	0.00		1	163	0.01	
BK02	Off	0	155	0.00		0	174	0.00	
BK04	Off	94	57	1.65		107	46	2.33	0.15 (for 3:1)
BK05	On+	69	96	0.72		51	96	0.53	
BK06	Off	0	159	0.00		0	158	0.00	
BK07	Off	106	53	2.00		0	151	0.00	
BK08	Off	8	148	0.05		8	152	0.05	
BK09	On+	66	84	0.79	0.14 (for 1:1)	68	78	0.87	0.41 (for 1:1)
BK11	On+	77	37	2.08	0.07 (for 3:1)	118	25	4.72	
BK17	On+	89	41	2.17	0.09 (for 3:1)	77	16	4.81	0.48 (for 3:1)
BK18	Off	83	53	1.57		90	46	1.96	
BK19	Off	83	52	1.60		94	45	2.09	
BK20	On+	110	32	3.44	0.50 (for 3:1)	91	14	6.50	

BK21	Off	101	39	2.59	0.43 (for 3:1)	124	14	8.86	0.06 (for 15:1)
BK22	Off	51	82	0.62		73	72	1.01	0.93 (for 1:1)
BK23	Off	0	132	0.00		0	137	0.00	
BK24	Off	0	139	0.00		0	144	0.00	

Arabidopsis transformed with the TA29:Cre (pBE855) were selected on bar and selfed. Few lines that showing single insert by 3:1 segregation in T2 were identified and these will be crossed to reporter lines to test the specificity of germline excision. Importantly these results show that there is no toxicity associated with TA29:Cre expression.

EXAMPLE 3

Identifying Promoters for Marker Excision in Common Germline in Tobacco

The Xba I-Bam HI fragment of pBE894 carrying the SCP1 promoter was replaced by an Xba I-Bam HI fragment carrying one of several different developmentally-regulated promoters. These promoters and the resultant binary plasmids are:

Promoter for Heat Shock (HSP) gene from *Arabidopsis thaliana* Col. (pBE917);

Safener inducible promoter, IN 2 (pBE927) described in PCT application WO 99/22003. Promoter for *Apetala 1* (AP1) gene from *Arabidopsis thaliana* Col. (pBE913).

	<u>Promoter</u>	<u>Length</u>	<u>Nucleotide positions</u> (<u>Genbank Accession #</u>)	<u>PCR Primers</u>	
				<u>UP</u>	<u>LP</u>
20	TA29	1525 bp PH815	(from plasmid pTZALG)	PH795	
	HSP18.2 (HSP)	926 bp PH807	50050-50975 (AB006705)	PH806	
25	<i>Apetala 1</i> (AP1)	1850 bp	27937 29807 (AC008262)	P355	P356
	<i>Agamous</i> (AG)	2999 bp	48943-49528 (AL021711)	P353	P354
	<i>Leafy</i> (LFY)	2287 bp	465-2752 (M91208)		

Oligos used for PCR of promoters from *Arabidopsis* genomic DNA.

AP1 PCR primers

P355:

5'-CGT CTA GAC CCG GGA TGT TGT CTT CAA GGC-3' (SEQ ID NO:20)

P356:

5'-ATG GAT CCC GGG TTC TCA CAC ACA GTC TTA-3' (SEQ ID NO:21)

5 TA29 PCR primers

PH795:

5'-CCT CTA GAC CCG GGA-TTA TAT TAG GGA TTT-3' (SEQ ID NO:14)

PH815:

5'-GCG GAT CCC GGG TAG CTA ATT TCT TTA AC-3' (SEQ ID NO:15)

10

AG PCR primers

P353:

5'-CTG CCT AGG TTT CTT CTT CTT CTC GTG CTC TG-3' (SEQ ID NO:22)

P354:

15 5'-GAC CCT AGG CAA TAA TTT TTT TAA AGG AAT TAA TAA GT-3'
(SEQ ID NO:23)

PCR was performed as described above and the desired PCR products were cloned into a PCR cloning vector pGEM-T (Promega Corp., Madison, WI) according to the manufacturer's protocols. After the presence of a cloned insert was established, the ends of the insert were sequenced to confirm that the desired promoter had been amplified and cloned. The cloned promoter fragments were isolated as Xba I- Bam HI fragments and cloned into binary plasmid pBE894.

The binary plasmids were transformed into *Agrobacterium tumefaciens* strain C58 and the resulting transformed agrobacterium, alongwith the earlier ones with AP3:Cre, Bcp1:Cre, ER:Cre used to transform Arabidopsis, were used to transform tobacco (*Nicotiana tabacum* var. Xanthi) by the leaf-disc method as described in PCT application WO 99/22003.

HSP:Cre in pBE894 (pBE917): Transformed leaf discs were selected on kanamycin 300. After 6 weeks, twenty four shoots were excised, placed on rooting medium containing 10 mg/L bar. The shoots were imaged for luciferase expression immediately upon transfer to rooting medium and immediately after a heat shock treatment, which consisted of placing the petri plates containing the shoots in an air incubator at 40°C for 2 hrs for 7 d. The results of luciferase expression is shown in Table 10.

35

Table 10

Luciferase expression in tobacco shoots before and after heat treatment.

Shoots expressing luciferase before heat shock treatment	8/24 (33%)
Shoots expressing luciferase after heats shock treatment	22/24 (91%)
Shoots that showed increased luciferase upon heat shock	19/24 (79%)

- 5 AP3:Cre (pBE913)k: Several shoots were selected on Kanamycin and then rooted. Some of these were tested for luc expression as a reporter for excision. Some rooted plants were not expressing any luciferase and some expressed very high levels of Luciferase expression. All of these plants will be selfed and the progeny used to confirm common or male germline excisions.

- 10 Bcp 1:Cre and ER:Cre: several tobacco shoots were selected on Kan. While Bcp 1 shoots have rooted, those with ER have not. The latter may suggest that ER:Cre expression may be detrimental to root development. All rooted plants will be selfed and analyzed for excision.

CLAIMS

What is claimed is:

1. A trait removal construct comprising:
 - a) A first recombinase element having the general structure P1-R
 - 5 and;
 - b) A second recombinase element selected from the group of general structures consisting of K-TG, and RS-K-TG-RS;

wherein,

 - (i) P1 is a first promoter;
 - 10 (ii) R is a recombinase coding sequence and 3' region
 - (iii) TG is a transgene;
 - (iv) RS is a recombinase site responsive to the recombinase
 - (v) K is selected from the group consisting of;
 - 1) P2-RS-STP-RS and
 - 15 2) P2

wherein P2 is a second promoter, RS is a recombinase site responsive to the recombinase and STP is a stop fragment;

wherein P1 and P2 are operably linked to their down stream elements and wherein P1 is activated prior to P2 in the plant life cycle and wherein

 - 20 expression of the recombinase results in excision of any element contained between the recombinase site responsive to the recombinase.
2. A trait removal construct comprising:
 - a) A first recombinase element selected from the group of general structures consisting of: P1-R1 and P1-R2, and;
 - 25 b) A second recombinase element having the general structure RS2-K-TG-RS2;

wherein,

 - (i) P1 is a first promoter;
 - (ii) R1 is a first recombinase coding sequence and 3' region;
 - 30 (iii) R2 is a second recombinase coding sequence and 3' region;
 - (iv) RS2 is a second recombinase site responsive to the second recombinase;
 - (v) TG is a transgene sequence and 3' region ;
 - (vi) K is selected from the group consisting of:
 - 35 1) P2-RS1-STP-RS1 and
 - 2) P2-RS1-TG

wherein P2 is a second promoter, RS1 is a first recombinase site responsive to the first recombinase, STP is a stop fragment and TG is a transgene sequence and 3' region ;

5 wherein P1 and P2 are operably linked to their down stream elements, and wherein P1 is activated earlier than P2 in the plant life cycle, and wherein expression of the first recombinase results in excision of any element contained between the first recombinase sites responsive to the first recombinase and expression of the second recombinase results in excision of any element contained between the second recombinase sites responsive to the second recombinase.

10 3. A trait removal construct comprising:

- a) a first recombinase element selected from the group of general structures consisting of P1-R1 and RS2 -P1-R1 -RS2 ;
- b) a second recombinase element selected from the group of general structures Z-Y and RS2 -Z-Y-RS2 ;
- 15 c) a third recombinase element selected from the group of general structures Q-X and RS2 -Q-X-RS2 ;

wherein:

- (i) P1 is a first promoter;
- (ii) R1 is a first recombinase coding sequence and 3' region;
- 20 (iii) RS2 is a second recombinase site responsive to a second recombinase;
- (iv) Z has the general formula, P2-RS1-STP-RS1, wherein P2 is a second promoter, RS1 is a first recombinase site responsive to a first recombinase, and STP is a stop fragment;
- 25 (v) Y is selected from the group consisting of R2 and TG wherein R2 is a second recombinase coding sequence and 3' region and TG is a transgene sequence and 3' region ;
- (vi) Q has the general formula P3-RS-STP-RS,
- 30 wherein P3 is a third promoter, RS is a recombinase site selected from the group consisting of RS1 and RS2 , and STP is a stop fragment;
- (vii) X is selected from the group consisting of TG and R wherein TG is a transgene sequence and 3' region and R is a recombinase coding sequence and 3' region selected from the group consisting of R1 and R2;
- 35

wherein P1, P2 and P3 are operably linked to their down stream elements and wherein P1 is activated earlier than P2 in the plant life cycle, and wherein P2 is activated earlier than P3 in the plant life cycle, and wherein

expression of the first recombinase results in excision of any element contained between the first recombinase sites responsive to the first recombinase and expression of the second recombinase results in excision of any element contained between the second recombinase sites responsive to the second recombinase.

5 4. A trait removal construct according to Claim 1, 2 or 3 wherein the first promoter is a germline promoter.

5. A trait removal construct according to Claim 4 wherein the germline promoter is selected from the group consisting of:

- 10 a) constitutive plant promoters;
- b) plant tissue-specific promoters;
- c) plant developmental stage-specific promoters;
- d) inducible plant promoters;
- e) viral promoters;
- f) male germline promoters;
- 15 g) female germline promoters;
- h) common germline promoters;
- i) floral common germline promoters;
- j) vegetative shoot apical meristem promoters; and
- k) floral shoot apical meristem promoters.

20 6. A trait removal construct according to Claim 5 wherein the male germline promoter is derived from genes selected from the group consisting of genes specific to anther primordia and anther sporophyte and to pollen gametophyte.

25 7. A trait removal construct according to Claim 5 wherein the common germline promoter is derived from genes selected from the group consisting of *Apetala 3* (AP3), *Pistillata* (PI), synthetic anther promoter, TA29, and BCP1 and orthologs thereof.

30 8. A trait removal construct according to Claim 5 wherein the common germline promoter is derived from genes selected from the group consisting vegetative and floral shoot apical meristems.

9. A trait removal construct according to Claim 8 wherein the common germline promoter is derived from genes selected from the group consisting of *Leafy* (LFY), *Apetala 3* (AP3), *Pistillata* (PI), *Apetala 1* (AP1), *Agamous* (AG) and *Pistillata* (PI) and orthologs thereof.

35 10. A trait removal construct according to Claim 5 wherein the floral common germline promoter is derived from genes selected from the group consisting of *Agamous* (AG), *Apetala 1* (AP1), *Apetala 3* (AP3), *Leafy* (LFY) and orthologs thereof.

11. A trait removal construct according to Claim 5 wherein the vegetative shoot apical meristem promoter is selected from the group consisting of *Agamous* (AG), *Apetala 1* (AP1), *Apetala 3* (AP3), *Leafy* (LFY), *Aintegumenta* (ANT), *Clavata 3* (CLV3), *Wushel* (WUS), and *Meristemless* (STM) and orthologs thereof.
12. A trait removal construct according to Claim 1, 2 or 3 wherein the second promoter is selected from the group consisting of:
- constitutive plant promoters;
 - plant tissue-specific promoters;
 - plant development stage-specific promoters;
 - inducible plant promoters; and
 - viral promoters.
13. A trait removal construct according to Claim 1, 2 or 3 wherein the transgene is selected from the group consisting of:
- genes encoding a transformation marker;
 - genes encoding a morphological trait;
 - genes conveying sterility;
 - genes conveying specific phenotype on a plant or plant cell; and
 - hormone biosynthetic genes
14. A trait removal construct according to Claim 13 wherein the genes encoding a transformation marker are selected from the group consisting of:
- antibiotic resistance genes;
 - herbicide resistance genes;
 - green fluorescent protein genes;
 - luciferase genes;
 - tumorigenic genes;
 - shoot inducing genes
 - root inducing genes; and
 - selectable genes.
15. A trait removal construct according to Claim 13 wherein the genes encoding a morphological trait are selected from the group consisting of:
- rol C,
 - IPT,
 - Ti plasmid genes,
 - Ri plasmid genes,
 - STM,
 - KNAT,
 - AINTEGUMATA,

- h) Lec1,
- i) *Brassica* "Babyboom"
- j) OSHI, and
- k) Kn1

5 16. A trait removal construct according to Claim 13 wherein the genes encoding hormone biosynthetic genes are cytokinin biosynthesis genes.

 17. A trait removal construct according to Claims 1,2 or 3 wherein the recombinase coding sequences and 3' region R1 and R2 are independently selected from the group consisting of Cre and Flp.

10 18. A trait removal construct according to Claims 1, 2 or 3 wherein the first and second recombinase sites are independently selected from the group consisting of Lox and Frt.

 19. A trait removal construct according to Claims 1, 2 or 3 wherein the first, second and third recombinase elements may be genetically linked or
15 unlinked.

 20. A trait removal construct according to Claim 19 wherein the first, second and third recombinase elements may be genetically unlinked and reside in different plants.

 21. A method for removing a transgene encoding a genetic trait from a
20 plant cell comprising introducing into the plant cell a trait removal construct of Claims 1, 2 or 3 wherein expression of the recombinase results in excision of any element contained between the recombinase site responsive to the recombinase.

 22. A method according to Claim 21 wherein P1 and P2 are selected from the group consisting of:

- 25 a) constitutive plant promoters;
 b) plant tissue-specific promoters;
 c) plant development-stage-specific promoters;
 d) inducible promoters; and
 e) viral promoters.

30 23. A method according to Claim 21 wherein P1 is selected from the group consisting of:

- a) male germline-specific promoters;
 b) female germline-specific promoters;
 c) common germline-specific promoters;
35 d) flower-specific promoters;
 e) vegetative shoot apical meristem-specific promoters; and
 f) floral shoot apical meristem promoters.

24. A method according to Claim 23 wherein P1 is a male germline-specific promoter and the transgene is excised only from the male germline cells to result in pollen without the transgene.

5 25. A method according to Claim 23 wherein P1 is a female germline-specific and the transgene is excised only from the female germline cells to result in the egg cell without the transgene.

26. A method according to Claim 23 wherein P1 is a common germline promoter and the transgene is excised from both male and female germline cells to result in progeny seeds without the transgene.

10 27. A method according to Claim 23 wherein P1 is a floral SAM or flower promoter.

28. A method according to Claim 23 wherein P1 is a vegetative SAM promoter such that the transgene is excised from germline and somatic cells or tissues derived from the vegetative SAM.

15 29. A method according to Claim 22 wherein P1 is an inducible promoter responsive to heat shock.

30. A method according to Claim 22 wherein P1 is an inducible promoter responsive to a chemical safener.

20 31. A method according to Claim 21 wherein the first, second recombinase and third elements are genetically unlinked and reside in different plants.

32. A method for removing a transgene encoding a genetic trait from a plant cell comprising introducing into the plant cell a trait removal construct of Claim 3 wherein expression of the recombinase results in excision of any element
25 contained between the recombinase site responsive to the recombinase.

33. A method according to Claim 21 wherein TG is a transgene selected from the group consisting of:

- 30
- a) genes encoding a transformation marker;
 - b) genes encoding a morphological trait;
 - c) genes conveying sterility;
 - d) genes conveying specific phenotype on a plant or plant cell; and
 - e) hormone biosynthetic genes.

34. A method according to Claim 33 wherein the transgene is selected from the group consisting of:

- 35
- a) antibiotic resistance genes;
 - b) herbicide resistance genes;
 - c) green fluorescent protein genes;
 - d) luciferase genes;

- e) tumorigenic genes;
- f) root inducing genes; and
- g) selectable genes.

35. A method according to Claim 34 wherein the transgene is selected
 5 from the group consisting of:

- a) IPT;
- b) Ti plasmid genes; and
- c) Ri plasmid genes.

36. A method for conditionally activating a transgene in a second
 10 generation plant comprising:

- 1) providing a construct comprising:
 - a) a first recombinase element having the general structure P1-R1;
 - b) a second recombinase element having general structures
 15 P2-RS1-STP-RS1-R2;
 - c) a third recombinase element having the general structure P3-RS2-STP-RS2-TG;

wherein:

- (i) P1 is a first promoter;
- (ii) R1 is a first recombinase coding sequence and 3' region;
- (iii) RS1 is a first recombinase site responsive to a first recombinase;
- (iv) P2 is a second promoter,
- (v) RS2 is a second recombinase site responsive to a second
 25 recombinase;
- (vi) STP is a stop fragment;
- (vii) R2 is a second recombinase coding sequence and 3' region
- (viii) TG is a transgene sequence and 3' region ; and
- (vi) P3 is a third promoter;

wherein P1, P2 and P3 are operably linked to their down stream elements;

- 2) providing a transgenic plant comprising the first, second and third recombinase elements;
- 3) activating P1 such that the R1 recombinase coding sequence is
 35 expressed in the common germline of a first generation plant, wherein expression of R1 excises the stop fragment from the second recombinase element;

- 4) activating P2 such that R2 is expressed either in the common germline of the first generation plant or in the second generation plant, wherein expression of R2 excises the stop fragment from the third recombinase element allowing expression of the transgene in the second and all subsequent generations of plants.

37. A method for conditionally activating a transgene in a second generation plant comprising:

- 1) providing a construct comprising:
 - a) a first recombinase element having the general structure P1-R1;
 - b) a second recombinase element having general structures P2-RS1-STP-RS1-R2;
 - c) a third recombinase element having the general structure P3-RS2-STP-RS2-TG;

wherein:

- (i) P1 is a first promoter;
- (ii) R1 is a first recombinase coding sequence and 3' region;
- (iii) RS1 is a first recombinase site responsive to a first recombinase;
- (iv) P2 is a second promoter,
- (v) RS2 is a second recombinase site responsive to a second recombinase;
- (vi) STP is a stop fragment;
- (vii) R2 is a second recombinase coding sequence and 3' region;
- (viii) TG is a transgene sequence and 3' region ; and
- (vi) P3 is a third promoter;

wherein P1, P2 and P3 are operably linked to their down stream elements;

- 2) providing a transgenic plant comprising the third recombinase element;
- 3) transforming the transgenic plant of (2) with either the second recombinase element to generate a first plant or the third recombinase element to generate a second plant;
- 4) crossing the first and second plants such that expression of R1 under the control of P1 in the common germline of the first generation, excises the stop fragment from the second recombinase element allowing expressing of R2 under the control of P2 which in turn, excises the stop fragment from the

third recombinase element, permitting expression of the trait gene(s) under the control of P3 in the second and subsequent generation(s).

38. A method for conditionally activating a transgene in a second generation plant comprising:

- 1) providing a construct comprising:
 - a) a first recombinase element having the general structure P1-R1;
 - b) a second recombinase element having general structures P2-RS1-STP-RS1-R2;
 - c) a third recombinase element having the general structure P3-RS2-STP-RS2-TG;

wherein:

- (i) P1 is a first promoter;
- (ii) R1 is a first recombinase coding sequence and 3' region;
- (iii) RS1 is a first recombinase site responsive to a first recombinase;
- (iv) P2 is a second promoter,
- (v) RS2 is a second recombinase site responsive to a second recombinase;
- (vi) STP is a stop fragment;
- (vii) R2 is a second recombinase coding sequence and 3' region;
- (viii) TG is a transgene sequence and 3' region ; and
- (vi) P3 is a third promoter;

wherein P1, P2 and P3 are operably linked to their down stream elements and wherein, P1 is a germline promoter that can be induced in the first generation of plant life cycle;

- 2) providing a transgenic plant comprising the first, second and third recombinase elements;
- 3) inducing the first promoter such that expression of R1 under the control P1 in the common germline of the first generation, excises the stop fragment from the second recombinase element allowing expressing of R2 under the control of P2, which, in turn, excises the stop fragment from the third recombinase element, permitting expression of the trait gene(s) under the control of P3 promoter in in the second and subsequent generation(s).

39. A method for conditionally activating a transgene in a second generation plant comprising:

- 1) providing a construct comprising:
 - a) a first recombinase element having the general structure P1-R1;
 - b) a second recombinase element having general structures P2-RS1-STP-RS1-R2;
 - c) a third recombinase element is selected from the group consisting of the general structures P3-RS2-STP-RS2-TG1 and P4-RS2-STP-RS2-TG2;
- wherein:
- (i) P1 is a first promoter;
 - (ii) R1 is a first recombinase coding sequence and 3' region;
 - (iii) RS1 is a first recombinase site responsive to a first recombinase;
 - (iv) P2 is a second promoter,
 - (v) RS2 is a second recombinase site responsive to a second recombinase;
 - (vi) STP is a stop fragment;
 - (vii) R2 is a second recombinase coding sequence and 3' region;
 - (viii) TG1 is a first transgene sequence and 3' region ;
 - (ix) TG2 is a second transgene sequence and 3' region;
 - (ix) P3 is a third promoter; and
 - (x) P4 is a fourth promoter;

wherein P1, P2, P3 and P4 are operably linked to their down stream elements and wherein TG1 and TG2 are different trait transgenes and wherein P3 and P4 are activated in a second generation plant;

- 2) providing a first plant comprising the first and third recombinase elements;
- 3) providing a second plant comprising the second and third recombinase elements;
- 4) crossing the first and second plants to produce a first generation plant wherein conditional expression of the first recombinase coding sequence (R1) under the control of the P1 promoter in the common germline of the first generation, excises the stop fragment from the second recombinase element allowing expressing of the second recombinase coding sequence and 3'

region (R2) under the control of P2 promoter, which recombinate, in turn, excises the stop fragments from the two third recombinate elements, permitting expression of the trait gene(s) TG1 and TG2 under the control of P3 and P4 promoter, respectively, in the second generation.

40. A method according to Claim 39 wherein, TG1 is a trait gene, TG2 is a lethal gene that blocks plant development, and the third promoter (P3) is expressed earlier in the plant life cycle than the fourth promoter (P4) in the second generation.

41. A method according to Claim 39 wherein TG1 is a trait gene, TG2 is a sterility gene that prevents pollen formation and seed set, and the third promoter (P3) is expressed earlier than P4 in the next generation

42. A method according to Claim 36 wherein the first promoter is a constitutive and common germline promoter, and the second promoter is floral common germline promoter; and the third promoter is seed-specific; wherein conditional expression of the first recombinate coding sequence in common germline under the control of the first promoter results in expression of the second recombinate coding sequence under the control of the second promoter in floral common germline, and wherein expression of the second recombinate results in the expression of the trait in the progeny seed.

43. A method according to Claim 37 where the first promoter is inducible and responsive to an inducing agent.

44. A method for conditional and transient expression of a trait transgene comprising:

- 1) providing a construct comprising:
 - a) a first recombinate element having the general structure P1-R1;
 - b) a second recombinate element having the general structure S2-P2- RS1-STP-RS1-TG-RS2 ;
 - c) a third recombinate element having the general structure P3-RS1-STP-RS1-R2

wherein:

- (i) R1 is a first recombinate coding sequence and 3' region;
- (ii) RS1 is a first recombinate site responsive to a first recombinate;
- (iii) R2 is a second recombinate coding sequence and 3' region;

- (iv) RS2 is a second recombinase site responsive to a second recombinase
- (iv) STP is a stop fragment;
- (v) TG is trait transgene sequence and 3' region;
- 5 (vi) P1 is a first promoter which will be activated in the germline of the first generation;
- (vii) P2 is a second promoter which will be activated in somatic and/or germline cells and may be activated prior to or at the same time as P3;
- 10 (viii) P3 is a third promoter which is activated in the germline;
- wherein P1, P2 and P3 are operably linked to their down stream elements, and
- 2) providing a first generation transgenic plant comprising all three recombinase elements;
 - 15 3) activating the first promoter (P1) in the first generation plant of (2) wherein the first recombinase coding sequence is expressed priming both the second and third recombinase elements;
 - 4) activating the second promoter wherein the transgene is expressed; and
 - 20 5) activating the third promoter wherein the second recombinase coding sequence is expressed excising the second recombinase element.

45. A method for conditional and transient expression of a trait transgene comprising:
- 25 1) providing a construct comprising:
- a) a first recombinase element having the general structure P1-R1;
 - b) a second recombinase element having the general structure RS2-P2- RS1-STP-RS1-TG-RS2 ;
 - 30 c) a third recombinase element having the general structure P3-RS1-STP-RS1-R2

- wherein:
- (i) R1 is a first recombinase coding sequence and 3' region;
 - (ii) RS1 is a first recombinase site responsive to a first
 - 35 recombinase;
 - (iii) R2 is a second recombinase coding sequence and 3' region;

- (iv) RS2 is a second recombinase site responsive to a second recombinase
- (iv) STP is a stop fragment;
- (v) TG is trait transgene sequence and 3' region;
- 5 (vi) P1 is a first promoter which will be activated in the germline of the first generation;
- (vii) P2 is a second promoter which will be activated in somatic and/or germline cells and may be activated prior to or at the same time as P3;
- 10 (viii) P3 is a third promoter which is activated in the germline

wherein P1, P2 and P3 are operably linked to their down stream elements, and

- 2) providing a first plant comprising the first recombinase element;
- 3) providing a second plant comprising the second and third
15 recombinase elements, wherein the second plant is made homozygous for the elements;
- 4) crossing the first and second plants to produce a first generation plant wherein the first promoter (P1) is activated, expressing the first recombinase coding sequence (R1) which excises the stop
20 fragments from the second and third recombinase elements sequentially activating the trait (TG) gene and the second (R2) recombinase coding sequence, which in turn removes the second recombinase element from the chromosome, permitting transient expression of the transgene.

25 46. A method according to Claims 44 wherein P1 is a first germline promoter that is induced in the first generation of plant life cycle; and wherein all three recombinase elements are introduced into a plant either by co-transformation, sequential transformations, or genetic crosses after single transformation in a plant.

30 47. A method according to any one of Claims 44-46 wherein the first promoter (P1) is selected from the group consisting of constitutive promoter, seed promoter, a vegetative or floral shoot apical meristem promoter, and floral common germline promoter.

35 48. A method according to Claim 47 wherein the P3 promoter is male germline-specific promoter such that the TG is removed from only the pollen of the first generation.

49. A method according to Claim 47 wherein the P3 promoter is expressed in the floral common germline to result in TG removal from both pollen and the progeny seed of the first generation.

50. A method according to Claim 44 wherein:

- 5 a) P1 is a floral common germline promoter that expresses the first recombinase coding sequence (R1) in the common germline of the first generation and is not expressed in seed or germinating seed;
- 10 b) P2 expresses the transgene (TG) in the seed of the second generation; and
- c) P3 expresses the second recombinase coding sequence (R2) in the common germline of germinating seeds of the second generation to result in the transgene (TG) removal from both pollen and the seed of the second generation.

15 51. A method for conditional expression of male sterility in the first generation plant comprising:

- 1) providing a construct comprising:
 - a) a first recombinase element having the general structure P1-R1;
 - 20 b) a second recombinase element having the general structure RS2-P2- RS1-STP-RS1-TG-RS2 ;
 - c) a third recombinase element having the general structure P3-RS1-STP-RS1-R2;

wherein:

- 25 (i) R1 is a first recombinase coding sequence and 3' region;
- (ii) RS1 is a first recombinase site responsive to a first recombinase;
- (iii) R2 is a second recombinase coding sequence and 3' region;
- 30 (iv) RS2 is a second recombinase site responsive to a second recombinase;
- (v) STP is a stop fragment;
- (vi) TG is transgene sequence and 3' region encoding male sterility;
- 35 (vii) P1 promoter is a common germline promoter which is activated in vegetative SAM, floral SAM, or flower of the first generation and is not expressed in seed;

(viii) P2 is a second sporophytic anther or anther primordia promoter;

(ix) P3 is a third seed-specific promoter that is activated in the germline of the second generation and

5 wherein P1, P2 and P3 are operably linked to their down stream elements, and

2) providing a first transgenic plant comprising the first recombinase element;

3) providing a second transgenic plant comprising the second and third recombinase elements in a homozygous state;

10 4) crossing the first and second plants to produce a first generation plant wherein the expression of the first recombinase under the control of the first promoter excises the stop fragment from both the second and third recombinase elements, causing the male sterility transgene to be expressed;

15 5) crossing the first generation plant of step (4) with a male fertile inbred to produce a second generation plant, wherein activation of the seed specific third promoter expresses the second recombinase, excising the male sterility transgene and restoring male fertility.

20 52. A method for expressing conditional male sterility in the first and second generations comprising:

1) providing a construct comprising:

a) a first recombinase element having the general structure P1-R1 ;

25 b) a second recombinase element having the general structure P2- RS1-STP-RS1-MS ;

c) a third recombinase element having the general structure P2-RS2-STP-RS2-MS;

30 d) a fourth recombinase element having the general structure P1-R2;

wherein:

(i) R1 is a first recombinase;

(ii) RS1 is a first recombinase site responsive to a first recombinase

35 (iii) R2 is a second recombinase;

(iv) RS2 is a second recombinase site responsive to a second recombinase;

(v) STP is a stop fragment;

(vi) TG is a transgene encoding male sterility with 3' region;
and

(vii) P1 is germline promoter that is activated in the plant
life cycle earlier than, or at the same time as, P2 and is
selected from the group consisting of; a constitutive
promoter, seed-specific promoter, and a vegetative shoot
apical meristem promoter;

(viii) P2 is a promoter which expresses the transgene,

wherein P1 and P2 are operably linked to their down stream elements;

- 2) providing a first plant comprising the first and third recombinase elements;
- 3) providing a second plant comprising the second and third recombinase elements;
- 4) providing a third plant comprising the fourth recombinase element;
- 5) crossing the first and second plants to obtain first progeny in which expression of the first recombinase under the control of the P1 promoter of the first recombinase element excises the stop fragment from the second recombinase element, allowing the expression of the transgene encoding male sterility;
- 6) crossing the progeny of step (5) with the third plant to obtain second progeny in which expression of the second recombinase under the control of the P1 promoter in the fourth recombinase element excises the stop fragment from the third recombinase element, allowing the expression of the male sterility gene.

53. A method according to Claim 49 wherein the P2 promoter is an anther- or anther primordia-specific promoter that is selected from the group consisting of a tapetum-specific promoter, and a male germline-specific promoter.

54. A method for expressing conditional male sterility in a plant comprising:

- 1) providing constructs comprising:
 - a) a first recombinase element having the general structure P1-R1 ;
 - b) a second recombinase element having the general structure RS2-P2- RS1-STP-RS1-TG -RS2-SG;
 - c) a third recombinase element having the general structure P1-R2 ;

wherein:

- (i) R1 is a first recombinase;
- (ii) RS1 is a first recombinase site responsive to a first recombinase;
- (iii) RS2 is a second recombinase site responsive to a second recombinase;
- (iv) STP is a stop fragment;
- (v) SG is a plant selectable gene; and
- (vi) TG is a transgene sequence and 3' region encoding male sterility;
- (vii) P1 is a common germline promoter that is activated in the plant life cycle earlier than or at the same time as P2 promoter and is selected from the group consisting of; a constitutive promoter, seed-specific promoter, and a vegetative or floral shoot apical meristem promoter;
- (viii) P2 is an anther- or anther primordia-specific promoter that is activated in the life cycle at the same time as or later than P1 promoter and is selected from the group consisting of a tapetum-specific promoter, male germline-specific promoter;
- wherein P1, and P2 are operably linked to their down stream elements;
- 2) providing a first plant comprising the first recombinase element;
 - 3) providing a second plant comprising the second recombinase element;
 - 4) providing a third plant comprising the third recombinase element;
 - 5) crossing the first and second plants to produce progeny in which expression of the first recombinase under the control of P1 promoter excises the stop fragment from the second recombinase element permitting the expression of the male sterility gene under the control of the P2 promoter;
 - 6) crossing the progeny of step (5) with a male fertile plant to produce progeny seeds;
 - 7) selecting progeny seeds of step (6) on the basis of the gene product of the selectable gene which ensures male sterility; and
 - 8) crossing the male sterile progeny of step (7) with the third plant of step (4) to produce a progeny in which expression of the second recombinase under the control of the P1 promoter

excises the male sterility trait from the third recombinase element restoring male fertility in the hybrid progeny.

55. A method for transgene removal comprising:

1) providing a construct comprising:

- 5 a) a first recombinase element having the general structure P1-R1;
b) a second recombinase element having the general structure RS1-P2-TG-RS1 ;

wherein:

- 10 (i) R1 is a first recombinase coding sequence and 3' region;
(ii) RS1 is a first recombinase site responsive to a first recombinase;
(iii) TG is transgene sequence and 3' region encoding either a trait or a transformation marker;
15 (iv) P1 is a first promoter that expresses R1 in the germline of the first generation;
(vi) P2 is a second promoter, wherein P2 is activated before or at the same as P1;

wherein P1, and P2 are operably linked to their down stream elements;

- 20 2) providing a first plant comprising the first recombinase element;
3) providing a second plant comprising the second recombinase element;
4) crossing the first and second plants to create a first generation plant wherein the first promoter (P2) is activated, expressing the transgene;
25 5) activating the first promoter (P1) wherein the recombinase is expressed, resulting in the removal of the transgene from the chromosome.

56. A method according to Claim 55 wherein P1 is a common germline selected from the group consisting of *Apetala 3*, *Pistillata*, *Bcp I*, *Apetala 1*, *Leafy*, and *Agamous* genes .

57. A method according to Claim 55 wherein P1 is a male germline promoter which excises the transgene from the pollen.

58. A method according to Claim 55 wherein P1 is a floral common germline promoter that expresses the recombinase to excises the transgene from the progeny seed.

59. A method according to Claim 57 wherein P1 is a promoter derived from *Arabidopsis* selected from the group consisting of *Apetala* 3 (AP3), *Pistillata* (PI), *Bcp* I genes or a synthetic anther promoter.

60. A method according to Claim 58 wherein P1 is a promoter derived from *Arabidopsis* selected from the group consisting of *Apetala* 3 (AP3), *Pistillata* (PI), *Leafy*, *Agamous*, *Apetala* 1.

61. A method for transformation marker excision comprising:

- 1) providing a construct comprising a recombinase element having the general structure RS1-P1-R1-P2-TG1-RS1 P3-TG2;

wherein:

- (i) R1 is a recombinase coding sequence and 3' region;
- (ii) RS1 is a recombinase site responsive to recombinase R1;
- (iii) TG1 is transgene sequence and 3' region encoding a transformation marker and 3' region;
- (iv) TG2 is trait transgene sequence and 3' region;
- (v) P1 is a first promoter that is activated in the germline of the primary transformant;
- (vi) P2 is a second promoter that is activated in the primary transformant;
- (vii) P3 is a third promoter;

wherein P1, P2 and P3 are operably linked to their downstream elements;

- 2) providing a transgenic plant comprising the recombinase element;
- 3) activating the second promoter (P2) whereby the transgene encoding a marker (TG1) is expressed;
- 4) activating the first promoter (P1) whereby the recombinase (R1) is expressed resulting in the excision of the transgene encoding a marker (TG1); and
- 5) activating the third promoter (P3) whereby the trait transgene (TG2) is expressed.

62. A method according to Claim 61, wherein P1 is a common germline promoter that is inducible.

63. A method according to Claim 61, wherein P1 is a common germline promoter from *Arabidopsis* heat shock promoter.

64. A method according to Claim 61, wherein P1 is a common germline promoter from safener inducible promoter IN-2.

65. A method according to Claim 61, wherein P1 is a common germline promoter from developmentally regulated genes of shoot apical meristem, floral germline, or male germline cells.

5 66. A method according to Claim 61, wherein P1 is a common germline promoter derived from vegetative and floral shoot apical meristem, flower, or male germline wherein the vegetative and floral shoot apical meristem promoter is selected from the group consisting of *Leafy*, *Apetala 1*, *Apetala 3* (AP3), *Pistillata* (PI) or their orthologs from other species, the flower promoter is selected from the group consisting of synthetic anther promoter and Bcp I genes or their orthologs
10 from other species, and the male germline promoter is selected from the group consisting of *Apetala 3* (AP3), *Pistillata* (PI) or their orthologs from other species.

67. A method according to Claim 57, wherein the transformation marker is a morphological trait, hormone biosynthetic gene, selectable gene.

15 68. A method for the conditional gametophytic expression of male sterility comprising:

- 1) providing a construct comprising:
 - a) a first recombinase element having the general structure P1-R1;
 - b) a second recombinase element having the general structure
20 P1-RS1-STP-RS1-R2; and
 - c) a third recombinase element having the general structure P2-RS2-STP-RS2-TG1;
 - d) a linked gene construct having the general formula, TG2-TG3;

25 wherein:

- (i) P1 is a first promoter;
- (ii) R1 is a first recombinase coding sequence and 3' region;
- (iii) RS1 is a first recombinase sequence responsive to a first recombinase;
- (iv) P2 is a second pollen specific promoter,
- (v) RS2 is a second recombinase sequence responsive to a second recombinase;
- (vi) STP is a stop fragment;
- (vii) R2 is a second recombinase coding sequence and
35 3' region;
- (viii) TG1 is a transgene sequence and 3' region encoding gametophytic male sterility;

(ix) TG2 is a transgene sequence and 3' region encoding a plant trait;

(x) TG3 is a gametophytic male fertility restorer gene;

wherein P1 and P2 are operably linked to their down stream elements;

- 5 2) providing a transgenic plant comprising the construct of step (1);
- 3) activating the first promoter (P1) whereby the recombinase (R1) is expressed resulting in the excision of the stop fragment (STP) from the second recombinase element, expression of the second recombinase and excision of the stop fragment (STP) from the
- 10 third recombinase element;
- 4) activation of the pollen specific promoter (P2) where in the male sterility transgene (TG1) is expressed in the pollen;
- 5) selfing the plant of step (4) wherein TG2 and TG3 are expressed in the resulting second generation plant.

15 69. A method according to Claim 68, wherein TG1 is barnase and TG3 is barstar.

70. A method for the conditional expression or excision of a transgene in a plant comprising:

- 20 (i) providing a multiplicity of recombinase elements, each recombinase element comprising:
 - a) at least one promoter;
 - b) a DNA fragment;
- wherein the DNA fragment comprises at least one genetic element selected from the group consisting of, a recombinase coding sequence, a
- 25 site-specific recombinase sequence responsive to a recombinase; a stop fragment and a transgene;
- (ii) introducing at least two of the recombinase elements of step (i) into at least one plant wherein the at least two recombinase elements are selected from the group consisting of
- 30 a) a recombinase element having a first recombinase under the control of a promoter; and
- b) a recombinase element having a second recombinase under the control of a promoter whose expression is dependent on the expression of the first recombinase;
- 35 (iii) activating at the promoter of step (ii)(a) wherein the expression of the recombinase is effected by the expression of the first recombinase.

71. A method for the germline specific expression of a transgene comprising:

- (i) providing a construct comprising a homeotic promoter operably linked to a transgene;
- 5 (ii) introducing the construct of step (i) into at least one plant; and
- (iii) activating the promoter whereby the transgene is expressed.

71. A method according to Claim 71 wherein the homeotic promoter is selected from the group consisting of AP3, BCP1, PI, SAP and TA29.


72. A trait removal construct comprising:

- 10 a) a first recombinase element comprising a first promoter operably linked to a sequence encoding a first recombinase;
- b) a second recombinase element comprising a second promoter, a stop fragment bounded by site specific sequences responsive to the first recombinase and a sequence encoding a second
- 15 recombinase wherein the presence of the stop fragment inhibits expression of the second recombinase, and wherein the first and second recombinases are different; and
- c) a DNA molecule bounded by site specific sequences responsive to the second recombinase;


20 wherein expression of the first recombinase excises the stop fragment from the second recombinase element, operably linking the second promoter and the sequence encoding the second recombinase, and wherein expression of the second recombinase results in site specific recombination within the DNA molecule bounded by site specific sequences responsive to the second recombinase.

25

FIGURE 1



Stage 1: R1 recombinease expression occurs under promoter (P1) primes trait gene (TG)



Stage 2: TG expression occurs under P2 promoter, e.g. mature leaf promoter



FIGURE 2









FIGURE 3











RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

MS

male sterile F1 hybrid

X

RS2

P2 promoter

MS

male sterile F1 hybrid

↓

Top-Cross[®] hybrid

RS1

P2 promoter

SEQUENCE LISTING

<110> E.I. DuPont de Nemours Company, Inc.

<120> METHODS FOR CONDITIONAL TRANSGENE EXPRESSION AND TRAIT REMOVAL IN PLANTS

<130> CL1127 PCT2

<160> 23

<170> Microsoft Office 97

<210> 1

<211> 46

<212> DNA

<213> primer

<400> 1

tagcatacat tatacgaagt tattagaaga cgccaaaaac ataaag

46

<210> 2

<211> 44

<212> DNA

<213> primer

<400> 2

cgacgcactc cttctttagg taccgaatta cacggcgatc tttc

44

<210> 3

<211> 20

<212> DNA

<213> primer

<400> 3

ccaaaagaga tctcctttgc

20

<210> 4

<211> 40

<212> DNA

<213> primer

<400> 4

gtaccctagg taccgttcgt ataattgtatg ctatacgaag

40

<210> 5

<211> 20

<212> DNA

<213> primer

<400> 5

ttcacacaaa cggtgatacg

20

<210> 6
<211> 56
<212> DNA
<213> primer

<400> 6
cttcgtatag catacattat acgaagttat cctaggaaaa ggagaggggcc aagagg 56

<210> 7
<211> 60
<212> DNA
<213> primer

<400> 7
cttcgtataa tgtatgctat acgaagttat ttacctaggc atatgatcca atatcaaagg 60

<210> 8
<211> 30
<212> DNA
<213> primer

<400> 8
agtctagacc cgggatggaa gtgacgatta 30

<210> 9
<211> 28
<212> DNA
<213> primer

<400> 9
gaggatcccg ggtcttctct ctttgttt 28

<210> 10
<211> 30
<212> DNA
<213> primer

<400> 10
tatctagacc cgggtctcga tccgatcgaa 30

<210> 11
<211> 30
<212> DNA
<213> primer

<400> 11
ttggatcccg ggttctctct ctctttctta 30

<210> 12
<211> 29
<212> DNA
<213> primer

<400> 12
ggtctagacc cgggactttt tgagaaaag 29

<210> 13
<211> 30
<212> DNA
<213> primer

<400> 13
atggatcccg ggttctcaca cacagtctta 30

<210> 14
<211> 30
<212> DNA
<213> primer

<400> 14
cctctagacc cgggattata ttagggattt 30

<210> 15
<211> 29
<212> DNA
<213> primer

<400> 15
gcggatcccg ggtagctaatttctttaac 29

<210> 16
<211> 30
<212> DNA
<213> primer

<400> 16
cgtctagacc cgggatgttg tcttcaaggc 30

<210> 17
<211> 30
<212> DNA
<213> primer

<400> 17
atggatcccg ggttctcaca cacagtctta 30

<210> 18
<211> 29
<212> DNA
<213> primer

<400> 18
tttctagacc cggggaaaag agaccaagc 29

<210> 19
<211> 28
<212> DNA
<213> primer

<400> 19
ttgatcccc ggtgttcgt tgcttttc 28

<210> 20
<211> 30
<212> DNA
<213> primer

<400> 20
cgtctagacc cgggatgttg tcttcaaggc 30

<210> 21
<211> 30
<212> DNA
<213> primer

<400> 21
atggatcccg ggttctcaca cacagtctta 30

<210> 22
<211> 32
<212> DNA
<213> primer

<400> 22
ctgcctaggt ttcttcttct tctcgtgctc tg 32

<210> 23
<211> 38
<212> DNA
<213> primer

<400> 23
gaccctaggc aataattttt ttaaaggaat taataagt 38