



(86) Date de dépôt PCT/PCT Filing Date: 1999/11/10
(87) Date publication PCT/PCT Publication Date: 2000/05/18
(85) Entrée phase nationale/National Entry: 2001/05/08
(86) N° demande PCT/PCT Application No.: US 99/26592
(87) N° publication PCT/PCT Publication No.: WO 00/28031
(30) Priorités/Priorities: 1998/11/10 (60/107,911) US;
1999/08/17 (60/149,332) US; 1999/08/27 (60/151,242) US


(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/12, A61K 48/00, A61K 38/17,
A61K 38/44, C07K 14/47, C07K 16/40, C07K 16/18,
C12N 9/02, C12N 15/53, G01N 33/50

(71) Demandeur/Applicant:
EMORY UNIVERSITY, US

(72) Inventeurs/Inventors:
GUANGJIE, CHENG, US;
ARNOLD, REBECCA S., US;
LAMBETH, J. DAVID, US;
GRIENDLING, KATHY K., US;
LASSEGUE, BERNARD P., US

(74) Agent: BERESKIN & PARR

(54) Titre : NOUVEAUX REGULATEURS MITOGENIQUES
(54) Title: NOVEL MITOGENIC REGULATORS



(57) **Abrégé/Abstract:**

The present invention relates to new genes encoding for the production of novel proteins involved in generation of reactive oxygen intermediates that affect cell division. The present invention also provides vectors containing these genes, cells transfected with these vectors, antibodies raised against these novel proteins, kits for detection, localization and measurement of these genes and proteins, and methods to determine the activity of drugs to affect the activity of the proteins of the present invention.





PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, 15/53, C07K 16/40, 16/18, 14/47, C12N 9/02, A61K 38/17, 38/44, 48/00, G01N 33/50	A3	(11) International Publication Number: WO 00/28031 (43) International Publication Date: 18 May 2000 (18.05.00)
--	-----------	---


(21) International Application Number: **PCT/US99/26592**(22) International Filing Date: **10 November 1999 (10.11.99)**

(30) Priority Data:

60/107,911	10 November 1998 (10.11.98)	US
60/149,332	17 August 1999 (17.08.99)	US
60/151,242	27 August 1999 (27.08.99)	US

(71) Applicant: **EMORY UNIVERSITY [US/US]; 2009 Ridgewood Drive, Atlanta, GA 30322 (US).**(72) Inventors: **LAMBETH, J., David; 461 Emory Drive, Decatur, GA 30207 (US). LASSEGUE, Bernard, P.; 564 Emory Oaks Way, Decatur, GA 30033 (US). GRIENDLING, Kathy, K.; 1819 Kanawha Trail, Stone Mountain, GA 30087 (US). ARNOLD, Rebecca, S.; 4344 Bridle Path Court, Tucker, GA 30084 (US). GUANGJIE, Cheng; Apt 1, 2751 Briarcliff Road, Atlanta, GA 30329 (US).**(74) Agents: **ELSEVIER, Lisa, C. et al.; Jones & Askew, LLP, 2400 Monarch Tower, 3424 Peachtree Road, N.E., Atlanta, GA 30326 (US).**(81) Designated States: **AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**

Published

With international search report.(88) Date of publication of the international search report:
23 November 2000 (23.11.00)(54) Title: **MITOGENIC REGULATORS**

(57) Abstract

The present invention relates to new genes encoding for the production of novel proteins involved in generation of reactive oxygen intermediates that affect cell division. The present invention also provides vectors containing these genes, cells transfected with these vectors, antibodies raised against these novel proteins, kits for detection, localization and measurement of these genes and proteins, and methods to determine the activity of drugs to affect the activity of the proteins of the present invention.

5

NOVEL MITOGENIC REGULATORS

The U.S. Government has a paid-up license in this invention and the right in limited circumstances to require the patent owner to license others on reasonable terms as provided for by the terms of National Institutes of Health grants HL38206 and HL58000.

10

TECHNICAL FIELD

15

The present invention relates to the field of normal and abnormal cell growth, in particular mitogenic regulation. The present invention provides the following: nucleotide sequences encoding for the production of enzymes that are mitogenic regulators; amino acid sequences of these enzymes; vectors containing these nucleotide sequences; methods for transfecting cells with vectors that produce these enzymes; transfected cells; methods for administering these transfected cells to animals to induce tumor formation; and antibodies to these enzymes that are useful for detecting and measuring levels of these enzymes, and for binding to cells possessing extracellular epitopes of these enzymes.

20

25

BACKGROUND OF THE INVENTION

30

Reactive oxygen intermediates (ROI) are partial reduction products of oxygen: 1 electron reduces O_2 to form superoxide (O_2^-), and 2 electrons reduce O_2 to form hydrogen peroxide (H_2O_2). ROI are generated as a byproduct of aerobic metabolism and by toxicological mechanisms. There is growing evidence for regulated enzymatic generation of O_2^- and its conversion to H_2O_2 in a variety of cells. The conversion of O_2^- to H_2O_2 occurs spontaneously, but is markedly accelerated by superoxide dismutase (SOD). High levels of ROI are associated with damage to biomolecules such as DNA, biomembranes and proteins. Recent evidence indicates generation of ROI under

35

normal cellular conditions and points to signaling roles for O_2^- and H_2O_2 .

5 Several biological systems generate reactive oxygen. Phagocytic cells such as neutrophils generate large quantities of ROI as part of their battery of bactericidal mechanisms. Exposure of neutrophils to bacteria or to various soluble mediators such as formyl-Met-Leu-Phe or phorbol esters activates a massive consumption of oxygen, termed the respiratory burst, to initially generate superoxide, with
10 secondary generation of H_2O_2 , HOCl and hydroxyl radical. The enzyme responsible for this oxygen consumption is the respiratory burst oxidase (nicotinamide adenine dinucleotide phosphate-reduced form (NADPH) oxidase).

15 There is growing evidence for the generation of ROI by non-phagocytic cells, particularly in situations related to cell proliferation. Significant generation of H_2O_2 , O_2^- , or both have been noted in some cell types. Fibroblasts and human endothelial cells show increased release of superoxide in response to cytokines such as interleukin-1 or tumor necrosis
20 factor (TNF) (Meier et al. (1989) *Biochem J.* 263, 539-545.; Matsubara et al. (1986) *J. Immun.* 137, 3295-3298). Ras-transformed fibroblasts show increased superoxide release compared with control fibroblasts (Irani, et al. (1997) *Science*
25 275, 1649-1652). Rat vascular smooth muscle cells show increased H_2O_2 release in response to PDGF (Sundaresan et al. (1995) *Science* 270, 296-299) and angiotensin II (Griendling et al. (1994) *Circ. Res.* 74, 1141-1148; Fukui et al. (1997) *Circ. Res.* 80, 45-51; Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321), and H_2O_2 in these cells is associated with
30 increased proliferation rate. The occurrence of ROI in a variety of cell types is summarized in Table 1 (adapted from Burdon, R. (1995) *Free Radical Biol. Med.* 18, 775-794).

Table 1

	<u>Superoxide</u>	<u>Hydrogen Peroxide</u>
	human fibroblasts	Balb/3T3 cells
	human endothelial cells	rat pancreatic islet cells
5	human/rat smooth muscle cells	murine keratinocytes
	human fat cells	rabbit chondrocytes
	human osteocytes	human tumor cells
	BHK-21 cells	fat cells, 3T3 L1 cells
	human colonic epithelial cells	
10		
15	ROI generated by the neutrophil have a cytotoxic function. While ROI are normally directed at the invading microbe, ROI can also induce tissue damage (e.g., in inflammatory conditions such as arthritis, shock, lung disease, and inflammatory bowel disease) or may be involved in tumor initiation or promotion, due to damaging effects on DNA. Nathan (Szatrowski et al. (1991) <i>Canc. Res.</i> 51, 794-798) proposed that the generation of ROI in tumor cells may contribute to the hypermutability seen in tumors, and may therefore contribute to tumor heterogeneity, invasion and metastasis.	
20		
25	In addition to cytotoxic and mutagenic roles, ROI have ideal properties as signal molecules: 1) they are generated in a controlled manner in response to upstream signals; 2) the signal can be terminated by rapid metabolism of O_2^- and H_2O_2 by SOD and catalase/peroxidases; 3) they elicit downstream effects on target molecules, e.g., redox-sensitive regulatory proteins such as NF kappa B and AP-1 (Schreck et al. (1991) <i>EMBO J.</i> 10, 2247-2258; Schmidt et al. (1995) <i>Chemistry & Biology</i> 2, 13-22). Oxidants such as O_2^- and H_2O_2 have a relatively well defined signaling role in bacteria, operating via the SoxI/II regulon to regulate transcription.	
30		
35	ROI appear to have a direct role in regulating cell division, and may function as mitogenic signals in pathological conditions related to growth. These conditions include cancer and cardiovascular disease. O_2^- is generated in endothelial cells	

in response to cytokines, and might play a role in angiogenesis (Matsubara et al. (1986) *J. Immun.* 137, 3295-3298). O_2^- and H_2O_2 are also proposed to function as "life-signals", preventing cells from undergoing apoptosis (Matsubara et al. (1986) *J. Immun.* 137, 3295-3298). As discussed above, many cells respond to growth factors (e.g., platelet derived growth factor (PDGF), epidermal derived growth factor (EGF), angiotensin II, and various cytokines) with both increased production of O_2^- / H_2O_2 and increased proliferation. Inhibition of ROI generation prevents the mitogenic response. Exposure to exogenously generated O_2^- and H_2O_2 results in an increase in cell proliferation. A partial list of responsive cell types is shown below in Table 2 (adapted from Burdon, R. (1995) *Free Radical Biol. Med.* 18, 775-794).

Table 2

	<u>Superoxide</u>	<u>Hydrogen peroxide</u>
	human, hamster fibroblasts	mouse osteoblastic cells
	Balb/3T3 cells	Balb/3T3 cells
20	human histiocytic leukemia	rat, hamster fibroblasts
	mouse epidermal cells	human smooth muscle cells
	rat colonic epithelial cells	rat vascular smooth muscle cells
	rat vascular smooth muscle cells	

While non-transformed cells can respond to growth factors and cytokines with the production of ROI, tumor cells appear to produce ROI in an uncontrolled manner. A series of human tumor cells produced large amounts of hydrogen peroxide compared with non-tumor cells (Szatrowski et al. (1991) *Canc. Res.* 51, 794-798). Ras-transformed NIH 3T3 cells generated elevated amounts of superoxide, and inhibition of superoxide generation by several mechanisms resulted in a reversion to a "normal" growth phenotype.

O_2^- has been implicated in maintenance of the transformed phenotype in cancer cells including melanoma, breast carcinoma, fibrosarcoma, and virally transformed tumor

cells. Decreased levels of the manganese form of SOD (MnSOD) have been measured in cancer cells and *in vitro*-transformed cell lines, predicting increased O_2^- levels (Burdon, R. (1995) *Free Radical Biol. Med.* 18, 775-794). MnSOD is encoded on chromosome 6q25 which is very often lost in melanoma. Overexpression of MnSOD in melanoma and other cancer cells (Church et al. (1993) *Proc. of Natl. Acad. Sci.* 90, 3113-3117; Fernandez-Pol et al. (1982) *Canc. Res.* 42, 609-617; Yan et al. (1996) *Canc. Res.* 56, 2864-2871) resulted in suppression of the transformed phenotype.

ROI are implicated in growth of vascular smooth muscle associated with hypertension, atherosclerosis, and restenosis after angioplasty. O_2^- generation is seen in rabbit aortic adventitia (Pagano et al. (1997) *Proc. Natl. Acad. Sci.* 94, 14483-14488). Vascular endothelial cells release O_2^- in response to cytokines (Matsubara et al. (1986) *J. Immun.* 137, 3295-3298). O_2^- is generated by aortic smooth muscle cells in culture, and increased O_2^- generation is stimulated by angiotensin II which also induces cell hypertrophy. In a rat model system, infusion of angiotensin II leads to hypertension as well as increased O_2^- generation in subsequently isolated aortic tissue (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321.; Yu et al. (1997) *J. Biol. Chem.* 272, 27288-27294). Intravenous infusion of a form of SOD that localizes to the vasculature or an infusion of an O_2^- scavenger prevented angiotensin II induced hypertension and inhibited ROI generation (Fukui et al. (1997) *Circ. Res.* 80, 45-51).

The neutrophil NADPH oxidase, also known as phagocyte respiratory burst oxidase, provides a paradigm for the study of the specialized enzymatic ROI-generating system. This extensively studied enzyme oxidizes NADPH and reduces oxygen to form O_2^- . NADPH oxidase consists of multiple proteins and is regulated by assembly of cytosolic and membrane components. The catalytic moiety consists of flavocytochrome b_{558} , an integral plasma membrane enzyme comprised of two components: gp91phox (gp refers to

glycoprotein; phox is an abbreviation of the words phagocyte and oxidase) and p22phox (p refers to protein). gp91phox contains 1 flavin adenine dinucleotide (FAD) and 2 hemes as well as the NADPH binding site. p22phox has a C-terminal
5 proline-rich sequence which serves as a binding site for cytosolic regulatory proteins. The two cytochrome subunits, gp91phox and p22phox appear to stabilize one another, since the genetic absence of either subunit, as in the inherited disorder chronic granulomatous disease (CGD), results in the absence of
10 the partner subunit (Yu et al. (1997) *J. Biol. Chem.* 272, 27288-27294). Essential cytosolic proteins include p47phox, p67phox and the small GTPase Rac, of which there are two isoforms. p47phox and p67phox both contain SH₃ regions and proline-rich regions which participate in protein interactions
15 governing assembly of the oxidase components during activation. The neutrophil enzyme is regulated in response to bacterial phagocytosis or chemotactic signals by phosphorylation of p47phox, and perhaps other components, as well as by guanine nucleotide exchange to activate the GTP-
20 binding protein Rac.

The origin of ROI in non-phagocytic tissues is unproven, but the occurrence of phagocyte oxidase components has been evaluated in several systems by immunochemical methods, Northern blots and reverse transcriptase-polymerase
25 chain reaction (RT-PCR). The message for p22phox is expressed widely, as is that for Rac1. Several cell types that are capable of O₂⁻ generation have been demonstrated to contain all of the phox components including gp91phox, as summarized below in Table 3. These cell types include endothelial cells,
30 aortic adventitia and lymphocytes.

Table 3

	<u>Tissue</u>	<u>gp91phox</u>	<u>p22phox</u>	<u>p47phox</u>	<u>p67phox</u>
	neutrophil	+ ^{1,2}	+ ^{1,2}	+ ^{1,2}	+ ^{1,2}
	aortic adventitia	+ ¹	+ ¹	+ ¹	+ ¹
5	lymphocytes	+ ²	+ ²	+ ^{1,2}	+ ^{1,2}
	endothelial cells	+ ²	+ ²	+ ^{1,2}	+ ^{1,2}
	glomerular mesangial cells	-	+ ^{1,2}	+ ^{1,2}	+ ^{1,2}
	fibroblasts	-	+ ²	+ ^{1,2}	+ ²
10	aortic sm. muscle	-	+ ^{1,2}	?	?

1= protein expression shown. 2= mRNA expression shown.

15 However, a distinctly different pattern is seen in several other cell types shown in Table 3 including glomerular mesangial cells, rat aortic smooth muscle and fibroblasts. In these cells, expression of gp91phox is absent while p22phox and in some cases cytosolic phox components have been demonstrated to be present. Since gp91phox and p22phox stabilize one another in the neutrophil, there has been much speculation that some molecule, possibly related to gp91phox, accounts for ROI generation in glomerular mesangial cells, rat aortic smooth muscle and fibroblasts (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321). Investigation of fibroblasts from a patient with a genetic absence of gp91phox provides proof that the gp91phox subunit is not involved in ROI generation in these cells (Emmendorffer et al. (1993) *Eur. J. Haematol.* 51, 223-227). Depletion of p22phox from vascular smooth muscle using an antisense approach indicated that this subunit participates in ROI generation in these cells, despite the absence of detectable gp91phox (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321). At this time the molecular candidates possibly related to gp91phox and involved in ROI generation in these cells are unknown.

35 Accordingly, what is needed is the identity of the proteins involved in ROI generation, especially in non-phagocytic tissues and cells. What is also needed are the

5 nucleotide sequences encoding for these proteins, and the
primary sequences of the proteins themselves. Also needed are
vectors designed to include nucleotides encoding for these
proteins. Probes and PCR primers derived from the nucleotide
10 sequence are needed to detect, localize and measure nucleotide
sequences, including mRNA, involved in the synthesis of these
proteins. In addition, what is needed is a means to transfect
cells with these vectors. What is also needed are expression
systems for production of these molecules. Also needed are
15 antibodies directed against these molecules for a variety of uses
including localization, detection, measurement and passive
immunization.

SUMMARY OF THE INVENTION

15 The present invention solves the problems
described above by providing a novel family of nucleotide
sequences and proteins encoded by these nucleotide sequences
termed mox proteins and duox proteins. In particular the
present invention provides compositions comprising the
20 nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID
NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, and
fragments thereof, which encode for the expression of proteins
comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ
ID NO:42, SEQ ID NO:46, SEQ ID NO:48, respectively, and
25 fragments thereof. While not wanting to be bound by the
following statement, it is believed that these proteins are
involved in ROI production. The present invention also
provides vectors containing these nucleotide sequences, cells
transfected with these vectors which produce the proteins
30 comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ
ID NO:42, SEQ ID NO:46, SEQ ID NO:48, and fragments
thereof, and antibodies to these proteins and fragments thereof.
The present invention also provides methods for stimulating
cellular proliferation by administering vectors encoded for
35 production of the proteins comprising SEQ ID NO:2, SEQ ID
NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ
ID NO:48 and fragments thereof. The present invention also

provides methods for stimulating cellular proliferation by administering the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The nucleotides and antibodies of the present invention are useful for the detection, localization and measurement of the nucleic acids encoding for the production of the proteins of the present invention, and also for the detection, localization and measurement of the proteins of the present invention. These nucleotides and antibodies can be combined with other reagents in kits for the purposes of detection, localization and measurement.

Most particularly, the present invention involves a method for regulation of cell division or cell proliferation by modifying the activity or expression of the proteins described as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof. These proteins, in their naturally occurring or expressed forms, are expected to be useful in drug development, for example for screening of chemical and drug libraries by observing inhibition of the activity of these enzymes. Such chemicals and drugs would likely be useful as treatments for cancer, prostatic hypertrophy, benign prostatic hypertrophy, hypertension, atherosclerosis and many other disorders involving abnormal cell growth or proliferation as described below. The entire expressed protein may be useful in these assays. Portions of the molecule which may be targets for inhibition or modification include but are not limited to the binding site for pyridine nucleotides (NADPH or NADH), the flavoprotein domain (approximately the C-terminal 265 amino acids), and/or the binding or catalytic site for flavin adenine dinucleotide (FAD).

The method of the present invention may be used for the development of drugs or other therapies for the treatment of conditions associated with abnormal growth including, but not limited to the following: cancer, psoriasis, prostatic hypertrophy, benign prostatic hypertrophy, cardiovascular disease, proliferation of vessels, including but

not limited to blood vessels and lymphatic vessels, arteriovenous malformation, vascular problems associated with the eye, atherosclerosis, hypertension, and restenosis following angioplasty. The enzymes of the present invention are excellent
5 targets for the development of drugs and other agents which may modulate the activity of these enzymes. It is to be understood that modulation of activity may result in enhanced, diminished or absence of enzymatic activity. Modulation of the activity of these enzymes may be useful in treatment of
10 conditions associated with abnormal growth.

Drugs which affect the activity of the enzymes represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, may also be combined with other therapeutics in the
15 treatment of specific conditions. For example, these drugs may be combined with angiogenesis inhibitors in the treatment of cancer, with antihypertensives for the treatment of hypertension, and with cholesterol lowering drugs for the treatment of atherosclerosis.

20 Accordingly, an object of the present invention is to provide nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production.

Another object of the present invention is to
25 provide vectors containing these nucleotide sequences, or fragments thereof.

Yet another object of the present invention is to provide cells transfected with these vectors.

Still another object of the present invention is to
30 administer cells transfected with these vectors to animals and humans.

Another object of the present invention is to provide proteins, or fragments thereof, that are involved in ROI production.

35 Still another object of the present invention is to provide antibodies, including monoclonal and polyclonal

antibodies, or fragments thereof, raised against proteins, or fragments thereof, that are involved in ROI production.

5 Another object of the present invention is to administer genes containing nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans and also to cells obtained from animals and humans.

10 Another object of the present invention is to administer antisense complimentary sequences of genes containing nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans and also to cells obtained from animals and humans.

15 Yet another object of the present invention is to provide a method for stimulating or inhibiting cellular proliferation by administering vectors containing nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans. It is also an object of the present invention to provide a method for stimulating or inhibiting cellular proliferation by administering vectors containing antisense complimentary sequences of nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production, to animals and humans. These methods of stimulating cellular proliferation are useful for a variety of purposes, including but not limited to, developing animal models of tumor formation, stimulating cellular proliferation of blood marrow cells following chemotherapy or radiation, or in cases of anemia.

30 Still another object of the present invention is to provide antibodies useful in immunotherapy against cancers expressing the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof.

35

Yet another object of the present invention is to provide nucleotide probes useful for the detection, localization and measurement of nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production.

Another object of the present invention is to provide antibodies useful for the detection, localization and measurement of nucleotide sequences, or fragments thereof, encoding for the production of proteins, or fragments thereof, that are involved in ROI production.

Another object of the present invention is to provide kits useful for detection of nucleic acids including the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins, or fragments thereof, that are involved in ROI production.

Yet another object of the present invention is to provide kits useful for detection and measurement of nucleic acids including the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins, or fragments thereof, that are involved in ROI production.

Still another object of the present invention is to provide kits useful for the localization of nucleic acids including the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins, or fragments thereof that are involved in ROI production.

Another object of the present invention is to provide kits useful for detection of proteins, including the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, that are involved in ROI production.

Yet another object of the present invention is to provide kits useful for detection and measurement of proteins,

including the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, that are involved in ROI production.

5 Still another object of the present invention is to provide kits useful for localization of proteins, including the proteins represented in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, that are involved in ROI production.

10 Yet another object of the present invention is to provides kits useful for the detection, measurement or localization of nucleic acids, or fragments thereof, encoding for proteins, or fragments thereof, that are involved in ROI production, for use in diagnosis and prognosis of abnormal
15 cellular proliferation related to ROI production.

Another object of the present invention is to provides kits useful for the detection, measurement or localization of proteins, or fragments thereof, that are involved in ROI production, for use in diagnosis and prognosis of
20 abnormal cellular proliferation related to ROI production.

These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiments and the appended drawings.

25

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1(a-d). Comparison of amino acid sequences of the human mox1 protein (labeled mox1.human, SEQ ID NO:2), rat mox1 protein (labeled mox1.rat, SEQ ID NO:21),
30 human mox2 protein (labeled mox2.human., SEQ ID NO:4) of the present invention to human (gp 91phox/human.pep, SEQ ID NO:12) bovine (gp 91 phox/bovine.pep, SEQ ID NO:37), and murine (gp 91 phox/mouse.pep, SEQ ID NO:38) proteins. Also included are related plant enzyme proteins cytb
35 558.arabidopsis.pep (SEQ ID NO:39) and cytb558.rice.pep,

(SEQ ID NO:40). Enclosed in boxes are similar amino acid residues.

5 Fig. 2. Sequence similarities among proteins related to gp91phox including human mox1 (SEQ ID NO:2), human mox2 (SEQ ID NO:4), and rat mox1 (SEQ ID NO:21). The dendrogram indicates the degree of similarity among this family of proteins, and also includes the related plant enzymes.

10 Fig. 3. Cell free assay for mox-1 activity. Superoxide generation was measured using the chemiluminescent reaction between lucigenin and superoxide in cell lysates from vector control NEF2 and mox1 transfected NIH3T3 cells.

15 Fig. 4. Superoxide generation by human mox1. Reduction of NBT in mox1 transfected and control fibroblasts was measured in the absence (filled bars) or presence (open bars) or superoxide dismutase.

20 Fig. 5. Aconitase (filled bars), lactate dehydrogenase (narrow hatching) and fumarase (broad hatching) were determined in lysates of cells transfected with vector alone (NEF2) or with mox1 (YA26, YA28 and YA212).

DETAILED DESCRIPTION OF THE INVENTION

25 The present invention solves the problems described above by providing a novel family of nucleotide sequences and proteins, encoded by these nucleotide sequences, termed mox proteins and duox proteins. The term "mox" refers to "mitogenic oxidase" while the term "duox" refers to "dual oxidase". In particular, the present invention provides novel compositions comprising the nucleotide sequences SEQ ID
30 NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, and fragments thereof, which encode, respectively, for the expression of proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof.

35 Both the mox and duox proteins described herein have homology to the gp91phox protein involved in ROI

generation, however, the mox and duox proteins comprise a novel and distinct family of proteins. The mox proteins included in the present invention have a molecular weight of approximately 65 kDa as determined by reducing gel electrophoresis and are capable of inducing ROI generation in cells. As described in more detail below, the mox proteins of the present invention also function in the regulation of cell growth, and are therefore implicated in diseases involving abnormal cell growth such as cancer. The present invention describes mox proteins found in human and rat, however, it is likely that the mox family of genes/proteins is widely distributed among multicellular organisms.

The duox proteins described herein are larger than the mox proteins and have three distinct regions: the amino terminal region having homology to peroxidase proteins, the internal region having homology to calmodulin (CAM) proteins and the carboxy-terminal region having homology to mox proteins. Human duox1 is shown in SEQ ID NO:46 and a portion of human duox2 is shown in SEQ ID NO:48. Nucleotides encoding duox1 and duox2 proteins are also shown in SEQ ID NO: 45 and SEQ ID NO:47, respectively. In addition to the human duox proteins, comparison of the sequence of human duox1 and human duox2 with genomic databases using BLAST searching resulted in the identification of two homologs of duox in *C. elegans* (Ce-duox1 and Ce-duox2). *Drosophila* also appears to have at least one duox homolog. Thus, the duox family of genes/proteins is widely distributed among multicellular organisms.

Although not wanting to be bound by the following statement, it is believed that duox1 and duox2 have dual enzymatic functions, catalyzing both the generation of superoxide and peroxidative type reactions. The latter class of reactions utilize hydrogen peroxide as a substrate (and in some cases have been proposed to utilize superoxide as a substrate). Since hydrogen peroxide is generated spontaneously from the dismutation of superoxide, it is believed that the NAD(P)H

oxidase domain generates the superoxide and/or hydrogen peroxide which can then be used as a substrate for the peroxidase domain. In support of this hypothesis, a model for the duox1 protein in *C. elegans* has been developed that has an
5 extracellular N-terminal peroxidase domain, a transmembrane region and a NADPH binding site located on the cytosolic face of the plasma membrane. By analogy with the neutrophil NADPH-oxidase which generates extracellular superoxide, human duox1 is predicted to generate superoxide and its
10 byproduct hydrogen peroxide extracellularly where it can be utilized by the peroxidase domain.

While the ROI generated by duox1 and duox2 may function as does mox1 in regulation of cell growth, the presence of the peroxidase domain is likely to confer additional
15 biological functions. Depending upon the co-substrate, peroxidases can participate in a variety of reactions including halogenation such as the generation of hypochlorous acid (HOCl) by myeloperoxidase and the iodination of tyrosine to form thyroxine by thyroid peroxidase. Peroxidases have also
20 been documented to participate in the metabolism of polyunsaturated fatty acids, and in the chemical modification of tyrosine in collagen (by sea urchin ovoperoxidase). Although not wanting to be bound by this statement, it is believed that the predicted transmembrane nature of duox1 facilitates its function
25 in the formation or modification of extracellular matrix or basement membrane. Since the extracellular matrix plays an important role in tumor cell growth, invasion and metastasis, it is believed that the duox type enzymes play a pathogenic role in such conditions.

30 In addition to the nucleotide sequences described above, the present invention also provides vectors containing these nucleotide sequences and fragments thereof, cells transfected with these vectors which produce the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ
35 ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof, and antibodies to these proteins and fragments thereof.

5 The present invention also provides methods for stimulating cellular proliferation by administering vectors, or cells containing vectors, encoded for production of the proteins comprising SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The nucleotides and antibodies of the present invention are useful for the detection, localization and measurement of the nucleic acids encoding for the production of the proteins of the present invention, and also for the detection, localization and measurement of the proteins of the present invention. These nucleotides and antibodies can be combined with other reagents in kits for the purposes of detection, localization and measurement. These kits are useful for diagnosis and prognosis of conditions involving cellular proliferation associated with production of reactive oxygen intermediates.

10 The present invention solves the problems described above by providing a composition comprising the nucleotide sequence SEQ ID NO:1 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:3 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:22 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:41 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:45 and fragments thereof. The present invention also provides a composition comprising the nucleotide sequence SEQ ID NO:47 and fragments thereof.

25 The present invention provides a composition comprising the protein SEQ ID NO:2 encoded by the nucleotide sequence SEQ ID NO:1. The present invention provides a composition comprising the protein SEQ ID NO:4 encoded by the nucleotide sequence SEQ ID NO:3. The present invention provides a composition comprising the protein SEQ ID NO:21 encoded by the nucleotide sequence SEQ ID NO:22. The present invention provides a composition comprising the protein

5 SEQ ID NO:42 encoded by the nucleotide sequence SEQ ID NO:41. The present invention provides a composition comprising the protein SEQ ID NO:46 encoded by the nucleotide sequence SEQ ID NO:45. The present invention provides a composition comprising the protein SEQ ID NO:48 encoded by the nucleotide sequence SEQ ID NO:47.

10 The present invention provides a composition comprising the protein SEQ ID NO:2 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:1 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:4 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:3 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:21 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:22 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:42 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:41 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:46 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:45 or fragments thereof. The present invention also provides a composition comprising the protein SEQ ID NO:48 or fragments thereof, encoded by the nucleotide sequence SEQ ID NO:47 or fragments thereof.

20 The present invention also provides vectors containing the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. The present invention also provides cells transfected with these vectors. In addition, the present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:1 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof. The

present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:47 or fragments thereof.

The present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:1 or fragments thereof, which produce the protein SEQ ID NO:2 or fragments thereof. In addition, the present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof which produce the protein SEQ ID NO:4 or fragments thereof. In addition, the present invention provides cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof which produce the protein SEQ ID NO:21 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof which produce the protein SEQ ID NO:42 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof which produce the protein SEQ ID NO:46 or fragments thereof. The present invention also provides cells stably transfected with the nucleotide sequence SEQ ID NO:47 or fragments thereof which produce the protein SEQ ID NO:48 or fragments thereof.

The present invention provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:1 which produce the protein SEQ ID NO:2 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof, which produce the protein SEQ ID NO:4 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof, which produce

the protein SEQ ID NO:21 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof, which produce
5 the protein SEQ ID NO:42 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof, which produce
10 the protein SEQ ID NO:46 or fragments thereof. The present invention also provides a method for stimulating growth by administering cells stably transfected with the nucleotide sequence SEQ ID NO:47 or fragments thereof, which produce
the protein SEQ ID NO:48 or fragments thereof.

Specifically, the present invention provides a
15 method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:1 or fragments thereof, which produce the protein SEQ ID NO:2 or fragments thereof. The present invention also provides a
20 method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:3 or fragments thereof, which produce the protein SEQ ID NO:4 or fragments thereof. The present invention also provides a
25 method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:22 or fragments thereof, which produce the protein SEQ ID NO:21 or fragments thereof. The present invention also provides a
30 method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:41 or fragments thereof, which produce the protein SEQ ID NO:42 or fragments thereof. The present invention also provides a
35 method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:45 or fragments thereof, which produce the protein SEQ ID NO:46 or fragments thereof. The present invention also provides a
method for stimulating tumor formation by administering cells stably transfected with the nucleotide sequence SEQ ID NO:47

or fragments thereof, which produce the protein SEQ ID NO:48 or fragments thereof.

5 The present invention may also be used to develop anti-sense nucleotide sequences to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. These anti-sense molecules may be used to interfere with translation of nucleotide sequences, such as SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, that encode for proteins such as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof. Administration of these anti-sense molecules, or vectors encoding for anti sense molecules, to humans and animals, would interfere with production of proteins such as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, thereby decreasing production of ROIs and inhibiting cellular proliferation. These methods are useful in producing animal models for use in study of tumor development and vascular growth, and for study of the efficacy of treatments for affecting tumor and vascular growth *in vivo*.

15 The present invention also provides a method for high throughput screening of drugs and chemicals which modulate the proliferative activity of the enzymes of the present invention, thereby affecting cell division. Combinatorial chemical libraries may be screened for chemicals which modulate the proliferative activity of these enzymes. Drugs and chemicals may be evaluated based on their ability to modulate the enzymatic activity of the expressed or endogenous proteins, including those represented by SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof. Endogenous proteins may be obtained from many different tissues or cells, such as colon cells. Drugs may also be evaluated based on their ability to bind to the expressed or endogenous proteins represented by SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID

NO:46, SEQ ID NO:48 or fragments thereof. Enzymatic activity may be NADPH- or NADH-dependent superoxide generation catalyzed by the holoprotein. Enzymatic activity may also be NADPH- or NADH-dependent diaphorase activity catalyzed by either the holoprotein or the flavoprotein domain.

5

10

15

20

25

30

35

By flavoprotein domain, is meant approximately the C-terminal half of the enzymes shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, or fragments thereof, and the C-terminal end of the enzymes shown in SEQ ID NO:46 and SEQ ID NO:48 (approximately the C-terminal 265 amino acids). This fragment of gp91phox has NADPH-dependent reductase activity towards cytochrome c, nitrobluetetrazolium and other dyes. Expressed proteins or fragments thereof can be used for robotic screens of existing combinatorial chemical libraries. While not wanting to be bound by the following statement, it is believed that the NADPH or NADH binding site and the FAD binding site are useful for evaluating the ability of drugs and other compositions to bind to the mox and duox enzymes or to modulate their enzymatic activity. The use of the holoprotein or the C-terminal half or end regions are preferred for developing a high throughput drug screen. Additionally, the N-terminal one-third of the duox domain (the peroxidase domain) may also be used to evaluate the ability of drugs and other compositions to inhibit the peroxidase activity, and for further development of a high throughput drug screen.

The present invention also provides antibodies directed to the proteins SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The antibodies of the present invention are useful for a variety of purposes including localization, detection and measurement of the proteins SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. The antibodies may be employed in kits to accomplish these purposes. These antibodies may also be linked to cytotoxic agents for selected killing of cells. The

term antibody is meant to include any class of antibody such as IgG, IgM and other classes. The term antibody also includes a completely intact antibody and also fragments thereof, including but not limited to Fab fragments and Fab + Fc fragments.

5 The present invention also provides the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 and fragments thereof. These nucleotides are useful for a variety of purposes including localization, detection, and measurement of messenger
10 RNA involved in synthesis of the proteins represented as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. These nucleotides may also be used in the construction of labeled probes for the localization, detection, and measurement of
15 nucleic acids such as messenger RNA or alternatively for the isolation of larger nucleotide sequences containing the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. These nucleotide sequences may
20 be used to isolate homologous strands from other species using techniques known to one of ordinary skill in the art. These nucleotide sequences may also be used to make probes and complementary strands. In particular, the nucleotide sequence shown in SEQ ID NO:47 may be used to isolate the complete
25 coding sequence for duox2. The nucleotides may be employed in kits to accomplish these purposes.

 Most particularly, the present invention involves a method for modulation of growth by modifying the proteins represented as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof.
30

 The term "mitogenic regulators" is used herein to mean any molecule that acts to affect cell division.

 The term "animal" is used herein to mean humans and non-human animals of both sexes.
35

The terms "a", "an" and "the" as used herein are defined to mean one or more and include the plural unless the context is inappropriate.

5 "Proteins", "peptides," "polypeptides" and "oligopeptides" are chains of amino acids (typically L-amino acids) whose alpha carbons are linked through peptide bonds formed by a condensation reaction between the carboxyl group of the alpha carbon of one amino acid and the amino group of the alpha carbon of another amino acid. The terminal amino acid at one end of the chain (*i.e.*, the amino terminal) has a free amino group, while the terminal amino acid at the other end of the chain (*i.e.*, the carboxy terminal) has a free carboxyl group. As such, the term "amino terminus" (abbreviated N-terminus) refers to the free alpha-amino group on the amino acid at the amino terminal of the protein, or to the alpha-amino group (imino group when participating in a peptide bond) of an amino acid at any other location within the protein. Similarly, the term "carboxy terminus" (abbreviated C-terminus) refers to the free carboxyl group on the amino acid at the carboxy terminus of a protein, or to the carboxyl group of an amino acid at any other location within the protein.

Typically, the amino acids making up a protein are numbered in order, starting at the amino terminal and increasing in the direction toward the carboxy terminal of the protein. Thus, when one amino acid is said to "follow" another, that amino acid is positioned closer to the carboxy terminal of the protein than the preceding amino acid.

25 The term "residue" is used herein to refer to an amino acid (D or L) or an amino acid mimetic that is incorporated into a protein by an amide bond. As such, the amino acid may be a naturally occurring amino acid or, unless otherwise limited, may encompass known analogs of natural amino acids that function in a manner similar to the naturally occurring amino acids (*i.e.*, amino acid mimetics). Moreover, an amide bond mimetic includes peptide backbone modifications well known to those skilled in the art.

Furthermore, one of skill will recognize that, as mentioned above, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 1%) in an encoded sequence are conservatively modified variations where the alterations result in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

When the peptides are relatively short in length (*i.e.*, less than about 50 amino acids), they are often synthesized using standard chemical peptide synthesis techniques. Solid phase synthesis in which the C-terminal amino acid of the sequence is attached to an insoluble support followed by sequential addition of the remaining amino acids in the sequence is a preferred method for the chemical synthesis of the antigenic epitopes described herein. Techniques for solid phase synthesis are known to those skilled in the art.

Alternatively, the antigenic epitopes described herein are synthesized using recombinant nucleic acid methodology. Generally, this involves creating a nucleic acid sequence that encodes the peptide or protein, placing the nucleic acid in an expression cassette under the control of a particular promoter, expressing the peptide or protein in a host, isolating the expressed peptide or protein and, if required, renaturing the

peptide or protein. Techniques sufficient to guide one of skill through such procedures are found in the literature.

5 When several desired protein fragments or peptides are encoded in the nucleotide sequence incorporated into a vector, one of skill in the art will appreciate that the protein fragments or peptides may be separated by a spacer molecule such as, for example, a peptide, consisting of one or more amino acids. Generally, the spacer will have no specific biological activity other than to join the desired protein fragments or peptides together, or to preserve some minimum distance or other spatial relationship between them. However, the constituent amino acids of the spacer may be selected to influence some property of the molecule such as the folding, net charge, or hydrophobicity. Nucleotide sequences encoding for the production of residues which may be useful in purification of the expressed recombinant protein may be built into the vector. Such sequences are known in the art. For example, a nucleotide sequence encoding for a poly histidine sequence may be added to a vector to facilitate purification of the expressed recombinant protein on a nickel column.

20 Once expressed, recombinant peptides, polypeptides and proteins can be purified according to standard procedures known to one of ordinary skill in the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like. Substantially pure compositions of about 50 to 99% homogeneity are preferred, and 80 to 95% or greater homogeneity are most preferred for use as therapeutic agents.

25 One of skill in the art will recognize that after chemical synthesis, biological expression or purification, the desired proteins, fragments thereof and peptides may possess a conformation substantially different than the native conformations of the proteins, fragments thereof and peptides. In this case, it is often necessary to denature and reduce protein and then to cause the protein to re-fold into the preferred conformation. Methods of reducing and denaturing proteins

and inducing re-folding are well known to those of skill in the art.

5 The genetic constructs of the present invention include coding sequences for different proteins, fragments thereof, and peptides. The genetic constructs also include epitopes or domains chosen to permit purification or detection of the expressed protein. Such epitopes or domains include DNA sequences encoding the glutathione binding domain from glutathione S-transferase, hexa-histidine, thioredoxin, hemagglutinin antigen, maltose binding protein, and others commonly known to one of skill in the art. The preferred genetic construct includes the nucleotide sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, SEQ ID NO:47 or fragments thereof. It is to be understood that additional or alternative nucleotide sequences may be included in the genetic constructs in order to encode for the following: a) multiple copies of the desired proteins, fragments thereof, or peptides; b) various combinations of the desired proteins, fragments thereof, or peptides; and c) conservative modifications of the desired proteins, fragments thereof, or peptides, and combinations thereof. Preferred proteins include the human mox1 protein and human mox2 protein shown as SEQ ID NO:2 and SEQ ID NO:4, respectively, and fragments thereof. Some preferred fragments of the human mox1 protein (SEQ ID NO:2) include but are not limited to the proteins shown as SEQ ID NO:23, SEQ ID NO:24, and SEQ ID NO:25. The protein mox1 is also called p65mox in this application. Another preferred protein of the present invention is rat mox1 protein shown as SEQ ID NO:21 and fragments thereof. Another preferred protein of the present invention is rat mox1B protein shown as SEQ ID NO:42 and fragments thereof. Yet another preferred protein of the present invention is duox1 protein shown as SEQ ID NO:46 and fragments thereof. Still another preferred protein of the present invention is duox2 protein. A partial amino acid sequence of the duox2 protein is shown as SEQ ID NO:48.

5 The nucleotide sequences of the present invention may also be employed to hybridize to nucleic acids such as DNA or RNA nucleotide sequences under high stringency conditions which permit detection, for example, of alternately spliced messages.

10 The genetic construct is expressed in an expression system such as in NIH 3T3 cells using recombinant sequences in a pcDNA-3 vector (Invitrogen, Carlsbad, CA) to produce a recombinant protein. Preferred expression systems include but are not limited to Cos-7 cells, insect cells using recombinant baculovirus, and yeast. It is to be understood that other expression systems known to one of skill in the art may be used for expression of the genetic constructs of the present invention. The preferred proteins of the present invention are the proteins referred to herein as human mox1 and human mox2 or fragments thereof which have the amino acid sequences set forth in SEQ ID NO:3 and SEQ ID NO:4, respectively, or an amino acid sequence having amino acid substitutions as defined in the definitions that do not significantly alter the function of the recombinant protein in an adverse manner. Another preferred protein of the present invention is referred to herein as rat mox1 and has the amino acid sequence set forth in SEQ ID NO:21. Yet another preferred protein of the present invention is referred to herein as rat mox1B and has the amino acid sequence set forth in SEQ ID NO:42. Two other preferred proteins of the present invention are referred to herein as human duox1 and human duox2, or fragments thereof, which have the amino acid sequences set forth in SEQ ID NO:46 and SEQ ID NO:48, respectively, or an amino acid sequence having amino acid substitutions as defined in the definitions that do not significantly alter the function of the recombinant protein in an adverse manner.

Terminology

35 It should be understood that some of the terminology used to describe the novel mox and duox proteins

5 contained herein is different from the terminology in U.S. Provisional Application Serial No. 60/107,911 and U.S. Provisional Application Serial No. 60/149,332 upon which this application claims priority in part. As described herein, the term "human mox1" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:2, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:1, or a fragment thereof. As described herein, the term "human mox2" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:4, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:3, or a fragment thereof. As described herein, the term "human duox1" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:46, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:45, or a fragment thereof. As described herein, the term "human duox2" refers to a protein comprising an amino acid sequence as set forth in SEQ ID NO:48, or a fragment thereof, and encoded by the nucleotide sequence as set forth in SEQ ID NO:47, or a fragment thereof.

Construction of the Recombinant Gene

25 The desired gene is ligated into a transfer vector, such as pcDNA3, and the recombinants are used to transform host cells such as Cos-7 cells. It is to be understood that different transfer vectors, host cells, and transfection methods may be employed as commonly known to one of ordinary skill in the art. Six desired genes for use in transfection are shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45 and SEQ ID NO:47. For example, lipofectamine-mediated transfection and *in vivo* homologous recombination was used to introduce the mox1 gene into NIH 3T3 cells.

35 The synthetic gene is cloned and the recombinant construct containing mox or duox gene is produced and grown in confluent monolayer cultures of a Cos-7 cell line. The

expressed recombinant protein is then purified, preferably using affinity chromatography techniques, and its purity and specificity determined by known methods.

5 A variety of expression systems may be employed for expression of the recombinant protein. Such expression methods include, but are not limited to the following: bacterial expression systems, including those utilizing *E. coli* and *Bacillus subtilis*; virus systems; yeast expression systems; cultured insect and mammalian cells; and other expression systems known to
10 one of ordinary skill in the art.

Transfection of Cells

It is to be understood that the vectors of the present invention may be transfected into any desired cell or cell line.
15 Both *in vivo* and *in vitro* transfection of cells are contemplated as part of the present invention. Preferred cells for transfection include but are not limited to the following: fibroblasts (possibly to enhance wound healing and skin formation), granulocytes (possible benefit to increase function in a
20 compromised immune system as seen in AIDS, and aplastic anemia), muscle cells, neuroblasts, stem cells, bone marrow cells, osteoblasts, B lymphocytes, and T lymphocytes.

Cells may be transfected with a variety of methods known to one of ordinary skill in the art and include but are not
25 limited to the following: electroporation, gene gun, calcium phosphate, lipofectamine, and fugene, as well as adenoviral transfection systems.

Host cells transfected with the nucleic acids represented in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22,
30 SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, or fragments thereof, are used to express the proteins SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 and SEQ ID NO:48, respectively, or fragments thereof.

These expressed proteins are used to raise
35 antibodies. These antibodies may be used for a variety of applications including but not limited to immunotherapy against

cancers expressing one of the mox or duox proteins, and for detection, localization and measurement of the proteins shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48 or fragments thereof.

5

Purification and Characterization of the Expressed Protein

The proteins of the present invention can be expressed as a fusion protein with a poly histidine component, such as a hexa histidine, and purified by binding to a metal affinity column using nickel or cobalt affinity matrices. The protein can also be expressed as a fusion protein with glutathione S-transferase and purified by affinity chromatography using a glutathione agarose matrix. The protein can also be purified by immunoaffinity chromatography by expressing it as a fusion protein, for example with hemagglutinin antigen. The expressed or naturally occurring protein can also be purified by conventional chromatographic and purification methods which include anion and cation exchange chromatography, gel exclusion chromatography, hydroxylapatite chromatography, dye binding chromatography, ammonium sulfate precipitation, precipitation in organic solvents or other techniques commonly known to one of skill in the art.

10

15

20

25

Methods of Assessing Activity of Expressed Proteins

Different methods are available for assessing the activity of the expressed proteins of the present invention, including but not limited to the proteins represented as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48 substituted analogs thereof, and fragments thereof.

30

1. Assays of the holoprotein and fragments thereof for superoxide generation:

35

A. General considerations.

These assays are useful in assessing efficacy of drugs designed to modulate the activity of the enzymes of the present invention. The holoprotein may be expressed in COS-7 cells, NIH 3T3 cells, insect cells (using baculoviral technology) or other cells using methods known to one of skill in the art. Membrane fractions or purified protein are used for the assay. The assay may require or be augmented by other cellular proteins such as p47phox, p67phox, and Rac1, as well as potentially other unidentified factors (e.g., kinases or other regulatory proteins).

B. Cytochrome c reduction.

NADPH or NADH is used as the reducing substrate, in a concentration of about 100 μ M. Reduction of cytochrome c is monitored spectrophotometrically by the increase in absorbance at 550 nm, assuming an extinction coefficient of 21 $\text{mM}^{-1}\text{cm}^{-1}$. The assay is performed in the absence and presence of about 10 μ g superoxide dismutase. The superoxide-dependent reduction is defined as cytochrome c reduction in the absence of superoxide dismutase minus that in the presence of superoxide dismutase (Uhlinger et al. (1991) *J. Biol. Chem.* 266, 20990-20997). Acetylated cytochrome c may also be used, since the reduction of acetylated cytochrome c is thought to be exclusively via superoxide.

C. Nitroblue tetrazolium reduction.

For nitroblue tetrazolium (NBT) reduction, the same general protocol is used, except that NBT is used in place of cytochrome c. In general, about 1 mL of filtered 0.25 % nitrotetrazolium blue (Sigma, St. Louis, MO) is added in Hanks buffer without or with about 600 Units of superoxide dismutase (Sigma) and samples are incubated at approximately 37°C. The oxidized NBT is clear, while the reduced NBT is blue and insoluble. The insoluble product is collected by centrifugation, and the pellet is re-suspended in about 1 mL of pyridine (Sigma) and heated for about 10 minutes at 100°C to solubilize

the reduced NBT. The concentration of reduced NBT is determined by measuring the absorbance at 510 nm, using an extinction coefficient of $11,000 \text{ M}^{-1}\text{cm}^{-1}$. Untreated wells are used to determine cell number.

5

D. Luminescence.

Superoxide generation may also be monitored with a chemiluminescence detection system utilizing lucigenin (bis-N-methylacridinium nitrate, Sigma, St. Louis, MO). The sample is mixed with about $100 \mu\text{M}$ NADPH (Sigma, St. Louis, MO) and $10 \mu\text{M}$ lucigenin (Sigma, St. Louis, MO) in a volume of about $150 \mu\text{L}$ Hanks solution. Luminescence is monitored in a 96-well plate using a LumiCounter (Packard, Downers Grove, IL) for 0.5 second per reading at approximately 1 minute intervals for a total of about 5 minutes; the highest stable value in each data set is used for comparisons. As above, superoxide dismutase is added to some samples to prove that the luminescence arises from superoxide. A buffer blank is subtracted from each reading (Ushio-Fukai et al. (1996) *J. Biol. Chem.* 271, 23317-23321).

10

15

20

E. Assays in intact cells.

Assays for superoxide generation may be performed using intact cells, for example, the mox-transfected NIH 3T3 cells. In principle, any of the above assays can be used to evaluate superoxide generation using intact cells, for example, the mox-transfected NIH 3T3 cells. NBT reduction is a preferred assay method.

25

30

2. Assays of truncated proteins comprised of approximately the C-terminal 265 amino acid residues

While not wanting to be bound by the following statement, the truncated protein comprised of approximately the C-terminal 265 amino acid residues is not expected to generate superoxide, and therefore, superoxide dismutase is not added in assays of the truncated protein. Basically, a similar assay is

35

established and the superoxide-independent reduction of NBT, cytochrome c, dichlorophenolindophenol, ferricyanide, or another redox-active dye is examined.

5 *Nucleotides and Nucleic Acid Probes*

10 The nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof and PCR primers therefor, may be used, respectively, for localization, detection and measurement of nucleic acids related to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof. The nucleotide sequences SEQ ID NO:1 and SEQ ID NO:3 are also called the human mox1 gene and the human mox2 gene in this application. 15 SEQ ID NO:22 is also known as the rat mox1 gene in this application. SEQ ID NO:41 is also known as the rat mox1B gene in this application. SEQ ID NO:45 is also known as the human duox1 gene in this application. SEQ ID NO:47 is also known as the human duox2 gene in this application.

20 The nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof, may be used to create probes to isolate larger nucleotide sequences containing the nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, respectively. The nucleotide sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47, as well as fragments thereof, may also be used to create probes to identify and isolate mox and duox proteins in 25 other species. 30

35 The nucleic acids described herein include messenger RNA coding for production of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. Such nucleic acids include but are not limited to cDNA probes. These probes may be labeled in a variety of ways known to one of ordinary skill in

5 the art. Such methods include but are not limited to isotopic and non-isotopic labeling. These probes may be used for *in situ* hybridization for localization of nucleic acids such as mRNA encoding for SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. Localization may be performed using *in situ* hybridization at both ultrastructural and light microscopic levels of resolution using techniques known to one of ordinary skill in the art.

10 These probes may also be employed to detect and quantitate nucleic acids and mRNA levels using techniques known to one of ordinary skill in the art including but not limited to solution hybridization.

15 *Antibody Production*

20 The proteins shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48, or fragments thereof, are combined with a pharmaceutically acceptable carrier or vehicle to produce a pharmaceutical composition and administered to animals for the production of polyclonal antibodies using methods known to one of ordinary skill in the art. The preferred animals for antibody production are rabbits and mice. Other animals may be employed for immunization with these proteins or fragments thereof. Such animals include, but are not limited to the following; sheep, horses, pigs, donkeys, cows, monkeys and rodents such as guinea pigs and rats.

25 The terms "pharmaceutically acceptable carrier or pharmaceutically acceptable vehicle" are used herein to mean
30 any liquid including but not limited to water or saline, oil, gel, salve, solvent, diluent, fluid ointment base, liposome, micelle, giant micelle, and the like, which is suitable for use in contact with living animal or human tissue without causing adverse physiological responses, and which does not interact with the
35 other components of the composition in a deleterious manner.

5 The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers.

10 Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. 15 20 Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets commonly used by one of ordinary skill in the art.

25 Preferred unit dosage formulations are those containing a dose or unit, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients, particularly mentioned above, the formulations of the present invention may include other agents commonly used by one of ordinary skill in the art.

30 The pharmaceutical composition may be administered through different routes, such as oral, including buccal and sublingual, rectal, parenteral, aerosol, nasal, intramuscular, subcutaneous, intradermal, and topical. The pharmaceutical composition of the present invention may be administered in different forms, including but not limited to 35 solutions, emulsions and suspensions, microspheres, particles, microparticles, nanoparticles, and liposomes. It is expected that

from about 1 to 7 dosages may be required per immunization regimen. Initial injections may range from about 0.1 μg to 1 mg, with a preferred range of about 1 μg to 800 μg , and a more preferred range of from approximately 25 μg to 500 μg .
5 Booster injections may range from 0.1 μg to 1 mg, with a preferred range of approximately 1 μg to 800 μg , and a more preferred range of about 10 μg to 500 μg .

The volume of administration will vary depending on the route of administration and the size of the recipient. For
10 example, intramuscular injections may range from about 0.1 ml to 1.0 ml.

The pharmaceutical composition may be stored at temperatures of from about 4°C to -100°C. The pharmaceutical composition may also be stored in a lyophilized state at different
15 temperatures including room temperature. The pharmaceutical composition may be sterilized through conventional means known to one of ordinary skill in the art. Such means include, but are not limited to filtration, radiation and heat. The pharmaceutical composition of the present invention may also
20 be combined with bacteriostatic agents, such as thimerosal, to inhibit bacterial growth.

Adjuvants

A variety of adjuvants known to one of ordinary
25 skill in the art may be administered in conjunction with the protein in the pharmaceutical composition. Such adjuvants include, but are not limited to the following: polymers, co-polymers such as polyoxyethylene-polyoxypropylene copolymers, including block co-polymers; polymer P1005;
30 Freund's complete adjuvant (for animals); Freund's incomplete adjuvant; sorbitan monooleate; squalene; CRL-8300 adjuvant; alum; QS 21, muramyl dipeptide; trehalose; bacterial extracts, including mycobacterial extracts; detoxified endotoxins; membrane lipids; or combinations thereof.

35 Monoclonal antibodies can be produced using hybridoma technology in accordance with methods well known

to those skilled in the art. The antibodies are useful as research or diagnostic reagents or can be used for passive immunization. The composition may optionally contain an adjuvant.

5 The polyclonal and monoclonal antibodies useful as research or diagnostic reagents may be employed for detection and measurement of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 and fragments thereof. Such antibodies may be used to detect these proteins in a biological sample, including but not limited to
10 samples such as cells, cellular extracts, tissues, tissue extracts, biopsies, tumors, and biological fluids. Such detection capability is useful for detection of disease related to these proteins to facilitate diagnosis and prognosis and to suggest possible treatment alternatives.

15 Detection may be achieved through the use of immunocytochemistry, ELISA, radioimmunoassay or other assays as commonly known to one of ordinary skill in the art. The mox1, mox2, duox1 and duox2 proteins, or fragments thereof, may be labeled through commonly known approaches,
20 including but not limited to the following: radiolabeling, dyes, magnetic particles, biotin-avidin, fluorescent molecules, chemiluminescent molecules and systems, ferritin, colloidal gold, and other methods known to one of skill in the art of labeling proteins.

25

Administration of Antibodies

The antibodies directed to the proteins shown as SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48, or directed to fragments
30 thereof, may also be administered directly to humans and animals in a passive immunization paradigm. Antibodies directed to extracellular portions of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46 or SEQ ID NO:48 bind to these extracellular epitopes. Attachment of
35 labels to these antibodies facilitates localization and visualization of sites of binding. Attachment of molecules such as ricin or

other cytotoxins to these antibodies helps to selectively damage or kill cells expressing SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, SEQ ID NO:48 or fragments thereof.

5

Kits

The present invention includes kits useful with the antibodies, nucleic acids, nucleic acid probes, labeled antibodies, labeled proteins or fragments thereof for detection, localization and measurement of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48 or combinations and fragments thereof.

10

Kits may be used for immunocytochemistry, *in situ* hybridization, solution hybridization, radioimmunoassay, ELISA, Western blots, quantitative PCR, and other assays for the detection, localization and measurement of these nucleic acids, proteins or fragments thereof using techniques known to one of skill in the art.

15

20

The nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45, SEQ ID NO:47, or fragments thereof, may also be used under high stringency conditions to detect alternately spliced messages related to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45, SEQ ID NO:47 or fragments thereof, respectively.

25

As discussed in one of the Examples, rat mox1 protein (SEQ ID NO: 21) is similar to mouse gp91 protein (SEQ ID NO: 38), whereas rat mox1B protein (SEQ ID NO:42) is similar to human gp91 protein (SEQ ID NO:12). This observation suggests that other isoforms of mouse and human gp91 may exist. In addition, another subtype of human mox1, similar to rat mox1B (SEQ ID NO:42), also exists. The presence of two isoforms of rat mox1 protein in vascular smooth muscle may have important physiological consequences

30

35

and biomedical applications. For example, the two isoforms may have different biological activities, different tissue distributions and may be regulated differently in physiological and/or pathological conditions. The fact that mox1B (SEQ ID NO:42) was isolated from cells exposed to angiotensin II, known to promote oxidative stress and vascular growth, suggests that it may be upregulated by this hormone and may be overexpressed in disease. Therefore, the diagnostic kits of the present invention can measure the relative expression of the two mox isoforms. The diagnostic kits may also measure or detect the relative expression of the mox proteins described herein (i.e. human mox1 and/or human mox2) and duox proteins described herein (i.e. human duox1 and/or human duox2).

Fragments of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41 SEQ ID NO:45 and SEQ ID NO:47 containing the relevant hybridizing sequence can be synthesized onto the surface of a chip array. RNA samples, e.g., from tumors, are then fluorescently tagged and hybridized onto the chip for detection. This approach may be used diagnostically to characterize tumor types and to tailor treatments and/or provide prognostic information. Such prognostic information may have predictive value concerning disease progression and life span, and may also affect choice of therapy.

The present invention is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof, which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention.

EXAMPLE 1

Sequence Analysis and Cloning of the Human mox1 cDNA (SEQ ID NO:1) Encoding for Production of the Human mox1 Protein p65mox (SEQ ID NO:2)

5 Using gp91phox as a query sequence, a 334 base sequenced portion of expressed sequence tag (EST) 176696 (GenBank Accession number AA305700) showed 68.8% sequence identity at the predicted amino acid level with human (h) gp91phox. The bacterial strain number 129134 containing the EST sequence in the pBluescript SK⁻ vector, was purchased from American Tissue Type Culture Collection (ATCC, Rockville, MD). The EST sequence was originally cloned from a Caco-2 human colon carcinoma cell line. The EST176696 DNA was further sequenced using the T7 and T3 vector promoters and primers designed to match the known 3' sequence. Internal primers used for sequencing were as follows: 5'-AAC AAG CGT GGC TTC AGC ATG-3' SEQ ID NO:5 (251S, numbering is based on the nucleotides from the 5' end of EST176696, and S indicates the sense direction), 5'-AGC AAT ATT GTT GGT CAT-3' SEQ ID NO:6 (336S), 5'-GAC TTG ACA GAA AAT CTA TAA GGG-3' SEQ ID NO:7 (393S), 5'-TTG TAC CAG ATG GAT TTC AA-3' SEQ ID NO:8 (673A, A indicates the antisense direction), 5'-CAG GTC TGA AAC AGA AAA CCT-3' SEQ ID NO:9 (829S), 5'-ATG AAT TCT CAT TAA TTA TTC AAT AAA-3' SEQ ID NO:10 (1455A). The coding sequence in EST176696 showed homology to a 250 amino acid stretch corresponding to the N-terminal 44% of human gp91phox, and contained a stop codon corresponding to the location in human gp91phox. 5' Rapid amplification of cDNA ends (RACE) was carried out using a human colon cDNA library and Marathon cDNA Amplification Kit (ClonTech) using 5'-ATC TCA AAA GAC TCT GCA CA-3' SEQ ID NO:11 (41A) as an internal gene-specific primer (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85, 8998-9002). 5' RACE resulted in a 1.1 kb fragment representing the complete 5' sequence, based on homology with gp91phox.

Reamplification was performed with primers spanning the putative start and stop codons, using the 1.1 kb 5' RACE product and pSK-EST176696 for primer design. The amplified 1.7kb fragment was TA cloned into the PCR2.1 vector (Invitrogen, Carlsbad, CA). This recombinant vector is referred to as PCR-*mox*.

Figure 1(a-d) presents a comparison of the present amino acid sequences of human, bovine and murine gp91 phox with the human and rat mox1 proteins of the present invention and the human duox2 protein of the present invention. Also shown are the amino acid sequences for related plant enzyme proteins.

The encoded hp65mox ("mox" referring to mitogenic oxidase and "65" referring to its predicted molecular weight) is listed as SEQ ID NO:2. h-gp91phox (SEQ ID NO:12) and SEQ ID NO:2 differ in length by 3 residues and are 70% identical in their amino acid sequence. h-gp91phox and SEQ ID NO:2 show a greater percentage identity in the C-terminal half of the molecule which contains the putative NADPH and FAD binding sites, and there are several relatively long stretches of complete identity within this region.

A dendrogram (Figure 2) comparing the amino acid sequences of mouse and human gp91phox with that of mox1 SEQ ID NO:2 shows that the latter probably represents a distinct isoform of gp91phox. Two plant homologs of cytochrome b₅₅₈ large subunit are also indicated and represent more distant relatives of the human sequences. Human (and rat mox1 described more fully below) lack asparagine-linked glycosylation sites, which are seen in the highly glycosylated human and mouse gp91phox. Additionally, the hydropathy profiles of human gp91phox and mox1 are nearly identical and include five very hydrophobic stretches in the amino-terminal half of the molecules which are predicted to be membrane-spanning regions.

EXAMPLE 2*Expression of Mox1*

Human multiple tissue northern (MTN) Blot I and Human MTN Blot IV (ClonTech) membranes were hybridized with the putative coding region of the PCR-mox vector at 68°C for several hours. The mox coding region was labeled by random priming with [α -³²P]dCTP (10 μ Ci) using the Prime-It II kit (Stratagene). For analysis of mox1 expression in cell lines, total RNA was prepared from 10⁶ cells using the High Pure RNA Isolation Kit (Boehringer Mannheim) or RNeasy kit (Quiagen). Total RNA (10-20 μ g) was separated on a 1% agarose formaldehyde mini-gel and transferred to a Nytran filter (Biorad) and immobilized by ultraviolet cross-linking.

Northern blotting revealed that the major location of mRNA coding for the mox1 protein was colon. The message was also detected in prostate and uterus. The human colon-carcinoma cell line, Caco-2, also expressed large quantities of mox1 message. Northern blotting of mRNA from rat aortic smooth muscle cells also showed strong hybridization, which increased roughly two-fold within 12 hours after treatment with platelet-derived growth factor. This increase in the expression of rat mox1 is consistent with the idea that mox1 contributes to the growth-stimulatory effects of PDGF.

25

EXAMPLE 3*Transfection of NIH3T3 Cells with SEQ ID NO:1*

The nucleotide sequence (SEQ ID NO:1) encoding for production of the mox1 protein (SEQ ID NO:2) was subcloned into the *NotI* site of the pEF-PAC vector (obtained from Mary Dinanuer, Indiana University Medical School, Indianapolis, IN) which has a puromycin resistance gene. Transfection was carried out as described in Sambrook et al., Molecular Cloning, A Laboratory Manual, Volumes 1-3, 2nd edition, Cold Spring Harbor Laboratory Press, N.Y., 1989. The SEQ ID NO:1 in pEF-PAC and the empty vector were separately transfected into NIH 3T3 cells using Eugene 6

(Boeringer Mannheim). About 2×10^6 cells maintained in DMEM containing 10% calf serum were transfected with 10 μ g of DNA. After 2 days, cells were split and selected in the same medium containing 1mg/ml puromycin. Colonies that survived in selection media for 10 to 14 days were subcultured continuously in the presence of puromycin.

Transfected cells exhibited a "transformed"-like morphology, similar to that seen with (V12)Ras-transfected cells, characterized by long spindle-like cells. The parent NIH 3T3 cells or cells transfected with the empty vector showed a normal fibroblast-like morphology.

EXAMPLE 4

Expression of Mox1 (SEQ ID NO:1) in Transfected NIH3T3 Cells

To verify the expression of *mox1* mRNA after transfection, RT-PCR and Northern blotting were performed. Total RNAs were prepared from 10^6 cells using the High Pure RNA Isolation Kit (Boeringer Mannheim) or RNeasy kit (Qiagen). cDNAs for each colony were prepared from 1-2 μ g of total RNA using Advantage RT-PCR Kit (ClonTech). PCR amplification was performed using primers, 5'-TTG GCT AAA TCC CAT CCA-3' SEQ ID NO:13 (NN459S, numbering containing NN indicates numbering from the start codon of *mox1*) and 5'-TGC ATG ACC AAC AAT ATT GCT G-3' SEQ ID NO:14 (NN1435A). For Northern blotting, 10-20 μ g of total RNA was separated on a 1% agarose formaldehyde gel and transferred to a nylon filter. After ultraviolet (UV) cross-linking, filters were used for Northern blotting assay as described in Example 2.

Colonies expressing large amounts of *mox1* mRNA were chosen for further analysis. The expression of mRNA for glyceraldehyde 3 phosphate dehydrogenase in the various cell lines was normal.

EXAMPLE 5

Colony Formation on Soft Agar

5 10^5 to 10^3 cells stably transfected with human mox1 gene SEQ ID NO:1 and with empty vector were prepared in 0.3% warm (40°C) agar solution containing DMEM and 10% calf serum. Cells were distributed onto a hardened 0.6% agar plate prepared with DMEM and 10% calf serum. After three weeks in culture (37°C, 5% CO₂) colony formation was observed by microscopy.

10 Cells which were stably transfected with the empty vector and cultured in soft agar for 3 weeks as above did not display anchorage independent growth. In contrast, NIH 3T3 cells which had been stably transfected with the mox1 (SEQ ID NO:1) and cultured for 3 weeks in soft agar demonstrated anchorage independent growth of colonies.

15

EXAMPLE 6*NADPH-Dependent Superoxide Generation Assay*

20 In one embodiment of the present invention, NIH 3T3 cells stably transfected with the human mox1 gene (SEQ ID NO:1) were analyzed for superoxide generation using the lucigenin (Bis-N-methylacridinium luminescence assay (Sigma, St. Louis, MO, Li et al. (1998) *J. Biol. Chem.* 273, 2015-2023). Cells were washed with cold HANKS' solution and homogenized on ice in HANKS' buffer containing 15% sucrose using a Dounce homogenizer. Cell lysates were frozen immediately in a dry ice/ethanol bath. For the assay, 30 µg of cell lysate was mixed with 200 µM NADPH and 500 µM lucigenin. Luminescence was monitored using a LumiCounter (Packard) at three successive one minute intervals and the highest value was used for comparison. Protein concentration was determined by the Bradford method.

25

30

35 Superoxide generation was monitored in lysates from some of the stably transfected cell lines and was compared with superoxide generation by the untransfected NIH 3T3 cell lysates. The results are shown in Table 4. Cell lines 26, 27, and 28 gave the highest degree of morphological changes by

microscopic examination corresponding to the highest degree of superoxide generation. The luminescent signal was inhibited by superoxide dismutase and the general flavoprotein inhibitor diphenylene iodonium, but was unaffected by added recombinant human p47phox, p67phox and Rac1(GTP- γ S), which are essential cytosolic factors for the phagocyte respiratory-burst oxidase.

Table 4

10	<u>Cell Line Name</u>	<u>Superoxide Generation</u> (RLU)
	Control (untransfected)	6045
	mox1-26	17027
	mox1-27	14670
15	mox1-28	18411
	mox1-65	5431
	mox1-615	11331
	mox1-+3	8645
	mox1-+10	5425
20	mox1-pcc16	8050

In an alternate and preferred embodiment of the present invention, cells that had been stably transfected with mox1 (YA28) or with empty vector (NEF2) were grown in 10 cm tissue culture plates in medium containing DMEM, 10% calf serum, 100 units/ml penicillin, 100 μ g/ml streptomycin, and 1 μ g/ml puromycin to approximately 80% confluency. Cells (five tissue culture plates of each cell type) were washed briefly with 5 ml phosphate buffered saline (PBS) then dissociated from the plates with PBS containing 5 mM EDTA. Cells were pelleted by centrifuging briefly at 1000 x g.

To permeabilize the cells, freeze thaw lysis was carried out and this was followed by passage of the cell material through a small bore needle. The supernatant was removed and the cells were frozen on dry ice for 15 minutes. After cells were thawed, 200 μ l lysis buffer (HANKS' Buffered Salt

Solution - HBBS) containing a mixture of protease inhibitors from Sigma (Catalog # P2714) was added. Cells on ice were passed through an 18 guage needle 10 times and 200 μ l of HBSS buffer containing 34% sucrose was added to yield a final concentration of 17% sucrose. Sucrose appeared to enhance stability upon storage. The combination of freeze-thawing and passage through a needle results in lysis of essentially all of the cells, and this material is referred to as the "cell lysate."

The cell lysates were assayed for protein concentration using the BioRad protein assay system. Cell lysates were assayed for NADPH-dependent chemiluminescence by combining HBSS buffer, arachidonic acid, and 0.01 – 1 μ g protein in assay plates (96 well plastic plates). The reaction was initiated by adding 1.5 mM NADPH and 75 μ M lucigenin to the assay mix to give a final concentration of 200 μ M NADPH and 10 μ M lucigenin, and the chemiluminescence was monitored immediately. The final assay volume as 150 μ l. The optimal arachidonic acid concentration was between 50-100 μ M. A Packard Lumicount luminometer was used to measure chemiluminescence of the reaction between lucigenin and superoxide at 37°C. The plate was monitored continuously for 60 minutes and the maximal relative luminescence unit (RLU) value for each sample was used for the graph.

Figure 3 shows the RLU at various concentrations of cell lysates from mox1-transfected (YA28) and vector control (NEF2) cells. The presence of NaCl or KCl within a concentration range of 50-150 μ M is important for optimal activity. $MgCl_2$ (1-5 mM) further enhanced activity by about 2-fold. This cell-free assay for mox1 NADPH-oxidase activity is useful for screening modulators (inhibitors or stimulators) of the mox1 enzyme. The assay may also be used to detect mox and duox NADPH-oxidase activity in general and to screen for modulators (inhibitors or stimulators) of the mox and duox family of enzymes.

EXAMPLE 7*Nitro Blue Tetrazolium Reduction by Superoxide Generated by NIH 3T3 cells Transfected with the Mox1 cDNA (SEQ ID NO:1)*

5 Superoxide generation by intact cells was monitored by using superoxide dismutase-sensitive reduction of nitroblue tetrazolium. NEF2 (vector alone control), YA26 (mox1 (SEQ ID NO:1)-transfected) and YA28 (mox1 (SEQ ID NO:1)-transfected) cells were plated in six well plates at 10 500,000 cells per well. About 24 hours later, medium was removed from cells and the cells were washed once with 1 mL Hanks solution (Sigma, St. Louis, MO). About 1 mL of filtered 0.25% Nitro blue tetrazolium (NBT, Sigma) was added in Hanks without or with 600 units of superoxide dismutase 15 (Sigma) and cells were incubated at 37°C in the presence of 5% CO₂. After 8 minutes the cells were scraped and pelleted at more than 10,000g. The pellet was re-suspended in 1 mL of pyridine (Sigma) and heated for 10 minutes at 100°C to solubilize the reduced NBT. The concentration of reduced NBT 20 was determined by measuring the absorbance at 510 nm, using an extinction coefficient of 11,000 M⁻¹cm⁻¹. Some wells were untreated and used to determine cell number.

25 The data are presented in Table 5 and Figure 4 and indicate that the mox1 (SEQ ID NO:1)-transfected cells generated significant quantities of superoxide.

Table 5

	<u>NBT Reduction (nmols/10⁶ cells)</u>	<u>- SOD</u>	<u>+ SOD</u>
30	vector control cells	2.5 ± 0.5	2.1 ± 0.5
	YA26 (mox1) cells	6.4 ± 0.2	3.4 ± 0.1
	YA28 (mox1) cells	5.2 ± 0.6	3.4 ± 0.3

-SOD, and +SOD mean in the absence or presence of added superoxide dismutase, respectively.

35

Because superoxide dismutase is not likely to penetrate cells, superoxide must be generated extracellularly. The amount of

superoxide generated by these cells is about 5-10% of that generated by activated human neutrophils.

EXAMPLE 8

5 *Modification of Intracellular Components in Mox1 Transfected Cells*

To test whether superoxide generated by *mox1* can affect intracellular "targets," aconitase activity in control and *mox*-transfected cell lines was monitored as described in Suh et al. (1999) *Nature* 401, 79-82. Aconitase contains a four-iron-sulphur cluster that is highly susceptible to modification by superoxide, resulting in a loss of activity, and has been used as a reporter of intra-cellular superoxide generation. Aconitase activity was determined as described in Gardner et al. (1995) *J. Biol. Chem.* 270, 13399-13405. Aconitase activity was significantly diminished in all three *mox*-transfected cell lines designated YA26, YA28 and YA212 as compared to the transfected control (Figure 5). Approximately 50% of the aconitase in these cells is mitochondrial, based on differential centrifugation, and the cytosolic and mitochondrial forms were both affected. Control cytosolic and mitochondrial enzymes that do not contain iron-sulfur centres were not affected. Superoxide generated in *mox1*-transfected cells is therefore capable of reacting with and modifying intracellular components.

EXAMPLE 9

30 *Tumor Generation in Nude Mice Receiving Cells Transfected with the Human *mox1* cDNA (SEQ ID NO:1)*

About 2×10^6 NIH 3T3 cells (either *mox1*-transfected with SEQ ID NO:1 or cells transfected using empty vector) were injected subdermally into the lateral aspect of the neck of 4-5 week old nude mice. Three to six mice were injected for each of three *mox1*-transfected cell lines, and 3 mice were injected with the cells transfected with empty vector (control). After 2 to 3 weeks, mice were sacrificed. The tumors were fixed in 10% formalin and characterized by

5 histological analysis. Tumors averaged 1.5 x 1 x 1 cm in size and showed histology typical of sarcoma type tumors. In addition, tumors appeared to be highly vascularized with superficial capillaries. Eleven of twelve mice injected with mox1 gene-transfected cells developed tumors, while none of the three control animals developed tumors.

10 In another study, 15 mice were injected with mox1-transfected NIH 3T3 cells. Of the 15 mice injected, 14 showed large tumors within 17 days of injection, and tumors showed expression of mox1 mRNA. Histologically, the tumors resembled fibrosarcomas and were similar to ras-induced tumors. Thus, ras and mox1 were similarly potent in their ability to induce tumorigenicity of NIH 3T3 cells in athymic mice.

15

EXAMPLE 10

Demonstration of the Role of Mox1 in Non-Cancerous Growth

20 A role in normal growth was demonstrated in rat aortic vascular smooth-muscle cells by using antisense to rat mox1. Transfection with the antisense DNA resulted in a decrease in both superoxide generation and serum-dependent growth. Mox1 is therefore implicated in normal growth in this cell type.

25

EXAMPLE 11

Expression of Human Mox1 Protein (SEQ ID NO:2) in a Baculovirus Expression System

30 SEQ ID NO:2 was also expressed in insect cells using recombinant baculovirus. To establish the p65mox1 expressing virus system, the mox1 gene (SEQ ID NO:1) was initially cloned into the pBacPAK8 vector (Clontech, Palo Alto, CA) and recombinant baculovirus was constructed using standard methods according to manufacturer's protocols. Briefly, PCR amplified mox1 DNA was cloned into the KpnI and EcoRI site of the vector. Primers used for PCR amplification were: 5'-CAA GGT ACC TCT TGA CCA TGG

35

5 GAA ACT-3' , SEQ ID NO:15, and 5'-ACG AAT TCA AGT
AAA TTA CTG AAG ATA C-3' , SEQ ID NO:16. Sf9 insect
cells (2×10^6 cells) were infected with 0.5 mg of linearized
baculovirus DNA sold under the trademark BACULOGOLD®
10 (PharMingen, San Diego, CA) and 5 mg pBacPAC8-p65mox1
using Transfection Buffers A and B (PharMingen, San Diego,
CA). After 5 days, the supernatants containing recombinant
viruses were harvested and amplified by infecting fresh sf9 cells
for 7 days. Amplification was carried out three times and the
15 presence of the recombinant virus containing mox1 DNA was
confirmed by PCR using the same primers. After three times
amplification of viruses, plaque purification was carried out to
obtain the high titer viruses. Approximately 2×10^8 sf9 cells in
agar plates were infected for 5 days with serial dilutions of
virus and were dyed with neutral red for easy detection of virus
plaques. Selected virus plaques were extracted and the presence
of the human mox1 DNA was confirmed again by PCR.

EXAMPLE 12

20 *Cloning of a Rat Homolog of p65mox (SEQ ID NO:2)*

cDNA clones of p65mox from a rat aortic smooth
muscle cell have been obtained. RT-PCR (reverse transcription
polymerase chain reaction) was carried out as follows: first
strand cDNA synthesis was performed using total RNA from rat
25 aortic vascular smooth muscle cells, oligo dT primer and
superscript II reverse transcriptase, and followed by incubation
with RNase H. Degenerate PCR primers were designed to
anneal to conserved areas in the coding regions of h-mox1 and
gp91phox of human (X04011), mouse (U43384) and porcine
30 (SSU02476) origin. Primers were: sense 5'-
CCIGTITGTCGIAATCTGCTSTCCTT-3', SEQ ID NO:17 and
antisense 5'-TCCCIGCAIAICCAGTAGAARTAGATCTT-3',
SEQ ID NO:18. A major PCR product of the expected 1.1 kb
size was purified by agarose electrophoresis and used as
35 template in a second PCR amplification reaction.

An aliquot of the RT-PCR product was blunt-

ended, ligated into a modified Litmus 29 vector and used to transform XL10 competent *E. coli*. Approximately 120 bacterial colonies were screened for the presence of a full-length insert by direct PCR using vector primers and Taq polymerase. Plasmids were purified from 25 positive colonies and mapped by digestion with *Bam* HI. Representative plasmids from each digestion pattern were partially sequenced. Five out of 25 clones contained non-specific amplification products and 20 contained identical inserts similar to human (h)-mox1. One of the latter clones was fully sequenced and found to be 83% identical to h-mox1 over 1060 nucleotides. A 1.1 kb probe was generated by PCR amplification of the insert of a rat mox1 clone with the degenerate primers described above and used to hybridize to a Northern blot of rat vascular smooth muscle cell RNA. A single band, migrating between 28S rRNA and 18S rRNA, indicated the presence of a message with a size compatible to that of human mox-1 (2.6 kb).

To obtain full-length rat mox1, 3' and 5' rapid amplification of cDNA ends (RACE) reactions were performed as describe above, using the gene-specific primers 5'-TTGGCACAGTCAGTGAGGATGTCTTC-3', SEQ ID NO:19 and 5'-CTGTTGGCTTCTACTGTAGCGTTCAAAGTT-3', SEQ ID NO:20 for 3' and 5' RACE, respectively. Single major 1.5 kb and 850 bp products were obtained for 3' and 5' RACE, respectively. These products were purified by agarose gel eletrophoresis and reamplified with Taq polymerase. Both products were cloned into the pCR 2.1 vector and used to transform electrocompetent XL1 blue *E. coli*. The RACE products were sequenced and new terminal primers were designed:

sense 5'-TTCTGAGTAGGTGTGCATTTGAGTGTCATAAAGAC-3' (SEQ ID NO:43), and antisense 5'-TTTTCCGTCAAAATTATAACTTTTTATTTTCTTTTATAACACAT-3' (SEQ ID NO:44). PCR amplification of rat VSMC cDNA was performed using these primers.

5 A single 2.6 kb product was obtained, ligated into
pCR 2.1 and used to transform electrocompetent XL1 blue *E.*
coli. The insert was sequenced with 12 sense and 14 antisense
primers. Its length is 2577 bp (including primer sequences),
10 comprising a 1692 bp open reading frame, 127 bp 5' and 758
bp 3' untranslated regions. The presence of six in-frame stop
codons in the 5' untranslated region suggests that the full length
coding region has been obtained. Consensus polyadenylation
sequences are present at nucleotides 2201 and 2550. Conceptual
translation yields a 563 amino acid peptide, one residue shorter
than the human deduced sequence. This new amino acid
sequence is more similar to human mox1 SEQ ID NO:3 (82%
identity) than to mouse gp91phox SEQ ID NO:38 (55%
identity), suggesting that it is indeed rat mox1 (SEQ ID NO:21).
15 This rat (r) homolog of p65mox protein is called r-p65mox or
p65mox/rat.pep and is shown as SEQ ID NO:21. The
nucleotide sequence encoding for r-p65mox is shown as SEQ ID
NO:22.

20

EXAMPLE 13

*Expression of rat (r)-p65mox mRNA in Vascular Smooth
Muscle and Induction by Angiotensin II, Platelet-Derived
Growth Factor (PDGF), and Phorbol Myristic Acid (PMA)*

25

30

35

Using the partial cDNA clone from rat, we have
examined cultured rat aortic smooth muscle cells for expression
of message for r-p65mox. We have observed the mRNA for r-
p65mox in these cells. It has been previously reported
(Griendling et al. (1994) *Circ. Res.* 74, 1141-1148; Fukui et al.
(1997) *Circ. Res.* 80, 45-51; Ushio-Fukai et al. (1996) *J. Biol.*
Chem. 271, 23317-23321) that *in vitro* or *in vivo* treatment
with angiotensin II (AII) is a growth stimulus for vascular
smooth muscle cells, and that AII induces increased superoxide
generation in these cells. Platelet-derived growth factor
(PDGF) and PMA are proliferative signals for vascular smooth
muscle cells. We observed that the mRNA for r-p65mox was
induced approximately 2-3 fold by angiotensin II (100 nM),

corresponding to the increased level of superoxide generation. Thus, the increased superoxide generation in these cells correlates with increased expression of the mRNA for this enzyme. The mRNA for r-p65mox also increased 2 or more fold in response to the growth stimulus PDGF (20 ng/ml), and 2-3 fold in response to PMA. Quantitation by densitometry revealed that rat mox1 message was induced nearly 4-fold at the 6 and 12 hour time points in response to PDGF, and about 2-fold at the 12 hour time point in response to AII. 28S RNA was used as a control for RNA recovery.

EXAMPLE 14

Antibodies to Fragments of Human (h)-p65mox (SEQ ID NO:2)

Polyclonal antibodies were raised in rabbits against the C-terminal half of h-p65mox (residues 233 through 564, SEQ ID NO:23) which is predicted to fold into a cytosolic domain containing FAD and the NADPH or NADH binding site. This domain was expressed in *E. coli* as an N-terminal GST-fusion protein and was purified on glutathione agarose by standard methods. Two antipeptide antibodies were also made against h-p65mox (residues 243-256, referred to as Peptide A, SEQ ID NO:24) and h-65mox (residues 538-551, referred to as Peptide B, SEQ ID NO:25). Peptides were conjugated to keyhole limpet hemocyanin (KLH) using glutaraldehyde.

Antigens were injected into different rabbits initially in complete Freund's adjuvant, and were boosted 4 times with antigen in incomplete Freund's adjuvant at intervals of every three weeks. Approximately 0.5 mg to 1 mg of peptide was administered at each injection. Blood was drawn 1 week after each boost and a terminal bleed was carried out 2 weeks after the final boost. Antibodies to Peptide A and Peptide B were affinity purified by column chromatography through peptide A or peptide B conjugated to Affigel 15 (Bio-Rad, Richmond, CA). 10 mg of peptide was covalently crosslinked to 2 ml of Affigel 15 resin and the gel was washed with 20 ml of binding buffer (20 mM Hepes/NaOH, pH 7.0, 200 mM NaCl,

and 0.5 % Triton X-100). The remaining functional N-hydrosuccinimide was blocked with 100 μ l of 1 M ethanolamine. After washing with 20 ml of binding buffer, 5 ml of the antiserum was incubated with the pep A-conjugated Affigel 15 resin overnight at 4°C. Unbound protein was washed away with 20 ml of binding buffer. Elution of the antibodies from the gel was performed with 6 ml of elution buffer (100 mM glycine/HCl, pH 2.5, 200 mM NaCl, and 0.5% Triton X-100). The eluate was then neutralized by adding 0.9 ml of 1 M Tris/HCl, pH 8.0. The GST-fusion form of truncated p65mox1 protein (residues 233-566, SEQ ID NO:23) was expressed in *E. coli*. Samples (20 μ g each) were run on 12 % SDS-PAGE either before or 1 or 4 hours after induction with 100 μ M IPTG (isopropyl β -thiogalactoside).

The extracted proteins were subjected to immunoprob- ing with affinity purified antiserum to peptide A at a 1:1000 dilution. The detection of antigens was performed using an enhanced chemiluminescence kit (Amersham, Buckinghamshire, UK). The affinity purified antibody to mox1 (243-256, SEQ ID NO:24) was used at a dilution of 1:1000 in a Western blot in which a total of 10 μ g of protein was added to each lane. The major band observed at 4 hours after IPTG induction corresponded to the size of the GST-mox1 expressed in bacteria containing the pGEX-2T vector encoding the GST-mox1 fusion protein.

Example 15

Presence of an NAD(P)H Oxidase in Ras-Transformed Fibroblasts

A superoxide-generating NADPH oxidase activity was detected in homogenates from NIH 3T3 cells, and this activity increased about 10-15 fold in Ras-transformed NIH 3T3 cells (Table 6). To establish the stable Ras-transformed cell lines, the DNA for human Ras encoding an activating mutation at amino acid number 12 (Valine, referred to as V12-Ras) was subcloned into BamH1 and EcoR1 sites of pCDNA3 vector

which has a neomycin resistance gene. V12-Ras in pCDNA3 and empty vector were transfected into NIH 3T3 cells using Lipofectamine Plus (Gibco). 2×10^6 cells were maintained with DMEM containing 10% calf serum and transfected with 1 mg of DNA. After 2-days, cells were split and selected with the same medium but containing 1 mg/ml neomycin. Colonies surviving in selection media for 10 to 14 days were sub-cultured and characterized by immunoblot analysis using antibody against human H-Ras.

The expression of Ras in cells transfected with pcDNA-3 vector alone or in three cell lines transfected with V12-Ras in the same vector was analyzed on a Western blot. The three cell lines were named V12-Ras-7, V12-Ras-4, and V12-Ras-8. The expression of V12-Ras varied widely among the three cell lines tested. The V12-Ras-4 cell line expressed the highest level of Ras followed by the V12-Ras-8 cell line. The V12-Ras-7 cell line expressed the lowest level of Ras.

Lysates from each of these lines were then prepared and tested for their ability to generate superoxide. For each cell line, cells were washed with cold HANKS' balanced salt solution (HBSS), collected by centrifugation, kept on dry-ice for more than 30 min, and disrupted by suspending in low salt buffer (LSB; 50 mM Tris/HCl, pH 7.5, 1 mM PMSF, and protease cocktail from Sigma) and passing through a syringe needle (18 gauge) ten times. Cell lysates were frozen in dry-ice immediately after determining the protein concentration.

Table 6 shows superoxide generation in the transfected cells measured using the lucigenin luminescence assay. For the assay, 5 μ g of cell lysates were incubated with the reaction mixture containing 10 μ M lucigenin (luminescent probe) and 100 μ M NADPH (substrate) in the presence or absence of 100 μ M arachidonate in the absence or presence of 100 U of superoxide dismutase (SOD) or 1 μ M diphenyleneiodonium (DPI). Luminescence of the reaction mixture was monitored for 0.5 second by LumiCounter (Packard) for four times at 3 second intervals. RLU in Table 5

refers to relative luminescence units.

As shown in Table 6, the luminescence was partially inhibited by superoxide dismutase indicating that the signal was due at least in part to the generation of superoxide. DPI, a known inhibitor of both neutrophil and non-neutrophil NADPH oxidase activities, completely inhibited activity. The generation of superoxide correlated with the expression of Ras in the three cell lines. Thus, oncogenic Ras appears to induce an NADPH-dependent superoxide generating activity that is similar to the activity catalyzed by p65mox1.

Table 6

		RLU/5 μ g protein		
		no additions	plus SOD	plus DPI
15	Vector Control (1)	465	154	48
	V12-Ras-7 (2)	1680	578	39
	V12-Ras-4 (3)	5975	2128	36
	V12-Ras-8 (4)	4883	2000	35

20

EXAMPLE 16

Molecular Cloning of Another Rat mox1 cDNA Called Rat mox1B

A rat cDNA library was screened in an effort to identify new rat mox sequences. The library was constructed in a ZAP express lambda phage vector (Stratagene, La Jolla, CA) using RNA isolated from rat vascular smooth muscle cells which had been exposed to 100 nM angiotensin II for 4 hours. The library was screened using standard blot hybridization techniques with the rat mox1 probe described previously. Fifteen individual clones were obtained that were characterized by PCR and restriction mapping. Two different types of clones were thus identified and representatives of each type were sequenced. A clone of the first type (representative of 13) was found to be similar to the previously identified rat mox1 and

35

was thus named rat *mox1B*. Clones of the second type (representative of 2) were incomplete rat *mox* sequences.

5 The length of the rat *mox1B* nucleotide sequence is 2619 bp and is listed as SEQ ID NO:41. The single longest 1497 bp open reading frame encompasses nucleotides 362 to 1858. The presence of two in-frame stop codons in the 5' untranslated region at nucleotides 74 and 257 indicates that the full-length coding region has been isolated. Two putative polyadenylation sites are present at positions 2243 and 2592. 10 Alignment of the rat *mox1* nucleotide sequence (SEQ ID NO:22) and the rat *mox1B* nucleotide sequence (SEQ ID NO:41) shows that the two nucleotides sequences are identical except at their 5' ends, suggesting that they may represent two alternatively spliced messages from the same gene. Sequence 15 identity starts at nucleotides 269 and 311, for rat *mox1* and rat *mox1B*, respectively.

Conceptual translation of the rat *mox1B* nucleotide sequence (SEQ ID NO:41) yields a 499 amino acid sequence with a predicted molecular weight of 58 kDa. This amino acid 20 sequence for rat *mox1B* protein is shown in SEQ ID NO:42. Alignment of the deduced amino acid sequences for rat *mox1* (SEQ ID NO:21) and rat *mox1B* (SEQ ID NO:42) indicates that rat *mox1B* is identical to rat *mox1A*, except for a missing stretch of 64 residues at the N-terminus. Therefore, rat *mox1B* 25 appears to be a splicing variant derived from the same gene as rat *mox1*.

EXAMPLE 17

30 *Sequence Analysis and Cloning of the Human Mox2 cDNA (SEQ ID NO:3) Encoding for Production of the Human Mox2 Protein (SEQ ID NO:4)*

Note that the *mox2* protein as described herein, was described in U.S. Provisional Application Serial No. 60/149,332 as *mox3*.

35 A blast search was carried out using the sequence of *mox1* as a query sequence. The sequence identified by this

5 search was a sequence present in the GenBank database that
contains regions of homology with *mox1* and *gp91phox*. The
GenBank sequence located in the search was a 90.6 kb
sequenced region of human chromosome 6 (6q25.1-26) that was
10 reported as a GenBank direct submission dated February 9,
1999 and given the Accession No. AL031773. Sequencing was
carried out as part of the human genome sequencing project by
S. Palmer, at Sanger Centre, in Hinxton, Cambridgeshire, UK.
The GenBank sequence was reported as being similar to
15 "Cytochrome B" and was not reported as having any homology
or relation to a *mox* protein. The sequence contained a
theoretical amino acid sequence that was derived by computer
using an algorithm that predicted intron/exon boundaries and
coding regions. This predicted region contained a 545 amino
acid sequence that was 56% identical to *mox1* and 58% identical
to *gp91phox*.

In the present invention, based on the GenBank
genomic sequence and the homologies described above, several
specific primers were designed and used to determine the tissue
20 expression patterns of a novel *mox* protein, *mox2*, using Human
Multiple Tissue PCR Panels (Clontech, Palo Alto, CA). The
primers were as follows: Primer 1: 5'-
CCTGACAGATGTATTTCACTACCCAG-3' (SEQ ID NO:49);
Primer 2: 5'-GGATCGGAGTCACTCCCTTCGCTG-3' (SEQ
25 ID NO:50); Primer 3: 5'-
CTAGAAGCTCTCCTTGTTGTAATAGA-3' (SEQ ID NO:51);
Primer 4: 5'-ATGAACACCTCTGGGGTCAGCTGA-3' (SEQ
ID NO:52). It was determined that *mox2* is expressed primarily
in fetal tissues, with highest expression in fetal kidney, with
30 expression also seen in fetal liver, fetal lung, fetal brain, fetal
spleen and fetal thymus. Among 16 adult tissues tested, *mox2*
expression was seen in brain, kidney, colon and lung, although
levels of expression appeared to be very low.

35 Additionally, the 5' RACE (RACE = Rapid
Amplification of cDNA Ends) and 3' RACE techniques were
used to complete the sequence of the 5' and 3' regions of *mox2*.

(5' RACE kit and 3' RACE kit were from Clontech, Palo Alto, CA and are more fully described in Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85, 8998-9002. The 5' RACE and 3'-RACE techniques were carried out using a human fetal kidney library (Marathon-Ready cDNA library, Cat. #7423-1), using the following specific primers: 5'-RACE: Primer 4: 5'-ATGAACACCTCTGGGGTCAGCTGA-3' (SEQ ID NO:53); Primer 5: 5'-GTCCTCTGCAGCATTGTTCTCTTA-3' (SEQ ID NO:54); 3'-RACE: Primer 1: 5'-CCTGACAGATGTATTTCACTACCCAG-3' (SEQ ID NO:55); Primer 2: 5'-GGATCGGAGTCACTCCCTTCGCTG-3' (SEQ ID NO:56). The RACE procedures were successful in completing the 5' sequence and in confirming the 3' sequence. The complete coding sequence of *mox2* is shown in SEQ ID NO:2, while the predicted amino acid sequence of *mox2* is shown in SEQ ID NO:4.

In comparing the sequences of the present invention to the predicted coding regions of the GenBank sequence, the GenBank sequence did not contain a start codon, appeared to be missing approximately 45 base pairs at the N-terminus, and contained one other major difference in the predicted coding region which could have been due to inaccurate computer prediction of intron/exon boundaries.

25

EXAMPLE 18

Sequence Analysis and Partial Cloning of the Human Duox2 cDNA (SEQ ID NO:47) Encoding for Production of the Human Duox2 Protein (SEQ ID NO:48)

A partial cDNA clone of *duox2* was obtained as follows. A 535-base portion of an expressed sequence tag (EST *zc92h03.rl*; Genbank accession no. W52750) from human pancreatic islet was identified using the human *gp91phox* amino-acid sequence as a query in a Blast search. The bacterial strain #595758 containing the EST sequence *zc92h03.rl* in the pBluescript SK-vector was purchased from ATCC (Rockville, MD). The DNA inserted into the pBluescript SK-vector was

further sequenced using T7 and T3 vector promoters as well as sequence specific internal primers. The EST encoded 440 amino acids showing a 24.4% identity to gp91phox, including a stop codon corresponding to the C-terminus of gp91phox. 5'-RACE was carried out using mRNA obtained from human colon carcinoma cells (CaCo2) and the Marathon cDNA Amplification Kit (ClonTech, Palo Alto). The following gene-specific primers were used for this procedure: 5'-GAAGTGGTGGGAGGCGAAGACATA-3' (SEQ ID NO:26) and 5'-CCTGTCATACCTGGGACGGTCTGG-3' (SEQ ID NO:27).

The results of the 5'-RACE yielded an additional 2 kilobase of sequenced DNA but this region did not contain the start codon. To complete the sequence of the 5' and 3' regions of duox2, 5'-RACE and 3'-RACE were carried out using a human adult pancreas mRNA (Clontech, Palo Alto, CA) with the kit of 5' RACE System for Rapid Amplification of cDNA Ends version 2.0 (Gibco BRL, Gaithersburg, MD). PCR done using the following specific primers resulted in a total predicted amino acid sequence of about 1000 residues: 5'-RACE: Primer 3: 5'-GAGCACAGTGAGATGCCTGTTCAG-3' (SEQ ID NO:28); Primer 4: 5'-GGAAGGCAGCAGAGAGCAATGATG-3' (SEQ ID NO:29) (for nested PCR); 3'-RACE Primer 5: 5'-ACATCTGCGAGCGGCACTTCCAGA-3' (SEQ ID NO:30) Primer 6: 5'-AGCTCGTCAACAGGCAGGACCGAGC-3' (SEQ ID NO:31) (for nested PCR).

EXAMPLE 19

Sequence Analysis and Cloning of the Human Duox1 cDNA (SEQ ID NO:45) Encoding for Production of the Human Duox1 Protein (SEQ ID NO:46)

A cDNA clone of duox1 was obtained as follows. A homologous 357-base portion of an expressed sequence tag (EST nr80d12.s1; Genbank accession no. AA641653) from an invasive human prostate was identified by using the partial

duox2 predicted amino-acid sequence described above as a query in a Blast search. The bacterial strain #1441736 containing the EST sequence nr80d12.s1 in the pBluescript SK-vector was purchased from ATCC (Rockville, MD). The DNA inserted into the pBluescript SK-vector was further sequenced using T7 and T3 vector promoters as well as sequence specific internal primers. The EST insert encoded 673 amino acids with no start or stop codons present. Northern Blot analysis of duox1 indicated the gene was about 5.5 kilobase pairs. To complete the sequence of 5' and 3' regions of duox1, 5' RACE and 3'-RACE were carried out using a human adult lung mRNA (Clontech, Palo Alto, CA) with the kit of 5' RACE System for Rapid Amplification of cDNA Ends version 2.0 (Gibco BRL, Gaithersburg, MD). The RACE procedure was carried out using the following specific primers: 5'-RACE: Primer 5: 5'-GCAGTGCATCCACATCTTCAGCAC-3'(SEQ ID NO:32); Primer 6: 5'-GAGAGCTCTGGAGACACTTGAGTTC-3' (SEQ ID NO:33) (for nested PCR); 3'-RACE Primer 7: 5'-CATGTTCTCTCTGGCTGACAAG-3' (SEQ ID NO:34); Primer 8: 5'-CACAATAGCGAGCTCCGCTTCACGC-3' (SEQ ID NO:35) (for nested PCR). RACE procedures were successful in completing the 5' sequence and the 3' sequence of duox1. The open reading frame is approximately 4563 base pairs.

25

EXAMPLE 20

Tissue Expression of Duox1 and Duox2

Based on the duox1 sequence data, several specific primers were designed (Primer 1a: 5'-GCAGGACATCAACCCTGCACTCTC-3' (SEQ ID NO:36); Primer 2a: 5'-AATGACACTGTACTGGAGGCCACAG-3' (SEQ ID NO:57); Primer 3a: 5'-CTGCCATCTACCACACGGATCTGC-3' (SEQ ID NO:58); Primer 4a: 5'-CTTGCCATTCCAAAGCTTCCATGC-3' (SEQ ID NO:59) and used these to determine the tissue expression patterns of duox1 using Human Multiple Tissue PCR Panels

30

35

(Clontech, Palo Alto, CA). It was determined that duox1 is expressed primarily in lung, testis, placenta, prostate, pancreas, fetal heart, fetal kidney, fetal liver, fetal lung, fetal skeletal muscle and thymus, with highest expression in adult and fetal lung. Among 16 adult tissues and 8 fetal tissues tested, duox1 expression in brain, heart, kidney, colon, ovary, thymus, fetal brain and fetal spleen appeared to be low.

Two duox2 specific primers were also used to determine the tissue expression patterns of duox2 using Human Multiple Tissue PCR (polymerase chain reaction) Panels (Clontech, Palo Alto, CA). (Primer 1b: 5'-GTACAAGTCAGGACAGTGGGTGCG-3' (SEQ ID NO:60); Primer 2b: 5'-TGGATGATGTCAGCCAGCCACTCA-3' (SEQ ID NO:61)). Duox2 is expressed primarily in lung, pancreas, placenta, colon, prostate, testis and fetal tissues, with highest expression in adult lung and fetal tissues. Among 16 adult tissues and 8 fetal tissues tested, duox2 expression in brain, heart, kidney, liver, skeletal muscle, thymus and fetal brain appeared to be low.

EXAMPLE 21

Role of Duox1 and Duox2 in Collagen Crosslinking

To investigate a possible role for the human duox1 and duox2, the model organism *Caenorhabditis elegans* and a new reverse genetic tool, RNA interference (RNAi), were used to "knock out" the homologues of duox in this organism (Fire et al. (1998) *Nature* 391, 806-811). This technique involved injection of double stranded RNA encoding a segment of Ce-duox1 or Ce-duox2 into gonads of *C. elegans* N2 hermaphrodites. Injected worms were then allowed to lay eggs, and the harvested eggs were allowed to develop and the F1 progeny were scored for phenotypes. This procedure has been documented to "knock-out" the expression of the gene of interest (Fire et al. (1998) *Nature* 391, 806-811).

In the case of Ce-duox1 and Ce-duox2, the knockout animals resulted in a complex phenotype including

worms with large superficial blisters, short or "dumpy" worms, worms with locomotion disorders, and worms with retained eggs and/or larvae. Because of the high identity between Ce-duox1 and Ce-duox2, three different RNA constructs were predicted to knock out either both genes or Ce-duox2 alone. In all cases, essentially the same group of phenotypes was obtained. Most or all of these phenotypes had been described previously in *C. elegans* mutated in the collagen biosynthetic pathway. *C. elegans* has an extracellular structure known as the cuticle, a complex sheath composed largely of cross-linked collagen, which functions as the exoskeleton of the nematode. Cross-linking of collagen in nematodes occurs in part by cross-linking tyrosine residues, and peroxidases such as sea urchin ovoperoxidase and human myeloperoxidase have previously been shown to be capable of carrying out this reaction.

Based upon the similarities of the phenotypes obtained, the Ce-duox1/2 knockout worms were examined for the presence of dityrosine linkages, using an HPLC methodology (Andersen, S.O. (1966) *Acta Physiol. Scand.* 66, Suppl. 263-265; Abdelrahim et al. (1997) *J. Chromatogr. B Biomed. Sci. Appl.* 696, 175-182). It was determined that dityrosine linkages, while easily detected in the wild type worms, were almost completely lacking in the knockout worms. Thus, an inability to catalyze dityrosine cross-linking accounts for the phenotype of *C. elegans* failing to express Ce-duox1/2. These data support the concept that the duox enzymes in higher organisms can probably function in a similar manner to modulate the extracellular milieu, possibly the extracellular matrix and/or the basement membrane.

All patents, publications and abstracts cited above are incorporated herein by reference in their entirety. It should be understood that the foregoing relates only to preferred embodiments of the present invention and that numerous modifications or alterations may be made therein without

WO 00/28031

PCT/US99/26592

65

departing from the spirit and the scope of the present invention
as defined in the following claims.

WO 00/28031

PCT/US99/26592

SEQUENCE LISTING

<110> Emory University

<120> Novel Mitogenic Regulators

<130> 05501-0103wp

<140>

<141>

<160> 61

<170> PatentIn Ver. 2.0

<210> 1

<211> 2609

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (207)..(1901)

<400> 1

gctgatagca cagttctgtc cagagaagga aggcggaata aacttattca ttcccaggaa 60

ctcttgggggt aggtgtgtgt ttttcacatc ttaaaggctc acagaccctg cgctggacaa 120

atgttccatt cctgaaggac ctctccagaa tccggattgc tgaatcttcc ctgttgcccta 180

gaagggtccc aaaccacctc ttgaca atg gga aac tgg gtg gtt aac cac tgg 233

Met Gly Asn Trp Val Val Asn His Trp

1

5

ttt tca gtt ttg ttt ctg gtt gtt tgg tta ggg ctg aat gtt ttc ctg 281

Phe Ser Val Leu Phe Leu Val Val Trp Leu Gly Leu Asn Val Phe Leu

10

15

20

25

ttt gtg gat gcc ttc ctg aaa tat gag aag gcc gac aaa tac tac tac 329

Phe Val Asp Ala Phe Leu Lys Tyr Glu Lys Ala Asp Lys Tyr Tyr Tyr

30

35

40

aca aga aaa atc ctt ggg tca aca ttg gcc tgt gcc cga gcg tct gct 377

Thr Arg Lys Ile Leu Gly Ser Thr Leu Ala Cys Ala Arg Ala Ser Ala

45

50

55

ctc tgc ttg aat ttt aac agc acg ctg atc ctg ctt cct gtg tgt cgc 425

WO 00/28031

PCT/US99/26592

Leu	Cys	Leu	Asn	Phe	Asn	Ser	Thr	Leu	Ile	Leu	Leu	Pro	Val	Cys	Arg		
		60					65					70					
aat	ctg	ctg	tcc	ttc	ctg	agg	ggc	acc	tgc	tca	ttt	tgc	agc	cgc	aca	473	
Asn	Leu	Leu	Ser	Phe	Leu	Arg	Gly	Thr	Cys	Ser	Phe	Cys	Ser	Arg	Thr		
		75				80					85						
ctg	aga	aag	caa	ttg	gat	cac	aac	ctc	acc	ttc	cac	aag	ctg	gtg	gcc	521	
Leu	Arg	Lys	Gln	Leu	Asp	His	Asn	Leu	Thr	Phe	His	Lys	Leu	Val	Ala		
		90				95				100					105		
tat	atg	atc	tgc	cta	cat	aca	gct	att	cac	atc	att	gca	cac	ctg	ttt	569	
Tyr	Met	Ile	Cys	Leu	His	Thr	Ala	Ile	His	Ile	Ile	Ala	His	Leu	Phe		
				110					115					120			
aac	ttt	gac	tgc	tat	agc	aga	agc	cga	cag	gcc	aca	gat	ggc	tcc	ctt	617	
Asn	Phe	Asp	Cys	Tyr	Ser	Arg	Ser	Arg	Gln	Ala	Thr	Asp	Gly	Ser	Leu		
			125					130					135				
gcc	tcc	att	ctc	tcc	agc	cta	tct	cat	gat	gag	aaa	aag	ggg	ggg	tct	665	
Ala	Ser	Ile	Leu	Ser	Ser	Leu	Ser	His	Asp	Glu	Lys	Lys	Gly	Gly	Ser		
		140					145					150					
tgg	cta	aat	ccc	atc	cag	tcc	cga	aac	acg	aca	gtg	gag	tat	gtg	aca	713	
Trp	Leu	Asn	Pro	Ile	Gln	Ser	Arg	Asn	Thr	Thr	Val	Glu	Tyr	Val	Thr		
		155				160					165						
ttc	acc	agc	gtt	gct	ggg	ctc	act	gga	gtg	atc	atg	aca	ata	gcc	ttg	761	
Phe	Thr	Ser	Val	Ala	Gly	Leu	Thr	Gly	Val	Ile	Met	Thr	Ile	Ala	Leu		
					175					180					185		
att	ctc	atg	gta	act	tca	gct	act	gag	ttc	atc	cgg	agg	agt	tat	ttt	809	
Ile	Leu	Met	Val	Thr	Ser	Ala	Thr	Glu	Phe	Ile	Arg	Arg	Ser	Tyr	Phe		
				190					195					200			
gaa	gtc	ttc	tgg	tat	act	cac	cac	ctt	ttt	atc	ttc	tat	atc	ctt	ggc	857	
Glu	Val	Phe	Trp	Tyr	Thr	His	His	Leu	Phe	Ile	Phe	Tyr	Ile	Leu	Gly		
			205					210					215				
tta	ggg	att	cac	ggc	att	ggg	gga	att	gtc	cgg	ggg	caa	aca	gag	gag	905	
Leu	Gly	Ile	His	Gly	Ile	Gly	Gly	Ile	Val	Arg	Gly	Gln	Thr	Glu	Glu		
			220				225					230					
agc	atg	aat	gag	agt	cat	cct	cgc	aag	tgt	gca	gag	tct	ttt	gag	atg	953	
Ser	Met	Asn	Glu	Ser	His	Pro	Arg	Lys	Cys	Ala	Glu	Ser	Phe	Glu	Met		
			235			240					245						
tgg	gat	gat	cgt	gac	tcc	cac	tgt	agg	cgc	cct	aag	ttt	gaa	ggg	cat	1001	

WO 00/28031

PCT/US99/26592

Trp Asp Asp Arg Asp Ser His Cys Arg Arg Pro Lys Phe Glu Gly His	
250 255 260 265	
ccc cct gag tct tgg aag tgg atc ctt gca ccg gtc att ctt tat atc	1049
Pro Pro Glu Ser Trp Lys Trp Ile Leu Ala Pro Val Ile Leu Tyr Ile	
270 275 280	
tgt gaa agg atc ctc cgg ttt tac cgc tcc cag cag aag gtt gtg att	1097
Cys Glu Arg Ile Leu Arg Phe Tyr Arg Ser Gln Gln Lys Val Val Ile	
285 290 295	
acc aag gtt gtt atg cac cca tcc aaa gtt ttg gaa ttg cag atg aac	1145
Thr Lys Val Val Met His Pro Ser Lys Val Leu Glu Leu Gln Met Asn	
300 305 310	
aag cgt ggc ttc agc atg gaa gtg ggg cag tat atc ttt gtt aat tgc	1193
Lys Arg Gly Phe Ser Met Glu Val Gly Gln Tyr Ile Phe Val Asn Cys	
315 320 325	
ccc tca atc tct ctc ctg gaa tgg cat cct ttt act ttg acc tct gct	1241
Pro Ser Ile Ser Leu Leu Glu Trp His Pro Phe Thr Leu Thr Ser Ala	
330 335 340 345	
cca gag gaa gat ttc ttc tcc att cat atc cga gca gca ggg gac tgg	1289
Pro Glu Glu Asp Phe Phe Ser Ile His Ile Arg Ala Ala Gly Asp Trp	
350 355 360	
aca gaa aat ctc ata agg gct ttc gaa caa caa tat tca cca att ccc	1337
Thr Glu Asn Leu Ile Arg Ala Phe Glu Gln Gln Tyr Ser Pro Ile Pro	
365 370 375	
agg att gaa gtg gat ggt ccc ttt ggc aca gcc agt gag gat gtt ttc	1385
Arg Ile Glu Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe	
380 385 390	
cag tat gaa gtg gct gtg ctg gtt gga gca gga att ggg gtc acc ccc	1433
Gln Tyr Glu Val Ala Val Leu Val Gly Ala Gly Ile Gly Val Thr Pro	
395 400 405	
ttt gct tct atc ttg aaa tcc atc tgg tac aaa ttc cag tgt gca gac	1481
Phe Ala Ser Ile Leu Lys Ser Ile Trp Tyr Lys Phe Gln Cys Ala Asp	
410 415 420 425	
cac aac ctc aaa aca aaa aag atc tat ttc tac tgg atc tgc agg gag	1529
His Asn Leu Lys Thr Lys Lys Ile Tyr Phe Tyr Trp Ile Cys Arg Glu	
430 435 440	
aca ggt gcc ttt tcc tgg ttc aac aac ctg ttg act tcc ctg gaa cag	1577

WO 00/28031

PCT/US99/26592

Thr Gly Ala Phe Ser Trp Phe Asn Asn Leu Leu Thr Ser Leu Glu Gln
 445 450 455

gag atg gag gaa tta ggc aaa gtg ggt ttt cta aac tac cgt ctc ttc 1625
 Glu Met Glu Glu Leu Gly Lys Val Gly Phe Leu Asn Tyr Arg Leu Phe
 460 465 470

ctc acc gga tgg gac agc aat att gtt ggt cat gca gca tta aac ttt 1673
 Leu Thr Gly Trp Asp Ser Asn Ile Val Gly His Ala Ala Leu Asn Phe
 475 480 485

gac aag gcc act gac atc gtg aca ggt ctg aaa cag aaa acc tcc ttt 1721
 Asp Lys Ala Thr Asp Ile Val Thr Gly Leu Lys Gln Lys Thr Ser Phe
 490 495 500 505

ggg aga cca atg tgg gac aat gag ttt tct aca ata gct acc tcc cac 1769
 Gly Arg Pro Met Trp Asp Asn Glu Phe Ser Thr Ile Ala Thr Ser His
 510 515 520

ccc aag tct gta gtg gga gtt ttc tta tgt ggc cct cgg act ttg gca 1817
 Pro Lys Ser Val Val Gly Val Phe Leu Cys Gly Pro Arg Thr Leu Ala
 525 530 535

aag agc ctg cgc aaa tgc tgt cac cga tat tcc agt ctg gat cct aga 1865
 Lys Ser Leu Arg Lys Cys Cys His Arg Tyr Ser Ser Leu Asp Pro Arg
 540 545 550

aag gtt caa ttc tac ttc aac aaa gaa aat ttt tga gttataggaa 1911
 Lys Val Gln Phe Tyr Phe Asn Lys Glu Asn Phe
 555 560 565

taaggacggt aatctgcatt ttgtctcttt gtatcttcag taattgagtt ataggaataa 1971

ggacggtaat ctgcattttg tctctttgta tcttcagtaa tttacttggt ctcttcaggt 2031

ttgancagtc actttaggat aagaatgtgc ctctcaagcc ttgactccct ggtattcttt 2091

ttttgattgc attcaacttc gttacttgag cttcagcaac ttaagaactt ctgaagttct 2151

taaagttctg aanttcttaa agcccatgga tcctttctca gaaaaataac tgtaaacttt 2211

tctggacagc catgactgta gcaaggcttg atagcagaag tttggtggtt canaattata 2271

caactaatcc caggtgattt tatcaattcc agtggtacca tctcctgagt tttggtttgt 2331

aatcttttgt cctcccacc cccacagaag attttaagta ggggtgacttt ttaaataaaa 2391

atttattgaa taattaatga taaaacataa taataaacat aaataataaa caaaattacc 2451

WO 00/28031

PCT/US99/26592

gagaacccca tcccatata acaccaacag tgtacatgtt tactgtcact tttgatatgg 2511

tttatccagt gtgaacagca atttattatt tttgctcatc aaaaaataaa ggattttttt 2571

tcacttgaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2609

<210> 2

<211> 564

<212> PRT

<213> Homo sapiens

<400> 2

Met Gly Asn Trp Val Val Asn His Trp Phe Ser Val Leu Phe Leu Val
1 5 10 15

Val Trp Leu Gly Leu Asn Val Phe Leu Phe Val Asp Ala Phe Leu Lys
20 25 30

Tyr Glu Lys Ala Asp Lys Tyr Tyr Tyr Thr Arg Lys Ile Leu Gly Ser
35 40 45

Thr Leu Ala Cys Ala Arg Ala Ser Ala Leu Cys Leu Asn Phe Asn Ser
50 55 60

Thr Leu Ile Leu Leu Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg
65 70 75 80

Gly Thr Cys Ser Phe Cys Ser Arg Thr Leu Arg Lys Gln Leu Asp His
85 90 95

Asn Leu Thr Phe His Lys Leu Val Ala Tyr Met Ile Cys Leu His Thr
100 105 110

Ala Ile His Ile Ile Ala His Leu Phe Asn Phe Asp Cys Tyr Ser Arg
115 120 125

Ser Arg Gln Ala Thr Asp Gly Ser Leu Ala Ser Ile Leu Ser Ser Leu
130 135 140

Ser His Asp Glu Lys Lys Gly Gly Ser Trp Leu Asn Pro Ile Gln Ser
145 150 155 160

Arg Asn Thr Thr Val Glu Tyr Val Thr Phe Thr Ser Val Ala Gly Leu
165 170 175

Thr Gly Val Ile Met Thr Ile Ala Leu Ile Leu Met Val Thr Ser Ala

WO 00/28031

PCT/US99/26592

180					185					190						
Thr	Glu	Phe	Ile	Arg	Arg	Ser	Tyr	Phe	Glu	Val	Phe	Trp	Tyr	Thr	His	
195					200					205						
His	Leu	Phe	Ile	Phe	Tyr	Ile	Leu	Gly	Leu	Gly	Ile	His	Gly	Ile	Gly	
210					215					220						
Gly	Ile	Val	Arg	Gly	Gln	Thr	Glu	Glu	Ser	Met	Asn	Glu	Ser	His	Pro	
225					230					235					240	
Arg	Lys	Cys	Ala	Glu	Ser	Phe	Glu	Met	Trp	Asp	Asp	Arg	Asp	Ser	His	
245					250					255						
Cys	Arg	Arg	Pro	Lys	Phe	Glu	Gly	His	Pro	Pro	Glu	Ser	Trp	Lys	Trp	
260					265					270						
Ile	Leu	Ala	Pro	Val	Ile	Leu	Tyr	Ile	Cys	Glu	Arg	Ile	Leu	Arg	Phe	
275					280					285						
Tyr	Arg	Ser	Gln	Gln	Lys	Val	Val	Ile	Thr	Lys	Val	Val	Met	His	Pro	
290					295					300						
Ser	Lys	Val	Leu	Glu	Leu	Gln	Met	Asn	Lys	Arg	Gly	Phe	Ser	Met	Glu	
305					310					315					320	
Val	Gly	Gln	Tyr	Ile	Phe	Val	Asn	Cys	Pro	Ser	Ile	Ser	Leu	Leu	Glu	
325					330					335						
Trp	His	Pro	Phe	Thr	Leu	Thr	Ser	Ala	Pro	Glu	Glu	Asp	Phe	Phe	Ser	
340					345					350						
Ile	His	Ile	Arg	Ala	Ala	Gly	Asp	Trp	Thr	Glu	Asn	Leu	Ile	Arg	Ala	
355					360					365						
Phe	Glu	Gln	Gln	Tyr	Ser	Pro	Ile	Pro	Arg	Ile	Glu	Val	Asp	Gly	Pro	
370					375					380						
Phe	Gly	Thr	Ala	Ser	Glu	Asp	Val	Phe	Gln	Tyr	Glu	Val	Ala	Val	Leu	
385					390					395					400	
Val	Gly	Ala	Gly	Ile	Gly	Val	Thr	Pro	Phe	Ala	Ser	Ile	Leu	Lys	Ser	
405					410					415						
Ile	Trp	Tyr	Lys	Phe	Gln	Cys	Ala	Asp	His	Asn	Leu	Lys	Thr	Lys	Lys	
420					425					430						
Ile	Tyr	Phe	Tyr	Trp	Ile	Cys	Arg	Glu	Thr	Gly	Ala	Phe	Ser	Trp	Phe	

WO 00/28031

PCT/US99/26592

435	440	445
Asn Asn Leu Leu Thr Ser Leu Glu Gln Glu Met Glu Glu Leu Gly Lys 450	455	460
Val Gly Phe Leu Asn Tyr Arg Leu Phe Leu Thr Gly Trp Asp Ser Asn 465	470	475 480
Ile Val Gly His Ala Ala Leu Asn Phe Asp Lys Ala Thr Asp Ile Val 485	490	495
Thr Gly Leu Lys Gln Lys Thr Ser Phe Gly Arg Pro Met Trp Asp Asn 500	505	510
Glu Phe Ser Thr Ile Ala Thr Ser His Pro Lys Ser Val Val Gly Val 515	520	525
Phe Leu Cys Gly Pro Arg Thr Leu Ala Lys Ser Leu Arg Lys Cys Cys 530	535	540
His Arg Tyr Ser Ser Leu Asp Pro Arg Lys Val Gln Phe Tyr Phe Asn 545	550	555 560
Lys Glu Asn Phe		

<210> 3

<211> 2044

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (104)..(1810)

<400> 3

caaagacaaa ataatttact aggggaagccc ttactaacga cccaacatcc agacacaggt 60

gagggagaag aaatttcctg acagccgaag agcaacaagt atc atg atg ggg tgc 115
Met Met Gly Cys

1

tgg att ttg aat gag ggt ctc tcc acc ata tta gta ctc tca tgg ctg 163
Trp Ile Leu Asn Glu Gly Leu Ser Thr Ile Leu Val Leu Ser Trp Leu
5 10 15 20

gga ata aat ttt tat ctg ttt att gac acg ttc tac tgg tat gaa gag 211
Gly Ile Asn Phe Tyr Leu Phe Ile Asp Thr Phe Tyr Trp Tyr Glu Glu

WO 00/28031

PCT/US99/26592

25	30	35	
gag gag tct ttc cat tac aca cga gtt att ttg ggt tca aca ctg gct Glu Glu Ser Phe His Tyr Thr Arg Val Ile Leu Gly Ser Thr Leu Ala			259
40	45	50	
tgg gca cga gca tcc gca ctg tgc ctg aat ttt aac tgc atg cta att Trp Ala Arg Ala Ser Ala Leu Cys Leu Asn Phe Asn Cys Met Leu Ile			307
55	60	65	
cta ata cct gtc agt cga aac ctt att tca ttc ata aga gga aca agt Leu Ile Pro Val Ser Arg Asn Leu Ile Ser Phe Ile Arg Gly Thr Ser			355
70	75	80	
att tgc tgc aga gga ccg tgg agg agg caa tta gac aaa aac ctc aga Ile Cys Cys Arg Gly Pro Trp Arg Arg Gln Leu Asp Lys Asn Leu Arg			403
85	90	95	100
ttt cac aaa ctg gtc gcc tat ggg ata gct gtt aat gca acc atc cac Phe His Lys Leu Val Ala Tyr Gly Ile Ala Val Asn Ala Thr Ile His			451
105	110	115	
atc gtg gcg cat ttc ttc aac ctg gaa cgc tac cac tgg agc cag tcc Ile Val Ala His Phe Phe Asn Leu Glu Arg Tyr His Trp Ser Gln Ser			499
120	125	130	
gag gag gcc cag gga ctt ctg gcc gca ctt tcc aag ctg ggc aac acc Glu Glu Ala Gln Gly Leu Leu Ala Ala Leu Ser Lys Leu Gly Asn Thr			547
135	140	145	
cct aac gag agc tac ctc aac cct gtc cgg acc ttc ccc aca aac aca Pro Asn Glu Ser Tyr Leu Asn Pro Val Arg Thr Phe Pro Thr Asn Thr			595
150	155	160	
acc act gaa ttg cta agg aca ata gca ggc gtc acc ggt ctg gtg atc Thr Thr Glu Leu Leu Arg Thr Ile Ala Gly Val Thr Gly Leu Val Ile			643
165	170	175	180
tct ctg gct tta gtc ttg atc atg acc tcg tca act gag ttc atc aga Ser Leu Ala Leu Val Leu Ile Met Thr Ser Ser Thr Glu Phe Ile Arg			691
185	190	195	
cag gcc tcc tat gag ttg ttc tgg tac aca cac cat gtt ttc atc gtc Gln Ala Ser Tyr Glu Leu Phe Trp Tyr Thr His His Val Phe Ile Val			739
200	205	210	
ttc ttt ctc agc ctg gcc atc cat ggg acg ggt cgg att gtt cga ggc Phe Phe Leu Ser Leu Ala Ile His Gly Thr Gly Arg Ile Val Arg Gly			787

WO 00/28031

PCT/US99/26592

215	220	225	
caa acc caa gac agt ctc tct ctg cac aac atc acc ttc tgt aga gac			835
Gln Thr Gln Asp Ser Leu Ser Leu His Asn Ile Thr Phe Cys Arg Asp			
230	235	240	
cgc tat gca gaa tgg cag aca gtg gcc caa tgc ccc gtg cct caa ttt			883
Arg Tyr Ala Glu Trp Gln Thr Val Ala Gln Cys Pro Val Pro Gln Phe			
245	250	255	260
tct ggc aag gaa ccc tcg gct tgg aaa tgg att tta ggc cct gtg gtc			931
Ser Gly Lys Glu Pro Ser Ala Trp Lys Trp Ile Leu Gly Pro Val Val			
265	270	275	
ttg tat gca tgt gaa aga ata att agg ttc tgg cga ttt caa caa gaa			979
Leu Tyr Ala Cys Glu Arg Ile Ile Arg Phe Trp Arg Phe Gln Gln Glu			
280	285	290	
gtt gtc att acc aag gtg gta agc cac ccc tct gga gtc ctg gaa ctt			1027
Val Val Ile Thr Lys Val Val Ser His Pro Ser Gly Val Leu Glu Leu			
295	300	305	
cac atg aaa aag cgt ggc ttt aaa atg gcg cca ggg cag tac atc ttg			1075
His Met Lys Lys Arg Gly Phe Lys Met Ala Pro Gly Gln Tyr Ile Leu			
310	315	320	
gtg cag tgc cca gcc ata tct tcg ctg gag tgg cac ccc ttc acc ctt			1123
Val Gln Cys Pro Ala Ile Ser Ser Leu Glu Trp His Pro Phe Thr Leu			
325	330	335	340
acc tct gcc ccc cag gaa gac ttt ttc agc gtg cac atc cgg gca gca			1171
Thr Ser Ala Pro Gln Glu Asp Phe Phe Ser Val His Ile Arg Ala Ala			
345	350	355	
gga gac tgg aca gca gcg cta ctg gag gcc ttt ggg gca gag gga cag			1219
Gly Asp Trp Thr Ala Ala Leu Leu Glu Ala Phe Gly Ala Glu Gly Gln			
360	365	370	
gcc ctc cag gag ccc tgg agc ctg cca agg ctg gca gtg gac ggg ccc			1267
Ala Leu Gln Glu Pro Trp Ser Leu Pro Arg Leu Ala Val Asp Gly Pro			
375	380	385	
ttt gga act gcc ctg aca gat gta ttt cac tac cca gtg tgt gtg tgc			1315
Phe Gly Thr Ala Leu Thr Asp Val Phe His Tyr Pro Val Cys Val Cys			
390	395	400	
gtt gcc gcg ggg atc gga gtc act ccc ttc gct gct ctt ctg aaa tct			1363
Val Ala Ala Gly Ile Gly Val Thr Pro Phe Ala Ala Leu Leu Lys Ser			

WO 00/28031

PCT/US99/26592

405	410	415	420	
ata tgg tac aaa tgc agt gag gca cag acc cca ctg aag ctg agc aag				1411
Ile Trp Tyr Lys Cys Ser Glu Ala Gln Thr Pro Leu Lys Leu Ser Lys				
	425	430	435	
gtg tat ttc tac tgg att tgc cgg gat gca aga gct ttt gag tgg ttt				1459
Val Tyr Phe Tyr Trp Ile Cys Arg Asp Ala Arg Ala Phe Glu Trp Phe				
	440	445	450	
gct gat ctc tta ctc tcc ctg gaa aca cgg atg agt gag cag ggg aaa				1507
Ala Asp Leu Leu Leu Ser Leu Glu Thr Arg Met Ser Glu Gln Gly Lys				
	455	460	465	
act cac ttt ctg agt tat cat ata ttt ctt acc ggc tgg gat gaa aat				1555
Thr His Phe Leu Ser Tyr His Ile Phe Leu Thr Gly Trp Asp Glu Asn				
	470	475	480	
cag gct ctt cac ata gct tta cac tgg gac gaa aat act gac gtg att				1603
Gln Ala Leu His Ile Ala Leu His Trp Asp Glu Asn Thr Asp Val Ile				
	485	490	500	
aca ggc tta aag cag aag acc ttc tat ggg agg ccc aac tgg aac aat				1651
Thr Gly Leu Lys Gln Lys Thr Phe Tyr Gly Arg Pro Asn Trp Asn Asn				
	505	510	515	
gag ttc aag cag att gcc tac aat cac ccc agc agc agt att ggc gtg				1699
Glu Phe Lys Gln Ile Ala Tyr Asn His Pro Ser Ser Ser Ile Gly Val				
	520	525	530	
ttc ttc tgt gga cct aaa gct ctc tcg agg aca ctt caa aag atg tgc				1747
Phe Phe Cys Gly Pro Lys Ala Leu Ser Arg Thr Leu Gln Lys Met Cys				
	535	540	545	
cac ttg tat tca tca gct gac ccc aga ggt gtt cat ttc tat tac aac				1795
His Leu Tyr Ser Ser Ala Asp Pro Arg Gly Val His Phe Tyr Tyr Asn				
	550	555	560	
aag gag agc ttc tag actttggagg tcaagtccag gcattgtgtt ttcaatcaag				1850
Lys Glu Ser Phe				
	565			
ttattgattc caaagaactc caccaggaat tcctgtgacg gcctgttgat atgagctccc				1910
agttgggaac tgggtgaataa taattaacta ttgtgaacag tacactatac catacttcct				1970
tagcttataa ataacatgtc atatacaaca gaacaaaaac atttactgaa attaaaatat				2030

WO 00/28031

PCT/US99/26592

attatgtttc tcca

2044

<210> 4

<211> 568

<212> PRT

<213> Homo sapiens

<400> 4

Met	Met	Gly	Cys	Trp	Ile	Leu	Asn	Glu	Gly	Leu	Ser	Thr	Ile	Leu	Val
1				5					10					15	

Leu	Ser	Trp	Leu	Gly	Ile	Asn	Phe	Tyr	Leu	Phe	Ile	Asp	Thr	Phe	Tyr
			20					25					30		

Trp	Tyr	Glu	Glu	Glu	Glu	Ser	Phe	His	Tyr	Thr	Arg	Val	Ile	Leu	Gly
		35					40					45			

Ser	Thr	Leu	Ala	Trp	Ala	Arg	Ala	Ser	Ala	Leu	Cys	Leu	Asn	Phe	Asn
	50					55					60				

Cys	Met	Leu	Ile	Leu	Ile	Pro	Val	Ser	Arg	Asn	Leu	Ile	Ser	Phe	Ile
65					70					75					80

Arg	Gly	Thr	Ser	Ile	Cys	Cys	Arg	Gly	Pro	Trp	Arg	Arg	Gln	Leu	Asp
				85					90					95	

Lys	Asn	Leu	Arg	Phe	His	Lys	Leu	Val	Ala	Tyr	Gly	Ile	Ala	Val	Asn
		100						105					110		

Ala	Thr	Ile	His	Ile	Val	Ala	His	Phe	Phe	Asn	Leu	Glu	Arg	Tyr	His
		115					120					125			

Trp	Ser	Gln	Ser	Glu	Glu	Ala	Gln	Gly	Leu	Leu	Ala	Ala	Leu	Ser	Lys
	130					135					140				

Leu	Gly	Asn	Thr	Pro	Asn	Glu	Ser	Tyr	Leu	Asn	Pro	Val	Arg	Thr	Phe
145					150					155					160

Pro	Thr	Asn	Thr	Thr	Thr	Glu	Leu	Leu	Arg	Thr	Ile	Ala	Gly	Val	Thr
			165						170					175	

Gly	Leu	Val	Ile	Ser	Leu	Ala	Leu	Val	Leu	Ile	Met	Thr	Ser	Ser	Thr
		180						185						190	

Glu	Phe	Ile	Arg	Gln	Ala	Ser	Tyr	Glu	Leu	Phe	Trp	Tyr	Thr	His	His
		195					200					205			

WO 00/28031

PCT/US99/26592

Val Phe Ile Val Phe Phe Leu Ser Leu Ala Ile His Gly Thr Gly Arg
 210 215 220

Ile Val Arg Gly Gln Thr Gln Asp Ser Leu Ser Leu His Asn Ile Thr
 225 230 235 240

Phe Cys Arg Asp Arg Tyr Ala Glu Trp Gln Thr Val Ala Gln Cys Pro
 245 250 255

Val Pro Gln Phe Ser Gly Lys Glu Pro Ser Ala Trp Lys Trp Ile Leu
 260 265 270

Gly Pro Val Val Leu Tyr Ala Cys Glu Arg Ile Ile Arg Phe Trp Arg
 275 280 285

Phe Gln Gln Glu Val Val Ile Thr Lys Val Val Ser His Pro Ser Gly
 290 295 300

Val Leu Glu Leu His Met Lys Lys Arg Gly Phe Lys Met Ala Pro Gly
 305 310 315 320

Gln Tyr Ile Leu Val Gln Cys Pro Ala Ile Ser Ser Leu Glu Trp His
 325 330 335

Pro Phe Thr Leu Thr Ser Ala Pro Gln Glu Asp Phe Phe Ser Val His
 340 345 350

Ile Arg Ala Ala Gly Asp Trp Thr Ala Ala Leu Leu Glu Ala Phe Gly
 355 360 365

Ala Glu Gly Gln Ala Leu Gln Glu Pro Trp Ser Leu Pro Arg Leu Ala
 370 375 380

Val Asp Gly Pro Phe Gly Thr Ala Leu Thr Asp Val Phe His Tyr Pro
 385 390 395 400

Val Cys Val Cys Val Ala Ala Gly Ile Gly Val Thr Pro Phe Ala Ala
 405 410 415

Leu Leu Lys Ser Ile Trp Tyr Lys Cys Ser Glu Ala Gln Thr Pro Leu
 420 425 430

Lys Leu Ser Lys Val Tyr Phe Tyr Trp Ile Cys Arg Asp Ala Arg Ala
 435 440 445

Phe Glu Trp Phe Ala Asp Leu Leu Leu Ser Leu Glu Thr Arg Met Ser
 450 455 460

WO 00/28031

PCT/US99/26592

Glu Gln Gly Lys Thr His Phe Leu Ser Tyr His Ile Phe Leu Thr Gly
 465 470 475 480

Trp Asp Glu Asn Gln Ala Leu His Ile Ala Leu His Trp Asp Glu Asn
 485 490 495

Thr Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Phe Tyr Gly Arg Pro
 500 505 510

Asn Trp Asn Asn Glu Phe Lys Gln Ile Ala Tyr Asn His Pro Ser Ser
 515 520 525

Ser Ile Gly Val Phe Phe Cys Gly Pro Lys Ala Leu Ser Arg Thr Leu
 530 535 540

Gln Lys Met Cys His Leu Tyr Ser Ser Ala Asp Pro Arg Gly Val His
 545 550 555 560

Phe Tyr Tyr Asn Lys Glu Ser Phe
 565

<210> 5
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 5
 aacaagcgtg gcttcagcat g

21

<210> 6
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 6
 agcaatattg ttgggtcat

18

<210> 7
 <211> 24

WO 00/28031

PCT/US99/26592

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 7

gacttgacag aaaatctata aggg

24

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 8

ttgtaccaga tggatttcaa

20

<210> 9

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 9

caggtctgaa acagaaaacc t

21

<210> 10

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 10

atgaattctc attaattatt caataaa

27

<210> 11

<211> 20

WO 00/28031

PCT/US99/26592

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 11

atctcaaaag actctgcaca

20

<210> 12

<211> 569

<212> PRT

<213> Homo sapiens

<400> 12

Gly	Asn	Trp	Ala	Val	Asn	Glu	Gly	Leu	Ser	Ile	Phe	Ala	Ile	Leu	Val
1				5					10					15	

Trp	Leu	Gly	Leu	Asn	Val	Phe	Leu	Phe	Val	Trp	Tyr	Tyr	Arg	Val	Tyr
			20					25					30		

Asp	Ile	Pro	Pro	Lys	Phe	Phe	Tyr	Thr	Arg	Lys	Leu	Leu	Gly	Ser	Ala
		35					40					45			

Leu	Ala	Leu	Ala	Arg	Ala	Pro	Ala	Ala	Cys	Leu	Asn	Phe	Asn	Cys	Met
	50					55					60				

Leu	Ile	Leu	Leu	Pro	Val	Cys	Arg	Asn	Leu	Leu	Ser	Phe	Leu	Arg	Gly
65					70					75					80

Ser	Ser	Ala	Cys	Cys	Ser	Thr	Arg	Val	Arg	Arg	Gln	Leu	Asp	Arg	Asn
			85						90					95	

Leu	Thr	Phe	His	Lys	Met	Val	Ala	Trp	Met	Ile	Ala	Leu	His	Ser	Ala
			100					105					110		

Ile	His	Thr	Ile	Ala	His	Leu	Phe	Asn	Val	Glu	Trp	Cys	Val	Asn	Ala
		115					120					125			

Arg	Val	Asn	Asn	Ser	Asp	Pro	Tyr	Ser	Val	Ala	Leu	Ser	Glu	Leu	Gly
	130					135					140				

Asp	Arg	Gln	Asn	Glu	Ser	Tyr	Leu	Asn	Phe	Ala	Arg	Lys	Arg	Ile	Lys
145					150					155					160

Asn	Pro	Glu	Gly	Gly	Leu	Tyr	Leu	Ala	Val	Thr	Leu	Leu	Ala	Gly	Ile
			165						170					175	

WO 00/28031

PCT/US99/26592

Thr Gly Val Val Ile Thr Leu Cys Leu Ile Leu Ile Ile Thr Ser Ser
 180 185 190

Thr Lys Thr Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr His
 195 200 205

His Leu Phe Val Ile Phe Phe Ile Gly Leu Ala Ile His Gly Ala Glu
 210 215 220

Arg Ile Val Arg Gly Gln Thr Ala Glu Ser Leu Ala Val His Asn Ile
 225 230 235 240

Thr Val Cys Glu Gln Lys Ile Ser Glu Trp Gly Lys Ile Lys Glu Cys
 245 250 255

Pro Ile Pro Gln Phe Ala Gly Asn Pro Pro Met Thr Trp Lys Trp Ile
 260 265 270

Val Gly Pro Met Phe Leu Tyr Leu Cys Glu Arg Leu Val Arg Phe Trp
 275 280 285

Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Thr His Pro Phe
 290 295 300

Lys Thr Ile Glu Leu Gln Met Lys Lys Lys Gly Phe Lys Met Glu Val
 305 310 315 320

Gly Gln Tyr Ile Phe Val Lys Cys Pro Lys Val Ser Lys Leu Glu Trp
 325 330 335

His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser Ile
 340 345 350

His Ile Arg Ile Val Gly Asp Trp Thr Glu Gly Leu Phe Asn Ala Cys
 355 360 365

Gly Cys Asp Lys Gln Glu Phe Gln Asp Ala Trp Lys Leu Pro Lys Ile
 370 375 380

Ala Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe Ser Tyr
 385 390 395 400

Glu Val Val Met Leu Val Gly Ala Gly Ile Gly Val Thr Pro Phe Ala
 405 410 415

Ser Ile Leu Lys Ser Val Trp Tyr Lys Tyr Cys Asn Asn Ala Thr Asn
 420 425 430

WO 00/28031

PCT/US99/26592

Leu Lys Leu Lys Lys Ile Tyr Phe Tyr Trp Leu Cys Arg Asp Thr His
 435 440 445
 Ala Phe Glu Trp Phe Ala Asp Leu Leu Gln Leu Leu Glu Ser Gln Met
 450 455 460
 Gln Glu Arg Asn Asn Ala Gly Phe Leu Ser Tyr Asn Ile Tyr Leu Thr
 465 470 475 480
 Gly Trp Asp Glu Ser Gln Ala Asn His Phe Ala Val His His Asp Glu
 485 490 495
 Glu Lys Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Leu Tyr Gly Arg
 500 505 510
 Pro Asn Trp Asp Asn Glu Phe Lys Thr Ile Ala Ser Gln His Pro Asn
 515 520 525
 Thr Arg Ile Gly Val Phe Leu Cys Gly Pro Glu Ala Leu Ala Glu Thr
 530 535 540
 Leu Ser Lys Gln Ser Ile Ser Asn Ser Glu Ser Gly Pro Arg Gly Val
 545 550 555 560
 His Phe Ile Phe Asn Lys Glu Asn Phe
 565

<210> 13
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 13
 ttggctaaat cccatcca

18

<210> 14
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

WO 00/28031

PCT/US99/26592

<400> 14

tgcattgacca acaatattgc t

21

<210> 15

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 15

caagggtacct cttgacctg ggaaact

27

<210> 16

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 16

acgaattcaa gtaaattact gaagata

27

<210> 17

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<221> modified_base

<222> ()..)

<223> n at position 3 = inosine

<220>

<221> modified_base

<222> ()

<223> n at position 6 = inosine

<220>

<221> modified_base

<222> ()

<223> n at position 12 = inosine

WO 00/28031

PCT/US99/26592

<220>

<223> Description of Artificial Sequence: primer

<400> 17

ccngtntgtc gnaatctgct stcctt

26

<210> 18

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<221> modified_base

<222> (5)

<223> n at position 5 = inosine

<220>

<221> modified_base

<222> (9)

<223> n at position 9 = inosine

<220>

<221> modified_base

<222> (11)

<223> n at position 11 = inosine

<220>

<223> Description of Artificial Sequence: primer

<400> 18

tcccngcana nccagtagaa rtagatctt

29

<210> 19

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 19

ttggcacagt cagtgaggat gtcttc

26

<210> 20

WO 00/28031

PCT/US99/26592

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 20

ctgttggtt ctactgtagc gttcaaagtt

30

<210> 21

<211> 563

<212> PRT

<213> Rat

<400> 21

Met	Gly	Asn	Trp	Leu	Val	Asn	His	Trp	Leu	Ser	Val	Leu	Phe	Leu	Val
1				5					10					15	

Ser	Trp	Leu	Gly	Leu	Asn	Ile	Phe	Leu	Phe	Val	Tyr	Val	Phe	Leu	Asn
		20						25					30		

Tyr	Glu	Lys	Ser	Asp	Lys	Tyr	Tyr	Tyr	Thr	Arg	Glu	Ile	Leu	Gly	Thr
		35					40					45			

Ala	Leu	Ala	Leu	Ala	Arg	Ala	Ser	Ala	Leu	Cys	Leu	Asn	Phe	Asn	Ser
	50					55					60				

Met	Val	Ile	Leu	Ile	Pro	Val	Cys	Arg	Asn	Leu	Leu	Ser	Phe	Leu	Arg
65					70					75					80

Gly	Thr	Cys	Ser	Phe	Cys	Asn	His	Thr	Leu	Arg	Lys	Pro	Leu	Asp	His
			85						90					95	

Asn	Leu	Thr	Phe	His	Lys	Leu	Val	Ala	Tyr	Met	Ile	Cys	Ile	Phe	Thr
		100						105					110		

Ala	Ile	His	Ile	Ile	Ala	His	Leu	Phe	Asn	Phe	Glu	Arg	Tyr	Ser	Arg
		115					120					125			

Ser	Gln	Gln	Ala	Met	Asp	Gly	Ser	Leu	Ala	Ser	Val	Leu	Ser	Ser	Leu
	130					135					140				

Phe	His	Pro	Glu	Lys	Glu	Asp	Ser	Trp	Leu	Asn	Pro	Ile	Gln	Ser	Pro
145				150						155					160

Asn	Val	Thr	Val	Met	Tyr	Ala	Ala	Phe	Thr	Ser	Ile	Ala	Gly	Leu	Thr
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

WO 00/28031

PCT/US99/26592

165	170	175
Gly Val Val Ala Thr Val Ala Leu Val Leu Met Val Thr Ser Ala Met 180	185	190
Glu Phe Ile Arg Arg Asn Tyr Phe Glu Leu Phe Trp Tyr Thr His His 195	200	205
Leu Phe Ile Ile Tyr Ile Ile Cys Leu Gly Ile His Gly Leu Gly Gly 210	215	220
Ile Val Arg Gly Gln Thr Glu Glu Ser Met Ser Glu Ser His Pro Arg 225	230	235 240
Asn Cys Ser Tyr Ser Phe His Glu Trp Asp Lys Tyr Glu Arg Ser Cys 245	250	255
Arg Ser Pro His Phe Val Gly Gln Pro Pro Glu Ser Trp Lys Trp Ile 260	265	270
Leu Ala Pro Ile Ala Phe Tyr Ile Phe Glu Arg Ile Leu Arg Phe Tyr 275	280	285
Arg Ser Arg Gln Lys Val Val Ile Thr Lys Val Val Met His Pro Cys 290	295	300
Lys Val Leu Glu Leu Gln Met Arg Lys Arg Gly Phe Thr Met Gly Ile 305	310	315 320
Gly Gln Tyr Ile Phe Val Asn Cys Pro Ser Ile Ser Phe Leu Glu Trp 325	330	335
His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Glu Phe Phe Ser Ile 340	345	350
His Ile Arg Ala Ala Gly Asp Trp Thr Glu Asn Leu Ile Arg Thr Phe 355	360	365
Glu Gln Gln His Ser Pro Met Pro Arg Ile Glu Val Asp Gly Pro Phe 370	375	380
Gly Thr Val Ser Glu Asp Val Phe Gln Tyr Glu Val Ala Val Leu Val 385	390	395 400
Gly Ala Gly Ile Gly Val Thr Pro Phe Ala Ser Phe Leu Lys Ser Ile 405	410	415
Trp Tyr Lys Phe Gln Arg Ala His Asn Lys Leu Lys Thr Gln Lys Ile		

WO 00/28031

PCT/US99/26592

420	425	430
Tyr Phe Tyr Trp Ile Cys Arg Glu Thr Gly Ala Phe Ala Trp Phe Asn 435 440 445		
Asn Leu Leu Asn Ser Leu Glu Gln Glu Met Asp Glu Leu Gly Lys Pro 450 455 460		
Asp Phe Leu Asn Tyr Arg Leu Phe Leu Thr Gly Trp Asp Ser Asn Ile 465 470 475 480		
Ala Gly His Ala Ala Leu Asn Phe Asp Arg Ala Thr Asp Val Leu Thr 485 490 495		
Gly Leu Lys Gln Lys Thr Ser Phe Gly Arg Pro Met Trp Asp Asn Glu 500 505 510		
Phe Ser Arg Ile Ala Thr Ala His Pro Lys Ser Val Val Gly Val Phe 515 520 525		
Leu Cys Gly Pro Pro Thr Leu Ala Lys Ser Leu Arg Lys Cys Cys Arg 530 535 540		
Arg Tyr Ser Ser Leu Asp Pro Arg Lys Val Gln Phe Tyr Phe Asn Lys 545 550 555 560		
Glu Thr Phe		

<210> 22

<211> 2577

<212> DNA

<213> Rat

<400> 22

ttctgagtag gtgtgcattt gagtgtcata aagacatata tcttgagcta gacagaagtt 60

cctatcctga aggatcccat cagagaaacc agattgctcc taagaggctc cagacctcca 120

tttgacaatg ggaaactggc tggttaacca ctggctctca gttttgtttc tggtttcttg 180

gttggggctg aacatttttc tgtttgtgta cgtcttcctg aattatgaga agtctgacaa 240

gtactattac acgagagaaa ttctcggaac tgccttggcc ttggccagag catctgcttt 300

gtgcctgaat ttaacagca tggtgatcct gattcctgtg tgctgaaatc tgctctcctt 360

WO 00/28031

PCT/US99/26592

cctgaggggc acctgctcat ttgcaacca cacgctgaga aagccattgg atcacaacct 420
 caccttccat aagctggtgg catatatgat ctgcatattc acagctattc atatcattgc 480
 acatctattt aactttgaac gctacagtag aagccaacag gccatggatg gatctcttgc 540
 ctctgttctc tccagcctat tccatcccga gaaagaagat tcttggttaa atcccatcca 600
 gtctccaaac gtgacagtga tgtatgcagc atttaccagt attgctggcc ttactggagt 660
 ggtcgccact gtggctttgg ttctcatggt aacttcagct atggagttta tccgcaggaa 720
 ttattttgag ctcttctggt atacacatca ccttttcatc atctatatca tctgcttagg 780
 gatccatggc ctggggggga ttgtccgggg tcaaacagaa gagagcatga gtgaaagtca 840
 tccccgcaac tgttcatact ctttccacga gtgggataag tatgaaagga gttgcaggag 900
 tcctcatttt gtggggcaac cccctgagtc ttggaagtgg atcctcgccg cgattgcttt 960
 ttatatcttt gaaaggatcc ttcgctttta tcgctcccgg cagaaggteg tgattaccaa 1020
 ggttgctatg caccatgta aagttttgga attgcagatg aggaagcggg gctttactat 1080
 gggaatagga cagtatatat tcgtaaattg cccctcgatt tccttcctgg aatggcatcc 1140
 ctttactctg acctctgctc cagaggaaga atttttctcc attcatattc gagcagcagg 1200
 ggactggaca gaaaatctca taaggacatt tgaacaacag cactcaccaa tgcccaggat 1260
 cgaggtggat ggtccctttg gcacagtcag tgaggatgct ttccagtagc aagtggctgt 1320
 actggttggg gcagggattg gcgtcactcc ctttgcttcc ttcttgaaat ctatctggta 1380
 caaattccag cgtgcacaca acaagctgaa aacacaaaag atctatttct actggatttg 1440
 tagagagacg ggtgcctttg cctgggtcaa caacttattg aattccctgg aacaagagat 1500
 ggacgaatta ggcaaaccgg atttcctaaa ctaccgactc ttctcactg gctgggatag 1560
 caacattgct ggtcatgcag cattaaactt tgacagagcc actgacgtcc tgacaggtct 1620
 gaaacagaaa acctcctttg ggagaccaat gtgggacaat gagttttcta gaatagctac 1680
 tgcccacccc aagtctgtgg tgggggtttt cttatgcggc cctccgactt tggcaaaaag 1740
 cctgcgcaaa tgctgtcggc ggtactcaag tctggatcct aggaaggttc aattctactt 1800

WO 00/28031

PCT/US99/26592

caacaaagaa acgttctgaa ttggaggaag ccgcacagta gtacttctcc atcttccttt 1860
 tcactaacgt gtgggtcagc tactagatag tccgttgctg cacaaggact tcactcccat 1920
 cttaaagttg actcaactcc atcattcttg ggctttggca acatgagagc tgcataactc 1980
 acaattgcaa aacacatgaa ttattattgg ggggattgta aatccttctg ggaaacctgc 2040
 ctttagctga atcttgctgg ttgacacttg cacaatttaa cctcaggtgt cttggttgat 2100
 acctgataat ctccctccc acctgtccct cacagaagat ttctaagtag ggtgatttta 2160
 aaatatttat tgaatccacg acaaaacaat aatcataaat aataaacata aaattaccaa 2220
 gattcccact cccatatcat acccactaag aacatcgta tacatgagct tatcatccag 2280
 tgtgaccaac aatttatact ttactgtgcc aaaataatct tcatctttgc ttattgaaca 2340
 attttgctga ctcccttag taatatctta agtatattaa ctggaatcaa atttgtatta 2400
 tagttagaag ccaactatat tgccagtttg tattgtttga aataactgga aaggcctgac 2460
 ctacatcgtg gggtaattta acagaagctc tttccatttt ttgttggtgt tgttaaagag 2520
 ttttgtttat gaatgtgta taaaagaaa ataaaagtt ataattttga cggaaaa 2577

<210> 23

<211> 332

<212> PRT

<213> Homo sapiens

<400> 23

Glu Ser Met Asn Glu Ser His Pro Arg Lys Cys Ala Glu Ser Phe Glu
 1 5 10 15

Met Trp Asp Asp Arg Asp Ser His Cys Arg Arg Pro Lys Phe Glu Gly
 20 25 30

His Pro Pro Glu Ser Trp Lys Trp Ile Leu Ala Pro Val Ile Leu Tyr
 35 40 45

Ile Cys Glu Arg Ile Leu Arg Phe Tyr Arg Ser Gln Gln Lys Val Val
 50 55 60

Ile Thr Lys Val Val Met His Pro Ser Lys Val Leu Glu Leu Gln Met
 65 70 75 80

WO 00/28031

PCT/US99/26592

Asn	Lys	Arg	Gly	Phe	Ser	Met	Glu	Val	Gly	Gln	Tyr	Ile	Phe	Val	Asn
				85					90					95	
Cys	Pro	Ser	Ile	Ser	Leu	Leu	Glu	Trp	His	Pro	Phe	Thr	Leu	Thr	Ser
			100					105					110		
Ala	Pro	Glu	Glu	Asp	Phe	Phe	Ser	Ile	His	Ile	Arg	Ala	Ala	Gly	Asp
		115					120					125			
Trp	Thr	Glu	Asn	Leu	Ile	Arg	Ala	Phe	Glu	Gln	Gln	Tyr	Ser	Pro	Ile
	130					135					140				
Pro	Arg	Ile	Glu	Val	Asp	Gly	Pro	Phe	Gly	Thr	Ala	Ser	Glu	Asp	Val
145					150				155					160	
Phe	Gln	Tyr	Glu	Val	Ala	Val	Leu	Val	Gly	Ala	Gly	Ile	Gly	Val	Thr
			165						170					175	
Pro	Phe	Ala	Ser	Ile	Leu	Lys	Ser	Ile	Trp	Tyr	Lys	Phe	Gln	Cys	Ala
			180					185					190		
Asp	His	Asn	Leu	Lys	Thr	Lys	Lys	Ile	Tyr	Phe	Tyr	Trp	Ile	Cys	Arg
		195					200					205			
Glu	Thr	Gly	Ala	Phe	Ser	Trp	Phe	Asn	Asn	Leu	Leu	Thr	Ser	Leu	Glu
	210					215					220				
Gln	Glu	Met	Glu	Glu	Leu	Gly	Lys	Val	Gly	Phe	Leu	Asn	Tyr	Arg	Leu
225					230				235						240
Phe	Leu	Thr	Gly	Trp	Asp	Ser	Asn	Ile	Val	Gly	His	Ala	Ala	Leu	Asn
			245						250					255	
Phe	Asp	Lys	Ala	Thr	Asp	Ile	Val	Thr	Gly	Leu	Lys	Gln	Lys	Thr	Ser
			260					265					270		
Phe	Gly	Arg	Pro	Met	Trp	Asp	Asn	Glu	Phe	Ser	Thr	Ile	Ala	Thr	Ser
		275					280					285			
His	Pro	Lys	Ser	Val	Val	Gly	Val	Phe	Leu	Cys	Gly	Pro	Arg	Thr	Leu
		290				295					300				
Ala	Lys	Ser	Leu	Arg	Lys	Cys	Cys	His	Arg	Tyr	Ser	Ser	Leu	Asp	Pro
305					310					315					320
Arg	Lys	Val	Gln	Phe	Tyr	Phe	Asn	Lys	Glu	Asn	Phe				
			325						330						

WO 00/28031

PCT/US99/26592

<210> 24

<211> 14

<212> PRT

<213> Homo sapiens

<400> 24

Cys Ala Glu Ser Phe Glu Met Trp Asp Asp Arg Asp Ser His
1 5 10

<210> 25

<211> 14

<212> PRT

<213> Homo sapiens

<400> 25

Lys Ser Leu Arg Lys Cys Cys His Arg Tyr Ser Ser Leu Asp
1 5 10

<210> 26

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 26

gaagtgggtgg gaggcgaaga cata

24

<210> 27

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 27

cctgtcatatc ctgggacggt ctgg

24

<210> 28

<211> 24

<212> DNA

WO 00/28031

PCT/US99/26592

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 28

gagcacagtg agatgcctgt tcag

24

<210> 29

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 29

ggaaggcagc agagagcaat gatg

24

<210> 30

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 30

acatctgcga gcggcacttc caga

24

<210> 31

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 31

agctcgtcaa caggcaggac cgagc

25

<210> 32

<211> 24

<212> DNA

WO 00/28031

PCT/US99/26592

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 32

gcagtgcattc cacatcttca gcac

24

<210> 33

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 33

gagagctctg gagacacttg agttc

25

<210> 34

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 34

catgttctct ctggctgaca ag

22

<210> 35

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 35

cacaatagcg agctccgctt cacgc

25

<210> 36

<211> 24

<212> DNA

WO 00/28031

PCT/US99/26592

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 36

gcaggacatc aaccctgcac tctc

24

<210> 37

<211> 570

<212> PRT

<213> Bovine

<400> 37

Met	Gly	Asn	Trp	Val	Val	Asn	Glu	Gly	Ile	Ser	Ile	Phe	Val	Ile	Leu
1				5					10					15	

Val	Trp	Leu	Gly	Met	Asn	Val	Phe	Leu	Phe	Val	Trp	Tyr	Tyr	Arg	Val
		20						25						30	

Tyr	Asp	Ile	Pro	Asp	Lys	Phe	Phe	Tyr	Thr	Arg	Lys	Leu	Leu	Gly	Ser
		35					40					45			

Ala	Leu	Ala	Leu	Ala	Arg	Ala	Pro	Ala	Ala	Cys	Leu	Asn	Phe	Asn	Cys
	50					55						60			

Met	Leu	Ile	Leu	Leu	Pro	Val	Cys	Arg	Asn	Leu	Leu	Ser	Phe	Leu	Arg
65					70					75					80

Gly	Ser	Ser	Ala	Cys	Cys	Ser	Thr	Arg	Ile	Arg	Arg	Gln	Leu	Asp	Arg
			85						90					95	

Asn	Leu	Thr	Phe	His	Lys	Met	Val	Ala	Trp	Met	Ile	Ala	Leu	His	Thr
		100						105					110		

Ala	Ile	His	Thr	Ile	Ala	His	Leu	Phe	Asn	Val	Glu	Trp	Cys	Val	Asn
	115					120						125			

Ala	Arg	Val	Asn	Asn	Ser	Asp	Pro	Tyr	Ser	Ile	Ala	Leu	Ser	Asp	Ile
	130					135					140				

Gly	Asp	Lys	Pro	Asn	Glu	Thr	Tyr	Leu	Asn	Phe	Val	Arg	Gln	Arg	Ile
145					150					155					160

Lys	Asn	Pro	Glu	Gly	Gly	Leu	Tyr	Val	Ala	Val	Thr	Arg	Leu	Ala	Gly
			165						170					175	

WO 00/28031

PCT/US99/26592

Ile Thr Gly Val Val Ile Thr Leu Cys Leu Ile Leu Ile Ile Thr Ser
 180 185 190

Ser Thr Lys Thr Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr
 195 200 205

His His Leu Phe Val Ile Phe Phe Ile Gly Leu Ala Ile His Gly Ala
 210 215 220

Gln Arg Ile Val Arg Gly Gln Thr Ala Glu Ser Leu Leu Lys His Gln
 225 230 235 240

Pro Arg Asn Cys Tyr Gln Asn Ile Ser Gln Trp Gly Lys Ile Glu Asn
 245 250 255

Cys Pro Ile Pro Glu Phe Ser Gly Asn Pro Pro Met Thr Trp Lys Trp
 260 265 270

Ile Val Gly Pro Met Phe Leu Tyr Leu Cys Glu Arg Leu Val Arg Phe
 275 280 285

Trp Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Thr His Pro
 290 295 300

Phe Lys Thr Ile Glu Leu Gln Met Lys Lys Lys Gly Phe Lys Met Glu
 305 310 315 320

Val Gly Gln Tyr Ile Phe Val Lys Cys Pro Val Val Ser Lys Leu Glu
 325 330 335

Trp His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser
 340 345 350

Ile His Ile Arg Ile Val Gly Asp Trp Thr Glu Gly Leu Phe Lys Ala
 355 360 365

Cys Gly Cys Asp Lys Gln Glu Phe Gln Asp Ala Trp Lys Leu Pro Lys
 370 375 380

Ile Ala Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe Ser
 385 390 395 400

Tyr Glu Val Val Met Leu Val Gly Ala Gly Ile Gly Val Thr Pro Phe
 405 410 415

Ala Ser Ile Leu Lys Ser Val Trp Tyr Lys Tyr Cys Asn Lys Ala Pro
 420 425 430

WO 00/28031

PCT/US99/26592

Asn Leu Arg Leu Lys Lys Ile Tyr Phe Tyr Trp Leu Cys Arg Asp Thr
 435 440 445
 His Ala Phe Glu Trp Phe Ala Asp Leu Leu Gln Leu Leu Glu Thr Gln
 450 455 460
 Met Gln Glu Lys Asn Asn Thr Asp Phe Leu Ser Tyr Asn Ile Cys Leu
 465 470 475 480
 Thr Gly Trp Asp Glu Ser Gln Ala Ser His Phe Ala Met His His Asp
 485 490 495
 Glu Glu Lys Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Leu Tyr Gly
 500 505 510
 Arg Pro Asn Trp Asp Asn Glu Phe Lys Thr Ile Gly Ser Gln His Pro
 515 520 525
 Asn Thr Arg Ile Gly Val Phe Leu Cys Gly Pro Glu Ala Leu Ala Asp
 530 535 540
 Thr Leu Asn Lys Gln Cys Ile Ser Asn Ser Asp Ser Gly Pro Arg Gly
 545 550 555 560
 Val His Phe Ile Phe Asn Lys Glu Asn Phe
 565 570

<210> 38

<211> 570

<212> PRT

<213> murine

<400> 38

Met Gly Asn Trp Ala Val Asn Glu Gly Leu Ser Ile Phe Val Ile Leu
 1 5 10 15
 Val Trp Leu Gly Leu Asn Val Phe Leu Phe Ile Asn Tyr Tyr Lys Val
 20 25 30
 Tyr Asp Asp Gly Pro Lys Tyr Asn Tyr Thr Arg Lys Leu Leu Gly Ser
 35 40 45
 Ala Leu Ala Leu Ala Arg Ala Pro Ala Ala Cys Leu Asn Phe Asn Cys
 50 55 60
 Met Leu Ile Leu Leu Pro Val Cys Arg Asn Leu Leu Ser Phe Leu Arg
 65 70 75 80

WO 00/28031

PCT/US99/26592

Gly Ser Ser Ala Cys Cys Ser Thr Arg Ile Arg Arg Gln Leu Asp Arg
 85 90 95

Asn Leu Thr Phe His Lys Met Val Ala Trp Met Ile Ala Leu His Thr
 100 105 110

Ala Ile His Thr Ile Ala His Leu Phe Asn Val Glu Trp Cys Val Asn
 115 120 125

Ala Arg Val Gly Ile Ser Asp Arg Tyr Ser Ile Ala Leu Ser Asp Ile
 130 135 140

Gly Asp Asn Glu Asn Glu Glu Tyr Leu Asn Phe Ala Arg Glu Lys Ile
 145 150 155 160

Lys Asn Pro Glu Gly Gly Leu Tyr Val Ala Val Thr Arg Leu Ala Gly
 165 170 175

Ile Thr Gly Ile Val Ile Thr Leu Cys Leu Ile Leu Ile Ile Thr Ser
 180 185 190

Ser Thr Lys Thr Ile Arg Arg Ser Tyr Phe Glu Val Phe Trp Tyr Thr
 195 200 205

His His Leu Phe Val Ile Phe Phe Ile Gly Leu Ala Ile His Gly Ala
 210 215 220

Glu Arg Ile Val Arg Gly Gln Thr Ala Glu Ser Leu Glu Glu His Asn
 225 230 235 240

Leu Asp Ile Cys Ala Asp Lys Ile Glu Glu Trp Gly Lys Ile Lys Glu
 245 250 255

Cys Pro Val Pro Lys Phe Ala Gly Asn Pro Pro Met Thr Trp Lys Trp
 260 265 270

Ile Val Gly Pro Met Phe Leu Tyr Leu Cys Glu Arg Leu Val Arg Phe
 275 280 285

Trp Arg Ser Gln Gln Lys Val Val Ile Thr Lys Val Val Thr His Pro
 290 295 300

Phe Lys Thr Ile Glu Leu Gln Met Lys Lys Lys Gly Phe Lys Met Glu
 305 310 315 320

Val Gly Gln Tyr Ile Phe Val Lys Cys Pro Lys Val Ser Lys Leu Glu
 325 330 335

WO 00/28031

PCT/US99/26592

Trp His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Asp Phe Phe Ser
 340 345 350
 Ile His Ile Arg Ile Val Gly Asp Trp Thr Glu Gly Leu Phe Asn Ala
 355 360 365
 Cys Gly Cys Asp Lys Gln Glu Phe Gln Asp Ala Trp Lys Leu Pro Lys
 370 375 380
 Ile Ala Val Asp Gly Pro Phe Gly Thr Ala Ser Glu Asp Val Phe Ser
 385 390 395 400
 Tyr Glu Val Val Met Leu Val Gly Ala Gly Ile Gly Val Thr Pro Phe
 405 410 415
 Ala Ser Ile Leu Lys Ser Val Trp Tyr Lys Tyr Cys Asp Asn Ala Thr
 420 425 430
 Ser Leu Lys Leu Lys Lys Ile Tyr Phe Tyr Trp Leu Cys Arg Asp Thr
 435 440 445
 His Ala Phe Glu Trp Phe Ala Asp Leu Leu Gln Leu Leu Glu Thr Gln
 450 455 460
 Met Gln Glu Arg Asn Asn Ala Asn Phe Leu Ser Tyr Asn Ile Tyr Leu
 465 470 475 480
 Thr Gly Trp Asp Glu Ser Gln Ala Asn His Phe Ala Val His His Asp
 485 490 495
 Glu Glu Lys Asp Val Ile Thr Gly Leu Lys Gln Lys Thr Leu Tyr Gly
 500 505 510
 Arg Pro Asn Trp Asp Asn Glu Phe Lys Thr Ile Ala Ser Glu His Pro
 515 520 525
 Asn Thr Thr Ile Gly Val Phe Leu Cys Gly Pro Glu Ala Leu Ala Glu
 530 535 540
 Thr Leu Ser Lys Gln Ser Ile Ser Asn Ser Glu Ser Gly Pro Arg Gly
 545 550 555 560
 Val His Phe Ile Phe Asn Lys Glu Asn Phe
 565 570

<210> 39

WO 00/28031

PCT/US99/26592

<211> 944

<212> PRT

<213> Arabidopsis sp.

<400> 39

Met Lys Pro Phe Ser Lys Asn Asp Arg Arg Arg Trp Ser Phe Asp Ser
 1 5 10 15

Val Ser Ala Gly Lys Thr Ala Val Gly Ser Ala Ser Thr Ser Pro Gly
 20 25 30

Thr Glu Tyr Ser Ile Asn Gly Asp Gln Glu Phe Val Glu Val Thr Ile
 35 40 45

Asp Leu Gln Asp Asp Asp Thr Ile Val Leu Arg Ser Val Glu Pro Ala
 50 55 60

Thr Ala Ile Asn Val Ile Gly Asp Ile Ser Asp Asp Asn Thr Gly Ile
 65 70 75 80

Met Thr Pro Val Ser Ile Ser Arg Ser Pro Thr Met Lys Arg Thr Ser
 85 90 95

Ser Asn Arg Phe Arg Gln Phe Ser Gln Glu Leu Lys Ala Glu Ala Val
 100 105 110

Ala Lys Ala Lys Gln Leu Ser Gln Glu Leu Lys Arg Phe Ser Trp Ser
 115 120 125

Arg Ser Phe Ser Gly Asn Leu Thr Thr Thr Ser Thr Ala Ala Asn Gln
 130 135 140

Ser Gly Gly Ala Gly Gly Gly Leu Val Asn Ser Ala Leu Glu Ala Arg
 145 150 155 160

Ala Leu Arg Lys Gln Arg Ala Gln Leu Asp Arg Thr Arg Ser Ser Ala
 165 170 175

Gln Arg Ala Leu Arg Gly Leu Arg Phe Ile Ser Asn Lys Gln Lys Asn
 180 185 190

Val Asp Gly Trp Asn Asp Val Gln Ser Asn Phe Glu Lys Phe Glu Lys
 195 200 205

Asn Gly Tyr Ile Tyr Arg Ser Asp Phe Ala Gln Cys Ile Gly Met Lys
 210 215 220

Asp Ser Lys Glu Phe Ala Leu Glu Leu Phe Asp Ala Leu Ser Arg Arg

WO 00/28031

PCT/US99/26592

225	230	235	240
Arg Arg Leu Lys Val Glu Lys Ile Asn His Asp Glu Leu Tyr Glu Tyr	245	250	255
Trp Ser Gln Ile Asn Asp Glu Ser Phe Asp Ser Arg Leu Gln Ile Phe	260	265	270
Phe Asp Ile Val Asp Lys Asn Glu Asp Gly Arg Ile Thr Glu Glu Glu	275	280	285
Val Lys Glu Ile Ile Met Leu Ser Ala Ser Ala Asn Lys Leu Ser Arg	290	295	300
Leu Lys Glu Gln Ala Glu Glu Tyr Ala Ala Leu Ile Met Glu Glu Leu	305	310	315
Asp Pro Glu Arg Leu Gly Tyr Ile Glu Leu Trp Gln Leu Glu Thr Leu	325	330	335
Leu Leu Gln Lys Asp Thr Tyr Leu Asn Tyr Ser Gln Ala Leu Ser Tyr	340	345	350
Thr Ser Gln Ala Leu Ser Gln Asn Leu Gln Gly Leu Arg Gly Lys Ser	355	360	365
Arg Ile His Arg Met Ser Ser Asp Phe Val Tyr Ile Met Gln Glu Asn	370	375	380
Trp Lys Arg Ile Trp Val Leu Ser Leu Trp Ile Met Ile Met Ile Gly	385	390	395
Leu Phe Leu Trp Lys Phe Phe Gln Tyr Lys Gln Lys Asp Ala Phe His	405	410	415
Val Met Gly Tyr Cys Leu Leu Thr Ala Lys Gly Ala Ala Glu Thr Leu	420	425	430
Lys Phe Asn Met Ala Leu Ile Leu Phe Pro Val Cys Arg Asn Thr Ile	435	440	445
Thr Trp Leu Arg Ser Thr Arg Leu Ser Tyr Phe Val Pro Phe Asp Asp	450	455	460
Asn Ile Asn Phe His Lys Thr Ile Ala Gly Ala Ile Val Val Ala Val	465	470	475
Ile Leu His Ile Gly Asp His Leu Ala Cys Asp Phe Pro Arg Ile Val			480

WO 00/28031

PCT/US99/26592

485	490	495
Arg Ala Thr Glu Tyr Asp Tyr Asn Arg Tyr Leu Phe His Tyr Phe Gln 500	505	510
Thr Lys Gln Pro Thr Tyr Phe Asp Leu Val Lys Gly Pro Glu Gly Ile 515	520	525
Thr Gly Ile Leu Met Val Ile Leu Met Ile Ile Ser Phe Thr Leu Ala 530	535	540
Thr Arg Trp Phe Arg Arg Asn Leu Val Lys Leu Pro Lys Pro Phe Asp 545	550	555 560
Arg Leu Thr Gly Phe Asn Ala Phe Trp Tyr Ser His His Leu Phe Val 565	570	575
Ile Val Tyr Ile Leu Leu Ile Leu His Gly Ile Phe Leu Tyr Phe Ala 580	585	590
Lys Pro Trp Tyr Val Arg Thr Thr Trp Met Tyr Leu Ala Val Pro Val 595	600	605
Leu Leu Tyr Gly Gly Glu Arg Thr Leu Arg Tyr Phe Arg Ser Gly Ser 610	615	620
Tyr Ser Val Arg Leu Leu Lys Val Ala Ile Tyr Pro Gly Asn Val Leu 625	630	635 640
Thr Leu Gln Met Ser Lys Pro Thr Gln Phe Arg Tyr Lys Ser Gly Gln 645	650	655
Tyr Met Phe Val Gln Cys Pro Ala Val Ser Pro Phe Glu Trp His Pro 660	665	670
Phe Ser Ile Thr Ser Ala Pro Glu Asp Asp Tyr Ile Ser Ile His Ile 675	680	685
Arg Gln Leu Gly Asp Trp Thr Gln Glu Leu Lys Arg Val Phe Ser Glu 690	695	700
Val Cys Glu Pro Pro Val Gly Gly Lys Ser Gly Leu Leu Arg Ala Asp 705	710	715 720
Glu Thr Thr Lys Lys Ser Leu Pro Lys Leu Leu Ile Asp Gly Pro Tyr 725	730	735
Gly Ala Pro Ala Gln Asp Tyr Arg Lys Tyr Asp Val Leu Leu Leu Val		

WO 00/28031

PCT/US99/26592

740	745	750
Gly Leu Gly Ile Gly Ala Thr	Pro Phe Ile Ser Ile	Leu Lys Asp Leu
755	760	765
Leu Asn Asn Ile Val Lys Met	Glu Glu His Ala Asp	Ser Ile Ser Asp
770	775	780
Phe Ser Arg Ser Ser Glu Tyr	Ser Thr Gly Ser Asn Gly	Asp Thr Pro
785	790	800
Arg Arg Lys Arg Ile Leu Lys	Thr Thr Asn Ala Tyr Phe	Tyr Trp Val
805	810	815
Thr Arg Glu Gln Gly Ser Phe	Asp Trp Phe Lys Gly	Val Met Asn Glu
820	825	830
Val Ala Glu Leu Asp Gln Arg	Gly Val Ile Glu Met	His Asn Tyr Leu
835	840	845
Thr Ser Val Tyr Glu Glu Gly	Asp Ala Arg Ser Ala	Leu Ile Thr Met
850	855	860
Val Gln Ala Leu Asn His Ala	Lys Asn Gly Val Asp	Ile Val Ser Gly
865	870	875
Thr Arg Val Arg Thr His Phe	Ala Arg Pro Asn Trp	Lys Lys Val Leu
885	890	895
Thr Lys Leu Ser Ser Lys His	Cys Asn Ala Arg Ile	Gly Val Phe Tyr
900	905	910
Cys Gly Val Pro Val Leu Gly	Lys Glu Leu Ser Lys	Leu Cys Asn Thr
915	920	925
Phe Asn Gln Lys Gly Ser Thr	Lys Phe Glu Phe His	Lys Glu His Phe
930	935	940

<210> 40
 <211> 590
 <212> PRT
 <213> Rice

<400> 40

WO 00/28031

PCT/US99/26592

Asn Leu Ala Gly Leu Arg Lys Lys Ser Ser Ile Arg Lys Ile Ser Thr
 1 5 10 15
 Ser Leu Ser Tyr Tyr Phe Glu Asp Asn Trp Lys Arg Leu Trp Val Leu
 20 25 30
 Ala Leu Trp Ile Gly Ile Met Ala Gly Leu Phe Thr Trp Lys Phe Met
 35 40 45
 Gln Tyr Arg Asn Arg Tyr Val Phe Asp Val Met Gly Tyr Cys Val Thr
 50 55 60
 Thr Ala Lys Gly Ala Ala Glu Thr Leu Lys Leu Asn Met Ala Ile Ile
 65 70 75 80
 Leu Leu Pro Val Cys Arg Asn Thr Ile Thr Trp Leu Arg Ser Thr Arg
 85 90 95
 Ala Ala Arg Ala Leu Pro Phe Asp Asp Asn Ile Asn Phe His Lys Thr
 100 105 110
 Ile Ala Ala Ala Ile Val Val Gly Ile Ile Leu His Ala Gly Asn His
 115 120 125
 Leu Val Cys Asp Phe Pro Arg Leu Ile Lys Ser Ser Asp Glu Lys Tyr
 130 135 140
 Ala Pro Leu Gly Gln Tyr Phe Gly Glu Ile Lys Pro Thr Tyr Phe Thr
 145 150 155 160
 Leu Val Lys Gly Val Glu Gly Ile Thr Gly Val Ile Met Val Val Cys
 165 170 175
 Met Ile Ile Ala Phe Thr Leu Ala Thr Arg Trp Phe Arg Arg Ser Leu
 180 185 190
 Val Lys Leu Pro Arg Pro Phe Asp Lys Leu Thr Gly Phe Asn Ala Phe
 195 200 205
 Trp Tyr Ser His His Leu Phe Ile Ile Val Tyr Ile Ala Leu Ile Val
 210 215 220
 His Gly Glu Cys Leu Tyr Leu Ile His Val Trp Tyr Arg Arg Thr Thr
 225 230 235 240
 Trp Met Tyr Leu Ser Val Pro Val Cys Leu Tyr Val Gly Glu Arg Ile
 245 250 255

WO 00/28031

PCT/US99/26592

Leu Arg Phe Phe Arg Ser Gly Ser Tyr Ser Val Arg Leu Leu Lys Val
 260 265 270

Ala Ile Tyr Pro Gly Asn Val Leu Thr Leu Gln Met Ser Lys Pro Pro
 275 280 285

Thr Phe Arg Tyr Lys Ser Gly Gln Tyr Met Phe Val Gln Cys Pro Ala
 290 295 300

Val Ser Pro Phe Glu Trp His Pro Phe Ser Ile Thr Ser Ala Pro Gly
 305 310 315 320

Asp Asp Tyr Leu Ser Ile His Val Arg Gln Leu Gly Asp Trp Thr Arg
 325 330 335

Glu Leu Lys Arg Val Phe Ala Ala Ala Cys Glu Pro Pro Ala Gly Gly
 340 345 350

Lys Ser Gly Leu Leu Arg Ala Asp Glu Thr Thr Lys Lys Ile Leu Pro
 355 360 365

Lys Leu Leu Ile Asp Gly Pro Tyr Gly Ser Pro Ala Gln Asp Tyr Ser
 370 375 380

Lys Tyr Asp Val Leu Leu Leu Val Gly Leu Gly Ile Gly Ala Thr Pro
 385 390 395 400

Phe Ile Ser Ile Leu Lys Asp Leu Leu Asn Asn Ile Ile Lys Met Glu
 405 410 415

Glu Glu Glu Asp Ala Ser Thr Asp Leu Tyr Pro Pro Met Gly Arg Asn
 420 425 430

Asn Pro His Val Asp Leu Gly Thr Leu Met Thr Ile Thr Ser Arg Pro
 435 440 445

Lys Lys Ile Leu Lys Thr Thr Asn Ala Tyr Phe Tyr Trp Val Thr Arg
 450 455 460

Glu Gln Gly Ser Phe Asp Trp Phe Lys Gly Val Met Asn Glu Ile Ala
 465 470 475 480

Asp Leu Asp Gln Arg Asn Ile Ile Glu Met His Asn Tyr Leu Thr Ser
 485 490 495

Val Tyr Glu Glu Gly Asp Ala Arg Ser Ala Leu Ile Thr Met Leu Gln
 500 505 510

WO 00/28031

PCT/US99/26592

Ala Leu Asn His Ala Lys Asn Gly Val Asp Ile Val Ser Gly Thr Lys
 515 520 525

Val Arg Thr His Phe Ala Arg Pro Asn Trp Arg Lys Val Leu Ser Lys
 530 535 540

Ile Ser Ser Lys His Pro Tyr Ala Lys Ile Gly Val Phe Tyr Cys Gly
 545 550 555 560

Ala Pro Val Leu Ala Gln Glu Leu Ser Lys Leu Cys His Glu Phe Asn
 565 570 575

Gly Lys Cys Thr Thr Lys Phe Asp Phe His Lys Glu His Phe
 580 585 590

<210> 41
 <211> 2619
 <212> DNA
 <213> Rat

<400> 41
 gtgctgtcag agctttacag agcctctggg catgcgcatg gctacccatt tcattgattt 60
 acagaagtca tgctaaaatc tctttcatgc atgtcttctt ttttcagtct ctcctttccc 120
 aaagcttttc agtttgccct ttgcttgtac caactgctat cctcctcaa aggctgctgc 180
 aaaaggatg cctttttctt ggaggctttc agcaaatact acctgggaac ctgcttcagc 240
 tcttggaata tttaagtga gagaacattt catagcattt gtatctttct ttgaaggagc 300
 caccagacag actgccttgg ccttggccag agcatctgct ttgtgcctga attttaacag 360
 catggtgatc ctgattcctg tgtgtcgaaa tctgctctcc ttctgaggg gcacctgctc 420
 attttgcaac cacacgctga gaaagccatt ggatcacaac ctcaccttc ataagctgg 480
 ggcatatatg atctgcatat tcacagctat tcatatcatt gcacatctat ttaactttga 540
 acgctacagt agaagccaac aggccatgga tggatctctt gcctctgttc tctccagcct 600
 attccatccc gagaaagaag attcttggct aaatcccatc cagtctccaa acgtgacagt 660
 gatgtatgca gcatttacca gtattgctgg ccttactgga gtggtcgcca ctgtggcttt 720
 ggttctcatg gtaacttcag ctatggagtt tatccgcagg aattattttg agctcttctg 780

WO 00/28031

PCT/US99/26592

gtatacacat caccttttca tcatctatat catctgctta gggatccatg gcctgggggg 840
 gattgtccgg ggtcaaacag aagagagcat gagtgaaagt catccccgca actgttcata 900
 ctctttccac gagtgggata agtatgaaag gagttgcagg agtcctcatt ttgtggggca 960
 accccctgag tcttggaagt ggatcctcgc gccgattgct tttatatct ttgaaaggat 1020
 ccttcgcttt tategctccc ggcagaagggt cgtgattacc aaggttgtca tgcacccatg 1080
 taaagttttg gaattgcaga tgaggaagcg gggctttact atgggaatag gacagtatat 1140
 attcgtaaatt tgcacctcga tttccttcct ggaatggcat ccctttactc tgacctctgc 1200
 tccagaggaa gaatttttct ccattcatat tcgagcagca ggggactgga cagaaaatct 1260
 cataaggaca tttgaacaac agcactcacc aatgcccagg atcgagggtg atggtccctt 1320
 tggcacagtc agtgaggatg tcttcagta cgaagtggct gtactgggtg gggcagggat 1380
 tggcgctact ccctttgctt ccttcttgaa atctatctgg tacaaattcc agcgtgcaca 1440
 caacaagctg aaaacacaaa agatctatct ctactggatt tgtagagaga cgggtgcctt 1500
 tgcttggttc aacaacttat tgaattccct ggaacaagag atggacgaat taggcaaacc 1560
 ggatttccta aactaccgac tcttcctcac tggctgggat agcaacattg ctggtcatgc 1620
 agcattaaac tttgacagag ccactgacgt cctgacaggt ctgaaacaga aaacctcctt 1680
 tgggagacca atgtgggaca atgagttttc tagaatagct actgcccacc ccaagtctgt 1740
 ggtggggggtt ttcttatgcg gccctccgac tttggcaaaa agcctgcgca aatgctgtcg 1800
 gcggtactca agtctggatc ctaggaagggt tcaattctac ttcaacaaag aaacgttctg 1860
 aattggagga agccgcacag tagtacttct ccacttcct tttcactaac gtgtgggtca 1920
 gctactagat agtccgttgt cgcacaagga cttcactccc atcttaaagt tgactcaact 1980
 ccactattct tgggcttttg caacatgaga gctgcataac tcacaattgc aaaacacatg 2040
 aattattatt ggggggattg taaatccttc tgggaaacct gccttttagct gaatcttgct 2100
 ggttgacact tgcacaattt aacctcaggt gtcttggttg atacctgata atcttcctc 2160
 ccacctgtcc ctcacagaag atttctaagt agggtgattt taaaatattt attgaatcca 2220

WO 00/28031

PCT/US99/26592

cgacaaaaca ataatcataa ataataaaca taaaattacc aagattccca ctcccatatc 2280
 ataccacta agaacatcgt tatacatgag cttatcatcc agtgtgacca acaatttata 2340
 ctttactgtg ccaaaataat cttcatcttt gcttattgaa caattttgct gactttccct 2400
 agtaatatct taagtatatt aactggaatc aaatttgat tatagttaga agccaactat 2460
 attgccagtt tgtattgttt gaaataactg gaaaggcctg acctacatcg tggggtaatt 2520
 taacagaagc tctttccatt ttttggtgtt gttgttaaag agttttgttt atgaatgtgt 2580
 tataaaaaga aaataaaaag ttataatttt gacggaaaa 2619

<210> 42

<211> 499

<212> PRT

<213> Rat

<400> 42

Met	Val	Ile	Leu	Ile	Pro	Val	Cys	Arg	Asn	Leu	Leu	Ser	Phe	Leu	Arg
1				5					10					15	
Gly	Thr	Cys	Ser	Phe	Cys	Asn	His	Thr	Leu	Arg	Lys	Pro	Leu	Asp	His
			20					25					30		
Asn	Leu	Thr	Phe	His	Lys	Leu	Val	Ala	Tyr	Met	Ile	Cys	Ile	Phe	Thr
			35				40					45			
Ala	Ile	His	Ile	Ile	Ala	His	Leu	Phe	Asn	Phe	Glu	Arg	Tyr	Ser	Arg
	50					55					60				
Ser	Gln	Gln	Ala	Met	Asp	Gly	Ser	Leu	Ala	Ser	Val	Leu	Ser	Ser	Leu
65					70					75					80
Phe	His	Pro	Glu	Lys	Glu	Asp	Ser	Trp	Leu	Asn	Pro	Ile	Gln	Ser	Pro
				85					90					95	
Asn	Val	Thr	Val	Met	Tyr	Ala	Ala	Phe	Thr	Ser	Ile	Ala	Gly	Leu	Thr
			100					105					110		
Gly	Val	Val	Ala	Thr	Val	Ala	Leu	Val	Leu	Met	Val	Thr	Ser	Ala	Met
		115					120					125			
Glu	Phe	Ile	Arg	Arg	Asn	Tyr	Phe	Glu	Leu	Phe	Trp	Tyr	Thr	His	His
	130					135					140				

WO 00/28031

PCT/US99/26592

Leu Phe Ile Ile Tyr Ile Ile Cys Leu Gly Ile His Gly Leu Gly Gly	
145	150 155 160
Ile Val Arg Gly Gln Thr Glu Glu Ser Met Ser Glu Ser His Pro Arg	
	165 170 175
Asn Cys Ser Tyr Ser Phe His Glu Trp Asp Lys Tyr Glu Arg Ser Cys	
	180 185 190
Arg Ser Pro His Phe Val Gly Gln Pro Pro Glu Ser Trp Lys Trp Ile	
	195 200 205
Leu Ala Pro Ile Ala Phe Tyr Ile Phe Glu Arg Ile Leu Arg Phe Tyr	
	210 215 220
Arg Ser Arg Gln Lys Val Val Ile Thr Lys Val Val Met His Pro Cys	
	225 230 235 240
Lys Val Leu Glu Leu Gln Met Arg Lys Arg Gly Phe Thr Met Gly Ile	
	245 250 255
Gly Gln Tyr Ile Phe Val Asn Cys Pro Ser Ile Ser Phe Leu Glu Trp	
	260 265 270
His Pro Phe Thr Leu Thr Ser Ala Pro Glu Glu Glu Phe Phe Ser Ile	
	275 280 285
His Ile Arg Ala Ala Gly Asp Trp Thr Glu Asn Leu Ile Arg Thr Phe	
	290 295 300
Glu Gln Gln His Ser Pro Met Pro Arg Ile Glu Val Asp Gly Pro Phe	
	305 310 315 320
Gly Thr Val Ser Glu Asp Val Phe Gln Tyr Glu Val Ala Val Leu Val	
	325 330 335
Gly Ala Gly Ile Gly Val Thr Pro Phe Ala Ser Phe Leu Lys Ser Ile	
	340 345 350
Trp Tyr Lys Phe Gln Arg Ala His Asn Lys Leu Lys Thr Gln Lys Ile	
	355 360 365
Tyr Phe Tyr Trp Ile Cys Arg Glu Thr Gly Ala Phe Ala Trp Phe Asn	
	370 375 380
Asn Leu Leu Asn Ser Leu Glu Gln Glu Met Asp Glu Leu Gly Lys Pro	
	385 390 395 400

WO 00/28031

PCT/US99/26592

Asp Phe Leu Asn Tyr Arg Leu Phe Leu Thr Gly Trp Asp Ser Asn Ile
 405 410 415

Ala Gly His Ala Ala Leu Asn Phe Asp Arg Ala Thr Asp Val Leu Thr
 420 425 430

Gly Leu Lys Gln Lys Thr Ser Phe Gly Arg Pro Met Trp Asp Asn Glu
 435 440 445

Phe Ser Arg Ile Ala Thr Ala His Pro Lys Ser Val Val Gly Val Phe
 450 455 460

Leu Cys Gly Pro Pro Thr Leu Ala Lys Ser Leu Arg Lys Cys Cys Arg
 465 470 475 480

Arg Tyr Ser Ser Leu Asp Pro Arg Lys Val Gln Phe Tyr Phe Asn Lys
 485 490 495

Glu Thr Phe

<210> 43
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: synthetic
 primer

<400> 43
 ttctgagtag gtgtgcattt gagtgtcata aagac 35

<210> 44
 <211> 45
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: synthetic
 primer

<400> 44
 ttttccgtca aaattataac tttttatattt ctttttataa cacat 45

WO 00/28031

PCT/US99/26592

<210> 45

<211> 5508

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (155)..(4810)

<400> 45

```

1   GCAGAGCTGC AGAGGCACCG GACGAGAGAG GGCTCCGCGG GCCCAGCTGG CAGCCAGGCC
61  GGAGACAAGT TGCAGTCCCG GGCTCTGGTG ACGCCGTGGC CGCAGGGTCT CCATTTTGGG
121 ACATTCTAAT CCCTGAGCCC CTATTATTTT CATCATGGGC TTCTGCCTGG CTCTAGCATG
181 GACACTTCTG GTTGGGGCAT GGACCCCTCT GGGAGCTCAG AACCCCATTT CGTGGGAGGT
241 GCAGCGATTT GATGGGTGGT ACAACAACCT CATGGAGCAC AGATGGGGCA GCAAAGGCTC
301 CCGGCTGCAG CGCCTGGTCC CAGCCAGCTA TGCAGATGGC GTGTACCAGC CCTTGGGAGA
361 ACCCCACCTG CCCAACCCCC GAGACCTTAG CAACACCATC TCAAGGGGCC CTGCAGGGCT
421 GGCTTCCCTG AGAAACCGCA CAGTGTGGG GGTCTTCTTT GGCTATCACG TGCTTTCAGA
481 CCTGGTGAGC GTGGAAACTC CCGGCTGCCC CGCCGAGTTC CTCAACATTC GCATCCCGCC
541 CGGAGACCCC ATGTTTCGACC CCGACCAGCG CGGGGACGTG GTGCTGCCCT TCCAGAGAAG
601 CCGCTGGGAC CCCGAGACCG GACGGAGTCC CAGCAATCCC CGGGACCCGG CCAACCAGGT
661 GACGGGCTGG CTGGACGGCA GCGCCATCTA TGGTTCCTCG CATTCCCTGA GCGACGCGCT
721 GCGGAGCTTC TCCAGGGGAC AGCTGGCGTC GGGGCCCGAC CCCGCTTTTC CCCGAGACTC
781 GCAGAACCCC CTGCTCATGT GGGCGGCGCC CGACCCCGCC ACCGGGCAGA ACGGGCCCCG
841 GGGGCTGTAC GCCTTCGGGG CAGAGAGAGG GAACCGGGAA CCCTTCCTGC AGGCGCTGGG
901 CCTGCTCTGG TTCCGCTACC ACAACCTGTG GGCGCAGAGG CTGGCCCGCC AGCACCAGCA
961 CTGGGAGGAC GAGGAGCTGT TCCAGCACGC ACGCAAGAGG GTCATCGCCA CCTACCAGAA
1021 CATCGCTGTG TATGAGTGGC TGCCCAGCTT CCTGCAGAAA AACTCCCGG AGTATACAGG
1081 ATACCGGCCA TTTCTGGACC CCAGCATCTC CTCAGAGTTC GTGGCGGCCT CTGAGCAGTT
1141 CCTGTCCACC ATGGTGCCCC CTGGCGTCTA CATGAGAAAT GCCAGCTGCC ACTTCCAGGG
1201 GGTCATCAAT CGGAACTCAA GTGTCTCCAG AGCTCTCCGG GTCTGCAACA GCTACTGGAG
1261 CCGTGAGCAC CCAAGCCTAC AAAGTGCTGA AGATGTGGAT GCACTGCTGC TGGGCATGGC
1321 CTCCCAGATC GCAGAGCGAG AGGACCATGT GTTGGTTGAA GATGTGCGGG ATTTCTGGCC
1381 TGGGCCACTG AAGTTTCCC GCACAGACCA CCTGGCCAGC TGCTGCAGC GGGGCCGGGA
1441 TCTGGGCCTG CCCTCTTACA CCAAGGCCAG GGCAGCACTG GGCTTGTCTC CCATTACCCG
1501 CTGGCAGGAC ATCAACCCTG CACTCTCCCG GAGCAATGAC ACTGTACTGG AGGCCACAGC
1561 TGCCCTGTAC AACCAGGACT TATCCTGGCT AGAGCTGCTC CCTGGGGGAC TCCTGGAGAG
1621 CCACCGGGAC CCTGGACCTC TGTTACGAC CATCGTCCTT GAACAATTTG TGCGGCTACG
1681 GGATGGTGAC CGCTACTGGT TTGAGAACAC CAGGAATGGG CTGTTCTCCA AGAAGGAGAT
1741 TGAAGAAATC CGAAATACCA CCCTGCAGGA CGTGCTGGTC GCTGTTATCA ACATTGACCC
1801 CAGTGCTCTG CAGCCCAATG TCTTTGTCTG GCATAAAGGA GACCCCTGTC CGCAGCCGAG
1861 ACAGCTCAGC ACTGAAGGCC TGCCAGCGTG TGCTCCCTCT GTTGTTCGTG ACTATTTTGA
1921 GGGCAGTGGA TTTGGCTTCG GGGTCACCAT CGGGACCCTC TGTTGCTTCC CTTTGGTGAG
1981 CCTGCTCAGT GCCTGGATTG TTGCCC GGCT CCGGATGAGA AATTTCAAGA GGCTCCAGGG
2041 CCAGGACCGC CAGAGCATCG TGTCTGAGAA GCTCGTGGGA GGCATGGAAG CTTTGGAAATG
2101 GCAAGGCCAC AAGGAGCCCT GCCGGCCCGT GCTTGTGTAC CTGCAGCCCG GGCAGATCCG
2161 TGTGGTAGAT GGCAGGCTCA CCGTGCTCCG CACCATCCAG CTGCAGCCTC CACAGAAGGT
2221 CAACTTCGTC CTGTCCAGCA ACCGTGGACG CCGCACTCTG CTGCTCAAGA TCCCAAGGA
2281 GTATGACCTG GTGCTGCTGT TTAAC TTGGA GGAAGAGCGG CAGGCGCTGG TGGAAAATCT
2341 CCGGGGAGCT CTGAAGGAGA GCGGGTTGAG CATCCAGGAG TGGGAGCTGC GGGAGCAGGA

```

WO 00/28031

PCT/US99/26592

2401 GCTGATGAGA GCAGCTGTGA CACGGGAGCA GCGGAGGCAC CTCCTGGAGA CCTTTTTCAG
 2461 GCACCTTTTC TCCCAGGTGC TGGACATCAA CCAGGCCGAC GCAGGGACCC TGCCCTGGA
 2521 CTCCTCCCAG AAGGTGCGGG AGGCCCTGAC CTGTGAGCTG AGCAGGGCCG AGTTTGCCGA
 2581 GTCCCTGGGC CTCAAGCCCC AGGACATGTT TGTGGAGTCC ATGTTCTCTC TGGCTGACAA
 2641 GGATGGCAAT GGCTACCTGT CCTTCCGAGA GTTCCTGGAC ATCCTGGTGG TCTTCATGAA
 2701 AGGCTCTCCT GAGGAAAAGT CTCGCCTTAT GTTCCGCATG TACGACTTTG ATGGGAATGG
 2761 CCTCATTTCC AAGGATGAGT TCATCAGGAT GCTGAGATCC TTCATCGAGA TCTCCAACAA
 2821 CTGCCTGTCC AAGGCCCAGC TGGCTGAGGT GGTGGAGTCC ATGTTCCGGG AGTCGGGATT
 2881 CCAGGACAAG GAGGAACTGA CATGGGAAGA TTTTCACTTC ATGCTGCGGG ACCACAATAG
 2941 CGAGCTCCGC TTCACGCAGC TCTGTGTCAA AGGGGTGGAG GTGCCTGAAG TCATCAAGGA
 3001 CCTCTGCCGG CGAGCCTCCT ACATCAGCCA GGATATGATC TGTCCCTCTC CCAGAGTGAG
 3061 TGCCCGCTGT TCCCGCAGCG ACATTGAGAC TGAGTTGACA CCTCAGAGAC TGCAGTGCCC
 3121 CATGGACACA GACCCTCCCC AGGAGATTTCG GCGGAGGTTT GGCAAGAAGG TAACGTCATT
 3181 CCAGCCCTTG CTGTTCACTG AGGCGCACCG AGAGAAGTTC CAACGCAGCT GTCTCCACCA
 3241 GACGGTGCAA CAGTTCAAGC GCTTCATTGA GAACTACCGG CGCCACATCG GCTGCGTGGC
 3301 CGTGTCTTAC GCCATCGCTG GGGGGCTTTT CCTGGAGAGG GCCTACTACT ACGCCTTTGC
 3361 CGCACATCAC ACGGGCATCA CGGACACCAC CCGCGTGGGA ATCATCCTGT CGCGGGGCAC
 3421 AGCAGCCAGC ATCTCTTTCA TGTCTCTCTA CATCTTGCTC ACCATGTGCC GCAACCTCAT
 3481 CACCTTCCTG CGAGAAACCT TCCTCAACCG CTACGTGCCC TTCGACGCCG CCGTGGACTT
 3541 CCATCGCCTC ATTGCCTCCA CCGCCATCGT CCTCACAGTC TTACACAGTG TGGGCCATGT
 3601 GGTGAATGTG TACCTGTTCT CCATCAGCCC CCTCAGCGTC CTCTCTTGCC TCTTTCCTGG
 3661 CCTCTTCCAT GATGATGGGT CTGAGTTCCC CCAGAAGTAT TACTGGTGGT TCTTCCAGAC
 3721 CGTACCAGGC CTCACGGGGG TTGTGCTGCT CCTGATCCTG GCCATCATGT ATGTCTTTGC
 3781 CTCCCACCAC TTCCGCCGCC GCAGTTTCCG GGGCTTCTGG CTGACCCACC ACCTCTACAT
 3841 CCTGCTCTAT GTCTGCTCA TCATCCATGG TAGCTTTGCC CTGATCCAGC TGCCCCGTTT
 3901 CCACATCTTC TTCCTGGTCC CAGCAATCAT CTATGGGGGC GACAAGCTGG TGAGCCTGAG
 3961 CCGGAAGAAG GTGGAGATCA GCGTGGTGAA GCGGAGCTG CTGCCCTCAG GAGTGACCCA
 4021 CCTGCGGTTT CAGCGGCCCC AGGGCTTTGA GTACAAGTCA GGGCAGTGGG TGCGGATCGC
 4081 TTGCCTGGCT CTGGGGACCA CCGAGTACCA CCCCTTCACA CTGACCTCTG CGCCCCATGA
 4141 GGACACGCTT AGCCTGCACA TCCGGGCAGC AGGGCCCTGG ACCACTCGCC TCAGGGAGAT
 4201 CTACTCAGCC CCGACGGGTG ACAGATGTGC CAGATACCCA AAGCTGTACC TTGATGGACC
 4261 ATTTGGAGAG GGCCACCAGG AGTGGCATAA GTTTGAGGTG TCAGTGTTAG TGGGAGGGGG
 4321 CATTGGGGTC ACCCCTTTTG CCTCCATCCT CAAAGACCTG GTCTTCAAGT CATCCGTCAG
 4381 CTGCCAAGTG TTCTGTAAGA AGATCTACTT CATCTGGGTG ACGCGGACCC AGCGTCAGTT
 4441 TGAGTGCTG GCTGACATCA TCCGAGAGGT GGAGGAGAAT GACCACCAGG ACCTGGTGTC
 4501 TGTGCACATC TACATCACCC AGCTGGCTGA GAAGTTCGAC CTCAGGACCA CTATGCTGTA
 4561 CATCTGTGAG CGGCACTTCC AGAAGGTCTT GAACCGGAGT CTATTCACAG GCCTGCGCTC
 4621 CATCACCCAC TTTGGCCGTC CCCCCTTTGA GCCCTTCTTC AACTCCCTGC AGGAGGTCCA
 4681 CCCCCAGGTC CGGAAGATCG GGGTGTTTAG CTGTGGCCCC CCTGGCATGA CCAAGAATGT
 4741 GGAAAAGGCC TGTCAGCTCA TCAACAGGCA GGACCGGACT CACTTCTCCC ACCATTATGA
 4801 GAACTTCTAG GCCCCTGCCC GGGGGTTCTG CCCACTGTCC AGTTGAGCAG AGGTTTGAGC
 4861 CCACACCTCA CCTCTGTTCT TCCTATTTCT GGCTGCCTCA GCCTTCTCTG ATTTCCCACC
 4921 TCCCAACCTT GTTCCAGGTG GCCATAGTCA GTCACCATGT GTGGGCTCAG GGACCCCCAG
 4981 GACCAGGATG TGTCTCAGCC TGGAGAAATG GTGGGGGGGC AGTGTCTAGG GACTAGAGTG
 5041 AGAAGTAGGG GAGCTACTGA TTTGGGGCAA AGTGAAACCT CTGCTTCAGA CTTCAGAAAC
 5101 AAATCTCAGA AGACAAGCTG ACCTGACAAG TACTATGTGT GTGCATGTCT GTATGTGTGT
 5161 TGGGGCGGTG AGTGTAAGGA TGCAGTGGGA GCATGGATGC TGGCATCTTA GAACCCCTCC
 5221 TACTCCCATC CCTCCTCCTC TTCTGGGCTC CCCACTGTCA GACGGGCTGG CAAATGCCTT
 5281 GCAGGAGGTA GAGGCTGGAC CCATGGCAAG CCATTTACAG AAACCCACTC GGCACCCAG

WO 00/28031

PCT/US99/26592

5341 TCTAACACCA CAACTAATTT CACCCAAGGT TTTAAGCACG TTCTTTCATC AGACCCTGGC
5401 CCAATACCTA TGTATGCAAT GCTCCTCAGC CCTCTTCTCC CTGCTCCAGT AGTCTCCCTT
5461 CCAAATAAAT CACTTTTCTG CCAAAAAAAAA AAAA

WO 00/28031

PCT/US99/26592

<210> 46
<211> 1521
<212> PRT
<213> Homo sapiens

<400> 46

WO 00/28031

PCT/US99/26592

1 MGFCALAWT LLVGAWTPLG AQNPISWEVQ RFDGWYNNLM EHRWGSKGSR LQRLVPASYA
 61 DGVYQPLGEP HLPNPRDLSN TISRGPAOLA SLRNRTVLGV FFGYHVLSL VSVETPGCPA
 121 EFLNIRIPPG DPMFDPDQRG DVVLPFQSR WDPETGRSPS NPRDPANQVT GWLDGSAIYG
 181 SSHSWSDALR SFSRGQLASG PDPAFPRDSQ NPLLMWAAPD PATGQNGPRG LYAFGAERGN
 241 REPFLQALGL LWFRYHNLWA QRLARQHPDW EDEELFQHAR KRVIATYQNI AVYEWLPSFL
 301 QKTLPEYTG Y RPFLDPSISS EFVAASEQFL STMVPPGVYM RNASCHFQGV INRNSSVSRA
 361 LRVCSYWSR EHPSLQSAED VDALLLGMA S QIAEREDHVL VEDVRDFWPG PLKFSRTDHL
 421 ASCLQRGRDL GLPSYTKARA ALGLSPITRW QDINPALSRS NDTVLEATAA LYNQDL SWLE
 481 LLPGGLLESH RDPGPLFSTI VLEQFVRLRD GDRYWFENTR NGLFSKKEIE EIRNTTLQDV
 541 LVAVINIDPS ALQPNVFWH KGDPCPQPRQ LSTEGLPACA PSVVRDYFEG SGFGFGVTIG
 601 TLCCFPLVSL LSAWIVARLR MRNFKRLQGO DRQSIVSEKL VGGMEALEWQ GHKEPCRPVL
 661 VYLQPGQIRV VDGRLTVLRT IQLQPPQKVN FVLSSNRGR TLLKIPKEY DLVLLFNLEE
 721 ERQALVENLR GALKESGLSI QEWELEQEL MRAAVTREQR RHLLETFFRH LFSQVLDINQ
 781 ADAGTLPLDS SQKVREALTC ELSRAEFAES LGLKPQDMFV ESMFSLADKD GNGYLSFREF
 841 LDILVVFMDG SPEEKSRMLF RMYDFDGNL ISKDEFIRML RSFIEISNNC LSKAQLAEVV
 901 ESMFRESGFQ DKEELTWEDF HFMLRDHNSE LRFTQLCVKG VEVPEVIKDL CRRASYISQD
 961 MICPSPRVSA RCSRSDIETE LTPQRLQCPM DTDPPQEIRR RFGKKVTSFQ PLLFTEAHRE
 1021 KFQRSCLHQT VQQFKRFIEN YRRHIGCVAV FYAIAGGLFL ERAYYYAFAA HHTGITDTTR
 1081 VGIILSRGTA ASISFMFSYI LLTMCRLNLT FLRETFLNRY VPFDAAVDFH RLIATAIVL
 1141 TVLHSVGHVV NVYLFSSISPL SVLSCLFPGL FHDDGSEFPQ KYWFFQTV PGLTGVVLLL
 1201 ILAIMYVFAS HHFRRSFRG FWLTHLYIL LYVLLIIHGS FALIQLPRFH IFFLVPAIY
 1261 GGDKLVSLSR KKVEISVKA ELLPSGVTHL RFQRPQGFY KSGQWVRIAC LALGTTEYHP
 1321 FTLTSAPHED TSLHIRAAG PWTTRLREIY SAPTGDRCAR YPKLYLDGPF GEGHQEWHKF
 1381 EVSVLVGGGI GVTPFASILK DLVFKSSVSC QVFCKKIYFI WVTTRTQRQFE WLADIIREVE
 1441 ENDHQDLVSV HIYITQLAEK FDLRTTMLYI CERHFQKVLN RSLFTGLRSI THFGRPPFEP
 1501 FFNSLQEVHP QVRKIGVFSC GPPGMTKNVE KACQLINRQD RTHFSHHYEN F.

WO 00/28031

PCT/US99/26592

<210> 47

WO 00/28031

PCT/US99/26592

<211> 3453
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (438) .. (3134)

<400> 47

```

gtcctcgacc agtttgtacg gctgcgggat ggtgaccgct actggtttga gaacaccagg 60
aatgggctgt tctccaagaa ggagattgag acatccgaaa taccaccgtg cgggacgtgc 120
tggtcgctgt tatcaacatt gaccccagtg ccctgcagcc caatgtcttt gtctggcata 180
aagggtgcacc ctgccctcaa cctaagcagc tcacaactga cggcctgccc cagtgtgcac 240
ccctgactgt gcttgacttc tttgaaggca gcagccctgg ttttgccatc accatcattg 300
ctctctgctg ccttccttta gtgagtctgc ttctctctgg agtgggtggcc tatttccggg 360
gccgagaaca caagaagcta caaaagaaac tcaaagagag cgtgaagaag gaagcagcca 420
aagatggagt gccagcg atg gag tgg cca ggc ccc aag gag agg agc agt      470
          Met Glu Trp Pro Gly Pro Lys Glu Arg Ser Ser
              1              5              10
ccc atc atc atc cag ctg ctg tca gac agg tgt ctg cag gtc ctg aac      518
Pro Ile Ile Ile Gln Leu Leu Ser Asp Arg Cys Leu Gln Val Leu Asn
          15              20              25
agg cat ctc act gtg ctc cgt gtg gtc cag ctg cag cct ctg cag cag      566
Arg His Leu Thr Val Leu Arg Val Val Gln Leu Gln Pro Leu Gln Gln
          30              35              40
gtc aac ctc atc ctg tcc aac aac cga gga tgc cgc acc ctg ctg ctc      614
Val Asn Leu Ile Leu Ser Asn Asn Arg Gly Cys Arg Thr Leu Leu Leu
          45              50              55
aag atc cct aag gag tat gac ctg gtg ctg ctg ttt agt tct gaa gag      662
Lys Ile Pro Lys Glu Tyr Asp Leu Val Leu Leu Phe Ser Ser Glu Glu
          60              65              70              75
gaa cgg ggc gcc ttt gtg cag cag cta tgg gac ttc tgc gtg cgc tgg      710
Glu Arg Gly Ala Phe Val Gln Gln Leu Trp Asp Phe Cys Val Arg Trp
          80              85              90
gct ctg ggc ctc cat gtg gct gag atg agc gag aag gag cta ttt agg      758

```

WO 00/28031

PCT/US99/26592

Ala	Leu	Gly	Leu	His	Val	Ala	Glu	Met	Ser	Glu	Lys	Glu	Leu	Phe	Arg		
			95					100					105				
aag	gct	gtg	aca	aag	cag	cag	cgg	gaa	cgc	atc	ctg	gag	atc	ttc	ttc	806	
Lys	Ala	Val	Thr	Lys	Gln	Gln	Arg	Glu	Arg	Ile	Leu	Glu	Ile	Phe	Phe		
		110					115					120					
aga	cac	ctt	ttt	gct	cag	gtg	ctg	gac	atc	aac	cag	gcc	gac	gca	ggg	854	
Arg	His	Leu	Phe	Ala	Gln	Val	Leu	Asp	Ile	Asn	Gln	Ala	Asp	Ala	Gly		
		125					130					135					
acc	ctg	ccc	ctg	gac	tcc	tcc	cag	aag	gtg	cgg	gag	gcc	ctg	acc	tgc	902	
Thr	Leu	Pro	Leu	Asp	Ser	Ser	Gln	Lys	Val	Arg	Glu	Ala	Leu	Thr	Cys		
		140				145				150					155		
gag	ctg	agc	agg	gcc	gag	ttt	gcc	gag	tcc	ctg	ggc	ctc	aag	ccc	cag	950	
Glu	Leu	Ser	Arg	Ala	Glu	Phe	Ala	Glu	Ser	Leu	Gly	Leu	Lys	Pro	Gln		
				160					165					170			
gac	atg	ttt	gtg	gag	tcc	atg	ttc	tct	ctg	gct	gac	aag	gat	ggc	aat	998	
Asp	Met	Phe	Val	Glu	Ser	Met	Phe	Ser	Leu	Ala	Asp	Lys	Asp	Gly	Asn		
			175					180						185			
ggc	tac	ctg	tcc	ttc	cga	gag	ttc	ctg	gac	atc	ctg	gtg	gtc	ttc	atg	1046	
Gly	Tyr	Leu	Ser	Phe	Arg	Glu	Phe	Leu	Asp	Ile	Leu	Val	Val	Phe	Met		
		190					195					200					
aaa	ggc	tcc	cca	gag	gat	aag	tcc	cgt	cta	atg	ttt	acc	atg	tat	gac	1094	
Lys	Gly	Ser	Pro	Glu	Asp	Lys	Ser	Arg	Leu	Met	Phe	Thr	Met	Tyr	Asp		
		205					210				215						
ctg	gat	gag	aat	ggc	ttc	ctc	tcc	aag	gac	gaa	ttc	ttc	acc	atg	atg	1142	
Leu	Asp	Glu	Asn	Gly	Phe	Leu	Ser	Lys	Asp	Glu	Phe	Phe	Thr	Met	Met		
		220				225				230					235		
cga	tcc	ttc	atc	gag	atc	tcc	aac	aac	tgc	ctg	tcc	aag	gcc	cag	ctg	1190	
Arg	Ser	Phe	Ile	Glu	Ile	Ser	Asn	Asn	Cys	Leu	Ser	Lys	Ala	Gln	Leu		
				240					245					250			
gcc	gag	gtg	gtg	gag	tct	atg	ttc	cgg	gag	tcg	gga	ttc	cag	gac	aag	1238	
Ala	Glu	Val	Val	Glu	Ser	Met	Phe	Arg	Glu	Ser	Gly	Phe	Gln	Asp	Lys		
				255				260					265				
gag	gag	ctg	aca	tgg	gag	gat	ttt	cac	ttc	atg	ctg	cgg	gac	cat	gac	1286	
Glu	Glu	Leu	Thr	Trp	Glu	Asp	Phe	His	Phe	Met	Leu	Arg	Asp	His	Asp		
			270				275					280					
agc	gag	ctc	cgc	ttc	acg	cag	ctc	tgt	gtc	aaa	ggg	gga	ggg	gga	ggg	1334	

WO 00/28031

PCT/US99/26592

Ser Glu Leu Arg Phe Thr Gln Leu Cys Val Lys Gly Gly Gly Gly Gly	
285	290 295
gga aat ggt att aga gat atc ttt aaa caa aac atc agc tgt cga gtc	1382
Gly Asn Gly Ile Arg Asp Ile Phe Lys Gln Asn Ile Ser Cys Arg Val	
300	305 310 315
tcg ttc atc act cgg aca cct ggg gag cgc tcc cac ccc cag gga ctg	1430
Ser Phe Ile Thr Arg Thr Pro Gly Glu Arg Ser His Pro Gln Gly Leu	
	320 325 330
ggg ccc cct gtc cca gaa gcc cca gag ctg gga ggc cct gga ctg aag	1478
Gly Pro Pro Val Pro Glu Ala Pro Glu Leu Gly Gly Pro Gly Leu Lys	
	335 340 345
aag agg ttt ggc aaa aag gca gca gtg ccc act ccc cgg ctg tac aca	1526
Lys Arg Phe Gly Lys Lys Ala Ala Val Pro Thr Pro Arg Leu Tyr Thr	
	350 355 360
gag gcg ctg caa gag aag atg cag cga ggc ttc cta gcc caa aag ctg	1574
Glu Ala Leu Gln Glu Lys Met Gln Arg Gly Phe Leu Ala Gln Lys Leu	
	365 370 375
cag cag tac aag cgc ttc gtg gag aac tac cgg agg cac atc gtg tgt	1622
Gln Gln Tyr Lys Arg Phe Val Glu Asn Tyr Arg Arg His Ile Val Cys	
380	385 390 395
gtg gca atc ttc tcg gcc atc tgt gtt ggc gtg ttt gca gat cgt gct	1670
Val Ala Ile Phe Ser Ala Ile Cys Val Gly Val Phe Ala Asp Arg Ala	
	400 405 410
tac tac tat ggc ttt gcc ttg cca ccc tcg gac att gca cag acc acc	1718
Tyr Tyr Tyr Gly Phe Ala Leu Pro Pro Ser Asp Ile Ala Gln Thr Thr	
	415 420 425
ctc gtg ggc atc atc ctg tca cga ggc acg gcg gcc agc gtc tcc ttc	1766
Leu Val Gly Ile Ile Leu Ser Arg Gly Thr Ala Ala Ser Val Ser Phe	
	430 435 440
atg ttc tct tat atc ttg ctc acc atg tgc cgc aac ctc ata acc ttc	1814
Met Phe Ser Tyr Ile Leu Leu Thr Met Cys Arg Asn Leu Ile Thr Phe	
	445 450 455
ctg cga gag act ttc ctc aac cgc tat gtg cct ttt gat gcc gca gtg	1862
Leu Arg Glu Thr Phe Leu Asn Arg Tyr Val Pro Phe Asp Ala Ala Val	
460	465 470 475
gac ttc cac cgc tgg atc gcc atg gct gct gtt gtc ctg gcc att ttg	1910

WO 00/28031

PCT/US99/26592

Asp Phe His Arg Trp Ile Ala Met Ala Ala Val Val Leu Ala Ile Leu	
480	485 490
cac agt gct ggc cac gca gtc aat gtc tac atc ttc tca gtc agc cca	1958
His Ser Ala Gly His Ala Val Asn Val Tyr Ile Phe Ser Val Ser Pro	
495	500 505
ctc agc ctg ctg gcc tgc ata ttc ccc aac gtc ttt gtg aat gat ggg	2006
Leu Ser Leu Leu Ala Cys Ile Phe Pro Asn Val Phe Val Asn Asp Gly	
510	515 520
tcc aag ctt ccc cag aag ttc tat tgg tgg ttc ttc cag acc gtc cca	2054
Ser Lys Leu Pro Gln Lys Phe Tyr Trp Trp Phe Phe Gln Thr Val Pro	
525	530 535
ggt atg aca ggt gtg ctt ctg ctc ctg gtc ctg gcc atc atg tat gtc	2102
Gly Met Thr Gly Val Leu Leu Leu Val Leu Ala Ile Met Tyr Val	
540	545 550 555
ttc gcc tcc cac cac ttc cgc cgc cgc agc ttc cgg ggc ttc tgg ctg	2150
Phe Ala Ser His His Phe Arg Arg Arg Ser Phe Arg Gly Phe Trp Leu	
560	565 570
acc cac cac ctc tac atc ctg ctc tat gcc ctg ctc atc atc cat ggc	2198
Thr His His Leu Tyr Ile Leu Leu Tyr Ala Leu Leu Ile Ile His Gly	
575	580 585
agc tat gct ctg atc cag ctg ccc act ttc cac atc tac ttc ctg gtc	2246
Ser Tyr Ala Leu Ile Gln Leu Pro Thr Phe His Ile Tyr Phe Leu Val	
590	595 600
ccg gca atc atc tat gga ggt gac aag ctg gtg agc ctg agc cgg aag	2294
Pro Ala Ile Ile Tyr Gly Gly Asp Lys Leu Val Ser Leu Ser Arg Lys	
605	610 615
aag gtg gag atc agc gtg gtg aag gcg gag ctg ctg ccc tca gga gtg	2342
Lys Val Glu Ile Ser Val Val Lys Ala Glu Leu Leu Pro Ser Gly Val	
620	625 630 635
acc tac ctg caa ttc cag agg ccc caa ggc ttt gag tac aag tca gga	2390
Thr Tyr Leu Gln Phe Gln Arg Pro Gln Gly Phe Glu Tyr Lys Ser Gly	
640	645 650
cag tgg gtg cgg atc gcc tgc ctg gct ctg ggg acc acc gag tac cac	2438
Gln Trp Val Arg Ile Ala Cys Leu Ala Leu Gly Thr Thr Glu Tyr His	
655	660 665
ccc ttc aca ctg acc tcc gcg ccc cat gag gac aca ctc agc ctg cac	2486

WO 00/28031

PCT/US99/26592

Pro Phe Thr Leu Thr Ser Ala Pro His Glu Asp Thr Leu Ser Leu His	
670 675 680	
atc cgg gca gtg ggg ccc tgg acc act cgc ctc agg gag atc tac tca	2534
Ile Arg Ala Val Gly Pro Trp Thr Thr Arg Leu Arg Glu Ile Tyr Ser	
685 690 695	
tcc cca aag ggc aat ggc tgt gct gga tac cca aag ctg tac ctt gat	2582
Ser Pro Lys Gly Asn Gly Cys Ala Gly Tyr Pro Lys Leu Tyr Leu Asp	
700 705 710 715	
gga ccg ttt gga gag ggc cat cag gag tgg cat aaa ttt gag gtg tca	2630
Gly Pro Phe Gly Glu Gly His Gln Glu Trp His Lys Phe Glu Val Ser	
720 725 730	
gtg ttg gtg gga ggg ggc att ggg gtc acc ccc ttt gcc tcc atc ctc	2678
Val Leu Val Gly Gly Gly Ile Gly Val Thr Pro Phe Ala Ser Ile Leu	
735 740 745	
aaa gac ctg gtc ttc aag tca tcc ttg ggc agc caa atg ctg tgt aag	2726
Lys Asp Leu Val Phe Lys Ser Ser Leu Gly Ser Gln Met Leu Cys Lys	
750 755 760	
aag atc tac ttc atc tgg gtg aca cgg acc cag cgt cag ttt gag tgg	2774
Lys Ile Tyr Phe Ile Trp Val Thr Arg Thr Gln Arg Gln Phe Glu Trp	
765 770 775	
ctg gct gac atc atc caa gag gtg gag gag aac gac cac cag gac ctg	2822
Leu Ala Asp Ile Ile Gln Glu Val Glu Glu Asn Asp His Gln Asp Leu	
780 785 790 795	
gtg tct gtg cac att tat gtc acc cag ctg gct gag aag ttc gac ctc	2870
Val Ser Val His Ile Tyr Val Thr Gln Leu Ala Glu Lys Phe Asp Leu	
800 805 810	
agg acc acc atg cta tac atc tgc gag cgg cac ttc cag aaa gtg ctg	2918
Arg Thr Thr Met Leu Tyr Ile Cys Glu Arg His Phe Gln Lys Val Leu	
815 820 825	
aac cgg agt ctg ttc acg ggc ctg cgc tcc atc acc cac ttt ggc cgt	2966
Asn Arg Ser Leu Phe Thr Gly Leu Arg Ser Ile Thr His Phe Gly Arg	
830 835 840	
ccc ccc ttc gag ccc ttc ttc aac tcc ctg cag gag gtc cac cca cag	3014
Pro Pro Phe Glu Pro Phe Phe Asn Ser Leu Gln Glu Val His Pro Gln	
845 850 855	
gtg cgc aag atc ggg gtg ttc agc tgc ggc cct cca gga atg acc aag	3062

WO 00/28031

PCT/US99/26592

Val Arg Lys Ile Gly Val Phe Ser Cys Gly Pro Pro Gly Met Thr Lys
 860 865 870 875

aat gta gag aag gcc tgt cag ctc gtc aac agg cag gac cga gcc cac 3110
 Asn Val Glu Lys Ala Cys Gln Leu Val Asn Arg Gln Asp Arg Ala His
 880 885 890

ttc atg cac cac tat gag aac ttc tgagcctgtc ctccctggct gctgcttcca 3164
 Phe Met His His Tyr Glu Asn Phe
 895

gtatcctgcc ttctcttctg tgcacctaag ttgcccagcc ctgctggcaa tctctccatc 3224

agaatccacc ttaggcctca gctggagggc tgcagagccc ctcccaatat tgggagaata 3284

ttgaccaga caattatata aatgagaaaa ggcattaaaa ttacgtttc tgatgatggc 3344

aaagctcatt tttctattag taactctgct gaagatccat ttattgcaat tcatgctgaa 3404

tctaaattgt aaaatttaaa attaaatgca tgtcctcaaa aaaaaaaaaa 3453

<210> 48

<211> 899

<212> PRT

<213> Homo sapiens

<400> 48

Met Glu Trp Pro Gly Pro Lys Glu Arg Ser Ser Pro Ile Ile Ile Gln
 1 5 10 15

Leu Leu Ser Asp Arg Cys Leu Gln Val Leu Asn Arg His Leu Thr Val
 20 25 30

Leu Arg Val Val Gln Leu Gln Pro Leu Gln Gln Val Asn Leu Ile Leu
 35 40 45

Ser Asn Asn Arg Gly Cys Arg Thr Leu Leu Leu Lys Ile Pro Lys Glu
 50 55 60

Tyr Asp Leu Val Leu Leu Phe Ser Ser Glu Glu Glu Arg Gly Ala Phe
 65 70 75 80

Val Gln Gln Leu Trp Asp Phe Cys Val Arg Trp Ala Leu Gly Leu His
 85 90 95

Val Ala Glu Met Ser Glu Lys Glu Leu Phe Arg Lys Ala Val Thr Lys
 100 105 110

WO 00/28031

PCT/US99/26592

Gln Gln Arg Glu Arg Ile Leu Glu Ile Phe Phe Arg His Leu Phe Ala
 115 120 125

Gln Val Leu Asp Ile Asn Gln Ala Asp Ala Gly Thr Leu Pro Leu Asp
 130 135 140

Ser Ser Gln Lys Val Arg Glu Ala Leu Thr Cys Glu Leu Ser Arg Ala
 145 150 155 160

Glu Phe Ala Glu Ser Leu Gly Leu Lys Pro Gln Asp Met Phe Val Glu
 165 170 175

Ser Met Phe Ser Leu Ala Asp Lys Asp Gly Asn Gly Tyr Leu Ser Phe
 180 185 190

Arg Glu Phe Leu Asp Ile Leu Val Val Phe Met Lys Gly Ser Pro Glu
 195 200 205

Asp Lys Ser Arg Leu Met Phe Thr Met Tyr Asp Leu Asp Glu Asn Gly
 210 215 220

Phe Leu Ser Lys Asp Glu Phe Phe Thr Met Met Arg Ser Phe Ile Glu
 225 230 235 240

Ile Ser Asn Asn Cys Leu Ser Lys Ala Gln Leu Ala Glu Val Val Glu
 245 250 255

Ser Met Phe Arg Glu Ser Gly Phe Gln Asp Lys Glu Glu Leu Thr Trp
 260 265 270

Glu Asp Phe His Phe Met Leu Arg Asp His Asp Ser Glu Leu Arg Phe
 275 280 285

Thr Gln Leu Cys Val Lys Gly Gly Gly Gly Gly Gly Asn Gly Ile Arg
 290 295 300

Asp Ile Phe Lys Gln Asn Ile Ser Cys Arg Val Ser Phe Ile Thr Arg
 305 310 315 320

Thr Pro Gly Glu Arg Ser His Pro Gln Gly Leu Gly Pro Pro Val Pro
 325 330 335

Glu Ala Pro Glu Leu Gly Gly Pro Gly Leu Lys Lys Arg Phe Gly Lys
 340 345 350

Lys Ala Ala Val Pro Thr Pro Arg Leu Tyr Thr Glu Ala Leu Gln Glu
 355 360 365

WO 00/28031

PCT/US99/26592

Lys Met Gln Arg Gly Phe Leu Ala Gln Lys Leu Gln Gln Tyr Lys Arg
 370 375 380

Phe Val Glu Asn Tyr Arg Arg His Ile Val Cys Val Ala Ile Phe Ser
 385 390 395 400

Ala Ile Cys Val Gly Val Phe Ala Asp Arg Ala Tyr Tyr Tyr Gly Phe
 405 410 415

Ala Leu Pro Pro Ser Asp Ile Ala Gln Thr Thr Leu Val Gly Ile Ile
 420 425 430

Leu Ser Arg Gly Thr Ala Ala Ser Val Ser Phe Met Phe Ser Tyr Ile
 435 440 445

Leu Leu Thr Met Cys Arg Asn Leu Ile Thr Phe Leu Arg Glu Thr Phe
 450 455 460

Leu Asn Arg Tyr Val Pro Phe Asp Ala Ala Val Asp Phe His Arg Trp
 465 470 475 480

Ile Ala Met Ala Ala Val Val Leu Ala Ile Leu His Ser Ala Gly His
 485 490 495

Ala Val Asn Val Tyr Ile Phe Ser Val Ser Pro Leu Ser Leu Leu Ala
 500 505 510

Cys Ile Phe Pro Asn Val Phe Val Asn Asp Gly Ser Lys Leu Pro Gln
 515 520 525

Lys Phe Tyr Trp Trp Phe Phe Gln Thr Val Pro Gly Met Thr Gly Val
 530 535 540

Leu Leu Leu Leu Val Leu Ala Ile Met Tyr Val Phe Ala Ser His His
 545 550 555 560

Phe Arg Arg Arg Ser Phe Arg Gly Phe Trp Leu Thr His His Leu Tyr
 565 570 575

Ile Leu Leu Tyr Ala Leu Leu Ile Ile His Gly Ser Tyr Ala Leu Ile
 580 585 590

Gln Leu Pro Thr Phe His Ile Tyr Phe Leu Val Pro Ala Ile Ile Tyr
 595 600 605

Gly Gly Asp Lys Leu Val Ser Leu Ser Arg Lys Lys Val Glu Ile Ser
 610 615 620

WO 00/28031

PCT/US99/26592

Val	Val	Lys	Ala	Glu	Leu	Leu	Pro	Ser	Gly	Val	Thr	Tyr	Leu	Gln	Phe	625	630	635	640
Gln	Arg	Pro	Gln	Gly	Phe	Glu	Tyr	Lys	Ser	Gly	Gln	Trp	Val	Arg	Ile	645	650	655	
Ala	Cys	Leu	Ala	Leu	Gly	Thr	Thr	Glu	Tyr	His	Pro	Phe	Thr	Leu	Thr	660	665	670	
Ser	Ala	Pro	His	Glu	Asp	Thr	Leu	Ser	Leu	His	Ile	Arg	Ala	Val	Gly	675	680	685	
Pro	Trp	Thr	Thr	Arg	Leu	Arg	Glu	Ile	Tyr	Ser	Ser	Pro	Lys	Gly	Asn	690	695	700	
Gly	Cys	Ala	Gly	Tyr	Pro	Lys	Leu	Tyr	Leu	Asp	Gly	Pro	Phe	Gly	Glu	705	710	715	720
Gly	His	Gln	Glu	Trp	His	Lys	Phe	Glu	Val	Ser	Val	Leu	Val	Gly	Gly	725	730	735	
Gly	Ile	Gly	Val	Thr	Pro	Phe	Ala	Ser	Ile	Leu	Lys	Asp	Leu	Val	Phe	740	745	750	
Lys	Ser	Ser	Leu	Gly	Ser	Gln	Met	Leu	Cys	Lys	Lys	Ile	Tyr	Phe	Ile	755	760	765	
Trp	Val	Thr	Arg	Thr	Gln	Arg	Gln	Phe	Glu	Trp	Leu	Ala	Asp	Ile	Ile	770	775	780	
Gln	Glu	Val	Glu	Glu	Asn	Asp	His	Gln	Asp	Leu	Val	Ser	Val	His	Ile	785	790	795	800
Tyr	Val	Thr	Gln	Leu	Ala	Glu	Lys	Phe	Asp	Leu	Arg	Thr	Thr	Met	Leu	805	810	815	
Tyr	Ile	Cys	Glu	Arg	His	Phe	Gln	Lys	Val	Leu	Asn	Arg	Ser	Leu	Phe	820	825	830	
Thr	Gly	Leu	Arg	Ser	Ile	Thr	His	Phe	Gly	Arg	Pro	Pro	Phe	Glu	Pro	835	840	845	
Phe	Phe	Asn	Ser	Leu	Gln	Glu	Val	His	Pro	Gln	Val	Arg	Lys	Ile	Gly	850	855	860	
Val	Phe	Ser	Cys	Gly	Pro	Pro	Gly	Met	Thr	Lys	Asn	Val	Glu	Lys	Ala	865	870	875	880

WO 00/28031

PCT/US99/26592

Cys Gln Leu Val Asn Arg Gln Asp Arg Ala His Phe Met His His Tyr
 885 890 895

Glu Asn Phe

<210> 49

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 49

cctgacagat gtatttcact acccag

26

<210> 50

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 50

ggatcggagt cactcccttc gctg

24

<210> 51

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 51

ctagaagctc tccttggtgt aataga

26

<210> 52

<211> 24

<212> DNA

<213> Artificial Sequence

WO 00/28031

PCT/US99/26592

<220>

<223> Description of Artificial Sequence: Primer

<400> 52

atgaacacct ctggggtcag ctga

24

<210> 53

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 53

atgaacacct ctggggtcag ctga

24

<210> 54

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 54

gtcctctgca gcattgttcc tctta

25

<210> 55

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 55

cctgacagat gtatttcact acccag

26

<210> 56

<211> 24

<212> DNA

<213> Artificial Sequence

WO 00/28031

PCT/US99/26592

<220>

<223> Description of Artificial Sequence: Primer

<400> 56

ggatcggagt cactcccttc gctg

24

<210> 57

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 57

aatgacactg tactggaggc cacag

25

<210> 58

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 58

ctgccatcta ccacacggat ctgc

24

<210> 59

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 59

cttgccattc caaagcttcc atgc

24

<210> 60

<211> 24

<212> DNA

<213> Artificial Sequence

WO 00/28031

PCT/US99/26592

<220>

<223> Description of Artificial Sequence: Primer

<400> 60

gtacaagtca ggacagtggg tgcg

24

<210> 61

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 61

tggatgatgt cagccagcca ctca

24

1. A protein capable of stimulating superoxide production, wherein the protein comprises the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, a fragment thereof, or a conservative substitution thereof, wherein the protein, the fragment thereof or the conservative substitution thereof is capable of generating reactive oxygen intermediates.
2. A protein capable of stimulating superoxide production, wherein the protein comprises the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, or a conservative substitution thereof.
3. A protein capable of stimulating superoxide production, wherein the protein comprises the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48, a deletion thereof or an addition thereto of no more than about 5% of the amino acid sequence, or a conservative substitution thereof, wherein the conservative substitution comprises substitution of
 - a) alanine, serine, or threonine for each other;
 - b) aspartic acid or glutamic acid for each other;
 - c) asparagine or glutamine for each other;
 - d) arginine or lysine for each other;
 - e) isoleucine, leucine, methionine, or valine for each other; and
 - f) phenylalanine, tyrosine, or tryptophan for each other.
4. A protein comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48.
5. A nucleotide sequence encoding for the protein, the fragment thereof or the conservative substitution thereof, as recited in any one of Claims 1 to 4.

6. The nucleotide sequence of Claim 5, wherein the nucleotide sequence comprises SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, or SEQ ID NO:47.
7. A nucleotide sequence comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, or SEQ ID NO:47.
8. A vector, wherein the vector comprises a nucleotide sequence encoding for the protein, the fragment thereof or the conservative substitution thereof, as recited in any one of Claims 1 to 4.
9. The vector of Claim 8 wherein the nucleotide sequence comprises SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, or SEQ ID NO:47.
10. A vector, wherein the vector comprises a nucleotide sequence comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:22, SEQ ID NO:41, SEQ ID NO:45, or SEQ ID NO:47.
11. A cell containing the vector of Claim 8.
12. A cell containing the vector of Claim 10.
13. An antibody generated against the protein of any one of Claims 1 to 4.
14. An antibody generated against the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:21, SEQ ID NO:42, SEQ ID NO:46, or SEQ ID NO:48.
15. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the protein of any one of Claims 1 to 4 in a pharmaceutically acceptable carrier.

16. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the protein of Claim 4 in a pharmaceutically acceptable carrier.
17. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the vector of Claim 8 in a pharmaceutically acceptable carrier.
18. A method of stimulating superoxide formation comprising administration, in vitro or in vivo, of a composition comprising the vector of Claim 9 in a pharmaceutically acceptable carrier.
19. A method for determining activity of a drug or a chemical comprising measuring enzymatic activity of the protein, the fragment thereof or the conservative substitution thereof, as recited in any one of Claims 1 to 4, to stimulate superoxide production following administration of the drug or the chemical.
20. A method for determining activity of a drug or a chemical comprising measuring enzymatic activity of the protein recited in Claim 4 to stimulate superoxide production following administration of the drug or the chemical.
21. The method of Claim 19, wherein the enzymatic activity is NADPH- dependent or NADH-dependent superoxide generation.
22. The method of Claim 19, wherein the enzymatic activity is NADPH- dependent or NADH-dependent diaphorase activity.
23. A method for determining activity of a drug or a chemical comprising measuring binding of the drug or the chemical to the protein, the fragment thereof or the conservative substitution thereof, as recited in any one of Claims 1 to 4.

24. A method for determining activity of a drug or a chemical comprising measuring binding of the drug or the chemical to the protein recited in Claim 4.

25. A method for determining activity of a drug or a chemical comprising measuring the activity of the drug or the chemical to modulate proliferative activity of the protein, the fragment thereof or the conservative substitution thereof, as recited in any one of Claims 1 to 4.

26. A method for determining activity of a drug or a chemical comprising measuring the activity of the drug or the chemical to modulate proliferative activity of the protein recited in Claim 4.

27. The method of Claim 19, wherein the enzymatic activity is assessed using intact cells, transfected cells or cell lysates thereof.

28. The method of Claim 20, wherein the enzymatic activity is assessed using intact cells, transfected cells or cell lysates thereof.

1	- - G N W A V - - - - -	N E G	gp91phox.human
1	M - G N W V V - - - - -	N E G	gp91phox.bovine
1	M - G N W A V - - - - -	N E G	gp91phox.mouse
1	M - G N W V V - - - - -	N H W	Mox1.human
1	M - G N W L V - - - - -	N H W	Mox1.rat
1	M M G C W I L - - - - -	N E G	mox2.human
1	M L S Q K L I P T K D R N P V K R F A M N I S Y F F L E N W		cytb558/arabidopsis.pep
1	- - - - N L A G L R K K S S I R K I S T S L S Y Y F E D N W		cytb558.rice.pep
9	L S I F A I L V W L G L N V F L F V - W Y Y R V Y D I P P K		gp91phox.human
10	I S I F V I L V W L G M N V F L F V - W Y Y R V Y D I P D K		gp91phox.bovine
10	L S I F V I L V W L G L N V F L F I - N Y Y K V Y D D G P K		gp91phox.mouse
10	F S V L F L V V W L G L N V F L F V D A F L K - Y E K A D K		Mox1.human
10	L S V L F L V S W L G L N I F L F V Y V F L N - Y E K S D K		Mox1.rat
11	L S T I L V L S W L G I N F Y L F I D T F Y W - Y E E E E S		mox2.human
31	K R I W V L T L W I S I C I T L F T W K F L Q - Y K R K T V		cytb558/arabidopsis.pep
27	K R L W V L A L W I G I M A G L F T W K F M O - Y R N R Y V		cytb558.rice.pep
38	F F Y T R K L L G S A L A L A R A P A A C L N F N C M L I L		gp91phox.human
39	F F Y T R K L L G S A L A L A R A P A A C L N F N C M L I L		gp91phox.bovine
39	Y N Y T R K L L G S A L A L A R A P A A C L N F N C M L I L		gp91phox.mouse
39	Y Y Y T R K I L G S T L A C A R A S A L C L N F N S T L I L		Mox1.human
39	Y Y Y T R E I L G T A L A L A R A S A L C L N F N S M V I L		Mox1.rat
40	F H Y T R V I L G S T L A W A R A S A L C L N F N C M L I L		mox2.human
60	F - - - - E V M G Y C V T V A K G S A E T L K E N M A L I L		cytb558/arabidopsis.pep
56	F - - - - D V M G Y C V T T A K G A A E T L K L N M A I I L		cytb558.rice.pep
68	L P V C R N L L S F L R G S S A C C S T R V R R Q L D R N L		gp91phox.human
69	L P V C R N L L S F L R G S S A C C S T R I R R Q L D R N L		gp91phox.bovine
69	L P V C R N L L S F L R G S S A C C S T R I R R Q L D R N L		gp91phox.mouse
69	L P V C R N L L S F L R G T C S F C S R T L R K Q L D H N L		Mox1.human
69	I P V C R N L L S F L R G T C S F C N H T L R K P L D H N L		Mox1.rat
70	I P V S R N L I S F I R G T S I C C R G P W R R O L D K N L		mox2.human
86	L P V C R N T I T W L R T K S K L I G S V V - - P F D D N I		cytb558/arabidopsis.pep
82	L P V C R N T I T W L R S T R A - - A R A L - - P F D D N I		cytb558.rice.pep
98	T F H K M V A W M I A L H S A I H T I A H L F - N V E W C V		gp91phox.human
99	T F H K M V A W M I A L H T A I H T I A H L F - N V E W C V		gp91phox.bovine
99	T F H K M V A W M I A L H T A I H T I A H L F - N V E W C V		gp91phox.mouse
99	T F H K L V A Y M I C L H T A I H I A H L F - N F D C Y S		Mox1.human
99	T F H K L V A Y M I C I F T A I H I A H L F - N F E R Y S		Mox1.rat
100	R F H K L V A Y G I A V N A T I H I V A H F F - N L E R Y H		mox2.human
114	N F H K V V A F G I A V G I G L H A I S H L A C D F P R L L		cytb558/arabidopsis.pep
108	N F H K T I A A A I V V G I I L H A G N H L V C D F P R L I		cytb558.rice.pep
127	N A R V N N S D P Y S V A L S E L - - G D R O N E - S Y L N		gp91phox.human
128	N A R V N N S D P Y S I A L S D I - - G D K P N E - T Y L N		gp91phox.bovine
128	N A R V G I S D R Y S I A L S D I - - G D N E N E - E Y L N		gp91phox.mouse
128	R S R Q A T D G S L A S I L S S L S H D E K K G G - S W L N		Mox1.human
128	R S Q Q A M D G S L A S V L S S L F H P E K E D - - S W L N		Mox1.rat
129	W S Q S E E A Q G L L A A L S K L - - G N T P N E - S Y L N		mox2.human
144	H A K N V E F E P - - - - M K K F F G D E R P E N Y G W F -		cytb558/arabidopsis.pep
138	K S S D E K Y A P - - - - L G Q Y F G E I K P T Y F T - L -		cytb558.rice.pep

FIGURE 1(a)

154	F	A	R	K	R	I	K	N	P	E	G	G	L	Y	L	A	V	T	L	L	A	G	I	T	G	V	V	I	T	L		gp91phox.human		
155	F	V	R	Q	R	I	K	N	P	E	G	G	L	Y	V	A	V	T	R	L	A	G	I	T	G	V	V	I	T	L		gp91phox.bovine		
155	F	A	R	E	K	I	K	N	P	E	G	G	L	Y	V	A	V	T	R	L	A	G	I	T	G	I	V	I	T	L		gp91phox.mouse		
157	P	I	Q	S	R	-	-	-	N	T	T	V	E	Y	V	T	F	T	S	V	A	G	L	T	G	V	I	M	T	I		Mox1.human		
156	P	I	Q	S	P	-	-	-	N	V	T	V	M	Y	A	A	F	T	S	I	A	G	L	T	G	V	V	A	T	V		Mox1.rat		
156	P	V	R	T	F	P	T	N	T	T	T	E	L	L	R	T	I	-	-	-	A	G	V	T	G	L	V	I	S	L		mox2.human		
169	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	M	K	G	T	D	G	W	T	G	V	T	M	V		cytb558/arabidopsis.pep
162	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	V	K	G	V	E	G	I	T	G	V	I	M	V		cytb558.rice.pep
184	C	L	I	L	I	T	S	S	T	K	T	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	gp91phox.human	
185	C	L	I	L	I	T	S	S	T	K	T	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	gp91phox.bovine
185	C	L	I	L	I	T	S	S	T	K	T	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	gp91phox.mouse
184	A	L	I	L	M	V	T	S	A	T	E	F	I	R	R	S	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	Mox1.human
183	A	L	V	L	M	V	T	S	A	M	E	F	I	R	R	N	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	F	Mox1.rat
183	A	L	V	L	I	M	T	S	S	T	E	F	I	R	Q	A	S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Y	mox2.human
183	L	M	L	V	A	Y	V	L	A	Q	S	W	F	R	R	N	R	A	N	L	P	K	S	L	K	R	L	T	G	F			cytb558/arabidopsis.pep	
176	C	M	I	I	A	F	T	L	A	T	R	W	F	R	R	S	L	V	K	L	P	R	P	F	D	K	L	T	G	F			cytb558.rice.pep	
202	E	V	F	W	Y	T	H	H	L	F	V	I	F	F	I	G	L	A	I	H	G	A	E	R	I	V	R	G	Q	T			gp91phox.human	
203	E	V	F	W	Y	T	H	H	L	F	V	I	F	F	I	G	L	A	I	H	G	A	Q	R	I	V	R	G	Q	T			gp91phox.bovine	
203	E	V	F	W	Y	T	H	H	L	F	V	I	F	F	I	G	L	A	I	H	G	A	E	R	I	V	R	G	Q	T			gp91phox.mouse	
202	E	V	F	W	Y	T	H	H	L	F	I	F	Y	I	L	G	L	G	I	H	G	I	G	G	I	V	R	G	Q	T			Mox1.human	
201	E	L	F	W	Y	T	H	H	L	F	I	I	Y	I	I	C	L	G	I	H	G	L	G	G	I	V	R	G	Q	T			Mox1.rat	
201	E	L	F	W	Y	T	H	H	V	F	I	V	F	F	L	S	L	A	I	H	G	T	G	R	I	V	R	G	Q	T			mox2.human	
213	N	A	F	W	Y	S	H	H	L	F	V	I	V	Y	V	L	L	I	V	H	G	-	-	-	-	-	-	-	-	-	-	-		cytb558/arabidopsis.pep
206	N	A	F	W	Y	S	H	H	L	F	I	I	V	Y	I	A	L	I	V	H	G	-	-	-	-	-	-	-	-	-	-	-		cytb558.rice.pep
232	A	E	S	L	A	V	H	N	I	T	V	C	E	Q	K	I	S	E	W	G	K	I	K	-	E	C	P	I	P	O			gp91phox.human	
233	A	E	S	L	L	K	H	Q	P	R	N	C	Y	Q	N	I	S	Q	W	G	K	I	E	-	N	C	P	I	P	E			gp91phox.bovine	
233	A	E	S	L	E	E	H	N	L	D	I	C	A	D	K	I	E	E	W	G	K	I	K	-	E	C	P	V	P	K			gp91phox.mouse	
232	E	E	S	M	N	E	S	H	P	R	K	C	A	E	S	F	E	M	W	D	D	R	D	S	H	C	R	R	P	K			Mox1.human	
231	E	E	S	M	S	E	S	H	P	R	N	C	S	Y	S	F	H	E	W	D	K	Y	E	R	S	C	R	S	P	H			Mox1.rat	
231	Q	D	S	L	S	L	H	N	I	T	F	C	R	D	R	Y	A	E	W	Q	T	V	A	-	Q	C	P	V	P	O			mox2.human	
234	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		cytb558/arabidopsis.pep
227	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		cytb558.rice.pep
261	F	A	G	N	P	P	M	T	W	K	W	I	V	G	P	M	F	L	Y	L	C	E	R	L	V	R	F	W	R	S			gp91phox.human	
262	F	S	G	N	P	P	M	T	W	K	W	I	V	G	P	M	F	L	Y	L	C	E	R	L	V	R	F	W	R	S			gp91phox.bovine	
262	F	A	G	N	P	P	M	T	W	K	W	I	V	G	P	M	F	L	Y	L	C	E	R	L	V	R	F	W	R	S			gp91phox.mouse	
262	F	E	G	H	P	P	E	S	W	K	W	I	L	A	P	V	I	L	Y	I	C	E	R	I	L	R	F	Y	R	S			Mox1.human	
261	F	V	G	Q	P	P	E	S	W	K	W	I	L	A	P	I	A	F	Y	I	F	E	R	I	L	R	F	Y	R	S			Mox1.rat	
260	F	S	G	K	E	P	S	A	W	K	W	I	L	G	P	V	V	L	Y	A	C	E	R	I	I	R	F	W	R	F			mox2.human	
243	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		cytb558/arabidopsis.pep
236	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		cytb558.rice.pep
291	Q	Q	K	V	V	-	I	T	K	V	V	T	H	P	F	K	T	I	E	L	Q	M	K	K	-	K	G	F	K	M			gp91phox.human	
292	Q	Q	K	V	V	-	I	T	K	V	V	T	H	P	F	K	T	I	E	L	Q	M	K	K	-	K	G	F	K	M			gp91phox.bovine	
292	Q	Q	K	V	V	-	I	T	K	V	V	T	H	P	F	K	T	I	E	L	Q	M	K	K	-	K	G	F	K	M			gp91phox.mouse	
292	Q	Q	K	V	V	-	I	T	K	V	V	M	H	P	S	K	V	L	E	L	Q	M	N	K	-	R	G	F	S	M			Mox1.human	
291	R	Q	K	V	V	-	I	T	K	V	V	M	H	P	C	K	V	L	E	L	Q	M	R	K	-	R	G	F	T	M			Mox1.rat	
290	Q	Q	E	V	V	-	I	T	K	V	V	S	H	P	S	G	V	L	E	L	H	M	K	K	-	R	G	F	K	M			mox2.human	
270	G	A	K	A	V	K	V	L	K	V	A	V	Y	P	G	N	V	L	S	L	Y	M	S	K	P	K	G	F	K	Y			cytb558/arabidopsis.pep	
263	G	S	Y	S	V	R	L	L	K	V	A	I	Y	P	G	N	V	L	T	L	Q	M	S	K	P	P	T	F	R	Y			cytb558.rice.pep	

FIGURE 1(b)

3/8

319	E	V	G	Q	Y	I	F	V	K	C	P	K	V	S	K	L	E	W	H	P	F	T	L	T	S	A	P	E	E	D	gp91phox.human			
320	E	V	G	Q	Y	I	F	V	K	C	P	V	V	S	K	L	E	W	H	P	F	T	L	T	S	A	P	E	E	D	gp91phox.bovine			
320	E	V	G	Q	Y	I	F	V	K	C	P	K	V	S	K	L	E	W	H	P	F	T	L	T	S	A	P	E	E	D	gp91phox.mouse			
320	E	V	G	Q	Y	I	F	V	N	C	P	S	I	S	L	L	E	W	H	P	F	T	L	T	S	A	P	E	E	D	Mox1.human			
319	G	I	G	Q	Y	I	F	V	N	C	P	S	I	S	F	L	E	W	H	P	F	T	L	T	S	A	P	E	E	E	Mox1.rat			
318	A	P	G	Q	Y	I	L	V	Q	C	P	A	I	S	S	L	E	W	H	P	F	T	L	T	S	A	P	Q	E	D	mox2.human			
300	T	S	G	Q	Y	I	Y	I	N	C	S	D	V	S	P	L	Q	W	H	P	F	S	I	T	S	A	S	G	D	D	cytb558/arabidopsis.pep			
293	K	S	G	Q	Y	M	F	V	Q	C	P	A	V	S	P	F	E	W	H	P	F	S	I	T	S	A	P	G	D	D	cytb558.rice.pep			
349	F	F	S	I	H	I	R	I	V	G	D	W	T	E	G	L	F	N	A	C	G	-	-	C	-	-	-	-	-	-	gp91phox.human			
350	F	F	S	I	H	I	R	I	V	G	D	W	T	E	G	L	F	K	A	C	G	-	-	C	-	-	-	-	-	-	gp91phox.bovine			
350	F	F	S	I	H	I	R	I	V	G	D	W	T	E	G	L	F	N	A	C	G	-	-	C	-	-	-	-	-	-	gp91phox.mouse			
350	F	F	S	I	H	I	R	A	A	G	D	W	T	E	N	L	I	R	A	F	E	-	-	-	-	-	-	-	-	-	Mox1.human			
349	F	F	S	I	H	I	R	A	A	G	D	W	T	E	N	L	I	R	T	F	E	-	-	-	-	-	-	-	-	-	Mox1.rat			
348	F	F	S	V	H	I	R	A	A	G	D	W	T	A	A	L	L	E	A	F	G	-	-	A	-	-	-	-	-	-	mox2.human			
330	Y	L	S	V	H	I	R	T	L	G	D	W	T	S	Q	L	K	S	L	Y	S	K	V	C	Q	L	P	S	T	S	cytb558/arabidopsis.pep			
323	Y	L	S	I	H	V	R	Q	L	G	D	W	T	R	E	L	K	R	V	F	A	A	A	C	E	P	P	A	G	G	cytb558.rice.pep			
371	-	-	-	-	-	-	-	-	-	D	K	Q	E	F	Q	D	A	W	K	L	P	K	I	A	V	D	G	P	F	G	T	A	gp91phox.human	
372	-	-	-	-	-	-	-	-	-	D	K	Q	E	F	Q	D	A	W	K	L	P	K	I	A	V	D	G	P	F	G	T	A	gp91phox.bovine	
372	-	-	-	-	-	-	-	-	-	D	K	Q	E	F	Q	D	A	W	K	L	P	K	I	A	V	D	G	P	F	G	T	A	gp91phox.mouse	
371	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Q	Q	Y	S	P	I	P	R	I	E	V	D	G	P	F	G	T	A	Mox1.human
370	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Q	Q	H	S	P	M	P	R	I	E	V	D	G	P	F	G	T	V	Mox1.rat
370	-	-	-	-	-	-	-	-	-	E	G	Q	A	L	Q	E	P	W	S	L	P	R	L	A	V	D	G	P	F	G	T	A	mox2.human	
360	Q	S	G	L	F	I	A	D	I	G	Q	A	N	N	I	T	R	F	P	R	L	L	I	D	G	P	Y	G	A	P	cytb558/arabidopsis.pep			
353	K	S	G	L	L	R	A	D	-	E	T	T	K	K	I	-	-	L	P	K	L	L	I	D	G	P	Y	G	S	P	cytb558.rice.pep			
394	S	E	D	V	F	S	Y	E	V	V	M	L	V	G	A	G	I	G	V	T	P	F	A	S	I	L	K	S	V	W	gp91phox.human			
395	S	E	D	V	F	S	Y	E	V	V	M	L	V	G	A	G	I	G	V	T	P	F	A	S	I	L	K	S	V	W	gp91phox.bovine			
395	S	E	D	V	F	S	Y	E	V	V	M	L	V	G	A	G	I	G	V	T	P	F	A	S	I	L	K	S	V	W	gp91phox.mouse			
389	S	E	D	V	F	Q	Y	E	V	A	V	L	V	G	A	G	I	G	V	T	P	F	A	S	I	L	K	S	I	W	Mox1.human			
388	S	E	D	V	F	Q	Y	E	V	A	V	L	V	G	A	G	I	G	V	T	P	F	A	S	F	L	K	S	I	W	Mox1.rat			
393	L	T	D	V	F	H	Y	P	V	C	V	C	V	A	A	G	I	G	V	T	P	F	A	A	L	L	K	S	I	W	mox2.human			
390	A	Q	D	Y	R	N	Y	D	V	L	L	V	G	L	G	I	G	A	T	P	L	I	S	I	I	R	D	V	L	cytb558/arabidopsis.pep				
380	A	Q	D	Y	S	K	Y	D	V	L	L	V	G	L	G	I	G	A	T	P	F	I	S	I	L	K	D	L	L	cytb558.rice.pep				
424	Y	K	Y	C	N	N	A	T	N	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	gp91phox.human			
425	Y	K	Y	C	N	K	A	P	N	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	gp91phox.bovine			
425	Y	K	Y	C	D	N	A	T	S	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	gp91phox.mouse			
419	Y	K	F	Q	C	A	D	H	N	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mox1.human			
418	Y	K	F	Q	R	A	H	N	K	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Mox1.rat			
423	Y	K	C	S	E	A	Q	T	P	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	mox2.human			
420	N	N	I	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	cytb558/arabidopsis.pep			
410	N	N	I	I	K	M	E	E	E	D	A	S	T	D	L	Y	P	P	M	G	R	N	N	P	H	V	D	L	G	cytb558.rice.pep				
433	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	L	K	L	K	K	I	Y	F	Y	W	L	C	R	D	T	H	A	F	gp91phox.human
434	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	L	R	L	K	K	I	Y	F	Y	W	L	C	R	D	T	H	A	F	gp91phox.bovine
434	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	L	K	L	K	K	I	Y	F	Y	W	L	C	R	D	T	H	A	F	gp91phox.mouse
428	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	L	K	T	K	K	I	Y	F	Y	W	I	C	R	E	T	G	A	F	Mox1.human
427	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	L	K	T	Q	K	I	Y	F	Y	W	I	C	R	E	T	G	A	F	Mox1.rat
432	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	L	K	L	S	K	V	Y	F	Y	W	I	C	R	D	A	R	A	F	mox2.human
432	T	N	Q	H	I	-	-	-	-	K	N	Y	V	A	T	K	R	A	Y	F	Y	W	V	T	R	E	Q	G	S	L	cytb558/arabidopsis.pep			
440	T	L	M	T	I	T	S	R	P	K	K	I	L	K	T	T	N	A	Y	F	Y	W	V	T	R	E	Q	G	S	F	cytb558.rice.pep			

FIGURE 1(c)

4/8

451	EWFADLLQLLESQMQRN	NAGFLSY	-	-	NIY	gp91phox.human			
452	EWFADLLQLLETQMQRN	NTDFLSY	-	-	NIC	gp91phox.bovine			
452	EWFADLLQLLETQMQRN	NANFLSY	-	-	NIY	gp91phox.mouse			
446	SWFNNLLTSL	EQEMEELGKVGFLNY	-	-	RLF	Mox1.human			
445	AWFNNLLNSL	EQEMDELGKPDFLNY	-	-	RLF	Mox1.rat			
450	EWFADLLLSL	ETRMSEQGKTHFLSY	-	-	HIF	mox2.human			
458	EWFSEVMNEV	-	AEYDSEGMIELHNY	CTSVY		cytb558/arabidopsis.pep			
470	DWFKGVMNEI	-	ADLDQRNIIEMHNY	LTSVY		cytb558.rice.pep			
479	LTG	-	WDESQANHF	AVHHDEEK	DVITGLK	gp91phox.human			
480	LTG	-	WDESQASHF	AMHHDEEK	DVITGLK	gp91phox.bovine			
480	LTG	-	WDESQANHF	AVHHDEEK	DVITGLK	gp91phox.mouse			
474	LTG	-	WDSNIVGH	AALNFDKAT	DIVTGLK	Mox1.human			
473	LTG	-	WDSNIAGH	AALNFDRA	T	DVITGLK	Mox1.human		
478	LTG	-	WDENOA	LHIALHWD	ENT	DVITGLK	Mox1.rat		
487	EEGDARSALI	ITMLQSLHHA	KSGID	IVSGTR		mox2.human			
499	EEGDARSALI	ITMLQALNH	AKNGVD	IVSGTK		cytb558/arabidopsis.pep			
						cytb558.rice.pep			
506	QKTL	YGRPNWDNEFK	TIASQHPN	TRIGVFL		gp91phox.human			
507	QKTL	YGRPNWDNEFK	TIASQHPN	TRIGVFL		gp91phox.bovine			
507	QKTL	YGRPNWDNEFK	TIASQHPN	TRIGVFL		gp91phox.mouse			
501	QKT	SFGRPMWDNEF	STIATSHPK	SVVG	VFL	Mox1.human			
500	QKT	SFGRPMWDNEF	SRIATAHPK	SVVG	VFL	Mox1.human			
505	QKT	FYGRPNWNNEF	KQIAYNHP	SSSIG	VFF	Mox1.rat			
517	VRT	HFA	RPNWR	SVF	KHVA	VNHVNQR	VGVFY	mox2.human	
529	VRT	HFA	RPNWR	KVLSK	IS	SKHP	YAKIG	VFF	cytb558/arabidopsis.pep
									cytb558.rice.pep
536	CGPEALA	ETLSKQ	SISNS	ESGPRGV	HFI	FN		gp91phox.human	
537	CGPEALA	ETLSKQ	SISNS	DSGPRGV	HFI	FN		gp91phox.bovine	
537	CGPEALA	ETLSKQ	SISNS	ESGPRGV	HFI	FN		gp91phox.mouse	
531	CGPRT	LAKSLRK	CC	CHRYSS	SLDPRK	VQFY	FN	Mox1.human	
530	CGPPT	LAKSLRK	CC	CHRYSS	SLDPRK	VQFY	FN	Mox1.human	
535	CGPKA	LSRTLQ	KMCHLYSS	ADPRGV	HFI	FN		Mox1.rat	
547	CGNTC	IIGEL	KRLAQDF	SRKT	-	-	TTKFE	FH	mox2.human
559	CGAPV	LAQELSK	LC	HEFNGKC	-	-	TTKFE	DFH	cytb558/arabidopsis.pep
									cytb558.rice.pep
566	KENF								gp91phox.human
567	KENF								gp91phox.bovine
567	KENF								gp91phox.mouse
561	KEN	-	F						Mox1.human
560	KET	F							Mox1.rat
565	KE	S	F						mox2.human
575	KENF								cytb558/arabidopsis.pep
587	KEH	F							cytb558.rice.pep

Decoration 'Decoration #1': Box residues that match the Consensus exactly.

FIGURE 1(d)

5/8




FIGURE 2

6/8




FIGURE 3

7/8




FIGURE 4

8/8




FIGURE 5

