

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
10 December 2020 (10.12.2020)



(10) International Publication Number
WO 2020/247341 A1

(51) International Patent Classification:

A61K 35/28 (2015.01) C12N 5/0789 (2010.01)
C12N 5/078 (2010.01)

(21) International Application Number:

PCT/US2020/035648

(22) International Filing Date:

02 June 2020 (02.06.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/858,142 06 June 2019 (06.06.2019) US

(71) Applicant: **MEDEOR THERAPEUTICS, INC.**
[US/US]; 611 Gateway Blvd., Suite 120, South San Francisco, CA 94080 (US).

(72) Inventors: **BATTY, D., Scott, Jr.**; 1001 Laurel Street, Apt. 419, San Carlos, CA 94070 (US). **CHEN, Corinna, X.**; 611 Gateway Blvd., Suite 120, South San Francisco, CA 94080 (US).

(74) Agent: **SCHOEN, Adam, M.**; Brown Rudnick LLP, One Financial Center, Boston, MA 02111 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: METHODS OF MAKING CELLULAR PRODUCTS BY POST-MORTEM MOBILIZATION AND HARVESTING OF HEMATOPOIETIC CELLS




FIG. 1

(57) Abstract: The invention provides methods of preparing cellular products to support organ transplantation using hematopoietic cells from deceased donors. The methods include providing to a deceased subject an agent that promotes mobilization of hematopoietic cells from bone marrow to peripheral blood, harvesting hematopoietic cells from the subject, and preparing a cellular product comprising the harvested hematopoietic cells.



WO 2020/247341 A1

METHODS OF MAKING CELLULAR PRODUCTS BY POST-MORTEM MOBILIZATION
AND HARVESTING OF HEMATOPOIETIC CELLS

Cross-Reference to Related Applications

5 This application claims the benefit of, and priority to, U.S. Provisional Patent Application No. 62/858,142, filed June 6, 2019, the contents of which are incorporated by reference.

Field of the Invention

The invention generally relates to methods of making therapeutic compositions
10 containing cells derived from deceased donors.

Background

Nearly 35,000 organ transplants are performed in the United States each year. The primary complication of organ transplantation is rejection of the organ by the recipient's immune
15 system. To avoid organ rejection, almost all transplant recipients must take immunosuppressive drugs for the rest of their lives. Immunosuppressive therapy, however, carries its own set of risks, including increased risk of infection, cancer, hypertension, and liver damage. In addition, immunosuppression does not guarantee that the recipient will tolerate the graft.

Long-term graft tolerance without immunosuppression can be achieved by reconstruction
20 of the recipient's immune system to comprise a mixture of donor-derived and recipient-derived cells, a state called mixed chimerism. Accordingly, establishing mixed chimerism in organ transplant recipients has become very desirable.

Mixed chimerism can be achieved by providing the recipient with donor blood cells, including hematopoietic stem and progenitor cells (HSPCs) and T cells. Unfortunately, the
25 supply of blood cells from adult donors is limited by the availability and complexity of HSPCs obtained from living donors. Not all patients who require an organ transplant have an available and willing living donor. Also, HSPCs may be collected from living donors by surgical extraction, but surgery typically involves general anesthesia, can be painful, and carries a risk of infection or damage to nerves and muscles. Alternatively, HSPCs can be recovered non-
30 surgically by administering to the donor an agent that mobilizes HSPCs from the bone marrow into the blood and collecting cells from peripheral blood by apheresis. However, apheresis can

take up to six hours, may need to be repeated for several days to obtain a sufficient quantity of HSPCs, and poses a risk of infection and blood clotting. Furthermore, regardless of the method, donation of HSPCs provides no medical benefit to the donor, so many individuals are unwilling to donate HSPCs unless they have a direct familial or personal relationship with the recipient.

5

Summary

The invention overcomes the difficulties associated with harvesting HSPCs from living donors by providing methods of obtaining HSPCs and T cells from deceased donors. The methods include post-mortem administration of an agent that mobilizes hematopoietic cells from bone marrow to peripheral blood. By providing a mobilizing agent to the donor shortly after death, the number of HSPCs in the peripheral blood is increased. Consequently, using simple exsanguination and processing steps, HSPCs can be obtained from a single donor in quantities sufficient to promote mixed chimerism in one or more recipients. The methods of the invention also allow improved yield of HSPCs extracted from bone marrow. Thus, the invention provides methods of using cells or tissue from deceased donors to make products that support organ transplantation without the need for recipients to undergo immunosuppressive therapy.

In an aspect, the invention provides methods of preparing a cellular product by providing to a deceased subject one or more agents that promote mobilization of hematopoietic cells from bone marrow to peripheral blood, harvesting hematopoietic cells from the subject, and preparing a cellular product comprising the harvested hematopoietic cells.

The agent that promotes mobilization of hematopoietic cells from bone marrow to peripheral blood may be provided to the subject within a defined period following the subject's death. For example, the agent may be provided within 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 6 hours, 8 hours, 12 hours, 18 hours, 24 hours, 30 hours, 36 hours, 48 hours, or 72 hours of the subject's death. When multiple agent a provided, each may be independently be provided within 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 6 hours, 8 hours, 12 hours, 18 hours, 24 hours, 30 hours, 36 hours, 48 hours, or 72 hours of the subject's death.

Each agent that promotes mobilization of hematopoietic cells from bone marrow to peripheral blood may independently be an adenosine receptor antagonist, BIO5192, a CCR1 antagonist, a CCR2 antagonist, a CXCR2 antagonist, a CXCR4 antagonist, cyclophosphamide, defibrotide, EphA3-Fc, erythropoietin (EPO), filgrastim, glycosaminoglycan (GAG) mimetic,

granulocyte colony stimulating factor (G-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), growth-regulated oncogene beta (GRO-beta), human growth hormone, IL-8, lenograstim, macrophage inflammatory protein-1 alpha (MIP-1 alpha), met-SDF-1 beta, NSC23766, parathyroid hormone, pertussis toxin, plerixafor, a poly-[1-6]-D-glucopyranosyl-[1-3]-D-glucopyranose (PGG) glucan, a Rac1 inhibitor, a retinoic acid receptor agonist, SB290157, a SDF-1 alpha peptide analog, stem cell factor (SCF), sulfated colominic acid, a sulfated polysaccharide, T134, T140, thrombopoietin (TPO), a TPO receptor agonist, a VCAM-1 inhibitor, a VLA-1 inhibitor, a VLA-4 inhibitor, or an analog or derivative of any of the aforementioned agents.

10 The hematopoietic cells may be harvested from a particular tissue or source from the subject's body. The hematopoietic cells may be harvested from peripheral blood. The hematopoietic cells may be harvested from bone marrow. The hematopoietic cells may be harvested from multiple tissues or source from the subject's body. The hematopoietic cells may be harvested from peripheral blood and bone marrow. The cellular product may contain
15 hematopoietic cells harvested from one or more tissues or sources from the subject's body.

The cellular product may include various amounts of CD34⁺ cells and/or CD3⁺ cells. The amount may be specified as a number of cells relative to the body mass of the recipient.

The cellular product may contain at least 1 x 10⁵ CD34⁺ cells/kg recipient weight, at least 2 x 10⁵ CD34⁺ cells/kg recipient weight, at least 4 x 10⁵ CD34⁺ cells/kg recipient weight, at least 5 x 10⁵ CD34⁺ cells/kg recipient weight, at least 1 x 10⁶ CD34⁺ cells/kg recipient weight, at least 2 x 10⁶ CD34⁺ cells/kg recipient weight, at least 4 x 10⁶ CD34⁺ cells/kg recipient weight, at least 5 x 10⁶ CD34⁺ cells/kg recipient weight, at least 1 x 10⁷ CD34⁺ cells/kg recipient weight, at least 2 x 10⁷ CD34⁺ cells/kg recipient weight, at least 4 x 10⁷ CD34⁺ cells/kg recipient weight, at least 1 x 10⁸ CD34⁺ cells/kg recipient weight, at least 2 x 10⁸ CD34⁺ cells/kg recipient weight, at least
25 4 x 10⁸ CD34⁺ cells/kg recipient weight, or at least 5 x 10⁸ CD34⁺ cells/kg recipient weight.

The cellular product may contain at least 1 x 10⁵ CD3⁺ cells/kg recipient weight, at least 2 x 10⁵ CD3⁺ cells/kg recipient weight, at least 4 x 10⁵ CD3⁺ cells/kg recipient weight, at least 5 x 10⁵ CD3⁺ cells/kg recipient weight, at least 1 x 10⁶ CD3⁺ cells/kg recipient weight, at least 2 x 10⁶ CD3⁺ cells/kg recipient weight, at least 4 x 10⁶ CD3⁺ cells/kg recipient weight, at least 5 x 10⁶ CD3⁺ cells/kg recipient weight, at least 1 x 10⁷ CD3⁺ cells/kg recipient weight, at least 2 x 10⁷ CD3⁺ cells/kg recipient weight, at least 4 x 10⁷ CD3⁺ cells/kg recipient weight, at least 1 x
30 10⁷ CD3⁺ cells/kg recipient weight, at least 4 x 10⁷ CD3⁺ cells/kg recipient weight, at least 1 x

10^8 CD3⁺ cells/kg recipient weight, at least 2×10^8 CD3⁺ cells/kg recipient weight, at least 4×10^8 CD3⁺ cells/kg recipient weight, or at least 5×10^8 CD3⁺ cells/kg recipient weight.

The cellular product may contain about 1×10^5 CD3⁺ cells/kg recipient weight, about 2×10^5 CD3⁺ cells/kg recipient weight, about 4×10^5 CD3⁺ cells/kg recipient weight, about 5×10^5 CD3⁺ cells/kg recipient weight, about 1×10^6 CD3⁺ cells/kg recipient weight, about 2×10^6 CD3⁺ cells/kg recipient weight, about 4×10^6 CD3⁺ cells/kg recipient weight, about 5×10^6 CD3⁺ cells/kg recipient weight, about 1×10^7 CD3⁺ cells/kg recipient weight, about 2×10^7 CD3⁺ cells/kg recipient weight, about 4×10^7 CD3⁺ cells/kg recipient weight, about 1×10^8 CD3⁺ cells/kg recipient weight, about 2×10^8 CD3⁺ cells/kg recipient weight, about 4×10^8 CD3⁺ cells/kg recipient weight, or about 5×10^8 CD3⁺ cells/kg recipient weight.

The cellular product may contain at least 1×10^5 CD34⁺ cells/kg recipient weight and at least 1×10^5 CD3⁺ cells/kg recipient weight, at least 1×10^6 CD34⁺ cells/kg recipient weight and at least 1×10^6 CD3⁺ cells/kg recipient weight, at least 1×10^7 CD34⁺ cells/kg recipient weight and at least 1×10^7 CD3⁺ cells/kg recipient weight, at least 1×10^7 CD34⁺ cells/kg recipient weight and at least 1×10^8 CD3⁺ cells/kg recipient weight, at least 1×10^7 CD34⁺ cells/kg recipient weight and at least 1.25×10^8 CD3⁺ cells/kg recipient weight, at least 1×10^7 CD34⁺ cells/kg recipient weight and about 1.25×10^8 CD3⁺ cells/kg recipient weight, or at least 4×10^6 CD34⁺ cells/kg recipient weight and about 1×10^8 CD3⁺ cells/kg recipient weight.

The CD34⁺ cells and CD3⁺ cells may be harvested from the same tissue or source from the subject's body. The CD34⁺ cells and CD3⁺ cells may both be harvested from peripheral blood or from bone marrow. The CD34⁺ cells and CD3⁺ cells may be harvested from different tissues or sources from the subject's body. The CD34⁺ cells may be harvested from bone marrow, and the CD3⁺ cells may be harvested from peripheral blood.

The cellular product may contain facilitating cells. The cellular product may contain facilitating cells at $\geq 1 \times 10^4$ cells/kg of body weight, $\geq 2 \times 10^4$ cells/kg of body weight, $\geq 4 \times 10^4$ cells/kg of body weight, $\geq 5 \times 10^4$ cells/kg of body weight, $\geq 1 \times 10^5$ cells/kg of body weight, $\geq 2 \times 10^5$ cells/kg of body weight, $\geq 4 \times 10^5$ cells/kg of body weight, $\geq 5 \times 10^5$ cells/kg of body weight, $\geq 1 \times 10^6$ cells/kg of body weight, $\geq 2 \times 10^6$ cells/kg of body weight, $\geq 4 \times 10^6$ cells/kg of body weight, $\geq 5 \times 10^6$ cells/kg of body weight, $\geq 1 \times 10^7$ cells/kg of body weight, $\geq 2 \times 10^7$ cells/kg of body weight, $\geq 4 \times 10^7$ cells/kg of body weight, $\geq 5 \times 10^7$ cells/kg of body weight, ≥ 1

x 10^8 cells/kg of body weight, $\geq 2 \times 10^8$ cells/kg of body weight, $\geq 4 \times 10^8$ cells/kg of body weight, or $\geq 5 \times 10^8$ cells/kg of body weight.

Preparing the cellular product may include enriching a portion of the harvested hematopoietic cells for CD34⁺ cells and/or CD3⁺ cells. Preparing the cellular product may include enriching one portion of the harvested hematopoietic cells for CD34⁺ cells and enriching another portion of the harvested hematopoietic cells for CD3⁺ cells.

The enriching step may include one or more of affinity purification, buoyancy-activated cell separation, cell lysis, column filtration, hetastarch sedimentation, immunomagnetic depletion, immunomagnetic purification, size-based centrifugal separation, and spinning membrane filtration. The enriching step may include depleting the harvested hematopoietic cells of red blood cells, platelets, or both. The enriching step may include separating the peripheral blood into a cellular fraction and a plasma fraction.

Preparing the cellular product may include assessing whether the hematopoietic cells are suitable for use in preparing the cellular product. The assessing step may include analyzing in the harvested hematopoietic cells one or more of T cell activation, T cell exhaustion, T cell anergy, T cell proliferation, T cell viability, T cell apoptosis, hematopoietic stem and progenitor cell (HSPC) proliferation, HSPC viability, HSPC apoptosis, chemokine levels, and cytokine levels.

The methods may include cryopreserving the cellular product. For example, cryopreservation may include addition of a cryoprotectant, such as DMSO or dextran having a molecular weight of about 40,000 Da. The cryoprotectant may be present at a concentration of about 1%, 2%, 3%, 4%, 5%, 7.5%, or 10%. The cellular product may be cryopreserved in a single container, or it may be cryopreserved in multiple containers.

The subject and a recipient of the cellular product may be HLA-matched. The subject and a recipient of the cellular product may be HLA-mismatched. The CD34⁺ cells, the CD3⁺ cells, or both may be HLA-matched. The CD34⁺ cells, the CD3⁺ cells, or both may be HLA-mismatched. The subject and recipient may be HLA-matched at six, eight, ten, or twelve alleles among the HLA-A, HLA-B, HLA-C, HLA-DP, HLA-DQ, and HLA-DR genes. The subject and recipient may be HLA-mismatched at one, two, three, four, five, six, or more alleles among the HLA-A, HLA-B, HLA-C, HLA-DP, HLA-DQ, and HLA-DR genes.

Brief Description of the Drawings

FIG. 1 is an illustration of an external cortical cutting trocar of a marrow extracting device that may be used in methods of the invention.

5 FIG. 2 is an illustration of an internal trocar of a marrow extracting device that may be used in methods of the invention.

FIG. 3 shows a blood collection device that may be used in methods of the invention.

FIG. 4 shows a blood collection device that may be used in methods of the invention.

10 FIG. 5 is a flow diagram of the tissue collection step according to a method of the invention.

FIG. 6 is a flow diagram of the bone marrow preparation step according to a method of the invention.

FIG. 7 is a flow diagram of the step of the blood preparation step according to a method of the invention.

15 FIG. 8 is a flow diagram of the buffer preparation step according to a method of the invention.

FIG. 9 is a flow diagram of the CD34⁺ cell-enrichment step according to a method of the invention.

FIG. 10 is a flow diagram of the dividing step according to a method of the invention.

20 FIG. 11 is a flow diagram of the cryopreservation step according to a method of the invention.

FIG. 12 is a flow diagram illustrating a method of preparing a cellular composition according to an embodiment of the invention.

25 Detailed Description

The primary challenge in organ transplantation is getting the recipient's immune system to tolerate the donor's tissue. If the recipient's immune system detects the donated organ as foreign, it attacks the tissue, leading to graft rejection. To avoid graft rejection, most transplant recipients must take drugs that suppress the immune system, but such drugs increase the
30 recipient's risk of infection and cancer. As another means to prevent graft rejection, transplantation of solid organs may be accompanied by transfer of donor-derived blood cell

progenitors. Providing donor blood cells allows reconstitution of the recipient's immune system to include cells from the donor of the organ that have been educated to recognize the organ as non-foreign tissue. Consequently, the donated organ is not attacked, and the recipient tolerates the graft.

5 One strategy for reconstructing the recipient's immune system entails complete replacement of the recipient's hematopoietic system with exclusively donor-derived cells to achieve a state of full chimerism. A risk associated with full chimerism, however, is that the completely donor-derived immune system may identify the recipient's tissue as foreign and attack it, a condition called graft-versus-host disease (GVHD). *See, e.g.,* Sachs et al., Induction
10 of Tolerance through Mixed Chimerism, *Cold Spring Harb Perspect Med* 2014;4:a015529, doi: 10.1101/cshperspect.a015529, the contents of which are incorporated herein by reference. As a result, fully chimeric patients must remain on immunosuppressive therapy indefinitely.

 Another strategy is to repopulate the recipient's immune system with a mixture of donor-derived cells and recipient-derived cells to attain a state called mixed chimerism. Compared to
15 full chimerism, mixed chimerism is associated with lower rates of GVHD. In addition, mixed chimeric regimens require lower doses of immunosuppressive therapy initially and allow complete discontinuation of immunosuppression after the stability of the recipient's mixed chimerism has been established. To date, induction of mixed chimerism is the only method of producing graft tolerance in humans without maintaining immunosuppressive therapy.

20 Current methods of establishing mixed chimerism require transfer of two different populations of hematopoietic cells from the donor. Mature T cells recognize the transplanted organ as "self" tissue and prevent the immune system from attacking it. Mature T cells express the cell-surface marker CD3, and different sub-populations may be found in the blood and lymph nodes. Although CD3⁺ T cells promote tolerance upon transfer, they have a finite lifespan and
25 are unable to regenerate themselves. Consequently, transfer of hematopoietic stem and progenitor cells (HSPCs), pluripotent cells that can differentiate into T cells, is also necessary to allow continual replenishment of the donor-derived T cell population. HSPCs express the cell-surface marker CD34 and reside primarily in the bone marrow.

 To give a transplant recipient the chance to achieve long-term graft tolerance without
30 lifelong immunosuppressive therapy, the sacrifice a living donor must make is substantial. Thus, a living donor must be highly motivated by altruism to undertake the procedures necessary to

provide the tissues and cells necessary to support immunosuppression-free graft tolerance. Consequently, few individuals are willing to become living donors of solid organs and hematopoietic cells, particularly if they have no familial or personal relationship with the prospective recipient.

5 By comparison, a much higher percentage of individuals are willing to donate organs and blood cells posthumously. However, obtaining hematopoietic cells suitable for transfer to a living recipient is more challenging when the donor is deceased. In particular, it is difficult to harvest HSPCs in sufficient quantities to promote mixed chimerism in an organ transplant recipient. HSPCs reside predominantly in the bone marrow and ordinarily are only present at
10 low numbers in the circulating blood. Within the bone marrow, HSPCs exist in a complex milieu containing a variety of cell types and extracellular matrix proteins and form various molecular interactions with those other cells and proteins. During extraction of HSPCs, such interactions hinder isolation of HSPCs from other cell types and non-cellular material, so procedures to disaggregate and fractionate cells are necessary to recover populations of HSPCs
15 suitable to support transplantation.

The invention provides preparative methods that overcome the difficulties of obtaining hematopoietic cells from deceased donors. The methods entail treating the donor's body with an agent that mobilizes hematopoietic cells from bone marrow to peripheral blood prior to harvesting the cells from the donor's body. Because the agent increases the number of HSPCs in
20 the peripheral blood, HSPCs can simply be recovered by exsanguination and purification of HSPCs from the blood. Consequently, the invention unlocks the therapeutic potential of tissue donated from deceased donors to extend and improve the lives of patients who need organ transplants.

25 **Cellular products derived from deceased donors for transfer to living recipients**

The invention provides methods of making cellular products that contain hematopoietic cells obtained from deceased donors. All hematopoietic cells are derived from HSPCs, multipotent cells that can differentiate into various specialized cells and also reproduce to generate new HSPCs. HSPCs that differentiate form either lymphoid progenitors or myeloid
30 progenitors. Lymphoid progenitors give rise to lymphocytes and natural killer cells. Myeloid progenitors produce cells of the myeloid and erythroid lineages, such as erythrocytes, platelets,

basophils, neutrophils, eosinophils, monocytes, macrophages, and antigen-presenting cells, such as dendritic cells. In adults, most hematopoietic development occurs in the bone marrow, although maturation and activation of some lymphoid cells occurs in the spleen, thymus, and lymph nodes.

5 In certain embodiments, the invention provides methods of making cellular compositions that include two populations of cells that allow donor HSPCs to develop into mature cells of the immune system in the recipient's body. One population includes CD34⁺ cells. CD34 is a cell surface marker that is expressed in stem cells and their immediate descendants, multipotent progenitor cells, which have not committed to either the myeloid or lymphoid lineage.

10 Consequently, CD34 expression is a useful measure for identifying populations of cells that contain HSPCs. In adults, CD34⁺ cells reside predominantly in the bone marrow.

The cellular compositions may also include CD3⁺ cells. CD3 comprises a group of polypeptides that interact with the two polypeptide chains of the T cell receptor to form the T cell receptor complex. The CD3 complex includes a gamma chain, delta chain, and two epsilon
15 chains. CD3 is expressed on the surface of mature T cells and is thus useful as a marker for T cells. CD3⁺ cells are abundant in the circulating blood.

To promote establishment of mixed chimerism in the recipient, the cellular products include CD34⁺ cells and CD3⁺ cells in appropriate quantities. The cellular products may contain CD34⁺ cells and CD3⁺ cells in defined amounts.

20 For example and without limitation, the cellular product may contain CD34⁺ cells at $\geq 1 \times 10^4$ cells/kg of body weight, $\geq 2 \times 10^4$ cells/kg of body weight, $\geq 4 \times 10^4$ cells/kg of body weight, $\geq 5 \times 10^4$ cells/kg of body weight, $\geq 1 \times 10^5$ cells/kg of body weight, $\geq 2 \times 10^5$ cells/kg of body weight, $\geq 4 \times 10^5$ cells/kg of body weight, $\geq 5 \times 10^5$ cells/kg of body weight, $\geq 1 \times 10^6$ cells/kg of body weight, $\geq 2 \times 10^6$ cells/kg of body weight, $\geq 4 \times 10^6$ cells/kg of body weight, $\geq 5 \times 10^6$
25 cells/kg of body weight, $\geq 1 \times 10^7$ cells/kg of body weight, $\geq 2 \times 10^7$ cells/kg of body weight, $\geq 4 \times 10^7$ cells/kg of body weight, $\geq 5 \times 10^7$ cells/kg of body weight, $\geq 1 \times 10^8$ cells/kg of body weight, $\geq 2 \times 10^8$ cells/kg of body weight, $\geq 4 \times 10^8$ cells/kg of body weight, or $\geq 5 \times 10^8$ cells/kg of body weight.

For example and without limitation, the cellular product may contain CD3⁺ cells at $\geq 1 \times$
30 10^4 cells/kg of body weight, $\geq 2 \times 10^4$ cells/kg of body weight, $\geq 4 \times 10^4$ cells/kg of body weight, $\geq 5 \times 10^4$ cells/kg of body weight, $\geq 1 \times 10^5$ cells/kg of body weight, $\geq 2 \times 10^5$ cells/kg of body

weight, $\geq 4 \times 10^5$ cells/kg of body weight, $\geq 5 \times 10^5$ cells/kg of body weight, $\geq 1 \times 10^6$ cells/kg of body weight, $\geq 2 \times 10^6$ cells/kg of body weight, $\geq 4 \times 10^6$ cells/kg of body weight, $\geq 5 \times 10^6$ cells/kg of body weight, $\geq 1 \times 10^7$ cells/kg of body weight, $\geq 2 \times 10^7$ cells/kg of body weight, $\geq 4 \times 10^7$ cells/kg of body weight, $\geq 5 \times 10^7$ cells/kg of body weight, $\geq 1 \times 10^8$ cells/kg of body weight, $\geq 2 \times 10^8$ cells/kg of body weight, $\geq 4 \times 10^8$ cells/kg of body weight, or $\geq 5 \times 10^8$ cells/kg of body weight.

For example and without limitation, the cellular product may contain CD3⁺ cells at about 1×10^4 cells/kg of body weight, about 2×10^4 cells/kg of body weight, about 4×10^4 cells/kg of body weight, about 5×10^4 cells/kg of body weight, about 1×10^5 cells/kg of body weight, about 2×10^5 cells/kg of body weight, about 4×10^5 cells/kg of body weight, about 5×10^5 cells/kg of body weight, about 1×10^6 cells/kg of body weight, about 2×10^6 cells/kg of body weight, about 4×10^6 cells/kg of body weight, about 5×10^6 cells/kg of body weight, about 1×10^7 cells/kg of body weight, about 2×10^7 cells/kg of body weight, about 4×10^7 cells/kg of body weight, about 5×10^7 cells/kg of body weight, about 1×10^8 cells/kg of body weight, about 2×10^8 cells/kg of body weight, about 4×10^8 cells/kg of body weight, or about 5×10^8 cells/kg of body weight.

The cellular product may contain at least 1×10^5 CD34⁺ cells/kg recipient weight and at least 1×10^5 CD3⁺ cells/kg recipient weight, at least 1×10^6 CD34⁺ cells/kg recipient weight and at least 1×10^6 CD3⁺ cells/kg recipient weight, or at least 4×10^6 CD34⁺ cells/kg recipient weight and about 1×10^8 CD3⁺ cells/kg recipient weight. Other concentrations are exemplified in each of Strober et al., U.S. patent number 9,504,717 and Strober et al., U.S. patent number 9,561,253, the content of each of which is incorporated by reference herein in its entirety.

Facilitating cells are non-stem cells that improve long-term engraftment of stem cells. Facilitating cells and methods of preparing them are known in the art and described in, for example, U.S. Patent Nos. 5,772,994; 8,632,768; 9,452,184; and 9,678,062 the contents of which are incorporated herein by reference.

Facilitating cells may be characterized based on presence, absence, or level of expression of phenotypic markers. Typically, facilitating cells are CD8⁺ and alpha beta TCR⁺. However, alternative or additional markers may be used to characterize facilitating cells. For example and without limitation, facilitating cells may be characterized by the presence, absence, or level of one or more of B220, CD11a, CD11b, CD11c, CD123, CD162, CD19, CD3 epsilon, CD52, CD56, CD62L, CXCR4, FoxP3, gamma delta TCR, HLADR, NKp30, NKp44, and NKp46.

The cellular product may contain facilitating cells at $\geq 1 \times 10^4$ cells/kg of body weight, $\geq 2 \times 10^4$ cells/kg of body weight, $\geq 4 \times 10^4$ cells/kg of body weight, $\geq 5 \times 10^4$ cells/kg of body weight, $\geq 1 \times 10^5$ cells/kg of body weight, $\geq 2 \times 10^5$ cells/kg of body weight, $\geq 4 \times 10^5$ cells/kg of body weight, $\geq 5 \times 10^5$ cells/kg of body weight, $\geq 1 \times 10^6$ cells/kg of body weight, $\geq 2 \times 10^6$ cells/kg of body weight, $\geq 4 \times 10^6$ cells/kg of body weight, $\geq 5 \times 10^6$ cells/kg of body weight, $\geq 1 \times 10^7$ cells/kg of body weight, $\geq 2 \times 10^7$ cells/kg of body weight, $\geq 4 \times 10^7$ cells/kg of body weight, $\geq 5 \times 10^7$ cells/kg of body weight, $\geq 1 \times 10^8$ cells/kg of body weight, $\geq 2 \times 10^8$ cells/kg of body weight, $\geq 4 \times 10^8$ cells/kg of body weight, or $\geq 5 \times 10^8$ cells/kg of body weight.

The cellular product may contain CD34⁺ cells at a designated level of purity. For example, the cellular product may contain CD34⁺ cells that are at least 0.1%, at least 0.2%, at least 0.3%, at least 1%, at least 3%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% pure. The cellular product may contain CD34⁺ cells that are from about 0.1% to about 2%, from about 0.2% to about 2%, from about 0.3% to about 2%, from about 0.1% to about 3%, from about 0.2% to about 3%, from about 0.3% to about 3%, from about 0.1% to about 4%, from about 0.2% to about 4%, from about 0.3% to about 4%, from about 0.1% to about 5%, from about 0.2% to about 5%, from about 0.3% to about 5%, from about 1% to about 3%, from about 1% to about 10%, from about 1% to about 20%, or from about 1% to about 40% pure. Other purities are exemplified in each of Strober et al., U.S. patent number 9,504,717 and Strober et al., U.S. patent number 9,561,253, the content of each of which is incorporated by reference herein in its entirety.

The CD34⁺ cells and CD3⁺ cells may be provided as a mixture in one or more containers. The CD34⁺ cells and CD3⁺ cells may be provided in separate containers. Any commercially available container approved to hold cellular products may be used.

The cellular product may be provided frozen. Consequently, the cellular product may contain a cryoprotectant. Any cryoprotectant known in the art may be used. For example and without limitation, the cryoprotectant may be DMSO, dextran having an average molecular weight of 40 kDa, serum, e.g., bovine serum, albumin, e.g., human serum albumin, or cell culture medium. The cryoprotectant may be present at a defined concentration. For example, the cellular product may contain about 1% DMSO, about 2% DMSO, about 5% DMSO, about 7.5% DMSO, about 10% DMSO, about 12.5% DMSO, about 15% DMSO, or about 20% DMSO. The

cellular product may contain about 1% dextran, about 2% dextran, about 5% dextran, about 7.5% dextran, about 10% dextran, about 12.5% dextran, about 15% dextran, or about 20% dextran.

The cryoprotectant may be a commercially available freezing medium, such as the medium sold under the trade name CryoStor 10 by BioLife Solutions (Bothell, WA). Cryoprotection is
5 discussed in U.S. Patent No. 9,504,717 and U.S. Patent No. 9,561,253, the contents of each of which are incorporated by reference herein in its entirety.

The cellular product may contain agents that enhance engraftment or functional mobilization of the hematopoietic cells in the recipient. The cellular product may contain agents that prevent a negative reaction of the recipient to the hematopoietic cells. For example and
10 without limitation, the pharmaceutical composition may contain a cytokine, chemokine, growth factor, excipient, carrier, antibody or a fragment thereof, small molecule, drug, agonist, antagonist, matrix protein, or complementary cell type.

The cellular product may contain a buffer. The cellular product may be buffer to maintain physiologically compatible pH. For example, the cellular product may be buffered to a
15 neutral pH, such as from about 6.0 to about 8.0.

The cellular product may be supplied in the form of a pharmaceutical composition, comprising an isotonic excipient prepared under sufficiently sterile conditions for human administration. Choice of the cellular excipient and any accompanying elements of the composition are adapted in accordance with the route and device used for administration. For
20 general principles in medicinal formulation, see Cell Therapy: Stem Cell Transplantation, Gene Therapy, and Cellular Immunotherapy, by G. Morstyn & W. Sheridan. eds., Cambridge University Press, 1996; and Hematopoietic Stem Cell Therapy, E. D. Ball, J. Lister & P. Law, Churchill Livingstone, 2000.

The CD34⁺ cells, CD3⁺ cells, or both may be HLA-matched or HLA-mismatched to the
25 recipient. Human leukocyte antigens (HLAs), also called major histocompatibility complex (MHC) antigens, are protein molecules expressed on the surface of cells that confer a unique antigenic identity to these cells. MHC/HLA antigens are target molecules that are recognized by T-cells and natural killer (NK) cells as being derived from the same source of hematopoietic stem cells as the immune effector cells (“self”) or as being derived from another source of
30 hematopoietic reconstituting cells (“non-self”). Two main classes of HLA antigens are recognized: HLA class I and HLA class II. HLA class I antigens (A, B, and C in humans)

render each cell recognizable as “self,” whereas HLA class II antigens (DR, DP, and DQ in humans) are involved in reactions between lymphocytes and antigen presenting cells.

A key aspect of the HLA gene system is its polymorphism. Each gene exists in different alleles. Allelic gene products differ in one or more amino acids in the alpha and/or beta
5 domain(s). An individual has two alleles of each gene, for a total of twelve alleles among the HLA-A, HLA-B, HLA-C, HLA-DP, HLA-DQ, and HLA-DR genes. An HLA-matched donor may have a match with the recipient at six, eight, ten, or twelve alleles selected from any combination of the HLA-A, HLA-B, HLA-C, HLA-DP, HLA-DQ, and HLA-DR genes. The genes most important for HLA typing are HLA-A, HLA-B, and HLA-DR, so the donor and
10 recipient may be matched at all six alleles of the HLA-A, HLA-B, and HLA-DR genes. An HLA-mismatched donor may have a mismatch at one, two, three, four, five, six, or more alleles among the HLA-A, HLA-B, HLA-C, HLA-DP, HLA-DQ, and HLA-DR genes. HLA typing may be performed by any method known in the art. Examples of HLA typing methods include serological cytotoxicity, flow cytometry, and DNA typing. Such methods are described in, for
15 example, U.S. Patent No. 9,561,253, the contents of which are incorporated herein by reference.

The HLA genes are clustered in a super-locus present on chromosome position 6p21. Consequently, the set of alleles present on a single chromosome, i.e., a haplotype, tends to be inherited as a group. Identifying a patient’s haplotypes can help predict the probability of finding matching donors and assist in developing a search strategy. Haplotypes vary in how
20 common they are among the general population and in their frequency within different racial and ethnic groups.

Manufacture of cellular products from deceased donors may include additional characterization of the products. For example, a profile of one or more secreted molecules, such as cytokines or chemokines, may be established for compositions containing CD34⁺ cells and/or
25 CD3⁺ cells. Alternatively or additionally, a profile of expression of cell surface markers, such as CD3, CD34, or CD45, may be established for the composition. Any suitable method may be used to characterize the compositions, including one or more of the methods described below in relation to analysis of blood from deceased donors. Cellular products may also be tested for the presence of pathogens, such as mycoplasma, or endotoxins that evidence the presence of
30 pathogens.

Providing an agent that promotes mobilization of hematopoietic cells

The methods of the invention include administering to the body of a deceased donor one or more agents that promote mobilization of hematopoietic cells from bone marrow to peripheral blood. As discussed above, the agent facilitates recovery of HSPCs from the blood.

5 The agent should be provided to the donor's body shortly after death to minimize the detrimental effects of cytokines released during the brain death process. For example, the agent may be provided within 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 6 hours, 8 hours, 12 hours, 18 hours, 24 hours, 30 hours, 36 hours, 48 hours, or 72 hours of the subject's death. When multiple agent a provided, each may be independently be provided within 30 minutes, 1 hour, 2
10 hours, 3 hours, 4 hours, 6 hours, 8 hours, 12 hours, 18 hours, 24 hours, 30 hours, 36 hours, 48 hours, or 72 hours of the subject's death.

A variety of agents that mobilize hematopoietic cells from bone marrow to peripheral blood are known in the art. Many such agents disrupt HSPC cell-cell and/or cell-matrix interactions within the bone marrow to allow the release of HSPCs into the peripheral blood.
15 Classes of agents that mobilize hematopoietic cells, such as HSPCs, from bone marrow include chemotherapeutics, hematopoietic growth factors, chemokines, inhibitors of chemokine receptors, and inhibitors of integrins. Specific examples of agents that promote mobilization of hematopoietic cells include adenosine receptor antagonists, BIO5192, CCR1 antagonists, CCR2 antagonists, CXCR2 antagonists, CXCR4 antagonists, cyclophosphamide, defibrotide, EphA3-
20 Fc, erythropoietin (EPO), filgrastim, glycosaminoglycan (GAG) mimetic, granulocyte colony stimulating factor (G-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), growth-regulated oncogene beta (GRO-beta), human growth hormone, IL-8, lenograstim, macrophage inflammatory protein-1 alpha (MIP-1 alpha), met-SDF-1 beta, NSC23766, parathyroid hormone, pertussis toxin, plerixafor, a poly-[1-6]--D-glucopyranosyl-[1-3]--D-
25 glucopyranose (PGG) glucan, Rac1 inhibitors, retinoic acid receptor agonists, SB290157, a SDF-1 alpha peptide analog, stem cell factor (SCF), sulfated colominic acid, sulfated polysaccharides, T134, T140, thrombopoietin (TPO), TPO receptor agonists, VCAM-1 inhibitors, VLA-1 inhibitors, VLA-4 inhibitors, and analogs and derivatives of any of the aforementioned agents.

The agent may be provided to the deceased donor by any suitable means. For example,
30 the agent may be provided by injection using a needle, catheter, central line or the like. The agent may be delivered intravascularly, intravenously, intraarterially, or directly to the bone.

The hematopoietic cells may be harvested at a defined time point. The time of harvesting may be defined in relation to the time of death or in relation to the time of administration of one or more agents. For example and without limitation, hematopoietic cells may be harvested about 8 hours, about 12 hours, about 18 hours, about 24 hours, about 30 hours, about 36 hours, about 48 hours, about 72 hours, about 4 days, about 5 day, about 6 days, or about 7 days after death. Hematopoietic cells may be harvested about 8 hours, about 12 hours, about 18 hours, about 24 hours, about 30 hours, about 36 hours, about 48 hours, about 72 hours, about 4 days, about 5 day, about 6 days, or about 7 days after administration of an agent that mobilizes hematopoietic cells.

The method may include additional steps to preserve the subject's body or to facilitate recovery of the hematopoietic cells from the subject's body. For example, the method may include regulating the temperature, e.g., heating or cooling, of the subject's body or of a component of the subject's body, e.g., the blood. The additional steps may be performed prior to administration of the agent, following administration of the agent, or both.

Preparation of cellular products from deceased donors for transfer to living recipients

The invention provides methods of preparing cellular products that contain of CD34⁺ cells and CD3⁺ cells obtained from deceased donors. Given the predominant anatomical locations of CD34⁺ cells and CD3⁺ cells, as indicated above, different tissues or fluids may be preferred sources for the two cell types. For example, methods may include obtaining CD34⁺ cells from bone marrow of a deceased donor and obtaining CD3⁺ cells from a different source, such as blood, liver, lymph nodes, spleen, or thymus, from the donor. Methods of harvesting CD34⁺ cells and CD3⁺ cells from deceased donors generally are described in co-owned, co-pending U.S. Application Nos. 62/732,823; 62/732,827; 62/732,830; and 62/732,832, the contents of each of which are incorporated herein.

Bones include a hard outer layer, called cortical bone or compact bone, and an internal spongy portion, called cancellous bone, which contains the bone marrow. Bone marrow may be obtained from the cancellous bone material of large bones, such as the pelvis, vertebrae, ribs, femur, tibia, and sternum. Preferred sources of bone marrow are the iliac crests of the pelvis and vertebral bodies of the vertebrae.

The methods may include removal of bone marrow and a source of CD3⁺ cells, such as blood, from the body of the deceased donor in either order. Preferably, the blood is removed

first, i.e., the body is exsanguinated, and then the bone marrow is obtained. The methods may include removal of the solid organ of interest, such as a kidney, lung, pancreas, pancreatic islet cells, heart, intestine, colon, liver, skin, muscle, gum, eye, or tooth. The solid organ of interest, bone marrow, and source of CD3⁺ cells, such as blood, may be removed in any order.

5 Preferably, removal occurs in the following sequence: blood, solid organ of interest, and bone marrow.

Bone marrow may be removed from the body of the deceased donor by any suitable method. In some methods, bone marrow is removed by aspiration. Aspiration involves inserting a needle into the bone and withdrawing the bone marrow. In some methods, bone marrow is
10 removed by trephination. A trephine is a tube with a circular end that cuts into the bone to extract a cylindrical portion of bone and bone marrow.

FIG. 1 is an illustration of an external cortical cutting trocar 101 of a marrow extracting device that may be used in methods of the invention. The external trocar 101 includes a hollow shaft 113, which may be made of surgical steel. At the distal end of the external trocar 101 is a
15 cutting tip 115, such as a toothed saw. The cutting tip 115 may be driven manually, pneumatically, or electrically. The cutting tip 115 and may be replaceable. At a proximal end of the external trocar 101 is a removable pneumatic drive adapter 117. The pneumatic drive adapter 117 couples the external trocar 101 with a pneumatic drive and may be removable from the external trocar 101. The external trocar 101 may include a suction adapter that couples the
20 external trocar 101 to a vacuum pump or other suction source to facilitate removal of bone fragments. The suction adaptor may be fitted to the proximal end of the external trocar 101 when the drive adapter 117 is removed, or it may fit directly into the drive adapter 117.

FIG. 2 is an illustration of an internal trocar 201 of a marrow extracting device that may be used in methods of the invention. The internal trocar 201 includes a shaft 223 that has one or
25 more deployable rongeur blades 219 positioned at the distal end. The rongeur blades can be positioned at different angles to achieve different circular cutting radii. Adjusting the cutting radius is useful for cutting into bones of different sizes. For example, a smaller cutting radius is needed for removal of vertebral bodies, whereas a larger cutting radius is optimal for extracting bone segments from the femur or tibia. The internal trocar 201 also includes one or more
30 handles 217, or wings, at its proximal end. The handles 217 can be rotated by hand to allow

manual extraction of bone segments. Alternatively or additionally, the internal trocar 201 may be coupled to a pneumatic drive for low-speed operation.

The following sequence of steps may be used to remove a portion of bone from a deceased donor. The following sequence is for illustrative purposes only; one of skill in the art will understand that other methods of bone removal are possible within the scope of the invention. First, a cortical surface of target bone is exposed by standard surgical procedure. Next, an external cutting trocar 101 with a cutting tip 115 and a pneumatic drive adapter 117 placed in the chuck of a pneumatic drive is used to cut through the cortex of the bone and into the medullary space. The pneumatic drive adapter 117 is then removed from the drive. Next, a solid internal trocar 201 is inserted through the shaft 113 of the external trocar 101, and the rongeur blades 219 are deployed. The handles 217 are used to twist the rongeur blades 219 and disrupt the medullary trabeculae. Alternatively, the internal trocar 201 can be engaged to a pneumatic drive at low speed. The internal trocar 201 is then removed from the shaft 113, and a suction adapter is affixed to the hollow trocar. Finally, a suction device is attached via the suction adapter, and suction is applied to evacuate the medullary space into a collection bag.

Because portions of bone obtained by trephination typically contain bone shards, methods of obtaining CD34⁺ cells from portions of bone may include procedures to separate bone marrow cells from bone shards. A variety of methods may be used to isolate bone marrow cells from bone shards, such as physical agitation, enzymatic disaggregation, washing, and filtration. Methods may include treating the bone shards with one or more agents that mobilize hematopoietic cells, such as those agents described above, to improve the yield of HPSCs from the portions of bone. Treatment may include immersing bone shards in a preservation solution or liquid that contains one or more mobilizing agents. Treatment may occur within a storage container so that release of HPSCs occurs during the transport or shipping of extracted material. Thus, the portions of bone may be extracted from a donor at a first site and shipped to a recipient at a second site, and mobilization of hematopoietic cells may occur, at least in part, while the tissue is in transit.

Although bone marrow is the native environment for HPSCs, HPSCs represent only a small fraction of bone marrow cells. CD34⁺ cells make up only about 1% of all nucleated cells in the bone marrow. Therefore, methods of the invention may include steps to enrich bone marrow extracts from deceased donors for CD34⁺ cells.

One form of enrichment is to separate mononuclear cells from enucleated cells, such as red blood cells and platelets, and cells with multi-lobed nuclei, such as granulocytes, including neutrophils, basophils, and eosinophils. Several methods for enriching for isolating or enriching for mononuclear cells are known in the art. For example and without limitation, mononuclear
5 cells may be isolated or enriched by buoyancy-activated cell separation, cell lysis, hetastarch sedimentation, immunomagnetic depletion, size-based centrifugal separation, and spinning membrane filtration. Systems for buoyancy-activated cell separation are commercially available from Cesca Therapeutics, Inc. (Rancho Cordova, CA) and described in, for example, U.S. Patent No. 9,695,394, the contents of which are incorporated herein by reference. Systems for size-
10 based centrifugal separation are commercially available from Sepax Technologies, Inc. (Newark, DE). Systems for spinning membrane filtration are commercially available, such as the system sold under the trade name Lovo Automate Cell Processing System by Fresenius Kabi USA, LLC (Lake Zurich, IL).

CD34⁺ cells may be purified based on qualitative or quantitative expression of one or
15 more cell surface markers. Examples of suitable cell surface markers include AC133, CD3, CD34, CD38, CD45, and Thy-1. CD34⁺ cells may be purified based on the presence or absence of a marker or on the level of expression of a marker, e.g., high vs. low. Purification of CD34⁺ cells may include comparison of marker expression, complete blood cell counts, and/or mononuclear cell counts between starting material and material that has been enriched for CD34⁺
20 cells.

CD34⁺ cells may be purified by selectively binding a suitable affinity reagent to CD34 or another marker. The affinity reagent may be an antibody, a full-length antibody, a fragment of an antibody, a naturally occurring antibody, a synthetic antibody, an engineered antibody, a full-length affibody, a fragment of an affibody, a full-length affilin, a fragment of an affilin, a full-
25 length anticalin, a fragment of an anticalin, a full-length avimer, a fragment of an avimer, a full-length DARPIn, a fragment of a DARPIn, a full-length fynomer, a fragment of a fynomer, a full-length kunitz domain peptide, a fragment of a kunitz domain peptide, a full-length monobody, a fragment of a monobody, a peptide, a polyaminoacid, or the like. The affinity reagent may be directly conjugated to a detection reagent and/or purification reagent. The detection reagent and
30 purification reagent may be the same, or they may be different. For example, the detection reagent and/or purification reagent may be fluorescent, magnetic, or the like. The detection

reagent and/or purification reagent may be a magnetic particle for column purification. For example, magnetic column purification may be performed using the Miltenyi system of columns, antibodies, buffers, preparation materials and reagents, etc. known to those of skill in the art. Methods of affinity purification of hematopoietic cells, including CD34⁺ and CD3⁺ cells, and analysis of purified populations are described in, for example, U.S. Patent Nos. 9,561,253; and 9,452,184, the contents of which are incorporated herein by reference.

CD34⁺ cells may be isolated, enriched, or purified by any method. For example, CD34⁺ cells may be isolated, enriched, or purified by column purification, flow cytometry, cell sorting, or immunoadsorption column separation. Preferably, CD34⁺ cells are purified using an immunomagnetic column system, such as those sold under the trade name CliniMACS by Miltenyi Biotec Inc. (Auburn, CA), Methods of affinity purification of hematopoietic cells, including CD34⁺ cells, and analysis of purified populations are described in, for example, U.S. Patent No. 9,561,253; U.S. Patent No. 9,452,184; Ng et al., Isolation of human and mouse hematopoietic stem cells, *Methods Mol Biol.* (2009) 506:13-21. doi: 10.1007/978-1-59745-409-4_2; and Spohn et al., Automated CD34⁺ cell isolation of peripheral blood stem cell apheresis product, *Cytotherapy* (2015) Oct;17(10):1465-71. doi: 10.1016/j.jcyt.2015.04.005, the contents of each of which are incorporated herein by reference. The methods may include positive selection, negative selection, or both.

The methods may include various other treatments of bone marrow, blood, or other tissue sources that facilitate the recovery of CD34⁺ cells and/or CD3⁺ cells for use in the manufacture of product to administer to a living recipient. For example, blood, bone marrow, or other tissue may be treated to remove clots and/or cell clumps, i.e., agglutination. Clots and clumps may be removed by any suitable method. Non-limiting examples for removal of clots and clumps include filtration, e.g., spinning membrane filtration, as described above; treatment with thrombolytic drugs, such as alteplase, anistreplase, kabikinase, recombinant tissue plasminogen activators, reteplase, streptokinase, tenecteplase, and urokinase; ultrasound; and mechanical rubbing, as described in, for example, Khalil, et al., Rubbing Against Blood Clots Using Helical Robots: Modeling and In Vitro Experimental Validation, *IEEE Robotics and Automation Letters* (2(2):927-934, April 2017, DOI: 10.1109/LRA.2017.2654546, the contents of which are incorporated herein by reference.

Bone marrow, blood, or other tissue may be treated with one or more anticoagulants to prevent or minimize clotting. For example and without limitation, anticoagulants include acenocoumarol, antithrombin III, apixaban, argatroban, atromentin, betrixaban, bivalirudin, brodifacoum, dabigatran, dalteparin, difenacoum, edoxaban, EDTA, enoxaparin, fondaparinux, 5 heparin, idraparinux, phenindione, phenprocoumon, rivaroxaban, and warfarin. The anticoagulant may be administered to the body of the deceased donor. Alternatively or additionally, the bone marrow, blood, or other tissue may be treated with the anticoagulant after removal from the body.

Bone marrow, blood, or other tissue may be treated to deplete red blood cells and/or 10 platelets. For example and without limitation, red blood cells and/or platelets may be depleted by buoyancy-activated cell separation, cell lysis, hetastarch sedimentation, immunomagnetic depletion, size-based centrifugal separation, or spinning membrane filtration, as described above.

Methods for obtaining CD3⁺ cells from blood may include separating blood into different constituents, such as a cellular fraction and a plasma fraction, as described in detail below.

15 Obtaining CD3⁺ cells may include other methods of enriching blood or other non-bone marrow sources for CD3⁺ cells. Enrichment may include positive selection of CD3⁺ cells, depletion of non-CD3⁺ cells, or a combination thereof. For example and without limitation, CD3⁺ cells may be positively selected by use of an antibody or other agent that binds a marker on the surface of CD3⁺ cells, such as CD3, CD4, or CD8. For example and without limitation, 20 depletion of non-CD3⁺ cells may be depleted by use of an antibody agent that that binds a marker absent from the surface of CD3⁺ cells, such as CD10, CD14, CD15, CD33, CD41, CD71, CD209, or CD235. Positive selection or depletion may be performed by binding antibodies conjugated to particles or beads to subpopulations of cells and sorting cell sub-populations by methods known in the art, such as those described in U.S. Patent No. 9,090,871; U.S. Patent 25 Publication No. 2010/0310588; and International Patent Publication No. WO 2017/005647, the contents of each of which are incorporated herein by reference.

During preparation of CD34⁺ cells and/or CD3⁺ cells for use in cellular products, cells may be frozen, i.e., cryopreserved, at any stage. Cryopreservation may include addition of one or more cryoprotectants, such as those described above in relation to cellular products of the 30 invention. Cryopreservation typically involves reducing the temperature of the cell-containing sample at a controlled rate. Cryopreservation may include thawing the cell-containing sample

and washing the sample to remove one or more cryoprotectants. Methods and reagents for cryopreservation, including freezing, thawing, and washing samples, are known in the art and described in, for example, U.S. Patent No. 9,561,253, the contents of which are incorporated herein by reference.

5 In some circumstances, the number of CD34⁺ cells and/or CD3⁺ cells initially obtained from a deceased donor may be insufficient to generate a product to promote mixed chimerism in a living recipient. Therefore, the methods may include expanding CD34⁺ cells and/or CD3⁺ cells ex vivo. Any desired cell type or population of cells may be expanded. For example and without limitation, the population expanded may include HSPCs, T cells, T regulatory (T_{reg})
10 cells, memory T cells, effector memory T cells, central memory T cells, stem memory T cells, naïve T cells, cytotoxic T cells, gamma delta T cells, B cells, natural killer (NK) cells, natural killer T (NKT) cells, megakaryocytes, myeloblasts, monoblasts, monocytes, macrophages, dendritic cells, CD34⁺ cells, CD3⁺ cells, or CD4⁺ cells. Expansion may occur prior to, or
15 subsequent to, freezing. Expansion may include providing one or more growth factors, and it may include culturing cells in the presence of another cell type, e.g., feeder cells. Methods for expanding hematopoietic cells are described in, for example, U.S. Patent No. 9,561,253, the contents of which are incorporated herein by reference.

 CD34⁺ cells and/or CD3⁺ cells may be genetically modified ex vivo. For example, in autologous transfer of donor cells, a genetic defect may be corrected using gene therapy.
20 Methods of gene therapy are described in, for example, Mali, Delivery systems for gene therapy, Indian J Hum Genet. 2013 Jan-Mar; 19(1): 3–8, doi: 10.4103/0971-6866.112870; Gennady Ermak (2015) FRONT MATTER. Emerging Medical Technologies, ISBN: 978-981-4675-80-2, doi.org/10.1142/9789814675826_fmatter; and Bakhuraysah et al., Hematopoietic stem cell transplantation for multiple sclerosis: is it a clinical reality? Stem Cell Res Ther. 2016; 7:12, doi:
25 10.1186/s13287-015-0272-1, the contents of each of which are incorporated herein by reference.

Ex vivo extraction of hematopoietic cells from bone marrow

 Methods of the invention may include extracting hematopoietic cells, such as CD34⁺ cells, from the bone marrow of a deceased donor for use in manufacture of cellular products to
30 transfer to a living recipient. In certain embodiments, hematopoietic cells are extracted from bone marrow ex vivo. Ex vivo extraction may include contacting a bone sample with an agent

that mobilize hematopoietic cells, such as one described above, while the bone sample is in a container.

Such methods involve obtaining a bone sample from a deceased donor. The sample may be from any large bone that contains the cancellous material, such as the pelvis, vertebrae, ribs, femur, tibia, and sternum. Preferably, the sample is from the iliac crests of the pelvis and vertebral bodies of the vertebrae.

The methods may further involve obtaining bone marrow cells from the bone sample after it has been removed from the body. Bone marrow cells may be obtained from the bone sample by any method. For example, bone marrow cells may be obtained by trephination or aspiration, as described above. Samples obtained by trephination may be treated to remove bone shards, as described above.

The methods may include treating the bone sample with an agent that mobilizes CD34⁺ cells from bone marrow, such as one of the mobilizing agents described above.

The methods may be used to obtain any type of hematopoietic cells. For example and without limitation, the cells may be B cells, basophils, eosinophils, hematopoietic cells, hematopoietic stem and progenitor cells (HSPCs), lymphocytes, lymphoid progenitor cells, macrophages, mast cells, megakaryocytes, monocytes, myeloblasts, myeloid progenitor cells, natural killer (NK) cells, neutrophils, platelets/thrombocytes, T cells, T regulatory (T_{reg}) cells, memory T cells, effector memory T cells, central memory T cells, stem memory T cells, naïve T cells, cytotoxic T cells, gamma delta T cells, natural killer T cells, CD34⁺ cells, CD4⁺ cells, or CD3⁺ cells. Preferably, the cells are HSPCs or CD34⁺ cells.

The methods may include one or more additional steps, such as those described above in relation to preparation of cellular products. For example, the methods may include any of the following: treating the hematopoietic cells with an anticoagulant, such as one described above; depleting the hematopoietic cells or red blood cells and/or platelets; and enriching the hematopoietic cells for a cell type or population of cells, such as CD34⁺ cells.

The methods may include using the hematopoietic cells extracted from bone marrow ex vivo for use in manufacture of a product for transfer into a living recipient, such as one of the cellular compositions described above.

The invention also includes materials useful for ex vivo extraction of hematopoietic cells, such as CD34⁺ cells, from bone marrow of deceased donors. For example, the invention

provides receptacles that are pre-loaded with one or more agents that mobilize CD34⁺ cells from bone marrow, such as one of the mobilizing agents described above. The invention also includes systems that include a container that contains one or more such agents and one or more portions of bone derived from a deceased donor.

5 The materials and systems for ex vivo extraction of hematopoietic cells may be characterized by their ability to release particular cell type or population of cells into suspension over time. For example, the materials and systems may increase the release of CD34⁺ cells from bone marrow as compared to a reference storage system by 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 20-fold, 50-fold, 100-fold, 200-fold, 500-fold, 1000-fold, or more.

10 The materials and systems for ex vivo extraction of hematopoietic cells may achieve release of a particular cell type or population of cells into suspension in a defined period, such as 1 hour, 2 hours, 4 hours, 8 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, or more.

 The materials and systems may achieve release into suspension of a particular cell type or population of cells having characteristics that improve the utility of the cells for manufacture of products to promote mixed chimerism. For example and without limitation, the cells may have superior viability, proliferation potential, marker expression, apoptotic characteristics, and/or apoptotic characteristics as compared to cells from a reference storage system.

15

Fractionation of blood from deceased donors

20 Methods of the invention may include fractionating blood from deceased donors into two or more constituents and using one or more of the constituents in the manufacture of products for transfer into living recipients. Apheresis of blood from live donors involves separation of blood into different constituents, isolating one or more constituents, and returning the remainder to circulation. For example, leukapheresis entails the isolation of white blood cells from the blood of a donor and returning the remaining cells and plasma to the donor's body. Apheresis methods and devices are known in the art and described in, for example, U.S. Publication No. 2002/0107469; U.S. Patent No. 5,607,579; U.S. Patent No. 9,364,600; and U.S. Patent No. 6,743,192, the contents of each of which are incorporated herein by reference.

25

 The invention includes methods in which procedures analogous to apheresis are used to separate components of blood from deceased donors. An advantage of using blood from deceased donors is that the unused portion of blood need not be returned to the donor's body after

30

isolation of the component of interest. For example, blood from a deceased donor may be separated into a cellular fraction and a plasma fraction or into two or more different cellular fractions. The blood may be fractionated sequentially. For example, after initial separation into cellular and non-cellular fractions, the cellular fraction may be further processed to enrich for a particular cell type of population of cells, such as CD3⁺ cells. Any method known in the art may be used to enrich for CD3⁺ cells, including any of the methods described above.

Separating blood from a deceased donor into different fractions is useful for obtaining fractions that are enriched for particular components, such as cell types or populations of cells, to be included in cellular products of the invention. The cells may be any type of hematopoietic cells, such as B cells, basophils, eosinophils, hematopoietic cells, hematopoietic stem and progenitor cells (HSPCs), lymphocytes, lymphoid progenitor cells, macrophages, mast cells, megakaryocytes, monocytes, myeloblasts, myeloid progenitor cells, natural killer (NK) cells, neutrophils, platelets/thrombocytes, T cells, T regulatory (T_{reg}) cells, memory T cells, effector memory T cells, central memory T cells, stem memory T cells, naïve T cells, cytotoxic T cells, gamma delta T cells, natural killer T cells, CD34⁺ cells, CD4⁺ cells, or CD3⁺ cells. In particular, blood fractionation is useful for obtaining fractions enriched for CD3⁺ cells for use in manufacture of cellular products. Fractionation of blood is also useful to obtain fractions enriched for CD34⁺ cells, which can be used to supplement the bone marrow-derived CD34⁺ cells in products of the invention.

The methods of fractionation methods may be performed on blood that has been supplemented with additional components from the deceased donor. For example, cells extracted from bone marrow or other tissues, as described above, may be combined with blood to enrich for a desired cell type of population of cells in the mixture. For example and without limitation, the population enriched may include HSPCs, T cells, T regulatory (T_{reg}) cells, memory T cells, effector memory T cells, central memory T cells, stem memory T cells, naïve T cells, cytotoxic T cells, gamma delta T cells, B cells, natural killer (NK) cells, natural killer T (NKT) cells, megakaryocytes, myeloblasts, monoblasts, monocytes, macrophages, dendritic cells, CD34⁺ cells, CD3⁺ cells, or CD4⁺ cells. The cells extracted from bone marrow or other tissue may have been subjected to any of the procedures described above to enrich for CD34⁺ cells and CD3⁺ cells. The supplemented blood may then be separated into fractions, as described above.

Analysis of blood from deceased donors

Methods of the invention may include analytical methods to determine whether blood from deceased donors to determine whether the blood is suitable for use in manufacture of a product to administer to a living recipient. The methods may involve analysis of the blood itself, or they may involve analysis of other tissue from the deceased donor. The analytical methods may be combined with any of the preparative methods described above.

The methods are useful for determining whether material in blood is suitable for use in manufacture of products for transfer to a living recipient. Preferably, the material from the blood is a cell type or population of cells. For example and without limitation, the cells may be B cells, basophils, eosinophils, hematopoietic cells, hematopoietic stem and progenitor cells (HSPCs), lymphocytes, lymphoid progenitor cells, macrophages, mast cells, megakaryocytes, monocytes, myeloblasts, myeloid progenitor cells, natural killer (NK) cells, neutrophils, platelets/thrombocytes, T cells, T regulatory (T_{reg}) cells, memory T cells, effector memory T cells, central memory T cells, stem memory T cells, naïve T cells, cytotoxic T cells, gamma delta T cells, natural killer T cells, CD34⁺ cells, CD4⁺ cells, or CD3⁺ cells. The analytical methods may be combined with any of the preparative methods described above. Alternatively or additionally, the material from the blood may be a non-cellular component, such as a molecule. For example and without limitation, the molecule may be a cytokine, chemokine, antibody, or immunoglobulin.

Analysis of several characteristics of T cells from deceased donors is useful for determining whether T cells will promote mixed chimerism in a transplant recipient. For example and without limitation, analyses of T cell activation, anergy, apoptosis, exhaustion, proliferation, viability, and cytokine secretion provide information that may be used to assess whether the T cells may provide a therapeutic benefit to a living recipient.

Characterization of T cells or CD3⁺ cells from a deceased donor may include analysis of cells in the blood. Any suitable assay may be used. For example and without limitation, the methods may include proliferation assays, DNA synthesis assay, apoptosis assays, colony formation assays, marker expression assays, and the like. Methods of analysis of T cells are known in the art and described in, for example, Brousseau, et al., eds., *Manual of Immunological Methods*, CRC Press, Boca Raton, 1998, ISBN 9780849385582; Detrick, et al., eds., *Manual of*

Molecular and Clinical Laboratory Immunology, Eighth Edition, American Society for Microbiology Press, Washington D.C., USA, 2016, ISBN-13: 978-1555818715; and Balakrishnan, et al., Practical Immunology: A Laboratory Manual, Lap Lambert Academic Publishing, 2017, ISBN-13: 978-3330352032, the contents of each of which are incorporated
5 herein by reference.

Characterization of T cells or CD3⁺ cells from a deceased donor may include analysis of a non-cellular component of the blood. For example and without limitation, the non-cellular component may be a cytokine, e.g., a pro-inflammatory or anti-inflammatory cytokine, a chemokine, an antibody, or an immunoglobulin. Non-cellular components may be analyzed in
10 serum or plasma from the deceased donor. Non-cellular components may be analyzed by immunoassays, such as an enzyme-linked immunoassay (ELISA), radioimmunoassay, immunoPCR, and immunoassay systems sold under the trade names Luminex xMAP by ThermoFisher Scientific (Waltham, MA); Meso Scale Discovery by Meso Scale Diagnostics (Rockville, MD); AlphaLISA and DELFIA by Perkin Elmer (Waltham, MA); Gyrolab by Gyros
15 Protein Technologies (Uppsala, Sweden); and Erenna by MilliporeSigma (Burlington, MA). Other methods of analysis of non-cellular components are known in the art and described in, for example, Brousseau, et al., eds., Manual of Immunological Methods, CRC Press, Boca Raton, 1998, ISBN 9780849385582; Detrick, et al., eds., Manual of Molecular and Clinical Laboratory Immunology, Eighth Edition, American Society for Microbiology Press, Washington D.C., USA,
20 2016, ISBN-13: 978-1555818715; and Balakrishnan, et al., Practical Immunology: A Laboratory Manual, Lap Lambert Academic Publishing, 2017, ISBN-13: 978-3330352032, the contents of each of which are incorporated herein by reference.

The methods may include analysis of whether HSPCs or CD34⁺ cells in the blood from a deceased donor are suitable for use in manufacture of products for transfer into living recipients.
25 Properties such as HSPC proliferation, HSPC viability, and HSPC apoptosis may be analyzed to determine whether HSPCs in the blood may provide a therapeutic benefit may be used to assess whether HSPCs may provide a therapeutic benefit to a living recipient. For example and without limitation, analysis may include colony-forming assays or long-term culture assays that measure cell growth in response to exogenous growth factors. Analysis may include assays for the ability
30 of HSPCs to repopulate themselves in a mouse model, such as a severe-combined immunodeficiency (SCID) mouse. These and other methods are known in the art and described

in, for example, Loring and Peterson, eds., *Human Stem Cell Manual: A Laboratory Guide*, Second Edition, Academic Press, 2012, ISBN: 978-0-12-385473-5; Stein, et al., eds., *Human Stem Cell Technology and Biology: A Research Guide and Laboratory Manual*, Wiley-Blackwell, 2011, ISBN 978-0-470-59545-9; Frisch and Calvi, *Hematopoietic Stem Cell Cultures and Assays*, *Methods Mol Biol.* 2014; 1130: 315–324, doi: 10.1007/978-1-62703-989-5_24; 5 Dick JE, et al., Assay of human stem cells by repopulation of NOD/SCID mice, *Stem Cells*, 1997;15 Suppl 1:199-203; discussion 204-7, DOI: 10.1002/stem.5530150826; and T Tatekawa, et.a., (2006) A novel direct competitive repopulation assay for human hematopoietic stem cells using NOD/SCID mice, *Cytotherapy*, 8:4, 390-398, DOI: 10.1080/14653240600847191, the 10 contents of each of which are incorporated herein by reference.

The analysis may include providing a report of the characteristics of the blood or non-blood tissue from the deceased donor. For example and without limitation, the report may include percentages of cells that have any of the following properties: expression of a marker, viability, proliferation potential, apoptotic characteristics, anergy, exhaustion, activation, and 15 cytokine secretion. The report may describe characteristics of sub-populations of cells. For example and without limitation, the report may include percentages of viability, proliferation potential, apoptotic characteristics, anergy, exhaustion, activation, or cytokine secretion of cells that express a particular marker, such as CD34⁺ cells, or CD3⁺ cells.

Determination of whether material in the blood is suitable for use in manufacture of 20 product for transfer into living recipients may include analysis of non-blood tissue from the deceased donor. For example and without limitation, the non-blood tissue may be bone marrow, spleen, liver, lymph nodes, or thymus. Cellular components, e.g., hematopoietic cells, such as those described above, or non-cellular components, such as the molecules described above, of non-blood tissue may be analyzed. Any analytical methods, including those described above 25 may be used.

The content and location of hematopoietic cells in the body vary during growth and development. For example, the spleen is a rich source of lymphocytes in adults, whereas the liver contains many lymphocytes in the developing fetus. Therefore, the tissue source of hematopoietic cells may account for the age of the donor at the time of death.

30 The methods may include analysis of multiple components. The multiple components may come from the same source, or they may come from different sources. For example, the

methods may include analysis of multiple cellular components of the blood, multiple non-cellular components of the blood, multiple components of a non-blood tissue, or any combination of a cellular blood component, a non-cellular blood component, and a component of non-blood tissue.

5 To better determine whether components from the blood of a deceased donor are suitable for use in manufacture of products for transfer to a living recipient, analysis may include comparison of one or more parameters from a tissue or fluid of a deceased donor to the corresponding material from a living donor. Analysis may include a comparison of any characteristic described above. For example and without limitation, analysis may include
10 comparison of one or more of T cell activation, T cell anergy, T cell apoptosis, T cell exhaustion, T cell proliferation, T cell viability, T cell cytokine secretion, HSPC proliferation, HSPC viability, HSPC apoptosis, cytokine levels, chemokine levels, antibody levels, and immunoglobulin levels.

The determination of whether blood or a blood component from a deceased donor is
15 suitable for use in manufacture of products for transfer to a living recipient may rely on a comparison of one or more parameters between the deceased donor and a living donor. For example, when the value of one more parameters from a deceased donor is comparable to the corresponding values from a living donor or population of living donors, the blood or blood component from a deceased donor may be deemed suitable for use in manufacture of products
20 for transfer to a living recipient. Comparative analysis may include establishing a profile of characteristics for living donors and for donors at various temporal stages following brain death. Values may be compared between a deceased donor and living donor or population of living donors that meet one or more criteria. For example, the living donor or population of living donors may have done one or more of the following: previously donated a solid organ;
25 previously donated hematopoietic cells, such as HSPCs or T cells; or previously donated tissue that supported establishment of mixed chimerism in a recipient. The living donor or population of living donors may be healthy, within a particular age range, of a certain sex, or the like.

The methods may include using one or materials from the blood of a deceased donor in the manufacture of a product for transfer to a living recipient. The materials may be any
30 materials described above, such as cellular or non-cellular components.

The methods may include expansion of a cell type or population of cells for use in the manufacture of a product for transfer to a living recipient. For example, the cells may be B cells, basophils, eosinophils, hematopoietic cells, hematopoietic stem and progenitor cells (HSPCs), lymphocytes, lymphoid progenitor cells, macrophages, mast cells, megakaryocytes, monocytes, myeloblasts, myeloid progenitor cells, natural killer (NK) cells, neutrophils, platelets/thrombocytes, T cells, T regulatory (T_{reg}) cells, memory T cells, effector memory T cells, central memory T cells, stem memory T cells, naïve T cells, cytotoxic T cells, gamma delta T cells, natural killer T cells, CD34⁺ cells, CD4⁺ cells, or CD3⁺ cells.

The methods may include treating blood from a deceased donor to mitigate damage to material in the blood that may be used in manufacture of a product for transfer into living recipients. For example, the methods may include adding an anticoagulant, buffer, cryoprotectant, or stabilizer to the blood. The stabilizer may prevent degradation of blood component or may promote osmotic stability. For example, the stabilizer may be human serum albumin (HSA) or the electrolyte solution sold under the trade name Plasma-Lyte by Baxter International Inc. (Deerfield, IL).

The methods may include cryopreserving the products. The products may be frozen at a temperature suitable for long-term storage, such as about -150 °C. The methods may include gradually reducing the temperature of the product to the long-term storage temperature.

20 Devices and methods for collection of blood from deceased donors

Methods of the invention may include specific devices and methods for collection of blood from a deceased donor. Blood must be expeditiously processed for use as a source of material for transfer into a living recipient. In particular, it must be removed from the body rapidly after brain death, preferably within hours, before the release of cytokines that can alter the function of blood cells. In addition, blood must be rapidly chilled to a suitable short-term storage temperature, typically 2-8 °C, upon removal from the body, but freezing should be avoided until the blood or blood products are ready for cryopreservation. Given that the total blood volume of an adult human may be 5 liters or more and that maximal recovery is needed to obtain sufficient quantities of CD34⁺ and CD3⁺ cells, collection of blood from deceased donors presents unique challenges.

FIG. 3 shows a blood collection device 301 that may be used in methods of the invention. The device 301 includes one or more receptacles 303a and 303b coupled to a cooling system. The receptacles may have a combined capacity of about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 liters, and preferably from about 5 to about 10 liters. The devices may have 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,
5 or more receptacles. The receptacles may be rigid and have a fixed shape, or they may be flexible and expandable. For example and without limitation, the receptacles may be cups, bottles, cylinders, bags, pouches, or the like. The receptacles may be made from a flexible or pliable plastic. In an embodiment, each receptacle is a rectangular prism of 25 cm x 20 cm x 2 cm and has a capacity of 1 liter.

10 The device 301 may include one or more cannulae 305 that can be inserted into a vein or artery of the deceased donor. The cannula may be connected to one or more receptacles 303a and 303b by a connector 307a, such as tubing. In a device 301 having multiple receptacles 303a and 303b, blood may be collected into the receptacles 303a and 303b in parallel, sequentially, or some combination thereof. In devices that have receptacles arranged in sequence, the receptacles
15 303a and 303b may be joined by connectors 307b. One or more of the connectors 307a and 307b may include a one-way valve to prevent reverse flow of blood through the device 301, thereby minimizing contamination of the blood and maintaining its sterility.

FIG. 4 shows a blood collection device 401 that may be used in methods of the invention. The device includes receptacle 403a, 403b, 403c, and 403d disposed within a refrigeration
20 compartment 409. The refrigeration compartment 409 includes a coolant 411 that cools blood after its removal from the donor. Any suitable coolant 411 may be used. For example and without limitation, the coolant 411 may include ammonium chloride, ammonium nitrate, brine, calcium ammonium nitrate, diethylene glycol, hydroxyethyl cellulose, ice, propylene glycol silica gel, sodium polyacrylate, urea, or water.

25 The cooling system is configured to cool blood in the receptacles to a target temperature. The target temperature may from about 2 °C to about 8 °C. The cooling system may prevent blood from freezing or formation of ice crystals within the blood. The cooling system may cool blood to the target temperature within a specified period. For example, the cooling system may cool blood to the target temperature in about 4 hours, about 5 hours, about 6 hours, about 7
30 hours, about 8 hours, about 9 hours, about 10 hours, not less than 4 hours, not less than 5 hours,

not less than 6 hours, not less than 7 hours, not less than 8 hours, not more than 6 hours, not more than 7 hours, not more than 8 hours, not more than 9 hours, or not more than 10 hours.

The device may include a vacuum system configured to apply a vacuum to remove blood from the deceased donor. The vacuum system may be a pump. Vacuum pumps for blood
5 collection are known in the art and described in, for example, U.S. Patent Publication No. 2010/0042015; and U.S. Patent Publication No. 2008/0199949, the contents of each of which are incorporated herein by reference. The vacuum system may be coupled to one or more of the needle, tubing, and receptacle.

The methods may include transferring blood to one or more receptacles coupled to a
10 cooling system. The methods may include inserting a needle into a blood vessel, artery, or vein of the deceased donor. The methods may include applying a vacuum to the blood vessel, artery, or vein of the deceased donor. The methods may include contacting the blood with a stabilizing agent or cryoprotectant, such as one described above.

15 **Treating organ transplant recipients using cellular products derived from deceased donors**

The cellular products made by methods of the invention may be provided to the recipient of a solid organ transplant. The cellular product may be provided by any suitable means. For example and without limitation, the CD34⁺ cells and/or CD3⁺ cells may be delivered to the recipient by injection using a needle, catheter, central line, or the like. In some cases, the cells
20 may be delivered intravascularly, intravenously, intraarterially, subcutaneously, intramuscularly, directly to the bone, or through any source which permits the hematopoietic cells to home to an appropriate site in the recipient such that the hematopoietic cells persist, regenerate and differentiate in the recipient. The CD34⁺ cells and/or CD3⁺ cells may be provided by infusion. The CD34⁺ cells and/or CD3⁺ cells may be provided in an inpatient procedure or in an outpatient
25 procedure. An inpatient procedure requires admission to a hospital, and the patient may spend one or more nights in the hospital. An outpatient procedure does not require admission to a hospital and may be performed in a non-hospital setting, such as a clinic, doctor's office, home, or other location.

The compositions made by methods of the invention may be used in conjunction with
30 transplantation of any solid organ. For example and without limitation, the solid organ may be a kidney, lung, pancreas, pancreatic, islet cells, heart, intestine, colon, liver, skin, muscle, gum,

eye, or tooth. The transplant may include a complete organ, a portion of an organ, or cells from a tissue of an organ. The cellular product may be provided prior to, during, or subsequent to the solid organ transplant. For example and without limitation, the cellular product may be provided one, two, three, four, five, or six days or one, two, three, or four weeks prior to the solid organ
5 transplant, or it may be provided one, two, three, four, five, or six days or one, two, three, or four weeks after the solid organ transplant.

To facilitate establishment of mixed chimerism in the recipient, the recipient's immune system may be conditioned in conjunction with providing the cellular product. For example, non-myeloablative conditioning may be used. In non-myeloablative conditioning, the recipient
10 is exposed to drugs, antibodies, irradiation, or some combination thereof at a dose that is too low to eradicate all the bone marrow cells. Typically, the conditioning regimen includes treatment with anti-thymocyte globulin (ATG), total lymphoid irradiation, and corticosteroids (e.g. prednisone) for a period of from about 10 to 12 days (e.g. for about 11 days). The irradiation may be targeted to a particular location of the recipient's body. For example, irradiation may be
15 targeted to a tissue, an organ, a region of the body or the whole body. Irradiation may be targeted to the lymph nodes, the spleen, or the thymus or any other area known to a person of skill in the art. When multiple doses of irradiation are administered, the doses may be targeted to the same location or to different locations. Non-myeloablative conditioning may include the use of a T cell depleting agent, such as a monoclonal antibody or drug, e.g., fludarabine. Regimens
20 for non-myeloablative conditioning are known in the art and are described in, for example, U.S. Patent No. 9,561,253, the contents of which are incorporated herein by reference.

Methods of treatment may include immunosuppressive therapy. Immunosuppressive therapy, or immunosuppression, involves treatment of the graft recipient with agents that diminish the response of the host immune system against the donor cells, which can lead to graft
25 rejection. Primary immunosuppressive agents include calcineurin inhibitors, such as tacrolimus, cyclosporin A. Adjuvant agents are usually combined with a calcineurin inhibitor. Adjuvant agents include steroids, azathioprine, mycophenolic acid (MPA) agents, such as mycophenolate mofetil, mTOR inhibitors, such as sirolimus, and belatacept. The use of adjuvant agents allows clinicians to achieve adequate immunosuppression while decreasing the dose and toxicity of
30 individual agents. Antibody-based therapy may use monoclonal (e.g., muromonab-CD3) or polyclonal antibodies or anti-CD25 antibodies (e.g., basiliximab, daclizumab). Antibody-based

therapy allows for avoidance or dose reduction of calcineurin inhibitors, possibly reducing the risk of nephrotoxicity. Regimens for immunosuppressive therapy are known in the art and are described in, for example, U.S. Patent No. 9,561,253, the contents of which are incorporated herein by reference.

5 Immunosuppression may also diminish the response of the donor immune cells against recipient tissue, which can lead to GVHD. GVHD may be acute or chronic. Acute GVHD typically occurs in the first 3 months after graft and may involve the skin, intestine, or the liver. Treatment for acute GVHD usually includes high-dose corticosteroids such as prednisone. Chronic GVHD typically occurs after the first 3 months following transplant and is the major
10 source of late treatment-related complications. Chronic GVHD may cause functional disability and require prolonged immunosuppressive therapy.

 Immunosuppressive therapy may occur in multiple phases. For example, the immunosuppressive regimen may have an induction phase and a maintenance phase. Induction and maintenance phase strategies may use different medicines at doses adjusted to achieve target
15 therapeutic levels to enhance establishment of mixed chimerism in the recipient.

 Immunosuppressive therapy may be withdrawn after stable mixed chimerism has been established in the recipient. The chimeric status of the recipient may be monitored as described below and deemed stable after a certain period, for example, 3 months, 6 months 12 months, 18 months, 24 months, or longer. Thus, immunosuppression may be discontinued for the recipients
20 after a certain period, for example, 3 months, 6 months 12 months, 18 months, 24 months, or longer. Withdrawal of immunosuppressive therapy may include tapering, i.e., progressively reducing the dosage or frequency of treatment.

 A determination of whether an individual is a full chimera, mixed chimera, or non-chimera made be made by an analysis of a hematopoietic cell sample from the solid organ
25 transplant recipient, e.g. peripheral blood, bone marrow, etc. as known in the art. Analysis may be done by any convenient method of typing. Analysis may be performed on hematopoietic cells or a subset thereof, such as all mononuclear cells, T cells, B cells, CD56⁺ NK cells, and CD15⁺ neutrophils. Chimerism can be assessed by PCR analysis of microsatellites. For example, commercial kits that distinguish polymorphisms in short terminal repeat lengths of donor and
30 host origin are available. Automated readers provide the percentage of donor type cells based on standard curves from artificial donor and host cell mixtures.

Recipients may be categorized as fully chimeric, mixed chimeric, or non-chimeric based on the fraction of cells that are derived from the donor. For example, recipients can be deemed fully chimeric if they have at least 90%, at least 95%, at least 98%, or at least 99% donor-derived cells. Recipients can be deemed mixed chimeric if they have too few donor-derived cells to be
5 categorized as fully chimeric but a fraction of donor-derived cells that exceeds a certain threshold, such as at least 0.5%, at least 1%, at least 2%, at least 3%, at least 5%, at least 7.5%, at least 10% donor-derived cells. Recipients can be deemed non-chimeric if the fraction of donor-derived cells falls below the threshold required to be categorized as mixed chimeric.

10 **Transfer of cellular products derived from a single deceased donor to multiple recipients**

Methods of the invention may be used to make multiple cellular products derived from a single deceased donor for transfer to multiple living recipients. Pre-treatment of the deceased subject's body with one or more agents that mobilize hematopoietic cells from bone marrow to peripheral blood improves recovery of HSPCs, thus providing quantities of HSPCs sufficient to
15 make multiple cellular products. The generation of multiple cellular products containing CD34⁺ cells and CD3⁺ cells from a single donor allows the products to be transferred to different recipients. Thus, two or more solid organs from the deceased donor may be transplanted into different recipients, and each recipient may also receive donor CD34⁺ cells and CD3⁺ cells that promote establishment mixed chimerism. For example, the products and methods of the
20 invention allow a single deceased donor to provide a kidney and product containing CD34⁺ cells and CD3⁺ cells to each of two recipients. Consequently, the products and methods of the invention increase the number of patients who can benefit from organ transplantation procedures without requiring additional organ donors.

A single deceased donor may be the source of material for 1, 2, 3, 4, 5, or more cellular
25 products that promote mixed chimerism in a solid organ transplant recipient. The multiple cellular products from a single deceased donor may comprise any element of the cellular products described above, such as cell type, cell number, degree of HLA matching or mismatching, cell purity, cryoprotectants, stabilizing agents, or graft-enhancement agents.

The invention also provides methods of preparing multiple cellular products from a single
30 deceased donor for transfer to multiple living recipients. The methods include obtaining CD34⁺ cells from bone marrow of a deceased donor and CD3⁺ cells from non-bone marrow of the

donor. The methods may include any element described above in relation to preparation of cellular products, ex vivo extraction of CD3⁺ cells from non-bone marrow, fractionation of blood, or analysis of blood. For example and without limitation, the methods may include one or more of removal of bone marrow, separation of cells from bone shards, treatment with an agent
5 that mobilizes CD34⁺ cells, separation of mononuclear cells from enucleated or multinucleated cells, depletion of RBCs and platelets, enrichment of CD34⁺ cells and/or CD3⁺ cells, removal of clots and/or cell clumps, treatment with an anticoagulant, cryopreservation, ex vivo expansion of cells, genetic modification of cells, and analysis of T cell activation, T cell anergy, T cell apoptosis, T cell exhaustion, T cell proliferation, T cell viability, T cell cytokine secretion, HSPC
10 proliferation, HSPC viability, HSPC apoptosis, cytokine levels, chemokine levels, antibody levels, and/or immunoglobulin levels.

Examples

Example 1

15 A method of preparing a cellular composition containing CD34⁺ cells derived from bone marrow of a deceased donor and CD3⁺ cells derived from blood of the deceased donor according to an embodiment of the invention is described below. The method includes a series of steps, which are described in more detail below. The method includes the following steps: collection of tissue; preparation of bone marrow; preparation of blood; preparation of buffer; enrichment of
20 CD34⁺ cells from bone marrow; dividing the combined material into individual doses; and cryopreservation. The sequence of steps described herein is for illustrative purposes only. Although it will be understood by one of skill in the art that certain steps must be performed prior to others, the methods are not limited to a particular sequence of steps.

25 FIG. 5 is a flow diagram of the tissue collection step according to a method of the invention. Blood, bone marrow, and one or more solid organs of interest are harvested from the body of a deceased donor. Bone marrow is extracted by trephination, as described above. Blood is collected into one or more receptacles of devices of the invention, as described above. Blood or other material from the body is tested for infectious disease markers (IDMs).

30 FIG. 6 is a flow diagram of the bone marrow preparation step according to a method of the invention. The extracted bone marrow is passed through a sieve to remove bone shards, and it is depleted of red blood cells by one of the methods described above. The bone marrow is also

analyzed for cell number, sterility, viability, colony-forming ability, and expression of CD3, CD34, and CD45 markers. Viability is assayed by exclusion of 7-aminoactinomycin D (7-AAD), and marker expression is analyzed by flow cytometry.

5 FIG. 7 is a flow diagram of the step of the blood preparation step according to a method of the invention. The blood is depleted of red blood cells and platelets by one of the methods described above. The blood is also analyzed for cell number, sterility, viability, and expression of CD3, CD34, and CD45 markers. Viability is assayed by exclusion of 7-AAD, and marker expression is analyzed by flow cytometry.

10 FIG. 8 is a flow diagram of the buffer preparation step according to a method of the invention. Two buffers are prepared, one that contains HSA and another that contains a balanced electrolyte solution.

FIG. 9 is a flow diagram of the CD34⁺ cell-enrichment step according to a method of the invention. The buffers are used with the cells obtained from the bone marrow preparation step to enrich for CD34⁺ cells by spinning membrane filtration. The sample is further enriched for CD34⁺ cells by immunomagnetic selection. Cell number, viability, colony-forming ability, and marker expression is analyzed as described above, either prior to or following spinning membrane filtration. Cells from the blood preparation step are analyzed for cell number, viability, marker expression, T cell activation, T cell proliferation, and T cell apoptosis, as described above. If the bone marrow-derived cells enriched for CD34⁺ cells and the blood-derived cells containing CD3⁺ cells are both determined to be suitable for use in a cellular product, the two cellular fractions are combined.

15
20

FIG. 10 is a flow diagram of the dividing step according to a method of the invention. Pooled cells are filtered to remove clots and cell clumps, i.e., agglutination. Pooled cells are then divided into two individual doses, each of which contains sufficient quantities of CD34⁺ cells and CD3⁺ cells to promote mixed chimerism in a separate organ transplant recipient. During this step, cell number, viability, and marker expression is analyzed, as described above. Cells may also be analyzed for sterility, endotoxins, mycoplasma, colony-forming ability, release of cytokines, e.g., IL-2.

25

FIG. 11 is a flow diagram of the cryopreservation step according to a method of the invention. Doses may be further divided into aliquots, or they may be maintained in a single container. Doses are frozen by in a freezer that reduces the temperature of the cellular products

30

to a target temperature at a controlled rate. After the cellular products have reached the target temperature, they are transferred to liquid nitrogen for long-term storage.

Example 2

5 An exemplary process for preparing a cellular product, MDR-104, containing CD34⁺ cells and CD3⁺ cells is described below. The process provides two doses, each of which contains 100 x 10⁶ T cells/kg recipient weight and > 4 x 10⁶ CD34⁺ cells/kg recipient weight, from a single deceased donor. Each dose supports transplantation of one kidney from the deceased donor into a different recipient. The product also includes > 500 colony-forming units (CFUs) per 10⁵ cells. The product may be further characterized to determine IL-2 release and percentage of viable CD34⁺ cells and CD3⁺ cells.

FIG. 12 is a flow diagram illustrating a method of preparing a cellular composition according to an embodiment of the invention. As described in more detail below, CD3⁺ cells are prepared from whole blood in MDR-T processing, CD34⁺ are prepared from bone marrow in 15 MDR-E processing, and the prepared CD3⁺ cells and HSPCs are combined in MDR-34T processing.

Whole blood (WB) and bone marrow (BM) are collected as starting material to supply the requisite CD3⁺ T cells and CD34⁺ HSPCs, respectively, in the cellular compositions of the final drug product. The exsanguinated blood is collected following systemic heparinization of 20 the deceased donor into either a single collection device or several smaller collection devices to permit ease of transport from the collection site to the manufacturing site. Whole blood is subsequently processed using an apheresis instrument for platelet and red blood cell depletion, then washed for additional platelet removal either prior to or following the addition of the CD34⁺-selected, in-process intermediate.

25 Bone marrow is harvested via one or both of two methods. One method employs standard bone marrow aspiration from the exposed iliac crests of the deceased donor. In the second method, marrow is recovered via the trephine orthopedic device, which is capable of harvesting cores of cortical and cancellous bone containing BM. Bone marrow may also be recovered from deceased donor vertebral bodies. The bone marrow harvested by each of the two 30 methods is placed in an appropriate shipping container to support transport of the recovered BM to the manufacturing site.

The isolated CD34⁺ HSPCs are combined with the leukapherised blood product and separated as required to fulfill dose requirements for each recipient of the deceased donor kidneys. Each dose is formulated and cryopreserved as separate final drug product lots.

Whole blood collected from the deceased donor is utilized to supply the needed dose of CD3⁺ T cells for each recipient of the deceased donor kidneys. CD34⁺ HPSCs are isolated from harvested bone marrow to supply the needed dose of CD34⁺ cells for each recipient of the donor kidney.

Whole blood is collected from the exsanguinated deceased donor. Exsanguination occurs following systemic heparinization and in conjunction with cold flush of preservative solution. The exsanguinated whole blood is egressed from the deceased donor via two cannulae from both arterial and venous supply and connected through sterile connections to a collection device for transport to the manufacturing site.

Deceased donor bone marrow is collected by one or both of two independent methods: standard bone marrow aspiration and marrow recovery via an orthopedic device, the trephine. Bone marrow aspiration is performed through repeated aspirations using a heparinized syringe and Jamshedie needle. The trephine device permits taking cores of bone marrow containing cancellous bone shards.

Exsanguinated blood is required to arrive at the manufacturing site with minimal hemagglutination, red cell lysis, and cell clumping, and with high leukocyte viability to facilitate downstream processing of red blood cell and platelet depletion. Whole blood is collected in devices that allow blood to be maintained at a controlled temperature and shipped in a stable sterile transport container. The collection devices are also compatible with closed system bioprocessing.

Bone marrow collected by either standard aspiration practices or trephination is required to arrive at the manufacturing site with minimal hemagglutination, red cell lysis, and cell clumping, and with high leukocyte viability to facilitate downstream processing of red blood cell, platelet depletion and CD34⁺ HSPC isolation. Total CD34⁺ yields and recoveries are tracked as an extrapolated metric. Total volume of harvested bone marrow is calculated for each method and extrapolated for each deceased donor.

Harvested bone marrow from the deceased donor undergoes downstream processing prior to being used as a starting material for CD34⁺ HPSC isolation. Several methods and/or

technologies may be employed to establish optimal methods to separate the hematopoietic marrow from cancellous bone shards and red blood cell depletion in order to obtain bone marrow mononuclear cells (BMMNCs).

The deceased donor blood is removed prior to harvesting the bone marrow.

5 Incoming bone marrow collected by needle aspiration is diluted with buffer and filtered with an appropriate filtration device to remove bone shards. One or more of the following methods may be used to obtain maximal recovery of mononuclear cells (MNCs): size-based centrifugal separation, hetastarch sedimentation, immunomagnetic depletion or lysis followed by a spinning membrane filtration wash step or buoyancy-activated cell separation.

10 Incoming bone marrow cores obtained using the trephine are processed by one more of physical agitation, enzymatic disaggregation, or washing and filtration to recover the hematopoietic marrow compartment embedded inside the cancellous cores.

15 Incoming bone marrow is analyzed for complete blood cell counts, sterility, and leukocyte viability via flow cytometric analysis using specific antigens on the cell surface of target cell populations including but not limited to CD45, CD34 and CD3. Data from flow cytometry analysis is used to determine CD34⁺ viability, CD34⁺ HSPC frequency, and quantification of CD34⁺ cell counts as reference values from deceased donors.

The exsanguinated blood collected from the deceased donor is processed by apheresis into an appropriate starting material prior to formulation and cryopreservation.

20 The incoming whole blood receiving into the manufacturing facility processed to enrich for peripheral blood mononuclear cells to achieve an appropriate starting material for the MDR-104 manufacturing process. Incoming whole blood starting material first undergoes apheresis to reduce volume and obtain a comparable leukapheresis in-process material. Filtration of whole blood products is performed as necessary to remove clots and cell clumps from the incoming
25 material as well as the resulting leukapheresis. The resulting leukapheresis in-process material is used to formulate the CD3⁺ T cells dose for each recipient of the deceased donor's kidneys.

30 Incoming whole blood is analyzed for complete blood cell counts, sterility, and leukocyte viability via flow cytometric analysis using specific antigens on the cell surface of target cell populations including, but not limited to, CD45, CD34 and CD3. The potential to meet T cell dose expectations and T cell recoveries is calculated prior to and post apheresis.

The red blood cell-depleted, in-process material is utilized for the immunoselection of CD34⁺ HSPCs. First, platelets are depleted, and then the RBC- and platelet-reduced material undergoes CD34 immunoselection.

5 The red blood cell-depleted in-process material obtained from the bone marrow preparation sub-process is further depleted of platelets using spinning membrane filtration. Next, the RBC- and platelet-reduced sample is incubated with a CD34-capture reagent, washed using spinning membrane filtration to remove unbound beads, and loaded onto the immunomagnetic column system for immunoselection of the CD34⁺ HSPCs.

10 Incoming bone marrow is analyzed for complete blood cell counts, sterility, and leukocyte viability via flow cytometric analysis using specific antigens on the cell surface of target cell populations, including, but not limited to, CD45, CD34 and CD3. The potential to meet CD34⁺ cell dose expectations and CD34⁺ recoveries is calculated prior to and post immunoselection.

15 MDR-104 is a combined product dose formulation in which the CD34⁺ HSPCs are combined with the required T cell number to fulfill the CD3⁺ dose requirements. First, the MDR-E subprocess is divided into two separate bags in sufficient CD34⁺ cell numbers to meet the minimum dose requirements of $> 4 \times 10^6$ CD34⁺ cells/kg for each deceased donor kidney recipient. CD3⁺ T cells quantified in the whole blood preparation sub-process are then be added to each MDR-E preformulation bag to meet the required CD3⁺ T cell dose of 100×10^6 CD3 T
20 cells/kg.

MDR-T34 represents the sub-process step of combining the MDR-E derived from the CD34⁺ immunoselection with the requisite T cell dose derived from the resulting leukapheresis of the whole blood preparation unit of operation.

25 The MDR-E process intermediate is quantified for viable CD34⁺ HSPCs and then separated into pre-formulation bags based on the required patient dose of $> 4 \times 10^6$ viable CD34⁺ cells/kg. Viable CD3 T cells are quantified from the leukapheresis process intermediate, and 100×10^6 cells/kg of viable CD3 T cells are added to each of the MDR-E preformulation bags based on the T cell dose requirements for each respective patient. Each pre-formulated intermediate then undergoes an additional spinning membrane filtration step to further reduce the platelet
30 concentration.

Platelet burden and efficiency to platelet reduction are monitored pre and post spinning membrane filtration. CD34⁺ HSPC and CD3⁺ T cell loss are also monitored pre and post spinning membrane filtration. Complete cell counts and quantitative flow cytometry are used to monitor frequencies and viability of target cell populations. Functional assays are employed as necessary to ensure that sufficient T cells and CD34⁺ HSPCs with expected functional characteristics are maintained at dose requirements.

The resulting pre-formulated process intermediates are combined at a ratio of 1:2 with CryoStor 10 (CS10) to generate the drug substance and dispensed into 3-4 final drug product containers just prior to cryopreservation. The pre-formulated MDR-E and T cells are combined with cryoprotectant to generate the final product.

The preformulated MDR-T34 process intermediates for each recipient are combined with an equal volume of CS10 in the collection containers. Next, each product is divided across final product containers each at 50 mL total volume. The need for filtration is evaluated as needed prior to container closure and cryopreservation. Following the completion and container closure for both MDR-104 products, the final drug products undergo controlled cryopreservation using control rate freezers and transferred to vapor phase liquid N₂ for long-term storage at ≤ -150 °C.

Incorporation by Reference

References and citations to other documents, such as patents, patent applications, patent publications, journals, books, papers, web contents, have been made throughout this disclosure. All such documents are hereby incorporated herein by reference in their entirety for all purposes.

Equivalents

Various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full contents of this document, including references to the scientific and patent literature cited herein. The subject matter herein contains important information, exemplification, and guidance that can be adapted to the practice of this invention in its various embodiments and equivalents thereof.

Claims

What is claimed is:

1. A method for preparing a cellular product, the method comprising:
providing to a deceased subject one or more agents that promote mobilization of hematopoietic cells from bone marrow to peripheral blood;
harvesting hematopoietic cells from the subject; and
preparing a cellular product comprising the harvested hematopoietic cells.
2. The method of claim 1, wherein the one or more agents are provided to the deceased subject within 24 hours of the subject's death.
3. The method of claim 1, wherein the hematopoietic cells are harvested from peripheral blood.
4. The method of claim 1, wherein the hematopoietic cells are harvested from bone marrow.
5. The method of claim 1, wherein:
a first portion of the hematopoietic cells is harvested from peripheral blood;
a second portion of the hematopoietic cells is harvested from peripheral blood; and
the cellular product comprises hematopoietic cells from the first portion and hematopoietic cells from the second portion.
6. The method of claim 1, wherein the one or more agents are selected from the group consisting of adenosine receptor antagonist, BIO5192, a CCR1 antagonist, a CCR2 antagonist, a CXCR2 antagonist, a CXCR4 antagonist, cyclophosphamide, defibrotide, EphA3-Fc, erythropoietin (EPO), filgrastim, glycosaminoglycan (GAG) mimetic, granulocyte colony stimulating factor (G-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), growth-regulated oncogene beta (GRO-beta), human growth hormone, IL-8, lenograstim, macrophage inflammatory protein-1 alpha (MIP-1 alpha), met-SDF-1 beta, NSC23766, parathyroid hormone, pertussis toxin, plerixafor, a poly-[1-6]-D-glucopyranosyl-[1-3]-D-

glucopyranose (PGG) glucan, a Rac1 inhibitor, a retinoic acid receptor agonist, SB290157, a SDF-1 alpha peptide analog, stem cell factor (SCF), sulfated colominic acid, a sulfated polysaccharide, T134, T140, thrombopoietin (TPO), a TPO receptor agonist, a VCAM-1 inhibitor a, VLA-1 inhibitor, a VLA-4 inhibitor, and analogs and derivatives thereof.

7. The method of claim 6, wherein the one or more agents are plerixafor and G-CSF.
8. The method of claim 1, wherein the cellular product comprises CD34⁺ cells and CD3⁺ cells.
9. The method of claim 8, wherein the cellular product comprises:
at least 1×10^5 CD34⁺ cells/kg recipient weight; and
at least 1×10^5 CD3⁺ cells/kg recipient weight.
10. The method of claim 8, wherein the cellular product comprises:
at least 1×10^6 CD34⁺ cells/kg recipient weight; and
at least 1×10^6 CD3⁺ cells/kg recipient weight.
11. The method of claim 8, wherein the cellular product comprises:
at least 4×10^6 CD34⁺ cells/kg recipient weight; and
about 1×10^8 CD3⁺ cells/kg recipient weight.
12. The method of claim 1, wherein the cellular product comprises facilitating cells.
13. The method of claim 1, wherein the preparing the cellular product comprises enriching a portion of the harvested hematopoietic cells for CD34⁺ cells.
14. The method of claim 13, wherein the enriching step comprises at least one selected from the group consisting of affinity purification, buoyancy-activated cell separation, cell lysis, column filtration, hetastarch sedimentation, immunomagnetic depletion, immunomagnetic purification, size-based centrifugal separation, and spinning membrane filtration.

15. The method of claim 13, wherein the portion of the harvested hematopoietic cells is harvested from peripheral blood.
16. The method of claim 15, wherein the enriching step comprises depleting the harvested hematopoietic cells of red blood cells, platelets, or both.
17. The method of claim 15, wherein the enriching step comprises separating the peripheral blood into a cellular fraction and a plasma fraction.
18. The method of claim 13, wherein the portion of the harvested hematopoietic cells is harvested from bone marrow.
19. The method of claim 1, wherein the preparing the cellular product comprises enriching a portion of the harvested hematopoietic cells for CD3⁺ cells.
20. The method of claim 19, wherein the portion of the harvested hematopoietic cells is harvested from peripheral blood.
21. The method of claim 1, wherein preparing the cellular product comprises:
enriching a first portion of the harvested hematopoietic cells for CD34⁺ cells;
enriching a second portion of the harvested hematopoietic cells for CD3⁺ cells; and
combining the enriched first portion and the enriched second portion.
22. The method of claim 21, wherein the first portion comprises hematopoietic cells harvested from peripheral blood.
23. The method of claim 21, wherein the first portion comprises hematopoietic cells harvested from bone marrow.

24. The method of claim 21, wherein the second portion comprises hematopoietic cells harvested from peripheral blood.
25. The method of claim 1, wherein the preparing step comprises cryopreserving the cellular product.
26. The method of claim 25, wherein the cellular product is cryopreserved in a single container.
27. The method of claim 25, wherein the cellular product is cryopreserved in a multiple containers.
28. The method of claim 1, wherein the preparing step comprises assessing whether the hematopoietic cells are suitable for use in preparing the cellular product.
29. The method of claim 28, wherein the assessing step comprises analyzing in the harvested hematopoietic cells one or more of T cell activation, T cell exhaustion, T cell anergy, T cell proliferation, T cell viability, T cell apoptosis, hematopoietic stem and progenitor cell (HSPC) proliferation, HSPC viability, HSPC apoptosis, chemokine levels, and cytokine levels.
30. The method of claim 1, wherein the subject is HLA-matched to a recipient of the cellular product.
31. The method of claim 1, wherein the subject is HLA-mismatched to a recipient of the cellular product.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8

9/12




FIG. 9




FIG. 10




FIG. 11




FIG. 12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2020/035648

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(8) - A61K 35/28; C12N 5/078; C12N 5/0789 (2020.01)
 CPC - A01N 1/0221; A61K 35/28; C12N 5/0623 (2020.08)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 see Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 see Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 see Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2007/0190023 A1 (BATTISTA et al) 16 August 2007 (16.08.2007) entire document	1-31
Y	US 2017/0086453 A1 (INSTITUT PASTEUR et al) 30 March 2017 (30.03.2017) entire document	1-31
Y	WO 2016/164502 A1 (PRESIDENT AND FELLOWS OF HARVARD COLLEGE et al) 13 October 2016 (13.10.2016) entire document	7
Y	US 2017/0000825 A1 (UNIVERSITY OF LOUISVILLE RESEARCH FOUNDATION, INC.) 05 January 2017 (05.01.2017) entire document	8-12, 19-24, 30, 31
Y	WO 2018/201065 A1 (FRED HUTCHINSON CANCER RESEARCH CENTER) 01 November 2018 (01.11.2018) entire document	14, 16, 17
Y	WO 2006/047569 A2 (CELLERANT THERAPEUTICS, INC. et al) 04 May 2006 (04.05.2006) entire document	25-27
P, X	KRAFT et al. "The MarrowMiner: A Novel Minimally Invasive and Effective Device for the Harvest of Bone Marrow," Biology of Blood and Marrow Transplantation, 03 September 2019 (03.09.2019), Vol. 26, Iss. 2, Pgs. 219-229. entire document	1-31
P, X	US 2019/0275085 A1 (MEDEOR THERAPEUTICS, INC.) 12 September 2019 (12.09.2019) entire document	1-31

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"I." document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 August 2020

Date of mailing of the international search report

03 SEP 2020

Name and mailing address of the ISA/US
 Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
 P.O. Box 1450, Alexandria, VA 22313-1450
 Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2020/035648

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 2020/047236 A1 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA et al) 05 March 2020 (05.03.2020) entire document	1-31
A	US 2014/0193376 A1 (CHILDREN'S MEDICAL CENTER CORPORATION) 10 July 2014 (10.07.2014) entire document	1-31
A	US 2014/0030232 A1 (SHOEMAKER et al) 30 January 2014 (30.01.2014) entire document	1-31