



(51) International Patent Classification:

C07D 209/48 (2006.01) **C07D 235/04** (2006.01)
C07D 213/26 (2006.01) **C07D 239/34** (2006.01)
C07D 213/56 (2006.01) **C07D 239/42** (2006.01)
C07D 213/65 (2006.01) **C07D 277/30** (2006.01)
C07D 213/74 (2006.01) **C07D 307/68** (2006.01)
C07D 217/00 (2006.01) **C07D 317/68** (2006.01)
C07D 231/10 (2006.01) **C07D 319/18** (2006.01)
C07D 233/90 (2006.01)

(21) International Application Number:

PCT/US2010/032347

(22) International Filing Date:

26 April 2010 (26.04.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/172,873 27 April 2009 (27.04.2009) US
 61/319,482 31 March 2010 (31.03.2010) US

(71) **Applicant** (for all designated States except US):
BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; Binger Strasse 173, 55216 Ingelheim (DE).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **KOWALSKI, Jennifer A.** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **MARSHALL, Daniel Richard** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **PROKOPOWICZ, Anthony S. III** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **SCHLYER, Sabine** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **SIBLEY, Robert** [US/US];

Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **SORCEK, Ronald John** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **WU, Di** [CN/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **WU, Frank** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **YOUNG, Erick Richard Roush** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US).


(74) **Agents:** **MORRIS, Michael P.** et al.; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: CXCR3 RECEPTOR ANTAGONISTS



(57) **Abstract:** The present invention relates to compounds of formula (I), and pharmaceutically acceptable salts thereof, wherein R^1 to R^5 , A, B, D and X are as defined herein. The invention also relates to pharmaceutical compositions comprising these compounds, methods of using these compounds in the treatment of various diseases and disorders, processes for preparing these compounds and intermediates useful in these processes.

WO 2010/126811 A1



Published:

— *with international search report (Art. 21(3))*

CXCR3 RECEPTOR ANTAGONISTS

FIELD OF THE INVENTION

This invention relates to heterocyclic compounds that are useful as antagonists of CXCR3 and are thus useful for treating a variety of diseases and disorders that are mediated or sustained through the interaction of CXCR3 and its ligands including multiple sclerosis, psoriasis, rheumatoid arthritis, inflammatory bowel disease and atherosclerosis. This invention also relates to pharmaceutical compositions comprising these compounds, methods of using these compounds in the treatment of various diseases and disorders, processes for preparing these compounds and intermediates useful in these processes.

BACKGROUND OF THE INVENTION

Chemokine receptors, a subclass of the G-protein coupled receptors (GPCRs) are expressed on the surface of T-cells and other leukocytes. The interaction of chemokine receptors with their ligands plays an important role in the migration of leukocytes to sites of inflammation (A.D. Luster, New Engl. J. Med., 1998, 338, 436). The chemokine receptor CXCR3 is preferentially expressed on T helper (Th1) cells but is also found on natural killer cells and subsets of dendritic cells. Three major chemokine ligands for CXCR3 have been identified: Mig (Monokine Induced by γ -IFN/CXCL9), IP-10 (γ -interferon inducible protein) and I-TAC (IFN-Inducible T Cell α Chemoattractant /CXCR11) (K.E. Cole et al., J. Exp. Med., 1998, 187, 2009; Y. Weng et al., J. Biol.Chem., 1998, 273, 18288).

Histological evaluations of numerous inflammatory lesions, including those from patients with multiple sclerosis (T.L. Sorenson et al., J. Clin. Invest., 1999, 103, 807), rheumatoid arthritis (S. Qin et al., J. Clin. Invest., 1998, 101, 746), psoriasis (J. Flier et al., J. Pathol., 2001, 194, 398) and inflammatory bowel disease (Y.H. Yuan et al., Inflamm. Bowel Dis.,

2001, 7, 281) have shown elevated expression of CXCR3 ligands accompanied by an increased frequency of T cells bearing CXCR3. This is in marked contrast to what is found in most normal tissues, where expression of CXCR3 and its ligands is extremely low. This correlative evidence suggests a role of CXCR3 in Th1-mediated chronic inflammation.

Studies with CXCR3 and IP-10 deficient mice also suggest a role for CXCR3 and IP-10 in Th1 mediated disease. For example, in one study CXCR3 ^{-/-} mice showed significant resistance to allograft rejection (W.W. Hancock et al., J. Exp. Med., 2000, 192, 1515). In another study, IP-10 deficient mice showed protection against the development of colitis (U.P. Singh et al., J. Immunol., 2003, 171, 1401). Further evidence of a role for CXCR3 and IP-10 as mediators of disease is provided by studies utilizing blocking antibodies. For example in a rat model of adjuvant induced arthritis (I. Salomon et al., J. Immunol., 2002, 169, 2685) a DNA vaccine approach to overexpress self IP-10 was used to induce the production of self-IP-10 antibodies. These Abs are specific for IP-10 and do not cross react with other proinflammatory cytokines or chemokines including Mig and I-TAC. Pretreatment with this vaccine protected rats from the development of severe arthritis and reduced the time to remission of symptoms. In addition, affinity purified anti-IP-10 from vaccinated rats could therapeutically transfer protection to newly diseased rats. In another study, this vaccine approach was successful in suppressing disease in a mouse model of multiple sclerosis (G. Wildbaum et al., J. Immunol., 2002, 168, 5885).

In a study of pulmonary inflammation, (N. Li et al., Acta Pharmacol. Sinica, 2008, 29, 14) CXCR3 knockout mice showed alleviated inflammation compared to wild type mice in cigarette smoke induced pulmonary injury as well as lower influx of inflammatory T cells. Similarly, in a model of nephrotoxic nephritis, CXCR3 knockout mice showed reduced influx of T cells, less severe nephritis and improved renal function compared to wild type mice (U. Panzer et al., J. Am. Soc. Nephrol., 2007, 18, 2071). Thus CXCR3

may play a role in inflammatory pulmonary diseases such as COPD and inflammatory kidney disease.

Studies such as those cited above suggest that inhibitors of CXCR3 may be useful in treating inflammatory and autoimmune diseases in which CXCR3-mediated cellular recruitment plays a role, including multiple sclerosis, psoriasis, rheumatoid arthritis, inflammatory bowel disease, COPD and kidney disease.

Recent work has also implicated CXCR3 in the pathogenesis of atherosclerosis. In one study (F. Mach et al., *J. Clin. Invest.*, 1999, 104, 1041) CXCR3 was found expressed in all T lymphocytes within human atherosclerotic lesions. The ligands IP10, Mig and I-TAC were all found within lesion-associated cells including endothelial and smooth muscle cells (Mig and I-TAC) and macrophages (IP10), suggesting these ligands play a role in recruitment of activated T lymphocytes within vascular wall lesions in atherogenesis. Left untreated and allowed to progress, atherosclerosis can result in narrowing of the lumen of the artery and plaque rupture which can lead to coronary heart disease, myocardial infarction and stroke (J. Sanz and Z. A. Fayad, *Nature*, 2008, 451, 953).

Further evidence has come from genetic deletion studies in mice. CXCR3 deletion on an ApoE^{-/-} background resulted in a significant reduction in atherosclerotic lesion formation following ten weeks on a high cholesterol diet (N.R. Veillard et al., *Circulation*, 2005, 112, 870). Moreover, deletion of the CXCR3 ligand, IP-10 on an ApoE^{-/-} background similarly reduced atherosclerotic lesion load (E. Heller et al., *Circulation*, 2006, 113, 2301). More recently, NBI-74330 a CXCR3 antagonist was dosed prophylactically in a LDL receptor knockout model. Similar to the CXCR3 deletion studies in the ApoE^{-/-} results, NBI-74330 significantly attenuated atherosclerotic lesion formation (E.J.A. van Wanrooij et al., *Arterioscler. Thromb. Vasc. Biol.*, 2008, 28,,251-257).


As a result of studies such as those cited above implicating the interaction of CXCR3 and its ligands in the etiology of various inflammatory and autoimmune diseases as well as atherosclerosis, considerable effort has been directed towards discovering antagonists of this interaction. A number of inhibitors have been reported in the scientific literature, including small molecule antagonists, antibodies and modified ligands (see for example J.C. Medina et al., Ann. Rep. Med. Chem., 2005, 40, 215). However, to date, no CXCR3 antagonist has been approved as a marketed drug.

BRIEF SUMMARY OF THE INVENTION

The present invention provides novel compounds which block the interaction of CXCR3 and its ligands and are thus useful for treating diseases and disorders that are mediated or sustained through the activity of CXCR3 including multiple sclerosis, psoriasis, rheumatoid arthritis, inflammatory bowel disease, COPD, kidney disease and atherosclerosis, myocardial infarction and stroke. This invention also relates to pharmaceutical compositions comprising these compounds, methods of using these compounds in the treatment of various diseases and disorders, processes for preparing these compounds and intermediates useful in these processes.

DETAILED DESCRIPTION OF THE INVENTION

In its broadest embodiment, the present invention relates compounds of formula (I):



wherein:

A is C or N;

B is C or N;

D is C or N;

X is -NH-, -NHC(O)-, -N(CH₃)C(O)- or absent;

R¹ is H, -CN, halogen, -CF₃, -OCF₃, C₁₋₃alkyl, C₁₋₃alkoxy, -S(O)_nCH₃, amino, mono- or dimethylamino, -NHC(O)C₁₋₃alkyl, -NO₂, -C(O)NH₂, -C(O)NHC₁₋₃alkyl or -C(O)C₁₋₃alkyl;

R² is aryl or heteroaryl, each optionally substituted with one to three R⁶;

R³ is heteroaryl, heterocyclyl, aryl or C₃₋₁₀cycloalkyl each optionally substituted with one to three R⁷;

R⁴ and R⁵ are each independently selected from H and C₁₋₂alkyl; or R⁴ and R⁵ may join to form an ethyl bridge;

each R⁶ is independently -OH, oxo, hydroxyC₁₋₆alkyl, halogen, -(CH₂)_m-CN, nitro, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₁₀cycloalkyl, C₁₋₆alkoxy, phenoxy, heteroaryloxy, C₁₋₆alkoxycarbonyl, carboxyl, -C(O)C₁₋₆alkyl, -(CH₂)_m-NR₈R₉, -S(O)_nC₁₋₆alkyl, -NHS(O)_nC₁₋₆alkyl, -NR⁸C(O)C₁₋₆alkyl, -S(O)₂NR₈R₉, -C(O)NR₈R₉, heteroaryl, heterocyclyl, phenyl or benzyl, wherein each alkyl, alkenyl, alkynyl or alkoxy of said R⁶ is optionally partially or fully halogenated and each heterocyclyl, heteroaryl, phenyl or benzyl of said R⁶ is optionally substituted with one to three C₁₋₆alkyl, C₁₋₆alkoxy(CH₂)_m, halogen, -CN, -CF₃, C₁₋₆acyl, -NR⁸R⁹, C(O)NR⁸R⁹, -OH, hydroxyC₁₋₆alkyl, pyrrolidinyl or -S(O)_nC₁₋₆alkyl;

each R^7 is independently C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, C_{1-6} alkoxy, halogen, oxo, -CN, carboxy, $-(CH_2)_m-NR_8R_9$, phenyl or heteroaryl, wherein each alkyl, alkenyl, alkynyl or alkoxy of said R^7 is optionally partially or fully halogenated and wherein two R^7 on adjacent carbon atoms of R^3 may join to form a three to seven member ring fused to R^3 ;

R^8 and R^9 are each independently hydrogen, C_{1-6} alkyl, C_{1-6} acyl, C_{3-10} cycloalkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyl C_{1-6} alkoxy, C_{1-6} alkylsulfonyl or C_{1-6} alkoxycarbonyl;

m is 0-3;

n is 0-2; and

----- is a single bond if A is N and may be a single or double bond if A is C;

or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described above and wherein R^3 is aryl or heteroaryl, or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described in the broadest embodiment above wherein R^2 is aryl, and R^3 is aryl or heteroaryl, or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described in the broadest embodiment above wherein R^2 is heteroaryl, and R^3 is aryl or heteroaryl, or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described in the broadest embodiment above and wherein:

A is N; and

D is C;

or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described above and wherein:

B is C;

X is absent;

or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described in the broadest embodiment above and wherein:

B is C;

D is C;

R¹ is H, -CN, -F, -Cl, -CF₃, -CH₃, -OCH₃, -C(O)NHCH₃ or -S(O)₂CH₃;

R² is benzoimidazolyl, benzooxazolyl, benzo[b]thiophenyl, dibenzofuranyl, dibenzothiophenyl, furanyl, imidazolyl, 1H-imidazo[4,5-c]pyridinyl, indolyl, isoindolyl, isoquinolinyl, isoxazolyl, oxazolyl, phenyl, pyranal, pyrazinyl, pyrazolyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, quinolinyl, thiadiazolyl, thiazolyl, thienyl or triazinyl, wherein each of the foregoing is optionally substituted by one to three R⁶;

R³ is benzoimidazolyl, benzo[d]isothiazolyl, benzo[b]thiophenyl, benzotriazolyl, furanyl, imidazo[4,5-b]pyridinyl, imidazolyl, indazolyl, indolyl, isoindolyl, isoquinolinyl, isothiazolyl, phenyl, phthalazinyl, pyrazolyl, pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-c]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl, pyrrolo[2,3-d]pyrimidinyl, pyrrolyl, quinazolinyl, thiadiazolyl or thiazolyl wherein each of the foregoing is optionally substituted with one to three R⁷;

R⁴ and R⁵ are H;

each R⁶ is independently -OH, oxo, hydroxyC₁₋₆alkyl, halogen, -(CH₂)_m-CN, nitro, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₁₀cycloalkyl, C₁₋₆alkoxy, phenoxy, pyridyloxy, C₁₋₆alkoxycarbonyl, carboxyl, -C(O)C₁₋₆alkyl, -(CH₂)_m-NR₈R₉, -S(O)_nC₁₋₆alkyl, -NHS(O)₂C₁₋₆alkyl, -NHC(O)C₁₋₆alkyl, -S(O)₂NR₈R₉, -C(O)NR₈R₉, thienyl, morpholinyl, pyrrolidinyl, piperidinyl, [1,3]-oxazepan-1-yl, piperazinyl, azabicyclo[2.2.1]heptyl, 2-oxa-5-azabicyclo[2.2.1]heptyl, octahydroisoquinolinyl, [1,4]-oxazepanyl, azetidiny phenyl or benzyl, wherein each alkyl, alkenyl or alkynyl of said R⁶ is optionally partially or fully halogenated and each heterocyclyl or phenyl of said R⁶ is optionally substituted with one to three CH₃, -OCH₃, halogen, -CN, -CF₃, -C(O)CH₃, NR⁸R⁹, -C(O)NH₂, -C(O)NHCH₃, -OH, -CH₂OCH₃, -C(OH)(CH₃)CH₃, -SCH₃, pyrrolinyl or -S(O)₂CH₃;

each R⁷ is independently C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₁₀cycloalkyl, C₁₋₆alkoxy, halogen, oxo, -CN, carboxy, -(CH₂)_m-NR₈R₉, phenyl or pyridyl, wherein each alkyl, alkenyl or alkynyl of said R⁷ is optionally partially or fully halogenated;

R⁸ and R⁹ are each independently hydrogen, C₁₋₆alkyl, C₁₋₆acyl or C₃₋₁₀cycloalkyl;

or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described in the embodiment immediately above and wherein:

A is N;

X is -NH- or absent;

R¹ is H, -CN, -F, -Cl, -CF₃, -CH₃ or -OCH₃;

R² is benzoimidazolyl, benzooxazolyl, imidazolyl, 1H-imidazo[4,5-c]pyridinyl, indolyl, isoindolyl, isoquinolyl, isoxazolyl, oxazolyl, phenyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridinyl, pyrimidinyl, quinolyl, thiadiazolyl, thiazolyl or thienyl, wherein each of the foregoing is optionally substituted by one to three R⁶;

R³ is benzoimidazolyl, benzotriazolyl, imidazolyl, imidazo[4,5-b]pyridinyl, indazolyl, indolyl, isoindolyl, isothiazolyl, pyrazolyl, pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl, pyrrolo[2,3-d]pyrimidinyl, pyrrolyl, thiadiazolyl or thiazolyl wherein each of the foregoing is optionally substituted with one to three R⁷;

each R⁶ is independently -CH₂OH, -Cl, -F, -CN, oxo, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₂ alkoxy, phenoxy, pyridyloxy, -C(O)CH₃, -(CH₂)_m-NR₈R₉, -S(O)_nCH₃, -NHS(O)₂CH₃, -NH(CO)CH₃, -S(O)₂NR₈R₉, -C(O)NR₈R₉, thienyl, morpholyl, pyrrolidinyl, piperidinyl, [1,3]-oxazepan-1-yl, piperazinyl, phenyl or benzyl, wherein each alkyl of said R⁶ is optionally partially or fully halogenated and each heterocyclyl or phenyl of said R⁶ is optionally substituted with one to three CH₃, -OCH₃, halogen, -CN, -CF₃, N(CH₃)₂, -C(O)NH₂, -C(O)NHCH₃, -OH, -CH₂OCH₃, -C(OH)(CH₃)CH₃, pyrrolidinyl or S(O)₂CH₃;

each R⁷ is independently C₁₋₃ alkyl, -OCH₃, CF₃, oxo, -CN, -Cl or -F;

R⁸ and R⁹ are each independently hydrogen, C₁₋₆ alkyl, C₁₋₃ acyl or C₃₋₆ cycloalkyl;

or the pharmaceutically acceptable salts thereof.

In another embodiment there is provided a compound of formula (I) as described in the embodiment immediately above and wherein:



X is absent;







R³ is benzoimidazolyl, benzotriazolyl, imidazolyl, imidazo[4,5-b]pyridinyl, indazolyl, indolyl, isoindolyl, pyrazolyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl or pyrrolyl, wherein each of the foregoing is optionally substituted with one to three R⁷;







or the pharmaceutically acceptable salts thereof.






The following are representative compounds of the invention which can be made by the methods described in the general synthetic schemes, the synthetic examples, and known methods in the art.






Table I






MOLECULAR STRUCTURE	CHEMICAL NAME	Observed MH ⁺
	3-(2-Methylsulfanyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	471
	1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	446






	4-Amino-N-[2-(4-biphenyl-2-yl)-piperazin-1-yl]-2-oxo-ethyl]-3-methoxy-benzamide	446
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-naphthalen-1-yl-phenyl)-piperazin-1-yl]-ethanone	426
	1-Methyl-1H-imidazole-2-carboxylic acid [2-(4-biphenyl-2-yl)-piperazin-1-yl]-2-oxo-ethyl]-amide	404
	5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl)-acetyl]-piperazin-1-yl}-phenyl}-pyridine-2-carbonitrile	423
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	492
	1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl}-ethanone	497

	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(3,4-dimethoxybenzylamino)-ethanone	447
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-piperidin-1-yl-ethanone	365
	Isoquinoline-6-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	451
	Thiazole-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	408
	1-(4-{2-[2-(2,6-Dimethylmorpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	491
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	429






	1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	443
	1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	485
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	447
	4,5-Dimethyl-furan-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	418
	1-(2-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-imidazolidin-2-one	452






	1-[4-(4'-Dimethylaminomethyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	433
	Isoquinoline-7-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	451
	5-Methoxy-pyridine-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	432
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	475
	1-{3-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-3,8-diaza-bicyclo[3.2.1]oct-8-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	455






	1-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-1H-benzoimidazole-5-carboxylic acid	442
	1-[3-(2-Morpholin-4-yl-pyrimidin-5-yl)-3',6'-dihydro-2'H-[2,4']bipyridinyl-1'-yl]-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	483
	2,3-Dihydro-benzo[1,4]dioxine-6-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	459
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-piperidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	461
	3-(2-Methoxy-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	455

	1-(4-{2-[2-((2R,6R)-2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	513
	1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	429
	1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	421
	3-(2-Methylamino-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	454
	3-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	510






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	531
	N-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-3-pyridin-3-yl-benzamide	477
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-3-(3,5-dimethyl-pyrazol-1-yl)-propan-1-one	390
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(5-methanesulfonyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	454
	1-Biphenyl-2-yl-4-[2-(3,5-dimethyl-pyrazol-1-yl)-ethyl]-piperazine	362
	3-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-3H-benzimidazole-5-carboxylic acid	442






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	475
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	497
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	463
	1-{4-[4-Methoxy-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	515
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-c]pyridin-1-yl-ethanone	429






	1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	514
	1-{4-[2-(2-[1,4]Oxazepan-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	499
	1-{4-[2-(2-Ethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	443
	1-{4-[2-(2-Pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	469
	1-(2-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1,3-dihydro-benzoimidazol-2-one	501






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	424
	1-(4-{2-[2-((2R,6S)-2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	513
	1-(4-{2-[2-(4-Methyl-piperazin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	498
	1-{4-[2-(2-Piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	483
	1-{4-[2-(2-Morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperidin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	484






	1-{4-[2-(2-Methanesulfonyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	546
	1-{4-[4-Chloro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	519
	5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyrimidine-2-carbonitrile	424
	2-Imidazo[4,5-b]pyridin-3-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	430
	1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	553






	5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	401
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-ethoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	422
	4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-morpholin-4-yl-pyrimidin-5-yl)-benzonitrile	488
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1-methyl-1H-indol-6-yl)-phenyl]-piperazin-1-yl}-ethanone	429
	1-{4-[2-(2-Morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-3,6-dihydro-2H-pyridin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	482






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-3,6-dihydro-2H-pyridin-1-yl}-ethanone	460
	1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-5-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	553
	5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyridine-2-carboxylic acid amide	441
	1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-4-ylamino)-ethanone	461
	1-{4-[2-Fluoro-6-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	503






	2-Indazol-2-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	429
	2-(2,4-Dimethyl-imidazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	463
	1-{4-[2-(6-Chloro-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	433
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-trifluoromethyl-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone	444
	1-{4-[2-(6-Methoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	429





	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	407
	2-Indazol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	429
	1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(thiazol-2-ylamino)-ethanone	467
	4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methylsulfanyl-pyrimidin-5-yl)-benzonitrile	449
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	443






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-phenoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	470
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methyl-piperazin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	476
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperidin-1-yl}-ethanone	462
	5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidine-2-carbonitrile	402
	2-Indol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	428






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	407
	5-(2-{4-[2-(2,4-Dimethyl-imidazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	402
	5-(2-{4-[2-(Thiazol-2-ylamino)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	405
	1-[4-(2-Benzo[b]thiophen-2-yl)-phenyl]-piperazin-1-yl-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	432
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methanesulfonyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	524






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-methoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	406
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(pyridin-3-yloxy)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	471
	1-[4-(2-Benzothiazol-5-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	433
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-methoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone	406
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(thiazol-2-ylamino)-ethanone	411






	2-(2,4-Dimethyl-imidazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	407
	1-[4-(4'-Benzoyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	480
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	391
	3,4-Dimethoxy-N-(2-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-benzamide	548
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	390






	5-(2-{4-[2-(5-Methyl-3-trifluoromethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	455
	1-{4-[2-(9H-Carbazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	465
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[3'-(5-methyl-[1,3,4]oxadiazol-2-yl)-biphenyl-2-yl]-piperazin-1-yl}-ethanone	458
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-naphthalen-2-yl-phenyl)-piperazin-1-yl]-ethanone	426






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-indol-6-yl)-phenyl]-piperazin-1-yl}-ethanone	415
	1-[4-(4'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	410
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-trifluoromethoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	459
	5-(2-{4-[2-(Pyridin-4-ylamino)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	399
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	406






	2'-[4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl]-biphenyl-4-carbonitrile	400
	1-[4-(4'-tert-Butyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	432
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methyl-3-trifluoromethyl-pyrazol-1-yl)-ethanone	461
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-isopropyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	418
	1-[4-(3'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	410






	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,4,5-trimethyl-pyrazol-1-yl)-ethanone	421
	1-{4-[2-(2-Methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	478
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-[1,1';4',1'']terphenyl-2-yl)-piperazin-1-yl-ethanone	452
	1-[2-(4-Biphenyl-2-yl)-piperazin-1-yl]-2-oxo-ethyl-1,3-dihydro-benzoimidazol-2-one	413
	1-{4-[2-(2-Cyclopropylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	433






	4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methoxy-pyrimidin-5-yl)-benzonitrile	432
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	397
	1-{4-[2-(2-tert-Butyl-thiazol-4-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	439
	3-(2-Methanesulfonyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	503
	2-(3-tert-Butyl-[1,2,4]thiadiazol-5-ylamino)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	468







	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-fluoro-biphenyl-2-yl)-piperazin-1-yl]-ethanone	393
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	443
	1-{4-[4-Methanesulfonyl-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	562
	1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-3-ylamino)-ethanone	461
	2-Benzotriazol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	430







	N-(2-{4-[2-(6-Cyano-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3,4-dimethoxy-benzamide	487
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-4-ylamino)-ethanone	405
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(2,5-dimethyl-imidazol-1-yl)-ethanone	375
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-indol-5-yl)-phenyl]-piperazin-1-yl}-ethanone	415
	N-(2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-yl)-methanesulfonamide	469








	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(pyridin-4-ylamino)-ethanone	373
	5-(2-{4-[2-(5-Methyl-3-trifluoromethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid amide	473
	1-{4-[3-(2-Morpholin-4-yl-pyrimidin-5-yl)-pyridin-2-yl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	486
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-pyrimidin-5-yl-phenyl)-piperazin-1-yl]-ethanone	377
	1-[4-(3',5'-Bis-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	511







	1-{4-[2-(2-Amino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	392
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methanesulfonyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	454
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-sulfonic acid amide	455
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-imidazo[4,5-b]pyridin-1-yl-ethanone	398
	1-[4-(2'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	410

	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-hydroxymethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	406
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(thiazol-2-ylamino)-ethanone	379
	3,4-Dimethoxy-N-(2-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-benzamide	493
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-hydroxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	391
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-[1,1';3',1'']terphenyl-2-yl)-piperazin-1-yl)-ethanone	452







	N-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-3,4-dimethoxy-benzamide	461
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-carboxylic acid amide	419
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(pyridin-3-ylamino)-ethanone	373
	2-((S)-Indan-1-ylamino)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	444
	2-Methyl-2H-pyrazole-3-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	405
	N-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-4-fluoro-3-methoxy-benzamide	449






	N-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-benzamide	401
	2-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-isoindole-1,3-dione	426
	N-(2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-yl)-acetamide	433
	2-Benzoimidazol-1-yl-1-(4-biphenyl-2-yl-piperazin-1-yl)-ethanone	397
	2-(2,4-Dimethyl-pyrrol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	406
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-pyridin-3-yl-phenyl)-piperazin-1-yl]-ethanone	376

	1-Methyl-1H-pyrrole-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	403
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(4-methyl-pyrazol-1-yl)-ethanone	393
	1-(4-Biphenyl-2-yl)-3,6-dihydro-2H-pyridin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	372
	N-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-2,3-dimethoxy-benzamide	461
	1H-Pyrazole-3-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	390
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(3-methyl-pyrazol-1-yl)-ethanone	361
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-pyridin-4-yl-phenyl)-piperazin-1-yl]-ethanone	376






	1-(4-Biphenyl-2-yl-piperidin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	374
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(5-methyl-3-trifluoromethyl-pyrazol-1-yl)-ethanone	429
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1-methyl-1H-pyrazol-4-yl)-phenyl]-piperazin-1-yl}-ethanone	379
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	456
	2,5-Dimethyl-2H-pyrazole-3-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	419
	2-Ethyl-5-methyl-2H-pyrazole-3-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	432





	2-(4,5-Dimethyl-thiazol-2-ylamino)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	439
	2,4-Dimethyl-thiazole-5-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	435
	1,5-Dimethyl-1H-pyrazole-3-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	419
	Benzo[1,3]dioxole-5-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	445
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(4-methyl-thiazol-2-ylamino)-ethanone	425
	1-{4-[4-Methanesulfonyl-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	507





	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2'-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	443
	1-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-2-one	567
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-4-yl)-ethanone	400
	Adamantane-1-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide	459
	N-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-4-methoxy-3-methyl-benzamide	445
	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(4-phenyl-imidazol-1-yl)-ethanone	423






	1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-imidazo[4,5-b]pyridin-3-yl-ethanone	398
	2-(3-tert-Butyl-isothiazol-5-ylamino)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	467
	1-[4-(3',4'-Dimethoxy-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	435
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-quinolin-3-yl-phenyl)-piperazin-1-yl]-ethanone	426
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-ethoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone	420






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(5-methoxy-pyrazin-2-yl)-phenyl]-piperazin-1-yl}-ethanone	407
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-morpholin-4-yl-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone	461
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-ethoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	419
	1-[4-(2-Benzo[1,3]dioxol-5-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	419
	1-[4-(3'-Acetyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	417






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1-methyl-1H-indol-5-yl)-phenyl]-piperazin-1-yl}-ethanone	428
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-quinolin-6-yl-phenyl)-piperazin-1-yl]-ethanone	426
	1-[4-(4'-Acetyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	417
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-ethoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	419
	4-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester	434





	5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-nicotinic acid ethyl ester	448
	1-{4-[2-(1-Benzyl-1H-pyrazol-4-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	455
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-thiophen-2-yl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	457
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-sulfonic acid dimethylamide	482





	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(5-phenyl-thiophen-2-yl)-phenyl]-piperazin-1-yl}-ethanone	457
	1-[4-(2-Dibenzothiophen-2-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	481
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-3-carboxylic acid methylamide	432
	1-[4-(4'-Benzyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	465






	N-(2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-3-yl)-acetamide	432
	5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester	434
	1-[4-(3'-Benzyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	465
	1-[4-(2-Benzo[b]thiophen-3-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	431
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-carboxylic acid dimethylamide	446





	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-3-sulfonic acid dimethylamide	482
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-carboxylic acid methylamide	432
	1-[4-(2-Dibenzofuran-4-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	465
	2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-3-carboxylic acid dimethylamide	446
	1-{4-[2-(1H-Benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	415





	1-[4-(2-Benzooxazol-2-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	416
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-imidazo[4,5-c]pyridin-2-yl)-phenyl]-piperazin-1-yl}-ethanone	416
	1-{4-[2-(5-Chloro-1H-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	449
	2-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-1H-benzoimidazole-5-carbonitrile	440
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-ethanone	483





 Chiral	1-(4-{2-[2-((R)-3-Dimethylamino-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	489
	1-(4-{2-[(1R,4S)-2-(2-Aza-bicyclo[2.2.1]hept-2-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	472
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[4-fluoro-2-(2-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl}-ethanone	481
	1-{4-[4-Chloro-2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl}-ethanone	462





	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methanesulfonyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	538
	1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-4-carboxylic acid amide	503
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxy-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	476
	1-{4-[4-Chloro-2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	459





	1-{4-[4-Chloro-2-(2-methoxypyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	463
	1-(4-{2-[2-(3,3-Difluoro-azetidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethylpyrazol-1-yl)-ethanone	468
	2-(3,5-Dimethylpyrazol-1-yl)-1-(4-{2-[2-(3-fluoropiperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	478
	2-(5-Fluoropyrrolo[2,3-b]pyridin-1-yl)-1-{4-[2-(2-methoxypyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	447
	1-{4-[4-Chloro-2-(2-methylsulfanylpirimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	479






	1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-3-carboxylic acid amide	504
	1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-4-carboxylic acid methylamide	517
	1-(4-{2-[2-(3,3-Difluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	496
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-hydroxy-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	462





	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-fluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	479
	2-(4-Chloro-3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	442
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-ethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	420
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-[1,4]oxazepan-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	476






	1-{4-[4-Chloro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	496
	1-{3-Methyl-4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	497
	1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3-methyl-1,3-dihydro-benzoimidazol-2-one	460
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-trifluoromethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	528






	1-(4-{2-[2-(4,4-Dimethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	488
 Chiral	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	490
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-{2-[4-(1-hydroxy-1-methyl-ethyl)-piperidin-1-yl]-pyrimidin-5-yl}-phenyl)-piperazin-1-yl]-ethanone	518
	1-{4-[4-Chloro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	441







	1-{2-Methyl-4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	497
	1-{4-[4-Chloro-2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	440
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[(1S,4S)-2-(2-oxa-5-azabicyclo[2.2.1]hept-5-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	474
	1-(4-{2-[2-(3,5-Dimethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	488






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[4-fluoro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	425
	2-(7-Chloro-pyrrolo[2,3-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	464
	1-{4-[2-(2-Morpholin-4-yl-pyridin-4-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	483
	2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-2,3-dihydro-isoindol-1-one	445
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(octahydro-isoquinolin-2-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	514






	1-(4-{2-[2-(4,4-Difluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	496
	1-{4-[4-Chloro-2-(2-methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	511
	1-{4-[2-(2-Methoxy-pyridin-4-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	428
	1-(4-{2-[2-(4-Dimethylamino-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	503






	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methoxy-4-methyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	504
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-pyrrolidin-1-yl)-piperidin-1-yl]-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	529
	2-(4-Chloro-pyrrolo[2,3-d]pyrimidin-7-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	465
	2-(4-Chloro-pyrrolo[3,2-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	464
	4-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methylamide	434






	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[3,2-c]pyridin-1-yl-ethanone	429
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methyl-3-nitro-pyrazol-1-yl)-ethanone	438
	2-(5-Chloro-pyrrolo[3,2-b]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	464
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methoxy-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	490
	4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methylamino-pyrimidin-5-yl)-benzonitrile	431






	2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-2H-phthalazin-1-one	458
	1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1H-indole-2,3-dione	458
	3-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3H-quinazolin-4-one	458
	2-(6-Bromo-pyrrolo[3,2-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	508
Chiral 	1-((S)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	390
	N-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-N-methyl-acetamide	448

	1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1H-pyrrolo[2,3-b]pyridine-4-carbonitrile	455
	(R)-1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-pyrrolidine-2-carboxylic acid amide	489
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methoxy-pyrrolo[2,3-b]pyridin-1-yl)-ethanone	460
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3-methyl-benzo[b]thiophen-2-yl)-ethanone	460
	2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-2H-isoquinolin-1-one	464






	2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1,1-dioxo-1,2-dihydro-116-benzo[d]isothiazol-3-one	495
	2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(pyrrolidine-1-carbonyl)-pyridin-4-yl]-phenyl}-piperazin-1-yl)-ethanone	491
	2-(4,5-Dichloro-imidazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	448
	1-{4-[3-(2-Morpholin-4-yl)-pyrimidin-5-yl]-pyridin-4-yl}-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	486
 Chiral	1-((R)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	390





Chiral 	1-((S)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	412
Chiral 	(S)-1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-pyrrolidine-2-carboxylic acid methylamide	503
	2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-isoindole-1,3-dione	466
Chiral 	1-((R)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	412
	5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methylamide	434






	2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-3,4-dihydro-2H-isoquinolin-1-one	466
	1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methoxy-pyrrolo[3,2-b]pyridin-1-yl)-ethanone	460
	2-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-2,3-dihydro-isoindol-1-one	430
	4-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid dimethylamide	448
	5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-N-methyl-nicotinamide	434






 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide	503
	1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one	594
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone	620
	1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	580
 Chiral	5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-yl)acetyl]-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide	504

	1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	642
	2-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-2,3-dihydro-1H-isindol-1-one	657
	5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isindol-2-yl)acetyl]piperazin-1-yl}benzamide	518
	2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone	643

	1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	574
	1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one	658
 Chiral	2-[(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl]-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide	481
	1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	516
	1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	517

	1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	578
	1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	532
	1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	592
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)ethanone	558

	1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	592
 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl]benzamide	484
	1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	528
	1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	512
 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl]benzamide	519

 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide	503
 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl}benzamide	487
	1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	578
	1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	512
	2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one	593






	1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	574
	1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one	608
	5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide	503
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	494
	2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isindol-1-one	595






	2-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	531
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone	552
	1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	596
	5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide	519
	1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	516






	2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	527
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]ethanone	570
	2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one	607
	5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide	484
	1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	580






	1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	579
	1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	447
	1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one	561
	5-fluoro-2-[(3S)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide	504
	1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)piperidin-2-one	623






	1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	581
	1-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	518
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone	556
	1-{2-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one	590
Chiral	N-{2-[(2S)-4-{4-fluoro-2-[(2-methoxypyridin-4-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	525






 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-((3S)-3-methyl-4-(N-methyl-N-(2-methylpropanoyl)glycyl)piperazin-1-yl)benzamide	486
 Chiral	1-(4-(2-(2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl)phenyl)-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	513
 Chiral	5-fluoro-2-((3S)-3-methyl-4-((2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl)piperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	564
 Chiral	2-((3S)-4-((3,5-dimethyl-1H-pyrazol-1-yl)acetyl)-3-methylpiperazin-1-yl)-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide	481
 Chiral	N-(2-((2R)-4-(4-fluoro-2-((2-methoxypyridin-4-yl)carbamoyl)phenyl)-2-methylpiperazin-1-yl)-2-oxoethyl)-N,5-dimethyl-1,3-oxazole-4-carboxamide	525






 Chiral	2-((3S)-4-((3,5-dimethyl-1H-pyrazol-1-yl)acetyl)-3-methylpiperazin-1-yl)-5-fluoro-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	526
	1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	593
	1-(2-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one	463
 Chiral	5-fluoro-N-(2-methoxypyridin-4-yl)-2-((3S)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl)benzamide	518
	1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	448






	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)ethanone	490
	1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	502
	1-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	531
	1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	487
	1-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	502






 Chiral	5-fluoro-2-((3S)-3-methyl-4-(N-methyl-N-(2-methylpropanoyl)glycyl)piperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	531
	2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone	575
	1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	486
	1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}piperidin-2-one	573
	1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	502






	5-fluoro-2-((3S)-3-methyl-4-((1-oxo-1,3-dihydro-2H-isindol-2-yl)acetyl)piperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	563
	5-fluoro-2-((3S)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	549
	1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}piperidin-2-one	559
	2-((3R)-4-((3,5-dimethyl-1H-pyrazol-1-yl)acetyl)-3-methylpiperazin-1-yl)-5-fluoro-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	526
	5-fluoro-2-((3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	549






	2-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	501
	1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	486
	5-fluoro-2-{(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	563
	1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	497
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	464

	5-fluoro-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	548
	5-fluoro-2-{(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	529
	5-fluoro-2-{(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	564
	5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide	503
	1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one	493

 Chiral	5-fluoro-2-((3R)-3-methyl-4-((N-methyl-N-(2-methylpropanoyl)glycyl)piperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	531
	1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	575
 Chiral	5-fluoro-2-[(3R)-3-methyl-4-((1H-pyrrolo[2,3-b]pyridin-1-yl)acetyl)piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	548
	1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	515
 Chiral	5-fluoro-2-((3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	529






 Chiral	N-{2-[(2S)-4-{4-fluoro-2-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	570
	2-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	517
	1-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	467
	1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	503
	2-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	530

	1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	491
	1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	559
	1-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	483
	1-{2-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}piperidin-2-one	555
 Chiral	N-{2-[(2R)-4-{4-fluoro-2-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	570

	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	493
	1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	516
	1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	511
	1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	509
	1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	448

	2-(3,5-dimethyl-1H-pyrazol-1-yl)- 1-(2-methyl-4-{2-[2-((methylsulfonyl)pyrimidin-5-yl]- 4-((trifluoromethyl)phenyl}piperazin- 1-yl)ethanone	537
	1-[2-(4-{4-fluoro-2-[2-((methylsulfonyl)pyrimidin-5- yl]phenyl}-2-methylpiperazin-1- yl)-2-oxoethyl]-1,3-dihydro-2H- benzimidazol-2-one	525
	1-[2-(2-methyl-4-{2-[2-((methylsulfonyl)pyrimidin-5- yl]phenyl}piperazin-1-yl)-2- oxoethyl]-1,3-dihydro-2H- benzimidazol-2-one	507
	1-(4-{4-fluoro-2-[2-((methylsulfonyl)pyrimidin-5- yl]phenyl}piperazin-1-yl)-2-(1H- pyrrolo[2,3-b]pyridin-1- yl)ethanone	495
	1-[2-(4-{4-fluoro-2-[2-(4- methylpiperazin-1-yl)pyrimidin-5- yl]phenyl}piperazin-1-yl)-2- oxoethyl]piperidin-2-one	496

	2-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	574
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)ethanone	487
	1-(2-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-oxoethyl)piperidin-2-one	428
	2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)ethanone	560
	2-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	524

	2-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	510
	2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	473
	2-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	506
	1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	540
	2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	492

	1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	496
	1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one	490
	1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	476
	1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	472
	1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	510

In one embodiment, the invention relates to a compound selected from compounds described in Table 1, or the pharmaceutically acceptable salts thereof.

In another embodiment there is a compound selected from

3-(2-Methylsulfanyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile;

1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyridine-2-carbonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(4-{2-[2-(2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

1-{3-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-3,8-diaza-bicyclo[3.2.1]oct-8-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-piperidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

3-(2-Methoxy-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile;

1-(4-{2-[2-((2R,6R)-2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

3-(2-Methylamino-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile;

3-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[4-Methoxy-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-c]pyridin-1-yl-ethanone;

1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-[1,4]Oxazepan-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Ethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(2-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1,3-dihydro-benzoimidazol-2-one;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-(4-{2-[2-((2R,6S)-2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(4-{2-[2-(4-Methyl-piperazin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Piperidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperidin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Methanesulfonyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[4-Chloro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyrimidine-2-carbonitrile;

2-Imidazo[4,5-b]pyridin-3-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-ethoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-morpholin-4-yl-pyrimidin-5-yl)-benzonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1-methyl-1H-indol-6-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-3,6-dihydro-2H-pyridin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-3,6-dihydro-2H-pyridin-1-yl}-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-5-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyridine-2-carboxylic acid amide;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-4-ylamino)-ethanone;

1-{4-[2-Fluoro-6-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-Indazol-2-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(2,4-Dimethyl-imidazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(6-Chloro-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-trifluoromethyl-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(6-Methoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-Indazol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(thiazol-2-ylamino)-ethanone;

4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methylsulfanyl-pyrimidin-5-yl)-benzonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-phenoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methyl-piperazin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperidin-1-yl}-ethanone;

5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidine-2-carbonitrile;

2-Indol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

5-(2-{4-[2-(2,4-Dimethyl-imidazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile;

5-(2-{4-[2-(Thiazol-2-ylamino)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile;

1-[4-(2-Benzo[b]thiophen-2-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methanesulfonyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-methoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(pyridin-3-yloxy)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-[4-(2-Benzothiazol-5-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-methoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(thiazol-2-ylamino)-ethanone;

2-(2,4-Dimethyl-imidazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-[4-(4'-Benzoyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

3,4-Dimethoxy-N-(2-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-benzamide;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

5-(2-{4-[2-(5-Methyl-3-trifluoromethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile;

1-{4-[2-(9H-Carbazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[3'-(5-methyl-[1,3,4]oxadiazol-2-yl)-biphenyl-2-yl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-naphthalen-2-yl-phenyl)-piperazin-1-yl]-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-indol-6-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-[4-(4'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-trifluoromethoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

5-(2-{4-[2-(Pyridin-4-ylamino)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-carbonitrile;

1-[4-(4'-tert-Butyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methyl-3-trifluoromethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-isopropyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

1-[4-(3'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,4,5-trimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-[1,1';4',1'']terphenyl-2-yl-piperazin-1-yl)-ethanone;

1-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-1,3-dihydro-benzoimidazol-2-one;

1-{4-[2-(2-Cyclopropylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methoxy-pyrimidin-5-yl)-benzonitrile;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-tert-Butyl-thiazol-4-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

3-(2-Methanesulfonyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl)-acetyl]-piperazin-1-yl]-benzonitrile;

2-(3-tert-Butyl-[1,2,4]thiadiazol-5-ylamino)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-fluoro-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

1-{4-[4-Methanesulfonyl-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-3-ylamino)-ethanone;

2-Benzotriazol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

N-(2-{4-[2-(6-Cyano-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3,4-dimethoxy-benzamide;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-4-ylamino)-ethanone;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(2,5-dimethyl-imidazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-indol-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

N-(2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-yl)-methanesulfonamide;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(pyridin-4-ylamino)-ethanone;

5-(2-{4-[2-(5-Methyl-3-trifluoromethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid amide;

1-{4-[3-(2-Morpholin-4-yl-pyrimidin-5-yl)-pyridin-2-yl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-pyrimidin-5-yl-phenyl)-piperazin-1-yl]-ethanone;

1-[4-(3',5'-Bis-trifluoromethyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Amino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methanesulfonyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-sulfonic acid amide;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-imidazo[4,5-b]pyridin-1-yl-ethanone;

1-[4-(2'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-hydroxymethyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(thiazol-2-ylamino)-ethanone;

3,4-Dimethoxy-N-(2-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-benzamide;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-hydroxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-[1,1';3',1'']terphenyl-2-yl-piperazin-1-yl)-ethanone;

1-[4-(3',4'-Dimethoxy-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-quinolin-3-yl-phenyl)-piperazin-1-yl]-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-ethoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(5-methoxy-pyrazin-2-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-morpholin-4-yl-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-ethoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

1-[4-(2-Benzo[1,3]dioxol-5-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-[4-(3'-Acetyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1-methyl-1H-indol-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-quinolin-6-yl-phenyl)-piperazin-1-yl]-ethanone;

1-[4-(4'-Acetyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-ethoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

4-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester;

5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-nicotinic acid ethyl ester;

1-{4-[2-(1-Benzyl-1H-pyrazol-4-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-thiophen-2-yl-biphenyl-2-yl)-piperazin-1-yl]-ethanone;

2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-sulfonic acid dimethylamide;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(5-phenyl-thiophen-2-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-[4-(2-Dibenzothiophen-2-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-3-carboxylic acid methylamide;

1-[4-(4'-Benzyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

N-(2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-3-yl)-acetamide;

5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester;

1-[4-(3'-Benzyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-[4-(2-Benzo[b]thiophen-3-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(1H-Benzimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-[4-(2-Benzooxazol-2-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-imidazo[4,5-c]pyridin-2-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(5-Chloro-1H-benzimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-1H-benzimidazole-5-carbonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-(4-{2-[2-((R)-3-Dimethylamino-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-(4-{2-[(1R,4S)-2-(2-Aza-bicyclo[2.2.1]hept-2-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[4-fluoro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[4-Chloro-2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methanesulfonyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-4-carboxylic acid amide;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxy-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-{4-[4-Chloro-2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[4-Chloro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(4-{2-[2-(3,3-Difluoro-azetidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-fluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(5-Fluoro-pyrrolo[2,3-b]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[4-Chloro-2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-3-carboxylic acid amide;

1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-4-carboxylic acid methylamide;

1-(4-{2-[2-(3,3-Difluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-hydroxy-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-fluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(4-Chloro-3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-ethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-[1,4]oxazepan-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[4-Chloro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{3-Methyl-4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3-methyl-1,3-dihydro-benzoimidazol-2-one;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-trifluoromethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-(4-{2-[2-(4,4-Dimethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-{2-[4-(1-hydroxy-1-methyl-ethyl)-piperidin-1-yl]-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-{4-[4-Chloro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{2-Methyl-4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[4-Chloro-2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[(1S,4S)-2-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-(4-{2-[2-(3,5-Dimethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-[4-fluoro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl]-ethanone;

2-(7-Chloro-pyrrolo[2,3-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyridin-4-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-2,3-dihydro-isoindol-1-one;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(octahydro-isoquinolin-2-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

1-(4-{2-[2-(4,4-Difluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[4-Chloro-2-(2-methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Methoxy-pyridin-4-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(4-{2-[2-(4-Dimethylamino-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methoxy-4-methyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-pyrrolidin-1-yl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(4-Chloro-pyrrolo[2,3-d]pyrimidin-7-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-2,3-dihydro-isoindol-1-one;

2-(4-Chloro-pyrrolo[3,2-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

4-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methylamide;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[3,2-c]pyridin-1-yl-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methyl-3-nitro-pyrazol-1-yl)-ethanone;

2-(5-Chloro-pyrrolo[3,2-b]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methoxy-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone;

4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methylamino-pyrimidin-5-yl)-benzonitrile;

2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-2H-phthalazin-1-one;

1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1H-indole-2,3-dione;

3-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3H-quinazolin-4-one;

2-(6-Bromo-pyrrolo[3,2-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-((S)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

N-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-N-methyl-acetamide;

1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1H-pyrrolo[2,3-b]pyridine-4-carbonitrile;

(R)-1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-pyrrolidine-2-carboxylic acid amide;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(5-methoxy-pyrrolo[2,3-b]pyridin-1-yl)-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3-methyl-benzo[b]thiophen-2-yl)-ethanone;

2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl)-acetyl]-piperazin-1-yl}-phenyl}-2H-isoquinolin-1-one;

2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1,1-dioxo-1,2-dihydro-1H-benzo[d]isothiazol-3-one;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(pyrrolidine-1-carbonyl)-pyridin-4-yl]-phenyl}-piperazin-1-yl)-ethanone;

2-(4,5-Dichloro-imidazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[3-(2-Morpholin-4-yl-pyrimidin-5-yl)-pyridin-4-yl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-((R)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-((S)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

(S)-1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-pyrrolidine-2-carboxylic acid methylamide;

2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-isoindole-1,3-dione;

1-((R)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carboxylic acid methylamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide;

1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-2,3-dihydro-1H-isoindol-1-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}benzamide;

2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone;

1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one;

2-{(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl}-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide;

1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl}benzamide;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl)piperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one;

1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone;
1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide;
1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl})piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;
2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]ethanone;
2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide;
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;
1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone
1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;
1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one
5-fluoro-2-[(3S)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide;
1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)piperidin-2-one;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone;

1-{2-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

N-{2-[(2S)-4-{4-fluoro-2-[(2-methoxypyridin-4-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl}benzamide;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

5-fluoro-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

2-{(3S)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl}-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide;

N-{2-[(2R)-4-{4-fluoro-2-[(2-methoxypyridin-4-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide;

2-{(3S)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl}-5-fluoro-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-(2-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}benzamide;

1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-2-{(3S)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}piperidin-2-one;

1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

5-fluoro-2-{(3S)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

5-fluoro-2-[(3S)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl}piperidin-2-one;

2-[(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl]-5-fluoro-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

2-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

5-fluoro-2-[(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

5-fluoro-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

5-fluoro-2-[(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

5-fluoro-2-[(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one;

5-fluoro-2-[(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

5-fluoro-2-{(3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide;

N-{2-[2-(2S)-4-{4-fluoro-2-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)carbonyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide;

2-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

1-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

2-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

1-{2-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

N-{2-[(2R)-4-{4-fluoro-2-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

2-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)ethanone;

1-(2-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-oxoethyl)piperidin-2-one;

2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)ethanone;

2-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

2-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one;

1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one;

1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

and the pharmaceutically acceptable salts thereof.

In another embodiment, the invention relates to a compound selected from:

3-(2-Methylsulfonyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile;

1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

4-Amino-N-[2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-3-methoxy-benzamide;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-naphthalen-1-yl-phenyl)-piperazin-1-yl]-ethanone;

1-Methyl-1H-imidazole-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyridine-2-carbonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(3,4-dimethoxy-benzylamino)-ethanone

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-piperidin-1-yl-ethanone;

Isoquinoline-6-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

Thiazole-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

1-(4-{2-[2-(2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

4,5-Dimethyl-furan-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

1-(2-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-imidazolidin-2-one;

1-[4-(4'-Dimethylaminomethyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

Isoquinoline-7-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide;

1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;
2-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-2,3-dihydro-1H-isoindol-1-one;
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}benzamide;

2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone;

1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one;

2-{(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl}-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide;

1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-yl)acetyl]piperazin-1-yl}benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl}benzamide;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one;

1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone;

1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]ethanone;

2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone
and the pharmaceutically acceptable salts thereof.

For all compounds disclosed hereinabove in this application, in the event the nomenclature is in conflict with the structure, it shall be understood that the compound is defined by the structure.

The invention also relates to pharmaceutical preparations, containing as active substance one or more compounds of the invention, or the pharmaceutically acceptable derivatives thereof, optionally combined with conventional excipients and/or carriers.

Compounds of the invention also include their isotopically-labelled forms. An isotopically-labelled form of an active agent of a combination of the present invention is identical to said active agent but for the fact that one or more atoms of said active agent have been replaced by an atom or atoms having an atomic mass or mass number different from the atomic mass or mass number of said atom which is usually found in nature. Examples of isotopes which are readily available commercially and which can be incorporated into an active agent of a combination of the present invention in accordance with well established procedures, include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, *e.g.*, ²H, ³H, ¹³C, ¹⁴C, ¹⁵N, ¹⁸O, ¹⁷O, ³¹P, ³²P, ³⁵S, ¹⁸F, and ³⁶Cl, respectively. An active agent of a combination of the present invention, a prodrug thereof, or a pharmaceutically acceptable salt of either which contains one or more of the above-mentioned isotopes and/or other isotopes of other atoms is contemplated to be within the scope of the present invention.

The invention includes the use of any compounds of described above containing one or more asymmetric carbon atoms may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. Isomers shall be defined as being enantiomers and diastereomers. All such isomeric forms of these compounds are expressly included in the present invention. Each stereogenic carbon may be in the R or S configuration, or a combination of configurations.

Some of the compounds of the invention can exist in more than one tautomeric form.

The invention includes methods using all such tautomers.

All terms as used herein in this specification, unless otherwise stated, shall be understood in their ordinary meaning as known in the art. For example, "C₁₋₆ alkoxy" is a C₁₋₆ alkyl with a terminal oxygen, such as methoxy, ethoxy, propoxy, butoxy. All alkyl, alkenyl, and alkynyl groups shall be understood as being branched or unbranched where structurally possible and unless otherwise specified. Other more specific definitions are as follows:

The term "alkyl" refers to both branched and unbranched alkyl groups. It should be understood that any combination term using an "alk" or "alkyl" prefix refers to analogs according to the above definition of "alkyl". For example, terms such as "alkoxy", "alkythio" refer to alkyl groups linked to a second group via an oxygen or sulfur atom. "Alkanoyl" refers to an alkyl group linked to a carbonyl group (C=O).

In all alkyl groups or carbon chains, one or more carbon atoms can be optionally replaced by heteroatoms such as O, S or N. It shall be understood that if N is not substituted then it is NH. It shall also be understood that the heteroatoms may replace either terminal carbon atoms or internal carbon atoms within a branched or unbranched carbon chain.

Such groups can be substituted as herein above described by groups such as oxo to result in definitions such as but not limited to: alkoxy carbonyl, acyl, amido and thioxo.

As used herein, "nitrogen" and "sulfur" include any oxidized form of nitrogen and sulfur and the quaternized form of any basic nitrogen. For example, for a -S-C₁₋₆ alkyl radical, unless otherwise specified, shall be understood to include -S(O)-C₁₋₆ alkyl and -S(O)₂-C₁₋₆ alkyl.

The term "C₃₋₁₀ cycloalkyl" refers to a nonaromatic 3 to 10-membered (but preferably, 3 to 6-membered) monocyclic carbocyclic radical or a nonaromatic 6 to 10-membered fused bicyclic, bridged bicyclic, or spirocyclic carbocyclic radical. The C₃₋₁₀ cycloalkyl

may be either saturated or partially unsaturated, and the carbocycle may be attached by any atom of the cycle which results in the creation of a stable structure. Non-limiting examples of 3 to 10-membered monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptanyl, cycloheptenyl, and cyclohexanone. Non-limiting examples of 6 to 10-membered fused bicyclic carbocyclic radicals include bicyclo[3.3.0]octane, bicyclo[4.3.0]nonane, and bicyclo[4.4.0]decanyl (decahydronaphthalenyl). Non-limiting examples of 6 to 10-membered bridged bicyclic carbocyclic radicals include bicyclo [2.2.2]heptanyl, bicyclo[2.2.2]octanyl, and bicyclo[3.2.1]octanyl. Non-limiting examples of 6 to 10-membered spirocyclic carbocyclic radicals include but are not limited to spiro[3,3]heptanyl, spiro[3,4]octanyl and spiro[4,4]heptanyl.

As used herein, the term “aryl” refers to an aromatic hydrocarbon rings containing from six to ten carbon ring atoms (e.g., a C₆₋₁₀ aryl).. The term C₆₋₁₀ aryl includes monocyclic rings and bicyclic rings where at least one of the rings is aromatic. Non-limiting examples of C₆₋₁₀ aryls include phenyl, indanyl, indenyl, benzocyclobutanyl, dihydronaphthyl, tetrahydronaphthyl, naphthyl, benzocycloheptanyl and benzocycloheptenyl.

As used herein, the term “heterocyclyl” refers to a “5 to 11-membered heterocycle” and includes stable nonaromatic 4-8 membered monocyclic heterocyclic radical or a stable nonaromatic 6 to 11-membered fused bicyclic, bridged bicyclic or spirocyclic heterocyclic radical. The 5 to 11-membered heterocycle consists of carbon atoms and one or more, preferably from one to four heteroatoms chosen from nitrogen, oxygen and sulfur. The heterocycle may be either saturated or partially unsaturated. Non-limiting examples of nonaromatic 4-8 membered monocyclic heterocyclic radicals include tetrahydrofuranyl, azetidiny, pyrrolidinyl, pyranly, tetrahydropyranly, dioxanyl, thiomorpholinyl, 1,1-dioxo-1λ⁶-thiomorpholinyl, morpholinyl, piperidinyl, piperazinyl, and azepinyl. Non-limiting examples of nonaromatic 6 to 11-membered fused bicyclic radicals include octahydroindolyl, octahydrobenzofuranyl, and octahydrobenzothiophenyl. Non-limiting examples of nonaromatic 6 to 11-membered

bridged bicyclic radicals include 2-azabicyclo[2.2.1]heptanyl, 3-azabicyclo[3.1.0]hexanyl, and 3-azabicyclo[3.2.1]octanyl. Non-limiting examples of nonaromatic 6 to 11-membered spirocyclic heterocyclic radicals include 7-aza-spiro[3,3]heptanyl, 7-spiro[3,4]octanyl, and 7-aza-spiro[3,4]octanyl.

As used herein, the term “heteroaryl” refers to a “5 to 11-membered heteroaryl” and includes aromatic 5 to 6-membered monocyclic heteroaryls and aromatic 7 to 11-membered heteroaryl bicyclic rings where at least one of the rings is aromatic, wherein the heteroaryl ring contains 1-4 heteroatoms such as N, O and S. Non-limiting examples of 5 to 6-membered monocyclic heteroaryl rings include furanyl, oxazolyl, isoxazolyl, oxadiazolyl, pyranlyl, thiazolyl, pyrazolyl, pyrrolyl, imidazolyl, tetrazolyl, triazolyl, thienyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, and purinyl. Non-limiting examples of 7 to 11-membered heteroaryl bicyclic rings include benzimidazolyl, 1,3-dihydrobenzoimidazol-2-one, quinolinyl, dihydro-2*H*-quinolinyl, isoquinolinyl, quinazolinyl, indazolyl, thieno[2,3-*d*]pyrimidinyl, indolyl, isoindolyl, indazolyl, benzotriazolyl, benzofuranyl, benzopyranlyl, benzodioxolyl, benzoxazolyl, benzothiazolyl, pyrrolo[2,3-*b*]pyridinyl, , and imidazo[4,5-*b*]pyridinyl.

It will be understood that when a heterocyclyl or heteroaryl contains a S ring atom, such S ring atom can be present in the ring in its divalent, tetravalent, or hexavalent form, i.e., -S-, -S(O)- or -S(O)₂-.

Each aryl or heteroaryl unless otherwise specified includes its partially or fully hydrogenated derivatives. For example, quinolinyl may include decahydroquinolinyl and tetrahydroquinolinyl, naphthyl may include its hydrogenated derivatives such as tetrahydronaphthyl. Other partially or fully hydrogenated derivatives of the aryl and heteroaryl compounds described herein will be apparent to one of ordinary skill in the art. It will be understood that one to three carbon ring moieties in each of the C₃₋₁₀ carbocyclic rings, the 5 to 11-membered heterocyclic rings, the nonaromatic portion of the bicyclic aryl or heteroaryl rings, and the nonaromatic portion of the bicyclic heteroaryl rings can independently be replaced with a carbonyl, thiocarbonyl, or iminyl

moiety, i.e., -C(=O)- , -C(=S)- and $\text{-C(=NR}^8\text{)-}$, respectively, where R^8 is as defined above. The term “heteroatom” as used herein shall be understood to mean atoms other than carbon such as O, N, and S.

The term “halogen” as used in the present specification shall be understood to mean bromine, chlorine, fluorine or iodine. The definitions “halogenated”, “partially or fully halogenated”; partially or fully fluorinated; “substituted by one or more halogen atoms”, includes for example, mono, di or tri halo derivatives on one or more carbon atoms. For alkyl, a non-limiting example would be $\text{-CH}_2\text{CHF}_2$, -CF_3 etc.

Each alkyl, carbocycle, heterocycle or heteroaryl, or the analogs thereof, described herein shall be understood to be optionally partially or fully halogenated.

The compounds of the invention are only those which are contemplated to be ‘chemically stable’ as will be appreciated by those skilled in the art. For example, a compound which would have a ‘dangling valency’, or a ‘carbanion’ are not compounds contemplated by the inventive methods disclosed herein.

The invention includes pharmaceutically acceptable derivatives of compounds of formula (I). A “pharmaceutically acceptable derivative” refers to any pharmaceutically acceptable salt or ester, or any other compound which, upon administration to a patient, is capable of providing (directly or indirectly) a compound useful for the invention, or a pharmacologically active metabolite or pharmacologically active residue thereof. A pharmacologically active metabolite shall be understood to mean any compound of the invention capable of being metabolized enzymatically or chemically. This includes, for example, hydroxylated or oxidized derivative compounds of the invention.

Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acids include hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycolic, lactic, salicylic, succinic, toluene-p-sulfuric, tartaric, acetic, citric,

methanesulfonic, formic, benzoic, malonic, naphthalene-2-sulfuric and benzenesulfonic acids. Other acids, such as oxalic acid, while not themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds and their pharmaceutically acceptable acid addition salts. Salts derived from appropriate bases include alkali metal (*e.g.*, sodium), alkaline earth metal (*e.g.*, magnesium), ammonium and N-(C₁-C₄ alkyl)₄⁺ salts.

In addition, within the scope of the invention is use of prodrugs of compounds of the invention. Prodrugs include those compounds that, upon simple chemical transformation, are modified to produce compounds of the invention. Simple chemical transformations include hydrolysis, oxidation and reduction. Specifically, when a prodrug is administered to a patient, the prodrug may be transformed into a compound disclosed hereinabove, thereby imparting the desired pharmacological effect.

The compounds of formula (I) may be made using the general synthetic methods described below, which also constitute part of the invention.

GENERAL SYNTHETIC METHODS


The compounds of the invention may be prepared by the methods described below. In each of the schemes below, the groups R¹ to R⁵, A, B, D and X are as defined above for general formula (I) unless noted otherwise. Optimum reaction conditions and reaction times may vary depending on the particular reactants used. Unless otherwise specified, solvents, temperatures, pressures and other reaction conditions may be readily selected by one of ordinary skill in the art. Amide bond formations may be carried out by standard coupling conditions well-known in the art (see, for example, M. Bodanszky, The Practice of Peptide Synthesis (Springer-Verlag: 1984), which is hereby incorporated by reference in its entirety), for example, by reacting a carboxylic acid and an amine in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) and 1-hydroxybenzotriazole. Specific procedures are provided in the Synthetic Examples section. Typically, reaction

progress may be monitored by thin layer chromatography (TLC) or HPLC-MS if desired. Intermediates and products may be purified by chromatography on silica gel, recrystallization, HPLC and/or reverse phase HPLC.

Starting materials and reagents are either commercially available or may be prepared by one skilled in the art using methods described in the chemical literature. Initial products of formula (I) may be modified further by methods known in the art to produce additional compounds of formula (I).

Compounds of formula (I) where A is N and where ----- is a single bond may be prepared as shown in Scheme 1

Scheme 1




As illustrated in Scheme 1, intermediate II, where P is a suitable amine protecting group such as a t-Boc (*tert*-butoxycarbonyl) group, is reacted with intermediate III, a boronic acid bearing R², under Suzuki-Miyaura coupling conditions (see for example, F. Bellina et al., *Synthesis*, 2004, 2419-2440) to provide intermediate IV. Removal of the protecting group, for example by treatment with acid if P is a t-Boc group, provides intermediate V. Coupling intermediate V with carboxylic acid intermediate VI under

peptide coupling conditions known in the art, for example by reacting in the presence of O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU) and a suitable base such as di-isopropylethylamine in a suitable solvent such as methylene chloride provides the desired compound of formula (I).

Compounds of formula (I) may also be made by a variation of this method illustrated in Scheme 2.


Scheme 2



As illustrated above, the peptide coupling with carboxylic acid may be carried out with intermediate VII bearing a Br substituent or similar functionality capable of undergoing the Suzuki-Miyaura coupling, for example a Cl, I or triflate substituent, providing intermediate VIII. Intermediate VIII is then coupled with intermediate III to provide the desired compound of formula (I).

Compounds of formula (I) where A is C may be prepared as described in Scheme 3.


Scheme 3



As illustrated above 1,2-dibromobenzene, optionally substituted with R^1 is reacted with a boron ester bearing R^2 such as the pinacol ester shown, under Suzuki-Miyaura coupling conditions to provide intermediate XI. Intermediate XI then undergoes a second Suzuki-Miyaura coupling reaction with intermediate XII, where P is an amine protecting group such as a t-Boc group, to provide intermediate XIII. Deprotection, as described above in Scheme 1, provides intermediate XIV. Intermediate XIV then undergoes peptide coupling with intermediate VI as described in Scheme 1 to provide the compound of formula (I) where A is C and ----- is a double bond. Reduction of the double bond by methods well known in the art, for example by treating with hydrogen in the presence of Pd on carbon, provides the compound of formula (I) where A is C and ----- is a single bond.

Scheme 4 illustrates an alternate procedure, useful for preparing compounds of formula (I) where R^2 is benzoimidazole, benzoxazole or benzothiazole.

Scheme 4




As illustrated above, intermediate VI is coupled with intermediate XIV, where R is an alkyl group such as methyl or ethyl, as described in Scheme 2, followed by hydrolysis of the ester, for example, by treatment with aqueous NaOH in THF/MeOH, to provide XVII. Intermediate XVII is then coupled with intermediate XVIII, where Y is NH₂, OH, or SH, to provide intermediate XIX. Cyclization, for example by heating in a suitable solvent such as dichloroethane in the presence of a suitable catalyst such as p-TsOH, provides the desired compound of formula (I) where Y is NH, O, or S.

All of the compounds in Table I were prepared by the methods illustrated above and in the Synthetic Examples section below.

SYNTHETIC EXAMPLES

Example 1: Synthesis of 1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone




1

4-(2-Bromophenyl)-piperazine-1-carboxylic acid, t-butyl ester (10 g, 29.30 mmol) and 2-methoxypyrimidine-5-boronic acid (5 g, 32.48 mmol) are dissolved in DMF (200 mL) under a stream of nitrogen. To this mixture is added a 2M aqueous solution of sodium carbonate (73.26 mL, 150 mmol) followed by bis(triphenylphosphine) palladium (II) chloride (2.06 g, 2.93 mmol). The resulting mixture is allowed to stir at 100 °C for 30 min. A precipitate forms. Water is added and the resulting mixture is filtered to provide a gray colored solid which is air dried. The solid is then re-dissolved in dichloromethane (20 mL) and loaded on a 340g Biotage SNAP column and eluted with 35% ethyl acetate in hexanes to provide 4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazine-1-carboxylic acid t-butyl ester (5.87 g, 54% yield).

To a solution of the above ester (5.67 g, 15.31 mmol) dissolved in dichloromethane (50 mL) is added TFA (10 mL, 129.80 mmol). The reaction is stirred at ambient temperature for 20 h. The reaction is then concentrated by rotary evaporation and the residue is reconstituted with EtOAc (200 mL) and washed with 1N aqueous NaOH solution (2 x 150 mL). The organic solution is dried with magnesium sulfate, filtered and concentrated by rotary evaporation to provide 2-methoxy-5-(2-piperazin-1-yl-phenyl)-pyrimidine (3.85 g, 14.24 mmol, 93% yield) as a brown solid.

To 2-methoxy-5-(2-piperazin-1-yl-phenyl)-pyrimidine (7.4 g, 27.37 mmol) dissolved in dichloromethane (100 mL) is added pyrrolo[2,3-b]pyridin-1-yl-acetic acid (4.82 g, 27.37 mmol), TBTU (8.79 g, 27.37 mmol), and DIPEA (14.66 mL, 82.12 mmol). The reaction is stirred for 16 h at ambient temperature. The product is purified on a 100 g Biotage SNAP column with 40% EtOAc in hexanes to provide the title compound (8.3 g, 19.37 mmol, 71% yield).

Example 2: Synthesis of 1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone



To a solution of 2-fluoro-4-trifluoromethylnitrobenzene (1 g, 4.78 mmol) and Hunig's base (2.45 mL, 14.35 mmol) in DMF (10 mL) is added 1-Boc-piperazine (1.14 g, 7.17 mmol). The resulting reaction is allowed to stir at 80 °C in an oil bath for 1 h. Water (30 mL) is added and mixture is extracted with EtOAc. The organic layer is washed with water, brine, dried (Na₂SO₄) and concentrated. The resulting 4-(2-nitro-4-trifluoromethyl-

phenyl)-piperazine-1-carboxylic acid t-butyl ester is isolated as an orange oil (1.80g, 100%) and is used in the next step without further purification.

To a solution of the above intermediate (1.80 g, 4.80 mmol) in EtOH (25 mL) is added Pd/C (5% on activated carbon, 100 mg) and the mixture is purged and back filled with H₂ three times via a H₂ balloon. The resulting reaction is allowed to stir at ambient temperature overnight. The reaction mixture is then filtered through a thin pad of diatomaceous earth and washed with MeOH (10mL). The filtrate is concentrated to give a dark oil. The crude oil is purified using biotage chromatography eluting with 0-30% EtOAc/hexane. Removal of the solvent provides 4-(2-amino-4-trifluoromethyl-phenyl)-piperazine-1-carboxylic acid t-butyl ester as a colorless solid (1.65g, 99%).

Isoamyl nitrite (0.76 mL, 5.68 mmol) is added dropwise to an ice-bath cooled solution of the above amine intermediate (1.40 g, 4.05 mmol) in dried acetonitrile (20 mL) under nitrogen. The resulting mixture turns light yellow during addition. The reaction is allowed to stir at 0 °C for 20 min. Then CuBr (872 mg, 6.08 mmol) is added in small portions. The reaction gradually turns to dark green. It is allowed to warm up to ambient temperature in 30 min and stirred overnight.

Water (20 mL) is added and mixture is extracted with EtOAc. The organic layer is washed with water, brine, dried (Na₂SO₄) and concentrated. The resulting dark red oil is purified using biotage chromatography eluting with 0-25% EtOAc/hexane. Removal of the solvent provides 4-(2-bromo-4-trifluoromethyl-phenyl)-piperazine-1-carboxylic acid t-butyl ester red oil (1.03 g, 62%).


TFA (1 mL) is added dropwise to a solution of the above intermediate (550mg, 1.34 mmol) in methylene chloride (5 mL). The reaction is allowed to stir at ambient temperature for 2 h. Solvent is removed by rotavap and to the residue is added 2N NaOH to adjust pH>12. The resulting solution is extracted with EtOAc. The organic layer is washed with water, brine, dried (Na₂SO₄) and concentrated to provide 1-(2-bromo-4-

trifluoromethyl-phenyl)-piperazine as a red oil (415mg, 99%). The product is used in the next step without further purification.

1-(2-Bromo-4-trifluoromethyl-phenyl)-piperazine (225 mg, 0.73 mmol), pyrrolo[2,3-b]pyridin-1-yl-acetic acid (154 mg, 0.87 mmol), TBTU (304 mg, 0.95 mmol) and triethylamine (304 μ L, 2.18 mmol) are combined in dichloromethane (3 mL). The resulting reaction is allowed to stir at ambient temperature for 2 h. Water is added and the mixture is extracted with EtOAc. The organic layer is washed with water, brine, dried (Na_2SO_4) and concentrated. The crude product is purified using biotage chromatography eluting with 0-50% EtOAc/hexane. Removal of the solvent provides 1-[4-(2-bromo-4-trifluoromethyl-phenyl)-piperazin-1-yl]-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone as a colorless foam (268 mg, 79%).

The above intermediate (150 mg, 0.32 mmol), 2-(methoxy) pyrimidine-5-boronic acid (70 mg, 0.38 mmol), bis(triphenylphosphine)palladium (II) chloride (23 mg, 0.032 mmol), aqueous Na_2CO_3 solution (2N, 803 μ L, 1.6mmol) and dimethylformamide (2mL) are added to a 2-5 mL microwave tube. The reaction is carried out in a microwave oven at 100 °C for 30 min. The mixture turns black. The reaction mixture is filtered and washed with a bit MeOH/water (1/0.1mL). The clear filtrate is purified through prep-HPLC. Removal of the solvent gives the title compound as a colorless foam (110mg, 69%).

Example 3: Synthesis of 2-(3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperidin-1-yl}-ethanone



2-(4-Morpholino)pyrimidine-5-boronic acid pinacol ester (300 mg, 1.03 mmol), 1,2-dibromobenzene (243 mg, 1.03 mmol), Bis(triphenylphosphine)palladium (II) chloride (72 mg, 0.10 mmol), aqueous Na_2CO_3 (2N, 2.58 mL, 5.15 mmol) and DMF (5 mL) are added to a 10 mL microwave tube. Reaction is carried out in a microwave oven at 100 °C for 30 min. The reaction mixture turns black. The mixture is poured into water and extracted with EtOAc. The organic layers are washed with water, brine, dried (Na_2SO_4) and concentrated. The crude product is purified using biotage eluting with 0-30% EtOAc/Hexane. The desired mono-coupled product elutes at 30% EtOAc. The di-coupled by-product elutes earlier. Removal of the solvent gives 4-[5-(2-bromo-phenyl)-pyrimidin-2-yl]-morpholine as a white solid (130 mg, 39%).


The above intermediate (150 mg, 0.47 mmol), 3,6-dihydro-2H-pyridine-1-tert-butoxycarbonyl-4-boronic acid, pinacol ester (174 mg, 0.56 mmol), bis(triphenyl phosphine)palladium (II) chloride (33 mg, 0.05 mmol), aqueous Na₂CO₃ (2N, 1.17 mL, 2.34 mmol) and DMF (4 mL) are added to a 10 mL microwave tube. The reaction is carried out in a microwave oven at 100°C for 30 min. The mixture turns black. The reaction mixture is poured into water and a precipitate forms. The mixture is filtered and the cake is washed with water. The solid is dissolved and washed with dichloromethane/MeOH (1/1) and the dark solution is filtered and collected. The solvent is removed from the filtrate by rotavap and the residue, dissolved with a bit of methylene chloride is purified using biotage chromatography eluting with 0-30% EtOAc/hexane to provide 4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester as a colorless foam (195 mg, 98%).

TFA (1mL) is added dropwise to a solution of the above t-butyl ester (195 mg, 0.46 mmol) in methylene chloride (5 mL). The reaction is allowed to stir at ambient temperature for 2 h. Solvent is removed by rotavap and to the residue is added 2N NaOH to adjust pH>12. The resulting solution is extracted with EtOAc. The organic layer is washed with water, brine, dried (Na₂SO₄) and concentrated. The crude 4-{5-[2-(1,2,3,6-tetrahydro-pyridin-4-yl)-phenyl]-pyrimidin-2-yl}-morpholine is isolated as a red oil (149 mg, 100%) and is used in the next step without further purification.

The above intermediate (60 mg, 0.19 mmol), pyrrolo[2,3-b]pyridin-1-yl-acetic acid (36 mg, 0.21 mmol), TBTU (78 mg, 0.24 mmol) and triethylamine (78μl, 0.56mmol) are combined in dichloromethane (3 mL). The resulting reaction is allowed to stir at ambient temperature for 2 h. Water is added and the mixture is extracted with EtOAc. The organic layer is washed with water, brine, dried (Na₂SO₄) and concentrated. The crude product is purified using biotage chromatography eluting with 0-100% EtOAc/hexane. Removal of the solvent provides 1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-3,6-dihydro-2H-pyridin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone as a colorless foam (55mg, 61%).

To the solution of the above intermediate (55 mg, 0.11mmol) in EtOH (5 mL) is added Pd/C (5% on activated carbon, 5 mg) and the mixture is purged and back filled with H₂ three times using a H₂ balloon. The resulting reaction is allowed to stir at ambient temperature overnight then is filtered and the solid is washed with a bit MeOH (5mL). The filtrate is collected and concentrated on a rotavap. The residue is re-dissolved in a minimum amount of DMSO/water. It was then purified using prep-HPLC. Removal of the solvent gave the title compound as a white solid (25 mg, 45%).

Example 4: Synthesis of 1-{4-[2-(1*H*-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone



2-Piperazin-1-yl-benzoic acid (1.00 g, 4.89 mmol) is suspended in 10 mL of MeOH.. To this is added 5 mL of conc. H₂SO₄. The mixture is stirred for 16 h resulting in a white

precipitate. The reaction volume is increased by an additional 210 mL of MeOH and 65 mL of H₂SO₄. The mixture stirred for 2 h at room temperature and then heated at reflux for 12 h. The reaction volume is concentrated to approximately 175 mL and applied to an ion exchange column, eluting with 5 x 200 mL of 10% NH₃/MeOH. The product fractions are concentrated and then co-evaporated with toluene to remove residual water providing 0.667 g of 2-piperazin-1-yl-benzoic acid methyl ester.

To a vial containing the 2-piperazin-1-yl-benzoic acid methyl ester (0.667 g, 3.03 mmol) is added a pre-mixed solution containing (3,5-dimethyl-pyrazol-1-yl)-acetic acid (0.428 g, 2.78 mmol) and carbonyl diimidazole (0.567 g, 3.50 mmol). The mixture is stirred overnight and then diluted with 200 mL of EtOAc followed by 200 mL of sat. NH₄Cl. The organic phase is washed with 2 x 200 mL of H₂O and 1 x 200 mL of brine. The organic phase is dried with MgSO₄, filtered and concentrated to give 0.849 g of 2-{4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzoic acid methyl ester.

2-{4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzoic acid methyl ester (0.848 g, 2.38 mmol) is dissolved in 10 mL of 1:1 THF/MeOH. 2 mL of 15% aq NaOH is added and the mixture is stirred overnight. The mixture is concentrated to dryness and suspended in 50 mL of H₂O. The pH of the solution is adjusted to 4 by careful addition of conc. HCl. The aqueous phase is extracted with 3 x 100 mL of CH₂Cl₂. The organic phase is dried with MgSO₄, filtered and concentrated to give 0.730 g of {4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzoic acid in 90% yield.


2-{4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzoic acid (0.107 g, 0.296 mmol) is dissolved into 5 mL of dry CH₂Cl₂ under argon. The mixture is cooled to 4⁰ C and a 2 M solution of oxalyl chloride (0.150 mL, 0.300 mmol) is added in a dropwise fashion resulting in gas evolution. The reaction is stirred for 30 min at 4⁰ C and then 1 drop of DMF is added. The mixture is stirred at room temperature for 30 min, cooled to 4⁰ C and 1,2-diaminobenzene (0.048 g, 0.444 mmol) is added. The reaction is warmed to room temperature and stirred for 1h. The reaction is then diluted with 50 mL CH₂Cl₂ followed by 20 mL of sat. NH₄Cl. The mixture is extracted with 3 x 20 mL of CH₂Cl₂.

The organic phase is dried with MgSO_4 , filtered and concentrated. The crude product is applied to a SiO_2 prep plate and eluted with (2.5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$) to give 58.4 mg of *N*-(2-amino-phenyl)-2-{4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzamide.

The *N*-(2-amino-phenyl)-2-{4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzamide (0.050 g, 0.116 mmol) is dissolved into 2 mL of dichloroethane and placed in a microwave tube. To this is added a catalytic amt. of *p*-TsOH. The mixture is heated in a microwave at 130°C for 20 min and then 160°C for 30 min. The reaction mixture is concentrated to dryness and the product is purified using a Gilson Prep HPLC system eluting with 20%-100% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ to give 43 mg of desired product. Impure fractions are concentrated and purified using a SiO_2 prep plate eluting with (5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$) to give an additional 17.5 mg of 1-{4-[2-(1*H*-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone.

1-[4-(2-Benzooxazol-2-yl-phenyl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone is prepared in an analogous fashion.

Example 5: Synthesis of intermediate 4-(2-carboxy-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester




Method A.

Piperazin-1-yl-benzoic acid (2.00 g, 9.79 mmol) is suspended in 100 mL of CH₂Cl₂. To this is added diisopropylethyl amine (2.22 mL, 12.00 mmol) and BOC anhydride (2.07 g, 9.50 mmol). The mixture remains heterogeneous after 1h. An additional 100 mL of anhydrous CH₃CN is added and the mixture stirred overnight. The mixture is diluted with 250 mL of EtOAc followed by 250 mL of sat. NH₄Cl. The organic phase is washed with 2 x 250 mL of H₂O and 1 x 250 mL of brine and then is dried with MgSO₄, filtered and concentrated to give 2.37 g of 4-(2-carboxy-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester.

Method B

4-(2-Formyl-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester (0.50 g, 1.72 mmol) is dissolved in 80 mL of dioxane and 20 mL of H₂O. The mixture is cooled to 4⁰ C and then sulfamic acid (1.36 g, 14.00 mmol) is added in one portion. The mixture is stirred for an additional 30 min and then 3 mL of a solution of NaClO₂ (0.343 g, 3.80 mmol) is added in a drop-wise portion. The reaction is quenched by the addition of 50 mL of H₂O and 50 mL of brine. The resulting mixture is extracted with 3x100 mL of CH₂Cl₂. The combined organic phase is washed with 2x50 mL of brine, dried with MgSO₄, filtered and concentrated to give 500 mg of 4-(2-carboxy-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester.

Example 6: Synthesis of 2-(3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-ethanone



4-(2-Carboxy-phenyl)-piperazine-1-carboxylic acid *tert*-butyl ester, (0.060 g, 0.195 mmol) is dissolved in 5 mL of dry CH₂Cl₂ under argon. The solution is cooled to 4° C and a 2 M solution of oxalyl chloride (0.10 mL, 0.21 mmol) is added in a drop-wise fashion resulting in gas evolution. After stirring the mixture for 30 min, 1 drop of DMF is added and the mixture is stirred at room temperature for 30 min. The reaction mixture is cooled to 4° C and 4-trifluoromethyl-benzene-1,2-diamine (0.069 g, 0.390 mmol) and diisopropylethyl amine (0.092 mL, 0.500mmol) are added. The reaction is warmed to room temperature and stirred for 1h. The mixture is diluted with 50 mL EtOAc followed by 20 mL of sat. NH₄Cl. The organic phase is washed with 2 x 20 mL of H₂O and 1 x 20 mL of brine. The organic phase is dried with MgSO₄, filtered and concentrated. The residue is applied to a SiO₂ prep plate and eluted with 50%EtOAc/hexanes to give 0.045 g of 4-[2-(2-amino-4-trifluoromethyl-phenylcarbamoyl)-phenyl]-piperazine-1-carboxylic acid *tert*-butyl ester.

The above *tert*-butyl ester (0.045 g, 0.097 mmol) in a vial is dissolved in 1 mL of CH₂Cl₂ and 1 mL of TFA. The mixture is stirred for 1h. and then concentrated to dryness to give crude *N*-(2-amino-4-trifluoromethyl-phenyl)-2-piperazin-1-yl-benzamide trifluoroacetamide. To a vial containing the amine salt is added 1.0 mL of a premixed (1 hr) solution containing (3,5-dimethyl-pyrazol-1-yl)-acetic acid (0.016 g, 0.104 mmol) and carbonyl diimidazole (0.017 g, 0.104 mmol). The mixture is stirred overnight and then 20 mL of sat. NH₄Cl is added followed by 50 mL of CH₂Cl₂. The mixture is washed with 2 x 20 mL of H₂O and 1 x 20 mL of brine. The organic phase is dried with MgSO₄, filtered and concentrated to give 52 mg of *N*-(2-amino-4-trifluoromethyl-phenyl)-2-{4-[2-(3,5-dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-benzamide.

The above benzamide (0.049 g, 0.090 mmol) is dissolved in 2 mL of CH₂Cl₂ and placed in a microwave vial. A catalytic amount of *p*-TsOH is added and then the mixture is heated in the microwave oven for 20 min at 160⁰ C. The mixture is then concentrated under a stream of N₂. The residue is dissolved in MeOH and purified using a Gilson Prep HPLC system eluting with 20%-80% CH₃CN/H₂O to give 36.3 mg of 2-(3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(5-trifluoromethyl-1*H*-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-ethanone in 77% yield.


The following compounds are prepared analogously:

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1*H*-imidazo[4,5-*c*]pyridin-2-yl)-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(5-Chloro-1*H*-benzoimidazol-2-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone; and

2-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-1*H*-benzoimidazole-5-carbonitrile.


Example 7: Synthesis of 2-(3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone



To a solution of 2-(3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone (210 mg, 0.497 mmol) in THF (5mL) is added a solution of oxone (611 mg, 0.994 mmol) in water (5 mL). The mixture is allowed to stir at room temperature for 4 h. Water (10 mL) is added and the mixture is extracted with EtOAc 3x. The organic extracts are washed with water, brine, dried (Na₂SO₄) and concentrated. The residue is re-dissolved in dichloromethane (3mL) and loaded on a 10g Biotage SNAP column and eluted with 100% EtOAc to provide 2-(3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone (102 mg).

The above sulfone (40 mg, 0.088mmol) is added to a 0.5-2mL Biotage microwave tube followed by methylamine (2M in THF, 0.22mL, 0.44mmol), Hunig's base (0.045 mL, 0.264 mmol) and isopropanol (1 mL). The reaction is carried out in a microwave oven at 175°C for 15 min. After cooling, the mixture is filtered and purified via prep- HPLC. Removal of the solvent gives the title compound as colorless foam (15mg).

Example 8: Synthesis of 4-amino-N-[2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-3-methoxy-benzamide



A solution of N-Boc-glycine (1.31 g, 7.33 mmol) and carbonyl diimidazole (1.21 mg, 7.33 mmol) in DMF (50 mL) is stirred at room temperature for 45 min. To the solution is added the biphenylpiperazine intermediate (2.28 g, 7.33 mmol) and triethylamine (5.13 mL, 36.63 mmol) and the solution stirred at room temperature for 24 h. The reaction is diluted with EtOAc (600 mL) and washed with sat. NH₄Cl (600 mL), NaHCO₃ (600 mL), then brine (100 mL). The aqueous layers are extracted again with more EtOAc (600 mL). The organics are dried (Na₂SO₄) and concentrated to give a solid. Trituration in diethyl ether and hexane followed by filtration afforded 2.32 g of the desired amide intermediate.


To a solution of the above amide intermediate (5.17 g, 12.94 mmol) in dichloromethane (200 mL) is added 4M HCl/dioxane (33 mL, 129.4 mmol) and the suspension stirred vigorously at room temperature for 20 h with periodic venting. The suspension is diluted with Et₂O (350 mL) and hexane (350 mL) and stirred 1h. The hygroscopic white solid is

filtered washing with Et₂O and hexane, then dried in vacuo under P₂O₅ to afford 4.84 g of the desired amine intermediate

A solution of Boc-4-amino-3-methoxybenzoic acid (56 mg, 0.204 mmol) and carbonyl diimidazole (33 mg, 0.204 mmol) in DMF (2 mL) is stirred at 50° C for 1 h. To the solution is added the above amine intermediate (75 mg, 0.204 mmol) and triethylamine (0.143 mL, 1.02 mmol) and the solution stirred at room temperature 24 h. The reaction is concentrated to a crude gum by nitrogen flow and used in the next reaction without further purification.

The crude coupling product from above is dissolved in trifluoroacetic acid (4 mL) and stirred at ambient temperature for 20 h. The reaction is concentrated in vacuo at 40°C to a residue. The reaction is diluted with DMF (2 mL) and water (1 mL) and purified by reverse-phase preparative HPLC (10-100% CH₃CN/H₂O) to give 24 mg of the title compound.


Example 9: Synthesis of isoquinoline-6-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide



A solution of isoquinoline-6-carboxylic acid (24 mg, 0.136 mmol) in excess thionyl chloride (3 mL) is stirred in a sealed vial at 85 °C for 24 h. The thionyl chloride is blown off by argon flow and to the solid residue is added the glycine amine intermediate (50 mg, 0.136 mmol) followed by DMF (1.5 mL) and triethylamine (0.250 mL, 1.79 mmol). The reaction is stirred at 50 °C for 18 h. The reaction is diluted with DMF (1 mL) and

water (1 mL), quenched with trifluoroacetic acid (0.2 mL) and purified by reverse-phase preparative HPLC (20-100% CH₃CN/H₂O) to give 51 mg of the title compound.

Example 10: Synthesis of 3,4-dimethoxy-N-(2-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-benzamide




A solution of 3,4-dimethoxybenzoic acid (2.80 g, 15.25 mmol) and carbonyl diimidazole (2.50 g, 15.25 mmol) in DMF (20 mL) is stirred at room temperature for 30 min. To this is added glycine *t*-butyl ester HCl (2.00 g, 15.25 mmol) at room temperature and Hunig's base (2.7 mL, 15.25 mmol) and the solution stirred at room temperature for 72 h. The reaction is diluted with EtOAc (500 mL) and washed with 0.05M HCl (500 mL), sat. NaHCO₃ (500 mL) then water (500 mL). The aqueous layers are extracted again with more EtOAc (500 mL), and the organics dried (Na₂SO₄) and concentrated to give the 3.92 g of the desired amide intermediate as a gum in 84% yield.

To a solution of the above amide intermediate (3.92 g, 13.14 mmol) in dichloromethane (100 mL) at 0° C is added trifluoroacetic acid (50 mL) and the reaction stirred at room temperature for 72 h with periodic venting. The reaction is concentrated at 40 °C and the resulting oil dissolved in dichloromethane and concentrated again. The oil is slurried in EtOAc (150 mL). The resulting precipitate is filtered, washing with EtOAc to give 2.72 g of the carboxylic acid intermediate in 85% yield

A solution of the above carboxylic acid intermediate (38 mg, 0.154 mmol) and carbonyl diimidazole (26 mg, 0.154 mmol) in DMF (2 mL) is stirred at room temperature for 1h. To the solution is added the substituted piperazine intermediate (50 mg, 0.154 mmol) and triethylamine (0.151 mL, 1.08 mmol) and the solution stirred at room temperature 18 h. The reaction is diluted with DMF (1 mL) and water (1 mL), quenched with trifluoroacetic acid (0.2 mL) and purified by reverse-phase preparative HPLC (15-100% CH₃CN/H₂O) to give 70 mg of the title compound.

Example 11: Synthesis of 1-((S)-4-Biphenyl-2-yl-2-methyl-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone



To a microwave tube is added 2-bromobiphenyl (1.0 g, 4.16 mmol), (S)-N-1-t-Boc-2-methylpiperazine (868 mg, 4.16 mmol), 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (X-Phos) (198 mg, 0.416 mmol), $\text{Pd}_2(\text{dba})_3$ (381 mg, 0.461 mmol), sodium tert-butoxide (400 mg, 4.16 mmol) and toluene (5 mL). The mixture is microwaved at 80° C for 60 min. The reaction is filtered through paper to remove solids, diluted with EtOAc (200 mL), and washed with water (200 mL) then brine (50 mL). The combined organics are dried (Na_2SO_4) and concentrated in vacuo at 45° C to give the desired *t*-Boc-piperazine intermediate as a crude brown gum. This is reacted without further purification.

To a solution of the above crude *t*-Boc-piperazine in dichloromethane (10 mL) is added TFA (50 mL) and the solution is stirred for 2 h. The reaction is concentrated in vacuo at 45° C to give a brown oil, which is diluted in EtOAc (200 mL) and washed with saturated NaHCO_3 (200 mL), water (200 mL) and brine (50 mL). The organic layer is

concentrated to a brown oil at 45 °C and purified by silica gel chromatography eluting with a gradient of 100% dichloromethane to 25% MeOH/dichloromethane. Fractions containing product are pooled and concentrated in vacuo to afford 128 mg of the desired piperazine intermediate as a brown solid.

(3,5-Dimethyl-1H-pyrazol-1-yl)acetic acid (21 mg, 0.125 mmol) is dissolved in dichloromethane (1 mL) and cooled to -5 °C. To this is added oxalyl chloride solution (2 M in dichloromethane, 0.063 mL, 0.125 mmol), followed by 0.02 mL DMF. The mixture is stirred for 2 h warming to room temperature. To the acid chloride solution is then added a solution of the above piperazine intermediate (30 mg, 0.119 mmol) and Hunig's base (0.104 mL, 0.595 mmol) in dichloromethane (1 mL) and the reaction is stirred at ambient temp for 20 h. The reaction is concentrated in vacuo at 40 °C to a residue. The residue is diluted with DMF (2 mL), water (1 mL) and TFA (0.2 mL) and purified by reverse-phase preparative HPLC (10-100% CH₃CN/H₂O) to give 21 mg of the title compound.

ASSESSMENT OF BIOLOGICAL PROPERTIES

Compounds are assessed for the ability to block the interaction of CXCR3 and IP-10 in a functional cellular assay measuring calcium flux in CXCR3 transfected cells.

Cyno-CHO cells, stably expressing recombinant CXCR3 and G-alpha-16 are grown in F12 medium (Mediatech #45000-360) supplemented with 10% (V/V) FBS (Mediatech #35-01500), 1% Geneticin (Invitrogen # 10131-027) and 0.2% Zeocin (Invitrogen #R250-05). The cells are spun down and re-suspended in growth media to a concentration of 4.8 E5 cells/mL. 25 microL of cell suspension is added to each well of a BD-384-well TC treated plate, providing 12,000 cells/well. The plate is incubated at 37 °C/5% CO₂ overnight. On the day of the assay, the plates are removed, the media is flicked out and 25 microL of Ca-4 dye in assay buffer (HBSS, 10 mM HEPES ph 7.4), containing 2mM probenacid is added to each well. The cell assay plates are then incubated at 37 °C/5% CO₂ for one hour.

Test compounds are dissolved in DMSO and diluted to 1.045 mM in DMSO. Just prior to assay, 2.75 microL of appropriately diluted test compound are added to each well of a 384 well plate containing 45 microL of HBSS buffer. After mixing, 5 microL of diluted compound are added to each well of the cell assay plate for a final assay concentration of 10 microM. The plate is incubated at room temperature for 15 min. 10 microL of IP-10 stock solution in HBSS (4X EC80 concentration) are added to each well of the cell assay plate except those cells reserved as blank wells containing buffer only. Intracellular calcium flux is recorded on the HAMAMTSU FDSS6000, using excitation at 480 nm and emission at 540 nm. Data are analyzed using Activity Base software.

In general, the preferred potency range (IC_{50}) of compounds in the above assay is between 1 nM to 3 μ M, and the most preferred potency range is 1 nM to 200 nM. The following table shows IC_{50} s for representative compounds of the invention in the above assay.

Table II

Compound	IC_{50} , nM
1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	9
1-{4-[2-(2-Piperidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	11
1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperidin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	11
1-{4-[2-(2-Methanesulfonyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	13
3-(2-Methylsulfanyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	4
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	5
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	8

3,4-Dimethoxy-N-(2-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-benzamide	68
5-(2-{4-[2-(Thiazol-2-ylamino)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	47
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methanesulfonyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	54
2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(3'-methoxy-biphenyl-2-yl)-piperazin-1-yl]-ethanone	56
1-{4-[2-(2-Ethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	9
5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyridine-2-carbonitrile	5
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	7
1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	5
1-(4-{2-[2-(2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	6
2-Indol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	44
2'-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-biphenyl-4-carbonitrile	100
1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	6
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	7
1-{4-[2-(6-Chloro-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	31
1-{3-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-3,8-diaza-bicyclo[3.2.1]oct-8-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	7
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1-methyl-1H-indol-6-yl)-phenyl]-piperazin-1-yl}-ethanone	19
1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-3,6-dihydro-2H-pyridin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	20
1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(pyridin-4-ylamino)-ethanone	27

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-piperidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	7
3-(2-Methoxy-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	7
1-{4-[2-Fluoro-6-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	27
2-Indazol-2-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	28
1-(4-{2-[2-((2R,6R)-2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	7
1-[4-(3'-Chloro-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	111
1-{4-[2-(2-Methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	115
3-(2-Methylamino-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	8
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(6-methoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-ethanone	58
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	67
1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-c]pyridin-1-yl-ethanone	9
1-{4-[2-(2-Pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	9
1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(thiazol-2-ylamino)-ethanone	35
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	10
1-(4-{2-[2-(4-Methyl-piperazin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	11
1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	7
1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	7
1-{4-[4-Chloro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	13

5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyrimidine-2-carbonitrile	13
2-Imidazo[4,5-b]pyridin-3-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	15
1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	15
5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	18
1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	4
5-(2-{4-[2-(5-Methyl-3-trifluoromethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	71
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(1H-indol-6-yl)-phenyl]-piperazin-1-yl}-ethanone	81
1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	6
1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	6
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-ethoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	18
4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-morpholin-4-yl-pyrimidin-5-yl)-benzonitrile	18
1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	8
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	9
3-(2-Morpholin-4-yl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile	8
1-{4-[2-(6-Methoxy-pyridin-3-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	34
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	34
2-Indazol-1-yl-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	35
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	46

1-(4-{2-[2-((2R,6S)-2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	10
4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methylsulfanyl-pyrimidin-5-yl)-benzonitrile	36
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-phenoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	39
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methyl-piperazin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	40
5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidine-2-carbonitrile	43
1-{4-[2-(2-[1,4]Oxazepan-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	9
1-(2-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-1,3-dihydro-benzoimidazol-2-one	10
5-(2-{4-[2-(2,4-Dimethyl-imidazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyridine-2-carbonitrile	47
4-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-3-(2-methoxy-pyrimidin-5-yl)-benzonitrile	150
1-{4-[4-Methoxy-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	9
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(pyridin-3-yloxy)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	56
1-[2-(4-Biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-1,3-dihydro-benzoimidazol-2-one	140
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone	8
2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(4'-methyl-biphenyl-2-yl)-piperazin-1-yl]-ethanone	70
1-(4-{2-[2-(3,5-Dimethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	14
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxy-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	7
1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-3-carboxylic acid amide	9
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(octahydro-isoquinolin-2-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	21

1-(4-{2-[2-(4-Dimethylamino-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	31
1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-4-carboxylic acid amide	6
1-(4-{2-[2-((R)-3-Dimethylamino-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	4
1-[5-(2-{4-[2-(3,5-Dimethyl-pyrazol-1-yl)-acetyl]-piperazin-1-yl}-phenyl)-pyrimidin-2-yl]-piperidine-4-carboxylic acid methylamide	9
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-fluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	10
1-(4-{2-[2-(3,3-Difluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	9
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-trifluoromethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	12
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-ethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	10
1-(4-{2-[2-(4,4-Difluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	22
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-fluoro-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	8
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methoxy-4-methyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	33
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(4-methanesulfonyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	6
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-(3-hydroxy-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	9
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-[1,4]oxazepan-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	11
2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-{2-[4-(1-hydroxy-1-methyl-ethyl)-piperidin-1-yl]-pyrimidin-5-yl}-phenyl)-piperazin-1-yl]-ethanone	12
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[2-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-ethanone	12
1-(4-{2-[2-(4,4-Dimethyl-piperidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	12
2-(3,5-Dimethyl-pyrazol-1-yl)-1-(4-{2-[(1S,4S)-2-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-	14

ethanone

1-(4-{2-[(1R,4S)-2-(2-Aza-bicyclo[2.2.1]hept-2-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	4
1-(4-{2-[2-(3,3-Difluoro-azetidin-1-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	7
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[4-fluoro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	5
1-{2-Methyl-4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	13
1-{3-Methyl-4-[2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	11
2-(5-Fluoro-pyrrolo[2,3-b]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	8
1-{4-[4-Chloro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	7
1-{4-[2-(2-Methoxy-pyridin-4-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	24
1-{4-[2-(2-Morpholin-4-yl-pyridin-4-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	16
2-(7-Chloro-pyrrolo[2,3-c]pyridin-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	16
2-(4-Chloro-pyrrolo[2,3-d]pyrimidin-7-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	48
1-{4-[4-Chloro-2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	8
2-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-2,3-dihydro-isoindol-1-one	17
1-{4-[4-Chloro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	12
1-(2-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-3-methyl-1,3-dihydro-benzoimidazol-2-one	11
1-{4-[4-Chloro-2-(2-methanesulfonyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	24
1-{4-[4-Chloro-2-(2-morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	11

1-{4-[4-Chloro-2-(2-methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	7
1-{4-[4-Chloro-2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone	6
1-{4-[4-Chloro-2-(2-methylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone	13
2-(4-Chloro-3,5-dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	10
2-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-2,3-dihydro-isoindol-1-one	48
2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[4-fluoro-2-(2-methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone	14
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-ylacetyl)piperazin-1-yl]benzamide	6
1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one	7
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone	7
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	7
5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide	7
1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	7
2-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-2,3-dihydro-1H-isoindol-1-one	7
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}benzamide	7
2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone	7
1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	8
1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-	8

2H-benzimidazol-2-one

2-[(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl]-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide	8
1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	8
1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	8
1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	8
1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	8
1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	8
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)ethanone	8
1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	8
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl]benzamide	8
1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	8
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	8
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl]benzamide	8
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide	9
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl]benzamide	9
1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	9
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	9

2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one	9
1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	9
1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one	9
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-yl)acetyl]piperazin-1-yl]benzamide	9
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	9
2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	9
2-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	9
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone	9
1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	9
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide	9
1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	10
2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	10
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]ethanone	10
2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one	10
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide	10
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	10

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	10
1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	11
1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one	11
5-fluoro-2-[(3S)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide	11
1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)piperidin-2-one	11
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	11
1-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	11
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone	11
1-{2-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one	11
N-{2-[(2S)-4-{4-fluoro-2-[(2-methoxypyridin-4-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	11
5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl}benzamide	11
1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	11
5-fluoro-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	12
2-{(3S)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl}-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide	12
N-{2-[(2R)-4-{4-fluoro-2-[(2-methoxypyridin-4-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	12
2-{(3S)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl}-5-fluoro-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	12

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	13
1-(2-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one	13
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl]benzamide	13
1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	13
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)ethanone	13
1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	13
1-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	14
1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	14
1-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	14
5-fluoro-2-[(3S)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	14
2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone	14
1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone	14
1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}piperidin-2-one	14
1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	14
5-fluoro-2-[(3S)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	15
5-fluoro-2-[(3S)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	15
1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}piperidin-2-one	15

2-[(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl]-5-fluoro-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	15
5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	15
2-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	15
1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	15
5-fluoro-2-[(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	15
1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	16
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	16
5-fluoro-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	17
5-fluoro-2-[(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	17
5-fluoro-2-[(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	18
5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]benzamide	18
1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]piperidin-2-one	19
5-fluoro-2-[(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	19
1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	20
5-fluoro-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	20
1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	20

5-fluoro-2-((3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl)-N-(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)benzamide	20
N-{2-[(2S)-4-{4-fluoro-2-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	21
2-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	23
1-[2-(4-{4-fluoro-2-[2-(pyrrolidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	23
1-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	24
2-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	26
1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	27
1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	30
1-[2-(4-{4-fluoro-2-[2-(morpholin-4-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	30
1-{2-[2-methyl-4-(2-[2-(4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl]phenyl)piperazin-1-yl]-2-oxoethyl}piperidin-2-one	34
N-{2-[(2R)-4-{4-fluoro-2-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)carbamoyl]phenyl}-2-methylpiperazin-1-yl]-2-oxoethyl}-N,5-dimethyl-1,3-oxazole-4-carboxamide	35
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	40
1-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	43
1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	47
1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	48
1-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	53
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)ethanone	56

1-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methyl)piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	60
1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one	63
1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone	67
1-[2-(4-{4-fluoro-2-[2-(4-methylpiperazin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	94
2-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	101
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)ethanone	157
1-(2-{4-[4-fluoro-2-(2-methoxypyrimidin-5-yl)phenyl]piperazin-1-yl}-2-oxoethyl)piperidin-2-one	174
2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)ethanone	189
2-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	245
2-[2-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	249
2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	278
2-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one	293
1-[2-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}piperazin-1-yl)-2-oxoethyl]piperidin-2-one	310
2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-(2-methyl-4-{2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone	434
1-(4-{4-fluoro-2-[2-(methylsulfonyl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone	488

METHODS OF USE

The compounds of the invention are effective antagonists of the interaction of CXCR3 and its ligands and thus inhibit CXCR3 activation. Therefore, in one embodiment of the invention, there is provided methods of treating CXCR3-mediated disorders using compounds of the invention. In another embodiment, there is provided methods of treating inflammatory, autoimmune and cardiovascular diseases using compounds of the invention.

Without wishing to be bound by theory, by inhibiting the activity of CXCR3 the compounds of the invention block the migration of T-cells and other leukocytes that express CXCR3. Thus, the inhibition of CXCR3 activity is an attractive means for preventing and treating a variety of autoimmune and immunological diseases exacerbated by the influx of these leukocytes. These include multiple sclerosis, rheumatoid arthritis, psoriasis inflammatory bowel disease, chronic obstructive pulmonary disease (COPD) and kidney disease. Furthermore, a genetic deletion study and a study in LDL receptor KO mice with a CXCR3 antagonist have both shown that inhibition of CXCR3 activity attenuates atherosclerotic lesion formation. Thus inhibition of CXCR3 activity is also an attractive means for treating and preventing atherosclerosis and secondary atherothrombotic events such as myocardial infarction and stroke.

For treatment of the above-described diseases and conditions, a therapeutically effective dose will generally be in the range of approximately 0.01 mg to about 100 mg/kg of body weight per dosage of a compound of the invention; preferably, from about 0.1 mg to about 20 mg/kg of body weight per dosage. For example, for administration to a 70 kg person, the dosage range would be approximately 0.7 mg to about 7000 mg per dosage of a compound of the invention, preferably from about 7.0 mg to about 1400 mg per dosage. Some degree of routine dose optimization may be required to determine an optimal dosing level and pattern. The active ingredient may be administered from 1 to 6 times a day.

General Administration and Pharmaceutical Compositions

When used as pharmaceuticals, the compounds of the invention are typically administered in the form of a pharmaceutical composition. Such compositions can be prepared using procedures well known in the pharmaceutical art and comprise at least one compound of the invention. The compounds of the invention may also be administered alone or in combination with adjuvants that enhance stability of the compounds of the invention, facilitate administration of pharmaceutical compositions containing them in certain embodiments, provide increased dissolution or dispersion, increased antagonist activity, provide adjunct therapy, and the like. The compounds according to the invention may be used on their own or in conjunction with other active substances according to the invention, optionally also in conjunction with other pharmacologically active substances. In general, the compounds of this invention are administered in a therapeutically or pharmaceutically effective amount, but may be administered in lower amounts for diagnostic or other purposes.

Administration of the compounds of the invention, in pure form or in an appropriate pharmaceutical composition, can be carried out using any of the accepted modes of administration of pharmaceutical compositions. Thus, administration can be, for example, orally, buccally (e.g., sublingually), nasally, parenterally, topically, transdermally, vaginally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as, for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages. The pharmaceutical compositions will generally include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, vehicles, or combinations thereof. Such pharmaceutically acceptable excipients, carriers, or additives as well as methods of making pharmaceutical compositions for various modes of administration are well-known to those of skill in the art. The state of the art is evidenced, e.g., by *Remington: The Science and Practice of Pharmacy*, 20th Edition, A. Gennaro (ed.), Lippincott Williams & Wilkins, 2000; *Handbook of Pharmaceutical*


Additives, Michael & Irene Ash (eds.), Gower, 1995; *Handbook of Pharmaceutical Excipients*, A.H. Kibbe (ed.), American Pharmaceutical Ass'n, 2000; H.C. Ansel and N.G. Popovich, *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 5th ed., Lea and Febiger, 1990; each of which is incorporated herein by reference in their entireties to better describe the state of the art.

As one of skill in the art would expect, the forms of the compounds of the invention utilized in a particular pharmaceutical formulation will be selected (e.g., salts) that possess suitable physical characteristics (e.g., water solubility) that are required for the formulation to be efficacious.

Claims

What is claimed is:

1. A compound of formula (I):



(I)

wherein:

A is C or N;

B is C or N;

D is C or N;

X is -NH-, -NHC(O)-, -N(CH₃)C(O)- or absent;

R¹ is H, -CN, halogen, -CF₃, -OCF₃, C₁₋₃alkyl, C₁₋₃alkoxy, -S(O)_nCH₃, amino, mono- or dimethylamino, -NHC(O)C₁₋₃alkyl, -NO₂, -C(O)NH₂, -C(O)NHC₁₋₃alkyl or -C(O)C₁₋₃alkyl;

R² is aryl or heteroaryl, each optionally substituted with one to three R⁶;

R³ is heteroaryl, heterocyclyl, aryl or C₃₋₁₀cycloalkyl each optionally substituted with one to three R⁷;

R⁴ and R⁵ are each independently selected from H and C₁₋₂alkyl; or R⁴ and R⁵ may join to form an ethyl bridge;

each R⁶ is independently -OH, oxo, hydroxyC₁₋₆alkyl, halogen, -(CH₂)_m-CN, nitro, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, C₁₋₆ alkoxy, phenoxy, heteroaryloxy, C₁₋₆ alkoxy carbonyl, carboxyl, -C(O)C₁₋₆alkyl, -(CH₂)_m-NR₈R₉, -S(O)_nC₁₋₆alkyl, -NHS(O)_nC₁₋₆alkyl, -NR⁸C(O)C₁₋₆alkyl, -S(O)₂NR₈R₉, -C(O)NR₈R₉, heteroaryl, heterocyclyl, phenyl or benzyl, wherein each alkyl, alkenyl, alkynyl or alkoxy of said R⁶ is optionally partially or fully halogenated and each heterocyclyl, heteroaryl, phenyl or benzyl of said R⁶ is optionally substituted with one to three C₁₋₆

alkyl, C₁₋₆alkoxy(CH₂)_m, halogen, -CN, -CF₃, C₁₋₆ acyl, -NR⁸R⁹, C(O)NR⁸R⁹, -OH, hydroxyC₁₋₆alkyl, pyrrolidinyl or -S(O)_nC₁₋₆alkyl;

each R⁷ is independently C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, C₁₋₆ alkoxy, halogen, oxo, -CN, carboxy, -(CH₂)_m-NR₈R₉, phenyl or heteroaryl, wherein each alkyl, alkenyl, alkynyl or alkoxy of said R⁷ is optionally partially or fully halogenated and wherein two R⁷ on adjacent carbon atoms of R³ may join to form a three to seven member ring fused to R³;

R⁸ and R⁹ are each independently hydrogen, C₁₋₆ alkyl, C₁₋₆ acyl, C₃₋₁₀ cycloalkyl, hydroxyC₁₋₆ alkyl, C₁₋₆ alkylC₁₋₆ alkoxy, C₁₋₆ alkylsulfonyl or C₁₋₆ alkoxycarbonyl;

m is an integer from 0 to 3;

n is an integer from 0 to 2; and

----- is a single bond if A is N and may be a single or double bond if A is C; or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein R³ is aryl or heteroaryl, or a pharmaceutically acceptable salt thereof.
3. The compound of claim 1, wherein R² is aryl, and R³ is aryl or heteroaryl, or a pharmaceutically acceptable salt thereof.
4. The compound of claim 1, wherein R² is heteroaryl, and R³ is aryl or heteroaryl, or a pharmaceutically acceptable salt thereof.
5. The compound of claim 1, wherein A is N, and D is C, or a pharmaceutically acceptable salt thereof.
6. The compound of claim 1, wherein B is C, X is absent, or a pharmaceutically acceptable salt thereof.
7. The compound of claim 1, wherein:
B is C, D is C;

R^1 is H, -CN, -F, -Cl, -CF₃, -CH₃, -OCH₃, -C(O)NHCH₃ or -S(O)₂CH₃;

R^2 is benzoimidazolyl, benzooxazolyl, benzo[b]thiophenyl, dibenzofuranyl, dibenzothiophenyl, furanyl, imidazolyl, 1H-imidazo[4,5-c]pyridinyl, indolyl, isoindolyl, isoquinolinyl, isoxazolyl, oxazolyl, phenyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, quinolinyl, thiadiazolyl, thiazolyl, thienyl or triazinyl, wherein each of the foregoing is optionally substituted by one to three R^6 ;

R^3 is benzoimidazolyl, benzo[d]isothiazolyl, benzo[b]thiophenyl, benzotriazolyl, furanyl, imidazo[4,5-b]pyridinyl, imidazolyl, indazolyl, indolyl, isoindolyl, isoquinolinyl, isothiazolyl, phenyl, phthalazinyl, pyrazolyl, pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-c]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl, pyrrolo[2,3-d]pyrimidinyl, pyrrolyl, quinazolinyl, thiadiazolyl or thiazolyl wherein each of the foregoing is optionally substituted with one to three R^7 ;

R^4 and R^5 are H;

each R^6 is independently -OH, oxo, hydroxyC₁₋₆alkyl, halogen, -(CH₂)_m-CN, nitro, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, C₁₋₆ alkoxy, phenoxy, pyridyloxy, C₁₋₆ alkoxy carbonyl, carboxyl, -C(O)C₁₋₆alkyl, -(CH₂)_m-NR₈R₉, -S(O)_nC₁₋₆alkyl, -NHS(O)₂C₁₋₆alkyl, -NHC(O)C₁₋₆alkyl, -S(O)₂NR₈R₉, -C(O)NR₈R₉, thienyl, morpholinyl, pyrrolidinyl, piperidinyl, [1,3]-oxazepan-1-yl, piperazinyl, azabicyclo[2.2.1]heptyl, 2-oxa-5-azabicyclo[2.2.1]heptyl, octahydroisoquinolinyl, [1,4]-oxazepanyl, azetidiny phenyl or benzyl, wherein each alkyl, alkenyl or alkynyl of said R^6 is optionally partially or fully halogenated and each heterocyclyl or phenyl of said R^6 is optionally substituted with one to three CH₃, -OCH₃, halogen, -CN, -CF₃, -C(O)CH₃, NR⁸R⁹, -C(O)NH₂, -C(O)NHCH₃, -OH, -CH₂OCH₃, -C(OH)(CH₃)CH₃, -SCH₃, pyrrolinyl or -S(O)₂CH₃;

each R^7 is independently C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, C₁₋₆ alkoxy, halogen, oxo, -CN, carboxy, -(CH₂)_m-NR₈R₉, phenyl or pyridyl, wherein each alkyl, alkenyl or alkynyl of said R^7 is optionally partially or fully halogenated;

R^8 and R^9 are each independently hydrogen, C₁₋₆ alkyl, C₁₋₆ acyl or C₃₋₁₀ cycloalkyl;

or a pharmaceutically acceptable salt thereof.

8. The compound of claim 1, wherein:

A is N;

X is -NH- or absent;

R¹ is H, -CN, -F, -Cl, -CF₃, -CH₃ or -OCH₃;

R² is benzoimidazolyl, benzooxazolyl, imidazolyl, 1H-imidazo[4,5-c]pyridinyl, indolyl, isoindolyl, isoquinolinyl, isoxazolyl, oxazolyl, phenyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridinyl, pyrimidinyl, quinolinyl, thiadiazolyl, thiazolyl or thienyl, wherein each of the foregoing is optionally substituted by one to three R⁶;

R³ is benzoimidazolyl, benzotriazolyl, imidazolyl, imidazo[4,5-b]pyridinyl, indazolyl, indolyl, isoindolyl, isothiazolyl, pyrazolyl, pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl, pyrrolo[2,3-d]pyrimidinyl, pyrrolyl, thiadiazolyl or thiazolyl wherein each of the foregoing is optionally substituted with one to three R⁷;

each R⁶ is independently -CH₂OH, -Cl, -F, -CN, oxo, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₂ alkoxy, phenoxy, pyridyloxy, -C(O)CH₃, -(CH₂)_m-NR₈R₉, -S(O)_nCH₃, -NHS(O)₂CH₃, -NH(CO)CH₃, -S(O)₂NR₈R₉, -C(O)NR₈R₉, thienyl, morpholinyl, pyrrolidinyl, piperidinyl, [1,3]-oxazepan-1-yl, piperazinyl, phenyl or benzyl, wherein each alkyl of said R⁶ is optionally partially or fully halogenated and each heterocyclyl or phenyl of said R⁶ is optionally substituted with one to three CH₃, -OCH₃, halogen, -CN, -CF₃, N(CH₃)₂, -C(O)NH₂, -C(O)NHCH₃, -OH, -CH₂OCH₃, -C(OH)(CH₃)CH₃, pyrrolidinyl or S(O)₂CH₃;

each R⁷ is independently C₁₋₃ alkyl, -OCH₃, CF₃, oxo, -CN, -Cl or -F;

R⁸ and R⁹ are each independently hydrogen, C₁₋₆ alkyl, C₁₋₃ acyl or C₃₋₆ cycloalkyl;

or a pharmaceutically acceptable salt thereof.

9. The compound of claim 1 wherein:

X is absent;

R³ is benzoimidazolyl, benzotriazolyl, imidazolyl, imidazo[4,5-b]pyridinyl, indazolyl, indolyl, isoindolyl, pyrazolyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-b]pyridinyl,

pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl or pyrrolyl, wherein each of the foregoing is optionally substituted with one to three R⁷;

or a pharmaceutically acceptable salt thereof.

10. A compound selected from:

3-(2-Methylsulfanyl-pyrimidin-5-yl)-4-[4-(2-pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-benzonitrile;

1-{4-[2-(2-Methylsulfanyl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

4-Amino-N-[2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-3-methoxy-benzamide;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-[4-(2-naphthalen-1-yl-phenyl)-piperazin-1-yl]-ethanone;

1-Methyl-1H-imidazole-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

5-{2-[4-(2-Pyrrolo[2,3-b]pyridin-1-yl-acetyl)-piperazin-1-yl]-phenyl}-pyridine-2-carbonitrile;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-methylsulfanyl-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-ethanone;

1-{4-[2-(2-Methylamino-pyrimidin-5-yl)-4-trifluoromethyl-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-(3,4-dimethoxy-benzylamino)-ethanone

1-(4-Biphenyl-2-yl-piperazin-1-yl)-2-piperidin-1-yl-ethanone;

Isoquinoline-6-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

Thiazole-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

1-(4-{2-[2-(2,6-Dimethyl-morpholin-4-yl)-pyrimidin-5-yl]-phenyl}-piperazin-1-yl)-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

1-{4-[2-(2-Methoxy-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Dimethylamino-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

1-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-pyrrolo[2,3-b]pyridin-1-yl-ethanone;

2-(3,5-Dimethyl-pyrazol-1-yl)-1-{4-[2-(2-pyrrolidin-1-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-ethanone;

4,5-Dimethyl-furan-2-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

1-(2-{4-[2-(2-Morpholin-4-yl-pyrimidin-5-yl)-phenyl]-piperazin-1-yl}-2-oxo-ethyl)-imidazolidin-2-one;

1-[4-(4'-Dimethylaminomethyl-biphenyl-2-yl)-piperazin-1-yl]-2-(3,5-dimethyl-pyrazol-1-yl)-ethanone;

Isoquinoline-7-carboxylic acid [2-(4-biphenyl-2-yl-piperazin-1-yl)-2-oxo-ethyl]-amide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

5-fluoro-2-[(3R)-4-(3H-imidazo[4,5-b]pyridin-3-ylacetyl)-3-methylpiperazin-1-yl]-N-(2-methoxypyridin-4-yl)benzamide;

1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-2,3-dihydro-1H-isoindol-1-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(1-oxo-1,3-dihydro-2H-isoindol-2-yl)acetyl]piperazin-1-yl}benzamide;

2-(3H-imidazo[4,5-b]pyridin-3-yl)-1-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}ethanone;

1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-(2-{2-methyl-4-[2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2-oxoethyl)-1,3-dihydro-2H-benzimidazol-2-one;

2-[(3R)-4-[(3,5-dimethyl-1H-pyrazol-1-yl)acetyl]-3-methylpiperazin-1-yl]-5-fluoro-N-(2-methoxypyridin-4-yl)benzamide;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

1-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3S)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl]benzamide;

1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl]benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-c]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-[N-methyl-N-(2-methylpropanoyl)glycyl]piperazin-1-yl]benzamide;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one;

1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-[(3R)-3-methyl-4-(1H-pyrrolo[2,3-b]pyridin-1-ylacetyl)piperazin-1-yl]benzamide;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)ethanone;

2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-[2-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[2-methyl-4-(2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]ethanone;

1-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-benzimidazol-2-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3S)-3-methyl-4-[(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)acetyl]piperazin-1-yl}benzamide;

1-(4-{4-fluoro-2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}piperazin-1-yl)-2-(1H-pyrrolo[2,3-c]pyridin-1-yl)ethanone;

2-[2-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]phenyl}-2-methylpiperazin-1-yl)-2-oxoethyl]-2,3-dihydro-1H-isoindol-1-one;

2-(3,5-dimethyl-1H-pyrazol-1-yl)-1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]ethanone;

2-{2-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)-2-methylpiperazin-1-yl]-2-oxoethyl}-2,3-dihydro-1H-isoindol-1-one;

5-fluoro-N-(2-methoxypyridin-4-yl)-2-{(3R)-3-methyl-4-[(2-oxopiperidin-1-yl)acetyl]piperazin-1-yl}benzamide;

1-(4-{2-[2-(4-hydroxypiperidin-1-yl)pyrimidin-5-yl]-4-(trifluoromethyl)phenyl}-2-methylpiperazin-1-yl)-2-(1H-pyrrolo[2,3-b]pyridin-1-yl)ethanone;

1-[4-(4-fluoro-2-{2-[4-(methylsulfonyl)piperidin-1-yl]pyrimidin-5-yl}phenyl)piperazin-1-yl]-2-(3H-imidazo[4,5-b]pyridin-3-yl)ethanone

or a pharmaceutically acceptable salt thereof.

11. A method of treating a CXCR3-mediated disorder comprising administering a therapeutically effective amount of a compound of claim 1, or a pharmaceutically acceptable salt thereof, to patient in need thereof.

12. A method of treating an inflammatory, autoimmune or cardiovascular disease comprising administering a therapeutically effective amount of a compound of claim 1, or a pharmaceutically acceptable salt thereof, to patient in need thereof.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/032347

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D209/48	C07D213/26	C07D213/56	C07D213/65	C07D213/74
C07D217/00	C07D231/10	C07D233/90	C07D235/04	C07D239/34
C07D239/42	C07D277/30	C07D307/68	C07D317/68	C07D319/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2007/127635 A2 (SQUIBB BRISTOL MYERS CO [US]; WANG TAO [US]; KADOW JOHN F [US]; ZHANG) 8 November 2007 (2007-11-08) page 184, column 185; compounds G-3a, G-3b -----	1-12
A	PEASE JAMES E ET AL: "Chemokine receptor antagonists: part 2." EXPERT OPINION ON THERAPEUTIC PATENTS FEB 2009 LNKD- PUBMED:19441918, vol. 19, no. 2, February 2009 (2009-02), pages 199-221, XP002591790 ISSN: 1744-7674 page 207 - page 212; figure 3 -----	1-12



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

14 July 2010

Date of mailing of the international search report

23/07/2010

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Skulj, Primoz

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2010/032347

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007127635 A2	08-11-2007	AR 060658 A1	02-07-2008
		AU 2007242989 A1	08-11-2007
		CA 2650382 A1	08-11-2007
		EA 200802203 A1	28-04-2009
		EP 2010525 A2	07-01-2009
		JP 2009535352 T	01-10-2009
		KR 20090005393 A	13-01-2009
		US 2007249579 A1	25-10-2007
		US 2008139572 A1	12-06-2008
