



(86) Date de dépôt PCT/PCT Filing Date: 2011/02/18
(87) Date publication PCT/PCT Publication Date: 2011/08/25
(85) Entrée phase nationale/National Entry: 2012/08/17
(86) N° demande PCT/PCT Application No.: US 2011/025459
(87) N° publication PCT/PCT Publication No.: 2011/103446
(30) Priorités/Priorities: 2010/02/18 (US61/338,464);
2010/02/18 (US61/338,489); 2010/07/30 (US61/369,562)

(51) Cl.Int./Int.Cl. *C12N 5/073* (2010.01),
A61F 2/08 (2006.01), *A61K 35/50* (2006.01),
A61P 17/02 (2006.01), *C12N 5/07* (2010.01)

(71) Demandeur/Applicant:
OSIRIS THERAPEUTICS, INC., US

(72) Inventeurs/Inventors:
TOM, SAMSON, US;
DANILKOVITCH, ALLA, US;
YOO, DANA, US;
JANSEN, TIMOTHY, US;
KUANG, JIN-QIANG, US;
MARCONI, JENNIFER MICHELLE, US

(74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(54) Titre : PRODUITS DE MEMBRANE CHORIONIQUE IMMUNOCOMPATIBLE
(54) Title: IMMUNOCOMPATIBLE CHORIONIC MEMBRANE PRODUCTS

(57) **Abrégé/Abstract:**

Provided herein is a placental product comprising an immunocompatible chorionic membrane. Such placental products can be cryopreserved and contain viable therapeutic cells after thawing. The placental product of the present invention is useful in treating a patient with a tissue injury (e.g. wound or burn) by applying the placental product to the injury. Similar application is useful with ligament and tendon repair and for engraftment procedures such as bone engraftment.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
25 August 2011 (25.08.2011)

PCT

(10) International Publication Number
WO 2011/103446 A1(51) International Patent Classification:
C12N 5/073 (2010.01) *A61K 35/50* (2006.01)(21) International Application Number:
PCT/US2011/025459(22) International Filing Date:
18 February 2011 (18.02.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/338,489 18 February 2010 (18.02.2010) US
61/338,464 18 February 2010 (18.02.2010) US
61/369,562 30 July 2010 (30.07.2010) US(71) Applicant (for all designated States except US): **OSIRIS THERAPEUTICS, INC.** [US/US]; 7015 Albert Einstein Dr., Columbia, Maryland 21046 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **TOM, Samson** [US/US]; 5200 Leavers Court, Baltimore, Maryland 21237 (US). **DANILKOVITCH, Alla** [RU/US]; 9489 Good Lion Road, Columbia, Maryland 21045 (US). **YOO, Dana** [US/US]; 7154 Harlan Lane, Sykesville, Maryland 21784 (US). **JANSEN, Timothy** [US/US]; 1420 Light Street, Apt. 1, Baltimore, Maryland 21230 (US). **KUANG, Jin-Qiang** [CA/US]; 10679 Hillingdon Road, Woodstock, Maryland 21163 (US). **MARCONI, Jennifer Michelle** [US/US]; 308 Congressional Court, Glen Burnie, Maryland 21061 (US).(74) Agents: **BABEL, Angeline** et al.; c/o McAndrews, Held & Malloy Ltd., 500 West Madison Street, 34th Floor, Chicago, Illinois 60661 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: IMMUNOCOMPATIBLE CHORIONIC MEMBRANE PRODUCTS

(57) Abstract: Provided herein is a placental product comprising an immunocompatible chorionic membrane. Such placental products can be cryopreserved and contain viable therapeutic cells after thawing. The placental product of the present invention is useful in treating a patient with a tissue injury (e.g. wound or burn) by applying the placental product to the injury. Similar application is useful with ligament and tendon repair and for engraftment procedures such as bone engraftment.



WO 2011/103446 A1

IMMUNOCOMPATIBLE CHORIONIC MEMBRANE PRODUCTS

RELATED APPLICATIONS

001 This application claims priority to:

002 U.S. Provisional Applications Ser. No, 61/338,464 entitled "Selectively Immunodepleted Chorionic Membranes" , filed on February 18, 2010 bearing Docket No. 22924US01,

003 U.S. Provisional Applications Ser. No, 61/338,489 entitled "Selectively Immunodepleted Amniotic Membranes" , filed on February 18, 2010 bearing Docket No. 22925US01, and

004 U.S. Provisional Applications Ser. No, 61/369,562 entitled "Therapeutic Products Comprising Vitalized Placental Dispersions" filed on July 30, 2010 bearing Docket No 23498US01, the contents of which are hereby incorporated by reference in their entireties.

005 This application is being co-filed on February 18, 2011 with, and incorporates by reference, applications entitled:

006 "Methods of Manufacture of Immunocompatible Chorionic Membrane Products",

007 "Immunocompatible Amniotic Membrane Products",

008 "Methods of Manufacture of Immunocompatible Amniotic Membrane Products",

009 "Therapeutic Products Comprising Vitalized Placental Dispersions" , and

0010 "Methods of Manufacture of Therapeutic Products Comprising Vitalized Placental Dispersions"

FIELD OF THE INVENTION

0011 The present technology relates to products to facilitate wound healing such as placenta membrane-derived products and biologic skin substitutes. The present technology relates to products to protect injured or damaged tissue, or as a covering to prevent adhesions, to exclude bacteria, to inhibit bacterial activity, or to promote healing or growth of tissue. The field also relates to methods of manufacturing and methods of use of such membrane-derived products.

BACKGROUND OF THE INVENTION

0012 Fresh or decellularized placental membranes have been used topically in surgical applications since at least 1910 when Johns Hopkins Hospital reported the use of placental membrane for dermal applications. Subsequently unseparated amnion and chorion were used as skin substitutes to treat burned or ulcerated surfaces. During the 1950's and 60's Troensegaard-Hansen applied boiled amniotic membranes to chronic leg ulcers.

0013 The human chorionic membrane (CM) is one of the membranes that exists during pregnancy between the developing fetus and mother. It is formed by extraembryonic mesoderm and the two layers of trophoblast and surrounds the embryo and other membranes. The chorionic villi emerge from the chorion, invade the endometrium, and allow transfer of nutrients from maternal blood to fetal blood.

0014 Both fresh and frozen CMs have been used for wound healing therapy. When fresh CM is used, there is increased risk of disease transmission. According to published reports, fresh placental tissue, for example, chorionic tissue exhibits cell viability of 100%, however within 28 days of storage above 0°C diminished cell viability to 15 to 35%. Freezing over a time of 3 weeks reduced cell viability to 13 to 18%, regardless of the temperature or medium. As the CM is believed to be immunogenic, it has not been used in commercial wound healing products.

0015 Two placental tissue graft products containing living cells, Apligraf and Dermagraft, are currently commercially available. Both Apligraf and Dermagraft comprise *ex vivo* cultured cells. Neither Apligraf nor Dermagraft comprise stem cells. Furthermore, neither Apligraf nor Dermagraft comprise Insulin-like Growth Factor Binding Protein-1 (IGFBP-1) and adiponectin, which are key factors in the natural wound healing process. In addition, neither Apligraf nor Dermagraft exhibit a protease-to-protease inhibitor ratio favorable for wound healing. As wound healing is a multi-factorial biological process, many factors are needed to properly treat a wound; products having non-native cellular populations are less capable of healing wounds relative to a product having an optimal population of cells representing the native array. It would represent an advance in the art to provide a chorion-derived biologic skin substitute comprising a population of cells representing the native array of factors, including, for example, growth factors and cytokines.

0016 Apligraf is a living, bi-layered skin substitute manufactured using neonatal foreskin keratinocytes and fibroblasts with bovine Type I collagen. As used in this application, Apligraf refers to the product available for commercial sale in November 2009.

0017 Dermagraft is cryopreserved human fibroblasts derived from newborn foreskin tissue seeded on extracellular matrix. According to its product literature, Dermagraft requires three washing steps before use which limits the practical implementation of Dermagraft as a skin substitute relative to products that require less than three washing steps. As used in this application, Dermagraft refers to the product available for commercial sale in November 2009.

0018 Engineered skin substitutes such as Apligraf and Dermagraft do not provide the best potential for wound healing because they comprise sub-optimal cellular compositions and therefore do not provide proper wound healing. For example, neither Apligraf nor Dermagraft comprises stem cells and, as a result, the ratio between different factors secreted by cells does not enable efficient wound healing. Additionally, some factors that are important for wound healing, including EGF, IGFBP1, and adiponectin are absent from both Apligraf and Dermagraft. Additionally, some factors, including MMPs and TIMPs, are present in proportions that differ greatly from the proportions found in the natural wound healing process; this difference significantly alters, among other things, upstream inflammatory cytokine pathways which in turn allows for sub-optimal micro-environments at the wound site. The present inventors have identified a need for the development of chorionic membrane products that more closely resemble natural tissue.

0019 Paquet-Fifield et al. report that mesenchymal stem cells and fibroblasts are important for wound healing (J Clin Invest, 2009, 119: 2795). No product has yet been described that comprise mesenchymal stem cells and fibroblasts.

0020 Both MMPs and TIMPs are among the factors that are important for wound healing. However, expression of these proteins must be highly regulated and coordinated. Excess of MMPs versus TIMPs is a marker of poor chronic wound healing (Liu et al, Diabetes Care, 2009, 32: 117; Mwaura et al, Eur J Vasc Endovasc Surg, 2006, 31: 306; Trengove et al, Wound Rep Reg, 1999, 7: 442; Vaalamo et al, Hum Pathol, 1999, 30: 795).

0021 α 2-macroglobulin is known as a plasma protein that inactivates proteinases from all 4 mechanistic classes: serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases (Borth et al., FASEB J, 1992,6: 3345; Baker et al., J Cell Sci, 2002, 115:3719). Another important function of this protein is to serve as a reservoir for cytokines and growth factors, examples of which include TGF, PDGF, and FGF (Asplin et al, Blood, 2001, 97: 3450; Huang et al, J Biol Chem, 1988; 263: 1535). In chronic wounds like diabetic ulcers or venous ulcers, the presence of high amount of proteases leads to rapid degradation of growth factors and delays in wound healing. Thus, a placental membrane skin substitute comprising α 2-macroglobulin would constitute an advance in the art.

0022 bFGF modulates a variety of cellular processes including angiogenesis, tissue repair, and wound healing (Presta et al., 2005, Reuss et al., 2003, and Su et al., 2008). In wound healing models, bFGF has been shown to increase wound closure and enhance vessel formation at the site of the wound (Greenhalgh et al., 1990).

0023 An *in vitro* cell migration assay is important for assessing the wound healing potential of a skin substitute. The process of wound healing is highly complex and involves a series of structured events controlled by growth factors (Goldman, Adv Skin Wound Care, 2004, 1:24). These events include increased vascularization, infiltration by inflammatory immune cells, and increases in cell proliferation. The beginning stages of wound healing revolve around the ability of individual cells to polarize towards the wound and migrate into the wounded area in order to close the wound area and rebuild the surrounding tissue. Keratinocytes are the primary cell type of the epithelial layer. Upon proper stimulation, they are implicated in the wound healing process (Pastar et al, 2008 and Bannasch et al., 2000). Specifically, they proliferate and migrate into the wound area to promote healing. An assay capable of evaluating the wound healing potential of skin substitutes by examining the correlation between cell migration and wound healing would represent an advance in the art.

SUMMARY OF THE INVENTION

0024 The present invention provides a pharmaceutically acceptable placental product.

0025 A placental product according to the present invention comprises an immunocompatible chorionic membrane in a cryopreservation medium (optionally cryopreserved) and viable native therapeutic cells and native therapeutic factors.

0026 In some embodiments, the placental product further comprises an amniotic membrane that is selectively devitalized.

0027 There is now provided a placental product that is selectively depleted of substantially all immunogenic cells.

0028 There is now provided a placental product that does not contain ex vivo cultured cells.

0029 There is now provided a placental product that comprises at least one of Epidermal Growth Factor, IGFBP1, and Adiponectin.

0030 Optionally, the therapeutic factors include one or more of IGFBP1, adiponectin, α 2-macroglobulin, bFGF, and EGF. Optionally, the therapeutic factors include MMP-9 and TIMP1, wherein the ratio of MMP-9:TIMP1 is from about 7 to about 10. Optionally, the therapeutic factors include IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9, and TIMP1. Optionally, the therapeutic factors include IGFBP1, adiponectin, α 2-macroglobulin, bFGF, MMP-9, and TIMP1, wherein the ratio of MMP-9:TIMP1 is from about 7 to about 10. Optionally, the therapeutic factor is present in a substantial amount in comparison to the equivalent unprocessed human placental membrane. Optionally, each placental product embodiment optionally is devoid of *ex-vivo expanded* cultured cells.

0031 The present invention also provides a method of manufacturing a placental product comprising: obtaining a placenta, wherein the placenta comprises a chorionic membrane, selectively depleting the placenta of immunogenicity, and cryopreserving the placenta, thereby providing a placental product. According to the present invention, the selective depletion step comprises removing immunogenic cells (e.g. CD14+ macrophages and/or trophoblasts) and/or immunogenic factors (e.g. TNF α). Optionally, the selective depletion step comprises selectively immunodepleting the placenta, whereby the placental product is purified from immunogenic cells and/or immunogenic factors. Optionally, the selective depletion step comprises removing a layer of trophoblasts, for example, by treatment with a digestive enzyme and/or mechanical removal. Optionally, the selective depletion step comprises removing CD14+ macrophages by a cryoprocess wherein the placental product is incubated for a period of time (e.g. about 30-60 mins.) at a temperature above freezing (e.g. at 2-8°C), and then freezing, whereby CD14+ macrophages are selectively killed relative to therapeutic cells.

0032 The present invention also provides a method of screening a placental product for therapy comprising assaying the placental product for immunogenicity and/or therapeutic value. Optionally, the step of assaying the placental product for immunogenicity comprises a Mixed Lymphocyte Reaction (MLR) and/or Lipopolysaccharide (LPS)-induced Tumor Necrosis Factor (TNF)- α secretion. Optionally, the step of assaying the placental product for therapeutic value comprises assaying the placental product for cell migration induction.

0033 The present invention also provides a method of treating a subject comprising administering a placental product to the subject. Optionally, the step of administering comprises applying the placental product to a wound, for example, topically applying the placental product to a skin wound. In one embodiment, a placental product is used in a tendon or ligament surgery to promote healing of a tendon or ligament.

0034 The present inventors have identified a need for the development of chorionic membrane products comprising at least one of IGFBP1, and adiponectin, providing superior wound healing properties.

BRIEF DESCRIPTION OF THE DRAWINGS

0035 FIG. 1 depicts freezing rates of various freezing methods.

0036 FIG. 2 depicts process cell recovery as a function of cryo volume.

0037

- 0038 FIG. 3 depicts process cell recovery as a function of refrigeration time.
- 0039 FIG. 4 shows representative images of the live/dead staining of the epithelial layer of fresh amniotic membrane.
- 0040 FIG. 5 depicts IL-2sR concentrations of various manufacturing intermediates.
- 0041 FIG. 6 depicts IL-2sR concentrations of various manufacturing intermediates.
- 0042 FIG. 7 depicts TNF α concentrations from LPS-induced secretion by placental tissues.
- 0043 FIG. 8 shows representative images of the live/dead staining of the epithelial layer of fresh amniotic membrane.
- 0044 FIG. 9 depicts a correlation between IL-2sR release and the number of CD45+ cells.
- 0045 FIG. 10 depicts a correlation between the amount of CD45+ cells present in amnion-derived cell suspensions and immunogenicity in MLR *in vitro*.
- 0046 FIG. 11 depicts expression of EGF (A), IGFBP1 (B), and Adiponectin (C) in amniotic and/or chorionic membranes.
- 0047 FIG. 12 depicts expression of IFN-2 α and TGF- β 3 in amniotic membrane homogenates.
- 0048 FIG. 13 depicts expression of BMP-2, BMP-4, PLAB, PIGF (A), and IGF-1 (B) in amniotic membrane homogenates.
- 0049 FIG. 14 depicts the ratio of MMPs to TIMPs in various membrane products.
- 0050 FIG. 15 depicts bFGF levels in amniotic and chorionic membranes (CM).
- 0051 FIG. 16 depicts representative expression of bFGF in chorionic tissue samples derived from two separate placenta donors.
- 0052 FIG. 17 depicts a schematic of the cell migration assay.
- 0053 FIG. 18 depicts the results of cell migration assay of various membrane preparations.
- 0054 FIG. 19 depicts growth factor and adiponectin expression in protein extracts of various membrane preparations.

DETAILED DESCRIPTION OF THE INVENTION

0055 As used herein, the following definitions apply:

0056 “Exemplary” (or “e.g.” or “by example”) means a non-limiting example.

0057 “hCMSCs: means human chorionic membrane stromal cells . hCMSCs are generally positive for CD73, CD70, CD90, CD105, and CD166; negative for CD45 and CD34,. hCMSCs differentiate to mesodermal lineages (osteogenic, chondrogenic, and adipogenic).

0058 “Selective depletion of immunogenicity” or “selective depletion of immunogenic cells or factors” or “selective depletion” means a placental product that retains live therapeutic cells and/or retains therapeutic efficacy for the treatment of tissue injury yet is free, substantially free, or depleted of at least one of immune cell type (e.g. CD14+ macrophages, trophoblasts, and/or vascular-tissue derived cells) and/or immunogenic factor that are otherwise present in a native placenta or chorionic membrane.

0059 “MSC” means mesenchymal stem cells and include fetal, neonatal, adult, or post-natal. “MSCs” include chorionic MSCs (CMSCs). MSCs generally express one or more of CD73, CD90, CD105, and CD166.

0060 “Native cells” means cells that are native, resident, or endogenous to the placental membrane, i.e. cells that are not exogenously added to the placental membrane.

0061 “Native factors” means placental membrane factors that are native, resident, or endogenous to the placental membrane, i.e. factors that are not exogenously added to the placental membrane.

0062 “Placental products” means the instant placental products disclosed herein.

0063 “Substantially free” means present in only a negligible amount or not present at all. For example, when a cell is abundant at least than about 20% or less than about 10% or less than about 1% of the amount in an unprocessed sample.

0064 “Substantial amount” of an element of the present invention, e.g. native factors, therapeutic factors, or selective depletion, means a value at least about 2% or at least 10% in comparison to an unprocessed, not cryopreserved, fresh membrane sample. A substantial amount can optionally be at least about 50%.

0065 “Therapeutic cells” or “beneficial cells” means stromal cells, MSCs, and/or fibroblasts.

0066 “Therapeutic factors” means placenta- or chorionic membrane- derived factors that promote wound healing. Examples include IGFBP1, adiponectin, α 2-macroglobulin, and/or bFGF. Other examples include MMP-9 and TIMP1.

0067 “Stromal cells” refers to a mixed population of cells present (optionally in native proportions) composed of neonatal mesenchymal stem cells and neonatal fibroblasts. Both neonatal mesenchymal stem cells and neonatal fibroblasts are immunoprivileged; neither express surface proteins present on immunogenic cell types.

0068 In some embodiments, the present technology discloses placental products for clinical use, including use in wound healing such as diabetic foot ulcers, venous leg ulcers, and burns. The manufacturing process optionally eliminates essentially all potentially immunogenic cells from the placental membrane while preserving of specific cells that play an important role in wound healing.

0069 In some embodiments, the present technology discloses a placental product that is selectively devitalized. There is now provided a placental product that is selectively depleted of substantially all immunogenic cells. There is now provided a placental product that does not contain ex vivo cultured cells. There is now provided a placental product that comprises at least one of IGFBP1, and adiponectin. There is now provided a placental product that comprises IGFBP1. There is now provided a placental product that comprises adiponectin.

0070 In some embodiments, the present technology discloses a method of cryopreserving a placental product that preserves the viability of specific beneficial cells that are the primary source of factors for the promotion of healing to the wound healing process while selectively depleting immunogenic cells (e.g. killing or rendering non-immunogenic) from the chorionic membranes.

0071 In some embodiments, the present technology discloses a bioassay to test immunogenicity of manufactured placental products.

0072 In some embodiments, the present technology discloses a placental product exhibiting a ratio of MMP:TIMP comparable to that exhibited *in vivo*. The present inventors have identified a need for the development of placental products exhibiting a ratio of MMP-9 and TIMP1 of about 7-10 to one.

0073 In some embodiments, the present technology discloses a placental product that comprises α 2-macroglobulin.

0074 The present inventors have identified a need for the development of placental products that comprise α 2-macroglobulin.

0075 There is now provided a placental product that inactivates substantially all serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases present in the chorionic membrane. There is now provided a placental product that inactivates substantially all serine proteinases present in the chorionic membrane. There is now provided a placental product that inactivates substantially all cysteine proteinases present in the chorionic membrane. There is now provided a placental product that inactivates substantially all aspartic proteinases present in the chorionic membrane. There is now provided a placental product that inactivates substantially all metalloproteinases present in the chorionic membrane.

0076 In some embodiments, the present technology discloses a placental product that comprises bFGF, optionally in a substantial amount.

0077 In some embodiments, the present technology discloses a placental product exhibiting a protease-to-protease inhibitor ratio favorable for wound healing, optionally in a substantial amount.

0078 In some embodiments, the present technology discloses a cell migration assay capable of evaluating the wound-healing potential of a placental product.

0079 IGFBP1 and adiponectin are among the factors that are important for wound healing. Evaluation of proteins derived from placental products prepared according to the presently disclosed technology reveal that bFGF is one of the major factors secreted in substantial higher quantities by the chorionic membrane. Additionally, the importance of EGF for wound healing together with high levels of bFGF detected in the presently disclosed chorionic membranes support selection of bFGF as a potency marker for evaluation of membrane products manufactured for clinical use pursuant to the present disclosure.

0080 The present technology discloses a cryopreservation procedure for a placental products that selectively depletes immunogenic cells from the a chorionic membranes and preserves the viability of other beneficial cells (including at least one of mesenchymal stem cells, and fibroblasts in some embodiments and all of mesenchymal stem cells and fibroblasts in some embodiments) that are the primary

source of factors for the promotion of healing. During the development of cryopreservation methodology for chorionic membranes, the inventors of the present application evaluated key parameters of cryopreservation including volume of cryopreservative solution, effect of tissue equilibration prior to freezing, and cooling rates for a freezing procedures.

0081 Placental products, their usefulness, and their immunocompatibility are surprisingly enhanced by depletion of maternal trophoblast and selective elimination of CD14+ fetal macrophages. Immunocompatibility can be demonstrated by any means commonly known by the skilled artisan, such demonstration can be performed by the mixed Lymphocyte Reaction (MLR) and by lipopolysaccharide (LPS)-induced Tumor Necrosis Factor (TNF)- α secretion.

0082 The instant placental products contain bFGF, optionally at a substantial concentration.

0083 The instant placental products optionally secrete factors that stimulate cell migration and/or wound healing. The presence of such factors can be demonstrated by any commonly recognized method. Optionally, the factors are in a substantial amount.

0084 For example, commercially available wound healing assays exist (Cell Biolabs) and cell migration can be assessed by cell line (HMVEC, Lonza Inc.). In one embodiment, conditioned medium from the present placental products enhance cell migration.

0085 The placental products disclosed herein are useful in treating a number of wounds including: tendon repair, cartilage repair (e.g. femoral condyle, tibial plateau), ACL replacement at the tunnel/bone interface, dental tissue augmentation, fistulas (e.g. Crohn's disease, G-tube, tracheoesophageal), missing tissue at adhesion barriers (e.g. nasal septum repair, vaginal wall repair, abdominal wall repair, tumor resection), dermal wounds (e.g. partial thickness burns, toxic epidermal necrolysis, epidermolysis bullosa, pyoderma gangrenosum, ulcers e.g. diabetic ulcers (e.g. foot), venous leg ulcers), surgical wounds, hernia repair, tendon repair, bladder repair, periosteum replacement, keloids, organ lacerations, epithelial defects, and repair or replacement of a tympanic membrane.

0086 The placental products disclosed herein exhibit one or more of the following properties beneficial to the wound healing process:

- a. approximate number of cells per cm^2 being about 20,000 to about 200,000,
- b. thickness of about 40 to about 400 μm ,
- c. a thin basement membrane,
- d. low immunogenicity,
- e. cryopreserved/cryopreserveable, and
- f. human Chorionic Membrane Stromal Cells (hCMSC).

0087 The present inventors have now identified a need for the development of placental products that do not contain *ex vivo* cultured cells.

0088 The present inventors have now identified a need for the development of placental products comprising IGFBP1.

0089 The present inventors have now identified a need for the development of placental products comprising adiponectin.

0090 The present inventors have now identified a need for the development of placental products exhibiting a protease-to-protease inhibitor ratio favorable for wound healing.

0091 The present inventors have now identified a need for the development of a method of cyropreserving placental products that preserves the viability of specific cells that are other beneficial cells that are the primary source of factors for the promotion of healing to the wound healing process while selectively depleting immunogenic cells from chorionic membranes.

0092 The present inventors have now identified a need for the development of a bioassay to test immunogenicity of manufactured placental products.

0093 The present inventors have now identified a need for the development of placental products exhibiting a ratio of MMP to TIMP comparable to that exhibited *in vivo*. The present inventors have now identified a need for the development of placental products exhibiting a ratio of MMP-9 and TIMP1 of about 7-10 to one.

0094 The present inventors have now identified a need for the development of placental products that comprise $\alpha 2$ -macroglobulin.

0095 The present inventors have now identified a need for the development of placental products that inactivate serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases. The present inventors have now identified a need for the development of placental products that inactivate serine proteinases. The present inventors have now identified a need for the development of placental products that inactivate cysteine proteinases. The present inventors have now identified a need for the development of placental products that inactivate aspartic proteinases. The present inventors have now identified a need for the development of placental products that inactivate metalloproteinases.

0096 The present inventors have now identified a need for the development of placental products that comprise bFGF.

0097 The present inventors have now identified a need for the development of a cell migration assay to evaluate the potential of placental membrane products.

0098 The present inventors have now identified a need for the development of a placental product for wound healing that comprises mesenchymal stem cells and fibroblasts.

Placental Product

Overview

0099 One embodiment of the present invention provides a placental product comprising a cryopreservation medium and a chorionic membrane, wherein the chorionic membrane comprises viable therapeutic native cells and native therapeutic factors, and wherein the cryopreservation medium comprises a cryopreserving amount of a cryopreservative. According to this embodiment, the chorionic membrane is substantially free of at least one or 2 or 3 immunogenic cell types such as: trophoblasts, CD14+ macrophages, and vascularized tissue-derived cells.

00100 In one embodiment, the chorionic membrane comprises one or more layers which exhibit the architecture of the native chorionic membrane (e.g. has not been homogenized or treated with collagenase).

00101 In one embodiment, the placental product is suitable for dermal application to a wound.

00102 With the teachings provided herein, the skilled artisan can now produce the present placental products. The present disclosure provides methods of manufacture

that produce the technical features of the present placental products. Accordingly, in one embodiment, the placental product is manufactured by steps taught herein. The present placental products are not limited to products manufactured by the methods taught here. For example, products of the present invention could be produced through methods that rely on screening steps; e.g. steps to screen for preparations with the described technical features and superior properties.

00103 The present placental products comprises one or more of the following technical features:

- a. the viable therapeutic native cells are capable of differentiating into cells of more than one lineage (e.g. osteogenic, adipogenic and/or chondrogenic lineages),
- b. the native therapeutic factors include IGFBP1, optionally present in a substantial amount,
- c. the native therapeutic factors include adiponectin, optionally present in a substantial amount,
- d. the native therapeutic factors include α 2-macroglobulin, optionally present in a substantial amount,
- e. the native therapeutic factors include bFGF, optionally present in a substantial amount,
- f. the native therapeutic factors include EGF, optionally present in a substantial amount,
- g. the native therapeutic factors include MMP-9 and TIMP1, optionally present in a substantial amount,
- h. the native therapeutic factors include MMP-9 and TIMP1 in a ratio of about 7 to about 10,
- i. the placental product does not comprise ex-vivo cultured cells,
- j. the cryopreservative medium is present in an amount of greater than about 20 ml or greater than about 50 ml,
- k. the cryopreservative comprises DMSO,
- l. cryopreservative comprises DMSO in a majority amount,

- m. the cryopreservation medium further comprises albumin, optionally wherein the albumin is HSA,
- n. the cryopreservative comprises DMSO and albumin (e.g. HSA),
- o. the chorionic membrane comprises about 5,000 to about 240,000 cells/cm² or about 20,000 to about 60,000 cells/cm²,
- p. the chorionic membrane comprises 20,000 to about 200,000 cells/cm², with a cell viability of at least about 70%,
- q. comprises at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 of stromal cells per cm² of the chorionic membrane,
- r. comprises about 5,000 to about 50,000 of stromal cells per cm² of the chorionic membrane,
- s. comprises about 4% to about 46% of viable non-culturally expanded fibroblasts per cm² of the placental product,
- t. comprises stromal cells wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle,
- u. comprises stromal cells wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle,
- v. the chorionic membrane has a thickness of about 40 μ m to about 400 μ m,
- w. secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours,
- x. cryopreservation and thawing, secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours,
- y. after refrigeration, cryopreservation and thawing, secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium

upon placing a 2 cm x 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours,

- z. the maternal side of the chorionic membrane comprises fragments of extracellular matrix proteins in a concentration substantially greater than that of a native, unprocessed chorion, optionally wherein the chorionic membrane has been treated with Dispase II or wherein a substantial portion of the protein fragments comprises terminal leucine or phenylalanine,
- aa. further comprises an amniotic membrane,
- bb. further comprises an amniotic membrane, wherein the amniotic membrane comprises a layer of amniotic epithelial cells,
- cc. further comprises an amniotic membrane, wherein the amniotic membrane and the chorionic membrane are associated to one another in the native configuration,
- dd. further comprises an amniotic membrane, wherein the amniotic membrane and the chorionic membrane are not attached to one another in the native configuration,
- ee. further comprises an amniotic membrane wherein the chorionic membrane comprises about 2 to about 4 times more stromal cells relative to the amniotic membrane,
- ff. does not comprise an amniotic membrane,
- gg. the chorionic membrane comprises about 2 to about 4 times more stromal cells relative to an amniotic membrane of the same area derived from the same placenta, and
- hh. is suitable for dermal application to a wound.

Cells

00104 According to the present invention, a placental product comprises native therapeutic cells of the chorionic membrane. The cells comprise one or more of stromal cells, MSCs, and fibroblasts.

00105 In one embodiment, the native therapeutic cells comprise viable stromal cells.

00106 In one embodiment, the native therapeutic cells comprise viable MSCs.

00107 In one embodiment, the native therapeutic cells comprise viable fibroblasts.

00108 In one embodiment, the native therapeutic cells comprise viable MSCs and viable fibroblasts.

00109 In one embodiment, the native therapeutic cells comprise viable MSCs and viable fibroblasts.

00110 In one embodiment, the native therapeutic cells comprise viable stromal cells and viable epithelial cells.

00111 In one embodiment, the therapeutic native cells are viable cells capable of differentiating into cells of more than one lineage (e.g. osteogenic, adipogenic and/or chondrogenic lineages).

00112 In one embodiment, the chorionic membrane comprises about 10,000 to about 360,000 cells/cm² or about 40,000 to about 90,000 cells/cm².

00113 In one embodiment, the chorionic membrane comprises at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 of stromal cells per cm² of the chorionic membrane.

00114 In one embodiment, the chorionic membrane comprises about 5,000 to about 50,000 of stromal cells per cm² of the chorionic membrane.

00115 In one embodiment, the chorionic membrane comprises stromal cells wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle.

00116 In one embodiment, the chorionic membrane comprises stromal cells wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle.

00117 In one embodiment, the chorionic membrane (of the placental product) comprises fibroblasts in about 50% to about 90% of the total cells.

00118 In one embodiment, the chorionic membrane comprises CD14+ macrophage in an amount of less than about 5% or less than about 1% or less than about 0.5%, optionally as demonstrated by a substantial decrease in LPS stimulation of TNF α release.

00119 In one embodiment, the placental product comprises about 2 to about 4 times more stromal cells relative to an amniotic membrane of the same area derived from the same placenta.

00120 In one embodiment, the placental product further comprises an amniotic membrane, wherein the placental product contains about 2 to about 4 times more stromal cells relative to the amniotic membrane.

00121 In one embodiment, the placental product further comprises an amniotic membrane, wherein the amniotic membrane comprises a layer of amniotic epithelial cells.

00122 In one embodiment, the placental product is substantially free of trophoblasts.

00123 In one embodiment, the placental product is substantially free of functional CD14+ macrophages.

00124 In one embodiment, the placental product is substantially free of vascularized tissue-derived cells.

In one embodiment, the placental product is substantially free of trophoblasts, functional CD14+ macrophages, and vascularized tissue-derived cells. Optionally, the placental product comprises viable stromal cells. Optionally, the placental product comprises viable MSCs. Optionally, the placental product comprises viable fibroblasts. Optionally, the placental product comprises viable MSCs and viable fibroblasts.

00125 In one embodiment, the placental product is substantially free of maternal decidual cells.

00126 In one embodiment, the placental product is substantially free of maternal leukocytes and/or trophoblast cells.

00127 In one embodiment, the chorionic membrane (of the placental product) comprises MSCs in an amount of about 5% to about 30%, about 5% to about 25%, about 5% to about 20%, about 5% to about 15%, about 3% to about 12%, at least about 5%, at least about 10%, or at least about 15%, relative to the total number of cells in the chorionic membrane. Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the MSCs are viable after a freeze-thaw cycle.

00128 In one embodiment, the chorionic membrane (of the placental product) comprises fibroblasts in an amount of about 50% to about 95%, about 60% to about

90%, or about 70% to about 85%, relative to the total number of cells in the chorionic membrane. Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the fibroblasts are viable after a freeze-thaw cycle.

00129 In one embodiment, the chorionic membrane (of the placental product) comprises functional macrophages in an amount of less than about any of: 5%, 4%, 3%, 2%, 1%, or 0.1%.

00130 In one embodiment, the chorionic membrane (of the placental product) comprises MSCs and functional macrophages in a ratio of greater than about any of: 3:1, 4:1, 5:1, 7:1, 10:1, 12:1, or 15:1.

00131 In one embodiment, the chorionic membrane comprises fibroblasts and functional macrophages in a ratio of greater than about any of: 14:1, 15:1, 16:1, 17:1, 28:1, 30:1, 35:1, 45:1, or 50:1.

00132 In one embodiment, the chorionic membrane (of the placental product) comprises fibroblasts and MSCs in a ratio of: about 9:2 to about 17:3.

00133 In one embodiment, the chorionic membrane (of the placental product) comprises MSCs in an amount of at least about 1,500 cells/cm², at least about 3,000 cells/cm², about 15,000 to about 9,000 cells/cm², or about 3,000 to about 9,000 cells/cm². Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the MSCs are viable after a freeze-thaw cycle.

00134 In one embodiment, the chorionic membrane (of the placental product) comprises fibroblasts in an amount of at least about 7,000 cells/cm², at least about 14,000 cells/cm², about 7,000 to about 51,000 cells/cm², or about 14,000 to about 51,000 cells/cm². Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the fibroblasts are viable after a freeze-thaw cycle.

00135 In one embodiment, the chorionic membrane (of the placental product) comprises functional macrophages in an amount of less than about 3,000 cells/cm², or less than about 1,000 cells/cm².

00136 In one embodiment, the placental product is substantially free of ex-vivo cultured cells.

Placental Factors

00137 According to the present invention, a placental product comprises native therapeutic factors of the chorionic membrane.

00138 In one embodiment, the factors include one or more of: IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9, and TIMP1. Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native chorionic membrane or layer thereof (e.g. $\pm$ 10% or 20%).

00139 In one embodiment, the factors include IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9, and TIMP1. Optionally, the factors are present in ratios that are substantially similar to that of a native chorionic membrane or layer thereof. Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native chorionic membrane or layer thereof (e.g. $\pm$ 10% or 20%).

00140 In one embodiment, the factors include MMP-9 and TIMP1 in a ratio of about 7 to about 10 (e.g. about 7). Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native chorionic membrane or layer thereof (e.g. $\pm$ 10% or 20%).

00141 In one embodiment, the factors include one or more (e.g. a majority or all) of the factors listed in Table 15. Optionally, the factors are present in ratios that are substantially similar to that of a native chorionic membrane or layer thereof. Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native chorionic membrane or layer thereof (e.g. $\pm$ 10% or 20%).

00142 In one embodiment, the placental product secretes substantially less TNF- α /cm² than a native, unprocessed chorionic membrane.

00143 In one embodiment, the placental product secretes substantially less TNF- α /cm² than a native placental product upon stimulation by LPS or CT.

00144 In one embodiment, the placental product secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.

00145 In one embodiment, after cryopreservation and thawing, the placental product secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.

00146 In one embodiment, after refrigeration, cryopreservation and thawing, the placental product secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.

00147 In one embodiment, the placental product further comprises an exogenously added inhibitor of TNF- α . Optionally, the inhibitor of TNF- α is IL-10.

00148 In one embodiment, the product has been treated with an antibiotic

Architecture

00149 A placental product of the present invention comprises one or more non-trophoblast layers which exhibit the architecture of the native chorionic membrane. With the teachings provided herein, the skilled artisan will recognize placental layers that exhibit native architecture, for example, layers that have not been homogenized or treated with collagenase or other enzyme that substantially disrupts the layer.

00150 In one embodiment, the placental product comprises a stromal layer with native architecture.

00151 In one embodiment, the placental product comprises a basement membrane with native architecture.

00152 In one embodiment, the placental product comprises a reticular layer with native architecture.

00153 In one embodiment, the placental product comprises a reticular layer and a basement layer with native architecture.

00154 In one embodiment, the placental product comprises a stromal layer, a basement layer, and a reticular layer with native architecture.

00155 In one embodiment, the placental product is substantially free of trophoblasts. In one embodiment, the placental product comprises a basement membrane with native architecture and the chorionic membrane is substantially free of trophoblasts. Optionally, the maternal side of the placental product comprises fragments of extracellular matrix proteins in a concentration substantially greater than that of a native chorionic membrane. Optionally, the placental product has been treated with Dispase (e.g. Dispase II) and/or a substantial portion of the extracellular matrix protein fragments comprises terminal leucine or phenylalanine.

00156 In one embodiment, the placental product has a thickness of about 40 μm to about 400 μm .

00157 In one embodiment, the placental product further comprises an amniotic membrane. Optionally, the amniotic membrane and the chorionic membrane in the placental product are associated to one another in the native configuration. Alternatively, the amniotic membrane and the chorionic membrane are not attached to one another in the native configuration.

00158 In one embodiment, the placental product does not comprise an amniotic membrane.

Formulation

00159 According to the present invention, the placental product can be formulated with a cryopreservation medium.

00160 In one embodiment, the cryopreservation medium comprising one or more cell-permeating cryopreservatives, one or more non cell-permeating cryopreservatives, or a combination thereof.

00161 Optionally, the cryopreservation medium comprises one or more cell-permeating cryopreservatives selected from DMSO, a glycerol, a glycol, a propylene glycol, an ethylene glycol, or a combination thereof.

00162 Optionally, the cryopreservation medium comprises one or more non cell-permeating cryopreservatives selected from polyvinylpyrrolidone, a hydroxyethyl starch, a polysaccharide, a monosaccharides, a sugar alcohol, an alginate, a trehalose, a raffinose, a dextran, or a combination thereof.

00163 Other examples of useful cryopreservatives are described in "Cryopreservation" (BioFiles Volume 5 Number 4 –Sigma-Aldrich® datasheet).

00164 In one embodiment, the cryopreservation medium comprises a cell-permeating cryopreservative, wherein the majority of the cell-permeating cryopreservative is DMSO. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

00165 In one embodiment, the cryopreservation medium comprises DMSO. Optionally, the cryopreservation medium does not comprise glycerol in a majority

amount. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

00166 In one embodiment, the cryopreservation medium comprises additional components such as albumin (e.g. HSA or BSA), an electrolyte solution (e.g. Plasma-Lyte), or a combination thereof.

00167 In one embodiment, the cryopreservation medium comprises 1% to about 15% albumin by weight and about 5% to about 20% cryopreservative by volume (e.g. about 10%). Optionally, the cryopreservative comprises DMSO (e.g. in a majority amount).

00168 In one embodiment, the placental product is formulated in greater than about 20 ml or greater than about 50 ml of cryopreservation medium. Optionally, the cryopreservative comprises DMSO (e.g. in a majority amount). Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

00169 In one embodiment, the placental product is placed on nitrocellulose paper.

00170 In one embodiment, the placenta is cut into a plurality of sections. Optionally, the sections are less than about 10 cm x 10 cm. Optionally, the sections are between about 2 cm x 2 cm and 5 cm x 5 cm.

Manufacture

Overview

00171 A placental product of the present invention can be manufactured from a placenta in any suitable manner that provides the technical features taught herein. According to the present invention, a placental product comprises at least an immunocompatible chorionic membrane.

00172 In one embodiment, a placental product is manufactured by a method comprising:

- a. obtaining a placenta,
- b. selectively depleting the placenta of immunogenicity; and
- c. cryopreserving the placenta.

00173 In one embodiment, a placental product is manufactured by a method comprising:

- a. obtaining a placenta;

- b. removing a substantial portion of trophoblasts from the placenta; and
- c. cryopreserving the placenta.

00174 Optionally, the method comprises a step of removing the amniotic membrane or portion thereof ('an amniotic membrane') from the placenta. Optionally, the method comprises a step of removing an amniotic membrane from the placenta without removing a substantial portion of amniotic epithelial cells from the placenta.

00175 Optionally, the step of removing a substantial portion of trophoblasts from the placenta comprises treating the placenta with a digestive enzyme such as a protease (e.g. dispase or dispase II), mechanically removing trophoblasts from the placenta (e.g. by scraping), or a combination thereof.

00176 Optionally, the method comprises a step of removing vascularized tissue from the placenta, for example, by lysing red blood cells, by removing blood clots, or a combination thereof.

00177 Optionally, the method comprises a step of treating the placenta with one or more antibiotics.

00178 Optionally, the method comprises a step of selective depletion of CD14+ macrophages.

00179 Optionally, the step of cryopreserving the placenta comprises freezing the placenta in a cryopreservation medium which comprises one or more cell-permeating cryopreservatives, one or more non cell-permeating cryopreservatives, or a combination thereof.

00180 Optionally, the step of cryopreserving the placenta comprises refrigerating for a period of time and then freezing, thereby selectively depleting CD14+ macrophages.

00181 An exemplary placental product of the present invention can be manufactured or provided with a bandage or skin substitute.

Immunocompatibility and Selective Depletion

00182 In one embodiment, the invention the placental product is immunocompatible. Immunocompatibility can be accomplished by any selective depletion step that removes immunogenic cells or factors or immunogenicity from the placenta (or chorionic membrane thereof).

00183 In one embodiment, the placental product is made immunocompatible by selectively depleting it of functional immunogenic cells. A placenta can be made immunocompatible by selectively removing immunogenic cells from the placenta (or chorionic membrane thereof) relative to therapeutic cells. For example, immunogenic cells can be removed by killing the immunogenic cells or by purification of the placenta there from.

00184 In one embodiment, the placenta is made immunocompatible by selectively depleting trophoblasts, for example, by removal of the trophoblast layer.

00185 In one embodiment, the placenta is made immunocompatible by selective depletion of functional CD14+ macrophages, optionally resulting in depletion of TNF α upon stimulation, or a combination thereof.

00186 In one embodiment, the placenta is made immunocompatible by selective depletion of vascularized tissue-derived cells.

00187 In one embodiment, the placenta is made immunocompatible by selective depletion of functional CD14+ macrophages, trophoblasts, and vascularized tissue-derived cells.

00188 In one embodiment, the placenta product is made immunocompatible by selective depletion of trophoblasts and/or CD14+ macrophages, optionally resulting in depletion of TNF α upon stimulation.

Trophoblast Removal

00189 In one embodiment, trophoblasts are depleted or removed from the placental product. Surprisingly, such a placental product has one or more of the following superior features:

- a. is substantially non-immunogenic;
- b. provides remarkable healing time; and
- c. provides enhanced therapeutic efficacy.

00190 Trophoblasts can be removed in any suitable manner which substantially diminishes the trophoblast content of the placental product. Optionally, the trophoblasts are selectively removed or otherwise removed without eliminating a substantial portion of one or more therapeutic components from the placenta (e.g.

MSCs, placental factors, etc). Optionally, a majority (e.g. substantially all) of the trophoblasts are removed.

00191 One method of removing trophoblasts comprises treating the placenta (e.g. chorion or amnio-chorion) with a digestive enzyme such as dispase (e.g. dispase II) and separating the trophoblasts from the placenta. Optionally, the step of separating comprises mechanical separation such as peeling or scraping. Optionally, scraping comprises scraping with a soft instrument such as a finger.

00192 One method of removing trophoblasts comprises treating the chorionic membrane with dispase for about 30 to about 45 minutes separating the trophoblasts from the placenta. Optionally, the dispase is provided in a solution of about less than about 1% (e.g. about 0.5%). Optionally, the step of separating comprises mechanical separation such as peeling or scraping. Optionally, scraping comprises scraping with a soft instrument such as a finger.

00193 Useful methods of removing trophoblasts from a placenta (e.g. chorion) are described by Portmann-Lanz et al. ("Placental mesenchymal stem cells as potential autologous graft for pre- and perinatal neuroregeneration"; American Journal of Obstetrics and Gynecology (2006) 194, 664–73), ("Isolation and characterization of mesenchymal cells from human fetal membranes"; Journal Of Tissue Engineering And Regenerative Medicine 2007; 1: 296–305.), and (Concise Review: Isolation and Characterization of Cells from Human Term Placenta: Outcome of the First International Workshop on Placenta Derived Stem Cells").

00194 In one embodiment, trophoblasts are removed before cryopreservation.

Macrophage Removal

00195 In one embodiment, functional macrophages are depleted or removed from the placental product. Surprisingly, such a placental product has one or more of the following superior features:

- a. is substantially non-immunogenic;
- b. provides remarkable healing time; and
- c. provides enhanced therapeutic efficacy.

00196 Functional macrophages can be removed in any suitable manner which substantially diminishes the macrophage content of the placental product. Optionally, the macrophages are selectively removed or otherwise removed without eliminating a

Page 26

substantial portion of one or more therapeutic components from the placenta (e.g. MSCs, placental factors, etc). Optionally, a majority (e.g. substantially all) of the macrophages are removed.

00197 One method of removing immune cells such as macrophages comprises killing the immune cells by rapid freezing rates such as 60-100°C/min.

00198 Although immune cells can be eliminated by rapid freezing rates, such a method can also be detrimental to therapeutic cells such as stromal cells (e.g. MSCs). The present inventors have discovered a method of selectively killing CD14+ macrophages can be selectively killed by refrigerating the placenta for a period of time (e.g. for at least about 10 min such as for about 30-60 mins) at a temperature above freezing (e.g. incubating at 2-8°C) and then freezing the placenta (e.g. incubating at -80°C ± 5°C). Optionally, the step of freezing comprises freezing at a rate of less than 10°/min (e.g. less than about 5°/min such as at about 1°/min).

00199 In one embodiment, the step of refrigerating comprises soaking the placenta in a cryopreservation medium (e.g. containing DMSO) for a period of time sufficient to allow the cryopreservation medium to penetrate (e.g. equilibrate with) the placental tissues. Optionally, the step of freezing comprises reducing the temperature at a rate of about 1°/min. Optionally, the step of freezing comprises freezing at a rate of less than 10°/min (e.g. less than about 5°/min such as at about 1°/min).

00200 In one embodiment, the step of refrigerating comprises soaking the placenta in a cryopreservation medium (e.g. containing DMSO) at a temperature of about -10-15°C (e.g. at 2-8°C) for at least about any of: 10 min, 20 min, 30 min, 40 min, or 50 min. In another embodiment, the step of refrigerating comprises soaking the placenta in a cryopreservation medium (e.g. containing DMSO) at a temperature of about -10-15°C (e.g. at 2-8°C) for about any of: 10-120, 20-90 min, or 30-60 min. Optionally, the step of freezing comprises freezing at a rate of less than 10°/min (e.g. less than about 5°/min such as at about 1°/min).

Removal of Vascularized Tissue-Derived Cells

00201 In one embodiment, vascularized tissue-derived cells (or vascularized tissue) are depleted or removed from the placental product. Surprisingly, such a placental product has one or more of the following superior features:

- a. is substantially non-immunogenic;

- b. provides remarkable healing time; and
- c. provides enhanced therapeutic efficacy.

00202 Vascularized tissue-derived cells can be removed in any suitable manner which substantially diminishes such cell content of the placental product. Optionally, the vascularized tissue-derived cells are selectively removed or otherwise removed without eliminating a substantial portion of one or more therapeutic components from the placenta (e.g. MSCs, placental factors, etc).

00203 In one embodiment, removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord. The chorion on the umbilical side of the placenta is not removed due to the vascularization on this side.

00204 In one embodiment, removal of vascularized tissue-derived cells comprises rinsing the chorionic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells.

00205 In one embodiment, removal of vascularized tissue-derived cells comprises treating the chorionic membrane with an anticoagulant (e.g. citrate dextrose solution).

00206 In one embodiment, removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord and rinsing the chorionic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells.

00207 In one embodiment, removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord and treating the chorionic membrane with an anticoagulant (e.g. citrate dextrose solution).

00208 In one embodiment, removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord, rinsing the chorionic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells, and treating the chorionic membrane with an anticoagulant (e.g. citrate dextrose solution).

Selective depletion of immunogenicity as demonstrated by a substantial decrease in LPS stimulation of TNF α release.

00209 In one embodiment, the placental product is selectively depleted of immunogenicity as demonstrated by a reduction in LPS stimulated TNF- α release. depletion d of TNF- α depleted or removed from the placental product.

00210 In one embodiment, TNF- α is depleted by killing or removal of macrophages.

00211 In one embodiment, TNF- α is depleted by treatment with an anti- TNF- α antibody.

00212 In one embodiment, TNF- α is functionally depleted by treatment with IL-10, which suppresses TNF- α secretion.

Preservation

00213 A placental product of the present invention may be used fresh or may be preserved for a period of time. Surprisingly, cryopreservation results in immunocompatible placental products.

00214 In one embodiment, a placental product is cryopreserved. A placental product may be cryopreserved by incubation at freezing temperatures (e.g. a $-80^{\circ}\text{C} \pm 5^{\circ}\text{C}$) in a cryopreservative medium.

00215 Cryopreservation can comprise, for example, incubating the placental product at 4°C for 30-60 min, and then incubating at -80°C until use. The placental product may then be thawed for use. Optionally, the placental product is cryopreserved in a manner such that cell viability is retained surprisingly well after a freeze-thaw cycle.

00216 In one embodiment, cryopreservation comprises storage in a cryopreservation medium comprising one or more cell-permeating cryopreservatives, one or more non cell-permeating cryopreservatives, or a combination thereof. Optionally, the cryopreservation medium comprises one or more cell-permeating cryopreservatives selected from DMSO, a glycerol, a glycol, a propylene glycol, an ethylene glycol, or a combination thereof. Optionally, the cryopreservation medium comprises one or more non cell-permeating cryopreservatives selected from polyvinylpyrrolidone, a hydroxyethyl starch, a polysaccharide, a monosaccharides, a sugar alcohol, an alginate, a trehalose, a raffinose, a dextran, or a combination thereof. Other examples of useful cryopreservatives are described in "Cryopreservation" (BioFiles Volume 5 Number 4 –Sigma-Aldrich® datasheet).

00217 In one embodiment, the cryopreservation medium comprises a cell-permeating cryopreservative, wherein the majority of the cell-permeating

cryopreservative is DMSO. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

00218 In one embodiment, the cryopreservation medium comprises DMSO. Optionally, the cryopreservation medium does not comprise glycerol in a majority amount. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

00219 In one embodiment, the cryopreservation medium comprises additional components such as albumin (e.g. HSA or BSA), an electrolyte solution (e.g. Plasma-Lyte), or a combination thereof.

00220 In one embodiment, the cryopreservation medium comprises 1% to about 15% albumin by weight and about 5% to about 20% cryopreservative by volume (e.g. about 10%). Optionally, the cryopreservative comprises DMSO (e.g. in a majority amount).

00221 In one embodiment, cryopreservation comprises placing the placenta on nitrocellulose paper.

00222 In one embodiment, the placenta is cut into a plurality of sections before cryopreservation. Optionally, the sections are placed on nitrocellulose paper before refrigeration.

Methods of Use

00223 The placental products (e.g. derived from chorionic tissue) of the present invention may be used to treat any tissue injury. A method of treatment may be provided, for example, by administering to a subject in need thereof, a placental product of the present invention.

00224 A typical administration method of the present invention is topical administration. Administering the present invention can optionally involve administration to an internal tissue where access is gained by a surgical procedure.

00225 Placental products can be administered autologously, allogeneically or xenogeneically.

00226 In one embodiment, a present placental product is administered to a subject to treat a wound. Optionally, the wound is a laceration, scrape, thermal or chemical

burn, incision, puncture, or wound caused by a projectile. Optionally, the wound is an epidermal wound, skin wound, chronic wound, acute wound, external wound, internal wounds, congenital wound, ulcer, or pressure ulcer. Such wounds may be accidental or deliberate, e.g., wounds caused during or as an adjunct to a surgical procedure. Optionally, the wound is closed surgically prior to administration.

00227 In one embodiment, a present placental product is administered to a subject to treat a burn. Optionally, the burn is a first-degree burn, second-degree burn (partial thickness burns), third degree burn (full thickness burns), infection of burn wound, infection of excised and unexcised burn wound, loss of epithelium from a previously grafted or healed burn, or burn wound impetigo.

00228 In one embodiment, a present placental product is administered to a subject to treat an ulcer, for example, a diabetic ulcer (e.g. foot ulcer).

00229 In one embodiment, a placental product is administered by placing the placental product directly over the skin of the subject, e.g., on the stratum corneum, on the site of the wound, so that the wound is covered, for example, using an adhesive tape. Additionally or alternatively, the placental product may be administered as an implant, e.g., as a subcutaneous implant.

00230 In one embodiment, a placental product is administered to the epidermis to reduce rhtids or other features of aging skin. Such treatment is also usefully combined with so-called cosmetic surgery (e.g. rhinoplasty, rhytidectomy, etc.).

00231 In one embodiment, a placental product is administered to the epidermis to accelerate healing associated with a dermal ablation procedure or a dermal abrasion procedure (e.g. including laser ablation, thermal ablation, electric ablation, deep dermal ablation, sub-dermal ablation, fractional ablation, and microdermal abrasion).

00232 Other pathologies that may be treated with placental products of the present invention include traumatic wounds (e.g. civilian and military wounds), surgical scars and wounds, spinal fusions, spinal cord injury, avascular necrosis, reconstructive surgeries, ablations, and ischemia.

00233 In one embodiment, a placental product of the present invention is used in a tissue graft procedure. Optionally, the placental product is applied to a portion of the graft which is then attached to a biological substrate (e.g. to promote healing and/or attachment to the substrate). By way of non-limiting example, tissues such as skin,

cartilage, ligament, tendon, periosteum, perichondrium, synovium, fascia, mesenter and sinew can be used as tissue graft.

00234 In one embodiment, a placental product is used in a tendon or ligament surgery to promote healing of a tendon or ligament. Optionally, the placental product is applied to portion of a tendon or ligament which is attached to a bone. The surgery can be any tendon or ligament surgery, including, e.g. knee surgery, shoulder, leg surgery, arm surgery, elbow surgery, finger surgery, hand surgery, wrist surgery, toe surgery, foot surgery, ankle surgery, and the like. For example, the placental product can be applied to a tendon or ligament in a grafting or reconstruction procedure to promote fixation of the tendon or ligament to a bone.

00235 Through the insight of the inventors, it has surprisingly been discovered that placental products of the present invention provide superior treatment (e.g. healing time and/or healing strength) for tendon and ligament surgeries. Tendon and ligament surgeries can involve the fixation of the tendon or ligament to bone. Without being bound by theory, the present inventors believe that osteogenic and/or chondrogenic potential of MSCs in the present placental products promotes healing process and healing strength of tendons or ligaments. The present inventors believe that the present placental products provide an alternative or adjunctive treatment to periosteum-based therapies. For example, useful periosteum based treatments are described in Chen et al. ("Enveloping the tendon graft with periosteum to enhance tendon-bone healing in a bone tunnel: A biomechanical and histologic study in rabbits"; Arthroscopy. 2003 Mar;19(3):290-6), Chen et al. ("Enveloping of periosteum on the hamstring tendon graft in anterior cruciate ligament reconstruction"; Arthroscopy. 2002 May-Jun;18(5):27E), Chang et al. ("Rotator cuff repair with periosteum for enhancing tendon-bone healing: a biomechanical and histological study in rabbits"; Knee Surgery, Sports Traumatology, Arthroscopy Volume 17, Number 12, 1447-1453), each of which are incorporated by reference.

00236 As non-limiting example of a method of tendon or ligament surgery, a tendon is sutured to and/or wrapped or enveloped in a placental membrane and the tendon is attached to a bone. Optionally, the tendon is placed into a bone tunnel before attached to the bone.

00237 In one embodiment, the tendon or ligament surgery is a graft procedure, wherein the placental product is applied to the graft. Optionally, the graft is an allograft, xenograft, or an autologous graft.

00238 In one embodiment, the tendon or ligament surgery is repair of a torn ligament or tendon, wherein the placental product is applied to the torn ligament or tendon.

00239 Non-limiting examples of tendons to which a placental product can be applied include a digitorum extensor tendon, a hamstring tendon, a bicep tendon, an Achilles Tendon, an extensor tendon, and a rotator cuff tendon.

00240 In one embodiment, a placental product of the present invention is used to reduce fibrosis by applying the placental product to a wound site.

00241 In one embodiment, a placental product of the present invention is used as an anti-adhesion wound barrier, wherein the placental product is applied to a wound site, for example, to reduce fibrosis (e.g. postoperative fibrosis).

00242 Non-limiting examples of wound sites to which the placental product can be applied include those that are surgically induced or associated with surgery involving the spine, laminectomy, knee, shoulder, or child birth, trauma related wounds or injuries, cardiovascular procedures, angiogenesis stimulation, brain/neurological procedures, burn and wound care, and ophthalmic procedures. For example, optionally, the wound site is associated with surgery of the spine and the stromal side of the placental product is applied to the dura (e.g. the stromal side facing the dura). Direction for such procedures, including the selection of wound sites and/or methodologies, can be found, for example, in WO 2009/132186 and US 2010/0098743, which are hereby incorporated by reference.

00243 A placental product of the present invention can optionally be used to reduce adhesion or fibrosis of a wound. Postoperative fibrosis is a natural consequence of all surgical wound healing. By example, postoperative peridural adhesion results in tethering, traction, and compression of the thecal sac and nerve roots, which cause a recurrence of hyperesthesia that typically manifests a few months after laminectomy surgery. Repeated surgery for removal of scar tissue is associated with poor outcome and increased risk of injury because of the difficulty of identifying neural structures that are surrounded by scar tissue. Therefore, experimental and clinical studies have primarily focused on preventing the adhesion of scar tissue to the dura matter and

nerve roots. Spinal adhesions have been implicated as a major contributing factor in failure of spine surgery. Fibrotic scar tissue can cause compression and tethering of nerve roots, which can be associated with recurrent pain and physical impairment.

00244 Without being bound by theory, the present inventors believe that placental products taught herein are useful to reduce adhesion or fibrosis of a wound, at least in part, because the placental products can perform the very critical function in-situ of providing a immunoprivileged environment (i.e. relatively high resistance against immune responses) in the human development process. One advantage of the wound dressings and processes of the present invention is that an anti-adhesion barrier is provided which can be used to prevent adhesions following surgery, and in particular following back surgery.

00245 In the preceding paragraphs, use of the singular may include the plural except where specifically indicated. As used herein, the words “a,” “an,” and “the” mean “one or more,” unless otherwise specified. In addition, where aspects of the present technology are described with reference to lists of alternatives, the technology includes any individual member or subgroup of the list of alternatives and any combinations of one or more thereof.

00246 The disclosures of all patents and publications, including published patent applications, are hereby incorporated by reference in their entireties to the same extent as if each patent and publication were specifically and individually incorporated by reference.

00247 It is to be understood that the scope of the present technology is not to be limited to the specific embodiments described above. The present technology may be practiced other than as particularly described and still be within the scope of the accompanying claims.

00248 Likewise, the following examples are presented in order to more fully illustrate the present technology. They should in no way be construed, however, as limiting the broad scope of the technology disclosed herein.

00249 The presently described technology and its advantages will be better understood by reference to the following examples. These examples are provided to describe specific embodiments of the present technology. By providing these specific examples, it is not intended limit the scope and spirit of the present technology. It will be understood by those skilled in the art that the full scope of the presently described

technology encompasses the subject matter defined by the claims appending this specification, and any alterations, modifications, or equivalents of those claims.

EXAMPLES

00250 Other features and embodiments of the present technology will become apparent from the following examples which are given for illustration of the present technology rather than for limiting its intended scope.

Example 1 Characterization of Placental Membranes

00251 Characterization of cells in placental membranes by Fluorescence Activated Cell Sorting (FACS) demonstrated the presence of stromal cells (Mesenchymal Stem Cell-like cells) in addition to fetal epithelial cells and fibroblasts in amniotic and/or chorionic membranes.

00252 One unique characteristic of the presently disclosed placental products is the presence of MSCs, which have been shown to be one of three types of cells (in addition to epithelial cells and fibroblasts) that are important for wound healing. Placental membranes secrete a variety of factors involved in wound healing such as angiogenic factors, factors supporting proliferation and migration of epithelial cells and fibroblasts, factors attracting endothelial stem cells from blood circulation to the wound site, antibacterial factors, and others.

00253 Evaluation of proteins secreted by exemplary placental products of the invention in comparison to Apligraf and Dermagraft demonstrated a number of growth factors present in the tested products that are involved in wound healing. Examples are Vascular Endothelial Growth Factor (VEGF), Platelet-Derived Growth Factor (PDGF), Transforming Growth Factor (TGF) and others. However, several unique factors including Epidermal Growth Factor (EGF), which is one of the key factors for wound healing, are present in placental membranes and absent in Apligraf and Dermagraft. Also, placental membranes have a favorable protease-to-protease inhibitor ratio for wound healing. In an *in vitro* model of wound healing (cell migration assay, disclosed herein), the present inventors have demonstrated that placental membranes secrete factors promoting cell migration that will support wound closure.

Example 2 Exemplary Manufacturing Process of a Placental Product

00254 In one embodiment, the present application discloses a procedure for manufacturing chorionic membranes from placenta post partum.

Example 2.1 Exemplary Manufacturing Process of Chorionic Membrane Product

00255 One method of manufacturing a placental product comprising a chorionic membrane according to the presently disclosed manufacturing procedure is as follows:

- a. Remove umbilical cord close to placental surface,
- b. Blunt dissect of the amnion to placental skirt,
- c. Flip placenta over and completely remove amnion,
- d. Remove chorion by cutting around placental skirt,
- e. Rinse the chorionic membrane in PBS to remove red blood cells,
- f. Rinse the chorionic membrane once with 11% ACD-A solution to assist in red blood cell removal,
- g. Rinse the chorionic membrane PBS to remove ACD-A solution,
- h. Treat chorion in 0.5% dispase solution at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 30-45 minutes, optionally, during dispase incubation period, use PBS to remove any remaining blood from the amnion,
- i. When dispase treatment is complete, rinse chorion with PBS to remove dispase solution,
- j. Gently remove trophoblast layer from the chorion, for example, by scraping (e.g. with finger),
- k. Place chorion into a bottle containing antibiotic solution and incubate at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 24-28 hrs,
- l. Remove bottle from the incubator and rinse each membrane with PBS to remove antibiotic solution,
- m. Mount chorion on reinforced nitrocellulose paper and cut to size,
- n. Place each piece into an FP-90 cryobag and heat seal,
- o. Add 50 mL cryopreservation solution to the bag through a syringe and remove any air trapped within the bag with the syringe,
- p. Tube seal the solution line on the FP-90 bag,
- q. Place filled bag into secondary bag and heat seal,
- r. Place unit into packaging carton,

- s. Refrigerate at 2-8°C for 30-60 minutes, Freeze at -80°C $\pm$ 5°C inside a Styrofoam container.

00256

Example 2.2 Exemplary Manufacturing Process of Product Comprising Chorionic Membrane and Amniotic Membrane

00257 One method of manufacturing a placental product comprising a chorionic membrane and an amniotic membrane according to the presently disclosed manufacturing procedure is as follows:

- a. Remove umbilical cord close to placental surface,
- b. Blunt dissect of the amnion to placental skirt,
- c. Flip placenta over and completely remove amnion,
- d. Remove chorion by cutting around placental skirt,
- e. Rinse both membranes in PBS to remove red blood cells,
- f. Rinse both membranes once with 11% ACD-A solution to assist in red blood cell removal,
- g. Rinse both membranes with PBS to remove ACD-A solution,
- h. Treat chorion in 0.5% dispase solution at 37°C $\pm$ 2°C for 30-45 minutes, optionally, during dispase incubation period, use PBS to remove any remaining blood from the amnion,
- i. Gently remove the connective tissue layer from the amnion,
- j. Place the amnion in PBS and set aside,
- k. When dispase treatment is complete, rinse chorion with PBS to remove dispase solution,
- l. Gently remove trophoblast layer from the chorion,
- m. Place the amnion and chorion each into a bottle containing antibiotic solution and incubate at 37°C $\pm$ 2°C for 24-28 hrs,
- n. Remove bottles from the incubator and rinse each membrane with PBS to remove antibiotic solution,
- o. Mount amnion (epithelial side up) or chorion on reinforced nitrocellulose paper and cut to size,
- p. Place each piece into an FP-90 cryobag and heat seal,
- q. Add 50 mL cryopreservation solution to the bag through a syringe and remove any air trapped within the bag with the syringe,

- r. Tube seal the solution line on the FP-90 bag,
- s. Place filled bag into secondary bag and heat seal,
- t. Place unit into packaging carton,
- u. Refrigerate at 2-8°C for 30-60 minutes, Freeze at -80°C $\pm$ 5°C inside a Styrofoam container.

00258

Example 2.3 Exemplary Placental Product Manufacturing Process

00259 One method manufacturing a placental product comprising a chorionic membrane according to the presently disclosed manufacturing procedure was as follows:

00260 The placenta was processed inside a biological safety cabinet. The umbilical cord was first removed, and the amniotic membrane was peeled from the underlying chorionic membrane using blunt dissection. Subsequently, the chorion was removed by cutting around the placental skirt on the side opposite of the umbilical cord. The chorion on the umbilical side of the placenta was not removed due to the vascularization on this side. The chorionic membrane was rinsed with phosphate buffered saline (PBS) (Gibco Invitrogen, Grand Island, NY) to remove gross blood clots and any excess blood cells. The membrane was then washed with 11% anticoagulant citrate dextrose solution (USP) formula A (ACD-A) (Baxter Healthcare Corp., Deerfield, IL) in saline (Baxter Healthcare Corp., Deerfield, IL) to remove remaining blood cells.

00261 The chorion was then incubated in 200 mL of a 0.5% dispase (BD Biosciences, Bedford, MA) solution in Dulbecco's Modified Eagles media (DMEM) (Lonza, Walkersville, MD) at 37°C $\pm$ 2°C for 30-45 minutes to digest the connective tissue layer between the chorion and adjacent trophoblast layer. Once the chorion incubation period was complete, the chorion was rinsed with PBS to remove the dispase solution. Subsequently, the trophoblast layer was removed by gently peeling or scraping away these maternal decidual cells.

00262 The chorion was then disinfected in 500 mL of antibiotic solution consisting of gentamicin sulfate (50 µg/mL) (Abraxis Pharmaceutical Products, Schaumburg, IL), vancomycin HCl (50 µg/mL) (Hospira Inc., Lake Forest, IL), and amphotericin B (2.5 µg/mL) (Sigma Aldrich, St. Louis, MO) in DMEM at 37°C $\pm$ 2°C for 24-28 hours.

Vented caps were used with the incubation flasks. After the incubation period, the membrane was washed with PBS to remove any residual antibiotic solution.

00263 The membrane was mounted on Optitran BA-S 85 reinforced nitrocellulose paper (Whatman, Dassel, Germany) and cut to the appropriate size prior to packaging into an FP-90 cryobag (Charter Medical Ltd., Winston-Salem, NC). Once the membrane unit was placed into the FP-90 cryobag and the cryobag was heat sealed, 50 mL of a cryopreservation solution containing 10% dimethyl sulfoxide (DMSO) (Bioniche Teo. Inverin Co., Galway, Ireland) and 5% human serum albumin (HSA) (Baxter, West Lake Village, CA) in PlasmaLyte-A (Baxter Healthcare Corp., Deerfield, IL) were added through the center tubing line. Any excess air was removed, and the tubing line was subsequently sealed.

00264 The FP-90 cryobag was placed into a mangar bag (10 in. x 6 in.) (Mangar Industries, New Britain, PA), which was then heat sealed. The mangar bag was placed into a packaging carton (10.5 in. x 6.5 in. x 0.6 in.) (Diamond Packaging, Rochester, NY). All cartons were refrigerated at 2-8°C for 30-60 minutes prior to freezing at -80°C ± 5°C inside a Styrofoam container.

00265

Example 2.4 Exemplary Manufacturing Process of a Placental Product Comprising Chorionic Membrane and Amniotic Membrane

00266 One method of manufacturing a placental product comprising a chorionic membrane product and an amniotic membrane product according to the presently disclosed manufacturing procedure was as follows:

00267 The placenta was processed inside a biological safety cabinet. The umbilical cord was first removed, and the amniotic membrane was peeled from the underlying chorionic membrane using blunt dissection. Subsequently, the chorion was removed by cutting around the placental skirt on the side opposite of the umbilical cord. The chorion on the umbilical side of the placenta was not removed due to the vascularization on this side. Both membranes were rinsed with phosphate buffered saline (PBS) (Gibco Invitrogen, Grand Island, NY) to remove gross blood clots and any excess blood cells. The membranes were then washed with 11% anticoagulant citrate dextrose solution (USP) formula A (ACD-A) (Baxter Healthcare Corp.,

Deerfield, IL) in saline (Baxter Healthcare Corp., Deerfield, IL) to remove remaining blood cells.

00268 The chorion was then incubated in 200 mL of a 0.5% dispase (BD Biosciences, Bedford, MA) solution in Dulbecco's modified eagles media (DMEM) (Lonza, Walkersville, MD) at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 30-45 minutes to digest the connective tissue layer between the chorion and adjacent trophoblast layer. During this incubation period, the stromal side of the amnion was cleaned by gently scraping away any remaining connective tissue. Once the chorion incubation period was complete, the chorion was rinsed with PBS to remove the dispase solution. Subsequently, the trophoblast layer was removed by gently peeling or scraping away these maternal decidual cells.

00269 The amnion and chorion were then each disinfected in 500 mL of antibiotic solution consisting of gentamicin sulfate (50 $\mu\text{g/mL}$) (Abraxis Pharmaceutical Products, Schaumburg, IL), vancomycin HCl (50 $\mu\text{g/mL}$) (Hospira Inc., Lake Forest, IL), and amphotericin B (2.5 $\mu\text{g/mL}$) (Sigma Aldrich, St. Louis, MO) in DMEM at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 24-28 hours. Vented caps were used with the incubation flasks. After the incubation period, the membranes were washed with PBS to remove any residual antibiotic solution.

00270 The membranes were mounted on Optitran BA-S 85 reinforced nitrocellulose paper (Whatman, Dassel, Germany) and cut to the appropriate size prior to packaging into an FP-90 cryobag (Charter Medical Ltd., Winston-Salem, NC). For the amnion, the stromal side was mounted towards the nitrocellulose paper. Once a membrane unit was placed into the FP-90 cryobag and the cryobag was heat sealed, 50 mL of a cryopreservation solution containing 10% dimethyl sulfoxide (DMSO) (Bioniche Teo. Inverin Co., Galway, Ireland) and 5% human serum albumin (HSA) (Baxter, West Lake Village, CA) in PlasmaLyte-A (Baxter Healthcare Corp., Deerfield, IL) were added through the center tubing line. Any excess air was removed, and the tubing line was subsequently sealed.

00271 The FP-90 cryobag was placed into a mangar bag (10 in. x 6 in.) (Mangar Industries, New Britain, PA), which was then heat sealed. The mangar bag was placed into a packaging carton (10.5 in. x 6.5 in. x 0.6 in.) (Diamond Packaging, Rochester, NY). All cartons were refrigerated at $2-8^{\circ}\text{C}$ for 30-60 minutes prior to freezing at $-80^{\circ}\text{C} \pm 5^{\circ}\text{C}$ inside a Styrofoam container.

Example 3 Quantitative Evaluation of Cell Number and Cell Viability after Enzymatic Digestion of Placental Membranes

00272 Amnion and chorion membranes and present placental products (from above) were evaluated for cell number and cell viability throughout the process. These analyses were performed on fresh placental tissue (prior to the antibiotic treatment step), placental tissue post antibiotic treatment, and product units post thaw. Cells were isolated from the placental membranes using enzymatic digestion. For the frozen product units, the FP-90 cryobags were first removed from the packaging cartons and mangar bags. Then the FP-90 cryobags were thawed for 2-3 minutes in a room temperature water bath. Early experiments involved the use of a $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ water bath. After thaw, the placental membranes were removed from the FP-90 cryobag and placed into a reservoir containing saline (Baxter Healthcare Corp., Deerfield, IL) for a minimum of 1 minute and a maximum of 60 minutes. Each membrane was detached from the reinforced nitrocellulose paper prior to digestion.

00273 Amniotic membranes were digested with 40 mL of 0.75% collagenase (Worthington Biochemical Corp., Lakewood, NJ) solution at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 20-40 minutes on a rocker. After collagenase digestion, the samples were centrifuged at 2000 rpm for 10 minutes. The supernatant was removed, and 40 mL of 0.05% trypsin-EDTA (Lonza, Walkersville, MD) were added and incubated at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for an additional 5-15 minutes on a rocker. The trypsin was warmed to $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ in a water bath prior to use. After trypsin digestion, the suspension was filtered through a 100 μm cell strainer nylon filter to remove any debris. Centrifugation at 2000 rpm for 10 minutes was performed, and supernatant was removed. Cell pellets were reconstituted with a volume of PlasmaLyte-A that was proportional to the pellet size, and 20 μL of the resuspended cell suspension were mixed with 80 μL of trypan blue (Sigma Aldrich, St. Louis, MO) for counting. The cell count sample was placed into a hemocytometer and evaluated using a microscope.

00274 Chorionic membranes were digested with 25 mL of 0.75% collagenase solution at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 20-40 minutes on a rocker. After collagenase digestion, the suspension was filtered through a 100 μm cell strainer nylon filter to remove any debris. Centrifugation at 2000 rpm for 10 minutes was performed, and supernatant was removed. Cell pellets were reconstituted with a volume of PlasmaLyte-A that was proportional to the pellet size, and 20 μL of the resuspended cell suspension were

mixed with 80 μ L of trypan blue for counting. The cell count sample was placed into a hemocytometer and evaluated using a microscope.

00275 Placenta membranes were analyzed prior to any processing to determine the initial characteristics of the membranes. Table 1 contains the average cell count per cm^2 and cell viability values for the amniotic and chorionic membranes from 32 placenta lots.

00276 The average cell count per cm^2 for the amniotic membrane was 91,381 cells with a corresponding average cell viability of 84.5%. For the chorionic membrane, the average cell count per cm^2 was 51,614 cells with a corresponding cell viability of 86.0%.

00277 These data illustrate cell numbers that are useful with certain embodiments of the present invention; e.g. a placental product comprising a chorionic membrane containing about 20,000 to about 200,000 cells/ cm^2 .

00278 Since the amniotic membrane consists of epithelial cells and stromal cells, experiments were conducted to determine the ratio of epithelial cells to stromal cells. Amniotic membranes from 3 placenta lots were analyzed. First, a 5 cm x 5 cm piece of amniotic membrane was digested with approximately 25 mL of 0.05% Trypsin-EDTA (Lonza, Walkersville, MD) at $37^\circ\text{C} \pm 2^\circ\text{C}$ in a water bath for 30 minutes. After the incubation step, epithelial cells were removed by gently scraping the cells from the membrane. After rinsing with PBS (Gibco Invitrogen, Grand Island, NY), the membrane was subsequently digested in the same manner as chorionic membrane (described above). In addition, another intact 5 cm x 5 cm piece of amniotic membrane was digested using the standard procedure (described above) to determine the total number of cells. The percentage of stromal cells was then determined by dividing the cell count from the amniotic membrane with the epithelial cells removed with the cell count from the intact membrane.

00279 Results indicate that 19% of the total cells were stromal cells. Therefore, approximately 17,362 stromal cells were present in amniotic membrane with approximately 74,019 epithelial cells. These data indicated that there are approximately 3 times more stromal cells in chorionic membranes as compared to amniotic membranes. This ratio is consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane and an

amniotic membrane, wherein the chorionic membrane comprises about 2 to about 4 times more stromal cells relative to the amniotic membrane.

Table 1 Cell count per cm² and cell viability values from fresh placental tissue from 32 donors.

Membrane	Statistics	Cell Count per cm ²	Cell Viability
Amnion	Average	91,381	84.5%
	SD	49,597	3.7%
Chorion	Average	51,614	86.0%
	SD	25,478	4.7%

00280 The second point in the manufacturing process where cell count and cell viability values were assessed was after the antibiotic treatment step. Table 2 provides the results from these analyses. Cell recoveries from this step for the amniotic membrane and the chorionic membrane were 87.7% and 70.3%, respectively.

Table 2 Cell count per cm², cell viability, and process (antibiotic treatment) cell recovery values for post antibiotic placental tissue from 28 donors.

Membrane	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery
Amnion	Average	75,230	84.4%	87.7%
	SD	46,890	4.2%	49.4%
Chorion	Average	33,028	85.6%	70.3%
	SD	18,595	4.4%	31.1%

Example 4 Development of a Placental Product Cryopreservation Procedure

00281 Cryopreservation is a method that provides a source of tissues and living cells. A main objective of cryopreservation is to minimize damage to biological materials during low temperature freezing and storage. Although general cryopreservation rules are applicable to all cells, tissues, and organs, optimization of the cryopreservation procedure is required for each type of biological material. The present application discloses a cryopreservation procedure for placental membrane products that can selectively deplete immunogenic cells from the placental membranes; and preserve viability of other beneficial cells that are the primary source of factors for the promotion of healing.

00282 During cryopreservation method development for placental membranes, the present inventors evaluated key parameters of cryopreservation including volume of cryopreservative solution, effect of tissue equilibration prior to freezing, and cooling rates for a freezing procedures.

00283 Acceptance of tissue allografts in the absence of immunosuppression will depend on the number of satellite immune cells present in the tissue.

Cryopreservation is an approach which can be utilized to reduce tissue immunogenicity. This approach is based on differential susceptibility of different cell types to freezing injury in the presence of DMSO; leukocytes are sensitive to fast cooling rates. The freezing rate of 1°C/min is considered optimal for cells and tissues including immune cells. Rapid freezing rates such as 60-100°C/min eliminate immune cells. However, this type of procedure is harmful to other tissue cells, which are desirable for preservation according to the present invention. The developed cryopreservation procedure utilized a cryopreservation medium containing 10% DMSO, which is a key component protecting cells from destruction when water forms crystals at low temperatures. The second step of cryopreservation was full equilibration of placental membrane in the cryopreservation medium, which was achieved by soaking membranes in the cryopreservation medium for 30-60 min at 4°C. This step allowed DMSO to penetrate the placental tissues. Although there are data in the literature showing that tissue equilibration prior to freezing affects survival of immune cells (Taylor & Bank, Cryobiology, 1988, 25:1), it was an unexpected finding that 30-60 min placental membrane equilibration in a DMSO-containing solution at 2-8°C selectively increases sensitivity of immune cells to freezing (in comparison to therapeutic cells) so that these type of cells are selectively depleted during the freezing process (e.g. 1°C/min freezing rate).

00284 For example, CD14+ macrophages are selectively killed relative to therapeutic cells such as hMSCs and/or fibroblasts.

00285 Temperature mapping experiments were performed to analyze the temperature profiles of potential cryopreservation conditions for the membrane products. These results are illustrated in FIG. 1. Eight (8) FP-90 cryobags were filled with either 20 mL or 50 mL of cryopreservation solution, and temperature probes were placed inside each cryobag. The first set of parameters (conditions 1 through 4 of FIG. 1a through FIG. 1d, respectively) involved a 30-minute refrigeration ($2-8^{\circ}\text{C}$) step prior to freezing ($-80^{\circ}\text{C} \pm 5^{\circ}\text{C}$). In addition, the analysis involved freezing of the cryobags either inside a Styrofoam container or on the freezer shelf. The second set of parameters (conditions 5 through 8 of FIG. 1e through FIG. 1h, respectively) involved direct freezing ($-80^{\circ}\text{C} \pm 5^{\circ}\text{C}$) of the cryobags either inside a Styrofoam container or on the freezer shelf. The results indicated that condition 6 and condition 2 exhibited the most gradual temperature decreases. Gradual temperature decreases are typically desired in order to preserve cell viability. The difference between condition 6 and condition 2 was that condition 2 included a 30-minute refrigeration step. Therefore, the decrease in temperature from the start of freezing to -4°C , where latent heat evolution upon freezing occurs, was examined further. For condition 6, the rate of cooling was approximately $-1^{\circ}\text{C}/\text{minute}$ during this period. The rate of cooling for condition 2 was approximately $-0.4^{\circ}\text{C}/\text{minute}$ during the same timeframe. Therefore, condition 2 was selected for incorporation into a non-limiting cryopreservation process since slower rates of cooling are generally desired to maintain optimal cell viability.

00286 FIG. 2 depicts the effects of cryopreservation solution volume on process (cryopreservation) cell recovery for the chorionic membrane. The analysis of the 10 mL cryopreservation solution volume involved 5 placenta lots, and the analysis of the 20 mL cryopreservation solution volume included 3 lots. For the 50 mL cryopreservation solution volume, 16 placenta lots were analyzed.

00287 As depicted in FIG. 2, the 50mL volume of cryopreservation solution volume provided superior cell recovery compared to that of the 10ml and 20ml. These data indicate that a cryopreservation medium volume of greater than 20mL such as about 50 mL or more can provide superior placental product according to the present invention.

00288 Experiments were conducted to evaluate different potential freezing conditions to maximize cell recovery after the cryopreservation process.

00289 FIG. 3 includes these results, depicting the effects of refrigeration time and freezing parameters on process (cryopreservation) cell recovery for the chorionic membrane. Three conditions were analyzed. These conditions were also linked to the temperature mapping studies. The first condition involved directly freezing the product unit on a shelf within the freezer ($-80^{\circ}\text{C} \pm 5^{\circ}\text{C}$). The second condition also contained a direct freeze, but the product unit was placed into a Styrofoam container within the freezer. The third condition included a refrigeration ($2-8^{\circ}\text{C}$) period of 30 minutes prior to the freezing step. For the amniotic membrane, 3 placenta lots were evaluated. Two (2) placenta lots were analyzed for the chorionic membrane. Results indicated that the third condition was optimal for both membrane types. As depicted in

00290 FIG. 3, a refrigeration period at least about 30 min provided the best cell recovery.

00291 Cryopreservation parameters are assessed for the amniotic and chorionic membranes and summarized in Table 3 and Table 4. The evaluation of the cell recoveries and cell viabilities from these experiments resulted in the selection of the final parameters for the manufacturing process. In addition, all average cell viability values were $\geq 70\%$.

Table 3 Post thaw cell count per cm², cell viability, and process (cryopreservation) cell recovery values for the chorionic membrane.

Parameter	Condition Tested	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery	
Refrigerate at 2-8°C for 30-60 min and freeze at -80°C $\pm$ 5°C	All conditions	Average	23,217	87.3%	102.8%	
		SD	9,155	4.1%	65.5%	
		N	27	27	27	
Dispase treatment	30 min	Average	22,354	85.7%	81.1%	No decrease in process cell recovery for the 45 min treatment. A 30-45 min range was established.
		SD	9,505	5.1%	32.4%	
		N	24	24	24	
	45 min	Average	27,125	90.6%	172.6%	
		SD	7,963	2.2%	101.2%	
		N	6	6	6	
Refrigeration time interval	30 min	Average	23,815	86.8%	102.2%	The process recovery value
		SD	9,681	5.2%	68.8%	

	60 min	N	25	25	25	was > 80% for the 60 min time interval. A 30-60 min range was established.
		Average	20,773	85.8%	84.9%	
		SD	7,356	4.7%	14.4%	
		N	5	5	5	
Thawing temperature	37°C ± 2°C water bath	Average	33,360	85.9%	114.7%	No significant difference found in process cell recovery. The room temp condition was selected for logistical reasons.
		SD	8,497	4.0%	38.1%	
		N	5	5	5	
	Room temp water bath	Average	21,298	86.8%	96.3%	
		SD	8,189	5.3%	67.2%	
		N	25	25	25	
Holding period after transfer into saline	1-15 min	Average	23,733	86.6%	100.6%	No significant difference found in process cell recovery. Membranes can be held in saline for up to 1 hr.
		SD	9,674	5.1%	67.0%	
		N	26	26	26	
	1 hr	Average	20,550	87.0%	91.4%	
		SD	6,575	4.8%	32.0%	
		N	4	4	4	
Tissue size	5 cm x 5 cm	Average	23,391	86.1%	99.6%	No decrease in process cell recovery from the 5 cm x 5 cm product to the 2 cm x 2 cm product.
		SD	8,865	5.0%	58.7%	
		N	23	23	23	
	2 cm x 2 cm	Average	23,036	88.4%	98.7%	
		SD	11,362	5.0%	81.3%	

		N	7	7	7	Both sizes were acceptable for use.
--	--	---	---	---	---	-------------------------------------

00292 Notes: cm=centimeter; min=minutes; temp=temperature; hr=hour, SD=standard deviation; N=number

Table 4 Post thaw cell count per cm², cell viability, and process (cryopreservation) cell recovery values for the amniotic membrane

Parameter	Condition Tested	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery	Comments/Conclusions
Refrigerate at 2-8°C for 30-60 min and freeze at -80°C ± 10°C	All conditions	Average	55,709	83.4%	64.2%	Overall assessment
		SD	45,210	4.4%	22.5%	
		N	32	32	32	
Refrigeration time interval	30 min	Average	52,173	83.1%	63.7%	No significant difference found in process cell recovery. A 30-60 min range was established.
		SD	39,750	4.5%	21.4%	
		N	26	26	26	
	60 min	Average	71,033	85.0%	66.5%	
		SD	66,525	3.9%	29.3%	
		N	6	6	6	
Thawing temperature	37°C ± 2°C water bath	Average	48,524	83.3%	64.0%	No significant difference found in process cell
		SD	27,804	1.7%	34.4%	
		N	7	7	7	

	Room temp water bath	Average	57,721	83.5%	64.3%	recovery. The room temp condition was selected for logistical reasons.
		SD	49,271	4.9%	19.0%	
		N	25	25	25	
Holding period after transfer into saline	1-15 min	Average	50,873	83.1%	65.0%	No significant difference found in process cell recovery. Membranes can be held in saline for up to 1 hr.
		SD	38,969	3.9%	24.2%	
		N	26	26	26	
	1 hr	Average	76,667	85.1%	61.0%	
		SD	66,565	6.2%	14.3%	
		N	6	6	6	
Tissue size	5 cm x 5 cm	Average	58,431	83.3%	62.8%	No decrease in process cell recovery from the 5 cm x 5 cm product to the 2 cm x 2 cm product. Both sizes were acceptable for use.
		SD	47,603	4.5%	21.7%	
		N	28	28	28	
	2 cm x 2 cm	Average	36,656	84.4%	73.9%	
		SD	13,175	3.4%	29.5%	
		N	4	4	4	

00293 These data are consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane containing about 20,000 to about 60,000 or to about 200,000 cells/cm².

Example 5 Qualitative Evaluation of Cell Viability by Tissue Staining

00294 The amniotic and chorionic membranes were stained using a LIVE/DEAD[®] Viability/Cytotoxicity kit (Molecular Probes Inc., Eugene, OR) to qualitatively assess cell viability. Staining was performed as per the manufacturer's protocol. Membrane segments of approximately 0.5 cm x 0.5 cm were used. Evaluation of stained membranes was performed using a fluorescent microscope. An intense uniform green fluorescence indicated the presence of live cells, and a bright red fluorescence indicated the presence of dead cells. Images of fresh amniotic and chorionic membranes as well as cryopreserved amniotic and chorionic membranes demonstrated that the manufacturing process did not alter the phenotypic characteristics of the membranes and the proportion of viable cell types (epithelial and stromal cells) in the membranes post thaw.

00295 FIG. 4 shows representative images of the live/dead staining of the epithelial layer of fresh amniotic membrane (A); epithelial layer of cryopreserved amniotic membrane (B); stromal layer of fresh amniotic membrane (C); stromal layer of cryopreserved amniotic membrane (D); fresh chorionic membrane (E); and cryopreserved chorionic membrane (F). Live cells are green, and dead cells are red.

Example 6 Placental Tissue Immunogenicity Testing

00296 One unique feature of the human chorion is the absence of fetal blood vessels that prevent mobilization of leukocytes from fetal circulation. On the fetal side, macrophages resident in the chorioamniotic mesodermal layer represent the only population of immune cells. Thus, fetal macrophages present in the chorion are a major source of tissue immunogenicity, as such the chorion is considered immunogenic. In a study where the amnion was used together with the chorion for plastic repair of conjunctival defects, the success rate was low (De Roth Arch Ophthalmol, 1940, 23: 522). Without being bound by theory, the present inventors believe that removal of CD14⁺ cells from placental membranes eliminates activation of lymphocytes *in vitro*. In addition to the presence of fetal macrophages, the present inventors believe that immunogenicity of chorion can be mediated by contamination of blood cells coming from the maternal trophoblast, which contains blood vessels. Thus, the processing of placental membrane for clinical use can be enhanced by purification of the chorion from maternal trophoblasts and selective elimination of all CD14⁺ fetal macrophages. Immunogenicity testing can be used to characterize a chorion-derived product as safe clinical therapeutics. For example, two bioassays can be used to test

immunogenicity of manufactured placental products: Mixed Lymphocyte Reaction (MLR) and Lipopolysaccharide (LPS)-induced Tumor Necrosis Factor (TNF)- α secretion.

Example 7 Mixed Lymphocyte Reaction (MLR)

00297 An MLR is a widely used *in vitro* assay to test cell and tissue immunogenicity. The assay is based on the ability of immune cells (responders) derived from one individual to recognize allogeneic Human Leukocyte Antigen (HLA) and other antigenic molecules expressed on the surface of allogeneic cells and tissues (stimulators) derived from another individual when mixed together in a well of an experimental tissue culture plate. The response of immune cells to stimulation by allogeneic cells and tissues can be measured using a variety of methods such as secretion of particular cytokines (e.g., Interleukin (IL-2), expression of certain receptors (e.g., IL-2R), or cell proliferation, all of which are characteristics of activated immune cells.

00298 Placental tissue samples representing different steps of the presently disclosed manufacturing process were used for immunogenicity testing. These samples included amnion with chorion and trophoblast as a starting material and separated choriotrophoblast, chorion, trophoblast, and amnion. Both freshly purified and cryopreserved (final products) tissues were tested.

00299 For the MLR assay, cells from placental tissues were isolated using 280 U/mL of collagenase type II (Worthington, Cat No. 4202). Tissues were treated with enzyme for 60-90 min at 37°C $\pm$ 2°C, and the resulting cell suspension was filtered through a 100 μ m filter to remove tissue debris. Single cell suspensions were then centrifuged using a Beckman, TJ-6 at 2000 rpm for 10 min and washed twice with DPBS. Supernatant was discarded after each wash, and cells were resuspended in 2 mL of DMEM (Invitrogen, Cat No. 11885) and evaluated for cell number and cell viability by counting cells in the presence of Trypan blue dye (Invitrogen, Cat No. 15250-061). For the MLR, placental-derived cells were mixed with allogeneic hPBMCs at a 1:5 ratio in 24-well culture plates in DMEM supplemented with 5% fetal bovine serum (FBS) and incubated for 4 days in the incubator containing 5% CO₂, 95% humidity at 37°C $\pm$ 2°C. Human Peripheral Blood Mononuclear Cells (hPBMCs) alone were used as a negative control, and a mixture of two sets of hPBMCs derived from two different donors was used as a positive MLR control. After 4 days of incubation, cells were collected from

wells, lysed using a lysis buffer (Sigma, Cat No. C2978) supplemented with protease inhibitor cocktail (Roche, Cat No. 11836153001), and IL-2R was measured in cell lysates using the sIL-2R ELISA kit (R&D Systems, Cat No. SR2A00) generally following the manufacturer's protocol. The level of IL-2R is a measure of activation of T-cells in response to immunogenic molecules expressed by allogeneic cells. Results of 2 out of 12 representative experiments are shown in FIG. 5 and FIG. 6. Results presented in these figures demonstrated that the present application discloses a process for manufacturing of placental membranes that result in low immunogenicity of the final chorionic membrane products.

00300 As depicted in FIG. 5, the manufacturing process serially reduces immunogenicity of the placental product. Samples representing different steps of the manufacturing process Chorion+Trophoblast (CT), Trophoblast (T), Amnion (AM), and Chorion (CM) were co-cultured with hPBMCs for 4 days. IL-2sR was measured in cell lysates as a marker of T-cell activation. Negative control shows a basal level of immune cell activation: PBMCs derived from one donor were cultured alone. Positive control: a mixture of PBMCs derived from 2 different donors.

00301 As depicted in FIG. 6, selective depletion of immunogenicity results from the present cryopreservation process of producing the present placental products, as evidenced by the significant decrease in immunogenicity upon cryopreservation.

Example 8 LPS-Induced TNF- α Secretion by Placental Membrane Cells

00302 As described herein, fetal macrophages present in the amnion and chorion are a major source of tissue immunogenicity. Without being bound by theory, the present inventors believe that removal of CD14+ cells from placental membrane eliminates activation of lymphocytes and that depletion of allogeneic donor tissue macrophages decreases the level of inflammatory cytokine secretion and tissue immunogenicity. The inventors also believe that reduction of tissue immunogenicity can also be reached by depletion of TNF- α with anti-TNF- α antibodies or suppression of TNF- α secretion by IL-10. Macrophages in fetal placental membranes respond to bacteria by secretion of inflammatory cytokines. The secretion of TNF- α by fresh placental membranes *in vitro* in response to bacterial LPS is significantly higher in the chorionic membrane. Thus, the present inventors believe that immunogenicity of placental membranes is mediated by macrophages, the amount and/or activity of which is higher in the chorionic membrane.

00303 According to the present invention, selective depletion of macrophages is an optional approach to selectively deplete immunogenicity of the amniotic and chorionic membranes, allowing the use of both allogeneic membranes for clinical applications. The assay of functional macrophages in a placental product is used here as an assay for immunogenicity testing (e.g. in production or prior to clinical use) based on the facts that: macrophages are the source of immunogenicity in chorionic membranes. Macrophages in placenta-derived membranes respond to bacterial LPS by secretion of high levels of TNF- α ; and TNF- α is a critical cytokine involved in immune response and allograft tissue rejection. Therefore, secretion of TNF- α by placenta-derived membranes in response to LPS is used here to characterize tissue immunogenicity and for pre-use screening.

Example 9 Establishment of Allowed LPS-Induced TNF- α Secretion Level by Chorionic Membranes

00304 Data from published reports regarding the level of TNF- α , which is associated with the absence or an insignificant immune response in a variety of experimental systems, are presented in Table 5. These data indicate that a TNF- α level below 100 pg/mL correlates with a low immune response. The ability of amniotic and chorionic membranes to produce TNF- α spontaneously and in response to bacteria or bacterial LPS *in vitro* has been shown by a number of investigators. Table 6 summarizes such data. The lowest spontaneous TNF- α secretion by amniotic membrane of about 70 pg/cm² of the membrane was reported by Fortunato et al. (Am J Reprod Immunol, 1994, 32:184). All reports also showed that fresh placental membranes secrete large amounts of TNF- α in response to bacteria or bacterial LPS (Table 6), which is attributed to the presence of viable functional macrophages.

Table 5

Description of experimental system	TNF-α levels associated with the absence/reduction of immune response	Comments	References
IL-10-induced inhibition of MLR <i>in vitro</i> . TNF was measured in	Mean 260 pg/mL		Wang et al., Transplantation, 2002, 74:772

tissue culture supernatant by ELISA.			
MLR using skin tissue explants (0.02 cm ² per well) as stimulators in the presence or absence of IL-10 (skin explant assay). Skin tissue destruction was assessed microscopically, and severity was assigned based on histopathological tissue damage.	Mean 100 pg/mL		
Endogeneous TNF production in MLR in the presence or absence of anti-TNF antibodies. TNF levels were assessed using the WEHI-164 cytotoxicity assay.	~0.04 U/mL for the negative control and MLR in the presence of anti-TNF antibodies, which correlated with no or significant inhibition of lymphocyte proliferation	TNF activity per mg is not provided.	Shalaby et al., J Immunol, 1988, 141:499
TNF levels in BAL fluid of lung isografts, unmodified allograft, and alveolar macrophages (AM) depleted allograft in rats.	Isograft: below detection; AM-depleted allograft: ~15 pg/mL of BAL (total 75 pg/ 5 ml of BAL)	Unmodified allograft: ~45 pg/mL (immunogenic)	Sekine et al., J Immunol, 1997, 159:4084

TNF levels in MLR after 48 hours in the presence or absence of advanced glycation end products (MLR inhibitors).	~<200 pg/mL TNF correlated with a complete inhibition of MLR		Ohashi et al, Clin Immunol, 2010, 134:345
TNF levels in MLR.	<100 pg/mL TNF in MLR with HLA-matched donors (control, no stimulation)		Toungouz et al., Hum Immunol, 1993, 38:221
TNF activity in MLR when pieces of cryopreserved skin allografts (~0.2 cm ²) were incubated with hPBMCs for 24 hours. Positive control: hPBMC+LPS; negative: hPBMC alone.	Negative control ~20 U of TNF activity; MLR with skin explants: 0-40 U; Positive control: 600 U	Unit of activity was calculated as TNF in ng/mL divided by OD at 570 nm for the same experimental well	Lomas et al., Cell Tissue Bank, 2004, 5:23.
Cytokine time course in MLR, including TNF.		Optimal TNF after 24 hours: ~150 pg/mL	Jordan & Ritter, J Immunol Meth, 2002, 260:1

MLR using skin tissue explants (0.02 cm ² per well) as stimulators in the presence or absence of anti-TNF antibodies (skin explant assay). Skin tissue destruction was assessed microscopically, and severity was assigned based on histopathological tissue damage.	For no skin destruction: 0.5-1.1 pg/mL for HLA compatible responders, and 2.6-1376 pg/mL for unmatched MLR	Recalculation per 1 cm ² of skin tissue: lowest TNF non-immunogenic level is 100 pg/cm ²	Dickinson et al., Cytokine, 1994, 6:141
---	--	--	---

Table 6

Description of experimental system	TNF levels secreted by fresh placental membranes in culture	Comments/ recalculations of the lowest TNF levels per cm²	References
TNF secretion by "fresh" amnion and chorion tissues (1.44 cm ²) incubated for 24 hours in the presence or absence of LPS (500 ng/mL).	Chorion: basal 3.3±0.46 ng/cm ² , LPS-induced: 150-250 ng/cm ² Amnion: basal 2.5±1.3 ng/cm ² , LPS-induced: ~50 ng/cm ²	Lowest TNF level for amnion is 1200 pg/cm ²	Zaga et al., Biol Reprod, 2004, 71:1296

TNF secretion by "fresh" amnion and chorion tissues (1.8 cm diameter disks: 2.5 cm ²) incubated for 24 hours in the presence or absence of <i>E. Coli</i> in 1 mL medium.	Basal ~1-2.5 pg/μg total protein in the medium for both amnion and chorion; <i>E. Coli</i> -induced: amnion → 29.2 (14.5-35.3) pg and chorion → 53.15 (40-94.2) pg per μg total protein	Lowest TNF level for amnion is 800 pg/cm ²	Zaga-Clavellina et al., Reprod Biol Endocrinol, 2007, 5:46
TNF secretion by "fresh" amnion and chorion tissues (chorion 8-10 mg tissue/mL; amnion 5-7 mg/mL, 0.02-0.04cm ²) incubated for 20 hours in the presence or absence of LPS (5 μg/mL).	Basal: ~ 2-64 U/mL or 8-10 mg chorion; <1U/mL for 5-7 mg amnion; LPS-induced: >100 U/10mg for chorion and ~15-17 U/10 mg for amnion	1 unit=~100-200 pg/mL; Lowest TNF level for amnion is <100 pg/mL corresponding to <2500 pg/cm ²	Paradowska et al., Placenta, 1997, 18:441
TNF secretion by "fresh" amnion (0.57 cm ²) in 0.8 mL incubated for 24 hours in the presence or absence of LPS (50 ng/mL).	Amnion: Basal → 40 pg/mL, LPS-induced → 410 pg/mL	Lowest TNF level for fresh amnion is ~ 70 pg/cm²	Fortunato et al., Am J Obstet Gynecol, 1996, 174:1855

TNF secretion by "fresh" amnion and chorion tissues (4 cm ²) incubated for 24 hours in the presence or absence of LPS (1-1000 ng/mL)	Basal: Amnion ~7-13 ng/mL/g tissue); Chorion ~18 ng/mL/g tissue LPS-induced (1000 ng/mL): Amnion ~14 ng/mL/g), Chorion ~27 ng/mL/g	Amnion is 5-7 mg corresponds ~0.02-0.04 cm ² ; 1 g is ~6 cm ² ; Lowest TNF level for amnion is ~1000 pg/cm ²	Thiex et al., Reprod Biol Endocrinol, 2009, 7:117
--	---	---	---

Example 10 LPS-Induced TNF- α Secretion Immunogenicity Assay

00305 2 cm x 2 cm pieces of placental derived membranes representing production intermediates and final placental products were placed in tissue culture medium and exposed to bacterial LPS (1 μ g/mL) for 20-24 hr. After 24 hours, tissue culture supernatant were collected and tested for the presence of TNF- α using a TNF- α ELISA kit (R&D Systems) according to the manufacturer's protocol. Human hPBMCs (SeraCare) known to contain monocytes responding to LPS by secretion of high levels of TNF- α were used as a positive control in the assay. hPBMCs and placental tissues without LPS were also included as controls in the analysis. In this assay, TNF detected in the culture medium from greater than 70 pg/cm² (corresponding to 280 pg/mL) for both spontaneous and LPS-induced TNF- α secretion was considered immunogenic.

00306 The low levels of TNF- α and the absence of the response to LPS by AM and CM indicates the absence of viable functional macrophages that are the major source of immunogenicity for amniotic and chorionic membranes. Results of this assay showed a correlation with the MLR data: tissues that produce high levels of TNF- α in response to LPS are immunogenic in the MLR assay (FIG. 7A and FIG. 7B for TNF- α secretion; Figure 9, C – MLR).

00307 As depicted in FIG. 7A and FIG. 7B, the manufacturing process serially reduces immunogenicity of the placental product. Samples representing different steps of the manufacturing process (Amnion+Chorion+Trophoblast (ACT),

Chorion+Trophoblast (CT), Amnion (AM), and Chorion (CM)) were incubated in the presence of LPS for 24 hr, and after that tissue culture supernatants were tested for the TNF- α by ELISA. Tissues cultured in medium without LPS show the basal level of TNF α secretion. PBMCs, which are known to secrete high levels of TNF, were used as a positive control.

00308 Choriotrophoblast (CT), which secreted high levels of TNF- α (FIG. 7 B), was tested in MLR against two different PBMC donors. CT cells were co-cultured with PBMCs for 4 days. IL-2 α R was measured in cell lysates as a marker of T-cell activation. Positive control: a mixture of PBMCs derived from 2 different donors.

00309 FIG. 7C shows that preparations producing high levels of TNF- α are immunogenic. Choriotrophoblast (CT), which secreted high levels of TNF- α (FIG. 7, B), was tested in MLR against two different PBMC donors. CT cells were co-cultured with PBMCs for 4 days. IL-2 α R was measured in cell lysates as a marker of T-cell activation. Positive control: a mixture of PBMCs derived from 2 different donors.

00310

Example 11 Analysis of Placental Cells by FACS

00311 Knowing the cellular composition of chorionic membranes is important for developing a thorough understanding of potential functional roles in wound healing and immunogenicity. Previous reports demonstrated that the chorion contains multiple cell types. In addition to fibroblasts, stromal cells were identified in the chorion. Although there are no fetal blood vessels within the chorionic membranes, it comprises resident fetal macrophages. The close proximity to maternal blood circulation and decidua provide a potential source of immunogenic cells (maternal leukocytes and trophoblast cells) and therefore are a potential source of immunogenicity. To investigate the cellular composition of the chorion, FACS analysis was performed.

Example 11.1 FACS Procedure: Single Cell Suspension Preparation

00312 Purified chorionic membranes were used for cellular phenotypic analysis via FACS. Cells from chorion were isolated using 280 U/mL collagenase type II (Worthington, Cat No. 4202). Tissues were treated with enzyme for 60-90 min at 37°C $\pm$ 2°C, and the resulting cell suspension was filtered through a 100 μ m filter to

remove tissue debris. Single cell suspensions were then centrifuged using a Beckman TJ-6 at 2000 rpm for 10 min and washed twice with DPBS. Supernatant was discarded after each wash, and cells were resuspended in 2 mL of FACS staining buffer (DPBS + 0.09% NaN₃ + 1% FBS).

Example 11.2 Immunolabeling Cells for Specific Cellular Markers

00313 Once the single cell suspension was prepared according to Example 10.1, a minimum of 1×10^5 cells in 100 μ L of FACS staining buffer was treated with antibodies labeled with fluorescent dye. Table 7 provides descriptions of the antibodies and the amounts used. For cell surface markers, cells were incubated for 30 min at room temperature in the dark with antibodies followed by washing twice with FACS staining buffer by centrifugation at 1300 rpm for 5 min using a Beckman TJ-6 centrifuge. Cells were then resuspended in 400 μ L of FACS staining buffer and analyzed using a BD FACSCalibur flow cytometer. To assess cell viability, 10 μ L of 7-AAD reagent (BD, Cat No. 559925) was added just after the initial FACS analysis and analyzed again. For intracellular staining, cells were permeabilized and labeled following the manufacturer's recommendations (BD Cytotfix/Cytoperm, Cat No. 554714) and analyzed using a BD FACSCalibur flow cytometer.

Table 7 Description of reagents used for placental cell characterization by FACS.

Cell marker antibody and label type	Cat No.	Volume of antibody solution used	Cell marker type	Cell marker specificity
IgG1 isotype-PE	BD 559320	5 μ L	Cell surface	Isotype control
CD105-PE	Caltag MHCD10504	20 μ L	Cell surface	MSC marker
CD166-PE	BD 559263	80 μ L	Cell surface	MSC marker
CD45-PE	BD 555483	10 μ L	Cell surface	Hematopoietic cell marker
IgG2a isotype-PE	BD 555574	2 μ L	Cell surface	Isotype control

CD14-PE	BD 555398	20 µL	Cell surface	Monocyte marker
HLA-DR-PE	BD 556644	20 µL	Cell surface	HLA class II specific for antigen- presenting cells
IgG1 isotype- FITC	BD555748	5 µL	Cell surface	Isotype control
CD86-FITC	BD 557343	20 µL	Cell surface	Immune co- stimulatory marker
CD40-FITC	BD 556624	20 µL	Cell surface	Immune co- stimulatory marker
IgG1 isotype- unlabeled	Dako X0931	10 µL	Intracellular	Isotype control
Cytokeratin 7- unlabeled	Dako M7018	2 µL	Intracellular	Trophoblast marker
Rabbit anti- mouse FITC	Dako F0261	5 µL	Intracellular	Secondary antibody

00314

Example 12 Phenotypic Analysis of Placental Cells

00315 FACS analysis of single cell suspensions of chorionic membranes demonstrates that both membranes contain cells expressing markers specific for mesenchymal stem cells (refer to Table 8), implicating the presence of stromal cells. In addition, several immunogenic markers, which are more likely expressed on CD14+ placental macrophages, were detected. The % ranges for different markers are wide. It can be explained by: 1) high variability in cell number between placenta donors; and 2) technical issues, which include the presence of the high and variable cellular and tissue debris in the cellular suspension. Although debris can be gated out, debris particles that are comparable with cells by size will affect the accuracy of the

calculated % for each tested marker. In addition, Table 9 provides a FACS analysis of cells from the chorionic membranes that were cultured in 10% FBS in DMEM at 37°C±2°C until confluency (passage 0 cells). These data demonstrated that cells derived from chorionic membranes retained a phenotype similar to MSCs after culturing. In conclusion, the presence of stromal cells in placental tissues was confirmed by FACS analysis.

00316 These data are consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane containing MSCs.

Table 8 Characterization of the cellular composition of placental membranes based on selective CD markers.

	Marker	Chorion (% range)
MSC Markers	CD105	6.4-78.5
	CD166	4.8-51.5
Hematopoietic Cell Markers	CD14	0.9-6.1
	CD45	4.6-14.7
Immune co-stimulatory markers	HLA-DR	0-14.7
	CD86	4.9-22.5
	CD40	2-5.8
Trophoblast marker	Cytokeratin-7	2.71-23.07

00317

Table 9 FACS analysis of cultured cells (passage 0) from placenta lot D16.

Cell Surface Marker	Chorion (%)
CD45	0.53
CD166	82.62
CD105	86.73

CD49a	92.26
CD73	94.57
CD41a	-0.05
CD34	-0.25
HLA-DR	-0.19
CD19	-0.22
CD14	-0.27
CD90	98.00

Example 13 Differentiation Capacity of Cells Derived from the Chorionic Membrane

00318 Therapeutic cells, in optional embodiments of the present invention, are adherent, express specific cellular markers such as CD105 and lack expression of other markers such as CD45, and demonstrate the ability to differentiate into adipocytes, osteoblasts, and chondroblasts.

00319 The expression of specific cellular markers has already been described in Example 12. To determine if the cells within the placental product derived from the chorionic membrane can adhere to plastic and differentiate into one of the lineages, cells were isolated from the placental product derived from the chorion as described in this invention and cultured at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ and expanded.

00320 FIG. 8-A shows a representative image of passage 2 cells, demonstrating the ability of the cells to adhere to tissue culture plastic. As a comparison, a representative image of MSCs isolated and expanded from human bone marrow aspirate is shown in FIG. 8-B.

00321 Osteogenic differentiation capacity was demonstrated by staining the cultured cells with alkaline phosphatase labeling following the manufacturer's recommendations (BCIP/NBT Alkaline Phosphatase Substrate Kit IV, Vector Laboratories Cat. No. SK-5400). Alkaline phosphatase is an enzyme involved in bone mineralization (Allori et al., Tissue Engineering: Part B, 2008, 8:275), and its

expression within cells is indicative of osteo-precursor cells (Majors et al., J Orthopaedic Res, 1997, 15:546). Staining for alkaline phosphatase is carried out through an enzymatic reaction with Bromo-4-Chloro-3'-Indolylphosphate p-Toluidine Salt (BCIP) and Nitro-Blue Tetrazolium Chloride (NTP). BCIP is hydrolyzed by alkaline phosphatase to form an intermediate that undergoes dimerization to produce an indigo dye. The NBT is reduced to the NBT-formazan by the two reducing equivalents generated by the dimerization. Together these reactions produce an intense, insoluble black-purple precipitate when reacted with alkaline phosphatase. FIG. 8-C shows a representative image of passage 2 cells staining positively for alkaline phosphatase.

Example 14 Live CD45+ FACS Analysis

00322 As CD45 is a general marker for hematopoietic cells and therefore a marker for the presence immunogenic cells, the presence of CD45+ cells may correlate well with how immunogenic a tissue may be. An initial study indeed showed a correlation between amount of immunogenicity as measured via an *in vitro* MLR assay of placental tissue at various stages within the manufacturing process (as described previously), and the amount of CD45+ cells was determined via FACS analysis. As FIG. 9 demonstrates, membranes that trigger the expression of high levels of IL-2sR on hPBMC responders in MLR also contained a high percentage of CD45+ cells, indicating that immunogenicity of placental membranes can be correlated with the number of CD45+ cells. Further studies revealed, however, that quantifying CD45+ cells via FACS alone showed high variability that did not allow for the establishment of a safety threshold for CD45+ cells in placental membranes. Accordingly, the inventors evaluated whether or not viability of CD45+ cells is correlated with immunogenicity.

00323 To eliminate some of the variability in CD45+ measurements via FACS, viability of CD45+ cells was assessed, as dead CD45+ cells do not contribute to immunogenicity. To ensure an accurate assessment of live CD45+ cells, a pilot experiment was conducted in which a single cell suspension of amnion membrane was spiked in with a known concentration of live CD45+ cells (hPBMCs) ranging from a theoretical 1.25% to 20% (0.75-12% - actual % of the spiked cells) of the total cell concentration in suspension. Cells were stained with CD45-PE antibody at determined concentrations (refer to Table 10), incubated with 7-AAD cell viability test reagent, and analyzed using a BD FACSCalibur. Table 10 demonstrates that recovery of known amounts of CD45+ cells was not correct (4th column in the table). For example,

although 12% of PBMCs was spiked into a single-cell suspension of amnion membrane, only 4.26% of CD45+ cells were recovered according to FACS analysis (>60% difference from the actual spike). To correlate with immunogenicity, MLR was also performed in parallel. Briefly, single cell suspensions of amniotic membrane spiked with various amounts of live hPBMCs were co-cultured with another donor of PBMCs in the MLR. FIG. 10 depicts a correlation between the amount of CD45+ cells present in amnion-derived cell suspensions and immunogenicity in MLR in vitro. Table 10 and FIG. 10 show that the suspensions spiked with higher amounts of live CD45+ cells resulted in higher immunogenicity as measured by IL-2sR expression on the hPBMC responder donor.

Table 10 %CD45+ recovery experiments.

Sample Description (in % of cell types in the mixture)	% CD45+ cells (detected by FACS)	Actual spike (%, based on 60% CD45+ cells in this hPBMC batch)	% Difference from actual spike	Cell suspension immunogenicity (tested in MLR and expressed as IL-2R in pg/mL)
100% amnion	0.65	N/A	N/A	20.23
0% PBMC	N/A	N/A	N/A	15.6 (negative control)
100% PBMC	61.51	N/A	N/A	86.31 (positive control)
20% PBMC + 80% Amnion	4.26	12	64.5%	24.38
10% PBMC + 90% Amnion	2.24	6	62.7%	21.17
5% PBMC + 95% Amnion	1.7	3	43.3%	16.75

2.5% PBMC +97.5% Amnion	1.36	1.5	Not calculated*	15.9
1.25% PBMC + 98.75% Amnion	1.06	0.75	Not calculated*	12.27

00324 Notes: N/A- not applicable; *Not calculated – values are close to the method detection limits.

Example 15 Protein Array Analyses

00325 The protein profiles of amniotic and chorionic membranes were investigated using a SearchLight Multiplex chemiluminescence array. The presence of proteins in tissue membrane extracts and secreted by tissues in culture medium was investigated. For comparison, two commercially available products containing living cells, Apligraf and Dermagraft, were assayed.

Example 15.1 Dermagraft

00326 Dermagraft membrane was thawed and washed according to the manufacturer's instructions. Dermagraft membrane was cut into 7.5 cm² pieces. For tissue lysates, one 7.5 cm² piece of membrane was snap frozen in liquid nitrogen followed by pulverization using a mortar and pestle. Crushed tissue was transferred to a 1.5 mL microcentrifuge tube and 500 µL of Lysis buffer (Cell Signaling Technologies, Cat No. 9803) with protease inhibitor (Roche, Cat No. 11836153001) was added and incubated on ice for 30 min with frequent vortexing. The sample was then centrifuged at 16000 g for 10 min. The supernatant was collected and sent for protein array analysis by Aushon Biosystems. For tissue culture, one 7.5 cm² piece of membrane was plated onto a well of a 12-well dish and 2 mL of DMEM +1% HSA+ antibiotic/antimycotic were added and incubated at 37°C±2°C for 3, 7, or 14 days. After incubation, tissue and culture media were transferred to a 15 mL conical tube and centrifuged at 2000 rpm for 5 min. Culture supernatant was collected and sent for protein array analysis by Aushon Biosystems.

Example 15.2 Apligraf

00327 Apligraf membrane was cut into 7.3 cm² pieces. For tissue lysates, one 7.3 cm² piece of membrane was snap frozen in liquid nitrogen followed by pulverization using a mortar and pestle. Crushed tissue was transferred to a 1.5 mL microcentrifuge tube and 500 µL of Lysis buffer (Cell Signaling Technologies, Cat No. 9803) with protease inhibitor (Roche, Cat No. 11836153001) was added and incubated on ice for 30 min with frequent vortexing. The sample was then centrifuged at 16000 g for 10 min. The supernatant was collected and sent for protein array analysis by Aushon Biosystems. For tissue culture, one 7.3 cm² piece of membrane was plated onto a well of a 12-well dish and 2 mL of DMEM +1% HSA+ antibiotic/antimycotic were added and incubated at 37°C±2°C for 3, 7, or 14 days. After incubation, tissue and culture media were transferred to a 15 mL conical tube and centrifuged at 2000 rpm for 5 min. Culture supernatant was collected and sent for protein array analysis by Aushon Biosystems.

Example 15.3 Chorionic Membranes

00328 Chorionic membranes were isolated and packaged at -80°C ±5°C according to the manufacturing protocols disclosed herein in Example 2. Packaged membranes were then thawed in a 37°C±2°C water bath and washed 3 times with DPBS. Membranes were cut into 8 cm² pieces. For tissue lysates, one 8 cm² piece of membrane was snap frozen in liquid nitrogen followed by pulverization using a mortar and pestle. Crushed tissue was transferred to a 1.5 mL microcentrifuge tube and 500 µL of Lysis buffer (Cell Signaling Technologies, Cat No. 9803) with protease inhibitor (Roche, Cat No. 11836153001) was added and incubated on ice for 30 min with frequent vortexing. Tissue lysate was then centrifuged at 16000 g for 10 min. The supernatant was collected and sent for protein array analysis by Aushon Biosystems. For tissue culture, one 8 cm² piece of membrane was plated onto a well of a 12-well dish and 2 mL of DMEM +1% HSA+ antibiotic/antimycotic were added and incubated at 37°C±2°C for 3, 7, or 14 days. After incubation, tissue and culture media were transferred to a 15 mL conical tube and centrifuged at 2000 rpm for 5 min. Culture supernatant was collected and sent for protein array analysis by Aushon Biosystems.

00329 Initial testing consisted of an analysis of 36 proteins that are important for wound healing. The list of identified proteins is described in Table 11.

Table 11 List of selected proteins for analysis.

Protein Group Based on Functionality		Comments
Metalloproteases	Matrix Metalloproteinase 1 (MMP1), MMP2,3,7,8,9,10,13	Matrix and growth factor degradation; facilitate cell migration.
MMP Inhibitors	Tissue Inhibitors of MMPs (TIMP1 and 2)	Have angiogenic activity; can be placed in the “angiogenic factors” group.
Angiogenic Factors	Angiotensin-2 (Ang-2); basic Fibroblast Growth Factor (bFGF); heparin-bound Epidermal Growth Factor (HB-EGF); EGF; FGF-7 (also known as Keratinocyte Growth Factor-KGF); Platelet derived Growth Factors (PDGF) AA, AB, and BB; Vascular Endothelial Growth Factor (VEGF), VEGF-C and VEGF-D; Neutrophil gelatinase-associated lipocalin (NGAL); Hepatocyte Growth Factor (HGF); Placenta Growth Factor (PIGF); Pigment Epithelium Derived Factor (PEGF); Thrombopoietin (TPO)	Majority of these factors also have growth and migration stimulatory activities and can be placed in a group of growth factors.
Protease Inhibitor/ Protein Carrier	Alpha-2-macroglobulin	Inhibit protease activity; regulate growth factor activity.

Growth Factors	See “angiogenic factors” + Transforming Growth Factor alpha (TGF-a)	See “angiogenic factors.”
Cytokines	Adiponectin (Acrp-30)	Affect keratinocyte functions.
	Granulocyte Colony-Stimulating Factor (G-CSF)	Protection from infections.
	Interleukin1 Receptor Antagonist (IL-1 RA)	Regulate activity of inflammatory cytokine IL-1.
	Leukemia Inhibitory Factor (LIF)	Support angiogenic growth factors.
Chemokines	SDF-1beta	Attracts endothelial and other stem cells from circulation to wound site.
Regulators of IGF	Insulin-like growth factor binding protein (IGFBP1,2,3)	Regulate IGF activity.

00330

Example 15.4 Protein Expression in Present Placental Products

00331 Preliminary protein array data analyses showed that the majority of selected testing factors (see Table 11) were expressed in amniotic membrane, chorionic membrane, Apligraf, and Dermagraft.

00332 Three proteins were identified as unique for the chorionic membrane which are undetectable in Apligraf and Dermagraft. These proteins are EGF, IGFBP1, and Adiponectin). FIG. 11 depicts expression of EGF (A), IGFBP1 (B), and Adiponectin (C) in amniotic or chorionic membranes. CM75 and CM 78 are placental products of the present invention (e.g. cryopreserved), AM75 and AM78 are cryopreserved amniotic membrane products. These proteins are believed by the inventors to facilitate the therapeutic efficacy of the present placental products for wound healing.

00333 These data are consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane containing EGF, IGFBP1, and/or adiponectin.

Example 16 Wound Healing Proteins are Secreted for a Minimum of 14 days

00334 Placental products of the present invention demonstrate a durable effect, desirable for wound healing treatments. The extracellular matrix and presence of viable cells within the amniotic membrane described in this invention allow for a cocktail of proteins that are known to be important for wound healing to be present for at least 14 days. Amniotic membranes were thawed and plated onto tissue culture wells and incubated at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 3, 7, and 14 days. At each time point, a sample of the culture supernatant was collected and measured through protein array analysis as described in Example 15. Table 12 illustrates the level of various secreted factors in tissue culture supernatants from two donors of chorionic membranes at 3, 7 and 14 days as measured through protein array analysis.

Table 12 Levels of proteins secreted in chorion tissue culture supernatants at different time points (pg/ml).

	Day 3	Day 7
hACRP30	298.4	614.3
hAlpha2Macroglobulin	34,480.5	6,952.5
hANG2	0.0	2.0
hEGF	0.7	0.4
hFGF	84.3	13.5
hFibronectin	37,510.9	41,871.4
hHBEGF	102.6	40.0
hHGF	1,382.9	1,715.4
hIGFBP1	201.6	201.0
hIGFBP2	62.7	172.9
hIGFBP3	778.1	812.4
hIL1ra	30,037.4	556.1
hKGF	4.2	2.4
hMMP1	32,388.5	67,665.6
hMMP10	4,016.4	4,140.1
hMMP13	13.3	0.0
hMMP2	768.8	1,230.5
hMMP3	1,294.7	2,646.0
hMMP7	14.7	43.7

hMMP8	95.9	249.4
hMMP9	10,034.6	29,201.5
hNGAL	1,968.1	2,608.9
hPDGFAA	18.6	21.8
hPDGFAB	6.2	55.5
hPDGFBB	15.1	5.2
hPEDF	9,216.2	576,962.0
hSDF1b	85.9	15.3
hTGFA	0.0	0.0
hTGFB1	377.5	410.9
hTGFB2	11.2	20.7
hTIMP1	12,279.0	15,562.7
hTIMP2	216.7	419.6
hTSP1	223.1	0.0
hTSP2	42.7	210.7
hVEGF	53.1	45.9
hVEGFC	197.4	182.7

Example 17 Interferon 2 α (IFN-2 α) and Transforming Growth Factor- β 3 (TGF- β 3)

00335 Placental products described in this invention have been analyzed for the presence of IFN-2 α and TGF- β 3. Briefly, after thawing, the membranes were homogenized and centrifuged at 16,000g to collect the resulting supernatants. Supernatants were analyzed on a commercially available ELISA kit from MabTech (IFN-2 α) and R&D Systems (TGF- β 3).

00336 FIG. 12 shows significant expression of IFN-2 α (A) and TGF- β 3 (B) in cellular chorionic membrane homogenates.

00337 Without being bound by theory, interferon-2 α and TGF- β 3 may aid in the prevention of scar and contracture formation. IFN-2 α may serve a role to decrease collagen and fibronectin synthesis and fibroblast-mediated wound contracture.

Example 18 Tissue Reporative Proteins in Chorionic Membranes

00338 Chorionic membrane homogenates were analyzed for the presence of proteins that are important in tissue repair.

00339 Chorionic membranes described in this invention have been analyzed for the presence of tissue reparative proteins. Briefly, amniotic membranes were incubated in DMEM + 10% FBS for 72 hrs. The membranes were then homogenized in a bead

homogenizer with the culture media. The homogenates were centrifuged, and the supernatants were analyzed on commercially available ELISA kits from R&D Systems. **Error! Not a valid bookmark self-reference.** shows significant expression of BMP-2, BMP-4, PLAB, PIGF, and IGF-1 in several donors of chorionic membranes.

00340 Without being bound by theory, the inventors believe that efficacy of the present placental products for wound repair are due, in part, to the role of BMPs, IGF-1, and PIGF in the development and homeostasis of various tissues by regulating key cellular processes. BMP-2 and BMP-4 may stimulate differentiation of MSCs to osteoblasts in addition to promote cell growth; placental BMP or PLAB is a novel member of the BMP family that is suggested to mediate embryonic development. Insulin-like growth factor 1 (IGF-1) may promotes proliferation and differentiation of osteoprogenitor cells. Placental derived growth factor (PIGF) may acts as a mitogen for osteoblasts.

Example 19 MMPs and TIMPs

00341 Both MMPs and TIMPs are among the factors that are important for wound healing. However, expression of these proteins must be highly regulated and coordinated. Excess of MMPs versus TIMPS is a marker of poor chronic wound healing. We investigated expression of MMPs and TIMPs and its ratio in amniotic membrane and chorionic membrane and compared it to the expression profile in Apligraf and Dermagraft.

00342 Results showed that all membranes express MMPs and TIMPs; the ratio in the thawed placental products and amniotic membranes is significantly lower. Therefore, the placental products (optionally including chorionic membranes) will be more beneficial for wound healing.

00343 Accumulated data indicate that the MMP to TIMP ratio is higher in cases of non-healing wounds. For example, the ratio between MMP-9 and TIMP1 is approximately 7-10 to one or good healing and 18-20 to one or higher for poor healing.

00344 As shown in FIG. 14, analysis of the ratio between MMPs and TIMPs secreted by placental tissues, Apligraf, and Dermagraft showed that the chorionic membrane products contain MMPs and TIMPs at an approximate ratio of 7, which is favorable for wound healing. In contrast, Dermagraft had a ratio > 20, and Apligraf had a ratio >200.

00345 These data are consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane containing MMP-9 and TIMP1 at a ratio of about 7-10 to one.

Example 20 α 2-Macroglobulin

00346 α 2-macroglobulin is known as a plasma protein that inactivates proteinases from all 4 mechanistic classes. Another important function of this protein is to serve as a reservoir for cytokines and growth factors, examples of which include TGF, PDGF, and FGF. In the chronic wounds like diabetic ulcers or venous ulcers, the presence of high amount of proteases leads to rapid degradation of growth factors and delays in wound healing. Thus, the presence of α 2-macroglobulin in products designed for chronic wound healing will be beneficial. Results of the protein array analysis showed that amniotic and chorionic membranes contain α 2-macroglobulin (Table 13).

00347 These data are consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane containing α 2-macroglobulin.

Table 13 Expression of α 2-macroglobulin in placental tissue protein extracts.

Sample	α 2-macroglobulin (pg/mL/cm ²)
AM75	7
CM75	790
AM78	53042
CM78	1014

Example 21 Establishment of bFGF as a Marker for Chorionic Tissue Potency

00348 bFGF modulates a variety of cellular processes including angiogenesis, tissue repair, and wound healing (Presta et al., 2005, Reuss et al., 2003, and Su et al., 2008). In wound healing models, bFGF has been shown to increase wound closure and enhance vessel formation at the site of the wound (Greenhalgh et al., 1990). Evaluation of proteins derived from chorionic membranes prepared pursuant to the

presently disclosed manufacturing process revealed that bFGF is one of the major factors in placental tissue protein extracts (FIG. 15). FIG. 15 depicts expression of bFGF by amniotic membranes (AM) and chorionic membranes (CM) detected during the protein profile evaluation of placental membranes.

00349 The importance of bFGF for wound healing supports selection of bFGF as a potency marker for evaluation of chorionic membrane products manufactured for clinical use pursuant to the present disclosure. A commercially available ELISA kit from R&D Systems was selected for evaluation of its suitability to measure bFGF secreted by placental membranes. ELISA method qualification experiments were designed according to FDA and ICH guidances for bioanalytical assay validation (Validation of Analytical Procedures: Text and Methodology Q2 (R1), 1994; ICH Harmonized Tripartite Guideline and Guidance for Industry Bioanalytical Method Validation, 2001).

00350 The ELISA procedure was performed according to the manufacturer's instructions (bFGF ELISA brochure). The evaluation of the kit was performed prior to measurement of bFGF in placental tissue samples. Assay performance was assessed by analyzing linearity, range, lower and upper limits of quantitation (LLOQ and ULOQ), precision, and accuracy. Experimental data suggested that the quantitation range of this assay was 40-1280 pg/mL bFGF. The intra- and inter-assay CVs ranged from 2.42 to 6.23% and 0.59 to 7.02%, respectively. Additionally, sample recovery analysis demonstrated accuracy within 20%. This assay showed dilutional linearity and specificity. Ruggedness was demonstrated by assay insensitivity to variations introduced by different analysts. The analytical performance of the bFGF ELISA indicated that this assay was suitable for the measurement of bFGF secreted by placental membranes. bFGF ELISA parameters are summarized in Table 14.

Table 14 Established ELISA parameters for measuring bFGF in placenta homogenates.

Calibration Standard Range	20-1280 pg/mL
Assay Quantitation Range	40-1280 pg/mL

LLOQ	40 pg/mL
LOD	20 pg/mL
ULOQ	1280 pg/mL

bFGF Expression in Chorionic Membranes

00351 Measurement of bFGF in chorionic membrane preparations has proven to be both reliable and reproducible. The placental tissue homogenates were prepared using the “bead” method as described above. Also, secretion of bFGF in tissue culture media was evaluated. Measurement of bFGF in multiple donors showed that this method of quantification was a valuable means of evaluating potency the presently disclosed tissue products prepared for use in a clinical setting. FIG. 16 shows representative expression of bFGF in chorionic tissue samples derived from two separate placenta donors. Results have been reproduced in multiple tissue preparations.

00352 These data are consistent with certain embodiments of the present invention that provide a placental product comprising a chorionic membrane containing bFGF.

Example 22 Placental Tissues Enhance Cell Migration and Wound Healing

00353 The process of wound healing is highly complex and involves a series of structured events controlled by growth factors (Goldman, 2004). These events include increased vascularization, infiltration by inflammatory immune cells, and increases in cell proliferation. The beginning stages of wound healing revolve around the ability of individual cells to polarize towards the wound and migrate into the wounded area in order to close the wound area and rebuild the surrounding tissue. Upon proper stimulation, several different types of cells including epithelial, endothelial, mesenchymal, and fibroblastic cells are implicated in the wound healing process (Pastar et al, 2008 and Bannasch et al., 2000). Specifically, they proliferate and migrate into the wound area to promote healing. Therefore, experiments were conducted to determine if factors secreted from amniotic and chorionic membranes produced pursuant to the present disclosure promote cell migration and wound field closure. To accomplish this, a commercially available wound healing assay (Cell Biolabs) and a highly accepted human microvascular endothelial cell line (HMVEC, Lonza Inc.) were utilized. Results indicated that cell migration was enhanced by treatment with conditioned media from the placental membranes.

In Vitro Cell Migration

00354 Human microvascular endothelial cells (HMVECs) were grown under normal cell culture conditions in defined complete media (Lonza Inc.). To assess migration and wound field closure, a commercially available wound healing assay was used (Cell Biolab). The assay principle is outlined in FIG. 17.

00355 FIG. 17 depicts the Cell Biolabs 24-well Cytoselect wound healing assay. (Figure reproduced from Cell Biolabs).

00356 Cells were collected via trypsinization, pelleted, and counted before being resuspended in complete media at a density of 2×10^5 cells/mL. 250 μ L (5×10^4 cells) of cell suspension was then pipetted into each side of a well containing a wound healing insert (Cytoselect 24-well Wound Healing Assay Plate, Cell Biolabs). The cells were grown for 24 hours in complete media. After 24 hours, the wound inserts were removed. At the same time, complete media was removed and replaced with experimental media. Complete media and basal media were used as positive and negative controls, respectively. To generate experimental media, placental membranes were incubated for 3 days in DMEM with 1% human serum albumin (HSA) in a tissue culture incubator. The resulting tissue and media were then placed in eppendorf tubes and spun at high speed in a microcentrifuge. The supernatants were collected and stored at $-80^\circ\text{C} \pm 5^\circ\text{C}$ until use. For migration and wound healing studies, conditioned media from placental membranes was diluted 1:20 in basal media before being added to experimental wells. After 18 hours, the media was removed, and the cells were fixed for 20 min in 4% paraformaldehyde and stained with crystal violet. The wound field in each well was then photographed. Wound healing was determined by the amount of wound field still visible at the end of the experiment when compared to control pictures taken before conditioned media was added to the wells.

Placental Membrane Conditioned Media Supports Cell Migration and Wound Field Closure

00357 Conditioned media from amniotic and chorionic membranes was used to assess the potential for these membranes to promote cell migration and wound field closure. Conditioned media from placental chorionic membranes supported migration of cells into the experimental wound field. FIG. 18 depicts representative images of *HMVECs* treated with 5% conditioned media from amniotic, chorionic, or a

combination of amniotic/chorionic tissue as well as positive and negative controls. Wound field is 0.9 mm in width.

00358 The ability of factors from placental membranes produced pursuant to the present disclosure to promote HMVEC migration indicated that these tissues have the ability to enhance wound healing. Additionally, based on the insight of the inventors, it has been surprisingly discovered that these tissues also enhance revascularization since the HMVEC cell line is derived from vascular endothelial cells.

00359 These data demonstrate that placental products of the present invention produce unexpectedly superior levels of factors that promote wound healing therapies.

Example 23 Analysis of Factors in Exemplary Placental Tissue Products

00360 Table 15 depicts the biochemical profile of exemplary placental products of the invention (results adjusted per cm^2 after subtraction of the negative background).

Table 15 Factors in placental tissue product (pg/cm^2)

	Units	Apligraf	Dermagraft	AM75	CM75	AM78	CM78
hMMP1	$\text{pg}/\text{ml}/\text{cm}^2$	1964945.37	14818.20	2821.85	3531.81	117326.89	95.46
hMMP7	$\text{pg}/\text{ml}/\text{cm}^2$	911.54	0.00	0.00	0.00	3.96	0.00
hMMP10	$\text{pg}/\text{ml}/\text{cm}^2$	0.00	0.00	113.94	0.00	0.00	0.00
hMMP13	$\text{pg}/\text{ml}/\text{cm}^2$	21.61	0.00	0.00	0.00	0.71	0.00
hMMP3	$\text{pg}/\text{ml}/\text{cm}^2$	208281.70	180721.52	170.26	161.52	8325.17	0.00
hMMP9	$\text{pg}/\text{ml}/\text{cm}^2$	8872.28	19321.39	214.78	1455.11	630.56	57.59
hMMP2	$\text{pg}/\text{ml}/\text{cm}^2$	153341.77	19712.21	287.14	37.93	3823.38	24.44
hMMP8	$\text{pg}/\text{ml}/\text{cm}^2$	36.92	12.19	0.00	0.00	0.00	0.00
hTIMP1	$\text{pg}/\text{ml}/\text{cm}^2$	2487.18	10909.84	569.23	883.05	28743.48	97.94
hTIMP2	$\text{pg}/\text{ml}/\text{cm}^2$	7285.53	1796.56	89.29	13.72	424.06	4.83
MMP/TIMP		239.26	19.72	6.81	6.26	4.50	2.62

Example 24 Factors in Exemplary Placental Products as Measured Through Protein Array Analysis by Aushon Biosystems

00361 Table 16 depicts the biochemical profile of the lysates of exemplary placental products of the invention (results adjusted per cm² after subtraction of the negative background).

Table 16

	AM75 lysate	AM78 lysate	CM75 lysate	CM78 lysate
	pg/ml	pg/ml	pg/ml	pg/ml
hACRP30	50.8	1154.6	1213.7	225.3
hAlpha2-Macroglobulin	1910.6	426191.6	8174.4	9968.6
hEGF	127.3	361.4	0.0	0.8
hbFGF	119.1	821.5	375.0	351.3
hGCSF	0.7	3.2	1.2	0.7
hHBEGF	127.5	168.0	15.4	84.5
hHGF	3943.7	15060.0	29979.6	50392.8
hIGFBP1	5065.0	9456.6	934.0	1443.6
hIGFBP2	12460.8	5569.7	135.9	134.6
hIGFBP3	50115.7	41551.4	4571.5	11970.2
hIL1ra	3881.0	32296.9	5168.2	525.5
hKGF	1.4	8.8	3.1	1.5
hLIF	0.0	4.2	0.0	0.0
hMMP1	9144.1	20641.2	2882.9	6582.3
hMMP10	0.0	15.5	79.3	87.5
hMMP2	2067.3	4061.9	949.5	748.8
hMMP3	0.0	36.2	0.0	0.0

hMMP7	5.1	11.4	4.5	9.1
hMMP8	0.0	0.0	0.0	0.0
hMMP9	92.2	2878.1	2676.2	1259.3
hNGAL	6900.1	6175.9	938.5	229.7
hPDGFAA	0.0	12.5	39.8	35.2
hPDGFAB	11.2	31.3	14.4	14.0
hPDGFbb	4.6	13.4	4.0	1.3
hPEDF	0.0	652.6	0.0	0.0
hTIMP1	7958.1	35955.6	50712.3	17419.9
hTIMP2	3821.8	7443.2	640.7	780.0
hVEGF	3.3	11.8	125.2	8.4
hVEGFC	46.5	150.0	123.5	51.7
hVEGFD	25.7	31.0	15.0	20.4

CLAIMS

What is claimed is:

1. A placental product comprising: a chorionic membrane, wherein the chorionic membrane:
 - a. comprises viable therapeutic cells; and
 - b. is substantially free of CD14+ macrophages; andoptionally, wherein the viable therapeutic cells are native to the chorionic membrane.
2. The placental product of Claim 1, wherein the chorionic membrane is substantially free of vascularized tissue or vascularized tissue-derived immunogenic cells.
3. The placental product of Claim 1, wherein the chorionic membrane is substantially free of trophoblasts.
4. The placental product of Claim 3, wherein the chorionic membrane is substantially free of vascularized tissue or vascularized tissue-derived immunogenic cells.
5. The placental product of Claim 1, wherein the chorionic membrane is substantially free of immunogenic maternal cells.
6. The placental product of Claim 1, wherein the chorionic membrane is substantially free of maternal dendritic cells.
7. The placental product of Claim 1, wherein the chorionic membrane is substantially free of maternal leukocytes.
8. The placental product of Claim 1, wherein the viable therapeutic cells comprise one or more of:
 - a. MSCs;
 - b. fibroblasts; and
 - c. stromal cells.
9. The placental product of Claim 8, wherein the chorionic membrane is substantially free of vascularized tissue or vascularized tissue-derived immunogenic cells.

10. The placental product of Claim 8, wherein the chorionic membrane is substantially free of trophoblasts.
11. The placental product of Claim 10, wherein the chorionic membrane is substantially free of vascularized tissue or vascularized tissue-derived immunogenic cells.
12. The placental product of Claim 8, wherein the chorionic membrane is substantially free of immunogenic maternal cells.
13. The placental product of Claim 8, wherein the chorionic membrane is substantially free of maternal dendritic cells.
14. The placental product of Claim 8, wherein the chorionic membrane is substantially free of maternal leukocytes.
15. The placental product of any of Claims 8-14, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable.
16. The placental product of Claim 15, wherein the stromal cells are about 5,000 to about 50,000 in number per cm^2 of the chorionic membrane.
17. The placental product of Claim 15, wherein the stromal cells are present at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 per cm^2 .
18. The placental product of any of Claims 8-14, wherein about 40% to about 92.5% of the stromal cells are viable.
19. The placental product of Claim 18, wherein the stromal cells are about 5,000 to about 50,000 in number per cm^2 of the chorionic membrane.
20. The placental product of Claim 19, wherein the stromal cells are at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 in number per cm^2 .
21. The placental product of any of Claims 1-14, wherein the chorionic membrane comprises about 5,000 to about 50,000 stromal cells per cm^2 of the chorionic membrane, optionally, wherein the stromal cells are native to the chorionic membrane.

22. The placental product of Claim 21, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable.
23. The placental product of Claim 21, wherein about 40% to about 92.5% of the stromal cells are viable.
24. The placental product of any of Claims 1-14, wherein the chorionic membrane comprises at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 stromal cells per cm² of the chorionic membrane, optionally, wherein the stromal cells are native to the chorionic membrane.
25. The placental product of Claim 24, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable.
26. The placental product of Claim 24, wherein about 40% to about 92.5% of the stromal cells are viable.
27. The placental product of any of Claims 1-14, wherein the chorionic membrane comprises fibroblasts in about 50% to about 90% of the total cells
28. The placental product of any of Claims 1-14, wherein the chorionic membrane comprises CD14+ macrophage in an amount of less than about 5% or less than about 1% or less than about 0.5%.
29. The placental product of any of Claims 1-14, wherein the placental product or chorionic membrane releases between about 0 to about 420 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the placental product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.
30. The placental product of Claim 29, wherein said about between about 0 to about 420 pg/mL TNF- α is: about 0 to about 350 pg/mL TNF- α or 0 to about 280 pg/mL TNF- α .
31. The placental product of any of Claims 1-14, wherein the chorionic membrane further comprises one or more native factors consisting of the group of IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9 and TIMP1;
wherein native factor is present in an amount:

- i. substantially greater per gram of chorionic membrane than that of a fresh unprocessed chorionic membrane;
 - ii. substantially greater per native chorionic membrane cell than that of a fresh unprocessed chorionic membrane; or
 - iii. substantially greater per μm of chorionic membrane.
- 32. The placental product of any of Claims 1-14, wherein the chorionic membrane has a thickness of about 40 μm to about 400 μm .
- 33. The placental product of Claims 30, wherein the chorionic membrane has a thickness of about 40 μm to about 400 μm .
- 34. The placental product of any of Claims 1-14, further comprising a basement membrane, a reticular layer, or a combination thereof.
- 35. The placental product of Claims 31, further comprising a basement membrane, a reticular layer, or a combination thereof
- 36. A placental product comprising: a cryopreservation medium and a chorionic membrane, wherein:
 - a. the chorionic membrane is substantially free of trophoblasts, CD14+ macrophages, or both trophoblasts and CD14+ macrophage; and
 - b. wherein the cryopreservation medium comprises a cryopreserving amount of a cryopreservative; andoptionally, wherein the chorionic membrane is substantially free of vascularized tissue or vascularized tissue-derived immunogenic cells.
- 37. The placental product of Claim 35, wherein the chorionic membrane comprises viable therapeutic cells, optionally, wherein the viable therapeutic cells are native to the chorionic membrane and of one or more of the types.
 - a. MSCs;
 - b. fibroblasts; and
 - c. stromal cells.
- 38. The placental product of Claim 36, wherein the cryopreservative comprises a cell-permeating cryopreservative, a non-cell-permeating cryopreservative, or a combination thereof, optionally wherein the permeating cryopreservative comprises:
 - a. DMSO; or
 - b. DMSO in a majority amount.
- 39. The placental product of Claim 37, wherein the cell-permeating cryopreservative:
 - a. does not comprise glycerol in a majority amount; or

- b. does not comprise a substantial amount of glycerol.
40. The placental product of Claim 37, wherein the cell-permeating cryopreservative comprises a DMSO in amount of: about 2% to about 20%, about 5% to about 20%, about 5% to about 15%, or about 7% to about 13%.
41. The placental product of Claim 37, wherein the cryopreservation medium is present in an amount of at least: about 20 ml, about 30 ml, about 35 ml, about 40 ml, about 45 ml, or about 50 ml.
42. The placental product of Claim 39, wherein the cryopreservation medium is present in an amount of at least: about 20 ml, about 30 ml, about 35 ml, about 40 ml, about 45 ml, or about 50 ml.
43. The placental product of Claim 37, wherein the cryopreservation medium further comprises albumin, optionally wherein the albumin is HSA.
44. The placental product of Claim 39, wherein the cryopreservation medium further comprises albumin, optionally wherein the albumin is HSA.
45. The placental product of any of Claims 35-43, wherein the cryopreservative comprises a non-cell-permeating cryopreservative selected from is a polyvinylpyrrolidone, a starch, a hydroxyethyl starch, a polysaccharide, a monosaccharides, a sugar alcohol, an alginate, a trehalose, a raffinose, a dextran, or a combination thereof.
46. The placental product of any of Claims 35-43, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
47. The placental product of Claim 45, wherein the stromal cells are about 5,000 to about 50,000 in number per cm^2 of the chorionic membrane.
48. The placental product of Claim 45, wherein the stromal cells are at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 in number per cm^2 .
49. The placental product of any of Claims 35-43, wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
50. The placental product of Claim 48, wherein the stromal cells are about 5,000 to about 50,000 in number per cm^2 of the chorionic membrane.

51. The placental product of any of Claims 48, wherein the stromal cells are at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 in number per cm².
52. The placental product of any of Claims 35-43, wherein the chorionic membrane comprises about 5,000 to about 50,000 stromal cells per cm² of the chorionic membrane, optionally, wherein the stromal cells are native to the chorionic membrane.
53. The placental product of Claim 51, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
54. The placental product of Claim 51, wherein about 40% to about 92.5% of the stromal cells are viable.
55. The placental product of any of Claims 35-43, wherein the chorionic membrane comprises at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 stromal cells per cm² of the chorionic membrane, optionally, wherein the stromal cells are native to the chorionic membrane.
56. The placental product of Claim 54, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
57. The placental product of Claim 54, wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
58. The placental product of any Claims 35-43, wherein the chorionic membrane comprises about comprises fibroblasts in about 50% to about 90% of the total cells.
59. The placental product of any of Claims 35-43, wherein the chorionic membrane comprises CD14+ macrophages in an amount of less than about 5% or less than about 1%.
60. The placental product of any of Claims 35-43, wherein the chorionic membrane is substantially free of:
- a. maternal dendritic cells, maternal leukocytes, or a combination thereof, or
 - b. immunogenic maternal cells,

61. The placental product of any of Claims 35-43, wherein the product comprises about 7,500 to about 48,000 stromal cells per cm^2 and wherein the viability of said stromal cells after a freeze-thaw cycle is about 70% to about 95% relative to fresh placental or chorionic membrane.
62. The placental product of any of Claims 35-43, wherein after a freeze-thaw cycle, the placental product or chorionic membrane release between about 0 to about 420 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the placental product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.
63. The placental product of any of Claims 35-43, wherein after a refrigeration-freeze-thaw cycle, the placental product or chorionic membrane releases between about 0 to about 420 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm x 2 cm piece of the placental product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.
64. The placental product of Claim 61, wherein said about between about 0 to about 420 pg/mL TNF- α is: about 0 to about 350 pg/mL TNF- α or 0 to about 280 pg/mL TNF- α .
- a. The placental product of any of Claims 35-53, wherein the chorionic membrane further comprises one or more native factors selected from the group consisting of IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9 and TIMP1; wherein the one or more native factors are present in an amount, wherein the amount is:
 - i. substantially greater per gram of chorionic membrane than that of a fresh unprocessed chorionic membrane;
 - ii. substantially greater per native chorionic membrane cell than that of a fresh unprocessed chorionic membrane; or
 - iii. substantially greater per μm of chorionic membrane.
65. The placental product of any of Claims 1-14 or 35-43, further comprising an amniotic membrane.
66. The placental product of Claim 65, wherein the amniotic membrane comprises a layer of amniotic epithelial cells.
67. The placental product of Claim 66, wherein the amniotic membrane and the chorionic membrane are associated to one another in the native architecture.


68. The placental product of Claim 66, wherein the amniotic membrane and the chorionic membrane are not attached to one another in the native architecture.
69. The placental product of any of Claims 35-43, wherein the chorionic membrane comprises a basement membrane, a reticular layer, or a combination thereof.
70. The placental product of any of Claims 35-43, wherein the chorionic membrane has a thickness of about 40 μm to about 400 μm .
71. The placental product of any of Claims 1-14 or 35-43, wherein the maternal side of the chorionic membrane comprises fragments of extracellular matrix proteins in a concentration substantially greater than that of a native, unprocessed chorion.
72. The placental product of Claim 70, wherein a substantial portion of the protein fragments comprises terminal leucine or phenylalanine.
73. The placental product of any of Claims 1-14 or 35-43, wherein the chorionic membrane has been treated with a protease, optionally wherein the protease is a dispase
74. The placental product of any of Claims 1-14 or 35-43, wherein the chorionic membrane:
- a. comprises a basement membrane, a reticular layer, or a combination thereof; and
 - b. has a thickness of about 40 μm to about 400 μm .
75. A placental product comprising: a cryopreservation medium and a chorionic membrane, wherein:
- a. the chorionic membrane:
 - i. comprises a reticular layer and a stromal layer, wherein each of said layers comprises viable therapeutic cells; and
 - ii. is substantially free of vascularized tissue, functional CD14+ macrophages, and trophoblasts; and
 - b. the cryopreservation medium comprises a cryopreserving amount of a cryopreservative.
76. The placental product of Claim 74, wherein the chorionic membrane comprises about 5,000 to about 50,000 stromal cells per cm^2 of the amniotic membrane.
77. The placental product of Claim 75, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable.


78. The placental product of any Claim 75, wherein about 40% to about 92.5% of the stromal cells are viable.
79. The placental product of Claim 74, wherein the chorionic membrane comprises at least: about 7,400 or about 15,000 or about 23,217, or about 35,000, or about 40,000 or about 47,800 stromal cells per cm² of the chorionic membrane.
80. The placental product of Claim 75, wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
81. The placental product of any Claim 75, wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle.
82. The placental product of any of Claims 74-80, wherein the chorionic membrane further comprises a basement membrane.
83. The placental product of any of Claims 74-80, wherein the amniotic membrane has a thickness of about 40 um to about 400 um.
84. The placental product of any of Claims 74-80, wherein the cryopreservative comprises DMSO.
85. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein the chorionic membrane comprises MSCs in an amount of about 5% to about 30%, about 5% to about 25%, about 5% to about 20%, about 5% to about 15%, about 3% to about 12%, at least about 5%, at least about 10%, or at least about 15%, relative to the total number of cells in the chorionic membrane.
86. The placental product of Claim 104, wherein at least: about 40%, about 50%, about 60%, or about 70% of the MSCs are viable after a freeze-thaw cycle.
87. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein the chorionic membrane comprises fibroblasts in an amount of about 50% to about 95%, about 60% to about 90%, or about 70% to about 85%, relative to the total number of cells in the chorionic membrane.
88. The placental product of Claim 86, wherein at least: about 40%, about 50%, about 60%, or about 70% of the fibroblasts are viable after a freeze-thaw cycle.
89. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein, the chorionic membrane comprises functional macrophages in an amount of less than about any of 5%, 4%, 3%, 2%, 1%, or 0.1%.


90. The placental product of any of Claims 1-14, 35-43, and 74, wherein the chorionic membrane comprises MSCs and functional macrophages in a ratio of greater than about any of 3:1, 4:1, 5:1, 7:1, 10:1, 12:1, or 15:1.
91. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein, the chorionic membrane comprises fibroblasts and functional macrophages in a ratio of greater than about any of 14:1, 15:1, 16:1, 17:1, 28:1, 30:1, 35:1, 45:1, or 50:1
92. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein the chorionic membrane comprises fibroblasts and MSCs in a ratio of: about 9:2 to about 17:3.
93. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein the chorionic membrane comprises MSCs in an amount of at least about 1,500 cells/cm², at least about 3,000 cells/cm², about 15,000 to about 9,000 cells/cm², or about 3,000 to about 9,000 cells/cm².
94. The placental product of Claim 92, wherein at least: about 40%, about 50%, about 60%, or about 70% of the MSCs are viable after a freeze-thaw cycle.
95. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein the chorionic membrane comprises fibroblasts in an amount of at least about 7,000 cells/cm², at least about 14,000 cells/cm², about 7,000 to about 51,000 cells/cm², or about 14,000 to about 51,000 cells/cm².
96. The placental product of Claim 94, wherein at least: about 40%, about 50%, about 60%, or about 70% of the fibroblasts are viable after a freeze-thaw cycle.
97. The placental product of any of Claims 1-14, 35-43, or 74 - 80, wherein the chorionic membrane comprises functional macrophages in an amount of less than about 3,000 cells/cm², or less than about 1,000 cells/cm².
98. A method of treating a wound of a patient comprising administering the placental product of any of Claims 1-14, 35-43, or 74 - 80 to the wound.
99. The method of Claim 97, wherein the wound is an ulcer.
100. The method of Claim 98, wherein the ulcer is a foot ulcer, optionally a diabetic foot ulcer.
101. The method of Claim 98, wherein the ulcer is a diabetic foot ulcer.
102. The method of Claim 97, wherein the patient has diabetes.


103. A method of ligament or tendon surgery comprising applying the placental product of any of Claims 1-14, 35-43, or 74 - 80 to a ligament or tendon.
104. The method of Claim 102, wherein the ligament or tendon surgery is a grafting surgery, a reconstruction surgery, or a repair of a torn ligament or tendon.
105. The method of Claim 103, wherein the grafting surgery comprises the attachment of a graft to a biological substrate, optionally, wherein the biological substrate is bone.
106. The method of Claim 104, wherein the graft is an allograft, a xenograft, or an autologous graft.
107. The method of Claim 105, wherein the ligament or tendon surgery is selected from: knee surgery, shoulder surgery, leg surgery, and arm surgery, elbow surgery, finger surgery, hand surgery, wrist surgery, toe surgery, foot surgery, ankle surgery.
108. The method of Claims 105, wherein the tendon is hamstring tendon, a bicep tendon, an Achilles Tendon, an extensor tendon, or a rotator cuff tendon.
109. The method of Claims 105, wherein the step of applying the placental product comprises: suturing the placental product to the tendon or ligament, enveloping the tendon or ligament with the placental product, or a combination thereof.
110. A method of tissue grafting comprising applying the placental product of any of Claims 1-14, 35-43, or 74 - 80 to a tissue graft, and attaching the tissue graft to a biological substrate.
111. The method of Claim 109, wherein the placental product is applied to a portion of the tissue graft which is attached to the biological substrate.
112. The method of Claim 110, wherein the step of applying the placental product comprises: suturing the placental product to the graft, enveloping graft with the placental product, or a combination thereof.
113. The method of Claim 111, wherein the tissue graft is comprises a tissue selected from: skin, cartilage, ligament, tendon, periosteum, perichondrium, synovium, fascia, and mesenter.
114. A method of inhibiting fibrosis or adhesion at a wound site comprising applying the placental product of any of Claims 1-14, 35-43, or 74 - 80 to the wound site.


115. The method of Claim 113, wherein the wound site is surgery induced or associated with surgery.
116. The method of Claim 114, wherein the surgery is a surgery selected from: spine, laminectomy, knee, shoulder, or child birth, eye, cardiovascular, angiogenesis stimulation, burn, and brain/neurological procedures.
117. The method of Claim 115, wherein the wound site is a trauma related wound.
118. A method of promoting migration or proliferation of at least one cell type in a wound of a subject wherein the at least one cell type selected from the group consisting of fibroblasts, epithelial cells, and endothelial cells in a subject comprising administering the placental product of any of Claims 1-14, 35-43, or 74 - 80 to a subject in need thereof.
119. A method of reorganizing extracellular matrix in a subject comprising administering the placental product of any of Claims 1-14, 35-43, or 74 - 80 to the subject.
120. A method of providing at least one factor to a wound of a subject comprising administering the placental product of any of Claims 1-14, 35-43, or 74 - 80 to the wound and wherein the at least one factor is a chemotactic factor or a growth factor.
121. A method of providing stem cells to tissue of a subject comprising administering the placental product of any of Claims 1-14, 35-43, or 74 - 80 to the subject.




















FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5
Expression of IL-2R on T-cells stimulated by Placental Derived Cells (donor 37 fresh)




FIGURE 6




FIGURE 7A




FIGURE 7B




FIGURE 7C




FIGURE 8A




FIGURE 8B






FIGURE 8C



C

FIGURE 9









FIGURE 11A**FIGURE 11B****FIGURE 11C**




FIGURE 12A**FIGURE 12B**

FIGURE 13A

Expression of bone reparative proteins in chorionic membrane homogenates

**FIGURE 13B**

Expression of IGF-1 in chorionic membrane homogenates




FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17




FIGURE 18




FIGURE 19A




FIGURE 19B




FIGURE 19C

