



(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**  
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/10/24  
(87) Date publication PCT/PCT Publication Date: 2018/05/03  
(85) Entrée phase nationale/National Entry: 2019/04/23  
(86) N° demande PCT/PCT Application No.: IB 2017/001427  
(87) N° publication PCT/PCT Publication No.: 2018/078442  
(30) Priorité/Priority: 2016/10/24 (US62/411,783)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),  
*A61K 39/00* (2006.01), *A61K 31/00* (2006.01),  
*C07K 16/24* (2006.01), *C07K 16/28* (2006.01)


(71) Demandeurs/Applicants:  
NOVIMMUNE S.A., CH;  
DE MIN, CRISTINA, CH;  
FERLIN, WALTER, CH;  
DE BENEDETTI, FABRIZIO, CH

(72) Inventeurs/Inventors:  
DE MIN, CRISTINA, CH;  
FERLIN, WALTER, CH;  
DE BENEDETTI, FABRIZIO, CH

(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : PROCEDES, COMPOSITIONS ET SCHEMAS POSOLOGIQUES POUR TRAITER OU PREVENIR DES INDICATIONS ASSOCIEES A L'INTERFERON GAMMA  
(54) Title: METHODS, COMPOSITIONS AND DOSING REGIMENS FOR TREATING OR PREVENTING INTERFERON-GAMMA RELATED INDICATIONS

FIGURE 19



(57) **Abrégé/Abstract:**

The disclosure relates generally to methods compositions and dosing regimens for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with elevated IFN- $\gamma$  levels.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau

(43) International Publication Date  
03 May 2018 (03.05.2018)



(10) International Publication Number  
**WO 2018/078442 A3**

(51) International Patent Classification:

A61K 39/395 (2006.01) C07K 16/28 (2006.01)  
A61K 31/00 (2006.01) A61K 39/00 (2006.01)  
C07K 16/24 (2006.01)

(21) International Application Number:

PCT/IB2017/001427

(22) International Filing Date:

24 October 2017 (24.10.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/411,783 24 October 2016 (24.10.2016) US

(71) Applicant: NOVIMMUNE SA [CH/CH]; 14 Chemin des  
Aulx, 1228 Plane-les-Ouates (CH).

(72) Inventors; and

(71) Applicants: DE MIN, Cristina [CH/CH]; 14 Chemin  
des Aulx, 1228 Plane-les-Ouates (CH). FERLIN, Wal-  
ter [CH/CH]; 14 Chemin des Aulx, 1228 Plane-les-Ouates  
(CH). DE BENEDETTI, Fabrizio [CH/CH]; 14 Chemin  
des Aulx, 1228 Plane-les-Ouates (CH).

(74) Agent: WRAIGE, David Leslie Charles et al.; Cooley  
(UK) LLP, Dashwood, 69 Old Broad Street, London EC2M  
1QS (GB).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,  
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,  
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,  
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,  
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,  
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,  
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,  
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,  
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,  
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

23 August 2018 (23.08.2018)

(54) Title: METHODS, COMPOSITIONS AND DOSING REGIMENS FOR TREATING OR PREVENTING INTERFERON-GAM-  
MA RELATED INDICATIONS

(57) Abstract: The disclosure relates generally to methods compositions and dosing regimens for treating, preventing and/or delaying  
the onset or progression of, or alleviating a symptom associated with elevated IFN- $\gamma$  levels.

WO 2018/078442 A3

## **METHODS, COMPOSITIONS AND DOSING REGIMENS FOR TREATING OR PREVENTING INTERFERON-GAMMA RELATED INDICATIONS**

### **Related Applications**

**[0001]** This application claims the benefit of and priority to U.S. Provisional Application No. 62/411,783, filed October 24, 2016 the contents of which are incorporated herein by reference in its entirety

### **Field of the Invention**

**[0002]** The disclosure relates generally to methods and compositions for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with elevated levels interferon gamma (IFN $\gamma$ , IFN-gamma) such as for example, hemophagocytic lymphohistiocytosis (HLH), hemorrhagic fever, CAR-T cell therapy, transplant failure, transplant rejection, Graft-versus-host disease (GvHD), and/or an inflammatory disorder associated with transplant rejection. The disclosure also relates to methods and compositions for prolonging survival of transplanted biological material.

### **Background of the Invention**

**[0003]** Human interferon gamma (IFN $\gamma$ , IFN-gamma) is a lymphokine produced by activated T-lymphocytes and natural killer cells. It manifests anti-proliferative and immunomodulatory activities and binds to IFN $\gamma$ -R, a heterodimeric receptor on most primary cells of the immune system, and triggers a cascade of events leading to inflammation. The immunomodulatory activity of IFN $\gamma$  is known to have beneficial effects in a number of clinical conditions. However, there are many clinical settings in which IFN $\gamma$ -activity is known to have deleterious effects. For example, autoimmune diseases are associated with high levels of IFN $\gamma$  in the blood and diseased tissue from autoimmune patients. IFN $\gamma$ -activity has also been linked to such disease states as cachexia and septic shock.

**[0004]** IFN $\gamma$  has been implicated in a number of disorders; and anti-IFN $\gamma$  agents are being developed as therapeutic agents..

### **Summary of the Invention**

**[0005]** In various aspects the invention provides multiple variable dose treatment regimen

for the treatment of diseases, disorder or conditions associated with elevated IFN- $\gamma$  levels.

[0006] In one aspect, the invention provides a method for the treatment of primary hemophagocytic lymphohistiocytosis (HLH) in a human by intravenously administering to a subject a first dose and second dose of an antibody that binds interferon gamma (IFN $\gamma$ ). The subject is an adult subject or a pediatric subject. The first dose is 1.0 or 3.0 mg/kg of the subject's body weight and the second dose is 3.0, 6.0 or 10.0 mg/kg of the subject's body weight. Optionally, a third dose of 1.0 mg/kg of the subject's body weight is administered.

[0007] In another aspect, the invention provides a method for the treatment of secondary hemophagocytic lymphohistiocytosis (HLH) in a human pediatric subject by intravenously administering to a subject a first dose and second dose of an antibody that binds interferon gamma (IFN $\gamma$ ). The first dose is 6.0 mg/kg of the subject's body weight and the second dose is 3.0 mg/kg of the subject's body weight. Optionally, a third dose of 6.0 mg/kg of the subject's body weight is administered.

[0008] In a further aspect, the invention provides a method for the treatment of secondary hemophagocytic lymphohistiocytosis (HLH) in a human adult subject by intravenously administering to a subject a first dose and second dose of an antibody that binds interferon gamma (IFN $\gamma$ ). The first dose is 3.0 mg/kg or 6.0 mg/kg of the subject's body weight and the second dose is no more than 10 mg/kg of the subject's body weight. For example, the second dose is 1.0, 3.0, 6.0 or 10.0 mg/kg. Optionally, a third dose of less than 10.0 mg/kg of the subject's body weight is administered. For example the third dose is 1.0, 3.0, or 6.0 mg/kg of the subject's body weight.

[0009] In yet another aspect, the invention provides a method for the treatment of a condition in a human subject by intravenously administering to a subject a first dose and second dose of an antibody that binds interferon gamma (IFN $\gamma$ ). The subject is an adult subject or a pediatric subject. The condition is transplant rejection such as solid organ transplant disorder or bone marrow acute graft rejection. The condition is graft vs. host disease, paraneoplastic cerebellar degeneration, hemorrhagic fever, sarcoidosis, or adult onset Still's disease. Alternatively, the method is administered to a subject after receiving CART cell therapy. The first dose is between 1.0 to 10 mg/kg of the subject's body weight and the second dose is between 1.0 to 10 mg/kg of the subject's body weight e subject's body weight. For example, the second dose is 1.0, 3.0, 6.0 or 10.0 mg/kg. Preferably, the second dose is higher or lower than the first dose. Optionally, a third dose of between 1.0 to 10 mg/kg of the subject's body weight is administered. For example, the first, second, or third dose is 1.0, 3.0, 6.0 or 10.0 mg/kg of the subject's body

weight.

**[0010]** The antibody that binds interferon gamma (IFN $\gamma$ ) includes: a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6). For example the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

**[0011]** The dose of the antibody is administered within 1, hr, 6 hrs or 12 hrs.

**[0012]** The second dose is administered for a first treatment period of every three days after the first dose. Additionally, the second dose is administered for a second treatment period after completion of the first treatment period. The the second treatment period is for example twice weekly.

**[0013]** The antibody dose is administered as a single injection.

**[0014]** The antibody is administered as a monotherapy or a co-therapy.

**[0015]** Optionally, the methods of the invention further includes administering dexamethasone immediately prior to the dosing of the antibody. the dexamethasone is administered at a dose of at least 10 mg/m<sup>2</sup> or at least 5 mg/m<sup>2</sup>.

**[0016]** The subject has not previously been treated for HLH.

**[0017]** In various aspects, the method further includes comprises administering at least a second agent to the subject. The second agent is a therapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent.

**[0018]** Also included in the invention is an injectable pharmaceutical formulation comprising per mL: 5 mg or 25 mg of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody; 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is between 5.8 and 6.2.

[0019] In a further aspect, the invention provides a unit dose vial containing 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution suitable for injection where the concentration of antibody is 5 mg/ml or 25 mg/ml and the pH of the solution is between 5.8 and 6.2. The antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate.

[0020] In another aspect, the invention provides a unit dose vial containing 10 ml or 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution suitable for injection where the concentration of antibody is at 25 mg/ml, and the pH of the solution is between 5.8 and 6.2. The antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate.

[0021] In yet another aspect the invention provides a unit dose vial containing 2 ml or 10 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution suitable for injection where the concentration of antibody is 5 mg/ml and the pH of the solution is between 5.8 and 6.2. The antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate.

[0022] The antibody that binds interferon gamma (IFN $\gamma$ ) includes: a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6). For example the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

[0001] Any of the above aspects or embodiments may be combined with any other aspect or embodiment.

[0002] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention

pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are expressly incorporated by reference in their entirety. In cases of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples described herein are illustrative only and are not intended to be limiting.

**[0023]** Other features and advantages of the invention will be apparent from and encompassed by the following detailed description and claims

### **Brief Description of the Drawings**

**[0024]** Figure 1 is a graph depicting correlations between predose serum CXCL9 levels and total IFN $\gamma$  levels at 24 h post infusion with the NI-0501 antibody in an ongoing phase 2 pilot study in primary HLH patients.

**[0025]** Figure 2 is a graph depicting correlations between serum CXCL9 levels and total IFN $\gamma$  predose levels at 24 h post infusion with the NI-0501 antibody in an ongoing phase 2 pilot study in primary HLH patients.

**[0026]** Figures 3A and 3B are a series of graphs depicting correlations between serum CXCL9 levels and IFN $\gamma$  levels in patients with Macrophage Activation Syndrome (MAS) secondary to systemic Juvenile Idiopathic Arthritis (sJIA) and in patients with active sJIA.

**[0027]** Figures 4A-1, 4A-2, 4B-1, 4B-2, 4C-1, 4C-2, 4D-1, and 4D-2 are a series of graphs depicting correlations between IFN $\gamma$  and serum CXCL9 levels and clinical parameters in patients with active sJIA and MAS secondary to sJIA.

**[0028]** Figure 5 is a graph depicting that IFN $\gamma$  was fully neutralized as shown by undetectable levels of IFN $\gamma$ -inducible chemokines.

**[0029]** Figure 6 is a graph depicting improvement of HLH disease activity during NI-0501 treatment (2 weeks and end of treatment): percent of patients with Platelet count  $>100 \times 10^9/L$ , Neutrophil count  $>1 \times 10^9/L$ , Fibrinogen  $>1.5 \text{ g/L}$  and ferritin decrease of at least 25%.

**[0030]** Figures 7A and 7B are a series of graphs depicting the correlation between pre-dose CXCL9 and total IFN $\gamma$  levels at 24h after NI-0501 infusion. The insert shown in Figure 7B, depicts an example of individual IFN $\gamma$  and CXCL9 profile during NI-0501 treatment.

**[0031]** Figures 8A, 8B, 8C, and 8D are a series of graphs depicting serum levels of IFN $\gamma$  and of CXCL9, CXCL10 and CXCL11 in individual patients from whom paired samples were

available during active MAS and during active sJIA without MAS at sampling (Act sJIA).

Significance levels (p) were obtained using the Wilcoxon rank test for paired samples.

**[0032]** Figures 9A and 9B are a series of graphs depicting Changes in white blood cell (WBC) and platelet (PLT) counts and in ferritin levels (Fig. 9A) and changes in serum levels of IFN $\gamma$ , CXCL9, CXCL10 and CXCL11 (Fig. 9B) in one patient who presented, during the course of his sJIA, 3 episodes of MAS.

**[0033]** Figures 10A, 10B, 10C, 10D, 10E, 10F, 10G, 10H, 10I, and 10J are a series of graphs depicting correlation of levels of IFN $\gamma$  and CXCL9 with ferritin levels, neutrophil and platelet count and with LDH and ALT levels in patients with active MAS at sampling (red circles) and in patients with active sJIA without MAS at sampling (black triangles). Spearman correlation coefficient (Rs) and significance level (p) of each correlation are shown in Table 3.

**[0034]** Figures 11A, 11B, 11C, 11D, 11E, and 11F are a series of graphs depicting the relation of IFN $\gamma$  with CXCL9 and CXCL10 production in MAS. Panel A: Correlations of the levels of IFN $\gamma$  with the levels of CXCL9 and CXCL10 in patients with MAS at sampling Spearman correlation coefficient (Rs) and significance level (p) of each correlation are shown in Table 3.

**[0035]** Figure 12 is a schematic representation of the screening, treatment, and follow-up portions of the studies presented in Example 7.

**[0036]** Figures 13A and 13B are graphs depicting the effect of NI-0501 administration on body temperature in two patients having body temperature  $> 37.5^{\circ}\text{C}$  at initiation of NI-0501 treatment.

**[0037]** Figure 14 is a series of graphs and a table depicting the effect of NI-0501 administration on neutrophil count in patients.

**[0038]** Figure 15 is a series of graphs and a table depicting the effect of NI-0501 administration on platelet count in patients.

**[0039]** Figure 16 is a series of graphs and a table depicting the effect of NI-0501 administration on serum levels of ferritin in patients.

**[0040]** Figure 17 is a series of graphs and a table depicting the effect of NI-0501 administration on glucocorticoid tapering in patients.

**[0041]** Figure 18 is a graph depicting that administration of NI-0510 maintained IFN $\gamma$  neutralization until the time of HSCT. HLH response to NI-0501 treatment also persisted until

transplantation.

**[0042]** Figure 19 is a schematic representation of the screening, treatment, and follow-up portions of the studies presented in Example 8.

### **Detailed Description of the Invention**

**[0043]** The compositions and methods provided herein use a fully human IgG1 anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody (mAb) referred to herein as NI-0501, which binds and neutralizes IFN $\gamma$ . NI-0501 binds to soluble and receptor (IFN $\gamma$ R1)-bound forms of IFN $\gamma$ . Since NI-0501 is a human IgG1, it retains the characteristics of this immunoglobulin isotype, including the capacity to engage Fc $\gamma$  receptors and bind complement. IFN $\gamma$  is one of the most potent and pleiotropic cytokines of the immune system. It is critical for innate and adaptive immunity against viral and intracellular bacterial infections. After binding to its receptor, IFN $\gamma$  acts to produce a variety of physiological and cellular responses. Numerous studies over the last 20 years have associated IFN $\gamma$  with the pathogenesis and the maintenance of inflammatory diseases (see e.g., Billiau A. "Interferon-gamma: biology and role in pathogenesis." *Adv. Immunol.* 1996; 62:61-130; Schoenborn JR, Wilson CB. "Regulation of interferon-gamma during innate and adaptive immune responses." *Adv. Immunol.* 2007; 96:41-101; and Zhang SY, Boisson-Dupuis S, Chapgier A et al. "Inborn errors of interferon (IFN)-mediated immunity in humans: insights into the respective roles of IFN-alpha/beta, IFN-gamma, and IFN-lambda in host defense." *Immunol. Rev.* 2008; 226:29-40. IFN $\gamma$  is produced predominantly by natural killer (NK) and natural killer T (NKT) cells, as part of the innate immune response, and by CD4 Th1 and CD8 cytotoxic T lymphocyte (CTL) effector T cells, once antigen-specific immunity develops.

**[0044]** The compositions and methods provided herein are useful in the treatment of diseases, disorders and conditions associated with elevated IFN $\gamma$ . Diseases, disorders and conditions amenable to treatment or prevention using the compositions and methods of the invention include for example, hemophagocytic lymphohistiocytosis (HLH), graft vs. host disease, paraneoplastic cerebellar degeneration, hemorrhagic fever, sarcoidosis, adult onset Still's disease, transplant failure, transplant rejection, and/or an inflammatory disorder associated with transplant rejection. Transplant rejection included solid organ transplant disorder, bone marrow, acute graft rejection. In addition, the composition and methods are also useful in the treatment or alleviation of a side effect of CAR-T cell therapy.

**[0045]** HLH is a syndrome characterized by a severe impairment or absence of cytotoxic function

by NK and CD8+ T cells with striking activation of the immune system. HLH comprises primary (genetic/familial) HLH and secondary HLH, both clinically described by a dysregulation of the immune system leading to a profound hypercytokinemia with deleterious consequences on various tissues and organs (Henter JI, Elinder G, Soder O et al. “Hypercytokinemia in familial hemophagocytic lymphohistiocytosis.” Blood 1991; 78:2918-2922). HLH Classification is shown below in Table 9. HLH occurs in both adults and pediatric patients

Table 9. HLH classification

|                      |                                                     |                 |                |                                                   |
|----------------------|-----------------------------------------------------|-----------------|----------------|---------------------------------------------------|
| <b>Primary HLH</b>   | <b>Familial HLH (FLH)</b>                           |                 |                |                                                   |
|                      |                                                     | <b>Gene</b>     | <b>Protein</b> | <b>Function</b>                                   |
|                      | FLH-1                                               | Unknown         | Unknown        |                                                   |
|                      | FLH-2                                               | PRF1            | Perforin       | Pore-forming protein                              |
|                      | FLH-3                                               | UNC13D          | Munc13-4       | Vesicle priming                                   |
|                      | FLH-4                                               | STX11           | Syntaxin11     | Vesicle transport and fusion                      |
|                      | FLH-5                                               | STXBP2 (UNC18B) | Munc18-2       | Vesicle transport and fusion                      |
|                      | <b>Immune deficiency syndromes</b>                  |                 |                |                                                   |
|                      |                                                     | <b>Gene</b>     | <b>Protein</b> | <b>Function</b>                                   |
|                      | CHS                                                 | LYST            | Lyst           | Vesicle transport                                 |
|                      | GS-2                                                | RAB27A          | Rab27a         | Vesicle docking                                   |
|                      | XLP-1, XLP-2                                        | SH2DIA, BIRC4   | SAP, XIAP      | Signal transduction and activation of lymphocytes |
| <b>Secondary HLH</b> | Infections                                          |                 |                |                                                   |
|                      | Rheumatic diseases (Macrophage Activation Syndrome) |                 |                |                                                   |
|                      | Metabolic diseases                                  |                 |                |                                                   |
|                      | Malignancies                                        |                 |                |                                                   |

[0046] Primary HLH is a heterogeneous autosomal recessive disorder. Primary HLH is mostly seen in infancy and early childhood with an estimated prevalence in Europe of 1/50,000 live births (Henter JI, Elinder G, Soder O, Ost A. Incidence in Sweden and clinical features of familial hemophagocytic lymphohistiocytosis. Acta Paediatr. Scand. 1991; 80:428-435). The disease is invariably fatal with a median survival of less than 2 months after onset of symptoms, if untreated (Janka GE. Familial hemophagocytic lymphohistiocytosis. Eur. J.

Pediatr. 1983; 140:221-230; and Aricò M, Janka G, Fischer A, Henter JJ, Blanche S, Elinder G, Martinetti M, Rusca MP Hemophagocytic lymphohistiocytosis Report of 122 children from the International Registry. FHL Study Group of the Histiocyte Society. Leukemia. 1996 Feb; 10(2):197-203).

**[0047]** The impaired cytotoxic function present in HLH leads to hypercytokinemia and hemophagocytosis. These in turn cause all the typical symptoms of HLH (Dhote R, Simon J, Papo T et al. Reactive hemophagocytic syndrome in adult systemic disease: report of twenty-six cases and literature review. Arthritis Rheum. 2003; 49:633-639; Risdall RJ, McKenna RW, Nesbit ME et al. Virus-associated hemophagocytic syndrome: a benign histiocytic proliferation distinct from malignant histiocytosis. Cancer 1979; 44:993-1002; and Risdall RJ, Brunning RD, Hernandez JJ, Gordon DH. Bacteria-associated hemophagocytic syndrome. Cancer 1984; 54:2968-2972). Typical symptoms of HLH include, for example, prolonged fever, splenomegaly, hepatomegaly, cytopenia, hyperferritinemia, hypertriglyceridemia, hypofibrinogenemia, hemophagocytosis, hypercytokinemia, and/or lymphohistiocytic infiltrate, bone marrow hypoplasia, meningeal infiltrate.

**[0048]** Among the cytokines elevated in HLH patients are: IFN $\gamma$ , interleukin 6 (IL-6), IL-10, tumor necrosis factor (TNF)  $\alpha$ , IL-8, macrophage colony stimulating factor (M-CSF) and granulocyte-macrophage colony-stimulating factor (GM-CSF).

**[0049]** HLH can also occur during the course of an infection, a rheumatic or a neoplastic disease, and in this case, it is referred to as secondary HLH. Secondary HLH presents with the same signs and symptoms of primary forms and can be equally severe. The current treatment of secondary HLH is aimed at addressing the cause of the underlying disease. This is certainly the case for HLH caused by infections such as Leishmaniasis. Of note, the presence of certain infections, in particular viral infections such as those due to CMV or EBV, is very often the trigger for the manifestation of primary forms of HLH. This observation is also supported by the evidence that in animal models of primary HLH, infection with lymphocytic choriomeningitis virus (LCMV) is required for the development of the disease (Jordan MB, Hildeman D, Kappler J, Marrack P. An animal model of hemophagocytic lymphohistiocytosis (HLH): CD8<sup>+</sup> T cells and interferon gamma are essential for the disorder. Blood 2004; 104:735-743; Pachlopnik SJ, Ho CH, Chretien F et al. Neutralization of IFN $\gamma$  defeats haemophagocytosis in LCMV- infected perforin- and Rab27a-deficient mice. EMBO Mol. Med. 2009; 1:112-124; Kögl T, Müller J, Jessen B et al. Hemophagocytic lymphohistiocytosis in

syntaxin-11-deficient mice: T- cell exhaustion limits fatal disease. *Blood*. 2013; 121:604-613; and Sepulveda FE, Debeume F, Menasché G et al. Distinct severity of HLH in both human and murine mutants with complete loss of cytotoxic effector PRF1, RAB27A, and STX11 *Blood*. 2013;121:595-603).

**[0050]** When HLH manifests during a neoplastic disease, in particular a hematological malignancy, often the severity of the patient condition requires the immediate treatment of HLH, prior to specifically addressing the underlying disease.

**[0051]** The presence of signs and symptoms of HLH in patients suffering from a rheumatic disease, such as systemic Juvenile Idiopathic Arthritis (sJIA) and Systemic Lupus Erythematosus (SLE), is often referred to by rheumatologists as Macrophage Activation Syndrome (MAS) and can precede the appearance of the rheumatic disease itself. The majority of patients with MAS have impaired NK and perforin functional tests and a significant number of patients show polymorphisms or heterozygous mutations in PRF1 and UNC13D. Although it is an extremely severe and life threatening condition, usually it resolves when an adequate treatment is initiated, consisting in most cases of corticosteroids and cyclosporine. However, in approximately 15% of patients developing MAS, the disease can be difficult to control and the use of etoposide may be considered (Minoia F, Davi S, Horne AC et al. Clinical Features, Treatment, and Outcome of Macrophage Activation Syndrome Complicating Systemic Juvenile Idiopathic Arthritis: A Multinational, Multicenter Study of 362 Patients. *Arthritis & Rheumatism* 2014; 66: 3160–3169).

**[0052]** While primary HLH is recognized as predominantly a childhood disease, HLH is a condition that can be found in adults, and increased awareness indicates this may happen more often than recognized in the past. In the majority of adult patients the disease develops during malignancies (mainly non-Hodgkin lymphomas), infections, auto-inflammatory or autoimmune diseases and iatrogenic immune deficiencies.

**[0053]** There are currently no approved drugs for the treatment of HLH. However, experts in the field have established guidelines for the management HLH patients (Henter JI, Horne AC, Arico' M, Egeler RM, Filipovich AH, Imashuku S, Ladisch S, McClain K, Webb D, Winiarski J, and Janka Diagnostic and Therapeutic Guidelines for Hemophagocytic Lymphohistiocytosis *Blood Cancer* 2007; 48:124–13.1; Henter JI, Samuelsson-Horne A, Arico

M et al. Treatment of hemophagocytic lymphohistiocytosis with HLH-94 immunochemotherapy and bone marrow transplantation. *Blood* 2002; 100:2367-2373; and Jordan MB, Allen CE, Weitzman S, Filipovich AH, McClain KL. How I treat hemophagocytic lymphohistiocytosis. *Blood* 2011; 118:4041-4052).

**[0054]** The management of primary HLH patients currently comprises of the following steps (Henter et al., *Blood Cancer* 2007): (i) induction therapy of 8 weeks with a combination of corticosteroids and immunosuppressive drugs (e.g. etoposide, CsA, alemtuzumab, anti-thymocyte globulin); (ii) maintenance therapy up to transplantation; and (iii) transplantation for all patients with an identified genetic deficiency and eventually in very severe HLH cases with no disease-associated mutations.

**[0055]** The main goal of induction therapy is to suppress the life-threatening inflammatory process that characterizes HLH, enabling transplantation in those patients who require it (Horne A, Janka G, Maarten ER et al. Haematopoietic stem cell transplantation in haemophagocytic lymphohistiocytosis. *Br. J. Haematol.* 2005; 129:622-630). Transplantation is the only curative treatment for HLH associated with high penetrance genetic mutations (Henter et al., *Blood* 2002).

**[0056]** Despite the adoption of such guidelines the overall mortality rate for primary HLH remains around 40 to 50% (Henter et al., *Blood* 2002; Trottestam H, Horne A, Arico M et al. Chemoimmunotherapy for hemophagocytic lymphohistiocytosis: long-term results of the HLH-94 treatment protocol. *Blood* 2011; 118:4577-4584).

**[0057]** The need to use, during the induction period, drugs associated with severe short and long term-safety issues further contributes to the already high mortality. The compositions and methods provided herein were developed as a targeted treatment ensuring efficacy with less toxicity.

**[0058]** During the last years, growing evidence of the pivotal role of IFN $\gamma$  in the development of HLH has been generated (Henter JJ, Elinder G, Soder O et al. Hypercytokinemia in familial hemophagocytic lymphohistiocytosis. *Blood* 1991; 78:2918-2922; Jordan MB, Hildeman D, Kappler J, Marrack P. An animal model of hemophagocytic lymphohistiocytosis (HLH): CD8+ T cells and interferon gamma are essential for the disorder. *Blood* 2004; 104:735- 743; Pachlopnik SJ, Ho CH, Chretien F et al. Neutralization of IFN $\gamma$  defeats haemophagocytosis in LCMV-infected perforin- and Rab27a-deficient mice. *EMBO Mol. Med.* 2009; 1:112-124; Behrens EM, Canna SW, Slade K et al. Repeated TLR9 stimulation results in macrophage activation syndrome-

like disease in mice. *J. Clin. Invest* 2011; 121:2264-2277; Xu XJ, Tang YM, Song H, MD, Yang SL, Xu WQ, Zhao N, Shi SW, Shen HP, Mao JQ, Zhang LY, and Pan BH, Diagnostic Accuracy of a Specific Cytokine Pattern in Hemophagocytic Lymphohistiocytosis in Children *J Pediatr* 2011; and Risma K, Jordan MB. Hemophagocytic lymphohistiocytosis: updates and evolving concepts. *Curr. Opin. Pediatr.* 2012; 24:9-15).

**[0059]** The mutations of genes which characterize primary forms of HLH all affect proteins involved in the same process, ultimately impairing cytotoxic activity. Perforin mutations were the first identified in HLH patients.

**[0060]** Perforin knocked out (KO) mice are considered a relevant model for the human disease. In fact, these mice, once infected with LCMV, develop all the diagnostic and many of the clinical and laboratory characteristic features of the human disease, and they die if untreated. For these reasons, perforin KO mice have been used to study the pathophysiology of HLH. The HLH-like pathology that they develop is dependent on CD8<sup>+</sup> T cells and IFN $\gamma$  produced in response to antigen stimulation.

**[0061]** It was demonstrated that when the high circulating levels of IFN $\gamma$  are neutralized, with the administration of an anti-IFN $\gamma$  antibody, not only are the clinical and laboratory abnormalities reverted, but also survival rate is dramatically improved. On the contrary, the ablation of any other cytokine had no impact on survival (Jordan et al., *Blood* 2004; Pachlounik et al., *EMBO Mol. Med.* 2009).

**[0062]** Two models of secondary HLH have been investigated in the context of the NI-0501 development program. In one model, repeated administration of CpG (causing TLR9 stimulation) has been used to mimic a chronic severe hyperstimulation in healthy mice (i.e. with normal genetics of the cytotoxic pathway) as a model of HLH secondary to infection. Although these mice do not necessarily die, they develop typical clinical and laboratory features of HLH. When IFN $\gamma$  is neutralized, with the administration of an anti-IFN $\gamma$  antibody, clinical and laboratory features of the disease are reverted. Interestingly, in this model it has been demonstrated that administration of the anti-IFN $\gamma$  antibody leads to full neutralization of IFN $\gamma$  effects also in relevant target tissues, such as liver and spleen (manuscript in preparation).

**[0063]** To study the physiopathology of secondary HLH occurring in the context of rheumatic diseases, an animal model has been generated using IL-6 transgenic mice expressing high levels of IL-6, similarly to what occurs in patients with sJIA, the rheumatic disease most frequently associated with secondary forms of HLH. When triggered with Toll Like Receptor

(TLR) ligands, these mice die with many of the features of the human disease (Strippolli R, Carvalho F, Scianaro R et al. Amplification of the response to Toll-like receptor ligands by prolonged exposure to interleukin-6 in mice: Implication for the pathogenesis of macrophage activation syndrome. *Arthritis & Rheumatism* 2012; 64: 1680–1688). In these mice, when IFN $\gamma$  is neutralized with the administration of an anti-IFN $\gamma$  antibody, survival is markedly improved and laboratory parameters reverted (Prencipe G et al, manuscript in preparation).

**[0064]** Further strengthening the importance of IFN $\gamma$  in HLH are the high concentrations of circulating IFN $\gamma$  levels in primary HLH patients (Henter et al., *Blood* 1991; Xu et al., *J Pediatr* 2011). In a series of 71 patients monitored from HLH diagnosis to treatment and follow-up, IFN $\gamma$  levels were above the upper limit of normal (17.3 pg/mL) in all patients, and in particular 53.5% had levels above 1000 pg/mL. It was also reported that IFN $\gamma$  levels rise early and quickly, and can fall from > 5000 pg/mL to normal in 48 hours upon effective treatment of HLH.

**[0065]** More recently, in an observational study in patients with secondary forms of HLH, high levels of IFN $\gamma$  were demonstrated both in patients with HLH secondary to infections and in patients with HLH occurring in the context of sJIA. The levels of CXCL9, CXCL10 and CXCL11, three chemokines that are known to be induced by IFN $\gamma$ , were also significantly elevated. Noteworthy, levels of IFN $\gamma$ , and of the three IFN $\gamma$  chemokines, were found to be significantly correlated with laboratory parameters of disease severity, such as ferritin, platelet count and transaminases (Bracaglia et al., manuscript submitted).

**[0066]** As hypercytokinemia and organ infiltration by activated lymphocytes and histiocytes are responsible for all HLH symptoms and are dependent on CD8<sup>+</sup> T cells hyperactivity and high IFN $\gamma$  levels, the neutralization of IFN $\gamma$  constitutes a rational therapeutic approach. In fact, no agents specifically targeting CD8<sup>+</sup> T cells are available at the moment, and targeting individual cytokines downstream of IFN $\gamma$  would not necessarily be feasible.

**[0067]** Therefore, based on the data from animal models of primary and secondary HLH and from the observation made in patients with both primary and secondary HLH, confirming the critical role played by IFN $\gamma$  in the pathogenesis of this disease, the neutralization of IFN $\gamma$  offers a robust rationale to develop a targeted therapy for HLH, which must be effective with no or limited toxicity.

**[0068]** The disclosure also provides compositions and methods that are useful in identifying or otherwise refining a patient population suffering from a disorder, where the patient has an elevated level of CXCL9, alone or in combination with one or more additional interferon

(IFN $\gamma$ ) related biomarkers. In particular, the disclosure provides compositions and methods for detecting CXCL9 levels as a biomarker for IFN $\gamma$  production in hemophagocytic lymphohistiocytosis (HLH), in second HLH, and/or in macrophage activation syndrome (MAS).

**[0069]** A body of evidence in animal models points to a pivotal pathogenic role of IFN $\gamma$ , in primary hemophagocytic lymphohistiocytosis (HLH). High levels of IFN $\gamma$  are also found in humans with HLH. It has been previously reported that high levels of IFN $\gamma$  and of the three IFN $\gamma$ -related chemokine, CXCL9, CXCL10 and CXCL11, are observed in patients with active MAS, a form of secondary HLH that occurs in the context of systemic Juvenile Idiopathic Arthritis (sJIA) (*See e.g.*, Bracaglia C., Caiello I, De Graaf K., et al. *Pediatric Rheumatology* 2014,12(Suppl 1):O3). Indirect evidence in mice suggests that IFN $\gamma$  is mostly produced in peripheral tissues and blood concentrations may be relatively low.

**[0070]** The term macrophage activation syndrome (MAS) refers to a severe potentially fatal complication of chronic inflammatory rheumatic diseases. It occurs typically in the context of systemic juvenile idiopathic arthritis (sJIA) with 10-20% of patients developing this syndrome during the course of disease. It may occur also, albeit more rarely, in systemic lupus erythematosus, Kawasaki disease, as well as other autoimmune and autoinflammatory disorders. In sJIA, MAS occurs typically during active disease phases, including at disease onset. An infectious trigger can be identified in a high proportion of patients. Typical features of MAS include fever, splenomegaly, hemorrhages, and signs of liver, central nervous system and kidney involvement that may lead to multiple organ failure. Laboratory abnormalities include decrease in white blood cells, platelet and hemoglobin, hypertransaminasemia, marked increase in ferritin, and evidence for intravascular activation of the coagulation system (Ravelli, A., et al., *Macrophage activation syndrome as part of systemic juvenile idiopathic arthritis: diagnosis, genetics, pathophysiology and treatment*. *Genes Immun.* 13(4): p. 289-98). MAS causes significant morbidity and mortality accounting for a relevant portion of the deaths due to sJIA (Minoia, F., et al., *Clinical features, treatment, and outcome of macrophage activation syndrome complicating systemic juvenile idiopathic arthritis: a multinational, multicenter study of 362 patients*. *Arthritis Rheumatol*, 2014. 66(11): p. 3160-9; Hashkes, P.J., et al., *Mortality outcomes in pediatric rheumatology in the US*. *Arthritis Rheum*, 2010. 62(2): p. 599-608). A better understanding of disease pathogenesis, with the consequent identification of new therapeutic targets and the possible development of targeted therapies, may lead to significant improvements in the management and the outcome of MAS.

[0071] MAS shares the majority of the clinical features and laboratory abnormalities of haemophagocytic lymphohistiocytoses (HLH), and it is indeed currently classified among secondary or reactive HLH (sec-HLH) (Jordan, M.B., et al., *How I treat hemophagocytic lymphohistiocytosis*. Blood, 2011. 118(15): p. 4041-52). Primary forms of HLH (p-HLH) are caused by mutations of genes coding for proteins involved in granule exocytosis, including PRF1, UNC13D, STXBP2, STX11, RAB27A and XIAP, typically leading to defective cytotoxic activity of CD8<sup>+</sup> lymphocytes and NK cells. According to the current classification, in the absence of an identifiable genetic cause and/or of familial inheritance, HLH is defined as secondary or reactive. Sec-HLH can occur in the absence of a demonstrable trigger or in the context of infections, malignancies or rheumatic diseases, the latter being commonly referred to as MAS. The genetic basis for the development of MAS is being progressively unraveled, with a number of studies pointing to the association of MAS, and in general of sec-HLH, with heterozygosity for low penetrance variants or mutations of the same causative genes of p-HLH (Kaufman, K.M., et al., *Whole-exome sequencing reveals overlap between macrophage activation syndrome in systemic juvenile idiopathic arthritis and familial hemophagocytic lymphohistiocytosis*. Arthritis Rheumatol, 2014. 66(12): p. 3486-95; Vastert, S.J., et al., *Mutations in the perforin gene can be linked to macrophage activation syndrome in patients with systemic onset juvenile idiopathic arthritis*. Rheumatology (Oxford), 2010. 49(3): p. 441-9.; Zhang, K., et al., *Macrophage activation syndrome in patients with systemic juvenile idiopathic arthritis is associated with MUNC13-4 polymorphisms*. Arthritis Rheum, 2008. 58(9): p. 2892-6; and Zhang, M., et al., *Genetic defects in cytotoxicity in macrophage activation syndrome*. Curr Rheumatol Rep, 2014. 16(9): p. 439; and Bracaglia C, Sieni E, Da Ros M, et al. Mutations of familial hemophagocytic lymphohistiocytosis (FHL) related genes and abnormalities of cytotoxicity function tests in patients with macrophage activation syndrome (MAS) occurring in systemic juvenile idiopathic arthritis (sJIA). Pediatric Rheumatology 2014, 12(Suppl 1):P53). These similarities in the genetic background between p-HLH and MAS further support a shared pathogenic mechanism.

[0072] Studies in patients with p-HLH, as well as in murine models of p-HLH, support the hypothesis that defective cytotoxic activity and abnormalities in antigen-presenting cell (APC)-CD8<sup>+</sup> T cell crosstalk leads to defective silencing of the immune response and abnormal T cell activation. This results in uncontrolled immune activation and production of pro-inflammatory cytokines by T lymphocytes and macrophages, leading to organ damage. Studies in animal models of p-HLH performed in perforin and in Rab27 deficient mice point to a critical role of interferon-

gamma (IFN $\gamma$ ) produced by activated CD8<sup>+</sup> T cells. In perforin deficient mice, neutralization of IFN $\gamma$  leads to survival of an otherwise lethal syndrome, with reversal of biochemical and hematological abnormalities (Jordan, M.B., et al., *An animal model of hemophagocytic lymphohistiocytosis (HLH): CD8<sup>+</sup> T cells and interferon gamma are essential for the disorder*. Blood, 2004. 104(3): p. 735-43; Pachlopnik Schmid, J., et al., *Neutralization of IFN $\gamma$  defeats haemophagocytosis in LCMV-infected perforin- and Rab27a-deficient mice*. EMBO Mol Med, 2009. 1(2): p. 112-24). In Rab27 deficient mice, in which the disease does not lead to death, neutralization of IFN $\gamma$  causes a marked improvement of the involvement of peripheral organs, including the central nervous system (Pachlopnik 2009). High circulating IFN $\gamma$  levels are also found in patients with HLH diagnosed according to the HLH 2004 diagnostic guidelines (My, L.T., et al., *Comprehensive analyses and characterization of haemophagocytic lymphohistiocytosis in Vietnamese children*. Br J Haematol, 2010. 148(2): p. 301-10; Takada, H., et al., *Increased serum levels of interferon-gamma-inducible protein 10 and monokine induced by gamma interferon in patients with haemophagocytic lymphohistiocytosis*. Clin Exp Immunol, 2003. 133(3): p. 448-53; Tang, Y., et al., *Early diagnostic and prognostic significance of a specific Th1/Th2 cytokine pattern in children with haemophagocytic syndrome*. Br J Haematol, 2008. 143(1): p. 84-91; Xu, X.J., et al., *Diagnostic accuracy of a specific cytokine pattern in hemophagocytic lymphohistiocytosis in children*. J Pediatr, 2012. 160(6): p. 984-90 e1.), and therefore not necessarily on the basis of the presence of a genetic mutation. It should be noted that these studies included a significant, although variable, proportion of patients without a demonstrable genetic cause (Ibid).

**[0073]** The studies provided herein were designed to evaluate the correlation between serum levels of IFN $\gamma$  and of the three IFN $\gamma$  related chemokines with themselves and with laboratory parameters of disease activity in patients with active MAS in order to search for a biomarker of IFN $\gamma$  *in vivo* production. In particular, circulating levels of IFN $\gamma$ , CXCL9, CXCL10, CXCL11 and IL-6 were measured in patients with sJIA where about 37% (20 out of 54) of the patients had MAS at time of sampling. The relation of the circulating levels to disease activity parameters was also evaluated, as were the correlations of the levels of IFN $\gamma$  with those of CXCL9, CXCL10 and CXCL11. In some embodiments, the biomarker is total IFN $\gamma$  level, which is useful as a pharmacodynamic biomarker.

**[0074]** As demonstrated herein, levels of IFN $\gamma$  and of the 3 IFN $\gamma$ -related chemokines CXCL9, CXCL10, and CXCL11 were significantly elevated in active MAS as compared to active sJIA without MAS at sampling. In active MAS laboratory parameters of disease severity, such as

ferritin, neutrophils, platelets, alanine aminotransferase and lactate dehydrogenase, were significantly correlated with IFN $\gamma$  and CXCL9, and to a lesser extent with CXCL10 and CXCL11; no correlation with IL-6 levels was found. In patients with active sJIA without MAS there was no significant correlation between laboratory parameters and cytokine levels. In active MAS I IFN $\gamma$  levels were significantly correlated with levels of CXCL9, to a lesser extent with levels of CXCL10, and not with levels of CXCL11.

[0075] The high levels of IFN $\gamma$  and of CXCL9 present in patients with active MAS are significantly correlated with laboratory parameters of disease severity. In patients with active MAS IFN $\gamma$  and CXCL9 are tightly correlated. Since CXCL9 has been shown to be induced only by IFN $\gamma$  and not by other interferons (*see e.g.*, Groom J.R. and Luster A.D. Immunol Cell Biol 2011, Feb; 89(2):207-15), the findings disclosed herein demonstrate that CXCL9 is a biomarker of IFN $\gamma$  production in MAS.

[0076] The studies provided herein also demonstrate that levels of IFN $\gamma$  and of Chemokine (C-X-C Motif) Ligand 9 (CXCL9), CXCL10 and CXCL11, three chemokines that are known to be induced by IFN $\gamma$ , are elevated in patients with MAS complicating sJIA, but not in patients with active sJIA without MAS. Moreover, in these patients levels of IFN $\gamma$ , CXCL9, CXCL10 and CXCL11 were correlated with laboratory parameters of disease severity.

[0077] The disclosure provides compositions and methods using a neutralizing anti-IFN $\gamma$  antibody or antigen-binding fragment thereof to treat, prevent and/or delay the onset or progression of, or alleviate a symptom associated with transplant failure, transplant rejection, and/or an inflammatory disorder associated with transplant rejection such as solid organ transplant disorder, bone marrow acute graft rejection. The disclosure provides compositions and methods using a neutralizing anti-IFN $\gamma$  antibody or antigen-binding fragment thereof to treat, inhibit, delay the progression of, or otherwise ameliorate a symptom of Graft- versus-host disease (GvHD) in a subject who has received or is receiving a transplant comprising biological material or a series of transplants comprising biological material. The disclosure provides compositions and methods using a neutralizing anti-IFN $\gamma$  antibody or antigen-binding fragment thereof to prolong survival of transplanted biological material. The disclosure provides compositions and methods using a neutralizing anti-IFN $\gamma$  antibody or antigen-binding fragment thereof to treat, prevent and/or delay the onset or progression of, or alleviate a symptom associated paraneoplastic cerebellar degeneration, hemorrhagic fever, sarcoidosis, adult onset Still's disease or CART-T cell therapy.

[0078] The compositions and methods provided herein are useful in transplanting any biological

material, including, for example, cells, tissue(s), bone marrow, and/or organ(s), including, by way of non-limiting example, heart, kidney, pancreas, liver, and/or intestine. In some embodiments, the biological material to be transplanted is allogeneic biological material. In some embodiments, the transplanted biological material is bone marrow. In some embodiments, the transplanted biological material is a population of hematopoietic stem cells. In some embodiments, the biological material to be transplanted is or is derived from one or more hepatocytes.

**[0079]** In the compositions and methods provided herein, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with transplant failure, transplant rejection, and/or an inflammatory disorder associated with transplant rejection. In some embodiments, the transplant rejection, also referred to herein as transplant failure, is acute. In some embodiments, the transplant rejection is hyperacute.

**[0080]** Neutralizing anti-IFN $\gamma$  antibodies of the invention include, for example, the heavy chain complementarity determining regions (CDRs) shown below in Table 1A, the light chain CDRs shown in Table 1B, and combinations thereof. The amino acids encompassing the complementarity determining regions (CDR) as defined by Chothia et al. 1989, E.A. Kabat et al., 1991 are highlighted in underlined and italicized text below. (*See Chothia, C, et al., Nature 342:877-883 (1989); Kabat, EA, et al., Sequences of Protein of immunological interest, Fifth Edition, US Department of Health and Human Services, US Government Printing Office (1991).*)

Table 1A. VH CDR sequences from antibody clones that bind and neutralize IFN $\gamma$

| Clone Name    | VH CDR1                 | VH CDR2                             | VH CDR3                          |
|---------------|-------------------------|-------------------------------------|----------------------------------|
| NI-0501       | SYAMS<br>(SEQ ID NO: 1) | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2) | DGSSGWYVPHWFDP<br>(SEQ ID NO: 3) |
| AC1.2R3P2__A6 | SYAMS<br>(SEQ ID NO: 1) | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2) | DGSSGWYVPHWFDP<br>(SEQ ID NO: 3) |
| AC1R3P2__B4   | SYAMS<br>(SEQ ID NO: 1) | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2) | DHSSGWYVISGMDV<br>(SEQ ID NO: 9) |
| AD14R4P1__B9  | SYAMS<br>(SEQ ID NO: 1) | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2) | DLTVGGPWYYFDY<br>(SEQ ID NO: 11) |
| AD14R4P2__C9  | SYAMS<br>(SEQ ID NO: 1) | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2) | DGWNALGWLES<br>(SEQ ID NO: 14)   |

|               |                          |                                      |                                  |
|---------------|--------------------------|--------------------------------------|----------------------------------|
| AC1.4R4P2_C10 | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | DGSSGWYVPHWFDP<br>(SEQ ID NO: 3) |
| AC1.2R3P7_D3  | SNAMS<br>(SEQ ID NO: 20) | TLTGSGGTAYYADSVKG<br>(SEQ ID NO: 21) | GTELVGGGLDN<br>(SEQ ID NO: 22)   |
| AD1R2P2_D6    | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | DGWNALGWLES<br>(SEQ ID NO: 14)   |
| AC1.2R3P2_D8  | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | DGSSGWYVPHWFDP<br>(SEQ ID NO: 3) |
| AD1.3R3P6_E1  | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | RSFDSGGSFY<br>(SEQ ID NO: 28)    |
| AD1.3R3P5_F8  | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | VGSWYLEDFDI<br>(SEQ ID NO: 31)   |
| AD1.3R3P6_F9  | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | GGNYGDYFDYFDY<br>(SEQ ID NO: 34) |
| AD14R4P2_G7   | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | DGWNALGWLES<br>(SEQ ID NO: 14)   |
| AD1.1R3P3_G9  | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | DFWVITSGNDY<br>(SEQ ID NO: 39)   |
| AD1.3R3P6_G10 | SYAMS<br>(SEQ ID NO: 1)  | AISGSGGSTYYADSVKG<br>(SEQ ID NO: 2)  | DGWNALGWLES<br>(SEQ ID NO: 14)   |

Table 1B. VL CDR sequences from antibody clones that bind and neutralize IFN $\gamma$ 

| Clone Name    | VL CDR1                           | VL CDR2                   | VL CDR3                        |
|---------------|-----------------------------------|---------------------------|--------------------------------|
| NI-0501       | TRSSGSIASNYVQ<br>(SEQ ID NO: 4)   | EDNQRP<br>(SEQ ID NO: 5)  | QSYDGSNRWM<br>(SEQ ID NO: 6)   |
| AC1.2R3P2_A6  | TRSSGSIVSNYVQ<br>(SEQ ID NO: 7)   | EDNRFP<br>(SEQ ID NO: 8)  | QSYDGSNRWM<br>(SEQ ID NO: 6)   |
| AC1R3P2_B4    | TRSSGSIASNYVQ<br>(SEQ ID NO: 4)   | EDNQRP<br>(SEQ ID NO: 5)  | QSNDSNVV<br>(SEQ ID NO: 10)    |
| AD14R4F1_B9   | TRSSGSIVSNYVQ<br>(SEQ ID NO: 7)   | DDNQRP<br>(SEQ ID NO: 12) | QSYDSSNVV<br>(SEQ ID NO: 13)   |
| AD14R4P2_C9   | TRSGGSIGSYYVQ<br>(SEQ ID NO: 15)  | DDKKRP<br>(SEQ ID NO: 16) | QSYDSNNLVV<br>(SEQ ID NO: 17)  |
| AC1.4R4P2_C10 | TRSSGTIASNYVQ<br>(SEQ ID NO: 18)  | EDNQRP<br>(SEQ ID NO: 5)  | QSYDNSNHVV<br>(SEQ ID NO: 19)  |
| AC1.2R3P7_D3  | TGSGGSIAATNYVQ<br>(SEQ ID NO: 23) | EDNQRP<br>(SEQ ID NO: 5)  | QSYDSNNHHVV<br>(SEQ ID NO: 24) |
| AD1R2P2_D6    | TGSSGSIASNYVQ<br>(SEQ ID NO: 25)  | EDNQRP<br>(SEQ ID NO: 5)  | QSYDSSNQEVV<br>(SEQ ID NO: 26) |

|               |                                   |        |                                 |
|---------------|-----------------------------------|--------|---------------------------------|
| AC1.2R3P2_D8  | TRSSGSIVSNYVQ<br>(SEQ ID NO: 7)   | EDNQRP | QSYDSNFWV<br>(SEQ ID NO: 27)    |
| AD1.3R3P6_E1  | TRSSGYIASSYVQ<br>(SEQ ID NO: 105) | EDDRRP | QSYDDTTPWV<br>(SEQ ID NO: 30)   |
| AD1.3R3P5_F8  | TRSSGSIASNYVH<br>(SEQ ID NO: 32)  | EDNRRP | QSSDTTYHGGVV<br>(SEQ ID NO: 33) |
| AD1.3R3P6_F9  | TRSSGSIASNYVQ<br>(SEQ ID NO: 4)   | EDNQRP | QSYEGF<br>(SEQ ID NO: 35)       |
| AD14R4P2_G7   | TGRNGNIASNYVQ<br>(SEQ ID NO: 36)  | EDTQRP | QSSDSNRVL<br>(SEQ ID NO: 38)    |
| AD1.1R3P3_G9  | TRSSGSIASNYVQ<br>(SEQ ID NO: 4)   | EDNRRP | QSFSTNLVV<br>(SEQ ID NO: 40)    |
| AD1.3R3P6_G10 | AGSSGSIASNYVQ<br>(SEQ ID NO: 41)  | EDNQRP | QSYSYNNQVV<br>(SEQ ID NO: 42)   |

**[0081]** Exemplary antibodies of the invention include, for example, the anti-IFN $\gamma$  antibodies described in PCT Publication No. WO 2006/109191, the contents of which are hereby incorporated by reference in their entirety.

**[0082]** Exemplary antibodies of the invention include, for example, the antibody referred to herein as NI-0501, which binds the human IFN $\gamma$ . The heavy chain, light chain, variable heavy (VH) chain, and variable light (VL) chain sequences of the NI-0501 antibody are shown below, with the CDR sequences underlined in the VH and VL amino acid sequences:

NI-0501 Heavy chain nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTAT  
ATTACTGTGCGAAAGATGGTAGCAGTGGCTGGTACGTACCACACTGGTTCGACCCCTGGGGCCAGGGAAC  
CCTGGTCACCGTCTCCTCAGCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGC  
ACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGT  
GGAAGTCAAGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCCTACAGTCCTCAGGACTCTACTC  
CCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCAC  
AAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGTGACAAAACCTCACACATGCCCAC  
CGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCCTCTTCCCCCAAACCCAAGGACACCCT  
CATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAG

TTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACA  
GCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTG  
CAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA  
GAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGACCTGCC  
TGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAATA  
CAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTATAGCAAGCTCACCGTGGACAAG  
AGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGC  
AGAAGAGCCTCTCCCTGTCTCCGGGTAAATAG (SEQ ID NO: 43)

NI-0501 Heavy chain amino acid sequence:

EVQLLESGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAKDGSSGWYVPHWFDPWGQGLTVTVSSASTKGPSVFPLAPSSKS  
TSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNH  
KPSNTKVDKRVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVK  
FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR  
EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDK  
SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 44)

NI 0501 Light chain nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACTC  
GCAGCAGTGGCAGCATTGCCAGCAACTATGTGCAGTGGTACCAACAGCGCCCGGGCAGTCCCCCACCAC  
TGTCATCTATGAGGATAACCAGAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACAGCTCC  
TCCAATTCTGCCTCCCTCACCATCTCTGGGCTGAAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
ATGATGGCAGCAATCGTTGGATGTTCTGGCGGAGGGACCAAGCTGACCGTCCTAGGTCAGCCCAAGGCTGC  
CCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTC  
ATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAG  
TGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAGCAGCTACCTGAGCCTGACGCC  
TGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGACA  
GTGGCCCCTACAGAATGTTTCATAG (SEQ ID NO: 45)

NI 0501 Light chain amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVIYEDNQRP SGVPDRFSGSIDSS  
SNSASLTISGLKTEDEADYYCQSYDGSNRWMFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCL  
ISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKT  
VAPTECS (SEQ ID NO: 46)

NI-0501 Heavy chain variable region amino acid sequence:

EVQLLES GGGLVQPGGSLRLSCAASGFTFS SYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAK DGSSGWYVPHWFDPWGQGTILTVSS (SEQ ID NO: 47)

NI 0501 Light chain amino variable region acid sequence

NFMLTQPHSVSESPGKTVTIS CTRSSGSIASNYVQWYQQRPGSSPTTVIYEDNQRPSGVPDRFSGSIDSS  
SNSASLTISGLKTEDEADYYC QSYDGSNRWMFGGGTKLTVL (SEQ ID NO: 48)

**[0083]** Suitable anti-IFN $\gamma$  antibodies include the antibodies described in U.S. Patent 7,700,098, which is hereby incorporated by reference in its entirety. Several exemplary antibodies include the antibodies referred to therein as ARC1.2R3P2\_A6 (“A6”), ARC1.2R3P2\_B4 (“B4”), ARC1.2R3P2\_B9 (“B9”), ARC1.2R3P2\_C9 (“C9”), ARC1.2R3P2\_C10 (“C10”), ARC1.2R3P2\_D3 (“D3”), ARC1.2R3P2\_D6 (“D6”), ARC1.2R3P2\_D8 (“D8”), ARC1.2R3P2\_E1 (“E1”), ARC1.2R3P2\_F8 (“F8”), ARC1.2R3P2\_F9 (“F9”), ARC1.2R3P2\_G7 (“G7”), ARC1.2R3P2\_G9 (“G9”), and ARC1.2R3P2\_G10 (“G10”).

The sequences of these antibodies are shown below.

A6 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTAT  
ATTACTGTGCGAAAGATGGTAGCAGTGGCTGGTACGTACCACACTGGTTCGACCCCTGGGGCCGGGGCAC  
CCTGGTCACCGTCTCGAGT (SEQ ID NO: 49)

A6 VH amino acid sequence:

EVQLLES GGGLVQPGGSLRLSCAASGFTFS SYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAK DGSSGWYVPHWFDPWGRGTLTVSS (SEQ ID NO: 50)

A6 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACTC  
GCAGCAGTGGCAGCATTGTCAGCAACTATGTGCAGTGGTACCAACAGCGCCCGGGCAGTGCCCCACCAC  
TGTCATCTATGAGGATAACCGGAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACAGCTCC

TCCAATACTGCCTCCCTCACCATCTCTGGGCTGGAGGCTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
ATGATGGCAGCAATCGTTGGATGTTCTGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
NO: 51)

A6 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGSIVSNYVQWYQQRPGSAPTTVIYEDNRRPSGVPDRFSGSIDSS  
SNTASLTISGLEAEDEADYYCQSYDGSNRWFMFGGKLTVLG (SEQ ID NO: 52)

B4 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTAT  
ATTACTGTGCGAAAGATCATAGCAGTGGCTGGTACGTAATCTCCGGTATGGACGTCTGGGGCCGAGGGAC  
AATGGTCACCGTCTCGAGT (SEQ ID NO: 53)

B4 VH amino acid sequence:

EVQLLESGLLVQPGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAKDHSSGWYVISGMDVWGRGTMVTVSS (SEQ ID NO: 54)

B4 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACCC  
GCAGCAGTGGCAGCATTGCCAGCAACTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTCCCCCACCAC  
TGTGATCTCTGAGGATAACCAAAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCGTCGACAGCTCC  
TCCAACCTCTGCCTCCCTCACCATTTCTGGACTGAGGACTGAGGACGAGGCTGACTATTACTGTCAGTCTA  
ATGATTCCGACAATGTGGTTTTCTGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID NO: 55)

B4 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVISEDNQRPSGVPDRFSGSVDS  
SNSASLTISGLRTEDEADYYCQSNDSNDNVVFGGKLTVLG (SEQ ID NO: 56)

B9 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC

TCCAGAGACAATCCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
ATTACTGTGCGAAGGACCTAACAGTGGGTGGTCCCTGGTACTACTTTGACTACTGGGGCCAAGGAACCCT  
GGTCACCGTCTCGAGT (SEQ ID NO: 57)

B9 VH amino acid sequence:

EVQLLES~~GGGLVQPGGSLRL~~SCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNPKN~~TLYLQMNSL~~RAEDTAVYYCAKDLTVGGPWYYFDYWGQGT~~LVTVSS~~ (SEQ ID NO: 58)

B9 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACCC  
GCAGCAGTGGCAGCATTGTCAGCAACTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTGGCCCCACCAC  
TGTGATCTTTGACGATGACCAAAGACCCTCTGGGGTCCCTGGTTCGGTTCTCTGGCTCCCTCGACAGCTCC  
TCCAAC~~TCTGCCTCCCTC~~ACCATCTCTGGGCTGCAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
ATGATAGCAGCAATGTGGTATTCGGCGGGGGGACCAAGGTCACCGTCCTAGGT (SEQ ID NO: 59)

B9 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGSIVSNYVQWYQQRPGSAPTTVIFDD~~DQRPS~~GVPGRFSGSLDSS  
SNSASLTISGLQTEDEADYYCQSYDSSNVVFGGGTKVTVLG (SEQ ID NO: 60)

C9 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
ATTACTGTGCGAAAGATGGATGGAACGCGCTGGGATGGCTTGAATCCTGGGGCCGGGGCACCCTGGTCAC  
CGTCTCGAGT (SEQ ID NO: 61)

C9 VH amino acid sequence:

EVQLLES~~GGGLVQPGGSLRL~~SCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAKDGW~~NALGWLES~~WGRGTLTVTVSS (SEQ ID NO: 62)

C9 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAGGACGATAACCATCTCCTGCACCC  
GCAGTGGTGGCAGCATTGGCAGCTACTATGTGCAGTGGTACCAGCAGCGCCCGGGCACTGGCCCCACCAC  
TGTGATCTATGACGATAAAAAAAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACAGCTCC

TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTATTGTCAGTCTT  
ATGATAGCAACAATCTTGTGGTTTTTCGGCGGAGGGACCAAGGTCACCGTCCTAGGT (SEQ ID  
NO: 63)

C9 VL amino acid sequence:

NFMLTQPHSVSESPGRTITISCTRSGGSIGSYVQWYQQRPGTAPTTVIYDDKKRPSGVPDRFSGSIDSS  
SNSASLTISGLKTEDEADYYCQSYDSNNLVVFGGGTKVTVLG (SEQ ID NO: 64)

C10 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTAT  
ATTACTGTGCGAAAGATGGTAGCAGTGGCTGGTACGTACCACACTGGTTCGACCCCTGGGGCAGGGGGAC  
AATGGTCACCGTCTCGAGT (SEQ ID NO: 65)

C10 VH amino acid sequence:

EVQLLESGLLVQPGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAKDGS SGWYVPHWFDPWGRGTMVTVSS (SEQ ID NO: 66)

C10 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACCC  
GCAGCAGTGGCACCATTGCCAGCAACTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTCCCCCACCAC  
TGTGATCTATGAGGATAACCAAAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACAGCTCC  
TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
ATGATAACAGCAATCATTTGGGTGTTTCGGCGGAGGGACCAAGGTCACCGTCCTAGGT (SEQ ID  
NO: 67)

C10 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGTIASNYVQWYQQRPGSSPTTVIYEDNQRP SGVPDRFSGSIDSS  
SNSASLTISGLKTEDEADYYCQSYDNSNHVWVFGGGTKVTVLG (SEQ ID NO: 68)

D3 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCAGGGGGGTCCCTGAAACTCTCCTGTGCAG  
CCTCTGGATTCACCTTTAGCAGCAATGCCATGAGTTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG

GGTCTCAACTCTTACTGGTAGTGGTGGTACCGCATACTACGCAGACTCCGTGGAGGGCCGGTTCAGCATC  
TCCAGAGACAATTCCAAGAACACACTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTAT  
ATTACTGTGCGAAGGGCACGGAACCTCGTGGGAGGAGGACTTGACAACCTGGGGCCAAGGCACCCTGGTCAC  
CGTCTCGAGT (SEQ ID NO: 69)

D3 VH amino acid sequence:

EVQLLES~~GGGLVQPGGSLKLS~~CAASGFTFS~~SNAMSWVRQAPGKGLEWVSTLTGSGGTAYYADS~~VEGRFSI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAK~~GT~~ELVGGGLDNWGQGTILVTVSS (SEQ ID NO: 70)

D3 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTCTGTCTGGAGTCTCCGGGGAAGACGGTGACGATCTCCTGCACCG  
GCAGCGGAGGCAGCATTGCCACCAACTATGTGCAGTGGTATCAGCAGCGCCCGGGCAGTGCCCCCACCAC  
TGTGATCCATGAGGATAACCAAAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACGGCTCC  
TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGCAGCCTGAGGACGAGGCTGATTACTACTGTCAGTCTT  
ATGATAGTGACAATCATCATGTGGTATTCGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
NO: 71)

D3 VL amino acid sequence:

NFMLTQPHSLSESPGKTVTISCT~~SGSGSIATNYVQWYQQR~~PGSAPTTVIHEDN~~QRPSGVPDR~~FSGSIDGS  
SNSASLTISGLQPEDEADYYC~~QSYDS~~DNHHVVF~~GGG~~TKLTVLG (SEQ ID NO: 72)

D6 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
ATTACTGTGCGAAAGATGGATGGAACGCGCTGGGATGGCTTGAATCCTGGGGCAAGGGGACAATGGTCAC  
CGTCTCGAGT (SEQ ID NO: 73)

D6 VH amino acid sequence:

EVQLLES~~GGGLVQPGGSLRL~~SCAASGFTFS~~SYAMSWVRQAPGKGLEWVSAISGSGGSTYYADS~~VKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAK~~DGWNALGWLES~~WGKGTMTVTVSS (SEQ ID NO: 74)

D6 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCTGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACCG

GCAGCAGTGGCAGCATTGCCAGCAACTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTGCCCCCACCAC  
TGTGATCTATGAGGATAACCAAAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACAGCTCC  
TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
ATGATAGCAGCAATCAAGAGGTGGTATTCGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
NO: 75)

D6 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTGSSGSIASNYVQWYQQRPGSAPTTVIYEDNQRP SGVPDRFSGSIDSS  
SNSASLTISGLKTEDEADYYCQSYDSSNQEVVF GGGGTKLTVLG (SEQ ID NO: 76)

D8 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTAT  
ATTACTGTGCGAAAGATGGTAGCAGTGGCTGGTACGTACCACACTGGTTCGACCCCTGGGGCCAGGGAAC  
CCTGGTCACCGTCTCGAGT (SEQ ID NO: 77)

D8 VH amino acid sequence:

EVQLLES GGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
SRDNSKNTLYLQMNSLRAEDTAVYYCAKDGSSGWYVPHWFD PWGQGLTVTVSS (SEQ ID NO: 78)

D8 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTTACCATCTCCTGCACCC  
GCAGCAGTGGCAGCATTGTCAGCAACTATGTACAGTGGTACCAGCAGCGCCCGGGCAGTCCCCCACCAC  
TGTGATCTATGAGGATAACCAAAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACAGCTCC  
TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
ATGATAGCAACAATTTTTGGGTGTTTCGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
NO: 79)

D8 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGSIVSNYVQWYQQRPGSSPTTVIYEDNQRP SGVPDRFSGSIDSS  
SNSASLTISGLKTEDEADYYCQSYDSNNFWVF GGGGTKLTVLG (SEQ ID NO: 80)

E1 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
 CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
 GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
 TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
 ATTACTGTGTGAAAAGGTCCTTTGATAGTGGTGGGTCTTTGAGTACTGGGGCCAGGGGACAATGGTCAC  
 CGTCTCGAGT (SEQ ID NO: 81)

E1 VH amino acid sequence:

EVQLLES~~GGGLVQPGGSLRL~~SCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
 SRDNSKNTLYLQMNSLRAEDTAVYYCVKRSFDSGGSFEYWGQGTMTVTVSS (SEQ ID NO: 82)

E1 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTGCGAGTCTCCGGGGAAGACGGTCACCATCTCCTGCACCC  
 GCAGCAGTGGCTACATTGCCAGCTCCTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTCCCCCACCAC  
 TGTAACTCTTTGAGGATGACCGGAGACCCTCTGGGGTCCCTGATCGGTTCTCTGGCTCCATCGACGGCTCC  
 TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAGGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
 ATGATGACACCACTCCCTGGGTGTTCTGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
 NO: 83)

E1 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGYIASSYVQWYQQRPGSSPTTVIFEDDRRPSGVPDRFSGSIDGS  
 SNSASLTISGLRTEDEADYYCQSYDDTTPWVFGGGKLTVLG (SEQ ID NO: 84)

F8 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
 CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
 GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
 TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
 ATTACTGTGCGAGAGTCGGCAGCTGGTACCTGGAAGATTTTGATATCTGGGGCCGGGGGACAATGGTCAC  
 CGTCTCGAGT (SEQ ID NO: 85)

F8 VH amino acid sequence:

EVQLLES~~GGGLVQPGGSLRL~~SCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
 SRDNSKNTLYLQMNSLRAEDTAVYYCARVGSWYLEDFDIWGRGTMVTVSS (SEQ ID NO: 86)

F8 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCTGGAGTCTCCGGGGAAGACGGTTACCATCTCCTGCACCC  
 GCAGCAGTGGCAGCATTGCCAGCAACTATGTTCACTGGTATCAGCAGCGCCCGGGCAGTTCACCCACCAC  
 TGTGATCTATGAGGATAACCGAAGACCCTCTGGGGTCCCTGCTCGGTTCTCTGGCTCCATCGACAGCTCC  
 TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGGAGACTGACGACGAGGCTGACTACTACTGTCAGTCTT  
 CTGATAACCACCTATCATGGAGGTGTGGTATTCGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
 NO: 87)

F8 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTTRSSGSIASNYVHWYQQRPGSSPTTVIYEDNRRPSGVPARFSGSIDSS  
 SNSASLTISGLETDDEADYYCQSSDTTYHGGVVFGGGTKLTVLG (SEQ ID NO: 88)

F9 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
 CCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
 GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
 TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
 ATTACTGTGCGAAAGGCGGTAACTACGGTGATTACTTCGACTACTTTGACTACTGGGGCAGAGGGACAAT  
 GGTCACCGTCTCGAGT (SEQ ID NO: 89)

F9 VH amino acid sequence:

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
 SRDNSKNTLYLQMNSLRAEDTAVYYCAKGGNYGDFDYFDYWGRGTMVTVSS (SEQ ID NO: 90)

F9 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCTGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCACCC  
 GCAGCAGTGGCAGCATTGCCAGCAATTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTGCCCCCACCAT  
 TGTGATCTATGAAGATAACCAAAGACCCTCTGGGGTCCCTCATCGGTTCTCTGGCTCCATCGACAGCTCC  
 TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
 ATGAGGGGTTCGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID NO: 91)

F9 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTTRSSGSIASNYVQWYQQRPGSAPTIVIIYEDNQRPSTGVPHRFSGSIDSS  
 SNSASLTISGLKTEDEADYYCQSYEGFFGGGTKLTVLG (SEQ ID NO: 92)

G7 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
 CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
 GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACATC  
 TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
 ATTACTGTGCGAAAGATGGATGGAACGCGCTGGGATGGCTTGAATCCTGGGGCCAGGGGACAATGGTCAC  
 CGTCTCGAGT (SEQ ID NO: 93)

G7 VH amino acid sequence:

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
 SRDNSKNTLYLQMNSLRAEDTAVYYCAKDGNALGWLESWGQGTMTVTVSS (SEQ ID NO: 94)

G7 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACGCTGTGTCTGGAGTCTCCGGGGAAGACGGTGACCATTTCCTGCACCG  
 GCAGAAATGGCAACATTGCCAGCAACTATGTGCAGTGGTACCAGCAGCGCCCGGACAGTGCCCCACCCCT  
 TATAATCTTTGAAGATACCCAAAGACCCTCTGGGGTCCCTACTCGGCTCTCAGGCTCCATCGACACCTCC  
 TCCAATTCTGCCTCCCTCATCATCTCTTCATTGAGGACTGAGGACGAGGCTGATTACTACTGTCAATCTT  
 CTGATTCCAACAGGGTGCTGTTCTGGCGGAGGGACCAAGGTCACCGTCCTAGGT (SEQ ID NO: 95)

G7 VL amino acid sequence:

NFMLTQPHAVSESPGKTVTISCTGRNGNIASNYVQWYQQRPD SAPTLIIIFEDTQRPSGVPTRLSGSIDTS  
 SNSASLIISLRTED EADYYCQSSDSNRVLFGGGTKVTVLG (SEQ ID NO: 96)

G9 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
 CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
 GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
 TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
 ATTACTGTGCGAAAGATTTTTGGGTTATTACGAGTGGGAATGACTACTGGGGGCGGGGGACCACGGTCAC  
 CGTCTCGAGT (SEQ ID NO: 97)

G9 VH amino acid sequence:

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
 SRDNSKNTLYLQMNSLRAEDTAVYYCAKDFWVITSGNDYWGRGTTVTVSS (SEQ ID NO: 98)

G9 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTGACCATCTCCTGCACCC  
 GCAGCAGTGGCAGCATTGCTAGCAATTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTTCACCCACCAC  
 TGTGATCTTTGAAGATAACCGAAGACCCTCTGGGGTCCCTGATCGGTTTTCTGGCTCCATCGACACCTCC  
 TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTACTGTCAGTCTT  
 TTGATAGCACCAATCTTGTGGTGTTCGGCGGAGGGACCAAGCTGACCGTCCTAGGT (SEQ ID  
 NO: 99)

G9 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVIFEDNRRPSGVPDRFSGSIDTS  
 SNSASLTISGLKTEDEADYYCQSF DSTNLVVFGGGTKLTVLG (SEQ ID NO: 100)

G10 VH nucleic acid sequence:

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAG  
 CCTCTGGATTCACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG  
 GGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATC  
 TCCAGAGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGT  
 ATTACTGTGCGAAAGATGGATGGAACGCGCTGGGATGGCTTGAATCCTGGGGGAAGGGGACCACGGTCAC  
 CGTCTCGAGT (SEQ ID NO: 101)

G10 VH amino acid sequence:

EVQLLES GGGLVQPGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTI  
 SRDNSKNTLYLQMNSLRAEDTAVYYCAKD GWNALGWLESWGKGTTTVTVSS (SEQ ID NO: 102)

G10 VL nucleic acid sequence:

AATTTTATGCTGACTCAGCCCCACTCTGTGTCGGAGTCTCCGGGGAAGACGGTAACCATCTCCTGCGCCG  
 GCAGCAGTGGCAGCATTGCCAGCAACTATGTGCAGTGGTACCAGCAGCGCCCGGGCAGTGCCCCCACC GC  
 TGTGATCTATGAGGATAACCAAAGACCCTCTGGGGTCCCTGATCGATTCTCTGGCTCCATCGACAGCTCC  
 TCCAACCTCTGCCTCCCTCACCATCTCTGGACTGAAGACTGAGGACGAGGCTGACTACTACTGTCAATCTT  
 ACTCTTACAACAATCAGGTCGTGTTCGGCGGAGGGACCAAGGTCACCGTCCTAGGT (SEQ ID  
 NO: 103)

G10 VL amino acid sequence:

NFMLTQPHSVSESPGKTVTISCAGSSGSIASNYVQWYQQRPGSAPTAVIYEDNQRP SGVPDRFSGSIDSS

SNSASLTISGLKTEDEADYYCQSYSYNNQVVFGGGTKVTVLG (SEQ ID NO: 104)

**[0084]** In some embodiments, the IFN $\gamma$  antibodies are formatted in an IgG isotype.  
In some embodiments, the IFN $\gamma$  antibodies are formatted in an IgG1 isotype.

**[0085]** In some embodiments, IFN $\gamma$  antibodies of the invention specifically bind human and/or cynomolgus IFN $\gamma$ , wherein the antibody binds to the same epitope as the NI-0501 antibody, the A6 antibody, the B4 antibody, the B9 antibody, the C9 antibody, the C10 antibody, the D3 antibody, the D6 antibody, the D8 antibody, the E1 antibody, the F8 antibody, the F9 antibody, the G7 antibody, the G9 antibody, and/or the G10 antibody.

#### Methods of Treatment

**[0086]** The compositions and methods are useful in the treatment of any of a variety of disorders associated with interferon-gamma (IFN $\gamma$ ) expression and/or activity, including aberrant IFN $\gamma$  expression and/or activity. The compositions and methods of the disclosure are useful in the treatment of hemophagocytic lymphohistiocytosis (HLH). HLH is a rare, serious and life threatening disease of pathologic immune activation, characterized by clinical signs and symptoms of extreme inflammation (fever, splenomegaly, cytopenias, coagulopathy), leading to the development of abnormal immune-mediated pathologies which, through tissue damage, ultimately may cause multi-organ failure and death (Henter JJ, Elinder G, Söder O, Hansson M, et al: Hypercytokinemia in familial hemophagocytic lymphohistiocytosis. Blood 1991, 78:2918–2922). HLH comprises primary (genetic/familial) HLH and secondary HLH.

**[0087]** Primary HLH is a heterogeneous autosomal recessive disorder, mostly seen in infancy and early childhood with an estimated prevalence in Europe of 1/50,000 live births (Janka GE: Familial hemophagocytic lymphohistiocytosis. Eur. J. Pediatr. 1983, 140:221–230). The disease is invariably fatal with a median survival of less than 2 months after onset of symptoms, if untreated (Filipovich AH: Hemophagocytic lymphohistiocytosis (HLH) and related disorders. Hematology Am Soc Hematol Educ Program 2009:127–131).

**[0088]** The genetic defects in primary HHL all affect genes involved in cytotoxic pathway of NK-cells and/or cytotoxic lymphocytes required to eliminate activated macrophages, encoding proteins for perforin synthesis, cytolytic granule maturation, granule exocytosis and release granule exocytosis or function (Filipovich, A., K. McClain, and A. Grom. 2010. Histiocytic disorders: recent insights into pathophysiology and practical guidelines. Biol. Blood Marrow Transplant. 16(1 Suppl):S82–S89). In about 20-40% of primary HLH patients, the impaired

cytotoxic function characterizing the HLH syndrome is due to mutations in the gene encoding perforin (PRF1), a cytolytic protein of cytotoxic granules which is a key regulator for T-cell- and natural killer-cell-mediated cytotoxicity<sup>10</sup>. In about 10% of patients, the disease is caused by mutations in the UNC13D gene, encoding a protein which is involved in the release of perforin into the target cell. In addition, some immunodeficiency syndromes, e.g., Griscelli syndrome type 2 (GS-2) and Chediak-Higashi syndrome (CHS), present frequently with HLH (Janka GE, Lehmborg K: Hemophagocytic lymphohistiocytosis: pathogenesis and treatment. Hematology Am Soc Hematol Educ Program 2013, 2013:605–611).

[0089] Secondary forms of HLH can occur during the course of an infection, an autoimmune/rheumatic disease or in association to a malignancy. Secondary forms present with the same signs and symptoms of primary HLH and can be equally severe.

[0090] The compositions and methods of the disclosure are useful in the treatment of secondary HLH. The compositions and methods of the disclosure are useful in the treatment of macrophage activation syndrome (MAS).

[0091] MAS is a severe, potentially life-threatening complication of rheumatic diseases which is caused by excessive activation and expansion of T lymphocytes and macrophages. The uncontrolled expansion of these immune cells results in a marked hypercytokinemia and a hyperinflammatory state associated with fever, cytopenias, hepatosplenomegaly, liver dysfunction, coagulation abnormalities and hyperferritinemia, and may progress to multiple organ failure and death (Schulert GS, Grom AA: Pathogenesis of macrophage activation syndrome and potential for cytokine- directed therapies. Annu. Rev. Med. 2015, 66:145–159). [00161] Because of its strong clinical and pathological similarity to HLH, MAS is classified among the secondary or acquired forms of HLH. In fact, it has been recently demonstrated that the majority of patients with MAS have impaired NK and perforin functional tests and that a significant number of MAS patients show polymorphisms or heterozygous mutations in PRF1 and UNC13D (Zhang M, Behrens EM, Atkinson TP, Shakoory B, et al: Genetic defects in cytotoxicity in macrophage activation syndrome. Curr Rheumatol Rep 2014, 16:439).

[0092] MAS occurs most frequently in patients with sJIA and, more rarely with systemic lupus erythematosus (SLE), but is also described, though more rarely, in patients with vasculitis, particularly with Kawasaki disease. Approximately 7–17% of patients with SJIA develop overt MAS (Sawhney S, Woo P, Murray KJ: Macrophage activation syndrome: a potentially fatal complication of rheumatic disorders. Arch. Dis. Child. 2001, 85:421–426; Moradinejad MH, Ziaee

V: The incidence of macrophage activation syndrome in children with rheumatic disorders.

Minerva Pediatr. 2011, 63:459–466), some evidence suggests that subclinical MAS may be seen in as many as one third of patients with active systemic disease (Behrens EM, Beukelman T, Paessler M, Cron RQ: Occult macrophage activation syndrome in patients with systemic juvenile idiopathic arthritis. J. Rheumatol. 2007, 34:1133–1138).

[0093] Because MAS is potentially fatal, a timely diagnosis and immediate therapeutic intervention are essential for appropriate management of the disease. The reported mortality rates in MAS reach 20-30%, and it remains the major source of mortality in pediatric rheumatology (Grom AA, Horne A, De Benedetti F: Macrophage activation syndrome in the era of biologic therapy. Nat Rev Rheumatol. 2016 Mar 24. doi: 10.1038/nrrheum.2015.179).

[0094] Different sets of criteria have been proposed for the diagnosing of MAS in patients with sJIA. The HLH-2004 diagnostic guidelines (Henter J, Horne A, Aricó M, Egeler RM, et al: HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. Pediatr Blood Cancer 2007, 48:124–131), primarily developed for primary (genetic) forms of HLH, have sometimes been recommended. However, they present several limitations and may not apply to patients with sJIA. For example criteria such as cytopenias and hypofibrinogenemia below the thresholds required by HLH-2004 become evident only in the later stages of MAS, as these patients often have increased white blood cell and platelet counts as well as elevated serum levels of fibrinogen as a part of the sJIA inflammatory response (Schulert GS, Grom AA: Pathogenesis of macrophage activation syndrome and potential for cytokine- directed therapies. Annu. Rev. Med. 2015, 66:145–159). Hemophagocytosis may not be present in a significant proportion of patients with MAS at presentation (Minoia F, Davì S, Horne A, Demirkaya E, et al: Clinical features, treatment, and outcome of macrophage activation syndrome complicating systemic juvenile idiopathic arthritis: a multinational, multicenter study of 362 patients. Arthritis & rheumatology (Hoboken, N.J.) 2014, 66:3160–3169). Moreover, hemophagocytosis, NK cell activity and sCD25 are not routinely assessed in the context of MAS.

[0095] An alternative approach is based on the application of the preliminary diagnostic guidelines (PDG) for MAS complicating sJIA, which were created through the analysis of a cohort of patients with MAS compared with a group of patients with a flare of sJIA1.

[0096] Recently, the HLH-2004 diagnostic guidelines and the preliminary diagnostic guidelines for sJIA-associated MAS were compared for their capacity to discriminate sJIA/ MAS from sJIA (in the absence of MAS) and systemic infection in a

large patient population (Davì S, Minoia F, Pistorio A, Horne A, et al: Performance of current guidelines for diagnosis of macrophage activation syndrome complicating systemic juvenile idiopathic arthritis. *Arthritis & rheumatology* (Hoboken, N.J.) 2014, 66:2871–2880). Although with some limitations due to its retrospective nature, this study seems to indicate that the preliminary MAS guidelines achieve the best balance between sensitivity and specificity, and the best concordance with the diagnosis made by the treating physician. The sensitivity of the HLH-2004 set of criteria was <30%. Nevertheless, it has also been reported that the proportion of patients fulfilling each single criterion of the PDG is highly variable, and some clinical features (e.g. CNS dysfunction and hemorrhages) may manifest at a late stage of MAS, rendering their sensitivity low in incipient MAS (Lehmberg K, Pink I, Eulenburg C, Beutel K, et al: Differentiating macrophage activation syndrome in systemic juvenile idiopathic arthritis from other forms of hemophagocytic lymphohistiocytosis. *The Journal of pediatrics* 2013, 162:1245–1251).

[0097] More recently, a diagnostic score (HScore) has been developed and validated in a retrospective cohort of 312 patients, of whom 162 were judged to have reactive hemophagocytic syndrome (Fardet L, Galicier L, Lambotte O, Marzac C, Aumont C, Chahwan D, Coppo P, Hejblum G: Development and validation of the HScore, a score for the diagnosis of reactive hemophagocytic syndrome. *Arthritis & rheumatology* (Hoboken, N.J.) 2014, 66:2613–2620).

[0098] Nine variables (3 clinical [i.e., known underlying immunosuppression, high temperature, organomegaly], 5 biologic [i.e., triglyceride, ferritin, serum glutamic oxaloacetic transaminase, fibrinogen levels, and cytopenia], and 1 cytologic [i.e., hemophagocytosis features on bone marrow aspirate]), were retained in the HScore, and the probability of having hemophagocytic syndrome ranged from <1% with an HScore of  $\leq 90$  to >99% with an HScore of  $\geq 250$ .

[0099] Until a final consensus on validated diagnostic criteria for MAS is achieved, the clinical diagnosis by an expert physician is still key in the challenge to distinguish MAS from conditions presenting with overlapping features such as flares of SJIA or sepsis-like syndromes.

[0100] There are currently no approved drugs for the treatment of MAS. Usually, high-dose glucocorticoids are the first-line treatment for MAS. In patients failing to respond

to glucocorticoids, Cyclosporine A (CsA) has been proposed as additional treatment (Stéphan JL, Koné-Paut I, Galambrun C, Mouy R, Bader-Meunier B, Prieur AM: Reactive haemophagocytic syndrome in children with inflammatory disorders. A retrospective study of 24 patients. *Rheumatology (Oxford, England)* 2001, 40:1285–1292).

[0101] Being part of the HLH-94 treatment protocol developed for treating pHLH, the administration of etoposide is also considered in patients failing high dose glucocorticoids. However, the potential toxicity of the drug remains a major concern. Other current first line HLH treatments include dexamethasone. However, treatments such as etoposide and/or dexamethasone are myelosuppressive and/or broadly immune suppressive. There is currently no standard of care for second line HLH treatment, and treatments such as alemtuzumab/ATG are profoundly immunosuppressive, and survival is thought to be very poor with these treatments.

[0102] The utility of biologics inhibiting the IL-1, IL-6R or TNF $\alpha$  pathways in the treatment of MAS still remains unclear. Although biologics inhibiting these pathways have been reported to be effective in isolated cases, there are also reports of patients developing MAS in the setting of these treatments (Stern A, Riley R, Buckley L: Worsening of macrophage activation syndrome in a patient with adult onset Still's disease after initiation of etanercept therapy. *J Clin Rheumatol* 2001, 7:252–256; Ramanan AV, Schneider R: Macrophage activation syndrome following initiation of etanercept in a child with systemic onset juvenile rheumatoid arthritis. *J. Rheumatol.* 2003, 30:401–403; De Benedetti F, Brunner HI, Ruperto N, Kenwright A, et al: Randomized trial of tocilizumab in systemic juvenile idiopathic arthritis. *N. Engl. J. Med.* 2012, 367:2385–2395; Ruperto N, Brunner HI, Quartier P, Constantin T, et al: Two randomized trials of canakinumab in systemic juvenile idiopathic arthritis. *N. Engl. J. Med.* 2012, 367:2396–2406), as well as patients who do not respond to these treatments, indicating that inhibition of IL-1, IL- 6R or TNF $\alpha$  does not provide full protection against MAS development nor an efficacious treatment of the full blown syndrome.

[0103] A large retrospective, multicenter study has investigated the clinical, laboratory, and histopathological characteristics as well as current treatment and outcome of MAS/sJIA in a total of 362 patients (Minoia F, Davì S, Horne A, Demirkaya E, et al:

Clinical features, treatment, and outcome of macrophage activation syndrome complicating systemic juvenile idiopathic arthritis: a multinational, multicenter study of 362 patients. *Arthritis & rheumatology* (Hoboken, N.J.) 2014, 66:3160–3169). In approximately half of the patients, MAS occurred in the context of active sJIA or during a sJIA flare in 30% of them at disease onset. An infectious trigger was identified in one third of the patients. Among the 24 patients for whom the type of infection was reported, EBV was the most common causative agent (25%). In 11 patients (3.8%), MAS was believed to be related to a treatment side effect: 8 of these involved a biologic agent targeting the IL-6 (N=4), IL-1 (N=3) or TNF $\alpha$  (N=1) pathway. Nearly all patients were given glucocorticoids. Cyclosporine, biologic medications and etoposide were given to 61%, 15% and 12% of the patients respectively.

**[0104]** The identification of effective therapeutic regimens for MAS therefore represents an area of unmet high medical need. More than 50% of patients with sJIA and MAS do not respond to systemic glucocorticoids alone, or may require prolonged treatment at high doses with associated significant morbidity. When patients fail to respond to glucocorticoids, no good evidence-based data is available on the effectiveness of additional treatments such as CsA or etoposide. The course of MAS may become rapidly irreversible leading to a fatal outcome.

Current data suggest that the mortality of sJIA-associated MAS is 8%, with about one third of the patients requiring ICU admission. Recent findings on the pivotal role of IFN $\gamma$  in the pathogenesis of the disease suggest IFN $\gamma$  blockage to potentially represent a novel therapeutic target.

**[0105]** The compositions, including NI-0501 compositions, and methods of the disclosure are advantageous over current therapies for primary and secondary HLH.

**[0106]** MAS and HLH are characterized by sustained immune cell activation and an associated cytokine storm of proinflammatory cytokines with overproduction of IFN $\gamma$ , TNF $\alpha$ , IL-1 and IL-6 (Henter JI, Elinder G, Söder O, Hansson M, et al: Hypercytokinemia in familial hemophagocytic lymphohistiocytosis. *Blood* 1991, 78:2918–2922; Imashuku S, Hibi S, Fujiwara F, Todo S: Hyper-interleukin (IL)-6-naemia in haemophagocytic lymphohistiocytosis. *Br. J. Haematol.* 1996, 93:803–807; Xu X, Tang Y, Song H, Yang S, et al: Diagnostic accuracy of a specific cytokine pattern in hemophagocytic lymphohistiocytosis in children. *J. Pediatr.* 2012, 160:984-90.e1; Put K, Avau A, Brisse E, Mitera T, et al: Cytokines in systemic juvenile idiopathic arthritis and haemophagocytic lymphohistiocytosis: tipping the balance between interleukin-18 and interferon-

$\gamma$ . Rheumatology (Oxford) 2015). During the last years, evidence has been accumulating in support of the pivotal role of IFN $\gamma$  in the development of both HLH (Jordan MB, Hildeman D, Kappler J, Marrack P: An animal model of hemophagocytic lymphohistiocytosis (HLH): CD8<sup>+</sup> T cells and interferon gamma are essential for the disorder. Blood 2004, 104:735–743; Pachlopnik Schmid J, Ho C, Chrétien F, Lefebvre JM, et al: Neutralization of IFN $\gamma$  defeats haemophagocytosis in LCMV-infected perforin- and Rab27a-deficient mice. EMBO Mol Med 2009, 1:112–124; Zoller EE, Lykens JE, Terrell CE, Aliberti J, et al: Hemophagocytosis causes a consumptive anemia of inflammation. J. Exp. Med. 2011, 208:1203–1214) and MAS (Behrens EM, Canna SW, Slade K, Rao S, et al: Repeated TLR9 stimulation results in macrophage activation syndrome-like disease in mice. J. Clin. Invest. 2011, 121:2264–2277).

[0107] For primary HLH, perforin knock-out mice are considered a relevant model as these mice, once infected with LCMV, develop all the diagnostic and many of the clinical and laboratory characteristic features of the human disease. The HLH-like disease that they develop is dependent on CD8<sup>+</sup> T cells and IFN $\gamma$  produced in response to antigen stimulation (Imashuku S, Hibi S, Fujiwara F, Todo S: Hyper-interleukin (IL)-6-naemia in haemophagocytic lymphohistiocytosis. Br. J. Haematol. 1996, 93:803–807). It was demonstrated that when the high circulating levels of IFN $\gamma$  are neutralized with the administration of an anti-IFN $\gamma$  antibody, not only are the clinical and laboratory abnormalities reverted, but also survival rate is dramatically improved. On the contrary, the ablation of many other cytokines had no impact on survival (Imashuku S, Hibi S, Fujiwara F, Todo S: Hyper-interleukin (IL)-6-naemia in haemophagocytic lymphohistiocytosis. Br. J. Haematol. 1996, 93:803–807; Xu X, Tang Y, Song H, Yang S, et al: Diagnostic accuracy of a specific cytokine pattern in hemophagocytic lymphohistiocytosis in children. J. Pediatr. 2012, 160:984-90.e1). Further strengthening the importance of IFN $\gamma$  in HLH are the high concentrations of circulating IFN $\gamma$  levels found in these patients (Henter JJ, Elinder G, Söder O, Hansson M, et al: Hypercytokinemia in familial hemophagocytic lymphohistiocytosis. Blood 1991, 78:2918–2922; Xu X, Tang Y, Song H, Yang S, et al: Diagnostic accuracy of a specific cytokine pattern in hemophagocytic lymphohistiocytosis in children. J. Pediatr. 2012, 160:984-90.e1). In a series of 71 patients monitored from HLH diagnosis to treatment and follow-up, IFN $\gamma$  levels were above the upper limit of normal (17.3 pg/mL) in all patients, and in particular 53.5% had levels above 1000 pg/mL. It was also reported that IFN $\gamma$  levels rise early and quickly, and can fall from > 5000 pg/mL to normal in 48 hours upon effective treatment of HLH.

[0108] Two animal models of secondary HLH have been investigated in the context of the NI-

0501 development program to elucidate the potential pathogenetic role of IFN $\gamma$ . First, in a murine model that mimics an infection-driven HLH, repeated administrations of CpG via activation of TLR9 triggered a hypercytokinemia that led to clinical (e.g. body weight loss, splenomegaly) and laboratory (e.g. cytopenia, hyperferritinemia) features of HLH33. When IFN $\gamma$  was neutralized by the administration of an anti-IFN $\gamma$  antibody, clinical and laboratory features of the disease were reverted. The neutralization of IFN $\gamma$  was shown to be complete also in relevant target tissues, such as the liver and the spleen. Interestingly, the administration of the anti-IFN $\gamma$  antibody unveiled an amount of IFN $\gamma$  500- to 2,000-fold higher than that measured in blood, likely to better reflect the IFN $\gamma$  production in tissues. The two IFN $\gamma$ -inducible chemokines (CXCL9 and CXCL10) were upregulated after TLR9 stimulation both in blood and in liver, and a significant correlation was observed between serum levels of IFN $\gamma$  with CXCL9 and CXCL10 serum concentrations. The neutralization of IFN $\gamma$  induced a significant decrease of serum CXCL9 and CXCL10, and of their mRNA levels in the liver (Buatois V, Chatel L, Cons L, Lory S, et al: IFN $\gamma$  drives disease in the TLR9-mediated secondary HLH in mice: rationale for a new therapeutic target in secondary HLH, in preparation).

[0109] Second, an animal model of IL-6 transgenic mice expressing high levels of IL-6 has been studied, since it mimics the condition of patients with sJIA, the rheumatic disease most frequently associated with secondary forms of HLH. When triggered with Toll-Like Receptor (TLR) ligands, increased lethality, increased inflammatory cytokine production and hyperactivation of inflammatory signaling pathways was observed. Moreover, these mice showed a drop in platelet and neutrophil counts, increased sCD25, ferritin and LDH levels, resembling many of the features typically present in patients with MAS (Strippoli R, Carvello F, Scianaro R, De Pasquale L, et al: Amplification of the response to Toll-like receptor ligands by prolonged exposure to interleukin-6 in mice: implication for the pathogenesis of macrophage activation syndrome. *Arthritis Rheum.* 2012, 64:1680-1688). In these mice, when IFN $\gamma$  is neutralized with the administration of an anti-IFN $\gamma$  antibody, survival is markedly improved and laboratory parameters reverted (Prencipe G et al, manuscript in preparation).

[0110] Similar evidence has been recently gathered in an observational study conducted in patients with secondary forms of HLH, either secondary to infections, or of unknown origin (pHLH having been excluded by normal cytotoxic activity, absence of mutation in known genes causing pHLH and absence of family history) or with MAS occurring in the context of sJIA.

[0111] In 14 patients with secondary HLH (in 7 of whom an underlying infection was identifiable),

serum samples were analyzed during active full blown disease and during disease remission. Levels of IFN $\gamma$ , CXCL9 and CXCL10 were markedly higher in the active phase compared to disease remission (IFN $\gamma$ : 34.7 vs. <3.5 pg/ml; CXCL9: 33598 vs. 745 pg/ml; CXCL10: 4420 vs. 132 pg/ml; median values). IFN $\gamma$  levels significantly correlated with the levels of CXCL9 ( $p=0.0018$ ) and, to a lesser extent, of CXCL10 ( $p=0.014$ ). The levels of IFN $\gamma$  and chemokines (in particular CXCL9) correlated significantly with parameters of disease severity, such as neutrophil and platelet counts, ferritin and ALT, further supporting the pathogenic role of IFN $\gamma$  in secondary HLH and the potential use of chemokines as relevant biomarkers of the disease (Buatois V, Chatel L, Cons L, Lory S, et al: IFN $\gamma$  drives disease in the TLR9-mediated secondary HLH in mice: rationale for a new therapeutic target in secondary HLH).

[0112] Similar findings have been shown in patients with MAS occurring in patients with sJIA. Serum concentrations of IFN $\gamma$ , IFN $\gamma$ -inducible chemokines (CXCL9, CXCL10, CXCL11) and IL-6 were measured in 54 patients with sJIA, of whom 20 had MAS. The levels of IL-6 were comparable in patients with full-blown MAS and those with active sJIA but without MAS at the time of sampling. On the contrary, circulating IFN $\gamma$  and chemokine levels were significantly higher in MAS, particularly for CXCL9, whose median levels were approximately 15-fold higher compared to patients with active sJIA without MAS (13392 vs. 837 pg/mL;  $p=0.005$ ). Noteworthy, a significant correlation was demonstrated only in patients with MAS between CXCL9 levels and parameters typically abnormal such as ferritin ( $p=0.041$ ), neutrophil ( $p=0.010$ ) and platelet ( $p=0.022$ ) counts, ALT ( $p=0.044$ ) and LDH ( $p=0.013$ ). Levels of IFN $\gamma$  also correlated with laboratory parameters of disease severity, with the exception of LDH for which statistical significance was not achieved (Bracaglia et al., manuscript in preparation).

[0113] All together these data provide a robust rationale for the neutralization of IFN $\gamma$  as targeted therapy for secondary HLH and MAS, and for its investigation in the clinical setting.

[0114] The compositions, including NI-0501 compositions, and methods of the disclosure are advantageous over current therapies for sJIA. For example, the compositions, including NI-0501 compositions, and methods of the disclosure are useful in treating MAS/sHLH in sJIA patients, with the primary objective of achieving MAS remission.

[0115] The rationale for the identification of this patient population as benefiting from a treatment with NI-0501 and for evaluating the efficacy of NI-0501 in MAS/sHLH is based on a number of factors. First, pre-clinical data obtained in an animal model relevant for MAS in sJIA has shown that IFN $\gamma$  neutralization markedly improved survival and reverted alterations of

laboratory parameters. Next, the observational data in patients with MAS/sHLH demonstrate the presence of high levels of IFN $\gamma$  and, more importantly, extremely elevated levels of IFN $\gamma$ -induced chemokines CXCL9, CXCL10 and CXCL11. Third, in patients with MAS/sHLH the concentrations of IFN $\gamma$  and CXCL9 significantly correlate with disease parameters such as ferritin, platelet count and transaminases. Next, the favorable tolerability profile and absence of relevant safety concerns observed in pHLH patients in previous studies in which all infusions administered were well tolerated, confirming the observations made in Healthy Volunteers, no infections caused by pathogens known to be favored by the neutralization of IFN $\gamma$  were reported, and none of the infections that occurred in some of the pHLH patients were considered related to NI-0501 treatment, rather to their immune status, disease duration and previous or concomitant treatments. Fifth, the preliminary data of the previous clinical studies shows a favorable impact on disease parameters, with appreciable onset of effects within the first days of treatment: typical clinical signs and symptoms of HLH started to improve rapidly after the first administration of NI-0501 (fever within hours, spleno/hepatomegaly within days); and of the 18 evaluable patients at the cut-off, treatment with NI-0501 enabled 10 patients to move to HSCT. Next, evidence from the PK modeling and simulation approach shows a predictable pharmacokinetic profile of NI-0501, and that neutralization of IFN $\gamma$  is achieved and maintained. Finally, conventional therapy (e.g. CsA) can immediately be initiated without the need for a wash-out period, in case NI-0501 would not control the disease adequately.

[0116] In conclusion, there is a strong rationale for neutralizing IFN $\gamma$  in MAS/sHLH secondary to rheumatic diseases based on pre-clinical and clinical evidence, and the preliminary data in pHLH patients indicates a favorable benefit risk profile of NI-0501 with a significant improvement to normalization of HLH features.

[0117] Thus, NI-0501 represents an innovative and effective therapeutic approach in the management of this severe, life-threatening complication of rheumatic diseases, potentially limiting side effects from long-term high dose glucocorticoid treatment.

#### Administration of anti-IFN $\gamma$ antibodies

[0118] It will be appreciated that administration of therapeutic entities in accordance with the invention will be administered with suitable carriers, excipients, and other agents that are incorporated into formulations to provide improved transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences (15th ed., Mack Publishing Company, Easton,

PA (1975)), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as Lipofectin™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in treatments and therapies in accordance with the present invention, provided that the active ingredient in the formulation is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. *See also* Baldrick P. “Pharmaceutical excipient development: the need for preclinical guidance.” *Regul. Toxicol Pharmacol.* 32(2):210-8 (2000), Wang W. “Lyophilization and development of solid protein pharmaceuticals.” *Int. J. Pharm.* 203(1-2):1-60 (2000), Charman WN “Lipids, lipophilic drugs, and oral drug delivery- some emerging concepts.” *J Pharm Sci.* 89(8):967-78 (2000), Powell *et al.* “Compendium of excipients for parenteral formulations” *PDA J Pharm Sci Technol.* 52:238-311 (1998) and the citations therein for additional information related to formulations, excipients and carriers well known to pharmaceutical chemists.

**[0119]** Efficaciousness of treatment is determined in association with any known method for diagnosing or treating the particular immune-related disorder. Alleviation of one or more symptoms of the immune-related disorder indicates that the antibody confers a clinical benefit.

**[0120]** Antibodies of the invention, including polyclonal, monoclonal, humanized and fully human antibodies, may be used as therapeutic agents. Such agents will generally be employed to treat or prevent a disease or pathology associated with aberrant expression or activation of a given target in a subject. An antibody preparation, preferably one having high specificity and high affinity for its target antigen, is administered to the subject and will generally have an effect due to its binding with the target. Administration of the antibody may abrogate or inhibit or interfere with the signaling function of the target. Administration of the antibody may abrogate or inhibit or interfere with the binding of the target with an endogenous ligand to which it naturally binds.

**[0121]** A therapeutically effective amount of an antibody of the invention relates generally to the amount needed to achieve a therapeutic objective. As noted above, this may be a binding interaction between the antibody and its target antigen that, in certain cases, interferes with the functioning of the target. The amount required to be administered will furthermore depend on the binding affinity of the antibody for its specific antigen, and will also depend on

the rate at which an administered antibody is depleted from the free volume other subject to which it is administered. Common ranges for therapeutically effective dosing of an antibody or antibody fragment of the invention may be, by way of nonlimiting example, from about 0.1 mg/kg body weight to about 50 mg/kg body weight. Preferred doses include 1, 3, 6, 10 mg/kg body weight. Common dosing frequencies may range, for example, from once daily to twice a week. Treatment may last 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or more weeks.

[0122] Antibodies or a fragment thereof of the invention can be administered for the treatment of a variety of diseases and disorders in the form of pharmaceutical compositions. Principles and considerations involved in preparing such compositions, as well as guidance in the choice of components are provided, for example, in Remington: The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa.: 1995; Drug Absorption Enhancement: Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4), 1991, M. Dekker, New York.

[0123] The formulation can also contain more than one active compound, *e.g.*, anti-IFN $\gamma$  antagonist as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[0124] In one embodiment, the active compound, *e.g.*, an anti-IFN $\gamma$  antagonist, is administered in combination therapy, *i.e.*, combined with one or more additional agents that are useful for treating pathological conditions or disorders. The term “in combination” in this context means that the agents are given substantially contemporaneously, either simultaneously or sequentially. If given sequentially, at the onset of administration of the second compound, the first of the two compounds is preferably still detectable at effective concentrations at the site of treatment.

[0125] For example, the combination therapy can include one or more neutralizing anti-IFN $\gamma$  antibodies of the invention coformulated with, and/or coadministered with, one or more additional therapeutic agents, *e.g.*, one or more cytokine and growth factor inhibitors, immunosuppressants, anti-inflammatory agents, metabolic inhibitors, enzyme inhibitors, and/or

cytotoxic or cytostatic agents, as described in more detail below. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

[0126] In some embodiments, the additional agent is an immunosuppressive agent. In some embodiments, the immunosuppressive agent is Cyclosporin A (CsA). In some embodiments, the subject has been receiving CsA prior to administration of NI-0501. In some embodiments, the additional agent includes at least etoposide. In some embodiments, the subject has been receiving etoposide prior to administration of NI-0501.

[0127] In some embodiments, the additional agent is intrathecal methotrexate and/or glucocorticoids. In some embodiments, the subject has been receiving intrathecal methotrexate and/or glucocorticoids prior to administration of NI-0501.

[0128] In some embodiments, the additional agent is IV immunoglobulins (IVIG). In some embodiments, the IVIG is administered as replacement treatment in a subject with a documented immunoglobulin deficiency. In some embodiments where the subject has a documented immunoglobulin deficiency, IVIG is administered given at a dose of 0.5 g/kg, every 4 weeks or more frequently in order to maintain adequate IgG levels.

[0129] In some embodiments, the one or more additional agents is analgesic treatment, transfusion of blood products, electrolyte and glucose infusions, antibiotics, anti-fungal and anti-viral treatment and/or general supportive care.

[0130] Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein and/or the smallest inhibitory fragment that interferes with or otherwise antagonizes IFN $\gamma$  signaling is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. (*See, e.g.,* Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993)). The formulation can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

### Dosing Regimens

**[0131]** The invention further provides dosing regimen for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with elevated IFN- $\gamma$ . The dosing regimens are multiple variable dose regimens. The dose between 1.0 to 10 mg/kg of the subject's body weight of an antibody that binds interferon gamma (IFN $\gamma$ ). The dose is administered in at least a first and second dose. The second dose is lower or higher than the first dose.

**[0132]** A multiple variable dose regimen for the treatment of primary hemophagocytic lymphohistiocytosis (HLH) in a human subject includes an induction or a first dose and a treatment or second dose of an anti- IFN-g monoclonal antibody. The subject is an adult subject or a pediatric subject.

**[0133]** The first dose is 1.0 or 3.0, mg/kg body weight. The second dose is 3.0, 6.0 or 10.0 mg/kg body weight. The first dose is 1.0 mg/kg body weight an antibody and the second does is 3.0, 6.0 or 10.0 mg/kg body weight. Alternatively, the first dose is 3.0 mg/kg body weight an antibody and the second does is 6.0 or 10.0 mg/kg body weight.

**[0134]** The first dose is administered once as a single dose. Alternatively the first dose is administer more than once for an induction treatment period.

**[0135]** The second dose is administered for one or more treatment periods. The second dose is administered for a first treatment period. During the first treatment period the second dose is administered every three days after the first dose. The first treatment period last about 1, 2, 3, 4, 5, 6 or more weeks. Preferably the first treatment period is 2 weeks. Optionally, the second dose is administered for a second treatment period after completion of the first treatment period. During the second treatment period the second dose is administered twice a week. The second treatment period lasts about 1-20 weeks, 2-20 weeks, 3- 20 weeks, 4-20 weeks, 5-20 weeks, 6- 20 weeks, 1-10 weeks, 2-10 weeks, 3- 10 weeks, 4-10 weeks, 5-10 weeks, 6- 10 weeks. The second treatment period is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,11, 12 or more weeks.

**[0136]** In some embodiments, the second dose is increased or decreased during the course of the first or second treatment period. The increased or decreased dose is referred to as a third dose. The third dose may 1.0, 3.0 or 6.0 mg/kg body weight. The third dose is administered for the remainder of the first and second treatment period. Alternatively, the third

dose is administered for a third treatment period. In some embodiments the third dose is referred to as a maintenance dose and the third treatment period is referred to as a maintenance period.

[0137] A multiple variable dose regimen for the treatment of secondary hemophagocytic lymphohistiocytosis (HLH) in a human pediatric includes an induction or a first dose and a treatment or second dose of an anti- IFN-g monoclonal antibody .

[0138] The first dose is 6.0, mg/kg body weight. The second dose is 3.0, mg/kg body weight. 10.0 mg/kg body weight.

[0139] The first dose is administered once as a single dose. Alternatively, the first dose is administered more than once for a induction treatment period.

[0140] The second dose is administered for one or more treatment periods. The second dose is administered for a first treatment period. During the first treatment period the second dose is administered every three days after the first dose. The first treatment period last about 1, 2, 3, 4, 5, 6 or more weeks. Preferably the first treatment period is 2 weeks. Optionally, the second dose is administered for a second treatment period after completion of the first treatment period. During the second treatment period the second dose is administered twice a week. The second treatment period lasts about 1-20 weeks, 2-20 weeks, 3- 20 weeks, 4-20 weeks, 5-20 weeks, 6- 20 weeks, 1-10 weeks, 2-10 weeks, 3- 10 weeks, 4-10 weeks, 5-10 weeks, 6- 10 weeks. The second treatment period is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,11, 12 or more weeks.

[0141] In some embodiments, the second dose is increased or decreased during the course of the first or second treatment period. The increased or decreased dose is referred to as a third dose. Preferably, the third dose is 6.0 mg/kg body weight. The third dose is administered for the remainder of the first and second treatment period. Alternatively, the third dose is administered for a third treatment period. In some embodiments the third dose is referred to as a maintenance dose and the third treatment period is referred to as a maintenance period.

[0142] A multiple variable dose regimen for the treatment of secondary hemophagocytic lymphohistiocytosis (HLH) in a human adult includes an induction or a first dose and a treatment or second dose of an anti- IFN-g monoclonal antibody .

[0143] The first dose is 3.0 or 6.0 mg/kg body weight. Preferably, the first dose is 6.0

mg/kg body weight. The second dose is 6.0 or 10.0 mg/kg body weight. Preferably, the second dose is 10.0 mg/kg body weight. The first dose is 6.0 mg/kg body weight an antibody and the second does is 10.0 mg/kg body weight. The first dose is administered once as a single dose. Alternatively the first dose is administer more than once for an induction treatment period.

**[0144]** The second dose is administered for one or more treatment periods. The second dose is administered for a first treatment period. During the first treatment period the second dose is administered every three days after the first dose. The first treatment period last about 1, 2, 3, 4, 5, 6 or more weeks. Preferably the first treatment period is 2 weeks. Optionally, the second dose is administered for a second treatment period after completion of the first treatment period. During the second treatment period the second dose is administered twice a week. The second treatment period lasts about 1-20 weeks, 2-20 weeks, 3- 20 weeks, 4-20 weeks, 5-20 weeks, 6- 20 weeks, 1-10 weeks, 2-10 weeks, 3- 10 weeks, 4-10 weeks, 5-10 weeks, 6- 10 weeks. The second treatment period is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,11, 12 or more weeks.

**[0145]** In some embodiments, the second dose is increased or decreased during the course of the first or second treatment period. The increased or decreased dose is referred to as a third dose. The third dose may 1.0, 3.0 or 6.0 mg/kg body weight. The third dose is administered for the remainder of the first and second treatment period. Alternatively, the third dose is administered for a third treatment period. In some embodiments the third dose is referred to as a maintenance dose and the third treatment period is referred to as a maintenance period.

**[0146]** A multiple variable dose regimen for the treatment of a condition in a human subject includes an induction or a first dose and a treatment or second dose of an anti- IFN-g monoclonal antibody. The subject is an adult subject or a pediatric subject. The condition is associated with elevated IFNg levels. The condition is transplant rejection such as solid organ transplant disorder or bone marrow acute graft rejection graft vs. host disease, paraneoplastic cerebellar degeneration, hemorrhagic fever, sarcoidosis, adult onset Still's disease. In other embodiments dosing regimen is administered to a subject after receiving CAR-T cell therapy.

**[0147]** The first dose is between 1.0- 10 mg/kg body weight. For example, first dose is 1.0, 3.0, 6.0, or 10 mg/kg body weight an antibody. The second dose is higher or lower that the first doses. The second dose is between 1.0- 10 mg/kg body weight. For example, second

dose is 1.0, 3.0, 6.0 or 10, mg/kg body weight an antibody. The first dose is 1.0 mg/kg body weight and the second does is 3.0, 6.0 or 10.0 mg/kg body weight. Alternatively, the first dose is 3.0 mg/kg body weight an antibody and the second does is 6.0 or 10.0 mg/kg body weight. The first dose is 6.0 mg/kg body weight an antibody and the second does is 10.0 mg/kg body weight.

**[0148]** The first dose is administered once as a single dose. Alternatively the first dose is administer more than once for an induction treatment period.

**[0149]** The second dose is administered for one or more treatment periods. The second dose is administered for a first treatment period. During the first treatment period the second dose is administered every three days after the first dose. The first treatment period last about 1, 2, 3, 4, 5, 6 or more weeks. Preferably the first treatment period is 2 weeks. Optionally, the second dose is administered for a second treatment period after completion of the first treatment period. During the second treatment period the second dose is administered twice a week. The second treatment period lasts about 1-20 weeks, 2-20 weeks, 3- 20 weeks, 4-20 weeks, 5-20 weeks, 6- 20 weeks, 1-10 weeks, 2-10 weeks, 3- 10 weeks, 4-10 weeks, 5-10 weeks, 6- 10 weeks. The second treatment period is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10,11, 12 or more weeks.

**[0150]** In some embodiments, the second dose is increased or decreased during the course of the first or second treatment period. The increased or decreased dose is referred to as a third dose. The third dose may 1.0, 3.0, 6.0 or 10.0 mg/kg body weight. The third dose is administered for the remainder of the first and second treatment period. Alternatively, the third dose is administered for a third treatment period. In some embodiments the third dose is referred to as a maintenance dose and the third treatment period is referred to as a maintenance period

**[0151]** In the multiple variable dosing regimen's according to the invention , the first and second time period for the second dose may be adjusted from time to time based upon for example, the subject's heath status. For example, adjustments to the time between doses may be adjusted  $\pm 1, 2, 3,$  or 4 days. Preferably  $\pm 2$  days.

**[0152]** In the multiple variable dosing regimen's according to the invention the dose is administered with 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 hrs. Preferably, the doses is administered with an hour. The time period for infusion of the dose is based upon a number of factors know in

the art, for example, age, weight, physical condition or adverse reaction to the therapeutic. The time period for the infusion of the dose is typically the fasted infusion rate that is tolerated (i.e., without causing an adverse reaction) by the subject.

[0153] The dose is administered as a single injection. The antibody is administered as a monotherapy or a co-therapy for the disease or condition being treated. For example the subject is administered a second agent. The second agent is for example an anti-inflammatory agent, and/or an immunosuppressive agent

[0154] Optionally, the subject has been administered dexamethasone immediately prior to the dosing of the antibody. The dexamethasone is administered at a dose of at least 10 mg/m<sup>2</sup>. Alternatively, the dexamethasone is administered at a dose of at least 5 mg/m<sup>2</sup>.

[0155] In some embodiments, subject has not previously been treated for HLH.

#### Pharmaceutical compositions

[0156] The antibodies or soluble chimeric polypeptides of the invention (also referred to herein as “active compounds”), and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the antibody or soluble chimeric polypeptide and a pharmaceutically acceptable carrier. As used herein, the term “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington’s Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, ringer’s solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0157] A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal,

or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

**[0158]** The preferred route of administration is by injection such as intravenous injection. The intravenous injection can be rapid or slow infusion. For example, the infusions is over about a 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 hour period. The time period for infusion is based upon a number of factors know in the art, for example, age, weight, physical condition or adverse reaction to the therapeutic. The time period for the infusion is typically the fasted infusion rate that is tolerated (i.e., without causing an adverse reaction) by the subject.

**[0159]** Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[0160] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0161] Preferred injectable pharmaceutical compositions include excipients such as L-histidine, 3 L-histidine monohydrochloride, monohydrate, sodium chloride (NaCl), Polysorbate 80.

[0162] The pH of the injectable pharmaceutical composition is between about 5.8 and 6.2. Preferably the pH of the injectable pharmaceutical composition is pH 6.0.

[0163] In some embodiments, the anti-IFN- $\gamma$  antibody, such as NI-0501 is formulated (per ml) as follows: 5 mg of NI-051, 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is  $6.0 \pm 0.2$ .

[0164] In some aspects the pharmaceutical composition is packaged in a unit dose. The unit dose is packaged in a container. The container is a glass or plastic container. The container is a syringe, vial, infusion bottle, ampoule or carpoule. The container can hold a volume of 1-25 ml. For example, the container can hold a volume of 2, 5, 10 or 20 ml.

[0165] The unit dose of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody (e.g., NI-501) is between 5- 25 mg/ ml. Preferably the unit dose is 5 mg/ml or 25 mg/ml. The antibody is in a solution (e.g., water for injection) having per ml: 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80.

[0166] In some embodiment the invention provides a unit dose container having 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution at a concentration of 5 mg/ml or 25 mg/ml antibody, wherein the pH of the solution is between 5.8 and 6.2. The antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate. Preferably, the antibody is in a solution (e.g., water for injection) having per ml: 1.55 mg L-

histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80.

**[0167]** In other embodiments the invention provides a unit dose container having 10 ml or 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution at a concentration of 25 mg/ml antibody, wherein the pH of the solution is between 5.8 and 6.2. The antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate. Preferably, the antibody is in a solution (e.g., water for injection) having per ml: 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80.

**[0168]** In other embodiments the invention provides a unit dose container having 2 ml or 10 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution at a concentration of 5 mg/ml antibody, wherein the pH of the solution is between 5.8 and 6.2. The antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate. Preferably, the antibody is in a solution (e.g., water for injection) having per ml: 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80.

**[0169]** Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

**[0170]** For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

[0171] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[0172] The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[0173] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[0174] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

[0175] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

### Detection of CXCL9 and other Biomarkers

[0176] Levels of CXCL9 and other biomarkers are detecting using any of a variety of standard detection techniques. Detection agents can be used for detecting the presence of a given target (or a protein fragment thereof) in a sample. In some embodiments, the detection agent contains a detectable label. In some embodiments, the detection agent is an antibody (or fragment thereof) or a probe. In some embodiments, the agent or probe is labeled. The term “labeled”, with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin.

[0177] The term “biological sample” is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. Included within the usage of the term “biological sample”, therefore, is blood and a fraction or component of blood including blood serum, blood plasma, or lymph. The bodily fluids can be fluids isolated from anywhere in the body of the subject, preferably a peripheral location, including but not limited to, for example, blood, plasma, serum, synovial fluid, urine, sputum, spinal fluid, cerebrospinal fluid, pleural fluid, fluid of the respiratory, intestinal, and genitourinary tracts, saliva, intra-organ system fluid, ascitic fluid, tumor cyst fluid, amniotic fluid and combinations thereof. The biological sample also includes experimentally separated fractions of all of the preceding fluids. Biological samples also include solutions or mixtures containing homogenized solid material, such as feces, tissues, and biopsy samples. The detection method of the invention can be used to detect an analyte mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of an analyte mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of an analyte protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of an analyte genomic DNA include Southern hybridizations. Procedures for conducting immunoassays are described, for example in “ELISA: Theory and Practice: Methods in Molecular Biology”, Vol. 42, J. R. Crowther (Ed.) Human Press, Totowa, NJ, 1995; “Immunoassay”, E. Diamandis and T. Christopoulos, Academic Press, Inc., San

Diego, CA, 1996; and “Practice and Theory of Enzyme Immunoassays”, P. Tijssen, Elsevier Science Publishers, Amsterdam, 1985. Furthermore, *in vivo* techniques for detection of an analyte protein include introducing into a subject a labeled anti-analyte protein antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

Definitions:

[0178] Unless otherwise defined, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures utilized in connection with, and techniques of; cell and tissue culture, molecular biology, and protein and oligo-or polynucleotide chemistry and hybridization described herein are those well-known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (*e.g.*, electroporation, lipofection). Enzymatic reactions and purification techniques are performed according to manufacturer’s specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. *See e.g.*, Sambrook *et al.* Molecular Cloning: A Laboratory Manual (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). The nomenclatures utilized in connection with, and the laboratory procedures and techniques of; analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0179] The term “dose,” as used herein, refers to an amount of an anti- IFN-gamma antibody which is administered to a subject.

[0180] The term “multiple-variable dose” includes different doses of an anti- IFN-gamma antibody which are administered to a subject for therapeutic treatment. “Multiple-variable dose regimen” or “multiple-variable dose therapy” describe a treatment schedule which is based on

administering different amounts of an anti- IFN-gamma antibody at various time points throughout the course of treatment. In one embodiment, the invention describes a multiple-variable dose method of treatment comprising an induction phase and a treatment phase, wherein an anti- FN-gamma antibody is administered at a higher or lower dose during the induction phase than the treatment phase. In another embodiment, the invention describes a multiple-variable dose method of treatment comprising an induction phase, a first a treatment phase, and a second treatment phase wherein an anti- IFN-gamma antibody is administered at a higher or lower dose during the induction phase than the treatment phase. The anti- IFN-gamma antibody may be administered at a higher or lower dose during the first treatment phase than the second treatment phase. Preferably, the dose is the same in the first treatment phases and the second treatment phase. .

**[0181]** The term “induction phase” or “loading phase”, as used herein, refers to a period of treatment comprising administration of an IFN-gamma antibody to a subject in order to attain a threshold level. During the induction phase, at least one induction dose of IFN-gamma is administered to a subject suffering from a disorder in which IFN-gamma is detrimental.

**[0182]** The term “threshold level”, as used herein, refers to a therapeutically effective level of an IFN-gamma antibody in a subject. A threshold level is achieved by administering at least one induction dose during the induction phase of treatment. Any number of induction doses may be administered to achieve a threshold level of IFN-gamma. Once a threshold level is achieved, the treatment phase is initiated.

**[0183]** The term “induction dose” or “loading dose,” used interchangeably herein, refers to the first dose of an anti- IFN-gamma antibody, which is either larger or smaller in comparison to the maintenance or treatment dose. The induction dose can be a single dose or, alternatively, a set of doses. The induction dose is often used to bring the drug in the body to a steady state amount, and may be used to which to achieve maintenance drug levels quickly. An induction dose is subsequently followed by administration of smaller or larger doses of an anti- IFN-gamma antibody, i.e., the treatment dose. The induction dose is administered during the induction phase of therapy.

**[0184]** The term “treatment phase” or “maintenance phase”, as used herein, refers to a period of treatment comprising administration of an anti- IFN-gamma antibody to a subject in order to maintain a desired therapeutic effect. The treatment phase follows the induction phase, and, therefore, is initiated once a threshold level is achieved. There may be more than one

treatments phases, that is a treatment phases shortened or extended to maintain a certain threshold or achieve a desired clinical response. Treatment phase may be Preferably, the substance is administered every 3, days following the induction dose. (e.g. initial dose) days. Alternatively, the substance is administered once or twice a week.

[0185] The term “treatment dose” or “maintenance dose” is the amount of an anti- IFN-gamma antibody taken by a subject to maintain or continue a desired therapeutic effect. A treatment dose is administered subsequent to the induction dose. A treatment dose can be a single dose or, alternatively, a set of doses. A treatment dose is administered during the treatment phase of therapy. Treatment doses are smaller or larger than the induction dose and can be equal to each other when administered in succession. When there is more than one treatment phases there may also be more than one treatment doses.

[0186] A “dosage regimen” or “dosing regimen” includes a treatment regimen based on a determined set of doses.

[0187] The term “dosing”, as used herein, refers to the administration of a substance (e.g., an an anti- IFN-gamma antibody) to achieve a therapeutic objective (e.g., the treatment of a IFN-gamma-associated disorder).

[0188] The terms “biweekly dosing regimen”, “biweekly dosing”, and “biweekly administration”, as used herein, refer to the time course of administering a substance (e.g., an anti- IFN-gamma antibody) to a subject to achieve a therapeutic objective (e.g., the treatment of a IFN-gamma-associated disorder). The biweekly dosing regimen is not intended to include a weekly dosing regimen.

[0189] The term “combination” as in the phrase “a first agent in combination with a second agent” includes co administration of a first agent and a second agent, which for example may be dissolved or intermixed in the same pharmaceutically acceptable carrier, or administration of a first agent, followed by the second agent, or administration of the second agent, followed by the first agent. The present invention, therefore, includes methods of combination therapeutic treatment and combination pharmaceutical compositions.

[0190] The term “concomitant” as in the phrase “concomitant therapeutic treatment” includes administering an agent in the presence of a second agent. A concomitant therapeutic treatment method includes methods in which the first, second, third, or additional agents are co-administered. A concomitant therapeutic treatment method also includes methods in which

the first or additional agents are administered in the presence of a second or additional agents, wherein the second or additional agents, for example, may have been previously administered. A concomitant therapeutic treatment method may be executed step-wise by different actors. For example, one actor may administer to a subject a first agent and a second actor may to administer to the subject a second agent, and the administering steps may be executed at the same time, or nearly the same time, or at distant times, so long as the first agent (and additional agents) are after administration in the presence of the second agent (and additional agents). The actor and the subject may be the same entity (e.g., human).

[0191] The term “combination therapy”, as used herein, refers to the administration of two or more therapeutic substances, e.g., an anti- IFN-gamma antibody and another drug, such as a DMARD or NSAID. The other drug(s) may be administered concomitant with, prior to, or following the administration of an anti- IFN-gamma antibody.

#### Embodiments

[0192] The compositions and methods provided herein, NI-0501 is formulated as a sterile concentrate for infusion (per mL). In some embodiments, NI-0501 is formulated as follows: 5 mg of NI-051, 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is between 5.8 and 6.2. In some embodiments, NI-0501 is formulated as follows: 5 mg of NI-051, 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is 6.0.

[0193] In the compositions and methods provided herein, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with HLH. In some embodiments, NI-0501 is administered to a subject in need thereof by IV infusion over a period of one hour at an initial dose of 1 mg/kg. In certain patient populations, e.g., those with low body weight and/or the very young, the IV infusion may last more than one hour, for example, at least 90 minutes, at least 2 hours, or at least 3 hours or greater.

[0194] In some embodiments, NI-0501 is administered to a subject in need thereof by at least one additional IV infusion after the initial IV infusion over a period of one hour at an initial dose of 1 mg/kg. In some embodiments, the at least one additional IV infusion is at a dose that is higher than the initial dose of 1 mg/kg. In some embodiments, the at least one additional IV infusion dosage is 3 mg/kg. In some embodiments, the at least one additional

IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the at least one additional IV infusion is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the at least one additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion.

**[0195]** In some embodiments, NI-0501 is administered to a subject in need thereof by at least one series additional IV infusions after the initial IV infusion over a period of one hour at an initial dose of 1 mg/kg, where the series of additional IV infusions includes at least one series of twice weekly IV infusions. In some embodiments, the at least one series of twice weekly IV infusions is administered at a dose higher than the initial dose of 1 mg/kg. In some embodiments, the at least one series of twice weekly IV infusions is administered at a dose of 3 mg/kg. In some embodiments, the at least one additional IV infusion is administered at least three weeks after the initial IV infusion. In some embodiments, the at least one additional IV infusion is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the at least one additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

**[0196]** In some embodiments, NI-0501 is administered to a subject in need thereof by at least two additional IV infusions after the initial IV infusion. In some embodiments, the at least two additional IV infusions are at a dose that is higher than the initial dose of 1 mg/kg. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at the same dosage. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at the same dosage that is higher than the initial dose. In some embodiments, at least one of the first and second additional IV infusions is administered at a dosage of 3mg/kg. In some embodiments, the first additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the first

additional IV infusion is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the second additional IV infusion is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion, and the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

[0197] In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at different dosages. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at different dosages, where the second additional IV infusion dosage is higher than the first additional IV infusion. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at different dosages, where the second additional IV infusion dosage is higher than the first additional IV infusion, and where both the first and the second additional IV infusion dosages are higher than the initial dosage. In some embodiments, at least one of the first and second additional IV infusions is administered at a dosage of 3mg/kg. In some embodiments, the first additional IV infusion is administered at a dosage of 3 mg/kg, and the second additional IV infusion is administered at a dosage of 6 mg/kg. In some embodiments, the first additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the first additional IV infusion is administered at a time selected from the group consisting of at 3

days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the second additional IV infusion is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion, and the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

[0198] In some embodiments, the first additional IV infusion includes at least a first series of twice weekly IV infusions and the second additional IV infusion includes at least a second series of twice weekly IV infusions. In some embodiments, the first series of twice weekly IV infusions and the second series of twice weekly IV infusions are administered at a dose higher than the initial dose of 1 mg/kg. In some embodiments, the first series of twice weekly IV infusions is administered at a dose of 3 mg/kg, and the second series of twice weekly IV infusions is administered at a dose of 6 mg/kg. In some embodiments, the first series of additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the first series of additional IV infusions is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the first series of additional IV infusions is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the

second series of additional IV infusions is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the second series of additional IV infusions is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the first series of additional IV infusions is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion, and the second series of additional IV infusions is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

[0199] In some embodiments, infusions are performed every 3 days after the initial dose for up to 15 days after the initial dose. In some embodiments, infusions are performed every 3 days after the initial dose for up to 15 days after the initial dose, followed by infusions twice per week starting at least 15 days after the initial dose. In some embodiments, the infusion dosage is increased to 3 mg/kg at any point after the initial dose. In some embodiments, after a minimum of two infusions at 3 mg/kg, the dose of NI-0501 is increased to 6 mg/kg for up to four infusions.

[0200] In the compositions and methods provided herein, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with HLH. In some embodiments, NI-0501 is administered to a subject in need thereof by IV infusion over a period of one hour at an initial dose of 1 mg/kg. In some embodiments, infusions are performed every 3 days after the initial dose for up to 15 days after the initial dose. In some embodiments, infusions are performed every 3 days after the initial dose for up to 15 days after the initial dose, followed by infusions twice per week starting at least 15 days after the initial dose. In some embodiments, the infusion dosage is increased to 3 mg/kg at any point after the initial dose. In some embodiments, after a minimum of two infusions at 3 mg/kg, the dose of NI-0501 is increased to 6 mg/kg for up to four infusions.

[0201] In the compositions and methods provided herein, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with HLH. In some embodiments, NI-0501 is administered to a subject in

need thereof by IV infusion for a dosage of greater than 6 mg/kg. In some embodiments, NI-0501 is administered to a subject in need thereof by IV infusion, after an initial dosage, for a second dosage of greater than 6 mg/kg. In some embodiments, the second dosage is at least 10 mg/kg. In some embodiments, the second dosage is 10 mg/kg. In some embodiments, the second dosage is 10 mg/kg, repeated daily. In some embodiments, the second dosage is 10 mg/kg, repeated daily for 1 week. In some embodiments, the second dosage is 10 mg/kg, repeated daily for 2 weeks. In some embodiments, the second dosage is 10 mg/kg, repeated daily for more than 2 weeks.

[0202] In some embodiments, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with secondary HLH. In some embodiments, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with secondary HLH on a background of sJIA. In some embodiments, NI-0501 is administered to a subject in need thereof as an initial dose of 6 mg/kg. In some embodiments, the NI-0501 treatment is continued with a subsequent NI-0501 dose. In some embodiments, the NI-0501 treatment is continued with a subsequent NI-0501 dose of 3 mg/kg every 3 days for at least 4 weeks (i.e., up to SD27).

[0203] In some embodiments, NI-0501 treatment is reduced, stopped, or otherwise shortened upon achievement of a desired clinical outcome. In some embodiments, NI-0501 treatment is shortened upon evidence of complete clinical response, i.e., MAS remission.

[0204] In some embodiments, after 4 weeks, NI-0501 treatment is continued for up to an additional 4 weeks (i.e., up to SD56) as maintenance as needed until MAS remission is achieved. In some embodiments, after 4 weeks, NI-0501 treatment is continued for up to an additional 4 weeks (i.e., up to SD56) as maintenance as needed until MAS remission is achieved, with the possibility of decreasing the dose to 1 mg/kg and elongating the interval between infusion to weekly administration.

[0205] In the compositions and methods provided herein, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with HLH, where the subject has been administered a background of dexamethasone. In some embodiments, the subject is a treatment-naïve patient (i.e., has not previously been treated for HLH), and the dexamethasone is administered at a dose of at least 10 mg/m<sup>2</sup>. In some embodiments, the subject is receiving NI-0501 as a second line HLH treatment, and the

dexamethasone is administered at a dose of in the range of 10 mg/m<sup>2</sup> to 5 mg/m<sup>2</sup>. In some embodiments, the subject is receiving NI-0501 as a second line HLH treatment, and the dexamethasone is administered at a dose of at least 5 mg/m<sup>2</sup>. In some embodiments, the subject is receiving NI-0501 as a second line HLH treatment, and the dexamethasone is administered at a dose of less than 5 mg/m<sup>2</sup>.

[0206] In some embodiments, NI-0501 is administered before and/or during and/or after treatment in combination with one or more additional agents such as, by way of non-limiting example, a therapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent. In some embodiments, the second agent is an agent known to be used in the treatment of HLH. In some embodiments, the additional agent includes at least etoposide. In some embodiments, NI-0501 and the additional agent are formulated into a single therapeutic composition, and NI-0501 and additional agent are administered simultaneously. Alternatively, NI-0501 and additional agent are separate from each other, *e.g.*, each is formulated into a separate therapeutic composition, and NI-0501 and the additional agent are administered simultaneously, or NI-0501 and the additional agent are administered at different times during a treatment regimen. For example, NI-0501 is administered prior to the administration of the additional agent, NI-0501 is administered subsequent to the administration of the additional agent, or NI-0501 and the additional agent are administered in an alternating fashion. As described herein, NI-0501 and additional agent are administered in single doses or in multiple doses.

[0207] In some embodiments, NI-0501 and the additional agent(s) are administered simultaneously. For example, NI-0501 and the additional agent(s) can be formulated in a single composition or administered as two or more separate compositions. In some embodiments, NI-0501 and the additional agent(s) are administered sequentially, or NI-0501 and the additional agent are administered at different times during a treatment regimen.

[0208] In some embodiments, the additional agent is an immunosuppressive agent. In some embodiments, the immunosuppressive agent is Cyclosporin A (CsA). In some embodiments, the subject has been receiving CsA prior to administration of NI-0501. In some embodiments, the additional agent includes at least etoposide. In some embodiments, the subject has been receiving etoposide prior to administration of NI-0501.

[0209] In some embodiments, the additional agent is intrathecal methotrexate and/or glucocorticoids. In some embodiments, the subject has been receiving intrathecal

methotrexate and/or glucocorticoids prior to administration of NI-0501.

[0210] In some embodiments, the additional agent is IV immunoglobulins (IVIG). In some embodiments, the IVIG is administered as replacement treatment in a subject with a documented immunoglobulin deficiency. In some embodiments where the subject has a documented immunoglobulin deficiency, IVIG is administered given at a dose of 0.5 g/kg, every 4 weeks or more frequently in order to maintain adequate IgG levels.

[0211] In some embodiments, the one or more additional agents is analgesic treatment, transfusion of blood products, electrolyte and glucose infusions, antibiotics, anti-fungal and anti-viral treatment and/or general supportive care.

[0212] The compositions and methods provided herein are useful in treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with transplant failure, transplant rejection, and/or an inflammatory disorder associated with transplant rejection. The compositions and methods provided herein are useful in treating, inhibiting, delaying the progression of, or otherwise ameliorating a symptom of Graft-versus-host disease (GvHD) in a subject who has received or is receiving a transplant comprising biological material or a series of transplants comprising biological material. The compositions and methods provided herein are useful in prolonging survival of transplanted biological material.

[0213] The compositions and methods provided herein are useful in transplanting any biological material, including, for example, cells, tissue(s), bone marrow, and/or organ(s) or at least a portion of an organ, including, by way of non-limiting example, heart, kidney, pancreas, liver, and/or intestine. In some embodiments, the biological material to be transplanted is allogeneic biological material. In some embodiments, the transplanted biological material is bone marrow. In some embodiments, the transplanted biological material is a population of hematopoietic stem cells. In some embodiments, the biological material to be transplanted is or is derived from one or more hepatocytes.

[0214] In the compositions and methods provided herein, NI-0501 is administered to a subject in need thereof for treating, preventing and/or delaying the onset or progression of, or alleviating a symptom associated with transplant failure, transplant rejection, and/or an inflammatory disorder associated with transplant rejection. In some embodiments, the transplant rejection, also referred to herein as transplant failure, is acute. In some embodiments, the transplant rejection is hyperacute.

[0215] In the compositions and methods provided herein, NI-0501 is formulated as a sterile

concentrate for infusion (per mL). In some embodiments, NI-0501 is formulated as follows: 5 mg of NI-051, 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is between 5.8 and 6.2. In some embodiments, NI-0501 is formulated as follows: 5 mg of NI-051, 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is 6.0.

[0216] In some embodiments, the transplant rejection is chronic. In some embodiments, NI-0501 is administered to a subject in need thereof by IV infusion over a period of one hour at an initial dose of 1 mg/kg. In certain patient populations, e.g., those with low body weight and/or the very young, the IV infusion may last more than one hour, for example, at least 90 minutes, at least 2 hours, or at least 3 hours or greater.

[0217] In some embodiments, NI-0501 is administered to a subject in need thereof by at least one additional IV infusion after the initial IV infusion over a period of one hour at an initial dose of 1 mg/kg. In some embodiments, the at least one additional IV infusion is at a dose that is higher than the initial dose of 1 mg/kg. In some embodiments, the at least one additional IV infusion dosage is 3 mg/kg. In some embodiments, the at least one additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the at least one additional IV infusion is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the at least one additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion.

[0218] In some embodiments, NI-0501 is administered to a subject in need thereof by at least one series additional IV infusions after the initial IV infusion over a period of one hour at an initial dose of 1 mg/kg, where the series of additional IV infusions includes at least one series of twice weekly IV infusions. In some embodiments, the at least one series of twice weekly IV infusions is administered at a dose higher than the initial dose of 1 mg/kg. In some embodiments, the at least one series of twice weekly IV infusions is administered at a dose of 3 mg/kg. In some embodiments, the at least one additional IV infusion is administered at least three weeks after the initial IV infusion. In some embodiments, the at least one additional IV infusion is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks

after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the at least one additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

[0219] In some embodiments, NI-0501 is administered to a subject in need thereof by at least two additional IV infusions after the initial IV infusion. In some embodiments, the at least two additional IV infusions are at a dose that is higher than the initial dose of 1 mg/kg. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at the same dosage. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at the same dosage that is higher than the initial dose. In some embodiments, at least one of the first and second additional IV infusions is administered at a dosage of 3mg/kg. In some embodiments, the first additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the first additional IV infusion is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the second additional IV infusion is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion, and the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion,

at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

[0220] In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at different dosages. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at different dosages, where the second additional IV infusion dosage is higher than the first additional IV infusion. In some embodiments, the first additional IV infusion and the second additional IV infusion are administered at different dosages, where the second additional IV infusion dosage is higher than the first additional IV infusion, and where both the first and the second additional IV infusion dosages are higher than the initial dosage. In some embodiments, at least one of the first and second additional IV infusions is administered at a dosage of 3mg/kg. In some embodiments, the first additional IV infusion is administered at a dosage of 3 mg/kg, and the second additional IV infusion is administered at a dosage of 6 mg/kg. In some embodiments, the first additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the first additional IV infusion is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the second additional IV infusion is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the first additional IV infusion is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion, and the second additional IV infusion is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the

initial infusion.

[0221] In some embodiments, the first additional IV infusion includes at least a first series of twice weekly IV infusions and the second additional IV infusion includes at least a second series of twice weekly IV infusions. In some embodiments, the first series of twice weekly IV infusions and the second series of twice weekly IV infusions are administered at a dose higher than the initial dose of 1 mg/kg. In some embodiments, the first series of twice weekly IV infusions is administered at a dose of 3 mg/kg, and the second series of twice weekly IV infusions is administered at a dose of 6 mg/kg. In some embodiments, the first series of additional IV infusion is administered at least three days after the initial IV infusion. In some embodiments, the first series of additional IV infusions is administered at a time selected from the group consisting of at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the first series of additional IV infusions is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion. In some embodiments, the second series of additional IV infusions is administered at a time selected from the group consisting of at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the second series of additional IV infusions is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion. In some embodiments, the first series of additional IV infusions is administered at 3 days after the initial IV infusion, at 6 days after the initial IV infusion, at 9 days after the initial IV infusion, at 12 days after the initial infusion, and at 15 days after the initial infusion, and the second series of additional IV infusions is administered at 3 weeks after the initial IV infusion, at 4 weeks after the initial IV infusion, at 5 weeks after the initial IV infusion, at 6 weeks after the initial infusion, at 7 weeks after the initial infusion, and at 8 weeks after the initial infusion.

[0222] In some embodiments, infusions are performed every 3 days after the initial dose for up to 15 days after the initial dose. In some embodiments, infusions are performed every 3 days after the initial dose for up to 15 days after the initial dose, followed by infusions twice per week starting at least 15 days after the initial dose. In some embodiments, the infusion dosage is

increased to 3 mg/kg at any point after the initial dose. In some embodiments, after a minimum of two infusions at 3 mg/kg, the dose of NI-0501 is increased to 6 mg/kg for up to four infusions.

[0223] The disclosure also provides compositions and methods that are useful in identifying or otherwise refining a patient population suffering from a disorder, where the patient has an elevated level of CXCL9, alone or in combination with one or more additional interferon  $\gamma$  (IFN $\gamma$ ) related biomarkers. In particular, the disclosure provides compositions and methods for detecting CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from hemophagocytic lymphohistiocytosis (HLH). In particular, the disclosure provides compositions and methods for detecting CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from secondary hemophagocytic lymphohistiocytosis (HLH). In some embodiments, the compositions and methods are used to detect CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from macrophage activation syndrome (MAS). In some embodiments, the compositions and methods are used to detect CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from MAS in the context of an autoimmune disease or inflammatory disorder. In some embodiments, the compositions and methods are used to detect CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from MAS in the context of a systemic autoimmune disease or inflammatory disorder. In some embodiments, the compositions and methods are used to detect CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from MAS in the context of systemic Juvenile Idiopathic Arthritis (sJIA). In some embodiments, the compositions and methods are used to detect CXCL9 levels as a biomarker for IFN $\gamma$  production in patients suffering from or suspected of suffering from MAS in the context of Systemic Lupus Erythematosus (SLE).

[0224] Patients identified as having elevated levels of CXCL9 are identified as suitable candidates for treatment with an agent (*e.g.*, antibodies or other polypeptide-based therapeutics, peptide-based therapeutics, small molecule inhibitors, nucleic acid-based therapeutics and derivatives thereof) that interferes with or otherwise antagonizes one or more biological activities of IFN $\gamma$  such as, for example, IFN $\gamma$  signaling, and neutralizes at least one biological activity of IFN $\gamma$ .

[0225] In some patients suffering from or suspected of suffering from a disorder, fluids and other

biological samples contain elevated levels of CXCL9, alone or in combination with other IFN $\gamma$ -related biomarkers such as, for example, CXCL10 and/or CXCL11.

[0226] CXCL9 and these other biomarkers are indicators of *in vivo* IFN $\gamma$  production. Thus, use of an anti-IFN $\gamma$  antagonist that interferes with, inhibits, reduces or otherwise antagonizes IFN $\gamma$  signaling, *e.g.*, a neutralizing anti-IFN $\gamma$  antibody or other polypeptide-based therapeutic, a peptide-based therapeutic, a small molecule inhibitor, a nucleic acid-based therapeutic and derivatives thereof, blocks or otherwise inhibits IFN $\gamma$  activity. Thus, the compositions and methods are useful in treating, delaying the progression of or otherwise ameliorating a symptom of a disorder that is dependent on, driven by, associated with, or otherwise impacted by aberrant, *e.g.*, elevated, IFN $\gamma$  expression and/or activity, aberrant pro- inflammatory cytokine production and/or combinations thereof, by administering an anti-IFN $\gamma$  antagonist, *e.g.*, a neutralizing anti-IFN $\gamma$  antibody or other polypeptide-based therapeutic, a peptide-based therapeutic, a small molecule inhibitor, a nucleic acid-based therapeutic and derivatives thereof, to patients exhibiting an elevated level of expression CXCL9 and/or other biomarkers. Patients that are likely suitable candidates for treatment with the anti-IFN $\gamma$  antagonist, *e.g.*, neutralizing anti-IFN $\gamma$  antibody such as those described herein, are identified by detecting the level of CXCL9, alone or in combination with one or more IFN $\gamma$ -related ligands or other biomarkers. In some embodiments, patients that do not have elevated levels of CXCL9, alone or in combination with other IFN $\gamma$ -related biomarkers may still be treated with an anti- IFN $\gamma$  antagonist, including any of the neutralizing anti-IFN $\gamma$  antibodies described herein or other polypeptide-based therapeutic, a peptide-based therapeutic, a small molecule inhibitor, a nucleic acid-based therapeutic and derivatives thereof.

[0227] Patients with elevated levels of CXCL9, alone or in combination with one or more additional IFN $\gamma$ -related biomarkers are identified as suitable candidates for therapy with one or more anti-IFN $\gamma$  antagonists, *e.g.*, a neutralizing anti-IFN $\gamma$  antibody described herein. As used herein, the phrase “elevated level of expression” refers to a level of expression that is greater than a baseline level of expression of CXCL9, alone or in combination with one or more additional biomarkers, in a sample from a patient that is not suffering from or suspected of suffering from primary or secondary HLH or HLH-related disorder, or from another control sample. In some embodiments, the elevated level of expression of CXCL9 and/or other biomarker is a significant level of elevation.

[0228] The detected level of CXCL9, alone or in combination with one or more other IFN $\gamma$ -related biomarkers, is useful to refine or otherwise stratify a patient population. In some embodiments, the detected level is used to determine the dosage of anti-IFN $\gamma$  antagonist that should be administered to a given patient. In some embodiments, the detected level is used to categorize or otherwise stratify a patient population. For example, patients can be classified as having “severe” or high grade MAS, or conversely, not severe or low grade MAS, based on the detected level of CXCL9.

[0229] The sample is, for example, blood or a blood component, *e.g.*, serum, plasma. In some embodiments, the sample is another bodily fluid such as, by way of non-limiting example, urine, synovial fluid, bronchial alveolar fluid, cerebrospinal fluid, broncho-alveolar lavage (BAL), and/or saliva. In some embodiments, the biological sample is CSF. In some embodiments, the biological sample is CSF from an HLH patient.

[0230] In addition to detecting the level of IFN $\gamma$  and/or other IFN $\gamma$ -related biomarkers, suitable patients for treatment with an anti-IFN $\gamma$  antagonist can also be identified by evaluating any of a number of additional biological and clinical parameters that will improve the sensitivity and specificity of the biomarker for identifying or otherwise refining the patient population. Alternatively, these additional biological and clinical parameters can be used alone as a means for identifying patients that are suitable candidates for treatment with an anti-IFN $\gamma$  antagonist or other suitable therapy. These biological and clinical parameters include, by way of non-limiting example, any of the following: ferritin levels, neutrophil count, platelet count, alanine aminotransferase levels, and/or lactate dehydrogenase levels.

[0231] Disorders that are useful with the compositions and methods of the invention include any disorder where aberrant, *e.g.*, elevated, IFN $\gamma$  expression and/or activity, particularly HLH, including secondary HLH, MAS, and/or sJIA.

[0232] By way of non-limiting examples, the methods and compositions provided herein are suitable for diagnosing and/or treating disorders such as primary and/or secondary HLH disorders. Suitable autoimmune and/or inflammatory disorders include, by way of non-limiting example, primary and/or secondary HLH disorders associated with aberrant IFN $\gamma$  activity and/or expression.

[0233] Once patients are identified as having an elevated level of CXCL9, alone or in combination with one or more IFN $\gamma$ -related biomarkers, they are then treated with an anti-IFN $\gamma$

antagonist. For example, the anti-IFN $\gamma$  antagonist is a neutralizing anti-IFN $\gamma$  antibody or an immunologically active (*e.g.*, antigen binding) fragment thereof. Suitable neutralizing anti-IFN $\gamma$  antibodies include any of the anti-IFN $\gamma$  antibodies described herein.

**[0234]** In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a variable heavy chain complementarity determining region 1 (VH CDR1) comprising an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) comprising an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) comprising an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

**[0235]** In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a VH CDR1 comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a VH CDR2 region comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a VH CDR3 region comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) region comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a VL CDR2 region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a VL CDR3 region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

**[0236]** In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment

thereof comprises a heavy chain that comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 1A.

[0237] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a light chain that comprises a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 1B.

[0238] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a heavy chain that comprises a combination of a VH CDR1 sequence, a VH CDR2 sequence, and a VH CDR3 sequence, wherein the combination is a combination of the three heavy chain CDR sequences (VH CDR1, VH CDR2, VH CDR3) shown in a single row in Table 1A, and a light chain that comprises a combination of a VL CDR1 sequence, a VL CDR2 sequence, and a VL CDR3 sequence, wherein the combination is a combination of the three light chain CDR sequences (VL CDR1, VL CDR2, VL CDR3) shown in a single row in Table 1B.

[0239] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47.

[0240] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

[0241] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

[0242] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof

comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47.

[0243] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

[0244] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

[0245] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the heavy chain amino acid sequence to the amino acid sequence of SEQ ID NO: 44.

[0246] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the light chain amino acid sequence to the amino acid sequence of SEQ ID NO: 46.

[0247] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the heavy chain amino acid sequence to the amino acid sequence of SEQ ID NO: 44, and an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to the light chain amino acid sequence to the amino acid sequence of SEQ ID NO: 46.

[0248] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises the heavy chain amino acid sequence to the amino acid sequence of SEQ ID NO: 44.

[0249] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises the light chain amino acid sequence to the amino acid sequence of SEQ ID NO: 46.

[0250] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises the heavy chain amino acid sequence to the amino acid sequence of SEQ ID NO: 44, and the light chain amino acid sequence to the amino acid sequence of SEQ ID NO: 46.

[0251] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to a heavy chain variable amino acid sequence selected from the group consisting of

SEQ ID NOs: 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, and 102.

[0252] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to a light chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, and 104.

[0253] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to a heavy chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, and 102, and an amino acid sequence at least 90%, 92%, 95%, 96%, 97% 98%, 99% or more identical to a light chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, and 104. In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a heavy chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, and 102.

[0254] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a light chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, and 104.

[0255] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof comprises a heavy chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, and 102, and a light chain variable amino acid sequence selected from the group consisting of SEQ ID NOs: 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, and 104.

[0256] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof is administered in a therapeutically effective amount. A therapeutically effective amount of an antibody of the invention relates generally to the amount needed to achieve a therapeutic objective. This therapeutic objective may be a binding interaction between the antibody and its target antigen that, in certain cases, interferes with the functioning of the target. Common ranges for therapeutically effective dosing of an antibody or antibody fragment of the invention may be, by way of nonlimiting example, from about 0.1 mg/kg body weight to about 50 mg/kg body weight. Common dosing frequencies may range, for example, from twice daily to once a week.

[0257] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof

is administered at an initial dose, i.e., load dose, in the range from about 0.5 mg/kg to about 2 mg/kg, for example, in a range from about 0.5 mg/kg to about 1.5 mg/kg, and/or from about 0.5 mg/kg to about 1.0 mg/kg. In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof is administered at an initial dose of about 1.0 mg/kg.

[0258] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof is administered as an initial load dose followed by one or more maintenance doses. In some embodiments, the one or more maintenance dose(s) is a dosage that is substantially similar to the initial load dose. In some embodiments, the one or more maintenance dose(s) is a dosage that is less than the initial load dose. In some embodiments, the one or more maintenance dose(s) is a dosage that is greater than the initial load dose.

[0259] In some embodiments, the one or more maintenance dose(s) comprises at least two or more dosages, wherein each maintenance dosage is the same dosage. In some embodiments, the two or more maintenance dosages are substantially similar to the initial load dose. In some embodiments, the two or more maintenance dosages are greater than the initial load dose. In some embodiments, the two or more maintenance dosages are less than the initial load dose.

[0260] In some embodiments, the one or more maintenance dose(s) comprises at least two or more dosages, wherein each maintenance dosage is not the same dosage. In some embodiments, the two or more maintenance dosages are administered in an increasing dosage amount. In some embodiments, the two or more maintenance dosages are administered in a decreasing dosage amount.

[0261] In some embodiments, the one or more maintenance dose(s) comprises at least two or more dosages, wherein each maintenance dosage is administered at a periodic time interval. In some embodiments, two or more dosages are administered at increasing time intervals. In some embodiments, two or more dosages are administered at decreasing time intervals.

[0262] In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof is administered at an initial load dose in the range from about 0.5 mg/kg to about 2 mg/kg, for example, in a range from about 0.5 mg/kg to about 1.5 mg/kg, and/or from about 0.5 mg/kg to about 1.0 mg/kg, followed by at least one, e.g., two or more, three or more, four or more, or five or more maintenance doses. In some embodiments, the anti-IFN $\gamma$  antibody or immunologically active fragment thereof is administered at an initial load dose of about 1.0 mg/kg, followed by at least one, e.g., two or more, three or more, four or more, or five or more maintenance doses.

[0263] Pharmaceutical compositions according to the invention can include an anti-IFN $\gamma$  antibody of the invention and a carrier. These pharmaceutical compositions can be included in kits, such as, for example, diagnostic kits.

[0264] The invention also provides kits for practicing any of the methods provided herein. For example, in some embodiments, the kits include a detection reagent specific for CXCL9, alone or in combination with one or more IFN $\gamma$ -related biomarkers and a means for detecting the detection reagent.

[0265] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

### EXAMPLES

#### **Example 1. CXCL9 LEVELS AS A BIOMARKER FOR IFN $\gamma$ PRODUCTION IN MACROPHAGE ACTIVATION SYNDROME (MAS)**

[0266] The studies presented herein were designed to evaluate the correlation between serum levels of IFN $\gamma$  and of the three IFN $\gamma$  related chemokines with themselves and with laboratory parameters of disease activity in patients with active MAS in order to search for a potential biomarker of IFN $\gamma$  *in vivo* production.

[0267] Circulating levels of IFN $\gamma$ , CXCL9, CXCL10, CXCL11 and IL-6 were measured using a Luminex multiplexing assay in patients with sJIA (n=54) of whom 20 had MAS at time of sampling. The relation of these circulating levels to disease activity parameters was evaluated, along with correlations of the levels of IFN $\gamma$  with those of CXCL9, CXCL10 and CXCL11.

[0268] Levels of IFN $\gamma$  and of the 3 IFN $\gamma$ -related chemokines (CXCL9, CXCL10 and CXCL11) were significantly elevated in active MAS compared to active sJIA without MAS at sampling (all p-values <0.005). In active MAS laboratory parameters of disease severity (ferritin, neutrophils, platelets, alanine aminotransferase and lactate dehydrogenase) were significantly correlated with IFN $\gamma$  and CXCL9, and to a lesser extent with CXCL10 and CXCL11; no correlation with IL-6 levels was found. In patients with active sJIA without MAS there was no significant correlation between laboratory parameters and cytokine levels as shown in Table 7 below. In active MAS IFN $\gamma$  levels were significantly correlated with levels of CXCL9 ( $r=0.69$ ;  $r^2=0.47$ ;  $p=0.001$ ), to a lesser extent with levels of CXCL10 ( $r=0.53$ ;

$r^2=0.28$ ;  $p=0.015$ ), and not with levels of CXCL11 ( $r=-0.04$ ;  $p=0.886$ ). Table 7. Correlation of laboratory parameters of disease activity with IFN $\gamma$ , CXCL9, CXCL10, CXCL11, and IL-6 in patients with MAS and in patients with active sJIA.

|              | Macrophage<br>Activation Syndrome         | IFN $\gamma$ |       | CXCL9 |       | CXCL10 |       | CXCL11 |       | IL-6  |       |
|--------------|-------------------------------------------|--------------|-------|-------|-------|--------|-------|--------|-------|-------|-------|
|              |                                           | r*           | p     | r*    | p     | r*     | p     | r*     | p     | r*    | p     |
| Neutrophil N | 8000 (3159 - 13174)                       | 0.57         | 0.014 | 0.49  | 0.041 | 0.66   | 0.002 | 0.62   | 0.023 | 0.17  | >0.1  |
| PLT          | 6.9 (3.4 - 13.9)                          | -0.64        | 0.005 | -0.61 | 0.010 | -0.37  | >0.1  | -0.08  | >0.1  | 0.09  | >0.1  |
| ALT          | 198 (115 - 392)                           | -0.53        | 0.017 | -0.52 | 0.022 | -0.58  | 0.008 | -0.22  | >0.1  | -0.02 | >0.1  |
| LDH          | 46 (18 - 164)                             | 0.49         | 0.045 | 0.49  | 0.044 | 0.51   | 0.038 | 0.06   | >0.1  | -0.44 | 0.080 |
|              | 1152 (722 - 2135)                         | 0.45         | 0.095 | 0.62  | 0.013 | 0.64   | 0.010 | 0.64   | 0.048 | 0.08  | >0.1  |
|              | Systemic Juvenile<br>Idiopathic Arthritis | IFN $\gamma$ |       | CXCL9 |       | CXCL10 |       | CXCL11 |       | IL-6  |       |
|              |                                           | r*           | p     | r*    | p     | r*     | p     | r*     | p     | r*    | p     |
| Neutrophil N | 215 (38 - 1669)                           | -0.27        | >0.1  | 0.28  | >0.1  | 0.27   | >0.1  | 0.29   | >0.1  | -0.12 | >0.1  |
| PLT          | 8.4 (5.2 - 14.5)                          | 0.30         | >0.1  | 0.40  | 0.061 | 0.32   | >0.1  | 0.40   | 0.067 | 0.28  | >0.1  |
| ALT          | 444 (353 - 544)                           | 0.21         | >0.1  | -0.14 | >0.1  | -0.13  | >0.1  | 0.27   | >0.1  | 0.35  | 0.064 |
| LDH          | 16 (11 - 24)                              | 0.29         | >0.1  | 0.42  | 0.049 | 0.50   | 0.011 | 0.44   | 0.039 | 0.04  | >0.1  |
|              | 506 (456 - 851)                           | 0.07         | >0.1  | 0.49  | >0.1  | 0      | >0.1  | 0.26   | >0.1  | 0     | >0.1  |

N = neutrophil count; PLT = platelet count; ALT = alanine aminotransferase, <sup>1</sup> = Median (IQR); r\* = Spearman r

[0269] The high levels of IFN $\gamma$  and of CXCL9 present in patients with active MAS are significantly correlated with laboratory parameters of disease severity. In patients with active MAS IFN $\gamma$  and CXCL9 are tightly correlated. Since CXCL9 has been shown to be induced only by IFN $\gamma$  and not by other interferons (*see e.g.*, Groom J.R. and Luster A.D. Immunol Cell Biol 2011, Feb; 89(2):207-15), these findings demonstrate that CXCL9 is a biomarker of IFN $\gamma$  production in MAS.

**Example 2. CXCL9 AND IFN $\gamma$  LEVEL CORRELATION IN PRIMARY HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH) PATIENTS**

[0270] The studies presented herein are from an ongoing phase 2 pilot study in primary HLH patients who were administered the NI-0501 antibody and from patients who received the NI-0501 antibody in compassionate use.

[0271] As shown in Figure 1, serum levels of CXCL9 and IFN $\gamma$  were measured by

Luminex and Meso Scale Discovery (MSD) technology, respectively, in samples obtained from 6 primary HLH patients and from 3 compassionate use patients. Correlations were performed between CXCL9 and total IFN $\gamma$  concentrations. Statistics were performed and p values were obtained using the Spearman test.

[0272] As shown in Figure 2, predose serum levels of CXCL9 and IFN $\gamma$  were measured by Luminex and MSD technology, respectively, in samples obtained from 6 primary HLH patients and 3 compassionate use patients. Correlations were performed between CXCL9 and total IFN $\gamma$  concentrations. Statistics were performed and p values obtained using the Spearman test.

### **Example 3. CXCL9 AND IFN $\gamma$ LEVEL CORRELATION IN SECONDARY HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH) PATIENTS**

[0273] The studies presented herein are from an observational study in secondary HLH patients who were administered the NI-0501 antibody and from patients who received the NI-0501 antibody in compassionate use.

[0274] In particular, these are patients with systemic Juvenile Idiopathic Arthritis (sJIA) who developed Macrophage Activation Syndrome (MAS, a form of secondary HLH). For these patients, there are also correlations between CXCL9 or IFN $\gamma$  and disease parameters such as Ferritin, Platelet count (PLT), Neutrophil count (Neu), and Alanine Aminotransferase (ALT).

[0275] As shown in Figures 3A and 3B, serum levels of CXCL9 and IFN $\gamma$  were measured with by multiplex assay using the Luminex technology from samples obtained from 19 patients with MAS secondary to sJIA and 24 patients with active sJIA at the time of sampling. Correlations were performed between CXCL9 and IFN $\gamma$  concentrations. Statistics were performed and p values obtained using the Spearman test.

[0276] As shown in Figures 4A-1, 4A-2, 4B-1, 4B-2, 4C-1, 4C-2, 4D-1, and 4D-2, serum levels of CXCL9 and IFN $\gamma$  were measured by multiplex assay using the Luminex technology from patients with MAS secondary to sJIA and patients with active sJIA at the time of sampling. Correlations were performed between IFN $\gamma$  or CXCL9 levels and ferritin, platelet count, neutrophil count or ALT (alanine aminotransferase). Statistics were performed and p values obtained using the Spearman

test.

**Example 4. CXCL9 AND IFN $\gamma$  LEVEL CORRELATION IN SEVERE HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH) PATIENT**

[0277] The study presented herein is from a patient who received the NI-0501 antibody in compassionate use. This patient exhibited symptoms of NLRC4-related disease and severe hemophagocytic lymphohistiocytosis (HLH). Mutations in *NLRC4* gene have recently been reported to cause recurrent macrophage activation syndrome and increased production of IL-18, which is known to induce IFN $\gamma$ .

[0278] The patient in this study had the following characteristics: Onset at 20 days of age with fever, rash, marked hepatosplenomegaly, pancytopenia, hypofibrinogenemia, hypertriglyceridemia, marked ferritin and sCD25 increase. Followed by multiorgan failure, required ICU admission. HLH diagnosis was based on 6 out of the 8 HLH-2004 criteria. Gene causing primary-HLH (PRF1, UNC13D, STXBP2, STX11, RAB27A, XIAP) and functional tests (perforin expression, degranulation and cytotoxicity) were negative. High-dose i.v. glucocorticoids and i.v. cyclosporine-A with progressive improvement of general conditions and laboratory abnormalities. HLH reactivation triggered by infections (*Candida Albicans* and *Klebsiella Pneumoniae* sepsis), rapid worsening of general conditions, and a new ICU admission. Treatment with etoposide and/or ATG was not considered because of the presence of active infections in an already immunocompromised subject.

[0279] Measurable serum levels of IFN $\gamma$  and high serum levels of the IFN $\gamma$ -induced chemokines CXCL9 and CXCL10 were documented as well as markedly elevated serum levels of IL-18 (Table 8).

Table 8. Levels of IFN $\gamma$ , IFN $\gamma$ -related chemokines and IL-18 when NI-0501 treatment was started and during treatment with NI-0501

|                               | Before treatment | 1 mo after treatment | 2 mo after treatment | 3 mo after treatment | 4 mo after treatment | sJIA* Inactive |
|-------------------------------|------------------|----------------------|----------------------|----------------------|----------------------|----------------|
| Free IFN $\gamma$ (pg/ml)     | 6.02             | nd                   | nd                   | nd                   | nd                   | 4.2 (3.2-9.3)  |
| CXCL9 (pg/ml)                 | 5670             | 495.16               | 207.4                | 207.4                | 207.4                | 901 (466-1213) |
| CXCL10 (pg/ml)                | 4400             | 529.54               | 201.12               | 147.84               | 138                  | 235 (172-407)  |
| CXCL11 (pg/ml)                | 188.68           | nd                   | nd                   | nd                   | nd                   | 111 (63-187)   |
| IL18 (pg/ml)                  | >300000          | 95000                | 34000                | 27000                | 32000                | -              |
| *Median (interquartile range) |                  |                      |                      |                      |                      |                |

[0280] Compassionate use treatment with NI-0501 was started on a background of dexamethasone (13.6 mg/m<sup>2</sup>) and i.v. cyclosporine-A. NI-0501 was administered every 3 and subsequently every 7 days according to pharmacokinetics. No infusion reaction was observed. NI0501 was well tolerated. HLH clinical features and laboratory abnormalities progressively improved. Active ongoing infections were rapidly cleared. After 5 months of treatment, the patient remained in excellent conditions. The patient continued to receive oral cyclosporine-A (6 mg/kg) and prednisone (0.3 mg/kg equivalent to 0.9 mg/m<sup>2</sup> of dexamethasone). All HLH parameters have normalized.

[0281] The subject still presented unexplained episodes of inflammation Analysis of *NLRC4* showed a *de novo* missense mutation (T337N). Elevated serum IL-18 was documented, confirming the relevance of the *NLRC4* mutation. High production of IFN $\gamma$  was demonstrated by high levels IFN $\gamma$  complexed with NI-0501. IFN $\gamma$  was fully neutralized as shown by undetectable levels of IFN $\gamma$ -inducible chemokines (Figure 5 and Table 8). Circulating levels of were IL-18 persistently elevated.

[0282] Thus, this study demonstrates that in a patient with severe recalcitrant HLH (due to *NLRC4* mutation), blocking IFN $\gamma$  with NI-0501 was well tolerated with no safety concerns, allowed control of all HLH features, allowed rapid glucocorticoid tapering, and was associated with resolution of ongoing active infections.

**Example 5. TARGETED APPROACH TO THE TREATMENT OF  
HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH) WITH NI-0501**

[0283] The study presented herein is from a pilot phase 2 study in children with primary HLH. Primary HLH (pHLH) is a rare immune regulatory disorder, which is invariably lethal if untreated. It is driven by a pathologic immune activation, leading to the development of fever, splenomegaly, cytopenias and coagulopathy, which may cause multi-organ failure and death. Based on data from murine models of primary and secondary HLH (sHLH) treated with an anti-IFN $\gamma$  antibody, and observational studies in patients with HLH, the high production of IFN $\gamma$  is thought to be a critical factor driving development of the disease. Immune-chemotherapy, primarily etoposide-based regimens, are at present the only pharmacological approaches to control HLH and bring patients to curative allogeneic hematopoietic stem cell transplant (allo-HSCT). In spite of recent attempts to further intensify treatment regimens, mortality and morbidity remain high, in part due to drug-related toxicities.

[0284] As described above, NI-0501 is a fully human, high affinity, anti-IFN $\gamma$  mAb that binds to and neutralizes human IFN $\gamma$ , offering a novel and targeted approach for the control of HLH.

[0285] *Methods:* An open-label Phase 2 study has been conducted in United States and Europe to evaluate the safety and efficacy of NI-0501 in children with confirmed or suspected pHLH. NI-0501 was administered at the initial dose of 1 mg/kg every 3 days, with possible dose increase guided by PK data and/or clinical response in each patient, on initial background dexamethasone 5-10 mg/m<sup>2</sup>. Treatment duration ranged from 4 to 8 weeks. Ability to move to allo-HSCT, relevant HLH disease parameters, and 8-week survival were assessed.

[0286] *Study Population:* A total of 13 patients were enrolled: 8F/5M, median age 1.0y (range 2.5mo-13y). Twelve pts received NI-0501 as a second line treatment after having received conventional therapy and either reactivating, obtaining an unsatisfactory response, or being intolerant to therapy. One patient was treated with NI-0501 in 1<sup>st</sup> line. Nine patients carried a known HLH genetic defect (3 FHL2, 2 FHL3, 2 GS-2, 1 XLP1, 1 XLP2). The majority of patients were at the severe end of HLH spectrum, in compromised general condition, carrying significant toxicities from previous HLH treatments. Ferritin was elevated in 12/13 patients and sCD25 in 8, cytopenias were present in 10 patients, splenomegaly in 8, hypofibrinogenemia and hypertriglyceridemia in 9. Liver impairment and CNS involvement were present in 7 and 3 patients, respectively.

[0287] *Results:* Overall, NI-0501 treatment significantly improved parameters of HLH disease activity (Fig. 6), and 9 of 13 patients achieved a satisfactory response. Six patients have proceeded to HSCT. Two patients with good HLH control are planning to proceed to HSCT upon identification of an appropriate donor. In one patient (who achieved disease control with 1<sup>st</sup> line NI-0501) HSCT is not yet planned given the absence of a causative HLH gene mutation. Eleven of 13 patients were alive at 8 weeks. CNS signs and symptoms resolved in the 2 evaluable patients. More than 50% reduction of dexamethasone dose was possible in 50% of the patients during the first 4 weeks of NI-0501 treatment.

[0288] Biomarker assessment in particular CXCL9, a chemokine known to be induced exquisitely by IFN $\gamma$ , not only allowed demonstrating full IFN $\gamma$  neutralization, but appears as a new parameter for the diagnosis of HLH, correlating with IFN $\gamma$  production (Figs. 7A and 7B).

[0289] NI-0501 was well tolerated and no safety concern was identified. None of the infections known to be favored by IFN $\gamma$  neutralization was reported, and no infections occurred in patients who did not receive previous chemotherapy. Seven patients reported at least one SAE, all assessed by the DMC as not related to NI-0501 administration. No unexpected events attributable to “off target” effects of NI-0501 (e.g. myelotoxicity, hemodynamic effects) were observed.

[0290] *Conclusions:* Targeted neutralization of IFN $\gamma$  by NI-0501 offers an innovative and potentially less toxic approach to HLH management. The results of this study show that NI- 0501 is a safe and effective therapeutic option in patients with primary HLH who have unsatisfactorily responded to conventional therapy or shown intolerance to it. Furthermore, therapy with NI-0501 was not associated with any of the typical short or long-term toxicities associated with etoposide-based regimens. Assessment of NI-0501 as 1<sup>st</sup> line treatment in patients with pHLH is ongoing, anticipating that similar significant clinical benefit can be achieved.

**Example 6. ELEVATED CIRCULATING LEVELS OF INTERFERON- $\gamma$  AND INTERFERON- INDUCED CHEMOKINES CHARACTERIZE PATIENTS WITH MACROPHAGE ACTIVATION SYNDROME COMPLICATING SYSTEMIC JIA**

[0291] Interferon gamma (IFN $\gamma$ ) is the pivotal mediator in murine models of primary haemophagocytic lymphohistiocytosis (HLH). Given the similarities between primary and secondary HLH (sec-HLH), including macrophage activation syndrome (MAS), IFN $\gamma$  levels and its biologic activity in patients with systemic juvenile idiopathic arthritis (sJIA) and MAS were analyzed.

[0292] In the studies provided herein, the Luminex multiplexing assay was used to assess serum levels of IL-1 $\beta$ , IL-6, IFN $\gamma$ , and of the IFN-induced and/or IFN-related chemokines CXCL9, CXCL10, and CXCL11 in patients with sec-HLH (n=11), and in patients with sJIA (n=54) of whom 20 had MAS at sampling. Expression of IFN $\gamma$ -induced chemokines (CXCL9 and CXCL10 mRNA levels in liver and spleen), as well as their correlation with serum ferritin levels were assessed in an IL-6 transgenic mouse model in which MAS features are induced by TLR4 stimulation with LPS.

[0293] As will be shown in more detail below, circulating levels of IFN $\gamma$  and of IFN-induced chemokines were markedly elevated during MAS, also referred to herein as active MAS, and sec-HLH. Levels of IFN $\gamma$  and IFN induced chemokines were markedly higher in patients with MAS compared to those with active sJIA without MAS. In this latter group, IFN $\gamma$  and IFN $\gamma$  induced chemokines were comparable to those of patients with clinically inactive sJIA. During MAS, the laboratory abnormalities characterizing this syndrome, including ferritin and alanine transferase levels and neutrophil and platelet count, were significantly correlated with levels of IFN $\gamma$  and CXCL9. In a murine model of MAS, serum levels of ferritin were significantly correlated with mRNA levels of CXCL9 in liver and spleen.

[0294] Thus, the studies presented below demonstrate that the high levels of IFN $\gamma$  and of IFN-induced chemokines and their correlation particularly for CXCL9 with the severity of laboratory abnormalities of MAS suggest that IFN $\gamma$  plays a pivotal role in MAS. Elevated circulating levels of interferon- $\gamma$  and interferon- induced chemokines characterize patients with macrophage activation syndrome complicating systemic JIA.

[0295] *Materials and Methods:* Patients and Samples. Peripheral blood samples were collected from patients with sJIA with or without MAS in 3 Paediatric Rheumatology Centres:

the Ospedale Pediatrico Bambino Gesù in Rome, the Istituto Giannina Gaslini in Genoa and the Cincinnati Children's Hospital Medical Centre. Fifty-four patients with sJIA (age at onset 7.9 years, interquartile range 4.6-13.6 years; female 48%) who met the ILAR classification criteria for systemic arthritis were studied (Petty, R.E., et al., *International League of Associations for Rheumatology classification of juvenile idiopathic arthritis: second revision, Edmonton, 2001*. J Rheumatol, 2004. **31**(2): p. 390-2). For twenty of the SJIA patients, samples were collected during episodes of active full-blown MAS, as diagnosed by the treating physicians at each of the three centers. An *a posteriori* analysis showed that 17 of these 20 episodes (85%) met the newly proposed MAS classification criteria (Minoia F, Davì S, Bovis F, et al. Development of new classification criteria for macrophage activation syndrome complicating systemic juvenile idiopathic arthritis. Pediatric Rheumatology 2014, 12(Suppl 1):O1). Twenty-eight patients with active SJIA without evidence of MAS had samples available. Thirty-five samples were available from 35 sJIA patients (both with or without MAS in their disease history) during clinically inactive disease, defined according to Wallace's criteria (Wallace, C.A., et al., *Preliminary criteria for clinical remission for select categories of juvenile idiopathic arthritis*. J Rheumatol, 2004. **31**(11): p. 2290-4).

[0296] Since IFN $\gamma$  has been shown to be increased in patients with sec-HLH (a rheumatic disease was excluded), samples were collected also from 11 patients (age at onset 8.6 years, interquartile range 4.1-12.9 years; female 36%) with sec-HLH, seen at the Ospedale Pediatrico Bambino Gesù, and used as positive controls. All sec-HLH patients met the 2004-HLH diagnostic guidelines (Henter, J.I., et al., *HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis*. Pediatr Blood Cancer, 2007. 48(2): p. 124-31): 6 patients met 5 criteria and 5 patients met 4 criteria. It should be noted that levels of sCD25 in U/ml were not available as the test is not performed routinely in the institution where these patients were recruited. A diagnosis of primary HLH was excluded based on the absence of family history, the absence of pathogenic mutations in the genes known to cause HLH and the presence of normal functional studies (including NK activity, perforin expression and CD107 degranulation). All 11 patients with sec-HLH contributed one sample each obtained during active disease.

[0297] Clinical and laboratory features of all patients concerning diagnosis and at time of sampling were collected in a centralized web database by the investigators of each center. Of the

20 MAS patients sampled during active disease, 6 were not receiving any treatment at time of sampling, while the remaining 14 patients had already received one of the treatments specific for MAS, including glucocorticoids pulses, cyclosporine A, anakinra or cyclophosphamide. Six out of 11 patients with sec-HLH in active disease were not yet receiving specific treatment at time of sampling, while the remaining 5 patients had already received at least one among the above mentioned treatments. The Ethical Committee of the Ospedale Pediatrico Bambino Gesù approved the study. Written consent was collected for all participants.

**[0298]** Quantification of Cytokines. Levels of IL-6, IL-1 $\beta$ , IFN $\gamma$ , CXCL9, CXCL10 and CXCL11 were analyzed by Luminex® multiplexing beads technology. Reagents were purchased from Millipore, and all reagents were provided with the Milliplex® MAP kits. Reagents were prepared according to the manufacturers' protocol. 25  $\mu$ l/well of standards, blank and Quality Check samples were added in duplicate in the Milliplex MAP 96-well plate, followed by an addition of 25  $\mu$ l of Serum Matrix. 25  $\mu$ l of Assay Buffer was added to each sample well followed by the addition of 25  $\mu$ l sample. Samples are added in duplicate or triplicate, depending on the available volume of sample. The plate was measured on the Luminex 200® system (Luminex Corp.). Raw data were acquired using xPONENT software version 3.1 (Luminex Corp.), and data were analyzed using Milliplex Analyst software version 3.5.5.0 (Millipore). Raw data obtained in Milliplex Analyst software were then further analyzed in dedicated macro for Luminex analysis (NI-Sc-ESM-MAC-012-v01 and Sc-ESM-MAC-013-v01).

**[0299]** Animal Experiments. The generation and the phenotype of the IL-6 transgenic mice, as well as the features of the MAS-like syndrome induced by administration of TLR ligands, have been described previously (Strippoli, R., et al., *Amplification of the response to Toll-like receptor ligands by prolonged exposure to interleukin-6 in mice: implication for the pathogenesis of macrophage activation syndrome*. Arthritis Rheum, 2012. 64(5): p. 1680-8). Mice were maintained under specific pathogen-free conditions and handled in accordance with the national policies. The study protocol was approved by the local ethics committee. All experiments have been performed on mice between 10 and 14 weeks of age. Mice were administered intraperitoneally with a single dose of 5  $\mu$ g/g body weight of lipopolysaccharide (LPS, E. coli serotype 055:B5; Sigma-Aldrich). Mice were sacrificed after 30 hours. Total RNA was extracted from spleen and liver tissues using Trizol (Life technologies). cDNA was obtained using the Superscript Vilo kit (Invitrogen). Real-time PCR assays were performed using the TaqMan Universal PCR Master Mix (Applied Biosystems) with the mouse *Cxcl9* and *Cxcl10* gene-expression assays (Applied Biosystems). Gene

expression data were normalized using mouse *Hprt* (Applied Biosystems). Data are expressed as arbitrary units (AU), determined using the  $2^{-\Delta\text{ct}}$  method. Serum ferritin concentrations were determined using a commercially available ELISA kit (ALPCO Diagnostics), according to the manufacturer's instructions.

**[0300]** Statistical Analysis. Statistical analysis was performed using the GraphPad Prism 5 software. Continuous variables (quantitative demographic, clinical and laboratory data) were expressed as medians and interquartile ranges (IQR) and were compared using the Mann-Whitney U test. The Wilcoxon signed rank test was used to compare two paired groups without assuming that the distribution of the before-after differences follow a Gaussian distribution. Spearman rank correlation was used to assess the relation with laboratory parameters. A p value <0.05 was considered statistically significant.

**[0301]** *Results:* Increased levels of IFN $\gamma$  and IFN $\gamma$ -induced chemokines in patients with MAS. When patients with active sJIA without MAS at sampling were compared with patients sampled during clinically inactive disease, it was found, as expected (de Benedetti, F., et al., *Correlation of serum interleukin-6 levels with joint involvement and thrombocytosis in systemic juvenile rheumatoid arthritis*. Arthritis Rheum, 1991. 34(9): p. 1158-63), that IL-6 levels were significantly higher in patients with active sJIA (p<0.01) compared to those of patients with clinically inactive disease. As has been reported in several previous studies of active SJIA, serum IL-1 $\beta$  levels were below the limit of detection in the majority of the patients, independently from the disease activity state. It is noteworthy that there were no differences in the levels of IFN $\gamma$  and of three IFN $\gamma$ -induced chemokines among patients with clinically active sJIA and patients with clinically inactive disease.

**[0302]** When patients with MAS at time of sampling were compared with patients with active sJIA without MAS at sampling, levels of IL-1 $\beta$  and of IL-6 were comparable, suggesting that the levels of the two cytokines that are known to play a pivotal role in active sJIA do not increase during full-blown MAS. It should be noted that circulating IL-1 $\beta$  levels were below the limit of quantification (i.e., 3.5 pg/ml) in the majority of patients with sJIA with or without MAS. In contrast, circulating IFN $\gamma$  levels were significantly higher in patients with active MAS compared to patients with active sJIA without MAS at sampling. The levels of the three IFN $\gamma$  related chemokines CXCL9, CXCL10 and CXCL11 were also markedly higher in patients with active MAS compared to patients with active sJIA without MAS at sampling. This difference was particularly evident for CXCL9 of which median levels were approximately 15 fold higher in patients with MAS compared to patients with active sJIA without MAS.

[0303] In patients with sec-HLH, levels of IFN $\gamma$  as well as levels of the three IFN $\gamma$ - related chemokines were markedly increased. The levels of IFN $\gamma$  and of the IFN $\gamma$ -related chemokines were largely indistinguishable from those of patients with MAS and the differences were not statistically significant. Incidentally, in patients with active MAS and in patients with active sec-HLH levels of IFN $\gamma$  and of the three IFN $\gamma$ -induced chemokines were comparable in patients receiving no treatment and in patients already receiving treatment.

[0304] Levels of IFN $\gamma$  and of CXCL9, CXCL10 and CXCL11 are related to the presence of MAS in individual patients. Figures 8A-8D show the levels of IFN $\gamma$  and of CXCL9, CXCL10 and CXCL11 in individual patients from whom paired samples were available during active MAS and during active sJIA without MAS. In concordance with the results obtained in the cross- sectional analysis, the levels of IFN $\gamma$  and of the three IFN $\gamma$ -induced chemokines were significantly higher in samples obtained during MAS by paired sample analysis. Additionally, in several patients samples were available both prior to and after MAS episodes, and demonstrated that IFN $\gamma$  and IFN $\gamma$ -induced chemokine levels return to normal upon resolution of MAS clinical symptoms. For example, one patient in this study experienced three episodes of MAS, with serum samples obtained during these episodes as well as during disease phases without MAS at sampling. Further confirming the relation between increased production of IFN $\gamma$  and of the three IFN $\gamma$ -induced chemokines with active MAS, in this patient elevated levels of IFN $\gamma$  and of the three IFN $\gamma$ -related chemokines were found only at time of the MAS episodes (Figures 9A-9B).

[0305] Levels of IFN $\gamma$  and of the IFN $\gamma$  related chemokines correlate with laboratory abnormalities of MAS. The correlation of the levels of IFN $\gamma$  and of the three IFN $\gamma$ -induced chemokines with laboratory parameters of MAS at time of sampling was then examined. In patients with active sJIA without MAS, levels of IFN $\gamma$  and of the three IFN $\gamma$ -induced chemokines were not associated with laboratory parameters of MAS with one exception: levels of CXCL9, CXCL10 and CXCL11 were weakly correlated with ALT levels with  $r^2$  ranging from 0.17 to 0.25 (Table 2). The significance of this association is unclear; however it should be noted that ALT levels were within normal range in all patients with active sJIA without MAS. In patients with MAS at sampling, no significant correlation with laboratory features of MAS with IL-1 and IL-6 were found. In contrast, in patients with MAS at sampling, levels of IFN $\gamma$  and of the IFN $\gamma$ -induced chemokines were associated with levels of ferritin, with neutrophil and platelet counts, and with increased LDH and ALT, all typically abnormal in patients with MAS (Table 2). The correlations with laboratory abnormalities

were particularly evident for IFN $\gamma$  and for CXCL9 with the only exception of the correlation of IFN $\gamma$  with LDH, which did not reach statistical significance (Table 2 and Figures 10A-10J). Again, as mentioned above, these correlations were not present in patients with active s-JIA without MAS at sampling. One patient in this group had markedly high levels of IFN $\gamma$  (336.2 pg/ml), CXCL9 (549400 pg/ml) and CXCL10 (35066 pg/ml). This patient had particularly severe MAS and was admitted to the intensive care unit with severe central nervous system involvement. This observation provides further support for the hypothesis that there is a strong association between levels of IFN $\gamma$  and CXCL9 and disease severity. Taken together, these results show that increased production of IFN $\gamma$  and of the IFN $\gamma$ -related chemokines is a feature of active MAS that strongly correlates with the severity of the laboratory abnormalities of MAS.

Table 2. Serum levels of IL-1 $\beta$ , IL-6, IFN $\gamma$  and of the three IFN $\gamma$  related chemokines CXCL9, CXCL10 and CXCL11 in patients with active secondary HLH, with active MAS at sampling, with active sJIA without MAS at sampling, and with clinically inactive sJIA.

|                      | Active sJIA             |                        |                        |                        | sec-HLH | MAS                     |
|----------------------|-------------------------|------------------------|------------------------|------------------------|---------|-------------------------|
|                      | sec-HLH                 | MAS at sampling        | without MAS            | inactive sJIA          | vs MAS  | Active sJIA without MAS |
|                      | (n=11)                  | (n=20)                 | (n=28)                 | (n=35)                 | p value | p value                 |
| IL-1 $\beta$ (pg/ml) | < 3.5<br>(< 3.5 – 10.7) | < 3.5<br>(< 3.5 – 6.1) | < 3.5<br>(< 3.5 – 3.8) | < 3.5<br>(< 3.5 – 3.5) | 0.69    | 0.86                    |
| IL-6 (pg/ml)         | 11.4<br>(3.2 – 49.3)    | 22.9<br>(5.5 – 45.6)   | 20.3* (5.9 – 54.9)     | 3.2<br>(3.2 – 7.9)     | 0.56    | 0.43                    |
| IFN $\gamma$ (pg/ml) | 34.7<br>(23.9 –         | 15.4<br>(5.1 – 52.6)   | 4.9<br>(3.2 – 8.6)     | 4.2<br>(3.2 – 9.3)     | 0.12    | 0.03                    |

|                |                          |                         |                     |                     |      |        |
|----------------|--------------------------|-------------------------|---------------------|---------------------|------|--------|
|                | 170.1)                   |                         |                     |                     |      |        |
| CXCL9 (pg/ml)  | 33598<br>(3083 – 127687) | 13392<br>(2163 – 35452) | 837<br>(471 – 2505) | 901<br>(466 – 1213) | 0.23 | 0.005  |
| CXCL10 (pg/ml) | 4420<br>(799 – 8226)     | 1612<br>(425 – 4309)    | 307<br>(199 – 694)  | 235<br>(172 – 407)  | 0.19 | 0.0016 |
| CXCL11 (pg/ml) | 1327<br>(189 – 2000)     | 565<br>(198 – 1007)     | 122<br>(62 – 197)   | 111<br>(63 – 187)   | 0.30 | 0.003  |

Values are shown as median (interquartile range)  
\*Active sJIA versus clinical inactive sJIA: p < 0.01

Table 3. Correlation of laboratory parameters of disease activity with levels of IFN $\gamma$ , CXCL9, CXCL10, CXCL11, and IL-6 in patients with MAS and in patients with active sJIA without MAS at sampling.

| MAS                     |                                  | IFN $\gamma$ |       | CXCL9 |       | CXCL10 |       | CXCL11 |       | IL-6  |       |
|-------------------------|----------------------------------|--------------|-------|-------|-------|--------|-------|--------|-------|-------|-------|
|                         |                                  | r*           | p     | r*    | p     | r*     | p     | r*     | p     | r*    | p     |
| Ferritin                | 8000 (3158 – 13174) <sup>1</sup> | 0.57         | 0.014 | 0.49  | 0.041 | 0.66   | 0.002 | 0.62   | 0.023 | 0.17  | >0.1  |
| NEU                     | 6.9 (3.4 – 13.9) <sup>1</sup>    | -0.64        | 0.005 | -0.61 | 0.010 | -0.37  | >0.1  | -0.08  | >0.1  | 0.09  | >0.1  |
| PLT                     | 197 (114 – 392) <sup>1</sup>     | -0.53        | 0.017 | -0.52 | 0.022 | -0.53  | 0.008 | -0.22  | >0.1  | -0.02 | >0.1  |
| ALT                     | 46 (18 – 164) <sup>1</sup>       | 0.49         | 0.045 | 0.49  | 0.044 | 0.51   | 0.038 | 0.06   | >0.1  | -0.44 | 0.089 |
| LDH                     | 1152 (722 – 2135) <sup>1</sup>   | 0.45         | 0.095 | 0.62  | 0.013 | 0.64   | 0.001 | 0.64   | 0.048 | 0.08  | >0.1  |
| Active sJIA without MAS |                                  | r*           | p     | r*    | p     | r*     | p     | r*     | p     | r*    | p     |
| Ferritin                | 214 (37 – 1669) <sup>1</sup>     | -0.27        | >0.1  | 0.28  | >0.1  | 0.27   | >0.1  | 0.29   | >0.1  | -0.12 | >0.1  |
| N                       | 8.4 (5.2 – 14.5) <sup>1</sup>    | 0.39         | >0.1  | 0.40  | 0.061 | 0.32   | >0.1  | 0.40   | 0.067 | 0.28  | >0.1  |
| PLT                     | 444 (353 – 544) <sup>1</sup>     | 0.21         | >0.1  | -0.14 | >0.1  | -0.13  | >0.1  | 0.27   | >0.1  | 0.35  | 0.064 |
| ALT                     | 16 (11 – 24) <sup>1</sup>        | 0.29         | >0.1  | 0.42  | 0.049 | 0.50   | 0.011 | 0.44   | 0.039 | 0.04  | >0.1  |
| LDH                     | 506 (453 – 851) <sup>1</sup>     | 0.07         | >0.1  | 0.49  | >0.1  | 0      | >0.1  | 0.26   | >0.1  | 0     | >0.1  |

NEU=neutrophils count; PLT=platelets count; ALT=alanine aminotransferase; LDH=lactate dehydrogenase;  
1= Median (IQR); r\*=  
Spearman r.  
Correlation of laboratory parameters of disease activity with IFN- $\gamma$ , CXCL9, CXCL10, CXCL11 and IL-6 in patients with MAS and in patients with active sJIA

**[0306]** Correlation of IFN $\gamma$  with the levels of IFN $\gamma$ -induced chemokines in patients with MAS. To further characterize the relationship between IFN $\gamma$  and the three IFN $\gamma$ -induced chemokines in patients with MAS, the correlation of IFN $\gamma$  levels with the levels of each individual chemokine was evaluated. Notably, CXCL9 appears to be primarily and specifically induced by IFN $\gamma$ , while CXCL10 and CXCL11 are also induced by type I interferons. In agreement with this, in patients with active MAS, circulating levels of IFN $\gamma$  were significantly correlated with CXCL9 ( $r=0.693$ ;  $r^2=0.48$ ;  $p=0.001$ ), but had a weaker correlation with CXCL10 levels ( $r=0.535$ ;  $r^2=0.29$ ;  $p=0.015$ ) (Figures 11A-11F). The correlation with CXCL11 levels was also weaker and did not reach statistical significance ( $r=0.447$ ;  $r^2=0.20$ ;  $p=0.08$ ) (not shown).

**[0307]** IFN $\gamma$ -induced chemokines correlate with disease activity in mouse model of MAS. In order to further investigate the association of IFN $\gamma$ -induced chemokine production with MAS, the expression of these chemokines in target tissues (liver and spleen) was investigated in a murine model of MAS. In this model, MAS clinical and laboratory features are induced by mimicking acute infection with the TLR4 agonist lipopolysaccharide (LPS) on a background of high levels of IL-6 in IL-6 transgenic mice (Strippoli et al., Arthritis Rheum 2012). This approach recapitulates what occurs in patients with sJIA: an infection may trigger MAS/HLH in the presence of active disease, which is indeed characterized by high levels of IL-6. Following induction with LPS, high mRNA levels of CXCL9 and CXCL10 were present in liver and spleen in IL-6 transgenic mice. Notably, serum levels of ferritin were significantly correlated with the levels of expression of CXCL9 in spleen and liver and of CXCL10 in liver, showing a relation among IFN $\gamma$ -related upstream events in target tissues, (i.e. CXCL9 and CXCL10 production in liver and spleen) and typical downstream laboratory abnormalities, such as high ferritin levels. All together, the data in patients with MAS and in the MAS murine model point to a clear relation of increased production of IFN $\gamma$  with increased expression of CXCL9 and, to a lesser extent of CXCL10, and laboratory abnormalities of MAS.

**[0308]** Studies in both patients and animal models of p-HLH have demonstrated a central role for IFN $\gamma$  in disease pathogenesis. However, the role of IFN $\gamma$  in sec-HLH, including MAS in the setting of sJIA, has remained unclear. This study demonstrates conclusively that high levels of IFN $\gamma$  and of the IFN $\gamma$ -induced chemokines were present in patients with MAS occurring in sJIA. Additionally, levels of IFN $\gamma$ , of CXCL9 and of CXCL10 strongly correlated with laboratory parameters of MAS

severity. This study found that serum levels of IFN $\gamma$  and of the three IFN $\gamma$ -related chemokines were comparable between patients with active sJIA and patients with clinically inactive disease. This result argues against a pathogenic role of IFN $\gamma$  in sJIA and is, indeed, consistent with a number of observations by other authors. Three gene expression studies have failed to find a prominent IFN $\gamma$ -induced signature in peripheral blood mononuclear cells (PBMCs) of patients with active sJIA without MAS at sampling (Fall, N., et al., *Gene expression profiling of peripheral blood from patients with untreated new-onset systemic juvenile idiopathic arthritis reveals molecular heterogeneity that may predict macrophage activation syndrome*. Arthritis Rheum, 2007. 56(11): p. 3793-804; Ogilvie, E.M., et al., *Specific gene expression profiles in systemic juvenile idiopathic arthritis*. Arthritis Rheum, 2007. 56(6):p. 1954-65; Pascual, V., et al., *Role of interleukin-1 (IL-1) in the pathogenesis of systemic onset juvenile idiopathic arthritis and clinical response to IL-1 blockade*. J Exp Med, 2005. 201(9): p. 1479-86). After *ex vivo* stimulation of PBMCs, the number of cells producing IFN $\gamma$  in patients with active sJIA was similar to that of controls (Lasigle, D., et al., *Role of IL-1 beta in the development of human T(H)17 cells: lesson from NLPR3 mutated patients*. PLoS One, 2011. 6(5): p. e20014). Consistently, patients with both active and inactive SJIA do not exhibit increased serum or synovial fluid levels of IFN $\gamma$  (de Jager, W., et al., *Blood and synovial fluid cytokine signatures in patients with juvenile idiopathic arthritis: a cross-sectional study*. Ann Rheum Dis, 2007. 66(5): p. 589-98). Supporting the absence of a role of IFN $\gamma$  in the joint inflammation of sJIA, CXCL9 and CXCL10 are almost undetectable in the synovial tissues of sJIA patients, while high levels of these chemokines can be found in synovial tissues from patients with oligoarticular or polyarticular JIA (Sikora, K.A., et al., *The limited role of interferon-gamma in systemic juvenile idiopathic arthritis cannot be explained by cellular hyporesponsiveness*. Arthritis Rheum, 2012. 64(11): p. 3799-808). Recent data in mice show that immune stimulation of IFN $\gamma$  knock-out mice with Freund's complete adjuvant produces a systemic inflammatory syndrome that includes features of sJIA, further supporting the limited role of IFN $\gamma$  in sJIA (Avau, A., et al., *Systemic juvenile idiopathic arthritis-like syndrome in mice following stimulation of the immune system with Freund's complete adjuvant: regulation by interferon-gamma*. Arthritis Rheumatol, 2014. 66(5): p. 1340-51).

[0309] In sharp contrast, this study has shown markedly higher levels of IFN $\gamma$  and of IFN $\gamma$ -related chemokines in patients with active MAS at sampling compared with those of patients with active sJIA without MAS at sampling. This was also confirmed in individual patients with serial samples obtained during both active MAS and during active sJIA without MAS. Incidentally, this study did

not find, in patients sampled during MAS, a significant increase in the levels of IL-6 or IL-1 $\beta$  nor any association with laboratory parameters of MAS, suggesting that these cytokines, albeit critically involved in the pathogenic mechanism of sJIA (De Benedetti, F., et al., *Randomized trial of tocilizumab in systemic juvenile idiopathic arthritis*. N Engl J Med, 2012. 367(25): p. 2385-95; Ruperto, N., et al., *Two randomized trials of canakinumab in systemic juvenile idiopathic arthritis*. N Engl J Med, 2012. 367(25): p. 2396-406) may not be crucial in maintaining MAS. This finding of elevated levels of IFN $\gamma$  and of IFN $\gamma$ -related chemokines is consistent with some previous observations. Shimizu et al reported that levels of neopterin, a catabolite of guanosine triphosphate synthesized by human macrophages upon stimulation with IFN $\gamma$ , were higher in patients with MAS during sJIA compared with patients with active sJIA without MAS (Shimizu, M., et al., *Distinct cytokine profiles of systemic-onset juvenile idiopathic arthritis-associated macrophage activation syndrome with particular emphasis on the role of interleukin-18 in its pathogenesis*. Rheumatology (Oxford), 2010. 49(9): p. 1645-53). More recently, Put et al have reported elevated levels of IFN $\gamma$  and CXCL10 in 5 patients with both primary and secondary HLH, 3 of whom had MAS in the course of sJIA (Put, K., et al., *Cytokines in systemic juvenile idiopathic arthritis and haemophagocytic lymphohistiocytosis: tipping the balance between interleukin-18 and interferon-gamma*. Rheumatology (Oxford), 2015). Consistently with these results, 5 patients with active sJIA without MAS at sampling had markedly lower levels of IFN $\gamma$  and CXCL10 (Put et al., Rheumatology 2015).

[0310] Interestingly, this study found that not only were the levels of IFN $\gamma$  and of the IFN $\gamma$ -related chemokines markedly elevated, but also that their levels, particularly those of CXCL9, were strictly correlated with laboratory features of MAS, indicating association with disease severity. Further supporting an association with disease severity, this study found markedly higher levels of IFN $\gamma$  and of CXCL9 and CXCL10 in one patient with severe disease with multiple organ failure and central nervous system involvement with generalized seizures requiring prolonged intensive care unit admission.

[0311] In patients with MAS, of the three IFN $\gamma$ -induced chemokines, CXCL9 was found to have the strongest correlation with IFN $\gamma$  levels. This observation is consistent with the established notion that CXCL9 production appears to be induced specifically and only by IFN $\gamma$  in contrast to the production of CXCL10 and CXCL11 that can also be induced by type I interferons (Groom, J.R. and A.D. Luster, *CXCR3 ligands: redundant, collaborative and antagonistic functions*. Immunol Cell Biol, 2011. 89(2): p. 207-15). This suggests that CXCL9 levels could serve as a sensitive and

specific biomarker for MAS activity. Indeed, using a murine model of MAS that mimics the triggering of MAS by an infectious stimulus on a background of high IL-6 levels (Strippoli et al., Arthritis Rheum 2012), this study has also found that the expression levels of CXCL9 in liver and spleen was significantly correlated to the circulating levels of ferritin. For CXCL10 expression level this correlation was present only for liver, but not for spleen levels. This is also supported by the findings in patients with MAS, where CXCL9 levels were strictly correlated with all laboratory parameters of MAS. Taken together, these observations in humans and in mice show that CXCL9 strongly correlates with MAS features and IFN $\gamma$  production further supporting the hypothesis that excessive production of IFN $\gamma$  plays a major pathogenic role in MAS. These observations are also consistent with the immunohistochemistry data generated by Put et al using serial lymph node biopsies from the same SJIA patient obtained during active sJIA without MAS, as well as during MAS. They report that CXCL10 and indoleamine 2,3-dioxygenase, both IFN $\gamma$  inducible proteins, were detected at high levels by immunohistochemistry in the tissue obtained during MAS, but not in that obtained during active sJIA without MAS (Put et al., Rheumatology 2015).

[0312] These results in MAS and in sec-HLH, together with the observations available in the literature in patients with p-HLH, support the hypothesis that an increase in IFN $\gamma$  and in IFN $\gamma$ -related chemokines, particularly of CXCL9, is a characteristic feature of HLH, independently from the underlying cause. In this respect it is interesting to note that high levels of CXCL9 were detected in a patient with relapsing MAS induced by an NLRC4 gain of function mutation (Canna, S.W., et al., *An activating NLRC4 inflammasome mutation causes autoinflammation with recurrent macrophage activation syndrome*. Nat Genet, 2014. 46(10): p. 1140-6), suggesting that even in the setting of an HLH induced only by inflammasome dysregulation IFN $\gamma$  hyperproduction may be in place.

[0313] Data in animal models of p-HLH, both in perforin and in Rab27a knock-out mice, unequivocally demonstrate the pathogenic role of IFN $\gamma$ . Similarly, recent data in the TLR9 induced model of HLH, a model of HLH secondary to infection, have also shown a major role for increased IFN $\gamma$  production (Behrens, E.M., et al., *Repeated TLR9 stimulation results in macrophage activation syndrome-like disease in mice*. J Clin Invest, 2011. 121(6): p. 2264-77 and (Bautois et al., in progress). Additional studies have recently demonstrated in the above mentioned murine model of MAS that treatment with an anti- IFN $\gamma$  antibody led to increase in survival and reverted clinical and laboratory features of MAS (Prencipe et al., in progress). All together the results of this study and these observations in animals provide the rational for IFN $\gamma$  neutralization as a

therapeutic approach in MAS.

**Example 7. Safety, Tolerability, Pharmacokinetics and Efficacy Assessment of Intravenous Multiple Administrations of Anti-interferon Gamma Anti-IFN $\gamma$  Monoclonal Antibodies in Pediatric Patients with Primary Hemophagocytic Lymphohistiocytosis (HLH)**

[0314] The studies presented herein were designed to determine the safety and tolerability profile of multiple intravenous (IV) administrations of the anti-IFN $\gamma$  antibody referred to herein as NI-0501; to determine NI-0501 efficacy and benefit/risk profile in HLH patients; to describe the pharmacokinetics (PK) profile of NI-0501 in HLH patients; to define an appropriate NI-0501 therapeutic dose regimen for HLH; and to assess the immunogenicity of NI-0501.

[0315] *Preclinical studies:* Previous studies have demonstrate that NI-0501 shows similar binding affinity and blocking activity for IFN $\gamma$  from non-human primate species, including rhesus and cynomolgus monkeys, but not from dogs, cats, pigs, rabbits, rats or mice. Toxicology and safety studies in cynomolgus monkeys demonstrated that there was no off-target toxicity attributed to administration of NI-0501, weekly administrations of NI-0501 were well tolerated and did not require the need for antibiotic prophylaxis, and no abnormal histopathological or behavioral findings were observed during these prior studies.

[0316] Due to NI-0501 capacity to bind free and IFN $\gamma$ R1-bound IFN $\gamma$ , studies were performed to investigate the potential of NI-0501 to mediate ADCC and CDC activities, in the presence of target. A lack of ADCC activity was demonstrated, and no induction of CDC activity was observed.

[0317] *Phase I Clinical Studies:* A Phase 1 randomized double-blinded placebo- controlled single ascending dose study in 20 healthy adult volunteers investigating the safety, tolerability and pharmacokinetic profiles of single intravenous (IV) administrations of NI-0501. During this study, 6 subjects received placebo, while 3, 3, 4, and 4 subjects (in total 14 subjects) received NI-0501 doses of 0.01, 0.1, 1, and 3 mg/kg, respectively.

[0318] The PK analysis of NI-0501 revealed the expected profile for an IgG1 with a long half-life (around 22 days), a slow clearance ( $\leq 0.007$  L/h) and a low volume of distribution ( $< 6$  L on average).

[0319] A total of 41 adverse events (AEs) were observed after start of drug infusion in 14 out of 20 subjects (70%), 10 of which were reported by 4 subjects having received placebo. Thirty-six (87.8%) AEs were of mild intensity and 5 (12.2%) were of moderate intensity. No severe or life-threatening AEs were reported. Twenty-three AEs (56.1%) in 10 of the 14 subjects who experienced an AE were

reported as drug-related (at least with a reasonable possibility). Most AEs were singular occurrences and no trend in relation to increasing NI-0501 dosage was observed. All NI-0501 infusions were uneventful.

[0320] In summary, the infusion of NI-0501 was well tolerated, and the effects observed during the 8 week monitoring after drug infusion did not reveal any serious or unexpected off- target safety or immunogenicity concerns.

[0321] *Phase 2/3 Clinical Study Materials and Methods:* These studies are performed on primary HLH patients. The studies are divided into three parts: screening, treatment, and follow- up. An overview is presented in Figure 12.

[0322] In these studies, suitable patients include patients naïve to HLH treatment (also referred to herein as “first line patients”), or patients who may have already received conventional HLH therapy (also referred to herein as “second line patients”) without having obtained a satisfactory response, *e.g.*, according to the treating physician, or having shown signs of intolerance to it. Patients who receive NI-0501 after having failed conventional HLH therapy or having shown intolerance to it represent the pivotal cohort of the study, to demonstrate the efficacy of NI-0501 as second line treatment of primary HLH. Treatment-naïve patients are enrolled for collection of efficacy and safety data in the first line setting.

[0323] The following patients are excluded from this study: patients who have had a diagnosis of secondary HLH consequent to a proven rheumatic or neoplastic disease; patients who are previously treated with any T-cell depleting agents (such as, for example, anti- thymocyte globulin (ATG), anti-CD52 therapy) during the previous 2 weeks prior to screening or treated with any other biologic drug within 5 times their defined half-life period (with the exception for rituximab in case of documented B-cell EBV infection); patients having active mycobacteria, *Histoplasma Capsulatum*, *Shigella*, *Salmonella*, *Campylobacter* and *Leishmania* infections; patients with evidence of past history of tuberculosis or latent tuberculosis; patients with positive serology for HIV antibodies, hepatitis B surface antigen or hepatitis C antibodies; patients with presence of malignancy; patients with patients who have another concomitant disease or malformation severely affecting the cardiovascular, pulmonary, liver or renal function; patients with history of hypersensitivity or allergy to any component of the study regimen; patients with receipt of a live or attenuated live (including BCG) vaccine within the previous 12 weeks from screening; and/or pregnant or lactating female patients.

[0324] The studies presented herein use the anti-interferon gamma antibody NI-0501, a fully human IgG1 monoclonal antibody (mAb) directed against human IFN $\gamma$ . NI-0501 is provided as a sterile

concentrate for infusion (per mL) as shown below in Table 4.

Table 4. NI-0501 Formulation

| Ingredient                                 | Quantity (per mL) |
|--------------------------------------------|-------------------|
| NI-0501                                    | 5 mg              |
| L-Histidine                                | 1.55 mg           |
| L-Histidine monohydrochloride, monohydrate | 3.14 mg           |
| Sodium chloride (NaCl)                     | 7.31 mg           |
| Polysorbate 80                             | 0.05 mg           |
| pH                                         | 6.0 ± 0.2         |

[0325] In these studies, NI-0501 is administered by IV infusion over a period of one hour at an initial dose of 1 mg/kg. This dose is predicted to inhibit for 3 days at least 99% of IFN $\gamma$  effect in patients with baseline IFN $\gamma$  concentrations lower or equal to 3400 pg/mL. Infusions are performed every 3 days until Study Day 15 (SD15) (infusion #6), and twice per week thereafter. NI-0501 dose increase to 3 mg/kg is possible according to pre-defined criteria guided by clinical and laboratory response in each patient (as described in Table 5 below) at any time during the study. After a minimum of two infusions at 3 mg/kg if, upon re-assessment, the same clinical and laboratory criteria qualifying the patient to receive 3 mg/kg of NI-0501 are found to still apply, the dose of NI-0501 may be increased to 6 mg/kg for up to four infusions, with a regular monitoring of the clinical and laboratory HLH parameters. Based on the evolution of these parameters, the dose of NI-0501 may either i) be decreased back to 3 mg/kg, or ii) remain at 6 mg/kg for additional IV infusions (or be increased above 6 mg/kg), if PK and PD evidence indicates excessively high IFN $\gamma$  production and, consequently, fast NI-0501 elimination. Dose increase may occur any time during the study, if the clinical and laboratory criteria set forth herein are met.

Table 5. Clinical and laboratory criteria to guide dose increase

| Study Day (SD)                | NI-0501 dose                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-------------------------------|----------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| On SD0                        | Starting dose of 1 mg/kg         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| On SD3                        | Increase to 3 mg/kg              | <i>Criteria to be met:</i><br>- if fever persists or reoccurs (when present at baseline)<br><u>or</u><br>if significant worsening of clinical conditions                                                                                                                                                                                                                                                                                                                                                             |
| From SD6 onwards <sup>a</sup> | Increase to 3 mg/kg <sup>b</sup> | <i>Criteria to be met:</i><br>- if no satisfactory improvement in clinical conditions<br><u>and</u><br>at least 1 of the following<br><i>Platelet counts</i> (x10 <sup>3</sup> /mcl)<br>If<br>bsl. counts < 50 → no improvement to > 50<br>bsl. counts 50-100 → less than 30% improvement bsl.<br>counts > 100 → any decrease to < 100<br><i>ANC</i> (count/mcl)<br>If<br>bsl. counts < 500 → no improvement to > 500<br>bsl. counts 500-1000 → any decrease to < 500<br>bsl. counts > 1000 → any decrease to < 1000 |

|                                       |                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|---------------------------------------|----------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                       |                                  | <p><i>Ferritin</i> (ng/ml)</p> <p>If</p> <p>bsl. levels <math>\geq 3000 \rightarrow</math> no improvement (<math>&lt; 20\%</math> decrease)</p> <p>bsl. levels <math>&lt; 3000 \rightarrow</math> any decrease to <math>&gt; 3000</math></p> <p><i>Splenomegaly</i> <math>\rightarrow</math> worsening (at clinical or US examination)</p> <p><i>Coagulopathy</i> (both D-Dimer and Fibrinogen have to apply)</p> <p>D-Dimer</p> <p>If abnormal at bsl. <math>\rightarrow</math> no improvement</p> <p>Fibrinogen</p> <p>If</p> <p>bsl. levels <math>\leq 100 \rightarrow</math> no improvement</p> <p>bsl. levels <math>\geq 100 \rightarrow</math> any decrease to <math>&lt; 100</math></p> |
| From SD9 or SD12 onwards <sup>c</sup> | Increase to 3 mg/kg <sup>d</sup> | <p><i>Criteria to be met:</i></p> <ul style="list-style-type: none"> <li>- in case, after a minimum of two infusions at 3 mg/kg, the criteria above reported have been reassessed and found to be still met</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |

<sup>a</sup> NI-0501 dose has to be increased from 1 to 3 mg/kg if these criteria apply after SD6.

<sup>b</sup> If NI-0501 dose has been already increased on SD3, at least two infusions at the dose of 3 mg/kg have to be performed before criteria re-assessment.

<sup>c</sup> Depending on whether dose increase to 3 mg/kg has occurred on SD3 or SD6.

<sup>d</sup> For a maximum of four infusions.

Abbreviations: bsl. = baseline; ANC = absolute neutrophil count; US = ultrasound

**[0326]** In these studies, NI-0501 is administered for 8 weeks, and the treatment period will be divided in 2 separate periods: Treatment Period 1 and 2 as shown in Figure 12.

**[0327]** After NI-0501 is administered for 8 weeks, the conditioning regimen in preparation for Hematopoietic Stem Cell Transplantation (HSCT) may be initiated. The anticipated duration of treatment can be shortened, although not to less than 4 weeks, if the patient's condition and donor availability allow the performance of a transplant. In the event that an appropriate donor has not been identified by Week 8 or in case of the need to delay the schedule for transplantation for reasons unrelated to the administration of NI-0501, NI-0501 treatment can be continued in the context of a long-term follow-up study, provided that

a favorable benefit/risk has been established for the patient.

[0328] In these studies, NI-0501 is administered on a background of dexamethasone, which can be tapered depending on patient condition. In treatment-naïve patients, NI-0501 will be administered on a background of 10 mg/m<sup>2</sup> of dexamethasone. In patient receiving NI-0501 as second line HLH treatment, dexamethasone has to be administered at the dose of at least 5mg/m<sup>2</sup>, or at the same dose administered prior to screening if higher. Patients are required to have received dexamethasone from SD-1.

[0329] Dexamethasone can be tapered depending on patient condition, according to the judgment of the treating physician. The tapering scheme can be selected by the treating physician, provided that the dexamethasone dose, at each step, is not more than halved and frequency of change is not more than weekly.

[0330] In the event of disease worsening after tapering of dexamethasone, the dose of dexamethasone can be increased and maintained until a satisfactory response is achieved according to the treating physician.

[0331] As recommended in HLH treatment guidelines, patients receive prophylactic treatment for *Pneumocystis jiroveci*, fungal and *Herpes Zoster* virus infection from the day before initiation of NI-0501 treatment until the end of the study. Patients receive prophylactic treatments starting from the day prior to initiation of NI-0501 treatment (i.e. SD-1) until the end of the study. For example, for *Pneumocystis jiroveci* prevention, patients may receive, e.g. 750 mg/m<sup>2</sup>/day sulfamethoxazole with 150 mg/m<sup>2</sup>/day trimethoprim given orally in equally divided doses twice a day, on 3 consecutive days per week. For fungal infection prevention, patients may receive, e.g. Fluconazole 12 mg/kg daily with a maximum of 400 mg daily dose. For HZ virus prevention, patients may receive, e.g. Acyclovir 200 mg four times daily for children over two years, for children under two years 100 mg four times daily. These treatments will be given orally, whenever possible, otherwise intravenously.

[0332] Patients can also receive any of a variety of concomitant therapies, such as, for example, Cyclosporin A, intrathecal methotrexate and glucocorticosteroids, and others. Cyclosporin A (CsA) can be continued if already being administered to the patient prior to screening. CsA can be withdrawn at any time. CsA is not to be introduced de novo during the course of the study once NI-0501 administration has started.

[0333] If the patient is receiving intrathecal methotrexate and glucocorticoids at the time of NI-0501

treatment initiation, this treatment will be continued as required. If the appearance of CNS symptoms occurs before the initiation of NI-0501 treatment, therapy with intrathecal methotrexate and glucocorticoids must be initiated prior to the first administration of NI-0501.

**[0334]** IV immunoglobulins (IVIG) are only allowed as replacement treatment in case of a documented immunoglobulin deficiency. For example, in case of a documented immunoglobulin deficiency justifying replacement, IVIG can be given at a dose of 0.5 g/kg, every 4 weeks or more frequently in order to maintain adequate IgG levels. Any infusion within the previous 4 weeks prior to screening, as well as any infusion during NI-0501 treatment is acceptable.

**[0335]** Analgesic treatment, transfusion of blood products, electrolyte and glucose infusions, antibiotics, anti-fungal and anti-viral treatment and general supportive care are allowed. Additional HLH treatments may be allowed in case of unsustained or limited HLH improvement once the maximum NI-0501 dose level is achieved. As used herein, unsustained HLH improvement refers to patients who are unable to maintain at least 50% improvement from baseline for 3 HLH parameters (see Table 6 below). At least two consecutive measurements must document the loss of HLH improvement. As used herein, limited HLH improvement refers to less than 50% change from baseline in a minimum of 3 HLH clinical and laboratory criteria. Etoposide should be administered as additional HLH treatment, unless clear evidence of lack of response or intolerance to the drug is derived from previous medical history.

**[0336]** The following therapies may not be used concomitantly with NI-0501 administration: etoposide, T-cell depleting agents, or any other biologic drug is generally not allowed, except for the following: G-CSF, in case of prolonged neutropenia; Rituximab, in case of documented B-cell EBV infection; and additional HLH treatments, in case of unsustained or limited HLH improvement (as defined herein) at the maximum NI-0501 dose level. Etoposide should be administered, unless a clear evidence of lack of response or intolerance to the drug is derived from previous medical history. Vaccination with a live or attenuated (including BCG) vaccine must be avoided during the whole study including the 4 week follow-up period. In the event that NI-0501 concentrations remain at therapeutic levels after the end of the study, the period with no vaccinations should be extended until measurable concentration of NI-0501 are no longer detectable.

**[0337]** Evolution of clinical signs (fever, splenomegaly, CNS symptoms) and laboratory parameters (CBC, fibrinogen, ferritin, sCD25 levels), which characterize the disease, are used to assess the achievement of response and time to response. The primary efficacy endpoint includes overall

response rate, i.e. achievement of either Complete or Partial Response or HLH Improvement, at End of Treatment (EoT), as defined in Table 6 below. The secondary efficacy endpoints include time to response any time during the study; durability of response, i.e., maintenance of response achieved any time during the study until EoT and beyond (including data collected in any long-term follow-up study); number of patients able to reduce glucocorticoids by 50% or more of baseline dose; number of patients able to proceed to HSCT, when deemed indicated; survival at Week 8 (or EoT) and at the end of the study; serum concentration of NI-0501 to determine NI-0501 pharmacokinetic (PK) profile; determination of pharmacodynamic (PD) effects, including levels of circulating total IFN $\gamma$  and markers of its neutralization, namely CXCL9 and CXCL10; and determination of other biomarkers, e.g. sCD25, IL-10.

Table 6. Definition of response

| <b>Overall Response Rate</b> |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Complete Response</b>     | <p>Complete Response is adjudicated if:</p> <p>No fever = body temperature &lt; 37.5°C</p> <p>Normal spleen size as measured by 3D abdominal ultrasound</p> <p>No cytopenia = Absolute Neutrophil Counts <math>\geq 1.0 \times 10^9/L</math> and platelet count <math>\geq 100 \times 10^9/L</math> [absence of G-CSF and transfusion support must be documented for at least 4 days to report no cytopenia]</p> <p>No hyperferritinemia = serum level is &lt; 2000 <math>\mu g/L</math></p> <p>No evidence of coagulopathy, i.e., normal D-Dimer and/or normal (&gt; 150 mg/dL) fibrinogen levels</p> <p>No neurological and CSF abnormalities attributed to HLH</p> <p>No sustained worsening of sCD25 (as indicated by at least two consecutive measurements that are &gt; 2-fold higher than baseline)</p> |
| <b>Partial Response</b>      | Partial Response is adjudicated if:                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | <p>At least 3 of the HLH clinical and laboratory abnormalities (including CNS abnormalities) meet the above mentioned criteria for “Complete Response”. In the case of “reactivated patients” who enter the study with 3 abnormal HLH features, at least 2 criteria should meet the definition given</p> <p>There is no progression of other aspects of HLH disease pathology (e.g., jaundice, liver size, oedema, CNS clinical alterations)</p> |
| <b>HLH Improvement</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | <p>Improvement (&gt;50% change from baseline) of at least 3 HLH clinical and laboratory abnormalities (including CNS involvement). In the case of “reactivated patients” who enter the study with 2 abnormal HLH features, a change from baseline greater than 50% for both will define HLH as improved.</p>                                                                                                                                     |
| <b>Limited Improvement/Lack of Improvement/No Response</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <p>Less than 50% change from baseline of 3 or more of the above mentioned HLH clinical and laboratory abnormalities [in the case of “reactivated patients” who enter the study with 2 abnormal HLH features, less than 50% change from baseline in both will be sufficient to define limited improvement]</p> <p><u>and</u></p> <p>No apparent improvement in other aspects of disease pathology</p>                                                                                                                                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Reactivation</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <p>Deterioration of two or more HLH d clinical and laboratory criteria with the following specifications:</p> <p>numerical laboratory values* must become abnormal and worsen by more than 30% compared to the previous evaluation, on two sequential assessments performed with an interval of minimum 1 day and maximum 1 week</p> <p>deterioration of clinical criteria must be confirmed by consistent observations of worsening over three consecutive days</p> <p>The development of new or recurrent CNS symptoms counts as a single criterion for reactivation.</p> <p><i>* The following laboratory parameters are specifically considered for determination of reactivation:</i></p> |                                                                                                                                                                                                                                                                                                                                                                                                                                                  |

- *platelets*
- *neutrophils*
- *fibrinogen*
- *ferritin*
- *soluble CD25 (sCD25; i.e. soluble IL-2 receptor).*

*The assessment of NK function, red blood cells/hemoglobin and triglyceride levels cannot be considered for the determination of reactivation.*

**[0338]** Safety parameters to be collected and assessed include incidence, severity, causality and outcomes of Adverse Events (AEs) (serious and non-serious), with particular attention being paid to infections; evolution of laboratory parameters such as complete blood cell count (CBC), with a focus on red cells (hemoglobin), neutrophils and platelets, liver tests, renal function tests and coagulation; number of patients withdrawn for safety reasons; and other parameters, such as the level (if any) of circulating antibodies against NI-0501 to determine immunogenicity (ADA).

**[0339]** The primary endpoint (Overall Response Rate) is evaluated using the exact binomial test at the one-sided 0.025 level. Time to Response, durability of Response and Survival time are presented using Kaplan-Meier curves with medians calculated if available. 95% confidence intervals are calculated for the median for each of these endpoints. Additional endpoints based on binary outcomes including number of patients who reduce glucocorticoids by 50% or more, and number of patients able to proceed to HSCT will be converted to proportions and associated 95% confidence intervals calculated. Statistical significance in terms of p-values are only obtained for the primary endpoint. All other endpoints are viewed as supportive for the primary endpoint and as a consequence no formal hierarchy of endpoints is declared.

**[0340]** Administration of NI-0501 in patients lead to rapid normalization of fever within hours after the first infusion of NI-0501. Figures 13A and 13B depict the effect of NI-0501 infusion on body temperature in two patients having body temperature > 37.5 °C at initiation of NI-0501 treatment. Figure 14 is a series of graphs and a table depicting the effect of NI-0501 administration on neutrophil count in patients. Figure 15 is a series of graphs and a table depicting the effect of NI-0501 administration on platelet count in patients. Figure 16 is a series of graphs and a table depicting the effect of NI-0501 administration on serum levels of ferritin in patients.

Figure 17 is a series of graphs and a table depicting the effect of NI-0501 administration on glucocorticoid tapering in patients. Figure 18 is a graph depicting that administration of NI-0510 maintained IFN $\gamma$  neutralization until the time of HSCT. HLH response to NI-0501 treatment also persisted until transplantation. Patients were also evaluated for any CNS involvement following NI-0501 administration. A summary of the baseline CNS involvement and the status by the end of treatment (EOT) is shown below in Table 11.

Table 11. Response to NI-0501 treatment - CNS involvement

|                      | Baseline (SD0)                                               | EOT                        |
|----------------------|--------------------------------------------------------------|----------------------------|
| Pt. #6               | Obtundation; Hemiparesis<br>Loss of developmental milestones | Resolved<br>Fully Regained |
|                      | Elevated protein and neopterin in CSF; Pleocytosis           | Resolved                   |
|                      | Abnormal enhancement on MRI                                  | Improved                   |
| Pt. #3               | Elevated protein in CSF; Pleocytosis                         | Resolved                   |
| Pt. #15              | Loss of walking ability                                      | Regained                   |
|                      | Elevated protein and neopterin in CSF; Pleocytosis           | Resolved                   |
| Pt. #18*             | Pleocytosis; Elevated protein in CSF                         | Improved                   |
|                      | Axial hypotonia                                              | Improved                   |
| Pt. #20 <sup>§</sup> | Elevated protein and neopterin in CSF; Pleocytosis           | Initial improvement        |
|                      | 6 <sup>th</sup> nerve palsy; clonus at ankles                | Initial improvement        |
| Pt. #4 <sup>^</sup>  | Elevated protein in CSF; Pleocytosis                         | Not evaluable              |
|                      | Abnormal infiltrates at MRI                                  | Not evaluable              |

Note: patients received IT therapy, except pt. #4 in whom regular medicated LP was not performed

\* Treatment ongoing

<sup>§</sup> Treatment started since 2 weeks

<sup>^</sup> Control at EOT was not performed

[0341] Of ten patients who have undergone hematopoietic stem cell transplantation (HSCT), all patients engrafted; in 1 patient CD34 stem cell boost was required due to mixed chimerism on D+145 post-HSCT. Secondary graft failure occurred in 1 patient, followed by HLH reactivation. This patient died at D+68 post-HSCT due to acute respiratory failure and bacterial infections. Another patient died at D+47 post-HSCT (septic shock in a context of severe GvHD). Mild GvHD was reported in other 3 patients and resolved/is resolving.

[0342] Neutralizing serum concentrations of NI-0501 at the time of HSCT were measured in 8 of the 10 patients undergoing transplant, as reflected by levels of CXCL9 (a

chemokine exquisitely induced by IFN $\gamma$ ) below the limit of quantification. Thus, these data show that NI-0501 can spare short- or long-term toxicities reported for etoposide-based regimens. This translates into a reduced risk of allo-HSCT-related complications.

[0343] These data demonstrate that NI-0501 treatment improves and/or can resolve relevant clinical and laboratory abnormalities of HLH, including CNS signs and symptoms. Response to NI-0501 is independent of the presence and the type of causative mutations and/or the presence and the type of an infectious trigger. NI-0501 was well tolerated. No safety concerns emerged to date (e.g., no myelotoxicity, no broad immunosuppression). No infections caused by pathogens known to be promoted by IFN $\gamma$  neutralization have been observed. The neutralization of IFN $\gamma$  by NI-0501 can offer an innovative and targeted approach to the management of HLH.

**Example 8. Safety, tolerability, pharmacokinetics and efficacy of short term intravenous administrations of NI-0501, an anti-interferon gamma (anti-IFN $\gamma$ ) monoclonal antibody, in patients with systemic Juvenile Idiopathic Arthritis (sJIA) developing Macrophage Activation Syndrome/secondary HLH (MAS/sHLH)**

[0344] The studies provided herein are designed to demonstrate the efficacy and safety of NI-0501 for the treatment of MAS/sHLH in patients with sJIA, divided in two parts (i) a pilot study to evaluate the PK profile of NI-0501 and the dosing strategy, and to preliminarily assess the NI-0501 benefit/risk in this patient population; and (ii) a pivotal study to demonstrate the efficacy and safety of NI-0501 (study to be continued upon confirmation of the dosing regimen and the positive benefit/risk profile of NI-0501). An overview of this study design is shown in Figure 19.

[0345] The main objectives of the pilot study are: (i) to define an appropriate NI-0501 therapeutic dose regimen for sJIA patients with MAS/sHLH; (ii) to assess the benefit/risk profile of NI-0501 in sJIA patients with MAS/sHLH; and (iii) to describe the pharmacokinetics (PK) profile of NI-0501 in sJIA patients with MAS/sHLH. The main objectives of the pivotal study are: (i) to determine NI-0501 efficacy in sJIA patients with MAS/sHLH; (ii) to evaluate the safety and tolerability profile of short-term intravenous (i.v.) administrations of NI-0501 in sJIA patients with MAS/sHLH; (iii) to confirm the positive benefit/risk profile of NI-0501 in sJIA patients with MAS/sHLH; (iv) to perform an exploratory evaluation of chemokines CXCL9 and CXCL10 as MAS/sHLH diagnostic biomarkers and as predictors of response to NI-0501 treatment; and (v) to assess the immunogenicity of NI-0501 in sJIA patients with MAS/sHLH.

**[0346]** The study population includes sJIA patients with MAS/sHLH having shown inadequate response to high dose glucocorticoid treatment. The inclusion criteria include the following: (i) gender: male and female; (ii) age: < 16 years at the time of sJIA diagnosis; (iii) diagnosis of active MAS/sHLH confirmed by the treating rheumatologist, in the presence of at least 2 of the following laboratory and clinical criteria<sup>1</sup>: (a) laboratory criteria: platelet count  $\leq 262 \times 10^9/L$ , WBC count  $\leq 4.0 \times 10^9/L$ , AST levels  $> 59 U/L$ , and/or fibrinogen levels  $\leq 2.5 g/L$ ; (b) clinical criteria: hepatomegaly, haemorrhagic manifestations, and/or CNS dysfunction; (iv) patient presenting an inadequate response to high dose i.v. glucocorticoid treatment for at least 3 days (including but not limited to pulses of 30 mg/kg mPDN on 3 consecutive days), as per local standard of care; (v) high dose i.v. glucocorticoid should not be lower than 2 mg/Kg/day of mPDN equivalent in 2 separate daily doses up to 60 mg/day. In case of rapid worsening of the patient's condition and/or lab parameters, inclusion may occur within less than 3 days from starting high dose i.v. glucocorticoids; (vi) patient consent (or consent of legally authorized representative(s)); and (vii) having accepted contraceptive measures when the patient is post-pubescent.

**[0347]** Exclusion criteria include: (i) diagnosis of suspected or confirmed primary HLH or HLH consequent to a neoplastic disease; (ii) patients treated with: Anakinra, Tocilizumab, Canakinumab, TNF inhibitors, rituximab or any other biologic drug within 5 times of their defined half-life; (iii) active mycobacteria (typical and atypical), Histoplasma Capsulatum, Shigella, Salmonella, Campylobacter and Leishmania infections; (iv) evidence of latent tuberculosis; (v) positive serology for HIV antibodies; (vi) presence of malignancy; (vii) patients who have another concomitant disease or malformation severely affecting the cardiovascular, pulmonary, CNS, liver or renal function that in the opinion of the Investigator may significantly affect likelihood to respond to treatment and/or assessment of NI-0501 safety; (viii) history of hypersensitivity or allergy to any component of the study regimen; (ix) receipt of a BCG vaccine within 12 weeks prior to screening; (x) receipt of other live or attenuated live vaccines within 6 weeks prior to screening; and/or (xi) pregnant or lactating female patients.

**[0348]** *Dosing Regimen, Frequency of Administration & Treatment Duration:* In these studies, NI-0501 is used in the formulation shown in Example 7. In Part 1, NI-0501 is administered at the initial dose of 6 mg/kg by infusion over a period of one hour on SD0. NI-0501 treatment is continued at the dose of 3 mg/kg every 3 days for 4 weeks (i.e. up to SD27). NI-0501 treatment may be shortened upon achievement of complete clinical response (i.e. MAS remission). After 4

weeks, NI-0501 treatment may be continued for up to additional 4 weeks (i.e. up to SD56) as maintenance as needed until MAS remission is achieved, with the possibility of decreasing the dose to 1 mg/kg and elongating the interval between infusion to weekly administration. If the PK profile shows an unanticipated TMDD (thus signaling an exceptionally high IFN $\gamma$  production) the dose of NI-0501 may be increased to 10 mg/kg guided by clinical and PK evidence. This dose increase is approved only upon careful assessment of the benefit/risk profile in that individual patient.

[0349] In Part 2, upon confirmation that the proposed dosing regimen is appropriate, and demonstration of the positive benefit/risk of NI-0501, the study will be continued. Minor modifications of the dosing regimen might be applied, if required based on evidence obtained in Part 1.

[0350] *Background Therapy & Concomitant Medication:* NI-0501 is administered on a background of at least 2 mg/kg of methylprednisolone (mPDN) equivalent up to 60 mg/day (in patients of 30 kg or more), which can be tapered during the treatment depending on patient conditions. Patients receive prophylactic treatment for Herpes Zoster infections starting preferably the day before (and in any case prior to initiation of NI-0501 treatment) until serum NI-0501 levels are no longer detectable. Cyclosporine A (CsA) can be continued if started at least 3 days prior to initiation of NI-0501 treatment. CsA dose adjustments are allowed in order to maintain therapeutic levels. CsA can be withdrawn at any time during the study, upon judgment of the Investigator. CsA cannot be introduced de novo once NI-0501 administration has started. If the patient is receiving intrathecal methotrexate and glucocorticoids at the time of NI-0501 treatment initiation, this treatment may be continued as required. Vaccination with a live or attenuated (including BCG) vaccine must be avoided during the whole study and, in any case, until serum NI-0501 levels are no longer detectable. Analgesic treatment, transfusion of blood products, electrolyte and glucose infusions, antibiotics, anti-fungal and anti-viral treatments, and general supportive care are allowed.

[0351] *Sample size:* In Part 1, at least 5 evaluable patients will be enrolled. In Part 2, upon continuation of the study, at least 10 evaluable patients will be enrolled to achieve a total of 15 evaluable patients. The sample size of 15 has not been formally justified given the rare orphan nature of the disease and the lack of any approved treatment. Nonetheless, based on the assumption that at least 50% of patients inadequately respond to systemic glucocorticoids alone, i.e. 50% of patients on glucocorticoids achieve MAS remission by Week 8 after start of

treatment, this study will have 70% power to detect an improvement from 50% to 77% using a one-sided significance level of 5%.

**[0352]** Study Duration and Study End Definition: The duration of the study will be 8 weeks for each patient (plus up to 1 week screening period). End of the study is defined as last patient last visit. All patients who have received at least one dose of NI-0501 will be asked to enter the NI-0501-05 study for a long-term follow-up.

**[0353]** *Study endpoints:* In the Part 1 (pilot) of the study, the following are assessed to confirm the dosing regimen in this patient population: (i) benefit/risk profile of NI-0501; (ii) PK profile of NI-0501; (iii) levels of chemokines known to be induced by IFN $\gamma$  (e.g., CXCL9, CXCL10, CXCL11); (iv) evolution of MAS distinct features of cytopenia, liver dysfunction and coagulopathy at 2, 4, 6 and 8 weeks after NI-0501 initiation; and (v) dose and duration of NI-0501 treatment. In the Part 2 (pivotal) of the study, the efficacy study endpoints are as follows: (a): primary efficacy endpoint: number of patients achieving MAS remission by Week 8 after initiation of NI-0501 treatment; and (b) secondary efficacy endpoints: time to MAS remission; time to initial response, according to the Investigator's assessment; number of patients for whom at any time during the study glucocorticoids can be tapered to the same (or lower) dose being administered before the occurrence of MAS; time to achievement of glucocorticoids tapering; survival at the end of the study; and number of patients withdrawn from the study due to lack of efficacy. In the Part 2 (pivotal) of the study, the safety study endpoints are as follows: (a) incidence, severity, causality and outcomes of AEs (serious and non-serious), with particular attention being paid to infections; evolution of laboratory parameters, in particular CBC (with a focus on hemoglobin, neutrophils and platelets), LFTs, and coagulation parameter; number of patients withdrawn from the study due to safety reasons; and levels (if any) of circulating antibodies against NI-0501 to determine immunogenicity (ADA).

**[0354]** Pharmacokinetics and Pharmacodynamics are evaluated by the PK profile of NI-0501; levels of circulating free IFN $\gamma$  at predose, and total IFN $\gamma$  (free IFN $\gamma$ +bound to NI-0501) after initiation of NI-0501; levels of chemokines known to be induced by IFN $\gamma$  (e.g., CXCL9, CXCL10, CXCL11); correlation between chemokine levels (CXCL9, CXCL10) and levels of free NI-0501, free IFN $\gamma$  (pre-dose) and total IFN $\gamma$ ; correlation of chemokine and total IFN $\gamma$  levels, and laboratory parameters of MAS severity, e.g. ferritin, platelet count, LFTs (exploratory analysis); and levels of other potential disease biomarkers (e.g., sCD25, IL-10, IL-6, IL-18, TNF $\alpha$ , neopterin).

### **Other Embodiments**

**[0355]** While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

**What is claimed is:**

1. A multiple variable dose method for the treatment of primary hemophagocytic lymphohistiocytosis (HLH) in a human subject in need thereof, comprising intravenously administering to a subject:
  - a) a first dose of 1.0 or 3.0 mg/kg of the subject's body weight an antibody that binds interferon gamma (IFN $\gamma$ ) administered to the subject within 12 hrs; and
  - b) a second dose of 3.0, 6.0 or 10.0 mg/kg of the subject's body weight the antibody within 12 hrs,
 wherein the antibody comprises: a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).
2. The method of claim 1, wherein the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.
3. The method of claim 1, wherein the dose of the antibody is administered within 6 hrs.
4. The method of claim 3, wherein the dose of the antibody is administered within 1 hr.
5. The method of claim 1, wherein the second dose is administered for a first treatment period of every three days after the first dose.
6. The method of claim 5, wherein the second dose is administered for a second treatment

period after completion of the first treatment period 1, 2, 3, 4, 5, 6, 7, 8, weeks after the first dose.

7. The method of claim 6, wherein the second treatment period is twice weekly.
8. The method of claim 1, further comprising administering a third dose of 1.0 mg/kg of the subject's body weight the antibody administered to the subject within 12 hrs
9. The method of claim 1, wherein the antibody dose is administered as a single injection.
10. The method of claim 1, wherein the antibody is administered as a monotherapy or a co-therapy.
11. The method of claim 1, wherein the subject is an adult subject or a pediatric subject.
12. A multiple variable dose method for the treatment of secondary hemophagocytic lymphohistiocytosis (HLH) in a human pediatric subject in need thereof, comprising intravenously administering to a subject:
  - a) a first dose of 6.0 mg/kg of the subject's body weight an antibody that binds interferon gamma (IFN $\gamma$ ) administered to the subject within 12 hrs; and
  - b) a second dose of 3.0 mg/kg of the subject's body weight the antibody within 12 hrs, wherein the antibody comprises: a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

13. The method of claim 12, wherein the dose of the antibody is administered within 6 hrs.
14. The method of claim 13, wherein the dose of the antibody is administered within 1 hr.
15. The method of claim 12, wherein the second dose is administered for a first treatment period of every three days after the first dose.
16. The method of claim 15, wherein the second dose is administered for a second treatment period after completion of the first treatment period.
17. The method of claim 16, wherein the second treatment period is twice weekly.
18. The method of claim 12, further comprising administering after the second dose an additional dose of 6.0 mg/kg of the subject's body weight the antibody administered within 12 hrs.
19. The method of claim 12, wherein the antibody dose is administered as a single injection.
20. The method of claim 12, wherein the antibody is administered as a monotherapy or a co-therapy.
21. A multiple variable dose method for the treatment of secondary hemophagocytic lymphohistiocytosis (HLH) in a human adult subject in need thereof, comprising intravenously administering to a subject:
  - a) a first dose of 3 mg/kg or 6.0 mg/kg of the subject's body weight an antibody that binds interferon gamma (IFN $\gamma$ ) administered to the subject within 12 hrs; and
  - b) a second dose of no more than 10.0 mg/kg of the subject's body weight the antibody within 12 hrs,
 wherein the antibody comprises: a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of

DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

22. The method of claim 21, wherein the dose of the antibody is administered within 6 hrs.

23. The method of claim 22, wherein the dose of the antibody is administered within 1 hr.

24. The method of claim 22, wherein the second dose is administered for a first treatment period of every three days after the first dose.

25. The method of claim 24, wherein the second dose is administered for a second treatment period after completion of the first treatment period.

26. The method of claim 25, wherein the second treatment period is twice weekly.

27. The method of claim 21, further comprising administering after the second dose an additional dose of 1.0, 3.0, or 6.0 mg/kg of the subject's body weight the antibody administered within 12 hrs.

28. The method of claim 21, wherein the antibody dose is administered as a single injection.

29. The method of claim 21, wherein the antibody is administered as a monotherapy or a co-therapy.

30. A multiple variable dose method for the treatment of a condition in a human subject in need thereof, comprising intravenously administering to a subject:

a) a first dose of an antibody that binds interferon gamma (IFN $\gamma$ ) administered to the subject within 12 hrs; and

b) a second dose of the subject's body weight the antibody within 12 hrs,  
 wherein the antibody comprises: a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

31. The method of claim 30, wherein the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

32. The method of claim 30, wherein the dose of the antibody is administered within 6 hrs.

33. The method of claim 32, wherein the dose of the antibody is administered within 1 hr.

34. The method of claim 30, wherein the second dose is administered for a first treatment period of every three days after the first dose.

35. The method of claim 34, wherein the second dose is administered for a second treatment period after completion of the first treatment period.

36. The method of claim 35, wherein the second treatment period is twice weekly.

37. The method of claim 30, wherein the first dose is between 1.0 to 10 mg/kg of the subject's body weight.

38. The method of claim 30, wherein the dose is between 1.0 to 10 mg/kg of the subject's body weight, wherein the second dose is lower or higher than the first dose.
39. The method of claim 30, further comprising administering a third dose of between 1.0 to 10 mg/kg of the subject's body weight the antibody administered to the subject within 12 hrs
40. The method of claim 30, wherein the antibody dose is administered as a single injection.
41. The method of claim 30, wherein the antibody is administered as a monotherapy or a co-therapy.
42. The method of claim 30, wherein the subject is an adult subject or a pediatric subject.
43. The method of claim 30, wherein the condition is transplant rejection
44. The method of claim 43, wherein the transplant rejection is solid organ transplant disorder, bone marrow acute graft rejection.
45. The method of claim 30, wherein the condition is graft vs. host disease, paraneoplastic cerebellar degeneration, hemorrhagic fever, sarcoidosis, adult onset Still's disease
46. The method of claim 30, wherein the method is administered to a subject after receiving CART cell therapy.
47. The method of anyone of the proceeding claims, wherein the subject has been administered dexamethasone immediately prior to the dosing of the antibody.
48. The method of claim 30, wherein the dexamethasone is administered at a dose of at least 10 mg/m<sup>2</sup>.
49. The method of claim 30, wherein subject he dexamethasone is administered at a dose of at least 5 mg/m<sup>2</sup>.

50. The method of claim anyone of the proceeding claims, wherein the subject has not previously been treated for HLH.

51. The method of claim anyone of the proceeding claims , wherein the method further comprises administering at least a second agent to the subject.

52. The method of claim 523, wherein the second agent is a therapeutic agent, an anti-inflammatory agent, and/or an immunosuppressive agent.

53. An injectable pharmaceutical formulation comprising per mL:

- a) 5 mg or 25 mg of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody; and
- b) 1.55 mg L-histidine, 3.14 mg L-histidine monohydrochloride, monohydrate, 7.31 mg sodium chloride (NaCl), and 0.05 mg Polysorbate 80, where the pH is between 5.8 and 6.2.

54. The formulation of claim 53, wherein the antibody comprises a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

55. The formulation of claim 54, wherein the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

56. A unit dose vial comprising, 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ )

monoclonal antibody solution suitable for injection wherein the concentration of antibody is 5 mg/ml or 25 mg/ml, wherein the pH of the solution is between 5.8 and 6.2.

57 The unit dose vial of claim 56, wherein the antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate.

58. The unit dose vial of claim 56, wherein the antibody comprises a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

59. The unit dose vial of claim 58, wherein the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

60. A unit dose vial comprising, 10 ml or 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution suitable for injection wherein the concentration of antibody is at 25 mg/ml, wherein the pH of the solution is between 5.8 and 6.2.

61. The unit dose vial of claim 60, wherein the antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate.

62. The unit dose vial of claim 60, wherein the antibody comprises a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH

CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

63. The unit dose vial of claim 62, wherein the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

64. A unit dose vial comprising, 2 ml or 20 ml of a fully human anti-interferon gamma (IFN $\gamma$ ) monoclonal antibody solution suitable for injection wherein the concentration of antibody is at 5 mg/ml, wherein the pH of the solution is between 5.8 and 6.2.

65. The unit dose vial of claim 64, wherein the antibody is solubilized in the solution such that the solution is clear, colorless, and without precipitate.

66. The unit dose vial of claim 64, wherein the antibody comprises a variable heavy chain complementarity determining region 1 (VH CDR1) comprising the amino acid sequence of SYAMS (SEQ ID NO: 1); a variable heavy chain complementarity determining region 2 (VH CDR2) comprising the amino acid sequence of AISGSGGSTYYADSVKG (SEQ ID NO: 2); and a variable heavy chain complementarity determining region 3 (VH CDR3) comprising the amino acid sequence of DGSSGWYVPHWFDP (SEQ ID NO: 3); a variable light chain complementarity determining region 1 (VL CDR1) comprising the amino acid sequence of TRSSGSIASNYVQ (SEQ ID NO: 4); a variable light chain complementarity determining region 2 (VL CDR2) region comprising the amino acid sequence of EDNQRPS (SEQ ID NO: 5); and a variable light chain complementarity determining region 3 (VL CDR3) region comprising the amino acid sequence of QSYDGSNRWM (SEQ ID NO: 6).

67. The unit dose vial of claim 66, wherein the antibody comprises the heavy chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 47, and the light chain variable amino acid sequence to the amino acid sequence of SEQ ID NO: 48.

FIGURE 1




FIGURE 2






FIGURE 3A



Number of XY Pairs 19  
Spearman r 0,6930  
95% confidence interval 0.3356 - 0.8760  
P value (two-tailed) 0,0010

FIGURE 3B



Number of XY Pairs 24  
Spearman r 0,4469  
95% confidence interval 0.04039 - 0.7265  
P value (two-tailed) 0,0286

FIGURE 4A-1





FIGURE 4A-2



3/15

FIGURE 4B-1




FIGURE 4B-2




FIGURE 4C-1




FIGURE 4C-2




FIGURE 4D-1





FIGURE 4D-2



4/15

FIGURE 5





FIGURE 6



5/15

FIGURE 7A





FIGURE 7B  
Pt#1



|                      |                 |
|----------------------|-----------------|
| Number of XY pairs   | 17              |
| Spearman r           | 0.7885          |
| 95% CI               | 0.4839 – 0.9227 |
| p-value (two tailed) | 0.0003          |

6/15

FIGURE 8A




FIGURE 8B




FIGURE 8C




FIGURE 8D




FIGURE 9A




FIGURE 9B




FIGURE 10A




FIGURE 10B




FIGURE 10C




FIGURE 10D




FIGURE 10E





FIGURE 10F



9/15

FIGURE 10G




FIGURE 10H




FIGURE 10I




FIGURE 10J




FIGURE 11A





FIGURE 11B



10/15

FIGURE 11C




FIGURE 12




FIGURE 13A




FIGURE 13B






FIGURE 14



EOT = end of treatment with NI-0501  
Solid lines = pts. with favorable response  
Dashed lines = pts. with insufficient responses


| No. of patients with ANC < 1000/mcl |                  |
|-------------------------------------|------------------|
| Baseline (SD0)                      | End of Treatment |
| 11/18                               | 3/18             |
| p < 0.001 (*)                       |                  |

\* One-sided single proportion test



Median (Q1;Q3)

FIGURE 15



| No. of patients with $\text{PLT} < 100 \times 10^3/\text{mcl}$ |                  |
|----------------------------------------------------------------|------------------|
| Baseline (SD0)                                                 | End of Treatment |
| 13/18                                                          | 6/18             |
| $p < 0.001 (*)$                                                |                  |

\* One-sided single proportion test






FIGURE 16




| Median serum levels<br>(pts. with baseline $> 1000 \text{ ng/ml}$ ; $n=13$ ) |                     |
|------------------------------------------------------------------------------|---------------------|
| Baseline (SD0)                                                               | End of Treatment    |
| 5880 $\text{ng/ml}$                                                          | 1909 $\text{ng/ml}$ |
| $p = 0.006 (*)$                                                              |                     |

\* Paired t-test on log transformed data



14/15

FIGURE 17



| Median dose                     |                       |
|---------------------------------|-----------------------|
| Baseline (SD0)                  | End of Treatment      |
| 10.0 mg/m <sup>2</sup>          | 4.5 mg/m <sup>2</sup> |
| p = 0.02 (*)                    |                       |
| No. of pts with tapering ≥ 50%: |                       |
| 13/17^                          |                       |

\* two-sided Wilcoxon Signed rank test  
^ data for 1 patient not yet available

FIGURE 18




FIGURE 19




**FIGURE 19**

