

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 August 2011 (25.08.2011)

(10) International Publication Number
WO 2011/103105 A1

PCT

(51) International Patent Classification:

C07K 14/78 (2006.01) C12N 15/10 (2006.01)
A61K 38/39 (2006.01) A61K 38/00 (2006.01)
C40B 30/04 (2006.01)

(21) International Application Number:

PCT/US2011/024959

(22) International Filing Date:

16 February 2011 (16.02.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/305,566 18 February 2010 (18.02.2010) US
61/330,706 3 May 2010 (03.05.2010) US

(71) Applicant (for all designated States except US): **BRISTOL-MYERS SQUIBB COMPANY** [US/US]; P.O. Box 4000, Route 206 and ProvinceLine Road, Princeton, New Jersey 08543-4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BUSH, Alex** [US/US]; c/o Bristol-Myers Squibb Company, 100 Beaver Street, Waltham, Massachusetts 02453 (US). **DASGUPTA, Ruchira** [US/US]; c/o Bristol-Myers Squibb Company, 100 Beaver Street, Waltham, Massachusetts 02453 (US). **ENGLE, Linda** [US/US]; c/o

Bristol-Myers Squibb Company, 100 Beaver Street, Waltham, Massachusetts 02453 (US). **SCHNEEWEIS, Lumelle** [US/US]; c/o Bristol-Myers Squibb Company, Route 206 and Province Line Road, Princeton, New Jersey 08543 (US).

(74) Agents: **PARLET, Nickki L.** et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, New Jersey 08543-4000 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GE, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) Title: FIBRONECTIN BASED SCAFFOLD DOMAIN PROTEINS THAT BIND IL-23


FIG. 3

Wildtype VSDVPRDLVVAATPTSLLSWDAPAVTVRYRITVGETGNSPVQETVFGSKCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1434A08 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPRKTYCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1437G04 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPRKTYCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1437A09 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1438E05 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1438D01 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1438B02 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPRKTYCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1438A09 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1486G03 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1486C04 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1486D04 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1486B05 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1486D05 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1487C03 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1487G03 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1487D09 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1487H04 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490E02 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490G02 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490H05 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490B03 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490H06 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490A07 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490C07 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1490H08 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1491A05 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572H03 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1571G04 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1571G06 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1571F10 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT

FIG. 3 (continued)

1572D04 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572F05 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572G06 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572H01 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572C09 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572H05 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
1572H06 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
155C07 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
155C05 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
155C03 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
155C306 VSDVPRDLVVAATPTSLLSWGHYPMHVRYYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT
155C05 VSDVPRDLVVAATPTSLLSWGHYPLHRYRITVGETGNSPVQETVPSRHCATISGLKPGVDYTTVAVTYGRDSSPASSKPIISINVRT

(57) Abstract: The present invention relates to fibronectin based scaffold domain protein that bind interleukin 23 (IL-23), specifically the p19 subunit of IL-23. The invention also relates to the use of the innovative proteins in therapeutic applications to treat autoimmune diseases. The invention further relates to cells comprising such proteins, polynucleotide encoding such proteins or fragments thereof, and to vectors comprising the polynucleotides encoding the innovative proteins.



WO 2011/103105 A1

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG). **Published:**

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *of inventorship (Rule 4.17(iv))*

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

FIBRONECTIN BASED SCAFFOLD DOMAIN PROTEINS THAT BIND IL-23

FIELD OF THE INVENTION

5 [0001] The present invention relates to fibronectin based scaffold domain protein that bind interleukin 23 (IL-23), specifically the p19 subunit of IL-23. The invention also relates to the use of the innovative proteins in therapeutic applications to treat autoimmune diseases. The invention further relates to cells comprising such proteins, polynucleotide encoding such proteins or fragments thereof, and to vectors comprising the polynucleotides encoding the innovative proteins.

10

INTRODUCTION

[0002] IL-23 is a member of the IL-12 heterodimeric cytokine family. It contains the p40 subunit, which is common to IL-12, and a unique p19 subunit. IL-23 sends signals through a heterodimeric receptor complex consisting of IL-12R β 1 and IL-23R (Aggarwal, 15 S et al., "Interleukin-23 promotes a distinct CD4 T cell activation state characterized by the production of interleukin-17", *J. Biol. Chem.*, 278:1910-1914 (2003)). IL-23 is a potential target for the treatment of chronic inflammatory disorders such as multiple sclerosis, rheumatoid arthritis, psoriasis and Crohn's disease.

[0003] Fibronectin based scaffolds are a family of proteins capable of evolving to 20 bind any compound of interest. These proteins, which generally make use of a scaffold derived from a fibronectin type III (Fn3) or Fn3-like domain, function in a manner characteristic of natural or engineered antibodies (that is, polyclonal, monoclonal, or single-chain antibodies) and, in addition, possess structural advantages. Specifically, the structure of these antibody mimics has been designed for optimal folding, stability, and 25 solubility, even under conditions that normally lead to the loss of structure and function in antibodies. An example of fibronectin-based scaffold proteins are Adnectins (Adnexus, a Bristol-Myers Squibb R&D Company).

[0004] Fibronectin type III (Fn3) domains comprise, in order from N-terminus to C-terminus, a beta or beta-like strand, A; a loop, AB; a beta or beta-like strand, B; a loop, 30 BC; a beta or beta-like strand C; a loop CD; a beta or beta-like strand D; a loop DE; a beta or beta-like strand, E; a loop, EF; a beta or beta-like strand F; a loop FG; and a beta or beta-like strand G. Any or all of loops AB, BC, CD, DE, EF and FG may participate in

target binding. The BC, DE, and FG loops are both structurally and functionally analogous to the complementarity determining regions (CDRs) from immunoglobulins. U.S. Patent No. 7,115,396 describes Fn3 domain proteins wherein alterations to the BC, DE, and FG loops result in high affinity TNF α binders. U.S. Publication No.

5 2007/0148126 describes Fn3 domain proteins wherein alterations to the BC, DE, and FG loops result in high affinity VEGFR2 binders.

[0005] It would be advantageous to obtain improved fibronectin domain scaffold proteins for therapeutic treatment of autoimmune disorders. A subset of effector T cells that produce interleukin 17 (IL-17; "Th17 cells") are highly proinflammatory and induce
10 severe autoimmunity. Th17 cells express a distinct subset of cytokines and chemokines compared to Th1 and Th2 cells, including IL-6, tumor necrosis factor (TNF), IL-22, IL-17A and IL-17F as well as the chemokine receptor CCR6. IL-23 promotes the production of IL-17 by activated T cells (Aggarwal, S et al., "Interleukin-23 promotes a distinct CD4 T cell activation state characterized by the production of interleukin-17", *J. Biol. Chem.*,
15 278:1910-1914 (2003)) and is a key cytokine to induce expansion of IL-17-producing CD4+ T cells. Exposure to IL-23 seems to be the key feature that determines the pathogenicity of Th17 cells.

SUMMARY OF THE INVENTION

20 **[0006]** The application provides Adnectins™ against human IL-23-specific p19 subunit. One aspect of the invention provides for polypeptides comprising Fn3 domain in which one or more of the solvent accessible loops has been randomized or mutated. In some embodiments, the Fn3 domain is a Fn3 domain derived from the wild-type tenth module of the human fibronectin type III domain (¹⁰Fn3). In some embodiments, the
25 ¹⁰Fn3 polypeptide of the invention is at least 40%, 50%, 60%, 65%, 70%, 75%, 80%, 85%, or 90% identical to the human ¹⁰Fn3 domain.

[0007] In some embodiments, one or more loops selected from BC, DE, and FG may be extended or shortened in length relative to the corresponding human fibronectin loop.

[0008] In some embodiments, the polypeptides of the invention comprises a tenth
30 fibronectin type III (¹⁰Fn3) domain, wherein the ¹⁰Fn3 domain comprises a loop, AB; a loop, BC; a loop, CD; a loop, DE; a loop EF; and a loop FG; and has at least one loop

selected from loop BC, DE, and FG with an altered amino acid sequence relative to the sequence of the corresponding loop of the human ¹⁰Fn3 domain.

[0009] In some embodiments, the polypeptide of the invention comprises a Fn3 domain that comprises an amino acid sequence at least 80, 85, 90, 95, 98, 99 or 100% identical to the non-loop regions.

[0010] In some embodiments, the BC loop of the protein of the invention comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 2-6.

[0011] In some embodiments, the DE loop of the protein of the invention comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 7-48.

10 [0012] In some embodiments, the FG loop of the protein of the invention comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 49-59.

[0013] In some embodiments, the ¹⁰Fn3 domain may begin and/or end with amino acid substitutions, insertions or deletions.

[0014] In some embodiments, the protein of the invention comprises one loop sequence from the BC loop sequences shown in SEQ ID NOs: 2-6, one DE loop sequence shown in SEQ ID NOs: 7-48 and one FG loop sequence shown in SEQ ID NOs: 49-59.

[0015] In some embodiments, the protein of the invention comprises a BC, DE and FG loop amino acid sequence at least 70, 75, 80, 85, 90, 95, 98, 99 or 100% identical to of any one of SEQ ID NOS:2-59.

20 [0016] In some embodiments, the anti-IL-23 Adnectin comprises the amino acid sequence of any one of SEQ ID NOS:60-100.

[0017] In some embodiments, the anti-IL-23 Adnectin comprises the Fn3 domain amino acid sequence from position 3-96 of any one of SEQ ID NOS:60-100.

[0018] In some embodiments, the anti-IL-23 Adnectin comprises the amino acid sequence at least 70, 75, 80, 85, 90, 95, 98, 99 or 100% identical to of any one of SEQ ID NOS:60-100.

[0019] In one aspect, the anti-IL-23 Adnectin further comprising a pharmacokinetic (PK) moiety. In some embodiments, the PK moiety comprises polyethylene glycol (PEG).

30 [0020] In one aspect, the application provides an anti-IL-23 Adnectin useful in the treatment of autoimmune diseases.

[0021] In one aspect, the present invention provides a fusion polypeptide comprising a fibronectin type III tenth (¹⁰Fn3) domain and anti-IL-23 Adnectin, wherein the ¹⁰Fn3 domain binds to HSA with a Kd of 1 uM or less. In certain embodiments, the ¹⁰Fn3 domain comprises an amino acid sequence at least 70% identical to SEQ ID NO: 103. In one embodiment, the ¹⁰Fn3 domain comprises a BC loop having the amino acid sequence set forth in SEQ ID NO: 104, a DE loop having the amino acid sequence set forth in SEQ ID NO: 105, and an FG loop having the amino acid sequence set forth in SEQ ID NO:106. In another embodiment, the ¹⁰Fn3 domain comprises one or more of a BC loop having the amino acid sequence set forth in SEQ ID NO: 104, a DE loop having the amino acid sequence set forth in SEQ ID NO: 105, and an FG loop having the amino acid sequence set forth in SEQ ID NO: 106.

[0022] In one embodiment, the ¹⁰Fn3 domain of the fusion polypeptide also binds to one or more of rhesus serum albumin (RhSA), cynomolgus monkey serum albumin (CySA), or murine serum albumin (MuSA). In other embodiments, the ¹⁰Fn3 domain does not cross-react with one or more of RhSA, CySA or MuSA.

[0023] In certain embodiments, the ¹⁰Fn3 domain of the fusion polypeptide binds to HSA with a Kd of 1 uM or less. In some embodiments, the ¹⁰Fn3 domain binds to HSA with a Kd of 500 nM or less. In other embodiments, the ¹⁰Fn3 domain binds to HSA with a Kd of at least 200 nM, 100 nM, 50 nM, 20 nM, 10 nM, or 5 nM.

[0024] In other embodiments, the ¹⁰Fn3 domain of the fusion polypeptide binds to domain I or II of HSA. In one embodiment, the ¹⁰Fn3 domain binds to both domains I and II of HSA. In some embodiments, the ¹⁰Fn3 domain binds to HSA at a pH range of 5.5 to 7.4. In other embodiments, the ¹⁰Fn3 domain binds to HSA with a Kd of 200 nM or less at pH 5.5. In another embodiment, the ¹⁰Fn3 domain binds to HSA with a Kd of at least 500 nM, 200 nM, 100 nM, 50 nM, 20 nM, 10 nM, or 5 nM at a pH range of 5.5 to 7.4. In one embodiment, the ¹⁰Fn3 domain binds to HSA with a Kd of at least 500 nM, 200 nM, 100 nM, 50 nM, 20 nM, 10 nM, or 5 nM at pH 5.5.

[0025] In some embodiments, the serum half-life of the fusion polypeptide in the presence of serum albumin is at least 5-fold greater than the serum half-life of the polypeptide in the absence of serum albumin. In certain embodiments, the serum half-life of the fusion polypeptide in the presence of serum albumin is at least 2-fold, 5-fold, 7-fold, 10-fold, 12-fold, 15-fold, 20-fold, 22-fold, 25-fold, 27-fold, or 30-fold greater than

the serum half-life of the polypeptide in the absence of serum albumin. In some embodiments, the serum albumin is any one of HSA, RhSA, CySA, or MuSA.

- [0026]** In certain embodiments, the serum half-life of the fusion polypeptide in the presence of serum albumin is at least 20 hours. In certain embodiments, the serum half-life of the fusion polypeptide in the presence of serum albumin is at least 10 hours, 12 hours, 15 hours, 20 hours, 25 hours, 30 hours, 40 hours, 50 hours, 75 hours, 90 hours, 100 hours, 110 hours, 120 hours, 130 hours, 150 hours, 170 hours, or 200 hours. In some embodiments, the half-life of the fusion polypeptide is observed in a primate (*e.g.*, human or monkey) or a murine.
- 10 **[0027]** In any of the foregoing aspects and embodiments, the ¹⁰Fn3 domain comprises a sequence selected from SEQ ID NO: 107, 111, 115, 119, and 123-143.

BRIEF DESCRIPTION OF THE FIGURES

- [0028]** Figure 1 shows the full length DNA sequence alignment of the anti-IL23 Adnectin of the invention.
- 15 **[0029]** Figure 2 shows pBMS2008/ATI001044 protein expression vector as described in Example 2.
- [0030]** Figure 3 shows the full length amino acid sequence alignment of the anti-IL23 adnectin of the invention.
- 20 **[0031]** Figure 4 shows a representative IC₅₀ curves from PBMC pSTAT3 inhibition by anti-IL-23 adnectin as described in Example 4.
- [0032]** Figure 5 shows a representative IC₅₀ curves for inhibition of IL-23-dependent IL-17A by anti-IL-23 adnectins and anti-p40 monoclonal antibody (MAB1510) as described in Example 4.
- 25 **[0033]** Figure 6 shows ATI001045 inhibition of IL-23-induced IL-17 production by PBMCs of donor 228 (one of 4 donors tested as described in Example 4).
- [0034]** Figure 7 shows representative selectivity data for the anti-IL-23 adnectins. Buffer subtracted sensorgrams illustrating the association and dissociation phases of 10 nM IL-23 and 1 uM IL-12 binding to captured ATI001016 as described in Example 4 are
- 30 shown.
- [0035]** Figure 8 shows that anti-IL-23 adnectins do not inhibit IL-12 induced IFN- γ production in NK-92 cells as described in Example 4.

- [0036] Figure 9A shows that ATI001045 Inhibits Serum IL-17 Levels in Mouse Pharmacodynamic Model as described in Example 4.
- [0037] Figure 9B shows a comparison of inhibitory activities of anti-IL-23 adnectins in a mouse pharmacodynamic model as described in Example 4.
- 5 [0038] Figure 10A shows ATI000934 dose response in human IL-23 induced acanthosis as described in Example 4.
- [0039] Figure 10B shows ATI001045 Dose Response in Human IL-23-Induced acanthosis as described in Example 4.
- [0040] Figure 11 shows *in vivo* HSA half-life in mice. HSA was injected into mice at
10 20 mg/kg (Figure 11A) or 50 mg/kg (Figure 11B).
- [0041] Figures 12A-D show the half-life determination of SABA1-SABA4 in mice.
- [0042] Figure 13A shows a graph summary of half-life enhancement in mice of SABA1-4 when co-injected with HSA. Figure 13b compares data from cynomolgus monkey and mice.
- 15 [0043] Figures 14A-B show the half-life determination for SABA1.1 and SABA5.1 in cynomolgus monkey.
- [0044] Figure 15 shows SABA1.2 binding to albumins from human, mouse and rat by direct binding ELISA assay.
- [0045] Figure 16 shows the determination of SABA1.1 and HSA stoichiometry.
- 20 [0046] Figure 17 shows Biacore analysis of SABA1.2 binding to recombinant domain fragments of HSA.
- [0047] Figure 18 shows the pharmacokinetic profile for SABA1.2 in monkeys dosed at 1 mpk and 10 mpk.
- [0048] Figure 19 shows the pharmacokinetic profile for SABA1.2 in monkeys dosed
25 intravenously or subcutaneously at 1 mpk.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

- [0049] By a “polypeptide” is meant any sequence of two or more amino acids,
30 regardless of length, post-translation modification, or function. “Polypeptide,” “peptide,” and “protein” are used interchangeably herein. Polypeptides can include natural amino acids and non-natural amino acids such as those described in U.S. Patent No. 6,559,126,

incorporated herein by reference. Polypeptides can also be modified in any of a variety of standard chemical ways (*e.g.*, an amino acid can be modified with a protecting group; the carboxy-terminal amino acid can be made into a terminal amide group; the amino-terminal residue can be modified with groups to, *e.g.*, enhance lipophilicity; or the polypeptide can be chemically glycosylated or otherwise modified to increase stability or *in vivo* half-life). Polypeptide modifications can include the attachment of another structure such as a cyclic compound or other molecule to the polypeptide and can also include polypeptides that contain one or more amino acids in an altered configuration (*i.e.*, R or S; or, L or D). The peptides of the invention are proteins derived from the tenth type III domain of fibronectin that have been modified to bind specifically to the p19 subunit of IL-23 and are referred to herein as “Adnectin” or “anti-IL-23 Adnectin”.

[0050] The term “PK” is an acronym for “pharmokinetic” and encompasses properties of a compound including, by way of example, absorption, distribution, metabolism, and elimination by a subject. A “PK modulation protein” or “PK moiety” refers to any protein, peptide, or moiety that affects the pharmacokinetic properties of a biologically active molecule when fused to or administered together with the biologically active molecule. Examples of a PK modulation protein or PK moiety include PEG, human serum albumin (HSA) binders (as disclosed in U.S. Publication Nos. 2005/0287153 and 2007/0003549), human serum albumin, Fc or Fc fragments, and sugars (*e.g.*, sialic acid).

[0051] “Percent (%) amino acid sequence identity” herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in a selected sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR®) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared.

[0052] An “isolated” polypeptide is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to greater than 95% by weight of polypeptide as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing condition using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes the polypeptide in situ within recombinant cells since at least one component of the polypeptide’s natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

[0053] The “half-life” of an amino acid sequence or compound can generally be defined as the time taken for the serum concentration of the polypeptide to be reduced by 50%, *in vivo*, for example due to degradation of the sequence or compound and/or clearance or sequestration of the sequence or compound by natural mechanisms. The half-life can be determined in any manner known per se, such as by pharmacokinetic analysis. Suitable techniques will be clear to the person skilled in the art, and may for example generally involve the steps of suitably administering to the primate a suitable dose of the amino acid sequence or compound of the invention; collecting blood samples or other samples from said primate at regular intervals; determining the level or concentration of the amino acid sequence or compound of the invention in said blood sample; and calculating, from (a plot of) the data thus obtained, the time until the level or concentration of the amino acid sequence or compound of the invention has been reduced by 50% compared to the initial level upon dosing. Reference is for example made to the standard handbooks, such as Kenneth, A. et al., *Chemical Stability of Pharmaceuticals: A Handbook for Pharmacists* and in Lee, P.I.D. et al., *Pharmacokinetic Analysis: A Practical Approach* (1996). Reference is also made to Gibaldi, M. et al., *Pharmacokinetics*, 2nd Rev. Edition, Marcel Dekker, publ. (1982).

[0054] Half-life can be expressed using parameters such as the $t_{1/2}$ -alpha, $t_{1/2}$ -beta and the area under the curve (AUC). In the present specification, an “increase in half-life”

refers to an increase in any one of these parameters, any two of these parameters, or all three these parameters. An “increase in half-life” in particular refers to an increase in the $t_{1/2}$ -beta, either with or without an increase in the $t_{1/2}$ -alpha and/or the AUC or both.

5 Overview

[0055] The application provides Adnectins against human IL-23-specific p19 subunit. In order to identify IL-23 specific antagonist, IL-23 was presented to large synthetic libraries of Adnectin using anti-p40 mAbs. Adnectins that bound to IL-23 p19 subunit were screened for binding to human IL-23, competition of the IL-23/IL-23R interaction and inhibition of IL-23 induced signaling in a T-cell line. The anti-IL-23 Adnectins were subjected to further selective pressure by lowering the target concentration and selecting for anti-IL-23 Adnectins with slow off-rates. From this optimization process a family of Adnectins were identified as IL-23 specific inhibitors with favorable biochemical and biophysical properties.

15

Fibronectin Based Scaffolds

[0056] One aspect of the application provides for polypeptides comprising Fn3 domain in which one or more of the solvent accessible loops has been randomized or mutated. In some embodiments, the Fn3 domain is an Fn3 domain derived from the wild-type tenth module of the human fibronectin type III domain (10 Fn3):

20 VSDVPRDLEVVAATPTSLLISWDAPAVTVRYYRITYGETGGNSPVQEFTVPGSKS
TATISGLKPGVDYTITVYAVTGRGDSPASSKPIISINYRT (SEQ ID NO: 1). In the 10 Fn3 sequence above, the BC, DE and FG loops are underlined.

[0057] A variety of mutant 10 Fn3 scaffolds have been reported. In one aspect, one or more of Asp 7, Glu 9, and Asp 23 is replaced by another amino acid, such as, for example, a non-negatively charged amino acid residue (*e.g.*, Asn, Lys, etc.). These mutations have been reported to have the effect of promoting greater stability of the mutant 10 Fn3 at neutral pH as compared to the wild-type form (See, PCT Publication No. WO02/04523). A variety of additional alterations in the 10 Fn3 scaffold that are either beneficial or neutral have been disclosed. See, for example, Batori et al., *Protein Eng.*, 15(12):1015-1020 (Dec. 2002); Koide et al., *Biochemistry*, 40(34):10326-10333 (Aug. 28, 2001).

30

[0058] Both variant and wild-type ¹⁰F_n3 proteins are characterized by the same structure, namely seven beta-strand domain sequences designated A through G and six loop regions (AB loop, BC loop, CD loop, DE loop, EF loop, and FG loop) which connect the seven beta-strand domain sequences. The beta strands positioned closest to the N- and C-termini may adopt a beta-like conformation in solution. In SEQ ID NO:1, the AB loop corresponds to residues 15-16, the BC loop corresponds to residues 21-30, the CD loop corresponds to residues 39-45, the DE loop corresponds to residues 51-56, the EF loop corresponds to residues 60-66, and the FG loop corresponds to residues 76-87 (Xu et al., *Chemistry & Biology*, 9:933-942 (2002)).

[0059] In some embodiments, the ¹⁰F_n3 polypeptide may be at least 40%, 50%, 60%, 65%, 70%, 75%, 80%, 85%, or 90% identical to the human ¹⁰F_n3 domain, shown in SEQ ID NO:1. Much of the variability will generally occur in one or more of the loops. Each of the beta or beta-like strands of a ¹⁰F_n3 polypeptide may consist essentially of an amino acid sequence that is at least 80%, 85%, 90%, 95% or 100% identical to the sequence of a corresponding beta or beta-like strand of SEQ ID NO:1, provided that such variation does not disrupt the stability of the polypeptide in physiological conditions.

[0060] In some embodiments, the disclosure provides polypeptides comprising a tenth fibronectin type III (¹⁰F_n3) domain, wherein the ¹⁰F_n3 domain comprises a loop, AB; a loop, BC; a loop, CD; a loop, DE; a loop EF; and a loop FG; and has at least one loop selected from loop BC, DE, and FG with an altered amino acid sequence relative to the sequence of the corresponding loop of the human ¹⁰F_n3 domain. In some embodiments, the BC and FG loops are altered, in some embodiments, the BC, DE, and FG loops are altered, *i.e.*, the F_n3 domains comprise non-naturally occurring loops. By “altered” is meant one or more amino acid sequence alterations relative to a template sequence (corresponding human fibronectin domain) and includes amino acid additions, deletions, and substitutions. Altering an amino acid sequence may be accomplished through intentional, blind, or spontaneous sequence variation, generally of a nucleic acid coding sequence, and may occur by any technique, for example, PCR, error-prone PCR, or chemical DNA synthesis.

[0061] In some embodiments, one or more loops selected from BC, DE, and FG may be extended or shortened in length relative to the corresponding human fibronectin loop. In some embodiments, the length of the loop may be extended by 2-25 amino acids. In

some embodiments, the length of the loop may be decreased by 1-11 amino acids. To optimize antigen binding, therefore, the length of a loop of ¹⁰Fn3 may be altered in length as well as in sequence to obtain the greatest possible flexibility and affinity in antigen binding.

5 **[0062]** In some embodiments, the polypeptide comprises a Fn3 domain that comprises an amino acid sequence at least 80, 85, 90, 95, 98, 99 or 100% identical to the non-loop regions of SEQ ID NO:1, wherein at least one loop selected from BC, DE, and FG is altered. In some embodiments, the altered BC loop has up to 10 amino acid substitutions, up to 4 amino acid deletions, up to 10 amino acid insertions, or a
10 combination thereof. In some embodiments, the altered DE loop has up to 6 amino acid substitutions, up to 4 amino acid deletions, up to 13 amino acid insertions or a combination thereof. In some embodiments, the FG loop has up to 12 amino acid substitutions, up to 11 amino acid deletions, up to 25 amino acid insertions or a combination thereof.

15 **[0063]** In some embodiments, the BC loop of the protein of the invention comprises an amino acid sequence selected from the group consisting of GHYPMHV (SEQ ID NO: 2), GHYPLHV (SEQ ID NO: 3), GHYPMHI (SEQ ID NO:4), GHYPLHI (SEQ ID NO:5) and GHYPLHL (SEQ ID NO:6).

[0064] In some embodiments, the DE loop of the protein of the invention comprises
20 an amino acid sequence selected from the group consisting of HIRTH(SEQ ID NO:7), YYHY(SEQ ID NO:8), SKQH (SEQ ID NO:9), SNVH (SEQ ID NO:10), NRAH (SEQ ID NO:11), RKTY(SEQ ID NO:12), RSRV (SEQ ID NO:13), SRYV (SEQ ID NO:14), PHRY (SEQ ID NO:15), RSTH (SEQ ID NO:16), SRIY (SEQ ID NO:17), HQRY (SEQ ID NO:18), KQVY (SEQ ID NO:19), AHRY (SEQ ID NO:20), RSRH (SEQ ID NO:21),
25 ARQY (SEQ ID NO:22), RTQY (SEQ ID NO:23), PRYH (SEQ ID NO:24), MRQH (SEQ ID NO:25), SRKY (SEQ ID NO:26), RQKY (SEQ ID NO:27), HAKY(SEQ ID NO:28), SNRY (SEQ ID NO:29), NTSH (SEQ ID NO:30), SQVY (SEQ ID NO:31), NRVY (SEQ ID NO:32), PRSH (SEQ ID NO:33), RTKY (SEQ ID NO:34), SRYH (SEQ ID NO:35), PRRY(SEQ ID NO:36), RQKY (SEQ ID NO:37), RYKY (SEQ ID NO:38),
30 VPRH (SEQ ID NO:39), TPKH (SEQ ID NO:40), RSKY (SEQ ID NO:41), SRKY (SEQ ID NO:42), VPRY (SEQ ID NO:43), PRRY (SEQ ID NO:44), RMRH (SEQ ID NO:45), PPRH (SEQ ID NO:46), RQIY (SEQ ID NO:47), and MRQH(SEQ ID NO:48).

[0065] In some embodiments, the FG loop of the protein of the invention comprises an amino acid sequence selected from the group consisting of YYNEADYSQI (SEQ ID NO:49), YYQEYERYI (SEQ ID NO:50), YMEEEKYAVI (SEQ ID NO:51), YYAQENYKEI (SEQ ID NO:52), YYKEANYREI (SEQ ID NO:53), YYAQEEYHII (SEQ ID NO:54), YYKEADYSQI (SEQ ID NO:55), YYEQVEYREI (SEQ ID NO:56), YYEQPIYATI (SEQ ID NO:57), YYEQVEYREI (SEQ ID NO:58) and YYSEELYKYI (SEQ ID NO:59).

[0066] The ¹⁰F_{n3} domain may begin with amino acid alterations. For example, an additional MG sequence may be placed at the N-terminus of an F_{n3} domain. The M will usually be cleaved off, leaving a G at the N-terminus. In some embodiments, sequences may be placed at the C-terminus of the ¹⁰F_{n3} domain. For example, in site directed PEGylation where a cysteine containing linker such as GSGC (SEQ ID NO: 101) is added to the C-terminus. Alternatively, PEGylation of the naturally occurring C-terminus tail that has been mutated by changing the Ser to a Cys for a cysteine containing linker EIDKPCQ (SEQ ID NO: 102). Examples of the anti-IL-23 adnectin of the invention comprising the GSGC linker include ATI001014, ATI001015, ATI001016, ATI001044, ATI001045 and ATI001047. ATI000934 is an example of the anti-IL-23 adnectin of the invention comprising the EIDKPCQ linker.

[0067] In some embodiments, the protein of the invention comprises one loop sequence from the BC loop sequences shown in SEQ ID NOS: 2-6, one DE loop sequence shown in SEQ ID NOS: 7-48 and one FG loop sequence shown in SEQ ID NOS: 49-59. In some embodiments, the protein of the invention comprises a BC, DE and FG loop amino acid sequence at least 70, 75, 80, 85, 90, 95, 98, 99 or 100% identical to of any one of SEQ ID NOS:2-59.

[0068] Further, one skilled in the art will recognize that BC loop sequences shown in SEQ ID NO: 2-6 share a common sequence motif GHYPX₁HX₂ (SEQ ID NO:257) where X₁ is either M or L, and X₂ is either I or V, and the FG loop sequences shown in SEQ ID NO: 49-59 share a common sequence motif YYX₃X₃X₃X₃YX₃X₃I (SEQ ID NO: 258) where X₃ can be any amino acid. It would therefore be possible to generate additional Adnectins that bind IL-23 with BC loops that fit the consensus sequence GHYPX₁HX₂ and/or with other FG loops, beyond those explicitly listed in SEQ ID NOS:49-59, that fit the pattern YYX₃X₃X₃X₃YX₃X₃I.

[0069] In some embodiments, the anti-IL-23 Adnectin comprises the amino acid sequence of any one of SEQ ID NOS:60-100. In some embodiments, the anti-IL-23 Adnectin comprises the Fn3 domain amino acid sequence from position 3-96 of any one of SEQ ID NOS:60-100. In some embodiments, the anti-IL-23 Adnectin comprises the amino acid sequence at least 70, 75, 80, 85, 90, 95, 98, 99 or 100% identical to any one of SEQ ID NOS: 60-100. In some embodiments, the anti-IL-23 Adnectin comprises the amino acid sequence at least 70, 75, 80, 85, 90, 95, 98, 99 or 100% identical to amino acid sequence from position 3-96 any one of SEQ ID NOS:60-100.

[0070] In some embodiments, the anti-IL-23 Adnectin may be pegylated and/or contain a his-tag. As used herein, ATI000934 refers to a protein wherein the loop sequences are identical to those of construct 1571G06 (Seq ID 87), and the protein contains the residues EIDKPCQ at the C-terminus where the protein is pegylated and contains a his-tag. ATI001014 refers to a protein wherein the loop sequences are identical to those of construct 1571G04 (Seq ID 86), and the protein contains a GSGC linker at the C-terminus where the protein is pegylated and contains a his-tag. ATI001015 refers to a protein wherein the loop sequences are identical to those of construct 1572G06 (Seq ID 91), and the protein contains a GSGC linker at the C-terminus where the protein is pegylated and contains a his-tag. ATI001016 refers to a protein wherein the loop sequences are identical to those of construct 1490B03 (Seq ID 79), and the protein contains a GSGC linker at the C-terminus where the protein is pegylated and contains a his-tag. ATI001044 refers to a protein wherein the loop sequences are identical to those of construct 1490B03 (Seq ID 79), and the protein contains a GSGC linker at the C-terminus, but protein is not pegylated and there is no his tag. ATI001045 refers to a protein wherein the loop sequences are identical to those of construct 1490B03 (Seq ID 79), and the protein contains a GSGC linker at the C-terminus where the protein is pegylated; and there is no his tag. ATI001047 refers to a protein wherein the loop sequences are identical to those of construct 1571G04 (Seq ID 86), and the protein contains a GSGC linker at the C-terminus where the protein is pegylated, and there is no his tag

[0071] Fibronectin naturally binds certain types of integrins through its integrin-binding motif, "arginine-glycine-aspartic acid" (RGD). In some embodiments, the polypeptide comprises a ¹⁰F_n3 domain that lacks the (RGD) integrin binding motif.

Pharmacokinetic Moieties

[0072] In one aspect, the application provides for anti-IL-23 Adnectin further comprising a pharmacokinetic (PK) moiety. Improved pharmacokinetics may be assessed according to the perceived therapeutic need. Often it is desirable to increase bioavailability and/or increase the time between doses, possibly by increasing the time that a protein remains available in the serum after dosing. In some instances, it is desirable to improve the continuity of the serum concentration of the protein over time (e.g., decrease the difference in serum concentration of the protein shortly after administration and shortly before the next administration). The anti-IL-23 Adnectin may be attached to a moiety that reduces the clearance rate of the polypeptide in a mammal (e.g., mouse, rat, or human) by greater than three-fold relative to the unmodified Adnectin. Other measures of improved pharmacokinetics may include serum half-life, which is often divided into an alpha phase and a beta phase. Either or both phases may be improved significantly by addition of an appropriate moiety.

[0073] Moieties that tend to slow clearance of a protein from the blood, herein referred to as "PK moieties", include polyoxyalkylene moieties, e.g., polyethylene glycol, sugars (e.g., sialic acid), and well-tolerated protein moieties (e.g., Fc, Fc fragments, transferrin, or serum albumin). The Adnectin may be fused to albumin or a fragment (portion) or variant of albumin as described in U.S. Publication No. 2007/0048282.

[0074] In some embodiments, the PK moiety is a serum albumin binding protein such as those described in U.S. Publication Nos. 2007/0178082 and 2007/0269422.

[0075] In some embodiments, the PK moiety is a serum immunoglobulin binding protein such as those described in U.S. Publication No. 2007/0178082.

[0076] In some embodiments, the Adnectin comprises polyethylene glycol (PEG). One or more PEG molecules may be attached at different positions on the protein, and such attachment may be achieved by reaction with amines, thiols or other suitable reactive groups. The amine moiety may be, for example, a primary amine found at the N-terminus of a polypeptide or an amine group present in an amino acid, such as lysine or arginine. In some embodiments, the PEG moiety is attached at a position on the polypeptide selected from the group consisting of: a) the N-terminus; b) between the N-terminus and the most N-terminal beta strand or beta-like strand; c) a loop positioned on a

face of the polypeptide opposite the target-binding site; d) between the C-terminus and the most C-terminal beta strand or beta-like strand; and e) at the C-terminus.

[0077] Pegylation may be achieved by site-directed pegylation, wherein a suitable reactive group is introduced into the protein to create a site where pegylation

5 preferentially occurs. In some embodiments, the protein is modified to introduce a cysteine residue at a desired position, permitting site directed pegylation on the cysteine. PEG may vary widely in molecular weight and may be branched or linear.

[0078] In some embodiments, the Adnectin comprises an Fn3 domain and a PK moiety. In some embodiments, the Fn3 domain is a ¹⁰Fn3 domain. In some embodiments, 10 the PK moiety increases the serum half-life of the polypeptide by more than 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 150, 200, 400, 600, 800, 1000% or more relative to the Fn3 domain alone.

[0079] In some embodiments, the PK moiety is a polymeric sugar. In some 15 embodiments, the PK moiety is a polyethylene glycol moiety. In some embodiments the PK moiety is a serum albumin binding protein. In some embodiments the PK moiety is human serum albumin. In some embodiments the PK moiety is a serum immunoglobulin binding protein. In some embodiments, the PK moiety is transferrin. In some 20 embodiments the PK moiety is another Adnectin specific for a serum protein.

20 Biophysical and Biochemical Characterization

[0080] The application provides Adnectin comprising a Fn3 domain that binds to the p19 subunit of IL-23. As shown in Table 1 and Example 4, polypeptide binding to a target molecule may be assessed in terms of equilibrium constants (*e.g.*, dissociation, K_D) and in terms of kinetic constants (*e.g.*, on-rate constant, K_{on} and off-rate constant, k_{off}).

25 An Adnectin will generally bind to a target molecule with a K_D of less than 500 nM, 100 nM, 10 nM, 1 nM, 500 pM, 200 pM, 100 pM, although higher K_D values may be tolerated where the K_{off} is sufficiently low or the K_{on} , is sufficiently high.

[0081] The BC, DE and FG loop sequences of the family of anti-IL-23 Adnectin of the invention are presented in Table 1 below, as well as the corresponding full length 30 SEQ ID NO.

Table 1: Anti-IL-23 Adnectin Family

Clone ID	BC Loop	DE Loop	FG loop	On-rate (k_a , $M^{-1}s^{-1}$)	Off-rate (k_d , s^{-1})	Affinity (K_D , M)	Kit 225 pSTAT3 IC50 (nM)	SEQ ID NO
1434A08	GHYPMHV	HRTH	YYNEADYSQI	n.d.	n.d.	n.d.	4.9	60
1437G04	GHYPLHV	YYHY	YYNEADYSQI	1.34E+05	4.46E-04	3.3E-09	3.3	61
1437A09	GHYPMHV	SKQH	YYNEADYSQI	8.06E+04	4.60E-04	5.7E-09	20.2	62
1438E05	GHYPMHV	SNVH	YYNEADYSQI	n.d.	n.d.	n.d.	39.6	63
1438D01	GHYPMHV	NRAH	YYNEADYSQI	n.d.	n.d.	n.d.	11.4	64
1438B02	GHYPMHV	RKTY	YYNEADYSQI	n.d.	n.d.	n.d.	8.6	65
1438A09	GHYPMHV	RSRY	YYNEADYSQI	n.d.	n.d.	n.d.	10.8	66
1486G03	GHYPMHV	SRYY	YYNEADYSQI	8.62E+04	4.25E-04	4.9E-09	3	67
1486C04	GHYPMHV	PHRY	YYNEADYSQI	1.12E+05	3.79E-04	3.4E-09	31.9	68
1486D04	GHYPLHI	RSTH	YYNEADYSQI	1.51E+05	3.52E-04	2.3E-09	2	69
1486B05	GHYPMHV	SRIY	YYNEADYSQI	1.34E+05	3.81E-04	2.8E-09	4	70
1486D05	GHYPLHV	HQRY	YYNEADYSQI	1.20E+05	3.44E-04	2.9E-09	4	71
1487C03	GHYPLHI	KQVY	YYNEADYSQI	1.61E+05	3.82E-04	2.4E-09	3.7	72
1487G03	GHYPLHV	AHRY	YYNEADYSQI	1.03E+05	3.16E-04	3.1E-09	3	73
1487D09	GHYPMHI	RSRH	YYNEADYSQI	1.50E+05	2.50E-04	1.7E-09	2	74
1487H04	GHYPMHV	ARQY	YYQEYERYI	5.96E+04	too slow to measure	<nM	2	75
1490 E02	GHYPMHV	RTQY	YYNEADYSQI	9.73E+04	4.65E-04	4.8E-09	n.d.	76

Clone ID	BC Loop	DE Loop	FG loop	On-rate (k_{on} , $M^{-1}s^{-1}$)	Off-rate (k_{off} , s^{-1})	Affinity (K_D , M)	Kit 225 pSTAT3 IC50 (nM)	SEQ ID NO
1490G02	GHYPMHV	PRYH	YYMEEKYAVI	1.97E+05	3.32E-04	1.7E-09	0.4	77
1490H05	GHYPLHV	MRQH	YYAQENYKEI	1.56E+05	3.22E-04	2.1E-09	0.4	78
1490B03	GHYPLHV	SRKY	YYKEANYREI	1.53E+05	too slow to measure	<nM	0.3	79
1490H06	GHYPLHI	RQKY	YYNEADYSQI	9.35E+04	too slow to measure	<nM	2.3	80
1490A07	GHYPLHI	HAKY	YYAQENYKEI	1.63E+05	too slow to measure	<nM	0.6	81
1490C07	GHYPLHV	SNRY	YYNEADYSQI	9.56E+04	4.26E-04	4.5E-09	5.6	82
1490H08	GHYPLHI	NTSH	YYNEADYSQI	1.67E+05	5.87E-04	3.5E-09		83
1491A05	GHYPLHV	SQVY	YYAQENYKEI	1.98E+05	4.08E-04	2.1E-09	0.3	84
1571H03	GHYPLHV	NRVY	YYAQEEYHII	1.18E+05	3.95E-04	3.4E-09	0.4	85
1571G04	GHYPLHV	PRSH	YYAQENYKEI	1.70E+05	3.61E-04	2.1E-09	0.1	86
1571G06	GHYPLHL	RTKY	YYKEADYSQI	1.31E+05	2.38E-04	1.8133E-09	0.3	87
1571F10	GHYPLHI	SRYH	YYEQVEYREI	2.68E+05	too slow to measure	<nM	0.1	88
1572D04	GHYPMHV	PRRY	YYEQPIYATI	1.03E+05	2.94E-04	2.9E-09	4	89
1572F05	GHYPLHI	RQKY	YYNEADYSQI	1.32E+05	2.85E-04	2.2E-09	3	90
1572G06	GHYPLHV	RYKY	YYAQENYKEI	1.05E+05	too slow to measure	<nM	0.05	91
1572B10	GHYPMHV	VPRH	YYNEADYSQI	6.53E+04	too slow to measure	<nM	0.8	92
1572C09	GHYPMHI	TPKH	YYNEADYSQI	6.07E+04	too slow to measure	<nM	0.5	93
1572H05	GHYPLHI	RSKY	YYEQVEYREI	3.75E+05	2.30E-04	6.1E-10	0.01	94

Clone ID	BC Loop	DE Loop	FG loop	On-rate (k_{on} , $M^{-1}s^{-1}$)	Off-rate (k_{off} , s^{-1})	Affinity (K_D , M)	Kit 225 pSTAT3 IC50 (nM)	SEQ ID NO
1572H08	GHYPLHV	SRKY	YYNEADYSQI	5.73E+04	too slow to measure	<nM	2.8	95
1550A07	GHYPMHV	VPRY	YYAQENYKEI	1.17E+05	1.33E-04	1.1E-09	1.0	96
1550C05	GHYPMHV	PRRY	YYNEADYSQI	6.59E+04	1.35E-04	2.1E-09	16.8	97
1550 E03	GHYPLHI	RMRH	YYSEELYKYI	9.95E+04	1.86E-04	1.9E-09	5.3	98
1550 E06	GHYPMHV	PPRH	YYAQENYKEI	5.54E+04	1.16E-04	2.1E-09	0.4	99
1550H05	GHYPLHV	RQIY	YYNEADYSQI	6.52E+04	2.12E-04	3.3E-09	1.5	100

***Method for affinity determinations:** The anti-His antibody, mAb050 (RnD Systems, MN) was diluted to 20 ug/mL in acetate 5.0 and immobilized to ~9000 RU on flow cells 1 and 2 of a CM5 chip surface (GE Healthcare, Piscataway, NJ) according to the manufacturer's instructions. All surface plasmon experiments were conducted in HBS-EP (10 mM Hepes 150 mM NaCl 3 mM EDTA 0.05% Surfactant P20) at 25 °C. IL-23 was injected over anti-His mAb captured Adnectins for 2 minutes followed by a 10 minute dissociation phase.

Evaluation of the binding specificity was completed using Biacore T100 evaluation software. Additional detailed methods are described in Example 4.

[0082] Additional anti-IL-23 Adnectin characterization is described in Table 2.

Table 2: Anti-IL-23 Adnectin IC50/EC50

clone ID	BC	DE	FG	PBMNC pSTAT3 IC50 (nM)	IL-17 EC50 (nM)	IL-22 EC50 (nM)
1571G04	GHYPLHV	PRSH	YYAQENYKEI	0.23 ± .05	1.4 ± 0.3	1.3 ± 0.7
1490B03	GHYPLHV	SRKY	YYKEANYREI	.09 ± .01	1.4 ± 0.1	1.7 ± 0.8
1572G06	GHYPLHV	RYKY	YYAQENYKEI	.21 ± .03	1.6 ± 0.3	1.9 ± 0.3
1550E06	GHYPMHV	PPRH	YYAQENYKEI	1.15 ± .5	1.5 ± 0.6	2.1 ± 0.8
1571H03	GHYPLHV	NRVY	YYAQEEYHII	n.d.	2.9 ± 0.8	2.3 ± 0.4
1490H05	GHYPLHV	MRQH	YYAQENYKEI	n.d.	1.8 ± 2.1	2.9 ± 1.0
1571G06	GHYPLHL	RTKY	YYKEADYSQI	0.93 ± .5	3.5 ± 1.4	5.1 ± 4.3
1572C09	GHYPMHI	TPKH	YYNEADYSQI	n.d.	7.9 ± 6.1	5.3 ± 4.5

(n.d. not determined) (Detailed methods described in Example 4).

5

Nucleic Acid-Protein Fusion Technology

[0083] In one aspect, the application provides Adnectin comprising fibronectin type III domains that bind p19 subunit of IL-23. One way to rapidly make and test Fn3 domains with specific binding properties is the nucleic acid-protein fusion technology of Adnexus, a Bristol-Myers Squibb R&D Company. This disclosure utilizes the *in vitro* expression and tagging technology, termed PROfusion, which exploits nucleic acid-protein fusions (RNA- and DNA-protein fusions) to identify novel polypeptides and amino acid motifs that are important for binding to proteins. Nucleic acid-protein fusion technology is a technology that covalently couples a protein to its encoding genetic information. For a detailed description of the RNA-protein fusion technology and fibronectin-based scaffold protein library screening methods see .Szostak et al., U.S. Patent Nos. 6,258,558, 6,261,804, 6,214,553, 6,281,344, 6,207,446, 6,518,018, 6,818,418; and Roberts et al., *Proc. Natl., Acad. Sci.*, 94:12297-12302 (1997), herein incorporated by reference.

20

Vectors and Polynucleotides Embodiments

[0084] Nucleic acids encoding any of the various proteins or polypeptides disclosed herein may be synthesized chemically. Codon usage may be selected so as to improve expression in a cell. Such codon usage will depend on the cell type selected. Specialized
5 codon usage patterns have been developed for *E. coli* and other bacteria, as well as mammalian cells, plant cells, yeast cells and insect cells. See for example: Mayfield et al., *Proc. Natl. Acad. Sci. USA*, 100(2):438-442 (Jan. 21, 2003); Sinclair et al., *Protein Expr. Purif.*, 26(1):96-105 (Oct. 2002); Connell, N.D., *Curr. Opin. Biotechnol.*, 12(5):446-449 (Oct. 2001); Makrides et al., *Microbiol. Rev.*, 60(3):512-538 (Sept. 1996); and Sharp et
10 al., *Yeast*, 7(7):657-678 (Oct. 1991).

[0085] General techniques for nucleic acid manipulation are described for example in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Vols. 1-3, Cold Spring Harbor Laboratory Press, publ. (1989), or Ausubel, F. et al., *Current Protocols in Molecular Biology*, Green Publishing and Wiley-Interscience, New York, publ. (1987)
15 and periodic updates, herein incorporated by reference. Generally, the DNA encoding the polypeptide is operably linked to suitable transcriptional or translational regulatory elements derived from mammalian, viral, or insect genes. Such regulatory elements include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and sequences that control
20 the termination of transcription and translation. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene, to facilitate recognition of transformants, are additionally incorporated.

[0086] The proteins described herein may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which is
25 preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. The heterologous signal sequence selected preferably is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell.

[0087] For prokaryotic host cells that do not recognize and process a native signal
30 sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders.

[0088] For yeast secretion the native signal sequence may be substituted by, *e.g.*, the yeast invertase leader, a factor leader (including *Saccharomyces* and *Kluyveromyces* alpha-factor leaders), or acid phosphatase leader, the *C. albicans* glucoamylase leader, or the signal described in U.S. Patent 5,631,144. In mammalian cell expression, mammalian
5 signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available. The DNA for such precursor regions may be ligated in reading frame to DNA encoding the protein.

[0089] Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning
10 vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 micron plasmid origin is suitable for yeast, and various viral origins
15 (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

[0090] Expression and cloning vectors may contain a selection gene, also termed a
20 selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*.

[0091] Expression and cloning vectors usually contain a promoter that is recognized
25 by the host organism and is operably linked to the nucleic acid encoding the protein of the invention, *e.g.*, a fibronectin-based scaffold protein. Promoters suitable for use with prokaryotic hosts include the *phoA* promoter, beta-lactamase and lactose promoter systems, alkaline phosphatase, a tryptophan (*trp*) promoter system, and hybrid promoters such as the *tan* promoter. However, other known bacterial promoters are suitable.
30 Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the protein of the invention. Promoter sequences are also known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region

located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

[0092] Examples of suitable promoter sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

[0093] Transcription from vectors in mammalian host cells can be controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter, from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[0094] Transcription of a DNA encoding proteins of the invention by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, .alpha.-fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, *Nature*, 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the peptide-encoding sequence, but is preferably located at a site 5' from the promoter.

[0095] Expression vectors used in eukaryotic host cells (*e.g.*, yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3',

untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of mRNA encoding the protein of the invention. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO 94/11026 and
5 the expression vector disclosed therein.

[0096] The recombinant DNA can also include any type of protein tag sequence that may be useful for purifying the protein. Examples of protein tags include but are not limited to a histidine tag, a FLAG tag, a myc tag, an HA tag, or a GST tag. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian
10 cellular hosts can be found in *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, publ. (1985), the relevant disclosure of which is hereby incorporated by reference.

[0097] The expression construct is introduced into the host cell using a method appropriate to the host cell, as will be apparent to one of skill in the art. A variety of methods for introducing nucleic acids into host cells are known in the art, including, but
15 not limited to, electroporation; transfection employing calcium chloride, rubidium chloride, calcium phosphate, DEAE-dextran, or other substances; microprojectile bombardment; lipofection; and infection (where the vector is an infectious agent).

[0098] Suitable host cells include prokaryotes, yeast, mammalian cells, or bacterial cells. Suitable bacteria include gram negative or gram positive organisms, for example, *E. coli* or *Bacillus spp.* Yeast, preferably from the *Saccharomyces* genus, such as *S. cerevisiae*, may also be used for production of polypeptides. Various mammalian or
20 insect cell culture systems can also be employed to express recombinant proteins. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow et al. (*Bio/Technology*, 6:47 (1988)). Examples of suitable mammalian host
25 cell lines include endothelial cells, CO8-7 monkey kidney cells, CV-1, L cells, C127, 3T3, Chinese hamster ovary (CHO), human embryonic kidney cells, HeLa, 293, 293T, and BHK cell lines. Purified polypeptides are prepared by culturing suitable host/vector systems to express the recombinant proteins. For many applications, the small size of
30 many of the polypeptides disclosed herein would make expression in *E. coli* the preferred method for expression. The protein is then purified from culture media or cell extracts.

Protein Production

[0099] Host cells are transformed with the herein-described expression or cloning vectors for protein production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. In the examples shown here, the host cells used for high-throughput protein production (HTPP) and mid-scale production was the BL21 DE3
5 plysS-bacterial strain. The host cells used to produce the proteins of this invention may be cultured in a variety of media, such as those described in Ham et al., *Meth. Enzymol.*, 58:44 (1979), Barites et al., *Anal. Biochem.*, 102:255 (1980), U.S. Patent Nos. 4,767,704, 4,657,866, 4,927,762, 4,560,655, 5,122,469, 6,048,728, 5,672,502, or U.S. Patent No.
10 RE30,985. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

[00100] Proteins disclosed herein can also be produced using cell-translation systems.
15 For such purposes the nucleic acids encoding the polypeptide must be modified to allow *in vitro* transcription to produce mRNA and to allow cell-free translation of the mRNA in the particular cell-free system being utilized (eukaryotic such as a mammalian or yeast cell-free translation system or prokaryotic such as a bacterial cell-free translation system).

[00101] Proteins of the invention can also be produced by chemical synthesis (*e.g.*, by
20 the methods described in *Solid Phase Peptide Synthesis*, 2nd Edition, The Pierce Chemical Co., Rockford, IL, publ. (1984). Modifications to the protein can also be produced by chemical synthesis.

[00102] The proteins of the present invention can be purified by isolation/purification methods for proteins generally known in the field of protein chemistry. Non-limiting
25 examples include extraction, recrystallization, salting out (*e.g.*, with ammonium sulfate or sodium sulfate), centrifugation, dialysis, ultrafiltration, adsorption chromatography, ion exchange chromatography, hydrophobic chromatography, normal phase chromatography, reversed-phase chromatography, gel filtration, gel permeation chromatography, affinity chromatography, electrophoresis, countercurrent distribution or any combinations of
30 these. After purification, polypeptides may be exchanged into different buffers and/or concentrated by any of a variety of methods known to the art, including, but not limited to, filtration and dialysis.

[00103] The purified polypeptide is preferably at least 85% pure, or preferably at least 95% pure, and most preferably at least 98% pure. Regardless of the exact numerical value of the purity, the polypeptide is sufficiently pure for use as a pharmaceutical product.

[00104] A platform manufacturing process was used to prepare anti-IL-23 Adnectin.

5 Example 1 describes an example of the manufacturing process. The Adnectin is produced in *Escherichia coli* (*E. coli*). *E. coli* MG1655 cells were transformed with expression vector (pBMS2008/ATI001044) which produces the protein in an insoluble form as inclusion bodies. The recombinant strain is grown in stirred tank fermentors. At the end of fermentation the inclusion bodies are collected, solubilized, and refolded in preparation
10 for purification. The purified Adnectin is conjugated to a 40 kDa branched methoxyPEG using a maleimide linker. The conjugated material is subsequently repurified to remove free PEG, free Adnectin and product related impurities. Quality control testing is performed on the bulk drug substance.

15 Therapeutic *In Vivo* Uses

[00105] In one aspect, the application provides anti-IL-23 Adnectin useful in the treatment of autoimmune diseases such as lupus (*e.g.*, lupus erythematosus, lupus nephritis), Hashimoto's thyroiditis, primary myxedema, Graves' disease, pernicious anemia, autoimmune atrophic gastritis, Addison's disease, diabetes (*e.g.*, insulin
20 dependent diabetes mellitus, type I diabetes mellitus), Goodpasture's syndrome, myasthenia gravis, pemphigus, Crohn's disease, sympathetic ophthalmia, autoimmune uveitis, multiple sclerosis, autoimmune hemolytic anemia, idiopathic thrombocytopenia, primary biliary cirrhosis, chronic action hepatitis, ulceratis colitis, Sjögren's syndrome, rheumatic diseases (*e.g.*, rheumatoid arthritis), polymyositis, scleroderma, and mixed
25 connective tissue disease.

[00106] The application also provides methods for administering anti-IL-23 Adnectins to a subject. In some embodiments, the subject is a human. In some embodiments, the anti-IL-23 Adnectins are pharmaceutically acceptable to a mammal, in particular a human. A "pharmaceutically acceptable" polypeptide refers to a polypeptide that is
30 administered to an animal without significant adverse medical consequences, such as essentially endotoxin free or having very low endotoxin levels.

Formulation and Administration

[00107] The application further provides pharmaceutically acceptable compositions comprising the anti-IL-23 Adnectin described herein, wherein the composition is essentially endotoxin free. Therapeutic formulations comprising anti-IL-23 Adnectin are prepared for storage by mixing the described Adnectin having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (Osol, A., ed., *Remington's Pharmaceutical Sciences*, 16th Edition (1980)), in the form of aqueous solutions, lyophilized or other dried formulations. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as Tween, PLURONIC® or polyethylene glycol (PEG).

[00108] The formulations herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[00109] The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[00110] The skilled artisan will understand that the dosage of each therapeutic agent will be dependent on the identity of the agent.

[00111] For therapeutic applications, the anti-IL-23 Adnectin is administered to a subject, in a pharmaceutically acceptable dosage form. It can be administered

intravenously as a bolus or by continuous infusion over a period of time, or by subcutaneous routes. Suitable pharmaceutically acceptable carriers, diluents, and excipients are well known and can be determined by those of skill in the art as the clinical situation warrants. Examples of suitable carriers, diluents and/or excipients include: (1) 5 Dulbecco's phosphate buffered saline, (2) 0.9% saline (0.9% w/v NaCl), and (3) 5% (w/v) dextrose.

[00112] The method of the present invention can be practiced *in vitro*, *in vivo*, or *ex vivo*.

[00113] Administration of anti-IL-23 Adnectin, and one or more additional therapeutic 10 agents, whether co-administered or administered sequentially, may occur as described above for therapeutic applications. Suitable pharmaceutically acceptable carriers, diluents, and excipients for co-administration will be understood by the skilled artisan to depend on the identity of the particular therapeutic agent being administered.

[00114] When present in an aqueous dosage form, rather than being lyophilized, the 15 protein typically will be formulated at a concentration of about 0.1 mg/ml to 100 mg/ml, although wide variation outside of these ranges is permitted. For the treatment of disease, the appropriate dosage of anti-IL-23 Adnectin will depend on the type of disease to be treated, the severity and course of the disease, whether the Adnectin is administered for preventive or therapeutic purposes, the course of previous therapy, the patient's clinical 20 history and response to the Adnectin, and the discretion of the attending physician. The protein is suitably administered to the patient at one time or over a series of treatments.

Fusions of Serum Albumin Binding Adnectin (SABA)

[00115] In certain aspects, the application provides fusion proteins comprising anti- 25 IL23-Adnectin fused to a ¹⁰F_n3 domains that binds to human serum albumin (a Serum Albumin Binding Adnectin (¹⁰F_n3 domain) or SABA). Such fusion proteins have extended serum half lives in the presence of albumin relative to anti-IL23-Adnectin alone.

[00116] In certain aspects, the application provides fusion proteins comprising ¹⁰F_n3 30 domains that bind specifically to serum albumin, *e.g.*, human serum albumin (HSA) to prolong the $t_{1/2}$ of the fusion protein.

[00117] In certain embodiments, the serum half-life of the anti-IL23-Adnectin fused to the SABA is increased relative to the serum half-life of the anti-IL23-Adnectin when not

conjugated to the SABA. In certain embodiments, the serum half-life of the SABA fusion is at least 20, 40, 60, 80, 100, 120, 150, 180, 200, 400, 600, 800, 1000, 1200, 1500, 1800, 1900, 2000, 2500, or 3000% longer relative to the serum half-life of the anti-IL23-Adnectin when not fused to the SABA. In other embodiments, the serum half-life of the SABA fusion is at least 1.5-fold, 2-fold, 2.5-fold, 3-fold, 3.5 fold, 4-fold, 4.5-fold, 5-fold, 6-fold, 7-fold, 8-fold, 10-fold, 12-fold, 13-fold, 15-fold, 17-fold, 20-fold, 22-fold, 25-fold, 27-fold, 30-fold, 35-fold, 40-fold, or 50-fold greater than the serum half-life of the anti-IL23-Adnectin when not fused to the SABA. In some embodiments, the serum half-life of the SABA fusion is at least 10 hours, 15 hours, 20 hours, 25 hours, 30 hours, 35 hours, 40 hours, 50 hours, 60 hours, 70 hours, 80 hours, 90 hours, 100 hours, 110 hours, 120 hours, 130 hours, 135 hours, 140 hours, 150 hours, 160 hours, or 200 hours.

[00118] Accordingly, the SABA fusion molecules described herein are useful for increasing the half-life of anti-IL23-Adnectin by creating a fusion between anti-IL23-Adnectin and the SABA. Such fusion molecules may be used to treat conditions which respond to the biological activity of IL23. The present invention contemplates the use of the SABA fusion molecules in diseases caused by the dysregulation of IL-23.

[00119] The fusion may be formed by attaching anti-IL23-Adnectin to either end of the SABA molecule, *i.e.*, SABA-anti-IL23-Adnectin or anti-IL23-Adnectin-SABA arrangements.

[00120] In one aspect, the disclosure provides fusion proteins comprising anti-IL23-Adnectin comprising a serum albumin binding ¹⁰F_n3 domain. In exemplary embodiments, the serum albumin binding ¹⁰F_n3 proteins described herein bind to HSA with a K_D of less than 3 uM, 2.5 uM, 2 uM, 1.5 uM, 1 uM, 500 nM, 100 nM, 50 nM, 10 nM, 1 nM, 500 pM, 200 pM, 100 pM, 50 pM or 10 pM. In certain embodiments, the serum albumin binding ¹⁰F_n3 proteins described herein bind to HSA with a K_D of less than 3 uM, 2.5 uM, 2 uM, 1.5 uM, 1 uM, 500 nM, 100 nM, 50 nM, 10 nM, 1 nM, 500 pM, 200 pM, 100 pM, 50 pM or 10 pM at a pH range of 5.5 to 7.4 at 25 °C or 37 °C. In some embodiments, the serum albumin binding ¹⁰F_n3 proteins described herein bind more tightly to HSA at a pH less than 7.4 as compared to the binding affinity for HSA at a pH of 7.4 or greater.

[00121] In certain embodiments, the fusion proteins comprising HSA binding ¹⁰F_n3 domains described herein may also bind serum albumin from one or more of monkey, rat,

or mouse. In certain embodiments, the serum albumin binding ¹⁰F_n3 proteins described herein bind to rhesus serum albumin (RhSA) or cynomolgus monkey serum albumin (CySA) with a K_D of less than 3 uM, 2.5 uM, 2 uM, 1.5 uM, 1 uM, 500 nM, 100 nM, 50 nM, 10 nM, 1 nM, 500 pM or 100 pM.

5 **[00122]** In certain embodiments, the fusion proteins comprising serum albumin binding ¹⁰F_n3 domains described herein bind to domain I and/or domain II of HSA. In one embodiment, the fusion proteins comprising serum albumin binding ¹⁰F_n3 domains described herein do not bind to domain III of HSA.

[00123] In certain embodiments, the serum albumin binding ¹⁰F_n3 (SABA) portion of
10 the fusion proteins comprises a sequence having at least 40%, 50%, 60%, 70%, 75%, 80% or 85% identity to the wild-type ¹⁰F_n3 domain (SEQ ID NO: 1). In one embodiment, at least one of the BC, DE, or FG loops is modified relative to the wild-type ¹⁰F_n3 domain. In another embodiment, at least two of the BC, DE, or FG loops are modified relative to the wild-type ¹⁰F_n3 domain. In another embodiment, all three of the
15 BC, DE, and FG loops are modified relative to the wild-type ¹⁰F_n3 domain. In other embodiments, a SABA comprises a sequence having at least 40%, 50%, 60%, 70%, 75%, 80%, 85%, 90%, or 95% identity to any one of the 26 core SABA sequences shown in Table 3 (*i.e.*, SEQ ID NO: 103, 107, 111, 115, 119, and 123-143) or any one of the extended SABA sequences shown in Table 3 (*i.e.*, SEQ ID NO: 188-215, minus the
20 6xHIS tag).

[00124] In certain embodiments, the core amino acid residues are fixed and any substitutions, conservative substitutions, deletions or additions occur at residues other than the core amino acid residues. In exemplary embodiments, the BC, DE, and FG loops are replaced with polypeptides comprising the BC, DE and FG loop sequences from any
25 of the HSA binders shown in Table 3 below (*i.e.*, SEQ ID NOs: 103, 107, 111, 115, 119, and 123-143 in Table 3).

[00125] In certain embodiments, a SABA (*e.g.*, a SABA core sequence or a sequence based thereon as described above) may be modified to comprise an N-terminal extension sequence and/or a C-terminal extension sequence. Exemplary extension sequences are
30 shown in Table 3. For example, SEQ ID NO: 188 designated as SABA1.1 comprises the core SABA 1 sequence (SEQ ID NO: 103) with an N-terminal sequence MGVSDVPRDLE (SEQ ID NO: 144, designated as AdNT1), and a C-terminal sequence

EIDKPSQ (SEQ ID NO: 153). SABA1.1 further comprises a His6 tag at the C-terminus, however, it should be understood that the His6 tag is completely optional and may be placed anywhere within the N- or C-terminal extension sequences. Further, any of the exemplary N- or C-terminal extension sequences provided in Table 3 (SEQ ID NO: 144-163), and any variants thereof, can be used to modify any given SABA core sequence provided in Table 3.

[00126] In other embodiments, the tail sequences may be combined with other known linker sequences (*e.g.*, SEQ ID NO: 164-187 in Table 3) as necessary when designing a SABA fusion molecule.

10

Conjugation Linkers

[00127] SABA fusions may be covalently or non-covalently linked. In some embodiments, a serum albumin binding ¹⁰F_n3 may be directly or indirectly linked to an anti-IL23-Adnectin via a polypeptide linker. Suitable linkers for joining F_n3 are those which allow the separate domains to fold independently of each other forming a three dimensional structure that permits high affinity binding to a target molecule.

[00128] The disclosure provides a number of suitable linkers that meet these requirements, including glycine-serine based linkers, glycine-proline based linkers, as well as the linker having the amino acid sequence PSTSTST (SEQ ID NO: 184). The Examples described herein demonstrate that F_n3 domains joined via polypeptide linkers retain their target binding function. In some embodiments, the linker is a glycine-serine based linker. These linkers comprise glycine and serine residues and may be between 8 and 50, 10 and 30, and 10 and 20 amino acids in length. Examples include linkers having an amino acid sequence (GS)₇ (SEQ ID NO: 171), G(GS)₆ (SEQ ID NO: 166), and G(GS)₇G (SEQ ID NO: 168). Other linkers contain glutamic acid, and include, for example, (GSE)₅ (SEQ ID NO: 173) and GGSE GGSE (SEQ ID NO: 177). Other exemplary glycine-serine linkers include (GS)₄ (SEQ ID NO: 170), (GGGS)₇ (SEQ ID NO: 179), (GGGS)₅ (SEQ ID NO: 180), and (GGGS)₃G (SEQ ID NO: 181). In some embodiments, the linker is a glycine-proline based linker. These linkers comprise glycine and proline residues and may be between 3 and 30, 10 and 30, and 3 and 20 amino acids in length. Examples include linkers having an amino acid sequence (GP)₃G (SEQ ID NO: 182) and (GP)₅G (SEQ ID NO: 183). In other embodiments, the linker may be a

proline-alanine based linker having between 3 and 30, 10 and 30, and 3 and 20 amino acids in length. Examples of proline alanine based linkers include, for example, (PA)₃ (SEQ ID NO: 185), (PA)₆ (SEQ ID NO: 186) and (PA)₉ (SEQ ID NO: 187). It is contemplated, that the optimal linker length and amino acid composition may be determined by routine experimentation by methods well known in the art.

[00129] In some embodiments, the fusions described herein are linked via a polypeptide linker having a protease site that is cleavable by a protease in the blood or target tissue. Such embodiments can be used to release a therapeutic protein for better delivery or therapeutic properties or more efficient production.

[00130] Additional linkers or spacers, may be introduced at the C-terminus of a Fn3 domain between the Fn3 domain and the polypeptide linker. Additional linkers or spacers may be introduced at the N-terminus of a Fn3 domain between the Fn3 domain and the polypeptide linker.

[00131] In some embodiments, a therapeutic moiety may be directly or indirectly linked to a SABA via a polymeric linker. Polymeric linkers can be used to optimally vary the distance between each component of the fusion to create a protein fusion with one or more of the following characteristics: 1) reduced or increased steric hindrance of binding of one or more protein domains when binding to a protein of interest, 2) increased protein stability or solubility, 3) decreased protein aggregation, and 4) increased overall avidity or affinity of the protein.

[00132] In some embodiments, a therapeutic moiety is linked to a SABA via a biocompatible polymer such as a polymeric sugar. The polymeric sugar can include an enzymatic cleavage site that is cleavable by an enzyme in the blood or target tissue. Such embodiments can be used to release a therapeutic proteins for better delivery or therapeutic properties or more efficient production.

Summary of Serum Albumin-Binding Adnectins (SABA) Sequences

[00133] Many of the SABA sequences referenced in this application are summarized in Table 3 below. Unless otherwise specified, all N-terminal extensions are indicated with a single underline, all C-terminal tails/extensions are indicated with a double underline, and linker sequences are boxed. Loop regions BC, DE and FG are shaded for each core SABA sequence.

Table 3: Summary of SABA Exemplary Sequences

SEQ ID NO:	Sequence Name	Description	Sequence
103	SABA1	Core 1 Adnectin	EVVAATPTSL LISWHSY EQNSYYRITYGE TGGNSPVQEF TVPSQT TATISGLKPGVDY TITVYAV YGSKYYY PISINYRT
104	SABA1BC	Core 1 BC Loop	HSY EQNS
105	SABA1DE	Core 1 DE Loop	YSQT
106	SABA1FG	Core 1 FG Loop	YGSKYYY
107	SABA2	Core 2 Adnectin	EVVAATPTSL LISWPKYDKTGH YYRITYGE TGGNSPVQEF TVPTRQT TATISGLKPGVDY TITVYAV SKDDYYPHEHR PISINYRT
108	SABA2BC	Core 2 BC Loop	PKYDKTGH
109	SABA2DE	Core 2 DE Loop	TRQT
110	SABA2FG	Core 2 FG Loop	SKDDYYPHEHR
111	SABA3	Core 3 Adnectin	EVVAATPTSL LISWSNDGPGLS YYRITYGE TGGNSPVQEF TVPSST TATISGLKPGVDY TITVYAV SYTTKKAYSAG PISINYRT
112	SABA3BC	Core 3 BC Loop	SNDGPGLS
113	SABA3DE	Core 3 DE Loop	SSQT
114	SABA3FG	Core 3 FG Loop	SYTTKKAYSAG
115	SABA4	Core 4 Adnectin; contains a scaffold mutation (bolded); scaffold-perfect version is SABA5	EMVAATPTSL LISWEDDSYYSR YYRITYGE TGGNSPVQEF TVPSDLY TATISGLKPGVDY TITVYAV TYDVTDLIMHE PISINYRT
116	SABA4BC	Core 4 BC Loop	EDDSYYSR
117	SABA4DE	Core 4 DE Loop	SDLY
118	SABA4FG	Core 4 FG Loop	YDVTDLIMHE

SEQ ID NO:	Sequence Name	Description	Sequence
119	SABA5	Core 5 Adnectin; see description for SAVA4; corrected residue is bolded	EVVAATPTSLLLISW E DDSYYSRYYRITYGE TGGNSPVQEFTVPS D LYTATISGLKPGVDY TITVYAVTYD V TDLIMHEPISINYRT
120	SABA5BC	Core 5 BC Loop	EDDSYYSR
121	SABA5DE	Core 5 DE Loop	SDLY
122	SABA5FG	Core 5 FG Loop	YDVTDLIMHE
123	SABA6	Core 6 Adnectin	EVVAATPTSLLLISW M DEYDVRYYRITYGE TGGNSPVQEFTV P NYNTATISGLKPGVDY TITVYAV T RIKANNMYGPISINYRT
124	SABA7	Core 7 Adnectin	EVVAATPTSLLLISW N HLEHVARYYRITYGE TGGNSPVQEFTV P EYPTTATISGLKPGVDY TITVYAV T ITMLKYPTQSPISINYRT
125	SABA8	Core 8 Adnectin	EVVAATPTSLLLISW G HYRPSGHYYRITYGE TGGNSPVQEFTV D PSSYTATISGLKPGVDY TITVYAV S KDDYYPHEHRPISINYRT
126	SABA9	Core 9 Adnectin	EVVAATPTSLLLISW D ASHYERRYYRITYGE TGGNSPVQEFTV P RYHHTATISGLKPGVDY TITVYAV T QAEHYQPPISINYRT
127	SABA10	Core 10 Adnectin	EVVAATPTSLLLISW N SYHHSADYYRITYGE TGGNSPVQEFTV P YPPTTATISGLKPGVDY TITVYAV S AKSYYPISINYRT
128	SABA11	Core 11 Adnectin	EVVAATPTSLLLISW S KYSKHGHYYRITYGE TGGNSPVQEFTV P SGNATATISGLKPGVDY TITVYAV E DTNDYPHTRPISINYRT
129	SABA12	Core 12 Adnectin	EVVAATPTSLLLISW H GEPDQTRYRITYGE TGGNSPVQEFTV P YRRATATISGLKPGVDY TITVYAV T SGYTGHYQPISINYRT

SEQ ID NO:	Sequence Name	Description	Sequence
130	SABA13	Core 13 Adnectin	EVVAATPTSLLISW SKYSKHHG HYRITYGE TGGNSPVQEFV DPSSY TATISGLKPGVDY TITVYAV SKDDYPHEHR PISINVRT
131	SABA14	Core 14 Adnectin	EVVAATPTSLLISW YEPYTPH HYRITYGE TGGNSPVQEFV PGYYG TATISGLKPGVDY TITVYAV YGYQYTP PISINVRT
132	SABA15	Core 15 Adnectin	EVVAATPTSLLISW SKYSKHHG HYRITYGE TGGNSPVQEFV PSGNA TATISGLKPGVDY TITVYAV SDDNKYHQHR PISINVRT
133	SABA16	Core 16 Adnectin	EVVAATPTSLLISW GHYRRSGH HYRITYGE TGGNSPVQEFV DPSSY TATISGLKPGVDY TITVYAV SKDDYPHEHR PISINVRT
134	SABA17	Core 17 Adnectin	EVVAATPTSLLISW SKYSKHHG HYRITYGE TGGNSPVQEFV PSGNA TATISGLKPGVDY TITVYAV EDTNDYPHTR PISINVRT
135	SABA18	Core 18 Adnectin	EVVAATPTSLLISW YEPGASV YYYRITYGE TGGNSPVQEFV PSYYHT TATISGLKPGVDY TITVYAV YGYEYEP PISINVRT
136	SABA19	Core 19 Adnectin	EVVAATPTSLLISW QSYAHSD YYRITYGE TGGNSPVQEFV YPPT TATISGLKPGVDY TITVYAV YAGSSYP PISINVRT
137	SABA20	Core 20 Adnectin	EVVAATPTSLLISW GHYRRSGH HYRITYGE TGGNSPVQEFV DPSSY TATISGLKPGVDY TITVYAV SKDDYPHEHR PISINVRT
138	SABA21	Core 21 Adnectin	EVVAATPTSLLISW PEPGTPV YYYRITYGE TGGNSPVQEFV PAYYG TATISGLKPGVDY TITVYAV YGYDYS PISINVRT
139	SABA22	Core 22 Adnectin	EVVAATPTSLLISW YRYEKTQ HYRITYGE TGGNSPVQEFV PESGT TATISGLKPGVDY TITVYAV YAGYEYPHTR PISINVRT

SEQ ID NO:	Sequence Name	Description	Sequence
140	SABA23	Core 23 Adnectin	EVVAATPTSLLISWVKSEEEYRYRITYGE TGGNSPVQEFTVPPYYVHTATISGLKPGVDY TITVYAVTEYYYAGAVVSVPI SINYRT
141	SABA24	Core 24 Adnectin	EVVAATPTSLLISWYDPYTYGSYYRITYGE TGGNSPVQEFTVGPYTTTATISGLKPGVDY TITVYAVSYYYSTQPI SINYRT
142	SABA25	Core 25 Adnectin	EVVAATPTSLLISWSDNGPGLSYYRITYGE TGGNSPVQEFTVPSSTTATISGLKPGVDY TITVYAVSYTTKKAYSAGPI SINYRT
143	SABA26	Core 26 Adnectin	EVVAATPTSLLISWPDFYYKPDYYRITYGE TGGNSPVQEFTVPRDYTTATISGLKPGVDY TITVYAVYSYYGYPI SINYRT
Exemplary Adnectin N-Terminal Extension Sequences			
144	AdNT1	Exemplary leader	MGVSDVPRDL
145	AdNT2	Exemplary leader	GVSDVPRDL
146	AdNT3	Exemplary leader	VSDVPRDL
147	AdNT4	Exemplary leader	SDVPRDL
148	AdNT5	Exemplary leader	DVPRDL
149	AdNT6	Exemplary leader	VPRDL
150	AdNT7	Exemplary leader	PRDL
151	AdNT8	Exemplary leader	RDL
152	AdNT9	Exemplary leader	DL
Exemplary Adnectin C-Terminal Extension Sequences			
153	AdCT1	Exemplary tail	EIDKPSQ
154	AdCT2	Exemplary tail	EIDKPS
155	AdCT3	Exemplary tail	EIDKPC
156	AdCT4	Exemplary tail	EIDKP
157	AdCT5	Exemplary tail	EIDK
158	AdCT6	Exemplary tail	EI

SEQ ID NO:	Sequence Name	Description	Sequence
159	AdCT7	Exemplary tail	EIEKPSQ
160	AdCT8	Exemplary tail	EIDKPSQLE
161	AdCT9	Exemplary tail	EIEDEDEDEDED
162	AdCT10	Exemplary tail	EIEKPSQEDEDEDEDED
163	AdCT11	Exemplary tail	EGSGS
164	L1	G(GS) ₂	GGSGS
165	L2	G(GS) ₄	GGSGSGSGS
166	L3	G(GS) ₆	GGSGSGSGSGSGS
167	L4	G(GS) ₇	GGSGSGSGSGSGSGS
168	L5	G(GS) ₇ G	GGSGSGSGSGSGSGSG
169	L6	GSGS	GSGS
170	L7	(GS) ₄	GSGSGSGS
171	L7	(GS) ₇	GSGSGSGSGSGSGS
172	L9	GS(A) ₉ GS	GSAAAAAAAAAAGS
173	L10	(GSE) ₅	GSEGSEGSEGSEGSE
174	L11	(PAS) ₅	PASPASPASPASPAS
175	L12	(GSP) ₅	GSPGSPGSPGSPGSP
176	L13	GS(TVAAPS) ₂	GSTVAAPSTVAAPS
177	L14	(GGSE) ₂	GGSEGGSE
178	L15	(ST) ₃ G	STSTSTG
179	L16	(GGGGS) ₇	GGGGSGGGSGGGSGGGSGGGSGGGSGGGGS GGGGS
180	L17	(GGGGS) ₅	GGGGSGGGSGGGSGGGSGGGSGGGSGGGGS
181	L18	(GGGGS) ₃ G	GGGGSGGGSGGGSG
182	L19	(GP) ₃ G	GPGPGPG
183	L20	(GP) ₅ G	GPGPGPGPGPG
184	L21	P(ST) ₃	PSTSTST
185	L22	(PA) ₃	PAPAPA

SEQ ID NO:	Sequence Name	Description	Sequence
186	L23	(PA) ₆	PAPAPAPAPAPA
187	L24	(PA) ₉	PAPAPAPAPAPAPAPAPA
Exemplary Extensions to Adnectin Core Sequences			
188	SABA1.1	Adnectin core 1 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWHSYYEQ</u> NSYYRITYGETGGNSPVQEFVTPYSQTTAT ISGLKPGVDYTITVYAVYGSKYYYPI SINY RTE <u>IDKPSQH</u> HHHHH
189	SABA1.2	Adnectin core 1 sequence having AdNT1 and AdCT8 terminal sequences	<u>MGVSDVPRDLEVVAATPTSLLISWHSYYEQ</u> NSYYRITYGETGGNSPVQEFVTPYSQTTAT ISGLKPGVDYTITVYAVYGSKYYYPI SINY RTE <u>IEDEDEDEDEDE</u>
190	SABA1.3	Adnectin core 1 sequence having AdNT1 and AdCT9 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWHSYYEQ</u> NSYYRITYGETGGNSPVQEFVTPYSQTTAT ISGLKPGVDYTITVYAVYGSKYYYPI SINY RTE <u>IEDEDEDEDEDE</u> HHHHH
191	SABA2.1	Adnectin core 2 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWPKYDKT</u> GHYYRITYGETGGNSPVQEFVTPTRQTTAT ISGLKPGVDYTITVYAVSKDDYYPHEHRPI SINYRTE <u>IDKPSQH</u> HHHHH
192	SABA3.1	Adnectin core 3 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWSNDGPG</u> LSYYRITYGETGGNSPVQEFVTPSSQTTAT ISGLKPGVDYTITVYAVSYTTKAYSAGPI SINYRTE <u>IDKPSQH</u> HHHHH

SEQ ID NO:	Sequence Name	Description	Sequence
193	SABA4.1	Adnectin core 4 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEMVAATPTSLLISWEDDSYY</u> SRYRITYGETGGNSPVQEFTVPSDLYTAT ISGLKPGVDYTTITVYAVTYDVTDLIMHEPI SINYRTE <u>IDKPSQ</u> HHHHHH
194	SABA5.1	Adnectin core 5 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWEDDSYY</u> SRYRITYGETGGNSPVQEFTVPSDLYTAT ISGLKPGVDYTTITVYAVTYDVTDLIMHEPI SINYRTE <u>IDKPSQ</u> HHHHHH
195	SABA6.1	Adnectin core 6 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWYMDEYD</u> VRYRITYGETGGNSPVQEFTVPNYYNTAT ISGLKPGVDYTTITVYAVTRIKANNMYGPI SINYRTE <u>IDKPSQ</u> HHHHHH
196	SABA7.1	Adnectin core 7 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWNHLEHV</u> ARYRITYGETGGNSPVQEFTVPEYPTTAT ISGLKPGVDYTTITVYAVTITMLKYPTQSPI SINYRTE <u>IDKPSQ</u> HHHHHH
197	SABA8.1	Adnectin core 8 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWGHYRRS</u> GHYYRITYGETGGNSPVQEFTVDPSSYTAT ISGLKPGVDYTTITVYAVSKDDYYPHEHRPI SINYRTE <u>IDKPSQ</u> HHHHHH

SEQ ID NO:	Sequence Name	Description	Sequence
198	SABA9.1	Adnectin core 9 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWDASHYE</u> RRYYRITYGETGGNSPVQEFTVPRYHHTAT ISGLKPGVDYTTITVYAVTQAQEHYQPPISI NYRTE <u>IDKPSQ</u> HHHHHH
199	SABA10.1	Adnectin core 10 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWNSYYHS</u> ADYYRITYGETGGNSPVQEFTVPYPPTTAT ISGLKPGVDYTTITVYAVYSAKSYPISINY RTE <u>IDKPSQ</u> HHHHHH
200	SABA11.1	Adnectin core 11 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWSKYSKH</u> GHYYRITYGETGGNSPVQEFTVPSGNATAT ISGLKPGVDYTTITVYAVEDTNDYPHTRPI SINYRTE <u>IDKPSQ</u> HHHHHH
201	SABA12.1	Adnectin core 12 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWHGEPDQ</u> TRYRITYGETGGNSPVQEFTVPPYRRATAT ISGLKPGVDYTTITVYAVTSGYTGHYQPISI NYRTE <u>IDKPSQ</u> HHHHHH
202	SABA13.1	Adnectin core 13 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWSKYSKH</u> GHYYRITYGETGGNSPVQEFTVDPSSYTAT ISGLKPGVDYTTITVYAVSKDDYYPHEHRPI SINYRTE <u>IDKPSQ</u> HHHHHH

SEQ ID NO:	Sequence Name	Description	Sequence
203	SABA14.1	Adnectin core 14 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWYEPYTP</u> IHYYRITYGETGGNSPVQEFTVPGYYGTAT ISGLKPGVDYTITVYAVYGYQYTPISINY RTE <u>IDKPSQH</u> HHHHH
204	SABA15.1	Adnectin core 15 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWSKYSKH</u> GHYYRITYGETGGNSPVQEFTVPSGNATAT ISGLKPGVDYTITVYAVSDDNKYYHQHRPI SINYRTE <u>IDKPSQH</u> HHHHH
205	SABA16.1	Adnectin core 16 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWGHYRRS</u> GHYYRITYGETGGNSPVQEFTVDPSSYTAT ISGLKPGVDYTITVYAVSKDDYYPHEHRPI SINYRTE <u>IDKPSQH</u> HHHHH
206	SABA17.1	Adnectin core 17 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWSKYSKH</u> GHYYRITYGETGGNSPVQEFTVPSGNATAT ISGLKPGVDYTITVYAVEDTNDYPHTHRPI SINYRTE <u>IDKPSQH</u> HHHHH
207	SABA18.1	Adnectin core 18 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWYEPGAS</u> VYYRITYGETGGNSPVQEFTVPSYYHTAT ISGLKPGVDYTITVYAVYGYEYEPISINY RTE <u>IDKPSQH</u> HHHHH

SEQ ID NO:	Sequence Name	Description	Sequence
208	SABA19.1	Adnectin core 19 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWQSYAH</u> SDYYRITYGETGGNSPVQEFTVPYPPQTAT ISGLKPGVDYTITVYAVYAGSSYPISINY RTE <u>IDKPSQH</u> HHHHH
209	SABA20.1	Adnectin core 20 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWGHYRRS</u> GHYYRITYGETGGNSPVQEFTVDPSSYTAT ISGLKPGVDYTITVYAVSKDDYPHEHRPI SINYRTE <u>IDKPSQH</u> HHHHH
210	SABA21.1	Adnectin core 21 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWPEPGTP</u> VYYRITYGETGGNSPVQEFTVPAYYGTAT ISGLKPGVDYTITVYAVYGYDYSPISINY RTE <u>IDKPSQH</u> HHHHH
211	SABA22.1	Adnectin core 22 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWRYEKT</u> QHYYRITYGETGGNSPVQEFTVPPESGTAT ISGLKPGVDYTITVYAVYAGYEPHTHRPI SINYRTE <u>IDKPSQH</u> HHHHH
212	SABA23.1	Adnectin core 23 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWVKSEY</u> YRYRITYGETGGNSPVQEFTVPYYVHTAT ISGLKPGVDYTITVYAVTEYYYAGAVVSP ISINYRTE <u>IDKPSQH</u> HHHHH

SEQ ID NO:	Sequence Name	Description	Sequence
213	SABA24.1	Adnectin core 24 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWYDPYTY</u> GSYYRITYGETGGNSPVQEFTVGPYTTTAT ISGLKPGVDYTITVYAVSYYYSTQPISINY <u>RTEIDKPSQH</u> HHHHH
214	SABA25.1	Adnectin core 25 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWSNDGPG</u> LSYYRITYGETGGNSPVQEFTVPSSQTTAT ISGLKPGVDYTITVYAVSYYTKKAYSAGPI SINYR <u>TEIDKPSQH</u> HHHHH
215	SABA26.1	Adnectin core 26 sequence having AdNT1 and AdCT1 terminal sequences with His6 tag	<u>MGVSDVPRDLEVVAATPTSLLISWPDPYK</u> PDYYRITYGETGGNSPVQEFTVPRDYTTAT ISGLKPGVDYTITVYAVYSYYGYYPISINY <u>RTEIDKPSQH</u> HHHHH

EXAMPLES

Example 1

5

Manufacturing Process

Fermentation and Harvest

[00134] A production fermentation is prepared with sterile basal medium. A vial is thawed and used to inoculate a transfer vessel containing growth medium. The inoculum is immediately transferred to the production fermentation. The culture is maintained at a temperature of 34 °C with agitation and allowed to grow to an OD₆₀₀ of 5-10 (one OD unit is approximately 1x10⁹ cells/mL) is reached. The addition of feed medium is initiated at this OD. The fermentation proceeds to OD₆₀₀ = 25 at which point the culture is induced to produce the Adnectin by the addition of isopropyl β-D-1-thiogalactopyranoside (IPTG). The temperature of the vessel is increased from 34 °C to

39 °C at the time of induction. Samples are taken aseptically every hour and tested for cell density.

5 [00135] After 9-12 hrs of induced fermentation the vessel is prepared for harvest by reducing the temperature to 25 °C, addition of ethylenediaminetetraacetic acid (EDTA) to a final concentration of 10 mM, pH increase to 7.8 by the addition of sodium hydroxide and reduction of agitation. After a one hour hold period the fermentor content is drained into a collection vessel.

Preparation of Inclusion Bodies

10 [00136] Cell disruption of the harvest pool is done by passing the material through a MICROFLUIDIZER® which disrupts the cells and releases their contents. Following cell disruption the inclusion bodies are collected using a disc stack centrifuge to separate solids and liquid phases in a continuous process by extremely high centrifugal forces. Inclusion bodies are then washed twice with buffer (20-25 °C) and twice with water (20-
15 25 °C). Each time the washed inclusion bodies are collected by centrifugation. The washed inclusion bodies are recovered as a slurry.

Solubilization of Inclusion Bodies and Protein Refolding

20 [00137] Solubilization buffer is added to the inclusion body slurry followed by stirring at room temperature for 1 hr. An $OD_{280}=20$ (total protein) is targeted during this process.

[00138] The protein refolding is performed using a two step dilution process. Dilution buffer is added to the solubilized inclusion bodies at a ratio of one part solubilized inclusion bodies to one half part dilution buffer (v/v). A second dilution is carried out by adding solubilized inclusion bodies to refold buffer to target an $OD_{280}=0.7$ (total protein).
25 The dilutions are carried out while stirring at room temperature. Following thorough mixing for one hour, the stirring is stopped and the protein solution is held at room temperature overnight. The solubilized and refolded Adnectin is passed through a 0.8 μm -0.22 μm filter and tested for protein content by A_{280} and RP-HPLC.

30 Purification and Conjugation to PEG

[00139] Refolded and filtered Adnectin is directly loaded onto a cation exchange (CEX1) column for initial capture. The bound material is washed with wash buffer and

eluted with 50 mM sodium acetate, 500 mM sodium chloride, 1.5% propylene glycol, pH 5.5. The eluate pool is assayed for purity, identity, concentration, and endotoxin.

5 [00140] The eluate from the capture chromatography is further purified using hydrophobic interaction chromatography (HIC). The CEX1 eluate is directly loaded on the HIC column, washed and subsequently eluted with 50 mM sodium acetate, 30% propylene glycol, pH 5.5. The eluate pool is assayed for purity, identity and concentration.

10 [00141] The purified Adenctin is then formatted directly with a maleimide derivative of a 40 kDa branched PEG (mPEG2-MAL). The HIC eluate is stirred at room temperature and the mPEG2-MAL is added. After 1 hr of mixing at room temperature, the reaction mixture is allowed to incubate overnight at the same temperature. The PEGylation solution is then processed on the final CEX column (CEX2). Samples are taken for protein content, purity and endotoxin.

15 [00142] The pH and conductivity of the PEGylation solution are adjusted to 4.0 and 1.0 mS/cm respectively, with 75 mM acetic acid prior to loading on the final cation exchange column (CEX2) for repurification. Once loaded, the bound material is washed with buffer and subsequently eluted with 50 mM sodium acetate, 25 mM sodium chloride, pH 5.0. Samples are taken for protein content, purity and endotoxin.

20 [00143] The CEX2 eluate is concentrated to 15 mg/mL in a tangential flow filtration unit equipped with a 30 kDa nominal molecular weight cut off membrane with a V-screen. The bulk drug substance in 50 mM sodium acetate, 25 mM sodium chloride, pH 5.0. is passed through a 0.22 μ m filter and frozen at -80 °C.

Example 2

25 Gene, Vector and Host Cell

[00144] A plasmid encoding the protein under the control of the T7 promoter was generated for use in strain construction. This plasmid DNA was used to transform competent *E. coli* K-12 MG1655 cells (F-lambda-, *ilvG-rfb-50 rph-1*). The host strain was designed to allow induction of expression from genes upon addition of IPTG. The transformed MG1655 strain is resistant to kanamycin. The protein expression vector is shown in Figure 2. A single colony selection from plates is used to inoculate a

30

fermentation culture which is then aliquoted and frozen away to be used as a research cell bank.

Example 3

5 Biophysical and Biochemical Characterization

[00145] The structure and quality of the protein of the invention were examined by several comprehensive analytical methods.

MALDI-MS

10 **[00146]** Mass spectral profiles were analyzed by MALDI. To evaluate precision of MALDI analysis on the samples, 20 individual spots were placed onto the steel plate for each sample and analyzed sequentially. A total of 20 spectra were generated.

Peptide Mapping

15 **[00147]** Peptide mapping was used to confirm correct expression of the amino acid sequence (primary structure) predicted from the cDNA sequence for the protein of the invention as well as the corresponding unPEGylated protein. In order to obtain complete sequence coverage, trypsin (cleavage to C-terminal side of Lys and Arg residues) and endoproteinase Glu-C (cleavage to C-terminal side of Glu residues) were employed to
20 yield two overlapping sets of peptide fragments. Peptide mapping was also used to determine covalent post-translational modifications including residual N-terminal methionine, disulfide-bridging, deamidation of asparagine, methionine oxidation (etc.). Peptides were identified and characterized by liquid chromatography mass spectrometry (LC-MS) via molecular weight and tandem mass spectrometry (MSMS) which provides
25 partial sequence information via collision-induced dissociation (CID).

SDS-PAGE

[00148] Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) was used to visualize molecular weight banding patterns of unPEGylated and PEGylated anit-
30 IL-23 Adnectins. The samples were prepared in a sample buffer with or without a reducing agent. After heating in SDS, the samples and molecular weight markers were electrophoretically analyzed on pre-cast, gradient (4-20%) polyacrylamide gels. After

electrophoresis, the gels were fixed and stained using Coomassie Blue. The equivalence of the banding patterns of samples was assessed visually.

Size-Exclusion Chromatography/Multi-Angle Light Scattering (SECMALS)

5 **[00149]** Size-exclusion chromatography (SEC) was used for the quantitative analysis of monomer, High Molecular Weight (HMW), and Low Molecular Weight (LMW) species. Following SEC separation, the molecular mass of separated species was determined by multi-angle light scattering in tandem with a differential refractometer.

10

Example 4

In Vitro Nonclinical Pharmacology

K_D by SPR

[00150] The binding characteristics were characterized by Surface Plasmon Resonance (SPR). Human IL-23 was immobilized at two to four levels in one dimension of a
15 ProteOn XPR (Bio-Rad) chip surfaces and exposed to 6 different concentrations of anti-IL-23 adnectins in the other dimension of the same SPR chip surface. This allowed kinetic determination in the absence of regeneration. Duplicate chips were used for kinetic determinations at 25 °C and 37 °C. Evaluation of the kinetic parameters was performed using the Langmuir interaction model and constant parameter fitting with the
20 ProteOn Manager software.

[00151] As shown in Table 4 below, the off-rates for these anti-IL-23 adnectins are slow (on the order of 10^{-5} s^{-1}) at 25 °C. Even at 37 °C the off rates were close to the limit of detection for SPR technologies so it is possible that the reported dissociation constant measurements are under-estimates.

25

Table 4: Kinetic Parameters of Anti-IL-23 Adnectin Against Directly Immobilized Human IL-23

Anti-IL-23 adnectin	Analysis temp (° C)	k_{on} ($M^{-1} s^{-1}$)	k_{off} (s^{-1})	K_D (nM)
1490B03	25	$2.8 \pm 0.6 \text{ E}+04$	$8.2 \pm 1.1 \text{ E}-06$	0.03 ± 0.002
1571G04	25	$5.7 \pm 0.6 \text{ E}+04$	$1.2 \pm 0.2 \text{ E}-05$	0.2 ± 0.05

Anti-IL-23 adnectin	Analysis temp (° C)	k_{on} ($M^{-1} s^{-1}$)	k_{off} (s^{-1})	K_D (nM)
ATI000934	25	9.4 E + 03	1.8 E-05	1.9
ATI001014	25	9.4 ± 0.2 E+03	1.7 ± 0.3 E-05	1.8 ± 0.2
ATI001047	25	1.3 ± 0.03 E+04	2 ± 0.1 E-05	1.6 ± 0.1
ATI001045	25	1.5 ± 0.2 E+05	2.5 ± 0.4 E-05	0.17 ± 0.01
ATI001045	37	2.03 ± 0.01 E+05	5.5 ± 0.6 E-05	0.27 ± 0.03

Solution Phase Affinity

[00152] The solution affinity of ATI001045 for human IL-23 was measured using a Kinetic Exclusion Assay (KinExA). In one format duplicate titrations of hIL-23 were performed for each of three concentrations. The relative unbound ATI001045 concentration was measured by capture on a human IL-23 solid matrix followed by detection with a fluorescently labeled antibody that recognizes the Adnectin scaffold. Due to technical limitations, the lowest concentration that could be tested was 0.75 nM. Hence, while the global K_D analysis shown in Table 5, gives an estimate of 51 pM for the K_D , the affinity could be as low as single digit pM or as high as 150 pM within a 95% confidence interval.

Table 5: Solution Phase Affinity Measurements for ATI001045

K_D	51 pM
95% confidence interval:	
K_D high	153 pM
K_D low	1 pM

[00153] The solution affinity of ATI001045 and ATI001047 for human IL-23 was also measured using an alternate format in the KinExA. Duplicate titrations of adnectins were performed for each of three (ATI001045) or single (ATI001047) concentrations of human IL-23 (quadruplicate for the lowest concentration). The relative unbound human IL-23 concentration was measured by capture on a non-PEGylated ATI001045 solid matrix followed by detection with a fluorescently labeled antibody that recognizes the p40

subunit of hIL-23. The global K_D analysis shown in Table 6 gives a K_D of 9.4 pM with a 95% confidence interval of 22 – 2.4 pM for ATI001045 and a K_D of 36.3 pM with a 95% confidence interval of 60.1 to 19.4 pM.

5

Table 6: Solution Phase Affinity Measurements

	ATI001045	ATI001047
K_D	9.4 pM	36.33 pM
95% confidence interval:		
K_D high	22 pM	60.07 pM
K_D low	2.4 pM	19.44 pM

STAT3 Phosphorylation on Kit225 Cells

[00154] Parham et al. (“A receptor for the heterodimeric cytokine IL-23 is composed of IL-12Rbeta1 and a novel cytokine receptor subunit, IL-23R”, *J. Immunol.*, 10 168(11):5699-5708 (Jun. 1, 2002)) cloned the IL-23R from the human IL-2 dependent T-cell line, Kit225. These cells have been characterized for expression of both IL-12RB1 and IL-23R by FACS analysis and responded to IL-23 by stimulation of pSTAT3 and to IL-12 by stimulation of pSTAT4. Kit225 cells were seeded into 96 well plates and quiesced in the absence of FBS and IL-2 for 3 hrs at 37 °C. Following this incubation, 15 human recombinant IL-23 (or IL-23 preincubated with antagonist for 1 hr) was applied and the cells returned to the incubator for 15 minutes at 37 °C to stimulate the phosphorylation of STAT3 (abbreviated as p-STAT3). Each condition was assayed in duplicate in 96-well plates. Stimulation was stopped by placing the cells on ice and addition of ice-cold PBS. Finally, the cells were pelleted and lysed following standard 20 protocols and pSTAT3 production detected by ELISA.

[00155] The optimal concentration of IL-23 for stimulation was 35 pM. Inhibition of the IL-23 induced pSTAT3 was demonstrated by a titration of anti-p40 monoclonal antibody (mAb1510) as well as an anti-p19 polyclonal antibody (AF1716). ATI001045, ATI001047, ATI001014 and ATI001016 had equivalent activity with an IC_{50} of ~300 25 pM, approximately 150 fold more potent than the anti-p19 polyclonal antibody while ATI001015 had an IC_{50} of ~1.2 nM, approximately 40 fold more potent than the anti-p19

polyclonal antibody. Adnectin ATI000934 is 1/3rd the potency of ATI001045, with an IC_{50} of 1 nM (Table 7).

Table 7: Inhibition of IL-23 Induced STAT3 Phosphorylation
by Anti-IL-23 Antagonists

5

	pSTAT3 $IC_{50} \pm SD$ (nM)
ATI001045	0.28 \pm 0.14
ATI001047	0.36
ATI001014	0.3 \pm 0.1
ATI001015	1.24
ATI001016	0.3 \pm 0.1
ATI000934	0.8 \pm 0.2
Anti-p40 (mAb1510)	0.19 + 0.04
anti-p19 (AF1716)	52 \pm 13

STAT3 Phosphorylation on Human PBMCs

[00156] A secondary cell-based confirmatory assay was developed with the goal of evaluating phosphorylation of STAT3 as a mechanism of action in primary human cells.

10 Peripheral blood mononuclear cells (PBMC) from healthy donors consist primarily of naïve and quiescent T-cells that nominally express low levels of IL-23R and do not appreciably respond when stimulated with exogenous IL-23. However, polyclonal activation of naïve PBMC with IL-2 results in activation and differentiation of naïve T-cells with subsequent increased expression of IL-23R. These activated cells are then
15 susceptible to stimulation with exogenous IL-23 which activates the STAT pathway, resulting in phosphorylation of STAT3.

[00157] Commercially available antibodies (AF1716, an anti-p19 pAb and mAb1510, an anti-p40 mAb, both from R&D Systems) were used as positive controls for inhibition of IL-23 induced STAT3 phosphorylation. The inhibitory activity of six adnectins was
20 compared in ten separate experiments using blood from multiple donors (summarized in Table 8). Exemplary data for a subset are shown in Figure 4. The anti-IL-23 adnectins

were significantly (>150-fold) more potent than the anti-p19 in inhibiting STAT3 phosphorylation but similar to 5 fold less potent than the anti-p40 monoclonal antibody.

Table 8: Inhibition of PBMC pSTAT3 by Anti-IL-23 Antagonists

	pSTAT3 IC ₅₀ ± SD (nM)
1490B03	0.03 ± 0.02
1571G04	0.09 ± 0.06
1572G06	0.07±0.02
ATI 934	0.14 ±0.12
ATI 1016	0.06 ± 0.03
ATI 1045	0.07 ± 0.03
MAB1510	0.03 ± 0.04
AF1716	21.4 ± 9.4

5

IL-23 Induced Cytokine Production by Mouse Splenocytes

[00158] Initial cellular assays with primary cells were designed to evaluate the capacity of anti-IL-23 adnectins to inhibit IL-23-dependent cytokine secretion from murine Th17 cells. To differentiate murine Th17 cells for analysis, CD4⁺ T cells were enriched with magnetic beads, co-cultured with irradiated splenocytes, and activated with anti-CD3 in presence of TGF- β and IL-6 and neutralizing antibodies for IL-4 and IFN- γ . After 6 days in culture, the polarized Th17 cells were harvested, re-seeded in a 96-well plate and stimulated with 100 ng/ml human IL-23 and 5 ng/ml murine IL-2. The addition of IL-2 was required to maintain cell viability and enable robust cytokine production in response to IL-23 but did not strongly induce IL-17A or IL-22 production alone. Because IL-2 induces a low level of cytokine secretion, each sample set included cells stimulated with IL-2 alone to control for baseline levels of cytokine produced in the absence of IL-23. The IL-23-dependent response was evaluated by calculating the difference between the level of cytokine induced by the combination of IL-2 and IL-23 and the baseline level induced by IL-2 alone. A dose range of adnectins were added during re-stimulation of the Th17 cells with IL-2 and IL-23 to test their inhibitory potential. A dose range of human anti-p40 antibody (R&D Systems MAB1510) was run in parallel as positive controls for

10
15
20

assessing IL-23 inhibition. Each condition was tested in triplicate wells of a 96-well plate. After 4 days, the conditioned media from the triplicates was pooled, cleared of cellular debris, and assayed for both IL-17A and IL-22 concentrations by ELISA.

[00159] Stimulation of Th17 cells with IL-2 and IL-23 induced a 2- to 3-fold increase of IL-17A and at least a 5-fold enhancement of IL-22 compared to the levels induced by IL-2 alone. ATI000934, ATI001014, ATI001015, ATI001016, ATI001045 and the positive control anti-p40 monoclonal antibody mediated dose-dependent decreases in IL-23-dependent IL-17A and IL-22 secretion. IC₅₀ values for inhibition of both IL-17A and IL-22 secretion were calculated for each adnectin as well as the anti-p40 control and these data were summarized in Table 9. All of the adnectins tested were within 2-fold as potent as the anti-p40 control for inhibition of IL-23-dependent IL-17A secretion and within 2- to 3-fold as potent for inhibition of IL-23-dependent IL-22 production.

Table 9: Inhibition of IL-23-Dependent Cytokines by Anti-IL-23 Adnectins

Adnectin/Ab	IC ₅₀ ± S.D. IL-17 (nM)	IC ₅₀ ± S.D. IL-22 (nM)
anti-p40 (MAB1510)	2.3 ± 0.7 (n=5)	1.9±0.7 (n=5)
ATI000934	5.3 ± 1.6 (n=2)	N.D.
ATI001045	1.3 ± 0.3 (n=3)	2.2 ± 1.1 (n=3)
ATI001014	3.7±0.0 (n=2)	6.5±2.2 (n=4)
ATI001015	2.0±0.1 (n=2)	5.5±2.4 (n=4)
ATI001016	2.0±2.0 (n=4)	3.0±1.8 (n=5)

15

IL-23 Induced Cytokine Production by Human T Cells

[00160] PBMCs were obtained by density-gradient separation of EDTA-treated whole blood from normal healthy donors. T cells were prepared from E+ fractions of PBMC rosetted with sheep red blood cells (SRBC). The T cells were plated at 100,000 cells per well into 96-well flat bottom plates that were coated with anti-CD3 (OKT at 10 µg/ml) for 1 hour at 37 °C and washed with PBS. Mixtures of RPMI-FCS media containing anti-CD28 (9.3 at 1 µg/ml) and IL-1β (10 ng/ml) or IL-1β + IL-23 (1 ng/ml) were prepared. This combination of cytokines has been shown to promote the differentiation of human T

20

cells into IL-17-secreting T cells. ATI001045, starting concentration of 1 µg/ml was added to the mixture containing IL-1β + IL-23. IL-17 was detected in supernatants using DUOSET® ELISA development kits (R&D Systems). ATI001045 inhibited IL-17 production with an EC₅₀ of 2.0 ± 1.6 nM (n = 4 different donors), using the IL-1β alone as background. The commercial anti-p40 antibody (MAB1510) was used as an internal control and inhibited IL-17 production with an EC₅₀ of 2.2 ± 1.4 nM (n = 3). Exemplary data from donor 228 is shown in Figure 6.

Selectivity of Anti-IL-23 Adnectin for IL-23 Over IL-12

10 [00161] Adnectins listed in Table 2 as well as ATI001016 were used to examine the biochemical selectivity towards IL-23/IL-12. The binding analysis involved the capture of anti-IL-23 Adnectins on immobilized anti-His antibody followed by flow of IL-23 or IL-12 over the Adnectin. The selectivity of the Adnectins for IL-23 was assessed by comparing the binding signal for a 100 fold higher concentration of IL-12 over IL-23. Exemplary data in Figure 7 shows that ATI001016 displayed robust binding (~40 RU) towards 10 nM human IL-23 while no detectable binding was observed for 1 µM human IL-12.

[00162] NK-92 cells are a human natural killer cell line known to respond to IL-12 in an IL-2 dependent fashion by secreting IFN-γ. Cells are typically washed to remove IL-2 then seeded into 96-well plates, then treated with 25 pM recombinant human IL-12 (or IL-12 preincubated with antagonists) and incubated for an additional 20 hours. Clarified supernatants are assayed for IFN-γ by ELISA.

[00163] A 4 point, 5 fold dilution series starting at 5µM was prepared of each of the adnectin clones listed in Table 2 and incubated with 25pM IL-12 for 30 minutes at 37 °C prior to the addition to NK-92 cells. A 12 point, 5 fold dilution series starting at 5 µM of ATI001045 and ATI001016 were incubated with 25 pM IL-12 for 30 minutes at 37 °C prior to the addition to NK-92 cells. None of the clones listed in Table 2 nor ATI001045 or ATI001016 detectably inhibited IFN-γ secretion at any of the concentrations tested demonstrating that these anti-IL-23 adnectins do not inhibit the interaction of IL-12 with the receptors on the surface of NK-92 cells. They appear equivalent to a negative control and 100 nM anti-p19 polyclonal antibody. As a positive control, anti-p40 monoclonal

antibody (mAb1510) inhibited IL-12 induced IFN- γ secretion with an IC₅₀ of 0.07 nM (Figure 8).

Anti-IL-23 Adnectin Block IL-23 Induced IL-17 in a Pharmacodynamic Model

- 5 [00164] Female C57Bl/6 mice were injected intraperitoneally (IP) with recombinant murine IL-2 and human IL-23 according with the following schedule.

Table 10: Dosing and Injection Schedule

	Time = -24h	Time = 0h	Time = 7h	Time = 23h
Murine IL-2	5 μ g	5 μ g	10 μ g	5 μ g
Human IL-23	0	10 μ g	10 μ g	10 μ g

- 10 [00165] All mice were euthanized 7 - 8 hours following the final dose of IL-2 and IL-23 at Time = 30h. Serum was collected and assayed for IL-17 and IL-23 by ELISA.
- [00166] Human IL-23 binds to the mouse receptor and induces the production of cytokines such as IL-17 and IL-22. Splenocytes from animals dosed intraperitoneally (IP) with IL-2 and human IL-23 secrete IL-17 when stimulated in culture *ex vivo* with
- 15 anti-mouse CD3e. Significant levels of IL-17 can be detected in the serum of animals that were subjected to the treatment regimen described in Table 10 in which C57Bl/6 mice are primed with IL-2 24 hours prior to 3 dual injections of IL-2 + IL-23 over an additional span of 24 hours. Presumably, IL-2 polyclonally activates and expands Th populations *in situ* and up-regulates the expression of IL-23 receptor. This provides a
- 20 method where the mechanisms of drug action and the relationship between drug concentration and effect in an *in vivo* setting can be investigated. The model was validated with an anti-p40 monoclonal antibody, mAb1510 (data not shown). In eight separate experiments, five anti-IL-23 adnectins were tested for their ability to inhibit the production of murine IL-17 when dosed SC at 0.5, 0.15, 0.05 and 0.015 mg/kg 2 hours
- 25 prior to the initial dose of IL-2 + IL-23 in eight separate experiments. Exemplary dose response data for ATI001045 is shown in Fig 9a (calculated average ED₅₀ of 0.03 mg/kg). All anti-IL-23 adnectins tested showed dose dependent inhibition of human IL-

23 murine IL-17 production in serum though the extent of inhibition was variable across adnectins.

Activity of Anti-IL-23 Adnectin in the IL-23 Induced Skin Acanthosis Model

- 5 [00167] The intra-dermal injection of IL-23 into the skin of the back or into the external ear pinna of mice induces dermal inflammation and hyperplasia of the epidermis (acanthosis) (Zheng, Y., "Interleukin-22, a TH17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis", *Nature*, Vol. 445/8 (Feb. 2007)). In these studies, recombinant human IL-23 (rHuIL-23) was injected into mouse ears to explore the
- 10 downstream consequences of aberrant cutaneous IL-23 exposure.
- [00168] Six to eight week old C57BL/6 female mice were injected with 5 ug of dual chain, recombinant, human IL-23 into the right ear every other day until Day 12. PBS was injected into the contra-lateral ear as a control. In one study, treatment with ATI001045 began approximately 2 hours before the first IL-23 injection and continued 3
- 15 times per week until Day 12. ATI001045 was administered SC at doses of 0.1, 0.3, 1, 3 mg/kg. In a second study vehicle or ATI000934-123 (1753E02) was administered IP at 1, 3, or 10 mg/kg approximately 1 hour prior to IL-23 administration and 3 times per week thereafter until Day 10. Anti-HuIL-12/IL-23 p40 Antibody (R&D mAb1510) at 10 mg/kg was given IP on Day 0 and 4 as a positive control. Ear thickness (in thousandths of an
- 20 inch) was measured every-other-day, prior to the next ear injection, using a MITUTOYO® (#2412F) dial caliper. Ear thickness was calculated by subtracting the value of the control ear from the measurement for the IL-23 injected ear for each animal. At the end of the study (Day 14 for ATI001045 and Day 12 for ATI000934), following euthanasia with CO₂ gas, ears were excised at the hairline and formalin fixed/paraffin-
- 25 embedded tissues were examined histologically on H&E stained slides.
- [00169] Overall, doses of 1, 3, and 10 mg/kg of ATI000934 provided a similar level of inhibition of IL-23-induced ear thickening in this study (Figure 10). Ear thickness in all treatment groups was significantly ($p < 0.01$ ANOVA/Dunnett's) less than Vehicle, including the anti-p40 group, from Day 5 through the end of the study on Day 12. On
- 30 Day 12, terminal plasma samples were obtained 48 hours post last dose and analyzed for circulating levels of ATI000934 which were determined to be 11, 18, 36 ug/ml respectively.

[00170] Following the last measurement on Day 12, ears were collected at necropsy for routine histologic examination from 10 animals per group. The majority of animals administered ATI000934 had acanthosis and dermal infiltrates, but the histologic severity score was reduced from that observed in vehicle treated animals. There was no apparent dose response. All of the animals administered anti-p40 also had acanthosis and dermal infiltrates, but the histologic severity score was also reduced from that observed in vehicle treated animals.

[00171] ATI001045 (1 mg/kg and 3 mg/kg) dose-dependently reduced ear thickness compared to Vehicle (PBS) treated animals from Day 5 through Day 14 ($p < 0.01$ vs. Vehicle ANOVA/Dunnett's, Figure 10). In contrast, the 0.1 mg/kg dose level was not statistically different ($p > 0.05$) from Vehicle treatment on any study day. Treatment with 0.3 mg/kg provided intermediate reduction that was statistically less than Vehicle on Days, 5, 7, 9. Serum samples collected 48 hours post last dose were evaluated for circulating levels of ATI001045 which were determined to be 0.698, 2.72, 8, 22.5 ug/ml for doses of 0.1, 0.3, 1, 3 mg/kg respectively. Histological analysis revealed that administration of ATI001045 resulted in a dose dependent reduction of IL-23 induced cellular infiltrates and acanthosis which correlated with the ear thickness score.

Example 5

Material and Methods Used Herein

High Throughput Protein Production (HTPP)

[00172] Selected binders were cloned into pET9d vector and transformed into *E. coli* BL21 DE3 plysS cells were inoculated in 5 ml LB medium containing 50 μ g/mL kanamycin in a 24-well format and grown at 37 °C overnight. Fresh 5 ml LB medium (50 μ g/mL kanamycin) cultures were prepared for inducible expression by aspiration 200 μ l from the overnight culture and dispensing it into the appropriate well. The cultures were grown at 37 °C until A600 0.6-0.9. After induction with 1mM isopropyl- β -thiogalactoside (IPTG) the culture was expressed for 6 hours at 30 °C and harvested by centrifugation for 10 minutes at 2750 g at 4 °C.

[00173] Cell pellets (in 24-well format) were lysed by resuspension in 450 μ l of Lysis buffer (50mM NaH₂PO₄, 0.5 M NaCl, 1x Complete Protease Inhibitor Cocktail-EDTA free (Roche), 1mM PMSF, 10mM CHAPS, 40mM Imidazole, 1 mg/ml lysozyme,

30µg/ml DNase, 2µg/ml aprotonin, pH 8.0) and shaken at room temperature for 1-3 hours. Lysates were clarified and re-racked into a 96-well format by transfer into a 96-well Whatman GF/D UNIFILTER® fitted with a 96-well, 1.2 ml catch plate and filtered by positive pressure. The clarified lysates were transferred to a 96-well Ni-Chelating Plate that had been equilibrated with equilibration buffer (50mM NaH₂PO₄, 0.5 M NaCl, 40mM Imidazole, pH 8.0) and was incubated for 5 min. Unbound material was removed by positive pressure. The resin was washed 2 x 0.3 ml/well with Wash buffer #1 (50mM NaH₂PO₄, 0.5 M NaCl, 5 mM CHAPS, 40mM Imidazole, pH 8.0) with each wash removed by positive pressure. Prior to elution each well was washed with 50µl Elution buffer (PBS + 20mM EDTA), incubated for 5 min and this wash was discarded by positive pressure. Protein was eluted by applying an additional 100µl of Elution buffer to each well. After a 30 minute incubation at room temperature the plate(s) were centrifuged for 5 minutes at 200 g and eluted protein is collected in 96-well catch plates containing 5µl of 0.5M MgCl₂ added to the bottom of elution catch plate prior to elution. Eluted protein was quantified using a total protein assay (BCA) with SGE as the protein standard.

Midscale Expression and Purification of Insoluble Fibronectin-Based Scaffold Protein Binders

[00174] For expression, selected clone(s), followed by the HIS6tag, were cloned into a pET9d vector and were expressed in *E. coli* BL21 DE3 plysS cells. Twenty ml of an inoculum culture (generated from a single plated colony) was used to inoculate 1 liter of LB medium or TB-Overnight Expression Media (auto induction) containing 50µg/ml Kanamycin and 34 µg/ml chloramphenicol. Cultures in LB medium were incubated at 37 °C until A600 0.6-1.0 at which time they then induced with 1mM isopropyl-β-thiogalactoside (IPTG) and grown for 4 hours at 30 °C. Cultures grown in TB-Overnight Expression Media were incubated at 37 °C for 5 hours at which time the temperature was lowered to 18 °C grown fir 19 hours. Cultures were harvested by centrifugation for 30 minutes at ≥10,000 g at 4 °C. Cell pellets were frozen at -80 °C. the cell pellet was resuspended in 25 ml of lysis buffer (20mM NaH₂PO₄, 0.5 M NaCl, 1x Complete Protease Inhibitor Cocktail-EDTA free (Roche),pH 7.4) using an ULTRA-TURRAX® homogenizer (IKA works) on ice. Cell lysis was achieved by high pressure

homogenization ($\geq 18,000$ psi) using a Model M-110S MICROFLUIDIZER® (Microfluidics). The insoluble fraction was separated by centrifugation for 30 minutes at $\geq 23,300$ g at 4 °C. The insoluble pellet recovered from centrifugation of the lysate was washed with 20mM sodium phosphate/500mM NaCl, pH7.4. The pellet was

5 resolubilized in 6.0M guanidine hydrochloride in 20mM sodium phosphate/500 mM NaCl pH 7.4 with sonication followed by incubation at 37 degrees for 1-2 hours. The resolubilized pellet was filtered to 0.45 μ m and loaded onto a HISTRAP® column equilibrated with the 20mM sodium phosphate/500 mM NaCl/6.0M guanidine pH7.4 buffer. After loading, the column was washed for an additional 25 CV with the same

10 buffer. Bound protein was eluted with 50mM Imidazole in 20mM sodium phosphate/500mM NaCl/6.0M guan-HCl pH7.4. The purified protein was refolded by dialysis against 50mM sodium acetate/150mM NaCl pH 4.5 or PBS pH 7.2.

Midscale Expression and Purification of Soluble Fibronectin-Base Scaffold Protein

15 Binders

[00175] As an alternative to purification of insoluble binders, the purification of soluble binders may also be used. For expression, selected clone(s), followed by the HIS6tag, were cloned into a pET9d vector and were expressed in *E. coli* BL21 DE3 plysS cells. Twenty ml of an inoculum culture (generated from a single plated colony) was

20 used to inoculate 1 liter of LB medium or TB-Overnight Expression Media (auto induction) containing 50 μ g/ml Kanamycin and 34 μ g/ml chloramphenicol. Cultures in LB medium were incubated at 37 °C until A600 0.6-1.0 at which time they were then induced with 1mM isopropyl- β -thiogalactoside (IPTG) and grown for 4 hours at 30 °C. Cultures grown in TB-Overnight Expression Media were incubated at 37 °C for 5 hours

25 at which time the temperature was lowered to 18 °C grown fir 19 hours. Cultures were harvested by centrifugation for 30 minutes at $\geq 10,000$ g at 4 °C. Cell pellets are frozen at -80 °C. The cell pellet is resuspended in 25 ml of lysis buffer (20mM NaH₂PO₄, 0.5 M NaCl, 1x Complete Protease Inhibitor Cocktail-EDTA free (Roche), pH 7.4) using an ULTRA-TURRAX® homogenizer (IKA works) on ice. Cell lysis is achieved by high

30 pressure homogenization ($\geq 18,000$ psi) using a Model M-110S MICROFLUIDIZER® (Microfluidics). The soluble fraction is separated by centrifugation for 30 minutes at $\geq 23,300$ g at 4 °C. The supernatant is clarified via 0.45 μ m filter. The clarified lysate is

loaded onto a HISTRAP® column (GE) pre-equilibrated with the 20mM sodium phosphate/500 mM NaCl pH 7.4. The column is then washed with 25 column volumes of the same buffer, followed by 20 column volumes of 20mM sodium phosphate/500 mM NaCl/ 25mM Imidazole, pH 7.4 and then 35 column volumes of 20mM sodium phosphate/500 mM NaCl/ 40mM Imidazole, pH 7.4. Protein is eluted with 15 column volumes of 20mM sodium phosphate/500 mM NaCl/ 500mM Imidazole, pH 7.4, fractions are pooled based on absorbance at A280 and are dialyzed against 1x PBS, 50mM Tris, 150mM NaCl. pH 8.5 or 50mM NaOAc; 150mM NaCl; pH4.5. Any precipitate is removed by filtering at 0.22µm.

10 **[00176]** Fibronectin-based scaffold proteins (Adnectins) can be pegylated with various sizes and types of PEG. To allow for pegylation, the naturally occurring residues EIDKPSQ, found at the C-terminus end of 10FN3 proteins can be modified by a single point mutation of an amino acid, typically a serine, to a cysteine. PEGylation of the protein at the single cysteine residue is accomplished by conjugating various maleimide-

15 derivatized PEG forms, combining the PEG reagent with the protein solution and incubating. An alternative method is to replace the EIDKPSQ tail with a GSGC linker, and similarly use the cysteine residue for PEGylation. Adnectins containing an engineered cysteine residue were conjugated with PEG via Michael-addition chemistry between the thiol group on the cysteine and the maleimide functional group of the PEG

20 reagent. Briefly, 40kDa PEG is added in a molar excess to protein solution under slightly acidic to neutral conditions. The reaction is allowed to proceed at room temperature for 2 hours to overnight. The reaction is then applied to an ion exchange column to separate the PEGylated Adnectin from the unreacted PEG-maleimide and non-PEGylated Adnectin. SE/HPLC methods may also be used. The purified PEGylated

25 Adnectin is typically analyzed by SDS-PAGE and size exclusion chromatography.

Example 6

Screening and Selection of Candidate Serum Albumin-Binding Adnectin (SABA)

[00177] A selection technique known as PROFusion (see, *e.g.*, Roberts et al., *Proc. Natl. Acad. Sci. USA*, 94(23):12297-12302 (1997) and WO 2008/066752) was applied to a DNA library with variable regions designed into the BC, DE and FG loops of ¹⁰Fn3. A random library of greater than 10¹³ molecules was created from this design, and selection

pressure was applied against a biotinylated form of HSA to isolate candidate serum albumin-binding Adnectin (SABA) with desirable binding properties.

High Throughput Protein Production (HTPP) Process

- 5 [00178] The various HSA binding Adnectins were purified using a high throughput protein production process (HTPP). Selected binders were cloned into pET9d vector containing a HIS6 tag and transformed into *E. coli* BL21(DE3)pLysS cells. Transformed cells were inoculated in 5 ml LB medium containing 50 µg/mL Kanamycin in a 24-well format and grown at 37 °C overnight. Fresh 5 ml LB medium (50 µg/mL Kanamycin)
- 10 cultures were prepared for inducible expression by aspirating 200 µl from the overnight culture and dispensing it into the appropriate well. The cultures were grown at 37 °C until A₆₀₀ 0.6-0.9. After induction with 1 mM isopropyl-β-thiogalactoside (IPTG), the culture was grown for another 4 hours at 30 °C and harvested by centrifugation for 10 minutes at 3220 x g at 4 °C. Cell Pellets were frozen at -80 °C.
- 15 [00179] Cell pellets (in 24-well format) were lysed by resuspension in 450 µl of Lysis buffer (50 mM NaH₂PO₄, 0.5 M NaCl, 1x Complete Protease Inhibitor Cocktail-EDTA free (Roche), 1 mM PMSF, 10 mM CHAPS, 40 mM Imidazole, 1 mg/ml lysozyme, 30 µg/ml DNase, 2 µg/ml aprotinin, pH 8.0) and shaken at room temperature for 1 hour. Lysates were clarified and re-racked into a 96-well format by transfer into a 96-well
- 20 Whatman GF/D UNIFILTER® fitted with a 96-well, 650 µl catch plate and centrifuged for 5 minutes at 200 x g. The clarified lysates were transferred to a 96-well Ni-Chelating Plate that had been equilibrated with equilibration buffer (50 mM NaH₂PO₄, 0.5 M NaCl, 10 mM CHAPS, 40 mM Imidazole, pH 8.0) and incubated for 5 min. Unbound material was removed. The resin was washed 2 x 0.3 ml/well with Wash buffer #1 (50
- 25 mM NaH₂PO₄, 0.5 M NaCl, 5 mM CHAPS, 40 mM Imidazole, pH 8.0). Next the resin was washed with 3 x 0.3 ml/well with PBS. Prior to elution each well was washed with 50 µl Elution buffer (PBS + 20 mM EDTA), incubated for 5 min and this wash discarded by vacuum. Protein was eluted by applying an additional 100 µl of Elution buffer to each well. After 30 minute incubation at room temperature the plate(s) were centrifuged for 5
- 30 minutes at 200 x g and eluted protein collected in 96-well catch plates containing 5 µl of 0.5M MgCl₂ affixed to the bottom of the Ni-plates. Eluted protein was quantified using a BCA Protein assay with SGE (control Adnectin) as the protein standard. The SGE

Adnectin is a wild-type ¹⁰Fn3 domain (SEQ ID NO: 1) in which integrin binding domain (amino acids RGD at positions 78-80) have been replaced with SGE.

HSA, RhSA and MuSA Direct Binding ELISA

5 [00180] For assaying direct binders to HSA, MaxiSorp plates (Nunc International, Rochester, NY) were coated with 10 ug/mL HSA (Sigma, St. Louis, MO) in PBS at 4 °C overnight followed by blocking in casein block buffer (Thermo Scientific, Rockford, IL) for 1-3 hours at room temperature. For single-point screening assays, purified HTPP Adnectin were diluted 1:20 in casein block buffer and allowed to bind to HSA in each
10 well for 1 hour at room temperature. For dose response assays, concentrations ranging from 0.1 nM up to 1 μM were used. After washing in PBST to remove unbound Adnectins, anti-His mAb-HRP conjugate (R&D Systems, MN) diluted 1:2500 in casein block buffer was added to the bound His-tagged Adnectin for 1 hour at room temperature. Excess conjugate was removed by washing with PBST and bound Adnectins detected
15 using TMB detection reagents (BD Biosciences) according to the manufacturer's instructions.

Identification of Candidate Serum Albumin-Binding Adnectin (SABA)

[00181] As a result of the screening for HSA/RhSA/MuSA binding and biophysical
20 criteria, four unique serum albumin-binding Adnectins (SABA) were identified and chosen to have their half-lives evaluated in mice. In order to carry out *in vitro* and *in vivo* characterization, midscales were undertaken for the four SABAs. Table 3 provides the sequences of twenty-six unique SABA core sequences identified from PROfusion, designated as SABA 1-26. SABA4 had a scaffold mutation that was fixed prior to
25 midscaling. The scaffold-perfect version of SABA4 is SABA5. SABA4 and SABA5 have identical sequences in the BC, DE, and FG loops.

Example 7

Production and Formulation of Candidate SABAs

30 Midscale Protein Production of SABAs

[00182] The selected SABAs followed by the His₆ tag, were cloned into a pET 9d vector and expressed in *E. coli* BL21(DE3)pLysS cells (see Table 3 for each His-tagged

SABA sequence designated SABA1.1, SABA2.1, SABA3.1, and SABA5.1). 20 ml of an inoculum culture (generated from a single plated colony) was used to inoculate 1 liter of LB medium containing 50 µg/mL Kanamycin. The culture was grown at 37 °C until A₆₀₀ 0.6-1.0. After induction with 1 mM isopropyl-β-thiogalactoside (IPTG) the culture was grown for another 4 hours at 30 °C and harvested by centrifugation for 30 minutes at ≥10,000 x g at 4 °C. Cell Pellets were frozen at -80 °C. The cell pellet was resuspended in 25 mL of lysis buffer (20 mM NaH₂PO₄, 0.5 M NaCl, 1x Complete Protease Inhibitor Cocktail-EDTA free (Roche), pH 7.4) using an ULTRA-TURRAX® homogenizer (IKA works) on ice. Cell lysis was achieved by high pressure homogenization (≥18,000 psi) using a Model M-110S MICROFLUIDIZER® (Microfluidics). The soluble fraction was separated by centrifugation for 30 minutes at 23,300 x g at 4 °C. The supernatant was clarified via 0.45 µm filter. The clarified lysate was loaded onto a HISTRAP® column (GE) pre-equilibrated with 20 mM NaH₂PO₄, 0.5 M NaCl, pH 7.4. The column was then washed with 25 column volumes of 20 mM NaH₂PO₄, 0.5 M NaCl, pH 7.4, followed by 20 column volumes of 20 mM NaH₂PO₄, 0.5 M NaCl, 25mM imidazole pH 7.4, and then 35 column volumes of 20 mM NaH₂PO₄, 0.5 M NaCl, 40 mM imidazole pH 7.4. Protein was eluted with 15 column volumes of 20 mM NaH₂PO₄, 0.5 M NaCl, 500 mM imidazole pH 7.4, fractions pooled based on absorbance at A₂₈₀ and dialyzed against 1x PBS, 50 mM Tris, 150 mM NaCl pH 8.5 or 50 mM NaOAc; 150 mM NaCl; pH 4.5. Any precipitate was removed by filtering at 0.22 µm .

[00183] Mid-scale expression and purification yielded highly pure and active Adnectins that were expressed in a soluble form and purified from the soluble fraction of the bacterial cytosol. SEC analysis on a SUPERDEX® 200 or SUPERDEX® 75 10/30GL in a mobile phase of 100 mM NaPO₄, 100 mM NaSO₄, 150 mM NaCl, pH 6.8 (GE Healthcare) demonstrated predominantly monomeric Adnectins.

Formulation of SABA1.2

[00184] One specific SABA, SABA1.2 (SEQ ID NO: 180), was chosen for a preliminary formulation screen. SABA1.2 comprises an (ED)₅ extension on the “core 1” sequence of ¹⁰Fn3. For SABA1.2, a stable formulation of 10 mM succinic acid, 8% sorbitol, 5% glycine at pH 6.0 and at a product concentration of 5 mg/mL was identified. In this formulation the protein melting temperature was 75 °C as determined by

Differential Scanning Calorimetry (DSC) using a protein concentration of 1.25 mg/mL. The formulation provided satisfactory physical and chemical stability at 4 °C and 25 °C, with an initial aggregate level at 1.2%. After one month of stability, the level of aggregation was very low (1.6% at 4 °C and 3.8% at 25 °C). The protein was also stable in this formulation after five cycles of freeze-thaw as transitioned from -80 °C and -20 °C to ambient temperature. In addition, in this formulation SABA1.2 was soluble to at least 20 mg/mL protein concentration at 4 °C and ambient temperature with no precipitation or increase in aggregation.

10

Example 8

Biophysical Characterization of Candidate SABAs

Size Exclusion Chromatography

[00185] Standard size exclusion chromatography (SEC) was performed on the candidate SABAs resulting from the midscale process. SEC of midscaled material was performed using a SUPERDEX® 200 10/30 or on a SUPERDEX® 75 10/30 column (GE Healthcare) on an Agilent 1100 or 1200 HPLC system with UV detection at A₂₁₄ nm and A₂₈₀ nm and with fluorescence detection (excitation = 280 nm, emission = 350 nm). A buffer of 100 mM sodium sulfate, 100 mM sodium phosphate, 150 mM sodium chloride, pH 6.8 at appropriate flow rate of the SEC column employed. Gel filtration standards (Bio-Rad Laboratories, Hercules, CA) were used for molecular weight calibration.

[00186] The results of the SEC on the midscaled purified SABAs showed predominantly monomeric Adnectin and elution in the approximate range of 10 kDa vs. globular Gel Filtration standards (BioRad) as showed.

25 Thermostability

[00187] Differential Scanning Calorimetry (DSC) analyses of the midscaled SABAs were performed to determine their respective T_m's. A 1 mg/ml solution was scanned in a N-DSC II calorimeter (Calorimetry Sciences Corp) by ramping the temperature from 5 °C to 95 °C at a rate of 1 degree per minute under 3 atm pressure. The data was analyzed vs. a control run of the appropriate buffer using a best fit using Origin Software (OriginLab Corp). The results of the SEC and DSC analyses are summarized in Table 11.

Table 11: Summary of SEC and DSC Analyses on Candidate SABAs

Clone	SEC		DSC (Tm)
	Monomer (%)	Dimer (%)	
SABA1.1	92.3	7.7	63.9 °C
SABA5.1	88	12	70.1 °C
SABA2.1	91	9	58.5 °C/78.2 °C
SABA3.1	99	BLD	65.2 °C

Example 9

Characterization of Candidate SABA1 Binding to Serum Albumin

- 5 **[00188]** The kinetics of selected SABA clones purified from HTPP and/or midscaled material were determined by capturing the respective serum albumin (HSA/RhSA/MuSA) on the surface of a Biosensor CM5 chip and flowing a concentration series of SABAs over both the reference flow cell and the captured albumins. In addition, binding to albumin was carried out under various pH conditions ranging from pH 5.5 to pH 7.4.
- 10 HSA-binding Adnectins SABA2.1, SABA3.1, SABA4.1 (SABA5.1) & SABA1.1 cross reacted with RhSA but did not cross react with MuSA. SABA2 and SABA4 binding is pH sensitive whereas clone SABA3 demonstrated pH resistance binding to HSA down to pH 6.0. SABA1.1 fits biochemical criteria for pH resistance and affinity/kinetics down to pH 5.5.
- 15 **[00189]** Domain mapping was determined by Biacore. Selected SABA clones purified from HTPP and/or midscaled material were determined by capturing HSA or a construct consisting of just HSA-domain I & II or HSA-domain III on the surface of a Biosensor CM5 chip and flowing a concentration series of the SABAs over both the reference flow cell and the captured albumins. Clones SABA2 & SABA1 bound to HSA
- 20 and the HSA-domain I-II construct but not the HSA-domain III construct. Clones SABA3 & SABA4 bound to HSA but not to either the HSA-domain I-II or HSA-domain III constructs. The results are summarized in Table 12.

Table 12: Binding Affinity and Kinetics of Candidate SABAs

25 (SABA1.1, 2.1, 3.1 and 4.1)

Adnectin	Target	K _D (nM)	K _{off} (s ⁻¹)	Resistant to pH 7.4→5.5	Epitope on HSA
SABA2	HSA	33.8 +/- 20.5 (n=6)	1.71E-04	--	Domain I-II
	RhSA	63.6	4.42E-04		
SABA3	HSA	863	6.82E-02	+++ (down to pH 6.0)	Neither domain I- II nor III (interfacial?)
	RhSA	431	3.37E-02		
SABA4	HSA	412 +/- 8 (n=4)	7.82E-04	--	Neither domain I- II nor III (interfacial?)
	RhSA	>1000	3.83E-03		
SABA1	HSA	47.2 +/- 18.2 (n=9)	4.57E-04	+++	Domain I-II
	RhSA	778 +/- 313 (n=4)	5.45E-03		

Example 10

Examination of the *In Vivo* t_{1/2} of Candidate SABAs

[00190] The half-life of HSA in mice was determined to allow for evaluation of HSA-binding Adnectins in mice as the HSA-binding Adnectins do not cross react with MuSA. HSA was injected into the tail vein of approximately 6 week old Ncr nude female mice at a 20 mg/kg (Figure 11A) and 50 mg/kg dose (Figure 11B), and the concentration of HSA in blood samples taken at intervals post-injection was determined by ELISA. The t_{1/2} of HSA injected into mice at 20 mg/kg and 50 mg/kg were determined to be ~24 hrs and ~20 hrs, respectively.

Half-Life Determination of SABA1-4 in Mice

[00191] One liter *E. coli* growth of HSA binding clones SABA1.1, SABA2.1, SABA3.1, and SABA4.1 were prepared, purified and endotoxin removed. Each SABA variant was injected into the tail vein of mice, and the concentration in blood samples taken at intervals post-injection was determined by ELISA.

[00192] The pharmacokinetic profiles of each SABA were compared in the presence or absence of HSA in approximately 6 week old Ncr nude female mice. The mice that were co-injected with HSA had the HSA premixed with each SABA (HSA in a 3-4 molar

excess) because the binding clone was selective for HSA and RhSA and did not bind the mouse serum albumin. The half-life of SABA1.1 in mice plasma was 0.56 hours whereas the half-life of SABA1.1 co-injected with HSA was 5.6 hours, a ~10-fold increase in half life (Figure 12A). The half-life of SABA2.1 in mice plasma was 0.24 hours whereas the half-life of SABA2.1 co-injected with HSA was 2.8 hours, a ~12-fold increase in half life (Figure 12B). The half-life of SABA3.1 in mice plasma was 0.28 hours whereas the half-life of SABA3.1 co-injected with HSA was 0.53 hours, a ~2-fold increase in half life (Figure 12C). The half-life of SABA4.1 in mice plasma was 0.66 hours whereas the half-life of SABA4 co-injected with HSA was 4.6 hours, a ~7-fold increase in half life (Figure 12D). A summary of the present example is shown in Figure 13A.

Half-Life Determination of SABA1.1 and SABA5.1 in Cynomolgus Monkeys

[00193] A three week single dose proof of concept study of SABA1.1 and SABA5.1 was conducted in cynomolgus monkeys to assess pharmacokinetics at a 1 mg per kg (mpk) dose IV in 2 cynomolgus monkeys. The pharmacokinetics were evaluated using a quantitative ELISA-based assay that was developed to detect the Adnectin in plasma samples. SABA1.1 has a half-life in the range of 96-137 hours. SABA5.1 has a half-life of approximately 12 hours and was only measureable in the ELISA up to 120 hours. Figure 14 A and B summarizes data for these clones and compares data from cynomolgus monkey.

Example 11

Characterization of SABA1 Binding To Serum Albumin

SABA1.1 and 1.2 Binds to HSA and RhSA

[00194] SABA1.2, a “core 1” ¹⁰F_n3 comprising an (ED)₅ extension (SEQ ID NO: 190) bound to human serum albumin (HSA) at neutral pH and 25 °C with an average association rate constant (k_a) of 8.21E+03 M⁻¹s⁻¹, and an average dissociation rate constant (k_d) of 4.43E-04 s⁻¹, for a calculated average K_d of 55.3 nM (Table 13). For rhesus serum albumin (RhSA), the measured average association rate constant was 6.6E+03 M⁻¹s⁻¹, and the dissociation rate constant was 3.78E-03 s⁻¹, giving a calculated average K_d of 580 nM. No measurable interaction between SABA1.2 and mouse or rat serum albumin could be observed up to 1 μM (Table 13 and Figure 15). At 37 °C, the k_a

and k_d increased between 2 to 5-fold, leading to a ~2-fold increase in affinity for HSA and 1/2 the affinity for RhSA (Table 13).

Table 13. Kinetic Parameters for SABA1.2 Binding to Albumins, in HBS-P Buffer

Albumin	Temp (°C)	k_a (1/Ms)	k_d (1/s)	KD (nM)
Human	25	$8.21 \pm 1.19 \text{ E}+03$	$4.43 \pm 0.65 \text{ E}-04$	55.3 ± 13.7
Rhesus		$6.60 \pm 1.18 \text{ E}+03$	$3.78 \pm 0.45 \text{ E}-03$	580 ± 62.6
Mouse		no observable binding		
Human	37	$3.38\text{E}+04$	$8.15\text{E}-04$	24.1
Rhesus		$1.89\text{E}+04$	$1.85\text{E}-02$	977.4
Mouse		no observable binding		

5

[00195] Additionally, a calorimetric titration was performed to determine the stoichiometry between SABA1 and HSA. For this study, SABA1.1, a “core 1” ¹⁰Fn3 comprising a His6 extension (SEQ ID NO: 189), was used. HSA (10 µl per injection of 115 µM protein solution) was injected into the calorimetric cell containing SABA1.1 at a concentration of 8.1 µM. The experiment was performed at 37 °C in PBS buffer pH 7.4. Figure 16 shows that SABA1.1 binds to HSA with 1:1 stoichiometry.

10

SABA1.2 Binds Potently to HSA at Low pH

[00196] The long half-life of albumins (*e.g.*, $t_{1/2}$ of HSA is 19 days) is due in large part to the fact that they are recycled from an endocytic pathway by binding to the neonatal Fc receptor, FcRn, under the low pH conditions that exist inside the endosome. As shown in Table 14 SABA1.2 potently bound HSA at the endosomal pH of 5.5, suggesting that the $t_{1/2}$ of SABA1, once bound to HSA, would also benefit from the FcRn recycling mechanism.

20

Table 14. Comparison of Albumin Binding Kinetics at pH 7.4 and 5.5, in MES Buffer

albumin	pH	k_a (1/Ms)	k_d (1/s)	KD (nM)
---------	----	--------------	-------------	---------

albumin	pH	ka (1/Ms)	kd (1/s)	KD (nM)
Human	7.4	9.26E+03	3.88E-04	41.9
	5.5	9.44E+03	2.70E-04	28.6
Rhesus	7.4	6.16E+03	2.95E-03	479
	5.5	7.57E+03	2.72E-03	359

SABA1.2 Binds to Domains I and II of HSA, but Not Domain III

[00197] The binding site SABA1.2 on albumin was mapped to the N-terminal domains I or II using recombinant HSA fragments and has no detectable binding to domain III (Figure 17). Because domain III is the domain of HSA that primarily interacts with FcRn, it is less likely that SABA1.2 would compete for HSA binding to FcRn, again increasing the possibility of fully leveraging the recycling mechanism for enhanced half-life.

Example 12

10

In Vivo Pharmacology of SABA1.2

[00198] A four week single dose pre-toxicology study of SABA1.2 was conducted in cynomolgus monkeys to assess pharmacokinetics and immunogenicity at two different dose levels. The pharmacokinetics and immunogenicity were also evaluated in a three-week, single-dose pre-toxicology study that included both intravenous and subcutaneous administration arms. Additionally, the pharmacokinetics of SABA1.2 was evaluated in two separate, single dose pre-toxicology studies in cynomolgus monkeys using a quantitative ELISA-based assay that was developed to detect SABA1.2 in plasma samples.

[00199] SABA1.2 was administered to monkeys at 1 mpk and 10 mpk IV. As shown in Figure 18 and the parameters described below, the C_{mx} and AUC increased approximately linear with dose. Non-compartmental analyses using WINNONLIN® software were performed to evaluate pharmacokinetic parameters. The clearance (CL) for SABA1.2 at 10 mpk was 0.15 ml/hr/kg, the beta phase half-life (t_{1/2}) was 143 hours, the volume of distribution (V_z) was 30 mL/kg, and total drug exposure (AUC_{0-∞}) was 5,609,457 hr*nmol/L (Table 15). The clearance (CL) for SABA1.2 at 1 mpk was 0.4

25

ml/hr/kg, the half-life ($t_{1/2}$) was 124 hours, the volume of distribution (V_z) was 72 mL/kg, and total drug exposure (AUC_{call}) was 214,636 hr*nmol/L (Table 15).

[00200] After SC or IV administration of SABA1.2, the beta-phase pharmacokinetic profiles were similar (Figure 19). Non-compartmental analyses using WINNONLIN® software were performed to evaluate pharmacokinetic parameters. The clearance (CL) for SABA1.2 at 1 mpk IV was 0.22 ml/hr/kg, the beta phase half-life ($t_{1/2}$) was 125 hours, the volume of distribution (V_z) was 40 mL/kg, and total drug exposure (AUC_{call}) was 357,993 hr*nmol/L (Table 15). The clearance (CL) for SABA1.2 at 1 mpk SC was 0.32 ml/hr/kg, the beta phase half-life ($t_{1/2}$) was 134 hours, the volume of distribution (V_z) was 62 mL/kg, and total drug exposure (AUC_{call}) was 251,339 hr*nmol/L (Table 15). The SC relative bioavailability (F) compared to IV was 0.7.

Table 15. Pharmacokinetic Parameters for SABA1.2 in Monkeys

Study #	1		2	
Dose (mg/kg)	1	10	1	1
Route of administration	i.v.	i.v.	i.v.	s.c.
N	3	3	1	2
CL (mL/hr/kg)	0.4	0.15	0.22	0.32
V_z (mL/kg)	72	30	40	62
AUC _{call} (hr*nmol/L)	214,636	5,609,457	357,993	251,339
beta $T_{1/2}$ (h)	124	143	125	134
Bioavailability (F)	n/a	n/a	n/a	0.7

We claim:

1. A polypeptide comprising a fibronectin type III tenth domain (¹⁰Fn3) wherein the ¹⁰Fn3 has at least one loop selected from loop BC, DE and FG with an altered amino acid sequence relative to the sequence of the corresponding loop of the human
5 ¹⁰Fn3 domain, and wherein the polypeptide binds the p19 subunit of IL-23 with a K_D of less than 500nM.
2. A polypeptide comprising a fibronectin type III tenth domain (¹⁰Fn3) wherein the ¹⁰Fn3 has at least one loop selected from loop BC, DE and FG with an altered amino acid sequence relative to the sequence of the corresponding loop of the human
10 ¹⁰Fn3 domain, and wherein the polypeptide binds the structural epitope of the p19 subunit of IL-23.
3. The polypeptide of claims 1 or 2 wherein the BC loop is selected from SEQ ID NO:2-6.
4. The polypeptide of claims 1 or 2 wherein the DE loop is selected from
15 SEQ ID NO:7-48.
5. The polypeptide of claims 1 or 2 wherein the FG loop is selected from SEQ ID NO:49-59.
6. The polypeptide of claim 1 or 2 wherein the BC, DE or FG loop amino acid sequence is at least 80% identical to any one of SEQ ID NOs: 2-59.
- 20 7. The polypeptide of claim 1 or 2 wherein the polypeptide amino acid sequence is at least 90% identical to any one of SEQ ID NOs: 60-100.
8. The polypeptide of claim 1 or 2 wherein the polypeptide amino acid sequence is at least 90% identical to amino acids 3-96 of SEQ ID NOs: 60-100.
9. The polypeptide of claims 1 or 2 wherein the BC loop sequence motif is
25 GHYPX₁HX₂ shown in SEQ ID NO: 257, wherein X₁ is either methionine or leucine, and X₂ is either isoleucine or valine.
10. The polypeptide of claims 1 or 2 wherein the FG loop sequence motif is YXX₃X₃X₃YX₃X₃I shown in SEQ ID NO: 258, wherein X₃ can be any amino acid.
11. The polypeptide of claims 1 or 2 further comprising one or more
30 pharmacokinetic (PK) moieties selected from the group consisting of polyethylene glycol, sialic acid, Fc, Fc fragment, transferrin, serum albumin, a serum albumin binding protein and a serum immunoglobulin binding protein.

12. The polypeptide of claim 11 wherein the PK moiety is polyethylene glycol.
13. The polypeptide of claim 1 or 2 further comprising a cysteine linker.
14. The polypeptide of claim 13 wherein the cysteine linker is selected from
5 the group consisting of the amino acids GSGC shown in SEQ ID NO: 101 and EIDKPCQ shown in SEQ ID NO: 102.
15. A pharmaceutically acceptable composition comprising the polypeptide of any one of claims 1-14, wherein the composition is essentially endotoxin free.
16. A method for regulating the pathogenicity of Th17 cells comprising
10 contacting the polypeptide of claim 1-14 with IL-23 in an amount effective to interfere with the reaction of endogenous IL-23 with Th17 cells.

1/33

FIG. 1

1434A08 (SEQ ID NO:216)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCATCGTACTCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1437G04 (SEQ ID NO:217)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTACTACCATTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1437A09 (SEQ ID NO:218)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTAAACAGCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1438E05 (SEQ ID NO:219)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTAACGTTTCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1438D01 (SEQ ID NO:220)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTAACCGTGCTCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1438B02 (SEQ ID NO:221)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTAAAACCTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1438A09 (SEQ ID NO:222)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTAAACAGCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

FIG. 1 (continued)

1486G03 (SEQ ID NO:223)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTCGTTACTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1486C04 (SEQ ID NO:224)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCCGCATCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1486D04 (SEQ ID NO:225)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTTCTACTCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1486B05 (SEQ ID NO:226)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTCGTATCTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1486D05 (SEQ ID NO:227)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCATCAGCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1487C03 (SEQ ID NO:228)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTAAACAGGTTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1487G03 (SEQ ID NO:229)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTGCTCATCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

FIG. 1 (continued)

1487D09 (SEQ ID NO:230)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATATTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTTCGTACATACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1487H04 (SEQ ID NO:231)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTGCTCGTACGTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACCAGGAATACGAATACCGTTACATACCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490E02 (SEQ ID NO:232)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTACTCAGTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490G02 (SEQ ID NO:233)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCCGCGTTACCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACATGGAAGAAAAATACGCTGTTATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490H05 (SEQ ID NO:234)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTATGCGTCAGCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490B03 (SEQ ID NO:235)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTCGTAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAAAGAAGCTAACTATCGTGAATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490H06 (SEQ ID NO:236)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTCAGAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

FIG. 1 (continued)

1490A07 (SEQ ID NO:237)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCATGCTAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490C07 (SEQ ID NO:238)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTAACCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1490H08 (SEQ ID NO:239)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTAACACTTCTCATAACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1491A05 (SEQ ID NO:240)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTCAGGTTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1571H03 (SEQ ID NO:241)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTAACCGTGTTCACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAGAAATACCATATCATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1571G04 (SEQ ID NO:242)

ATGGGAGTTTCTGATGTGCCSCGCGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCCGCGTTCTCACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1571G06 (SEQ ID NO:243)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCACCTGCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTACTAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAAGAAGCTGACTACTCTCAGATCCCAATTTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

FIG. 1 (continued)

1571F10 (SEQ ID NO:244)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTCGTTACCATACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGAACAGGTTGAATACCGTGAATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1572D04 (SEQ ID NO:245)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCCGCGTTCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGAACAGCCGATCTACGCCACTATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1572F05 (SEQ ID NO:246)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTTCAGAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1572G06 (SEQ ID NO:247)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTTACAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1572B10 (SEQ ID NO:248)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTGTTCCGCGTCATACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1572C09 (SEQ ID NO:249)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTACTCCGAAACATACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1572H05 (SEQ ID NO:250)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTTCTAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGAACAGGTTGAATACCGTGAATCCCAATTT
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

FIG. 1 (continued)

1572H08 (SEQ ID NO:251)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTTCTCGTAAATACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1550A07 (SEQ ID NO:252)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTGTTCCGCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1550C05 (SEQ ID NO:253)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCCGCGTTCGTTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1550E03 (SEQ ID NO:254)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATATCCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTATGCGTCATACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACTCCGAAGAACTGTACAAATACATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1550E06 (SEQ ID NO:255)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGATGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCCGCGCGTCATACCGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACGCTCAGGAAAACTACAAAGAAATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

1550H05 (SEQ ID NO:256)

ATGGGAGTTTCTGATGTGCCGCGGACCTGGAAGTGGTTGCTGCCACCCCACCAGCCTGCTGATCAG
CTGGGGTCATTACCCGCTGCATGTTTCGATATTACCGCATCACTTACGGCGAAACAGGAGGCAATAGCC
CTGTCCAGGAGTTCACTGTGCCTCGTCAGATCTACACAGCTACCATCAGCGGCCTTAAACCTGGCGTT
GATTATACCATCACTGTGTATGCTGTCACTTACTACAACGAAGCTGACTACTCTCAGATCCCAATTTTC
CATTAATTACCGCACAGAAATTGACAAACCATCCCAGCACCATCACCACCACCCTGA

FIG. 2





FIG. 3

Wildtype VSDVPRDLEVVAATPTSLLISWDAPAVTVRYRITYGETGGNSPVQEFVTPGSKSTATISGLKPGVDYTIITVYAVTGRGDSPASSKPIISINYRT
 1434A08 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPHRTHTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1437G04 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPYHYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1437A09 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPSKQHTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1438E05 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPSNVHTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1438D01 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPNRAHTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1438B02 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPRKYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1438A09 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPRSRYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1486G03 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPSRYYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1486C04 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPPHRYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1486D04 VSDVPRDLEVVAATPTSLLISWGHYPLHIRYRITYGETGGNSPVQEFVTPRSTHTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1486B05 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPSRIYATATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1486D05 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPHQRYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1487C03 VSDVPRDLEVVAATPTSLLISWGHYPLHIRYRITYGETGGNSPVQEFVTPKQVYATATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1487G03 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPAHRYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1487D09 VSDVPRDLEVVAATPTSLLISWGHYPMHIRYRITYGETGGNSPVQEFVTPSRRHTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1487H04 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPARQYATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1490E02 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPRTQYATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1490G02 VSDVPRDLEVVAATPTSLLISWGHYPMHVRYRITYGETGGNSPVQEFVTPRYHTATISGLKPGVDYTIITVYAVTYMEEKYAVIPIISINYRT
 1490H05 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPMRQHTATISGLKPGVDYTIITVYAVTYAQENYKEIPIISINYRT
 1490B03 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPSRKYTATISGLKPGVDYTIITVYAVTYKEANYREIPIISINYRT
 1490H06 VSDVPRDLEVVAATPTSLLISWGHYPLHIRYRITYGETGGNSPVQEFVTPRQKYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1490A07 VSDVPRDLEVVAATPTSLLISWGHYPLHIRYRITYGETGGNSPVQEFVTPHAKYTATISGLKPGVDYTIITVYAVTYAQENYKEIPIISINYRT
 1490C07 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPSNRYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1490H08 VSDVPRDLEVVAATPTSLLISWGHYPLHIRYRITYGETGGNSPVQEFVTPNTHATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1491A05 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPSQVYATATISGLKPGVDYTIITVYAVTYAQENYKEIPIISINYRT
 1571H03 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPNRVYATATISGLKPGVDYTIITVYAVTYAQEYH IPIISINYRT
 1571G04 VSDVPRDLEVVAATPTSLLISWGHYPLHVRYRITYGETGGNSPVQEFVTPRSHATISGLKPGVDYTIITVYAVTYAQENYKEIPIISINYRT
 1571G06 VSDVPRDLEVVAATPTSLLISWGHYPLLRYYRITYGETGGNSPVQEFVTPRTKYTATISGLKPGVDYTIITVYAVTYNEADYSQIPIISINYRT
 1571F10 VSDVPRDLEVVAATPTSLLISWGHYPLHIRYRITYGETGGNSPVQEFVTPSRYHTATISGLKPGVDYTIITVYAVTYEVEYREIPIISINYRT

FIG. 3 (continued)

1572D04 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPPRRYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1572F05 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPPRQKYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1572G06 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPRYKYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1572B10 VSDVPRDLEVVAAATPTSLLLISWGHYPMHVRYYRITYGETGGNSPVQEFVPPRHTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1572C09 VSDVPRDLEVVAAATPTSLLLISWGHYPMHIRYYRITYGETGGNSPVQEFVPTPKHTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1572H05 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPRSKYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1572H08 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPSRKYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1550A07 VSDVPRDLEVVAAATPTSLLLISWGHYPMHVRYYRITYGETGGNSPVQEFVPPRRYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1550C05 VSDVPRDLEVVAAATPTSLLLISWGHYPMHVRYYRITYGETGGNSPVQEFVPPRRYTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1550E03 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPRMRHTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1550E06 VSDVPRDLEVVAAATPTSLLLISWGHYPMHVRYYRITYGETGGNSPVQEFVPPRHTATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT
1550H05 VSDVPRDLEVVAAATPTSLLLISWGHYPLHVRYYRITYGETGGNSPVQEFVPRQIYATISGLKPGVDYTIITVYAVTYEQQIYATIPISINYRT

FIG. 4






FIG. 6




FIG. 7





FIG. 8






FIG. 9A

FIG. 9B





FIG. 10A



#. 3 mg/kg and Anti-p40 p<0.05 on Day 3
All treatments p<0.01 vs. Vehicle from Day 5




FIG. 11A

Determination of HSA *in vivo* half-life in mice

20 mg/kg




FIG. 11B

50 mg/kg




FIG. 12B

SABA2.1




FIG. 12D
SABA4.1




FIG. 13A
Summary of half-life enhancement in mice of SABAI-4 when co-injected with HSA





FIG. 13B


CLONE	PK (T1/2)		Comments
	Mice	Cyno	
SABA1.1	5.6hrs	96-137hrs	T1/2 = 96-137hrs
SABA5.1	4.6hrs	12hrs	Poor binding affinity for RhSA. 2-fold decrease in KD observed at pH<6.0
SABA2.1	2.8hrs	NA	Loss of binding at pH <6.5
SABA3.1	32min	NA	Poor T1/2 observed in mice

FIG. 14A
SABA1.1 and SABA5.1 $t_{1/2}$ in Cynomolgus monkey
SABA1.1



Monkey	$t_{1/2}$ (hrs)	C_{max} ($\mu\text{g/mL}$)	AUC _{0-t} ($\text{hr} \cdot \mu\text{g/mL}$)	CL _{obs} (mL/hr/kg)	V_z _{obs} (mL/kg)
#1	95.8	9.03	673.7	1.45	200.8
#2	136.6	7.24	625.1	1.60	315.2

FIG. 14B
SABA5.1
SABA5.1



	HL_Lambda_z (hr)	Cmax (ug/mL)	AUCcall (hr*ug/mL)	Cl_obs (mL/hr/kg)	Vz_obs (mL/kg)
N	2	2	2	2	2
Mean	12.186	17.358	246.882	4.089	72.507
SD	1.451	3.088	36.246	0.596	19.045
Min	11.16	16.18	221.26	3.67	59.04
Max	13.21	19.54	272.51	4.51	85.97
CV%	11.9	17.7	14.7	14.6	26.3

FIG. 15
SABA1.2 binding to albumins from human, mouse and rat by direct binding ELISA assay




FIG. 16

SABA1.1 binds to HSA with 1:1 stoichiometry




FIG. 17
Biacore Analysis of SABAI.2 Binding to Recombinant Domain Fragments of HSA




FIG. 18

Pharmacokinetic Profile for SABA1.2 in Monkeys Dosed at 1mpk and 10mpk

Dose linearity SABA1.2 monkey PK study





FIG. 19
Pharmacokinetic Profile for SABAI.2 in Monkeys Dosed Intravenously or Subcutaneously at 1mpk



INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/024959

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07K14/78 A61K38/39 C40B30/04 C12N15/10
 ADD. A61K38/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 C07K A61K C40B C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
 EPO-Internal, BIOSIS, Sequence Search, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2009/133208 A1 (NOVARTIS AG [CH]; HASTEWELL JOHN [US]; LOEW ANDREAS [US]) 5 November 2009 (2009-11-05)	1,2,4, 6-8, 10-15
Y	page 42; figures 1,8; sequence 68 the whole document	1-15
X	WO 2009/102421 A2 (SQUIBB BRISTOL MYERS CO [US]; CAMPHAUSEN RAY [US]; MORSE BRENT [US]; E) 20 August 2009 (2009-08-20)	1,2,4, 6-8, 10-15
Y	claims; sequence 43	1-15
Y	WO 2009/068649 A2 (GLAXO GROUP LTD [GB]; ASHMAN CLAIRE [GB]; BATUWANGALA THIL [GB]; BURDE) 4 June 2009 (2009-06-04) pages 7, 22-23; claims; example 17; sequences 132, 24 page 37 the whole document	1-16
	----- -/--	

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search 4 May 2011	Date of mailing of the international search report 26/05/2011
---	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Madruga, Jaime
--	--

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/024959

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	WO 2010/112458 A1 (NOVARTIS AG [CH]; BARDROFF MICHAEL [DE]; CARBALLIDO HERRERA JOSE M [CH] 7 October 2010 (2010-10-07) page 23; claims the whole document	1-16
Y	----- US 7 115 396 B2 (LIPOVSEK DASA [US] ET AL) 3 October 2006 (2006-10-03) cited in the application the whole document	1,2
Y	----- ALIAHMADI EHSAN ET AL: "TLR2-activated human langerhans cells promote Th17 polarization via IL-1beta, TGF-beta and IL-23.", EUROPEAN JOURNAL OF IMMUNOLOGY MAY 2009 LNKD- PUBMED:19350551, vol. 39, no. 5, May 2009 (2009-05), pages 1221-1230, XP002635193, ISSN: 1521-4141 the whole document	16
A	----- TAN Z Y ET AL: "Interleukin-23: Immunological roles and clinical implications", INTERNATIONAL JOURNAL OF BIOCHEMISTRY AND CELL BIOLOGY, EXETER, GB, vol. 41, no. 4, 1 April 2009 (2009-04-01), pages 733-735, XP025982131, ISSN: 1357-2725, DOI: DOI:10.1016/J.BIOCEL.2008.04.027 [retrieved on 2008-08-03] the whole document	1-16
A	----- KOIDE AKIKO ET AL: "Monobodies: antibody mimics based on the scaffold of the fibronectin type III domain", METHODS IN MOLECULAR BIOLOGY, HUMANA PRESS INC, NJ, US, vol. 352, 1 January 2007 (2007-01-01), pages 95-109, XP009102789, ISSN: 1064-3745 the whole document	1-16
A	----- BATORI V ET AL: "Exploring the potential of the monobody scaffold: effects of loop elongation on the stability of a fibronectin type III domain", PROTEIN ENGINEERING, OXFORD UNIVERSITY PRESS, SURREY, GB, vol. 15, no. 12, 1 January 2002 (2002-01-01), pages 1015-1020, XP002996047, ISSN: 0269-2139, DOI: DOI:10.1093/PROTEIN/15.12.1015 the whole document	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2011/024959

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009133208	A1	05-11-2009	AU 2009242038 A1 05-11-2009
			CA 2723277 A1 05-11-2009
			EP 2274331 A1 19-01-2011
			US 2011038866 A1 17-02-2011

WO 2009102421	A2	20-08-2009	AU 2009213141 A1 20-08-2009
			CN 102007145 A 06-04-2011
			EP 2247615 A2 10-11-2010
			KR 20100128291 A 07-12-2010
			US 2011053842 A1 03-03-2011

WO 2009068649	A2	04-06-2009	AR 069495 A1 27-01-2010
			AU 2008328726 A1 04-06-2009
			CA 2706419 A1 04-06-2009
			CN 101932608 A 29-12-2010
			DE 112008003232 T5 24-02-2011
			EA 201000704 A1 30-12-2010
			EP 2222709 A2 01-09-2010
			GB 2468232 A 01-09-2010
			JP 2011504742 T 17-02-2011
			KR 20100097720 A 03-09-2010
			PE 12342009 A1 07-09-2009
			US 2009148905 A1 11-06-2009
			US 2011008345 A1 13-01-2011
			UY 31504 A1 17-07-2009

WO 2010112458	A1	07-10-2010	NONE

US 7115396	B2	03-10-2006	US 2005038229 A1 17-02-2005
