




(86) **Date de dépôt PCT/PCT Filing Date:** 2011/05/26
(87) **Date publication PCT/PCT Publication Date:** 2012/01/05
(45) **Date de délivrance/Issue Date:** 2016/11/01
(85) **Entrée phase nationale/National Entry:** 2012/12/21
(86) **N° demande PCT/PCT Application No.:** JP 2011/062643
(87) **N° publication PCT/PCT Publication No.:** 2012/002096
(30) **Priorité/Priority:** 2010/06/29 (JP2010-148286)

(51) **Cl.Int./Int.Cl. C07D 253/06** (2006.01),
A01N 43/707 (2006.01), **A01N 43/80** (2006.01),
A01N 43/84 (2006.01), **A01N 47/02** (2006.01),
A01P 13/02 (2006.01), **C07D 401/04** (2006.01),
C07D 403/04 (2006.01), **C07D 403/06** (2006.01),
C07D 405/04 (2006.01), **C07D 409/04** (2006.01),
C07D 413/04 (2006.01), **C07D 417/04** (2006.01)

(72) **Inventeurs/Inventors:**
SHIBAYAMA, ATSUSHI, JP;
KAJIKI, RYU, JP;
KOBAYASHI, MASAMI, JP;
MITSUNARI, TAKASHI, JP;
NAGAMATSU, ATSUSHI, JP

(73) **Propriétaire/Owner:**

(54) **Titre : DERIVE 6-ACYL-1,2,4-TRIAZINE-3,5-DIONE ET HERBICIDES**
(54) **Title: 6-ACYL-1,2,4-TRIAZINE-3,5-DIONE DERIVATIVE AND HERBICIDES**



(57) **Abrégé/Abstract:**

Disclosed are compounds exhibiting sufficient herbicidal activity at low application dosage when they are applied to soils and foliage, and an agrochemical composition using the same, in particular herbicides. The compounds are triazine derivatives



(73) **Propriétaires(suite)/Owners(continued):**FMC CORPORATION, US

(74) **Agent:** RIDOUT & MAYBEE LLP

(57) **Abrégé(suite)/Abstract(continued):**

represented by following Formula 1 or salts thereof, and the herbicides containing them: [Chem. 28] wherein in the formula, R¹ represents a hydrogen atom; a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group, etc., R² represents a C₁-C₁₂ alkyl group, etc., Y and Z represent an oxygen atom or a sulfur atom, and A represents a 5- or 6-membered cyclic group which may contain a nitrogen atom, an oxygen atom, or a sulfur atom.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 January 2012 (05.01.2012)

PCT

(10) International Publication Number
WO 2012/002096 A1

(51) International Patent Classification:

C07D 253/06 (2006.01) **C07D 403/04** (2006.01)
A01N 43/707 (2006.01) **C07D 403/06** (2006.01)
A01N 43/80 (2006.01) **C07D 405/04** (2006.01)
A01N 43/84 (2006.01) **C07D 409/04** (2006.01)
A01N 47/02 (2006.01) **C07D 413/04** (2006.01)
A01P 13/02 (2006.01) **C07D 417/04** (2006.01)
C07D 401/04 (2006.01)

[JP/JP]; c/o KUMIAI CHEMICAL INDUSTRY CO., LTD., 4-26, Ikenohata 1-chome, Taito-ku, Tokyo, 1108782 (JP). **NAGAMATSU, Atsushi** [JP/JP]; c/o KUMIAI CHEMICAL INDUSTRY CO., LTD., 4-26, Ikenohata 1-chome, Taito-ku, Tokyo, 1108782 (JP).

(74) **Agent: SAEKI, Norio**; 4th Floor, Aminosan Kaikan Building, 15-8, Nihonbashi 3-chome, Chuo-ku, Tokyo, 1030027 (JP).

(21) International Application Number:

PCT/JP2011/062643

(22) International Filing Date:

26 May 2011 (26.05.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2010-148286 29 June 2010 (29.06.2010) JP

(71) **Applicants (for all designated States except US): KUMIAI CHEMICAL INDUSTRY CO., LTD.** [JP/JP]; 4-26, Ikenohata 1-chome, Taito-ku, Tokyo, 1108782 (JP). **IHARA CHEMICAL INDUSTRY CO., LTD.** [JP/JP]; 4-26, Ikenohata 1-chome, Taito-ku, Tokyo, 1100008 (JP).

(72) Inventors; and


(75) **Inventors/Applicants (for US only): SHIBAYAMA, Atsushi** [JP/JP]; c/o K.I CHEMICAL RESEARCH INSTITUTE CO., LTD., 408-1, Shioshinden, Iwata-shi, Shizuoka, 4371213 (JP). **KAJIKI, Ryu** [JP/JP]; c/o K.I CHEMICAL RESEARCH INSTITUTE CO., LTD., 408-1, Shioshinden, Iwata-shi, Shizuoka, 4371213 (JP). **KOBAYASHI, Masami** [JP/JP]; c/o KUMIAI CHEMICAL INDUSTRY CO., LTD., 4-26, Ikenohata 1-chome, Taito-ku, Tokyo, 1108782 (JP). **MITSUNARI, Takashi**

(81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** 6-ACYL-1,2,4-TRIAZINE-3,5-DIONE DERIVATIVE AND HERBICIDES

[1]

(57) **Abstract:** Disclosed are compounds exhibiting sufficient herbicidal activity at low application dosage when they are applied to soils and foliage, and an agrochemical composition using the same, in particular herbicides. The compounds are triazine derivatives represented by following Formula 1 or salts thereof, and the herbicides containing them: [Chem. 28] wherein in the formula, R¹ represents a hydrogen atom; a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group, etc., R² represents a C₁-C₁₂ alkyl group, etc., Y and Z represent an oxygen atom or a sulfur atom, and A represents a 5- or 6-membered cyclic group which may contain a nitrogen atom, an oxygen atom, or a sulfur atom.



WO 2012/002096 A1

DESCRIPTION

Title of Invention

6-ACYL-1,2,4-TRIAZINE-3,5-DIONE DERIVATIVE AND HERBICIDES

5 Technical Field

The present invention relates to a novel triazine derivative or its salt, and herbicides containing it as an effective component.

Background Art

10 Triazine derivatives are known from "Collection of Czechoslovak Chemical Communications (1969), 34(6), 1673-83.," etc., for example. However, no herbicidal activity is described for the compounds disclosed in these literatures. Although various compounds are reported as triazine-based herbicides (for example, see "The Pesticide Manual 15th Edition, 2009, published by BCPC"), they all have a 1,3,5-triazine ring. Specific examples of the
15 1,3,5-triazine-based agrochemicals include 2-chloro-4,6-bis-(ethylamino)-1,3,5-triazine (Simazine), 2-chloro-4-ethylamino-6-isopropylamino-1,3,5-triazine (Atrazin), 2,4-bis(ethylamino)-6-methylthio-1,3,5-triazine (Simetryn), 2,4-bis(isopropylamino)-6-methylthio-1,3,5-triazine (Prometryn), and 2-(1,2-dimethylpropylamino)-4-ethylamino-6-methylthio-1,3,5-triazine (Dimethametryn).

20 Further, as a 1,2,4-triazine-based agrochemical, there are known 4-amino-3-methyl-6-phenyl-1,2,4-triazine-5(4H)-one (Metamitron), 4-amino-6-tert-butyl-3-methylthio-1,2,4-triazine-5(4H)-one (Metribuzin), etc. It is disclosed in Japanese Patent Application Laid-Open (JP-A) No. 8-259546 that 4-(2,4-dihalogeno-5-alkoxyphenyl)-1,2,4-triazine-3,5-dione derivatives having a hydrocarbon substituent group at
25 6-position have a herbicidal activity. It is disclosed in JP-A No. 5-51369 that 3,5-diaryl-6-amino-1,2,4-triazine derivatives have a herbicidal activity. It is disclosed in JP-A No. 5-32641 that 3-mercapto-1,2,4-triazine derivatives have a herbicidal activity.

However, it is not known from any literatures that 6-acyl-1,2,4-triazine-3,5-dione derivatives represented by Formula 1 below have a herbicidal activity.

Citation List

Patent Literature

PLT 1: Japanese Patent Application Laid-Open No. 8-259546

PLT 2: Japanese Patent Application Laid-Open No. 5-51369

5 PLT 3: Japanese Patent Application Laid-Open No. 5-32641

Non Patent Literature

NPL 1: Collection of Czechoslovak Chemical Communications (1969), 34(6), 1673-83.

NPL 2: The Pesticide Manual 15th Edition (2009, published by BCPC)

10

Summary of Invention

Technical Problem

Herbicides used for useful crops and useful plants are required to be a chemical preparation
15 which can be applied to soils or leaves and exhibit a sufficient herbicidal effect with low chemical dosage. Further, as there is an increasing need concerning safety and effect on environment of a chemical substance, development of safer herbicides is waited for. The invention is devised to cope with such problems.


20 Solution to Problem

In order to achieve the object above, inventors of the invention synthesized many triazine compounds to study the herbicidal activity of various triazine derivatives, and intensively determined the herbicidal activity and usefulness of the compounds. As a result, it is found that, when triazine derivatives of the invention are applied to weeds or soils wherein weeds thrive, an
25 excellent herbicidal effect is obtained for a long period of time, and therefore the invention is completed accordingly.

Thus, the present invention relates to the following (1) to (43).

(1) A triazine derivative or a salt thereof represented by following Formula 1:

[Chem. 1]



[in the formula, R¹ represents a hydrogen atom; a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₃-C₆ halocycloalkyl C₁-C₆ alkyl group; an amino C₁-C₆ alkyl group; a nitro C₁-C₆ alkyl group; a C₁-C₆ alkylamino C₁-C₆ alkyl group; a di(C₁-C₆ alkyl)amino C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ haloalkylthio C₁-C₆ alkyl group; a C₁-C₆ haloalkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ haloalkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a hydroxy C₁-C₆ alkyl group; a phenyl C₁-C₆ alkoxy C₁-C₆ alkyl group (phenyl in the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α); a C₁-C₆ alkoxy C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyloxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyloxy C₁-C₆ alkyl group; a phenyloxy C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylthio C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfinyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfonyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a C₁-C₆ haloalkoxy C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α; a phenyl C₂-C₆ alkenyl group which may be substituted with one or more substituents selected from the Substituent group α; a

phenyl C₂-C₆ alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a phenoxyimino C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a di(C₁-C₆ alkoxy)C₁-C₆ alkyl group; a (R³¹R³²N-C=O)C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyloxy C₁-C₆ alkyl group; a C₁-C₆ alkylidene aminooxy C₁-C₆ alkyl group; a formyl C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkoxy C₁-C₆ alkyl group; a cyano C₁-C₆ alkoxy C₁-C₆ alkyl group; a cyano C₁-C₆ alkyl group; a C₂-C₆ alkylidene amino group; a di(C₁-C₁₀ alkyl)amino C₁-C₆ alkylidene amino group; a NR³¹R³² group; a C₁-C₆ alkoxy group; a C₂-C₆ alkenyloxy group; a C₂-C₆ alkynyloxy group; a C₃-C₆ cycloalkyloxy group; a C₃-C₆ cycloalkyl C₁-C₆ alkyloxy group; a C₁-C₆ haloalkoxy group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone]; a C₁-C₆ alkyl group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α]; a C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α]; or a C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic-oxy group in which the heterocyclic group in the heterocyclic-oxy group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α];


R² represents a hydrogen atom; a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl

group; a C₃-C₆ cycloalkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyloxy C₁-C₆ alkyl group; a di(C₁-C₆ alkoxy) C₁-C₆ alkyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a phenyl C₂-C₆ alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ,


Y and Z represent an oxygen atom or a sulfur atom,

"A" represents any one of the following formula A-1 to A-5,


[Chem. 2]




A-1




A-2



A-3



A-4



A-5


R⁴ represents a hydroxyl group; O⁻M⁺ (M⁺ represents an alkali metal cation or an ammonium cation); an amino group; a halogen atom; a cyano group; an isothiocyanate group; an isocyanate group; a hydroxycarbonyloxy group; a C₁-C₆ alkoxy carbonyloxy group; a benzyloxy carbonyloxy group which may be substituted with a substituent group selected from Substituent group α ; a C₁-C₆ alkoxy group; a C₂-C₆ alkenyloxy group; a C₂-C₆ alkynyloxy group; a C₃-C₆ cycloalkyloxy group; a cyanomethylene oxy group; a C₃-C₆ cycloalkyl C₁-C₆ alkyloxy group; a C₁-C₆ alkyl carbonyloxy group; a C₁-C₆ haloalkyl carbonyloxy group; a C₂-C₆ alkenyl carbonyloxy group; a C₂-C₆ haloalkenyl carbonyloxy group; a C₂-C₆ alkynyl carbonyloxy group; a C₂-C₆ haloalkynyl carbonyloxy group; a C₁-C₆ alkoxy carbonyl C₁-C₆ alkoxy group; a phenoxy group which may be substituted with one or more substituents selected from the Substituent group α ; a benzyloxy group which may be substituted with one or more substituents

selected from the Substituent group α ; a phenylcarbonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylcarbonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a phenylcarbonyl C₁-C₆ alkyloxy group which may be substituted with one or more substituents
5 selected from the Substituent group α ; a C₁-C₁₀ alkylsulfonyloxy group; a C₁-C₆ haloalkylsulfonyloxy group; a phenylsulfonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₁₀ alkylthio group; a C₁-C₁₀ alkylsulfinyl group; a C₁-C₁₀ alkylsulfonyl group; a C₁-C₆ haloalkylthio
10 group; a C₁-C₆ haloalkylsulfinyl group; a C₁-C₆ haloalkylsulfonyl group; a C₂-C₆ alkenylthio group; a C₂-C₆ alkenylsulfinyl group; a C₂-C₆ alkenylsulfonyl group; a C₂-C₆ alkynylthio group; a C₂-C₆ alkynylsulfinyl group; a C₂-C₆ alkynylsulfonyl group; a phenylthio group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylthio group which may be substituted with one or more substituents selected from the Substituent
15 group α ; a phenylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent
20 group α ; a C₁-C₁₀ alkylamino group; a di(C₁-C₁₀ alkyl)amino group; a C₁-C₆ alkoxy carbonylamino group; a C₁-C₆ alkoxy group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5
25 identical or different substituents selected from the Substituent group α); a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); or a heterocyclic-oxy group in which the heterocyclic group in the heterocyclic-oxy group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur

atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α),


A₁ represents a group represented by the following formula

[Chem. 3]




A₂ represents a group represented by the following formula

[Chem. 4]



A₃ represents a group represented by the following formula

[Chem. 5]



n represents 0, 1, or 2,

R⁵, R⁶, R⁸, R⁹, R³⁵ and R³⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, herein, R⁵ and R⁸ may be joined together to form a C₂-C₅ alkylene chain or a C₂-C₅ alkenylene chain, and may form a ring together with adjacent carbon atoms, and R⁵ and R³⁵ may be joined together to form a C₁-C₅ alkylene chain to form a ring with adjacent carbon atoms,

R⁷, R³³, and R³⁴ each independently represent a hydrogen atom, a C₁-C₆ alkyl group, a C₁-C₆ haloalkyl group, a C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, or a C₁-C₆ alkoxy group,

R¹⁴, R¹⁵, R¹⁶, and R¹⁷ each independently represent a hydrogen atom, a C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, or a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ,

R¹⁸ represents a hydrogen atom, a C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a C₂-C₆ alkynyl

group, a cyanomethyl group, or a benzyl group,

R^{20} represents a C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a C_3 - C_6 cycloalkyl group, or a C_3 - C_6 cycloalkyl C_1 - C_6 alkyl group,

R^{21} represents a hydrogen atom, a C_1 - C_6 alkyl group, or a halogen atom,

5 R^{23} represents a C_1 - C_6 alkyl group, a C_1 - C_6 haloalkyl group, a C_3 - C_6 cycloalkyl group, a C_1 - C_{10} alkylthio group, a C_1 - C_{10} alkylsulfinyl group, a C_1 - C_{10} alkylsulfonyl group, a phenylthio group which may be substituted with one or more substituents selected from the Substituent group α , a benzylthio group which may be substituted with one or more substituents selected from the Substituent group α , a phenylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α , a benzylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α , a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α , or a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ,

15 R^{24} represents a hydrogen atom, a halogen atom, a cyano group, a C_1 - C_6 alkyl group, a C_3 - C_6 cycloalkyl group, or a C_1 - C_6 alkoxycarbonylamino group,

R^{25} represents a C_1 - C_6 alkoxycarbonyl group, a cyano group, or a nitro group,

R^{31} and R^{32} each independently represent a hydrogen atom; a C_1 - C_6 alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylcarbonyl group; a C_1 - C_{10} alkylthio carbonyl group; a C_1 - C_6 alkoxycarbonyl group; a C_1 - C_6 haloalkyl group; a C_3 - C_6 cycloalkyl group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl group; a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); or a C_1 - C_6 alkyl group substituted with a

heterocyclic group in which the heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α), herein, R^{31} and R^{32} may be joined together to form a 5- to 6-membered ring with adjacent nitrogen atom, and the one or more carbon atoms in the ring may be substituted with a sulfur atom and/or an oxygen atom.

Herein, "Substituent group α " represents a group selected from a group consisting of:

a halogen atom; a hydroxyl group; a C_1 - C_6 alkyl group; a C_3 - C_6 cycloalkyl group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_2 - C_6 haloalkynyl group; a C_3 - C_6 halocycloalkyl group; a C_3 - C_6 halocycloalkyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy group; a C_3 - C_6 cycloalkyloxy group; a C_2 - C_6 alkenyloxy group; a C_2 - C_6 alkynyloxy group; a C_1 - C_6 alkylcarbonyloxy group; a C_1 - C_6 haloalkoxy group; a C_1 - C_6 alkylthio group; a C_1 - C_6 alkylsulfinyl group; a C_1 - C_6 alkylsulfonyl group; a C_1 - C_6 haloalkylthio group; a C_1 - C_6 haloalkylsulfinyl group; a C_1 - C_6 haloalkylsulfonyl group; an amino group; a C_1 - C_6 alkylcarbonylamino group; a mono(C_1 - C_6 alkyl)amino group; a di(C_1 - C_6 alkyl)amino group; a hydroxy C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylthio C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylthio C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylsulfonyl C_1 - C_6 alkyl group; a cyano C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkoxy group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy group; a C_1 - C_6 haloalkoxy C_1 - C_6 alkoxy group; a cyano C_1 - C_6 alkoxy group; a C_1 - C_6 acyl group; a C_1 - C_6 alkoxyimino C_1 - C_6 alkyl group; a carboxyl group; a C_1 - C_6 alkoxycarbonyl group; a carbamoyl group; a mono(C_1 - C_6 alkyl)aminocarbonyl group; a di(C_1 - C_6 alkyl)aminocarbonyl group; a nitro group; a cyano group; a phenyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic group comprising 2 to 10 carbon atoms and 1 to 5 identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic oxy group comprising 2 to 10 carbon atoms and 1 to 5 identical or different heteroatoms selected from an oxygen atom, a sulfur

atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); and a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting an carbonyl group; and

"Substituent group β " represents a group selected from a group consisting of : a halogen atom, a nitro group, a cyano group, a C₁-C₆ alkyl group, a C₁-C₆ haloalkyl group, a C₁-C₆ alkoxy group, and a C₁-C₆ haloalkoxy group.].

(2) The triazine derivative or the salt thereof according to (1), wherein A in Formula 1 is A-1.

(3) The triazine derivative or the salt thereof according to (1) or (2), wherein in A-1, A₁ is [X₁], A₂ is [X₃], and A₃ is [X₉].

(4) The triazine derivative or the salt thereof according to (3), wherein R⁵ and R⁶ in [X₁] is a hydrogen atom or a C₁-C₆ alkyl group, R⁸ and R⁹ in [X₃] is a hydrogen atom or a C₁-C₆ alkyl group, and R³⁵ and R³⁶ in [X₉] is a hydrogen atom or a C₁-C₆ alkyl group, or R⁵ and R³⁵ may bind to each other via a C₁-C₅ alkylene chain to form a ring.

(5) The triazine derivative or the salt thereof according to (1), wherein A in Formula 1 is A-3.

(6) The triazine derivative or the salt thereof according to (5), wherein R²⁰ in A-3 is a C₁-C₆ alkyl group, and R₂₁ in A-3 is a hydrogen atom or a C₁-C₆ alkyl group.

(7) The triazine derivative or the salt thereof according to any one of (1) to (6), wherein R⁴ in A-1 is a hydroxyl group or an O⁻M⁺ (M⁺ represents an alkali metal cation or an ammonium cation).

(8) The triazine derivative or the salt thereof according to any one of (1) to (7), wherein Y in Formula 1 is an oxygen atom.

(9) The triazine derivative or the salt thereof according to any one of (1) to (8), wherein R¹ in

Formula 1 is the group selected from the group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a phenyl group which may be substituted

with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone].

(10) The triazine derivative or the salt thereof according to any one of (1) to (9), wherein R² in Formula 1 is the group selected from the group consisting of a C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α).

(11) The triazine derivative or the salt thereof according to (1), in which the groups in Formula 1 are as follows: R¹ represents a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyloxy C₁-C₆ alkyl group; a phenyloxy C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylthio C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfinyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfonyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a

phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ; a

5 C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a di(C₁-C₆ alkoxy) C₁-C₆ alkyl group; a C₁-C₆ alkoxy carbonyl C₁-C₆ alkyl group; a C₁-C₆ alkyl carbonyl C₁-C₆ alkyl group; a C₁-C₆ alkyl carbonyloxy C₁-C₆ alkyl group; a NR³¹R³² group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different

10 substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); or a C₁-C₆ alkyl group substituted with a heterocyclic group in which the heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different

15 substituents selected from the Substituent group α);

R² represents a hydrogen atom; a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen

20 atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ;

Y and Z represent an oxygen atom or a sulfur atom,

25 A represents any one of A-1, A-3, and A-5,

A₁ is [X₁],

A₂ is [X₃] or [X₄], and

A₃ is [X₉],

in [X₁], R⁵ and R⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group,

in [X₃], R⁸ and R⁹ each independently represent a hydrogen atom or a C₁-C₆ alkyl group,
 in [X₉], R³⁵ and R³⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group,
 herein, R⁵ and R⁸ may be joined together to form a C₂-C₅ alkylene chain or a C₂-C₅
 alkenylene chain, and may form a ring together with adjacent carbon atoms, and R⁵ and R³⁵ may

be joined together to form a C₁-C₅ alkylene chain to form a ring with adjacent carbon atoms,

in A-3, R²⁰ is a C₁-C₆ alkyl group,

R²¹ is a hydrogen atom or a C₁-C₆ alkyl group,

in A-5, R²⁴ represents a hydrogen atom, a C₁-C₆ alkyl group, or a C₃-C₆ cycloalkyl group,

R²⁵ represents a C₁-C₆ alkoxycarbonyl group, a cyano group, or a nitro group,

R⁴ represents a hydroxyl group; O⁻M⁺ (M⁺ represents an alkali metal cation or an ammonium cation); or a C₁-C₁₀ alkylsulfonyloxy group;

R³¹ and R³² each independently represent a hydrogen atom; a C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ; herein, R³¹ and R³² may be joined together to form a 5- to 6-membered ring with adjacent nitrogen atom, and the one or more carbon atoms in the ring may be substituted with a sulfur atom and/or an oxygen atom,

herein, "Substituent group α " represents a group selected from a group consisting of:

a halogen atom; a C₁-C₆ alkyl group; a C₃-C₆ cycloalkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₁-C₆ alkoxy group; a C₃-C₆ cycloalkyloxy group; a C₂-C₆ alkenyloxy group; a C₂-C₆ alkynyloxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a nitro group; a cyano group; a phenyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic oxy group comprising 2 to 10 carbon atoms and 1 to 5 heteroatoms that are optionally selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); and a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be

substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting an carbonyl group.

(12) The triazine derivative or the salt thereof according to (1), in which the groups in Formula 1 are as follows:

5 R^1 is a group selected from a group consisting of a C_1 - C_{12} alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_3 - C_6 cycloalkenyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_1 - C_6 alkylthio C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a phenyl group which may be substituted with one or more substituents selected
10 from the Substituent group α ; a phenyl C_1 - C_6 alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C_1 - C_6 alkoxyimino C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy carbonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkyl carbonyl C_1 - C_6 alkyl group; a $NR^{31}R^{32}$ group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the
15 group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); and, a C_1 - C_6 alkyl group substituted with a heterocyclic group in which the heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a
20 nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α);

R^{31} and R^{32} each independently represent a group selected from a group consisting of a hydrogen atom; a C_1 - C_6 alkyl group; and, a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ;

25 R^2 represents a group selected from a group consisting of a C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_1 - C_6 haloalkyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α);

and, a phenyl group which may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α ;

Y and Z represent an oxygen atom or a sulfur atom,

A represents any one of A-1, A-3, and A-5,

5 R^4 in A-1 represents a hydroxyl group;

OM^+ (M^+ represents an alkali metal cation or an ammonium cation);

or a C_1 - C_{10} alkylsulfonyloxy group;

in A-1, A_1 is $[X_1]$,

A_2 is $[X_3]$ or $[X_4]$, and

10 A_3 is $[X_9]$,

in $[X_1]$, R^5 and R^6 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_3]$, R^8 and R^9 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_9]$, R^{35} and R^{36} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group,

15 herein, R^5 and R^8 may bind to each other via a C_2 - C_5 alkylene chain or a C_2 - C_5 alkenylene chain to form a ring, and R^5 and R^{35} may bind to each other via a C_1 - C_5 alkylene chain to form a ring,

in A-3, R^{20} is a C_1 - C_6 alkyl group,

R^{21} is a hydrogen atom or a C_1 - C_6 alkyl group, and

20 R^4 represents a hydroxyl group; OM^+ (M^+ represents an alkali metal cation or an ammonium cation); or a C_1 - C_{10} alkylsulfonyloxy group;

"Substituent group α " represents a group selected from a group consisting of : a halogen atom; a C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_1 - C_6 haloalkyl group; a C_1 - C_6 alkoxy group; a C_1 - C_6 haloalkoxy group; a C_1 - C_6 alkylthio group; a C_1 - C_6 alkylsulfinyl group; a C_1 - C_6 alkylsulfonyl group; a nitro group; a cyano group; a phenyl group; 25 and a C_3 - C_6 alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting an carbonyl group.

(13) The triazine derivative or the salt thereof according to (1), in which the groups in

Formula 1 are as follows:

R^1 represents a group selected from a group consisting of a C_1 - C_{12} alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_3 - C_6 cycloalkenyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_1 - C_6 alkylthio C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxyimino C_1 - C_6 alkyl group; a C_1 - C_6 alkoxycarbonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylcarbonyl C_1 - C_6 alkyl group; a $NR^{31}R^{32}$ group; a heterocyclic group selected from the group consisting of pyridyl group, pyrimidinyl group, pyridazinyl group, thienyl group, isoxazolyl group, pyrazolyl group, morpholinyl group, thiomorpholinyl group, pyrazinyl group, piperidinyl group, and piperazinyl group (the heterocyclic group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); and, a tetrahydrofuryl-methyl group;

R^{31} and R^{32} each independently represent a group selected from a group consisting of a hydrogen atom; a C_1 - C_6 alkyl group; and a phenyl group;

R^2 represents a group selected from a group consisting of a C_1 - C_6 alkyl group; a C_1 - C_6 haloalkyl group; a pyridyl group; and a phenyl group;

Y and Z represent an oxygen atom or a sulfur atom,

A represents any one of A-1 and A-3,

R^4 in A-1 represents a hydroxyl group; or a C_1 - C_{10} alkylsulfonyloxy group,

in A-1, A_1 is $[X_1]$, A_2 is $[X_3]$ or $[X_4]$, and A_3 is $[X_9]$,

in $[X_1]$, R^5 and R^6 are a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_3]$, R^8 and R^9 are a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_9]$, R^{35} and R^{36} are a hydrogen atom or a C_1 - C_6 alkyl group,

herein, R^5 and R^8 may be joined together to form a C_2 - C_5 alkylene chain and to form a ring,

and R^5 and R^{35} may be joined together to form a C_1 - C_5 alkylene chain and to form a ring,

in A-3, R^{20} is a C_1 - C_6 alkyl group, R^{21} is a hydrogen atom or a C_1 - C_6 alkyl group, and R^4

represents a hydroxyl group or a C₁-C₁₀ alkylsulfonyloxy group, and

"Substituent group α " represents a group selected from a group consisting of : a halogen atom; a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₁-C₆ alkoxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a nitro group; a cyano group; a phenyl group; and a methylenedioxy group.

(14) An agrochemical composition comprising the triazine derivative or the salt thereof described in any one of (1) to (13), and an agriculturally acceptable carrier.

(15) The agrochemical composition according to (14), in which the agrochemical composition further comprises a surface active agent.

(16) A herbicide comprising the triazine derivative or the salt thereof described in any one of (1) to (13) as an active component.

(17) The herbicide according to (16), in which the herbicide has a herbicidal activity for weeds in a field or a paddy field in which agrohorticultural plants are cultivated.

(18) The herbicide according to (17), in which the agrohorticultural plants are agrohorticultural plants given with resistance by a breeding method or a genetic recombination technique.


(19) A method of eliminating weeds in soils by applying an effective amount of herbicides comprising the triazine derivative or the salt thereof described in any one of (16) to (18).

(20) The method according to (19), in which the soils are a farmland.

(21) The method according to (19), in which the farmland is a field or a paddy field in which agrohorticultural plants are cultivated.

(22) A triazine derivative or a salt thereof represented by following Formula 2:

[Chem. 6]



[in the formula, B represents a hydroxyl group or a C₁-C₆ alkoxy group and R¹, R², Y, and Z have the same definitions as those described in above Formula 1].

(23) The triazine derivative or the salt thereof according to (22), wherein Y in Formula 2 is an oxygen atom.

(24) The triazine derivative or the salt thereof according to (22) or (23), wherein R¹ in Formula 2 represents a group selected from a group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone).

(25) The triazine derivative or the salt thereof according to any one of (22) to (24), wherein R² in Formula 2 represents a group selected from a group consisting of a C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α).

(26) The triazine derivative or the salt thereof according to (22) or (23), wherein B is a hydroxyl group and R² is a C₁-C₆ alkyl group.

(27) The triazine derivative or the salt thereof according to (26), wherein R¹ represents a group selected from a group consisting of a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆

alkylcarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyloxy C₁-C₆ alkyl group; a C₁-C₆ alkylidene aminooxy C₁-C₆ alkyl group; a NR³¹R³² group; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone].

(28) The triazine derivative or the salt thereof according to (26), wherein R¹ represents a group selected from a group consisting of a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a NR³¹R³² group; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone].

(29) The triazine derivative or the salt thereof according to (27) or (28), wherein a heterocyclic group is 5- or 6-membered aromatic heterocyclic group having 1 to 3 nitrogen atoms as a heteroatom.

(30) The triazine derivative or the salt thereof according to any one of (26) to (29), wherein R³¹ and R³² each independently represent a hydrogen atom; a C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkylcarbonyl group; a C₁-C₆ alkoxycarbonyl group; a C₁-C₆ haloalkyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; or R³¹ and R³² may be joined together to form a 5- to 6-membered ring with adjacent nitrogen atom, and in such case, one or more carbon atom in the ring may be substituted with a sulfur atom and/or an oxygen atom.

(31) The triazine derivative or the salt thereof according to (30), wherein R³¹ and R³² each independently represent a hydrogen atom; a C₁-C₆ alkyl group; or a phenyl group which may be substituted with one or more substituents selected from the Substituent group α .

(32) The triazine derivative or the salt thereof according to any one of (26) to (31), wherein "Substituent group α " represents a group selected from a group consisting of a halogen atom; a C₁-C₆ alkyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₃-C₆ halocycloalkyl group; a C₃-C₆ halocycloalkyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy group; a C₃-C₆ cycloalkyloxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ haloalkylthio group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; or a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting an carbonyl group.

(33) The triazine derivative or the salt thereof according to (32), wherein "Substituent group α " represents a group selected from a group consisting of a halogen atom; a C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₁-C₆ alkoxy group; or a C₁-C₆ alkylthio group.

(34) The triazine derivative or the salt thereof according to any one of (22) to (33), wherein

Y in Formula 2 is an oxygen atom,

R¹ in Formula 2 represents a group selected from a group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); and

R² in Formula 2 represents a group selected from a group consisting of a C₁-C₆ alkyl group;

a C₁-C₆ haloalkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α).

(35) The triazine derivative or the salt thereof according to any one of (22) to (34), wherein

Y in Formula 2 is an oxygen atom,

R¹ in Formula 2 represents a group selected from a group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group selected from the group consisting of pyridyl group, pyrimidinyl group, pyrazinyl group, pyridazinyl group, thienyl group, thiazolyl group, isoxazolyl group, pyrazolyl group, morpholinyl group, thiomorpholinyl group, and piperazinyl group (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone) ;

R² is a group selected from a group consisting of a C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; and a pyridyl group; and,

"Substituent group α " represents a group selected from a group consisting of a halogen atom; a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₁-C₆ alkoxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a nitro group; a cyano group; a phenyl group; and a methylenedioxy group.

(36) An agrochemical composition comprising the triazine derivative or the salt thereof described

in any one of (22) to (35), and an agriculturally acceptable carrier.

(37) The agrochemical composition according to (36), in which the agrochemical composition further comprises a surface active agent.

(38) A herbicide comprising the triazine derivative or the salt thereof described in any one of (22) to (35) as an active component.

(39) The herbicide according to (38), in which the herbicide has a herbicidal activity for weeds in a field or a paddy field in which agrohorticultural plants are cultivated.

(40) The herbicide according to (39), in which the agrohorticultural plants are agrohorticultural plants given with resistance by a breeding method or a genetic recombination technique.

(41) A method of eliminating weeds in soils by applying an effective amount of herbicides comprising the triazine derivative or the salt thereof described in any one of (22) to (35).

(42) The method according to (41), in which the soils are a farmland.

(43) The method according to (41), in which the farmland is a field or a paddy field in which agrohorticultural plants are cultivated.

Advantageous Effects of Invention

The invention provides the novel triazine derivative represented by Formula 1 or its salt which can effectively control weeds. The triazine derivative of the invention or its salt exhibits an excellent herbicidal effect against various weeds, which cause a problem particularly in an agricultural field over a long period of time from a pre-germination stage to a growing stage, for example, a broad-leaf weed like white pepper, *Amaranthus viridis*, white goosefoot, *Stellaria media*, chamomile, China jute, *Sida spinosa*, sesbania, hogweed, red poppy, morning glory, and cocklebur, annual and perennial weeds of *Cyperus microiria* family including coco grass, edible galingale, *Kyllinga brevifolia* var. *leiolepis*, java galingale, and *Cyperus iria*, and gramineous weeds like barnyard millet, finger grass, foxtail, spear grass, Syrian sorghum *nitidum*, short awn, and wild oat. In addition, it can control rice paddy weeds including annual weeds like *Echinochloa oryzicola*, *Cyperus difformis*, and *Monochoria vaginalis* and perennial weeds like *Sagittaria pygmaea*, *Sagittaria trifolia*, *Cyperus serotinus*, *Eleocharis kuroguwai*, *Scirpus hotarui*, and *Alisma canaliculatum*.

Further, the compound of the invention is highly safe to useful crops and useful plants, in particular, to rice, wheat, barley, corn, grain sorghum, soybean, cotton, sugar beet, etc.

Thus, the invention provides an agrochemical composition having an excellent effect as herbicides.

5

Description of Embodiments

The definitions of the terms used in the present Description are given below.

Halogen atom refers to a fluorine atom, a chlorine atom, a bromine atom, or an iodine atom.

The descriptions like C₁-C₆ indicate the number of carbon atoms in a substituent group described hereinbelow. For example, C₁-C₆ means 1 to 6 carbon atoms.

The C₁-C₆ alkyl group represents, unless specified otherwise, a linear or branched alkyl group having 1 to 6 carbon atoms, and examples thereof include a group like methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1-ethylpropyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, neopentyl, n-hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1-ethylbutyl, 2-ethylbutyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, and 1-ethyl-2-methylpropyl.

The C₁-C₁₂ alkyl group represents, unless specified otherwise, a linear or branched alkyl group having 1 to 12 carbon atoms, and examples thereof include, in addition to those exemplified above for the C₁-C₆ alkyl group, a group like heptyl, 1-methylhexyl, 5-methylhexyl, 1,1-dimethylpentyl, 2,2-dimethylpentyl, 4,4-dimethylpentyl, 1-ethylpentyl, 2-ethylpentyl, 1,1,3-trimethylbutyl, 1,2,2-trimethylbutyl, 1,3,3-trimethylbutyl, 2,2,3-trimethylbutyl, 2,3,3-trimethylbutyl, 1-propylbutyl, 1,1,2,2-tetramethylpropyl, octyl, 1-methylheptyl, 3-methylheptyl, 6-methylheptyl, 2-ethylhexyl, 5,5-dimethylhexyl, 2,4,4-trimethylpentyl, 1-ethyl-1-methylpentyl, nonyl, 1-methyloctyl, 2-methyloctyl, 3-methyloctyl, 7-methyloctyl, 1-ethylheptyl, 1,1-dimethylheptyl, 6,6-dimethylheptyl, decyl, 1-methylnonyl, 2-methylnonyl, 6-methylnonyl, 1-ethyloctyl, 1-propylheptyl, n-nonyl, and n-decyl.

The C₃-C₆ cycloalkyl group represents, unless specified otherwise, a cycloalkyl group

having 3 to 6 carbon atoms, and examples thereof include a group like cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

The C₃-C₆ cycloalkenyl group represents, unless specified otherwise, a cycloalkenyl group having 3 to 6 carbon atoms, and examples thereof include a group like cyclopentenyl and cyclohexenyl.

The C₃-C₆ cycloalkyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a cycloalkyl having 3 to 6 carbon atoms, wherein the cycloalkyl moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like cyclopropylmethyl, 1-cyclopropylethyl, 2-cyclopropylethyl, 1-cyclopropylpropyl, 2-cyclopropylpropyl, 3-cyclopropylpropyl, cyclobutylmethyl, cyclopentylmethyl, and cyclohexylmethyl.

The C₃-C₆ cycloalkyl C₁-C₆ alkyloxy group represents an (alkyl)-O- group (i.e., alkoxy group) having 1 to 6 carbon atoms substituted with a cycloalkyl having 3 to 6 carbon atoms, wherein the cycloalkyl moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like cyclopropylmethoxy, 1-cyclopropylethoxy, 2-cyclopropylethoxy, 1-cyclopropylpropoxy, 2-cyclopropylpropoxy, 3-cyclopropylpropoxy, cyclobutylmethoxy, cyclopentylmethoxy, and cyclohexylmethoxy.

The C₃-C₆ halocycloalkyl group represents, unless specified otherwise, a cycloalkyl group having 3 to 6 carbon atoms substituted with 1 to 5, or preferably 1 to 3 halogen atoms, wherein the cycloalkyl moiety and the halogen atom have the same definitions as above, and examples thereof include a group like 2,2-difluorocyclopropyl and 2,2-dichlorocyclopropyl.

The C₃-C₆ halocycloalkyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a cycloalkyl group having 3 to 6 carbon atoms substituted with 1 to 5, or preferably 1 to 3 halogen atoms, wherein the cycloalkyl moiety, the alkyl moiety, and the halogen atom have the same definitions as above, and examples thereof include a group like 2,2-difluorocyclopropylmethyl and 2,2-dichlorocyclopropylmethyl.

The amino C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an amino group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like 2-aminoethyl and 3-aminopropyl.

The nitro C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a nitro group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like nitromethyl and 2-nitroethyl.

The C₁-C₆ haloalkyl group represents a linear or branched alkyl group having 1 to 6 carbon atoms substituted with a halogen atom, and examples thereof include a group like fluoromethyl, chloromethyl, bromomethyl, difluoromethyl, dichloromethyl, trifluoromethyl, trichloromethyl, chlorodifluoromethyl, bromodifluoromethyl, 2-fluoroethyl, 1-chloroethyl, 2-chloroethyl, 1-bromoethyl, 2-bromoethyl, 2,2-difluoroethyl, 1,2-dichloroethyl, 2,2-dichloroethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, 1,1,2,2-tetrafluoroethyl, pentafluoroethyl, 2-bromo-2-chloroethyl, 2-chloro-1,1,2,2-tetrafluoroethyl, 1-chloro-1,2,2,2-tetrafluoroethyl, 1-chloropropyl, 2-chloropropyl, 3-chloropropyl, 2-bromopropyl, 3-bromopropyl, 2-bromo-1-methylethyl, 3-iodopropyl, 2,3-dichloropropyl, 2,3-dibromopropyl, 3,3,3-trifluoropropyl, 3,3,3-trichloropropyl, 3-bromo-3,3-difluoropropyl, 3,3-dichloro-3-fluoropropyl, 2,2,3,3-tetrafluoropropyl, 1-bromo-3,3,3-trifluoropropyl, 2,2,3,3,3-pentafluoropropyl, 2,2,2-trifluoro-1-trifluoromethylethyl, heptafluoropropyl, 1,2,2,2-tetrafluoro-1-trifluoromethylethyl, 2,3-dichloro-1,1,2,3,3-pentafluoropropyl, 2-chlorobutyl, 3-chlorobutyl, 4-chlorobutyl, 2-chloro-1,1-dimethylethyl, 4-bromobutyl, 3-bromo-2-methylpropyl, 2-bromo-1,1-dimethylethyl, 2,2-dichloro-1,1-dimethylethyl, 2-chloro-1-chloromethyl-2-methylethyl, 4,4,4-trifluorobutyl, 3,3,3-trifluoro-1-methylpropyl, 3,3,3-trifluoro-2-methylpropyl, 2,3,4-trichlorobutyl, 2,2,2-trichloro-1,1-dimethylethyl, 4-chloro-4,4-difluorobutyl, 4,4-dichloro-4-fluorobutyl, 4-bromo-4,4-difluorobutyl, 2,4-dibromo-4,4-difluorobutyl, 3,4-dichloro-3,4,4-trifluorobutyl, 3,3-dichloro-4,4,4-trifluorobutyl, 4-bromo-3,3,4,4-tetrafluorobutyl, 4-bromo-3-chloro-3,4,4-trifluorobutyl, 2,2,3,3,4,4-hexafluorobutyl, 2,2,3,4,4,4-hexafluorobutyl, 2,2,2-trifluoro-1-methyl-1-trifluoromethylethyl, 3,3,3-trifluoro-2-trifluoromethylpropyl, 2,2,3,3,4,4,4-heptafluorobutyl, 2,3,3,3-tetrafluoro-2-trifluoromethylpropyl, 1,1,2,2,3,3,4,4-octafluorobutyl, nonafluorobutyl, 4-chloro-1,1,2,2,3,3,4,4-octafluorobutyl, 5-fluoropentyl, 5-chloropentyl, 5,5-difluoropentyl, 5,5-dichloropentyl, 5,5,5-trifluoropentyl, 6,6,6-trifluorohexyl, and 5,5,6,6,6-pentafluorohexyl.

The C₂-C₆ alkenyl group represents, unless specified otherwise, a linear or branched alkenyl group having 2 to 6 carbon atoms, and examples thereof include a group like vinyl, 1-propenyl, isopropenyl, 2-propenyl, 1-butenyl, 1-methyl-1-propenyl, 2-butenyl, 1-methyl-2-propenyl, 3-butenyl, 2-methyl-1-propenyl, 2-methyl-2-propenyl, 1,3-butadienyl, 1-pentenyl, 1-ethyl-2-propenyl, 2-pentenyl, 1-methyl-1-butenyl, 3-pentenyl, 1-methyl-2-butenyl, 4-pentenyl, 1-methyl-3-butenyl, 3-methyl-1-butenyl, 1,2-dimethyl-2-propenyl, 1,1-dimethyl-2-propenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1,2-dimethyl-1-propenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,3-pentadienyl, 1-vinyl-2-propenyl, 1-hexenyl, 1-propyl-2-propenyl, 2-hexenyl, 1-methyl-1-pentenyl, 1-ethyl-2-butenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-4-pentenyl, 1-ethyl-3-butenyl, 1-(isobutyl)vinyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-2-propenyl, 1-(isopropyl)-2-propenyl, 2-methyl-2-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1,3-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1, 5-hexadienyl, 1-vinyl-3-butenyl, and 2,4-hexadienyl.

The C₂-C₆ alkynyl group represents, unless specified otherwise, a linear or branched alkynyl group having 2 to 6 carbon atoms, and examples thereof include a group like ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 1-methyl-2-propynyl, 2-butyne, 3-butyne, 1-pentyne, 1-ethyl-2-propynyl, 2-pentyne, 3-pentyne, 1-methyl-2-butyne, 4-pentyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 1-hexynyl, 1-(n-propyl)-2-propynyl, 2-hexynyl, 1-ethyl-2-butyne, 3-hexynyl, 1-methyl-2-pentyne, 1-methyl-3-pentyne, 4-methyl-1-pentyne, 3-methyl-1-pentyne, 5-hexynyl, 1-ethyl-3-butyne, 1-ethyl-1-methyl-2-propynyl, 1-(isopropyl)-2-propynyl, 1,1-dimethyl-2-butyne, and 2,2-dimethyl-3-butyne.

The C₂-C₆ halolalkenyl group represents, unless specified otherwise, a linear or branched alkenyl group having 2 to 6 carbon atoms substituted with 1 to 11 halogen atoms that are the same or different from each other, and examples thereof include 2-chlorovinyl, 2-bromovinyl, 2-iodovinyl, 3-chloro-2-propenyl, 3-bromo-2-propenyl, 1-chloromethylvinyl, 2-bromo-1-methylvinyl, 1-trifluoromethylvinyl, 3,3,3-trichloro-1-propenyl, 3-bromo-3,3-difluoro-1-propenyl, 2,3,3,3-tetrachloro-1-propenyl, 1-trifluoromethyl-2,2-difluorovinyl, 2-chloro-2-propenyl, 3,3-difluoro-2-propenyl,

2,3,3-trichloro-2-propenyl, 4-bromo-3-chloro-3,4,4-trifluoro-1-butenyl,
 1-bromomethyl-2-propenyl, 3-chloro-2-butenyl, 4,4,4-trifluoro-2-butenyl,
 4-bromo-4,4-difluoro-2-butenyl, 3-bromo-3-butenyl, 3,4,4-trifluoro-3-butenyl,
 3,4,4-tribromo-3-butenyl, 3-bromo-2-methyl-2-propenyl, 3,3-difluoro-2-methyl-2-propenyl,
 5 3,3,3-trifluoro-2-methylpropenyl, 3-chloro-4,4,4-trifluoro-2-butenyl,
 3,3,3-trifluoro-1-methyl-1-propenyl, 3,4,4-trifluoro-1,3-butadienyl, 3,4-dibromo-1-pentenyl,
 4,4-difluoro-3-methyl-3-butenyl, 3,3,4,4,5,5,5-heptafluoro-1-pentenyl, 5,5-difluoro-4-pentenyl,
 4,5,5-trifluoro-4-pentenyl, 3,4,4,4-tetrafluoro-3-trifluoromethyl-1-butenyl,
 4,4,4-trifluoromethyl-3-methyl-2-butenyl, 3,5,5-trifluoro-2,4-pentadienyl,
 10 4,4,5,5,6,6,6-heptafluoro-2-hexenyl, 3,4,4,5,5,5-hexafluoro-3-trifluoromethyl-1-pentenyl,
 4,5,5,5-tetrafluoro-4-trifluoromethyl-2-pentenyl, and
 5-bromo-4,5,5-trifluoro-4-trifluoromethyl-2-pentenyl.

The C₂-C₆ halolalkynyl group represents, unless specified otherwise, a linear or branched
 alkynyl group having 2 to 6 carbon atoms substituted with 1 to 9 halogen atoms that are the same
 15 or different from each other, and examples thereof include 3-chloro-2-propynyl,
 3-bromo-2-propynyl, 3-iodo-2-propynyl, 3-chloro-1-propynyl, and 5-chloro-4-pentynyl.

The C₁-C₆ alkoxy group represents an (alkyl)-O- group having 1 to 6 carbon atoms, wherein
 the alkyl moiety has the same definition as above, and examples thereof include a group like
 methoxy, ethoxy, propoxy, isopropoxy, butoxy, pentyloxy, and hexyloxy.

20 The C₁-C₆ haloalkoxy group represents a linear or branched alkyl-O- group having 1 to 6
 carbon atoms substituted with 1 to 13 halogen atoms that are the same or different from each
 other, wherein the haloalkyl moiety has the same definition as above, and examples thereof
 include a group like chloromethoxy, difluoromethoxy, chlorodifluoromethoxy, trifluoromethoxy,
 and 2,2,2-trifluoroethoxy.

25 The C₁-C₆ alkoxy C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms
 substituted with an alkoxy group having 1 to 6 carbon atoms, wherein the alkyl moiety and
 alkoxy moiety have the same definitions as above, and examples thereof include a group like
 methoxymethyl, ethoxymethyl, isopropoxymethyl, pentyloxymethyl, methoxyethyl, and
 butoxyethyl.

The hydroxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a hydroxy group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like 2-hydroxyethyl and 3-hydroxypropyl.

5 The C₁-C₆ alkoxy C₁-C₆ alkoxy C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy having 1 to 6 carbon atoms substituted with an alkoxy having 1 to 6 carbon atoms, wherein the alkyl moiety and alkoxy moiety have the same definitions as above, and examples thereof include a group like 2-(2-methoxyethoxy)ethyl and 2-(2-ethoxyethoxy)ethyl.

10 The phenyl C₁-C₆ alkoxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with a phenyl, wherein the alkyl moiety and alkoxy moiety have the same definitions as above, and examples thereof include a group like benzyloxymethyl and benzyloxyethyl.

15 The C₁-C₆ haloalkoxy C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with a haloalkoxy group having 1 to 6 carbon atoms, wherein the haloalkoxy moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like chloromethoxymethyl, difluoromethoxymethyl, chlorodifluoromethoxymethyl, trifluoromethoxymethyl, and 2,2,2-trifluoroethoxymethyl.

20 The C₁-C₆ haloalkoxy C₁-C₆ alkoxy group represents, unless specified otherwise, an alkoxy group having 1 to 6 carbon atoms substituted with a haloalkoxy group having 1 to 6 carbon atoms, wherein the haloalkoxy moiety and alkoxy moiety have the same definitions as above, and examples thereof include a group like chloromethoxymethoxy, difluoromethoxymethoxy, chlorodifluoromethoxymethoxy, trifluoromethoxymethoxy, and 2,2,2-trifluoroethoxymethoxy.

25 The C₃-C₆ cycloalkyloxy group represents, unless specified otherwise, a (cycloalkyl)-O- group having 3 to 6 carbon atoms, wherein the cycloalkyl moiety has the same definition as above, and examples thereof include a group like cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, and cyclohexyloxy.

The C₃-C₆ cycloalkyloxy C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with a (cycloalkyl)-O- group having 3 to 6 carbon atoms, wherein the alkyl

moiety and cycloalkyl moiety have the same definitions as above, and examples thereof include a group like cyclopropyloxymethyl, cyclobutyloxymethyl, cyclopentyloxymethyl, and cyclohexyloxymethyl.

The C₃-C₆ cycloalkyl C₁-C₆ alkyloxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with a cycloalkyl group having 3 to 6 carbon atoms, wherein the alkyl moiety, alkoxy moiety, and cycloalkyl moiety have the same definitions as above, and examples thereof include a group like cyclopropylmethyloxymethyl, cyclobutylmethyloxymethyl, cyclopentylmethyloxymethyl, and cyclohexylmethyloxymethyl.

The (R³¹R³²N-C=O) C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (R³¹R³²N-OC-) group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like N,N-dimethylaminocarbonylmethyl, N,N-dimethylaminocarbonylethyl, and N-methyl-N-ethylaminocarbonylmethyl.

The C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxycarbonyl group having 1 to 6 carbon atoms, wherein the alkoxy moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like 2-methoxy-2-oxoethyl, 2-ethoxy-2-oxoethyl, and 2-tert-butoxy-2-oxoethyl.

The C₁-C₆ alkoxycarbonyl C₁-C₆ alkoxy group represents, unless specified otherwise, an alkoxy group having 1 to 6 carbon atoms substituted with an alkoxycarbonyl group having 1 to 6 carbon atoms, wherein the alkoxy moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like a 2-methoxy-2-oxoethoxy group, a 2-ethoxy-2-oxoethoxy group, and a 2-tert-butoxy-2-oxoethoxy group.

The C₁-C₆ alkylcarbonyl group represents an (alkyl (having 1 to 6 carbon atoms))-C(=O)-group, wherein the alkyl moiety has the same definition as above, and examples thereof include acetyl and propionyl.

The C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkylcarbonyl group having 1 to 6 carbon

atoms, wherein the alkylcarbonyl moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like 2-oxopropyl, 3-oxopropyl, and 2-oxobutyl.

The C₁-C₆ alkylcarbonyloxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an (alkyl (having 1 to 6 carbon
5 atoms))-C(=O)O- group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like acetoxymethyl, propionyloxymethyl, isopropionyloxymethyl, and pivaloyloxymethyl.

The C₁-C₆ alkylidene group represents, unless specified otherwise, a divalent alkylidene group having 1 to 6 carbon atoms, wherein a single carbon carries a divalent charge and the alkyl
10 moiety has the same definition as above, and examples thereof include a group like a methylene group, an ethylidene group, and an isopropylidene group.

The C₁-C₆ alkylidene aminooxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with (alkylidene (having 1 to 6 carbon
atoms))=N-O-, wherein the alkylidene moiety and alkyl moiety have the same definitions as
15 above, and examples thereof include a group like methyleneaminooxymethyl, 2-(ethylidene aminooxy)ethyl, and 2-(isopropylidene aminooxy)ethyl.

The C₂-C₆ alkenyloxy group represents, unless specified otherwise, an (alkenyl)-O- group having 2 to 6 carbon atoms, wherein the alkenyl moiety has the same definition as above, and examples thereof include a group like 2-propenyloxy.

20 The C₂-C₆ alkynyloxy group represents, unless specified otherwise, an (alkynyl)-O- group having 2 to 6 carbon atoms, wherein the alkynyl moiety has the same definition as above, and examples thereof include 2-propynyloxy.

The phenyloxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (phenyl)-O- group, wherein the alkyl moiety has
25 the same definition as above, and examples thereof include a group like phenoxymethyl, 2-phenoxyethyl, and 3-phenoxypropyl.

The phenylthio C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (phenyl)-S- group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like phenylthiomethyl,

2-phenylthioethyl, and 3-phenylthiopropyl.

The phenylsulfinyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (phenyl)-SO- group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like phenylsulfinylmethyl,

5 2-phenylsulfinylethyl, and 3-phenylsulfinylpropyl.

The phenylsulfonyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (phenyl)-SO₂- group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like 2-phenylsulfonylethyl, 3-phenylsulfonylpropyl, and 4-phenylsulfonylbutyl.

10 The C₁-C₆ alkoxyimino group represents, unless specified otherwise, an (alkoxy)-N= group having 1 to 6 carbon atoms, wherein the alkoxy moiety has the same definition as above, and examples thereof include methoxyimino and ethoxyimino.

The C₁-C₆ alkoxyimino C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an alkoxyimino group having 1 to 6 carbon atoms, wherein the
15 alkoxyimino moiety and alkyl moiety have the same definitions as above, and examples thereof include methoxyiminomethyl and ethoxyiminomethyl.

The phenoxyimino group represents, unless specified otherwise, a (substituted) (phenoxy)-N= group, and examples thereof include phenoxyimino.

The phenoxyimino C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms
20 substituted with a phenoxyimino group, wherein the phenoxyimino moiety and alkyl moiety have the same definitions as above, and examples thereof include phenoxyiminomethyl.

The di(C₁-C₆ alkoxy) C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms di-substituted with an alkoxy group having 1 to 6 carbon atoms, and examples thereof
25 include (2,2-dimethoxy)ethyl, (3,3-dimethoxy)propyl, (2,2-diethoxy)ethyl group, and a (3,3-diethoxy)propyl.

The formyl C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with a formyl group, wherein the alkyl moiety has the same definition as above, and examples thereof include (2-formyl)ethyl and (3-formyl)propyl.

The C₁-C₆ alkylthio group represents an (alkyl)-S- group having 1 to 6 carbon atoms,

wherein the alkyl moiety has the same definition as above, and examples thereof include methylthio, ethylthio, n-propylthio, and isopropylthio.

The C₁-C₁₀ alkylthio group represents an (alkyl)-S- group having 1 to 10 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include, in addition to those exemplified above for the C₁-C₆ alkylthio group, n-heptylthio, n-octylthio, n-nonylthio, and n-decylthio.

The C₁-C₆ alkylsulfinyl group represents an (alkyl)-SO- group having 1 to 6 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include methylsulfinyl, ethylsulfinyl, n-propylsulfinyl, and isopropylsulfinyl.

The C₁-C₁₀ alkylsulfinyl group represents an (alkyl)-S- group having 1 to 10 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include, in addition to those exemplified above for the C₁-C₆ alkylsulfinyl group, n-heptylsulfinyl, n-octylsulfinyl, n-nonylsulfinyl, and n-decylsulfinyl.

The C₁-C₆ alkylsulfonyl group represents an (alkyl)-SO₂- group having 1 to 6 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, and isopropylsulfonyl.

The C₁-C₁₀ alkylsulfonyl group represents an (alkyl)-SO₂- group having 1 to 10 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include, in addition to those exemplified above for the C₁-C₆ alkylsulfonyl group, n-heptylsulfonyl, n-octylsulfonyl, n-nonylsulfonyl, and n-decylsulfonyl.

The C₂-C₆ alkenylthio group represents an (alkenyl)-S- group having 2 to 6 carbon atoms, wherein the alkenyl moiety has the same definition as above, and examples thereof include a group like allylthio.

The C₂-C₆ alkenylsulfinyl group represents an (alkenyl)-SO- group having 2 to 6 carbon atoms, wherein the alkenyl moiety has the same definition as above, and examples thereof include a group like allylsulfinyl.

The C₂-C₆ alkenylsulfonyl group represents an (alkenyl)-SO₂- group having 2 to 6 carbon atoms, wherein the alkenyl moiety has the same definition as above, and examples thereof include a group like allylsulfonyl.

The C₂-C₆ alkynylthio group represents an (alkynyl)-S- group having 2 to 6 carbon atoms, wherein the alkynyl moiety has the same definition as above, and examples thereof include a group like 2-propynylthio.

5 The C₂-C₆ alkynylsulfinyl group represents an (alkynyl)-SO- group having 2 to 6 carbon atoms, wherein the alkynyl moiety has the same definition as above, and examples thereof include a group like 2-propynylsulfinyl.

The C₂-C₆ alkenylsulfonyl group represents an (alkynyl)-SO₂- group having 2 to 6 carbon atoms, wherein the alkynyl moiety has the same definition as above, and examples thereof include a group like 2-propynylsulfonyl.

10 The C₁-C₁₀ alkylsulfonyloxy group represents an (alkyl)SO₂-O- group having 1 to 10 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include methylsulfonyloxy and ethylsulfonyloxy.

The C₁-C₆ alkylthio C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an alkylthio group having 1 to 6 carbon atoms, wherein the alkyl moiety and
15 alkylthio moiety have the same definitions as above, and examples thereof include methylthiomethyl and ethylthiomethyl.

The C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an alkylsulfinyl group having 1 to 6 carbon atoms, wherein the alkyl moiety and alkylsulfinyl moiety have the same definitions as above, and examples thereof include
20 methylsulfinylmethyl and ethylsulfinylmethyl.

The C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an alkylsulfonyl group having 1 to 6 carbon atoms, wherein the alkyl moiety and alkylsulfonyl moiety have the same definitions as above, and examples thereof include methylsulfonylmethyl and ethylsulfonylmethyl.

25 The C₁-C₆ alkoxy C₁-C₆ alkoxy group represents an alkoxy group having 1 to 6 carbon atoms substituted with an alkoxy having 1 to 6 carbon atoms, wherein the alkoxy moiety has the same definition as above, and examples thereof include a group like methoxymethoxy, ethoxymethoxy, 2-methoxyethoxy, and 2-ethoxyethoxy.

The C₁-C₆ haloalkylthio C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl

group having 1 to 6 carbon atoms substituted with a (haloalkyl)-S- group having 1 to 6 carbon atoms, wherein the alkyl moiety and haloalkyl moiety have the same definitions as above, and examples thereof include a group like difluoromethylthiomethyl and trifluoromethylthiomethyl.

The C₁-C₆ haloalkylsulfinyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (haloalkyl)-SO- group having 1 to 6 carbon atoms, wherein the alkyl moiety and haloalkyl moiety have the same definitions as above, and examples thereof include a group like difluoromethylsulfinylmethyl and trifluoromethylsulfinylmethyl.

The C₁-C₆ haloalkylsulfonyl C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with a (haloalkyl)-SO₂- group having 1 to 6 carbon atoms, wherein the alkyl moiety and haloalkyl moiety have the same definitions as above, and examples thereof include a group like difluoromethylsulfonylmethyl and trifluoromethylsulfonylmethyl.

The C₁-C₆ alkylthio C₁-C₆ alkoxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with an alkylthio group having 1 to 6 carbon atoms, wherein the alkylthio moiety, alkoxy moiety, and alkyl moiety have the same definitions as above, and examples thereof include a group like 2-methylthioethoxymethyl and 2-ethylthioethoxymethyl.

The C₁-C₆ alkylsulfinyl C₁-C₆ alkoxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with an alkynylsulfinyl group having 1 to 6 carbon atoms, wherein the alkylsulfinyl moiety, alkoxy moiety, and alkyl moiety have the same definitions as above, and examples thereof include a group like 2-methylsulfinyl ethoxymethyl and 2-ethylsulfinyl ethoxymethyl.

The C₁-C₆ alkylsulfonyl C₁-C₆ alkoxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with an alkylsulfonyl group having 1 to 6 carbon atoms, wherein the alkylsulfonyl moiety, alkoxy moiety, and alkyl moiety have the same definitions as above, and examples thereof include a group like 2-methylsulfonylethoxymethyl and

2-ethylsulfonylethoxymethyl.

The C₁-C₆ acyl group represents an acyl group derived from C₁-C₆ carboxylic acid, and examples thereof include an acetyl group and a propionyl group.

5 The C₁-C₆ alkylcarbonyl group represents an (alkyl (having 1 to 6 carbon atoms))-C(=O)- group, wherein the alkyl moiety has the same definition as above, and examples thereof include an acetyl group and a propionyl group.

The C₁-C₆ alkylcarbonyloxy group represents an (alkyl (having 1 to 6 carbon atoms))-C(=O)-O- group, wherein the alkyl moiety has the same definition as above, and examples thereof include acetoxy and propionyloxy.

10 The C₁-C₆ haloalkylcarbonyloxy group represents a (haloalkyl (having 1 to 6 carbon atoms))-C(=O)-O- group, wherein the haloalkyl moiety has the same definition as above, and examples thereof include a group like chloromethylcarbonyloxy, difluoromethylcarbonyloxy, chlorodifluoromethylcarbonyloxy, trifluoromethylcarbonyloxy, and 2,2,2-trifluoroethylcarbonyloxy.

15 The C₂-C₆ alkenylcarbonyloxy group represents an (alkenyl (having 2 to 6 carbon atoms))-C(=O)-O- group, wherein the alkenyl moiety has the same definition as above, and examples thereof include a group like 1-propenylcarbonyloxy, 2-propenylcarbonyloxy, 1-butenylcarbonyloxy, and 1-methyl-1-propenylcarbonyloxy.

20 The C₂-C₆ halolalkenylcarbonyloxy group represents a (haloalkenyl (having 2 to 6 carbon atoms))-C(=O)-O- group, wherein the haloalkenyl moiety has the same definition as above, and examples thereof include a group like 3-chloro-2-propenylcarbonyloxy and 3-bromo-2-propenylcarbonyloxy.

The C₂-C₆ alkynylcarbonyloxy group represents an (alkynyl (having 2 to 6 carbon atoms))-C(=O)-O- group, wherein the alkynyl moiety has the same definition as above, and 25 examples thereof include a group like 1-propynylcarbonyloxy and 2-propynylcarbonyloxy.

The C₂-C₆ haloalkynylcarbonyloxy group represents a (haloalkynyl (having 2 to 6 carbon atoms))-C(=O)-O- group, wherein the haloalkynyl moiety has the same definition as above, and examples thereof include a group like 3-chloro-1-propynylcarbonyloxy and 3,3,3-trifluoro-1-propynylcarbonyloxy.

The C₂-C₆ alkylidene amino group represents an alkyl (having 1 to 5 carbon atoms)-CH=N-group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like ethylideneamino and propylideneamino.

5 The di(C₁-C₁₀ alkyl)amino C₁-C₆ alkylidene amino group represents an amino group substituted with an alkylidene group having 1 to 6 carbon atoms substituted with an amino group di-substituted with an alkyl group having 1 to 10 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like a dimethylamino methylidene amino group and a diethylamino methylidene amino group.

10 The C₁-C₁₀ alkylamino group represents an (alkyl)-NH- group having 1 to 10 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include methylamino and ethylamino.

The di(C₁-C₁₀ alkyl)amino group represents an (alkyl)₂N- group, wherein the alkyl moiety has the same definition as above, and examples thereof include dimethylamino, diethylamino, methylethylamino, dipropylamino, and dibutylamino.

15 The mono(C₁-C₆ alkyl)amino group represents an (alkyl)-NH- group having 1 to 6 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like methylamino and ethylamino.

20 The di(C₁-C₆ alkyl)amino group represents an (alkyl (having 1 to 6 carbon atoms))₂N- group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like dimethylamino, diethylamino, methylethylamino, dipropylamino, and dibutylamino.

The C₁-C₆ alkylamino C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an alkylamino group having 1 to 6 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include N-methylaminomethyl and N-methylaminoethyl.

25 The di(C₁-C₆ alkyl)amino C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with an (alkyl (having 1 to 6 carbon atoms))₂N- group, wherein the alkyl moiety has the same definition as above, and examples thereof include N,N-dimethylaminomethyl and N,N-dimethylaminoethyl.

The C₁-C₆ alkoxycarbonyl amino group represents an amino group substituted with an

(alkoxy (having 1 to 6 carbon atoms))-C(=O)- group, wherein the alkoxy moiety has the same definition as above, and examples thereof include methoxycarbonyl amino and ethoxycarbonyl amino.

5 The C₁-C₆ alkylcarbonyl amino group represents, unless specified otherwise, an amino group substituted with an alkylcarbonyl group having 1 to 6 carbon atoms, wherein the alkylcarbonyl moiety has the same definition as above, and examples thereof include a group like formamide, acetamide, and propionamide.

10 The C₁-C₆ alkoxycarbonyl group represents an (alkyl (having 1 to 6 carbon atoms))-O-C(=O)- group, wherein the alkyl moiety has the same definition as above, and examples thereof include methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, and isopropoxycarbonyl.

The C₁-C₁₀ alkylthiocarbonyl group represents an (alkyl (having 1 to 10 carbon atoms))-S-C(=O)- group, wherein the alkyl moiety has the same definition as above, and examples thereof include methylthiocarbonyl and ethylthiocarbonyl.

15 The C₁-C₆ alkoxycarbonyloxy group represents an oxy group substituted with an (alkoxy (having 1 to 6 carbon atoms))-C(=O)- group, wherein the alkoxycarbonyl moiety has the same definition as above, and examples thereof include methoxycarbonyloxy and ethoxycarbonyloxy.

20 The C₁-C₆ haloalkylcarbonyl group represents a (haloalkyl (having 1 to 6 carbon atoms))-C(=O)- group, wherein the haloalkyl moiety has the same definition as above, and examples thereof include chloroacetyl, trifluoroacetyl, pentafluoropropionyl, and difluoromethylthio.

The C₁-C₆ haloalkylthio group represents a (haloalkyl (having 1 to 6 carbon atoms))-S-group, wherein the haloalkyl moiety has the same definition as above, and examples thereof include difluoromethylthio and trifluoromethylthio.

25 The C₁-C₆ haloalkylsulfinyl group represents a (haloalkyl (having 1 to 6 carbon atoms))-SO-group, wherein the haloalkyl moiety has the same definition as above, and examples thereof include trifluoromethylsulfinyl and difluoromethylsulfinyl.

The C₁-C₆ haloalkylsulfonyl group represents a (haloalkyl (having 1 to 6 carbon atoms))-SO₂- group, wherein the haloalkyl moiety has the same definition as above, and

examples thereof include chloromethylsulfonyl, difluoromethylsulfonyl, and trifluoromethylsulfonyl.

The C₁-C₆ haloalkylsulfonyloxy group represents a (haloalkyl (having 1 to 6 carbon atoms))-SO₂-O- group, wherein the haloalkyl moiety has the same definition as above, and examples thereof include chloromethylsulfonyloxy and trifluoromethylsulfonyloxy.

The mono(C₁-C₆ alkyl)aminocarbonyl group represents an (alkyl (having 1 to 6 carbon atoms))-NH-C(=O)- group, wherein the alkyl moiety has the same definition as above, and examples thereof include methylaminocarbonyl and ethylaminocarbonyl.

The di(C₁-C₆ alkyl)aminocarbonyl group represents an (alkyl (having 1 to 6 carbon atoms))₂N-C(=O)- group, wherein the alkyl moiety has the same definition as above, and examples thereof include a group like dimethylaminocarbonyl, diethylaminocarbonyl, methylethylaminocarbonyl, dipropylaminocarbonyl, and dibutylaminocarbonyl.

The cyano C₁-C₆ alkyl group represents a cyano alkyl group having 1 to 6 carbon atoms, wherein the alkyl moiety has the same definition as above, and examples thereof include cyanomethyl and cyanoethyl.

The cyano C₁-C₆ alkoxy group represents an alkoxy group having 1 to 6 carbon atoms substituted with a cyano group, wherein the alkoxy moiety has the same definition as above, and examples thereof include a group like 2-cyanoethoxy and 3-cyanopropoxy.

The cyano C₁-C₆ alkoxy C₁-C₆ alkyl group represents, unless specified otherwise, an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with a cyano group, wherein the alkoxy moiety and alkyl moiety have the same definitions as above, and examples thereof include a group like 2-cyanoethoxymethyl and 3-cyanopropoxymethyl.

The phenyl C₁-C₆ alkyl group represents an alkyl group having 1 to 6 carbon atoms substituted with a phenyl group, wherein the alkyl moiety has the same definition as above, and examples thereof include benzyl, phenethyl, and phenylpropyl.

The phenyl C₂-C₆ alkenyl group represents an alkenyl group having 2 to 6 carbon atoms substituted with a phenyl group, wherein the alkenyl moiety has the same definition as above, and examples thereof include styryl and cinnamyl.

The phenyl C₂-C₆ alkynyl group represents an alkynyl group having 2 to 6 carbon atoms substituted with a phenyl group, wherein the alkynyl moiety has the same definition as above, and examples thereof include (2-phenyl)ethynyl and 2-(3-phenyl)ethynyl.

5 The phenylcarbonyloxy group represents a (phenyl)-C(=O)-O- group and examples thereof include a phenylcarbonyloxy group.

The phenylcarbonyl C₁-C₆ alkoxy group represents an alkoxy group having 1 to 6 carbon atoms substituted with a (phenyl)-C(=O) group and examples thereof include phenylcarbonylmethoxy.

The phenylthio group represents a phenyl-S- group.

10 The phenylsulfinyl group represents a phenyl-SO- group.

The phenylsulfonyl group represents a phenyl-SO₂- group.

The phenylsulfonyloxy group represents a phenyl-SO₂-O- group.

The benzylthio group represents a benzyl-S- group.

The benzylsulfinyl group represents a benzyl-SO- group.

15 The benzylsulfonyl group represents a benzyl-SO₂- group.

The benzylsulfonyloxy group represents a benzyl-SO₂-O- group.

As a group constituting a C₃-C₆ alkylene group, 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting a carbonyl group, and the C₃-C₆ alkylene group is a linear or branched divalent alkylene group having 3 to 6 carbon atoms, and 1 to 3 carbon atoms in the alkylene group may be substituted with an atom or a group of atoms selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting a carbonyl group, and examples thereof include a trimethylene group, a propylene group, a butylene group, a methylenedioxy group, and an ethylenedioxy group. Preferred
20 examples of the alkylene group include a C₁-C₃ alkylendioxy group.

25 Examples of the heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom include furan, thiophene, pyrrole, pyrazole, imidazole, pyridine, pyrimidine, pyrazine, pyridazine, pyrrolidine, piperidine, piperazine, morpholine, thiomorpholine,

benzofuran, benzothiophene, indole, benzoxazole, benzothiazole, benzimidazole, isoxazole, isoxazoline, oxazole, oxazoline, isothiazole, isothiazoline, thiazole, tetrahydrofuran, and thiazoline. Preferred examples of the heterocyclic group include pyridine, pyrimidine, pyrazine, thiophene, pyrazole, isoxazole, morpholine, thiomorpholine (sulfur atom of thiomorpholine may be bonded with one or two oxygen atoms), piperidine, pyridazine, piperazine, and tetrahydrofuran. More preferred examples of the heterocyclic group include pyridine, pyrimidine, pyrazine, thiophene, pyrazole, isoxazole, morpholine, thiomorpholine (sulfur atom of thiomorpholine may be bonded with one or two oxygen atoms), and piperidine.

The heterocyclic oxy group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and optionally selected from an oxygen atom, a sulfur atom, and a nitrogen atom represents, unless specified otherwise, a group in which the oxygen atom is substituted with a heterocycle having the same definition as above, and examples thereof include (tetrahydrofuran-2-yl)oxy, (4,5-dihydroisoxazol-5-yl)oxy, (isoxazol-5-yl)oxy, and a (thiophen-2-yl)oxy group.

The C₁-C₆ alkyl group substituted with a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom represents an alkyl group having 1 to 6 carbon atoms substituted with a heterocycle wherein the alkyl moiety and heterocyclic moiety have the same definitions as above, and examples thereof include (2-furan)methyl, (3-furan)methyl, (2-thiophene)methyl, and (3-thiophene)methyl.

The C₁-C₆ alkyl group substituted with a heterocyclic oxy group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom represents an alkyl group having 1 to 6 carbon atoms substituted with a heterocyclic oxy group wherein the alkyl moiety and heterocyclic moiety have the same definitions as above, and examples thereof include (tetrahydrofuran-2-yl)oxymethyl, (4,5-dihydroisoxazol-5-yl)oxymethyl, (isoxazol-5-yl)oxymethyl, and (thiophen-2-yl)oxymethyl.

The C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic oxy group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and

selected from an oxygen atom, a sulfur atom, and a nitrogen atom represents an alkyl group having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with a heterocyclic oxy group wherein the alkyl moiety, alkoxy moiety, and heterocyclic moiety have the same definitions as above, and examples thereof include

5 (tetrahydrofuran-2-yl)oxymethoxymethyl, (4,5-dihydroisoxazol-5-yl)oxyethoxymethyl, (isoxazol-5-yl)oxymethoxymethyl, and (thiophen-2-yl)oxyethoxymethyl.

The C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom represents an alkyl group
10 having 1 to 6 carbon atoms substituted with an alkoxy group having 1 to 6 carbon atoms substituted with a heterocyclic group wherein the alkyl moiety, alkoxy moiety, and heterocyclic moiety have the same definitions as above, and examples thereof include tetrahydrofurfuryloxyethyl and tetrahydrofurfuryloxymethyl.

The C₁-C₆ alkoxy group substituted with a heterocyclic group having 3 to 10 carbon atoms
15 and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom represents an alkoxy group having 1 to 6 carbon atoms substituted with a heterocyclic group wherein the heterocyclic moiety and alkoxy moiety have the same definitions as above, and examples thereof include a 6-methyl-2-pyridinemethoxy group and a tetrahydrofurfuryloxy group.

20 Alkali metal includes sodium, potassium, and the like.


Next, specific examples of the compound of the invention represented by Formula 1 are described in Table 1 to Table 43. However, the invention is not limited to those compounds.

In the present Description, the following descriptions included in the tables indicate the corresponding group, respectively, as shown below.

25 For example, Me represents a methyl group, Et represents an ethyl group, Pr-n represents a n-propyl group, Pr-i represents an isopropyl group, Pr-c represents a cyclopropyl group, Bu-n represents a n-butyl group, Bu-s represents a secondary butyl group, Bu-i represents an isobutyl group, Bu-t represents a tertiary butyl group, Bu-c represents a cyclobutyl group, Pen-n represents a n-pentyl group, Pen-c represents a cyclopentyl group, Hex-n represents a n-hexyl group, Hex-c

represents a cyclohexyl group, Ac represents an acetyl group, Ph represents a phenyl group, Bn represents a benzyl group, Ts represents a p-toluene sulfonyl group, pyridyl represents a pyridyl group, and pyrimidinyl represents a pyrimidinyl group. Further, Ph(2-OMe) represents a 2-methoxyphenyl group, CH₂Ph(2-OMe) represents a 2-methoxybenzyl group, and Ph(3,4-Cl₂) represents a 3,4-dichlorophenyl group.

[Table 1]

					
Compound No.	R ¹	R ²	Y	Z	R ⁴
I-1	Me	Me	O	O	OH
I-2	Et	Me	O	O	OH
I-3	Pr ⁿ	Me	O	O	OH
I-4	Pr ⁱ	Me	O	O	OH
I-5	Bu ⁿ	Me	O	O	OH
I-6	Bu ⁱ	Me	O	O	OH
I-7	Bu ^s	Me	O	O	OH
I-8	Bu ^t	Me	O	O	OH
I-9	Hex ⁿ	Me	O	O	OH
I-10	CH ₂ CF ₃	Me	O	O	OH
I-11	CH ₂ CH=CH ₂	Me	O	O	OH
I-12	CH ₂ C(Me)=CH ₂	Me	O	O	OH
I-13	CH ₂ CH ₂ CH=CMe ₂	Me	O	O	OH
I-14	CH ₂ C≡CH	Me	O	O	OH
I-15	CH ₂ C≡CCH ₃	Me	O	O	OH
I-16	Pr ^c	Me	O	O	OH
I-17	Bu ^c	Me	O	O	OH
I-18	Pen ^c	Me	O	O	OH
I-19	Hex ^c	Me	O	O	OH
I-20	CH ₂ Pr ^c	Me	O	O	OH
I-21	CH ₂ Bu ^c	Me	O	O	OH
I-22	CH ₂ Pen ^c	Me	O	O	OH
I-23	CH ₂ Hex ^c	Me	O	O	OH
I-24	CH ₂ CH=CCl ₂	Me	O	O	OH
I-25	CH ₂ CCl=CHCl	Me	O	O	OH
I-26	CH ₂ CH ₂ CH=CCl ₂	Me	O	O	OH
I-27	CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O	OH
I-28	CH ₂ CH ₂ CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O	OH
I-29	CH ₂ CH=CF ₂	Me	O	O	OH
I-30	CH ₂ CH ₂ OMe	Me	O	O	OH
I-31	CH ₂ CH ₂ OEt	Me	O	O	OH
I-32	CH(Me)CH ₂ OMe	Me	O	O	OH
I-33	CH ₂ CH ₂ OCH ₂ CH ₂ OMe	Me	O	O	OH
I-34	CH ₂ CH ₂ OPr ⁿ	Me	O	O	OH
I-35	CH ₂ CH ₂ OPr ⁱ	Me	O	O	OH
I-36	CH ₂ CH ₂ OPr ^c	Me	O	O	OH
I-37	CH ₂ CH ₂ OBu ^c	Me	O	O	OH
I-38	CH ₂ CH ₂ OPen ^c	Me	O	O	OH
I-39	CH ₂ CH ₂ OHex ^c	Me	O	O	OH
I-40	CH ₂ CH ₂ OCH ₂ CF ₃	Me	O	O	OH
I-41	CH ₂ CH ₂ CH ₂ OMe	Me	O	O	OH

[Table 2]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-42	CH=CHMe	Me	O	O	OH
I-43	CH ₂ SMe	Me	O	O	OH
I-44	CH ₂ SP _n	Me	O	O	OH
I-45	CH ₂ CH ₂ SMe	Me	O	O	OH
I-46	CH ₂ SOMe	Me	O	O	OH
I-47	CH ₂ SO ₂ Me	Me	O	O	OH
I-48	CH ₂ CH ₂ CH ₂ SMe	Me	O	O	OH
I-49	CH ₂ CH ₂ CH ₂ SO ₂ Me	Me	O	O	OH
I-50	Ph	Me	O	O	OH
I-51	Ph(2-Cl)	Me	O	O	OH
I-52	Ph(3-Cl)	Me	O	O	OH
I-53	Ph(4-Cl)	Me	O	O	OH
I-54	Ph(2-F)	Me	O	O	OH
I-55	Ph(3-F)	Me	O	O	OH
I-56	Ph(4-F)	Me	O	O	OH
I-57	Ph(2-Me)	Me	O	O	OH
I-58	Ph(3-Me)	Me	O	O	OH
I-59	Ph(4-Me)	Me	O	O	OH
I-60	Ph(2-OMe)	Me	O	O	OH
I-61	Ph(3-OMe)	Me	O	O	OH
I-62	Ph(4-OMe)	Me	O	O	OH
I-63	Ph(2-CF ₃)	Me	O	O	OH
I-64	Ph(3-CF ₃)	Me	O	O	OH
I-65	Ph(4-CF ₃)	Me	O	O	OH
I-66	Ph(2-NO ₂)	Me	O	O	OH
I-67	Ph(3-NO ₂)	Me	O	O	OH
I-68	Ph(4-NO ₂)	Me	O	O	OH
I-69	Ph(2-OCF ₃)	Me	O	O	OH
I-70	Ph(3-OCF ₃)	Me	O	O	OH
I-71	Ph(4-OCF ₃)	Me	O	O	OH
I-72	Ph(2-CN)	Me	O	O	OH
I-73	Ph(3-CN)	Me	O	O	OH
I-74	Ph(4-CN)	Me	O	O	OH
I-75	Ph(3,4-F ₂)	Me	O	O	OH
I-76	Ph(3,5-F ₂)	Me	O	O	OH
I-77	Ph(2,3-F ₂)	Me	O	O	OH
I-78	Ph(2,4-F ₂)	Me	O	O	OH
I-79	Ph(2,5-F ₂)	Me	O	O	OH
I-80	Ph(2,6-F ₂)	Me	O	O	OH
I-81	Ph(3,4-Cl ₂)	Me	O	O	OH
I-82	Ph(3,5-Cl ₂)	Me	O	O	OH
I-83	Ph(2,3-Cl ₂)	Me	O	O	OH
I-84	Ph(2,4-Cl ₂)	Me	O	O	OH
I-85	Ph(2,5-Cl ₂)	Me	O	O	OH














[Table 3]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-86	Ph(2,6-Cl ₂)	Me	O	O	OH
I-87	Ph(3,4-Me ₂)	Me	O	O	OH
I-88	Ph(3,5-Me ₂)	Me	O	O	OH
I-89	Ph(2,3-Me ₂)	Me	O	O	OH
I-90	Ph(2,4-Me ₂)	Me	O	O	OH
I-91	Ph(2,5-Me ₂)	Me	O	O	OH
I-92	Ph(2,6-Me ₂)	Me	O	O	OH
I-93	Ph(3,4-OMe ₂)	Me	O	O	OH
I-94	Ph(3,5-OMe ₂)	Me	O	O	OH
I-95	Ph(2,3-OMe ₂)	Me	O	O	OH
I-96	Ph(2,4-OMe ₂)	Me	O	O	OH
I-97	Ph(2,5-OMe ₂)	Me	O	O	OH
I-98	Ph(2,6-OMe ₂)	Me	O	O	OH
I-99	Ph(3-F-4-OMe)	Me	O	O	OH
I-100	Ph(3-F-5-OMe)	Me	O	O	OH
I-101	Ph(2-F-3-OMe)	Me	O	O	OH
I-102	Ph(2-F-4-OMe)	Me	O	O	OH
I-103	Ph(2-F-5-OMe)	Me	O	O	OH
I-104	Ph(2-F-6-OMe)	Me	O	O	OH
I-105	Ph(3-F-4-Me)	Me	O	O	OH
I-106	Ph(3-F-5-Me)	Me	O	O	OH
I-107	Ph(2-F-3-Me)	Me	O	O	OH
I-108	Ph(2-F-4-Me)	Me	O	O	OH
I-109	Ph(2-F-5-Me)	Me	O	O	OH
I-110	Ph(2-F-6-Me)	Me	O	O	OH
I-111	Ph(3-OMe-4-F)	Me	O	O	OH
I-112	Ph(2-OMe-3-F)	Me	O	O	OH
I-113	Ph(2-OMe-4-F)	Me	O	O	OH
I-114	Ph(2-OMe-5-F)	Me	O	O	OH
I-115	Ph(3-Me-4-F)	Me	O	O	OH
I-116	Ph(2-Me-3-F)	Me	O	O	OH
I-117	Ph(2-Me-4-F)	Me	O	O	OH
I-118	Ph(2-Me-5-F)	Me	O	O	OH
I-119	Ph(3-Cl-4-OMe)	Me	O	O	OH
I-120	Ph(3-Cl-5-OMe)	Me	O	O	OH
I-121	Ph(2-Cl-3-OMe)	Me	O	O	OH
I-122	Ph(2-Cl-4-OMe)	Me	O	O	OH
I-123	Ph(2-Cl-5-OMe)	Me	O	O	OH
I-124	Ph(2-Cl-6-OMe)	Me	O	O	OH
I-125	Ph(3-Cl-4-Me)	Me	O	O	OH
I-126	Ph(3-Cl-5-Me)	Me	O	O	OH
I-127	Ph(2-Cl-3-Me)	Me	O	O	OH
I-128	Ph(2-Cl-4-Me)	Me	O	O	OH
I-129	Ph(2-Cl-5-Me)	Me	O	O	OH

















[Table 4]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-130	Ph(2-Cl-6-Me)	Me	O	O	OH
I-131	Ph(3-OMe-4-Cl)	Me	O	O	OH
I-132	Ph(2-OMe-3-Cl)	Me	O	O	OH
I-133	Ph(2-OMe-4-Cl)	Me	O	O	OH
I-134	Ph(2-OMe-5-Cl)	Me	O	O	OH
I-135	Ph(3-Me-4-Cl)	Me	O	O	OH
I-136	Ph(2-Me-3-Cl)	Me	O	O	OH
I-137	Ph(2-Me-4-Cl)	Me	O	O	OH
I-138	Ph(2-Me-5-Cl)	Me	O	O	OH
I-139	Ph(3-F-4-Cl)	Me	O	O	OH
I-140	Ph(3-F-5-Cl)	Me	O	O	OH
I-141	Ph(2-F-3-Cl)	Me	O	O	OH
I-142	Ph(2-F-4-Cl)	Me	O	O	OH
I-143	Ph(2-F-5-Cl)	Me	O	O	OH
I-144	Ph(2-F-6-Cl)	Me	O	O	OH
I-145	Ph(3-Cl-4-F)	Me	O	O	OH
I-146	Ph(2-Cl-3-F)	Me	O	O	OH
I-147	Ph(2-Cl-4-F)	Me	O	O	OH
I-148	Ph(2-Cl-5-F)	Me	O	O	OH
I-149	Ph(3-Me-4-OMe)	Me	O	O	OH
I-150	Ph(3-Me-5-OMe)	Me	O	O	OH
I-151	Ph(2-Me-3-OMe)	Me	O	O	OH
I-152	Ph(2-Me-4-OMe)	Me	O	O	OH
I-153	Ph(2-Me-5-OMe)	Me	O	O	OH
I-154	Ph(2-Me-6-OMe)	Me	O	O	OH
I-155	Ph(3-OMe-4-Me)	Me	O	O	OH
I-156	Ph(2-OMe-3-Me)	Me	O	O	OH
I-157	Ph(2-OMe-4-Me)	Me	O	O	OH
I-158	Ph(2-OMe-5-Me)	Me	O	O	OH
I-159	Ph(3-CN-4-OMe)	Me	O	O	OH
I-160	Ph(3-OMe-4-CN)	Me	O	O	OH
I-161	Ph(3-Me-4-CN)	Me	O	O	OH
I-162	Ph(3-CN-4-Me)	Me	O	O	OH
I-163	Ph(3-NO ₂ -4-OMe)	Me	O	O	OH
I-164	Ph(3-OMe-4-NO ₂)	Me	O	O	OH
I-165	Ph(3-Me-4-NO ₂)	Me	O	O	OH
I-166	Ph(3-NO ₂ -4-Me)	Me	O	O	OH
I-167	Ph(3,5-F ₂ -4-OMe)	Me	O	O	OH
I-168	Ph(3,5-F ₂ -4-Me)	Me	O	O	OH
I-169	Ph(3,4,5-(OMe) ₃)	Me	O	O	OH




[Table 5]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-170		Me	O	O	OH
I-171		Me	O	O	OH
I-172		Me	O	O	OH
I-173		Me	O	O	OH
I-174		Me	O	O	OH
I-175		Me	O	O	OH
I-176		Me	O	O	OH
I-177		Me	O	O	OH
I-178		Me	O	O	OH
I-179		Me	O	O	OH
I-180		Me	O	O	OH
I-181		Me	O	O	OH
I-182		Me	O	O	OH
I-183		Me	O	O	OH
I-184		Me	O	O	OH






[Table 6]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-185		Me	O	O	OH
I-186		Me	O	O	OH
I-187		Me	O	O	OH
I-188		Me	O	O	OH
I-189		Me	O	O	OH
I-190		Me	O	O	OH
I-191		Me	O	O	OH
I-192		Me	O	O	OH
I-193		Me	O	O	OH
I-194		Me	O	O	OH
I-195		Me	O	O	OH
I-196		Me	O	O	OH
I-197		Me	O	O	OH
I-198		Me	O	O	OH
I-199		Me	O	O	OH
I-200		Me	O	O	OH













[Table 7]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-201		Me	O	O	OH
I-202		Me	O	O	OH
I-203		Me	O	O	OH
I-204		Me	O	O	OH
I-205	CH ₂ Ph	Me	O	O	OH
I-206	CH ₂ CH ₂ Ph	Me	O	O	OH
I-207	CH ₂ CH ₂ CH ₂ Ph	Me	O	O	OH
I-208	CH ₂ CH=CHPh	Me	O	O	OH
I-209	CH ₂ C≡CPh	Me	O	O	OH
I-210	CH ₂ CH=NOMe	Me	O	O	OH
I-211	CH ₂ CH=NOEt	Me	O	O	OH
I-212	CH ₂ CH=NOPr-n	Me	O	O	OH
I-213	CH ₂ CH=NOPh	Me	O	O	OH
I-214	CH ₂ CH(OMe) ₂	Me	O	O	OH
I-215	CH ₂ CHO	Me	O	O	OH
I-216	NH ₂	Me	O	O	OH
I-217	NHMe	Me	O	O	OH
I-218	NHEt	Me	O	O	OH
I-219	NHPr-n	Me	O	O	OH
I-220	NHPr-i	Me	O	O	OH
I-221	NHBu-n	Me	O	O	OH
I-222	NHBu-i	Me	O	O	OH
I-223	NHBu-s	Me	O	O	OH
I-224	NHCH ₂ Pr-c	Me	O	O	OH
I-225	NHPen-n	Me	O	O	OH
I-226	NHHex-n	Me	O	O	OH
I-227	NHCH ₂ CH ₂ CH ₂ Cl	Me	O	O	OH
I-228	NHCH ₂ CH ₂ CH ₂ F	Me	O	O	OH
I-229	NHCH ₂ CH ₂ OMe	Me	O	O	OH
I-230	NMe ₂	Me	O	O	OH
I-231	NEt ₂	Me	O	O	OH
I-232	N(Pr-n) ₂	Me	O	O	OH
I-233	N(Bu-n) ₂	Me	O	O	OH
I-234	N(Me)Et	Me	O	O	OH
I-235	N(Me)CH ₂ CH ₂ OMe	Me	O	O	OH
I-236	NHPh	Me	O	O	OH
I-237	NHCH ₂ Ph	Me	O	O	OH
I-238	N=CMe ₂	Me	O	O	OH
I-239	N=CEt ₂	Me	O	O	OH






[Table 8]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-240	N=CHNMe ₂	Me	O	O	OH
I-241	NHC(=O)Me	Me	O	O	OH
I-242	N[C(=O)Me] ₂	Me	O	O	OH
I-243	NHC(=O)OMe	Me	O	O	OH
I-244	N[C(=O)OMe] ₂	Me	O	O	OH
I-245	NHSO ₂ Me	Me	O	O	OH
I-246	NHSO ₂ Ph	Me	O	O	OH
I-247	NHSO ₂ CH ₂ Ph	Me	O	O	OH
I-248	OMe	Me	O	O	OH
I-249	OE _t	Me	O	O	OH
I-250	OP _{r-n}	Me	O	O	OH
I-251	OP _{r-i}	Me	O	O	OH
I-252	OCH ₂ P _{r-c}	Me	O	O	OH
I-253	OCH ₂ Cl	Me	O	O	OH
I-254	OCHCl ₂	Me	O	O	OH
I-255	OCCl ₃	Me	O	O	OH
I-256	OCH ₂ F	Me	O	O	OH
I-257	OCHF ₂	Me	O	O	OH
I-258	OCF ₃	Me	O	O	OH
I-259	Ph	Et	O	O	OH
I-260	Ph	P _{r-i}	O	O	OH
I-261	Ph	CHF ₂	O	O	OH
I-262	Ph	Ph	O	O	OH
I-263	Ph	Me	O	S	OH
I-264	Ph	Me	S	S	OH
I-265	Me	Me	O	S	OH
I-266	Me	Me	S	S	OH
I-267	Ph	Me	O	O	SPh
I-268	Ph(4-OEt)	Me	O	O	OH
I-269	Ph(2-Ph)	Me	O	O	OH
I-270	Ph(3-Ph)	Me	O	O	OH
I-271	Ph(4-Ph)	Me	O	O	OH
I-272		Me	O	O	OH
I-273		Me	O	O	OH
I-274	Me		O	O	OH
I-275	Et		O	O	OH
I-276		Me	O	O	OH








[Table 9]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-277		Me	O	O	OH
I-278		Me	O	O	OH
I-279		Me	O	O	OH
I-280		Me	O	O	OH
I-281		Me	O	O	OH
I-282	Ph(2-Me-4-Br)	Me	O	O	OH
I-283	Ph(2-Me-4-I)	Me	O	O	OH
I-284	Ph(2-Me-5-CF ₃)	Me	O	O	OH
I-285	Ph(2-Me-6-OCF ₃)	Me	O	O	OH
I-286	Ph(2-Pr-i)	Me	O	O	OH
I-287		Me	O	O	OH
I-288	Ph(2-Et)	Me	O	O	OH
I-289		Me	O	O	OH
I-290		Me	O	O	OH
I-291		Me	O	S	OH
I-292		Me	O	O	OH
I-293		Me	O	O	OH
I-294	CH ₂ COOBu-t	Me	O	O	OH
I-295	(C ₇ H ₁₄)CH ₃	Me	O	O	OH
I-296	(C ₉ H ₁₈)CH ₃	Me	O	O	OH
I-297	Ph(2-F,4-Cl,5-OMe)	Me	O	O	OH
I-298	Ph(2,3,4-(OMe) ₃)	Me	O	O	OH
I-299	Ph(3,5-Cl ₂ -4-OMe)	Me	O	O	OH
I-300	Ph(3,5-Cl ₂ -4-SMe)	Me	O	O	OH
I-301	Ph(3,5-Cl ₂ -4-SO ₂ Me)	Me	O	O	OH
I-302	Ph(3,4,5-F ₃)	Me	O	O	OH
I-303		Me	O	O	OH



[Table 10]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-304		Me	O	O	OH
I-305		Me	O	O	OH
I-306	Bu-n		O	O	OH
I-307	CH ₂ CH(CH ₃) ₂		O	O	OH
I-308	Ph	Pen-n	O	O	OH
I-309	H	Me	O	O	OH
I-310	CH ₂ C≡CF	Me	O	O	OH
I-311		Me	O	O	OH
I-312		Me	O	O	OH
I-313	CH ₂ NH ₂	Me	O	O	OH
I-314	CH ₂ NO ₂	Me	O	O	OH
I-315	CH ₂ NHCH ₃	Me	O	O	OH
I-316	CH ₂ N(CH ₃) ₂	Me	O	O	OH
I-317	CH ₂ SCH ₂ CF ₃	Me	O	O	OH
I-318	CH ₂ SOCH ₂ CF ₃	Me	O	O	OH
I-319	CH ₂ SO ₂ CH ₂ CF ₃	Me	O	O	OH
I-320	CH ₂ OH	Me	O	O	OH
I-321	CH ₂ OBn	Me	O	O	OH
I-322	CH ₂ OCH ₂ Pr-c	Me	O	O	OH
I-323	CH ₂ OPh	Me	O	O	OH
I-324	CH ₂ SPh	Me	O	O	OH
I-325	CH ₂ SOPh	Me	O	O	OH
I-326	CH ₂ SO ₂ Ph	Me	O	O	OH
I-327	CH ₂ CON(CH ₃) ₂	Me	O	O	OH
I-328	CH ₂ COCH ₃	Me	O	O	OH
I-329	CH ₂ OCOCH ₃	Me	O	O	OH
I-330	CH ₂ ON=CHCH ₃	Me	O	O	OH
I-331	C ₂ H ₄ OC ₂ H ₄ SCH ₃	Me	O	O	OH
I-332	C ₂ H ₄ OC ₂ H ₄ SOCH ₃	Me	O	O	OH
I-333	C ₂ H ₄ OC ₂ H ₄ SO ₂ CH ₃	Me	O	O	OH
I-334	CH ₂ OCH ₂ CN	Me	O	O	OH
I-335	CH ₂ CN	Me	O	O	OH
I-336	OCH ₂ CH=CH ₂	Me	O	O	OH
I-337	OCH ₂ C≡CH	Me	O	O	OH
I-338	OPr-c	Me	O	O	OH












[Table 11]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-339		Me	O	O	OH
I-340		Me	O	O	OH
I-341		Me	O	O	OH
I-342		Me	O	O	OH
I-343		Me	O	O	OH
I-344	Ph	H	O	O	OH
I-345	Ph	CH ₂ CH=CH ₂	O	O	OH
I-346	Ph	CH ₂ C≡CH	O	O	OH
I-347	Ph	Pr-c	O	O	OH
I-348	Ph	CH ₂ CH=CF ₂	O	O	OH
I-349	Ph	CH ₂ C≡CF	O	O	OH
I-350	Ph	C ₂ H ₄ OCH ₃	O	O	OH
I-351	Ph	C ₂ H ₄ OC ₂ H ₅	O	O	OH
I-352	Ph	CH(Me)OEt	O	O	OH
I-353	Ph	CH ₂ OPr-c	O	O	OH
I-354	Ph	CH(OCH ₃) ₂	O	O	OH
I-355	Ph	CH ₂ Ph	O	O	OH
I-356	Ph		O	O	OH
I-357	Ph		O	O	OH
I-358	Ph	Me	O	O	NH ₂
I-359	Ph	Me	O	O	Cl
I-360	Ph	Me	O	O	CN
I-361	Ph	Me	O	O	NCS
I-362	Ph	Me	O	O	NCO
I-363	Ph	Me	O	O	OCO ₂ H
I-364	Ph	Me	O	O	OCO ₂ CH ₃
I-365	Ph	Me	O	O	OCO ₂ CH ₂ Ph
I-366	Ph	Me	O	O	OMe
I-367	Ph	Me	O	O	OEt
I-368	Ph	Me	O	O	OPr
I-369	Ph	Me	O	O	OCH ₂ CH=CH ₂
I-370	Ph	Me	O	O	OCH ₂ C≡CH
I-371	Ph	Me	O	O	OPr-c
I-372	Ph	Me	O	O	OBu-c
I-373	Ph	Me	O	O	OPen-c
















[Table 12]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-374	Ph	Me	O	O	OH _{Hex-c}
I-375	Ph	Me	O	O	OCH ₂ CN
I-376	Ph	Me	O	O	OCH ₂ Pr-c
I-377	Ph	Me	O	O	OCOCH ₃
I-378	Ph	Me	O	O	OCOCCl ₃
I-379	Ph	Me	O	O	OCOCH=CH ₂
I-380	Ph	Me	O	O	OCOCH=CF ₂
I-381	Ph	Me	O	O	OCOCH ₂ C≡CH
I-382	Ph	Me	O	O	OCOCH ₂ C≡CF
I-383	Ph	Me	O	O	OCH ₂ CO ₂ CH ₃
I-384	Ph	Me	O	O	OPh
I-385	Ph	Me	O	O	OCH ₂ Ph
I-386	Ph	Me	O	O	OCOPh
I-387	Ph	Me	O	O	OCOCH ₂ Ph
I-388	Ph	Me	O	O	OCH ₂ COPh
I-389	Ph	Me	O	O	OSO ₂ CH ₂ CF ₃
I-390	Ph	Me	O	O	OSO ₂ CH ₂ Ph
I-391	Ph	Me	O	O	SCH ₃
I-392	Ph	Me	O	O	SOCH ₃
I-393	Ph	Me	O	O	SO ₂ CH ₃
I-394	Ph	Me	O	O	SCH ₂ CF ₃
I-395	Ph	Me	O	O	SOCH ₂ CF ₃
I-396	Ph	Me	O	O	SO ₂ CH ₂ CF ₃
I-397	Ph	Me	O	O	SCH ₂ CH=CH ₂
I-398	Ph	Me	O	O	SOCH ₂ CH=CH ₂
I-399	Ph	Me	O	O	SO ₂ CH ₂ CH=CH ₂
I-400	Ph	Me	O	O	SCH ₂ CH≡CH
I-401	Ph	Me	O	O	SOCH ₂ CH≡CH
I-402	Ph	Me	O	O	SO ₂ CH ₂ CH≡CH
I-403	Ph	Me	O	O	SCH ₂ Ph
I-404	Ph	Me	O	O	SOPh
I-405	Ph	Me	O	O	SOCH ₂ Ph
I-406	Ph	Me	O	O	SO ₂ Ph
I-407	Ph	Me	O	O	SO ₂ CH ₂ Ph
I-408	Ph	Me	O	O	NHCH ₃
I-409	Ph	Me	O	O	N(CH ₃) ₂
I-410	Ph	Me	O	O	NHCOCH ₃
I-411	Ph	Me	O	O	OCH ₂ CH ₂ - 
I-412	Ph	Me	O	O	- 











[Table 13]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-413	Ph	Me	O	O	
I-414	Ph	Me	O	O	
I-415	Ph	Me	O	O	
I-416	Ph	Me	O	O	
I-417	(4-Pr-c)Ph	Me	O	O	OH
I-418	(4-CH ₂ Pr-c)Ph	Me	O	O	OH
I-419	(4-CH ₂ =CHCH ₂)Ph	Me	O	O	OH
I-420	(4-CH≡CCH ₂)Ph	Me	O	O	OH
I-421	(4-CH ₂ CH=CF ₂)Ph	Me	O	O	OH
I-422	(4-CH ₂ CH≡CF)Ph	Me	O	O	OH
I-423		Me	O	O	OH
I-424		Me	O	O	OH
I-425		Me	O	O	OH
I-426		Me	O	O	OH
I-427		Me	O	O	OH
I-428		Me	O	O	OH
I-429	(4-OCHF ₂)Ph	Me	O	O	OH
I-430	(4-SMe)Ph	Me	O	O	OH
I-431	(4-SOMe)Ph	Me	O	O	OH
I-432	(4-SO ₂ Me)Ph	Me	O	O	OH
I-433	(4-SCF ₃)Ph	Me	O	O	OH
I-434	(4-SOCF ₃)Ph	Me	O	O	OH
I-435	(4-SO ₂ CF ₃)Ph	Me	O	O	OH
I-436		Me	O	O	OH
I-437		Me	O	O	OH


[Table 14]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-438		Me	O	O	OH
I-439		Me	O	O	OH
I-440		Me	O	O	OH
I-441		Me	O	O	OH
I-442		Me	O	O	OH
I-443		Me	O	O	OH
I-444		Me	O	O	OH
I-445		Me	O	O	OH
I-446		Me	O	O	OH
I-447		Me	O	O	OH
I-448		Me	O	O	OH
I-449		Me	O	O	OH
I-450		Me	O	O	OH
I-451		Me	O	O	OH
I-452		Me	O	O	OH
I-453		Me	O	O	OH
I-454		Me	O	O	OH
I-455		Me	O	O	OH

[Table 15]

Compound No.	R ¹	R ²	Y	Z	R ⁴
I-456		Me	O	O	OH
I-457		Me	O	O	OH
I-458		Me	O	O	OH
I-459		Me	O	O	OH
I-460		Me	O	O	OH
I-461		Me	O	O	OH
I-462		Me	O	O	OH
I-463		Me	O	O	OH
I-464		Me	O	S	OH
I-465	Ph(3,4,5-Cl)	Me	O	O	OH
I-466	N(Me)Ph	Me	O	O	OH
I-467		Me	O	O	OH
I-468	CH ₂ CO(Bu-t)	Me	O	O	OH
I-469	Ph(2,3,5,6-F ₄)	Me	O	O	OH
I-470	Ph[(3,5-(CF ₃) ₂)]	Me	O	O	OH
I-471	CH ₂ C(Me)=NOMe	Me	O	O	OH
I-472	Ph(2,4,6-Me ₃)	Me	O	O	OH
I-473	Ph(2,3,4,5,6-F ₅)	Me	O	O	OH
I-474	N(Et)Ph	Me	O	O	OH
I-475	N(Pr-i)Ph	Me	O	O	OH
I-476	N(Me)Ph(4-F)	Me	O	O	OH
I-477	Ph	CH ₂ CF ₃	O	O	OH
I-478	CH ₂ C(Me)=NOEt	Me	O	O	OH
I-479	CH ₂ C(Me)=NO(Pr-i)	Me	O	O	OH
I-480	Ph(4-F)	Me	O	S	OH

[Table 16]

							
Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-1	Me	Me	O	O	Me	H	OH
II-2	Et	Me	O	O	Me	Me	OH
II-3	Pr ⁿ	Me	O	O	Me	H	OH
II-4	Pr ⁱ	Me	O	O	Me	H	OH
II-5	Bu ⁿ	Me	O	O	Me	H	OH
II-6	Bu ⁱ	Me	O	O	Me	H	OH
II-7	Bu ^s	Me	O	O	Me	Me	OH
II-8	Bu ^t	Me	O	O	Me	H	OSO ₂ Pr
II-9	Hex ⁿ	Me	O	O	Me	H	OH
II-10	CH ₂ CF ₃	Me	O	O	Me	H	OH
II-11	CH ₂ CH=CH ₂	Me	O	O	Et	H	OH
II-12	CH ₂ C(Me)=CH ₂	Me	O	O	Me	H	OH
II-13	CH ₂ CH ₂ CH=CMe ₂	Me	O	O	Me	H	OH
II-14	CH ₂ C≡CH	Me	O	O	Me	Me	OH
II-15	CH ₂ C≡CCH ₃	Me	O	O	Me	H	OSO ₂ Ph
II-16	Pr ^c	Me	O	O	Me	H	OH
II-17	Bu ^c	Me	O	O	i-Pr	H	OH
II-18	Pen ^c	Me	O	O	Me	H	OH
II-19	Hex ^c	Me	O	O	Et	H	OH
II-20	CH ₂ Pr ^c	Me	O	O	Me	H	OH
II-21	CH ₂ Bu ^c	Me	O	O	Me	H	OH
II-22	CH ₂ Pen ^c	Me	O	O	Me	Me	OH
II-23	CH ₂ Hex ^c	Me	O	O	Me	H	OH
II-24	CH ₂ CH=CCl ₂	Me	O	O	Me	H	OSO ₂ Pr
II-25	CH ₂ CCl=CHCl	Me	O	O	Me	H	OH
II-26	CH ₂ CH ₂ CH=CCl ₂	Me	O	O	Et	H	OH
II-27	CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O	Me	H	OH
II-28	CH ₂ CH ₂ CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O	Me	H	OH
II-29	CH ₂ CH=CF ₂	Me	O	O	Me	Me	OH
II-30	CH ₂ CH ₂ OMe	Me	O	O	Me	H	OH
II-31	CH ₂ CH ₂ OEt	Me	O	O	Me	H	OH
II-32	CH(Me)CH ₂ OMe	Me	O	O	Pr ⁱ	H	OH
II-33	CH ₂ CH ₂ OCH ₂ CH ₂ OMe	Me	O	O	Me	H	OH
II-34	CH ₂ CH ₂ OPr ⁿ	Me	O	O	Et	H	OH
II-35	CH ₂ CH ₂ OPr ⁱ	Me	O	O	Me	H	OH
II-36	CH ₂ CH ₂ OPr ^c	Me	O	O	Me	Me	OH
II-37	CH ₂ CH ₂ OBu ^c	Me	O	O	Me	H	OH
II-38	CH ₂ CH ₂ OPen ^c	Me	O	O	Me	H	OH
II-39	CH ₂ CH ₂ OHex ^c	Me	O	O	Me	H	OH

[Table 17]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-40	CH ₂ CH ₂ OCH ₂ CF ₃	Me	O	O	Et	Me	OH
II-41	CH ₂ CH ₂ CH ₂ OMe	Me	O	O	Me	H	OH
II-42	CH=CHMe	Me	O	O	Me	H	OSO ₂ Ph
II-43	CH ₂ SMe	Me	O	O	Me	H	OH
II-44	CH ₂ SPr-n	Me	O	O	Me	H	OH
II-45	CH ₂ CH ₂ SMe	Me	O	O	Pr-i	H	OH
II-46	CH ₂ CH ₂ SOMe	Me	O	O	Me	H	OH
II-47	CH ₂ CH ₂ SO ₂ Me	Me	O	O	Me	Me	OH
II-48	CH ₂ CH ₂ CH ₂ SMe	Me	O	O	Et	H	OH
II-49	CH ₂ CH ₂ CH ₂ SO ₂ Me	Me	O	O	Me	H	OH
II-50	Ph	Me	O	O	Me	H	OH
II-51	Ph(2-Cl)	Me	O	O	Me	H	OH
II-52	Ph(3-Cl)	Me	O	O	Me	H	OH
II-53	Ph(4-Cl)	Me	O	O	Me	H	OSO ₂ Pr
II-54	Ph(2-F)	Me	O	O	Pr-i	H	OH
II-55	Ph(3-F)	Me	O	O	Me	H	OH
II-56	Ph(4-F)	Me	O	O	Me	H	OH
II-57	Ph(2-Me)	Me	O	O	Me	Me	OH
II-58	Ph(3-Me)	Me	O	O	Me	H	OH
II-59	Ph(4-Me)	Me	O	O	Et	H	OH
II-60	Ph(2-OMe)	Me	O	O	Me	H	OH
II-61	Ph(3-OMe)	Me	O	O	Me	H	OH
II-62	Ph(4-OMe)	Me	O	O	Me	H	OH
II-63	Ph(2-CF ₃)	Me	O	O	Me	H	OH
II-64	Ph(3-CF ₃)	Me	O	O	Pr-i	H	OSO ₂ Ph
II-65	Ph(4-CF ₃)	Me	O	O	Me	H	OH
II-66	Ph(2-NO ₂)	Me	O	O	Me	H	OH
II-67	Ph(3-NO ₂)	Me	O	O	Me	H	OH
II-68	Ph(4-NO ₂)	Me	O	O	Me	Me	OH
II-69	Ph(2-OCF ₃)	Me	O	O	Me	H	OH
II-70	Ph(3-OCF ₃)	Me	O	O	Et	H	OH
II-71	Ph(4-OCF ₃)	Me	O	O	Me	H	OH
II-72	Ph(2-CN)	Me	O	O	Me	H	OH
II-73	Ph(3-CN)	Me	O	O	Me	H	OH
II-74	Ph(4-CN)	Me	O	O	Me	H	OH
II-75	Ph(3,4-F ₂)	Me	O	O	Pr-i	H	OH
II-76	Ph(3,5-F ₂)	Me	O	O	Me	H	OH
II-77	Ph(2,3-F ₂)	Me	O	O	Me	Me	OSO ₂ Pr
II-78	Ph(2,4-F ₂)	Me	O	O	Me	H	OH
II-79	Ph(2,5-F ₂)	Me	O	O	Me	H	OH












[Table 18]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-80	Ph(2,6-F ₂)	Me	O	O	Et	Me	OH
II-81	Ph(3,4-Cl ₂)	Me	O	O	Me	H	OH
II-82	Ph(3,5-Cl ₂)	Me	O	O	Me	H	OH
II-83	Ph(2,3-Cl ₂)	Me	O	O	Me	H	OH
II-84	Ph(2,4-Cl ₂)	Me	O	O	Me	H	OH
II-85	Ph(2,5-Cl ₂)	Me	O	O	Pr-i	H	OH
II-86	Ph(2,6-Cl ₂)	Me	O	O	Me	H	OH
II-87	Ph(3,4-Me ₂)	Me	O	O	Me	H	OH
II-88	Ph(3,5-Me ₂)	Me	O	O	Me	H	OH
II-89	Ph(2,3-Me ₂)	Me	O	O	Me	Me	OH
II-90	Ph(2,4-Me ₂)	Me	O	O	Me	H	OH
II-91	Ph(2,5-Me ₂)	Me	O	O	Me	H	OH
II-92	Ph(2,6-Me ₂)	Me	O	O	Me	H	OH
II-93	Ph(3,4-(OMe) ₂)	Me	O	O	Me	H	OH
II-94	Ph(3,5-(OMe) ₂)	Me	O	O	Me	H	OH
II-95	Ph(2,3-(OMe) ₂)	Me	O	O	Me	H	OH
II-96	Ph(2,4-(OMe) ₂)	Me	O	O	Pr-i	H	OH
II-97	Ph(2,5-(OMe) ₂)	Me	O	O	Me	H	OSO ₂ Ph
II-98	Ph(2,6-(OMe) ₂)	Me	O	O	Me	H	OH
II-99	Ph(3-F-4-OMe)	Me	O	O	Me	Me	OH
II-100	Ph(3-F-5-OMe)	Me	O	O	Me	H	OH
II-101	Ph(2-F-3-OMe)	Me	O	O	Me	H	OH
II-102	Ph(2-F-4-OMe)	Me	O	O	Me	H	OH
II-103	Ph(2-F-5-OMe)	Me	O	O	Me	H	OH
II-104	Ph(2-F-6-OMe)	Me	O	O	Me	H	OH
II-105	Ph(3-F-4-Me)	Me	O	O	Me	H	OSO ₂ Pr
II-106	Ph(3-F-5-Me)	Me	O	O	Me	H	OH
II-107	Ph(2-F-3-Me)	Me	O	O	Me	H	OH
II-108	Ph(2-F-4-Me)	Me	O	O	Me	H	OH
II-109	Ph(2-F-5-Me)	Me	O	O	Me	Me	OH
II-110	Ph(2-F-6-Me)	Me	O	O	Me	H	OH
II-111	Ph(3-OMe-4-F)	Me	O	O	Me	H	OH
II-112	Ph(2-OMe-3-F)	Me	O	O	Pr-i	H	OSO ₂ Ph(4-Me)
II-113	Ph(2-OMe-4-F)	Me	O	O	Me	H	OH
II-114	Ph(2-OMe-5-F)	Me	O	O	Me	H	OH
II-115	Ph(3-Me-4-F)	Me	O	O	Me	H	OH
II-116	Ph(2-Me-3-F)	Me	O	O	Me	H	OH
II-117	Ph(2-Me-4-F)	Me	O	O	Me	H	OH
II-118	Ph(2-Me-5-F)	Me	O	O	Me	H	OH
II-119	Ph(3-Cl-4-OMe)	Me	O	O	Me	Me	OH
II-120	Ph(3-Cl-5-OMe)	Me	O	O	Me	H	OH















[Table 19]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-121	Ph(2-Cl-3-OMe)	Me	O	O	Me	H	OH
II-122	Ph(2-Cl-4-OMe)	Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-123	Ph(2-Cl-5-OMe)	Me	O	O	Pr-i	H	OH
II-124	Ph(2-Cl-6-OMe)	Me	O	O	Me	H	OH
II-125	Ph(3-Cl-4-Me)	Me	O	O	Me	H	OH
II-126	Ph(3-Cl-5-Me)	Me	O	O	Me	Me	OH
II-127	Ph(2-Cl-3-Me)	Me	O	O	Me	H	OH
II-128	Ph(2-Cl-4-Me)	Me	O	O	Me	H	OH
II-129	Ph(2-Cl-5-Me)	Me	O	O	Me	H	OH
II-130	Ph(2-Cl-6-Me)	Me	O	O	Me	H	OH
II-131	Ph(3-OMe-4-Cl)	Me	O	O	Me	H	OH
II-132	Ph(2-OMe-3-Cl)	Me	O	O	Me	H	OSO ₂ Ph
II-133	Ph(2-OMe-4-Cl)	Me	O	O	Pr-i	H	OH
II-134	Ph(2-OMe-5-Cl)	Me	O	O	Me	H	OH
II-135	Ph(3-Me-4-Cl)	Me	O	O	Me	Me	OH
II-136	Ph(2-Me-3-Cl)	Me	O	O	Me	H	OH
II-137	Ph(2-Me-4-Cl)	Me	O	O	Me	H	OH
II-138	Ph(2-Me-5-Cl)	Me	O	O	Me	H	OH
II-139	Ph(3-F-4-Cl)	Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-140	Ph(3-F-5-Cl)	Me	O	O	Me	H	OH
II-141	Ph(2-F-3-Cl)	Me	O	O	Me	H	OH
II-142	Ph(2-F-4-Cl)	Me	O	O	Me	H	OH
II-143	Ph(2-F-5-Cl)	Me	O	O	Me	H	OH
II-144	Ph(2-F-6-Cl)	Me	O	O	Me	H	OH
II-145	Ph(3-Cl-4-F)	Me	O	O	Me	H	OH
II-146	Ph(2-Cl-3-F)	Me	O	O	Me	Me	OH
II-147	Ph(2-Cl-4-F)	Me	O	O	Me	H	OH
II-148	Ph(2-Cl-5-F)	Me	O	O	Pr-i	H	OH
II-149	Ph(3-Me-4-OMe)	Me	O	O	Me	H	OH
II-150	Ph(3-Me-5-OMe)	Me	O	O	Me	H	OH
II-151	Ph(2-Me-3-OMe)	Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-152	Ph(2-Me-4-OMe)	Me	O	O	Me	H	OH
II-153	Ph(2-Me-5-OMe)	Me	O	O	Me	H	OH
II-154	Ph(2-Me-6-OMe)	Me	O	O	Me	H	OH
II-155	Ph(3-OMe-4-Me)	Me	O	O	Me	Me	OH
II-156	Ph(2-OMe-3-Me)	Me	O	O	Me	H	OH
II-157	Ph(2-OMe-4-Me)	Me	O	O	Me	H	OH
II-158	Ph(2-OMe-5-Me)	Me	O	O	Me	Me	OH
II-159	Ph(3-CN-4-OMe)	Me	O	O	Me	H	OH
II-160	Ph(3-OMe-4-CN)	Me	O	O	Me	H	OH
II-161	Ph(3-Me-4-CN)	Me	O	O	Pr-i	H	OSO ₂ Ph(4-Me)











[Table 20]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-162	Ph(3-CN·4-Me)	Me	O	O	Me	H	OH
II-163	Ph(3-NO ₂ ·4-OMe)	Me	O	O	Me	H	OH
II-164	Ph(3-OMe·4-NO ₂)	Me	O	O	Me	H	OH
II-165	Ph(3-Me·4-NO ₂)	Me	O	O	Me	H	OH
II-166	Ph(3-NO ₂ ·4-Me)	Me	O	O	Me	H	OH
II-167	Ph(3,5-F ₂ ·5-OMe)	Me	O	O	Me	H	OH
II-168	Ph(3,5-F ₂ ·5-Me)	Me	O	O	Me	Me	OH
II-169	Ph(3,4,5-(OMe) ₃)	Me	O	O	Me	H	OH
II-170		Me	O	O	Me	H	OH
II-171		Me	O	O	Me	H	OH
II-172		Me	O	O	Pri	H	OSO ₂ Ph(4-Me)
II-173		Me	O	O	Me	H	OH
II-174		Me	O	O	Me	H	OH
II-175		Me	O	O	Me	Me	OH
II-176		Me	O	O	Me	H	OH
II-177		Me	O	O	Me	H	OH
II-178		Me	O	O	Me	H	OH
II-179		Me	O	O	Me	H	OH
II-180		Me	O	O	Me	H	OH

[Table 21]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-181		Me	O	O	Me	H	OH
II-182		Me	O	O	Me	H	OH
II-183		Me	O	O	Me	H	OH
II-184		Me	O	O	Pri	H	OH
II-185		Me	O	O	Me	Me	OH
II-186		Me	O	O	Me	H	OH
II-187		Me	O	O	Me	H	OH
II-188		Me	O	O	Me	H	OH
II-189		Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-190		Me	O	O	Me	H	OH
II-191		Me	O	O	Me	H	OH
II-192		Me	O	O	Me	H	OH
II-193		Me	O	O	Me	H	OH
II-194		Me	O	O	Me	H	OH







[Table 22]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-195		Me	O	O	Me	H	OH
II-196		Me	O	O	Me	H	OH
II-197		Me	O	O	Me	H	OH
II-198		Me	O	O	Me	H	OH
II-199		Me	O	O	Pr-i	Me	OH
II-200		Me	O	O	Me	H	OH
II-201		Me	O	O	Me	H	OH
II-202		Me	O	O	Me	H	OH
II-203		Me	O	O	Me	H	OH
II-204		Me	O	O	Me	Me	OH
II-205	CH ₂ Ph	Me	O	O	Me	H	OH
II-206	CH ₂ CH ₂ Ph	Me	O	O	Me	H	OH
II-207	CH ₂ CH ₂ CH ₂ Ph	Me	O	O	Me	H	OH
II-208	CH ₂ CH=CHPh	Me	O	O	Me	H	OH
II-209	CH ₂ C≡CPh	Me	O	O	Me	H	OH
II-210	CH ₂ CH=NOMe	Me	O	O	Me	H	OH
II-211	CH ₂ CH=NOEt	Me	O	O	Me	H	OH
II-212	CH ₂ CH=NOPr-n	Me	O	O	Me	H	OH
II-213	CH ₂ CH=NOPh	Me	O	O	Me	Me	OH
II-214	CH ₂ CH(OMe) ₂	Me	O	O	Me	H	OH
II-215	CH ₂ CHO	Me	O	O	Et	H	OH
II-216	NH ₂	Me	O	O	Me	H	OH
II-217	NHMe	Me	O	O	Me	H	OH
II-218	NHEt	Me	O	O	Me	H	OH






[Table 23]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-219	NHPr-n	Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-220	NHPr-i	Me	O	O	Me	H	OH
II-221	NHBu-n	Me	O	O	Pr-i	H	OH
II-222	NHBu-i	Me	O	O	Me	H	OH
II-223	NHBu-s	Me	O	O	Me	Me	OH
II-224	NHCH ₂ Pr-c	Me	O	O	Me	H	OH
II-225	NHPen-n	Me	O	O	Me	H	OH
II-226	NHHex-n	Me	O	O	Me	H	OH
II-227	NHCH ₂ CH ₂ CH ₂ Cl	Me	O	O	Me	H	OH
II-228	NHCH ₂ CH ₂ CH ₂ F	Me	O	O	Me	H	OH
II-229	NHCH ₂ CH ₂ OMe	Me	O	O	Me	H	OH
II-230	NMe ₂	Me	O	O	Me	H	OH
II-231	NEt ₂	Me	O	O	Me	H	OH
II-232	N(Pr-n) ₂	Me	O	O	Me	H	OH
II-233	N(Bu-n) ₂	Me	O	O	Me	H	OH
II-234	N(Me)Et	Me	O	O	Me	Me	OH
II-235	N(Me)CH ₂ CH ₂ OMe	Me	O	O	Et	H	OH
II-236	NHPh	Me	O	O	Me	H	OH
II-237	NHCH ₂ Ph	Me	O	O	Me	H	OH
II-238	N=CMe ₂	Me	O	O	Me	H	OH
II-239	N=CEt ₂	Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-240	N=CHNMe ₂	Me	O	O	Me	H	OH
II-241	NHC(=O)Me	Me	O	O	Me	H	OH
II-242	N[C(=O)Me] ₂	Me	O	O	Me	H	OH
II-243	NHC(=O)OMe	Me	O	O	Pr-i	H	OH
II-244	N[C(=O)OMe] ₂	Me	O	O	Me	H	OH
II-245	NHSO ₂ Me	Me	O	O	Me	H	OH
II-246	NHSO ₂ Ph	Me	O	O	Me	Me	OH
II-247	NHSO ₂ CH ₂ Ph	Me	O	O	Me	H	OH
II-248	OMe	Me	O	O	Et	H	OH
II-249	OEt	Me	O	O	Me	H	OH
II-250	OPr-n	Me	O	O	Me	H	OH
II-251	OPr-i	Me	O	O	Me	H	OH
II-252	OCH ₂ Pr-c	Me	O	O	Me	H	OH
II-253	OCH ₂ Cl	Me	O	O	Me	H	OH
II-254	OCHCl ₂	Me	O	O	Me	H	OH
II-255	OCCL ₃	Me	O	O	Me	Me	OH
II-256	OCH ₂ F	Me	O	O	Me	H	OH
II-257	OCHF ₂	Me	O	O	Me	H	OH
II-258	OCF ₃	Me	O	O	Me	H	OSO ₂ Ph(4-Me)
II-259	Ph	Et	O	O	Et	H	OH
II-260	Ph	Pr-i	O	O	Me	H	OH
II-261	Ph	CHF ₂	O	O	Me	H	OH



[Table 24]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-262	Ph	Ph	O	O	Me	H	OH
II-263	Ph	Me	O	S	Me	Me	OH
II-264	Ph	Me	S	S	Me	H	OH
II-265	Me	Me	O	S	Me	H	OH
II-266	Me	Me	S	S	Me	H	OH
II-267	Ph	Me	O	O	Pr-i	H	OSO ₂ Pr
II-268	Ph(4-OEt)	Me	O	O	Me	H	OH
II-269	Ph(2-Ph)	Me	O	O	Me	H	OH
II-270	Ph(3-Ph)	Me	O	O	Me	H	OH
II-271	Ph(4-Ph)	Me	O	O	Me	H	OH
II-272		Me	O	O	Me	Me	OH
II-273		Me	O	O	Et	H	OSO ₂ Ph(4-Me)
II-274	Me		O	O	Me	H	OH
II-275	Et		O	O	Me	H	OH
II-276	H	Me	O	O	Me	H	OH
II-277	CH ₂ C≡CF	Me	O	O	Me	H	OH
II-278		Me	O	O	Me	H	OH
II-279		Me	O	O	Me	H	OH
II-280	CH ₂ NH ₂	Me	O	O	Me	H	OH
II-281	CH ₂ NO ₂	Me	O	O	Me	H	OH
II-282	CH ₂ NHCH ₃	Me	O	O	Me	H	OH
II-283	CH ₂ N(CH ₃) ₂	Me	O	O	Me	H	OH
II-284	CH ₂ SCH ₂ CF ₃	Me	O	O	Me	H	OH
II-285	CH ₂ SOCH ₂ CF ₃	Me	O	O	Me	H	OH
II-286	CH ₂ SO ₂ CH ₂ CF ₃	Me	O	O	Me	H	OH
II-287	CH ₂ OH	Me	O	O	Me	H	OH
II-288	CH ₂ OBn	Me	O	O	Me	H	OH
II-289	CH ₂ OCH ₂ Pr-c	Me	O	O	Me	H	OH







[Table 25]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-290	CH ₂ OPh	Me	O	O	Me	H	OH
II-291	CH ₂ SPh	Me	O	O	Me	H	OH
II-292	CH ₂ SOPh	Me	O	O	Me	H	OH
II-293	CH ₂ SO ₂ Ph	Me	O	O	Me	H	OH
II-294	CH ₂ CON(CH ₃) ₂	Me	O	O	Me	H	OH
II-295	CH ₂ COCH ₃	Me	O	O	Me	H	OH
II-296	CH ₂ OCOCH ₃	Me	O	O	Me	H	OH
II-297	CH ₂ ON=CHCH ₃	Me	O	O	Me	H	OH
II-298	C ₂ H ₄ OC ₂ H ₄ SCH ₃	Me	O	O	Me	H	OH
II-299	C ₂ H ₄ OC ₂ H ₄ SOCH ₃	Me	O	O	Me	H	OH
II-300	C ₂ H ₄ OC ₂ H ₄ SO ₂ CH ₃	Me	O	O	Me	H	OH
II-301	CH ₂ OCH ₂ CN	Me	O	O	Me	H	OH
II-302	CH ₂ CN	Me	O	O	Me	H	OH
II-303	OCH ₂ CH=CH ₂	Me	O	O	Me	H	OH
II-304	OCH ₂ C≡CH	Me	O	O	Me	H	OH
II-305	OPr-c	Me	O	O	Me	H	OH
II-306		Me	O	O	Me	H	OH
II-307		Me	O	O	Me	H	OH
II-308		Me	O	O	Me	H	OH
II-309		Me	O	O	Me	H	OH
II-310		Me	O	O	Me	H	OH
II-311	Ph	H	O	O	Me	H	OH
II-312	Ph	CH ₂ CH=CH ₂	O	O	Me	H	OH
II-313	Ph	CH ₂ C≡CH	O	O	Me	H	OH
II-314	Ph	Pr-c	O	O	Me	H	OH
II-315	Ph	CH ₂ CH=CF ₂	O	O	Me	H	OH
II-316	Ph	CH ₂ C≡CF	O	O	Me	H	OH
II-317	Ph	C ₂ H ₄ OCH ₃	O	O	Me	H	OH
II-318	Ph	C ₂ H ₄ OC ₂ H ₅	O	O	Me	H	OH
II-319	Ph	CH(Me)OEt	O	O	Me	H	OH
II-320	Ph	CH ₂ OPr-c	O	O	Me	H	OH
II-321	Ph	CH(OCH ₃) ₂	O	O	Me	H	OH
II-322	Ph	CH ₂ Ph	O	O	Me	H	OH


[Table 26]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-323	Ph		O	O	Me	H	OH
II-324	Ph		O	O	Me	H	OH
II-325	Ph	Me	O	O	Me	H	NH ₂
II-326	Ph	Me	O	O	Me	H	Cl
II-327	Ph	Me	O	O	Me	H	CN
II-328	Ph	Me	O	O	Me	H	NCS
II-329	Ph	Me	O	O	Me	H	NCO
II-330	Ph	Me	O	O	Me	H	OCO ₂ H
II-331	Ph	Me	O	O	Me	H	OCO ₂ CH ₃
II-332	Ph	Me	O	O	Me	H	OCO ₂ CH ₂ Ph
II-333	Ph	Me	O	O	Me	H	OMe
II-334	Ph	Me	O	O	Me	H	OE _t
II-335	Ph	Me	O	O	Me	H	OP _r
II-336	Ph	Me	O	O	Me	H	OCH ₂ CH=CH ₂
II-337	Ph	Me	O	O	Me	H	OCH ₂ C≡CH
II-338	Ph	Me	O	O	Me	H	OP _{r-c}
II-339	Ph	Me	O	O	Me	H	OBu _c
II-340	Ph	Me	O	O	Me	H	OPen _c
II-341	Ph	Me	O	O	Me	H	OHex _c
II-342	Ph	Me	O	O	Me	H	OCH ₂ CN
II-343	Ph	Me	O	O	Me	H	OCH ₂ Pr _c
II-344	Ph	Me	O	O	Me	H	OCOCH ₃
II-345	Ph	Me	O	O	Me	H	OCOCCl ₃
II-346	Ph	Me	O	O	Me	H	OCOCH=CH ₂
II-347	Ph	Me	O	O	Me	H	OCOCH=CF ₂
II-348	Ph	Me	O	O	Me	H	OCOCH ₂ C≡CH
II-349	Ph	Me	O	O	Me	H	OCOCH ₂ C≡CF
II-350	Ph	Me	O	O	Me	H	OCH ₂ CO ₂ CH ₃
II-351	Ph	Me	O	O	Me	H	OPh
II-352	Ph	Me	O	O	Me	H	OCH ₂ Ph
II-353	Ph	Me	O	O	Me	H	OCOPh
II-354	Ph	Me	O	O	Me	H	OCOCH ₂ Ph
II-355	Ph	Me	O	O	Me	H	OCH ₂ COPh
II-356	Ph	Me	O	O	Me	H	OSO ₂ CH ₂ CF ₃
II-357	Ph	Me	O	O	Me	H	OSO ₂ CH ₂ Ph
II-358	Ph	Me	O	O	Me	H	SCH ₃
II-359	Ph	Me	O	O	Me	H	SOCH ₃
II-360	Ph	Me	O	O	Me	H	SO ₂ CH ₃
II-361	Ph	Me	O	O	Me	H	SCH ₂ CF ₃
II-362	Ph	Me	O	O	Me	H	SOCH ₂ CF ₃
II-363	Ph	Me	O	O	Me	H	SO ₂ CH ₂ CF ₃

[Table 27]

Compound No.	R ¹	R ²	Y	Z	R ²⁰	R ²¹	R ⁴
II-364	Ph	Me	O	O	Me	H	SCH ₂ CH=CH ₂
II-365	Ph	Me	O	O	Me	H	SOCH ₂ CH=CH ₂
II-366	Ph	Me	O	O	Me	H	SO ₂ CH ₂ CH=CH ₂
II-367	Ph	Me	O	O	Me	H	SCH ₂ CH≡CH
II-368	Ph	Me	O	O	Me	H	SOCH ₂ CH≡CH
II-369	Ph	Me	O	O	Me	H	SO ₂ CH ₂ CH≡CH
II-370	Ph	Me	O	O	Me	H	SCH ₂ Ph
II-371	Ph	Me	O	O	Me	H	SOPh
II-372	Ph	Me	O	O	Me	H	SOCH ₂ Ph
II-373	Ph	Me	O	O	Me	H	SO ₂ Ph
II-374	Ph	Me	O	O	Me	H	SO ₂ CH ₂ Ph
II-375	Ph	Me	O	O	Me	H	NHCH ₃
II-376	Ph	Me	O	O	Me	H	N(CH ₃) ₂
II-377	Ph	Me	O	O	Me	H	NHCOCH ₃
II-378	Ph	Me	O	O	Me	H	OCH ₂ CH ₂ - 
II-379	Ph	Me	O	O	Me	H	
II-380	Ph	Me	O	O	Me	H	
II-381	Ph	Me	O	O	Me	H	
II-382	Ph	Me	O	O	Me	H	
II-383	Ph	Me	O	O	Me	H	

[Table 28]

					
Compound No.	R ¹	R ²	Y	Z	R ⁴
III-1	Me	Me	O	O	OH
III-2	Et	Me	O	O	OH
III-3	Pr-n	Me	O	O	OH
III-4	Pr-i	Me	O	O	OH
III-5	Bu-n	Me	O	O	OH
III-6	Bu-i	Me	O	O	OH
III-7	Bu-s	Me	O	O	OH
III-8	Bu-t	Me	O	O	OH
III-9	Hex-n	Me	O	O	OH
III-10	CH ₂ CF ₃	Me	O	O	OH
III-11	CH ₂ CH=CH ₂	Me	O	O	OH
III-12	CH ₂ C(Me)=CH ₂	Me	O	O	OH
III-13	CH ₂ CH ₂ CH=CMe ₂	Me	O	O	OH
III-14	CH ₂ C≡CH	Me	O	O	OH
III-15	CH ₂ C≡CCH ₃	Me	O	O	OH
III-16	Pr-c	Me	O	O	OH
III-17	Bu-c	Me	O	O	OH
III-18	Pen-c	Me	O	O	OH
III-19	Hex-c	Me	O	O	OH
III-20	CH ₂ Pr-c	Me	O	O	OH
III-21	CH ₂ Bu-c	Me	O	O	OH
III-22	CH ₂ Pen-c	Me	O	O	OH
III-23	CH ₂ Hex-c	Me	O	O	OH
III-24	CH ₂ CH=CCl ₂	Me	O	O	OH
III-25	CH ₂ CCl=CHCl	Me	O	O	OH
III-26	CH ₂ CH ₂ CH=CCl ₂	Me	O	O	OH
III-27	CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O	OH
III-28	CH ₂ CH ₂ CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O	OH
III-29	CH ₂ CH=CF ₂	Me	O	O	OH
III-30	CH ₂ CH ₂ OMe	Me	O	O	OH
III-31	CH ₂ CH ₂ OEt	Me	O	O	OH
III-32	CH(Me)CH ₂ OMe	Me	O	O	OH
III-33	CH ₂ CH ₂ OCH ₂ CH ₂ OMe	Me	O	O	OH
III-34	CH ₂ CH ₂ OPr-n	Me	O	O	OH
III-35	CH ₂ CH ₂ OPr-i	Me	O	O	OH
III-36	CH ₂ CH ₂ OPr-c	Me	O	O	OH
III-37	CH ₂ CH ₂ OBu-c	Me	O	O	OH
III-38	CH ₂ CH ₂ OPen-c	Me	O	O	OH

[Table 29]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-39	CH ₂ CH ₂ OH _{ex-c}	Me	O	O	OH
III-40	CH ₂ CH ₂ OCH ₂ CF ₃	Me	O	O	OH
III-41	CH ₂ CH ₂ CH ₂ OMe	Me	O	O	OH
III-42	CH=CHMe	Me	O	O	OH
III-43	CH ₂ SMe	Me	O	O	OH
III-44	CH ₂ SP _{r-n}	Me	O	O	OH
III-45	CH ₂ CH ₂ SMe	Me	O	O	OH
III-46	CH ₂ CH ₂ SOMe	Me	O	O	OH
III-47	CH ₂ CH ₂ SO ₂ Me	Me	O	O	OH
III-48	CH ₂ CH ₂ CH ₂ SMe	Me	O	O	OH
III-49	CH ₂ CH ₂ CH ₂ SO ₂ Me	Me	O	O	OH
III-50	Ph	Me	O	O	OH
III-51	Ph(2-Cl)	Me	O	O	OH
III-52	Ph(3-Cl)	Me	O	O	OH
III-53	Ph(4-Cl)	Me	O	O	OH
III-54	Ph(2-F)	Me	O	O	OH
III-55	Ph(3-F)	Me	O	O	OH
III-56	Ph(4-F)	Me	O	O	OH
III-57	Ph(2-Me)	Me	O	O	OH
III-58	Ph(3-Me)	Me	O	O	OH
III-59	Ph(4-Me)	Me	O	O	OH
III-60	Ph(2-OMe)	Me	O	O	OH
III-61	Ph(3-OMe)	Me	O	O	OH
III-62	Ph(4-OMe)	Me	O	O	OH
III-63	Ph(2-CF ₃)	Me	O	O	OH
III-64	Ph(3-CF ₃)	Me	O	O	OH
III-65	Ph(4-CF ₃)	Me	O	O	OH
III-66	Ph(2-NO ₂)	Me	O	O	OH
III-67	Ph(3-NO ₂)	Me	O	O	OH
III-68	Ph(4-NO ₂)	Me	O	O	OH
III-69	Ph(2-OCF ₃)	Me	O	O	OH
III-70	Ph(3-OCF ₃)	Me	O	O	OH
III-71	Ph(4-OCF ₃)	Me	O	O	OH
III-72	Ph(2-CN)	Me	O	O	OH
III-73	Ph(3-CN)	Me	O	O	OH
III-74	Ph(4-CN)	Me	O	O	OH
III-75	Ph(3,4-F ₂)	Me	O	O	OH
III-76	Ph(3,5-F ₂)	Me	O	O	OH
III-77	Ph(2,3-F ₂)	Me	O	O	OH
III-78	Ph(2,4-F ₂)	Me	O	O	OH










[Table 30]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-79	Ph(2,5-F ₂)	Me	O	O	OH
III-80	Ph(2,6-F ₂)	Me	O	O	OH
III-81	Ph(3,4-Cl ₂)	Me	O	O	OH
III-82	Ph(3,5-Cl ₂)	Me	O	O	OH
III-83	Ph(2,3-Cl ₂)	Me	O	O	OH
III-84	Ph(2,4-Cl ₂)	Me	O	O	OH
III-85	Ph(2,5-Cl ₂)	Me	O	O	OH
III-86	Ph(2,6-Cl ₂)	Me	O	O	OH
III-87	Ph(3,4-Me ₂)	Me	O	O	OH
III-88	Ph(3,5-Me ₂)	Me	O	O	OH
III-89	Ph(2,3-Me ₂)	Me	O	O	OH
III-90	Ph(2,4-Me ₂)	Me	O	O	OH
III-91	Ph(2,5-Me ₂)	Me	O	O	OH
III-92	Ph(2,6-Me ₂)	Me	O	O	OH
III-93	Ph(3,4-(OMe) ₂)	Me	O	O	OH
III-94	Ph(3,5-(OMe) ₂)	Me	O	O	OH
III-95	Ph(2,3-(OMe) ₂)	Me	O	O	OH
III-96	Ph(2,4-(OMe) ₂)	Me	O	O	OH
III-97	Ph(2,5-(OMe) ₂)	Me	O	O	OH
III-98	Ph(2,6-(OMe) ₂)	Me	O	O	OH
III-99	Ph(3-F-4-OMe)	Me	O	O	OH
III-100	Ph(3-F-5-OMe)	Me	O	O	OH
III-101	Ph(2-F-3-OMe)	Me	O	O	OH
III-102	Ph(2-F-4-OMe)	Me	O	O	OH
III-103	Ph(2-F-5-OMe)	Me	O	O	OH
III-104	Ph(2-F-6-OMe)	Me	O	O	OH
III-105	Ph(3-F-4-Me)	Me	O	O	OH
III-106	Ph(3-F-5-Me)	Me	O	O	OH
III-107	Ph(2-F-3-Me)	Me	O	O	OH
III-108	Ph(2-F-4-Me)	Me	O	O	OH
III-109	Ph(2-F-5-Me)	Me	O	O	OH
III-110	Ph(2-F-6-Me)	Me	O	O	OH
III-111	Ph(3-OMe-4-F)	Me	O	O	OH
III-112	Ph(2-OMe-3-F)	Me	O	O	OH
III-113	Ph(2-OMe-4-F)	Me	O	O	OH
III-114	Ph(2-OMe-5-F)	Me	O	O	OH
III-115	Ph(3-Me-4-F)	Me	O	O	OH
III-116	Ph(2-Me-3-F)	Me	O	O	OH
III-117	Ph(2-Me-4-F)	Me	O	O	OH














[Table 31]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-118	Ph(2-Me-5-F)	Me	O	O	OH
III-119	Ph(3-Cl-4-OMe)	Me	O	O	OH
III-120	Ph(3-Cl-5-OMe)	Me	O	O	OH
III-121	Ph(2-Cl-3-OMe)	Me	O	O	OH
III-122	Ph(2-Cl-4-OMe)	Me	O	O	OH
III-123	Ph(2-Cl-5-OMe)	Me	O	O	OH
III-124	Ph(2-Cl-6-OMe)	Me	O	O	OH
III-125	Ph(3-Cl-4-Me)	Me	O	O	OH
III-126	Ph(3-Cl-5-Me)	Me	O	O	OH
III-127	Ph(2-Cl-3-Me)	Me	O	O	OH
III-128	Ph(2-Cl-4-Me)	Me	O	O	OH
III-129	Ph(2-Cl-5-Me)	Me	O	O	OH
III-130	Ph(2-Cl-6-Me)	Me	O	O	OH
III-131	Ph(3-OMe-4-Cl)	Me	O	O	OH
III-132	Ph(2-OMe-3-Cl)	Me	O	O	OH
III-133	Ph(2-OMe-4-Cl)	Me	O	O	OH
III-134	Ph(2-OMe-5-Cl)	Me	O	O	OH
III-135	Ph(3-Me-4-Cl)	Me	O	O	OH
III-136	Ph(2-Me-3-Cl)	Me	O	O	OH
III-137	Ph(2-Me-4-Cl)	Me	O	O	OH
III-138	Ph(2-Me-5-Cl)	Me	O	O	OH
III-139	Ph(3-F-4-Cl)	Me	O	O	OH
III-140	Ph(3-F-5-Cl)	Me	O	O	OH
III-141	Ph(2-F-3-Cl)	Me	O	O	OH
III-142	Ph(2-F-4-Cl)	Me	O	O	OH
III-143	Ph(2-F-5-Cl)	Me	O	O	OH
III-144	Ph(2-F-6-Cl)	Me	O	O	OH
III-145	Ph(3-Cl-4-F)	Me	O	O	OH
III-146	Ph(2-Cl-3-F)	Me	O	O	OH
III-147	Ph(2-Cl-4-F)	Me	O	O	OH
III-148	Ph(2-Cl-5-F)	Me	O	O	OH
III-149	Ph(3-Me-4-OMe)	Me	O	O	OH
III-150	Ph(3-Me-5-OMe)	Me	O	O	OH
III-151	Ph(2-Me-3-OMe)	Me	O	O	OH
III-152	Ph(2-Me-4-OMe)	Me	O	O	OH
III-153	Ph(2-Me-5-OMe)	Me	O	O	OH
III-154	Ph(2-Me-6-OMe)	Me	O	O	OH
III-155	Ph(3-OMe-4-Me)	Me	O	O	OH
III-156	Ph(2-OMe-3-Me)	Me	O	O	OH













[Table 32]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-157	Ph(2-OMe-4-Me)	Me	O	O	OH
III-158	Ph(2-OMe-5-Me)	Me	O	O	OH
III-159	Ph(3-CN-4-OMe)	Me	O	O	OH
III-160	Ph(3-OMe-4-CN)	Me	O	O	OH
III-161	Ph(3-Me-4-CN)	Me	O	O	OH
III-162	Ph(3-CN-4-Me)	Me	O	O	OH
III-163	Ph(3-NO ₂ -4-OMe)	Me	O	O	OH
III-164	Ph(3-OMe-4-NO ₂)	Me	O	O	OH
III-165	Ph(3-Me-4-NO ₂)	Me	O	O	OH
III-166	Ph(3-NO ₂ -4-Me)	Me	O	O	OH
III-167	Ph(3,5-F ₂ -5-OMe)	Me	O	O	OH
III-168	Ph(3,5-F ₂ -5-Me)	Me	O	O	OH
III-169	Ph(3,4,5-(OMe) ₃)	Me	O	O	OH
III-170		Me	O	O	OH
III-171		Me	O	O	OH
III-172		Me	O	O	OH
III-173		Me	O	O	OH
III-174		Me	O	O	OH
III-175		Me	O	O	OH
III-176		Me	O	O	OH
III-177		Me	O	O	OH
III-178		Me	O	O	OH

[Table 33]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-179		Me	O	O	OH
III-180		Me	O	O	OH
III-181		Me	O	O	OH
III-182		Me	O	O	OH
III-183		Me	O	O	OH
III-184		Me	O	O	OH
III-185		Me	O	O	OH
III-186		Me	O	O	OH
III-187		Me	O	O	OH
III-188		Me	O	O	OH
III-189		Me	O	O	OH
III-190		Me	O	O	OH
III-191		Me	O	O	OH
III-192		Me	O	O	OH







[Table 34]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-193		Me	O	O	OH
III-194		Me	O	O	OH
III-195		Me	O	O	OH
III-196		Me	O	O	OH
III-197		Me	O	O	OH
III-198		Me	O	O	OH
III-199		Me	O	O	OH
III-200		Me	O	O	OH
III-201		Me	O	O	OH
III-202		Me	O	O	OH
III-203		Me	O	O	OH
III-204		Me	O	O	OH
III-205	CH ₂ Ph	Me	O	O	OH
III-206	CH ₂ CH ₂ Ph	Me	O	O	OH
III-207	CH ₂ CH ₂ CH ₂ Ph	Me	O	O	OH
III-208	CH ₂ CH=CHPh	Me	O	O	OH
III-209	CH ₂ C≡CPh	Me	O	O	OH
III-210	CH ₂ CH=NOMe	Me	O	O	OH
III-211	CH ₂ CH=NOEt	Me	O	O	OH







[Table 35]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-212	CH ₂ CH=NOPr-n	Me	O	O	OH
III-213	CH ₂ CH=NOPh	Me	O	O	OH
III-214	CH ₂ CH(OMe) ₂	Me	O	O	OH
III-215	CH ₂ CHO	Me	O	O	OH
III-216	NH ₂	Me	O	O	OH
III-217	NHMe	Me	O	O	OH
III-218	NHEt	Me	O	O	OH
III-219	NHPr-n	Me	O	O	OH
III-220	NHPr-i	Me	O	O	OH
III-221	NHBu-n	Me	O	O	OH
III-222	NHBu-i	Me	O	O	OH
III-223	NHBu-s	Me	O	O	OH
III-224	NHCH ₂ Pr-c	Me	O	O	OH
III-225	NHPen-n	Me	O	O	OH
III-226	NHHex-n	Me	O	O	OH
III-227	NHCH ₂ CH ₂ CH ₂ Cl	Me	O	O	OH
III-228	NHCH ₂ CH ₂ CH ₂ F	Me	O	O	OH
III-229	NHCH ₂ CH ₂ OMe	Me	O	O	OH
III-230	NMe ₂	Me	O	O	OH
III-231	NEt ₂	Me	O	O	OH
III-232	N(Pr-n) ₂	Me	O	O	OH
III-233	N(Bu-n) ₂	Me	O	O	OH
III-234	N(Me)Et	Me	O	O	OH
III-235	N(Me)CH ₂ CH ₂ OMe	Me	O	O	OH
III-236	NHPh	Me	O	O	OH
III-237	NHCH ₂ Ph	Me	O	O	OH
III-238	N=CMe ₂	Me	O	O	OH
III-239	N=CEt ₂	Me	O	O	OH
III-240	N=CHNMe ₂	Me	O	O	OH
III-241	NHC(=O)Me	Me	O	O	OH
III-242	N[C(=O)Me] ₂	Me	O	O	OH
III-243	NHC(=O)OMe	Me	O	O	OH
III-244	N[C(=O)OMe] ₂	Me	O	O	OH
III-245	NHSO ₂ Me	Me	O	O	OH
III-246	NHSO ₂ Ph	Me	O	O	OH
III-247	NHSO ₂ CH ₂ Ph	Me	O	O	OH
III-248	OMe	Me	O	O	OH
III-249	OEt	Me	O	O	OH
III-250	OPr-n	Me	O	O	OH
III-251	OPr-i	Me	O	O	OH
III-252	OCH ₂ Pr-c	Me	O	O	OH
III-253	OCH ₂ Cl	Me	O	O	OH
III-254	OCHCl ₂	Me	O	O	OH


[Table 36]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-255	OCCL ₃	Me	O	O	OH
III-256	OCH ₂ F	Me	O	O	OH
III-257	OCHF ₂	Me	O	O	OH
III-258	OCF ₃	Me	O	O	OH
III-259	Ph	Et	O	O	OH
III-260	Ph	Pr-i	O	O	OH
III-261	Ph	CHF ₂	O	O	OH
III-262	Ph	Ph	O	O	OH
III-263	Ph	Me	O	S	OH
III-264	Ph	Me	S	S	OH
III-265	Me	Me	O	S	OH
III-266	Me	Me	S	S	OH
III-267	Ph	Me	O	O	SPh
III-268	Ph(4-OEt)	Me	O	O	OH
III-269	Ph(2-Ph)	Me	O	O	OH
III-270	Ph(3-Ph)	Me	O	O	OH
III-271	Ph(4-Ph)	Me	O	O	OH
III-272		Me	O	O	OH
III-273		Me	O	O	OH
III-274	Me		O	O	OH
III-275	Et		O	O	OH
III-276	H	Me	O	O	OH
III-277	CH ₂ C≡CF	Me	O	O	OH
III-278		Me	O	O	OH
III-279		Me	O	O	OH
III-280	CH ₂ NH ₂	Me	O	O	OH
III-281	CH ₂ NO ₂	Me	O	O	OH
III-282	CH ₂ NHCH ₃	Me	O	O	OH
III-283	CH ₂ N(CH ₃) ₂	Me	O	O	OH
III-284	CH ₂ SCH ₂ CF ₃	Me	O	O	OH
III-285	CH ₂ SOCH ₂ CF ₃	Me	O	O	OH
III-286	CH ₂ SO ₂ CH ₂ CF ₃	Me	O	O	OH
III-287	CH ₂ OH	Me	O	O	OH
III-288	CH ₂ OBn	Me	O	O	OH
III-289	CH ₂ OCH ₂ Pr-c	Me	O	O	OH







[Table 37]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-290	CH ₂ OPh	Me	O	O	OH
III-291	CH ₂ SPh	Me	O	O	OH
III-292	CH ₂ SOPh	Me	O	O	OH
III-293	CH ₂ SO ₂ Ph	Me	O	O	OH
III-294	CH ₂ CON(CH ₃) ₂	Me	O	O	OH
III-295	CH ₂ COCH ₃	Me	O	O	OH
III-296	CH ₂ OCOCH ₃	Me	O	O	OH
III-297	CH ₂ ON=CHCH ₃	Me	O	O	OH
III-298	C ₂ H ₄ OC ₂ H ₄ SCH ₃	Me	O	O	OH
III-299	C ₂ H ₄ OC ₂ H ₄ SOCH ₃	Me	O	O	OH
III-300	C ₂ H ₄ OC ₂ H ₄ SO ₂ CH ₃	Me	O	O	OH
III-301	CH ₂ OCH ₂ CN	Me	O	O	OH
III-302	CH ₂ CN	Me	O	O	OH
III-303	OCH ₂ CH=CH ₂	Me	O	O	OH
III-304	OCH ₂ C≡CH	Me	O	O	OH
III-305	OPr-c	Me	O	O	OH
III-306		Me	O	O	OH
III-307		Me	O	O	OH
III-308		Me	O	O	OH
III-309		Me	O	O	OH
III-310	CH ₂ CH ₂ OCH ₂ CH ₂ O- 	Me	O	O	OH
III-311	Ph	H	O	O	OH
III-312	Ph	CH ₂ CH=CH ₂	O	O	OH
III-313	Ph	CH ₂ C≡CH	O	O	OH
III-314	Ph	Pr-c	O	O	OH
III-315	Ph	CH ₂ CH=CF ₂	O	O	OH
III-316	Ph	CH ₂ C≡CF	O	O	OH
III-317	Ph	C ₂ H ₄ OCH ₃	O	O	OH
III-318	Ph	C ₂ H ₄ OC ₂ H ₅	O	O	OH
III-319	Ph	CH(Me)OEt	O	O	OH
III-320	Ph	CH ₂ OPr-c	O	O	OH
III-321	Ph	CH(OCH ₃) ₂	O	O	OH
III-322	Ph	CH ₂ Ph	O	O	OH
III-323	Ph		O	O	OH




[Table 38]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-324	Ph		O	O	OH
III-325	Ph	Me	O	O	NH ₂
III-326	Ph	Me	O	O	Cl
III-327	Ph	Me	O	O	CN
III-328	Ph	Me	O	O	NCS
III-329	Ph	Me	O	O	NCO
III-330	Ph	Me	O	O	OCO ₂ H
III-331	Ph	Me	O	O	OCO ₂ CH ₃
III-332	Ph	Me	O	O	OCO ₂ CH ₂ Ph
III-333	Ph	Me	O	O	OMe
III-334	Ph	Me	O	O	OE _t
III-335	Ph	Me	O	O	OPr
III-336	Ph	Me	O	O	OCH ₂ CH=CH ₂
III-337	Ph	Me	O	O	OCH ₂ C≡CH
III-338	Ph	Me	O	O	OPr ^c
III-339	Ph	Me	O	O	OBu ^c
III-340	Ph	Me	O	O	OPen ^c
III-341	Ph	Me	O	O	OHex ^c
III-342	Ph	Me	O	O	OCH ₂ CN
III-343	Ph	Me	O	O	OCH ₂ Pr ^c
III-344	Ph	Me	O	O	OCOCH ₃
III-345	Ph	Me	O	O	OCOCCl ₃
III-346	Ph	Me	O	O	OCOCH=CH ₂
III-347	Ph	Me	O	O	OCOCH=CF ₂
III-348	Ph	Me	O	O	OCOCH ₂ C≡CH
III-349	Ph	Me	O	O	OCOCH ₂ C≡CF
III-350	Ph	Me	O	O	OCH ₂ CO ₂ CH ₃
III-351	Ph	Me	O	O	OPh
III-352	Ph	Me	O	O	OCH ₂ Ph
III-353	Ph	Me	O	O	OCOPh
III-354	Ph	Me	O	O	OCOCH ₂ Ph
III-355	Ph	Me	O	O	OCH ₂ COPh
III-356	Ph	Me	O	O	OSO ₂ CH ₂ CF ₃
III-357	Ph	Me	O	O	OSO ₂ CH ₂ Ph
III-358	Ph	Me	O	O	SCH ₃
III-359	Ph	Me	O	O	SOCH ₃
III-360	Ph	Me	O	O	SO ₂ CH ₃
III-361	Ph	Me	O	O	SCH ₂ CF ₃
III-362	Ph	Me	O	O	SOCH ₂ CF ₃
III-363	Ph	Me	O	O	SO ₂ CH ₂ CF ₃
III-364	Ph	Me	O	O	SCH ₂ CH=CH ₂
III-365	Ph	Me	O	O	SOCH ₂ CH=CH ₂
III-366	Ph	Me	O	O	SO ₂ CH ₂ CH=CH ₂















[Table 39]

Compound No.	R ¹	R ²	Y	Z	R ⁴
III-367	Ph	Me	O	O	SCH ₂ CH≡CH
III-368	Ph	Me	O	O	SOCH ₂ CH≡CH
III-369	Ph	Me	O	O	SO ₂ CH ₂ CH≡CH
III-370	Ph	Me	O	O	SCH ₂ Ph
III-371	Ph	Me	O	O	SOPh
III-372	Ph	Me	O	O	SOCH ₂ Ph
III-373	Ph	Me	O	O	SO ₂ Ph
III-374	Ph	Me	O	O	SO ₂ CH ₂ Ph
III-375	Ph	Me	O	O	NHCH ₃
III-376	Ph	Me	O	O	N(CH ₃) ₂
III-377	Ph	Me	O	O	NHCOCH ₃
III-378	Ph	Me	O	O	OCH ₂ CH ₂ - 
III-379	Ph	Me	O	O	- 
III-380	Ph	Me	O	O	- 
III-381	Ph	Me	O	O	- 
III-382	Ph	Me	O	O	- 
III-383	Ph	Me	O	O	O- 









[Table 40]

				
Compound No.	R ¹	A ₁	A ₂	A ₃
VI-1	Ph	C(CH ₃) ₂	CO	C(CH ₃) ₂
VI-2	Ph	CHCH ₃	CH ₂	CH ₂
VI-3	Ph	CH ₂	CHCH ₃	CH ₂
VI-4	Ph	CHCH ₃	CHCH ₃	CHCH ₃
VI-5	Ph	C(CH ₃) ₂	CH ₂	CH ₂
VI-6	Ph	CH ₂	C(CH ₃) ₂	CH ₂
VI-7	Ph	CHCH ₃	CH ₂	C(CH ₃) ₂
VI-8	Ph	CHCH ₃	CH ₂	CHCH ₃
VI-9	Ph	CHCH ₃	CHCH ₃	CH ₂
VI-10	Ph	NCH ₃	CO	CH ₂
VI-11	Ph	C(CH ₃) ₂	C(CH ₃) ₂	C(CH ₃) ₂
VI-12	Ph	C(CH ₃) ₂	S	C(CH ₃) ₂
VI-13	Ph	C(CH ₃) ₂	SO	C(CH ₃) ₂
VI-14	Ph	C(CH ₃) ₂	SO ₂	C(CH ₃) ₂
VI-15	Ph	C(CH ₃) ₂	O	C(CH ₃) ₂
VI-16	Ph	C(CH ₃) ₂	NCH ₃	C(CH ₃) ₂
VI-17	Me	C(CH ₃) ₂	CO	C(CH ₃) ₂
VI-18	Me	CHCH ₃	CH ₂	CH ₂
VI-19	Me	CH ₂	CHCH ₃	CH ₂
VI-20	Me	CHCH ₃	CHCH ₃	CHCH ₃
VI-21	Me	C(CH ₃) ₂	CH ₂	CH ₂
VI-22	Me	CH ₂	C(CH ₃) ₂	CH ₂
VI-23	Me	CHCH ₃	CH ₂	C(CH ₃) ₂
VI-24	Me	CHCH ₃	CH ₂	CHCH ₃
VI-25	Me	CHCH ₃	CHCH ₃	CH ₂
VI-26	Me	NCH ₃	CO	CH ₂
VI-27	Me	C(CH ₃) ₂	C(CH ₃) ₂	C(CH ₃) ₂
VI-28	Me	C(CH ₃) ₂	S	C(CH ₃) ₂
VI-29	Me	C(CH ₃) ₂	SO	C(CH ₃) ₂
VI-30	Me	C(CH ₃) ₂	SO ₂	C(CH ₃) ₂
VI-31	Me	C(CH ₃) ₂	O	C(CH ₃) ₂
VI-32	Me	C(CH ₃) ₂	NCH ₃	C(CH ₃) ₂
VI-33		C(CH ₃) ₂	CO	C(CH ₃) ₂
VI-34		CHCH ₃	CH ₂	CH ₂









[Table 41]

Compound No.	R ¹	A ₁	A ₂	A ₃
VI-35		CH ₂	CHCH ₃	CH ₂
VI-36		CHCH ₃	CHCH ₃	CHCH ₃
VI-37		C(CH ₃) ₂	CH ₂	CH ₂
VI-38		CH ₂	C(CH ₃) ₂	CH ₂
VI-39		CHCH ₃	CH ₂	C(CH ₃) ₂
VI-40		CHCH ₃	CH ₂	CHCH ₃
VI-41		CHCH ₃	CHCH ₃	CH ₂
VI-42		NCH ₃	CO	CH ₂
VI-43		C(CH ₃) ₂	C(CH ₃) ₂	C(CH ₃) ₂
VI-44		C(CH ₃) ₂	S	C(CH ₃) ₂
VI-45		C(CH ₃) ₂	SO	C(CH ₃) ₂
VI-46		C(CH ₃) ₂	SO ₂	C(CH ₃) ₂
VI-47		C(CH ₃) ₂	O	C(CH ₃) ₂
VI-48		C(CH ₃) ₂	NCH ₃	C(CH ₃) ₂
VI-49	Ph(4-OMe)	C(CH ₃) ₂	CO	C(CH ₃) ₂
VI-50	Ph(4-OMe)	CHCH ₃	CH ₂	CH ₂
VI-51	Ph(4-OMe)	CH ₂	CHCH ₃	CH ₂
VI-52	Ph(4-OMe)	CHCH ₃	CHCH ₃	CHCH ₃
VI-53	Ph(4-OMe)	C(CH ₃) ₂	CH ₂	CH ₂
VI-54	Ph(4-OMe)	CH ₂	C(CH ₃) ₂	CH ₂
VI-55	Ph(4-OMe)	CHCH ₃	CH ₂	C(CH ₃) ₂
VI-56	Ph(4-OMe)	CHCH ₃	CH ₂	CHCH ₃
VI-57	Ph(4-OMe)	CHCH ₃	CHCH ₃	CH ₂

[Table 42]

Compound No.	R ¹	A ₁	A ₂	A ₃
VI-58	Ph(4-OMe)	NCH ₃	CO	CH ₂
VI-59	Ph(4-OMe)	C(CH ₃) ₂	C(CH ₃) ₂	C(CH ₃) ₂
VI-60	Ph(4-OMe)	C(CH ₃) ₂	S	C(CH ₃) ₂
VI-61	Ph(4-OMe)	C(CH ₃) ₂	SO	C(CH ₃) ₂
VI-62	Ph(4-OMe)	C(CH ₃) ₂	SO ₂	C(CH ₃) ₂
VI-63	Ph(4-OMe)	C(CH ₃) ₂	O	C(CH ₃) ₂
VI-64	Ph(4-OMe)	C(CH ₃) ₂	NCH ₃	C(CH ₃) ₂
VI-65	Ph(2,4-Me ₂)	C(CH ₃) ₂	CO	C(CH ₃) ₂
VI-66	Ph(2,4-Me ₂)	CHCH ₃	CH ₂	CH ₂
VI-67	Ph(2,4-Me ₂)	CH ₂	CHCH ₃	CH ₂
VI-68	Ph(2,4-Me ₂)	CHCH ₃	CHCH ₃	CHCH ₃
VI-69	Ph(2,4-Me ₂)	C(CH ₃) ₂	CH ₂	CH ₂
VI-70	Ph(2,4-Me ₂)	CH ₂	C(CH ₃) ₂	CH ₂
VI-71	Ph(2,4-Me ₂)	CHCH ₃	CH ₂	C(CH ₃) ₂
VI-72	Ph(2,4-Me ₂)	CHCH ₃	CH ₂	CHCH ₃
VI-73	Ph(2,4-Me ₂)	CHCH ₃	CHCH ₃	CH ₂
VI-74	Ph(2,4-Me ₂)	NCH ₃	CO	CH ₂
VI-75	Ph(2,4-Me ₂)	C(CH ₃) ₂	C(CH ₃) ₂	C(CH ₃) ₂
VI-76	Ph(2,4-Me ₂)	C(CH ₃) ₂	S	C(CH ₃) ₂
VI-77	Ph(2,4-Me ₂)	C(CH ₃) ₂	SO	C(CH ₃) ₂
VI-78	Ph(2,4-Me ₂)	C(CH ₃) ₂	SO ₂	C(CH ₃) ₂
VI-79	Ph(2,4-Me ₂)	C(CH ₃) ₂	O	C(CH ₃) ₂
VI-80	Ph(2,4-Me ₂)	C(CH ₃) ₂	NCH ₃	C(CH ₃) ₂
VI-81		C(CH ₃) ₂	CO	C(CH ₃) ₂
VI-82		CHCH ₃	CH ₂	CH ₂
VI-83		CH ₂	CHCH ₃	CH ₂
VI-84		CHCH ₃	CHCH ₃	CHCH ₃
VI-85		C(CH ₃) ₂	CH ₂	CH ₂
VI-86		CH ₂	C(CH ₃) ₂	CH ₂
VI-87		CHCH ₃	CH ₂	C(CH ₃) ₂
VI-88		CHCH ₃	CH ₂	CHCH ₃

[Table 43]

Compound No.	R ¹	A ₁	A ₂	A ₃
VI-89		CHCH ₃	CHCH ₃	CH ₂
VI-90		NCH ₃	CO	CH ₂
VI-91		C(CH ₃) ₂	C(CH ₃) ₂	C(CH ₃) ₂
VI-92		C(CH ₃) ₂	S	C(CH ₃) ₂
VI-93		C(CH ₃) ₂	SO	C(CH ₃) ₂
VI-94		C(CH ₃) ₂	SO ₂	C(CH ₃) ₂
VI-95		C(CH ₃) ₂	O	C(CH ₃) ₂
VI-96		C(CH ₃) ₂	NCH ₃	C(CH ₃) ₂
VI-97	Ph(3,4,5-(OMe) ₃)	C(CH ₃) ₂	CO	C(CH ₃) ₂

Preferred examples of the triazine derivative represented by Formula 1 of the invention or salt thereof include the followings.

5 A in Formula 1 is preferably A-1, A-3, or A-5, and more preferably A-1 or A-3.

In A-1, A₁ is preferably [X₁], A₂ is preferably [X₃] or [X₄], and A₃ is preferably [X₉].

10 In [X₁], R⁵ and R⁶ are preferably a hydrogen atom or a C₁-C₆ alkyl group. In [X₃], R⁸ and R⁹ are preferably a hydrogen atom or a C₁-C₆ alkyl group. In [X₉], R³⁵ and R³⁶ are preferably a hydrogen atom or a C₁-C₆ alkyl group. Further, according to the preferred example of the invention, R⁵ in [X₁] and R³⁵ in [X₉] bind to each other via a C₁-C₅ alkylene chain, preferably an ethylene chain, to form a ring.

In A-3, R²⁰ is preferably a C₁-C₆ alkyl group and R²¹ is preferably a hydrogen atom or a C₁-C₆ alkyl group.

15 In A-1 and A-3, R⁴ is preferably a hydroxyl group, an O⁺M⁺ (M⁺ represents an alkali metal cation or an ammonium cation), or a C₁-C₁₀ alkylsulfonyloxy group.

In Formula 1, Y is preferably an oxygen atom.

In Formula 1, R¹ is preferably a group selected from a group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ halolalkenyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a phenyl group which may be substituted with a substituent group selected from Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with a substituent group selected from Substituent group α ; and a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α , and when the heterocyclic group contains a sulfur atom, it may be oxidized to be present as sulfoxide or sulfone).


In Formula 1, R² is preferably a group selected from a group consisting of a C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a phenyl group which may be substituted with a substituent group selected from Substituent group α ; and a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α).

The triazine derivative compounds represented by Formula 1, i.e., the compounds of the invention, and their salts can be produced according to various methods. Representative examples of the production method are given below, but the invention is not limited to them.

<Production method 1>

The compound represented by following Formula 1a, which is one of the compounds of the invention, can be produced according to the method with the reaction scheme shown below.

[Chem. 7]



(in the formula, R¹, R², A₁, A₂, A₃, Y and Z have the same definitions as above and Q represents a leaving group like a halogen atom, an alkylcarbonyloxy group, an alkoxy carbonyloxy group, a haloalkylcarbonyloxy group, a haloalkoxy carbonyloxy group, a benzoyloxy group, a pyridyl group, and an imidazolyl group).

(Process 1)

By reacting the compound of Formula 3 and the compound of Formula 4a in a solvent in the presence of a base, the enolester compound of Formula 5a and/or Formula 5b can be produced.

Herein, the use amount of Formula 4a can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 3. Preferably, it is from 1.0 to 1.2 mol.

Examples of the base which can be used for the present process include organic amines like triethylamine, pyridine, 4-dimethylaminopyridine, N,N-dimethylaniline, and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU); metal carbonates like sodium carbonate, potassium carbonate, magnesium carbonate, and calcium carbonate; metal hydrogen carbonates like sodium hydrogen carbonate and potassium hydrogen carbonate; metal carboxylate salts represented by metal acetate salts like sodium acetate, potassium acetate, calcium acetate, and magnesium

acetate; metal alkoxides like sodium methoxide, sodium ethoxide, sodium tertiary butoxide, potassium methoxide, and potassium tertiary butoxide; metal hydroxides like sodium hydroxide, potassium hydroxide, calcium hydroxide, and magnesium hydroxide, and; metal hydrides like lithium hydride, sodium hydride, potassium hydride, and calcium hydride. The use amount of the base is appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 3.

Preferably, it is from 1.0 to 1.2 mol.

The solvent that can be used for the present process can be any solvent if it does not inhibit the progress of the reaction. Solvents including nitriles like acetonitrile; ethers like diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, monoglyme, and diglyme; halogenated hydrocarbons like dichloroethane, chloroform, carbon tetrachloride, and tetrachloroethane; aromatic hydrocarbons like benzene, chlorobenzene, nitrobenzene, and toluene; amides like N,N-dimethylformamide and N,N-dimethylacetamide; imidazolinones like 1,3-dimethyl-2-imidazolinone, and; sulfur compounds like dimethyl sulfoxide can be used. Further, their mixture solvent can be also used.

The reaction temperature may be selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. By using a phase transfer catalyst like quaternary ammonium salt, the reaction can be carried out in a two-phase system.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 5a and/or Formula 5b, which is the target compound of the reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

(Process 2)

Compound of Formula 5a and/or Formula 5b can be also produced by reacting the compound of Formula 3 and the compound of Formula 4b with a dehydrating condensing agent in a solvent, in the presence or absence of a base.

The use amount of Formula 4b that is used for the present process can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 3. Preferably, it is from 1.0 to 1.2

mol.

Examples of the dehydrating condensing agent include dicyclohexyl carbodiimide (DCC), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide (EDC or WSC), N,N-carbonyldiimidazole, 2-chloro-1,3-dimethylimidazolium chloride, and 2-chloro-1-pyridinium iodide.

5 Examples of the base and the solvent which can be used for the present process include those described above for Process 1.

The reaction temperature may be selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

10 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

The compound of Formula 5a and/or Formula 5b, which is the target compound of the reaction, can be separated and purified in the same manner as Process 1.

(Process 3)

15 Compound of Formula 1a can be produced by reacting the compound of Formula 5a and/or Formula 5b produced by Process 1 or Process 2 with a cyano compound in the presence of a base.

Examples of the base which can be used for the present process include those described above for Process 1. The use amount of the base can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 5a and Formula 5b. Preferably, it is from 1.0 to 1.2 mol.

20 Examples of the cyano compound which can be used for the present process include potassium cyanide, sodium cyanide, acetone cyanohydrin, hydrogen cyanide, and a polymer supported with hydrogen cyanide. The use amount of the cyano compound can be appropriately selected from the range of 0.01 to 1.0 mol per 1 mol of Formula 5a and Formula 5b. Preferably, it is from 0.05 to 0.2 mol.

25 For the present process, a small amount of a phase transfer catalyst like crown ether can be also used.

Examples of the solvent which can be used for the present process include those described above for Process 1. The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from

0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Further, according to the present process, compound of Formula 1a can be produced while using Formula 5a and/or Formula 5b produced by Process 1 or Process 2 without any separation.

5 (Process 4)

Compound of Formula 1a can be also produced by reacting the compound of Formula 3 and the compound of Formula 4c in the presence of a base or a Lewis acid.

The use amount of Formula 4c that is used for the present process can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 3. Preferably, it is from 1.0 to 1.2
10 mol.

Examples of the Lewis acid include zinc chloride and aluminum chloride.

Examples of the base which can be used for the present process include those described above for Process 1. The use amount of the base that can be used for the present process can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 3. Preferably, it is
15 from 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described above for Process 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

20 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.


After the completion of the reaction, compound of Formula 1a, which is produced according to Process 3 or Process 4, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

25 <Production method 2>

With regard to Formula 1a produced by Production method 1, the hydroxyl group in the cyclohexane ring can be converted to other substituent group according to the method with the following reaction scheme.

[Chem. 8]

30



(in the formula, R^1 , R^2 , A_1 , A_2 , A_3 , Y and Z each have the same definitions as above, R^{4a}

represents an amino group, a cyano group, an isothiocyanate group, an isocyanate group, a

hydroxycarbonyloxy group, a C_1 - C_6 alkoxy carbonyloxy group, a benzyloxy carbonyloxy group

5 which may be substituted with a substituent group selected from Substituent group α , a C_1 - C_6

alkoxy group, a C_2 - C_6 alkenyloxy group, a C_2 - C_6 alkynyloxy group, a C_3 - C_6 cycloalkyloxy group,

a cyanomethyleneoxy group, a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy group, a C_1 - C_6 alkyl carbonyloxy

group, a C_1 - C_6 haloalkyl carbonyloxy group, a C_2 - C_6 alkenyl carbonyloxy group, a C_2 - C_6

halolalkenyl carbonyloxy group, a C_2 - C_6 alkynyl carbonyloxy group, a C_2 - C_6

10 halolalkynyl carbonyloxy group, a C_1 - C_6 alkoxy carbonyl C_1 - C_6 alkoxy group, a phenyloxy group

which may be substituted with a substituent group selected from Substituent group α , a

benzyloxy group which may be substituted with a substituent group selected from Substituent

group α , a phenyl carbonyloxy group which may be substituted with a substituent group selected

from Substituent group α , a benzyl carbonyloxy group which may be substituted with a

15 substituent group selected from Substituent group α , a phenyl carbonyl C_1 - C_6 alkyloxy group

which may be substituted with a substituent group selected from Substituent group α , a C_1 - C_{10}

alkylsulfonyloxy group, a phenylsulfonyloxy group which may be substituted with a substituent

group selected from Substituent group α , a benzylsulfonyloxy group which may be substituted

with a substituent group selected from Substituent group α , a C_1 - C_{10} alkylthio group, a C_1 - C_{10}

20 alkylsulfinyl group, a C_1 - C_{10} alkylsulfonyl group, a C_1 - C_6 haloalkylthio group, a C_1 - C_6

haloalkylsulfinyl group, a C_1 - C_6 haloalkylsulfonyl group, a C_2 - C_6 alkenylthio group, a C_2 - C_6

alkenylsulfinyl group, a C_2 - C_6 alkenylsulfonyl group, a C_2 - C_6 alkynylthio group, a C_2 - C_6

alkynylsulfinyl group, a C_2 - C_6 alkynylsulfonyl group, a phenylthio group which may be

substituted with a substituent group selected from Substituent group α , a benzylthio group which

25 may be substituted with a substituent group selected from Substituent group α , a phenylsulfinyl

group which may be substituted with a substituent group selected from Substituent group α , a

benzylsulfinyl group which may be substituted with a substituent group selected from Substituent group α , a phenylsulfonyl group which may be substituted with a substituent group selected from Substituent group α , a benzylsulfonyl group which may be substituted with a substituent group selected from Substituent group α , a C_1 - C_{10} alkylamino group, a di(C_1 - C_{10} alkyl)amino group, a C_1 - C_6 alkoxycarbonyl amino group, a C_1 - C_6 alkoxy group substituted with a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α], a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α], or a heterocyclic oxy group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α], and X represents a halogen atom).

Specifically, the compound of Formula 1b can be produced by reacting the compound of Formula 1a and a halogenating agent, and Formula 1c can be produced by reacting the compound of Formula 1b and a nucleophilic reagent in the presence of a base.

Examples of the halogenating agent that can be used for preparation of the compound of Formula 1b from the compound of Formula 1a include thionyl chloride, thionyl bromide, phosphorus oxy chloride, phosphorus oxy bromide, phenyltrimethyl ammonium tribromide, and Meldrum's acid tribromide. The use amount of the halogenating agent can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 1a. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature may be selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

5 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Examples of the nucleophilic reagent for the process for obtaining Formula 1c from Formula 1b, which is a compound represented by the formula $R^{4a}-H$, include alcohols like methanol, ethanol, and benzyl alcohol; mercaptans like methyl mercaptan and ethyl mercaptan; amines like ammonia, methyl amine, and ethyl amine; phenols like p-cresol and phenol; thiophenols like
10 p-chlorothiophenol; a C_1 - C_6 alkyl acids like acetic acid, and benzoic acids. The use amount of the nucleophilic reagent can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 1b. Preferably, it is from 1.0 to 1.2 mol.

Examples of the base which can be used for the present process include those described above for Process 1 of Production method 1.

15 Examples of the solvent which can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.


20 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 1c, which is produced according to this method, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

<Production method 3>

25 Compound of Formula 1c can be also produced by the method with the following reaction scheme.

[Chem. 9]



(in the formula, R^1 , R^2 , A_1 , A_2 , A_3 , Y and Z each have the same definitions as above, R^{4a} represents a hydroxycarbonyl group, a C_1 - C_6 alkoxy carbonyl group, a benzyloxycarbonyl group which may be substituted with a substituent group selected from Substituent group α , a C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a C_3 - C_6 cycloalkyl group, a cyanomethylene group, a C_3 - C_6 cycloalkyl C_1 - C_6 alkyl group, a C_1 - C_6 alkyl carbonyl group, a C_1 - C_{10} alkylthiocarbonyl group, a C_1 - C_6 haloalkyl carbonyl group, a C_2 - C_6 alkenyl carbonyl group, a C_2 - C_6 halolalkenyl carbonyl group, a C_2 - C_6 alkynyl carbonyl group, a C_2 - C_6 halolalkynyl carbonyl group, a C_1 - C_6 alkoxy carbonyl C_1 - C_6 alkyl group, a C_1 - C_{10} alkylsulfonyl group, a phenyl group which may be substituted with a substituent group selected from Substituent group α , a benzyl group which may be substituted with a substituent group selected from Substituent group α , a phenyl carbonyl group which may be substituted with a substituent group selected from Substituent group α , a benzyl carbonyl group which may be substituted with a substituent group selected from Substituent group α , a phenylsulfonyl group which may be substituted with a substituent group selected from Substituent group α , a phenyl carbonyl C_1 - C_6 alkyl group which may be substituted with a substituent group selected from Substituent group α , or a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α]).

Specifically, the compound of Formula 1c can be produced by reacting the compound of Formula 1a and an electrophilic reagent in a solvent in the presence or absence of a base.

The electrophilic reagent indicates a compound represented by the formula R^{4b} - L_a (L_a represents a leaving group), and examples thereof include C_1 - C_6 alkyl halide like methyl iodide

and propyl chloride; benzyl halide like benzyl bromide; C₁-C₆ alkylcarbonyl halide like acetyl chloride and propionyl chloride; benzoyl halide like benzoyl chloride; C₂-C₆ alkenylcarbonyl halide like methacryl chloride and crotonyl chloride; C₂-C₆ alkynylcarbonyl halide like 4-pentynoyl chloride; C₁-C₆ alkyl sulfonyl halide like methane sulfonyl chloride and ethane sulfonyl chloride; benzene sulfonyl halide like benzene sulfonyl chloride and p-toluene sulfonyl chloride; and di C₁-C₆ alkyl sulfate ester like dimethyl sulfate and diethyl sulfate. The use amount of the electrophilic reagent can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 1a. Preferably, it is from 1.0 to 1.2 mol.

Examples of the base which can be used for the present process include those described above for Process 1 of Production method 1. The use amount of the base can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 1a. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described above for Process 1 of Production method 1.


The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 1c, which is a target compound of this process, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

Formula 1c of the invention has many tautomers shown below, and they are all included in the invention.


[Chem.10]



<Production method 4>

Compound of Formula 1d can be also produced by the method with the following reaction scheme.

5 [Chem. 11]



(in the formula, R^1 , R^2 , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , Y and Z each have the same definitions as

above and Q represents a leaving group like a halogen atom, alkylcarbonyloxy group, an alkoxy carbonyloxy group, a haloalkylcarbonyloxy group, a haloalkoxy carbonyloxy group, a benzoyloxy group, a pyridyl group, and an imidazolyl group, as described above).

Specifically, compound of Formula 1d can be produced by reacting the compound of Formula 6a and the compound of Formula 4a in a solvent, in the presence of a Lewis acid.

The use amount of Formula 4a can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 6a. Preferably, it is from 1.0 to 1.2 mol.


Examples of the Lewis acid that can be used include organo lithium compounds like methyl lithium, ethyl lithium, n-butyl lithium, sec-butyl lithium, tert-butyl lithium, and benzyl lithium; Grignard's reagent like methyl magnesium iodide and ethyl magnesium bromide; metal compounds like lithium, potassium and sodium; organo copper compounds produced from Grignard's reagent or organometallic compound and monovalent copper salt; alkali metal amides like lithium diisopropyl amide (LDA), and; organic amines like triethylamine, pyridine, 4-dimethylaminopyridine, N,N-dimethylaniline, and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). n-Butyl lithium and lithium diisopropyl amide (LDA) are particularly preferable. The use amount of Lewis acid can be appropriately selected from the range of 0.5 to 10 mol per 1 mol of Formula 5a. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described above for Process 1 of Production method 1. Diethyl ether and tetrahydrofuran are particularly preferable. The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 1d, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

Formula 1d of the invention has many tautomers shown below, and they are all included in the invention.


[Chem. 12]



<Production method 5>

Compound of Formula 1e can be also produced by the method with the following reaction scheme.

5 [Chem. 13]



(in the formula, R^1 , R^2 , R^{20} , R^{21} , Y and Z each have the same definitions as above and Q represents a leaving group like a halogen atom, alkylcarbonyloxy group, an alkoxycarbonyloxy group, a haloalkylcarbonyloxy group, a haloalkoxycarbonyloxy group, a benzoyloxy group, a pyridyl group, and an imidazolyl group, as described above).

Specifically, the compound of Formula 5c can be produced by reacting the compound of Formula 6 and the compound of Formula 4a in a solvent in the presence of a base, and the

compound of Formula 1e can be produced by reacting the compound of Formula 5c and a cyano compound in the presence of a base.

In the above reaction, use amount of Formula 4a for preparing Formula 5c from Formula 6 can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 6.

5 Preferably, it is from 1.0 to 1.2 mol.

Examples of the base and solvent that can be used include those described above for Process 1 of Production method 1. The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the
10 reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Examples of the cyano compound which can be used for the reaction above for obtaining Formula 1e from Formula 5c include potassium cyanide, sodium cyanide, acetone cyanohydrin, hydrogen cyanide, and a polymer supported with hydrogen cyanide. The use amount of the cyano compound can be appropriately selected from the range of 0.01 to 1.0 mol per 1 mol of
15 Formula 6. Preferably, it is 0.05 to 0.2 mol.

Examples of the base that can be used include those described above for Process 1 of Production method 1. The use amount of the base can be appropriately selected from the range of 0.1 to 1.0 mol per 1 mol of Formula 6. Preferably, it is 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described
20 above for Process 1 of Production method 1.


The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

25 After the completion of the reaction, the compound of Formula 1e, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

Formula 1e of the invention has many tautomers shown below, and they are all included in the invention.


[Chem.14]



<Production method 6>

Compound of Formula 1g in which the substituent group in the pyrazole ring is modified
 5 can be also produced from the compound of Formula 1e by the method with the following reaction scheme.

[Chem. 15]



10 (in the formula, R^1 , R^2 , R^{20} , R^{21} , Y and Z each have the same definitions as above, R^{22a} represents an amino group, a cyano group, an isothiocyanate group, an isocyanate group, a hydroxycarbonyloxy group, a C_1 - C_6 alkoxy carbonyloxy group, a benzyloxycarbonyloxy group which may be substituted with a substituent group selected from Substituent group α , a C_1 - C_6 alkoxy group, a C_2 - C_6 alkenyloxy group, a C_2 - C_6 alkynyloxy group, a C_3 - C_6 cycloalkyloxy group,
 15 a cyanomethyleneoxy group, a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy group, a C_1 - C_6 alkylcarbonyloxy

group, a C₁-C₆ haloalkylcarbonyloxy group, a C₂-C₆ alkenylcarbonyloxy group, a C₂-C₆ halolalkenylcarbonyloxy group, a C₂-C₆ alkynylcarbonyloxy group, a C₂-C₆ halolalkynylcarbonyloxy group, a C₁-C₆ alkoxycarbonyl C₁-C₆ alkoxy group, a phenyloxy group which may be substituted with a substituent group selected from Substituent group α , a

5 benzyloxy group which may be substituted with a substituent group selected from Substituent group α , a phenylcarbonyloxy group which may be substituted with a substituent group selected from Substituent group α , a benzylcarbonyloxy group which may be substituted with a substituent group selected from Substituent group α , a phenylcarbonyl C₁-C₆ alkyloxy group which may be substituted with a substituent group selected from Substituent group α , a C₁-C₁₀

10 alkylsulfonyloxy group, a phenylsulfonyloxy group which may be substituted with a substituent group selected from Substituent group α , a benzylsulfonyloxy group which may be substituted with a substituent group selected from Substituent group α , a C₁-C₁₀ alkylthio group, a C₁-C₁₀ alkylsulfinyl group, a C₁-C₁₀ alkylsulfonyl group, a C₁-C₆ haloalkylthio group, a C₁-C₆ haloalkylsulfinyl group, a C₁-C₆ haloalkylsulfonyl group, a C₂-C₆ alkenylthio group, a C₂-C₆

15 alkenylsulfinyl group, a C₂-C₆ alkenylsulfonyl group, a C₂-C₆ alkynylthio group, a C₂-C₆ alkynylsulfinyl group, a C₂-C₆ alkynylsulfonyl group, a phenylthio group which may be substituted with a substituent group selected from Substituent group α , a benzylthio group which may be substituted with a substituent group selected from Substituent group α , a phenylsulfinyl group which may be substituted with a substituent group selected from Substituent group α , a

20 benzylsulfinyl group which may be substituted with a substituent group selected from Substituent group α , a phenylsulfonyl group which may be substituted with a substituent group selected from Substituent group α , a benzylsulfonyl group which may be substituted with a substituent group selected from Substituent group α , a C₁-C₁₀ alkylamino group, a di(C₁-C₁₀ alkyl)amino group, a C₁-C₆ alkoxycarbonyl amino group, a C₁-C₆ alkoxy group substituted with a heterocyclic group

25 having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α], a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or

different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α], or a heterocyclic oxy group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α], and X represents a halogen atom).

Specifically, the compound of Formula 1f can be produced by reacting the compound of Formula 1e and a halogenating agent and Formula 1g can be produced by reacting it with a nucleophilic reagent.

Examples of the halogenating agent that can be used for the process of producing the compound of Formula 1f from the compound of Formula 1e include thionyl chloride, thionyl bromide, phosphorus oxychloride, phosphorus oxybromide, phenyltrimethyl ammonium tribromide, and Meldrum's acid tribromide.

The use amount of the halogenating agent can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 1e. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described above for Process 1 of Production method 1. The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

The nucleophilic reagent for the process for obtaining Formula 1g from Formula 1f is, for example, a compound represented by the formula $R^{22a}-H$, and examples thereof include alcohols like methanol, ethanol, and benzyl alcohol; mercaptans like methyl mercaptan and ethyl mercaptan; amines like ammonia, methyl amine, and ethyl amine; phenols like p-cresol and phenol; thiophenols like p-chlorothiophenol; C_1 - C_6 alkyl acids like acetic acid, and benzoic acids. The use amount of the nucleophilic reagent can be appropriately selected from the range of 0.1 to

10 mol per 1 mol of Formula 1f. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent which can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.


The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 1g, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

<Production method 7>

Compound of Formula 1g can be also produced by the method with the following reaction scheme.

[Chem. 16]



(in the formula, R^1 , R^2 , R^{20} , R^{21} , Y and Z each have the same definitions as above, R^{22b} represents a hydroxycarbonyl group, a C_1 - C_6 alkoxy carbonyl group, a benzyloxycarbonyl group which may be substituted with a substituent group selected from Substituent group α , a C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a C_3 - C_6 cycloalkyl group, a cyanomethylene group, a C_3 - C_6 cycloalkyl C_1 - C_6 alkyl group, a C_1 - C_6 alkyl carbonyl group, a C_1 - C_{10} alkylthiocarbonyl group, a C_1 - C_6 haloalkyl carbonyl group, a C_2 - C_6 alkenyl carbonyl group, a C_2 - C_6 halolalkenyl carbonyl group, a C_2 - C_6 alkynyl carbonyl group, a C_2 - C_6 halolalkynyl carbonyl group, a C_1 - C_6 alkoxy carbonyl C_1 - C_6 alkyl group, a C_1 - C_{10} alkylsulfonyl group, a phenyl group which may be substituted with a substituent group selected from Substituent group α , a benzyl group which may be substituted with a substituent group selected from Substituent group α , a phenyl carbonyl group which may be substituted with a substituent

20

25

group selected from Substituent group α , a benzylcarbonyl group which may be substituted with a substituent group selected from Substituent group α , a phenylsulfonyl group which may be substituted with a substituent group selected from Substituent group α , a phenylcarbonyl C₁-C₆ alkyl group which may be substituted with a substituent group selected from Substituent group α ,
5 or a heterocyclic group having 3 to 10 carbon atoms and one or more heteroatoms that are the same or different from each other and selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α]).

10 Specifically, the compound of Formula 1g can be produced by reacting the compound of Formula 1e and an electrophilic reagent in a solvent, in the presence or absence of a base.

The electrophilic reagent that can be used indicates a compound represented by the formula R^{22b}-L_a (L_a represents a leaving group), and examples thereof include C₁-C₆ alkyl halide like methyl iodide and propyl chloride; benzyl halide like benzyl bromide; C₁-C₆ alkylcarbonyl halide
15 like acetyl chloride and propionyl chloride; benzoyl halide like benzoyl chloride; C₂-C₆ alkenylcarbonyl halide like methacryl chloride and crotonyl chloride; C₂-C₆ alkynylcarbonyl halide like 4-pentynoyl chloride; C₁-C₆ alkyl sulfonyl halide methane sulfonyl chloride and ethane sulfonyl chloride; benzene sulfonyl halide like benzene sulfonyl chloride and p-toluene sulfonyl chloride; and di C₁-C₆ alkyl sulfate ester like dimethyl sulfate and diethyl sulfate. The use
20 amount of the electrophilic reagent can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 1e. Preferably, it is from 1.0 to 1.2 mol.

Examples of the base and the solvent which can be used for the present process include those described above for Process 1 of Production method 1.

The use amount of the base can be appropriately selected from the range of 0.1 to 10 mol per
25 1 mol of Formula 1e. Preferably, it is from 1.0 to 1.2 mol.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.


The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 1g, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

<Production method 8>

5 Compound of Formula 1h can be also produced by the method with the following reaction scheme.

[Chem. 17]



10 (in the formula, R¹, R², R²⁴, R²⁵, Y and Z each have the same definitions as above and Q represents a leaving group like a halogen atom, alkylcarbonyloxy group, an alkoxycarbonyloxy group, a haloalkylcarbonyloxy group, a haloalkoxycarbonyloxy group, a benzoyloxy group, a pyridyl group, and an imidazolyl group, as described above).

Specifically, the compound of Formula 5d can be produced by reacting the compound of
15 Formula 7 and the compound of Formula 4a in a solvent, in the presence of a base, and the compound of Formula 1h can be produced by reacting the compound of Formula 5d and a cyano compound in the presence of a base.

In the above reaction, use amount of Formula 4a for preparing Formula 5d from Formula 7 can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 7.

20 Preferably, it is from 1.0 to 1.2 mol.

Examples of the base that can be used include those described above for Process 1 of Production method 1. Use amount of the base can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 7. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used include those described above for Process 1 of
25 Production method 1.

Examples of the cyano compound which can be used for the reaction above for obtaining Formula 1h from Formula 5d include potassium cyanide, sodium cyanide, acetone cyanohydrin, hydrogen cyanide, and a polymer supported with hydrogen cyanide. The use amount of the cyano compound can be appropriately selected from the range of 0.01 to 1.0 mol per 1 mol of Formula 5d. Preferably, it is 0.05 to 0.2 mol.

Examples of the base that can be used include those described above for Process 1 of Production method 1. The use amount of the base can be appropriately selected from the range of 0.1 to 1.0 mol per 1 mol of Formula 5d. Preferably, it is 1.0 to 1.2 mol.


Examples of the solvent that can be used include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 1h, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

Formula 1h of the invention has many tautomers shown below, and they are all included in the invention.


[Chem. 18]



<Production method 9>

Compound of Formula 1i can be produced by the method with the following reaction scheme.

[Chem. 19]



(in the formula, R^1 , R^2 , R^{24} , Y and Z each have the same definitions as above, R^{25} represents a C_1 - C_6 alkoxy carbonyl group, R^{26} represents an alkoxy group, a haloalkoxy group, a cycloalkoxy group, or a dimethylamino group, and R^{27} represents an alkyl group or a benzyl group).

(Process 1)

10 In this process, Formula 8a can be prepared by reacting Formula 1h and acid with or without using a solvent.

Examples of the acid that can be used for the present process include sulfonic acids like p-toluene sulfonic acid. Use amount of the acid can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 1h. Preferably, it is from 1.0 to 1.2 mol.

15 Examples of the solvent that can be used include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C .

The reaction time varies depending on the reaction temperature, the reaction substrates, the

20 reaction amount, etc. In general, it is from 10 minutes to 48 hours.

(Process 2)

By reacting Formula 8a and an ortho formic acid ester compound in N,N-dimethylacetamide dimethyl acetal compound or acetic anhydride, Formula 8b can be obtained. Use amount of N,N-dimethylacetamide dimethyl acetal and ortho formic acid ester can be appropriately selected
5 from the range of 0.1 to 10 mol per 1 mol of Formula 8a. Preferably, it is from 1.0 to 3.0 mol.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 150°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

10 (Process 3)

Formula 8c can be obtained by reacting Formula 8a and carbon disulfide, and without isolation, adding with alkyl halide like methyl iodide or benzyl halide like benzyl bromide. Use amount of carbon disulfide can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 8a. Preferably, it is from 1.0 to 1.2 mol. Use amount of the halide can be
15 appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 8a. Preferably, it is 2.0 to 2.4 mol. Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

20 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

(Process 4 & Process 5)

Formula 1i can be obtained by reacting Formula 8b or Formula 8c obtained from Process 2 or Process 3 above and hydroxylamine chloride in a solvent.

25 Use amount of hydroxylamine chloride can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 8b or Formula 8c. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an

inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.


After the completion of the reaction, the compound of Formula 1i, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

Hereinbelow, a method of producing synthetic intermediates of the compounds of the invention is given.

<Production method 10>

Compound of Formula 3b can be produced by the method with the following reaction scheme.

[Chem. 20]



(in the formula, R^1 , R^2 , Y and Z each have the same definitions as above, R^{30} represents a phenyl group or an alkyl group, and M^1 represents sodium, potassium or trimethylsilyl).

(Route a)

Specifically, compound of Formula 10 can be obtained by reacting the compound of Formula 9 and diethyl ketomalonate. In addition, compound of Formula 13a can be obtained by reacting the compound of Formula 10 and the compound of Formula 11 or the compound of

Formula 12 in the presence of a base.

Use amount of diethyl ketomalonate for the process of producing Formula 10 from Formula 9 can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 9.

Preferably, it is from 1.0 to 1.2 mol.

5 Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

10 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Use amount of the compound of Formula 11 or the compound of Formula 12 for the process of producing Formula 13a from Formula 10 can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 10. Preferably, it is from 1.0 to 1.2 mol.

15 Examples of the base that can be used for the present process include those described above for Process 1 of Production method 1. Use amount of the base can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 10. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

20 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

(Route b)

25 Specifically, compound of Formula 15 can be obtained by reacting the compound of Formula 9 and the compound of Formula 14. In addition, compound of Formula 16 can be obtained by reacting the compound of Formula 15 and diethyl ketomalonate. In addition, compound of Formula 13a can be obtained by reacting the compound of Formula 16 and an alkylating agent in the presence of a base.

Use amount of the compound of Formula 14 for the process of producing Formula 15 from

Formula 9 can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 9. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

5 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Use amount of diethyl ketomalonate for the process of producing Formula 16 from Formula
10 15 can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 15. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

15 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Use amount of the alkylating agent for the process of producing Formula 13a from Formula
16 can be appropriately selected from the range of 1.0 to 3.0 mol per 1 mol of Formula 16.
20 Preferably, it is from 1.0 to 1.5 mol.

Examples of the alkylating agent that can be used include alkyl sulfates like dimethyl sulfate and diethyl sulfate; alkyl halides like methyl iodide, ethyl iodide, benzyl chloride, benzyl bromide, propargyl bromide, ethyl bromoacetate, and chloroacetonitrile, and; sulfonic acid esters like ethoxyethyl p-toluene sulfonate and cyclopentylmethane sulfonate.

25 Examples of the base that can be used for the present process include those described above for Process 1 of Production method 1. Use amount of the base can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 16. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

5 (Route c)

Specifically, compound of Formula 17a can be obtained by reacting the compound of Formula 11a and hydrazine hydrate. In addition, compound of Formula 18 can be obtained by reacting the compound of Formula 17 and diethyl ketomalonate. In addition, compound of Formula 13a can be obtained by reacting the compound of Formula 18 and an alkylating agent in
10 the presence of a base.

Use amount of hydrazine hydrate for the process of producing Formula 17a from Formula 11a can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 9. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described
15 above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

20 Use amount of diethyl ketomalonate for the process of producing Formula 18 from Formula 17a can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 17a. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

25 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C. The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Use amount of the alkylating agent for the process of producing Formula 13a from Formula

18 can be appropriately selected from the range of 1.0 to 3.0 mol per 1 mol of Formula 18.

Preferably, it is from 1.0 to 1.5 mol.

Examples of the alkylating agent that can be used include alkyl sulfates like dimethyl sulfate and diethyl sulfate; alkyl halides like methyl iodide, ethyl iodide, benzyl chloride, benzyl bromide, propargyl bromide, ethyl bromoacetate, and chloroacetonitrile, and; sulfonic acid esters like ethoxyethyl p-toluene sulfonate and cyclopentylmethane sulfonate.

Examples of the base that can be used for the present process include those described above for Process 1 of Production method 1. Use amount of the base can be appropriately selected from the range of 0.1 to 10 mol per 1 mol of Formula 18. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

(Route d)

Specifically, compound of Formula 17b can be obtained by reacting the compound of Formula 11a and the compound of Formula 9. In addition, compound of Formula 13a can be obtained by reacting the compound of Formula 17b and diethyl ketomalonate, using an acid or a base depending on the condition.

Use amount of the compound of Formula 9 for the process of producing Formula 17b from Formula 11a can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 9. Preferably, it is from 1.0 to 1.2 mol.

Examples of the acid that can be used include organic acids represented by organic sulfonic acid like p-toluene sulfonic acid, methane sulfonic acid, and benzene sulfonic acid; hydrogen halide acids represented by hydrochloric acid and hydrogen bromic acid, and; inorganic acids like sulfuric acid and phosphoric acid. These acids can be used either singly or in combination of two or more.

Examples of the base that can be used for the present process include those described above

for Process 1 of Production method 1.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

5 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

10 Use amount of diethyl ketomalonate for the process of producing Formula 13a from Formula 17b can be appropriately selected from the range of 1.0 to 1.5 mol per 1 mol of Formula 17b. Preferably, it is from 1.0 to 1.2 mol.

Examples of the solvent that can be used for the present process include those described above for Process 1 of Production method 1.

15 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

Examples of the acid include organic acids like p-toluene sulfonic acid.

20 Examples of the base include organic bases like triethylamine and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) and inorganic bases like sodium hydride, sodium methoxide, and sodium ethoxide.

After the completion of the reaction, the compound of Formula 13a, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

(Route e)

25 Specifically, compound of Formula 3b can be obtained by hydrolyzing the compound of Formula 13a.

With regard to the process of obtaining the compound of Formula 3b from the compound of Formula 13a, the production can be carried out by hydrolysis in water, organic solvent, or a mixture solvent in the presence of an acid or a base.

Examples of the base that can be used include those described above for Process 1 of Production method 1.

Use amount of the base can be appropriately selected from the range of 0.01 to 100 mol per 1 mol of Formula 13a. Preferably, it is 0.1 to 10 mol.

5 Examples of the acid that can be used include inorganic acids like hydrochloric acid, hydrobromic acid, and sulfuric acid, and organic acids like acetic acid and trifluoroacetic acid.

Use amount of the acid can be appropriately selected from the range of 1 mol to excess amount per 1 mol of Formula 13a. Preferably, it is from 1 to 100 mol.

10 Examples of the organic solvent that can be used include a mixture solvent of water and an organic solvent. Examples of the organic solvent include alcohols like methanol and ethanol, ether like tetrahydrofuran, ketones like acetone and methyl isobutyl ketone, amides like N,N-dimethyl formamide and N,N-dimethyl acetamide, sulfur compounds like dimethyl sulfoxide and sulfolane, acetonitrile, and their mixture.

15 Use amount of the solvent is 0.01 to 100 L per 1 mol of Formula 13a. Preferably, it is 0.1 to 10 L.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

20 (Route f)

Specifically, compound of Formula 13b can be obtained by reacting the compound of Formula 13a and a sulfurizing agent. In addition, the compound of Formula 3b can be obtained by hydrolyzing the compound of Formula 13b.

25 Use amount of the compound of the sulfurizing agent for the process of producing Formula 13b from Formula 13a can be appropriately selected from the range of 1.0 to 8.0 mol per 1 mol of Formula 13a. Preferably, it is from 1.0 to 4.0 mol.

Examples of the sulfurizing agent that can be used include diphosphorus pentoxide and 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide.

Use amount of the compound of the sulfurizing agent can be appropriately selected from the

range of 1.0 to 8.0 mol per 1 mol of Formula 13a. Preferably, it is 0.1 to 4.0 mol.

Examples of the solvent that can be used include those described above for Process 1 of Production method 1.

5 The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.

The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

10 With regard to the process of obtaining the compound of Formula 3b from the compound of Formula 13b, the production can be carried out by hydrolysis in water, organic solvent, or a mixture solvent in the presence of an acid or a base.

Examples of the base that can be used include those described above for Process 1 of Production method 1.

Use amount of the base can be appropriately selected from the range of 0.01 to 100 mol per 1 mol of Formula 13b. Preferably, it is 0.1 to 10 mol.

15 Examples of the acid that can be used include inorganic acids like hydrochloric acid, hydrobromic acid, and sulfuric acid, and organic acids like acetic acid and trifluoroacetic acid.

Use amount of the acid can be appropriately selected from the range of 1 mol to excess amount per 1 mol of Formula 13b. Preferably, it is from 1 to 100 mol.

20 Examples of the organic solvent that can be used include a mixture solvent of water and an organic solvent. Examples of the organic solvent include alcohols like methanol and ethanol, ether like tetrahydrofuran, ketones like acetone and methyl isobutyl ketone, amides like N,N-dimethyl formamide and N,N-dimethyl acetamide, sulfur compounds like dimethyl sulfoxide and sulfolane, acetonitrile, and their mixture.

25 Use amount of the solvent is 0.01 to 100 L per 1 mol of Formula 13b. Preferably, it is 0.1 to 10 L.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.


The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 3b, i.e., the target compound of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

<Intermediate synthesis method 1>

5 Compound of Formula 3a can be produced according to the method with the following reaction scheme.

[Chem. 21]



10 (in the formula, R¹, R², Y and Z each have the same definitions as above and X represents a chlorine or a bromine).

Specifically, Formula 3a can be produced by reacting Formula 3b and an appropriate halogenating agent with or without a solvent.

Examples of the halogenating agent that can be used include oxalyl chloride and thionyl chloride.

15 Use amount of the halogenating agent can be appropriately selected from the range of 0.01 to 20 mol per 1 mol of Formula 3b. Preferably, it is from 1 to 10 mol.

Examples of the solvent include halogenated hydrocarbons like dichloromethane and chloroform, ethers like diethyl ether and tetrahydrofuran, and aromatic hydrocarbons like benzene and toluene.

20 Use amount of the solvent is 0.01 to 100 L per 1 mol of Formula 3b. Preferably, it is 0.1 to 10 L.

The reaction temperature is selected from the range of from -20°C to the boiling point of an inert solvent used. Preferably, the reaction is carried out in the range of from 0°C to 100°C.


25 The reaction time varies depending on the reaction temperature, the reaction substrates, the reaction amount, etc. In general, it is from 10 minutes to 48 hours.

After the completion of the reaction, the compound of Formula 3a, i.e., the target compound

of this reaction, can be collected from the reaction system by general method, and if necessary, purified by a process like column chromatography and recrystallization.

Examples of the production intermediates [13a] and [3b], that can be described for the Production method 10, are shown in Table 44 to Table 67.

[Table 44]

				
Compound No.	R ¹	R ²	Y	Z
IV-1	Me	Me	O	O
IV-2	Et	Me	O	O
IV-3	Pr-n	Me	O	O
IV-4	Pr-i	Me	O	O
IV-5	Bu-n	Me	O	O
IV-6	Bu-i	Me	O	O
IV-7	Bu-s	Me	O	O
IV-8	Bu-t	Me	O	O
IV-9	Hex-n	Me	O	O
IV-10	CH ₂ CF ₃	Me	O	O
IV-11	CH ₂ CH=CH ₂	Me	O	O
IV-12	CH ₂ C(Me)=CH ₂	Me	O	O
IV-13	CH ₂ CH ₂ CH=CMe ₂	Me	O	O
IV-14	CH ₂ C≡CH	Me	O	O
IV-15	CH ₂ C≡CCH ₃	Me	O	O
IV-16	Pr-c	Me	O	O
IV-17	Bu-c	Me	O	O
IV-18	Pen-c	Me	O	O
IV-19	Hex-c	Me	O	O
IV-20	CH ₂ Pr-c	Me	O	O
IV-21	CH ₂ Bu-c	Me	O	O
IV-22	CH ₂ Pen-c	Me	O	O
IV-23	CH ₂ Hex-c	Me	O	O
IV-24	CH ₂ CH=CCl ₂	Me	O	O
IV-25	CH ₂ CCl=CHCl	Me	O	O
IV-26	CH ₂ CH ₂ CH=CCl ₂	Me	O	O
IV-27	CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O
IV-28	CH ₂ CH ₂ CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O
IV-29	CH ₂ CH=CF ₂	Me	O	O
IV-30	CH ₂ CH ₂ OMe	Me	O	O
IV-31	CH ₂ CH ₂ OEt	Me	O	O
IV-32	CH(Me)CH ₂ OMe	Me	O	O
IV-33	CH ₂ CH ₂ OCH ₂ CH ₂ OMe	Me	O	O
IV-34	CH ₂ CH ₂ OPr-n	Me	O	O
IV-35	CH ₂ CH ₂ OPr-i	Me	O	O
IV-36	CH ₂ CH ₂ OPr-c	Me	O	O

[Table 45]

Compound No.	R ¹	R ²	Y	Z
IV-37	CH ₂ CH ₂ OBu ^c	Me	O	O
IV-38	CH ₂ CH ₂ OPen ^c	Me	O	O
IV-39	CH ₂ CH ₂ OHex ^c	Me	O	O
IV-40	CH ₂ CH ₂ OCH ₂ CF ₃	Me	O	O
IV-41	CH ₂ CH ₂ CH ₂ OMe	Me	O	O
IV-42	CH=CHMe	Me	O	O
IV-43	CH ₂ SMe	Me	O	O
IV-44	CH ₂ SPr ⁿ	Me	O	O
IV-45	CH ₂ CH ₂ SMe	Me	O	O
IV-46	CH ₂ SOMe	Me	O	O
IV-47	CH ₂ SO ₂ Me	Me	O	O
IV-48	CH ₂ CH ₂ CH ₂ SMe	Me	O	O
IV-49	CH ₂ CH ₂ CH ₂ SO ₂ Me	Me	O	O
IV-50	Ph	Me	O	O
IV-51	Ph(2-Cl)	Me	O	O
IV-52	Ph(3-Cl)	Me	O	O
IV-53	Ph(4-Cl)	Me	O	O
IV-54	Ph(2-F)	Me	O	O
IV-55	Ph(3-F)	Me	O	O
IV-56	Ph(4-F)	Me	O	O
IV-57	Ph(2-Me)	Me	O	O
IV-58	Ph(3-Me)	Me	O	O
IV-59	Ph(4-Me)	Me	O	O
IV-60	Ph(2-OMe)	Me	O	O
IV-61	Ph(3-OMe)	Me	O	O
IV-62	Ph(4-OMe)	Me	O	O
IV-63	Ph(2-CF ₃)	Me	O	O
IV-64	Ph(3-CF ₃)	Me	O	O
IV-65	Ph(4-CF ₃)	Me	O	O
IV-66	Ph(2-NO ₂)	Me	O	O
IV-67	Ph(3-NO ₂)	Me	O	O
IV-68	Ph(4-NO ₂)	Me	O	O
IV-69	Ph(2-OCF ₃)	Me	O	O
IV-70	Ph(3-OCF ₃)	Me	O	O
IV-71	Ph(4-OCF ₃)	Me	O	O
IV-72	Ph(2-CN)	Me	O	O
IV-73	Ph(3-CN)	Me	O	O
IV-74	Ph(4-CN)	Me	O	O
IV-75	Ph(3,4-F ₂)	Me	O	O









[Table 46]

Compound No.	R ¹	R ²	Y	Z
IV-76	Ph(3,5-F ₂)	Me	O	O
IV-77	Ph(2,3-F ₂)	Me	O	O
IV-78	Ph(2,4-F ₂)	Me	O	O
IV-79	Ph(2,5-F ₂)	Me	O	O
IV-80	Ph(2,6-F ₂)	Me	O	O
IV-81	Ph(3,4-Cl ₂)	Me	O	O
IV-82	Ph(3,5-Cl ₂)	Me	O	O
IV-83	Ph(2,3-Cl ₂)	Me	O	O
IV-84	Ph(2,4-Cl ₂)	Me	O	O
IV-85	Ph(2,5-Cl ₂)	Me	O	O
IV-86	Ph(2,6-Cl ₂)	Me	O	O
IV-87	Ph(3,4-Me ₂)	Me	O	O
IV-88	Ph(3,5-Me ₂)	Me	O	O
IV-89	Ph(2,3-Me ₂)	Me	O	O
IV-90	Ph(2,4-Me ₂)	Me	O	O
IV-91	Ph(2,5-Me ₂)	Me	O	O
IV-92	Ph(2,6-Me ₂)	Me	O	O
IV-93	Ph(3,4-(OMe) ₂)	Me	O	O
IV-94	Ph(3,5-(OMe) ₂)	Me	O	O
IV-95	Ph(2,3-(OMe) ₂)	Me	O	O
IV-96	Ph(2,4-(OMe) ₂)	Me	O	O
IV-97	Ph(2,5-(OMe) ₂)	Me	O	O
IV-98	Ph(2,6-(OMe) ₂)	Me	O	O
IV-99	Ph(3-F-4-OMe)	Me	O	O
IV-100	Ph(3-F-5-OMe)	Me	O	O
IV-101	Ph(2-F-3-OMe)	Me	O	O
IV-102	Ph(2-F-4-OMe)	Me	O	O
IV-103	Ph(2-F-5-OMe)	Me	O	O
IV-104	Ph(2-F-6-OMe)	Me	O	O
IV-105	Ph(3-F-4-Me)	Me	O	O
IV-106	Ph(3-F-5-Me)	Me	O	O
IV-107	Ph(2-F-3-Me)	Me	O	O
IV-108	Ph(2-F-4-Me)	Me	O	O
IV-109	Ph(2-F-5-Me)	Me	O	O
IV-110	Ph(2-F-6-Me)	Me	O	O
IV-111	Ph(3-OMe-4-F)	Me	O	O
IV-112	Ph(2-OMe-3-F)	Me	O	O
IV-113	Ph(2-OMe-4-F)	Me	O	O
IV-114	Ph(2-OMe-5-F)	Me	O	O















[Table 47]

Compound No.	R ¹	R ²	Y	Z
IV-115	Ph(3-Me-4-F)	Me	O	O
IV-116	Ph(2-Me-3-F)	Me	O	O
IV-117	Ph(2-Me-4-F)	Me	O	O
IV-118	Ph(2-Me-5-F)	Me	O	O
IV-119	Ph(3-Cl-4-OMe)	Me	O	O
IV-120	Ph(3-Cl-5-OMe)	Me	O	O
IV-121	Ph(2-Cl-3-OMe)	Me	O	O
IV-122	Ph(2-Cl-4-OMe)	Me	O	O
IV-123	Ph(2-Cl-5-OMe)	Me	O	O
IV-124	Ph(2-Cl-6-OMe)	Me	O	O
IV-125	Ph(3-Cl-4-Me)	Me	O	O
IV-126	Ph(3-Cl-5-Me)	Me	O	O
IV-127	Ph(2-Cl-3-Me)	Me	O	O
IV-128	Ph(2-Cl-4-Me)	Me	O	O
IV-129	Ph(2-Cl-5-Me)	Me	O	O
IV-130	Ph(2-Cl-6-Me)	Me	O	O
IV-131	Ph(3-OMe-4-Cl)	Me	O	O
IV-132	Ph(2-OMe-3-Cl)	Me	O	O
IV-133	Ph(2-OMe-4-Cl)	Me	O	O
IV-134	Ph(2-OMe-5-Cl)	Me	O	O
IV-135	Ph(3-Me-4-Cl)	Me	O	O
IV-136	Ph(2-Me-3-Cl)	Me	O	O
IV-137	Ph(2-Me-4-Cl)	Me	O	O
IV-138	Ph(2-Me-5-Cl)	Me	O	O
IV-139	Ph(3-F-4-Cl)	Me	O	O
IV-140	Ph(3-F-5-Cl)	Me	O	O
IV-141	Ph(2-F-3-Cl)	Me	O	O
IV-142	Ph(2-F-4-Cl)	Me	O	O
IV-143	Ph(2-F-5-Cl)	Me	O	O
IV-144	Ph(2-F-6-Cl)	Me	O	O
IV-145	Ph(3-Cl-4-F)	Me	O	O
IV-146	Ph(2-Cl-3-F)	Me	O	O
IV-147	Ph(2-Cl-4-F)	Me	O	O
IV-148	Ph(2-Cl-5-F)	Me	O	O
IV-149	Ph(3-Me-4-OMe)	Me	O	O
IV-150	Ph(3-Me-5-OMe)	Me	O	O
IV-151	Ph(2-Me-3-OMe)	Me	O	O
IV-152	Ph(2-Me-4-OMe)	Me	O	O
IV-153	Ph(2-Me-5-OMe)	Me	O	O













[Table 48]

Compound No.	R ¹	R ²	Y	Z
IV-154	Ph(2-Me-6-OMe)	Me	O	O
IV-155	Ph(3-OMe-4-Me)	Me	O	O
IV-156	Ph(2-OMe-3-Me)	Me	O	O
IV-157	Ph(2-OMe-4-Me)	Me	O	O
IV-158	Ph(2-OMe-5-Me)	Me	O	O
IV-159	Ph(3-CN-4-OMe)	Me	O	O
IV-160	Ph(3-OMe-4-CN)	Me	O	O
IV-161	Ph(3-Me-4-CN)	Me	O	O
IV-162	Ph(3-CN-4-Me)	Me	O	O
IV-163	Ph(3-NO ₂ -4-OMe)	Me	O	O
IV-164	Ph(3-OMe-4-NO ₂)	Me	O	O
IV-165	Ph(3-Me-4-NO ₂)	Me	O	O
IV-166	Ph(3-NO ₂ -4-Me)	Me	O	O
IV-167	Ph(3,5-F ₂ -5-OMe)	Me	O	O
IV-168	Ph(3,5-F ₂ -5-Me)	Me	O	O
IV-169	Ph(3,4,5-(OMe) ₃)	Me	O	O
IV-170		Me	O	O
IV-171		Me	O	O
IV-172		Me	O	O
IV-173		Me	O	O
IV-174		Me	O	O
IV-175		Me	O	O
IV-176		Me	O	O
IV-177		Me	O	O

[Table 49]

Compound No.	R ¹	R ²	Y	Z
IV-178		Me	O	O
IV-179		Me	O	O
IV-180		Me	O	O
IV-181		Me	O	O
IV-182		Me	O	O
IV-183		Me	O	O
IV-184		Me	O	O
IV-185		Me	O	O
IV-186		Me	O	O
IV-187		Me	O	O
IV-188		Me	O	O
IV-189		Me	O	O
IV-190		Me	O	O
IV-191		Me	O	O






[Table 50]

Compound No.	R ¹	R ²	Y	Z
IV-192		Me	O	O
IV-193		Me	O	O
IV-194		Me	O	O
IV-195		Me	O	O
IV-196		Me	O	O
IV-197		Me	O	O
IV-198		Me	O	O
IV-199		Me	O	O
IV-200		Me	O	O
IV-201		Me	O	O
IV-202		Me	O	O
IV-203		Me	O	O
IV-204		Me		O












[Table 51]

Compound No.	R ¹	R ²	Y	Z
IV-205	CH ₂ Ph	Me	O	O
IV-206	CH ₂ CH ₂ Ph	Me	O	O
IV-207	CH ₂ CH ₂ CH ₂ Ph	Me	O	O
IV-208	CH ₂ CH=CHPh	Me	O	O
IV-209	CH ₂ C≡CPh	Me	O	O
IV-210	CH ₂ CH=NOMe	Me	O	O
IV-211	CH ₂ CH=NOEt	Me	O	O
IV-212	CH ₂ CH=NOPr ⁿ	Me	O	O
IV-213	CH ₂ CH=NOPh	Me	O	O
IV-214	CH ₂ CH(OMe) ₂	Me	O	O
IV-215	CH ₂ CHO	Me	O	O
IV-216	NH ₂	Me	O	O
IV-217	NHMe	Me	O	O
IV-218	NHEt	Me	O	O
IV-219	NHPr ⁿ	Me	O	O
IV-220	NHPr ⁱ	Me	O	O
IV-221	NHBu ⁿ	Me	O	O
IV-222	NHBu ⁱ	Me	O	O
IV-223	NHBu ^s	Me	O	O
IV-224	NHCH ₂ Pr ^c	Me	O	O
IV-225	NHPen ⁿ	Me	O	O
IV-226	NHHex ⁿ	Me	O	O
IV-227	NHCH ₂ CH ₂ CH ₂ Cl	Me	O	O
IV-228	NHCH ₂ CH ₂ CH ₂ F	Me	O	O
IV-229	NHCH ₂ CH ₂ OMe	Me	O	O
IV-230	NMe ₂	Me	O	O
IV-231	NEt ₂	Me	O	O
IV-232	N(Pr ⁿ) ₂	Me	O	O
IV-233	N(Bu ⁿ) ₂	Me	O	O
IV-234	N(Me)Et	Me	O	O
IV-235	N(Me)CH ₂ CH ₂ OMe	Me	O	O
IV-236	NHPh	Me	O	O
IV-237	NHCH ₂ Ph	Me	O	O
IV-238	N=CMe ₂	Me	O	O
IV-239	N=CEt ₂	Me	O	O
IV-240	N=CHNMe ₂	Me	O	O
IV-241	NHC(=O)Me	Me	O	O
IV-242	N[C(=O)Me] ₂	Me	O	O
IV-243	NHC(=O)OMe	Me	O	O
IV-244	N[C(=O)OMe] ₂	Me	O	O
IV-245	NHSO ₂ Me	Me	O	O







[Table 52]

Compound No.	R ¹	R ²	Y	Z
IV-246	NHSO ₂ Ph	Me	O	O
IV-247	NHSO ₂ CH ₂ Ph	Me	O	O
IV-248	OMe	Me	O	O
IV-249	OEt	Me	O	O
IV-250	OPr-n	Me	O	O
IV-251	OPr-i	Me	O	O
IV-252	OCH ₂ Pr-c	Me	O	O
IV-253	OCH ₂ Cl	Me	O	O
IV-254	OCHCl ₂	Me	O	O
IV-255	OCCL ₃	Me	O	O
IV-256	OCH ₂ F	Me	O	O
IV-257	OCHF ₂	Me	O	O
IV-258	OCF ₃	Me	O	O
IV-259	Ph	Et	O	O
IV-260	Ph	Pr-i	O	O
IV-261	Ph	CHF ₂	O	O
IV-262	Ph	Ph	O	O
IV-263	Ph	Me	O	S
IV-264	Ph	Me	S	S
IV-265	Me	Me	O	S
IV-266	Me	Me	S	S
IV-267	Ph	Me	O	O
IV-268	Ph(4-OEt)	Me	O	O
IV-269	Ph(2-Ph)	Me	O	O
IV-270	Ph(3-Ph)	Me	O	O
IV-271	Ph(4-Ph)	Me	O	O
IV-272		Me	O	O
IV-273		Me	O	O
IV-274	Me		O	O
IV-275	Et		O	O
IV-276		Me	O	O






[Table 53]

Compound No.	R ¹	R ²	Y	Z
IV-277		Me	O	O
IV-278		Me	O	O
IV-279		Me	O	O
IV-280		Me	O	O
IV-281		Me	O	O
IV-282	Ph(2-Me-4-Br)		O	O
IV-283	Ph(2-Me-4-I)	Me	O	O
IV-284	Ph(2-Me-4-CF ₃)	Me	O	O
IV-285	Ph(2-Me-4-OCF ₃)	Me	O	O
IV-286	Ph(2-Pr-i)	Me	O	O
IV-287		Me	O	O
IV-288	Ph(2-Et)	Me	O	O
IV-289		Me	O	O
IV-290		Me	O	O
IV-291		Me	O	S
IV-292		Me	O	O
IV-293		Me	O	O
IV-294	CH ₂ COOBu-t	Me	O	O
IV-295	(C ₇ H ₁₄)CH ₃	Me	O	O
IV-296	(C ₉ H ₁₈)CH ₃	Me	O	O
IV-297	Ph(2-F,4-Cl,5-OMe)	Me	O	O
IV-298	Ph(2,3,4-(OMe) ₃)	Me	O	O
IV-299	Ph(3,5-Cl ₂ -4-OMe)	Me	O	O
IV-300	Ph(3,5-Cl ₂ -4-SMe)	Me	O	O


[Table 54]

Compound No.	R ¹	R ²	Y	Z
IV-301	Ph(3,5-Cl ₂ -4-SO ₂ Me)	Me	O	O
IV-302	Ph(3,4,5-F ₃)	Me	O	O
IV-303		Me	O	O
IV-304		Me	O	O
IV-305		Me	O	O
IV-306	Bu-n		O	O
IV-307	CH ₂ CH(CH ₃) ₂		O	O
IV-308	Ph	Pen-n	O	O
IV-309	H	Me	O	O
IV-310	CH ₂ C≡CF	Me	O	O
IV-311		Me	O	O
IV-312		Me	O	O
IV-313	CH ₂ NH ₂	Me	O	O
IV-314	CH ₂ NO ₂	Me	O	O
IV-315	CH ₂ NHCH ₃	Me	O	O
IV-316	CH ₂ N(CH ₃) ₂	Me	O	O
IV-317	CH ₂ SCH ₂ CF ₃	Me	O	O
IV-318	CH ₂ SOCH ₂ CF ₃	Me	O	O
IV-319	CH ₂ SO ₂ CH ₂ CF ₃	Me	O	O
IV-320	CH ₂ OH	Me	O	O
IV-321	CH ₂ OBn	Me	O	O
IV-322	CH ₂ OCH ₂ Pr-c	Me	O	O
IV-323	CH ₂ OPh	Me	O	O
IV-324	CH ₂ SPh	Me	O	O
IV-325	CH ₂ SOPh	Me	O	O
IV-326	CH ₂ SO ₂ Ph	Me	O	O
IV-327	CH ₂ CON(CH ₃) ₂	Me	O	O
IV-328	CH ₂ COCH ₃	Me	O	O
IV-329	CH ₂ OCOCH ₃	Me	O	O

[Table 55]

Compound No.	R ¹	R ²	Y	Z
IV-330	CH ₂ ON=CHCH ₃	Me	O	O
IV-331	C ₂ H ₄ OC ₂ H ₄ SCH ₃	Me	O	O
IV-332	C ₂ H ₄ OC ₂ H ₄ SOCH ₃	Me	O	O
IV-333	C ₂ H ₄ OC ₂ H ₄ SO ₂ CH ₃	Me	O	O
IV-334	CH ₂ OCH ₂ CN	Me	O	O
IV-335	CH ₂ CN	Me	O	O
IV-336	OCH ₂ CH=CH ₂	Me	O	O
IV-337	OCH ₂ C≡CH	Me	O	O
IV-338	OPr-c	Me	O	O
IV-339		Me	O	O
IV-340		Me	O	O
IV-341		Me	O	O
IV-342		Me	O	O
IV-343		Me	O	O
IV-344	Ph	H	O	O
IV-345	Ph	CH ₂ CH=CH ₂	O	O
IV-346	Ph	CH ₂ C≡CH	O	O
IV-347	Ph	Pr-c	O	O
IV-348	Ph	CH ₂ CH=CF ₂	O	O
IV-349	Ph	CH ₂ C≡CF	O	O
IV-350	Ph	C ₂ H ₄ OCH ₃	O	O
IV-351	Ph	C ₂ H ₄ OC ₂ H ₅	O	O
IV-352	Ph	CH(Me)OEt	O	O
IV-353	Ph	CH ₂ OPr-c	O	O
IV-354	Ph	CH(OCH ₃) ₂	O	O
IV-355	Ph	CH ₂ Ph	O	O
IV-356	Ph	CH=CH·Ph	O	O
IV-357	Ph	C≡C·Ph	O	O

[Table 56]

				
Compound No.	R ¹	R ²	Y	Z
V-1	Me	Me	O	O
V-2	Et	Me	O	O
V-3	Pr-n	Me	O	O
V-4	Pr-i	Me	O	O
V-5	Bu-n	Me	O	O
V-6	Bu-i	Me	O	O
V-7	Bu-s	Me	O	O
V-8	Bu-t	Me	O	O
V-9	Hex-n	Me	O	O
V-10	CH ₂ CF ₃	Me	O	O
V-11	CH ₂ CH=CH ₂	Me	O	O
V-12	CH ₂ C(Me)=CH ₂	Me	O	O
V-13	CH ₂ CH ₂ CH=CMe ₂	Me	O	O
V-14	CH ₂ C≡CH	Me	O	O
V-15	CH ₂ C≡CCH ₃	Me	O	O
V-16	Pr-c	Me	O	O
V-17	Bu-c	Me	O	O
V-18	Pen-c	Me	O	O
V-19	Hex-c	Me	O	O
V-20	CH ₂ Pr-c	Me	O	O
V-21	CH ₂ Bu-c	Me	O	O
V-22	CH ₂ Pen-c	Me	O	O
V-23	CH ₂ Hex-c	Me	O	O
V-24	CH ₂ CH=CCl ₂	Me	O	O
V-25	CH ₂ CCl=CHCl	Me	O	O
V-26	CH ₂ CH ₂ CH=CCl ₂	Me	O	O
V-27	CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O
V-28	CH ₂ CH ₂ CH ₂ CH ₂ C(Me)=CF ₂	Me	O	O
V-29	CH ₂ CH=CF ₂	Me	O	O
V-30	CH ₂ CH ₂ OMe	Me	O	O
V-31	CH ₂ CH ₂ OEt	Me	O	O
V-32	CH(Me)CH ₂ OMe	Me	O	O
V-33	CH ₂ CH ₂ OCH ₂ CH ₂ OMe	Me	O	O
V-34	CH ₂ CH ₂ OPr-n	Me	O	O
V-35	CH ₂ CH ₂ OPr-i	Me	O	O
V-36	CH ₂ CH ₂ OPr-c	Me	O	O
V-37	CH ₂ CH ₂ OBu-c	Me	O	O
V-38	CH ₂ CH ₂ OPen-c	Me	O	O

[Table 57]

Compound No.	R ¹	R ²	Y	Z
V-39	CH ₂ CH ₂ OH _{hex-c}	Me	O	O
V-40	CH ₂ CH ₂ OCH ₂ CF ₃	Me	O	O
V-41	CH ₂ CH ₂ CH ₂ OMe	Me	O	O
V-42	CH=CHMe	Me	O	O
V-43	CH ₂ SMe	Me	O	O
V-44	CH ₂ SPr-n	Me	O	O
V-45	CH ₂ CH ₂ SMe	Me	O	O
V-46	CH ₂ SOMe	Me	O	O
V-47	CH ₂ SO ₂ Me	Me	O	O
V-48	CH ₂ CH ₂ CH ₂ SMe	Me	O	O
V-49	CH ₂ CH ₂ CH ₂ SO ₂ Me	Me	O	O
V-50	Ph	Me	O	O
V-51	Ph(2-Cl)	Me	O	O
V-52	Ph(3-Cl)	Me	O	O
V-53	Ph(4-Cl)	Me	O	O
V-54	Ph(2-F)	Me	O	O
V-55	Ph(3-F)	Me	O	O
V-56	Ph(4-F)	Me	O	O
V-57	Ph(2-Me)	Me	O	O
V-58	Ph(3-Me)	Me	O	O
V-59	Ph(4-Me)	Me	O	O
V-60	Ph(2-OMe)	Me	O	O
V-61	Ph(3-OMe)	Me	O	O
V-62	Ph(4-OMe)	Me	O	O
V-63	Ph(2-CF ₃)	Me	O	O
V-64	Ph(3-CF ₃)	Me	O	O
V-65	Ph(4-CF ₃)	Me	O	O
V-66	Ph(2-NO ₂)	Me	O	O
V-67	Ph(3-NO ₂)	Me	O	O
V-68	Ph(4-NO ₂)	Me	O	O
V-69	Ph(2-OCF ₃)	Me	O	O
V-70	Ph(3-OCF ₃)	Me	O	O
V-71	Ph(4-OCF ₃)	Me	O	O
V-72	Ph(2-CN)	Me	O	O
V-73	Ph(3-CN)	Me	O	O
V-74	Ph(4-CN)	Me	O	O
V-75	Ph(3,4-F ₂)	Me	O	O
V-76	Ph(3,5-F ₂)	Me	O	O
V-77	Ph(2,3-F ₂)	Me	O	O
V-78	Ph(2,4-F ₂)	Me	O	O










[Table 58]

Compound No.	R ¹	R ²	Y	Z
V-79	Ph(2,5-F ₂)	Me	O	O
V-80	Ph(2,6-F ₂)	Me	O	O
V-81	Ph(3,4-Cl ₂)	Me	O	O
V-82	Ph(3,5-Cl ₂)	Me	O	O
V-83	Ph(2,3-Cl ₂)	Me	O	O
V-84	Ph(2,4-Cl ₂)	Me	O	O
V-85	Ph(2,5-Cl ₂)	Me	O	O
V-86	Ph(2,6-Cl ₂)	Me	O	O
V-87	Ph(3,4-Me ₂)	Me	O	O
V-88	Ph(3,5-Me ₂)	Me	O	O
V-89	Ph(2,3-Me ₂)	Me	O	O
V-90	Ph(2,4-Me ₂)	Me	O	O
V-91	Ph(2,5-Me ₂)	Me	O	O
V-92	Ph(2,6-Me ₂)	Me	O	O
V-93	Ph(3,4-(OMe) ₂)	Me	O	O
V-94	Ph(3,5-(OMe) ₂)	Me	O	O
V-95	Ph(2,3-(OMe) ₂)	Me	O	O
V-96	Ph(2,4-(OMe) ₂)	Me	O	O
V-97	Ph(2,5-(OMe) ₂)	Me	O	O
V-98	Ph(2,6-(OMe) ₂)	Me	O	O
V-99	Ph(3-F-4-OMe)	Me	O	O
V-100	Ph(3-F-5-OMe)	Me	O	O
V-101	Ph(2-F-3-OMe)	Me	O	O
V-102	Ph(2-F-4-OMe)	Me	O	O
V-103	Ph(2-F-5-OMe)	Me	O	O
V-104	Ph(2-F-6-OMe)	Me	O	O
V-105	Ph(3-F-4-Me)	Me	O	O
V-106	Ph(3-F-5-Me)	Me	O	O
V-107	Ph(2-F-3-Me)	Me	O	O
V-108	Ph(2-F-4-Me)	Me	O	O
V-109	Ph(2-F-5-Me)	Me	O	O
V-110	Ph(2-F-6-Me)	Me	O	O
V-111	Ph(3-OMe-4-F)	Me	O	O
V-112	Ph(2-OMe-3-F)	Me	O	O
V-113	Ph(2-OMe-4-F)	Me	O	O
V-114	Ph(2-OMe-5-F)	Me	O	O
V-115	Ph(3-Me-4-F)	Me	O	O
V-116	Ph(2-Me-3-F)	Me	O	O
V-117	Ph(2-Me-4-F)	Me	O	O















[Table 59]

Compound No.	R ¹	R ²	Y	Z
V-118	Ph(2-Me-5-F)	Me	O	O
V-119	Ph(3-Cl-4-OMe)	Me	O	O
V-120	Ph(3-Cl-5-OMe)	Me	O	O
V-121	Ph(2-Cl-3-OMe)	Me	O	O
V-122	Ph(2-Cl-4-OMe)	Me	O	O
V-123	Ph(2-Cl-5-OMe)	Me	O	O
V-124	Ph(2-Cl-6-OMe)	Me	O	O
V-125	Ph(3-Cl-4-Me)	Me	O	O
V-126	Ph(3-Cl-5-Me)	Me	O	O
V-127	Ph(2-Cl-3-Me)	Me	O	O
V-128	Ph(2-Cl-4-Me)	Me	O	O
V-129	Ph(2-Cl-5-Me)	Me	O	O
V-130	Ph(2-Cl-6-Me)	Me	O	O
V-131	Ph(3-OMe-4-Cl)	Me	O	O
V-132	Ph(2-OMe-3-Cl)	Me	O	O
V-133	Ph(2-OMe-4-Cl)	Me	O	O
V-134	Ph(2-OMe-5-Cl)	Me	O	O
V-135	Ph(3-Me-4-Cl)	Me	O	O
V-136	Ph(2-Me-3-Cl)	Me	O	O
V-137	Ph(2-Me-4-Cl)	Me	O	O
V-138	Ph(2-Me-5-Cl)	Me	O	O
V-139	Ph(3-F-4-Cl)	Me	O	O
V-140	Ph(3-F-5-Cl)	Me	O	O
V-141	Ph(2-F-3-Cl)	Me	O	O
V-142	Ph(2-F-4-Cl)	Me	O	O
V-143	Ph(2-F-5-Cl)	Me	O	O
V-144	Ph(2-F-6-Cl)	Me	O	O
V-145	Ph(3-Cl-4-F)	Me	O	O
V-146	Ph(2-Cl-3-F)	Me	O	O
V-147	Ph(2-Cl-4-F)	Me	O	O
V-148	Ph(2-Cl-5-F)	Me	O	O
V-149	Ph(3-Me-4-OMe)	Me	O	O
V-150	Ph(3-Me-5-OMe)	Me	O	O
V-151	Ph(2-Me-3-OMe)	Me	O	O
V-152	Ph(2-Me-4-OMe)	Me	O	O
V-153	Ph(2-Me-5-OMe)	Me	O	O
V-154	Ph(2-Me-6-OMe)	Me	O	O
V-155	Ph(3-OMe-4-Me)	Me	O	O
V-156	Ph(2-OMe-3-Me)	Me	O	O













[Table 60]

Compound No.	R ¹	R ²	Y	Z
V-157	Ph(2-OMe-4-Me)	Me	O	O
V-158	Ph(2-OMe-5-Me)	Me	O	O
V-159	Ph(3-CN-4-OMe)	Me	O	O
V-160	Ph(3-OMe-4-CN)	Me	O	O
V-161	Ph(3-Me-4-CN)	Me	O	O
V-162	Ph(3-CN-4-Me)	Me	O	O
V-163	Ph(3-NO ₂ -4-OMe)	Me	O	O
V-164	Ph(3-OMe-4-NO ₂)	Me	O	O
V-165	Ph(3-Me-4-NO ₂)	Me	O	O
V-166	Ph(3-NO ₂ -4-Me)	Me	O	O
V-167	Ph(3,5-F ₂ -4-OMe)	Me	O	O
V-168	Ph(3,5-F ₂ -4-Me)	Me	O	O
V-169	Ph(3,4,5-(OMe) ₃)	Me	O	O
V-170		Me	O	O
V-171		Me	O	O
V-172		Me	O	O
V-173		Me	O	O
V-174		Me	O	O
V-175		Me	O	O
V-176		Me	O	O
V-177		Me	O	O
V-178		Me	O	O

[Table 61]

Compound No.	R ¹	R ²	Y	Z
V-179		Me	$\overset{\circ}{\text{O}}$	O
V-180		Me	O	O
V-181		Me	O	O
V-182		Me	O	O
V-183		Me	O	O
V-184		Me	O	O
V-185		Me	O	O
V-186		Me	O	O
V-187		Me	O	O
V-188		Me	O	O
V-189		Me	O	O
V-190		Me	O	O
V-191		Me	O	O
V-192		Me	O	O









[Table 62]

Compound No.	R ¹	R ²	Y	Z
V-193		Me	O	O
V-194		Me	O	O
V-195		Me	O	O
V-196		Me	O	O
V-197		Me	O	O
V-198		Me	O	O
V-199		Me	O	O
V-200		Me	O	O
V-201		Me	O	O
V-202		Me	O	O
V-203		Me	O	O
V-204		Me	O	O
V-205	CH ₂ Ph	Me	O	O
V-206	CH ₂ CH ₂ Ph	Me	O	O
V-207	CH ₂ CH ₂ CH ₂ Ph	Me	O	O
V-208	CH ₂ CH=CHPh	Me	O	O
V-209	CH ₂ C≡CPh	Me	O	O
V-210	CH ₂ CH=NOMe	Me	O	O










[Table 63]

Compound No.	R ¹	R ²	Y	Z
V-211	CH ₂ CH=NOEt	Me	O	O
V-212	CH ₂ CH=NOPr-n	Me	O	O
V-213	CH ₂ CH=NOPh	Me	O	O
V-214	CH ₂ CH(OMe) ₂	Me	O	O
V-215	CH ₂ CHO	Me	O	O
V-216	NH ₂	Me	O	O
V-217	NHMe	Me	O	O
V-218	NHEt	Me	O	O
V-219	NHPr-n	Me	O	O
V-220	NHPr-i	Me	O	O
V-221	NHBu-n	Me	O	O
V-222	NHBu-i	Me	O	O
V-223	NHBu-s	Me	O	O
V-224	NHCH ₂ Pr-c	Me	O	O
V-225	NHPen-n	Me	O	O
V-226	NHHex-n	Me	O	O
V-227	NHCH ₂ CH ₂ CH ₂ Cl	Me	O	O
V-228	NHCH ₂ CH ₂ CH ₂ F	Me	O	O
V-229	NHCH ₂ CH ₂ OMe	Me	O	O
V-230	NMe ₂	Me	O	O
V-231	NEt ₂	Me	O	O
V-232	N(Pr-n) ₂	Me	O	O
V-233	N(Bu-n) ₂	Me	O	O
V-234	N(Me)Et	Me	O	O
V-235	N(Me)CH ₂ CH ₂ OMe	Me	O	O
V-236	NHPh	Me	O	O
V-237	NHCH ₂ Ph	Me	O	O
V-238	N=CMe ₂	Me	O	O
V-239	N=CEt ₂	Me	O	O
V-240	N=CHNMe ₂	Me	O	O
V-241	NHC(=O)Me	Me	O	O
V-242	N[C(=O)Me] ₂	Me	O	O
V-243	NHC(=O)OMe	Me	O	O
V-244	N[C(=O)OMe] ₂	Me	O	O
V-245	NHSO ₂ Me	Me	O	O
V-246	NHSO ₂ Ph	Me	O	O
V-247	NHSO ₂ CH ₂ Ph	Me	O	O
V-248	OMe	Me	O	O
V-249	OEt	Me	O	O
V-250	OPr-n	Me	O	O
V-251	OPr-i	Me	O	O
V-252	OCH ₂ Pr-c	Me	O	O
V-253	OCH ₂ Cl	Me	O	O
V-254	OCHCl ₂	Me	O	O






[Table 64]

Compound No.	R ¹	R ²	Y	Z
V-255	OCCL ₃	Me	O	O
V-256	OCH ₂ F	Me	O	O
V-257	OCHF ₂	Me	O	O
V-258	OCF ₃	Me	O	O
V-259	Ph	Et	O	O
V-260	Ph	Pr-i	O	O
V-261	Ph	CHF ₂	O	O
V-262	Ph	Ph	O	O
V-263	Ph	Me	O	S
V-264	Ph	Me	S	S
V-265	Me	Me	O	S
V-266	Me	Me	S	S
V-267	Ph	Me	O	O
V-268	Ph(4-OEt)	Me	O	O
V-269	Ph(2-Ph)	Me	O	O
V-270	Ph(3-Ph)	Me	O	O
V-271	Ph(4-Ph)	Me	O	O
V-272		Me	O	O
V-273		Me	O	O
V-274	Me		O	O
V-275	Et		O	O
V-276		Me	O	O
V-277		Me	O	O
V-278		Me	O	O
V-279		Me	O	O








[Table 65]

Compound No.	R ¹	R ²	Y	Z
V-280		Me	O	O
V-281		Me	O	O
V-282	Ph(2-Me-4-Br)	Me	O	O
V-283	Ph(2-Me-5-I)	Me	O	O
V-284	Ph(2-Me-5-CF ₃)	Me	O	O
V-285	Ph(2-Me-6-OCF ₃)	Me	O	O
V-286	Ph(2-Pr-i)	Me	O	O
V-287		Me	O	O
V-288	Ph(2-Et)	Me	O	O
V-289		Me	O	O
V-290		Me	O	O
V-291		Me	O	S
V-292		Me	O	O
V-293		Me	O	O
V-294	CH ₂ COOBu ^t	Me	O	O
V-295	(C ₇ H ₁₄)CH ₃	Me	O	O
V-296	(C ₉ H ₁₈)CH ₃	Me	O	O
V-297	Ph(2-F,4-Cl,5-OMe)	Me	O	O
V-298	Ph(2,3,4-(OMe) ₃)	Me	O	O
V-299	Ph(3,5-Cl ₂ -4-OMe)	Me	O	O
V-300	Ph(3,5-Cl ₂ -4-SMe)	Me	O	O
V-301	Ph(3,5-Cl ₂ -4-SO ₂ Me)	Me	O	O
V-302	Ph(3,4,5-F ₃)	Me	O	O
V-303		Me	O	O

[Table 66]

Compound No.	R ¹	R ²	Y	Z
V-304		Me	O	O
V-305		Me	O	O
V-306	Bu-n		O	O
V-307	CH ₂ CH(CH ₃) ₂		O	O
V-308	Ph	Pen-n	O	O
V-309	H	Me	O	O
V-310	CH ₂ C≡CF	Me	O	O
V-311		Me	O	O
V-312		Me	O	O
V-313	CH ₂ NH ₂	Me	O	O
V-314	CH ₂ NO ₂	Me	O	O
V-315	CH ₂ NHCH ₃	Me	O	O
V-316	CH ₂ N(CH ₃) ₂	Me	O	O
V-317	CH ₂ SCH ₂ CF ₃	Me	O	O
V-318	CH ₂ SOCH ₂ CF ₃	Me	O	O
V-319	CH ₂ SO ₂ CH ₂ CF ₃	Me	O	O
V-320	CH ₂ OH	Me	O	O
V-321	CH ₂ OBn	Me	O	O
V-322	CH ₂ OCH ₂ Pr-c	Me	O	O
V-323	CH ₂ OPh	Me	O	O
V-324	CH ₂ SPh	Me	O	O
V-325	CH ₂ SOPh	Me	O	O
V-326	CH ₂ SO ₂ Ph	Me	O	O
V-327	CH ₂ CON(CH ₃) ₂	Me	O	O
V-328	CH ₂ COCH ₃	Me	O	O
V-329	CH ₂ OCOCH ₃	Me	O	O
V-330	CH ₂ ON=CHCH ₃	Me	O	O
V-331	C ₂ H ₄ OC ₂ H ₄ SCH ₃	Me	O	O
V-332	C ₂ H ₄ OC ₂ H ₄ SOCH ₃	Me	O	O
V-333	C ₂ H ₄ OC ₂ H ₄ SO ₂ CH ₃	Me	O	O

[Table 67]

Compound No.	R ¹	R ²	Y	Z
V-334	CH ₂ OCH ₂ CN	Me	O	O
V-335	CH ₂ CN	Me	O	O
V-336	OCH ₂ CH=CH ₂	Me	O	O
V-337	OCH ₂ C≡CH	Me	O	O
V-338	OPr-c	Me	O	O
V-339		Me	O	O
V-340		Me	O	O
V-341		Me	O	O
V-342		Me	O	O
V-343	CH ₂ CH ₂ OCH ₂ CH ₂ O- 	Me	O	O
V-344	Ph	H	O	O
V-345	Ph	CH ₂ CH=CH ₂	O	O
V-346	Ph	CH ₂ C≡CH	O	O
V-347	Ph	Pr-c	O	O
V-348	Ph	CH ₂ CH=CF ₂	O	O
V-349	Ph	CH ₂ C≡CF	O	O
V-350	Ph	C ₂ H ₄ OCH ₃	O	O
V-351	Ph	C ₂ H ₄ OC ₂ H ₅	O	O
V-352	Ph	CH(Me)OEt	O	O
V-353	Ph	CH ₂ OPr-c	O	O
V-354	Ph	CH(OCH ₃) ₂	O	O
V-355	Ph	CH ₂ Ph	O	O
V-356	Ph	CH=CH-Ph	O	O
V-357	Ph	C≡C-Ph	O	O
V-358	Ph(3,4,5-Cl)	Me	O	O
V-359	N(Me)Ph	Me	O	O
V-360		Me	O	O
V-361		Me	O	O
V-362	CH ₂ CO(Bu-t)	Me	O	O
V-363	Ph(2,3,5,6-F ₄)	Me	O	O
V-364	Ph[(3,5-(CF ₃) ₂)]	Me	O	O
V-365	CH ₂ C(Me)=NOMe	Me	O	O
V-366	Ph(2,4,6-Me ₃)	Me	O	O
V-367	Ph(2,3,4,5,6-F ₅)	Me	O	O
V-368	N(Et)Ph	Me	O	O
V-369	N(Pr-i)Ph	Me	O	O
V-370	N(Me)Ph(4-F)	Me	O	O
V-371	CH ₂ C(Me)=NOEt	Me	O	O

Compounds of the invention have an excellent herbicidal activity and some of them show

excellent selectivity between crops and weeds and are useful as an agrochemical composition for farmland, especially as herbicides. In other words, the compounds of the invention have a herbicidal activity for various weeds during foliage treatment, soil treatment, seed dressing treatment, soil blending treatment, soil treatment before sowing, treatment at the time of sowing, 5 soil treatment after sowing, soil covering and blending treatment at the time of sowing, and soil treatment before and after sowing for no-tillage farming of a field for cultivating agrohorticultural plants.

Hereinbelow, examples of the weeds are given, but the invention is not limited to them;

weeds of Onagraceae family: *Oenothera erythrosepala*, *Oenothera laciniata*;

10 weeds of Ranunculaceae family: *Ranunculus muricatus*, *Ranunculus sardous*;

weeds of Polygonaceae family: *Polygonum convolvulus*, *Polygonum lapathifolium*, *Polygonum pensylvanicum*, *Polygonum persicaria*, *Rumex crispus*, *Rumex obtusifolius*, *Poligonum cuspidatum*, *Polygonum pensylvanicum*, *Persicaria longiseta*, *Persicaria lapathifolia*, *Persicaria nepalensis*;

15 weeds of Portulacaceae family: *Portulaca oleracea*;

weeds of Caryophyllaceae family: *Stellaria media*, *Cerastium glomeratum*, *Stellaria alsine*, *Spergula arvensis*, *Stellaria aquatica*;

weeds of Chenopodiaceae family: *Chenopodium album*, *Kochia scoparia*, *Chenopodium album*, *Chenopodium ficifolium*;

20 weeds of Amaranthaceae family: *Amaranthus retroflexus*, *Amaranthus hybridus*, *Amaranthus palmeri*, *Amaranthus spinosus*, *Amaranthus rudis*, *Amaranthus albus*, *Amaranthus viridus*, *Amaranthus lividus*;

weeds of Brassicaceae family: *Raphanus raphanistrum*, *Sinapis arvensis*, *Capsella bursa-pastoris*, *Lepidium virginicum*, *Thlaspi arvense*, *Descurarinia sophia*, *Rorippa indica*, 25 *Rorippa islandica*, *Sisymnrium officinale*, *Cardamine flexuosa*, *Nasturtium officinale*, *Draba nemorosa*;

weeds of Fabaceae family: *Sesbania exaltata*, *Cassia obtusifolia*, *Desmodium tortuosum*, *Trifolium repens*, *Vicia sativa*, *Medicago lupulina*, *Vicia hirsuta*; *Kummerowia striata*, *Medicago polymorpha*, *Vicia angustifolia*, *Aeschynomene indica*;

weeds of Malvaceae family: *Abutilon theophrasti*, *Sida spinosa*;

weeds of Violet family: *Viola arvensis*, *Viola tricolor*;

weeds of Rubiaceae family: *Galium aparine*;

weeds of Convolvulaceae family: *Ipomoea hederacea*, *Ipomoea purpurea*, *Ipomoea*

5 *hederacea* var *integriuscula*, *Ipomoea lacunosa*, *Convolvulus arvensis*, *Ipomoea indica*, *Ipomoea coccinea*, *Ipomoea triloba*;

weeds of Lamiaceae family: *Lamium purpureum*, *Lamium amplexicaule*, *Stachys arvensis*;

weeds of Solanaceae family: *Datura stramonium*, *Solanum nigrum*, *Physalis angulata*,
Solanum americanum, *Solanum carolinense*;

10 weeds of Scrophulariaceae family: *Veronica persica*, *Veronica arvensis*, *Veronica hederifolia*;

weeds of Asteraceae family: *Xanthium pensylvanicum*, *Helianthus annuus*, *Matricaria chamomilla*, *Matricaria perforata* or *inodora*, *Chrysanthemum segetum*, *Matricaria matricarioides*,
Ambrosia artemisiifolia, *Ambrosia trifida*, *Erigeron canadensis*, *Artemisia princeps*, *Solidago*
15 *altissima*, *Taraxacum officinale*, *Anthemis cotula*, *Brexa setosa*, *Sonchus oleraceus*, *Helianthus tuberosus*, *Cirsium arvense*, *Bidens frondosa*, *Bidens pilosa*, *Centurea cyanus*, *Cirsium vulgare*,
Lactuca scariola, *Rudbeckia hirta*, *Rudbeckia laciniata*, *Rudbeckia laciniata* var. *hortensis* Bailey,
Senecio vulgaris, *Silybum marianum*, *Sonchus asper*, *Sonchus arvensis*, *Salsola kali*, *Bidens*
frondosa, *Eclipta prostrata*, *Bidens bipartita*, *Senecio madagascariensis*, *Coreopsis lanceolata*,

20 *Rudbeckia laciniata*;

weeds of Boraginaceae family: *Myosotis arvensis*;

weeds of Asclepiadaceae family: *Asclepias syriaca*;

weeds of Euphorbiaceae family: *Euphorbia helioscopia*, *Euphorbia maculata*, *Acalypha*
australis;

25 weeds of Geraniaceae family: *Geranium carolinianum*;

weeds of Oxalidaceae family: *Oxalis corymbosa*;

weeds of Cucurbitaceae family: *Sicyos angulatus*;

weeds of Poaceae family: *Echinochloa crus-galli*, *Setaria viridis*, *Setaria faberi*, *Digitaria sanguinalis*, *Eleusine indica*, *Poa annua*, *Alopecurus myosuroides*, *Avena fatua*, *Sorghum*

halepense, Agropyron repens, Bromus tectorum, Cynodone dactylon, Panicum dichotomiflorum, Panicum texanum, Sorghum vulgare, Alopecurus geniculatus, Lolium multiflorum, Lolium rigidum, Setaria glauca, Beckmannia syzigachne;

weeds of Commelinaceae family: Commelina communis;

5 weeds of Equisetaceae family: Equisetum arvense;

weeds of Papaveraceae family: Papaver rhoeas;

weeds of Cyperaceae family: Cyperus iria, Cyperus rotundus, Cyperus esculentus.

Compounds of the invention do not exhibit any phytotoxicity which causes a problem in major crops like Zea mays, Triticum aestivum, Hordeum vulgare, Oryza sativa, Sorghum bicolor, 10 Glycine max, Gossypium spp., Beta vulgaris, Arachis hypogaea, Helianthus annuus, Brassica napus, buck wheat, sugar cane, and tobacco, and horticultural crops like flowers and vegetables.

Further, the compounds of the invention are useful for effective elimination of various weeds which cause a trouble in no-tillage farming of soybean, corn, and wheat, and they do not exhibit any problematic phytotoxicity to crops.

15 According to many treatment methods like soil treatment before cultivation; soil treatment after cultivation but before or after sowing; soil treatment after harrowing but before or after sowing, or treatment before or after transplanting a seedling; treatment at the time of transplanting a seedling; desalination treatment after transplanting a seedling; and foliage treatment, the compounds of the invention can exhibit an herbicidal activity for many problematic weeds in 20 paddy field that are described below.

Hereinbelow, examples of the weeds are given, but the invention is not limited to them:

weeds of Poaceae family: Echinochloa oryzicola; Echinochloa crus-galli, Leptochloa chinensis, Isachne globosa, Paspalum distichum, Leersia sayanuka, Leersia oryzoides;

weeds of Scrophulariaceae family: Lindernia procumbens, Lindernia dubia, Dopatrium 25 junceum, Gratiola japonica, Lindernia angustifolia, Limnophila sessiliflora;

weeds of Lythraceae family: Rotala indica, Ammannia multiflora;

weeds of Elatinaceae family: Elatine triandra;

weeds of Cyperaceae family: Cyperus difformis, Scirpus hotarui, Eleocharis acicularis, Cyperus serotinus, Eleocharis kuroguwai, Fimbristylis miliacea, Cyperus flaccidus, Cyperus

globosus, *Scirpus juncoides*, *Scirpus wallichii*, *Scirpus nipponicus*, *Fimbristylis autumnalis*, *Scirpus tabernaemontani*, *Scirpus juncoides* Roxb., *Scirpus lineolatus* Franch. et Savat., *Cyperus orthostachyus* Franch. et Savat., *Cyperus orthostachyus* Franch. et Savat., *Eleocharis congesta* D. Don, *Scirpus planiculmis* Fr. Schm.;

5 weeds of Pontederiaceae family: *Monochoria vaginalis*, *Monochoria korsakowii*, *Heteranthera limosa*;

weeds of Alismataceae family: *Sagittaria pygmaea*, *Sagittaria trifolia*, *Alisma canaliculatum*, *Sagittaria aginashi*;

weeds of Potamogetonaceae family: *Potamogeton distinctus*;

10 weeds of Eruocaulaceae family: *Eriocaulon cinereum*;

weeds of Apiaceae family: *Oenanthe javanica*;

weeds of Asteraceae family: *Eclipta prostrata*, *Bidens tripartita*;

weeds of Commelinaceae family: *Murdannia keisak*;

weeds of Characeae family: *Chara braunii*;

15 weeds of Lemnaceae family: *Spirodela polyrhiza*;

Hepaticae: *Ricciocarpus natans*;

Zygnemataceae: *Spirogyra arcla*.

Further, the compounds of the invention show no phytotoxicity to paddy rice according to any cultivation method including direct sowing or transplanting of paddy rice followed by cultivation.

20 Further, the compounds of the invention can be used for controlling a wide spectrum of weeds thriving in a lot for industrial facilities like a slope of a levee, a riverbed, a shoulder and a slope of a road, a railway site, park spaces, grand, a parking lot, an airport, a factory and a storage facility, a non-crop land like fallow fields, and vacant lots in city, which needs the weed control, or an orchard, a pasture land, a grass land, a forest land, etc.

25 Moreover, according to foliage treatment, water-surface application, etc., the compounds of the invention can exhibit a herbicidal activity for water weeds which occur in river, waterway, canal, reservoir, etc., wherein the water weeds include Pontederiaceae family: *Eichhornia crassipes*; *Salvinia natans* family: *Azolla imbricata*, *Azolla japonica*, *Salvinia natans*; Araceae family: *Pistia stratiotes*; Haloragaceae family: *Myriophyllum brasiliense*, *Myriophyllum*

verticillatum; Myriophyllum spicatum; Myriophyllum matogrossense; Azollaceae family: Azolla cristata; Scrophulariaceae family: Veronica anagallis-aquatica; Amaranthaceae family: Alternanthera philoxeroides; Gymnocoronis spilanthoides; Poaceae family: Spartina anglica; Apiaceae family: Hydrocotyle ranunculoides; Hydrocharitaceae family: Hydrilla verticillata, 5 Egeria densa; Cabombaceae family: Cabomba caroliniana; and Lemnaceae family: Wolffia globosa.

The agrohorticultural plants described in the invention include crops like corn, rice, wheat, barley, rye, sorghum, cotton, soybean, peanuts, buck wheat, sugar beet, rapeseed, sun flower, sugar cane, and tobacco; vegetable like vegetables of Solanaceae (eggplant, tomato, bell pepper, 10 pepper, potato, etc.), vegetables of Cucurbitaceae (cucumber, pumpkin, zucchini, water melon, melon, etc.), vegetables of Cruciferae (daikon, turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, mustard, broccoli, cauliflower, etc.), vegetables of Compositae (burdock, crown daisy, artichoke, lettuce, etc.), vegetables of Liliaceae (scallion, onion, garlic, asparagus, etc.), vegetables of Apiaceae (carrot, parsley, celery, parsnip, etc.), vegetables of Chenopodiaceae 15 (spinach, leaf beet, etc.), vegetables of Lamiaceae (beefsteak plant, mint, basil, etc.), vegetables like strawberry, sweet potato, yam, and taro; kernel fruits (apple, western pear, Japanese pear, Chinese quince, quince, etc.), stone fruits (peach, plum, nectarine, Japanese apricot, cherry, apricot, prune, etc.), mandarins (tangerine, orange, lemon, lime, grape fruits, etc.), nuts (chestnut, walnut, hazelnut, almond, pistachio, cashewnut, macadamia nut, etc.), berries (blueberry, 20 cranberry, blackberry, raspberry, etc.), fruits like grape, persimmon, olive, loquat, banana, coffee, date, coconut, and oil nut; trees other than fruit tree like tea, mulberry, roadside trees (ash tree, birch, American flowering dogwood, eucalyptus, ginkgo, lilac, maple tree, oak tree, poplar tree, redbud tree, liquidambar, sycamore, zelkova, Japanese arborvitae, Japanese fir, hemlock spruce, juniper, pine tree, spruce, yew, elm, a horse chestnut, etc.), coral, Buddhist pine, cedar, Japanese 25 cypress, croton, spindle tree, Photinia glabra, etc.; grasses like turf (turf, gold turf, etc.), Bermuda grasses (Cynodon dactylon, etc.), bentgrasses (creeping bentgrass, Agrostis alba L., Agrostis capillaries, etc.), bluegrasses (Kentucky bluegrass, Poa trivialis L., etc.), fescues (tall fescue, chewings fescue, Festuca rubra L., etc.), rye grasses (Lolium temulentum L., Lolium perenne L., etc.), orchard grass, timothy, etc.; oil crops like oil coconut, Jatropha curcas, etc.; flowers (rose,

carnation, mum, prairie gentian, common gypsophila, gerbera, marigold, salvia, petunia, verbena, tulip, Chinese aster, *Gentiana scabra* var. *buergeri*, lily, pansy, cyclamen, orchid, lily of the valley, lavender, stock, cauliflower, primula, poincetia, gladiolus, cattleya, daisy, verbena, cymbidium, begonia, etc.); a foliage plant, etc., but the invention is not limited thereto.

5 The agrohorticultural plant described in the invention includes a plant given with resistance to HPPD inhibitor like Isoxaflutole, ALS inhibitor like Imazetaphyr and tifensulfuron methyl, EPSP synthase inhibitor like glifosate, glutamine synthase inhibitor like glufosinate, acetyl CoA carboxylase inhibitor like sethoxydim, PPO inhibitor like flumioxazin, and herbicides like bromoxinil, dicamba and 2,4-D according to classic breeding method or genetic recombination
10 method.

Examples of the "agrohorticultural plant" given with resistance according to classic breeding include rapeseed, wheat, sun flower, rice, and corn that are resistant to imidazolinone-based ALS inhibitor like Imazetaphyr, and they are already commercially available in the name of Clearfield
<trade name>.

15 Similarly, there is soybean resistant to sulfonyleurea-based ALS inhibitor like tifensulfuron metil as produced by classic breeding method, and it is already commercially available in the trade name of STS Soybean. Similarly, examples of the "agrohorticultural plant" given with resistance to an acetyl CoA carboxylase inhibitor like trione-oxime based or aryloxyphenoxy propionic acid-based herbicides according to classic breeding include SR Corn. The
20 horticultural plant given with resistance to acetyl CoA carboxylase is described in Proceedings of the National Academy of Sciences of the United States of America (Proc. Natl. Acad. Sci. USA), Vol 87, pages 7175 to 7179 (1990), etc. Further, a mutant acetyl CoA carboxylase which is resistant to acetyl CoA carboxylase inhibitor is reported in Weed Science Vol. 53, pages 728 to 746 (2005), and by introducing a mutant gene of acetyl CoA carboxylase to a plant by genetic
25 recombination technique or by introducing mutation for giving resistance to acetyl CoA carboxylase of crops, a plant which is resistant to an acetyl CoA carboxylase inhibitor can be produced. Further, by having site-specific amino acid substitution mutation on a gene of crops based on introduction of a nucleic acid with base substitution mutation to a plant cell as represented by chimera-plasty technique (Gura T. 1999. Repairing the Genome's Spelling

Mistakes. Science 285: 316-318), a plant which is resistant to acetyl CoA carboxylase inhibitor/herbicides can be produced.

Examples of the "agrohorticultural plant" given with resistance according to genetic recombination technique include corn, soybean, cotton, rapeseed, and sugar beet that are resistant to glyphosate, and they are already commercially available in the name of RoundupReady <trade name>, AgrisureGT, etc. Similarly, there are corn, soybean, cotton, and rapeseed varieties that are produced to be resistant to glufosinate by genetic recombination technique, and they are already commercially available in the name of LibertyLink <trade name>, etc. Similarly, cotton having resistance to bromoxinil is also made available by genetic recombination technique and is already commercially available in the trade name of BXN.

The "agrohorticultural plant" includes a plant which is engineered by genetic recombination technique to synthesize a selective toxin like *Bacillus* spp., for example.

Examples of the insecticidal toxin expressed in a genetically engineered plant include an insecticidal protein originating from *Bacillus cereus* or *Bacillus popilliae*; δ -endotoxin originating from *Bacillus thuringiensis* like Cry1Ab, Cry1Ac, Cry1F, Cry1Fa2, Cry2Ab, Cry3A, Cry3Bb1, and Cry9C, and insecticidal proteins like VIP1, VIP2, VIP3, and VIP3A; insecticidal proteins originating from a nematode; animal-produced toxins like scorpion toxin, spider toxin, bee toxin, and insect specific neurotoxin; filamentous fungus toxin; plant lectin; agglutinin; protease like trypsin inhibitor, serine protease, patatin, cistatin, and papain inhibitor; ribosome inactivating proteins (RIP) like lysine, corn-RIP, abrin, saporin, and briodin; enzymes for steroid metabolism like 3-hydroxysteroid oxidase, ecdysteroid-UDP-glycosyl transferase, and cholesterol oxidase; ecdysone inhibitor; HMG-CoA reductase; ion channel inhibitors like sodium channel inhibitor and potassium channel inhibitor; juvenile hormone esterase; natriuretic hormone receptor; stilbene synthase; bibenzyl synthase; chitinase; and glucanase.

Examples of the toxins expressed in a genetically engineered plant include a hybrid toxin, a partially deleted toxin, and a modified toxin of an insecticidal protein like δ -endotoxin including Cry1Ab, Cry1Ac, Cry1F, Cry1Fa2, Cry2Ab, Cry3A, Cry3Bb1, and Cry9C, and insecticidal proteins including VIP1, VIP2, VIP3, and VIP3A. The hybrid toxin is produced by new combination of domains having different proteins based on recombination technique. Examples

of the partially deleted toxin include Cry1Ab in which part of amino acid sequence is deleted. In the modified toxin, one or more amino acids of a natural type toxin are replaced with other amino acids.

Examples of the toxins and recombinant plants capable of producing the toxins are described
5 in EP-A-0374753, WO93/07278, WO95/34656, EP-A-0427529, EP-A-451878, and WO03/052073, and the like.

The toxins contained in such recombinant plant can provide a plant with a resistance to harmful insects of Coleoptera, harmful insects of Diptera, and harmful insects of Lepidoptera.

A genetically engineered plant which contains one or more pesticidal harmful
10 insect-resistant gene and expresses one or more toxins is known, and some are already commercially available. Examples of the genetically engineered plant include YieldGard <trade name> (corn variety which expresses Cry1Ab toxin), YieldGard Rootworm <trade name> (corn variety which expresses Cry3Bb1 toxin), YieldGard Plus <trade name> (corn variety which expresses Cry1Ab and Cry3Bb1 toxin), Herculex I <trade name> (corn variety which expresses phosphinotricine N-acetyl transferase (PAT) to give resistance to Cry1Fa2 toxin and glufosinate),
15 NuCOTN33B <trade name> (cotton variety which expresses Cry1Ac toxin), Bollgard I <trade name> (cotton variety which expresses Cry1Ac toxin), Bollgard II <trade name> (cotton variety which expresses Cry1Ac and Cry2Ab toxin), VIPCOT <trade name> (cotton variety which expresses VIP toxin), NewLeaf <trade name> (potato variety which expresses Cry3A toxin),
20 NatureGard <trade name> Agrisure <trade name> GT Advantage (GA21 glyphosate resistant trait), Agrisure <trade name> CB Advantage (Bt11 Corn Borer (CB) trait), and Protecta <trade name>.

The "agrohorticultural plant" includes a plant which is genetically engineered to have an ability of producing an anti-pathogenic substance having selective activity.

Examples of the anti-pathogenic substance include PR proteins (PRPs, described in
25 EP-A-0392225); ion channel inhibitors like sodium channel inhibitor and calcium channel inhibitor (KP1, KP4, KP6 toxin that are produced by virus are known); stilbene synthase; bibenzyl synthase; chitinase; glucanase; and a substance produced by a microorganism like peptide antibiotics, antibiotics having heterocycle, and a protein factor related to resistance to plant disease (referred to as "plant disease resistant gene", and described in WO03/000906).

Such anti-pathogenic substances and plants genetically engineered to produce the substances are described in EP-A-0392225, WO95/33818, and EP-A-0353191, etc.

The "agrohorticultural plant" includes a plant which is given with useful traits like a trait of having modified oil components or a trait for producing enhanced amount of amino acid
5 according to genetic recombination technique. Examples thereof include VISTIVE <trade name> (low-linolenic soybean having reduced linolen content) or high-lysine (high oil) corn (corn having enhanced amount of lysine or oil).

Further, there is also a stack variety in which multiple traits of classic herbicidal trait or herbicides-resistant gene, pesticidal insect-resistant gene, anti-pathogenic substance-producing
10 gene, and useful traits like a trait of having modified oil components or a trait for producing enhanced amount of amino acid are combined.

The agrochemical composition of the invention contains the triazine derivative of the invention or a salt thereof, and an agriculturally acceptable carrier. The agrochemical composition of the invention may contain additive components that may be normally employed
15 for agrochemical formulations, as needed.

Examples of the additive components include carriers such as solid carrier and liquid carrier, surface active agent, binder, tackifier, thickener, coloring agent, spreader, sticker, antifreezing agent, anticaking agent, collapsing agent, decomposition inhibitor and the like.

If necessary, an antiseptic agent, a piece of plant (soybean powder, tobacco powder, walnut
20 powder, wheat powder, wood powder, hulls, wheat hulls, outer hulls, sawdust, pulp flock, corn stalk, nut shell, fruit core chips, etc.) and the like may also be employed as additive components.

These additive components may be used alone or in combination of two or more kinds.

The above additive components will be described.

Examples of the solid carrier include natural minerals such as quartz, clay, kaolinite,
25 pyrophyllite, sericite, talc, bentonite, acid clay, attapulgite, zeolite and diatomite; inorganic salts such as calcium carbonate, ammonium sulfate, sodium sulfate and potassium chloride; organic solid carriers such as synthetic silicic acid, synthetic silicate, starch, cellulose and plant powder; plastic carriers such as polyethylene, polypropylene and polyvinylidene chloride; and the like. These may be used alone or in combination of two or more kinds.

Examples of the liquid carrier include alcohols classified broadly into monohydric alcohols such as methanol, ethanol, propanol, isopropanol and butanol, and polyhydric alcohols such as ethylene glycol, diethylene glycol, propylene glycol, hexylene glycol, polyethylene glycol, polypropylene glycol and glycerin; polyhydric alcohol derivatives such as propylene based glycol ether; ketones such as acetone, methylethyl ketone, methylisobutyl ketone, diisobutyl ketone, and cyclohexanone; ethers such as ethyl ether, dioxane, cellosolve, dipropyl ether and tetrahydrofuran; aliphatic hydrocarbons such as n-paraffin, naphthene, isoparaffin, kerosene and mineral oil; aromatic hydrocarbons such as benzene, toluene, xylene, solvent naphtha and alkylnaphthalene; halogenated hydrocarbons such as dichloroethane, chloroform and carbon tetrachloride; esters such as ethyl acetate, diisopropyl phthalate, dibutyl phthalate, dioctyl phthalate and dimethyl adipate; lactones such as γ -butyrolactone; amides such as N,N-dimethylformamide, N,N-diethylformamide, N,N-dimethylacetamide and N-alkyl pyrrolidinone; nitriles such as acetonitrile; sulfur compounds such as dimethylsulfoxide; vegetable oils such as soybean oil, canola oil, cottonseed oil and castor oil; water; and the like.

These may be use alone or in combination of two or more kinds.

The surface active agent is not particularly limited, but preferred are those either turning into a gel in water or exhibiting swelling property. Examples thereof include non-ionic surface active agents such as sorbitan fatty acid ester, polyoxyethylene sorbitan fatty acid ester, sucrose fatty acid ester, polyoxyethylene fatty acid ester, polyoxyethylene resin acid ester, polyoxyethylene fatty acid diester, polyoxyethylene alkyl ether, polyoxyethylene alkylphenyl ether, polyoxyethylene dialkylphenyl ether, polyoxyethylene alkylphenyl ether formaldehyde condensate, polyoxyethylene polyoxypropylene block polymer, alkylpolyoxyethylene polypropylene block polymer ether, polyoxyethylene alkylamine, polyoxyethylene fatty acid amide, polyoxyethylene fatty acid bisphenyl ether, polyalkylene benzylphenyl ether, polyoxyalkylene styrylphenyl ether, acetylene diol, polyoxyalkylene-added acetylene diol, polyoxyethylene ether silicone, ester silicone, fluorine-based surface active agent, polyoxyethylene castor oil, and polyoxyethylene hydrogenated castor oil; anionic surface active agents such as alkyl sulfate, polyoxyethylene alkyl ether sulfate, polyoxyethylene alkylphenyl ether sulfate, polyoxyethylene styrylphenyl ether sulfate, alkyl benzene sulfonate, lignin sulfonate,

alkyl sulfosuccinate, naphthalene sulfonate, alkyl naphthalene sulfonate, naphthalenesulfonic acid formaldehyde condensate salt, alkyl naphthalenesulfonic acid formaldehyde condensate salt, fatty acid salt, polycarboxylate, N-methyl-fatty acid sarcosinate, resin acid salt, polyoxyethylene alkyl ether phosphate, and polyoxyethylene alkylphenyl ether phosphate; cationic surface active agents
5 such as laurylamine hydrochloride, stearylamine hydrochloride, oleylamine hydrochloride, stearylamine acetate, stearylaminopropylamine acetate, alkyltrimethylammonium chloride, and alkyl dimethylbenzalkonium chloride; amino acid or betaine type amphoteric surface active agents; and the like.

These surface active agents may be used alone or in combination of two or more kinds.

10 Examples of the binder or tackifier include carboxymethyl cellulose and a salt thereof, dextrin, water-soluble starch, xanthan gum, guar gum, sucrose, polyvinylpyrrolidone, gum arabic, polyvinyl alcohol, polyvinyl acetate, sodium polyacrylate, polyethylene glycol having an average molecular weight of 6,000 to 20,000, polyethylene oxide having an average molecular weight of 100,000 to 5,000,000 natural phospholipids (for instance, cephalic acid, lecithin) and the like.

15 Examples of the thickener include water-soluble polymers such as xanthan gum, guar gum, carboxymethyl cellulose, polyvinylpyrrolidone, carboxy vinyl polymer, acrylic polymer, starch derivative and polysaccharide; fine inorganic powders such as high purity bentonite and white carbon; and the like.

20 Examples of the coloring agent include inorganic pigments such as iron oxide, titanium oxide and Prussian blue; organic dyes such as alizarin dye, azo dye and metal phthalocyanine dye; and the like.

Examples of the extender agent include silicone surface active agent, cellulose powder, dextrin, processed starch, polyaminocarboxylic acid chelate compound, crosslinked polyvinylpyrrolidone, maleic acid-styrenes-methacrylic acid copolymer, half ester of polyhydric
25 alcohol polymer with dicarboxylic anhydride, water-soluble salt of polystyrene sulfonate and the like.

Examples of the spreader include various surface active agents such as dialkyl sodium sulfosuccinate, polyoxyethylene alkyl ether, polyoxyethylene alkylphenyl ether and polyoxyethylene fatty acid ester, paraffin, terpene, polyamide resin, polyacrylate,

polyoxyethylene, wax, polyvinylalkyl ether, alkylphenol-formaldehyde condensate, synthetic resin emulsion and the like.

Examples of the antifreezing agent include polyhydric alcohols such as ethylene glycol, diethylene glycol, propylene glycol, glycerin, and the like.

5 Examples of the anticaking agent include polysaccharides such as starch, alginic acid, mannose and galactose, polyvinylpyrrolidone, white carbon, ester gum, petroleum resin and the like.

Examples of the collapsing agent include sodium tripolyphosphate, sodium hexametaphosphate, metal stearate, cellulose powder, dextrin, copolymer of methacrylic acid
10 ester, polyvinylpyrrolidone, polyaminocarboxylic chelate compound, sulfonated styrene-isobutylene-maleic anhydride copolymer, starch-polyacrylonitrile graft copolymer and the like.

Examples of the decomposition inhibitor include desiccants such as zeolite, quicklime and magnesium oxide; antioxidants that are based on phenol, amine, sulfur and phosphoric acid;
15 ultraviolet absorbers that are based on salicylic acid, benzophenone or the like; and the like.

Examples of the antiseptic agent include potassium sorbate, 1,2-benzthiazolin-3-one and the like.

According to the agrochemical composition of the invention, in the case where the additive components described above are included, the content ratio of the carrier (weight base) is
20 generally selected from 5 to 95%, preferably from 20 to 90%, the content ratio of the surface active agent is generally selected from 0.1 to 30%, preferably from 0.5 to 10%, and the content ratio of other additives are selected from 0.1 to 30%, preferably from 0.5 to 10%.

The agrochemical composition of the invention can be used in any forms such as liquid formulation, emulsifiable concentrate, wettable powder, dust, oil solution, water dispersible
25 granule, flowable, granule, Jumbo formulation, and suspoemulsion.

On the occasion of use, the agrochemical composition can be sprayed after being diluted in an adequate concentration or be used directly.

The agrochemical composition of the invention can be used for foliage application, soil application, water-surface application or the like. The agrochemical composition of the

invention, in particular the herbicides, is used for soils, i.e., farmland of fields and paddy fields in which agrohorticultural plants are cultivated.

For the agrochemical composition of the invention, the blending ratio of active component according to the invention is arbitrarily selected as needed. In the case of dust, granule or the like, the ratio should be arbitrarily selected from 0.01 to 10% (by weight), preferably from 0.05 to 5% (by weight). In the case of emulsifiable concentrate, wettable powder or the like, the ratio should be arbitrarily selected from 1 to 50% (by weight), preferably from 5 to 30% (by weight). In addition, in the case of a flowable agent or the like, the ratio should be arbitrarily selected from 1 to 40% (by weight), preferably from 5 to 30% (by weight).

The application amount of the agrochemical composition according to the invention varies depending on a kind of a compound to be used, target weed, growth pattern, environmental conditions, formulation for use or the like. In the case of a direct use of dust, granule or the like, the amount should be arbitrarily selected from 1 g to 50 kg, preferably from 10 g to 10 kg per hectare as an active component. Further, in the case of using in a liquid form, for example, in the case of emulsifiable concentrate, wettable powder, flowable agent or the like, the amount should be arbitrarily selected from 0.1 to 50,000 ppm, preferably from 10 to 10,000 ppm.

The agrochemical composition of the invention has an excellent herbicidal activity, and therefore is useful as herbicides in particular.

According to purpose of use, the agrochemical composition of the invention may be formulated, mixed or used in combination with at least one additional agrochemically active component, for example, a plant disease control component, a pesticidal component, an acaricidal component, an nematocidal component, a synergistic agent component, an attracting component, a repellent component, a herbicidal component, a safener component, a microbial pesticidal component, a plant growth control component, a fertilizer component, a soil improving agent, etc.

When the composition is used in combination with other agrochemically active component or fertilizer, the preparation of each individual component may be mixed with others at the time of use. Further, each preparation of an individual component may be used in order, or used with an interval of some days. When the preparations are used with an interval of some days, they may be applied with an interval of 1 day to 40 days, for example, although it may vary depending

on each component to be used.

According to the agrochemical composition of the invention, when a mixture of at least one compound selected from the triazine derivatives represented by Formula 1 and their salt and at least one kind selected from other agrochemically active components is used, they are generally
5 used in weight ratio of 100 : 1 to 1 : 100, preferably 20 : 1 to 1 : 20, and particularly 10 : 1 to 1 : 10.

Among other agrochemically active components that may be mixed or used in combination with the compound of the invention in the agrochemical composition of the invention, examples of known herbicides or plant growth control agents are described below, but the invention is not
10 limited thereto.

[Herbicides]

A1. Acetyl CoA carboxylase (ACCase) inhibition type herbicides

(A1-1) Aryl oxy phenoxy propionic acid-based compound: clodinafop-propargyl, cyhalofop-butyl, diclofop-methyl, diclofop-P-methyl, fenoxaprop-P-ethyl, fluazifop-butyl,
15 fluazifop-P-butyl, haloxyfop, haloxyfop-etotyl, haloxyfop-P, metamifop, propaquizafop, quizalofop-ethyl, quizalofop-P-ethyl, quizalofop-P-tefuryl, and fenthiaprop-ethyl

(A1-2) Cyclohexane dione-based compound: alloxymid, butoxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxymid, and tralkoxydim

(A1-3) Phenyl pyrazoline-based compound: aminopyralid, and pinoxaden

20 B. Acetolactate synthase (ALS) inhibition type herbicides

(B-1) Imidazolinone-based compound: imazamethabenz-methyl, imazamox, imazapic (including salts with amine or the like), imazapyr (including salts with isopropylamine or the like), imazaquin, and imazethapyr

(B-2) Pyrimidinyloxy benzoic acid-based compound: bispyribac-sodium, pyribenzoxim, pyriftalid, pyriminobac-methyl, pyriothiobac-sodium, and pyrimisulfan
25

(B-3) Sulfonylamino carbonyl triazolinone-based compound: flucarbazone-sodium, thienicarbazone (including sodium salt, methyl ester, or the like), propoxycarbazone-sodium, procarbazone-sodium

(B-4) Sulfonylurea-based compound: amidosulfuron, azimsulfuron, bensulfuron-methyl,

chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, flupyrsulfuron-methyl-sodium, foramsulfuron, halosulfuron-methyl, imazosulfuron, iodosulfuron-methyl-sodium, mesosulfuron-methyl, metsulfuron-methyl, nicosulfuron, oxasulfuron, primisulfuron-methyl, prosulfuron, pyrazosulfuron-ethyl, rimsulfuron, sulfometuron-methyl, sulfosulfuron, thifensulfuron-methyl, triasulfuron, tribenuron-methyl, trifloxysulfuron-sodium, triflusulfuron-methyl, tritosulfuron, orthosulfamuron, propyrisulfuron, metazosulfuron, and flucetosulfuron

(B-5) Triazolopyrimidine-based compound: cloransulam-methyl, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, pyroxsulam, and HNPC-C-9908 (code number)

10 C1. Herbicides 1 for photosystem II photosynthesis inhibition

(C1-1) Phenylcarbamate-based compound: desmedipham and phenmedipham

(C1-2) Pyridazinone-based compound: chloridazon and brompyrazon

(C1-3) Triazine-based compound: ametryn, atrazine, cyanazine, desmetryne, dimethametryn, eglazine-ethyl, prometon, prometryn, propazine, simazine, simetryn, terbumeton, terbuthylazine, terbutryn, and trietazine

(C1-4) Triazinone-based compound: metamitron and metribuzin

(C1-5) Triazolinone-based compound: amicarbazone

(C1-6) Uracil-based compound: bromacil, lenacil, and terbacil

C2. Herbicides 2 for photosystem II photosynthesis inhibition

20 (C2-1) Amide-based compound: pentanochlor and propanil

(C2-2) Urea-based compound: chlorbromuron, chlorotoluron, chloroxuron, dimefuron, diuron, ethidimuron, fenuron, fluometuron, isoproturon, isouron, linuron, methabenzthiazuron, metobromuron, metoxuron, monolinuron, neburon, siduron, tebuthiuron, and metobenzuron

C3. Herbicides 3 for photosystem II photosynthesis inhibition

25 (C3-1) Benzothiadiazone-based compound: bentazone

(C3-2) Nitrile-based compound: bromofenoxim, bromoxynil (including ester form with butyric acid, octanoic acid and heptanoic acid), and ioxynil

(C3-3) Phenyl pyrazine-based herbicide compound: pyridafol, and pyridate

D. Photosystem I radical generation type herbicides

(D-1) Bipyridinium-based compound: diquat and paraquat dichloride

E. Protoporphyrinogen oxidase (PPO) inhibition herbicides

(E-1) Diphenyl ether-based compound: acifluorfen-sodium, bifenox, chlomethoxyfen, ethoxyfen-ethyl, fluoroglycofen-ethyl, fomesafen, lactofen, and oxyfluorfen

5 (E-2) N-Phenylphthalimide-based compound: cinidon-ethyl, flumiclorac-pentyl, flumioxazin, and chlorphthalim

(E-3) Oxy diazole-based compound: oxadiargyl and oxadiazon

(E-4) Oxazolidinedione-based compound: pentoxazone

(E-5) Phenylpyrazole-based compound: fluazolate and pyraflufen-ethyl

10 (E-6) Pyrimidinedione-based compound: benzfendizone, butafenacil, and saflufenacil

(E-7) Thiadiazole-based compound: fluthiacet-methyl and thidiazimin

(E-8) Triazolinone-based compound: azafenidin, carfentrazone-ethyl, sulfentrazone, and bencarbazone

(E-9) Other compounds: flufenpyr-ethyl, profluazol, pyraclonil, SYP-298 (code number),
15 and SYP-300 (code number)

F1. Phytoene desaturase (PDS) inhibition herbicides

(F1-1) Pyridazinone-based compound: norflurazon

(F1-2) Pyrimidine carboxamide-based compound: diflufenican and picolinafen

(F1-3) Other compounds: beflubutamid, fluridone, flurochloridone, and flurtamone

20 F2. 4-Hydroxyphenylpyruvate deoxygenase (HPPD) inhibition herbicides

(F2-1) Callistemon-based compound: mesotrione

(F2-2) Isoxazole-based compound: pyrasulfotole, isoxaflutole, and isoxachlortole

(F2-3) Pyrazole-based compound: benzofenap, pyrazolynate, and pyrazoxyfen

(F2-4) Triketone-based compound: sulcotrione, tefuryltrion, tembotrione, pyrasulfotole,
25 topramezone, bicyclopyrone, and 4-chloro-5-(1,3-dioxocyclohexa-2-yl)
carbonyl-2,3-dihydrobenzothiophene-1,1-dioxide

F3. Carotenoid biosynthesis inhibition (unknown target) herbicides

(F3-1) Diphenyl ether-based compound: aclonifen

(F3-2) Isoxazolidinone-based compound: clomazone

(F3-3) Triazole-based compound: amitrole

G. EPSP synthase synthesis inhibition (aromatic amino acid biosynthesis inhibition) type herbicides

(G-1) Glycine-based compound: glyphosate (including salts with sodium, amine, propylamine, isopropylamine, dimethylamine, and trimesium)

H. Glutamine synthesis inhibition herbicides

(H-1) Phosphinic acid-based compound: bilanafos, glufosinate (including salts with amine or sodium)

I. Dihydropteroic acid (DHP) inhibition herbicides

(I-1) Carbamate-based compound: asulam

K1. Microtubule association inhibition type herbicides

(K1-1) Benzamide-based compound: propyzamide and tebutam

(K1-2) Benzoic acid-based compound: chlorthal-dimethyl

(K1-3) Dinitroaniline-based compound: benfluralin, butralin, dinitramine, ethalfluralin, fluchloralin, oryzalin, pendimethalin, prodiamine, and trifluralin

(K1-4) Phosphoroamidate-based compound: amiprofos-methyl and butamifos

(K1-5) Pyridine-based compound: dithiopyr and thiazopyr

K2. Mitosis/Microtubule tissue formation inhibition herbicides

(K2-1) Carbamate-based compound: carbetamide, chlorpropham, propham, sweep, and karbutilate

K3. Very long-chain fatty acid (VLCFA) synthase inhibition herbicides

(K3-1) Acetamide-based compound: diphenamid, napropamide, and naproanilide

(K3-2) Chloroacetamide-based compound: acetochlor, alachlor, butachlor, butenachlor, diethatyl-ethyl, dimethachlor, dimethenamid, dimethenamid-P, metazachlor, metolachlor, pethoxamid, pretilachlor, propachlor, propisochlor, S-metolachlor, and thenylchlor

(K3-3) Oxyacetamide-based compound: flufenacet and mefenacet

(K3-4) Tetrazolinone-based compound: fentrazamide

(K3-5) Other compounds: anilofos, bromobutide, cafenstrole, indanofan, piperophos, fenoxasulfone, pyroxasulfone, and ipfencarbazone

L. Cellulose synthesis inhibition herbicides

(L-1) Benzamide-based compound: isoxaben

(L-2) Nitrile-based compound: dichlobenil, chlorthiamid

(L-3) Triazolocarboxamide-based compound: flupoxame

5 M. Uncoupler (cell membrane distruction) type herbicides

(M-1) Dinitrophenol-based compound: dinoterb and DNOC (including salts with amine or sodium)

N. Lipid bioynthesis (excluding ACCase inhibition) inhibition herbicides

(N-1) Benzofuran-based compound: benfuresate and ethofumesate

10 (N-2) Halogenated carboxylic acid-based compound: dalapon, flupropanate, and TCA (including salts with sodium, potassium, or ammonia)

(N-3) Phosphorodithioate-based compound: bensulide

(N-4) Thiocarbamate-based compound: butylate, cycloate, dimepiperate, EPTC, esprocarb, molinate, orbencarb, pebulate, prosulfocarb, thiobencarb, tiocarbazil, tri-allate, and vernolate

15 O. Auxin synthesis inhibition herbicides

(O-1) Benzoic acid-based compound: chloramben, 2,3,6-TBA, and dicamba (including salts with amine, diethyl amine, triethanolamine, isopropylamine, sodium, or lithium)

(O-2) Phenoxy carboxylic acid-based compound: 2,4,5-T, 2,4-D (including salts with amine, diethyl amine, isopropylamine, diglycolamine, sodium, or lithium), 2,4-DB, clomeprop, 20 dichlorprop, dichlorprop-P, MCPA, MCPA-thioethyl, MCPB (including sodium salt and ethyl ester), mecoprop (including salts with sodium, potassium, isopropylamine, triethanol amine, and dimethylamine), and mecoprop-P

(O-3) Pyridine carboxylic acid-based compound: clopyralid, fluroxypyr, picloram, triclopyr, and triclopyr-butotyl

25 (O-4) Quinoline carboxylic acid-based compound: quinclorac and quinmerac

(O-5) Other compounds: benazolin

P. Auxin transport inhibition type herbicides

(P-1) Phthalamates-based compound: naptalam (including salts with sodium)

(P-2) Semicarbazone-based compound: diflufenzopyr

Z. Herbicides with unknown mode of action

Flamprop-M (including methyl, ethyl, and isopropyl ester), flamprop (including methyl, ethyl, and isopropyl ester), chlorflurenol-methyl, cinmethylin, cumyluron, daimuron, methyldymuron, difenzoquat, etobenzanid, fosamine, pyributicarb, oxaziclomefone, acrolein,
 5 AE-F-150944 (code number), aminocyclopyrachlor, cyanamide, heptamaloxyloglucan, indaziflam, triaziflam, quinoclamine, endothal-disodium, phenisopham, BDPT, BAU-9403 (code number), SYN-523 (code number, SYP-249 (code number), JS-913 (code number), IR-6396 (code number), metiozolin, Triafamone, HW-02 (code number), and BCS-AA10579 (code number)

10 [Plant growth controlling compounds]

1-Methylcyclopropene, 1-naphthylacetamide, 2,6-diisopropyl-naphthalene, 4-CPA, benzylaminopurine, ancymidol, aviglycine, carvone, chlormequat, cloprop, cloxyfonac, cloxyfonac-potassium, cyclanilide, cytokinins, daminodide, dikegulac, dimethipin, ethephon, ethychlozate, flumetralin, flurenol, flurprimidol, forchlorfenuron, gibberellin acid, inabenfide,
 15 indol acetic acid, indol butyric acid, maleic hydrazide, mefluidide, mepiquat chloride, n-decanol, paclobutrazol, prohexadione-calcium, prohydrojasmon, sintofen, thidiazuron, triacontanol, trinexapac-ethyl, uniconazole, uniconazole-P, and ecolyst

Hereinbelow, known safeners which may be mixed or used in combination with the compound of the invention are exemplified, but the invention is not limited thereto: benoxacor,
 20 furilazole, dichlormid, dicyclonone, DKA-24 (N1,N2-diallyl-N2-dichloroacetyl-glycinamide), AD-67(4-dichloroacetyl-1-oxa-4-azaspiro[4.5]decane), PPG-1292 (2,2-dichloro-N-(1,3-dioxan-2-yl methyl)-N-(2-propenyl)acetamide), R-29148 (3-dichloroacetyl-2,2,5-trimethyl-1,3-oxazolidine), cloquintcet-mexyl, naphthalic anhydride (1,8-naphthalic anhydride), mefenpyr-diethyl, mefenpyr, mefenpyr-ethyl, fenchlorazole O ethyl,
 25 fenclorim, MG-191 (2-dichloromethyl-2-methyl-1,3-dioxane), cyometrinil, flurazole, fluxofenim, isoxadifen, isoxadifen-ethyl, mecoprop, MCPA, daimuron, 2,4-D, MON4660 (code number), oxabetrinil, cyprosulfamide, lower alkyl substituted benzoic acid, and TI-35 (code number).

Among other herbicidically active components that may be mixed or used in combination with the compound of the invention, known plant disease control agents are described below, but

the invention is not limited thereto.

1. Nucleic acid biosynthesis inhibitor

acyl alanine compound: benalaxyl, benalaxyl-M, furalaxyl, metalaxyl, and metalaxyl-M;

oxazolidinone-based compound: oxadixyl;

5 butylol lactone-based compound: clozylacon and ofurace;

hydroxy-(2-amino)pyrimidine-based compound: bupirimate, dimethirimol, and ethirimol;

isoxazole-based compound: hymexazol;

isotahiazolone-based compound: octhilinone;

carboxylic acid-based compound: oxolinic acid

10 2. Mitosis and cell differentiation inhibitor

benzimidazole-based compound: benomyl, carbendazim, fuberidazole, and thiabendazole;

thiophanate-based compound: thiophanate and thiophanate-methyl;

N-phenylcarbamate-based compound: diethofencarb;

toluamide-based compound: zoxamide;

15 phenylurea-based compound: pencycuron;

pyridinylmethyl benzamide-based compound: fluopicolide

3. Respiration inhibitor

pyrimidine amine-based compound: diflumetorim;

carboxamide-based compound: benodanil, flutolanil, mepronil, fluopyram, fenfuram,

20 carboxin, oxycarboxin, thifluzamide, bixafen, furametpyr, isopyrazam, penflufen, penthiopyrad, sedaxane, and boscalid;

methoxy acrylate-based compound: azoxystrobin, enestroburin, picoxystrobin, and pyraoxystrobin;

methoxycarbamate-based compound: pyraclostrobin, pyrametostrobin;

25 oxyiminoacetate compound: kresoxim-methyl and trifloxystrobin;

oxyiminoacetamide-based compound: dimoxystrobin, metominostrobin, and orysastrobin;

oxazolidinedione-based compound: famoxadone;

dihydrodioxadine-based compound: fluoxastrobin;

imidazolinone-based compound: fenamidone;

benzylcarbamate-based compound: pyribencarb;

cyanoimidazole-based compound: cyazofamid;

sulfamoyltriazole-based compound: amisulbrom;

dinitrophenylcrotonic acid-based compound: binapacryl, meptyldinocap, and dinocap;

5 2,6-dinitroaniline-based compound: fluazinam;

pyrimidinone hydrazone-based compound: ferimzone;

triphenyl tin-based compound: TPTA, TPTC, TPTH;

thiophenecarboxamide-based compound: silthiofam;

triazolopyrimidyl amine-based compound: ametoctradin

10 4. Amino acid and protein synthesis inhibitor

anilino pyrimidine-based compound: cyprodinil, mepanipyrin, and pyrimethanil;

enopyranuronic acid-based antibiotics: blasticidin-S and mildiomyacin;

hexopyranosyl-based antibiotics: kasugamycin;

glucopyranosyl-based antibiotics: streptomycin;

15 tetracycline-based antibiotics: oxytetracycline;

other antibiotics: gentamycin

5. Preparation acting on signal transduction pathway

quinoline-based compound: quinoxifen;

quinazoline-based compound: proquinazid;

20 phenylpyrrol-based compound: fenpiclonil and fludioxonil;

dicarboxyimide-based compound: chlozolate, iprodione, procymidone, and vinclozolin

6. Lipid and cell membrane synthesis inhibitor

phosphorothiorate-based compound: edifenphos, iprobenfos, and pyrazophos;

dithiolane-based compound: isoprothiolane;

25 aromatic hydrocarbon-based compound: biphenyl, chloroneb, dicloran, quintozone,
tecnazene, and tolclufos-methyl;

1,2,4-thiadiazole-based compound: etridiazole;

carbamate-based compound: iodocarb, propamocarb-hydrochloride, and prothiocarb;

cinnamic amide-based compound: dimethomorph and flumorph;

valine amide carbamate-based compound: benthiavalicarb-isopropyl, iprovalicarb, and valifenalate;

mandelic amide-based compound: mandipropamid;

Bacillus subtilis and bactericidal lipopeptide product: Bacillus subtilis (strain: QST 713)

5 7. Sterol biosynthesis inhibitor

piperazine-based compound: triforine;

pyridine-based compound: pyrifenox;

pyrimidine-based compound: fenarimol and nuarimol;

10 imidazole-based compound: imazalil, oxpoconazole-fumarate, pefurazoate, prochloraz, and triflumizole;

triazole-based compound: azaconazole, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, etaconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, 15 tetraconazole, triadimefon, triadimenol, triticonazole, furconazole, furconazole-cis, and quinconazole;

morpholine-based compound: aldimorph, dodemorph, fenpropimorph, and tridemorph;

piperidine-based compound: fenpropidin and piperalin;

spiroketal amine-based compound: spiroxamine;

20 hydroxy anilide-based compound: fenhexamid;

thiocarbamate-based compound: pyributicarb;

aryl amine-based compound: naftifine and terbinafine

8. Glucan biosynthesis inhibitor

glucopyranosyl-based antibiotics: validamycin;

25 peptidyl pyridine nucleotide compound: polyoxin

9. Melanine synthesis inhibitor

isobenzofuranone-based compound: phthalide;

pyrroloquinoline-based compound: pyroquilon;

triazolobenzothiazole-based compound: tricyclazole;

carboxamide-based compound: carpropamid, diclocymet;

propionamide-based compound: fenoxanil

10. Preparation for inducing resistance to plant disease

benzothiadiazole-based compound: acibenzolar-S-methyl;

5 benzoisothiazole-based compound: probenazole;

thiadiazole carboxamide-based compound: tiadinil and isotianil;

natural product: laminarin

11. Preparation with unknown mode of action or multiple mode of action

copper compound: copper hydroxide, copper dioctanoate, copper oxychloride, copper

10 sulfate, cuprous oxide, oxine-copper, Bordeaux mixture, and copper nonyl phenol sulphonate;

sulfur compound: sulfur;

dithiocarbamate-based compound: ferbam, mancozeb, maneb, metiram, propineb, thiram, zineb, ziram, and cufraneb;

phthalimide-based compound: captan, folpet, and captafol;

15 chloronitrile-based compound: chlorothalonil;

sulfamide-based compound: dichlofluanid, tolylfluanid;

guanidine-based compound: guazatine, iminoctadine-albesilate, and iminoctadine-triacetate, dodine;

other compounds: anilazine, dithianon, cymoxanil, vfosetyl (aluminium, calcium, and

20 sodium), phosphorous acid and salts, tecloftalam, triazoxide, flusulfamide, diclomezine,

methasulfocarb, ethaboxam, cyflufenamid, metrafenone, potassium bicarbonate, sodium

bicarbonate, BAF-045 (code number), BAG-010 (code number), benthiazole, bronopol, carvone,

chinomethionat, dazomet, DBEDC, debacarb, dichlorophen, difenzoquat-methyl sulfate, dimethyl

disulfide, diphenylamine, ethoxyquin, flumetover, fluoroimide, flutianil, fluxapyroxad,

25 furancarboxylic acid, metam, nabam, natamycin, nitrapyrin, nitrothal-isopropyl, o-phenylphenol,

oxazinylazole, oxyquinoline sulfate, phenazine oxide, polycarbamate, pyriofenone, S-2188 (code

number), silver, SYP-Z-048 (code number), tebufloquin, tolnifanide, trichlamide, mineral oils,

and organic oils

12. Microorganisms and products of microorganisms

Agrobacterium radiobacter, Fermented product from Aspergillus spp., Bacillus spp., Harpin protein, Erwinia carotovora, Fusarium oxysporum, Gliocladium spp., Laccase, Pseudomonas spp., Talaromyces spp., Trichoderma spp., Extract from mushroom, and Bacteriophage

Among other herbicidically active components that may be mixed or used in combination
 5 with the compound of the invention, known pesticides, acaricides, nematocides, and synergistic agents are described below, but the invention is not limited thereto.

[Pesticides, acaricides & nematocides]

1. Acetylcholine esterase inhibitor:

(1A) carbamate compound: alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb,
 10 butocarboxim, butoxycarboxim, carbaryl, carbofuran, carbosulfan, ethiofencarb, fenobucarb, formetanate, furathiocarb, isoprocarb, methiocarb, methomyl, metolcarb, oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, triazamate, trimethacarb, 3,5-xylyl methylcarbamate(XMC), and xylylcarb

(1B) organo phosphorus compound: acephate, azamethiphos, azinphos-ethyl,
 15 azinphos-methyl, cadusafos, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos, chlorpyrifos-methyl, coumaphos, cyanophos, demeton-S-methyl, diamidafos, diazinon, dichlorvos, dicrotophos, dimethoate, dimethylvinphos, dioxabenzofos, disulfoton, DSP, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fenthion, fonofos, fosthiazate, fosthietan, heptenophos, isamidofos, isazophos, isofenphos-methyl, isopropyl
 20 O-(methoxyaminothio-phosphoryl) salicylate, isoxathion, malathion, mecarbam, methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, oxydeprofos, parathion, parathion-methyl, phenthoate, phorate, phosalone, phosmet, phosphamidon, phoxim, pirimiphos-methyl, profenofos, propaphos, propetamphos, prothiofos, pyraclofos, pyridaphenthion, quinalphos, sulfotep, tebupirimfos, temephos, terbufos, tetrachlorvinphos,
 25 thiometon, thionazin, triazophos, trichlorfon, vamidothion, dichlofenthion, imicyafos, isocarbophos, mesulfenfos, and flupyrazofos

2. GABA receptor (chloride channel) inhibitor

(2A) cyclodiene organic chloride-based compound: chlordane, endosulfan, and gamma-BCH

(2B) phenylpyrazole-based compound: acetoprol, ethiprole, fipronil, pyrafluprole, pyriprole, and RZI-02-003 (code number)

3. Preparation acting on sodium channel

(3A) pyrethroid-based compound: acrinathrin, allethrin [including d-cis-trans and d-trans],
 5 bifenthrin, bioallethrin, bioallethrin S-cyclopentenyl, bioresmethrin, cycloprothrin, and cyfluthrin [including beta-], cyhalothrin [including gamma- and lambda-], cypermethrin [including alpha-, beta-, theta-, and zeta-], cyphenothrin [including (1R)-trans-isomers], deltamethrin, empenthrin, esfenvalerate, etofenprox, fenpropathrin, fenvalerate, flucythrinate, flumethrin, and
 10 tau-fluvalinate [including tau-], halfenprox, imiprothrin, metofluthrin, permethrin, and phenothrin [including (1R)-trans-isomer], prallethrin, profluthrin, pyrethrine, resmethrin, RU15525 (code number), silafluofen, tefluthrin, tetramethrin, tralomethrin, transfluthrin, ZXI8901 (code number), fluvalinate, tetramethylfluthrin, and meperfluthrin

(3B) DDT-based compound: DDT, methoxychlor

4. Nicotinic acetylcholine receptor agonist/antagonist

15 (4A) neonicotinoid-based compound: acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, thiacloprid, and thiamethoxam

(4B) nicotine-based compound: nicotine-sulfate

5. Nicotinic acetylcholine receptor allosteric activator

spinosyn-based compound: spinetoram and spinosad

20 6. Chloride channel activating preparation

avermectin, milbemycin-based compound: abamectin, emamectin benzoate, lepimectin, milbemectin, ivermectin, and polynactins

7. Juvenile hormone preparation

diofenolan, hydroprene, kinoprene, methothrin, fenoxycarb, and pyriproxyfen

25 8. Preparation with non-specific mode of action (multiple mode of action)

1,3-dichloropropene, DCIP, ethylene dibromide, methyl bromide, chloropicrin, and sulfuryl fluoride

9. Feeding inhibitor

pymetrozine, flonicamid and pyrifluquinazon

10. Mite growth controlling agent

clofentezine, diflovidazin, hexythiazox, and etoxazole

11. Preparation for disrupting insect intima

BT preparation:

5 12. ATP biosynthesis enzyme inhibitor

diafenthiuron;

organo tin compound: azocyclotin, cyhexatin, and fenbutatin oxide;

propargite, tetradifon

13. Uncoupler

10 chlorfenapyr and DNOC

14. Preparation for blocking nicotinic acetylcholine channel

nereistoxin-based compound: bensultap, cartap, thiocyclam, and thiosultap

15. Chitin biosynthesis inhibitor (type 0)

15 benzoylurea-based compound: bistrifluron, chlorfluazuron, diflubenzuron, flucycloxuron,
flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron, teflubenzuron, triflumuron,
and fluazuron

16. Chitin biosynthesis inhibitor (type 1)

buprofezin

17. Molting inhibitor (for Diptera)

20 cyromazine

18. Ecdysone agonist (for promoting molting)

diacylhydrazine-based compound: chromafenozide, halofenozide, methoxyfenozide, and
tebufenozide

19. Octopamine agonist

25 amitraz

20. Mitochondrial electron transport chain (complex III) inhibitor

cyflumetofen, hydramethylnon, acequinocyl, fluacrypyrim, and cyenopyrafen

21. Mitochondrial electron transport chain (complex I) inhibitor

METI acaricides: fenazaquin, fenpyroximate, pyridaben, pyrimidifen, tebufenpyrad, and

tolfenpyrad

others: rotenone

22. Sodium channel inhibitor

indoxacarb and metaflumizon

5 23. Lipid biosynthesis inhibitor

tetronic-based insecticides/acaricides: spiroticlofen, spiromesifen, and spirotetramat

24. Mitochondrial electron transport chain (complex IV) inhibitor

aluminium phosphide, phosphine, zinc phosphide, calcium cyanide, and phosphine

25. Neuronal inhibitor preparation (unknown mode of action)

10 bifenazate

26. Aconitase inhibitor

sodium fluoroacetate

27. Preparation acting on ryanodine receptor

chlorantraniliprole, flubendiamide and cyantraniliprole

15 28. Other preparations (unknown mode of action)

azadirachtin, amidoflumet, benclonthiaz, benzoximate, bromopropylate, chinomethionat, CL900167 (code number), cryolite, dicofol, dicyclanil, dienochlor, dinobuton, fenbutatin oxide, fenothiocarb, fluensulfone, flufenerim, flusulfamide, karanjin, metham, methoprene, methoxyfenozide, methyl isothiocyanate, pyridalyl, pyrifluquinazon, sulcofuron-sodium, 20 sulflamid, and sulfoxaflor

29. Synergistic agent

piperonyl butoxide and DEF.

Hereinafter, production methods of the compound of Formula 1 according to the compound of the invention, formulation examples, and applications will be described in detail with reference 25 to Examples below. However, the invention is not limited to these Examples in any way. In the description below, "%" means "percent by weight" and "parts" means "parts by weight".

[Example 1]

Production of 6-(2-hydroxy-6-oxo cyclohexa-1-enecarbonyl)-2-methyl-4-phenyl-1,2,4-triazine-3,5(2H, 4H)-dione (Compound No.

I-50)

(1) Production of 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride

0.93 g (3.76 mmol) of

5 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid and 0.72 g (5.64 mmol) of oxalyl chloride were dissolved in dichloromethane (20 ml). To the mixture, a drop of N,N-dimethylformamide was added and stirred at room temperature for 2 hours. The reaction solution was concentrated to obtain
2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride as a pale
10 yellow oily substance.

(2) Production 6-(2-hydroxy-6-oxo
cyclohexa-1-enecarbonyl)-2-methyl-4-phenyl-1,2,4-triazine-3,5(2H, 4H)-dione

0.63 g (5.64 mmol) of 1,3-cyclohexanedione and 0.57 g (5.64 mmol) of triethylamine were dissolved in dichloromethane (20 ml) under ice cooling. To the mixture, the dichloromethane
15 solution (10 ml) of 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride produced from the above (1) was slowly added dropwise, and stirred for 30 minutes under ice cooling. The reaction mixture was extracted with chloroform, and the organic layer was washed with water, dried over magnesium sulfate, and concentrated under reduced pressure. The residues obtained were dissolved in acetonitrile (20 ml), added with 0.57 g (5.64 mmol) of
20 triethylamine and 0.03 g (0.38 mmol) of acetone cyanohydrin, and refluxed for 30 minutes under heating. After concentration under reduced pressure, the residues were dissolved in water and washed with ethyl acetate. The aqueous layer was acidified by using citric acid, extracted with chloroform, dried over magnesium sulfate, and concentrated under reduced pressure. The crystals obtained were washed with methanol to obtain 0.36 g of the target compound (yield
25 28%).

Melting point: 182 to 185°C

[Example 2]

Production of

6-(5-hydroxy-1-methyl-1H-pyrazole-4-carbonyl)-2-methyl-4-phenyl-1,2,4-triazine-3,5(2H,

4H)-dione (Compound No. II-50)

1.50 g (6.07 mmol) of
2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid and 1.16 g (9.10
mmol) of oxalyl chloride were dissolved in dichloromethane (30 ml). To the mixture, a drop of
5 N,N-dimethylformamide was added and stirred at room temperature for 2 hours. The reaction
solution was concentrated under reduced pressure to obtain
2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride as a pale
yellow oily substance.

Next, 1.22 g (9.10 mmol) of 1-methyl-5-hydroxypyrazole hydrochloride and 1.53 g (15.17
10 mmol) of triethylamine were added to dichloromethane (30 ml) under ice cooling. To the
mixture, the dichloromethane solution (15 ml) of
2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride was slowly
added dropwise, and stirred for 30 minutes. The reaction mixture was extracted with
chloroform, and the organic layer was washed with water, dried over magnesium sulfate, and
15 concentrated under reduced pressure. The residues obtained were dissolved in acetonitrile (30
ml), added with 0.92 g (9.10 mmol) of triethylamine and 0.05 g (0.61 mmol) of acetone
cyanohydrin, and refluxed for 30 minutes under heating. The reaction mixture was concentrated
under reduced pressure, and then the residues were dissolved in water and washed with ethyl
acetate. The aqueous layer was acidified by using citric acid, extracted with chloroform, dried
20 over magnesium sulfate, and concentrated under reduced pressure. The crystals obtained were
washed with methanol to obtain 0.40 g of the target compound (yield 20%).

Melting point: 197 to 199°C

[Example 3]

Production of 6-(2-hydroxy-4-oxobicyclo[3.2.1]octa-2-en-yl
25 carbonyl]-2-methyl-4-phenyl-1,2,4-triazine-3,5(2H, 4H)-dione (Compound No. III-50)

1.00 g (4.04 mmol) of
2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid and 1.03 g (8.09
mmol) of oxalyl chloride were dissolved in dichloromethane (20 ml). To the mixture, a drop of
N,N-dimethylformamide was added and stirred at room temperature for 2 hours. The reaction

solution was concentrated under reduced pressure to obtain 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride as a pale yellow oily substance.

Next, 0.83 g (6.07 mmol) of bicyclo[3.2.1]octane-2,4-dione and 0.61 g (6.07 mmol) of triethylamine were dissolved in dichloromethane (20 ml) under ice cooling. To the solution, the dichloromethane solution (10 ml) of 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carbonyl chloride that is prepared previously was slowly added dropwise. After stirring for 30 minutes under ice cooling, the reaction mixture was extracted with chloroform, and the organic layer was washed with water, dried over magnesium sulfate, and concentrated under reduced pressure. The residues obtained were dissolved in acetonitrile (20 ml), added with 0.61 g (6.07 mmol) of triethylamine and 0.03 g (0.4 mmol) of acetone cyanohydrin, and refluxed for 30 minutes under heating. The reaction mixture was concentrated under reduced pressure, and then the residues were dissolved in water and washed with ethyl acetate. The aqueous layer was acidified by using citric acid, extracted with chloroform, dried over magnesium sulfate, and concentrated under reduced pressure. The crystals obtained were washed with methanol to obtain 0.70 g of the target compound (yield 47%).

Melting point: 163 to 165°C

[Example 4]

20 Production of

1-isopropyl-4-(2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-yl carbonyl)-1H-pyrazole-5-yl propane-1-sulfonate (Compound No. II-267)

0.85 g (2.60 mmol) of 6-(5-hydroxy-1-isopropyl-1H-pyrazol-4-yl carbonyl)-2-methyl-4-phenyl-1,2,4-triazine-3,5(2H, 4H)-dione was dissolved in 20 ml of dichloromethane. To the solution, 0.27 g (2.60 mmol) of triethylamine and 0.37 g (2.60 mmol) of 1-propane sulfonyl chloride were added at room temperature and stirred overnight. The reaction mixture was concentrated under reduced pressure, and the residues were purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1) to obtain 0.71 g of the target compound (yield 63%).

Melting point: 51 to 53°C

[Example 5]

Production of

2-methyl-3,5-dioxo-4-(4-chlorophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

5 (Compound No. V-53)

(1) Production of diethyl 2-(2-methylhydrazono) malonate

5.00 g (0.0287 mol) of diethyl ketomalonate was dissolved in 30 ml ethanol. To the solution, 1.45 g (0.0316 mol) of methyl hydrazine was added and stirred for 7 hours at 60°C followed by further stirring overnight at room temperature. The reaction mixture was
10 concentrated under reduced pressure and extracted with ethyl acetate. The organic layer was washed with water, dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residues were purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1) to obtain 5.28 g of the target compound (yield 91%).

(2) Production of ethyl

15 4-(4-chlorophenyl)-2-methyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

2.00 g (9.89 mmol) of diethyl 2-(2-methylhydrazono) malonate and 1.50 g (9.89 mmol) of DBU were dissolved in 50 ml of tetrahydrofuran. To the solution, the tetrahydrofuran (10 ml) solution of 4-chlorophenyl isocyanate (3.34 g, 21.7 mmol) was slowly added dropwise at room temperature, and stirred over night. The reaction mixture was concentrated under reduced
20 pressure, and the residues were extracted with ethyl acetate, washed with water, dried over magnesium sulfate, and concentrated under reduced pressure. The resulting residues were purified by silica gel column chromatography (hexane : ethyl acetate = 7 : 1) to obtain 2.00 g of the target compound (yield 65%).

(3) Production of

25 2-methyl-3,5-dioxo-4-(4-chlorophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

2.00 g (6.46 mmol) of ethyl

2-methyl-4-(4-chlorophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester was stirred at room temperature for 2 days in a mixed solvent of acetic acid (30 ml) and conc. hydrochloric acid (30 ml). The reaction mixture was concentrated under reduced pressure to obtain 1.88 g of

the target compound (yield, quantitative).

Melting point: 234 to 236°C

[Example 6]

Production of 2,4-dimethyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

5 (Compound No. V-1)

(1) Production of 2-methylsemicarbazide

13 g (282.1 mmol) of methyl hydrazine was dissolved in 60 ml of tetrahydrofuran. To the solution, 25 g (217 mmol) of trimethylsilyl isocyanate was slowly added dropwise at 0°C and further stirred for 1 hour. To the reaction mixture, 40 ml of methanol was added and stirred for 10 5 hours at 40°C. The reaction mixture was concentrated to obtain 18 g of 2-methyl semicarbazide as a pale yellow solid (yield 93%).

¹H-NMR(CDCl₃,TMS) δ(ppm):

3.15(3H,s), 3.80(2H,br), 5.61(2H,br)

(2) Production of ethyl 2-methyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic
15 acid ester

35.2 g (202 mmol) of diethyl ketomalonate and 18 g (202 mmol) of 2-methyl semicarbazide were dissolved in 200 ml ethanol, and then refluxed under heating for 36 hours. The reaction solution was concentrated to obtain 31 g of ethyl 2-methyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester as a white solid
20 (yield 78%).

¹H-NMR(CDCl₃,TMS) δ(ppm):

1.39(3H,t,J=7.1Hz), 3.72(3H,s), 4.42(2H,q,J=7.1Hz), 9.38(1H,br)

(3) Production of ethyl

2,4-dimethyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

25 2.0 g (10.0 mmol) of ethyl 2-methyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester, 1.9 g (13.5 mmol) of potassium carbonate, and 1.8 g (12.5 mmol) of methyl iodide were added to 20 ml of N,N-dimethylformamide, and stirred for 2 hours at 60°C. Upon the completion of the reaction, the reaction solution was added with water, and then extracted with ethyl acetate. The organic layer obtained was dried over anhydrous magnesium sulfate and

concentrated to obtain 1.8 g of ethyl

2,4-dimethyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester (yield 86%).

$^1\text{H-NMR}(\text{CDCl}_3, \text{TMS}) \delta(\text{ppm})$:

1.40(3H,t,J=7.1Hz), 3.38(3H,s), 3.74(3H,s), 4.42(2H,q,J=7.1Hz)

- 5 (4) Production of 2,4-dimethyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid
1.8 g (8.41 mmol) of ethyl

2,4-dimethyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester was stirred at room temperature for 24 hours in a mixed solvent of acetic acid (30 ml) and conc. hydrochloric acid (30 ml). The reaction solution was concentrated to obtain 1.40 g of

- 10 2,4-dimethyl-3,5-dioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid as a white solid (yield 90%).

Melting point: 220 to 223°C

$^1\text{H-NMR}(\text{CDCl}_3, \text{TMS}) \delta(\text{ppm})$:

3.48(3H,s), 3.88(3H,s)

- 15 [Example 7]

Production of 2-ethyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid (Compound No. V-259)

(1) Production of ethyl 3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

- 20 9.0 g (0.0517 mol) of diethyl ketomalonate and 7.81 g (0.0517 mol) of 2-phenyl semicarbazide were stirred in 50 ml xylene for 1 hour at 100°C. The reaction mixture was refluxed under heating, and by adding sodium methoxide (8.37 g, 0.155 mol) in small portions, the reaction was completed. After cooling to room temperature, the reaction mixture was neutralized with 1 N aqueous hydrochloric acid solution, extracted with ethyl acetate, and dried
25 over magnesium sulfate. The reaction mixture was concentrated under reduced pressure and the residues were isolated and purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1) to obtain 6.18 g of the target compound (yield 46%).

(2) Production of ethyl

2-ethyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

1.50 g (5.74 mmol) of ethyl

3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester was dissolved in 30 ml of N,N-dimethylformamide, added with 60% sodium hydride (0.23 g, 5.74 mmol) under ice cooling, and further stirred for 30 minutes. The mixture was added with ethyl iodide (0.90 g, 5.74 mmol) and stirred. After raising to room temperature, an aqueous solution of ammonium chloride was added to terminate the reaction. The resultant was extracted with diethyl ether, dried over magnesium chloride, and concentrated under reduced pressure. The residues were purified by silica gel column chromatography to obtain 1.33 g of the target compound (yield 80%).

(3) Production of 2-ethyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

1.30 g (4.49 mmol) of ethyl

2-ethyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester was dissolved in 30 ml ethanol, added with a 25% aqueous solution of sodium hydroxide (1.29 g, 8.09 mmol), and stirred overnight. After dilution by adding water, the aqueous layer was washed with diethyl ether. The aqueous layer was acidified by adding 6 N aqueous hydrochloric acid solution, and then extracted with ethyl acetate. After drying over magnesium sulfate and concentration under reduced pressure, 1.10 g of the target compound was obtained (yield 94%).

[Example 8]

Production of 2,4-dimethyl-5-oxo-3-thioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid (Compound No. V-265)

(1) Production of ethyl

2,4-dimethyl-5-oxo-3-thioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

2.00 g (9.89 mmol) of diethyl 2-(2-methylhydrazono) malonate and 1.50 g (9.89 mmol) of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) were dissolved in 50 ml of tetrahydrofuran. To the solution, the tetrahydrofuran (10 ml) of methylisothiocyanate (1.58 g, 21.7 mmol) was slowly added dropwise and stirred overnight. The reaction mixture was concentrated under reduced pressure, extracted with ethyl acetate, washed with water, and dried over magnesium sulfate. The residues obtained after concentration under reduced pressure were purified by silica gel

column chromatography (hexane : ethyl acetate = 3 : 1) to obtain 2.20 g of the target compound (yield 97%).

(2) Production of

2,4-dimethyl-5-oxo-3-thioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

5 2.30 g (0.01 mol) of ethyl

2,4-dimethyl-5-oxo-3-thioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester was stirred overnight at room temperature in a mixed solvent of acetic acid (30 ml) and conc. hydrochloric acid (30 ml). The reaction mixture was concentrated under reduced pressure to obtain 2.01 g of the target compound (yield; quantitative).

10 [Example 9]

Production of

2-methyl-3,5-dioxo-4-(2-cyanophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid
(Compound No. V-72)

(1) Production of ethyl

15 2-methyl-3,5-dioxo-4-(2-cyanophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

2.0 g (9.89 mmol) of diethyl 2-(2-methylhydrazono) malonate and 3.3 g (21.8 mmol) of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) were dissolved in 20 ml of tetrahydrofuran. To the solution, 4.9 g (20.8 mmol) of phenyl-2-cyanophenylcarbamate was added at room temperature and stirred for 1 hour at the same temperature. After that, the mixture was refluxed under
20 heating for 3 hours. The reaction solution was concentrated and the residues were extracted with ethyl acetate. The organic layer obtained was washed with water and an aqueous solution of citric acid in order, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residues were purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1) to obtain 2.3 g of ethyl

25 2-methyl-3,5-dioxo-4-(2-cyanophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester (yield 78%).

¹H-NMR(CDCl₃, TMS) δ(ppm):

1.40(3H,t,J=7.1Hz), 3.81(3H,s), 4.45(2H,q,J=7.1Hz), 7.39(1H,d,J=8.0Hz),

7.60-7.64(1H,m), 7.75-7.80(1H,m), 7.85(1H,d,J=7.6Hz)

(2) Production of

2-methyl-3,5-dioxo-4-(2-cyanophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

2.3 g (7.65 mmol) of ethyl

2-methyl-3,5-dioxo-4-(2-cyanophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

5 was stirred for 24 hours at room temperature in a mixed solvent of acetic acid (30 ml) and conc. hydrochloric acid (30 ml). The reaction solution was concentrated under reduced pressure to obtain 1.8 g of

2-methyl-3,5-dioxo-4-(2-cyanophenyl)-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid as a white solid (yield 90%).

10 Melting point: 213 to 215°C

$^1\text{H-NMR}$ (DMSO- d_6 , TMS) δ (ppm):

3.65(3H,s), 7.67(1H,d,J=8.0Hz), 7.70-7.75(1H,m), 7.90-7.96(1H,m),

8.09(1H,d,J=7.4Hz), 14.02(1H,br)

[Example 10]

15 Production of 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid (Compound No. V-50)

(1) Production of ethyl

2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

20 2.0 g (9.89 mmol) of diethyl 2-oxomalonate and 0.04 g (0.2 mmol) of p-toluene sulfonic acid were dissolved in 50 ml of toluene. To the solution, 2.5 g (15.2 mmol) of

1-methyl-N-phenylhydrazine carboxamides was added at room temperature, and then stirred for 2 hours with reflux under heating. The reaction mixture was cooled to room temperature and added with 0.08 g (0.5 mmol) of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) followed by stirring at room temperature for two hours. The reaction solution was washed with water and dried over

25 magnesium sulfate. The solvent was distilled off to obtain ethyl

2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester.

(2) Production of

2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid

Ethyl 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid ester

produced from the above (1) was stirred for 24 hours at room temperature in a mixed solvent of acetic acid (30 ml) and conc. hydrochloric acid (30 ml). The reaction mixture was concentrated under reduced pressure, extracted with a saturated aqueous solution of sodium hydrogen carbonate, washed with ethyl acetate, and then adjusted to be weakly acidic by using diluted hydrochloric acid. After that, the mixture was extracted with ethyl acetate and dried over magnesium sulfate, and the solvent was distilled off to obtain 2.6 g of 2-methyl-3,5-dioxo-4-phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylic acid as a white solid (2-step yield 70%).

Melting point: 195 to 198°C

¹H-NMR(DMSO-d₆, TMS) δ(ppm):

3.59(3H,s), 7.29-7.31(2H,m), 7.43-7.54(3H,m), 13.64(1H,bs)

Physical property values (melting point or refractive index) of the compound of the invention represented by Formula 1, which has been synthesized according to the above Examples, are shown in Table 68 to Table 70 including above Examples. Herein, * means refractive index.

[Table 68]

Compound No.	Melting Point(°C) or Refractive Index(n_D^{20})	Compound No.	Melting Point(°C) or Refractive Index(n_D^{20})
I-2	87-89	I-83	191-194
I-3	1.5530*	I-84	124-127
I-5	1.5630*	I-85	235-238
I-9	1.5380*	I-86	199-202
I-10	124-125	I-87	197-198
I-11	97-98	I-88	160-163
I-14	126-129	I-89	190-193
I-16	116-118	I-90	164-166
I-19	132-134	I-91	89-91
I-27	1.5460*	I-92	245-247
I-41	1.5495*	I-93	168-169
I-43	98-101	I-94	155-157
I-47	155-157	I-96	151-153
I-50	182-185	I-98	155-157
I-51	184-185	I-99	178-181
I-52	187-190	I-105	186-188
I-53	182-183	I-106	228-231
I-54	174-176	I-107	212-215
I-55	209-212	I-108	167-169
I-56	181-183	I-109	166-168
I-57	135-136	I-110	151-152
I-58	198-199	I-111	196-199
I-59	190-193	I-115	144-147
I-60	190-191	I-116	176-179
I-61	186-187	I-117	140-143
I-62	137-139	I-118	140-143
I-63	166-169	I-119	191-194
I-64	89-92	I-120	191-194
I-65	184-187	I-125	148-151
I-66	151-152	I-126	126-129
I-67	174-177	I-127	237-240
I-68	208-210	I-128	217-220
I-71	130-131	I-129	155-158
I-72	166-169	I-131	204-205
I-73	181-184	I-134	215-217
I-74	108-111	I-135	152-154
I-75	173-176	I-136	156-157
I-76	242-245	I-137	154-157
I-77	192-194	I-138	123-126
I-78	149-151	I-149	175-178
I-79	161-163	I-155	196-199
I-80	98-101	I-167	183-185
I-81	158-161	I-169	178-180
I-82	212-215	I-170	213-215

[Table 69]

Compound No.	Melting Point(°C) or Refractive Index(n_D^{20})	Compound No.	Melting Point(°C) or Refractive Index(n_D^{20})
I-179	215-218	I-293	158-160
I-182	159-161	I-294	113-115
I-183	138-141	I-295	1.5360*
I-184	100-103	I-296	1.5300*
I-185	108-111	I-297	89-92
I-187	180-183	I-298	148-150
I-189	190-193	I-299	212-215
I-198	135-137	I-300	203-205
I-199	169-170	I-301	274-277
I-202	161-162	I-302	222-224
I-203	188-191	I-303	62-65
I-204	201-204	I-304	148-151
I-205	87-90	I-307	58-61
I-259	150-153	I-328	58-61
I-260	152-154	I-463	131-134
I-261	190-193	I-464	168-170
I-262	103-106	I-465	211-213
I-263	174-176	I-466	89-92
I-265	164-167	I-467	211-214
I-268	201-204	I-468	128-130
I-269	112-115	I-469	172-174
I-270	172-175	I-470	147-148
I-271	251-254	I-471	1.5620*
I-272	204-207	I-472	162-164
I-274	101-103	I-473	143-146
I-275	89-92	I-474	70-73
I-276	167-170	I-475	83-86
I-277	96-99	I-476	191-193
I-278	98-101	I-477	149-151
I-279	218-220	I-478	1.5270*
I-280	168-171	I-479	1.5450*
I-281	146-147	I-480	179-181
I-282	148-151	II-50	197-199
I-283	172-175	II-267	51-53
I-284	160-162	III-50	163-165
I-285	149-152	III-62	158-159
I-286	88-91	VI-1	151-154
I-287	155-158	VI-5	145-148
I-288	94-97	VI-6	145-146
I-289	215-218	VI-7	163-166
I-290	138-141	VI-65	93-96
I-291	194-197	VI-97	158-160
I-292	167-169		

Compound number and ^1H -NMR data (standard; TMS, δ (ppm) value) are given below.

Data without a name of solvent are measured by using CDCl_3 .

5 Compound No. I-1:

2.04-2.10(2H,m), 2.45-2.49(2H,m), 2.76-2.80(2H,m), 3.56(3H,s),

3.65(3H,s), 16.05(1H,br)

Compound No. I-3:

0.92(3H,t,J=6.00Hz), 1.69(2H,q,J=6.00Hz), 2.03-2.11(2H,m),
2.45-2.49(2H,m), 2.75-2.79(2H,m), 3.64(3H,s), 3.89(2H,t,J=6.00Hz),

5 16.05(1H,br)

Compound No. I-4:

1.49(6H,d,J=6.00Hz), 2.03-2.11(2H,m), 2.44-2.49(2H,m), 2.74-2.79(2H,m),
3.61(3H,s), 5.07(1H,sept,J=6.00Hz), 16.08(1H,br)

Compound No. I-5:

10 0.95(3H,t,J=7.2Hz), 1.32-1.43(2H,m), 1.59-1.68(2H,m), 2.03-2.10(2H,m),
2.45-2.49(2H,m), 2.75-2.79(2H,m), 3.64(3H,s), 3.92(2H,t,J=6.9Hz),
16.05(1H,br)

Compound No. I-9:

15 0.88(3H,t,J=6.6Hz), 1.20-1.40(6H,m), 1.58-1.64(2H,m), 2.03-2.12(2H,m),
2.44-2.48(2H,m), 2.75-2.79(2H,m), 3.64(3H,s), 3.89-3.94(2H,m),
16.04(1H,br)

Compound No. I-27:

1.65(3H,t,J=3.00Hz), 2.03-2.09(2H,m), 2.31-2.36(2H,m), 2.44-2.49(2H,m),
2.74-2.79(2H,m), 3.64(3H,s), 4.01(2H,t,J=6.00), 16.00(1H,br)

20 Compound No. I-41:

[0251]

1.89-1.97(2H,m), 2.04-2.11(2H,m), 2.44-2.48(2H,m), 3.31(3H,s),
3.44(2H,t,J=6.0Hz), 3.64(3H,s), 4.03(2H,t,J=7.0Hz), 16.04(1H,br)

Compound No. I-75:

25 2.05-2.11(2H,m), 2.45-2.49(2H,m), 2.75-2.80(2H,m), 3.69(3H,s),
7.05-7.09(1H,m), 7.14-7.21(1H,m), 7.24-7.33(1H,m), 15.99(1H,s)

Compound No. I-76:

2.04-2.09(2H,m), 2.46-2.50(2H,m), 2.75-2.80(2H,m), 3.69(3H,s),
6.88-6.96(3H,m), 15.97(1H,s)

Compound No. I-77:

2.03-2.09(2H,m), 2.45-2.49(2H,m), 2.75-2.78(2H,m), 3.71(3H,s),
7.11-7.14(1H,m), 7.18-7.33(2H,m), 15.95(1H,s)

Compound No. I-79:

5 2.04-2.10(2H,m), 2.45-2.50(2H,m), 2.75-2.79(2H,m), 3.70(3H,s),
7.10-7.24(3H,m), 15.96(1H,s)

Compound No. I-80:

2.01-2.08(2H,m), 2.46-2.49(2H,m), 2.75-2.78(2H,m), 3.71(3H,s),
7.05-7.08(2H,m), 7.40-7.48(1H,m), 15.93(1H,s)

10 Compound No. I-81:

[0252]

2.05-2.08(2H,m), 2.45-2.50(2H,m), 2.75-2.80(2H,m), 3.69(3H,s),
7.14-7.19(1H,m), 7.43(1H,d,J=2.5), 7.57(1H,d,J=8.5), 15.97(1H,s)

Compound No. I-295:

15 0.85-0.89(3H,m), 1.26-1.32(10H,m), 1.57-1.65(2H,m), 2.05-2.12(2H,m),
2.44-2.49(2H,m), 2.75-2.79(2H,m), 3.64(3H,s), 3.88-3.93(2H,m),
16.04(1H,br)

Compound No. I-296:

0.85-0.90(3H,m), 1.25-1.36(14H,m), 1.59-1.69(2H,m), 2.05-2.09(2H,m),
20 2.44-2.49(2H,m), 2.74-2.79(2H,m), 3.64(3H,s), 3.88-3.93(2H,m),
16.04(1H,br)

Compound No. I-306:

0.96(3H,t,J=7.15), 1.39-1.46(2H,m), 1.69-1.71(2H,m), 2.05-2.09(2H,m),
2.44-2.48(2H,m), 4.01(2H,t,J=7.69), 7.32-7.36(2H,m), 7.56-7.59(1H,m),
25 7.83-7.88(1H,m), 8.61-8.63(1H,m), 16.05(1H,br)

Compound No. I-308:

0.88-0.92(3H,m), 0.35-0.37(4H,m), 0.79-1.82(2H,m), 2.03-2.07(2H,m),
2.44-2.49(2H,m), 2.73-2.78(2H,m), 4.01(2H,t,J=7.69), 7.28-7.30(2H,m),
7.43-7.53(3H,m), 16.06(1H,br)

Compound No. I-339:

1.84-2.11(4H,m), 2.44-2.48(2H,m), 2.74-2.78(2H,m), 3.64(3H,s),
3.69-3.92(3H,m), 4.07-4.34(2H,m), 16.04(1H,br)

Compound No. I-462:

5 1.30(3 H,t,J=7.66), 2.03-2.07(2H,m), 2.45-2.49(2H,m), 2.69-2.77(4H,m),
3.68(3H,s), 7.28-7.30(1H,m), 7.77-7.73(1H,m), 8.51(1H,s), 16.03(1H,br)

Physical property values of the production intermediate [13a] and [3b] are given in Table 70 and Table 71.

[Table 70]

Compound No.	Melting Point (°C)
IV-116	111-114
IV-117	100-102
IV-118	118-121
IV-136	131-133
IV-137	102-105
IV-138	122-125
IV-182	107-108
IV-185	50-53
IV-197	122-125
IV-259	84-86
IV-260	107-109
IV-261	132-135
IV-275	102-103
IV-276	46-49
IV-278	171-172
IV-280	137-140
IV-284	136-137
IV-285	112-114
IV-287	140-142
IV-288	101-102
IV-290	124-127
IV-291	137-138

[Table 71]

Compound No.	Melting Point(°C)	Compound No.	Melting Point(°C)
V-1	220-223	V-131	201-204
V-2	165-168	V-135	224-227
V-3	113-115	V-149	216-218
V-4	122-125	V-155	229-231
V-5	98-100	V-167	211-212
V-9	99-102	V-169	199-202
V-10	127-129	V-170	177-180
V-11	82-84	V-179	237-240
V-14	142-144	V-184	158-161
V-16	155-158	V-189	200-201
V-27	114-117	V-202	200-203
V-41	90-91	V-203	164-167
V-43	145-146	V-204	199-202
V-47	144-147	V-268	201-204
V-50	195-198	V-269	155-157
V-51	154-157	V-270	184-187
V-52	118-120	V-271	208-211
V-53	234-236	V-272	100-102
V-54	95-98	V-273	202-205
V-55	95-98	V-275	166-169
V-56	212-215	V-282	193-196
V-57	150-152	V-283	186-189
V-58	196-199	V-291	175-178
V-60	145-146	V-294	204-207
V-61	173-174	V-295	105-107
V-66	164-166	V-296	106-108
V-67	200-203	V-297	176-179
V-68	206-209	V-298	145-146
V-72	213-215	V-299	241-244
V-73	221-224	V-300	245-248
V-87	162-165	V-301	259-261
V-88	227-230	V-302	211-212
V-89	184-186	V-303	152-155
V-90	156-159	V-304	140-143
V-91	179-181	V-305	166-167
V-92	207-210	V-328	143-146
V-93	220-223	V-358	240-243
V-99	166-169	V-359	91-94
V-105	169-171	V-360	240-242
V-106	231-234	V-361	155-158
V-107	166-169	V-362	148-151
V-108	153-156	V-363	189-192
V-109	197-198	V-364	213-216
V-110	194-197	V-365	75-78
V-111	187-190	V-366	218-221
V-115	188-191	V-367	192-195
V-119	205-208	V-368	153-156
V-125	173-175	V-369	111-113
V-127	135-138	V-370	100-103
V-128	186-188	V-371	80-83
V-129	198-201		

Compound number and ¹H-NMR data (standard; TMS, δ (ppm) value) for the production

intermediates are given below. Data without a name of solvent are measured by using CDCl₃.

Compound No. IV-19:

1.19-1.41(3H,m), 1.39(3H,t,J=5.3Hz), 1.56-1.66(3H,m), 1.83-1.87(2H,m),
2.37(2H,dq,J=3.3Hz,12.1Hz), 3.68(3H,s), 4.41(2H,q,J=7.1Hz),
5 4.73(1H,tt,J=3.3Hz,12.1Hz)

Compound No. IV-50:

1.39(3H,t,J=7.1Hz), 3.71(3H,s), 4.43(2H,q,J=7.1Hz), 7.24-7.26(2H,m),
7.49-7.57(3H,m)

Compound No. IV-53:

10 1.39(3H,t,J=5.3Hz), 3.77(3H,s), 4.43(2H,q,J=5.3Hz), 7.18(2H,d,J=6.4Hz),
7.49(2H,d,J=6.4Hz)

Compound No. IV-56:

1.39(3H,t,J=7.1Hz), 3.77(3H,s), 4.43(2H,q,J=7.1Hz), 7.20-7.22(4H,m)

Compound No. IV-59:

15 1.39(3H,t,J=7.1Hz), 2.41(3H,s), 3.77(3H,s), 4.42(2H,q,J=7.1Hz),
7.10(2H,d,J=8.3Hz), 7.31(2H,d,J=8.3Hz)

Compound No. IV-62:

1.39(3H,t,J=7.1Hz), 3.76(3H,s), 3.84(3H,s), 4.43(2H,q,J=7.1Hz),
7.01(2H,d,J=9.0Hz), 7.14(2H,d,J=9.0Hz)

20 Compound No. IV-63:

1.39(3H,t,J=7.1Hz), 3.78(3H,s), 4.43(2H,q,J=7.1Hz), 7.30(1H,d,J=7.7Hz),
7.67(1H,t,J=7.7), 7.74(1H,dt,J=1.1Hz,7.7Hz), 7.84(1H,dd,J=1.1Hz,7.7Hz)

Compound No. IV-64:

25 1.40(3H,t,J=7.1Hz), 3.78(3H,s), 4.44(2H,q,J=7.1Hz), 7.44(1H,d,J=8.0Hz),
7.54(1H,s), 7.66(1H,t,J=8.0Hz), 7.75(1H,d,J=8.0Hz)

Compound No. IV-65:

1.40(3H,t,J=5.3Hz), 3.79(3H,s), 4.44(2H,q,J=5.3Hz), 7.39(2H,d,J=6.2Hz),
7.79(2H,d,J=6.2Hz)

Compound No. IV-71:

1.39(3H,t,J=7.1Hz), 3.78(3H,s), 4.43(2H,q,J=7.1Hz), 7.28(2H,d,J=8.5Hz),
7.36(2H,d,J=8.5Hz)

Compound No. IV-74:

1.39(3H,t,J=7.1Hz), 3.78(3H,s), 4.44(2H,q,J=7.1Hz),
5 7.39(2H,dd,J=1.9Hz,6.6Hz), 7.82(2H,dd,J=1.9Hz,6.6Hz)

Compound No. IV-78:

1.40(3H,t,J=7.1Hz), 3.79(3H,s), 4.43(2H,q,J=7.1Hz), 6.99-7.05(2H,m),
7.22-7.28(1H,m)

Compound No. IV-93:

10 1.39(3H,t,J=7.1Hz), 3.77(3H,s), 3.78(6H,s), 4.43(2H,q,J=7.1Hz),
6.35(2H,d,J=2.2Hz), 6.55(1H,t,J=2.2Hz)

Compound No. IV-96:

1.39(3H,t,J=7.1Hz), 3.76(6H,s), 3.83(3H,s), 4.42(2H,q,J=7.1Hz),
6.55-6.59(2H,m), 7.05(1H,d,J=9.1Hz)

15 Compound No. IV-134:

1.40(3H,t,J=5.3Hz), 3.77(3H,s), 3.79(3H,s), 4.43(2H,q,J=5.3Hz),
6.97(1H,d,J=6.8Hz), 7.17(1H,d,J=2.0Hz), 7.41(1H,dd,J=2.0Hz,6.8Hz)

Compound No. IV-179:

1.39(3H,t,J=5.3Hz), 3.77(3H,s), 4.43(2H,q,J=5.3Hz), 7.32(1H,d,J=5.7Hz),
20 7.46(1H,dd,J=5.7Hz,3.7Hz), 7.92(1H,dt,J=1.1Hz,5.7Hz),
8.68(1H,dt,J=3.7Hz,1.1Hz)

Compound No. IV-198:

1.40(3H,t,J=5.3Hz), 3.78(3H,s), 4.43(2H,q,J=5.3Hz), 7.07-7.12(2H,m),
7.42(1H,dd,J=1.1Hz,4.0Hz)

25 Compound No. IV-259:

1.39(3H,t,J=7.1Hz), 1.43(3H,t,J=7.1Hz), 4.17(2H,q,J=7.1Hz),
4.43(2H,q,J=7.1Hz), 7.21-7.26(2H,m), 7.44-7.55(3H,m)

Compound No. IV-260:

1.39(3H,t,J=7.1Hz), 1.43(6H,d,J=6.8Hz), 4.42(2H,q,J=7.1Hz),

5.01(1H,p,J=6.8Hz), 7.22-7.26(2H,m), 7.46-7.55(3H,m)

Compound No. IV-261:

1.40(3H,t,J=7.1Hz), 4.46(2H,q,J=7.1Hz), 7.23-7.26(2H,m),

7.47(1H,t,J=57.8Hz), 7.51-7.66(3H,m)

5 Compound No. IV-262:

1.39(3H,t,J=7.1Hz), 4.44(2H,q,J=7.1Hz), 7.26-7.60(10H,m)

Compound No. IV-265:

1.40(3H,t,J=7.1Hz), 3.71(3H,s), 4.05(3H,s), 4.44(2H,q,J=7.1Hz)

Compound No. IV-286:

10 1.19-1.17(6H,dd,J=7.0 Hz,J=2.2 Hz), 1.41-1.37(3H,t,J=7.0Hz),

2.65-2.58(1H,sept.,J=7.0Hz), 3.78(3H,s), 4.46-4.39(2H,q,J=7.0Hz),

7.05-7.03(1H,d,J=8.0Hz), 7.33-7.29(1H,m), 7.47-7.46(2H,d,J=4.0 Hz)

Compound No. V-19: (solvent for measurement: DMSO-d₆)

1.09-1.34(3H,m), 1.59-1.64(2H,m), 1.76-1.80(2H,m),

15 2.22(2H,dq,J=3.3Hz,12.3Hz), 3.51(3H,s), 4.54(1H,t,J=3.3Hz,12.3Hz),

13.53(1H,bs)

Compound No. V-50:(solvent for measurement:DMSO-d₆)

3.59(3H,s), 7.29-7.31(2H,m), 7.43-7.54(3H,m), 13.64(1H,bs)

Compound No. V-53:(solvent for measurement:DMSO-d₆)

20 3.59(3H,s), 7.35(2H,dd,J=1.6Hz,5.0Hz), 7.59(2H,dd,J=1.6Hz,5.0Hz),

13.66(1H,bs)

Compound No. V-56:(solvent for measurement:DMSO-d₆)

3.59(3H,s), 7.34-7.37(4H,m), 13.65(1H,bs)

Compound No. V-59:(solvent for measurement:DMSO-d₆)

25 2.36(3H,s), 3.58(3H,s), 7.17(2H,d,J=8.3Hz), 7.30(2H,d,J=8.3Hz),

13.62(1H,bs)

Compound No. V-62:(solvent for measurement:DMSO-d₆)

3.39(3H,s), 3.74(3H,s), 6.93(2H,d,J=9.0), 7.39(2H,d,J=9.0Hz),

9.54(1H,bs)

Compound No. V-63:(solvent for measurement:DMSO-d₆)

3.62(3H,s), 7.64(1H,d,J=7.7Hz), 7.75(1H,t,J=7.68Hz), 7.87-7.94(2H,m),
13.90(1H,bs)

Compound No. V-64:(solvent for measurement:DMSO-d₆)

5 3.41(3H,s), 7.46(1H,d,J=6.0Hz), 7.60(1H,t,J=6.0Hz), 7.82(1H,d,J=6.0Hz),
7.97(1H,s), 9.90(1H,bs)

Compound No. V-65:(solvent for measurement:DMSO-d₆)

3.60(3H,s), 7.58(2H,d,J=8.3Hz), 7.92(2H,d,J=8.3Hz), 13.69(1H,bs) [0259]

Compound No. V-71: (solvent for measurement: DMSO-d₆)

10 3.59(3H,s), 7.47(2H,dt,J=9.3Hz,2.2Hz), 7.54(2H,d,J=9.3Hz), 13.67(1H,bs)

Compound No. V-75:

3.92(3H,s), 7.03-7.06(1H,m), 7.13-7.18(1H,m), 7.35-7.41(1H,m)

Compound No. V-76:

3.92(3H,s), 7.85-7.87(2H,m), 7.00-7.12(1H,m)

15 Compound No. V-77:

3.94(3H,s), 7.07-7.11(1H,m), 7.29-7.31(1H,m), 7.38-7.42(1H,m)

Compound No. V-78:(solvent for measurement:DMSO-d₆)

3.61(3H,s), 7.25-7.31(1H,m), 7.49-7.58(2H,m), 13.79(1H,bs)

Compound No. V-79:

20 3.94(3H,s), 7.05-7.07(1H,m), 7.27-7.32(2H,m)

Compound No. V-80:

3.94(3H,s), 7.12-7.18(2H,m), 7.52-7.61(1H,m)

Compound No. V-81:(solvent for measurement:DMSO-d₆)

3.60(3H,s), 7.37(1H,d,J=8.5Hz), 7.69(1H,s), 7.82(1H,d,J=7.7Hz)

25 Compound No. V-82:

3.92(3H,s), 7.20(2H,s), 7.56(1H,s)

Compound No. V-83:

3.93(3H,s), 7.25(1H,d,J=10.4), 7.44(1H,t,J=8.0), 7.68(1H,d,J=11.7)

Compound No. V-84:

3.93(3H,s), 7.21(1H,d,J=15.6), 7.45-7.48(1H,m), 7.68(1H,d,J=2.4Hz)

Compound No. V-85:

3.93(3H,s), 7.33(1H,d,J=5.7), 7.49-7.58(2H,m)

Compound No. V-86:

5 3.95(3H,s), 7.45-7.56(2H,m)

Compound No. V-93:(solvent for measurement:DMSO-d₆)

3.58(3H,s), 3.74(6H,s), 7.52(2H,d,J=2.2Hz), 6.59(1H,t,J=2.2Hz),

13.63(1H,bs)

Compound No. V-96:(solvent for measurement:DMSO-d₆)

10 3.59(3H,s), 3.73(3H,s), 3.82(3H,s), 7.62(1H,dd,J=2.5Hz,8.8Hz),

6.71(1H,s), 7.16(1H,d,J=8.5Hz), 13.76(1H,bs)

Compound No. V-134: (solvent for measurement: DMSO-d₆)

3.60(3H,s), 3.76(3H,s), 7.23(1H,d,J=9.1Hz), 7.43(1H,d,J=2.8Hz),

7.54(1H,dd,J=2.8Hz,9.1Hz), 13.84(1H,bs)

15 Compound No. V-170:(solvent for measurement:DMSO-d₆)

3.58(3H,s), 6.10(2H,s), 6.78(1H,dd,J=1.0Hz,6.2Hz), 6.89(1H,d,J=1.0Hz),

7.01(1H,d,J=6.2Hz), 13.63(1H,bs)

Compound No. V-179:(solvent for measurement:DMSO-d₆)

3.60(3H,s), 7.49(1H,d,J=7.7Hz), 7.55(1H,ddd,J=1.1Hz,5.0Hz,7.7Hz),

20 8.05(1H,dt,J=1.9Hz,7.7Hz), 8.62(1H,dd,J=1.1Hz,5.0Hz)

Compound No. V-198:(solvent for measurement:DMSO-d₆)

3.57(3H,s), 7.07-7.10(2H,m), 7.63(1H,dd,J=1.9Hz,5.2Hz)

Compound No. V-259:(solvent for measurement:DMSO-d₆)

1.09(3H,t,J=5.3Hz), 3.96(2H,q,J=5.3Hz), 7.32-7.37(2H,m),

25 7.45-7.54(3H,m), 9.51(1H,bs)

Compound No. V-261:(solvent for measurement:DMSO-d₆)

7.36-7.53(5H,m), 7.82(1H,t,J=42.9Hz)

Compound No. V-265:(solvent for measurement:DMSO-d₆)

3.53(3H,s), 3.90(3H,s)

Compound No. V-268:(solvent for measurement:DMSO-d₆)

1.45(3H,t), 3.91(3H,s), 4.09(2H,q), 7.04(2H,d), 7.15(2H,d) [0261]

<Formulation example 1> Wettable powder

10 parts of the compound (I-1), 0.5 parts of polyoxyethylene octylphenyl ether, 0.5 parts of
5 sodium β-naphthalene sulfonate formalin condensate, 20 parts of diatomaceous earth, and 69
parts of clay were mixed and pulverized to give a wettable powder.

<Formulation example 2> Flowable agent

20 parts of roughly crushed compound (I-1) were dispersed in 69 parts of water, and added
with 200 ppm of silicone AF-118N (trade name, manufactured by Asahi Kasei Corporation) while
10 simultaneously adding 4 parts of polyoxyethylene styryl phenyl ether sulfonate and 7 parts of
ethylene glycol. After mixing for 30 minutes by high-speed mixer, the mixture was pulverized
using a wet-type pulverizer to give a flowable agent.

<Formulation example 3> Emulsifiable concentrate

30 parts of the compound (I-1), 60 parts of a mixture of xylene and isophorone (1: 1
15 mixture), and 10 parts of a mixture of polyoxyethylene sorbitan alkylate, polyoxyethylene
alkylaryl polymer, and alkylaryl sulfonate were mixed well to give an emulsifiable concentrate.

<Formulation example 4> Granules

10 parts of the compound (I-1), 80 parts of extender in which talc and bentonite are mixed in
ratio of 1 to 3, 5 parts of white carbon, and 5 parts of a mixture of polyoxyethylene sorbitan
20 alkylate, polyoxyethylene alkylaryl polymer, and alkylaryl sulfonate were added with 10 parts of
water. After kneading well, the resulting paste was extruded through a sieve (diameter; 0.7 mm)
followed by drying. By cutting it to have length of 0.5 to 1 mm, granules were obtained.

Effect of the compounds of the invention is explained by way of following test examples.

<Test example 1> Test for determining herbicidal activity by paddy field soil treatment

25 A 100 cm² wide plastic pot was filled with a paddy field soil and, after watering and
shuffling, seeds of each of *Echinochloa oryzicola*, *Monochoria vaginalis*, and *Scirpus juncoides*
Rocxb. were sowed and watered to a depth of 3 cm. On the next day, the wettable powder
obtained in view of Formulation example 1 was diluted with water and applied on water surface.
The application amount was 1000 g of effective component per hectare. After that, the plants

were cultivated in a greenhouse, and on day 21 after the treatment, evaluation was made by the criteria of Table 72 for determining the herbicidal effects. The results are shown in Table 73 to Table 76.

[Table 72]

5

Index Number	Herbicidal Effects
10	100% of herbicidal effects (complete death)
9	90% or more and less than 100% of herbicidal effects
8	80% or more and less than 90% of herbicidal effects
7	70% or more and less than 80% of herbicidal effects
6	60% or more and less than 70% of herbicidal effects
5	50% or more and less than 60% of herbicidal effects
4	40% or more and less than 50% of herbicidal effects
3	30% or more and less than 40% of herbicidal effects
2	20% or more and less than 30% of herbicidal effects
1	10% or more and less than 20% of herbicidal effects
0	0% or more and less than 10% of herbicidal effects

[Table 73]

Compound No.	<i>Echinochloa oryzicola</i>	Compound No.	<i>Echinochloa oryzicola</i>	Compound No.	<i>Echinochloa oryzicola</i>	Compound No.	<i>Echinochloa oryzicola</i>
I-1	10	I-83	10	I-187	10	I-328	10
I-2	10	I-84	10	I-198	8	I-339	10
I-3	10	I-85	10	I-199	9	I-463	10
I-4	9	I-86	10	I-202	10	I-464	10
I-5	10	I-87	9	I-203	10	I-465	10
I-9	8	I-88	10	I-205	7	I-466	8
I-10	10	I-89	10	I-259	10	I-468	10
I-11	10	I-90	9	I-260	10	I-469	10
I-14	10	I-91	10	I-261	8	I-470	10
I-16	9	I-92	10	I-263	10	I-471	10
I-19	10	I-93	8	I-265	10	I-473	8
I-27	10	I-96	8	I-268	10	I-474	10
I-41	8	I-99	10	I-269	8	I-475	10
I-43	10	I-105	10	I-270	8	I-476	10
I-50	10	I-106	10	I-271	8	I-477	10
I-51	10	I-107	10	I-272	7	I-478	10
I-52	10	I-108	10	I-273	9	I-479	10
I-53	10	I-109	10	I-274	8	I-480	10
I-54	10	I-110	10	I-275	9	III-50	10
I-55	10	I-111	10	I-276	8	III-62	8
I-56	10	I-115	10	I-277	9	VI-1	10
I-57	10	I-116	10	I-278	8	VI-5	10
I-58	10	I-117	10	I-279	8	VI-6	10
I-59	10	I-118	10	I-280	10	VI-7	10
I-60	10	I-119	9	I-281	10	VI-65	10
I-61	8	I-120	8	I-282	10	VI-97	10
I-63	10	I-125	10	I-283	10	V-300	10
I-64	10	I-126	10	I-284	10	V-358	10
I-65	10	I-127	10	I-285	10	V-359	8
I-66	10	I-128	8	I-286	10	V-362	10
I-67	10	I-129	10	I-287	10	V-363	10
I-68	10	I-131	9	I-288	10	V-364	10
I-71	10	I-134	10	I-289	10	V-365	10
I-72	10	I-135	10	I-292	10	V-367	8
I-73	10	I-136	9	I-294	9	V-368	10
I-74	10	I-137	10	I-297	10	V-369	10
I-75	10	I-138	10	I-298	10	V-370	10
I-76	10	I-149	9	I-299	10	V-371	10
I-77	10	I-155	10	I-300	10		
I-78	10	I-169	10	I-301	10		
I-79	10	I-170	10	I-302	10		
I-80	10	I-179	10	I-303	10		
I-81	10	I-184	10	I-304	10		
I-82	8	I-185	8	I-307	8		

[Table 74]

Compound No.	<i>Monochoria vaginalis</i>	Compound No.	<i>Monochoria vaginalis</i>	Compound No.	<i>Monochoria vaginalis</i>
I-1	10	I-82	10	I-183	10
I-2	10	I-83	10	I-184	10
I-3	10	I-84	10	I-185	9
I-4	9	I-85	10	I-187	10
I-5	10	I-86	10	I-189	10
I-9	8	I-87	10	I-198	10
I-10	10	I-88	10	I-199	8
I-11	10	I-89	10	I-202	10
I-14	10	I-90	10	I-203	10
I-16	9	I-91	10	I-204	8
I-19	10	I-92	10	I-205	10
I-27	10	I-93	10	I-259	10
I-41	8	I-94	10	I-260	10
I-43	10	I-96	10	I-261	10
I-47	10	I-99	10	I-262	8
I-50	10	I-105	10	I-263	10
I-51	10	I-106	10	I-265	10
I-52	10	I-107	10	I-268	10
I-53	7	I-108	10	I-269	8
I-54	10	I-109	10	I-270	8
I-55	10	I-110	10	I-271	8
I-56	10	I-111	10	I-272	8
I-57	10	I-115	10	I-273	9
I-58	10	I-116	10	I-274	8
I-59	10	I-117	10	I-275	10
I-60	10	I-118	10	I-276	8
I-61	10	I-119	10	I-277	9
I-62	8	I-120	10	I-278	9
I-63	10	I-125	10	I-279	8
I-64	10	I-126	10	I-280	10
I-65	10	I-127	10	I-281	10
I-66	10	I-128	9	I-282	10
I-67	10	I-129	10	I-283	10
I-68	10	I-131	10	I-284	10
I-71	10	I-134	10	I-285	10
I-72	10	I-135	10	I-286	10
I-73	10	I-136	10	I-287	10
I-74	10	I-137	10	I-288	10
I-75	10	I-138	10	I-289	10
I-76	10	I-149	10	I-290	10
I-77	10	I-155	10	I-291	10
I-78	10	I-169	10	I-292	10
I-79	10	I-170	10	I-293	10
I-80	10	I-179	10	I-294	9
I-81	10	I-182	8	I-297	10

[Table 75]

Compound No.	<i>Monochoria vaginalis</i>	Compound No.	<i>Monochoria vaginalis</i>
I-298	10	V-361	10
I-299	10	V-362	10
I-300	10	V-363	10
I-301	10	V-364	10
I-302	10	V-365	10
I-303	10	V-366	10
I-304	10	V-367	10
I-306	9	V-368	10
I-307	9	V-369	10
I-308	8	V-370	10
I-328	10	V-371	10
I-339	10		
I-462	10		
I-463	10		
I-464	10		
I-465	10		
I-466	10		
I-467	10		
I-468	10		
I-469	10		
I-470	10		
I-471	10		
I-472	10		
I-473	10		
I-474	10		
I-475	10		
I-476	10		
I-477	10		
I-478	10		
I-479	10		
II-50	8		
II-267	8		
III-50	10		
III-62	10		
VI-1	10		
VI-5	10		
VI-6	10		
VI-7	10		
VI-65	10		
VI-97	10		
V-291	8		
V-300	10		
V-358	10		
V-359	10		
V-360	10		

[Table 76]

Compound No.	<i>S. juncoides</i> Rocxb.	Compound No.	<i>S. juncoides</i> Rocxb.	Compound No.	<i>S. juncoides</i> Rocxb.	Compound No.	<i>S. juncoides</i> Rocxb.
I-1	10	I-84	10	I-187	10	I-304	10
I-2	10	I-85	10	I-189	10	I-307	8
I-3	10	I-86	10	I-198	10	I-328	10
I-4	10	I-87	10	I-199	9	I-339	10
I-5	10	I-88	10	I-202	10	I-462	10
I-9	8	I-89	10	I-203	10	I-463	10
I-10	10	I-90	10	I-205	10	I-464	10
I-11	10	I-91	10	I-259	10	I-465	10
I-14	10	I-92	10	I-260	10	I-466	10
I-16	10	I-93	10	I-261	8	I-467	10
I-19	10	I-94	10	I-263	10	I-468	10
I-27	10	I-96	10	I-265	10	I-469	10
I-41	10	I-99	10	I-268	10	I-470	10
I-43	10	I-105	10	I-269	10	I-471	10
I-47	10	I-106	10	I-270	8	I-472	8
I-50	10	I-107	10	I-271	10	I-473	8
I-51	10	I-108	10	I-272	8	I-474	10
I-52	10	I-109	10	I-273	10	I-475	10
I-53	10	I-110	10	I-274	4	I-476	10
I-54	10	I-111	10	I-275	8	I-477	10
I-55	10	I-115	10	I-276	9	I-478	9
I-56	10	I-116	10	I-277	10	I-479	10
I-57	10	I-117	10	I-278	10	I-480	10
I-58	10	I-118	10	I-279	10	II-50	7
I-59	10	I-119	10	I-280	10	III-50	10
I-60	10	I-120	10	I-281	10	III-62	10
I-61	10	I-125	10	I-282	10	VI-1	10
I-63	10	I-126	10	I-283	10	VI-5	10
I-64	10	I-127	10	I-284	10	VI-6	10
I-65	10	I-128	10	I-285	10	VI-7	10
I-66	10	I-129	10	I-286	9	VI-65	10
I-67	10	I-131	10	I-287	10	VI-97	10
I-68	10	I-134	10	I-288	10	V-300	10
I-71	10	I-135	10	I-289	10	V-358	10
I-72	10	I-136	10	I-290	10	V-359	10
I-73	10	I-137	10	I-291	10	V-360	10
I-74	10	I-138	10	I-292	10	V-361	10
I-75	10	I-149	10	I-293	10	V-362	10
I-76	10	I-155	10	I-294	9	V-363	10
I-77	10	I-169	10	I-297	10	V-364	10
I-78	10	I-170	10	I-298	10	V-365	10
I-79	10	I-179	10	I-299	10	V-366	8
I-80	10	I-182	9	I-300	10	V-367	8
I-81	10	I-183	10	I-301	10	V-368	10
I-82	10	I-184	10	I-302	10	V-369	10
I-83	10	I-185	9	I-303	10	V-370	10
						V-371	9

<Test example 2> Test for determining herbicidal activity by field soil treatment

A 80 cm² wide plastic pot was filled with a field soil and seeds of each of *Echinochloa*

5 *crus-galli*, foxtail, Indian millet, and *A. retroflexus* were sowed and then covered with soil. The

wettable powder produced with reference to the Formulation example 1 was diluted water, and applied on the soil surface by using a small sprayer in an amount of 1000 liters per hectare so that the effective component is 1000 g per hectare. After that, the plants were cultivated in a greenhouse, and on day 21 after the treatment, evaluation was made by the criteria described in

5 Table 72 for determining the herbicidal effects. The results are shown in Table 77 to Table 80.

[Table 77]

Compound No.	<i>Echinochloa crus-galli</i>	Compound No.	<i>Echinochloa crus-galli</i>	Compound No.	<i>Echinochloa crus-galli</i>
I-1	8	I-92	9	I-282	10
I-2	10	I-93	7	I-283	9
I-3	10	I-98	7	I-284	10
I-4	9	I-99	8	I-285	10
I-5	10	I-105	9	I-286	10
I-9	7	I-106	10	I-287	10
I-10	10	I-107	8	I-288	10
I-11	10	I-109	9	I-289	9
I-14	10	I-110	7	I-292	7
I-16	9	I-111	9	I-294	9
I-19	8	I-115	9	I-297	10
I-27	10	I-116	10	I-298	7
I-41	9	I-117	10	I-299	9
I-43	10	I-118	10	I-302	7
I-50	10	I-119	8	I-303	9
I-51	10	I-120	8	I-304	10
I-52	10	I-125	7	I-307	7
I-53	8	I-127	10	I-339	8
I-54	10	I-128	8	I-471	7
I-55	10	I-129	9	I-474	7
I-56	10	I-131	9	I-475	7
I-57	10	I-134	10	I-476	7
I-58	10	I-135	9	I-477	9
I-60	10	I-137	10	I-478	9
I-61	8	I-138	9	I-479	9
I-63	10	I-149	8	I-480	8
I-64	10	I-167	8	VI-5	8
I-65	10	I-169	10	VI-7	10
I-66	10	I-179	10	V-300	7
I-67	10	I-182	7	V-365	7
I-68	10	I-184	8	V-368	7
I-71	10	I-185	9	V-369	7
I-72	9	I-187	7	V-370	7
I-73	9	I-198	7	V-371	9
I-74	10	I-199	9		
I-75	9	I-202	10		
I-76	10	I-203	9		
I-77	9	I-259	10		
I-78	10	I-260	10		
I-79	9	I-265	10		
I-80	9	I-269	8		
I-81	9	I-270	8		
I-82	9	I-271	10		
I-83	9	I-273	9		
I-84	9	I-274	7		
I-85	9	I-275	8		
I-86	10	I-276	9		
I-87	9	I-277	8		
I-88	8	I-278	9		
I-89	9	I-279	7		
I-90	8	I-280	9		
I-91	9	I-281	9		

[Table 78]

Compound No.	<i>Setaria viridis</i>	Compound No.	<i>Setaria viridis</i>	Compound No.	<i>Setaria viridis</i>
I-1	7	I-84	9	I-280	8
I-2	7	I-85	9	I-281	9
I-3	10	I-86	9	I-282	10
I-4	9	I-87	6	I-283	8
I-5	7	I-89	8	I-284	8
I-10	10	I-91	10	I-285	10
I-11	7	I-92	9	I-286	9
I-14	10	I-93	6	I-288	9
I-16	9	I-98	7	I-289	7
I-19	8	I-99	6	I-294	7
I-41	7	I-105	7	I-297	9
I-50	10	I-109	6	I-298	7
I-51	10	I-111	7	I-299	10
I-52	10	I-116	9	I-303	9
I-54	10	I-117	7	I-304	9
I-55	10	I-118	9	VI-7	10
I-56	10	I-127	8	VI-65	7
I-57	10	I-128	9		
I-58	8	I-129	10		
I-63	10	I-131	7		
I-66	10	I-134	10		
I-67	10	I-136	8		
I-68	10	I-137	9		
I-71	6	I-155	7		
I-72	10	I-169	10		
I-73	8	I-179	10		
I-74	7	I-202	9		
I-75	7	I-260	5		
I-76	9	I-265	10		
I-77	9	I-269	9		
I-79	10	I-270	7		
I-80	9	I-271	10		
I-81	7	I-276	9		
I-82	9	I-277	8		
I-83	9	I-278	9		

[Table 79]

Compound No.	<i>Abutilon theophrasti</i>	Compound No.	<i>Abutilon theophrasti</i>	Compound No.	<i>Abutilon theophrasti</i>	Compound No.	<i>Abutilon theophrasti</i>
I-1	9	I-83	10	I-167	10	I-294	9
I-2	10	I-84	10	I-169	9	I-297	9
I-3	10	I-85	10	I-170	10	I-298	10
I-4	9	I-86	9	I-179	10	I-299	10
I-5	10	I-87	9	I-182	10	I-300	10
I-10	10	I-88	10	I-183	10	I-302	10
I-11	9	I-89	10	I-184	10	I-303	9
I-14	10	I-90	10	I-185	10	I-304	10
I-16	9	I-91	10	I-187	9	I-306	7
I-27	10	I-92	10	I-189	10	I-307	9
I-41	8	I-93	10	I-198	10	I-339	10
I-50	10	I-94	10	I-199	10	I-462	10
I-51	10	I-96	10	I-202	9	I-463	10
I-52	10	I-98	7	I-259	8	I-465	10
I-53	10	I-99	9	I-260	10	I-470	7
I-54	10	I-105	9	I-261	10	I-471	10
I-55	10	I-106	10	I-263	8	I-474	10
I-56	10	I-107	10	I-265	10	I-475	7
I-57	10	I-108	8	I-268	8	I-476	10
I-58	10	I-109	10	I-269	9	I-477	10
I-59	10	I-110	10	I-271	10	I-478	10
I-60	10	I-111	9	I-273	7	I-479	10
I-61	10	I-115	9	I-274	8	I-480	10
I-63	10	I-116	10	I-275	10	VI-1	10
I-64	10	I-117	10	I-276	10	VI-5	10
I-65	10	I-118	9	I-277	7	VI-6	9
I-66	10	I-119	9	I-279	10	VI-7	10
I-67	10	I-120	9	I-280	9	VI-65	10
I-68	10	I-125	8	I-281	9	V-61	8
I-71	10	I-126	10	I-282	9	V-300	9
I-72	10	I-127	10	I-283	9	V-358	10
I-73	10	I-128	10	I-284	9	V-361	10
I-74	10	I-129	10	I-285	10	V-364	7
I-75	10	I-131	9	I-286	9	V-365	10
I-76	10	I-134	10	I-287	9	V-368	10
I-77	10	I-135	9	I-288	9	V-369	7
I-78	10	I-136	9	I-289	9	V-370	10
I-79	10	I-137	10	I-290	10	V-371	10
I-80	9	I-138	9	I-291	10		
I-81	10	I-149	10	I-292	9		
I-82	9	I-155	10	I-293	10		

[Table 80]

Compound No.	<i>Amaranthus retroflexus</i>	Compound No.	<i>Amaranthus retroflexus</i>	Compound No.	<i>Amaranthus retroflexus</i>	Compound No.	<i>Amaranthus retroflexus</i>
I-1	10	I-82	10	I-179	9	I-302	10
I-2	10	I-83	10	I-182	10	I-303	10
I-3	10	I-84	10	I-183	8	I-304	10
I-4	10	I-85	10	I-184	10	I-306	10
I-5	10	I-86	10	I-185	10	I-307	10
I-9	10	I-87	9	I-187	10	I-308	10
I-10	10	I-88	10	I-189	10	I-339	10
I-11	10	I-89	10	I-198	10	I-462	9
I-14	10	I-90	10	I-199	10	I-463	7
I-16	10	I-91	10	I-202	10	I-464	7
I-19	8	I-92	10	I-203	7	I-465	10
I-27	10	I-93	10	I-259	10	I-468	7
I-41	10	I-94	10	I-260	10	I-470	8
I-43	10	I-96	10	I-263	10	I-471	10
I-47	10	I-99	10	I-265	10	I-474	10
I-50	10	I-105	10	I-268	10	I-475	7
I-51	10	I-106	10	I-269	10	I-476	10
I-52	10	I-107	10	I-270	10	I-477	10
I-53	10	I-108	10	I-271	10	I-478	10
I-54	10	I-109	10	I-272	8	I-479	10
I-55	10	I-110	10	I-273	10	I-480	10
I-56	9	I-111	10	I-274	7	VI-1	10
I-57	10	I-115	9	I-275	10	VI-5	10
I-58	10	I-116	10	I-276	10	VI-6	10
I-59	10	I-117	10	I-277	8	VI-7	10
I-60	10	I-118	10	I-278	10	VI-65	10
I-61	10	I-119	10	I-279	10	V-300	10
I-63	10	I-120	10	I-280	10	V-358	10
I-64	10	I-125	10	I-281	10	V-361	8
I-65	10	I-126	10	I-282	10	V-362	7
I-66	10	I-127	10	I-283	10	V-364	8
I-67	10	I-128	10	I-284	10	V-365	10
I-68	10	I-129	10	I-285	10	V-368	10
I-71	10	I-131	10	I-286	10	V-369	7
I-72	10	I-134	10	I-287	10	V-370	10
I-73	10	I-135	10	I-288	10	V-371	10
I-74	10	I-136	9	I-289	10		
I-75	10	I-137	10	I-290	8		
I-76	10	I-138	10	I-291	8		
I-77	10	I-149	10	I-294	10		
I-78	10	I-155	10	I-297	10		
I-79	10	I-167	10	I-298	10		
I-80	10	I-169	10	I-299	10		
I-81	10	I-170	10	I-300	10		

<Test example 3> Test for determining herbicidal activity by field foliage treatment

A 80 cm² wide plastic pot was filled with a field soil and seeds of each of Indian millet and

A. retroflexus were sowed and then incubated for 2 weeks in a green house. The wettable powder produced with reference to the Formulation example 1 was diluted water, and applied from the air to entire body of the plant as foliage treatment by using a small sprayer in an amount of 1000 liters per hectare so that the effective component is 1000 g per hectare. After that, the
5 plants were cultivated in a greenhouse, and on day 14 after the treatment, evaluation was made by the criteria described in Table 72 for determining the herbicidal effects. The results are shown in Table 81 to Table 84.

[Table 81]

Compound No.	<i>Echinochloa crus-galli</i>	Compound No.	<i>Echinochloa crus-galli</i>	Compound No.	<i>Echinochloa crus-galli</i>	Compound No.	<i>Echinochloa crus-galli</i>
I-1	8	I-81	10	I-169	9	I-300	9
I-2	9	I-82	10	I-170	10	I-302	10
I-3	9	I-83	10	I-179	10	I-303	8
I-4	9	I-84	9	I-182	8	I-304	10
I-5	10	I-85	9	I-184	10	I-328	7
I-9	10	I-86	9	I-185	10	I-339	9
I-10	8	I-87	9	I-187	8	I-463	7
I-11	9	I-88	8	I-198	10	I-465	8
I-14	9	I-89	9	I-199	10	I-467	8
I-16	9	I-90	8	I-202	9	I-468	9
I-19	10	I-91	9	I-203	6	I-469	10
I-27	9	I-92	10	I-259	10	I-470	8
I-41	10	I-93	9	I-260	8	I-471	9
I-43	8	I-96	7	I-263	9	I-474	7
I-50	10	I-98	7	I-265	8	I-475	9
I-51	10	I-99	9	I-268	7	I-476	7
I-52	10	I-105	10	I-269	10	I-477	9
I-53	8	I-106	10	I-270	9	I-478	8
I-54	10	I-107	7	I-271	10	I-479	9
I-55	10	I-109	10	I-272	6	I-480	9
I-56	10	I-110	9	I-273	10	III-50	10
I-57	10	I-111	10	I-274	9	VI-1	10
I-58	10	I-115	10	I-275	9	VI-5	10
I-59	7	I-116	9	I-276	10	VI-6	8
I-60	10	I-117	9	I-277	10	VI-7	10
I-61	9	I-118	9	I-278	10	VI-65	9
I-63	10	I-119	9	I-279	9	VI-97	7
I-64	10	I-120	9	I-280	9	V-300	8
I-65	8	I-125	10	I-281	9	V-358	8
I-66	10	I-126	9	I-282	8	V-360	8
I-67	10	I-127	10	I-283	8	V-362	9
I-68	10	I-128	9	I-284	9	V-363	10
I-71	10	I-129	10	I-285	9	V-364	8
I-72	10	I-131	10	I-286	10	V-365	9
I-73	10	I-134	10	I-287	8	V-368	7
I-74	9	I-135	10	I-288	9	V-369	9
I-75	10	I-136	9	I-289	9	V-370	7
I-76	10	I-137	10	I-292	8	V-371	8
I-77	10	I-138	8	I-294	9		
I-78	10	I-149	8	I-297	9		
I-79	10	I-155	9	I-298	10		
I-80	10	I-167	10	I-299	8		

[Table 82]

Compound No.	<i>Setaria viridis</i>	Compound No.	<i>Setaria viridis</i>	Compound No.	<i>Setaria viridis</i>
I-1	8	I-86	9	I-285	9
I-2	10	I-89	10	I-286	10
I-3	9	I-90	7	I-288	10
I-4	9	I-91	10	I-289	7
I-5	10	I-92	10	I-297	9
I-10	7	I-93	9	I-298	8
I-11	9	I-105	7	I-302	10
I-14	10	I-109	7	I-303	10
I-16	9	I-116	9	I-304	10
I-19	10	I-117	9	I-328	7
I-27	6	I-118	10	I-463	7
I-41	10	I-126	7	I-464	7
I-43	8	I-127	10	I-465	10
I-50	10	I-128	10	I-468	9
I-51	10	I-129	9	I-469	10
I-52	10	I-134	10	I-470	9
I-54	10	I-136	9	I-471	10
I-55	10	I-137	10	I-475	7
I-56	10	I-138	8	I-479	7
I-57	10	I-155	7	VI-1	10
I-58	10	I-167	8	VI-5	10
I-60	9	I-169	9	VI-6	10
I-63	10	I-179	10	VI-7	10
I-66	10	I-184	8	VI-65	7
I-67	10	I-185	9	VI-97	10
I-68	8	I-187	8	V-300	8
I-71	9	I-199	8	V-358	10
I-72	10	I-202	9	V-362	9
I-73	10	I-261	7	V-363	10
I-74	9	I-263	9	V-364	9
I-75	10	I-265	9	V-365	10
I-76	9	I-269	8	V-369	7
I-77	10	I-271	7		
I-78	10	I-274	8		
I-79	10	I-275	8		
I-80	7	I-276	10		
I-81	10	I-277	9		
I-82	10	I-278	10		
I-83	10	I-281	7		
I-84	10	I-282	7		
I-85	10	I-284	7		

[Table 83]

Compound No.	<i>Abutilon theophrasti</i>	Compound No.	<i>Abutilon theophrasti</i>	Compound No.	<i>Abutilon theophrasti</i>	Compound No.	<i>Abutilon theophrasti</i>
I-1	9	I-85	10	I-189	9	I-307	10
I-2	10	I-86	10	I-198	10	I-308	8
I-3	9	I-87	9	I-199	9	I-328	10
I-4	9	I-88	9	I-202	9	I-339	10
I-5	10	I-89	10	I-203	9	I-462	10
I-9	10	I-90	9	I-205	10	I-463	9
I-10	9	I-91	10	I-259	10	I-464	10
I-11	9	I-92	10	I-260	10	I-465	10
I-14	10	I-93	10	I-261	10	I-466	10
I-16	9	I-94	10	I-263	10	I-467	7
I-19	10	I-96	9	I-265	9	I-468	10
I-27	9	I-98	9	I-268	9	I-469	10
I-41	10	I-99	10	I-269	9	I-470	10
I-43	9	I-105	10	I-270	9	I-471	10
I-47	9	I-106	9	I-271	9	I-472	7
I-50	10	I-107	9	I-272	9	I-473	10
I-51	10	I-108	9	I-273	9	I-474	9
I-52	10	I-109	9	I-274	8	I-475	8
I-53	10	I-110	9	I-275	9	I-476	9
I-54	10	I-111	10	I-276	9	I-477	9
I-55	10	I-115	10	I-277	9	I-478	9
I-56	10	I-116	9	I-278	9	I-479	9
I-57	10	I-117	9	I-279	10	I-480	10
I-58	10	I-118	10	I-280	9	II-50	8
I-59	10	I-119	9	I-281	9	II-267	9
I-60	10	I-120	9	I-282	9	III-50	10
I-61	10	I-125	10	I-283	8	III-62	10
I-62	9	I-126	9	I-284	8	VI-1	10
I-63	10	I-127	9	I-285	7	VI-5	10
I-64	10	I-128	10	I-286	9	VI-6	10
I-65	10	I-129	9	I-287	9	VI-7	10
I-66	10	I-131	10	I-288	10	VI-65	10
I-67	10	I-134	10	I-289	9	VI-97	10
I-68	10	I-135	10	I-290	9	V-300	9
I-71	9	I-136	9	I-291	9	V-358	10
I-72	10	I-137	10	I-292	10	V-359	10
I-73	10	I-138	9	I-293	8	V-360	7
I-74	9	I-149	10	I-294	7	V-361	9
I-75	10	I-155	10	I-295	9	V-362	10
I-76	10	I-167	10	I-297	8	V-363	10
I-77	10	I-169	9	I-298	9	V-364	10
I-78	10	I-170	10	I-299	9	V-365	10
I-79	10	I-179	10	I-300	8	V-366	7
I-80	9	I-182	9	I-301	7	V-367	10
I-81	10	I-183	9	I-302	10	V-368	9
I-82	10	I-184	9	I-303	9	V-369	8
I-83	10	I-185	9	I-304	9	V-370	9
I-84	10	I-187	9	I-306	9	V-371	9


[Table 84]

Compound No.	<i>Amaranthus retroflexus</i>	Compound No.	<i>Amaranthus retroflexus</i>	Compound No.	<i>Amaranthus retroflexus</i>	Compound No.	<i>Amaranthus retroflexus</i>
I-1	10	I-85	10	I-189	10	I-306	7
I-2	10	I-86	10	I-198	10	I-307	9
I-3	9	I-87	10	I-199	9	I-328	10
I-4	9	I-88	10	I-202	9	I-339	10
I-5	10	I-89	10	I-203	10	I-462	10
I-9	10	I-90	10	I-204	7	I-463	10
I-10	10	I-91	10	I-205	10	I-464	10
I-11	10	I-92	10	I-259	10	I-465	10
I-14	10	I-93	10	I-260	10	I-466	10
I-16	10	I-94	10	I-261	10	I-467	10
I-19	10	I-96	10	I-263	10	I-468	10
I-27	9	I-98	8	I-265	10	I-469	10
I-41	10	I-99	10	I-268	10	I-470	10
I-43	10	I-105	10	I-269	10	I-471	10
I-47	10	I-106	10	I-270	10	I-472	10
I-50	10	I-107	10	I-271	10	I-473	3
I-51	10	I-108	10	I-272	10	I-474	9
I-52	10	I-109	9	I-273	10	I-475	9
I-53	10	I-110	9	I-274	9	I-476	10
I-54	10	I-111	10	I-275	9	I-477	10
I-55	10	I-115	10	I-276	10	I-478	10
I-56	10	I-116	8	I-277	10	I-479	10
I-57	10	I-117	8	I-278	10	I-480	10
I-58	10	I-118	10	I-279	10	II-50	10
I-59	10	I-119	10	I-280	10	III-50	10
I-60	10	I-120	10	I-281	9	III-62	10
I-61	10	I-125	10	I-282	8	VI-1	10
I-62	10	I-126	10	I-283	8	VI-5	10
I-63	10	I-127	10	I-284	9	VI-6	10
I-64	10	I-128	10	I-285	8	VI-7	10
I-65	10	I-129	10	I-286	10	VI-65	10
I-66	10	I-131	10	I-287	10	VI-97	10
I-67	10	I-134	10	I-288	10	V-300	10
I-68	10	I-135	10	I-289	10	V-358	10
I-71	10	I-136	10	I-290	10	V-359	10
I-72	10	I-137	10	I-291	10	V-360	10
I-73	10	I-138	10	I-292	10	V-361	10
I-74	10	I-149	10	I-293	10	V-362	10
I-75	10	I-155	10	I-294	8	V-363	10
I-76	10	I-167	10	I-295	8	V-364	10
I-77	10	I-169	8	I-297	8	V-365	10
I-78	10	I-170	10	I-298	10	V-366	10
I-79	10	I-179	10	I-299	10	V-368	9
I-80	10	I-182	9	I-300	10	V-369	9
I-81	10	I-183	10	I-301	10	V-370	10
I-82	10	I-184	10	I-302	10	V-371	10
I-83	10	I-185	10	I-303	10		
I-84	10	I-187	10	I-304	10		

As a result of the tests, it was found that the compounds of the invention have an excellent herbicidal activity.

Claims

1. A triazine derivative or a salt thereof represented by following Formula 1:



[in the formula, R¹ represents a hydrogen atom; a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₃-C₆ halocycloalkyl C₁-C₆ alkyl group; an amino C₁-C₆ alkyl group; a nitro C₁-C₆ alkyl group; a C₁-C₆ alkylamino C₁-C₆ alkyl group; a di(C₁-C₆ alkyl)amino C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ haloalkylthio C₁-C₆ alkyl group; a C₁-C₆ haloalkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ haloalkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a hydroxy C₁-C₆ alkyl group; a phenyl C₁-C₆ alkoxy C₁-C₆ alkyl group (phenyl in the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α); a C₁-C₆ alkoxy C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyloxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyloxy C₁-C₆ alkyl group; a phenyloxy C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylthio C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfinyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfonyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a C₁-C₆ haloalkoxy C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected

from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a phenoxyimino C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a di(C₁-C₆ alkoxy)C₁-C₆ alkyl group; a (R³¹R³²N-C=O)C₁-C₆ alkyl group; a C₁-C₆ alkoxy carbonyl C₁-C₆ alkyl group; a C₁-C₆ alkyl carbonyl C₁-C₆ alkyl group; a C₁-C₆ alkyl carbonyloxy C₁-C₆ alkyl group; a C₁-C₆ alkylidene aminooxy C₁-C₆ alkyl group; a formyl C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkoxy C₁-C₆ alkyl group; a cyano C₁-C₆ alkoxy C₁-C₆ alkyl group; a cyano C₁-C₆ alkyl group; a C₂-C₆ alkylidene amino group; a di(C₁-C₁₀ alkyl)amino C₁-C₆ alkylidene amino group; a NR³¹R³² group; a C₁-C₆ alkoxy group; a C₂-C₆ alkenyloxy group; a C₂-C₆ alkynyloxy group; a C₃-C₆ cycloalkyloxy group; a C₃-C₆ cycloalkyl C₁-C₆ alkyloxy group; a C₁-C₆ haloalkoxy group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone;


a C₁-C₆ alkyl group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α]; a C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α]; or a C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic-oxy group in which the heterocyclic group in the heterocyclic-oxy group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents

selected from the Substituent group α];


R^2 represents a C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_2 - C_6 haloalkynyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_3 - C_6 cycloalkyloxy C_1 - C_6 alkyl group; a di(C_1 - C_6 alkoxy) C_1 - C_6 alkyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_1 - C_6 alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_2 - C_6 alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a phenyl C_2 - C_6 alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ,

Y and Z represent an oxygen atom or a sulfur atom,


A represents any one of the following formula A-1 to A-5,




A-1




A-2



A-3



A-4




A-5

R^4 represents a hydroxyl group; O^-M^+ (M^+ represents an alkali metal cation or an ammonium cation); an amino group; a halogen atom; a cyano group; an isothiocyanate group; an isocyanate group; a hydroxycarbonyloxy group; a C_1 - C_6 alkoxy carbonyloxy group; a benzyloxycarbonyloxy group which may be substituted with a substituent group selected from Substituent group α ; a C_1 - C_6 alkoxy group; a C_2 - C_6 alkenyloxy group; a C_2 - C_6 alkynyloxy group; a C_3 - C_6 cycloalkyloxy group; a cyanomethylene oxy group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy group; a C_1 - C_6 alkyl carbonyloxy group; a C_1 - C_6 haloalkyl carbonyloxy group; a C_2 - C_6


alkenylcarbonyloxy group; a C₂-C₆ haloalkenylcarbonyloxy group; a C₂-C₆ alkynylcarbonyloxy group; a C₂-C₆ haloalkynylcarbonyloxy group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkoxy group; a phenoxy group which may be substituted with one or more substituents selected from the Substituent group α ; a benzyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a phenylcarbonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylcarbonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a phenylcarbonyl C₁-C₆ alkyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₁₀ alkylsulfonyloxy group; a C₁-C₆ haloalkylsulfonyloxy group; a phenylsulfonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyloxy group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₁₀ alkylthio group; a C₁-C₁₀ alkylsulfinyl group; a C₁-C₁₀ alkylsulfonyl group; a C₁-C₆ haloalkylthio group; a C₁-C₆ haloalkylsulfinyl group; a C₁-C₆ haloalkylsulfonyl group; a C₂-C₆ alkenylthio group; a C₂-C₆ alkenylsulfinyl group; a C₂-C₆ alkenylsulfonyl group; a C₂-C₆ alkynylthio group; a C₂-C₆ alkynylsulfinyl group; a C₂-C₆ alkynylsulfonyl group; a phenylthio group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylthio group which may be substituted with one or more substituents selected from the Substituent group α ; a phenylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₁₀ alkylamino group; a di(C₁-C₁₀ alkyl)amino group; a C₁-C₆ alkoxycarbonylamino group; a C₁-C₆ alkoxy group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to

5 identical or different substituents selected from the Substituent group α); or a heterocyclic-oxy group in which the heterocyclic group in the heterocyclic-oxy group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α),


A₁ represents a group represented by the following formula



A₂ represents a group represented by the following formula



A₃ represents a group represented by the following formula



n represents 0, 1, or 2,

R⁵, R⁶, R⁸, R⁹, R³⁵ and R³⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, wherein, R⁵ and R⁸ may be joined together to form a C₂-C₅ alkylene chain or a C₂-C₅ alkenylene chain, and may form a ring together with adjacent carbon atoms, and R⁵ and R³⁵ may be joined together to form a C₁-C₅ alkylene chain to form a ring with adjacent carbon atoms,

R⁷, R³³, and R³⁴ each independently represent a hydrogen atom, a C₁-C₆ alkyl group, a

C₁-C₆ haloalkyl group, a C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, or a C₁-C₆ alkoxy group,

R¹⁴, R¹⁵, R¹⁶, and R¹⁷ each independently represent a hydrogen atom, a C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, or a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ,

R¹⁸ represents a hydrogen atom, a C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a cyanomethyl group, or a benzyl group,

R²⁰ represents a C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, or a C₃-C₆ cycloalkyl C₁-C₆ alkyl group,

R²¹ represents a hydrogen atom, a C₁-C₆ alkyl group, or a halogen atom,

R²³ represents a C₁-C₆ alkyl group, a C₁-C₆ haloalkyl group, a C₃-C₆ cycloalkyl group, a C₁-C₁₀ alkylthio group, a C₁-C₁₀ alkylsulfinyl group, a C₁-C₁₀ alkylsulfonyl group, a phenylthio group which may be substituted with one or more substituents selected from the Substituent group α , a benzylthio group which may be substituted with one or more substituents selected from the Substituent group α , a phenylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α , a benzylsulfinyl group which may be substituted with one or more substituents selected from the Substituent group α , a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α , or a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ,

R²⁴ represents a hydrogen atom, a halogen atom, a cyano group, a C₁-C₆ alkyl group, a C₃-C₆ cycloalkyl group, or a C₁-C₆ alkoxy-carbonylamino group,

R²⁵ represents a C₁-C₆ alkoxy-carbonyl group, a cyano group, or a nitro group,

R³¹ and R³² each independently represent a hydrogen atom; a C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl group; a C₁-

C₁₀ alkylthio carbonyl group; a C₁-C₆ alkoxy carbonyl group; a C₁-C₆ haloalkyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl group; a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); or a C₁-C₆ alkyl group substituted with a heterocyclic group in which the heterocyclic group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α), wherein, R³¹ and R³² may be joined together to form a 5- to 6-membered ring with adjacent nitrogen atom, and the one or more carbon atoms in the ring may be substituted with a sulfur atom and/or an oxygen atom ,

wherein, "Substituent group α " represents a group selected from a group consisting of:

a halogen atom; a hydroxyl group; a C₁-C₆ alkyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₃-C₆ halocycloalkyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy group; a C₃-C₆ cycloalkyloxy group; a C₂-C₆ alkenyloxy group; a C₂-C₆ alkynyloxy group; a C₁-C₆ alkylcarbonyloxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a C₁-C₆ haloalkylthio group; a C₁-C₆ haloalkylsulfinyl group; a C₁-C₆ haloalkylsulfonyl group; an amino group; a C₁-C₆ alkylcarbonylamino group; a mono(C₁-C₆ alkyl)amino group; a di(C₁-C₆ alkyl)amino group; a hydroxy C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ haloalkylthio C₁-C₆ alkyl group; a C₁-C₆ haloalkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ haloalkylsulfonyl C₁-C₆ alkyl group; a cyano C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkoxy group; a C₃-C₆ cycloalkyl C₁-C₆ alkyloxy group; a C₁-C₆ haloalkoxy C₁-C₆ alkoxy group; a cyano C₁-C₆ alkoxy group; a C₁-C₆ acyl group; a C₁-C₆

alkoxyimino C₁-C₆ alkyl group; a carboxyl group; a C₁-C₆ alkoxycarbonyl group; a carbamoyl group; a mono(C₁-C₆ alkyl)aminocarbonyl group; a di(C₁-C₆ alkyl)aminocarbonyl group; a nitro group; a cyano group; a phenyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic group comprising 2 to 10 carbon atoms and 1 to 5 identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic oxy group comprising 2 to 10 carbon atoms and 1 to 5 identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); and a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting a carbonyl group; and

"Substituent group β " represents a group selected from a group consisting of: a halogen atom, a nitro group, a cyano group, a C₁-C₆ alkyl group, a C₁-C₆ haloalkyl group, a C₁-C₆ alkoxy group, and a C₁-C₆ haloalkoxy group.

2. The triazine derivative or the salt thereof according to claim 1, wherein

R¹ represents a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₃-C₆ cycloalkyloxy C₁-C₆ alkyl group; a phenyloxy C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylthio C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfinyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfonyl C₁-C₆ alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different

substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₂-C₆ alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a di(C₁-C₆ alkoxy) C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyloxy C₁-C₆ alkyl group; a NR³¹R³² group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone) ; or a C₁-C₆ alkyl group substituted with a heterocyclic group in which the heterocyclic group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α);

R² represents a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ;

Y and Z represent an oxygen atom or a sulfur atom,

A represents any one of A-1, A-3, and A-5,

A₁ is [X₁],

A₂ is [X₃] or [X₄], and

A₃ is [X₉],

in [X₁], R⁵ and R⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group,

in [X₃], R⁸ and R⁹ each independently represent a hydrogen atom or a C₁-C₆ alkyl group,

in [X₉], R³⁵ and R³⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group,

wherein, R⁵ and R⁸ may be joined together to form a C₂-C₅ alkylene chain or a C₂-C₅ alkenylene chain, and may form a ring together with adjacent carbon atoms, and R⁵ and R³⁵ may be joined together to form a C₁-C₅ alkylene chain to form a ring with adjacent carbon atoms,

in A-3, R²⁰ is a C₁-C₆ alkyl group,

R²¹ is a hydrogen atom or a C₁-C₆ alkyl group,

in A-5, R²⁴ represents a hydrogen atom, a C₁-C₆ alkyl group, or a C₃-C₆ cycloalkyl group, R²⁵ represents a C₁-C₆ alkoxy carbonyl group, a cyano group, or a nitro group,

R⁴ represents a hydroxyl group; O⁻M⁺ (M⁺ represents an alkali metal cation or an ammonium cation); or a C₁-C₁₀ alkylsulfonyloxy group;

R³¹ and R³² each independently represent a hydrogen atom; a C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ; wherein, R³¹ and R³² may be joined together to form a 5- to 6-membered ring with adjacent nitrogen atom, and the one or more carbon atoms in the ring may be substituted with a sulfur atom and/or an oxygen atom,

wherein, "Substituent group α " represents a group selected from a group consisting of:

a halogen atom; a C₁-C₆ alkyl group; a C₃-C₆ cycloalkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl

group; a C₃-C₆ halocycloalkyl group; a C₁-C₆ alkoxy group; a C₃-C₆ cycloalkyloxy group; a C₂-C₆ alkenyloxy group; a C₂-C₆ alkynyloxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a nitro group; a cyano group; a phenyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a heterocyclic oxy group comprising 2 to 10 carbon atoms and 1 to 5 heteroatoms that are optionally selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); and a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting a carbonyl group.

3. The triazine derivative or the salt thereof according to claim 1, wherein

R¹ is a group selected from a group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkoxy carbonyl C₁-C₆ alkyl group; a C₁-C₆ alkyl carbonyl C₁-C₆ alkyl group; a NR³¹R³² group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); and, a C₁-C₆ alkyl group substituted with a heterocyclic group in which the heterocyclic group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α);

R^{31} and R^{32} each independently represent a group selected from a group consisting of a hydrogen atom; a C_1 - C_6 alkyl group; and, a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ;

R^2 represents a group selected from a group consisting of a C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_1 - C_6 haloalkyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); and, a phenyl group which may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α ;

Y and Z represent an oxygen atom or a sulfur atom,

A represents any one of A-1, A-3, and A-5,

R^4 in A-1 represents a hydroxyl group; O^-M^+ (M^+ represents an alkali metal cation or an ammonium cation); or a C_1 - C_{10} alkylsulfonloxy group;

in A-1, A_1 is $[X_1]$,

A_2 is $[X_3]$ or $[X_4]$, and

A_3 is $[X_9]$,

in $[X_1]$, R^5 and R^6 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_3]$, R^8 and R^9 each independently represent a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_9]$, R^{35} and R^{36} each independently represent a hydrogen atom or a C_1 - C_6 alkyl group,

wherein, R^5 and R^8 may bind to each other via a C_2 - C_5 alkylene chain or a C_2 - C_5 alkenylene chain to form a ring, and R^5 and R^{35} may bind to each other via a C_1 - C_5 alkylene chain to form a ring,

in A-3, R²⁰ is a C₁-C₆ alkyl group,

R²¹ is a hydrogen atom or a C₁-C₆ alkyl group, and

"Substituent group α " represents a group selected from a group consisting of : a halogen atom; a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₁-C₆ alkoxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a nitro group; a cyano group; a phenyl group; and a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting a carbonyl group.

4. The triazine derivative or the salt thereof according to claim 1, wherein

R¹ represents a group selected from a group consisting of a C₁-C₁₂ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a NR³¹R³² group; a heterocyclic group selected from the group consisting of pyridyl group, pyrimidinyl group, pyridazinyl group, thienyl group, isoxazolyl group, pyrazolyl group, morpholinyl group, thiomorpholinyl group, pyrazinyl group, piperidinyl group, and pyperazinyl group (the heterocyclic group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); and, a tetrahydrofurylmethyl group;

R³¹ and R³² each independently represent a group selected from a group consisting of a hydrogen atom; a C₁-C₆ alkyl group; and a phenyl group;

R^2 represents a group selected from a group consisting of a C_1 - C_6 alkyl group; a C_1 - C_6 haloalkyl group; a pyridyl group; and a phenyl group;

Y and Z represent an oxygen atom or a sulfur atom,

A represents any one of A-1 and A-3,

R^4 in A-1 represents a hydroxyl group; or a C_1 - C_{10} alkylsulfonyloxy group,

in A-1, A_1 is $[X_1]$, A_2 is $[X_3]$ or $[X_4]$, and A_3 is $[X_9]$,

in $[X_1]$, R^5 and R^6 are a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_3]$, R^8 and R^9 are a hydrogen atom or a C_1 - C_6 alkyl group,

in $[X_9]$, R^{35} and R^{36} are a hydrogen atom or a C_1 - C_6 alkyl group,

wherein, R^5 and R^8 may be joined together to form a C_2 - C_5 alkylene chain and to form a ring,

and R^5 and R^{35} may be joined together to form a C_1 - C_5 alkylene chain and to form a ring,


in A-3, R^{20} is a C_1 - C_6 alkyl group, R^{21} is a hydrogen atom or a C_1 - C_6 alkyl group, and R^4 represents a hydroxyl group or a C_1 - C_{10} alkylsulfonyloxy group, and

"Substituent group α " represents a group selected from a group consisting of: a halogen atom; a C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_1 - C_6 haloalkyl group; a C_1 - C_6 alkoxy group; a C_1 - C_6 haloalkoxy group; a C_1 - C_6 alkylthio group; a C_1 - C_6 alkylsulfinyl group; a C_1 - C_6 alkylsulfonyl group; a nitro group; a cyano group; a phenyl group; and a methylenedioxy group.

5. An agrochemical composition comprising the triazine derivative or the salt thereof described in any one of claims 1 to 4, and an agriculturally acceptable carrier.

6. The agrochemical composition according to claim 5, in which the agrochemical composition further comprises a surface active agent.

7. A herbicide comprising the triazine derivative or the salt thereof described in any one of claims 1 to 4 as active component.
8. The herbicide according to claim 7, in which the herbicide has a herbicidal activity against weeds in a field or a paddy field in which agrohorticultural plants are cultivated.
9. The herbicide according to claim 8, in which the agrohorticultural plants are agrohorticultural plants given with resistance by a breeding method or a genetic recombination technique.
10. A method of eliminating weeds in soils by applying an effective amount of herbicides comprising the triazine derivative or the salt thereof described in any one of claims 1 to 4.
11. The method according to claim 10, in which the soils are a farmland.
12. The method according to claim 10, in which the soils are farmland of fields or paddy fields in which agrohorticultural plants are cultivated.
13. A triazine derivative or a salt thereof represented by following Formula 2:



[in the formula, B represents a hydroxyl group or a C₁-C₆ alkoxy group;

Y and Z represent an oxygen atom or a sulfur atom,

R¹ represents a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₂-C₆ haloalkynyl group; a C₃-C₆ halocycloalkyl group; a C₃-C₆ halocycloalkyl C₁-C₆ alkyl group; an amino C₁-C₆ alkyl group; a nitro C₁-C₆ alkyl group; a C₁-C₆ alkylamino C₁-C₆ alkyl group; a di(C₁-C₆ alkyl)amino C₁-C₆ alkyl group; a C₁-C₆ alkylthio

C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylthio C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a hydroxy C_1 - C_6 alkyl group; a phenyl C_1 - C_6 alkoxy C_1 - C_6 alkyl group (phenyl in the group may be substituted with one substituent group selected from Substituent group α or 2 to 5 substituent groups that are the same or different from each other and selected from Substituent group α); a C_1 - C_6 alkoxy C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_3 - C_6 cycloalkyloxy C_1 - C_6 alkyl group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy C_1 - C_6 alkyl group; a phenyloxy C_1 - C_6 alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylthio C_1 - C_6 alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfinyl C_1 - C_6 alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenylsulfonyl C_1 - C_6 alkyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a C_1 - C_6 haloalkoxy C_1 - C_6 alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_1 - C_6 alkyl group; a phenyl C_2 - C_6 alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_2 - C_6 alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C_1 - C_6 alkoxyimino C_1 - C_6 alkyl group; a phenoxyimino C_1 - C_6 alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a di(C_1 - C_6 alkoxy) C_1 - C_6 alkyl group; a ($R^{31}R^{32}N$ - $C=O$) C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy carbonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkyl carbonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkyl carbonyloxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylidene aminooxy C_1 - C_6 alkyl group; a formyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylthio C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a cyano C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a cyano C_1 - C_6 alkyl group; a C_2 - C_6 alkylidene amino group; a di(C_1 - C_{10} alkyl)amino C_1 - C_6 alkylidene amino group; a $NR^{31}R^{32}$ group; a C_1 - C_6 alkoxy group; a C_2 - C_6 alkenyloxy group; a C_2 - C_6 alkynyloxy group; a C_3 - C_6 cycloalkyloxy group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy group; a C_1 - C_6 haloalkoxy group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected

from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone]; a C₁-C₆ alkyl group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α]; a C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α]; or a C₁-C₆ alkoxy C₁-C₆ alkyl group substituted with a heterocyclic-oxy group in which the heterocyclic group in the heterocyclic-oxy group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom [the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α];

R³¹ and R³² each independently represent a C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzyl group which may be substituted with one or more substituents selected from the Substituent group α ; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl group; a C₁-C₁₀ alkylthio carbonyl group; a C₁-C₆ alkoxycarbonyl group; a C₁-C₆ haloalkyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl group; a phenylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a benzylsulfonyl group which may be substituted with one or more substituents selected from the Substituent group α ; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); or a C₁-C₆ alkyl group substituted with a heterocyclic group in which the heterocyclic group comprises 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α),

wherein, R^{31} and R^{32} may be joined together to form a 5- to 6-membered ring with adjacent nitrogen atom, and the one or more carbon atoms in the ring may be substituted with a sulfur atom and/or an oxygen atom,

R^2 represents a hydrogen, a C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_2 - C_6 haloalkynyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_3 - C_6 cycloalkyloxy C_1 - C_6 alkyl group; a di(C_1 - C_6 alkoxy) C_1 - C_6 alkyl group; a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α); a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_1 - C_6 alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_2 - C_6 alkenyl group which may be substituted with one or more substituents selected from the Substituent group α ; or a phenyl C_2 - C_6 alkynyl group which may be substituted with one or more substituents selected from the Substituent group α ,

wherein, "Substituent group α " represents a group selected from a group consisting of:

a halogen atom; a hydroxyl group; a C_1 - C_6 alkyl group; a C_3 - C_6 cycloalkyl group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyl group; a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_2 - C_6 haloalkynyl group; a C_3 - C_6 halocycloalkyl group; a C_3 - C_6 halocycloalkyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy group; a C_3 - C_6 cycloalkyloxy group; a C_2 - C_6 alkenyloxy group; a C_2 - C_6 alkynyloxy group; a C_1 - C_6 alkylcarbonyloxy group; a C_1 - C_6 haloalkoxy group; a C_1 - C_6 alkylthio group; a C_1 - C_6 alkylsulfinyl group; a C_1 - C_6 alkylsulfonyl group; a C_1 - C_6 haloalkylthio group; a C_1 - C_6 haloalkylsulfinyl group; a C_1 - C_6 haloalkylsulfonyl group; an amino group; a C_1 - C_6 alkylcarbonylamino group; a mono(C_1 - C_6 alkyl)amino group; a di(C_1 - C_6 alkyl)amino group; a hydroxy C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylthio C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylthio C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 haloalkylsulfonyl C_1 - C_6 alkyl group; a cyano C_1 - C_6 alkyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkoxy group; a C_3 - C_6 cycloalkyl C_1 - C_6 alkyloxy group; a C_1 -

C₆ haloalkoxy C₁-C₆ alkoxy group; a cyano C₁-C₆ alkoxy group; a C₁-C₆ acyl group; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a carboxyl group; a C₁-C₆ alkoxycarbonyl group; a carbamoyl group; a mono(C₁-C₆ alkyl)aminocarbonyl group; a di(C₁-C₆ alkyl)aminocarbonyl group; a nitro group; a cyano group; a phenyl group (the phenyl in the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic group comprising 2 to 10 carbon atoms and 1 to 5 identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); a heterocyclic oxy group comprising 2 to 10 carbon atoms and 1 to 5 identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group β); and a C₃-C₆ alkylene group formed with two adjacent substituent groups, wherein 1 to 3 carbon atoms in the alkylene group may be substituted with an atom selected from a group consisting of an oxygen atom, a sulfur atom, a nitrogen atom, and a carbon atom constituting a carbonyl group; and

"Substituent group β " represents a group selected from a group consisting of : a halogen atom, a nitro group, a cyano group, a C₁-C₆ alkyl group, a C₁-C₆ haloalkyl group, a C₁-C₆ alkoxy group, and a C₁-C₆ haloalkoxy group.

14. The triazine derivative or the salt thereof according to claim 13, wherein

Y in Formula 2 is an oxygen atom,

R¹ in Formula 2 represents a group selected from a group consisting of a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₃-C₆ cycloalkyl group; a C₃-C₆ cycloalkenyl group; a C₁-C₆ haloalkyl group; a C₂-C₆ haloalkenyl group; a C₁-C₆ alkoxy C₁-C₆ alkyl group; a C₁-C₆ alkylthio C₁-C₆ alkyl group; a C₁-C₆ alkylsulfinyl C₁-C₆ alkyl group; a C₁-C₆ alkylsulfonyl C₁-C₆ alkyl group; a C₁-C₆ alkoxyimino C₁-C₆ alkyl group; a C₁-C₆ alkoxycarbonyl C₁-C₆ alkyl group; a C₁-C₆ alkylcarbonyl C₁-C₆ alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C₁-C₆ alkyl group; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be

substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone); and

R^2 in Formula 2 represents a group selected from a group consisting of a C_1 - C_6 alkyl group; a C_1 - C_6 haloalkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group comprising 3 to 10 carbon atoms and one or more identical or different heteroatoms selected from an oxygen atom, a sulfur atom, and a nitrogen atom (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α).

15. The triazine derivative or the salt thereof according to claim 14, wherein


Y in Formula 2 is an oxygen atom,

R^1 in Formula 2 represents a group selected from a group consisting of a C_2 - C_6 alkenyl group; a C_2 - C_6 alkynyl group; a C_3 - C_6 cycloalkyl group; a C_3 - C_6 cycloalkenyl group; a C_1 - C_6 haloalkyl group; a C_2 - C_6 haloalkenyl group; a C_1 - C_6 alkoxy C_1 - C_6 alkyl group; a C_1 - C_6 alkylthio C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfinyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylsulfonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkoxyimino C_1 - C_6 alkyl group; a C_1 - C_6 alkoxycarbonyl C_1 - C_6 alkyl group; a C_1 - C_6 alkylcarbonyl C_1 - C_6 alkyl group; a phenyl group which may be substituted with one or more substituents selected from the Substituent group α ; a phenyl C_1 - C_6 alkyl group which may be substituted with one or more substituents selected from the Substituent group α ; and a heterocyclic group selected from the group consisting of pyridyl group, pyrimidinyl group, pyrazinyl group, pyridazinyl group, thienyl group, thiazolyl group, isoxazolyl group, pyrazolyl group, morpholinyl group, thiomorpholinyl group, and piperazinyl group (the group may be substituted with 1 to 5 identical or different substituents selected from the Substituent group α , and when the heteroatom in the heterocyclic group is a sulfur atom, the sulfur atom may be oxidized to sulfoxide or sulfone) ;

R^2 is a group selected from a group consisting of a C_1 - C_6 alkyl group; a C_1 - C_6 haloalkyl group; and a pyridyl group; and,

"Substituent group α " represents a group selected from a group consisting of a halogen atom; a C₁-C₆ alkyl group; a C₂-C₆ alkenyl group; a C₂-C₆ alkynyl group; a C₁-C₆ haloalkyl group; a C₁-C₆ alkoxy group; a C₁-C₆ haloalkoxy group; a C₁-C₆ alkylthio group; a C₁-C₆ alkylsulfinyl group; a C₁-C₆ alkylsulfonyl group; a nitro group; a cyano group; a phenyl group; and a methylenedioxy group.

16. A triazine derivative or a salt thereof having a formula:



wherein:

R¹ and R² are independently selected from C₁-C₆ alkyl, or phenyl substituted with 1 to 5 identical or different substituents selected from the substituent group α , wherein substituent group α is a halogen atom;

R⁴ is hydroxyl; and

Y and Z are both oxygen.

17. The triazine derivative or a salt thereof according to claim 16, wherein R¹ is a phenyl substituted with 1 to 5 halogen atoms selected from chloro or fluoro, and R² is methyl, ethyl or propyl.

18. The triazine derivative or a salt thereof according to claim 16, wherein R¹ is phenyl, substituted at any of carbons 2, 3 or 4 of the phenyl ring with chloro or fluoro and R² is methyl.

19. The triazine derivative or a salt thereof according to claim 16, wherein R¹ is phenyl substituted with 2-chloro and R² is methyl.

20. The triazine derivative or a salt thereof according to claim 16, wherein R^1 is phenyl substituted with 3-chloro and R^2 is methyl.
21. The triazine derivative or a salt thereof according to claim 16, wherein R^1 is phenyl substituted with 4-chloro and R^2 is methyl.
22. The triazine derivative or a salt thereof according to claim 16, wherein R^1 is phenyl substituted with 2-fluoro and R^2 is methyl.
23. The triazine derivative or a salt thereof according to claim 16, wherein R^1 is phenyl substituted with 3-fluoro and R^2 is methyl.
24. The triazine derivative or a salt thereof according to claim 16, wherein R^1 is phenyl substituted with 4-fluoro and R^2 is methyl.
25. An agrochemical composition comprising the triazine derivative or a salt thereof according to any one of claims 16 to 24, and an agriculturally acceptable carrier.
26. An agrochemical composition comprising at least one triazine derivative or a salt thereof according to claim 1 or claim 16, and one additional agrochemically active component selected from the group consisting of a plant disease control component, a pesticidal component, an acaricidal component, an nematocidal component, a synergistic agent component, an attracting component, a repellent component, a herbicidal component, a safener component, a microbial pesticidal component, a plant growth control component, a fertilizer component, and a soil improving agent.
27. The agrochemical composition of claim 26, wherein said triazine derivative or a salt thereof and said additional agrochemically active component are in a weight ratio of 100:1 to 1:100.
28. The agrochemical composition of claim 26, wherein said additional agrochemically active component is a herbicide selected from the group consisting of clodinafop-propargyl, cyhalofop-butyl, diclofop-methyl, diclofop-P-methyl, fenoxaprop-P-ethyl, fluazifop-butyl, fluazifop-P-butyl, haloxyfop, haloxyfop-etotyl, haloxyfop-P, metamifop, propaquizafop,

quizalofop-ethyl, quizalofop-P-ethyl, quizalofop-P-tefuryl, fenthiaprop-ethyl, alloxydim, butroxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxym, tralkoxydim, aminopyralid, pinoxaden, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, bispyribac-sodium, pyribenzoxim, pyriftalid, pyriminobac-methyl, pyriithiobac-sodium, pyrimisulfan, flucarbazone-sodium, thiencazone, propoxycarbazone-sodium, procarbazone-sodium, amidosulfuron, azimsulfuron, bensulfuron-methyl, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, flupyrsulfuron-methyl-sodium, foramsulfuron, halosulfuron-methyl, imazosulfuron, iodosulfuron-methyl-sodium, mesosulfuron-methyl, metsulfuron-methyl, nicosulfuron, oxasulfuron, primisulfuron-methyl, prosulfuron, pyrazosulfuron-ethyl, rimsulfuron, sulfometuron-methyl, sulfosulfuron, thifensulfuron-methyl, triasulfuron, tribenuron-methyl, trifloxysulfuron-sodium, triflusulfuron-methyl, tritosulfuron, orthosulfamuron, propyrisulfuron, metazosulfuron, flucetosulfuron, cloransulam-methyl, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, pyroxsulam, HNPC-C-9908, desmedipham, phenmedipham, chloridazon, brompyrazon, ametryn, atrazine, cyanazine, desmetryne, dimethametryn, eglinazine-ethyl, prometon, prometryn, propazine, simazine, simetryn, terbumeton, terbuthylazine, terbutryn, trietazine, metamitron, metribuzin, amicarbazone, bromacil, lenacil, terbacil, pentanochlor, propanil, chlorbromuron, chlorotoluron, chloroxuron, dimefuron, diuron, ethidimuron, fenuron, fluometuron, isoproturon, isouron, linuron, methabenzthiazuron, metobromuron, metoxuron, monolinuron, neburon, siduron, tebuthiuron, metobenzuron, bentazone, bromfenoxim, bromoxynil, ioxynil, pyridafol, pyridate, diquat, paraquat dichloride, acifluorfen-sodium, bifenox, chlomefenox, ethoxyfen-ethyl, fluoroglycofen-ethyl, fomesafen, lactofen, oxyfluorfen, cinidon-ethyl, flumiclorac-pentyl, flumioxazin, chlorphthalim, oxadiargyl, oxadiazon, pentoxazone, fluazolate, pyraflufen-ethyl, benzfendazole, butafenacil, saflufenacil, fluthiacet-methyl, thidiazimin, azafenidin, carfentrazone-ethyl, sulfentrazone, bencarbazone, flufenpyr-ethyl, proflumazone, pyraclonil, SYP-298, SYP-300, norflurazon, diflufenican, picolinafen, beflubutamid, fluridone, fluorchloridone, flurtamone, mesotrione, pyrasulfotole, isoxaflutole, isoxachlortole, benzofenap, pyrazolynate, pyrazoxyfen, sulcotrione, tefuryltrion, terbotrione, pyrasulfotole, topramezone, bicyclopyrone, 4-chloro-5-(1,3-dioxocyclohexa-2-yl) carbonyl-2,3-dihydrobenzothiophene-1,1-dioxide, aclonifen, clomazone. Amitrole, glyphosate, bilanafos, glufosinate, asulam, propyzamide, tebutam, chlorthal-dimethyl, benfluralin, butralin,

dinitramine, ethalfluralin, fluchloralin, oryzalin, pendimethalin, prodiamine, trifluralin, amiprofos-methyl, butamifos, dithiopyr, thiazopyr, carbetamide, chlorpropham, propham, swep, karbutilate. diphenamid, napropamide, naproanilide, acetochlor, alachlor, butachlor, butenachlor, diethatyl-ethyl, dimethachlor, dimethenamid, dimethenamid-P, metazachlor, metolachlor, pethoxamid, pretilachlor, propachlor, propisochlor, S-metolachlor, thenylchlor, flufenacet, mefenacet, fentrazamide, anilofos, bromobutide, cafenstrole, indanofan, piperophos, fenoxasulfone, pyroxasulfone, ipfencarbazone, isoxaben, dichlobenil, chlorthiamid, flupoxame, dinoterb, DNOC, benfuresate, ethofumesate. dalapon, flupropanate, TCA, bensulide. butylate, cycloate, dimepiperate, EPTC, esprocarb, molinate, orbencarb, pebulate, prosulfocarb, thiobencarb, tiocarbazil, tri-allate, vernolate, chloramben, 2,3,6-TBA, dicamba, 2,4,5-T, 2,4-D, 2,4-DB, clomeprop, dichlorprop, dichlorprop-P, MCPA, MCPA-thioethyl, MCPB, mecoprop, mecoprop-P, clopyralid, fluoroxypyr, picloram, triclopyr, triclopyr-butotyl, quinclorac, quinmerac, benazolin, naptalam, diflufenzopyr, Flamprop-M, flamprop, chlorflurenol-methyl, cinmethylin, cumyluron, daimuron, methyldymuron, difenzoquat, etobenzanid, fosamine, pyributicarb, oxaziclomefone, acrolein, AE-F-150944, aminocyclopyrachlor, cyanamide, heptamaloxyloglucan, indaziflam, triaziflam, quinoclamine, endothal-disodium, phenisopham, BDPT, BAU-9403, SYN-523, SYP-249, JS-913, IR-6396, metiozolin, Triafamone, HW-02, and BCS-AA10579.

29. The agrochemical composition of claim 26, wherein said additional agrochemically active component is a plant growth controlling compound selected from the group consisting of 1-Methylcyclopropene, 1-naphthylacetamide, 2,6-diisopropyl-naphthalene, 4-CPA, benzylaminopurine, ancymidol, aviglycine, carvone, chlormequat, cloprop, cloxyfonac, cloxyfonac-potassium, cyclanilide, cytokinins, daminodide, dikegulac, dimethipin, ethephon, ethychlozate, flumetralin, flurenol, flurprimidol, forchlorfenuron, gibberellin acid, inabenfide, indol acetic acid, indol butyric acid, maleic hydrazide, mefluidide, mepiquat chloride, n-decanol, paclobutrazol, prohexadione-calcium, prohydrojasmon, sintofen, thidiazuron, triacontanol, trinexapac-ethyl, uniconazole, uniconazole-P, and ecolyst, benoxacor, furilazole, dichlormid, dicyclonone, DKA-24 (N1,N2-diallyl-N2-dichloroacetyl-glycinamide), AD-67 (4-dichloroacetyl-1-oxa-4-azaspiro[4.5]decane), PPG-1292 (2,2-dichloro-N-(1,3-dioxan-2-ylmethyl)-N-(2-propenyl)acetamide), R-29148 (3-dichloroacetyl-2,2,5-trimethyl-1,3-oxazolidine), cloquintcet-


mexyl, naphthalic anhydride (1,8-naphthalic anhydride), mefenpyr-diethyl, mefenpyr, mefenpyr-ethyl, fenclorazole 0 ethyl, fenclorim, MG-191 (2-dichloromethyl-2-methyl-1,3-dioxane), cyometrinil, flurazole, fluxofenim, isoxadifen, isoxadifen-ethyl, mecoprop, MCPA, daimuron, 2,4-D, MON4660, oxabetrinil, cyprosulfamide, lower alkyl substituted benzoic acid, TI-35, benalaxyl, benalaxyl-M, furalaxyl, metalaxyl, metalaxyl-M,, oxadixyl, clozylacon, ofurace, bupirimate, dimethirimol, ethirimol,, hymexazol,, octhilinone,, oxolinic acid, benomyl, carbendazim, fuberidazole, thiabendazole, thiophanate, thiophanate-methyl, diethofencarb, zoxamide, pencycuron, fluopicolide diflumetorim, benodanil, flutolanil, mepronil, fluopyram, fenfuram, carboxin, oxycarboxin, thifluzamide, bixafen, furametpyr, isopyrazam, penflufen, penthiopyrad, sedaxane, boscalid,, azoxystrobin, enestroburin, picoxystrobin, pyraoxystrobin, pyraclostrobin, pyrametostrobin,, kresoxim-methyl, trifloxystrobin, dimoxystrobin, metominostrobin, orysastrobin,, famoxadone,, fluoxastrobin,, fenamidone,, pyribencarb, cyazofamid, amisulbrom, binapacryl, meptyldinocap, dinocap,, fluazinam, ferimzone, TPTA, TPTC, TPTH, silthiofam, ametocetradin, cyprodinil, mepanipyrim, pyrimethanil, blasticidin-S, mildiomicin, kasugamycin, streptomycin, oxytetracycline, gentamycin quinoxifen, proquinazid, fenpiclonil, fludioxonil, chlozolate, iprodione, rocymidone, vinclozolin, edifenphos, iprobenfos, pyrazophos, isoprothiolane, biphenyl, chloroneb, dicloran, quintozone, tecnazene, and tolclofos-methyl, etridiazole, iodocarb, propamocarb-hydrochloride, prothiocarb, dimethomorph, flumorph, benthiavalicarb-isopropyl, iprovalicarb, and valifenalate, mandipropamid, *Bacillus subtilis* (strain: QST 713), triforine, pyrifenoxy, fenarimol, nuarimol, imazalil, oxpoconazole-fumarate, pefurazoate, prochloraz, triflumizole, azaconazole, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, etaconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, triticonazole, furconazole, furconazole-cis, quinconazole, aldimorph, dodemorph, fenpropimorph, tridemorph, fenpropidin, piperalin, spiroxamine, fenhexamid, pyributicarb, naftifine, terbinafine validamycin, polyoxin, phthalide, pyroquilon, tricyclazole, carpropamid, diclocymet, fenoxanil, acibenzolar-S-methyl, probenazole, tiadinil, isotianil, laminarin, copper hydroxide, copper dioctanoate, copper oxychloride, copper, sulfate, cuprous oxide, oxine-copper, Bordeaux mixture, and copper nonyl phenol sulphonate, sulfur, ferbam, mancozeb, maneb,

metiram, propineb, thiram, zineb, ziram, cufraneb, captan, folpet, captafol, chlorothalonil, dichlofluanid, tolylfluanid, guazatine, iminoctadine-albesilate, iminoctadine-triacetate, dodine, anilazine, dithianon, cymoxanil, vfosetyl (aluminium, calcium, and sodium), phosphorous acid and salts, tecloftalam, triazoxide, flusulfamide, diclomezine, methasulfocarb, ethaboxam, cyflufenamid, metrafenone, potassium bicarbonate, sodium bicarbonate, BAF-045, BAG-010, benthiazole, bronopol, carvone, chinomethionat, dazomet, DBEDC, debacarb, dichlorophen, difenzoquat-methyl sulfate, dimethyl disulfide, diphenylamine, ethoxyquin, flumetover, fluoroimide, flutianil, fluxapyroxad, furancarboxylic acid, metam, nabam, natamycin, nitrapyrin, nitrothal-isopropyl, o-phenylphenol, oxazinylazole, oxyquinoline sulfate, phenazine oxide, polycarbamate, pyriofenone, S-2188, silver, SYP-Z-048, tebufloquin, tolnifanide, trichlamide, mineral oils, organic oils, *Agrobacterium radiobacter*, Fermented product from *Aspergillus* spp., *Bacillus* spp., Harpin protein, *Erwinia carotovora*, *Fusarium oxysporum*, *Gliocladium* spp., Laccase, *Pseudomonas* spp., *Talaromyces* spp., *Trichoderma* spp., Extract from mushroom, and Bacteriophage.

30. The agrochemical composition of claim 26, wherein said additional agrochemically active component is selected from the group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, butocarboxim, butoxycarboxim, carbaryl, carbofuran, carbosulfan, ethiofencarb, fenobucarb, formetanate, furathiocarb, isoprocarb, methiocarb, methomyl, metolcarb, oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, triazamate, trimethacarb, 3,5-xylyl methylcarbamate(XMC), xylylcarb, acephate, azamethiphos, azinphos-ethyl, azinphos-methyl, cadusafos, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos, chlorpyrifos-methyl, coumaphos, cyanophos, demeton-S-methyl, diamidafos, diazinon, dichlorvos, dicrotophos, dimethoate, dimethylvinphos, dioxabenzofos, disulfoton, DSP, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fenthion, fonofos, fosthiazate, fosthietan, heptenophos, isamidofos, isazophos, isofenphos-methyl, isopropyl O-(methoxyaminothio-phosphoryl) salicylate, isoxathion, malathion, mecarbam, methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, oxydeprofos, parathion, parathion-methyl, phenthoate, phorate, phosalone, phosmet, phosphamidon, phoxim, pirimiphos-methyl, profenofos, propaphos, propetamphos, prothiofos, pyraclofos, pyridaphenthion, quinalphos, sulfotep, tebupirimfos, temephos, terbufos, tetrachlorvinphos,

thiometon, thionazin, triazophos, trichlorfon, vamidothion, dichlofenthion, imicyafos, isocarbophos, mesulfenfos, flupyrzofos, chlordane, endosulfan, gamma-BCH, acetoprol, ethiprole, fipronil, pyrafluprole, pyriprole, RZI-02-003 , acrinathrin, allethrin [including d-cis-trans and d-trans], bifenthrin, bioallethrin, bioallethrin S-cyclopentenyl, bioresmethrin, cycloprothrin, and cyfluthrin [including beta-], cyhalothrin [including gamma- and lambda-], cypermethrin [including alpha-, beta-, theta-, and zeta-], cyphenothrin [including (1R)-trans-isomers], deltamethrin, empenthrin, esfenvalerate, etofenprox, fenpropathrin, fenvalerate, flucythrinate, flumethrin, and tau-fluvalinate [including tau-], halfenprox, imiprothrin, metofluthrin, permethrin, and phenothrin [including (1R)-trans-isomer], prallethrin, profluthrin, pyrethrine, resmethrin, RU15525 , silafluofen, tefluthrin, tetramethrin, tralomethrin, transfluthrin, ZXI8901 , fluvalinate, tetramethylfluthrin, meperfluthrin, DDT, methoxychlor, acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, thiacloprid, thiamethoxam, spinetoram, spinosad, abamectin, emamectin benzoate, lepimectin, milbemectin, ivermectin, polynactins, diofenolan, hydroprene, kinoprene, methothrin, fenoxycarb, and pyriproxyfen, 1,3-dichloropropene, DCIP, ethylene dibromide, methyl bromide, chloropicrin, and sulfuryl fluoride, pymetrozine, flonicamid and pyrifluquinazon, clofentezine, diflovidazin, hexythiazox, and etoxazole, diafenthiuron,, azocyclotin, cyhexatin, fenbutatin oxide, propargite, tetradifon, chlorfenapyr and DNOC, bensultap, cartap, thiocyclam, thiosultap, bistrifluoron, chlorfluazuron, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron, teflubenzuron, triflumuron, fluazuron, buprofezin, cyromazine, chromafenozide, halofenozide, methoxyfenozide, tebufenozide, amitraz, cyflumetofen, hydramethylnon, acequinocyl, fluacrypyrim, and cyenopyrafen, fenazaquin, fenpyroximate, pyridaben, pyrimidifen, tebufenpyrad, tolfenpyrad, rotenone, indoxacarb and metaflumizon, spirotetramat, aluminium phosphide, phosphine, zinc phosphide, calcium cyanide, and phosphine, bifenazate, sodium fluoroacetate, chlorantraniliprole, flubendiamide and cyantraniliprole, azadirachtin, amidoflumet, benclothiaz, benzoximate, bromopropylate, chinomethionat, CL900167 , cryolite, dicofol, dicyclanil, dienochlor, dinobuton, fenbutatin oxide, fenothiocarb, fluensulfone, flufenerim, flusulfamide, karanjin, metham, methoprene, methoxyfenozide, methyl isothiocyanate, pyridalyl, pyrifluquinazon, sulcofuron-sodium, sulflamid, and sulfoxaflo, piperonyl butoxide and DEF.

31. The agrochemical composition of claim 26, wherein said additional agrochemically active component is selected from the group consisting of pyroxasulfone, pethoxamid, bromoxynil , flufenacet, sulfentrazone, and S-metolachlor.
32. The agrochemical composition comprising the triazine derivative or a salt thereof according to claim 26 , further comprising an agriculturally acceptable carrier.
33. The agrochemical composition according to claim 26, further comprising a surface active agent.
34. A method of controlling the growth of weeds for useful crops or plants, comprising applying the agrochemical composition of claim 26.
35. The method of claim 34, wherein the agrochemical composition is applied within an interval of 1 day to 40 days.
36. The method of claim 34 or claim 35, wherein the useful crops or plants are selected from the group consisting of rice, wheat, barley, corn, grain sorghum, soybean, cotton, and sugar beet.



[1]