

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(10) International Publication Number

WO 2017/214170 A2

(43) International Publication Date
14 December 2017 (14.12.2017)

(51) International Patent Classification:

C07K 16/28 (2006.01)

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(22) International Filing Date:

06 June 2017 (06.06.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/346,324 06 June 2016 (06.06.2016) US

(71) Applicants: CITY OF HOPE [US/US]; 1500 East Duarte Road, Duarte, California 91010 (US). BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 West 7th Street, Austin, Texas 78701 (US).

(72) Inventors: QIN, Hong; c/o City of Hope, 1500 East Duarte Road, Duarte, California 91010 (US). KWAK, Larry W.; c/o City of Hope, 1500 East Duarte Road, Duarte, California 91010 (US). LI, Jingxing; c/o City of Hope, 1500 East Duarte Road, Duarte, California 91010 (US). HUANG, Kexin; c/o City of Hope, 1500 East Duarte Road, Duarte, California 91010 (US).

(74) Agent: MCKEON, Tina Williams et al.; Kilpatrick Townsend & Stockton LLP, 1100 Peachtree Street, Suite 2800, Atlanta, Georgia 30309 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: BAFF-R ANTIBODIES AND USES THEREOF

(57) Abstract: Provided herein are BAFF-R antibodies as well as compositions and methods of making and using the same. The antibodies provided herein are, inter alia, useful for the treatment of cancer and autoimmune diseases.

BAFF-R ANTIBODIES AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority to U.S. Provisional Application No. 62/346,324, filed June 6, 2016, which is incorporated by reference herein in its entirety.

5

BACKGROUND

Antibody therapy is one of the most successful immunotherapies available in the clinic to treat hematological malignancies. An exemplary case is rituximab, which targets CD20 and elicits a cytotoxic effect against B cell lymphomas. However, a major concern 10 regarding rituximab is the emergence of rituximab-resistance thought to be due to the down-regulation of CD20, thus hindering antibodies from binding the target cell.

BRIEF SUMMARY

Provided herein are B cell activating factor receptor (BAFF-R) antibodies including a light chain variable region and a heavy chain variable region. The light chain variable 15 region includes a CDR L1 as set forth in SEQ ID NO:1, a CDR L2 as set forth in SEQ ID NO:2 and a CDR L3 as set forth in SEQ ID NO:3. And the heavy chain variable region includes a CDR H1 as set forth in SEQ ID NO:4, a CDR H2 as set forth in SEQ ID NO:5, and a CDR H3 as set forth in SEQ ID NO:6. In another aspect, the light chain variable region includes a CDR L1 as set forth in SEQ ID NO:7, a CDR L2 as set forth in SEQ ID NO:8 and 20 a CDR L3 as set forth in SEQ ID NO:9. And the heavy chain variable region includes a CDR H1 as set forth in SEQ ID NO:10, a CDR H2 as set forth in SEQ ID NO:11, and a CDR H3 as set forth in SEQ ID NO:12. Optionally, the antibody is a humanized antibody. Also provided are functional fragments of the disclosed antibodies.

A humanized B cell activating factor receptor (BAFF-R) antibody capable of 25 binding BAFF-R with a K_D of less than about 4 nM or a functional fragment thereof is provided.

A humanized B cell activating factor receptor (BAFF-R) antibody bound to a BAFF-R at a K_D of less than about 4 nM is also provided.

Also provided are chimeric antigen receptors (CAR) including an antibody provided 30 herein or a functional fragment thereof.

Isolated nucleic acids encoding BAFF-R antibodies or functional fragments of the antibodies are provided herein.

Also provided are pharmaceutical compositions including a therapeutically effective amount of a BAFF-R antibody or functional fragment thereof as disclosed herein and a 5 pharmaceutically acceptable excipient.

A mouse fibroblast cell expressing a human BAFF-R protein or functional fragment thereof is provided and the human BAFF-R protein or functional fragment thereof is expressed on the cell surface of the cell.

Methods of treating cancer in a subject in need thereof are provided. The methods 10 include administering to a subject a therapeutically effective amount of a chimeric antigen receptor provided herein, thereby treating cancer in the subject.

Also provided are methods of treating cancer in a subject in need thereof including administering to a subject a therapeutically effective amount of an antibody or functional fragment thereof disclosed herein, thereby treating cancer in the subject.

15 Methods of treating an autoimmune disease in a subject in need thereof are provided. The methods include administering to the subject a therapeutically effective amount of an antibody or functional fragment thereof as disclosed herein, thereby treating an autoimmune disease in the subject.

Also provided are methods of inhibiting proliferation of a cell. The methods include 20 contacting a cell with a BAFF-R antibody or functional fragment thereof as disclosed herein, thereby forming a contacted cell. The BAFF-R antibody or functional fragment thereof is allowed to bind a BAFF-R protein on the contacted cell, thereby inhibiting proliferation of the cell. Optionally, the cell is a lymphoid cell.

Methods of producing an anti-human BAFF-R antibody or functional fragment 25 thereof are provided. The methods include administering a mouse fibroblast cell that expresses a BAFF-R protein or fragment thereof as provided herein to a mouse, thereby forming an immunized BAFF-R mouse. A splenic cell from the immunized BAFF-R mouse is fused with a human myeloma cell, thereby forming a BAFF-R hybridoma cell. The BAFF-R hybridoma cell is then allowed to express a BAFF-R antibody, thereby producing an anti- 30 BAFF-R antibody.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A, 1B, 1C, 1D and 1E are FACS images showing generation and specificity of novel monoclonal antibodies against human BAFF-R. Figure 1A is a FACS analysis of cell surface expression of hBAFF-R-GFP fusion protein in mouse fibroblast L cells. Gated on 5 GFP-positive cells, engineered L cell clone (right plot) is compared to parental L cells (left plot). Clone D2C was selected for further studies. Figures 1B, 1C, 1D and 1E are FACS traces of fluorescent counts of anti-BAFF-R antibodies binding cell lines and patient samples. Figure 1B shows affinity purified hybridoma mAb (C90, C67, C55, and C53) binding BAFF-R-positive, human MCL lines including Mino, JeKo-1, REC-1, JVM-13, and Z-138 at a 10 concentration of 0.05 µg mAb/10⁶ cells. BAFF-R-negative 293T embryonic kidney cell line was used as a control. Figure 1C shows chimeric antibodies C55 and C90 at high and low concentration binding hBAFF-R-expressing L cells. Parental L cells and secondary anti-hIgG-APC antibodies only were used as controls. Figure 1D shows alexa fluor 488-conjugated chimeric antibodies binding a panel of NHL cell lines. Figure 1E shows chimeric 15 antibodies binding three types of NHL primary patient samples. The data are representative of three independent experiments. For all of Figures 1B-1E, the traces from top to bottom as shown in the figures correlate with the variables (e.g., antibody type or cell type) used from top to bottom shown below or next to the figures.

Figures 2A, 2B and 2C are graphs showing BAFF-R monoclonal antibodies exhibited 20 specific *in vitro* cytotoxicity against B-cell tumor lines. Antibody-induced cytotoxicities measured by chromium-51 release after incubation with C55, C90, or rituximab and effectors (NK cells or complement containing serum). NK effector cell to target ratio (E:T) of 20:1. Percentage of cell specific lysis of target cells: the first panel shows BAFF-R-expressing L cell or control parental L cells; the second and third panels show BAFF-R positive JeKo-1 25 MCL or BAFF-R negative U266 multiple myeloma cells with varying antibody concentrations shown as dose response curves. Figure 2B shows specific lysis of antibodies mixed with active complement-containing human serum (1:3 dilution) against CDC-sensitive (Raji) and -resistant (Raji-2P). Figure 2C shows ADCC effects by BAFF-R chimeric antibodies with or without NK effector cells (E:T = 20:1) on NHL lines JeKo-1, SU-DHL-6, 30 Raji, and RL. Data are shown as the mean ± s.d. of triplicate samples. * P < 0.05 compared with NK cells by two-tailed Student's t-test.

Figures 3A and 3B are graphs showing BAFF-R monoclonal antibodies induce *in* 30 *vitro* Antibody Dependent Cell-Mediated Cytotoxicity (ADCC) against primary B-cell

tumors. Antibody-dependent cell-mediated cytotoxicity (ADCC) effects were measured by chromium-51 release after incubation with C55, C90, or rituximab and effectors (NK cells). Percentage of cell specific lysis of target cells: Figure 3A shows NHL patient samples (E:T = 20:1 or 10:1); Figure 3B shows primary MCL and CLL samples from rituximab-treated refractory patients (E:T = 20:1). Data are shown as the mean \pm s.d. of triplicate samples. * P < 0.05 compared with NK cells by two-tailed Student's t-test.

Figure 4A is a schematic showing treatment schedule following Day 0 tumor challenge with minimum lethal dose of tumors. Treatments were given by IV tail vein injections. Figures 4B and 4C are images showing chimeric antibodies targeting human BAFF-R elicited *in vivo* therapeutic effects against B-cell tumors. Bioluminescence images of mice challenged with luciferase-expressing tumors: JeKo-1 (MCL) (Fig. 4B) or RS4;11 (ALL) (Fig. 4C). Experimental groups received treatment of chimeric BAFF-R mAbs (C55 or C90, as indicated). Control group mice received PBS, NK cells alone, or rituximab on the same schedule. Data are representative of three independent experiments.

Figures 5A, 5B and 5C are images or graphs showing chimeric BAFF-R antibodies induce ADCC on drug resistant lymphoma models *in vitro*. Figure 5A is a scatterplot of FACS analysis showing CD20 binding on JeKo-1 cells following CRISPR/HDR knock-out of CD20 gene. CD20 expression on selected CD20-/-clone #25 compared to WT JeKo-1. ADCC effects measured by chromium-51 release after incubation with C55, C90, or rituximab and effectors NK cells (E:T = 20:1). Percentage of cell specific lysis of target cells: rituximab-resistant JeKo-1-CD20-KO (Fig. 5B) and ibrutinib-resistant Z-138 and SP49-IR (Fig. 5C). All data are representative of two or more identical experiments. Data are shown as the mean \pm s.d. of triplicate samples. * P < 0.05 compared with NK cells by two-tailed Student's t-test.

Figures 6A and 6B are images and Figure 6C are graphs showing chimeric antibodies targeting human BAFF-R elicited *in vivo* therapeutic effects against drug resistant B-cell tumors. Bioluminescence images of mice challenged with luciferase-expressing tumors JeKo-1-CD20-KO cells (Fig. 6A) or ibrutinib-resistant Z-138 cells (Fig. 6B) followed by antibody treatments as in Figure 4. Control group mice received PBS, NK cells alone, or rituximab on the same schedule. Figure 6C shows 80-day tumor-free and overall survival curves of the mice shown in (A) and (B), respectively. Tumor free rate and survival differences between experimental and all control groups were analyzed by log-rank test (** P<0.001). Data are representative of three independent experiments.

Figures 7A and 7B show anti-human BAFF-R monoclonal antibody generation and clone selection. Figure 7A is a schematic showing L cell clone D2C, which stably expressed human hBAFF-R with a C-terminal GFP tag on the intracellular domain, was used to immunize BALB/c mice according to the schedule shown. Splenic tissue was harvested on 5 day 20 and B-cell hybridoma clones were established. Figure 7B is a table showing ELISA results from five hybridoma supernatants using anti-mouse IgG-HRP. Clones 53, 55, 67, and 90 produced BAFF-R- specific mAbs, whereas Clone 37 did not (representative of other negative clones).

Figure 8 are flow cytometry results showing verification that selected hybridoma 10 clones bind MCL cells. Binding of hybridoma Clone 53, 55, 67, and 90 supernatants (1/10, 1/50, and 1/200 dilutions) to Mino (mantle cell lymphoma) and 293T (negative control) cell lines assessed by flow cytometry performed with anti-mouse IgG-APC.

Figure 9 are graphs showing dose-dependent binding of purified mAbs to human 15 BAFF-R. Mouse mAb from hybridoma Clones 53, 55, 67, 90 were purified by protein A affinity chromatography. Binding of serially diluted (1 μ g/10⁶ cells – 1.6 ng/10⁶ cells) purified mouse mAb to Mino cells was assessed by flow cytometry with anti-mouse IgG-APC secondary antibody.

Figure 10 are results of FACS analyses showing hBAFF-R mAbs recognized non- 20 Hodgkin lymphoma cell lines *in vitro*. Mouse mAb Clones 55 and 90 bound additional cell lines at high (2 μ g mAb/ 10⁶ cells) and low (0.05 μ g mAb/ 10⁶ cells) doses: JeKo-1 (mantle cell lymphoma), SU-DHL-6 (diffuse large B cell lymphoma), Raji (Burkitt lymphoma) and RL (follicular lymphoma). Flow cytometry analysis was performed with anti-mouse IgG-APC. The traces from top to bottom as shown in the figures correlate with the variables (e.g., antibody type or cell type) used from top to bottom shown next to the figures.

Figure 11 are graphs showing hBAFF-R mAbs recognized lymphoma patient 25 samples. Mantle cell lymphoma, diffuse large B cell lymphoma, and follicular lymphoma patient samples were stained with mouse mAbs C55 and C90 at high (2 μ g/10⁶ cells) and low (0.05 μ g/10⁶ cells) doses. Flow cytometry analysis was performed with anti-mouse IgG-APC. The traces from top to bottom as shown in the figures correlate with the variables (e.g., antibody type or cell type) used from left to right shown below the figures.

Figure 12 is a graph showing chimeric antibodies induced ADCC against BAFF-R expressing L cells. BAFF-R-expressing D2C L cells (targets) were labeled with chromium-51

followed by incubation overnight with chimeric mAb + NK cells (effector to target ratio, 20:1). Culture supernatant was analyzed for released chromium.

Figure 13 are graphs showing chimeric antibodies required NK cells for cytotoxicity against tumor cells. JeKo-1 cells (target) were labeled with chromium-51. Cells were 5 incubated with chimeric mAb (C55, C90, or rituximab) and with or without NK cells (effector) at effector to target ratio of 20:1. Chimeric antibodies were added at concentrations from 50 to 0.005 μ g/mL. Culture supernatant was analyzed for released chromium.

Figure 14 are FACS results showing hBAFF-R mAb blocked BAFF/BAFF-R interaction. BAFF-R-expressing D2C L cell clones were incubated with C90 at 4 °C for 45 10 min (0-1000 ng/10⁶ cells) followed by incubation with recombinant BAFF ligand (0.5 μ g/ 10⁶ cells) at 4 °C for 90 min. Flow cytometry was performed and gated for anti-BAFF-PE. The signal plot shows BAFF/BAFF-R binding signal in the presence of each mAb concentration. Concentrations shown in the signal plot are shown at the top of each of the FACS results.

Figure 15 are FACS results showing limited internalization was observed with BAFF-R mAbs. Mino cells were incubated with mAb C90 (0.05 μ g/10⁶ cells) at 4 °C for 20 minutes 15 followed by incubation at 37 °C for 1 hour. Flow cytometry analysis was performed with anti-mouse IgG-APC. Cells were gated for surface localized antibodies (OUT) and loss of cell surface staining (IN).

Figure 16A are FACS results and 16B is a gel image showing CD20 knock out clones 20 generated with CRISPR. CD20 knock out clones of JeKo-1 were generated with a commercial CRISPR/HDR system substituting RFP at the CD20 locus. For Figure 16A, clones were screened and sorted by flow cytometry for CD20-/RFP+ expression. For Figure 16B, Western blotting with anti-CD20 antibodies was performed on total cell lysate from CD20-/RFP+ clones. β -actin was blotted as a loading control. Figure 16C shows FACS 25 results for clones as in Fig. 16A screened for BAFF-R/RFP+ expression to confirm that BAFF-R expression had not been affected by the CRISPR/HDR manipulation of CD20.

Figure 17 are FACS results showing characterization of BAFF-R binding against 30 normal B cells. PBMC from healthy donors were co-stained with APC-conjugated C90 chimeric antibody and (A) a lymphocyte marker panel (anti-CD20-PE, anti-CD3-PacificBlue, and anti-CD56-FITC) or (B) a myeloid cell marker panel (anti-CD45-PE, anti-CD15-PerCP-Cy5.5, and anti-CD14-PacificBlue). Each specific immune cell sub-population was gated and analyzed for binding with BAFF-R antibodies.

Figures 18A and 18B are FACS results showing characterization of hBAFF-R mAbs in normal immune cells from peripheral blood. Mouse mAb Clones 55 and 90 were tested for binding to isolated human immune cell sub-populations. B cells, T cells, and NK cells were isolated with commercial specific cell type isolation kits and stained with C55 and C90 (0.05 μ g mAb/ 10^6 cells). Flow cytometry analysis was performed with anti-mouse IgG. Myeloid cells from PBMC were gated for CD66b+ and analyzed for mAb C55 and C90 staining. The traces from top to bottom as shown in the figures correlate with the variables (e.g., antibody type or cell type) used from top to bottom shown next to the figures.

Figures 19A and 19B are immunohistochemistry images. For Figure 19A, immunohistochemistry was performed to identify the tissue specificity of the anti-BAFF-R antibodies. 1:150 dilution of 1 mg/mL antibodies were used to stain tissue samples. Tissue specificity of C55 mAb against human BAFF-R (20x objective lens): 1:150 dilution of the stock at 1 mg/ml. For Figure 19B, immunohistochemistry was performed on additional tonsil tissue and breast tissue to identify the tissue specificity of the anti-BAFF-R antibodies. 1:150 dilution of 1 mg/mL antibodies were used to stain tissue samples. Tissue specificity of mAb against human BAFF-R (upper panel: tonsil tissue; lower panel: breast tissue; 20x objective lens).

Figures 20A and 20B are graphs showing functional *in vitro* assays performed on the humanized variants. For Figure 20A, an ELISA assay was performed on the nine humanized variants of C90. The recombinant extracellular domain of human BAFF-R was used as the antigen. The antibodies were administered at concentrations varying from 0.78 to 100 ng/mL and their absorbance taken at 450 nm. For Figure 20B, the humanized variants were tested against JeKo-1 cells in a chromium release assay. The cells were allowed to uptake chromium followed by treatment with a humanized C90 variant and effector NK cells. The cells were incubated for 6 hours and their supernatants were sampled for their chromium content.

Figures 21A and 21B are graphs showing humanized antibodies C90-4 and C90-5 analyzed for their specific cytotoxicity of various lymphoma lines. For Figure 21A, JeKo-1, Z138, and RS4 were subjected to a chromium release assay with humanized antibodies C90-4 and C90-5. Antibodies were administered to the cell lines at concentrations between 0 to 5 μ g/mL and incubated for 6 hours with NK cells at an E:T ratio of 20:1. The cell supernatants were analyzed for their chromium content. For Figure 21B, LY-10, MEC-2, RL, and Raji lymphoma lines were subjected to a chromium release assay with humanized antibodies C90-

4 and C90-5. Antibodies were administered to the cell lines at 5 μ g/mL and incubated for 6 hours with NK cells at an E:T ratio of 20:1. The cell supernatants were analyzed for their chromium content.

Figures 22A and 22B are FACS results and graphs showing humanized C90 antibody 5 lead candidates tested for binding and cytotoxicity against primary MCL samples. For Figure 22A, three primary MCL tumor samples were co-stained with CD20-APC and biotinylated humanize C90 followed by signal detection using PE-conjugated streptavidin. For Figure 22B, cytotoxicity of humanized C90 against primary tumor samples were evaluated with a 10 chromium release assay. Cells were incubated with chromium-51 followed by treatment with antibodies and effector NK cells. Following overnight incubation, supernatants were sampled and the chromium contents were determined.

Figure 23 are FACS results showing flow cytometry analysis of biotinylated 15 humanized C90-4 and C90-5. The antibodies were used to stain PBMCs followed by detection with fluorescent PE streptavidin probe. The PBMCs were also labeled with granulocyte marker CD66b-PerCP-Cy5.5, monocyte marker CD14-PE-Cy7, B cell marker CD20-APC, T cell marker CD3-PE-Cy5, and NK cell marker CD56-FITC. The PBMCs were analyzed by flow cytometry.

Figure 24 are images showing BAFF-R chimeric antigen receptor (CAR) T cell *in* 20 *vivo* tumor treatments. Donor T cells were engineered to express chimeric C55 anti-BAFF-R single chain (sFv) onto a T cell receptor signaling domain with a 4-1BB motif. NSG mice were challenged with the minimum lethal dose of NHL JeKo-1-Luci cells (1×10^6 cells). The tumor cells were allowed to engraft until a tumor was detectable by bioluminescent imaging (Day 9). Mice were administered either a T cell therapy (5×10^6 CAR-T cells) or controls on days 9 and 15 post tumor challenge. The mice were monitored closely and imaged every 25 three days to track the tumor development.

DETAILED DESCRIPTION

Provided herein are, *inter alia*, BAFF-R antibodies including a light chain variable 30 region and a heavy chain variable region. Functional fragments of the antibodies are also provided. The BAFF-R antibodies and functional fragments thereof provided herein are capable of binding to human BAFF-R protein and induce antibody-dependent cellular cytotoxicity (ADCC) on BAFF-R-expressing cells (e.g., B cells). Optionally, the light chain variable region and the heavy chain variable region of the antibodies provided herein form

part of a chimeric antigen receptor (CAR). Thus, the compositions and methods provided herein may, *inter alia*, be used for the treatment of cancer (e.g., B cell malignancies) or autoimmune diseases.

A BAFF-R, BAFF receptor or BAFF-R protein as referred to herein includes any of the recombinant or naturally-occurring forms of the B-cell activating factor receptor (BAFF-R) also known as tumor necrosis factor receptor superfamily member 13C (TNFRSF13C) or variants or homologs thereof that maintain BAFF-R activity (e.g. within at least 50%, 80%, 90%, 95%, 96%, 97%, 98%, 99% or 100% activity compared to BAFF-R). Optionally, the variants or homologs have at least 90%, 95%, 96%, 97%, 98%, 99% or 100% amino acid sequence identity across the whole sequence or a portion of the sequence (e.g. a 50, 100, 150 or 200 continuous amino acid portion) compared to a naturally occurring BAFF-R. Optionally, the BAFF-R is substantially identical to the protein identified by the UniProt reference number Q96RJ3 or a variant or homolog having substantial identity thereto. Optionally, the BAFF-R is substantially identical to the protein identified by the UniProt reference number Q9D8D0 or a variant or homolog having substantial identity thereto. Optionally, the BAFF-R is substantially identical to the protein identified by the NCBI reference number GI:16445027 or a variant or homolog having substantial identity thereto. Optionally, the BAFF-R is substantially identical to the protein identified by the NCBI reference number GI:16306481 or a variant or homolog having substantial identity thereto.

A B cell activating factor receptor (BAFF-R) antibody including a light chain variable region and a heavy chain variable region is provided. The light chain variable region includes a CDR L1 as set forth in SEQ ID NO:1, a CDR L2 as set forth in SEQ ID NO:2 and a CDR L3 as set forth in SEQ ID NO:3. And the heavy chain variable region includes a CDR H1 as set forth in SEQ ID NO:4, a CDR H2 as set forth in SEQ ID NO:5, and a CDR H3 as set forth in SEQ ID NO:6. Optionally, the light chain variable region includes a CDR L1 as set forth in SEQ ID NO:7, a CDR L2 as set forth in SEQ ID NO:8 and a CDR L3 as set forth in SEQ ID NO:9. And the heavy chain variable region includes a CDR H1 as set forth in SEQ ID NO:10, a CDR H2 as set forth in SEQ ID NO:11, and a CDR H3 as set forth in SEQ ID NO:12. Optionally, the antibody is a humanized antibody. Also provided are functional fragments of the disclosed antibodies.

The humanized antibodies as provided herein are capable of binding a BAFF-R protein and include at least one mouse CDR or a functional fragment or variant thereof of the BAFF-R antibody provided herein (e.g., CDR L1 of SEQ ID NO:1 or 7, CDR L2 of SEQ ID

NO:2 or 8, CDR L3 of SEQ ID NO:3 or 9, CDR H1 of SEQ ID NO:4 or 10, CDR H2 of SEQ ID NO:5 or 11, CDR H3 of SEQ ID NO:7 or 13). A functional fragment of a CDR is a portion of a complete CDR amino acid sequence yet the antibody or fragment thereof containing the functional fragment is still capable of binding to an antigen (e.g., BAFF-R). A

5 functional variant of a CDR is a CDR with one or more changes to the CDR sequence yet the antibody or functional fragment thereof containing the functional variant is still capable of binding to an antigen (e.g., BAFF-R). For example, a functional variant of a nucleic acid sequence encoding a CDR can include one or more changes yet still encode the same amino acid sequence of the CDR. Further, a functional variant of a polypeptide sequence of a CDR 10 can include one or more amino acid changes as long as the antibody or functional fragment thereof bind to the antigen. Thus, a functional fragment or variant of a CDR typically includes the amino acid residues required for antibody binding to the antigen (e.g., BAFF-R). Where a humanized antibody includes at least one CDR, the at least one CDR or a functional fragment thereof is derived from a donor antibody. Optionally, the donor antibody is a 15 mouse antibody. A person of skill in the art will immediately recognize that a humanized antibody including at least one mouse CDR is a humanized antibody with at least one mouse CDR derived from a donor antibody and the additional CDRs are derived from the acceptor antibody (e.g. where the light chain includes a total of three CDRs and the heavy chain includes a total of three CDRs).

20 Where the BAFF-R antibody provided herein is a humanized antibody, the antibody may include a humanized heavy chain variable region and/or a humanized light chain variable region. Optionally, the humanized light chain variable region and the humanized heavy chain variable region include combined one mouse CDR or functional fragment or variant of a mouse CDR. Thus, the humanized light chain variable region and the humanized 25 heavy chain variable region can include combined six CDRs wherein at least one of the six CDRs is a mouse CDR. Where the humanized light chain variable region and the humanized heavy chain variable region include combined one mouse CDR, the humanized light chain variable region or the humanized heavy chain variable region include one mouse CDR. For example, a humanized antibody may include CDR L3 derived from the donor antibody (e.g. 30 mouse, also referred to herein as a mouse CDR L3) and CDR L1, CDR L2, CDR H1, CDR H2, and CDR H3 derived from the acceptor antibody (i.e. human).

Optionally, the humanized light chain variable region and the humanized heavy chain variable region include combined two mouse CDRs. Where the humanized light chain

variable region and the humanized heavy chain variable region include combined two mouse CDRs, the humanized light chain variable region and the humanized heavy chain variable region each include one mouse CDR (i), the humanized light chain variable region includes two mouse CDRs (ii), or the humanized heavy chain variable region includes two mouse CDRs (iii). For example, a humanized antibody may include CDR L3 and CDR H3 derived from the donor antibody (also referred to herein as a mouse CDR L3 and a mouse CDR H3, respectively), and CDR L1, CDR L2, CDR H1, and CDR H2 derived from the acceptor antibody (i.e., human).

10 Optionally, the humanized light chain variable region and the humanized heavy chain variable region include combined three mouse CDRs. Where the humanized light chain variable region and the humanized heavy chain variable region include combined three mouse CDRs, the humanized light chain variable region may include one mouse CDR and the humanized heavy chain variable region may include two mouse CDRs (i), the humanized light chain variable region includes two mouse CDRs and the humanized heavy chain variable region includes one mouse CDR (ii), the humanized light chain variable region includes three mouse CDRs (iii), or the humanized heavy chain variable region includes three mouse CDRs (iv). For example, a humanized antibody may include CDR L3, CDR H3 and CDR L2 derived from the donor antibody (e.g. mouse, also referred to herein as a CDR L3, mouse CDR H3, and mouse CDR L2 respectively) and CDR L1, CDR H1, and CDR H2 derived from the acceptor antibody (i.e., human).

15 The humanized light chain variable region and the humanized heavy chain variable region can include combined four mouse CDRs. Where the humanized light chain variable region and the humanized heavy chain variable region include combined four mouse CDRs, the humanized light chain variable region includes one mouse CDR and the humanized heavy chain variable region includes three mouse CDRs (i), the humanized light chain variable region includes three mouse CDRs and the humanized heavy chain variable region includes one mouse CDR (ii), or the humanized light chain variable region includes two mouse CDRs and the humanized heavy chain variable region includes two mouse CDRs (iii). For example, a humanized antibody may include CDR L3, CDR H3, CDR L2 and CDR L1 derived from the donor antibody (e.g. mouse, also referred to herein as a mouse CDR L3, mouse CDR H3, mouse CDR L2 and mouse CDR L1 respectively) and CDR H1 and CDR H2 derived from the acceptor antibody (i.e. human).

The humanized light chain variable region and the humanized heavy chain variable region each can include at least one mouse CDR. Where the humanized light chain variable region and the humanized heavy chain variable region each include at least one mouse CDR, the humanized light chain variable region includes at least one mouse CDR and the

5 humanized heavy chain variable region includes at least one mouse CDR. Thus, the humanized light chain variable region can include mouse CDR L1 and the humanized heavy chain includes mouse CDR H1. Optionally, mouse CDR L1 includes the amino acid sequence of SEQ ID NO:1 and mouse CDR H1 includes the amino acid sequence of SEQ ID NO:4. Optionally, mouse CDR L1 is the amino acid sequence of SEQ ID NO:1 and mouse

10 CDR H1 is the amino acid sequence of SEQ ID NO:4. Optionally, the humanized light chain variable region includes mouse CDR L2 and the humanized heavy chain variable region includes mouse CDR H2. Optionally, mouse CDR L2 includes the amino acid sequence of SEQ ID NO:2 and mouse CDR H2 includes the amino acid sequence of SEQ ID NO:5. Optionally, mouse CDR L2 is the amino acid sequence of SEQ ID NO:2 and mouse CDR H2

15 is the amino acid sequence of SEQ ID NO:5. Optionally, the humanized light chain variable region includes mouse CDR L3 and the humanized heavy chain variable region includes mouse CDR H3. Optionally, mouse CDR L3 includes the amino acid sequence of SEQ ID NO:3 and mouse CDR H3 includes the amino acid sequence of SEQ ID NO:6. Optionally, CDR L3 is the amino acid sequence of SEQ ID NO:3 and mouse CDR H3 is the amino acid

20 sequence of SEQ ID NO:6.

Optionally, mouse CDR L1 includes the amino acid sequence of SEQ ID NO:7 and mouse CDR H1 includes the amino acid sequence of SEQ ID NO:10. Optionally, mouse CDR L1 is the amino acid sequence of SEQ ID NO:7 and mouse CDR H1 is the amino acid sequence of SEQ ID NO:10. Optionally, the humanized light chain variable region includes mouse CDR L2 and the humanized heavy chain variable region includes mouse CDR H2.

25 Optionally, mouse CDR L2 includes the amino acid sequence of SEQ ID NO:8 and mouse CDR H2 includes the amino acid sequence of SEQ ID NO:11. Optionally, mouse CDR L2 is the amino acid sequence of SEQ ID NO:8 and mouse CDR H2 is the amino acid sequence of SEQ ID NO:11. Optionally, the humanized light chain variable region includes mouse CDR

30 L3 and the humanized heavy chain variable region includes mouse CDR H3. Optionally, mouse CDR L3 includes the amino acid sequence of SEQ ID NO:9 and mouse CDR H3 includes the amino acid sequence of SEQ ID NO:12. Optionally, CDR L3 is the amino acid

sequence of SEQ ID NO:9 and mouse CDR H3 is the amino acid sequence of SEQ ID NO:12.

The presence of mouse CDR L3 and mouse CDR H3 may be sufficient for binding of a humanized antibody to BAFF-R. Thus, the humanized antibody may not include mouse

5 CDR L1, mouse CDR L2, CDR H1 or mouse CDR H2. Where the humanized antibody does not include mouse CDR L1, mouse CDR L2, mouse CDR H1 or mouse CDR H2, the humanized antibody includes CDR L1, CDR L2, CDR H1 or CDR H2 derived from the acceptor antibody (i.e. human). Thus, a humanized antibody that does not include mouse CDR L1, mouse CDR L2, mouse CDR H1 or mouse CDR H2, does not include CDR L1,

10 CDR L2, CDR H1 or CDR H2 from a donor antibody (e.g. mouse, rat, rabbit), but includes CDR L1, CDR L2, CDR H1 or CDR H2 from the acceptor antibody (i.e. human). Thus, the humanized light chain variable region may not include mouse CDR L1 or mouse CDR L2 and the humanized heavy chain variable region does not include mouse CDR H1 or mouse CDR H2. Optionally, the humanized light chain variable region does not include mouse

15 CDR L1 and mouse CDR L2 and the humanized heavy chain variable region does not include mouse CDR H1 and mouse CDR H2.

Optionally, the humanized light chain variable region includes mouse CDR L2 and mouse CDR L3 and the humanized heavy chain variable region includes mouse CDR H2 and mouse CDR H3. Optionally, the humanized light chain variable region includes mouse CDR

20 L1, mouse CDR L2 and mouse CDR L3 and the humanized heavy chain variable region includes mouse CDR H1, mouse CDR H2 and mouse CDR H3. Optionally, the humanized light chain variable region includes mouse CDR L1 as set forth in SEQ ID NO:1, mouse CDR L2 as set forth in SEQ ID NO:2 and mouse CDR L3 as set forth in SEQ ID NO:3, and the humanized heavy chain variable region includes mouse CDR H1 as set forth in SEQ ID

25 NO:4, mouse CDR H2 as set forth in SEQ ID NO:5, and mouse CDR H3 as set forth in SEQ ID NO:6. Optionally, the humanized light chain variable region includes mouse CDR L1 as set forth in SEQ ID NO:7, mouse CDR L2 as set forth in SEQ ID NO:8 and mouse CDR L3 as set forth in SEQ ID NO:9, and the humanized heavy chain variable region includes mouse CDR H1 as set forth in SEQ ID NO:10, mouse CDR H2 as set forth in SEQ ID NO:11, and

30 mouse CDR H3 as set forth in SEQ ID NO:12.

The position of CDRs and FRs may be defined by the Kabat numbering system (Kabat et al., Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, U.S. Government Printing Office (1991)).

Likewise, the positions occupied by individual residues within the light or the heavy chain of an antibody may be defined by the Kabat numbering system. Therefore, the location of residues required for binding within a humanized light chain and a humanized heavy chain of a humanized antibody may be defined by the position of the residue according to the Kabat numbering system as is well known in the art. As described above, a humanized antibody may be an antibody having CDRs from a donor antibody (e.g. mouse) and variable region framework (FR) from a human antibody. The framework regions (FRs) are said to hold the CDRs in place in a humanized antibody. Proceeding from the amino-terminus, these regions are designated FR L1, FR L2, FR L3, and FR L4 for the light chain and FR H1, FR H2, FR 5 H3, and FR H4, for the heavy chain, respectively. Provided herein are humanized antibodies that include one or more residues within the framework regions. Optionally, these residues are important for epitope binding of the humanized antibody. A framework region residue involved in (or important for) epitope binding (e.g. BAFF-R binding) is referred to herein as a binding framework region residue. The binding framework region residues may reside in 10 the framework region of a humanized light chain variable region (i.e. FR L1, FR L2, FR L3, FR L4) or they may reside in the framework of a humanized heavy chain variable region (i.e. FR H1, FR H2, FR H3, FR H4). A binding framework residue residing in the FR L3 region of a humanized light chain is referred to herein as a FR L3 binding framework region residue. Thus, a binding framework region residue residing in the FR H3 region of a humanized heavy 15 chain is referred to herein as a FR H3 binding framework region residue.

20

Optionally, the humanized antibody includes at least one binding framework region residue. Optionally, the humanized light chain variable region includes at least one binding framework region residue. Optionally, the humanized light chain variable region includes one or more FR L1, FR L2, FR L3 or FR L4 binding framework region residues. Optionally, the humanized light chain variable region includes one or more FR L1 binding framework 25 region residues. Optionally, the humanized light chain variable region includes one or more FR L2 binding framework region residues. Optionally, the humanized light chain variable region includes one or more FR L3 binding framework region residues. Optionally, the humanized light chain variable region includes one or more FR L4 binding framework region residues. Optionally, the humanized heavy chain variable region includes one or more FR 30 H1, FR H2, FR H3 or FR H4 binding framework region residues. Optionally, the humanized heavy chain variable region includes one or more FR H1 binding framework region residues. Optionally, the humanized heavy chain variable region includes one or more FR H2 binding

framework region residues. Optionally, the humanized heavy chain variable region includes one or more FR H3 binding framework region residues. Optionally, the humanized heavy chain variable region includes one or more FR H4 binding framework region residues.

The humanized light chain variable region can include at least one binding framework region residue (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 or more residues) and the humanized heavy chain variable region includes at least one binding framework region residue (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 or more residues). The position of a binding framework region residue within a humanized antibody may be defined by the Kabat numbering system similar to the positions of CDR residues.

Optionally, the light chain variable region includes a serine at a position corresponding to Kabat position 7. Optionally, the light chain variable region includes a proline at a position corresponding to Kabat position 8. Optionally, the light chain variable region includes a valine at a position corresponding to Kabat position 15. Optionally, the light chain variable region includes a threonine at a position corresponding to Kabat position 22. Optionally, the light chain variable region includes a glutamine at a position corresponding to Kabat position 24. Optionally, the light chain variable region includes a glycine at a position corresponding to Kabat position 41. Optionally, the light chain variable region includes a lysine at a position corresponding to Kabat position 42. Optionally, the light chain variable region includes an alanine at a position corresponding to Kabat position 43. Optionally, the light chain variable region includes a proline at a position corresponding to Kabat position 44. Optionally, the light chain variable region includes a threonine at a position corresponding to Kabat position 56. Optionally, the light chain variable region includes a threonine at a position corresponding to Kabat position 72. Optionally, the light chain variable region includes a phenylalanine at a position corresponding to Kabat position 73. Optionally, the light chain variable region includes a glutamine at a position corresponding to Kabat position 79. Optionally, the light chain variable region includes a valine at a position corresponding to Kabat position 104.

Optionally, the light chain variable region includes a serine at a position corresponding to Kabat position 7, a proline at a position corresponding to Kabat position 8, a valine at a position corresponding to Kabat position 15, a threonine at a position

corresponding to Kabat position 22, a glutamine or a serine at a position corresponding to Kabat position 24, a glycine at a position corresponding to Kabat position 41, a lysine at a position corresponding to Kabat position 42, an alanine or a threonine at a position corresponding to Kabat position 43, a proline at a position corresponding to Kabat position 5 44, a threonine at a position corresponding to Kabat position 56, a threonine at a position corresponding to Kabat position 72, a phenylalanine or a lysine at a position corresponding to Kabat position 73, a glutamine at a position corresponding to Kabat position 79 or a valine at a position corresponding to Kabat position 104.

10 Optionally, the light chain variable region includes a serine at a position corresponding to Kabat position 7, a proline at a position corresponding to Kabat position 8, a valine at a position corresponding to Kabat position 15, a threonine at a position corresponding to Kabat position 22, a glutamine or a serine at a position corresponding to Kabat position 24, a glycine at a position corresponding to Kabat position 41, a lysine at a position corresponding to Kabat position 42, an alanine or a threonine at a position 15 corresponding to Kabat position 43, a proline at a position corresponding to Kabat position 44, a threonine at a position corresponding to Kabat position 56, a threonine at a position corresponding to Kabat position 72, a phenylalanine or a lysine at a position corresponding to Kabat position 73, a glutamine at a position corresponding to Kabat position 79 and a valine at a position corresponding to Kabat position 104.

20 Optionally, the light chain variable region includes a binding framework region residue that is a serine at a position corresponding to Kabat position 7, a proline at a position corresponding to Kabat position 8, a valine at a position corresponding to Kabat position 15, a threonine at a position corresponding to Kabat position 22, a glutamine or a serine at a position corresponding to Kabat position 24, a glycine at a position corresponding to Kabat 25 position 41, a lysine at a position corresponding to Kabat position 42, an alanine or a threonine at a position corresponding to Kabat position 43, a proline at a position corresponding to Kabat position 44, a threonine at a position corresponding to Kabat position 56, a threonine at a position corresponding to Kabat position 72, a phenylalanine or a lysine at a position corresponding to Kabat position 73, a glutamine at a position corresponding to Kabat 30 position 79 or a valine at a position corresponding to Kabat position 104.

Optionally, the heavy chain variable region includes a threonine or an alanine at a position corresponding to Kabat position 10. Optionally, the heavy chain variable region includes a lysine at a position corresponding to Kabat position 11. Optionally, the heavy

chain variable region includes a valine at a position corresponding to Kabat position 12.

Optionally, the heavy chain variable region includes a threonine at a position corresponding to Kabat position 15. Optionally, the heavy chain variable region includes a threonine at a position corresponding to Kabat position 19. Optionally, the heavy chain variable region

5 includes a threonine at a position corresponding to Kabat position 23. Optionally, the heavy chain variable region includes a proline at a position corresponding to Kabat position 41.

Optionally, the heavy chain variable region includes an alanine at a position corresponding to Kabat position 44. Optionally, the heavy chain variable region includes a proline or a threonine at a position corresponding to Kabat position 61. Optionally, the heavy chain

10 variable region includes an arginine at a position corresponding to Kabat position 66.

Optionally, the heavy chain variable region includes a threonine at a position corresponding to Kabat position 70. Optionally, the heavy chain variable region includes a lysine at a position corresponding to Kabat position 75. Optionally, the heavy chain variable region includes a valine at a position corresponding to Kabat position 79. Optionally, the heavy

15 chain variable region includes a threonine at a position corresponding to Kabat position 81.

Optionally, the heavy chain variable region includes a methionine at a position corresponding to Kabat position 82. Optionally, the heavy chain variable region includes an asparagine at a position corresponding to Kabat position 82B. Optionally, the heavy chain variable region includes a methionine at a position corresponding to Kabat position 82C. Optionally, the

20 heavy chain variable region includes a proline at a position corresponding to Kabat position 84. Optionally, the heavy chain variable region includes a valine at a position corresponding to Kabat position 85. Optionally, the heavy chain variable region includes a lysine at a position corresponding to Kabat position 108. Optionally, the heavy chain variable region includes a valine at a position corresponding to Kabat position 109.

25 Optionally, the heavy chain variable region includes a threonine or an alanine at a position corresponding to Kabat position 10, a lysine at a position corresponding to Kabat position 11, a valine at a position corresponding to Kabat position 12, a threonine at a position corresponding to Kabat position 15, a threonine at a position corresponding to Kabat position 19, a threonine at a position corresponding to Kabat position 23, a proline at a position corresponding to Kabat position 41, an alanine at a position corresponding to Kabat position 44, a proline, a serine or a threonine at a position corresponding to Kabat position 61, an arginine at a position corresponding to Kabat position 66, a threonine at a position corresponding to Kabat position 70, a lysine at a position corresponding to Kabat position 75,

a valine at a position corresponding to Kabat position 79, a threonine or a lysine at a position corresponding to Kabat position 81, a methionine at a position corresponding to Kabat position 82, an asparagine at a position corresponding to Kabat position 82B, a methionine at a position corresponding to Kabat position 82C, a proline at a position corresponding to Kabat position 84, a valine at a position corresponding to Kabat position 85, a lysine at a position corresponding to Kabat position 108 or a valine at a position corresponding to Kabat position 109.

5 Optionally, the heavy chain variable region includes a threonine or an alanine at a position corresponding to Kabat position 10, a lysine at a position corresponding to Kabat position 11, a valine at a position corresponding to Kabat position 12, a threonine at a position corresponding to Kabat position 15, a threonine at a position corresponding to Kabat position 19, a threonine at a position corresponding to Kabat position 23, a proline at a position corresponding to Kabat position 41, an alanine at a position corresponding to Kabat position 44, a proline, a serine or a threonine at a position corresponding to Kabat position 10 61, an arginine at a position corresponding to Kabat position 66, a threonine at a position corresponding to Kabat position 70, a lysine at a position corresponding to Kabat position 75, a valine at a position corresponding to Kabat position 79, a threonine or a lysine at a position corresponding to Kabat position 81, a methionine at a position corresponding to Kabat position 82, an asparagine at a position corresponding to Kabat position 82B, a methionine at a position corresponding to Kabat position 82C, a proline at a position corresponding to Kabat position 84, a valine at a position corresponding to Kabat position 85, a lysine at a position corresponding to Kabat position 108 and a valine at a position corresponding to Kabat position 109.

15 Optionally, the heavy chain variable region includes a binding framework region residue that is a threonine or an alanine at a position corresponding to Kabat position 10, a lysine at a position corresponding to Kabat position 11, a valine at a position corresponding to Kabat position 12, a threonine at a position corresponding to Kabat position 15, a threonine at a position corresponding to Kabat position 19, a threonine at a position corresponding to Kabat position 23, a proline at a position corresponding to Kabat position 41, an alanine at a position corresponding to Kabat position 44, a proline, a serine or a threonine at a position corresponding to Kabat position 61, an arginine at a position corresponding to Kabat position 66, a threonine at a position corresponding to Kabat position 70, a lysine at a position corresponding to Kabat position 75, a valine at a position corresponding to Kabat position 79, 20 30

a threonine or a lysine at a position corresponding to Kabat position 81, a methionine at a position corresponding to Kabat position 82, an asparagine at a position corresponding to Kabat position 82B, a methionine at a position corresponding to Kabat position 82C, a proline at a position corresponding to Kabat position 84, a valine at a position corresponding to Kabat position 85, a lysine at a position corresponding to Kabat position 108 or a valine at a position corresponding to Kabat position 109.

5 Provided is a humanized BAFF-R antibody including a humanized light chain variable region including a mouse CDR L1, mouse CDR L2, or mouse CDR L3 and a humanized heavy chain variable region including a mouse CDR H1, mouse CDR H2, or mouse CDR H3. The humanized light chain variable region may include a mouse CDR L1 as set forth in SEQ ID NO:1, a mouse CDR L2 as set forth in SEQ ID NO:2, or a mouse CDR L3 as set forth in SEQ ID NO:3. The humanized light chain variable region may include a mouse CDR L1 as set forth in SEQ ID NO:1, a mouse CDR L2 as set forth in SEQ ID NO:2, and a mouse CDR L3 as set forth in SEQ ID NO:3. The humanized heavy chain variable region may include a mouse CDR H1 as set forth in SEQ ID NO:4, a mouse CDR H2 as set forth in SEQ ID NO:5, or a mouse CDR H3 as set forth in SEQ ID NO:6. The humanized heavy chain variable region may include a mouse CDR H1 as set forth in SEQ ID NO:4, a mouse CDR H2 as set forth in SEQ ID NO:5, and a mouse CDR H3 as set forth in SEQ ID NO:6. Optionally, the humanized light chain variable region includes a mouse CDR L1 as set forth in SEQ ID NO:1. Optionally, the humanized light chain variable region includes a mouse CDR L2 as set forth in SEQ ID NO:2. Optionally, the humanized light chain variable region includes a mouse CDR L3 as set forth in SEQ ID NO:3. Optionally, the humanized heavy chain variable region includes a mouse CDR H1 as set forth in SEQ ID NO:4. Optionally, the humanized heavy chain variable region includes a mouse CDR H2 as set forth in SEQ ID NO:5. Optionally, the humanized light chain variable region includes a mouse CDR H3 as set forth in SEQ ID NO:6. In further embodiments, the humanized light chain variable region includes at least one binding framework region residue. In other further embodiments, the humanized heavy chain variable region includes at least one binding framework region residue.

20 25 30 Provided is a humanized BAFF-R antibody including a humanized light chain variable region including a mouse CDR L1, mouse CDR L2, or mouse CDR L3 and a humanized heavy chain variable region including a mouse CDR H1, mouse CDR H2, or mouse CDR H3. The humanized light chain variable region may include a mouse CDR L1 as

set forth in SEQ ID NO:7, a mouse CDR L2 as set forth in SEQ ID NO:8, or a mouse CDR L3 as set forth in SEQ ID NO:9. The humanized light chain variable region may include a mouse CDR L1 as set forth in SEQ ID NO:7, a mouse CDR L2 as set forth in SEQ ID NO:8, and a mouse CDR L3 as set forth in SEQ ID NO:9. The humanized heavy chain variable

5 region may include a mouse CDR H1 as set forth in SEQ ID NO:10, a mouse CDR H2 as set forth in SEQ ID NO:11, or a mouse CDR H3 as set forth in SEQ ID NO:12. The humanized heavy chain variable region may include a mouse CDR H1 as set forth in SEQ ID NO:10, a mouse CDR H2 as set forth in SEQ ID NO:11, and a mouse CDR H3 as set forth in SEQ ID NO:12. Optionally, the humanized light chain variable region includes a mouse CDR L1 as
10 set forth in SEQ ID NO:7. Optionally, the humanized light chain variable region includes a mouse CDR L2 as set forth in SEQ ID NO:8. Optionally, the humanized light chain variable region includes a mouse CDR L3 as set forth in SEQ ID NO:9. Optionally, the humanized heavy chain variable region includes a mouse CDR H1 as set forth in SEQ ID NO:10.

15 Optionally, the humanized heavy chain variable region includes a mouse CDR H2 as set forth in SEQ ID NO:11. Optionally, the humanized light chain variable region includes a mouse CDR H3 as set forth in SEQ ID NO:12. In further embodiments, the humanized light chain variable region includes at least one binding framework region residue. In other further embodiments, the humanized heavy chain variable region includes at least one binding framework region residue.

20 Optionally, the light chain variable region includes the sequence of SEQ ID NO:18, SEQ ID NO:20 or SEQ ID NO:22. Optionally, the light chain variable region includes the sequence of SEQ ID NO:18. Optionally, the light chain variable region includes the sequence of SEQ ID NO:20. Optionally, the light chain variable region includes the sequence of SEQ ID NO:22. Optionally, the light chain variable region is the sequence of SEQ ID NO:18. Optionally, the light chain variable region is the sequence of SEQ ID
25 NO:20. Optionally, the light chain variable region is the sequence of SEQ ID NO:22. Optionally, the heavy chain variable region includes the sequence of SEQ ID NO:24, SEQ ID NO:26 or SEQ ID NO:28. Optionally, the heavy chain variable region includes the sequence of SEQ ID NO:24. Optionally, the heavy chain variable region includes the sequence of SEQ ID NO:26. Optionally, the heavy chain variable region includes the sequence of SEQ ID NO:28. Optionally, the heavy chain variable region is the sequence of SEQ ID NO:24. Optionally, the heavy chain variable region is the sequence of SEQ ID NO:26. Optionally, the heavy chain variable region is the sequence of SEQ ID NO:28. Thus, in another aspect,

provided is a humanized BAFF-R antibody including a humanized light chain variable region and a humanized heavy chain variable region, wherein the humanized light chain variable region includes the sequence of SEQ ID NO:18 and the heavy chain variable region includes the sequence of SEQ ID NO:24. In another aspect, provided is a humanized BAFF-R antibody including a humanized light chain variable region and a humanized heavy chain variable region, wherein the humanized light chain variable region includes the sequence of SEQ ID NO:20 and the heavy chain variable region includes the sequence of SEQ ID NO:26. In another aspect, provided is a humanized BAFF-R antibody including a humanized light chain variable region and a humanized heavy chain variable region, wherein the humanized light chain variable region includes the sequence of SEQ ID NO:22 and the heavy chain variable region includes the sequence of SEQ ID NO:28.

5 Optionally, the antibody is a chimeric antibody. Optionally, the light chain variable region includes the sequence of SEQ ID NO:14. Optionally, the heavy chain variable region includes the sequence of SEQ ID NO:16. Optionally, the light chain variable region is the sequence of SEQ ID NO:14. Optionally, the heavy chain variable region is the sequence of SEQ ID NO:16. Thus, in another aspect, provided is a chimeric BAFF-R antibody including a light chain variable region and a heavy chain variable region, wherein the light chain variable region includes the sequence of SEQ ID NO:14 and the heavy chain variable region includes the sequence of SEQ ID NO:16.

10 20 Optionally, the light chain variable region includes the sequence of SEQ ID NO:30. Optionally, the heavy chain variable region includes the sequence of SEQ ID NO:32. Optionally, the light chain variable region is the sequence of SEQ ID NO:30. Optionally, the heavy chain variable region is the sequence of SEQ ID NO:32. Thus, in another aspect, provided is a chimeric BAFF-R antibody including a light chain variable region and a heavy chain variable region, wherein the light chain variable region includes the sequence of SEQ ID NO:30 and the heavy chain variable region includes the sequence of SEQ ID NO:32.

15 25 30 In each case where an antibody is recited herein a functional fragment can be used. Thus, for example, provided are Fab' fragments can include a heavy chain (e.g. including a constant and a variable region) and a light chain (e.g. including a constant and a variable region). Optionally, the Fab' fragment includes a humanized heavy chain (e.g. including a constant and a variable region) and a humanized light chain (e.g. including a constant and a variable region).

5 Optionally, the BAFF-R antibody or fragment thereof includes a human constant region. Optionally, the BAFF-R antibody or fragment thereof is an IgG. Optionally, the BAFF-R antibody or fragment thereof is an IgG1. Optionally, the BAFF-R antibody or fragment thereof is an IgG2. Optionally, the BAFF-R antibody or fragment thereof is an IgG3. Optionally, the BAFF-R antibody or fragment thereof is an IgG4. Optionally, the BAFF-R antibody or fragment thereof is an IgA. Optionally, the BAFF-R antibody or fragment thereof is an IgM.

10 Optionally, the BAFF-R antibody or fragment thereof is a single chain antibody. A single chain antibody includes a variable light chain and a variable heavy chain. A person of skill in the art will immediately recognize that a single chain antibody includes a single light chain and a single heavy chain, in contrast to an immunoglobulin antibody, which includes two identical pairs of polypeptide chains, each pair having one light chain and one heavy chain. Each light chain and heavy chain in turn consists of two regions: a variable (“V”) region (i.e. variable light chain and variable heavy chain) involved in binding the target 15 antigen, and a constant (“C”) region that interacts with other components of the immune system. The variable light chain and the variable heavy chain in a single chain antibody may be linked through a linker peptide. Examples for linker peptides of single chain antibodies are described in Bird, R. E., et al., *Science*. 242(4877):423-6 (1988). Methods of making scFv antibodies have been described. *See*, , Huse *et al.*, *Science* 246:1275-1281 (1989); 20 Ward *et al.*, *Nature* 341:544-546 (1989); and Vaughan *et al.*, *Nature Biotech.* 14:309-314 (1996). Briefly, mRNA from B-cells from an immunized animal is isolated and cDNA is prepared. The cDNA is amplified using primers specific for the variable regions of heavy and light chains of immunoglobulins. The PCR products are purified and the nucleic acid sequences are joined. If a linker peptide is desired, nucleic acid sequences that encode the 25 peptide are inserted between the heavy and light chain nucleic acid sequences. The nucleic acid which encodes the scFv is inserted into a vector and expressed in the appropriate host cell.

30 The ability of an antibody or functional fragment thereof to bind a specific epitope (e.g., BAFF-R) can be described by the equilibrium dissociation constant (K_D). The equilibrium dissociation constant (K_D) as defined herein is the ratio of the dissociation rate (K-off) and the association rate (K-on) of a BAFF-R antibody to a BAFF-R protein. It is described by the following formula: $K_D = K\text{-off}/K\text{-on}$. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less

than about 5 nM. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 4.5 nM. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 4 nM. Optionally, the BAFF-R antibody is capable of 5 binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 3.5 nM. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 3 nM. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 2.5 nM. Optionally, the BAFF-R antibody is capable of binding a 10 BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 2 nM. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 1.5 nM. Optionally, the BAFF-R antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 1 nM. Optionally, the BAFF-R antibody is capable of binding a 15 BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 0.5 nM.

BAFF-R protein with an equilibrium dissociation constant (K_D) of about 5 nM. Optionally, the BAFF-R antibody or functional fragment thereof is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, or 5 nM.

5 Optionally, the provided humanized B cell activating factor receptor (BAFF-R) antibody is capable of binding BAFF-R with a K_D of less than about 4 nM is provided. Optionally, the humanized B cell activating factor receptor (BAFF-R) antibody bound to a BAFF-R at a K_D of less than about 4 nM is provided. Optionally, the antibody does not induce BAFF-R activity.

10 Optionally, the BAFF-R antibody is bound to a BAFF-R protein. Optionally, the BAFF-R protein is a human BAFF-R protein. Optionally, the BAFF-R protein is encoded by a nucleic acid sequence identified by NCBI Gene ID number 115650. Optionally, the BAFF-R protein forms part of a cell. Optionally, the BAFF-R protein is expressed on the surface of said cell. Optionally, the cell is a lymphoid cell. Optionally, the cell is a B cell. Optionally, 15 the cell is a cancer cell. Optionally, the cancer cell is a lymphoma cell.

 A large variety of diagnostic and therapeutic moieties and combinations thereof may be conjugated to the BAFF-R antibody or functional fragment thereof provided herein including embodiments thereof, thereby, providing for highly stable and/or versatile drug delivery and/or diagnostic compositions. Optionally, the BAFF-R antibody or functional 20 fragment thereof includes a therapeutic moiety or a diagnostic moiety. Optionally, the therapeutic moiety or the diagnostic moiety is bound to the BAFF-R antibody or functional fragment thereof through a chemical linker. Optionally, the chemical linker is a covalent linker or a non-covalent linker. Techniques for conjugating therapeutic moieties to antibodies are well known (see, e.g., Arnon et al., *Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy*, in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., 25 *Antibodies For Drug Delivery in Controlled Drug Delivery* (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, *Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review* in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); and Thorpe et al., *The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates*, *Immunol. Rev.*, 62:119-58 (1982)). As used herein, the term antibody-drug conjugate or ADC refers to a therapeutic

moiety conjugated or otherwise covalently bound to to an antibody or functional fragment thereof.

The term therapeutic moiety as provided herein is used in accordance with its plain ordinary meaning and refers to a monovalent compound having a therapeutic benefit (e.g., prevention, eradication, amelioration of the underlying disorder being treated) when given to a subject in need thereof. Therapeutic moieties as provided herein may include, without limitation, peptides, proteins, nucleic acids, nucleic acid analogs, small molecules, antibodies, enzymes, prodrugs, cytotoxic agents (e.g. toxins) including, but not limited to ricin, doxorubicin, daunorubicin, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin D, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, and glucocorticoid. Optionally, the therapeutic moiety is an anti-cancer agent or chemotherapeutic agent as described herein. Optionally, the therapeutic moiety is a nucleic acid moiety, a peptide moiety or a small molecule drug moiety. Optionally, the therapeutic moiety is a nucleic acid moiety.

15 Optionally, the therapeutic moiety is an antibody moiety. Optionally, the therapeutic moiety is a peptide moiety. Optionally, the therapeutic moiety is a small molecule drug moiety. Optionally, the therapeutic moiety is a nuclease. Optionally, the therapeutic moiety is an immunostimulator. Optionally, the therapeutic moiety is a toxin. Optionally, the therapeutic moiety is a nuclease.

20 Also provided herein are chimeric antigen receptors (CAR) including an antibody provided herein or a functional fragment thereof.

An isolated nucleic acid encoding a BAFF-R antibody or functional fragment thereof provided herein including embodiments thereof is provided. The BAFF-R antibody or functional fragment thereof encoded by the isolated nucleic acid is described in detail throughout this application (including the description above and in the examples section). For example, the nucleic acid may encode at least one CDR, specific residues involved in binding the epitope, or binding framework residues. For instance, the nucleic acid may encode a light chain including a sequence of SEQ ID NO:1.

30 Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:29 or SEQ ID NO:31. Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:13 and the sequence of SEQ ID NO:15. Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:29 and the sequence of SEQ ID NO:31.

5 Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, or SEQ ID NO:27. Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:17 and the sequence of SEQ ID NO:23. Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:19 and the sequence of SEQ ID NO:25. Optionally, the isolated nucleic acid includes the sequence of SEQ ID NO:21 and the sequence of SEQ ID NO:27.

A pharmaceutical composition including a therapeutically effective amount of a BAFF-R antibody or functional fragment thereof provided herein and a pharmaceutically acceptable excipient is provided.

10 A therapeutically effective amount as provided herein refers to an amount effective to achieve its intended purpose. The actual amount effective for a particular application will depend, *inter alia*, on the condition being treated. When administered in methods to treat a disease, the pharmaceutical compositions described herein will contain an amount of active humanized antibody effective to achieve the desired result, e.g., modulating the activity of a 15 target molecule (e.g., BAFF-R), and/or reducing, eliminating, or slowing the progression of disease symptoms (e.g., cancer, autoimmune disease). Determination of a therapeutically effective amount of a BAFF-R antibody provided herein is well within the capabilities of those skilled in the art, especially in light of the detailed disclosure herein.

20 Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or acetate at a pH typically of 5.0 to 8.0, optionally 6.0 to 7.0; salts such as sodium chloride, potassium chloride, and the like to make isotonic; antioxidants; preservatives; low molecular weight polypeptides; proteins; hydrophilic polymers such as polysorbate 80; amino acids such as glycine; carbohydrates; chelating agents; sugars; and other standard ingredients known to 25 those skilled in the art (*Remington: The Science and Practice of Pharmacy*, 22nd Edition, Loyd V. Allen et al., editors, Pharmaceutical Press (2012)). The mAb can be present at a concentration of 0.1 - 100 mg/ml, e.g., 1 - 10 mg/ml or 10 - 50 mg/ml, for example 5, 10, 20, 30, 40, 50 or 60 mg/ml.

30 A pharmaceutical composition including an antibody, e.g., a humanized antibody, or a functional fragment thereof as described herein can be administered by a variety of methods known in the art. The route and/or mode of administration vary depending upon the desired results. Optionally, administration is intravenous, intramuscular, intraperitoneal, or subcutaneous, or administered proximal to the site of the target. Pharmaceutically acceptable

excipients can be suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion).

Pharmaceutical compositions of the antibody or functional fragment thereof can be prepared in accordance with methods well known and routinely practiced in the art. See, e.g., 5 *Remington: The Science and Practice of Pharmacy*, 22nd Edition, Loyd V. Allen et al., editors, Pharmaceutical Press (2012); and *Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978. Pharmaceutical compositions are preferably manufactured under GMP conditions. Typically, a therapeutically effective dose or efficacious dose of the humanized antibody is employed in 10 the pharmaceutical compositions. The humanized antibodies provided can be formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art. Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased 15 as indicated by the exigencies of the therapeutic situation. It may be advantageous to formulate the humanized antibodies in combination with other therapies or agents. It can be advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit 20 contains a predetermined quantity of humanized antibody calculated to produce the desired therapeutic effect in association with the required pharmaceutical excipient.

Actual dosage levels of the active ingredients in the pharmaceutical compositions can be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of 25 administration, without being toxic to the patient. The selected dosage level depends upon a variety of pharmacokinetic factors including the activity of the particular compositions employed, the route of administration, the time of administration, the rate of excretion of the particular antibody being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, 30 sex, weight, condition, general health and prior medical history of the patient being treated, and like factors.

A physician or veterinarian can start doses of the antibodies or functional fragments thereof employed in the pharmaceutical composition at levels lower than that required to

achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, effective doses of the compositions may vary depending upon many different factors, including the specific disease or condition to be treated, means of administration, target site, physiological state of the patient, whether the patient is human or 5 an animal, other medications administered, and whether treatment is prophylactic or therapeutic. Treatment dosages need to be titrated to optimize safety and efficacy. For administration with an antibody, the dosage ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 5 mg/kg, of the host body weight. For example dosages can be 1 mg/kg 10 body weight or 10 mg/kg body weight or within the range of 1-10 mg/kg. An exemplary treatment regime entails administration once per every two or three weeks or once a month or once every 3 to 6 months.

The BAFF-R antibody or functional fragment thereof provided herein can be administered on multiple occasions. Intervals between single dosages can be weekly, monthly or yearly. Intervals can also be irregular as indicated by measuring blood levels of 15 the humanized antibody in the patient. In some methods, dosage is adjusted to achieve a plasma antibody concentration of 1–1000 µg/ml and in some methods 25–300 µg/ml. Alternatively, antibody can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life 20 of the antibody in the patient. In general, humanized antibodies show longer half-life than that of chimeric antibodies and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest 25 of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

Mouse fibroblast cells expressing a human BAFF-R protein or fragment thereof are provided and the human BAFF-R protein or fragment thereof is expressed on the cell surface 30 of the cell. Optionally, the human BAFF-R protein or fragment thereof includes a detectable moiety. Optionally, the detectable moiety is a fluorescent moiety. Optionally, the detectable moiety is an enhanced green fluorescent protein (eGFP).

Methods of treating cancer in a subject in need thereof are provided. The method includes administering to a subject a therapeutically effective amount of a chimeric antigen receptor provided herein, thereby treating cancer in the subject.

In another aspect, a method of treating cancer in a subject in need thereof is provided including administering to a subject a therapeutically effective amount of an antibody or functional fragment thereof provided herein, thereby treating cancer in the subject.

5 Optionally, the cancer is lymphoma, leukemia or myeloma. Optionally, the cancer is lymphoma. Optionally, the lymphoma is mantle cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, marginal zone lymphoma or Burkitt's lymphoma.

10 Optionally, the lymphoma is mantle cell lymphoma. Optionally, the lymphoma is follicular lymphoma. Optionally, the lymphoma is diffuse large B-cell lymphoma. Optionally, the lymphoma is marginal zone lymphoma. Optionally, the lymphoma is Burkitt's lymphoma.

15 Optionally, the cancer is leukemia. Optionally, the leukemia is lymphoblastic leukemia, chronic lymphocytic leukemia or hairy cell leukemia. Optionally, the leukemia is lymphoblastic leukemia. Optionally, the leukemia is chronic lymphocytic leukemia. Optionally, the leukemia is hairy cell leukemia.

20 Optionally, the cancer is myeloma. Optionally, the myeloma is multiple myeloma.

25 Optionally, the method further includes administering to the subject a second therapeutic agent. Optionally, the therapeutic agent is a chimeric monoclonal antibody capable of binding a CD 20 antigen. Optionally, the therapeutic agent is rituximab. The term "rituximab" refers in a customary sense to the monoclonal antibody against the protein CD20 identified by the ATC code L01XC02.

Also provided are methods of treating an autoimmune disease in a subject in need thereof. The method includes administering to the subject a therapeutically effective amount of an antibody or functional fragment thereof as provided herein, thereby treating an autoimmune disease in the subject. Optionally, the autoimmune disease is rheumatoid arthritis, systemic Lupus erythematosus, multiple sclerosis, glomerulonephritis, Sjögren's Syndrome or autoimmune hemolytic anemia. Optionally, the autoimmune disease is rheumatoid arthritis. Optionally, the autoimmune disease is systemic Lupus erythematosus.

30 Optionally, the autoimmune disease is multiple sclerosis. Optionally, the autoimmune disease is glomerulonephritis. Optionally, the autoimmune disease is Sjögren's Syndrome. Optionally, the autoimmune disease is autoimmune hemolytic anemia. Optionally, the method further includes administering to the subject a second therapeutic agent.

5 In another aspect, a method of inhibiting proliferation of a cell is provided. The method includes contacting a cell with a BAFF-R antibody or functional fragment thereof as provided herein including embodiments thereof, thereby forming a contacted cell. The BAFF-R antibody or functional fragment thereof is allowed to bind a BAFF-R protein on the contacted cell, thereby inhibiting proliferation of the cell. Optionally, the cell is a lymphoid cell. Optionally, the cell is a B cell. Optionally, the cell is a cancer cell. Optionally, the cell is a lymphoma cell.

10 In another aspect, a method of producing an anti-human BAFF-R antibody is provided. The method includes administering a mouse fibroblast cell as provided herein to a mouse, thereby forming an immunized BAFF-R mouse. A splenic cell from the immunized BAFF-R mouse is fused with a human myeloma cell, thereby forming a BAFF-R hybridoma cell. The BAFF-R hybridoma cell is then allowed to express a BAFF-R antibody, thereby producing an anti-BAFF-R antibody. Optionally, the anti-BAFF-R antibody is an antibody as provided herein.

15 While various embodiments and aspects are shown and described herein, it will be obvious to those skilled in the art that such embodiments and aspects are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art. It should be understood that various alternatives to the embodiments described herein may be employed.

20 Antibodies are large, complex molecules (molecular weight of ~150,000 Da or about 1320 amino acids) with intricate internal structure. A natural antibody molecule contains two identical pairs of polypeptide chains, each pair having one light chain and one heavy chain. Each light chain and heavy chain in turn consists of two regions: a variable (“V”) region involved in binding the target antigen, and a constant (“C”) region that interacts with other components of the immune system. The light and heavy chain variable regions come together in 3-dimensional space to form a variable region that binds the antigen (for example, a receptor on the surface of a cell). Within each light or heavy chain variable region, there are three short segments (averaging 10 amino acids in length) called the complementarity determining regions (“CDRs”). The six CDRs in an antibody variable domain (three from the light chain and three from the heavy chain) fold up together in 3-dimensional space to form the actual antibody binding site (paratope), which docks onto the target antigen (epitope). The position and length of the CDRs have been precisely defined by Kabat, E. *et al.*, Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human

Services, 1983, 1987. The part of a variable region not contained in the CDRs is called the framework ("FR"), which forms the environment for the CDRs.

The term antibody is used according to its commonly known meaning in the art.

Antibodies exist, e.g., as intact immunoglobulins. However, a functional antibody fragment or fragments can be used whenever the terms antibody or antibodies are recited herein. For example, a number of well-characterized functional antibody fragments can be produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce $F(ab)'$ ₂, a dimer of Fab which itself is a light chain joined to V_H - C_H1 by a disulfide bond. The $F(ab)'$ ₂ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the $F(ab)'$ ₂ dimer into a Fab' monomer. The Fab' monomer is essentially a Fab with part of the hinge region (see *Fundamental Immunology* (Paul ed., 3d ed. (1993))). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, is exemplary and antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (e.g., single chain Fv) or those identified using phage display libraries (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990)) can be used as described for antibodies.

For preparation of monoclonal or polyclonal antibodies, any technique known in the art can be used (see, e.g., Kohler & Milstein, *Nature* 256:495-497 (1975); Kozbor *et al.*, *Immunology Today* 4:72 (1983); Cole *et al.*, pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy* (1985)). Monoclonal antibodies (mAb) refer to antibodies derived from a single clone. Techniques for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce antibodies to polypeptides described herein. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies. Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990); Marks *et al.*, *Biotechnology* 10:779-783 (1992)).

The epitope of a mAb is the region of its antigen to which the mAb binds. Two antibodies bind to the same or overlapping epitope if each competitively inhibits (blocks) binding of the other to the antigen. That is, a 1x, 5x, 10x, 20x or 100x excess of one antibody inhibits binding of the other by at least 30% but preferably 50%, 75%, 90% or even 99% as

measured in a competitive binding assay (see, e.g., Junghans *et al.*, *Cancer Res.* 50:1495, 1990). Alternatively, two antibodies have the same epitope if essentially all amino acid mutations in the antigen that reduce or eliminate binding of one antibody reduce or eliminate binding of the other. Two antibodies have overlapping epitopes if some amino acid 5 mutations that reduce or eliminate binding of one antibody reduce or eliminate binding of the other.

A ligand refers to an agent, e.g., a polypeptide or other molecule, capable of binding to a receptor molecule (e.g., an antibody).

A label or a detectable moiety is a composition detectable by spectroscopic, 10 photochemical, biochemical, immunochemical, chemical, or other physical means. For example, useful labels include ³²P, fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins or other entities which can be made detectable, e.g., by incorporating a radiolabel into a peptide or antibody 15 specifically reactive with a target peptide. Any appropriate method known in the art for conjugating an antibody to the label may be employed, e.g., using methods described in Hermanson, *Bioconjugate Techniques* 1996, Academic Press, Inc., San Diego.

Contacting is used in accordance with its plain ordinary meaning and refers to the process of allowing at least two distinct species (e.g. chemical compounds including biomolecules or cells) to become sufficiently proximal to react, interact or physically touch. 20 It should be appreciated; however, the resulting reaction product can be produced directly from a reaction between the added reagents or from an intermediate from one or more of the added reagents which can be produced in the reaction mixture.

The term contacting may include allowing two species to react, interact, or physically touch, wherein the two species may be, for example, an antibody as described herein and a 25 BAFF-R protein. Contacting includes, for example, allowing a humanized antibody as described herein to interact with BAFF-R.

As used herein, treating or treatment of a condition, disease or disorder or symptoms associated with a condition, disease or disorder refers to an approach for obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results 30 can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of extent of condition, disorder or disease, stabilization of the state of condition, disorder or disease, prevention of development of condition, disorder or disease, prevention of spread of condition, disorder or disease, delay or slowing of condition, disorder

or disease progression, delay or slowing of condition, disorder or disease onset, amelioration or palliation of the condition, disorder or disease state, and remission, whether partial or total. Treating can also mean prolonging survival of a subject beyond that expected in the absence of treatment. Treating can also mean inhibiting the progression of the condition, disorder or disease, slowing the progression of the condition, disorder or disease temporarily, although in some instances, it involves halting the progression of the condition, disorder or disease permanently. As used herein the terms treatment, treat, or treating refers to a method of reducing the effects of one or more symptoms of a disease or condition characterized by expression of the protease or symptom of the disease or condition characterized by expression of the protease. Thus in the disclosed method, treatment can refer to a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% reduction in the severity of an established disease, condition, or symptom of the disease or condition. For example, a method for treating a disease is considered to be a treatment if there is a 10% reduction in one or more symptoms of the disease in a subject as compared to a control. Thus the reduction can be a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or any percent reduction in between 10% and 100% as compared to native or control levels. It is understood that treatment does not necessarily refer to a cure or complete ablation of the disease, condition, or symptoms of the disease or condition. Further, as used herein, references to decreasing, reducing, or inhibiting include a change of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater as compared to a control level and such terms can include but do not necessarily include complete elimination.

The terms polypeptide, peptide, and protein are used interchangeably herein to refer to a polymer of amino acid residues, wherein the polymer may be conjugated to a moiety that does not consist of amino acids. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymers. A fusion protein refers to a chimeric protein encoding two or more separate protein sequences that are recombinantly expressed as a single moiety. The term peptidyl and peptidyl moiety means a monovalent peptide.

The term amino acid refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, γ -

carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, *i.e.*, an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified

5 R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid. The terms non-naturally occurring amino acid and unnatural amino acid refer to amino acid

10 analogs, synthetic amino acids, and amino acid mimetics which are not found in nature.

Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

15 Conservatively modified variants applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids that encode identical or essentially identical amino acid sequences. Because of the degeneracy of the genetic code, a number of nucleic acid sequences will encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are silent variations, which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One

20 of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

25 As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a conservatively modified variant where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables

providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles.

The following eight groups each contain amino acids that are conservative

5 substitutions for one another:

- 1) Alanine (A), Glycine (G);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 10 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);
- 7) Serine (S), Threonine (T); and
- 8) Cysteine (C), Methionine (M)

(*see, e.g.*, Creighton, *Proteins* (1984)).

15 Percentage of sequence identity is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the 20 number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

25 The terms identical or percent identity in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (*i.e.*, 60% identity, optionally 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99% identity over a specified region, *e.g.*, of the entire polypeptide sequences or individual domains of the polypeptides), when compared and aligned for maximum correspondence over a comparison 30 window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Such sequences are then said to be substantially identical. This definition also refers to the complement of a test sequence. Optionally, the identity exists over a region that is at least about 50 nucleotides in length, or

more preferably over a region that is 100 to 500 or 1000 or more nucleotides in length. The present description includes polypeptides that are substantially identical to any of SEQ ID NOs:30-51.

For sequence comparison, typically one sequence acts as a reference sequence, to 5 which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences 10 relative to the reference sequence, based on the program parameters.

A comparison window, as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of, e.g., a full length sequence or from 20 to 600, about 50 to about 200, or about 100 to about 150 amino acids or nucleotides in which a sequence may be compared to a reference sequence of the same 15 number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1970) *Adv. Appl. Math.* 2:482c, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of 20 Pearson and Lipman (1988) *Proc. Nat'l. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., Ausubel *et al.*, *Current Protocols in Molecular Biology* (1995 supplement)).

25 An example of an algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.* (1977) *Nuc. Acids Res.* 25:3389-3402, and Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). 30 This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*).

These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a word length of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below. Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequence.

An amino acid residue in an antibody corresponds to a given residue when it occupies the same essential structural position within the antibody as the given residue. For example, a selected residue in a comparison antibody corresponds to position 48 (according to the Kabat numbering system as described herein) in an antibody provided herein when the 5 selected residue occupies the same essential spatial or structural relationship to Kabat position 48 as assessed using applicable methods in the art. For example, a comparison antibody may be aligned for maximum sequence homology with the antibody provided herein and the position in the aligned comparison antibody that aligns with Kabat position 48 may be determined to correspond to it. Alternatively, instead of (or in addition to) a primary 10 sequence alignment as described above, a three dimensional structural alignment can also be used, e.g., where the structure of the comparison antibody is aligned for maximum correspondence with an antibody provided herein and the overall structures compared. In this case, an amino acid that occupies the same essential position as Kabat position 48 in the structural model may be said to correspond.

15 The term isolated, when applied to a protein, denotes that the protein is essentially free of other cellular components with which it is associated in the natural state. It is preferably in a homogeneous state although it can be in either a dry or aqueous solution. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein 20 that is the predominant species present in a preparation is substantially purified. The term purified denotes that a protein gives rise to essentially one band in an electrophoretic gel. Particularly, it means that the protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure.

25 The phrase specifically (or selectively) binds to an antibody or specifically (or selectively) immunoreactive with, when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Typically a 30 specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

A cell, as used herein, refers to a cell carrying out metabolic or other function sufficient to preserve or replicate its genomic DNA. A cell can be identified by well-known

methods in the art including, for example, presence of an intact membrane, staining by a particular dye, ability to produce progeny or, in the case of a gamete, ability to combine with a second gamete to produce a viable offspring. Cells may include prokaryotic and eukaryotic cells. Prokaryotic cells include but are not limited to bacteria. Eukaryotic cells include but 5 are not limited to yeast cells and cells derived from plants and animals, for example mammalian, insect (e.g., *spodoptera*) and human cells.

As defined herein, the term inhibition, inhibit, inhibiting and the like in reference to a protein-inhibitor (e.g., BAFF-R antibody provided herein) interaction means negatively affecting (e.g., decreasing) the activity or function of the protein (e.g., decreasing the activity 10 of BAFF-R) relative to the activity or function of the protein in the absence of the inhibitor (e.g., BAFF-R antibody). Inhibition includes reduction of a disease or symptoms of disease (e.g., cancer or an autoimmune disease). Thus, inhibition includes, at least in part, partially or totally blocking stimulation, decreasing, preventing, delaying activation, inactivating, desensitizing, or down-regulating signal transduction or enzymatic activity or the amount of a 15 protein. Similarly an inhibitor is a compound or protein that inhibits BAFF-R activity, e.g., by binding, partially or totally blocking, decreasing, preventing, delaying, inactivating, desensitizing, or down-regulating activity (e.g., BAFF-R signaling activity).

Agents provided herein, e.g., antibodies, are often administered as pharmaceutical compositions comprising an active therapeutic agent and a variety of other pharmaceutically acceptable components. See, *Remington: The Science and Practice of Pharmacy*, 22nd 20 Edition, Loyd V. Allen et al., editors, Pharmaceutical Press (2012). The preferred form depends on the intended mode of administration and therapeutic application. The compositions can also include, depending on the formulation desired, pharmaceutically-acceptable, non-toxic carriers or diluents, which are defined as vehicles commonly used to 25 formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the combination. Examples of such diluents are distilled water, physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may also include other carriers, adjuvants, or nontoxic, nontherapeutic, 30 nonimmunogenic stabilizers and the like.

The compositions can be administered for therapeutic or prophylactic treatments. In therapeutic applications, compositions are administered to a patient suffering from a disease (e.g., cancer) in a therapeutically effective dose. Amounts effective for this use will depend

upon the severity of the disease and the general state of the patient's health. Single or multiple administrations of the compositions may be administered depending on the dosage and frequency as required and tolerated by the patient. A patient or subject includes both humans and other animals, particularly mammals. Thus, the methods are applicable to both 5 human therapy and veterinary applications. Optionally, the patient is a mammal, a primate, or human.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the antibodies provided herein suspended in diluents, such as water, saline or PEG 400; (b) capsules, sachets or tablets, each containing a predetermined amount 10 of the active ingredient, as liquids, solids, granules or gelatin; (c) suspensions in an appropriate liquid; and (d) suitable emulsions. Tablet forms can include one or more of lactose, sucrose, mannitol, sorbitol, calcium phosphates, corn starch, potato starch, microcrystalline cellulose, gelatin, colloidal silicon dioxide, talc, magnesium stearate, stearic acid, and other excipients, colorants, fillers, binders, diluents, buffering agents, moistening 15 agents, preservatives, flavoring agents, dyes, disintegrating agents, and pharmaceutically compatible carriers. Lozenge forms can comprise the active ingredient in a flavor, e.g., sucrose, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin or sucrose and acacia emulsions, gels, and the like containing, in addition to the active ingredient, carriers known in the art.

20 Pharmaceutical compositions can also include large, slowly metabolized macromolecules such as proteins, polysaccharides such as chitosan, polylactic acids, polyglycolic acids and copolymers (such as latex functionalized sepharose(TM), agarose, cellulose, and the like), polymeric amino acids, amino acid copolymers, and lipid aggregates (such as oil droplets or liposomes). Additionally, these carriers can function as 25 immunostimulating agents (*i.e.*, adjuvants).

Suitable formulations for rectal administration include, for example, suppositories, which consist of the packaged nucleic acid with a suppository base. Suitable suppository bases include natural or synthetic triglycerides or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the compound 30 of choice with a base, including, for example, liquid triglycerides, polyethylene glycols, and paraffin hydrocarbons.

Formulations suitable for parenteral administration, such as, for example, by intraarticular (in the joints), intravenous, intramuscular, intratumoral, intradermal,

intraperitoneal, and subcutaneous routes, include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. Compositions can be administered, for example, by intravenous infusion, orally, topically, intraperitoneally, intravesically or intrathecally. Parenteral administration, oral administration, and intravenous administration are the preferred methods of administration. The formulations of compounds can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials.

10 Injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. Cells transduced by nucleic acids for *ex vivo* therapy can also be administered intravenously or parenterally as described above.

15 The pharmaceutical preparation can be in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form. The composition can, if desired, also contain other compatible therapeutic agents.

20 The combined administrations contemplates co-administration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities.

25 Effective doses of the compositions provided herein vary depending upon many different factors, including means of administration, target site, physiological state of the patient, whether the patient is human or an animal, other medications administered, and whether treatment is prophylactic or therapeutic. However, a person of ordinary skill in the art would immediately recognize appropriate and/or equivalent doses looking at dosages of approved compositions for treating and preventing cancer for guidance.

30 The terms disease or condition refer to a state of being or health status of a patient or subject capable of being treated with a compound, pharmaceutical composition, or method provided herein. Optionally, the disease is cancer (e.g. lung cancer, ovarian cancer, osteosarcoma, bladder cancer, cervical cancer, liver cancer, kidney cancer, skin cancer (e.g.,

Merkel cell carcinoma), testicular cancer, leukemia, lymphoma, head and neck cancer, colorectal cancer, prostate cancer, pancreatic cancer, melanoma, breast cancer, neuroblastoma). The disease may be an autoimmune, inflammatory, cancer, infectious, metabolic, developmental, cardiovascular, liver, intestinal, endocrine, neurological, or other 5 disease.

As used herein, the term cancer refers to all types of cancer, neoplasm or malignant tumors found in mammals, including leukemias, lymphomas, melanomas, neuroendocrine tumors, carcinomas and sarcomas. Exemplary cancers that may be treated with a compound, pharmaceutical composition, or method provided herein include lymphoma, sarcoma, bladder 10 cancer, bone cancer, brain tumor, cervical cancer, colon cancer, esophageal cancer, gastric cancer, head and neck cancer, kidney cancer, myeloma, thyroid cancer, leukemia, prostate cancer, breast cancer (e.g. triple negative, ER positive, ER negative, chemotherapy resistant, herceptin resistant, HER2 positive, doxorubicin resistant, tamoxifen resistant, ductal carcinoma, lobular carcinoma, primary, metastatic), ovarian cancer, pancreatic cancer, liver 15 cancer (e.g., hepatocellular carcinoma), lung cancer (e.g. non-small cell lung carcinoma, squamous cell lung carcinoma, adenocarcinoma, large cell lung carcinoma, small cell lung carcinoma, carcinoid, sarcoma), glioblastoma multiforme, glioma, melanoma, prostate cancer, castration-resistant prostate cancer, breast cancer, triple negative breast cancer, glioblastoma, ovarian cancer, lung cancer, squamous cell carcinoma (e.g., head, neck, or 20 esophagus), colorectal cancer, leukemia, acute myeloid leukemia, lymphoma, B cell lymphoma, or multiple myeloma. Additional examples include, cancer of the thyroid, endocrine system, brain, breast, cervix, colon, head & neck, esophagus, liver, kidney, lung, non-small cell lung, melanoma, mesothelioma, ovary, sarcoma, stomach, uterus or Medulloblastoma, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, 25 neuroblastoma, glioma, glioblastoma multiforme, ovarian cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, primary brain tumors, cancer, malignant pancreatic insuloma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, endometrial cancer, adrenal cortical 30 cancer, neoplasms of the endocrine or exocrine pancreas, medullary thyroid cancer, medullary thyroid carcinoma, melanoma, colorectal cancer, papillary thyroid cancer, hepatocellular carcinoma, Paget's Disease of the Nipple, Phyllodes Tumors, Lobular

Carcinoma, Ductal Carcinoma, cancer of the pancreatic stellate cells, cancer of the hepatic stellate cells, or prostate cancer.

The term leukemia refers broadly to progressive, malignant diseases of the blood-forming organs and is generally characterized by a distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow. Leukemia is generally clinically classified on the basis of (1) the duration and character of the disease-acute or chronic; (2) the type of cell involved; myeloid (myelogenous), lymphoid (lymphogenous), or monocytic; and (3) the increase or non-increase in the number abnormal cells in the blood-leukemic or aleukemic (subleukemic). Exemplary leukemias that may be treated with a compound, pharmaceutical composition, or method provided herein include, for example, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocytemic leukemia, basophylic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia, hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, multiple myeloma, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, or undifferentiated cell leukemia.

As used herein, the terms metastasis, and metastatic cancer can be used interchangeably and refer to the spread of a proliferative disease or disorder, e.g., cancer, from one organ or another non-adjacent organ or body part. Cancer occurs at an originating site, e.g., breast, which site is referred to as a primary tumor, e.g., primary breast cancer. Some cancer cells in the primary tumor or originating site acquire the ability to penetrate and infiltrate surrounding normal tissue in the local area and/or the ability to penetrate the walls of the lymphatic system or vascular system circulating through the system to other sites and tissues in the body. A second clinically detectable tumor formed from cancer cells of a primary tumor is referred to as a metastatic or secondary tumor. When cancer cells metastasize, the metastatic tumor and its cells are presumed to be similar to those of the

original tumor. Thus, if lung cancer metastasizes to the breast, the secondary tumor at the site of the breast consists of abnormal lung cells and not abnormal breast cells. The secondary tumor in the breast is referred to a metastatic lung cancer. Thus, the phrase metastatic cancer refers to a disease in which a subject has or had a primary tumor and has 5 one or more secondary tumors. The phrases non-metastatic cancer or subjects with cancer that is not metastatic refers to diseases in which subjects have a primary tumor but not one or more secondary tumors. For example, metastatic lung cancer refers to a disease in a subject with or with a history of a primary lung tumor and with one or more secondary tumors at a second location or multiple locations, e.g., in the breast.

10 The term associated or associated with in the context of a substance or substance activity or function associated with a disease (e.g., cancer (e.g. leukemia, lymphoma, B cell lymphoma, or multiple myeloma)) means that the disease (e.g. cancer, (e.g. leukemia, lymphoma, B cell lymphoma, or multiple myeloma)) is caused by (in whole or in part), or a symptom of the disease is caused by (in whole or in part) the substance or substance activity 15 or function.

As used herein, an autoimmune disease refers to a disease or disorder that arises from altered immune reactions by the immune system of a subject, e.g., against substances tissues and/or cells normally present in the body of the subject. Autoimmune diseases include, but are not limited to, arthritis, rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic 20 arthritis, scleroderma, systemic scleroderma, multiple sclerosis, systemic lupus erythematosus (SLE), myasthenia gravis, juvenile onset diabetes, diabetes mellitus type 1, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, ankylosing spondylitis, psoriasis, Sjogren's syndrome, vasculitis, glomerulonephritis, auto-immune thyroiditis, Behcet's disease, Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, 25 psoriasis, ichthyosis, Graves ophthalmopathy, inflammatory bowel disease, Addison's disease, Vitiligo, asthma, and allergic asthma.

As used herein, an inflammatory disease refers to a disease or disorder associated with abnormal or altered inflammation. Inflammation is a biological response initiated by the immune system as part of the healing process in response to a pathogen, damaged cells or 30 tissues or irritants. Chronic inflammation can lead to a variety of diseases. Inflammatory diseases include, but are not limited to, atherosclerosis, allergies, asthma, rheumatoid arthritis, transplant rejection, celiac disease, chronic prostatitis, inflammatory bowel diseases, pelvic inflammatory diseases, and inflammatory myopathies.

A humanized antibody is a genetically engineered antibody in which at least one CDR (or functional fragment or variant thereof) from a mouse antibody (“donor antibody,” which can also be rat, hamster or other non-human species) are grafted onto a human antibody framework (“acceptor antibody”). Optionally, more than one mouse CDR is grafted (e.g. all 5 six mouse CDRs are grafted). The sequence of the acceptor antibody can be, for example, a mature human antibody sequence (or fragment thereof), a consensus sequence of a human antibody sequence (or fragment thereof), or a germline region sequence (or fragment thereof). Thus, a humanized antibody may be an antibody having one or more CDRs from a donor antibody and a variable region framework (FR). The FR may form part of a constant region 10 and/or a variable region within a human antibody. In addition, in order to retain high binding affinity, amino acids in the human acceptor sequence may be replaced by the corresponding amino acids from the donor sequence, for example, where (1) the amino acid is in a CDR or (2) the amino acid is in the human framework region (e.g., the amino acid is immediately adjacent to one of the CDRs). See, U.S. Patent No. 5,530,101 and 5,585,089, incorporated 15 herein by reference, which provide detailed instructions for construction of humanized antibodies. Although humanized antibodies often incorporate all six CDRs (e.g., as defined by Kabat, but often also including hypervariable loop H1 as defined by Chothia) from a mouse antibody, they can also be made with fewer mouse CDRs and/or less than the complete mouse CDR sequence (e.g., a functional fragment of a CDR) (e.g., Pascalis *et al.*, J. 20 Immunol. 169:3076, 2002; Vajdos *et al.*, Journal of Molecular Biology, 320: 415-428, 2002; Iwahashi *et al.*, Mol. Immunol. 36:1079-1091, 1999; Tamura *et al.*, Journal of Immunology, 164:1432-1441, 2000).

Typically a humanized antibody as provided herein may include (i) a light chain 25 variable region comprising at least one CDR (often three CDRs) from a mouse antibody (also referred to herein as a mouse CDR) and a human variable region framework; and (ii) a heavy chain variable region comprising at least one CDR (often three CDRs) from the mouse antibody and a human variable region framework (FR). The light and heavy chain variable region frameworks (FRs) may each be a mature human antibody variable region framework sequence (or fragment thereof), a germline variable region framework sequence (combined with a J region sequence) (or fragment thereof), or a consensus sequence of a human antibody variable region framework sequence (or fragment thereof). Optionally, the humanized antibody includes a light chain variable region as described in (i), a heavy chain variable 30

region as described in (ii) together with a light chain human constant region and a heavy chain human constant region.

A chimeric antibody is an antibody in which the variable region of a mouse (or other rodent) antibody is combined with the constant region of a human antibody; the construction of a chimeric antibody by means of genetic engineering is well-known. Such antibodies retain the binding specificity of the mouse antibody, while being about two-thirds human. The proportion of nonhuman sequence present in mouse, chimeric and humanized antibodies suggests that the immunogenicity of chimeric antibodies is intermediate between mouse and humanized antibodies. Other types of genetically engineered antibodies that may have reduced immunogenicity relative to mouse antibodies include human antibodies made using phage display methods (Dower et al., WO91/17271; McCafferty et al., WO92/001047; Winter, WO92/20791; and Winter, FEBS Lett. 23:92, 1998, each of which is incorporated herein by reference) or using transgenic animals (Lonberg et al., WO93/12227; Kucherlapati WO91/10741, each of which is incorporated herein by reference).

Other approaches to design humanized antibodies may also be used to achieve the same result as the methods in U.S. Patent No. 5,530,101 and 5,585,089. For example, superhumanization as described in Tan et al. J. Immunol. 169: 1119, 2002, and U.S. Patent No. 6,881,557) or the method of Studnicak et al., Protein Eng. 7:805, 1994. Moreover, other approaches to produce genetically engineered, reduced-immunogenicity mAbs include reshaping, hyperchimerization and veneering/ resurfacing, as described, e.g., in Vaswami et al., Annals of Allergy, Asthma and Immunology 81:105, 1998; Roguska et al. Protein Eng. 9:895, 1996; and US Patent Nos. 6,072,035 and 5,639,641.

Disclosed are materials, compositions, and components that can be used for, can be used in conjunction with, can be used in preparation for, or are products of the disclosed methods and compositions. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each various individual and collective combinations and permutations of these compounds may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a method is disclosed and discussed and a number of modifications that can be made to a number of molecules including the method are discussed, each and every combination and permutation of the method, and the modifications that are possible are specifically contemplated unless specifically indicated to the contrary. Likewise, any subset or

combination of these is also specifically contemplated and disclosed. This concept applies to all aspects of this disclosure including, but not limited to, steps in methods using the disclosed compositions. Thus, if there are a variety of additional steps that can be performed, it is understood that each of these additional steps can be performed with 5 any specific method steps or combination of method steps of the disclosed methods, and that each such combination or subset of combinations is specifically contemplated and should be considered disclosed.

The examples below are intended to further illustrate certain aspects of the methods and compositions described herein, and are not intended to limit the scope of the 10 claims.

EXAMPLES

Example 1. Novel BAFF-receptor antibody to natively folded recombinant protein eliminates drug resistant human B-cell malignancies *in vivo*.

Conventional recombinant immunogen proteins produced in bacteria for developing 15 mAbs lack post-translational modifications and are simplistically folded because compared with eukaryotes, prokaryotes lack chaperone proteins and oxidizing environments. As a result, such proteins may differ in conformational structure from the corresponding plasma membrane-anchored native proteins. Furthermore, antibodies may be raised against off-target domains such as transmembrane or intracellular domains of the target protein. As described 20 herein, a strategy of generating mAbs against a natively folded, glycosylated immunogen expressed on eukaryotic cells was applied. Specifically, human BAFF-R as a native protein on mouse fibroblast cells, and used the engineered cell clone as an immunogen in mice. Described herein is the generation of novel mAbs that specifically bound and lysed human 25 malignant B cell lines and primary lymphomas in vitro, and inhibited growth of drug-resistant lymphoma cell lines in xenogenic tumor models *in vivo*.

Materials and Methods

Animals, cell lines, and primary human tumor samples. BALB/c mice for antibody development and NOD scid gamma (NSG) breeding pairs were purchased from The Jackson 30 Laboratory (Bar Harbor, ME). The NSG breeding colony was maintained by the Animal Resource Center at City of Hope. Mice were housed in a pathogen-free animal facility according to institutional guidelines. All animal studies were approved by the Institutional Animal Care and Use Committee (IACUC: 15020). JeKo-1, SU-DHL-6, Raji, U266 and RL

were purchased from ATCC (Manassas, VA). Z-138 line was provided by Dr. Michael Wang (MD Anderson Cancer Center). Ibrutinib-resistant SP49-IR line was developed and provided by Dr. Jianguo Tao (University of South Florida). Ibrutinib-resistant SP49 cell lines (SP49-IR) were established by treating cells with escalating doses of ibrutinib. IC50 was 5 nM for 5 parental SP49 compared to >100 nM for SP49-IR. At 100 nM ibrutinib ~5% of SP49 cells were viable compared with >90% of SP49-IR cells. Human NK-92 176V cells were obtained from Conkwest Inc. (San Diego, CA). For human blood and tumor samples, non-cultured, primary human lymphomas were obtained as cryopreserved, viable single cell suspensions in 10% DMSO from the Lymphoma Satellite Tissue Bank at MD Anderson Cancer Center 10 under an Institutional Review Board approved protocol (IRB: 2005-0656). Primary patient samples included leukapheresis or blood from patients with mantle cell lymphoma (MCL) or chronic lymphocytic leukemia (CLL), and excised lymph nodes from patients with diffuse large B-cell lymphoma (DLBCL) or follicular lymphoma (FL). Tumor cells in each sample ranged from 80% to 98% for leukapheresis or blood, and from 50% to 60% for lymph node 15 biopsies. Peripheral blood mononuclear cells (PBMC) was provided by the Michael Amini Transfusion Medicine Center at City of Hope (IRB: 15283).

Generation of human-BAFF-R expressing mouse fibroblast cells. Human BAFF-R (hBAFF-R) cDNA was from human B cells and cloned in-frame with GFP gene on pEGFP-N1 vector (Takara/Clotech, Mountain View, CA). hBAFF-R cDNA sequence was confirmed 20 against the NCBI gene sequence database (Gene ID: 115650). The cDNA encoding hBAFF-R-GFP fusion was subsequently cloned into a lentivirus gene delivery system (pLenti6/V5-DEST Gateway Vector kit, Life Technologies, Grand Island, NY) to produce hBAFF-R-GFP fusion proteins when transduced into mouse fibroblast (L) cells. Single cell clones were established from sorted GFP-positive L cells, and (h)BAFF-R-GFP-expressing L cell clone 25 D2C was used in further studies.

Antibody-producing hybridomas. Two 6-week-old BALB/c mice were immunized with D2C cells by five subcutaneously injections at the foot pad once every three days. Blood samples were obtained from both mice to measure serum antibodies against D2C by ELISA. Splenic tissue was harvested on day 20. Harvested splenocytes were fused with Sp2/0 30 myeloma to establish hybridomas and ELISA screened for antibodies using D2C or parental L cell-coated plates. Immunization and hybridoma procedures were conducted at the Antibody Core Facility at MD Anderson Cancer Center.

Chimeric antibody production. cDNA from selected hybridomas encoding the variable regions of antibody light and heavy chains were engineered onto expression vectors containing respective human IgG1 constant regions. Vectors were co-transfected into the FreeStyle 293 Expression System (Life Technologies, Carlsbad, CA) according to manufacturer's directions. Antibodies in culture supernatant were purified by HiTrap Protein A affinity chromatography columns (GE Healthcare, Marlborough, MA) according to the manufacturer's directions.

5 Cytotoxicity assays. Target cells (L cells, human tumor lines, primary patient samples) were labeled with chromium-51 (51Cr, Perkin Elmer, Waltham, MA) for a 51Cr release assay. Briefly, antibodies and effectors (NK cells or complement serum standard [Sigma Aldrich, St. Louis, MO]), were added to labeled target cells and incubated up to 18 hours. NK cells were enriched from PBMC (NK cell enrichment kit, Stemcell Technologies, Vancouver, Canada). 51Cr released into supernatant was detected with a Wizard Automatic Gamma Counter (Perkin Elmer).

10 15 Generation of JeKo-1-CD20-KO. FACS-sorted, stable JeKo-1-CD20-KO were generated using CD20-CRISPR/Cas9 and HDR Plasmid Systems (Santa Cruz Biotechnology, Santa Cruz, CA) according to manufacturer's directions. CD20 knock-out was verified by flow cytometry and Western blots.

20 In vivo studies. For tumor models, stable, luciferase-expressing tumors lines were established for bioluminescent imaging in mouse models. Briefly, a luciferase gene was introduced into tumor lines by a lentivirus gene delivery system (pLenti7.3/V5-DEST Gateway Vector Kit, Life Technologies, Carlsbad, CA). The minimum lethal dose per mouse was determined for each tumor cell line by dose titration. Tumor cells were injected intravenously (IV) and mice were monitored by in vivo bioluminescence imaging for 25 minimum tumor dose to ensure engraftment. Minimum lethal tumor doses were: 1×10^6 JeKo-1, 5×10^5 RS4;11, 5×10^5 JeKo-1-CD20-KO, or 2.5×10^4 Z-138 cells.

30 Bioluminescent Imaging: Mice were anesthetized with isoflurane and administered 150 mg/kg D-luciferin (Life Technologies, Carlsbad, CA) via intraperitoneal (IP) injection 10 minutes prior to imaging. Imaging was performed on an AmiX imaging system (Spectral Instruments Imaging, Tucson, AZ).

Antibody Studies: Mice (n=5 per group) were IV tumor challenged three days prior to four treatments once every five days. Treatments were 300 μ L IV injection: 200 μ g treatment antibody, 10×10^6 effector human NK-92-176V cells, and 5×10^4 IU IL-2 (Prometheus

Laboratories, San Diego, CA). Control groups received the same volume injections with a control antibody or without the antibody and/or NK cells. Bioluminescent imaging was performed weekly up to 80 days. Survival was tracked up to 100 days post tumor challenge.

Results

5 Generation of monoclonal antibodies against human BAFF-R. In order to generate a therapeutic antibody to a biologically relevant epitope of BAFF-R, a eukaryotic cell-surface expression system was used in which endogenous cell-surface proteins are presented in their native conformation with appropriate post-translational modifications. A mouse fibroblast (L) cell clone was engineered expressing cell-surface GFP-tagged, human BAFF-R. BAFF-R-
10 expressing L cell clones were generated and characterized for GFP expression (Figure 1A). Clone D2C was expanded and successfully used to immunize BALB/c mice according to Methods and immunization schedule in Figure 7A.

15 After generating and screening hybridoma clones, four clones (53, 55, 67, and 90) were identified as producing antibodies that specifically bound BAFF-R-expressing, but not parental, L cells (Figure 7B). Supernatants of all four clones contained antibodies that bound BAFF-R-expressing Mino cell line (MCL) in a dose-dependent manner. No antibody binding was detected in BAFF-R-negative control cell line, 293T (Figure 8).

20 Antibodies from the four hybridoma supernatants were purified by protein A affinity chromatography. Purified antibodies bound Mino cells in a dose-dependent manner (Figure 9), as well as other human MCL lines, including JeKo-1, REC-1, and ibrutinib-resistant JVM-13 and Z-138 (Figure 1B).

25 An analysis of the complementarity determining regions (CDRs) on the four antibodies revealed that clones 53, 55, and 67 had nearly identical sequences, whereas clone 90 was unique. Therefore, clones 55 and 90 were selected for further investigation. Both clones 55 and 90 effectively bound JeKo-1 (MCL), SU-DHL-6 (DLBCL), Raji (Burkitt lymphoma), and RL (FL) at both high ($2 \mu\text{g}/10^6$ cells) and low ($0.05 \mu\text{g}/10^6$ cells) concentrations (Figure 10).

30 Chimeric mAb against human BAFF-R induced anti-tumor effects both in vitro and in vivo. Clones 55 and 90 were further developed into their respective chimeric mAbs containing human IgG1 constant regions (termed C55 and C90). The chimeric antibodies retained specific dose-dependent binding to BAFF-R-expressing L cells (Figure 1C). C55 and C90 were conjugated to Alexa Fluor 488 and exhibited direct binding to non-Hodgkin lymphoma (NHL) lines JeKo-1, SU-DHL-6, Raji, and RL (Figure 1D). Importantly, chimeric

mAbs readily bound MCL, DLBCL, and FL patient primary tumor samples (Figure 1E and Figure 11).

5 C55 and C90 elicited antibody-dependent cell-mediated cytotoxicity (ADCC) specifically against BAFF-R-expressing L cells and JeKo-1, but not BAFF-R negative L cells nor the BAFF-R negative human multiple myeloma line, U266 (Figure 2A, Figure 12). In contrast, antibodies did not elicit in vitro complement dependent cytotoxicity (CDC, Figure 2B). Cytotoxicity required the addition of NK cells, as shown for SU-DHL-6, Raji, and RL lymphoma cell lines (Figure 2C and Figure 13), suggesting ADCC as a principal mechanism of antibody-mediated cytotoxicity. Importantly, chimeric antibodies elicited ADCC against 10 primary patient tumor samples (Figure 3A).

The antibodies inhibited BAFF/BAFF-R binding in a dose-dependent manner (Figure 14), suggesting potential disruption of BAFF/BAFF-R survival signaling in tumor cells. Furthermore, C55 and C90 exhibited limited internalization upon binding BAFF-R (Figure 15).

15 In vivo, NSG mice were challenged with luciferase knock-in JeKo-1 MCL cell line followed by antibody treatments. Treatment followed the schedule in Figure 4A. Mice receiving either C55 or C90 demonstrated significant retardation of tumor growth, compared with PBS or NK cells alone control groups (Figure 4B). Similarly, C55 and C90 also markedly retarded tumor growth in RS4;11 (acute lymphoblastic leukemia, ALL) challenged 20 NSG mice, compared with no inhibition by rituximab or controls (Figure 4C).

25 Chimeric mAb induced potent antitumor effects against drug-resistant lymphomas in vitro and in vivo. The antibodies were further tested against primary CLL (n=3) and MCL (n=2) samples from patients who had been previously treated with rituximab. All five primary samples were sensitive to killing by ADCC with C55 and C90, suggesting their effectiveness against tumors which progressed clinically after exposure to rituximab (Figure 3B).

30 In order to create a model of drug-resistant lymphomas, a stable CD20 knock-out (KO) clone of JeKo-1 was generated using a CRISPR/HDR system. CD20-KO clones were confirmed for absence of CD20 surface expression by flow cytometry and Western blotting (Figure 5A and Figures 16A and 16B) and the presence of BAFF-R surface expression by flow cytometry (Figure 16C). JeKo-1-CD20-KO clone 25, selected for further studies, retained sensitivity to C55- and C90-mediated ADCC, but became insensitive to cytotoxicity mediated by anti-CD20 rituximab (Figure 5B).

As a second model of drug-resistant lymphomas, the chimeric BAFF-R mAb was tested for ADCC against the naturally ibrutinib resistant human MCL line, Z-138, and the induced ibrutinib resistant MCL line, SP49-IR, which had been induced in vitro for resistance to ibrutinib (see Methods). Significant in vitro ADCC was observed with the antibodies 5 against both ibrutinib resistant lines (Figure 5C).

Finally, three days following IV challenge with JeKo-1-CD20-KO tumor cells in vivo, NSG mice (n=5 per group) received BAFF-R antibody treatments (C55 or C90) or rituximab as described in Methods and according to the schedule in Figure 4A.

Bioluminescent imaging on Day 20 revealed substantial tumor burden in controls and 10 rituximab treated mice, but no visible tumors in BAFF-R antibody treatment groups (Figure 6A). Monitoring tumor free and long-term overall survival confirmed the significant antitumor effects of both BAFF-R antibodies, but not rituximab (Figure 6C). Similarly, significant effects were observed following treatment of ibrutinib-resistant Z-138 tumor-bearing mice with either BAFF-R antibody, compared with controls (PBS or NK only) 15 (Figure 6B-C).

BAFF-R mAbs also bind normal B cells. When tested against normal PBMC, anti-BAFF-R antibody C90 exhibited specific binding against B cells, as expected, without staining any T cells, NK cells, granulocytes, or monocytes (Figure 17). The positive staining results were verified on purified B cells (Figure 18). Again, purified T cells, NK cells, and 20 gated myeloid cells showed no binding.

Expanding our scope, immunohistochemistry studies showed positive staining of our antibodies on tonsil and spleen samples, leaving all other vital organs, including heart, lung, kidney, and brain, untouched (Figures 19A and 19B).

Discussion

25 The provided BAFF-R mAbs elicited robust in vivo antitumor effects as a single agent against multiple B-cell tumor types, including NHL, CLL, and ALL. Furthermore, the antibodies eradicated established tumors, which led to long-term, tumor-free survival in vivo.

The distinctive features of the BAFF-R mAbs may be due to the approach used to 30 generate them. The provided approach was to express human BAFF-R as a native surface protein on mouse fibroblast cells for immunization, increasing the likelihood of presenting a natively folded, glycosylated immunogen. Therefore, it is very likely that the antibodies are binding an accessible human BAFF-R epitope distinct from the other antibodies described. Thus, a technical strategy was demonstrated for generating monoclonal antibodies against a

natively folded, eukaryotically glycosylated human BAFF-R that is able to specifically bind, lyse, and inhibit B-cell tumors in vivo. The results suggest the main anti-tumor mechanism of our mAbs is ADCC, as NK cells were required in addition to mAbs for in vitro activity (Figure 2); no evidence of CDC was observed. Both antibodies were able to competitively 5 inhibit BAFF ligand binding to BAFF-R (Figure 14).

One clinically relevant mechanism of resistance to rituximab is down-regulation of CD20. This phenomenon of drug resistance was modeled with a CRISPR edited MCL line, JeKo-1, which is deficient in CD20. The significant in vivo antitumor effects of C55 or C90, but not rituximab treatment, against this line and similarly against the naturally 10 ibrutinib resistant Z-138 MCL suggests efficacy against drug-resistant lymphomas (Figure 5). Taken together with the in vitro cytotoxicity of these antibodies against primary tumors from lymphoma patients who were previously treated with, and progressed in response to rituximab, these data suggest C55 and C90 as a potential treatment strategy to overcome drug 15 resistance (Figure 3).

15 **Example 2. Humanizing BAFF-R mAb.**

The chimeric antibody, clone 90, was humanized while retaining its binding specificity and cytotoxic effects. Through computational analysis of the CDRs and predicted structure, three variants of the heavy and three variants of the light chain were produced with varying degrees of likeness to human antibodies. A total of nine combinational variants was 20 constructed from the humanized heavy and light chains. These variants all proved comparable in binding affinity to the parental chimeric antibody with K_D values ranging from 2.6 to 5.0 nM (Table 1).

TABLE 1. Binding Affinity of Humanized BAFF-R Antibody Variants

Loading Sample ID	Sample ID	KD (M)	kon(1/Ms)	kdis(1/s)	Full R ²	Full X ²
Chimeric Parental	BAFF-R	3.0E-09	9.9E+05	2.0E-03	0.0049	0.9809
Humanized HC1 + LC1	BAFF-R	4.0E-09	6.1E+05	2.4E-03	0.0114	0.9619
Humanized HC1 + LC2	BAFF-R	3.7E-09	6.2E+05	2.3E-03	0.005	0.9800
Humanized HC1 + LC3	BAFF-R	5.0E-09	4.1E+05	2.1E-03	0.0125	0.9565

Humanized HC2 + LC1	BAFF-R	2.6E-09	7.7E+05	2.0E-03	0.0051	0.9790
Humanized HC2 + LC2	BAFF-R	3.6E-09	6.1E+05	2.2E-03	0.0047	0.9799
Humanized HC2 + LC2	BAFF-R	3.2E-09	8.2E+05	3.1E-03	0.0045	0.9821
Humanized HC3 + LC1	BAFF-R	3.8E-09	8.2E+05	3.1E-03	0.0045	0.9821
Humanized HC3 + LC2	BAFF-R	3.1E-09	1.2E+06	3.7E-03	0.0074	0.9793
Humanized HC3 + LC3	BAFF-R	3.4E-09	6.8E+05	2.3E-03	0.0025	0.9900

The nine candidate antibodies were further assessed and a lead candidate was determined. The binding of the humanized antibodies were observed to have specificity to BAFF-R and all had similar relative binding in a dose dependent manner (FIG. 20A).

5 Additionally, the humanized antibodies were assessed for their ADCC effects. Again, it was found that the humanized candidates maintained their specific cytotoxicity, and performed equally well compared to the chimeric controls and rituximab (FIG. 20B).

Humanized clone 90 variants 4 and 5 were selected for further *in vitro* testing. The humanized antibody variants were biotinylated and visualized with a fluorescent streptavidin 10 probe. Their binding against various non-Hodgkin's lymphoma, lymphoblastic leukemia, and multiple myeloma lines were assessed, including JeKo-1, Ly-10, MEC-2, RL, RS4, Raji, Z138, and U266 (FIG. 22A). The flow cytometry results reveal a significant binding to each 15 of these cell lines. Further flow analysis with the humanized variants against normal PBMCs show specificity in the binding. When assessed for the binding of granulocytes, monocytes, B cells, T cells, and NK cells in normal healthy PBMCs, the antibodies only bind the B cells population (FIG. 22B).

The two variants were further assessed in their ability to initiate ADCC. The antibodies were administered in varying concentration to JeKo-1, Z138, and RS4 lines after a period of chromium uptake. The cells and antibodies were incubated with effector NK cells. 20 The supernatant was analyzed 6 hours post treatment (FIG. 21A). The antibodies have a clear cytotoxic effect against tumor lines and demonstrate a dose dependency with each 10 fold

dilution. The results are comparable to that of rituximab but can also be seen in RS4 acute lymphoblastic lymphoma, where rituximab is not active. Further assays with LY-10, MEC-2, RL, and Raji (FIG. 21B) continue to demonstrate the potency of the humanized antibody treatment. All results found were comparable to that of the current conventional treatment 5 with rituximab.

Example 3. Chimeric Antigen Receptor T Cell.

Antibodies with high binding affinity and bioactivity were used to construct chimeric antigen receptor (CAR) T cells for *in vivo* studies. DNA sequences for heavy and light chain variable domains were arranged into a single chain (sFv) format and engineered to a T cell 10 signaling domain (δ chain) along with a 4-1BB motif. The engineered CAR gene was introduced into purified healthy donor derived CD8+ T cells along with a co-expressing GFP via a lentivirus. CAR-T cells were cell sorted for their expression of GFP and expanded in vitro with CD3 and CD28 beads for animal studies. NSG mice were challenged with a luciferase expressing JeKo-1 MCL line (JeKo-1-luci). The tumor was allowed to develop 15 and monitored by bioluminescent imaging until a visible population of tumor cells were observed; approximate 9 days post tumor challenge. Mice were administered two dose of 5×10^6 CAR-T cells (anti-BAFF-R and anti-CD19) on days 9 and 15 post tumor challenge. Control groups received untreated T cells or saline (PBS). Mice were monitored closely and imaged every 3 days to track the tumor development in order to evaluate the therapeutic anti- 20 tumor effects of CAR-T therapy.

Humanized anti-BAFF-R mAbs were further assessed against primary patient tumor samples for their binding and cytotoxicity. Three mantel cell lymphoma patient samples were characterized with a majority of the tumor cells expressing BAFF-R. Flow cytometry results reveal distinct populations of these primary tumor cells that were bound by our 25 humanized antibodies (FIG. 22A). Furthermore, chromium release cytotoxicity assays on the same primary tumor samples revealed high specific killing compared to controls. The results were comparable to the effects of rituximab and consistent with the chimeric antibodies developed earlier (FIG. 22B). The cell type specificity of the humanized antibodies was determined by assessing their binding to normal PBMC. No appreciable binding was noted 30 for the major groups of PBMCs including granulocytes, monocytes, T cells and NK cells. The B cell population was the only detectable population that was bound by the antibody (FIG. 23). The results of the assay is also consistent with the previously characterized chimeric antibodies.

The anti-BAFF-R mAbs were further used to create chimeric antigen receptor (CAR) T cells. Experiments utilized the chimeric C55 variable region engineered into a single chain (sFv) format. The anti-BAFF-R C55 sFv was attached to a T cell receptor signaling domain containing a 4-1BB motif and successfully introduced into healthy normal human donor 5 CD8+ T cells isolated from PBMC. The CAR-T cells were administered to tumor bearing mice with an appreciable tumor burden (FIG. 24). Mice treated with the anti-BAFF-R CAR-T cells had significant tumor clearance when compared to either saline or non-engineered T cell control groups. Additionally, the anti-tumor effects of our CAR-T cells are comparable to those of the anti-CD-19 CAR-T treated group.

10 The chimeric anti-BAFF-R antibody C90 was humanized with several variants. The humanization process took into consideration and analysis of the variable region and specifically the CDRs of the chimeric antibody. From there, three variants for each heavy and light chain was developed with varying degrees of human-likeness ranging from 1 the most human to 3 the most conservative to the chimeric. The variants were combined to produce 9 15 variants. Biacore analysis was performed on each variant as well as the chimeric parental C90 to determine their equilibrium dissociation constant K_D . The antigen was commercial, recombinant extracellular domain of human BAFF-R.

20

WHAT IS CLAIMED IS:

1. A B cell activating factor receptor (BAFF-R) antibody comprising a light chain variable region and a heavy chain variable region,
wherein said light chain variable region comprises:
a CDR L1 as set forth in SEQ ID NO:1, a CDR L2 as set forth in SEQ ID NO:2 and a CDR L3 as set forth in SEQ ID NO:3; and
wherein said heavy chain variable region comprises:
a CDR H1 as set forth in SEQ ID NO:4, a CDR H2 as set forth in SEQ ID NO:5, and a CDR H3 as set forth in SEQ ID NO:6.
2. The antibody of claim 1, wherein said antibody is a humanized antibody.
3. The antibody of one of claims 1-2, wherein said light chain variable region comprises the sequence of SEQ ID NO:18, SEQ ID NO:20 or SEQ ID NO:22.
4. The antibody of one of claims 1-5, wherein said heavy chain variable region comprises the sequence of SEQ ID NO:24, SEQ ID NO:26 or SEQ ID NO:28.
5. The antibody of claim 1, wherein said antibody is a chimeric antibody.
6. The antibody of claim 5, wherein said light chain variable region comprises the sequence of SEQ ID NO:14.
7. The antibody of claim 5 or 6, wherein said heavy chain variable region comprises the sequence of SEQ ID NO:16.
8. A B cell activating factor receptor (BAFF-R) antibody comprising a light chain variable region and a heavy chain variable region,
wherein said light chain variable region comprises:
a CDR L1 as set forth in SEQ ID NO:7, a CDR L2 as set forth in SEQ ID NO:8 and a CDR L3 as set forth in SEQ ID NO:9; and
wherein said heavy chain variable region comprises:
a CDR H1 as set forth in SEQ ID NO:10, a CDR H2 as set forth in SEQ ID NO:11, and a CDR H3 as set forth in SEQ ID NO:12.

9. The antibody of claim 8, wherein said antibody is a humanized antibody.
10. The antibody of claim 8, wherein said antibody is a chimeric antibody.
11. The antibody of claim 10, wherein said light chain variable region comprises the sequence of SEQ ID NO:30.
12. The antibody of claim 10 or 11, wherein said heavy chain variable region comprises the sequence of SEQ ID NO:32.
13. The antibody of one of claims 1-12, wherein said antibody is an IgG.
14. The antibody of one of claims 1-13, wherein said antibody is an IgG1.
15. The antibody of one of claims 1-12, wherein said antibody is a Fab' fragment.
16. The antibody of one of claims 1-12, wherein said antibody is a single chain antibody (scFv).
17. The antibody of one of claims 1-16, wherein said antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 5 nM.
18. The antibody of one of claims 1-17, wherein said antibody is capable of binding a BAFF-R protein with an equilibrium dissociation constant (K_D) of less than about 4 nM.
19. The antibody of one of claims 1-18, wherein said antibody is bound to a BAFF-R protein.
20. The antibody of claim 19, wherein said BAFF-R protein is a human BAFF-R protein.
21. The antibody of claim 18 or 19, wherein said BAFF-R protein forms part of a cell.

22. The antibody of claim 21, wherein said BAFF-R protein is expressed on the surface of said cell.
23. The antibody of claim 21 or 22, wherein said cell is a lymphoid cell.
24. The antibody of one of claims 21-23, wherein said cell is a B cell.
25. The antibody of one of claims 21-24, wherein said cell is a cancer cell.
26. The antibody of claim 25, wherein said cancer cell is a lymphoma cell.
27. An isolated nucleic acid encoding an antibody of one of claims 1-26.
28. A pharmaceutical composition comprising a therapeutically effective amount of an antibody of one of claims 1-26 and a pharmaceutically acceptable excipient.
29. A chimeric antigen receptor (CAR) comprising an antibody of one of claims 1-26 or a functional fragment thereof.
30. A method of treating cancer in a subject in need thereof, said method comprising administering to a subject a therapeutically effective amount of the chimeric antigen receptor of claim 29, thereby treating cancer in said subject.
31. A method of treating cancer in a subject in need thereof, said method comprising administering to a subject a therapeutically effective amount of an antibody of one of claims 1-26, thereby treating cancer in said subject.
32. The method of claim 30 or 31, wherein said cancer is lymphoma, leukemia or myeloma.
33. The method of claim 32, wherein said lymphoma is mantle cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, marginal zone lymphoma or Burkitt's lymphoma.
34. The method of claim 32, wherein said leukemia is lymphoblastic leukemia, chronic lymphocytic leukemia or hairy cell leukemia.
35. The method of claim 32, wherein said myeloma is multiple myeloma.

36. The method of one of claim 30-32, said method further comprising administering to said subject a second therapeutic agent.

37. The method of claim 36, wherein said therapeutic agent is a chimeric monoclonal antibody capable of binding a CD 20 antigen.

38. The method of claim 36 or 37, wherein said therapeutic agent is rituximab.

39. A method of treating an autoimmune disease in a subject in need thereof, said method comprising administering to said subject a therapeutically effective amount of an antibody of one of claims 1-26, thereby treating an autoimmune disease in said subject.

40. The method of claim 39, wherein said autoimmune disease is rheumatoid arthritis, systemic Lupus erythematosus, multiple sclerosis, glomerulonephritis, Sjögren's Syndrome or autoimmune hemolytic anemia.

41. The method of claim 40, said method further comprising administering to said subject a second therapeutic agent.

42. A method of inhibiting proliferation of a cell, said method comprising:

- (i) contacting a cell with a BAFF-R antibody of one of claims 1-26, thereby forming a contacted cell; and
- (ii) allowing said BAFF-R antibody to bind a BAFF-R protein on said contacted cell, thereby inhibiting proliferation of said cell.

43. The method of claim 42, wherein said cell is a lymphoid cell.

44. The method of claim 42 or 43, wherein said cell is a B cell.

45. The method of one of claims 42-44, wherein said cell is a cancer cell.

46. The method of one of claims 42-45, wherein said cell is a lymphoma cell.

47. A humanized B cell activating factor receptor (BAFF-R) antibody capable of binding BAFF-R with a K_D of less than about 4 nM.

48. A humanized B cell activating factor receptor (BAFF-R) antibody bound to a BAFF-R at a K_D of less than about 4 nM.

49. The humanized antibody of claim 47 or 48, wherein said antibody does not induce BAFF-R activity.

50. A mouse fibroblast cell expressing a human BAFF-R protein, wherein said human BAFF-R protein is expressed on the cell surface of said cell.

51. The mouse fibroblast cell of claim 50, wherein said human BAFF-R protein comprises a detectable moiety.

52. The mouse fibroblast cell of claim 51, wherein said detectable moiety is a fluorescent moiety.

53. The mouse fibroblast cell of claim 51 or 52, wherein said detectable moiety is an enhanced green fluorescent protein (eGFP).

54. A method of producing an anti-human BAFF-R antibody, said method comprising:

(i) administering a mouse fibroblast cell of one of claims 50-53 to a mouse, thereby forming an immunized BAFF-R mouse;

(ii) fusing a splenic cell from said immunized BAFF-R mouse with a human myeloma cell, thereby forming a BAFF-R hybridoma cell; and

(iii) allowing said BAFF-R hybridoma cell to express a BAFF-R antibody, thereby producing an anti-BAFF-R antibody.

55. The method of claim 54, wherein said anti-BAFF-R antibody is an antibody of claim 1 or claim 8.

(h) BAFF-R-expressing L
cell clone D2C

Parental L cells

1/45





FIG. 1A

2/45

Mino Jeko-1 REC-1 JVM-13 Z-138 293T



SUBSTITUTE SHEET (RULE 26)

FIG. 1B

C55 C90




FIG. 1C

4/45






FIG. 1D

5/45





EIG. 2A

7/45




FIG. 2B

8/45





FIG. 2C



10/45




FIG. 3B

11/45




FIG. 4A




FIG. 4B

13/45





FIG. 4C



15/45




FIG. 5B




FIG. 5C

17/45




FIG. 6A




FIG. 6B

19/45




FIG. 6C

20/45




FIG. 7A

21/45

Hybridoma clones screened by ELISA for antibody production			
Clone	(h)BAFF-R-expressing cells (OD 450 nm)	Parental L cell (OD 450 nm)	
C53	0.552	0.095	
C55	1.067	0.102	
C67	0.615	0.093	
C90	0.645	0.116	
C39	0.137	0.091	

FIG. 7B




FIG. 8

23/45




FIG. 9

24/45




FIG. 10




FIG. 11

26/45




FIG. 12

27/45




FIG. 13

28/45




FIG. 14





FIG. 15

30/45



31/45





FIG. 16B



33/45




FIG. 17A

34/45




FIG. 17B




FIG. 18




FIG. 19A




FIG. 19B

38/45




FIG. 20A

39/45




FIG. 20B

40/45




FIG. 21A




FIG. 21B

42/45




FIG. 22A




FIG. 22B




FIG. 23

45/45




FIG. 24