




(86) Date de dépôt PCT/PCT Filing Date: 2003/05/16
(87) Date publication PCT/PCT Publication Date: 2003/11/27
(45) Date de délivrance/Issue Date: 2012/02/28
(85) Entrée phase nationale/National Entry: 2004/11/09
(86) N° demande PCT/PCT Application No.: GB 2003/002102
(87) N° publication PCT/PCT Publication No.: 2003/097834
(30) Priorités/Priorities: 2002/05/17 (GB0211369.4);
2002/12/17 (US10/321,195)

(51) Cl.Int./Int.Cl. *C12N 15/10* (2006.01),
C12Q 1/68 (2006.01)
(72) Inventeurs/Inventors:
FUREBRING, CHRISTINA, SE;
CARLSSON, ROLAND, SE;
BORREBAECK, CARL ARNE KRISTER, SE;
MALMBORG HAGER, ANN-CHRISTIN, SE
(73) Propriétaire/Owner:
ALLIGATOR BIOSCIENCE AB, SE
(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : METHODE DE DEVELOPPEMENT MOLECULAIRE IN VITRO D'UNE FONCTION DE PROTEINE
(54) Title: A METHOD FOR IN VITRO MOLECULAR EVOLUTION OF PROTEIN FUNCTION



(57) Abrégé/Abstract:

A method for in vitro molecular evolution of protein function The invention provides a method for generating a polynucleotide sequence or population of sequences from parent single-stranded polynucleotide sequences encoding one or more protein motifs,

(57) Abrégé(suite)/Abstract(continued):

comprising the steps of (a) providing a first population of single-stranded polynucleotide molecules and a second population of single-stranded polynucleotide molecules, the first and second populations together constituting plus and minus strands of parent polynucleotide sequences, (b) carrying out a reaction for digesting the first and second populations of single-stranded polynucleotide molecules with an exonuclease to generate corresponding populations of single-stranded polynucleotide fragments, (c) contacting said fragments generated from the plus strands with fragments generated from the minus strands and optionally, adding primer sequences that anneal to the 3' and 5' ends of at least one of the parent polynucleotides under annealing conditions, and (d) amplifying the fragments that anneal to each other to generate at least one polynucleotide sequence encoding one or more protein motifs having altered characteristics as compared to the one or more protein motifs encoded by said parent polynucleotides, wherein, in step (b), at least one parameter of the reaction used for digestion of the first population of single-stranded polynucleotide molecules is different from the equivalent parameter(s) used in the reaction for digestion of the second population of single-stranded polynucleotide molecules. Preferably, the reaction parameter is selected from exonuclease type, exonuclease concentration, reaction volume, duration of the digestion reaction, temperature of the reaction mixture, pH of the reaction mixture, length of parent single-stranded polynucleotide sequences, amount of single-stranded polynucleotide molecules and buffer composition of the reaction mixture.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 November 2003 (27.11.2003)

PCT

(10) International Publication Number
WO 2003/097834 A3

(51) International Patent Classification⁷: **C12N 15/10**,
C12Q 1/68

(21) International Application Number:
PCT/GB2003/002102

(22) International Filing Date: 16 May 2003 (16.05.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0211369.4 17 May 2002 (17.05.2002) GB
10/321,195 17 December 2002 (17.12.2002) US

(71) Applicant (for all designated States except MN, US): **ALLIGATOR BIOSCIENCE AB** [SE/SE]; Scheelevägen 19a, S-223 70 Lund (SE).

(71) Applicant (for MN only): **THOMAS, Philip, John, Duval** [GB/GB]; Park View House, 58 The Ropewalk, Nottingham NG1 5DD (GB).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **FUREBRING, Christina** [SE/SE]; Iliongränd 147, S-224 72 Lund (SE). **CARLSSON, Roland** [SE/SE]; Stenåldersvägen 99, S-226 54 Lund (SE). **BORREBAECK, Carl, Arne, Krister** [SE/SE]; Åttevägen 8A, S-245 62 Hjärup (SE). **MALMBORG, Hager, Ann-Christin** [SE/SE]; Jönköpingsgatan 83, 252 50 Helsingborg (SE).

(74) Agent: **THOMAS, Philip, J., D.**; Eric Potter Clarkson, Park View House, 58 The Ropewalk, Nottingham, NG1 5DD (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

[Continued on next page]

(54) Title: A METHOD FOR IN VITRO MOLECULAR EVOLUTION OF PROTEIN FUNCTION



(57) Abstract: A method for *in vitro* molecular evolution of protein function. The invention provides a method for generating a polynucleotide sequence or population of sequences from parent single-stranded polynucleotide sequences encoding one or more protein motifs, comprising the steps of (a) providing a first population of single-stranded polynucleotide molecules and a second population of single-stranded polynucleotide molecules, the first and second populations together constituting plus and minus strands of parent polynucleotide sequences, (b) carrying out a reaction for digesting the first and second populations of single-stranded polynucleotide molecules with an exonuclease to generate corresponding populations of single-stranded polynucleotide fragments, (c) contacting said fragments generated from the plus strands with fragments generated from the minus strands and optionally, adding primer sequences that anneal to the 3' and 5' ends of at least one of the parent polynucleotides under annealing conditions, and (d) amplifying the fragments that anneal to each other to generate at least one polynucleotide sequence encoding one or more protein motifs

having altered characteristics as compared to the one or more protein motifs encoded by said parent polynucleotides, wherein, in step (b), at least one parameter of the reaction used for digestion of the first population of single-stranded polynucleotide molecules is different from the equivalent parameter(s) used in the reaction for digestion of the second population of single-stranded polynucleotide molecules. Preferably, the reaction parameter is selected from exonuclease type, exonuclease concentration, reaction volume, duration of the digestion reaction, temperature of the reaction mixture, pH of the reaction mixture, length of parent single-stranded polynucleotide sequences, amount of single-stranded polynucleotide molecules and buffer composition of the reaction mixture.

WO 2003/097834 A3

WO 2003/097834 A3



(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) **Date of publication of the international search report:**
19 February 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *with international search report*

A METHOD FOR IN VITRO MOLECULAR EVOLUTION OF PROTEIN FUNCTION

5

The present invention relates to a method for *in vitro* molecular evolution of protein function, in particular by shuffling of single-stranded DNA segments obtained using a nuclease.

10 Protein function can be modified and improved *in vitro* by a variety of methods, including site directed mutagenesis (Alber *et al.*, Nature, 5; 330(6143):41-46, 1987) combinatorial cloning (Huse *et al.*, Science, 246:1275-1281, 1989; Marks *et al.*, Biotechnology, 10: 779-783, 1992) and random mutagenesis combined with appropriate selection systems (Barbas
15 *et al.*, PNAS. USA, 89: 4457-4461, 1992).

The method of random mutagenesis together with selection has been used in a number of cases to improve protein function and two different strategies exist. Firstly, randomisation of the entire gene sequence in combination
20 with the selection of a variant (mutant) protein with the desired characteristics, followed by a new round of random mutagenesis and selection. This method can then be repeated until a protein variant is found which is considered optimal (Schier R. *et al.*, J. Mol. Biol. 1996 263 (4): 551-567). Here, the traditional route to introduce mutations is by error
25 prone PCR (Leung *et al.*, Technique, 1: 11-15, 1989) with a mutation rate of approximately 0.7%. Secondly, defined regions of the gene can be mutagenised with degenerate primers, which allows for mutation rates up to 100% (Griffiths *et al.*, EMBO. J, 13: 3245-3260, 1994; Yang *et al.*, J. Mol.

Biol. 254: 392-403, 1995). The higher the mutation rate used, the more limited the region of the gene that can be subjected to mutations.

Random mutation has been used extensively in the field of antibody engineering. *In vivo* formed antibody genes can be cloned *in vitro* (Larrick *et al.*, Biochem. Biophys. Res. Commun. 160: 1250-1256, 1989) and random combinations of the genes encoding the variable heavy and light genes can be subjected to selection (Marks *et al.*, Biotechnology, 10: 779-783, 1992). Functional antibody fragments selected can be further improved using random mutagenesis and additional rounds of selections (Schier R. *et al.*, J. Mol. Biol. 1996 263 (4): 551-567).

The strategy of random mutagenesis is followed by selection. Variants with interesting characteristics can be selected and the mutated DNA regions from different variants, each with interesting characteristics, are combined into one coding sequence (Yang *et al.*, J. Mol. Biol. 254: 392-403, 1995). This is a multi-step sequential process, and potential synergistic effects of different mutations in different regions can be lost, since they are not subjected to selection in combination. Thus, these two strategies do not include simultaneous mutagenesis of defined regions and selection of a combination of these regions.

Another process involves combinatorial pairing of genes which can be used to improve *e.g.* antibody affinity (Marks *et al.*, Biotechnology, 10: 779-783, 1992). Here, the three CDR-regions in each variable gene are fixed and this technology does not allow for shuffling of individual gene segments in the gene for the variable domain, for example, including the CDR regions, between clones.

The concept of DNA shuffling (Stemmer, Nature 370: 389-391, 1994) utilises random fragmentation of DNA and assembly of fragments into a functional coding sequence. In this process, it is possible to introduce chemically synthesised DNA sequences and in this way target variation to defined places in the gene which DNA sequence is known (Cramer *et al.*, Biotechniques, 18: 194-196, 1995). Stemmer and coworkers developed this *in vitro* method, which resembles the normal evolution process of protein in nature. The DNA shuffling generates diversity by recombination, combining useful mutations from individual genes. It has been used successfully for artificial evolution of different proteins, *e.g.* enzymes and cytokines (Chang *et al.* Nature Biotech. 17, 793-797, 1999; Zhang *et al.* Proc. Natl. Acad. Sci. USA 94, 4504-4509, 1997; Christians *et al.* Nature Biotech. 17, 259-264, 1999). The genes are randomly fragmented using DNase I and then reassembled by recombination with each other. The starting material can be either a single gene (first randomly mutated using error-prone PCR) or naturally occurring homologous sequences (so-called family shuffling). DNase I hydrolyses DNA preferentially at sites adjacent to pyrimidine nucleotides, therefore it is a suitable choice for random fragmentation of DNA. However, the activity is dependent on Mg or Mn ions, Mg ions restrict the fragment size to 50bp, while the Mn ions will give fragment sizes less than 50bp. Therefore, in order to have all possible sizes for recombination the gene in question needs to be treated at least twice with DNase I in the presence of either of the two different ions, followed by removal of these very same ions.

25

In theory, it is possible to shuffle DNA between any clones. However, if the resulting shuffled gene is to be functional with respect to expression and activity, the clones to be shuffled have preferably to be related or even identical, with the exception of a low level of random mutations. DNA

WU 03/097834

PCT/GB03/02102

4

shuffling between genetically different clones will generally produce non-functional genes. However, it has been proven by the methodology of ITCHY that interspecies fusion libraries can be created between fragments of the *E. coli* and human glycinamide ribonucleotide transformylase genes, which have only 50% identity on the DNA level (Ostermeier *et al.*, Nat Biotechnol 17, 1205-9, 1999).

A successful recombination of two different genes requires formation of hetero-duplex molecules. In some cases the family shuffling almost only form homo-duplexes resulting in a low frequency of recombination. This problem has been addressed by using DNase I-digested single-stranded DNA (Kikuchi *et al.* Gene 243,133-137 2000).

Single-stranded DNA can be obtained using methods known in the art. For example, biotinylated primers may be used in the PCR reactions in combination with e.g. Dynabeads[®] (Dynal, Norway) or AffiniTip[™] Streptavidin Capture Micro-columns (Genosys Biotechnologies Inc., The Woodlands, USA). Alternatively, single-stranded DNA can be obtained by utilising bacteriophage that are able to pack single-stranded DNA (Viruses and Related Entities in Modern Microbiology, Principles and Applications pp.171-192, Ed. E.A. Birge, Wm. C. Brown Publishers 1992; Sambrook *et al.* Molecular Cloning, A laboratory manual 2nd edition. Cold Spring Harbor Laboratory Press, 1989). In addition, asymmetric PCR methods may be used (see Example 1).

25

Selection of enzymes with altered and improved properties is often based on the actual function of the enzyme. For example, increased thermostability of an enzyme can be selected for by incubating transformed colonies at temperatures that cause inactivation of wild type enzyme. In addition,

improved β -glucosidase activity can be identified by using PNPG as the substrate (Arrizubieta *et al* J Biol Chem Jun 27, 2000).

Selection of functional proteins from molecular libraries has been
5 revolutionised by the development of the phage display technology
(Parmley *et al.*, Gene, 73: 305-391 1988; McCafferty *et al.*, Nature, 348:
552-554, 1990; Barbas *et al.*, PNAS. USA, 88: 7978-7982, 1991). Here, the
phenotype (protein) is directly linked to its corresponding genotype (DNA)
and this allows for direct cloning of the genetic material, which can then be
10 subjected to further modifications in order to improve protein function.
Phage display has been used to clone functional binders from a variety of
molecular libraries with up to 10^{11} transformants in size (Griffiths *et al.*,
EMBO. J. 13: 3245-3260, 1994). Thus, phage display can be used to
directly clone functional binders from molecular libraries, and can also be
15 used to improve further the clones originally selected. Other types of
viruses that have been used for surface expression of protein libraries and
selections thereof are baculovirus (Boublik *et al* Biotechnol 13:1079-1084.
1995; Mottershead *et al* Biochem Biophys Res Com 238:717-722, 1997;
Grabherr *et al* Biotechniques 22:730-735, 1997) and retrovirus (Buchholz *et*
20 *al* Nature Biotechnol 16:951-954, 1998).

Selection of functional proteins from molecular libraries can also be
performed by cell surface display. Also here, the phenotype is directly
linked to its corresponding genotype. Bacterial cell surface display has been
25 used for *e.g.* screening of improved variants of carboxymethyl cellulase
(CMCase) (Kim *et al* Appl Environ Microbiol 66:788-93, 2000). Other cells
that can be used for this purpose are yeast cells (Boder and Wittrup Nat.
Biotechnol 15:553-557, 1997), COS cells (Higuchi *et al* J Immunol Meth

202:193-204, 1997) and insect cells (Granzerio *et al* J Immunol Meth
203:131-139, 1997; Ernst *et al* Nucleic Acids Res 26:1718-1723, 1998).

Random combination of DNA from different mutated clones in combination
5 with selection of desired function is a more efficient way to search through
sequence space as compared to sequential selection and combination of
selected clones.

The present invention seeks to provide improved methods for *in vitro*
10 protein evolution. In particular, the invention aims to provide more
efficient recombination and shuffling methods, which will give rise to more
altered molecules and thereby improve the probability of finding molecules
with desirable properties.

15 According to a first aspect of the present invention, there is provided a
method for generating a polynucleotide sequence or population of
sequences from parent single-stranded (ss) polynucleotide sequences
encoding one or more protein motifs, comprising the steps of

- 20 a) providing a first population of single-stranded polynucleotide
molecules and a second population of single-stranded
polynucleotide molecules, the first and second populations
together constituting plus and minus strands of parent
polynucleotide sequences;
- 25 b) carrying out a reaction for digesting the first and second
populations of single-stranded polynucleotide molecules with an
exonuclease to generate corresponding populations of single-
stranded polynucleotide fragments;

5 c) contacting said fragments generated from the plus strands with fragments generated from the minus strands and optionally, adding primer sequences that anneal to the 3' and 5' ends of at least one of the parent polynucleotides under annealing conditions;

d) amplifying the fragments that anneal to each other to generate at least one polynucleotide sequence encoding one or more protein motifs having altered characteristics as compared to the one or more protein motifs encoded by said parent polynucleotides.

10 wherein, in step (b), at least one parameter of the reaction used for digestion of the first population of single-stranded polynucleotide molecules is different from the equivalent parameter(s) used in the reaction for digestion of the second population of single-stranded polynucleotide molecules.

15 Thus, the invention provides a method for generating a variant polynucleotide sequence or population of variants from parent single-stranded polynucleotide sequences.

20 The use of different parameters of the reaction used for digestion of the first and second populations of single-stranded polynucleotide molecules provides the advantage of increased variability in the variant polynucleotides produced by the method of the invention.

25 Preferably, the polynucleotide molecules of step (a) are DNA molecules.

By 'corresponding populations of single-stranded polynucleotide fragments' we mean the population of fragments produced by digestion of the first and

second populations of single-stranded polynucleotide molecules with an exonuclease.

By 'equivalent parameter' we mean the same parameter used in the reaction
5 for digestion of the other population of single-stranded polynucleotide molecules. For example, the exonuclease used for digestion of the first population of single-stranded polynucleotide molecules may differ from the exonuclease used for digestion of the second population of single-stranded polynucleotide molecules.

10

By 'exonuclease' we mean a polypeptide, *e.g.* enzyme or fragment thereof, having exonucleolytic activity. Preferably, the exonucleolytic activity of the polypeptide is greater than the endonucleolytic activity of the polypeptide. More preferably, the polypeptide has exonucleolytic activity
15 but is substantially free of endonucleolytic activity.

Advantageously, the parameter of the digestion reaction which differs is selected from exonuclease type, exonuclease concentration, reaction volume, duration of the digestion reaction, temperature of the reaction
20 mixture, pH of the reaction mixture, length of parent single-stranded polynucleotide sequences, amount of single-stranded polynucleotide molecules and buffer composition of the reaction mixture.

In a preferred embodiment of the method of the first aspect of the invention,
25 the exonuclease used for digestion of the first population of single-stranded polynucleotide molecules is different from the exonuclease used for digestion of the second population of single-stranded polynucleotide molecules. Preferably, the exonuclease used for digestion of the first population of single-stranded polynucleotide molecules is a 3' exonuclease

(*i.e.* which preferentially or exclusively removes nucleotides from 3' terminus of ss polynucleotides) and the exonuclease used for digestion of the second population of single-stranded polynucleotide molecules is a 5' exonuclease (*i.e.* which preferentially or exclusively removes nucleotides from 5' terminus of ss polynucleotides).

In a further embodiment of the method of the first aspect of the invention, the exonuclease concentration used for digestion of the first population of single-stranded polynucleotide molecules is different from the exonuclease concentration used for digestion of the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the reaction volume used for digestion of the first population of single-stranded polynucleotide molecules is different from the reaction volume used for digestion of the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the duration of the digestion reaction used for digestion of the first population of single-stranded polynucleotide molecules is different from the duration of the digestion reaction used for digestion of the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the temperature of the reaction mixture used for digestion of the first population of single-stranded polynucleotide molecules is different from the temperature of the reaction mixture used for digestion of the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the pH of the reaction mixture used for digestion of the first population of single-stranded polynucleotide molecules is different from the pH of the reaction mixture used for digestion of the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the length of the polynucleotides in the first population of single-stranded polynucleotide molecules is different from the length of the polynucleotides in the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the buffer composition of the reaction mixture used for digestion of the first population of single-stranded polynucleotide molecules is different from the buffer composition of the reaction mixture used for digestion of the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the amount of single-stranded polynucleotide molecules in the first population of single-stranded polynucleotide molecules is different from the amount of single-stranded polynucleotide molecules in the second population of single-stranded polynucleotide molecules.

In a further embodiment of the method of the first aspect of the invention, the first population of single-stranded polynucleotide molecules constitutes the plus strands of parent polynucleotide sequences and the second population of single-stranded polynucleotide molecules constitutes the minus strands of parent polynucleotide sequences.

Conveniently, step c) further comprises adding primer sequences that anneal to the 3' and/or 5' ends of at least one of the parent polynucleotides under annealing conditions.

5

Thus, the invention provides a method of combining polynucleotide fragments to generate a polynucleotide sequence or population of sequences of desired characteristics, which method comprises the steps of:

10 a) digesting a linear parent single-stranded polynucleotide encoding one or more protein motifs with a nuclease other than DNase I to generate a population of single-stranded fragments of varying lengths;

 b) assembling a polynucleotide sequence from the sequences derived from step (a).

15

Preferably the method further comprises the step of (c) expressing the resulting protein encoded by the assembled polynucleotide sequence and d) screening the protein for desired characteristics.

20 By controlling the parameters of the exonuclease digestion reaction, the size of the polynucleotide fragments may be controlled. Determining the lengths of the polynucleotide fragments in this way avoids the necessity of having to provide a further step such as purifying the fragments of desired length from a gel.

25

In order to generate a polynucleotide sequence of desired characteristics the parent polynucleotides encoding one or more protein motifs may be subjected to mutagenesis to create a plurality of differently mutated

derivatives thereof. Likewise, a parent polynucleotide may be obtained already encoding a plurality of variant protein motifs of unknown sequence.

Random mutation can be accomplished by any conventional method as
5 described above, but a suitable method is error-prone PCR.

It is preferable to use PCR technology to assemble the single-stranded polynucleotide fragments into a double-stranded (ds) polynucleotide sequence.

10

The polynucleotide sequence is preferably DNA although RNA may be used. For simplicity the term polynucleotide will now be used in the following text in relation to DNA but it will be appreciated that the present invention is applicable to both RNA and DNA.

15

Preferably, any exonuclease that digests polynucleotide from the 5' prime end to the 3' prime, from the 3' to the 5' end or from both the 3' and the 5' ends may be used. Examples of suitable exonucleases which may be used in accordance with the present invention include BAL 31, exonuclease I,
20 exonuclease V, exonuclease VII, exonuclease T7 gene 6, bacteriophage lambda exonuclease and exonuclease Rec J_f.

Using BAL 31 nuclease in the DNA shuffling process of the invention provides a fast, easy and controllable system. This enzyme can give all
25 sizes of gene fragments and the activity of the enzyme can be easily controlled by stopping the digestion at various time points. BAL 31 is predominately a 3' prime exonuclease that removes mononucleotides from both 3' termini of the two strands of a linear DNA. BAL 31 is also an endonuclease; thus the single-stranded DNA generated by the 3' prime

exonuclease activity is degraded by the endonuclease. The 3' prime exonuclease activity of the enzyme works about 20-fold more efficiently than the endonuclease. The enzyme concentrations are therefore important for the obtained DNA fragments. High concentration of enzyme favours
5 blunt-ended DNA whereas at low concentrations the single-stranded DNA termini may be very long. BAL 31 consists of two kinetically distinct forms of the enzyme, a fast (F) and a slow (S) form. The S form is a proteolytic degradation product of the F form. Furthermore, BAL 31 works asynchronously, generating a population of DNA molecules whose termini
10 have been resected to various extents and whose single-stranded tails vary in length. Both forms also act on ssDNA in an exonucleolytic fashion in a highly processive manner. The direction of attack is from the 5' end, in contrast to the mode of digestion of duplex DNA. It has been suggested that the nuclease molecules initially are non-productively bound away from the
15 5'ends and undergo facilitated diffusion to yield productive (terminally bound) enzyme-substrate complexes (Lu T and Gray jr. *HB Biochimica et Biophysica Acta* 1995, vol. 1251, p125-138). The enzyme uses Ca^{2+} as a co-factor which can be bound in complex with EGTA (Ethylene Glycol bis (β -amino ethyl Ether) N,N,N',N'-tetra acetic acid). Linear DNA sequences are
20 digested with BAL31 and the reaction stopped at different time points by the addition of EGTA.

The individual digested fragments are purified, mixed and reassembled with PCR technology. The assembled (reconstituted) gene may then be cloned
25 into an expression vector for expressing the protein. The protein may then be analysed for improved characteristics.

The method of the present invention provides several advantages over known shuffling techniques, including increased rates of recombination, increased variability and control of fragment size.

- 5 The method of the present invention produces a set of progressively shortened DNA fragments for each time point a DNA sample is taken from the BAL31 treatment. The DNA samples may be collected and pooled together or, optionally, individual samples may be chosen and used in the method. Thus the present invention allows a selection of what DNA
10 samples are to be used in the recombination system and thereby offers a further degree of control.

The method of the present invention may be carried out on any polynucleotide which codes for a particular product, for example any
15 protein having binding or catalytic properties *e.g.* antibodies or parts of antibodies, enzymes or receptors. Furthermore, any polynucleotide that has a function that may be altered, such as catalytic RNA, may be shuffled in accordance with the present invention. It is preferable that the parent polynucleotide encoding one or more protein motif is at least 12 nucleotides
20 in length, more preferably at least 20 nucleotides in length, even more preferably more than 50 nucleotides in length. Polynucleotides being at least 100 nucleotides in length or even at least 200 nucleotides in length may be used. Where parent polynucleotides are used that encode large proteins such as enzymes or antibodies, these may be many hundreds or
25 thousands of bases in length. The present invention may be carried out on any size of parent polynucleotide.

The present invention also provides polynucleotide sequences generated by the method described above having desired characteristics. These sequences

may be used for generating gene therapy vectors and replication-defective gene therapy constructs or vaccination vectors for DNA-based vaccinations. In addition, the polynucleotide sequences may be used as research tools.

- 5 The present invention also provides a polynucleotide library of sequences generated by the method described above from which a polynucleotide may be selected which encodes a protein having the desired characteristics. It is preferable that the polynucleotide library is a DNA or cDNA library.
- 10 The present inventions also provides proteins such as enzymes, antibodies, and receptors having characteristics different to that of the wild type produced by the method described above. These proteins may be used individually or within a pharmaceutically acceptable carrier as vaccines or medicaments for therapy, for example, as immunogens, antigens or
- 15 otherwise in obtaining specific antibodies. They may also be used as research tools.

The desired characteristics of a polynucleotide generated by the present invention or a protein encoded by a polynucleotide generated by the present

20 invention may be any variation or alteration in the normal activity of the wild type (parent) polynucleotide or the polypeptide, protein or protein motifs it encodes. For example, it may be desirable to reduce or increase the catalytic activity of an enzyme, or improve or reduce the binding specificity of an antibody. Furthermore, if the protein or polynucleotide is

25 an immunogen, it may be desirable to reduce or increase its ability to obtain specific antibodies against it.

The parent polynucleotide preferably encodes one or more protein motifs. These are defined as regions or elements of polynucleotide sequence that

encode a polypeptide (*i.e.* amino acid) sequence which has, or potentially has, characteristic protein function. For example, a protein motif may define a portion of a whole protein, such as an epitope, a cleavage site or a catalytic site *etc.* However, within the scope of the present invention, an
5 expressed protein motif does not have to display activity, or be "correctly" folded.

Several searchable databases of protein motifs and potential protein motifs are available, such as MOTIF, PROSITE, SMART and BLOCKS
10 (www.blocks.fhcrc.org).

It may be desirable to modify a protein so as to alter the conformation of certain epitopes, thereby improving its antigenicity and/or reducing cross-reactivity. For example, should such a protein be used as an antigen, the
15 modification may reduce any cross-reaction of raised antibodies with similar proteins.

Although the term "enzyme" is used, this is to be interpreted as also including any polypeptide having enzyme-like activity, *i.e.* a catalytic
20 function. For example, polypeptides being part of an enzyme may still possess catalytic function. Furthermore, proteins such as interferons and cytokines are included. Likewise, the term "antibody" should be construed as covering any binding substance having a binding domain with the required specificity. This includes antibody fragments, derivatives,
25 functional equivalents and homologues of antibodies, including synthetic molecules and molecules whose shape mimics that of an antibody enabling it to bind an antigen or epitope. Examples of antibody fragments, capable of binding an antigen or other binding partner are Fab fragment consisting of the VL, VH, Cl and CH1 domains, the Fd fragment consisting of the VH

and CH1 domains; the Fv fragment consisting of the VL and VH domains of a single arm of an antibody; the dAb fragment which consists of a VH domain; isolated CDR regions and F(ab')₂ fragments, a bivalent fragment including two Fab fragments linked by a disulphide bridge at the hinge
5 region. Single chain Fv fragments are also included.

In order to obtain expression of the generated polynucleotide sequence, the sequence may be incorporated in a vector having control sequences operably linked to the polynucleotide sequence to control its expression.

10 The vectors may include other sequences such as promoters or enhancers to drive the expression of the inserted polynucleotide sequence, further polynucleotide sequences so that the protein encoded for by the polynucleotide is produced as a fusion and/or nucleic acid encoding secretion signals so that the protein produced in the host cell is secreted
15 from the cell. The protein encoded for by the polynucleotide sequence can then be obtained by transforming the vectors into host cells in which the vector is functional, culturing the host cells so that the protein is produced and recovering the protein from the host cells or the surrounding medium. Prokaryotic and eukaryotic cells are used for this purpose in the art,
20 including strains of *E. coli*, yeast, and eukaryotic cells such as COS or CHO cells. The choice of host cell can be used to control the properties of the protein expressed in those cells, *e.g.* controlling where the protein is deposited in the host cells or affecting properties such as its glycosylation.

25 The protein encoded by the polynucleotide sequence may be expressed by methods well known in the art. Conveniently, expression may be achieved by growing a host cell in culture, containing such a vector, under appropriate conditions which cause or allow expression of the protein.

Systems for cloning and expression of a protein in a variety of different host cells are well known. Suitable host cells include bacteria, eukaryotic cells such as mammalian and yeast, and baculovirus systems. Also, utilising the retrovirus system for cloning and expression is a good alternative, since this virus can be used together with a number of cell types. Mammalian cell lines available in the art for expression of a heterologous polypeptide include Chinese hamster ovary cells, HeLa cells, baby hamster kidney cells, COS cells and many others. A common, preferred bacterial host is *E. coli*.

Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. Vectors may be plasmids, viral *e.g.* phage, or phagemid, as appropriate. For further details see, for example, Molecular Cloning: a Laboratory Manual: 3rd edition, Sambrook and Russell, 2001, Cold Spring Harbor Laboratory Press. Many known techniques and protocols for manipulation of polynucleotide sequences, for example in preparation of polynucleotide constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins, are described in detail in Current Protocols in Molecular Biology, Ausubel *et al.* eds., John Wiley & Sons, 1992.

The system can be used for the creation of DNA libraries comprising variable sequences which can be screened for the desired protein function in a number of ways. Enzyme function can be screened for with methods specific for the actual enzyme function *e.g.* CMCase activity, β -glucosidase activity and also thermostability. Furthermore, phage display and cell surface display may be used for screening for enzyme function (Cramer *et al.*, Nature 1998 15; 391 (6664):288-291; Zhang J. H. *et al.*, PNAS. USA

1997 94 (9): 4504-4509; Warren M.S. *et al.*, Biochemistry 1996, 9; 35(27): 8855-8862; Kim *et al.*, Appl Environ Microbiol 66:788-93, 2000) as well as for altered binding properties of *e.g.* antibodies (Griffith *et al.*, EMBO J. 113: 3245-3260, 1994).

5

A protein provided by the present invention may be used in screening for molecules which affect or modulate its activity or function. Such molecules may be useful in a therapeutic (possibly including prophylactic) context.

- 10 The present invention also provides vectors comprising polynucleotide sequences generated by the method described above.

The present inventions also provides compositions comprising either polynucleotide sequences, vectors comprising the polynucleotide sequences
15 or proteins generated by the method described above and a pharmaceutically acceptable carrier or a carrier suitable for research purposes.

The present invention further provides a method comprising, following the
20 identification of the polynucleotide or polypeptide having desired characteristics by the method described above, the manufacture of that polypeptide or polynucleotide in whole or in part, optionally in conjunction with additional polypeptides or polynucleotides.

- 25 Thus, a further aspect of the invention provides a method for making a polypeptide having desired properties, the method comprising the following steps:

- (a) generating variant forms of a parent polynucleotide using a method according to the first aspect of the invention;
- (b) expressing the variant polynucleotides produced in step (a) to produce variant polypeptides;
- 5 (c) screening the variant polypeptides for desired properties; and
- (d) selecting a polypeptide having desired properties from the variant polypeptides.

10 The invention further provides a polypeptide obtained by the above method.

Following the identification of a polynucleotide or polypeptide having desired characteristics, these can then be manufactured to provide greater numbers by well-known techniques such as PCR, cloning and expression within a host cell.

15 The resulting polypeptides or polynucleotides may be used in the preparation of industrial enzymes, *e.g.* laundry detergent enzymes where an increased activity is preferred at lower temperatures. Alternatively, the manufactured polynucleotide or polypeptide may be used as a research tool,
20 *i.e.* antibodies may be used in immunoassays, and polynucleotides may be used as hybridization probes or primers. Alternatively, the resulting polypeptides or polynucleotides may be used in the preparation of medicaments for diagnostic use, pharmaceutical use, therapy etc. as discussed as follows.

25 The polypeptides or polynucleotides generated by the methods of the invention and identified as having desirable characteristics can be formulated in pharmaceutical compositions. These compositions may comprise, in addition to one of the above substances, a pharmaceutically

acceptable excipient, carrier, buffer, stabilizer or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of administration,
5 *e.g.* oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal routes.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as
10 gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

15

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to
20 prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilizers, buffers, antioxidants and/or other additives may be included, as required.

25 Thus, the invention further provides a polypeptide produced by the methods of the invention for use in medicine and the use of provides a polypeptide produced by the methods of the invention in the preparation of a medicament for use in the treatment, therapy and/or diagnosis of a disease.

Whether it is a polypeptide, *e.g.* an antibody or fragment thereof, an enzyme, a polynucleotide or nucleic acid molecule, identified following generation by the present invention that is to be given to an individual, administration is preferably in a "prophylactically effective amount" or a
5 "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, *e.g.* decisions on dosage
10 etc, is within the responsibility of general practitioners and other medical doctors, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of the techniques and protocols mentioned above can be found in Remington's
15 Pharmaceutical Sciences, 16th edition, Osol, A. (ed), 1980.

Alternatively, targeting therapies may be used to deliver the active agent more specifically to certain types of cell, by the use of targeting systems such as antibody or cell specific ligands. Targeting may be desirable for a
20 variety of reasons; for example if the agent is unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.

Instead of administering these agents directly, they could be produced in the
25 target cells by expression from an encoding gene introduced into the cells, *e.g.* in a viral vector (a variant of the VDEPT technique *i.e.* the activating agent, *e.g.* an enzyme, is produced in a vector by expression from encoding DNA in a viral vector). The vector could be targeted to the specific cells to

be treated, or it could contain regulatory elements which are switched on more or less selectively by the target cells.

Alternatively, the agent could be administered in a precursor form, for
5 conversion to the active form by an activating agent produced in, or targeted to, the cells to be treated. This type of approach is sometimes known as ADEPT or VDEPT; the former involving targeting the activating agent to the cells by conjugation to a cell-specific antibody, while the latter involves producing the activating agent, *e.g.* an enzyme, in a vector by expression
10 from encoding DNA in a viral vector (see for example, EP-A-415731 and WO 90/07936).

A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the
15 condition to be treated.

As a further alternative, the polynucleotide identified as having desirable characteristics following generation by the method of the present invention could be used in a method of gene therapy, to treat a patient who is unable
20 to synthesize the active polypeptide encoded by the polynucleotide or unable to synthesize it at the normal level, thereby providing the effect provided by the corresponding wild-type protein.

Vectors such as viral vectors have been used in the prior art to introduce
25 polynucleotides into a wide variety of different target cells. Typically the vectors are exposed to the target cells so that transfection can take place in a sufficient proportion of the cells to provide a useful therapeutic or prophylactic effect from the expression of the desired polypeptide. The transfected nucleic acid may be permanently incorporated into the genome

of each of the targeted tumour cells, providing long lasting effect, or alternatively the treatment may have to be repeated periodically.

5 A variety of vectors, both viral vectors and plasmid vectors, are known in the art, see US Patent No. 5,252,479 and WO 93/07282. In particular, a number of viruses have been used as gene transfer vectors, including papovaviruses, such as SV40, vaccinia virus, herpes viruses, including HSV and EBV, and retroviruses. Many gene therapy protocols in the prior art have used disabled murine retroviruses.

10

As an alternative to the use of viral vectors other known methods of introducing nucleic acid into cells includes electroporation, calcium phosphate co-precipitation, mechanical techniques such as microinjection, transfer mediated by liposomes and direct DNA uptake and receptor-mediated DNA transfer.

15

As mentioned above, the aim of gene therapy using nucleic acid encoding a polypeptide, or an active portion thereof, is to increase the amount of the expression product of the nucleic acid in cells in which the level of the wild-type polypeptide is absent or present only at reduced levels. Such treatment may be therapeutic in the treatment of cells which are already cancerous or prophylactic in the treatment of individuals known through screening to have a susceptibility allele and hence a predisposition to, for example, cancer.

20

25 The present invention also provides a kit for generating a polynucleotide sequence or population of sequences of desired characteristics comprising reagents for ssDNA preparation, an exonuclease and components for

carrying out a PCR technique, for example, thermostable DNA (nucleotides) and a stopping device, for example, EGTA.

As outlined above the present invention conveniently provides for the
5 creation of mutated enzyme gene sequences and their random combination to functional enzymes having desirable characteristics. As an example of this aspect of the invention, the enzyme genes are mutated by error prone PCR which results in a mutation rate of approximately 0.7%. The resulting pool of mutated enzyme genes are then digested with an exonuclease,
10 *e.g.* BAL31, and the reaction inhibited by the addition of EGTA or by heat inactivation at different time points, resulting in a set of DNA fragments of different sizes. These may then be subjected to PCR based reassembly as described above. The resulting reassembled DNA fragments are then cloned and a gene library constructed. Clones may then be selected from this
15 library and sequenced.

A further application of this technology is the generation of a population of variable DNA sequences which can be used for further selections and analyses. Besides encoding larger proteins, *e.g.* antibody fragments and
20 enzymes, the DNA may encode peptides where the molecules functional characteristics can be used for the design of different selection systems. Selection of recombined DNA sequences encoding peptides has previously been described (Fisch *et al.*, PNAS. USA 1996 Jul 23; 93 (15): 7761-7766). In addition, the variable DNA population can be used to produce a
25 population of RNA molecules with *e.g.* catalytic activities. Vaish *et al.*, (PNAS. USA 1998 Mar 3; 95 (5): 2158-2162) demonstrated the design of functional systems for the selection of catalytic RNA and Eckstein F (Ciba Found. Symp. 1997; 209; 207-212) has outlined the applications of catalytic RNA by the specific introduction of catalytic RNA in cells. The system may

be used to further search through the sequence space in the selection of functional peptides/molecules with catalytic activities based on recombined DNA sequences.

- 5 Aspects and embodiments of the present invention will now be illustrated, by way of example, with reference to the accompanying figures. Further aspects and embodiments will be apparent to those skilled in the art.

Figure 1 shows the principle of the method from template molecule to
10 improved molecule.

Figure 2 shows the principle steps in preparation of single-stranded DNA using biotin.

- 15 Figure 3 shows the principle steps in the preparation of single-stranded DNA using phage.

Figure 4 shows the principle steps generating single-stranded DNA fragments using exonuclease treatment.
20

Figure 5 shows the principle steps for assembly of single-stranded DNA fragments using PCR.

- Figure 6 shows the % of recombinants formed having one cross-over
25 following digestion of dsDNA with 20 U/ml BAL31 for varying periods of time.

Figure 7 shows the % of recombinants formed having two cross-overs following digestion of dsDNA with 20 U/ml BAL31 for varying periods of time.

- 5 Figure 8 shows the % of recombinants formed having one cross-over following digestion of ssDNA with 1.25 U/ml BAL31 for varying periods of time.

- 10 Figure 9 shows the % of recombinants formed having two cross-overs following digestion of ssDNA with 1.25 U/ml BAL31 for varying periods of time.

- 15 Figure 10 shows the % of recombinants formed having one cross-over following digestion of ssDNA with 11 U/ml BAL31 for varying periods of time.

- 20 Figure 11 shows the % of recombinants formed having two cross-overs following digestion of ssDNA with 11 U/ml BAL31 for varying periods of time.

- Figure 12 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng ssDNA with BAL31 for 50 minutes (lane 1), 30 minutes (lane 2) and 10 minutes (lane 3). Untreated ssDNA is shown in lane 4. Molecular weight markers are shown in lane 5.

- 25 Figure 13 shows the corresponding gel chromatograms for lane 4 in Figure 12.

Figure 14 shows the corresponding gel chromatograms for lane 3 in Figure 12.

Figure 15 shows the corresponding gel chromatograms for lane 2 in Figure 12.

Figure 16 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng DNA with exonuclease VII for 10 minutes (lane 3), 20 minutes (lane 4) and 30 minutes (lane 5). Untreated ssDNA is shown in lane 2. Molecular weight markers are shown in lane 1.

Figure 17 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng DNA with exonuclease Rec J_f (9 U/mg ss/DNA) for 10 minutes (lane 2), 20 minutes (lane 3) and 30 minutes (lane 4). Untreated ssDNA is shown in lane 1. Molecular weight markers are shown in lane 5.

Figure 18 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng DNA with exonuclease Rec J_f (36 U/mg ss/DNA) for 10 minutes (lane 3), 20 minutes (lane 4) and 30 minutes (lane 5). Untreated ssDNA is shown in lane 2. Molecular weight markers are shown in lanes 1 and 6.

Figure 19 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng DNA with DNase I (0.15 U enzyme/mg DNA). Lane samples were as follows:

Lane 1:	Molecular weight markers
Lane 2:	Untreated ssDNA in Mg buffer

29

- 5
- | | |
|---------|--|
| Lane 3: | ssDNA fragmented with DNase I in Mg buffer |
| Lane 4: | Untreated ssDNA in Mn buffer |
| Lane 5: | ssDNA fragmented with DNase I in Mn buffer |
| Lane 6: | (empty) |
| Lane 7: | (empty) |

Figure 20 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng DNA with DNase I. Lane samples were as follows:

10

- | | | |
|----------|--|--|
| Lane 1: | Molecular weight markers | |
| Lane 2: | Untreated dsDNA in Mg buffer | |
| Lane 3: | Untreated dsDNA in Mg buffer | |
| Lane 4: | Untreated ssDNA (forward strand) in Mg buffer | |
| 15 | Lane 5: | Untreated ssDNA (forward strand) in Mg buffer |
| Lane 6: | Untreated ssDNA (reverse strand) in Mg buffer | |
| Lane 7: | Untreated ssDNA (reverse strand) in Mg buffer | |
| Lane 8: | dsDNA fragmented with DNase I (0.24 U enzyme/ μ g DNA) in Mg buffer | |
| 20 | Lane 9: | dsDNA fragmented with DNase I (1.3 U enzyme/ μ g DNA) in Mg buffer |
| Lane 10: | ssDNA (forward strand) fragmented with DNase I (0.24 U enzyme/ μ g DNA) in Mg buffer | |
| Lane 11: | ssDNA (forward strand) fragmented with DNase I (1.3 U enzyme/ μ g DNA) in Mg buffer | |
| 25 | Lane 12: | ssDNA (reverse strand) fragmented with DNase I (0.24 U enzyme/ μ g DNA) in Mg buffer |
| Lane 13: | ssDNA (reverse strand) fragmented with DNase I (1.3 U enzyme/ μ g DNA) in Mg buffer | |

Figure 21 shows the corresponding gel chromatograms for lane 6 in Figure 20.

5 Figure 21 shows the corresponding gel chromatograms for lane 6 in Figure 20.

Figure 22 shows the corresponding gel chromatograms for lane 12 in Figure 20.

10

Figure 23 shows the corresponding gel chromatograms for lane 13 in Figure 20.

15 Figure 24 shows an agarose electrophoresis gel image of fragments generated following digestion of 300 ng DNA with Mung bean nuclease. Lane samples were as follows:

20	Lane 1:	Untreated ssDNA in Mg buffer
	Lane 2:	ssDNA fragmented with Mung bean nuclease for 10 minutes
	Lane 3:	Molecular weight markers

25 Figure 25 shows the effect of duration of fragmentation on frequency of recombination of *tet*-resistance genes following fragmentation of single-stranded DNA with (a) BAL 31, (b) Exo I, (c) T7gene6 and (d) Exo V combined with Exo I.

Figure 26 shows the percentage of multiple cross-overs generated after treatment with different Exonucleases. Frequency of recombination was

evaluated for a) ExoI treated ssDNA, b) ExoI (10 min) treated ssDNA combined with ExoVII treated DNA, c) ExoI (10 min) treated ssDNA combined with ExoV treated ssDNA, and d) ExoI (10 min) treated ssDNA combined with ExoV and ExoVII treated ssDNA.

5

Figure 27 shows a comparison of the number of recombinations observed following fragmentation of ssDNA with an exonuclease and fragmentation of dsDNA with an endonuclease.

10

EXAMPLES

The DNA shuffling procedure can be illustrated by the steps shown in
5 Figures 1 to 5. The gene encoding the protein of interest (X) in the plasmid
pFab5chis is used in this example. Random mutations are introduced by
error prone PCR. Single-stranded DNA is prepared. This can be carried out
by either biotinylated primers or by the use of phage being able to pack
single-stranded DNA, as discussed above. The coding and the non-coding
10 ssDNA strands are prepared in different reactions (A and B). The ssDNA
strands from either reactions are subjected to separate enzymatic treatment
using *e.g.* BAL 31. By mixing the two pools of single-stranded DNA
fragments in equimolar amounts the gene can be reassembled in a shuffled
nature and in many versions by the use of two subsequent PCR reactions,
15 where the first reaction contains no primers. After cloning this library of
reassembled genes in pY, selections can be performed to achieve the
improved molecule of interest.

A more detailed description of examples of the present invention is given
20 below.

Example 1

25 *Reagents*

AmpliTaq® polymerase was purchased from Perkin-Elmer Corp., dNTPs
from Boehringer Mannheim Biochemica (Mannheim, Germany), and
BAL31 Nuclease from New England Biolabs Inc. (Beverly, USA). All
30 restriction enzymes were purchased from New England Biolabs Inc.

(Beverly, USA). Ethidium bromide was purchased from Bio-Rad Laboratories (Bio-Rad Laboratories, Hercules, CA, USA). T4 DNA Ligase was purchased from New England Biolabs Inc. (Beverly, USA). EDTA and EGTA was purchased from Kebo Lab (Sweden).

5

All primers were designed in the laboratory and obtained from Life Technologies (Täby, Sweden) and SGS-DNA (Köping, Sweden).

PCR

10

All Polymerase Chain Reactions (PCR) were carried out in a automatic thermocycler (Perkin-Elmer Cetus 480, Norwalk, CT, USA). PCR techniques for the amplification of nucleic acid are described in US Patent No. 4,683,195. References for the general use of PCR techniques include

15 Mullis *et al.*, Cold Spring Harbor Symp. Quant. Biol., 51:263, (1987), Ehrlich (ed), PCR technology, Stockton Press, NY, 1989, Ehrlich *et al.*, Science, 252:1643-1650, (1991), "PCR protocols; A Guide to Methods and Applications", Eds. Innis *et al.*, Academic Press, New York, (1990).

20 *Sequencing*

All constructs have been sequenced by the use of BigDye Terminator Cycle Sequencing kit (Perkin-Elmer, Elmervill, CA, USA). The sequencing was performed on a ABI Prism 377 DNA Sequencer.

25

Agarose electrophoresis

Agarose electrophoresis of DNA was performed with 2% agarose gels (AGAROSE (FMC Bioproducts, Rockland, ME, USA)) with 0.25µg/ml

ethidium bromide in Tris-acetate buffer (TAE-buffer 0.04M Tris-acetate, 0.001M EDTA). Samples for electrophoresis were mixed with a sterile filtrated loading buffer composed of 25% Ficoll[®] and Bromphenolic blue and loaded into wells in a the 2% agarose gel. The electrophoresis was run at 90 V for 45 minutes unless otherwise stated in Tris-acetate buffer with 0.25 µg/ml ethidium bromide. Bands of appropriate size were gel-purified using the Qiaquick Gel Extraction Kit (Qiagen GmbH, Hilden, Germany) when needed. As molecular weight standard, DNA molecular weight marker 1 kb ladder (Gibco BRL) was used. The DNA-concentration of the gel extracted products were estimated using a spectrophotometer.

Bacterial Strains

The *Escherichia coli*-strain TOP10F⁺ was used as a bacterial host for transformations. Chemically competent cells of this strain were produced basically as described Hanahan, D. 1983. Studies on transformation of *Escherichia coli* with plasmids. J. Mol. Biol. 166: 557-580. Electrocompetent cells of this bacterial strain were produced (Dower, W.J., J. F. Miller, and C.W. Ragsdale. 1988: High efficiency transformation of *E.coli* by high voltage electroporation. Nucleic Acids Res. 16:6127).

Plasmids

All genetic manipulations were performed in pFab5chis according to Molecular cloning; a laboratory manual (Second Edition, Cold Spring Harbor Laboratory Press, 1989). This vector is designed to harbour any scFv gene inserted between SfiI and NotI sites. The SfiI site is located in the pelB leader and the NotI site is located just after the VL region, such

that VH-linker-VL is inserted. In this case, an antibody directed to CD40 was used.

Primers

5

Two biotinylated primers surrounding the antibody gene of pFab5chis were designed with the following sequences including designated unique restriction sites:

10 1736 SfiI forward primer:

5'-ATT ACT CGC GGC CCA GCC GGC CAT GGC CCA CAG GTC
AAG CTC GA

15 and 1735 NotI reversed primer:

5'-TTA GAG CCT GCG GCC GCC TTG TCA TCG TCG TCC TT

Two non-biotinylated primers surrounding the antibody gene of pFab5chis
20 were designed with the following sequences including designated unique restriction sites:

1664 SfiI forward primer:

25 5'-ATT ACT CGC GGC CCA GCC GGC CAT GGC CCA CAG GTC
AAG CTC GA

and 1635 NotI reversed primer:

5'-TTA GAG CCT GCG GCC GCC TTG TCA TCG TCG TCC TT

Standard PCR

- 5 Standard PCR reactions were run at 25 cycles consisting of following profile: denaturation (94°C, 1 minute), primer annealing (55°C, 1 minute) and extension (72°C, 3 minutes). Each PCR reaction contained 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 200 µM dNTP, 1 µM forward primer, 1 µM reverse primer, 1.25 U AmpliTaq® thermostable
- 10 DNA polymerase (Perkin-Elmer Corp.), and 50 ng template in a final volume of 100 µl.

Error Prone PCR

- 15 The error prone PCR reactions were carried out in a 10 x buffer containing 500 mM NaCl, 100 mM Tris-HCl, pH 8.8, 5mM MgCl₂ 100 µg gelatine (according to Kuipers *et al.*, Nucleic Acids Res. 1991, Aug 25;19 (16):4558 but with MgCl₂ concentration increased from 2 mM to 5 mM).
- 20 For each 100 µl reaction the following was mixed:

	dATP 5 mM	5 µl
	dGTP 5 mM	5 µl
	dTTP 10 mM	10 µl
25	dCTP 10 mM	10 µl
	20 µM 3' primer	1.5µl
	20 µM 5'-primer	1.5 µl
	10x Kuipers buffer	10 µl
	sterile mp H ₂ O	46.3 µl

The template in pFab5chis vector was added at an amount of 50 ng. 10 µl of 10 mM MnCl₂ was added and the tube was checked that no precipitation of MnO₂ occurred. At last 5 Units of Taq enzyme was added. The error prone PCR was run at the following temperatures for 25 cycles without a hot start: 94°C 1', 45 °C 1', 72 °C 1' , + 72 °C for 7 minutes. The resulting product was an error pruned insert over the protein of approximately 750 bp. This insert was purified with Gibco PCR purification kit, before further treatment.

10

Generation of single-stranded DNA by biotinylated primers

The fragment of interest was amplified by two separate PCR reactions. These reactions can be standard PCR as described above or error prone PCR also as described above. The primers should be designed so that in one reaction the forward primer is biotinylated and in the other reaction the reverse primer is biotinylated. For example, PCR reactions with A) primers 1736 and 1635 and B) primers 1664 and 1735, with the above mentioned profile was performed for 25 cycles with pFab5chis-antibody as template. This yielded PCR-products of approximately 750 bp where in A the upper strand was biotinylated and in B the lower strand was biotinylated.

The non-biotinylated strands were retrieved by purification using a solid matrix coated with streptavidin e.g. Dynabeads[®]. The magnetic beads are washed and equilibrated with PBS/1% BSA and B&W buffer containing 5 mM Tris pH 7.5, 1 M NaCl, and 0.5 mM EGTA. 100 µl of each PCR product is mixed with 100 µl beads dissolved in 2 x B&W buffer and incubated at room temperature for 15 minutes with rotation. Unbound PCR products are removed by careful washing twice with B&W. The non-

biotinylated strand of the captured DNA is eluted by alkaline denaturation by letting the DNA incubate with 25 μ l 0.1 M NaOH for 10 minutes in room temperature. The solution is separated from the beads and neutralised with 7.5 μ l 0.33 M HCl and 2.5 μ l 1 M Tris pH 8.

5

Generation of single-stranded DNA using phage

The fragment of interest was cloned into bacteriophage M13 vectors M13mp18 and M13mp19 using PstI/HindIII restriction enzymes. The
10 bacteriophage were propagated using *Escherichia coli*-strain TOP10F' according to conventional methods. Single-stranded DNA for the upper strand was prepared from bacteriophage vector M13mp18 and single-stranded DNA for the lower strand was prepared from bacteriophage vector M13mp19. Briefly, 1.5 ml of an infected bacterial culture was centrifuged at
15 12 000g for 5 minutes at 4°C. The supernatant was precipitated with 200 μ l 20% PEG8000/2.5 M NaCl. The pelleted bacteriophage was resuspended in 100 μ l TE. 50 μ l phenol equilibrated with Tris-Cl (pH 8.0) was added and the sample was vortexed. After centrifugation at 12 000g for 1 minute at RT the upper phase, containing the DNA, was transferred and precipitated with
20 ethanol. The DNA pellet was dissolved in 50 μ l TE (pH 8.0) and stored at –20°C. (Sambrook *et al.* Molecular Cloning, A laboratory manual 2nd edition. Cold Spring Harbor Laboratory Press. 1989, chapter 4). Single-stranded DNA prepared from phage is circular and must be opened prior to BAL31 treatment. This can be performed with an endonuclease able to cleave
25 single-stranded DNA.

Generation of single-stranded DNA using asymmetric PCR

PCR products are purified using a spin column to remove excess primers from the previous PCR. 150 ng of the purified product is used as template in
5 a linear amplification carried out in 100 µl of 1xGeneAmp® 10 x PCR buffer containing 1.5 mM MgCl₂ (Applied Biosystems), 200 µM of each dNTP (New England BioLabs), 1,25 U AmpliTaq® DNA Polymerase (Applied Biosystems) and 1.0 µM of a single primer. PCR cycle conditions are: denaturation at 94°C for 1 minute, 35 cycles of 94°C for 30 seconds,
10 55°C for 30 seconds, 72°C for 1 minute followed by extension at 72°C for 7 minutes.

Asymmetric PCR products are size separated from double stranded template on a 1 % agarose gel and purified using Qiaquick Gel Extraction Kit
15 (Qiagen).

Generation of single-stranded fragmented DNA using BAL 31

The ssDNA strands (containing upper and lower strands, respectively) were
20 subjected to separate enzymatic treatment using *e.g.* BAL 31. Each digestion reaction contained 0.02 µg/µl ssDNA, 600 mM NaCl, 20 mM Tris-HCl, 12 mM CaCl₂, 12 mM MgCl₂, 1 mM EDTA pH 8.0 and BAL 31 at various enzyme concentrations ranging from 0.1 - 5 U/ml. The reactions were incubated at 30°C and fractions of digested ssDNA were collected
25 sequentially at 10, 30, 60 and 120 seconds or longer. The reactions were stopped by addition of EDTA and heat treatment at 65°C for 10 minutes. The ssDNA fragments were purified by phenol/chloroform extraction and ethanol precipitated. The ssDNA are resuspended in 10 mM Tris pH 8.0.

WO 03/097834

PCT/GB03/02102

40

The digestion pattern was evaluated by 1 % agarose gel electrophoresis.

Purification of digestion produced fragments:

- 5 Digested DNA fragments were purified by phenol/chloroform/
isoamylalcohol extraction. 50 µl of buffered phenol was added to each tube
of 100 µl sample together with 50 µl of a mixture of chloroform and
isoamylalcohol (24:1). The tubes were vortexed for 30 seconds and then
centrifuged for 1 minute in a microfuge at 14000 r.p.m. The upper phase
10 was then collected and mixed with 2.5 volumes of 99.5% Ethanol (1/10 was
3M Sodium Acetate, pH 5.2). The DNA was precipitated for 1 hour in -80
°C. The DNA was then pelleted by centrifugation for 30 minutes in a
microfuge at 14,000 r.p.m. The pellet was washed once with 70% ethanol
and then re-dissolved in 10 µl of sterile water.

15

Analysis of digestion produced purified fragments on agarose gel

- 5 µl of the dissolved pellet from each time point and from the blank were
mixed with 2.5 µl of loading buffer (25% Ficoll[®] and Bromphenolic blue)
20 and loaded into wells in a 2% agarose gel. The electrophoresis of the
different time points were performed as above.

Reassembly of full length fragments

- 25 Reassembly of the ssDNA fragments is achieved by two sequential PCR
reactions. The first PCR reaction should contain 10 mM Tris-HCl, pH 8.3,
50 mM KCl, 1.5 mM MgCl₂, 200 µM dNTP, 0.3 U Taq polymerase and 2
µl BAL31 treated sample, all in a final volume of 25 µl, and subjected to 5
cycles with the following profile: 94 °C for 1 minute, 50 °C for 1 minute

and 72 °C for 2 minutes + 72 °C for 5 minutes. The second PCR reaction should contain 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 200 µM dNTP, 0.6 U Taq polymerase, 1 µM forward primer, 1 µM reverse primer, and 5 µl sample from the first PCR reaction, all in a final volume of 50 µl, and subjected to 15 cycles with the following profile: 94 °C for 1 minute, 55 °C for 1 minute and 72 °C for 2 minutes + 72 °C for 7 minutes. The resulting products can be evaluated by agarose gel electrophoresis.

Restriction digestion of reassembled fragment and plasmid with SfiI and NotI

The reassembled fragment and the plasmid pFab5chis were first cleaved with SfiI by using NEB buffer 2 including BSA and 11 U enzyme/µg DNA. The reaction was carried out for 4 h at 50°C. After this the DNA was cleaved with NotI by adding conversion buffer and 6 U enzyme/µg DNA. This reaction was carried out for 37°C overnight.

Gel purification of restriction digested vector and restriction digested reassembled fragment

The cleavage reactions were analysed on a 1% agarose gel. The restriction digested insert showed a cleavage product of about 750 bp. This corresponds well with the expected size. The band of the cleaved insert and plasmid was cut out and gel-extracted as previously described.

Ligation of reassembled restriction digested fragment with restriction digested pFab5chis

Purified cleaved pFab5chis was ligated with purified reassembled restriction

digested fragment at 12°C water bath for 16 hours. 50 µl of the vector was mixed with 50 µl of the insert and 15 µl of 10x buffer (supplied with the enzyme), 7.5µl ligase (5 U/µl) and sterile water to a final volume of 150µl. A ligation of restriction digested pFab5chis without any insert was also
5 performed in the same manner.

Transformation of chemically competent E coli TOP10F' with the ligated reassembled insert and pFab5chis

10 The ligation reactions were purified by phenol/chloroform extraction as described above. The upper phase from the extraction was collected and mixed with 2.5 volumes of 99.5% Ethanol (1/10 was 3M Sodium Acetate, pH 5.2). The DNA was precipitated for 1 hour in -80 °C. The DNA was then pelleted by centrifugation for 30 minutes in a microfuge at 14.000
15 r.p.m. The pellet was washed once with 70% ethanol and then re-dissolved in 10 µl of sterile water. 5 µl of each ligation was separately mixed with 95 µl chemically competent *E coli* TOP10F' incubated on ice for 1 hour and then transformed (Sambrook *et al.* Molecular Cloning, A laboratory manual 2nd edition. Cold Spring Harbor Laboratory Press, 1989). After one hour's
20 growth the bacteria from the two transformations were spread onto ampicillin containing agar plates (100 µg/ml). The plates were grown upside-down in a 37°C incubator for 14 hours.

Example 2 – Recombination frequencies; comparison of dsDNA and ssDNA

In further comparable experiments, three scFv antibody fragments were
5 used in a recombination experiments, either as dsDNA or as ssDNA.

dsDNA

The three scFv genes were each amplified in PCR using forward and
10 reverse primers and standard PCR procedure. The size of the bands was confirmed with agarose electrophoresis and the rest of the amplified PCR products were purified using Concert PCR purification kit (Gibco). The dsDNA from the three scFv were mixed in equimolar amounts and treated with BAL31. Each digestion reaction contained dsDNA at a concentration
15 of 0.02 µg/µl reaction volume, 600 mM NaCl, 20 mM Tris-HCl, 12 mM CaCl₂, 12 mM MgCl₂, 1 mM EDTA pH 8.0 and BAL31 at various enzyme concentrations (using 4, 20 or 100 U enzyme/ml reaction volume). The reactions were incubated at 30°C and fractions of digested dsDNA were collected sequentially at 10, 30, and 50 minutes. The reactions were
20 stopped with EDTA and heat treatment (alternatively, an EDTA-free heat inactivation protocol may be used; see below) and purified using phenol/chloroform extraction and ethanol precipitation. The dsDNA samples were resuspended in 10 mM Tris pH 8.0.

25 Keeping each time point separate, the samples were subjected to reassembly PCR (for this reassembly, 60 ng DNA is used) and amplification PCR according to the protocol, and cloned in pGEM[®] (Product No A362A, Promega, Madison, USA). Eighteen clones from each time point were

sequenced and the number and frequency of recombinations were determined.

Heat inactivation of exonuclease digestions

5

A protocol to stop the BAL31 reaction without using EDTA has been established. This heat inactivation protocol avoids using phenol/chloroform extraction, which is hazardous to health and also causes loss of material.

- 10 In brief, the sample is incubated for 10 minutes at 95 C and then put directly in ice, to stop the enzymatic reaction. After this the sample can be directly precipitated using ethanol.

ssDNA

15

- The three scFv genes were each amplified in two PCR reactions using primer pairs forward/reverse-biotin and forward-biotin/reverse using standard PCR procedure. The size of the bands were confirmed with agarose electrophoresis and the rest of the amplified PCR products were
- 20 purified using Concert PCR purification kit (Gibco). Single-stranded DNA was obtained using magnetic beads according to the protocol, achieving three sense strands and three antisense strands. The sense strands and the antisense strands, respectively, from the three scFv were mixed in equimolar amounts and treated with BAL31 according to the protocol
- 25 (using 1.25 or 11 U enzyme/ml reaction volume and ssDNA at a concentration of 0.015 µg/µl reaction volume) and samples were taken out at 0 (*i.e.* undigested), 10, 30 and 50 minutes. The reactions were stopped with EDTA and heat treatment and purified using phenol/chloroform extraction and ethanol precipitation. Keeping each time point separate, but

mixing sense and antisense strands, the samples were subjected to reassembly PCR (for this reassembly 60 ng DNA is used) and amplification PCR according to the protocol, and cloned in pGEM. Eighteen clones from each time point were sequenced and the number and frequency of recombination 5 were determined.

Results

The highest frequency of recombination using dsDNA was achieved using 10 20 U enzyme/ml reaction volume (containing 0.02 µg/µl DNA) and treating for 10 minutes. This gave 39% of the clones with one cross-over (Figure 6) and 17% of the clones with two cross-overs (Figure 7). Using 4 U enzyme/ml gave no cross-overs independent of time for fragmentation and 100 U enzyme/ml resulted in complete fragmentation into very small 15 fragments, as indicated by the failure to regain the full-length gene during reassembly.

The results from the experiments using ssDNA are shown in Figures 8 to 10. Figure 8 shows 1.25 U/ml BAL31 and clones with one cross-over, 20 Figure 9 shows 1.25 U/ml BAL31 and clones with two cross-overs. Figure 10 shows 11 U/ml BAL31 and clones with one cross-over, and Figure 11 shows 11 U/ml BAL31 and clones with two cross-overs.

The highest frequency of recombination giving one cross-over using ssDNA 25 was achieved using 11 U enzyme/ml and treating for 10 minutes (Figure 10). 59% of the clones had one cross over. The highest frequency of recombination giving two cross-overs using ssDNA was achieved using 1.25 U enzyme/ml and treating for 30 minutes (Figure 9). 20% of the clones had two cross overs.

Conclusions and comments

These data clearly show that a higher frequency of recombination is achieved using ssDNA. The three scFv used have the same framework sequences, indicating that the number of cross overs reported may be higher due to cross overs in regions where no sequence difference will result. These experiments using ssDNA were carried out in a non-optimal fashion for showing maximum recombination, since all strands from all three molecules were mixed. Mixing the sense strand from one scFv with the antisense strand from another scFv would produce higher frequencies of cross overs, see Example 3 below. Also, each time point was here kept separate and it would be logical to estimate the frequency of cross overs to increase if different time points, i.e. different fragments sizes, are mixed.

15

Example 3 - Recombination frequencies; homology dependence using ssDNA

To investigate the homology required to achieve cross-over we set up experiments to recombine four scFv (designated SMUC159, CT17, AE11 and MO152) making up three pairs with different homologies, as follows:

	SMUC159 – CT17	92%
25	SMUC159 – AE11	70%
	SMUC159 – MO152	60%

The four scFv genes were each amplified in two PCR reactions using primer pairs forward/reverse-biotin and forward-biotin/reverse using standard PCR

procedure. The size of the bands were confirmed with agarose electrophoresis and the rest of the amplified PCR products were purified using Concert PCR purification kit (Gibco). Single-stranded DNA was obtained using magnetic beads according to the protocol, achieving four
5 sense strands and four antisense strands. Each strand was treated with BAL31 according to the protocol (using 4.2 or 12.5 U enzyme/ml) and samples were taken out at 0, 10, 30 and 50 minutes, or 0, 15, 30, 45 and 60 minutes. The reactions were stopped with EDTA and heat treatment and purified using phenol/chloroform extraction and ethanol precipitation.
10 Keeping each time point separate, but mixing sense and antisense strands forming the pairs as indicates above, the samples were subjected to reassembly PCR and amplification PCR according to the protocol, and cloned in pGEM. Fifteen clones from each time point were sequenced and the number and frequency of recombination were determined.

15

Results

Cross overs were identified in all combinations of scFv, indicating that as low as 60% homology is enough to achieve recombination.

20

Example 4 - Improved control of fragment size using exonucleases

(A) Exonucleases

25

We use exonucleases, *e.g.* BAL31, exonuclease I, exonuclease V, exonuclease VII, exonuclease T7 gene 6, bacteriophage lambda exonuclease, and exonuclease Rec J_f for fragmentation in the methods of the present invention. These enzymes cleave off one nucleotide at a time, either

from the 5' end or from the 3' end or from both ends. The reaction can be stopped using EDTA or heat inactivation (see above), depending on the enzyme used. This means that fragments of all possible sizes, differing with only one nucleotide, can be obtained.

5

The following examples demonstrate how exonuclease digestion allows the creation of fragments of various and controllable sizes depending on the conditions used.

10 *BAL 31*

Single-stranded DNA was digested with BAL31 according to the protocol in Example 1, with an enzyme concentration of 4.2 U/ml reaction volume and ssDNA concentration of 0.008 µg/µl reaction volume.

15

In a typical experiment, about 300 ng DNA is isolated at each time point of BAL31 treatment. Figure 12 shows an agarose electrophoresis gel image of such an experiment with untreated ssDNA in lane 4 and ssDNA treated for 10 minutes in lane 3, for 30 minutes in lane 2 and for 50 minutes in lane 1.

20 Lane 5 is the molecular weight (MW) standard.

Figures 13 to 15 shows the corresponding gel chromatograms of the lanes, respectively. Figure 13 is the untreated material and the multiple peaks refer to different conformations of the ssDNA. Figure 14 corresponds to lane 3 and material treated for 10 minutes. The material was heat treated to stop the enzymatic reaction, and thus resolving the different conformations, and one peak of a distinct size is shown. Figure 15 corresponds to lane 2 and material treated for 30 minutes.

25

Here it is clear that the peak corresponding to larger fragments is decreasing and a peak of smaller DNA fragments has appeared.

Exonuclease VII

5

Single-stranded DNA was digested with exonuclease VII using an enzyme concentration of 7.7 U/ml reaction volume and ssDNA concentration of 0.008 µg/µl reaction volume. The reaction buffer comprised 67 mM potassium phosphate (pH 7.9), 10 mM mercaptoethanol, 6.7 mM MgCl₂ and 10 8.3 mM EDTA.

The reaction was allowed to proceed at 37°C for 10, 20 and 30 minutes, before being stopped by heat inactivation (95°C for 5 minutes).

15 In Figure 16 the fragmentation pattern using exonuclease VII is shown. Lane 1 is MW standard, lane 2 is untreated ssDNA, lane 3 is ssDNA fragmented with exonuclease VII for 10 minutes, lane 4 is ssDNA fragmented with exonuclease VII for 20 minutes, and lane 5 is ssDNA fragmented with exonuclease VII for 30 minutes. This shows that the 20 fragment sizes are decreased by time.

Exonuclease Rec J_f

Single-stranded DNA was digested with exonuclease Rec J_f using an 25 enzyme concentration of either 2.5 U/ml reaction volume or 10 U/ml reaction volume and ssDNA at a concentration of 0.007 µg/µl reaction volume, corresponding to 0.36 U enzyme/µg DNA and 1.4 U enzyme/µg DNA, respectively. The reaction buffer comprised 50 mM NaCl, 10 mM Tris.HCl, 10 mM MgCl₂ and 1 mM dithiothreitol, at pH 7.9

The reaction was allowed to proceed at 37°C for 10, 20 and 30 minutes, before being stopped by heat inactivation (95°C for 5 minutes).

5 In Figure 17 the fragmentation pattern using exonuclease Rec J_f at 036 U/microgram ssDNA is shown. Lane 1 untreated ssDNA, lane 2 is ssDNA fragmented with exonuclease Rec J_f for 10 minutes, lane 3 is ssDNA fragmented with exonuclease Rec J_f for 20 minutes, and lane 4 is ssDNA fragmented with exonuclease Rec J_f for 30 minutes. This shows that
10 the fragment sizes are decreased by time. In Figure 18 the enzyme concentration is increased 4 times (1.4 U/microgram ssDNA) and the fragmentation pattern is shown from 0 to 30 minutes, showing a higher degree of fragmentation as compared to Figure 17. This shows that both time and enzyme concentration can be used to control the fragmentation.

15 (B) Endonucleases

Conventional DNA shuffling methods typically use DNase I for fragmentation (for example, see Stemmer, 1994, *Nature* 370:389-391). DNase I cleaves DNA in an endonucleolytic fashion at sites adjacent to
20 pyrimidines. Consequently, not all possible fragment sizes can be obtained.

Moreover, using magnesium in the reaction buffer, a homologous mix of mono- and oligomers is obtained. Hence, different methods such as gel agarose electrophoresis purification or gel filtration need to be used in order
25 to isolate fragments of different sizes. Often fragments of small size or a mix of small and larger fragments are desired to optimise recombination. However, these purification methods introduce single-stranded nicks in the double-stranded PCR products. Fragments of a particular size purified on a gel would thus consist of dsDNA with a large number of single-stranded

nicks, which would give rise to many smaller fragments upon denaturation. This means that many of the single-stranded fragments generated upon denaturation would be too short to function as primers during the annealing, resulting in a great loss of product.

5

Using manganese in the reaction buffer creates fragments of sizes smaller than 50 bp and no gel purification is needed. However, here you are restricted to use only small fragments and these can not be mixed with larger fragments, something that would probably increase the recombination frequency.

10

The problems associated with the use of endonucleases are demonstrated in the following experiments:

15 *DNase I*

DNA was digested for 5 minutes with DNase I at a concentration of 0.15 U/ μ g DNA.

20 Magnesium and manganese buffers were compared when fragmenting with DNase I and the result is shown in Figure 19. Lane 1 is MW standard, lane 2 is untreated ssDNA in Mg buffer, lane 3 is ssDNA fragmented with DNase I in Mg buffer according to Stemmer (1994) *Nature* **370**:389-391, lane 4 is untreated ssDNA in Mn buffer and lane 5 is ssDNA fragmented with DNase I in Mn buffer according to Kikuchi *et al.* (2000) *Gene* **243**:133-137. It is clear from Figure 19 that, when using Mg buffer and conditions according to the Stemmer and Kikuchi papers, no fragmentation occurs. Moreover, when using Mn buffer and conditions according to the

25

Stemmer and Kikuchi papers, all material is totally fragmented within only a few minutes.

In an attempt to obtain fragments of different sizes we decided to use Mg
5 buffer and increase the enzyme concentration. Figure 20 shows an agarose electrophoresis gel image of such an experiment using DNase I. Lane 1 is the MW standard. Lane 6 is untreated ssDNA. Lane 12 is ssDNA treated according to the Stemmer and Kikuchi papers, using 0.15 U enzyme/microgram DNA and lane 13 is the same material treated with 1 U
10 enzyme/microgram DNA (i.e. six times more enzyme).

Figures 21 to 23 shows the corresponding chromatograms. The untreated ssDNA has been heat-treated, therefore only one peak appears in Figure 21 (indicated by arrow). In figure 22, it is apparent that using the amount of
15 DNase I according to the Stemmer and Kikuchi papers the peak for untreated ssDNA is somewhat decreased (indicated by arrow) but no distinct peak is visible for the fragmented DNA, only a smear. Using 6 times more enzyme the untreated ssDNA is totally abolished (Figure 23) and neither here is any visible peak of the fragments.

20

Mung bean nuclease

Single-stranded DNA was digested with Mung bean nuclease (Product No MO250S, New England Biolabs) using an enzyme concentration of either
25 0.375 U/ml reaction volume and ssDNA at a concentration of 0.007 µg/µl reaction volume. The reaction buffer comprised 50 mM sodium acetate, 30 mM NaCl, 1 mM ZnSO₄, at pH 5.0.

The reaction was allowed to proceed at 25°C for 10 minutes, before being stopped by heat inactivation (95°C for 5 minutes).

Figure 24 shows fragmentation using another endonuclease, Mung bean nuclease. Lane 1 is the untreated ssDNA, lane 2 is the same material treated for 10 minutes. Lane 3 is the MW standard.

Results indicate that all DNA was totally fragmented after only 10 minutes digestion with Mung bean nuclease (see lane 2), despite using the enzyme at a concentration lower than that recommended by the manufacturer.

Conclusions and comments

The above examples show how the fragment sizes can be controlled using exonucleases and altering the reaction conditions, *i.e.* time, reaction volume, enzyme concentration. The different peaks are visualised using gel image chromatograms.

In contrast, using endonucleases, such as DNase I, gives a reaction which is hard to control. Using conditions as referred in the literature, either using Mg or Mn containing buffers, typically gives a situation when either everything or nothing is fragmented, see especially Figure 20. An experiment using another endonuclease (Mung bean nuclease) confirms these observations.

Example 5 – Digestion of sub-populations of single-stranded DNA starting material with different exonucleases

In further experiments, the single-stranded DNA starting material was split
5 into two populations, which were then digested using different exonucleases.

Materials and methods

10 *Plasmids*

A tetracycline-deleted variant of plasmid pBR322 was constructed by cleavage with Sall and BamHI (Roche, Basel, Switzerland) Klenow treatment (Amersham Biosciences AB, Uppsala, Sweden) and blunt-end ligation (New England Biolabs, MA, USA). The resulting plasmid was
15 checked for tetracycline sensitivity and is called pBR322dtet.

pBR322stop1 and pBR322stop3 were created by PCR amplification of the tetracycline gene of pBR322 using specific primers (Table 1). Each mutated tetracycline gene was cloned into pBR322.

20

Table 1
Primer sequences

pBR322 NheI forward stop:

25 5'-CACTATGGCGTGCTGCTAGCGCTATATGCGTTGATGCAATTT
CT ATGAGCACCCGTTCT-3'

pBR322 EagI reversed:

5'- CGTAGCCCAGCGCGTCGGCCGCCATGCCGGCGATAATG-3'

pBR322 HindIII forward:

5'- CAGCTTATCATCGATAAGCTTTAATGCGGTAGTTTAT-3'

5 pBR322 Sall reversed stop:

5'-TCTCAAGGGGCATCGGTCGACGCTCTCCCTTATGCGACTCCTGC
ATTAGGAATCAGCCCAGTAGTA-3'

PCR

10

Unless otherwise noted PCR reactions contained 4 μ M of each primer, 160 μ M dNTP (Roche, Basel, Switzerland), 1x AmpliTaq reaction buffer, 2.5 U AmpliTaq thermostable DNA polymerase (Applied Biosystems, CA, USA).

15 FIND PCR 1: 5 or 25 cycles of 94° C 30 s, 50° C 45 s, 72° C 1 minute and then 72° C for 7 minutes, no external primers were included.

FIND PCR 2: 15, 25 or 50 cycles of 94° C 30 s, 55° C 45s, 72° C 1 minute and then 72° C for 7 minutes with external primers included.

20

Single-stranded DNA preparation

The gene of interest, *i.e.* *tet-r*, was amplified using specific primers, one of the primer was biotinylated. SsDNA from sense and antisense strains was
25 purified using streptavidin-magnetic beads (purchased from either Dynal AS, Oslo, Norway or Miltenyi Biotec, Bergisch Gladbach, Germany) according to manufacturer's recommendations. The ssDNA hereby obtained was further purified either by ethanol precipitation or by using recochip (TaKaRa, Shiga, Japan) according to manufacturers recommendations.

FIND experiments

The FIND experiments were initiated by digesting DNA with an exonuclease. The DNA was single-stranded (prepared as above) and originated from the tetracycline resistance gene (pBR322stop1 or pBR322stop3, 945 bp). The exonucleases were BAL31 (0.08-1 U/ μ g DNA, New England Biolabs, MA, USA), exonuclease I (100 U/ μ g DNA, New England Biolabs, MA, USA), T7 gene 6 exonuclease (320 U/ μ g DNA, USB, Cleveland Ohio, USA) and exonuclease V (12.5 U/ μ g DNA, USB, Cleveland Ohio, USA). The time for digestion was in the range 2-90 minutes. The digestion reactions were stopped by adding EDTA to a final concentration of 20 mM and/or heat inactivation at 65 or 95° C for 10 minutes. When EDTA was used to stop the DNA fragmentation the DNA was further purified by phenol/chloroform extraction and ethanol precipitation. The fragments were recombined in a FIND PCR1 reaction for 5 or 25 cycles and the material was amplified in a FIND PCR 2 reaction for 15, 25 or 50 cycles. Finally the full length genes were cloned into pBR322dtet by the use of HindIII and EagI (New England Biolabs, MA, USA) for functionality evaluation or into pGEM (Promega, Madison, WI, USA) for sequencing.

Evaluation of functionality of tetracycline clones

The clones introduced into pBR322dtet were transformed into chemical competent TG1 *E. coli* and plated on LB agar plates containing 1 μ g/ml ampicillin. One to two hundred clones were then moved to LB agar plates containing 50 μ g/ml tetracycline and the frequency of tetracycline resistant clones could be calculated.

Results

As shown above, higher frequencies of recombination using the FIND
5 procedure of the present invention could be achieved using ssDNA in the
fragmentation. The exonuclease BAL 31 is predominately a 3' exonuclease
that removes mononucleotides from both 3' termini of the two strands of a
linear double stranded DNA. However BAL 31 can also degrade the single-
10 stranded DNA ends generated by the 3' exonuclease activity on the double
stranded DNA. The activity of BAL 31 on ssDNA is removal of
mononucleotides from the 5' termini only. Using BAL 31 for
fragmentation of ssDNA and then reassembly to full length genes will
theoretically result in one cross-over per gene. Experiments were therefore
performed to test different exonucleases for the fragmentation of ssDNA.
15 Exonuclease I has 3' activity only whereas BAL 31, T7gene6 and RecJ
exonucleases have 5' activity only. Exonuclease V and Exonuclease VII
have activity from both ends (5' and 3'). In order to show that these
exonucleases can be used in the fragmentation step in a FIND experiment
and yield functionally recombined genes a model system based on
20 tetracycline resistance genes was used.

It was found that BAL 31 (Figure 25a) as well as Exonuclease I (Figure
25b) and T7 gene 6 exonuclease (Figure 25c) all worked well in the FIND
procedure and a dependency on fragmentation time of recombination
25 frequency was observed.

If only one exonuclease, which digests ssDNA from only one end, is used
only one cross-over can in theory be achieved. However, it was found that
further cross-overs can be the obtained if DNA fragments from treatment by

different exonucleases were combined. Exonuclease V and Exonuclease VII treatment will result in small fragments without the 5' and 3' ends. These ends are necessary in order to amplify the recombined material in the last PCR reaction. These DNA fragments can therefore be combined with DNA
5 digested from 5' or 3' ends. The result from such a combination can be seen in Figure 25d where ssDNA treated with Exonuclease I for 10 minutes was combined with ssDNA treated with Exonuclease V for 40 and 50 minutes. Functional clones of up to 40 % were obtained, a result that should be compared to the maximum 25 % achieved in the same system using only
10 one enzyme (Figure 25a-c). Fragments have also combined from Exonuclease I and Exonuclease VII treatment, and fragments from T7 gene 6 exonuclease and Exonuclease VII treatment, at different time points, and functionally recombined clones could be obtained (data not shown).

15

Example 6 – Digestion of sub-populations of single-stranded DNA starting material with different exonucleases (further experiments)

In a separate set of experiments, multiple cross-overs were produced by
20 combining fragments obtained by digesting single-stranded DNA with Exonuclease I with fragment obtained by digesting single-stranded DNA Exonuclease V and/or Exonuclease VII.

Exonuclease I (Exo I) has only 3' activity whereas Exonuclease V (ExoV)
25 and Exonuclease VII (ExoVII) have activity from both ends (5' and 3'). Hence, ExoV and ExoVII treatment will result in small fragments without the 5' and 3' ends. These ends are necessary in order to amplify the recombined material in the last PCR reaction. These DNA fragments can therefore be combined with DNA digested from 5' or 3' ends only.

An equimolar mixture of the three scFv genes was used in fragmentation with ExoI, ExoV and ExoVII. Different combination of fragmented material was mixed and PCR reassembled into full length genes and finally
5 sequenced (Figure 26a-d). ExoI behaved similar to BAL 31 with up to 40% clones with one recombination (Figure 26a). The combination of ExoI with ExoV or ExoVII increased number of crossover up to five recombinations (Figure 26 a and c, respectively). ExoVII seems to be more efficient than ExoV, the ExoI/ExoVII combination generated recombinations in over 75%
10 of the clones. Finally, a combination of fragmented DNA from all three enzymes was mixed which generated recombinations (1 to 4) in almost all clones, only 7% of them were wild type genes. Furthermore, longer fragmentation times were most efficient (Figure 26d). In this particular case the scFv genes consist of large areas of complete homology (the frame work
15 regions), meaning that the number of recombinations identified (up to 4) are the ones that are possible to identify, *i.e.* the actual number of recombinations might be much higher.

Example 7 – Comparison of the effects of digestion of ss DNA with an exonuclease with the effects of digestion of ds DNA with an endonuclease

- 5 The experiments in Figure 27 were carried out by recombining genes coding for three different scFv's using either ssDNA and exonuclease treatment or dsDNA and endonuclease treatment.

Endonuclease digestion of dsDNA was performed using DNase I, as
10 described in Stemmer *et al.*, 1994, *Nature* **370**:389-91. Exonuclease digestion of ssDNA was performed using ExoI, ExoV and ExoVII, with equimolar mixes of fragments from ExoI (10 minutes reaction time), ExoV (30 minutes reaction time) and ExoVII (30 minutes reaction time) being used in recombination reactions.

15

The number of recombinations were evaluated by the use of sequencing. The result shows that ssDNA/exonuclease digestion yields fewer wild type clones and more recombinations.

SEQUENCE LISTING

<110> Alligator Bioscience AB
 <120> A Method for In Vitro Molecular Evolution of Protein Function

<130> ALLBA/P28382PC

<140> PCT/GB03/02102
 <141> 16 May 2003

<150> GB 0211369.4
 <151> 17 May 2002

<150> US 10/321195
 <151> 17 December 2002

<160> 8

<170> SeqWin99

<210> 1
 <211> 44
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide 1736 SfiI forward primer

<400> 1
 attactcgcg gccagccgg ccatggccca caggtcaagc tcga 44

<210> 2
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide 1735 reverse primer

<400> 2
 ttagagcctg cggccgcctt gtcacgtcg tcctt 35

<210> 3
 <211> 44
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide 1664 SfiI forward primer

<400> 3
 attactcgcg gccagccgg ccatggccca caggtcaagc tcga 44

<210> 4
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide 1664 NotI reverse primer

<400> 4
 ttagagcctg cggccgcctt gtcacgtcg tcctt 35

<210> 5
 <211> 59
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide pBR322 NheI forward stop primer

<400> 5
 cactatggcg tgctgctagc gctatatgcy ttgatgcaat ttctatgagc acccgttct 59

<210> 6
 <211> 38
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide pBR322 EagI reverse primer

<400> 6
 cgtagcccag cgcgtcggcc gccatgccgg cgataatg 38

<210> 7
 <211> 37
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide pBR322 HindIII forward primer

<400> 7
 cagcttatca tcgataagct ttaatgcggt agtttat 37

<210> 8
 <211> 65
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide pBR322 SalI reverse stop primer

<400> 8
 tctcaagggc atcggtcgac gctctccctt atgcgactcc tgcattagga atcagcccag 60
 tagta 65

CLAIMS:

1. A method for generating a polynucleotide sequence or population of sequences from parent single-stranded polynucleotide sequences encoding one or more protein motifs, the method comprising the steps of:

(a) providing a first population of single-stranded polynucleotide molecules and a second population of single-stranded polynucleotide molecules, the first and second populations together constituting plus and minus strands of parent polynucleotide sequences;

(b) carrying out a reaction for digesting the first and second populations of single-stranded polynucleotide molecules with an exonuclease to generate corresponding populations of single-stranded polynucleotide fragments;

(c) contacting said polynucleotide fragments generated from the plus strands with fragments generated from the minus strands; and

(d) amplifying the fragments that anneal to each other to generate at least one polynucleotide sequence encoding one or more protein motifs having altered characteristics as compared to the one or more protein motifs encoded by said parent polynucleotides,

wherein, in step (b), at least one parameter of the reaction used for digestion of the first population of single-stranded polynucleotide molecules is different from the equivalent parameter(s) used in the reaction for digestion of the second population of single-stranded polynucleotide molecules.

2. A method according to claim 1, wherein the reaction parameter is selected from exonuclease type, exonuclease concentration, reaction volume, duration of the digestion reaction, temperature of the reaction mixture, pH of the reaction mixture, length of parent single-stranded polynucleotide sequences, the amount of single-stranded polynucleotide molecules and the buffer composition of the reaction mixture.

3. A method according to claim 1 or 2, wherein the exonuclease used for digestion of the first population of single-stranded polynucleotide molecules is different from the exonuclease used for digestion of the second population of single-stranded polynucleotide molecules.
4. A method according to claim 3 wherein the exonuclease used for digestion of the first population of single-stranded polynucleotide molecules is a 3' exonuclease and the exonuclease used for digestion of the second population of single-stranded polynucleotide molecules is a 5' exonuclease.
5. A method according to any one of claims 1 to 4, wherein the exonuclease concentration used for digestion of the first population of single-stranded polynucleotide molecules is different from the exonuclease concentration used for digestion of the second population of single-stranded polynucleotide molecules.
6. A method according to any one of claims 1 to 5, wherein the reaction volume used for digestion of the first population of single-stranded polynucleotide molecules is different from the reaction volume used for digestion of the second population of single-stranded polynucleotide molecules.
7. A method according to any one of claims 1 to 6, wherein the duration of the digestion reaction used for digestion of the first population of single-stranded polynucleotide molecules is different from the duration of the digestion reaction used for digestion of the second population of single-stranded polynucleotide molecules.
8. A method according to any one of claims 1 to 7, wherein the temperature of the reaction mixture used for digestion of the first population of single-stranded polynucleotide molecules is different from the temperature of the reaction mixture

used for digestion of the second population of single-stranded polynucleotide molecules.

9. A method according to any one of claims 1 to 8, wherein the pH of the reaction mixture used for digestion of the first population of single-stranded polynucleotide molecules is different from the pH of the reaction mixture used for digestion of the second population of single-stranded polynucleotide molecules.

10. A method according to any one of claims 1 to 9, wherein the length of the polynucleotides in the first population of single-stranded polynucleotide molecules is different from the length of the polynucleotides in the second population of single-stranded polynucleotide molecules.

11. A method according to any one of claims 1 to 10, wherein the buffer composition of the reaction mixture used for digestion of the first population of single-stranded polynucleotide molecules is different from the buffer composition of the reaction mixture used for digestion of the second population of single-stranded polynucleotide molecules.

12. A method according to any one of claims 1 to 11, wherein the amount of single-stranded polynucleotide molecules in the first population of single-stranded polynucleotide molecules is different from the amount of single-stranded polynucleotide molecules in the second population of single-stranded polynucleotide molecules.

13. A method according to any one of claims 1 to 12, wherein the first population of single-stranded polynucleotide molecules constitutes the plus strands of parent polynucleotide sequences and the second population of single-stranded polynucleotide molecules constitutes the minus strands of parent polynucleotide sequences.

14. A method according to any one of claims 1 to 11, wherein the polynucleotide molecules of step (a) are DNA molecules.
15. A method according to any one of claims 1 to 14, wherein step (c) further comprises adding primer sequences that anneal to the 3' and/or 5' ends of at least one of the parent polynucleotides under annealing conditions.
16. A method according to any one of claims 1 to 15, wherein the exonuclease used to digest the first and/or second population of single-stranded polynucleotide molecules is selected from the group consisting of BAL 31, exonuclease I, exonuclease V, exonuclease VII, exonuclease T7 gene 6, bacteriophage lambda exonuclease and exonuclease Rec J_f.
17. A method according to any one of claims 1 to 16, wherein a parent polynucleotide sequence or sequences has been subjected to mutagenesis.
18. A method according to any one of claims 1 to 17, wherein one or both of the populations of fragments generated in step (b) are subjected to mutagenesis.
19. A method according to claim 17 or 18, wherein the mutagenesis is error prone PCR.
20. A method according to any one of claims 1 to 19 wherein step (b) is carried out to generate populations of single-stranded fragments of varying lengths.
21. A method according to claim 20, wherein step (b) is controlled to generate a population of single-stranded fragments having an average length of more than approximately 50 nucleotides.

22. A method according to any one of claims 1 to 21 further comprising the step of expressing at least one polynucleotide sequence generated in step (d) to produce the encoded polypeptide *in vitro*.

23. A method according to claim 22 further comprising the step of testing the encoded polypeptide for desired characteristics.

24. A method according to any one of claims 1 to 23, wherein the parent polynucleotide sequence encodes an antibody or fragment thereof.

25. A method according to any one of claims 1 to 23, wherein the parent polynucleotide sequence encodes an enzyme.

26. A method according to any one of claims 1 to 23, wherein the parent polynucleotide sequence encodes an antigen.

27. A method for making a polypeptide having desired properties, the method comprising the following steps:


(a) generating variant forms of a parent polynucleotide using a method according to any one of claims 1 to 21;

(b) expressing the variant polynucleotides produced in step (a) to produce variant polypeptides;

(c) screening the variant polypeptides for desired properties; and


(d) selecting a polypeptide having desired properties from the variant polypeptides.

28. A process for preparing a pharmaceutical composition which comprises, following the identification of an encoded polypeptide with desired characteristics by a method according to claim 27, adding said polynucleotide and/or encoded polypeptide to a pharmaceutically acceptable carrier.


*1/33***FIGURE 1**


2/33

FIGURE 2




3/33


FIGURE 3

4/33**FIGURE 4**

5/33


FIGURE 5



6/33**FIGURE 6**


7/33


FIGURE 7




8/33


FIGURE 8



9/33**FIGURE 9**

10/33

FIGURE 10

*11/33***FIGURE 11**

12/33

FIGURE 12

C:\Program Files\GeneSnap\Ank1ssDNA GFP BAL010626KB.sgd

lane 5

lane 4

lane 3

lane 2


lane 1

13/33


FIGURE 13



Track 4

Profile height



Profile height




14/33**FIGURE 14****Track 3**

15/33


FIGURE 15

Track 2

Profile height




Profile height



16/33

FIGURE 16


C:\Program Files\GeneSnap\karin\Tet\ExoVIIstop1 011123.sgd



1006524342 Mats 11/23/01 02:05:42 PM B=0.00 C=1.00 G=1.00 I=0.200s

*17/33***FIGURE 17**


C:\Program Files\GeneSnap\Helene\AE11F MO152 etc\RecJf fragmentering 011204.sgd



1007466528 Mats 12/04/01 11:48:48 AM B=0.00 C=1.00 G=1.00 I=0.120s

*18/33***FIGURE 18**

C:\Program Files\GeneSnap\Helene\AE11F MO152 etc\RecJf fragmentering 011205.sgd




1007543454 Mats 12/05/01 09:10:54 AM B=0.00 C=1.00 G=1.00 I=0.120s

19/33

FIGURE 19

C:\Program Files\GeneSnap\Tina\Mg Mn 010912.sgd




1000298690 User 09/12/01 01:44:50 PM B=0.00 C=1.00 G=1.00 I=1.360s

20/33

FIGURE 20

C:\Program Files\GeneSnap\Ank\DNase I beh 010721 histogram.sgd





998386880 User 08/21/01 10:41:20 AM B=0.00 C=1.00 G=1.00 I=0.280s

21/33



FIGURE 21

Track 6



Track 6			
Number	Mol. weight	Height	Raw vol.
1	0.00	13.300	5305.14
2	0.00	16.665	30111.09
3	0.00	34.741	18767.46
4	0.00	16.002	31320.35

22/33



FIGURE 22**Track 12**

Track 12			
Number	Mol. weight	Height	Raw vol.
1	0.00	13.846	2872.55
2	0.00	20.893	12279.14
3	0.00	15.181	25217.71

23/33

FIGURE 23

Track 13




Track 13			
Number	Mol. weight	Height	Raw vol.
1	0.00	9.094	2156.20
2	0.00	15.183	97844.64
3	0.00	13.462	14175.73

24/33


FIGURE 24

C:\Program Files\GeneSnap\Helene\AE11F MO152 etc\Mung Bean nuclease fragmenting 011205.sgd




1007542887 Mats 12/05/01 09:01:27 AM B=0.00 C=1.00 G=1.00 I=0.120s

25/33


FIGURE 25 (a)

26/33


FIGURE 25 (b)




27/28

FIGURE 25 (c)

28/33


FIGURE 25 (d)


29/33

FIGURE 26 (a)


30/33

FIGURE 26 (b)





*31/33***FIGURE 26 (c)**

32/33

FIGURE 26 (d)

33/33

FIGURE 27



Target gene cloned
into pFab5c



ssDNA preparation



Enzyme treatment



Reassemble PCR

