

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199736590 B2**  
**(10) Patent No. 738083**

(54) Title  
**Cationic amphiphile/DNA complexes**

(51)<sup>7</sup> International Patent Classification(s)  
**A61K 047/48**

(21) Application No: **199736590**

(22) Application Date: **1997.07.11**

(87) WIPO No: **WO98/02190**

(30) Priority Data

(31) Number	(32) Date	(33) Country
<b>08/679514</b>	<b>1996.07.12</b>	<b>US</b>

(43) Publication Date : **1998.02.09**

(43) Publication Journal Date : **1998.04.02**

(44) Accepted Journal Date : **2001.09.06**

(71) Applicant(s)  
**Genzyme Corporation**

(72) Inventor(s)  
**Ronald K Scheule; Rebecca G Bagley; Simon J Eastman; Seng H Cheng;  
John Marshall; Nelson S Yew; David J Harris; Edward R Lee; Craig S Siegel;  
Chau-Dung Chang; C. Shirley Hubbard**

(74) Agent/Attorney  
**DAVIES COLLISON CAVE, 1 Little Collins Street, MELBOURNE VIC 3000**

(56) Related Art  
**WO 96/18372**  
**WO 93/24640**  
**MORADPUR ET AL BIOCHEM. BIOPHYS. RES. 221, 82-88 (1996)**



36890/n

(51) International Patent Classification <sup>6</sup> : <b>A61K 47/48</b>	<b>A3</b>	(11) International Publication Number: <b>WO 98/02190</b> (43) International Publication Date: <b>22 January 1998 (22.01.98)</b>
(21) International Application Number: <b>PCT/US97/12105</b> (22) International Filing Date: <b>11 July 1997 (11.07.97)</b> (30) Priority Data: <b>08/679,514</b> <b>12 July 1996 (12.07.96)</b> <b>US</b> (71) Applicant: <b>GENZYME CORPORATION [US/US]; One Mountain Road, Framingham, MA 01701-9322 (US).</b> (72) Inventors: <b>SCHEULE, Ronald, K.; 26 East Street, Hopkinton, MA 01748 (US). BAGLEY, Rebecca, G.; 28 Walden Drive #17, Natick, MA 01760 (US). EASTMAN, Simon, J.; 13 Royal Crest Drive #10, Marlboro, MA 01752 (US). CHENG, Seng, H.; 50 Wall Street, Wellesley, MA 02181 (US). MARSHALL, John; 45 West Walnut Street, Milford, MA 01757 (US). YEW, Nelson, S.; 25 Rockdale Hill Circle, West Upton, MA 01568 (US). HARRIS, David, J.; Eastern Avenue, Lexington, MA 02173 (US). LEE, Edward, R.; 175-J Center Street, Lincoln Heights, Quincy, MA 02169 (US). SIEGEL, Craig, S.; 15 Bradford Road, Woburn, MA 01801 (US). CHANG, Chau-Dung; 7 Middleby Road, Lexington, MA 02173 (US). HUBBARD, C., Shirley; 26 Wiley Road, Belmont, MA 02178 (US).</b>	(74) Agents: <b>GARRETT, Arthur, S.; Finnegan, Henderson, Farabow, Garrett &amp; Dunner, L.L.P., 1300 I Street, N.W., Washington, DC 20005-3315 (US) et al.</b> (81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  Published <i>With international search report.</i> (88) Date of publication of the international search report: <b>22 October 1998 (22.10.98)</b>	

(54) Title: **CATIONIC AMPHIPHILE/DNA COMPLEXES**

## (57) Abstract

Novel cationic amphiphiles are provided that facilitate transport of biologically active (therapeutic) molecules into cells. The amphiphiles contain lipophilic groups derived from steroids, from mono or dialkylamines, or from alkyl or acyl groups; and cationic groups, protonatable at physiological pH, derived from amines, alkylamines or polyalkylamines. There are provided also therapeutic compositions prepared typically by contacting a dispersion of one or more cationic amphiphiles with the therapeutic molecules. Therapeutic molecules that can be delivered into cells according to the practice of the invention include DNA, RNA, and polypeptides. Representative uses of the therapeutic compositions of the invention include providing gene therapy, and delivery of antisense polynucleotides or biologically active polypeptides to cells. With respect to therapeutic compositions for gene therapy, the DNA is provided typically in the form of a plasmid for complexing with the cationic amphiphile. Novel and highly effective plasmid constructs are also disclosed, including those that are particularly effective at providing gene therapy for clinical conditions complicated by inflammation. Additionally, targeting of organs for gene therapy by intravenous administration of therapeutic compositions is described.



5

## Cationic Amphiphile / DNA Complexes

10           This application is a continuation-in-part of U.S. patent application  
08/545,473 filed October 19, 1995 and entitled " Organ-Specific Targeting of  
Cationic Amphiphile /DNA Complexes for Gene Therapy" , itself a continuation-  
in-part of U.S. patent application 08/ 540,867 filed October 11, 1995 and entitled  
"Cationic Amphiphiles Containing Steroid Lipophilic Groups for Intracellular  
15 Delivery of Therapeutic Molecules", itself a continuation-in-part of U.S.  
application No. 08/352,479 entitled "Cationic Amphiphiles for Intracellular  
Delivery of Therapeutic Molecules", as filed on December 9, 1994. This  
application also claims the priority of (1) United States provisional patent  
application identified as Express Mail Label TB798223107 US, filed September 26,  
20 1995 and entitled "Molecular Model of Cationic Lipid/DNA Complexes", and (2)  
United States provisional patent application identified as Express Mail Label  
EF109437051 US filed on September 27, 1995 and entitled "Intravenous Delivery  
of Therapeutic Compositions for Gene Therapy" .

          This application is also a continuation-in-part of U.S. patent application  
25 identified as Express Mail Label No. EM288778977 US filed June 20, 1996 and  
entitled " Organ-Specific Targeting of Cationic Amphiphile /DNA Complexes for  
Gene Therapy.

The complete text, claims and drawings of all of the above applications are incorporated herein by reference in their entirety.

### Background of the Invention

The present invention relates to novel cationic amphiphilic compounds  
5 that facilitate the intracellular delivery of biologically active (therapeutic)  
molecules. The present invention relates also to pharmaceutical compositions  
that comprise such cationic amphiphiles, and that are useful to deliver into the  
cells of patients therapeutically effective amounts of biologically active  
molecules. The novel cationic amphiphilic compounds of the invention are  
10 particularly useful in relation to gene therapy.

Effective therapeutic use of many types of biologically active molecules  
has not been achieved simply because methods are not available to cause  
delivery of therapeutically effective amounts of such substances into the  
particular cells of a patient for which treatment therewith would provide  
15 therapeutic benefit. Efficient delivery of therapeutically sufficient amounts of  
such molecules into cells has often proved difficult, if not impossible, since, for  
example, the cell membrane presents a selectively-permeable barrier.  
Additionally, even when biologically active molecules successfully enter targeted  
cells, they may be degraded directly in the cell cytoplasm or even transported to  
20 structures in the the cell, such as lysosomal compartments, specialized for  
degradative processes. Thus both the nature of substances that are allowed to  
enter cells, and the amounts thereof that ultimately arrive at targeted locations  
within cells, at which they can provide therapeutic benefit, are strictly limited.

Although such selectivity is generally necessary in order that proper cell  
25 function can be maintained, it comes with the disadvantage that many  
therapeutically valuable substances (or therapeutically effective amounts thereof)  
are excluded. Additionally, the complex structure, behavior, and environment

presented by an intact tissue that is targeted for intracellular delivery of biologically active molecules often interfere substantially with such delivery, in comparison with the case presented by populations of cells cultured *in vitro* .

5        Examples of biologically active molecules for which effective targeting to a patients' tissues is often not achieved: (1) numerous proteins including immunoglobulin proteins, (2) polynucleotides such as genomic DNA, cDNA, or mRNA (3) antisense polynucleotides; and (4) many low molecular weight compounds, whether synthetic or naturally occurring, such as the peptide hormones and antibiotics.

10        One of the fundamental challenges now facing medical practitioners is that although the defective genes that are associated with numerous inherited diseases (or that represent disease risk factors including for various cancers) have been isolated and characterized, methods to correct the disease states themselves by providing patients with normal copies of such genes (the technique of gene  
15        therapy) are substantially lacking. Accordingly, the development of improved methods of intracellular delivery therefor is of great medical importance.

      Examples of diseases that it is hoped can be treated by gene therapy include inherited disorders such as cystic fibrosis, Gaucher's disease, Fabry's disease, and muscular dystrophy. Representative of acquired disorders that can  
20        be treated are: (1) for cancers — multiple myeloma, leukemias, melanomas, ovarian carcinoma and small cell lung cancer; (2) for cardiovascular conditions — progressive heart failure, restenosis, and hemophilias; and (3) for neurological conditions — traumatic brain injury.

      Gene therapy requires successful transfection of target cells in a patient.  
25        Transfection may generally be defined as the process of introducing an expressible polynucleotide (for example a gene, a cDNA, or an mRNA patterned thereon) into a cell. Successful expression of the encoding polynucleotide leads



intracellular delivery (whether for *in vitro* or *in vivo* application) meet this definition. One particularly important class of such amphiphiles is the cationic amphiphiles. In general, cationic amphiphiles have polar groups that are capable of being positively charged at or around physiological pH, and this property is understood in the art to be important in defining how the amphiphiles interact with the many types of biologically active (therapeutic) molecules including, for example, negatively charged polynucleotides such as DNA.

Examples of cationic amphiphilic compounds that have both polar and non-polar domains and that are stated to be useful in relation to intracellular delivery of biologically active molecules are found, for example, in the following references, which contain also useful discussion of (1) the properties of such compounds that are understood in the art as making them suitable for such applications, and (2) the nature of structures, as understood in the art, that are formed by complexing of such amphiphiles with therapeutic molecules intended for intracellular delivery.

- (1) Felgner, et al., Proc. Natl. Acad. Sci. USA, 84, 7413-7417 (1987) disclose use of positively-charged synthetic cationic lipids including N-[1(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride ("DOTMA"), to form lipid/DNA complexes suitable for transfections. See also Felgner et al., The Journal of Biological Chemistry, 269(4), 2550-2561 (1994).
- (2) Behr et al., Proc. Natl. Acad. Sci. USA, 86, 6982-6986 (1989) disclose numerous amphiphiles including dioctadecylamidoglycylspermine ("DOGS").
- (3) U.S. Patent 5, 283,185 to Epand et al. describes additional classes and species of amphiphiles including 3 $\beta$ -[N-(N<sup>1</sup>,N<sup>1</sup>-dimethylaminoethane)-carbamoyl] cholesterol, termed "DC-chol".
- (4) Additional compounds that facilitate transport of biologically active molecules into cells are disclosed in U.S. Patent No. 5,264,618 to Felgner et al. See

also Felgner et al., The Journal Of Biological Chemistry, 269(4), pp. 2550- 2561 (1994) for disclosure therein of further compounds including "DMRIE" 1,2-dimyristyloxypropyl-3-dimethyl-hydroxyethyl ammonium bromide, which is discussed below.


- 5 (5) Reference to amphiphiles suitable for intracellular delivery of biologically active molecules is also found in U.S. Patent No. 5,334,761 to Gebeyehu et al., and in Felgner et al., Methods (Methods in Enzymology), 5, 67- 75 (1993).

Although the compounds mentioned in the above-identified references have been demonstrated to facilitate (although in many such cases only *in vitro* )  
10 the entry of biologically active molecules into cells, it is believed that the uptake efficiencies provided thereby are insufficient to support numerous therapeutic applications, particularly gene therapy. Additionally, since the above-identified compounds are understood to have only modest activity, substantial quantities thereof must be used leading to concerns about the toxicity of such compounds  
15 or of the metabolites thereof. Accordingly there is a need to develop a "second generation" of cationic amphiphiles whose activity is so sufficient that successful therapies can be achieved therewith.

Summary of the Invention

This invention provides for cationic amphiphiles that are particularly effective to facilitate transport of biologically active molecules into cells. The cationic amphiphiles of the invention are divided into four (4) groups, although it will be seen that there are certain structural and functional features that many of the amphiphiles share.

Accordingly, there are provided cationic amphiphiles of Group I (see Figure 1, panels A, B, and C) capable of facilitating transport of biologically active molecules into cells, said amphiphiles having the structure (I),



wherein:

- 15 Z is a steroid;  
 X is a carbon atom or a nitrogen atom;  
 Y is a short linking group, or Y is absent;  
 R<sup>3</sup> is H, or a saturated or unsaturated aliphatic group;  
 R<sup>1</sup> is -NH-, an alkylamine, or a polyalkylamine;  
 20 R<sup>4</sup> is H, or a saturated or unsaturated aliphatic group;  
 R<sup>2</sup> is -NH-, an alkylamine, or a polyalkylamine;  
 and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup>, except that both R<sup>1</sup> and R<sup>2</sup> cannot be -NH-.



In one preferred embodiment, the steroid component "Z" is selected from the group consisting of 3-sterols, wherein said sterol molecule is linked by the 3-O- group thereof, or by N- in replacement thereof, to Y (or directly to X, if Y is absent). According to this aspect of the invention, particularly effective


5 amphiphiles include, for example, spermidine cholesterol carbamate (N<sup>4</sup>-spermidine cholesteryl carbamate, amphiphile No. 53), and spermine cholesterol carbamate (N<sup>4</sup>-spermine cholesteryl carbamate, amphiphile No. 67), and amphiphiles patterned thereon.

In a further preferred embodiment, the steroid group is linked to Y (or directly to X, if Y is absent) from ring position 17 of the steroid nucleus (see Figures 1 and 22), or from the arm that normally extends from position 17 in many steroids (see the structure of cholesterol in Figure 1), or from any shortened form of said arm.

In other preferred embodiments, within linking group Y are contained no more than about three or four atoms that themselves form a bridge of covalent bonds between X and Z. In a specific preferred embodiment of the invention, Y is a linking group wherein no more than one atom of said group forms a bond with both X and Z, or Y is absent.


20 Representative amphiphiles provided according to Group I include:





No. 75


$N^1, N^8$ -Bis(3-aminopropyl)- $N^4$ -spermidine cholesteryl carbamate



No. 78

$N(N^4$ -3-aminopropyl)spermidine cholesteryl carbamate

Additionally there are provided cationic amphiphiles of Group II (see Figure 5) capable of facilitating transport of biologically active molecules into cells said amphiphiles having the structure (II),



(II)

10 wherein:

Z is a steroid;

X is a carbon atom or a nitrogen atom;

Y is a linking group or Y is absent;

$R^3$  is an amino acid, a derivatized amino acid, H or alkyl;

15  $R^1$  is  $-NH-$ , an alkylamine, or a polyalkylamine;

$R^4$  is an amino acid, a derivatized amino acid, H or alkyl;

$R^2$  is  $-NH-$ , an alkylamine, or a polyalkylamine;



Z is an alkylamine or a dialkylamine, linked by the N-atom thereof, to Y ( or directly to X, if Y is absent ), wherein if Z is a dialkylamine, the alkyl groups thereof can be the same or different;

X is a carbon atom or a nitrogen atom;

5 Y is a short linking group , or Y is absent;

R<sup>3</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>1</sup> is -NH- , an alkylamine, or a polyalkylamine;

R<sup>4</sup> is H, or a saturated or unsaturated aliphatic group;


R<sup>2</sup> is -NH- , an alkylamine, or a polyalkylamine;

10 and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup> , except that both R<sup>1</sup> and R<sup>2</sup> cannot be -NH-.


With respect to amphiphiles provided according to Structure (III), it is again preferred that within linking group Y there are contained no more than about three or four atoms that themselves form a bridge of covalent bonds  
 15 between X and Z. In a specific preferred embodiment of the invention, Y is a linking group, such as > C=O, wherein no more than one atom of said group forms a bond with both X and Z, or Y is absent.

Representative amphiphiles provided according to Group III include: .

20



Additionally there are provided cationic amphiphiles of Group IV (see Figure 7) capable of facilitating transport of biologically active molecules into cells said amphiphiles having the structure (IV),



wherein:

A and B are independently O, N or S;

10  $R^5$  and  $R^6$  are independently alkyl or acyl groups and may be saturated or contain sites of unsaturation;

C is selected from the group consisting of  $-CH_2-$ ,  $>C=O$ , and  $>C=S$ ;

E is a carbon atom or a nitrogen atom;

D is a linking group such as  $-NH(C=O)-$  or  $-O(C=O)-$ , or D is absent;

$R^3$  is H, or a saturated or unsaturated aliphatic group;


15  $R^1$  is  $-NH-$ , an alkylamine, or a polyalkylamine;

$R^4$  is H, or a saturated or unsaturated aliphatic group;

$R^2$  is  $-NH-$ , an alkylamine, or a polyalkylamine;

and wherein  $R^1$  is the same or is different from  $R^2$ , except that both  $R^1$  and  $R^2$  cannot be  $-NH-$ .

Representative amphiphiles of Group IV include:



The invention provides also for pharmaceutical compositions that  
 5 comprise one or more cationic amphiphiles, and one or more biologically active molecules, wherein said compositions facilitate intracellular delivery in the tissues of patients of therapeutically effective amounts of the biologically active molecules. The pharmaceutical compositions of the invention may be formulated to contain one or more additional physiologically acceptable  
 10 substances that stabilize the compositions for storage and/or contribute to the successful intracellular delivery of the biologically active molecules.

In a further aspect, the invention provides a method for facilitating the transfer of biologically active molecules into cells comprising the steps of:  
 preparing a dispersion of a cationic amphiphile of the invention; contacting said  
 15 dispersion with a biologically active molecule to form a complex between said amphiphile and said molecule, and contacting cells with said complex thereby facilitating transfer of said biologically-active molecule into the cells.

For pharmaceutical use, the cationic amphiphile(s) of the invention may be formulated with one or more additional cationic amphiphiles including those  
 20 known in the art, or with neutral co-lipids such as dioleoylphosphatidyl-ethanolamine, ("DOPE"), to facilitate delivery to cells of the biologically active molecules. Additionally, compositions that comprise one or more cationic

amphiphiles of the invention can be used to introduce biologically active molecules into plant cells, such as plant cells in tissue culture.

Additionally, the present application provides for novel plasmids suitable for complexing with the amphiphiles of the invention in order to treat patients by gene therapy, so that a high level of expression of the appropriate therapeutic transgene can be achieved. Representative examples thereof include the plasmid pCMVHI and pCFI. pCF1 plasmid contains the enhancer/promoter region from the immediate early gene of cytomegalovirus. The plasmid also contains a hybrid intron located between the promoter and the transgene cDNA. The polyadenylation signal of the bovine growth hormone gene was selected for placement downstream from the transgene. These and other features contribute substantially to the improved transgene expression possible with this plasmid.

Further enhancements in plasmid performance are made possible by the provision of replicating episomal plasmids. Additional therapeutic enhancements are made possible by providing plasmids in which expression of the therapeutic transgene is placed under the control of a transcriptional promoter that is sensitive to the concentration of inflammation-related substances in the target tissue. Such plasmids are of particular use for the treatment of clinical cases in which inflammation is a major complication.

In a still further embodiment of the invention, particular organs or tissues may be targeted for gene therapy, by intravenous administration of amphiphile/transgene complexes, by adjusting the ratio of amphiphile to DNA in such complexes, and by adjusting the apparent charge or zeta potential thereof.

Further additional and representative aspects of the invention are described according to the Detailed Description of the Invention which follows directly.

### Brief Description of the Drawings

- FIGURE 1 depicts representative Group I cationic amphiphiles.
- FIGURE 2 depicts representative steroid lipophilic groups.
- FIGURE 3 depicts representative steroid lipophilic groups.
- 5 FIGURE 4 depicts a transacylation reaction.
- FIGURE 5 depicts representative Group II cationic amphiphiles.
- FIGURE 6 depicts representative Group III cationic amphiphiles.
- FIGURE 7 depicts representative Group IV cationic amphiphiles.
- FIGURE 8 provides a route of synthesis for spermidine cholesterol carbamate.
- 10 FIGURE 9 provides a route of synthesis for spermine cholesterol carbamate.
- FIGURE 10 provides a comparison of in vivo transfection efficiency for certain cationic amphiphiles under particular conditions.
- FIGURE 11 is a depiction of in vivo transfection efficiency as a function of DNA concentration for a particular cationic amphiphile.
- 15 FIGURE 12 is a depiction of in vivo transfection efficiency as a function of concentration of a particular cationic amphiphile.
- FIGURE 13 provides relative transfection efficiencies for Group I amphiphiles.
- FIGURE 14 provides relative transfection efficiencies for Group II amphiphiles.
- FIGURE 15 provides relative transfection efficiencies for Group IV amphiphiles.
- 20 FIGURE 16 provides a map of pCMVHI-CAT plasmid.
- FIGURE 17 shows the hybrid intron of pCMVHI-CAT.
- FIGURE 18 (panel A) provides a map of pCF1/CAT plasmid.
- FIGURE 18 (panel B) provides a map of pCF2/CAT plasmid.
- FIGURE 19 (panel A) shows a plot of corrected chloride ion transport in pCMV-CFTR
- 25 transfected nasal polyp epithelial cells from a cystic fibrosis patient.
- FIGURE 19 (panel B) shows a plot of chloride ion transport using pCMV- $\beta$ -galactosidase control.
- FIGURE 20 provides a map of pMyc4-CFTR plasmid.
- FIGURE 21 demonstrates intravenous targeting of the heart and lung.

FIGURE 22 demonstrates expression of SEAP following intravenous administration in BALB/c mice.

### Detailed Description of the Invention

#### Information Concerning the Structure of Cationic Amphiphiles of the Invention

This invention provides for cationic amphiphile compounds, and compositions containing them, that are useful to facilitate transport of  
5 biologically active molecules into cells. The amphiphiles are particularly useful in facilitating the transport of biologically active polynucleotides into cells, and in particular to the cells of patients for the purpose of gene therapy.

Cationic amphiphiles according to the practice of the invention possess several novel features. These features may be seen in comparison with, for  
10 example, cationic amphiphile structures such as those disclosed in U.S. Patent No. 5, 283,185 to Epanand et al., a representative structure of which is 3 $\beta$  [N-(N<sup>1</sup>N<sup>1</sup> - dimethylaminoethane)-carbamoyl] cholesterol, commonly known as "DC-chol", and to those disclosed by Behr et al. Proc. Natl. Acad. Sci., USA, 86, 6982- 6986 (1989), a representative structure of which is dioctadecylamidolo-  
15 glycylspermine ("DOGS").

Cationic amphiphiles of the present invention contain distinctive structural features: (1) the presence of a lipophilic group which is connected directly, or through a linking group, to two cationic groups (see below) that themselves comprise amino, alkylamine or polyalkylamine groups, there  
20 resulting an overall and novel "T-shaped" structure; and (2) in many cases, and in comparison with numerous art-recognized amphiphiles, the use of a relatively short linking group to bring into close proximity the lipophilic and cationic regions of the amphiphile. Without being limited as to theory, it is believed that these features contribute substantially to the transfection-enhancing capability of  
25 these compounds. As an example of this, Figure 10 below demonstrates the very substantial in vivo transfection-enhancing capability of spermidine cholesterol

carbamate (a novel amphiphile of the invention) in comparison to DC-chol and DMRIE— two well recognized transfectants.

In connection with the practice of the present invention, it is noted that "cationic" means that the R groups, as defined herein, tend to have one or more  
5 positive charges in a solution that is at or near physiological pH. Such cationic character may enhance interaction of the amphiphile with therapeutic molecules (such as nucleic acids) or with cell structures (such as plasma membrane glycoproteins) thereby contributing to successful entry of the therapeutic molecules into cells, or processing within subcompartments (such as the nucleus)  
10 thereof. In this regard, the reader is referred to the numerous theories in the art concerning transfection-enhancing function of cationic amphiphiles, none of which is to be taken as limiting on the practice of the present invention.


Biological molecules for which transport into cells can be facilitated according to the practice of the invention include, for example, genomic DNA,  
15 cDNA, mRNA, antisense RNA or DNA, polypeptides and small molecular weight drugs or hormones. Representative examples thereof are mentioned below in connection with the description of therapeutic (pharmaceutical) compositions of the invention.

In an important embodiment of the invention the biologically active  
20 molecule is an encoding polynucleotide that is expressed when placed in the cells of a patient leading to the correction of a metabolic defect. In a particularly important example, the polynucleotide encodes for a polypeptide having an amino acid sequence sufficiently duplicative of that of human cystic fibrosis transmembrane regulator ("CFTR") to allow possession of the biological  
25 property of epithelial cell anion channel regulation.

As aforementioned, characteristic and novel features of the amphiphiles of the invention include first, that the linking group that connects the two cationic


amine groups to the lipophilic group is very short, or absent entirely, and second, that the resultant linking of the the two cationic R groups to the lipophilic group forms a T-shaped structure when viewed from the position of atom "X" (a carbon or nitrogen atom) as depicted, for example, in Structures (I), (II), (III) and  
 5 (IV, see atom "E").

As examples of the cationic amphiphiles of the invention, both spermidine cholesterol carbamate (N<sup>4</sup>-spermidine cholesteryl carbamate) and spermine cholesterol carbamate (N<sup>4</sup>-spermine cholesteryl carbamate) have been determined to be superior transfectants in vivo in comparison with non "T-  
 10 shaped" amphiphiles having otherwise equivalent amounts of cationic alkylamine structure. Superior performance (see also Example 3) has been determined for:




( spermidine cholesterol carbamate )

15 in comparison with, for example,




(N<sup>1</sup>-spermidine cholesteryl carbamate).


Additionally, superior performance has been determined for



5 in comparison with, for example,



and




Applicants have also noted that numerous of the cationic amphiphiles of  
 15 the invention have structural features in common with naturally occurring  
 polyamines such as spermine and spermidine (including N-atom spacing). In  
 this regard, the structures of amphiphiles 53, 67, 78, 90, and 91 are representative.  
 As can be seen by examination of the data in Figures 13, 14 and 15, the placement  
 of the nitrogen atoms in the polar head groups of the amphiphiles such that they  
 20 are separated by one or more combinations of 3 and 4 carbon atoms leads to high  
in vivo transfection efficiency for plasmid transgenes complexed therewith.  
 Applicants have also noted that these in-common structural features may have a

useful effect upon the binding of the amphiphiles to DNA, and on interaction with cell surface polyamine receptors. Interaction with cell polyamine receptors may be particularly important with respect to the treatment of cancer cells by gene therapy, since the DNA replication requirements of such cells may lead to high level expression of such receptors.

#### Group I Amphiphiles

In connection with the design of the Group I amphiphiles of the invention, the following considerations are of note. Many of these design features are then discussed in connection with the other amphiphiles of the invention, those classified under Groups II, III and IV.

Accordingly, there are provided cationic amphiphiles of Group I (see Figure 1, panels A, B, and C) capable of facilitating transport of biologically active molecules into cells, said amphiphiles having the structure (I),



wherein:

Z is a steroid;

X is a carbon atom or a nitrogen atom;

Y is a short linking group, or Y is absent;

R<sup>3</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>1</sup> is --NH-- , an alkylamine, or a polyalkylamine;

R<sup>4</sup> is H, or a saturated or unsaturated aliphatic group;

$R^2$  is  $-NH-$ , an alkylamine, or a polyalkylamine;

and wherein  $R^1$  is the same or is different from  $R^2$ , except that both  $R^1$  and  $R^2$  cannot be  $-NH-$ .

#### The Linking Group

5            Preferably the linking group that connects the lipophilic group to the two cationic R groups is relatively short. It is preferred that within linking group Y are contained no more than about three or four atoms that themselves form a bridge of covalent bonds between X and Z. Examples of Y groups include  $-(CH_2)_2-$ ;  $-(CH_2)_3-$ ;  $-(CH_2)-(C=O)-$ ;  $-(CH_2)_n-NH-(C=O)-$  where n is preferably  
10 4 or less. Additional linking groups useful in the practice of the invention are those patterned on small amino acids such as glyciny, alanyl, beta-alanyl, seriny, and the like.

              With respect to the above representations, the left hand side thereof-as depicted- is intended to bond to atom "X", and the right hand side thereof to  
15 group "Z" (see structure I).

              In certain preferred embodiments of the invention, Y is a linking group wherein no more than one atom of this group forms a bond with both "X" and "Z". Examples of preferred linking groups include  $--CH_2-$ ,  $>C=S$ , and  $>C=O$ . Alternatively, the linking group "Y" may be absent entirely.

20            As aforementioned (see Structure I, directly above), "X" forms a connecting point in the amphiphiles to which is also attached the two cationic R groups. As can be seen therein (see also Figure 1), the placement of the nitrogen atom that represents "X" clearly causes the molecule to assume a T-shape.

#### Steroid Lipophilic Groups

25            Cationic amphiphiles according to the practice of the invention may include a variety of structures as lipophilic group. Steroids represent a preferred group of such structures.

With respect to the design and orientation of steroids as lipophilic groups according to the practice of the invention, the following considerations are of note. Steroids are widely distributed in the animal, microbial and plant kingdoms. They may be defined as solid alcohols that typically contain, as their  
5 basic skeleton, 17 carbon atoms arranged in the form of a perhydrocyclopenteno-phenanthrene ring system. Accordingly, such compounds include bile acids, cholesterol and related substances, vitamin D, certain insect molting hormones, certain sex hormones, corticoid hormones, certain antibiotics, and derivatives of all of the above wherein additional rings are added or are deleted from the basic  
10 structure. [see Natural Products Chemistry, K. Nakanashi et al. eds., Academic Press, Inc., New York (1974), volume 1, at Chapter 6 for a further discussion of the broad classes of molecules that are understood in the art to be steroids]. Additionally, for the purposes of the invention, the term steroid is used broadly to include related molecules derived from multiple isoprenoid units, such as  
15 vitamin E. Steroids representative of those useful in the practice of the invention are shown in Figures 1, 2, 3 and 5.

As elaborated below, certain preferred amphiphiles of the invention include a steroid component "Z" that is selected from the group consisting of 3-sterols, wherein said sterol molecule is linked by the 3-O- group thereof, or by  
20 N- in replacement thereof, to Y (see Figure 1). Such structures include, for example, spermidine cholesterol carbamate, spermine cholesterol carbamate, spermidine 7-dehydrocholesteryl carbamate, lysine 3-N-dihydrocholesteryl carbamate, spermidine cholestamine urea, and N-3-aminopropyl-N-4-aminobutylcholestamine.

25 In a further preferred embodiment, the steroid group is linked to Y (or directly to X if Y is absent) from ring position 17 of the steroid nucleus (see

Figures 1 and 3), or from the arm that normally extends from position 17 in many steroids (see Figures 1 and 3), or from any shortened form of said arm.

In connection with the selection of steroids for inclusion in the amphiphiles of the invention, it is preferred that the molecules have structures  
5 which can be metabolized by the body and are nontoxic at the doses thereof that are used. Preferred are steroids such as cholesterol and ergosterol that are substantially non toxic and which possess biologically normal stereospecificity in order to facilitate their safe metabolism in patients. Additional steroids useful in the practice of the invention include, for example, ergosterol B1, ergosterol B2,  
10 ergosterol B3, androsterone, cholic acid, desoxycholic acid, chenodesoxycholic acid, lithocholic acid and, for example, various derivatives thereof as are shown in the panels of Figures 2 and 3.


With respect to the orientation of the steroid lipophilic group, that is, how the group is attached (with or without a linker) to the cationic (alkyl) amine  
15 groups of an amphiphile, the following further information is of note. Any ring position or substituent on the steroid can in general be used as point of attachment. It is preferred, however, to use a point of attachment that (1) minimizes the complexity of chemical syntheses, and (2) is positioned near either "end" of the steroid molecule, for example, a position near ring position  
20 3, or near ring position 17 (or the arm that typically extends therefrom). Such positions provide an orientation of the steroid with respect to the rest of the amphiphile structure that facilitates bilayer formation, and/or micelle formation, and/or stabilizes interaction with the biologically active molecules to be carried into the target cells. Representative structures showing attachment of the  
25 cationic (alkyl) amine groups to the steroid lipophilic group through the arm extending from ring position 17 thereof are shown in Figure 3 (panels A, B). With respect to this type of structure, it is further preferred that any polar groups on

the steroid, such as may be attached to ring position 3, be either removed or capped (for example, hydroxy as methoxy) to avoid potentially destabilizing bilayer or micelle structures.


The representation in Figure 3 of cationic amphiphiles in which the steroid lipophilic group thereof is linked to the cationic alkylamine groups through steroid ring position 17 is but an example of the invention. Similarly, the representation in Figures 1 to 4 of cationic amphiphiles in which the steroid lipophilic group thereof is linked to the cationic alkylamine groups through steroid ring position 3 is an example of the invention. As aforementioned, use of any steroid ring position (or moiety or branch extending therefrom) as point of attachment is within the practice of the invention.


Preferred steroids for use as group "Z" according to the practice of the invention include:

3- sterols (derived from cholesterol)




3-N steryl groups (patterned on cholesterol)




ergosterol and derivatives

Representative species of steroid that are patterned on ergosterol and that may be used to define the structure of cationic amphiphiles of the invention include: ergosterol (double bonds as shown); ergosterol B1 ( $\Delta$  8, 9;  $\Delta$  14, 15;  $\Delta$  22, 23); ergosterol B1 ( $\Delta$  6, 7;  $\Delta$  8, 14;  $\Delta$  22, 23); ergosterol B1 ( $\Delta$  7, 8;  $\Delta$  14, 15;  $\Delta$  22, 23); and lumisterol ( the 9b-H isomer of ergosterol).

cholic acid and derivatives

Representative species of steroid that are patterned on cholic acid and that may be used to define the structure of cationic amphiphiles of the invention include: cholic acid wherein r<sup>1</sup> and r<sup>2</sup> = OH; desoxycholic acid wherein r<sup>1</sup> = H and r<sup>2</sup> = OH; chenodesoxycholic acid wherein r<sup>1</sup> = OH and r<sup>2</sup> = H; and lithocholic acid wherein r<sup>1</sup> and r<sup>2</sup> = H.

androsterone and derivatives thereof5 Selection of Groups R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup>For R<sup>3</sup> and R<sup>4</sup>:

According to the practice of the invention R<sup>3</sup> and R<sup>4</sup> are, independently, H, or saturated or unsaturated aliphatic groups. The aliphatic groups can be  
 10 branched or unbranched. Representative groups include alkyl, alkenyl, and cycloalkyl.

For R<sup>1</sup> and R<sup>2</sup>:

R<sup>1</sup> and R<sup>2</sup> represent structures recognized in the art as being amine; alkylamines (including primary, secondary, and tertiary amines), or extended  
 15 versions thereof-herein termed "polyalkylamines". It is understood that both alkylamine and polyalkylamine groups as defined herein may include one or more carbon-carbon double bonds and the use of such alkenylamines is therefore within the practice of the invention.

Representative alkylamines include: (a) --NH-(CH<sub>2</sub>)<sub>z</sub> - where z is other  
 20 than 0; (b) - [[CH<sub>3</sub>(CH<sub>2</sub>)<sub>y</sub>N] -(CH<sub>2</sub>)<sub>z</sub> - where z is other than 0; and (c) - [[CH<sub>3</sub>(CH<sub>2</sub>)<sub>x</sub>][CH<sub>3</sub>(CH<sub>2</sub>)<sub>y</sub>]]N -(CH<sub>2</sub>)<sub>z</sub> - where z is other than 0.

With respect to the circumstance where one or both of R<sup>1</sup> and R<sup>2</sup> are tertiary amines, such as is represented in Structure (c) above, it is understood that a hydrogen atom corresponding to either R<sup>3</sup> or R<sup>4</sup>, as appropriate, may or may

not be present since such hydrogen atoms correspond to the N:H(+) structure whose level of protonation will vary according to pH.

The term "polyalkylamine" as referred to herein defines a polymeric structure in which at least two alkylamines are joined. The alkylamine units that are so joined may be primary or secondary, and the polyalkylamines that result may contain primary, secondary, or tertiary N-atoms. The alkylamine (sub)units may be saturated or unsaturated, and therefore the term "alkylamine" encompasses alkenylamines in the description of the invention.

Representative resultant polyalkylamines include: (d)  $-\text{[NH-(CH}_2\text{)}_z\text{]}_q$   $-\text{}$ , where  $z$  is other than 0, and  $q$  is 2 or higher; (e)  $-\text{[NH-(CH}_2\text{)}_y\text{]}_p - \text{[NH-(CH}_2\text{)}_z\text{]}_q - \text{}$ , where  $y$  and  $z$  are each other than 0, and  $p$  and  $q$  are each other than 0; (f)  $-\text{[NH-(CH}_2\text{)}_x\text{]}_n - \text{[NH-(CH}_2\text{)}_y\text{]}_p - \text{[NH-(CH}_2\text{)}_z\text{]}_q - \text{}$ , where  $x$ ,  $y$ , and  $z$  are each other than 0, and  $n$ ,  $p$  and  $q$  are each other than 0; (g)  $-\text{[NH-(CH}_2\text{)}_w\text{]}_m - \text{[NH-(CH}_2\text{)}_x\text{]}_n - \text{[NH-(CH}_2\text{)}_y\text{]}_p - \text{[NH-(CH}_2\text{)}_z\text{]}_q - \text{}$ , where  $w$ ,  $x$ ,  $y$ , and  $z$  are each other than 0, and  $m$ ,  $n$ ,  $p$ , and  $q$  are each other than 0; (h)  $-\text{[NH-(CH}_2\text{)}_w\text{]}_m - \text{[NH-(CH}_2\text{)}_x\text{]}_n - \text{[CH}_3\text{(CH}_2\text{)}_y\text{N]} - \text{(CH}_2\text{)}_z - \text{}$ , where  $x$ ,  $n$  and  $z$  are each other than 0; (i)  $-\text{[NH-(CH}_2\text{)}_w\text{]}_p - \text{[CH}_3\text{(CH}_2\text{)}_x\text{N]} - \text{(CH}_2\text{)}_y - \text{[NH-(CH}_2\text{)}_z\text{]}_q - \text{}$ , where  $w$ ,  $p$ ,  $y$ ,  $z$ , and  $p$  are each other than 0; and (j)  $-\text{[NH-(CH}_2\text{)}_v\text{]}_l - \text{[NH-(CH}_2\text{)}_w\text{]}_m - \text{[NH-(CH}_2\text{)}_x\text{]}_n - \text{[NH-(CH}_2\text{)}_y\text{]}_p - \text{[NH-(CH}_2\text{)}_z\text{]}_q - \text{}$ , where  $v$ ,  $w$ ,  $x$ ,  $y$ , and  $z$  are each other than 0, and  $l$ ,  $m$ ,  $n$ ,  $p$ , and  $q$  are each other than 0.

As mentioned above  $R^1$  and  $R^2$ , independently, can be  $-\text{NH}-$ , an alkylamine, or a polyalkylamine, and can be the same or different from each other, except that both  $R^1$  and  $R^2$  cannot be  $-\text{NH}-$  in order to (1) preserve the "T-shape" of the resultant compound, and (2) to provide for the stability thereof. It is preferred that - in combination- the combined backbone length of  $R^3R^1$  (or of  $R^4R^2$ ) be less than about 40 atoms of nitrogen and carbon, more

preferably less than about 30 atoms of nitrogen and carbon.

In the case where the R<sup>1</sup> group adjacent to R<sup>3</sup> (or R<sup>2</sup> adjacent to R<sup>4</sup>) includes a terminal nitrogen atom that defines a tertiary center, then a quaternary amine is formed (at that nitrogen atom of R<sup>1</sup>) if R<sup>3</sup> is an aliphatic group, and a  
 5 tertiary amine remains (at that nitrogen atom of R<sup>1</sup>) if R<sup>3</sup> is H. Accordingly, with respect to such resultant R<sup>3</sup>R<sup>1</sup> or R<sup>4</sup>R<sup>2</sup> structures, representative respective formulas are:

(k) H-(CH<sub>2</sub>)<sub>(w)</sub> — [[CH<sub>3</sub>(CH<sub>2</sub>)<sub>x</sub>][CH<sub>3</sub>(CH<sub>2</sub>)<sub>y</sub>]N] — (CH<sub>2</sub>)<sub>z</sub> — , where w and z are each other than zero; and (l) H — [[CH<sub>3</sub>(CH<sub>2</sub>)<sub>x</sub>][CH<sub>3</sub>(CH<sub>2</sub>)<sub>y</sub>]N] — (CH<sub>2</sub>)<sub>z</sub> — ,  
 10 where z is other than zero.

In connection with interpreting the structural diagrams described herein, it is intended that the attachment of R<sup>3</sup>R<sup>1</sup>— (or R<sup>4</sup>R<sup>2</sup>— ) structures to atom "X" is through the right hand side (as depicted) of the R<sup>3</sup>R<sup>1</sup>—, that is, through a CH<sub>2</sub>— moiety. The coefficients (i.e. v, w, x, y, and z and l, m, n, p, and q) as  
 15 depicted herein represent whole numbers. For the purposes of the invention, "whole number" means 0 and the natural numbers 1,2,3,4,5,6.....and up, unless specifically restricted.

With respect to the amphiphiles of the invention including those represented by formulas (a) to (l), it is noted that there are certain preferences  
 20 concerning the design of such groups depending on whether atom 'X' as it is shown according to structure (I) above, is a nitrogen atom or a carbon atom. If "X" is nitrogen, then amphiphiles containing R<sup>3</sup>-R<sup>1</sup> ( or R<sup>4</sup>-R<sup>2</sup> ) groups that end in an N atom [ i.e formula (e) where z equals 0 and q=1; formula (h) where z equals 0] are not preferred, since the resultant N-N linkage involving position X  
 25 results in an amphiphile that may be unstable and/or difficult to prepare. An additional group of structures that are difficult to prepare and/or are unstable is represented, for example, by the R sequence (whether in R<sup>1</sup>, or bridging R<sup>1</sup> and

$R^3$ ) -NH-CH<sub>2</sub>-NH-CH<sub>2</sub>- . Accordingly, use of such structures [ i.e. formula (a) where Z equals 1, formula (e) where one or both of y and z equals 1] in the practice of the invention is not preferred.

With respect to the design of structures (such as those depicted above) for  
5 inclusion in cationic amphiphiles, the following further considerations are of  
note. Any combination of alternating amine and alkyl moieties creates an R  
structure within the scope of the invention. A polyalkylamine may be  
represented, for example, by the formulas above, although many more structures  
(such structures being within the scope of the invention) can be depicted by  
10 extending the number of, or types or combinations of, alkylamine subunits  
within the amphiphile structure. That further such variations can be made is  
apparent to those skilled in the art.

It is noted that a polyalkylamine group (or resultant  $R^3R^1$  group) that is  
very long may interfere, for example, with the solubility of the resultant  
15 amphiphile, or interfere with its ability to stably interact with the biologically  
active molecule selected for intracellular delivery. In this regard,  
polyalkylamines (or resultant  $R^3R^1$  groups) having a backbone length of about  
40 nitrogen and carbon atoms, or more, may not be suitable for inclusion in  
amphiphiles. However, for each such proposed structure, its properties may be  
20 determined by experimentation, and its use is nonetheless within the practice of  
the invention.

Accordingly, specific alkylamine and polyalkylamine structures result as follows:

Table 1

<u>For R<sup>1</sup> and/or R<sup>2</sup></u>	
5	(1) -NH-
	(2) -NH-(CH <sub>2</sub> ) <sub>2</sub> -
	(3) -NH-(CH <sub>2</sub> ) <sub>3</sub> -
	(4) -NH-(CH <sub>2</sub> ) <sub>4</sub> -
	(5) -NH-(CH <sub>2</sub> ) <sub>6</sub> -
10	(6) -NH-(CH <sub>2</sub> ) <sub>3</sub> -NH-(CH <sub>2</sub> ) <sub>4</sub> -
	(7) -NH-(CH <sub>2</sub> ) <sub>2</sub> -NH-(CH <sub>2</sub> ) <sub>2</sub> -
	(8) -NH-(CH <sub>2</sub> ) <sub>4</sub> -NH-(CH <sub>2</sub> ) <sub>3</sub> -
	(9) -NH-(CH <sub>2</sub> ) <sub>y</sub> -NH-(CH <sub>2</sub> ) <sub>z</sub> -
	(10) -NH-(CH <sub>2</sub> ) <sub>x</sub> -NH-(CH <sub>2</sub> ) <sub>y</sub> -NH-(CH <sub>2</sub> ) <sub>z</sub> -
15	(11) -NH-(CH <sub>2</sub> ) <sub>w</sub> -NH-(CH <sub>2</sub> ) <sub>x</sub> -NH-(CH <sub>2</sub> ) <sub>y</sub> -NH-(CH <sub>2</sub> ) <sub>z</sub> -
	(12) -NH-(CH <sub>2</sub> ) <sub>v</sub> -NH-(CH <sub>2</sub> ) <sub>w</sub> -NH-(CH <sub>2</sub> ) <sub>x</sub> -NH-(CH <sub>2</sub> ) <sub>y</sub> -NH-(CH <sub>2</sub> ) <sub>z</sub> -
	(13) -[NH-(CH <sub>2</sub> ) <sub>w</sub> ] <sub>m</sub> - [NH-(CH <sub>2</sub> ) <sub>x</sub> ] <sub>n</sub> - [[CH <sub>3</sub> (CH <sub>2</sub> ) <sub>y</sub> N] - (CH <sub>2</sub> ) <sub>z</sub> -
	(14) -[NH-(CH <sub>2</sub> ) <sub>x</sub> ] <sub>n</sub> - [[CH <sub>3</sub> (CH <sub>2</sub> ) <sub>y</sub> N] - (CH <sub>2</sub> ) <sub>z</sub> -
	(15) -[NH-(CH <sub>2</sub> ) <sub>w</sub> ] <sub>m</sub> - [NH-(CH <sub>2</sub> ) <sub>x</sub> ] <sub>n</sub> - [[CH <sub>3</sub> (CH <sub>2</sub> ) <sub>y</sub> N] - (CH <sub>2</sub> ) <sub>z</sub> -
20	(16) - [[CH <sub>3</sub> (CH <sub>2</sub> ) <sub>x</sub> ][CH <sub>3</sub> (CH <sub>2</sub> ) <sub>y</sub> N] - (CH <sub>2</sub> ) <sub>z</sub> -
	(17) -NH-(CH <sub>2</sub> ) <sub>z</sub> - NH -
	(18) -NH-(CH <sub>2</sub> ) <sub>y</sub> - NH-(CH <sub>2</sub> ) <sub>z</sub> - NH -
	(19) -NH-(CH <sub>2</sub> ) <sub>y</sub> - CH=CH - (CH <sub>2</sub> ) <sub>z</sub> -
	(20) - [NH-(CH <sub>2</sub> ) <sub>w</sub> ] <sub>p</sub> - [[CH <sub>3</sub> (CH <sub>2</sub> ) <sub>x</sub> N] - (CH <sub>2</sub> ) <sub>y</sub> - [NH-(CH <sub>2</sub> ) <sub>z</sub> ] <sub>q</sub> -

For R<sup>3</sup> and/or R<sup>4</sup>


---

- (1) H-
- (2) CH<sub>3</sub>-
- (3) CH<sub>3</sub>-(CH<sub>2</sub>)<sub>2</sub>-
- 5 (4) CH<sub>3</sub>-(CH<sub>2</sub>)<sub>4</sub>-
- (5) CH<sub>3</sub>-(CH<sub>2</sub>)<sub>z</sub>-
- (6) CH<sub>3</sub>-[CH<sub>3</sub>-(CH<sub>2</sub>)<sub>z</sub>]CH -
- (7) CH<sub>3</sub>-[CH<sub>3</sub>-(CH<sub>2</sub>)<sub>2</sub>]CH -
- (8) CH<sub>3</sub>-[[CH<sub>3</sub>-(CH<sub>2</sub>)<sub>y</sub>][CH<sub>3</sub>-(CH<sub>2</sub>)<sub>z</sub>]C -
- 10 (9) CH<sub>3</sub>-(CH<sub>2</sub>)<sub>z</sub>-CH=CH-CH<sub>2</sub> -
- (10) CH<sub>3</sub>-[CH<sub>3</sub>-(CH<sub>2</sub>)<sub>y</sub>-CH=CH-(CH<sub>2</sub>)<sub>z</sub>]CH -
- (11) CH<sub>3</sub>-[[CH<sub>3</sub>-(CH<sub>2</sub>)<sub>w</sub>-CH=CH-(CH<sub>2</sub>)<sub>x</sub>][CH<sub>3</sub>-(CH<sub>2</sub>)<sub>y</sub>-CH=CH-(CH<sub>2</sub>)<sub>z</sub>]]CH -
- (12) CH<sub>3</sub>-[CH<sub>3</sub>-(CH<sub>2</sub>)<sub>y</sub>]CH-(CH<sub>2</sub>)<sub>z</sub> -

Group II Amphiphiles

Additionally there are provided cationic amphiphiles of Group II (see Figure 5) capable of facilitating transport of biologically active molecules into cells said amphiphiles having the structure (II),

5



wherein:

Z is a steroid;

10 X is a carbon atom or a nitrogen atom;

Y is a linking group or Y is absent;

R<sup>3</sup> is an amino acid, a derivatized amino acid, H or alkyl;

R<sup>1</sup> is --NH-- , an alkylamine, or a polyalkylamine;

R<sup>4</sup> is an amino acid, a derivatized amino acid, H or alkyl;

15 R<sup>2</sup> is --NH-- , an alkylamine, or a polyalkylamine;

and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup> , except that both R<sup>1</sup> and R<sup>2</sup> cannot be --NH--.

Representative amphiphiles provided according to Group II include amphiphiles 87, 91, 93, 95, 97, 99, 100, and 103. With respect to the structural  
 20 features of these amphiphiles, and the other amphiphiles of Group II, the following should be considered.

The steroid group may be selected according to the criteria defined above for the Group I amphiphiles. Accordingly, preferred amphiphiles include those selected from 3- sterols, wherein the sterol molecule is linked by the 3-O- group thereof, or by N in replacement thereof, to "Y".


5           The linking group Y of the Group II amphiphiles consists of an N-acylamino acid (or a derivative thereof), or consists of a group (such as  $>C=O$  or  $>C=S$ ) wherein no more than one atom of said group forms a bond with both "X" and "Z". Optionally, group Y may be absent. Representative N-acylamino groups include an N-Acyl serine ( No. 87), an N-Acyl glycine (No. 91), and an N-  
10 Acyl aspartic acid ( No. 103). With respect to the use of N-Acyl aspartic acid in amphiphile No. 103, it is noted that, as provided, the gamma carboxyl thereof is further derivatized to an additional alkylamine moiety.

The criteria for selection of  $R^1$  and  $R^2$  are as set forth for the Group I amphiphiles.  $R^3$  and  $R^4$  represent H or alkyl, or may be natural or artificial  
15 amino acids including derivatives of either. Representative examples of  $R^3$  or  $R^4$  amino acid groups include those derived from tryptophan ( No. 97) and from arginine ( No. 95).

Group III Amphiphiles

Additionally there are provided cationic amphiphiles of Group III (see Figure 6) capable of facilitating transport of biologically active molecules into cells said amphiphiles having the structure (III),

5



wherein:

Z is an alkylamine or a dialkylamine, linked by the N-atom thereof, to Y, or directly to X if Y is absent, wherein if Z is a dialkylamine, the alkyl groups thereof can be the same or different;

10 X is a carbon atom or a nitrogen atom;

Y is a short linking group, or Y is absent;

R<sup>3</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>1</sup> is --NH-- , an alkylamine, or a polyalkylamine;

15 R<sup>4</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>2</sup> is --NH-- , an alkylamine, or a polyalkylamine;

and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup>, except that both R<sup>1</sup> and R<sup>2</sup> cannot be --NH--.

Representative cationic amphiphiles according to the practice of the invention that contain an alkyl amine or dialkylamine as lipophilic group include, for example, N,N-dioctadecyllysineamide; N<sup>1</sup>, N<sup>1</sup>-dioctadecyl-1,2,6-triaminohexane; N,N-didodecyllysineamide; N,N- didecyllysineamide; spermidine- N,N- dioctadecyl urea; N-myristyllysineamide; and N-

(dioctyldecylaminoethyl)-lysineamide. Representative amphiphiles are depicted (Figure 6) as amphiphiles 43, 47, 56, 60, and 73. With respect to the structural features of these amphiphiles, and the other amphiphiles of Group III, the following should be considered.

- 5 With respect to the selection of the lipophilic alkylamine or dialkylamine group "Z", Table 2 below provides representative structures.

Table 2

For "Z"

- 
- (1)  $\text{CH}_3\text{-(CH}_2\text{)}_{13}\text{-NH-}$   
 10 (2)  $\text{CH}_3\text{-(CH}_2\text{)}_z\text{-NH-}$   
 (3)  $[[\text{CH}_3\text{(CH}_2\text{)}_{17}][\text{CH}_3\text{(CH}_2\text{)}_{17}]]\text{N-}$   
 (4)  $[[\text{CH}_3\text{(CH}_2\text{)}_{11}][\text{CH}_3\text{(CH}_2\text{)}_{11}]]\text{N-}$   
 (5)  $[[\text{CH}_3\text{(CH}_2\text{)}_9][\text{CH}_3\text{(CH}_2\text{)}_9]]\text{N-}$   
 (6)  $[[\text{CH}_3\text{(CH}_2\text{)}_x][\text{CH}_3\text{(CH}_2\text{)}_y]]\text{N-}$   
 15 (7)  $[[\text{CH}_3\text{(CH}_2\text{)}_x][\text{CH}_3\text{(CH}_2\text{)}_y\text{CH=CH(CH}_2\text{)}_z]]\text{N-}$   
 (8)  $[[\text{CH}_3\text{(CH}_2\text{)}_w][\text{CH}_3\text{(CH}_2\text{)}_x\text{CH=CH(CH}_2\text{)}_y\text{CH=CH(CH}_2\text{)}_z]]\text{N-}$


- In connection with the selection of suitable alkylamine or dialkylamine groups for inclusion at position Z in the amphiphiles of the invention, an alkyl chain(s) of the group should not be so large in molecular weight that it interferes  
 20 with the solubility of the amphiphile, or interferes with its ability to interact with plasmid DNA. Additionally, an alkyl chain of an alkylamine or dialkylamine may include one or more points of unsaturation.

- The selection of R groups  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ , and  $\text{R}^4$  follows that disclosed for the Group I amphiphiles, and these R groups may be selected, for example, from  
 25 Table I. Linking group Y may be selected as for the Group I amphiphiles, and preferred examples thereof include  $\text{—CH}_2\text{—}$ , and  $>\text{C=O}$ .

Group IV Amphiphiles

Additionally there are provided cationic amphiphiles of Group IV (see Figure 7) capable of facilitating transport of biologically active molecules into cells said amphiphiles having the structure (IV),

5



wherein:

A and B are independently O, N or S;

10  $R^5$  and  $R^6$  are independently alkyl or acyl groups and may be saturated or contain sites of unsaturation;

C is selected from the group consisting of  $-CH_2-$ ,  $>C=O$ , and  $>C=S$ ;

E (analogous to "X" in structures I, II, III) is a carbon atom or a nitrogen atom;

D is a linking group such as  $-NH(C=O)-$  or  $-O(C=O)-$ , or D is absent;

15  $R^3$  is H, or a saturated or unsaturated aliphatic group;

$R^1$  is  $--NH--$ , an alkylamine, or a polyalkylamine;

$R^4$  is H, or a saturated or unsaturated aliphatic group;

$R^2$  is  $--NH--$ , an alkylamine, or a polyalkylamine;

and wherein  $R^1$  is the same or is different from  $R^2$ , except that both  $R^1$  and  $R^2$

20 cannot be  $--NH--$ .

Representative amphiphiles of Group IV include Nos. 64, 76, 85, 89, 94, 98, 102, 105, 110, and 111. With respect to the structural features of these

amphiphiles, and the other amphiphiles of Group IV, the following should be considered.

With respect to the selection of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup>, the teachings provided for Group I, II, and III amphiphiles are applicable. As aforementioned, group  
5 "E" represents a carbon atom or a nitrogen atom.

R<sup>5</sup> and R<sup>6</sup> are independently alkyl or acyl groups, preferably containing about 8 to about 30 carbon atoms, and such groups may contain one or more points of unsaturation.

With respect to the selection of Group D, linkers such as -NH(C=O)- or  
10 -O(C=O)- are preferred, and are depicted such that the left side thereof is intended to bond to "C" and the right side thereof is intended to bond to "E".  
Optionally, group D may be absent (amphiphile No.94). Additional linkers may be selected based on the teachings provided with respect to Groups I, II, and III above, and based upon the in vivo test data derived ( Figure 15), it is preferred  
15 that the linker D be short or absent.

#### Co-lipids

Representative co-lipids that are useful according to the practice of the invention for mixing with one or more cationic amphiphiles include  
dioleoylphosphatidylethanolamine ("DOPE"), diphytanoylphosphatidyl-  
20 ethanolamine, lyso-phosphatidylethanolamines other phosphatidyl-  
ethanolamines, phosphatidylcholines, lyso-phosphatidylcholines and cholesterol.  
Typically, a preferred molar ratio of cationic amphiphile to colipid is about 1:1.  
However, it is within the practice of the invention to vary this ratio (see Example  
3 below), including also over a considerable range.

25 It is generally believed in the art that preparing cationic amphiphiles as complexes with co-lipids (particularly neutral co-lipids) enhances the capability of the amphiphile to facilitate transfections. Although colipid-enhanced

performance has been observed for numerous of the amphiphiles of the invention, the amphiphiles of the invention are active as transfectants without co-lipid. Accordingly, the practice of the present invention is neither to be considered limited by theories as to co-lipid participation in intracellular delivery mechanisms, nor to require the involvement of co-lipids.

#### Transacylation Reactions

Although heretofore unrecognized in the art, it has been determined also that certain co-lipids may react chemically with certain types of cationic amphiphiles under conditions of co-storage, there resulting new molecular species. Generation of such new species is believed to occur via mechanisms such as transacylation. In this regard, see Figure 4 which depicts a transacylation reaction involving spermine cholesterol carbamate(No.67) and DOPE, there resulting lyso PE species and multiple forms of particular acyl- cationic amphiphile ( designated No. 80).

With respect to such reactions, the following remarks are of interest. With respect to use of amphiphile No.67, it has been observed that a mixture of amphiphile and DOPE, in chloroform solvent, does not appear to participate in such reactions. However, preparing the amphiphile and co-lipid in an aqueous solution where bilayer-containing structures such as liposomes can form will permit transacylation. Additionally, if amphiphile and co-lipid are dried down to a thin film, such as from chloroform (thereby placing the 2 species in intimate contact), then transacylation also occurs, possibly as a result of entropic effects. It is expected that these phenomena would also apply to lyophilized amphiphile/DOPE preparations.

Accordingly, it is highly preferred to maintain such amphiphile /DOPE preparations at very cold temperatures, such as -70 degrees C. Preparation of

amphiphile No. 67 as a mono, di, or tri acetate salt has also been determined to slow transacylations.

It is to be understood that therapeutically-effective pharmaceutical compositions of the present invention may or may not contain such  
5 transacylation byproducts, or other byproducts, and that the presence of such byproducts does not prevent the therapeutic use of the compositions containing them. Rather use of such compositions is within the practice of the invention, and such compositions and the novel molecular species thereof are therefore specifically claimed.

#### 10 Preparation of Pharmaceutical Compositions and Administration Thereof

The present invention provides for pharmaceutical compositions that facilitate intracellular delivery of therapeutically effective amounts of biologically active molecules. Pharmaceutical compositions of the invention facilitate entry of biologically active molecules into tissues and organs such as the gastric mucosa,  
15 heart, lung, and solid tumors. Additionally, compositions of the invention facilitate entry of biologically active molecules into cells that are maintained *in vitro*, such as in tissue culture. The amphiphilic nature of the compounds of the invention enables them to associate with the lipids of cell membranes, other cell surface molecules, and tissue surfaces, and to fuse or to attach thereto. One type  
20 of structure that can be formed by amphiphiles is the liposome, a vesicle formed into a more or less spherical bilayer, that is stable in biological fluids and can entrap biological molecules targeted for intracellular delivery. By fusing with cell membranes, such liposomal compositions permit biologically active molecules carried therewith to gain access to the interior of a cell through one or  
25 more cell processes including endocytosis and pinocytosis. However, unlike the case for many classes of amphiphiles or other lipid-like molecules that have been proposed for use in therapeutic compositions, the cationic amphiphiles of the

invention need not form highly organized vesicles in order to be effective, and in fact can assume (with the biologically active molecules to which they bind) a wide variety of loosely organized structures. Any of such structures can be present in pharmaceutical preparations of the invention and can contribute to  
5 the effectiveness thereof.

Biologically active molecules that can be provided intracellularly in therapeutic amounts using the amphiphiles of the invention include:

- (a) polynucleotides such as genomic DNA, cDNA, and mRNA that encode for therapeutically useful proteins as are known in the art,
- 10 (b) ribosomal RNA;
- (c) antisense polynucleotides, whether RNA or DNA, that are useful to inactivate transcription products of genes and which are useful, for example, as therapies to regulate the growth of malignant cells; and
- (d) ribozymes.

15 In general, and owing to the potential for leakage of contents therefrom, vesicles or other structures formed from numerous of the cationic amphiphiles are not preferred by those skilled in the art in order to deliver low molecular weight biologically active molecules. Although not a preferred embodiment of the present invention, it is nonetheless within the practice of the invention to  
20 deliver such low molecular weight molecules intracellularly. Representative of the types of low molecular weight biologically active molecules that can be delivered include hormones and antibiotics.

Cationic amphiphile species of the invention may be blended so that two or more species thereof are used, in combination, to facilitate entry of  
25 biologically active molecules into target cells and/or into subcellular compartments thereof. Cationic amphiphiles of the invention can also be blended for such use with amphiphiles that are known in the art.

Dosages of the pharmaceutical compositions of the invention will vary, depending on factors such as half-life of the biologically-active molecule, potency of the biologically-active molecule, half-life of the amphiphile(s), any potential adverse effects of the amphiphile(s) or of degradation products thereof, the route  
5 of administration, the condition of the patient, and the like. Such factors are capable of determination by those skilled in the art.

A variety of methods of administration may be used to provide highly accurate dosages of the pharmaceutical compositions of the invention. Such preparations can be administered orally, parenterally, topically, transmucosally,  
10 or by injection of a preparation into a body cavity of the patient, or by using a sustained-release formulation containing a biodegradable material, or by onsite delivery using additional micelles, gels and liposomes. Nebulizing devices, powder inhalers, and aerosolized solutions are representative of methods that may be used to administer such preparations to the respiratory tract.

15 Additionally, the therapeutic compositions of the invention can in general be formulated with excipients (such as the carbohydrates lactose, trehalose, sucrose, mannitol, maltose or galactose) and may also be lyophilized (and then rehydrated) in the presence of such excipients prior to use. Conditions of optimized formulation for each amphiphile of the invention are capable of  
20 determination by those skilled in the pharmaceutical art. By way of example, for spermidine cholesterol carbamate (amphiphile No. 53), it has been determined that use of sucrose is preferred over mannitol in order to prevent formation of amphiphile/DNA aggregates, particularly as the concentration of DNA is increased therein. Addition of such excipients maintains the consistency of  
25 lyophilized pharmaceutical compositions during storage, and prevent difficulties such as aggregation, or insolubility, that may likely occur upon rehydration from the lyophilized state.

Accordingly, a principal aspect of the invention involves providing a composition that comprises a biologically active molecule (for example, a polynucleotide) and one or more cationic amphiphiles (including optionally one or more co-lipids), and then maintaining said composition in the presence of one  
5 ore more excipients as aforementioned, said resultant composition being in liquid or solid (preferably lyophilized) form, so that: (1) the therapeutic activity of the biologically active molecules is substantially preserved; (2) the transfection-enhancing nature of the amphiphile( or of amphiphile/ DNA complex) is maintained. Without being limited as to theory, it is believed that  
10 the excipients stabilize the interaction (complexes)of the amphiphile and biologically active molecule through one or more effects including:

- (1) minimizing interactions with container surfaces,
- (2) preventing irreversible aggregation of the complexes, and
- (3) maintaining amphiphile/DNA complexes in a chemically-stable state, i.e.,  
15 preventing oxidation and/or hydrolysis.

Although the presence of excipients in the pharmaceutical compositions of the invention stabilizes the compositions and facilitates storage and manipulation thereof, it has also been determined that moderate concentrations of numerous excipients may interfere with the transfection-enhancing capability of  
20 pharmaceutical formulations containing them. In this regard, an additional and valuable characteristic of the amphiphiles of the invention is that any such potentially adverse effect can be minimized owing to the greatly enhanced in vivo activity of the amphiphiles of the invention in comparison with amphiphilic compounds known in the art. Without being limited as to theory, it  
25 is believed that osmotic stress ( at low total solute concentration) may contribute positively to the successful transfection of polynucleotides into cells in vivo . Such a stress may occur when the pharmaceutical composition, provided in

unbuffered water, contacts the target cells. Use of such otherwise preferred compositions may therefore be incompatible with treating target tissues that already are stressed, such as has damaged lung tissue of a cystic fibrosis patient. Accordingly, and using sucrose as an example, selection of concentrations of this  
5 excipient that range from about 15 mM to about 200 mM provide a compromise between the goals of (1) stabilizing the pharmaceutical composition to storage and (2) mimizing any effects that high concentrations of solutes in the composition may have on transfection performance.

Selection of optimum concentrations of particular excipients for particular  
10 formulations is subject to experimentation, but can be determined by those skilled in the art for each such formulation.

An additional aspect of the invention concerns the protonation state of the cationic amphiphiles of the invention prior to their contacting plasmid DNA in order to form a therapeutic composition. It is within the practice of the invention  
15 to utilize fully protonated, partially protonated, or free base forms of the amphiphiles in order to form such therapeutic compositions. With respect to amphiphile No. 67 (spermine cholesterol carbamate), it has been observed that when providing this amphiphile for a transfecting composition with DOPE (itself provided as a zwitterion), transgene expression was best for the free base, but  
20 decreased if the amphiphile was prepared as an acetate salt. Activity decreased step-wise through the mono and di acetate salts and was minimal for the tri-acetate salt. Under the circumstances described, the plasmid DNA provided for contacting with the amphiphile was prepared (without buffer) as a sodium salt in water.

#### 25 Transfection of the Vascular System

A further aspect of the invention involves transfection of the vascular system. By transfection of the vascular system is meant that the

therapeutic composition (comprising one or more cationic amphiphiles, a therapeutic polynucleotide, and optionally, one or more co-lipids) is placed in a blood vessel of a patient through which it will be distributed to blood vessel cells. Blood vessels suitable for application in the practice of the invention include  
5 those of the arterial, venous, or capillary systems. Blood vessel cells that may be transfected according to the practice of the invention also include those of the arterial, venous, or capillary systems.

It is also within the practice of the invention to transfect cells of identifiable vessels of the lymphatic system.

10 The cationic amphiphiles of the invention (including those defined herein by Groups I, II, III and IV) can be formulated with co-lipids and polynucleotides for such therapeutic application.

The vascular system of a patient is contacted with a composition that comprises a cationic amphiphile and a polynucleotide that encodes a protein  
15 having therapeutic properties, such that cells of the vascular system are transfected thereby, and express said protein from said polynucleotide. In a preferred aspect, the protein is one normally secreted from cells, and the encoding polynucleotide includes, for example, sequences for pre- or pro-peptides, or for amino acids that are to be glycosylated, such that the encoded  
20 protein is secreted into the vascular circulation of a patient, after which the secreted protein provides therapeutic benefit at a site remote or adjacent to the transfected blood vessel cells. Examples of therapeutic proteins that can be expressed in patients according to this aspect of the invention include adenosine deaminase, glucocereb-  
25 idase, and further include numerous of the protein hormones such as growth hormones, insulin and the like. Efficient expression and secretion of such a protein is demonstrated in Example 11.

Methods of Syntheses

The following methods illustrate production of certain of the cationic amphiphiles of the invention. Those skilled in the art will recognize other methods to produce these compounds, and to produce also the other compounds  
5 of the invention.

Group I amphiphiles(A) N<sup>4</sup>-Spermidine cholesteryl carbamate

Spermidine cholesterol carbamate (Figure 1, No. 53) was synthesized according to the following procedure which is outlined in Figure 8.

10 Synthesis of N<sup>1</sup>, N<sup>8</sup>-DiCBZ -N<sup>4</sup>-Spermidine Cholesterol Carbamate

N<sup>1</sup>, N<sup>8</sup> - dicarbobenzoxy spermidine (61% yield, m.p. 104 - 105° C) was prepared according to the procedure of S. K. Sharma, M. J. Miller, and S. M. Payne, J. Med. Chem., 1989, 32, 357-367. The N<sup>1</sup>, N<sup>8</sup>-  
dicarbobenzoxy spermidine (25 g, 60.5 mmol) and triethylamine (25 ml, 178  
15 mmol) were dissolved in 625 ml of anhydrous methylene chloride, cooled to 0 - 4°C and stirred under N<sub>2</sub>. Cholesteryl chloroformate (27.2 g, 60.6 mmol) was dissolved in 250 ml of methylene chloride and added to the reaction over a 20 minute period. A white precipitate formed upon addition. After the addition was complete, the reaction was stirred at 0 - 4°C for 10 minutes and then at room  
20 temperature for 1.5 hr. At this point, the white precipitate completely dissolved. The reaction was followed by TLC with hexane / ethyl acetate 6 / 4 as eluent (product R<sub>f</sub> = 0.25). To this reaction mixture was added 625 ml of methylene chloride and 625 ml of water. The layers were then allowed to separate. The organic layer was dried over MgSO<sub>4</sub> and filtered. The filtrate was concentrated  
25 *in vacuo* to give an oil. Vacuum drying was then carried out overnight. This crude product had a glue-like consistency. The crude product was purified by column chromatography (2 kg silica gel, eluent - hexane / ethyl acetate 6 / 4) to

give 46.8 g of the 3-β-[N<sup>4</sup>-(N<sup>1</sup>,N<sup>8</sup>-dicarbobenzoxyspermidine)carbamoyl] cholesterol (also described herein as N<sup>1</sup>,N<sup>8</sup>-diCBZ-N<sup>4</sup>-spermidine cholesterol carbamate) in 93% yield.

#### Final Synthesis of Spermidine Cholesterol Carbamate

5 To 6.0 grams of 10% palladium on activated carbon under N<sub>2</sub> was added a solution of 30 grams of 3-β-[N<sup>4</sup>-(N<sup>1</sup>,N<sup>8</sup>-dicarbobenzoxyspermidine)carbamoyl] cholesterol in 1 liter of ethanol, see Figure 13. The reaction mixture was purged with N<sub>2</sub> and stirred under H<sub>2</sub> (atmospheric pressure) for 18 hr. The mixture was again purged with N<sub>2</sub> and filtered through a 10 g bed of celite. The filter cake  
10 was washed with 2 liters of 10% triethylamine in ethanol and the combined filtrates were concentrated *in vacuo* to a gel. The product was then dried under vacuum overnight to a sticky solid. This crude product was purified by column chromatography (2 kg of silica gel, eluent - 4 L of chloroform / methanol 95 / 5 followed by 30 L of chloroform / methanol / iso-propylamine 95 / 5 / 5, R<sub>f</sub> =  
15 0.24) to obtain 13.1 g of the desired spermidine cholesterol carbamate in 64% yield. HPLC (C-18 reversed phase column, linear gradient elution profile - methanol / iso-propanol / water / trifluoroacetic acid 60 / 20 / 20 / 0.1 to methanol / iso-propanol / trifluoroacetic acid 70 / 30 / 0.1 to methanol / iso-propanol / chloroform / trifluoroacetic acid 60 / 20 / 20 / 0.1) analysis of this  
20 material showed it to be 99.2% pure with the 7-dehydrocholesterol analog present at a level of 0.8%.

In connection with this example and those that follow, it is noted that all TLC plates were visualized with phosphomolybdic acid.

#### (B) N<sup>4</sup>-Spermine cholesteryl carbamate

25 Spermine cholesterol carbamate (Figure 1, No. 67) was prepared according to the following procedure which is outlined in Figure 9.

N<sup>1</sup>,N<sup>12</sup>-diCBZ-spermine

Benzylchloroformate (1.76g, 1.5 ml, 10.36 mmol) was dissolved in methylene chloride (5 ml) and placed in a three neck flask under a nitrogen atmosphere. Imidazole (1.4 g, 20.6 mmol) was dissolved in methylene chloride (20 ml) and placed in an addition funnel. The three neck flask was cooled to 0°C  
5 and the imidazole solution was added gradually over 20 min. The mixture was stirred at room temperature for 1 hour and then methylene chloride (25 mL) and citric acid (10%, 25 ml) were added. The layers were separated and the organic fraction was washed with citric acid (10%, 25 ml). The organic component was dried over magnesium sulfate and concentrated in vacuo. The residue was dried  
10 under high vacuum for 1 hour at ambient temperature.

To the residue was added dimethylaminopyridine (35 mg), methylene chloride (25 ml) and the mixture was cooled to 0°C, under a nitrogen atmosphere. To an addition funnel was added a solution of spermine (1g, 4.94 mmol) in methylene chloride (25 ml). The spermine solution was added  
15 gradually over 15 min. The reaction mixture was stirred overnight at ambient temperature and then concentrated in vacuo. The residue was dissolved in ethyl acetate (80 ml) and washed three times with water (15 ml). The organics were dried over magnesium sulfate, filtered and concentrated in vacuo to give a crude white solid. The material was purified by flash chromatography (65g silica gel,  
20 100:100:10 CHCl<sub>3</sub>: MeOH: NH<sub>4</sub>OH, product Rf.=0.33), to give after drying under high vacuum 1.01g (2.146 mmol, 43 % yield) of product.

N<sup>1</sup>N<sup>12</sup>-diCBZ-N<sup>4</sup>-spermine cholesteryl carbamate

Cholesteryl chloroformate (964 mg, 2.15 mmol) was dissolved in chloroform (10 ml) and added dropwise to a cooled (0°C) solution of N<sup>1</sup>,N<sup>12</sup>-  
25 diCBZ spermine (1.01g, 2.15 mmol), triethylamine (1 ml) in chloroform (10 ml). The reaction was allowed to warm to room temperature and stirred for 2 hours. To the reaction solution was added water (25 ml) and chloroform (25 ml). The

layers were separated and the organic fraction dried over magnesium sulfate.

The solution was concentrated in vacuo to give a crude material that was purified by flash chromatography (68g silica gel, MeOH / CHCl<sub>3</sub> 1/4, product Rf. =0.36) to give 1.23 g (1.39 mmol, 65% yield) of product.

5 final synthesis of N<sup>4</sup>-Spermine Cholesteryl Carbamate

N<sup>1</sup>N<sup>12</sup>-diCBZ-N<sup>4</sup>-spermine cholesteryl carbamate (262 mg, 0.300 mmol) was dissolved in 5 ml of acetic acid and 45 mg of 10% Pd on C was added. The solution was purged with nitrogen and stirred under hydrogen at atmospheric pressure. The hydrogenolysis was allowed to proceed for 7 hours. The reaction mixture was filtered and the catalyst was washed with 40 ml of ethyl acetate / acetic acid 9 / 1 and the filtrate will be concentrated *in vacuo* to give a residue. The crude product was dissolved in 35 mL of 1N NaOH and extracted three times with 40 ml of chloroform / methanol 9 / 1. The combined organic fractions were washed with 20 mL of water and dried over Na<sub>2</sub>SO<sub>4</sub>. The solution was filtered, concentrated *in vacuo* and dried under vacuum to give 125 mg of the desired product in 67% yield.

In connection with the above procedure, it is noted that the hydrogenolysis should be carried out under acidic conditions, in order to minimize the poisoning of the catalyst.

20 Urea analogs - such as spermine or spermidine cholestamine urea - can be prepared by a sequence of reactions well known to those versed in the art of organic synthesis. For example an amine can be treated with an equal molar amount of carbonyldiimidazole followed by the addition of a second amine to give the desired urea.

25 (C) N,N Bis (3-aminopropyl)-O-cholesteryl -3-carbamate

N,N Bis (3-aminopropyl)-O-cholesteryl-3-carbamate (Figure 1, No. 69) was prepared according to the following procedure.

Bis (3-CBZ aminopropyl) amine was prepared using the method described above for N<sup>1</sup>,N<sup>12</sup> -diCBZ-spermine, except that N-(3-aminopropyl)1,3-propanediamine was substituted for spermine as reactant. The pure product was isolated in 34 % yield by silica gel flash chromatography using as solvent CHCl<sub>3</sub>/ MeOH/ NH<sub>4</sub>OH 80/20/0.5.

The Bis (3-CBZ aminopropyl) amine so prepared was then reacted with cholesteryl chloroformate according to the method described above for the synthesis of N<sup>1</sup>, N<sup>8</sup>-DiCBZ -N<sup>4</sup>-spermidine cholesteryl carbamate. The pure product (N,N Bis ( 3-CBZ aminopropyl)-O-cholesteryl-3-carbamate) was obtained in 73% yield.

Synthesis of N,N Bis(3-aminopropyl)-O-cholesteryl-3-carbamate was completed by hydrogenolysis of the CBZ groups from N,N Bis(3-CBZ aminopropyl)-O-cholesteryl-3-carbamate following the procedure described above in relation to the synthesis of N<sup>4</sup>-spermidine cholesteryl carbamate. The product was obtained in 23% yield without silica gel chromatography purification.

(D) N,N Bis (6-aminohexyl)-O-cholesteryl -3- carbamate.

N,N Bis (6-aminohexyl)-O-cholesteryl-3-carbamate (Figure 1, No. 70) was prepared according to the following procedure.

First, Bis (6-CBZ aminoethyl) amine was prepared using the method described above for N<sup>1</sup>,N<sup>12</sup> -diCBZ-spermine, except that Bis(hexamethylene)triamine was substituted for spermine as reactant. Pure product was isolated in 24% yield by recrystallization from toluene.

Bis (6-CBZ aminoethyl) amine was then reacted with cholesteryl chloroformate according to the method described above for the synthesis of N<sup>1</sup>, N<sup>8</sup>-DiCBZ -N<sup>4</sup>-spermidine cholesteryl carbamate. Product N,N Bis(6-CBZ

aminohexyl)-O-cholesteryl-3-carbamate was isolated in 40% yield by silica gel flash chromatography using hexanes/ethyl acetate 7/3 .

(E) Lysine 3-N- dihydrocholesteryl carbamate

Lysine 3-N- dihydrocholesteryl carbamate (Figure 1, panel C) was  
5 prepared according to the following procedure.

To a solution of dihydrocholesterol (5.0 g, 12.9 mmol, Aldrich),  
phthalimide (2.0 g, 13.6 mmol, Aldrich), and triphenylphosphine (3.8 g, 13.6  
mmol, Aldrich) in THF (20 ml, Aldrich) stirred at 0° C under a nitrogen  
atmosphere was added dropwise diethylazodicarboxylate (2.3 ml, 14.5 mmol,  
10 Aldrich). Upon the completion of addition the reaction mixture was allowed to  
warm to ambient temperature and stirred overnight. The reaction mixture was  
concentrated *in vacuo* to a residue. This residue was dissolved in 50 ml hexane /  
ethyl acetate 95 / 5 and a precipitate formed. The mixture was filtered. The  
filtrate was concentrated to dryness *in vacuo*, dissolved in 25 ml of hexane / ethyl  
15 acetate 95 / 5 and chromatographed on 200 g silica gel (eluent 2 L hexane / ethyl  
acetate 95 / 5 then 1 L hexane / ethyl acetate 90 / 10). A 76% yield of the desired  
3-phthalimidocholestane (5.43 g) was obtained.

The 3-phthalimidocholestane (5.40 g, 9.75 mmol) was dissolved in 60 mL  
of methanol and anhydrous hydrazine (3.1 ml, 99 mmol) was added. The  
20 reaction mixture was stirred and heated at reflux under a nitrogen atmosphere  
for 4 hr. This mixture was then cooled to room temperature, 3.1 mL of  
concentrated HCl was added and the resulting mixture was heated at reflux  
overnight. Upon cooling to ambient temperature, 100 ml of diethyl ether and 50  
ml of 1 N NaOH were added (final pH of 10.1) and the layers were separated.  
25 The aqueous layer was extracted with 50 ml of diethyl ether and the combined  
organic fractions were filtered. The filtrate was concentrated *in vacuo* and the

residue was purified by silica gel chromatography (chloroform / methanol 90 / 10) to give 2.24 g of 3-aminocholestane in 59 % yield.

L-N $\alpha$ ,N $\epsilon$ -diBOClysine N-hydroxysuccinimide ester (286 mg, 0.644 mmol, Sigma) and 3-aminocholestane (250 mg, 0.644 mmol) were dissolved in 5 mL of methylene chloride, 0.1 mL of triethylamine was added and the resulting solution was stirred under a nitrogen atmosphere at ambient temperature overnight. To the reaction mixture was added 10 mL of water and 25 mL of methylene chloride and the layers were separated. The aqueous layer was extracted with 25 mL of methylene chloride and the combined organic fractions were dried over MgSO<sub>4</sub> and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by chromatography on 25 g of silica gel (eluent - hexane / ethyl acetate 6 / 4, sample applied in hexane / ethyl acetate 9 / 1). The purified material was dissolved in 25 mL of chloroform and HCl gas was bubbled through the solution for 2 hr. followed by nitrogen for 10 min. The solution was concentrated *in vacuo* to give 299 mg of the desired product in 79% yield as the dihydrochloride salt.

(F) N<sup>1</sup>N<sup>8</sup>-Bis(3-aminopropyl)-N<sup>4</sup>-spermidine cholesteryl carbamate

N<sup>1</sup>N<sup>8</sup>-Bis(3-aminopropyl)-N<sup>4</sup>-spermidine cholesteryl carbamate (Figure 1, No. 75) was prepared according to the following procedure.

N<sup>4</sup>-Spermidine cholesteryl carbamate (1.14g, 2.04 mmol) was dissolved in MeOH (5 mL). Freshly distilled acrylonitrile (0.28 mL, 4.29 mmol) was added and the solution was stirred at room temperature for 18 h. The solvent was concentrated *in vacuo* to give an oil. Vacuum drying was then carried out overnight. The crude product was purified by column chromatography (125 g silica gel, eluent - CHCl<sub>3</sub> MeOH 1/9) to give 1.15 g (85 %) of the N<sup>1</sup>N<sup>8</sup>-Bis (cyanoethyl) N<sup>4</sup>-Spermidine cholesteryl carbamate.

Raney Nickel 50% slurry (1.2 g, Aldrich) was placed in a Parr Bomb with 1M NaOH in 95% EtOH (50 mL). The N<sup>1</sup>,N<sup>8</sup>-Bis (cyanoethyl) N<sup>4</sup>-Spermidine cholesteryl carbamate. was dissolved in EtOH (35 mL) and added to the bomb. The vesicle was evacuated and placed under Argon pressure (80-100 psi), three  
5 times and then evacuated and placed under Hydrogen pressure (100 psi), three times. The reaction was stirred under hydrogen pressure (100 psi) at room temperature for 72h. The vesicle was evacuated and placed under argon pressure. The catalyst was removed by filtration. The filtrate was concentrated *in vacuo*. The resulting oil was dissolved in 2:1 CH<sub>2</sub>Cl<sub>2</sub>: MeOH (100 mL) and  
10 washed with H<sub>2</sub>O (35 and 25 mL). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by chromatography on 100 g of silica gel (eluent - CHCl<sub>3</sub>/MeOH/conc. NH<sub>4</sub>OH 40/25/10, sample applied in CHCl<sub>3</sub>/MeOH 40/25). The purified material was concentrated *in vacuo* with iPrOH (3 X 50 mL) and CH<sub>2</sub>Cl<sub>2</sub>(3X50 mL) and then  
15 vacuum dried to give 986 mg (85%) of N<sup>1</sup>,N<sup>8</sup>-Bis(3-aminopropyl)-N<sup>4</sup>-spermidine cholesteryl carbamate.

(G) N(N<sup>4</sup>-3-aminopropyl-spermidine) cholesteryl carbamate

N(N<sup>4</sup>-3-aminopropyl-spermidine) cholesteryl carbamate (Figure 1, No. 78) was prepared as follows:

20 N<sup>1</sup>,N<sup>8</sup>-dicarbobenzoxyspermidine (1.0 g, 2.4 mmol) was dissolved in MeOH (10 mL). Freshly distilled acrylonitrile (0.3 mL, 4.5 mmol) was added and the reaction was stirred at room temperature for 18 h. The solvent was concentrated *in vacuo* to give an oil. The crude product was purified by column chromatography (100 g silica gel, eluent - CHCl<sub>3</sub>/MeOH 1/19) to give 1.10 g (97  
25 %) of N<sup>4</sup>-2-Cyanoethyl-N<sup>1</sup>, N<sup>8</sup>- dicarbobenzoxyspermidine.

The N<sup>4</sup>-2-Cyanoethyl-N<sup>1</sup>, N<sup>8</sup>-dicarbobenzoxyspermidine (0.5 g, 1.07 mmol) was dissolved in MeOH (5 mL) and CoCl<sub>2</sub> (280 mg, 2.15 mmol, Aldrich) was

added. The blue solution was cooled in an ice bath and NaBH<sub>4</sub> (405 mg, 10.7 mmol, Aldrich) was added in portions over 15 min. The resulting black solution was stirred at room temperature for 1 h. The black solution turned blue over this period. To the reaction was added CH<sub>2</sub>Cl<sub>2</sub>/MeOH 2/1 (30 mL). A black ppt  
5 formed. To this was added H<sub>2</sub>O (20mL) and the mixture was filtered. The resulting layers were separated and the organic layer dried with MgSO<sub>4</sub>. The drying agent was filtered and the filtrate concentrated *in vacuo* to give an oil. The crude product was purified by column chromatography (50 g silica gel, eluent - CHCl<sub>3</sub>/MeOH/conc NH<sub>4</sub>OH 100/100/5) to give 309 mg (62 %) of the  
10 N<sup>4</sup>-3-aminopropyl-N<sup>1</sup>, N<sup>8</sup> - dicarbobenzoxy spermidine.

To the N<sup>4</sup>-3-aminopropyl-N<sup>1</sup>, N<sup>8</sup> - dicarbobenzoxy spermidine (300 mg, 0.66 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> was added Et<sub>3</sub>N under N<sub>2</sub>. Cholesteryl chloroformate (326 mg, 0.726 mmol, Aldrich) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and added to the reaction dropwise. The mixture was stirred for 2h at room temperature. After  
15 adding CH<sub>2</sub>Cl<sub>2</sub> (25 mL) and H<sub>2</sub>O (10 mL), the layers were separated. The organic layer was dried with MgSO<sub>4</sub> and filtered. The filtrate was concentrated *in vacuo* to give 640 mg of crude product. The residue was purified by chromatography on 80 g of silica gel (eluent - CHCl<sub>3</sub> / MeOH 90 /10, sample applied in CHCl<sub>3</sub> / MeOH 90/10). The purified material was concentrated *in*  
20 *vacuo* and then vacuum dried to give 329 mg (57%) of N-(N<sup>4</sup>-3-aminopropyl-N<sup>1</sup>, N<sup>8</sup> - dicarbobenzoxy spermidine) cholesteryl carbamate.

To 10% Pd on carbon (65 mg, Aldrich) was added a solution of N-(N<sup>4</sup>-3-aminopropyl-N<sup>1</sup>, N<sup>8</sup> - dicarbobenzoxy spermidine) cholesteryl carbamate (300 mg) in acetic acid (25 mL). The reaction was placed under H<sub>2</sub> and stirred at room  
25 temperature overnight. After being placed under N<sub>2</sub>, the reaction was filtered. The catalyst was washed with 10 % acetic acid in EtOAc (50 mL). The filtrate was concentrated *in vacuo* to give an oil. The oil was dissolved in 2/1

CH<sub>2</sub>Cl<sub>2</sub>/MeOH (35 mL) and washed with 1 M NaOH (15 mL). The organic layer was dried with MgSO<sub>4</sub> and filtered. The filtrate was concentrated *in vacuo* and vacuum dried to give 196 mg (93%) of N-(N<sup>4</sup>-3-aminopropylspermidine) cholesteryl carbamate.

5        (H) N-[N<sup>1</sup>,N<sup>4</sup>,N<sup>8</sup>-Tris(3-aminopropyl)spermidine] cholesteryl carbamate  
N-[N<sup>1</sup>,N<sup>4</sup>,N<sup>8</sup>-Tris(3-aminopropyl)spermidine] cholesteryl carbamate  
(Figure 1, No. 96) was prepared by reacting N-(N<sup>4</sup>-3-aminopropylspermidine)  
cholesteryl carbamate with acrylonitrile (90% yield) and subsequent reduction of  
the di adduct with Raney nickel (75 % yield) as described for the preparation of  
10    N<sup>1</sup>,N<sup>8</sup>Bis(3-aminopropyl)-N<sup>4</sup>-spermidine cholesteryl carbamate.

(I) N,N-Bis(4-aminobutyl) cholesteryl carbamate

N,N-Bis(4-aminobutyl) cholesteryl carbamate (Figure 1, No. 82) was prepared as follows.

To a mixture of Benzylamine (2.0 g, 18.6 mmol, Aldrich), Na<sub>2</sub>CO<sub>3</sub> (4.4g, 42  
15    mmol) and KI (1.4 g, 9.5 mmol) in n-butanol (50 mL) was added 4-  
Chlorobutyronitrile (4.0 mL, 95 mmol) under nitrogen. The reaction was stirred  
at reflux of 48 h under nitrogen. After cooling to room temperature, diethyl ether  
(50 mL) was added and the precipitate filtered off. The filtrate was concentrated  
*in vacuo* to an oil. Toluene (100 mL) was added and the solution was  
20    concentrated *in vacuo*. Chloroform (100 mL) was added and again the solution  
was concentrated *in vacuo* and then vacuum dried for 18 h. The resulting oil was  
dissolved in Chloroform (100 mL) filtered and concentrated *in vacuo*. The crude  
product was purified by column chromatography (250 g silica gel, eluent -  
hexanes/EtOAc 60/40) to give 3.75g (97 %) of N,N-Bis(3-cyanopropyl)  
25    benzylamine.

The N,N-Bis(3-cyanopropyl) benzylamine (3.7 g, 17.8 mmol) was dissolved in EtOH (150 mL) and Acetic acid (4 mL) was added. This solution was added to

10% Pd on carbon (400 mg) under N<sub>2</sub>. The mixture was placed under H<sub>2</sub> and the reaction stirred for 18 h at room temperature. The reaction was placed under N<sub>2</sub>. The catalyst was filtered off and washed with EtOH (150 mL). The filtrate was concentrated *in vacuo*, chloroform (50 mL) was added and again concentrated *in vacuo*. The resulting oil was vacuum dried for 0.5 h and used directly in the next reaction. To this oil dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) was added Et<sub>3</sub>N (5 mL, 35 mmol) under N<sub>2</sub> and the solution cooled in an ice bath. Cholesteryl chloroformate (6.2 g, 13.87 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and this solution was added to the reaction dropwise over 10 min. The cooling bath was removed and the reaction stirred at room temperature for 18 h under N<sub>2</sub>. CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and H<sub>2</sub>O (100 mL) was added and the resulting layers were separated. The organic layer was dried with MgSO<sub>4</sub> and filtered. The filtrate was concentrated *in vacuo* and vacuum dried for 1 h. The crude product was purified by column chromatography (600 g silica gel, eluent - hexanes/EtOAc 60/40) to give 1.05g (10 %) of N,N-Bis (3-cyanopropyl) cholesteryl carbamate.

Raney Nickel 50% slurry (1.2 g) was placed in a Parr Bomb with 1M NaOH in 95% EtOH (50 mL). The N,N-Bis (3-cyanopropyl) cholesteryl carbamate (1.0 g, 1.77 mmol) was dissolved in EtOH (100 mL) and added to the bomb. The vesicle was evacuated and placed under Argon pressure (80-100 psi), three times and then evacuated and placed under Hydrogen pressure (100 psi), three times. The reaction was stirred under hydrogen pressure (100 psi) at room temperature for four days. The vesicle was evacuated and placed under argon pressure. The catalyst was removed by filtration. The filtrate was concentrated *in vacuo*. The resulting oil was dissolved in 2:1 CH<sub>2</sub>Cl<sub>2</sub>: MeOH (250 mL) and washed twice with H<sub>2</sub>O (75 and 50 mL). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by chromatography on 110 g of silica gel (eluent - CHCl<sub>3</sub>/MeOH/iPrNH<sub>2</sub> 95/5/5,

sample applied in  $\text{CHCl}_3/\text{MeOH}$  95/5). The purified material was concentrated *in vacuo* and then vacuum dried to give 900 mg (85%) of *N,N*-Bis(4-aminobutyl) cholesteryl carbamate.

(J) *N,N*-Bis(*N'*-3-aminopropyl-4-aminobutyl) cholesteryl carbamate

5 *N,N*-Bis(*N'*-3-aminopropyl-4-aminobutyl) cholesteryl carbamate (Figure 1, No. 83) was prepared by reacting *N,N*-Bis(4-aminobutyl) cholesteryl carbamate with acrylonitrile (82% yield) and subsequent reduction of the di acrylonitrile adduct with Raney nickel (81 % yield) as described for the preparation of  $\text{N}^1, \text{N}^8$ -Bis(3-aminopropyl)- $\text{N}^4$ -spermidine cholesteryl carbamate.

10 (K)  $\text{N}^4$  Spermidine cholesteryl carboxamide

$\text{N}^4$  Spermidine cholesteryl carboxamide ( Figure 1, No. 90) was prepared as follows.

A solution of cholesteryl chloride (5.0 g, 12.3 mmol) in THF (50 mL) was added dropwise over 0.5 h under reflux to Magnesium turnings (390 mg) in THF  
15 (25 mL). Initially a pinch of Iodine and three drops of Iodomethane were added to initiate the reaction. After refluxing for 3 h. the reaction was cooled to room temperature. This mixture was poured onto Dry ice (10 g) and then stirred for 1h. This solution was cooled in an ice bath and added to ice cold 1 M  $\text{H}_2\text{SO}_4$  (100 mL). After stirring for 5 min., sodium chloride (1 g) and diethyl ether (100 mL)  
20 was added. The layers were separated and the aqueous layer was extracted with diethyl ether (100 mL). The combined organic layers were washed with a solution of Sodium thiosulfate pentahydrate(120 mg) in  $\text{H}_2\text{O}$  (30 mL). The organic layer was concentrated *in vacuo* and vacuum dried for 18 h. The crude solid was titrated with hexanes (25 mL). After filtration the solid was washed  
25 with ice cold hexanes (10 mL). The solid was vacuum dried for 1h. The cholesteryl carboxylic acid obtained (3.0 g, 59 %) was ca. 90 % pure and used without further purification.

Cholesteryl carboxylic acid (500 mg, 1.2 mmol) and N-hydroxysuccinimide (140 mg, 1.2 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$ . To this solution was added Dicyclohexylcarbodiimide (275 mg, 1.32 mmol) was added and the reaction was stirred under  $\text{N}_2$  for 2h.  $\text{N}^1, \text{N}^8$ -dicarbobenzoxy spermidine (474 mg, 1.2 mmol) and  $\text{Et}_3\text{N}$  (1.0 mL, 7.1 mmol) was added and the reaction was stirred under  $\text{N}_2$  for 72 h. The reaction was filtered and the precipitate was washed with  $\text{CH}_2\text{Cl}_2$  (50 mL). The filtrate was washed with  $\text{H}_2\text{O}$  (25mL). The separated organic layer was dried over  $\text{MgSO}_4$  and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by chromatography on 150 g of silica gel (eluent - hexanes / EtOAc 1/ 1). The purified material was concentrated *in vacuo* and then vacuum dried to give 680 mg (70%) of  $\text{N}^1, \text{N}^8$ -dicarbobenzoxy- $\text{N}^4$ -spermidine cholesteryl carboxamide.

The carbobenzoxy group were removed from  $\text{N}^1, \text{N}^8$ -dicarbobenzoxy- $\text{N}^4$ -spermidine cholesteryl carboxamide as described in the preparation of  $\text{N}^4$ -spermidine cholesteryl carbamate. The purified product,  $\text{N}^4$  Spermidine cholesteryl carboxamide, was obtained in 53 % yield.

#### Group II Amphiphiles

(A)  $\text{N}^1, \text{N}^8$ -Bis(Arginine carboxamide)- $\text{N}^4$ -spermidine cholesteryl carbamate  
 $\text{N}^1, \text{N}^8$ -Bis(Arginine carboxamide)- $\text{N}^4$ -spermidine cholesteryl carbamate (Figure 5, No. 95) was prepared as follows.

To  $\text{N}^{\text{(a)}}\text{N}^{\text{(e)}}\text{N}^{\text{(e)}}$  (alpha, epsilon, epsilon) -tricarbenzoxyArginine in  $\text{CH}_2\text{Cl}_2$  (25 mL) was added N-hydroxysuccinimide (100 mg, 0.89 mmol) and dicyclohexylcarbodiimide (240 mg, 0.89 mmol). The mixture was stirred under  $\text{N}_2$  at room temperature for 2.5 hours.  $\text{N}^4$ -Spermidine Cholesteryl Carbamate (250 mg, 0.448 mmol) and  $\text{Et}_3\text{N}$  (0.25 mL, 1.8 mmol) was added and the reaction stirred at room temperature under  $\text{N}_2$  for 72 h. The reaction was filtered and the precipitate was washed with  $\text{CH}_2\text{Cl}_2$  (20 mL). The filtrate was washed with  $\text{H}_2\text{O}$

(20 mL). The separated organic layer was dried over  $MgSO_4$  and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by chromatography on 70 g of silica gel (eluent -  $CHCl_3$  / MeOH 95/ 5). The purified material was concentrated *in vacuo* and then vacuum dried to give 533 mg (71%) of  $N^1, N^8$ -Bis (N(a), N(e), N(e)-tricarbobenzoxy Arginine carboxamide)- $N^4$ -spermidine cholesteryl carbamate.

The carbobenzoxy group were removed from  $N^1, N^8$ -Bis (N(a), N(e), N(e)-tricarbobenzoxy Arginine carboxamide)- $N^4$ -spermidine cholesteryl carbamate as described in the preparation of N-( $N^4$ -3-aminopropylspermidine) cholesteryl carbamate. The product,  $N^1, N^8$ -Bis(Arginine carboxamide)- $N^4$ -spermidine cholesteryl carbamate was obtained in 27 % yield.

### Group III Amphiphiles

#### (A) N,N-Dioctadecyllysineamide

N,N- dioctadecyllysineamide (Figure 6, No.73) was prepared according to the following procedure. N,N-dioctadecylamine (1.35 g, 2.58 mmol, Fluka) and L-N $\alpha$ ,N $\epsilon$ -diBOCllysine N-hydroxysuccinimide ester ( 1.00 g, 2.58 mmol, Sigma) were combined in 15 ml of methylene chloride and 2 ml triethylamine was added. The reaction mixture was heated briefly to effect complete dissolution and then stirred at ambient temperature overnight. Water (20 ml) and methylene chloride (50 ml) were added to the reaction mixture and the layers were separated. The aqueous fraction was extracted a second time with 50 ml methylene chloride. The combined organic fractions were dried over  $MgSO_4$ , filtered and concentrated *in vacuo*. The residue was purified by column chromatography (150 g silica gel , eluent - hexane/ethyl acetate 8/2). The purified material, N,N-dioctadecyl-N $\alpha$ ,N $\epsilon$ -diBOC lysineamide(1.59 g) was dissolved in 25 ml of chloroform and stirred for 2 hr. while HCl gas was bubbled through the solution. This solution was purged with  $N_2$  gas and concentrated *in*

*vacuo*. N,N-dioctadecyllysineamide (1.34 g) was obtained in 68% yield as the di HCl salt.

(B) N<sup>1</sup>N<sup>1</sup>-Dioctadecyl-1,2,6-triaminohexane

N<sup>1</sup>N<sup>1</sup>-Dioctadecyl-1,2,6-triaminohexane (Figure 6, No. 47) was prepared  
5 as follows. To N,N-Dioctadecyl-N $\alpha$ ,N $\epsilon$ -diBOClysineamide (760 mg, 0.823 mmol)  
in 30 ml anhydrous THF stirred at ambient temperature was added LiAlH<sub>4</sub> (185  
mg, 4.87 mmol) in portions. The reaction mixture was stirred at ambient  
temperature overnight under a nitrogen atmosphere. The reaction was quenched  
by the dropwise addition of 2 ml water and the resulting solution was  
10 concentrated *in vacuo*. To this residue was added in order 10 mL of 1 M HCl, 50  
ml of methylene chloride, and 10 ml of 1M NaOH (final pH 10). The layers were  
separated and the aqueous fraction was extracted a second time with 50 ml of  
methylene chloride. The combined organic layers were dried over MgSO<sub>4</sub> and  
filtered. The filter cake was washed with 50 ml of methylene chloride. The  
15 combined filtrates were concentrated *in vacuo* to give 700 mg of crude product.  
The crude product was purified by column chromatography (80 g silica gel,  
eluent - hexane/ethyl acetate 7/3). The fractions containing the purified product  
were combined and concentrated *in vacuo* to obtain 490 mg of the product  
protected as the diBOC derivative. To 200 mg of this diBOC derivative was  
20 added 4 ml of chloroform and 1 ml of TFA. The resulting reaction mixture was  
stirred at ambient temperature for 2 hr and concentrated *in vacuo*. The residue  
was dissolved in 25 ml of water and 25 mL of methylene chloride and adjusted to  
pH 10 with approximately 2 ml of concentrated ammonium hydroxide. The  
layers were separated and the aqueous layer was extracted a second time with 25  
25 ml of methylene chloride. The organic fractions were combined, dried over  
Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The resulting residue was dissolved in 10 ml  
of diethyl ether, HCl gas was bubbled through the solution for 2 minutes and the

solution was cooled at 4° C overnight. The precipitated product was collected by filtration, washed with cold (4° C) diethyl ether, and dried under vacuum to obtain 160 mg of the desired product in 67% yield.

#### Group IV Amphiphiles

5 (A) 1-(N<sup>4</sup>-spermine)-2,3-dilaurylglycerol carbamate

1-(N<sup>4</sup>-spermine)-2,3-dilaurylglycerol carbamate (Figure 7, No. 89) was prepared as follows. A solution of 3-benzyloxy-1,2-propanediol (1.00 g, 5.49 mmol) in THF (20 mL) was added to a suspension of sodium hydride (60% w/w in oil, 550 mg, 13.725 mmol) in THF (30 mL) and allowed to reflux overnight  
10 under dry nitrogen. A solution of dodecyl methane sulfonate (3.39 g, 12.078 mmol) in THF (20 mL) was added and the reaction was refluxed for another two days. After cooling to room temperature the reaction was filtered through a bed of Celite, rinsing with THF. The filtrate was reduced *in vacuo* to a yellow oil which was redissolved in diethyl ether (100 mL). The ether solution was washed  
15 with 0.1 N NaOH (30 mL) and dH<sub>2</sub>O (2 x 30 mL). The organic layer was dried over magnesium sulfate, filtered and reduced *in vacuo* to a red-brown oil. The crude material was purified by flash column chromatography (300 g silica gel) eluting with 3% ethyl acetate/ hexanes. The desired product was isolated as a pale yellow oil and characterized by <sup>1</sup>H NMR as 3-OBn-1,2-dilaurylglycerol (1.70  
20 g, 60%). 3-OBn-1,2-dilaurylglycerol (1.70 g, 3.28 mmol) in ethanol (100 mL) was stirred with 10% Pd/C (250 mg, 15 wt%) under a hydrogen atmosphere for 24 hours. The reaction was flushed with nitrogen and filtered through Celite, rinsing with ethanol, to remove the catalyst. The filtrate was reduced *in vacuo* to a solid. The crude material was purified by flash column chromatography (140 g  
25 silica gel) eluting with 10% ethyl acetate/ hexanes. The desired product was isolated as a white solid and characterized by <sup>1</sup>H NMR as 1,2-dilaurylglycerol (1.23g, 88%).

A 1.93 M solution of phosgene in toluene (0.77 mL, 1.49 mmol) was added to a solution of 1,2-dilaurylglycerol (580 mg, 1.35 mmol) and N,N-diisopropylethylamine (0.26 mL, 1.49 mmol) in methylene chloride (10 mL) and stirred overnight. A solution of N<sup>1</sup>,N<sup>12</sup>-di-CBz-spermine•2HCl (734 mg, 1.35 mmol) in 60: 25: 4 chloroform/ methanol/ water (80 mL) was added. After 3 hours another equivalent of N,N-diisopropylethylamine (0.26 mL, 1.49 mmol) was added. An additional 0.5 equivalents of N,N-diisopropylethylamine (0.13 mL, 0.75 mmol) was added three hours later and the reaction was allowed to stir overnight under nitrogen at ambient temperature. The reaction was washed with 1M NaOH (20 mL) and dH<sub>2</sub>O (15 mL). The organic layer was separated, dried over magnesium sulfate, filtered and reduced *in vacuo* to a white solid. The crude material was purified by flash column chromatography (125 g silica gel) eluting with 90: 10: 0.5 chloroform/ methanol/ ammonium hydroxide. The desired product was isolated as an oil and characterized by <sup>1</sup>H NMR as 1-(N<sup>4</sup>-(N<sup>1</sup>,N<sup>12</sup>-di-CBz-spermine))-2,3-dilaurylglycerol carbamate (188 mg, 15%).

The 1-(N<sup>4</sup>-(N<sup>1</sup>,N<sup>12</sup>-di-CBz-spermine))-2,3-dilaurylglycerol carbamate (188 mg, 0.203 mmol) was dissolved in glacial acetic acid (10 mL) and stirred with 10% Pd/C (45 mg, 24 wt %) under a hydrogen atmosphere for 5 hours. The catalyst was removed by vacuum filtration rinsing with 10% acetic acid/ ethyl acetate (10 mL). The filtrate was reduced to an oil by rotary evaporation. The resulting oil was dissolved in 10% methanol/ chloroform (85 mL) and was washed with 1M NaOH (15 mL) and dH<sub>2</sub>O (10 mL). The organic layer was separated, dried over magnesium sulfate, filtered and reduced *in vacuo* to an oil. The product was characterized by <sup>1</sup>H NMR as 1-(N<sup>4</sup>-spermine)-2,3-dilaurylglycerol carbamate (125 mg, 94%).

Other amphiphiles of the invention may be prepared according to procedures that are within the knowledge of those skilled in art.

### Examples

The following Examples are representative of the practice of the invention.

#### Example 1 - Cell Transfection Assay

Separate 3.35  $\mu$ mole samples of spermidine cholesterol carbamate  
5 (amphiphile No. 53) and the neutral lipid dioleoylphosphatidylethanolamine  
("DOPE") were each dissolved in chloroform as stock preparations. Following  
combination of the solutions, a thin film was produced by removing chloroform  
from the mixture by evaporation under reduced pressure (20 mm Hg). The film  
was further dried under vacuum (1 mm Hg) for 24 hours. As aforementioned,  
10 some of the amphiphiles of the invention participate in transacylation reactions  
with co-lipids such as DOPE, or are subject to other reactions which may cause  
decomposition thereof. Accordingly, it is preferred that amphiphile/co-lipid  
compositions be stored at low temperature, such as -70 degrees C, until use.

To produce a dispersed suspension, the lipid film was then hydrated with  
15 sterile deionized water (1 ml) for 10 minutes, and then vortexed for 1 minute (  
sonication for 10 to 20 seconds in a bath sonicator may also be used, and  
sonication has proved useful for other amphiphiles such as DC-chol). The  
resulting suspension was then diluted with 4 ml of water to yield a solution that  
is 670 $\mu$ M in cationic amphiphile and 670 $\mu$ M in neutral colipid.

20 Experiments were also performed using spermine cholesterol carbamate  
(amphiphile No. 67) and other amphiphiles of the invention. With respect to  
spermine cholesterol carbamate, the optimum molar ratio of amphiphile to  
DOPE under the conditions tested was determined to be 1:2, not 1:1. Optimized  
ratios for many of the amphiphiles of the invention are reported in Figures 13, 14  
25 and 15, and are readily determined by those skilled in the art.

For preparation of the transfecting solution, DNA encoding for  $\beta$ -  
galactosidase (pCMV $\beta$ , ClonTech., Palo Alto, CA) was dissolved in OptiMEM

culture medium (Gibco/ BRL No. 31885-013). The resulting solution had a DNA concentration of 960  $\mu\text{M}$  (assuming an average molecular weight of 330 daltons for nucleotides in the encoding DNA).

The following procedure was used to test a 1:1 molar mixture of the  
5 cationic amphiphile spermidine cholesterol carbamate in combination with  
DOPE. A 165  $\mu\text{l}$  aliquot of spermidine cholesterol carbamate (670  $\mu\text{M}$ ) containing  
also the colipid ( at 670  $\mu\text{M}$  ) was pipetted into 8 separate wells in a 96-well plate  
containing OptiMEM (165 $\mu\text{l}$ ) in each well. The resulting 335  $\mu\text{M}$  solutions were  
then serially diluted 7 times to generate 8 separate amphiphile-containing  
10 solutions having concentrations ranging from 335  $\mu\text{M}$  to 2.63  $\mu\text{M}$ , with each  
resultant solution having a volume of 165  $\mu\text{l}$ . Thus, 64 solutions were prepared  
in all, there being 8 wells each of 8 different concentrations of amphiphile/DOPE.

Independently, DNA solutions (165 $\mu\text{l}$ , 960 $\mu\text{M}$ ) were pipetted into 8 wells  
containing OptiMEM (165  $\mu\text{l}$ ), and the resulting 480 $\mu\text{M}$  solutions were then  
15 serially diluted 7 times to generate 8 separate 165  $\mu\text{l}$  solutions from each well,  
with the concentrations of DNA in the wells ranging from 480  $\mu\text{M}$  to 3.75  $\mu\text{M}$ .

The 64 test solutions (cationic amphiphile: neutral lipid) were then  
combined with the 64 DNA solutions to give separate mixtures in 64 wells, each  
having a volume of 330 $\mu\text{l}$ , with DNA concentrations ranging from 240  $\mu\text{M}$  to  
20 1.875  $\mu\text{M}$  along one axis, and lipid concentrations ranging from 167  $\mu\text{M}$  to 1.32  
 $\mu\text{M}$  along the other axis. Thus 64 solutions were prepared in all, each having a  
different amphiphile: DNA ratio and /or concentration. The solutions of DNA  
and amphiphile were allowed to stand for 15 to 30 minutes in order to allow  
complex formation.

25 A CFT-1 cell line (human cystic fibrosis bronchial epithelial cells  
immortalized with papillomavirus) provided by Dr. James Yankaskas, University  
of North Carolina, Chapel Hill, was used for the *in vitro* assay. The cells are

homozygous for a mutant allele (deletion of phenylalanine at position 508, hereinafter  $\Delta$  F508 ) of the gene encoding for cystic fibrosis transmembrane conductance regulator ("CFTR") protein. CFTR is a cAMP-regulated chloride ( $\text{Cl}^-$ ) channel protein. Mutation of the CFTR gene results typically in complete  
5 loss ( or at least substantial impairment) of  $\text{Cl}^-$  channel activity across, for example, cell membranes of affected epithelial tissues.

The  $\Delta$  F508 mutation is the most common mutation associated with cystic fibrosis disease. For a discussion of the properties of the  $\Delta$  F508 mutation and the genetics of cystic fibrosis disease see, in particular, Cheng et al., Cell, 63, 827-834  
10 (1990). See also Riordan et al., Science, 245, 1066-1073 (1989); published European Patent Application No. 91301819.8 of Gregory et al., bearing publication number 0 446 017 A1; and Gregory et al., Nature, 347, 382-385 (1990).

The cells were cultured in Hams F12 nutrient media (Gibco/ BRL No. 31765-027) supplemented with 2% fetal bovine serum ("FBS", Irvine Scientific,  
15 No. 3000) and 7 additional supplements. Cells were then plated into 96-well tissue culture plates at a density of approximately 7,500 cells/well. Before being used in the assay, cells were allowed to grow for periods of 5-7 days until a confluent pattern had been achieved.

Following the allotted time period, three 96-well plates with CFT-1 cells  
20 were aspirated in order to remove the growth medium. The various concentrations of DNA-lipid complex (in 100  $\mu\text{l}$  aliquots) were transferred to each of three 96-well plates bringing the DNA-lipid complexes in contact with the cells. DNA-only/cell and lipid-only/cell control wells were also prepared on one of the three plates.

25 The 100  $\mu\text{l}$  solutions of DNA-lipid complex were maintained over the cells for 6 hours, after which 50  $\mu\text{l}$  of 30% FBS (in OptiMEM) was added to each well. After a further 20-hour incubation period, an additional 100  $\mu\text{l}$  of 10% FBS in

OptiMEM was also added. Following a further 24-hour incubation period, cells were assayed for expression of protein and  $\beta$ -galactosidase.

For the assays, the resultant medium was removed from the plates and the cells washed with phosphate buffered saline. Lysis buffer (50  $\mu$ l, 250 mM Tris-HCl, pH 8.0, 0.15% Triton X-100) was then added, and the cells were lysed for 30 minutes. The 96-well plates were carefully vortexed for 10 seconds to dislodge the cells and cell debris, and 5  $\mu$ l volumes of lysate from each well were transferred to a plate containing 100 $\mu$ l volumes of Coomassie Plus® protein assay reagent (Pierce Company, No. 23236). The protein assay plates were read by a Bio-Rad Model 450 plate-reader containing a 595nm filter, with a protein standard curve included in every assay.

The level of  $\beta$ -galactosidase activity in each well was measured by adding phosphate buffered saline (50  $\mu$ l) to the remaining lysates, followed by addition of a buffered solution consisting of chlorophenol red galactopyranoside (100  $\mu$ l, 1 mg per ml, Calbiochem No. 220588), 60 mM disodium hydrogen phosphate pH 8.0, 1 mM magnesium sulfate, 10 mM potassium chloride, and 50 mM 2-mercaptoethanol. The chlorophenol red galactopyranoside, following enzymatic ( $\beta$ -galactosidase) hydrolysis, gave a red color which was detected by a plate-reader containing a 570 nm filter. A  $\beta$ -galactosidase (Sigma No. G6512) standard curve was included to calibrate every assay.

Following subtraction of background readings, optical data determined by the plate-reader allowed determination of  $\beta$ -galactosidase activity and protein content. In comparison to the amount of  $\beta$ -galactosidase expressed by known transfectants, for example, DMRIE (1,2-dimyristyloxypropyl-3-dimethyl-hydroxyethyl ammonium bromide), compounds of the invention are particularly effective in transfecting airway epithelial cells and inducing therein  $\beta$ -galactosidase expression. Relative to DMRIE:DOPE (1:1), the spermidine

cholesterol carbamate: DOPE mixture (also 1:1) demonstrated transfection efficiency improved by a factor of about 5 (see, for example, Figures 13, 14 and 15).

Example 2 - Transfection of the Gene Encoding for Human Cystic Fibrosis

5 Transmembrane Conductance Regulator Protein

The ability of the cationic amphiphiles of the invention to transfect cells and to induce therein biochemical corrections was demonstrated with a separate *in vitro* assay. Immortalized human cystic fibrosis airway cells (CFT-1, as above) were used.

10 In preparation for the assay, the cells were grown on glass coverslips until approximately 60% confluent. The cells were then transfected with a complex of spermidine cholesterol carbamate:DOPE (1:1) and a plasmid(pCMV- CFTR) containing a cDNA that encodes wild type human CFTR. pCMV-CFTR plasmid is a construct containing the encoding sequence for CFTR and the following  
15 regulatory elements, a CMV promoter and enhancer, and an SV40 polyadenylation signal. Additional constructs suitable for the practice of this example include pMT-CFTR, Cheng et al., Cell, 63, 827-834 (1990). The complex used was 10.5  $\mu$ molar of spermidine cholesterol carbamate (also of DOPE) and 30  $\mu$ molar of pCMV-CFTR based on nucleotide.

20 48 hours after amphiphile-mediated transfection, cells were tested for cAMP-stimulated Cl<sup>-</sup> channel activity using the 6-methoxy-N-(3-sulfopropyl)quinolinium ("SPQ") assay. See S. Cheng et al., Cell, 66, 1027-1036 (1991) for further information concerning assay methodology. In the assay, cAMP-dependent Cl<sup>-</sup> channel activity was assessed using "SPQ" (from  
25 Molecular Probes, Eugene, Oregon), a halide-sensitive fluorophore. Increases in halide permeability results in a more rapid increase in SPQ fluorescence, and the rate of change (rather than the absolute change in fluorescence) is the important

variable in assessing  $\text{Cl}^-$  permeability. See also Rich et al., Nature, 347, 358-363 (1990) for background information.

Fluorescence of the SPQ molecule in individual cells was measured using an inverted microscope, Nikon, , a digital imaging system from Universal  
5 Imaging, and an ICCD camera, Hamamatsu, Inc.. Cells were selected for analysis without prior knowledge of their expected rate-of-change- in-fluorescence characteristics.

In each experiment, up to five microscope fields of between 90 and 100 cells were examined on a given day, and studies under each condition were  
10 repeated on at least 3 different days. Since expression of CFTR is heterogenous (i.e. cells do not produce identical amounts of CFTR), the data presented were for the 20% of cells in each field exhibiting the greatest response.

As expected, cells that were mock transfected failed to exhibit any measurable increase in cAMP-stimulated halide fluorescence. In contrast, cells  
15 that had been transfected with the wild type CFTR cDNA displayed a rapid increase in SPQ fluorescence upon stimulation with cAMP agonist, indicating increased permeability to anions. Approximately 60% of the cells assayed exhibited measurable cAMP-stimulated  $\text{Cl}^-$  channel activity. Accordingly, spermidine cholesterol carbamate, and other cationic amphiphiles of the  
20 invention similarly tested, are effective in transferring CFTR-encoding plasmid into immortalized CF airway cells.

#### Example 3 - CAT Assay

##### part A

This assay was used to assess the ability of the cationic amphiphiles of the  
25 invention to transfect cells in vivo from live specimens. In the assay, the lungs of balb/c mice were instilled intra-nasally (the procedure can also be performed trans-tracheally) with 100  $\mu\text{l}$  of cationic amphiphile:DNA complex, which was

allowed to form during a 15-minute period prior to administration according to the following procedure. The amphiphile (premixed with co-lipid, see below) was hydrated in water for 10 minutes, a period sufficient to yield a suspension at twice the final concentration required. This was vortexed for two minutes and  
5 aliquoted to provide 55 microliter quantities for each mouse to be instilled. Similarly, DNA encoding the reporter (CAT) gene was diluted with water to a concentration twice the required final concentration, and then aliquoted at 55 microliters for each mouse to be instilled. The lipid was gently combined with the DNA (in a polystyrene tube), and the complex allowed to form for 15 minutes  
10 before the mice were instilled therewith.

The plasmid used (pCMVHI-CAT, see Example 4) provides an encoding DNA for chloramphenicol transferase enzyme. Specifics on the amphiphile:DNA complexes are provided below.

Two days following transfection, mice were sacrificed, and the lungs and  
15 trachea removed, weighed, and homogenized in a buffer solution (250 mM Tris, pH 7.8, 5mM EDTA). The homogenate was clarified by centrifugation, and the deacetylases therein were inactivated by heat treatment at 70 °C for 10 minutes. Lysate was incubated overnight with acetyl coenzyme A and C<sup>14</sup>--chloramphenicol. CAT enzyme activity was then visualized by thin layer  
20 chromatography ("TLC") following an ethyl acetate extraction. Enzyme activity was quantitated by comparison with a CAT standard curve.

The presence of the enzyme CAT will cause an acetyl group to be transferred from acetylcoenzyme A to C<sup>14</sup>--chloramphenicol. The acetylated/radiolabeled chloramphenicol migrates faster on a TLC plate and  
25 thus its presence can be detected. The amount of CAT that had been necessary to generate the determined amount of acetylated chloramphenicol can then be calculated from standards.

The activity of spermidine cholesterol carbamate (amphiphile No.53) was determined in the CAT assay in relation to the recognized transfection reagents DMRIE and DC-Chol. Figure 10 demonstrates dramatically (as ng CAT activity per 100 mg tissue) the enhanced ability of spermidine cholesterol carbamate (amphiphile No. 53) to transfect cells in vivo, which enhancement is about 20-fold, or greater, in this assay. In the assay, activity was measured as ng CAT enzyme per 100 mg lung tissue. As a comparison, it is generally observed that DMRIE, a well known transfectant, when prepared as a 1:1 molar mixture with DOPE and then complexed with plasmid DNA (1.7 mM DMRIE, 1.7 mM DOPE, 1.2 mM plasmid DNA measured as nucleotide) gives about 1 to 2 ng activity per 100 mg lung tissue in this assay.

With respect to the comparison provided by Figure 10, the following conditions are of note. The transfection solution for spermidine cholesterol carbamate contained 6mM DNA measured as concentration of nucleotide, and 1.5 mM of cationic amphiphile. Following generally the procedure of Example 1, each amphiphile had also been premixed with DOPE, in this case at 1:1 molar ratio. For transfection with DC-chol, the molar ratio of DC-chol to DOPE was 3:2, and the concentrations of cationic amphiphile and of DNA (as nucleotide) were 1.3 mM and 0.9 mM, respectively. For transfection with DMRIE, the molar ratio of DMRIE to DOPE was 1:1 and the concentrations of cationic amphiphile and of DNA were 1.7 mM and 1.2 mM, respectively. These concentrations (and concentration ratios) for each amphiphile, and colipid and DNA, had been determined to be optimal for transfection for that respective amphiphile, and accordingly were used as the basis for the comparison presented herein.

For spermidine cholesterol carbamate (amphiphile No. 53), optimization experiments were also performed to determine preferred concentrations of plasmid for a particular amphiphile concentration (see Figure 11), and also to

determine preferred concentrations of the same amphiphile in relation to a particular plasmid concentration (see Figure 12). Transfection efficiency was optimal at an amphiphile concentration of 1.5 mM (DOPE also being present at 1.5 mM), and about 6 mM (by nucleotide) of plasmid, or about at a ratio of 1:4. It was noted, however, that concentrations of about 0.75 mM of amphiphile, and 3.0 mM of plasmid were less toxic to the target cells.

Intra-nasal transfection with pCMVHI-CAT vector was also performed in mice using spermidine cholesterol carbamate as cationic amphiphile but with cholesterol as co-lipid. In this experiment, the concentrations of spermidine cholesterol carbamate tested were between 1.0 and 1.5 mM (cholesterol being present at a 1:1 molar ratio in each case, with the mixing of amphiphile and co-lipid being performed as above). The DNA concentration (measured as nucleotide concentration) was between 4.0 and 6.0 mM. Transfection efficiency (again measured as ng CAT/100 mg tissue) was less effective than with DOPE as co-lipid; however, the transfections were substantially more effective than those achieved using DC-Chol/DOPE.

#### part B

Additional experiments were performed to compare in vivo the transfection efficiency of cationic amphiphiles depicted in Figures 1, 5 and 7. Results therefor are reported in Figures 13, 14 and 15 respectively. The compounds were administered intra-nasally using between 12 and 15 mice per compound. As in part A above, ng CAT activity was measured per 100 mg of tissue. However, improved vectors (pCF1/CAT and its near equivalent pCF2/CAT) were used. In part resulting from improved vector performance, incubations of lysate with acetyl coenzyme A and C<sup>14</sup>-chloramphenicol were conducted for only 30 minutes. Construction of pCF1/CAT and pCF2/CAT is described below in Example 4.

The in vivo data reported in Figures 13, 14 and 15 were compiled generally as follows. As aforementioned, Figures 10 and 11 report data from the complete in vivo optimization of amphiphile No. 53. Amphiphile No. 67 was subjected to a similar partial optimization. With respect to all of the other cationic amphiphiles reported on, and taking advantage of numerous structural similarities, optimized compositions for in vivo testing were extrapolated from in vitro results. This facilitated the screening of large numbers of amphiphiles and produced broadly, if not precisely, comparable data. For all amphiphiles other than Nos. 53 and 67, the ratio, for in vivo testing, of amphiphile concentration to DOPE concentration was taken from the in vitro experiments, as was the optimized ratio of amphiphile concentration to DNA concentration (see Example 1). Accordingly, for such amphiphiles the in vivo test concentration was fixed at 1mM, thereby fixing also the co-lipid concentration. [Broadly, the molar ratio of the amphiphile to co-lipid DOPE ranged from 1:2 (for example, spermine cholesterol carbamate, No. 67) through 1:1 (for example, spermidine cholesterol carbamate, No. 53) to about 2:1 (for example, amphiphile No. 75)]. The concentration of plasmid DNA varied for each amphiphile species tested in order to duplicate the optimized amphiphile/DNA ratio that had been determined in vitro.

20 part C

That the novel amphiphiles of the invention are an important contribution to the art is immediately seen by comparing their performance - as in vivo transfection enhancers - to that of closely related cationic amphiphiles that lack the novel T-shape. It has been determined that spermidine cholesterol carbamate provides a much greater level of enhancement than N<sup>1</sup>-spermidine cholesteryl carbamate which contains the same number of carbon and nitrogen atoms in its cationic alkylamine component but which is linear and not "T-shaped".

Following generally the procedures of Example 3, part B, and using respectively 6mM (as nucleotide), 1.5 mM, and 1.5 mM concentrations of DNA, amphiphile and of co-lipid, the transfection enhancement provided by spermidine cholesterol carbamate (amphiphile No.53) , in relation to N<sup>1</sup>-spermidine  
5 cholesteryl carbamate, was determined to be about 30 fold.

Also following the procedures of Example 3, part B, and using respectively 4mM (as nucleotide), 1mM, and 2 mM concentrations of DNA, amphiphile and co-lipid, the transfection enhancement provided by spermine cholesterol carbamate (amphiphile No. 67) — in relation to N<sup>1</sup>-thermospermine  
10 cholesteryl carbamate and N<sup>1</sup>-spermine cholesteryl carbamate to which spermine cholesterol carbamate is similarly related — is at least about 30 fold.

#### Example 4- Construction of vectors

As aforementioned, numerous types of biologically active molecules can be transported into cells in therapeutic compositions that comprise one or more  
15 of the cationic amphiphiles of the invention. In an important embodiment of the invention, the biologically active macromolecule is an encoding DNA. There follows a description of novel vectors (plasmids) that are preferred in order to facilitate expression of such encoding DNAs in target cells.

#### part A — pCMVHI-CAT

20 pCMVHI-CAT is representative of plasmid constructs useful in the practice of the invention. Although the plasmid is provided in a form carrying a reporter gene (see Example 3), transgenes having therapeutic utility may also be included therein.

The pCMVHI-CAT vector is based on the commercially available vector  
25 pCMV $\beta$  (Clontech). The pCMV $\beta$  construct has a pUC19 backbone (J. Vieira, et al., Gene , 19, 259-268, 1982) that includes a procaryotic origin of replication derived originally from pBR322. Basic features of the pCMVHI-CAT plasmid (as

constructed to include a nucleotide sequence coding for CAT) are as follows. Proceeding clockwise — the human cytomegalovirus immediate early gene promoter and enhancer, a fused tripartite leader from adenovirus and a hybrid intron, a linker sequence, the CAT cDNA, an additional linker sequence, the late SV40 polyadenylation signal, and the pUC origin of replication and backbone that includes the gene for ampicillin resistance.

The human cytomegalovirus immediate early gene promoter and enhancer spans the region from nucleotides 1-639. This corresponds to the region from -522 to +72 relative to the transcriptional start site (+1) and includes almost the entire enhancer region from -524 to -118 as originally defined by Boshart et al, Cell 41:521-530, 1985. The CAAT box is located at nucleotides 487-491 and the TATA box is at nucleotides 522-526 in pCMVHI-CAT. The CAT transcript is predicted to initiate at nucleotide 549, which is the transcriptional start site of the CMV promoter. The tripartite leader-hybrid intron is composed of a fused tri-partite leader from adenovirus containing a 5' splice donor signal, and a 3' splice acceptor signal derived from an IgG gene. The elements in the intron are as follows: the first leader, the second leader, part of the third leader, the splice donor sequence and intron region from the first leader, and the mouse immunoglobulin gene splice donor sequence. The length of the intron is 230 nucleotides. The CAT coding region comprises nucleotides 1257-1913. The SV40 poly A signal extends from nucleotide 2020 to 2249.

Accordingly, construction of the pCMVHI-CAT plasmid proceeded as follows. The vector pCMV $\beta$  (Clontech, Palo Alto, CA) was digested with Not I to excise the  $\beta$ -galactosidase gene. The vector fragment lacking the  $\beta$ -galactosidase gene was isolated and ligated to form pCMV.

The hybrid intron (Figure 17) was obtained from the plasmid pADB (Clontech). The hybrid intron had been isolated from a 695 base pair XhoI-EcoRI

fragment of p91023(B), see Wong et al., *Science*, 228, 810-815 (1985). The hybrid intron contains the fused tripartite leader from adenovirus, the donor site from the first segment of the tripartite leader, and the acceptor site from an IgG gene, and has a length of 230 bp.

5 pAD $\beta$  was digested with Pml I and Not I, and the ~500 base-pair (bp) fragment was isolated, and then ligated into the Not I site of pBluescriptII KS(-) (Stratagene, La Jolla, CA) to form pBlueII-HI.

pBlueII-HI was digested with XhoI and NotI to excise the hybrid intron fragment. This fragment was ligated into the XhoI and NotI sites of pCMV, 10 replacing the SV40 intron to form pCMVHI.

The CAT gene was obtained from the Chloramphenicol Acetyltransferase GenBlock (Pharmacia, Piscataway, NJ). This 792 bp Hind III fragment was blunted with the Klenow fragment of DNA Polymerase I, then Not I linkers (New England Biolabs) were ligated to each end. After digestion with Not I to 15 expose the Not I sticky ends, the fragment was subcloned into the Not I site of pCMV to form pCMV-CAT. pCMV-CAT was digested with Not I to excise the CAT fragment. The CAT fragment was ligated into pCMVHI to form pCMVHI-CAT which is depicted in Figure 16.

part B — pCF1 and pCF2

20 Although pCMVHI is suitable for therapeutic transfections, further performance enhancements (including increased expression of transgenes) are provided by the pCF1 and pCF2 plasmids. A map of pCF1/CAT is shown in Figure 18, panel A, and a map of pCF2/CAT is shown in panel B.

Briefly, pCF1 contains the enhancer/promoter region from the immediate 25 early gene of cytomegalovirus (CMV). A hybrid intron is located between the promoter and the transgene cDNA. The polyadenylation signal of the bovine growth hormone gene was selected for placement downstream from the

transgene. The vector also contains a drug-resistance marker that encodes the aminoglycosidase 3'-phosphotransferase gene (derived from the transposon Tn903, A. Oka et al., Journal of Molecular Biology, 147, 217-226, 1981) thereby conferring resistance to kanamycin. Further details of pCF1 structure are  
5 provided directly below, including description of placement therein of a cDNA sequence encoding for cystic fibrosis transmembrane conductance regulator (CFTR) protein.

The pCF1 vector is based on the commercially available vector pCMV $\beta$  (Clontech). The pCMV $\beta$  construct has a pUC19 backbone (J. Vieira, et al., Gene,  
10 19, 259-268, 1982) that includes a procaryotic origin of replication derived originally from pBR322.

Basic features of the pCF1-plasmid (as constructed to include a nucleotide sequence coding for CFTR) are as follows. Proceeding clockwise — the human cytomegalovirus immediate early gene promoter and enhancer, a fused tripartite  
15 leader from adenovirus and a hybrid intron, a linker sequence, the CFTR cDNA, an additional linker sequence, the bovine growth hormone polyadenylation signal, pUC origin of replication and backbone, and the kanamycin resistance gene. The pCF1-CFTR plasmid has been completely sequenced on both strands.

The human cytomegalovirus immediate early gene promoter and  
20 enhancer spans the region from nucleotides 1-639. This corresponds to the region from -522 to +72 relative to the transcriptional start site (+1) and includes almost the entire enhancer region from -524 to -118 as originally defined by Boshart et al., Cell, 41, 521-530 (1985). The CAAT box is located at nucleotides 486-490 and the TATA box is at nucleotides 521-525 in pCF1-CFTR. The CFTR  
25 transcript is predicted to initiate at nucleotide 548, which is the transcriptional start site of the CMV promoter.

The hybrid intron is composed of a fused tri-partite leader from adenovirus containing a 5' splice donor signal, and a 3' splice acceptor signal derived from an IgG gene. The elements in the intron are as follows: the first leader (nucleotides 705-745), the second leader (nucleotides 746-816), the third leader (partial, nucleotides 817-877), the splice donor sequence and intron region from the first leader (nucleotides 878-1042), and the mouse immunoglobulin gene splice donor sequence (nucleotides 1043-1138). The donor site (G | GT) is at nucleotides 887-888, the acceptor site (AG | G) is at nucleotides 1128-1129, and the length of the intron is 230 nucleotides. The CFTR coding region comprises nucleotides 1183-5622.

Within the CFTR-encoding cDNA of pCF1-CFTR, there are two differences from the originally-published predicted cDNA sequence (J. Riordan et al., Science, 245, 1066-1073, 1989); (1) an A to C change at position 1990 of the CFTR cDNA which corrects an error in the original published sequence, and (2) a T to C change introduced at position 936. The change at position 936 was introduced by site-directed mutagenesis and is silent but greatly increases the stability of the cDNA when propagated in bacterial plasmids (R. J. Gregory et al. et al., Nature, 347, 382-386, 1990). The 3' untranslated region of the predicted CFTR transcript comprises 51 nucleotides of the 3' untranslated region of the CFTR cDNA, 21 nucleotides of linker sequence and 114 nucleotides of the BGH poly A signal.

The BGH poly A signal contains 90 nucleotides of flanking sequence 5' to the conserved AAUAAA and 129 nucleotides of flanking sequence 3' to the AAUAAA motif. The primary CFTR transcript is predicted to be cleaved downstream of the BGH polyadenylation signal at nucleotide 5808. There is a deletion in pCF1-CFTR at position +46 relative to the cleavage site, but the deletion is not predicted to effect either polyadenylation efficiency or cleavage

site accuracy, based on the studies of E.C. Goodwin et al., J. Biol. Chem., 267, 16330-16334 (1992). After the addition of a poly A tail, the size of the resulting transcript is approximately 5.1 kb.

pCF2 plasmid, Figure 18 (B), contains a second CMV enhancer, in tandem  
5 with the first. Enhanced expression of transgenes from pCF1 or pCF2 results from the combination of a strong promoter, the presence of a highly efficient polyadenylation signal, a leader sequence that enhances translation, and an intron to increase message stability.

Example 5- Correction of Chloride Ion Transport Defect in Nasal Polyp  
10 Epithelial Cells of a Cystic Fibrosis Patient by Cationic Amphiphile-Mediated Gene Transfer

Primary (non-immortalized) nasal polyp cells from an adult male cystic fibrosis patient (homozygous for the  $\Delta F508$  mutation) were grown on collagen-coated permeable filter supports (Millicells) to form a polarized and confluent  
15 epithelial monolayer. Once the monolayer was electrically tight (about 5 to 7 days post seeding, and as indicated by the development of resistance across the cell sheet), the apical surface can be exposed to formulations of cationic amphiphile: DNA complex.

In this case, the amphiphile (spermidine cholesterol carbamate) was  
20 provided as a 1:1 (by mole) mixture with DOPE, and this mixture was then complexed with pCMV-CFTR plasmid vector (a construct encoding wild type human cystic fibrosis transmembrane conductance regulator protein, see above). Concentrations in the final mixture were 42  $\mu$ molar of spermidine cholesterol carbamate (and also of DOPE) and 60  $\mu$ molar (based on molarity in nucleotides)  
25 of the plasmid expression vector.

Expression of CFTR was determined by measuring cAMP-stimulated transepithelial chloride secretion in a modified Ussing chamber, Zabner et al.,

Nature Genetics ,6, 75-83 (1984). The mucosal side of the epithelium was bathed in Ringer's bicarbonate solution bubbled with 95% O<sub>2</sub> and 5% CO<sub>2</sub>. The composition of the submucosal solution was similar to the mucosal solution with the exception that sodium gluconate replaced sodium chloride. Transepithelial  
5 voltage was clamped to 0 mV and short circuit current was recorded. Amiloride (10 μM) was applied into the apical bath, followed by the mucosal addition of forskolin and IBMX (at 100 μM each). 5-nitro-2-(3-phenylpropylamino) benzoic acid ("NPPB") , an inhibitor of CFTR chloride channels, was then added to the mucosal solution at 10 to 30 μM.

10 Chloride secretion (i.e. movement of chloride from the epithelial cells to the mucosal solution) is shown as an upward deflection (see Figure 19A). The same plasmid vector, but containing a reporter gene, was used as a negative control (Figure 19B) . A cAMP-stimulated current (0.5 to 2.5 μampere/cm<sup>2</sup>) was observed in monolayers transfected with wild type CFTR gene. Current was not detected with the  
15 pCMB-β-galactosidase control.

Example 6- Correction of Chloride Ion Transport Defect in Airway Epithelial Cells of a Cystic Fibrosis Patient by Cationic Amphiphile-Mediated Gene Transfer

A recommended procedure for formulating and using the pharmaceutical  
20 compositions of the invention to treat cystic fibrosis in human patients is as follows.

Following generally the procedures described in Example 1, a thin film (evaporated from chloroform) is produced wherein spermine cholesterol carbamate (amphiphile No. 67) and DOPE are present in the molar ratio of 1: 2.  
25 The amphiphile-containing film is rehydrated in water-for -injection with gentle vortexing to a resultant amphiphile concentration of about 3mM. However, in order to increase the amount of amphiphile/DNA complex that may be stably

delivered by aerosol as a homogeneous phase (for example, using a Puritan Bennett Raindrop nebulizer from Lenexa Medical Division, Lenexa, KS, or the PARI LC Jet™ nebulizer from PARI Respiratory Equipment, Inc., Richmond, VA), it may be advantageous to prepare the amphiphile-containing film to  
5 include also one or more further ingredients that act to stabilize the final amphiphile/DNA composition. Accordingly, it is presently preferred to prepare the amphiphile-containing film as a 1: 2 : 0.05 molar mixture of amphiphile No. 67, DOPE, and PEG(5000)-DMPE. [A suitable source of PEG-DMPE, polyethylene glycol 5000 - dimyristoylphosphatidyl ethanolamine, is Catalog No.  
10 880210 from Avanti Polar Lipids, Alabaster, AL]. Additional fatty acid species of PEG-PE may be used in replacement therefor.

Without being limited as to theory, PEG(5000)-DMPE is believed to stabilize the therapeutic compositions by preventing further aggregation of formed amphiphile/DNA complexes. Additionally it is noted that PEG(2000)-  
15 DMPE was found to be less effective in the practice of the invention.

pCF1-CFTR plasmid (containing an encoding sequence for human cystic fibrosis transmembrane conductance regulator, see Example 4) is provided in water-for-injection at a concentration, measured as nucleotide, of 4 mM. Complexing of the plasmid and amphiphile is then allowed to proceed by gentle  
20 contacting of the two solutions for a period of 10 minutes.

It is presently preferred to deliver aerosolized DNA to the lung at a concentration thereof of between about 2 and about 12 mM (as nucleotide). A sample of about 10 to about 40 ml is generally sufficient for one aerosol administration to the lung of an adult patient who is homozygous for the  $\Delta$  F508  
25 mutation in the CFTR-encoding gene.

It is expected that this procedure (using a freshly prepared sample of amphiphile/DNA) will need to be repeated at time intervals of about two weeks,

but depending considerably upon the response of the patient, duration of expression from the transfected DNA, and the appearance of any potential adverse effects such as inflammation, all of which can be determined for each individual patient and taken into account by the patient's physicians.

5           One important advantage of the cationic amphiphiles of the present invention is that they are substantially more effective — in vivo — than other presently available amphiphiles, and thus may be used at substantially lower concentrations than known cationic amphiphiles. There results the opportunity to substantially minimize side effects (such as amphiphile toxicity, inflammatory  
10 response) that would otherwise affect adversely the success of the gene therapy.

A further particular advantage associated with use of many of the amphiphiles of the invention should again be mentioned. Many of the amphiphiles of the invention were designed so that the metabolism thereof would rapidly proceed toward relatively harmless biologically-compatible  
15 components. In this regard, highly active amphiphiles 53, 67, and 75 are of particular note.

#### Alternate Procedure to Prepare an Amphiphile/Co-lipid Composition

In order to formulate material that is suitable for clinical administration, it may be preferable to avoid use of chloroform when the cationic amphiphile and  
20 the co-lipid are prepared together. An alternate method to produce such compositions is suggested using formulation of amphiphile 67 (N<sup>4</sup>-spermine cholesteryl carbamate, Figure 1A) as the example.

The cationic amphiphile, the neutral co-lipid DOPE, and PEG(5000)-DMPE are weighed into vials, and each is dissolved in t-butanol:water 9:1 with  
25 vortexing, followed by transfer to a single volumetric flask. An appropriate amount of each lipid is selected to obtain a molar ratio of cationic amphiphile to DOPE to DMPE-PEG of 1 : 2 : 0.05. The resultant solution is vortexed, and

further diluted as needed with t-butanol:water 9:1, to obtain the desired concentration. The solution is then filtered using a sterile filter (0.2 micron, nylon).

One mL of the resultant filtered 1 : 2 : 0.05 solution is then pipetted into  
5 individual vials. The vials are partially stoppered with 2-leg butyl stoppers and placed on a tray for lyophilization. The t-butanol:water 9:1 solution is removed by freeze drying over 2 to 4 days at a temperature of approximately -5°C. The lyophilizer is then backfilled with argon that is passed through a sterile 0.2 micron filter. The stoppers are then fully inserted into the vials, and the vials are  
10 then crimped shut with an aluminum crimp-top. The vials are then maintained at -70°C until use .

Example 7- Further Enhancements in Plasmid Design for Gene Therapy :

Replicating Episomal Plasmids

15 Although the above design features substantially enhance the performance of available plasmids, further modifications are desirable in order that therapeutic compositions comprising such plasmids and cationic amphiphiles have optimal performance for gene therapy.

It is desirable that plasmids for gene therapy also be able to replicate in the  
20 cells of patients, since continued presence of the plasmid will provide correction of the genetic defect (in the case of cystic fibrosis, lack of functioning CFTR protein in the cell membrane of lung epithelial cells or other cells) over an extended period of time. There is concern that plasmids representative of the current art (that is, those that cannot replicate in the targeted cells of a patient)  
25 may be degraded after only a relatively short period of maintenance in the patient, thus requiring excessive repeat administrations.

Long term correction could perhaps be achieved using a vector designed to integrate into chromosomes in the patient's targeted cells (for example, vectors patterned on retrovirus). Such a strategy, however, involves risks including (1) that the vector will integrate into an essential region of a chromosome, or (2) that the vector will integrate adjacent to an oncogene and activate it.

Accordingly, it would be desirable to provide for continued maintenance of gene therapy vectors (plasmids) in target cells by other means. One such strategy is to construct a plasmid capable of being maintained separately in the nucleus of a target cell, and that is also able to replicate there (i.e. an episome).

Plasmids provided according to this aspect of the invention can be constructed as follows. It has been determined (C. McWhinney et al., Nucleic Acids Research, 18, 1233-1242, 1990) that the 2.4 kb HindIII-XhoI fragment that is present immediately 5' to exon 1 of the human c-myc gene contains an origin of replication. The fragment was then cloned into a plasmid that if transfected into HeLa cells was shown to persist therein for more than 300 generations under drug selection. Replication was shown to be semiconservative (C. McWhinney et al.). Although approximately 5% of the plasmid population was lost per cell generation without drug selection in those experiments, this result nonetheless demonstrates substantial stabilization would be of benefit with respect to the design of therapeutic plasmids for gene therapy.

Accordingly, in one example of a replicating episomal vector, a variant of pCF1-CFTR (or pCF1-CAT) can be constructed in which a copy of the 2.4 kb HindIII-XhoI fragment is placed just 5' to the CMV enhancer/promoter region of the pCF1 backbone. Alternatively, between 2 and about 4 - in tandem - copies of the 2.4 kb fragment may be similarly positioned. The increase in plasmid size that results from insertion of the 2.4 kb fragment (or multiple copies thereof) is

predicted to provide an additional benefit, that is, to facilitate plasmid unwinding, thus facilitating the activity of DNA polymerase.

Use of this origin of replication, or multiple copies thereof, allows the resultant plasmid to replicate efficiently in human cells. Other DNA sequences  
5 containing other origins of replication may also be used (for example, as found in the human  $\beta$ -globin gene, or the mouse DHFR gene.

A plasmid that can be constructed according to this aspect of the invention and containing the cytomegalovirus promoter and enhancer, an intron, the CFTR cDNA, the bovine growth hormone polyadenylation signal, the kanamycin  
10 resistance transposon Tn903, and 4 copies of the 2.4 kb 5' flanking region of the human c-myc gene is shown in Figure 20.

Example 8- Further Enhancements in Plasmid Design for Gene Therapy :

15 Use of Cytokine Promoters to Modulate Expression  
of Transgenes in Gene Therapy

Chronic inflammation is associated with numerous of the disease states that can be treated by gene therapy. Representative of such disease states are cystic fibrosis (using CFTR), bronchitis, adult respiratory distress syndrome  
20 (using alpha-1 antitrypsin), and metastatic cancers (through upregulation of p53, TIMP-1, and TIMP-2). Inflammatory conditions typically involve many interrelated processes (for example, involvement by many types of immune system cells and liver proteins), whereby the body attempts to heal a damaged or infected tissue. However, chronic inflammation which persists as a result of an  
25 unresolved condition may lead to permanent tissue damage, as is the case with respect to lung tissue affected by cystic fibrosis and associated and unresolved lung infections. In fact, permanent damage to the lung tissue of cystic fibrosis patients is a leading cause of their mortality. It would be desirable to provide

gene therapy in such a manner as to treat inflammatory conditions associated with the targeted disease state.

Accordingly, a further aspect of the present invention involves construction of gene therapy vectors in which the therapeutic transgene is placed  
5 under control of an RNA polymerase promoter from a cytokine gene (or a gene that encodes another similar regulatory protein) such as, for example, the promoter for any of interleukin 2, interleukin 8, interleukin 1, interleukin 11, interleukin 6, endothelin -1, monocyte chemoattractant protein -1, IL-1ra (receptor agonist), or for GM-CSF.

10 Cytokines may be defined as hormone-like intercellular signal proteins that are involved in regulation of cell proliferation, differentiation, and function, such as concerning haematopoiesis and lymphopoiesis. The interleukins are a particular group of cytokines having promoters that are useful in the practice of the invention. The interleukins are proteins, typically of unrelated origin, which  
15 act as intercellular signals mediating reactions between immunoreactive cells. However, it is understood that many "interleukins" have effects upon additional cell types including endothelial cells, epithelial cells, and fibroblasts.

Since the concentration of many cytokines is upregulated at an affected site in response to the level of inflammation that is present, gene therapy vectors  
20 can be designed wherein the level of therapeutic transgene expressed therefrom is determined, in part, by the level of inflammation present. There follows hereafter description of how such vectors are designed using primarily properties of the interleukin 8 gene as an example.

It has been determined that numerous biologically active molecules are  
25 present in tissues at concentrations thereof that increase with the severity of an inflammatory condition (for example, tumor necrosis factor "TNF" and potentially transcription factors such as NF-kB, AP-1, NF-IL6 and octamer binding protein).

It has also been determined that interleukin 8, a polypeptide of 8,500 MW, is upregulated by inflammation and acts as a potent chemoattractant for T lymphocytes and neutrophil cells that are themselves involved in the inflammation response. The interleukin 8 gene is regulated primarily at the transcriptional level, and it has also been determined (H. Nakamura et al., Journal of Biological Chemistry , 266, 19611-19617, 1991) that TNF can increase interleukin 8 transcription by more than 30 -fold in vitro in bronchial epithelial cells. Accordingly, there follows description of gene therapy vectors which take advantage of the above.

10           A plasmid can be constructed that is substantially similar to pCF1, that is, derived from a pUC plasmid containing a bacterial-derived origin of replication and a gene conferring resistance to kanamycin. The resultant plasmid contains also, in sequence, a CMV enhancer, a promoter, a hybrid intron, a cDNA sequence encoding CFTR, and the bovine growth hormone polyadenylation signal. As RNA polymerase promoter there is selected the -335 to +54 region of the interleukin 8 promoter. This region gave the highest ratio in terms of promoter activity plus TNF over minus TNF (Nakamura, 1991)

Such a plasmid has particularly valuable performance attributes. As inflammation increases in a cystic fibrosis-affected lung (and therefore the need to treat the lung with gene therapy also increases), the concentration of various inflammation-related molecules ( such as TNF) will increase. By placing the CFTR-encoding cDNA of the therapeutic plasmid under the control of a transcriptional promoter (that of interleukin 8, for example) that is itself sensitive to the concentration of inflammation-related substances in contact with the cell, the promoter will function as a natural gene switch such that the amount of beneficial CFTR transcription will be tailored to the amount of inflammation. As

aforementioned, RNA polymerase promoter sequences derived from the other aforementioned genes are also useful in the practice of the invention.

#### Example 9 Intravenous Delivery of Transgenes

For some disease states, such as cystic fibrosis, it is desirable to  
5 deliver transgenes to the lung. Delivery by aerosol is the most direct  
approach to achieve this goal. However, given the difficulties inherent with  
the delivery of an aerosol together with the potential need to target organs  
other than the lung (for example, the pancreas for cystic fibrosis), it is  
important to evaluate the feasibility of lung delivery using non-aerosol  
10 delivery formats. Accordingly, intravenous delivery of a reporter  
transgene was performed using a mouse model and the feasibility of  
intravenous organ targeting was assessed. A comparison was made of  
feasibility of delivery to the lung and the heart.

The reporter plasmid pCF-1 CAT (Example 4) was used and was  
15 purified to minimize endotoxin (<1 EU/mg pDNA), and also chromosomal  
DNA contamination (< 2%). Amphiphile No. 53 (1:1 with DOPE) / DNA  
complex was prepared according to the procedures of Example 3. The  
amphiphile was provided as the free base, the plasmid was prepared as a  
sodium salt in water, and the DOPE was provided in zwitterionic form.

20 The animal model was the BALB/c mouse. Females 6-8 weeks old  
weighing 16-18 g were injected intravenously using the tail vein, using 5  
animals per group. The volume of lipid:pDNA complex used was 100 µl in  
all experiments. Unless noted otherwise, mice were sacrificed 48 h  
following administration of the complex. Organs were frozen immediately  
25 on dry ice to store for subsequent analysis.

Expression of chloramphenicol acetyl transferase (CAT) was  
quantitated using a radiochemical assay for CAT enzymatic activity.

Organs were weighed and homogenized on ice in a lysis buffer containing protease inhibitors. The lysate was freeze-thawed 3X, centrifuged, and heated to 65°C to inactivate deacetylases before adding it to a reaction mixture containing  $^{14}\text{C}$ -chloramphenicol. After an incubation at 37°C, the  
5 mixture was extracted with ethyl acetate, concentrated, spotted onto TLC plates and eluted with  $\text{CHCl}_3/\text{MeOH}$ . Spots corresponding to the acylated reaction products were quantitated (Betagen) and converted to ng CAT activity using authentic CAT standards.

It was surprisingly determined that targeting to the heart could be  
10 substantially improved by altering the molar ratio (at a constant DNA concentration of 0.9 mM, measured as nucleotide) of amphiphile/DNA in the therapeutic composition. This information is of value in connection with gene therapy for the heart, such as for coronary disease. However, targeting to the lung remained relatively constant over a range of  
15 amphiphile/DNA ratios, all at constant DNA concentration (Figure 21).

At molar ratios of less than about 0.5, the organ distribution was found to be strongly weighted toward the lung. At this molar ratio, the zeta potential of the complex is negative (about -30 mV) due, in part, to excess negative charge from the DNA relative to the amphiphile. At an  
20 amphiphile/DNA ratio of 1.25, however, where the complex has a positive zeta potential (about +30mV), organ distribution was remarkably altered and substantial expression was found in the heart (Figure 21).

Zeta potentials of the samples can be measured (using typically 5 measurements per sample) employing a Malvern Zetasizer 4 (Malvern  
25 Instruments, Southborough, MA.) and a zeta cell (AZ-104 cell, Malvern Instruments Co.). Dried lipid films containing the cationic lipid and DOPE are hydrated in distilled water ( $\text{dH}_2\text{O}$ ). DNA typically should be diluted to

a concentration of about 300  $\mu\text{M}$  in  $\text{dH}_2\text{O}$ . The DNA solution (1.5 mL) can then be added to an equal volume of cationic lipid vesicles and incubated at room temperature for 10 min. Enough NaCl (for example, 4 mM stock) may be added to result in a final concentration of 1 mM NaCl. If necessary, the  
5 sample can be diluted further with 1 mM NaCl (to maintain a photomultiplier signal below 4000 counts per second), and distilled water can be used in place of the NaCl solutions.

According to this aspect of the invention, amphiphiles No. 53 and No. 67 are among those preferred for use in intravenous targeting of the  
10 heart, as are many other amphiphiles selected from Groups I and II.

Example 10- Additional Experimental Procedures

(A) Additional synthesis procedure for N<sup>4</sup>-spermine cholesteryl carbamate, amphiphile No. 67

(Synthesis of N<sup>1</sup>N<sup>12</sup>-diCbz-spermine di-HCl salt)

15 Benzylchloroformate (15 mL, 105 mmol) was dissolved in methylene chloride (335 mL) and placed in a three neck flask under a nitrogen atmosphere. Imidazole (14 g, 206 mmol) was dissolved in methylene chloride (200 mL). The three neck flask was cooled to 0-2 °C using an ice-water bath and the imidazole solution was added gradually over 30 min. The cooling bath was removed and  
20 the mixture stirred at room temperature for 1 hour. Methylene chloride (250 mL) and aqueous citric acid (10%, 250 mL) were added to the mixture. The layers were separated and the organic layer was washed with aqueous citric acid (10%, 250 mL). The organic fraction was dried over magnesium sulfate and concentrated *in vacuo*. The resulting oil was vacuum dried for 2 hours at ambient  
25 temperature. To the oil was added dimethylaminopyridine (530 mg, 4.3 mmol) and methylene chloride (250 mL). The mixture was cooled to 0-2 °C and kept under a nitrogen atmosphere. A solution of spermine (10g, 49 mmol) in

methylene chloride (250 mL) was added gradually over 15 minutes, maintaining a reaction temperature of 0-2 °C. The reaction mixture was stirred overnight at ambient temperature and then concentrated *in vacuo*. To the resulting material was added 1M hydrochloric acid (67 mL) and methanol (400 mL). The solution  
5 was cooled overnight at 4 °C yielding a white precipitate. The precipitate was isolated by vacuum filtration using Whatman #1 filter paper. The N<sup>1</sup>,N<sup>12</sup>-diCbz-spermine di HCl salt (13.38g, 24.7 mmol, 50% yield) thus obtained was dried under vacuum at ambient temperature for 17 hours.

(Synthesis of N<sup>1</sup>,N<sup>12</sup>-diCbz-N<sup>4</sup>-spermine cholesteryl carbamate)

10 N<sup>1</sup>,N<sup>12</sup>-diCbz-spermine di HCl salt (13.38g, 24.7 mmol) was dissolved in a chloroform, methanol and water mixture in the ratio 65:25:4 (940 mL). The solution was stirred at room temperature and cholesteryl chloroformate (11g, 24.5 mmol) was added. The solution was stirred at ambient temperature for 1.5 hours and then diluted with 1M sodium hydroxide solution (165 mL). The organic and  
15 aqueous layers were separated and the organic layer containing the product was washed with water (110 mL). The organic fraction was dried over sodium sulfate, concentrated *in vacuo* and vacuum dried. The crude oil was purified by chromatography using silica gel (60Å, 1 Kg). The silica was packed in 10% MeOH / CHCl<sub>3</sub> and the column was eluted with 25% MeOH / CHCl<sub>3</sub>. Fractions of 900  
20 mL were collected and analyzed by thin layer chromatography. Fractions containing the product (R<sub>f</sub> = 0.5 in 20% MeOH / CHCl<sub>3</sub>) were combined and concentrated *in vacuo*. The resulting oil was dried under vacuum for 17 hours to give 8.5g (9.67 mmol, 39% yield) of product.

(Synthesis of N<sup>4</sup>-spermine cholesteryl carbamate)

N<sup>1</sup>N<sup>12</sup>-diCbz-N<sup>4</sup>-spermine cholesteryl carbamate (8.5g, 9.67 mmol) was dissolved in 200 mL of acetic acid and 1.66 g of 10% Pd on carbon was added. The solution was purged with nitrogen and stirred under hydrogen at atmospheric pressure. The hydrogen was supplied to the reaction flask using a balloon filled with hydrogen gas. The hydrogenolysis was allowed to proceed for 3 hours. The reaction mixture was filtered through Whatman #1 filter paper and the catalyst was washed with 250 mL of 10% acetic acid in ethyl acetate. The filtrate was concentrated *in vacuo* to give a residue, coevaporation with chloroform aids removal of the acetic acid. To the crude product was added 1M sodium hydroxide solution (400 mL) and the solution was extracted three times with 10% MeOH / CHCl<sub>3</sub> (700 mL). The combined organic fractions were washed with water (600 mL) and dried over sodium sulfate. The solution was filtered, concentrated *in vacuo* and vacuum dried at ambient temperature for 48 hours. The crude material was purified by chromatography on silica gel (500 g). The column was packed in 40:25 MeOH : CHCl<sub>3</sub> and eluted with 40:25 MeOH : CHCl<sub>3</sub> and then 40:25:10 MeOH : CHCl<sub>3</sub> : NH<sub>4</sub>OH. The fractions collected were analyzed by thin layer chromatography and the product containing fractions were combined and concentrated *in vacuo* (the evaporation was assisted by the addition of *iso*-propanol in order to azeotrope the residual water). The material was vacuum dried at ambient temperature for 48 hours to give N<sup>4</sup>-spermine cholesteryl carbamate (4g, 6.5 mmol, 67% yield).

(B) N<sup>4</sup>-(N'-cholesteryl carbamate glycineamide)-spermine (amphiphile No. 91)

N-t-BOC-glycine-N-hydroxysuccinimide ester (0.5 g, 1.83 mmol) was added to a solution of diCbz-spermine · 2HCl (1.0 g, 1.94 mmol) and N,N-diisopropylethylamine (0.3 mL, 1.72 mmol) in 65/ 25/ 4 chloroform/ methanol/ water (50 mL). The solution was stirred overnight at room temperature.

Analysis of the reaction by TLC (20% methanol/ chloroform) indicated the presence of a new spot. The reaction was washed first with 1M NaOH (10 mL) then with H<sub>2</sub>O (10 mL). The organic layer was separated, dried over sodium sulfate, vacuum filtered, and reduced *in vacuo* to an oil. The crude material was  
5 purified by flash column chromatography (85 g silica gel) eluting with 20% methanol/ chloroform. The desired product was isolated and characterized by <sup>1</sup>H NMR as N<sup>1</sup>,N<sup>12</sup>-diCbz-N<sup>4</sup>-(N'-t-BOC-glycineamide)-spermine (402 mg, 0.65 mmol, 35%).

Benzyl chloroformate (100 mg, 0.58 mmol) was added to a solution of  
10 N<sup>1</sup>,N<sup>12</sup>-diCbz-N<sup>4</sup>-(N'-t-BOC-glycineamide)-spermine (220 mg, 0.354 mmol) and triethylamine (4 drops) in methylene chloride (20 mL). The reaction was stirred overnight at room temperature. Analysis of the reaction by TLC (20% methanol/ chloroform) indicated the presence of a new, higher running spot. The reaction was quenched by the addition of 1M HCl (5 mL). The organic layer was isolated,  
15 washed with H<sub>2</sub>O (5 mL), dried over sodium sulfate, filtered, and reduced *in vacuo*.

The resulting crude material was dissolved in chloroform (30 mL) and anhydrous HCl gas was bubbled through the solution for 2 hours. Analysis of the reaction by TLC (10% methanol/ chloroform) indicated the complete  
20 disappearance of the starting material. The reaction was purged with dry nitrogen, and washed with 1M NaOH (2 x 10 mL) and dH<sub>2</sub>O (10 mL). The organic layer was isolated, dried over sodium sulfate, filtered, and reduced *in vacuo* to give N<sup>1</sup>,N<sup>9</sup>,N<sup>12</sup>-triCbz-N<sup>4</sup>-glycineamide-spermine (219 mg, 0.33 mmol, 93% yield for two steps).

25 Cholesteryl chloroformate (148 mg, 0.33 mmol) was added to a solution of N<sup>1</sup>,N<sup>9</sup>,N<sup>12</sup>-triCbz-N<sup>4</sup>-glycineamide-spermine (219 mg, 0.33 mmol) and triethylamine (0.3 mL, 2.15 mmol) in methylene chloride (30 mL). The reaction

was stirred at room temperature for 3 hours. The reaction was washed with H<sub>2</sub>O (10 mL). The organic layer was separated, dried over sodium sulfate, filtered, and reduced *in vacuo*. The crude material was purified by flash column chromatography (30 g silica gel) eluting with 65% ethyl acetate/ hexanes. The  
5 desired product was isolated and characterized by <sup>1</sup>H NMR as N<sup>1</sup>,N<sup>9</sup>,N<sup>12</sup>-triCbz-N<sup>4</sup>-(N'-cholesteryl carbamate glycineamide)-spermine (221 mg, 0.2 mmol, 62% yield).

N<sup>1</sup>,N<sup>9</sup>,N<sup>12</sup>-tri-Cbz-N<sup>4</sup>-(N'-cholesteryl carbamate glycineamide)-spermine (221 mg, 0.2 mmol) was stirred with 10% Pd/C (50 mg) in glacial acetic acid (10  
10 mL) under a hydrogen atmosphere for 2.5 hours. Analysis of the reaction by TLC (65% ethyl acetate/ hexanes) indicated the complete disappearance of the starting material. The flask was purged with nitrogen and the catalyst was removed by vacuum filtration through filter paper rinsing with 10% acetic acid/ ethyl acetate (20 mL). The filtrate was reduced *in vacuo* to an oil which was  
15 dissolved in 10% methanol/ chloroform (100 mL) and washed with 1M NaOH (20 mL) and H<sub>2</sub>O (15 mL). The organic layer was separated, dried over sodium sulfate, filtered, and reduced *in vacuo*. The isolated product was characterized by <sup>1</sup>H NMR as N<sup>4</sup>-(N'-cholesteryl carbamate glycineamide)-spermine (128 mg, 0.19 mmol, 95% yield).

20 (C) Synthesis of N<sup>4</sup>-spermidine-2,3-dilauryloxypropylamine, amphiphile No. 94.

2,3 Dimyristoylglycerol (600 mg, 1.4 mmol) was dissolved in pyridine and the solution cooled to 0°C. The solution was stirred under a nitrogen atmosphere and p-toluenesulfonyl chloride (300 mg, 1.57 mmol) was added. The solution was allowed to warm to room temperature and was then stirred  
25 overnight at ambient temperature. To the solution was added hydrochloric acid (2.5M, 20 mL) and the solution was extracted three times with methylene chloride (25 mL). The combined organic extracts were dried over sodium sulfate,

filtered and concentrated in *vacuo* to give a crude oil. The oil was purified by flash chromatography (50g of silica gel, 60Å) eluting with 5% ethyl acetate / hexane. The oil obtained by flash chromatography was dried under high vacuum at ambient temperature to give 2,3-Dimyristoylglycerol-tosylate(630 mg, 5 77% yield).

2,3-Dimyristoylglycerol-tosylate (300 mg, 0.51 mmol) and N<sup>1</sup>,N<sup>8</sup>-diCbz-spermidine (1.5g, 3.6 mmol) were dissolved in toluene (15 mL). The solution was stirred under a nitrogen atmosphere and heated at reflux (110°C). The reaction was heated for 5 days at reflux temperature. The reaction was cooled to room 10 temperature and then filtered through Whatman #1 filter paper. The filtrate was concentrated in *vacuo*. The residue was dissolved in chloroform (50 mL) and washed with sodium hydroxide solution (1 M, 10 mL) and water (10 mL). The organic fraction was dried over sodium sulfate, filtered and concentrated in *vacuo*. The crude material was purified by flash chromatography (30g silica gel, 60Å) 15 eluting with 5% methanol / chloroform. The product containing fractions were concentrated in *vacuo*. The material was purified by a second flash chromatography column (20 g silica, 60Å) eluting with 50% ethyl acetate / hexane. Chromatography gave, after drying the product under high vacuum at ambient temperature, N<sup>4</sup>-(N<sup>1</sup>,N<sup>8</sup>)-diCbz-spermidine-2,3- 20 dilauryloxypropylamine, as an oil (142 mg, 35% yield).

N<sup>4</sup>-(N<sup>1</sup>,N<sup>8</sup>)-diCbz-spermidine-2,3-dilauryloxypropylamine (142 mg, 0.18 mmol) in glacial acetic acid (5 mL) was stirred with 10% Pd/C (50 mg) under a hydrogen atmosphere, for 2 hours. The catalyst was removed by vacuum filtration through Whatman #1 filter paper. The catalyst was washed with ethyl 25 / acetate hexane (10%, 10 mL). The filtrate was concentrated in *vacuo* and dried for 2 hours under high vacuum. To the residue was added sodium hydroxide solution (1 M, 8 mL) and the solution was extracted three times with methanol /

chloroform (10%, 20 mL). The combined organic extracts were dried over sodium sulfate, filtered and concentrated in *vacuo* to give after drying under high vacuum N<sup>4</sup>-spermidine-2,3-dilauryloxypropylamine (52 mg, 52% yield).

(D) Synthesis of N<sup>4</sup>-spermine-2,3-dilauryloxypropylamine, amphiphile No. 102

5 N<sup>1</sup>,N<sup>12</sup>-diCbz-spermine (0.87g, 1.85 mmol) and 2,3-dimyristoylglycerol-tosylate (280mg, 0.48 mmol) were dissolved in toluene (25 mL) and heated at reflux temperature (110°C) for 3 days. The solution was concentrated in *vacuo* and the resulting material was purified by flash chromatography (30g silica gel, 60Å) eluting with 10% methanol /chloroform. The material isolated was  
10 dissolved in methanol / chloroform (10%, 85 mL) and washed twice with sodium hydroxide solution (1 M, 15 mL) and water (10 mL). The organic fraction was dried over sodium sulfate, filtered and concentrated in *vacuo* . The material was dried under high vacuum overnight, at ambient temperature, to yield N<sup>4</sup>-  
(N<sup>1</sup>,N<sup>12</sup>-diCbz-spermine)-2,3-dilauryloxypropylamine (180 mg, 43 % yield).

15 N<sup>4</sup>-(N<sup>1</sup>,N<sup>12</sup>-diCbz-spermine)-2,3-dilauryloxypropylamine (180 mg, 0.2 mmol) in glacial acetic acid (10 mL) was stirred with 10% Pd/C (50 mg) under a hydrogen atmosphere, for 3 hours. The catalyst was removed by vacuum filtration through Whatman #1 filter paper. The catalyst was washed with ethyl / acetate hexane (10%, 30 mL). The filtrate was concentrated in *vacuo* and dried  
20 for 2 hours under high vacuum. To the residue was added methanol / chloroform (10%, 85 mL) and the organic layer was washed twice with sodium hydroxide solution (1 M, 15 mL) and water (10 mL). The organic fraction was dried over sodium sulfate, filtered and concentrated in *vacuo* to give after drying under high vacuum N<sup>4</sup>-spermine-2,3-dilauryloxypropylamine (50 mg,  
25 40% yield).

Example 11- Expression of a Secreted Protein from Vascular Tissue

Human secreted alkaline phosphatase (SEAP) was detected in the serum of BALB/c mice following intravenous administration of a plasmid containing an encoding cDNA. Following generally the procedures of Examples 1 and 3, a cationic amphiphile plasmid composition was prepared. A cDNA encoding  
5 sequence for human SEAP was placed in pCF1 plasmid (see Example 4). The transfecting composition was prepared to contain 0.75 mM of amphiphile No. 67, 1.5 mM DOPE, and 2mM (as nucleotide) of pCF1 plasmid. Thus the amphiphile/DNA ratio was 1:4 thereby providing a negative zeta potential.

As demonstrated in Figure 22, substantial expression of SEAP protein was  
10 detected in the serum of BALB/c mice following tail vein administration of the amphiphile/DNA complex. pCF1 plasmid with the encoding cDNA was used as control and error bars are shown. Similar results were achieved using amphiphile No. 53 (see Example 3 for preferred procedures). The amphiphile/DNA complex was provided as 0.5 mM of amphiphile No. 53, 0.5  
15 mM DOPE, and 2mM (as nucleotide) of pCF1 plasmid.

---

The above descriptions of preferred embodiments of the invention have been presented to illustrate the invention to those skilled in the art. They are not  
20 intended to limit the invention to the precise forms disclosed.

Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.


The reference to any prior art in this specification is not, and should not be taken as, an acknowledgement or any form of suggestion that that prior art forms part of the common general knowledge in Australia.



## CLAIMS

1. A method of delivering a biologically active molecule to blood *in vivo*, comprising:  
administering a composition to vascular tissue of a mammal wherein said composition comprises:

- 5 (1) a nucleic acid molecule coding for said biologically active molecule; and  
(2) a cationic amphiphile effective for delivering said biologically active molecule into a blood vessel of said mammal, said cationic amphiphile being selected from:



10 wherein:

Z is a steroid;

X is a carbon atom or a nitrogen atom;

Y is a linking group, or Y is absent;

R<sup>3</sup> is H, or a saturated or unsaturated aliphatic group;

15 R<sup>1</sup> is -NH-, an alkylamine, or a polyalkylamine;

R<sup>4</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>2</sup> is -NH-, an alkylamine, or a polyalkylamine

and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup>, except that both R<sup>1</sup> and R<sup>2</sup> cannot be -NH-; and optionally


(3) a co-lipid.



2. A method of delivering a biologically active molecule to blood *in vivo*, comprising:  
administering a composition to vascular tissue of a mammal wherein said composition comprises:

(1) a nucleic acid molecule coding for said biologically active molecule; and

5 (2) a cationic amphiphile effective for delivering said biologically active molecule into a blood vessel of said mammal, said cationic amphiphile being selected from:



wherein:

10 Z is a steroid;

X is a carbon atom or a nitrogen atom;

Y is a linking group, or Y is absent;

R<sup>3</sup> is an amino group, a substituted amino acid, H or alkyl;

R<sup>1</sup> is -NH-, an alkylamine, or a polyalkylamine;

15 R<sup>4</sup> is an amino group, a substituted amino acid, H or alkyl;

R<sup>2</sup> is -NH-, an alkylamine, or a polyalkylamine

and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup>, except that both R<sup>1</sup> and R<sup>2</sup> cannot be -NH-; and optionally

(3) a co-lipid.


20 3. A method of delivering a biologically active molecule to blood *in vivo*, comprising:



administering a composition to vascular tissue of a mammal wherein said composition comprises:

(1) a nucleic acid molecule coding for said biologically active molecule; and

(2) a cationic amphiphile effective for delivering said biologically active molecule into a blood vessel of said mammal, said cationic amphiphile being selected from:



wherein:

Z is an alkylamine or a dialkylamine, linked by the N-atom thereof, to Y (or directly to X, if Y is absent), wherein if Z is a dialkylamine, the alkyl groups thereof can be the same or different;

X is a carbon atom or a nitrogen atom;

Y is selected from  $-\text{CH}_2-$ ;  $-(\text{CH}_2)_2-$ ;  $-(\text{CH}_2)_3-$ ;  $>\text{C}=\text{O}$ ;  $>\text{C}=\text{S}$ ;  $-(\text{CH}_2)_n\text{-NH-(C=O)-}$  where n is 1, 2, 3, or 4; and  $-\text{CH}_2\text{-(C=O)-}$ ; or Y is absent;

$R^3$  is H, or a saturated or unsaturated aliphatic group;

$R^1$  is  $-\text{NH-}$ , an alkylamine, or a polyalkylamine;

$R^4$  is H, or a saturated or unsaturated aliphatic group;

$R^2$  is  $-\text{NH-}$ , an alkylamine, or a polyalkylamine

and wherein  $R^1$  is the same or is different from  $R^2$ , except that both  $R^1$  and  $R^2$  cannot be  $-\text{NH-}$ ; and optionally

(3) a co-lipid.


4. A method of delivering a biologically active molecule to blood *in vivo*, comprising:



administering a composition to vascular tissue of a mammal wherein said composition comprises:

(1) a nucleic acid molecule coding for said biologically active molecule; and

(2) a cationic amphiphile effective for delivering said biologically active molecule into a blood vessel of said mammal, said cationic amphiphile being selected from:



wherein:

(A) and (B) are independently O, NH, or S;

10 R<sup>5</sup> and R<sup>6</sup> are independently alkyl or acyl groups and may be saturated or contain sites of unsaturation;

(C) is selected from -CH<sub>2</sub>-, >C=O, and >C=S;

(D) is a linking group or is absent;

(E) is a carbon atom or a nitrogen atom;

15 R<sup>3</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>1</sup> is -NH-, an alkylamine, or a polyalkylamine;

R<sup>4</sup> is H, or a saturated or unsaturated aliphatic group;

R<sup>2</sup> is -NH-, an alkylamine, or a polyalkylamine



and wherein R<sup>1</sup> is the same or is different from R<sup>2</sup>, except that both R<sup>1</sup> and R<sup>2</sup> cannot be -NH-; and optionally

(3) a co-lipid.

5 5. A method according to Claim 4, wherein said linking group (D) is -NH(C=O)- or -O(C=O)-.

6. A method according to Claim 1, wherein said composition further comprises a fatty acid species of PEG-PE.

7. A method according to Claim 1, wherein said biologically active molecule is a DNA molecule comprising an encoding sequence for a protein that is secreted from cells.

10 8. A method according to Claim 2, wherein said composition further comprises a fatty acid species of PEG-PE.

9. A method according to Claim 2, wherein said biologically active molecule is a DNA molecule comprising an encoding sequence for a protein that is secreted from cells.

15 10. A method according to Claim 3, wherein said composition further comprises a fatty acid species of PEG-PE.

11. A method according to Claim 3, wherein said biologically active molecule is a DNA molecule comprising an encoding sequence for a protein that is secreted from cells.

12. A method according to Claim 4, wherein said composition further comprises a fatty acid species of PEG-PE.


20 13. A method according to Claim 4, wherein said biologically active molecule is a DNA molecule comprising an encoding sequence for a protein that is secreted from cells.

14. The method according to any one of Claims 1-13, substantially as herein before described with reference to the figures and/or examples.

DATED this 20th day of July 2001


**Genzyme Corporation**  
DAVIES COLLISON CAVE  
Patent Attorneys for the applicant






**53**

N<sup>4</sup>-spermidine cholesteryl carbamate




**71**

N<sup>4</sup>-spermidine dihydrocholesteryl carbamate




**61**

N<sup>4</sup>-spermidine cholesteryl carbamate  
di HCl salt




**72**

N<sup>4</sup>-spermidine lithocholic carbamate  
methyl ester




**65**

N<sup>4</sup>-spermidine-7 dehydro cholesteryl  
carbamate




**75**

N<sup>1</sup>,N<sup>8</sup>-Bis (3-aminopropyl)- N<sup>4</sup> -  
spermidine cholesteryl carbamate




**67**

N<sup>4</sup>-spermine cholesteryl carbamate




**69**

N,N bis (3-aminopropyl) cholesteryl  
carbamate



**78**


N(N<sup>4</sup>-3-aminopropylspermidine)  
cholesteryl carbamate




**70**

N,N bis (6-aminohexyl) cholesteryl  
carbamate


**FIG. 1 A**





N,N-Bis(4-aminobutyl) cholesteryl carbamate




N,N-Bis(N'-3-aminopropyl-N''-4-aminobutyl) cholesteryl carbamate




N<sup>4</sup>-spermidine cholesteryl urea





N<sup>4</sup>-spermine cholesteryl urea




N<sup>4</sup> Spermidine cholesteryl carboxamide



N<sup>4</sup>-spermidine dihydro cholesteryl urea



N<sup>4</sup>-spermine dihydro cholesteryl urea



N-[N<sup>1</sup>,N<sup>4</sup>,N<sup>8</sup>-Tris(3-aminopropyl) spermidine] cholesteryl carbamate

FIG. 1B




FIG. 1C






FIG. 1D


4/23



Ergosterol (double bonds as shown)  
 Ergosterol B1 ( $\Delta 8, 9; \Delta 14, 15; \Delta 22, 23$ )  
 Ergosterol B1 ( $\Delta 6, 7; \Delta 8, 14; \Delta 22, 23$ )  
 Ergosterol B1 ( $\Delta 7, 8; \Delta 14, 15; \Delta 22, 23$ )  
 Lumisterol ( $9\beta$ -H isomer of ergosterol)





Cholic Acid  $r^1, r^2 = OH$   
 Desoxycholic Acid  $r^1 = H, r^2 = OH$   
 Chenodesoxycholic Acid  $r^1 = OH, r^2 = H$   
 Lithocholic Acid  $r^1, r^2 = H$



Androsterone

**FIG. 2**


5/23




Cholic Acid Derivative  $r^1, r^2 = \text{OH}$   
Desoxycholic Acid Derivative  $r^1 = \text{H}, r^2 = \text{OH}$   
Chenodesoxycholic Acid Derivative  $r^1 = \text{OH}, r^2 = \text{H}$   
Lithocholic Acid Derivative  $r^1, r^2 = \text{H}$

FIG. 3

**Transacylation Reaction.**




**FIG. 4**




N<sup>1</sup>,N<sup>8</sup>-Bis(arginine carboxamide)-  
N<sup>4</sup>-spermidine cholesteryl carbamate

FIG. 5


8/23




N,N-dioctadecyllysineamide diHCl salt




**60**



N<sup>1</sup>,N<sup>1</sup>-dioctadecyl-1,2,6-triaminohexane tri HCl salt







N,N-dioctadecyllysineamide




**FIG. 6**



9/23





1-(N<sup>4</sup>-spermine)-2,3-dilaurylglycerol carbamate




N<sup>4</sup>-spermidine-2,3-dilauryloxypropylamine



N<sup>4</sup>-spermine-2,3-dilauryloxypropylamine



**110**



**111**

FIG. 7

10/23




FIG. 8



12/23




FIG. 10




FIG. 11

14/23




FIG. 12

15/23

<u>Lipid #</u>	<u>Opt. Ratio</u>	<u>In Vitro</u>	<u>In Vivo</u>
53	1\1	4.2	20
65	1\2	2	20
67	1\2	4.8	150
69	2\1	2	2
70	1\1	1	ND
71	1\1	2.4	3
72	1\2	0.7	0.5
75	2\1	5	80
78	1\1	2.3	75
79	1\2	0.8	ND
81	1\2	0.7	ND
82	1\1	3.8	5
83	1\2	4	20
84	1\2	0.4	1
86	1\2	0.5	ND
88			ND
90	1\1	1.8	80
91	1\2	2.1	30
92	1\2	0.5	ND
96	1\2	5	45
101	2\1	1.5	1
104	2\1	0.7	0
106	1\2	1.6	6
107	1\1	1.3	

FIG. 13

16/23


<u>Lipid #</u>	<u>Opt. Ratio</u>	<u>In Vitro</u>	<u>In Vivo</u>
87			
91	1\2	2.1	30
93	1\2	2.7	
95	1\2	3.7	70
97	1\1	0.4	ND
99	1\1	0	ND
100	1\1	0.1	ND
103			

FIG. 14

<u>Lipid #</u>	<u>Opt. Ratio</u>	<u>In Vitro</u>	<u>In Vivo</u>
64	1\1	2.1	1
76	1\1	1	ND
85	1\1	1.5	1
89	1\1	9	15
94	1\1	2	60
98	1\1	1.9	
102	1\2	10	40
105	1\2	3	
110	2\1	0.9	
111			

FIG.15

17/23






FIG. 17

FIG. 18A







FIG. 18B





21/23




FIG. 20

22/23





FIG. 21

23/23

## Expression of SEAP in BALB/c Mice Following Intravenous Administration

**FIGURE 22**