



(86) Date de dépôt PCT/PCT Filing Date: 2006/01/09
(87) Date publication PCT/PCT Publication Date: 2006/07/20
(85) Entrée phase nationale/National Entry: 2007/05/09
(86) N° demande PCT/PCT Application No.: EP 2006/050098
(87) N° publication PCT/PCT Publication No.: 2006/074986
(30) Priorité/Priority: 2005/01/13 (EP05000542.0)

(51) Cl.Int./Int.Cl. *C12N 7/00* (2006.01),
A61K 39/12 (2006.01), *A61P 31/14* (2006.01)

(71) Demandeur/Applicant:
BOEHRINGER INGELHEIM VETMEDICA GMBH, DE


(72) Inventeurs/Inventors:
FETZER, CHRISTIANE, DE;
PESCH, STEFAN, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : VACCINS AMELIORES CONTRE LE SRRP
(54) Title: IMPROVED PRRS VACCINES

(57) **Abrégé/Abstract:**

The present invention is related to improved modified live PRRS vaccines of European genotype and new PRRSV strains which can be used for the manufacture of such vaccines.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 July 2006 (20.07.2006)

PCT

(10) International Publication Number
WO 2006/074986 A2

(51) International Patent Classification: **Not classified**

(21) International Application Number:
PCT/EP2006/050098

(22) International Filing Date: 9 January 2006 (09.01.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
05000542.0 13 January 2005 (13.01.2005) EP

(71) Applicant (*for all designated States except US*):
BOEHRINGER INGELHEIM VETMEDICA GMBH
[DE/DE]; Binger Str. 173, 55216 Ingelheim (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FETZER, Chris-
tiane** [DE/DE]; Tresckowstr. 31, 48163 Münster (DE).
PESCH, Stefan [DE/DE]; Mauritzdahl 6, 48145 Münster
(DE).

(74) Agents: **HAMMANN, Heinz** et al.; Binger Str. 173,
55216 Ingelheim (DE).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *without international search report and to be republished
upon receipt of that report*
- *with sequence listing part of description published sepa-
rately in electronic form and available upon request from
the International Bureau*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: IMPROVED PRRS VACCINES

(57) Abstract: The present invention is related to improved modified live PRRS vaccines of European genotype and new PRRSV strains which can be used for the manufacture of such vaccines.

WO 2006/074986 A2

Improved PRRS Vaccines

5 BACKGROUND OF THE INVENTION

1. TECHNICAL FIELD

The present invention belongs to the field of vaccines against infectious diseases. More
10 particularly, it relates to vaccines against Porcine Reproductive and Respiratory Syndrome (PRRS), a viral disease affecting pigs.

2. BACKGROUND INFORMATION

15 The Porcine Reproductive and Respiratory Syndrome Virus (PRRSV) is a (+)stranded ssRNA virus belonging to the family of Arteriviridae. The virus exists as two genotypes referred to as „US“ and „EU“ type which share about 50% sequence homology (Dea S et al. (2000). Arch Virol 145:659-88), and which can be also distinguished by their immunological properties. Most sequencing information on various isolates is based on the
20 structural proteins, namely the envelope protein GP5 which accounts for only ~ 4 % of the viral genome, while only little is known on the non-structural proteins (nsp).

The nsp2 protein has 1078 amino acids in the EU type prototype Lelystad Virus (LV), and shares only 32 % homology with PRRSV-US (Allende R et al. (1999). J Gen Virol 80:307-
25 15). It induces serum antibodies during natural infection (Oleksiewicz MB et al. (2001a). J Virol 75:3277-90, Oleksiewicz MB et al. (2001b). Vet Microbiol 81:109-25), seems to play a crucial role for viral replication (Snijder EJ et al. (2001). J Gen Virol 82:985-94, van der Meer Y et al. (1998). J Virol 72:6689-98) and is thought to have species-specific functions (de Vries AAF et al. (1997). Seminars in Virology 8:33-47). The nucleotide sequence
30 coding for the nsp2 amino acid sequence is represented in SEQ ID NO: 2. Length variations in the nsp2 coding region have so far only been described for one US-type isolate (Shen S et al. (2000). Arch Virol 145:871-83) and for EU type strains found in the US

(Fang Y et al. Virus Res. 2004 Mar 15;100(2):229-35, Ropp SL et al. J Virol. 2004 Apr;78(7):3684-703.

Isolation of PRRSV and manufacture of vaccines have been described in a number of publications (WO 92/21375 , WO 93/06211, WO93/03760, WO 93/07898, WO 96/36356, EP 0 676 467, EP 0 732 340, EP 0 835 930). There is still the need to provide vaccines with improved characteristics, in particular with respect to efficacy and safety.

BRIEF SUMMARY OF THE INVENTION

10

The present invention is related to improved modified live PRRS vaccines of European genotype and new PRRSV strains which can be used for the manufacture of such vaccines. In particular, the invention provides improved PRRS virus strains that have been deposited with the European Collection of Cell Cultures (ECACC) under the Accession Numbers
15 ECACC 04102703, ECACC 04102702, and ECACC 04102704, or any descendant or progeny of one of the aforementioned strains.

BRIEF DESCRIPTION OF THE DRAWINGS

20 Figure 1: Agarose gel (2% agarose, ethidium bromide stained) of RT-PCR products obtained from the nsp2 region (pos. 2068 – 2713) of a Lelystad-like isolate (lane 1), Porcilis[®] PRRS (lane 2) and negative control (lane 3). Lane 4: 100 bp ladder. The products from Porcilis[®] PRRS (double band at 434 and 416 bp and band at 347 bp) are smaller than the LV-like product (569 bp).

25

Figure 2: nsp2 PCR products obtained from a selection of plaques after 1 passage on Ma104 cell culture.

Figure 3: Nucleotide sequence of one representative plaque of each length variant compared to the nucleotide sequence of the prototype strain Lelystad Virus (LV).
30

Figure 4: nsp2 PCR products obtained from different samples of the two sows showing reproductive failure after Porcilis[®] PRRS vaccination and their offspring.

Figure 5: nsp2 PCR products obtained from clinical samples of piglets vaccinated with Porcilis[®] PRRS at 6 weeks of age. Circle shows samples of pigs diseased on PMWS.

Figure 6 A: Nucleotide sequence of nsp2 PCR products obtained from different field samples showing the same deletion of 222 nucleotides as one group of plaques derived from the Porcilis PRRS vaccine bottle (e.g. plaque P45), compared to the full-length (Lelystad-like) sequence of the Porcilis-derived plaque 36.

Figure 6B: Nucleotide sequence of nsp2 PCR products obtained from different field samples showing the same deletion of 135 nucleotides as one group of plaques derived from the Porcilis PRRS vaccine bottle (e.g. plaque P27), compared to the full-length (Lelystad-like) sequence of the Porcilis-derived plaque 36.

Figure 6 C: Nucleotide sequence of nsp2 PCR products obtained from different field samples showing the same deletion of 153 nucleotides as one group of plaques derived from the Porcilis PRRS vaccine bottle (e.g. plaque 31), compared to the full-length (Lelystad-like) sequence of the Porcilis-derived plaque 36.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to improved PRRS vaccines. In particular, the invention is related to modified live vaccines (MLV). A modified live vaccine is characterized in that it contains live virus which can replicate in pigs, but does not exert clinical symptoms of PRRS, or only few and moderate symptoms of the disease. Furthermore, upon administration it induces an immunological response in pigs which generally leads to a significant extent of protection against subsequent infection with pathogenic PRRS virus. Virus showing such characteristics is usually called attenuated virus. In general, attenuated virus may

be generated from pathogenic virus isolates by repeated passaging in suitable host cells until it shows the desired properties (WO 92/21375, WO 93/06211, WO93/03760, WO 93/07898, WO 96/36356, EP 0 676 467, EP 0 732 340, EP 0 835 930). Alternatively, it may be generated by genetic reengineering through use of an infectious clone, normally a full-length complementary DNA transcript of the viral genome (WO 98/18933, EP 1 018 557, WO 03/062407, Nielsen et al, J Virol 2003, 77:3702-3711). In a preferred embodiment, the present invention relates to a MLV containing attenuated PRRS virus of European genotype.

10 The present invention is based on the unexpected finding that safety and efficacy of a PRRS MLV may be improved if it is taken care that the virus used for the vaccine does not show certain deletions in the nsp2 sequence. More specifically, a virus in accordance with the invention should contain a nucleotide sequence of SEQ ID NO: 1, or a nucleotide sequence which differs from SEQ ID NO: 1 by 1, 2, 3, 4, or 5 nucleotides (either substitu-
15 tions, deletions, or insertions), as part of a full or partial nsp2 coding sequence. In other words, the nucleotides corresponding to nucleotides Nos. 145 to 163 of SEQ ID NO: 1 should not be deleted in said virus.

Attenuated virus strains which may be used for vaccines in accordance with the present
20 invention have been deposited by Boehringer Ingelheim Vetmedica GmbH according to the Budapest Treaty on 27 October 2004 with the European Collection of Cell Cultures (ECACC), Porton Down, Salisbury, Wiltshire, SP4 0JG, Great Britain, with Accession Numbers 04102703 (PRRSV bs104-P27), 04102702 (PRRSV bs104-P31), and 04102704 (PRRSV bs104-P36). A vaccine according to the present invention should however not
25 contain a virus strain deposited with the ECACC at the same day under the same conditions with Accession Number 04102701 (PRRSV bs104-P45). The deposited strains have been obtained by plaque-purification from a commercially obtainable vaccine product (Porcilis[®] PRRS, Intervet Deutschland GmbH, Unterschleißheim), as described in the examples. It unexpectedly turned out that said product comprised four sub-clones, with three
30 of these clones (ECACC04102703, ECACC04102702, or ECACC 04102704) having desirable characteristics in accordance with the present invention, making them suitable as

ingredients of improved vaccine preparations, while the fourth clone (ECACC04102701) has an undesirable nsp2 deletion

In one aspect, the present invention is a PRRS virus strain which is any one of the PRRS virus strains that have been deposited on 27 October 2004 with the European Collection of Cell Cultures (ECACC), Porton Down, Salisbury, Wiltshire, SP4 0JG, Great Britain, under the Accession Numbers ECACC 04102703, ECACC 04102702, and ECACC 04102704, or any descendant or progeny of one of the aforementioned strains. The invention therefore extends to PRRS virus strains which are derived from the deposited strains through propagation or multiplication in an identical or divergent form, in particular descendants which possess the essential characteristics of the deposited strains. Upon continued propagation, the strains may acquire mutations most of which will not alter the properties of these strains significantly.

It is clear to the expert that the strains of the invention may be modified to impart further desirable properties to them. This may be achieved by classical propagation and selection techniques, like continued propagation in suitable host cells to extend the attenuated phenotype. This may be furthermore achieved by directed mutation of the nucleic acid sequence of the genome of these strains by suitable genetic engineering techniques. As outlined above, such techniques usually employ construction of a full-length complementary DNA copies (infectious clones) of the viral genome which may then be modified by DNA recombination and manipulation methods (like site-directed mutagenesis etc.). This way, for example antigenic sites or enzymatic properties of viral proteins may be modified. Infectious clones of PRRS virus strains of European and North American genotype have been reported in the literature, see references cited above.

PRRS virus strains suitable for the vaccines of the invention can be grown and harvested by methods known in the art, e.g. by propagating in suitable host cells like the simian cell line MA-104, Vero cells, or porcine alveolar macrophages, as described in the examples.

Vaccines comprising any one of PRRSV strains ECACC04102703, ECACC04102702, or ECACC 04102704, as well as any combination of these strains or their descendants, but not PRRSV strain ECACC04102701 are thus preferred embodiments of the present invention. Preferably, vaccines according to the present invention are modified live vaccines comprising one or more of these strains alive in a suitable carrier, but inactivated virus may also be used to prepare killed vaccine (KV). MLV are typically formulated to allow administration of 10^1 to 10^7 viral particles per dose, preferably 10^3 to 10^5 particles per dose, more preferably 10^4 to 10^5 particles per dose (4.0 - $5.0 \log_{10}$ TCID₅₀). KV may be formulated based on a pre-inactivation titre of 10^3 to 10^{10} viral particles per dose. The vaccine may comprise a pharmaceutically acceptable carrier, for example a physiological salt-solution. The vaccine may or may not comprise an adjuvant. An example of a suitable adjuvant is a-tocopherol acetate which can be obtained under the trade name Diluvac Forte®. Alternatively, for example alum based adjuvants may be used.

A vaccine according to the present invention may be presented in form of a freeze-dried preparation of the live virus, to be reconstituted with a solvent, to result in a solution for injection. The solvent may e.g. be water, physiological saline, or buffer, or an adjuvanting solvent. The solvent may contain adjuvants, for example a-tocopherol acetate. The reconstituted vaccine may then be injected into the a pig, for example as an intramuscular or intradermal injection into the neck. For intramuscular injection, a volume of 2 ml may be applied, for an intradermal injection it is typically 0.2 ml. In a further aspect, the present invention therefore is a vaccine product, comprising in separate containers a freeze-dried composition of the virus, and a solvent for reconstitution, and optionally further containing a leaflet or label comprising instructions of use.

25

A vaccine according to the present invention may not only comprise one or more of the aforementioned strains, but may include further components active against PRRS or other porcine viral or bacterial diseases, like porcine circovirus or classical swine fever virus. Therefore, the invention further relates to a vaccine as described, characterised in that it contains at least one further antigen active against a porcine disease which is not PRRS.

30

EXAMPLES

1. Materials and Methods

5

*1.1. Experimental infection of pregnant sows with Porcilis[®] PRRS*Study design

10 Fourteen healthy pregnant sows from a confirmed PRRSV negative herd (tested virologically and serologically) were used in this study. Sows faced first or second parturition and were confirmed to be pregnant at the time of vaccination / challenge infection on day 94 (+/-3) of gestation. They were divided into three treatment groups (table 1). The first group was treated with a commercial dose of Porcilis[®] PRRS of 2 ml containing at least
15 $10^{4.0}$ TCID₅₀ i.m. at day 94 (+/- 3) of gestation. The challenge control group (group 2) received a dose of $10^{4.72}$ TCID₅₀ in 2 ml cell culture medium of the pathogenic European field isolate 92045 intranasally. Group 3 was vaccinated with the commercial dose of 2 ml containing $10^{4.0}$ TCID₅₀ i.m. Ingelvac PRRS MLV seven days before insemination and was challenged with the European field isolate 92045 ($10^{4.72}$ TCID₅₀ in 2 ml cell culture medium i.n.) at day 94 (+/- 3) of gestation.
20

Animals from group 1 were monitored until day 5 post-farrowing. Animals from group 2 and 3 were monitored until day 28 post farrowing.

25 Animal phase

All sows were accustomed to the animal facilities 1 week before vaccination. Sows and piglets were observed for their general health status by the investigator on a daily basis. Every animal that died or was euthanised was subjected to post-mortem examination and
30 subsequent laboratory analysis.

Pregnancy was confirmed with ultrasound examination. Serum from sows was obtained on study days 0, 7, 14 and at farrowing for PCR and ELISA investigations. Any material that was associated with abortion was subjected to laboratory investigations.

- 5 Routine gross pathology was performed on all deadborn piglets. Lung tissue samples from all lung lobes were collected from deadborn piglets and from mummies. Samples for PCR testing were stored at -70°C. 2 ml of precolostral blood from each piglet was collected on the day of birth. Serum was prepared and aliquots were stored at -70°C. Serum was used to test for viremia to evaluate the transplacental infection. All piglets of group 1 that survived
10 until day 5, were euthanized at 5 days of age.

Clinical and reproductive performance parameters

- The following main criteria (priority order) were investigated: number of live born piglets
15 per litter, number of stillborn piglets per litter, number of mummified fetuses per litter and number of piglets surviving through day 5 or 28 of age, respectively. The number of piglets born viremic was determined using pre-colostral serum. The frequency of PCR positive blood and tissue samples from sows and/or piglets was investigated to evaluate the epidemiology and course of infection.

20

1.2 Field samples

- The field samples investigated in this study were taken from routine PRRSV diagnostics and consisted of blood, serum and various organ materials, mostly lungs and lymph nodes,
25 from different European countries. The samples were stored at -20°C for a maximum of 3 days before RNA preparation and residual material was subsequently transferred to -70°C for long term storage. RNA and RT-PCR products were stored at -20°C.

1.3. Cell culturing

Ma104 cells (clone CL2621) were grown in MEM (Dulbecco, Germany) supplemented with 10% FCS and antibiotics.

5

Porcine alveolar macrophages were harvested using a method described by Wensvoort et al. (Wensvoort, G. et al. Vet. Quat. 1991, 13:121-130) and modified as follows: each lung lobe was infused with 50-100 ml PBS and subsequently massaged for 3 to 5 min. Then the fluid was rescued from the lobe and passed through a gaze filter. This procedure was repeated until the lavage fluid was clear. The pooled lavage fluid was centrifuged at 500 g for 15 min at room temperature. The pellet was washed in PBS and aliquots of 1×10^7 cells in 50% RPMI 1640 (Biochrom), 40% FCS and 10 % DMSO were frozen at -196°C . For further use the PAMs were cultured in RPMI 1640 medium supplemented with 10% FCS and antibiotics.

15

1.4. Preparation of organ material for virus isolation in cell culture

About $0,5 \text{ cm}^3$ of tissue material was transferred into a tube containing one steel homogenizer ballet in 1,8 ml of sterile PBS. The tubes were agitated for 10 min until the organ material was homogenized. Cell debris was pelleted by centrifugation for 2 min at 450 g and room temperature. The supernatant was passed through a $0,45 \mu\text{m}$ pore sterile filter and stored at -70°C . Aliquots of $30 \mu\text{l}$ were used to inoculate one semiconfluent cell culture monolayer using 24 well microtitre plates.

20

1.5. RNA isolation

25

RNA from organ material was extracted with the RNeasy Mini Kit and from serum, plasma, cell culture supernatant and vaccine solution with the QIAamp Viral RNA Mini Kit (both Qiagen) according to the manufacturer's recommendations, using approximately 100 mg organ material and $140 \mu\text{l}$ fluid material, respectively, for each preparation. The RNA was finally eluted in $65 \mu\text{l}$ buffer as recommended by the manufacturer.

30

1.6. Plaque purification of virus

Confluent monolayers of Ma104 cells in cell culture dishes of 10 cm Ø seeded 48 hours
5 before were infected with the respective virus at tenfold dilutions from 10^{-1} to 10^{-4} . The
cells were incubated for 1 hour with the virus dilutions which were then removed, and the
cells were overlaid with 30 ml of Ma104 medium containing 5% methylcellulose (Sigma).
Plaques were picked after five to seven days and were transferred to Ma104 monolayers in
24 well plates. Virus from these plates was harvested at about 50% CPE and was subjected
10 to further analysis.

1.7. Immunofluorescence assay

Cells were fixed at -20°C for 15 min using ice-cold acetone : methanol (1:1) and air-dried
15 thereafter. After rehydration in PBS, cells were incubated with the PRRSV specific mono-
clonal antibody SDOW17 (Rural Technologies Inc., USA) diluted 1 : 1000 in PBS for 1
hour. After 3 washes with PBS, cells were incubated with goat anti-mouse FITC conju-
gated secondary antibody (Dianova, Hamburg, Germany) (1 : 150 in PBS) for another
hour. After 3 final washes with PBS, cells were overlaid with glycerine : PBS solution
20 (1:1) and subjected to immunofluorescence microscopy.

1.8. Diagnostic nRT-PCR and nsp2 PCR

A diagnostic RT-nPCR was carried out to check the samples for the presence of PRRSV-
25 EU virus; positive samples were then subjected to nsp2 fragment amplification. The primer
sequences are listed in table 2.

The diagnostic RT-nPCR was carried out with the Titan One Tube Kit (Roche Molecular
Biochemicals) as follows: [5 µl total RNA preparation, 1*RT-PCR buffer, 0.4 mM dNTPs,
30 20 pmol of primers PLS and PLR, 5 mM dithiothreitol, 1 mM MgCl_2 , 2.5 – 5 U RNasin
(Promega Ltd), 1 - 2.5 U enzyme mixture, adjusted to a final volume of 25 µl with DEPC

treated aqua dest]. Cycling conditions were as follows: 45°C for 1 hour, 94°C for 2 min and 30 cycles of 94°C for 30 sec, 58°C for 45 sec and 68°C for 45 sec, final elongation step at 68°C for 5 min. The nested PCR reaction was carried out with Qiagen Taq (Qiagen AG) as follows: [1 µl RT-PCR product, 1*PCR buffer, 10 µl Q-solution, 3.5 mM MgCl₂, 0.3 mM dNTPs, 20 pmol of each EU-7-n-s and EU-7-n-as primers, 2.5 U Taq polymerase, adjusted to a final volume of 50 µl with aqua dest]. Cycling conditions were as follows: 7 cycles with 94°C for 1 min, 58°C for 1 min and 72°C for 1 min, followed by 30 cycles with 94°C for 1 min and 70°C for 1.5 min (no annealing step), final elongation step at 70°C for 5 min.

10

Positive samples were then subjected to nsp2 fragment amplification using a one-step reverse transcription PCR with primers EU-1a-2058-s and EU-1a-2683-as with the Qiagen One Step RT-PCR Kit (Qiagen) as follows: [2 µl RNA preparation, 5 µl RT-PCR buffer, 5 µl Q-solution, 0.4 mM dNTPs, 20 pmol of each primer, 2.5 – 5 U RNasin (Promega), 1 µl One Step enzyme mix, adjusted to a final volume of 25 µl with DEPC treated aqua dest]. Cycling conditions were as follows: 50°C for 1 hour, 95°C for 15 min and 40 cycles of 94°C for 60 sec, 60°C for 5 sec, 50°C for 5 sec and 72°C for 2 min, final elongation step at 72°C for 10 min.

20 The nested PCR reaction was carried out with Taq polymerase (Qiagen) and the primer pair EU-1a-2061-s + EU-1a-2574-as as follows: [1 µl RT-PCR product, 1*PCR buffer, 10 µl Q-solution, 3.5 mM MgCl₂, 0.6 mM dNTPs, 20 pmol of each primer, 2.5 U Taq polymerase, adjusted to a final volume of 50 µl with aqua dest]. Cycling conditions were as follows: 8 cycles with 94°C for 1 min, 60°C for 5 sec, 50°C for 5 sec and 68°C for 45 sec, followed by 50 cycles with 94°C for 1 min and 70°C for 1 min (no annealing step), final elongation step at 70°C for 5 min. The nested PCR primers were elongated with M13 tags to facilitate direct nucleotide sequencing of the PCR products.

25

1.9. Nucleotide sequencing

Nucleotide sequencing was performed on the nested PCR products which had been generated with primers that contained an M13 tag, either directly from the PCR reaction or from PCR products that had been excised from agarose gels and purified with the JETsorb gel extraction kit (Genomed). Sequencing was done using the automated sequencer LI-COR DNA Analyzer GENE READER 4200[®] (LI-COR Inc., Lincoln, Nebraska, USA). Nucleotide and deduced amino acid sequences were analyzed with AlignIR[®], vs1.1 (LI-COR Inc., Lincoln, Nebraska, USA) and the DNASIS[®] 2.6 software package (Hitachi Software Genetic Systems Inc., San Francisco, USA).

2. Results

2.1. PCR analysis of an nsp2 fragment generated from the Porcilis[®] PRRS vaccine

A vial containing 10 doses of freeze-dried Porcilis[®] PRRS modified live vaccine was reconstituted with 20 ml of sterile PBS. RNA was prepared from this solution and was subjected to amplification of the nsp2 fragment by RT-nPCR as described. RNA from another European Lelystad-like PRRSV strain of known nucleotide sequence was amplified as a positive control in the same experiment. An amplification product of 569 bp was expected for this fragment, and was obtained for the control strain. However, the Porcilis[®] PRRS sample did not yield such a product. Instead, it showed a double band in a regular 2% agarose gel with one band at about 430 bp and the other at about 350 bp (see figure 1).

Both bands were purified separately from the gel and then subjected to nucleotide sequencing. It became obvious that both PCR products had a deletion compared to the nucleotide sequence of Lelystad Virus; however, the sequences obtained were somewhat ambiguous, and the precise scope of the deletion could not be determined. It was therefore decided to plaque-purify the vaccine in order to rule out that the ambiguities were due to a non-homogeneous vaccine strain.

2.2. Characterization of plaques obtained from the Porcilis[®] PRRS vaccine virus

Again, a vial containing 10 doses of freeze-dried Porcilis[®] PRRS vaccine was reconstituted with 20 ml of sterile PBS. The reconstituted vaccine solution was incubated with confluent
5 Ma104 monolayers at different dilutions as described in *Materials and Methods* and then overlaid with methylcellulose containing medium. Plaques were picked after 5 to 7 days of incubation and were grown in 24 well plates on Ma104 until they showed a CPE of about 50%. The cell culture supernatants were then harvested and stored at -70°C.

10 A total of 47 plaques was subjected to further investigation. To this end, RNA was prepared from the supernatants of the first passage on 24 well plates and was used for RT-nPCR amplification of the nsp2 fragment. The resulting PCR products were analyzed on a 2% agarose gel. The result is summarized in table 3. Apart from the mixed populations which were probably due to overlapping plaques, four specific populations were found (in
15 the size range of about 569 bp, 434 bp, 416 bp and 347 bp, respectively). The PCR products of some representative plaques are shown in figure 2.

The nucleotide sequence of one representative plaque from each group was then determined by direct sequencing of the nsp2 RT-nPCR products and the following results were
20 obtained: plaque no. 36: no deletion in nsp2; plaque no. 27: deletion of 135 nt; plaque no. 31: deletion of 153 nt; and plaque no. 45: deletion of 222 nt. The nucleotide sequences of these plaques are shown in figure 3.

2.3. Experimental infection of pregnant sows with the Porcilis[®] PRRS vaccine

25

Reproductive parameters

Upon farrowing, sows 7062 and 6677 which had been vaccinated with the Porcilis[®] PRRS vaccine on day 90 of gestation, showed moderate to severe reproductive failure (see table
30 4). The proportion of dead and weak born animals was 62.5% in litter 7062 and 44.4% in litter 6677. In addition, until day 5 post farrowing the four live born piglets from sow 7062

and all weak born piglets from the sow 6677 died. The clinical observations are listed in table 5. In conclusion, 5 days after farrowing only 29.4% (5 piglets) of the piglets within both litters still survived.

- 5 Six sows from group 2 farrowed after infection with the PRRS field virus isolate 92045. 25.3% and 26.4% of the piglets were born dead or mummified, respectively. Out of 42 piglets born alive, 24 (27.6%) piglets were born weak and only 18 piglets (20.7%) were born normal. Similar to the litters from sows vaccinated with Porcilis[®] PRRS on day 94 (+/-3) of gestation, only 29.9% of the piglets born survived.

10

Sows vaccinated with Ingelvac PRRS[®] MLV and challenge infected like group 2 farrowed 72.9% normal healthy piglets. 16.7% and 6.2% of the piglets were born dead or mummified, respectively. Although sow 6930 lost all the piglets until day 28 of age, only 2 piglets from other litters died during that period. Thus, 60.4% of the pigs vaccinated with Ingelvac

- 15 PRRS[®] MLV and challenge infected with the highly pathogenic PRRS field virus isolate 92045 survived until day 28 of age.

Virological investigation of clinical samples

20

Sera from sows and piglets as well as lung tissue and body fluid from group 1 piglets were examined for the presence of PRRSV. PCR positive samples were submitted to nucleotide sequencing of the open reading frames 5 and 7 and also of the nsp2 fragment. The results obtained by PCR and virus isolation are listed in table 6.

25

In the two sows 6677 and 7062 that suffered reproductive failure, every PCR positive sample showed an ORF5 and 7 sequence identical with Porcilis[®] PRRS, meaning that no other PRRSV strain was detectable in these animals. While PRRSV was found in all dead born piglets from sow 7062, only one dead born piglet from sow 6677 (50%) reacted positive.

- 30 All weak born piglets from sow 7062 and 67% of weak born piglets as well as 50% of the healthy normal born piglets from sow 6677 yielded positive PCR results which were veri-

fied as Porcilis[®] PRRS by sequence analysis even from samples taken before colostrum uptake. Thus only 4 out of 17 piglets (23.5%) were found to be negative for Porcilis[®] PRRS. For the nsp2 fragment, every sample showed a strong band at about 347 bp which corresponds to the shortest variant of possible nsp2 fragments that can be amplified from the Porcilis[®] PRRS vaccine (figure 4). Most samples also had a very weak band at about 434 bp; only piglet 7062/4310 had a strong double band.

For differential diagnosis, organ material as well serum were investigated for the US genotype of PRRSV and for PPV, BVD, CSFV, BDV, PrV, PCV2, SIV, PEV/PTV, swParamyxoV, EMCV, swHEV as well as on *Leptospira* spp., *Chlamydophila* spp. and *Chlamydia* spp. All these investigations yielded negative results.

Virus isolation on Ma104 and on porcine alveolar macrophages

Lung material from five piglets of each sow was subjected to virus isolation attempts on Ma104 cells and on porcine alveolar macrophages, choosing two dead born and three live born piglets from each sow (see table 6). An aliquot of 30 µl of homogenized and sterile filtered organ suspension was used for inoculation of one well of Ma104 cells and PAM cells, respectively. Cells were incubated at 37°C and 5% CO₂ for 5 days and then stained in an immunofluorescence assay with the PRRSV specific monoclonal antibody SDOW17. Out of the ten organ suspensions, four yielded a cytopathic effect and positive staining results on both Ma104 and PAM cells. The remaining six samples yielded negative results on both cell types.

From each positive sample, RNA was prepared from the respective cell culture supernatants from both PAM and Ma104 cells. RNA was subjected to RT-nPCR for the ORF 5 and 7 as well as for the nsp2 fragment. Nucleotide sequencing of the ORF 5 and 7 PCR products confirmed the virus as Porcilis[®] PRRS. The nsp2 PCR product unambiguously yielded the Porcilis[®] PRRS sequence with the largest deletion of 222 nucleotides.

Resulting out of these experiments, it can be concluded that the variant represented by plaque 45 and deposited under ECACC04102701 is preferably responsible for transplacental infection and the reproductive performance reduction observed.

5 2.4. *Analysis of the nsp2 amplification product obtained from field samples of pigs previously vaccinated with the Porcilis[®] PRRS vaccine*

a) Samples from a farm vaccinated with Porcilis PRRS

10 A PRRSV field virus negative herd that regularly vaccinated the piglets with Porcilis[®] PRRS at the age of 6 weeks was investigated for PRRSV-EU. Ten samples each were taken from piglets aged 4, 7, 9.5 and 13.5 weeks, respectively. In the 9.5 weeks group, three piglets were suffering from the Postweaning Multisystemic Wasting Syndrome (PMWS), which is often correlated with porcine circovirus 2 and PRRS coinfection (Ellis
15 et al, Vet Microbiol 2004, 98:159-163). A PRRSV-EU specific RT-nPCR product was obtained for 1/10, 7/10, 9/10 and 7/10 animals before vaccination, at 1, 3.5 and 7.5 weeks after vaccination. Upon RT-PCR amplification of the nsp2 coding region from the PRRSV positive samples, only Porcilis[®] PRRS typical PCR products were obtained: 8 samples yielded only the ~347 bp product, 4 samples only the ~434 bp product and 4 samples the
20 double band of 434 and 347 bp. No sample gave the Lelystad-like product (see figure 5). The nsp2 region could be amplified in three of the four PMWS diseased piglets, two of which showed the 434 bp product and one the 347 bp product.

b) Nsp2 length variation in field samples from routine diagnostics

25

A total of 514 blood and tissue samples which had been sent in for diagnostics from PRRS suspected herds and had been PRRSV positive in routine RT-nPCR was subjected to nsp2 fragment amplification. Amplification products were obtained in 41.4 % out of 514 samples. Deletions of varying length were found in 30.7 % of the analyzed samples, which indicates a total frequency of at least 12.7% nsp 2 deletion variants in the field. In most
30 cases, the bands found showed the same pattern as found with the Porcilis[®] PRRS vaccine,

consisting of a (double) band of 434 and 416 bp which cannot easily be differentiated on a regular 2% agarose gel, or a band of 347 bp. A selection of samples was therefore subjected to nucleotide sequencing to check whether these PCR products were in fact derived from Porcilis® PRRS vaccine virus. The results are shown in figure 6 A-C.

5

Table 1: Study groups and treatment

Group	Sows n	Vaccination seven days before insemination	Vaccination / Infection at day 94 (+/- 3) of gestation	Dose
1	2	-	Porcilis® PRRS	$> 10^{4.0}$ TCID ₅₀ in 2ml i.m.
2	6	-	European field isolate 92045 passage 3	$10^{4.72}$ TCID ₅₀ in 2ml i.n.
3	6	Ingelvac® PRRS MLV	European field isolate 92045 passage 3	$10^{4.72}$ TCID ₅₀ in 2ml i.m.

n – number of sows; i.m. intramuscularly, i.n. intranasally

10 Table 2: Primer sequences for generating nsp2 RT-PCR products

Name	Sequence	Purpose
PLS	ATGGCCAGCCAGTCAATC	RT-PCR ORF7
PLR	TCGCCCTAATTGAATAGGTG	RT-PCR ORF7
EU-7-n-s	<i>TGTAAAACGACGGCCAGT</i> <u><i>TATGATAAAGTCCCAGCGCCAG</i></u>	nPCR ORF7
EU-7-n-as	<i>CAGGAAACAGCTATGACCC</i> <u><i>TGTATGAGCAACCGGCAGCAT</i></u>	nPCR ORF7
EU-1a-2058-s	TGTTGAAGGATTGTCCGAGCTCC	RT-PCR ORF1a
EU-1a-2683-as	AGATCCAAAGGCGAACTGCTG	RT-PCR ORF1a
EU-1a-2061-s	<i>TGTAAAACGACGGCCAGT</i> <u><i>TGAAGGATTGTCCGAGCTCC</i></u>	nPCR ORF1a
EU-1a-2574-as	<i>CAGGAAACAGCTATGACCAT</i> <u><i>CAGAACCTGGGTTGTCGG</i></u>	nPCR ORF1a

For the nested PCR primers, M13 tags are marked *italic*, while the PRRSV specific sequence of the primers is underlined.

Table 3: nsp2 PCR results from 47 plaques after 1 passage in Ma104 cell culture

PCR product length	Σ plaques	Plaque no.
~ 569 bp	2	36, 58
~ 434 bp	10	24, 27, 32, 43, 46, 49, 53, 55, 57, 59
~ 416 bp	1	31
~ 347 bp	20	8, 9, 10, 11, 12, 13, 25, 26, 28, 33, 37, 38, 40, 42, 44, 45, 47, 51, 54, 56
Mixed	14	1, 2, 4, 5, 6, 7, 15, 16, 18, 30, 34, 35, 41, 48, 19, 50, 52

Table 4: Comparison of the reproductive and production parameters of sows vaccinated with Porcilis® PRRS at day 94 (+/-) with challenge and vaccinated / challenge controls

Group	Sow ID	n	Piglets born alive			Piglets born	Piglets died	Total piglets alive
			Total	Normal	Weak	Dead/Mummified	until 5 / 28 DoA	at 5 / 28 DoA
1	7062	8	4	3	1	4/0	4	0
	6677	9	7	5	2	2/0	2	5
	mean	8.5	5,5	4.0	1.5	3.0	3.0	2.5
	total	17	11	8	3	6/0	6	5
	%		64.7	47.0	17.7	35.3/0	35.3	29.4
2	6746	16	13	2	11	3/0	7	6
	7008	13	8	5	3	3/2	2	6
	7076	15	7	2	5	4/4	2	5
	7086	13	9	8	1	4/0	3	6
	7091	15	3	1	2	4/8	1	2
	7145	15	2	0	2	4/9	1	1
	Mean	14.5	7.0	3.0	4.0	3.7/3.8	2,7	4,3
	Total	87	42	18	24	22/23	16	26
	%		48.3	20.7	27.6	25.3/26.4	38.1	29.9
3	6678	11	10	9	1	1/0	1	9
	6930	7	6	5	1	1/0	6	0
	6955	6	3	3	0	3/0	0	3
	7095	9	6	6	0	2/1	1	5
	7158	15	12	12	0	1/2	0	12
	Mean	9.6	7,4	7,0	0,4	1,6/0.8	1,6	5,8
	Total	48	37	35	2	8/3	8	29
	%		77.1	72.9	4.2	16.7/6.2	21.6	60,4

Table 5: Clinical assessment of the offspring of the sows vaccinated with Porcilis[®] PRRS on day 90 of gestation (group 1)

Sow ID	Pig ID	Day of age					
		0	1	2	3	4	5
6677	4300	-	-	W	W	W	W
	4301	-	W, Dep	Died			
	4302	-	-	W	W	W	-
	4303	-	-	-	-	-	-
	4304	W	Died				
	4305	-	Died ¹	-	-	-	-
	4306	W	-	W, A	W, A	W, Dep	W, A
	1	Born dead					
	2	Born dead					
7062	4307	-	W, Dep	W, Dep	W, Dep	Died ²	
	4308	W	Died ²				
	4309	-	W, Dep	Died			
	4310	-	W, Dep	Died ³			
	1	Born dead					
	2	Born dead					
	3	Born dead					
	4	Born dead					

W = weak, Dep = depressed, A = anorexia

¹Piglet 4305 died on day 1 due to a blood taking accident.

²Piglets 4307 and 4308 died due to agalactia.

³Piglet 4310 showed severe signs of pneumonia.

Table 6: Virological results from sows vaccinated with Porcilis[®] PRRS at day 94 (+/- 3) of gestation and their litters

Sow/Piglet ID	Material	PRRSV-EU (PCR)	Porcilis PRRS verified by sequencing*	Reisolates of Porcilis PRRS MLV on PAMs**	Reisolates of Porcilis PRRS MLV on Ma104**
6677	serum, day 0	neg	nd.	nd.	nd.
	serum, day 7	pos	yes	nd.	nd.
	serum, day 14	pos	yes	nd.	nd.
	serum, day of farrowing	pos	yes	nd.	nd.
7062	serum, day 0	neg	nd.	nd.	nd.
	serum, day 7	pos	yes	nd.	nd.
	serum, day 14	pos	yes	nd.	nd.
	serum, day of farrowing	pos	yes	nd.	nd.
6677 / 1	lung tissue	neg	nd.	neg	neg
6677 / 2	lung tissue	pos	yes	neg	neg
6677 / 4300	lung tissue	pos	yes	nd.	nd.
6677 / 4301	lung tissue	pos	yes	pos	pos
6677 / 4302	lung tissue	pos	yes	neg	neg
6677 / 4303	lung tissue	pos	yes	pos	pos
6677 / 4304	lung tissue	neg	nd.	nd.	nd.
6677 / 4305	lung tissue	neg	nd.	nd.	nd.
6677 / 4306	lung tissue	neg	nd.	nd.	nd.
7062 / 1	lung tissue	pos	yes	neg	neg
7062 / 2	lung tissue	pos	yes	neg	neg
7062 / 3	lung tissue	pos	yes	nd.	nd.
7062 / 4	lung tissue	pos	yes	nd.	nd.
7062 / 4307	lung tissue	pos	yes	pos	pos
7062 / 4308	lung tissue	pos	yes	pos	pos
7062 / 4309	lung tissue	pos	yes	neg	neg.
7062 / 4310	lung tissue	pos	yes	nd.	nd.

- 5 nd: not done
 *: The PCR fragment of the routinely performed diagnostic PCR was sequenced
 **: All reisolates obtained from MA 104 and PAMs were sequenced in ORF5 and ORF 7
 and were identical with Porcilis[®] PRRS

Table 7: History of samples showing an nsp2 deletion PCR product; cf. Figure 6A

Sample ID	Origin			History
115502	Hungary	2004	33 d.p.i.	animal vaccinated with Porcilis [®] PRRS in a controlled study experiment
159160	Great Britain	2003	non-vaccinated piglet, 4 weeks old, in a herd that is regularly vaccinated with Porcilis [®] PRRS	healthy piglet
159178	Great Britain	2003	piglet vaccinated with Porcilis [®] PRRS at the age of 6 weeks, 24 days post vaccination	piglet is showing respiratory disease

Table 8: History of samples showing an nsp2 deletion PCR product; cf. Figure 6B

Sample ID	Origin			History
115502	Hungary	2004	33 d.p.i.	animal vaccinated with Porcilis [®] PRRS in a controlled study experiment
159176	Great Britain	2003	piglet vaccinated with Porcilis [®] PRRS at the age of 6 weeks, 24 days post vaccination	piglet is showing respiratory disease

Table 9: History of samples showing an nsp2 deletion PCR product; cf. Figure 6C

Sample ID	Origin			History
159161	Great Britain	2003	piglet vaccinated with Porcilis [®] PRRS at the age of 6 weeks, 7 days post vaccination	healthy piglet

CLAIMS:

1. Vaccine against Porcine Respiratory Reproductive Syndrome (PRRS), comprising live PRRS virus, characterized in that it contains one or more of the PRRS virus strains
5 that have been deposited on 27 October 2004 with the European Collection of Cell Cultures (ECACC), Porton Down, Salisbury, Wiltshire, SP4 0JG, Great Britain, under the Accession Numbers ECACC 04102703, ECACC 04102702, and ECACC 04102704, or a descendant of one or more of the aforementioned strains, optionally together with a pharmaceutically acceptable carrier, but not the strain deposited under
10 the Accession Number ECACC 04102701.
2. Vaccine of claim 1, containing the PRRS virus strains that have been deposited under the Accession Numbers ECACC 04102703, ECACC 04102702, and ECACC 04102704.
15
3. Vaccine of claim 1 or claim 2, further containing an adjuvant.
4. Vaccine of claim 3, wherein the adjuvant is α -tocopherol acetate.
- 20 5. Vaccine of any one of claims 1 to 4, which comprises a freeze-dried composition of the virus.
6. A PRRS virus strain which is any one of the PRRS virus strains that have been deposited on 27 October 2004 with the European Collection of Cell Cultures (ECACC), Porton Down, Salisbury, Wiltshire, SP4 0JG, Great Britain, under the Accession Numbers
25 ECACC 04102703, ECACC 04102702, and ECACC 04102704, or a descendant of any one of the aforementioned strains.
7. Method of making a vaccine, comprising admixing one or more of a PRRS virus strain
30 according to claim 6 with a pharmaceutically acceptable carrier.

8. Use of a PRRS virus according to claim 6 for the manufacture of a vaccine.
9. Vaccine product, comprising in separate containers a freeze-dried composition according to claim 5, and a solvent for reconstitution, and optionally further containing a
5 leaflet or label comprising instructions of use.
10. Vaccine product of claim 9, wherein the solvent contains an adjuvant.
11. Method of vaccinating a pig, comprising administering a vaccine or vaccine product
10 according to any one of claims 1 to 5, 7, 9, or 10 to such pig.
12. Vaccine according to any one of the foregoing claims, characterised in that it contains at least one further antigen active against a porcine disease which is not PRRS.

1



Figure 1




Figure 2

```

      *      20      *      40      *      60      *      80
Lelystad : ..... :
Plaque27 : ..... :
Plaque31 : ..... :
Plaque36 : ..... :
Plaque45 : .....T..... :
      TGAAGGATTGTCGAGCTCCAAACAGGCCATGGCCCTTCTGGCAAAATTAAAGTTCCATCCTCAAAGGCCCGTCTGTGTCC

      *      100     *      120     *      140     *      160
Lelystad : .....A.....TT.....A..... :
Plaque27 : ..... :
Plaque31 : ..... :
Plaque36 : .....T..... :
Plaque45 : ..... :
      CTGGACGGGTGTTTCCCTACGGATGTTCCAGCCGACTTCGAGCCAGCATCTCCGAAAGGCCCAAAGTTCCGGCGCTGCTGT

      *      180     *      200     *      220     *      240
Lelystad : .....G.....G..... :
Plaque27 : ..... :
Plaque31 : ..... :
Plaque36 : .....A..... :
Plaque45 : ..... :
      TGTCTGTGTTCAACCGGATGCAAAAGAGTTTCGAGGAAACAGCCCCAGAAGAAGTTCAAGAGAGTGGCCACAAGGCCGTCCACT

      *      260     *      280     *      300     *      347     *
Lelystad : ..... :
Plaque27 : ..... :
Plaque31 : ..... :
Plaque36 : ..... :
Plaque45 : ..... :
      CTGCACTCCTTGCCGAGGGTCCTAACAATGAGCAGGTACAGGTGGTTGCCGGTGAGCAACTGAAGCTCGGCGGTTGTGGTTTG

      340      *      360      *      416      *      434      *
Lelystad : .....C..T.....C. :
Plaque27 : .....T..... :
Plaque31 : ..... :
Plaque36 : ..... :
Plaque45 : ..... :
      GCAGTCGGGAATGCTCATGAAGGTGTTCCGGTCTCAGCTGGTCTAATTAACCTGGTAGGCGGAATTTGTCCCCCTCAGACTC

      420      *      440      *      460      *      480      *      5
Lelystad : .....T..... :
Plaque27 : .....G..... :
Plaque31 : .....T..GCTGGTC..... :
Plaque36 : ..... :
Plaque45 : ..... :
      CATGAAAGAAAACATGCTCAATAGCCGGAAGACGAACCACTGGATTTGTCCAACCAGCACCAGCTGCCACAACGACCCTTG

      00      *      520      *
Lelystad : ..... :
Plaque27 : ..... :
Plaque31 : ..... :
Plaque36 : ..... :
Plaque45 : ..... :
      TGAGAGAGCAAACACCCGACAACCCAGGTTCTGAT
```

Figure 3

1 w p.vacc.

3.5 ww p.vacc.

7.5 ww p.vacc.

Porcilis® PRRS MLV

Lelystad-like isolate

Pigs diseased on PMWS

Figure 5

4

		*	20	*	40	*	60	*	80			
P36	:	TGAAGGAT	TGTCCGAGCT	CCAAACAGGCCAT	GGCCCTTCT	TGGCAAAAAT	TAAAGTTCCAT	CCTCAAAGGCCCCG	TCTGTGTCCCT			
159160	:											
159178	:								T..			
161009	:											
169736	:											
190146	:											
81984	:											
		*	100	*	120	*	140	*	160	*		
P36	:	GGACGGGT	GTTTCCCTAC	GGATGTTCC	AGCCGACTT	CGAGCCAGC	ATTTCCGGAA	AGGCCCAAG	TTCCGGCG	CTGCTGTTGTC		
159160	:					C.						
159178	:					C.						
161009	:					C.						
169736	:					C.						
190146	:					C.						
81984	:					A..C.						
		180	*	200	*	220	*	240	*			
P36	:	CTGTGTT	CACCGAAT	GCAAAAG	AGTT	CGAGGAA	ACAGCCCC	CAGAAGA	AGTTCAAG	GAGAGTGGCCACAAGGCCG	TCCACTCTGCAC	
159160	:											
159178	:											
161009	:											
169736	:											
190146	:											
81984	:											
		260	*	280	*	300	*	347	*	340		
P36	:	TCCTTGCC	GAGGGT	CCTAACA	ATGAGC	AGGTAC	AGGTGGT	TGCCGGT	GAGCAACT	GAAGCTCGGCGGT	TGTGGTTTGGCAGT	CGG
159160	:											
159178	:											
161009	:											
169736	:											
190146	:											
81984	:											
		*	360	*	416	*	434	*	420			
P36	:	GAATGCT	CATGAAG	GTGTTCC	GGTCTC	AGCTGGT	CTAATTA	ACCTGGT	AGGCGGGA	ATTTGTCCCCCT	CAGACTCCAT	GAAAGAA
159160	:											
159178	:											
161009	:										G.	
169736	:											
190146	:											
81984	:											
		*	440	*	460	*	480	*	500	*		
P36	:	AACATGCT	CAATAGCC	GGGAAGAC	GAACCACT	GGATTTGT	CCCAACC	AGCACC	AGCTGCC	ACAACG	ACCCTTGT	GAGAGAGCAAA
159160	:											
159178	:										T..T.	
161009	:											
169736	:											
190146	:											
81984	:											
		520	*									
P36	:	CACCCGACA	ACCCAGG	TTCTGAT								
159160	:	...										
159178	:											
161009	:											
169736	:	..										
190146	:											
81984	:											

Figure 6A




Figure 6B




Figure 6C