

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017355504 C1**

(54) Title
Anti-BCMA CAR T cell compositions

(51) International Patent Classification(s)
A61K 48/00 (2006.01) **C12N 15/62** (2006.01)

(21) Application No: **2017355504** (22) Date of Filing: **2017.11.03**

(87) WIPO No: **WO18/085690**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/417,840	2016.11.04	US
62/514,401	2017.06.02	US

(43) Publication Date: **2018.05.11**

(44) Accepted Journal Date: **2024.02.22**

(44) Amended Journal Date: **2024.06.20**

(71) Applicant(s)
2seventy bio, Inc.

(72) Inventor(s)
Quigley, Travis; Ross, Robert

(74) Agent / Attorney
Davies Collison Cave Pty Ltd, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000, AU

(56) Related Art
WO 2016/094304 A2
ANONYMOUS: "Study of bb2121 in Multiple Myeloma - Tabular View - ClinicalTrials.gov", 20 January 2016 (2016-01-20), XP055697969, Retrieved from the Internet [retrieved on 20200525]



(51) International Patent Classification:

A61K 48/00 (2006.01) C12N 15/62 (2006.01)

(21) International Application Number:

PCT/US2017/059989

(22) International Filing Date:

03 November 2017 (03.11.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/417,840 04 November 2016 (04.11.2016) US
62/514,401 02 June 2017 (02.06.2017) US

(71) Applicant: BLUEBIRD BIO, INC. [US/US]; 60 Binney Street, Cambridge, Massachusetts 02142 (US).

(72) Inventors: QUIGLEY, Travis; 49 Elm Court, Cohasset, Massachusetts 02025 (US). ROSS, Robert; 9 Somerset Road, Brookline, Massachusetts 02445 (US).

(74) Agent: MASHRUWALA, Mary Anne et al.; Cooley LLP, 1299 Pennsylvania Avenue, NW, Suite 700, Washington, District of Columbia 20004 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: ANTI-BCMA CAR T CELL COMPOSITIONS




FIGURE 3

(57) Abstract: The invention provides improved anti-BCMA CAR T cell compositions for adoptive T cell therapy for relapsed/refractory multiple myeloma.

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

ANTI-BCMA CAR T CELL COMPOSITIONS

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 62/514,401, filed June 2, 2017, and U.S. Provisional Application No. 62/417,840, filed November 4, 2016, each of which is incorporated by reference herein in its entirety.

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification.

The name of the text file containing the Sequence Listing is BLBD_079_02WO_ST25.txt. The text file is 27 KB, was created on November 3, 2017, and is being submitted electronically via EFS-Web, concurrent with the filing of the specification.

BACKGROUND

Technical Field

The present invention relates to improved compositions and methods for treating B cell related conditions. More particularly, the invention relates to improved compositions comprising therapeutic doses of anti-BCMA chimeric antigen receptor (CAR) T cells to treat relapsed/refractory multiple myeloma.

Description of the Related Art

Chimeric Antigen Receptor (CAR) T-cells are molecules that combine antibody-based specificity for a desired antigen with a T cell receptor-activating intracellular domain to generate a chimeric protein that exhibits a specific anti-cancer immune activity. CARs endow a patient's own T cells with the ability to recognize and kill cancer cells directly. CAR T cell technology has developed rapidly, but has yet to

realize its therapeutic potential. Several obstacles remain to the clinical application of CAR T cells to hematological malignancies.

Most CAR T cell therapies applied to hematological malignancies are associated with cytokine release syndrome (CRS) or cytokine storm. CRS is specific for adoptive
5 cellular immunotherapies, manifests with septic shock-like symptoms, and may ultimately result in patient death. In fact, a number of clinical trials were temporarily suspended by the FDA due to CRS-related deaths in CAR T-cell trials. CRS can be rapid, unpredictable and currently, there is no consensus on prevention and management of CRS. However, one characteristic that contributes to CRS is the
10 therapeutic cell dose of CAR T cells.

CAR T cells are “living drugs” that have the ability to multiply *in vivo* - after infusion and home to bone marrow. For example, in a seminal case study from UPenn, published in 2011, as low as 1.5×10^5 CAR T cells, given to chronic lymphocytic leukemia (CLL) patients, expanded more than 1000 times *in vivo* over time. Thus,
15 CAR T cell therapy presents a unique set of challenges with respect to dosing. There is no uniform consensus on CAR T cell infusion dosage and influence factors, including tumor load, efficacy and side effects. Small dose infusions may not obtain the ideal curative effect, but may instead induce tumor antigen deletion, and antigen escape. Large dose infusions and serious tumor load will increase CRS and tumor lysis
20 syndrome. Because it is a living drug, determining the correct therapeutic dose is unpredictable. Conventional drug-body interaction concepts of pharmacodynamics and pharmacokinetics can’t readily be applied to CAR T cell therapy because CAR T cells are a dynamic, living, and persistent drug.

Some challenges to dose selection include, but are not limited to, the paucity of
25 correlative animal models, first in human products have limited “a priori” information, *in-vivo* expansion of humans cell is unpredictable, the limitations in “borrowing” safety data from first generation CAR T products, extrapolating safety data from related products (TILs, “similar” TCR redirected cells, “similar” class of CAR T product, and extrapolating safety data using the same product in histologically different tumor
30 type(s).

Accordingly, current approaches to CAR T cell dosing do not appear to be sufficiently mature to predict therapeutically effective CAR T cell dose among T cells harboring different CARs for use in the same indication and among the same CAR T cells for use in different indications.

5 BRIEF SUMMARY

The invention generally provides improved anti-BCMA CAR T cell compositions and methods of using the same to treat relapsed/refractory multiple myeloma.

In various embodiments, a composition is contemplated comprising a
10 therapeutically effective amount of anti-B cell maturation antigen (BCMA) chimeric antigen receptor (CAR) T cells, wherein the therapeutically effective amount is greater than about 5.0×10^7 anti-BCMA CAR T cells, and the anti-BCMA CAR comprises the amino acid sequence set forth in SEQ ID NO: 9.

In particular embodiments, the composition further comprises a
15 pharmaceutically acceptable carrier.

In certain embodiments, the composition is formulated in a solution comprising 50:50 PlasmaLyte A to CryoStor CS10.

In various embodiments, the therapeutically effect amount is at least about 15.0×10^7 anti-BCMA CAR T cells.

20 In particular embodiments, the therapeutically effect amount is at least about 45.0×10^7 anti-BCMA CAR T cells.

In some embodiments, the therapeutically effect amount is at least about 80.0×10^7 anti-BCMA CAR T cells.

In certain embodiments, the therapeutically effect amount is at least about $12.0 \times$
25 10^7 anti-BCMA CAR T cells.

In particular embodiments, the therapeutically effect amount is between about 5.0×10^7 anti-BCMA CAR T cells and about 15.0×10^7 anti-BCMA CAR T cells.

In various embodiments, the therapeutically effect amount is between about 5.0×10^7 anti-BCMA CAR T cells and about 45.0×10^7 anti-BCMA CAR T cells.

In some embodiments, the therapeutically effect amount is between about 5.0×10^7 anti-BCMA CAR T cells and about 80.0×10^7 anti-BCMA CAR T cells.

In particular embodiments, the therapeutically effect amount is between about 5.0×10^7 anti-BCMA CAR T cells and about 12.0×10^8 anti-BCMA CAR T cells.

5 In certain embodiments, the therapeutically effect amount is between about 15.0×10^7 anti-BCMA CAR T cells and about 45.0×10^7 anti-BCMA CAR T cells.

In some embodiments, the therapeutically effect amount is between about 15.0×10^7 anti-BCMA CAR T cells and about 80.0×10^7 anti-BCMA CAR T cells.

In various embodiments, the therapeutically effect amount is between about 15.0×10^7 anti-BCMA CAR T cells and about 12.0×10^8 anti-BCMA CAR T cells.

In particular embodiments, the anti-BCMA CAR T cells are transduced with a lentiviral vector encoding the anti-BCMA CAR.

In certain embodiments, the lentiviral vector copy number (VCN) is about 2.0 copies per anti-BCMA CAR T cell.

15 In certain embodiments, the lentiviral vector is a human immunodeficiency virus 1 (HIV-1) vector.

In particular embodiments, a pharmaceutical composition is contemplated comprising a population of T cells transduced with a lentivirus encoding an anti-BCMA CAR, wherein population of T cells comprises greater than about 5.0×10^7 anti-BCMA CAR T cells, and the anti-BCMA CAR comprises the amino acid sequence set forth in SEQ ID NO: 9.

In some embodiments, the composition further comprising a therapeutically acceptable carrier.

In various embodiments, the composition comprises at least about 15.0×10^7 anti-BCMA CAR T cells.

In particular embodiments, the composition comprises at least about 45.0×10^7 anti-BCMA CAR T cells.

In some embodiments, the composition comprises at least about 80.0×10^7 anti-BCMA CAR T cells.

30 In certain embodiments, the composition comprises at least about 12.0×10^8 anti-BCMA CAR T cells.

In various embodiments, the composition comprises about 15.0×10^7 anti-BCMA CAR T cells.

In particular embodiments, the composition comprises about 45.0×10^7 anti-BCMA CAR T cells.

5 In some embodiments, the composition comprises about 80.0×10^7 anti-BCMA CAR T cells.

In certain embodiments, the composition comprises about 12.0×10^7 anti-BCMA CAR T cells.

10 In particular embodiments, the composition comprises between about 5.0×10^7 anti-BCMA CAR T cells and about 15.0×10^7 anti-BCMA CAR T cells.

In various embodiments, the composition comprises between about 5.0×10^7 anti-BCMA CAR T cells and about 45.0×10^7 anti-BCMA CAR T cells.

In particular embodiments, the composition comprises between about 5.0×10^7 anti-BCMA CAR T cells and about 80.0×10^7 anti-BCMA CAR T cells.

15 In some embodiments, the composition comprises between about 5.0×10^7 anti-BCMA CAR T cells and about 12.0×10^8 anti-BCMA CAR T cells.

In various embodiments, the composition comprises between about 15.0×10^7 anti-BCMA CAR T cells and about 45.0×10^7 anti-BCMA CAR T cells.

20 In certain embodiments, the composition comprises between about 15.0×10^7 anti-BCMA CAR T cells and about 80.0×10^7 anti-BCMA CAR T cells.

In some embodiments, the composition comprises between about 15.0×10^7 anti-BCMA CAR T cells and about 12.0×10^8 anti-BCMA CAR T cells.

In particular embodiments, the T cells comprise CD8⁺ T cells.

25 In various embodiments, a method of treating a subject that has been diagnosed with relapsed/refractory multiple myeloma is provided comprising administering the subject a composition contemplated herein.

In particular embodiments, a method of treating a subject that has relapsed/refractory multiple myeloma is provided, comprising administering the subject the composition contemplated herein.

30 In certain embodiments, the composition is administered in a single dose.

In certain embodiments, the composition is intravenously administered.

In some embodiments, the multiple myeloma was refractory to at least three treatment regimens, including a proteasome inhibitor and an immunomodulatory agent, prior to the administration of the composition.

In particular embodiments, the multiple myeloma was double-refractory to one
5 or more treatment regimens, prior to the administration of the composition.

In some embodiments, the subject was treated with daratumumab, lenalidomide, pomalidomide, bortezomib, and/or carfilzomib, prior to the administration of the composition.

In certain embodiments, the subject received an autologous hematopoietic stem
10 cell transplant, prior to the administration of the composition.

In particular embodiments, the subject was lymphodepleted with cyclophosphamide 300 mg/m² and fludarabine 30 mg/m².

BRIEF DESCRIPTION OF SEVERAL VIEWS OF THE DRAWINGS

Figure 1 shows a schematic of a murine B cell maturation antigen (muBCMA)
15 CAR construct.

Figure 2A shows comparable levels of expansion for anti-BCMA CAR T cells manufactured from PBMCs of 11 individual donors compared to a matched culture of untransduced donor T cells.

Figure 2B shows comparable levels of lentiviral transduction efficiency (VCN)
20 in anti-BCMA CAR T cells manufactured from PBMCs of 11 individual donors.

Figure 2C shows comparable anti-BCMA CAR expression by flow cytometry in anti-BCMA CAR T cells manufactured from PBMCs of 11 individual donors.

Figure 2D shows comparable levels of IFN γ release when anti-BCMA CAR T cells manufactured from PBMCs of 11 individual donors were exposed to BCMA-
25 expressing K562 cells.

Figure 3 shows clinical response over time of relapsed refractory multiple myeloma patients treated with anti-BCMA CAR T cells.

BRIEF DESCRIPTION OF THE SEQUENCE IDENTIFIERS

SEQ ID NOs: 1-3 set forth amino acid sequences of exemplary light chain CDR sequences for BCMA CARs contemplated herein.

SEQ ID NOs: 4-6 set forth amino acid sequences of exemplary heavy chain CDR sequences for BCMA CARs contemplated herein.

SEQ ID NO: 7 sets forth an amino acid sequence of an exemplary light chain sequences for BCMA CARs contemplated herein.

SEQ ID NO: 8 sets forth an amino acid sequence of an exemplary heavy chain sequences for BCMA CARs contemplated herein.

SEQ ID NO: 9 sets forth an amino acid sequence of an exemplary BCMA CAR contemplated herein.

SEQ ID NO: 10 set forth a polynucleotide sequence that encode an exemplary BCMA CAR contemplated herein.

SEQ ID NO: 11 sets forth the amino acid sequence of human BCMA.

SEQ ID NO: 12-22 set for the amino acid sequence of various linkers.

SEQ ID NOs: 23-35 set for the amino acid sequence of protease cleavage sites and self-cleaving polypeptide cleavage sites.

SEQ ID NO: 36 sets for the polynucleotide sequence of a vector encoding a BCMA CAR.

DETAILED DESCRIPTION**A. OVERVIEW**

Significant challenges exist in the art to predicting a therapeutically effective CAR T cell dose with respect to a particular CAR and its target indication. The present disclosure generally relates to improved anti-BCMA CAR T cell compositions and methods for treating relapsed/refractory multiple myeloma.

B cell maturation antigen (BCMA, also known as CD269 or tumor necrosis factor receptor superfamily, member 17; TNFRSF17 is a member of the tumor necrosis factor receptor superfamily (see, e.g., Thompson *et al.*, *J. Exp. Medicine*, 192(1): 129-135, 2000, and Mackay *et al.*, *Annu. Rev. Immunol.*, 21: 231-264, 2003. BCMA binds

B-cell activating factor (BAFF) and a proliferation inducing ligand (APRIL) (*see, e.g.,* Mackay *et al.*, 2003 and Kalled *et al.*, *Immunological Reviews*, 204: 43-54, 2005).

Among nonmalignant cells, BCMA has been reported to be expressed mostly in plasma cells and subsets of mature B-cells (*see, e.g.,* Laabi *et al.*, *EMBO J.*, 77(1): 3897-3904, 1992; Laabi *et al.*, *Nucleic Acids Res.*, 22(7): 1147-1154, 1994; Kalled *et al.*, 2005; O'Connor *et al.*, *J. Exp. Medicine*, 199(1): 91-97, 2004; and Ng *et al.*, *J. Immunol.*, 73(2): 807-817, 2004. Mice deficient in BCMA are healthy and have normal numbers of B cells, but the survival of long-lived plasma cells is impaired (*see, e.g.,* O'Connor *et al.*, 2004; Xu *et al.*, *Mol. Cell. Biol.*, 21(12): 4067-4074, 2001; and Schiemann *et al.*, *Science*, 293(5537): 2111-2114, 2001). BCMA RNA has been detected universally in multiple myeloma cells and in other lymphomas, and BCMA protein has been detected on the surface of plasma cells from multiple myeloma patients by several investigators (*see, e.g.,* Novak *et al.*, *Blood*, 103(2): 689-694, 2004; Neri *et al.*, *Clinical Cancer Research*, 73(19): 5903-5909, 2007; Bellucci *et al.*, *Blood*, 105(10): 3945-3950, 2005; and Moreaux *et al.*, *Blood*, 703(8): 3148-3157, 2004.

In various embodiments, the improved adoptive cell therapy compositions contemplated herein comprise therapeutically effective doses of anti-BCMA CAR T cells that unexpectedly expand and elicit clinical responses *in vivo*, in the absence of severe cytokine release syndrome CRS.

In particular embodiments, the improved compositions comprising therapeutically effective doses of anti-BCMA CAR T cells contemplated herein are used to treat relapsed/refractory multiple myeloma.

The practice of the invention will employ, unless indicated specifically to the contrary, conventional methods of chemistry, biochemistry, organic chemistry, molecular biology, microbiology, recombinant DNA techniques, genetics, immunology, and cell biology that are within the skill of the art, many of which are described below for the purpose of illustration. Such techniques are explained fully in the literature. *See, e.g.,* Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual* (3rd Edition, 2001); Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); Maniatis *et al.*, *Molecular Cloning: A Laboratory Manual* (1982); Ausubel *et al.*, *Current Protocols in Molecular Biology* (John Wiley and Sons, updated July 2008);

Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology, Greene Pub. Associates and Wiley-Interscience; Glover, *DNA Cloning: A Practical Approach*, vol. I & II (IRL Press, Oxford, 1985); Anand, *Techniques for the Analysis of Complex Genomes*, (Academic Press, New York, 5 1992); *Transcription and Translation* (B. Hames & S. Higgins, Eds., 1984); Perbal, *A Practical Guide to Molecular Cloning* (1984); Harlow and Lane, *Antibodies*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1998) *Current Protocols in Immunology* Q. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach and W. Strober, eds., 1991); *Annual Review of Immunology*; as well as monographs in journals 10 such as *Advances in Immunology*.

B. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to 15 those described herein can be used in the practice or testing of the present invention, preferred embodiments of compositions, methods and materials are described herein. For the purposes of the present invention, the following terms are defined below.

The articles “a,” “an,” and “the” are used herein to refer to one or to more than one (*i.e.*, to at least one, or to one or more) of the grammatical object of the article. By 20 way of example, “an element” means one element or one or more elements.

The use of the alternative (*e.g.*, “or”) should be understood to mean either one, both, or any combination thereof of the alternatives.

The term “and/or” should be understood to mean either one, or both of the alternatives.

25 As used herein, the term “about” or “approximately” refers to a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2% or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length. In one embodiment, the term “about” or “approximately” 30 refers a range of quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length $\pm 20\%$, $\pm 15\%$, $\pm 10\%$, $\pm 9\%$, $\pm 8\%$, $\pm 7\%$, $\pm 6\%$, $\pm 5\%$, $\pm 4\%$,

± 3%, ± 2%, or ± 1% about a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

Throughout this specification, unless the context requires otherwise, the words “comprise”, “comprises” and “comprising” will be understood to imply the inclusion of
5 a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements. By “consisting of” is meant including, and limited to, whatever follows the phrase “consisting of.” Thus, the phrase “consisting of” indicates that the listed elements are required or mandatory, and that no other elements may be present. By “consisting essentially of” is meant including any
10 elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the phrase “consisting essentially of” indicates that the listed elements are required or mandatory, but that no other elements are present that materially affect the activity or action of the listed elements.

Reference throughout this specification to “one embodiment,” “an embodiment,” “a particular embodiment,” “a related embodiment,” “a certain embodiment,” “an additional embodiment,” or “a further embodiment” or combinations thereof means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present
20 invention. Thus, the appearances of the foregoing phrases in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments. It is also understood that the positive recitation of a feature in one embodiment, serves as a basis for excluding the feature in
25 a particular embodiment.

“Polypeptide,” “polypeptide fragment,” “peptide” and “protein” are used interchangeably, unless specified to the contrary, and according to conventional meaning, *i.e.*, as a sequence of amino acids. Polypeptides are not limited to a specific length, *e.g.*, they may comprise a full-length protein sequence or a fragment of a full
30 length protein, and may include post-translational modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. In various embodiments, the CAR polypeptides contemplated herein comprise a signal

(or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. Illustrative examples of suitable signal sequences useful in CARs disclosed herein include, but are not limited to the IgG1 heavy chain signal sequence and the CD8 α signal sequence. Polypeptides can be prepared using any of a variety of well-known recombinant and/or synthetic techniques. Polypeptides contemplated herein specifically encompass the CARs of the present disclosure, or sequences that have deletions from, additions to, and/or substitutions of one or more amino acid of a CAR as disclosed herein. In particular embodiments, the term “polypeptide” further includes variants, fragments, and fusion polypeptides

10 An “isolated peptide” or an “isolated polypeptide” and the like, as used herein, refer to *in vitro* isolation and/or purification of a peptide or polypeptide molecule from a cellular environment, and from association with other components of the cell, *i.e.*, it is not significantly associated with *in vivo* substances. Similarly, an “isolated cell” refers to a cell that has been obtained from an *in vivo* tissue or organ and is substantially free of extracellular matrix.

Polypeptide variants may differ from a naturally occurring polypeptide in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the above polypeptide sequences. For example, in particular embodiments, it may be desirable to improve the binding affinity and/or other biological properties of the CARs by introducing one or more substitutions, deletions, additions and/or insertions into a binding domain, hinge, TM domain, co-stimulatory signaling domain or primary signaling domain of a CAR polypeptide. Preferably, polypeptides of the invention include polypeptides having at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% amino acid identity thereto.

Polypeptides include “polypeptide fragments.” Polypeptide fragments refer to a polypeptide, which can be monomeric or multimeric and that has an amino-terminal deletion, a carboxyl-terminal deletion, and/or an internal deletion or substitution of a naturally-occurring or recombinantly-produced polypeptide. In certain embodiments, a polypeptide fragment can comprise an amino acid chain at least 5 to about 500 amino acids long. It will be appreciated that in certain embodiments, fragments are at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30,

31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 150, 200, 250, 300, 350, 400, or 450 amino acids long.

Fusion polypeptides and fusion proteins refer to a polypeptide having at least two, three, four, five, six, seven, eight, nine, or ten or more polypeptide segments.

5 As used herein, the terms “polynucleotide” or “nucleic acid” refers to messenger RNA (mRNA), RNA, genomic RNA (gRNA), plus strand RNA (RNA(+)), minus strand RNA (RNA(-)), genomic DNA (gDNA), complementary DNA (cDNA) or recombinant DNA. Polynucleotides include single and double stranded polynucleotides. Preferably, polynucleotides of the invention include polynucleotides
10 or variants having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to any of the reference sequences described herein (*see, e.g.*, Sequence Listing), typically where the variant maintains at least one biological activity of the reference sequence. In various illustrative embodiments, the present invention contemplates, in part, polynucleotides
15 comprising expression vectors, viral vectors, and transfer plasmids, and compositions, and cells comprising the same.

As used herein, “isolated polynucleotide” refers to a polynucleotide that has been purified from the sequences which flank it in a naturally-occurring state, *e.g.*, a
DNA fragment that has been removed from the sequences that are normally adjacent to
20 the fragment. An “isolated polynucleotide” also refers to a complementary DNA (cDNA), a recombinant DNA, or other polynucleotide that does not exist in nature and that has been made by the hand of man.

The “control elements” or “regulatory sequences” present in an expression vector are those non-translated regions of the vector—origin of replication, selection
25 cassettes, promoters, enhancers, translation initiation signals (Shine Dalgarno sequence or Kozak sequence) introns, a polyadenylation sequence, 5' and 3' untranslated regions—which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation
30 elements, including ubiquitous promoters and inducible promoters may be used.

An “endogenous” control sequence is one which is naturally linked with a given gene in the genome. An “exogenous” control sequence is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological

techniques) such that transcription of that gene is directed by the linked enhancer/promoter. A “heterologous” control sequence is an exogenous sequence that is from a different species than the cell being genetically manipulated.

The term “promoter” as used herein refers to a recognition site of a
5 polynucleotide (DNA or RNA) to which an RNA polymerase binds. An RNA polymerase initiates and transcribes polynucleotides operably linked to the promoter. In particular embodiments, promoters operative in mammalian cells comprise an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated and/or another sequence found 70 to 80 bases upstream from
10 the start of transcription, a CNCAAT region where N may be any nucleotide.

The term “enhancer” refers to a segment of DNA which contains sequences capable of providing enhanced transcription and in some instances can function independent of their orientation relative to another control sequence. An enhancer can function cooperatively or additively with promoters and/or other enhancer elements.
15 The term “promoter/enhancer” refers to a segment of DNA which contains sequences capable of providing both promoter and enhancer functions.

The term “operably linked”, refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. In one embodiment, the term refers to a functional linkage between a nucleic acid
20 expression control sequence (such as a promoter, and/or enhancer) and a second polynucleotide sequence, *e.g.*, a polynucleotide-of-interest, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

The term “vector” is used herein to refer to a nucleic acid molecule capable
25 transferring or transporting another nucleic acid molecule.

The term “lentiviral vector” refers to a viral vector or plasmid containing structural and functional genetic elements, or portions thereof, including LTRs that are primarily derived from a lentivirus.

In particular embodiments, the terms “lentiviral vector,” “lentiviral expression
30 vector” may be used to refer to lentiviral transfer plasmids and/or infectious lentiviral particles. Where reference is made herein to elements such as cloning sites, promoters, regulatory elements, heterologous nucleic acids, *etc.*, it is to be understood that the

sequences of these elements are present in RNA form in the lentiviral particles and are present in DNA form in the DNA plasmids.

“Self-inactivating” (SIN) vectors refers to replication-defective vectors, *e.g.*, retroviral or lentiviral vectors, in which the right (3') LTR enhancer-promoter region, known as the U3 region, has been modified (*e.g.*, by deletion or substitution) to prevent viral transcription beyond the first round of viral replication. Self-inactivation is preferably achieved through in the introduction of a deletion in the U3 region of the 3' LTR of the vector DNA, *i.e.*, the DNA used to produce the vector RNA. Thus, during reverse transcription, this deletion is transferred to the 5' LTR of the proviral DNA. In particular embodiments, it is desirable to eliminate enough of the U3 sequence to greatly diminish or abolish altogether the transcriptional activity of the LTR, thereby greatly diminishing or abolishing the production of full-length vector RNA in transduced cells. In the case of HIV based lentivectors, it has been discovered that such vectors tolerate significant U3 deletions, including the removal of the LTR TATA box (*e.g.*, deletions from -418 to -18), without significant reductions in vector titers.

C. ANTI-BCMA CHIMERIC ANTIGEN RECEPTORS

In various embodiments, genetically engineered receptors that redirect cytotoxicity of T cells toward BCMA expressing cells are provided. These genetically engineered receptors referred to herein as anti-BCMA chimeric antigen receptors (CARs).

Anti-BCMA CARs contemplated in particular embodiments, comprise a signal peptide, an anti-BCMA scFv, a CD8 α hinge domain, a CD8 α transmembrane domain, a 4-1BB co-stimulatory domain, and a CD3 ζ primary signaling domain.

“Single-chain Fv” or “scFv” antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain and in either orientation (*e.g.*, VL-VH or VH-VL). Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding.

“V_H” or “VH” refer to the variable region of an immunoglobulin heavy chain. “V_L” or “VL” refer to the variable region of an immunoglobulin light chain. In various embodiments, the scFv comprises a light chain sequence set forth in SEQ ID NO: 7 and a heavy chain sequence set forth in SEQ ID NO: 8.

Light and heavy chain variable regions contain three hypervariable regions, also called “complementarity-determining regions” or “CDRs.” In particular embodiments, the scFv light chain comprises the CDRL1-3 sequences set forth in SEQ ID NOs: 1-3, and the scFv heavy chain comprises the CDRH1-3 sequences set forth in SEQ ID NOs:

5 4-6.

In certain embodiments, the CARs contemplated herein may comprise linker residues between the various domains, *e.g.*, added for appropriate spacing and conformation of the molecule. In particular embodiments, the linker comprises the following amino acid sequence: GSTSGSGKPGSGEGSTKG (SEQ ID NO: 22)

10 (Cooper *et al.*, *Blood*, 101(4): 1637-1644 (2003)).

In particular embodiments, the hinge domain comprises a CD8 α hinge region.

The “transmembrane domain” is the portion of the CAR that fuses the extracellular binding portion and intracellular signaling domain and anchors the CAR to the plasma membrane of the T cell. In particular embodiments, the transmembrane

15 domain comprises a CD8 α transmembrane region.

In particular embodiments, a CAR contemplated herein comprises a “co-stimulatory signaling domain” and a “primary signaling domain.” As used herein, the term, “co-stimulatory signaling domain,” or “co-stimulatory domain,” refers to an intracellular signaling domain of a co-stimulatory molecule. Primary signaling domains regulate primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary signaling domains that act in a stimulatory manner may contain signaling motifs which are known as immunoreceptor tyrosine-based activation motifs or ITAMs. In particular embodiments, the anti-BCMA CAR comprises a CD137 co-stimulatory signaling domain and a CD3 ζ primary signaling domain.

25 In preferred embodiments, the anti-BCMA CAR comprises the amino acid sequence set forth in SEQ ID NO: 9.

In particular embodiments, a polynucleotide encoding an anti-BCMA CAR polypeptides set forth in SEQ ID NO: 9 is provided.

The polynucleotides contemplated in particular embodiments, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters and/or enhancers, untranslated regions (UTRs), signal sequences, Kozak sequences, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, internal ribosomal entry sites (IRES), recombinase recognition sites (*e.g.*, LoxP,

FRT, and Att sites), termination codons, transcriptional termination signals, and polynucleotides encoding self-cleaving polypeptides, epitope tags, as disclosed elsewhere herein or as known in the art, such that their overall length may vary considerably. It is therefore contemplated that a polynucleotide fragment of almost any
5 length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

In particular embodiments, a vector encoding an anti-BCMA CAR contemplated herein is a viral vector, will include exogenous, endogenous, or heterologous control sequences such as promoters and/or enhancers.

10 In a particular embodiment, the vector comprises a myeloproliferative sarcoma virus enhancer, negative control region deleted, dl587rev primer-binding site substituted (MND) promoter (Challita *et al.*, *J Virol.* 69(2):748-55 (1995)) operably linked to a polynucleotide encoding an anti-BCMA CAR contemplated herein.

In particular embodiments, a cell (*e.g.*, a T cell) is transduced with a retroviral
15 vector, *e.g.*, a lentiviral vector, encoding an anti-BCMA CAR contemplated herein. In preferred embodiments, the lentiviral vector is HIV (human immunodeficiency virus; including HIV type 1). In particular preferred embodiments, the lentiviral vectors is a SIN HIV-1 vector.

D. ANTI-BCMA CAR T CELLS

20 The present disclosure contemplates, in particular embodiments, T cells comprising an anti-BCMA CAR, *e.g.*, anti-BCMA CAR T cells. The terms “T cell” or “T lymphocyte” are art-recognized and are intended to include thymocytes, immature T lymphocytes, mature T lymphocytes, resting T lymphocytes, or activated T lymphocytes. A T cell can be a T helper (Th) cell, for example a T helper 1 (Th1) or a
25 T helper 2 (Th2) cell. The T cell can be a helper T cell (HTL; CD4⁺ T cell) CD4⁺ T cell, a cytotoxic T cell (CTL; CD8⁺ T cell), CD4⁺CD8⁺ T cell, CD4⁺CD8⁻ T cell, or any other subset of T cells. Other illustrative populations of T cells suitable for use in particular embodiments include naïve T cells and memory T cells.

T cells of the invention can be autologous/autogeneic (“self”) or non-autologous
30 (“non-self,” *e.g.*, allogeneic, syngeneic or xenogeneic). “Autologous,” as used herein, refers to cells from the same subject. “Allogeneic,” as used herein, refers to cells of the same species that differ genetically to the cell in comparison. “Syngeneic,” as used

herein, refers to cells of a different subject that are genetically identical to the cell in comparison. "Xenogeneic," as used herein, refers to cells of a different species to the cell in comparison. In preferred embodiments, the cells of the invention are allogeneic.

T cells can be obtained from a number of sources including, but not limited to,
5 peripheral blood mononuclear cells, bone marrow, lymph nodes tissue, cord blood, thymus issue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In certain embodiments, T cells can be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled person, such as sedimentation, *e.g.*, FICOLL™ separation. In one embodiment, cells from the
10 circulating blood of an individual are obtained by apheresis. The apheresis product typically contains lymphocytes, including T cells, monocytes, granulocyte, B cells, other nucleated white blood cells, red blood cells, and platelets. In one embodiment, the cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing. The cells
15 can be washed with PBS or with another suitable solution that lacks calcium, magnesium, and most, if not all other, divalent cations. As would be appreciated by those of ordinary skill in the art, a washing step may be accomplished by methods known to those in the art, such as by using a semiautomated flowthrough centrifuge. For example, the Cobe 2991 cell processor, the Baxter CytoMate, or the like. After
20 washing, the cells may be resuspended in a variety of biocompatible buffers or other saline solution with or without buffer. In certain embodiments, the undesirable components of the apheresis sample may be removed in the cell directly resuspended culture media.

T cells can be genetically modified following isolation using known methods, or
25 the T cells can be activated and expanded (or differentiated in the case of progenitors) *in vitro* prior to being genetically modified. In a particular embodiment, the T cells are genetically modified with the chimeric antigen receptors contemplated herein (*e.g.*, transduced with a viral vector comprising a nucleic acid encoding a CAR) and then are activated and expanded *in vitro*. In various embodiments, T cells can be activated and
30 expanded before or after genetic modification to express a CAR.

In one embodiment, T cells expressing an anti-BCMA CAR are cryopreserved such that the cells remain viable upon thawing. As used herein, "cryopreserving," refers to the preservation of cells by cooling to sub-zero temperatures, such as

(typically) 77 K or -196° C. (the boiling point of liquid nitrogen). Cryoprotective agents are often used at sub-zero temperatures to prevent the cells being preserved from damage due to freezing at low temperatures or warming to room temperature.

Cryopreservative agents and optimal cooling rates can protect against cell injury.

- 5 Cryoprotective agents which can be used include but are not limited to dimethyl sulfoxide (DMSO) (Lovelock and Bishop, *Nature*, 1959; 183: 1394-1395; Ashwood-Smith, *Nature*, 1961; 190: 1204-1205), glycerol, polyvinylpyrrolidone (Rinfret, *Ann. N.Y. Acad. Sci.*, 1960; 85: 576), and polyethylene glycol (Sloviter and Ravdin, *Nature*, 1962; 196: 48). The preferred cooling rate is 1° to 3° C/minute. After at least two hours,
10 the T cells have reached a temperature of -80° C. and can be placed directly into liquid nitrogen (-196° C.) for permanent storage such as in a long-term cryogenic storage vessel.

E. COMPOSITIONS AND FORMULATIONS

- The compositions contemplated herein comprise a therapeutically effective
15 amount of anti-BCMA CAR T cells. Compositions include, but are not limited to pharmaceutical compositions. A “pharmaceutical composition” refers to a composition formulated in pharmaceutically-acceptable or physiologically-acceptable solutions for administration to a cell or an animal, either alone, or in combination with one or more other modalities of therapy. It will also be understood that, if desired, the compositions
20 may be administered in combination with other agents as well, such as, *e.g.*, cytokines, growth factors, hormones, small molecules, chemotherapeutics, pro-drugs, drugs, antibodies, or other various pharmaceutically-active agents. There is virtually no limit to other components that may also be included in the compositions, provided that the additional agents do not adversely affect the ability of the composition to deliver the
25 intended therapy.

- The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or
30 complication, commensurate with a reasonable benefit/risk ratio.

As used herein “pharmaceutically acceptable carrier” includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative,

dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, surfactant, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals. Exemplary pharmaceutically acceptable carriers
5 include, but are not limited to, to sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; tragacanth; malt; gelatin; talc; cocoa butter, waxes, animal and vegetable fats, paraffins, silicones, bentonites, silicic acid, zinc oxide; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil,
10 olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and any other compatible substances employed in
15 pharmaceutical formulations.

In particular embodiments, compositions comprise an amount, and more preferably a therapeutically effective amount, of anti-BCMA CAR-expressing T cells contemplated herein.

As used herein, the term “amount” or “dose” refers to “an amount effective,” “a
20 dose effective,” “an effective amount,” or “an effective dose” of an anti-BCMA CAR T cell sufficient to achieve a beneficial or desired prophylactic or therapeutic result, including clinical results.

A “therapeutically effective amount” or “therapeutically effective dose” of an anti-BCMA CAR T cell is also one in which any toxic or detrimental effects of an anti-
25 BCMA CAR T cell, *e.g.*, CRS, are outweighed by the therapeutically beneficial effects. The term “therapeutically effective amount” includes an amount that is effective to “treat” a subject (*e.g.*, a patient). In one embodiment, the therapeutically effective dose is the minimal effective dose (MED) of anti-BCMA CAR T cells to treat multiple myeloma in a subject. In one embodiment, the therapeutically effective dose is the maximum
30 tolerated dose (MTD) of anti-BCMA CAR T cells that does not lead to unresolvable CRS in a subject.

In particular embodiments, compositions are preferably formulated for parenteral administration, *e.g.*, intravascular (intravenous or intraarterial), administration. In a

preferred embodiment, the compositions contemplated herein are intravenously infused into the subject in a single dose.

In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is at least about 5.0×10^7 cells, at least about 15.0×10^7 cells, at
5 least about 45.0×10^7 cells, at least about 80.0×10^7 cells, or at least about 12.0×10^8 cells.

In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is greater than about 5.0×10^7 cells, greater than about 15.0×10^7 cells, greater than about 45.0×10^7 cells, greater than about 80.0×10^7 cells, or greater than about 12.0×10^8 cells.

10 In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is between about 5.0×10^7 cells to about 15.0×10^7 cells, between about 5.0×10^7 cells to about 45.0×10^7 cells, between about 5.0×10^7 cells to about 80.0×10^7 cells, or between about 5.0×10^7 cells to about 12.0×10^8 cells.

In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition
15 administered to a subject is between about 15.0×10^7 cells to about 45.0×10^7 cells, between about 15.0×10^7 cells to about 80.0×10^7 cells, or between about 15.0×10^7 cells to about 12.0×10^8 cells.

In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is at least 5.0×10^7 cells $\pm 20\%$, at least 15.0×10^7 cells $\pm 20\%$, at
20 least 45.0×10^7 cells $\pm 20\%$, at least 80.0×10^7 cells, $\pm 20\%$ or at least 12.0×10^8 cells $\pm 20\%$.

In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is greater than 5.0×10^7 cells $\pm 20\%$, at least 15.0×10^7 cells $\pm 20\%$, at least 45.0×10^7 cells $\pm 20\%$, at least 80.0×10^7 cells $\pm 20\%$, or at least 12.0×10^8
25 cells $\pm 20\%$.

In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is between 5.0×10^7 cells $\pm 20\%$ to 15.0×10^7 cells $\pm 20\%$, between 5.0×10^7 cells $\pm 20\%$ to 45.0×10^7 cells $\pm 20\%$, between 5.0×10^7 cells $\pm 20\%$ to 80.0×10^7 cells $\pm 20\%$, or between 5.0×10^7 cells $\pm 20\%$ to 12.0×10^8 cells $\pm 20\%$.

30 In one embodiment, the amount of anti-BCMA CAR⁺ T cells in a composition administered to a subject is between 15.0×10^7 cells $\pm 20\%$ to 45.0×10^7 cells $\pm 20\%$, between 15.0×10^7 cells $\pm 20\%$ to 80.0×10^7 cells $\pm 20\%$, or between 15.0×10^7 cells $\pm 20\%$ to 12.0×10^8 cells $\pm 20\%$.

For uses provided herein, the cells are generally in a volume of a liter or less, can be 500 mLs or less, even 250 mLs or 100 mLs or less.

In particular embodiments, pharmaceutical compositions comprise a therapeutically effective amount of anti-BCMA CAR T cells, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients.

Pharmaceutical compositions comprising a therapeutically effective dose of anti-BCMA CAR T cells may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (*e.g.*, aluminum hydroxide); and preservatives.

The liquid pharmaceutical compositions, whether they be solutions, suspensions or other like form, may include one or more of the following: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. An injectable pharmaceutical composition is preferably sterile.

In one embodiment, anti-BCMA CAR T cell compositions contemplated herein are formulated in a pharmaceutically acceptable cell culture medium. Such compositions are suitable for administration to human subjects. In particular embodiments, the pharmaceutically acceptable cell culture medium is a serum free medium.

Serum-free medium has several advantages over serum containing medium, including a simplified and better defined composition, a reduced degree of contaminants, elimination of a potential source of infectious agents, and lower cost. In various embodiments, the serum-free medium is animal-free, and may optionally be protein-free. Optionally, the medium may contain biopharmaceutically acceptable recombinant proteins. "Animal-free" medium refers to medium wherein the components are derived from non-

animal sources. Recombinant proteins replace native animal proteins in animal-free medium and the nutrients are obtained from synthetic, plant or microbial sources. “Protein-free” medium, in contrast, is defined as substantially free of protein.

Illustrative examples of serum-free media used in particular compositions includes, but is not limited to QBSF-60 (Quality Biological, Inc.), StemPro-34 (Life Technologies), and X-VIVO 10.

In one preferred embodiment, compositions comprising anti-BCMA CAR T cells contemplated herein are formulated in a solution comprising PlasmaLyte A.

In another preferred embodiment, compositions comprising anti-BCMA CAR T cells contemplated herein are formulated in a solution comprising a cryopreservation medium. For example, cryopreservation media with cryopreservation agents may be used to maintain a high cell viability outcome post-thaw. Illustrative examples of cryopreservation media used in particular compositions includes, but is not limited to, CryoStor CS10, CryoStor CS5, and CryoStor CS2.

In a more preferred embodiment, compositions comprising anti-BCMA CAR T cells contemplated herein are formulated in a solution comprising 50:50 PlasmaLyte A to CryoStor CS10.

F. THERAPEUTIC METHODS

Methods for treating multiple myeloma contemplated herein comprise administering a composition comprising a therapeutically effective amount of T cells that express an anti-BCMA CAR to a subject.

As used herein, the terms “individual” and “subject” are often used interchangeably and refer to a human that has multiple myeloma. In preferred embodiments, a subject refers to a human that has relapsed/refractory multiple myeloma.

As used herein, the term “patient” refers to a subject that has been diagnosed with multiple myeloma and more preferably, relapsed/refractory multiple myeloma.

As used herein “treatment” or “treating,” includes any beneficial or desirable effect on the symptoms or pathology of multiple myeloma and more preferably, relapsed/refractory multiple myeloma, and may include even minimal reductions in one or more measurable markers of multiple myeloma and more preferably, relapsed/refractory multiple myeloma. Treatment can involve optionally either the

reduction or amelioration of symptoms of a multiple myeloma, or the delaying of the progression of a multiple myeloma. "Treatment" does not necessarily indicate complete eradication or cure of a multiple myeloma, or associated symptoms thereof.

As used herein, "prevent," and similar words such as "prevented," "preventing" etc., indicate an approach for preventing, inhibiting, or reducing the likelihood of relapse of a multiple myeloma.

Multiple myeloma is a B cell malignancy of mature plasma cell morphology characterized by the neoplastic transformation of a single clone of these types of cells. These plasma cells proliferate in BM and may invade adjacent bone and sometimes the blood. Variant forms of multiple myeloma include overt multiple myeloma, smoldering multiple myeloma, plasma cell leukemia, non-secretory myeloma, IgD myeloma, osteosclerotic myeloma, solitary plasmacytoma of bone, and extramedullary plasmacytoma (see, for example, Braunwald, *et al.* (eds), *Harrison's Principles of Internal Medicine*, 15th Edition (McGraw-Hill 2001)).

In particular embodiments, compositions comprising a therapeutically effective amount of anti-BCMA CAR T cells are administered to a subject to treat relapsed/refractory multiple myeloma. "Relapse" refers to the diagnosis of return, or signs and symptoms of return, of a cancer after a period of improvement or remission. "Refractory" refers to a cancer that is resistant to, or non-responsive to, therapy with a particular therapeutic agent. A cancer can be refractory from the onset of treatment (*i.e.*, non-responsive to initial exposure to the therapeutic agent), or as a result of developing resistance to the therapeutic agent, either over the course of a first treatment period or during a subsequent treatment period.

In particular embodiments, compositions contemplated herein are administered to a subject with relapsed/refractory multiple myeloma that has been unsuccessfully treated with one, two, three or more treatments, including at least one proteasome inhibitor and/or an immunomodulatory drug (IMiD). In one embodiment, the subject's multiple myeloma is refractory to three treatment regimens, including at least one proteasome inhibitor and an IMiD. In one embodiment, the subject's multiple myeloma is double-refractory to one or more treatment regimens.

Illustrative examples of proteasome inhibitors to which subject's multiple myeloma is refractory include, but are not limited to, bortezomib, and carfilzomib.

Illustrative examples of IMiDs to which subject's multiple myeloma is refractory include, but are not limited to thalidomide, lenalidomide, and pomalidomide.

Illustrative examples of other treatments, to which multiple myeloma may be refractory include, but are not limited to, dexamethasone, and antibody-based therapies selected from the group consisting of elotuzumab, daratumumab, MOR03087, isatuximab, bevacizumab, cetuximab, siltuximab, tocilizumab, elsilimomab, azintrel, rituximab, tositumomab, milatuzumab, lucatumumab, dacetuzumab, figitumumab, dalotuzumab, AVE1642, tabalumab, pembrolizumab, pidilizumab, and nivolumab.

In one embodiment, the subject's multiple myeloma is refractory to treatment with daratumumab.

In particular embodiments, the subject's multiple myeloma is refractory to treatment with an IMiD, a proteasome inhibitor, and dexamethasone.

Methods contemplated herein, may further comprise treating a subject with relapsed/refractory multiple myeloma with an autologous hematopoietic stem cell transplant, prior to the administration of the anti-BCMA CAR T cell composition.

Methods contemplated herein, may further comprise lymphodepleting the subject prior to administration of an anti-BCMA CAR T cell composition contemplated herein, *e.g.*, for example, the lymphodepleting chemotherapy ends 1-4 days (*e.g.*, 1, 2, 3, or 4 days) prior to the administration. In particular embodiments, the lymphodepletion comprises administering one or more of melphalan, cytoxan, cyclophosphamide, and fludarabine. In one embodiment the subject is lymphodepleted with cyclophosphamide 300 mg/m² and fludarabine 30 mg/m² prior to administration of an anti-BCMA CAR T cell composition contemplated herein.

All publications, patent applications, and issued patents cited in this specification are herein incorporated by reference as if each individual publication, patent application, or issued patent were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims. The following examples are provided by way of

illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

EXAMPLES

EXAMPLE 1

CONSTRUCTION OF BCMA CARs

CARs containing murine anti-BCMA scFv antibodies were designed to contain

5 an MND promoter operably linked to anti-BMCA scFv, a hinge and transmembrane domain from CD8 α and a CD137 co-stimulatory domains followed by the intracellular signaling domain of the CD3 ζ chain. *See, e.g.*, Figure 1. The anti-BMCA CAR shown in Figure 1 comprises a CD8 α signal peptide (SP) sequence for the surface expression on T cells. The polynucleotide sequence of an exemplary anti-BMCA CAR is set forth

10 in SEQ ID NO: 10; an exemplary polypeptide sequences of a anti-BMCA CAR is set forth in SEQ ID NO: 9; and a vector map of an exemplary CAR construct is shown in Figure 1. Table 3 shows the Identity, Genbank Reference, Source Name and Citation for the various nucleotide segments of an anti-BMCA CAR lentiviral vector.

Table 3.

Nucleotides	Identity	GenBank Reference	Source Name	Citation
1-185	pUC19 plasmid backbone	Accession #L09137.2 nt 1 – 185	pUC19	New England Biolabs
185-222	Linker	Not applicable	Synthetic	Not applicable
223-800	CMV	Not Applicable	pHCMV	(1994) PNAS 91: 9564-68
801-1136	R, U5, PBS, and packaging sequences	Accession #M19921.2 nt 454-789	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43
1137-1139	Gag start codon (ATG) changed to stop codon (TAG)	Not Applicable	Synthetic	Not applicable
1140-1240	HIV-1 gag sequence	Accession #M19921.2 nt 793-893	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43

Nucleotides	Identity	GenBank Reference	Source Name	Citation
1241-1243	HIV-1 gag sequence changed to a second stop codon	Not Applicable	Synthetic	Not applicable
1244-1595	HIV-1 gag sequence	Accession #M19921.2 nt 897-1248	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43
1596-1992	HIV-1 pol cPPT/CTS	Accession #M19921.2 nt 4745-5125	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43
1993-2517	HIV-1, isolate HXB3 env region (RRE)	Accession #M14100.1 nt 1875-2399	PgTAT-CMV	Malim, M. H. Nature (1988) 335:181-183
2518-2693	HIV-1 env sequences S/A	Accession #M19921.2 nt 8290-8470	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43
2694-2708	Linker	Not applicable	Synthetic	Not applicable
2709-3096	MND	Not applicable	pccl-c- MNDU3c-x2	Challita et al. (1995) J. Virol. 69: 748- 755
3097-3124	Linker	Not applicable	Synthetic	Not applicable
3125-3187	Signal peptide	Accession # NM_001768	CD8a signal peptide	Not applicable
3188-3934	BCMA02 scFv (V _L 1- linker-V _H 0)	Not applicable	Synthetic	Not applicable
3935-4141	CD8a hinge and TM	Accession # NM_001768	CD8a hinge and TM	Milone et al (2009) Mol Ther 17(8):1453-64
4144-4269	CD137 (4-1BB) signaling domain	Accession # NM_001561	CD137 signaling domain	Milone et al (2009) Mol Ther 17(8):1453-64

Nucleotides	Identity	GenBank Reference	Source Name	Citation
4270-4606	CD3- ζ signaling domain	Accession # NM_000734	CD3- ζ signaling domain	Milone et al (2009) Mol Ther 17(8):1453-64
4607-4717	HIV-1 ppt and part of 3' U3	Accession #M19921.2 nt 9005-9110	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43
4718-4834	HIV-1 part of U3 (399bp deletion) and R	Accession #M19921.2 nt 9511-9627	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11):5732-43
4835-4858	Synthetic polyA	Not applicable	Synthetic	Levitt, N. Genes & Dev (1989) 3:1019-1025
4859-4877	Linker	Not applicable	Synthetic	Not Applicable
4878-7350	pUC19 backbone	Accession #L09137.2 nt 2636-2686	pUC19	New England Biolabs

EXAMPLE 2

ANTI-BCMA CAR T CELL MANUFACTURING PROCESS

Unique anti-BCMA02 CAR T cell products are manufactured for each patient
 5 treatment. The reliability of the manufacturing process for anti-BCMA02 CAR T cell products was evaluated by generating anti-BCMA02 CAR T cells from 11 individual normal donor PBMC. Anti-BCMA02 CAR T cell expansion from each donor was comparable to a matched untransduced culture performed in parallel (Figure 2A).

At the end of the culture period (day 10), T cell transduction efficiency was
 10 assessed by quantitating the number of integrated lentiviruses with qPCR and lentiviral-specific primer sets (vector copy number, VCN). Anti-BCMA02 CAR T cell cultures from the 11 donors showed comparable lentiviral transduction efficiency (Figure 2B). The frequency of anti-BCMA02 CAR positive T cells was measured by flow cytometry and BCMA expression was found to be comparable across all donors (Figure 2C).

The activity of each anti-BCMA02 CAR T cell product was assessed by IFN γ -release after co-culture with K562 cells engineered to express BCMA. All anti-BCMA CAR02 T cell products exhibited therapeutically relevant levels of IFN γ release when exposed to BCMA-expressing K562 cells (Figure 2D).

5

EXAMPLE 3

ANTI-BCMA02 CAR T CELLS SHOW THERAPEUTIC ACTIVITY

IN HUMAN CLINICAL TRIALS OF RELAPSED/REFRACTORY MULTIPLE MYELOMA

INTRODUCTION

Chimeric antigen receptor (CAR) T cell therapies have demonstrated robust and
10 sustained clinical responses observed in several hematologic malignancies. However,
outside of CD19, clinical data supporting the promise of CAR T cells has been limited.
The potential for CAR T cell safety and efficacy was tested in relapsed/refractory
multiple myeloma (MM), a patient population with limited treatment options. To
redirect T cells to MM we have targeted B cell maturation antigen (BCMA), a member
15 of the tumor necrosis factor superfamily that is near uniformly expressed only by
malignant myeloma cells, plasma cells, and some mature B cells. Initial proof of anti-
BCMA activity has recently been demonstrated using T cells transduced with a gamma-
retroviral vector encoding an anti-BCMA CAR with a CD28 costimulatory domain, but
significant cytokine release syndrome occurred in patients with high disease burden
20 (Ali *et al.*, *Blood* 2016). Because the therapeutically effective dose of the anti-
BCMA02 CAR T cells in humans is unpredictable, a dose-escalation trial was
conducted.

METHODS

Anti-BCMA02 CAR T cells were administered to patients with relapsed and/or
25 refractory BCMA-positive MM who received at least 3 prior regimens, including a
proteasome inhibitor and an immunomodulatory agent or who were double-refractory.
Briefly, peripheral blood mononuclear cells were collected via leukapheresis and
transferred to a centralized manufacturing facility for transduction and expansion.
Patients were lymphodepleted with cyclophosphamide 300 mg/m² and fludarabine 30

mg/m² on study Days -5, -4 and -3 and then received a single infusion of anti-BCMA02 CAR T cells on Day 0.

The average vector copy number (VCN) for the CAR T cells was determined for the nine patients. Table 4.

5 **Table 4. CAR T cell VCN**

Patient	VCN
1	2.08
2	1.97
3	2.11
4	5.98
5	3.01
6	4.13
7	3.38
8	4.01
9	ND

ND = not yet determined

RESULTS

Nine patients were infused with anti-BCMA02 CAR T cells in three dose cohorts of 5.0×10^7 CAR+ T cells, 15.0×10^7 CAR+ T cells, and 45.0×10^7 CAR+ T cells. Cells were collected and successfully manufactured and released in all patients. Median age at enrollment for the 9 infused subjects was 58 years (43-68) and 67% were male. The median time from diagnosis was 6 years (1.3-8.6). The median number of prior lines of therapy was 6 (4-10), 100% of patients received at least 1 prior autologous stem cell transplant, 67% received prior daratumumab or CD38 mab, 89% received prior lenalidomide, 78% received prior pomalidomide, 100% received prior bortezomib, and 78% prior carfilzomib.

To date no dose limiting toxicities, neurotoxicities > Grade 2, or Grade 3 or higher cytokine release syndrome (CRS) have been observed. Moreover, no patients have required treatment with vasopressors, tocilizumab or corticosteroids. Grade 1 or 2 CRS was reported in 6/9 (67%) treated patients. In addition to CRS, the most common treatment emergent adverse events were neutropenia (89%), leukopenia (67%) and anemia (44%).

- Clinical responses were observed in every dose cohort. After the initial cohort, all patients remained on study and the overall response rate in the 6 patients treated at 15.0×10^7 CAR+ T cells or higher was 83%, including 2 stringent complete responses (CRs) (*see* Table 5). Surprisingly, at dose levels above 5.0×10^7 CAR+ T cells, all
- 5 patients with bone marrow involvement at baseline have had no bone marrow disease detected at any point after day 14.

Table 5. Response by Dose Level & Duration

Dose	Patient	Response	Time since bb2121 infusion (weeks)
5.0×10^7	1	PR	13
	2*	SD	9
	3*	PD	8
15.0×10^7	4	sCR	28+
	5	sCR	21+
	6*	VGPR	19+
45.0×10^7	7	PR	13+
	8	SD	11+
	9*	PR	4+

** indicates patient had $\geq 50\%$ bone marrow involvement at baseline. Responses graded using IMWG Uniform Response Criteria for Multiple Myeloma. PR: partial response, SD: stable disease, PD: progressive disease, sCR: stringent complete response, VGPR: very good partial response, NE: not yet evaluated. + denotes ongoing response.*

- CAR+ T cell expansion was consistently demonstrated, and is similar to other published CAR T cell trials. In the 5 patients treated with $> 5.0 \times 10^7$ CAR+ T cells for
- 10 whom data was available, the range of peak CAR+ T cells by flow cytometry was 10 to 686 CAR+ cells/ μ L in peripheral blood and 4 to 527 CAR+ cells/ μ L in bone marrow, and the range of peak copies/ μ g genomic DNA in peripheral blood was 34,231 to 860,204 by PCR.

CONCLUSIONS

- 15 Anti-BCMA02 CAR T cells showed remarkable efficacy at dose levels above 5.0×10^7 CAR+ T cells, including 2 CRs and ongoing clinical response at 6 months. In contrast to results with other CAR T cell therapies, the efficacy of the anti-BCMA02 CAR T cells was accompanied by unexpectedly mild and manageable CRS, including

in patients with $\geq 50\%$ bone marrow involvement. These data support the therapeutic efficacy of anti-BCMA02 CAR T cells for relapsed/refractory multiple myeloma.

Additional dose cohorts are planned with dose levels of 80.0×10^7 and 12.0×10^8 CAR+ T cells.

5

EXAMPLE 4

ANTI-BCMA02 CAR T CELLS CONTINUE TO SHOW THERAPEUTIC ACTIVITY IN HUMAN CLINICAL TRIALS OF RELAPSED/REFRACTORY MULTIPLE MYELOMA

Twenty one patients have been infused with anti-BCMA02 CAR T cells in three dose cohorts of 5.0×10^7 CAR+ T cells (3 patients), 15.0×10^7 CAR+ T cells (6 patients), 45.0×10^7 CAR+ T cells (9 patients), and 80.0×10^7 CAR+ T cells (3 patients). Cells were collected and successfully manufactured and released in all patients. Median age at enrollment for the 9 infused subjects was 58 years (37-74) and 62% were male. The median time from diagnosis was 5 years (1.0-16.0). The median number of prior lines of therapy was 7 (3-14), 100% of patients received at least 1 prior autologous stem cell transplant, 100% previously treated with lenalidomide and bortezomib, 91% previously treated with pomalidomide and carfilzomib, 71% previously treated with daratumumab, and 29% of patients were penta-refractory (bortezomib, lenalidomide, carfilzomib, pomalidomide, daratumumab).

All patients in active dose cohorts achieved an objective response, duration up to 54 weeks. Figure 3. CAR T cell dose, number of evaluable patients, overall response rate, best response, median lines of prior therapy, and safety for each cohort are presented in Table 6.

Table 6: Cohort Data

Cohort	1	2	3	4
CAR+ T Cell Dose	50×10^6	150×10^6	450×10^6	800×10^6
Number of Patients Evaluable for Efficacy	3	4	8	3
Overall Response Rate in Cohort	33%	100%	100%	100%
Best Response	PD (1 patient) SD (1 patient) PR (1 patient)	CR (2 patients, 1 patient MRD negative)	CR (1 patient*) VGPR (5 patients; 1	VGPR (1 patient) PR (1 patient) CR (1 patient)

		VGPR (1 patient MRD negative) PR (1 patient)	patient MRD negative) PR (2 patients; 1 patient MRD negative) *Patient died of unrelated cardio pulmonary arrest	
Bone Marrow Involvement		All patients in cohorts 2, 3 and 4 with bone marrow involvement at baseline had no detectable multiple myeloma cells in their bone marrow on Day 14 or beyond. Of four patients evaluable for MRD status, all four were found to be MRD-negative.		
Median Prior Lines of Therapy		7 (range: 3-14); all patients had at least one prior autologous stem cell transplant, as well as prior exposure to a proteasome inhibitor and an immunomodulatory agent; 71% of patients had previously received daratumumab or CD38 antibody		
Safety		15/21 (71%) of patients had CRS, mostly Grade 1 & 2; 2 patients with Grade 3 CRS that resolved within 24 hours. 4 patients received tocilizumab, 1 (Grade 2 CRS) received steroids. The most common treatment-emergent Grade 3-4 AEs in 21 infused patients include cytopenias commonly associated with cy/flu lymphodepletion, as well as Grade 3 events of hyponatraemia (n=4), cytokine release syndrome (n=2), upper respiratory infection (n=2), and syncope (n=2).		

In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A composition comprising a therapeutically effective amount of anti-B cell maturation antigen (BCMA) chimeric antigen receptor (CAR) T cells, wherein the therapeutically effective amount is from about 15.0×10^7 to about 12.0×10^8 anti-human BCMA CAR T cells, and the anti-BCMA CAR comprises amino acids 22-493 of the amino acid sequence set forth in SEQ ID NO: 9.
2. A composition comprising a therapeutically effective amount of anti-B cell maturation antigen (BCMA) chimeric antigen receptor (CAR) T cells, wherein the therapeutically effective amount is from about 15.0×10^7 to about 12.0×10^8 anti-human BCMA CAR T cells, and the anti-BCMA CAR is encoded by the nucleotide sequence set forth in SEQ ID NO: 10.
3. The composition of claim 1 or claim 2, further comprising a pharmaceutically acceptable carrier.
4. The composition of any one of claims 1 to 3, formulated in a solution comprising a cryopreservation medium.
5. The composition of any one of claims 1 to 4, wherein the therapeutically effective amount is:
 - (a) about 15.0×10^7 anti-human BCMA CAR T cells;
 - (b) about 45.0×10^7 anti-human BCMA CAR T cells;
 - (c) about 80.0×10^7 anti-human BCMA CAR T cells; or
 - (d) about 12.0×10^8 anti-human BCMA CAR T cells.
6. The composition of any one of claims 1 to 4, wherein the therapeutically effective amount is:

- (a) from about 15.0×10^7 anti-human BCMA CAR T cells to about 45.0×10^7 anti-BCMA CAR T cells; or
- (b) from about 15.0×10^7 anti-human BCMA CAR T cells to about 80.0×10^7 anti-BCMA CAR T cells

7. The composition of any one of claims 1-6, wherein the anti-BCMA CAR T cells are transduced with a lentiviral vector encoding the anti-human BCMA CAR.

8. The composition of claim 7, wherein the lentiviral vector copy number (VCN) is about 2.0 copies per anti-human BCMA CAR T cell.

9. The composition of claim 7 or claim 8, wherein the lentiviral vector is a human immunodeficiency virus 1 (HIV-1) vector.

10. The composition of any one of claims 1 to 9, wherein the anti-human BCMA CAR comprises the amino acid sequence set forth in SEQ ID NO: 9.

11. A method of treating a human subject that has been diagnosed with relapsed/refractory multiple myeloma comprising administering to the subject the composition of any one of claims 1 to 10.

12. A method of treating a human subject that has relapsed/refractory multiple myeloma comprising administering to the subject the composition of any one of claims 1 to 10.

13. The method of claim 11 or claim 12, wherein the composition is administered in a single dose.

14. The method of any one of claims 11 to 13, wherein the composition is intravenously administered.

15. The method of any one of claims 11 to 14, wherein the multiple myeloma was refractory to at least three treatment regimens, including a proteasome inhibitor and an immunomodulatory agent, prior to the administration of the composition.

16. The method of any one of claims 11 to 14, wherein the multiple myeloma was double-refractory to one or more treatment regimens, prior to the administration of the composition.

17. The method of any one of claims 11 to 14, wherein:

- (a) the subject was treated with daratumumab, lenalidomide, pomalidomide, bortezomib, and/or carfilzomib, prior to the administration of the composition;
- (b) the subject received an autologous hematopoietic stem cell transplant, prior to the administration of the composition; and/or
- (c) the subject was lymphodepleted with cyclophosphamide 300 mg/m² and fludarabine 30 mg/m².

18. Use of the composition of any one of claims 1 to 10 in the manufacture of a medicament for the treatment of relapsed/refractory multiple myeloma in a human subject.

19. The use of claim 18, wherein the composition is administered in a single dose and/or intravenously.

20. The use of claim 18 or 19, wherein the multiple myeloma was refractory to at least three treatment regimens, including a proteasome inhibitor and an immunomodulatory agent, prior to the administration of the composition.

21. The use of claim 18 or 19, wherein the multiple myeloma was double-refractory to one or more treatment regimens, prior to the administration of the composition.

22. The use of claim 18 or 19, wherein:

(a) the subject was treated with daratumumab, lenalidomide, pomalidomide, bortezomib, and/or carfilzomib, prior to the administration of the composition;

(b) the subject received an autologous hematopoietic stem cell transplant, prior to the administration of the composition; and/or

(c) the subject was lymphodepleted with cyclophosphamide 300 mg/m² and fludarabine 30 mg/m².




FIGURE 1




FIGURE 2A




FIGURE 2B




FIGURE 2C




FIGURE 2D




FIGURE 3

BLBD_079_02WO_ST25.TXT
SEQUENCE LISTING

<110> bluebird bio, Inc.
Quigley, Travis
Ross, Robert

<120> ANTI-BCMA CAR T CELL COMPOSITIONS

<130> BLBD-079/02WO

<150> US 62/514,401

<151> 2017-06-02

<150> US 62/417,840

<151> 2016-11-04

<160> 36

<170> PatentIn version 3.5

<210> 1

<211> 15

<212> PRT

<213> Mus musculus

<400> 1

Arg	Ala	Ser	Glu	Ser	Val	Thr	Ile	Leu	Gly	Ser	His	Leu	Ile	His
1				5					10					15

<210> 2

<211> 7

<212> PRT

<213> mus musculus

<400> 2

Leu	Ala	Ser	Asn	Val	Gln	Thr
1				5		

<210> 3

<211> 9

<212> PRT

<213> mus musculus

<400> 3

Leu	Gln	Ser	Arg	Thr	Ile	Pro	Arg	Thr
1				5				

BLBD_079_02WO_ST25.TXT

<210> 4
 <211> 5
 <212> PRT
 <213> Mus musculus

<400> 4

Asp Tyr Ser Ile Asn
 1 5

<210> 5
 <211> 17
 <212> PRT
 <213> Mus musculus

<400> 5

Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe Arg
 1 5 10 15

Gly

<210> 6
 <211> 8
 <212> PRT
 <213> Mus musculus

<400> 6

Asp Tyr Ser Tyr Ala Met Asp Tyr
 1 5

<210> 7
 <211> 111
 <212> PRT
 <213> Mus musculus

<400> 7

Asp Ile Val Leu Thr Gln Ser Pro Pro Ser Leu Ala Met Ser Leu Gly
 1 5 10 15

Lys Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Thr Ile Leu
 20 25 30

BLBD_079_02WO_ST25.TXT

Gly Ser His Leu Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45

Thr Leu Leu Ile Gln Leu Ala Ser Asn Val Gln Thr Gly Val Pro Ala
50 55 60

Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp
65 70 75 80

Pro Val Glu Glu Asp Asp Val Ala Val Tyr Tyr Cys Leu Gln Ser Arg
85 90 95

Thr Ile Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 8
<211> 117
<212> PRT
<213> Mus musculus

<400> 8

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Ser Ile Asn Trp Val Lys Arg Ala Pro Gly Lys Gly Leu Lys Trp Met
35 40 45

Gly Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe
50 55 60

Arg Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80

Leu Gln Ile Asn Asn Leu Lys Tyr Glu Asp Thr Ala Thr Tyr Phe Cys
85 90 95

BLBD_079_02WO_ST25.TXT

Ala Leu Asp Tyr Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser
100 105 110

Val Thr Val Ser Ser
115

<210> 9
<211> 493
<212> PRT
<213> Artificial Sequence

<220>
<223> anti-BCMA02 CAR

<400> 9

Met Ala Leu Pro Val Thr Ala Leu Leu Leu Pro Leu Ala Leu Leu Leu
1 5 10 15

His Ala Ala Arg Pro Asp Ile Val Leu Thr Gln Ser Pro Pro Ser Leu
20 25 30

Ala Met Ser Leu Gly Lys Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu
35 40 45

Ser Val Thr Ile Leu Gly Ser His Leu Ile His Trp Tyr Gln Gln Lys
50 55 60

Pro Gly Gln Pro Pro Thr Leu Leu Ile Gln Leu Ala Ser Asn Val Gln
65 70 75 80

Thr Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe
85 90 95

Thr Leu Thr Ile Asp Pro Val Glu Glu Asp Asp Val Ala Val Tyr Tyr
100 105 110

Cys Leu Gln Ser Arg Thr Ile Pro Arg Thr Phe Gly Gly Gly Thr Lys
115 120 125

Leu Glu Ile Lys Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly
130 135 140

BLBD_079_02WO_ST25.TXT

Glu Gly Ser Thr Lys Gly Gln Ile Gln Leu Val Gln Ser Gly Pro Glu
145 150 155 160

Leu Lys Lys Pro Gly Glu Thr Val Lys Ile Ser Cys Lys Ala Ser Gly
165 170 175

Tyr Thr Phe Thr Asp Tyr Ser Ile Asn Trp Val Lys Arg Ala Pro Gly
180 185 190

Lys Gly Leu Lys Trp Met Gly Trp Ile Asn Thr Glu Thr Arg Glu Pro
195 200 205

Ala Tyr Ala Tyr Asp Phe Arg Gly Arg Phe Ala Phe Ser Leu Glu Thr
210 215 220

Ser Ala Ser Thr Ala Tyr Leu Gln Ile Asn Asn Leu Lys Tyr Glu Asp
225 230 235 240

Thr Ala Thr Tyr Phe Cys Ala Leu Asp Tyr Ser Tyr Ala Met Asp Tyr
245 250 255

Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Ala Ala Thr Thr
260 265 270

Thr Pro Ala Pro Arg Pro Pro Thr Pro Ala Pro Thr Ile Ala Ser Gln
275 280 285

Pro Leu Ser Leu Arg Pro Glu Ala Cys Arg Pro Ala Ala Gly Gly Ala
290 295 300

Val His Thr Arg Gly Leu Asp Phe Ala Cys Asp Ile Tyr Ile Trp Ala
305 310 315 320

Pro Leu Ala Gly Thr Cys Gly Val Leu Leu Leu Ser Leu Val Ile Thr
325 330 335

Leu Tyr Cys Lys Arg Gly Arg Lys Lys Leu Leu Tyr Ile Phe Lys Gln
340 345 350

BLBD_079_02WO_ST25.TXT

Pro Phe Met Arg Pro Val Gln Thr Thr Gln Glu Glu Asp Gly Cys Ser
355 360 365

Cys Arg Phe Pro Glu Glu Glu Glu Gly Gly Cys Glu Leu Arg Val Lys
370 375 380

Phe Ser Arg Ser Ala Asp Ala Pro Ala Tyr Gln Gln Gly Gln Asn Gln
385 390 395 400

Leu Tyr Asn Glu Leu Asn Leu Gly Arg Arg Glu Glu Tyr Asp Val Leu
405 410 415

Asp Lys Arg Arg Gly Arg Asp Pro Glu Met Gly Gly Lys Pro Arg Arg
420 425 430

Lys Asn Pro Gln Glu Gly Leu Tyr Asn Glu Leu Gln Lys Asp Lys Met
435 440 445

Ala Glu Ala Tyr Ser Glu Ile Gly Met Lys Gly Glu Arg Arg Arg Gly
450 455 460

Lys Gly His Asp Gly Leu Tyr Gln Gly Leu Ser Thr Ala Thr Lys Asp
465 470 475 480

Thr Tyr Asp Ala Leu His Met Gln Ala Leu Pro Pro Arg
485 490

<210> 10
<211> 1485
<212> DNA
<213> Artificial Sequence

<220>
<223> anti-BCMA02 CAR

<400> 10
atggcactcc ccgtcaccgc ctttctcttg cccctcgccc tgctgctgca tgctgccagg 60
cccgacattg tgctcactca gtcacctccc agcctggcca tgagcctggg aaaaagggcc 120
accatctcct gtagagccag tgagtccgtc acaatcttgg ggagccatct tattcactgg 180

BLBD_079_02WO_ST25.TXT

tatcagcaga agcccgggca gcctccaacc cttcttattc agctcgcgtc aaacgtccag	240
acgggtgtac ctgccagatt ttctggtagc ggggtcccgca ctgattttac actgaccata	300
gatccagtgg aagaagacga tgtggccgtg tattattgtc tgcagagcag aacgattcct	360
cgcacatttg gtgggggtac taagctggag attaagggaa gcacgtccgg ctcagggaag	420
ccgggctccg gcgagggaag cacgaagggg caaattcagc tgggtccagag cggacctgag	480
ctgaaaaaac ccggcgagac tgttaagatc agttgtaaag catctggcta taccttcacc	540
gactacagca taaattgggt gaaacgggcc cctggaaagg gcctcaaag gatgggttg	600
atcaataccg aaactagga gcctgcttat gcatatgact tccgcgggag attcgccttt	660
tcactcgaga catctgcctc tactgcttac ctcaaataa acaacctcaa gtatgaagat	720
acagccactt acttttgcgc cctcgactat agttacgcca tggactactg gggacaggga	780
acctccgtta ccgtcagttc cgcggccgca accacaacac ctgctccaag gccccccaca	840
cccgctcaa ctatagccag ccaaccattg agcctcagac ctgaagcttg caggcccgca	900
gcaggaggcg ccgtccatac gcgaggcctg gacttcgctg gtgatattta tatttgggcc	960
cctttggccg gaacatgtgg ggtgttgctt ctctcccttg tgatcactct gtattgtaag	1020
cgcgggagaa agaagctcct gtacatcttc aagcagcctt ttatgcgacc tgtgcaaacc	1080
actcaggaag aagatgggtg ttcattgccg tttcccgagg aggaagaagg aggggtgtgaa	1140
ctgagggtga aattttctag aagcgccgat gctcccgcat atcagcaggg tcagaatcag	1200
ctctacaatg aattgaatct cggcaggcga gaagagtacg atgttctgga caagagacgg	1260
ggcagggatc ccgagatggg gggaaagccc cggagaaaaa atcctcagga ggggttgtag	1320
aatgagctgc agaaggacaa gatggctgaa gcctatagcg agatcggaat gaaaggcgaa	1380
agacgcagag gcaaggggca tgacggtctg taccagggtc tctctacagc caccaaggac	1440
acttatgatg cgttgcatat gcaagccttg ccaccccgct aatga	1485

<210> 11
 <211> 184
 <212> PRT
 <213> Homo sapiens

<400> 11

BLBD_079_02WO_ST25.TXT

Met Leu Gln Met Ala Gly Gln Cys Ser Gln Asn Glu Tyr Phe Asp Ser
1 5 10 15

Leu Leu His Ala Cys Ile Pro Cys Gln Leu Arg Cys Ser Ser Asn Thr
20 25 30

Pro Pro Leu Thr Cys Gln Arg Tyr Cys Asn Ala Ser Val Thr Asn Ser
35 40 45

Val Lys Gly Thr Asn Ala Ile Leu Trp Thr Cys Leu Gly Leu Ser Leu
50 55 60

Ile Ile Ser Leu Ala Val Phe Val Leu Met Phe Leu Leu Arg Lys Ile
65 70 75 80

Asn Ser Glu Pro Leu Lys Asp Glu Phe Lys Asn Thr Gly Ser Gly Leu
85 90 95

Leu Gly Met Ala Asn Ile Asp Leu Glu Lys Ser Arg Thr Gly Asp Glu
100 105 110

Ile Ile Leu Pro Arg Gly Leu Glu Tyr Thr Val Glu Glu Cys Thr Cys
115 120 125

Glu Asp Cys Ile Lys Ser Lys Pro Lys Val Asp Ser Asp His Cys Phe
130 135 140

Pro Leu Pro Ala Met Glu Glu Gly Ala Thr Ile Leu Val Thr Thr Lys
145 150 155 160

Thr Asn Asp Tyr Cys Lys Ser Leu Pro Ala Ala Leu Ser Ala Thr Glu
165 170 175

Ile Glu Lys Ser Ile Ser Ala Arg
180

<210> 12
<211> 5
<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 12

Asp Gly Gly Gly Ser
1 5

<210> 13

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 13

Thr Gly Glu Lys Pro
1 5

<210> 14

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 14

Gly Gly Arg Arg
1

<210> 15

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 15

Gly Gly Gly Gly Ser
1 5

<210> 16
 <211> 14
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Flexible peptide linker

<400> 16

Glu Gly Lys Ser Ser Gly Ser Gly Ser Glu Ser Lys Val Asp
 1 5 10

<210> 17
 <211> 18
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Flexible peptide linker

<400> 17

Lys Glu Ser Gly Ser Val Ser Ser Glu Gln Leu Ala Gln Phe Arg Ser
 1 5 10 15

Leu Asp

<210> 18
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Flexible peptide linker

<400> 18

Gly Gly Arg Arg Gly Gly Gly Ser
 1 5

<210> 19
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 19

Leu Arg Gln Arg Asp Gly Glu Arg Pro
1 5

<210> 20

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 20

Leu Arg Gln Lys Asp Gly Gly Gly Ser Glu Arg Pro
1 5 10

<210> 21

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 21

Leu Arg Gln Lys Asp Gly Gly Gly Ser Gly Gly Gly Ser Glu Arg Pro
1 5 10 15

<210> 22

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Flexible peptide linker

<400> 22

Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser Thr
1 5 10 15

Lys Gly

<210> 23
 <211> 7
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> protease cleavage site

<220>
 <221> MISC_FEATURE
 <222> (2)..(3)
 <223> Xaa = Any amino acid

<220>
 <221> MISC_FEATURE
 <222> (5)..(5)
 <223> Xaa = Any amino acid

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> Xaa is Gly or Ser

<400> 23

Glu Xaa Xaa Tyr Xaa Gln Xaa
 1 5

<210> 24
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> protease cleavage site

<400> 24

Glu Asn Leu Tyr Phe Gln Gly
 1 5

<210> 25
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 25

Glu Asn Leu Tyr Phe Gln Ser
1 5

<210> 26

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 26

Leu Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn
1 5 10 15

Pro Gly Pro

<210> 27

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 27

Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn
1 5 10 15

Pro Gly Pro

<210> 28

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 28

Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro Gly Pro
1 5 10

<210> 29

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 29

Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro Gly
1 5 10 15

Pro

<210> 30

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 30

Gln Leu Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser
1 5 10 15

Asn Pro Gly Pro
20

<210> 31

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 31

BLBD_079_02WO_ST25.TXT

Ala Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly
1 5 10 15

Asp Val Glu Ser Asn Pro Gly Pro
20

<210> 32

<211> 40

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 32

Val Thr Glu Leu Leu Tyr Arg Met Lys Arg Ala Glu Thr Tyr Cys Pro
1 5 10 15

Arg Pro Leu Leu Ala Ile His Pro Thr Glu Ala Arg His Lys Gln Lys
20 25 30

Ile Val Ala Pro Val Lys Gln Thr
35 40

<210> 33

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 33

Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro
1 5 10 15

Gly Pro

<210> 34

<211> 40

<212> PRT

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> protease cleavage site

<400> 34

Leu Leu Ala Ile His Pro Thr Glu Ala Arg His Lys Gln Lys Ile Val
1 5 10 15

Ala Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly
20 25 30

Asp Val Glu Ser Asn Pro Gly Pro
35 40

<210> 35

<211> 33

<212> PRT

<213> Artificial Sequence

$\langle 220 \rangle$

<223> protease cleavage site

<400> 35

Glu Ala Arg His Lys Gln Lys Ile Val Ala Pro Val Lys Gln Thr Leu
1 5 10 15

Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro Gly
20 25 30

Pro

<210> 36

<211> 7350

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> anti-BCMA02 CAR vector

<400> 36

tcgcgcgttt cggtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60

BLBD_079_02W0_ST25.TXT

cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg	120
ttggcgggtg tcggggctgg cttactatg cggcatcaga gcagattgta ctgagagtgc	180
accatcatat gccagcctat ggtgacattg attattgact agttattaat agtaatcaat	240
tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa	300
tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt	360
tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta	420
aactgcccac ttggcagtac atcaagtgt tcatatgcc agtacgcccc ctattgacgt	480
caatgacggg aaatggcccg cctggcatta tgcccagtac atgaccttat gggactttcc	540
tacttggcag tacatctacg tattagtcac cgctattacc atggtgatgc ggttttggca	600
gtacatcaat gggcgtggat agcggtttga ctacgggga tttccaagtc tccaccccat	660
tgacgtcaat gggagtttgt tttggcacca aaatcaacgg gactttcaa aatgtcgtaa	720
caactccgcc ccattgacgc aaatgggcgg taggcgtgta cgggtggagg tctatataag	780
cagagctcgt ttagtgaacc gggctctctt ggtagacca gatctgagcc tgggagctct	840
ctggctaact agggaaacca ctgcttaagc ctcaataaag cttgccttga gtgctcaaag	900
tagtgtgtgc ccgtctgttg tgtgactctg gtaactagag atccctcaga cccttttagt	960
cagtgtggaa aatctctagc agtggcgccc gaacagggac ttgaaagcga aagtaaagcc	1020
agaggagatc tctcgacgca ggactcggct tgctgaagcg cgcacggcaa gaggcgaggg	1080
gcggcgactg gtgagtacgc caaaaatttt gactagcgga ggctagaagg agagagtagg	1140
gtgcgagagc gtcggtatta agcgggggag aattagataa atgggaaaaa attcggttaa	1200
ggccaggggg aaagaaacaa tataaactaa aacatatagt tagggcaagc agggagctag	1260
aacgattcgc agttaatcct ggccttttag agacatcaga aggctgtaga caaatactgg	1320
gacagctaca accatccctt cagacaggat cagaagaact tagatcatta tataatacaa	1380
tagcagtcct ctattgtgtg catcaaagga tagatgtaaa agacaccaag gaagccttag	1440
ataagataga ggaagagcaa aacaaaagta agaaaaaggc acagcaagca gcagctgaca	1500
caggaaacaa cagccaggtc agccaaaatt accctatagt gcagaacctc caggggcaaa	1560
tggtagatca ggccatatca cctagaactt taaattaaga cagcagtaca aatggcagta	1620

BLBD_079_02WO_ST25.TXT

ttcatccaca	attttaaaag	aaaagggggg	attggggggg	acagtgcagg	ggaaagaata	1680
gtagacataa	tagcaacaga	catacaaact	aaagaattac	aaaaacaaat	tacaaaaatt	1740
caaaattttc	gggtttatta	cagggacagc	agagatccag	tttggaagg	accagcaaag	1800
ctcctctgga	aaggtgaagg	ggcagtagta	atacaagata	atagtacat	aaaagtagtg	1860
ccaagaagaa	aagcaaagat	catcagggat	tatggaaaac	agatggcagg	tgatgattgt	1920
gtggcaagta	gacaggatga	ggattaacac	atggaaaaga	ttagtaaaac	accatagctc	1980
tagagcgatc	ccgatcttca	gacctggagg	aggagatatg	agggacaatt	ggagaagtga	2040
attatataaa	tataaagtag	taaaaattga	accattagga	gtagcaccca	ccaaggcaaa	2100
gagaagagt	gtgcagagag	aaaaaagagc	agtgggaata	ggagctttgt	tccttggggt	2160
cttgggagca	gcaggaagca	ctatgggcgc	agcgtcaatg	acgctgacgg	tacaggccag	2220
acaattattg	tctggtatag	tgacgcagca	gaacaatttg	ctgagggcta	ttgaggcgca	2280
acagcatctg	ttgcaactca	cagtctgggg	catcaagcag	ctccaggcaa	gaatcctggc	2340
tgtggaaaga	tacctaaagg	atcaacagct	cctgggggatt	tggggttgct	ctggaaaact	2400
catttgcacc	actgctgtgc	cttggaatgc	tagttggagt	aataaatctc	tggaacagat	2460
ttggaatcac	acgacctgga	tggagtggga	cagagaaatt	aacaattaca	caagcttggt	2520
aggtttaaga	atagtttttg	ctgtactttc	tatagtgaat	agagttaggc	agggatattc	2580
accattatcg	tttcagaccc	acctcccaac	cccgagggga	cccgcagggc	ccgaaggaat	2640
agaagaagaa	ggtggagaga	gagacagaga	cagatccatt	cgattagtga	acggatccat	2700
ctcgacggaa	tgaaagaccc	cacctgtagg	tttggcaagc	taggatcaag	gtaggaaca	2760
gagagacagc	agaatatggg	ccaaacagga	tatctgtggg	aagcagttcc	tgccccggct	2820
cagggccaaag	aacagttgga	acagcagaat	atgggccaaa	caggatatct	gtggtaagca	2880
gttcctgccc	cggctcaggg	ccaagaacag	atgggtccca	gatgcgggtc	cgccctcagc	2940
agtttctaga	gaaccatcag	atgtttccag	ggtgccccaa	ggacctgaaa	tgacctgtg	3000
ccttatttga	actaaccaat	cagttcgctt	ctcgcttctg	ttcgcgcgct	tctgctcccc	3060
gagctcaata	aaagagccca	caaccctca	ctcggcgcgga	ttcacctgac	gcgtctacgc	3120
caccatggca	ctccccgtca	ccgcccttct	cttgccccctc	gccctgctgc	tgcatgctgc	3180

BLBD_079_02WO_ST25.TXT

cagggccgac attgtgctca ctcaagtcacc tcccagcctg gccatgagcc tgggaaaaag	3240
ggccaccatc tcctgtagag ccagtgaagc cgtcacaatc ttggggagcc atcttattca	3300
ctggtatcag cagaagcccg ggcagcctcc aacccttctt attcagctcg cgtcaaacgt	3360
ccagacgggt gtacctgcca gattttcttg tagcgggtcc cgcactgatt ttacactgac	3420
catagatcca gtggaagaag acgatgtggc cgtgtattat tgtctgcaga gcagaacgat	3480
tcctcgcaca tttggtgggg gtactaagct ggagattaag ggaagcacgt ccggctcagg	3540
gaagccgggc tccggcgagg gaagcacgaa ggggcaaatt cagctgggtcc agagcggacc	3600
tgagctgaaa aaacccggcg agactgttaa gatcagttgt aaagcatctg gctatacctt	3660
caccgactac agcataaatt ggggtgaaacg ggcccctgga aagggcctca aatggatggg	3720
ttggatcaat accgaaacta gggagcctgc ttatgcatat gacttccgcg ggagattcgc	3780
cttttcactc gagacatctg cctctactgc ttacctcaa ataaacaacc tcaagtatga	3840
agatacagcc acttactttt gcgccctcga ctatagttac gccatggact actggggaca	3900
gggaacctcc gttaccgtca gttccgcggc cgcaaccaca acacctgctc caaggccccc	3960
cacacccgct ccaactatag ccagccaacc attgagcctc agacctgaag cttgcaggcc	4020
cgcagcagga ggcgccgtcc atacgcgagg cctggacttc gcgtgtgata tttatatattg	4080
ggcccccttg gccggaacat gtgggggtgtt gcttctctcc cttgtgatca ctctgtattg	4140
taagcgcggg agaaagaagc tcctgtacat cttcaagcag ctttttatgc gacctgtgca	4200
aaccactcag gaagaagatg ggtgttcatg ccgcttcccc gaggaggaag aaggagggtg	4260
tgaactgagg gtgaaatttt ctagaagcgc cgatgctccc gcatatcagc agggtcagaa	4320
tcagctctac aatgaattga atctcggcag gcgagaagag tacgatgttc tggacaagag	4380
acggggcagg gatcccagga tggggggaaa gccccggaga aaaaatcctc aggaggggtt	4440
gtacaatgag ctgcagaagg acaagatggc tgaagcctat agcgagatcg gaatgaaagg	4500
cgaaagacgc agaggcaagg ggcattgacgg tctgtaccag ggtctctcta cagccaccaa	4560
ggacacttat gatgcgttgc atatgcaagc cttgccaccc cgctaattgac aggtaccttt	4620
aagaccaatg acttacaagg cagctgtaga tcttagccac tttttaaaag aaaagggggg	4680
actggaaggg ctaattcact cccaaagaag acaagatctg ctttttgcct gtactgggtc	4740

BLBD_079_02WO_ST25.TXT

tctctggtta gaccagatct gagcctggga gctctctggc taactaggga acccactgct	4800
taagcctcaa taaagcttgc cttgagtgtc tcaatgtgtg tgttggtttt ttgtgtgtcg	4860
aaattctagc gattctagct tggcgtaatc atgggtcatag ctgtttcctg tgtgaaattg	4920
ttatccgctc acaattccac acaacatacg agccggaagc ataaagtgtg aagcctgggg	4980
tgcctaata gtagagctaac tcacattaat tgcgttgctc tctactgccc ctttccagtc	5040
gggaaacctg tcgtgccagc tgcattaatg aatcggccaa cgcgcgggga gaggcggttt	5100
gcgtattggg cgctcttccg cttcctcgct cactgactcg ctgcgctcgg tcgttcggct	5160
gcggcgagcg gtatcagctc actcaaaggc ggtaatacgg ttatccacag aatcagggga	5220
taacgcagga aagaacatgt gagcaaaagg ccagcaaaag gccaggaacc gtaaaaaggc	5280
cgcgttgctg gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg	5340
ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt ttccccctgg	5400
aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc tgtccgcctt	5460
tctcccttcg ggaagcgtgg cgcttttctc tagctcacgc tgtaggtatc tcagttcggt	5520
gtaggtcgtt cgctccaagc tgggctgtgt gcacgaacct ccggttcagc ccgaccgctg	5580
cgcttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact	5640
ggcagcagcc actggtaaca ggattagcag agcgaggatg gtaggcgggtg ctacagagtt	5700
cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttggtg tctgcgctct	5760
gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac	5820
cgctggtagc ggtgggtttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc	5880
tcaagaagat cttttgatct tttctacggg gtctgacgct cagtggaacg aaaactcacg	5940
ttaagggatt ttgggtcatg gattatcaaa aaggatcttc acctagatcc ttttaaatta	6000
aaaatgaagt tttaaataca tctaaagtat atatgagtaa acttgggtctg acagttacca	6060
atgcttaatc agtgaggcac ctatctcagc gatctgtcta tttcgttcat ccatagttgc	6120
ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg gccccagtgc	6180
tgcaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc	6240
agccggaagg gccgagcgca gaagtgggtc tgcaacttta tccgcctcca tccagtctat	6300

BLBD_079_02WO_ST25.TXT

taattgttgc cgggaagcta gagtaagtag ttcgccagtt aatagtttgc gcaacgttgt	6360
tgccattgct acaggcatcg tgggtgtcacg ctcgtcgttt ggtatggcctt cattcagctc	6420
cggttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag	6480
ctccttcggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat cactcatggt	6540
tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac	6600
tggtgagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg	6660
cccggcgtca atacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat	6720
tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc	6780
gatgtaacc actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc	6840
tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg cgacacggaa	6900
atgttgaata ctcatactct tcctttttca atattattga agcatttatc agggttattg	6960
tctcatgagc ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttccgcg	7020
cacatttccc cgaaaagtgc cacctgggac tagctttttg caaaagccta ggcctccaaa	7080
aaagcctcct cactacttct ggaatagctc agaggccgag gcggcctcgg cctctgcata	7140
aataaaaaaa attagtcagc catggggcgg agaatgggcg gaactgggcg gagttagggg	7200
cgggatgggc ggagttaggg gcgggactat ggttgctgac taattgagat gagcttgcat	7260
gccgacattg attattgact agtccctaag aaaccattct tatcatgaca ttaacctata	7320
aaaataggcg tatcacgagg ccctttcgtc	7350