



(86) Date de dépôt PCT/PCT Filing Date: 2010/07/15
(87) Date publication PCT/PCT Publication Date: 2011/01/20
(85) Entrée phase nationale/National Entry: 2012/01/16
(86) N° demande PCT/PCT Application No.: US 2010/042063
(87) N° publication PCT/PCT Publication No.: 2011/008912
(30) Priorité/Priority: 2009/07/16 (US61/225,985)

(51) Cl.Int./Int.Cl. *C09B 57/00* (2006.01),
C12Q 1/66 (2006.01), *G01N 33/535* (2006.01)

(71) Demandeur/Applicant:
BECKMAN COULTER, INC., US

(72) Inventeurs/Inventors:
AKHAVAN-TAFTI, HASHEM, US;
DE SILVA, RENUKA, US;
WANG, GUOPING, US;
EICKHOLT, ROBERT A., US;
GUPTA, RAVINDER K., US;
KAANUMALLE, LAKSHMI S., US

(74) Agent: SIM & MCBURNEY

(54) Titre : NOUVEAUX COLORANTS FLUORESCENTS ET LEURS UTILISATIONS
(54) Title: NOVEL FLUORESCENT DYES AND USES THEREOF

(57) **Abrégé/Abstract:**

The present invention provides fluorescent dyes that are based on firefly luciferin structure. These dyes are optimally excited at shorter wavelengths and have Stokes shift of at least 50 nm. The fluorescent dyes of the invention are useful for preparation of dye-conjugates, which can be used in detection of an analyte in a sample.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 January 2011 (20.01.2011)

(10) International Publication Number
WO 2011/008912 A1

(51) International Patent Classification:
C09B 57/00 (2006.01) *G01N 33/535* (2006.01)
C12Q 1/66 (2006.01)

(74) Agent: PANDE, Hema; Beckman Coulter, Inc., 250 S. Kraemer Blvd., Brea, California 92821 (US).

(21) International Application Number:
PCT/US2010/042063

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
15 July 2010 (15.07.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/225,985 16 July 2009 (16.07.2009) US

(71) Applicant (*for all designated States except US*): BECKMAN COULTER, INC. [US/US]; 250 S. Kraemer Boulevard, Brea, California 92821 (US).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): AKHAVANTAFTI, Hashem [US/US]; 4545 Vines Road, Howell, Michigan 48843 (US). DE SILVA, Renuka [US/US]; 40628 Delta Drive, Northville, Michigan 48167 (US). WANG, Guoping [CN/US]; 24723 Naples Drive, Novi, Michigan 48375 (US). EICKHOLT, Robert, A. [US/US]; 1171 Vermont, Troy, Michigan 48083 (US). GUPTA, Ravinder, K. [US/US]; 13770 NW 18th Court, Pembroke Pines, Florida 33028 (US). KAANUMALLE, Lakshmi, S. [US/US]; 603 Connor Ct., Schenectady, New York 12309 (US).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: NOVEL FLUORESCENT DYES AND USES THEREOF

(57) Abstract: The present invention provides fluorescent dyes that are based on firefly luciferin structure. These dyes are optically excited at shorter wavelengths and have Stokes shift of at least 50 nm. The fluorescent dyes of the invention are useful for preparation of dye-conjugates, which can be used in detection of an analyte in a sample.



WO 2011/008912 A1

WO 2011/008912

PCT/US2010/042063

NOVEL FLUORESCENT DYES AND USES THEREOF

FIELD OF THE INVENTION

5

The present invention relates to fluorescent dyes, more specifically, ultraviolet excitable fluorescent dye compositions, compounds and conjugates, as well as method of using and making the same.

10 BACKGROUND OF THE INVENTION

Fluorescent dyes are widely used for labeling, detecting, and/or quantifying components in a sample. The various approaches used for such detection and/or quantification include fluorescence microscopy, fluorescence immunoassay, flow cytometric analysis of cells, and various other applications. In general, for many applications that utilize fluorescent dyes as
15 detection tools, it is necessary to conjugate the fluorescent dye with a ligand such as a protein, antibody, enzyme, nucleotide, nucleic acid, and other biological and non-biological molecules to make a dye-labeled ligand. The dye-labeled ligand is an important reagent that confers specificity for a subsequent biochemical interaction in which the fluorescent dye provides a method for detection and/or quantification of the interaction.

20

The choice in fluorescent dyes is particularly important in applications that utilize multiplex, multicolor analysis, such as fluorescence microscopy, fluorescence immunoassay, flow cytometry, and various other applications. Notably, certain detection applications require an ultraviolet excitable fluorophore having a specific excitation range.

25

Specifically, there is a need for a fluorescent dye that can be efficiently excited by the 405 nm violet laser in multi-color flow cytometry instruments. The target dye for such applications should essentially have the following features: 1) a maximum in its excitation spectrum near 405 nm, 2) a strong spectrally resolvable emission maximum, 3) a large
30 Stokes' shift, preferably, at least 50 nm, and 4) the ability of the fluorescent dye to couple to a biomolecule through a reactive group. Heretofore, there is a shortage of ultraviolet excitable fluorophores that can be conjugated to different ligands to provide dye-labeled reagents having optically and electronically separable fluorescence spectral properties.

WO 2011/008912

PCT/US2010/042063

The fluorescent dyes of the invention are structurally similar to firefly luciferin compounds. Such compounds have been previously used as chemiluminescent reagents whereby light is generated by the oxidative catalysis of luciferin. Luciferin, a substrate for the enzyme
 5 luciferase, is oxidized in the presence of luciferase to produce oxyluciferin and energy which is released in the form of light. For these types of assays, the luciferase-luciferin reaction provides the basis for simple, rapid, and sensitive assays for a wide range of substances (Karicka, L.J., Analytical Biochemistry, 175, 14-21 (1988)).


10 While such chemiluminescent dyes based on the structure of firefly luciferin have gained widespread use, such use does not evidence the capability of compounds having a structure similar to that of firefly luciferin to function as fluorescent dyes excitable within a specific part of the ultraviolet spectrum. Such ultraviolet excitable fluorescent dyes are particularly advantageous for a wide range of applications, including without limitation, applications
 15 that utilize multiplex, multicolor fluorescence analysis.

SUMMARY OF THE INVENTION

The present invention relates to novel ultraviolet excitable fluorescent dyes based generally
 20 on the firefly Luciferin structure, compositions comprising the same, intermediates thereof and methods of their use.

In one aspect, compositions of fluorescent dyes having the following general formula (I) are provided:

25



wherein

X_1 and X_2 are independently S or O;

Y_1 and Y_2 are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl, or one of

30 Y_1 or Y_2 is H, and the other is a haloalkyl group;

WO 2011/008912

PCT/US2010/042063

wherein each W is independently H or alkyl;


Z₁ is H or alkyl;

wherein Z₁ and one of the W groups may be absent and, if absent, are replaced with an additional bond between C₄ and C₅ in ring C, which is indicated by a dotted line;

5 Z₂ comprises a labeling substituent of the formula – L-RG, wherein L is a bond or a linking group; and

RG is a reactive group which enables the fluorescent dye to be bound to another molecule.

10 In another aspect, synthetic intermediates for synthesizing the fluorescent dyes of the invention are provided. In one embodiment, the synthetic intermediates are represented by the general formula.



15

wherein

Y₁ and Y₂ are independently, halogen, alkyl, haloalkyl, alkoxy, or alkenyl,

or one of Y₁ or Y₂ is H, and the other is a haloalkyl group.

20 In another aspect, dye-conjugates comprising fluorescent dyes conjugated to biomolecules are provided.

In another aspect, methods of using fluorescent dyes and dye-conjugates to locate or detect the presence or interaction of analytes or ligands in a sample are provided. In specific
25 embodiments, methods of using the dye-conjugates for detecting the presence of complementary biomolecules that bind to the dye-conjugates are provided.

In other embodiments reagents and kits comprising the fluorescent dyes, dye intermediates, or dye-conjugates are provided for use in detecting the presence of complementary
30 biomolecules that bind to the dye-conjugates of the invention.

WO 2011/008912

PCT/US2010/042063

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a general structure of ultraviolet excitable fluorescent dyes of the invention.

5 Figure 2A-2D depicts a synthesis scheme for luciferin derivatives.

Figure 3A and 3B depict synthesis schemes for derivatives of dehydroluciferin acids.

Figure 4 depicts histograms illustrating the relative performance of CD4-Dye2 conjugate to
10 CD4-Pacific Orange conjugate using two-color flow cytometry.

Figure 5 depicts histograms illustrating the relative performance of CD4-Dye3 to CD4-Pacific Orange conjugate using ten-color flow cytometry.

WO 2011/008912

PCT/US2010/042063

DETAILED DESCRIPTION OF INVENTION

DEFINITIONS

5

All scientific and technical terms used in this application have meanings commonly used in the art unless otherwise specified. As used in this application, the following words or phrases have the meanings specified.

- 10 As used herein, the term “activated ester” means an ester which spontaneously reacts with an amino group. Activated esters generally have the formula –COR, where R is a good leaving group. Activated esters include, but are not limited to, succinimidyloxy (–OC₆H₄O₂) also known as N-hydroxy-succinimide ester, sulfosuccinimidyloxy (–OC₆H₃O₂–SO₃H), -1-oxybenzotriazolyl (–OC₆H₄N₃).

15

As used herein, the term “alkyl” means a branched, straight chain or cyclic hydrocarbon group containing from 1-20 carbons. The term “alkyl” includes a “lower alkyl” group which typically contains up to 8 carbons.

- 20 As used herein, the term “alkenyl” means a branched, straight chain or cyclic hydrocarbon group containing at least one C=C double bond and containing from 2-20 carbons. The term “alkenyl” includes a “lower alkenyl” group which typically contains up to 8 carbons.

- As used herein, the term “alkynyl” means a branched or straight chain hydrocarbon group
25 containing at least one C≡C triple bond and containing from 2-20 carbons. The term “alkynyl” includes a “lower alkynyl” group which typically contains up to 8 carbons.

- As used herein, the term “aryl” means an aromatic ring-containing group containing 1-5
30 carbocyclic aromatic rings, which can be substituted with 1 or more substituents other than H.

As used herein, the term “analyte” means a substance the presence or amount of which is to be measured in a sample by an assay. The analyte may include a reactive group, e.g., a group through which a dye of the invention can be conjugated to the analyte. Analytes

WO 2011/008912

PCT/US2010/042063

include organic and biological molecules to which a specific binding partner having a specific binding affinity exists. Exemplary analytes include, without limitation, single stranded or double stranded DNA, RNA, DNA-RNA complexes, oligonucleotides, antibodies, antibody fragments, antibody-DNA chimeras, antigens, haptens, proteins, lectins, 5 avidin, streptavidin and biotin. Other exemplary analytes also include drugs and hormones.

As used herein, the term “conjugated molecule” or “CM” means a biological or non-biological component that is associated with at least one dye of the present invention. Such components include without limitations antigens, antibodies, carbohydrates (e.g., mono-, 10 oligo-, and poly-saccharides), proteins, peptides, haptens, nucleosides, nucleotides, oligonucleotides, nucleic acids, polymers, virus, microorganism, or cell or cellular components. “Conjugated molecule” also includes solid supports (e.g., synthesis support, chromatographic support, membrane, or beads).

15 As used herein, the term “fluorescent dye” means a compound that absorbs light in the ultraviolet and violet regions of the electromagnetic spectrum, and reemits light in the blue region to produce a detectable signal. “Fluorescent Dye” also includes fluorescent compounds having a chemically reactive group that facilitates attachment of dye to a conjugate molecule.

20

As used herein the term “dye-conjugate” means a molecule in which a fluorescent dye is associated with a biological or non-biological component. Such biological or non-biological component is also referred to as “conjugated molecule”. In one embodiment, the association of a fluorescent dye to a conjugated molecule is via a covalent linkage. The examples of the 25 dye-conjugates include, but are not limited to, conjugates of antigens, antibodies, carbohydrates, proteins, peptides, haptens, nucleosides, nucleotides, oligonucleotides, nucleic acids, polymers, solid supports (e.g., synthesis supports, chromatographic supports, membranes, or beads), virus, microorganism, or cell or cellular components.

30 As used herein the term “fluorophore” means a molecule which is capable of absorbing energy at a wavelength range and releasing energy at a wavelength range other than the absorbance range. The term “excitation wavelength” means the range of wavelengths at which a fluorophore absorbs energy. The term “emission wavelength” means the range of wavelengths that the fluorophore releases energy or fluoresces.


As used herein the term “halogen” means fluorine, chlorine, bromine or iodine atoms. Similarly, the term “halo” means fluoro, chloro, bromo, or iodo group.

5 As used herein the term “haloalkyl” means an alkyl group in which one or more hydrogen atoms is replaced by one or more halogen atoms. In one embodiment, the haloalkyl can be substituted with 1, 2, or 3 halo groups. The term haloalkyl also includes perfluoro-alkyl groups. Examples of haloalkyls include trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, and the like.

10


As used herein the term “linking group” or “L” means a group for linking the dye to the reactive group or RG. The linking group can be a bond, an atom, or another divalent or polyvalent group. The linking group can comprise a straight or branched chain of atoms some of which can be part of a ring structure. The linking group usually contains from 1 to
 15 about 50 non-hydrogen atoms, more usually from 1 to about 30 non-hydrogen atoms. Atoms comprising the chain are selected from C, O, N, S, P, Si, B, and Se atoms, preferably from C, O, N, P and S atoms that covalently link the fluorescent dye to another moiety such as a chemically reactive group. Halogen atoms can be present as substituents on the chain or ring. Typical functional groups comprising the linking substituent include alkene, alkylene,
 20 arylene, alkenylene, ether, peroxide, carbonyl as a ketone, ester, carbonate ester, thioester, or amide group, amine, amidine, carbamate, urea, imine, imide, imidate, carbodiimide, hydrazine, diazo, phosphodiester, phosphotriester, phosphonate ester, thioether, disulfide, sulfoxide, sulfone, sulfonate ester, sulfate ester, and thiourca groups. Examples of linking groups include substituted or unsubstituted polymethylene, arylene, alkylarylene,
 25 arylenealkyl, or arylthio.

One linking group according to the present invention for linking dye to a reactive group has the general structure $R_1C(O)R_2$, where R_1 is a bond or C_{1-10} methylene $(CH_2)_n$ attached to the dye, $C(O)$ is a carbonyl group, and R_2 is OH or a bond attaching the carbonyl group to a
 30 reactive group RG. Examples of this linking group include:




WO 2011/008912

PCT/US2010/042063



- 5 In another embodiment, illustrated below, the linking group has the general structure $\text{R}_1\text{C}(\text{O})\text{A}_1\text{R}_3\text{C}(\text{O})\text{R}_2$, where R_1 is a bond or C_{1-10} methylene as described above, A_1 is either NH, S or O, R_3 is an alkenyl $(\text{CH}_2)_n$, a five or a six membered ring having at least one unsaturated bond, or a combination of C_{1-10} methylene $(\text{CH}_2)_n$ and a five or six member ring, R_2 is OH or a bond attaching the terminal carbonyl group to a reactive group RG. Examples
- 10 of this linking group include:



15

As used herein, the term “reactive group” or “RG” is an atom or group whose presence facilitates bonding to another molecule by covalent attachment or physical forces. Selected

examples of reactive groups are shown in Table 1. In some embodiments, attachment of a fluorescent compound of the present invention to another compound will involve loss of one or more atoms from the reactive group for example when the reactive group is a leaving group such as a halogen atom or a tosylate group and the fluorescent labeling compound is covalently attached to another compound by a nucleophilic displacement reaction. In other embodiments, attachment of a fluorescent labeling compound to another compound by covalent bond formation will involve reorganization of bonds within the reactive group as occurs in an addition reaction such as a Michael addition or a cycloaddition reaction such as the Diels-Alder reaction or when the reactive group is an isocyanate or isothiocyanate group.

In still other embodiments, attachment will not involve covalent bond formation, but rather physical forces in which case the reactive group remains unaltered. By physical forces is meant attractive forces such as hydrogen bonding, electrostatic or ionic attraction, hydrophobic attraction such as base stacking, and specific affinity interactions such as biotin-streptavidin, antigen-antibody and nucleotide-nucleotide interactions.

TABLE 1

Reactive Groups for Chemical Binding of Dyes to Conjugated Molecules

REACTIVE GROUP (RG) (attached to reactive dye)	FUNCTIONAL GROUP (attached to conjugated molecule)
carboxylic acid	amine
carboxyl ester	amine
activated ester	amine
acid anhydride	amine
acid halide	amine
acyl azide	amine
aldehyde	amine
chloroformate	amine
isocyanate	amine
isothiocyanate	amine
sulfonyl halide	Amine/hydroxyl
alkyl halide	thiol
Alkyl sulfonate	thiol

aziridine	thiol
haloacetamide	thiol
maleimide	thiols
amine	Carboxylic acid
carbodiimide	Carboxylic acid
diazoalkane	Carboxylic acid
acyl halide	hydroxyl
halotriazine	hydroxyl
hydrazine/hydrazide	hydroxyl
phosphoramidites	alcohols

When the reactive group is a moiety, such as a carboxylic acid or an activated ester of a carboxylic acid, the reactive dye is particularly useful for preparing dye conjugates with molecules containing one or more amino groups, such as proteins, peptides, nucleotides, oligonucleotides, or haptens. Where the reactive group is a maleimide or haloacetamide, the reactive dye is particularly useful for conjugation with thiol-containing molecules. Where the reactive group is a hydrazide, the reactive dye is particularly useful for conjugation to periodate-oxidized carbohydrates and glycoproteins.

10 In one embodiment, reactive groups include carboxylic acid, succinimidyl ester of a carboxylic acid, isothiocyanate, haloacetamide, hydrazine, aliphatic amine, and maleimide groups. Methods to prepare each of these reactive groups are well known in the art and their application for a particular purpose is within the ability of one having skill in the art.

15 Bifunctional coupling reagents can also be used to couple labels to organic and biological molecules with moderately reactive groups (see L. J. Kricka, Ligand-Binder Assays, Marcel Dekker, Inc., New York, 1985, pp. 18-20, Table 2.2 and T. H. Ji, "Bifunctional Reagents," Methods in Enzymology, 91, 580-609 (1983)). There are two
20 types of bifunctional reagents, those which become incorporated into the final structure and those which do not and serve only to couple the two reactants.

As used herein, the term “sample” means a fluid containing or suspected of containing one or more analytes to be assayed. Typical samples which are analyzed utilizing fluorescent dyes are biological samples including body fluids such as blood, plasma, serum, urine, semen, saliva, cell lysates, tissue extracts and the like. The sample can
 5 also include diluents, buffers, detergents, contaminants, and the other such components that are usually present in a body fluid.

As used herein the term “specific binding pair” means two substances which exhibit a mutual binding affinity. Examples include antigen-antibody, hapten-antibody,
 10 antibody-antibody pairs, complementary oligonucleotides or polynucleotides, avidin-biotin, streptavidin-biotin, enzyme-substrate, hormone-hormone receptor, lectin-carbohydrate, IgG-protein A, IgG-protein G, nucleic acid-nucleic acid binding protein, and nucleic acid-anti-nucleic acid antibody (Table 2).

15

TABLE 2

REPRESENTATIVE SPECIFIC BINDING PAIRS	
antigen	antibody
hapten	antibody
biotin	avidin or strepavidin
carbohydrate	lectin or carbohydrate receptor
DNA	complementary strand DNA
RNA	complementary strand RNA
drug	drug receptor
IgG	protein A/protein G
substrate	enzyme
Protein/peptide	protein/peptide receptor
hormone	hormone receptor
Nucleic acids	Anti-nucleic acid antibody

20 As used herein, the term “substituted” means replacement of at least one hydrogen atom on a group by another atom or a group having 1 to 50 atoms selected from C, O,

WO 2011/008912

PCT/US2010/042063

N, S, P, Si, F, Cl, Br, I. In references to substituted groups, it is intended that multiple points of substitution can be present.


COMPOSITIONS OF INVENTION

5

The invention provides a class of fluorescent dyes that are based on the structure of firefly luciferin, and possess certain functional advantages. For example, the fluorescent dyes of the invention are adapted for excitation at shorter wavelengths (between 340 and 450 nm), have emission maxima usually between 500 and 550 nm
10 wavelengths, narrow emission bandwidth, a large Stokes shift of at least about 50 nm, and other favorable fluorescent properties.

The fluorescent dyes of the present invention are particularly useful with violet 405 nm lasers. Compared to existing violet laser excitable dyes (e.g., Pacific Orange), the
15 fluorescent dyes of the invention, despite having comparable absorptivity, show surprisingly and unexpectedly brighter fluorescence. The fluorescent dyes of the invention are well adapted for coupling to a conjugated molecule through a reactive group. In addition, the dye-conjugates of the invention, in particular, the protein conjugates, exhibit bright fluorescence even at a relatively high degree of dye
20 substitution. These properties make the dyes of the invention particularly well-suited for multicolor, multiplexing application.

The fluorescent dyes of the general formula are provided:



25

(I)

wherein

X₁ and X₂ are independently S or O;

Y₁ and Y₂ are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl,

or

30

one of Y₁ or Y₂ is H, and the other is a haloalkyl group;

WO 2011/008912

PCT/US2010/042063


wherein each W is independently H or alkyl;

Z₁ is H or alkyl;

wherein Z₁ and one of the W groups may be absent and, if absent, are replaced with an additional bond between C₄ and C₅ in ring C, which is indicated by a dotted line;

Z₂ comprises a labeling substituent of the formula – L-RG, wherein L is a bond or a linking group which connects the dye to a reactive group RG, wherein the linking group can comprise a covalent linkage comprising 1-50 non-hydrogen atoms selected from C, N, O, S, P and halogen atoms and is composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and wherein RG includes a carboxylic acid, an activated ester of a carboxylic acid, acid anhydride, acid chloride, acyl azide, an acyl halide, aldehyde, chloroformate, amine, hydroxy, hydrazine, isocyanate, isothiocyanate, sulfonyl halide, tosyl, maleimide, N-hydroxy-succinimide ester, aziridine, imine, and disulfide groups.

In one embodiment of the fluorescent dye of formula (I), X₁ and X₂ are S having the formula:



(I-A)

wherein

Y₁ and Y₂ are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl,
or

one of Y₁ or Y₂ is H, and the other is a haloalkyl group;

Z₁ is H or alkyl;

wherein each W is independently H or alkyl;

in L-RG, L is a bond or a linking group which connects the dye to a reactive group RG, wherein the linking group can comprise a covalent linkage comprising 1-50 non-hydrogen atoms selected from C, N, O, S, P and halogen

WO 2011/008912

PCT/US2010/042063

atoms and is composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and
wherein RG can comprise a carboxylic acid, an activated ester of a carboxylic
5 acid, acid anhydride, acid chloride, acyl azide, an acyl halide, aldehyde, chloroformate, amine, hydroxy, hydrazine, isocyanate, isothiocyanate, sulfonyl halide, tosyl, maleimide, N-hydroxy-succinimide ester, aziridine, imine, and disulfide groups.

10 Selected embodiments of the dyes of formula (I-A) are illustrated in Table 3


TABLE 3

Dye	X1	X2	Y1	Y2	ZI	W	L	RG
I-A-1	S	S	H	H	H	Mc	C(O)	OH
I-A-2	S	S	H	H	H	Me	C(O)	NHS
I-A-3	S	S	H	H	H	Me	C(O)	NHBSA
I-A-4	S	S	Cl	H	H	Me	C(O)	OH
I-A-5	S	S	H	Cl	H	Me	C(O)	OH
I-A-6	S	S	Cl	Cl	H	Me	C(O)	OH
I-A-7 (Dye1)	S	S	Cl	Cl	H	Me	C(O)	NHS
I-A-8	S	S	Cl	Cl	H	Me	C(O)	NHBSA
I-A-9 (Dye2)	S	S	Cl	Cl	H	Me	C(O)NH(CH ₂) ₅ C(O)	NHS
I-A-10	S	S	Cl	Cl	H	Me	C(O)NH(CH ₂) ₅ C(O)	NHBSA
I-A-11	S	S	Cl	Cl	H	Me	C(O)NH(CH ₂) ₂ NHCS NHPh	NCS
I-A-12	S	S	Cl	Cl	H	Me	C(O)NH(CH ₂) ₂ NHCS NHPh	NHCSNHBSA
I-A-13	S	S	F	F	H	Me	C(O)	OH
I-A-14 (Dye3)	S	S	F	F	H	Me	C(O)	NHS
I-A-15	S	S	F	F	H	Me	C(O)	NHBSA
I-A-16	S	S	Me	Me	H	Me	C(O)	OH
I-A-17	S	S	Me	H	H	Me	C(O)	OH
I-A-18	S	S	CF ₃	H	H	Me	C(O)	OH
I-A-19 (Dye7)	S	S	CF ₃	H	H	Me	C(O)	NHS
I-A-20	O	S	H	H	H	Mc	C(O)	OH
I-A-21	S	S	F	F	H	Me	C(O)NH(CH ₂) ₂ C(O)	OH
I-A-22 (Dye5)	S	S	F	F	H	Me	C(O)NH(CH ₂) ₂ C(O)	NHS
I-A-23	S	S	F	F	H	Me	C(O)NHCH ₂ C(O)	OH
I-A-24	S	S	Cl	Cl	H	Me	C(O)NH(CH ₂) ₂ C(O)	OH
I-A-25	S	O	F	F	H	H	C(O)	OH
I-A-26	S	S	Cl	H	H	H	C(O)	OH
I-A-27	S	S	H	Cl	H	H	C(O)	OH
I-A-28	S	S	Cl	Cl	H	H	C(O)	OH
I-A-27	S	S	F	F	Me	H	C(O)	OH
I-A-29 (Dye4)	S	S	F	F	Me	H	C(O)	NHS
I-A-30	S	S	F	F	H	H	C(O)	OH
I-A-31	S	S	F	F	H	H	C(O)	NHS
I-A-32	S	S	H	H	H	H	C(O)	OH
I-A-33	S	S	F	F	H	H	(CH ₂) ₂ C(O)	OH
I-A-34 (Dye6)	S	S	F	F	H	H	(CH ₂) ₂ C(O)	NHS

WO 2011/008912

PCT/US2010/042063

In other embodiments, the fluorescent dyes have the following formula:



5

(I-B)

wherein

X₁ and X₂ are independently S or O;

Y₁ and Y₂ are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl;

or

10

one of Y₁ or Y₂ is H, and the other is a haloalkyl group;

W is H or alkyl;

L is a bond or a linking group which connects the dye to a reactive group RG,

wherein the linking group can comprise a covalent linkage comprising 1-50

non-hydrogen atoms selected from C, N, O, S, P and halogen atoms and is

15

composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and


wherein RG can comprise a carboxylic acid, an activated ester of a carboxylic acid, acid anhydride, acid chloride, acyl azide, an acyl halide, aldehyde,

20

chloroformate, amine, hydroxy, hydrazine, isocyanate, isothiocyanate, sulfonyl halide, tosyl, malcimide, N-hydroxy-succinimide ester, aziridine, imine, and disulfide groups.

In one embodiment of the fluorescent dye of formula (I-B), X₁ and X₂ are S, as provided in formula:

25



WO 2011/008912

PCT/US2010/042063

wherein,

Y_1 and Y_2 are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl

or

one of Y_1 or Y_2 is H, and the other is a haloalkyl;

5 W is H or alkyl;

L is a bond or a linking group, wherein the linking group comprises a covalent linkage comprising 1-50 non-hydrogen atoms selected from C, N, O, S, P and halogen atoms and is composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and

10 RG is a reactive group, wherein RG can comprise a carboxylic acid, an activated ester of a carboxylic acid, acid anhydride, acid chloride, acyl azide, an acyl halide, aldehyde, chloroformate, amine, hydroxy, hydrazine, isocyanate, isothiocyanate, sulfonyl halide, tosyl, maleimide, N-hydroxy-succinimide ester, aziridine, imine, and disulfide groups.

Selected embodiments of dyes of formula (I-B) are illustrated in Table 4.

Table 4

20


Dye	X1	X2	Y1	Y2	W	L	RG
I-B-1	S	S	Cl	Cl	H	C(O)	OH
I-B-2	S	S	Cl	Cl	H	C(O)	NHS
I-B-3	S	S	Cl	Cl	Me	C(O)	OH
I-B-4	S	S	Cl	Cl	Me	C(O)	NHS
I-B-5	S	S	F	F	Me	C(O)	OH
I-B-6	S	S	F	F	Me	C(O)	NHS
I-B-7	S	S	F	F	H	C(O)	OH
I-B-8	S	S	F	F	H	C(O)	NHS

The invention further provides synthetic intermediate compounds that may be used for synthesizing the fluorescent dyes of the invention. In one embodiment, the intermediate compounds are represented by the general formula (II):

25

WO 2011/008912

PCT/US2010/042063



(II)

wherein


Y_1 and Y_2 are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl

5

or

one of Y_1 or Y_2 is H, and the other is a haloalkyl group.

In another embodiment, the intermediate compounds are represented by the general formula (III):



10

(III)

wherein

X_2 is S or O;

wherein each W is independently H or alkyl;

15

Z_1 is H or alkyl; and

LH is OH or $\text{NH}(\text{CH}_2)_n\text{COOH}$ (n is 1-10).


The fluorescent dyes of the invention may be covalently or non-covalently associated with one or more conjugated molecules "CM". Covalent association may occur through various mechanisms, including attachment of a conjugated molecule CM to the fluorescent dye via a linking group L as described above, and may involve a covalent linkage.

Where the dye is associated non-covalently with one or more molecules, the association may occur through various mechanisms, including incorporation of the dye or synthetic intermediate compound into or onto a solid or semisolid matrix, such as a bead or a surface, or by nonspecific interactions, such as hydrogen bonding, ionic bonding, or hydrophobic interactions (such as Van der Waals forces). The associated molecules

WO 2011/008912

PCT/US2010/042063

- may be selected from the group consisting of polypeptides, polynucleotides, polysaccharides, beads, microplate well surfaces, metal surfaces, semiconductor and non-conducting surfaces, nano-particles, and other solid surfaces. The associated or conjugated molecule may be associated with or conjugated to more than one
- 5 fluorescent dye, which may be the same or different. Generally, methods for the preparation of dye-conjugates of biological substances are well-known in the art. See, for example, (Greg T. Hermanson, Bioconjugate Techniques, 1996 Edition Academic Press, Chapter 8, pages 298-362; Nataliya Panchuk, et al. J. Histochemistry & Cytochemistry, 1179-1188 (1999); S Bhattacharya, et al Bioconjugate Chemistry, 10 Volume 19, 1186-1193 (2008); B. Dworecki, et al <http://www.piercenet.com/files/DyLight%20Poster%206-7-04.pdf>). Typically, the association or conjugation of a fluorophore to a dye-conjugate imparts the spectral properties of the fluorophore to that molecule.
- 15 In one embodiment, a fluorescent dye of the invention is attached to a conjugated molecule CM via a covalent linkage, the dye-conjugate having the formula:



- 20 wherein
- X_1 and X_2 are independently S or O;
- Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl wherein each W is independently H or alkyl;
- Z_1 is H or alkyl;
- 25 wherein Z_1 and one of the W groups may be absent and, if absent, are replaced with an additional bond between C_4 and C_5 in ring C, which is indicated by a dotted line;
- L is independently a bond or a linking group, wherein the linking group comprises a covalent linkage comprising 1-50 non-hydrogen atoms selected
- 30 from C, N, O, S, P and halogen atoms and is composed of any combination

single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and wherein CM is a conjugated molecule.

- 5 Conjugated molecules CM that may be used for preparing dye-conjugates according to the present invention include, but are not limited to, amino acids, peptides, proteins, nucleosides, nucleotides, nucleic acids, carbohydrates, enzymes, lipids, non-biological polymers, cells, and cellular components. The molecules to be conjugated may be protected on one or more functional groups in order to facilitate the conjugation, or to
10 insure subsequent reactivity.

Where the molecule is a peptide, the peptide may be a dipeptide or larger, and typically includes 5 to 50 amino acids. Where the conjugated molecule is a protein, it may be an enzyme, an antibody, catalytic antibody, kinase, lectin, glycoprotein, histone, albumin,
15 lipoprotein, avidin, streptavidin, protein A, protein G, hormone, toxin, growth factor. Typically, the conjugated protein is an antibody, an antibody fragment, avidin, streptavidin, a lectin, or a growth factor.

The conjugated molecule may be a nucleic acid polymer, such as for example DNA
20 oligonucleotides, RNA oligonucleotides (or hybrids thereof), or single-stranded, double-stranded, triple-stranded, or quadruple-stranded DNA, or single-stranded or double-stranded RNA. The conjugated molecule may also include carbohydrates that are polysaccharides, such as dextrans.

- 25 The associated or conjugated molecule may be a member of a specific binding pair, and therefore useful as a detection reagent for the complementary member of that specific binding pair. The dye-conjugate of a specific binding pair member may be useful for detecting and optionally quantifying the presence of the complementary specific binding pair member in a sample, by methods that are well known in the art.

30

Representative specific binding pairs may include without limitation ligands and receptors, and may include but are not limited to the following pairs: antigen-antibody, biotin-avidin, biotin-streptavidin, IgG-protein A, IgG-protein G, carbohydrate-lectin,

WO 2011/008912


PCT/US2010/042063

enzyme-enzyme substrate; DNA-complementary strand DNA, and RNA-complementary strand RNA, hormone-hormone receptor (Table 2).

METHODS OF PREPARING FLUORESCENT DYE COMPOUNDS

5

In one aspect of the invention a method is provided for preparing the fluorescent dyes of formula (I).



wherein

10

X_1 and X_2 are independently S or O;

Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl wherein each W is independently H or alkyl;

Z_1 is H or alkyl;

15

wherein Z_1 and one of the W groups may be absent and, if absent, are replaced with an additional bond between C_4 and C_5 in ring C, which is indicated by a dotted line;

wherein Z_2 comprises a labeling substituent of the formula $-L-RG$, wherein

L is independently a bond or a linking group, wherein the linking group

comprises a covalent linkage comprising 1-50 non-hydrogen atoms selected

20

from C, N, O, S, P and halogen atoms and is composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and

wherein RG is a reactive group selected from a group consisting of a carboxylic acid, an activated ester of a carboxylic acid, acid anhydride, acid chloride, acyl


25

azide, an acyl halide, aldehyde, chloroformate, amine, hydroxyl, hydrazine, isocyanate, isothiocyanate, sulfonyl halide, tosyl, maleimide, N-hydroxy-succinimide ester, aziridine, imine, and disulfide groups.

In one embodiment, the fluorescent dyes of the invention having the formula:

WO 2011/008912

PCT/US2010/042063



wherein Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl;

X_2 is S or O;

5 wherein each W is independently H or alkyl;


Z_1 is H or alkyl; and

L is O or $\text{NH}(\text{CH}_2)_n\text{COO}$ (n is 1-10)

can be synthesized using the synthesis scheme illustrated in Figure 2A-2D. The method includes the steps of:

10


(a) forming cyanobenzothiazole intermediate of the dye using the method shown in Figure 2B having the formula;



wherein

15 Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl.

(b) forming a carboxyl derivative of the dye by reacting cyanobenzothiazole intermediate with modified amino acid of formula:



20

wherein

X_2 is S or O;

wherein each W is independently H or alkyl;


Z_1 is H or alkyl; and

LH is OH or $\text{NH}(\text{CH}_2)_n\text{COOH}$ (n is 1-10).

WO 2011/008912

PCT/US2010/042063

to produce luciferin derivative of formula:



5

wherein

X_2 is S or O;

Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl;

wherein each W is independently H or alkyl;

Z_1 is H or alkyl; and


10

LH is OH or $\text{NH}(\text{CH}_2)_n\text{COOH}$ (n is 1-10).

(c) replacing the hydroxyl function of the carboxyl group in LH with a reactive group.

In another embodiment, the fluorescent dyes of the invention having the formula:

15



wherein

Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl;

and

20


R is H or alkyl

are synthesized using the synthesis scheme illustrated in Figures 3A.

In another embodiment, the fluorescent dyes of the invention having the formula:

WO 2011/008912

PCT/US2010/042063



wherein

Y₁ and Y₂ separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl;

5 and

R is H or alkyl

can be synthesized using the synthesis scheme illustrated in Figure 3B (Bagley, MC. et al., *J. Am. Chem. Soc.*, 122, 3301-3313 (2000)).

- 10 In another aspect the invention provides methods for making dye-conjugates of invention. The method comprises mixing a fluorescent dye of the invention comprising a reactive group with a conjugated molecule. The conjugated molecule may be an amino acid, protein, peptide, antibody, antibody fragment, nucleoside, nucleotide, nucleic acid polymer. The conjugated molecule may include carbohydrates that are
- 15 polysaccharides, such as dextrans. Typically, the conjugation of a fluorophore to a conjugated molecule imparts the spectral properties of the fluorophore to that molecule.

The methods for the preparation of dye-conjugates of are well-known in the art. For preparation of peptide or protein conjugates, the method typically comprises first

20 dissolving the protein to be conjugated in aqueous buffer at about 1-10 mg/ml at room temperature or below (typically 4–25° C). Borate or carbonate/bicarbonate buffers (pH about 8.0-9) are especially suitable for reaction with succinimidyl esters, phosphate buffers (pH about 4.0-8) for reaction with thiol-reactive functional groups, and carbonate or borate buffers (pH about 9.0-9.8) for reaction with isothiocyanates and

25 dichlorotriazines. The appropriate reactive dye is dissolved in a nonaqueous solvent (usually DMSO or DMF) in an amount sufficient to give a suitable degree of conjugation and added to a solution of the protein to be conjugated. The appropriate amount of reactive dye to be used for any protein or other component is usually predetermined by experimentation in which variable amounts of the reactive dye are

30 added to the protein to be conjugated. The reactive dye is usually used in 5-100 fold excess of the amount of protein to be conjugated. Following addition of the reactive compound to the component solution, the mixture is incubated for a suitable period

(typically about 1 hour at room temperature to several hours on ice), the dye-conjugate is separated from unconjugated dye by gel filtration, dialysis, HPLC or other suitable methods. The dye-conjugate can be stored in solution or lyophilized, until tested in its desired application. This method is generally applicable for preparing dye-conjugates using antibodies, antibody fragments, avidins, lectins, enzymes, proteins A and G, cellular proteins, albumins, histones, growth factors, hormones, and other proteins.

METHODS OF USE

10 The present invention provides a method of detecting a complementary member of a specific binding pair in a sample, comprising mixing said sample with a dye-conjugate of the invention which specifically binds to the complementary member; and detecting the complex formed by said mixture to detect the complementary member. The methods of the invention may comprise using the fluorescent dyes, including reactive
15 dyes and dye-conjugates of the invention for detection and/or quantitation of an analyte in a sample. In certain embodiments, the sample comprises heterogeneous mixtures of components, including intact cells, cell extracts, bacteria, viruses, organelles, and mixtures thereof. In other embodiments, the sample comprises a single component or homogeneous group of components, e.g. biological polymers such as amino acid
20 polymers, nucleic acid polymers, carbohydrate polymers, or lipid membrane complexes.

In certain embodiments, the fluorescent dyes of the invention, including reactive dyes and dye-conjugates are used to stain or label an analyte in a sample so that the analyte can be identified or quantitated. For example, fluorescent dyes, including dye-
25 conjugates of the invention may be used as a detectable tracer element in an assay for a target analyte, in a biological or non-biological fluid.

In other embodiments, the fluorescent dyes, including reactive dyes and dye-conjugates are used to detect a sample that comprises a ligand for which the conjugated molecule
30 is a complementary member of a specific binding pair (e.g., Table 2). Thus, the ligand is one member and the conjugated molecule the other member of the specific binding pair. The fluorescent dyes, including the dye-conjugates are also used for identifying an interaction between a dye-conjugate and a complementary binding molecule.

In the first step of the method, the fluorescent dyes, including the reactive dyes and dye-conjugates of the invention are generally utilized by combining a dye of the invention as described above with the sample of interest under conditions selected to yield a detectable optical response. The dye-conjugate typically forms a covalent or
5 non-covalent association or complex with an element of the sample, or is simply present within the bounds of the sample or portion of the sample. The sample is then illuminated at a wavelength selected to elicit the optical response. Typically, staining the sample is used to determine a specified characteristic of the sample by further comparing the optical response with a standard.

10

In one aspect of the invention, the dye-conjugate is a labeled protein, such as an antibody, antibody fragment, avidin or strepavidin, and the like. In this embodiment the dye-conjugate is used to detect a complementary specific binding pair which is typically, an antigen, a hapten or a biotin. Such dye-conjugates of the invention can be
15 use for detection of an analyte in a sample, using a variety of applications. Primary applications include immunofluorescence, fluorescence in-situ hybridization (FISH), flow cytometry, labeling of receptors, and tracing of labeled cells. For certain detection methods e.g., those based on cytometry which inherently involve spatial separation/resolution of bound vs. unbound conjugate, it is not necessary to remove the
20 unbound dye-conjugate from the analysis mixture. However, for other detection methods, e.g. immunoassays not involving spatial separation/resolution of bound vs. unbound conjugate, it may be necessary to remove the unbound dye-conjugate from the analysis mixture prior to detecting the optical response.

25 In another aspect, the fluorescent dyes and dye-conjugates of the invention may have utility as laser dyes. Since the dyes and dye-conjugates of the invention are excited by an ultraviolet or violet laser, these dyes and dye-conjugates are particularly useful for multiplex assays with other dyes that are excited at longer wavelengths. In one embodiment, these dyes and dye-conjugates are excited at about 350-405 nm and when
30 used with dyes that are excited at longer wavelengths (e.g., 480 nm and longer), their emission spectra are distinguishable from the dyes that are excited at longer wavelengths.

In another aspect of the invention, the dye-conjugates of the invention can be used in a multi-color method for detecting one or more analytes in a sample. In one embodiment, the multiple dye-conjugates can be used to detect multiple analytes in a sample using a multi-color analysis. The method includes incubating the sample with a composition of the invention that includes multiple dye-conjugates, for example, a first dye-conjugate and second dye-conjugate. In this composition, the component of the first dye-conjugate is a binding partner for the first analyte and the component of the second dye-conjugate is a binding partner for the second analyte. The incubation continues under conditions appropriate to induce an interaction between the first analyte and the first dye-conjugate. During this incubation period, it is generally preferred that a similar interaction occurs between the second analyte and second dye-conjugate, however, it is within the scope of the invention to change the incubation conditions as necessary to drive the formation of a dye-conjugate-analyte complex between the second dye-conjugate and second analyte. Following the formation of at least the first dye-conjugate-analyte complex, the sample is illuminated with light of a wavelength appropriate to cause the complex to fluoresce, thereby detecting the first analyte. The second analyte is detected in a similar manner and may be detected simultaneously with the first analyte or by the sequential illumination of the sample with wavelengths appropriate to induce the fluorescence of each fluorescent dye-conjugate. In a particular aspect, the illumination and/or detection step comprises a flow cytometer.

Alternatively, the multiple dye-conjugates, preferably fluorescing at different wavelengths can be used to detect different features of an analyte. For example, a cell or epitopes of an analyte, can be labeled with different colored dye-conjugates, the target is detected and its identity is confirmed using the co-localization of each color on each target.

The dyes and dye-conjugates of the invention may be used for diagnostic applications. For example, the dyes and dye-conjugates of the invention may be useful for detecting the presence of a pathogenic organism (e.g., bacterial, viral, fungal) in a subject. Alternatively, presence of a particular analyte and/or amount of a particular analyte and/or variants of a particular analyte may be associated with a pathological condition in a subject. The dyes and dye-conjugates of the invention that bind to such analytes may have utility for determining the presence and/or amount of a particular analyte

WO 2011/008912

PCT/US2010/042063

and/or its variants, thus diagnosing the presence and/or extent of a pathological condition.

KITS OF INVENTION

5

One aspect of the present invention is the formulation of kits that facilitate the practice of various assays using any of the dyes of the invention, as described above. The kits of the invention typically comprise a fluorescent dye of the invention, either present as a chemically reactive label useful for preparing dye-conjugates, or present as a dye-conjugate where the conjugated molecule is a specific binding pair member. Selected
10 conjugated molecules include without limitation, polymers of biological molecules (e.g. proteins, nucleic acids or carbohydrates). In specific embodiments, the dyes of the present invention listed in Tables 3 and 4 are particularly suited for the preparation of such a kit.

15

Preferably, the kits comprise one or more of fluorescent dyes listed in Table 5.




20

25

30

Table 5

No.	Formula
Dye1	
Dye2	
Dye3	
Dye4	

Dye5	
Dye6	
Dye7	

The kits can also comprise one or more of the dye-conjugates of Dye1, Dye2, Dye3, Dye4, Dye5, Dye6, or Dye7.

5

In an exemplary embodiment, the kit includes a reactive dye of the invention and instructions for conjugating the dye to a molecule possessing an appropriate functional group, and optionally for recovering the dye-conjugate.

- 10 In another exemplary embodiment, the kit includes instructions for performing an assay that detects an analyte or ligand in a sample. For example, in one embodiment, instructions are provided for detecting cell surface receptors, or an enzyme, or other ligands that can bind antibodies.

WO 2011/008912

PCT/US2010/042063

The kit optionally further comprises one or more buffering agents, typically present as an aqueous solution. The kits of the invention optionally further comprise additional detection reagents, a purification medium for purifying the resulting labeled molecules, fluorescent standards, enzymes, enzyme inhibitors, organic solvent, or instructions for

5 carrying out a method of the invention.

The various embodiments of the invention are demonstrated further by the following illustrative examples. The examples are offered by way of illustration and are not intended to limit the scope of invention or claims in any manner.

10

EXAMPLE 1

The following example provides a description of a general method for the synthesis of dihalo derivatives of firefly luciferin and their respective NHS esters.

15


1. Preparation of Appel Salt

20 A 500-mL round-bottomed flask was charged with chloroacetonitrile (20 mL), sulfur monochloride (120 mL), and CH_2Cl_2 (110 mL) under argon. The reaction stirred under argon for 3 days while a brown precipitate formed. The precipitate was collected by suction filtration and washed with CH_2Cl_2 (300 mL) and hexanes (300 mL). The washed solid was vacuum dried to give the product as a brown solid (75% yield).

25

2. Preparation of 2-Cyanobenzothiazoles

The following procedure is for the preparation of 5,7-dichloro-6-hydroxy-2-cyanobenzothiazole. The preparation of 5,7-difluoro-6-hydroxy-2-cyanobenzothiazole followed the same procedure with very similar results.



10

A 5-L round-bottomed flask was charged with 1,3-dichloro-2-methoxynitrobenzene (0.33 moles), ammonium chloride (3.3 moles), and ethanol (1200 mL). Zinc dust (3.3 moles) is added rapidly to the stirred mixture. The reaction mixture warmed to near reflux after several minutes and was then allowed to cool to room temperature. After stirring overnight, the suspension was filtered to remove residual zinc and ammonium chloride. The filtrate was concentrated under reduced pressure and the resulting solid was resuspended in CH₂Cl₂ (1500 mL). The suspension was filtered and the filtrate concentrated under reduced pressure to give the substituted aniline as a white solid (0.33 moles).

20

1A Y = Cl ¹H NMR (400 MHz, CDCl₃): δ 6.601 (s, 2H), 3.811 (s, 3H), 3.6-3.7 (broad s, 2H)

WO 2011/008912

PCT/US2010/042063

1B Y = F ¹H NMR (400 MHz, CDCl₃): δ 6.22 (m, 2H), 3.860 (s, 3H), 3.6-3.7 (broad s, 2H)

A 2-L three-necked round-bottomed flask was oven-dried and cooled under an argon flow. The flask was charged with 1,3-dichloro-2-methoxyaniline (0.33 moles), pyridine (75 mL), and CH₂Cl₂ (550 mL). The appel salt (0.33 moles) was added in portions over 5 minutes causing the solution to reflux. TLC showed the complete disappearance of the aniline after 15 minutes. The reaction mixture was concentrated to an orange solid. The solid was dissolved in 2.5 L of ethyl acetate. The organic solution was washed with type 1 water (2 x 1000 mL) and brine (1000 mL). The washed organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the appel salt adduct as a solid.

2A Y = Cl ¹H NMR (400 MHz, CDCl₃): δ 7.203 (s, 2H), 3.933 (s, 3H)

2B Y = F ¹H NMR (400 MHz, CDCl₃): δ 6.87 (m, 2H), 4.030 (s, 3H)

A 500-mL three-necked round-bottomed flask was equipped with an argon inlet line and a water-cooled condenser with a bubbler connected to the outlet. The flask was charged with appel salt adduct (0.134 moles) and heated to a surface temperature of 140° C for 1 hour. Toluene (50 mL) was added and the reaction was cooled to 90° C. CH₂Cl₂ (100 mL) and silica (50 g) were added. The resulting slurry was filtered and the filter cake was washed with CH₂Cl₂ (200 mL). The combined filtrate and washes were concentrated and flash chromatography (CH₂Cl₂/hexanes) provided the 5,7-dichloro-6-methoxy-2-cyanobenzothiazole as a solid (0.031 moles).

3A Y = Cl ¹H NMR (400 MHz, CDCl₃): δ 8.194 (s, 1H), 4.027 (s, 3H)

3B Y = F ¹H NMR (400 MHz, CDCl₃): δ 7.981 (dd, J = 10 Hz & 1.6 Hz, 1H), 4.156 (s, 3H)

A 250-mL round-bottomed flask was charged with pyridinium hydrochloride (160 mmoles) and 5,7-dichloro-6-methoxy-2-cyanobenzothiazole (8.2 mmoles). The reaction was padded under argon and then heated at 195° C for 3 hours. After cooling to room temperature, type 1 water (150 mL) was added and the mixture was sonicated to dissolve the solids. The aqueous solution was extracted with ethyl acetate. The

WO 2011/008912


PCT/US2010/042063

ethyl acetate was dried over anhydrous Na_2SO_4 and concentrated to give the product as a yellow solid (2.8 mmol).

4A Y = Cl ^1H NMR (400 MHz, acetone- d_6): δ 8.313 (s)

4B Y = F ^1H NMR (400 MHz, acetone- d_6): δ 8.385 (dd, $J = 10.4$ Hz & 1.6 Hz)

3. Preparation of Amino Acids




5A-F

	W	Z ₁	LH	X ₂
5A	H	H	OH	O
5B	H	Me	OH	S
5C	Me	H	OH	S
5D	Me	H	NHCH ₂ CO ₂ H	S
5E	Me	H	NHCH ₂ CH ₂ CO ₂ H	S
5F	Me	H	NHCH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CO ₂ H	S

15


Synthesis of Modified Penicillamines



A 2000-mL round-bottomed flask was charged with DL-penicillamine (10 g) and acetone (1000 mL). The reaction was refluxed until all the solid had dissolved (approximately 24 hours). The reaction mixture was filtered while hot and cooled to room temperature overnight. A small amount of crystals had formed overnight. The reaction mixture was kept at -20°C overnight. The resulting solids were collected by suction filtration and washed with acetone (300 mL). The thiazole product was dried under argon (11.9 g). ^1H NMR (400 MHz, DMSO- d_6): δ 3.750 (s, 1H), 1.556 (s, 6H), 1.438 (s, 3H), 1.185 (s, 3H)

A 250-mL round-bottomed flask was charged with thiazole acid (11.9 g) and sodium formate (4.3 g) under argon. Formic acid (98%, 100 mL) was added and the reaction mixture stirred under argon. The reaction mixture was cooled to 0° C in an ice/water bath. Acetic anhydride (33.3 mL) was added drop wise over 45 minutes while taking
 5 care that the temperature remained below 5° C. The ice/water bath was removed and the reaction was stirred overnight at room temperature. The solvents were removed under vacuum at 35° C producing a white solid. The solid was stirred in type 1 water (200 mL) and collected by suction filtration. The solid was washed with type 1 water (100 mL) and air-dried (7.4 g). The combined filtrate and wash was extracted with
 10 ethyl acetate (3 x 300 mL). The extracts were combined, dried over Na₂SO₄, and filtered. The filtrate was concentrated to give a pale yellow solid (6.8 g). ¹H NMR showed the two solids to be an isomeric mixture (~85:15) of the target compound. ¹H NMR (400 MHz, DMSO-d₆ 12.916 (s, 1H), 8.446 (s, 0.86H), 8.195 (s, 0.14H), 4.641 (s, 0.15H), 4.502 (s, 0.85H), 1.828 (s, 6H), 1.590 (s, 3H), 1.362 (s, 3H)

15



20 A round-bottomed flask was charged with formylated thiazole acid (1 equivalent), triethylamine (2 equivalents), and CH₂Cl₂. The reaction mixture was stirred at 0° C in an ice/water bath under argon. Isobutylchloroformate (1 equivalent) was added via syringe. After the reaction mixture had stirred for 1 hour, the amino ester (1 equivalent) was added. The ice/water bath was removed and the reaction mixture
 25 stirred overnight at room temperature. The reaction mixture was washed with 1.0 M HCl (75 mL), 5% NaHCO₃ (75 mL), and type 1 water (75 mL). The organic layer was concentrated under reduced pressure and flash chromatography (50:50 ethyl acetate:hexanes) gave the target product (60-70 % yield). ¹H NMR showed an isomeric mixture of products.

30

WO 2011/008912

PCT/US2010/042063

$n = 0$ ^1H NMR (400 MHz, CDCl_3): 8.389 (s, 0.88H), 8.328 (s, 0.12H), 6.932 (m, 0.15H), 6.383 (m, 0.82), 4.620 (s, 0.82H), 4.367 (s, 0.15H), 4.030 (dd, $J = 18.4$ Hz & 5.2 Hz, 1H), 3.91 (m, 1H), 2.00-1.94 (m, 6H), 1.70-1.59 (m, 3H), 1.47 (m, 12H)


5 $n = 1$ ^1H NMR (400 MHz, CDCl_3): 8.358 (s, 0.79H), 8.291 (s, 0.15H), 6.973 (m, 0.15H), 6.486 (m, 0.78), 4.418 (s, 0.80H), 4.313 (s, 0.13H), 3.6-3.4 (m, 2H), 2.43 (m, 2H), 2.00-1.94 (m, 6H), 1.68-1.60 (m, 3H), 1.44 (m, 12H)

10 A round-bottomed flask was charged with the formylated thiazole ester, and dioxane:2.0 M HCl (50:50). The reaction mixture was heated at 85° C under argon overnight. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to give the modified penicillamine as a gummy solid.

15 $n = 0$ ^1H NMR showed an isomeric mixture of products. ^1H NMR (400 MHz, D_2O): 4.128 & 4.083 (s, 1H), 4.043 & 4.000 (s, 2H), 1.557 (s, 3H), 1.481 (s, 3H)

$n = 1$ ^1H NMR (400 MHz, D_2O): 3.702 (s, 1H), 3.45-3.38 (m, 1H), 3.27-3.19 (m, 1H), 2.441 (m, 2H), 1.295 (s, 3H), 1.233 (s, 3H)

20 Synthesis of Methylated Cysteine



25

Cysteine methyl ester (8.6 g), pivalaldehyde (11 mL), and triethylamine (8 mL) were stirred in pentane. The reaction mixture was refluxed while water was removed via a

Dean-Stark trap. After 36 hours, the reaction was cooled to room temperature and solids formed. The solids were collected by suction filtration and washed with ether.

A 100-mL round-bottomed flask was charged with thiazole ester (4.0 g), formic acid
5 (98%, 30 mL) and sodium formate (1.5 g). The reaction mixture was stirred under argon and cooled to 0° C in an ice/water bath. Acetic anhydride (5.7 mL) was added drop wise over 30 minutes while taking care that the temperature remained below 5° C. The ice/water bath was removed and the reaction was stirred overnight at room temperature. The solvents were removed under vacuum and the resulting solid was
10 stirred in type 1 water (100 mL). The pH was adjusted to 7-8 by the addition of solid NaHCO₃. The aqueous solution was extracted with ether (3 x 100 mL). The extracts were combined, dried over Na₂SO₄, and filtered. The filtrate was concentrated to give a solid. The solids were recrystallized from ethyl acetate/hexanes. ¹H NMR (400 MHz, CDCl₃): 8.317 (s, 1H), 4.858 (t, J = 8.8 Hz, 1H), 4.708 (s, 1H), 3.739 (s, 3H), 3.258 (m,
15 2H), 1.000 (s, 9H)

A solution of lithium diisopropylamine (12 mmoles) in anhydrous THF (100 mL) at -78° C was stirred under argon. 1,3-Dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU, 10 mL) was added and the reaction mixture stirred for 90 minutes.
20 Formylated thiazole ester was added (10 mmoles) and the reaction mixture was stirred at -78° C for 1 hour. MeI (15 mmoles) was added and the reaction mixture was stirred at -78° C for 2 hours. The dry ice/ acetone bath was removed and the reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure and the residue was taken up in brine. The aqueous solution
25 was extracted with ether (100 mL). The extract was dried over Na₂SO₄, and filtered. The filtrate was concentrated to give a solid. The solid was purified by flash chromatography with 10% ethyl acetate/hexanes. ¹H NMR showed a diastereomeric mixture of products. ¹H NMR (400 MHz, CDCl₃) major product: 8.252 (s, 1H), 5.263 (s, 1H), 3.738 (s, 3H), 3.293 (d, J = 11.6 Hz, 1H), 2.695 (d, J = 11.2 Hz, 1H), 1.038 (s,
30 9H); minor product: 8.377 (s, 0.4H), 5.274 (s, 0.4H), 3.790 (s, 1.3H), 3.612 (d, J = 12 Hz, 0.5H), 2.831 (d, J = 12 Hz, 1H), 0.929 (s, 4.6H)

The methylated thiazole ester (1.8 g) and 5 M HCl (30 mL) were heated at 105° C for 3 days. The reaction mixture was cooled to room temperature and extracted with ethyl


WO 2011/008912

PCT/US2010/042063

acetate (2 x 100 mL). The aqueous phase was concentrated under reduced pressure to give a gummy solid.

4. Preparation of 6-Hydroxy Luciferin Acids

This is a typical procedure for the preparation of the basic luciferin structure.



	W	Z ₁	LH	X ₂	Y ₁ /Y ₂
6A	H	H	OH	O	F
6B	H	Me	OH	S	F
6C	Me	H	OH	S	F
6D	Me	H	NHCH ₂ CO ₂ H	S	F
6E	Me	H	NHCH ₂ CH ₂ CO ₂ H	S	F
6F	Me	H	NHCH ₂ CH ₂ CO ₂ H	S	Cl
6G	H	H	OH	S	F
6H	Me	H	OH	S	Cl
6I	H	H	OH	S	Cl

- 15 A 100-mL round-bottomed flask was charged with the cyanobenzothiazole (0.5 mmole) and methanol (25 mL). The stirred reaction mixture was sparged with argon for 1 hour. An argon-sparged solution of the amino acid (1 mmole) in water with a pH of 9 (pH adjustment by the addition of Na₂CO₃) was prepared and added. The reaction mixture was stirred for 2 hours under argon and TLC indicated the reaction was completed.
- 20 The reaction was concentrated under reduced pressure to ~10 mL and diluted with type 1 water (40 mL). The aqueous solution was extracted with ethyl acetate (3 x 75 mL) and the extracts were set aside. The aqueous solution was acidified to pH 2 by the drop wise addition of concentrated HCl. The acidified aqueous solution was extracted again with ethyl acetate (3 x 75 mL). The six extracts were combined, dried over anhydrous
- 25 Na₂SO₄, and concentrated to give the product as a solid (40-90% yields).


WO 2011/008912

PCT/US2010/042063

- 6A ¹H NMR (400 MHz, CD₃OD): 7.688 (dd, J = 11 Hz & 1.4 Hz, 1H), 5.017 (dd, J = 10.2 Hz & 8.2 Hz, 1H), 4.774 (m, 2H)
- 5 6B ¹H NMR (400 MHz, CDCl₃): 7.795 (dd, J = 10 Hz & 1.6 Hz, 1H), 3.932 (d, J = 11.8 Hz, 1H), 3.434 (d, J = 11.8 Hz, 1H), 1.688 (s, 3H)
- 6C ¹H NMR (400 MHz, CD₃OD): 7.646 (dd, J = 10.4 Hz & 1.2 Hz, 1 H), 4.935 (s, 1H), 1.778 (s, 3H), 1.488 (s, 3H)
- 10 6D ¹H NMR (400 MHz, CD₃OD): 8.379 (broad s, 1H), 7.658 (d, J = 10 Hz, 1H), 4.758 (s, 1H), 4.000 (d, J = 6 Hz, 2H), 1.829 (s, 3H), 1.448 (s, 3H)
- 15 6E ¹H NMR (400 MHz, CD₃OD): 8.190 (broad s, 1H), 7.647 (d, J = 10.4 Hz, 1H), 4.705 (s, 1H), 3.56-3.50 (m, 2H), 2.558 (t, J = 6.6 Hz, 2H), 1.814 (s, 3H), 1.395 (s, 3H)
- 20 6F ¹H NMR (400 MHz, CD₃OD): 8.271 (broad s, 1H), 8.093 (s, 1H), 4.794 (s, 1H), 3.610 (m, 2H), 2.640 (t, J = 6.6 Hz, 2H), 1.898 (s, 3H), 1.478 (s, 3H)
- 6G ¹H NMR (400 MHz, acetone- d₆): 9.838 (s, 1H), 7.810 (dd, J = 10.8 Hz, & 1.6 Hz, 1H), 5.504 (t, J = 9.2 Hz, 1H), 3.860 (m, 2H)
- 25 6H ¹H NMR (400 MHz, DMSO- d₆): 8.266 (s, 1H), 5.007 (s, 1H), 1.725 (s, 1H), 1.447 (s, 1H)
- 6I ¹H NMR (400 MHz, acetone- d₆): 8.149 (s, 1H), 5.505 (t, J = 9.2 Hz, 1H), 3.855 (m, 2H)
- 30 Compound 6J (L= NH₂(CH₂)₅COOH) was prepared in three steps starting from compound 6H. Reaction of carboxylic acid 6H with N-hydroxysuccinimide and DCC produced NHS ester 7H which was coupled with methyl 6-aminohexanoate. The resulting methyl ester was hydrolyzed with base to produce 6J.

WO 2011/008912

PCT/US2010/042063

5. Preparation of Luciferin NHS esters

5

	W	Z ₁	L	Y ₁ /Y ₂
7A	H	H	O	F
7B	H	Me	O	F
7C	Me	H	O	F
7D	Me	H	NHCH ₂ CO ₂	F
7E	Me	H	NHCH ₂ CH ₂ CO ₂	F
7F	Me	H	NHCH ₂ CH ₂ CO ₂	Cl
7G	H	H	O	F
7H	Me	H	O	Cl
7J	Me	H	NH(CH ₂) ₅ CO ₂	Cl

10 This is a typical procedure for the preparation of the luciferin NHS esters.

A 10-mL round-bottomed flask was padded with argon and charged with the luciferin (0.3 mmole), N-hydroxysuccinimide (0.3 mmole), and DCC (0.305 mmole). Anhydrous THF (4 mL) was added and the reaction mixture was stirred for 3 hours.

15 The solvent was removed under reduced pressure and a small volume of CH₂Cl₂ was added. A white precipitate formed and was removed by filtration. The filtrate was concentrated under reduced pressure to form a gummy solid. The solid was dissolved in a minimum of THF and the addition of hexanes resulted in the solid reforming. The solvents were decanted and the procedure was repeated 2 additional times to give the

20 product as a yellow solid (60% yields).

7B ¹H NMR (400 MHz, CDCl₃): 7.650 (m, 1H), 4.093 (d, J = 11.4 Hz, 1H), 3.481 (d, J = 11.4 Hz, 1H), 1.814 (s, 3H)

25

WO 2011/008912


PCT/US2010/042063

- 7C ¹H NMR (400 MHz, CDCl₃): 7.660 (dd, J = 10.2 Hz & 1.4 Hz, 1 H), 5.153 (s, 1H), 2.858 (s, 4H), 1.874 (s, 3H), 1.653 (s, 3H)
- 5 7D ¹H NMR (400 MHz, CDCl₃): 7.678 (d, J = 10 Hz, 1H), 7.608 (broad s, 1H), 4.713 (s, 1H), 4.614 (dd, J = 18 Hz & 6.6 Hz, 1H), 4.389 (dd, J = 18 Hz & 5.4 Hz, 1H), 2.829 (s, 4H), 1.907 (s, 3H), 1.447 (s, 3H)
- 10 7E ¹H NMR (400 MHz, CDCl₃): , 7.670 (d, J = 10 Hz, 1H), 7.616 (broad s, 1H), 4.660 (s, 1H), 3.82-3.65 (m, 2H), 2.899 (t, J = 6 Hz, 2H), 2.789 (s, 4H), 1.899 (s, 3H), 1.412 (s, H)
- 15 7F ¹H NMR (400 MHz, CDCl₃): 8.073 (s, 1H), 7.653 (broad s, 1H), 4.699 (s, 1H), 3.9-3.7 (m, 2H), 2.932 (t, J = 6.0 Hz, 2H), 2.820 (s, 4H), 1.940 (s, 3H), 1.451 (s, 3H)
- 20 7H ¹H NMR (400 MHz, acetone- d₆): 8.13 (s, 1H), 5.59 (s, 1H), 2.33 (s, 4H), 1.89 (s, 3H), 1.64 (s, 3H)
- 25 7J ¹H NMR (400 MHz, acetone- d₆): 9.53 (broad s, 1H), 8.10 (s, 1H), 7.78 (m, 1H), 4.69 (s, 1H), 3.22-3.42 (m, 2H), 2.81 (s, 4H), 2.62 (t, 2H), 1.86 (s, 3H), 1.76-1.69 (m, 2H), 1.63-1.56 (m, 2H), 1.50-1.44 (m, 2H), 1.42 (s, 3H)

6. Preparation of Luciferin Maleimide Derivatives


WO 2011/008912

PCT/US2010/042063



- 5 The luciferin NHS ester (22 mg) was stirred in anhydrous THF. N-(2-aminoethyl)maleimide-trifluoroacetic acid (13 mg) was added followed by aqueous NaHCO₃. The mixture was stirred for 4 hours and then concentrated under reduced pressure. The mixture was dissolved in 5 mL of water and extracted with ethyl acetate. The extract was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure.
- 10 The solids were dissolved in acetone and hexanes was added to generate a precipitate that was air-dried (13 mg). ¹H NMR (400 MHz, DMSO- d₆): 8.118 (m, 1H), 7.886 (d, J = 10.8 Hz, 1H), 6.972 (s, 2H), 4.670 (s, 1H), 3.38 (m, 2H), 3.33 (m, 2H), 1.678 (s, 3H), 1.264 (s, 3H)

15 Preparation of fluorescent dye1 (7H):



- Fluorescent dye 1 (LD1; 7H) was synthesized using the synthesis scheme described above. Briefly, 2-cyano-5,7-dichloro-6-hydroxybenzothiazole (4A), synthesized using the method describe above was reacted with D,L-Penicillamine (5C) under conditions described in detail above to afford 5,7-dichloro-6-hydroxy luciferin acid (6H).
- 20


- For the synthesis of 5,7-dichloro-6-hydroxy luciferin NHS ester, the dichloro-6-hydroxy lucifern acid (6H) was reacted with N-hydroxysuccinimide in the presence of
- 25

DCC as described above. The resulting NHS ester (7H) was purified and characterized by NMR as described above.

Preparation of fluorescent dye2 (7J):

5

10




Fluorescent dye 2 (LD2; Compound 7J) was synthesized using synthesis scheme described above, except that 2-cyano-5,7-dichloro-6-hydroxybenzothiazole (4A), was reacted with compound 5F under conditions described in detail above to afford 5,7-dichloro-6-hydroxy luciferin acid (6J).

15

The dichloro-6-hydroxy luciferin acid (6J) was next reacted with N-hydroxysuccinimide in the presence of DCC as described above to yield the NHS ester (7J), which was purified and analyzed by NMR as described above.

20 Preparation of fluorescent dye3 (7C):




25 Fluorescent dye 3 (LD3; 7C) was synthesized using synthesis scheme as described above, except that 2-cyano-5,7-difluoro-6-hydroxybenzothiazole (4B), was reacted with D,L- Penicillamine (5C) under conditions described in detail above to afford 5,7-difluoro-6-hydroxy luciferin acid (6C).

The difluoro-6-hydroxy luciferin acid (6C) was next reacted with N-hydroxysuccinimide in the presence of DCC as described above to yield the 5,7-difluoro-6-hydroxy luciferin NHS ester (7C), which was purified and analyzed by NMR as described above.

Preparation of fluorescent dye4 (7B):


5



Fluorescent dye 4 (LD4; 7B) was synthesized using synthesis scheme as described
 10 above, except that 2-cyano-5,7-difluoro-6-hydroxybenzothiazole (4B), was reacted with
 compound 5B under conditions described in detail above to afford 5,7-difluoro-6-
 hydroxy luciferin acid (6C). The 5,7-difluoro-6-hydroxy luciferin acid (6C) was next
 reacted with N-hydroxysuccinimide in the presence of DCC as described above to yield
 the 5,7-difluoro-6-hydroxy luciferin NHS ester (7C), which was purified and analyzed
 15 by NMR as described above.


Preparation of fluorescent dye5 (7E)

20



Fluorescent dye 5 (LD5; 7E) was synthesized using synthesis scheme as described
 25 above, except that 2-cyano-5,7-difluoro-6-hydroxybenzothiazole (4B), was reacted with
 compound 5E under conditions described in detail above to afford 5,7-difluoro-6-
 hydroxy luciferin acid (6E). The 5,7-difluoro-6-hydroxy luciferin acid (6E) was next
 reacted with N-hydroxysuccinimide in the presence of DCC as described above to yield
 the 5,7-difluoro-6-hydroxy luciferin NHS ester (7E), which was purified and analyzed
 30 by NMR as described above.

5



CC(C)(C)OC(=O)CCCC(NC(=O)C(C)(C)C)C(=O)O
 $\xrightarrow[2) \text{NaBH}_4]{1) \text{ClCO}_2\text{Et}}$
CC(C)(C)OC(=O)CCCC(NC(=O)C(C)(C)C)CO
 $\xrightarrow[\text{pyridine}]{\text{TsCl}}$
CC(C)(C)OC(=O)CCCC(NC(=O)C(C)(C)C)COST(=O)(=O)c1ccc(C)cc1
 $\xrightarrow{\text{CH}_3\text{COSK}}$
CC(C)(C)OC(=O)CCCC(NC(=O)C(C)(C)C)CS(=O)C
 $\xrightarrow{\text{HCl}}$
CC(C)(C)OC(=O)CCCC([NH3+])CS

10

N-Boc-L-glutamic acid γ -t-butyl ester (4.3 g) was stirred in CH_2Cl_2 at 0°C . Triethylamine (2 mL) was added via syringe followed by the addition of ethyl chloroformate (1.4 mL). The reaction mixture was stirred for 60 minutes, filtered, and the solids were washed with anhydrous THF. The filtrate and washes were combined and transferred to an addition funnel. The solution was added drop wise over 30 minutes to a stirred solution of NaBH_4 in H_2O (20 mL) at 0°C . The reaction mixture was stirred at 0°C for 5 hours before stirring at room temperature overnight. The reaction mixture was concentrated under reduced pressure and the residue was diluted with ethyl acetate. The solution was washed with water and brine, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. Flash chromatography (20:80 ethyl acetate:hexanes) gave the product as an oil. ^1H NMR (400 MHz, CDCl_3): 4.814 (broad s, 1H), 3.62-3.49 (m, 3H), 2.496 (broad s, 1H), 2.290 (q, $J = 6.8\text{ Hz}$, 2H), 1.83-1.69 (m, 2H), 1.406 (s, 18H).

25

45


WO 2011/008912

PCT/US2010/042063

The solution was warmed to room temperature and stirred overnight. The reaction mixture was concentrated under reduced pressure and the resulting solids were dissolved in ethyl acetate. The solution was washed with saturated NaHCO_3 followed by brine. The washed solution was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. Flash chromatography gave the product as an oil. ^1H NMR (400 MHz, CDCl_3): 7.744 (d, $J = 8.8$ Hz, 2H), 7.311 (d, $J = 8.8$ Hz, 2H), 4.623 (d, $J = 8.4$ Hz, 1H), 4.00-3.91 (m, 2H), 3.739 (broad s, 1H), 2.411 (s, 3H), 2.224 (m, 2H), 1.719 (m, 2H), 1.390 (s, 9H), 1.350 (s, 9H)

10 A solution of the tosylated ester (3.9 g) and potassium thioacetate (2.9 g) in anhydrous DMF was stirred at room temperature for 3 days. The reaction mixture was diluted with ethyl acetate (300 mL) and washed with brine (3 x 100 mL). The washed solution was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. Flash chromatography (15:85 ethyl acetate:hexanes) gave the product as an oil. ^1H NMR (400 MHz, CDCl_3): 4.522 (d, $J = 8$ Hz, 1H), 3.704 (broad s, 1H), 3.07-2.94 (m, 2H), 2.311 (s, 3H), 2.264 (t, $J = 7.2$ Hz, 2H), 1.80-1.60 (m, 2H), 1.404 (s, 9H), 1.386 (s, 9H)

The thioacetate (2.5 g) was heated in degassed 6 N HCl at 105°C overnight. The reaction mixture was cooled to room temperature and concentrated under reduced pressure to give a red oil. ^1H NMR (400 MHz, D_2O):



25 A flask was charged with 2-cyano-5,7-difluoro-6-hydroxybenzothiazole (110 mg) and methanol (25 mL). The stirred reaction mixture was sparged with argon for 1 hour. An

WO 2011/008912

PCT/US2010/042063

argon-sparged solution of the amino acid (300 mg) in water with a pH of 9 (pH adjustment by the addition of Na_2CO_3) was prepared and added to the reaction mixture in 3 portions at 1 hour intervals. The reaction mixture was stirred overnight. The reaction was concentrated under reduced pressure to ~10 mL and diluted with type 1
5 water (40 mL). The aqueous solution was extracted with ethyl acetate (50 mL) and the extract was set aside. The aqueous solution was acidified to pH 2 by the drop wise addition of concentrated HCl. The acidified aqueous solution was extracted again with ethyl acetate (2 x 50 mL). The 2 extracts were combined, dried over anhydrous Na_2SO_4 , and concentrated to give the product as a solid (130 mg). ^1H NMR (400 MHz, CD_3OD): 7.641 (d, $J = 10.4$ Hz, 1H), 4.694 (m, 1H), 3.612 (t, $J = 8.8$ Hz, 1H), 3.179 (t,
10 $J = 9.4$ Hz, 1H), 2.550 (m, 2H), 2.037 (m, 2H)


A flask was padded with argon and charged with the deoxoluciferin acid (41 mg), N-hydroxysuccinimide (15 mg), and DCC (26 mg). Anhydrous THF (4 mL) was added
15 and the reaction mixture was stirred for 3 hours. The reaction solution was filtered to remove solids. Hexanes were added to precipitate solids which were collected by suction filtration. The solid was redissolved in THF and hexanes were added to precipitate solids. The solids were collected by suction filtration and air-dried. ^1H NMR (400 MHz, CDCl_3): 7.666 (dd, $J = 10.4$ Hz & 1.6 Hz, 1H), 4.772 (m, 1H), 3.612
20 (m, 1H), 3.158 (dd, $J = 11$ Hz & 8.6 Hz, 1H), 3.00-2.86 (m, 2H), 2.821 (s, 4H), 2.188 (m, 2H)

25

30

WO 2011/008912

PCT/US2010/042063

Preparation of dehydroluciferin (10).

- 5 The following reaction description is for the example **10A** and similar reaction conditions were used in the case of **10B**. ^1H NMR data are given for both cases.

2-cyano-5,7-dichloro-6-hydroxybenzothiazole (2.8 mmole) was stirred in pyridine (20 mL) and triethylamine (0.3 mL). Hydrogen sulfide was bubbled through the solution
 10 for 4.5 hours. The addition of hydrogen sulfide was stopped and the reaction mixture stirred at room temperature overnight. The reaction mixture was stripped under reduced pressure and the resulting solid was recrystallized from methanol. ^1H NMR (400 MHz, CD_3OD): 8.000 (s, 1H)

- 15 The benzothiazole product (55 mg) from above was suspended in methanol (5 mL). Ethyl bromopyruvate (100 mg) was added and the reaction mixture was stirred at room temperature overnight. The reaction mixture was heated at 80°C for 7 hours. The reaction mixture was cooled to room temperature and solids formed. The solids were collected by suction filtration, washed with methanol, and air-dried (35 mg).

20

R = H ^1H NMR (400 MHz, CDCl_3): 8.295 (s, 1H), 7.992 (s, 1H), 4.436 (q, $J = 7\text{ Hz}$, 2H), 1.414 (t, $J = 7\text{ Hz}$, 3H)

WO 2011/008912

PCT/US2010/042063

R = Me Methyl ester ^1H NMR (400 MHz, CDCl_3): 7.964 (s, 1H), 3.951 (s, 3H), 2.837 (s, 3H)

- 5 The dehydroluciferin ethyl ester (30 mg) was suspended in methanol and 1 N NaOH. The reaction was stirred at room temperature for 2 days. The reaction mixture was acidified to pH of 2 with 1 N HCl and extracted with ethyl acetate (2 x 50 mL). The combined extracts were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give the product as a solid.


10

10A ^1H NMR (400 MHz, CD_3OD): 8.518 (s, 1H), 8.001 (s, 1H)

10B ^1H NMR (400 MHz, CD_3OD): 7.965 (s, 1H), 2.810 (s, 3H)

15

8. Preparation of 5-Chloro-4-Hydroxy Luciferin Acid (**11A**) and NHS Ester (**11B**)



The synthetic route to the 4-hydroxy luciferin acids and NHS esters is identical to the route used to prepare the 6-hydroxy luciferin acids and NHS esters. NMR data is given for the above compounds.

25

3-Chloro-2-methoxy aniline appel salt: ^1H NMR (400 MHz, CDCl_3): 7.224 (m, 1H), 7.068 (t, J = 8.2 Hz, 1H), 6.988 (d, J = 8 Hz, 1H), 3.809 (s, 3H)

WO 2011/008912

PCT/US2010/042063

2-Cyano-5-chloro-4-methoxy benzothiazole: ^1H NMR (400 MHz, CDCl_3): 7.591 (d, J = 8.4 Hz, 1H), 7.521 (d, J = 8.8 Hz, 1H), 4.349 (s, 3H)


5 2-Cyano-5-chloro-4-hydroxy benzothiazole: ^1H NMR (400 MHz, CDCl_3): 7.576 (d, J = 8.4 Hz, 1H), 7.416 (d, J = 8.4 Hz, 1H), 6.702 (s, 1H)

5-Chloro-4-hydroxy dimethyl luciferin acid (**11A**): ^1H NMR (400 MHz, CDCl_3): 7.451 (d, J = 8.4 Hz, 1H), 7.370 (d, J = 9.2 Hz, 1H), 6.689 (s, 1H), 4.903 (s, 1H), 1.886 (s, 3H), 1.528 (s, 3H)

5-Chloro-4-hydroxy dimethyl luciferin NHS ester (**11B**): ^1H NMR (400 MHz, CDCl_3): 7.425 (d, J = 8.8 Hz, 1H), 7.354 (d, J = 8.4 Hz, 1H), 6.707 (s, 1H), 5.157 (s, 1H), 2.855 (s, 4H), 1.882 (s, 3H), 1.659 (s, 3H)

15

20 9. Preparation of 6-Chloro-5-Hydroxy Luciferin Acid (**12D**)



25 The synthetic route to the 5-hydroxy luciferin acid is identical to the route used to prepare the 6-hydroxy luciferin acids. NMR data is given for the above compounds.

WO 2011/008912

PCT/US2010/042063

4-Chloro-3-methoxy aniline appel salt (**12A**): ^1H NMR (400 MHz, CDCl_3): 7.396 (d, J = 8.8 Hz, 1H), 6.78-6.76 (m, 2H), 3.882 (s, 3H)


5 2-Cyano-6-chloro-5-methoxy benzothiazole (**12B**): ^1H NMR (400 MHz, CDCl_3):
7.929 (s, 1H), 7.636 (s, 1H), 3.988 (s, 3H)

2-Cyano-6-chloro-5-hydroxy benzothiazole (**12C**): ^1H NMR (400 MHz, CDCl_3):
7.927 (s, 1H), 7.792 (s, 1H), 5.852 (s, 1H)

10 6-Chloro-5-hydroxy dimethyl luciferin acid (**12D**): ^1H NMR (400 MHz, CD_3OD):
7.980 (s, 1H), 7.511 (s, 1H), 4.927 (s, 1H), 1.778 (s, 3H), 1.489 (s, 3H)

WO 2011/008912

PCT/US2010/042063

10. Preparation of 5,7-Dimethyl-6-Hydroxy Dimethyl Luciferin Acid (13E)

5

The synthetic route to the 5,7-dimethyl-6-hydroxy dimethyl luciferin acid is identical to the route used to prepare the 6-hydroxy luciferin acids. NMR data is given for the above compounds.

3,5-Dimethyl-4-methoxy aniline (**13A**): ^1H NMR (400 MHz, CDCl_3): 6.320 (s, 2H), 3.615 (s, 3H), 2.167 (s, 6H)

15 3,5-Dimethyl-4-methoxy aniline appel salt (**13B**): ^1H NMR (400 MHz, CDCl_3): 6.905 (s, 2H), 3.711 (s, 3H), 2.278 (s, 6H)

2-Cyano-5,7-dimethyl-6-methoxy benzothiazole (**13C**): ^1H NMR (400 MHz, CDCl_3): 7.845 (s, 1H), 3.782 (s, 3H), 2.488 (s, 3H), 2.429 (s, 3H)

20


2-Cyano-5,7-dimethyl-6-hydroxy benzothiazole (**13D**): ^1H NMR (400 MHz, CDCl_3): 7.805 (s, 1H), 5.2 (broad s, 1H), 2.443 (s, 3H), 2.401 (s, 3H)

5,7-Dimethyl-6-hydroxy dimethyl luciferin acid (**13E**): ^1H NMR (400 MHz, CD_3OD): 7.641 (s, 1H), 4.905 (s, 1H), 2.401 (s, 3H), 2.333 (s, 3H), 1.777 (s, 3H), 1.491 (s, 3H)

25

11. Preparation of Trifluoromethyl-substituted Luciferin Acids (14E) and NHS Esters (Fluorescence Dye7; 14F):

5



- 10 The synthetic route to the trifluoromethyl-substituted luciferin acids and NHS esters is identical to the route used to prepare the 6-hydroxy luciferin acids and NHS esters. NMR data is given for the above compounds.

15 4-methoxy-3-trifluoromethyl aniline appel salt (**14A**): ¹H NMR (400 MHz, CDCl₃): 7.522 (d, J = 2.4 Hz, 1H), 7.424 (dd, J = 8.6 Hz & 2.6 Hz, 1H), 7.064 (d, J = 9.2 Hz, 1H), 3.916 (s, 3H)

20 2-Cyano-5-trifluoromethyl-6-methoxy benzothiazole (**14B**): ¹H NMR (400 MHz, CDCl₃): 8.384 (s, 1H), 7.451 (s, 1H), 4.033 (s, 3H)

2-Cyano-7-trifluoromethyl-6-methoxy benzothiazole (**14C**): ¹H NMR (400 MHz, CDCl₃): 8.313 (d, J = 8.8 Hz, 1H), 7.374 (d, J = 9.2 Hz, 1H), 3.997 (s, 3H)

25 2-Cyano-5-trifluoromethyl-6-hydroxy benzothiazole (**14D**): ¹H NMR (400 MHz, CDCl₃): 8.368 (s, 1H), 7.492 (s, 1H)

WO 2011/008912

PCT/US2010/042063

5-Trifluoromethyl-6-hydroxy dimethyl luciferin acid (**14E**): ¹H NMR (400 MHz, CD₃OD): 8.168 (s, 1H), 7.479 (s, 1H), 4.927 (s, 1H), 1.779 (s, 3H), 1.492 (s, 3H)

5-Trifluoromethyl-6-hydroxy dimethyl luciferin NHS ester (**14F**): ¹H NMR (400 MHz, CDCl₃): 8.278 (s, 1H), 7.445 (s, 1H), 5.148 (s, 1H), 2.861 (s, 4H), 1.874 (s, 3H), 1.653 (s, 3H)

Fluorescence Measurement of Luciferin compounds

10 The fluorescence measurement experiments were conducted on a HORIBA Jobin Yvon Fluoromax®-3 spectrometer. Typically, the excitation wavelength was set at 405 nm unless otherwise noted. The excitation and emission slits were at 1 nm. A solution of the dye at 20 μM in 25 mM Tris pH 8.0 buffer was used to obtain the fluorescence spectra and determine the intensity at the wavelength of maximum emission shown in
15 Table 6.

Table 6

Luciferin compounds	Fluorescence Intensity at Fluorescence Maxima
Compound 6H	530 nm = 1.1e6 (ex = 405 nm)
Compound 6J	542 nm = 1.75e6 (ex = 405 nm)
Compound 6C	530 nm = 1.1e6 (ex = 405 nm)
Compound 6B	527 nm = 935490 (ex = 405 nm)
Compound 6E	541 nm = 1.3e6 (ex = 405 nm) 541 nm = 4.5e6 (ex = 396 nm)
Compound 9A	532 nm = 964121(ex = 405nm) 532 nm = 1.2e6 (ex = 396 nm)
Compound 14E	532 nm = 570975 (ex = 405 nm) 532 nm = 774112 (ex = 396 nm)
Compound 10A	546 nm = 1.6e6 (ex = 405 nm)

EXAMPLE 2

The following example provides the description of a method for conjugating the fluorescent dyes of the invention to an antibody molecule.

5

In a typical experiment, antibody conjugates of Luciferin derivatives are prepared as follows. The antibody of interest is prepared in 50mM Borate pH 9.0 buffer at 4mg/ml. The dye reagent is dissolved in anhydrous DMSO to give a concentration of 5mg/ml. Predetermined amounts of the dye in DMSO are slowly added to the antibody solution with mixing. The reaction is incubated at room temperature for 60 minutes. Thereafter the reaction is quenched with addition glycyl glycine (200 molar excess to the dye used) solution at 75mg/ml in PBS, 2mM EDTA. The dye-antibody conjugate is separated desalting on a Sephadex G-50 column equilibrated in PBS. The elution is monitored for 280nm absorbance and the antibody containing band is collected from the column. The degree of substitution (F/P Value) is determined by measuring absorbance values for the dye-conjugate at 280nm and at the absorption maxima of the dye peak. The degree of substitution is calculated using the extinction coefficient for the dye being used for the conjugation as shown in Table 7.

20

Table 7

Dyes of the invention	Extinction Coefficient $M^{-1}cm^{-1}$ at Dye absorption maxima
5,7-Dichloroluciferin NHS ester (Dye1)	14662
5,7 dichloro luciferin-Linker-NHS ester (Dye2)	18174
Dimethyl difluoroluciferin NHS ester (Dye3)	17665
Monomethyl difluoroluciferin NHS ester (Dye4)	14665
Difluoro dimethylene Luciferin NHS ester (Dye5)	11532
Difluoro amide Luciferin NHS ester (Dye6)	14990

The following table (Table 8) provides the data for the various conjugates for six of the dyes of the invention to representative antibodies.

5

Table 8

Antibody	Dye	Dye to antibody ratio used	Absorption maxima for conjugate	Degree of substitution	Emission maxima for conjugate
CD4	Dye1	100	400nm	13.5	525nm
CD4	Dye2	60	399nm	30.4	516nm
CD3	Dye2	10	399nm	5.91	516nm
CD45	Dye2	5	399nm	4.2	516nm
CD4	Dye3	100	398nm	11.4	528nm
CD3	Dye3	20	398nm	4.8	528nm
CD45	Dye3	40	398nm	8.24	528nm
CD3	Dye4	30	390nm	2.89	528nm
CD4	Dye5	60	390nm	20.8	527nm
CD3	Dye5	20	390nm	6.2	527nm
CD45	Dye5	15	390nm	8.31	527nm
CD4	Dye6	80	382nm	27.6	517nm
CD3	Dye6	15	382nm	6.8	517nm
CD45	Dye6	10	382nm	6.8	517nm
CD4	Dye7	100	391nm	ND ^a	524nm
CD4	Dye8	100	420nm	ND	465nm
CD4	Dye8	100	420nm	ND	465nm

10

^aND means not determined

EXAMPLE 3:

15 The following example provides method for using the antibody-dye conjugates in flow cytometry.

Flow Analysis

20 Optimum quantities (determined separately by titration) of various compositions of antibody-dye conjugates of the invention were independently combined with a biological specimen containing whole blood (0.1mL), incubated for 10-15 minutes, processed using the standard procedure using VersaLyseTM lysing reagent, and

analyzed using a flow cytometer. Briefly, the treated biological specimen was combined with the VersaLyseTM reagent/0.2% Formaldehyde (1mL) for 10 minutes, spinning down the treated specimen, aspirating the supernatant, resuspending the pellet in PBS (2mL) spinning down the cell suspension and resuspending the pellet in
 5 PBS/0.1% formaldehyde (1mL) for analysis. The lymphocyte population was selected by flow cytometry based on forward and side scatter characteristics. Subpopulations were identified using specific monoclonal antibodies. Fluorescence signals were collected using appropriate band pass filters chosen based on the conjugate's fluorescence emission spectrum.

10

Similarly, the antibody-Pacific OrangeTM conjugates were evaluated. In selected cases, the flow data results of various luciferin derivatives dyes – antibody conjugates were compared with the corresponding antibody-Pacific Orange dye-conjugates. Following table summarizes these results (Table 9). The conjugates of antibody-fluorescent dyes
 15 of the invention showed superior signal/noise values compared to antibody-Pacific Orange conjugate.

Table 9

Conjugate	Signal/Noise values
CD4-Pacific Orange	8.7
CD4-Dye1	20.5
CD4-Dye2	26.8
CD4-Dye3	26.4
CD4-Dye4	27.4
CD4-Dye5	31.0
CD4-Dye7	9.2

20

Two-color flow analysis:

In selected cases, using 405 nm laser line, CD8-Pacific BlueTM dye-conjugate and a
 25 CD4-luciferin derivative dye of the invention, a two color flow analysis was similarly performed. The results were compared with the corresponding performance for CD8-Pacific Blue dye conjugate and a CD4-Pacific Orange dye (Invitrogen Corporation). As shown in Figure 4, the antibody-Dye2 conjugate showed brighter signal when compared to that of antibody-Pacific OrangeTM conjugate.

30

WO 2011/008912

PCT/US2010/042063

Multi-color flow analysis:

For one of the dyes (Dye3) of the invention, a ten-color application was also demonstrated on a 3 laser 10 color PMT instrument equipped with a 405nm laser,
5 488nm laser and 635nm laser. A 550/40 nm band pass filter was used to collect the fluorescence output into the photomultiplier tube. Following antibody reagents were used:


1. CD45 (RA)-FITC
2. CD56-PE
- 10 3. CD45(RO)-ECD
4. CD25-PC5
5. CD19-PC7
6. CD3-APC
7. CD27-APCA700
- 15 8. CD5-APCCy7
9. CD8-Pacific Blue
10. CD4-Pacific Orange/CD4-Dye3.

The histograms shown in Figure 5 illustrate that in the ten-color flow assay, the CD4-
20 Dye3 conjugate gave a brighter signal when compared to that of CD4-Pacific Orange conjugate.

What is claimed is:

1. A fluorescent dye having the formula:

5



wherein

X_1 and X_2 are independently S or O;

10 Y_1 and Y_2 are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl, or one of Y_1 or Y_2 is H, and the other is a haloalkyl group;

wherein each W is independently H or alkyl;

Z_1 is H or alkyl;

15 wherein Z_1 and one of W groups may be absent and, if absent, are replaced with an additional bond between C_4 and C_5 in ring C;


L is independently a bond or linking group; and

RG is a reactive group.

- 20 2. The fluorescent dye of claim 1, wherein the linking group comprises a covalent linkage comprising 1-50 non-hydrogen atoms selected from C, N, O, S, P and halogen atoms and is composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and
- 25 wherein RG comprises a reactive group selected from a group consisting of a carboxylic acid, an activated ester of a carboxylic acid, acid anhydride, acid chloride, acyl azide, an acyl halide, aldehyde, chloroformate, amine, hydroxyl, hydrazine, isocyanate, isothiocyanate, sulfonyl halide, tosyl, maleimide, N-hydroxy-succinimide ester, aziridine, imine, and disulfide.

3. The fluorescent dye of claim 1 or 2, wherein the linking group comprises a formula:
 $R_1C(O)R_2$, wherein R_1 is a bond or C_{1-10} methylene $(CH_2)_n$ attached to the dye,
 $C(O)$ is a carbonyl group, and R_2 is OH or a bond attaching the carbonyl
 5 group to the reactive group RG; or
 $R_1C(O)A_1R_3C(O)R_2$, where R_1 is a bond or C_{1-10} methylene, $C(O)$ is a
 carbonyl group, A_1 is either NH, S or O, R_3 is an alkenyl $(CH_2)_n$, a five or a
 six membered ring having at least one unsaturated bond, or a combination of
 C_{1-10} methylene $(CH_2)_n$ and a five or six member ring, R_2 is OH or a bond
 10 attaching the terminal carbonyl group to the reactive group RG.
4. The fluorescent dye of any one of claims 1 to 3, wherein RG comprises a
 carboxylic acid, succinimidyl ester of a carboxylic acid, a hydrazide, an amine,
 an isothiocyanate, or a maleimide.


5. The fluorescent dye of claim 1, having the formula:




wherein

20 R represents OH, succinimidoxy, $NH(CH_2)_nCOOH$, $NH(CH_2)_nCO-$
 succinimide, or $NH(CH_2)_n$ -maleimide, wherein $n = 1-10$.

6. The fluorescent dye of claim 1, having the formula:




7. The fluorescent dye of claim 1, having the formula:




wherein

5 R represents OH, succinimidoxy, $\text{NH}(\text{CH}_2)_n\text{COOH}$, $\text{NH}(\text{CH}_2)_n\text{CO-succinimide}$, or $\text{NH}(\text{CH}_2)_n\text{-maleimide}$, wherein $n = 1-10$.


8. The fluorescent dye of claim 1, having the formula:



15 9. The fluorescent dye of claim 1, having the formula:




20 10. The fluorescent dye of claim 1, having the formula:



WO 2011/008912

PCT/US2010/042063

11. The fluorescent dye of claim 1, having the formula:




5

wherein

R represents OH, succinimidoxy, $\text{NH}(\text{CH}_2)_n\text{COOH}$, $\text{NH}(\text{CH}_2)_n\text{CO-succinimide}$, or $\text{NH}(\text{CH}_2)_n\text{-maleimide}$, wherein $n = 1-10$.

10

12. The fluorescent dye of claim 1 having the formula:



wherein

X_1 and X_2 are independently S or O;

Y_1 and Y_2 are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl, or

15

one of Y_1 or Y_2 is H, and the other is a haloalkyl group;

W is H or alkyl;

L is independently a bond or a linking group, wherein the linking group

comprises a covalent linkage comprising 1-50 non-hydrogen atoms selected

from C, N, O, S, P and halogen atoms and is composed of any combination

20

single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds,

carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and

wherein RG is a reactive group selected from a group consisting of a carboxylic

acid, an activated ester of a carboxylic acid, acid anhydride, acid chloride, acyl

azide, an acyl halide, aldehyde, chloroformate, amine, hydroxyl, hydrazine,

25

isocyanate, isothiocyanate, sulfonyl halide, tosyl, maleimide, N-hydroxy-


succinimide ester, aziridine, imine, and disulfide.

WO 2011/008912

PCT/US2010/042063

13. The fluorescent dye of claim 1, which can be excited with a wavelength from about 340 nm to about 450 nm.

14. A dye-conjugate having the formula:



wherein

X_1 and X_2 are independently S or O;

Y_1 and Y_2 separately represent H, halogen, alkyl, haloalkyl, alkoxy, or alkenyl wherein each W is independently H or alkyl;

Z_1 is H or alkyl;

wherein Z_1 and one of the W groups may be absent and, if absent, are replaced with an additional bond between C_4 and C_5 in ring C;

L is independently a bond or a linking group, wherein the linking group comprises a covalent linkage comprising 1-50 non-hydrogen atoms selected from C, N, O, S, P and halogen atoms and is composed of any combination single, double, triple, or aromatic carbon-carbon bonds, carbon-oxygen bonds, carbon-sulfur bonds, carbon-nitrogen bonds, and nitrogen-nitrogen bonds; and wherein CM is a conjugated molecule comprising a peptide, protein, polysaccharide, enzyme, lipid, nucleotide, oligonucleotide or nucleic acid polymer.

15. The dye-conjugate of claim 14, wherein the conjugated molecule is a peptide or a protein.

16. The dye-conjugate of claim 14 or 15, wherein the conjugated molecule is an antibody, or a fragment of an antibody.

17. A method of detecting a complementary member of a specific binding pair in a sample, comprising:

WO 2011/008912

PCT/US2010/042063

- a) mixing said sample with a dye-conjugate of any one of claims 14 to 16 which specifically binds to the complementary member; and
- b) detecting the complex formed by said mixture to detect the complementary member.

5

18. The method of claim 17, wherein the complementary member is a protein or peptide.

10

19. The method of claim 17 or 18, wherein the complementary member is present on a bacteria, virus or animal cell.

20. The method of any one of claims 17 to 19, wherein the complex is detected by its fluorescence response.

15

21. The method of claim 20, wherein the fluorescence response is detected using a flow cytometer, fluorometer, fluorescence microscope, or fluorescence plate reader.

20

22. The method of claim 20 or 21, further comprising distinguishing the fluorescence response from that of a second fluorophore having detectably different optical properties.

23. The method of any one of claims 20 to 22, further comprising separating said complex from other sample constituents based on the fluorescence response.


25

30

WO 2011/008912

PCT/US2010/042063

24. A compound having the formula:



5

wherein

Y_1 and Y_2 are independently halogen, alkyl, haloalkyl, alkoxy, or alkenyl

or

one of Y_1 or Y_2 is H, and the other is a haloalkyl group.

Application number / Numéro de demande: US2010042063

Figures: 4-5

Pages: _____

Unscannable items received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de la préparation
des dossiers au 10^{ième} étage)

WO 2011/008912

PCT/US2010/042063

1/5




FIG. 1

WO 2011/008912

PCT/US2010/042063

FIG 2A

2/5




FIG 2B




FIG 2C





FIG 2D



WO 2011/008912

PCT/US2010/042063

3/5

FIG 3A




FIG 3B

