

(19) **DANMARK**

(10) **DK 2013 70080 A1**



(12) **PATENTANSØGNING**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 K 16/28 (2006.01)** **A 61 K 39/395 (2006.01)** **A 61 K 41/00 (2006.01)**
- (21) Ansøgningsnummer: **PA 2013 70080**
- (22) Indleveringsdato: **2013-02-14**
- (24) Løbedag: **2011-07-15**
- (41) Alm. tilgængelig: **2013-02-14**
- (86) International ansøgning nr: **PCT/IB2011/002391**
- (86) International indleveringsdag: **2011-07-15**
- (30) Prioritet: **2010-07-15 RU 2010129292** **2010-07-15 RU 2010129291**
2010-07-15 RU 2010129290 **2011-03-17 RU 2011110106**
- (71) Ansøger: **Oleg Iliich Epshtein, 4 Samotyochny Per., d. 3, kv.72, Moscow 127473, Russiske Føderation**
- (72) Opfinder: **Oleg Iliich Epshtein, 4 Samotyochny Per., d. 3, kv.72, Moscow 127473, Russiske Føderation**
Svetlana Alexandrovna Sergeeva, Ul. Rossoshanskaya, d. 1, Korp. 1, Kv. 214, Moscow 117535, Russiske Føderation
Liudmila Fyodorovna Dolgovyh, Ul. Kurchatova, d. 16, Kv. 1, Chelyabinsk 454091, Russiske Føderation
Vladimir Ivanovich Petrov, Ul. Mira, d. 6, Kv. 46, Volgograd 400066, Russiske Føderation
- (74) Fuldmægtig: **Chas. Hude A/S, H.C. Andersens Boulevard 33, 1780 København V, Danmark**
- (54) Benævnelse: **A combination pharmaceutical composition and methods of treating diseases or conditions associated with the cardiovascular system**
- (57) Sammendrag:
The present application provides a pharmaceutical composition for administration to a patient suffering from at least one symptom of a cardiovascular condition, the 5 composition comprising a) an activated-potentiated form of an antibody to angiotensin II ATI receptor, and b) an activated-potentiated form of an antibody to endothelial NO synthase.

**A COMBINATION PHARMACEUTICAL COMPOSITION AND METHODS OF
TREATING DISEASES OR CONDITIONS ASSOCIATED WITH THE
CARDIOVASCULAR SYSTEM**

5

FIELD

The present invention relates to the field of medicine and can be used for the treatment and prevention of diseases of the cardiovascular system.

BACKGROUND

10

Nitric oxide (NO) is a cellular signaling molecule which plays a role in many biological processes including the relaxation of vascular and non-vascular tissue. Nitric oxide is synthesized from L-arginine by nitric oxide synthase (NO synthase). NO synthase occurs in different isoforms, including a constitutive form (cNOS) and an inducible form (iNOS). The constitutive form is present in normal endothelial cells, neurons and some other tissues. Formation of nitric oxide by the constitutive form in endothelial cells is thought to play an important role in normal blood pressure regulation.

15

Angiotensin is a protein that causes blood vessels to constrict, and drives blood pressure up and stimulates the release of aldosterone from the adrenal cortex. Angiotensin is part of the renin-angiotensin system (a hormone system that regulates blood pressure and water (fluid) balance) which is a major target for drugs that lower blood pressure. Angiotensin II receptor type 1 (AT1) is believed to mediate the key effects of angiotensin II.

20

25

The therapeutic effect of an extremely diluted (or ultra-low) form of antibodies potentized by homeopathic technology has been discovered by the inventor of the present patent application, Dr. Oleg I. Epshtein. U.S. Patent No. 7,582,294 discloses a medicament for treating Benign Prostatic Hyperplasia or prostatitis by administration of a homeopathically activated form of antibodies to prostate specific antigen (PSA). U. S. Patent Publication No. 2010/0260742 discloses a homeopathically potentized form of antibodies to a C-terminal fragment of the angiotensin II AT1 receptor. The

30

homeopathically potentized form of antibodies to a C-terminal fragment of the angiotensin II AT1 receptor is marketed in the Russian Federation and other countries under the name Kardos®. U.S. Patent No. 7,700,096 discloses and claims a homeopathically potentized form of antibodies to endothelial NO-synthase. The
5 homeopathically potentized form of antibodies to endothelial NO-synthase is marketed in the Russian Federation and other countries under the name Impaza®.

There is a continuing need for new drug products with desired therapeutic efficacy for treatment of diseases and disorder of the cardiovascular system.

10

SUMMARY

In one aspect the invention provides a pharmaceutical composition for administration to a patient suffering from at least one symptom of a cardiovascular condition, said composition comprising a) an activated-potentiated form of an antibody to angiotensin II AT1 receptor, and b) an activated-potentiated form of an antibody to
15 endothelial NO-synthase.

15

In one variant of this aspect of the invention a pharmaceutical composition for administration to a patient suffering from at least one symptom of a cardiovascular condition comprises a) an activated-potentiated form of an antibody to a C-terminal fragment of angiotensin II AT1 receptor, and b) an activated-potentiated form of an
20 antibody to endothelial NO-synthase.

20

Preferably, the pharmaceutical composition of this aspect is administered to patients with cardiovascular condition(s) associated with a reduced quality of life, wherein the administration of said pharmaceutical composition to said patient improves said quality of life of said patient.

25

Various variants and embodiments of the pharmaceutical composition are contemplated and provided. They may be used in reference to method aspects and embodiments of the invention. The specific variants and embodiments of this aspect of the invention are set forth in the appended claims. Preferably, the pharmaceutical composition of the invention comprises activated-potentiated forms of polyclonal
30 antibodies. It is also preferred that the process of preparing the pharmaceutical

30

composition of this aspect of the invention includes successive centesimal dilutions coupled with vertical shaking of every dilution.

In another aspect, the invention provides a method of treating a patient suffering from a reduced overall quality of life associated with at least one symptom of a cardiovascular condition, said method comprising administering to said patient a combination of a) an activated-potentiated form of an antibody to angiotensin II AT1 receptor, and b) an activated-potentiated form of an antibody to endothelial NO-synthase, thereby said administration improves said overall quality of life of said patient. Preferably, the invention provides a method of treating a patient suffering from a reduced overall quality of life associated with at least one symptom of a cardiovascular condition, said method comprising administering to said patient a combination of a) an activated-potentiated form of an antibody to a C-terminal fragment of angiotensin II AT1 receptor, and b) an activated-potentiated form of an antibody to endothelial NO-synthase, thereby said administration improves said overall quality of life of said patient. Various variants and embodiments are contemplated. In particular, it is contemplated that the combination of this aspect of the invention is administered concomitantly with administration of an additional therapeutic agent suitable for administration to patients suffering from said at least one symptom of a cardiovascular condition. The non-limiting list of suitable additional agents includes ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium antagonists, hypolipidemic agents, antiaggregants, antihypoxants, and anticoagulants. Bisoprol, enalapril or aspirin are specifically contemplated.

Particularly contemplated is a method of this aspect of the invention in which said patient is administered said composition in the form of a solid unit dosage form comprising said activated-potentiated form of an antibody to a C-terminal fragment of angiotensin II AT1 receptor and said activated-potentiated form of an antibody to endothelial NO-synthase. Preferably, said patient is administered one to two of said unit dosage forms, each administration carried out from once daily to four times daily.

In another aspect, the invention provides a method of treating a patient suffering from chronic heart failure, said method comprising administering to said patient a

combination of a) an activated-potentiated form of an antibody to a C-terminal fragment of angiotensin II AT1 receptor, and b) an activated-potentiated form of an antibody to endothelial NO-synthase. Various variants and embodiments are contemplated. In particular, it is contemplated that the combination of this aspect of the invention is administered concomitantly with administration of an additional therapeutic agent suitable for administration to patients suffering from said at least one symptom of a cardiovascular condition. The non-limiting list of suitable additional agents includes ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium antagonists, hypolipidemic agents, antiaggregants, antihypoxants, and anticoagulants. Bisoprol, enalapril or aspirin are specifically contemplated.

Particularly contemplated is a method of this aspect of the invention in which said patient is administered said composition in the form of a solid unit dosage form comprising said activated-potentiated form of an antibody to a C-terminal fragment of angiotensin II AT1 receptor and said activated-potentiated form of an antibody to endothelial NO-synthase. Preferably, said patient is administered one to two of said unit dosage forms, each administration carried out from once daily to four times daily. In accordance with this aspect of the invention, said patient exhibits statistically significant improvement in rigidity parameters of carotid radial artery segments upon said administration. In accordance with this aspect of the invention, said patient exhibits statistically significant improvement in rigidity parameters of carotid femoral artery segments upon said administration. In accordance with this aspect of the invention, said patient exhibits statistically significant reduction in anxiety associated with said chronic heart failure upon said administration. It is particularly contemplated that said administration of said combination leads to a statistically significant improvement in the The Minnesota Living with Heart Failure questionnaire (MLHFQ) score in a suitable population of said patients in reference to the baseline. It is particularly contemplated that said administration of said combination leads to a statistically significant reduction in the Kansas City Cardiomyopathy Questionnaire total Score in a suitable population of said patients in reference to the baseline. It is particularly contemplated that said administration of said combination leads to a statistically significant improvement in a 6-

minute walking test score in a suitable population of said patients. It is particularly contemplated that said administration of said combination leads to a statistically significant improvement in the Hospital Anxiety and Depression Scale (HADS) total score in a suitable population of said patients. It is particularly contemplated that said administration of said combination leads to said patient exhibiting statistically significant reduction in depression associated with said chronic heart failure upon said administration.

In another aspect, the invention provides a method of treating a patient suffering from asthenia and/or vegetative-vascular dystonia, said method comprising administering to said patient a combination of a) an activated-potentiated form of an antibody to a C-terminal fragment of angiotensin II AT1 receptor, and b) an activated-potentiated form of an antibody to endothelial NO-synthase. Various variants and embodiments are contemplated. In particular, it is contemplated that the combination of this aspect of the invention is administered concomitantly with administration of an additional therapeutic agent suitable for administration to patients suffering from said at least one symptom of a cardiovascular condition. The non-limiting list of suitable additional agents includes ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium antagonists, hypolipidemic agents, antiaggregants, antihypoxants, and anticoagulants. Bisoprol, enalapril or aspirin are specifically contemplated.

It is particularly contemplated in accordance with this aspect of the invention that said patient exhibits statistically significant improvement in rigidity parameters of carotid radial artery segments upon said administration. It is particularly contemplated that said patient exhibits statistically significant improvement in rigidity parameters of carotid femoral artery segments upon said administration. It is particularly contemplated that said patient exhibits statistically significant reduction in mental asthenia upon said administration. Preferably, said administration of said combination leads to a statistically significant reduction in the mental asthenia by the Multidimensional Fatigue Inventory (MFI-20) scale in a suitable population of said patients in reference to the baseline. Preferably, the administration of the combination drug or medicine in accordance with

this aspect of the invention leads to said patient exhibiting statistically significant reduction in general asthenia upon said administration. Preferably, said administration of said combination leads to a statistically significant reduction in the general asthenia by the MFI-20 scale in a suitable population of said patients in reference to the baseline.

5 Preferably, said patient exhibits statistically significant reduction in anxiety associated with said asthenia and/or vegetative vascular dystonia upon said administration. It is preferred that in accordance with this aspect of the invention, said administration of said combination drug or medicine leads to a statistically significant reduction in the trait anxiety as measured by the Spielberg test in a suitable population of said patients in
10 reference to the baseline. Preferably, said patient exhibits statistically significant reduction in depression associated with said asthenia and/or vegetative vascular dystonia upon said administration. It is particularly contemplated that said administration of said combination drug or medicine leads to a statistically significant reduction in depression as measured by the Beck test in a suitable population of said
15 patients in reference to the baseline. Preferably, said patient exhibits statistically significant improvement in brachial artery dilation level upon said administration.

In another aspect, the invention provides a method of treating hypertension comprising administering the combination described herein to a patient in need thereof. Concomitant administration of an additional therapeutic agent selected from the group
20 consisting of ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium antagonists, hypolipidemic agents, antiaggregants, antihypoxants, and anticoagulants is specifically contemplated.

DETAILED DESCRIPTION

25 The invention is defined with reference to the appended claims. With respect to the claims, the glossary that follows provides the relevant definitions.

The term "antibody" as used herein shall mean an immunoglobulin that specifically binds to, and is thereby defined as complementary with, a particular spatial and polar organization of another molecule. Antibodies as recited in the
30 claims may include a complete immunoglobulin or fragment thereof, may be natural,

polyclonal or monoclonal, and may include various classes and isotypes, such as IgA, IgD, IgE, IgG1, IgG2a, IgG2b and IgG3, IgM, etc. Fragments thereof may include Fab, Fv and F(ab')₂, Fab', and the like. The singular "antibody" includes plural "antibodies".

5 The term "activated-potentiated form" or "potentiated form", respectively, with respect to antibodies recited herein is used to denote a product of homeopathic potentization of any initial solution of antibodies. "Homeopathic potentization" denotes the use of methods of homeopathy to impart homeopathic potency to an initial solution of relevant substance. Although not so limited, 'homeopathic
10 potentization" may involve, for example, repeated consecutive dilutions combined with external treatment, particularly (mechanical) shaking. In other words, an initial solution of antibody is subjected to consecutive repeated dilution and multiple vertical shaking of each obtained solution in accordance with homeopathic technology. The preferred concentration of the initial solution of antibody in the
15 solvent, preferably, water or a water-ethyl alcohol mixture, ranges from about 0.5 to about 5.0 mg/ml. The preferred procedure for preparing each component, i.e. antibody solution, is the use of the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution (mother tincture) of antibodies diluted 100¹², 100³⁰ and 100²⁰⁰ times, respectively, which is equivalent to centesimal homeopathic
20 dilutions C12, C30 and C200. Examples of homeopathic potentization are described in U.S. Patent Nos. 7,572,441 and 7,582,294, which are incorporated herein by reference in their entirety and for the purpose stated. While the term "activated-potentiated form" is used in the claims, the term "ultra-low doses" is used in the examples. The term "ultra-low doses" became a term of art in the field of art created
25 by study and use of homeopathically diluted and potentized form of substance. The term "ultra-low dose" or "ultra-low doses" is meant as fully supportive and primarily synonymous with the term 'activated-potentiated" form used in the claims.

 In other words, an antibody is in the "activated-potentiated" form when three factors are present. First, the "activated-potentiated" form of the antibody is a
30 product of a preparation process well accepted in the homeopathic art. Second, the

“activated-potentiated” form of antibody must have biological activity determined by methods well accepted in modern pharmacology. And third, the biological activity exhibited by the “activated-potentiated” form of the antibody cannot be explained by the presence of the molecular form of the antibody in the final product of the homeopathic process.

For example, the activated potentiated form of antibodies may be prepared by subjecting an initial, isolated antibody in a molecular form to consecutive multiple dilutions coupled with an external impact, such as mechanical shaking. The external treatment in the course of concentration reduction may also be accomplished, for example, by exposure to ultrasonic, electromagnetic, or other physical factors. V. Schwabe "Homeopathic medicines", M., 1967, U.S. Patents Nos. 7,229,648 and 4,311,897, which are incorporated by reference in their entirety and for the purpose stated, describe such processes which are well accepted methods of homeopathic potentiation in the homeopathic art. This procedure gives rise to a uniform decrease in molecular concentration of the initial molecular form of the antibody. This procedure is repeated until the desired homeopathic potency is obtained. For the individual antibody, the required homeopathic potency can be determined by subjecting the intermediate dilutions to biological testing in the desired pharmacological model. Although not so limited, ‘homeopathic potentization’ may involve, for example, repeated consecutive dilutions combined with external treatment, particularly vertical (mechanical) shaking. In other words, an initial solution of antibody is subjected to consecutive repeated dilution and multiple vertical shaking of each obtained solution in accordance with homeopathic technology. The preferred concentration of the initial solution of antibody in the solvent, preferably, water or a water-ethyl alcohol mixture, ranges from about 0.5 to about 5.0 mg/ml. The preferred procedure for preparing each component, i.e. antibody solution, is the use of the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution (mother tincture) of antibodies diluted 100^{12} , 100^{30} and 100^{200} times, respectively, which is equivalent to centesimal homeopathic dilutions C12, C30 and C200 or the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution (mother tincture)

of antibodies diluted 100^{12} , 100^{30} and 100^{50} times, respectively, which is equivalent to centesimal homeopathic dilutions C12, C30 and C50. Examples of how to obtain the desired potency are also provided, for example, in U.S. Patents Nos. 7,229,648 and 4,311,897, which are incorporated by reference for the purpose stated. The procedure
5 applicable to the “activated potentiated” form of the antibodies described herein is described in more detail below.

There has been a considerable amount of controversy regarding homeopathic treatment of human subjects. While the present invention relies on accepted homeopathic processes to obtain the “activated-potentiated” form of antibodies, it does not rely solely on
10 homeopathy in human subjects for evidence of activity. It has been surprisingly discovered by the inventor of the present application and amply demonstrated in the accepted pharmacological models that the solvent ultimately obtained from consecutive multiple dilution of a starting molecular form of an antibody has definitive activity unrelated to the presence of the traces of the molecular form of the antibody in the target dilution. The “activated-
15 potentiated” form of the antibody provided herein are tested for biological activity in well accepted pharmacological models of activity, either in appropriate in vitro experiments, or in vivo in suitable animal models. The experiments provided further below provide evidence of biological activity in such models. The human clinical studies, also provided herein below, inter alia provide evidence that the activity observed in the animal model is well translated to
20 human therapy. The human study also provides evidence of availability of the “activated potentiated” forms described herein to treat specified human diseases or disorders well accepted as pathological conditions in the medical science.

Also, the claimed “activated-potentiated” form of antibody encompasses only solutions or solid preparations the biological activity of which cannot be explained by the presence of
25 the molecular form of the antibody remaining from the initial, starting solution. In other words, while it is contemplated that the “activated-potentiated” form of the antibody may contain traces of the initial molecular form of the antibody, one skilled in the art could not attribute the observed biological activity in the accepted pharmacological models to the remaining molecular form of the antibody with any degree of plausibility due to the extremely low
30 concentrations of the molecular form of the antibody remaining after the consecutive dilutions.

While the invention is not limited by any specific theory, the biological activity of the “activated-potentiated” form of the antibodies of the present invention is not attributable to the initial molecular form of the antibody. Preferred is the “activated-potentiated” form of antibody in liquid or solid form in which the concentration of the initial molecular form of the antibody is below the limit of detection of the accepted analytical techniques, such as capillary electrophoresis and High Performance Liquid Chromatography. Particularly preferred is the “activated-potentiated” form of antibody in liquid or solid form in which the concentration of the initial molecular form of the antibody is below the Avogadro number. In pharmacology of molecular forms of therapeutic substances, it is common practice to create a dose-response curve in which the level of pharmacological response is plotted against the concentration of the active drug administered to the subject or tested in vitro. The minimal level of the drug which produces any detectable response is known as a threshold dose. It is specifically contemplated and preferred that the “activated-potentiated” form of the antibodies contains molecular antibody, if any, at a concentration below the threshold dose for the molecular form of the antibody in the given biological model. The specific compositions described in the appended examples are described with the term ultra-low doses (ULD), which term is not intended to be used in construing the claims of the present application.

The term “cardiovascular condition” denotes disease or disorder of the cardiovascular system in human defined as such in the medical profession. Non-limiting examples of cardiovascular conditions include heart failure, arterial hypertension, and asthenia.

The term “symptom” with respect to “cardiovascular conditions” denotes those manifestations of patients suffering from diseases or disorders of the cardiovascular system that reduce quality of life of such patients. Non-limiting examples of symptoms of cardiovascular conditions that reduce quality of life include anxiety, depression, and walking difficulty.

The term “quality of life” denotes the subjective feeling of well-being of a patient. The quality of life of a patient increases with reduction in or improvement of symptoms. The term “quality of life” is meant to define the collection of symptoms associated with cardiovascular diseases or disorders in a patient population.

The term “MFI-20 scale” refers to “multidimensional fatigue inventory”, which is a 20-question inventory questionnaire used to measure the levels of physical, psychological and mental fatigue. It denotes the MFI-20 questionnaire commonly used by those skilled in the art. An example of use of the MFI-20 scale to assess fatigue in patients with chronic heart failure may be found in K. Falk et al., Fatigue and Anaemia in Patients with Chronic Heart Failure, European Journal of Heart Failure, 8, 744-749 (2006), which is incorporated herein by reference.

The term “The Minnesota Living with Heart Failure questionnaire” (MLHFQ), also referred to as the Minnesota questionnaire, refers to a well-established questionnaire commonly used to assess quality of life and levels of anxiety in people with cardiovascular conditions, and heart failure in particular.

The content of the Minnesota questionnaire is representative of the ways cardiovascular conditions, and particularly heart failure, together with appropriate treatments, affect the physical, emotional, social and mental dimensions of quality of life. The questionnaire focuses on questions that assess the impact of frequent physical symptoms - shortness of breath, fatigue, peripheral edema, and sleeping difficulty, and psychological symptoms of anxiety and depression. In addition, the effects of heart failure on physical/social functions including walking, climbing stairs, household work, need to rest, working to earn a living, going places away from home, doing things with family or friends, recreational activities, sexual activities, eating and mental and emotional functions of concentration, memory, loss of self-control, and being a burden to others were incorporated into the measure. Since treatments might have side effects in addition to ameliorating symptoms and functional limitations produced by heart failure, questions about side effects of medications, hospital stays and costs of care are included to help measure the overall impact of a treatment on quality of life. To measure the effects of symptoms, functional limitations, psychological distress on an individual’s quality of life, the questionnaire asks each person to indicate using a 6-point, zero to five, Likert scale how much each of 21 facets prevented them from living as they desired. This response format is consistent with the concept of quality of life and allows each individual to weigh each item using a common scale. Therefore, one can look at which items had the most effect and the sum of responses reflects the overall effects

of heart failure and treatments on the individual's quality of life. Although the Minnesota Living with Heart Failure Questionnaire incorporates relevant aspects of the key dimensions of quality of life, the questionnaire was not designed to measure any particular dimension separately. Given the conceptual basis for the questionnaire, items on the questionnaire are considered to be 'causal' indicators of quality of life in the sense that they can affect someone's quality of life when they occur, but may not be present when other aspects of heart failure are affecting an individual's quality of life. The total score is taken as the best measure of how cardiovascular condition, and particularly heart failure, and commensurate treatments impact the quality of life.

The term "Kansas City Cardiomyopathy Questionnaire" (KCCQ) or "Kansas Questionnaire" refers to a health-related quality-of-life measure for patients with congestive heart failure. It is a reliable, predictive tool that tracks how patients are doing if they have weakened heart muscle due to prior heart attacks, heart valve problems, viral infections, or other causes.

The answers patients give to the KCCQ's questions are used to calculate scores in ten scales:

1. **Physical Limitation:** a measure of how much a patient's condition is hampering his ability to do what he wants to do
2. **Symptom Stability:** a measure of whether a patient's symptoms are changing over time
3. **Symptom Frequency:** a measure of how often a patient has symptoms
4. **Symptom Burden:** a measure of what the impact of these symptoms are on the patient's well-being
5. **Total Symptom:** a combined measure of the symptom scales
6. **Social Limitation:** a measure of how much a patient's interpersonal relations are impacted by her condition
7. **Self-Efficacy:** a measure of how well a patient can manage her care, find answers and help
8. **Quality of Life:** a measure of the overall impact of a patient's condition on a patient's interpersonal relationships and state of mind

9. **Clinical Summary:** a combined measure of symptoms and social factors

10. **Overall Summary:** a combined measure of all the above

The term "Spielberg test" refers to the state-trait anxiety inventory for adults developed by Charles D. Spielberg which is well-accepted for measuring different forms of anxiety. The
5 sampler manuals for performing the Spielberg test are published by Mind Gardens, Inc.

The term "Beck Questionnaire Score" refers to a score obtained from the Beck Depression Inventory, a widely used method for measuring depression developed by Aaron Beck. The description of a modern Beck Questionnaire may be found for example in Beck
10 AT, Steer RA, Ball R, Ranieri W (December 1996). "*Comparison of Beck Depression Inventories -IA and -II in psychiatric outpatients*". *Journal of personality assessment* 67 (3): 588-97, incorporated herein by reference.

The term "in reference to the baseline" with respect to the score obtained from patient questionnaires denotes a comparison to the pre-treatment score of the same patient
15 population.

The present invention provides a pharmaceutical composition for
15 administration to a patient suffering from at least one symptom of a cardiovascular condition associated with a reduced quality of life of the patient, the composition comprising a) an activated-potentiated form of an antibody to the C-terminal fragment of angiotensin II AT1 receptor, and b) an activated-potentiated form of an
20 antibody to endothelial NO-synthase, wherein the administration of the pharmaceutical composition improves quality of life of the patient. As set forth herein above, each of the individual components of the combination is generally known for its own individual medical uses. However, the inventors of the present patent application surprisingly discovered that administration of the combination markedly
25 improves the quality of life of patients with cardiovascular conditions, such as heart failure, asthenia, dystonia, and hypertension.

The pharmaceutical composition in accordance with this aspect of the invention may be in the liquid form or in solid form. Each of the activated potentiated forms of the antibodies included in the pharmaceutical composition is prepared from
30 an initial molecular form of the antibody via a process accepted in homeopathic art.

The starting antibodies may be monoclonal, or polyclonal antibodies prepared in accordance with known processes, for example, as described in Immunotechniques, G. Frimel, M., "Meditsyna", 1987, p. 9-33; "Hum. Antibodies. Monoclonal and recombinant antibodies, 30 years after" by Laffly E., Sodoyer R. – 2005 – Vol. 14. – N 1-2. P.33-55, both incorporated herein by reference.

Monoclonal antibodies may be obtained, e.g., by means of hybridoma technology. The initial stage of the process includes immunization based on the principles already developed in the course of polyclonal antisera preparation. Further stages of work involve the production of hybrid cells generating clones of antibodies with identical specificity. Their separate isolation is performed using the same methods as in the case of polyclonal antisera preparation.

Polyclonal antibodies may be obtained via active immunization of animals. For this purpose, for example, suitable animals (e.g. rabbits) receive a series of injections of the appropriate antigen, either endothelial NO-synthase or angiotensin II AT1-receptor. The animals' immune system generates corresponding antibodies which are collected from the animals in a known manner. This procedure enables preparation of a monospecific antibody-rich serum. If desired, the serum containing antibodies may be purified, e.g. using affine chromatography, fractionation by salt precipitation, or ion-exchange chromatography. The resulting purified, antibody-enriched serum may be used as a starting material for preparation of the activated-potentiated form of the antibodies. The preferred concentration of the resulting initial solution of antibody in the solvent, preferably, water or a water-ethyl alcohol mixture, ranges from about 0.5 to about 5.0 mg/ml.

The preferred procedure for preparing each component of the combination drug according to the present invention is the use of the mixture of three aqueous-alcohol dilutions of the primary matrix solution of antibodies diluted 100^{12} , 100^{30} and 100^{200} times, respectively, which is equivalent to centesimal homeopathic dilutions C12, C30 and C200. To prepare a solid dosage form, a solid carrier is treated with the desired dilution obtained via the homeopathic process. To obtain a solid unit dosage form of the combination of the invention, the carrier mass is impregnated

with each of the dilutions. Both orders of impregnation are suitable to prepare the desired combination dosage form.

In a preferred embodiment, the starting material for the preparation of the activated potentiated form that comprise the combination of the invention is polyclonal, animal-raised antibody to the corresponding antigen, namely, the C-terminal fragment of angiotensin II AT1 receptor and endothelial NO- synthase. To obtain the activated-potentiated form of polyclonal antibodies to the C-terminal fragment of angiotensin II AT1 receptor, the desired antigen may be injected as immunogen into a laboratory animal, preferably, rabbits. The following sequences of the human angiotensin II AT1 receptor is specifically contemplated as suitable antigens. The use of a whole human angiotensin II AT1 receptor is specifically contemplated:

SEQ ID NO: 1

15	Ser	Pro	Pro	Ala	Gly	Thr	Arg	His	Met	Ala	Asn	Thr	Tyr	Pro	Glu
	1				5					10					15
	Ala	Asn	Gly	Ile	Thr	Glu	Asn	Ser	Ile	Asn	Ile	Ile	Arg	Glu	Cys
	16				20					25					30
	Glu	Pro	Thr	Arg	Ser	His	Met	Ser	Ala	Pro	Ile	Glu	Asn	Ser	Gly
20	31				35					40					45
	Asn	Ala	Gly	Thr	Arg	Pro	Glu	Ser	Val	Met	Ile	Leu	Asn	Ser	Ser
	46				50					55					60
	Thr	Glu	Asp	Gly	Ile	Lys	Arg	Ile	Gln	Asp	Asp	Cys	Pro	Lys	Ala
	61				65					70					75
25	Gly	Arg	His	Asn	Tyr	Ile	Phe	Val	Met	Ile	Pro	Thr	Leu	Tyr	Ser
	76				80					85					90
	Ile	Ile	Phe	Val	Val	Gly	Ile	Phe	Gly	Asn	Ser	Leu	Val	Val	Ile
	91				95					100					105
	Val	Ile	Tyr	Phe	Tyr	Met	Lys	Leu	Lys	Thr	Val	Ala	Ser	Val	Phe
30	106				110					115					120
	Leu	Leu	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Cys	Phe	Leu	Leu	Thr	Leu
	121				125					130					135
	Pro	Leu	Trp	Ala	Val	Tyr	Thr	Ala	Met	Glu	Tyr	Arg	Trp	Pro	Phe
	136				140					145					150
35	Gly	Asn	Tyr	Leu	Cys	Lys	Ile	Ala	Ser	Ala	Ser	Val	Ser	Phe	Asn
	151				155					160					165
	Leu	Tyr	Ala	Ser	Val	Phe	Leu	Leu	Thr	Cys	Leu	Ser	Ile	Asp	Arg
	166				170					175					180

	Tyr	Leu	Ala	Ile	Val	His	Pro	Met	Lys	Ser	Arg	Leu	Arg	Arg	Thr
	181				185					190					195
	Met	Leu	Val	Ala	Lys	Val	Thr	Cys	Ile	Ile	Ile	Trp	Leu	Leu	Ala
	196				200					205					210
5	Gly	Leu	Ala	Ser	Leu	Pro	Ala	Ile	Ile	His	Arg	Asn	Val	Phe	Phe
	211				215					220					225
	Ile	Glu	Asn	Thr	Asn	Ile	Thr	Val	Cys	Ala	Phe	His	Tyr	Glu	Ser
	226				230					235					240
	Gln	Asn	Ser	Thr	Leu	Pro	Ile	Gly	Leu	Gly	Leu	Thr	Lys	Asn	Ile
10	241				245					250					255
	Leu	Gly	Phe	Leu	Phe	Pro	Phe	Leu	Ile	Ile	Leu	Thr	Ser	Tyr	Thr
	256				260					265					270
	Leu	Ile	Trp	Lys	Ala	Leu	Lys	Lys	Ala	Tyr	Glu	Ile	Gln	Lys	Asn
	271				275					280					285
15	Lys	Pro	Arg	Asn	Asp	Asp	Ile	Phe	Lys	Ile	Ile	Met	Ala	Ile	Val
	286				290					295					300
	Leu	Phe	Phe	Phe	Phe	Ser	Trp	Ile	Pro	His	Gln	Ile	Phe	Thr	Phe
	301				305					310					315
	Leu	Asp	Val	Leu	Ile	Gln	Leu	Gly	Ile	Ile	Arg	Asp	Cys	Arg	Ile
20	316				320					325					330
	Ala	Asp	Ile	Val	Asp	Thr	Ala	Met	Pro	Ile	Thr	Ile	Cys	Ile	Ala
	331				335					340					345
	Tyr	Phe	Asn	Asn	Cys	Leu	Asn	Pro	Leu	Phe	Tyr	Gly	Phe	Leu	Gly
	346				350					355					360
25	Lys	Lys	Phe	Lys	Arg	Tyr	Phe	Leu	Gln	Leu	Leu	Lys	Tyr	Ile	Pro
	361				365					370					375
	Pro	Lys	Ala	Lys	Ser	His	Ser	Asn	Leu	Ser	Thr	Lys	Met	Ser	Thr
	376				380					385					390
	Leu	Ser	Tyr	Arg	Pro	Ser	Asp	Asn	Val	Ser	Ser	Ser	Thr	Lys	Lys
30	391				395					400					405
	Pro	Ala	Pro	Cys	Phe	Glu	Val	Glu							
	406				410					413					

The use of different fragments of C-terminal fragment of human angiotensin II AT1 receptor with as antigen is also contemplated. The suitable sequences for such antigen are as follow:

SEQ ID NO: 2

															Gly
40															360
	Lys	Lys	Phe	Lys	Arg	Tyr	Phe	Leu	Gln	Leu	Leu	Lys	Tyr	Ile	Pro
	361				365					370					375
	Pro	Lys	Ala	Lys	Ser	His	Ser	Asn	Leu	Ser	Thr	Lys	Met	Ser	Thr
	376				380					385					390

Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys
 391 395 400 405
 Pro Ala Pro Cys Phe Glu Val Glu
 406 410 413

5

SEQ ID NO: 3

Gln Leu Leu Lys Tyr Ile Pro
 369 370 375

Pro Lys Ala
 376 378

10

SEQ ID NO: 4

Ser Asn Leu Ser Thr Lys Met Ser Thr
 382 385 390

Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys
 391 395 400 405
 Pro Ala Pro Cys Phe
 406 410

15

20

The use of the fragments of C-terminal fragment of human angiotensin II AT1 receptor with N-terminal cysteine (Cys) as antigen is also contemplated. The suitable sequence for such antigen is as follow:

SEQ ID NO: 5

25

Cys Gly
 360 365

Lys Lys Phe Lys Arg Tyr Phe Leu Gln Leu Leu Lys Tyr Ile Pro
 361 365 370 375

30

Pro Lys Ala Lys Ser His Ser Asn Leu Ser Thr Lys Met Ser Thr
 376 380 385 390

Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys
 391 395 400 405

35

Pro Ala Pro Cys Phe Glu Val Glu
 406 410 413

SEQ ID NO: 6

Cys Gln Leu Leu Lys Tyr Ile Pro
 369 370 375

40

Pro Lys Ala
 376 378

SEQ ID NO: 7

					Cys	Ser	Asn	Leu	Ser	Thr	Lys	Met	Ser	Thr	
						382			385					390	
5	Leu	Ser	Tyr	Arg	Pro	Ser	Asp	Asn	Val	Ser	Ser	Ser	Thr	Lys	Lys
	391				395					400					405
	Pro	Ala	Pro	Cys	Phe										
	406				410										

The exemplary procedure for preparation of the starting polyclonal antibodies to C-terminal fragment of human angiotensin II AT1 receptor may be described as follows. In 7-9 days before blood sampling, 1-3 intravenous injections of the desired antigen are made to the rabbits to increase the level of polyclonal antibodies in the rabbit blood stream. Upon immunization, blood samples are taken to test the antibody level. Typically, the maximum level of immune reaction of the soluble antigen is achieved within 40 to 60 days after the first injection of the antigen. Upon completion of the first immunization cycle, rabbits have a 30-day rehabilitation period, after which re-immunization is performed with another 1-3 intravenous injections.

To obtain antiserum containing the desired antibodies, the immunized rabbits' blood is collected from rabbits and placed in a 50ml centrifuge tube. Product clots formed on the tube sides are removed with a wooden spatula, and a rod is placed into the clot in the tube center. The blood is then placed in a refrigerator for one night at the temperature of about 4°C. On the following day, the clot on the spatula is removed, and the remaining liquid is centrifuged for 10 min at 13,000 rpm. Supernatant fluid is the target antiserum. The obtained antiserum is typically yellow. 20% of NaN₃ (weight concentration) is added in the antiserum to a final concentration of 0.02% and stored before use in frozen state at the temperature of -20°C or without NaN₃ at the temperature of -70°C. To separate the target antibodies to the C-terminal fragment of angiotensin II AT1 receptor from the antiserum, the following solid phase absorption sequence is suitable:

10 ml of the antiserum of rabbits is diluted twofold with 0.15 M NaCl, after which 6.26g Na₂SO₄ is added, mixed and incubated for 12-16 hours at 4°C. The sediment is removed by centrifugation, diluted in 10ml of phosphate buffer and

dialyzed against the same buffer during one night at ambient temperature. After the sediment is removed, the solution is applied to a DEAE-cellulose column balanced by phosphate buffer. The antibody fraction is determined by measuring the optical density of eluate at 280 nm.

5 The isolated crude antibodies are purified using the affine chromatography method by attaching the obtained antibodies to a C-terminal fragment of angiotensin II AT1 receptor located on the insoluble matrix of the chromatography media, with subsequent elution by concentrated aqueous salt solutions.

10 The resulting buffer solution is used as the initial solution for the homeopathic dilution process used to prepare the activated potentiated form of the antibodies. The preferred concentration of the initial matrix solution of the antigen-purified polyclonal rabbit antibodies to a C-terminal fragment of angiotensin II AT1 receptor is 0.5 - 5.0 mg/ml, preferably, 2.0 - 3.0 mg/ml.

15 The polyclonal antibodies to endothelial NO-synthase are obtained by a similar methodology using an adjuvant. Preferably, the entire molecule of bovine endothelial NO-synthase is used as immunogen (antigen) for rabbits' immunization:

SEQ.ID. NO. 8

	Met	Gly	Asn	Leu	Lys	Ser	Val	Gly	Gln	Glu	Pro	Gly	Pro	Pro	Cys
	1				5					10					15
20	Gly	Leu	Gly	Leu	Gly	Leu	Gly	Leu	Gly	Leu	Cys	Gly	Lys	Gln	Gly
	16				20					25					30
	Pro	Ala	Ser	Pro	Ala	Pro	Glu	Pro	Ser	Arg	Ala	Pro	Ala	Pro	Ala
	31				35					40					45
25	Thr	Pro	His	Ala	Pro	Asp	His	Ser	Pro	Ala	Pro	Asn	Ser	Pro	Thr
	46				50					55					60
	Leu	Thr	Arg	Pro	Pro	Glu	Gly	Pro	Lys	Phe	Pro	Arg	Val	Lys	Asn
	61				65					70					75
	Trp	Glu	Leu	GLys	er	Ile	Thr	Tyr	Asp	Thr	Leu	Cys	Ala	Gln	Ser
	76				80					85					90
30	Gln	Gln	Asp	Gly	Pro	Cys	Thr	Pro	Arg	Cys	Cys	Leu	GLys	er	Leu
	91				95					100					105
	Val	Leu	Pro	Arg	Lys	Leu	Gln	Thr	Arg	Pro	Ser	Pro	Gly	Pro	Pro
	106				110					115					120
	Pro	Ala	Glu	Gln	Leu	Leu	Ser	Gln	Ala	Arg	Asp	Phe	Ile	Asn	Gln
35	121				125					130					135
	Tyr	Tyr	Ser	Ser	Ile	Lys	Arg	Ser	GLys	er	Gln	Ala	His	Glu	Glu
	136				140					145					150

	Arg	Leu	Gln	Glu	Val	Glu	Ala	Glu	Val	Ala	Ser	Thr	Gly	Thr	Tyr
	151				155					160					165
	His	Leu	Arg	Glu	Ser	Glu	Leu	Val	Phe	Gly	Ala	Lys	Gln	Ala	Trp
	166				170					175					180
5	Arg	Asn	Ala	Pro	Arg	Cys	Val	Gly	Arg	Ile	Gln	Trp	Gly	Lys	Leu
	181				185					190					195
	Gln	Val	Phe	Asp	Ala	Arg	Asp	Cys	Ser	Ser	Ala	Gln	Glu	Met	Phe
	196				200					205					210
	Thr	Tyr	Ile	Cys	Asn	His	Ile	Lys	Tyr	Ala	Thr	Asn	Arg	Gly	Asn
10	211				215					220					225
	Leu	Arg	Ser	Ala	Ile	Thr	Val	Phe	Pro	Gln	Arg	Ala	Pro	Gly	Arg
	226				230					235					240
	Gly	Asp	Phe	Arg	Ile	Trp	Asn	Ser	Gln	Leu	Val	Arg	Tyr	Ala	Gly
	241				245					250					255
15	Tyr	Arg	Gln	Gln	Asp	Gly	Ser	Val	Arg	Gly	Asp	Pro	Ala	Asn	Val
	256				260					265					270
	Glu	Ile	Thr	Glu	Leu	Cys	Ile	Gln	His	Gly	Trp	Thr	Pro	Gly	Asn
	271				275					280					285
	Gly	Arg	Phe	Asp	Val	Leu	Pro	Leu	Leu	Leu	Gln	Ala	Pro	Asp	Glu
20	286				290					295					300
	Ala	Pro	Glu	Leu	Phe	Val	Leu	Pro	Pro	Glu	Leu	Val	Leu	Glu	Val
	301				305					310					315
	Pro	Leu	Glu	His	Pro	Thr	Leu	Glu	Trp	Phe	Ala	Ala	Leu	Gly	Leu
	316				320					325					330
25	Arg	Trp	Tyr	Ala	Leu	Pro	Ala	Val	Ser	Asn	Met	Leu	Leu	Glu	Ile
	331				335					340					345
	Gly	Gly	Leu	Glu	Phe	Ser	Ala	Ala	Pro	Phe	Ser	Gly	Trp	Tyr	Met
	346				350					355					360
	Ser	Thr	Glu	Ile	Gly	Thr	Arg	Asn	Leu	Cys	Asp	Pro	His	Arg	Tyr
30	361				365					370					375
	Asn	Ile	Leu	Glu	Asp	Val	Ala	Val	Cys	Met	Asp	Leu	Asp	Thr	Arg
	376				380					385					390
	Thr	Thr	Ser	Ser	Leu	Trp	Lys	Asp	Lys	Ala	Ala	Val	Glu	Ile	Asn
	391				395					400					405
35	Leu	Ala	Val	Leu	His	Ser	Phe	Gln	Leu	Ala	Lys	Val	Thr	Ile	Val
	406				410					415					420
	Asp	His	His	Ala	Ala	Thr	Val	Ser	Phe	Met	Lys	His	Leu	Asp	Asn
	421				425					430					435
	Glu	Gln	Lys	Ala	Arg	Gly	Gly	Cys	Pro	Ala	Asp	Trp	Ala	Trp	Ile
40	436				440					445					450
	Val	Pro	Pro	Ile	Ser	Gly	Ser	Leu	Thr	Pro	Val	Phe	His	Gln	Glu
	451				455					460					465
	Met	Val	Asn	Tyr	Ile	Leu	Ser	Pro	Ala	Phe	Arg	Tyr	Gln	Pro	Asp
	466				470					475					480
45	Pro	Trp	Lys	Gly	Ser	Ala	Thr	Lys	Gly	Ala	Gly	Ile	Thr	Arg	Lys
	481				485					490					495

	Lys	Thr	Phe	Lys	Glu	Val	Ala	Asn	Ala	Val	Lys	Ile	Ser	Ala	Ser
	496				500					505					510
	Leu	Met	Gly	Thr	Leu	Met	Ala	Lys	Arg	Val	Lys	Ala	Thr	Ile	Leu
	511				515					510					525
5	Tyr	Ala	Ser	Glu	Thr	Gly	Arg	Ala	Gln	Ser	Tyr	Ala	Gln	Gln	Leu
	526				530					535					540
	Gly	Arg	Leu	Phe	Arg	Lys	Ala	Phe	Asp	Pro	Arg	Val	Leu	Cys	Met
	541				545					550					555
	Asp	Glu	Tyr	Asp	Val	Val	Ser	Leu	Glu	His	Glu	Ala	Leu	Val	Leu
10	556				560					565					570
	Val	Val	Thr	Ser	Thr	Phe	Gly	Asn	Gly	Asp	Pro	Pro	Glu	Asn	Gly
	571				575					580					585
	Glu	Ser	Phe	Ala	Ala	Ala	Leu	Met	Glu	Met	Ser	Gly	Pro	Tyr	Asn
	586				590					595					600
15	Ser	Ser	Pro	Arg	Pro	Glu	Gln	His	Lys	Ser	Tyr	Lys	Ile	Arg	Phe
	601				605					610					615
	Asn	Ser	Val	Ser	Cys	Ser	Asp	Pro	Leu	Val	Ser	Ser	Trp	Arg	Arg
	616				620					625					630
	Lys	Arg	Lys	Glu	Ser	Ser	Asn	Thr	Asp	Ser	Ala	Gly	Ala	Leu	Gly
20	631				635					640					645
	Thr	Leu	Arg	Phe	Cys	Val	Phe	Gly	Leu	Gly	Ser	Arg	Ala	Tyr	Pro
	646				650					655					660
	His	Phe	Cys	Ala	Phe	Ala	Arg	Ala	Val	Asp	Thr	Arg	Leu	Glu	Glu
	661				665					670					675
25	Leu	Gly	Gly	Glu	Arg	Leu	Leu	Gln	Leu	Gly	Gln	Gly	Asp	Glu	Leu
	676				680					685					690
	Cys	Gly	Gln	Glu	Glu	Ala	Phe	Arg	Gly	Trp	Ala	Lys	Ala	Ala	Phe
	691				695					700					705
	Gln	Ala	Ser	Cys	Glu	Thr	Phe	Cys	Val	Gly	Glu	Glu	Ala	Lys	Ala
30	706				710					715					720
	Ala	Ala	Gln	Asp	Ile	Phe	Ser	Pro	Lys	Arg	Ser	Trp	Lys	Arg	Gln
	721				725					730					735
	Arg	Tyr	Arg	Leu	Ser	Thr	Gln	Ala	Glu	Gly	Leu	Gln	Leu	Leu	Pro
	736				740					745					750
35	Gly	Leu	Ile	His	Val	His	Arg	Arg	Lys	Met	Phe	Gln	Ala	Thr	Val
	751				755					760					765
	Leu	Ser	Val	Glu	Asn	Leu	Gln	Ser	Ser	Lys	Ser	Thr	Arg	Ala	Thr
	766				770					775					780
	Ile	Leu	Val	Arg	Leu	Asp	Thr	Ala	Gly	Gln	Glu	Gly	Leu	Gln	Tyr
40	781				785					790					795
	Gln	Pro	Gly	Asp	His	Ile	Gly	Ile	Cys	Pro	Pro	Asn	Arg	Pro	Gly
	796				800					805					810
	Leu	Val	Glu	Ala	Leu	Leu	Ser	Arg	Val	Glu	Asp	Pro	Pro	Pro	Pro
	811				815					820					825
45	Thr	Glu	Ser	Val	Ala	Val	Glu	Gln	Leu	Glu	Lys	Gly	er	Pro	Gly
	826				830					835					840

	Gly	Pro	Pro	Pro	Ser	Trp	Val	Arg	Asp	Pro	Arg	Leu	Pro	Pro	Cys
	841				845					850					855
	Thr	Leu	Arg	Gln	Ala	Leu	Thr	Phe	Phe	Leu	Asp	Ile	Thr	Ser	Pro
	856				860					865					870
5	Pro	Ser	Pro	Arg	Leu	Leu	Arg	Leu	Leu	Ser	Thr	Leu	Ala	Glu	Glu
	871				875					880					885
	Pro	Ser	Glu	Gln	Gln	Glu	Leu	Glu	Thr	Leu	Ser	Gln	Asp	Pro	Arg
	886				890					895					900
10	Arg	Tyr	Glu	Glu	Trp	Lys	Trp	Phe	Arg	Cys	Pro	Thr	Leu	Leu	Glu
	901				905					910					915
	Val	Leu	Glu	Gln	Phe	Pro	Ser	Val	Ala	Leu	Pro	Ala	Pro	Leu	Leu
	916				920					925					930
	Leu	Thr	Gln	Leu	Pro	Leu	Leu	Gln	Pro	Arg	Tyr	Tyr	Ser	Val	Ser
	931				935					940					945
15	Ser	Ala	Pro	Asn	Ala	His	Pro	Gly	Glu	Val	His	Leu	Thr	Val	Ala
	946				950					955					960
	Val	Leu	Ala	Tyr	Arg	Thr	Gln	Asp	Gly	Leu	Gly	Pro	Leu	His	Tyr
	961				965					970					975
20	Gly	Val	Cys	Ser	Thr	Trp	Leu	Ser	Gln	Leu	Lys	Thr	Gly	Asp	Pro
	976				980					985					990
	Val	Pro	Cys	Phe	Ile	Arg	Gly	Ala	Pro	Ser	Phe	Arg	Leu	Pro	Pro
	991				995					1000					1005
	Asp	Pro	Tyr	Val	Pro	Cys	Ile	Leu	Val	Gly	Pro	Gly	Thr	Gly	Ile
	1006				1010					1015					1020
25	Ala	Pro	Phe	Arg	Gly	Phe	Trp	Gln	Glu	Arg	Leu	His	Asp	Ile	Glu
	1021				1025					1030					1035
	Ser	Lys	Gly	Leu	Gln	Pro	Ala	Pro	Met	Thr	Leu	Val	Phe	Gly	Cys
	1036				1040					1045					1050
30	Arg	Cys	Ser	Gln	Leu	Asp	His	Leu	Tyr	Arg	Asp	Glu	Val	Gln	Asp
	1051				1055					1060					1065
	Ala	Gln	Glu	Arg	Gly	Val	Phe	Gly	Arg	Val	Leu	Thr	Ala	Phe	Ser
	1066				1070					1075					1080
	Arg	Glu	Pro	Asp	Ser	Pro	Lys	Thr	Tyr	Val	Gln	Asp	Ile	Leu	Arg
	1081				1085					1090					1095
35	Thr	Glu	Leu	Ala	Ala	Glu	Val	His	Arg	Val	Leu	Cys	Leu	Glu	Arg
	1096				1100					1105					1110
	Gly	His	Met	Phe	Val	Cys	Gly	Asp	Val	Thr	Met	Ala	Thr	Ser	Val
	1111				1115					1120					1125
40	Leu	Gln	Thr	Val	Gln	Arg	Ile	Leu	Ala	Thr	Glu	Gly	Asp	Met	Glu
	1126				1130					1135					1140
	Leu	Asp	Glu	Ala	Gly	Asp	Val	Ile	Gly	Val	Leu	Arg	Asp	Gln	Gln
	1141				1145					1150					1155
	Arg	Tyr	His	Glu	Asp	Ile	Phe	Gly	Leu	Thr	Leu	Arg	Thr	Gln	Glu
	1156				1160					1165					1170
45	Val	Thr	Ser	Arg	Ile	Arg	Thr	Gln	Ser	Phe	Ser	Leu	Gln	Glu	Arg
	1171				1175					1180					1185

His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp Pro Pro Gly Pro
 1186 1190 1195 1200
 Asp Thr Pro Gly Pro
 1201 1205

5

Polyclonal antibodies to endothelial NO synthase may be obtained using the whole molecule of human endothelial NO synthase of the following sequence:

SEQ ID NO: 9

10 Met Gly Asn Leu Lys Ser Val Ala Gln Glu Pro Gly Pro Pro Cys
 1 5 10 15
 Gly Leu Gly Leu Gly Leu Gly Leu Gly Leu Cys Gly Lys Gln Gly
 16 20 25 30
 Pro Ala Thr Pro Ala Pro Glu Pro Ser Arg Ala Pro Ala Ser Leu
 31 35 40 45
 15 Leu Pro Pro Ala Pro Glu His Ser Pro Pro Ser Ser Pro Leu Thr
 46 50 55 60
 Gln Pro Pro Glu Gly Pro Lys Phe Pro Arg Val Lys Asn Trp Glu
 61 65 70 75
 20 Val GLys er Ile Thr Tyr Asp Thr Leu Ser Ala Gln Ala Gln Gln
 76 80 85 90
 Asp Gly Pro Cys Thr Pro Arg Arg Cys Leu GLys er Leu Val Phe
 91 95 100 105
 Pro Arg Lys Leu Gln Gly Arg Pro Ser Pro Gly Pro Pro Ala Pro
 106 110 115 120
 25 Glu Gln Leu Leu Ser Gln Ala Arg Asp Phe Ile Asn Gln Tyr Tyr
 121 125 130 135
 Ser Ser Ile Lys Arg Ser GLys er Gln Ala His Glu Gln Arg Leu
 136 140 145 150
 Gln Glu Val Glu Ala Glu Val Ala Ala Thr Gly Thr Tyr Gln Leu
 30 151 155 160 165
 Arg Glu Ser Glu Leu Val Phe Gly Ala Lys Gln Ala Trp Arg Asn
 166 170 175 180
 Ala Pro Arg Cys Val Gly Arg Ile Gln Trp Gly Lys Leu Gln Val
 181 185 190 195
 35 Phe Asp Ala Arg Asp Cys Arg Ser Ala Gln Glu Met Phe Thr Tyr
 196 200 205 210
 Ile Cys Asn His Ile Lys Tyr Ala Thr Asn Arg Gly Asn Leu Arg
 211 215 220 225
 Ser Ala Ile Thr Val Phe Pro Gln Arg Cys Pro Gly Arg Gly Asp
 40 226 230 235 240
 Phe Arg Ile Trp Asn Ser Gln Leu Val Arg Tyr Ala Gly Tyr Arg
 241 245 250 255
 Gln Gln Asp GLy Ser Val Arg Gly Asp Pro Ala Asn Val Glu Ile

	256				260					265				270	
	Thr	Glu	Leu	Cys	Ile	Gln	His	Gly	Trp	Thr	Pro	Gly	Asn	Gly	Arg
	271				275					280				285	
5	Phe	Asp	Val	Leu	Pro	Leu	Leu	Leu	Gln	Ala	Pro	Asp	Glu	Pro	Pro
	286				290					295				300	
	Glu	Leu	Phe	Leu	Leu	Pro	Pro	Glu	Leu	Val	Leu	Glu	Val	Pro	Leu
	301				305					310				315	
	Glu	His	Pro	Thr	Leu	Glu	Trp	Phe	Ala	Ala	Leu	Gly	Leu	Arg	Trp
	316				320					325				330	
10	Tyr	Ala	Leu	Pro	Ala	Val	Ser	Asn	Met	Leu	Leu	Glu	Ile	Gly	Gly
	331				335					340				345	
	Leu	Glu	Phe	Pro	Ala	Ala	Pro	Phe	Ser	Gly	Trp	Tyr	Met	Ser	Thr
	346				350					355				360	
15	Glu	Ile	Gly	Thr	Arg	Asn	Leu	Cys	Asp	Pro	His	Arg	Tyr	Asn	Ile
	361				365					370				375	
	Leu	Glu	Asp	Val	Ala	Val	Cys	Met	Asp	Leu	Asp	Thr	Arg	Thr	Thr
	376				380					385				390	
	Ser	Ser	Leu	Trp	Lys	Asp	Lys	Ala	Ala	Val	Glu	Ile	Asn	Val	Ala
	391				395					400				405	
20	Val	Leu	His	Ser	Tyr	Gln	Leu	Ala	Lys	Val	Thr	Ile	Val	Asp	His
	406				410					415				420	
	His	Ala	Ala	Thr	Ala	Ser	Phe	Met	Lys	His	Leu	Glu	Asn	Glu	Gln
	421				425					430				435	
25	Lys	Ala	Arg	Gly	Gly	Cys	Pro	Ala	Asp	Trp	Ala	Trp	Ile	Val	Pro
	436				440					445				450	
	Pro	Ile	Ser	GLys	er	Leu	Thr	Pro	Val	Phe	His	Gln	Glu	Met	Val
	451				455					460				465	
	Asn	Tyr	Phe	Leu	Ser	Pro	Ala	Phe	Arg	Tyr	Gln	Pro	Asp	Pro	Trp
	466				470					475				480	
30	Lys	GLys	er	Ala	Ala	Lys	Gly	Thr	Gly	Ile	Thr	Arg	Lys	Lys	Thr
	481				485					490				495	
	Phe	Lys	Glu	Val	Ala	Asn	Ala	Val	Lys	Ile	Ser	Ala	Ser	Leu	Met
	496				500					505				510	
35	Gly	Thr	Val	Met	Ala	Lys	Arg	Val	Lys	Ala	Thr	Ile	Leu	Tyr	Gly
	511				515					510				525	
	Ser	Glu	Thr	Gly	Arg	Ala	Gln	Ser	Tyr	Ala	Gln	Gln	Leu	Gly	Arg
	526				530					535				540	
	Leu	Phe	Arg	Lys	Ala	Phe	Asp	Pro	Arg	Val	Leu	Cys	Met	Asp	Glu
	541				545					550				555	
40	Tyr	Asp	Val	Val	Ser	Leu	Glu	His	Glu	Thr	Leu	Val	Leu	Val	Val
	556				560					565				570	
	Thr	Ser	Thr	Phe	Gly	Asn	Gly	Asp	Pro	Pro	Glu	Asn	Gly	Glu	Ser
	571				575					580				585	
45	Phe	Ala	Ala	Ala	Leu	Met	Glu	Met	Ser	Gly	Pro	Tyr	Asn	Ser	Ser
	586				590					595				600	
	Pro	Arg	Pro	Glu	Gln	His	Lys	Ser	Tyr	Lys	Ile	Arg	Phe	Asn	Ser

	601				605					610				615	
	Ile	Ser	Cys	Ser	Asp	Pro	Leu	Val	Ser	Ser	Trp	Arg	Arg	Lys	Arg
	616				620					625				630	
	Lys	Glu	Ser	Ser	Asn	Thr	Asp	Ser	Ala	Gly	Ala	Leu	Gly	Thr	Leu
5	631				635					640				645	
	Arg	Phe	Cys	Val	Phe	Gly	Leu	Gly	er	Arg	Ala	Tyr	Pro	His	Phe
	646				650					655				660	
	Cys	Ala	Phe	Ala	Arg	Ala	Val	Asp	Thr	Arg	Leu	Glu	Glu	Leu	Gly
	661				665					670				675	
10	Gly	Glu	Arg	Leu	Leu	Gln	Leu	Gly	Gln	Gly	Asp	Glu	Leu	Cys	Gly
	676				680					685				690	
	Gln	Glu	Glu	Ala	Phe	Arg	Gly	Trp	Ala	Gln	Ala	Ala	Phe	Gln	Ala
	691				695					700				705	
	Ala	Cys	Glu	Thr	Phe	Cys	Val	Gly	Glu	Asp	Ala	Lys	Ala	Ala	Ala
15	706				710					715				720	
	Arg	Asp	Ile	Phe	Ser	Pro	Lys	Arg	Ser	Trp	Lys	Arg	Gln	Arg	Tyr
	721				725					730				735	
	Arg	Leu	Ser	Ala	Gln	Ala	Glu	Gly	Leu	Gln	Leu	Leu	Pro	Gly	Leu
	736				740					745				750	
20	Ile	His	Val	His	Arg	Arg	Lys	Met	Phe	Gln	Ala	Thr	Ile	Arg	Ser
	751				755					760				765	
	Val	Glu	Asn	Leu	Gln	Ser	Ser	Lys	Ser	Thr	Arg	Ala	Thr	Ile	Leu
	766				770					775				780	
	Val	Arg	Leu	Asp	Thr	Gly	Gly	Gln	Glu	Gly	Leu	Gln	Tyr	Gln	Pro
25	781				785					790				795	
	Gly	Asp	His	Ile	Gly	Val	Cys	Pro	Pro	Asn	Arg	Pro	Gly	Leu	Val
	796				800					805				810	
	Glu	Ala	Leu	Leu	Ser	Arg	Val	Glu	Asp	Pro	Pro	Ala	Pro	Thr	Glu
	811				815					820				825	
30	Pro	Val	Ala	Val	Glu	Gln	Leu	Glu	Lys	Gly	Ser	Pro	Gly	Gly	Pro
	826				830					835				840	
	Pro	Pro	Gly	Trp	Val	Arg	Asp	Pro	Arg	Leu	Pro	Pro	Cys	Thr	Leu
	841				845					850				855	
	Arg	Gln	Ala	Leu	Thr	Phe	Phe	Leu	Asp	Ile	Thr	Ser	Pro	Pro	Ser
35	856				860					865				870	
	Pro	Gln	Leu	Leu	Arg	Leu	Leu	Ser	Thr	Leu	Ala	Glu	Glu	Pro	Arg
	871				875					880				885	
	Glu	Gln	Gln	Glu	Leu	Glu	Ala	Leu	Ser	Gln	Asp	Pro	Arg	Arg	Tyr
	886				890					895				900	
40	Glu	Glu	Trp	Lys	Trp	Phe	Arg	Cys	Pro	Thr	Leu	Leu	Glu	Val	Leu
	901				905					910				915	
	Glu	Gln	Phe	Pro	Ser	Val	Ala	Leu	Pro	Ala	Pro	Leu	Leu	Leu	Thr
	916				920					925				930	
	Gln	Leu	Pro	Leu	Leu	Gln	Pro	Arg	Tyr	Tyr	Ser	Val	Ser	Ser	Ala
45	931				935					940				945	
	Pro	Ser	Thr	His	Pro	Gly	Glu	Ile	His	Leu	Thr	Val	Ala	Val	Leu

	946			950					955				960		
	Ala	Tyr	Arg	Thr	Gln	Asp	Gly	Leu	Gly	Pro	Leu	His	Tyr	Gly	Val
	961			965					970				975		
5	Cys	Ser	Thr	Trp	Leu	Ser	Gln	Leu	Lys	Pro	Gly	Asp	Pro	Val	Pro
	976			980					985				990		
	Cys	Phe	Ile	Arg	Gly	Ala	Pro	Ser	Phe	Arg	Leu	Pro	Pro	Asp	Pro
	991			995					1000				1005		
	Ser	Leu	Pro	Cys	Ile	Leu	Val	Gly	Pro	Gly	Thr	Gly	Ile	Ala	Pro
	1006			1010					1015				1020		
10	Phe	Arg	Gly	Phe	Trp	Gln	Glu	Arg	Leu	His	Asp	Ile	Glu	Ser	Lys
	1021			1025					1030				1035		
	Gly	Leu	Gln	Pro	Thr	Pro	Met	Thr	Leu	Val	Phe	Gly	Cys	Arg	Cys
	1036			1040					1045				1050		
	Ser	Gln	Leu	Asp	His	Leu	Tyr	Arg	Asp	Glu	Val	Gln	Asn	Ala	Gln
15	1051			1055					1060				1065		
	Gln	Arg	Gly	Val	Phe	Gly	Arg	Val	Leu	Thr	Ala	Phe	Ser	Arg	Glu
	1066			1070					1075				1080		
	Pro	Asp	Asn	Pro	Lys	Thr	Tyr	Val	Gln	Asp	Ile	Leu	Arg	Thr	Glu
	1081			1085					1090				1095		
20	Leu	Ala	Ala	Glu	Val	His	Arg	Val	Leu	Cys	Leu	Glu	Arg	Gly	His
	1096			1100					1105				1110		
	Met	Phe	Val	Cys	Gly	Asp	Val	Thr	Met	Ala	Thr	Asn	Val	Leu	Gln
	1111			1115					1120				1125		
	Thr	Val	Gln	Arg	Ile	Leu	Ala	Thr	Glu	Gly	Asp	Met	Glu	Leu	Asp
25	1126			1130					1135				1140		
	Glu	Ala	Gly	Asp	Val	Ile	Gly	Val	Leu	Arg	Asp	Gln	Gln	Arg	Tyr
	1141			1145					1150				1155		
	His	Glu	Asp	Ile	Phe	Gly	Leu	Thr	Leu	Arg	Thr	Gln	Glu	Val	Thr
	1156			1160					1165				1170		
30	Ser	Arg	Ile	Arg	Thr	Gln	Ser	Phe	Ser	Leu	Gln	Glu	Arg	Gln	Leu
	1171			1175					1180				1185		
	Arg	Gly	Ala	Val	Pro	Trp	Ala	Phe	Asp	Pro	Pro	Gly	Ser	Asp	Thr
	1186			1190					1195				1200		
	Asn	Ser	Pro												
35	1201		1203												

To obtain polyclonal antibodies to endothelial NO synthase, it is also possible to use a fragment of endothelial NO synthase, selected, for example, from the following sequences:

40

SEQ ID NO: 10

Pro Trp Ala Phe
1192 1195

5 SEQ ID NO: 11

Gly Ala Val Pro
1189 1192

SEQ ID NO: 12

10 Arg
1185
His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp Pro Pro Gly Pro
1186 1190 1195 1200
Asp Thr Pro Gly Pro
1201 1205

15

SEQ. ID. NO. 13

Ala Phe Asp Pro Pro Gly Pro
11941195 1200
20 Asp Thr Pro Gly Pro
1201 1205

SEQ. NO. 14

25 His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp
1186 1190 11951196

SEQ ID NO: 15

30 His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp Pro Pro Gly Pro
1186 1190 1195 1200
Asp Thr Pro Gly Pro
1201 1205

The activated potentiated form of each component of the combination may be prepared from an initial solution by homeopathic potentization, preferably using the

method of proportional concentration decrease by serial dilution of 1 part of each preceding solution (beginning with the initial solution) in 9 parts (for decimal dilution), or in 99 parts (for centesimal dilution), or in 999 parts (for millesimal dilution) of a neutral solvent, starting with a concentration of the initial solution of antibody in the solvent, preferably, water or a water-ethyl alcohol mixture, in the range from about 0.5 to about 5.0 mg/ml, coupled with external impact. Preferably, the external impact involves multiple vertical shaking (dynamization) of each dilution. Preferably, separate containers are used for each subsequent dilution up to the required potency level, or the dilution factor. This method is well-accepted in the homeopathic art. See, e.g. V. Schwabe "Homeopathic medicines", M., 1967, p. 14-29, incorporated herein by reference for the purpose stated.

For example, to prepare a 12-centesimal dilution (denoted C12), one part of the initial matrix solution of antibodies, for example, to C-terminal fragment of angiotensin II AT1 receptor with the concentration of 3.0 mg/ml is diluted in 99 parts of neutral aqueous or aqueous-alcohol solvent (preferably, 15%-ethyl alcohol) and then vertically shaken many times (10 and more) to create the 1st centesimal dilution (denoted as C1). The 2nd centesimal dilution (C2) is prepared from the 1st centesimal dilution C1. This procedure is repeated 11 times to prepare the 12th centesimal dilution C12. Thus, the 12th centesimal dilution C12 represents a solution obtained by 12 serial dilutions of one part of the initial matrix solution of antibodies to C-terminal fragment of angiotensin II AT1 receptor with the concentration of 3.0 mg/ml in 99 parts of a neutral solvent in different containers, which is equivalent to the centesimal homeopathic dilution C12. Similar procedures with the relevant dilution factor are performed to obtain dilutions C30 and C 200. The intermediate dilutions may be tested in a desired biological model to check activity. The preferred activated potentiated forms for both antibodies comprising the combination of the invention are a mixture of C12, C30, and C200 dilutions. When using the mixture of various homeopathic dilutions (primarily centesimal) of the active substance as biologically active liquid component, each component of the composition (e.g., C12, C30, C200) is prepared separately according to the above-described procedure until

the next-to-last dilution is obtained (e.g., until C11, C29, and C199 respectively), and then one part of each component is added in one container according to the mixture composition and mixed with the required quantity of the solvent (e.g. with 97 parts for centesimal dilution).

5 It is possible to use the active substance as mixture of various homeopathic dilutions, e.g. decimal and/or centesimal (D20, C30, C100 or C12, C30, C50 etc.), the efficiency of which is determined experimentally by testing the dilution in a suitable biological model, for example, in models described in the examples herein.

10 In the course of potentiation and concentration decrease, the vertical shaking may be substituted for external exposure to ultrasound, electromagnetic field or any similar external impact procedure accepted in the homeopathic art.

15 Preferably, the pharmaceutical composition of the invention may be in the form of a liquid or in the solid unit dosage form. The preferred liquid form of the pharmaceutical composition is a mixture, preferably at a 1:1 ratio of the activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1 receptor and the activated potentiated form of antibodies to endothelial NO-synthase. The preferred liquid carrier is water or a water-ethyl alcohol mixture.

20 The solid unit dosage form of the pharmaceutical composition of the invention may be prepared by impregnating a solid, pharmaceutically acceptable carrier with the mixture of the activated potentiated form aqueous or aqueous-alcohol solutions of active components which are mixed, primarily in 1:1 ratio and used in liquid dosage form. Alternatively, the carrier may be impregnated consecutively with each requisite dilution. Both orders of impregnation are acceptable.

25 Preferably, the pharmaceutical composition in the solid unit dosage form is prepared from granules of the pharmaceutically acceptable carrier which were previously saturated with the aqueous or aqueous-alcoholic dilutions of the activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1 receptor and the activated potentiated form of antibodies to endothelial NO-synthase. The solid dosage form may be in any form known in the pharmaceutical art, including a tablet, a capsule, a lozenge, and others. As an inactive
30

pharmaceutical ingredients one can use glucose, sucrose, maltose, amyllum, isomaltose, isomalt and other mono- oligo- and polysaccharides used in manufacturing of pharmaceuticals as well as technological mixtures of the above mentioned inactive pharmaceutical ingredients with other pharmaceutically acceptable excipients, for example isomalt, crospovidone, sodium cyclamate, sodium saccharine, anhydrous citric acid etc), including lubricants, disintegrants, binders and coloring agents. The preferred carriers are lactose and isomalt. The pharmaceutical dosage form may further include standard pharmaceutical excipients, for example, microcrystalline cellulose, magnesium stearate and citric acid.

The example of preparation of the solid unit dosage form is set forth below. To prepare the solid oral form, 100-300 µm granules of lactose are impregnated with aqueous or aqueous-alcoholic solutions of the activated-potentiated form of antibodies to C-terminal fragment of angiotensin II AT1 receptor and the activated-potentiated form of antibodies to endothelial NO-synthase in the ratio of 1 kg of antibody solution to 5 or 10 kg of lactose (1:5 - 1:10). To effect impregnation, the lactose granules are exposed to saturation irrigation in the fluidized boiling bed in the boiling bed plant (e.g. "Hüttlin Pilotlab" by Hüttlin GmbH) with subsequent drying via heated air flow at a temperature below 40°C. The estimated quantity of the dried granules (10 to 34 weight parts) saturated with the activated potentiated form of antibodies is placed in the mixer, and mixed with 25 to 45 weight parts of "non-saturated" pure lactose (used for the purposes of cost reduction and simplification and acceleration of the technological process without decreasing the treatment efficiency), together with 0.1 to 1 weight parts of magnesium stearate, and 3 to 10 weight parts of microcrystalline cellulose. The obtained tablet mass is uniformly mixed, and tableted by direct dry pressing (e.g., in a Korsch – XL 400 tablet press) to form 150 to 500 mg round pills, preferably, 300 mg. After tableting, 300 mg pills are obtained that are saturated with aqueous-alcohol solution (3.0-6.0 mg/pill) of the combination of the activated-potentiated form of antibodies to C-terminal fragment of angiotensin II AT1 receptor and the activated potentiated form of antibodies to

endothelial NO-synthase. Each component of the combination used to impregnate the carrier is in the form of a mixture of centesimal homeopathic dilutions, preferably, C12, C30 and C200. While the invention is not limited to any specific theory, it is believed that the activated potentiated form of the antibodies described herein do not
5 contain the molecular form of the antibody in an amount sufficient to have biological activity attributed to such molecular form. The biological activity of the combination drug (combination pharmaceutical composition) of the invention is amply demonstrated in the appended examples.

The combination pharmaceutical composition of the invention may be used for
10 administration to patients having any cardiovascular condition specifically for the purpose of improving the quality of life of such patients. Such use is particularly advantageous because the side effect profile of the combination of the invention is highly favorable. Patients with cardiovascular conditions suffer from decreased quality of life due to their symptoms and decrease in their ability to engage in routine
15 tasks, such as walking, moving items in their environment, depression, and anxiety. As shown in the appended examples, the administration of the combination drug of the invention to such patients improves their overall quality of life, ability to walk, decreases anxiety and depression associated specifically with their primary condition of the cardiovascular system.

The population of patients suffering from cardiovascular conditions, such as,
20 for example, chronic heart failure, asthenia, dystonia, hypertension, and others, is routinely prescribed a variety of medications to treat their conditions. The use of the combination drug of the invention with such additional therapeutic agents to improve the quality of life, to treat chronic heart failure, dystonia, asthenia, and hypertension
25 is specifically contemplated. Non-limiting examples of suitable additional therapeutic agents include:

- ACE inhibitors, including combined ones (Enap, Enalapril, Capoten, Renitec, Prestarium (Berlipril, Diron, Capoten, Quadropril, Monopril, Renitec, Prestarium, Noliprel-Forte, Enap-N));

- diuretics (Furosemide, Veroshpiron, Hypothiazid, Arifon Retard, Indapamide, Hypothiazid, Diuver, Indap, Indapamide);

- β -adrenergic blockers (Egilok, Atenolol, Concor, Betaloc ZOK);

- nitrates (Dilasidom, Kardiket, Kardiket-Retord, Mitrolinate, MonoMak, 5 Monocinque, Nitroglycerin, Nitrosorbid, Olicard, Pectrol, Sydnopharm);

- cardiac glycosides (Digoxin);

- calcium antagonists (Normodipin, Cordaflex, Amlovas, Amlodipine, Amlovas, Amlotop, Cardilopin, Cordaflex, Cordipin XL);

- hypolipidemic agents (Vasilip, Liprimar, Liptonorm, Simvahexal, Simvastol, 10 Simvacard, Simgal, Tulip);

- antiaggregants (Acetylsalicylic acid, CardiASK, Cardiomagnyl, Thrombo ASS);

- antihypoxants (Preductal MB, Preductal, Trimectal);

- anticoagulants (Warfarin).

15 Preferably, for the purpose of improving the quality of life, the combination drug of the invention is administered from once daily to four times daily, each administration including one or two combination unit dosage forms.

It is known that a variety of cardiovascular conditions is accompanied with anxiety and/or depression in the relevant population of patients. The administration 20 of the combination drug described herein to such patients results in a statistically significant improvement in the levels of anxiety and/or depression as measured by the well-accepted tests described herein and exemplified in the examples.

The combination pharmaceutical composition of the invention is also useful for treating chronic heart failure, primarily at the 1st and 2nd stages of the disease. It 25 has been demonstrated experimentally that the administration of the combination pharmaceutical composition of the invention to patients that suffer from chronic heart failure leads to statistically significant improvement in rigidity parameters of carotid radial artery segments and carotid femoral artery segments. The combination of the invention provides an unexpected synergistic therapeutic effect and enhanced 30 influence on vascular remodeling and endothelium dysfunction that is significant for

the process and progression of treatment of chronic heart failure, as also on the improvement of the patients' life quality, on morphological parameters of the heart and tolerance to physical exercise, which is confirmed by clinical trials.

The administration of the combination pharmaceutical composition for
 5 treatment of patients with chronic heart failure improves the life quality parameters
 evaluated by such criteria as depression, anxiety, walking duration, increased
 tolerance to physical exercise, etc. The patients that suffer from chronic heart failure
 also often suffer from associated anxiety and/or depression. The administration of
 the combination pharmaceutical composition to patients that suffer from chronic
 10 heart failure leads to a statistically significant reduction in anxiety and depression
 associated with their primary condition. It has been demonstrated experimentally
 that the administration of the combination to patients with chronic heart failure leads
 to statistically significant improvement in The Minnesota Living with Heart Failure
 questionnaire Score, Kansas City Cardiomyopathy Questionnaire Total Score, HADS
 15 Total Score, and 6-minute walking test.

The population of patients suffering from chronic heart failure is routinely
 prescribed a variety of medications to treat their condition. The use of the
 combination drug of the invention with such additional therapeutic agents to improve
 the quality of life, to treat chronic heart failure, dystonia, asthenia, and hypertension
 20 is specifically contemplated. Non-limiting examples of suitable additional therapeutic
 agents include ACE inhibitors, including combined ones (Enap, Enalapril, Capoten,
 Renitec, Prestarium (Berlipril, Diroton, Capoten, Quadropril, Monopril, Renitec,
 Prestarium, Noliprel-Forte, Enap-N));

- diuretics (Furosemide, Veroshpiron, Hypothiazid, Arifon Retard, Indapamide,
 25 Hypothiazid, Diuver, Indap, Indapamide);

- β -adrenergic blockers (Egilok, Atenolol, Concor, Betaloc ZOK);

- nitrates (Dilasidom, Kardiket, Kardiket-Retord, Mitrolinate, MonoMak,
 Monocinque, Nitroglycerin, Nitrosorbid, Olicard, Pectrol, Sydnopharm);

- cardiac glycosides (Digoxin);

- calcium antagonists (Normodipin, Cordaflex, Amlovas, Amlodipine, Amlovas, Amlotop, Cardilopin, Cordaflex, Cordipin XL);

- hypolipidemic agents (Vasilip, Liprimar, Liptonorm, Simvahexal, Simvastol, Simvacard, Simgal, Tulip);

5 - antiaggregants (Acetylsalicylic acid, CardiASK, Cardiomagnyl, Thrombo ASS);

- antihypoxants (Preductal MB, Preductal, Trimectal);

- anticoagulants (Warfarin).

10 Particularly preferred are those additional therapeutic agents that are used in the medical art to treat chronic heart failure.

Preferably, for the purpose of treating chronic heart failure, the combination drug of the invention is administered from once daily to four times daily, each administration including one or two combination unit dosage forms. Each of the specific administration regiments within the described range is separately and specifically contemplated.

15 Separate administration of two independently prepared unit dosage forms, each containing one of the activated potentiated forms of antibodies of the combination is also contemplated.

20 The combination pharmaceutical composition of the invention is also useful for treating patients that suffer from asthenia and/or vegetative-vascular dystonia. It has been demonstrated experimentally that the administration of the combination pharmaceutical composition of the invention to patients that suffer from asthenia and/or vegetative-vascular dystonia leads to statistically significant improvement in rigidity parameters of carotid radial artery segments and carotid femoral artery segments. The combination pharmaceutical composition of the invention provides an unexpected synergistic therapeutic effect and enhanced influence on vascular remodeling and endothelium dysfunction that is significant for the process and progression of treatment of asthenia and/or vegetative-vascular dystonia, as also on the improvement of the patients' life quality, on morphological parameters of the heart and tolerance to physical exercise, which is confirmed by clinical trials.

The administration of the combination pharmaceutical composition for treatment of patients with asthenia and/or vegetative-vascular dystonia improves the life quality parameters evaluated by such criteria as depression, anxiety, walking duration, increased tolerance to physical exercise, etc. The patients that suffer from
 5 asthenia and/or vegetative-vascular dystonia also often suffer from associated anxiety and/or depression. The administration of the combination pharmaceutical composition to patients that suffer from asthenia and/or vegetative-vascular dystonia leads to a statistically significant reduction in anxiety and depression associated with their primary condition. It has been demonstrated experimentally that the
 10 administration of the combination pharmaceutical composition to patients with asthenia and/or vegetative-vascular dystonia leads to statistically significant reduction in MFI-20 scale score for mental asthenia and improvements in trait anxiety as measured by the Spielberg test.

The population of patients suffering from asthenia and/or vegetative-vascular
 15 dystonia is routinely prescribed a variety of medications to treat their condition. The use of the combination pharmaceutical composition of the invention with such additional therapeutic agents to improve the quality of life, to treat dystonia and asthenia is specifically contemplated. Non-limiting examples of suitable additional therapeutic agents include

- 20 - ACE inhibitors, including combined ones (Enap, Enalapril, Capoten, Renitec, Prestarium (Berlipril, Diroton, Capoten, Quadropril, Monopril, Renitec, Prestarium, Noliprel-Forte, Enap-N));
- diuretics (Furosemide, Veroshpiron, Hypothiazid, Arifon Retard, Indapamide, Hypothiazid, Diuver, Indap, Indapamide);
- 25 - β -adrenergic blockers (Egilok, Atenolol, Concor, Betaloc ZOK);
- nitrates (Dilasidom, Kardiket, Kardiket-Retord, Mitrolinate, MonoMak, Monocinque, Nitroglycerin, Nitrosorbid, Olicard, Pectrol, Sydnopharm);
- cardiac glycosides (Digoxin);
- calcium antagonists (Normodipin, Cordaflex, Amlovas, Amlodipine,
 30 Amlovas, Amlotop, Cardilopin, Cordaflex, Cordipin XL);

- hypolipidemic agents (Vasilip, Lipimar, Liptonorm, Simvahexal, Simvastol, Simvacard, Simgal, Tulip);

- antiaggregants (Acetylsalicylic acid, CardiASK, Cardiomagnyl, Thrombo ASS);

5 - antihypoxants (Preductal MB, Preductal, Trimectal);

- anticoagulants (Warfarin).

Particularly preferred are those additional therapeutic agents that are used in the medical art to treat asthenia and/or vegetative-vascular dystonia.

Preferably, for the purpose of treating asthenia and/or vegetative-vascular
10 dystonia, the combination pharmaceutical composition of the invention is administered from once daily to four times daily, each administration including one or two combination unit dosage forms. Each of the specific administration regimens within the described range is separately and specifically contemplated. Separate administration of two independently prepared unit dosage forms, each containing
15 one of the activated potentiated forms of antibodies of the combination drug is also contemplated.

The combination pharmaceutical composition of the invention is also useful for treating hypertension. It has been demonstrated experimentally that the administration of the combination pharmaceutical composition leads to statistically
20 significant reduction in systolic blood pressure. The population of patients suffering from hypertension is routinely prescribed a variety of medications to treat their condition. The use of the combination pharmaceutical composition of the invention with such additional therapeutic agents to treat hypertension is specifically contemplated. Non-limiting examples of suitable additional therapeutic agents
25 include

- ACE inhibitors, including combined ones (Enap, Enalapril, Capoten, Renitec, Prestarium (Berlipril, Diroton, Capoten, Quadropril, Monopril, Renitec, Prestarium, Noliprel-Forte, Enap-N));

- diuretics (Furosemide, Veroshpiron, Hypothiazid, Arifon Retard, Indapamide,
30 Hypothiazid, Diuver, Indap, Indapamide);

chronic heart failure (CHF) was conducted in 75 female Wistar rats (4-5 months old, weighing 220-250 g).

For chronic heart failure modeling all rats were receiving isadrine (isoproterenol hydrochloride, Sigma, Germany) 80 mg/kg twice with 24-hour interval.

5 Animals (15 per group) were receiving the following drugs by gastric gavage: group 1 – 7.5 mL/kg of ULD of antibodies against a C-terminal fragment of angiotensin II AT1 receptor (mixture of homeopathic dilutions C12, C30 and C200), group 2 – 7.5 mL/kg of ULD of anti-endothelial NO synthase antibodies (mixture of homeopathic dilutions C12, C30 and C200), group 3 – 15 mL/kg of combined drug (mixture of homeopathic dilutions C12, C30 and C200), group 4 – 7.5 mL/kg (dose of 10 mg/kg) of losartan as
10 a comparator, group 5 (control) – 7.5 mL/kg of distilled water.

The drug efficacy was assessed on Days 7, 14, 28 after repeated isadrine administration based on electrocardiogram and rheogram data, as well as the results of exercise tolerance test (swimming with weight load of 15% of the body weight at a
15 water temperature of 24°C).

In 7 days following the second isadrine injection animals from all experimental groups developed myocardial damage manifesting in statistically significantly decreased myocardial contractility (decrease in systolic volume and cardiac output), cardiac electrical activity impairment (T-wave amplitude increase), proven lowering
20 of exercise tolerance (swimming duration).

Following 14 days of the study drugs administration positive changes in evaluated parameters were noted. In 28 days significant improvement of the evaluated parameters was demonstrated compared to the results obtained on Day 7 (after the second isadrine injection) ($p < 0.05$) (see Table 1).

25

30

Table 1. Changes in efficacy endpoints of CHF treatment (in % on Day 28 vs. Day 7)

Evaluated parameter	ULD of antibodies against C-terminal fragment of human angiotensin II AT1 receptor	ULD of anti-endothelial NO synthase antibodies	Combined drug based on ULD of antibodies to C-terminal fragment of angiotensin II AT1 receptor and antibodies to endothelial NO synthase	Losartan	Distilled water
T-wave amplitude	-22.37%	-15.81%	-30.03%	-31.64%	-7.7%
Heart rate	-10.75%	-8.57%	-22.43%	-21.28%	-6.74%
Systolic volume (by Kubrick)	46.98%	22.62%	69.74%	64.01%	15.0%
Cardiac output	35.11%	12.43%	49.40%	55.31%	7.13%

Thus, concomitant administration of the study preparations of ULD antibodies (combined drug) proved to be more effective compared to monotherapies, increased exercise tolerance, helped normalize cardiac electrical activity and cardiac cycle, systemic and intracardiac hemodynamics in rats with CHF. ULD antibodies combination demonstrated non-inferiority in terms of efficacy compared to losartan.

Example 2. Double blind, placebo-controlled clinical study of a combination of activated potentiated forms of antibodies to the C-terminal fragment of the angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in human patients with chronic heart failure to evaluate key parameters of the CHF pathology.

80 patients (CHF of II-IV functional class (FC), left ventricular ejection fraction (LVEF) less than 40%) were divided in 4 equal treatment and control groups for a 6 months study. The background therapy was not discontinued (β -blocker bisoprolol, ACE inhibitor enalapril, aspirin (unless contraindicated); administration of diuretics, nitrates, digoxin was also admitted). Group 1 received the activated potentiated form of antibodies to a C-terminal fragment of the angiotensin II AT1 receptor (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 2 received the activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 3 received the combination pharmaceutical composition comprising both activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1-receptor (mixture of homeopathic dilutions C12, C30, C200) and activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 4 received placebo (3 tablets/day, n=20) The groups were comparable in the initial study parameters: in age and sex, and severity (class of CHF and LVEF) and duration of the disease.

Before and after treatment, the patients were evaluated for the effect of the administered medications on vascular remodeling and endothelium dysfunction that is important for the CHF process and progression. The effects of the medications on the processes of vascular remodeling were evaluated by pulse wave velocity (PWV) ("Colson" system) in the carotid-femoral (CF) (elastic type) and carotid-radial (CR) (muscle type) segments of arteries.

Table 2 shows the dynamics in the rates of pulse wave velocity in the carotid-femoral (CF) (elastic type) and carotid-radial (CR) (muscle type) segments of arteries.

Table 2

Groups /Parameters	ULDs ¹ of Abs ² to C-terminal fragment of AT1 receptor of angiotensin II	ULD of Abs to endothelial NO-synthase	Combination of ULDs of Abs to C-end fragment of AT1 receptor of angiotensine II and ULD of Abs to endothelial NO-synthase	Placebo

	^	&	Δ%	^	&	Δ%	^	&	Δ%	^	&	Δ%
CF, m/c	9,7± 0.5	8± 0.6	-14.8*	10.1± 0.5	9.8± 0.4	-2.97	10.8± 0.3	8.6± 0.6	-20.3*	8.2± 0.4	8.2± 0.5	0.1
CR, m/c	8.6± 0.2	8.9± 0.3	2.9	8.8± 0.1	8.3± 0.3	-5.7	8.9± 0.5	7.6± 0.7	-15.6 *#	9.1± 0.3	9.7± 0.3	6.4 *

(^) denotes initial value

(&) denotes 6 month after beginning of administration

(*) denotes difference from initial value is verifiable with p value < 0.05.

(#) denotes difference from the group receiving ULDs of Abs to C-terminal

5 fragment AT1 receptor angiotensin II with verifiable difference in p value of < 0.05.

(\$) denotes difference from the group receiving ULDs of Abs to endothelial

NO-synthase with significant difference in p value of < 0.05.

(1) ULD denotes ultra-low doses.

(2) Abs denotes antibodies.

10

After 6 months of treatment, only group 3 showed a proven effect of the claimed pharmaceutical composition on the stiffness of muscular type arteries. Group 1 which received ULD of antibodies to a C-terminal fragment of angiotensin II AT1 receptor, and group 3 which received the combination pharmaceutical composition of the invention showed a proven increase in the stiffness of elastic type

15 arteries.

Example 3.

Double blind, placebo-controlled clinical study of a combination of activated

20 potentiated forms of antibodies to the C-terminal fragment of angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in human patients with chronic heart failure to evaluate key measurement of quality of life.

25 80 patients (CHF of II-IV functional class (FC), left ventricular ejection fraction (LVEF) less than 40%) were divided in 4 equal treatment and control groups for a 6 months study. The background therapy was not discontinued (bisoprolol β-blocker,

ACE inhibitor enalapril, aspirin (unless contraindicated); administration of diuretics, nitrates, digoxin was also admitted). Group 1 received the activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1 receptor (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 2 received the
5 activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 3 received the combination pharmaceutical composition comprising both activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1 receptor (mixture of homeopathic dilutions C12, C30, C200) and activated potentiated form of antibodies
10 to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 4 received placebo (3 tablets/day, n=20). The groups were comparable in the initial study parameters: in age and sex, and severity (class of CHF and LVEF) and duration of the disease. Before and after treatment, the patients were evaluated for the life quality (Minnesota and Kansas questionnaires),
15 morphological parameters of the heart, and tolerance to physical exercise.

Table 3 shows the results of the study in the form of dynamics in the basic parameters of the treatment efficacy.

After 6 months of treatment, the patients in group 1 treated with ULD of antibodies to a C-terminal fragment of angiotensin II AT1 receptor showed a
20 significant improvement of the life quality, improvement of the left ventricular systolic function, and an increased tolerance to physical exercise. Group 2 showed a proven decrease in the anxiety and depression levels and in the life quality, which were evaluated using the Kansas questionnaire. The study confirmed that the maximum therapeutic effect was achieved with the combination pharmaceutical composition of
25 the invention in combination with the standard CHF therapy, which was administered to patients from group 3 that showed a proven positive dynamics in all parameters under study.

The combination of activated (potentiated) forms of antibodies to a C-terminal fragment of angiotensin II AT1 receptor and to endothelial nitric oxide synthase (NO-synthase) in the pharmaceutical composition of the invention (combination drug)
30

provides an unexpected synergistic therapeutic effect implying an enhanced influence on vascular remodeling and endothelium dysfunction that is critical for the CHF process and progression, as also on the improvement of the patients' life quality, on morphological parameters of the heart and tolerance to physical exercise, which is confirmed by clinical trials. The results are set forth in Table 3.

Table 3

Groups/ Parameters	ULD ¹ of Abs ² to C-terminal fragment of AT1 receptor of angiotensin II			ULD of Abs to endothelial NO-synthase			Combination of ULD of Abs to C-terminal fragment of AT1 receptor of angiotensin II and ULD of Abs to endothelial NO-synthase			Placebo		
	^	&	Δ%	^	&	Δ%	^	&	Δ%	^	&	Δ%
Minnesota ³	47.5 ± 2.8	39.1 ± 3.8 **	-17.6	48.1 ± 3.7	40.8 ± 3.8	-15.2	43.9 ± 2.8	32.0 ± 4.9 ***\$	-27.1	48.3 ± 3.7	42.4 ± 2.9 **	-12.2
Kansas ⁴	82.1 ± 2.3	70.1 ± 5.5 ***	-14.6	81.5 ± 2.5	72.0 ± 8.2 *	-11.7	87.7 ± 2.3	65.7 ± 7.3 ***\$	-25.1	83.8 ± 3.5	60.3 ± 6.8	-7.2

HADS ⁵	15.3± 1.0	12.5± 0.9**	-18.5	16.2± 1.7	11.34 ±2.1 ***	-30.3	16.2 ± 1.3	8.4 ±0.9 *** # \$\$	-48.1	17.3 ± 1.1	15.9 ±1.1	-8.1
FC CHF ⁶	2.7 ± 0.1	2.2± 0.1***	-17.3	2.9± 0.1	2.7±0 .2	-7.3	3.0 ± 0.2	1.9 ±0.1 *** # \$	-36.6	2.7 ± 0.1	2.5 ±0.1	-6.2
FF LV ⁷	27.1± 0.9	33.6± 1.5**	24.0	28.2 ±1.5	25.3± 1.7	10.3	25.3 ± 1.1	34.6 ±1.9 *** # \$	36.7	26.4 ± 1.1	28.0 ±1.4	6.3
6-minute walk test	378.7 ±12.4	419.6 ±13.7* **	10.8	383.1 ± 15.3	416.8 ±17.2	8.8	378.7 ±12.4	450.1 ±17.7 ** # \$	18.9	390.5 ± 11.9	409.1 ±11.5	4.8

*, **, *** - p values < 0.05, 0.01 and 0.001, respectively

#- difference from group receiving ULDs of Abs to C-terminal fragment AT1 of angiotensin receptor II with verifiable with p value < 0.05

5 \$, \$\$ - difference from the group receiving ULDs of Abs to endothelial NO-synthase is significant at p values of 0.05 and 0.01, respectively.

(1)-ULD means ultra low doses

(2) Abs means antibodies

(3) "Minnesota" denotes Minnesota Questionnaire

- (4) "Kansas" denotes Kansas Questionnaire
- (5) HADS denotes HADS total score
- (6) FC CHF denotes patients with chronic heart failure, functional class
- (7) FF LV denotes fraction of functioning of left ventricle.

5

Example 4. Double blind placebo-controlled randomized clinical study of combination of the activated potentiated forms of antibodies to a C-terminal fragment of angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in human patients with vegeto-vascular dystonia .

The clinical efficiency and safety of the combination pharmaceutical composition comprising activated-potentiated forms of antibodies to a C-terminal fragment of angiotensin II AT1 receptor and to endothelial NO-synthase (the mixtures of homeopathic dilutions C12, C30, C200) were tested on 60 male and female patients (male — 26.7% (n=16), female — 73.3% (n=44) of the total quantity of patients) with asthenia (more than 12 points according to "general asthenia" claim of the subjective asthenia scale MFI-20 (Multidimensional Fatigue Inventory)) and neurocirculatory asthenia (more than 15 points on the vegetative changes scale) aged 29-64 (average age 47.1 ± 1.77 years old).

The patients were randomized in two groups (n=30 each). The first group of patients received the combination pharmaceutical composition three times a day (1 pill each time). The second group patients took the placebo tablets three times a day (1 pill each time). The overall time of administration and patients monitoring lasted for 6 months.

The efficiency of treatment was evaluated by dynamics in the parameters of the subjective asthenia scale, questionnaires for estimating the anxiety and depression levels (Spielberger and Beck scale), questionnaire for evaluating the sleep and vegetative changes.

The dynamics in the vegetative and psychological state changes, in the patients' life quality during the treatment is shown in the Table 4.

Parameter	Combined Medication, n=30				Placebo, n=30			
	Visit 1	Visit 2 (4 weeks)	Visit 3 (12 weeks)	Visit 4 (24 weeks)	Visit 1	Visit 2 (4 weeks)	Visit 3 (12 weeks)	Visit 4 (24 weeks)
General asthenia (MFI-20 scale) (M±m)	12.9 ± 0.21	12.3 ± 0.29	11.5 ± 0.35 *** #	10.9 ± 0.27 *** ##	13.1 ± 0.26	12.9 ± 0.23	12.5 ± 0.17 ***	11.9 ± 0.19 ***
Physical asthenia (MFI-20 scale) (M±m)	12.3 ± 0.56	11.7 ± 0.49 **	11.2 ± 0.45 **	10.2 ± 0.39 *** #	12.3 ± 0.38	12.1 ± 0.29	11.7 ± 0.29 **	11.4 ± 0.21 **
Hypoactivity (MFI-20 scale) (M±m)	10.3 ± 0.50	10.1 ± 0.47	9.5 ± 0.42 *	9.1 ± 0.28 *	10.6 ± 0.63	10.1 ± 0.53 *	9.7 ± 0.50 **	9.5 ± 0.38 **
Motivation decrease (MFI-20 scale) (M±m)	9.60 ± 0.55	9.5 ± 0.47	8.6 ± 0.32 *	8.3 ± 0.27 *	9.9 ± 0.75	9.7 ± 0.57	9.4 ± 0.51	9.3 ± 0.48
Psychic asthenia (MFI-20 scale) (M±m)	10.2 ± 0.47	9.7 ± 0.44 *	9.5 ± 0.42 *	8.9 ± 0.41 ** #	10.8 ± 0.51	10.6 ± 0.43	10.5 ± 0.40	10.3 ± 0.38
Questionnaire of vegetative changes (total rate) (M±m)	28.9 ± 1.66	25.3 ± 1.26 **	22.1 ± 1.26 ***	17.6 ± 0.76 *** #	32.1 ± 3.36	29.5 ± 2.57 *	25.9 ± 2.42 ***	22.4 ± 2.08 ***
Questionnaire of sleep evaluation (total rate)	16.9 ± 0.44	18.4 ± 0.42 **	18.9 ± 0.34 ***	19.4 ± 0.29 *** #	16.6 ± 0.67	17.5 ± 0.56 **	18.1 ± 0.47 ***	18.2 ± 0.47 ***
Individual anxiety (Spielberger scale) (M±m)	37.6 ± 2.03	35.6 ± 1.42 *	35.1 ± 1.64 **	32.9 ± 1.23 ***	39.1 ± 3.05	38.7 ± 2.78	36.9 ± 2.45 *	36.9 ± 2.46 *
Reactive anxiety (Spielberger scale) (M±m)	39.3 ± 1.02	35.7 ± 0.72 *** #	34.5 ± 1.02 ***	32.4 ± 0.76 *** #	42.7 ± 2.65	40.3 ± 1.99**	38.3 ± 1.87 **	37.6 ± 1.81**
Beck questionnaire of depression (total rate) (M±m)	9.2 ± 0.44	8.4 ± 0.25 *	8.3 ± 0.27 *	7.8 ± 0.31 ** #	11.1 ± 1.43	10.5 ± 1.19 *	10.0 ± 0.93	9.9 ± 0.90 *

differences are significant vs the initial values at: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

differences are significant vs placebo group: # $p < 0.05$, ## $p < 0.01$, ### $p < 0.001$.

5 It is shown that the administration of the combination pharmaceutical composition during 6 months resulted in a statistically significant decrease in the asthenia level on all MFI-20 subscales. The most significant improvements were observed in the following areas: general asthenia (from 12.9 ± 0.21 to 10.9 ± 0.27 points, $p < 0.01$ of the placebo group), physical asthenia (from 12.3 ± 0.56 to 10.2 ± 0.39 points, $p < 0.05$ of the placebo group), psychic asthenia (from 10.2 ± 0.47 to 8.9 ± 0.41 points, $p < 0.05$ of the placebo group). After one month of treatment, basic group 1 showed a statistically proven decrease in the intensity of vegetative changes (according to the special-purpose questionnaire data), and after six months of treatment there was a statistically significant ($p < 0.05$) difference of the achieved results (decrease of the total rate from 28.9 ± 1.66 to 17.6 ± 0.76) in comparison with the placebo group (decrease of the total rate from 32.1 ± 3.36 to 22.4 ± 2.08).

10 In addition, the administration of the combination pharmaceutical composition contributed to improving the psychological state of the patients in the study. A statistically significant decrease in the anxiety and depression levels was observed after one month of administration, and when the treatment was continued the positive effect was enhanced. After six months, group 1 showed a statistically significant ($p < 0.05$) improvement in the reactive anxiety level on the Spielberger scale (decrease from 39.3 ± 1.02 to 32.4 ± 0.76) and the depression level according to the Beck questionnaire (decrease from 9.2 ± 0.44 to 7.8 ± 0.31) in comparison with the respective parameters of the placebo group (decrease from 42.7 ± 2.65 to 37.6 ± 1.81 and from 11.1 ± 1.43 to 9.9 ± 0.90 respectively). The results were confirmed by the positive dynamics ($p < 0.001$ of the initial rates) in the parameters of the sleep evaluation questionnaire, and after six months of treatment, group 1 showed a statistically significant ($p < 0.05$) difference in the total rate (increase from 16.9 ± 0.44

to 19.4 ± 0.29) in comparison with the placebo group (increase from 16.6 ± 0.67 to 18.2 ± 0.47).

Over the entire period of monitoring, the patients showed a high tolerability and no adverse effects.

5 The study showed that the combination pharmaceutical composition used in the study possesses antiasthenic effect, normalizes the vegetative state of patients with neurocirculatory dystonia and asthenia, and has anti-anxiety and anti-depression influence in the relevant patient population.

10 Example 5. Study of the combination of activated-potentiated forms of antibodies to the C-terminal fragment of the angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with the activated-potentiated form of antibodies to endothelial NO synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in SHR rats in a model of hypertension.

15 The combination of water solution of the activated-potentiated form of antibodies to a C-terminal fragment of the angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, and the activated-potentiated form of antibodies to endothelial NO synthase in a mixture of homeopathic dilutions of C12, C30, C200, was studied in the SHR rat hypertension model. Investigations were
20 conducted on 40 SHR line male rats from (weight 350 ± 50 g, age 4.5 - 5 months) with hypertension, which were divided into 4 groups of 10 animals each.

For 28 days, the animals were treated as follows. Group 1 – 2.5 ml/kg of the activated-potentiated form of antibodies to the C-terminal fragment of human angiotensin II AT1 receptor (a mixture of aqueous dilutions C12, C30, C200) in
25 combination with 2.5 ml/kg of distilled water, Group 2 – 2.5 ml/kg of the activated-potentiated form of antibodies to endothelial NO synthase (a mixture of aqueous dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water, Group 3 – 5 ml/kg of the combination pharmaceutical composition (a mixture of aqueous dilutions C12, C30, C200 for each component), and Group 4 – 5 ml/kg of distilled
30 water.

Systolic blood pressure (SBP) of awake rats was measured with the aid of an indirect method in a tail artery (using a cuff) once a week and 9 hours after the last administration of medicines.

5 All tested compositions demonstrated hypotensive effect ($p < 0.05$): by 28th day, systolic blood pressure (SBD) decreased in comparison with the initial level in Group 1 by – 20.6%; in Group 2 by 14.4%; in Group 3 by 27.6%. In the control Group 4, SBD changes were 1.6% in comparison with the initial values. The results demonstrate a clear synergistic hypotensive effect of the combination pharmaceutical composition.

10

Example 6. Study of the combination of the activated-potentiated forms of antibodies to a C-terminal fragment of angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with the activated-potentiated form of antibodies to endothelial NO synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in NISAG rats in a model of hypertension.

15 The combination water solution of the activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1 receptor, in a mixture of homeopathic dilutions of C12, C30, C200, and the activated potentiated form of antibodies to endothelial NO synthase in a mixture of homeopathic dilutions of C12, C30, C200, was studied in the NISAG rat hypertension model. Investigations were conducted on 20 50 NISAG line male rats (weight 300 g, age 4 months) with hereditary stipulated stress-sensitive arterial hypertension, which were divided into 5 groups by 10 animals each.

25 The animals were given per orally, once a day and for 28 days, the following medications: Group 1 – 2.5 ml/kg of the activated-potentiated form of antibodies to a C-terminal fragment of human angiotensin II AT1 receptor (a mixture of dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water; Group 2 – 2.5 ml/kg of the activated-potentiated form of antibodies to endothelial NO synthase (a mixture of dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water; Group 3 30 - 5 ml/kg of the combination pharmaceutical composition (a mixture of homeopathic

aqueous dilutions C12, C30, C200 of each component); Group 4 - 5 ml/kg (10 ml/kg dose) of the comparison drug (losartan); and Group 5 - 5 ml/kg of distilled water.

Two times a week, 2 to 6 hours after administration of ULD antibodies and losartan, systolic blood pressure (SBP) was measured by an indirect method in a tail artery (using a cuff). The Table 5 shows the dynamics of changes in systolic blood pressure in NISAG line rats, measured by indirect method.

Table 5.

	Initial SBP in mmHg	SBP after 28 days of medicine administration in mmHg	Δ in comparison with the initial level, in mmHg	% of the initial level
ULD antibodies to C-terminal fragment of angiotensin II AT1receptor	176	150	-26	- 14.7 %
ULD antibodies to endothelial NO synthase	175	164.5	-10.5	-6%
ULD antibodies to C-terminal fragment of angiotensin II AT1receptor and to endothelial NO synthase	179.5	140	-39.5	-22%
Losartan	173.5	140.5	-33	-19%
Control (distilled water)	181	178	-3	-1.6%

Example 7. Study of correction of endothelial dysfunction

10 Wistar rats weighed 250 - 300 g were divided into 4 groups. The 1st group received distilled water 9 ml/kg/day (every day, gastrointestinal introduction). The 2nd group received L-NAME (L-NG-nitroarginine methyl ester, which is known to inhibit endothelial isoform, and thus is believed to simulate endothelial dysfunction)

at 12.5 mg/kg. The third group received L-NAME (12.5 mg/kg) in combination with a mixture of C12, C30, and C200 homeopathic dilutions of polyclonal rabbit antibodies to human endothelial NO synthase at 9 ml/kg/day. The 4th group received L-NAME (12.5 mg/ml) in combination with a mixture of C12, C30, and C200 homeopathic dilutions of polyclonal rabbit antibodies to human endothelial NO synthase at 9 ml/kg/day and mixture of C12, C30, C200 homeopathic dilutions of polyclonal rabbit antibodies to a C-terminal fragment of AT1 receptor of angiotensin II at 9 ml/kg/day.

Endothelial dysfunction was simulated by introducing L-NAME at 12.5 mg/ml/day in the course of 28 days. On day 29 from the beginning of the experiment, under anesthesia (sodium thiopental at 50 mg/kg), a catheter was inserted into the left carotid artery to register parameter of arterial pressure (AP). Bolus introduction of the pharmacological agent was made into right femoral vein. The following parameters were measured: systolic arterial pressure (SAP), diastolic arterial pressure (DAP), and frequency of heart beat, which were measured in real time by sensor TSD104A measurement apparatus MP100 made by Biopac System, Inc., USA. The following functional probes were used: endothelium-dependent vasodilation (EDV), intra-vein introduction of acetylcholine (AC) in the dose of 40 µg/kg, endothelium-independent vasodilation (EIV) via intravein introduction of sodium nitroprusside at 30 µg/kg.

The degree of endothelial dysfunction was evaluated via the coefficient of endothelial dysfunction as the ratio of the area of the triangle under the trend of reaction of stabilization of arterial pressure (AP) as a response to introduction of nitroprusside to the area of the triangle under the trend of reaction of stabilization of arterial pressure as a response to introduction of acetylcholine.

The results of the experiments are presented in Table 6.

Table 6.

Groups	Functional Probe	SAP, mm, HG	DAP, mm HG	Vascular reaction observed with EDV with acetyl choline EIV with sodium nitroprusside	CED, (relative units)
Control (daily in-stomach introduction of distilled water at 9 ml/kg/day)	<i>Initial</i>	159.2 ± 5.4	124.2 ± 4.7		1.2 ± 0.1
	Acetyl choline	96.9 ± 6,7	52.0 ± 3.0	3071.2 ± 501.1	
	Sodium Nitroprusside	113.8 ± 6,1	55 ± 2,4	3617.2 ± 560.1	
L-NAME	<i>Initial</i>	204.8 ± 10*	164.2 ± 5.9*		3.5 ± 0.5*
	Acetyl choline	111.3 ± 7.4	64.7 ± 4.3*	3700.2 ± 536.9	
	Sodium Nitroprusside	118.2 ± 9,9	61.4 ± 4.4	11922.8 ± 1838.9*	
L-NAME in combination with a mixture of C12, C30, and C200 homeopathic dilutions of polyclonal rabbit antibodies to human endothelial NO-synthase	<i>Initial</i>	230.1 ± 8.8*	175.1 ± 6.4*		1.8 ± 0.2**
	Acetyl choline	116.4 ± 4.0*	73.5 ± 3.5*	5826.0 ± 801.2**	
	Sodium Nitroprusside	108.1 ± 4.6	60 ± 3.6	9628.5 ± 970.1*	
L-NAME in combination with a mixture of C12, C30, and C200 homeopathic dilutions of polyclonal rabbit antibodies to human endothelial NO-synthase and mixture of C12, C30, C200 homeopathic dilutions of polyclonal rabbit antibodies to C-terminal fragment of AT1 receptor of angiotensin II	<i>Initial</i>	213.6 ± 4.6*	166.8 ± 2.7*		1.4 ± 0.1**
	Acetyl choline	116.9 ± 5.5*	73.3 ± 3.4*	3295.3 ± 201.4	
	Sodium Nitroprusside	122.3 ± 4.6	70.5 ± 4.5*	4546.2 ± 299.4**	

* - p < 0,05 in comparison with the control group; ** - p < 0,05 in comparison with the L-NAME group. S – area under the curve of restoration of arterial hypertension per pharmacological probes, CED-coefficient of endothelial dysfunction.

CLAIMS

1. A combination pharmaceutical composition for administration to a patient suffering from at least one symptom of a cardiovascular condition, said composition comprising a) an activated-potentiated form of an antibody to
5 angiotensin II AT1 receptor, and b) an activated-potentiated form of an antibody to endothelial NO-synthase.

2. The combination pharmaceutical composition of claim 1, wherein said activated-potentiated form of an antibody to angiotensin II AT1 receptor is an activated-potentiated form of an antibody to the C-terminal fragment of the
10 angiotensin II AT1 receptor.

3. The combination pharmaceutical composition of claim 1, wherein said cardiovascular condition is associated with a reduced quality of life of said patient and wherein said administration of said pharmaceutical composition to said patient improves said quality of life of said patient.

15 4. The combination pharmaceutical composition of claim 1, wherein said activated-potentiated form of an antibody to angiotensin II AT1 receptor is in the form of a C12, C30, or C200 homeopathic dilution or a mixture thereof.

5. The combination pharmaceutical composition of claim 1, wherein said activated-potentiated form of an antibody to endothelial NO-synthase is in the form of
20 a C12, C30, or C200 homeopathic dilution or a mixture thereof.

6. The combination pharmaceutical composition of according to one of claims 1 to 4 and 5, which is prepared by impregnating a solid carrier with said activated-potentiated forms of the antibodies.

7. The combination pharmaceutical composition according to one of
25 claims 1 to 4, 5 and 6 for use in the prevention or treatment of the symptoms of a disease or condition associated with a cardiovascular condition, including a reduced overall quality of life ,

8. The combination pharmaceutical composition of claim 7, further comprising an additional therapeutic agent selected from the group consisting of
30 ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium

antagonists, hypolipidemic agents, antiaggregants, antihypoxants, bisoprol, enalapril, aspirin and anticoagulants.

9. The combination pharmaceutical composition of claim 7, wherein said cardiovascular condition is chronic heart failure.

5 10. The combination pharmaceutical composition of claim 7, wherein said patient exhibits statistically significant improvement in rigidity parameters of carotid radial artery segments upon said administration.

11. The combination pharmaceutical composition of claim 7, wherein said cardiovascular condition is asthenia and/or vegetative vascular dystonia.

10 12. The combination pharmaceutical composition of claim 11, wherein said administration of said combination leads to a statistically significant improvement in the mental asthenia by the MFI-20 scale in a suitable population of said patients in reference to the baseline.

15 13. The combination pharmaceutical composition of claim 11, wherein said administration of said combination leads to a statistically significant reduction in the general asthenia by the MFI-20 scale in a suitable population of said patients in reference to the baseline.

20 14. The combination pharmaceutical composition of claim 7, wherein said reduced overall quality of life is associated with anxiety related to said cardiovascular condition.

15 15. The combination pharmaceutical composition of claim 14, wherein said administration of said combination leads to a statistically significant improvement in the Minnesota Total Inventory Score in a suitable population of said patients in reference to the baseline.

25 16. The combination pharmaceutical composition of claim 7, wherein said reduced overall quality of life is associated with depression related to said cardiovascular condition.

17. The combination pharmaceutical composition of claim 16, wherein said administration of said combination leads to a statistically significant reduction in the

Beck Questionnaire Score in a suitable population of said patients in reference to the baseline.

18. The combination pharmaceutical composition according to one of claims 1 to 4, 5, 6, 8 and 9 to 17, wherein said patient is administered one to two of said unit dosage forms, each administration carried out from once daily to four times daily.

19. The combination pharmaceutical composition according to one of claims 1 to 4, 5, 6 and 10-14 for use in the prevention or treatment of the symptoms of a disease or condition associated with a asthenia and/or vegetative-vascular dystonia.

20. The pharmaceutical composition of claim 19, further comprising an additional therapeutic agent selected from the group consisting of ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium antagonists, hypolipidemic agents, antiaggregants, antihypoxants, bisoprol, enalapril, aspirin and anticoagulants.

21. The combination pharmaceutical composition according to one of claims 1-4, 5, 6, 7 and 9 to 17, wherein said activated-potentiated form is prepared by successive centesimal dilutions coupled with vertical shaking of every dilution.

22. The combination pharmaceutical composition according to claims 1-6 for use in treating hypertension.

23. The combination pharmaceutical composition of claim 22, further comprising concomitant administration to said patient of an additional therapeutic agent selected from the group consisting of ACE inhibitors, diuretics; β -adrenergic blockers, nitrates, cardiac glycosides, calcium antagonists, hypolipidemic agents, antiaggregants, antihypoxants, bisoprol, enalapril, aspirin and anticoagulants.

24. A combination pharmaceutical composition for use in treating at least one symptom of a cardiovascular condition, said composition having been obtained by providing a) a potentiated solution of an antibody to angiotensin II AT1 receptor, and b) a potentiated solution of an antibody to endothelial NO-synthase, each prepared by consecutive repeated dilution and multiple shaking of each obtained

solution in accordance with homeopathic technology, and then either combining the potentiated solutions by mixing them, or, alternatively, impregnating a carrier mass with said combined solution or with the solutions separately.

25. The combination pharmaceutical composition of claim 24, wherein said
5 potentiated solution of an antibody to a C-terminal fragment of the angiotensin II AT1 receptor is in the form of a C12, C30, or C200 homeopathic dilution or a mixture thereof.

26. The combination pharmaceutical composition of claim 24, wherein said
10 potentiated solution of an antibody to endothelial NO-synthase is in the form of a C12, C30, or C200 homeopathic dilution or a mixture thereof.

27. The combination pharmaceutical composition of 24, wherein said activated-potentiated form of an antibody to angiotensin II AT1 receptor is an activated-potentiated form of an antibody to the C-terminal fragment of the angiotensin II AT1 receptor.

15