



**AFRICAN REGIONAL INDUSTRIAL PROPERTY
ORGANISATION (ARIPO)**

1111

(11)

(A)

<p>(21) Application Number: AP/P/2000/001894</p> <p>(22) Filing Date: 19990422</p> <p>(24) Date of Grant & 20021023 (45) Publication</p> <hr/> <p>(30) Priority Data</p> <p>(33) Country: EP</p> <p>(31) Number: 98107397.6</p> <p>(32) Date: 19980423</p> <hr/> <p>(84) Designated States:</p> <p style="padding-left: 20px;">GH - GM KE LS MW SD</p> <p style="padding-left: 20px;">SL SZ UG ZW</p>	<p>(73) Applicant(s): AVENTIS PHARMA DEUTSCHLAND GmbH Bruningstrasse 50 D-65929 Frankfurt am Main Germany</p> <p>(72) Inventors: MUKHOPADHYAY, Triptikumar M-24, Hoechst Qtrs Amar Nagar, Mulund (West) Mumbai - 400 082 India (See Overleaf)</p> <p>(74) Representative FISHER, CORMACK & BOTHA P O BOX 74 BLANTYRE MALAWI</p>
--	--

(51) **International Patent Classification (Int. Cl. 7):** C07K 7/56

(54) **Title:** A Process For The Conversion Of Echinocandin Class Of Peptides To Their C4-Homotyrosine Monodeoxy Analogues

(57) **Abstract:**

The invention relates to a process for the conversion of echinocandin class of peptides to their C4-homotyrosine monodeoxy analogues, particularly mulundocandin to deoxy-mulundocandin, which consists of a single step selective reduction of C4-h Tyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues under neutral conditions without prior protection/deprotection of the equally facile C5-Orn (ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.

Inventors Continued


2. JAYVANTI, Kenia - 11
Dev Aashish Ganesh Gawde Road
Mulundi (West)
Mumbai - 400 080
INDIA

-
3. KUMAR, Erra Koteswara
Satya Vijaya - K - 3 Hoechst Qtrs
Amar Nagar
Mulund (West),
Mumbai - 400 080
INDIA

A process for the conversion of echinocandin class of peptides to their C4-homotyrosine monodeoxy analogues

5

This invention relates to a process for the conversion of echinocandin class of peptides of the formula I



(I)

10 wherein W, X, Y, Z, R and R' are as defined herein below :


		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
15	1. Echinocandin B	OH	OH	OH	OH	CH ₃	Linoleoyl
	2. Pneumocandin A ₀	OH	OH	OH	OH	CH ₂ -CONH ₂	10,12-Dimethyl-myristoyl
	3. Pneumocandin A ₁	H	OH	OH	OH	CH ₂ -CONH ₂	"
	4. Pneumocandin A ₂	OH	OH	H	H	CH ₂ -CONH ₂	"
	5. Pneumocandin B ₀	OH	OH	OH	OH	CH ₂ -CONH ₂	"
	6. Pneumocandin B ₂	OH	OH	H	H	CH ₂ -CONH ₂	"
20	7. Pneumocandin C ₀	OH	OH	OH	OH	CH ₂ -CONH ₂	"
	8. Mulundocandin	OH	OH	OH	OH	H	12-Methyl-tetradecanoyl

AP/P/00/01894

to their C4-homotyrosine monodeoxy analogues of the formula I, wherein W, X, Y, Z, R and R' are as defined herein below:

	<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>	
5							
1.	OH	H	OH	OH	CH ₃	Linoleoyl	
	(Echinocandin C)						
2.	OH	H	OH	OH	CH ₂ -CO-NH ₂	10,12-	
	Dimethyl-						
10						myristoyl	
3.	H	H	OH	OH	CH ₂ -CONH ₂	"	
4.	OH	H	H	H	CH ₂ -CONH ₂	"	
5.	OH	H	OH	OH	CH ₂ -CONH ₂	"	
6.	OH	H	H	H	CH ₂ -CONH ₂	"	
15	7.	OH	H	OH	OH	CH ₂ -CONH ₂	"
8.	OH	H	OH	OH	H	12-Methyl tetra-	
	decanoyl,						


particularly to a process for the conversion of mulundocandin (compound of the formula II)



(II)

to deoxymulundocandin (compound of the formula III)

5



(III)

1,3- β -glucan synthesis inhibitors are effective antifungal agents against *Candida albicans* and also *Pneumocystis carini*, an opportunistic organism responsible for an often fatal pneumonitis among HIV patients and other immunocompromised hosts. Of all the structural classes of 1,3- β -glucan synthesis inhibitors, only the echinocandins received considerable attention [Ref : J. Med. Chem. 35, 198-200 (1992)]. Echinocandin class of peptides are cyclic hexapeptides having a lipophilic side chain.

Several methods for the conversion of echinocandins to the corresponding deoxy analogues under acidic conditions have been reported [Ref : Tetrahedron Letts., 33, 4529-4532 (1992); US Patent Appl. No. 222157 dated April 4, 1994]. The above methods involve selective reduction of C4-hyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues with prior protection / deprotection of the equally facile C5-Orn (ornithine) hydroxyl group.

wherein W, X, Y, Z, R and R' are as defined herein below :

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
5	1. Echinocandin B	OH	OH	OH	OH	CH ₃	Linoleoyl
	2. Pneumocandin A ₀	OH	OH	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
	3. Pneumocandin A ₁	H	OH	OH	OH	CH ₂ -CO-NH ₂	"
	4. Pneumocandin A ₂	OH	OH	H	H	CH ₂ -CO-NH ₂	"
10	5. Pneumocandin B ₀	OH	OH	OH	OH	CH ₂ -CO-NH ₂	"
	6. Pneumocandin B ₂	OH	OH	H	H	CH ₂ -CO-NH ₂	"
	7. Pneumocandin C ₀	OH	OH	OH	OH	CH ₂ -CO-NH ₂	"
	8. Mulundocandin	OH	OH	OH	OH	H	12-Methyl- tetradecanoyl

15

to their C4-homotyrosine monodeoxy analogues of the formula I, wherein W, X, Y, Z, R and R' are as defined herein below:

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
20	1. Deoxyechinocandin B (Echinocandin C)	OH	H	OH	OH	CH ₃	Linoleoyl
	2. Deoxypneumocandin A ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
	3. Deoxypneumocandin A ₁	H	H	OH	OH	CH ₂ -CO-NH ₂	"
25	4. Deoxypneumocandin A ₂	OH	H	H	H	CH ₂ -CO-NH ₂	"
	5. Deoxypneumocandin B ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	"
	6. Deoxypneumocandin B ₂	OH	H	H	H	CH ₂ -CO-NH ₂	"
	7. Deoxypneumocandin C ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	"
	8. Deoxymulundocandin	OH	H	OH	OH	H	12-Methyl tetra- decanoyl

30

which consists of a single step selective reduction of C4-htyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues particularly under neutral conditions without prior protection / deprotection of the equally facile C5-Orn (ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.

The conversion of echinocandins to their monodeoxy analogues by selective reduction at C4-htyr may be effected by hydrogenolysis with Raney nickel in solvents such as methanol, ethanol, or dioxane at pH 3-9. Preferably, the selective reduction is carried out by hydrogenolysis with Raney nickel in ethanol at pH 7 and room temperature in the ratio of 6.8 ml Raney nickel per millimole of mulundocandin.

The monodeoxy compounds of the invention may, for example, be purified from the crude reaction mixture as follows :

By fractionation using normal phase chromatography (using alumina or silica gel as stationary phase and eluents such as petroleum ether, ethyl acetate, dichloromethane, chloroform, methanol or combinations thereof), reverse phase chromatography (using reverse phase silica gel like dimethyloctadecylsilylsilica gel, also called RP-18 or dimethyloctylsilylsilica gel also called RP-8 as stationary phase and eluents such as water, buffers such as phosphate, acetate, citrate (pH 2-8) and organic solvents such as methanol, acetonitrile, acetone, tetrahydrofuran or combination of the solvents), gel permeation chromatography - using resins such as "Sephadex LH-20sm" (Pharmacia Chemical Industries, Sweden), TSKgel Toyopearl HW (TosoHaas, Tosoh Corporation, Japan) in solvents such as methanol, chloroform or ethyl acetate or their combination or Sephadex G-10 and G-25 in water; or by counter-current chromatography using a biphasic eluent system made up of two or more solvents such as water, methanol, ethanol, *iso*-propanol, *n*-propanol, tetrahydrofuran, acetone, acetonitrile, methylene chloride, chloroform, ethylacetate, petroleum ether, benzene and toluene. These techniques may be used repeatedly or a combination of the different techniques may be used. Counter-

current chromatography (liquid-liquid⁸ chromatography) using a biphasic eluent system on ITO coil is preferred for purification of the compounds of the invention.

The following experimental example is illustrative of the present invention but not
5 limitative of the scope thereof.

Example 1

Mulundocandin (220 mg, 2.2 mM) in ethanol (8 ml)) was stirred with 15 ml of W-2
Raney nickel (pH 7) in ethanol (30 ml) for 3 hours at room temperature. After
10 standing for 15 minutes the supernatant solution was decanted and Raney nickel
washed with 3 x 30 ml. ethanol with stirring and filtered. Combined ethanolic
solutions were concentrated by distillation under a reduced pressure of 60-70
mm/Hg at 35° C to obtain 160 mg (75%) of crude deoxymulundocandin as a slightly
green solid.

15

The crude product was purified by liquid-liquid chromatography on ITO coil using
upper layer of CH₂Cl₂ : MeOH : *n*-PrOH : H₂O as the stationary phase and the lower
layer as the mobile phase in an ascending mode. The coils (15 + 25 + 215 ml) were
connected in series and a flow rate of 0.6 ml/min. at a piston stroke of 60 and
20 pressure 0.5 bars was maintained. The purification of deoxymulundocandin was
monitored both by bioactivity against *Candida albicans* and *Aspergillus niger* and by
analytical High Pressure Liquid Chromatography (HPLC) [column : (10 x 0.4 cm + 3
x 0.4 cm) ODS-Hypersil, 10μ; mobile phase: 50:50 CH₃CN : H₂O ; flow rate : 1
ml/min; Wavelength : 220 nm.) The fractions (4.5 ml each) containing
25 deoxymulundocandin were combined, concentrated by distillation under a reduced
pressure of 60-70 mm/Hg at 35°C and lyophilized to yield pure
deoxymulundocandin [65 mg (30% yield)]. Also recovered during the above
purification of deoxymulundocandin was unreacted mulundocandin in 10% yield.

The semi-synthetic deoxymulundocandin was identical in all respects to the
30 naturally isolated compound and the physico-chemical data is given in Table 1.

TABLE 1

5


Appearance :	White powder
Melting point:	170-172°C
$[\alpha]_D$:	-36.6° (c 0.25, MeOH)
HPLC RT :	4.42 min
10 FAB-MS (Fast Atom: Bombardment mass)	1014.7 (M + Na) ⁺
¹ H NMR (300 MHz, : CD ₃ OD)	Figure 1 of the accompanying drawings
¹³ C NMR (75 MHz, : 15 CD ₃ OD)	Figure 2 of the accompanying drawings

20

API/00/01894

Claims:

1. A process for the conversion of echinocandin class of peptides of the formula I



(I)

wherein W, X, Y, Z, R and R' are as defined herein below :

10

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
	1.	OH	OH	OH	OH	CH ₃	Linoleoyl
	2.	OH	OH	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl-myristoyl
15	3.	H	OH	OH	OH	CH ₂ -CO-NH ₂	"
	4.	OH	OH	H	H	CH ₂ -CO-NH ₂	"
	5.	OH	OH	OH	OH	CH ₂ -CO-NH ₂	"
	6.	OH	OH	H	H	CH ₂ -CO-NH ₂	"
	7.	OH	OH	OH	OH	CH ₂ -CO-NH ₂	"
20	8.	OH	OH	OH	OH	H	12-Methyl-tetradecanoyl

AP/P/00/01894

to their C4-homotyrosine monodeoxy analogues of the formula I, wherein W, X, Y, Z, R and R' are as defined herein below

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
5	1. Deoxyechinocandin B (Echinocandin C)	OH	H	OH	OH	CH ₃	Linoleoyl
	2. Deoxypneumocandin A ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
	3. Deoxypneumocandin A ₁	H	H	OH	OH	CH ₂ -CONH ₂	"
10	4. Deoxypneumocandin A ₂	OH	H	H	H	CH ₂ -CONH ₂	"
	5. Deoxypneumocandin B ₀	OH	H	OH	OH	CH ₂ -CONH ₂	"
	6. Deoxypneumocandin B ₂	OH	H	H	H	CH ₂ -CONH ₂	"
	7. Deoxypneumocandin C ₀	OH	H	OH	OH	CH ₂ -CONH ₂	"
15	8. Deoxymulundocandin	OH	H	OH	OH	H	12-Methyl tetra- decanoyl

which consists of a single step selective reduction of C4-htyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues under neutral conditions without prior protection / deprotection of the equally facile C5-Orn (ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.

2. A process as claimed in claim 1, wherein Mulundocandin is converted to Deoxymulundocandin.


3. A process as claimed in claims 1 or 2, wherein the reduction reaction is carried out by hydrogenolysis with Raney nickel in ethanol at pH 7 and room temperature.

4. A process as claimed in claims 1 to 3, wherein the hydrogenolysis is carried out in the ratio of 6.8 ml of Raney nickel per millimole of mulundocandin.

AP/P/00/01894

AP001111

1/2



(ppm)

Fig. 1

AP/P/00/01894

AP001111

2/2




Fig. 2

AP/P/00/01894