



(51) International Patent Classification:

A61K 31/47 (2006.01) C07D 217/04 (2006.01)
A61K 51/04 (2006.01) G21H 5/02 (2006.01)

(21) International Application Number:

PCT/US2016/029289

(22) International Filing Date:

26 April 2016 (26.04.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/156,534 4 May 2015 (04.05.2015) US

(71) Applicant: THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; 3160 Chestnut Street, Suite 200, Philadelphia, PA 19104 (US).

(72) Inventors: PRYMA, Daniel, A.; 714 North Ithan Avenue, Bryn Mawr, PA 19104 (US). LIEBERMAN, Brian, P.; 7901 Henry Avenue, D504, Philadelphia, PA 19104 (US). MAKVANDI, Mehran; 633 West Rittenhouse Street, Apt. B520, Philadelphia, PA 19144 (US). MACH, Robert, H.; 124 Dundee Mills Lane, Wallingford, PA 19086 (US).

(74) Agents: CALDWELL, John W. et al.; Baker & Hostetler LLP, Cira Center, 12th Floor, 2929 Arch Street, Philadelphia, PA 19104 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: 211-ASTATINE CONTAINING RADIOTHERAPEUTICS FOR THE TREATMENT OF CANCER



(57) Abstract: Described herein is an alpha-emitting radionuclide, ^{211}At , which has been incorporated into a selective sigma-2 ligand moiety to provide cytotoxic capabilities to a chemical compound. As described herein, a compound of formula (I), wherein R^1 - R^4 , m, and n are defined herein, was prepared and utilized in *in vitro* and *in vivo* tumor targeting of alpha-emitting sigma-2 ligand in a breast cancer model. In one embodiment, the compound is 5-(^{211}At)-N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide.

211-ASTATINE CONTAINING RADIOTHERAPEUTICS FOR THE TREATMENT OF CANCER

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of the priority of US Provisional Patent Application No. 62/156,534, filed May 4, 2015, which is incorporated by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0002] The subject matter disclosed herein was made with government support under grant number DE-SE0012476 awarded by the Department of Energy. The Government has certain rights in the invention.

TECHNICAL FIELD

[0003] This invention relates to the field of ²¹¹-astatine containing radiotherapeutics for treating cancer.


BACKGROUND

[0004] The sigma-2 receptor is a promising target for cancer therapy due to its overexpression in multiple malignancies including: breast, pancreatic, ovarian, lung, renal, melanoma, neuroendocrine and colon. The molecular function of the sigma-2 receptor has been defined through pharmacological studies that have validated it as a biomarker that can be utilized to non-invasively measure the proliferative status of tumors. During cell proliferation the sigma-2 receptor is overexpressed by a factor of 10 compared with quiescent or nonproliferative cells. Confocal microscopy studies have shown the subcellular localization of sigma-2 receptors on cell membrane, lysosomes, endoplasmic reticulum, and mitochondria.

[0005] Sigma-2 ligands, classified as azabicyclononane derivatives, have been shown to undergo rapid internalization and are capable of inducing caspase dependent and independent cell death pathways. More recently, the molecular identity of the sigma-2 receptor was identified as the progesterone receptor membrane component 1 (PGRMC1), providing new insight and potential for further defining its biological, pharmacological and molecular properties. Three

classes of sigma-2 ligands have been characterized pharmacologically through caspase-3/7 activation as agonist, partial agonist or antagonist. Azabicyclononane and tropane analogs induce or partially induce downstream caspase 3/7 activation, and therefore are considered agonist to the sigma-2 receptor. The benzamide class of ligands acts as antagonists, unable to induce caspase activity, and is believed to interact with binding sites on the surface membranes, although passive diffusion across the cell membrane has not been evaluated.

[0006] Sigma-2 antagonists have been historically developed as imaging agents due to their safe toxicity profile and high affinity and selectivity for the sigma-2 receptor, as well as fast tumor targeting and clearance *in vivo*. Numerous selective sigma-2 ligands have been reported, however, *N*-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1*H*)-yl)butyl)-2-(2-F-fluoroethoxy)-5-methylbenzamide is the only diagnostic imaging agent used clinically in a research setting. Other sigma-2 ligands have been applied in chemosensitization and the targeted delivery of cancer therapeutics showing a beneficial therapeutic response in pre-clinical models. Currently, only sigma-2 agonist conjugates have been used as therapeutics. This has been most likely due to the lack of cytotoxicity observed with sigma-2 antagonists and evidence to support whether this class of ligands enters the cell, which is an important characteristic for the therapeutics ability to reach the site of action.




N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1*H*)-yl)butyl)-2-(2-F-fluoroethoxy)-5-methylbenzamide

[0007] What are needed in the art are alternate chemotherapeutics for treating cancer.


SUMMARY OF THE INVENTION

[0008] In one aspect, a compound having the structure of formula (I), or a pharmaceutically acceptable salt or prodrug thereof, is provided, wherein R^1 - R^4 , n , and m are defined herein.




(I)

[0009] In another aspect, the compound has the structure of formula (II), or a pharmaceutically acceptable salt or prodrug thereof, is provided, wherein R¹-R⁴, n, and m are defined herein.




(II)


[0010] In a further aspect, the compound has the structure of formula (III), (IV), (V), (VI), (VII), or (VIII):




(III)




(IV)



(V)



(VI)



(VII)

(VIII)

[0011] In yet another aspect, 5-(²¹¹astatine)-N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide, derivative, salt, or prodrug thereof is provided.

[0012] In still a further aspect, a composition is provided and contains a compound described herein and a pharmaceutically acceptable carrier or diluent.

[0013] In another embodiment, a method of preventing or reducing proliferation of cancer cells is provided and includes contacting the cells with a compound or composition described herein.

[0014] In a further aspect, a method of sensitizing cancer cells to a chemotherapeutic is provided and includes comprising contacting the cells with a compound or composition described herein.

[0015] In still another aspect, a method of modulating expression of the sigma-2-receptor/progesterone receptor membrane component 1 on a cell is provided and includes contacting the cell with a compound or composition described herein.

[0016] In yet a further aspect, a method of treating cancer in a patient is provided and includes administering a compound or composition described herein to the patient.

[0017] In another aspect, a method of administering radiotherapy to a patient in need thereof is provided and includes administering a compound or composition described herein to the patient.

[0018] Other aspects and embodiments of the invention will be readily apparent from the following detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019] The present application is further understood when read in conjunction with the appended drawings. For the purpose of illustrating the subject matter, there are shown in the drawings exemplary embodiments of the subject matter; however, the presently disclosed subject matter is not limited to the specific compositions, methods, devices, and systems disclosed. In addition, the drawings are not necessarily drawn to scale.

[0020] FIG. 1 are examples of ^{211}At -containing compounds encompassed by the structure of formula (I).

[0021] FIG. 2 is a schematic of an apparatus for the dry distillation of ^{211}At astatine.

[0022] FIG. 3 is a graph illustrating *in vitro* sigma-2 receptor binding of ^{211}At -N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide alone (●) or in combination with a standard (⌘) to human-derived breast cancer cells. Error bars are presented as the standard of error measure (SEM). The time of half-maximal binding ($T_{1/2}$) is represented by a dashed line.

[0023] FIG. 4 is a bar graph illustrating the tumor:muscle (■) and tumor:blood ratios (⌘) over a period of 24 h. Error bars are represented as standard deviation (SD).

DETAILED DESCRIPTION OF THE INVENTION

[0024] In the present disclosure the singular forms "a", "an" and "the" include the plural reference, and reference to a particular numerical value includes at least that particular value, unless the context clearly indicates otherwise. Thus, for example, a reference to "a material" is a reference to at least one of such materials and equivalents thereof known to those skilled in the art, and so forth.

[0025] When a value is expressed as an approximation by use of the descriptor "about" or "substantially" it will be understood that the particular value forms another embodiment. In general, use of the term "about" or "substantially" indicates approximations that can vary depending on the desired properties sought to be obtained by the disclosed subject matter and is to be interpreted in the specific context in which it is used, based on its function. The person skilled in the art will be able to interpret this as a matter of routine. In some cases, the number of

significant figures used for a particular value may be one non-limiting method of determining the extent of the word "about" or "substantially". In other cases, the gradations used in a series of values may be used to determine the intended range available to the term "about" or "substantially" for each value. Where present, all ranges are inclusive and combinable. That is, references to values stated in ranges include every value within that range.

[0026] When a list is presented, unless stated otherwise, it is to be understood that each individual element of that list and every combination of that list is to be interpreted as a separate embodiment. For example, a list of embodiments presented as "A, B, or C" is to be interpreted as including the embodiments, "A," "B," "C," "A or B," "A or C," "B or C," or "A, B, or C."

[0027] It is to be appreciated that certain features of the invention which are, for clarity, described herein in the context of separate embodiments, may also be provided in combination in a single embodiment. That is, unless obviously incompatible or excluded, each individual embodiment is deemed to be combinable with any other embodiment(s) and such a combination is considered to be another embodiment. Conversely, various features of the invention that are, for brevity, described in the context of a single embodiment, may also be provided separately or in any sub-combination. It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as "solely," "only" and the like in connection with the recitation of claim elements, or use of a "negative" limitation. Finally, while an embodiment may be described as part of a series of steps or part of a more general structure, each said step may also be considered an independent embodiment in itself.

[0028] Alpha-emitting radionuclides have the potential for new therapeutic drug development and can result in new treatments for clinical utilization to combat cancer.


[0029] I. The Compounds

[0030] ²¹¹Astatine (²¹¹At) is a radionuclide that decays through the emission of a high-energy alpha particle and has a half-life of 7.21 h. Alpha-particles travel approximately 2-3 cell diameters (50-100 μm) and cause dense ionizations along the track resulting in clustered DNA damage capable of inducing cell death. The short path-length of an alpha particle also translates to a highly specific cell killing capability. Cells only within immediate proximity to the

radioactive decay event are affected. It is hypothesized that as few as 10 alpha-particles traversing a cell have a high probability of inducing cell death.

[0031] The compounds discussed herein are alpha-emitting radionuclides. The compounds modulate expression of the sigma-2-receptor/progesterone receptor membrane component 1 on a cell and are, thereby, effective in treating conditions, i.e., cancer, associated with the same. The compounds discussed herein contain a ^{211}At group as a substituent of the molecule. Such compounds are anticipated to be highly effective in treating a number of conditions which require applying internal radiation to a patient. In one embodiment, the compounds discussed herein may have a high therapeutic index. In another embodiment, the compounds have the potential to overcome resistance pathways through direct tumoricidal DNA damage. Accordingly, the compounds discussed herein have use as a radiopharmaceutical. Advantageously, it is anticipated that the cost of ^{211}At and compounds containing same will be less than the cost of commercially available, radioactive ^{123}I .

[0032] The compounds discussed herein have the structure of formula (I) or a pharmaceutically acceptable salt or prodrug thereof.




[0033] In this structure, m is 1 or 2 and n is 1 to 10. In one embodiment, n is 2 to 4.

[0034] R^1 to R^4 are attached to the benzene ring at carbon-atoms the 2-, 3-, 4-, 5-, or 6-position. R^1 to R^4 , independently, occupy any position of the benzene ring, with the proviso that two of R^1 - R^4 and ^{211}At do not occupy the same position on the benzene ring. R^1 to R^4 are, independently, H, C_1 to C_6 alkyl, C_1 to C_6 alkoxy, C_1 to C_6 haloalkyl, halogen, C_1 to C_6 fluoroalkoxy, C_1 to C_6 thioalkyl, C_1 to C_6 thio-haloalkyl, NH_2 , C_1 to C_6 aminoalkyl, or $-(\text{OCH}_2\text{CH}_2)_q\text{-F}$ and q is 2 to 6. In one embodiment, R^1 is C_1 to C_6 fluoroalkoxy. In another embodiment, R^1 is $-\text{O}(\text{CH}_2)_p\text{F}$ and p is 1 to 10. In a further embodiment, the halogen is a radioactive isotope.

[0035] Alternatively, any two of R^1 to R^4 together may form a 5- or 6-membered benzene ring. In one embodiment, R^1 and R^2 ; R^2 and R^3 ; or R^3 and R^4 together form a 5- or 6-membered benzene ring. This benzene ring may be substituted with H or 1 to 4 R^5 . R^5 is H, C_1 to C_6 alkyl, C_1 to C_6 alkoxy, C_1 to C_6 haloalkyl, halogen, C_1 to C_6 fluoroalkoxy, C_1 to C_6 thioalkyl, C_1 to C_6 thio-haloalkyl, NH_2 , C_1 to C_6 aminoalkyl, or $-(OCH_2CH_2)_q-F$, where q is defined above. In one embodiment, the halogen is a radioactive isotope.


[0036] The ^{211}At substituent may be any position, i.e., the 2-, 3-, 4-, 5- or 6-position, of the benzene ring. In one embodiment, the ^{211}At is 3-position of the benzene ring.

[0037] In one embodiment, the compound has the structure of formula (II), wherein R^1 - R^4 , m , and n are defined herein.




(II)

[0038] In another embodiment, the compound has the structure of formula (III), wherein R^1 , R^2 , m , and n are defined herein.




(III)

[0039] In a further embodiment, the compound has the structure of formula (IV), wherein R^1 - R^4 and n are defined herein.




(IV)

[0040] In yet another embodiment, the compound has the structure of formula (V), wherein R^1 - R^4 and n are defined herein.




(V)

[0041] In still a further embodiment, the compound has the structure of formula (VI), wherein R^1 - R^4 and m are defined herein.




(VI)

[0042] In another embodiment, the compound has the structure of formula (VII), wherein R^1 - R^4 are defined herein.



(VII)

[0043] In a further embodiment, the compound has the structure of formula (VIII), wherein R^1 - R^4 and m are defined herein.



(VIII)

[0044] In still another embodiment, the compound is a compound set forth in **Figure 1**, or a derivative, salt, or prodrug thereof.

[0045] In yet a further embodiment, the compound is 5-(²¹¹astatine)-N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide, derivative, salt, or prodrug thereof.

[0046] The term "alkyl" is used herein to refer to both straight- and branched-chain saturated aliphatic hydrocarbon groups. In one embodiment, an alkyl group has 1 to about 10 carbon atoms. In another embodiment, an alkyl group has 1 to about 6 carbon atoms. In a further embodiment, an alkyl group has 1 to about 4 carbon atoms.

[0047] The term "alkoxy" as used herein refers to the O-(alkyl) group, where the point of attachment is through the oxygen-atom and the alkyl group is defined above.

[0048] The term "thioalkyl" as used herein refers to the S-(alkyl) group, where the point of attachment is through the sulfur-atom and the alkyl group is defined above.

[0049] The term "aminoalkyl" as used herein refers to both secondary and tertiary amines where the point of attachment is through the nitrogen-atom and the alkyl groups are defined above. The alkyl groups can be the same or different.

[0050] The term "halogen" as used herein refers to Cl, Br, F, or I groups.

[0051] The term "haloalkyl" as used herein refers to an alkyl group that is substituted at one or more carbon atom of the alkyl chain with a halogen group. The point of attachment of the haloalkyl is through a carbon atom of the alkyl chain.

[0052] The term "fluoroalkoxy" as used herein refers to an alkoxy group that is substituted at one or more carbon atom of the alkyl chain with a fluorine atom. The point of attachment of the fluoroalkoxy is through a carbon atom of the alkoxy chain.

[0053] The term "thio-haloalkyl" as used herein refers to the -S(haloalkyl) group, where the point of attachment is through the sulfur-atom and the haloalkyl group is defined above.

[0054] II. Methods of Production

[0055] The compounds described above may be prepared by known chemical synthesis techniques. These compounds may also be purchased from commercial vendors, e.g., the Sigma-Aldrich Co. Among such preferred techniques known to one of skill in the art are included the synthetic methods described in conventional textbooks relating to the construction of synthetic compounds.

[0056] In one embodiment, ^{211}At is prepared as described by irradiation of a ^{209}Bi bismuth target in a cyclotron through the nuclear reaction $\text{Bi-209}(\alpha, 2n)\text{At-211}$ as described below. In one embodiment, the cyclotron produce alpha beams at energies of about 28.5 MeV. In one embodiment, the cyclotrons and/or systems utilized to prepare the ^{211}At or the compounds described herein would be readily accessible to a facility for treating a patient. In another embodiment, the systems and/or cyclotrons are located no more than about 24 hours from a facility for treating a patient with the compounds discussed herein. In a further embodiment, the systems and/or cyclotrons are located no more than about 12 hours from a facility for treating a patient with the compounds discussed herein. In yet another embodiment, the systems and/or cyclotrons are located no more than about 8 hours from a facility for treating a patient with the compounds discussed herein. In still a further embodiment, the systems and/or cyclotrons are located no more than about 4 hours from a facility for treating a patient with the compounds discussed herein.

[0057] III. Compositions Containing the Compound

[0058] Pharmaceutical compositions useful herein, in one embodiment, contain a compound discussed above in a pharmaceutically acceptable carrier or diluent with other optional suitable pharmaceutically inert or inactive ingredients. In another embodiment, a compound described above is present in a single composition. In a further embodiment, a compound described above is combined with one or more excipients and/or other therapeutic agents as described below.

[0059] (i) Salts

[0060] The compounds discussed above may encompass tautomeric forms of the structures provided herein characterized by the bioactivity of the drawn structures. Further, the compounds may also be used in the form of salts derived from pharmaceutically or physiologically acceptable acids, bases, alkali metals and alkaline earth metals.

[0061] In one embodiment, pharmaceutically acceptable salts can be formed from organic and inorganic acids including, e.g., acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic, malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable acids.

[0062] In another embodiment, pharmaceutically acceptable salts may also be formed from inorganic bases, desirably alkali metal salts including, e.g., sodium, lithium, or potassium, such as alkali metal hydroxides. Examples of inorganic bases include, without limitation, sodium hydroxide, potassium hydroxide, calcium hydroxide, and magnesium hydroxide. Pharmaceutically acceptable salts may also be formed from organic bases, such as ammonium salts, mono-, di-, and trimethylammonium, mono-, di- and triethylammonium, mono-, di- and tripropylammonium, ethyldimethylammonium, benzyldimethylammonium, cyclohexylammonium, benzyl-ammonium, dibenzylammonium, piperidinium, morpholinium, pyrrolidinium, piperazinium, 1-methylpiperidinium, 4-ethylmorpholinium, 1-isopropylpyrrolidinium, 1,4-dimethylpiperazinium, 1 n-butyl piperidinium, 2-methylpiperidinium, 1-ethyl-2-methylpiperidinium, mono-, di- and triethanolammonium, ethyl diethanolammonium, n-butylmonoethanolammonium, tris(hydroxymethyl)methylammonium, phenylmono-ethanolammonium, diethanolamine, ethylenediamine, and the like. In one example, the base is selected from among sodium hydroxide, lithium hydroxide, potassium hydroxide, and mixtures thereof.

[0063] (ii) Prodrugs

[0064] The salts, as well as other compounds, can be in the form of esters, carbamates and other conventional "pro-drug" forms, which, when administered in such form, convert to the active moiety *in vivo*. In one embodiment, the prodrugs are esters. In another embodiment, the prodrugs are carbamates. See, e.g., B. Testa and J. Caldwell, "Prodrugs Revisited: The "Ad Hoc" Approach as a Complement to Ligand Design", Medicinal Research Reviews, 16(3):233-241, ed., John Wiley & Sons (1996), which is incorporated by reference.

[0065] (iii) Carriers and Diluents

[0066] The pharmaceutical compositions include a compound described herein formulated neat or with one or more pharmaceutical carriers for administration, the proportion of which is

determined by the solubility and chemical nature of the compound, chosen route of administration and standard pharmacological practice. The pharmaceutical carrier may be solid or liquid.

[0067] The compound may be administered to a subject by any desirable route, taking into consideration the specific condition for which it has been selected. The compound may, therefore, be delivered orally, by injection, i.e., transdermally, intravenously, subcutaneously, intramuscularly, intravenous, intra-arterial, intraperitoneal, intracavitary, or epidurally, among others.

[0068] Although the compound may be administered alone, it may also be administered in the presence of one or more pharmaceutical carriers that are physiologically compatible. The carriers may be in dry or liquid form and must be pharmaceutically acceptable. Liquid pharmaceutical compositions are typically sterile solutions or suspensions.

[0069] When liquid carriers are utilized, they are desirably sterile liquids. Liquid carriers are typically utilized in preparing solutions, suspensions, emulsions, syrups and elixirs. In one embodiment, the compound is dissolved a liquid carrier. In another embodiment, the compound is suspended in a liquid carrier. One of skill in the art of formulations would be able to select a suitable liquid carrier, depending on the route of administration. In one embodiment, the liquid carrier includes, without limitation, water, organic solvents, oils, fats, or mixtures thereof. In another embodiment, the liquid carrier is water containing cellulose derivatives such as sodium carboxymethyl cellulose. In a further embodiment, the liquid carrier is water and/or dimethylsulfoxide. Examples of organic solvents include, without limitation, alcohols such as monohydric alcohols and polyhydric alcohols, e.g., glycols and their derivatives, among others. Examples of oils include, without limitation, fractionated coconut oil, arachis oil, corn oil, peanut oil, and sesame oil and oily esters such as ethyl oleate and isopropyl myristate.

[0070] Alternatively, the compound may be formulated in a solid carrier. In one embodiment, the composition may be compacted into a unit dose form, i.e., tablet or caplet. In another embodiment, the composition may be added to unit dose form, i.e., a capsule. In a further embodiment, the composition may be formulated for administration as a powder. The solid carrier may perform a variety of functions, i.e., may perform the functions of two or more of the excipients described below. For example, the solid carrier may also act as a flavoring agent, lubricant, solubilizer, suspending agent, filler, glidant, compression aid, binder, disintegrant, or

encapsulating material. Suitable solid carriers include, without limitation, calcium phosphate, dicalcium phosphate, magnesium stearate, talc, starch, sugars (including, e.g., lactose and sucrose), cellulose (including, e.g., microcrystalline cellulose, methyl cellulose, sodium carboxymethyl cellulose), polyvinylpyrrolidone, low melting waxes, ion exchange resins, and kaolin. The solid carrier can contain other suitable excipients, including those described below.

[0071] Examples of excipients which may be combined with the compound include, without limitation, adjuvants, antioxidants, binders, buffers, coatings, coloring agents, compression aids, diluents, disintegrants, emulsifiers, emollients, encapsulating materials, fillers, flavoring agents, glidants, granulating agents, lubricants, metal chelators, osmo-regulators, pH adjustors, preservatives, solubilizers, sorbents, stabilizers, sweeteners, surfactants, suspending agents, syrups, thickening agents, or viscosity regulators. See, the excipients described in the "Handbook of Pharmaceutical Excipients", 5th Edition, Eds.: Rowe, Sheskey, and Owen, APhA Publications (Washington, DC), December 14, 2005, which is incorporated herein by reference.

[0072] IV. Methods of Using the Compound

[0073] Since the compounds described herein, and compositions containing same, contain ²¹¹At, they are radioactive and, thereby effective in radiotherapy applications in a patient.

[0074] The terms "patient" or "subject" as used herein refer to a mammalian animal. In one embodiment, the patient or subject is a human. In another embodiment, the patient or subject is a veterinary or farm animal, a domestic animal or pet, or animal normally used for clinical research. In still a further embodiment, the subject or patient has cancer. The subject or patient has either been recognized as having or at risk of having cancer.

[0075] As noted above, the compounds here are capable of modulating expression of the sigma-2-receptor/progesterone receptor membrane component 1 on a cell. In one embodiment, the cell is a cancer cell. The compounds and compositions prevent and/or reduce the proliferation of cancer cells thereby curing a patient or putting a patient in remission. Accordingly, the compounds discussed herein are useful in treating cancer by contacting the cells with a compound or composition discussed herein.

[0076] As used herein, "treatment" encompasses treatment of a subject clinically diagnosed as having a disease or medical condition. In one embodiment, the subject is treated and the disease

or medical condition is eradicated, i.e., the subject is cured. As used herein, "prevention" encompasses prevention of symptoms in a subject who has been identified as at risk for the condition, but has not yet been diagnosed with the same and/or who has not yet presented any symptoms thereof.

[0077] The term "cancer" as used herein, refers to neoplastic cells in a patient which have abnormal cell group and invade or have the potential to invade one or more body parts of the patient. In one embodiment, the cancer is a neuroendocrine cancer. In another embodiment, the cancer is of the adrenal gland, appendix, bladder, blood, brain, bone, breast, bronchus, central nervous system, cervix, chest, colon, esophagus, eye, gallbladder, head, intestines, kidney, larynx, liver, lung, lymph nodes, mouth, neck, ovaries, pancreas, pharynx, pituitary, prostate, rectum, skin, stomach, testicles, throat, thymus, thyroid, uterus, urinary tract, or vagina, or is a leukemia. In a further embodiment, the cancer is breast cancer.

[0078] The compounds are also useful in sensitizing a cancer to treatment with a chemotherapeutic. In doing so, the compounds weaken some or all of the cancer cells to apoptosis by another chemotherapeutic agent or radiation. Alternatively, the compounds kill some of the cancer cells and a second chemotherapeutic or radiation may be utilized to kill the remaining cancer cells.

[0079] As described herein, a therapeutically or prophylactically effective amount of a compound is that amount of a compound which provides a sufficient amount of radiation. The sufficient amount of radiation may vary depending upon the formulation and route of delivery. In one embodiment, the amount (i.e., per unit) of the compound is that which does not exceed normal organ dose limits and delivers a tumoricidal dose to cancer cells. In one embodiment, the dose of the compound is dependent on the specific organ and cancer being treated. In another embodiment, the dose of the compound is the maximum dose tolerated by the patient. In a further embodiment, the compound delivers about 0.01 to about 100 mCi of radiation. In still another embodiment, the compound delivers about 0.05 to about 75 mCi of radiation. In still a further embodiment, the compound delivers about 0.1 to about 30 mCi of radiation. However, the effective amount to be used is subjectively determined by the attending physician and variables such as the size, age and response pattern of the patient.

[0080] These effective amounts may be provided on regular schedule, i.e., daily, weekly, monthly, or yearly basis or on an irregular schedule with varying administration days, weeks,

months, etc. Alternatively, the effective amount to be administered may vary. In one embodiment, the effective amount for the first dose is higher than the effective amount for one or more of the subsequent doses. In another embodiment, the effective amount for the first dose is lower than the effective amount for one or more of the subsequent doses.

[0081] The methods described herein may be performed by administering a compound described herein via a combination therapy in prior to, concurrently with, or subsequent to another medication such as a chemotherapeutic. Such combination treatment may occur by administering compositions containing multiple active ingredients, as described above. However, also encompassed is a method of administration of chemotherapeutics in conjunction with a composition containing a compound described herein. In one embodiment, the compound and chemotherapeutic are administered to the patient by one or more selected routes of administration sequentially. In another embodiment, a chemotherapeutic agent is administered before treatment with a compound described herein. In another embodiment, a chemotherapeutic agent is administered after treatment with a compound described herein. In still another embodiment, a chemotherapeutic agent is administered during treatment with a compound described herein.

[0082] In one embodiment, a method of preventing or reducing proliferation of cancer cells is provided and includes contacting the cells with a compound or composition described herein.

[0083] In a further embodiment, a method of treating cancer in a patient is provided and includes administering a compound or composition described herein to the patient.

[0084] In another embodiment, a method of sensitizing cancer cells to a chemotherapeutic is provided and includes contacting the cells with a compound or composition described herein.

[0085] In yet a further embodiment, a method of administering radiotherapy to a patient in need thereof is provided and includes administering a compound or composition to the patient.

[0086] V. Kits Containing the Compound

[0087] Also provided herein are kits or packages of pharmaceutical formulations containing a compound or composition described herein. The kits may be organized to indicate a single formulation or combination of formulations to be taken at each desired time. The composition may also be sub-divided to contain appropriate quantities of the compound. For example, the

unit dosage can be packaged compositions, e.g., packeted powders, vials, ampoules, prefilled syringes or sachets containing liquids.

[0088] Suitably, the kit contains packaging or a container with the compound formulated for the desired delivery route. Suitably, the kit contains instructions on dosing and an insert regarding the compound. Optionally, the kit may further contain instructions for monitoring circulating levels of product and materials for performing such assays including, e.g., reagents, well plates, containers, markers or labels, and the like. Such kits are readily packaged in a manner suitable for treatment of a desired indication. For example, the kit may also contain instructions for use of the delivery device. Other suitable components to include in such kits will be readily apparent to one of skill in the art, taking into consideration the desired indication and the delivery route. The doses are repeated daily, weekly, or monthly, for a predetermined length of time or as prescribed.

[0089] The compound or composition described herein can be a single dose or for continuous or periodic discontinuous administration. For continuous administration, a package or kit can include the compound in each dosage unit (e.g., solution, lotion, tablet, pill, or other unit described above or utilized in drug delivery). When the compound is to be delivered with periodic discontinuation, a package or kit can include placebos during periods when the compound is not delivered. When varying concentrations of a composition, or of the components of the composition, or of relative ratios of the compound or other agents within a composition over time is desired, a package or kit may contain a sequence of dosage units, so varying.

[0090] A number of packages or kits are known in the art for the use in dispensing pharmaceutical agents for oral use. In one embodiment, the package has indicators for each period. In another embodiment, the package is a labeled blister package, dial dispenser package, or bottle.

[0091] The packaging means of a kit may itself be geared for administration, such as an inhalant, syringe, pipette, eye dropper, or other such like apparatus, from which the formulation may be applied to an infected area of the body, such as the lungs, injected into a subject, or even applied to and mixed with the other components of the kit.

[0092] The compound or composition of these kits also may be provided in dried or lyophilized forms. When reagents or components are provided as a dried form, reconstitution

generally is by the addition of a suitable solvent. It is envisioned that the solvent also may be provided in another packaging means.

[0093] The kits may include a means for containing the vials in close confinement for commercial sale such as, e.g., injection or blow-molded plastic containers into which the desired vials are retained.

[0094] Irrespective of the number or type of packages, the kits also may include, or be packaged with a separate instrument for assisting with the injection/administration or placement of the ultimate complex composition within the body of an animal. Such an instrument may be an inhalant, syringe, pipette, forceps, measuring spoon, eye dropper or any such medically approved delivery means. Other instrumentation includes devices that permit the reading or monitoring of reactions *in vitro*.

[0095] In one embodiment, a pharmaceutical kit is provided and contains a compound of formula (I). The compound may be in the presence or absence of one or more of the carriers or excipients described above. The kit may optionally contain a chemotherapeutic and/or instructions for administering the chemotherapeutic and the compound to a subject having cancer.

[0096] In a further embodiment, a pharmaceutical kit is provided and contains a chemotherapeutic in a first dosage unit, one or more of a compound selected from those described herein in a second dosage unit, and one or more of the carriers or excipients described above in a third dosage unit. The kit may optionally contain instructions for administering the chemotherapeutic and/or compound to a subject having cancer.

[0097] The following Examples are provided to illustrate some of the concepts described within this disclosure. While each Example is considered to provide specific individual embodiments of composition, methods of preparation and use, none of the Examples should be considered to limit the more general embodiments described herein.

[0098] In the following examples, efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental error and deviation should be accounted for. Unless indicated otherwise, temperature is in degrees C, pressure is at or near atmospheric.

[0099] EXAMPLES**[00100] Example 1: ^{211}At Production and Distillation**

[00101] ^{211}At was produced via the nuclear reaction $^{209}\text{Bi}(\alpha, 2n)^{211}\text{At}$, on a Japan Steel Works BC3015 cyclotron using external solid target (8) positioned perpendicular to the beam line. The solid target (8) diameter was 1.6 cm and consisted of a 75 μm thick ^{209}Bi layer backed by 1100 aluminum alloy. The beam energy was degraded from 30 to 28.4 MeV by using a 75 μm thick aluminum degrader foil attached directly to the ^{209}Bi surface. A defocused beam profile was used to spread the beam across a larger surface area to mitigate pinpoint heating and subsequent melting due to the poor thermal properties of ^{209}Bi and the perpendicular incident beam angle.

[00102] Dry distillation was performed to isolate ^{211}At from the solid target (8). See, **Figure 2**. The foil was removed from solid target (8) and the backing with $^{209}\text{Bi}/^{211}\text{At}$ layer was cleaned to remove any thermal paste using alcohol wipes, hexane and acetone. The cleaned and dried target was placed in a quartz boat and was positioned inside the quartz furnace tube (7). The entire distillation apparatus was assembled and vacuum initiated through vacuum outlet (1). The apparatus also contained a vacuum trap (3) to trap extraneous liquids, a charcoal trap (2) to prevent volatilized ^{211}At being discharged, and a liquid NaOH trap (5) to collect the ^{211}At . Once the entire system was evacuated, a slow argon gas stream was turned on from argon inlet (10) along with furnace (6). A vent (9) was attached to the argon inlet (10) to for modulation of the pressure. Furnace (6), which is adjacent to the quartz furnace tube, was adjusted to a maximum temperature of 700 °C and was held for 10 minutes. Radioactivity in the collection vessel (5) was monitored in real time with dose calibrator (4; Capintec, Inc, Ramsey NJ) using the ^{133}Xe setting and a 2.3 conversion factor to calculate ^{211}At radioactivity.


[00103] ^{211}At production yields up to 340.4 MBq were achieved and are presented in Table 1. Dry distillation of ^{211}At resulted in decay corrected recovery yields from 50-89%, median=60% (Table 1). Monitoring the collection vessel radioactivity in real time provided instant feedback of the distillation process and helped adequately fine tune the parameters of temperature and argon flow. The entire distillation process lasted about 45 min.

[00104] Table 1: ^{211}At production and distillation - distillation yield presented as recovered radioactivity from measured solid target.

Average Current (μA)	$\mu\text{A} \cdot \text{h}$	Production Yield ($\text{MBq}/\mu\text{A} \cdot \text{h}$)	Distillation Yield (MBq)	Decay Corrected Distillation Yield (%)
10	30.1	10.4	238	89
10	20	9.8	107	61
6.8	20.1	9.2	104	64
9.9	13.3	8.6	70	66
7.5	16.95	7.5	85	76
9	45.4	8.7	196	58

[00105] Example 2: Precursor and Standard Synthesis

[00106] The precursor, *i.e.*, N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxy-5-(tributylstannyl)benzamide, and standard, *i.e.*, 5-iodo-N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide, were synthesized as described in Hou, "Characterization of a novel iodinated sigma-2 receptor ligand as a cell proliferation marker," Nuclear medicine and biology, 2006; 33(2):203-209 and Tu, "Fluorine-18-labeled benzamide analogues for imaging the sigma-2 receptor status of solid tumors with positron emission tomography," Journal of Medicinal Chemistry, 2007; 50(14):3194-3204, which documents are incorporated by reference.



standard

[00107] Example 3: Synthesis of ^{211}At -N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide

[00108] The title compound was synthesized by electrophilic radioastatodestannylation. Briefly, 0.1 M NaOH/ ^{211}At (1 mL) was added to precursor (100 μg), followed by 3:1, glacial acetic acid: hydrogen peroxide (100 μg). The reaction was vortexed periodically over 20 minutes, then quenched with ammonium formate (AMF; 3.5 mL) pH 4.5 and purified by High Performance Liquid Chromatography (HPLC) on a reversed phase C-18 Phenomenex® column (250 x 10mm, 5 μm). Isocratic mobile phase of 30:70 acetonitrile: 0.1 M AMF pH 4.5 was used

with UV wavelength of 288 nm and flow rate of 4 mL/min. The radioactive peak was collected at 23 minutes, diluted to 50 mL with water, and then concentrated on C-18 light Sep-Pak® cartridge (Waters, Milford MA). Product was eluted from cartridge using 200 proof ethanol in a volume of 100 µL. Quality control was performed using the same mobile phase as above on a reversed phase C-18 Agilent Zorbax® (4.6x150 mm, 5 µM) analytical column. Co-injection of the standard was used to identify ²¹¹At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide. There was no UV mass observed associated with this product so the specific activity of the title compound was assumed to be the same as the theoretical for ²¹¹At (435,000 curies/mmol).

[00109] ²¹¹At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide was synthesized with average recovery yield of 70% and is highly comparable to similar iodination chemistry. Radiochemical purity was >95% with no observable UV mass. Co-injection of the title compound and non-radioactive standard resulted in UV and radioactive peak elution at similar retention times.

[00110] Radiosynthesis of ²¹¹At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide was reproducible with high radiochemical purity.

[00111] Example 4: Further Compounds

[00112] The compounds of Figure 1 may be prepared using the synthetic steps of Example 3 and US Patent No. 7,390,902 which is incorporated by reference.

[00113] Example 5: *In Vitro* Binding Studies

[00114] ²¹¹At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide was evaluated in MDA-MB-231 breast cancer cells (ATCC-HTB-26) known to have high sigma-2 receptor density and compared with the well-characterized sigma-2 selective ligand standard. MDA-MB-231 were cultured in Minimum Essential Media (MEM) with added nutrients, 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin at 5% CO₂ and 37 °C. Twenty-four hours before assay cells were plated at 50,000 cell/well in 96-well format. Two solutions of freshly synthesized radioligand were diluted to a concentration of approximately 2,000 cpm/µL in phosphate buffered saline with calcium and magnesium (PBS w/ Ca²⁺ and Mg²⁺), with and without non-radioactive 100 µM of the standard to determine non-

specific binding. At the time of assay, culture media was removed and $^{211}\text{At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide}$ or $^{211}\text{At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide}$ + 100 μM of the standard was added to the wells. Total volume of treatment in each well was 50 μL . PBS w/ Ca^{2+} and Mg^{2+} without substrate was added to the last row of wells and was used for protein quantification at the end of the experiment. Treatment was removed at time points over the course of 2 h ($n=4/\text{time point}$). At each time of treatment removal, wells were washed with PBS x 3. At the end of the experiment wells were counted on an automatic gamma counter using a custom protocol with an energy window of 70-110 KeV (WizardTM 2470, Perkin Elmer, Waltham MA). Protein quantification was performed using a modified Lowry method. Independent experiments were repeated in triplicate. One-phase association kinetics were modeled using PrismTM software version 6.0 (La Holla, CA).

[00115] Live cellular binding studies with $^{211}\text{At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide}$ in MDA-MB-231 cells resulted in sigma-2 binding and demonstrated retained affinity to tumor cells. Fast kinetics were observed with 50% maximal binding occurring at 12.51 min. Selective sigma-2 competitive inhibition was shown with an 80% reduction in total binding of $^{211}\text{At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide}$ through the addition of the standard. **Figure 3** represents data from 3 independent experiments. Average protein quantification for experiments was $7.4 \pm 0.75 \mu\text{g}$.

[00116] Example 6: *In Vivo* Biodistribution

[00117] Tumors were grown by subcutaneously injecting 1×10^6 EMT-6 mouse breast cancer cells (ATCC #CRL-2755) into the flanks of female BALB/c mice (Taconic). Tumors were an appropriate size of approximately 200 mm^3 after 14 days of growth. On the day of experiment 10 μCi of newly synthesized $^{211}\text{At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide}$ in saline (ethanol <5%) was administered intravenously to mice under isoflurane anesthesia. Mice were sacrificed by cervical dislocation followed by immediate cardiac puncture post administration at time points of 5 min, 1, 2, 4, 8, and 24 h ($n=4$ mice/time point). Tissue harvested included; blood, heart, muscle, lung, kidney, pancreas, spleen, liver, skin, brain, thyroid, stomach and EMT-6 tumor. Tissues were weighed and counted on an automatic gamma counter as previously described. All animal experiments were conducted

under IACUC-approved protocols compliant with University of Pennsylvania guidelines for the care and use of research animals.

[00118] Biodistribution was evaluated over 24 h showing high initial kidney uptake at 5 min (47.7% ID/g) with rapid clearance over 1 h (5.83% ID/g). Similar results were also seen, although to a lesser extent, in the liver. Tumor accumulation of $^{211}\text{At-N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide}$ remained above 4% ID/g from 1 h to at least 8 h. At 24 h, the majority of decay corrected radioactivity was cleared from all tissues. Considerable uptake was noted in the stomach, thyroid and pancreas. High uptake of radioactivity in the stomach and thyroid might be due to *in vivo* deastatination. Future metabolite studies are needed to verify this and define what percentage of deastatination occurs. The high uptake in the pancreas is expected due to the high expression of the sigma-2 receptor. Biodistribution data is presented in Table 2 and tumor:muscle/blood ratios are presented in **Figure 4**.

[00119] Table 2: Decay corrected biodistribution data presented as percent injected dose/gram (%ID/g).

	5 min (N=4)	1 h (N=4)	2 h (N=4)	4 h (N=4)	8 h (N=4)	24 h (N=4)
Blood	4.06 ± 0.19	2.12 ± 0.15	2.43 ± 0.82	1.64 ± 0.11	1.82 ± 0.10	0.44 ± 0.10
Heart	10.2 ± 0.82	5.16 ± 1.21	4.70 ± 0.31	3.54 ± 0.17	4.01 ± 0.29	1.14 ± 0.27
Muscle	0.69 ± 0.16	1.24 ± 0.18	1.15 ± 0.07	0.56 ± 0.03	0.61 ± 0.04	0.19 ± 0.01
Lung	20.2 ± 1.39	7.14 ± 1.03	6.42 ± 0.39	5.74 ± 0.53	6.23 ± 0.57	2.21 ± 0.40
Kidney	47.7 ± 2.32	5.83 ± 0.53	5.59 ± 0.35	2.97 ± 0.13	3.24 ± 0.21	1.05 ± 0.23
Pancreas	8.81 ± 1.21	18.1 ± 1.60	18.0 ± 1.65	26.3 ± 2.11	21.1 ± 2.35	5.78 ± 0.89
Spleen	8.41 ± 2.67	13.2 ± 2.76	12.0 ± 1.47	7.59 ± 0.49	7.49 ± 1.38	2.14 ± 1.02
Liver	18.1 ± 2.11	6.39 ± 3.27	4.63 ± 1.01	1.87 ± 0.17	1.86 ± 0.19	0.59 ± 0.20
Skin	0.58 ± 0.05	2.30 ± 0.23	2.66 ± 0.30	1.79 ± 0.15	2.10 ± 0.42	0.85 ± 0.08
Brain	1.95 ± 0.14	0.46 ± 0.08	0.39 ± 0.01	0.21 ± 0.02	0.25 ± 0.02	0.07 ± 0.02
Thyroid	5.87 ± 0.85	11.6 ± 1.01	18.3 ± 6.12	14.0 ± 4.26	13.3 ± 2.96	4.87 ± 1.15
Stomach	5.87 ± 1.12	15.9 ± 2.53	16.3 ± 2.29	19.7 ± 1.79	28.2 ± 7.21	9.88 ± 1.62
Tumor	0.74 ± 0.07	4.40 ± 0.15	4.93 ± 0.78	5.05 ± 0.68	5.11 ± 0.09	1.34 ± 0.37

[00120] The results show retention of radioactivity in the tumor from 1-8 h post administration. The iodinated sigma-2 ligand standard was shown to have tumor clearance by 4 h with a lower tumor uptake of 2% ID/g. Tumor retention of $^{211}\text{At-N-(6,7-dimethoxy-3,4-}$

dihydroisoquinolin-2(1H-yl)butyl)-2,3-dimethoxybenzamide is highly desirable as a therapeutic and provides a tumor residence time approximately equal to the half-life of ^{211}At ($t_{1/2}=7.21$ h) providing at least 50% of total possible radiation dose to the tumor before washout. The results also illustrate a maximal tumor to muscle ratio of 9.02 at 4 h (**Figure 4**).


[00121] *In vivo* deastatination is the suspected cause of off-target localization in the thyroid and stomach. It is hypothesized that off-target localization might be due to sigma-2 receptor expression in the thyroid and stomach. The pancreas also had an appreciable amount of radioactivity localized. It is hypothesized that there is a known high expression of sigma-2 receptors on endocrine glands. At 24 h, most of the decay corrected ^{211}At -N-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H-yl)butyl)-2,3-dimethoxybenzamide was cleared from all tissues.

[00122] It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof, that the foregoing description and the examples that follow are intended to illustrate and not limit the scope of the invention. It will be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the scope of the invention, and further that other aspects, advantages and modifications will be apparent to those skilled in the art to which the invention pertains. In addition to the embodiments described herein, the present invention contemplates and claims those inventions resulting from the combination of features of the invention cited herein and those of the cited prior art references which complement the features of the present invention. Similarly, it will be appreciated that any described material, feature, or article may be used in combination with any other material, feature, or article, and such combinations are considered within the scope of this invention.

[00123] The disclosures of each patent, patent application, and publication cited or described in this document are hereby incorporated herein by reference, each in its entirety, for all purposes.

What is Claimed Is:

1. A compound having the structure of formula (I):



wherein:

m is 1 or 2;

n is 1 to 10;

R^1 to R^4 are at the 2-, 3-, 4-, 5-, or 6-position of the benzene ring and, independently, H, C_1 to C_6 alkyl, C_1 to C_6 alkoxy, C_1 to C_6 haloalkyl, halogen, C_1 to C_6 fluoroalkoxy, C_1 to C_6 thioalkyl, C_1 to C_6 thio-haloalkyl, NH_2 , C_1 to C_6 aminoalkyl, or $-(OCH_2CH_2)_q-F$;

q is 2 to 6;

or R^1 and R^2 ; R^2 and R^3 ; or R^3 and R^4 together form a 5- or 6-membered benzene ring substituted with H or 1 to 4 R^5 ;


^{211}At is at the 2-, 3-, 4-, 5- or 6-position of the benzene ring; and

R^5 is H, C_1 to C_6 alkyl, C_1 to C_6 alkoxy, C_1 to C_6 haloalkyl, halogen, C_1 to C_6 fluoroalkoxy, C_1 to C_6 thioalkyl, C_1 to C_6 thio-haloalkyl, NH_2 , C_1 to C_6 aminoalkyl, or $-(OCH_2CH_2)_q-F$;

with the proviso that two of R^1-R^4 and ^{211}At do not occupy the same position on said benzene ring;

or a pharmaceutically acceptable salt or prodrug thereof.


2. The compound of claim 1, having the structure of formula (II):



3. The compound of claim 1, wherein said halogen is a radioactive isotope.


4. The compound of claim 1, wherein R^1 is C_1 to C_6 fluoroalkoxy.

5. The compound of claim 3, wherein R^1 is $O(CH_2)_pF$ and p is 1 to 10.
6. The compound of claim 1, wherein n is 2 to 4.
7. The compound of claim 1, having the structure of formula (III):




(III).

8. The compound of claim 1, having the structure of formula (IV):




(IV).

9. The compound of claim 1, having the structure of formula (V):




(V).

10. The compound of claim 1, having the structure of formula (VI):




(VI).

11. The compound of claim 1, having the structure of formula (VII):




(VII).


12. The compound of claim 1, having the structure of formula (VIII):





(VIII).

13. The compound of claim 1, which is:










14. A compound which is 5-(211 astatine)-N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)butyl)-2,3-dimethoxybenzamide, derivative, salt, or prodrug thereof.
15. A composition comprising a compound of any one of claims 1 to 14 and a pharmaceutically acceptable carrier or diluent.
16. A method of preventing or reducing proliferation of cancer cells, said method comprising contacting said cells with a compound of any one of claims 1 to 14.
17. A method of sensitizing cancer cells to a chemotherapeutic, said method comprising contacting said cells with a compound of any one of claims 1 to 14.
18. The method of claim 17, further comprising administering radiation to said cells.
19. A method of modulating expression of the sigma-2-receptor/progesterone receptor membrane component 1 on a cell, said method comprising contacting said cell with a compound of any one of claims 1 to 14.
20. The method of claim 19, further comprising administering radiation to said cell.
21. A method of treating cancer in a patient, said method comprising administering a compound of any one of claims 1 to 14 to said patient.
22. The method of claim 21, wherein said cancer is a neuroendocrine cancer.
23. The method of claim 21, wherein said cancer is of the prostate, head, neck, eye, mouth, throat, esophagus, bronchus, larynx, pharynx, chest, bone, lung, colon, rectum, stomach, bladder, uterus, cervix, breast, ovaries, vagina, testicles, thyroid, blood, lymph nodes, kidney, liver, intestines, pancreas, brain, central nervous system, adrenal gland, neuroendocrine, or skin or a leukemia.
24. The method of claim 21, wherein said cancer is breast cancer.
25. The method of claim 21, further comprising administering radiation to said patient.

26. The method of claim 21, wherein said compound is administered to said patient through intravenous, intra-arterial, intraperitoneal or intracavitary injection.
27. The method of claim 21, comprising administering an amount of said compound having from about 0.01 to about 100 Curies of radiation.
28. A method of administering radiotherapy to a patient in need thereof, said method comprising administering a compound of any one of claims 1 to 14 to said patient.
29. The method of claim 28, further comprising administering radiation to said patient.
30. The method of claim 28, wherein said compound is administered to said patient through intravenous, intra-arterial, intraperitoneal or intracavitary injection.
31. The method of claim 28, comprising administering an amount of said compound having from about 0.01 to about 100 Curies of radiation.

**FIG. 1**

2/6




FIG. 1 (cont.)

3/6







FIG. 1 (cont.)



1. Vacuum Outlet
2. Charcoal Traps
3. Liquid Vacuum Trap
4. Dose Calibrator
5. Liquid NaOH Trap
6. Furnace
7. Quartz Furnace Tube
8. Solid Target
9. Vent
10. Argon Inlet

FIG. 2

**FIG. 3**

**FIG. 4**

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 16/29289

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/47; 51/04; C07D 217/04; G21H 5/02 (2016.01)

CPC - A61K51/04; G21H5/04; A61K31/47; C07D217/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): A61K 31/47; 51/04; C07D 217/04; G21H 5/02 (2016.01)

CPC: A61K51/04; G21H5/04; A61K31/47; C07D217/04

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC: 424/1.81; 424/1.11

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatBase, Google Scholar, PubWEST

Radionuclei, astatine-211, imaging, cancer/tumor, sigma-2 ligand, 5-(211astatine)-N-(4-(6,7-dimethoxy-3,4-dihydroisoquinolin-2(1H-yl)butyl)-2,3-dimethoxybenzamide, 5-bromo, neuroendocrine cancer, breast cancer, intravenous, radiation

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	US 2005/0107398 A1 (MACH et al.) 19 May 2005 (19.05.2005) para [0013], [0021], [0097], [0102], [0162], [0165] astatine-211	1-7, 9-10, 12-14, (15-16, 19, 21-24, 26-28, 30-31)/(1-7,9-10,12-14)
Y	FAN et al. 'Effect of structural modification in the amine portion of substituted aminobutylbenzamides as ligands for binding sigma1 and sigma2 receptors', Bioorg Med Chem. 2011, Vol.19(6), pp 1852-1859. abstract; pg 13, Fig 2, compound 3; pg 17, Table 1	8, 11, (15-16, 19, 21-24, 26-28, 30-31)/(8,11), 17-18, 20, 25, 29
Y	US 2009/0176705 A1 (MCDUNN et al.) 09 July 2009 (09.07.2009) abstract; para [0010]	8, 11, (15-31)/(8,11)
A	US 2014/0378460 A1 (CATALANO et al.) 25 December 2014 (25.12.2014) Entire Document	17-18, 20, 25, 29
A	US 2014/0378460 A1 (CATALANO et al.) 25 December 2014 (25.12.2014) Entire Document	1-31
A	US 5,919,934 A (JOHN et al.) 06 July 1999 (06.07.1999) Entire Document	1-31

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

15 June 2016 (15.06.2016)

Date of mailing of the international search report

27 JUL 2016

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774