




(86) **Date de dépôt PCT/PCT Filing Date:** 2009/03/23
(87) **Date publication PCT/PCT Publication Date:** 2009/10/01
(45) **Date de délivrance/Issue Date:** 2015/12/08
(85) **Entrée phase nationale/National Entry:** 2010/09/20
(86) **N° demande PCT/PCT Application No.:** IN 2009/000192
(87) **N° publication PCT/PCT Publication No.:** 2009/118759
(30) **Priorité/Priority:** 2008/03/26 (IN746/CHE/2008)

(51) **Cl.Int./Int.Cl.** **C07D 473/06** (2006.01),
A61K 31/522 (2006.01), **A61P 11/06** (2006.01),
A61P 3/10 (2006.01), **A61P 9/12** (2006.01),
C07D 473/08 (2006.01)

(72) **Inventeurs/Inventors:**
PALLE, VENKATA, IN;
BASU, SUJAY, IN;
WAMAN, YOGESH, IN;
RAMDAS, VIDYA, IN;
BARAWKAR, DINESH, IN;
PATEL, MEENA, IN;
PANMAND, ANIL, IN

(73) **Propriétaire/Owner:**

(54) **Titre : COMPOSES HETEROCYCLIQUES UTILES COMME ANTAGONISTES DES RECEPTEURS ADENOSINIQUES**
(54) **Title: HETEROCYCLIC COMPOUNDS AS ADENOSINE RECEPTOR ANTAGONIST**



I

(57) **Abrégé/Abstract:**

Compounds of the present disclosure are fused pyrimidine compounds of formula (I), its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof, as Adenosine receptor antagonists. Processes of their preparation are also described in the disclosure.



(73) **Propriétaires(suite)/Owners(continued):**ADVINUS THERAPEUTICS PVT. LTD., IN

(74) **Agent:** SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
1 October 2009 (01.10.2009)

PCT

(10) International Publication Number
WO 2009/118759 A3

(51) International Patent Classification:

C07D 473/06 (2006.01) *A61P 3/10* (2006.01)
C07D 473/08 (2006.01) *A61P 9/12* (2006.01)
A61K 31/522 (2006.01) *A61P 11/06* (2006.01)

(21) International Application Number:

PCT/IN2009/000192

(22) International Filing Date:

23 March 2009 (23.03.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

746/CHE/2008 26 March 2008 (26.03.2008) IN

(71) **Applicant** (for all designated States except US): **ADVINUS THERAPEUTICS PVT. LTD.**, [IN/IN]; 21 & 22, Peenya Industrial Area, Phase II, Bangalore 560 058 (IN).(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **PALLE, Venkata** [US/IN]; Advinus Therapeutics Pvt. Ltd., Quantum Towers, Plot No. 9, Rajiv Gandhi Infotech Park, Phase - I, Hinjewadi, Pune 411 057 (IN). **BASU, Sujay** [IN/IN]; Advinus Therapeutics Pvt.Ltd., Quantum Towers, Plot No. 9, Rajiv Gandhi Infotech Park, Phase - I, Hinjewadi, Pune 411 057 (IN). **WAMAN, Yogesh** [IN/IN]; Advinus Therapeutics Pvt.Ltd., Quantum Towers, Plot No. 9, Rajiv Gandhi Infotech Park, Phase - I, Hinjewadi, Pune 411 057 (IN). **RAMDAS, Vidya** [IN/IN]; Advinus Therapeutics Pvt.Ltd., Quantum Towers, Plot No. 9, Rajiv Gandhi Infotech Park, Phase - I, Hinjewadi, Pune 411 057 (IN). **BARAWKAR, Dinesh** [IN/IN]; Advinus Therapeutics Pvt. Ltd., Quantum Towers, Plot No. 9, Rajiv Gandhi Infotech Park, Phase - I, Hinjewadi, Pune 411 057 (IN).

(74) **Agents:** **VARADHACHARI, Lakshmikumaran** et al.; Lakshmikumaran & Sridharan, B-6/10, Safdarjung Enclave, New Delhi 110 029 (IN).


(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) **Date of publication of the international search report:**
10 December 2009

(54) **Title:** HETEROCYCLIC COMPOUNDS AS ADENOSINE RECEPTOR ANTAGONIST

(57) **Abstract:** Compounds of the present disclosure are fused pyrimidine compounds of formula (I), its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof, as Adenosine receptor antagonists. Processes of their preparation are also described in the disclosure.

54348-1

HETEROCYCLIC COMPOUNDS AS ADENOSINE RECEPTOR ANTAGONIST

Field of the invention

The present invention relates to a series of novel substituted fused pyrimidine compounds, their tautomers, polymorphs, stereoisomers, prodrugs, solvates, pharmaceutically acceptable salts, pharmaceutical compositions containing them and methods of treating conditions and diseases that are mediated by adenosine receptor (AR) activity. These compounds are useful in the treatment, prevention or suppression of diseases and disorders that may be susceptible to improvement by antagonism of the adenosine receptor, such as asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.

Background of the invention

Inflammatory and immunological reactions protect the host from invasion by microorganisms and eliminate debris at sites of tissue injury but they can also be responsible for significant tissue damage. Thus, regulatory mechanisms that limit damage from an overly exuberant immune response have evolved. It is increasingly apparent that adenosine, a purine nucleoside that is elaborated at injured and inflamed sites, has a central role in the regulation of inflammatory responses and in limiting inflammatory tissue destruction.

Adenosine is a ubiquitous purine nucleoside, playing a pivotal role in many biological processes such as energy generation, cell proliferation and proteins metabolism (European Journal of Pharmacology 533 (2006) 77–88). It acts on immune cells like mast cells, monocytes, macrophages, neutrophils, eosinophils, lymphocytes, airway smooth muscle cells, endothelial cells and airway epithelia. It is normally present in human tissues at low concentrations, but in response to metabolic stress, such as that encountered in the course of inflammatory events or during tissue hypoxia, a rapid increase in adenosine tissue levels takes place. Once generated, adenosine elicits its biological activities by interacting with its receptors (Jacobson KA, Gao ZG, Adenosine receptors as therapeutic targets. Nat Rev Drug Disc. 2006;5:247-264). There are four known subtypes of adenosine receptors (ARs) — referred to as A₁, A_{2A}, A_{2B} and A₃. It has been

54348-1

known for long time that intracellular signals activated by adenosine receptors include either stimulation or inhibition of adenylyl cyclase. In general, A₁ and A₃ receptors are coupled to pertussis toxin inhibited Gi coupled signal transduction proteins whereas A₂ receptors (A_{2A} and A_{2B}) are G_{α,s} linked receptors and stimulate adenylyl cyclase and cAMP.

5 Adenosine accumulation during ischemia and inflammation protects tissues from injury (Linden J, Molecular approach to adenosine receptors: receptor-mediated mechanisms of tissue protection. Rev. Pharmacol. Toxicol. 2001;41:775-787). Adenosine is related both structurally and metabolically to the bioactive nucleotides adenosine triphosphate (ATP), adenosine diphosphate (ADP), adenosine monophosphate (AMP) and cyclic adenosine mono-phosphate (cAMP); to the
10 biochemical methylating agent S-adenosyl-L-methione, and structurally to the coenzymes NAD, FAD and coenzyme A; and to RNA. Together adenosine and these related compounds are important in the regulation of many aspects of cellular metabolism (Poulsen SA, Quinn RJ, Adenosine receptors: new opportunities for future drugs. Bioorg Med Chem. 1998 Jun;6(6):619-641).

15 Adenosine signaling is known to serve apoptotic, angiogenic and pro-inflammatory functions and might be relevant to the pathogenesis of asthma and chronic obstructive pulmonary disease (Trends in Pharmacological Sciences, Vol.24, No.8, August 2003). Extracellular adenosine acts as a local modulator with a generally cytoprotective function in the body. Its effects on tissue protection and repair fall into four categories: increasing the ratio of oxygen
20 supply to demand; protecting against ischaemic damage by cell conditioning; triggering anti-inflammatory responses; and the promotion of angiogenesis.

The A_{2B} adenosine receptor subtype (see Feoktistov, I., Biaggioni, I. Pharmacol. Rev. 1997, 49, 381-402) has been identified in a variety of human and murine tissues and is involved in the regulation of vascular tone, smooth muscle growth, angiogenesis, hepatic glucose production,
25 bowel movement, intestinal secretion, and mast cell degranulation.

A_{2B} receptors have been implicated in mast cell activation and asthma, control of vascular tone, cardiac myocyte contractility, cell growth and gene expression, vasodilation, regulation of cell growth, intestinal function, and modulation of neurosecretion (Lin JH, Lu AYH. Role of Pharmacokinetics and Metabolism in Drug Discovery and Development, Pharmacological
30 Reviews, December 1, 1997 vol. 49 no. 4 403-449).

54348-1

A_{2B} receptors modulate mast cell function. Adenosine activates adenylate cyclase and protein kinase C, and potentiates stimulated mediator release in mouse bone marrow derived mast cells. (Feoktsov I, Biaggioni I, Polosa R, Holgate ST, Adenosine A_{2B} receptors: a novel therapeutic target in asthma? Trends in Pharmacological Sciences, April 1, 1998 vol 19 no 4 148-153). Activation of A_{2B} receptors in HMC-1 augments IL-8 release and potentiates PMA-induced secretion of IL-8. Thus, adenosine would contribute to the asthmatic response by acting on the mast

cell to enhance the release of proinflammatory mediators. (*Pulmonary Pharmacology & Therapeutics* 1999, 12, 111–114). In COPD, transformation of pulmonary fibroblasts into myofibroblasts is considered a major mechanism. Activation of the A_{2B} AR is involved in this process. Selective A_{2B} antagonists are expected to have beneficial effect on pulmonary fibrosis (*Curr. Drug Targets*, 2006, 7, 699-706; *Am. J. Resper. Cell. Mol. Biol.*, 2005, 32, 228). A_{2B} antagonists can be used as wound healing agents. Activation of the A_{2B} AR promotes angiogenesis by increasing the release of angiogenic factors and A_{2B} antagonists are useful to block angiogenesis (*Circ. Res.*, 2002, 90, 531-538). A_{2B} AR may be involved in the inhibition cardiac fibroblast (CF) proliferation (*Am.J. Physiol. Heart Circ. Physiol.*, 2004, 287, H2478-H2486). Adenosine stimulates Cl⁻ secretion in the intestinal epithelia pointing towards a possible treatment for cystic fibrosis patients with CFTR mutation (*Am. J. Respir. Cell Mol. Biol.*, 2008, 39, 190-197). High affinity A_{2B} antagonists are effective in hot plate model suggestive of the role of A_{2B} in nociception and can be used as potential analgesic agents (*The J. of Pharmacol. and Exp. Ther.*, 2004, 308, 358-366).

A_{2B} receptor is involved in release of IL-6. Increasing evidence suggests that IL-6 plays a role in Alzheimer's disease in the context of inflammatory process associated with disease. Hence A_{2B} receptor antagonist might be useful for Alzheimer's disease.

The A_{2B} ARs are involved in the stimulation of nitric oxide production during Na⁺-linked glucose or glutamine absorption. They are involved in glucose production in hepatocytes upon agonist stimulation. A_{2B}-receptor antagonists showed an anti-diabetic potential mainly by increasing plasma insulin levels under conditions when the adenosine tonus was elevated in-vivo and increased insulin release in-vitro (*J Pharm. Pharmacol.* 2006 Dec; 58(12):1639-45). Thus A_{2B} antagonists may serve as a novel target for the treatment of this metabolic disease.

A_{2B} receptor is a negative modulator of TNF- α hence A_{2B} agonists might have application in the management of sepsis.

The A_{2B} ARs are also important for adenosine-mediated inhibition of cardiac fibroblast functions. Adenosine receptor agonists inhibit rat cardiac fibroblasts with pharmacology suggestive of A_{2B} receptor indicating role of A_{2B} in cardiac remodelling and abnormal growth in cardiovascular diseases. Drugs that stimulate adenosine A_{2B} receptors or increase adenosine levels are new candidates for preventing

cardiac remodeling after MI (Circulation. 2006 Oct 31; 114(18): 1923-32. Epub 2006 Oct 16.).

Direct injections of adenosine into the corpus cavernosum of impotence patients produce a brief erection and if this effect is also mediated by A2B receptors in humans, it will be possible to develop stable and selective agonists that can be given locally.

An adenosine receptor-mediated signal-transduction pathway in the dermal papilla cells (DPCs) of hair contributes to minoxidil-induced hair growth through A2B receptor (J. Invest. Dermatol. 2007 Jun;127(6):1318-25. Epub 2007 Feb 15). Thus A2B agonists might stimulate hair growth through FGF-7 upregulation in DPCs.

In view of the physiological effects mediated by adenosine receptor, several A2B receptor antagonists have been recently disclosed for the treatment or prevention of asthma, bronchoconstriction, allergic diseases, hypertension, atherosclerosis, reperfusion injury, myocardial ischemia, retinopathy, inflammation, gastrointestinal tract disorders, cell proliferation diseases and/or diabetes mellitus. See for example WO2008002902, WO2007149277, WO2007017096, WO2007109547, WO2006091896, WO2006015357, WO2005042534, WO2005021548, WO2004106337, WO2003000694, WO2003082873, WO2003006465, WO2003053361, WO2003002566, WO2003063800, WO2003042214, WO2003035639, EP1283056, WO200073307, WO2000125210, WO2000073307, US20050119287, US20060281927.

It has now been found that compounds of the present invention are potent antagonists of the A2B adenosine receptor and can therefore be used in the treatment of the diseases mentioned herein above.

Under normal physiological conditions, A₁ ARs are quiescent; however, A₁ ARs are upregulated in conditions of stress, such as ischaemia, and in conditions of inflammation, typified by the inflammatory airway involvement in human asthmatics. A₁ ARs are upregulated in airway epithelium and bronchial smooth muscle in human asthmatics. A₁ ARs have been described on a number of different human cell types that are important in the pathophysiology of asthma, including APCs, human airway epithelial and bronchial smooth muscle cells, lymphocytes, mast cells, neutrophils, monocytes, macrophages, fibroblasts and endothelial cells. Activation of A₁ ARs on these different cell types induces the release of mediators and cytokines that lead to airway hyperreactivity, inflammation and airway remodelling. Activation of A₁ ARs on

54348-1

human asthmatic bronchial tissue produces bronchoconstriction. On human airway epithelial cells, activation of A₁ ARs causes an increase in expression of the MUC 2 gene responsible for mucus hypersecretion. Moreover, activation of A₁ ARs on a number of different human cells produces pro-inflammatory effects. Taken together, these effects of A₁ ARs in humans suggest
 5 that the A₁ AR antagonists could play potential therapeutic role in inflammatory diseases (C N Wilson, British J. of Pharm., 2008, 155, 475-86 and references cited therein). A₁ AR antagonists have been shown to have efficacy in rodent models of asthma and inflammation ((*J. Pharmacol. Exp. Ther.* 315, 329-336, 2005; *Eur. J. Pharmacol.*, 551, 116-124, 2006). A₁ antagonists have also been shown to have therapeutic potential in diseases such as hypertention,
 10 congestive heart failure where underlying mechanism is diuresis. There are several compounds in development for these indications (*J. Am. Soc. Nephrol.* 10, 714-720, 1999; *Circulation*, 105, 1348-1353, 2002; *J. Pharmacol. Exp. Ther.* 308, 846-856, 2004). A₁ AR antagonists are reported to reduce infarct size. It has been suggested that the ability of A₁ AR antagonists to reduce the the infarct size is also mediated by antagonism at A_{2B} AR (*Circulation*, 1996, 9,
 15 94; *J. Pharmacol. Exp. Ther.*, 2000, 292, 3, 929-938)

Activation of A₃ ARs induces the release of preformed mediators from basophils and produces bronchoconstriction, eosinophil migration into airways and mucus hypersecretion in animals, A₃ AR antagonists have been recommended for development as anti-asthma drugs (Fishman P, Bar-Yehuda S, Pharmacology and Therapeutic Applications of A₃ Receptor
 20 Subtype. Current Topics in Medicinal Chemistry, 2003, 3, 463-469; Nadeem A et al., Adenosine A₁ receptor antagonist versus montelukast on airway reactivity and inflammation. *Eur J Pharmacol.* 551(1-3): 116-124, Elsevier B.V. (2006)). A₃ AR antagonists have also been shown to play therapeutic role in various diseases including cardio-protection (*Vasc. Pharmacol.*, 2005, 42, 271; *J. Pharm. Exp. Ther.*, 2006, 319, 1200) and cancer
 25 (WO200010391).


Since several ARs have been implicated in asthma/COPD diseases pathophysiology, a pan AR antagonist may have therapeutic advantage.

54348-1

It has now been found that some of the compounds of the present invention are non-selective antagonists of ARs and can therefore be used in the treatment of above mentioned diseases.

Summary of the invention

- 5 In various embodiments, the present invention relates to a compound of formula (I):



I

or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, wherein,

- 10 Y is N or CH;

R^1 and R^2 are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxyalkyl, carboxyalkyl, cycloalkyl, cycloalkylalkyl, arylalkyl, aryl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl;

- wherein alkyl, alkenyl, alkynyl, alkoxyalkyl, carboxyalkyl, cycloalkyl, cycloalkylalkyl, arylalkyl, aryl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl are unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, $-SO_3H$, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycllyoxy, hydroxyamino, alkoxyamino, nitro, $-S(O)_2NR^aR^a$, $-NR^aS(O)_2R^a$ or $-S(O)_pR^b$;
- 15
20

54348-1

wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

- 5 wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2;

R³ is selected from the group consisting of hydrogen and alkyl;

- 10 wherein alkyl is unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, carboxyalkyloxy, alkylcarboxyalkyloxy, -SO₃H, aryl, 15 aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycllyoxy, hydroxyamino, alkoxyamino, CF₃, nitro, S(O)₂NR^aR^a, -NR^aS(O)₂R^a or -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl,

- 20 aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or

- 25 heteroaryl; and p is 0, 1 or 2;

X is either an unsubstituted or substituted arylene or an unsubstituted or substituted heteroarylene;

54348-1

A is selected from the group consisting of -O-, -C(O)NR⁴-, -NR⁴C(O)- and -(CR⁵R⁶)_q-;

wherein q is 1 or 2, and R⁴ is selected from the group consisting of hydrogen and alkyl,

wherein alkyl is either unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino,

- 5 arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxy-carbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycl-yloxy, hydroxyamino, alkoxyamino, CF₃, nitro, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a or -S(O)_pR^b;

- 10 wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

- 15 wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2;

R⁵ and R⁶ are independently selected from the group consisting of hydrogen, halogen, alkyl, hydroxyl, alkoxy and -C(O)R⁷;

- 20 wherein alkyl and alkoxy are unsubstituted, or substituted with halogen, hydroxy, hydroxyalkyl or CF₃;

or

R⁵ and R⁶ together represent O, S or cycloalkyl;

R⁷ is hydroxyl, or unsubstituted or substituted amino;

- 25 m and n are independently 0, 1, 2, 3, 4, 5 or 6;

54348-1

B is selected from the group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted alkenyl, unsubstituted or substituted cycloalkyl and unsubstituted or substituted heterocyclyl; and

5 D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted heterocyclylalkyl,

wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocyclylalkyl are substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, 10 monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, CF₃, OCF₃, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycliloxy, hydroxyamino, alkoxyamino, nitro, -C(O)R, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and 15 -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, -OCF₃, amino, substituted amino, cyano and -S(O)_pR^c;

20 wherein R is selected from the group consisting of hydrogen, hydroxyl, alkyl, alkoxy, amino, monoalkylamino, dialkylamino and heterocyclyl; each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl or heteroaryl; and p is 0, 1 or 2.

25 The present invention also relates to compounds of formula I or its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salt thereof that may have other AR antagonism such as A₁ A_{2A} and A₃ apart from A_{2B} AR antagonism.

54348-1

The present invention also relates to a process of preparation of compounds of formula-I. These and other features, aspects, and advantages of the present subject matter will become better understood with reference to the following description and appended claims. This summary is provided to introduce a selection of concepts in a simplified form. This
5 summary is not intended to identify key features or essential features of the claimed subject matter, nor is it intended to be used to limit the scope of the claimed subject matter.

Definitions

In the structural formulae given herein and throughout the present disclosure, the following terms have the indicated meaning, unless specifically stated otherwise.

10 The term "optionally substituted" as used herein means that the group in question is either unsubstituted or substituted with one or more of the substituents specified. When the group in question is substituted with more than one substituent, the substituent may be same or different.

The term "alkyl" refers to a monoradical branched or unbranched saturated
15 hydrocarbon chain having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 carbon atoms, preferably 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms, more preferably 1, 2, 3, 4, 5 or 6 carbon atoms. This term is exemplified by groups such as

methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, t-butyl, n-hexyl, n-decyl, tetradecyl, and the like.

The term "alkylene" refers to a diradical of a branched or unbranched saturated hydrocarbon chain, having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or
 5 20 carbon atoms, preferably 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms, more preferably 1, 2, 3, 4, 5 or 6 carbon atoms. This term is exemplified by groups such as methylene (-CH₂-), ethylene (-CH₂CH₂-), the propylene isomers (e.g., -CH₂CH₂CH₂- and -CH(CH₃)CH₂-) and the like.

The term "substituted alkyl" or "substituted alkylene" refers to: 1) an alkyl
 10 group or alkylene group as defined above, having 1, 2, 3, 4 or 5 substituents, preferably 1, 2 or 3 substituents, selected from the group consisting of alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, heteroaryl, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy,
 15 carboxyalkyl, -SO₃H, aryl, aryloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, nitro, -S(O)₂NRaRa, -NRaS(O)₂Ra and -S(O)_pRb, where each Ra is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; heterocyclyloxy where Rb is
 20 hydrogen, alkyl, aryl, heteroaryl or heterocyclyl. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1, 2, or 3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and -S(O)_pRc, where Rc is alkyl, aryl, or heteroaryl and p is 0, 1 or 2;

25 or 2) an alkyl group or alkylene group as defined above that is interrupted by 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 atoms independently chosen from oxygen, sulfur and NRd-, where Rd is chosen from hydrogen, alkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl and heterocyclyl or groups selected from carbonylalkyl, carboxyester, carboxyamide and sulfonyl. All substituents may be optionally further substituted by alkyl, alkoxy,
 30 halogen, CF₃, amino, substituted amino, cyano, or -S(O)_pRc, in which Rc is alkyl, aryl, or heteroaryl and p is 0, 1, or 2;

or 3) an alkyl group or alkylene as defined above that has 1, 2, 3, 4 or 5 substituents as defined above, as well as interrupted by 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 atoms as defined above.

The term "alkenyl" refers to a monoradical of a branched or unbranched unsaturated acyclic hydrocarbon group preferably having from 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 carbon atoms, more preferably 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms and even more preferably 2, 3, 4, 5 or 6 carbon atoms and having 1, 2, 3, 4, 5 or 6 double bond (vinyl), preferably 1 double bond. Preferred alkenyl groups include ethenyl or vinyl ($-\text{CH}=\text{CH}_2$), 1-propylene or allyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), isopropylene ($-\text{C}(\text{CH}_3)=\text{CH}_2$), bicyclo [2.2. 1] heptene, and the like.

The term "alkenylene" refers to a diradical of a branched or unbranched unsaturated acyclic hydrocarbon group preferably having from 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 carbon atoms, more preferably 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms and even more preferably 2, 3, 4, 5 or 6 carbon atoms and having 1, 3, 4, 5 or 6 double bond (vinyl), preferably 1 double bond.

The term "substituted alkenyl" refers to an alkenyl group as defined above having 1, 2, 3, 4 or 5 substituents, and preferably 1, 2, or 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, thiocarbonyl, carboxy, carboxyalkyl, SO_3H , aryl, aryloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, nitro, $-\text{S}(\text{O})_2\text{NR}^a\text{R}^a$, $-\text{NR}^a\text{S}(\text{O})_2\text{R}^a$ and $-\text{S}(\text{O})_p\text{R}^b$ where each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; heterocyclyloxy where R^b is alkyl, aryl, heteroaryl or heterocyclyl and p is 0, 1 or 2. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1, 2, or 3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF_3 , amino, substituted amino, cyano, and $-\text{S}(\text{O})_p\text{R}^c$, where R^c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

The term "alkynyl" refers to a monoradical of a branched or unbranched unsaturated acyclic hydrocarbon, preferably having from 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 carbon atoms, more preferably 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms and even more preferably 2, 3, 4, 5 or 6 carbon atoms and having 1, 2,

3, 4, 5 or 6 sites of acetylene (triple bond) unsaturation, preferably 1 triple bond. Preferred alkynyl groups include ethynyl, ($-C\equiv CH$), propargyl (or prop-1-yn-3-yl, $-CH_2C\equiv CH$), homopropargyl (or but-1-yn-4-yl, $-CH_2CH_2C\equiv CH$) and the like.

The term "alkynylene" refers to a diradical of a branched or unbranched unsaturated acyclic hydrocarbon group preferably having from 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 carbon atoms, more preferably 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms and even more preferably 2, 3, 4, 5 or 6 carbon atoms and having 1, 3, 4, 5 or 6 sites of acetylene (triple bond) unsaturation, preferably 1 triple bond.

The term "substituted alkynyl" refers to an alkynyl group as defined above having 1, 2, 3, 4 or 5 substituents, and preferably 1, 2, or 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, $-SO_3H$, aryl, aryloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, nitro, $-S(O)_2NR^aR^a$, $-NR^aS(O)_2R^a$ and $-S(O)_pR^b$, where each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl heteroarylalkyl, heterocyclyl and heterocyclylalkyl; heterocyclyloxy where R^b is alkyl, aryl, heteroaryl or heterocyclyl and p is 0, 1 or 2. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1, 2, or 3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF_3 , amino, substituted amino, cyano, and $-S(O)_pR^c$ where R^c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

The term "cycloalkyl" refers to carbocyclic groups of from 3 to 20 carbon atoms having a single cyclic ring or multiple condensed rings which may be partially unsaturated. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, bicyclo[2.2.1]heptane, 1,3,3-trimethylbicyclo[2.2.1]hept-2-yl, (2,3,3-trimethylbicyclo[2.2.1]hept-2-yl), or carbocyclic groups to which is fused an aryl group, for example indane, and the like. In the above mentioned cycloalkyl ring, one or two methylene groups may be replaced by C(O).

The term "substituted cycloalkyl" refers to cycloalkyl groups having 1, 2, 3, 4 or 5 substituents, and preferably 1, 2, or 3 substituents, selected from the group consisting of alkyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, thiocarbonyl, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, nitro, -C(O)R and -S(O)_pR^b, where R is hydrogen, hydroxyl, alkoxy, alkyl and cycloalkyl, heterocyclyloxy where R^b is alkyl, aryl, heteroaryl or heterocyclyl and p is 0, 1 or 2. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1, 2, or 3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and-S(O)_pR_c, where R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

"Halo" or "Halogen", alone or in combination with any other term means halogens such as chloro (Cl), fluoro (F), bromo (Br) and iodo (I).

"Haloalkyl" refers to a straight chain or branched chain haloalkyl group with 1 to 6 carbon atoms. The alkyl group may be partly or totally halogenated. Representative examples of haloalkyl groups include but are not limited to fluoromethyl, chloromethyl, bromomethyl, difluoromethyl, dichloromethyl, dibromomethyl, trifluoromethyl, trichloromethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, 3-fluoropropyl, 3-chloropropyl, 3-bromopropyl and the like.

The term "alkoxy" refers to the group R'''-O-, where R''' is optionally substituted alkyl or optionally substituted cycloalkyl, or optionally substituted alkenyl or optionally substituted alkynyl; or optionally substituted cycloalkenyl, where alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkenyl are as defined herein. Representative examples of alkoxy groups include but are not limited to methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, tert-butoxy, sec-butoxy, n-pentoxy, n-hexoxy, 1,2-dimethylbutoxy, trifluoromethoxy, and the like.

The term "aminocarbonyl" refers to the group -C(O)NR'R' where each R' is independently hydrogen, alkyl, aryl, heteroaryl, heterocyclyl or both R' groups are joined to form a heterocyclic group (e. g. morpholino). Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1-3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen,

CF₃, amino, substituted amino, cyano, and -S(O)_pR_c, where R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

The term "acylamino" refers to the group -NR"C(O)R" where each R" is independently hydrogen, alkyl, aryl, heteroaryl, or heterocyclyl. Unless otherwise
 5 constrained by the definition, all substituents may optionally be further substituted by 1-3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and -S(O)_pR_c, where R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

The term "acyloxy" refers to the groups -OC(O)-alkyl, -OC(O)-cycloalkyl, -
 10 OC(O)-aryl, -OC(O)-heteroaryl, and -OC(O)-heterocyclyl. Unless otherwise constrained by the definition, all substituents may be optionally further substituted by alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_pR_c, where R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

15 "Alkoxyalkyl" refers to alkyl groups as defined above wherein at least one of the hydrogen atoms of the alkyl group is replaced by an alkoxy group as defined above. Representative examples of alkoxyalkyl groups include but are not limited to methoxymethyl, methoxyethyl, ethoxymethyl and the like.

"Aryloxyalkyl" refers to the group -alkyl-O-aryl. Representative examples of
 20 aryloxyalkyl include but are not limited to phenoxymethyl, naphthyloxymethyl, phenoxyethyl, naphthyloxyethyl and the like.

"Di alkylamino" refers to an amino group, to which two same or different straight chain or branched chain alkyl groups with 1 to 6 carbon atoms are bound. Representative examples of di alkylamino include but are not limited to dimethylamino,
 25 diethylamino, methylethylamino, dipropylamino, dibutylamino and the like.

"Cycloalkylalkyl" refers to an alkyl radical as defined above which is substituted by a cycloalkyl radical as defined above. Representative examples of cycloalkylalkyl include but are not limited to cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, 1-cyclopentylethyl, 1-cyclohexylethyl, 2-
 30 cyclopentylethyl, 2-cyclohexylethyl, cyclobutylpropyl, cyclopentylpropyl, cyclohexylbutyl and the like.

"Aminoalkyl" refers to an amino group that is attached to (C₁₋₆)alkylene as defined herein. Representative examples of aminoalkyl include but are not limited to

aminomethyl, aminoethyl, 1-aminopropyl, 2-aminopropyl, and the like. The amino moiety of aminoalkyl may be substituted once or twice with alkyl to provide alkylaminoalkyl and dialkylaminoalkyl respectively. Representative examples of alkylaminoalkyl include but are not limited to methylaminomethyl, methylaminoethyl, 5 methylaminopropyl, ethylaminoethyl and the like. Representative examples of dialkylaminoalkyl include but are not limited to dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, N-methyl-N-ethylaminoethyl and the like.

The term "aryl" refers to an aromatic carbocyclic group of 6 to 20 carbon atoms having a single ring (e.g. phenyl) or multiple rings (e.g. biphenyl), or multiple 10 condensed (fused) rings (e.g. naphthyl or anthranyl). Preferred aryls include phenyl, naphthyl and the like.

The term "arylene" refers to a diradical of an aryl group as defined above. This term is exemplified by groups such as 1,4-phenylene, 1,3-phenylene, 1,2-phenylene, 1,4'-biphenylene, and the like.

15 Unless otherwise constrained the aryl or arylene groups may optionally be substituted with 1, 2, 3 4 or 5 substituents, preferably 1, 2 or 3 substituents, selected from the group consisting of alkyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, carboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, heteroaryl, aminosulfonyl, 20 aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, nitro, -S(O)₂NRaRa, -NRaS(O)₂Ra and -S(O)_pRb where each Ra is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl heteroarylalkyl, heterocyclyl and heterocyclylalkyl; where Rb is hydrogen, alkyl, aryl, heterocyclyl or heteroaryl and p is 25 0, 1 or 2. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1, 2 or 3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and -S(O)_pRc where Rc is hydrogen, alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

30 The term "arylalkyl" refers to an aryl group covalently linked to an alkylene group, where aryl and alkylene are defined herein.

"Optionally substituted arylalkyl" refers to an optionally substituted aryl group covalently linked to an optionally substituted alkylene group. Such arylalkyl groups are exemplified by benzyl, phenethyl, naphthylmethyl, and the like.

The term "aryloxy" refers to the group aryl-O- wherein the aryl group is as
5 defined above, and includes optionally substituted aryl groups as also defined above.

The term "arylthio" refers to the group -S-aryl, where aryl is as defined herein including optionally substituted aryl groups as also defined above.

The term "substituted amino" refers to the group -NR'R' where each R' is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl,
10 carboxyalkyl, alkoxycarbonyl, aryl, heteroaryl and heterocyclyl. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1, 2 or 3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and -S(O)_pR_c, where R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

15 The term "carboxyalkyl" refers to the groups -alkylene-C(O)OH.

The term "alkylcarboxyalkyl" refers to the groups -alkylene-C(O)OR_d where R_d is alkyl, cycloalkyl, where alkyl, cycloalkyl are as defined herein, and may be optionally further substituted by alkyl, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_pR_c, in which R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

20 The term "alkylcarboxy" refers to the group -C(O)O-alkyl, where the alkyl is as defined herein and may be optionally substituted by alkyl, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_pR_c, in which R_c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2.

The term "alkylcarboxyalkyloxy" refers to the group -alkylene-C(O)O-
25 alkylene-O-, where alkylene is as defined herein.

The term "heteroaryl" refers to an aromatic cyclic group having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 carbon atoms and 1, 2, 3 or 4 heteroatoms selected from oxygen, nitrogen and sulfur within at least one ring. Such heteroaryl groups can have a single ring (e.g. pyridyl or furyl) or multiple condensed rings (e.g. indoliziny, benzothiazolyl, or benzothienyl). Examples of heteroaryls include, but are not limited
30 to, [1,2,4] oxadiazole, [1,3,4] oxadiazole, [1,2,4] thiadiazole, [1,3,4] thiadiazole, pyrrole, imidazole, pyrazole, pyridine pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine,

quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, furan, thiophene, oxazole, thiazole, triazole, triazine and the like.

The term "heteroarylene" refers to a diradical of a heteroaryl group as defined above.

5 Unless otherwise constrained the heteroaryl or heterarylene groups can be optionally substituted with 1, 2, 3, 4 or 5 substituents, preferably 1, 2 or 3 substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, thiocarbonyl, carboxy, carboxyalkyl, -SO₃H, aryl, 10 aryloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxylamino, nitro, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and -S(O)_pR^b, where each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl heteroarylalkyl, heterocyclyl and heterocyclylalkyl; where R^b is hydrogen, alkyl, aryl, heterocyclyl or 15 heteroaryl, and p is 0, 1 or 2. Unless otherwise constrained by the definition, all substituents may optionally be further substituted by 1-3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and -S(O)_nR_c, where R_c is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

20 The term "heteroarylalkyl" refers to a heteroaryl group covalently linked to an alkylene group, where heteroaryl and alkylene are defined herein.

"Optionally substituted heteroarylalkyl" refers to an optionally substituted heteroaryl group covalently linked to an optionally substituted alkylene group. Such heteroarylalkyl groups are exemplified by 3-pyridylmethyl, quinolin-8-ylethyl, 4-methoxythiazol-2-ylpropyl, and the like. 25

The term "heterocyclyl" refers to a saturated or partially unsaturated group having a single ring or multiple condensed rings, having from 1 to 40 carbon atoms and from 1 to 10 hetero atoms, preferably 1, 2, 3 or 4 heteroatoms, selected from nitrogen, sulfur, phosphorus, and/or oxygen within the ring. Heterocyclic groups can have a 30 single ring or multiple condensed rings, and include tetrahydrofuranyl, morpholinyl, piperidinyl, piperazinyl, dihydropyridinyl, tetrahydroquinolinyl and the like. In the above mentioned heterocyclyl ring, one or two methylene groups may independently be replaced by one of C(O), S(O) or SO₂. Unless otherwise constrained by the definition

for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1, 2, 3, 4 or 5, and preferably 1, 2 or 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, 5 oxo, -C(O)R where R is hydrogen, hydroxyl, alkoxy, alkyl and cycloalkyl, thiocarbonyl, carboxy, carboxyalkyl, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, nitro, and -S(O)_pR_b, where R_b is hydrogen, alkyl, aryl, heterocyclyl or heteroaryl and p is 0, 1 or 2. Unless otherwise constrained by the definition, all 10 substituents may optionally be further substituted by 1-3 substituents chosen from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano, and -S(O)_nR_c, where R_c is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

The term "heterocyclylalkyl" refers to a heterocyclyl group covalently linked to 15 an alkylene group, where heterocyclyl and alkylene are defined herein.

"Optionally substituted heterocyclylalkyl" refers to an optionally substituted heterocyclyl group covalently linked to an optionally substituted alkylene group.

The term "heteroaryloxy" refers to the group heteroaryl-O-.

The term "thiol" refers to the group -SH.

20 The term "substituted alkylthio" refers to the group -S-substituted alkyl.

The term "heteroarylthio" refers to the group -S-heteroaryl wherein the heteroaryl group is as defined above including optionally substituted heteroaryl groups as also defined above.

The term "sulfoxide" refers to a group -S(O).

25 "Substituted sulfoxide" refers to a group -S(O)R, in which R is substituted alkyl, substituted aryl, or substituted heteroaryl, as defined herein.

The term "sulfone" refers to a group -S(O)₂.

The term "substituted sulfone" refers to a group -S(O)₂R, in which R is alkyl, aryl, or heteroaryl.

30 The compounds of the present invention may have the ability to crystallize in more than one form, a characteristic known as polymorphism, and all such polymorphic forms ("polymorphs") are encompassed within the scope of the invention. Polymorphism generally can occur as a response to changes in temperature or pressure

or both, and can also result from variations in the crystallization process. Polymorphs can be distinguished by various physical characteristics, and typically the x-ray diffraction patterns, solubility behavior, and melting point of the compound are used to distinguish polymorphs.

5 The compounds described herein may contain one or more chiral centers and/or double bonds and therefore, may exist as stereoisomers, such as double-bond isomers (i.e., geometric isomers), regioisomers, enantiomers or diastereomers. Accordingly, the chemical structures depicted herein encompass all possible enantiomers and stereoisomers of the illustrated or identified compounds including the
10 stereoisomerically pure form (e.g., geometrically pure, enantiomerically pure or diastereomerically pure) and enantiomeric and stereoisomeric mixtures. Enantiomeric and stereoisomeric mixtures can be resolved into their component enantiomers or stereoisomers using separation techniques or chiral synthesis techniques well known to the person skilled in the art. The compounds may also exist in several tautomeric forms
15 including the enol form, the keto form and mixtures thereof. Accordingly, the chemical structures depicted herein encompass all possible tautomeric forms of the illustrated or identified compounds.

Compounds may exist in unsolvated forms as well as solvated forms, including hydrated forms and as N-oxides. In general, compounds may be hydrated, solvated or
20 N-oxides. Certain compounds may exist in multiple crystalline or amorphous forms. Also contemplated within the scope of the invention are congeners, analogs, hydrolysis products, metabolites and precursor or prodrugs of the compound. In general, unless otherwise indicated, all physical forms are equivalent for the uses contemplated herein and are intended to be within the scope of the present invention.

25 “Prodrug” refers to a derivative of a drug molecule as, for example, esters, carbonates, carbamates, ureas, amides or phosphates that requires a transformation within the body to release the active drug. Prodrugs are frequently, although not necessarily, pharmacologically inactive until converted to the parent drug. Prodrugs may be obtained by bonding a promoiety (defined herein) typically via a functional
30 group, to a drug.

“Promoiety” refers to a group bonded to a drug, typically to a functional group of the drug, via bond(s) that are cleavable under specified conditions of use. The bond(s) between the drug and promoiety may be cleaved by enzymatic or non-


enzymatic means. Under the conditions of use, for example following administration to a patient, the bond(s) between the drug and promoiety may be cleaved to release the parent drug. The cleavage of the promoiety may proceed spontaneously, such as via a hydrolysis reaction, or it may be catalyzed or induced by another agent, such as by an enzyme, by light, by acid, or by a change of or exposure to a physical or environmental parameter, such as a change of temperature, pH, etc. The agent may be endogenous to the conditions of use, such as an enzyme present in the systemic circulation to which the prodrug is administered or the acidic conditions of the stomach or the agent may be supplied exogenously.

10 “Pharmaceutically acceptable salt” embraces salts with a pharmaceutically acceptable acid or base. Pharmaceutically acceptable acids include both inorganic acids, for example hydrochloric, sulphuric, phosphoric, diphosphoric, hydrobromic, hydroiodic and nitric acid and organic acids, for example citric, fumaric, maleic, malic, mandelic, ascorbic, oxalic, succinic, tartaric, benzoic, acetic, methanesulphonic, ethanesulphonic, benzenesulphonic or p-toluenesulphonic acid. Pharmaceutically acceptable bases include alkali metal (e.g. sodium or potassium) and alkali earth metal (e.g. calcium or magnesium) hydroxides and organic bases, for example alkyl amines, arylalkyl amines and heterocyclic amines.

Other preferred salts according to the invention are quaternary ammonium compounds wherein an equivalent of an anion (X-) is associated with the positive charge of the N atom. X- may be an anion of various mineral acids such as, for example, chloride, bromide, iodide, sulphate, nitrate, phosphate, or an anion of an organic acid such as, for example, acetate, maleate, fumarate, citrate, oxalate, succinate, tartrate, malate, mandelate, trifluoroacetate, methanesulphonate and p-toluenesulphonate. X- is preferably an anion selected from chloride, bromide, iodide, sulphate, nitrate, acetate, maleate, oxalate, succinate or trifluoroacetate. More preferably X- is chloride, bromide, trifluoroacetate or methanesulphonate.

Detailed description of the invention

The present invention provides compounds of formula I, or its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof, pharmaceutical compositions containing them and methods of treating conditions and diseases that are mediated by adenosine A_{2B} receptor activity,



I

or its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof, wherein,

5 Y is selected from N or CH;

R¹ and R² are independently selected from a group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxyalkyl, carboxyalkyl, cycloalkyl, cycloalkylalkyl, arylalkyl, aryl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl;

10 wherein alkyl, alkenyl, alkynyl, alkoxyalkyl, carboxyalkyl, cycloalkyl, cycloalkylalkyl, arylalkyl, aryl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl are unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, 15 halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycliloxy, hydroxyamino, alkoxyamino, nitro, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and -S(O)_pR^b;

20 wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

25 wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2;

R³ is selected from a group consisting of hydrogen and alkyl;

wherein alkyl is unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino,

monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, carboxyalkyloxy, alkylcarboxyalkyloxy, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, CF₃, nitro, S(O)₂NR^aR^a, - NR^aS(O)₂R^a and -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2;

X is either an unsubstituted or substituted arylene or an unsubstituted or substituted heteroarylene;

A is selected from a group consisting of -O-, -C(O)NR⁴-, -NR⁴C(O)- and -(CR⁵R⁶)_q-;

wherein q is 1 or 2, and R⁴ is selected from a group consisting of hydrogen and alkyl,

wherein alkyl is either unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, CF₃, nitro, -S(O)₂NR^aR^a, - NR^aS(O)₂R^a and -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

- wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2;
- 5 R^5 and R^6 are independently selected from a group consisting of hydrogen, halogen, alkyl, hydroxyl, alkoxy and $-C(O)R^7$;
- wherein alkyl and alkoxy are unsubstituted, or substituted with halogen, hydroxy, hydroxyalkyl, CF_3 ;
- or
- 10 R^5 and R^6 together represent O, S or cycloalkyl;
- R^7 is selected from hydroxyl, and unsubstituted or substituted amino;
- m and n are independently selected from 0, 1, 2, 3, 4, 5, and 6;
- B is selected from a group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted alkenyl, unsubstituted or substituted cycloalkyl and
- 15 unsubstituted or substituted heterocyclyl; and
- D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted heterocyclylalkyl,
- wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and
- 20 heterocyclylalkyl are substituted with 1, 2, or 3 substituents independently selected from alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, CF_3 , OCF_3 , keto, thiocarbonyl,
- 25 carboxy, alkylcarboxy, carboxyalkyl, $-SO_3H$, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyoxy, hydroxyamino, alkoxyamino, nitro, $-C(O)R$, $-S(O)_2NR^aR^a$, $-NR^aS(O)_2R^a$ and $-S(O)_pR^b$;
- wherein each substituent is unsubstituted or substituted with 1, 2, or 3
- 30 substituents independently selected from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF_3 , $-OCF_3$, amino, substituted amino, cyano and $-S(O)_pR^c$;

wherein R is selected from a group consisting of hydrogen, hydroxyl, alkyl, alkoxy, amino, monoalkylamino, dialkylamino and heterocyclyl; each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2.

In one embodiment, compounds of the invention comprise compounds of formula I excluding compounds:

wherein X is a heteroarylene, A is $-(CR^5R^6)_q-$, B is alkynylene or alkenylene, D is phenyl, pyridyl, pyrimidinyl, and R⁵ and R⁶ together is O.

As described herein above, the present invention provides novel fused pyrimidine compounds of formula (I), their tautomers, polymorphs, stereoisomers, prodrugs, solvates, pharmaceutically acceptable salts, pharmaceutical compositions containing them, methods for preparing said compounds and methods for the treatment, prevention or suppression of diseases and disorders that may be susceptible to improvement by antagonism of the adenosine receptor, such as asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.

According to an embodiment of the present invention, Y is N; R¹ and R² are independently selected from a group consisting of hydrogen, alkyl, alkenyl and alkynyl;

wherein alkyl, alkenyl and alkynyl are unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycliloxy, hydroxyamino, alkoxyamino, nitro, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and -S(O)_pR^b; wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl

heteroarylalkyl, heterocyclyl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; and p is 0, 1 or 2;

R^3 is selected from a group consisting of hydrogen and alkyl;

wherein alkyl is unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclalkylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, carboxyalkyloxy, alkylcarboxyalkyloxy $-SO_3H$, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclalkyl, heterocyclalkyloxy, hydroxyamino, alkoxylamino, nitro, $S(O)_2NR^aR^a$, $NR^aS(O)_2R^a$ and $-S(O)_pR^b$;

wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclalkyl and heterocyclalkylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclalkyl; and p is 0, 1 or 2;

X is either an unsubstituted or substituted arylene or an unsubstituted or substituted heteroarylene;

A is selected from a group consisting of $-O-$, $-C(O)NR^4-$, $-NR^4C(O)-$ and $-(CR^5R^6)_q-$;

wherein q is 1, or 2, and R^4 is selected from a group consisting of hydrogen and alkyl,

wherein alkyl is either unsubstituted or substituted with halogen, CF_3 and aryl;

R^5 and R^6 are independently selected from a group consisting of hydrogen, halogen, alkyl and hydroxyl;

or

R^5 and R^6 together represent O or cycloalkyl;

m and n are independently selected from 0, 1 and 2;

B is selected from a group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted cycloalkyl and unsubstituted or substituted heterocyclalkyl; and

D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted heterocyclalkyl,

5 wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocyclalkyl are substituted with 1, 2, or 3 substituents independently selected from alkyl, alkoxy, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, cyano, halogen, hydroxy, hydroxyalkyl, CF₃, OCF₃, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, heteroaryl, heterocycl, -C(O)R, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and -S(O)_pR^b;

10 wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, -OCF₃, amino, substituted amino, cyano and -S(O)_pR^c;

15 wherein R is selected from a group consisting of amino, monoalkylamino, dialkylamino and heterocycl; each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocycl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocycl; R^c is alkyl, aryl, 20 or heteroaryl; and p is 0, 1 or 2.

According to an embodiment of the present invention, wherein Y is N; R¹ and R² are independently selected from a group consisting of hydrogen and alkyl;

25 wherein alkyl is unsubstituted or substituted with alkenyl, alkoxy, acylamino, amino, monoalkylamino, dialkylamino, halogen, hydroxy, hydroxyalkyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H and aryl;

R³ is selected from a group consisting of hydrogen and alkyl;

X is either an unsubstituted or substituted arylene or an unsubstituted or substituted heteroarylene;

A is selected from a group consisting of -O-, -C(O)NR⁴-, -NR⁴C(O)- and -(CR⁵R⁶)_q-;

30 wherein q is 1 and R⁴ is selected from a group consisting of hydrogen and alkyl, wherein alkyl is either unsubstituted or substituted with halogen, CF₃ and aryl;

R^5 and R^6 are independently selected from a group consisting of hydrogen, halogen, alkyl and hydroxyl;

or

R^5 and R^6 together represent O or cycloalkyl;

5 m and n are independently selected from 0, 1 and 2;

B is selected from a group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted cycloalkyl and unsubstituted or substituted heterocyclyl; and

10 D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted heterocyclalkyl,

wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocyclalkyl are substituted with 1, 2, or 3 substituents independently selected from alkyl, alkoxy, acyl, acylamino, acyloxy, amino, monoalkylamino, 15 dialkylamino, arylamino, cycloalkylamino, cyano, halogen, hydroxy, hydroxyalkyl, CF_3 , OCF_3 , carboxy, alkylcarboxy, carboxyalkyl, $-SO_3H$, aryl, heteroaryl, heterocyclyl, $-C(O)R$, $-S(O)_2NR^aR^a$, $-NR^aS(O)_2R^a$ and $-S(O)_pR^b$;

20 wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF_3 , $-OCF_3$, amino, substituted amino, cyano and $-S(O)_pR^c$;

wherein R is selected from a group consisting of amino, monoalkylamino, dialkylamino and heterocyclyl; each R^a is independently selected from the group consisting of hydrogen, alkyl, 25 cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2.

30 According to an embodiment of the present invention, X is pyrazolyl, isoxazolyl, phenyl, pyridyl, oxazolyl or pyrimidyl.

According to another embodiment of the present invention, B is an alkynylene.

In another embodiment of the present invention, B is heterocyclyl or cycloalkyl.

Particular embodiments of the present invention are

- 1, 3-Dipropyl-8-[1-(3-p-tolyl-prop-2ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2, 6-dione,
- 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2, 6-dione,
- 5 8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-
10 purine-2,6-dione,
- 8-{1-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[3-(2,4-Difluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 15 8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione,
- 20 4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid ethyl ester,
- 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid ethyl ester,
- 25 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzonitrile,
- 8-{1-[3-(3-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 2-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid methyl ester,
- 30 8-{1-[4-(4-Fluoro-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

- 8-{1-[4-(3-Fluoro-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-[1-(4-p-tolyl-but-3-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,
 5 1,3-Dipropyl-8-{1-[4-(3-trifluoromethyl-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,
 1,3-Dipropyl-8-{1-[3-(2-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-
 10 dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-[1-(3-m-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,
 3-Ethyl-1-propyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 15 3-Ethyl-1-propyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 3-Ethyl-1-propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,
 3-Ethyl-8-{1-[3-(3-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-
 20 dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 25 3-Ethyl-8-{1-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
 3-{3-[4-(3-Ethyl-2,6-dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,
 3-Ethyl-1-propyl-8-[1-(3-m-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-
 30 2,6-dione,
 3-Ethyl-8-{1-[3-(4-methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

- 3-Ethyl-1-propyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-
3,7-dihydro-purine-2,6-dione,
3-Ethyl-8-{1-[3-(3-methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-
dihydro-purine-2,6-dione,
5 4-{3-[4-(3-Ethyl-2,6-dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-
prop-1-ynyl}-benzoic acid,
4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-
1-ynyl}-benzonitrile,
(3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-
10 prop-1-ynyl}-phenoxy)-acetic acid,
8-{1-[3-(3-tert-Butyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-
purine-2,6-dione,
4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-
1-ynyl}-benzoic acid,
15 (3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-
prop-1-ynyl}-phenyl)-acetic acid,
(4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-
prop-1-ynyl}-phenyl)-acetic acid,
8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-
20 purine-2,6-dione,
3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-
1-ynyl}-N-isopropyl-benzamide,
1,3-Dipropyl-8-(1-{3-[3-(pyrrolidine-1-carbonyl)-phenyl]-prop-2-ynyl}-1H-pyrazol-4-
yl)-3,7-dihydro-purine-2,6-dione,
25 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-
1-ynyl}-4-methyl-benzoic acid,
8-{1-[3-(3-Chloro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-
purine-2,6-dione,
3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-
30 1-ynyl}-4-methoxy-benzoic acid,
5-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-
1-ynyl}-pyridine-2-carboxylic acid methyl ester,

- 1,3-Dipropyl-8-{3-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,
 8-{3-[3-(2,4-Difluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 5 8-{3-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{3-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{3-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-
 10 dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-[3-(3-p-tolyl-prop-2-ynyloxy)-isoxazol-5-yl]-3,7-dihydro-purine-2,6-dione,
 8-{3-[3-(3-tert-Butyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 15 1,3-Dipropyl-8-{3-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,
 8-{1-Methyl-5-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{1-Methyl-5-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-1H-pyrazol-3-yl}-1,3-
 20 dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{5-[3-(3-Methoxy-phenyl)-prop-2-ynyloxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
 25 1,3-Dipropyl-8-[4-(3-p-tolyl-prop-2-ynyloxy)-phenyl]-3,7-dihydro-purine-2,6-dione,
 8-{4-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy]-prop-1-ynyl}-benzoic acid ethyl ester,
 30 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy]-prop-1-ynyl}-benzoic acid,
 1,3-Dipropyl-8-{4-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

- 8-{4-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-{4-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
- 5 8-{4-[3-(3-Methoxy-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{6-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-{3-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
- 10 1,3-Dipropyl-8-{3-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
- 8-{3-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 15 1,3-Dipropyl-8-{3-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-[3-(3-p-tolyl-prop-2-ynyloxy)-phenyl]-3,7-dihydro-purine-2,6-dione,
- 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 20 1-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-but-2-ynyl}-piperidine-3-carboxylic acid ethyl ester,
- 1-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-but-2-ynyl}-piperidine-3-carboxylic acid,
- 8-(1-{4-[4-(4-Chloro-phenyl)-4-hydroxy-piperidin-1-yl]-but-2-ynyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 25 1,3-Dipropyl-8-(1-{4-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-but-2-ynyl}-1H-pyrazol-4-yl)-3,7-dihydro-purine-2,6-dione,
- 1,3-Dimethyl-8-{1-[4-(4-methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
- 30 1-Propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,
- 1-Propyl-8-{1-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

- 1-Propyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
 5 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
 1-Propyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 1-Propyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 10 3-{3-[4-(2,6-Dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,
 8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 15 7-Methyl-1,3-dipropyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 7-Methyl-1,3-dipropyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,
 7-Methyl-8-{1-[4-(4-methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 20 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{1-[3-(3-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 25 8-{1-[3-(3-tert-Butyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 (3-{3-[4-(7-Methyl-2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenoxy)-acetic acid,
 8-{1-[3-(3-Hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 30 4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzenesulfonamide,

- 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzamide,
 4-(3-Trifluoromethyl-phenyl)-but-3-ynoic acid [5-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-2-methyl-2H-pyrazol-3-yl]-amide,
 5 4-(3-Fluoro-phenyl)-but-3-ynoic acid[5-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-2-methyl-2H-pyrazol-3-yl]-amide,
 4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-N-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-benzamide,
 8-{1-[4-(3-Fluoro-phenyl)-2-hydroxy-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-
 10 dihydro-purine-2,6-dione,
 2-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-4-(3-fluoro-phenyl)-but-3-ynoic acid,
 2-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-4-(3-fluoro-phenyl)-but-3-ynoic acid amide,
 15 8-{1-[4-(4-Fluoro-phenyl)-4-hydroxy-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{1-[4-(4-Fluoro-phenyl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{1-[3-(4-Fluoro-phenyl)-1,1-dimethyl-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-
 20 3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{1-[(E)-3-(3-trifluoromethyl-phenyl)-allyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{1-[(Z)-3-(3-trifluoromethyl-phenyl)-allyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
 25 1,3-dimethyl-6-[1-[3-(3-fluorophenyl)prop-2-ynyl]pyrazol-4-yl]-5H-pyrrolo[3,2-d]pyrimidine-2,4-dione,
 8-[1-[3-(3-fluorophenyl)prop-2-ynyl]pyrazol-4-yl]-1,3-dipropyl-7H-purine-2,6-dione,
 8-[1-[4-(3-fluorophenyl)-2-hydroxy-but-3-ynyl]pyrazol-4-yl]-1,3-dipropyl-7H-purine-2,6-dione,
 30 8-[1-[3-(4-fluorophenyl)-1,1-dimethyl-prop-2-ynyl]pyrazol-4-yl]-1,3-dipropyl-7H-purine-2,6-dione,
 1,3-Dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

- 8-{1-[2-(3-Fluoro-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[2-(4-Fluoro-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 5 1,3-Dipropyl-8-{1-[2-(4-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-
- 10 dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[5-Oxo-1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 15 8-[1-(5-Oxo-1-p-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-[1-(5-Oxo-1-m-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[5-Oxo-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-
- 20 1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(4-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 25 8-{1-[1-(3-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 3-Ethyl-8-{1-[1-(4-methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-
- 30 4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(4-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

- 4-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzonitrile, .
- 3-Ethyl-8-{1-[1-(4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 5 3-Ethyl-8-{1-[1-(3-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[5-Oxo-1-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 10 3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 3-Ethyl-8-{1-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 15 8-{1-[1-(3-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[5-Oxo-1-(2-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[5-Oxo-1-(4-trifluoromethoxy-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 20 1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(4-Methyl-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 4-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzoic acid,
- 25 8-{1-[1-(4-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 3-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzonitrile,
- 8-{1-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 30 1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(2,4-Difluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

- 8-{1-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(2-Chloro-4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
5. 8-{1-[1-(2-Chloro-4-trifluoromethyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 3-Ethyl-8-{1-[5-oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 10 8-{1-[1-(3-Fluoro-phenyl)-2-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[2-Oxo-1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 15 1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-{1-[1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-[1-(1-p-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,
- 20 8-{1-[1-(4-Methoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(3-Methoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 25 1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,
- 8-{4-[1-(4-Fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]phenyl}-1,3-dipropyl-3,7-dihydropurine-2,6-dione,
- 1,3-Dipropyl-8-{4-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
- 30 8-{3-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

- 8-{4-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{4-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 5 8-{6-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{3-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{3-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 10 8-{6-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{6-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 15 8-{4-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{4-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{6-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 20 8-{3-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{4-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 25 8-{3-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-{4-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
- 1,3-Dipropyl-8-{3-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,
- 30 8-{6-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

- 1,3-Dipropyl-8-{6-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,
 8-{6-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 5 1,3-Dipropyl-8-{6-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{6-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{4-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
 10 1,3-Dipropyl-8-{3-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{6-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,
 15 8-{6-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{6-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{3-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 20 8-{4-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 1,3-Dipropyl-8-{4-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,
 25 1,3-Dipropyl-8-{3-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,
 8-{3-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 8-{3-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
 30 1,3-Dipropyl-8-{3-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,


- 8-{4-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{3-[1-(2,4-Difluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 5 8-{5-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-Methyl-5-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethoxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{5-[1-(4-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 10 8-{5-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-Methyl-5-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 15 8-{5-[1-(3-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 20 1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-Oxo-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-Hydroxy-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 25 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-Oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-[4-(4-Chloro-phenyl)-4-hydroxy-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 30 8-(1-{2-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

- 8-{1-[2-Oxo-2-(4-p-tolyl-piperazin-1-yl)-ethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-Oxo-2-[4-(4-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 5 8-(1-{2-[4-(3-Chloro-4-fluoro-phenyl)-piperazin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 3-Ethyl-8-(1-{2-oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1-propyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-[4-(4-Fluoro-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-
- 10 dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-(1-{2-[4-(3-Fluoro-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid,
- 15 3-(2-Amino-ethyl)-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 7-Methyl-8-{1-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-
- 20 pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{1-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 7-Methyl-8-{1-[5-oxo-1-(2-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 25 8-{6-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{6-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,
- 8-{4-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-7-methyl-1,3-
- 30 dipropyl-3,7-dihydro-purine-2,6-dione, and
- 8-{4-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.

The present invention also relates to the process of preparation of compounds of formula (I), or pharmaceutically acceptable salts thereof.


The compounds of formula (I), may be prepared as outlined in the Scheme 1 and 3 below:

Scheme 1:




As exemplified in scheme 1 above, diamine of formula (II) wherein R¹ and R² are as defined herein above may be coupled with a carboxylic acid of formula (III) wherein all the symbols are as defined herein above and cyclised to provide compounds of formula (IV) which is reacted with R³-Hal wherein R³ is defined herein above to provide the desired compound of formula (I) wherein Y is N and all other symbols are as defined herein above.

Scheme 2:



As exemplified in scheme 2 above, nitro pyrimidine dione of formula (V) wherein R^1 and R^2 are as defined herein above may be condensed with an aldehyde of formula (VI) wherein all the symbols are as defined herein above, followed by reductive cyclisation to provide compound of formula (VII) which is reacted with R^3 -Hal wherein R^3 is defined herein above to provide the desired compound of formula (I) wherein Y is CH and all other symbols are as defined herein above.

Scheme 3:




As exemplified in scheme 3 above, compound of formula (VIII) wherein all the symbols are as defined herein above may be condensed with compound of formula (IX) wherein L_1 is a leaving group such as hydroxyl, mesylate or halo such as bromine, chlorine or iodine and all other symbols are as defined herein above to provide the desired compound of formula (I) wherein all symbols are as defined herein above.

15

Schemes 4 – 9 further describes synthesis of compounds of formula (I)

Scheme 4:



As exemplified in scheme 4, ethyl pyrazole-4-carboxylate of formula (1a) wherein all the symbols including R_a and R_b that are substituents on the pyrazole ring are defined herein above, may be reacted with compounds of formula (1g) wherein L_1 is hydroxyl or halo such as bromine, chlorine or iodine, in the presence of a suitable coupling agents such as triaryl phosphine/dialkyl azo dicarboxylate, for example, $PPh_3/DEAD$, $PPh_3/DIAD$ or K_2CO_3 /acetone and the like. Inert atmosphere may be maintained using nitrogen or argon. Solvents such as tetrahydrofuran, toluene, dioxane and the like may be used. The reaction temperature may range from room temperature to reflux temperature of the solvent(s) used, preferably at room temperature to $60^\circ C$. The reaction time may range from 1 to 48 hours, preferably 1 to 16 hours. After completion of reaction, the compounds of formula (1b) wherein all symbols are defined herein above may be isolated by conventional methods. Compounds of formula (1b) may be further reacted with $D-Hal$ wherein D is defined herein above to provide compounds of formula (1c) wherein all the symbols are defined herein above.

Alternatively, compounds of formula (1c) may be obtained by reacting, ethyl pyrazole-4-carboxylate of formula (1a) with compounds of formula (1h) wherein L_1 is a group such as hydroxyl, tosylate, mesylate or halogen such as bromine, chlorine or

iodine, in the presence of a base for example potassium carbonate in a polar solvent, for example N,N-dimethyl formamide, acetone and the like.

Compounds of formula (1c) may be hydrolysed by conventional methods for example in presence of a base, for example potassium hydroxide, sodium hydroxide, lithium hydroxide and the like in solvent such as methanol or ethanol, tetrahydrofuran or a mixture of ethanol, tetrahydrofuran and water, to obtain compound of formula (1d) wherein all symbols are defined herein above.

Compounds of formula (II) which are available commercially or may be prepared by procedures well known in the art may be reacted with carboxylic acids of formula (1d) wherein all symbols are defined herein above, to yield compounds of formula (1e). The reaction may be carried out with a suitable coupling agent for example EDCI, DCC, HBTU and the like in a protic solvent such as methanol, ethanol, propanol and the like or aprotic solvent such as DMF, CH₂Cl₂ and the like. The reaction temperature may range from 20-30°C. The reaction time may range from 4 to 16 hours. After completion of reaction, the product of formula (1e) wherein all symbols are defined herein above may be isolated by conventional methods.

Alternatively, compounds of formula (1e) may also be prepared from reaction of (II) and acid chloride (1d) to yield (1e). The reaction of acid halides of (1d) may be preferably carried out in the absence of solvent, using excess of halogenating agent such as thionyl chloride. The reaction temperature may range from 0°C to reflux temperature of the solvent or reagent used, preferably 60 to 70°C. The reaction time may range from 1 to 24 hours, preferably from 1 to 6 hours. After completion of the reaction the product may be isolated by removal of excess halogenating agent under reduced pressure. The isolated product may be treated with compounds of formula (II) in an inert solvent, for example acetonitrile, in the presence of tertiary base for example triethyl amine. The reaction temperature may range from 0 °C to reflux temperature of the solvent(s) used, preferably at 20-30 °C. The reaction time may range from 4 to 48 hours, preferably from 4 to 16 hours. After completion of reaction, the compounds of formula (1e) may be isolated by conventional methods.

Compounds of formula (1e) may be converted into compounds of formula (1f) by a cyclisation reaction. The reaction may be carried out in a protic solvent, such as methanol, ethanol, propanol and the like, preferably methanol, in presence of a base, such as alkali metal hydroxide such as potassium hydroxide, sodium hydroxide and the like, or sodium methoxide, sodium ethoxide, potassium tert-butoxide, preferable aqueous sodium hydroxide. The reaction temperature may range from 50-100°C, preferably at 80°C. The reaction time may range from 1 to 12 hours, preferably about 6-10 hours. After completion of reaction, the compounds of formula (1f) wherein all symbols are defined herein above may be isolated by conventional methods.


10

Compounds of formula (1f) may be reacted with R³-Hal wherein R³ is defined herein above. The reaction may be carried out in a solvent such as acetone or DMF, in the presence of a base such as K₂CO₃. The reaction temperature may range from room temperature to 80°C. The reaction time may range from 2 to 12 hours. After completion of reaction, the compounds of formula (I) wherein all symbols are defined herein above may be isolated by conventional methods.

15

Scheme 4 above can also be followed to obtain compounds wherein X of formula I is a aryl or heteroaryl rings such as phenyl, pyridyl, isoxazolyl, 3,5-pyrazolyl, oxazolyl and the like.

20




The compounds of formula (2b) may be converted to compounds of formula (2c) wherein all the symbols are defined herein above, by reaction with D-Hal wherein D is defined herein above by methods well known in the art.

5 Alternatively, compounds of formula (2a) may be converted to compounds of formula (2c) directly by reacting with compounds of formula (1h) wherein D is defined herein above and L2 is hydroxyl, tosylate, mesylate or halogen such as bromine, chlorine or iodine in the presence of a base for example potassium carbonate in a polar solvent, for example N,N-dimethyl formamide, acetone and the like. The reaction
10 temperature may range from 30⁰C to reflux temperature of the solvent(s) used, preferably at 20-60⁰C. The reaction time may range from 1 to 48 hours, preferably 1 to 16 hours. After completion of reaction, the product of formula (2c) wherein all symbols are defined herein above may be isolated by conventional methods.

15 The SEM (2-(trimethylsilyl)-ethoxymethyl chloride) protecting group of the compounds of formula (2c) may be deprotected by suitable acidic reagent to get compounds of formula (2d). In general, the compound of formula (2c) may be dissolved in ethanol and treated with HCl. The reaction temperature may range from 30⁰C to reflux temperature of the solvent(s) used, preferably at 80-100⁰C. The reaction
20 time may range from 1 to 6 hours. After completion of reaction, the product of general formula (2d) wherein all symbols are defined herein above may be isolated by conventional methods.

 Compounds of formula (2d) may be reacted with R³-Hal wherein R³ is defined
25 herein above. The reaction may be carried out in a solvent such as acetone or DMF, in the presence of a base such as K₂CO₃. The reaction temperature may range from room temperature to 80⁰C. The reaction time may range from 2 to 12 hours. After completion of reaction, the compounds of formula (I) wherein all symbols are defined herein above may be isolated by conventional methods.

30 Scheme 5 above can also be followed to obtain compounds wherein X of formula I is a aryl or heteroaryl rings such as phenyl, pyridyl, isoxazolyl, 3,5-pyrazolyl, oxazolyl and the like.



Compounds of formula (3b) may be reacted with N,N-dimethyl formamide, dimethyl acetal in a polar solvent, for example N,N-dimethyl formamide. The reaction temperature may range from 20 to 60°C, preferably at 40°C. The reaction time may range from 1 to 4 h hours, preferably 1 hour. After completion of reaction the compounds of formula (3c) may be reacted with R¹-Hal where Hal is chloro, bromo, or iodo in the presence of a base for example potassium carbonate in a polar solvent, for example N,N-dimethyl formamide, acetone and the like. The reaction temperature may range from 30°C to reflux temperature of the solvent(s) used, preferably at 70-80°C. The reaction time may range from 1 to 48 hours, preferably from 4 to 30 hours. After

completion of reaction, the compounds of formula (3d) wherein all symbols are defined herein above may be isolated by conventional methods.

Compounds of formula (3d) may be reacted with aqueous ammonia in a polar solvent, for example methanol. The reaction temperature may range from 20-30°C to
5 reflux temperature of the solvent(s) used, preferably at 20-30°C. The reaction time may range from 24 to 80 hours, preferably from 70 to 75 hours. After completion of reaction, the compounds of formula (3e) wherein all symbols are defined herein above may be isolated by conventional methods.


Compounds of formula (3e) may be reacted with sodium nitrite in aqueous
10 acidic solvent, for 50 % acetic acid in water. The reaction temperature may range from 50°C to reflux temperature of the solvent(s) used, preferably at about 75°C. The reaction time may range from 1 to 8 hours, preferably from 1 to 3 hours. After completion of reaction, the compounds of formula (3f) wherein all symbols are defined herein above may be isolated by conventional methods.

15 Compounds of formula (3f) may be reduced to (II) by procedures well known in the art.

Compounds of formula (II) may be converted to compounds of formula I wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above and R¹ is not equal to R², as described in scheme 3 herein
20 above.

Scheme 6 above can also be followed to obtain compounds wherein X of formula I is a aryl or heteroaryl rings such as phenyl, pyridyl, isoxazolyl, 3,5-pyrazolyl, oxazolyl and the like.

Scheme 7:



As exemplified in scheme 7, compounds of formula (3b) wherein R² is defined
 5 herein above, may be reacted with sodium nitrite in aqueous acidic solvent, for example
 50 % acetic acid in water. The reaction temperature may range from 50⁰C to reflux
 temperature of the solvent(s) used, preferably at about 75⁰C. The reaction time may
 range from 1 to 8 hours, preferably from 1 to 3 hours. After completion of reaction, the
 compounds of formula (3g) wherein R¹ is hydrogen and R² is defined herein above may
 10 be isolated by conventional methods.

Compounds of formula (3g) may be converted to diamine derivative by a
 reduction step. The reaction may be carried out by conventional reducing techniques
 for example sodium dithionate in aqueous ammonia solution; preferably, reduction is
 carried out with hydrogen and a metal catalyst. The reaction may be carried out in an
 15 inert solvent, for example methanol or ethanol, in the presence of a catalyst, for
 example 10% palladium on carbon catalyst, under an atmosphere of hydrogen,
 preferably under pressure for example about 20 to 40 psi for 2 to 18 hours. After
 completion of reaction, the compounds of formula (4a) wherein R² is defined herein
 above may be isolated by conventional methods.

20 Compound of formula (4a) may be reacted with carboxylic acids of formula
 (1d) wherein all symbols are defined herein above, to yield compounds of formula (4b)
 wherein all other symbols including Ra and Rb that are substituents on the pyrazole

ring are defined herein above. The reaction may be carried out with a suitable coupling agent for example EDCI, DCC, HOBt and the like, in protic solvent such as methanol, ethanol, propanol and the like or aprotic solvent such as DMF, CH₂Cl₂ and the like. The reaction temperature may range from 20-30°C. The reaction time may range from 4 to 16 hours. After completion of reaction, the compounds of formula (4b) wherein all other symbols are defined herein above may be isolated by conventional methods.

Compounds (4b) may also be prepared by reacting compounds of formula (4a) with the acid halide of the acid of formula (1d). The reaction of acid (1d) may be preferably carried out in the absence of solvent, using excess of halogenating agent (e.g. thionyl chloride). The reaction temperature may range from 0°C to reflux temperature of the solvent or reagent used, preferably 60 to 70°C. The reaction time may range from 1 to 24 hours, preferably from 1 to 6 hours. After completion of reaction, the product may be isolated by removal of excess halogenating agent under reduced pressure. The isolated product may be treated with compounds of formula (4a) in an inert solvent, for example acetonitrile, in the presence of tertiary base for example triethyl amine. The reaction temperature may range from 0 °C to reflux temperature of the solvent(s) used, preferably at 20-30°C. The reaction time may range from 4 to 48 hours, preferably from 4 to 16 hours. After completion of reaction the compounds of formula (4b) wherein all symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional methods.

The compounds of formula (4b) may be reacted with R¹-hal wherein R¹ is defined herein above and Hal is chloro, bromo, or iodo in the presence of a base for example potassium carbonate in a polar solvent, for example N,N-dimethyl formamide, acetone and the like. The reaction temperature may range from 30°C to reflux temperature of the solvent(s) used, preferably at 70-80°C. The reaction time may range from 1 to 48 hours, preferably from 4 to 30 hours. After completion of reaction, the product of formula (4c) wherein R¹ is different from R² wherein all symbols are defined herein above may be isolated by conventional method.


Compounds of formula (4c) may be then converted into compounds of general formula (4d) wherein R¹ is different from R², by a cyclisation reaction. The reaction may be carried out in a protic solvent, such as methanol, ethanol, propanol and the like, preferably methanol, in presence of a base, such as alkali metal hydroxide such as potassium hydroxide, sodium hydroxide and the like, or sodium methoxide, sodium

ethoxide, potassium tert-butoxide, preferable aqueous sodium hydroxide. The reaction temperature may range from 50-100°C, preferably at 80°C. The reaction time may range from 1 to 12 hours, preferably about 6-10 hours. After completion of reaction, the compounds of formula (4d) wherein all symbols are defined herein above may be isolated by conventional methods.

Compounds of formula (4d) may be reacted with R³-Hal wherein R³ is defined herein above. The reaction may be carried out in a solvent such as acetone or DMF, in the presence of a base such as K₂CO₃. The reaction temperature may range from room temperature to 80 °C. The reaction time may range from 2 to 12 hours. After completion of reaction, the compounds of formula (I) wherein all symbols are defined herein above may be isolated by conventional methods.

Scheme 7 above can also be followed to obtain compounds wherein X of formula I is a aryl or heteroaryl rings such as phenyl, pyridyl, isoxazolyl, 3,5-pyrazolyl, oxazolyl and the like.

15 Scheme- 8:



As exemplified in scheme 8, compounds of formula (1c) (obtained as mentioned in scheme 3) wherein all symbols are defined herein above may be converted into the analogous alcohol by a suitable reducing agent. In general, the compounds of formula (1c) may be dissolved in a suitable solvent, such as THF, and cooled to 0°C under inert atmosphere and treated with lithium aluminium hydride. The reaction temperature may

range from 0 to 25 °C. The reaction time may range from 1 to 6 hours, preferably 1 to 3 hours. After completion of reaction, the compounds of formula (5a) wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional methods.

5 Compounds of formula (5a) may be converted into the analogous aldehyde by a suitable oxidizing agent such as Dess-Martin Periodinane, manganese dioxide. In general, the compound of formula (5a) is dissolved in a suitable solvent, such as chloroform under inert atmosphere and treated with manganese dioxide portion wise. The reaction temperature may range from 30°C to reflux temperature of the solvent
10 used, preferably at 50-60°C. The reaction time may range from 1 to 24 hours, preferably 1 to 12 hours. After completion of reaction, compounds of formula (5b) wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional methods.

 The compound of formula (V) wherein all symbols are defined herein above
15 may be prepared by means well known in the art. Compounds of formula (5b) wherein all symbols are defined herein above may be condensed with compounds of formula (V) in the presence of a suitable base such as piperidine. The reaction may be carried out in an inert solvent, for example ethanol, 1,4-dioxane and the like. The reaction temperature may range from 60-80°C. The reaction time may range from 4 to 24 hours.
20 After completion of reaction, the compounds of formula (5c) wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional methods.

 The compounds of formula (5c) may be converted into compounds of formula (5d) by a reduction and cyclization reaction. The reaction may be carried out in a
25 suitable solvent such as N,N-dimethyl formamide (DMF) in presence of catalytic amount of SnCl₂. The reaction temperature may range from 100°C to reflux temperature of the solvent used, preferably 140-150°C. The reaction time may range from 1 to 10 hours, preferably from 1 to 6 hours. After cooling, acid solution such as 2N HCl is added to get compound of formula (5d) wherein all the symbols including
30 Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional method.

 Compounds of formula (5d) may be reacted with R³-Hal. The reaction may be carried out in a solvent such as acetone or DMF, in the presence of a base such as

K₂CO₃. The reaction temperature may range from room temperature to 80 °C. The reaction time may range from 2 to 12 hours. After completion of reaction, the compounds of formula (I) wherein all symbols are defined herein above may be isolated by conventional methods.

- 5 Scheme 8 above can also be followed to obtain compounds wherein X of formula I is a aryl or heteroaryl rings such as phenyl, pyridyl, isoxazolyl, 3,5-pyrazolyl, oxazolyl and the like.

Scheme 9:

- 10 As exemplified in scheme 9, compounds of formula (6a) wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above (may be prepared by means well known in the art J.Med.Chem. 2006, 49, 3682-3692), may be reacted with compound of formula (1g) wherein L1 is hydroxyl or halo such as bromine, chlorine or iodine, in the presence of suitable coupling agent such as triaryl phosphine/ dialkyl azo dicarboxylate, for example, PPh₃/DEAD, 15 PPh₃/DIAD, K₂CO₃ acetone and the like. Inert atmosphere may be maintained using nitrogen or argon. Solvents such as tetrahydrofuran, toluene, dioxane and the like may be used. The reaction temperature may range from 30°C to reflux temperature of the solvent(s) used, preferably at 20-60°C. The reaction time may range from 1 to 48 hours, preferably 1 to 16 hours. After completion of reaction compounds of formula (6b) 20 wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional methods.

Compounds of formula (6b) may be converted into compounds of formula (6c) by reaction with D-CH₂-Hal or D-Hal by well known methods in the art.

- 25 Alternatively, the compound of formula (6a) may be converted into a compound of formula (6c) directly by reacting with compounds of formula (1h) wherein L2 is a group such as hydroxyl, tosylate, mesylate or halogen such as bromine, chlorine or iodine, in the presence of a base for example potassium carbonate in a polar solvent, for example N,N-dimethyl formamide, acetone and the like. The reaction temperature may range from 30°C to reflux temperature of the solvent(s) used, preferably at 20-60°C. 30 The reaction time may range from 1 to 48 hours, preferably 1 to 16 hours. After completion of reaction, compounds of formula (6c) wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional methods.

The SEM (2-(trimethylsilyl)-ethoxymethyl chloride) protecting group of the compound of formula (6c) may be de-protected by suitable acidic reagent. In general, the compound of formula (6c) may be dissolved in ethanol and treated with 2N HCl. The reaction temperature may range from 30°C to reflux temperature of the solvent(s) used, preferably at 80-100°C. The reaction time may range from 1 to 6 hours, preferably 1 to 6 hours. After completion of reaction, compounds of formula (6d) wherein all the symbols including Ra and Rb that are substituents on the pyrazole ring are defined herein above may be isolated by conventional method.

Compounds of formula (6d) may be reacted with R³-Hal. The reaction may be carried out in a solvent such as acetone or DMF, in the presence of a base such as K₂CO₃. The reaction temperature may range from room temperature to 80°C. The reaction time may range from 2 to 12 hours. After completion of reaction, the compounds of formula (I) wherein all symbols are defined herein above may be isolated by conventional methods.

Scheme 9 above can also be followed to obtain compounds wherein X of formula I is a aryl or heteroaryl rings such as phenyl, pyridyl, isoxazolyl, 3,5-pyrazolyl, oxazolyl and the like.

Wherever desired or necessary, in any of the above mentioned processes, any of the compounds of formula (I) may be converted into a pharmaceutically acceptable salt or vice versa or converting one salt form into another pharmaceutically acceptable salt form.

According to an embodiment, the compounds of the present invention are adenosine A_{2B} receptor antagonists. Thus, the present invention provides a method for the modulation of adenosine A_{2B} receptor activity in mammals which method comprises administering to a mammal in need thereof a therapeutically effective amount of compound of formula I or its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof.

According to yet another embodiment, the compounds of the present invention are adenosine A₁ and A_{2B} antagonist or adenosine A_{2B} and A₃ antagonist or A₁, A_{2B} and A₃ antagonist thereby providing dual or pan antagonistic activity through additive/synergistic effect. Thus, the present invention provides a method for the modulation of adenosine A₁ and A_{2B} or A_{2B} and A₃ or A_{2B}, A₁ and A₃ receptor activity in mammals which method comprises administering to a mammal in need thereof a

therapeutically effective amount of compound of formula I or its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof.

The present invention also provides a method of prophylactic or therapeutic
5 treatment of disease or disorder susceptible to improvement by antagonism of adenosine receptor comprising administering an effective amount of compound of formula I or its tautomers, polymorphs, stereoisomers, prodrugs, solvate or a pharmaceutically acceptable salts thereof, to a mammal in need of such treatment.

As used throughout the specification and in the claims, the term "treatment"
10 embraces all the different forms or modes of treatment as known to those of the pertinent art and in particular includes preventive, curative, delay of progression and palliative treatment.

The term "therapeutically effective amount" as used herein refers to an amount of a drug or a therapeutic agent that will elicit the desired biological or medical
15 response of a tissue, system or an animal (including man) that is being sought by a researcher or clinician.

The term "mammal" or "patient" are used interchangeably herein and include, but are not limited to, humans, dogs, cats, horses, pigs, cows, sheep, monkeys, rabbits, mice and laboratory animals. The preferred mammals are humans.

20 The present invention further provides pharmaceutical compositions comprising a therapeutically effective amount of a compound of the present invention, alone or in combination with one or more pharmaceutically acceptable carriers.

The pharmaceutical compositions according to the invention are those suitable for enteral, such as oral or rectal, transdermal and parenteral administration to
25 mammals, including man, for the treatment of conditions mediated by the adenosine A_{2B} receptor. Such conditions include, but are not limited to, asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity,
30 inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.

Generally, the concentration of the compound(s) of the present invention in a liquid composition, such as a lotion, will be from about 0.01- about 25 wt %, preferably from about 0.1- about 10 wt %. The concentration in a semi-solid or a solid

composition such as a gel or a powder will be about 0.1- about 5 wt %, preferably about 0.5- about 25 wt %.

The amount of a compound of the present invention required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will be ultimately at the discretion of the administering physician or clinician. In general, a suitable dose will be in the range of from about 0.001 mg/kg/day to about 20 mg/kg/day For example, a dosage may be from about 0.002 mg/kg to about 10 mg/kg of body weight per day, from about 0.01 mg/kg/day to about 1 mg/kg/day, and from about 0.1 mg/kg/day to about 5 mg/kg/day.

The compound is conveniently administered in unit dosage form, e.g., containing 5 to 1000 µg, about 10 to about 750 µg, about 50 to about 500 µg of active ingredient per unit dosage form.

The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, e g , into a number of discrete loosely spaced administrations; such as multiple inhalations from an insufflator or by application of a plurality of drops into the eye Dosages above or below the range cited herein above are within the scope of the present invention and may be administered to the individual patient if desired and necessary.

Accordingly, in various embodiments, the present invention provides pharmaceutical compositions as described above for the treatment of conditions mediated by adenosine receptor, such as asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.

In various embodiments, the present invention provides pharmaceutical compositions comprising a therapeutically effective amount of a compound of the invention in combination with a therapeutically effective amount of another therapeutic agent, preferably selected from anti-inflammatory agents, anti-diabetic agents, anti-hypertensive agents and anti-dyslipidemic agents.

54348-1

According to an embodiment, the pharmaceutical compositions may contain a therapeutically effective amount of a compound of the invention as defined above, either alone or in a combination with another therapeutic agent, e.g., each at an effective therapeutic dose as reported in the art. Such therapeutic agents include: a) anti-inflammatory agents, such as anticholinergic or antimuscarinic agents; steroids; LTB₄ (leukotriene B₄) antagonists; dopamine receptor agonists; PDE₄ (phosphodiesterase 4) inhibitors; and beta-2 adrenergic receptor agonists; b) anti-diabetic agents, such as insulin, insulin derivatives and mimetics; insulin secretagogues; insulinotropic sulfonylurea receptor ligands; thiazolidone derivatives; GSK3 (glycogen synthase kinase-3) inhibitors; sodium-dependent glucose co-transporter inhibitors; glycogen phosphorylase A inhibitors; biguanides; alpha-glucosidase inhibitors; GLP-1 (glucagon like peptide-1), GLP-1 analogs and GLP-1 mimetics; modulators of PPARs (peroxisome proliferator-activated receptors); DPPIV (dipeptidyl peptidase IV) inhibitors; SCD-1 (stearoyl-CoA desaturase-1) inhibitors; DGAT1 and DGAT2 (diacylglycerol acyltransferase 1 and 2) inhibitors; ACC2 (acetyl CoA carboxylase 2) inhibitors; and breakers of AGE (advanced glycation end products); c) anti-hypertensive agents, such as loop diuretics; angiotensin converting enzyme (ACE) inhibitors; inhibitors of the Na-K-ATPase membrane pump such as digoxin; neutralendopeptidase (NEP) inhibitors; ACE/NEP inhibitors; angiotensin II antagonists; renin inhibitors; β -adrenergic receptor blockers; inotropic agents; calcium channel blockers; aldosterone receptor antagonists; and aldosterone synthase inhibitors; and d) anti-dyslipidemic agents such as 3-hydroxy-3-methyl-glutaryl coenzyme A (HMG-CoA) reductase inhibitors; HDL increasing compounds such as cholesterol ester transfer protein (CETP) inhibitors; squalene synthase inhibitors; FXR (farnesoid X receptor) and LXR (liver X receptor) ligands; cholestyramine; fibrates; nicotinic acid; and aspirinTM.

As described above, a compound of the present invention may be administered either simultaneously, before or after another active ingredient, either separately by the same or different route of administration or together in the same pharmaceutical formulation.

In another embodiment of the present invention, the compound of formula (I) is use in treatment conditions mediated by adenosine receptor.

The present invention further relates to use of compound of formula (I) for the treatment of asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.

The present invention further relates to use of compound of formula (I) for use in preparation of medicament useful in the treatment of conditions mediated by adenosine receptor.

The present invention further relates to use of compound of formula (I) in preparation of medicament useful in the treatment of asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.


The present invention further relates to pharmaceutical compositions as described above for use as a medicament.

The present invention further relates to use of pharmaceutical compositions or combinations as described above for the preparation of a medicament for the treatment of conditions mediated by adenosine receptor, such as asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, allergic diseases, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, and/or autoimmune diseases.

Examples

The invention is further illustrated by the following examples which in no way should be construed as being further limiting. One skilled in the art will readily appreciate that the specific methods and results described are merely illustrative. Structures of the intermediates as well as the final compounds were confirmed by nuclear magnetic resonance spectra for proton (^1H NMR) and LCMS.

Example A1: 1, 3-Dipropyl-8-[1-(3-p-tolyl-prop-2ynyl)-1H-pyrazol-4-yl]-3, 7-dihydro-purine-2, 6-dione



5 Step I: 1-(3-p-Tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid ethyl ester

A mixture of 1-prop-2-ynyl-1H-pyrazole-4-carboxylic acid ethyl ester obtained as given in example B1 (0.200g, 1.1mmol), 4-iodo toluene (0.254g, 1.1mol), copper iodide (0.021g, 0.11mmol), dichlorobis (triphenylphosphine)-palladium (II) (39mg, 0.06mmol), triethylamine (2ml), DMF (2ml) was degassed for 10min. and stirred for
 10 20hrs at 25-25 °C. Reaction mixture was diluted with water (10ml) and extracted with ethyl acetate. Organic layer was washed with brine solution and dried over Na₂SO₄. The solvent was evaporated and crude product was purified by column chromatography (Ethyl acetate: hexane-12:78) to obtain pure 1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid ethyl ester compound (0.226g, 75%).

15 ¹HNMR(400MHz, CDCl₃): δ 1.35 (t, J=6.8Hz, 3H); 2.37 (s, 3H); 4.31 (q, J=6.8Hz, 2H); 5.18 (s, 2H); 7.16 (d, J=7.6Hz, 2H); 7.38 (d, J=8Hz, 2H); 7.95 (s, 1H); 8.21 (s, 1H)

Step II: 1-(3-p-Tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid

1-(3-p-Tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid ethyl ester (0.226g, 0.84
 20 mmol) was dissolved in a mixture of solvents THF: methanol: water (3:1:1, 10ml) and LiOH (0.071g, 1.7mol) was added to the reaction mixture with stirring. The reaction mixture was then stirred at 20-25 °C for 2 hours. Solvents were evaporated and the residue was diluted with water (0.5 ml) and acidified with dil. HCl, filtered and dried to

obtain off white precipitate, 1-(3-p-Tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid (0.182g, 90%).


¹HNMR(400MHz, CDCl₃): δ 2.37 (s, 3H); 5.2 (s, 2H); 7.16 (d, J=7.6Hz, 2H); 7.38 (d, J=8Hz, 2H); 8.01 (s, 1H); 8.29 (s, 1H)

5 Step III: 1, 3-Dipropyl-8-[1-(3-p-tolyl-prop-2ynyl)-1H-pyrazol-4-yl]-3, 7-dihydro-purine-2, 6-dione

A mixture of 5,6-diamino-1,3-dipropyl-1H-pyrimidine-2,4-dione (0.075g, 0.33 mmol), 1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid (0.080gm, 0.33mmol), methanol (5ml), EDCI (0.089g, 0.46mmol) were taken and stirred for 12 hours at 20-25
 10 °C. The reaction mixture was concentrated to obtain intermediate 1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazole-4-carboxylic acid (6-amino-2, 4-dioxo-1, 3-dipropyl)-1, 2, 3, 4-tetrahydro-pyrimidine-5yl) amide (50mg, 34%) which was dissolved in hexamethyldisilazane (HMDS). To this reaction mixture ammonium sulphate (0.010g) was added. The reaction mixture was refluxed at 140 °C for 18hrs. The organic
 15 volatiles were evaporated and the residue was treated with crushed ice, the precipitate formed was filtered off. The product was then purified by column chromatography (1%MeOH in CHCl₃) to obtain 1, 3-dipropyl-8-[1-(3-p-tolyl-prop-2ynyl)-1H-pyrazol-4-yl]-3, 7-dihydro-purine-2, 6-dione (0.035g, 92%).

¹HNMR(400MHz, DMSO d₆): δ 0.76-0.87 (m, 6H); 1.51-1.57 (m, 2H); 1.68-1.74 (m, 2H); 2.29 (s, 3H); 3.82 (t, J=7.2Hz, 2H); 3.95 (t, J=7.2Hz, 2H); 5.36 (s, 2H); 7.18 (d, J=8Hz, 2H); 7.35 (d, J=8Hz, 2H); 8.08 (s, 1H); 8.49 (s, 1H); 13.9 (bs, 1H)
 20

Example B1: 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1, 3-dimethyl-3, 7-dihydro-purine-2, 6-dione



3-(3-Fluoro-phenyl)-prop-2-yn-1-ol

A mixture of propargyl alcohol (0.200g, 5.4mmol), 3-fluoro iodobenzene (0.63ml, 5.4mmol), copper iodide (0.103g, 0.54mol), dichlorobis (triphenylphosphine) palladium (II) (0.190g, 0.3mmol), diethylamine (10ml) was degassed for 10min. and stirred for 20hrs at 25-25 °C. Excess of diethyl amine was distilled off under vacuum. The residue was diluted with water (10ml) and extracted with ethyl acetate. The organic layer was washed with brine solution and dried over Na₂SO₄. The solvent was evaporated and the crude product was purified by column chromatography (10% Ethyl acetate in hexane) to obtain pure 3-(3-fluoro-phenyl)-prop-2-yn-1-ol (0.750g, 93%).

¹HNMR(400MHz, CDCl₃): δ 1.68 (t, J=4.8Hz, 1H); 4.50 (d, J=4.8Hz, 2H); 7.04-7.05 (m, 1H); 7.13-7.15 (m, 1H); 7.21-7.23 (m, 1H); 7.27-7.30 (m, 1H)

1-(3-Bromo-prop-1-ynyl)-3-fluoro benzene

3-(3-Fluoro-phenyl)-prop-2-yn-1-ol (0.750g, 0.005mol) was taken in diethyl ether (10ml) containing catalytic amount of pyridine (0.05ml). The reaction mixture was then cooled to -5-0 °C. Tribromo phosphine (0.26ml, 0.003mol) was added slowly at -5-0 °C and stirred at the same temperature for 3 hrs. The reaction mixture was quenched with saturated NaHCO₃ solution and extracted with ethyl acetate. The organic layer was washed with saturated brine solution, dried over Na₂SO₄, and evaporated under vacuum to get 1-(3-bromo-prop-1-ynyl)-3-fluoro benzene (0.805gm, 76%).

¹HNMR(400MHz, CDCl₃): δ 4.15 (s, 2H); 7.04-7.08 (m, 1H); 7.13-7.16 (m, 1H); 7.22-7.24 (m, 1H); 7.28-7.32 (m, 1H)

Step I: 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

A mixture of 1, 3-dimethyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione (obtained following literature procedure US6825349) (0.100g, 0.27mmol), 1-(3-Bromo-prop-1-ynyl)-3-fluoro benzene (0.057g, 0.27mmol), K₂CO₃ (0.086mg, 0.62mmol) and acetone (5ml) were refluxed at 80° C for 5hrs. The reaction mixture was cooled to room temperature and filtered off solid K₂CO₃ and evaporated excess of organic solvent under vacuum. The product was purified by column chromatography (2% methanol in dichloromethane) to get pure 8-{1-[3-(3-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.111g, 82%).

¹HNMR(400MHz, CDCl₃): δ 0.12 (s, 9H); 0.96 (t, J=7.2Hz, 2H); 3.45 (s, 3H); 3.66 (s, 3H); 3.82 (t, J=8.4Hz, 2H); 5.26 (s, 2H); 5.82 (s, 2H); 7.09-7.1 (m, 1H); 7.20 (m, 1H); 7.28-7.33 (m, 2H); 8.20 (s, 1H); 8.43 (s, 1H)

Step II: 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1, 3-dimethyl-3, 7-dihydro-purine-2, 6-dione

8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.100g, 0.19mmol) in EtOH (5 mL) was treated with 2N HCl(1.1mL) and heated at 80 °C for 2hrs. The reaction mixture was concentrated in vacuo, and the residue was triturated with diethyl ether to afford 8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1, 3-dimethyl-3, 7-dihydro-purine-2, 6-dione in (0.040mg, 54%) as a white solid.

¹HNMR(400 MHz, DMSO d₆): δ 3.26 (s, 3H); 3.47 (s, 3H); 5.42 (s, 2H); 7.29-7.38 (m, 3H); 7.43-7.47 (m, 1H); 8.13 (s, 1H); 8.53 (s, 1H); 13.6 (bs, 1H)

Examples B2 - B58 were prepared in an analogous manner of Example B1 from the appropriate intermediates.

Example	IUPAC name
B2	8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione
B3	8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B4	8-{1-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione


B5	8-{1-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B6	8-{1-[3-(2,4-Difluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B7	8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B8	8-{1-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B9	8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione
B10	4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid ethyl ester
B11	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid ethyl ester
B12	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzonitrile
B13	8-{1-[3-(3-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B14	2-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid methyl ester
B15	8-{1-[4-(4-Fluoro-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
B16	8-{1-[4-(3-Fluoro-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
B17	1,3-Dipropyl-8-[1-(4-p-tolyl-but-3-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione.
B18	1,3-Dipropyl-8-{1-[4-(3-trifluoromethyl-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
B19	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid.
B20	1,3-Dipropyl-8-{1-[3-(2-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.

B21	1,3-Dipropyl-8-[1-(3-m-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione.
B22	3-Ethyl-1-propyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
B23	3-Ethyl-1-propyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
B24	3-Ethyl-1-propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione.
B25	3-Ethyl-8-{1-[3-(3-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
B26	1,3-Dipropyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
B27	1,3-Dipropyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
B28	3-Ethyl-8-{1-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
B29	3-{3-[4-(3-Ethyl-2,6-dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid.
B30	3-Ethyl-1-propyl-8-[1-(3-m-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione.
B31	3-Ethyl-8-{1-[3-(4-methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
B32	3-Ethyl-1-propyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
B33	3-Ethyl-8-{1-[3-(3-methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
B34	4-{3-[4-(3-Ethyl-2,6-dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid.
B35	4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzonitrile.
B36	(3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenoxy)-acetic acid.

B37	8-{1-[3-(3-tert-Butyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
B38	4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid.
B39	(3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenyl)-acetic acid.
B40	(4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenyl)-acetic acid.
B41	8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B42	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-N-isopropyl-benzamide
B43	1,3-Dipropyl-8-(1-{3-[3-(pyrrolidine-1-carbonyl)-phenyl]-prop-2-ynyl}-1H-pyrazol-4-yl)-3,7-dihydro-purine-2,6-dione
B44	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-4-methyl-benzoic acid
B45	8-{1-[3-(3-Chloro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B46	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-4-methoxy-benzoic acid
B47	5-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-pyridine-2-carboxylic acid methyl ester
B48	1,3-Dipropyl-8-{3-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione
B49	8-{3-[3-(2,4-Difluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B50	8-{3-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B51	8-{3-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B52	1,3-Dipropyl-8-{3-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione

B53	1,3-Dipropyl-8-[3-(3-p-tolyl-prop-2-ynyloxy)-isoxazol-5-yl]-3,7-dihydro-purine-2,6-dione
B54	8-{3-[3-(3-tert-Butyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B55	1,3-Dipropyl-8-{3-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione
B56	8-{1-Methyl-5-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B57	8-{1-Methyl-5-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
B58	8-{5-[3-(3-Methoxy-phenyl)-prop-2-ynyloxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

Example C1: 1,3-Dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione



Step-I: 4-Benzyloxy-benzoic acid methyl ester

- 5 A mixture of methyl 4-hydroxy-benzoate (10.0 g, 0.066mol) and potassium carbonate (20.9 g, 0.098mol), were taken in acetone (50 ml) and heated at 50 °C for 1 hour. To the reaction mixture, benzyl bromide (8.6 ml, 0.072 mol) was added and heated at 80 °C for 5 hours. The mixture was cooled to room temperature and filtered, washed with acetone. Solvent was removed to obtain pure 4-benzyloxy-benzoic acid methyl ester as
- 10 white solid (16 g, 100 %).

¹HNMR(400MHz, CDCl₃): δ 3.85 (s, 3H); 5.17 (s, 2H); 6.99 (d, J=8.8Hz, 2H); 7.38-7.42 (m, 5H); 8.00 (d, J=8.8Hz, 2H)

Step II: 4-Benzyloxy-benzoic acid

4-Benzyloxy-benzoic acid methyl ester (8.0 g, 0.033mol) was dissolved in THF: methanol: water (3:2:1, 80ml) and NaOH (1.98g, 0.05mol) was added to the reaction mixture and stirred at 50-55 °C for 3 hours. Solvents were removed and residue was
5 diluted with water, washed with hexane and acidified with dil HCl to obtain 4-benzyloxy-benzoic acid (7.0 g, 92 %) as a white solid.

¹HNMR(400MHz, CDCl₃): δ 5.14 (s, 2H); 7.02 (d, J=8.8Hz, 2H); 7.4-7.42 (m, 5H); 8.05 (d, J=8.8Hz, 2H).

Step III: 8-(4-Benzyloxy-phenyl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.

10 A mixture of 4-benzyloxy-benzoic acid (0.868 g, 3.8 mmol), 5,6-diamino-1,3-dipropyl-1H-pyrimidine-2,4-dione (1.0g, 3.8 mmol), methanol (15ml), EDCI (0.95 g, 4.9 mmol), was stirred for 20 hours at room temperature. The reaction mixture was concentrated and water was added to obtain the intermediate N-(6-amino-2,4-dioxo-1,3-dipropyl-1,2,3,4-tetrahydro-pyrimidin-5-yl)-4-benzyloxy-benzamide. It was
15 dissolved in methanol (32 ml) and added 10% NaOH (24ml). The mixture was heated at 80-90 °C for 3 hours. The mixture was cooled and solvent was removed. The residue was dissolved in water (20 ml) acidified with dil HCl to provide 8-(4-benzyloxy-phenyl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (1.2 g , 75 %) as a white solid.

¹HNMR(400MHz, DMSO d₆): δ 0.82-0.89 (m, 6H); 1.52-1.57 (m, 2H); 1.68-1.73 (m, 2H); 3.83 (t, J=7.2Hz, 2H); 3.98 (t, J=6.8Hz, 2H); 5.14 (s, 2H); 7.09 (d, J=8.8Hz, 2H);
20 7.3-7.45 (m, 5H); 8.03 (d, J=8.8Hz, 2H); 13.58 (bs, 1H)

Step IV: 8-(4-Benzyloxy-phenyl)-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

A mixture of 8-(4-benzyloxy-phenyl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (1.2
25 g, 2.9 mmol) and potassium carbonate (1.19 g, 8.6 mmol), were taken in DMF (15 ml) and 2-(trimethylsilyl) ethoxymethyl chloride (1.52 ml, 8.6 mmol) was added drop wise at 0° C and mixture was stirred at room temperature for 20 hours. The mixture was cooled to 10 °C and diluted with water, extracted with ethyl acetate. The organic layer was washed with saturated brine solution and dried over Na₂SO₄. Solvent was removed
30 and residue obtained was further purified by column chromatography to obtain pure 8-(4-benzyloxy-phenyl)-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (1.12 g, 71 %).

¹HNMR(400MHz, CDCl₃): δ 0.12 (s, 9H); 0.94-1.00 (m, 8H); 1.66-1.72 (m, 2H); 1.81-1.86 (m, 2H); 3.89 (t, J=8.4Hz, 2H); 4.00 (t, J=7.6Hz, 2H); 4.13 (t, J=7.2Hz, 2H); 4.70 (s, 2H); 5.71 (s, 2H); 7.08 (d, J=8.8Hz, 2H); 7.34-7.45 (m, 5H); 7.93 (d, J=8.8Hz, 2H).

Step V: 8-(4-Hydroxy-phenyl)-1, 3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione

To a solution of 8-(4-benzyloxy-phenyl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.500 g, 0.91 mmol) in methanol (10ml), 10% Pd/C (0.150 g) was added and stirred at room temperature under hydrogen atmosphere for 2 hours. The mixture was filtered, solvent was removed and the residue was washed with hexane to obtain 8-(4-hydroxy-phenyl)-1, 3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione as a white solid (0.398 g, 95 %).

¹HNMR(400MHz, CDCl₃): δ 0.12 (s, 9H); 0.94-1.00 (m, 8H); 1.66-1.72 (m, 2H); 1.81-1.86 (m, 2H); 3.89 (t, J=8.4Hz, 2H); 4.00 (t, J=7.6Hz, 2H); 4.13 (t, J=7.2Hz, 2H); 5.71 (s, 2H); 7.08 (d, J=8.8Hz, 2H); 7.93 (d, J=8.8Hz, 2H).

Step VI: 1, 3-Dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione

A mixture of 8-(4-hydroxy-phenyl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.100 g, 0.22 mmol) and potassium carbonate (0.076g, 0.53mmol), 1-(3-bromo-prop-1-ynyl)-4-trifluoromethoxy-benzene (prepared as given in example E1) (0.073g, 0.26 mmol) in acetone (10 ml) was heated at 80 °C for 2 hours. The mixture was cooled to room temperature, filtered and washed with acetone. Solvent was evaporated to obtain a residue which was purified by preparative TLC to obtain 1, 3-dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione as a pale yellow solid (0.053 g, 37 %).

¹HNMR(400MHz, CDCl₃): δ 0.10 (s, 9H); 0.97-1.05 (m, 8H); 1.71-1.74 (m, 2H); 1.86-1.88 (m, 2H); 3.90 (t, J=8Hz, 2H); 4.03 (t, J=7.2Hz, 2H); 4.22 (t, J=7.2Hz, 2H); 5.01 (s, 2H); 5.75 (s, 2H); 7.18 (d, J=8.8Hz, 2H); 7.26 (d, J=8.8Hz, 2H); 7.49 (d, J=8.8Hz, 2H); 8.04 (d, J=8.8Hz, 2H)

Step VII: 1,3-Dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione.

A mixture of 1,3-dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.053 g, 0.055 mmol), 2N HCl (2 ml), ethanol (2 ml) was heated at 85 °C for 2 hours. The mixture was cooled and the solvent was evaporated. The residue was washed with n-pentane to obtain 1,3-dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione (0.038 g, 88 %).


¹HNMR(400MHz, DMSO d₆): δ 0.84-0.90 (m, 6H); 1.54-1.559 (m, 2H); 1.7-1.75 (m, 2H); 3.85 (t, J=7.6Hz, 2H); 3.99 (t, J=7.2Hz, 2H); 5.13 (s, 2H); 7.17 (d, J=8.8Hz, 2H); 7.37 (d, J=8.4Hz, 2H); 7.59 (d, J=8.4Hz, 2H); 8.09 (d, J=8.8Hz, 2H); 13.61 (s, 1H)

Examples C2 – C15 were prepared in an analogous manner of Example C1 from the appropriate intermediates.

Example	IUPAC name
C2	1,3-Dipropyl-8-[4-(3-p-tolyl-prop-2-ynyloxy)-phenyl]-3,7-dihydro-purine-2,6-dione
C3	8-{4-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
C4	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy]-prop-1-ynyl}-benzoic acid ethyl ester
C5	3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy]-prop-1-ynyl}-benzoic acid
C6	1,3-Dipropyl-8-{4-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione
C7	8-{4-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
C8	1,3-Dipropyl-8-{4-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione
C9	8-{4-[3-(3-Methoxy-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
C10	8-{6-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
C11	1,3-Dipropyl-8-{3-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione

C12	1,3-Dipropyl-8-{3-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione
C13	8-{3-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
C14	1,3-Dipropyl-8-{3-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione
C15	1,3-Dipropyl-8-[3-(3-p-tolyl-prop-2-ynyloxy)-phenyl]-3,7-dihydro-purine-2,6-dione

Example D1: 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.



5

Step I: 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyloxyethyl)-3,7-dihydro-purine-2,6-dione

A mixture of 1,3-dipropyl-8-(1-prop-2-ynyl-1H-pyrazol-4-yl)-7-(2-trimethylsilanyloxyethyl)-3,7-dihydro-purine-2,6-dione (0.1 g, 0.212 mmol), N-methyl piperazine (0.025 g, 0.254 mmol), formaldehyde (0.08 ml), CuI (0.008 g), and DMSO (2 ml) was stirred at 30 °C for 72 hours. The mixture was cooled to 25 °C and the residue was diluted with water (10 ml) and extracted with ethylacetate (2x10 ml). The organic layer was washed with saturated brine solution (50 ml) and dried over Na₂SO₄. The solvent was evaporated to obtain pure 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-

pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.105 g, 85 %).

¹HNMR(400MHz, DMSO d₆): δ 0.10 (s, 9H); 0.82–0.93 (m, 8H); 1.54–1.57 (m, 2H); 1.70–1.72 (m, 2H); 2.49–2.53 (m, 8H); 3.29 (s, 3H); 3.39–3.41 (m, 2H); 3.66 (t, J=8Hz, 2H); 3.86 (t, J=6.4Hz, 2H); 3.96–4.0 (m, 2H); 5.19 (s, 2H); 5.76 (s, 2H); 8.02 (s, 1H); 8.40 (s, 1H).

Step II: 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione


A mixture of 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.105 g, 0.180 mmol), 2N HCl (2 ml), ethanol (4 ml) was heated at 85 °C for 2 hours. The mixture was cooled and solvent was evaporated. The residue was triturated with diethyl ether and purified by preparative TLC to obtain pure 8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (0.042g, 52 %).

¹HNMR(400MHz, DMSO d₆): δ 0.82–0.89 (m, 6H); 1.51–1.53 (m, 2H); 1.68–1.70 (m, 2H); 2.12 (s, 2H); 2.44–2.50 (m, 8H); 3.31 (s, 3H); 3.81 (t, J=6.8Hz, 2H); 3.92 (t, J=6.8Hz, 2H); 5.07 (s, 2H); 7.82 (s, 1H); 8.09 (s, 1H).

Examples D2 – D6 were prepared in an analogous manner of Example D1 from the appropriate intermediates.

Example	IUPAC name
D2	1-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-but-2-ynyl}-piperidine-3-carboxylic acid ethyl ester.
D3	1-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-but-2-ynyl}-piperidine-3-carboxylic acid.
D4	8-(1-{4-[4-(4-Chloro-phenyl)-4-hydroxy-piperidin-1-yl]-but-2-ynyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
D5	1,3-Dipropyl-8-(1-{4-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-but-2-ynyl}-1H-pyrazol-4-yl)-3,7-dihydro-purine-2,6-dione.
D6	1,3-Dimethyl-8-{1-[4-(4-methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione

Example E1: 1-Propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione



Step I: 1-Propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-bis-(2-trimethylsilylanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

A mixture of 1-propyl-8-(1H-pyrazol-4-yl)-3,7-bis-(2-trimethylsilylanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (prepared as per literature procedure US20080194593) (0.1g, 0.18 mmol), 1-(3-bromo-prop-1-ynyl)-4-methyl-benzene (prepared according to example E1) (0.039g, 0.18 mmol), potassium carbonate (0.051g, 0.37 mmol), and acetone (10 ml) were heated at 50-55 °C for 16 hours. Reaction mixture was cooled to 20-25 °C and filtered off. The filtrate was evaporated and residue was purified by preparative TLC to afford 1-propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-bis-(2-trimethylsilylanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.035 g, 28%) as a brown sticky mass.

¹HNMR(400MHz, DMSO d₆): δ 0.12 (s, 18H); 0.91-1.02 (m, 7H); 1.68-1.69 (m, 2H); 2.36 (s, 3H); 3.75-3.80 (m, 4H); 3.99 (t, J=7.6Hz, 2H); 5.22 (s, 2H); 5.6 (s, 2H); 5.78 (s, 2H); 7.13 (d, J=7.6Hz, 2H); 7.37 (d, J=7.6Hz, 2H); 8.15 (s, 1H); 8.42 (s, 1H).

Step II: 1-Propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione.

A mixture of 1-propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-bis-(2-trimethylsilylanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione obtained in step 1 (0.035 g, 0.053 mmol;), 2 N HCl (1.5 ml), ethanol (3 ml) were heated at 85 °C for 6 hours.


The mixture was cooled to 10-15 °C. The solid obtained was filtered and washed with water (1ml), ethanol (1ml) to afford 1-propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione (0.009 g, 47 %) as a brown solid.

¹HNMR(400MHz, DMSO d6): δ 0.84 (t, J=7.2Hz, 3H); 1.50-1.55 (m, 2H); 2.24 (s, 3H); 3.76 (t, J=7.2Hz, 2H); 5.34 (s, 2H); 7.17-7.19 (m, 2H); 7.34-7.36 (m, 2H); 8.04 (s, 1H); 8.44 (s, 1H); 11.81 (bs, 1H); 13.39 (bs,1H)

Examples E2 - E8 were prepared in an analogous manner of Example E1 from the appropriate intermediate.

Example	IUPAC name
E2	1-Propyl-8-{1-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
E3	1-Propyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
E4	8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
E5	8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
E6	1-Propyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
E7	1-Propyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
E8	3-{3-[4-(2,6-Dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid.

10 **Example F1: 8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.**



A mixture of 8-{1-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (0.035g, 0.08 mmol), potassium carbonate (0.022g, 0.16


mmol), methyl iodide (0.005 ml, 0.088 mmol) and DMF (2 ml) were heated at 50 °C for 20 hour. The mixture was cooled to room temperature and diluted with water (10 ml). The aqueous layer was extracted with ethyl acetate. The organic layer was washed with water and sat. brine solution, and dried over Na₂SO₄. Solvent was evaporated, the residue obtained was triturated with hexane to obtain the title compound (0.022 g, 61 %).

¹HNMR(400MHz, CDCl₃): δ 0.94-0.99(m, 6H); 1.65-1.71 (m, 2H); 1.77-1.83 (m, 2H); 3.97 (t, J=7.6Hz, 2H); 4.05 (t, J=7.6Hz, 2H); 4.11 (s, 3H); 5.23 (s, 2H); 7.01-7.05 (m, 2H); 7.44-7.47 (m, 2H), 7.96 (s, 1H); 8.18 (s, 1H).

Examples F2 – F8 were prepared in an analogous manner of Example F1 from the appropriate intermediate.

Example	IUPAC name
F2	7-Methyl-1,3-dipropyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.
F3	7-Methyl-1,3-dipropyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione.
F4	7-Methyl-8-{1-[4-(4-methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
F5	8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
F6	8-{1-[3-(3-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
F7	8-{1-[3-(3-tert-Butyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
F8	(3-{3-[4-(7-Methyl-2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenoxy)-acetic acid

Example G1: 8-{1-[3-(3-Hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



5 Step I: 8-{1-[3-(3-Hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

To a solution of 3-(3-{4-[2,6-dioxo-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-prop-1-ynyl)-benzoic acid ethyl ester (50 mg, 0.08 mmol) in THF (5ml), NaBH₄ (5 mg, 0.16 mmol) was added slowly at 0 °C. It was stirred for 3 h at room temperature. Water was added to reaction mixture and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ and then filtered and concentrated under vacuum. The residue obtained was purified by column chromatography to obtain 8-{1-[3-(3-hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (25 mg, 54 %) as an yellow oil.

¹HNMR(400MHz, CDCl₃): δ 0.86-1.01 (m, 8H); 1.66-1.72 (m, 2H); 1.80-1.86 (m, 2H); 3.79-3.83 (t, J=8.4Hz, 2H); 3.98-4.01 (t, J= 7.2Hz, 2H); 4.09-4.13 (t, J=7.2Hz); 4.69 (s,


2H); 5.24 (s, 2H); 5.79 (s, 2H); 7.26-7.42 (m, 3H); 7.50 (s, 1H); 8.16 (s, 1H); 8.39 (s, 1H)

Step II: 8-{1-[3-(3-Hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

5 To a solution of 8-{1-[3-(3-hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (25 mg) in ethanol (3 ml), aq. HCl (2 N, 1 ml) was added and the reaction mixture was refluxed for 2 hrs. The volatiles were evaporated and the residue obtained was dissolved in ethyl acetate. The organic layer was washed with water, brine and dried
10 over Na₂SO₄ and concentrated under vacuum. The residue obtained was purified by column to obtain 8-{1-[3-(3-hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (10 mg, 52 %) as white solid.

¹HNMR(400MHz, CDCl₃): δ 0.96-1.00(m, 6H); 1.69-1.75(m, 2H); 1.80-1.86(m, 2H); 4.05-4.14(m, 4H); 4.68(s, 2H); 5.24(s, 2H); 7.25-7.40(m, 3H); 7.51(s, 1H); 8.3(s, 1H);
15 8.42(s, 1H).

Example H1: 4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzenesulfonamide




20 A mixture of 1,3-dipropyl-8-(1-prop-2-ynyl-1H-pyrazol-4-yl)-3,7-dihydro-purine-2,6-dione (200mg, 0.59mmol), 4-iodo-benzenesulfonamide (166mg, 0.59mmol), copper iodide (10mg, 0.059mmol), dichlorobis (triphenylphosphine)-palladium (II) (21mg, 0.03mmol), triethylamine (2ml), DMF (2ml) was degassed for 10min. and stirred for 20hrs at room temperature. The reaction mixture was diluted with water (10ml) and
25 extracted with DCM (3X10ml). Organic layer was washed with brine (2X15ml) and

dried over Na_2SO_4 . The organic volatile solvent was evaporated and the crude product was purified using HPLC to obtain pure 4-{3-[4-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzenesulfonamide (4.2 mg, 2%).

- 5 ^1H NMR(400MHz, DMSO d_6): δ 0.87-0.90 (m, 6H); 3.86 (t, $J=6.8\text{Hz}$, 2H); 3.99 (t, $J=6.8\text{Hz}$, 2H); 5.45 (s, 2H) 7.5 (s, 2H); 7.70 (d, $J=8.4\text{Hz}$, 2H); 7.85 (d, $J=8.4\text{Hz}$, 2H); 8.12 (s, 1H); 8.41 (s, 1H); 8.51 (s, 1H).

Example I1: 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzamide



10

Step I: 3-(3-{4-[2,6-Dioxo-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-prop-1-ynyl)-benzamide

- A mixture of 3-(3-{4-[2,6-dioxo-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-prop-1-ynyl)-benzoic acid (100mg, 0.17mmol), ammonium carbonate (33mg, 0.34mmol), HOBt (36mg, 0.24mmol), N-
- 15

5 methyl morpholine (34mg, 0.34mmol) were taken in DMF (2ml), EDCI (130mg, 0.68mmol) was added at cold condition and stirred for 18hrs at room temperature. The reaction mixture was diluted with water (10ml) and extracted with DCM (3X5ml), the organic layer was washed with brine (2X10ml), dried over Na₂SO₄. The organic solvent was evaporated and the crude product was purified using HPLC to obtain 3-(3-{4-[2,6-Dioxo-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-prop-1-ynyl)-benzamide (33mg, 18%).


10 ¹HNMR(400MHz, DMSO d₆): δ 0.12 (s, 9H); 0.84-0.91 (m, 8H); 1.58 (q, J=6.8Hz, 2H); 1.73 (q, J=7.2Hz, 2H); 3.69 (t, J=8.4Hz, 2H); 3.87 (t, J=7.2Hz, 2H); 4.01 (t, J=7.2Hz, 2H); 5.47 (s, 2H); 5.80 (s, 2H); 7.49-7.53 (m, 2H); 7.65 (d, J=7.6Hz, 1H); 7.92 (d, J=7.6Hz, 1H); 7.99 (s, 1H); 8.10 (s, 1H); 8.14 (s, 1H); 8.56 (s, 1H).

Step II: 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzamide

15 A solution of above product (33mg, 0.056mol) was taken in ethanol (1ml) and 2N HCl (0.5ml), refluxed for 3hrs at 80 °C. The reaction mixture was concentrated in vacuo and the residue was washed with ether to furnish 3-{3-[4-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzamide (15mg, 60%).

20 ¹HNMR(400MHz, DMSO d₆): δ 0.85-0.92 (m, 6H); 1.58 (q, J=6.8Hz, 2H); 1.73 (q, J=7.2Hz, 2H); 3.87 (t, J=7.6Hz, 2H); 3.4 (t, J=7.2Hz, 2H); 5.44 (s, 2H); 7.49-7.53 (m, 2H); 7.65 (d, J=7.6Hz, 1H); 7.92 (d, J=7.6Hz, 1H); 7.99 (s, 1H); 8.10 (s, 1H); 8.14 (s, 1H); 8.56 (s, 1H); 13.6 (bs, 1H).

Example J1: 1,3-Dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione



Step-1: [2-(3-Trifluoromethyl-phenyl)-cyclopropyl]-methanol

2-(3-Trifluoromethyl-phenyl)-cyclopropanecarboxylic acid ethyl ester (prepared as per
 5 literature procedure) (0.70 g, 2.7 mmol) in THF (5ml) was added to a solution of
 lithium aluminium hydride (0.123 g, 3.25 mmol) in THF (5ml) at 0 °C over a period of
 10 min. and stirred further at 20-25 °C for 2 hours. The reaction mixture was diluted
 with THF (15 ml) and quenched with 10 % NaOH solution (5 ml) and water (10 ml).
 The aqueous layer was extracted with ethyl acetate, washed with saturated brine
 10 solution and dried over Na₂SO₄. The organic layer was evaporated to afford [2-(3-
 trifluoromethyl-phenyl)-cyclopropyl]-methanol as a colorless oil (0.54g, 92 %).

¹HNMR(400MHz, CDCl₃): δ 1.01 (t, J=7.2Hz, 2H); 1.45-1.53 (m, 2H); 1.87-1.92 (m, 1H); 3.64-3.66 (m, 2H); 7.24 (m, 1H); 7.31 (s, 1H); 7.34-7.4 (m, 2H).

Step-2: 1-(2-Bromomethyl-cyclopropyl)-3-trifluoromethyl-benzene

15 [2-(3-Trifluoromethyl-phenyl)-cyclopropyl]-methanol obtained in step 1 (0.250g, 1.15
 mmol) was taken in diethyl ether (4ml) and cooled to 0 °C. Tribromo phosphine
 (0.06ml, 0.64 mmol) in diethyl ether (4ml) was added slowly at 0 °C and stirred at 15-
 20 °C for 2 hrs. Reaction mixture was quenched with saturated NaHCO₃ solution (20
 ml) and extracted with diethyl ether. Organic layer was washed with saturated brine
 20 solution, dried over Na₂SO₄, and evaporated under vacuum to get 1-(2-bromomethyl-
 cyclopropyl)-3-trifluoromethyl-benzene (0.26g, 80 %) as a sticky mass.

¹HNMR(400MHz, CDCl₃): δ 1.08-1.13 (m, 1H); 1.19-1.28 (m, 1H); 1.58-1.67 (m, 1H); 1.97-2.02 (m, 1H); 3.46 (d, J=7.6Hz, 2H); 7.26-7.28 (m, 1H); 7.33 (s, 1H); 7.37-7.39 (m, 1H); 7.42-7.44 (m, 1H).

Step-3: 1,3-Dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

A mixture of 1, 3-dipropyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione (0.100g, 0.23mmol), 1-(2-bromomethyl-cyclopropyl)-3-trifluoromethyl-benzene (0.64g, 0.23 mmol), K₂CO₃ (0.063g, 0.46 mmol) and acetone (10ml) were heated at 50-55 ° C for 16hrs. Reaction mixture was cooled to 20-25 ° C and filtered off. The filtrate was evaporated and residue was purified by column chromatography to afford 1,3-dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.55g, 58 %).

¹HNMR(400MHz, CDCl₃): δ 0.12 (s, 9H); (s, 9H); 0.93-0.99 (m, 8H); 1.16 (t, J=6.8Hz, 1H); 1.67-1.69 (m, 4H); 1.81-1.83 (m, 2H); 2.06-2.12 (m, 1H); 3.81 (t, J=8.4Hz, 2H); 3.99 (t, J=7.2Hz, 2H); 4.08-4.12 (m, 2H); 4.25 (d, J=6.8Hz, 2H); 5.76 (s, 2H); 7.22-7.25 (m, 1H); 7.30 (s, 1H); 7.33-7.37 (m, 1H); 7.40-7.42 (m, 1H); 8.13 (s, 1H); 8.17 (s, 1H).

Step-4: 1,3-Dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione


A mixture of 1,3-dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.55 g, 0.087 mmol), 2 N HCl (2 ml), ethanol (3 ml) were heated at 85 ° C for 3 hours. The mixture was cooled to 10-15 ° C and solid material was separated. It was filtered off and washed with water (1ml), diethyl ether (1ml) to obtain the title compound (0.016 g, 37 %).

¹HNMR(400MHz, CDCl₃): δ 0.97-1.01 (m, 6H); 1.18 (t, J=7.2Hz, 2H); 1.67-1.86 (m, 5H); 2.09-2.10 (m, 1H); 4.08-4.15 (m, 4H); 4.23-4.27 (m, 2H); 7.22-7.42 (m, 4H); 8.22 (s, 1H); 8.36 (s, 1H); 12.87 (bs, 1H).

Examples J2 - J4 were prepared in an analogous manner of Example J1 from the appropriate intermediate.

Example	IUPAC name
J2	8-{1-[2-(3-Fluoro-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}- 1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
J3	8-{1-[2-(4-Fluoro-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}- 1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
J4	1,3-Dipropyl-8-{1-[2-(4-trifluoromethyl-phenyl)- cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.

Example K1: 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



5 Step-1: 1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidine-3-carboxylic acid

A mixture of 4-isopropylaniline (0.76 g, 0.565 mmol) and itaconic acid (0.70g, 0.538 mmol) was heated at 120-130 °C in a sealed tube for 2.5 hrs. The mixture was cooled

and dissolved in 10 % NaOH solution (15 ml) and stirred for 20 min. The aqueous layer was washed with ethyl acetate and acidified with dil. HCl. The precipitate was filtered to obtain 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidine-3-carboxylic acid (1.0 g, 75 %).

¹HNMR(400MHz, CDCl₃): δ 1.25 (d, J=7.2Hz, 6H); 2.90-2.98 (m, 3H); 3.41-3.45 (m, 1H); 4.06-4.08 (m, 1H); 4.14-4.18 (m, 1H); 7.26 (d, J=8.4Hz, 2H); 7.49 (d, J=8.4Hz, 2H)

Step-2: 1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidine-3-carboxylic acid ethyl ester

A mixture of 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidine-3-carboxylic acid obtained in step 1 (0.6 g, 2.42 mmol), thionyl chloride (0.35g, 2.90 mmol), and ethanol (10 ml) was heated at 55-60 °C for 2 hours. The mixture was cooled and solvent was removed. The residue was dissolved in ethyl acetate (20 ml). The organic layer was washed with saturated NaHCO₃ and saturated brine solution (20 ml). The organic layer was dried over Na₂SO₄ and evaporated to obtain 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidine-3-carboxylic acid ethyl ester as yellow oil (0.7g, 100 %).

Step-3: 4-Hydroxymethyl-1-(4-isopropyl-phenyl)-pyrrolidin-2-one

A mixture of 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidine-3-carboxylic acid ethyl ester obtained in step 2 (0.7 g, 2.65 mmol) and ethanol were cooled to 10-15 °C. Sodium borohydride (0.25g, 6.6 mmol) was added portion wise over a period of 20 min and the reaction mixture was stirred for 3.5 hrs at 20-25 °C. The organic volatiles were evaporated and the residue was taken into brine solution (15 ml). The aqueous layer was extracted with ethyl acetate, dried over Na₂SO₄ and evaporated to obtain 4-hydroxymethyl-1-(4-isopropyl-phenyl)-pyrrolidin-2-one as an off white solid (0.5g, 81%).

¹HNMR(400MHz, CDCl₃): δ 1.21 (d, J=6.8 Hz, 6H); 2.40-2.44 (m, 1H); 2.68-2.73 (m, 2H); 2.85-2.89 (m, 1H); 3.67-3.74 (m, 3H); 3.91-3.95 (m, 1H); 7.21 (d, J=8.4Hz, 2H); 7.49 (d, J=8.4Hz, 2H).

Step-4: Methanesulfonic acid 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl ester

A mixture of 4-hydroxymethyl-1-(4-isopropyl-phenyl)-pyrrolidin-2-one obtained in step 3 (0.5 g, 2.14 mmol), dichloromethane (8 ml) and triethyl amine (0.30g, 3.0 mol) was cooled to 0 °C and stirred for 15 min at that temperature. Methane sulfonyl chloride (0.3g, 2.56 mmol) was added to the mixture over a period of 5 min and the reaction mixture was stirred for 1.5 hrs at 10-15 °C. To this reaction mixture saturated

NaHCO₃ solution (10 ml) was added and the organic layer was separated. The aqueous layer was extracted with dichloromethane (20 ml). The combined organic layer was washed with saturated brine solution, dried over Na₂SO₄ and evaporated to obtain methanesulfonic acid 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl ester as a brown oil (0.6g, 90 %).

¹HNMR(400MHz, CDCl₃): δ 1.21 (d, J=6.8Hz, 6H); 2.4-2.46 (m, 1H); 2.46-2.91 (m, 3H); 3.04 (s, 3H); 3.7-3.74 (m, 1H); 3.97-4.02 (m, 1H); 4.24-4.3 (m, 2H); 7.22 (d, J=8.4Hz, 2H); 7.46 (d, J=8.4Hz, 2H).

Step-5: 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

A mixture of 1, 3-dipropyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2, 6-dione (0.100g, 0.23mmol), methanesulfonic acid 1-(4-isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl ester (0.107g, 0.34mmol), K₂CO₃ (0.048g, 0.34mmol) and DMF (1ml) were heated at 75-80 °C for 16 hrs. Reaction mixture was cooled to 20-25 °C and water was added (10 ml). The aqueous layer was extracted with ethyl acetate, washed with saturated brine solution. The organic layer was dried over Na₂SO₄ and evaporated to obtain crude product (0.2g). The crude product was purified by preparative TLC to obtain pure 8-{1-[1-(4-isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.12g, 80 %) as a light yellow oil.

¹HNMR(400MHz, CDCl₃): δ 0.12 (s, 9H); (s, 9H); 0.83-0.87 (m, 8H); 1.16 (d, J=6.8Hz, 6H); 1.53-1.58 (m, 2H); 1.67-1.72 (m, 2H); 2.34-2.38 (m, 1H); 2.66-2.89 (m, 2H); 3.17-3.21 (m, 1H); 3.72-3.78 (m, 2H); 3.8-3.88 (m, 2H); 3.91-4.12 (m, 2H); 4.17-4.21 (m, 2H); 4.26-4.38 (m, 2H); 5.79 (m, 2H); 7.21 (d, J=8.4Hz, 2H); 7.46 (d, J=8.4Hz, 2H); 8.18 (s, 1H); 8.40 (s, 1H).

Step-6: 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

A mixture of 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione obtained in step 5 (0.1 g, 0.23 mmol), 2 N HCl (3 ml), ethanol (2 ml) were heated at 85 °C for 3 hours. The mixture was cooled to 10-15 °C. The solid obtained was filtered and washed with water (1ml), diethyl ether (1ml) to obtain 8-{1-[1-(4-isopropyl-

phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (0.071 g, 76 %) as a white solid.

¹HNMR(400MHz, CDCl₃): δ 0.83-0.87 (m, 6H); 1.15 (d, J=6.8Hz, 6H); 1.53-1.58 (m, 2H); 1.69-1.72 (m, 2H); 2.34-2.38 (m, 1H); 2.59-2.65 (m, 1H); 2.82-2.85 (m, 1H);
 5 2.89-3.01 (m, 1H); 3.61-3.65 (m, 1H); 3.83-3.91 (m, 3H); 3.96 (t, J=6.8Hz, 2H); 4.31 (d, J=6.8Hz, 2H); 7.19 (d, J=8.4Hz, 2H); 7.47 (d, J=8.4Hz, 2H); 8.08 (s, 1H); 8.43 (s, 1H); 13.46 (s, 1H).


Examples K2 - K33 were prepared in an analogous manner of Example K1 from the appropriate intermediate.

Example	IUPAC name
K2	8-{1-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K3	8-{1-[5-Oxo-1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K4	8-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K5	8-[1-(5-Oxo-1-p-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K6	8-[1-(5-Oxo-1-m-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K7	8-{1-[5-Oxo-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K8	8-{1-[1-(4-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K9	8-{1-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
K10	8-{1-[1-(3-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K11	3-Ethyl-8-{1-[1-(4-methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
K12	3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.

K13	8-{1-[1-(4-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K14	4-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzonitrile.
K15	3-Ethyl-8-{1-[1-(4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
K16	3-Ethyl-8-{1-[1-(3-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
K17	3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
K18	8-{1-[5-Oxo-1-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K19	3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
K20	3-Ethyl-8-{1-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione.
K21	8-{1-[1-(3-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K22	8-{1-[5-Oxo-1-(2-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
K23	8-{1-[5-Oxo-1-(4-trifluoromethoxy-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K24	8-{1-[1-(4-Methyl-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
K25	4-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzoic acid.
K26	8-{1-[1-(4-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
K27	3-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzonitrile
K28	8-{1-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

K29	8-{1-[1-(2,4-Difluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
K30	8-{1-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione
K31	8-{1-[1-(2-Chloro-4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
K32	8-{1-[1-(2-Chloro-4-trifluoromethyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
K33	3-Ethyl-8-{1-[5-oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione

Example L1: 8-{1-[2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



5 Step-1: 2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid

A mixture of 3-trifluoromethyl-phenylamine (2.8g, 17.5 mmol) and 6,6-dimethyl-5,7-dioxaspiro[2.5]octane-4,8-dione (1.0g, 5.8 mmol) was stirred at 20-25 °C for 18 hrs. The reaction mixture was diluted with dichloromethane and washed with 10 % NaOH solution (3 X 10 ml). The aqueous layer was cooled and acidified with dil HCl. The aqueous layer was extracted with dichloromethane, washed with saturated brine

solution. The organic layer was dried over Na_2SO_4 and evaporated to obtain 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid (1.1 g, 69 %) as an off white solid.

^1H NMR(400MHz, CDCl_3): δ 2.57-2.64 (m, 2H); 3.75 (t, $J=9.6\text{Hz}$, 1H); 3.99-4.03 (m, 2H); 7.51-7.61 (m, 2H); 7.88-7.9 (m, 2H).

Step-2: 2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid ethyl ester

A mixture of 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid obtained in step 1 (0.7 g, 2.56 mmol), thionyl chloride (0.36g, 3.0 mmol), and ethanol (10 ml) were heated at 55-60 °C for 2 hours. The mixture was cooled and solvent was removed. The residue was dissolved in ethyl acetate (20 ml). The organic layer was washed with sat. NaHCO_3 and saturated brine solution, dried over Na_2SO_4 and evaporated to obtain 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid ethyl ester as a yellow oil (0.5g, 65 %).

^1H NMR(400MHz, CDCl_3): δ 1.31 (t, $J=7.2\text{Hz}$, 3H); 2.46-2.49 (m, 1H); 2.60-2.65 (m, 1H); 3.71 (t, $J=7.6\text{Hz}$, 1H); 3.89-3.95 (m, 1H); 4.04-4.09 (m, 1H); 4.31 (q, $J=7.2\text{Hz}$, 2H); 7.45-7.47 (m, 1H); 7.51-7.55 (m, 1H); 7.92 (m, 1H).

Step-3: 3-Hydroxymethyl-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one

A mixture of 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid ethyl ester obtained in step 2 (0.5 g, 1.6 mmol) and ethanol were cooled to 10-15 °C. Sodium borohydride (0.126g, 3.3 mmol) was added portion wise over a period of 20 min and the reaction mixture was stirred for 3.5 hrs at 20-25 °C. The organic volatiles were evaporated and the residue was taken into brine solution (15 ml). The aqueous layer was extracted with ethyl acetate, dried over Na_2SO_4 and evaporated to obtain brown semi solid which was purified by preparative TLC to afford 3-Hydroxymethyl-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one as a colorless oil (0.177g, 43 %).

Step-4: Methanesulfonic acid 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl ester

A mixture of 3-hydroxymethyl-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one obtained in step 3 (0.17 g, 0.6 mmol), dichloromethane (8 ml) and triethyl amine (0.085g, 0.84 mmol) was cooled to 0 °C and stirred for 15 min at that temperature. Methane sulfonyl chloride (0.090g, 0.78 mmol) was added to the mixture over a period of 5 min and the

reaction mixture was stirred for 1.5 hrs at 10-15 °C. To this reaction mixture saturated NaHCO₃ solution (10 ml) was added and the organic layer was separated. The aqueous layer was extracted with dichloromethane. The combined organic layer was washed with saturated brine solution, dried over Na₂SO₄ and evaporated to obtain
 5 methanesulfonic acid 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl ester as a brown oil (0.23g, 100 %).

¹HNMR(400MHz, CDCl₃): δ 2.29-2.34 (m, 1H); 2.45-2.51 (m, 1H); 3.05-3.13 (m, 3H); 3.91-3.95 (m, 3H); 4.54-4.57 (m, 1H); 4.63-4.67 (m, 1H); 7.46-7.48 (m, 1H); 7.52-7.56 (m, 1H); 7.89-7.93 (m, 2H).

10 **Step-5: 8-{1-[2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione**

A mixture of 1, 3-dipropyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione (0.100g, 0.23mmol), methanesulfonic acid 2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl ester (0.10g, 0.32mmol), K₂CO₃ (0.045g, 0.32mmol) and DMF (1ml) was heated at 75-80 °C for 16hrs. Reaction mixture was
 15 cooled to 20-25 °C and diluted with water (10 ml). The aqueous layer was extracted with ethyl acetate, washed with saturated brine solution. The organic layer was dried over Na₂SO₄ and evaporated to obtain crude product (0.2g). The crude product was
 20 purified by preparative TLC to obtain pure 8-{1-[2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (0.10g, 65 %).

¹HNMR(400MHz, CDCl₃): δ 0.10 (s, 9H); 0.96-1.01 (m, 8H); 1.69-1.77 (m, 2H); 1.85-1.90 (m, 2H); 2.08-2.14 (m, 1H); 2.47-2.49 (m, 1H); 3.32-3.42 (m, 1H); 3.75-3.79 (m, 2H); 3.87 (t, J=8.4Hz, 2H); 4.03 (t, J=7.6Hz, 2H); 4.34 (t, J=7.6Hz, 2H); 4.60-4.63 (m, 1H); 4.74-4.78 (m, 1H); 5.83-5.90 (m, 2H); 7.44-7.53 (m, 1H); 7.51-7.55 (m, 1H); 7.85-7.92 (m, 2H); 8.35 (s, 1H); 8.84 (s, 1H).

Step-6: 8-{1-[2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

30 A mixture of 8-{1-[2-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione obtained in step 5 (0.1 g, 0.148 mmol), 2 N HCl (3 ml), ethanol (2 ml) was heated at 85 °C for 3 hours. The mixture was cooled to 10-15 °C. The solid obtained

was filtered and washed with water, diethyl ether to obtain the title compound as a white solid (0.067 g, 83 %).


¹HNMR(400MHz, DMSO d6): δ 0.87-0.92 (m, 6H); 1.56-1.62 (m, 2H); 1.70-1.76 (m, 2H); 1.86-1.91 (m, 1H); 2.17-2.21 (m, 1H); 3.28-3.31 (m, 1H); 3.72-3.76 (m, 1H);
 5 3.81-3.89 (m, 2H); 3.99 (t, J=7.2Hz, 2H); 4.47-4.6 (m, 2H); 7.53 (d, J=8Hz, 1H); 7.65 (dd, J=8Hz, 1H); 7.84 (d, J=8Hz, 1H); 8.11 (s, 1H); 8.21 (s, 1H); 8.42 (s, 1H); 13.59 (bs, 1H).

Examples L2 and L3 were prepared in an analogous manner of Example L1 from the appropriate intermediate.

Example	IUPAC name
L2	8-{1-[1-(3-Fluoro-phenyl)-2-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
L3	8-{1-[2-Oxo-1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

10

Example M1: 1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione



Step-1: [1-(3-Trifluoromethyl-phenyl)-pyrrolidin-3-yl]-methanol

5 5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid (0.35 g, 1.28 mmol) in diethyl ether (5 ml) was added to a solution of lithium aluminium hydride (0.097 g, 2.56 mmol) in THF (5 ml) at temperature 0 °C over a period of 10 min. The mixture was refluxed for 2 hours and then stirred at room temperature for 20 hours. The reaction mixture was diluted with diethyl ether (15 ml) and quenched with water (10 ml). The organic layer was separated and aqueous layer was extracted with diethyl ether. The combined organic layer was washed with saturated brine solution, dried over Na₂SO₄ and evaporated to obtain a residue which was purified by preparative TLC to afford [1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-methanol as a colorless mass (0.2g, 64 %).

¹HNMR(400MHz, CDCl₃): δ 1.85-1.90 (m, 1H); 2.16-2.21 (m, 1H); 2.60-2.63 (m, 1H); 3.17-3.21 (m, 1H); 3.36-3.50 (m, 4H); 3.65-3.75 (m, 2H); 6.77-6.80 (m, 2H); 6.94 (d, J=7.6Hz, 1H); 7.28-7.32 (m, 1H).

15 Step-2: Methanesulfonic acid 1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl ester

Prepared following the procedure similar to step 4 of Example K1.

¹HNMR(400MHz, CDCl₃): δ 1.86-1.95 (m, 1H); 2.19-2.27 (m, 1H); 2.78-2.85 (m, 1H); 3.00 (s, 3H); 3.18-3.23 (m, 1H); 3.33-3.50 (m, 3H); 4.19-4.23 (m, 1H); 4.27-4.31 (m, 1H); 6.69-6.74 (m, 2H); 6.93 (d, J=7.6Hz, 1H); 7.28-7.34 (m, 1H).

20 Step-3: 1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

Prepared following the procedure similar to step 5 of Example K1.

¹HNMR(400MHz, CDCl₃): δ 0.10 (s, 9H); 0.98-1.06 (m, 8H); 1.82-1.92 (m, 3H); 1.94-2.01 (m, 2H); 2.26-2.30 (m, 1H); 3.09-3.14 (m, 1H); 3.26-3.30 (m, 1H); 3.41-3.48 (m, 1H); 3.50-3.56 (m, 2H); 3.86 (t, J=8.0Hz, 2H); 4.03 (t, J=7.2Hz, 2H); 4.17 (t, J=7.6Hz, 2H); 4.34 (d, J=7.6Hz, 2H); 5.82-5.83 (m, 2H); 6.80-6.84 (m, 2H); 7.01-7.03 (m, 1H); 7.33-7.38 (m, 1H); 8.20 (s, 1H); 8.26 (s, 1H).

30 Step-4: 1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione


Prepared following the procedure similar to step 6 of Example K1.

¹HNMR(400MHz, DMSO d6): δ 0.83-0.88 (m, 6H); 1.52-1.58 (m, 2H); 1.67-1.78 (m, 3H); 2.01-2.04 (m, 1H); 2.85-2.88 (m, 1H); 3.08-3.12 (m, 1H); 3.23-3.25 (m, 1H); 3.34-3.38 (m, 2H); 3.83 (t, J=7.6Hz, 2H); 3.95 (t, J=7.6Hz, 2H); 4.27 (d, J=6.0Hz, 2H); 6.66 (s, 1H); 6.75 (d, J=8.4Hz, 1H); 6.85 (d, J=7.2Hz, 1H); 7.31-7.35 (m, 1H); 8.09 (s, 1H); 8.41 (s, 1H); 13.46 (bs, 1H).

Examples M2 - M6 were prepared in an analogous manner of Example M1 from the appropriate intermediate.

Example	IUPAC name
M2	1,3-Dipropyl-8-{1-[1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione
M3	1,3-Dipropyl-8-[1-(1-p-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione
M4	8-{1-[1-(4-Methoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
M5	8-{1-[1-(3-Methoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
M6	1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione.

Example N1: 8-{4-[1-(4-Fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]phenyl}-1,3-dipropyl-3,7-dihydropurine-2,6-dione



5 Step-1: 4-[1-(4-Fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzoic acid methyl ester

4-Hydroxybenzoic acid methyl ester (0.20 g, 1.314 mmol), 1-(4-fluoro-phenyl)-4-hydroxymethyl-pyrrolidin-2-one (0.33 g, 1.577 mmol), Ph_3P (0.345 g, 1.314 mmol) and triethyl amine (0.20 ml, 1.577 mmol) were mixed under argon atmosphere followed by the addition of the dry toluene (10 mL). The reaction mixture was cooled to 0 °C followed by addition of DIAD (0.3 mL, 0.788 mmol) drop wise. The reaction mixture was allowed to warm to room temperature and heated at 80 °C for 24h. The reaction was quenched with 5% HCl (10 mL), extracted with DCM. The organic layer was dried over Na_2SO_4 and concentrated to afford the crude product which was purified by flash

column chromatography to obtain 4-[1-(4-fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzoic acid methyl ester (0.33 g, 73 %) as a white solid.

¹HNMR(400MHz, DMSO d₆): δ 2.43 (dd, J=6.4, 6.4 Hz, 1H); 2.73 (dd, J=8.8, 9.2 Hz, 1H); 2.99-2.89 (m, 1H); 3.70 (dd, J=5.2, 5.6Hz, 1H); 3.79 (s, 3H); 4.02 (t, J=9 Hz, 1H); 4.14-4.00 (m, 2H); 7.05 (d, J=8.8Hz, 2H); 7.20 (t, J=8.8Hz, 2H); 7.69-7.66 (m, 2H); 7.90 (d, J=8.8 Hz, 2H).

Step-2: 4-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzoic acid

To a stirred solution of 4-[1-(4-fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzoic acid methyl ester obtained in step 1 (0.14g, 0.408 mmol) in THF:MeOH:H₂O (6 mL, 2:1:3), LiOH.H₂O was added at 20-25 °C. The reaction mixture was stirred for 16 hours at that temperature. The solvents were removed under reduced pressure and the residue was diluted with water (10 ml). The aqueous layer was washed with DCM, acidified with 2% HCl. The acidified mixture was extracted with ethyl acetate, washed with brine, dried over Na₂SO₄. The organic layer was evaporated under reduced pressure to obtain 4-[1-(4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzoic acid (0.130 g, quantitative) as a white solid.

¹HNMR(400MHz, DMSO d₆): δ 2.42 (dd, J=4.0, 8.0 Hz, 1H); 2.73 (dd, J=8.0, 8.0 Hz, 1H); 2.96-2.89 (m, 1H); 3.70 (dd, J=8.0, 4.0 Hz, 1H); 4.02 (t, J=8.0 Hz, 1H); 4.14-4.11 (m, 2H); 7.03 (d, J=8.0 Hz, 2H); 7.20 (t, J=8.0 Hz, 2H); 7.69-7.66 (m, 2H); 7.87 (d, J=8.0 Hz, 2H); 12.6 (br s, 1H).

Step-3: N-(6-Amino-2,4-dioxo-1,3-dipropyl-1,2,3,4-tetrahydro-pyrimidin-5-yl)-4-[1-(4-fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzamide

To a stirred solution of 4-[1-(4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzoic acid obtained in step 2 (0.140 g, 0.426 mmol) and HCl salt of 5,6-diamino-1,3-dipropyl-1H-pyrimidine-2,4-dione (0.112 g, 0.426 mmol) in MeOH (5 mL) was added EDCI (0.0136 g, 0.593 mmol). After stirring at 20-25 °C for 16 hours, the reaction mixture was quenched with water (5 mL). The solid obtained was filtered and washed with water to provide the crude product which was purified by column chromatography to afford N-(6-amino-2,4-dioxo-1,3-dipropyl-1,2,3,4-tetrahydro-pyrimidin-5-yl)-4-[1-(4-fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzamide (0.205 g, 90%) as a white solid.

¹HNMR(400MHz, DMSO d₆): δ 0.89-0.79 (m, 6H); 1.58-1.45 (m, 4H); 2.44 (dd, J=8, 8Hz 1H); 2.74 (dd, J=9.2, 8 Hz, 1H); 2.95-2.86 (m, 1H); 3.73-3.68 (m, 3H); 3.82 (t, J=16Hz, 2H); 4.03 (t, J=8 Hz, 1H); 4.11-4.13 (m, 2H); 6.68 (s, 2H); 7.02 (d, J=8.0Hz, 2H); 7.21 (t, J=8.0Hz, 2H); 7.70-7.67 (m, 2H); 7.94 (d, J=8.0Hz, 2H); 8.76 (s, 1H).

5 **Step-4: 8-{4-[1-(4-Fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]phenyl}-1,3-dipropyl-3,7-dihydropurine-2,6-dione**

10 *N*-(6-Amino-2,4-dioxo-1,3-dipropyl-1,2,3,4-tetrahydro-pyrimidin-5-yl)-4-[1-(4-fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-benzamide obtained in step 3 (0.90 g, 0.167 mmol) was dissolved in MeOH (4 mL) and 10% NaOH (0.020 g, 0.502 mmol) and stirred at 80 °C for 7 h. MeOH was distilled off and the residue was taken up in H₂O (2 mL), acidified with 2N HCl to pH 4-5. The precipitate was filtered, washed with water and methanol to provide the title compound as white solid (0.072 g, 83 %).

15 ¹HNMR(400MHz, DMSO d₆): δ 0.91-0.84 (m, 6H); 1.61-1.52 (m, 2H); 1.77-1.68 (m, 2H); 2.44 (dd, J=4 Hz, 1H); 2.74 (dd, J=8.0, 8.0 Hz, 1H); 2.97-2.89 (m, 1H); 3.71 (dd, J=8.0, 4.0Hz, 1H); 3.85 (t, J=8.0 Hz, 2H); 4.05-3.98 (m, 3H); 4.14-4.11 (m, 2H); 7.08 (d, J=8.8 Hz, 2H); 7.20 (t, J=8.8 Hz, 2H); 7.68 (m, 2H); 8.06 (d, J=8.8 Hz, 2H); 13.6 (br s, 1H).

Examples N2 - N43 were prepared in an analogous manner of Example N1 from the appropriate intermediate.


Example	IUPAC name
N2	1,3-Dipropyl-8-{4-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione.
N3	8-{3-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N4	8-{4-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N5	8-{4-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N6	8-{6-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N7	8-{3-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N8	8-{3-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-

	isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N9	8-{6-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N10	8-{6-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N11	8-{4-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N12	8-{4-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N13	8-{6-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N14	8-{3-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N15	8-{4-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N16	8-{3-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N17	1,3-Dipropyl-8-{4-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione.
N18	1,3-Dipropyl-8-{3-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione.
N19	8-{6-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N20	1,3-Dipropyl-8-{6-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione.
N21	8-{6-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N22	1,3-Dipropyl-8-{6-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione.
N23	1,3-Dipropyl-8-{6-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione.
N24	1,3-Dipropyl-8-{4-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-

	phenyl}-3,7-dihydro-purine-2,6-dione.
N25	1,3-Dipropyl-8-{3-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione.
N26	1,3-Dipropyl-8-{6-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione.
N27	8-{6-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N28	8-{6-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N29	8-{3-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N30	8-{4-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N31	1,3-Dipropyl-8-{4-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione.
N32	1,3-Dipropyl-8-{3-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione.
N33	8-{3-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N34	8-{3-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N35	1,3-Dipropyl-8-{3-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione.
N36	8-{4-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.
N37	8-{3-[1-(2,4-Difluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
N38	8-{5-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
N39	8-{1-Methyl-5-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethoxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
N40	8-{5-[1-(4-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-

	1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
N41	8-{5-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
N42	8-{1-Methyl-5-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
N43	8-{5-[1-(3-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

Example O1: 8-(1-{2-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



Step-1: [5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-acetonitrile

- 5 A mixture of methanesulfonic acid 5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl ester (prepared as given in Example B1)(0.9 g, 2.66 mmol), DMF (5 ml) and sodium cyanide (0.19g, 4.0 mol) were heated to 50-55 °C and stirred for 20hrs. To this reaction mixture water (50 ml) was added and extracted with ethyl acetate (3x20 ml), organic layer was washed with saturated brine solution (20 ml). The organic layer was

dried over Na₂SO₄ and evaporated to obtain [5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-acetonitrile (0.65g, 91 %) as brownish sticky mass.

¹HNMR(400MHz, CDCl₃): δ 1.51 (m, 1H); 2.62-2.65 (m, 2H); 2.89-2.93 (m, 2H); 3.70-3.74 (m, 1H); 4.09-4.13 (m, 1H); 7.42-7.44 (m, 1H); 7.48-7.52 (m, 1H); 7.81-7.86 (m, 1H); 8.02 (s, 1H).

Step-2: [5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-acetic acid

To a stirred solution of [5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-acetonitrile (0.25 g, 0.93 mmol) in ethanol (6 mL) was added sodium hydroxide solution (0.074g, 1.86 mmol, in 3ml water). The reaction mixture was refluxed for 3hrs and the solvents were removed under reduced pressure, diluted with water (3ml), washed with hexane (3x10 mL) and acidified with 2N HCl. The aqueous layer was extracted with DCM (3x15 mL) and the combined extract was washed with brine, dried with Na₂SO₄, evaporated under reduced pressure to obtain [5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-acetic acid (0.2g, 77 %) as an off white semi solid.

¹HNMR(400MHz, CDCl₃): δ 2.37-3.35 (m, 5H); 3.56-3.65 (m, 1H); 4.14-4.22 (m, 1H); 7.43-7.45 (m, 1H); 7.50-7.54 (m, 1H); 7.85-7.87 (m, 1H); 7.91-7.93 (m, 1H).

Step-3 to Step-7 were carried out as described in Example K1

Step-3: [5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-acetic acid methylester

¹HNMR(400MHz, CDCl₃): δ 2.23-2.37 (m, 1H); 2.59-2.61 (m, 2H); 2.85-2.95 (m, 2H); 3.61-3.66 (m, 1H); 3.74 (s, 3H); 4.09-4.15 (m, 1H); 7.43-7.45 (m, 1H); 7.50-7.54 (m, 1H); 7.85-7.87 (m, 1H); 7.91-7.93 (m, 1H).

Step-4: 4-(2-Hydroxy-ethyl)-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one

¹HNMR(400MHz, CDCl₃): δ 1.81-1.89 (m, 2H); 2.38-2.45 (m, 1H); 2.69-2.73 (m, 1H); 2.80-2.86 (m, 1H); 3.62-3.72 (m, 1H); 3.78-3.85 (m, 2H); 4.02-4.06 (m, 1H); 7.41-7.43 (m, 1H); 7.49-7.53 (m, 1H); 7.84 (s, 1H); 7.91-7.94 (m, 1H).

Step-5: Methanesulfonic acid 2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl ester

¹HNMR(400MHz, CDCl₃): δ 2.02-2.09 (m, 2H); 2.39-2.45 (m, 1H); 2.72-2.76 (m, 1H); 2.84-2.91 (m, 1H); 3.08 (s, 3H); 3.61-3.64 (m, 1H); 4.04-4.09 (m, 1H); 4.35-4.39 (m, 2H); 7.42-7.44 (m, 1H); 7.50-7.54 (m, 1H); 7.85-7.89 (m, 2H).


Step-6: 8-(1-{2-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

¹HNMR(400MHz, CDCl₃): δ 0.10 (s, 9H); 0.97-1.04 (m, 8H); 1.68-1.74 (m, 2H); 1.83-1.88 (m, 2H); 2.21-2.26 (m, 2H); 2.36-2.42 (m, 1H); 2.49-2.57 (m, 1H); 2.77-2.83 (m, 1H); 3.58-3.62 (m, 1H); 3.85 (t, J=8.0Hz, 2H); 3.99-4.04 (m, 3H); 4.13-4.18 (m, 2H); 4.35 (t, J=6.8Hz, 2H); 5.82 (s, 2H); 7.42 (d, J=7.6Hz, 1H); 7.49-7.53 (m, 1H); 7.79 (s, 1H); 7.90 (d, J=8.4Hz, 1H); 8.19 (s, 1H); 8.29 (s, 1H).

Step-7: 8-(1-{2-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

¹HNMR(400MHz, DMSO d₆): δ 0.84-0.91 (m, 6H); 1.55-1.60 (m, 2H); 1.69-1.75 (m, 2H); 2.03-2.05 (m, 2H); 2.32-2.38 (m, 2H); 2.63-2.68 (m, 1H); 3.62-3.66 (m, 1H); 3.86 (t, J=6.8Hz, 2H); 3.96-4.01 (m, 3H); 4.30 (t, J=6.4Hz, 2H); 7.48 (d, J=8.0Hz, 1H); 7.60-7.64 (m, 1H); 7.85 (d, J=7.6Hz, 1H); 8.11 (s, 2H); 8.45 (s, 1H); 13.55 (bs, 1H).

Example P1: 8-(1-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



5

Step-1: 5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid methoxy-methyl-amide

To a stirred solution of [5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid (1 g, 3.66 mmol) in DCM (15mL) was added 1,1'-carbonyldiimidazole (0.65g, 4.02 mmol) portion wise over a period of 10 min at 10-15 °C. N,O-

dimethylhydroxylamine hydrochloride (0.42g, 4.39 mmol) was added portion wise and stirred for 5 min at that temperature. To the reaction mixture triethylamine (0.66ml, 4.75 mmol) was added drop wise and it was stirred for 2 hr at 20-25 °C. The reaction mixture was diluted with water (10 ml) and aqueous layer was separated. The aqueous layer was extracted with DCM (2 X 25 ml). The combined organic layer was washed with 5 % HCl (25 ml) and saturated brine solution (25 ml). The organic layer was dried over Na₂SO₄, evaporated under reduced pressure to obtain 5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid methoxy-methyl-amide (0.95g, 83 %) as a colorless sticky mass.

¹HNMR(400MHz, CDCl₃): δ 2.84-2.91 (m, 1H); 3.0-3.06 (m, 1H); 3.29 (s, 3H); 3.72-3.76 (m, 1H); 3.79 (s, 3H); 4.02-4.07 (m, 1H); 4.15-4.19 (m, 1H); 7.42-7.46 (m, 1H); 7.50-7.53 (m, 1H); 7.86 (s, 1H); 7.91-7.93 (m, 1H).

Step-2: 4-Acetyl-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one

To a stirred solution of 5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid methoxy-methyl-amide (0.6g, 1.89 mmol) in THF (6mL) was added methylmagnesium bromide (1.6 ml, 4.74 mmol, 3.0M in ether) drop wise over a period of 10 min at 0 °C. The reaction mixture was stirred for 3 h at room temperature. The reaction mixture was quenched with saturated ammonium chloride solution (2ml) and extracted with ethyl acetate (2X15 ml). The combined organic layer was washed with saturated brine solution (25 ml). The organic layer was dried over Na₂SO₄, evaporated to obtain 4-Acetyl-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one as a brown oil (0.5g, 97%).

¹HNMR(400MHz, CDCl₃): δ 2.33 (s, 3H); 2.82-2.88 (m, 1H); 2.92-2.99 (m, 1H); 3.48-3.52 (m, 1H); 3.97-4.01 (m, 1H); 4.12-4.22 (m, 1H); 7.43-7.45 (m, 1H); 7.50-7.54 (m, 1H); 7.86-7.91 (m, 2H).

Step-3 to Step-6 were carried out similar to Example K1

Step-3: 4-(1-Hydroxy-ethyl)-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one

¹HNMR(400MHz, CDCl₃): δ 1.30-1.34 (m, 3H); 2.47-2.51 (m, 1H); 2.69-2.73 (m, 2H); 3.76-3.87 (m, 1H); 3.91-3.99 (m, 3H); 7.41-7.43 (m, 1H); 7.49-7.53 (m, 1H); 7.85-7.88 (m, 1H); 7.94-7.96 (m, 1H).

5 **Step-4: Methanesulfonic acid 1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl ester**

¹HNMR(400MHz, CDCl₃): δ 1.51-1.56 (m, 3H); 2.53-2.55 (m, 1H); 2.64-2.69 (m, 1H); 2.77-2.83 (m, 1H); 3.7 (s, 3H); 3.83-3.90 (m, 1H); 3.97-4.01 (m, 1H); 4.92-5.02 (m, 1H); 7.43-7.45 (m, 1H); 7.50-7.54 (m, 1H); 7.84-7.89 (m, 2H).

10 **Step-5: 8-(1-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione**

Crude material was taken as such to the next step.


15 **Step-6: 8-(1-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione**

¹HNMR(400MHz, DMSO d₆): δ 0.86-0.91 (m, 6H); 1.51 (d, J=6.8Hz, 3H); 1.55-1.61 (m, 2H); 1.69-1.74 (m, 2H); 2.53-2.59 (m, 1H); 2.98-3.04 (m, 2H); 3.66-3.69 (m, 2H); 3.86 (t, J=7.2Hz, 2H); 3.98 (t, J=6.4Hz, 2H); 4.58-4.62 (m, 1H); 7.45-7.47 (d, J=8.0Hz, 1H); 7.54-7.58 (m, 1H); 7.66 (d, J=8.4Hz, 1H); 8.11-8.13 (m, 2H); 8.46 (s, 1H); 13.51 (bs, 1H).

20

Example Q1: 8-(1-{2-Oxo-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

5



Step-1: 4-(2-bromo-acetyl)-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one

- 10 To a stirred solution of 5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidine-3-carboxylic acid (1.0 g, 3.66 mmol) in DCM (10 ml) was added thionyl chloride (0.9 ml, 12.44 mmol) and DMF (2 drops). The reaction mixture was stirred for 1hr at room temperature and the solvents were removed. The residue was taken in toluene (10 ml) and again concentrated to obtain brown oil. The resulting oil was taken in acetonitrile

(10 ml) and trimethylsilyldiazomethane (9.15 ml, 18.3 mmol). It was stirred at room temperature for 1hr. The reaction mixture was cooled to 0 °C and 33 % HBr in acetic acid (4.17 ml, 23.79 mmol) was added drop wise and stirred for 1 hr at room temperature. To the reaction mixture sodium bicarbonate (20 ml) was added and
 5 extracted with ethyl acetate (3 X 15 ml). The combined organic layer was washed with saturated brine solution (35 ml). The organic layer was dried over Na₂SO₄, evaporated to obtain crude product. The crude product was purified by column chromatography using silicagel (100-200) and 12 to 16 % ethyl acetate in hexane as an eluent to obtain
 10 4-(2-bromo-acetyl)-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one (0.9g, 72 %) as a brown mass.

¹HNMR(400MHz, CDCl₃): δ 2.28-3.0 (m, 2H); 3.89-3.96 (m, 2H); 3.99-4.06 (m, 2H); 4.13-4.17 (m, 1H); 4.42-4.44 (m, 1H); 7.48-7.52 (m, 1H); 7.82 (s, 1H); 7.84-7.86 (m, 1H).

Step-2: 8-(1-{2-Oxo-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione
 15

A mixture of 1, 3-dipropyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilanyl-ethoxymethyl)-3, 7-dihydro-purine-2, 6-dione (0.200g, 0.462mmol), 4-(2-Bromo-acetyl)-1-(3-trifluoromethyl-phenyl)-pyrrolidin-2-one (0.240g, 0.693mmol), Cs₂CO₃ (0.30g, 0.924mmol) and acetonitrile (2ml) were heated at 90-100 °C in a sealed tube for 6hrs. Reaction mixture was cooled to 20-25 °C and filtered off. The organic volatiles were evaporated. The crude product was purified by preparative TLC to obtain pure 8-(1-{2-oxo-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione
 25 (0.090g, 28 %) as a brown mass.


¹HNMR(400MHz, CDCl₃): δ 0.10 (s, 9H); 0.93-1.03 (m, 8H); 1.70-1.85 (m, 4H); 2.82-2.86 (m, 2H); 3.55-3.59 (m, 1H); 3.80-3.84 (m, 2H); 3.93-4.01 (m, 3H); 4.10-4.14 (m, 2H); 4.19-4.23 (m, 1H); 5.15 (s, 2H); 5.81 (d, J=2.0Hz, 2H); 7.42-7.44 (m, 1H); 7.48-7.52 (m, 1H); 7.80-7.85 (m, 2H); 8.23 (s, 1H); 8.28 (s, 1H).

Step-3: 8-(1-{2-Oxo-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
 30

The reaction was carried out as described before.

¹HNMR(400MHz, DMSO d6): δ 0.82-0.87 (m, 6H); 1.51-1.57 (m, 2H); 1.65-1.71 (m, 2H); 2.80-2.83 (m, 2H); 3.64-3.68 (m, 1H); 3.82 (t, J=7.2Hz, 2H); 3.94 (t, J=6.8Hz, 2H); 4.09-4.11 (m, 2H); 5.43 (d, J=1.6Hz, 2H); 7.48 (d, J=7.6Hz, 1H); 7.59-7.63 (m, 1H); 7.83 (d, J=8.4Hz, 1H); 8.08-8.11 (m, 2H); 8.34 (s, 1H); 13.56 (bs, 1H).


Example R1: 8-(1-{2-Hydroxy-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



10 The reaction was carried out by following the reaction conditions and procedure as described before in step-3 of Example K1

¹HNMR(400MHz, DMSO d6): δ 0.82-0.87 (m, 6H); 1.51-1.57 (m, 2H); 1.66-1.71 (m, 2H); 2.40-2.43 (m, 1H); 2.53-2.61 (m, 2H); 3.72-3.74 (m, 1H); 3.80-3.85 (m, 2H); 3.85-3.96 (m, 4H); 4.15-4.26 (m, 2H); 5.39-5.44 (m, 1H); 7.45 (d, J=7.6Hz, 1H); 7.57-7.61 (m, 1H); 7.83 (d, J=8.0Hz, 1H); 8.0 (s, 1H); 8.12 (s, 1H); 8.31(s, 1H); 13.45 (bs, 1H).

Example S1: 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl] 1Hpyrazol4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



Step-1: Chloro oxime-acetic acid ethyl ester

A mixture of solution of amino-acetic acid ethyl ester (5 g, 48.53 mmol) in water (5 ml) and conc. HCl (8 ml), NaNO₂ (4.91 g, 145.6 mmol) with water (5 ml) was added slowly at 0 °C stirred for 1 hr at room temperature. After completion of reaction, the compound was extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated under vacuum to get 1.5 g (20 %) of chloro oxime-acetic acid ethyl ester as white solid.

¹HNMR(400MHz, CDCl₃): δ 1.37-1.41 (t, J=7.2Hz, 3H); 4.38-4.43 (q, J=7.2Hz, 2H); 9.59-9.61 (m, 1H).

Step-2: 5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazole-3-carboxylic acid ethyl ester

To a mixture of 1-Allyl-4-fluoro-benzene (0.27 g, 0.198 mmol) and NaHCO₃ (0.166 gm, 0.198 mmol) in [Bmim] [BF₄] (0.2 ml), chloro oxime-acetic acid ethyl ester (0.3 g, 0.198 mmol) was added at room temperature. After stirring the reaction mixture for 6 h, the sticky mass obtained was washed with diethyl ether. The ether layer was concentrated to obtain a yellow residue which was further purified by column chromatography to obtain 5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazole-3-carboxylic acid ethyl ester (170 mg, 34 %) as a white solid.

54348-1

¹HNMR(400MHz, CDCl₃): δ 1.34-1.41 (m, 3H); 2.87-2.94 (m, 2H); 3.02-3.07 (m, 1H); 3.17-3.24 (m, 1H); 4.30-4.41 (m, 2H); 5.00-5.05 (m, 1H); 6.98-7.03 (m, 2H); 7.18-7.26 (m, 2H).

Step-3: [5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-yl]-methanol

To a solution of 5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazole-3-carboxylic acid ethyl ester (0.17 g, 0.677 mmol) in THF (10 ml), NaBH₄ (0.051 g, 0.81 mmol) was added slowly at 0 °C. The reaction mixture was stirred for 2 h at room temperature. Water was added to the reaction mixture and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over Na₂SO₄ and concentrated under vacuum. The residue obtained was purified by column chromatography to obtain [5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-yl]-methanol (140 mg, 100 %) as an yellow oil.

¹HNMR(400MHz, CDCl₃): δ 2.78-2.91 (m, 2H); 3.09-3.17 (m, 2H); 4.41 (s, 2H); 4.92-4.99 (m, 1H); 7.01-7.09 (m, 2H); 7.21-7.25 (m, 2H).

Step-4: Methanesulfonic acid 5-(4-fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl ester

To a mixture of solution of [5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-yl]-methanol (80 mg, 0.38 mmol), TEA (57 mg, 0.57 mmol) in DCM (10 ml), methanesulfonyl chloride (43 mg, 0.38 mmol) was added slowly at 0 °C and the mixture was stirred at room temperature for 3 hrs. Aqueous NaHCO₃ was added to the reaction mixture and extracted with methylene dichloride. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum. The residue obtained was purified by column chromatography to obtain Methanesulfonic acid 5-(4-fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl ester (82 mg, 75 %) as brown oil.

Step-5: 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione

A mixture of solution of Methanesulfonic acid 5-(4-fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl ester (73 mg, 0.25 mmol), 1,3-dipropyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (100 mg, 0.23 mmol) and K₂CO₃ (63 mg, 0.46 mmol) in acetonitrile (10 ml) was heated at 80 °C for 3 hrs. The reaction mixture was filtered through celite™ and washed with acetone. The organic layer was concentrated and the oil obtained was purified by column chromatography to obtain 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-

1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (85 mg, 60 %) as brown oil.


¹HNMR(400MHz, CDCl₃): δ 0.01 (s, 9 H); 0.98-1.08 (m 8H); 1.64-2.08 (m, 4H); 2.68-2.97 (m, 1H); 2.90-3.09 (m, 3H); 3.82-3.97 (m, 2H); 4.01-4.07 (m, 2H); 4.23-4.31 (m, 2H); 4.95-5.00 (m, 1H); 5.12 (s, 2H); 5.81-5.95 (m, 2H); 6.97-7.06 (m, 2H); 7.18-7.22 (m, 2H); 8.27 (s, 1H); 8.59 (s, 1H).

Step-6: 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

To a solution of 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (80 mg) in ethanol (5 ml) aq. HCl (2 N, 5 ml) was added and the reaction mixture was refluxed for 2 hrs. The volatiles were evaporated and the residue obtained was dissolved in ethyl acetate and washed with water and brine. The organic layer was dried over Na₂SO₄ and concentrated under vacuum. The residue obtained was purified by column chromatography to obtain 8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (100 mg, 98 %) as white solid.

¹HNMR(400MHz, CDCl₃): δ 0.82-0.98 (m, 6H); 1.56-1.64 (m, 2H); 1.71-1.8 (m, 2H); 2.58-2.62 (m, 2H); 2.78-2.98 (m, 3H); 3.82-3.97 (m, 2H); 4.01-4.07 (m, 2H); 4.79-4.84 (m, 1H); 5.19 (s, 1H); 7.07-7.16 (m, 2H); 7.24-7.28 (m, 2H); 8.17 (s, 1H); 8.41 (s, 1H).

Example T1: 8-(1-{2-Oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



Step-1: {4-[2,6-Dioxo-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-acetic acid

- 5 A mixture of solution of 1,3-dipropyl-8-(1H-pyrazol-4-yl)-7-(2-trimethylsilyl-ethoxymethyl)-3,7-dihydro-purine-2,6-dione (600 mg, 1.38 mmol), bromoacetic acid (289 mg, 2.08 mmol) and K_2CO_3 (575 mg, 4.16 mmol) in acetone (10 ml) was heated at $80^\circ C$ for 3 hrs. The reaction mixture was filtered through celite and washed with acetone. The organic layer was concentrated and the residue obtained was purified by
- 10 column chromatography to obtain {4-[2,6-dioxo-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-acetic acid (420 mg, 61%) as white solid.

Step-2: [4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-acetyl chloride

- 15 {4-[2,6-Dioxo-1,3-dipropyl-7-(2-trimethylsilyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-pyrazol-1-yl}-acetic acid (350 mg, 0.71 mmol) was taken with thionyl chloride (25 ml) and refluxed for 3 hrs. The volatiles were evaporated to obtain [4-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-acetyl chloride (400 mg) as a yellow solid.

Step-3: 8-(1-{2-Oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione


To a solution of [4-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-acetyl chloride (100 mg, 0.26 mmol) in THF (10 ml) was added TEA (53 mg, 0.53 mmol) and 1-(3-Trifluoromethyl-phenyl)-piperazine (66 mg, 0.29 mmol) at 0 °C and the mixture was stirred overnight. The reaction mixture was concentrated and purified by HPLC to obtain 8-(1-{2-Oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-

pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (30 mg, 21%) as a white solid.
¹HNMR(400MHz, DMSO d₆): δ 0.85-0.91 (m, 6H); 1.55-1.60 (m, 2H); 1.69-1.75 (m, 2H); 3.24-3.3 (m, 2H); 3.60-3.70 (m, 2H); 3.83-3.87 (t, J=7.6 Hz, 2H); 3.96 - 4.00 (t, J= 6.8 Hz, 2H); 5.31 (s, 2H); 7.10-7.12 (d, J= 7.2 Hz, 4H); 7.28 (s, 1H); 7.26-7.28 (d, J= 8.4, 1H); 7.43-7.47 (t, J= 8 Hz, 1H); 8.05 (s, 1H); 8.30 (s, 1H); 13.5(bs, 1H).

Examples T2 – T9 were prepared in an analogous manner of Example T1 from the appropriate intermediate.

Example	IUPAC name
T2	8-(1-{2-[4-(4-Chloro-phenyl)-4-hydroxy-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)1,3-dipropyl-3,7-dihydro-purine-2,6-dione
T3	8-(1-{2-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
T4	8-{1-[2-Oxo-2-(4-p-tolyl-piperazin-1-yl)-ethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
T5	8-(1-{2-Oxo-2-[4-(4-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
T6	8-(1-{2-[4-(3-Chloro-4-fluoro-phenyl)-piperazin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
T7	3-Ethyl-8-(1-{2-oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1-propyl-3,7-dihydro-purine-2,6-dione
T8	8-(1-{2-[4-(4-Fluoro-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
T9	8-(1-{2-[4-(3-Fluoro-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

Example U1: 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid



5

Step-1: 4-Oxo-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester

To a solution of 4-hydroxy-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester (8 g, 32.65 mmol) in DCM (200 ml), pyridinium dichromate (24.55 g, 65.3 mmol) was added at 0 °C. It was stirred at room temperature for overnight. Reaction mixture was filtered through celite pad and DCM layer was concentrated. Purification

10

by column chromatography offered 4-oxo-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester as a white solid (6 g, 76 %).

¹HNMR(400MHz, CDCl₃): δ 1.45 - 1.47 (Two s, 9H); 2.55-2.60 (m, 1H); 2.88-3.0 (m, 1H); 3.75 (s, 3H); 3.87-3.90 (d, 2H, J=13.2 Hz); 4.69-4.82 (m, 1H).

5 **Step-2: 4-Methylene-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester**

To a suspension of methyltriphenylphosphonium iodide (1.25 g, 3.08 mmol) in THF (20 ml) at 0 °C, 4-oxo-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester (500 mg, 2.05 mmol) and potassium ter-butoxide (460 mg, 4.11 mmol) was
10 added. It was stirred for overnight at room temperature. Water was added and the compound was extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over sodium sulphate, filtered and concentrated under vacuum. Purification by column chromatography offered 4-methylene-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester as a colorless oil (0.23 g, 46 %).

15 ¹HNMR(400MHz, CDCl₃): δ 1.45 - 1.50 (Two s, 9H); 2.62-2.68 (m, 1H); 2.93- 3.06 (m, 1H); 3.76 (s, 3H); 4.08-4.12 (d, 2H, J=15.2 Hz); 4.41-4.55 (m, 1H); 5.03-5.06 (m, 2H).

Step-3: 4-Hydroxymethyl-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester

20 To a solution of 4-methylene-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester (1g, 4.149 mmol) in THF (25 ml), 0.5 M solution of 9-BBN in THF (9.8 ml, 4.979 mmol) was added at 0 °C and stirred for 2hrs at room temperature. The reaction mixture was cooled to 0 °C, and then 2N NaOH (2.43 ml, 4.979 mmol) and 30% H₂O₂ (1.6 ml, 14.93 mmol) were added. The reaction mixture was stirred at room
25 temperature for 4hr and later it was poured into water. The mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with brine, filtered and concentrated under vacuum. Purification by column chromatography offered 4-hydroxymethyl-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester as a yellow oil (6 g, 56 %).

30 ¹HNMR(400MHz, CDCl₃): δ 1.43 - 1.48 (Two s, 9H); 1.74-1.81 (m, 1H); 2.37-2.58 (m, 2H); 3.20-3.29 (m, 1H); 3.58-3.72 (m, 3H); 3.74 (s, 3H).

Step-4: 4-Methanesulfonylmethyl-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester

To a solution of 4-hydroxymethyl-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester (0.5 g, 1.93 mmol) and TEA (0.33 gm, 2.89 mmol) in DCM (20 ml),
 5 methane sulphonyl chloride (0.38g, 3.86 mmol) was added at 0 °C. The reaction mixture was stirred at room temperature for 4h. Aqueous NaHCO₃ solution was added to it and extracted with DCM. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to offer 4-methanesulfonylmethyl-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester as a yellow solid
 10 (0.45 g, 69 %). ¹HNMR(400MHz, CDCl₃): δ 1.45 - 1.50 (Two s, 9H); 1.79-1.85 (m, 1H); 2.42-2.81 (m, 2H); 3.08 (s, 3H); 3.21-3.38 (m, 1H); 3.77 (s, 3H); 4.09-4.4 (m, 3H).

Step-5: 4-{4-[2, 6-Dioxo-1, 3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-2, 3, 6, 7-tetrahydro-1H-purin-8-yl]-phenoxy-methyl}-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester

A mixture of 8-(4-hydroxy-phenyl)-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-3,7-dihydro-purine-2,6 dione (100mg, 0.231 mmol), 4-methanesulfonyl methyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester (82 mg, 0.254 mmol) and K₂CO₃ (63 mg, 0.462 mmol) in DMF (8 ml) was heated at 80 °C for 3 hrs.
 20 The reaction mixture was filtered through celite and washed with ethyl acetate. The filtrate was washed with water, brine, dried over sodium sulphate, filtered and concentrated under vacuum. The residue obtained was purified by column chromatography to offer 4-{4-[2, 6-dioxo-1, 3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-2, 3, 6, 7-tetrahydro-1H-purin-8-yl]-phenoxy-methyl}-pyrrolidine-1, 2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester as a yellow oil (110 mg, 36 %).
 25

¹HNMR(400MHz, CDCl₃): δ 0.1 (s, 9H); 0.98-1.04 (m, 8H); 1.45 - 1.47 (Two s, 9H); 1.62-1.78 (m, 3H); 1.82-1.96 (m, 2H); 2.52-2.95 (m, 3H); 3.38-3.42 (m, 1H); 3.77 (s, 3H); 3.81-3.95 (m, 3H); 3.98-4.09 (m, 4H); 4.17-4.22 (m, 2H); 5.72 (s, 2H); 6.98-7.03 (m, 2H); 7.94-7.98 (m, 2H).

Step-6: 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-pyrrolidine-2-carboxylic acid methyl ester

To a solution of 4-{4-[2,6-Dioxo-1,3-dipropyl-7-(2-trimethylsilanyl-ethoxymethyl)-2,3,6,7-tetrahydro-1H-purin-8-yl]-phenoxy-methyl}-pyrrolidine-1,2-dicarboxylic acid 1-

tert-butyl ester 2-methyl ester (100 mg) in methanol (3 ml), methanolic HCl (3 N, 0.5 ml) was added at 0 °C, reaction mixture was stirred at room temperature for 2 hrs. The reaction mixture was evaporated and the residue obtained was purified by column chromatography to offer 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-pyrrolidine-2-carboxylic acid methyl ester as a yellow solid (50 mg, 74 %).

¹HNMR(400MHz, CDCl₃): δ 0.65-0.79 (m, 6H); 1.37-1.59 (m, 4H); 1.98-2.86 (m, 3H); 3.15-3.18 (m, 2H); 3.37-3.42 (m, 1H); 3.61-3.68 (m, 2H); 3.69-3.79 (m, 5H); 3.91-3.98 (m, 2H); 6.78-6.83 (m, 2H); 7.48-7.58 (m, 2H).

Step-7: 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid methyl ester

To a solution of 4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-pyrrolidine-2-carboxylic acid methyl ester (50 mg, 0.106 mmol) and TEA (32 mg, 0.319 mmol) in DCM (5 ml), 3-trifluoro methyl benzyl bromide (30 mg, 0.127 mmol) was added at 0 °C. Reaction mixture was stirred at room temperature for 4hr. Water was added into it and extracted with DCM. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to offer 4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid methyl ester as an off white solid (25 mg, 37 %).

¹HNMR(400MHz, CDCl₃): δ 0.98-1.14 (m, 6H); 1.62-1.78 (m, 3H); 1.82-1.96 (m, 2H); 2.84-3.15 (m, 2H); 3.38-3.78 (m, 2H); 3.81-3.98 (m, 5H); 3.98-4.21 (m, 5H); 4.51-4.77 (m, 2H); 6.98-7.03 (m, 2H); 7.62-7.78 (m, 3H); 8.07-8.12 (m, 1H); 8.31-8.42 (bs, 2H); 13.91 (bs, 1H).


Step-8: 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid.

A mixture of solution of 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid methyl ester (25 mg, 0.039 mmol) in THF (3 ml) and MeOH (1 ml) was added aq. solution of LiOH (3 mg, 0.079 mmol) in H₂O (1 ml) at 0 °C and it was stirred for 2 hrs. Reaction mixture was concentrated, residue was diluted with water and acidified with aq. HCl. The aqueous layer was extracted with ethyl acetate and washed with brine.

The ethyl acetate layer was concentrated and purified by column chromatography to offer 4-[4-(2, 6-Dioxo-1, 3-dipropyl-2, 3, 6, 7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid as viscous oil (18 mg, 75 %).

- 5 ¹HNMR(400MHz, CDCl₃): δ 0.82-0.99 (m, 6H); 1.58-1.62 (m, 2H); 1.62-1.78 (m, 2H); 2.64-3.87 (m, 1H); 2.98-3.08 (m, 1H); 3.23-3.38 (m, 1H); 3.61-3.68 (m, 1H); 3.81-3.98 (m, 2H); 3.98-4.18 (m, 5H); 4.27-4.34 (m, 1H); 4.75-4.87 (m, 2H); 6.98-7.03 (m, 2H); 7.52-7.78 (m, 2H); 7.81-8.92 (m, 2H); 8.11-8.14 (m, 2H); 13.62 (bs, 1H).

10 **Example V1: 3-(2-Amino-ethyl)-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione**



15 **Step-1: 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid ethyl ester**

- 20 In an oven dried two-necked round-bottomed flask, the methanesulfonic acid 5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl ester (3.2 g, 9.4 mmol), 1H-pyrazole-4-carboxylic acid ethyl ester (1.46 g, 10.0 mmol), and potassium carbonate (3.93 g, 28.0 mmol) were mixed under an argon atmosphere followed by the addition of the dry acetone (60 mL) at room temperature. The reaction mixture was refluxed at 80 °C for 24h. The reaction mixture was filtered, evaporated and extracted with DCM (100 x 3 mL), dried over Na₂SO₄ and concentrated on a rotary evaporator to afford the crude

product mixture. The product was isolated by flash column chromatography (eluting with hexane:EtOAc, 2:3) as a viscous oil (2.7 g, 75%).

¹HNMR(400MHz, CDCl₃): δ 1.35(t, J= 7.0 Hz, 3H), 2.46(dd, J= 17.0, 7.0Hz, 1H), 2.83(dd, J= 17.6, 8.0Hz, 1H), 3.14(m, 1H), 3.74(dd, J= 9.6, 6.0Hz, 1H), 3.97(t, J= 8.0Hz, 1H), 4.26-4.33(m, 4H), 7.41(d, J= 8.0Hz, 1H), 7.47(t, J= 8.0Hz, 1H), 7.81(br s, 2H), 7.94(d, J= 6.8Hz, 2H).

Step-2: 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid

To a stirred solution of 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid ethyl ester (2.6 g, 6.81 mmol) in 14 mL MeOH:H₂O (3:11), NaOH (0.4 g, 10.5 mmol) was added at room temperature. The mixture was heated at 100 °C for 2 h. The reaction mixture was cooled; MeOH was removed under reduced pressure, diluted with water, washed with DCM and acidified with 1% HCl. The acidified mixture was extracted with EtOAc (3x50 mL) and the combined extract was washed with brine, dried with Na₂SO₄, evaporated under reduced pressure to give the title compound in 88% yield (2.1 g).

¹HNMR(400MHz, DMSO d₆): δ 2.43(dd, J= 17.0, 6.0Hz, 1H), 2.69(dd, J= 17.2, 8.8Hz, 1H), 3.01(m, 1H), 3.72(dd, J= 10.0, 5.0Hz, 1H), 3.98(t, J= 8.0Hz, 1H), 4.31(d, J= 6.8Hz, 2H), 7.49(d, J= 7.6Hz, 1H), 7.612(t, J= 8.0Hz, 1H), 7.75(d, J= 8.0Hz, 1H), 7.82(s, 1H), 8.13(s, 1H), 8.36(s, 1H), 12.4(br s, 1H).

Step-3: 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid (6-amino-2,4-dioxo-3-propyl-1,2,3,4-tetrahydropyrimidin-5-yl)-amide

To a stirred solution of 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid (1 g, 2.83 mmol) and 5,6-Diamino-3-propyl-1H-pyrimidine-2,4-dione (521 mg, 2.83 mmol) in MeOH (40 mL) was added EDCI (813 mg, 4.24 mmol) under argon atmosphere. After stirring at overnight at room temperature the reaction mixture was quenched with water (10 mL) and white solid was precipitated out. The solid was filtered and washed with water to give the crude product which on purification by column chromatography (eluting with 5% MeOH:DCM) furnished the title compound as a white solid (1.3 g, 88 %).

¹HNMR(400MHz, DMSO d₆): δ 0.80(t, J= 7.0Hz, 3H), 1.47(m, 2H), 2.40(dd, J= 17.0, 6.4Hz, 1H), 2.66(dd, J= 16.8, 8.8Hz, 1H), 2.98(m, 1H), 3.62(t, J= 7.8Hz, 2H), 3.69(dd, J= 9.0, 5.0Hz, 1H), 3.96(t, J= 8.0Hz, 1H), 4.29(d, J= 6.4Hz, 2H), 6.00(br s, 2H), 7.47(d, J= 7.2Hz, 1H), 7.60(t, J= 8.0Hz, 1H), 7.74(d, J=8.0Hz, 1H), 7.97(s, 1H), 8.15(s, 1H),
5 8.29(s, 1H), 8.53(s, 1H), 10.4(br s, 1H).

Step-4: 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid {6-amino-1-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-2,4-dioxo-3-propyl-1,2,3,4-tetrahydro-pyrimidin-5-yl}-amide

A mixture of compound 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-
10 1H-pyrazole-4-carboxylic acid (6-amino-2,4-dioxo-3-propyl-1,2,3,4-tetrahydro-pyrimidin-5-yl)-amide (250 mg, 0.48 mmol), K₂CO₃ (133 mg, 0.96 mmol), dry DMF (4 ml) and 2-(2-bromo-ethyl)-isoindole-1,3-dione (183 mg, 0.72 mmol) was stirred at room temperature for over night. After that brine was added to the reaction mixture and extracted with EtOAc (5X10 ml). The combined extract was dried over Na₂SO₄,
15 filtered and concentrated under vacuum. The residue was purified by column chromatography using 3% MeOH and DCM as eluent to furnish the title compound (120 mg, 36 %).

¹HNMR(400MHz, DMSO d₆): δ 0.55 (t, J= 7.6Hz, 3H), 1.05(m, 2H), 2.45(m, 1H), 2.7(m, 1H), 3.0(m, 1H), 3.3-3.5(m, 2H), 3.76(m, 1H), 3.83(br t, 2H), 4.03(m, 1H),
20 4.25(br t, 2H), 4.35(m, 2H), 6.78(s, 2H), 7.50(m, 1H), 7.64(m, 1H), 7.78(m, 1H), 7.84(s, 4H), 8.06(s, 1H), 8.2(s, 1H), 8.36(s, 1H), 8.64(s, 1H).

Step-5: 3-[2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione

25 1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazole-4-carboxylic acid {6-amino-1-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-2,4-dioxo-3-propyl-1,2,3,4-tetrahydro-pyrimidin-5-yl}-amide (100 mg, 0.14 mmol) was taken in dry DMF (3 ml) under argon atmosphere, added P₂O₅ (61.4 mg, 0.43 mmol) and dipped the reaction flask in a oil bath at 100 °C and stirred the reaction mixture for 5
30 min. Then the reaction mixture was cooled to room temperature and diluted with water and filtered the solid (56 mg, 57%) as pure product.

¹HNMR(400MHz, DMSO d₆): δ 0.79(t, J= 7.6Hz, 3H), 1.42(m, 2H), 2.42(dd, J= 11.2Hz, 6.0Hz 1H), 2.7(m, 1H), 3.0(m, 1H), 3.75(m 3H), 3.9-4.04(m, 3H), 4.3(m, 4H),


7.50(d, J= 7.6Hz 1H), 7.62-7.67(m, 3H), 7.73-7.83(m, 4H), 8.06(s, 1H), 8.18(s, 1H), 13.4(s, 1H).

Step-6: 3-(2-Amino-ethyl)-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione

5 3-[2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione (56 mg, 0.083 mmol) was taken in EtOH (10 ml) to this monohydrated N_2H_4 (8.3 mg, 0.16 mmol) was added and stirred the reaction mixture for 1 h at room temperature. After completion of the reaction, solvent was evaporated under vacuum, solid was
10 filtered, washed with DCM and the filtrate was concentrated to give the pure product (20 mg, 44%) as white solid.

^1H NMR(400MHz, DMSO d_6): δ 0.89(t, J= 8.4Hz, 3H), 1.59(m, 2H), 2.45(m, 1H), 2.71(dd, J= 17.2, 8.8Hz 1H), 3.03(m, 3H), 3.74(m 1H), 3.85(t J= 7.6Hz, 2H), 4.02(t, J= 8.4Hz, 1H), 4.15(br t 2H), 4.36(d, J= 6.8Hz 2H), 7.49(d, J= 7.6Hz 1H), 7.62 (t, J= 8.0Hz 1H), 7.76(d, J= 8.0Hz 1H), 8.08(s, 1H), 8.14(s, 1H), 8.41(s, 1H).
15

Example W1: 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione



20 A mixture of 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione (0.030g, 0.057 mmol), potassium carbonate (0.016g, 0.115 mmol), methyl iodide (0.0039 ml, 0.0639 mmol) and DMF (1 ml) was heated at 50 °C for 20 hour. The mixture was cooled to room temperature and diluted with water (10 ml). The aqueous layer was extracted with ethyl acetate. The
25 organic layer was washed with water and saturated brine solution, and dried over

Na₂SO₄. The solvent was evaporated and the residue obtained was triturated with hexane to obtain the title compound (0.026 g, 84 %).

¹HNMR(400MHz, DMSO d₆): δ 0.85-0.90 (m, 6H); 1.24 (d, J=6.8Hz, 6H); 1.53-1.58 (m, 2H); 1.69-1.72 (m, 2H); 2.40-2.44 (m, 1H); 2.63-2.74 (m, 1H); 2.82-2.85 (m, 1H);
 5 2.89-3.01 (m, 1H); 3.61-3.65 (m, 1H); 3.83-3.91 (m, 3H); 3.96 (t, J=6.8Hz, 2H); 4.00 (s, 3H); 4.38 (d, J=6.8Hz, 2H); 7.22 (d, J=8.4Hz, 2H); 7.50 (d, J=8.4Hz, 2H); 8.05 (s, 1H); 8.54 (s, 1H); 13.46 (s, 1H).

10 Examples W2 – W8 were prepared in an analogous manner of Example W1 from the appropriate intermediate.

Example	IUPAC name
W2	7-Methyl-8-{1-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
W3	8-{1-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
W4	7-Methyl-8-{1-[5-oxo-1-(2-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
W5	8-{6-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
W6	8-{6-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
W7	8-{4-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione
W8	8-{4-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione

Biological activity

Radioligand Binding for A_{2B} Adenosine Receptor

Human A_{2B} adenosine receptor cDNA was stably transfected into HEK-293 cells.
 15 HEK-A_{2B} cells were harvested by trypsinization with 0.25% Trypsin-EDTA (Sigma),

and washed in 1X PBS at 1500 rpm for 5 minutes at room temperature. The cells were washed twice in wash buffer containing 150 mM NaCl, 1 mM EDTA, 50 mM Tris (pH-7.4) at 1500 rpm for 10 minutes at room temperature and incubated for 10 min at 4°C in sonication buffer containing 1 mM EDTA, 5 mM Tris (pH 7.4). The cells were
5 sonicated on ice for 6 min with six intermittent pulses of 9 second each and centrifuged at 1000Xg for 10 minutes at 4°C. The pellet was discarded and the supernatant was centrifuged at 49,000Xg for 45 minutes at 4°C. The protein pellet was resuspended in buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4), 1 Unit/ml adenosine deaminase (ADA) and incubated for 30 minutes at room temperature. The lysate was washed twice
10 with buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4) at 49,000Xg for 45 minutes at 4°C and the protein pellet was resuspended in 50 mM Tris, pH-7.4 supplemented with 1 Unit/ml ADA and 10% sucrose. Frozen aliquots were stored at -80°C.

Competition assays were started by mixing 1.6 nM [3H]-MRS-1754 with various concentrations of test compounds and 10 µg membrane protein in Reaction buffer (50
15 mM Tris pH 6.5, 5 mM MgCl₂, 1 mM EDTA) supplemented with 1U/ml Adenosine deaminase. The assay reactions were incubated for 90 minutes at room temperature and stopped by filtration using 96 well-plate harvester (Molecular Devices) and washed four times with ice cold 50 mM Tris (pH 7.4). Non specific binding was determined in presence of 200 µM NECA. Radioligand binding was read at Liquid scintillation
20 counter (Perkin Elmer) and the affinities of compounds (i.e. K_i values) were calculated using GraphPad software.

Compounds tested had micromolar to nanomolar activity.

Radioligand Binding for A₁ Adenosine Receptor

Human A₁ adenosine receptor cDNA was stably transfected into HEK-293 cells. HEK-
25 A_{2B} cells were harvested by trypsinization with 0.25% Trypsin-EDTA (Sigma), and washed in 1X PBS at 1500 rpm for 5 minutes at room temperature. The cells were washed twice in wash buffer containing 150 mM NaCl, 1 mM EDTA, 50 mM Tris (pH-7.4) at 1500 rpm for 10 minutes at room temperature and incubated for 10 min at 4°C in sonication buffer containing 1 mM EDTA, 5 mM Tris (pH 7.4). The cells were
30 sonicated on ice for 6 min with six intermittent pulses of 9 seconds each and centrifuged at 1000Xg for 10 minutes at 4°C. The pellet was discarded and the supernatant was centrifuged at 49,000Xg for 45 minutes at 4°C. The protein pellet was

resuspended in buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4), 1 Unit/ml adenosine deaminase (ADA) and incubated for 30 minutes at room temperature. The lysate was washed twice with buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4) at 49,000Xg for 45 minutes at 4°C and the protein pellet was resuspended in 50 mM Tris, (pH-7.4) supplemented with 1 Unit/ml ADA and 10% sucrose. Frozen aliquots were stored at -80°C.

Competition assays were started by mixing 1 nM [3H]-DPCPX with various concentrations of test compounds and 5 µg membrane protein in Reaction buffer (50 mM Tris pH 7.4, 1 mM EDTA) supplemented with 1 Unit/ml ADA. The assay reactions were incubated for 90 minutes at room temperature and stopped by filtration using 96 well-plate harvester (Molecular Devices) and washed four times with ice cold 50 mM Tris (pH 7.4). Non specific binding was determined in presence of 200 µM NECA. Radioligand binding was read at Liquid scintillation counter (Perkin Elmer) and the affinities of compounds (i.e. K_i values) were calculated using GraphPad software.

Compounds tested had micromolar to nanomolar activity.

Radioligand Binding for A_{2A} Adenosine Receptor

Human A_{2A} adenosine receptor cDNA was stably transfected into HEK-293 cells. HEK-A_{2B} cells were harvested by trypsinization with 0.25% Trypsin-EDTA (Sigma), and washed in 1X PBS at 1500 rpm for 5 minutes at room temperature. The cells were washed twice in wash buffer containing 150 mM NaCl, 1 mM EDTA, 50 mM Tris (pH-7.4) at 1500 rpm for 10 minutes at room temperature and incubated for 10 min at 4°C in sonication buffer containing 1 mM EDTA, 5 mM (Tris pH 7.4). The cells were sonicated on ice for 6 min with six intermittent pulses of 9 second each and centrifuged at 1000Xg for 10 minutes at 4°C. The pellet was discarded and the supernatant was centrifuged at 49,000Xg for 45 minutes at 4°C. The protein pellet was resuspended in buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4) supplemented with 1 Unit/ml adenosine deaminase (ADA) and incubated for 30 minutes at room temperature. The lysate was washed twice with buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4) at 49,000Xg for 45 minutes at 4°C and the protein pellet was resuspended in 50 mM Tris, pH-7.4 supplemented with 1 Unit/ml ADA and 10% sucrose. Frozen aliquots were stored at -80°C.

Competition assays were started by mixing 2 nM [3H]-ZM-241385 with various concentrations of test compounds and 5 µg membrane protein in Reaction buffer (50 mM Tris pH 7.4, 1 mM EDTA) supplemented with 1 Unit/ml ADA. The assay reactions were incubated for 90 minutes at room temperature and stopped by filtration using 96 well-plate harvester (Molecular Devices) and washed four times with ice cold 50 mM Tris (pH 7.4). Non specific binding was determined in presence of 200 µM NECA. Radioligand binding was read at Liquid scintillation counter (Perkin Elmer) and the affinities of compounds (i.e. K_i values) were calculated using GraphPad software.

Compounds tested had micromolar to nanomolar activity.

10 **Radioligand Binding for A₃ Adenosine Receptor**

Human A₃ adenosine receptor cDNA was stably transfected into HEK-293 cells. HEK-A_{2B} cells were harvested by trypsinization with 0.25% Trypsin-EDTA (Sigma), and washed in 1X PBS at 1500 rpm for 5 minutes at room temperature. The cells were washed twice in wash buffer containing 10 mM EDTA, 10mM HEPES (pH-7.4) at 1500 rpm for 10 minutes at room temperature and incubated for 10 min at 4°C in sonication buffer containing 1 mM EDTA, 10 mM HEPES (pH 7.4). The cells were sonicated on ice for 6 min with six intermittent pulses of 9 seconds each and centrifuged at 1000Xg for 10 minutes at 4°C. The pellet was discarded and the supernatant was centrifuged at 49,000Xg for 45 minutes at 4°C. The protein pellet was resuspended in buffer containing 1 mM EDTA, 10 mM HEPES (pH 7.4) supplemented with 1 Unit/ml adenosine deaminase (ADA) and incubated for 30 minutes at room temperature. The lysate was washed twice with buffer containing 1 mM EDTA, 10 mM HEPES (pH-7.4) at 49,000Xg for 45 minutes at 4°C and the protein pellet was resuspended in buffer containing 1 mM EDTA, 5 mM Tris (pH-7.4) supplemented with 1Unit/ml ADA and 10% sucrose. Frozen aliquots were stored at -80°C.

Competition assays were started by mixing 2nM [3H]-HEM-ADO with various concentrations of test compounds and 5 µg membrane protein in Reaction buffer (50 mM Tris pH 7.4, 1 mM EDTA, 10 mM MgCl₂) supplemented with 1Unit/ml ADA. The assay reactions were incubated for 90 minutes at room temperature and stopped by filtration using 96 well-plate harvester (Molecular Devices) and washed four times with ice cold 50 mM Tris pH 7.4. Non specific binding was determined in presence of 200 µM NECA. Radioligand binding was read at Liquid scintillation counter (Perkin

54348-1

Elmer) and the affinities of compounds (i.e. K_i values) were calculated using GraphPad software.


Compounds tested had micromolar to nanomolar activity.

Although the subject matter has been described in considerable detail with reference to certain
5 preferred embodiments thereof, other embodiments are possible. As such, the scope of the
appended claims should not be limited to the description of the preferred embodiment
contained therein.

54348-1

CLAIMS:

1. A compound of formula (I):



I

- 5 or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, wherein,

Y is N or CH;

R¹ and R² are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxyalkyl, carboxyalkyl, cycloalkyl, cycloalkylalkyl, arylalkyl, aryl, heteroaryl,

- 10 heteroarylalkyl, heterocyclyl and heterocyclalkyl;

wherein alkyl, alkenyl, alkynyl, alkoxyalkyl, carboxyalkyl, cycloalkyl, cycloalkylalkyl, arylalkyl, aryl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclalkyl are unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclalkylamino, aminocarbonyl, alkoxyaminocarbonyl, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclalkyl, heterocyclalkyloxy, hydroxyamino, alkoxyamino, nitro, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a or -S(O)_pR^b;

- 20 wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl,

54348-1

aminocarbonyl, hydroxy, alkoxy, halogen, CF_3 , amino, substituted amino, cyano and $-\text{S}(\text{O})_p\text{R}^c$;

wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and
 5 heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2;

R^3 is selected from the group consisting of hydrogen and alkyl;

wherein alkyl is unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino,
 10 cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, carboxyalkyloxy, alkylcarboxyalkyloxy, $-\text{SO}_3\text{H}$, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycliloxy, hydroxyamino, alkoxyamino, CF_3 , nitro, $\text{S}(\text{O})_2\text{NR}^a\text{R}^a$, $-\text{NR}^a\text{S}(\text{O})_2\text{R}^a$ or
 15 $-\text{S}(\text{O})_p\text{R}^b$;

wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF_3 , amino, substituted amino, cyano and $-\text{S}(\text{O})_p\text{R}^c$;

20 wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl; and p is 0, 1 or 2;

X is either an unsubstituted or substituted arylene or an unsubstituted or substituted
 25 heteroarylene;

A is selected from the group consisting of $-\text{O}-$, $-\text{C}(\text{O})\text{NR}^4-$, $-\text{NR}^4\text{C}(\text{O})-$ and $-(\text{CR}^5\text{R}^6)_q-$;

54348-1

wherein q is 1 or 2, and R⁴ is selected from the group consisting of hydrogen and alkyl,

wherein alkyl is either unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclylamino, aminocarbonyl, alkoxy carbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclyloxy, hydroxyamino, alkoxyamino, CF₃, nitro, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a or -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2, or 3 substituents

10 independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, amino, substituted amino, cyano and -S(O)_pR^c;

wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and

15 heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl, or heteroaryl and p is 0, 1 or 2;

R⁵ and R⁶ are independently selected from the group consisting of hydrogen, halogen, alkyl, hydroxyl, alkoxy and -C(O)R⁷;

wherein alkyl and alkoxy are unsubstituted, or substituted with halogen, hydroxy,
20 hydroxyalkyl or CF₃;

or

R⁵ and R⁶ together represent O, S or cycloalkyl;

R⁷ is hydroxyl, or unsubstituted or substituted amino;

m and n are independently 0, 1, 2, 3, 4, 5 or 6;

54348-1

B is selected from the group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted alkenyl, unsubstituted or substituted cycloalkyl and unsubstituted or substituted heterocyclyl; and

D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted
 5 arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted heterocyclalkyl,

wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocyclalkyl are substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino,
 10 monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroarylamino, heterocyclamino, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, CF₃, OCF₃, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclloxy, hydroxyamino, alkoxyamino, nitro, -C(O)R, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and
 15 -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, -OCF₃, amino, substituted amino, cyano and -S(O)_pR^c;

20 wherein R is selected from the group consisting of hydrogen, hydroxyl, alkyl, alkoxy, amino, monoalkylamino, dialkylamino and heterocyclyl; each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl or heteroaryl; and p is 0, 1 or 2.

25 2. A compound of formula (I) as claimed in claim 1, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, wherein Y is N;

54348-1

R^1 and R^2 are independently selected from the group consisting of hydrogen, alkyl, alkenyl and alkynyl;

wherein alkyl, alkenyl and alkynyl are unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, 5 dialkylamino, arylamino, cycloalkylamino, heteroaryl, heterocyclamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, $-SO_3H$, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocycl, heterocycl, hydroxyamino, alkoxyamino, nitro, $-S(O)_2NR^aR^a$, $-NR^aS(O)_2R^a$ or $-S(O)_pR^b$;

- 10 wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocycl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocycl; and p is 0, 1 or 2;

R^3 is selected from a group consisting of hydrogen and alkyl;

- wherein alkyl is unsubstituted or substituted with alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, 15 cycloalkenyl, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, heteroaryl, heterocyclamino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, hydroxyalkyl, keto, thiocarbonyl, carboxy, alkylcarboxy, carboxyalkyl, carboxyalkyloxy, alkylcarboxyalkyloxy, $-SO_3H$, aryl, aryloxy, cycloalkyloxy, heteroaryl, aminocarbonylamino, heteroaryloxy, heterocycl, 20 heterocycl, hydroxyamino, alkoxyamino, CF_3 , nitro, $S(O)_2NR^aR^a$, $-NR^aS(O)_2R^a$ or $-S(O)_pR^b$;

wherein each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocycl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocycl; and p is 0, 1 or 2;

- 25 X is either an unsubstituted or substituted arylene or an unsubstituted or substituted heteroarylene;

54348-1

A is selected from the group consisting of -O-, -C(O)NR⁴-, -NR⁴C(O)- and -(CR⁵R⁶)_q;

wherein q is 1 or 2, and R⁴ is selected from a group consisting of hydrogen and alkyl,

wherein alkyl is either unsubstituted or substituted with halogen, CF₃ or aryl;

R⁵ and R⁶ are independently selected from the group consisting of hydrogen, halogen, alkyl
5 and hydroxyl;

or

R⁵ and R⁶ together represent O or cycloalkyl;

m and n are independently 0, 1 or 2;

10 B is selected from the group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted cycloalkyl and unsubstituted or substituted heterocyclyl; and

D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted heterocyclalkyl,

15 wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocyclalkyl are substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, alkoxy, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, cyano, halogen, hydroxy, hydroxyalkyl, CF₃, OCF₃, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H, aryl, heteroaryl, heterocyclyl, -C(O)R, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and -S(O)_pR^b;

20 wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, -OCF₃, amino, substituted amino, cyano and -S(O)_pR^c;

54348-1

wherein R is selected from the group consisting of amino, monoalkylamino, dialkylamino and heterocyclyl; each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclylalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl or
 5 heteroaryl; and p is 0, 1 or 2.

3. A compound of formula (I) as claimed in claim 1, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof,

wherein Y is N;

R¹ and R² are independently selected from the group consisting of hydrogen and alkyl;

10 wherein alkyl is unsubstituted or substituted with alkenyl, alkoxy, acylamino, amino, monoalkylamino, dialkylamino, halogen, hydroxy, hydroxyalkyl, carboxy, alkylcarboxy, carboxyalkyl, -SO₃H or aryl;

R³ is selected from the group consisting of hydrogen and alkyl;

X is either an unsubstituted or substituted arylene or an unsubstituted or substituted
 15 heteroarylene;

A is selected from the group consisting of -O-, -C(O)NR⁴-, -NR⁴C(O)- and -(CR⁵R⁶)_q-;

wherein q is 1 and R⁴ is selected from the group consisting of hydrogen and alkyl,

wherein alkyl is either unsubstituted or substituted with halogen, CF₃ or aryl;

R⁵ and R⁶ are independently selected from the group consisting of hydrogen, halogen, alkyl
 20 and hydroxyl;

or

R⁵ and R⁶ together represent O or cycloalkyl;

m and n are independently 0, 1 or 2;

54348-1

B is selected from the group consisting of unsubstituted or substituted alkynyl, unsubstituted or substituted cycloalkyl and unsubstituted or substituted heterocyclyl; and

D is selected from the group consisting of substituted aryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, substituted cycloalkylalkyl and substituted
5 heterocyclalkyl,

wherein aryl, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkylalkyl and heterocyclalkyl are substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkyl, alkoxy, acyl, acylamino, acyloxy, amino, monoalkylamino, dialkylamino, arylamino, cycloalkylamino, cyano, halogen, hydroxy, hydroxyalkyl, CF₃, OCF₃, carboxy, alkylcarboxy,
10 carboxyalkyl, -SO₃H, aryl, heteroaryl, heterocyclyl, -C(O)R, -S(O)₂NR^aR^a, -NR^aS(O)₂R^a and -S(O)_pR^b;

wherein each substituent is unsubstituted or substituted with 1, 2 or 3 substituents independently selected from the group consisting of alkyl, carboxy, carboxyalkyl, aminocarbonyl, hydroxy, alkoxy, halogen, CF₃, -OCF₃, amino, substituted amino, cyano and
15 -S(O)_pR^c;

wherein R is selected from the group consisting of amino, monoalkylamino, dialkylamino and heterocyclyl; each R^a is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl and heterocyclalkyl; R^b is hydrogen, alkyl, aryl, heteroaryl or heterocyclyl; R^c is alkyl, aryl or
20 heteroaryl; and p is 0, 1 or 2.

4. A compound of formula (I) as claimed in claim 1, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, wherein X is pyrazolyl, isoxazolyl, phenyl, pyridyl, oxazolyl or pyrimidyl.

5. A compound of formula (I) as claimed in claim 1, or a tautomer, a polymorph,
25 a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, wherein B is an alkynylene.

54348-1

6. A compound of formula (I) as claimed in claim 1, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, wherein B is heterocyclyl or cycloalkyl.

7. A compound of formula (I) as claimed in claim 1, or a tautomer, a polymorph,
5 a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, which is:

1,3-Dipropyl-8-[1-(3-p-tolyl-prop-2ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1, 3-dimethyl-3, 7-dihydro-purine-2, 6-dione,

8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-
10 2,6-dione,

8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione,

15 8-{1-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(2,4-Difluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-
20 dihydro-purine-2,6-dione,

8-{1-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dimethyl-3,7-dihydro-purine-2,6-dione,

54348-1

4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid ethyl ester,

3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid ethyl ester,

5 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzonitrile,

8-{1-[3-(3-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

10 2-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid methyl ester,

8-{1-[4-(4-Fluoro-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[4-(3-Fluoro-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 1,3-Dipropyl-8-[1-(4-p-tolyl-but-3-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[4-(3-trifluoromethyl-phenyl)-but-3-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,

20 1,3-Dipropyl-8-{1-[3-(2-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-[1-(3-m-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

3-Ethyl-1-propyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

54348-1

3-Ethyl-1-propyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

3-Ethyl-1-propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

3-Ethyl-8-{1-[3-(3-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-
5 purine-2,6-dione,

1,3-Dipropyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

10 3-Ethyl-8-{1-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

3-{3-[4-(3-Ethyl-2,6-dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,

3-Ethyl-1-propyl-8-[1-(3-m-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-
15 dione,

3-Ethyl-8-{1-[3-(4-methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

3-Ethyl-1-propyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

20 3-Ethyl-8-{1-[3-(3-methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

4-{3-[4-(3-Ethyl-2,6-dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,

54348-1

4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzonitrile,

(3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenoxy)-acetic acid,

5 8-{1-[3-(3-tert-Butyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,

10 (3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenyl)-acetic acid,

(4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenyl)-acetic acid,

8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-N-isopropyl-benzamide,

1,3-Dipropyl-8-(1-{3-[3-(pyrrolidine-1-carbonyl)-phenyl]-prop-2-ynyl}-1H-pyrazol-4-yl)-3,7-dihydro-purine-2,6-dione,

20 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-4-methyl-benzoic acid,

8-{1-[3-(3-Chloro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-4-methoxy-benzoic acid,

54348-1

5-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-pyridine-2-carboxylic acid methyl ester,

1,3-Dipropyl-8-{3-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

5 8-{3-[3-(2,4-Difluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-
10 2,6-dione,

1,3-Dipropyl-8-{3-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-[3-(3-p-tolyl-prop-2-ynyloxy)-isoxazol-5-yl]-3,7-dihydro-purine-2,6-dione,

8-{3-[3-(3-tert-Butyl-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-
15 purine-2,6-dione,

1,3-Dipropyl-8-{3-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

8-{1-Methyl-5-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 8-{1-Methyl-5-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{5-[3-(3-Methoxy-phenyl)-prop-2-ynyloxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

1,3-Dipropyl-8-{4-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-[4-(3-p-tolyl-prop-2-ynyloxy)-phenyl]-3,7-dihydro-purine-2,6-dione,

5 8-{4-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy]-prop-1-ynyl}-benzoic acid ethyl ester,

3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy]-prop-1-ynyl}-benzoic acid,

10 1,3-Dipropyl-8-{4-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

8-{4-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 1,3-Dipropyl-8-{4-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

8-{4-[3-(3-Methoxy-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{6-[3-(4-Fluoro-phenyl)-prop-2-ynyloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 1,3-Dipropyl-8-{3-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{3-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

54348-1

8-{3-[3-(3-Fluoro-phenyl)-prop-2-ynyloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{3-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

5 1,3-Dipropyl-8-[3-(3-p-tolyl-prop-2-ynyloxy)-phenyl]-3,7-dihydro-purine-2,6-dione,

8-{1-[4-(4-Methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-but-2-ynyl}-piperidine-3-carboxylic acid ethyl ester,

10 1-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-but-2-ynyl}-piperidine-3-carboxylic acid,

8-(1-{4-[4-(4-Chloro-phenyl)-4-hydroxy-piperidin-1-yl]-but-2-ynyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-(1-{4-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-but-2-ynyl}-1H-pyrazol-

15 4-yl)-3,7-dihydro-purine-2,6-dione,

1,3-Dimethyl-8-{1-[4-(4-methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

1-Propyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

20 1-Propyl-8-{1-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

1-Propyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

54348-1

8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

1-Propyl-8-{1-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

5 1-Propyl-8-{1-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

3-{3-[4-(2,6-Dioxo-1-propyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzoic acid,

8-{1-[3-(4-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

7-Methyl-1,3-dipropyl-8-{1-[3-(3-trifluoromethoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

7-Methyl-1,3-dipropyl-8-[1-(3-p-tolyl-prop-2-ynyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

15 7-Methyl-8-{1-[4-(4-methyl-piperazin-1-yl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(3-Fluoro-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(3-Methoxy-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(3-tert-Butyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

(3-{3-[4-(7-Methyl-2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-phenoxy)-acetic acid,

54348-1

8-{1-[3-(3-Hydroxymethyl-phenyl)-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

4-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzenesulfonamide,

5 3-{3-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-prop-1-ynyl}-benzamide,

4-(3-Trifluoromethyl-phenyl)-but-3-ynoic acid [5-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-2-methyl-2H-pyrazol-3-yl]-amide,

10 4-(3-Fluoro-phenyl)-but-3-ynoic acid [5-(2,6-dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-2-methyl-2H-pyrazol-3-yl]-amide,

4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-N-[3-(3-trifluoromethyl-phenyl)-prop-2-ynyl]-benzamide,

8-{1-[4-(3-Fluoro-phenyl)-2-hydroxy-but-3-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 2-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-4-(3-fluoro-phenyl)-but-3-ynoic acid,

2-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-yl]-4-(3-fluoro-phenyl)-but-3-ynoic acid amide,

20 8-{1-[4-(4-Fluoro-phenyl)-4-hydroxy-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[4-(4-Fluoro-phenyl)-but-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[3-(4-Fluoro-phenyl)-1,1-dimethyl-prop-2-ynyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

1,3-Dipropyl-8-{1-[(E)-3-(3-trifluoromethyl-phenyl)-allyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[(Z)-3-(3-trifluoromethyl-phenyl)-allyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

5 1,3-dimethyl-6-[1-[3-(3-fluorophenyl)prop-2-ynyl]pyrazol-4-yl]-5H-pyrrolo[3,2-d]pyrimidine-2,4-dione,

8-[1-[3-(3-fluorophenyl)prop-2-ynyl]pyrazol-4-yl]-1,3-dipropyl-7H-purine-2,6-dione,

8-[1-[4-(3-fluorophenyl)-2-hydroxy-but-3-ynyl]pyrazol-4-yl]-1,3-dipropyl-7H-purine-2,6-dione,

10 8-[1-[3-(4-fluorophenyl)-1,1-dimethyl-prop-2-ynyl]pyrazol-4-yl]-1,3-dipropyl-7H-purine-2,6-dione,

1,3-Dipropyl-8-{1-[2-(3-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

15 8-{1-[2-(3-Fluoro-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[2-(4-Fluoro-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[2-(4-trifluoromethyl-phenyl)-cyclopropylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

20 8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

8-{1-[5-Oxo-1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

5 8-[1-(5-Oxo-1-p-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-[1-(5-Oxo-1-m-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

10 8-{1-[5-Oxo-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(4-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{1-[1-(3-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

3-Ethyl-8-{1-[1-(4-methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

20 3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(4-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

4-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzonitrile,

54348-1

3-Ethyl-8-{1-[1-(4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

3-Ethyl-8-{1-[1-(3-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

5 3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

8-{1-[5-Oxo-1-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

10 3-Ethyl-8-{1-[5-oxo-1-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

3-Ethyl-8-{1-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(3-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{1-[5-Oxo-1-(2-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[5-Oxo-1-(4-trifluoromethoxy-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 8-{1-[1-(4-Methyl-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

4-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzoic acid,

8-{1-[1-(4-Fluoro-benzyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

3-{4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-pyrazol-1-ylmethyl]-2-oxo-pyrrolidin-1-yl}-benzonitrile,

8-{1-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

5 8-{1-[1-(2,4-Difluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(2-Chloro-4-fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(2-Chloro-4-trifluoromethyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

3-Ethyl-8-{1-[5-oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

15 8-{1-[2-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(3-Fluoro-phenyl)-2-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[2-Oxo-1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[1-(4-trifluoromethoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

54348-1

1,3-Dipropyl-8-[1-(1-p-tolyl-pyrrolidin-3-ylmethyl)-1H-pyrazol-4-yl]-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(4-Methoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

5 8-{1-[1-(3-Methoxy-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{1-[1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-3,7-dihydro-purine-2,6-dione,

10 8-{4-[1-(4-Fluorophenyl)-5-oxo-pyrrolidin-3-ylmethoxy]phenyl}-1,3-dipropyl-3,7-dihydropurine-2,6-dione,

1,3-Dipropyl-8-{4-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

8-{3-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{4-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{4-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 8-{6-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

8-{6-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{6-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

5 8-{4-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{4-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

10 8-{6-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[1-(3-Fluoro-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{4-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{3-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{4-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

20 1,3-Dipropyl-8-{3-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

8-{6-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{6-[1-(4-trifluoromethyl-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,

54348-1

8-{6-[1-(4-Fluoro-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{6-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,

5 1,3-Dipropyl-8-{6-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{4-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

10 1,3-Dipropyl-8-{3-[1-(3-trifluoromethyl-phenyl)-piperidin-4-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{6-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,7-dihydro-purine-2,6-dione,

8-{6-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{6-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 8-{4-[5-Oxo-1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{4-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-phenyl}-3,7-dihydro-purine-2,6-dione,

1,3-Dipropyl-8-{3-[1-(4-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

54348-1

8-{3-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{3-[1-(4-Fluoro-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

5 1,3-Dipropyl-8-{3-[1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yloxy]-isoxazol-5-yl}-3,7-dihydro-purine-2,6-dione,

8-{4-[1-(3-Fluoro-phenyl)-pyrrolidin-3-yloxy]-phenyl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

10 8-{3-[1-(2,4-Difluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-isoxazol-5-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{5-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-Methyl-5-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethoxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{5-[1-(4-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{5-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 8-{1-Methyl-5-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethoxy]-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{5-[1-(3-Methoxy-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-1-methyl-1H-pyrazol-3-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-(1-{2-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

8-(1-{1-[5-Oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-(1-{2-Oxo-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

5 8-(1-{2-Hydroxy-2-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[5-(4-Fluoro-benzyl)-4,5-dihydro-isoxazol-3-ylmethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

10 8-(1-{2-Oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-(1-{2-[4-(4-Chloro-phenyl)-4-hydroxy-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-(1-{2-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{1-[2-Oxo-2-(4-p-tolyl-piperazin-1-yl)-ethyl]-1H-pyrazol-4-yl}-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-(1-{2-Oxo-2-[4-(4-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

20 8-(1-{2-[4-(3-Chloro-4-fluoro-phenyl)-piperazin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

3-Ethyl-8-(1-{2-oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-ethyl}-1H-pyrazol-4-yl)-1-propyl-3,7-dihydro-purine-2,6-dione,

8-(1-{2-[4-(4-Fluoro-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

54348-1

8-(1-{2-[4-(3-Fluoro-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-1H-pyrazol-4-yl)-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

4-[4-(2,6-Dioxo-1,3-dipropyl-2,3,6,7-tetrahydro-1H-purin-8-yl)-phenoxy-methyl]-1-(3-trifluoromethyl-benzyl)-pyrrolidine-2-carboxylic acid,

5 3-(2-Amino-ethyl)-8-{1-[5-oxo-1-(3-trifluoromethyl-phenyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-1-propyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(4-Isopropyl-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

7-Methyl-8-{1-[5-oxo-1-(4-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-
10 1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{1-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

7-Methyl-8-{1-[5-oxo-1-(2-trifluoromethyl-benzyl)-pyrrolidin-3-ylmethyl]-1H-pyrazol-4-yl}-
1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

15 8-{6-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{6-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-pyridin-3-yl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione,

8-{4-[1-(4-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-7-methyl-1,3-dipropyl-
20 3,7-dihydro-purine-2,6-dione, or

8-{4-[1-(3-Fluoro-phenyl)-5-oxo-pyrrolidin-3-ylmethoxy]-phenyl}-7-methyl-1,3-dipropyl-3,7-dihydro-purine-2,6-dione.

8. A compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable

54348-1

salt thereof, for the treatment of a disease or disorder susceptible to improvement by antagonism of A_{2B} receptor.

9. A compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, which is an adenosine A₁ and A_{2B} antagonist, an adenosine A_{2B} and A₃ antagonist, or an adenosine A₁, A_{2B} and A₃ antagonist thereby providing dual or pan antagonistic activity through additive or synergistic effect.
10. A compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, for the treatment of asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, an allergic disease, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, or an autoimmune disease.
11. Use of a compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, for the prophylactic or therapeutic treatment of a disease or disorder susceptible to improvement by antagonism of an A_{2B} receptor.
12. Use of a compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, for the modulation of an adenosine A_{2B}, A₁ and A_{2B}, A_{2B} and A₃ or A_{2B}, A₁ and A₃ receptor activity in a mammal.
13. A pharmaceutical composition comprising at least one compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable carriers or excipients.
14. A pharmaceutical composition comprising a compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a

54348-1

solvate or a pharmaceutically acceptable salt thereof, in combination with one or more therapeutically active agents.

15. The pharmaceutical composition as claimed in claim 14, wherein, the therapeutically active agent is an anti-inflammatory agent, an anti-diabetic agent, an
5 anti-hypertensive agent or an anti-dyslipidemic agent.


16. The pharmaceutical composition as claimed in claim 14, wherein the therapeutically active agent is selected from the group consisting of an anticholinergic agent, an antimuscarinic agent, a steroid, a LTB₄ (leukotriene B₄) antagonist, a dopamine receptor agonist, a phosphodiesterase 4 inhibitor, a beta-2 adrenergic receptor agonist, insulin, an
10 insulin derivative, an insulin mimetic, an insulin secretagogue, an insulinotropic sulfonylurea receptor ligand, a thiazolidone derivative, a glycogen synthase kinase-3 inhibitor, a sodium-dependent glucose co-transporter inhibitor, a glycogen phosphorylase A inhibitor, a biguanide, an alpha-glucosidase inhibitor, a glucagon like peptide-1 (GLP-1), a GLP-1 analog, a GLP-1 mimetic, a modulator of a peroxisome proliferator-activated receptor, a dipeptidyl
15 peptidase IV inhibitor, a stearyl-CoA desaturase-1 inhibitor, a diacylglycerol acyltransferase 1 and 2 inhibitor, an acetyl CoA carboxylase 2 inhibitor, a breaker of an advanced glycation end product, a loop diuretic, an angiotensin converting enzyme inhibitor, an inhibitor of the Na-K-ATPase membrane pump, a neutralendopeptidase (NEP) inhibitor, an ACE/NEP inhibitor, an angiotensin II antagonist, a renin inhibitor, a β -adrenergic receptor blocker, an
20 inotropic agent, a calcium channel blocker, an aldosterone receptor antagonist, an aldosterone synthase inhibitor, a 3-hydroxy-3-methyl-glutaryl coenzyme A reductase inhibitor, an HDL increasing compound, a squalene synthase inhibitor, a farnesoid X receptor ligand, a liver X receptor ligand, a cholestyramine, a fibrate, nicotinic acid and aspirinTM.

17. The pharmaceutical composition as claimed in claim 16, wherein the
25 Na-K-ATPase membrane pump is digoxin and the HDL increasing compound is a cholesterol ester transfer protein inhibitor.

54348-1


18. A process for the preparation of a compound of formula (I), as claimed in any one of claims 1 to 6, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, said process comprising:

reacting a diamine of formula (II):




5

with a carboxylic acid of formula (III):




to provide a compound of formula (IV):



10


which is reacted with $\text{R}^3\text{---Hal}$ to obtain the compound of formula (I):



54348-1


19. A process for the preparation of a compound of formula (I), as claimed in claim 1, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, said process comprising:

reacting a nitro pyrimidine dione of formula (V):




5

with an aldehyde of formula (VI):




followed by reductive cyclisation to provide a compound of formula (VII):



10


which is reacted with $\text{R}^3\text{-Hal}$ to obtain compound of formula (I):




54348-1

20. A process for the preparation of a compound of formula (I), as claimed in claim 1, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, said process comprising:


reacting compound of formula (VIII):



with a compound of formula (IX):



to obtain compound of formula (I):




21. A compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, for use in treatment of a condition mediated by an adenosine receptor.

22. Use of a compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, for the treatment of a condition mediated by an adenosine receptor.

15

54348-1

23. Use of a compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, for the treatment of asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, an allergic disease, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, or an autoimmune disease.
24. Use of a compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, in the preparation of medicament for the treatment of a condition mediated by an adenosine receptor.
25. Use of compound of formula (I), as claimed in any one of claims 1 to 7, or a tautomer, a polymorph, a stereoisomer, a prodrug, a solvate or a pharmaceutically acceptable salt thereof, in the preparation of medicament for the treatment of asthma, chronic obstructive pulmonary disorder, angiogenesis, pulmonary fibrosis, emphysema, an allergic disease, inflammation, reperfusion injury, myocardial ischemia, atherosclerosis, hypertension, congestive heart failure, retinopathy, diabetes mellitus, obesity, inflammatory gastrointestinal tract disorders, or an autoimmune disease.



I