



(21) (A1) **2,270,255**
(86) 1997/10/20
(87) 1998/05/07

(72) KEENAN, Robert Daniel, GB

(72) WIX, Loyd, GB

(72) YOUNG, David, GB

(71) UNILEVER PLC, GB

(51) Int.Cl.⁶ A23L 3/015, A23G 9/30, A23G 9/04, A23G 9/02

(30) 1996/10/29 (96307808.4) EP

(54) **PROCEDE DE PREPARATION D'UN PRODUIT ALIMENTAIRE**

(54) **METHOD FOR THE PREPARATION OF A FOODSTUFF**

(57) La présente invention porte sur un procédé de préparation d'une glace comprenant deux phases: une phase aqueuse et une phase corps gras, procédé dans lequel la glace est soumise à un traitement à pression ultra-élevée. Ladite glace contient, au cours de la phase aqueuse, au moins 1 % en poids de caséine micellaire et une quantité de sucre suffisante pour qu'un gel de protéines se forme à l'application de la pression ultra-élevée, étant entendu que si la quantité de caséine micellaire présente dans la phase aqueuse est inférieure à 2 % en poids, la composition de la glace devra inclure un agent stabilisant.

(57) A process for the preparation of an ice confection having an aqueous phase and a fat phase, wherein the ice confection is subjected to an ultra high pressure treatment, the ice confection comprising in the aqueous phase at least 1 wt.% micellar casein and sufficient total sugars such that a protein gel is formed on application of the ultra high pressure, providing that if micellar casein is present in the aqueous phase at less than 2 wt.%, the ice confection composition includes a stabiliser.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A23L 3/015, A23G 9/30, 9/04, 9/02	A1	(11) International Publication Number: WO 98/18350 (43) International Publication Date: 7 May 1998 (07.05.98)
(21) International Application Number: PCT/EP97/05917 (22) International Filing Date: 20 October 1997 (20.10.97) (30) Priority Data: 96307808.4 29 October 1996 (29.10.96) EP <i>(34) Countries for which the regional or international application was filed:</i> GB et al. (71) Applicant (for AU BB CA GB GH IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AU BB CA GB GH IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). (72) Inventors: KEENAN, Robert, Daniel; 29 Betony Walk, Rushden, Northampton NN10 0TL (GB). WIX, Loyd; 93 Hayway, Rushden, Northampton NN10 6AQ (GB). YOUNG, David; 13 Shurville Close, Earls Barton, Northampton NN6 0JB (GB).		(74) Agent: KIRSCH, Susan, Edith; Unilever plc, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHOD FOR THE PREPARATION OF A FOODSTUFF (57) Abstract A process for the preparation of an ice confection having an aqueous phase and a fat phase, wherein the ice confection is subjected to an ultra high pressure treatment, the ice confection comprising in the aqueous phase at least 1 wt.% micellar casein and sufficient total sugars such that a protein gel is formed on application of the ultra high pressure, providing that if micellar casein is present in the aqueous phase at less than 2 wt.%, the ice confection composition includes a stabiliser.		

Method for the Preparation of a Foodstuff**Technical Field of the Invention**

5 The invention relates to a method of preparation of an ice confection such as ice cream, wherein the ice confection is subjected to an ultra high pressure treatment.

Background to the Invention

10

Ultra high pressure (UHP) is a known method for killing spores and has been suggested as a suitable route to food product pasteurisation. In Japan a range of pressure de-contaminated products such as jellies, preserves, purees and sauces have been launched on the market (Byrne, M. (1993) Food Engineering International, 34-38).

15

Furthermore isolated, native proteins have been subjected to UHP. These proteins are in their native form, they have not been treated chemically or thermally before the pressure treatment by methods which significantly modify their protein structure (van Camp, J; Huyghebaert, A (1995) Food Chemistry 54(4) 357-364; Okamoto, M; Kawamura, Y; Hayashi, R; (1990) Agric Biol Chem 54(1) 183-189). It is generally believed that there would be no advantage in subjecting proteins which have already been substantially denatured by for example an initial heat-treatment prior to UHP.

20

25

30 DE 42 26 255 discloses the treatment of cream with ultra high pressure in order to crystallise the fat.

35

Surprisingly, we have noted that when an ice confection having an aqueous phase and a fat phase, such as ice cream, frozen yoghurt and the like, is subjected to UHP a number of advantages are achieved, providing that the micellar casein protein content and the total sugar content present

in the aqueous phase of the ice confection are within specified limits. Further these advantages are achieved even if the product has been heat treated prior to UHP treatment.

5

Disclosure of the Invention

Accordingly the invention provides a process for the preparation of an ice confection having an aqueous phase and a fat phase, wherein the ice confection is subjected to an ultra high pressure treatment, the ice confection comprising in the aqueous phase at least 1 wt% micellar casein and sufficient total sugars such that a protein gel is formed on application of the ultra high pressure, providing that if micellar casein is present in the aqueous phase at less than 2 wt%, the ice confection composition includes a stabiliser.

Preferably the aqueous phase comprises at least 2 wt% micellar casein.

By the term micellar casein it is meant casein protein present in the form of supramolecular structures, consisting of aggregated proteins and minerals, dispersed in the aqueous phase of the ice confection. These structures could have been formed naturally, ie native casein micelles as are present in milk, or may have been formed synthetically, such as those produced by the addition of certain minerals (eg Ca^{2+}) to casein.

30

Ice confections such as ice cream are usually produced by a continuous process comprising the following steps:

- (a) homogenisation of the ingredients
- (b) pasteurisation
- (c) cooling
- (d) freezing and aeration

35

- (e) extrusion; and
- (f) optionally deep freezing.

5 The applicants have surprisingly found that if the ice confection is subjected to an ultra high pressure treatment after step (b) and prior to step (d) the resulting product has a number of advantages as follows;

- 10 (i) the meltdown rate is decreased
- (ii) the product provided is perceived to be thicker, smoother and creamier and is less cold to eat;

15 Furthermore a particular advantage of such products is that they may be prepared in the absence of emulsifiers and/or stabilisers and/or have a zero or low fat content and/or a low milk content and yet retain high quality. With the proviso that at micellar casein levels of less than 2 wt% in the aqueous phase, stabiliser must be present.

20 Usually the protein content of the product will have been de-natured to some extent by the processing of the product prior to the application of the ultra high pressure treatment.

25 Optionally the conventional pasteurisation step (step b) may be omitted for ice confections which are subjected to the ultra high pressure treatment of the invention. However it is preferred that a conventional pasteurisation step is
30 conducted prior to the ultra high pressure treatment.

The pressure during the ultra high pressure treatment is typically greater than or equal to 250MPa, preferably approximately 400MPa. The foodstuff is subjected to this
35 pressure for typically greater than or equal to 1 minute, preferably from 1 to 80 minutes, most preferably from 5 to 40 minutes.

The ultra high pressure may be applied using any suitable apparatus. The process may be a batch process, a semi-continuous process, or preferably a continuous process.

5 The improved meltdown stability and sensory attributes of ice confections made from UHP treated premixes are believed to result from the formation of UHP-induced protein gels. In order to achieve the required protein gel on the application of the UHP a minimum content of total sugars
10 has been found to be required within the aqueous phase of the ice confection. This can be better illustrated in Example 8 below where a state diagram is provided for systems treated at an ultra high pressure of 400 MPa for 20 minutes. Essentially the lower the micellar casein content
15 in the aqueous phase the higher the total sugar content in the aqueous phase has to be. Thus at a micellar casein content in the aqueous phase of 2 wt% a total sugar content of at least 30 wt% is required. At a micellar casein content in the aqueous phase of 3 wt% a total sugar content
20 of at least 20 wt% is required. At a micellar casein content in the aqueous phase of 4 wt% a total sugar content of at least 15 wt% is required. At a micellar casein content in the aqueous phase of 5 wt% a total sugar content of at least 13 wt% is required.

25 Preferably the micellar casein content in the aqueous phase is from 3 to 5 wt%. Preferably the total sugar content in the aqueous phase is approximately 25 wt%.

30 The sugar content can be selected from monosaccharides, disaccharides, oligosaccharides and mixtures thereof. Suitable examples include sucrose, fructose, lactose, glucose and corn syrups.

35 D G Schmidt and W Buchheim in *Milchwissenschaft* 25, 596 (1970) showed that pressure causes the irreversible fragmentation of casein micelles in milk, probably through

pressure-induced dissolution of micellar calcium phosphate. However, we now believe that if this pressure induced fragmentation occurs in the presence of certain cosolutes (sugars) a network is formed from these fragments. This re-aggregation of the casein fragments may occur on decompression of the milk/sugar mixture. Examination of pressure-treated milk/sugar mixtures by transmission electron microscopy reveals that the gels are made from a particle network, the size of the individual particles being approximately an order of magnitude smaller than the casein micelles present in untreated mixtures. Therefore pressure treatment in the presence of appropriate levels of sugars provides a method of gelling casein at natural pH, room temperature and in the absence of coagulating enzymes.

Examples

Examples 1 to 3

An ice cream mixture having the following formulation

12.1% skimmed milk powder
25% dairy cream
15.44% sucrose
0.4% vanilla flavour
water to 100%

was prepared and pasteurised in the conventional way. The product was then subjected to a high pressure treatment as follows

Example 1 - 300 MPa for 40 min
Example 2 - 400 MPa for 5 min
Example 3 - 400 MPa for 40 min

at ambient temperature in a National Forge Cold Isostatic Press. The products were then frozen in the conventional

way in a standard ice cream freezer (scraped surface heat exchanger SSHE), extruded and deep frozen.

5 The percentage mass loss was determined at 37°C after at least 2 weeks frozen storage by measuring the weight of melted ice cream every minute over the required time period which passed through a grid having 3mm square holes and a 1mm mesh. The mass loss after 2 hours at 37°C was as follows;

10

Example 1 - 3%

Example 2 - 4%

Example 3 - 3%

15 Figure 1 shows the results of the percentage mass loss measurements. The mass loss of the ice cream composition prepared in Example 3 is shown by ----. This is compared to a control (—) which is the same ice cream composition prepared as in Example 3 except that the ice cream
20 composition is not subjected to an ultra high pressure treatment.

The percentage destabilised fat was measured using a solvent extraction technique. 10g of ice cream was melted
25 for 4 hours at ambient temperature before extraction with petroleum solvent. The solvent was evaporated and the extracted destabilised fat was weighed, this was expressed as a percent of the weight of the total fat in the ice cream.

30

Accordingly the percentage destabilised fat was measured to be 0.3%.

35 These results show that the gel has not been formed by a fat network.

Further the prepared ice cream of Example 3 was tested by

a trained sensory panel and found to be firmer, chewier, smoother, less icy, creamier, thicker and less cold than Comparative Example A.

5 Comparative Example A

An ice cream was prepared as in Example 1 above except that the ice cream pre-mix was not subjected to any UHP treatment.

10

Mass loss was measured as in Example 1. The mass loss after 2 hours at 37°C was 88%.

Example 4

15

Example 3 was repeated except that after pasteurisation the product was subjected to a heat treatment at 140°C for 5 seconds before the high pressure treatment.

20

Mass loss was measured as in Example 1. The mass loss after 2 hours at 37°C was 41%.

Example 5

25

An ice cream mixture having the following formulation

14.4% skimmed milk powder

0.5% butter fat

17.6% sucrose

30

0.4% vanilla flavour

water to 100%

35

was prepared and pasteurised in the conventional way. The product was then subjected to a high pressure treatment at 400 MPa for 40 min at ambient temperature in a National Forge Cold Isostatic Press. The product was then frozen in the conventional way in a standard ice cream freezer

WO 98/18350

PCT/EP97/05917

8

(scraped surface heat exchanger SSHE), extruded and deep frozen.

5 Mass loss was measured as in Example 1. The mass loss after 2 hours at 37°C was 43%.

10 Further the prepared ice cream was tested by a trained sensory panel and found to be firmer, chewier, smoother, less icy, creamier and thicker than Comparative Example B.

Comparative Example B

15 An ice cream was prepared as in Example 5 above except that the ice cream pre-mix was not subjected to any UHP treatment.

Mass loss was measured as in Example 1. The mass loss after 2 hours at 37°C was 98%.

20 Example 6

An ice cream mixture having the following formulation

25 9.1% skimmed milk powder
25% dairy cream
15.44% sucrose
0.4% vanilla flavour
water to 100%

30 was prepared and pasteurised in the conventional way. The product was then subjected to a high pressure treatment at 400 MPa for 40 min at ambient temperature in a National Forge Cold Isostatic Press. The product was then frozen in the conventional way in a standard ice cream freezer
35 (scraped surface heat exchanger SSHE), extruded and deep frozen.

Mass loss was measured as in Example 1. The mass loss after 2 hours at 37°C was 12%.

5 Further the prepared ice cream was tested by a trained sensory panel and found to be less icy, creamier and thicker than Comparative Example C.

Comparative Example C

10 An ice cream was prepared as in Example 6 above except that the ice cream pre-mix was not subjected to any UHP treatment.

15 Mass loss was measured as in Example 1. The mass loss after 2 hours at 37°C was 94%.

Example 7

An ice cream mixture having the following formulation

5 10% skimmed milk powder
 8% butteroil
 13% sucrose
 4% corn syrup
 0.144% Locust Bean Gum
10 0.016% Carageenan
 0.3% mono/di glyceride
 0.012% vanillin
 water to 100%

15 was prepared and pasteurised in the conventional way. The
 product was then subjected to a high pressure treatment at
 400 MPa for 40 min at ambient temperature in a National
 Forge Cold Isostatic Press. The product was then frozen in
20 the conventional way in a standard ice cream freezer
 (scraped surface heat exchanger SSHE), extruded and deep
 frozen.

 Mass loss was measured as in Example 1. The mass loss after
 2 hours at 37°C was 8%.

25 Further the prepared ice cream was tested by a trained
 sensory panel and found to be firmer and creamier than
 Comparative Example D.

30 Comparative Example D

 An ice cream was prepared as in Example 7 above except that
 the ice cream pre-mix was not subjected to any UHP
 treatment.

35 Mass loss was measured as in Example 1. The mass loss after
 2 hours at 37°C was 52%.

Example 8

5 A series of samples containing different amounts of sugars and micellar casein were prepared by dispersing skim milk powder (SMP), sucrose and/or lactose into water at room temperature. These samples had levels of micellar casein in the range 0-12% (w/w) and sugars in the range 5-50%. The final concentrations were calculated assuming that SMP contains approximately 50% lactose and 30% micellar casein.

10

Samples were sealed in 5 ml plastic bags and compressed at 400 MPa for 20 min using a cold isostatic press (Stansted Fluid Power, Stansted, UK) operating at room temperature. The press had a sample volume of 30 ml and the pressure-transmitting fluid was methanol. After decompression samples were stored at 5°C for 3 days prior to rheometric analysis.

15

In order to determine which samples had been gelled by the high pressure treatment, oscillatory rheometry was performed on both the pressure-treated samples and untreated controls. The rheological tests were performed at 5°C on a Carrimed CSL500 rheometer operating in oscillatory mode with a cone-and-plate geometry. A 6 cm diameter acrylic cone with a gap of 53 μm , operating at a stress of 1 Pa and a strain of 0.005, was used. Samples were considered to be gelled if the measured storage modulus (G') was greater than the loss modulus (G'') at a frequency of 1 Hz.

20

25

30

In all cases the untreated samples were liquids. Figure 2 shows the state diagram for systems treated at 400 MPa for 20 min. Samples with compositions above the line were found to be gelled (according to the criteria given above) while those below the line were liquids after pressure treatment.

35

Claims

- 5 1. A process for the preparation of an ice confection having an aqueous phase and a fat phase, wherein the ice confection is subjected to an ultra high pressure treatment, the ice confection comprising in the aqueous phase at least 1 wt% micellar casein and sufficient total sugars such that a protein gel is formed on application of
10 the ultra high pressure, providing that if micellar casein is present in the aqueous phase at less than 2 wt%, the ice confection composition includes a stabiliser.
- 15 2. A process according to claim 1 wherein the ice confection comprises at least 2 wt% micellar casein in the aqueous phase.
- 20 3. A process according to claim 1 wherein the ice confection comprises from 3 to 5 wt% micellar casein in the aqueous phase.
4. A process according to claim 3 wherein the ice confection comprises approximately 25 wt% total sugars.
- 25 5. A process according to claim 1 wherein the aqueous phase of the ice confection comprises 2 wt% micellar casein and at least 30 wt% total sugars.
- 30 6. A process according to claim 1 wherein the aqueous phase of the ice confection comprises 3 wt% micellar casein and at least 20 wt% total sugars.
- 35 7. A process according to claim 1 wherein the aqueous phase of the ice confection comprises 4 wt% micellar casein and at least 15 wt% total sugars.
8. A process according to claim 1 wherein the aqueous

13

phase of the ice confection comprises 5 wt% micellar casein and at least 13 wt% total sugars.

5 9. A process according to any preceding claim wherein the pressure during the ultra high pressure treatment is greater than or equal to 250MPa.

10 10. A process according to any preceding claim wherein the pressure during the ultra high pressure treatment is approximately 400MPa.

15 11. A process according to any preceding claim wherein the foodstuff is subjected to ultra high pressure for greater than or equal to 1 minute.

12. A process according to any preceding claim wherein the foodstuff is subjected to ultra high pressure for from 1 to 80 minutes.

20 13. A process according to any preceding claim wherein the foodstuff is subjected to ultra high pressure for from 5 to 40 minutes.

25 14. A process according to any one of claims 2 to 13 wherein the ice confection contains no added stabilisers.

15. A process according to any preceding claim wherein the ice confection contains no added emulsifiers.

30 16. A process according to any preceding claim wherein the ice confection has a zero or low fat content.

Fig.1.




Fig.2.

