



(86) **Date de dépôt PCT/PCT Filing Date:** 2010/07/13
(87) **Date publication PCT/PCT Publication Date:** 2011/06/30
(45) **Date de délivrance/Issue Date:** 2016/08/09
(85) **Entrée phase nationale/National Entry:** 2012/06/22
(86) **N° demande PCT/PCT Application No.:** US 2010/041848
(87) **N° publication PCT/PCT Publication No.:** 2011/078891
(30) **Priorité/Priority:** 2009/12/22 (US61/289,295)

(51) **Cl.Int./Int.Cl.** **A23K 20/20** (2016.01),
A23K 50/00 (2016.01), **A23K 50/10** (2016.01),
C05D 9/00 (2006.01), **C05F 11/00** (2006.01),
C05G 3/00 (2006.01)


(72) **Inventeurs/Inventors:**
KNOCHENMUS, BRIAN JON, US;
KNOCHENMUS, JON KENT, US;
LAMB, RICHARD DALE, US;
LAMB, MYRRA ARLENE, US

(73) **Propriétaire/Owner:**
RALCO NUTRITION, INC., US

(74) **Agent:** GOWLING WLG (CANADA) LLP


(54) **Titre : COMPOSITIONS MINERALES A CHELATION ORGANIQUE ET PROCEDES ASSOCIES**

(54) **Title: ORGANICALLY CHELATED MINERAL COMPOSITIONS AND METHODS THEREOF**



(57) **Abrégé/Abstract:**

Embodiments of the invention relate to a method of making a mineral product. The method includes contacting a carboxylic acid and an inorganic mineral compound sufficient to form a solution, reacting the solution over a period of time sufficient to provide a



(57) Abrégé(suite)/Abstract(continued):

mineral chelated compound, transferring the mineral chelated compound to one or more molds prior to the compound substantially solidifying and reducing the size of the mineral chelated compound sufficient to provide a rapidly soluble mineral chelated product.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
30 June 2011 (30.06.2011)(10) International Publication Number
WO 2011/078891 A1(51) International Patent Classification:
A23K 1/175 (2006.01) *A23K 1/18* (2006.01)Marshall, Minnesota 56258 (US). **LAMB, Myrra Arlene** [US/US]; 1107 DeVito Street, Marshall, Minnesota 56258 (US).(21) International Application Number:
PCT/US2010/041848
(22) International Filing Date:
13 July 2010 (13.07.2010)(74) Agents: **ARMITAGE, Benjamin C.** et al.; 605 US Highway 169 North, Suite 300, Plymouth, Minnesota 55441 (US).

(25) Filing Language: English

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
61/289,295 22 December 2009 (22.12.2009) US(71) Applicant (for all designated States except US): **RALCO NUTRITION, INC.** [US/US]; 1600 Hahn Road, Marshall, Minnesota 56258 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KNOCHENMUS, Brian Jon** [US/US]; 2196 190th Ave, Lynd, Minnesota 56157 (US). **KNOCHENMUS, Jon Kent** [US/US]; 101 Savannah Heights Blvd, Lynd, Minnesota 56157 (US). **LAMB, Richard Dale** [US/US]; 1107 DeVito Street,

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,

[Continued on next page]

(54) Title: ORGANICALLY CHELATED MINERAL COMPOSITIONS AND METHODS THEREOF

(57) Abstract: Embodiments of the invention relate to a method of making a mineral product. The method includes contacting a carboxylic acid and an inorganic mineral compound sufficient to form a solution, reacting the solution over a period of time sufficient to provide a mineral chelated compound, transferring the mineral chelated compound to one or more molds prior to the compound substantially solidifying and reducing the size of the mineral chelated compound sufficient to provide a rapidly soluble mineral chelated product.




FIG. 1

WO 2011/078891 A1

WO 2011/078891 A1



EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, **Published:**
LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, — *with international search report (Art. 21(3))*
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

ORGANICALLY CHELATED MINERAL COMPOSITIONS AND METHODS THEREOF

[0001] BACKGROUND

[0002] Cobalt and other inorganic minerals are useful to mammals in many ways. One important need for cobalt in animal nutrition is in regard to vitamin B12 production. Microorganisms require cobalt to produce vitamin B12. Digestive bacteria have a very strong affinity for cobalt. When a dissociable cobalt source is introduced into the rumen, for example, the uptake by bacteria is rapid- about 80-85% within 30-40 minutes. The plant form of cobalt supplied by ration ingredients is released with a considerable degree of variability. So, matching the form provided in a supplement to the metabolic demands by the rumen microorganisms and the animals' physiological needs is key to maximizing animal performance.

[0003] Vitamin B12 levels in an animal are directly proportional to the level of available cobalt supplied to the rumen. Vitamin B12 levels in the plasma and liver directly impact the ability to convert propionic acid to glucose and the ability to synthesize methionine. Products on the market to supply cobalt to animals or other living organisms are often in forms less available to the animal or organism than desirable, causing unnecessary waste and inefficiencies. Conventional processes of making such products are inefficient and impractical to provide a soluble form. There is a marked difference in solubility in water of cobalt lactate versus cobalt carbonate, for example.

[0004] Cobalt, transition metals and trace minerals are useful to microorganisms directly for metabolism, immune function and reproduction. Microorganisms in the soil that fix nitrogen to enhance plant growth are called Azotobacters. The form of the trace minerals added to soil or to other systems employing microorganisms, such as waste treatment systems, biofilters, anaerobic

digestors and the like, impacts the availability and therefore uptake of the trace element.

BRIEF DESCRIPTION OF THE DRAWINGS

[0005] In the drawings, which are not necessarily drawn to scale, like numerals describe substantially similar components throughout the several views. Like numerals having different letter suffixes represent different instances of substantially similar components. The drawings illustrate generally, by way of example, but not by way of limitation, various embodiments discussed in the present document.

[0006] FIG. 1 illustrates a block flow diagram of a method of making a rapidly soluble mineral chelated product, according to some embodiments.

[0007] FIG. 2 illustrates a block flow diagram of a method of applying a mineral lactate compound, according to some embodiment.

[0008] FIG. 3 illustrates a block flow diagram of a method of administering a mineral lactate compound, according to some embodiment.

SUMMARY

[0009] Embodiments of the invention relate to a method of making a mineral product. The method includes contacting a carboxylic acid and an inorganic mineral compound sufficient to form a solution, reacting the solution over a period of time sufficient to provide a mineral chelated compound, transferring the mineral chelated compound to one or more molds prior to the compound substantially solidifying and reducing the size of the mineral chelated compound sufficient to provide a rapidly soluble mineral chelated product.

DETAILED DESCRIPTION

[0010] The following detailed description includes references to the accompanying drawings, which form a part of the detailed description. The drawings show, by way of illustration, specific embodiments in which the invention may be practiced. These embodiments, which are also referred to herein as “examples,” are described in enough detail to enable those skilled in the art to practice the invention. The embodiments may be combined, other embodiments may be utilized, or

WO 2011/078891

PCT/US2010/041848

structural, and logical changes may be made without departing from the scope of the present invention. The following detailed description is, therefore, not to be taken in a limiting sense, and the scope of the present invention is defined by the appended claims and their equivalents.

[0011] In this document, the terms “a” or “an” are used to include one or more than one and the term “or” is used to refer to a nonexclusive “or” unless otherwise indicated. In addition, it is to be understood that the phraseology or terminology employed herein, and not otherwise defined, is for the purpose of description only and not of limitation.

[0012] Embodiments of the invention relate to inorganic mineral chelated compositions and methods of making and using such compositions. Embodiments describe a unique process including a reaction to form an organically bound cobalt or other mineral, optionally carrying the reacted material in an inert media and applying or administering the product. The process is substantially stoichiometric, in which there are little to no by-products that would otherwise need to be removed from the product. The process is very efficient, cost-effective and provides a readily available form of inorganic material for fertilization or nutrition or the benefit of microflora in a number of applications. The composition produced may be in a liquid form that is easily applied to plants, pastures, lawns or gardens or administering via a nutritional product or feed to animals or humans. Embodiments of the present invention utilize a carboxylic acid, such as lactic acid, to provide an inorganic mineral chelated composition that is easily administered or applied in any form or supplementation, liquid nutrition, agricultural use or industrial use.

[0013] In supplementation, the composition has a greater affinity to digestive microbials than other sources or structures of inorganic minerals. In ruminants, for example, the composition provides faster weight gain, heavier weaning weights and enhances rumen fermentation utilization. The composition increases cellulose breakdown and utilization and helps digest forage better. Additionally, the animals

utilize protein and calcium better, providing a decreased need for commercial proteins.

[0014] In addition to administering the chelated compound to mammals or other living organisms as part of feed or feed supplement, methods of applying the product for agricultural purposes are described. It is possible to transfer nutrients to a mammal through treatment of the soil or plants on which it grazes. This may be more cost effective and potentially more efficacious to the animal. Additionally, chelated trace minerals such as cobalt positively impact the health and growth of microorganisms, thus having a beneficial impact on the microorganisms' environment.

Definitions

[0015] As used herein, "fatty acid" refers to a carboxylic acid, often with a long unbranched aliphatic tail (chain), which may be either saturated or unsaturated. Examples of fatty acids may include lactic acid, propionic acid and butyric acid.

[0016] As used herein, "carboxylic acid" refers to organic acids characterized by the presence of a carboxyl group, which has the formula $-C(=O)OH$, usually written $-COOH$ or $-CO_2H$. Examples of carboxylic acids may include lactic acid, propionic acid and butyric acid.

[0017] As used herein "microflora" refers to living microorganisms that are so small that they can be seen only with a microscope and that maintain a more or less constant presence in a particular area, e.g. the pharynx or the rumen. Microflora includes bacteria, viruses, protozoa and fungi for example.

[0018] As used herein, "lactic acid" refers to a carboxylic acid with a chemical formula of $C_3H_6O_3$.

[0019] As used herein, "inorganic mineral compound" refers to an elemental or compound composition including one or more inorganic species. For example, an inorganic mineral compound may be cobalt, cobalt carbonate, zinc oxide, cupric oxide, manganese oxide or a combination thereof. Inorganic mineral compounds may include scandium, selenium, titanium, vanadium, chromium, manganese, iron, nickel, copper and zinc, for example. Transition metals may also be included and salts, oxides, hydroxides and carbonates of the above mentioned compounds may be used.

[0020] As used herein, “solution” refers to a homogeneous or substantially homogeneous mixture of two or more substances, which may be solids, liquids, gases or a combination thereof.

[0021] As used herein, “reacting” refers to undergoing a chemical change. Reacting may include a change or transformation in which a substance decomposes, combines with other substances, or interchanges constituents with other substances

[0022] As used herein, “mineral chelated compound” refers to chemical compound or mixture including at least one inorganic substance and a derivative of a carboxylic acid or reaction product of a carboxylic acid and an inorganic mineral compound.

[0023] As used herein, “transferring” refers to moving a component or substance from one place or location to another.

[0024] As used herein, “mold” refers to a hollow form or matrix for shaping a fluid, gel, semi-solid or plastic substance.

[0025] As used herein, “filtering” or “filtration” refers to a mechanical method to separate solids from liquids, or separate components by size or shape. This can be accomplished by gravity, pressure or vacuum (suction).

[0026] As used herein, “rapidly soluble mineral chelated product” refers to a mineral chelated compound that has been altered to increase solubility in a solvent. Altering may include reducing in size, filtering, screening or chemically reacting. An inorganic mineral compound may be organically chelated such that its solubility changes from insoluble to soluble in a chose solvent.

[0027] As used herein, “screening” refers to separating components by size by passing or refusing to pass components through a screen or mesh.

[0028] As used herein, “reducing in size” refers to physically or chemically reducing the size of one or more components, such as by grinding, crushing or milling, for example.

[0029] As used herein, “mesh size” refers to the number of openings in one inch of a screen or filter.

[0030] As used herein, “contacting” may refer to physically, chemically, electrically touching or bringing within sufficient close proximity.

[0031] As used herein, “carrier” refers to a substance physically or chemically bound with a target or active substance in order to facilitate the use or application of the target or active substance.

[0032] As used herein, “substrate” refers to a base layer or material on which an active or target material interacts with, is applied to or is carried by.

[0033] As used herein, “stoichiometric” or “stoichiometric amounts” refer to starting materials of a reaction having molar amounts or substantially molar amounts such that the reaction product is formed with little to no unused starting material or waste. A stoichiometric reaction is one in which all starting materials are consumed (or substantially consumed) and converted to a reaction product or products.

[0034] As used herein, “applying” refers to bringing one or more components into nearness or contact with another component.

[0035] As used herein, “administering” refers to giving or applying. More specifically, administering refers to providing a substance such a mammal ingests the substance, such as through feeding or medication.

[0036] As used herein, “feed” refers to food for animals, fish, reptiles, microflora, insects, birds or any living organism.

[0037] As used herein, “feed supplement” refers to something added to a feed or food to remedy a deficiency, strengthen or increase the value of the feed or food.

[0038] As used herein, “mixture” refers to a combination of two or more substances in physical contact with one another. For example, components of a mixture may be physically combined as opposed to chemically reacting.

[0039] Referring to FIG. 1 a block flow diagram 100 of a method of making a rapidly soluble mineral chelated product is shown, according to some embodiments. A carboxylic acid 102, such as lactic acid, may be contacted 106 with an inorganic mineral compound 104, sufficient to form a solution 108. The solution 108 may be reacted 110 over a period of time, sufficient to provide a mineral chelated compound 112. The mineral chelated compound 112 may then be transferred and reduced in size 114 to sufficient to provide a rapidly soluble mineral chelated product 116. Transferring may include transferring to one or more molds, prior to the compound substantially solidifying.

[0040] Carboxylic acid 102 may be contacted 106 with an inorganic mineral compound 104, such as by mixing. The molar amounts or stoichiometric amounts may be utilized. If the carboxylic acid 102 is lactic acid, the carboxylic acid content may be about 60% to about 80% of the mixture by weight. The inorganic mineral compound 104 may include about 20% to about 40% of the mixture by weight. More specifically, the lactic acid may include about 62% to about 76% and the inorganic mineral compound 104 may include about 24% to about 38% by weight of the mixture. The lactic acid 102 may be 88% strength lactic acid, for example.

[0041] When the carboxylic acid 102 is propionic acid, the carboxylic acid content may be about 55% to about 75% by weight and the inorganic mineral compound content about 25% to about 45% by weight. More specifically, the propionic acid may include about 57% to about 72% and the inorganic mineral compound 104 may include about 28% to about 43% by weight. When the carboxylic acid 102 is butyric acid, the carboxylic acid content may be about 60% to about 80% by weight and the inorganic mineral compound content about 20% to about 40% by weight. More specifically, the butyric acid may include about 61% to about 76% and the inorganic mineral compound 104 may include about 24% to about 39% by weight.


[0042] The carboxylic acid 102 and inorganic mineral compound 104 may be placed in a vessel, optionally with one or more catalysts. Examples of a catalyst include iron and alkaline earth metals. The vessel may be optionally agitated, such as by vibrating, shaking, turning or spinning. Water may be added to the vessel, before, during or after the contacting 106 of carboxylic acid 102 and inorganic mineral compound 104. Once a solution 108 is formed, it may be reacted 110 over a period of time. The reaction may initiate based solely on the contact 106 between carboxylic acid 102 and inorganic mineral compound 104, after addition or contact with a catalyst or similarly with the contact or addition of water of some combination thereof. Depending on the type of inorganic mineral compound utilized 104, carbon dioxide may be evolved as the solution 108 heats up. Both water vapor and optionally carbon dioxide may be generated and released from the vessel. No reflux process is needed or desired, as often used conventionally with regard to related reactions. All by-products may be passively and naturally removed, without the need for solvent or refluxing. Carbon dioxide and water may be released into the atmosphere, for example.

[0043] The reaction ultimately produces a mineral chelated compound 112. The mineral chelated compound 112 may form a porous, brittle rock if left to solidify. The mineral chelated compound 112 may then be transferred from the vessel to one or more molds, prior to the compound substantially solidifying. The molds may be of varying shapes or sizes, such that the compound may be easily handled and transported. Water vapor may be further driven off the compound as it solidifies within the one or more molds.

[0044] The mineral chelated compound 112 may be reduced in size 114. The compound 112 may be removed from the molds and placed in a “de-lumper” or single or double shaft disintegrator or crusher, which may reduce the size of the compound to small particles. The particles may be about 1 to about 2 inches in size, for example. The small particles may then be further reduced in size 114, such as by being contacted with a mill (i.e., hammer mill or roller mill). The small particles may then be reduced to a fine powder. Reducing the compound 112 to a fine powder may increase its solubility, providing a rapidly soluble mineral chelated product 116. After contacting with a mill, the particles may be screened to further separate larger particles from smaller ones. Any larger particles may be placed back in the mill for further reduction in size. Screening may include filtering with a mesh. The mesh size may be about 50 to about 70 or about 50, about 60 or about 70 size mesh. The mesh size may less than 50 for example.

[0045] The rapidly soluble mineral chelated product 116 may be further contacted with a carrier. The carrier may be a dry substrate or a liquid carrier, for example. The carrier may include one or more of diatomaceous earth, calcium carbonate, limestone, sugars, dextrose, water, ground corn cobs, starch and combinations thereof.

[0046] One example of the rapidly soluble mineral chelated product 116 may be organically chelated cobalt, with the following chemical formula:



[0047] Referring to FIG. 2, a block flow diagram 200 of a method of applying a mineral chelated compound is shown, according to some embodiment. A mineral chelated compound 202 may be applied 204 to a lawn, garden, pasture or field 206, for example. Alternatively, the compound 202 may be administered 302 to one or

more living organisms 304 (see view 300 of FIG. 3). The compound 202 may further include a carrier.

[0048] The compound 202 may include one or more of a cobalt lactate compound, zinc lactate compound, copper lactate compound or manganese lactate compound. The carrier may include diatomaceous earth. Applying 204 may include spraying, planting in a seed mix, planting in a fertilizer mix, or a combination thereof. Lawn, garden, pasture or field 206 may include sporting fields and golf courses, for example. Pasture or field may include a harvested field, bailed field, or field or pasture with crops cut. Applying 204 may also include applying while the crop is harvested or after the crop is harvested. Applying 204 may increase growth in a plant. Applying 204 may include applying the compound 202 in an amount between about 1 to about 100ppm or between about 1 to about 1000ppm, for example. The compound 202 may also be used as a benefit to any microflora, enzyme or biological industrial product, for example. The compound 202 may be used industrially or in human nutrition. Applying 204 may strengthen a root system of a plant.

[0049] The one or more living organisms may include mammals, such as monogastric or ruminant mammals. Administering 204 may include providing the product as a feed or feed supplement. Alternatively, administering 204 may include providing the product to the mammals through ingestion of a plant containing the product. Administering 304 may increase rumen activity in a mammal, for example. Increasing rumen activity may include increasing metabolism.

[0050] The mineral product discussed in embodiments of the current invention may include one or more mineral chelated lactates in addition to other components. The mineral product may include one or more metal sulfates, such as sulfates of manganese, zinc, copper or combinations thereof. The one or more mineral chelated lactates may be a cobalt lactate compound, zinc lactate compound, copper lactate compound or manganese lactate compound. A carrier may be utilized, such as dextrose. Additional components may include fibers, yucca and one or more enzymes.

[0051] The one or more mineral chelated lactates may be present in an amount of about 15% to about 20% of the product by weight. The one or more metal sulfates may be present in an amount of about 2% to about 10% of the product by weight. The fiber may be present in an amount of about 1% to about 5% of the product by

weight. The enzymes may include about 0.1% to about 2% by weight, the yucca about 1% to about 5% by weight and the carrier about 60% to about 80% by weight.

Example 1

[0052] In Table 1, the growth of grass when cobalt lactate was applied either in the soil (before seeding) or sprayed directly onto the grass once growing is shown for the various levels of cobalt (0, 1, 10, 100 and 1000 ppm) that was applied. Following growth 2, cuttings were made then the total weight of grass was shown for the various treatments. Growth roughly doubled when cobalt lactate was applied in the soil at 1 or 10 ppm or applied by spray at 100 ppm. All work was done with a fully complete soil with respect to nitrogen, phosphorus and potash (Miracle Gro Soil), so that the only difference in the greenhouse flats was the applied cobalt level and the manner in which it was applied. The uptake of cobalt correlated with the application rate and that was mirrored closely at the lower levels.

Table 1

Sample	Cobalt Medium	Soil Weight (gms)	Seed Weight (gms)	Target Cobalt Level (ppm)	Grass Cuttings Weight (gms, 1st)	Grass Cuttings Weight (gms, 2nd)	Sum Grass Cuttings Weight (gms, 1st + 2nd)
1	Soil	2550	2.6	0	19.94	3.31	23.25
2	Soil	2550	2.6	1	31.06	22.73	53.79
3	Soil	2550	2.6	10	22.8	31.41	54.21
4	Soil	2550	2.6	100	16.33	10.12	26.45
5	Soil	2550	2.6	1000	5.15	1.81	6.96
6	DI H ₂ O	2550	2.6	0	17.28	8.28	25.56
7	DI H ₂ O	2550	2.6	1	15.7	13.9	29.6
8	DI H ₂ O	2550	2.6	10	17.76	14.37	32.13
9	DI H ₂ O	2550	2.6	100	18.56	26.57	45.13
10	DI H ₂ O	2550	2.6	1000	24.03	15.44	39.47

Example 2

[0053] Sixty-four sample flats were analyzed. Empty flats were filled with about 0.25 cubic feet of Miracle Gro Soil per flat (a complete soil with enough macro-

nutrients, nitrogen, phosphorus and potash to feed the plants for 3-4 months). The seeds were planted and then the seeded soil treated. Flats were watered daily. Progress was documented weekly. This number of flats resulted from running a number of positive controls and several levels of the chelated cobalt lactate per plant selection or choice.

[0054] The makeup of 64 flats was 8 sections times 8 flats per section. The first 8 flats were planted with a mixture of clovers. Flats 9-16 were also planted with the clover mixture at 5.2 grams seed/flat versus 2.6 grams seed/flat in 1-8. Flats 17-24 were planted with Fast & Fine Seed at 2.6 grams/flat, equivalent to the seeding rate from 2008. Fast & Fine Premium Grass Seed is a commercial mixture of Indy Perennial Ryegrass, Silver Dollar Perennial Ryegrass, Boreal Creeping Red Fescue, Kelly Kentucky Bluegrass and Clearwater Kentucky Bluegrass. Flats 25-32 were planted with a mixture including green globe turnips, dwarf Essex forage rapeseed, Rangi forage and giant rapeseed. Flats 33-40 were planted with a mixture of seeds including hybrid sunflower, grain sorghum, Proso millet and buck-wheat, at 5.2 grams/flat. Flats 41-48 were planted with Vernal alfalfa at 5.2 grams/flat and 49-56 with the same seed but at 2.6 grams/flat.

[0055] Treatment sequence was the same within each of the above 7 sections of flats numbered 1-56. In each section the first flat was the true control, no additional soil additives. The second flat was treated with a commercial soil additive called Delt Ag, the third another product called Plot Max and the fourth with a Yucca extract. Delt Ag Seed Coat is a blend of organically complexed nutrients and is specifically designed to enhance seedling emergence and plant growth. It contains sulfur, soluble manganese and soluble zinc (from manganese sulfate and zinc sulfate. It is applied directly to the seed before planting. Plot Max is a commercial product sold by Antler King (Wisconsin). It comprises a liquid product diluted in water and applied to new or existing food plots in the spring or fall. Food plots are areas of forage that farmers keep for wild game such as deer. The active ingredient in Plot Max is humic acid (2%). The fifth, sixth and seventh flats were treated with cobalt lactate in combination with inorganic minerals (matching some of the positive control commercial product inorganic mineral selections) but at increasing levels. The last flat in each section was treated with chelated minerals, cobalt, zinc, copper and manganese lactate.

The flats were maintained daily and no other intervention was made, other than accommodating for plant sun/shade tolerance differences and weeding as necessary.

Table 2

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Compo- site								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	1	2	3	4	5	6	7	8
Seed Type		Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix
Seed Planted (grams)		2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6
Seed Planted (date)		7/11/2 009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009
Seeding Rate (#/acre)		125	125	125	125	125	125	125	125
Soil Treated (date)		7/11/2 009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009
Soil Treated (with)		Nothin g	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Comple x 1	Cobalt Lactate Minera l Comple x II
Treatment (grams per acre)		Nothin g	18	1920	1529	50	125	1238	140
DM Treat (grams/acre)		Nothin g	18	58	535	50	125	1238	140
Solution (grams/mls H2O)		Water only	0.083/ 1000	0.882/ 1000	0.5/10 00	0.23/ 1000	0.574/ 1000	0.5682/ 1000	0.574/ 1000
Applied (solution in H2O)		Water (place bo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H2O)									0.069/ 1000*
Applied (solution in H2O)									10 ml in 600
2nd Part Description									*3% min lactate

2nd Part Chelated Minerals									Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2
Organic manganese									5
Organic zinc									5
Organic copper									5
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									
Molybdenum (Mo)									
Sprouts (post planting days)		3	3	3	3	3	3	3	3
Plant Harvest Date		8/19/2 009	8/19/2 009	8/19/2 009	8/19/2 009	8/19/200 9	8/19/200 9	8/19/20 09	8/19/2 009
Age at Harvest (days)		39	39	39	39	39	39	39	39
Weight (gms, as-is)		66.5	201.66	268.57	261.55	203.43	223.55	302.69	247.87
% of Control		100	303.25	403.86	393.31	305.91	336.17	455.17	372.74
Soil Analysis (dry weight)		Not submit ted	Not submit ted	Not submit ted	Not submit ted	Not submitte d	Not submitte d	Not submitt ed	Not submit ted
Moisture (%)	52.91								
pH	7.1								
Aerobic Plate Count (Mcfu/g)	2.4								
Anaerobic Plate Count (Mcfu/g)	0.1								
Azotobacter (Mcfu/g)	3								

Microbial Activity (ug/10g soil/day)	245								
Total Nitrogen (N, %)	1.13								
Ammonia Nitrogen (%)	0								
Nitrate Nitrogen (%)	nd								
Organic Nitrogen (%)	1.06								
Phosphorus (P2O5, %)	0.23								
Potassium (K2O, %)	0.25								
Sulfur (S, %)	0.36								
Calcium (Ca, %)	4.44								
Magnesium (Mg, %)	1.83								
Sodium (Na, %)	0.02								
Copper (Cu, ppm)	nd								
Iron (Fe, ppm)	7539								
Manganese (Mn, ppm)	200								
Zinc (Zn, ppm)	65.8								
Total salts (per ton)	3.61								
Total Carbon (%)	25.63								
C/N Ratio	22.8:1								
Chloride	nd								
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen (mg/L)	246								
Phosphorus (P, mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								

Sodium (Na, mg/L)	91								
Plant Cuttings mixed forage (DW basis)									
Moisture (%)		85.22	85.35	85.89	85.82	85.75	83.38	84.88	83.96
Dry Matter (%)		14.78	14.65	14.11	14.18	14.25	16.62	15.12	16.04
Crude Protein (%)		12.6	12.4	12.9	12.4	13.5	12.6	11.8	12.3
Crude Fat (%)									
Acid Detergent Fiber (%)		35.2	35.1	35.3	37.5	40.4	33	35.5	40.3
Ash (%)									
Total digestible nutrients (%)		62.4	62.5	62.3	59.8	56.5	64.9	62.1	56.6
Net energy-lactation (Mcal/lb)		0.64	0.64	0.64	0.61	0.57	0.67	0.64	0.58
Net energy-maint. (Mcal/lb)		0.62	0.62	0.62	0.59	0.55	0.65	0.61	0.55
Net energy-gain (Mcal/lb)		0.34	0.34	0.34	0.35	0.32	0.38	0.34	0.32
Digestible Energy (Mcal/lb)									
Metabolizable energy (Mcal/lb)									
Sulfur (%)		0.57	0.55	0.59	0.5	0.54	0.5	0.55	0.47
Phosphorus (%)		0.52	0.34	0.36	0.29	0.32	0.29	0.33	0.29
Potassium (%)		4.61	5.2	5.37	4.39	4.45	4.26	4.96	4.7
Magnesium (%)		1.04	1	0.94	0.86	0.83	1.03	0.99	0.84
Calcium (%)		2.09	1.94	1.87	1.82	1.97	2.08	2.11	1.75
Sodium (%)		0.07	0.09	0.1	0.08	0.08	0.07	0.08	0.07
Iron (ppm)		69	64	65	59	64	62	60	63
Manganese (ppm)		18	19	17	23	23	23	24	19
Copper (ppm)		7	5	6	5	6	6	7	5
Zinc (ppm)		106	89	80	81	92	85	89	81
Cobalt (ppm)		1.8	0.9	0.51	0.49	0.48	0.36	0.38	0.26

[0056] The Delt Ag flat may be considered the control. Regardless, the flat with the highest growth was Number 7, treated with 1238 grams of a mix of chelated

cobalt and inorganic manganese, copper and zinc. Of all the treatments, it had the best response, about 50% ahead of the commercialized Delt Ag soil treatment product that contains minerals but no chelated cobalt. It along with all the other flats in this series was planted at 2.6 grams/flat, the same seeding rate that the 2008 grass was planted at.

[0057] Of interest in this series is Number 8 flat, treated with a combination of mineral lactates that include not only cobalt, but manganese, copper and zinc lactates yielding mineral concentrations as shown in the table. This combination is termed cobalt lactate mineral complex 2. Note that the growth was ahead of Number 2 flat, the Delt Ag product.

[0058] All of the plants harvested from each section were submitted for forage testing. Mineral uptake by the plants was measured and observed for possible compositional changes in the plants that might lead to differences in plant-available energy/nutrients. Flat 6, treated with chelated minerals, cobalt, zinc, copper and manganese lactate at 125 grams/acre appears to have a meaningful increase in total digestible nutrients.

Table 3

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Composite								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	9	10	11	12	13	14	15	16
Seed Type		Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix
Seed Planted (grams)		5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2
Seed Planted (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Seeding Rate (#/acre)		249	249	249	249	249	249	249	249
Soil Treated (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Soil Treated (with)		No-thing	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	COBALT LACTATE MINERAL COMPLEX 2
Treatment		No-	18	1920	1529	50	125	1238	140

(grams per acre)		thing							
DM Treat (grams/acre)		No-thing	18	58	535	50	125	1238	140
Solution (grams/mls H ₂ O)		Water only	0.083/1000	0.882/1000	0.5/1000	0.23/1000	0.574/1000	0.5682/1000	0.574/1000
Applied (solution in H ₂ O)		Water (placebo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H ₂ O)									0.069/1000*
Applied (solution in H ₂ O)									10 ml in 600
2nd Part Description									*3% min lactate
2nd Part Chelated Minerals									Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2
Organic manganese									5
Organic zinc									5
Organic copper									5
Total Nitrogen (N)									
Phosphate (P ₂ O ₅)									
Potash (K ₂ O)									
Chelated Iron									
Molybdenum (Mo)									
Sprouts (post planting days)		3	3	3	3	3	3	3	3
Plant Harvest Date		8/16/2009	8/16/2009	8/16/2009	8/16/2009	8/16/2009	8/16/2009	8/16/2009	8/16/2009
Age at Harvest		oi	36	36	36	36	36	36	36

(days)									
Weight (gms, as-is)		132.07	231.98	193.82	248.28	261.66	207.52	184.1	96.32
% of Control		100	175.65	146.76	187.99	198.12	157.13	139.40	72.93
Soil Analysis (dry weight)		Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted
Moisture (%)	52.91	65.3	53.7	59.26	56.3	54.22	55.7	51.94	57.64
pH	7.1	7.5	7.3	7.5	7.6	7.5	7.6	7.5	7.6
Aerobic Plate Count (Mcfu/g)	2.4	15	13	14	18	20	7.5	10	20
Anaerobic Plate Count (Mcfu/g)	0.1	1.2	0.32	0.63	1	0.51	0.48	1.3	0.63
Azotobacter (Mcfu/g)	3	125	17	81	31	44	22	62	92
Microbial Activity (ug/10g soil/day)	245	2156	1573	1754	1577	2043	1699	1823	1721
Total Nitrogen (N, %)	1.13	1.07	1.23	1.25	1.1	1.09	1.22	1.12	1.2
Ammonia Nitrogen (%)	0	0.01	0.01	0.02	nd	0.03	0.01	0.01	0.01
Nitrate Nitrogen (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd
Organic Nitrogen (%)	1.06	1.06	1.22	1.23	1.1	1.06	1.21	1.11	1.19
Phosphorus (P2O5, %)	0.23	0.32	0.3	0.27	0.25	nd	0.27	0.25	0.28
Potassium (K2O, %)	0.25	nd	nd	nd	nd	nd	nd	nd	Nd
Sulfur (S, %)	0.36	0.4	0.41	0.42	0.48	0.37	0.41	0.37	0.4
Calcium (Ca, %)	4.44	7.72	4.41	8.2	6	7.36	5.78	3.56	4.08
Magnesium (Mg, %)	1.83	3.6	1.64	3.53	2.43	3.6	2.51	1.31	1.49
Sodium (Na, %)	0.02	0.03	0.04	0.02	0.03	0.04	0.02	0.04	0.05
Copper (Cu, ppm)	nd	nd	nd	nd	nd	nd	nd	nd	Nd
Iron (Fe, ppm)	7539	7565	8475	8623	7952	7237	7226	7039	8050
Manganese (Mn, ppm)	200	245	326	280	220	199	208	198	224
Zinc (Zn, ppm)	65.8	86.5	90.7	85.9	107.6	67.7	74.5	81.1	111
Total salts (per ton)	3.61	4.3	3.39	5.3	4.17	5.54	4.22	2.9	2.89
Total Carbon (%)	25.63	26.6	25.01	25.85	24.87	21.91	25.37	23.85	27.41
C/N Ratio	22.8:1	24.9:1	20.3:1	20.6:1	22.6:1	20.1:1	20.8:1	21.2:1	22.8:1

Chloride	nd	nd	nd	nd	nd	nd	nd	nd	Nd
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen (mg/L)	246								
Phosphorus (P,mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								
Sodium (Na, mg/L)	91								
Plant Cuttings mixed forage (DW basis)									
Moisture (%)		86.21	83.4	87.54	85.68	87.21	88.2	88.11	86.78
Dry Matter (%)		13.79	16.6	12.46	14.32	12.79	11.8	11.89	13.22
Crude Protein (%)		14.5	14.5	15.4	15.6	14.1	19.8	17.9	16.4
Crude Fat (%)									
Acid Detergent Fiber (%)		26.7	17.5	24.5	28	24.5	37.1	33.1	34.8
Ash (%)									
Total digestible nutrients (%)		72.1	82.6	74.6	70.6	74.6	60.2	64.8	62.9
Net energy-lactation (Mcal/lb)		0.75	0.87	0.78	0.73	0.78	0.62	0.67	0.65
Net energy-maint. (Mcal/lb)		0.74	0.86	0.77	0.72	0.77	0.59	0.65	0.62
Net energy-gain (Mcal/lb)		0.48	0.58	0.51	0.46	0.51	0.35	0.37	0.35
Digestible Energy (Mcal/lb)									
Metabolizable energy (Mcal/lb)									

Sulfur (%)		0.6	0.52	0.58	0.67	0.5	0.58	0.7	0.69
Phosphorus (%)		0.3	0.29	0.41	0.34	0.32	0.36	0.4	0.35
Potassium (%)		3.76	5.48	5.67	4.74	5.75	6.47	5.73	4.9
Magnesium (%)		0.74	0.8	0.91	0.89	0.79	0.89	0.93	0.81
Calcium (%)		1.99	1.69	2.01	2.19	1.82	1.85	2.1	2.01
Sodium (%)		0.09	0.07	0.08	0.11	0.08	0.1	0.1	0.11
Iron (ppm)		311	73	71	105	87	84	93	112
Manganese (ppm)		39	24	25	25	23	20	25	20
Copper (ppm)		11	9	9	7	7	8	8	7
Zinc (ppm)		104	72	94	76	76	80	94	96
Cobalt (ppm)		0.34	0.14	0.14	0.18	0.18	0.14	0.29	0.17

[0059] The next series of flats also includes the mixture of clovers, but planted at 5.2 grams/flat versus 2.6 grams/flat. The written and qualitative description given by the Fast & Fine Grass seed manufacturer was followed to establish the 2.6 and 5.2 grams seed/flat. In this series Flat 13 utilized the chelated minerals, cobalt, zinc, copper and manganese lactate planted at 50 grams/acre, which had the best response, about 200% of the control growth weight at cutting. These flats were harvested at 36 days of age, compared to 39 days of age in Flats 1-8. Flats had to be staged in harvesting as there was a lot of work getting them prepared for shipment. All harvested plants after cutting in the greenhouse were weighed first in the lab, frozen in zip lock bags then forwarded frozen to the lab for testing. Soils if to be sent to the lab were separated from plant roots, collected, placed in zip lock bags and sent immediately to the lab for testing

[0060] The 50 grams/flat was chosen as a level because it matched pretty closely the per acre coverage of inorganic minerals presented with the Delt Ag product (for manganese and zinc). This demonstrates the power of the chelated cobalt when all other things are the same (13% ahead of that positive control).

[0061] Of interest in this series of flats was that Flat 13 apparently exhausted all the available phosphorus from the soil. The soil used for all flats was complete in nitrogen, phosphorus and potash and there was enough to last 3 to 4 months, but not for Flat 13.

Table 4

WO 2011/078891

PCT/US2010/041848

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Compo-site								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	17	18	19	20	21	22	23	24
Seed Type		Fast & Fine	Fast & Fine	Fast & Fine	Fast & Fine	Fast & Fine	Fast & Fine	Fast & Fine	Fast & Fine
Seed Planted (grams)		2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6
Seed Planted (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Seeding Rate (#/acre)		125	125	125	125	125	125	125	125
Soil Treated (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Soil Treated (with)		No-thing	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	COBALT LACTATE MINERAL COMPLEX 2
Treatment (grams per acre)		No-thing	18	1920	1529	50	125	1238	140
DM Treat (grams/acre)		No-thing	18	58	535	50	125	1238	140
Solution (grams/mls H2O)		Water only	0.083/1000	0.882/1000	0.5/1000	0.23/1000	0.574/1000	0.5682/1000	0.574/1000
Applied (solution in H2O)		Water (placebo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H2O)									0.069/1000*
Applied (solution in H2O)									10 ml in 600
2nd Part Description									*3% min lactate
2nd Part Chelated Minerals									Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2

WO 2011/078891

PCT/US2010/041848

Organic manganese									5
Organic zinc									5
Organic copper									5
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									
Molybdenum (Mo)									
Sprouts (post planting days)		5	5	5	5	5	5	5	5
Plant Harvest Date		8/29/-2009	8/29/2009	8/29/2009	8/29/2009	8/29/2009	8/29/2009	8/29/2009	8/29/2009
Age at Harvest (days)		49	49	49	49	49	49	49	49
Weight (gms, as-is)		42.95	52.68	49.27	52.93	49.07	98.69	107.77	45.38
% of Control		100	122.65	114.71	123.24	114.25	229.78	250.92	105.66
Soil Analysis (dry weight)		Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted
Moisture (%)	52.91	44.94	57.58	54.17	58.75	52.92	52.5	54.61	53
pH	7.1	7.4	7.5	7.5	7.5	7.5	7.7	7.7	7.7
Aerobic Plate Count (Mcfu/g)	2.4	9.8	8.3	5.6	3.3	7.9	10.4	12.3	10.1
Anaerobic Plate Count (Mcfu/g)	0.1	0.36	0.39	0.26	0.12	0.29	0.3	0.49	0.71
Azotobacter (Mcfu/g)	3	3.6	5.2	4.6	3.2	1.6	26	10.2	5.8
Microbial Activity (ug/10g soil/day)	245	1456	1771	1903	1926	1619	1692	1732	1556
Total Nitrogen (N, %)	1.13	1.04	1.32	1.11	1.36	1.08	1.26	1.17	1.15
Ammonia Nitrogen (%)	0	0	nd	nd	nd	nd	nd	nd	nd
Nitrate Nitrogen (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd
Organic	1.06	1.03	1.32	1.11	1.31	1.08	1.26	1.17	1.15

Nitrogen (%)									
Phosphorus (P ₂ O ₅ , %)	0.23	0.25	0.33	0.35	0.34	0.34	0.34	0.31	0.3
Potassium (K ₂ O, %)	0.25	nd	nd	nd	nd	nd	nd	nd	nd
Sulfur (S, %)	0.36	0.35	0.5	0.46	0.51	0.47	0.53	0.44	0.55
Calcium (Ca, %)	4.44	6.27	5.8	5.06	5.99	5.33	5.03	5.22	7.79
Magnesium (Mg, %)	1.83	2.74	2.31	1.7	2.3	2.08	1.73	1.96	3.49
Sodium (Na, %)	0.02	0.02	0.02	nd	0.02	0.02	0.02	0.02	0.02
Copper (Cu, ppm)	nd	nd	nd	nd	nd	nd	nd	nd	nd
Iron (Fe, ppm)	7539	7123	9974	8809	11576	10514	10322	9969	11702
Manganese (Mn, ppm)	200	232	264	253	272	268	288	251	353
Zinc (Zn, ppm)	65.8	70.8	103.7	87.3	106.7	99.8	113.7	94.7	106.4
Total salts (per ton)	3.61	5.54	4.01	3.61	3.99	4.01	3.82	3.8	5.85
Total Carbon (%)	25.63	20.21	26.9	24.15	28.9	23.13	26.55	25.47	25.09
C/N Ratio	22.8:1	19.5:1	20.4:1	21.7:1	21.3:1	21.4:1	21:01	21.8:1	21.8:1
Chloride	nd	nd	nd	nd	nd	nd	nd	nd	nd
Growth Media Extracts									
pH	7.5	7.9	7.8	7.8	7.6	7.8	7.8	8	7.2
Soluble salts (mS/cm)	4.1	2.01	1.58	2.1	2.19	2.41	2.64	2.35	1.32
Nitrate nitrogen (mg/L)	246	1	0	0	0	5	2	0	0
Phosphorus (P, mg/L)	0.7	0.3	0.3	0.4	1	0.4	0.7	0.4	0.8
Potassium (K, mg/L)	294	43	59	44	47	57	70	40	47
Calcium (Ca, mg/L)	578	230	306	315	282	358	397	361	329
Magnesium (mg, mg/L)	210	89	118	120	110	138	156	138	127
Sodium (Na, mg/L)	91	33	48	36	36	42	62	43	38
Plant Cuttings mixed forage (DW basis)									
Moisture (%)		77.64	77.9	77.09	75.85	75.27	76.99	77.16	74.4
Dry Matter (%)		22.36	22.1	22.91	24.15	24.73	23.01	22.84	25.6

Crude Protein (%)		12.4	12.8	11.7	10.6	9.6	11.1	12	10.7
Crude Fat (%)		2.15	1.4	2.11	1.35	1.58	2.27	2.03	1.39
Acid Detergent Fiber (%)		23	22.8	23.1	24.9	22.9	25.9	24.9	22.7
Ash (%)		13.8	10.4	14.9	10.7	10.5	10.9	11.2	10.2
Total digestible nutrients (%)		62	64.4	61.2	64.2	64.9	64	63.8	64.9
Net energy-lactation (Mcal/lb)		0.8	0.8	0.8	0.78	0.8	0.76	0.78	0.8
Net energy-maint. (Mcal/lb)		0.63	0.66	0.62	0.66	0.67	0.65	0.65	0.67
Net energy-gain (Mcal/lb)		0.35	0.39	0.34	0.38	0.39	0.38	0.38	0.4
Digestible Energy (Mcal/lb)		1.24	1.29	1.23	1.28	1.3	1.28	1.28	1.3
Metabolizable energy (Mcal/lb)		1.16	1.2	1.15	1.2	1.22	1.2	1.19	1.22
Sulfur (%)		0.35	0.37	0.32	0.35	0.31	0.34	0.46	0.36
Phosphorus (%)		0.38	0.39	0.36	0.41	0.35	0.34	0.35	0.32
Potassium (%)		3.45	3.5	3.32	2.9	2.73	3.19	3.43	2.77
Magnesium (%)		0.44	0.44	0.45	0.43	0.38	0.4	0.49	0.41
Calcium (%)		0.96	1.02	1.07	0.96	0.86	0.95	1.3	0.85
Sodium (%)		0.06	0.07	0.06	0.05	0.05	0.06	0.07	0.06
Iron (ppm)		61	58	55	56	72	63	79	54
Manganese (ppm)		37	38	37	45	53	57	63	45
Copper (ppm)		5	6	5	5	4	6	5	4
Zinc (ppm)		89	91	92	80	79	94	122	73
Cobalt (ppm)		6.6	4.1	3.5	3.7	3.5	5.92	5.43	1.1

[0062] The next series of flats, numbered 17 through 24 summarized in Table 4, contained the seeds used from the Fast & Fine Grass, noted in Table 1.

[0063] The results were the same with the cobalt lactate treated flats having the highest grass weights. Flats 23 and 22 had the best growth, over 200% of the control. Flat 23 was treated with 1238 grams/acre Cobalt Lactate Mineral Complex I and Flat 22 at 250 grams acre with Cobalt Lactate Mineral Complex I. Flat 23 was planted at exactly the same application rate of cobalt lactate as in 2008, 1 ppm based

upon soil, and led the series at 251% of the control flat. Cobalt Lactate Mineral Complex I contains cobalt lactate, zinc sulfate, manganese sulfate, copper sulfate, Yucca extract, hemicellulose extract, Enzyme W, Porzyme and dextrose.

[0064] In those two flats the total aerobic plate count and Azotobacter count surged in numbers over all other treatments, the control and the baseline soil composite soil sample. It is clear in this series of flats that the relationship between soil microbes and growth is cause-and-effect; hence the support for the relationship between cobalt lactate and microbial populations that support plant growth. This is the second year now that 1ppm cobalt lactate applied to the soil has resulted in double the grass growth.

[0065] The higher level of soil treatment minerals, at least in part, from the higher application rate of chelated minerals, cobalt, zinc, copper and manganese lactate in Flat 23 resulted in increased minerals being taken up by the grass in that flat. A look at the data shows that sulfur, magnesium, calcium, iron, manganese, zinc and cobalt were all higher or in the high-end range of plant mineral content for Flat 23 against all other treatments. The formulation includes chelated minerals, cobalt, zinc, copper and manganese lactate, hence the increased sulfur content (also important for plant growth), a small amount of Yucca extract, two enzymes and larch arabinogalactan. All of the combined effects may be adding to increased mineral transport from the soil to the plant, increasing available magnesium and calcium. Iron is also found in the cobalt carbonate raw material that is reacted with lactic acid to form cobalt lactate, hence the increased iron content, again, at least in part.

Table 5

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Composite								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	25	26	27	28	29	30	31	32
Seed Type		Food Plot Forage Mix	Food Plot Forage Mix	Food Plot Forage Mix	Food Plot Forage Mix	Food Plot Forage Mix	Food Plot Forage Mix	Food Plot Forage Mix	Food Plot Forage Mix
Seed Planted (grams)		5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2
Seed Planted (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Seeding Rate		249	249	249	249	249	249	249	249

WO 2011/078891

PCT/US2010/041848

(#/acre)									
Soil Treated (date)		7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009
Soil Treated (with)		No- thing	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	COBALT LACTATE MINERAL COMPLEX 2
Treatment (grams per acre)		No- thing	18	1920	1529	50	125	1238	140
DM Treat (grams/acre)		No- thing	18	58	535	50	125	1238	140
Solution (grams/mls H2O)		Water only	0.083/ 1000	0.882/ 1000	0.5/ 1000	0.23/ 1000	0.574/ 1000	0.5682/ 1000	0.574/ 1000
Applied (solution in H2O)		Water (pla- cebo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H2O)									0.069/ 1000*
Applied (solution in H2O)									10 ml in 600
2nd Part Description									*3% min lactate
2nd Part Chelated Minerals									Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2
Organic manganese									5
Organic zinc									5
Organic copper									5
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									
Molybdenum (Mo)									

WO 2011/078891

PCT/US2010/041848

Sprouts (post planting days)		2	2	2	2	2	2	2	2
Plant Harvest Date		8/12/2009	8/23/2009	8/23/2009	8/23/2009	8/23/2009	8/23/2009	8/23/2009	8/23/2009
Age at Harvest (days)		lost	43	43	43	43	43	43	43
Weight (gms, as-is)		vandalism	123.75	221.03	512.46	448.78	428.12	389.88	189.51
% of Control			100	178.61	414.11	362.65	345.96	315.05	153.14
Soil Analysis (dry weight)		Not submitted	Not submitted	Not submitted	Not submitted	Not submitted	Not submitted	Not submitted	Not submitted
Moisture (%)	52.91								
pH	7.1								
Aerobic Plate Count (Mcfu/g)	2.4								
Anaerobic Plate Count (Mcfu/g)	0.1								
Azotobacter (Mcfu/g)	3								
Microbial Activity (ug/10g soil/day)	245								
Total Nitrogen (N, %)	1.13								
Ammonia Nitrogen (%)	0								
Nitrate Nitrogen (%)	nd								
Organic Nitrogen (%)	1.06								
Phosphorus (P2O5, %)	0.23								
Potassium (K2O, %)	0.25								
Sulfur (S, %)	0.36								
Calcium (Ca, %)	4.44								
Magnesium (Mg, %)	1.83								
Sodium (Na, %)	0.02								
Copper (Cu, ppm)	nd								
Iron (Fe, ppm)	7539								
Manganese (Mn, ppm)	200								
Zinc (Zn, ppm)	65.8								
Total salts (per ton)	3.61								
Total Carbon (%)	25.63								
C/N Ratio	22.8:1								

Chloride	nd								
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen (mg/L)	246								
Phosphorus (P,mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								
Sodium (Na, mg/L)	91								
Plant Cuttings mixed forage (DW basis)		Vandal-ized							
Moisture (%)			78.97	79.11	83.75	84.94	84.38	84.64	82.41
Dry Matter (%)			21.03	20.89	16.25	15.06	15.62	15.54	17.59
Crude Protein (%)			8.39	8.1	8.57	8.34	9.21	8.93	9.69
Crude Fat (%)			2.28	2.36	3.42	3.21	2.64	2.4	2.5
Acid Detergent Fiber (%)			16.3	17.8	21.6	20.1	18.9	22.7	26.3
Ash (%)			11.9	13	11.9	13.1	13	12.9	13.7
Total digestible nutrients (%)			65	64	64.4	63.7	63.7	63.2	61.9
Net energy-lactation (Mcal/lb)			0.88	0.86	0.81	0.83	0.85	0.8	0.75
Net energy-maint. (Mcal/lb)			0.67	0.65	0.66	0.65	0.65	0.64	0.63
Net energy-gain (Mcal/lb)			0.4	0.38	0.39	0.38	0.38	0.37	0.35
Digestible Energy (Mcal/lb)			1.3	1.28	1.29	1.27	1.27	1.26	1.24
Metabolizable energy (Mcal/lb)			1.23	1.21	1.21	1.2	1.2	1.19	1.16
Sulfur (%)			1.13	1.03	0.88	0.92	0.92	0.83	1.04
Phosphorus (%)			0.38	0.28	0.35	0.35	0.38	0.36	0.4
Potassium (%)			2.14	2.18	2.91	2.92	3.02	2.89	3.02
Magnesium (%)			0.68	0.67	0.57	0.62	0.6	0.56	0.68
Calcium (%)			2.89	2.89	2.9	3.07	2.86	2.84	3.15
Sodium (%)			0.13	0.17	0.08	0.1	0.09	0.09	0.12
Iron (ppm)			42	36	80	43	43	52	56

Manganese (ppm)			20	20	27	27	30	33	22
Copper (ppm)			3	2	2	2	3	2	3
Zinc (ppm)			65	80	93	110	88	93	93
Cobalt (ppm)			3.12	5.9	4.35	4.67	4.3	2.75	4.08

[0066] Flats 25-32 are those in the next series, which include a combination of seeds that includes Green Globe Turnips, Dwarf Essex Forage Rapeseed, Rangi Forage and Giant Rapeseed. They are largely sold to land owners to support wild life.

[0067] The best performances in this series of flats were numbers 28, 29 and 30, respectively. Flat 26 treated with Delt Ag was used as the control. With that in mind, flats 28, 29 and 30 were about 300 to 400% of the control and looked pretty remarkable compared to it.

[0068] Flats 29 and 30 were treated with chelated minerals, cobalt, zinc, copper and manganese lactate at 50 and 125 grams/acre, respectively. However, the best flat in the series was treated with a Yucca extract. The literature on Yucca and improved plant performance relates to soil treatment and improved permeability of plant cells to growth media in the soil. Of the seven series of flats that were treated with Yucca extract and the chelated minerals, cobalt, zinc, copper and manganese lactate, Yucca performed best in two of those series.

Table 6

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Composite								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	33	34	35	36	37	38	39	40
Seed Type		Food Plot Grain Mix	Food Plot Grain Mix	Food Plot Grain Mix	Food Plot Grain Mix	Food Plot Grain Mix	Food Plot Grain Mix	Food Plot Grain Mix	Food Plot Grain Mix
Seed Planted (grams)		5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2
Seed Planted (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Seeding Rate		249	249	249	249	249	249	249	249

(#/acre)									
Soil Treated (date)		7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009	7/11/ 2009
Soil Treated (with)		No-thing	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	COBALT LACTATE MINERAL COMPLEX 2
Treatment (grams per acre)		No-thing	18	1920	1529	50	125	1238	140
DM Treat (grams/acre)		No-thing	18	58	535	50	125	1238	140
Solution (grams/mls H2O)		Water only	0.083/ 1000	0.882/ 1000	0.5/ 1000	0.23/ 1000	0.574/ 1000	0.5682/ 1000	0.574/ 1000
Applied (solution in H2O)		Water (placebo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H2O)									0.069/ 1000*
Applied (solution in H2O)									10 ml in 600
2nd Part Description									*3% min lactate
2nd Part Chelated Minerals									Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2
Organic manganese									5
Organic zinc									5
Organic copper									5
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									

Molybdenum (Mo)									
Sprouts (post planting days)		2	2	2	2	2	2	2	2
Plant Harvest Date		8/15/2009	8/15/2009	8/15/2009	8/15/2009	8/15/2009	8/12/2009	8/12/2009	8/12/2009
Age at Harvest (days)		35	35	35	35	35	lost	lost	lost
Weight (gms, as-is)		426.28	442.08	355.57	353.26	492.63	vandalism	vandalism	vandalism
% of Control		100	103.71	83.41	82.87	115.56			
Soil Analysis (dry weight)		Submitted	Submitted	Submitted	Submitted	Submitted	N/A	N/A	N/A
Moisture (%)	52.91	64.27	54.48	62.92	58.91	60.35			
pH	7.1	7.6	7.6	7.5	7.5	7.5			
Aerobic Plate Count (Mcfu/g)	2.4	19	25	21	13	5			
Anaerobic Plate Count (Mcfu/g)	0.1	0.83	0.73	0.53	0.7	0.91			
Azotobacter (Mcfu/g)	3	60	111	37	55	61			
Microbial Activity (ug/10g soil/day)	245	2255	2287	2216	2021	2069			
Total Nitrogen (N, %)	1.13	1.23	1.19	1.32	1.34	1.21			
Ammonia Nitrogen (%)	0	0.01	0	nd	nd	0			
Nitrate Nitrogen (%)	nd	nd	nd	nd	nd	nd			
Organic Nitrogen (%)	1.06	1.23	1.18	1.32	1.34	1.21			
Phosphorus (P2O5, %)	0.23	0.31	0.24	0.3	0.27	nd			
Potassium (K2O, %)	0.25	nd	nd	nd	nd	nd			
Sulfur (S, %)	0.36	0.45	0.37	0.51	0.44	0.48			
Calcium (Ca, %)	4.44	4.42	3.67	7.04	4.33	9.96			
Magnesium (Mg, %)	1.83	1.68	1.32	3.18	1.53	3.98			
Sodium (Na, %)	0.02	0.03	0.02	0.16	0.05	0.05			

(%)									
Copper (Cu, ppm)	nd	nd	nd	nd	nd	nd			
Iron (Fe, ppm)	7539	8741	7377	8862	8866	8971			
Manganese (Mn, ppm)	200	227	209	243	221	245			
Zinc (Zn, ppm)	65.8	98	96.7	94.4	94.9	90.8			
Total salts (per ton)	3.61	2.63	2.82	4.34	2.98	6.03			
Total Carbon (%)	25.63	26.14	24.63	26.86	27.77	25.3			
C/N Ratio	22.8:1	21.2:1	20.8:1	20.3:1	20.7:1	20.9:1			
Chloride	nd	nd	nd	nd	nd	nd			
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen (mg/L)	246								
Phosphorus (P, mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								
Sodium (Na, mg/L)	91								
Plant Cuttings mixed forage (DW basis)							Vandal-ism	Vandal-ism	Vandal-ism
Moisture (%)		84.7	84.01	83.35	84.67	83.6			
Dry Matter (%)		15.3	15.99	16.65	15.33	16.4			
Crude Protein (%)		6.96	5.81	7.14	6.39	7.08			
Crude Fat (%)									
Acid Detergent Fiber (%)		34.4	33.8	29.5	36.8	30.4			

Ash (%)									
Total digestible nutrients (%)		63.3	64	68.9	60.6	67.9			
Net energy-lactation (Mcal/lb)		0.65	0.66	0.71	0.62	0.7			
Net energy-maint. (Mcal/lb)		0.63	0.64	0.7	0.6	0.68			
Net energy-gain (Mcal/lb)		0.35	0.36	0.43	0.36	0.42			
Digestible Energy (Mcal/lb)									
Metabolizable energy (Mcal/lb)									
Sulfur (%)			0.39	0.29	0.33	0.3	0.28		
Phosphorus (%)		0.25	0.23	0.25	0.24	0.19			
Potassium (%)		3.43	3.14	3.38	3.25	3.07			
Magnesium (%)		0.55	0.47	0.48	0.43	0.47			
Calcium (%)		1.11	0.88	0.87	0.86	0.89			
Sodium (%)		0.05	0.03	0.04	0.03	0.03			
Iron (ppm)		101	61	54	35	56			
Manganese (ppm)		20	15	9	6	9			
Copper (ppm)		13	8	4	3	3			
Zinc (ppm)		50	39	42	34	36			
Cobalt (ppm)		0.9	0.29	0.27	0.15	0.2			

[0069] The next series of flats includes the Food Plot Grain Mix, a combination of seeds for game as described earlier. The plants were over three feet tall. The chelated minerals, cobalt, zinc, copper and manganese lactate at 50 grams per acre was significantly ahead of all the other treatments and control.

Table 7

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Com- posite								
Each Flat, 0.25 CUFT soil	Soil Analysis								

WO 2011/078891

PCT/US2010/041848

Flat Number	1 - 56	41	42	43	44	45	46	47	48
Seed Type		Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)
Seed Planted (grams)		5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2
Seed Planted (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Seeding Rate (#/acre)		249	249	249	249	249	249	249	249
Soil Treated (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Soil Treated (with)		No-thing	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	COBALT LACTATE MINERAL COMPLEX 2
Treatment (grams per acre)		No-thing	18	1920	1529	50	125	1238	140
DM Treat (grams/acre)		No-thing	18	58	535	50	125	1238	140
Solution (grams/mls H2O)		Water only	0.083/1000	0.882/1000	0.5/1000	0.23/1000	0.574/1000	0.5682/1000	0.574/1000
Applied (solution in H2O)		Water (pla-cebo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H2O)									0.069/1000*
Applied (solution in H2O)									10 ml in 600
2nd Part Description									*3% min lactate
2nd Part Chelated Minerals									Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2
Organic manganese									5
Organic zinc									5

WO 2011/078891

PCT/US2010/041848

Organic copper									5
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									
Molybdenum (Mo)									
Sprouts (post planting days)		1	1	1	1	1	1	1	1
Plant Harvest Date		8/25/2009	8/25/2009	8/25/2009	8/25/2009	8/25/2009	8/25/2009	8/25/2009	8/25/2009
Age at Harvest (days)		45	45	45	45	45	45	45	45
Weight (gms, as-is)		198.02	198.85	231.52	205.83	227.66	209.69	263.44	185.64
% of Control		100	100.42	116.92	103.94	114.97	105.89	133.04	93.75
Soil Analysis (dry weight)		Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted	Sub-mitted
Moisture (%)	52.91	43.14	49.94	47.12	48.71	47.35	46.91	50.51	53.48
pH	7.1	7.5	7.7	7.6	7.6	7.7	7.6	7.6	7.6
Aerobic Plate Count (Mcfu/g)	2.4	1.3	4.4	2.1	4	3.3	2.2	2.1	2
Anaerobic Plate Count (Mcfu/g)	0.1	0.15	0.24	0.07	0.074	0.082	0.21	0.07	0.042
Azotobacter (Mcfu/g)	3	33	55	33	43	12	9	29	26
Microbial Activity (ug/10g soil/day)	245	1891	2264	2149	2029	1995	1169	2387	2011
Total Nitrogen (N, %)	1.13	1.23	1.44	1.44	1.52	1.52	1.49	1.33	1.53
Ammonia Nitrogen (%)	0	nd	nd	nd	nd	nd	nd	nd	nd
Nitrate Nitrogen (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd

Organic Nitrogen (%)	1.06	1.23	1.44	1.44	1.52	1.52	1.49	1.33	1.53
Phosphorus (P ₂ O ₅ , %)	0.23	0.23	0.22	0.25	0.27	0.32	0.26	0.32	0.28
Potassium (K ₂ O, %)	0.25	nd	nd	nd	nd	nd	nd	nd	nd
Sulfur (S, %)	0.36	0.37	0.36	0.36	0.37	0.46	0.4	0.4	0.37
Calcium (Ca, %)	4.44	3.78	3.74	4.65	4.82	4.5	4.86	4.73	4.39
Magnesium (Mg, %)	1.83	1.42	1.32	1.93	1.93	1.67	1.88	1.82	1.68
Sodium (Na, %)	0.02	nd	nd	0.02	nd	0.02	nd	nd	nd
Copper (Cu, ppm)	nd	nd	nd	nd	nd	nd	nd	nd	nd
Iron (Fe, ppm)	7539	7355	7697	7685	9503	9299	8574	9028	7556
Manganese (Mn, ppm)	200	190	208	216	238	281	220	238	211
Zinc (Zn, ppm)	65.8	77.4	89.9	83.2	105.3	106.4	92.3	123.3	101
Total salts (per ton)	3.61	3.67	3.26	4.25	4.25	4.06	4.38	3.91	3.54
Total Carbon (%)	25.63	23.21	26.07	25.51	26.73	25.38	27.93	25.94	28.31
C/N Ratio	22.8:1	18.9:1	18.1:1	17.8:1	17.6:1	16.7:1	18.8:1	19.5:1	18.5:1
Chloride	nd	nd	nd	nd	nd	nd	nd	nd	nd
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen (mg/L)	246								
Phosphorus (P, mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								
Sodium (Na, mg/L)	91								
Plant Cuttings mixed									

forage (DW basis)									
Moisture (%)		79.76	76.07	80.24	80.83	80.07	78.91	81.08	83
Dry Matter (%)		20.24	23.93	19.76	19.17	19.93	21.09	18.92	17
Crude Protein (%)		22.2	20.4	21.5	21.7	17.8	16.2	17	23.4
Crude Fat (%)		1.67	1.7	1.67	1.69	2.2	1.48	1.11	1.76
Acid Detergent Fiber (%)		39.6	38.7	38.8	31.4	32.6	34.8	48.7	25.4
Ash (%)		12.2	10.8	11.1	12.1	12.1	11.2	11.8	13.4
Total digestible nutrients (%)		64.2	65.5	65.2	65	65.1	65.7	64.1	64.4
Net energy-lactation (Mcal/lb)		0.57	0.58	0.58	0.67	0.66	0.63	0.46	0.74
Net energy-maint. (Mcal/lb)		0.66	0.67	0.67	0.67	0.67	0.68	0.65	0.66
Net energy-gain (Mcal/lb)		0.39	0.4	0.4	0.4	0.4	0.41	0.38	0.39
Digestible Energy (Mcal/lb)		1.28	1.31	1.3	1.3	1.3	1.31	1.28	1.29
Metabolizable energy (Mcal/lb)		1.18	1.2	1.2	1.19	1.2	1.22	1.19	1.18
Sulfur (%)		0.42	0.52	0.53	0.59	0.66	0.49	0.47	0.62
Phosphorus (%)		0.26	0.25	0.26	0.33	0.32	0.24	0.26	0.34
Potassium (%)		3.21	3.28	3.67	4.46	4.53	3.7	3.91	4.58
Magnesium (%)		0.37	0.49	0.49	0.49	0.55	0.44	0.43	0.51
Calcium (%)		1.48	2	2	2.06	2.2	1.74	1.67	2.05
Sodium (%)		0.05	0.06	0.07	0.07	0.08	0.06	0.07	0.07
Iron (ppm)		65	64	67	86	96	62	84	107
Manganese (ppm)		22	28	28	29	27	23	24	35
Copper (ppm)		4	4	4	4	5	4	4	5
Zinc (ppm)		49	49	60	63	71	54	64	66
Cobalt (ppm)		0.45	0.28	0.16	0.23	0.26	0.18	0.21	0.47

[0070] This series of flats is the first of two planted with alfalfa. A Vernal variety was chosen because it was reported that this variety is accepted across the industry as a greenhouse standard.

[0071] As seen in Table 7, treatments followed the same pattern as in all of the preceding series of flats. The best response in this series was from Flat 47, the seed treated with the highest level of the chelated minerals, cobalt, zinc, copper and manganese lactate, at 1238 grams/acre. Growth at 133% of the control was significantly above any other treatment response. Nonetheless, the best response in this seemingly optimized alfalfa seed was with the chelated minerals, cobalt, zinc, copper and manganese lactate.

Table 8

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Composite								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	49	50	51	52	53	54	55	56
Seed Type		Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)	Alfalfa (Vernal)
Seed Planted (grams)		2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6
Seed Planted (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Seeding Rate (#/acre)		125	125	125	125	125	125	125	125
Soil Treated (date)		7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009	7/11/2009
Soil Treated (with)		No-thing	Delt Ag S. Coat	Plot Max	Yucca	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	Cobalt Lactate Mineral Complex 1	FP250X
Treatment (grams per acre)		No-thing	18	1920	1529	50	125	1238	140
DM Treat (grams/acre)		No-thing	18	58	535	50	125	1238	140
Solution (grams/mls H2O)		Water only	0.083/1000	0.882/1000	0.5/1000	0.23/1000	0.574/1000	0.5682/1000	0.574/1000
Applied (solution in H2O)		Water (placebo)	10 ml in 600	100 ml in 600	100 ml in 600	10 ml in 600	10 ml in 600	100 ml in 600	10 ml in 600
2nd Part (grams/mls H2O)									0.069/1000*
Applied (solution in H2O)									10 ml in 600
2nd Part Description									*3% min lactate
2nd Part Chelated									Mn, Zn, Cu

Minerals									
Mineral Application, gm/acre									
Organic cobalt						2.2	5.5	54.5	5.5
Manganese			0.46			0.43	1.08	10.7	1.08
Zinc			0.53			0.49	1.23	12.2	1.23
Copper						0.48	1.2	11.9	1.2
Organic manganese									
Organic zinc									
Organic copper									
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									
Molybdenum (Mo)									
Sprouts (post planting days)		1	1	1	1	1	1	1	1
Plant Harvest Date		9/27/2009	9/27/2009	9/27/2009	9/27/2009	9/27/2009	9/27/2009	9/27/2009	9/27/2009
Age at Harvest (days)		78	78	78	78	78	78	78	78
Weight (gms, as-is)		318.86	311.61	301.77	323.46	248.36	249.64	221.66	222.11
% of Control		100	97.73	94.64	101.44	77.89	78.29	69.52	69.66
Soil Analysis (dry weight)									
Moisture (%)	52.91								
pH	7.1								
Aerobic Plate Count (Mcfu/g)	2.4								
Anaerobic Plate Count (Mcfu/g)	0.1								
Azotobacter (Mcfu/g)	3								
Microbial Activity (ug/10g soil/day)	245								
Total Nitrogen (N, %)	1.13								
Ammonia Nitrogen (%)	0								
Nitrate Nitrogen (%)	nd								

Organic Nitrogen (%)	1.06								
Phosphorus (P ₂ O ₅ , %)	0.23								
Potassium (K ₂ O, %)	0.25								
Sulfur (S, %)	0.36								
Calcium (Ca, %)	4.44								
Magnesium (Mg, %)	1.83								
Sodium (Na, %)	0.02								
Copper (Cu, ppm)	nd								
Iron (Fe, ppm)	7539								
Manganese (Mn, ppm)	200								
Zinc (Zn, ppm)	65.8								
Total salts (per ton)	3.61								
Total Carbon (%)	25.63								
C/N Ratio	22.8:1								
Chloride	nd								
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen (mg/L)	246								
Phosphorus (P, mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								
Sodium (Na, mg/L)	91								

[0072] The next series is the second Vernal alfalfa set, this time planted at 2.6 grams/flat versus the previous section at 5.6 grams/flat. The most notable characteristic of this series is that all flats except Flat 52, treated with the Yucca extract, performed worse than the control. The seeding rate was low enough that the

treatments (except for Yucca) were too much to increase growth and instead inhibited plant development.

Table 9

64 Flats, 2 Sq Ft/Flat									
Miracle-Gro Garden Soil	Com- posite								
Each Flat, 0.25 CUFT soil	Soil Analysis								
Flat Number	1 - 56	57	58	59	60	61	62	63	64
Seed Type		Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix	Food Plot Clover Mix
Seed Planted (grams)		5.2	5.2	5.2	5.2	5.2	5.2	5.2	5.2
Seed Planted (date)		7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009
Seeding Rate (#/acre)		249	249	249	249	249	249	249	249
Soil Treated (date)		7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009	7/12/ 2009
Soil Treated (with)		No- thing	Color- burst	Color- burst	Cobalt Lactate Mineral Complex 1	125 + Che- lated	BK Che- lated	BK Che- lated	BK Che- lated
Treatment (grams per acre)		No- thing	10890000	21780000	2476	1586	8712	17424	26136
DM Treat (grams/acre)		No- thing	32670	65340	2476	1586	348	696	1045
Solution (grams/mls H2O)		Water only	12/ 4000	12/ 4000	0.5682/ 1000	0.5682/ 1000	0.4/100	0.8/ 100	1.2/ 100
Applied (solution in H2O)		Water	500 ml	1000 ml	200 ml in 600	100 ml in 600	100 ml in 600	100 ml in 600	100 ml in 600
2nd Part (grams/mls H2O)									
Applied (solution in H2O)						0.4 gm neat			
2nd Part Description						4% min lactate	4% min lactate	4% min lactate	4% min lactate

WO 2011/078891

PCT/US2010/041848

2nd Part Chelated Minerals						Co, Mn, Zn, Cu	Co, Mn, Zn, Cu	Co, Mn, Zn, Cu	Co, Mn, Zn, Cu
Mineral Application, gm/acre									
Organic cobalt					109	5.5	21.8	43.6	65.4
Manganese					21.4	1.08			
Zinc			19.6	39.2	24.4	1.23			
Copper			22.8	45.6	23.8	1.2			
Organic manganese			16.3	32.6		21.8	21.8	43.6	65.4
Organic zinc						21.8	21.8	43.6	65.4
Organic copper						21.8	21.8	43.6	65.4
Total Nitrogen (N)									
Phosphate (P2O5)									
Potash (K2O)									
Chelated Iron									
Molybdenum (Mo)									
Sprouts (post planting days)		3	3	3	3	3	3	3	3
Plant Harvest Date		8/20/ 2009	8/20/ 2009	8/20/ 2009	8/20/ 2009	8/20/ 2009	8/20/ 2009	8/20/ 2009	8/20/ 2009
Age at Harvest (days)		39	39	39	39	39	39	39	39
Weight (gms, as-is)		242.59	259.33	443.38	238.55	158.66	287.23	281.26	255.05
% of Control		100	106.90	182.77	98.33	65.40	118.40	115.94	105.14
Soil Analysis (dry weight)		Sub- mitted	Sub- mitted	Sub- mitted	Sub- mitted	Sub- mitted	Sub- mitted	Sub- mitted	Sub- mitted
Moisture (%)	52.91	64.56	60.88	63.86	64.16	63.71	60.7	61.84	63.97
pH	7.1	6.9	6.9	6.9	7.1	7.2	7.1	7.2	7.3
Aerobic Plate Count (Mcfu/g)	2.4	12.8	34	25	15	16	36	16	32

Anaerobic Plate Count (Mcfu/g)	0.1	0.42	0.98	0.55	0.85	1.4	2	0.53	1.3
Azotobacter (Mcfu/g)	3	67	380	240	240	480	260	420	380
Microbial Activity (ug/10g soil/day)	245	669	681	752	635	1145	873	1014	953
Total Nitrogen (N, %)	1.13	1.44	1.23	1.58	1.26	1.13	1.3	1.42	1.39
Ammonia Nitrogen (%)	0	nd	0.01	0	nd	nd	0.01	0.01	nd
Nitrate Nitrogen (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd
Organic Nitrogen (%)	1.06	1.44	1.22	1.57	1.26	1.13	1.29	1.41	1.39
Phosphorus (P2O5, %)	0.23	0.31	0.28	0.39	0.39	0.3	0.41	0.29	0.44
Potassium (K2O, %)	0.25	nd	nd	nd	nd	nd	nd	nd	nd
Sulfur (S, %)	0.36	0.48	0.33	0.44	0.31	0.33	0.41	0.34	0.5
Calcium (Ca, %)	4.44	2.91	2.43	2.57	2.23	2.09	2.52	2.23	2.78
Magnesium (Mg, %)	1.83	0.73	0.66	0.61	0.53	0.5	0.66	0.55	0.72
Sodium (Na, %)	0.02	nd	nd	0.03	nd	0.22	nd	nd	nd
Copper (Cu, ppm)	nd	nd	nd	nd	nd	nd	nd	nd	nd
Iron (Fe, ppm)	7539	12960	10583	11309	12162	10394	12382	11593	11035
Manganese (Mn, ppm)	200	330	258	340	307	278	313	275	275
Zinc (Zn, ppm)	65.8	90.3	109.9	119	86.5	68.9	96.7	83.9	88.8
Total salts (per ton)	3.61	1.81	1.7	1.73	1.45	1.43	1.77	1.61	1.77
Total Carbon (%)	25.63	21.05	21.8	25.71	18.75	18.32	21.02	22.25	22.56
C/N Ratio	22.8:1	14.6:1	17.8:1	16.3:1	14.9:1	16.2:1	16.2:1	15.7:1	16.3:1
Chloride	nd	nd	nd	nd	nd	nd	nd	nd	nd
Growth Media Extracts									
pH	7.5								
Soluble salts (mS/cm)	4.1								
Nitrate nitrogen	246								

(mg/L)									
Phosphorus (P,mg/L)	0.7								
Potassium (K, mg/L)	294								
Calcium (Ca, mg/L)	578								
Magnesium (mg, mg/L)	210								
Sodium (Na, mg/L)	91								
Plant Cuttings mixed forage (DW basis)									
Moisture (%)		87.22	88.88	86.93	87.6	88.61	85.95	87.1	86.36
Dry Matter (%)		12.78	11.12	13.07	12.4	11.39	14.05	12.9	13.64
Crude Protein (%)		16.3	23.3	14.2	21.1	26.5	23.2	19.9	23.5
Crude Fat (%)		2.14	2.3	2.38	2.85	3.04	3.69	2.55	2.89
Acid Detergent Fiber (%)		22.3	26.5	27	20.6	19.7	17.6	23.6	27.2
Ash (%)		14.8	15.5	12.7	14	15.3	14.3	14.3	13.5
Total digestible nutrients (%)		60.2	57.5	61.6	60.2	58.2	60.1	59.7	59.1
Net energy-lactation (Mcal/lb)		0.8	0.75	0.75	0.83	0.84	0.86	0.79	0.74
Net energy- maint. (Mcal/lb)		0.6	0.57	0.62	0.6	0.58	0.6	0.6	0.59
Net energy- gain (Mcal/lb)		0.32	0.34	0.35	0.32	0.34	0.32	0.36	0.35
Digestible Energy (Mcal/lb)		1.2	1.15	1.23	1.2	1.16	1.2	1.19	1.18
Metabolizable energy (Mcal/lb)		1.12	1.05	1.15	1.1	1.06	1.1	1.1	1.08
Sulfur (%)		0.57	0.58	0.55	0.58	0.54	0.57	0.52	0.52
Phosphorus (%)		0.31	0.36	0.29	0.33	0.36	0.38	0.31	0.36
Potassium (%)		4.81	5.6	4.2	4.67	5.33	4.74	4.75	4.34
Magnesium (%)		0.79	0.73	0.71	0.57	0.61	0.73	0.57	0.58

Calcium (%)		1.68	1.49	1.14	1.81	1.74	2.02	1.63	1.76
Sodium (%)		0.07	0.09	0.05	0.11	0.07	0.06	0.07	0.06
Iron (ppm)		77	128	93	90	118	167	81	94
Manganese (ppm)		15	14	12	18	13	14	13	14
Copper (ppm)		4	4	2	4	4	4	4	4
Zinc (ppm)		74	85	52	106	107	106	95	93
Cobalt (ppm)		0.52	0.45	0.17	0.47	0.39	0.36	0.3	0.28

[0073] This is the last series of flats. They were planted one day after all the others to test very high mineral levels, such as applied to plants by a product called “Colorburst.” Colorburst is an all purpose, dual action (fast acting, long lasting) plant food for promoting “bigger, more beautiful plants.” It was utilized because it applied much higher levels of trace minerals- higher than any of the other products that were tested against or considered for the BK 125 product (Cobalt lactate mineral product I or II). Unlike the other mineral products, Colorburst also contains a source of nitrogen, phosphate and potash that supplements the minerals. Minerals include; water soluble copper (copper sulfate), chelated iron (iron EDTA), chelated manganese (manganese EDTA), molybdenum, water soluble zinc (zinc sulfate). The mineral application rates of this product were very high, on a per acre basis (~65K grams/acre).

[0074] In this series, the order of magnitude higher mineral application rates were tested, but also wanted to test all of the chelated minerals in combination (zinc, copper, manganese and cobalt) that were produced in pilot batches in the lab. In most of the previous flats the organic chelated cobalt was used in combination with inorganic minerals.

[0075] Food Plot Clover Mix seed planted at 5.2 grams/flat was used. Of all the flats, Colorburst in Flat 59 at about 65K grams/acre performed the best. The most important results may be in Flats 62, 63 and 64, where increasing levels of all the chelated minerals used in combination produced growth results ahead of the control. Of further interest here is that as the total levels increased the response lessened, never less than the control, but definitely a trend down with increasing levels of the four (4) chelated materials.

[0076] In Flats 62-64 zinc, copper and manganese were used that had been mixed (reacted) together in industry recommended proportions with lactic acid to produce a single product that contained all the mineral lactates together. This solid reaction mass was allowed to partially dissolve in water, then that solution was diluted and applied to the soil in those flats (after having been mixed as well with cobalt lactate powder to yield a mixture of all four mineral lactates). The solution was tested for the level of each mineral lactate and that information is available.

Table 10

Effect of cobalt on methane, pH, ammonia-N, and digestibility in continuous cultures of ruminal microorganisms.

Item	Treatments						SE	Probability > F		
	Cobalt Carbonate (ppm)			CoMax New (ppm)		CoMax Old (ppm)				
	Control	0.2	1.0	0.2	1.0	1.0		Source	Level	Source X Level
Methane, nmol/ml										
Pre-feeding*	430.55	363.91	347.82	375.83	417.51	382.42	53.76	0.54	0.50	0.79
Post-feeding*	579.30	403.14	515.35	590.46	590.74	576.21	51.13	0.04	0.27	0.19
Culture pH										
Pre-feeding*	5.98	5.81	5.88	5.95	5.87	5.99	0.09	0.52	0.38	0.63
Post-feeding*	5.73	5.56	5.55	5.64	5.65	5.57	0.05	0.14	0.01	0.57
Ammonia-N, mg/dL	12.23	10.79	10.88	11.80	11.95	10.32	1.49	0.57	0.79	0.92
NDF Digestibility, %	23.90	23.16	17.09	20.63	27.36	15.69	2.23	0.06	0.42	0.001
IVDMTD**, %	60.52	59.35	59.17	59.50	60.67	57.94	1.16	0.42	0.43	0.62

*Pre-feeding samples taken at 0 hours: Post-feeding values are an average of samples taken at 1,2,4 and 6 h after AM feeding

**IVDMTD (in vitro dry matter true digestibility) is based on a 24 h incubation.

[0077] Addition of Co as Cobalt Lactate Mineral Complex II had no effect ($P > 0.10$) on methane concentration but providing Co as CoCo_3 had a significant negative impact (Table 10). Ammonia-N concentrations were not affected ($P > 0.10$) by dietary treatments. When compared to the control, supplemental Co resulted in a decrease ($P < 0.01$) in culture pH (Table 6). In vitro dry matter true digestibility

(IVDMTD) of the control diet did not differ from diets supplemented with Co (Table 10). There was a source x level interaction ($P < 0.001$) for neutral detergent fiber (NDF) digestibility (Table 6). Addition of CoCo_3 at the 1.0 ppm had a negative effect on NDF digestibility in contrast to the increase in NDF digestibility observed with Cobalt Lactate Mineral Complex II when added at the 1.0 ppm level. Earlier studies reported no effect on DM and NDF digestibility when Co was added at the 5.0 or 10.0 ppm levels.

Table 11

Sample	ID	Applied Cobalt Weight (gms)	Grass Cuttings Weight (gms, 1st)	Dry Matter Fraction	Grass Dry Weight (gms, 1st)	Cobalt Found (ppm)
1	1CoSo0	0	19.94	0.1597	3.184418	0.23
2	2CoSo1	0.0025	31.06	0.1525	4.73665	1.13
3	3CoSo10	0.025	22.8	0.1618	3.68904	9.92
4	4CoSo100	0.25	16.33	0.1688	2.756504	39.3
5	5CoSo1000	2.5	5.15	0.1868	0.96202	75.9
6	6CoAer0	0	17.28	0.1673	2.890944	0.49
7	7CoAer1	7.40E-05	15.7	0.1703	2.67371	0.78
8	8CoAer10	0.00067	17.76	0.1685	2.99256	1.77
9	9CoAer100	0.008	18.56	0.1695	3.14592	5.18
10	10CoAer1000	0.069	24.03	0.1695	4.073085	44.2
11	11CoXec0	0	17.41	0.1725	3.003225	0.23
12	12CoXec1	6.70E-05	24.42	0.1694	4.136748	0.6
13	13CoXec10	0.00071	23.52	0.1662	3.909024	1.43
14	14CoXec100	0.0068	12.25	0.1808	2.2148	118
15	15CoXec1000	0.084	12.97	0.1818	2.357946	154

[0078] In Table 11, the uptake of cobalt as measured on the grass cuttings that were harvested and submitted for testing. The uptake of cobalt correlated with the application rate and that was mirrored closely at the lower levels.

What is claimed is:

1. A method of making a mineral product, comprising:
 - contacting a lactic acid and an inorganic mineral compound to form a solution;
 - reacting the solution over a period of time to provide a mineral lactate compound and one or more gases and vapors;
 - passively releasing the one or more gases or vapors to provide a rapidly soluble mineral lactate product; and
 - wherein reacting occurs solely by contacting the lactic acid and the mineral compound and the lactic acid comprises about 62% to about 76% of the solution by weight.
2. The method of claim 1, further comprising screening the rapidly soluble mineral lactate product.
3. The method of claim 1, further comprising contacting the mineral lactate product with a carrier.
4. The method of claim 1, further comprising agitating the solution.
5. The method of claim 1, wherein the inorganic mineral compound comprises one or more of cobalt carbonate, zinc oxide, cupric oxide, manganese oxide, compounds of scandium, titanium, vanadium, chromium, manganese, iron, nickel, copper, selenium and zinc.
6. The method of claim 1, wherein the lactic acid and inorganic mineral compound comprise stoichiometric amounts.
7. A method of using the mineral lactate product produced by the method of claim 1, comprising:
 - applying the mineral lactate product to a lawn, garden, pasture or field;
 - wherein the mineral lactate product includes one or more of cobalt lactate, zinc lactate, copper lactate, and manganese lactate.
8. The method of claim 7 wherein the mineral lactate product further comprises a carrier.

REPLACEMENT SHEET

9. The method of claim 7, wherein the pasture or field comprises a harvested field, bailed field, or field or pasture with crops cut.
10. The method of claim 8, wherein the carrier comprises one or more of diatomaceous earth, calcium carbonate, limestone, sugars, dextrose, water, ground corn cobs, and starch.
11. The method of claim 7, wherein applying the mineral lactate product to a lawn, garden, pasture or field improves growth of plant life, impacts the health and growth of microorganisms, enhances emergence of a seedling, and strengthens a root system of a plant.
12. A method of using the mineral lactate product produced by the method of claim 1, comprising:


administering the mineral lactate product to one or more living organisms.
13. The method of claim 12, wherein the one or more living organisms comprise monogastric mammals.
14. The method of claim 12, wherein the one or more living organisms comprise ruminant mammals.
15. The method of claim 12, wherein the mineral lactate product comprises one or more of a cobalt lactate compound, a zinc lactate compound, a copper lactate compound or a manganese lactate compound.
16. The method of claim 12, wherein the mineral lactate product further comprises a carrier.
17. The method of claim 16, wherein the carrier comprises one or more of diatomaceous earth, calcium carbonate, limestone, sugars, dextrose, water, ground corn cobs, and starch.
18. The method of claim 12, wherein administering comprises providing the mineral lactate product as a feed or a feed supplement.
19. The method of claim 12, wherein administering comprises providing the mineral lactate product to the mammals through ingestion of a plant containing the mineral lactate product.
20. The method of claim 12, wherein administering the mineral lactate product to one or more living organisms increases rumen activity in mammals.


REPLACEMENT SHEET


21. The method of claim 20, wherein increasing rumen activity comprises increasing metabolism.
22. A mineral chelated composition for treating one or more of plants, animals, and soils, comprising:
- the mineral lactate product produced by the method of claim 1;
 - yucca;
 - fiber; and
 - one or more enzymes.
23. The mineral chelated composition of claim 22, further comprising one or more metal sulfates.
24. The mineral chelated composition of claim 22, wherein the mineral lactate product comprises one or more of cobalt lactate, zinc lactate, manganese lactate, iron lactate, and copper lactate.
25. The mineral chelated composition of claim 23, wherein the one or more metal sulfates comprises sulfates of manganese, zinc, copper or combinations thereof.
26. The mineral chelated composition of claim 22, wherein the mineral lactate product comprises about 15% to about 20% of the composition by weight.
27. The mineral chelated composition of claim 23, wherein the one or more metal sulfates comprise about 2% to about 10% of the composition by weight.
28. The mineral chelated composition of claim 22, wherein the fiber comprises about 1% to about 5% of the composition by weight.
29. The mineral chelated composition of claim 22, wherein the one or more enzymes comprise about 0.1% to about 2% of the composition by weight.
30. The mineral chelated composition of claim 22, further comprising a carrier.

REPLACEMENT SHEET

31. The mineral chelated composition of claim 22, wherein the yucca comprises about 1% to about 5% of the composition by weight.
32. The method of claim 1, wherein passively releasing comprises releasing the one or more gases and vapors without initiating further reactions with a solvent or utilizing a reflux.
33. The method of claim 1, wherein the one or more gases and vapors comprise one or more of water vapor and carbon dioxide.
34. The method of claim 7, wherein the mineral lactate product further comprises one or more metal sulfates.
35. The method of claim 34, wherein the one or more metal sulfates comprises sulfates of manganese, zinc, copper or combinations thereof.
36. A method of making a mineral chelated product, comprising:
- contacting a lactic acid and a cobalt mineral compound to form a solution;
 - reacting the solution over a period of time to provide a cobalt chelated compound and one or more gases and vapors;
 - passively releasing the one or more gases and vapors to provide a rapidly soluble cobalt chelated product;
 - wherein reacting occurs solely by contacting the lactic acid and the cobalt mineral compound and the lactic acid comprises about 62% to about 76% of the solution by weight.
37. The method of claim 36, wherein passively releasing comprises releasing the one or more gases and vapors without initiating further reactions with a solvent or utilizing a reflux.
38. The method of claim 36, wherein the lactic acid and cobalt mineral compound comprise stoichiometric amounts.
39. The method of claim 36, further comprising contacting the cobalt chelated product with a carrier.

1/3**FIG. 1**

2/3**FIG. 2**

3/3**FIG. 3**

