

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 353 450 B1

(12)

EUROPEAN PATENT SPECIFICATION

- (43) Date of publication of patent specification: **29.03.95** (51) Int. Cl.⁸: **C07D 257/02, A61K 31/395,**
A61K 51/00, A61K 47/00,
A61K 49/00
- (21) Application number: **89111497.7**
- (22) Date of filing: **23.06.89**

(54) **Macrocyclic bifunctional chelants, complexes thereof and their antibody conjugates.**

- (30) Priority: **24.06.88 US 211496**
21.06.89 US 370956
- (43) Date of publication of application:
07.02.90 Bulletin 90/06
- (45) Publication of the grant of the patent:
29.03.95 Bulletin 95/13
- (84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE
- (56) References cited:
EP-A- 0 232 751 EP-A- 0 238 196
EP-A- 0 255 471 EP-A- 0 292 689
EP-A- 0 296 522 WO-A-84/03698
WO-A-89/01475 US-A- 4 678 667
- (73) Proprietor: **THE DOW CHEMICAL COMPANY**
2030 Dow Center,
Abbott Road
Midland,
Michigan 48640 (US)
- (72) Inventor: **Cheng, Roberta C.**
3873 Old Pine Trail

Midland
Michigan 48640 (US)
Inventor: **Fordyce, William A.**
4103 Washington Street
Midland
Michigan 48640 (US)
Inventor: **Goeckeler, William F.**
4916 Sturgeon Creek Pkwy
Midland
Michigan 48640 (US)
Inventor: **Kruper, William F., Jr.**
230 Barden Road
Sanford
Michigan 48657 (US)
Inventor: **Baughman, Sharon**
111 Pinestone
Irvine
California 92714 (US)
Inventor: **Garlich, Joseph R.**
301 Southern Oaks
Lake Jackson
Texas 77566 (US)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Inventor: **Kiefer, Garry**
400 Timber Creek Dr. Nr. 913
Richwood
Texas 77531 (US)

Inventor: **McMillan, Kenneth**
405 Moore Street
Richwood
Texas 77531 (US)

Inventor: **Simon, Jaime J.**
Route 1, Box 120-G
Angleton
Texas 77515 (US)

Inventor: **Wilson, David A.**
229 San Saba
Richwood
Texas 77531 (US)

Inventor: **Frank, Richard Keith**
722 Sycamore
Lake Jackson
Texas 77566 (US)

⑦ Representative: **Huber, Bernhard, Dipl.-Chem.**
et al
Patentanwälte
H. Weickmann, Dr. K. Fincke
F.A. Weickmann, B. Huber
Dr. H. Liska, Dr. J. Prechtel, Dr. B.
Böhm
Postfach 86 08 20
D-81635 München (DE)

Description

Functionalized chelants, or bifunctional coordinators, are known to be capable of being covalently attached to an antibody having specificity for cancer or tumor cell epitopes or antigens. Radionuclide complexes of such antibody/chelant conjugates are useful in diagnostic and/or therapeutic applications as a means of conveying the radionuclide to a cancer or tumor cell. See, for example, Meares et al., *Anal. Biochem.* **142**, 68-78, (1984); and Krejcarek et al., *Biochem. and Biophys. Res. Comm.* **77**, 581-585 (1977).

Aminocarboxylic acid chelating agents have been known and studied for many years. Typical of the aminocarboxylic acids are nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), hydroxyethylthylenediaminetriacetic acid (HEDTA), diethylenetriaminepentaacetic acid (DTPA), trans-1,2-diaminocyclohexanetetraacetic acid (CDTA) and 1,4,7,10-tetraazacyclododecanetetraacetic acid (DOTA). Numerous bifunctional chelating agents based on aminocarboxylic acids have been proposed and prepared. For example the cyclic dianhydride of DTPA [Hnatowich et al. *Science* **220**, 613-615, (1983); U.S. Patent 4,479,930] and mixed carboxycarbonic anhydrides of DTPA [Gansow, U.S. Patents 4,454,106 and 4,472,509; Krejcarek et al., *Biochem. and Biophys. Res. Comm.* **77**, 581-585, (1977)] have been reported. When the anhydrides are coupled to proteins the coupling proceeds via formation of an amide bond thus leaving four of the original five carboxymethyl groups on the diethylenetriamine (DETA) backbone [Hnatowich et al. *Int. J. Appl. Isot.* **33**, 327-332, (1982)]. In addition, U.S. Patents 4,432,907 and 4,352,751 disclose bifunctional chelating agents useful for binding metal ions to "organic species such as organic target molecules or antibodies." As in the above, coupling is obtained via an amide group through the utilization of diaminotetraacetic acid dianhydrides. Examples of anhydrides include dianhydrides of EDTA, CDTA, propylenediaminetetraacetic acid and phenylene 1,2-diaminetetraacetic acid. A recent U.S. Patent 4,647,447 discloses several complex salts formed from the anion of a complexing acid for use in various diagnostic techniques. Conjugation via a carboxyl group of the complexing acid is taught which gives a linkage through an amide bond.

In the *J. of Radioanalytical Chemistry* **57** (12), 553-564 (1980), Paik et al. disclose the use of p-nitrobenzylbromide in a reaction with a "blocked" diethylenetriamine, i.e. bis-(2-phthalimidoethyl)amine followed by deblocking procedures and carboxymethylation using chloroacetic acid, to give N'-p-nitrobenzyl-diethylenetriamine N,N,N',N'-tetraacetic acid. Again, since the attachment is through a nitrogen, a tetraacetic acid derivative is obtained. Conjugation of the bifunctional chelating agent and chelation with indium is discussed. Substitution on the nitrogen atom is also taught by Eckelman, et al. in the *J. of Pharm. Sci.* **64**(4), 704-706 (1975) by reacting amines such as "ethylenediamine or diethylenetriamine with the appropriate alkyl bromide before carboxymethylation." The compounds are proposed as potential radiopharmaceutical imaging agents.

Another class of bifunctional chelating agents based on aminocarboxylic acid functionality is also well documented in the literature. Thus, Sundberg, Meares, et al. in the *J. of Med. Chem.* **17**(12), 1304 (1974), disclosed bifunctional analogs of EDTA. Representative of these compounds are 1-(p-aminophenyl)-ethylenediaminetetraacetic acid and 1-(p-benzene-diazonium)-ethylenediaminetetraacetic acid. Coupling to proteins through the para-substituent and the binding of radioactive metal ions to the chelating group is discussed. The compounds are also disclosed in *Biochemical and Biophysical Research Communications* **75**(1), 149 (1977), and in U.S. Patents 3,994,966 and 4,043,998. It is important to note that the attachment of the aromatic group to the EDTA structure is through a carbon of the ethylenediamine backbone. Optically active bifunctional chelating agents based on EDTA, HEDTA and DTPA are disclosed in U.S. 4,622,420. In these compounds an alkylene group links the aromatic group (which contains the functionality needed for attachment to the protein) to the carbon of the polyamine which contains the chelating functionality. Other references to such compounds include Brechbiel et al., *Inorg. Chem.* **25**, 2772-2781 (1986), U.S. Patent 4,647,447 and International Patent Publication No. WO 86/06384.

More recently, certain macrocyclic bifunctional chelating agents and the use of their copper chelate conjugates for diagnostic or therapeutic applications have been disclosed in U.S. Patent 4,678,667 and by Moi et al., *Inorg. Chem.* **26**, 3458-3463 (1987). Attachment of the aminocarboxylic acid functionality to the rest of the bifunctional chelating molecule is through a ring carbon of the cyclic polyamine backbone. Thus, a linker, attached at one end to a ring carbon of the cyclic polyamine, is also attached at its other end to a functional group capable of reacting with the protein.

Another class of bifunctional chelating agents, also worthy of note, consists of compounds wherein the chelating moiety, i.e. the aminocarboxylic acid, of the molecule is attached through a nitrogen to the functional group of the molecule containing the moiety capable of reacting with the protein. As an example, Mikola et al. in patent application (International Publication Number WO 84/03698, published 9/27/1984) disclose a bifunctional chelating agent prepared by reacting p-nitrobenzylbromide with DETA followed by

reaction with bromoacetic acid to make the aminocarboxylic acid. The nitro group is reduced to the corresponding amine group and is then converted to the isothiocyanate group by reaction with thiophosgene. These compounds are bifunctional chelating agents capable of chelating lanthanides which can be conjugated to bio-organic molecules for use as diagnostic agents. Since attachment of the linker portion of the molecule is through one of the nitrogens of the aminocarboxylic acid, then one potential aminocarboxyl group is lost for chelation. Thus, a DETA-based bifunctional chelant containing four (not five) acid groups is prepared. In this respect, this class of bifunctional chelant is similar to those where attachment to the protein is through an amide group with subsequent loss of a carboxyl chelating group.

Recently Carney, Rogers, and Johnson disclosed (3rd. International Conference on Monoclonal Antibodies For Cancer; San Diego, California - 2/4-6/88) abstracts entitled "Absence of Intrinsically Higher Tissue Uptake from Indium-111 Labeled Antibodies: Co-administration of Indium-111 and Iodine-125 Labeled B72.3 in a Nude Mouse Model" and "Influence of Chelator Denticity on the Biodistribution of Indium-111 Labeled B72.3 Immunoconjugates in Nude Mice". The biodistribution of indium-111 complexed with an EDTA and DTPA bifunctional chelating agent is disclosed. Attachment of the aromatic ring to the EDTA/DTPA moieties is through an acetate methylene. Also at a recent meeting D.K. Johnson et al. [Florida Conf. on Chem. in Biotechnology, April 26-29 (1988), Palm Coast, FL] disclosed bifunctional derivatives of EDTA and DTPA where a p-isothiocyanatobenzyl moiety is attached at the methylene carbon of one of the carboxymethyl groups. Previously Hunt et al. in U.S. Patents 4,088,747 and 4,091,088 (1978) disclosed ethylenediaminediacetic acid (EDDA) based chelating agents wherein attachment of an aromatic ring to the EDDA moiety is through the alkylene or acetate methylene. The compounds are taught to be useful as chelates for studying hepatobiliary function. The preferred metal is technetium-99m. Indium-111 and indium-113m are also taught as useful radionuclides for imaging.

EP-A-0 292 689, published on November 30, 1988, discloses substituted 1,4,7-triscarboxymethyl-1,4,7,10-tetraazacyclododecanes and analogues, complexes thereof with rare earth metal ions and their use in the radiotherapy.

EP-A-0 296 522, published on December 28, 1988, discloses functionalized polyamine chelants, which form complexes with rhodium. The rhodium complexes can be bound to an antibody or to an antibody fragment and can be employed for therapeutical or diagnostic purposes.

EP-A-0 232 751, published on August 19, 1987, discloses tris-carboxymethyl tetraazadodecane derivatives, wherein the fourth nitrogen substitute contains an aryl-alkyl group, if necessary. The term "aryl" according to EP-A-0 232 751 refers to phenyl and substituted phenyl. Preferred substituted phenyl groups are those substituted with one, two or three halogens, hydroxyl, alkyl, alkoxy, carbamoyl or carboxyl groups. The positions of these substituents is however not disclosed.

Consequently, it would be advantageous to provide a complex that does not readily dissociate, that exhibits rapid whole body clearance except from the desired tissue, and conjugates with an antibody to produce the desired results.

Brief Description of the Drawings

Figures 1-7 and 15-21 show the biodistribution of ^{153}Sm as administered as a conjugate containing ^{153}Sm of the present invention. The antibody, CC₄₉-IgG, was used in the conjugate of the present invention. The biodistribution was determined in nude mice bearing LS 174-T tumor.

Figures 8-14 and 22-28 show the biodistribution of ^{153}Sm as administered as a conjugate containing ^{153}Sm of the present invention. The conjugate of the present invention used CC₄₉-F(ab')₂ as the antibody fragment. The biodistribution was determined in nude mice bearing LS 174-T tumor.

Figures 29-34 show the biodistribution of ^{177}Lu as a conjugate containing ^{177}Lu (PA-DOTMA). The conjugate used CC₄₉-IgG as the antibody. The biodistribution was determined in nude mice (blab/C) bearing LS 174-T tumor.

Surprisingly, the complexes and/or conjugates of the invention are relatively stable (i.e. do not easily dissociate) and some display rapid clearance from the whole body except from the desired tissue.


The invention includes the design and synthesis of novel bifunctional chelants, each containing a chelating functionality, and a chemically reactive group for covalent attachment to biomolecules. Also forming part of the invention are methods for preparing various bifunctional coordinator (BFC)-metal complexes and the linking of the complexes to antibody to prepare radionuclide (such as samarium-153, lutetium-177 and yttrium-90) labeled antibody and/or fragments suitable for diagnostic and/or therapeutic applications.

The present invention is directed to novel bifunctional chelating agents that form complexes with metal ions, especially "radioactive" metal ions having rare earth-type chemistry. Preferred rare earth-type metal

ions include La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y and Sc. Preferred radioactive rare earth-type metal ions include ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc , and ^{142}Pr . Other radioactive metal ions which may be of interest are ^{47}Sc , $^{99\text{m}}\text{Tc}$, ^{186}Re , ^{188}Re , ^{97}Ru , ^{105}Rh , ^{109}Pd , ^{197}Pt , ^{67}Cu , ^{198}Au , ^{199}Au , ^{67}Ga , ^{68}Ga , ^{111}In , $^{113\text{m}}\text{In}$, $^{115\text{m}}\text{In}$, $^{117\text{m}}\text{Sn}$, and $^{212}\text{Pb}/^{212}\text{Bi}$. The complexes so formed can be attached (covalently bonded) to an antibody or fragment thereof and used for therapeutic and/or diagnostic purposes. The complexes and/or conjugates can be formulated for *in vivo* or *in vitro* uses. A preferred use of the formulated conjugates is the treatment of cancer in animals, especially humans.

Uses of the complexes and/or conjugates of this invention which contain a non-radioactive metal for diagnosis and/or treatment of disease states such as cancer are also possible. Such uses are known for non-radioactive metals using radio frequency to induce hyperthermia (Jpn. Kokai Tokkyo Koho JP 61,158,931) and fluorescent-immunoguided therapy (FIGS) [K. Pettersson et al., *Clinical Chemistry* 29 (1), 60-64 (1983) and C. Meares et al., *Acc. Chem. Res.* 17, 202-209 (1984)].

A subject-matter of the present invention is a compound of the formula:



wherein:

each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or C_1 - C_4 alkyl;

with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen, and that when Q^1 is

hydrogen, R^3 must be other than hydrogen;

each R^5 independently is hydrogen or C_1 - C_4 alkyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

p = 1 or 2;

r = 0 or 1;

w = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R^2 is hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido or carboxyl;

R^3 is C_1 - C_4 alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, hydroxy or hydrogen;

R^4 is hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

Preferred compounds of formula (IA) are subject-matter of claims 2-35.

Preferred features of the compounds of formula (IA) are those where: R is hydrogen; R^5 is H or methyl; n is 0; m is 0 through 5; r is 0.

A further subject-matter of the present invention is a complex comprising a compound as defined above or a pharmaceutically acceptable salt thereof, complexed with an ion of a metal selected from La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y and Sc. Preferred complexes of the present invention

are subject-matter of the claims 37-60.

Yet another subject-matter of the present invention is a conjugate comprising a complex as defined above with the proviso that R^2 and R^4 cannot be nitro, and covalently attached to an antibody or antibody fragment. Preferred conjugates are subject-matter of the claims 62-70.

Yet a further subject-matter of the present invention is a conjugate comprising a compound as defined above or a pharmaceutically acceptable salt thereof, covalently attached to an antibody or antibody fragment. Preferred conjugates are subject matter of the claims 72-75.

Yet a further subject-matter of the present invention is a process for preparing a compound as defined above according to claim 76.

Still a further subject-matter of the present invention is a process for preparing a mono-N-alkylated-polyazamacrocyclic according to claims 77 and 78.

When a conjugate of the present invention is desired R^2 and R^4 must be other than nitro. When R^2 or R^4 is nitro, then a precursor of the linker moiety (L) is present. This precursor moiety can be any moiety which is formed for R^2 or R^4 for the preparation of the compounds of formula (I) and which does not bind to an antibody or antibody fragment.

The present invention is also directed to rare-earth type metal ion complexes, especially radioactive neutral or charged rare-earth type metal ion complexes, and to conjugates formed with the aforementioned complexes and antibody or antibody fragments. In addition the present invention also includes formulations having the conjugates of the invention and a pharmaceutically acceptable carrier, especially formulations where the pharmaceutically acceptable carrier is a liquid. The invention also includes a method for the diagnosis or treatment of a disease state, especially cancer, in a mammal which comprises administering to the mammal an effective amount of the formulation.

Detailed Description of the Invention

As used herein, the following indicated terms have these meanings: with respect to the definition of R^1 , R^2 or R^4 , "electrophilic" moieties include, but are not limited to, isothiocyanate, bromoacetamide, mal-eimide, imidoester, thiophthalimide, N-hydroxysuccinimyl ester, pyridyl disulfide, and phenyl azide; suitable "nucleophilic" moieties include, but are not limited to, carboxyl, amino, acyl hydrazide, semicarbazide, and thiosemicarbazide; "synthetic linkers" include any synthetic organic or inorganic linkers which are capable of being covalently attached to an antibody or antibody fragment, preferred synthetic linkers are biodegradable synthetic linkers which are stable in the serum of a patient but which have a potential for cleavage within an organ of clearance for the radioisotope, for example biodegradable peptides or peptide containing groups. Of the electrophilic moieties isothiocyanate is preferred and of the nucleophilic moieties amino, semicarbazide and thiosemicarbazide are preferred. It is desirable that the nature and/or position of R^1 be such that it does not appreciably interfere with the chelation reaction.

The "X-C-Y" term in formula (I) represents the optional presence of a double or triple bond between adjacent carbon atoms. The unsaturated bonds may be present independently in the chain length of 0 to 10 carbon atoms inclusive as defined by the "m" term in formula (I).

As used herein, the term "mammal" means animals that nourish their young with milk secreted by mammary glands, preferably warm blooded mammals, more preferably humans. "Antibody" refers to any polyclonal, monoclonal, chimeric antibody or heteroantibody, preferably a monoclonal antibody; "antibody fragment" includes Fab fragments and $F(ab')_2$ fragments, and any portion of an antibody having specificity toward a desired epitope or epitopes. When using the term "metal chelate/antibody conjugate" or "conjugate", the "antibody" portion is meant to include whole antibodies and/or antibody fragments, including semisynthetic or genetically engineered variants thereof.


As used herein, "complex" refers to a compound of the invention, e.g. formula (I), complexed with a rare-earth type metal ion, especially a radioactive rare-earth type metal ion, wherein at least one metal atom is chelated or sequestered; "radioactive metal ion chelate/antibody conjugate" or "radioactive metal ion conjugate" refers to a radioactive metal ion conjugate that is covalently attached to an antibody or antibody fragment; "radioactive" when used in conjunction with the word "metal ion" refers to one or more isotopes of the rare-earth type elements that emit particles and/or photons, such as ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{158}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc , and ^{142}Pr ; the terms "bifunctional coordinator", "bifunctional chelating agent" and "functionalized chelant" are used interchangeably and refer to compounds that have a chelant moiety capable of chelating a metal ion and a linker/spacer moiety covalently bonded to the chelant moiety that is capable of serving as a means to covalently attach to an antibody or antibody fragment.

As used herein, "pharmaceutically acceptable salt" means any salt of a compound of formula (I) which is sufficiently non-toxic to be useful in therapy or diagnosis of mammals. Thus, the salts are useful in

accordance with this invention. Representative of those salts, which are formed by standard reactions, from both organic and inorganic sources include, for example, sulfuric, hydrochloric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, palmoic, mucic, glutamic, d-camphoric, glutaric, glycolic, phthalic, tartaric, formic, lauric, steric, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic, cinnamic acids and other suitable acids. Also included are salts formed by standard reactions from both organic and inorganic sources such as ammonium, alkali metal ions, alkaline earth metal ions, and other similar ions. Particularly preferred are the salts of the compounds of formula (I) where the salt is potassium, sodium, ammonium, or mixtures thereof.

Of course, the free acid of the compounds of formula (I) may be used, also the protonated form of the compounds, for example when the carboxylate is protonated and/or the nitrogen atoms i.e. when the HCl salt is formed.

Preferred compounds of formula (I) include those compounds where Q' is hydrogen and L is represented by formula A as shown by the following formula:



wherein:

each Q is independently hydrogen or $\text{CHR}^5\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least two of Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;

R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;


R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, hydroxy and hydrogen;

R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;

each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

In formula (II) when R^3 and R^4 are both hydrogen, the compounds are represented by the following formula:



wherein:

each Q is independently hydrogen or $\text{CHR}^5\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least two of Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;


R^2 is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;

each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;

or

a pharmaceutically acceptable salt thereof.

In formula (II) when R^2 is hydrogen, the compounds are represented by the formula:



wherein:

each Q is independently hydrogen or $\text{CHR}^5\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least two of Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;

R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$ and hydroxy;


R^4 is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido, and maleimido;

each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;

or

a pharmaceutically acceptable salt thereof.

Additional preferred compounds of formula (I) include those compounds where at least one Q is hydrogen and are represented by the formula:



wherein:

each Q is independently hydrogen or $\text{CHR}^5\text{CO}_2\text{R}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl;
with the proviso that at least two of the sum of Q and Q' must be other than hydrogen and one of Q is hydrogen;

m is an integer from 0 to 5 inclusive;

w is 0 or 1;

R² is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, maleimido and bromoacetamido;


R³ is selected from the group consisting of C₁-C₄ alkoxy, -OCH₂CO₂H, hydroxy and hydrogen;

R⁴ is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, maleimido and bromoacetamido;

each R⁵ independently is hydrogen or C₁-C₄ alkyl;

with the proviso that R² and R⁴ cannot both be hydrogen but one of R² and R⁴ must be hydrogen; or a pharmaceutically acceptable salt thereof.

Other preferred compounds of formula (I) include compounds where Q' is CO₂R (w=0) and are represented by the formula:



wherein:

each Q is independently hydrogen or CHR⁵CO₂R;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl;

with the proviso that at least one Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;

R² is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, maleimido and bromoacetamido;


R³ is selected from the group consisting of C₁-C₄ alkoxy, -OCH₂CO₂H, hydroxy and hydrogen;

R⁴ is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, maleimido and bromoacetamido;

each R⁵ independently is hydrogen or C₁-C₄ alkyl;

with the proviso that R² and R⁴ cannot both be hydrogen but one of R² and R⁴ must be hydrogen; or a pharmaceutically acceptable salt thereof.

Some preferred compounds of formula (VI) are those where R³ and R⁴ are both hydrogen, the compounds are represented by the formula:



(VII)

wherein:

each Q is independently hydrogen or CHR^5CO_2R ;

each R independently is hydrogen, benzyl or C_1-C_4 alkyl;

with the proviso that at least one Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;


R^2 is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;

each R^5 independently is hydrogen or C_1-C_4 alkyl;

or

a pharmaceutically acceptable salt thereof.

Other preferred compounds of formula (VI) are those where R^2 is hydrogen and are represented by the formula:



(VIII)

wherein:

each Q is independently hydrogen or CHR^5CO_2R ;

each R independently is hydrogen, benzyl or C_1-C_4 alkyl;

with the proviso that at least one Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;

R^3 is selected from the group consisting of C_1-C_4 alkoxy, $-OCH_2CO_2H$, and hydroxy;

R^4 is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, maleimido, and bromoacetamido;

each R^5 independently is hydrogen or C_1-C_4 alkyl;

or

a pharmaceutically acceptable salt thereof.

The bifunctional chelating agents described herein [represented by any one of formulas (I)-(VIII)] can be used to chelate or sequester the rare-earth type metal ions, particularly radioactive rare-earth type metal ions, so as to form metal ion chelates (also referred to herein as "complexes"). The complexes, because of the presence of the functionalizing moiety [represented by " R^1 " in formula (I)], can be attached to functionalized supports, such as functionalized polymeric supports, or preferably, when a complex of

formula (I) where L represents formula (A) and R^2 and R^4 must be other than nitro, can be covalently attached to proteins or more specifically to antibodies or antibody fragments. Thus the complexes described herein (represented by any one of formulas I-VIII complexed with rare-earth type metal ions, particularly radioactive rare-earth type metal ions) may be covalently attached to an antibody or antibody fragment and are referred to herein as "conjugates".

The antibodies or antibody fragments which may be used in the conjugates described herein can be prepared by techniques well known in the art. Highly specific monoclonal antibodies can be produced by hybridization techniques well known in the art, see for example, Kohler and Milstein [*Nature* 256, 495-497 (1975); and *Eur. J. Immunol.* 6, 511-519 (1976)]. Such antibodies normally have a highly specific reactivity. In the radioactive metal ion conjugates, antibodies directed against any desired antigen or hapten may be used. Preferably the antibodies which are used in the radioactive metal ion conjugates are monoclonal antibodies, or fragments thereof having high specificity for a desired epitope(s). Antibodies used in the present invention may be directed against, for example, tumors, bacteria, fungi, viruses, parasites, mycoplasma, differentiation and other cell membrane antigens, pathogen surface antigens, toxins, enzymes, allergens, drugs and any biologically active molecules. Some examples of antibodies or antibody fragments are CC-11, CC-46, CC-49, CC-49 F(ab')₂, CC-83, CC-83 F(ab')₂, and B72.3. [See D. Colcher et al., *Cancer Res.* 48, 4597-4603 (Aug. 15, 1988) for CC-49, CC-83 and B72.3 antibodies. The hybridoma cell line B72.3 is deposited in the American Type Culture Collection (ATCC) having the accession number HB 8108. The various CC antibodies are disclosed in United States patent application 7-073,685, filed July 15, 1987, which is available through NTIS. The other murine monoclonal antibodies bind to epitopes of TAG-72, a tumor associated antigen.] A more complete list of antigens can be found in U.S. Patent 4,193,983.

The radioactive metal ion conjugates of the present invention are particularly preferred for the diagnosis and treatment of various cancers.

The preferred rare-earth type (lanthanide or pseudo-lanthanide) complexes of the present invention are represented by the formula:



wherein: Ln is a rare-earth metal (lanthanide) ion, such as Ce³⁺, Pr³⁺, Nd³⁺, Pm³⁺, Sm³⁺, Eu³⁺, Gd³⁺, Tb³⁺, Dy³⁺, Ho³⁺, Er³⁺, Tm³⁺, Yb³⁺ and Lu³⁺, or pseudo-lanthanide metal ion, such as Sc³⁺, Y³⁺ and La³⁺; BFC represents a bifunctional chelant; and C represents a pharmaceutically acceptable ion or group of ions of sufficient charge to render the entire complex neutral. If the BFC contains four or more negatively charged moieties, then C is a cation or group of cations such as H⁺, Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺, Mg²⁺, Ca²⁺, Sr²⁺, Ba²⁺, NH₄⁺, N(CH₃)₄⁺, N(C₂H₅)₄⁺, N(C₃H₇)₄⁺, N(C₄H₉)₄⁺, As(C₆H₅)₄⁺, [(C₆H₅)₃P =]₂N⁺ and other protonated amines. If the BFC contains three negatively charged moieties, then C is not required. If the BFC contains two negatively charged moieties, then C is an anion such as F⁻, Cl⁻, Br⁻, I⁻, ClO₄⁻, BF₄⁻, H₂PO₄⁻, HCO₃⁻, HCO₂⁻, CH₃SO₃⁻, H₃C-C₆H₄-SO₃⁻, PF₆⁻, CH₃CO₂⁻ and B(C₆H₅)₄⁻.

The conjugates of this invention, and in some instances the complexes of this invention, may be employed as a formulation. The formulation comprises a compound of formula (I) with the antibody and/or metal ion and a physiologically acceptable carrier, excipient or vehicle therefore. Thus, the formulation may consist of a physiologically acceptable carrier with a complex (metal ion + ligand), conjugate (metal ion + ligand + antibody) or (ligand + antibody). The methods for preparing such formulations are well known. The formulation may be in the form of a suspension, injectable solution or other suitable formulation. Physiologically acceptable suspending media, with or without adjuvants, may be used.

The formulations of the present invention are in the solid or liquid form containing the active radionuclide complexed with the ligand. These formulations may be in kit form such that the two components (i.e. ligand and metal, complex and antibody, or ligand/antibody and metal) are mixed at the appropriate time prior to use. Whether premixed or as a kit, the formulations usually require a pharmaceutically acceptable carrier.

Injectable compositions of the present invention may be either in suspension or solution form. In the preparation of suitable formulations it will be recognized that, in general, the water solubility of the salt is greater than the acid form. In solution form the complex (or when desired the separate components) is dissolved in a physiologically acceptable carrier. Such carriers comprise a suitable solvent, preservatives such as benzyl alcohol, if needed, and buffers. Useful solvents include, for example, water, aqueous alcohols, glycols, and phosphonate or carbonate esters. Such aqueous solutions contain no more than 50 percent of the organic solvent by volume.

Injectable suspensions are compositions of the present invention that require a liquid suspending medium, with or without adjuvants, as a carrier. The suspending medium can be, for example, aqueous

polyvinylpyrrolidone, inert oils such as vegetable oils or highly refined mineral oils, or aqueous carboxymethylcellulose. Suitable physiologically acceptable adjuvants, if necessary to keep the complex in suspension, may be chosen from among thickeners such as carboxymethylcellulose, polyvinylpyrrolidone, gelatin, and the alginates. Many surfactants are also useful as suspending agents, for example, lecithin, alkylphenol, polyethylene oxide adducts, naphthalenesulfonates, alkylbenzenesulfonates, and the polyoxyethylene sorbitan esters.

Many substances which effect the hydrophilicity, density, and surface tension of the liquid suspension medium can assist in making injectable suspensions in individual cases. For example, silicone antifoams, sorbitol, and sugars are all useful suspending agents.

An "effective amount" of the formulation is used for therapy. The dose will vary depending on the disease being treated. Although *in vitro* diagnostics can be performed with the formulations of this invention, *in vivo* diagnostics are also contemplated using formulations of this invention. The conjugates and formulations of this invention can also be used in radioimmuno guided surgery (RIGS); however, other metals which could be used for this purpose also include ^{99m}Tc , ^{111}In , ^{113m}In , ^{67}Ga and ^{68}Ga .

Other uses of some of the chelants of the present invention may include magnetic resonance imaging, e.g. the complexes of formula I, especially complexes of formula VI, with Gd^{+3} , attachment to polymeric supports for various purposes, e.g. as diagnostic agents, and removal of lanthanide metal or pseudolanthanide metal ion by selective extraction.

The present invention provides chelants, complexes, and antibody conjugates some of which are more stable, and/or have improved biodistribution, and/or have more rapid clearance from the body, than those known in the art.

Detailed Description of the Process

A reasonable and general synthetic approach to a twelve-membered macrocyclic, bifunctional chelant of the present invention as represented by formula (I) involves monofunctionalization of a free-base macrocycle (e.g. 1,4,7,10-tetraazacyclododecane) at only one of the nitrogen atoms with an appropriate electrophile (eg. any appropriately substituted α -halocarboxylic acid ester). This electrophile must possess a suitable linker moiety or suitably protected linker moiety which would allow covalent attachment of the bifunctional ligand to a protein, antibody or antibody fragment.

It is recognized in the art that alkylative techniques for the production of mono-N-functional polyazamacrocycles may result in mixtures of difficult to separate products [T. Kaden, *Top. Curr. Chem.* **121**, 157-75 (1984)]. This problem has been overcome through the reported use of large excesses of macrocycle (5-10 equivalents relative to electrophile) which favors formation of the monoalkylation adduct [M. Studer and T.A. Kaden, *Helv.Chim.Acta* **69**, 2081-86 (1986); E. Kimura et al., *J.Chem.Soc.Chem.Comm.* 1158-59 (1986)].

Other routes to mono-N-functional polyazamacrocycles involve lengthy protection, functionalization, and deprotection schemes [P. S. Pallavicini et al., *J.Amer. Chem.Soc.* **109**, 5139-44 (1987); M. F. Tweedle et al., *Eur. Pat.Appl.No.* 0232-751]. A recent report of a reductive amination of substituted phenyl pyruvic acids and amines has issued by Abbott Labs. as an Abstract from a recent meeting (D. K. Johnson et al., Florida Conf. on Chem. in Biotechnology, April 26-29 (1988), Palm Coast, FL). A process that prepares mono-N-alkylated polyazamacrocycles by using an electrophile with between about one and five equivalents of a suitable macrocycle in a solvent which will not promote a proton transfer is disclosed in EP-A-0 374 929, Published on June 27, 1990.

General synthetic routes to the chelants of the present invention are disclosed in Synthesis Schemes I-IV hereinafter and involve the reaction of a suitable electrophile and the polyazamacrocycle in an appropriate organic solvent. Examples of suitable organic solvents are any hydrocarbon which supports solubility, such as acetonitrile, isopropanol, methylene chloride, toluene, chloroform, n-butanol, carbon tetrachloride, tetrahydrofuran, 5% ethanol in chloroform, with the most preferred being chloroform, methylene chloride, n-butanol, 1,4-dioxane and acetonitrile. Various stoichiometric amounts of reactants, temperatures and concentrations can be used. Nearly stoichiometric amounts of macrocycle and electrophile may be employed and yield the corresponding mono-N-alkylation product in a single step. The temperature range for the reaction is from about 0°C to about reflux, preferably from 0°C to 25°C. The time of the reaction is until completion about 1 to about 24 hours.

General access to substituted α -haloacid esters is desirable for the overall viability of Synthesis Schemes II and IV. One suitable approach involves bromination or chlorination of the acid halide generated *in situ*, e.g. D.N. Harpp et al., *J. Org. Chem.* **40**, 3420-27 (1975). This approach allows for exclusive alpha halogenation of alkanic acids which contain even reactive benzylic groups. A general method to substituted

acid halides involves reaction of the organic acid with thionyl chloride or sulfuryl chloride, e.g. E. Schwenk et al. *J. Amer. Chem. Soc.* 70, 3626-27 (1944). Both methods utilize the free carboxylic acid which is frequently available from commercial sources.


Polyazamacrocycles such as 1,4,7,10-tetraazacyclododecane may be prepared by documented methods such as T.J. Atkins et al., *J. Amer. Chem. Soc.* 96, 2268-70 (1974) and T.J. Richman et al., *Org. Synthesis* 58, 86-98 (1978).

Carboxymethylation of the mono-N-functional macrocycle may be performed by the method of Desreux using bromoacetic acid derivatives and a suitable base [J.F. Desreux, *Inorg. Chem.* 19, 1319-24 (1980)].

All of the starting materials required for preparing the compounds of this invention are either available from commercial sources or can be made from a known literature reference descriptions.


In the following Scheme I, the compounds of formula (I) are prepared where Q¹ is hydrogen. Although only one compound is indicated by the terms shown, other similar moieties within formula (I) where Q¹ is hydrogen, r = 0 or 1, n = 0 or 1, and m = 0 through 10 can also be prepared by this method.

Scheme I




In the following Scheme II, the compounds of formula (I) are prepared where Q' is hydrogen. Although only one compound is indicated by the terms shown, other similar moieties within formula (I) where Q' is hydrogen, $r = 0$ or 1 , $n = 0$ or 1 and $m = 0$ through 10 can also be prepared by this method.

Scheme II



In the following Scheme III, the compounds of formula (I) are prepared where Q^1 is CO_2R . Although only one compound is indicated by the terms shown, other similar moieties within formula (I) where Q^1 is CO_2R , including $m=0$ through 10, $n=0$ or 1 and $r=0$ can also be prepared by this method.

Scheme III




In the following Scheme IV, the compounds of formula (I) are prepared where Q' is $(\text{CHR}^5)_w\text{CO}_2\text{R}$. Although only one compound is indicated by the terms shown, other similar moieties within formula (I) where Q' is $(\text{CHR}^5)_w\text{CO}_2\text{R}$, including $m=0$ through 10, $n=0$ or 1 and $r=0$ can also be prepared by this method.

Use of an optically active alkylating agent has minimized the diastereomers and simplifies the synthesis. Isolation of a single diastereomer which would give rise to a single, easily purified lanthanide

complex would be desirable for radiochelation as well as antibody conjugation.


Scheme IV



In the following Scheme V, the compounds of formula (I) are prepared where Q¹ is hydrogen or (CHR⁵)_wCO₂R and R² is hydrogen. Although only two compounds are indicated by the terms shown, other similar moieties within formula (I) where Q¹ is hydrogen or (CHR⁵)_wCO₂R, Q is hydrogen or (CHR⁵)_pCO₂R, r = 0 or 1, n = 0 or 1, w = 0 or 1, m = 0 through 10, p = 1 or 2, L represents formula A wherein R³ is defined as before, R² and R⁴ are selected from the group consisting of hydrogen, amino, nitro, isothiocyanato,

semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, can also be prepared by this method.

Scheme V



The electrophilic moiety ("R'" in the formula) can also be prepared by methods known to the art. Such methods may be found in *Acc.Chem.Res.* 17, 202-209 (1984). A process that can prepare compounds of formula (I) where R' could be an isothiocyanato moiety (R² or R⁴ present are isothiocyanato when L = formula A) involves reacting the amino chelate (R² or R⁴ equal amino, present when L = formula A) with thiophosgene and is disclosed in EP-A-0 374 947, published on June 27, 1990.

Radionuclides can be produced in several ways. In a nuclear reactor a nuclide is bombarded with neutrons to obtain a radionuclide, e.g.

Sm-152 + neutron → Sm-153 + gamma

Another method of obtaining radionuclides is to bombard nuclides with particles in a linear accelerator or a cyclotron. Yet another way is to isolate the radionuclide from a mixture of fission products. The method of obtaining the nuclides employed in the present invention is not critical thereto.

The conjugates of the present invention can be prepared by first forming the complex and then binding the antibody or antibody fragment. Thus the process involves preparing or obtaining the ligand, forming the complex with the metal and then adding the antibody. Alternatively, a process for making labeled antibody conjugates can involve first conjugation of the BFC to the antibody and its subsequent chelation to yield the radionuclide-BFC labeled Ab. Any suitable process that results in the formation of the conjugates of this invention is within the scope of this invention.

In the following examples, the following terms and conditions were used unless otherwise specified.

General Experimental

Mass spectra were obtained on either a Finnigan TSQ mass spectrometer (Q¹MS mode) or a VG ZAB-MS high resolution mass spectrometer (fast atom bombardment with xenon, using 3:1 dithiothreitol:dithioerythritol).

¹H and ¹³C NMR spectra were obtained using a Varian VXR-300, Bruker APC 300, IBM/Bruker NR-80 or a Jeol FX400 spectrometer. All spectra were obtained at 30 °C unless otherwise noted. ¹H NMR was done at 300 MHz, 80 MHz or 400 MHz, respectively to the equipment listed above; ¹³C NMR was done at 75 MHz, 20 MHz or 100 MHz, respectively to the equipment listed above. The values for the NMR are δ versus TMS (tetramethylsilane) or when D₂O was the solvent versus DSS (2,2-dimethyl-2-silapentane-5-sulfonic acid, sodium salt).

Infrared spectra (IR) were recorded on a Nicolet 5SX FT/IR instrument.

For the chromatography procedures, most solvents were Fisher HPLC grade materials. Ammonium acetate was purchased from Aldrich. Water was purified using a Barnstead NANOpure™ water filtration system. Preparative chromatography of organic compounds was performed either by normal gravity chromatography using standard techniques or by flash chromatography as described by C.W. Still et al., *J. Org. Chem.* **43**, 2923-24 (1978). The following solvent systems were used:

Solvent System	Components
1	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 2:2:1 V:V:V
2	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 12:4:1 V:V:V
3	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 16:4:1 V:V:V
4	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 4:2:1 V:V:V
5	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 3:2:1 V:V:V
6	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 7:3:1 V:V:V
7	saline (0.85% of NaCl in distilled water):conc. NH ₄ OH 4:1 V:V
8	CHCl ₃ :CH ₃ OH:conc. NH ₄ OH 4:4:1 V:V:V

R_f values are reported using these solvent systems and commercially available, normal phase, silica TLC plates [GHLF 250 micron, Analtech Inc. or Merck Kiesel gel 60F₂₅₄] Preparative column chromatography was done using Merck grade 60, 60 Å silica gel.

All percentages are by weight unless stated otherwise.

Some solids were dried using a rotary evaporator (Buchi 461) and/or a vacuum oven at a temperature of about 55-60 °C for several hours. In addition, a Virtis model 10-010 automatic freezer dryer or Speed Vac™ concentrator was used for solvent removal.

Samarium-153 and lutetium-177 were produced by the Research Reactor, University of Missouri (Columbia, MO). Yttrium-90 was purchased from Oak Ridge National Laboratory.

1-(4-Isothiocyanatobenzyl)diethylenetriaminepentaacetic acid (SCN-Bz-DTPA) was prepared by a modification to the procedure of M. W. Brechbiel, et al., *Inorg. Chem.* **25**, 2772-2781 (1986), and purified to give a single species on anion exchange HPLC (Q-Sepharose™). Some of the chemicals used were obtained from the sources indicated: N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES), free acid and sodium salt were purchased from Behring Diagnostics (La Jolla, CA); sodium citrate (certified) was from Fisher Scientific; thiophosgene was from Aldrich Chemicals; ammonium acetate, sodium acetate, ethylenediaminetetraacetic acid (EDTA) and phosphate buffered saline (PBS) were from Sigma Diagnostics.

HPLC columns used were: Hand-packed Q-Sepharose™ (Pharmacia) either 1.5 cm x 25 cm or 2.5 cm x 25 cm; Zorbax™ BIO Series GF-250 (9.4 mm x 25 cm) from Du Pont Instruments; Vydac™ (Trademark of the Separations Group, Hesperia, CA) protein C-4 (4.6 mm x 25 cm) from the Separation Group (Hesperia, CA); and Mono-Q™ and SP-Sephadex™ (Tradename of Pharmacia Biotechnology Products) from Pharmacia. Sep-Pak™ (Tradename of Waters Associates) C-18 cartridge was purchased from Waters Associates (Milford, MA), Sephadex™ G-25 disposable columns (2.2 ml) from Isolab Inc. (Akron, OH), and Centricon™ -30 (Tradename of Amicon Division, W. R. Grace & Co., Danvers, MA) microconcentrators from Amicon.

Five HPLC systems were used for analyses and sample separations:

System I consisted of LKB 2150 pump, and 2152 controller, a UV detector - LKB 2238 UV Cord, a Berthold LB 506 A HPLC Radioactivity Monitor (of the International Berthold Group) and a Gilson Fraction collector 201-202 (Gilson International, Middleton, WI);

System II was equipped with an auto-sampler, WISP 710 B, two pumps (model 510) and an automated gradient controller (of Waters Associates), a Perkin-Elmer LC-75 spectrophotometric detector, a Beckman model 170 Radioisotope Detector, and a fraction collector (Frac-100 of Pharmacia);

System III consisted of two Waters M-6000A pumps with a Waters 660 solvent Programmer, a Waters Model 481 Variable Wavelength Detector, and a Waters Data Module, also an ISCO fraction collector was used preparatively;

System IV was Dionex BioLC™ System equipped with a variable wavelength UV detector; and

System V was a Waters Model 590 Pump with an ISCO Model 2360 gradient programmer and a Dionex variable wavelength detector.

For centrifugation and concentration, a Sorvall RT 6000B (refrigerated centrifuge of Du Pont) was used. A Speed Vac concentrator (Savant Instruments Inc., Hicksville, N.Y.) was employed for removal of volatile solvents. Radioactivity measurements were done on a Capintec™ Radioisotope Calibrator (Trademark of Capintec Inc.) CRC-12, or by liquid scintillation on Beckman LS 9800 (Beckman Instruments, Inc., Irvine, CA), or on a Canberra 35+ multichannel analyzer interfaced with a 7.62 cm (3 in.) thallium drifted sodium iodide well crystal.

In the examples concerning formation of complex, the percent complex determination was performed by the following general method where indicated.

Percent Complex Determination by Cation Exchange

A 10 ml plastic column was filled with 1 to 2 ml of SP-Sephadex™ C-25 resin which was swelled in water. The excess water was removed by applying pressure to the top of the column. The test solution (15 μ l) was added to the top of the resin, then 2 ml of 4:1 (V:V) of isotonic saline: conc. NH_4OH was used as an eluent. The eluent was collected in a counting tube. An additional 2 ml of eluent was used. After collection of eluent into a second counting tube, air pressure was applied to the top of the resin to bring it to dryness. The dried resin was transferred to a third counting tube. The activity of each counting tube was measured with a NaI well counter coupled to a Canberra multichannel analyzer. The amount of the metal as a complex was given as the percentage of the total activity that was found in the eluent. The amount of metal in the free uncomplexed state was given as the percentage of the total metal that remains in the resin.

Removal of Uncomplexed Metal by Ion Exchange

A 10 ml plastic column was filled with 0.5 ml of SP-Sephadex™ C-25 resin swelled in water. The excess water was removed by centrifuging the column at 3,000 rpm for 4 min. A volume of 200-500 μ l of the complex solution was placed at the top of the resin and the tube centrifuged again for 4 minutes at 3,000 rpm. The uncomplexed metal remained in the column and the complexed metal eluted with the solvent.

Yttrium Complex Preparation:

Complexes were made by preparing a 3×10^{-4} M yttrium solution in water ($\text{YCl}_3 \cdot 6\text{H}_2\text{O}$, 303.26 g/mole; or $\text{Y}(\text{OAc})_3$, 12.1% H_2O). Radioactive Y^{3+} solution (Oakridge National Laboratories) was added to give the desired level of radioactivity. Ten μ l of ligand solution (at 0.03M) was added to 990 μ l of the Y^{3+} solution, giving a 1:1 molar ratio of ligand:metal. Ten times the amount of ligand solution was used for a 10:1 ligand to metal ratio. The pH was adjusted to 7.4 using microliter quantities of HCl or NaOH. The solution was then tested for the amount of complexed yttrium using the cation exchange method given above.

Samarium Complex Preparation:

Samarium complexes were formed as described above for yttrium complexes except that 3×10^{-4} M samarium was prepared by dissolution of Sm_2O_3 (348.7 g/mole) in 0.1M HCl. Radioactive Sm-153 was obtained as an about 3×10^{-4} M solution in 0.1M HCl from the University of Missouri Research Reactor, Columbia, Missouri.

The following definitions are provided for some terms that are used throughout this text.

Glossary:

Conc. means concentrated;

^-OAc means the acetate moiety, $^-OCOCH_3$;

TLC means thin layer chromatography;

DI H₂O means deionized NANOpure™ water;

NANOpure™ water is pure water obtained from a Barnstead NANOpure™ water filtration system;

Ambient temperature means room temperature or about 20 to about 25 °C;

Overnight means from about 9 to 18 hours;

LC means liquid chromatography;

NBS means N-bromosuccinimide;

MES means 2-(N-morpholino)ethanesulfonic acid ;

HPLC means high performance liquid chromatography;

PBS means phosphate buffered saline from Sigma, containing 120 mM NaCl, 2.7 mM KCl and 10 mM phosphate buffer, pH 7.4;

SP-Sephadex™ C-25 resin is a cation exchange resin having sulfonic acid functionality, sold by Pharmacia, Inc.;

rpm = revolutions per minute;

pD means pH where the hydrogen is deuterated;

DOTA = 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid;

HEPES = N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid;

BFC = bifunctional chelant;

Cyclen = 1,4,7,10-tetraazacyclododecane;

PA-DOTA = α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid;

PA-DOTMA = α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-tris-(methylacetic) acid;

BA-DOTA = α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid;

OMe-BA-DOTA = α -(5-amino-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid;

EA-DO3A = 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid;

DTPA = diethylenetriaminepentaacetic acid;

SCN-Bz-DTPA = 1-(4-isothiocyanatobenzyl)-diethylenetriaminepentaacetic acid;

EDTA = ethylenediaminetetraacetic acid;

chelant is equivalent to ligand;

complex is equivalent to chelate, e.g. a chelant, linked to a suitable metal ion;

conjugate means a chelant or complex covalently attached to an antibody or antibody fragment; and

antibodies means CC-49, CC-83 and B72.3 and their fragments such as Fab and F(ab')₂. Other possible antibodies are given hereinbefore. The hybridoma cell line B72.3 is deposited in the American Type Culture Collection, having the accession number ATCC HB 8108, and the other named murine monoclonal antibodies bind to epitopes of TAG-72, a tumor associated antigen.

The invention will be further clarified by a consideration of the following examples, which are intended to be purely exemplary of the use of the invention.

Preparation of Starting MaterialsExample A

Preparation of *d,l*-2-bromo-4-(4-nitrophenyl)butanoic acid, methyl ester.

To a solution of 5 ml of carbon tetrachloride and 15 ml (0.2 mole) of thionyl chloride was added 10.46 g (0.05 mole) of 4-(4-nitrophenyl)butanoic acid under a nitrogen atmosphere. The solution was brought to reflux for 1 hour with initial liberation of hydrogen chloride gas and sulphur dioxide gas. To the warm solution was added 11.0 g (0.06 mole) of N-bromosuccinimide in 25 ml of carbon tetrachloride and three drops of 48 percent aqueous hydrogen bromide catalyst. Bromine gas liberation was noted. The dark red solution was refluxed for 35 minutes. The solution was cooled and poured into 100 ml of methanol with stirring. TLC analysis (60:40 ethylacetate:hexane v:v) indicated a new product (R_f = 0.69, silica plates). The excess solvent was removed by rotary evaporation and the dark red oil was filtered through a flash silica gel pad (2.54 x 12.7 cm) (1 in. x 5 in.) using methylene chloride as the eluent. Evaporation of the solvent gave a

clear oil, yield 15.43 g, which was a 85:15 mixture of the titled product: methyl ester of the starting butanoic acid derivative. The titled product was characterized by:

¹H NMR (CDCl₃)

8.16(d), 7.38(d), 4.20(dd), 3.79(s), 2.88 (m), 2.38(m);

¹³C NMR (CDCl₃, 75 MHz)

169.6, 147.5, 129.3, 123.7, 53.0, 44.4, 35.5, 33.0.

Example B

Preparation of α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester.

To a stirred solution of 1.72 g (10.0 mmole) of 1,4,7,10-tetraazacyclododecane in 17 ml of pentene stabilized chloroform was added 2.07 g (5.82 mmole) of *d,l*-2-bromo-4-(4-nitrophenyl)butanoic acid, methyl ester (prepared by the procedure of Example A) over five minutes under a nitrogen atmosphere with stirring. The reaction mixture was stirred for 48 hours at about 25 °C. TLC (using Solvent System 2, on Analtech silica gel plates) indicated the formation of the title product (*R_f* = 0.73). The yellow chloroform solution was applied to a 2.54 x 40.62 cm (1 inch x 16 inch) flash silica gel column (pre-eluted with 5 percent methanolic chloroform), eluted with 250 ml of 5 percent methanolic chloroform, followed by elution with Solvent System 2. Fractions containing pure title product were combined and evaporated to provide 2.15 g (5.46 mmole, 94 percent) of the title product as a light yellow glass. Trituration of a chloroform solution of this glass with diethyl ether resulted in precipitation of a white powder (MP = 156-59 °C.) of analytical purity and characterized by:

¹H NMR (CDCl₃)

8.14(d), 7.39(d), 3.71(s), 3.39(dd), 2.5-3.0(m), 2.08(m), 2.01(m);

¹³C NMR (CDCl₃, 75 MHz)

172.7, 149.3, 146.4, 129.2, 123.6, 62.3, 51.2, 48.9, 47.2, 45.8, 45.4, 32.8, 30.9.

Example C

Preparation of 2-bromo-2-(4-nitrophenyl)ethanoic acid, methyl ester.

A mixture of *p*-nitrophenylacetic acid (25.0 g, 0.14 mole) and thionyl chloride (15.1 ml, 0.21 mole) in dry benzene (100 ml) was stirred at reflux under a N₂ pad for three hours. The mixture was then evaporated to dryness *in vacuo*. The residue was dissolved in carbon tetrachloride and stirred at reflux under a nitrogen pad. Bromine (7.2 ml, 0.14 mole) was added in small portions over a period of three days to the refluxing mixture. The reaction mixture was allowed to cool and the acid chloride was quenched with methanol (50 ml, added slowly). The resulting bromoester (27 g, 71 percent) was recovered as a yellow oil by distillation at reduced pressure (BP = 168 °C. at 1.1 mm) through a six-inch column of glass helices. The structure of the title product was confirmed by:

¹H NMR (CDCl₃)

8.25(d), 7.81(d), 5.51(s), 3.86(s).

Example D

Preparation of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester.

To 529 mg (2.068 mmole) of 2-bromo-2-(4-nitrophenyl)ethanoic acid, methyl ester (prepared by the procedure of Example C) in 20 ml of acetonitrile was added all at once a solution of 354 mg (2.068 mmole) of 1,4,7,10-tetraazacyclododecane in 20 ml of acetonitrile containing just enough methanol to effect complete solution. The resulting pink solution was stirred for 1.5 hours then concentrated *in vacuo* at 35 °C to a low volume. This crude monoestertetraamine was purified by silica gel chromatography eluting with 20 percent (wt/v) NH₄OAc/CH₃OH. The purified monoestertetraamine thus isolated as the triacetate salt was then converted to the free amine. Thus, 270 mg in 3 ml of water was treated at 0 °C with aqueous K₂CO₃ - (10 percent wt/wt) to a pH of 11. The basic solution was then extracted with 10 ml of chloroform, five times, and the chloroform layers combined, dried over sodium sulfate and concentrated to give 160 mg (42 percent yield) of the desired monoestertetraamine, characterized by NMR and fast atom bombardment mass spectrometry ([M + H]⁺ = 366) and by:

¹H NMR (CDCl₃)

8.12(d), 7.44(d), 4.75(s), 3.76(s), 2.67-2.44(m);
¹³C NMR (CDCl₃)
 171.5, 147.6, 144.0, 130.0, 124.0, 67.0 49.8, 47.9, 46.9, 46.0.

5 Example E

Preparation of N-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane.

To a stirred chloroform solution (20 ml) containing 2.9 g of 1,4,7,10-tetraazacyclododecane (16.8 mmole) was added a chloroform solution (20 ml) containing 2.1 g of 2-methoxy-5-nitrobenzyl bromide (8.5 mmole) in one portion. After stirring at room temperature for three hours the reaction mixture was filtered and the filtrate concentrated (*in vacuo*) to give a residue which was chromatographed (silica, Solvent System 3). The monoalkylated product was isolated in 79 percent yield (MP = 154-156 °C), and characterized by:

15 ¹³C NMR (CDCl₃)
 162.47, 140.63, 127.92, 125.49, 124.53, 109.91, 55.88, 53.25, 50.90, 47.18, 45.45, 45.29.

Example F

20 Preparation of N-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane.

To a 1,4-dioxane solution (20 ml) of 1,4,7,10-tetraazacyclododecane (2.3 g, 14 mmole) was added a dioxane solution (15 ml) of 2-hydroxy-5-nitrobenzyl bromide (1.2 g, 7 mmole) in one portion with constant stirring. After several minutes K₂CO₃ (1 g) was added and stirring continued for 1 hour at room temperature. The reaction mixture was then filtered and the filtrate concentrated *in vacuo* to give a yellow semi-solid which was chromatographed (column) on silica gel eluting with Solvent System 5. The desired mono-alkylated product was isolated from the last 1/3 of bright yellow eluent which also contained unreacted amine. After concentration, the yellow oil was triturated with CHCl₃ (100 ml) and filtered to remove CHCl₃-soluble amine starting material. The final product was then isolated in pure form as a yellow solid (1.2 g, 55 percent); (MP = 142-145 °C) and further characterized by:

30 ¹³C NMR (D₂O)
 25.04, 25.28, 26.83, 31.80, 36.54, 101.88, 107.22, 110.92, 111.80, 115.26, 159.99.

Example G

35 Preparation of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane.

To a stirred solution of 3.50 g (20.3 mmole) of 1,4,7,10-tetraazacyclododecane in 50 ml of pentene stabilized chloroform was added dropwise over a five minute period with vigorous stirring under a nitrogen atmosphere 4.00 g (17.4 mmole) of 1-(2-bromoethyl)-4-nitrobenzene.

Stirring was continued for about 18 hours at room temperature (about 25 °C) whereupon crystals of amine hydrobromide precipitated from solution. The entire reaction mixture was applied to a flash silica gel column (2.54 x 45.72 cm) (1 in x 18 in) which had been pre-eluted with 5 percent methanol in chloroform; 200 ml of this solution was applied as an eluent, followed by elution with Solvent System 3. From the desired product was separated 1.45 g (9.7 mmole) of p-nitrostyrene (R_f = 0.98; Solvent System 1). The desired monofunctional title product was isolated as an orange-yellow oil (2.27 g, 7.06 mmole, 40.6 percent) which solidified upon standing (R_f = 0.73, Solvent System 1).

A sample of the titled product was recrystallized from CHCl₃/cyclohexane and showed the following characteristics: MP = 146.5-148.5 °C;

50 ¹H NMR (CDCl₃)
 8.135(d, m), 7.395(d, m), 2.91(t), 2.77(t), 2.72(t), 2.50(t), 2.60(s);
¹³C NMR (CDCl₃, 75 MHz)
 148.5, 146.7, 129.6, 123.4, 55.5, 51.4, 46.9, 45.9, 45.1, 33.7.

55

Example H

Preparation of 1-(4-nitrobenzyl)-1,4,7,10-tetraazacyclododecane.

In 50 ml of chloroform was added 3.5g (20.3 mmole) of 1,4,7,10-tetraazacyclododecane and 1.7g (7.87 mmole) p-nitrobenzyl bromide and the mixture stirred under nitrogen for 24 hours at 25 °C. The chloroform slurry of hydrobromide salt was then applied to a 1 in. x 17 in. column of flash silica gel (Solvent System 3). There was obtained 2.13 g (6.93 mmole) of the title product as a pale yellow solid of analytical purity in 88 percent yield (R_f = 0.58, Solvent System 3), MP = 128-129 °C, and further characterized by:

^1H NMR (CDCl_3)

8.19 (d), 7.49 (d), 3.69 (s), 2.82 (t), 2.70 (t), 2.59 (m);

^{13}C NMR (CDCl_3 , 75 MHz)

147.2, 128.4, 123.8, 58.8, 51.7, 47.1, 46.3, 45.1.

(No Example I; to avoid confusion with later biology example numbers).

Example J

Preparation of 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane.

In 10 ml of methanol was dissolved 690 mg (2.25 mmole) of 1-(4-nitrobenzyl)-1,4,7,10-tetraazacyclododecane (prepared by the procedure of Example H). To the solution was added 350 mg of 10 percent palladium on carbon catalyst. Excess hydrogen was purged through the solution at 25 °C. Within 45 minutes TLC indicated that the title product was prepared (R_f = 0.33, Solvent System 3). Chromatographic purification of the formed titled product was conducted on 1 in. x 16 in. flash silica column (Solvent System 3) and provided 480 mg (77 percent) of the title product as a pale yellow, clear oil.

The free base title product (103 mg, 0.37 mmole) was converted to the hydrochloride salt by bubbling anhydrous hydrogen chloride through ethanol solutions of the free base. The resulting tetrahydrochloride salt of the title product, after washing with cold ethanol and drying *in vacuo*, was obtained in a yield of 147 mg (0.347 mmole), MP = 255-260 °C (dec), and was further characterized by:

^1H (D_2O , pD = 1.9)

7.56 (d), 7.47 (d), 3.95 (s), 3.28 (t), 3.23 (t), 3.09 (t), 2.96 (t);

^{13}C NMR (D_2O , pD = 1.9, 75 MHz)

138.3, 134.2, 132.2, 126.0, 58.5, 50.3, 46.7, 44.6, 44.3.

Example K

Preparation of 2-methoxy-5-nitrobenzyl nitrile.

To a solution of sodium cyanide (6 g, 121.9 mmole) in water (5.3 ml) at 70 °C was added in small portions with stirring a hot solution of 2-methoxy-5-nitrobenzyl bromide (25 g., 101.6 mmole) in ethanol (15 ml). The reaction mixture was then refluxed for 1.5 hours, cooled, and filtered. The filter cake was washed with acetonitrile and the filtrate was evaporated *in vacuo* to yield 2-methoxy-5-nitrobenzyl nitrile (19.5 g., 100 percent) as a tan solid. The product was characterized by:

^{13}C NMR (CDCl_3)

161.4, 141.0, 125.7, 124.6, 119.8, 116.5, 110.2, 56.3, 18.8.

Example L

Preparation of 2-methoxy-5-nitrophenylacetic acid.

A slurry of 2-methoxy-5-nitrobenzyl nitrile (19.5 g) (prepared by the procedure of Example K) in conc. HCl (20 ml) was refluxed for three hours. After cooling, the crude product was recovered by filtration and washed with water. The solid was taken up in hot aqueous sodium hydroxide and filtered while hot. The orange solution was cooled and acidified with HCl. The precipitate was filtered, washed with water, and dried to yield 2-methoxy-5-nitrophenylacetic acid as a white solid (15 g, 70 percent). The product was characterized by:

^{13}C NMR ($\text{CDCl}_3\text{-CD}_3\text{OD}$)

172.8, 162.5, 140.7, 126.2, 124.7, 124.1, 109.7, 55.8, 35.0.

Example M

Preparation of α -bromo-(2-methoxy-5-nitrophenyl)acetic acid, methyl ester.

To a slurry of 2-methoxy-5-nitrophenylacetic acid (2.266 g, 10.74 mmole) (prepared by the procedure of Example L) in dry benzene (50 ml) was added thionyl chloride (4.0 ml, 54.8 mmole). A drying tube and condenser were placed on the flask and the mixture was refluxed for 2 hours. The resulting yellow solution was concentrated *in vacuo* to a small volume and taken up in dry carbon tetrachloride. Bromine (0.6 ml, 11.6 mmole) was added and the solution was refluxed for two days with the exclusion of atmospheric moisture. The reaction mixture was concentrated to a small volume and methanol (50 ml) was added. After evaporation of excess methanol *in vacuo* the resulting oil was purified chromatographically (silica gel, methylene chloride-hexane 4:1). The title product (2.399 g., 74 percent) was obtained as a yellow oil. The product was characterized by:

^1H NMR (CDCl_3)

8.45(dd), 8.15(dd), 6.95 (d), 5.75(s), 3.99(s), 3.78(s).

Example N

Preparation of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane, 1-acetic acid, methyl ester.

A solution of α -bromo (2-methoxy-5-nitrophenyl)-acetic (2.399 g, 7.89 mmole) (prepared by the procedure of Example M) in chloroform (10 ml) was added to a stirred solution of 1,4,7,10-tetraazacyclododecane (2.72 g, 15.79 mmole) in chloroform (50 ml). The mixture was stirred at room temperature for 2 hours. The cloudy mixture was then concentrated to a small volume *in vacuo* at room temperature. The crude product was purified by chromatography (silica gel, Solvent System 3). The title compound was recovered as a yellow solid (2.60 g, 83 percent) and characterized by:

^{13}C NMR (CDCl_3)

170.7, 161.6, 139.9, 125.0, 124.5, 124.2, 110.2, 59.5, 55.5, 50.5, 48.0, 47.3, 45.6, 44.3, 43.5.

Example O

Preparation of *d,l*-2-bromo-4-(4-nitrophenyl)butanoic acid isopropyl ester.

4-(4-Nitrophenyl)butanoic acid (21.0 g, 0.10 mole) was added to a solution of carbon tetrachloride (10 ml) and thionyl chloride (30 ml, 0.4 mole) under a nitrogen atmosphere using the procedure of Harpp et al., *J.Org.Chem* 40, 3420-27 (1975). The solution was brought to reflux for 1 hour with initial rapid liberation of hydrogen chloride and sulfur dioxide. At this point; N-bromosuccinimide (22.0 g, 0.12 mole) was added as a solution in carbon tetrachloride (50 ml) and 8 drops of 48% aqueous hydrogen bromide catalyst was added to the warm solution whereupon bromine liberation was noted. The dark red solution was refluxed for an additional 35 minutes. The solution was cooled and poured into isopropanol (400 ml) with stirring. TLC analysis (methylene chloride, silica gel plates) revealed a new product ($R_f=0.73$). The excess solvent was removed and the dark red oil was filtered through a flash silica gel pad (7.62 x 15.24 cm) (3 in. x 6 in.) using methylene chloride as an eluent. The solvent was removed *in vacuo* and the light yellow oil was applied to a flash silica gel column (3 in. x 18 in.) and eluted with methylene chloride to afford 25.0 g (0.076 mole) of the titled bromoester product as a clear oil in 75% yield as a isopropanol solvate which contained less than 5% of unbrominated ester and characterized by:

^1H NMR (CDCl_3)

8.16(d, 2H), 7.38(d, 2H), 5.05(septet, 1H), 4.14(dd, 1H), 2.88(m, 4H), 2.39(m, 4H), 1.29(d, 6H);

^{13}C NMR (CDCl_3)

168.7, 147.7, 129.3, 123.8, 69.9, 45.1, 35.6, 33.0, 21.5, 21.2.

Example P

Preparation of α -[3-(4-nitrophenyl)propyl]-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester.

To a stirred solution of 3.137 g (18.2 mmoles) of cyclen free base in 30 ml of pentene stabilized chloroform was added 4.81 g (14.6 mmoles corrected) of *d,l*-2-bromo-4-(4-nitrophenyl)-butanoic acid isopropyl ester (prepared by the procedure of Example O) over a 5 minute period under a nitrogen

atmosphere with stirring. The reaction solution was stirred for 24 hours at room temperature. TLC analysis (Solvent System 2) revealed conversion to the titled monoalkylation product ($R_f = 0.78$ detection with ninhydrin, iodine, and UV activity). The yellow chloroform solution was applied to a 2.54 x 43.18 cm (1 in. x 17 in.) flash silica gel column which had been pre-eluted with 5% methanolic chloroform. Elution with 300 ml of this solvent system was followed by elution with Solvent System 2 and provided fractions containing pure titled product which were combined and evaporated affording 4.85 g (5.46 mmoles) of the titled product free base as a light oil in 79% yield and further characterized by:

^1H NMR (CDCl_3)

8.15(d, 2H), 7.40(d, 2H), 5.07(p, 1H), 3.35(dd, 1H), 2.65-3.0(m, 13H), 2.5-2.64(m, 4H), 2.14(m, 1H), 2.00-

(m, 1H), 1.28(dd, 6H);

^{13}C NMR (CDCl_3)

171.6, 149.5, 146.5, 129.2, 123.6, 68.1, 62.7, 49.2, 47.5, 45.9, 45.7, 32.9, 31.0, 22.1, 22.0;

IR (CDCl_3) cm^{-1}

3231(N-H), 2978, 2933, 1721 (ester carbonyl), 1601, 1512, 1458, 1345, 1107;

Fast atom bombardment mass spectrum, m/e 422 ($\text{M} + \text{H}$) $^+$, 408, 392.

Preparation of Final Products - Ligands

Example 1

Preparation of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, monomethyl, triethyl ester.

In 46 ml of acetonitrile containing 4.6 g (33.3 mmole) of freshly powdered anhydrous potassium carbonate was dissolved 700 mg (1.91 mmole) of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester (prepared by the procedure of Example D). To this suspension was added all at once 1.022 g (6.12 mmole) of ethyl bromoacetate. After four hours of stirring at room temperature, the suspension was vacuum filtered, the filter cake washed with 15 ml of acetonitrile, two times, and the filtrate evaporated *in vacuo* at 38°C to give the crude titled tetraester as a purple solid. The tetraester was then purified by silica gel chromatography eluting with 5 percent $\text{C}_2\text{H}_5\text{OH}/\text{CHCl}_3$ then Solvent System 3, to give 620 mg (0.988 mmole, 52 percent) of the desired product. This product was characterized by fast atom bombardment mass spectrometry ($[\text{M} + \text{H}]^+ = 624$) and further characterized by:

^1H NMR (CDCl_3)

8.24(d), 7.45(d), 4.94(s), 4.35-4.10(m), 3.79(s), 3.75-1.84(m), 1.34-1.22(m);

^{13}C NMR (CDCl_3)

174.3, 173.8, 173.6, 171.5, 147.6, 138.9, 131.5, 123.4, 64.8, 61.5, 61.4, 60.2, 55.4, 55.0, 53.1, 52.9, 52.7, 52.4, 51.9, 51.7, 48.8, 48.3, 44.9, 19.1, 14.1.

Example 2

Preparation of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.

In 30 ml of 6N HCl was dissolved 300 mg (0.478 mmole) of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, monomethyl, triethyl ester (prepared by the procedure of Example 1) and the mixture was heated at reflux overnight. At the end of this reflux period, the solution was concentrated *in vacuo* at 70°C to give a yellow solid. This solid was dissolved in 3 ml of water, filtered, and the filtrate evaporated to give 253 mg (0.378 mmole, 79 percent) of the crude product. This product was purified by preparative TLC [silica gel, developed in 20 percent $\text{NH}_4\text{OAc}/\text{CH}_3\text{OH}$ (wt/v)] and characterized by fast atom bombardment mass spectrometry ($[\text{M} + \text{H}]^+ = 526$) and further characterized by:

^1H NMR (D_2O)

8.21(d), 7.42(d), 4.83(s), 3.83-2.19(m), 1.92(s);

^{13}C NMR (D_2O)

175.0, 173.6, 171.8, 167.0, 166.3, 146.9, 138.3, 130.1, 123.0, 62.2, 54.0, 53.0, 52.2, 50.4, 97.3, 46.8, 44.8, 42.8, 20.0.

Example 3

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid; (BA-DOTA).

5 Method A:

Catalytic hydrogenation of the nitro group of the tetraacid (prepared by the procedure of Example 2) was performed using Adams catalyst (PtO_2) and hydrogen in essentially quantitative yield to give the titled product. This product was characterized by fast atom bombardment ($[\text{M} + \text{H}]^+ = 496$) and anion exchange

10 HPLC (on Q-SepharoseTM) and further characterized by:

^1H NMR (D_2O)

8.12(d), 7.86(d), 5.66(s), 4.73-4.57(m), 4.27-3.78 (m), 3.39-2.91(m);

^{13}C NMR (D_2O)

176.54, 176.52, 176.46, 141.1, 128.9, 120.9, 112.5, 65.0, 55.9, 55.7, 53.6, 49.7, 49.5, 49.3, 45.7, 45.4,
15 44.9, 44.6, 41.4.

Method B:

Another method to synthesize this compound was to convert the monoester tetraamine compound from
20 Example D to the monoacid tetraamine compound (6N HCl, reflux overnight), followed by aqueous alkylation using bromoacetic acid, then reduction of the nitro group to the amine group (PtO_2 catalyst) to give a product identical to that described above.

Example 4

25

Preparation of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid.

In 60 ml of 6N HCl was dissolved 2.0g (5.48 mmole) of the monoester tetraamine compound (prepared by the procedure of Example D). The mixture was heated to 95 °C overnight (about 10 hours) after which it
30 was concentrated *in vacuo* at 75 °C to give 2.53 g (5.11 mmole, 96 percent) of the monoacid tetraamine tetrahydrochloride as a yellow solid.

The above monoacid tetraamine tetrahydrochloride (0.5 g, 1.0 mmole) was dissolved in 15 ml of water and adjusted to pH 9 with NaOH. A separate solution of bromoacetic acid was prepared (0.71 g, 5.13 mmole) and added to the water solution of the monoacid tetraamine. The pH of the solution was again
35 adjusted to 9 and maintained at this pH during the reaction by small additions of 1.0N NaOH. While the reaction was stirred, aliquots were removed at different time points and the progress of the alkylation was monitored by anion exchange HPLC. When the amount of dialkylation product (total of three acetate groups present) had reached a maximum, the whole reaction mixture was lyophilized to give a dark solid. This crude mixture of alkylated products was then purified by silica gel chromatography (eluting with Solvent System 1), followed by preparative anion exchange chromatography (eluting with a gradient of 0-100 percent 1M NH_4OAc), then by preparative TLC plates (developed in Solvent System 4) and finally by silica
40 gal chromatography (eluting with Solvent System 4). This extensive purification procedure gave, as a yellow solid, 30 mg (0.06 mmole, 6.0 percent) of the title product. This product was characterized by fast atom bombardment mass spectrometry ($[\text{M} + \text{H}]^+ = 468$), anion exchange HPLC and further characterized by:

45 ^1H NMR (D_2O)

8.12(bs), 7.35(bs), 4.76(s), 3.57-3.44(m), 2.97-1.83(m), 1.80(s);

^{13}C NMR (D_2O)

181.75, 180.68, 179.00, 175.37, 147.70, 141.22, 133.08, 123.53, 68.44, 59.72, 56.00, 52.80, 49.96, 49.29,
47.86, 45.36, 44.26, 42.78, 42.25, 23.72.

50

Example 5

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid pentahydrochloride and
55 α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid pentahydrochloride.

A crude mixture of alkylated products was prepared by the procedure described in Example 4. This crude mixture was chromatographed on a silica gel column eluting with 20 percent $\text{NH}_4\text{OAc}/\text{CH}_3\text{OH}$ (wt:v). The appropriate fractions from this column were combined and evaporated to dryness. The crude product,

having substantial amounts of NH_4OAc , was dissolved in 80 ml of NANOpure™ water and lyophilized to yield a light brown solid. This solid was dissolved in 20 ml of NANOpure™ water, shaken with PtO_2 catalyst under a H_2 atmosphere until hydrogen uptake stopped. The catalyst was separated by vacuum filtration, add the filtrate lyophilized to give a yellow solid. The solid was dissolved in 3 ml of Solvent System 4 and chromatographed on silica gel using Solvent System 4. The pooled fractions were then stripped *in vacuo* and lyophilized to yield 779.6 mg of light yellow solid. ^{13}C NMR and proton NMR suggests that this product is a 50/50 mixture of the two possible geometric isomers. The isomeric mixture was contacted with excess HCl and lyophilized to yield 548.4 mg (61 percent yield) of the title products. M.P. > 270 °C. Anion exchange chromatography (HPLC System III) showed one major peak (>94 percent purity); fast atom bombardment mass spectrum indicated the geometrical isomers $[\text{M} + \text{H}]^+ = 438$.

Example 6

Preparation of α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester.

To a solution of 1.43 g (3.63 mmole) of α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester (prepared by the procedure of Example B) in 40 ml of argon purged acetonitrile was added 1.71 g (12.34 mmole) of anhydrous potassium carbonate with vigorous stirring. To the reaction mixture was added 1.78 g (11.61 mmole) of methyl bromoacetate and the reaction mixture was stirred at about 25 °C for 48 hours. TLC analysis indicated formation of a new product ($R_f = 0.62$, Solvent System 2). To the slurry was added 6.0 g of flash silica gel and acetonitrile was removed on a rotary evaporator. The resulting material was applied to 2.54 x 40.64 cm (1 in. x 16 in.) silica column which had been pre-eluted with 5 percent methanolic chloroform. About 400 ml of the prepared solution was used to elute several nonpolar impurities and unreacted alkylating agent. Then Solvent System 2 was applied, fractions containing product ($R_f = 0.62$) were collected, combined and evaporated to provide 2.1 g (3.44 mmole, 95 percent) of the titled product as a pale yellow glass. ^1H NMR indicated this product to be a 2:1 mixture of conformational or geometric isomers.

^1H NMR (CDCl_3 , 50 °C)

8.137(d), 8.128(d), 7.398(d), 7.385(d), 3.82(s), 3.76(s), 3.75(s), 3.74(s), 3.68(s), 1.5-3.52(m);

^{13}C NMR (CDCl_3 , 50 °C, 75 MHz)

175.8, 174.2, 174.1, 174.0, 149.0, 129.5, 129.3, 123.6, 60.1, 55.1, 52.8, 52.7, 52.6, 52.4, 52.2, 52.1, 52.0, 51.2, 51.1, 51.0, 49.1, 48.8, 47.5, 45.2, 34.2, 32.6.

Example 7

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester.

In 25 ml of methanol containing 400 mg of 10 percent palladium-on-carbon catalyst under a nitrogen atmosphere was dissolved 1.41 g (2.31 mmole) of α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester (prepared by the procedure of Example 6). Excess hydrogen was purged through the solution at one atmosphere for three hours. TLC indicated the formation of a strongly ninhydrin positive material ($R_f = 0.62$, Solvent System 2). The methanolic catalyst slurry was filtered through celite and evaporation of solvent provided the title product, 1.21 g (2.08 mmole, 91 percent), as a white solid glass (as a 2:1 mixture of conformational isomers). The minor conformational or geometric isomer was separated from the major isomer by flash chromatography using 10 percent methanolic chloroform to yield the title product as an off white powder (MP = 89-95 °C) and characterized by:

^1H NMR (CDCl_3 , 50 °C)

6.94(d), 6.89(d), 6.69(d), 6.66(d), 3.91(s), 3.80(s), 3.78(s), 3.74(s), 3.73(s), 3.72(s), 3.71(s), 1.5-3.5(m);

^{13}C NMR (CDCl_3 , 50 °C, 75 MHz)

176.1, 174.0, 173.9, 172.2, 170.4, 169.6, 144.2, 144.1, 130.6, 130.5, 129.5, 129.1, 115.9, 115.8, 62.6, 58.7, 55.1, 54.3, 52.7, 52.5, 52.3, 52.2, 52.1, 52.0, 51.9, 51.8, 51.7, 50.2, 50.0, 47.3, 44.7, 32.8, 31.8, 30.0, 25.2;

Fast atom bombardment mass spectrum, m/e 602 $[\text{M} + \text{Na}]^+$, 618 $[\text{M} + \text{K}]^+$, $[624 \text{ M} + 2\text{Na}^+ - \text{H}]^+$.

Example 8

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid; (PA-DOTA).

In 50 ml of conc. hydrochloric acid was dissolved 1.5 g (2.59 mmole) of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester (as the crude 2:1 mixture prepared by the procedure of Example 7). The reaction mixture was refluxed at about 105 °C for six hours. TLC (Solvent System 1) indicated conversion of the ester (R_f = 0.88) to the title product as the hydrochloride salt (R_f = 0.43). Excess solvent was removed on a rotary evaporator and the resulting white solid was dried. The title product as the hydrochloride salt was obtained in a yield of 1.6 g and characterized by:

^1H NMR (D_2O , pD = 1.0, 90 °C)

7.48(d), 7.42(d), 2.8-4.3(m), 2.23(m), 2.03(m);

^{13}C NMR (D_2O , pD = 1.0, 90 °C, 75 MHz)

176.4, 174.9, 171.6, 144.7, 132.9, 132.8, 130.4, 125.7, 61.7, 56.8, 56.0, 53.7, 53.1, 51.6, 47.9, 34.3, 37.6,

Fast atom bombardment mass spectrum m/e 524 $[\text{M} + \text{H}]^+$, 546 $[\text{M} + \text{Na}]^+$, 568 $[\text{M} + 2\text{Na} - \text{H}]^+$.

Using flash chromatography and Solvent System 1, any trace impurities of the ester were removed from the title product as the hydrochloride salt. To a 2.54 x 58.42 cm (1 in. x 23 in.) flash silica gel column was applied 1.10 g of the title product, as the hydrochloride salt. Fractions containing only the title product, as the mixed ammonium potassium salt were combined to provide 580 mg.

HPLC analysis indicated the material to be greater than 98 percent area purity at 230 and 254 nm.

Fast atom bombardment mass spectrum, m/e 562 $[\text{M} + \text{K}]^+$, 584 $[\text{M} + \text{K} + \text{Na} - \text{H}]^+$, 600 $[\text{M} + 2\text{K} - \text{H}]^+$.

Example 9

Isolation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid, mixed ammonium, potassium salt.

A crude solution containing both the tetra and tri acids (2.05 g) (prepared by the procedure of Example 8) was dissolved in a minimum amount of Solvent System 1 and applied to a 30.48 x 7.62 cm (12 in. x 3 in.) column of flash silica gel which had been pre-eluted with this solvent. Elution of fractions containing the title product (R_f = 0.63 in Solvent System 1) were collected and afforded 200 mg of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid, mixed ammonium, potassium salt. ^1H and ^{13}C NMR analysis suggested that this triacid was the symmetrical isomer (1,4,10-triacid positional isomer):

^1H NMR (D_2O , pD = 0.5 with DCl, T = 90 °C)

7.52(d), 7.46(d), 3.60(m), 3.54(m), 3.19(m);

^{13}C NMR (D_2O , pD = 0.5, T = 90 °C)

176.1, 170.2, 140.0, 132.7, 131.1, 126.1, 57.6, 57.2, 56.1, 54.9, 53.2, 51.5, 51.2, 31.2.

Fast atom bombardment mass spectrum, m/e 466 $[\text{M} + \text{H}]^+$, 488 $[\text{M} + \text{Na}]^+$, 504 $[\text{M} + \text{K}]^+$, 526 $[\text{M} + \text{K} + \text{Na} - \text{H}]^+$, 542 $[\text{M} + 2\text{K} - \text{H}]^+$.

Example 10

Preparation of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetramethyl ester.

Methyl bromoacetate (565 μl , 5.97 mmole) was added to a stirred mixture of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane, 1-acetic acid, methyl ester (580 mg, 1.47 mmole) (prepared by the procedure of Example N) and pulverized potassium carbonate (1.15 g, 8.32 mmole) in acetonitrile (20 ml). The reaction mixture was stirred at room temperature for 2 hours. The mixture was then filtered and the filtrate was concentrated to an oil *in vacuo*. The crude product was purified by chromatography (silica gel, 8 percent methanol in methylene chloride). The title product was obtained as a yellow solid (469 mg, 52 percent), TLC R_f = 0.4 (silica gel, 10% CH_3OH in CHCl_3), and further characterized by:

^{13}C NMR (CDCl_3)

173.2, 172.6, 161.2, 139.1, 125.1, 124.6, 120.5, 110.7, 57.0, 55.6, 53.5, 52.6, 51.3, 50.8, 46.8, 46.0, 44.0.

Example 11

Preparation of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.

A solution of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetramethyl ester (prepared by the procedure of Example 10) in concentrated HCl (5 ml, J.T. Baker ULTREX) was refluxed under a nitrogen atmosphere for 5 hours. The solution was then concentrated to dryness *in vacuo* to leave a solid residue. This was purified by chromatography (silica gel, Solvent System 6) to yield the title product as an off-white solid (209 mg). This material was converted to the tetrahydrochloride salt and was characterized by:

^{13}C NMR (D_2O -DCI, 80°C)

171.0, 166.9, 161.2, 139.0, 125.5, 119.3, 110.7, 56.8, 54.5, 52.7, 51.0, 49.2, 49.0, 46.3, 42.9.

Example 12

Preparation of α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetraammonium salt.

To a solution of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (157 mg) (prepared by the procedure of Example 11) in water (20 ml) was added platinum oxide (20 mg). This mixture was hydrogenated (1 atmosphere hydrogen) for 1 hour at room temperature. After filtration, the material was further purified by chromatography (silica gel, Solvent System 5) to yield the title compound (141 mg) as an off-white solid, and characterized by:

^{13}C NMR (D_2O -DCI)

175.5, 172.6, 172.3, 159.9, 128.8, 127.5, 124.6, 123.8, 115.5, 61.2, 58.1, 57.3, 55.7, 53.4, 51.6, 49.6, 47.4.

Example 13

Preparation of 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-triacetic acid, 4,7,10-trimethyl ester.

To 4 ml of argon purged acetonitrile with stirring was added 100 mg (0.236 mmole) of 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane, tetrahydrochloride salt (prepared by the procedure of Example J), 260 mg (1.88 mmole) of potassium carbonate, and 108 mg (0.708 mmole) of methylbromoacetate. The mixture was stirred for 48 hours at 25°C . The salt containing solution was applied to a 1 cm x 4 cm flash silica gel column and was eluted with acetonitrile. Fractions containing the desired product were combined and the solvent removed by rotary evaporation to provide 65 mg (0.132 mmole, 56 percent) of the title product which was further characterized by:

^1H NMR (CDCl_3)

7.29 (d), 6.75 (d), 4.62 (s), 4.20 (broad s), 3.70 (s), 3.69 (s), 3.64 (s), 3.62 (t), 3.27 (s), 3.06 (t), 2.82 (broad t), 2.79 (broad t);

^{13}C NMR (CDCl_3 , 75 MHz)

171.4, 171.0, 148.4, 132.8, 117.5, 115.0, 55.9, 55.3, 54.7, 53.1, 51.7, 51.4, 50.6, 50.5, 47.4.

Example 14

Preparation of 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt.

In 2 ml of 6N hydrochloric acid, 42 mg (0.085 mmole) of 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester (prepared by the procedure of Example 13) was heated to 80°C for 2 hours. TLC indicated several products ($R_f = 0.60$ for the desired title product, Solvent System 1). Removal of solvent afforded 41 mg of crude title product.

Example 15

Preparation of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester.

To a solution of 2.24 g (6.97 mmole) of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane (prepared by the procedure of Example G) in 75 ml of argon purged acetonitrile was added with vigorous stirring 3.17 g of anhydrous potassium carbonate. To this reaction mixture was added dropwise over a five minute period 3.20 g (20.9 mmole) of methylbromoacetate. The reaction mixture was stirred for 17 hours under argon atmosphere. TLC analysis showed conversion of the starting material ($R_f = 0.73$, Solvent System 1) to a new product ($R_f = 0.46$, Solvent System 1). To the solution was added 70 ml of chloroform and the solution with the suspended salt was applied to a 2.54 x 17.78 cm (1 in. x 7 in.) flash silica column. The product was eluted using 10 percent methanol in chloroform to yield 2.95 g of the title product as an amber oil which formed a friable glass upon vacuum drying (78 percent). The title product was characterized by:

^1H NMR (CDCl_3)

8.17 (m,m), 7.4-7.66 (m), 2.38-3.95 (m).

^{13}C NMR (CDCl_3 , 75 MHz)

171.5, 171.2, 146.7, 144.5, 130.3, 123.9, 56.0, 55.2, 53.5, 53.4, 52.6, 51.9, 51.8, 50.6, 48.1, 29.5.

Example 16

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester.

In 50 ml of methanol was dissolved 2.8 g (5.18 mmole) of crude 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester (prepared by the procedure of Example 15). To the stirred solution which was purged with nitrogen was added 10 percent palladium on carbon catalyst (600 mg). The solution was maintained under an atmosphere of nitrogen, and then hydrogen (1 atm, 20-25 °C) was purged through the stirred solution for 2.5 hours. The solution was then purged for several minutes with nitrogen and the catalyst removed by filtration through a short bed of celite. TLC analysis (10 percent methanol in chloroform) revealed the title product ($R_f = 0.14$). The title product was eluted with chloroform from silica gel to yield 2.2 g (4.33 mmole, 83 percent) as a white glass and characterized by:

^1H NMR (CDCl_3)

7.03 (d,m), 6.63 (d), 3.4-3.6 (m), 2.7-3.2 (m);

^{13}C NMR (CDCl_3 , 75 MHz)

171.4, 171.2, 145.6, 129.7, 125.6, 115.5, 55.9, 54.4, 54.3, 53.0, 52.8, 51.8, 51.6, 50.3, 47.8, 28.6;

Fast atom bombardment mass spectrum, m/e 508 $[\text{M} + \text{H}]^+$.

Example 17

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt; (EA-D03A).

In 30 ml of conc. HCl was dissolved 850 mg (1.69 mmole) of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester (prepared by the procedure of Example 16). The solution was stirred at 70-80 °C for 2.5 hours and the solution allowed to cool with stirring to 25 °C and the stirring continued for about 18 hours. Solvent was removed on a rotary evaporator at reduced pressure (10 mm, 75 °C). Solvate and excess hydrogen chloride were removed from the resulting clear oil by vacuum drying (10^{-1} mm, 45 °C). The title product was provided as a white solid, in a yield of 890 mg, ($R_f = 0.46$, Solvent System 2) and further characterized by:

^1H NMR (D_2O , $pD = 1.9$, $T = 90^\circ\text{C}$)

7.50(d), 7.41(d), 4.26(s), 3.43-3.68(m), 3.0-3.3(m);

^{13}C NMR (D_2O , $pD = 1.9$, $T = 90^\circ\text{C}$)

176.7, 170.9 139.8, 133.0, 131.2, 125.9, 57.5, 57.1, 55.4, 54.4, 52.7, 51.0, 50.6, 31.3;

Fast atom bombardment mass spectrum, m/e 466 $[\text{M} + \text{H}]^+$, 488 $[\text{M} + \text{Na}]^+$, 510 $[\text{M} + 2\text{Na} + \text{H}]^+$.

HPLC analysis indicated greater than 92 percent area purity (254 nm) using a 10 cm Partisil®-5 OD53 RAC

II reverse phase column. The eluent was 10 percent acetonitrile in 0.05M pH = 7.0 potassium phosphate solution.

Example 18

Preparation of 1-[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid; (SCN-EA-DO3A).

1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt (prepared by the procedure of Example 17, 106 mg, 0.21 mmole) was dissolved in 2 ml of water under a nitrogen atmosphere with stirring. Sodium bicarbonate (105.6 mg, 1.26 mmole) was slowly added to prevent frothing from carbon dioxide evolution (resulting pH = 8.0). Thiophosgene (16.8 μ l, 0.22 mmole) was added and after one hour of vigorous stirring, TLC analysis (20 percent water in acetonitrile) indicated conversion of starting material (R_f = 0.12) to title product (R_f = 0.26). The solvent was removed from the crude product by rotary evaporation to afford 130 mg of title product which contained sodium chloride. Infrared analysis of this material (KBr pellet) confirmed the presence of the isothiocyanate moiety (SCN = 2150 cm^{-1}).

Example 19

Preparation of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, triethyl ester.

To 5 ml of a stirred acetonitrile solution containing 395 mg (1.12 mmole) of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane (prepared by the procedure of Example G) and 1.5 g (11 mmole) of K_2CO_3 was added 496 μ l (4.5 mmole) of ethyl bromoacetate in one portion. After heating at 68°C under a N_2 atmosphere for one hour, the resulting suspension was filtered. The filter cake was washed with CH_3CN (2 x 10 ml). The filtrate was then concentrated to give the crude product as a viscous oil, triturated with 30 ml of diethyl ether, and concentrated to give 650 mg (100% yield) of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, triethyl ester and characterized by:

^1H NMR (CDCl_3)

8.2(d), 7.6(d), 4.3(q), 2.6-3.8(m), 1.55(t);

Fast atom bombardment mass spectrum $[\text{M} + \text{H}]^+ = 588$.

Example 20

Preparation of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

To a suspension of 200 mg (0.35 mmole) of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, triethyl ester (prepared by the procedure of Example 19) in 15 ml of water was added 59 μ l of NaOH (50% wt/wt, 3 eq). The mixture was stirred at 80°C for 6 hours. The resulting homogeneous orange solution was then cooled to room temperature and freeze-dried to yield a brown solid which was purified by HPLC (Q-SepharoseTM, HPLC System III) using a linear gradient of 0-10% aqueous HOAc over 60 minutes to yield (50%) of the hydrolyzed product 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, and characterized by:

^1H NMR (D_2O)

8.22(d), 7.55(d), 3.60-3.90(m), 3.10-3.40(m).

Example 21

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid; (EA-DO3A).

In a Parr-hydrogenator was placed 50 ml of an aqueous solution of 214 mg (0.43 mmole) of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 20) and 40 mg of 10% Pd/C. The suspension was shaken until hydrogen uptake ceased. After filtration, the aqueous solution was freeze-dried yielding 197 mg (98%), as a tan solid, 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, and characterized by:

^1H NMR (D_2O)

7.35(d), 7.20(d), 3.10-3.70(m);

¹³C NMR (D₂O)

173.85, 172.50, 137.90, 131.90, 130.22, 121.55, 56.25, 55.14, 54.42, 50.12, 49.65, 49.31, 31.00.

Example 22

Preparation of 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester.

To a stirred acetonitrile solution (20 ml) containing 1.0g of 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane (3 mmole) (prepared by the procedure of Example E) was added 1.6g of methyl bromoacetate (11 mmole) in one portion. After one hour potassium carbonate (0.3 g) was added and stirring continued at room temperature for twelve hours. The reaction mixture was then filtered and the filtrate concentrated (*in vacuo*). The resulting residue was column chromatographed (silica, CH₃CN/CH₃OH, 9:1, V:V) yielding the triester as a white solid in 68 percent yield after concentration, and was further characterized by:

¹H NMR (CDCl₃)

8.19(d), 8.14(s), 7.18(d), 2.58-3.99(m);

¹³C NMR (CDCl₃)

173.53, 172.83, 164.59, 142.28, 128.51, 127.69, 126.49, 112.30, 57.25, 56.90, 55.03, 54.21, 53.06, 52.09, 52.00, 51.54, 50.68, 50.30, 49.85, 49.63, 49.31, 49.00, 48.68, 48.37, 48.05, 47.70, 47.39.

Example 23

Preparation of 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

A 6M hydrochloric acid solution (3 ml) containing 0.25 g of 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester (0.45 mmole) (prepared by the procedure of Example 22) was stirred and heated at 100 °C for 24 hours. After cooling to room temperature, 5 ml of water was added and the aqueous solution freeze-dried to give a tan solid. Following column chromatography (silica, Solvent System 5) the triacid was isolated in 60 percent yield as an off-white solid, MP = 228-230 °C (dec), and further characterized by:

¹H NMR (D₂O)

8.18(d), 8.09(s), 7.12(d), 3.87(s), 2.10-3.60(m);

¹³C NMR (D₂O)

180.45, 179.80, 163.66, 140.22, 128.60, 126.24, 122.66, 111.50, 59.91, 59.06, 56.44, 52.75, 50.92, 48.84.

Example 24

Preparation of 1-(2-methoxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

To a nitrogen purged aqueous (50 ml) solution containing 50 mg of PtO₂ was added 50 mg of 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 23). After hydrogenating in a Parr bomb for one hour, the solution was filtered and the aqueous filtrate freeze-dried to give the aniline derivative as a tan solid (95 percent yield), MP > 240 °C dec, and further characterized by:

¹H NMR (D₂O)

7.54(bs), 7.21(d), 7.09(d), 7.05(s), 6.88(s), 3.84(s), 3.78(s), 2.85-3.56(m);

¹³C NMR (D₂O)

180.59, 179.77, 153.38, 124.16, 124.03, 122.82, 120.18, 113.68, 112.43, 59.06, 57.02, 55.96, 53.67, 50.52, 50.08, 49.70, 49.04, 48.32.

Example 25

Preparation of 1-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

To an aqueous solution (1 ml) of 1-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane (200 mg, 0.62 mmole) (prepared by the procedure of Example F) was added bromoacetic acid (309 mg, 2.2 mmole, and NaOH (3.5 ml, ≈ 1M) with stirring at room temperature. The reaction progress was monitored by LC

(anion exchange, Q-Sepharose™) and the pH maintained at about 8 via the addition of NaOH as needed. After 12 hours the solution was freeze-dried and the solid residue chromatographed (column) on silica gel eluting with Solvent System 5. The major yellow fraction was concentrated, dissolved in H₂O and freeze dried to give the desired product as a bright yellow powder (150 mg, 49 percent); MP = 230-235 °C (dec),

and further characterized by:

¹³C NMR (D₂O)

166.41, 165.60, 160.00, 119.91, 116.02, 113.85, 111.71, 104.59, 45.20, 44.50, 43.96, 36.43.

Example 26

Preparation of 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

To an argon purged, aqueous solution (25 ml) of N-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (200 mg, 0.4 mmole) (prepared by the procedure of Example 25) was added PtO₂ (130 mg) followed by the introduction of H₂ (Parr hydrogenator). After total disappearance of yellow color, the solution was purged with argon and filtered. The aqueous solution was then freeze-dried to give the product as a tan solid (146 mg, 78 percent).

Example 27

Preparation of α-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester; (PN-DOTMA trimethylisopropyl ester).

To a solution of 1.00 g (2.37 mmole) of α-[3-(4-nitrophenyl)propyl]-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester (prepared by the procedure of Example P) in 35 ml dry acetonitrile was added 1.15 g (8.32 mmoles) of anhydrous potassium carbonate with vigorous stirring under nitrogen. Optically active α-benzene sulfonate of lactic acid methyl ester (1.79 g, 7.36 mmoles S:R=98.2) was added and the mixture was stirred at room temperature for 60 hours. TLC analysis indicated formation of new products (R_f=0.71, Solvent System 2 for titled tetraesters and R_f=0.89 for triester). The resulting solution with suspended carbonate salts was poured into 40 ml of chloroform and the precipitated inorganic salts were filtered. Solvent was removed from this filtrate and the crude orange oil was chromatographed twice on a 2.54 x 40.64 cm (1 in. x 16 in.) flash silica gel column using Solvent System 2 as the eluent to afford the titled product as a nonresolved mixture of tetraester diastereomers (1.00 g, 1.47 mmole, 62% yield) which were free of underalkylated products.

¹H NMR analysis of this material showed it to contain approximately 0.5 equivalents of unreacted benzene sulfonate derivative. The NMR spectra of this material were not coalescent at 60 °C in chloroform:

¹H NMR (CDCl₃, 60 °C)

7.9-8.1(m, 2H), 7.2-7.4(m, 2H), 5.06(m, 1H), 3.4-4.0(m, 9H), 1.7-3.4(m, 20H), 1.0-1.7(m, 15H);

IR (CDCl₃) cm⁻¹

2980, 2840, 1725, 1710 (ester carbonyl), 1590, 1510, 1440, 1340;

Fast atom bombardment mass spectrum, m/e 702 [M + Na]⁺, 687.

Example 28

Preparation of α-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester; (PA-DOTMA trimethylisopropyl ester).

α-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester (950 mg, 1.40 mmoles uncorrected) (prepared by the procedure of Example 27), which contained unreacted α-[3-(4-nitrophenyl)propyl]-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester, was dissolved in 10 ml of 30% aqueous methanol containing 1.00 g of 10% palladium on carbon catalyst under a nitrogen atmosphere. Excess hydrogen was purged through the solution for 7 hours. At this point, TLC inspection revealed formation of a strongly ninhydrin positive material (R_f=0.82 Solvent System 2, R_f=.71 for starting material). The methanolic catalyst slurry was filtered through celite and evaporation of solvent provided 800 mg of the crude title product as a clear oil which contained benzene sulfonic acid (R_f=0.09 Solvent System 2) from the concomitant hydrogenolysis of the benzene sulfonate ester. The crude oil was chromatographed on a 2.54 x 20.32 cm (1 in. x 8 in.) flash silica gel column with 10% methanol in chloroform which eluted a light yellow band in the void volume. Solvent

System 2 was then applied to elute the titled product (480 mg) in 52% yield as a mixture of unresolved diastereomers and geometric isomers. Three distinct ab quartets were observed in the aromatic region of the ^1H NMR:

^1H NMR (CDCl_3 , 30°C)

6.63-8.0(m (quartets), 4H), 5.07(m, 1H, isopropyl methine H), 3.53-3.9(m, 13 H with four singlets at 3.85, 3.73, 3.69 and 3.56), 3.46(m, 1H), 3.25(broad t, 3H), 2-3.1(m, 14H), 1.0-2.0(m, 15H);

^{13}C NMR (CDCl_3 , 30°C)

177.3, 177, 176.4, 176.2, 175.9, 174.3, 172.9, 170.6, 145.3, 144.9, 130.8, 129.7, 129.3, 129.1, 128.8, 128.4, 127.5, 126.3, 115.2, 115.1, 114.9, 69.0, 68.4, 67.3, 61.6, 57.8, 57.4, 56.7, 56.5, 56.4, 52.9, 52.7, 52.1, 50.8, 50.7, 50.6, 47.3, 47.1, 47.0, 46.9, 46.5, 46.3, 46.1, 44.8, 44.5, 44.3, 34.1, 32.9, 32.5, 32.2, 31.8, 24.8, 22.2, 22.1, 22.0, 21.8, 21.6, 15.8, 14.9, 7.7, 7.5, 7.4, 7.1;

IR (CDCl_3) cm^{-1}

2960, 2840, 1720, 1510, 1450, 1375;

Fast atom bombardment mass spectrum, m/e 672 $[\text{M} + \text{Na}^+]^+$, 686.

Example 209

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(R-methylacetic) acid; (PA-DOTMA).

The diastereomeric mixture of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester (400 mg, 0.59 mmol) (prepared by the procedure of Example 28) was dissolved in 50 ml of 6N concentrated hydrochloric acid and refluxed for 30 hours. TLC analysis (Solvent System 1) indicated conversion of ester ($R_f=0.98$ Solvent System 1) to two new spots ($R_f=0.6$, high diastereomer and $R_f=0.54$, low diastereomer Solvent System 1, both spots UV, ninhydrin, and iodine positive). Excess solvent was removed on a rotary evaporator and after drying, a mixture of crude diastereomers of the title product (403 mg) was obtained as the hydrochloride salt. Preparative separation of the two diastereomers of the title product was accomplished by applying 290 mg (0.51 mmol) of the crude salt to a 1×13 cm silica flash gel column which was eluted with Solvent System 8. The high R_f diastereomer (119 mg, 0.21 mmol) was obtained in purer form than the low R_f diastereomer (90 mg, 0.16 mmol) since high R_f diastereomer eluted from the column first:

^1H NMR for high R_f diastereomer:

^1H NMR (D_2O , 90°C , $\text{pD}=7.5$)

7.12(d, 2H), 6.80(d, 2H), 3.66(m, 1H), 3.57(broad t, 3H), 2-3.4(m, 19H), 1.89(m, 1H), 1.37(m, 1H), 1.15(d, 3H), 1.10(d, 3H), 1.06(d, 3H);

^{13}C NMR (D_2O , 90°C , $\text{pD}=7.5$)

184.4, 184.2, 184.0, 183.3, 135.9, 132.6, 131.7, 119.1, 78.2, 69.0, 61.7, 61.5, 61.1, 57.4, 54.1, 49.7, 49.6, 48.2, 47.7, 47.2, 34.8, 33.8, 9.6, 9.1, 9.0;

Fast atom bombardment mass spectrum, m/e 566 $[\text{M} + \text{H}^+]^+$, 588 $[\text{M} + \text{Na}^+]^+$, 604 $[\text{M} + \text{K}^+]^+$, 626 $[\text{M} + \text{K}^+ + \text{Na}^+ - \text{H}^+]^+$, 642 $[\text{M} + 2\text{K}^+ - \text{H}^+]^+$.

Preparation of Final Products - Complexes

Metal ligand complexes were prepared by various methods as shown below. The methods included mixing of the metal and ligand in aqueous solution and adjusting the pH to the desired value. Complexation was done in solutions containing salts and/or buffers as well as water. Sometimes heated solutions were found to give higher complex yields than when the complexation was done at ambient temperatures. Also the work-up used in the synthesis of BFC has been shown to effect the complexation.

Example 30

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, ammonium salt, samarium(III) complex; $\text{Sm}(\text{BA-DOTA})$.

A small sample, 53 mg, (0.094 mmol) of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 3) was dissolved in 100 μl of water and treated with 3 ml of 0.043M $\text{Sm}(\text{OAc})_3$ solution (48.8 mg, 0.128 mmol) at pH 6-7. This solution was heated at 100°C and the degree of complexation was determined by anion exchange HPLC. When complexation was

complete by anion exchange chromatography, HPLC System III, the solution of the complex was cooled to room temperature (about 25 °C) and freeze dried. The samarium complex was purified by silica gel chromatography (eluting with Solvent System 1). This procedure gave the title product in 82 percent yield. This complex was characterized by TLC, fast atom bombardment mass spectrometry, anion exchange

5 HPLC, and further characterized by:

^1H (D_2O)

8.94, 7.81, 7.21, 6.97, 6.80, 5.63, 5.01, 4.58, 4.01, 3.56, 1.78, 1.53, 1.24, 1.10, 0.77, -2.80, -2.95, -3.21, -3.91;

^{13}C NMR (D_2O)

10 190.3, 184.9, 181.4, 146.8, 134.3, 122.7, 116.4, 80.8, 72.6, 64.2, 57.2, 55.8, 54.7, 53.4, 51.5, 49.7, 45.5, 43.5, 23.6.

Example 31

15 Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex.

A solution of 150 μl Sm-153 ($3 \times 10^{-4}\text{M}$, in 0.1N HCl) and 600 μl SmCl_3 ($3 \times 10^{-4}\text{M}$) in 0.1N HCl was prepared by adding the Sm-153 solution to the SmCl_3 solution. Then 5.0 μl of 2.0M NaOAc was added
20 followed by 45 μl of 50 mM ligand (prepared by the procedure of Example 3) in HEPES buffer. The pH of the solution was adjusted to 7 with 275 μl of 0.5M HEPES. This solution was split into two fractions of 500 μl s each, and one fraction was heated for 1 hour at 100 °C.

The solutions were passed through SP-Sephadex™ cation exchange resin. The percent of metal as a complex was determined using the procedure previously described. The results showed 99.6% of the metal
25 as a complex for the heated sample and 96.6% for the non-heated sample.

Stability of chelates by pH profile.

The stability of metal ligand complexes was measured by subjecting them to various pH values and
30 analysing for complex in solution. At low pH, protonation of the ligand can cause the release of metal. At high pH, metal hydroxides compete with chelate formation. Thus, when looking for inert chelates, a desirable inert chelate will remain as a chelate when exposed to high and low pH values.

Profiles of the inertness at various pH values for some of the chelates of this invention were generated using the methods described in the following examples. In some cases free metal was removed by passing
35 the solution through a cation exchange resin. Dilute sodium hydroxide and hydrochloric acid solutions were used to adjust the pH of the complex solutions from 1 to 14. The percent metal as a complex was then determined by the methods previously described.

In a similar manner, several bifunctional chelate conjugates were prepared and subjected to pH 2.8, 4 and 6. The amount of conjugate remaining in solution was determined by HPLC using a radiometric
40 detector. The results are shown in Example XX in the biology section.

Example 32

pH Stability of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III)
45 complex.

The heated and non-heated samples from Example 30 were each split into 2 x 250 μl aliquots. One 250 μl aliquot was adjusted with HCl to give the pH's represented in the table following. The second 250 μl aliquot was adjusted with μl quantities of NaOH to the desired pH's. Once the desired pH was reached, a
50 25 μl aliquot was removed and the percent of metal as complex determined as previously described. The results are as follows:

pH	% Complex	
	Heated	Not heated
1	98.9	92.7
3	99.6	93.3
5	99.6	92.9
7	99.6	95.7
9	99.9	95.8
11	99.6	95.7
13	99.6	94.5

Example 33

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, yttrium(III) complex.

Ten μ l of a solution containing 1.4 mg/100 μ l of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 3) was added to 100 μ l of $Y(OAc)_3$ - (0.003M) spiked with Y-90. To this solution was added 500 μ l of water followed by 125 μ l of NaOAc (0.5M). Water (265 μ l) was added to the solution to bring the total volume to 1 ml. This solution was divided into two aliquots. One aliquot was heated to 100°C for 1 hour. Both heated and non-heated samples were tested to determine the percent of metal as a complex using the cation exchange procedure previously described. The results showed 84% and 4% of the metal as a complex for the heated sample and the non-heated sample, respectively.

Example 34

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, lutetium(III) complex.

A solution of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 3) was dissolved in distilled water to yield a concentration of 1.63 mg/ml.

A lutetium chloride solution was prepared by dissolving lutetium chloride in 0.1N HCl to result in a concentration of 3.9 mg/ml (0.01M). Lutetium-177 (0.3 mmolar) was used as a tracer.

A volume of 30 μ l of 0.01M lutetium solution was added to 2 μ l of Lu-177 solution. The appropriate amount of ligand from above was added and HEPES buffer (0.1M, pH = 7.6) was added to give a total volume of 1.0 ml. The resultant solutions were 0.3 mmolar in ligand and lutetium. The solution was then heated to 100°C for one hour and the percent of Lu as a complex determined to be 86% by the cation exchange method described previously.

Example 35

Preparation of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex.

A sample, 7 mg, (10.8 μ mole) of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex (prepared by the procedure of Example 30) was dissolved in 400 μ l water. Excess thiophosgene (50 μ l) was added, followed by 400 μ l $CHCl_3$ and the two-phase reaction stirred vigorously for 30 minutes. At the end of this time, the water layer was extracted with 500 μ l $CHCl_3$ four times, and the water layer then was lyophilized to give the desired titled product in quantitative yield.

The UV showed this compound to have a band at 272 and 282 nm. The TLC, silica gel developed by 75:25 V:V $CH_3CN:H_2O$, gave R_f = 0.38. The starting material has an R_f = 0.19. IR showed -SCN stretch at 2100 cm^{-1} ; fast atom bombardment mass spectrum $[M + H]^+$ = 687.

Example 36

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, ammonium salt, yttrium(III) complex.

A sample, 90 mg (160 mmole), of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 3) was dissolved in 1 ml water. To this solution was added 3 ml of water containing 51 mg (192 mmole) of $Y(OAc)_3$. The reaction mixture, pH = 6 to 7, was then heated at 100 °C for two hours. The resulting solution was then passed through glass wool and lyophilized to give 126 mg of yellow solid. The solid was chromatographed on silica gel (Solvent System 1) to yield 71 mg (76 percent) of the desired complex product. Analysis by anion exchange showed the same retention time as was found for the analogous Sm complex. The title product was characterized by:

^{13}C NMR (D_2O)

180.8, 180.5, 179.9, 146.1, 133.4, 122.0, 116.1, 74.9, 66.3, 56.3, 55.8, 55.4, 55.1, 54.8, 52.2, 46.0, 44.0, 23.6;

1H NMR (D_2O)

6.90(d), 6.70(d), 4.40(s), 3.45-3.06(m), 2.71-2.10(m), 1.75(s).

Fast atom bombardment mass spectrum, $[M + H]^+ = 582$.

Example 37

Preparation of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, sodium salt, yttrium(III) complex.

A sample of the α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, yttrium complex (prepared by the procedure of Example 36) (10 mg, 17 μ mole) was dissolved in 400 μ l H_2O . To this solution was added 64 μ l thiophosgene (excess) and 400 μ l $CHCl_3$ and the resulting mixture stirred vigorously for 40 minutes. During this time several small additions of solid $NaHCO_3$ were made to keep the pH at about 8. At the end of the reaction, the water layer was separated and extracted with 1 ml of $CHCl_3$, four times, and lyophilized. The title product was characterized by TLC and UV spectroscopy.

Example 38

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, yttrium(III) complex, ammonium salt; $NH_4[Y(PA-DOTA)]$.

The ligand, α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 8) was converted to the mixed protonated-ammonium salt by chromatography on a strong cation exchange resin (SP-SephadexTM C-25, Pharmacia). The column was in the protonated form and washed with distilled water. A concentrated aqueous solution of the ligand was applied to the column, followed by washing the column with distilled water. The ligand was eluted from the column with 0.5M NH_4OH . The eluent was reduced to dryness.

A solution of $Y(OAc)_3 \cdot 4H_2O$ (0.0728 g, 0.215 mmole) in 5 ml of distilled water was added to a warm solution of the mixed protonated-ammonium salt ligand (0.120 g, 0.215 mmole) in 5 ml of water. The solution was brought to reflux and the pH adjusted to about 7.0 with 0.1M NH_4OH . After 1 hour at reflux, the solution was cooled and reduced to dryness. Excess NH_4OAc was removed by heating the white solid in an oil bath at about 105 °C under vacuum. The title product (which contained about one equivalent of ammonium acetate) yield was 0.142 g (94 percent) and was characterized by:

^{13}C NMR (D_2O , 88 °C, 75 MHz)

23.8, 27.6, 35.4, 49.6, 55.0, 56.2, 56.9, 57.1, 65.7, 68.0, 121.7, 132.6, 138.8, 180.7, 181.4, 182.0, 183.1;

Fast atom bombardment mass spectrum, m/e 610 (positive ion, $[M^- + 2H^+]^+$), 608 (negative ion, M^-).

Example 39

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex, ammonium salt; $\text{NH}_4[\text{Sm}(\text{PA-DOTA})]$.

When the procedure of Example 38 was repeated, using $\text{Sm}(\text{OAc})_3 \cdot 3\text{H}_2\text{O}$ (0.082 g, 0.215 mmole) in place of $\text{Y}(\text{OAc})_3 \cdot 4\text{H}_2\text{O}$, the title product (which contained about one equivalent of ammonium acetate) was prepared in a yield of 0.155 g (94 percent) and was characterized by:

^{13}C NMR (D_2O , 88°C , 75 MHz)

23.5, 27.2, 35.6, 50.5, 54.5, 56.0, 57.2, 57.9, 69.5, 71.5, 122.3, 133.0, 139.7, 181.2, 188.4, 189.3, 190.9;

Fast atom bombardment mass spectrum, m/e 673 (positive ion, $[\text{M}^- + 2\text{H}^+]^+$, Sm isotope pattern), m/e 671 (negative ion, M^- , Sm isotope pattern).

Example 40

Preparation of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex, ammonium salt; $\text{NH}_4[\text{Sm}(\text{SCN-PA-DOTA})]$.

Solutions of $\text{Sm}(\text{OAc})_3 \cdot 3\text{H}_2\text{O}$ (27.9 mg, 381.56 g/mole, 73.1 μmole) in 10 ml of distilled water and 42 mg of PA-DOTA, mixed ammonium-potassium salt (632.97 g/mole, 66.4 μmole) (prepared by the procedure of Example 8) in 10 ml of distilled water were mixed and heated on a steam bath. After 30 min., the reaction to form $[\text{Sm}(\text{PA-DOTA})]$ was complete as determined by HPLC using the method described in Example 41. The solution was reacted with a solution of 45.4 mg of CSCl_2 (114.98 g/mole, 395 μmole) in 20 ml of chloroform by shaking in a separatory funnel. The reaction was complete after about one min. as determined by HPLC and by the absence of a positive test with ninhydrin (0.2 % in ethanol) when the aqueous solution was spotted on a silica gel TLC plate (the starting chelate, $[\text{Sm}(\text{PA-DOTA})]$, tested positive). The chloroform layer was removed and the aqueous layer was washed with two 20 ml portions of chloroform. The aqueous layer was reduced to dryness by the addition of 100 ml of acetonitrile and evaporated at ambient temperature under a stream of nitrogen to yield 56 mg of the title product.

Fast atom bombardment mass spectrum, m/e 715 (positive ion, $[\text{M}^- + 2\text{H}^+]^+$, Sm isotope pattern), m/e 713 (negative ion, M^- , Sm isotope pattern).

Example 41

HPLC Analysis of the rate of chelation of Y^{3+} with PA-DOTA.

The rate of PA-DOTA chelation with Y^{3+} was studied as a function of the work-up used in the synthesis of PA-DOTA. The extent of chelation was monitored by HPLC using an Alltech Econosphere™ C18 100 mm column. The gradient used was: A) 95:5 of pH 6.0, 0.05M NaOAc buffer: CH_3CN , B) 30:70 of pH 6.0, 0.05M NaOAc buffer: CH_3CN , A to B in 15 min. A solution of PA-DOTA $\cdot n\text{HCl}$ (retention time = 1.31 min) and of PA-DOTA mixed ammonium-potassium salt (retention time = 4.55 min), both prepared by the procedure of Example 8, were prepared as 1.6 mM, pH 6.0 0.5M NaOAc buffer. One ml (1.6 μmole) of each of the two solutions was reacted with 0.2 ml (8 μmole) of $\text{Y}(\text{OAc})_3 \cdot 4\text{H}_2\text{O}$ solution (40 mM in pH 6.0 0.5M NaOAc buffer). Chelate formation was determined by the appearance of a peak at a retention time = 3.96 min (confirmed by comparison to a sample from Example 38). Chelate formation results were as follows.

PA-DOTA Salt	% Complex	Time(min)/Temp.
PA-DOTA $\cdot n\text{HCl}$	>90	<1/ room temp.
PA-DOTA mixed ammonium-potassium salt	<10 >85	15/ room temp. 10/ 90°C

Clearly, the method of purification of the BFC has an effect on the rate of chelation.

Example 42

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(methylacetic) acid, samarium(III) complex, ammonium salt; $\text{NH}_4[\text{Sm}(\text{PA-DOTMA})]$.

A solution of $\text{Sm}(\text{OAc})_3 \cdot 3\text{H}_2\text{O}$ (13.2 mg in 2 ml of distilled water) was added to a solution of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(methylacetic) acid (30 mg in 2 ml of distilled water), (prepared by the procedure of Example 29, high R_f diastereomer). The pH 6.5 solution was heated on a steam bath and the reaction was monitored by HPLC. After 30 min. of heating the solution

was reduced to dryness on a rotary evaporator and dried in a vacuum oven. Fast atom bombardment mass spectrum, m/e 715 (positive ion, $[\text{M} + 2\text{H}^+]^+$, Sm isotope pattern), 737 (positive ion, $[\text{M} + \text{H}^+ + \text{Na}^+]^+$, Sm isotope pattern) 713 (negative ion, (M^-) , Sm isotope pattern).

In the following examples the complex solutions were passed through an ion exchange column to remove any excess free metal from solution. Labile systems would return to equilibrium and similar amounts of the free metal would be in solution. Inert systems would not re-equilibrate and the amount of free metal should decrease. Thus even though a complex can be formed in low yield, a purification by passing it through an ion exchange resin could result in a usable stable complex without uncomplexed metal.

Example 43

Preparation of α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex.

A $3 \times 10^{-2}\text{M}$ solution of α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 12) was prepared by dissolving 5.8 mg in 325 μl NANOpure™ water. A 10 μl aliquot of this solution was added to 990 μl of $3 \times 10^{-4}\text{M}$ SmCl_3 (in 0.1N HCl) spiked with Sm-153. The pH was then brought to 7.0 with NaOH. This solution was split into two fractions of 500 μl each, and one fraction was heated for 1 hour at 100 °C. Both heated and non-heated samples were tested to determine the percent of metal as a complex by the cation exchange method described previously. The solution was then passed through SP-Sephadex™ resin. Complexation was again determined as the purified samples. The results are shown in the following table.

	Heated		Non-Heated	
	Purified	Not Purified	Purified	Not Purified
% Complex	96	95	97	89

Example 44

pH Stability of α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex.

The heated and non-heated purified samples from Example 43 were each split into 2 x 250 μl aliquots. One aliquot was adjusted with HCl and the other with NaOH to give the pH's represented in the table following. After the desired pH was reached the sample was allowed to stand for 10 min and the amount of metal as a complex was determined by the cation exchange method described previously. The results are shown in the following table.

pH	% Complex	
	Heated	Not heated
2	95	94
3	97	95
5	98	97
7	96	97
9	98	96
11	97	98
13	100	99

Example 45

Preparation of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium(III) complex.

A 3×10^{-2} M solution of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (prepared by the procedure of Example 11) was prepared by dissolving 7.9 mg in 484 μ l NANOpure™ water. A 10 μ l aliquot of this solution was added to 950 μ l of 3×10^{-4} M SmCl_3 (in 0.1N HCl) spiked with Sm-153. The pH was then brought to 7.0 with NaOH. This solution was split into two fractions of 500 μ l each, and one fraction was heated for 1 hour at 100 °C. Both heated and non-heated samples were tested to determine the percent of metal as a complex using the cation exchange method described previously. The solutions were passed through SP-Sephadex™ resin. Complexation was then determined on the purified samples using the general procedure described hereinbefore. The results are shown in the following table.

	Heated		Non-Heated	
	Purified	Not Purified	Purified	Not Purified
% Complex	100	72	100	46

Example 46

pH Stability of α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, Sm(III) complex.

The heated and non-heated purified sample from Example 45 were each divided into two aliquots. One aliquot was adjusted with dilute HCl and the other with NaOH to give the pH's represented in the table following. After the desired pH was reached the sample was allowed to stand for 10 min and the amount of metal as a complex was determined by the cation exchange method described previously. The results are shown in the following table:

pH	Purified	
	Heat	No heat
2	99	99
4	100	100
5	100	100
7	100	100
9	100	100
11	100	100
13	100	100

Example 47

Preparation of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, samarium(III) complex.

A 0.0219M (100 μ l, 2.195 mmole) solution of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid (prepared by the procedure of Example 4) was contacted with a 0.043M (25 μ l, 1.075 mmole) solution of Sm(OAc)₃. The reaction mixture was subjected to anion exchange HPLC (Q-Sepharose™ column, 1 cm x 25 cm, flow rate = 2 ml/min, eluted with 0-1M NH₄OAc gradient over 30 min). After one hour, the complex (retention time = 11.1 min) had formed in 77 percent yield (area percent) and was completely resolved from the ligand peak (retention time = 15.5 min). Additional evidence that the title complex had formed was obtained on silica gel TLC (Solvent System 4) whereby the ligand has an R_f = 0.59 and the complex has an R_f = 0.00.

Example 48

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, samarium(III) complex.

A solution, 0.022M (100 μ l, 2.2 μ mole), of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid (prepared by the procedure of Example 5) was contacted with a 0.043M (6.4 μ l, 0.275 μ mole) solution of Sm(OAc)₃. The reaction mixture was subjected to anion exchange HPLC (Q-Sepharose™ column, 1 cm X 25 cm, flow rate = 2 ml/min, eluted with 0-0.25M NH₄OAc gradient over 30 min). After forty minutes, the complex (retention time = 9.0 min) had formed in 66 percent yield (area percent) and was completely resolved from the ligand peak (retention time = 18.5 min).

Example 49

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, samarium(III) complex and α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid, samarium(III) complex.

A 3 x 10⁻²M solution of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid pentahydrochloride and α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid pentachloride (prepared by the procedure of Example 5) was prepared by dissolving 4.2 mg in 300 μ l NANOpure™ water. A 10 μ l aliquot of this solution was added to 15 μ l of 2 x 10⁻²M SmCl₃ (in 0.1N HCl) spiked with Sm-153. The volume was brought to 1 ml by addition of water. The pH was then brought to 7.0 with NaOH. This solution was split into two fractions of 500 μ l each, and one fraction was heated for 1 hour at 100 °C.

Both heated and non-heated samples were tested to determine the percent of metal as a complex by the cation exchange method described previously. The results showed 89% and 96% of the metal as a complex for the heated and non-heated samples, respectively.

Example 50

pH Stability of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, samarium(III) complex, and α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid, samarium(III) complex.

The heated sample from Example 49 and the non-heated sample were each divided into 2 aliquots. One aliquot was adjusted with HCl and the other with NaOH to give the pH's represented in the table following. Once the desired pH was reached, the samples were allowed to stand for 10 minutes and the percent of metal as complex was determined by the cation exchange method described previously. The results are shown in the following table:

pH	% Complex	
	Heated	Not heated
1	86	83
3	99	99
5	99	99
7	89	96
9	99	98
11	96	89
13	97	97

Example 51

Preparation of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, samarium(III) complex.

Thirteen microliters of 2×10^{-2} M solution of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid (prepared by the procedure of Example 4) 15 μ l of SmCl_3 (0.02M 0.1N HCl) and 2 μ l of Sm-153, had the volume brought to 1 ml with NANOpure™ water. The pH was adjusted to 7.0 with NaOH. This solution was split into two fractions of 500 μ l each, and one fraction was heated for 1 hour at 100 °C. Both heated and non-heated samples were tested to determine the percent of metal as a complex using the cation exchange procedure described previously. The results showed 74% and 42% of the metal as a complex for the heated and non-heated samples, respectively.

Example 52

pH Stability of α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, samarium(III) complex.

The heated sample of Example 51 was divided into 2 aliquots. One aliquot was adjusted with HCl and the other with NaOH to give the pH values represented in the following table. After the desired pH was reached the sample was allowed to stand for 10 minutes and the amount of metal as a complex was determined by the cation exchange method described previously. The results are shown in the following table.

pH	% Complex
1	67
2	70
4	70
5	72
7	74
9	76
10	76
12	75
13	76

Example 53

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, lutetium(III) complex.

A solution of EA-DO3A (prepared by the procedure of Example 17) was made by dissolving the solid in water. The final concentration was 0.01M.

A lutetium chloride solution was prepared by dissolving lutetium chloride in 0.1N HCl. The resultant concentration was 3.9 mg/ml of lutetium chloride (0.01M Lu). Lutetium-177 was used as the tracer.

A volume of 30 μ l of 0.01M Lu solution was added to 2 μ l of Lu-177 solution. A volume of 30 μ l of ligand was added and HEPES buffer (0.1M, pH = 7.6) was added to give a total volume of 1.0 ml. The resultant solution was 0.3 mM in ligand and lutetium.

The amount of Lu as a complex was determined to be 98 percent by the cation exchange method previously described.

Example 54

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, yttrium (III) complex; [Y(EA-DO3A)]

The ligand, 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 17) was converted to the mixed protonated-ammonium salt by chromatography on a strong cation exchange resin (SP-SephadexTM C-25, Pharmacia). The column was in the protonated form and washed with distilled water. A concentrated aqueous solution of the ligand was applied to the column, followed by washing the column with distilled water. The ligand was eluted from the column with 0.5M NH₄OH. The eluate was reduced to dryness on a rotary evaporator and dried in a vacuum oven.

A solution of Y(OAc)₃·4H₂O (0.203 g, 338.101 g/mole, 0.600 mmole) in 10 ml of distilled water was added to a warm solution of the mixed protonated-ammonium salt ligand (0.300 g, 499.615 g/mole, 0.600 mmole) in 10 ml of distilled water. The solution was brought to reflux and the pH was adjusted to about 7.0 with 0.1M NH₄OH. After 15 minutes at reflux, the solution was cooled and reduced to dryness on a rotary evaporator. Excess ammonium acetate was removed by heating the white solid in an oil bath at about 105°C under vacuum. The title product (which contained about one equivalent of ammonium acetate) was provided in a yield of 373 mg (99 percent) and was characterized by:

¹³C NMR (88°C, D₂O, 75 MHz)

24.4, 28.4, 51.2, 56.5, 57.2 (2C), 57.7, 67.2, 67.6, 120.5, 132.3, 135.2, 182.1, 182.5, 183.0;

Fast atom bombardment mass spectrum, m/e 552 (positive ion, [M+H]⁺), m/e 610 (negative ion, [M+OAc]⁻).

Example 55

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium complex; [Sm(EA-DO3A)].

When the procedure of Example 54 was repeated, using $\text{Sm}(\text{OAc})_3 \cdot 3\text{H}_2\text{O}$ (0.229 g, 381.531 g/mole, 0.600 mmole) in place of the $\text{Y}(\text{OAc})_3 \cdot 4\text{H}_2\text{O}$, the title product (which contained about one equivalent of ammonium acetate) was prepared in a yield of 408 mg (99 percent) and was characterized by:

^{13}C NMR (88 °C, D_2O , 75 MHz)

23.8, 27.9, 51.4, 57.5, 58.1 (3C), 68.7, 73.4, 120.3, 132.2, 134.7, 185.0, 186.8, 192.1;

Fast atom bombardment mass spectrum, m/e 615 (positive ion $[\text{M} + \text{H}^+]^+$, Sm isotope pattern), m/e 673 (negative ion, $[\text{M} + \text{OAc}^-]^-$, Sm isotope pattern).

Example 56

Preparation of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium(III) complex.

Ten microliters of 1.48 mg/100 μl (0.03M) solution of 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 20) was added to 15 μl of SmCl_3 solution (0.02M in 0.1N HCl) and spiked with Sm-153. The volume was brought to 1 ml with NANOpure™ water. This solution was divided into two 500 μl fractions, and one fraction was heated for 1 hour at 100 °C. Both heated and non-heated samples were tested to determine the percent of metal as a complex using the cation exchange procedure described hereinbefore. The results showed 81% and 43% of the metal as a complex for the heated and non-heated samples, respectively.

Example 57

pH Stability 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium(III) complex.

The heated sample of Example 56 was divided into two aliquots. One aliquot was adjusted with HCl and the other with NaOH to give the pH's represented in the following table. After the desired pH was reached the sample was allowed to stand for 10 min and the amount of metal as a complex was determined by the cation exchange method described previously. The results are shown in the following table.

pH	% Complex
1	20
2	37
4	49
5	58
7	81
9	62
10	64
12	64
13	54

Example 58

Preparation of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium(III) complex.

Three μl of 4.65 mg/100 ml solution of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 20) was added to a solution of 15 μl of SmCl_3 - (0.02M in 0.1N HCl) previously spiked with Sm-153. The volume was brought to 1 ml with NANOpure™ water. The pH of the solution was adjusted to 7.3 using NaOH. The solution was divided into two aliquots, and one aliquot was heated at 100°C for 1 hour. Both heated and non-heated samples were tested to determine the percent of metal as a complex by the cation exchange method described previously. The results showed 96% and 93% of the metal as a complex for the heated and non-heated samples, respectively.

Example 59

pH Stability of 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium(III) complex.

The heated and non-heated samples of Example 58 were each divided into two aliquots. One aliquot of each sample was adjusted with HCl and the other with NaOH to give the pH's shown in the following table. After the desired pH was reached, the sample was allowed to stand for 10 minutes and the amount of metal as a complex was determined by the cation exchange method described previously. The results are shown in the following table.

pH	% Complex	
	Heated	Not Heated
1	44	13
3	50	42
5	73	55
7	96	93
9	98	98
11	89	90
13	28	37

Example 60

Preparation of 1-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium(III) complex.

A $3 \times 10^{-2}\text{M}$ solution of 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 25) was prepared by dissolving 3.9 mg in 260 μl NANOpure™ water. A 10 μl aliquot of this solution was added to 15 μl of $2 \times 10^{-2}\text{M}$ SmCl_3 (in H_2O) spiked with 10 μl of Sm-153 in 0.1N HCl ($3 \times 10^{-4}\text{M}$). The volume was brought to one milliliter by addition of 965 μl DI H_2O . The pH was then brought to 7.0 with NaOH. This solution was split into two fractions of 500 μl each, and one fraction was heated for 1 hour at 100°C. Both heated and non-heated samples were tested to determine the percent of metal as a complex by the previously described cation exchange method. The results showed 71 percent and 74 percent of the metal as a complex for the heated and non-heated samples, respectively.

Example 61

pH Stability of 1-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium-(III) complex.

The heated and non-heated sample of Example 60 were each split into two 250 μ l aliquots. One aliquot was adjusted with HCl and the other with NaOH to give the pH's represented in the table following. After the desired pH was reached the sample was allowed to stand for 10 minutes and the amount of metal as a complex was determined by the cation exchange method described previously. The results are shown in the following table.

pH	% Complex	
	Heated	Not heated
1	65	51
3	98	93
5	99	99
7	100	100
9	100	99
11	100	96
13	100	98

Example 62

Preparation of 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium-(III) complex.

A 3×10^{-2} M solution of 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid (prepared by the procedure of Example 26) was prepared by dissolving 3.2 mg in 100 μ l NANOpure™ water. A 10 μ l aliquot of this solution (diluted to 20 μ l) was added to 15 μ l of 2×10^{-2} M SmCl_3 (in H_2O) spiked with Sm-153. The volume was brought to 1 ml by addition of 950 μ l DI H_2O and 20 μ l 0.1N NaOH. This solution was split into two fractions of 500 μ l each, and one fraction was heated for 1 hour at 100 °C. Both heated and non-heated samples, were tested to determine the percent of metal as a complex by the cation exchange method described previously. When less than 95 percent of the metal was complexed, the solution was passed through SP-Sephadex™ resin. The percent of metal as a complex was again determined. The results are shown in the following table.

	Heated		Non-Heated	
	Purified	Not Purified	Purified	Not Purified
% Complex	99	68	99	21

Example 63

pH Stability of 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium-(III) complex.

The heated and non-heated purified chromatographed samples from Example 62 were each split into two 250 μ l aliquots. One 250 μ l aliquot was adjusted with HCl and the other with NaOH to give the pH's represented in the table following. Once the desired pH was reached, samples were allowed to stand for 10 minutes and the percent of metal as complex was determined by the cation exchange method described

previously. The results are as shown in the following table:

pH	% Complex	
	Heated	Not heated
1	60	56
2	74	76
3	84	87
5	97	98
7	99	99
9	99	98
11	99	99
13	99	99

Methods of Use-Biology

Bifunctional chelants having within the same molecule a metal chelating group such as DTPA and a reactive linker (the aryl amine) are known to be capable of being covalently attached to various target directed biomolecules having specificity for cancer or tumor cell epitopes or antigens. Radionuclide complexes of such conjugates are useful in diagnostic and/or therapeutic applications as means of conveying the radionuclide to a cancer or tumor cell. [Reference, Meares et al., *Anal. Biochem.* 142, 68-78 (1984); see also discussion on pp. 215-216 of *J. Protein Chem.*, 3(2) (1984) by Meares and Goodwin, more recent references are Meares, U.S. Patent 4,678,677, issued July, 7, 1987, Warshawsky et al. U.S. Patent 4,652,519, issued March 24, 1987 and Brechbiel, et al., *Inorg. Chem.*, 25, 2772-2781 (1986)]. Conceivably, the product, employing radioactive or luminescent metals, may also be used for *in vitro* immunoassays.

Background Information

The utility of the labeled antibodies depends on a number of factors, for example: 1) the specificity of the antibody, 2) the inertness or stability of the complex under use conditions (i.e., serum stability), and 3) the labeling technique such that the integrity of the antibody may have the same specificity and immunoreactivity as that of the native molecule.

Stability or inertness of the complex is of utmost importance to the effectiveness of the radionuclide-antibody conjugate as a diagnostic and/or therapeutic agent. Kinetic lability can lead to instability, e.g. in the serum, and the dissociation of the radionuclide from the complex. Thus it diminishes the diagnostic and therapeutic effectiveness. In addition, it poses a greater potential for general radiation damage to normal tissue. [Cole, et al., *J. Nucl. Med.*, 28, 83-90 (1987)].

Linking of Radionuclide to Antibody

Attachment of radionuclide to antibody can be carried out by either linking the BFC to the antibody via procedures well known in the art, followed by chelation of the radionuclide under conditions compatible with the antibody, or alternatively, conjugation of the antibody to preformed (ambient or elevated temperature) metal - BFC complex. [Meares et al. *Acc. Chem. Res.* 17, 202-209 (1984)]. Examples are provided for the latter.

Example ZA (Comparative)

Conjugation of 1-(4-isothiocyanatobenzyl)diethylenetriaminepentaacetic acid, samarium-153 complex to IgG and F(ab')₂ of CC-49; [¹⁵³Sm(SCN-Bz-DTPA)]-IgG and [¹⁵³Sm(SCN-Bz-DTPA)]-F(ab')₂ fragment.

The 1-(4-isothiocyanatobenzyl)diethylenetriaminepentaacetic acid, samarium-153 complex [¹⁵³Sm(SCN-Bz-DTPA)] was prepared by mixing 150 μ l ¹⁵³Sm in 0.1N HCl with 9 μ l of 1-(4-isothiocyanatobenzyl)-

diethylenetriaminepentaacetic acid to which was added HEPES buffer (0.5M pH 8.9, approximately 30 μ l) to bring the pH to about 6. To conjugate, 22.5×10^{-9} moles of IgG or F(ab')₂ of CC-49 (about 1×10^{-4} M concentration of protein in 50 mmole HEPES, pH 8.5) was mixed with the ¹⁵³Sm complex, and the pH was adjusted to 8.9 by addition of a sodium carbonate solution (1.0M, 12-15 μ l). Conjugation was carried out at room temperature (about 25°C) for about 3 hours. The ¹⁵³Sm complex labeled IgG or F(ab')₂ was isolated and characterized similarly as described in Example XII.

Experimental for Biology

Examples I, & II and Comparative Examples A-D IN VIVO SCREENING OF BIFUNCTIONAL CHELATES.

The stability of certain rare earth chelates has been examined by *in vivo* testing in animals. For example, Rosoff, et al. in the *International Journal of Applied Radiation and Isotopes* 14, 129-135 (1963) report on the distribution of radioactive rare earth chelates in mice for certain aminocarboxylic acids. It was found that *in vivo* "the competition between the chelating agent and body constituents (inorganic and organic) for the rare-earth ion, determines its deposition and excretion." The strong rare-earth chelates are believed to dissociate very little and be excreted, while the weak and intermediate strength chelates dissociate more readily and thus are deposited in organs such as the liver. However, concentration of radionuclide in the liver is not always due to weak complex formation, but in some cases is due to the affinity that the metal chelate has for the liver. Chelates have, in fact, been prepared and utilized for the evaluation of liver function [Fritzberg, Alan R., *Radiopharmaceuticals: Progress and Clinical Perspectives*, Vol. 1, 1986; U.S. Patents 4,088,747 and 4,091,088].

The biodistribution of the yttrium and samarium chelates of the compound of Example 3 were determined and the percent dose in the liver was used as an *in vivo* screening procedure to qualitatively estimate the stability of the chelates. Chelates of NTA and EDTA are included for comparison. Also samarium was injected as samarium chloride in unchelated form as a control.

Sprague-Dawley rats weighing from 150 to 200 g were purchased from Charles River Laboratories. These animals were placed in cages and fed water and food *ad libitum*. The animals were acclimated for at least five days before use. Prior to injection of complex, the animals were placed under a heat lamp (15 to 30 minutes) to dilate the tail vein. The animal was then placed in a restraining cage, the tail cleaned with an alcohol swipe, and the animal injected (50 to 200 μ l) via the tail vein. After injection, the animal was placed in another cage for two hours after which time the animal was sacrificed by cervical dislocation. The animal was then dissected, the parts rinsed with deionized H₂O, patted dry, and weighed into a tared counting vial. At least three standards of the same material as injected were prepared and counted with the animal tissues. Percent of dose is the number of counts in the organ divided by the number of counts in the standard multiplied by 100 (see the following Table).

Table

Biodistribution Data			
Example No.	Ligand of Ex. *	Metal	% Injected Dose in Liver
I	3	Y	0.17
II	3	Sm	0.29
(A)	EDTA	Sm	8.4
(B)	EDTA	Sm	4.4
(C)	NTA	Sm	8.6
(D)	SmCl ₃	Sm	39

*Complexes were prepared at ligand/metal ratios of 1:1 for Examples I and II; at 5:1 for Example A; and at about 300:1 for Examples B and C.

Examples III and E.

The 1:1 complexes of yttrium (spiked with a tracer amount of ^{90}Y) with the ligand of Example 3, which is BA-DOTA, and EDTA (Comparative E) were prepared by methods described previously. Several 100 μL aliquots were then transferred to separate centrifuge tubes. Excess Y(III) was added such that the total volume change is minimized and the time noted. One-half hour after metal addition, the percent complex was determined by the cation exchange method described previously and this was compared to the original amount of complex. The percent complex versus added metal gives an indication as to the lability of the ligand-metal complex. The results are given in the Table.

TABLE

Complex Study		
Metal/Ligand Molar Ratio	% Complex	
	BA-DOTA	EDTA
1	80	98
10	--	86
100	--	78
250	88	48
500	80	16

Example IV

Preparation of 1-[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium-153 complex; [$^{153}\text{Sm}(\text{SCN-EA-DO3A})$] complex.

To 100 μl of ^{153}Sm in 0.1N HCl was added 5 μl of SCN-EA-DO3A (5mM in 50 mM HEPES, pH 8.2) (prepared in Example 18). This was mixed on a vortex mixer, and HEPES buffer (0.5M, pH 8.9) was added gradually (about 25 μl total) to adjust the pH to around 7. The progress of the chelation was monitored by HPLC on GF-250 column (HPLC System I). Yields around 50% were obtained.

Example V

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex; [$^{153}\text{Sm}(\text{PA-DOTA})$] complex.

To 150 microliters (μl) of ^{153}Sm solution in 0.1N HCl (approximately 4.6 mCi) were added 1 μl of sodium acetate (2.0M) and 9 μl of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid mixed ammonium, potassium salt (5 mmole in 50 mmole HEPES, pH 8.5, prepared by the procedure of Example 8). The mixture was mechanically shaken and titrated gradually with a HEPES buffer (0.5M, pH 8.9; approximately 31 μl added) to pH 7. This was then heated at 98 °C in a sand bath for 1 hour. Upon termination, 5 μl of the mixture was used for analysis on a Mono-QTM column on HPLC System II, eluted with a gradient solvent system (0-15 minutes, from 0 to 100% B; where A = water, and B = 1.0M ammonium acetate and 0.1 mmole EDTA). Yields of 85-95% based on ^{153}Sm were obtained. The α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex thus prepared was characterized by comparison with the identical non-radioactive sample, by their respective chromatographic behavior on Mono-QTM and GF-250 columns. Further evidence of the presence of the radioactive complex was determined by the conversion to the isothiocyanato derivative and its subsequent conjugation to antibody. The complex has also been prepared without heating (at ambient temperature) by incubation for 6 to 18 hours to result in 70-80% yield.

Example VI

Preparation of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex; [$^{153}\text{Sm}(\text{SCN-PA-DOTA})$] complex.

To the reaction mixture obtained in Example V was added 2 μl of HEPES buffer (0.5M, pH 8.9), 2 μl of thiophosgene and 0.2 ml of chloroform. The mixture was mechanically shaken vigorously 2 or 3 times for a few seconds each time. The chloroform layer was discarded and the aqueous layer which contained mainly the desired product was saved and further purified. The yield of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex, based on ^{153}Sm activity measurement by HPLC on GF-250 column using HPLC System I, was 85-90%. To purify, the aqueous layer was passed through a Sep-PakTM C-18 cartridge and eluted with 90% acetonitrile in water. The first 300 μl of effluent was discarded, and the SCN-derivative which came off in the next 900 μl was characterized by HPLC on GF-250. The recovery of the ^{153}Sm activity was better than 90%. The bulk of the solvent was then evaporated over a period of 1.5 to 2 hours, and the residue was used for conjugation to antibody.

Example VII

Preparation of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex; [^{153}Sm (BA-DOTA)] complex.

To 200 μl of ^{153}Sm solution in 0.1N HCl were added 1 μl of sodium acetate (2.0M) and 12 μl of α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (5 mmole in 50 mmole HEPES, pH 8.5) (prepared by the procedure of Example 3). The mixture was mechanically shaken and titrated gradually with a HEPES buffer (0.5M, pH 8.9; approximately 38 μl added) to pH 7. This was then heated at 98 °C in a sand bath for 1 hour. Upon termination, 5 μl of the mixture was used for analysis on a Mono-QTM column (HPLC System II, eluted with a gradient solvent system: 0-15 minutes, from 0 to 100% B; where A = water, and B = 1.0M ammonium acetate and 0.1 mmole EDTA). In general, yields of 85-95% based on ^{153}Sm were obtained. The complex has also been prepared by incubation of the mixture at room temperature for 12-18 hours which resulted in a yield of the title product of 80-90%. The α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex thus prepared was characterized by comparison with an authentic sample by their chromatographic behavior on the Mono-QTM column, conversion to the isothiocyanato derivative and its subsequent conjugation to antibody.

Example VIII

Preparation of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex; [$^{153}\text{Sm}(\text{SCN-BA-DOTA})$] complex.

To the reaction mixture obtained in Example VII were added 2 μl of HEPES buffer (0.5M, pH 8.9), 2 μl of thiophosgene and 0.2 ml of chloroform. It was mechanically shaken vigorously 2 or 3 times for a few seconds each time. The chloroform layer was discarded and the aqueous layer which contained mainly the desired product was saved and further purified. The yield of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex, as analyzed by HPLC on GF-250 column based on the ^{153}Sm activity using the HPLC System I was over 90%. To purify, the aqueous layer was passed through a Sep-PakTM C-18 cartridge and eluted with 90 percent acetonitrile in water. The first 0.3 ml of effluent was discarded, and the desired product came off in the next 1.2 ml, with 86-93% recovery. The bulk of the solvent was then evaporated over a period of about 2 hours, and the residue was used for conjugation to antibody.

Example IX

Conjugation of 1-[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium-153 complex to antibody; [$^{153}\text{Sm}(\text{SCN-EA-DO3A})$]-IgG conjugate.

The antibody used was CC-49, a murine monoclonal IgG that binds to an epitope of TAG-72, a tumor associated antigen. To conjugate, 187 μl of the antibody solution ($1.20 \times 10^{-4}\text{M}$ in 50 mM HEPES, pH 8.5) was mixed with 4.5×10^{-8} moles of $^{153}\text{Sm}(\text{SCN-EA-DO3A})$ prepared as described in Example IV followed

by addition of a sodium carbonate solution (1.0M) to raise the pH to 8.9. The reaction was allowed to continue for 2 hours at room temperature. Upon termination, the ^{153}Sm labeled IgG was isolated by centrifugal gel filtration on Sephadex G-25 (2.2 ml) disposable columns, and further purified by HPLC on GF-250 column, eluted with a citrate buffer (0.25M, pH 7.4). The fractions contained the labeled IgG were
 5 pooled, concentrated and exchanged (3x) into PBS by use of CentriconTM concentrators. The specific activity of ^{153}Sm labeled IgG thus prepared was about 0.16 $\mu\text{Ci}/\mu\text{g}$ of protein. The integrity of the [^{153}Sm -(EA-DO3A)]-IgG preparation was verified by HPLC (Sivakoff, S.I., *BioChrom.* 1(1), 42-48 (1986)) and standard biochemical procedures, e.g. sodium dodecyl sulfate:polyacrylamide gel electrophoresis and autoradiography, and solid phase radioimmunoassay (RIA). [See David Colcher et al., *Cancer Res.* 43, 736-
 10 742 (1983).]

Example X

Conjugation of 1-[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, samarium-153 complex to Fragment F(ab')₂ of CC-49; [^{153}Sm (SCN-EA-DO3A)]-Fragment F(ab')₂ of CC-49.
 15

The F(ab')₂ fragment of CC-49, prepared by enzymatic digestion according to the procedure described by Lamoyi and Nisonoff [E. Lamoyi and A Nisonoff, *J. Immunol. Methods*, 56, 235-243 (1983)], (225 μl of 1×10^{-4} M in 50 mM HEPES, pH 8.5) was mixed with 2.9×10^{-8} moles of ^{153}Sm (SCN-EA-DO3A) prepared as
 20 described in Example IV. Sodium carbonate (1.0M, about 5 μl) was added to bring the pH to 8.9, and the reaction was continued for about 2.5 hours. Upon termination, the ^{153}Sm labeled fragment was isolated and characterized as described in Example IX.

Example XI (A and B)

In vivo localization of [^{153}Sm (EA-DO3A)]-IgG and [^{153}Sm (EA-DO3A)]-F(ab')₂.
 25

The utility of the ^{153}Sm (EA-DO3A) labeled IgG (from Example IX) and F(ab')₂ of CC-49 (from Example X) was demonstrated by the uptake of the labeled materials by human tumor xenograft in athymic mice. Thus female athymic (Nu/Nu, CD1 background) mice were inoculated subcutaneously (S.C.) (0.1 ml/source) with the human colon carcinoma cell line, LS-174 T (approximately 1×10^6 cells/animal). Approximately 2 weeks after inoculation, each animal was injected via the tail vein with about 2 μCi (15-30 μg) of ^{153}Sm labeled antibody in PBS. The mice were sacrificed at various time intervals. After exsanguination, the tumor and selected tissues were excised and weighed, and radioactivity was measured in a gamma counter. The
 30 counts per minute (CPM) of ^{153}Sm in each tissue was determined and expressed as CPM per gram of tissue per injected dose multiplied by 100 (% injected dose/gram). Results are shown in Figures 1-14 and Tables IA and IB. The ^{153}Sm (SCN-Bz-DTPA) labeled IgG and F(ab')₂ of CC-49 (from Example ZA) were included in the study for comparison. Results are shown in Table IC and ID.

Example XII

Conjugation of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex to antibody; [^{153}Sm (SCN-PA-DOTA)]-IgG conjugate.

The antibody used was CC-49, a murine monoclonal IgG that binds to an epitope of TAG-72, a tumor associated antigen. To conjugate, 178 μl of the antibody solution (1.26×10^{-4} M in 50 mmole HEPES, pH 8.5) was mixed with 3.4×10^{-8} moles of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium complex (prepared as described in Example VI), followed by addition of a sodium carbonate solution (1.0M, about 17 μl) to raise the pH to about 8.9. The
 45 reaction was allowed to continue for 2 hours at room temperature. Upon termination, the ^{153}Sm labeled IgG was isolated by centrifugal gel filtration on SephadexTM G-25 (2.2 ml) disposable columns, and further purified by HPLC on GF-250 column, eluted with a citrate buffer (0.25M, pH 7.4). The fractions which contained the labeled IgG were pooled, concentrated and exchanged (3 times) into PBS by use of CentriconTM concentrators. The specific activity of the ^{153}Sm labeled IgG thus prepared was about 0.16
 50 $\mu\text{Ci}/\mu\text{g}$ of protein. The homogeneity and integrity of the α -[2-(4-isothiocyanatophenyl)-ethyl]-1,4,7,10-tetraazacyclo dodecane-1,4,7,10-tetraacetic acid, samarium-153 complex-IgG preparation was verified by HPLC and standard biochemical procedures as Example IX.

The following Example is an alternative for making labeled antibody conjugates, which involves first conjugation of the BFC to the antibody, and the subsequent chelation to yield the radionuclide-BFC labeled Ab.

5 Example XIIA

Preparation of α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid - Ab CC₄₉ conjugate (IgG CC₄₉-PA-DOTA); and the subsequent chelation with ¹⁵³Sm to form ¹⁵³Sm-labeled antibody (IgG CC₄₉-PA-DOTA-¹⁵³Sm).

10

The ¹⁵³Sm(PA-DOTA) labeled antibody can be prepared by first coupling the BFC, e.g. α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (SCN-PA-DOTA), to an antibody at pH 8-9, followed by chelation with ¹⁵³Sm at pH 6 at room temperature for several hours.

15

In a typical experiment, CC-49 was concentrated and exchanged three times into a carbonate buffer (50 mM, pH 9.1) in an Amicon Concentrator (30 K molecular weight cut off) to leave a solution with antibody concentration greater than 1.5×10^{-4} M. To the formed conjugate was added 161 μ l of the antibody solution containing 25×10^{-9} mole of IgG CC-49 mixed with 5 μ l of the SCN-PA-DOTA (5mM concentration in the same carbonate buffer, prepared by the procedure of Example 18). The mixture was allowed to stand at room temperature for 5 hours, and terminated by filtration through the 30 K membrane in the Amicon
20 Concentrator at 5000 rpm on the Sorvall RT centrifuge. The concentrate was further washed with 2 ml of a 0.25M DTPA solution in PBS at pH 7.4 and five times (2 ml each time) with MES buffer (20 mM, pH 5.8); centrifuged for 30 min. at each wash. At the end, the PA-DOTA-IgG conjugate was concentrated to a minimum volume (about 100 μ l) and its integrity confirmed by HPLC analysis on a GF-250 column. To the purified conjugate was added a mixture of ¹⁵³Sm in 0.1N HCl (50 μ l) and 20 μ l of a MES buffer (1.0M, pH
25 6), mixed on a vortex mixer, and allowed to stand at room temperature overnight. Upon termination, the amount of ¹⁵³Sm incorporated, estimated by HPLC analysis, was 0.22 BFC-Sm/antibody. That ¹⁵³Sm was associated with the antibody through chelation with the BFC, which was linked covalently to the antibody, and not due to non-specific binding was demonstrated by the control experiment. In the control experiment, IgG CC-49 solution which was mixed with ¹⁵³Sm mixture under identical conditions, and isolated according
30 to the same procedure, showed no appreciable amount of ¹⁵³Sm associated with it.

Example XIII

35

Conjugation of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex to fragment F(ab')₂ of CC-49; [¹⁵³Sm(SCN-PA-DOTA)]-F(ab')₂ fragment.

40

The F(ab')₂ fragment of CC-49 [prepared by enzymatic digestion according to the procedure described by (E. Lamoyi et al., *J. Immunol. Methods* 56, 235-243 (1983)), (225 μ l of 1×10^{-4} M in 50 mmole HEPES, pH 8.5) was mixed with 2.9×10^{-8} moles of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex prepared as described in Example VI. Sodium carbonate (1.0M, about 9 μ l) was added to bring the pH to around 8.9, and the reaction was continued for about 2.5 hours. The ¹⁵³Sm labeled fragment was isolated and characterized as described in Example IX. The specific activity was around 0.4 μ Ci/ μ g.

45

Example XIV(A and B)

In vivo localization of the conjugate of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex labeled IgG and F(ab')₂; [¹⁵³Sm(SCN-PA-DOTA)]-IgG and [¹⁵³Sm(SCN-PA-DOTA)]-F(ab')₂.

50

The utility of the conjugate of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex labeled IgG and F(ab')₂ of CC-49 (from Example XII and XIII) was demonstrated by the uptake of the labeled materials by human tumor xenograft in athymic mice. The biolocalization was determined using the procedure described in Example XI. Results are shown in
55 Figures 1 to 7 for IgG and Figures 8 to 14 for F(ab')₂, also Tables IIA and IIB.

The conjugate of ¹⁵³Sm(SCN-Bz-DTPA) with IgG and F(ab')₂ labeled materials (prepared from Example ZA) were included in the study for comparison. (See Tables IC and ID.)

Example XV

Conjugation of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex to Antibody; [$^{153}\text{Sm}(\text{SCN-BA-DOTA})$]-IgG.

Whole IgG of CC-49 (174 μl of $1.2 \times 10^{-4}\text{M}$ in 0.25M HEPES, pH 8.7) was mixed with 2.0×10^{-8} moles of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 (2.8 mCi) prepared per Example VIII, followed by addition of a sodium carbonate solution (1.0M, about 2 μl) to maintain the pH about 8.7. The reaction was allowed to continue for 2.5 hours at room temperature. Upon termination, the ^{153}Sm labeled IgG was isolated by centrifugal gel filtration on SephadexTM G-25 (2.2 ml) disposable columns, and further purified by HPLC on GF-250 column, eluted with a citrate buffer (0.025M, pH 7.4). The fractions which contained the labeled IgG were pooled, concentrated and exchanged (3 times) into PBS by use of CentriconTM concentrators. Homogeneity and integrity of the samarium-153 labeled IgG preparation was verified by HPLC and standard biochemical procedures as described in Example IX.

Example XVI

Conjugation of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex to Fragment F(ab')₂ of CC-49; [$^{153}\text{Sm}(\text{SCN-BA-DOTA})$]-fragment F(ab')₂ of CC-49.

The F(ab')₂ of CC-49 (84 μl of $2.4 \times 10^{-4}\text{M}$ in 0.25M HEPES buffer, pH 8.7), prepared by enzymatic digestion according to the procedure described by Lamoyi et al. was mixed with 2.1×10^{-8} moles of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex (prepared by the procedure of Example VIII). Sodium carbonate (1.0M, about 2 μl) was added to bring the pH to 8.7, and the reaction was continued for about 2 hours. Upon termination, the ^{153}Sm labeled fragment was isolated and characterized as described in Example XII.

Example XVII (A and B)

In vivo localization of α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex labeled IgG (Example XV) and F(ab')₂ (Example XVI); [$^{153}\text{Sm}(\text{BA-DOTA})$]-IgG and [$^{153}\text{Sm}(\text{BA-DOTA})$]-F(ab')₂.

The *in vivo* study was conducted according to the protocol described in Example XIV or XI, and results are shown in Figures 1 to 14 and Tables IIIA and IIIB.

Example XVIII (A and B)

In vivo localization of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid ^{177}Lu -complex labeled IgG and F(ab')₂; [$^{177}\text{Lu}(\text{PA-DOTA})$]-IgG and [$^{177}\text{Lu}(\text{PA-DOTA})$]-F(ab')₂.

The title compounds were prepared and *in vivo* studies were conducted according to protocol described in Examples V, VI, XII and XIII and results are shown in Tables IVA and IVB.

Example XIX (A and B)

In vivo localization of α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, yttrium-90 complex labeled IgG and F(ab')₂; [$^{90}\text{Y}(\text{PA-DOTA})$]-IgG and [$^{90}\text{Y}(\text{PA-DOTA})$]-F(ab')₂.

The titled compounds were prepared and *in vivo* studies were conducted according to protocol described in Example V, VI, XII and XIII and results are shown in Tables VA and VB.

Example XX

In vitro pH stability of [$^{153}\text{Sm}(\text{BFC})$]-CC-49-IgG and [$^{177}\text{Lu}(\text{BFC})$]-CC-49-IgG.

5 The [$^{153}\text{Sm}(\text{BFC})$]-CC-49-IgG or [$^{177}\text{Lu}(\text{BFC})$]-CC-49-IgG was allowed to stand in a 0.2M NaOAc buffer at pH 6.0, 4.0 and 2.8, at room temperature, at a protein concentration of about 5×10^{-6} M with about 0.5 complex /antibody. Samples were withdrawn at certain time intervals and analyzed by HPLC (GF-250 column) for the dissociation of ^{153}Sm or ^{177}Lu activity from the protein. In general, the study was carried out for five days or until 90% of the radioisotopes have become dissociated. Results are expressed as the initial
10 rate of loss of the radioisotope per day. The results demonstrated the superior stability of $^{153}\text{Sm}(\text{PA-DOTA})$, $^{177}\text{Lu}(\text{PA-DOTA})$, $^{153}\text{Sm}(\text{BA-DOTA})$, and $^{153}\text{Sm}(\text{OMe-BA-DOTA})$ as compared with the standard [$^{153}\text{Sm}(\text{Bz-DTPA})$] complex at acidic pH. Results are shown in the table below.

This greater stability of the complexes also correlates well with the more rapid clearance of ^{153}Sm and ^{177}Lu from the body and non-target tissue (e.g. kidney, liver); see Figs. 1-14.

BFC	Isotope	% Loss of Isotope/Day pH		
		6.0	4.0	2.8
Bz-DTPA	^{153}Sm	<2	35	35
PA-DOTA	^{153}Sm	<2	<2	<2
BA-DOTA	^{153}Sm	<2	<2	<2
PA-DOTA	^{177}Lu	<2	<2	<2
MeOBA-DOTA	^{153}Sm	<2	<2	<2
EA-DO3A	^{153}Sm	<2	90	95

Example XXI (A and B)

In vivo localization of α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, samarium-153 complex labeled IgG and F(ab')₂; [$^{153}\text{Sm}(\text{MeO-BA-DOTA})$]-IgG and [$^{153}\text{Sm}(\text{MeO-BA-DOTA})$]-F(ab')₂.
35

The titled compounds were prepared according to the procedures described in Examples V, VI, and XII, and the *in vivo* studies were conducted by the procedures of Example XI. Results are shown in Tables VIA and VIB.

Example XXII

Preparation of $^{153}\text{Sm}(\text{PA-DOTMA})$ complex.

45 To 100 microliters (μl) of ^{153}Sm (0.3×10^{-3} M in 0.01N HCl; 5 mCi) were added 6 μl of PA-DOTMA (high R_f diastereomer prepared by the procedure of Example 29; 5 mM in Milli-Q™ water), 20 μl of a MES buffer (1.0M, pH 6) and 1 μl of a HEPES buffer (0.5M, pH 8.7). This solution was mixed on a vortex mixer and the final pH of the mixture was about 6. After heating the mixture at 90 °C for 30 min, it was analyzed by HPLC on GF-250 column for chelation efficiency, and yields of 90 percent or better, were generally
50 obtained. Characterization of the ^{153}Sm -PA-DOTMA complex was provided by comparison with the non-radioactive-Sm complex, which had been synthesized and characterized independently in Example 42.

Example XXIII

55 Preparation of $^{153}\text{Sm}(\text{SCN-PA-DOTMA})$.

To the ^{153}Sm -PA-DOTMA complex prepared in Example XXII and having been cooled for 20 min. at room temperature was added 10 μl of a 1 percent thiophosgene solution in 90 percent acetonitrile-water medium. The mixture was mixed vigorously on a vortex mixer, and allowed to stand at room temperature for

between 5 to 20 min. The reaction took place instantaneously as monitored by HPLC analysis. To purified the activated ^{153}Sm complex, it was extracted 3 times (200 μl each) with chloroform and the aqueous layer was passed through as PRP cartridge (PRP cartridge = Mini-cleanTM cartridge from Alltech Associates, Deerfield, IL) pretreated with 5 ml of methanol and 5 ml of a MES buffer (20 mM, pH 5.8). After being washed with 5 ml of the MES buffer and 5 ml of Milli-QTM water, it was extracted with 900 μl of 90 percent acetonitrile to recover approximately 90 percent of the ^{153}Sm activity, with the first 100 μl discarded. The mixture was evaporated to dryness under reduced pressure at temperatures below 40°C, and the residue which contained mostly the title product was used for conjugation to the antibody.

10 Example XXIV

Conjugation of $^{153}\text{Sm}(\text{SCN-PA-DOTMA})$ to IgG and $\text{F(ab')}_2 \text{CC}_{49}$.

In general, the antibody was concentrated and exchanged into a carbonate buffer (pH 9.1 or 9.5, 50 mM) in a CentriconTM concentrator (30 K molecular weight cut off) to result in a protein concentration of $1.5 \times 10^{-4} \text{ M}$ or greater. To conjugate, a small volume of the carbonate buffer, that which is required to bring the final protein concentration to $1.5 \times 10^{-4} \text{ M}$, is added to the isothiocyanato derivative of $^{153}\text{Sm}(\text{PA-DOTMA})$ (prepared in Example XXIII), followed by the concentrated antibody, of equimolar to the BFC- ^{153}Sm complex. The mixture was mixed on a vortex mixer and allowed to react at room temperature for 1 hour or until 40 to 50 percent of ^{153}Sm become bound to the antibody, as shown by HPLC analysis on a GF-250 column. The labeled antibody was isolated by two consecutive centrifugal gel filtrations on SephadexTM G-25 columns (2.2 ml). The homogeneity, integrity and immunoreactivity of the labeled antibody was analyzed by HPLC analysis and standard biochemical techniques described previously.

25 Example XXV

Biodistribution Studies of $^{153}\text{Sm}(\text{PA-DOTMA})$ labeled IgG and $\text{F(ab')}_2 \text{CC}_{49}$.

The studies were conducted in the similar fashion as described in Example XI. Results are shown in Tables VIIA and VIIB. See Figures 15-28.

Example XXVI

Preparation of $^{177}\text{Lu}(\text{PA-DOTMA})$ Complex.

To 30 μl of ^{177}Lu ($6 \times 10^{-3} \text{ M}$ in 1.1N HCl; 4 mCi) was added 36 μl of PA-DOTMA (prepared by the procedure of Example 29); 5 mM solution in milli-QTM water). The solution was mixed on a vortex mixer, and 115 μl of a MES buffer (1.0M, pH 6.0) was added to neutralize the solution to about pH 5.5 to 6. This was heated at 90°C for 30 minutes to result in better than 90% chelation. The mixture was passed through a PRP cartridge pretreated with 5 ml of methanol and 5 ml of a MES buffer (20 mM, pH 5.8). It was washed with 5 ml of Milli-QTM waste. The complex extracted in 1000 μl of 90% acetonitrile accounted for 78% of the starting ^{177}Lu activity (discarding the first 100 μl of the extract).

Example XXVII

Preparation of $^{177}\text{Lu}(\text{SCN-PA-DOTMA})$.

To the $^{177}\text{Lu}(\text{PA-DOTMA})$ complex in 90% acetonitrile obtained in Example XXVI was added 6 μl of a 10% thiophosgen in 90% acetonitrile. This solution was mixed on a vortex mixer, and after 20 minutes, analyzed for the formation of the isothiocyanate derivative by HPLC. The reaction is, in general, quantitative. Removal of the solvent and the excess thiophosgene was accomplished by evaporation under reduced pressure at temperatures below 40°C for 1 to 2 hours. The residue was used for conjugation with the antibody.

Example XXVIII

Conjugation of $^{177}\text{Lu}(\text{PA-DOTMA})$ to IgG CC49.

- 5 The antibody IgG CC49 in a carbonate buffer (50 mM, pH 9.1) was mixed with an equimolar quantity of the isothiocyanato derivative (prepared by the procedure of from Example XXVII) in the same buffer. The reaction was continued for 70 min at an antibody concentration of $1.5 \times 10^{-4}\text{M}$, and the labeled antibody was isolated and characterized according to procedures described in Example XXIV.

10 Example XXIX

Long Term Biodistribution Studies of $^{177}\text{Lu}(\text{PA-DOTMA})$ Labeled IgG CC49.

- 15 The long term animal test was conducted with the ^{177}Lu labeled antibody, taking advantage of its long half-life time (161 hours). It was carried out in Balb/c mice over a three week period, according to protocols described in Example XI. Results are shown in Table VIIA. See Figures 29-34.

In the Figures, which plot the data from the following tables, the symbols used were:

Conjugates	Symbol	Table	Figures	Bio. Example
$^{153}\text{Sm}(\text{EA-DO3A})\text{-IgG-CC-49}$	Δ	IA	1-7	IX
$^{153}\text{Sm}(\text{EA-DO3A})\text{-F(ab'2)-CC-49}$	Δ	IB	8-14	X
$^{153}\text{Sm}(\text{Bz-DTPA})\text{-IgG-CC-49}$	$\bigcirc$	IC	1-7	ZA
$^{153}\text{Sm}(\text{Bz-DTPA})\text{-F(ab'2)-CC-49}$	$\bigcirc$	ID	8-14	ZA
$^{153}\text{Sm}(\text{PA-DOTA})\text{-IgG-CC-49}$	$\square$	IIA	1-7	XII
$^{153}\text{Sm}(\text{PA-DOTA})\text{-F(ab'2)-CC-49}$	$\square$	IIB	8-14	XIII
$^{153}\text{Sm}(\text{BA-DOTA})\text{-IgG-CC-49}$	$\blacksquare$	IIIA	1-7	XV
$^{153}\text{Sm}(\text{BA-DOTA})\text{-F(ab'2)-CC-49}$	$\blacksquare$	IIIB	8-14	XVI
$^{153}\text{Sm}(\text{PA-DOTMA})\text{-IgG-CC-49}$	∇	VIIA	15-21	XXIV
$^{153}\text{Sm}(\text{PA-DOTMA})\text{-F(ab'2)-CC-49}$	∇	VIIIB	22-28	XXV
$^{177}\text{Lu}(\text{PA-DOTMA})\text{-IgG-CC-49}$	$\bullet$	VIIIA	29-34	XXIX

In the accompanying Figures the data represents the following:

FIGURE	ORGAN	TITLE
1	Blood	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
2	Liver	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
3	Spleen	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
4	Kidney	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
5	Tumor	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
6	Femur	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
7	Whole Body Retention	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
8	Blood	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
9	Liver	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
10	Kidney	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
11	Spleen	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
12	Tumor	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
13	Femur	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
14	Whole Body Retention	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
15	Blood	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
16	Liver	Biodistribution of ^{153}Sm -BFC-CC49-IgG*

* in nude mice bearing LS174-T tumor

** in Balb/c mice

FIGURE	ORGAN	TITLE
17	Spleen	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
18	Kidney	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
19	Tumor	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
20	Femur	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
21	Whole Body Retention	Biodistribution of ^{153}Sm -BFC-CC49-IgG*
22	Blood	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
23	Liver	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
24	Spleen	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
25	Kidney	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
26	Tumor	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
27	Femur	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
28	Whole Body Retention	Biodistribution of ^{153}Sm -BFC-CC49-F(ab') ₂ *
29	Blood	Biodistribution of ^{177}Lu -BFC-CC49-IgG**
30	Liver	Biodistribution of ^{177}Lu -BFC-CC49-IgG**
31	Spleen	Biodistribution of ^{177}Lu -BFC-CC49-IgG**
32	Kidney	Biodistribution of ^{177}Lu -BFC-CC49-IgG**

* in nude mice bearing LS174-T tumor

** in Balb/c mice

FIGURE	ORGAN	TITLE
33	Femur	Biodistribution of ^{177}Lu -BFC-CC49-IgG**
34	Whole Body Retention	Biodistribution of ^{177}Lu -BFC-CC49-IgG**

* in nude mice bearing LS174-T tumor

** in Blab/c mice

Table IA
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{EA-D03A})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	23.08	2.36	17.25	2.40	11.32	*1.06	5.78	*1.69
Liver	6.95	1.28	6.58	0.56	6.58	0.62	7.62	0.98
Spleen	5.17	0.59	5.06	0.96	4.62	0.72	4.92	1.72
Kidney	4.59	0.68	4.01	0.53	3.74	0.76	3.20	0.59
Tumor	11.44	4.13	27.19	2.12	53.01	9.10	59.01	13.23
Femur	-	-	2.58	0.52	2.78	0.20	4.22	0.17
Tumor Wt. g	0.16	0.09	0.20	0.09	0.30	0.17	0.52	0.34

*n=4

Table IB
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{EA}-\text{DO3A})$]-F(ab')₂ CC-49
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	13.36	0.97	1.52	0.42	0.20	0.06	0.05	0.01
Liver	7.13	0.65	8.69	1.3	8.08	0.8	8.11	0.92
Spleen	5.94	1.24	4.96	1.27	4.81	0.99	5.08	1.43
Kidney	41.60	5.60	64.32	7.88	64.12	12.59	37.63	6.57
Tumor	12.95	1.73	16.84	3.71	11.50	4.56	5.53	1.46
Femur	2.74	0.65	2.45	0.26	3.41	0.75	4.99	0.29
Tumor Wt. g	0.25	0.20	0.34	0.25	0.28	0.14	0.44	0.27

*n = 4

TABLE IC
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{Bz-DTPA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 (n=10)

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	24.97	3.07	16.05	2.45	12.81	2.69	5.84	1.76
Liver	7.74	1.35	6.01	0.95	5.68	1.07	5.43	1.45
Spleen	5.69	1.04	4.86	1.36	4.72	0.81	3.78	0.63
Kidney	4.52	1.04	3.96	0.88	3.81	0.61	3.76	0.31
Tumor	13.90	3.18	42.86	14.03	46.33	8.30	61.32	19.80
Femur	2.36	0.12	1.95	0.38	1.97	0.75	2.25	0.50
Whole Body Retention	85.45	4.74	80.16	3.58	77.36	5.32	70.60	3.32

TABLE ID
 BIODISTRIBUTION of ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{Bz-DTPA})$]-F(ab') $_2$ -CC-49
 % INJECTED DOSE/GRAM
 (n=10)

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	15.03	1.84	2.13	0.48	0.29	0.08	0.04	0.02
Liver	7.07	0.88	7.03	1.24	6.14	0.67	5.80	1.0
Spleen	4.50	0.97	4.22	1.15	3.65	0.62	3.24	0.42
Kidney	46.74	8.69	80.37	12.76	61.42	9.85	30.17	5.17
Tumor	15.95	3.88	18.68	5.10	15.55	2.80	6.54	1.14
Femur	2.48	0.24	1.77	0.28	2.48	0.40	2.85	0.35
Whole Body Retention	83.23	4.64	72.17	6.20	58.89	4.40	40.53	2.48

Table IIA
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{PA-DOTA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 (n = 5)

Organ	5.5 hr		25 hr		49 hr		121 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	24.65	2.89	16.18	2.10	13.22	1.08	7.34	4.23
Liver	8.25	1.14	5.92	0.41	5.59	0.52	4.16	1.03
Spleen	6.98	1.70	5.05	0.20	4.27	0.52	4.31	1.95
Kidney	4.01	0.86	3.16	0.55	3.44	0.39	3.26	0.57
Tumor	14.31	2.00	42.81	8.83	72.59	21.53	78.36	33.29
Femur	2.69	0.41	2.02	0.32	1.69	0.37	1.44	0.69
Tumor Wt. g	0.27	0.29	0.34	0.26	0.16	0.13	0.28	0.15

*n = 4

Table IIB
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [^{153}Sm (PA-DOTA)]-F(ab')₂-CC-49
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	14.74	0.76	2.25	0.45	0.33	0.08	0.02	*0.01
Liver	7.83	0.82	5.04	*0.32	4.48	0.41	1.61	0.26
Spleen	4.80	0.84	3.27	0.45	3.33	0.82	1.41	0.10
Kidney	51.28	5.83	78.78	8.40	61.91	8.17	19.79	5.75
Tumor	17.99	6.53	22.09	6.07	14.93	1.91	5.81	1.21
Femur	2.43	0.17	1.34	0.25	1.17	0.22	0.51	0.18
Tumor Wt. g	0.20	0.08	0.19	0.13	0.29	0.10	0.45	0.23

*n = 4

Table IIIA
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{BA-DOTA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5.5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	24.89	1.29	17.53	1.79	13.61	0.53	8.68	2.94
Liver	7.53	0.79	5.46	*0.62	5.35	*0.40	4.26	*0.58
Spleen	5.52	0.61	5.83	1.07	4.78	0.35	3.76	1.28
Kidney	3.84	0.50	3.96	*0.36	3.17	*0.57	3.49	*0.43
Tumor	14.41	2.96	38.65	6.45	55.84	11.43	85.26	25.70
Femur	2.54	0.34	1.81	0.35	2.08	0.30	1.17	0.48
Tumor Wt. g	0.34	0.22	0.13	0.04	0.20	0.06	0.20	0.15

*n = 4

Table IIIB
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{BA-DOTA})\text{]-F(ab')}_2\text{-CC-49}$
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5 hr		24 hr		49 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	15.62	40.86	2.42	0.42	0.28	0.08	0.03	0.01
Liver	7.69	0.71	6.18	0.82	4.22	0.88	1.56	0.25
Spleen	4.93	0.63	3.94	0.52	2.88	*0.36	1.43	*0.30
Kidney	50.29	4.99	77.19	*3.22	60.44	7.71	24.98	2.67
Tumor	14.58	2.24	24.28	1.81	17.65	3.25	7.39	2.54
Femur	2.13	0.18	1.36	0.15	1.03	0.28	0.68	0.35
Tumor Wt. g	0.17	0.02	0.12	0.07	0.17	0.06	0.12	0.10

*n = 4

TABLE IVA
 BIODISTRIBUTION OF ^{177}Lu INJECTED AS [$^{177}\text{Lu}(\text{PA-DOTA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5 hr		24 hr		48 hr		121 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	26.50	4.32	*19.43	*1.59	14.89	1.72	11.73	3.14
Liver	7.47	1.34	*6.74	*0.31	5.11	0.41	4.71	0.63
Spleen	6.29	1.56	*5.47	*1.24	4.43	0.31	5.35	1.82
Kidney	3.86	*0.85	*4.13	*0.45	3.13	*0.30	3.57	1.15
Tumor	11.94	3.39	*49.03	*3.21	56.36	*4.24	92.05	*15.32
Femur	2.79	1.08	*2.15	*0.30	2.11	0.39	1.32	*0.19
Tumor Wt. g	0.13	0.09	*0.09	*0.02	0.08	*0.02	0.10	*0.04

n = 4

TABLE IVB
 BIODISTRIBUTION OF ^{177}Lu INJECTED AS [$^{177}\text{Lu}(\text{PA-DOTA})$]-F(ab')₂-CC-49
 % INJECTED DOSE/GRAM
 (n=5)

Organ	5 hr		24 hr		48 hr		121 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	12.65	*0.87	*2.40	*0.18	0.47	0.08	0.04	*0.02
Liver	5.39	*0.27	*5.54	*0.31	3.25	0.45	1.30	*0.24
Spleen	4.21	0.95	*3.60	*0.55	2.60	0.74	1.33	0.27
Kidney	38.82	6.42	*74.40	*6.13	48.45	6.47	15.88	2.13
Tumor	10.79	2.69	*19.10	*2.08	15.44	2.35	4.69	0.78
Femur	2.13	0.83	*1.45	*0.33	0.95	0.43	0.41	*0.25
Tumor Wt. g	0.10	0.03	*0.10	*0.03	0.07	0.06	0.19	0.13

*n = 4

Table VA
 BIODISTRIBUTION OF ^{90}Y INJECTED AS [^{90}Y (PA-DOTA)]-IgG-CC-49
 % INJECTED DOSE/GRAM

Organ	n=3 5 hr		n=5 24 hr		n=5 48 hr		n=5 120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	25.15	1.96	18.22	2.69	15.85	*0.81	6.62	2.94
Liver	8.25	0.42	7.61	0.77	6.52	0.96	5.68	1.26
Spleen	5.13	0.26	5.50	0.72	5.66	*0.38	4.19	0.88
Kidney	3.04	0.58	4.35	1.06	3.66	0.60	2.47	0.33
Tumor	12.45	0.79	44.73	**1.43	68.41	*5.48	70.32	21.34
Femur	2.05	0.15	1.81	0.54	1.79	*0.11	1.04	0.45

*n = 4

**n = 3

TABLE VB
 BIODISTRIBUTION OF ^{90}Y INJECTED AS [^{90}Y (PA-DOTA)]-F(ab')₂-CC-49
 % INJECTED DOSE/GRAM
 n=5

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	13.87	1.19	2.22	0.46	0.31	0.03	0.04	0.02
Liver	6.98	0.84	4.13	*0.42	3.01	0.33	1.56	0.32
Spleen	4.27	0.35	3.44	*0.49	2.54	0.71	1.33	0.54
Kidney	44.60	5.52	65.40	5.92	42.31	4.67	12.36	2.60
Tumor	13.08	*0.88	22.44	4.77	17.84	5.09	5.83	2.09
Femur	1.42	0.22	1.12	0.14	0.39	0.18	0.38	0.12

*n = 4

Table VIA
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{MeO-BA-DOTA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM

Organ	n=5 5 hr		n=5 24 hr		n=5 48 hr		n=4 120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	21.06	*0.75	14.57	1.18	13.98	0.78	9.49	1.06
Liver	9.00	1.93	6.22	0.68	6.32	0.41	4.34	0.27
Spleen	5.39	1.40	4.00	0.64	4.69	0.93	4.53	0.41
Kidney	3.48	*0.24	3.23	0.41	3.47	0.45	3.47	0.19
Tumor	18.00	8.91	41.64	3.37	58.34	5.78	69.26	21.92
Femur	2.54	0.27	1.94	0.32	2.05	0.29	1.29	0.16

*n = 4

TABLE VIB
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{MeO}-\text{BA}-\text{DOTA})$]-F(ab')₂-CC-49
 % INJECTED DOSE/GRAM
 n=5

Organ	n=5 5 hr		n=5 24 hr		n=4 48 hr		n=5 120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	12.98	*0.36	2.13	0.37	0.31	0.03	0.05	0.03
Liver	6.58	**0.16	6.77	1.13	4.62	0.27	2.46	0.55
Spleen	3.95	0.57	3.53	0.76	2.59	0.40	2.18	0.68
Kidney	47.07	5.85	74.16	8.03	47.68	6.29	24.25	2.74
Tumor	12.59	3.96	17.30	1.68	12.33	*0.80	5.21	*0.63
Femur	2.61	0.30	1.61	0.31	0.90	0.03	0.68	**0.10

*n = 3

**n = 4

Table VIIA
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{PA-DOTMA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 n=5

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	23.00	1.11	15.58	1.37	11.78	0.36	8.98	0.91
Liver	7.71	0.51	6.68	1.16	4.98	0.32	4.72	0.69
Spleen	6.05	1.03	5.28	0.45	3.66	0.35	3.72	0.53
Kidney	3.68	*0.23	4.07	0.41	3.95	*0.15	3.17	0.86
Tumor	10.40	2.10	31.85	*2.52	44.70	10.03	66.45	13.60
Femur	1.94	0.20	1.66	0.11	1.49	0.10	1.22	0.08

*n = 4

TABLE VIIB
 BIODISTRIBUTION OF ^{153}Sm INJECTED AS [$^{153}\text{Sm}(\text{PA-DOTMA})$]-F(ab')₂-CC-49
 % INJECTED DOSE/GRAM
 n=5

Organ	5 hr		24 hr		48 hr		120 hr	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	12.85	1.06	1.84	0.13	0.38	0.09	0.07	0.02
Liver	6.12	0.52	5.36	0.49	3.65	0.65	1.93	0.32
Spleen	4.08	0.37	2.88	0.31	2.09	0.62	0.92	*0.13
Kidney	50.28	5.26	58.59	5.63	45.29	7.04	14.85	*0.66
Tumor	11.85	2.23	14.27	1.02	15.26	2.83	4.90	2.01
Femur	1.77	0.06	1.06	0.13	0.96	0.28	0.79	0.24

*n = 4

Table VIIIA
 BIODISTRIBUTION OF ^{177}Lu INJECTED AS [$^{177}\text{Lu}(\text{PA-DOTMA})$]-IgG-CC-49
 % INJECTED DOSE/GRAM
 n=5


Organ	24 hr		48 hr		7 day		14 day		21 day	
	AVG	STD	AVG	STD	AVG	STD	AVG	STD	AVG	STD
Blood	31.35	2.66	30.25	*0.93	22.47	0.71	14.13	1.05	9.88	0.83
Liver	10.01	0.46	11.22	1.47	8.61	0.79	6.76	0.67	5.41	1.39
Spleen	10.13	1.16	10.91	*0.67	10.79	*0.36	8.09	0.98	6.03	0.55
Kidney	6.52	2.06	6.14	0.76	5.21	0.97	3.44	0.43	2.17	0.22
Femur	3.97	0.45	4.53	0.38	2.84	0.16	2.36	0.23	1.67	0.30

*n = 4

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB IT, LI, LU, NL, SE

1. A compound of the formula:



(IA)

wherein:

each Q is independently hydrogen or (CHR⁵)_pCO₂R;

Q¹ is hydrogen or (CHR⁵)_wCO₂R;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl;

with the proviso that at least two of the sum of Q and Q¹ must be other than hydrogen, and that when Q¹ is hydrogen, R³ must be other than hydrogen;

each R⁵ independently is hydrogen or C₁-C₄ alkyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

p = 1 or 2;

r = 0 or 1;

w = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R² is hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido or carboxyl;

R³ is C₁-C₄ alkoxy, -OCH₂CO₂H, hydroxy or hydrogen;

R⁴ is hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

with the proviso that R² and R⁴ cannot both be hydrogen but one of R² and R⁴ must be hydrogen; or a pharmaceutically acceptable salt thereof.

2. A compound of Claim 1 wherein the pharmaceutically acceptable salt is selected from the group consisting of potassium, sodium, lithium, ammonium, calcium, magnesium, and hydrogen chloride adducts.


3. A compound of Claim 1 wherein R and R⁵ are both hydrogen.

4. A compound of Claim 1 wherein R is hydrogen, R⁵ is methyl and w is 0.

5. A compound of Claim 1 wherein n and r are both 0 and m is 0 through 5.

6. A compound of Claim 1 wherein R³ and R⁴ are hydrogen.


7. A compound of Claim 1 having the formula



wherein:

- each Q is independently hydrogen or $\text{CHR}^5\text{CO}_2\text{R}$;
 each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;
 with the proviso that at least two of Q must be other than hydrogen;
 m is an integer from 0 to 5 inclusive;
 R^2 , R^3 , R^4 and R^5 are defined as in claim 1;
 or a pharmaceutically acceptable salt thereof.

8. A compound of Claim 7 having the formula




wherein:

- Q and m are defined as in claim 7;
 R^2 is selected from the group consisting of nitro, amino, isothiocyanato, and semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl; or
 a pharmaceutically acceptable salt thereof.

9. The compound of Claim 7 which is 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester.
- 50 10. The compound of Claim 8 which is 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt.
11. The compound of Claim 8 which is 1-[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.
- 55 12. The compound of Claim 8 which is 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt.

13. A compound of Claim 7 having the formula



wherein:

Q and m are defined as in claim 7;

R³ is selected from the group consisting of C₁-C₄ alkoxy, -OCH₂CO₂H and hydroxy;

R⁴ is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;


or

a pharmaceutically acceptable salt thereof.

14. The compound of Claim 13 which is 1-(2-methoxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

15. The compound of Claim 13 which is 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid.

16. A compound of Claim 1 having the formula



wherein:

each Q is independently hydrogen or CHR⁵CO₂R;

Q¹ is hydrogen or (CHR⁵)_wCO₂R;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl;

with the proviso that at least two of the sum of Q and Q¹ must be other than hydrogen and one of Q is hydrogen;

m is an integer from 0 to 5 inclusive;

w is 0 or 1;


R², R³, R⁴ and R⁵ are defined as in claim 1;

or

a pharmaceutically acceptable salt thereof.

17. The compound of Claim 16 wherein R is hydrogen and R⁵ is C₁₋₄ alkyl.

18. A compound of Claim 16 having the formula



20 wherein:

each Q is independently hydrogen or CHR⁵CO₂R;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl;


with the proviso that at least one Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;

25 R², R³, R⁴ and R⁵ are defined as in claim 1;

or a pharmaceutically acceptable salt thereof.

19. A compound of Claim 18 having the formula



45 wherein:

Q and m are defined as in claim 18;

R² is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

or

50 a pharmaceutically acceptable salt thereof.

20. The compound of Claim 19 which is α-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic 4,7,10-tris-(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester.

21. The compound of Claim 19 which is α-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic 4,7,10-tris-(R-methylacetic) acid.

22. The compound of Claim 19 which is α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(R-methylacetic) acid.

23. The compound of Claim 19 which is α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid, mixed ammonium, potassium salt.

24. The compound of Claim 19 which is α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid.

25. The compound of Claim 19 which is α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid.

26. The compound of Claim 19 which is α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.

27. The compound of Claim 19 which is α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.


28. The compound of Claim 19 which is α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester.

29. The compound of Claim 19 which is α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester.

30. The compound of Claim 19 which is α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.

31. The compound of Claim 19 which is α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.

32. A compound of Claim 18 having the formula



wherein:

each Q is independently hydrogen or $\text{CHR}^5\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least one Q must be other than hydrogen;

m is an integer from 0 to 5 inclusive;

R^3 and R^5 are defined as in claim 1;

R^4 is selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

or

a pharmaceutically acceptable salt thereof.

33. The compound of Claim 32 which is α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.
34. The compound of Claim 32 which is α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetraamonium salt.
35. The compound of Claim 32 which is α -(2-methoxy-5-isothiocynatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetraamonium salt.
36. A complex comprising a compound according to claim 1 or a pharmaceutically acceptable salt thereof; complexed with an ion of a metal selected from La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y and Sc.
37. A complex of Claim 36 wherein the metal is Sm, Lu or Y.
38. A complex of Claim 36 wherein the metal is ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc , or ^{142}Pr .
39. A complex of Claim 38 wherein the metal is ^{153}Sm , ^{177}Lu or ^{90}Y .
40. A complex of any one of Claims 36 through 39 wherein the metal ion is complexed with the compound of Claims 7, 8, 13, 16, 18, 19 or 32.
41. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 26.
42. A complex of Claim 41 wherein the metal is ^{153}Sm or ^{90}Y .
43. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 27.
44. A complex of Claim 43 wherein the metal is ^{153}Sm .
45. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 30 or 31.
46. A complex of Claim 45 wherein the metal is ^{153}Sm , ^{90}Y or ^{177}Lu .
47. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 32.
48. A complex of Claim 47 wherein the metal is ^{153}Sm , ^{90}Y or ^{177}Lu .
49. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 10.
50. A complex of Claim 49 wherein the metal is ^{153}Sm .
51. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 11.
52. A complex of Claim 51 wherein the metal is ^{153}Sm .
53. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 34.
54. A complex of Claim 53 wherein the metal is ^{153}Sm .
55. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 35.
56. A complex of Claim 55 wherein the metal is ^{153}Sm .
57. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 21.
58. A complex of Claim 57 wherein the metal is ^{153}Sm , ^{90}Y or ^{177}Lu .

59. A complex of Claim 38 wherein the metal ion is complexed with the compound of Claim 22.

60. A complex of Claim 73 wherein the metal is ^{153}Sm , ^{90}Y or ^{177}Lu .

61. A conjugate comprising a complex according to claim 36 with the proviso that R^2 and R^4 cannot be nitro, and covalently attached to an antibody or antibody fragment.

62. The conjugate of Claim 61 wherein the metal is ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc or ^{142}Pr .

63. The conjugate of Claim 62 wherein the metal is ^{153}Sm , ^{90}Y or ^{177}Lu .

64. A conjugate of any one of Claims 61 through 63 wherein the antibody or antibody fragment is a monoclonal antibody or fragment thereof.

65. A conjugate of Claim 64 wherein the antibody or antibody fragment is CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ or B72.3 (ATCC HB 8108).

66. A conjugate of any one of Claims 61 through 63 wherein the compound is as defined in Claims 7, 8, 13, 16, 18, 19 or 32.

67. A conjugate of any one of Claims 62 through 65 wherein the metal ion is ^{153}Sm complexed with the compound of Claim 27.

68. A conjugate of any one of Claims 62 through 65 wherein the metal ion is ^{153}Sm , ^{90}Y or ^{177}Lu complexed with the compound of Claim 32.

69. A conjugate of any one of Claims 62 through 65 wherein the metal ion is ^{153}Sm complexed with the compound of Claim 35.

70. A conjugate of any one of Claims 62 through 65 wherein the metal ion is ^{153}Sm , ^{90}Y or ^{177}Lu complexed with the compound of Claim 22.

71. A conjugate comprising a compound according to claim 1 or a pharmaceutically acceptable salt thereof; covalently attached to an antibody or antibody fragment.


72. A conjugate of Claim 71 wherein the antibody or antibody fragment is a monoclonal antibody or fragment thereof.

73. A conjugate of Claim 72 wherein the antibody or antibody fragment is CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ or B72.3 (ATCC HB 8108).

74. A conjugate of any one of Claims 71 through 73 wherein the compound is as defined in Claims 7, 8, 13, 16, 18, 19 or 32.

75. A pharmaceutical formulation comprising a conjugate as claimed in any one of Claims 61 through 70 with a pharmaceutically acceptable carrier.

76. A process for preparing a compound of the formula



wherein:

each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen, and that when Q^1 is hydrogen, R^3 must be other than hydrogen;

each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

p = 1 or 2;

r = 0 or 1;

w = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, hydroxy and hydrogen;


R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or

a pharmaceutically acceptable salt thereof;

which comprises any one of the following steps:

(A) reacting a compound of the formula



wherein:


X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond;
with an α -halocarboxylic acid ester of the formula



where each R independently is benzyl or $\text{C}_1\text{-C}_4$ alkyl;
in the presence of an organic solvent and base, at a temperature of from 0°C to reflux,
to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is $\text{C}_1\text{-C}_4$ alkyl or benzyl; and

R^2 is nitro; or

(B) reacting the product of step (A) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is $\text{C}_1\text{-C}_4$ alkyl or benzyl; and

R^2 is amino; or


(C) reacting a compound of the formula (IIIA) as defined above with a hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions, followed by hydrogen chloride gas in the presence of a solvent to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 are hydrogen; and

R^2 is amino; or

(D) reacting the product of step (C) with an α -halocarboxylic acid ester of the formula



where each R independently is benzyl or $\text{C}_1\text{-C}_4$ alkyl;
in the presence of an organic solvent and base, at a temperature of from 0°C to reflux,
to provide the compounds of formula (IA) wherein:


X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is $\text{C}_1\text{-C}_4$ alkyl or benzyl; and

R^2 is amino; or

(E) reacting a compound of the formula



(VA)

wherein:

Q^1 is $(CHR^5)_wCO_2R$;

each R independently is benzyl or C_1-C_4 alkyl;

each R^5 independently is hydrogen or C_1-C_4 alkyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;


n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

w = 0 or 1;

with the proviso that the sum of r and w is 0 or 1;
with an α -halocarboxylic acid ester of the formula



where each R independently is benzyl or C_1-C_4 alkyl;
in the presence of an organic-solvent and base, at a temperature of from $0^\circ C$ to reflux,
to provide the compounds of formula (IA) wherein:

X, Y, Q^1 , m, n, r are defined as before;

R^3 , R^4 , R^5 are hydrogen;

R is C_1-C_4 alkyl or benzyl; and

R^2 is nitro; or

(F) reacting the product of step (E) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions, to provide the compounds of formula (IA) wherein:

X, Y, Q^1 , m, n, r are defined as before;

R^3 , R^4 , R^5 are hydrogen;

R is C_1-C_4 alkyl or benzyl; and

R^2 is amino; or

(G) reacting the product of either step (B) or (D) with a desterifying agent, such as a strong acid, to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is hydrogen; and

R^2 is amino; or

(H) reacting the product of step (F) with a desterifying agent, such as a strong acid, to provide the compounds of formula (IA) wherein:


X, Y, Q^1 , m, n, r are defined as before;

R^3 , R^4 , R^5 are hydrogen;

R is hydrogen; and

R^2 is amino; or

(I) reacting a compound of formula (VA) as defined above with an optically active α -benzene sulfonate of the formula



wherein R and R^5 are defined as before,

to provide the compounds of formula (IA) wherein:

X, Y, Q¹, R⁵, m, n, r are defined as before;

R³ and R⁴ are hydrogen;

R is C₁₋₄ alkyl or benzyl; and

R² is nitro; or

(J) reacting the product of step (I) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions, to provide the compounds of formula (IA) wherein:

X, Y, Q¹, R⁵, m, n, r are defined as before;

R³ and R⁴ are hydrogen;

R is C₁₋₄ alkyl or benzyl; and

R² is amino; or


(K) reacting the product of step (J) with a deesterifying agent, such as a strong acid, to provide the compounds of formula (IA) wherein.

X, Y, Q¹, R⁵, m, n, r are defined as before;

R³, R⁴ and R are hydrogen; and

R² is amino; or

(L) reacting a compound of the formula



(IVA)

wherein:

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;


R³ is C₁₋₄ alkoxy, -OCH₂CO₂H or hydroxy;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond; with an α -halocarboxylic acid of the formula



in the presence of an organic solvent and base, at a temperature of from 0°C to reflux, to provide the compounds of formula (IA) wherein:

X, Y, R³, m, n, r are defined as before;

Q¹ and R² are hydrogen; and

R⁴ is nitro; or


(M) reacting the product of step (L) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions to provide the compounds of formula (IA) wherein:

X, Y, R³, m, n, r are defined as before;

Q¹, R, R² and R⁵ are hydrogen; and

R² is amino; or

(N) reacting a compound of the formula



wherein:

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

Q¹ is CH₂CO₂R;


R is benzyl or C₁₋₄ alkyl;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond; with an α-halocarboxylic acid of the formula



where R is defined as before;

in the presence of an organic solvent and base, at a temperature of from 0 °C to reflux, to provide the compounds of formula (IA) wherein:

X, Y, Q¹, Q, m, n, r are defined as before;

R² is hydrogen; and

R⁴ is nitro; or

(O) reacting the product of step (N) with a desterifying agent, such as a strong acid, followed by a hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions to provide the compounds of formula (IA) wherein:

X, Y, Q¹, Q, m, n, r are defined as before;

R, R² and R⁵ are hydrogen; and

R² is amino; or


(P) reacting the product of step (G), (F), (K), (M) or (O) with thiophosgene to provide the corresponding compound wherein R² or R⁴ is isothiocyanato.

77. A process for preparing a mono-N-alkylated polyazamacrocyclic which comprises reacting an α-halocarboxylic-acid-ester with between one to five equivalents of a 1,4,7,10-tetraazacyclododecane in a solvent which will not promote a proton transfer.

78. The process of claim 77 wherein the α-halocarboxylic-acid-ester is d,l-2-bromo-4-(4-nitrophenyl)butanoic acid methylester, 2-bromo-2-(4-nitrophenyl)ethanoic acid-methylester, α-bromo-(2-methoxy-5-nitrophenyl)acetic-acid-methylester, or d,l-2-bromo-4-(4-nitrophenyl)-butanoic acid isopropylester.

Claims for the following Contracting States : ES, GR

1. A process for preparing a compound of the formula



wherein:

each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$; each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen, and that when Q^1 is hydrogen, R^3 must be other than hydrogen;

each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

p = 1 or 2;

r = 0 or 1;

w = 0 or 1;


with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl; R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, hydroxy and hydrogen;

R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof; which comprises any one of the following steps:

(A) reacting a compound of the formula



wherein:


X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond;
with an α -halocarboxylic acid ester of the formula



where each R independently is benzyl or C_1 - C_4 alkyl;
in the presence of an organic solvent and base, at a temperature of from 0°C to reflux,
to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is C_1 - C_4 alkyl or benzyl; and

R^2 is nitro; or

(B) reacting the product of step (A) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is C_1 - C_4 alkyl or benzyl; and

R^2 is amino; or


(C) reacting a compound of the formula (IIIA) as defined above with a hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions, followed by hydrogen chloride gas in the presence of a solvent to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 are hydrogen; and

R^2 is amino; or

(D) reacting the product of step (C) with an α -halocarboxylic acid ester of the formula



where each R independently is benzyl or C_1 - C_4 alkyl;
in the presence of an organic solvent and base, at a temperature of from 0°C to reflux,
to provide the compounds of formula (IA) wherein:


X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is C_1 - C_4 alkyl or benzyl; and

R^2 is amino; or

(E) reacting a compound of the formula



(VA)

wherein:

Q^1 is $(CHR^5)_wCO_2R$;

each R independently is benzyl or C_1-C_4 alkyl;

each R^5 independently is hydrogen or C_1-C_4 alkyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;


n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

w = 0 or 1;

with the proviso that the sum of r and w is 0 or 1;
with an α -halocarboxylic acid ester of the formula



where each R independently is benzyl or C_1-C_4 alkyl;

in the presence of an organic solvent and base, at a temperature of from $0^\circ C$ to reflux, to provide the compounds of formula (IA) wherein:

X, Y, Q^1 , m, n, r are defined as before;

R^3 , R^4 , R^5 are hydrogen;

R is C_1-C_4 alkyl or benzyl; and

R^2 is nitro; or

(F) reacting the product of step (E) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions, to provide the compounds of formula (IA) wherein:

X, Y, Q^1 , m, n, r are defined as before;

R^3 , R^4 , R^5 are hydrogen;

R is C_1-C_4 alkyl or benzyl; and

R^2 is amino; or

(G) reacting the product of either step (B) or (D) with a deesterifying agent, such as a strong acid, to provide the compounds of formula (IA) wherein:

X, Y, m, n, r are defined as before;

Q^1 , R^3 , R^4 , R^5 are hydrogen;

R is hydrogen; and

R^2 is amino; or

(H) reacting the product of step (F) with a deesterifying agent, such as a strong acid, to provide the compounds of formula (IA) wherein:


X, Y, Q^1 , m, n, r are defined as before;

R^3 , R^4 , R^5 are hydrogen;

R is hydrogen; and

R^2 is amino; or

(I) reacting a compound of formula (VA) as defined above with an optically active α -benzene sulfonate of the formula



wherein R and R^5 are defined as before,

to provide the compounds of formula (IA) wherein:

X, Y, Q¹, R⁵, m, n, r are defined as before;

R³ and R⁴ are hydrogen;

R is C₁₋₄ alkyl or benzyl; and

R² is nitro; or

(J) reacting the product of step (I) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions, to provide the compounds of formula (IA) wherein:

X, Y, Q¹, R⁵, m, n, r are defined as before;

R³ and R⁴ are hydrogen;

R is C₁₋₄ alkyl or benzyl; and

R² is amino; or


(K) reacting the product of step (J) with a deesterifying agent, such as a strong acid, to provide the compounds of formula (IA) wherein:

X, Y, Q¹, R⁵, m, n, r are defined as before;

R³, R⁴ and R are hydrogen; and

R² is amino; or

(L) reacting a compound of the formula



(IVA)

wherein:

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;


R³ is C₁₋₄ alkoxy, -OCH₂CO₂H or hydroxy;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond; with an α-halocarboxylic acid of the formula



in the presence of an organic solvent and base, at a temperature of from 0 °C to reflux, to provide the compounds of formula (IA) wherein:

X, Y, R³, m, n, r are defined as before;

Q¹ and R² are hydrogen; and

R⁴ is nitro; or


(M) reacting the product of step (L) with hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions to provide the compounds of formula (IA) wherein:

X, Y, R³, m, n, r are defined as before;

Q¹, R, R² and R⁵ are hydrogen; and

R² is amino; or

(N) reacting a compound of the formula



(VIII A)

wherein:

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

Q¹ is CH₂CO₂R;


R is benzyl or C₁₋₄ alkyl;

n is 0 or 1;

m is an integer from 0 to 10 inclusive;

r = 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond; with an α-halocarboxylic acid of the formula



where R is defined as before;

in the presence of an organic solvent and base, at a temperature of from 0 °C to reflux, to provide the compounds of formula (IA) wherein:

X, Y, Q¹, Q, m, n, r are defined as before;

R² is hydrogen; and

R⁴ is nitro; or

(O) reacting the product of step (N) with a desterifying agent, such as a strong acid, followed by a hydrogenation agent, such as Pd/C with hydrogen, under standard reaction conditions to provide the compounds of formula (IA) wherein:

X, Y, Q¹, Q, m, n, r are defined, as before;

R, R² and R⁵ are hydrogen; and

R² is amino; or

(P) reacting the product of step (G), (F), (K), (M) or (O) with thiophosgene to provide the corresponding compound wherein R² or R⁴ is isothiocyanato.

2. A process of Claim 1 for preparing a pharmaceutically acceptable salt of the compound of formula (IA) which comprises reacting the compound of formula (IA) with a reagent, by known methods, to obtain potassium, sodium, lithium, ammonium, calcium, magnesium, or hydrogen chloride adducts as the pharmaceutically acceptable salt.

3. The process of Claim 1(B) for preparing 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester which comprises reacting 1-[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester with hydrogen and palladium on carbon.

4. The process of Claim 1(G) for preparing 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt which comprises reacting 1-[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, 4,7,10-trimethyl ester with hydrochloric acid.

5. The process of Claim 1(P) for preparing 1-[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid which comprises reacting 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt with thiophosgene.
- 5 6. The process of Claim 1(A) for preparing 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid, hydrochloride salt which comprises reacting 1-(4-aminobenzyl)-1,4,7,10-tetraazacyclododecane, tetrahydrochloride salt with methylbromoacetate.
- 10 7. The process of Claim 1(M) for preparing 1-(2-methoxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid which comprises reacting 1-(2-methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid with hydrogen in a Parr bomb.
- 15 8. The process of Claim 1(M) for preparing 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid which comprises reacting N-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecane-4,7,10-triacetic acid with hydrogen and PtO_2 .
- 20 9. The process of Claim 1(J) for preparing α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic 4,7,10-tris-(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester which comprises reacting α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester with hydrogen and palladium on carbon.
- 25 10. The process of Claim 1(K) for preparing α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic 4,7,10-tris-(R-methylacetic) acid which comprises reacting α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(R-methylacetic) acid, 1-isopropyl-4,7,10-trimethyl ester with hydrochloric acid.
- 30 11. The process of Claim 1(P) for preparing α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic-4,7,10-tris-(R-methylacetic) acid which comprises reacting α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-(R,S)-acetic 4,7,10-tris-(R-methylacetic) acid with thiophosgene.
- 35 12. The process of Claim 1(F) for preparing α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid which comprises reacting α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid with hydrogen and PtO_2 .
- 40 13. The process of Claim 1(F) for preparing α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid which comprises reacting α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,10-triacetic acid with hydrogen and PtO_2 .
- 45 14. The process of Claim 1(F) for preparing α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid which comprises reacting α -(4-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid with hydrogen and PtO_2 .
- 50 15. The process of Claim 1(P) for preparing α -(4-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid which comprises reacting α -(4-aminophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid with thiophosgene.
- 55 16. The process of Claim 1(E) for preparing α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester which comprises reacting α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1-acetic acid, 1-methyl ester with methyl bromoacetate.
17. The process of Claim 1(F) for preparing α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester which comprises reacting α -[2-(4-nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester with hydrogen and palladium-on-carbon.
18. The process of Claim 1(H) for preparing α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid which comprises reacting α -[2-(4-aminophenyl)ethyl]-1,4,7,10-

tetraazacyclododecane-1,4,7,10-tetraacetic acid, 1,4,7,10-tetramethyl ester with hydrochloric acid.


19. The process of Claim 1(P) for preparing α -[2-(4-isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid which comprises reacting α -[2-(4-aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid with thiophosgene.
20. The process of Claim 1(O) for preparing α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7,10-tetraacetic acid which comprises reacting α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetramethyl ester with hydrochloric acid.
21. The process of Claim 1(O) for preparing α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7,10-tetraacetic acid, tetraammonium salt which comprises reacting α -(2-methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid with hydrogen and PtO_2 .
22. The process of Claim 1(P) for preparing α -(2-methoxy-5-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, tetraammonium salt which comprises reacting α -(2-methoxy-5-aminophenyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7,10-tetraacetic acid with thiophosgene.
23. A process for preparing a complex of a compound as claimed in any one of Claims 1 through 22 with an ion of a metal selected from La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y and Sc which comprises reacting the compound with the metal ion.
24. The process of Claim 23 wherein the metal is ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc , or ^{142}Pr .
25. The process of Claim 24 wherein the metal is ^{153}Sm , ^{177}Lu or ^{90}Y .
26. A process of any one of Claims 23 through 25 wherein the reaction is heated.
27. A process for preparing a conjugate containing a complex of a compound as claimed in any one of Claims 1 through 22 complexed with an ion of a metal selected from La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y and Sc, which comprises covalently attaching the complex to an antibody or antibody fragment.
28. The process of Claim 27 wherein the metal is ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc , or ^{142}Pr .
29. The process of Claim 28 wherein the metal is ^{153}Sm , ^{177}Lu or ^{90}Y .
30. A conjugate of any one of Claims 27 through 29 wherein the antibody or antibody fragment is a monoclonal antibody or fragment thereof.
31. A conjugate of Claim 30 wherein the antibody or antibody fragment is CC-49 (ATCC HB 9459), CC-49 F(ab')_2 , CC-83 (ATCC HB 9453), CC-83 F(ab')_2 or B72.3 (ATCC HB 8108).
32. A process for preparing a conjugate containing a compound as claimed in any one of Claims 1 through 22 covalently attached to an antibody or antibody fragment which comprises reacting the compound with an antibody or antibody fragment.
33. A conjugate of Claim 32 wherein the antibody or antibody fragment is a monoclonal antibody or fragment thereof.
34. A conjugate of Claim 33 wherein the antibody or antibody fragment is CC-49 (ATCC HB 9459), CC-49 F(ab')_2 , CC-83 (ATCC HB 9453), CC-83 F(ab')_2 or B72.3 (ATCC HB 8108).
35. A process for preparing a mono-N-alkylated polyazamacrocyclic compound which comprises reacting an α -halocarboxylic-acid-ester with between one to five equivalents of a 1,4,7,10-tetraazacyclododecane in a solvent which will not promote a proton transfer.

36. The process of claim 35 wherein the α -halocarboxylic-acid-ester is d,l-2-bromo-4(4-nitrophenyl)butanoic acid methylester, 2-bromo-2-(4-nitrophenyl)ethanoic acid-methylester, α -bromo-(2-methoxy-5-nitrophenyl)acetic-acid-methylester, or d,l-2-bromo-4-(4-nitrophenyl)-butanoic acid isopropylester.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Verbindung der Formel:



(IA)

worin:

jedes Q unabhängig Wasserstoff oder $(\text{CHR}^5)_p\text{CO}_2\text{R}$ ist,

Q^1 Wasserstoff oder $(\text{CHR}^5)_w\text{CO}_2\text{R}$ ist,

jedes R unabhängig Wasserstoff, Benzyl oder C_1 - C_4 -Alkyl ist,

mit der Maßgabe, daß mindestens zwei aus der Summe von Q und Q^1 von Wasserstoff verschieden sein müssen und, daß wenn Q^1 Wasserstoff ist R^3 von Wasserstoff verschieden sein muß,

jedes R unabhängig Wasserstoff oder C_1 - C_4 -Alkyl ist,

X und Y unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

n 0 oder 1 ist

m eine ganze Zahl von 0 bis einschließlich 10 ist,

p = 1 oder 2,

r = 0 oder 1,

w = 0 oder 1,

mit der Maßgabe, daß n nur dann 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden und die Summe von r und w 0 oder 1 ist,

R^2 Wasserstoff, Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido oder Carboxyl ist,

R^3 C_1 - C_4 -Alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, Hydroxy oder Wasserstoff ist,

R^4 Wasserstoff, Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl ist,

mit der Maßgabe, daß R^2 und R^4 nicht gleichzeitig Wasserstoff sein können, aber eines von R^2 und R^4 Wasserstoff sein muß oder

ein pharmazeutisch verträgliches Salz davon.

2. Verbindung nach Anspruch 1, worin das pharmazeutisch verträgliche Salz ausgewählt ist aus der Gruppe, bestehend aus Kalium, Natrium, Lithium, Ammonium, Calcium, Magnesium und Chlorwasserstoffaddukten.


3. Verbindung nach Anspruch 1, worin R und R^5 Wasserstoff sind.

4. Verbindung nach Anspruch 1, worin R Wasserstoff ist, R^5 Methyl ist und w 0 ist.

5. Verbindung nach Anspruch 1, worin n und r 0 sind und m 0 bis einschließlich 5 ist.

6. Verbindung nach Anspruch 1, worin R^3 und R^4 Wasserstoff sind.


7. Verbindung nach Anspruch 1 mit der Formel



worin:

20 jedes Q unabhängig Wasserstoff oder $\text{CHR}^5\text{CO}_2\text{R}$ ist,
 jedes R unabhängig Wasserstoff, Benzyl oder $\text{C}_1\text{-C}_4$ Alkyl ist,
 mit der Maßgabe, daß mindestens zwei von Q von Wasserstoff verschieden sein müssen,
 m eine ganze Zahl von 0 bis einschließlich 5 ist,
 R^2 , R^3 , R^4 und R^5 wie in Anspruch 1 definiert sind,
 25 oder ein pharmazeutisch verträgliches Salz davon.

8. Verbindung nach Anspruch 7 mit der Formel



worin:

45 Q und m wie in Anspruch 7 definiert sind,
 R^2 ausgewählt ist aus der Gruppe, bestehend aus Nitro, Amino, Isothiocyanato und Semicarbazido,
 Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl oder
 ein pharmazeutisch verträgliches Salz davon.


50 9. Verbindung nach Anspruch 7, welche 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure-4,7,10-trimethylester ist.

10. Verbindung nach Anspruch 8, welche 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäurehydrochloridsalz ist.

55 11. Verbindung nach Anspruch 8, welche 1-[2-(4-Isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure ist.

12. Verbindung nach Anspruch 8, welche 1-(4-Aminobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäurehydrochloridsalz ist.

13. Verbindung nach Anspruch 7 mit der Formel



worin:

Q und m wie in Anspruch 7 definiert sind,

R³ ausgewählt ist aus der Gruppe, bestehend aus C₁-C₄-Alkoxy, -OCH₂CO₂H und Hydroxy,

R⁴ ausgewählt ist aus der Gruppe, bestehend aus Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl


oder

ein pharmazeutisch verträgliches Salz davon.

14. Verbindung nach Anspruch 13, welche 1-(2-Methoxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure ist.

15. Verbindung nach Anspruch 13, welche 1-(2-Hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure ist.

16. Verbindung nach Anspruch 1 mit der Formel



worin:

jedes Q unabhängig Wasserstoff oder CHR⁵CO₂R ist,

Q¹ Wasserstoff oder (CHR⁵)_wCO₂R ist,

jedes R unabhängig Wasserstoff, Benzyl oder C₁-C₄-Alkyl ist,

mit der Maßgabe, daß mindestens zwei aus der Summe von Q und Q¹ von Wasserstoff verschieden sein müssen und eines von Q Wasserstoff ist,


m eine ganze Zahl von 0 bis einschließlich 5 ist,

w 0 oder 1 ist,

R^2 , R^3 , R^4 und R^5 wie in Anspruch 1 definiert sind,
oder
ein pharmazeutisch verträgliches Salz davon.

17. Verbindung nach Anspruch 16, worin R Wasserstoff und R^5 C_{1-4} -Alkyl ist.

18. Verbindung nach Anspruch 16 mit der Formel




worin:

jedes Q unabhängig Wasserstoff oder CHR^5CO_2R ist,
jedes R unabhängig Wasserstoff, Benzyl oder C_1-C_4 -Alkyl ist,
mit der Maßgabe, daß mindestens ein Q von Wasserstoff verschieden sein muß,
m eine ganze Zahl von 0 bis einschließlich 5 ist,
 R^2 , R^3 , R^4 und R^5 wie in Anspruch 1 definiert sind

oder
ein pharmazeutisch verträgliches Salz davon.

19. Verbindung nach Anspruch 18 mit der Formel



worin:

Q und m wie in Anspruch 18 definiert sind,
 R^2 ausgewählt ist aus der Gruppe, bestehend aus Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl,
oder
ein pharmazeutisch verträgliches Salz davon.

20. Verbindung nach Anspruch 19, welche α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure-1-isopropyl-4,7,10-trimethylester ist.

21. Verbindung nach Anspruch 19, welche α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure ist.

22. Verbindung nach Anspruch 19, welche α -[2-(4-Isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure ist.

23. Verbindung nach Anspruch 19, welche das gemischte Ammonium-Kaliumsalz von α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,10-triessigsäure ist.

24. Verbindung nach Anspruch 19, welche α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7-triessigsäure ist.

25. Verbindung nach Anspruch 19, welche α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,10-triessigsäure ist.

26. Verbindung nach Anspruch 19, welche α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure ist.

27. Verbindung nach Anspruch 19, welche α -(4-Isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure ist.


28. Verbindung nach Anspruch 19, welche α -[2-(4-Nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-1,4,7,10-tetramethylester ist.

29. Verbindung nach Anspruch 19, welche α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-1,4,7,10-tetramethylester ist.

30. Verbindung nach Anspruch 19, welche α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure ist.

31. Verbindung nach Anspruch 19, welche α -[2-(4-Isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure ist.

32. Verbindung nach Anspruch 18 mit der Formel



(VIII)

worin:

jedes Q unabhängig Wasserstoff oder $\text{CHR}^5\text{CO}_2\text{R}$ ist,

jedes R unabhängig Wasserstoff, Benzyl oder C_1 - C_4 -Alkyl ist,

mit der Maßgabe, daß mindestens ein Q von Wasserstoff verschieden ist,

m eine ganze Zahl von 0 bis einschließlich 5 ist,

R^3 und R^5 wie in Anspruch 1 definiert sind,


R^4 ausgewählt ist aus der Gruppe, bestehend aus Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl,

oder
ein pharmazeutisch verträgliches Salz davon.

- 5 33. Verbindung nach Anspruch 32, welche α -(2-Methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure ist.
34. Verbindung nach Anspruch 32, welche α -(2-Methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-tetraammoniumsalz ist.
- 10 35. Verbindung nach Anspruch 32, welche α -(2-Methoxy-5-isothiocyanatophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-tetraammoniumsalz ist.
- 15 36. Komplex, umfassend eine Verbindung gemäß Anspruch 1 oder ein pharmazeutisch verträgliches Salz davon, komplexiert mit einem Ion eines Metalls, ausgewählt von La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y und Sc.
37. Komplex nach Anspruch 36, worin das Metall Sm, Lu oder Y ist.
- 20 38. Komplex nach Anspruch 36, worin das Metall ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc oder ^{142}Pr ist.
39. Komplex nach Anspruch 38, worin das Metall ^{153}Sm , ^{177}Lu oder ^{90}Y ist.
- 25 40. Komplex nach einem der Ansprüche 36 bis 39, worin das Metallion komplexiert ist mit der Verbindung der Ansprüche 7, 8, 13, 16, 18, 19 oder 32.
41. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 26.
42. Komplex nach Anspruch 41, worin das Metall ^{153}Sm oder ^{90}Y ist.
- 30 43. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 27.
44. Komplex nach Anspruch 43, worin das Metall ^{153}Sm ist.
- 35 45. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 30 oder 31.
46. Komplex nach Anspruch 45, worin das Metall ^{153}Sm , ^{90}Y oder ^{177}Lu ist.
- 40 47. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 32.
48. Komplex nach Anspruch 47, worin das Metall ^{153}Sm , ^{90}Y oder ^{177}Lu ist.
49. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 10.
- 45 50. Komplex nach Anspruch 49, worin das Metall ^{153}Sm ist.
51. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 11.
- 50 52. Komplex nach Anspruch 51, worin das Metall ^{153}Sm ist.
53. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 34.
54. Komplex nach Anspruch 53, worin das Metall ^{153}Sm ist.
- 55 55. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 35.
56. Komplex nach Anspruch 55, worin das Metall ^{153}Sm ist.

57. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung nach Anspruch 21.
58. Komplex nach Anspruch 57, worin das Metall ^{153}Sm , ^{90}Y oder ^{177}Lu ist.
- 5 59. Komplex nach Anspruch 38, worin das Metallion komplexiert ist mit der Verbindung von Anspruch 22.
60. Komplex nach Anspruch 59, worin das Metall ^{153}Sm , ^{90}Y oder ^{177}Lu ist.
61. Konjugat, umfassend einen Komplex gemäß Anspruch 36, mit der Maßgabe, daß R^2 und R^4 nicht Nitro sein können und kovalent gebunden an einen Antikörper oder ein Antikörperfragment.
- 10 62. Konjugat nach Anspruch 61, worin das Metall ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc oder ^{142}Pr ist.
- 15 63. Konjugat nach Anspruch 62, worin das Metall ^{153}Sm , ^{90}Y oder ^{177}Lu ist.
64. Konjugat nach einem der Ansprüche 61 bis 63, worin der Antikörper oder das Antikörperfragment ein monoklonaler Antikörper oder ein Fragment davon ist.
- 20 65. Konjugat nach Anspruch 64, worin der Antikörper oder das Antikörperfragment CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ oder B72.3 (ATCC HB 8108) ist.
66. Konjugat nach einem der Ansprüche 61 bis 63, worin die Verbindung wie in den Ansprüchen 7, 8, 13, 16, 18, 19 oder 32 definiert ist.
- 25 67. Konjugat nach einem der Ansprüche 62 bis 65, worin das Metallion ^{153}Sm ist, welches mit der Verbindung von Anspruch 27 komplexiert ist.
68. Konjugat nach einem der Ansprüche 62 bis 65, worin das Metallion ^{153}Sm , ^{90}Y oder ^{177}Lu ist, welches mit der Verbindung von Anspruch 32 komplexiert ist.
- 30 69. Konjugat nach einem der Ansprüche 62 bis 65, worin das Metallion ^{153}Sm ist, welches mit der Verbindung von Anspruch 35 komplexiert ist.
- 35 70. Konjugat nach einem der Ansprüche 62 bis 65, worin das Metallion ^{153}Sm , ^{90}Y oder ^{177}Lu ist, welches mit der Verbindung von Anspruch 22 komplexiert ist.
71. Konjugat, umfassend eine Verbindung nach Anspruch 1 oder ein pharmazeutisch verträgliches Salz davon, kovalent gebunden an einen Antikörper oder ein Antikörperfragment.
- 40 72. Konjugat nach Anspruch 71, worin der Antikörper oder das Antikörperfragment ein monoklonaler Antikörper oder ein Fragment davon ist.
73. Konjugat nach Anspruch 72, worin der Antikörper oder das Antikörperfragment CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ oder B72.3 (ATCC HB 8108) ist.
- 45 74. Konjugat nach einem der Ansprüche 71 bis 73, worin die Verbindung wie in den Ansprüchen 7, 8, 13, 16, 18, 19 oder 32 definiert ist.
- 50 75. Pharmazeutische Formulierung, umfassend ein Konjugat gemäß einem der Ansprüche 61 bis 70, mit einem pharmazeutisch verträglichen Träger.

76. Verfahren zum Herstellen einer Verbindung der Formel



(IA)

worin:

jedes Q unabhängig Wasserstoff oder $(\text{CHR}^5)_p\text{CO}_2\text{R}$ ist,

Q^1 Wasserstoff oder $(\text{CHR}^5)_w\text{CO}_2\text{R}$ ist,

jedes R unabhängig Wasserstoff, Benzyl oder C_1 - C_4 -Alkyl ist,

mit der Maßgabe, daß mindestens zwei aus der Summe von Q und Q^1 von Wasserstoff verschieden sein müssen und, daß wenn Q^1 Wasserstoff ist R^3 von Wasserstoff verschieden sein muß,

jedes R^5 unabhängig Wasserstoff oder C_1 - C_4 Alkyl ist,

X und Y unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

n 0 oder 1 ist

m eine ganze Zahl von 0 bis einschließlich 10 ist,

p = 1 oder 2,

r = 0 oder 1,

w = 0 oder 1,

mit der Maßgabe, daß n nur dann 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden und die Summe von r und w 0 oder 1 ist,

R^2 ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl,

R^3 ausgewählt ist aus der Gruppe, bestehend aus C_1 - C_4 -Alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, Hydroxy und Wasserstoff,


R^4 ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl,

mit der Maßgabe, daß R^2 und R^4 nicht gleichzeitig Wasserstoff sein können, aber eines von R^2 und R^4 Wasserstoff sein muß oder

ein pharmazeutisch verträgliches Salz davon

umfassend einen der folgenden Schritte:

(A) Umsetzen einer Verbindung der Formel



(IIIA)

worin:

X und Y unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,


n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

r = 0 oder 1,

mit der Maßgabe, daß n nur 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

mit einem α -Halocarbonsäureester der Formel



worin R unabhängig Benzyl oder C₁-C₄-Alkyl ist,

in der Gegenwart eines organischen Lösungsmittels und einer Base, bei einer Temperatur von 0 °C bis Rückfluß,

um die Verbindungen der Formel (IA) bereitzustellen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q¹, R³, R⁴, R⁵ Wasserstoff sind,

R C₁-C₄-Alkyl oder Benzyl ist, und

R² Nitro ist oder

(B) Umsetzen des Produkts von Schritt (A) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q¹, R³, R⁴, R⁵ Wasserstoff ist,

R C₁-C₄-Alkyl oder Benzyl ist, und

R² Amino ist oder


(C) Umsetzen einer Verbindung der Formel (IIIA) wie vorstehend definiert, mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, gefolgt von Chlorwasserstoffgas in der Gegenwart eines Lösungsmittels, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q¹, R³, R⁴, Wasserstoff sind, und

R² Amino ist oder

(D) Umsetzen des Produkts von Schritt (C) mit einem α -Halocarbonsäureester der Formel



worin:

jedes R unabhängig Benzyl oder C₁-C₄-Alkyl ist,

in der Gegenwart eines organischen Lösungsmittels und einer Base, bei einer Temperatur von 0 °C bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin


X, Y, m, n, r wie vorstehend definiert sind,

Q¹, R³, R⁴, R⁵ Wasserstoff ist,

R C₁-C₄-Alkyl oder Benzyl ist, und

R² Amino ist oder

(E) Umsetzen einer Verbindung der Formel



(VA)

worin

 $\text{Q}^1 (\text{CHR}^5)_w \text{CO}_2 \text{R}$ ist,jedes R unabhängig Benzyl oder C_1 - C_4 -Alkyl ist,jedes R^5 unabhängig Wasserstoff oder C_1 - C_4 -Alkyl ist,

X und Y jeweils unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

r = 0 oder 1,

w = 0 oder 1,

mit der Maßgabe, daß die Summe von r und w 0 oder 1 ist,

mit einem α -Halocarbonsäureester der Formel $\text{BrCH}_2 \text{CO}_2 \text{R}$ worin jedes R unabhängig Benzyl oder C_1 - C_4 -Alkyl ist,in der Gegenwart eines organischen Lösungsmittels und einer Base, bei einer Temperatur von 0°C bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q^1 , m, n, r wie vorstehend definiert sind, R^3 , R^4 , R^5 Wasserstoff sind,R C_1 - C_4 -Alkyl oder Benzyl ist, und R^2 Nitro ist oder

(F) Umsetzen des Produkts von Schritt (E) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q^1 , m, n, r wie vorstehend definiert sind, R^3 , R^4 , R^5 Wasserstoff sind,R C_1 - C_4 Alkyl oder Benzyl ist, und R^2 Amino ist oder

(G) Umsetzen des Produkts entweder aus Schritt (B) oder (D) mit einem Entesterungsmittel, wie etwa einer starken Säure, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

 Q^1 , R^3 , R^4 , R^5 Wasserstoff sind,

R Wasserstoff ist und

 R^2 Amino ist oder


(H) Umsetzen des Produkts von Schritt (F) mit einem Entesterungsmittel, wie etwa einer starken Säure, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q^1 , m, n, r wie vorstehend definiert sind, R^3 , R^4 , R^5 Wasserstoff sind,

R Wasserstoff ist, und

R² Amino ist oder

(I) Umsetzen einer Verbindung der Formel (VA), wie vorstehend definiert, mit einem optisch aktiven α -Benzolsulfonat der Formel



worin

R und R⁵ wie vorstehend definiert sind,
um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, R⁵, m, n, r wie vorstehend definiert sind,

R³ und R⁴ Wasserstoff sind,

R C₁₋₄-Alkyl oder Benzyl ist, und

R² Nitro ist oder

(J) Umsetzen des Produkts von Schritt (I) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, R⁵, m, n, r wie vorstehend definiert sind,

R³ und R⁴ Wasserstoff sind,

R C₁₋₄-Alkyl oder Benzyl ist, und

R² Amino ist oder


(K) Umsetzen des Produkts von Schritt (J) mit einem Entesterungsmittel, wie etwa einer starken Säure, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, R⁵, m, n, r wie vorstehend definiert sind,

R³, R⁴ und R Wasserstoff sind und

R² Amino ist oder

(L) Umsetzen einer Verbindung der Formel



(IVA)

worin:

X und Y jeweils unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoff-Bindung bilden,


R³ C₁₋₄-Alkoxy, -OCH₂CO₂H oder Hydroxy ist,

n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

$r = 0$ oder 1

mit der Maßgabe, daß n nur dann 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,
mit einer α -Halocarbonsäure der Formel



in der Gegenwart eines organischen Lösungsmittels und einer Base bei einer Temperatur von 0°C bis Rückfluß,


um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, R^3, m, n, r wie vorstehend definiert sind,
 Q^1 und R^2 Wasserstoff sind und
 R^4 Nitro ist oder

(M) Umsetzen des Produkts von Schritt (L) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) bereitzustellen, worin:

X, Y, R^3, m, n, r wie vorstehend definiert sind,
 Q^1, R, R^2 und R^5 Wasserstoff sind und
 R^2 Amino ist oder

(N) Umsetzen einer Verbindung der Formel



(VIII A)

worin:

X und Y jeweils unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

$Q^1 \text{ CH}_2\text{CO}_2\text{R}$ ist,

R Benzyl oder C_{1-4} -Alkyl ist,


n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

$r = 0$ oder 1 ,

mit der Maßgabe, daß n nur 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoff-Bindung bilden,

mit einer α -Halocarbonsäure der Formel



worin R wie vorstehend definiert ist,

In der Gegenwart eines organischen Lösungsmittels und einer Base bei einer Temperatur von 0°C bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q^1, Q, m, n, r wie vorstehend definiert sind,

R^2 Wasserstoff ist, und

R^4 Nitro ist oder

(O) Umsetzen des Produkts von Schritt (N) mit einem Entesterungsmittel, wie etwa einer starken Säure, gefolgt von einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff unter Standardreaktions-

bedingungen, um die Verbindungen von Formel (IA) vorzusehen, worin:

X, Y, Q¹, Q, m, n, r wie vorstehend definiert sind,

R, R² und R⁵ Wasserstoff sind, und

R² Amino ist oder


(P) Umsetzen des Produkts von Schritt (G), (F), (K), (M) oder (O) mit Thiophosgen, um die entsprechende Verbindung bereitzustellen, worin R² oder R⁴ Isothiocyanato ist.

77. Verfahren zum Herstellen eines Mono-N-alkylierten Polyazamacrozyklus, umfassend das Umsetzen eines α -Halocarbonsäureesters mit zwischen eins bis fünf Äquivalenten eines 1,4,7,10-Tetraazacyclodecans in einem Lösungsmittel, welches einen Protonentransfer nicht fördert.

78. Verfahren nach Anspruch 77, worin der α -Halocarbonsäureester d,1-2-Brom-4(4-nitrophenyl)butansäuremethylester, 2-Brom-2-(4-nitrophenyl)ethansäuremethylester, α -Brom-(2-methoxy-5-nitrophenyl)essigsäuremethylester oder d,1-2-Brom-4-(4-nitrophenyl)butansäureisopropylester ist.

Patentansprüche für folgende Vertragsstaaten : ES, GR

1. Verfahren zum Herstellen einer Verbindung der Formel



worin:

jedes Q unabhängig Wasserstoff oder (CHR⁵)_pCO₂R ist,

Q¹ Wasserstoff oder (CHR⁵)_wCO₂R ist,

jedes R unabhängig Wasserstoff, Benzyl oder C₁-C₄-Alkyl ist,

mit der Maßgabe, daß mindestens zwei aus der Summe von Q und Q¹ von Wasserstoff verschieden sein müssen und, daß wenn Q¹ Wasserstoff ist R³ von Wasserstoff verschieden sein muß,

jedes R⁵ unabhängig Wasserstoff oder C₁-C₄ Alkyl ist,

X und Y unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

n 0 oder 1 ist

m eine ganze Zahl von 0 bis einschließlich 10 ist,

p = 1 oder 2,

r = 0 oder 1,

w = 0 oder 1,

mit der Maßgabe, daß n nur dann 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden und die Summe von r und w 0 oder 1 ist,

R² ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl,


R³ ausgewählt ist aus der Gruppe, bestehend aus C₁-C₄-Alkoxy, -OCH₂CO₂H, Hydroxy und Wasserstoff,

R⁴ ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Nitro, Amino, Isothiocyanato, Semicarbazido, Thiosemicarbazido, Maleimido, Bromacetamido und Carboxyl,

mit der Maßgabe, daß R² und R⁴ nicht gleichzeitig Wasserstoff sein können, aber eines von R² und R⁴ Wasserstoff sein muß oder

ein pharmazeutisch verträgliches Salz davon umfassend einen der folgenden Schritte:

(A) Umsetzen einer Verbindung der Formel



(IIIA)

worin:

X und Y unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,


n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

r = 0 oder 1,

mit der Maßgabe, daß n nur 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

mit einem α -Halocarbonsäureester der Formel



worin R unabhängig Benzyl oder C_1 - C_4 -Alkyl ist,

in der Gegenwart eines organischen Lösungsmittels und einer Base, bei einer Temperatur von 0°C bis Rückfluß,

um die Verbindungen der Formel (IA) bereitzustellen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q^1 , R^3 , R^4 , R^5 Wasserstoff sind,

R C_1 - C_4 -Alkyl oder Benzyl ist, und

R^2 Nitro ist oder

(B) Umsetzen des Produkts von Schritt (A) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q^1 , R^3 , R^4 , R^5 Wasserstoff ist,

R C_1 - C_4 -Alkyl oder Benzyl ist, und

R^2 Amino ist oder


(C) Umsetzen einer Verbindung der Formel (IIIA) wie vorstehend definiert, mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, gefolgt von Chlorwasserstoffgas in der Gegenwart eines Lösungsmittels, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q^1 , R^3 , R^4 , Wasserstoff sind, und

R^2 Amino ist oder

(D) Umsetzen des Produkts von Schritt (C) mit einem α -Halocarbonsäureester der Formel



worin:

jedes R unabhängig Benzyl oder C_1 - C_4 -Alkyl ist,

in der Gegenwart eines organischen Lösungsmittels und einer Base, bei einer Temperatur von 0°C bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin


X, Y, m, n, r wie vorstehend definiert sind,

Q¹, R³, R⁴, R⁵ Wasserstoff ist,

R C₁₋₄-Alkyl oder Benzyl ist, und

R² Amino ist oder

(E) Umsetzen einer Verbindung der Formel



(VA)

worin

Q¹ (CHR⁵)_wCO₂R ist,

jedes R unabhängig Benzyl oder C₁-C₄-Alkyl ist,

jedes R⁵ unabhängig Wasserstoff oder C₁-C₄-Alkyl ist,

X und Y jeweils unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

n 0 oder 1 ist,


m eine ganze Zahl von 0 bis einschließlich 10 ist,

r = 0 oder 1,

w = 0 oder 1,

mit der Maßgabe, daß die Summe von r und w 0 oder 1 ist,

mit einem α-Halocarbonsäureester der Formel



worin jedes R unabhängig Benzyl oder C₁-C₄-Alkyl ist,

in der Gegenwart eines organischen Lösungsmittels und einer Base, bei einer Temperatur von 0 °C bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, m, n, r wie vorstehend definiert sind,

R³, R⁴, R⁵ Wasserstoff sind,

R C₁₋₄-Alkyl oder Benzyl ist, und

R² Nitro ist oder

(F) Umsetzen des Produkts von Schritt (E) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, m, n, r wie vorstehend definiert sind,

R³, R⁴, R⁵ Wasserstoff sind,

R C₁₋₄ Alkyl oder Benzyl ist, und

R² Amino ist oder

(G) Umsetzen des Produkts entweder aus Schritt (B) oder (D) mit einem Entesterungsmittel, wie etwa einer starken Säure, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, m, n, r wie vorstehend definiert sind,

Q¹, R³, R⁴, R⁵ Wasserstoff sind,

R Wasserstoff ist und

R² Amino ist oder

(H) Umsetzen des Produkts von Schritt (F) mit einem Entesterungsmittel, wie etwa einer starken Säure, um die Verbindungen der Formel (IA) vorzusehen, worin:


X, Y, Q¹, m, n, r wie vorstehend definiert sind,

R³, R⁴, R⁵ Wasserstoff sind,

R Wasserstoff ist, und

R² Amino ist oder

(I) Umsetzen einer Verbindung der Formel (VA), wie vorstehend definiert, mit einem optisch aktiven α -Benzolsulfonat der Formel



worin

R und R⁵ wie vorstehend definiert sind,

um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, R⁵, m, n, r wie vorstehend definiert sind,

R³ und R⁴ Wasserstoff sind,

R C₁₋₄-Alkyl oder Benzyl ist, und

R² Nitro ist oder

(J) Umsetzen des Produkts von Schritt (I) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, R⁵, m, n, r wie vorstehend definiert sind,

R³ und R⁴ Wasserstoff sind,

R C₁₋₄-Alkyl oder Benzyl ist, und

R² Amino ist oder


(K) Umsetzen des Produkts von Schritt (J) mit einem Entesterungsmittel, wie etwa einer starken Säure, um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, R⁵, m, n, r wie vorstehend definiert sind,

R³, R⁴ und R Wasserstoff sind und

R² Amino ist oder

(L) Umsetzen einer Verbindung der Formel



(IVA)

worin:

X und Y jeweils unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoff-Bindung bilden,

$R^{3'}$ C_1 - C_4 -Alkoxy, $-OCH_2CO_2H$ oder Hydroxy ist,


n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

r = 0 oder 1

mit der Maßgabe, daß n nur dann 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

mit einer α -Halocarbonsäure der Formel



in der Gegenwart eines organischen Lösungsmittels und einer Base bei einer Temperatur von $0^\circ C$ bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, $R^{3'}$, m, n, r wie vorstehend definiert sind,

Q^1 und R^2 Wasserstoff sind und

R^4 Nitro ist oder


(M) Umsetzen des Produkts von Schritt (L) mit einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff, unter Standardreaktionsbedingungen, um die Verbindungen der Formel (IA) bereitzustellen, worin:

X, Y, $R^{3'}$, m, n, r wie vorstehend definiert sind,

Q^1 , R, R^2 und R^5 Wasserstoff sind und

R^2 Amino ist oder

(N) Umsetzen einer Verbindung der Formel



(VIII A)

worin:

X und Y jeweils unabhängig Wasserstoff sind oder zusammen mit einem benachbarten X und Y eine zusätzliche Kohlenstoff-Kohlenstoffbindung bilden,

Q^1 CH_2CO_2R ist,

R Benzyl oder C_1 - C_4 -Alkyl ist,


n 0 oder 1 ist,

m eine ganze Zahl von 0 bis einschließlich 10 ist,

r = 0 oder 1,

mit der Maßgabe, daß n nur 1 ist, wenn X und/oder Y eine zusätzliche Kohlenstoff-Kohlenstoff-Bindung bilden,

mit einer α -Halocarbonsäure der Formel



worin R wie vorstehend definiert ist,

in der Gegenwart eines organischen Lösungsmittels und einer Base bei einer Temperatur von $0^\circ C$ bis Rückfluß,

um die Verbindungen der Formel (IA) vorzusehen, worin:

X, Y, Q¹, Q, m, n, r wie vorstehend definiert sind,

R² Wasserstoff ist, und

R⁴ Nitro ist oder

(O) Umsetzen des Produkts von Schritt (N) mit einem Entesterungsmittel, wie etwa einer starken Säure, gefolgt von einem Hydrierungsmittel, wie etwa Pd/C mit Wasserstoff unter Standardreaktionsbedingungen, um die Verbindungen von Formel (IA) vorzusehen, worin:

X, Y, Q¹, Q, m, n, r wie vorstehend definiert sind,

R, R² und R⁵ Wasserstoff sind, und

R² Amino ist oder

(P) Umsetzen des Produkts von Schritt (G), (F), (K), (M) oder (O) mit Thiophosgen, um die entsprechende Verbindung bereitzustellen, worin R² oder R⁴ Isothiocyanato ist.

2. Verfahren nach Anspruch 1, zum Herstellen eines pharmazeutisch verträglichen Salzes der Verbindung von Formel (IA), umfassend das Umsetzen der Verbindung von Formel (IA) nach bekannten Verfahren mit einem Reagenz, um Kalium-, Natrium-, Lithium-, Ammonium-, Calcium-, Magnesium- oder Chlorwasserstoffaddukte als das pharmazeutisch verträgliche Salz zu erhalten.

3. Verfahren nach Anspruch 1(B), zum Herstellen von 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure-4,7,10-trimethylester, umfassend das Umsetzen von 1-[2-(4-Nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure-4,7,10-trimethylester mit Wasserstoff und Palladium auf Kohle.

4. Verfahren nach Anspruch 1(G), zum Herstellen von 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure-Hydrochloridsalz, umfassend das Umsetzen von 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure-4,7,10-trimethylester mit Chlorwasserstoffsäure.

5. Verfahren nach Anspruch 1(P), zum Herstellen von 1-[2-(4-Isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure, umfassend das Umsetzen von 1-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure-Hydrochloridsalz mit Thiophosgen.

6. Verfahren nach Anspruch 1(A), zum Herstellen von 1-(4-Aminobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäurehydrochloridsalz, umfassend das Umsetzen von 1-(4-Aminobenzyl)-1,4,7,10-tetraazacyclododecan-tetrahydrochloridsalz mit Methylbromacetat.

7. Verfahren nach Anspruch 1(M), zum Herstellen von 1-(2-Methoxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure, umfassend das Umsetzen von 1-(2-Methoxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure mit Wasserstoff in einer Parr-Bombe.

8. Verfahren nach Anspruch 1(M), zum Herstellen von 1-(2-Hydroxy-5-aminobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure, umfassend das Umsetzen von N-(2-Hydroxy-5-nitrobenzyl)-1,4,7,10-tetraazacyclododecan-4,7,10-triessigsäure mit Wasserstoff und PtO₂.

9. Verfahren nach Anspruch 1(J), zum Herstellen von α-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure-1-isopropyl-4,7,10-trimethylester, umfassend das Umsetzen von α-[2-(4-Nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure-1-isopropyl-4,7,10-trimethylester mit Wasserstoff und Palladium auf Kohle.

10. Verfahren nach Anspruch 1(K), zum Herstellen von α-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure, umfassend das Umsetzen von α-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure-1-isopropyl-4,7,10-trimethylester mit Chlorwasserstoffsäure.

11. Verfahren nach Anspruch 1(P), zum Herstellen von α-[2-(4-Isothiocyanatophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure, umfassend das Umsetzen von α-[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-(R,S)-essigsäure-4,7,10-tris-(R-methyl)essigsäure mit Thiophosgen.

12. Verfahren nach Anspruch 1(F), zum Herstellen von α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7-triessigsäure, umfassend das Umsetzen von α -(4-Nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7-triessigsäure mit Wasserstoff und PtO_2 .
- 5 13. Verfahren nach Anspruch 1(F), zum Herstellen von α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,10-triessigsäure, umfassend das Umsetzen von α -(4-Nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,10-triessigsäure mit Wasserstoff und PtO_2 .
- 10 14. Verfahren nach Anspruch 1(F), zum Herstellen von α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure, umfassend das Umsetzen von α -(4-Nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure mit Wasserstoff und PtO_2 .
- 15 15. Verfahren nach Anspruch 1(P), zum Herstellen von α -(4-Isothiocyantophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure, umfassend das Umsetzen von α -(4-Aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure mit Thiophosgen.
- 20 16. Verfahren nach Anspruch 1(E), zum Herstellen von α -[2-(4-Nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-1,4,7,10-tetramethylester, umfassend das Umsetzen von α -[2-(4-Nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1-essigsäure-1-methylester mit Methylbromacetat.
- 25 17. Verfahren nach Anspruch 1(F), zum Herstellen von α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-1,4,7,10-tetramethylester, umfassend das Umsetzen von α -[2-(4-Nitrophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-1,4,7,10-tetramethylester mit Wasserstoff und Palladium auf Kohle.
- 30 18. Verfahren nach Anspruch 1(H), zum Herstellen von α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure, umfassend das Umsetzen von α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-1,4,7,10-tetramethylester mit Chlorwasserstoffsäure.
- 35 19. Verfahren nach Anspruch 1(P), zum Herstellen von α -[2-(4-Isothiocyantophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure, umfassend das Umsetzen von α -[2-(4-Aminophenyl)ethyl]-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure mit Thiophosgen.
- 40 20. Verfahren nach Anspruch 1(O), zum Herstellen von α -(2-Methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure, umfassend das Umsetzen von α -(2-Methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-tetramethylester mit Chlorwasserstoffsäure.
- 45 21. Verfahren nach Anspruch 1(O), zum Herstellen von α -(2-Methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-tetraammoniumsalz, umfassend das Umsetzen von α -(2-Methoxy-5-nitrophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure mit Wasserstoff und PtO_2 .
- 50 22. Verfahren nach Anspruch 1(P), zum Herstellen von α -(2-Methoxy-5-isothiocyantophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure-tetraammoniumsalz, umfassend das Umsetzen von α -(2-Methoxy-5-aminophenyl)-1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraessigsäure mit Thiophosgen.
- 55 23. Verfahren zum Herstellen eines Komplexes einer Verbindung nach einem der Ansprüche 1 bis 22 mit einem Ion eines Metalls, ausgewählt aus La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y und Sc, umfassend das Umsetzen der Verbindung mit dem Metallion.
24. Verfahren nach Anspruch 23, worin das Metall ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc oder ^{142}Pr ist.
25. Verfahren nach Anspruch 24, worin das Metall ^{153}Sm , ^{177}Lu oder ^{90}Y ist.
26. Verfahren nach einem der Ansprüche 23 bis 25, worin die Reaktion erhitzt wird.
27. Verfahren zum Herstellen eines Konjugats, enthaltend einen Komplex von einer Verbindung nach einem der Ansprüche 1 bis 22, welche mit einem Ion eines Metalls komplexiert ist, ausgewählt aus La, Ce, Pr,


Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y und Sc, umfassend das kovalente Binden des Komplexes an einen Antikörper oder ein Antikörperfragment.

28. Verfahren nach Anspruch 27, worin das Metall ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc oder ^{142}Pr ist.
29. Verfahren nach Anspruch 28, worin das Metall ^{153}Sm , ^{177}Lu oder ^{90}Y ist.
30. Konjugat nach einem der Ansprüche 27 bis 29, worin der Antikörper oder das Antikörperfragment ein monoklonaler Antikörper oder ein Fragment davon ist.
31. Konjugat nach Anspruch 30, worin der Antikörper oder das Antikörperfragment CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ oder B72.3 (ATCC HB 8108) ist.
32. Verfahren zum Herstellen eines Konjugats, welches eine Verbindung nach einem der Ansprüche 1 bis 22 enthält, die kovalent an einen Antikörper oder ein Antikörperfragment gebunden ist, umfassend das Umsetzen der Verbindung mit einem Antikörper oder einem Antikörperfragment.
33. Konjugat nach Anspruch 32, worin der Antikörper oder das Antikörperfragment ein monoklonaler Antikörper oder ein Fragment davon ist.
34. Konjugat nach Anspruch 33, worin der Antikörper oder das Antikörperfragment CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ oder B72.3 (ATCC HB 8108) ist.
35. Verfahren zum Herstellen eines mono-N-alkylierten Polyazamakrocyclus, umfassend das Umsetzen eines α -Halocarbonsäureesters mit zwischen eins bis fünf Äquivalenten eines 1,4,7,10-Tetraazacyclodecans in einem Lösungsmittel, welches einen Protonentransfer nicht fördert.
36. Verfahren nach Anspruch 35, worin der α -Halocarbonsäureester d,1-2-Brom-4-(4-nitrophenyl)-butansäuremethylester, 2-Brom-2-(4-nitrophenyl)ethansäuremethylester, α -Brom-(2-methoxy-5-nitrophenyl)essigsäuremethylester oder d,1-2-Brom-4-(4-nitrophenyl)butansäureisopropylester ist.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Composé de formule :



dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe $(\text{CHR}^5)_p\text{CO}_2\text{R}$;

Q' représente un atome d'hydrogène ou un groupe $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C₁₋₄;

sous réserve qu'au moins deux de tous les Q et Q' représentent autre chose qu'un atome d'hydrogène,

et que, lorsque Q' représente un atome d'hydrogène, R³ représente autre chose qu'un atome d'hydrogène;

chaque R⁵ représente indépendamment un atome d'hydrogène ou un groupe alkyle en C₁₋₄;

X et Y représentent chacun indépendamment un atome d'hydrogène, ou peuvent constituer, conjointement avec un X ou Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

p vaut 1 ou 2;

r vaut 0 ou 1;

w vaut 0 ou 1;

sous réserve que n vaille seulement 1 lorsque X et/ou Y forment une liaison carbone-carbone supplémentaire, et que la somme de r et w vaille 0 ou 1;

R² représente un atome d'hydrogène ou un groupe nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido ou carboxy;

R³ représente un groupe alcoxy en C₁₋₄, hydroxy ou carboxyméthoxy, ou un atome d'hydrogène;

R⁴ représente un atome d'hydrogène ou un groupe nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido ou carboxy;

sous réserve que R² et R⁴ ne représentent pas tous les deux des atomes d'hydrogène, mais que l'un de R² et R⁴ représente un atome d'hydrogène; ou

un sel d'un tel composé, acceptable en pharmacie.

2. Composé conforme à la revendication 1, dans lequel le sel acceptable en pharmacie est choisi dans l'ensemble constitué par les sels de potassium, sodium, lithium, ammonium, calcium et magnésium, et les chlorhydrates.


3. Composé conforme à la revendication 1, dans lequel R et R⁵ représentent tous les deux des atomes d'hydrogène.

4. Composé conforme à la revendication 1, dans lequel R représente un atome d'hydrogène, R⁵ représente un groupe méthyle et w vaut 0.

5. Composé conforme à la revendication 1, dans lequel n et r valent tous les deux 0, et m vaut de 0 à 5.

6. Composé conforme à la revendication 1, dans lequel R³ et R⁴ représentent des atomes d'hydrogène.

7. Composé conforme à la revendication 1, de formule :



dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe CHR⁵CO₂R;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C₁₋₄;


sous réserve qu'au moins deux des Q représentent autre chose qu'un atome d'hydrogène;

m représente un nombre entier valant de 0 à 5, limites comprises;

R², R³, R⁴ et R⁵ ont les définitions indiquées dans la revendication 1;

ou un sel d'un tel composé, acceptable en pharmacie.

8. Composé conforme à la revendication 7, de formule :



dans laquelle :

Q et m ont les définitions indiquées dans la revendication 7;

R² est choisi dans l'ensemble constitué par les groupes nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido et carboxy;
ou un sel d'un tel composé, acceptable en pharmacie.


9. Composé conforme à la revendication 7, qui est l'ester 4,7,10-triméthyllique de l'acide 1-[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique.

10. Composé conforme à la revendication 8, qui est le chlorhydrate de l'acide 1-[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique.

11. Composé conforme à la revendication 8, qui est l'acide 1-[2-(4-isothiocyanatophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique.

12. Composé conforme à la revendication 8, qui est le chlorhydrate de l'acide 1-(4-aminobenzyl)-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique.

13. Composé conforme à la revendication 7, de formule :



dans laquelle :

Q et m ont les définitions indiquées dans la revendication 7;


R³ est choisi parmi les groupes alcoxy en C₁-₄, carboxyméthoxy et hydroxy;

R⁴ est choisi parmi les groupes nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido et carboxy;
ou un sel d'un tel composé, acceptable en pharmacie.

14. Composé conforme à la revendication 13, qui est l'acide 1-(2-méthoxy-5-aminobenzyl)-1,4,7,10-tétraa-
zacyclododécane-4,7,10-triacétique.

15. Composé conforme à la revendication 13, qui est l'acide 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tétraa-
cyclo-dodécane-4,7,10-triacétique.

16. Composé conforme à la revendication 1, de formule :



dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe $\text{CHR}^5\text{CO}_2\text{R}$;

Q^1 représente un atome d'hydrogène ou un groupe $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C_{1-4} ;

sous réserve qu'au moins deux de tous les Q et Q^1 représentent autre chose que des atomes d'hydrogène, et que l'un des Q représente un atome d'hydrogène;

m représente un nombre entier valant de 0 à 5, limites comprises;


w vaut 0 ou 1;

R^2 , R^3 , R^4 et R^5 ont les définitions indiquées dans la revendication 1;

ou un sel d'un tel composé, acceptable en pharmacie.

17. Composé conforme à la revendication 16, dans lequel R représente un atome d'hydrogène et R^5 représente un groupe alkyle en C_{1-4} .

18. Composé conforme à la revendication 16, de formule :




dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe $\text{CHR}^5\text{CO}_2\text{R}$;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C_{1-4} ;

sous réserve qu'au moins l'un des Q représente autre chose qu'un atome d'hydrogène;
 m représente un nombre entier valant de 0 à 5, limites comprises;
 R², R³, R⁴ et R⁵ ont les définitions indiquées dans la revendication 1;
 ou un sel d'un tel composé, acceptable en pharmacie.

19. Composé conforme à la revendication 18, de formule :



dans laquelle :

Q et m ont les définitions indiquées dans la revendication 18;

R² est choisi parmi les groupes nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido et carboxy;
 ou un sel d'un tel composé, acceptable en pharmacie.

20. Composé conforme à la revendication 19, qui est l'ester 1-isopropylique-4,7,10-triméthylque de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthyl-acétique).

21. Composé conforme à la revendication 19, qui est l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthyl-acétique).

22. Composé conforme à la revendication 19, qui est l'acide α -[2-(4-isothiocyanatophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthyl-acétique).

23. Composé conforme à la revendication 19, qui est un sel mixte d'ammonium et de potassium de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,10-triacétique.

24. Composé conforme à la revendication 19, qui est l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7-triacétique.

25. Composé conforme à la revendication 19, qui est l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,10-triacétique.

26. Composé conforme à la revendication 19, qui est l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.

27. Composé conforme à la revendication 19, qui est l'acide α -(4-isothiocyanatophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.


28. Composé conforme à la revendication 19, qui est l'ester 1,4,7,10-tétraméthylque de l'acide α -[2-(4-nitrophényl)éthyl]-1,4,7, 10-tétraazacyclododécane-1,4,7,10-tétraacétique.

29. Composé conforme à la revendication 19, qui est l'ester 1,4,7,10-tétraméthylque de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7, 10-tétraazacyclododécane-1,4,7,10-tétraacétique.

30. Composé conforme à la revendication 19, qui est l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.

31. Composé conforme à la revendication 19, qui est l'acide α -[2-(4-isothiocyanatophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.

32. Composé conforme à la revendication 18, de formule :



dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe $\text{CHR}^5\text{CO}_2\text{R}$;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C_{1-4} ;

sous réserve qu'au moins l'un des Q représente autre chose qu'un atome d'hydrogène;

m représente un nombre entier valant de 0 à 5, limites comprises;

R^3 et R^5 ont les définitions indiquées dans la revendication 1;

R^4 est choisi parmi les groupes nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido et carboxy;

ou un sel d'un tel composé, acceptable en pharmacie.

33. Composé conforme à la revendication 32, qui est l'acide α -(2-méthoxy-5-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.

34. Composé conforme à la revendication 32, qui est le sel tétraammonique de l'acide α -(2-éthoxy-5-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.

35. Composé conforme à la revendication 32, qui est le sel tétraammonique de l'acide α -(2-méthoxy-5-isothiocyanatophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique.

36. Complexe comprenant un composé conforme à la revendication 1 ou un sel d'un tel composé, acceptable en pharmacie, complexé avec un ion d'un métal choisi parmi La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y et Sc.

37. Complexe conforme à la revendication 36, dans lequel le métal est Sm, Lu ou Y.

38. Complexe conforme à la revendication 36, dans lequel le métal est ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc ou ^{142}Pr .

39. Complexe conforme à la revendication 38, dans lequel le métal est ^{153}Sm , ^{177}Lu ou ^{90}Y .

40. Complexe conforme à l'une des revendications 36 à 39, dans lequel l'ion métallique est complexé par un composé conforme à l'une des revendications 7, 8, 13, 16, 18, 19 et 32.

41. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 26.
42. Complexe conforme à la revendication 41, dans lequel le métal est ^{153}Sm ou ^{90}Y .
- 5 43. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 27.
44. Complexe conforme à la revendication 43, dans lequel le métal est ^{153}Sm .
- 10 45. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 30 ou 31.
46. Complexe conforme à la revendication 45, dans lequel le métal est ^{153}Sm , ^{90}Y ou ^{177}Lu .
- 15 47. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 32.
48. Complexe conforme à la revendication 47, dans lequel le métal est ^{153}Sm , ^{90}Y ou ^{177}Lu .
- 20 49. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 10.
50. Complexe conforme à la revendication 49, dans lequel le métal est ^{153}Sm .
- 25 51. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 11.
52. Complexe conforme à la revendication 51, dans lequel le métal est ^{153}Sm .
- 30 53. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 34.
54. Complexe conforme à la revendication 53, dans lequel le métal est ^{153}Sm .
- 35 55. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 35.
56. Complexe conforme à la revendication 55, dans lequel le métal est ^{153}Sm .
- 40 57. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 21.
58. Complexe conforme à la revendication 57, dans lequel le métal est ^{153}Sm , ^{90}Y ou ^{177}Lu .
- 45 59. Complexe conforme à la revendication 38, dans lequel l'ion métallique est complexé par un composé conforme à la revendication 22.
60. Complexe conforme à la revendication 59, dans lequel le métal est ^{153}Sm , ^{90}Y ou ^{177}Lu .
- 50 61. Conjugué comprenant un complexe conforme à la revendication 36, sous réserve que R^2 et R^4 ne représentent pas des groupes nitro, et lié par liaison covalente à un anticorps ou à un fragment d'anticorps.
- 55 62. Conjugué conforme à la revendication 61, dans lequel le métal est ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc ou ^{142}Pr .
63. Conjugué conforme à la revendication 62, dans lequel le métal est ^{153}Sm , ^{90}Y ou ^{177}Lu .

64. Conjugué conforme à l'une des revendications 61 à 63, dans lequel l'anticorps ou le fragment d'anticorps est un anticorps monoclonal ou un fragment d'un tel anticorps.

65. Conjugué conforme à la revendication 64, dans lequel l'anticorps ou le fragment d'anticorps est CC-49 (ATCC HB 9459), CC-49-F(ab')₂, CC-83 (ATCC HB 9453), CC-83-F(ab')₂ ou B72.3 (ATCC HB 8108).

66. Conjugué conforme à l'une des revendications 61 à 63, dans lequel le composé est un composé défini dans l'une des revendications 7, 8, 13, 16, 18, 19 et 32.

67. Conjugué conforme à l'une des revendications 62 à 65, dans lequel l'ion métallique est ¹⁵³Sm, complexé avec le composé conforme à la revendication 27.

68. Conjugué conforme à l'une des revendications 62 à 65, dans lequel l'ion métallique est ¹⁵³Sm, ⁹⁰Y ou ¹⁷⁷Lu, complexé avec un composé conforme à la revendication 32.

69. Conjugué conforme à l'une des revendications 62 à 65, dans lequel l'ion métallique est ¹⁵³Sm, complexé avec le composé conforme à la revendication 35.

70. Conjugué conforme à l'une des revendications 62 à 65, dans lequel l'ion métallique est ¹⁵³Sm, ⁹⁰Y ou ¹⁷⁷Lu, complexé avec le composé conforme à la revendication 22.

71. Conjugué comprenant un composé conforme à la revendication 1 ou un sel d'un tel composé, acceptable en pharmacie, lié par liaison covalente à un anticorps ou un fragment d'anticorps.


72. Conjugué conforme à la revendication 71, dans lequel l'anticorps ou le fragment d'anticorps est un anticorps monoclonal ou un fragment d'un tel anticorps.

73. Conjugué conforme à la revendication 72, dans lequel l'anticorps ou le fragment d'anticorps est CC-49 (ATCC HB 9459), CC-49-F(ab')₂, CC-83 (ATCC HB 9453), CC-83-F(ab')₂ ou B72.3 (ATCC HB 8108).

74. Conjugué conforme à la revendication 71, dans lequel le composé est un composé défini dans l'une des revendications 7, 8, 13, 16, 18, 19 et 32.

75. Formulation pharmaceutique comprenant un conjugué conforme à l'une des revendications 61 à 70, et un véhicule acceptable en pharmacie.

76. Procédé de préparation d'un composé de formule :



(IA)

dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe (CHR⁵)_pCO₂R;

Q¹ représente un atome d'hydrogène ou un groupe (CHR⁵)_wCO₂R;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C₁₋₄;

sous réserve qu'au moins deux de tous les Q et Q¹ représentent autre chose qu'un atome d'hydrogène,

et que, lorsque Q¹ représente un atome d'hydrogène, R³ représente autre chose qu'un atome d'hydrogène;

chaque R⁵ représente indépendamment un atome d'hydrogène ou un groupe alkyle en C₁₋₄;

X et Y représentent chacun indépendamment un atome d'hydrogène, ou peuvent constituer, conjointement avec un X ou Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

p vaut 1 ou 2;

r vaut 0 ou 1;

w vaut 0 ou 1;

sous réserve que n vaille seulement 1 lorsque X et/ou Y forment une liaison carbone-carbone supplémentaire, et que la somme de r et w vaille 0 ou 1;

R² représente un atome d'hydrogène ou un groupe nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido ou carboxy;

R³ représente un groupe alcoxy en C₁₋₄, hydroxy ou carboxyméthoxy, ou un atome d'hydrogène;


R⁴ représente un atome d'hydrogène ou un groupe nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido ou carboxy;

sous réserve que R² et R⁴ ne représentent pas tous les deux des atomes d'hydrogène, mais que l'un de R² et R⁴ représente un atome d'hydrogène; ou

d'un sel d'un tel composé, acceptable en pharmacie;

lequel procédé comprend l'une des étapes suivantes :

(A) réaction d'un composé de formule :



(IIIA)

dans laquelle :


X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

sous réserve que n vaille 1 quand X et/ou Y forment une liaison supplémentaire carbone-carbone; et d'un ester d'acide α -halogénocarboxylique, de formule :



dans laquelle R représente indépendamment un groupe benzyle ou un groupe alkyle en C₁₋₄;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe nitro; ou

(B) réaction du produit de l'étape (A) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de

formule (IA), dans laquelle :

X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou

(C) réaction d'un composé de formule (IIIA) définie plus haut avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, puis avec du chlorure d'hydrogène gazeux, en présence d'un solvant, pour obtenir les composés de formule (IA), dans laquelle :


X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³ et R⁴ représentent des atomes d'hydrogène; et

R² représente un groupe amino; ou

(D) réaction du produit de l'étape (C)

avec un ester d'acide α-halogénocarboxylique, de formule :



dans laquelle R représente indépendamment un groupe benzyle ou un groupe alkyle en C₁₋₄;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :


X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou

(E) réaction d'un composé de formule :



(VA)

dans laquelle :

Q¹ représente (CHR⁵)_wCO₂R;

chaque R représente indépendamment un groupe benzyle ou un groupe alkyle en C₁₋₄;

chaque R⁵ représente indépendamment un atome d'hydrogène ou un groupe alkyle en C₁₋₄;

X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;


m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

w vaut 0 ou 1;

sous réserve que la somme de r et w vaille 0 ou 1;

avec un ester d'acide α-halogénocarboxylique, de formule :



dans laquelle R représente indépendamment un groupe benzyle ou un groupe alkyle en C₁₋₄;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la

température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, m, n et r ont les définitions indiquées ci-dessus;

R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe nitro; ou

(F) réaction du produit de l'étape (E) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, m, n et r ont les définitions indiquées ci-dessus;

R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou

(G) réaction du produit de l'étape (B) ou (D) avec un agent de désestérification, comme un acide fort, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un atome d'hydrogène; et

R² représente un groupe amino; ou

(H) réaction du produit de l'étape (F) avec un agent de désestérification, comme un acide fort, pour obtenir les composés de formule (IA), dans laquelle :


X, Y, Q¹, m, n et r ont les définitions indiquées ci-dessus;

R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un atome d'hydrogène; et

R² représente un groupe amino; ou

(I) réaction d'un composé de formule (VA) définie plus haut, avec un α -benzènesulfonate optiquement actif, de formule :



dans laquelle R et R⁵ ont les définitions indiquées ci-dessus,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, R⁵, m, n et r ont les définitions indiquées ci-dessus;

R³ et R⁴ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou benzyle; et

R² représente un groupe nitro; ou

(J) réaction du produit de l'étape (I) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, R⁵, m, n et r ont les définitions indiquées ci-dessus;


R³ et R⁴ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou

(K) réaction du produit de l'étape (J) avec un agent de désestérification, comme un acide fort, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, R⁵, m, n et r ont les définitions indiquées ci-dessus;
 R³, R⁴ et R représentent des atomes d'hydrogène; et
 R² représente un groupe amino; ou
 (L) réaction d'un composé de formule :



(IVA)

dans laquelle :

X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;


R^{3'} représente un groupe alcoxy en C₁₋₄, carboxyméthoxy ou hydroxy;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

sous réserve que n vaille 1 quand X et/ou Y forment une liaison supplémentaire carbone-carbone; et d'un acide α-halogénocarboxylique, de formule :



en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, R^{3'}, m, n et r ont les définitions indiquées ci-dessus;

Q¹ et R² représentent des atomes d'hydrogène; et

R⁴ représente un groupe nitro; ou


(M) réaction du produit de l'étape (L) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, R^{3'}, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R, R² et R⁵ représentent des atomes d'hydrogène; et

R⁴ représente un groupe amino; ou

(N) réaction d'un composé de formule :



(VIIIA)

dans laquelle :

X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;

Q¹ représente CH₂CO₂R;


R représente un groupe benzyle ou alkyle en C₁₋₄;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

sous réserve que n vaille 1 quand X et/ou Y forment une liaison supplémentaire carbone-carbone; et d'un ester d'acide α-halogénocarboxylique, de formule :



dans laquelle R a la définition indiquée ci-dessus;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, Q, m, n et r ont les définitions indiquées ci-dessus;

R² représente un atome d'hydrogène; et

R⁴ représente un groupe nitro; ou

(O) réaction du produit de l'étape (N) avec un agent de désestérification, comme un acide fort, puis avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, Q, m, n et r ont les définitions indiquées ci-dessus;

R, R² et R⁵ représentent des atomes d'hydrogène; et

R² représente un groupe amino; ou


(P) réaction du produit de l'étape (G), (F), (K), (M) ou (O) avec du thiophosgène, pour obtenir le composé correspondant dans lequel R² ou R⁴ représente un groupe isothiocyanato.

77. Procédé de préparation d'un polyazamacrocyclé mono-N-alkylé, qui comporte la réaction d'un ester d'acide α-halogénocarboxylique avec de un à cinq équivalents d'un 1,4,7,10-tétraazacyclododécane, dans un solvant qui ne favorise pas le transfert des protons.

78. Procédé conforme à la revendication 77, dans lequel l'ester d'acide α-halogénocarboxylique est du d,l-2-bromo-4-(4-nitrophényl) butanoate de méthyle, du 2-bromo-2-(4-nitrophényl)éthanoate de méthyle, de l'α-bromo-(2-méthoxy-5-nitrophényl)acétate de méthyle ou du d,l-2-bromo-4-(4-nitrophényl)butanoate d'isopropyle.

Revendications pour les Etats contractants suivants : ES, GR

1. Procédé de préparation d'un composé de formule :



dans laquelle :

chaque Q représente indépendamment un atome d'hydrogène ou un groupe (CHR⁵)_pCO₂R;

Q' représente un atome d'hydrogène ou un groupe $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

chaque R représente indépendamment un atome d'hydrogène, un groupe benzyle ou un groupe alkyle en C_{1-4} ;

sous réserve qu'au moins deux de tous les Q et Q' représentent autre chose qu'un atome d'hydrogène, et que, lorsque Q' représente un atome d'hydrogène, R³ représente autre chose qu'un atome d'hydrogène;

chaque R⁵ représente indépendamment un atome d'hydrogène ou un groupe alkyle en C_{1-4} ;

X et Y représentent chacun indépendamment un atome d'hydrogène, ou peuvent constituer, conjointement avec un X ou Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

p vaut 1 ou 2;

r vaut 0 ou 1;

w vaut 0 ou 1;

sous réserve que n vaille seulement 1 lorsque X et/ou Y forment une liaison carbone-carbone supplémentaire, et que la somme de r et w vaille 0 ou 1;

R² représente un atome d'hydrogène ou un groupe nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido ou carboxy;

R³ représente un groupe alcoxy en C_{1-4} hydroxy ou carboxyméthoxy, ou un atome d'hydrogène;


R⁴ représente un atome d'hydrogène ou un groupe nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maléimido, bromoacétamido ou carboxy;

sous réserve que R² et R⁴ ne représentent pas tous les deux des atomes d'hydrogène, mais que l'un de R² et R⁴ représente un atome d'hydrogène; ou

d'un sel d'un tel composé, acceptable en pharmacie;

lequel procédé comprend l'une des étapes suivantes :

(A) réaction d'un composé de formule :



(IIIA)

dans laquelle :


X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

sous réserve que n vaille 1 quand X et/ou Y forment une liaison supplémentaire carbone-carbone; et d'un ester d'acide α -halogénocarboxylique, de formule :



dans laquelle R représente indépendamment un groupe benzyle ou un groupe alkyle en C_{1-4} ;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q', R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe nitro; ou

(B) réaction du produit de l'étape (A) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou

(C) réaction d'un composé de formule (IIIA) définie plus haut avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, puis avec du chlorure d'hydrogène gazeux, en présence d'un solvant, pour obtenir les composés de formule (IA), dans laquelle :


X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³ et R⁴ représentent des atomes d'hydrogène; et

R² représente un groupe amino; ou

(D) réaction du produit de l'étape (C)

avec un ester d'acide α-halogénocarboxylique, de formule :



dans laquelle R représente indépendamment un groupe benzyle ou un groupe alkyle en C₁₋₄;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :


X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R³, R⁴ et R⁵ représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou

(E) réaction d'un composé de formule :



(VA)

dans laquelle :

Q¹ représente (CHR⁵)_wCO₂R;

chaque R représente indépendamment un groupe benzyle ou un groupe alkyle en C₁₋₄;

chaque R⁵ représente indépendamment un atome d'hydrogène ou un groupe alkyle en C₁₋₄;

X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;

n vaut 0 ou 1;


m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

w vaut 0 ou 1;

sous réserve que la somme de r et w vaille 0 ou 1;

avec un ester d'acide α-halogénocarboxylique, de formule :



dans laquelle R représente indépendamment un groupe benzyle ou un groupe alkyle en C_{1-4} ;
en présence d'un solvant organique et d'une base, à une température valant de 0°C à la
température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q^1 , m, n et r ont les définitions indiquées ci-dessus;

R^3 , R^4 et R^5 représentent des atomes d'hydrogène;

R représente un groupe alkyle en C_{1-4} ou un groupe benzyle; et

R^2 représente un groupe nitro; ou

(F) réaction du produit de l'étape (E) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q^1 , m, n et r ont les définitions indiquées ci-dessus;

R^3 , R^4 et R^5 représentent des atomes d'hydrogène;

R représente un groupe alkyle en C_{1-4} ou un groupe benzyle; et

R^2 représente un groupe amino; ou

(G) réaction du produit de l'étape (B) ou (D) avec un agent de désestérification, comme un acide fort, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, m, n et r ont les définitions indiquées ci-dessus;

Q^1 , R^3 , R^4 et R^5 représentent des atomes d'hydrogène;

R représente un atome d'hydrogène; et

R^2 représente un groupe amino; ou

(H) réaction du produit de l'étape (F) avec un agent de désestérification, comme un acide fort, pour obtenir les composés de formule (IA), dans laquelle :


X, Y, Q^1 , m, n et r ont les définitions indiquées ci-dessus;

R^3 , R^4 et R^5 représentent des atomes d'hydrogène;

R représente un atome d'hydrogène; et

R^2 représente un groupe amino; ou

(I) réaction d'un composé de formule (VA) définie plus haut, avec un α -benzènesulfonate optiquement actif, de formule :



dans laquelle R et R^5 ont les définitions indiquées ci-dessus,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q^1 , R^5 , m, n et r ont les définitions indiquées ci-dessus;

R^3 et R^4 représentent des atomes d'hydrogène;

R représente un groupe alkyle en C_{1-4} ou benzyle; et

R^2 représente un groupe nitro; ou

(J) réaction du produit de l'étape (I) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q^1 , R^5 , m, n et r ont les définitions indiquées ci-dessus;

R^3 et R^4 représentent des atomes d'hydrogène;

R représente un groupe alkyle en C₁₋₄ ou un groupe benzyle; et

R² représente un groupe amino; ou


(K) réaction du produit de l'étape (J) avec un agent de désestérification, comme un acide fort, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q¹, R³, m, n et r ont les définitions indiquées ci-dessus;

R³, R⁴ et R représentent des atomes d'hydrogène; et

R² représente un groupe amino; ou

(L) réaction d'un composé de formule :



(IVA)

dans laquelle :

X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;


R^{3'} représente un groupe alcoxy en C₁₋₄, carboxyméthoxy ou hydroxy;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

sous réserve que n vaille 1 quand X et/ou Y forment une liaison supplémentaire carbone-carbone; et d'un acide α -halogénocarboxylique, de formule :



en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, R^{3'}, m, n et r ont les définitions indiquées ci-dessus;

Q¹ et R² représentent des atomes d'hydrogène; et

R⁴ représente un groupe nitro; ou


(M) réaction du produit de l'étape (L) avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, R^{3'}, m, n et r ont les définitions indiquées ci-dessus;

Q¹, R, R² et R⁵ représentent des atomes d'hydrogène; et

R⁴ représente un groupe amino; ou

(N) réaction d'un composé de formule :



(VIII A)

dans laquelle :

X et Y représentent chacun indépendamment un atome d'hydrogène ou peuvent former, avec un X ou un Y adjacent, une liaison carbone-carbone supplémentaire;

Q' représente $\text{CH}_2\text{CO}_2\text{R}$;


R représente un groupe benzyle ou alkyle en C_{1-4} ;

n vaut 0 ou 1;

m représente un nombre entier valant de 0 à 10, limites comprises;

r vaut 0 ou 1;

sous réserve que n vaille 1 quand X et/ou Y forment une liaison supplémentaire carbone-carbone; et d'un ester d'acide α -halogénocarboxylique, de formule :



dans laquelle R a la définition indiquée ci-dessus;

en présence d'un solvant organique et d'une base, à une température valant de 0°C à la température de reflux,

pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q', Q, m, n et r ont les définitions indiquées ci-dessus;

R² représente un atome d'hydrogène; et

R⁴ représente un groupe nitro; ou

(O) réaction du produit de l'étape (N) avec un agent de désestérification, comme un acide fort, puis avec un agent d'hydrogénation, comme de l'hydrogène en présence de Pd/C, dans les conditions habituelles de réaction, pour obtenir les composés de formule (IA), dans laquelle :

X, Y, Q', Q, m, n et r ont les définitions indiquées ci-dessus;

R, R² et R⁵ représentent des atomes d'hydrogène; et

R² représente un groupe amino; ou

(P) réaction du produit de l'étape (G), (F), (K), (M) ou (O) avec du thiophosgène, pour obtenir le composé correspondant dans lequel R² ou R⁴ représente un groupe isothiocyanato.

2. Procédé conforme à la revendication 1, permettant de préparer un sel d'un composé de formule (IA), acceptable en pharmacie, lequel procédé comporte le fait de faire réagir un composé de formule (IA) avec un réactif, selon des procédés connus, pour obtenir, en tant que sel acceptable en pharmacie, un sel de potassium, sodium, lithium, ammonium, calcium ou magnésium ou un chlorhydrate.
3. Procédé conforme à la revendication 1 (B), permettant de préparer de l'ester 4,7,10-triméthylque de l'acide 1-[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique, lequel procédé comporte le fait de faire réagir de l'ester 4,7,10-triméthylque de l'acide 1-[2-(4-nitrophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique avec de l'hydrogène et du charbon palladié.
4. Procédé conforme à la revendication 1 (G), permettant de préparer du chlorhydrate de l'acide 1-[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique, lequel procédé comporte le fait de faire réagir de l'ester 4,7,10-triméthylque de l'acide 1-[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacy-

clododécane-4,7,10-triacétique avec de l'acide chlorhydrique.

- 5 5. Procédé conforme à la revendication 1 (P), permettant de préparer de l'acide 1-[2-(4-isothiocyaziatophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique, lequel procédé comporte le fait de faire réagir du chlorhydrate de l'acide 1-[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique avec du thiophosgène.
- 10 6. Procédé conforme à la revendication 1 (A), permettant de préparer du chlorhydrate de l'acide 1-(4-aminobenzyl)-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique, lequel procédé comporte le fait de faire réagir du tétrachlorhydrate de 1-(4-aminobenzyl)-1,4,7,10-tétraazacyclododécane avec du bromoacétate de méthyle.
- 15 7. Procédé conforme à la revendication 1 (M), permettant de préparer de l'acide 1-(2-méthoxy-5-aminobenzyl)-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique, lequel procédé comporte le fait de faire réagir de l'acide 1-(2-méthoxy-5-nitrobenzyl)-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique avec de l'hydrogène, dans une bombe de Parr.
- 20 8. Procédé conforme à la revendication 1 (M), permettant de préparer de l'acide 1-(2-hydroxy-5-aminobenzyl)-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique, lequel procédé comporte le fait de faire réagir de l'acide N-(2-hydroxy-5-nitrobenzyl)-1,4,7,10-tétraazacyclododécane-4,7,10-triacétique avec de l'hydrogène et du PtO₂.
- 25 9. Procédé conforme à la revendication 1 (J), permettant de préparer de l'ester 1-isopropylique-4,7,10-triméthylque de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthylacétique), lequel procédé comporte le fait de faire réagir, avec de l'hydrogène et du charbon palladié, de l'ester 1-isopropylique-4,7,10-triméthylque de l'acide α -[2-(4-nitrophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthyl-acétique).
- 30 10. Procédé conforme à la revendication 1 (K), permettant de préparer de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthylacétique), lequel procédé comporte le fait de faire réagir de l'ester 1-isopropylique-4,7,10-triméthylque de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthyl-acétique) avec de l'acide chlorhydrique.
- 35 11. Procédé conforme à la revendication 1 (P), permettant de préparer de l'acide α -[2-(4-isothiocyantophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthylacétique), lequel procédé comporte le fait de faire réagir de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-(R,S)-acétique-4,7,10-tris-(R-méthyl-acétique) avec du thiophosgène.
- 40 12. Procédé conforme à la revendication 1 (F), permettant de préparer de l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7-triacétique, lequel procédé comporte le fait de faire réagir de l'acide α -(4-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7-triacétique avec de l'hydrogène et du PtO₂.
- 45 13. Procédé conforme à la revendication 1 (F), permettant de préparer de l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,10-triacétique, lequel procédé comporte le fait de faire réagir de l'acide α -(4-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,10-triacétique avec de l'hydrogène et du PtO₂.
- 50 14. Procédé conforme à la revendication 1 (F), permettant de préparer de l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'acide α -(4-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec de l'hydrogène et du PtO₂.
- 55 15. Procédé conforme à la revendication 1 (P), permettant de préparer de l'acide α -(4-isothiocyantophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'acide α -(4-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec du thiophosgène.

16. Procédé conforme à la revendication 1 (E), permettant de préparer de l'ester 1,4,7,10-tétraméthylrique de l'acide α -[2-(4-nitrophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'ester 1-méthylrique de l'acide α -[2-(4-nitrophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1-acétique avec du bromoacétate de méthyle.
17. Procédé conforme à la revendication 1 (F), permettant de préparer de l'ester 1,4,7,10-tétraméthylrique de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'ester 1,4,7,10-tétraméthylrique de l'acide α -[2-(4-nitrophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec de l'hydrogène et du charbon palladié.
18. Procédé conforme à la revendication 1 (H), permettant de préparer de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'ester 1,4,7,10-tétraméthylrique de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec de l'acide chlorhydrique.
19. Procédé conforme à la revendication 1 (P), permettant de préparer de l'acide α -[2-(4-isothiocyanatophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'acide α -[2-(4-aminophényl)éthyl]-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec du thiophosgène.
20. Procédé conforme à la revendication 1 (O), permettant de préparer de l'acide α -(2-méthoxy-5-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'ester tétraméthylrique de l'acide α -(2-méthoxy-5-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec de l'acide chlorhydrique.
21. Procédé conforme à la revendication 1 (O), permettant de préparer du sel tétraammonique de l'acide α -(2-méthoxy-5-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'acide α -(2-méthoxy-5-nitrophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec de l'hydrogène et du PtO_2 .
22. Procédé conforme à la revendication 1 (P), permettant de préparer du sel tétraammonique de l'acide (2-méthoxy-5-isocyanatophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique, lequel procédé comporte le fait de faire réagir de l'acide α -(2-méthoxy-5-aminophényl)-1,4,7,10-tétraazacyclododécane-1,4,7,10-tétraacétique avec du thiophosgène.
23. Procédé de préparation d'un complexe d'un composé conforme à l'une des revendications 1 à 22 et d'un ion d'un métal choisi parmi La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y et Sc, lequel procédé comporte le fait de faire réagir le composé avec l'ion métallique.
24. Procédé conforme à la revendication 23, dans lequel le métal est ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc ou ^{142}Pr .
25. Procédé conforme à la revendication 24, dans lequel le métal est ^{153}Sm , ^{177}Lu ou ^{90}Y .
26. Procédé conforme à l'une des revendications 23 à 25, dans lequel la réaction s'effectue à chaud.
27. Procédé de préparation d'un conjugué contenant un complexe d'un composé conforme à l'une des revendications 1 à 22 et d'un ion d'un métal choisi parmi La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Y et Sc, lequel procédé comporte le fait de lier le complexe, par liaison covalente, à un anticorps ou à un fragment d'anticorps.
28. Procédé conforme à la revendication 27, dans lequel le métal est ^{153}Sm , ^{166}Ho , ^{90}Y , ^{149}Pm , ^{159}Gd , ^{140}La , ^{177}Lu , ^{175}Yb , ^{47}Sc ou ^{142}Pr .
29. Procédé conforme à la revendication 28, dans lequel le métal est ^{153}Sm , ^{177}Lu ou ^{90}Y .

30. Conjugué préparé selon l'une des revendications 27 à 29, dans lequel l'anticorps ou le fragment d'anticorps est un anticorps monoclonal ou un fragment d'un tel anticorps.

31. Conjugué conforme à la revendication 30, dans lequel l'anticorps ou le fragment d'anticorps est CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ ou B72.3 (ATCC HB 8108).

32. Procédé de préparation d'un conjugué contenant un composé conforme à l'une des revendications 1 à 22, lié par liaison covalente à un anticorps ou à un fragment d'anticorps, lequel procédé comporte le fait de faire réagir le composé avec un anticorps ou un fragment d'anticorps.

33. Conjugué conforme la revendication 32, dans lequel l'anticorps ou le fragment d'anticorps est un anticorps monoclonal ou un fragment d'un tel anticorps.

34. Conjugué conforme à la revendication 33, dans lequel l'anticorps ou le fragment d'anticorps est CC-49 (ATCC HB 9459), CC-49 F(ab')₂, CC-83 (ATCC HB 9453), CC-83 F(ab')₂ ou B72.3 (ATCC HB 8108).

35. Procédé de préparation d'un polyazamacrocyclole mono-N-alkylé, qui comporte la réaction d'un ester d'acide α -halogénocarboxylique avec de un à cinq équivalents d'un 1,4,7,10-tétraazacyclododécane, dans un solvant qui ne favorise pas le transfert des protons.

36. Procédé conforme à la revendication 35, dans lequel l'ester d'acide α -halogénocarboxylique est du d,l-2-bromo-4-(4-nitrophényl) butanoate de méthyle, du 2-bromo-2-(4-nitrophényl)éthanoate de méthyle, de l' α -bromo-(2-méthoxy-5-nitrophényl)acétate de méthyle ou du d,l-2-bromo-4-(4-nitrophényl)butanoate d'isopropyle.




FIG.1




FIG.2




FIG.3




FIG. 4




FIG. 5




FIG.6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG.11




FIG.12




FIG.13




FIG.14




FIG.15




FIG.16




FIG.17




FIG.18




FIG. 19




FIG. 20




FIG. 21




FIG.22




FIG. 23




FIG. 24




FIG. 25




FIG.26




FIG.27




FIG. 28




FIG. 29




FIG.30




FIG. 31




FIG. 32




FIG. 33




FIG. 34

RE

REGISTER ENTRY FOR EP0353450

European Application No EP89111497.7 filing date 23.06.1989

Priorities claimed:

24.06.1988 in United States of America - doc: 211496

21.06.1989 in United States of America - doc: 370956

Designated States BE CH DE ES FR GB GR IT LI LU NL SE AT

Title MACROCYCLIC BIFUNCTIONAL CHELANTS, COMPLEXES THEREOF AND THEIR ANTIBODY CONJUGATES.

Applicant/Proprietor

THE DOW CHEMICAL COMPANY, 2030 Dow Center, Abbott Road, Midland MI 48640,
United States of America [ADP No. 50336619004]

Inventors

ROBERTA C. CHENG, 3873 Old Pine Trail, Midland Michigan 48640, United
States of America [ADP No. 54286604001]WILLIAM A. FORDYCE, 4103 Washington Street, Midland Michigan 48640, United
States of America [ADP No. 55507594001]WILLIAM F. GOECKELER, 4916 Sturgeon Creek Pkwy, Midland Michigan 48640,
United States of America [ADP No. 57192676001]JR. WILLIAM F. KRUPER, 230 Barden Road, Sanford Michigan 48657, United
States of America [ADP No. 57192684001]SHARON BAUGHMAN, 111 Pinestone, Irvine California 92714, United States of
America [ADP No. 57192692001]JOSEPH R. GARLICH, 301 Southern Oaks, Lake Jackson Texas 77566, United
States of America [ADP No. 57192700001]GARRY KIEFER, 400 Timber Creek Dr. Nr. 913, Richwood Texas 77531, United
States of America [ADP No. 57192718001]KENNETH MCMILLAN, 405 Moore Street, Richwood Texas 77531, United States of
America [ADP No. 57192726001]JAIME J. SIMON, Route 1, Box 120-G, Angleton Texas 77515, United States of
America [ADP No. 57192734001]DAVID A. WILSON, 229 San Saba, Richwood Texas 77531, United States of
America [ADP No. 53238580001]RICHARD KEITH FRANK, 722 Sycamore, Lake Jackson Texas 77566, United States
of America [ADP No. 57192742001]

Classified to

C07D A61K

Address for Service

W H BECK, GREENER & CO, 7 Stone Buildings, Lincoln's Inn, LONDON, WC2A
3SZ, United Kingdom [ADP No. 00000323001]

EPO Representative

DIPL.-CHEM. BERNHARD HUBER, Patentanwälte H. Weickmann, Dr. K. Fincke F.A.
Weickmann, B. Huber Dr. H. Liska, Dr. J. Prechtel Möhlstrasse 22 Postfach
860 820, D-8000 München 86, Federal Republic of Germany

[ADP No. 50771450001]

Publication No EP0353450 dated 07.02.1990

Publication in English

Examination requested 12.07.1990

Patent Granted with effect from 29.03.1995 (Section 25(1)) with title
MACROCYCLIC BIFUNCTIONAL CHELANTS, COMPLEXES THEREOF AND THEIR ANTIBODY
CONJUGATES.

11.10.1994 W H BECK, GREENER & CO, 7 Stone Buildings, Lincoln's Inn, LONDON,
WC2A 3SZ, United Kingdom [ADP No. 00000323001]
registered as address for service

Entry Type 8.11 Staff ID. SS1 Auth ID. AA

02.12.1994 Notification from EPO of change of EPO Representative details from
DIPL.-CHEM. BERNHARD HUBER, Patentanwälte H. Weickmann, Dr. K.
Fincke F.A. Weickmann, B. Huber Dr. H. Liska, Dr. J. Prechtel
Möhlstrasse 22 Postfach 860 820, D-8000 München 86, Federal
Republic of Germany [ADP No. 50771450001]

to
DIPL.-CHEM. BERNHARD HUBER, Patentanwälte H. Weickmann, Dr. K.
Fincke F.A. Weickmann, B. Huber Dr. H. Liska, Dr. J. Prechtel, Dr.
B. Böhm Postfach 86, D-81635 München, Federal Republic of Germany
[ADP No. 50067172001]

Entry Type 25.14 Staff ID. RD06 Auth ID. EPT

24.02.1995 Notification from EPO of change of Applicant/Proprietor details
from
THE DOW CHEMICAL COMPANY, 2030 Dow Center, Abbott Road, Midland MI
48640, United States of America [ADP No. 50336619004]

to
THE DOW CHEMICAL COMPANY, 2030 Dow Center, Abbott Road, Midland,
Michigan 48640, United States of America [ADP No. 50336619004]

Entry Type 25.14 Staff ID. RD06 Auth ID. EPT

**** END OF REGISTER ENTRY ****

OA80-01
EP

OPTICS - PATENTS

21/07/95 11:33:11
PAGE: 1

RENEWAL DETAILS

PUBLICATION NUMBER EP0353450 /

PROPRIETOR(S)

THE DOW CHEMICAL COMPANY, 2030 Dow Center, Abbott Road, Midland,
Michigan 48640, United States of America

DATE FILED 23.06.1989 /

DATE GRANTED 29.03.1995 /

DATE NEXT RENEWAL DUE 23.06.1996

DATE NOT IN FORCE

DATE OF LAST RENEWAL 30.05.1995

YEAR OF LAST RENEWAL 07

STATUS PATENT IN FORCE /

**** END OF REPORT ****