

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 February 2002 (21.02.2002)

PCT

(10) International Publication Number
WO 02/14355 A2

(51) International Patent Classification⁷: **C07K 14/00**

(21) International Application Number: PCT/EP01/09109

(22) International Filing Date: 7 August 2001 (07.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00117425.9 11 August 2000 (11.08.2000) EP

(71) Applicant (for all designated States except US): **MERCK PATENT GMBH** [DE/DE]; Frankfurter Strasse 250, 64293 Darmstadt (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **SCHARM, Burkhard** [DE/DE]; Mailänderstrasse 12/1570, 60598 Frankfurt/Main (DE).

(74) Common Representative: **MERCK PATENT GMBH**; Frankfurter Strasse 250, 64293 Darmstadt (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL MITOGEN ACTIVATED KINASE

(57) Abstract: MEKK3b polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing MEKK3b polypeptides and polynucleotides in diagnostic assays.



WO 02/14355 A2

Novel mitogen activated kinase

Field of the Invention

5 This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides sometimes hereinafter referred to as „novel mitogen activated kinase (MEKK3b)“, to their use in diagnosis and in identifying compounds that may be agonists, antagonists that are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

10

Background of the Invention

15 The drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics", that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superceding earlier approaches based on "positional cloning". A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

20 Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug
25 discovery.

30 It is one aspect of this invention, that the protein function relates to a mitogen-activated protein kinase (MAPK). It is well documented that the MAPK and stress activated protein kinase signal transduction pathways are responsible for triggering biological effects across a variety of pathophysiological conditions including conditions manifested by dysfunctional leukocytes, T-lymphocytes, acute and chronic inflammatory disease, auto-immune disorders, rheumatoid arthritis, osteoarthritis, transplant rejection, macrophage regulation, endothelial cell regulation,

angiogenesis, atherosclerosis fibroblasts regulation, pathological fibrosis, asthma, allergic response, ARDS, atheroma, osteoarthritis, heart failure, cancer, diabetes, obesity, cachexia, Alzheimers disease, sepsis, and neurodegeneration. As MAP kinases play a central role in signaling events which mediate cellular responses, controlling the activity of MAPK is key to the attenuation of the response. N. J. Holbrook, et al., Stress-Inducible Cellular Responses, 273, Feige, U., et al., Eds., Birkhauser Verlag (1996).

Summary of the Invention

The present invention relates to MEKK3b, in particular MEKK3b polypeptides and MEKK3b polynucleotides, recombinant materials and methods for their production. Such polypeptides and polynucleotides are of interest in relation to methods of treatment of certain diseases, including, but not limited to, to cell cycle control and/or apoptosis, including, but not limited to cancer, metabolic disorders, heart diseases and inflammatory diseases as multiple sclerosis herein after referred to as "diseases of the invention". In a further aspect, the invention relates to methods for identifying agonists and antagonists (e.g., inhibitors) using the materials provided by the invention, and treating conditions associated with MEKK3b imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate MEKK3b activity or levels.

Description of the Invention

In a first aspect, the present invention relates to MEKK3b polypeptides. Such polypeptides include:

(a) a polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3;

(b) a polypeptide comprising a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;

- (c) a polypeptide comprising the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;
- (d) a polypeptide having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;
- 5 (e) the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4; and
- (f) a polypeptide having or comprising a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;
- 10 (g) fragments and variants of such polypeptides in (a) to (f).

Polypeptides of the present invention are believed to be members of the serine/threonine kinase family of polypeptides. They are therefore of interest because MAPKKK (Garrington T. Curr Opinion Cell Biol. 1999, 11: 211-218-192) kinases are important regulatory proteins in signal transduction especially in ras induced pathways. They are involved in cell cycle control, cell growth, differentiation as angiogenesis (Yang J. nat genetics 2000, 24th march: 309-313), and metabolic pathways and cause in case of impaired control cancer and metabolic diseases..

15

The biological properties of the MEKK3b are hereinafter referred to as "biological activity of MEKK3b" or "MEKK3b activity". Preferably, a polypeptide of the present invention exhibits at least one biological activity of MEKK3b.

20

Polypeptides of the present invention also includes variants of the aforementioned polypeptides, including all allelic forms and splice variants. Such polypeptides vary from the reference polypeptide by insertions, deletions, and substitutions that may be conservative or non-conservative, or any combination thereof. Particularly preferred variants are those in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acids are inserted, substituted, or deleted, in any combination.

25

30

Preferred fragments of polypeptides of the present invention include a polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID

5 NO: 2 and/or SEQ ID NO: 4, or a polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4. Preferred fragments are biologically active fragments that mediate the biological activity of MEKK3b, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also preferred are those fragments that are antigenic or immunogenic in an animal, especially in a human.

10 Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention. The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is
15 often advantageous to include an additional amino acid sequence that contains secretory or leader sequences, pro-sequences, sequences that aid in purification, for instance multiple histidine residues, or an additional sequence for stability during recombinant production.

20 Polypeptides of the present invention can be prepared in any suitable manner, for instance by isolation from naturally occurring sources, from genetically engineered host cells comprising expression systems (*vide infra*) or by chemical synthesis, using for instance automated peptide synthesizers, or a combination of such methods.. Means for preparing such polypeptides are well understood in the art.

25

In a further aspect, the present invention relates to MEKK3b polynucleotides. Such polynucleotides include:

30 (a) a polynucleotide comprising a polynucleotide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3;

(b) a polynucleotide comprising the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3;

(c) a polynucleotide having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3;

(d) the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3;

5 (e) a polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;

(f) a polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4;

10 (g) a polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;

15 (h) a polynucleotide encoding the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4;

(i) a polynucleotide having or comprising a polynucleotide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polynucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3;

20 (j) a polynucleotide having or comprising a polynucleotide sequence encoding a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4; and

25 polynucleotides that are fragments and variants of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.

30 Preferred fragments of polynucleotides of the present invention include a polynucleotide comprising an nucleotide sequence having at least 15, 30, 50 or 100 contiguous nucleotides from the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3, or a polynucleotide comprising an sequence having at least 30, 50 or 100 contiguous nucleotides truncated or deleted from the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3.

Preferred variants of polynucleotides of the present invention include splice variants, allelic variants, and polymorphisms, including polynucleotides having one or more single nucleotide polymorphisms (SNPs).

5 Polynucleotides of the present invention also include polynucleotides encoding polypeptide variants that comprise the amino acid sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4 and in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acid residues are substituted, deleted or added, in
10 any combination.

In a further aspect, the present invention provides polynucleotides that are RNA transcripts of the DNA sequences of the present invention. Accordingly, there is provided an RNA polynucleotide that:

15 (a) comprises an RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4;

(b) is the RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4;

(c) comprises an RNA transcript of the DNA sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3; or

20 (d) is the RNA transcript of the DNA sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3;

and RNA polynucleotides that are complementary thereto.

25 The polynucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3 shows homology with MEKK3 (ACC NM_002401). The polynucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3 is a cDNA sequence that encodes the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4. The polynucleotide sequence encoding the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4 may be identical to the polypeptide encoding
30 sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3 or it may be a sequence other than SEQ ID NO: 1 and/or SEQ ID NO: 3, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes

the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4. The polypeptide of the SEQ ID NO: 2 and/or SEQ ID NO: 4 is related to other proteins of the serine/threonine kinase family, having homology and/or structural similarity with MEKK3 (NM_002401).

5 Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one MEKK3b activity.

10

Polynucleotides of the present invention may be obtained using standard cloning and screening techniques from a cDNA library derived from mRNA in cells of human testis, (see for instance, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). Polynucleotides of the invention
15 can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

When polynucleotides of the present invention are used for the
20 recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence,
25 or other fusion peptide portions. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, Proc Natl Acad Sci USA (1989) 86:821-824,
30 or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Polynucleotides that are identical, or have sufficient identity to a
35 polynucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3, may be

used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification reaction (for instance, PCR). Such probes and primers may be used to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO: 1 and/or SEQ ID NO: 3, typically at least 95% identity. Preferred probes and primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50, if not at least 100 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process comprising the steps of screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3 or a fragment thereof, preferably of at least 15 nucleotides; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes isolated polynucleotides, preferably with a nucleotide sequence of at least 100, obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3 or a fragment thereof, preferably of at least 15 nucleotides.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide does not extend all the way through to the 5' terminus. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during first strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., Proc Nat Acad Sci USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon (trade mark) technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon (trade mark) technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems comprising a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Polynucleotides may be introduced into host cells by methods described in many standard laboratory manuals, such as Davis et

al., Basic Methods in Molecular Biology (1986) and Sambrook *et al.*(*ibid*). Preferred methods of introducing polynucleotides into host cells include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, *e.g.*, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector that is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate polynucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, (*ibid*). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to

recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and/or purification.

Polynucleotides of the present invention may be used as diagnostic reagents, through detecting mutations in the associated gene. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3 in the cDNA or genomic sequence and which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques well known in the art.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or it may be amplified enzymatically by using PCR, preferably RT-PCR, or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled MEKK3b nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence difference may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, for instance, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations

may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, Proc Natl Acad Sci USA (1985) 85: 4397-4401).

5 An array of oligonucleotides probes comprising MEKK3b polynucleotide sequence or fragments thereof can be constructed to conduct efficient screening of *e.g.*, genetic mutations. Such arrays are preferably high density arrays or grids. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage,
10 and genetic variability, see, for example, M.Chee *et al.*, Science, 274, 610-613 (1996) and other references cited therein.

Detection of abnormally decreased or increased levels of polypeptide or mRNA expression may also be used for diagnosing or determining susceptibility of a subject to a disease of the invention. Decreased or
15 increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a
20 protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a
25 diagnostic kit comprising:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3, or a fragment or an RNA transcript thereof;
- (b) a nucleotide sequence complementary to that of (a);
- 30 (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly diseases of the invention, amongst others.

5

The polynucleotide sequences of the present invention are valuable for chromosome localisation studies. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (co-inheritance of physically adjacent genes). Precise human chromosomal localisations for a genomic sequence (gene fragment etc.) can be determined using Radiation Hybrid (RH) Mapping (Walter, M. Spillett, D., Thomas, P., Weissenbach, J., and Goodfellow, P., (1994) A method for constructing radiation hybrid maps of whole genomes, Nature Genetics 7, 22-28). A number of RH panels are available from Research Genetics (Huntsville, AL, USA) e.g. the GeneBridge4 RH panel (Hum Mol Genet 1996 Mar;5(3):339-46 A radiation hybrid map of the human genome. Gyapay G, Schmitt K, Fizames C, Jones H, Vega-Czarny N, Spillett D, Muselet D, Prud'Homme JF, Dib C, Auffray C, Morissette J, Weissenbach J, Goodfellow PN). To determine the chromosomal location of a gene using this panel, 93 PCRs are performed using primers designed from the gene of interest on RH DNAs. Each of these DNAs contains random human genomic fragments maintained in a hamster background (human / hamster hybrid cell lines). These PCRs result in 93 scores indicating the presence or absence of the PCR product of the gene of interest. These scores are compared with scores created using PCR products from genomic sequences of known location. This comparison is conducted at

<http://www.genome.wi.mit.edu/>. The gene of the present invention maps to human chromosome 2 (AC016725; pos. 48267-48965).

5 The polynucleotide sequences of the present invention are also valuable tools for tissue expression studies. Such studies allow the determination of expression patterns of polynucleotides of the present invention which may give an indication as to the expression patterns of the encoded polypeptides in tissues, by detecting the mRNAs that encode them. The techniques used are well known in the art and include in situ hybridisation techniques to clones arrayed on a grid, such as cDNA microarray hybridisation (Schena *et al*, Science, 270, 467-470, 1995 and Shalon *et al*, Genome Res, 6, 639-645, 1996) and nucleotide amplification techniques such as PCR. A preferred method uses the TAQMAN (Trade mark) technology available from Perkin Elmer. Results from these studies can provide an indication of the normal function of the polypeptide in the organism. In addition, comparative studies of the normal expression pattern of mRNAs with that of mRNAs encoded by an alternative form of the same gene (for example, one having an alteration in polypeptide coding potential or a regulatory mutation) can provide valuable insights into the role of the polypeptides of the present invention, or that of inappropriate expression thereof in disease. Such inappropriate expression may be of a temporal, spatial or simply quantitative nature.

25 A further aspect of the present invention relates to antibodies. The polypeptides of the invention or their fragments, or cells expressing them, can be used as immunogens to produce antibodies that are immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

30 Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C.,

35

Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, 77-96, Alan R. Liss, Inc., 1985).

5 Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

10 The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against polypeptides of the present invention may also be employed to treat diseases of the invention, amongst others.

15 Polypeptides and polynucleotides of the present invention may also be used as vaccines. Accordingly, in a further aspect, the present invention relates to a method for inducing an immunological response in a mammal that comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response,
20 including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said animal from disease, whether that disease is already established within the individual or not. An immunological response in a mammal may also be induced by a method comprises delivering a polypeptide of the present invention *via* a vector directing expression of
25 the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases of the invention. One way of administering the vector is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a
30 modified nucleic acid, or a DNA/RNA hybrid. For use a vaccine, a polypeptide or a nucleic acid vector will be normally provided as a vaccine formulation (composition). The formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance,
35 subcutaneous, intramuscular, intravenous, or intradermal injection).

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions that may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions that may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Polypeptides of the present invention have one or more biological functions that are of relevance in one or more disease states, in particular the diseases of the invention hereinbefore mentioned. It is therefore useful to to identify compounds that stimulate or inhibit the function or level of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those that stimulate or inhibit the function or level of the polypeptide. Such methods identify agonists or antagonists that may be employed for therapeutic and prophylactic purposes for such diseases of the invention as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, collections of chemical compounds, and natural product mixtures. Such agonists or antagonists so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; a structural or functional mimetic thereof (see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991)) or a small molecule.

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof, by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve measuring or detecting (qualitatively or quantitatively) the competitive binding of a candidate compound to the

polypeptide against a labeled competitor (e.g. agonist or antagonist). Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring a MEKK3b activity in the mixture, and comparing the MEKK3b activity of the mixture to a control mixture which contains no candidate compound.

Polypeptides of the present invention may be employed in conventional low capacity screening methods and also in high-throughput screening (HTS) formats. Such HTS formats include not only the well-established use of 96- and, more recently, 384-well micotiter plates but also emerging methods such as the nanowell method described by Schullek et al, Anal Biochem., 246, 20-29, (1997).

Fusion proteins, such as those made from Fc portion and MEKK3b polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

25

Screening techniques

The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents that may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

A polypeptide of the present invention may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide that compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

Examples of antagonists of polypeptides of the present invention include antibodies or, in some cases, oligonucleotides or proteins that are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or a small molecule that bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Screening methods may also involve the use of transgenic technology and MEKK3b gene. The art of constructing transgenic animals is well established. For example, the MEKK3b gene may be introduced through microinjection into the male pronucleus of fertilized oocytes, retroviral transfer into pre- or post-implantation embryos, or injection of genetically modified, such as by electroporation, embryonic stem cells into host blastocysts. Particularly useful transgenic animals are so-called "knock-in" animals in which an animal gene is replaced by the human equivalent within the genome of that animal. Knock-in transgenic animals are useful in the drug discovery process, for target validation, where the compound is specific for the human target. Other useful transgenic animals are so-called "knock-out" animals in which the expression of the animal ortholog of a polypeptide of the present invention and encoded by an endogenous DNA sequence in a cell is partially or completely annulled. The gene knock-out may be targeted to specific cells or tissues, may occur only in certain cells or tissues as a consequence of the limitations of the technology, or may occur in all, or substantially all, cells in the animal.

Transgenic animal technology also offers a whole animal expression-cloning system in which introduced genes are expressed to give large amounts of polypeptides of the present invention

5 Screening kits for use in the above described methods form a further aspect of the present invention. Such screening kits comprise:

- (a) a polypeptide of the present invention;
- (b) a recombinant cell expressing a polypeptide of the present invention;
- (c) a cell membrane expressing a polypeptide of the present invention; or
- (d) an antibody to a polypeptide of the present invention;

10 which polypeptide is preferably that of SEQ ID NO: 2 and/or SEQ ID NO: 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

15 Glossary

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

20 "Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an

Fab or other immunoglobulin expression library.

25 "Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method

is "isolated" even if it is still present in said organism, which organism may be living or non-living.

5 "Polynucleotide" generally refers to any polyribonucleotide (RNA) or polydeoxribonucleotide (DNA), which may be unmodified or modified RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. 10 "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides. 20

"Polypeptide" refers to any polypeptide comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. 30 It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. 35

Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, Proteins - Structure and Molecular Properties, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., Post-translational Protein Modifications: Perspectives and Prospects, 1-12, in Post-translational Covalent Modification of Proteins, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol, 182, 626-646, 1990, and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", Ann NY Acad Sci, 663, 48-62, 1992).

"Fragment" of a polypeptide sequence refers to a polypeptide sequence that is shorter than the reference sequence but that retains essentially the same biological function or activity as the reference polypeptide. "Fragment" of a polynucleotide sequence refers to a polynucleotide sequence that is shorter than the reference sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3..

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains the essential properties thereof. A typical variant of a polynucleotide differs in nucleotide sequence from the reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the

reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from the reference polypeptide. Generally, alterations are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, insertions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Typical conservative substitutions include Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe and Tyr. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allele, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Also included as variants are polypeptides having one or more post-translational modifications, for instance glycosylation, phosphorylation, methylation, ADP ribosylation and the like. Embodiments include methylation of the N-terminal amino acid, phosphorylations of serines and threonines and modification of C-terminal glycines.

"Allele" refers to one of two or more alternative forms of a gene occurring at a given locus in the genome.

"Polymorphism" refers to a variation in nucleotide sequence (and encoded polypeptide sequence, if relevant) at a given position in the genome within a population.

"Single Nucleotide Polymorphism" (SNP) refers to the occurrence of nucleotide variability at a single nucleotide position in the genome, within a population. An SNP may occur within a gene or within intergenic regions of the genome. SNPs can be assayed using Allele Specific Amplification (ASA). For the process at least 3 primers are required. A common primer is used in reverse complement to the polymorphism being assayed. This common primer can be between 50 and 1500 bps from the polymorphic base. The other two (or more) primers are identical to each other except that the final 3' base wobbles to match one of the two (or more) alleles that make up the polymorphism. Two (or more) PCR reactions are then conducted on sample DNA, each using the common primer and one of the Allele Specific Primers.

"Splice Variant" as used herein refers to cDNA molecules produced from RNA molecules initially transcribed from the same genomic DNA sequence but which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript undergoes splicing, generally for the removal of introns, which results in the production of more than one mRNA molecule each of that may encode different amino acid sequences. The term splice variant also refers to the proteins encoded by the above cDNA molecules.

"Identity" reflects a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, determined by comparing the sequences. In general, identity refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of the two polynucleotide or two polypeptide sequences, respectively, over the length of the sequences being compared.

"% Identity" - For sequences where there is not an exact correspondence, a "% identity" may be determined. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. This may include inserting "gaps" in either one or both sequences, to enhance the degree of alignment. A % identity may be determined over the whole length of each of the sequences being compared (so-called global alignment), that is particularly suitable for sequences of the same or very similar length, or over shorter, defined lengths (so-called local alignment), that is more suitable for sequences of unequal length.

"Similarity" is a further, more sophisticated measure of the relationship between two polypeptide sequences. In general, "similarity" means a comparison between the amino acids of two polypeptide chains, on a residue by residue basis, taking into account not only exact correspondences between a between pairs of residues, one from each of the sequences being compared (as for identity) but also, where there is not an exact correspondence, whether, on an evolutionary basis, one residue is a likely substitute for the other. This likelihood has an associated "score" from which the "% similarity" of the two sequences can then be determined.

Methods for comparing the identity and similarity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J et al, Nucleic Acids Res, 12, 387-395, 1984, available from Genetics Computer Group, Madison, Wisconsin, USA), for example the programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity and the % similarity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (J Mol Biol, 147,195-197, 1981, Advances in Applied Mathematics, 2, 482-489, 1981) and finds the best single region of similarity between two sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences that are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison, GAP aligns two sequences, finding a "maximum similarity", according to the algorithm of Neddleman and Wunsch (J Mol Biol, 48, 443-453, 1970). GAP is more suited to comparing sequences that are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3, for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively. Preferably, % identities and similarities are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S F et al, J Mol Biol, 215, 403-410, 1990, Altschul S F et al, Nucleic Acids Res., 25:389-3402, 1997, available from the National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at www.ncbi.nlm.nih.gov) and FASTA (Pearson W R, Methods in Enzymology, 183, 63-99, 1990; Pearson W R and Lipman D J, Proc Nat Acad Sci USA, 85, 2444-2448,1988, available as part of the Wisconsin Sequence Analysis Package).

Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S and Henikoff J G, Proc. Nat. Acad Sci. USA, 89, 10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide

sequences are first translated into amino acid sequences before comparison.

Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a reference polynucleotide or a polypeptide sequence, the query and the reference sequence being optimally aligned and the parameters of the program set at the default value, as hereinbefore described.

"Identity Index" is a measure of sequence relatedness which may be used to compare a candidate sequence (polynucleotide or polypeptide) and a reference sequence. Thus, for instance, a candidate polynucleotide sequence having, for example, an Identity Index of 0.95 compared to a reference polynucleotide sequence is identical to the reference sequence except that the candidate polynucleotide sequence may include on average up to five differences per each 100 nucleotides of the reference sequence. Such differences are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion. These differences may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between these terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polynucleotide sequence having an Identity Index of 0.95 compared to a reference polynucleotide sequence, an average of up to 5 in every 100 of the nucleotides of the in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

Similarly, for a polypeptide, a candidate polypeptide sequence having, for example, an Identity Index of 0.95 compared to a reference polypeptide sequence is identical to the reference sequence except that the polypeptide sequence may include an average of up to five differences per each 100 amino acids of the reference sequence. Such differences are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion. These differences may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or

anywhere between these terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polypeptide sequence having an Identity Index of 0.95 compared to a reference polypeptide sequence, an average of up to 5 in every 100 of the amino acids in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

The relationship between the number of nucleotide or amino acid differences and the Identity Index may be expressed in the following equation:

$$n_a \leq x_a - (x_a \bullet I),$$

in which:

n_a is the number of nucleotide or amino acid differences,

x_a is the total number of nucleotides or amino acids in SEQ ID NO: 1 and/or SEQ ID NO: 3 or SEQ ID NO: 2 and/or SEQ ID NO: 4, respectively,

I is the Identity Index ,

$\bullet$ is the symbol for the multiplication operator, and

in which any non-integer product of x_a and I is rounded down to the nearest integer prior to subtracting it from x_a .

"Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a reference sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the two sequences as hereinbefore defined. Falling within this generic term are the terms "ortholog", and "paralog". "Ortholog" refers to a polynucleotide or polypeptide that is the functional equivalent of the polynucleotide or polypeptide in another species. "Paralog" refers to a polynucleotide or polypeptide that within the same species which is functionally similar.

"Fusion protein" refers to a protein encoded by two, unrelated, fused genes or fragments thereof. Examples have been disclosed in US 5541087, 5726044. In the case of Fc-MEKK3b, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for performing the functional expression of Fc-MEKK3b or fragments of -
5 MEEK3b, to improve pharmacokinetic properties of such a fusion protein when used for therapy and to generate a dimeric MEKK3b. The Fc-MEKK3b DNA construct comprises in 5' to 3' direction, a secretion cassette, i.e. a signal sequence that triggers export from a mammalian cell, DNA encoding an immunoglobulin Fc region fragment, as a fusion
10 partner, and a DNA encoding MEKK3b or fragments thereof. In some uses it would be desirable to be able to alter the intrinsic functional properties (complement binding, Fc-Receptor binding) by mutating the functional Fc sides while leaving the rest of the fusion protein untouched or delete the Fc part completely after expression.
15

All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by
20 reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

Further examples

25 Cloning of the full length gene:

A marathon lung cDNA from clontech Laboratories GmbH, Heidelberg Germany was subjected to PCR using gene-specific primer 1 (SEQ ID NO: 6) and 2 (SEQ ID NO: 5) in reverse orientation. The conditions for PCR were 90 sec. at 94°C, 30 sec at 94°C and 1 min 62°C for 5 cycles, 30 sec
30 at 94°C and 1 min 60°C for 5 cycles, 30 sec at 94°C and 1 min 58°C for 32 cycles followed by an extension step 3 min at 72° using the advantage polymerase (clontech). The cDNA amplification product was cloned and sequenced. The 5' and 3' cDNA terminus of the cDNA was completed by the 5'-RACE (rapid amplification of cDNA ends) method (Frohmann et al (1988), Proc. Natl. Acad. Sci. USA (85), 8998-9002.) using the system of
35 clontech (clontech Laboratories GmbH, Heidelberg Germany). A marathon

human lung cDNA (clontech) was subjected to PCR using an anchor primer supplied with the marathon cDNA kit and a gene specific primer 2 (SEQ ID NO:5) in reverse orientation. The conditions for PCR were 1 min at 94°C, 30 sec at 94°C and 30 sec 68°C for 30 cycles using the advantage polymerase (clontech). The 3'-end was cloned in the same way using a gene specific primer 1 (SEQ ID NO:6) in forward direction and a primer supplied by the kit. The resulting products were cloned and sequenced.

Tissue distribution:

A set of normalised human cDNAs derived from heart, liver, skeletal muscle, brain, placenta, lung, kidney and pancreas was used to amplify a short gene fragment to examine the tissue distribution of MEKK3b. For this purpose the clontech Multiple Tissue cDNA Panel I (clontech Laboratories GmbH, Heidelberg Germany) was used with two MEKK3b gene-specific primers 1 and primer 2 (SEQ ID NO: 6 and SEQ ID NO: 5) in reverse orientation. Using the advantage polymerase mixture purchased from clontech a 1.0 kb long PCR fragment could be amplified as indicated in the gel photo. The PCR conditions were 90 sec at 94°C followed by five cycles 30 sec at 94°C, 60 sec. at 62°C and another five cycles 30 sec at 94°C, 60 sec. at 60°C and finally 30 sec 94°C and 60 sec. 58°C for 30 cycles using the advantage polymerase (clontech). A G3PDH 5' specific primer 3 combined with G3PDH 3' specific primer 4 (SEQ ID NO: 7 and SEQ ID NO: 8) served as the positive control for equal amounts of normalized CDNA templates and resulted in a 1,0 kb PCR product.

Figure legend

Fig.1: 1.1% agarose gel of multiple tissue cDNA panels. Human tissues, are indicated. 20 µl of each PCR reaction were loaded on the gel. In the upper panel MEKK3b gene specific primer 1 and 2 (SEQ ID NO: 6 and SEQ ID NO: 5) were used for PCR and in the lower panel G3PDH 5' specific primer 3 and primer 4 (SEQ ID NO: 7 and SEQ ID NO: 8).

Claims

1. A polypeptide selected from the group consisting of:
 - (a) a polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3;
 - 5 (b) a polypeptide comprising a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;
 - c) a polypeptide having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;
 - d) the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4 and
 - 10 (e) fragments and variants of such polypeptides in (a) to (d).
2. The polypeptide of claim 1 comprising the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4.
- 15 3. The polypeptide of claim 1 which is the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4.
4. A polynucleotide selected from the group consisting of:
 - (a) a polynucleotide comprising a polynucleotide sequence having at least 95%
20 identity to the polynucleotide sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3;
 - (b) a polynucleotide having at least 95% identity to the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3;
 - (c) a polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:
25 2 and/or SEQ ID NO: 4;


- (d) a polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 and/or SEQ ID NO: 4;
- (e) a polynucleotide with a nucleotide sequence of at least 100 nucleotides
5 obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 3 or a fragment thereof having at least 15 nucleotides;
- (f) a polynucleotide which is the RNA equivalent of a polynucleotide of (a) to (e);
- (g) a polynucleotide sequence complementary to said polynucleotide of any one of
10 (a) to (f), and
- (h) polynucleotides that are variants or fragments of the polynucleotides of any one of (a) to (g) or that are complementary to above mentioned polynucleotides, over the entire length thereof.
- 15 5. A polynucleotide of claim 4 selected from the group consisting of:
- (a) a polynucleotide comprising the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3;
- (b) the polynucleotide of SEQ ID NO: 1 and/or SEQ ID NO: 3;
- (c) a polynucleotide comprising a polynucleotide sequence encoding the
20 polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4; and
- (d) a polynucleotide encoding the polypeptide of SEQ ID NO: 2 and/or SEQ ID NO: 4.
- 25 6. An expression system comprising a polynucleotide capable of producing a polypeptide of any one of claim 1-3 when said expression vector is present in a compatible host cell.
7. A recombinant host cell comprising the expression vector of claim 6 or a membrane thereof expressing the polypeptide of any one of claim 1-3.

8. A process for producing a polypeptide of any one of claim 1-3 comprising the step of culturing a host cell as defined in claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.
9. A fusion protein consisting of the Immunoglobulin Fc-region and a polypeptide any one one of claims 1-3.
10. An antibody immunospecific for the polypeptide of any one of claims 1 to 3.
11. A method for screening to identify compounds that stimulate or inhibit the function or level of the polypeptide of any one of claim 1-3 comprising a method selected from the group consisting of:
- (a) measuring or, detecting, quantitatively or qualitatively, the binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;
 - (b) measuring the competition of binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;
 - (c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes expressing the polypeptide;
 - (d) mixing a candidate compound with a solution containing a polypeptide of any one of claims 1-3, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a control mixture which contains no candidate compound; or

(e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide or said polypeptide in cells, using for instance, an ELISA assay, and

(f) producing said compound according to biotechnological or chemical standard
5 techniques.

Figure 1:
MTC Panel Human I



SEQUENCE LISTING

<110> Merck Patent GmbH

5 <120> Novel mitogen activated protein kinase

<130> MEKK3bBSWS

<140>

10 <141>

<160> 8

<170> PatentIn Ver. 2.1

15 <210> 1

<211> 1896

<212> DNA

<213> Homo sapiens

20 <220>

<221> CDS

<222> (900)..(1469)

25 <400> 1

atttccaagt ctccacccca ttgacgtcaa tggagtttgt tttggcacca aaatcaacgg 60

actttccaaa atgtcgtacc aactccgccc cattgacgca aatgggcgggt aggcgtgtac 120

30 ggtgggaggt ctatataagc agagctcggt tagtgaaccg tcagatcgcc tggagacgcc 180

atccacgctg ttttgacctc catagaagac accgggaccg atccagcctc cggactctag 240

cctaggccgc gggacggata acaatttcac acaggaaaca gctatgacca ttaggcctat 300

35 ttaggtgaca ctatagaaca agtttgtaca aaaaagcagg ctggtaccgg tccggaattc 360

ccgggatggg aaaggacagt tactgcttta tctgactgaa aggtctggaa taggtctggc 420

40 tctgggtcca ggaggcttca gggatcagac aatgtcatca ggatctggtc tctctctctc 480

tttgcttggc tttttctcag gcacatatag tgactcaatg gccactgcat ttctaacctc 540

tcatcctccc aggttcaagt ccaatgggaa agaaatatct tccttcaaca tctgaatatg 600

45 ttactggaag tttggagaat cactactaga tggcaaaaac aaaagatgtt ccttccatct 660

tgtgaactgc ataagagatc ttggggggtg ggcgatgaag agaggtatac tgtggtctca 720

50 ctagtcaagg acagctaata gctgtaaaac aggtggcctt ggatacctct aataaattag 780

ctgctgaaaa ggaataccgg aaactacagg aagaagtaga tttgctcaaa gactgaaac 840

atgtcaacat tgtggcctat ttggggacat gcttgcaaga gaacactgtg agcattttc 899

55 atg gag ttt gtt cct ggt ggc tca atc tct agt att ata aac cgt ttt 947

Met Glu Phe Val Pro Gly Gly Ser Ile Ser Ser Ile Ile Asn Arg Phe

1 5 10 15

60 ggg cca ttg cct gag atg gtg ttc tgt aaa tat acg aaa caa ata ctt 995

Gly Pro Leu Pro Glu Met Val Phe Cys Lys Tyr Thr Lys Gln Ile Leu

2/6

	20	25	30	
5	caa ggt gtt gct tat ctc cat gag aac tgt gtg gta cat cgc gat atc Gln Gly Val Ala Tyr Leu His Glu Asn Cys Val Val His Arg Asp Ile 35 40 45	1043		
10	aaa gga aat aat gtt atg ctc atg cca act gga ata ata aag ctg att Lys Gly Asn Asn Val Met Leu Met Pro Thr Gly Ile Ile Lys Leu Ile 50 55 60	1091		
	gac ttt ggc tgt gcc agg cgt ttg gcc tgg gca ggt tta aat ggc acc Asp Phe Gly Cys Ala Arg Arg Leu Ala Trp Ala Gly Leu Asn Gly Thr 65 70 75 80	1139		
15	cac agt gac atg ctt aag tcc atg cat ggg act cca tat tgg atg gcc His Ser Asp Met Leu Lys Ser Met His Gly Thr Pro Tyr Trp Met Ala 85 90 95	1187		
20	cca gaa gtc atc aat gag tct ggc tat gga cgg aaa tca gat atc tgg Pro Glu Val Ile Asn Glu Ser Gly Tyr Gly Arg Lys Ser Asp Ile Trp 100 105 110	1235		
25	agc att ggt tgt act gtg ttt gag atg gct aca ggg aag cct cca ctg Ser Ile Gly Cys Thr Val Phe Glu Met Ala Thr Gly Lys Pro Pro Leu 115 120 125	1283		
30	gct tcc atg gac agg atg gcc gcc atg ttt tac atc gga gca cac cga Ala Ser Met Asp Arg Met Ala Ala Met Phe Tyr Ile Gly Ala His Arg 130 135 140	1331		
	ggg ctg atg cct cct tta cca gac cac ttc tca gaa aat gca gca gac Gly Leu Met Pro Pro Leu Pro Asp His Phe Ser Glu Asn Ala Ala Asp 145 150 155 160	1379		
35	ttt gtg cgc atg tgc ctg acc agg gac cag cat gag cga cct tct gct Phe Val Arg Met Cys Leu Thr Arg Asp Gln His Glu Arg Pro Ser Ala 165 170 175	1427		
40	ctc cag ctc ctg aag cac tcc ttc ttg gag aga agt cac tga Leu Gln Leu Leu Lys His Ser Phe Leu Glu Arg Ser His 180 185 190	1469		
45	atatacatca agactttctt cccagttcca ctgcagatgc tcccttgctt aattgtgggg	1529		
	aatgatggct aagggatctt tgtttcccca ctgaaaattc agtctaacc agtttaagca	1589		
	gacccatagg agtcattaac tgaaagttgc agttacatat tagcctcctc aagtgtcaga	1649		
50	cattattact catagtatca gaaaacatgt tcttaataac aacaaaaaac tatttcagtg	1709		
	tttacagttt tgattgtcca ggaactacat tctctattgt tttatatgac atttcttttt	1769		
	atttttggcc tgtcctgtca actttaatgt tgtagtttta aaataaattg taaaaacaac	1829		
55	ttaaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa	1889		
	aaaaaaaa	1896		
60	<210> 2 <211> 189			

3/6

<212> PRT

<213> Homo sapiens

<400> 2

```

5 Met Glu Phe Val Pro Gly Gly Ser Ile Ser Ser Ile Ile Asn Arg Phe
   1           5           10           15
   Gly Pro Leu Pro Glu Met Val Phe Cys Lys Tyr Thr Lys Gln Ile Leu
           20           25           30
10 Gln Gly Val Ala Tyr Leu His Glu Asn Cys Val Val His Arg Asp Ile
   35           40           45
   Lys Gly Asn Asn Val Met Leu Met Pro Thr Gly Ile Ile Lys Leu Ile
   50           55           60
   Asp Phe Gly Cys Ala Arg Arg Leu Ala Trp Ala Gly Leu Asn Gly Thr
   65           70           75           80
15 His Ser Asp Met Leu Lys Ser Met His Gly Thr Pro Tyr Trp Met Ala
   85           90           95
   Pro Glu Val Ile Asn Glu Ser Gly Tyr Gly Arg Lys Ser Asp Ile Trp
   100          105          110
   Ser Ile Gly Cys Thr Val Phe Glu Met Ala Thr Gly Lys Pro Pro Leu
   115          120          125
20 Ala Ser Met Asp Arg Met Ala Ala Met Phe Tyr Ile Gly Ala His Arg
   130          135          140
   Gly Leu Met Pro Pro Leu Pro Asp His Phe Ser Glu Asn Ala Ala Asp
   145          150          155          160
25 Phe Val Arg Met Cys Leu Thr Arg Asp Gln His Glu Arg Pro Ser Ala
   165          170          175
   Leu Gln Leu Leu Lys His Ser Phe Leu Glu Arg Ser His
   180          185

```

30

<210> 3

<211> 702

<212> DNA

35 <213> Homo sapiens

<220>

<221> CDS

<222> (1)..(702)

40

<400> 3

```

gta tac tgt ggt ctc act agt caa gga cag cta ata gct gta aaa cag 48
Val Tyr Cys Gly Leu Thr Ser Gln Gly Gln Leu Ile Ala Val Lys Gln
   1           5           10           15
45 gtg gct ttg gat acc tct aat aaa tta gct gct gaa aag gaa tac cgg 96
Val Ala Leu Asp Thr Ser Asn Lys Leu Ala Ala Glu Lys Glu Tyr Arg
           20           25           30
50 aaa cta cag gaa gaa gta gat ttg ctc aaa gca ctg aaa cat gtc aac 144
Lys Leu Gln Glu Glu Val Asp Leu Leu Lys Ala Leu Lys His Val Asn
           35           40           45
55 att gtg gcc tat ttg ggg aca tgc ttg caa gag aac act gtg agc att 192
Ile Val Ala Tyr Leu Gly Thr Cys Leu Gln Glu Asn Thr Val Ser Ile
           50           55           60
ttc atg gag ttt gtt cct ggt ggc tca atc tct agt att ata aac cgt 240
Phe Met Glu Phe Val Pro Gly Gly Ser Ile Ser Ser Ile Ile Asn Arg
   65           70           75           80
60

```

4/6

```

      ttt ggg cca ttg cct gag atg gtg ttc tgt aaa tat acg aaa caa ata   288
      Phe Gly Pro Leu Pro Glu Met Val Phe Cys Lys Tyr Thr Lys Gln Ile
                               85                               90                               95

5    ctt caa ggt gtt gct tat ctc cat gag aac tgt gtg gta cat cgc gat   336
      Leu Gln Gly Val Ala Tyr Leu His Glu Asn Cys Val Val His Arg Asp
                               100                               105                               110

      atc aaa gga aat aat gtt atg ctc atg cca act gga ata ata aag ctg   384
10   Ile Lys Gly Asn Asn Val Met Leu Met Pro Thr Gly Ile Ile Lys Leu
                               115                               120                               125

      att gac ttt ggc tgt gcc agg cgt ttg gcc tgg gca ggt tta aat ggc   432
15   Ile Asp Phe Gly Cys Ala Arg Arg Leu Ala Trp Ala Gly Leu Asn Gly
                               130                               135                               140

      acc cac agt gac atg ctt aag tcc atg cat ggg act cca tat tgg atg   480
20   Thr His Ser Asp Met Leu Lys Ser Met His Gly Thr Pro Tyr Trp Met
                               145                               150                               155                               160

      gcc cca gaa gtc atc aat gag tct ggc tat gga cgg aaa tca gat atc   528
      Ala Pro Glu Val Ile Asn Glu Ser Gly Tyr Gly Arg Lys Ser Asp Ile
                               165                               170                               175

25   tgg agc att ggt tgt act gtg ttt gag atg gct aca ggg aag cct cca   576
      Trp Ser Ile Gly Cys Thr Val Phe Glu Met Ala Thr Gly Lys Pro Pro
                               180                               185                               190

30   ctg gct tcc atg gac agg atg gcc gcc atg ttt tac atc gga gca cac   624
      Leu Ala Ser Met Asp Arg Met Ala Ala Met Phe Tyr Ile Gly Ala His
                               195                               200                               205

      cga ggg ctg atg cct cct tta cca gac cac ttc tca gaa aat gca gca   672
35   Arg Gly Leu Met Pro Pro Leu Pro Asp His Phe Ser Glu Asn Ala Ala
                               210                               215                               220

      gac ttt gtg cgc atg tgc ctg acc agg taa                               702
40   Asp Phe Val Arg Met Cys Leu Thr Arg
      225                               230

      <210> 4
      <211> 233
      <212> PRT
45   <213> Homo sapiens

      <400> 4
      Val Tyr Cys Gly Leu Thr Ser Gln Gly Gln Leu Ile Ala Val Lys Gln
           1           5           10           15
50   Val Ala Leu Asp Thr Ser Asn Lys Leu Ala Ala Glu Lys Glu Tyr Arg
           20           25           30
      Lys Leu Gln Glu Glu Val Asp Leu Lys Ala Leu Lys His Val Asn
           35           40           45
      Ile Val Ala Tyr Leu Gly Thr Cys Leu Gln Glu Asn Thr Val Ser Ile
55   50           55           60
      Phe Met Glu Phe Val Pro Gly Gly Ser Ile Ser Ser Ile Ile Asn Arg
           65           70           75           80
      Phe Gly Pro Leu Pro Glu Met Val Phe Cys Lys Tyr Thr Lys Gln Ile
           85           90           95
60   Leu Gln Gly Val Ala Tyr Leu His Glu Asn Cys Val Val His Arg Asp
           100           105           110

```

5/6

	Ile	Lys	Gly	Asn	Asn	Val	Met	Leu	Met	Pro	Thr	Gly	Ile	Ile	Lys	Leu	
			115					120					125				
	Ile	Asp	Phe	Gly	Cys	Ala	Arg	Arg	Leu	Ala	Trp	Ala	Gly	Leu	Asn	Gly	
		130					135					140					
5	Thr	His	Ser	Asp	Met	Leu	Lys	Ser	Met	His	Gly	Thr	Pro	Tyr	Trp	Met	
		145				150					155					160	
	Ala	Pro	Glu	Val	Ile	Asn	Glu	Ser	Gly	Tyr	Gly	Arg	Lys	Ser	Asp	Ile	
				165						170					175		
	Trp	Ser	Ile	Gly	Cys	Thr	Val	Phe	Glu	Met	Ala	Thr	Gly	Lys	Pro	Pro	
10			180						185					190			
	Leu	Ala	Ser	Met	Asp	Arg	Met	Ala	Ala	Met	Phe	Tyr	Ile	Gly	Ala	His	
		195					200						205				
	Arg	Gly	Leu	Met	Pro	Pro	Leu	Pro	Asp	His	Phe	Ser	Glu	Asn	Ala	Ala	
		210					215					220					
15	Asp	Phe	Val	Arg	Met	Cys	Leu	Thr	Arg								
	225					230											

20	<210> 5	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Description of Artificial Sequence:Primer01	
	<400> 5	
	gatcccttag ccatcattcc	20
30		
	<210> 6	
	<211> 22	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
	<223> Description of Artificial Sequence:Primer02	
40	<400> 6	
	taatagctgt aaaacaggtg gc	22
45		
	<210> 7	
	<211> 26	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
50	<223> Description of Artificial Sequence:Primer04	
	<400> 7	
	tgaaggtagg agtcaacgga tttggt	26
55		
	<210> 8	
	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	
60	<220>	

6/6

<223> Description of Artificial Sequence:Primer04

<400> 8

catgtgggcc atgaggtcca ccac

24

5