




(86) Date de dépôt PCT/PCT Filing Date: 1991/10/10
(87) Date publication PCT/PCT Publication Date: 1992/04/30
(45) Date de délivrance/Issue Date: 2001/08/21
(85) Entrée phase nationale/National Entry: 1993/04/08
(86) N° demande PCT/PCT Application No.: US 91/07467
(87) N° publication PCT/PCT Publication No.: WO 92/06974
(30) Priorité/Priority: 1990/10/12 (596,889) US

(51) Cl.Int.⁵/Int.Cl.⁵ C07D 471/10, A61K 31/47
(72) Inventeur/Inventor:
Malamas, Michael S., US
(73) Propriétaire/Owner:
AMERICAN HOME PRODUCTS CORPORATION, US
(74) Agent: RIDOUT & MAYBEE

(54) Titre : ANALOGUES ALKYLIDENIQUES DE 1'-AMINOSPIRO[ISOQUINOLEINE-4(1H),3'-PYRROLIDINE]-1,2',3,5'(2H)-TETRONES, UTILES COMME INHIBITEURS DE L'ALDOSE-REDUCTASE
(54) Title: ALKYLIDENE ANALOGS OF 1'-AMINOSPIRO[ISOQUINOLINE-4(1H),-3'-PYRROLIDINE]1,2',3,5'(2H)-TETRONES USEFUL AS ALDOSE REDUCTASE INHIBITORS



(57) Abrégé/Abstract:

This invention relates to alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrones (I), to processes for their preparation, to methods for using the compounds; and to pharmaceutical preparations thereof. The compounds have pharmaceutical properties which render them beneficial for the prevention or treatment of diabetes mellitus associated complications.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :

C07D 471/10, A61K 31/47

A1

(11) International Publication Number:

WO 92/06974

(43) International Publication Date:

30 April 1992 (30.04.92)

2093793

(21) International Application Number: PCT/US91/07467

(22) International Filing Date: 10 October 1991 (10.10.91)

(30) Priority data: 596,889 2 1 October 1990 (1.10.90) 2 US

(71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; 685 Third Avenue, New York, NY 10017 (US).

(72) Inventor: MALAMAS, Michael, Sotirios ; 2443 Oleander Circle, Jamison, PA 18929 (US).


(74) Agents: ALICE, Ronald, W.; American Home Products Corporation, 685 Third Avenue, New York, NY 10017 (US) et al.

(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent).

Published

With international search report.

(54) Title: ALKYLIDENE ANALOGS OF 1'-AMINOSPIRO[ISOQUINOLINE-4(1H),3'-PYRROLIDINE]-1,2',3,5'(2H)-TETRONES USEFUL AS ALDOSE REDUCTASE INHIBITORS



(57) Abstract

This invention relates to alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone (I), to processes for their preparation, to methods for using the compounds, and to pharmaceutical preparations thereof. The compounds have pharmaceutical properties which render them beneficial for the prevention or treatment of diabetes mellitus associated complications.

ALKYLIDENE ANALOGS OF 1'-AMINOSPIRO[ISOQUINOLINE-
4(1H),3'-PYRROLIDINE]-1,2',3,5'(2H)-TETRONES

Background of the Invention

5 This invention relates to alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrones, to processes for their preparation, to methods for using the compounds, and to pharmaceutical preparations thereof. The compounds have pharmaceutical properties which render them beneficial for the prevention or treatment of complications associated with diabetes mellitus.


10 The use of insulin and/or oral hypoglycemic agents in the treatment of diabetes mellitus has prolonged the life of many of these patients. However, their use has not had a demonstrable impact on the development of diabetic complications such as neuropathy, nephropathy, retinopathy, cataracts and vascular disease which accompany the underlying metabolic disorder. There is little question that chronic hyperglycemia
15 plays a major role in the genesis of these complications, and that complete normalization of blood glucose would likely prevent most if not all complications. For a number of reasons, though, chronic normalization of blood glucose has not been achieved with the currently available therapies.

20 The long-term complications of diabetes develop in tissues where glucose uptake is independent of insulin. In these tissues, which include the lens, retina, kidney and peripheral nerves, the systemic hyperglycemia of diabetes is rapidly transposed into high tissular concentrations of glucose. In all of these tissues this excess glucose is rapidly metabolized by the sorbitol pathway. The intense diabetes-induced flux of glucose through this pathway appears to initiate a cascade of
25 biochemical alterations which slowly progress to cell dysfunction and structural damage. Aldose reductase, the key enzyme in the sorbitol pathway, reduces glucose to sorbitol at the expense of the cofactor NADPH. In animal models of diabetes, compounds which inhibit aldose reductase have been shown to prevent the biochemical, functional and morphological changes induced by hyperglycemia. Early
30 studies by J. H. Kinoshita and collaborators implicated aldose reductase in the etiology of diabetic cataracts. More recent studies have provided compelling evidence that aldose reductase also plays a significant role in the initiation of diabetic nephropathy,

retinopathy and neuropathy (cf McCaleb et al, J. Diab. Comp., 2, 16, 1989; Robison et al, Invest. Ophthalmol. Vis. Sci., 30,, 2285, 1989; Notvest and Inserra, Diabetes, 36, 500, 1987.

Prior Art

- 5 The closest prior art is Malamas U. S. Patent 4,927,831, May 22, 1990, which discloses the spiro-isoquinoline-pyrrolidine tetrones of formula




(R¹ is hydrogen or fluorine)

useful as aldose reductase inhibitors for treating complications of diabetes and galactosemia.

10

Summary of Invention

The alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrones of the present invention are represented by formula (I):





wherein:


5 R^1 and R^2 are independently hydrogen, alkyl containing 1 to 6 carbon atoms, halogen, lower alkoxy containing 1 to 6 carbon atoms, trifluoromethyl, nitro, aryl or aryl (lower alkyl)oxy wherein aryl contains 6 to 10 carbon atoms and lower alkyl contains 1 to 6 carbon atoms;

R^3 is lower alkyl containing 1 to 6 carbon atoms, aryl, aryl (lower alkyl) or dihalogen substituted aryl (lower alkyl) wherein aryl contains 6 to 10 carbon atoms and lower alkyl contains 1 to 6 carbon atoms;

10 R^4 and R^5 are independently hydrogen, alkyl containing 1 to 12 carbon atoms, aryl, aryl (lower alkyl) wherein aryl contains 6 to 10 carbon atoms and lower alkyl contains 1 to 6 carbon atoms, or R^4 and R^5 are joined to form alicyclic or heterocyclic rings selected from the group consisting of




A more preferred group of compounds of the present invention is represented by formula (I) :



wherein:

- 5 R^1 and R^2 are hydrogen or halogen; R^3 is a dihalogen substituted benzyl; R^4 and R^5 are lower alkyl containing 1 to 3 carbon atoms or R^4 and R^5 are joined to form alicyclic or heterocyclic rings selected from the group consisting of



The most preferred compounds of the present invention are set forth below:

- 10 2-[(4-bromo-2-fluorophenyl)methyl]-1'-[(1-methylethylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone;
- 2-[(4-bromo-2-fluorophenyl)methyl]-6-fluoro-1'-[(1-methylethylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone;
- 15 2-[(4-bromo-2-fluorophenyl)methyl]-1'-[(1-ethylpropylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone;
- 2-[(4-bromo-2-fluorophenyl)methyl]-1'-(cyclopentylideneamino)spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone; and

2-[(4-bromo-2-fluorophenyl)methyl]-1'-[(tetrahydro-4H-pyran-4-ylidene)-amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.

5 The compounds of formula (I) all possess at least one asymmetric carbon atom, namely the spiro carbon atom at position 3' of the pyrrolidine ring. The compounds of formula (I) therefore exist, and may be isolated, in two or more stereoisomeric forms. This invention encompasses the compounds of formula (I) in racemic form or in any optically active form.

The alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone can be prepared by the processes described hereinafter.

10 A method is provided for preventing or relieving diabetes mellitus associated complications in a diabetic mammal by administering to said mammal a prophylactic or alleviating amount of the compounds of formula (I). Such complications include neuropathy, nephropathy, retinopathy, keratopathy, diabetic uveitis, cataracts and limited joint mobility.

15 The compounds of formula (I), when admixed with a pharmaceutically acceptable carrier, form a pharmaceutical composition which can be used according to the preceding method.

20 The alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone of this invention may be administered to mammals, for example, man, cattle, or rabbits, either alone or in dosage forms, i.e., capsules or tablets, combined with pharmacologically acceptable excipients.

25 The compounds of this invention may be given orally. However, the method of administering the present active ingredients of this invention is not to be construed as limited to a particular mode of administration. For example, the compounds may be administered topically directly to the eye in the form of drops of sterile, buffered ophthalmic solutions, preferably of pH 7.2-7.6. Also, they may be administered orally in solid form containing such excipients as starch, milk sugar, certain types of clay and so forth. They may also be administered orally in the form of solutions or they may be injected parenterally. For parenteral administration, they may be used in the form of a
30 sterile solution, preferably of pH 7.2-7.6, containing a pharmaceutically acceptable buffer.

2093793

- 6 -

The dosage of the alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone will vary with the form of administration and the particular compound chosen. Furthermore, it will vary with the particular host under treatment. Generally, treatment is initiated with small dosages substantially less than the optimal dose of the compound. Thereafter, the dosage is increased by small increments until efficacy is obtained. In general, the compounds of this invention are most desirably administered at a concentration level that will generally afford effective results without causing any harmful or deleterious side effects. For topical administration, a 0.05-1.0% solution may be administered dropwise in the eye. The frequency of instillation varies with the subject under treatment from a drop every two or three days to once daily. For oral or parenteral administration a preferred level of dosage ranges from about 0.1 mg to about 1.0 mg per kilo of body weight per day, although aforementioned variations will occur. However, a dosage level that is in the range of from about 0.1 mg to about 1.0 mg per kilo of body weight per day is most satisfactory.

Unit dosage forms such as capsules, tablets, pills and the like may contain from about 5.0 mg to about 25.0 mg of the active ingredients of this invention with a pharmaceutical carrier. Thus, for oral administration, capsules can contain from between about 5.0 mg to about 25.0 mg of the active ingredients of this invention with or without a pharmaceutical diluent. Tablets, either effervescent or noneffervescent, can contain between about 5.0 to 25.0 mg of the active ingredients of this invention together with conventional pharmaceutical carriers. Thus tablets, which may be coated and either effervescent or noneffervescent, may be prepared according to the known art. Inert diluents or carriers, for example, magnesium carbonate or lactose, can be used together with conventional disintegrating agents for example, magnesium stearate.

The alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone also can be used in combination with insulin or oral hypoglycemic agents to produce a beneficial effect in the treatment of diabetes mellitus. In this instance, commercially available insulin preparations or oral hypoglycemic agents, exemplified by acetohexamide, chlorpropamide, tolazamide, tolbutamide and phenformin, are suitable. The compounds hereof can be administered sequentially or simultaneously with insulin or the oral hypoglycemic agent. Suitable methods of administration, compositions and doses of the insulin preparation or oral hypoglycemic agent are described in medical textbooks; for instance, *Physicians' Desk Reference*, 42 ed., Medical Economics Co., Oradell, N.J., U.S.A., 1988.

SUBSTITUTE SHEET


The utilization of the compounds in preventing, diminishing and alleviating diabetic complications is demonstrable in experiments using galactosemic rats, see Dvornik *et al.*, Science, 182, 1146 (1973). Such experiments are exemplified
5 hereinbelow after the listing of the following general comments pertaining to these experiments:

(a) Four or more groups of six male rats, 50-70 g, Sprague-Dawley strain, were used. The first group, the control group, was fed a mixture of laboratory chow (rodent Laboratory Chow, Purina) and glucose at 20% (w/w %) concentration. An
10 untreated galactosemic group was fed a similar diet in which galactose was substituted for glucose. The third group was fed a diet prepared by mixing a given amount of the test compound with the galactose containing diet. The concentration of galactose in the diet of the treated groups was the same as that for the untreated galactosemic group.

(b) After four days, the animals were killed by euthanization. Both the lens
15 and sciatic nerve were removed, weighed and stored frozen for polyol determination.


(c) The polyol determination was performed by a modification of the procedure of M. Kraml and L. Cosyns, Clin. Biochem., 2, 373 (1969). Only two minor reagent changes were made: (a) the rinsing mixture was an aqueous 5% (w/v) trichloroacetic acid solution and (b) the stock solution was prepared by dissolving 25
20 mg of dulcitol in 100 mL of an aqueous trichloroacetic acid solution. [N.B.: For each experiment the average value found in the tissue from rats fed the glucose diet was subtracted from the individual values found in the corresponding tissue in galactose-fed rats to obtain the amount of polyol accumulated.] The aldose reductase inhibiting effects of the compounds of formula (I) were also tested by employing an *in vitro*
25 testing procedure similar to that described by S. Hayman and J.H. Kinoshita, J. Biol. Chem., 240, 877 (1965). In the present case the procedure of Hayman and Kinoshita was modified in that the final chromatography step was omitted in the preparation of the enzyme from bovine lens.

The following tabulated results show that the alkylidene analogs of 1'-
30 aminospino[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone of this invention show the property that they are active *in vivo* and diminish the accumulation of dulcitol in the lenses, sciatic nerves and diaphragm of rats fed galactose. The figures under L, N, and D represent the percentage decrease of dulcitol accumulation in the tissues of the lens, sciatic nerve, and diaphragm, respectively, for treated rats as compared to
35 untreated rats.




R ¹	R	% Inhibition of Aldose Reductase		% Lowering Galactitol Accumulation		
		<i>In Vitro</i>	Dose	<i>In Vivo</i>		
		10 ⁻⁵ M	mg/kg/day	% (L)	% (N)	% (D)
H		8	56	53	89	95
			10.8	30	82	86
			0.9	NS	60	66
F		5	27	58	84	97
			1.1	NS	47	85
H		0	51	71	98	93
H		0	48	76	82	95
H		36	56	44	57	89

The invention also includes a method of preparation of a compound having formula I, in which the compound having formula Ia



where R^1 , R^2 , and R^3 are as defined above is reacted with a carbonyl compound
5 having the formula R^4COR^5 , where R^4 and R^5 are as defined above, or a reactive derivative thereof, for instance, an acetal derivative. The method can be carried in conventional manner for the preparation of Schiff bases.

The invention also includes compounds having formula Ia as illustrated and defined above as new substances useful as chemical intermediates. The invention also
10 includes a method of preparation of a compound having formula Ia, as illustrated and defined above, in which a dicarboxylic acid having the formula Ib



where R^1 , R^2 and R^3 are as defined above, or a reactive derivative thereof is converted into an imide by reaction with hydrazine. This method is preferably carried out by
15 reacting a mono (lower alkyl) ester of the dicarboxylic acid with a reagent that activates the carboxyl group of the monoester and reacting the activated compound with hydrazine.


The aforesaid methods of preparation are illustrated by the following process.

2093793

- 10 -

The Process:

The alkylidene analogs of 1'-aminospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'-(2H)-tetrones of the present invention were prepared by the following reaction scheme:



wherein:

- 10 R¹ is halogen or hydrogen, R³ is disubstituted aralkyl and R⁴, R⁵ are alkyl, hydrogen or joined to form alicyclic and heterocyclic rings.

SUBSTITUTE SHEET

Step a) Reacting either 2-bromobenzoic acid or 2-chlorobenzoic acid of formula (III) wherein R^1 is as defined above with dimethyl malonate and NaH in the presence of a catalytic amount of CuBr to produce the propanedioic acid dimethyl ester of formula (IV) wherein R^1 is as defined above.

5 The 2-bromobenzoic acids or 2-chlorobenzoic acids of formula (III) required for the present invention are commercially available compounds or can be prepared by known methods.

Step b) The propanedioic acid dimethyl ester of formula (IV) can be reacted with thionyl chloride under refluxing conditions to produce the corresponding acid chloride
10 which upon treatment with Et_3N in a conventional solvent which does not adversely influence the reaction, for example, tetrahydrofuran, can produce the compound of formula (V), wherein R^1 is as defined above.

Step c) The compound of formula (V), wherein R^1 is as defined above, is reacted with R^3-NH_2 in the presence of Et_3N in a conventional solvent which does not
15 adversely influence the reaction, for example, DMF, produces the compound of the formula (VI), wherein R^1 and R^3 are as defined above.

Step d) The compound of formula (VI), wherein R^1 and R^3 are as defined above, is reacted with an inorganic base such as potassium carbonate in a conventional solvent which does not adversely influence the reaction, for example, N,N-dimethylformamide
20 and subsequent addition of the tert-butyl bromoacetate produces the compound of formula (VII), wherein R^1 and R^3 are as defined above.


Step e) The compound of formula (VII), wherein R^1 and R^3 are as defined above, can be reacted with an organic acid such as trifluoroacetic acid in a conventional solvent which does not adversely influence the reaction, for example, methylene
25 chloride, to produce the compound of formula (VIII), wherein R^1 and R^3 are as defined above.

Step f) The compound of formula (VIII), wherein R^1 and R^3 are as defined above, can be reacted with a coupling agent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (DCC)/1-hydroxybenzotriazole (HOBt) in a conventional solvent
30 which does not adversely influence the reaction, for example, N, N-dimethylformamide, and subsequent addition of hydrazine and Et_3N , produces the compound of formula (IX), wherein R^1 and R^3 are as defined above.

2093793

- 12 -

Step g) The compound of formula (IX), wherein R^1 and R^3 are as defined above

can be reacted with R^4  R^5 in the presence of a catalytic amount of any acid, for example, 10-camphorsulfonic acid, to produce the compound of formula (X), wherein R^1 , R^3 , R^4 and R^5 are as defined above.

5 The following examples further illustrate this invention:

EXAMPLE 1

**2-[(4-Bromo-2-fluorophenyl)methyl]-1'-[(1-ethylpropylidene)amino]-
spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone**

Step a) (2-Carboxyphenyl)propanedioic Acid Dimethyl Ester

10 To a rapidly stirred cold suspension (0°C) of 2-bromobenzoic acid (30.0 g, 149.32 mmol), cuprous bromide (2.14 g, 14.93 mmol) and dimethyl malonate (300 mL) was added NaH (80% in mineral oil, 10.75 g, 358.37 mmol) over a 30 minute period, while a stream of dry N_2 was passed over the mixture. After the addition of the NaH had been completed, the mixture was stirred for 10 minutes at room temperature
15 and 30 minutes at 70°C (external oil bath temperature). At this point, the suspension had turned to a solid mass, which was dissolved in H_2O (1000 mL). The aqueous layer was extracted with diethyl ether (3 x 500 mL) and was acidified with HCl (2N). The mixture was extracted with EtOAc and dried over $MgSO_4$. Evaporation gave an off-white solid which was recrystallized from Et_2O /hexane (after cooling to -20°C) to
20 give a white solid (34.2 g, 90.9%, m.p. 119-120°C).

^1H NMR ($DMSO-d_6$, 400 MHz): δ 3.67 [s, 6H, $-CH(CO_2CH_3)_2$], 5.72 [s, 1H, $-CH(CO_2CH_3)_2$], 7.3 (d, $J = 7.76$ Hz, 1H, Ar-H), 7.45 (dt, $J = 7.66$ Hz, 1.12 Hz, 1H, Ar-H), 7.6 (dt, $J = 7.66$ Hz, 1.45 Hz, 1H, Ar-H), 7.94 (dd, $J = 7.8$ Hz, 1.33 Hz, 1H, Ar-H), 13.2 (s, 1H, $-CO_2H$)

25 IR (KBr, cm^{-1}): 3300-2700 (CO_2H), 1750 (CO), 1730 (CO), 1680 (CO)

MS (m/e): 252 (M^+), 220 ($M^+ - CH_3OH$), 188 ($M^+ - 2xCH_3OH$)

Anal. Calcd.: C, 57.14; H, 4.80

Found: C, 57.05; H, 4.78.

SUBSTITUTE SHEET

The following compounds were prepared in substantially the same manner as that of Example 1, Step a):

(2-Carboxy-6-fluorophenyl)propanedioic Acid Dimethyl Ester

¹H NMR (DMSO-d₆, 400 MHz): δ 3.68 [s, 6H, (-CO₂Me)₂], 5.79 [s, 1H, Ar-
5 CH(CO₂Me)₂], 7.12 (dd, J = 10.06 Hz, 2.61 Hz, 1H, Ar-H), 7.33 (dt, J = 8.48 Hz, 2.64 Hz, 1H, Ar-H), 8.03 (dd, 8.77 Hz, 6.17 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 3400-2700 (CO₂H), 1730 (CO), 1680 (CO)

MS (m/e): 270 (M⁺), 238 (M⁺-CH₃OH), 210 (M⁺-CH₃OH, -CO), 151 (M⁺-CH₃OH, -CO, -CO₂CH₃)

10 Anal. Calcd.: C, 53.34; H, 4.10

Found: C, 53.36; H, 3.93

m.p. 121.5-123.0°C.

(2-Carboxy-6-chlorophenyl)propanedioic Acid Dimethyl Ester

¹H NMR (DMSO-d₆, 200 MHz): δ 3.69 [s, 6H, (-CO₂Me)₂], 5.78 [s, 1H, Ar-
15 CH(CO₂Me)₂], 7.38 (d, J = 1.8 Hz, 1H, Ar-H), 7.58 (dd, J = 7.8 Hz, 1.8 Hz, 1H, Ar-H), 7.96 (d, J = 8.2 Hz, 1H, Ar-H), 13.5 (br s, 1H, -CO₂H)

IR (KBr, cm⁻¹): 3200-2700 (CO₂H), 1760 (CO), 1740 (CO), 1690 (CO)

MS (m/e): 286 (20 M⁺), 254 (64, M⁺-CH₃OH), 222 (60, M⁺-2 x CH₃OH)

Anal. Calcd.: C, 50.28; H, 3.87

20 Found: C, 50.40; H, 3.87

m.p. 125-127°C.

(2-Carboxy-6-bromophenyl)propanedioic Acid Dimethyl Ester

¹H NMR (DMSO-d₆, 400 MHz): δ 3.68 [s, 6H, -(CO₂CH₃)₂], 5.74 (s, 1H, Ar-CH-),
7.5 (d, J = 2.02 Hz, 1H, Ar-H), 7.70 (dd, J = 8.4 Hz, 1.98 Hz, 1H, Ar-H), 7.87 (d,
25 J = 8.41 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 3400-2300 (CO₂H), 1745 (CO), 1720 (CO), 1695 (CO)

MS (m/e): 330 (M⁺), 298 (M⁺-CH₃OH)

Anal. Calcd.: C, 43.53; H, 3.35

Found: C, 43.56; H, 3.23

30 m.p. 127-128°C.

2093793

- 14 -

Step b) 3-Methoxy-1-oxo-1H-2-benzopyran-4-carboxylic Acid Methyl Ester

A mixture of (2-carboxyphenyl)propanedioic acid dimethyl ester (10.0 g, 39.68 mmol) and SOCl₂ (100 g) was refluxed for 2 hours. The volatiles were removed *in vacuo* and the crude product (acid chloride) was dissolved in THF (200 mL). Triethylamine (27.64 mL, 198.4 mmol) was added and the mixture was stirred for 30 minutes. The yellowish suspension was poured into HCl (1N, 1000 mL), extracted with EtOAc and the organic extracts were dried over MgSO₄. Evaporation and crystallization from acetone/ether/hexane (after cooling to -20°C) gave a white solid (87.6 g, 94.4%, m.p. 129-130°C).

¹H NMR (DMSO-d₆, 400 MHz): δ 3.82, (s, 3H, -CO₂Me), 4.03 (s, 3H, -OMe), 7.42 (t, J = 7.26 Hz, 1H, Ar-H), 7.8 (t, J = 8.2 Hz, 1H, Ar-H), 7.9 (d, J = 8.3 Hz, 1H, Ar-H), 8.1 (d, J = 7.26 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 1740 (CO), 1685 (CO)

MS (m/e): 234 (15, M⁺), 206 (38.5, M⁺-CO), 203 (12, M⁺-OMe)

Anal. Calcd.: C, 61.59; H, 4.30

Found: C, 61.82; H, 4.29.

The following compounds were prepared in substantially the same manner as that of Example 1, Step b):

20 6-Fluoro-3-methoxy-1-oxo-1H-2-benzopyran-4-carboxylic Acid Methyl Ester

¹H NMR (DMSO-d₆, 400 MHz): δ 3.81 (s, 3H, -CO₂CH₃), 4.06 (s, 3H, -OCH₃-), 7.27 (dt, J = 8.3 Hz, 1H, Ar-H), 7.8 (dd, J = 11.83 Hz, 2.49 Hz, 1H, Ar-H), 8.16 (dd, J = 8.92 Hz, 6.2 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 1750 (CO), 1685 (CO)

MS (m/e): 252 (24, M⁺), 224 (54, M⁺-CO)

Anal. Calcd.: C, 57.15; H, 3.60

Found: C, 57.19; H, 3.57

m.p. 142-143°C.

SUBSTITUTE SHEET

6-Chloro-3-methoxy-1-oxo-1H-2-benzopyran-4-carboxylic Acid Methyl Ester

^1H NMR (DMSO- D_6 , 400 MHz): δ 3.81 (s, 3H, $-\text{CO}_2\text{CH}_3$), 4.05 (s, 3H, $-\text{OCH}_3$), 7.44 (dd, $J = 8.56$ Hz, 1.99 Hz, 1H, Ar-H), 8.06 (m, 2H, Ar-H)

5 IR (KBr, cm^{-1}): 1750 (CO), 1690 (CO)

MS (m/e): 268 (34, M^+), 240 (86, $\text{M}^+ - \text{CO}$)

Anal. Calcd.: C, 53.65; H, 3.38

Found: C, 53.59; H, 3.35

m.p. 194-195°C.

10 **6-Bromo-3-methoxy-1-oxo-1H-2-benzopyran-4-carboxylic Acid Methyl Ester**

^1H NMR (DMSO- d_6 , 400 MHz): δ 3.81 (s, 3H, $-\text{CO}_2\text{CH}_3$), 4.05 (s, 3H, $-\text{OCH}_2$), 7.6 (dd, $J = 8.38$ Hz, 1.77 Hz, 1H, Ar-H), 8.0 (d, $J = 8.39$ Hz, 1H, Ar-H), 8.23 (d, $J = 1.95$ Hz, 1H, Ar-H)

15 IR (KBr, cm^{-1}): 1740 (CO), 1680 (CO)

MS (m/e): 312 (17 M^+), 284 (45, $\text{M}^+ - \text{CO}$)

Anal. Calcd.: C, 46.03; H, 2.90

Found: C, 46.12; H, 2.62

m.p. 200-201°C.

20 **Step c) 2-[(4-Bromo-2-fluorophenyl)methyl]-1,2,3,4-tetrahydro-1,3-dioxo-4-isoquinolinecarboxylic Acid Methyl Ester**

To a solution of 3-methoxy-1-oxo-1H-2-benzopyran-4-carboxylic acid methyl ester (5.0 g, 21.37 mmol) in DMF (100 mL) were added 4-bromo-2-fluorobenzylamine (4.36 g, 21.37 mmol) and Et_3N (5.96 mL, 42.74 mmol). The mixture was stirred at 25 80°C for 30 minutes, poured into H_2O (1500 mL), acidified with HCl (2N) and extracted with EtOAc. The organic extracts were dried over MgSO_4 . Evaporation and crystallization from acetone/hexane (after cooling to -20°C) gave a white solid (7.6 g, 87.7 %, m.p. 149-150°C).

2093793

- 16 -

^1H NMR (DMSO- d_6 , 400 MHz): δ [3.67 (s), 4.0 (s), 3H, $-\text{CO}_2\text{Me}$, tautomeric], [5.06 (q), $J = 15.4$ Hz, 5.30 (s), 2H, $-\text{NCH}_2-$, tautomeric], 5.4 (s), 1H, $\text{CH}-\text{CO}_2\text{Me}$, tautomeric], 7.07-8.43 (m, 7H, Ar-H, tautomeric)

IR (KBr, cm^{-1}): 1670 (CO), 1605 (CO)

5 MS (+ FAB): 406 (80, $\text{M}^+ + \text{H}$), 374 (40, $\text{M}^+ - \text{OCH}_3$)

Anal. Calcd.: C, 53.22; H, 3.23; N, 3.45

Found: C, 53.19; H, 2.98; N, 3.40.

The following compounds were prepared in substantially the same manner as that of Example 1, Step c):

10 **2-[(4-Bromo-2-fluorophenyl)methyl]-6-fluoro-1,2,3,4-tetrahydro-1,3-dioxo-4-isoquinolinecarboxylic Acid Methyl Ester**

^1H NMR (DMSO- d_6 , 400 MHz): δ 3.98 (s, 3H, $-\text{CO}_2\text{CH}_3$), 5.27 (s, 2H, $-\text{NCH}_2-$), 7.08 (t, $J = 7.95$ Hz, 2H, Ar-H), 7.2 (m, 1H, Ar-H), 7.34 (m, 2H, Ar-H, -OH), 7.54 (m, 1H, Ar-H), 8.1-8.26 (m, 2H, Ar-H)

15 IR (KBr, cm^{-1}): 1680 (CO), 1660 (CO), 1610 (CO)

MS (m/e): 423 (M^+), 391 ($\text{M}^+ - \text{CH}_3\text{OH}$)

Anal. Calcd.: C, 50.97; H, 2.85; N, 3.30

Found: C, 50.86; H, 2.86; N, 3.33

m.p. 157-158°C.

20 **6-Chloro-1,2,3,4-tetrahydro-2-methyl-1,3-dioxo-4-isoquinoline-carboxylic Acid Methyl Ester**

^1H NMR](DMSO- d_6 , 200 MHz): δ [3.23 (s), 3.44 (s), tautomeric, 3H, $-\text{NCH}_3$], [3.71 (s), 4.03 (s), tautomeric, 3H, $-\text{CO}_2\text{CH}_3$], 7.3-8.4 (tautomeric, Ar-H, -OH, 4H)

IR (KBr, cm^{-1}): 3440 (OH), 1680 (CO), 1600 (CO)

25 MS (m/e): 267 (M^+), 235 ($\text{M}^+ - \text{OMe}$)

Anal. Calcd.: C, 53.85; H, 3.77; N, 5.23

Found: C, 53.66; H, 3.63; N, 5.14

m.p. 166-167°C.

6-Bromo-1,2,3,4-tetrahydro-2-methyl-1,3-dioxo-4-isoquinoline-carboxylic Acid Methyl Ester

¹H NMR (DMSO-d₆, 400 MHz): δ [3.2 (s), 3.42 (s), 3H, tautomeric, N-CH₃], [3.7 (s), 4.01 (s), 3H, tautomeric, -CO₂CH₃], [5.33 (s), 1H, tautomeric, Ar-CH-], [7.5 (dd), 7.8 (dd), tautomeric, 1H, Ar-H], [8.0 (d), 8.08 (d), tautomeric, 1H, Ar-H], [8.51 (d), 7.63 (d), tautomeric, 1H, Ar-H]

IR (KBr, cm⁻¹): 1665 (CO), 605 (CO)

MS (m/e): 311 (M⁺)

Anal. Calcd.: C, 46.18; H, 3.23; N, 4.49

10 Found: C, 45.83; H, 2.77; N, 4.38

m.p. 190-191°C.

Step d) 2-[(4-Bromo-2-fluorophenyl)methyl]-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-1,3-dioxo-4-isoquinolineacetic Acid 1,1-Dimethylethyl Ester

15 To a suspension of 2-[(4-bromo-2-fluorophenyl)methyl]-1,2,3,4-tetrahydro-1,3-dioxo-4-isoquinolinecarboxylic acid methyl ester (4.79 g, 11.58 mmol), K₂CO₃ (3.19 g, 23.16 mmol) in DMF (100 mL) was added tert-butyl bromoacetate (2.81 mL, 17.37 mmol). After stirring at 75°C for 1 hour, the mixture was poured into H₂O, extracted with EtOAc and dried over MgSO₄. Evaporation and purification by flash
20 chromatography (hexane/EtOAc 4/1) gave a clear oil (5.69 g, 94.5%).

¹H NMR (DMSO-d₆, 400 MHz): δ 1.04 [s, 9H, -C(CH₃)₃], 3.53 s, 3H, -CO₂CH₃), 3.60 [dd, J = 17.7 Hz, 2H, -CH₂CO₂(CH₃)₃], 5.14 (s, 2H, NCH₂-), 7.17 (t, J = 8.25 Hz, 1H, Ar-H), 7.36 (dd, J = 8.36 Hz, 1.75 Hz, 1H, Ar-H), 7.6 (m, 3H, Ar-H), 7.77 (dt, J = 7.2 Hz, 1.27 Hz, 1H, Ar-H), 8.19 (dd, J = 8.25 Hz, 1.54 Hz, 1H, Ar-
25 H)

IR (CHCl₃, cm⁻¹): 1720 (CO), 1675 (CO)

MS (m/e): 520 (M + H)⁺, 464 [M⁺-C(CH₃)₃].

2093793

- 18 -

The following compounds were prepared in substantially the same manner as that of Example 1, Step d):

2-[(4-Bromo-2-fluorophenyl)methyl]-6-fluoro-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-1,3-dioxo-4-isoquinolineacetic Acid 1,1-Dimethylethyl Ester

¹H NMR (DMSO-d₆, 200 MHz): δ 1.10 (s, 9H, -CMe₃), 3.55 (s, 3H, -CO₂CH₃), 3.62 (d, J = 17.5 Hz, 1H, -CH₂CO₂CMe₃), 3.75 (d, J = 17.5 Hz, 1H, -CH₂CO₂CMe₃), 5.15 (s, 2H, -NCH₂-), 7.15 (t, J = 8.2 Hz, 1H, Ar-H), 7.35 (d, J = 8.2 Hz, 1H, Ar-H), 7.45-7.70 (m, 3H, Ar-H), 8.38 (dd, J = 8.16 Hz, 5.70 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 1750 (CO), 1720 (CO), 1675 (CO)

MS (m/e): 538 (M + H)⁺, 481 (M⁺ + H-CMe₃)

Anal. Calcd.: C, 53.55; H, 4.12; N, 2.60

Found: C, 53.49; H, 4.00; N, 2.63.

6-Chloro-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-2-methyl-1,3-dioxo-4-isoquinolineacetic Acid 1,1-Dimethylethyl Ester

¹H NMR (DMSO-d₆, 200 MHz): δ 1.06 (s, 9H, -CO₂CMe₃), 3.3 (s, 3H, -NCH₃), 3.6(s,3H, -CO₂CH₃), 3.67 (q, J = 17.5 Hz, 2H, -CH₂CO₂CMe₃), 7.68 (dd, J = 9.0 Hz, 1.6 Hz, 1H, Ar-H), 7.77 (d, J = 2.0 Hz, 1H, Ar-H), 8.21 (d, J = 8.2 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 1740 (CO), 1720 (CO), 1680 (CO)

MS (m/e): 381 (M⁺)

Anal. Calcd.: C, 56.82; H, 5.28; N, 3.67

Found: C, 57.00; H, 5.41; N, 3.66

m.p. 135-136°C.

6-Bromo-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-2-methyl-1,3-dioxo-4-isoquinolineacetic Acid 1,1-Dimethylethyl Ester

¹H NMR (DMSO-d₆, 200 MHz): δ 1.05[s, 9H, -C(CH₃)₃], 3.28 (s, 3H, -NCH₃), 3.59 (s,3H, -CO₂CH₃), 3.58 (d, J = 17.03 Hz, 1H, -CH₂CO₂-), 3.67 (d, J = 17.03

SUBSTITUTE SHEET

Hz, 1H, $-\underline{\text{CH}_2}\text{CO}_2-$), 7.81 (dd, $J = 8.4$ Hz, 1.85 Hz, 1H, Ar-H), 7.88 (d, $J = 1.81$ Hz, 1H, Ar-H), 8.08 (d, $J = 8.4$ Hz, 1H, Ar-H)

IR (KBr, cm^{-1}): 1740 (CO), 1710 (CO), 1670 (CO)

MS (m/e): 425 (M^+), 370 ($M^+ - \text{C}_4\text{H}_7$), 352 ($M^+ - \text{C}_4\text{H}_9\text{O}$)

5 Anal. Calcd.: AC, 50.72; H, 4.73; N, 3.29

 Found: C, 50.47; H, 4.68; N, 3.12

m.p. 152-153°C.

Step e) 2-[(4-Bromo-2-fluorophenyl)methyl]-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-1,3-dioxo-4-isoquinolineacetic Acid

10 A mixture of 2-[(4-bromo-2-fluorophenyl)methyl]-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-1,3-dioxo-4-isoquinolineacetic acid 1,1-dimethylethyl ester (5.19 g, 9.81 mmol), CH_2Cl_2 (100 mL) and $\text{CF}_3\text{CO}_2\text{H}$ (20 mL) was stirred at room temperature for 5 hours. The volatiles were removed in vacuo and the residue was purified by flash chromatography on acid washed silica gel (5% H_3PO_4 in MeOH), to
15 give a white solid (4.12 g, 90.5%, m.p. 139-140°C).

^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 3.54 (s, 3H, $\text{CO}_2\underline{\text{CH}_3}$), 3.64 (q, $J = 17.67$ Hz, 2H, $\underline{\text{CH}_2}\text{CO}_2\text{H}$), 5.12 (q, $J = 15.34$ Hz, 2H, $-\text{N}\underline{\text{CH}_2}-$), 7.14 (t, $J = 8.22$ Hz, 1H, Ar-H), 7.3 (d, $J = 8.3$ Hz, 1H, Ar-H), 7.5-7.6 (m, 3H, Ar-H), 7.76 (d, $J = 7.4$ Hz, 1H, Ar-H), 8.16 (d, $J = 7.8$ Hz, 1H, Ar-H), 12.35 (s, 1H, $-\text{CO}_2\underline{\text{H}}$)

20 IR (KBr, cm^{-1}): 3280 (OH), 3200-2700 (CO_2H), 1750 (CO), 1720 (CO), 1675 (CO)

MS (m/e): 463 (M^+), 445 ($M^+ - \text{H}$, $-\text{OH}$)

Anal. Calcd.: C, 51.28; H, 3.30; N, 2.99

 Found: C, 51.26; H, 3.48; N, 2.95.

25 The following compound was obtained in substantially the same manner as that of Example 1, Step e):

2-[(4-Bromo-2-fluorophenyl)methyl]-6-fluoro-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-1,3-dioxo-4-isoquinolineacetic Acid

^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 3.56 (s, 3H, $-\text{CO}_2\underline{\text{CH}_3}$), 3.6 (d, $J = 17.9$ Hz, 1H, $-\underline{\text{CH}_2}\text{CO}_2\text{H}$), 3.8 (d, $J = 17.9$ Hz, 1H, $-\underline{\text{CH}_2}\text{CO}_2\text{H}$), 5.1 (dd, $J = 15.5$ Hz, 2H, $-\text{N}\underline{\text{CH}_2}-$), 7.12 (s, $J = 8.23$ Hz, 1H, Ar-H), 7.31 (dd, $J = 8.28$ Hz, 1.68 Hz, 1H, Ar-H), 7.45 (dt, $J = 8.56$ Hz, 2.5 Hz, 1H, Ar-H), 7.54 (dd, $J = 9.77$ Hz, 1.89 Hz, 1H, Ar-H),
30 7.76 (d, $J = 7.4$ Hz, 1H, Ar-H), 8.16 (d, $J = 7.8$ Hz, 1H, Ar-H), 12.35 (s, 1H, $-\text{CO}_2\underline{\text{H}}$)

2093793

- 20 -

Ar-H), 7.64 (dd, J = 9.61 Hz, 2.46 Hz, 1H, Ar-H), 8.23 (dd, J = 8.79 Hz, 5.81 Hz, 1H, Ar-H), 12.67 (br s, 1H, -CO₂H)

IR (KBr, cm⁻¹): 3400-2700 (CO₂H), 1745 (CO), 1710 (CO), 1670 (CO)

MS (m/e): 481 (M⁺), 405 (M⁺-CO₂, -CH₃OH)

5 Anal. Calcd.: C, 49.81; H, 2.93; N, 2.90

Found: C, 49.94; H, 3.03; N, 2.84

m.p. 132-133.5°C.

6-Chloro-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-2-methyl-1,3-dioxo-4-isoquinolineacetic Acid

10 ¹H NMR (DMSO-d₆, 200 MHz): δ 3.27 (s, 3H, -CH₃), 3.59 (s, 3H, -CO₂CH₃), 3.64 (q, J = 17.5 Hz, 2H, -CH₂CO₂H), 7.65 (dd, J = 8.6 Hz, 2.0 Hz, 1H, Ar-H), 7.78 (d, J = 2.0 Hz, 1H, Ar-H), 8.18 (d, J = 8.0 Hz, 1H, Ar-H)

IR (KBr, cm⁻¹): 3440 (OH), 3200-2700 (CO₂H), 1750 (CO), 1710 (CO), 1675 (CO)

MS (m/e): 325 (M⁺)

15 Anal. Calcd.: C, 51.63; H, 3.71; N, 4.30

Found: C, 51.73; H, 2.70; N, 4.28

m.p. 195-196°C.

6-Bromo-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-2-methyl-1,3-dioxo-4-isoquinolineacetic Acid

20 ¹H NMR (DMSO-d₆, 200 MHz): δ 3.26 (s, 3H, N-CH₃), 3.53 (d, J = 17.2 Hz, 1H, -CH₂H), 3.58 (s, 3H, -CO₂CH₃), 3.74 (d, J = 17.2 Hz, 1H, -CH₂CO₂H), 7.77 (dd, J = 8.2 Hz, 2.2 Hz, 1H, Ar-H), 7.87 (d, J = 2.2 Hz, 1H, Ar-H), 8.0 (d, J = 8.2 Hz, 1H, Ar-H), 12.64 (s, 1H, -CO₂H)

IR (KBr, cm⁻¹): 3450-2600 (CO₂H), 1735 (CO), 1700 (CO), 1660 (CO)

25 MS (m/e): 369 (M⁺), 324 (M⁺-CO₂H)

Anal. Calcd.: C, 45.43; H, 3.27; N, 3.78

Found: C, 45.04; H, 3.16; N, 3.62

m.p. 194-195°C.

SUBSTITUTE SHEET

Step f) 1'-Amino-2-[(4-bromo-2-fluorophenyl)methyl]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone

To a solution of 2-[(4-bromo-2-fluorophenyl)methyl]-1,2,3,4-tetrahydro-4-(methoxycarbonyl)-1,3-dioxo-4-isoquinolineacetic acid (2.5 g, 5.39 mmol) in DMF
5 (60 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (DCC', 1.34 g, 7.0 mmol) and 1-hydroxybenzotriazole hydrate (HOBT, 1.09 g, 8.08 mmol). After stirring for 2 hours, anhydrous hydrazine (0.22 mL, 7.0 mmol) was added dropwise, followed by Et₃N (1.5 mL, 10.77 mmol) addition. The mixture was stirred for 30 minutes, poured into H₂O, neutralized with HCl (2N) and extracted with
10 EtOAc. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography (hexane/EtOAc 1:1) and subsequent crystallization from ether/hexane (after cooling to -20°C) gave a white solid (1.68 g, 70.0%, m.p. 95-97°C).

¹H NMR (DMSO-d₆, 400 MHz): δ 3.33 (d, J = 18.2 Hz, 1H, -HCHCO-), 3.51 (d, J = 18.2 Hz, 1H, -HCCHCO-), 5.07 (s, 2H, -NCH₂-), 5.23 (s, 2H, -NH₂), 7.17 (t, J = 8.3 Hz, 1H, Ar-H), 7.33 (dd, J = 8.3 Hz, 1.7 Hz, 1H, Ar-H), 7.54 (m, 2H, Ar-H), 7.63 (t, J = 8.51 Hz, 1H, Ar-H), 7.79 (dt, J = 8.7 Hz, 1.25 Hz, 1H, Ar-H) 8.18 (dd, J = 7.7 Hz, 1.25 Hz, 1H, Ar-H)

IR (KBr cm⁻¹): 3340 (NH), 1720 (C=O), 1670 (C=O)

20 MS (m/e): 445(4, M⁺)

Anal. Calcd.: C, 51.14; H, 2.94; N, 9.42

Found: C, 51.04; H, 2.94; N, 9.30.

The following compound was prepared in substantially the same manner as that of Example 1, Step f)

25 1'-Amino-2-[(4-bromo-2-fluorophenyl)methyl]-6-fluorospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone

¹H NMR (DMSO-d₆, 400 MHz): δ 3.42 (q, J = 18.2 Hz, 2H, -CH₂CO-), 5.05 (s, 2H, -NCH₂-), 5.18 (s, 2H, -NH₂), 7.15 (t, J = 8.3 Hz, 1H, Ar-H), 7.33 (dd, J = 8.1 Hz, 1.66 Hz, 1H, Ar-H), 7.49 (dt, J = 8.5 Hz, 2.29 Hz, 1H, Ar-H), 7.55 (dd, J =
30 9.96 Hz, 1.87 Hz, 1H, Ar-H), 7.6 (dd, J = 9.75 Hz, 2.49 Hz, 1H, Ar-H), 8.24 (dd, J = 8.9 Hz, 5.8 Hz, 1H, Ar-H)

2093793

- 22 -

IR (KBr, cm^{-1}): 3350 (NH), 3280 (NH), 1730 (C=O), 1710 (C=O), 1670 (C=O)

MS (m/e): 463 (94, M^+)

Anal. Calcd.: C, 49.16; H, 2.61; N, 9.05

Found: C, 49.19; H, 2.66; N, 8.96

5 m.p. 232-234°C.

Step g) 2-[(4-Bromo-2-fluorophenyl)methyl]-1'-[(1-ethylpropylidene)-amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone

A mixture of 1'-amino-2-[(4-bromo-2-fluorophenyl)methyl]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone (2.0 g, 4.48 mmol), 3-pentanone (20 mL) and 10-camphorsulfonic acid (5 mg) was stirred at 75°C for 20 minutes. Next, the volatiles were removed in vacuo and the residue was purified by flash chromatography on silica gel (hexane/EtOAc 2:1) to give a white solid (1.69 g, 73.7%, m.p. 144-146°C).

^1H NMR (DMSO- d_6 , 400 MHz): δ 0.83 (t, $J = 7.47$ Hz, 3H, $-\text{CH}_2\text{CH}_3$), 1.08 (t, $J = 7.26$ Hz, 3H, $-\text{CH}_2\text{CH}_3$), 2.0 (q, $J = 7.47$ Hz, 2H, $-\text{CH}_2\text{CH}_3$), 2.5 (q, $J = 7.26$ Hz, 2H, $-\text{CH}_2\text{CH}_3$), 3.59 (s, 2H, $-\text{CH}_2\text{CO}-$), 5.06 (q, $J = 15.15$ Hz, 2H, $-\text{NCH}_2-$), 7.15 (t, $J = 8.1$ Hz, 1H, Ar-H), 7.3 (dd, $J = 8.1$ Hz, 1.66 Hz, 1H, Ar-H), 7.53 (dd, $J = 9.96$ Hz, 1.87 Hz, 1H, Ar-H), 7.65 (dt, $J = 8.3$ Hz, 2.07 Hz, 1H, Ar-H), 7.8 (m, 2H, Ar-H), 8.2 (dd, $J = 8.3$ Hz, 1.25 Hz, 1H, Ar-H)

20 IR (KBr cm^{-1}): 1700 (C=O), 1670 (C=O)

MS (m/e): 513 (11, M^+), 387 [28, $\text{M}^+ - \text{CONN} = \text{C}(\text{CH}_2\text{CH}_3)_2$]

Anal. Calcd.: C, 56.04; H, 4.12; N, 8.17

Found: C, 56.44; H, 4.10; N, 8.06.

The following compounds were prepared in substantially the same manner as that of Example 1, Step g):

2-[(4-Bromo-2-fluorophenyl)methyl]-1'-[(1-methylethylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'-(2H)-tetrone

^1H NMR (DMSO- d_6 , 400 MHz): δ 1.72 (s, 3H, $-\text{CH}_3$), 2.14 (s, 3H, $-\text{CH}_3$), 3.57 (s, 2H, $-\text{CH}_2\text{CO}-$), 5.07 (dd, $J = 15.4$ Hz, 2H, $-\text{NCH}_2-$), 7.14 (t, $J = 8.24$ Hz, 1H, Ar-H), 7.3 (dd, $J = 8.24$ Hz, 1.5 Hz, 1H, Ar-H), 7.52 (dd, $J = 9.81$ Hz, 1.9 Hz, 1H, Ar-H), 7.63 (m, 1H, Ar-H), 7.81 (m, 2H, Ar-H), 8.17 (d, $J = 7.63$ Hz, 1H, Ar-H)

SUBSTITUTE SHEET

IR (KBr, cm^{-1}): 1710 (C=O), 1670 (C=O)

MS (m/e): 485 (78 M^+), 387 [100, M^+ -CONNC(CH₃CH₃)₂]

Anal. Calcd.: C, 54.34; H, 3.52; N, 8.64

Found: C, 54.35; H, 3.62; N, 8.34

5 m.p. 127-129°C.

2-[(4-Bromo-2-fluorophenyl)methyl]-6-fluoro-1'-[(1-methylethylidene)-amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone

¹H NMR (DMSO-d₆, 400 MHz): δ 1.71 (s, 3H, -CH₃), 2.15 (s, 3H, -CH₃), 3.55 (dd, J = 18.0 Hz, 2H, -CH₂CO-), 5.05 (dd, J = 15.4 Hz, 2H, -NCH₂-), 7.14 (t, J = 8.3 Hz, 1H, Ar-H), 7.35 (dd, J = 8.3 Hz, 1.6 Hz, 1H, Ar-H), 7.52 (m, 2H, Ar-H), 7.88 (dd, J = 9.96 Hz, 2.5 Hz, 1H, Ar-H), 8.27 (dd, J = 8.9 Hz, 6.1 Hz, 1H, Ar-H)

IR (KBr, cm^{-1}): 1710 (C=O), 1665 (C=O)

MS (m/e): 503 (10, M^+), 405 [14, M^+ -CONN=C(CH₃)₂]

Anal. Calcd.: C, 52.40; H, 3.20; N, 8.33

15 Found: C, 52.25; H, 3.08; N, 8.31

m.p. 152-154°C.

2-[(4-Bromo-2-fluorophenyl)methyl]-1'-(cyclopentylideneamino)spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone

¹H NMR (DMSO-d₆, 400 MHz): δ 1.7-1.78 (m, 4H, -CH₂CH₂CH₂CH₂-), 1.99 (t, J = 7.68 Hz, 2H, -CH₂CH₂CH₂CH₂-), 2.5 (t, J = 7.26 Hz, 2H, -CH₂CH₂CH₂CH₂-), 3.56 (s, 2H, -CH₂CO-), 5.0 (d, J = 15.36 Hz, 1H, -NHCH-), 5.13 (d, J = 15.36 Hz, 1H, -NHCH-), 7.16 (t, J = 8.3 Hz, 1H, Ar-H), 7.35 (dd, J = 8.3 Hz, 1.87 Hz, 1H, Ar-H), 7.53 (dd, J = 9.75 Hz, 1.87 Hz, 1H, Ar-H), 7.65 (dt, J = 8.3 Hz, 2.07 Hz, 1H, Ar-H), 7.81 (m, 2H, Ar-H), 8.2 (dd, J = 7.9 Hz, 0.85 Hz, 1H, Ar-H)

IR (KBr, cm^{-1}): 1705 (C=O), 1670 (C=O)

M/S (m/e): 511 (8, M^+)

Anal. Calcd.: C, 56.26; H, 3.74; N, 8.20

Found: C, 56.60; H, 3.74; N, 8.11

30 m.p. 142-144°C.

2093793

- 24 -

2-[(4-Bromo-2-fluorophenyl)methyl]-1'-[(tetrahydro-4H-pyran-4-yl-idene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone

¹H NMR (DMSO-d₆, 400 MHz): δ 2.16 (t, J = 5.6 Hz, 2H, -CH₂CH₂O-), 2.57 (t, J = 5.8 Hz, 2H, -CH₂CH₂O-), 3.5-3.6 (m, 4H, -CH₂CO-, -CH₂CH₂O-), 3.76-3.83 (m, 2H, -CH₂CH₂O-), 5.06 (dd, J = 15.15 Hz, 2H, -NCH₂-), 7.15 (t, J = 8.1 Hz, 1H, Ar-H), 7.36 (dd, J = 8.3 Hz, 1.87 Hz, 1H, Ar-H), 7.55 (dd, J = 9.96 Hz, 2.07 Hz, 1H, Ar-H), 7.65 (m, 1H, Ar-H), 7.82 (m, 2H, Ar-H), 8.2 (dd, J = 7.68 Hz, 0.83 Hz, 1H, Ar-H)

10 IR (KBr, cm⁻¹): 1705 (C=O), 1655 (C=O)

MS (m/e): 527 (62, M⁺)

Anal. Calcd.: C, 54.56; H, 3.62; N, 7.95

Found: C, 54.40; H, 3.65; N, 7.91


m.p. 161-163°C.

SUBSTITUTE SHEET

CLAIMS

We claim:

1. The compounds of structural formula (I)





5 wherein:


R^1 and R^2 are independently hydrogen, alkyl containing 1 to 6 carbon atoms, halogen, lower alkoxy containing 1 to 6 carbon atoms, trifluoromethyl, nitro, aryl or aryl (lower alkyl)oxy wherein aryl contains 6 to 10 carbon atoms and lower alkyl contains 1 to 6 carbon atoms;

10 R^3 is lower alkyl containing 1 to 6 carbon atoms, aryl, aryl (lower alkyl) or dihalogen substituted aryl (lower alkyl) wherein aryl contains 6 to 10 carbon atoms and lower alkyl contains 1 to 6 carbon atoms;

15 R^4 and R^5 are independently hydrogen, alkyl containing 1 to 12 carbon atoms, aryl, aryl (lower alkyl) wherein aryl contains 6 to 10 carbon atoms and lower alkyl contains 1 to 6 carbon atoms, or R^4 and R^5 are joined to form alicyclic or heterocyclic rings selected from the group consisting of




2. The compounds according to claim 1 of structural formula (I)



wherein:


- 5 R¹ and R² are hydrogen or halogen; R³ is dihalogen substituted benzyl; R⁴ and R⁵ are lower alkyl containing 1 to 3 carbon atoms or R⁴ and R⁵ are joined to form alicyclic or heterocyclic rings selected from the group consisting of



- 10 3. The compound according to claim 2 2-[(4-bromo-2-fluorophenyl)methyl]-1'-[(1-methylethylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.
4. The compound according to claim 2 2-[(4-bromo-2-fluorophenyl)methyl]-6-fluoro-1'-[(1-methylethylidene)amino]spiro-[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.
- 15 5. The compound according to claim 2 2-[(4-bromo-2-fluorophenyl)methyl]-1'-[(1-ethylpropylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.
6. The compound according to claim 2 2-[(4-bromo-2-fluorophenyl)methyl]-1'-(cyclopentylideneamino)spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.
- 20 7. The compound according to claim 2 2-[(4-bromo-2-fluorophenyl)methyl]-1'-[(tetrahydro-4H-pyran-4-ylidene)amino]spiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.

8. A pharmaceutical composition comprising a compound as claimed in any one of Claims 1 to 7 in combination or association with a pharmaceutically acceptable carrier.

9. A method of preparation of a compound as claimed in Claim 1, wherein a compound having the formula Ia




5

wherein R^1 , R^2 and R^3 are as defined in Claim 1 is reacted with a carbonyl compound having the formula R^4COR^5 where R^4 and R^5 are as defined in Claim 1 or a reactive derivative thereof.

10. A method of making a pharmaceutical composition claimed in Claim 8 wherein
10 a compound as claimed in any one of Claims 1 to 7 is brought into combination or association with a pharmaceutically acceptable carrier.

11. A compound having the formula Ia as illustrated in Claim 9 where R^1 , R^2 and R^3 are as defined in Claim 1.

12. A method of making a compound as claimed in Claim 11 wherein a dicarboxylic
15 acid having the formula



wherein R^1 , R^2 and R^3 are as defined in Claim 1 or a reactive derivative thereof is converted into an imide by reaction with hydrazine.

