

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2016/014240 A9

(43) International Publication Date
28 January 2016 (28.01.2016)

- (51) International Patent Classification:
C12Q 1/68 (2006.01) *C12N 15/09* (2006.01)
G06F 19/22 (2011.01)
- (21) International Application Number:
PCT/US2015/039362
- (22) International Filing Date:
7 July 2015 (07.07.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/028,527 24 July 2014 (24.07.2014) US
- (71) Applicant: **ABBOTT MOLECULAR INC.** [US/US];
1300 E. Touhy Avenue, Des Plaines, Illinois 60018 (US).
- (72) Inventors: **TANG, Ning**; 1300 E. Touhy Avenue, Des
Plaines, Illinois 60018 (US). **LECKIE, Gregor**; 1300 E.
Touhy Avenue, Des Plaines, Illinois 60018 (US). **PA-
HALAWATTA, Vihanga**; 1300 E. Touhy Avenue, Des
Plaines, Illinois 60018 (US). **FRANK, Andrea**; 1300 E.
Touhy Avenue, Des Plaines, Illinois 60018 (US). **LAMP-
INEN, John**; 1300 E. Touhy Avenue, Des Plaines, Illinois
60018 (US).
- (74) Agents: **ARENSON, Tanya A.** et al.; Casimir Jones, S.C.,
2275 Deming Way, Suite 310, Middleton, Wisconsin
53562 (US).
- (81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

- (88) Date of publication of the international search report:
17 March 2016
- (48) Date of publication of this corrected version:
6 May 2016
- (15) Information about Correction:
see Notice of 6 May 2016



WO 2016/014240 A9

(54) Title: COMPOSITIONS AND METHODS FOR THE DETECTION AND ANALYSIS OF MYCOBACTERIUM TUBERCULOSIS

(57) Abstract: Provided herein are compositions and methods useful for the detection of MTB. In particular, provided herein are kits, reagents, reaction mixtures, and methods involving such for nucleic acid amplification and detection procedures, which specifically and sensitively detect MTB in samples.

COMPOSITIONS AND METHODS FOR THE DETECTION AND ANALYSIS OF MYCOBACTERIUM TUBERCULOSIS

FIELD OF THE INVENTION

5 The present invention relates to mycobacterium tuberculosis. In particular, the invention relates to compositions and method for detecting mycobacterium tuberculosis.

BACKGROUND OF THE INVENTION

10 Mycobacterium tuberculosis (MTB) constitutes a serious threat to public health in the world and is second only to HIV/AIDS as the greatest killer worldwide due to a single infectious agent (Warren et al, Differentiation of *Mycobacterium tuberculosis* complex by PCR amplification of genomic regions of difference, 2006 Jul, Int J Tuberc Lung Dis.10 (7):818–822). The CDC reports that in 2011, there were an estimated 8.7 million new cases of MTB (13% co-infected with HIV); 1.4 million people died from MTB, including almost
15 one million deaths among HIV-negative individuals and 430,000 among people who were HIV-positive. The World Health Organization (WHO) reports that MTB is one of the top killers of women, with 300,000 deaths among HIV-negative women and 200,000 deaths among HIV-positive women in 2011. It is among the top three causes of death for women aged 15 to 44. MTB is also a leading killer of people living with HIV causing one quarter of
20 all deaths. There were an estimated 0.5 million cases and 64,000 deaths among children in 2011. Multi-drug resistant MTB (MDR-TB) is increasing and is present in virtually all countries surveyed. Geographically, the burden of MTB is highest in Asia and Africa. WHO reported that overall MTB case detection is still less than 60% in low-income countries (LICs) and only 66% globally. That is, of an estimated 8.7 million people who become ill
25 with MTB in 2011, 2.9 million with active disease were not diagnosed and notified to national MTB control programs. In addition, only 19% of MDR-MTB cases were appropriately diagnosed and notified. Fewer than 1 in 20 new MTB patients have access to drug susceptibility testing. Due to the risk of spread of MTB, the potential for the emergence of drug-resistant strains, and the severity of the disease in patients infected with HIV-1, a low
30 price, prompt and accurate MTB molecular test is extremely important. Routine cultures are time-consuming and can take up to six weeks. Microscopic examination of acid-fast smears is the most rapid method for the detection of mycobacteria, but it is insensitive and non-specific.

SUMMARY OF THE INVENTION

Provided herein are compositions and methods useful for the detection of MTB. In particular, provided herein are kits, reagents, reaction mixtures, and methods involving such
5 for nucleic acid amplification and detection procedures, which specifically and sensitively detect MTB in samples. Such compositions and method include primers, probes, primer sets, primer and probe sets, and methods for detecting MTB complex in different human samples such as sputum, bronchial alveolar lavage (BAL) and N-acetyl-L-cysteine (NALC)-NaOH sediments of sputum and BAL samples.

10 In some embodiments, two or more of the polynucleotide reagents provided herein as SEQ ID NOs: 1-9 are combined in a composition (e.g., reagent set, kit, reaction mixture, etc.). In some embodiments, one or more or all of the nucleic acid reagents comprise a detectable moiety (e.g., synthetic label). In some embodiments, the compositions comprise one or more primers SEQ ID NOs: 1-4 or 7-8. In some embodiments, the compositions
15 comprise one or more primer pairs SEQ ID NOs: 1 and 2, 3 and 4, or 7 and 8. In some embodiments, the compositions comprise one or more probes (e.g., labeled probes) of SEQ ID NOs: 5, 6, or 9. In some embodiments, the compositions comprise primer and probe sets: SEQ ID NOs 1-2 and 5, 3-4 and 6, or 7-9. In some embodiments, the compositions comprise internal control reagents, such as SEQ ID NOs: 7-9. In some embodiments, the compositions
20 comprise a dual probe system comprising SEQ ID NOs: 5 and 6.

In some embodiments, the compositions and methods employ reagents sets comprising a polynucleotide component having primers, probes, primers sets, and/or probe sets. In some embodiments, the polynucleotide component of the composition consists of the primer, probe, primer set, or probe set combinations described above. As reaction mixtures,
25 the compositions may consist of such polynucleotides as well as any polynucleotides included in a sample (i.e., the only non-sample nucleic acid molecules are the polynucleotides represented by SEQ ID NOs: 1-9, individually or in combinations (e.g., the combinations described above).

The primer sets herein provided comprise two primers, and are useful for the
30 amplification of target sequences, e.g., in PCR. In some embodiments, the compositions comprise at least two primers and one or more (e.g., two or more) probes that detect amplicons generated by the primers.

Also provided herein are methods for detecting MTB in a sample. In some embodiments, the methods comprise (a) forming a reaction mixture comprising nucleic acid

amplification reagents, at least polynucleotide primer or probe described herein, and a test sample potentially containing at least one target sequence; and (b) subjecting the mixture to amplification conditions to generate at least one copy of a nucleic acid sequence complementary to the target sequence. In some embodiments the method further comprises
5 detecting generated amplicons. In some embodiments, the detecting comprises (c) hybridizing a probe to the nucleic acid sequence complementary to the target sequence so as to form a hybrid comprising the probe and the nucleic acid sequence complementary to the target sequence; and (d) detecting, directly or indirectly, the hybrid as an indication of the presence of MTB in the test sample.

10 Further, when the amplification is PCR, or a similar thermocycling amplification process, step (b) can be repeated multiple times to increase the number of target sequence copies.

According to another embodiment, both MTB and one or more additional infectious agents (e.g., HIV) or other nucleic acid molecules (e.g., human sequences) are detected.

15 Accordingly, in some embodiments, compositions comprise reagents for detecting such other agents or nucleic acid molecules.

In some embodiments, the compositions and methods further employ control reagents or kit components (e.g., positive controls, negative controls). In some embodiments, the control reagents include a synthetic target nucleic acid. In some embodiments, the control
20 reagents include reagents for detecting an MTB, human, or other sequence expected to be present in a sample. In some embodiments, a control target nucleic acid, whether synthetic or endogenous in a sample, is selected such that amplification primers that amplify the MTB target nucleic acid also amplify the control target nucleic acid. In some such embodiments, a
25 probe that detects the MTB target nucleic acid or an amplicon generated therefrom does not detect the control target or an amplicon generated therefrom. In some embodiments, a control probe is provided that detects the control target nucleic acid or an amplicon generated therefrom but does not detect the MTB target nucleic acid or an amplicon generated therefrom. In some embodiments, internal standards are provided for quantitation.

In some embodiments, kits, in addition to the reagents discussed above, include one or
30 more suitable containers, instructions for use, software (e.g., data analysis software), and the like. In some embodiments, kits include reagents for labeling polynucleotides. In some embodiments, one or more components in the kit is in lyophilized form.

Embodiments of the present disclosure provide compositions, kits, systems, and methods for identifying MTB in complex biological samples such as sputum or

bronchoalveolar lavage and sediments thereof. In some embodiments, the compositions and methods provide inactivation reagents, and single probe or multiple probe real time detection methods that are able to specifically and accurately isolate and identify MTB.

For example, in some embodiments, the present disclosure provides a composition, comprising: at least one (e.g., one, two, or three) primer pair(s) selected from SEQ ID NOs: 1 and 2, SEQ ID NOs: 3 and 4, or SEQ ID NOs: 7 and 8. In some embodiments, the composition comprises SEQ ID NOs: 1-4 and 7-8. In some embodiments, the composition further comprises at least one probe selected from SEQ ID NOs: 5, 6, or 9.

Further embodiments provide a composition, comprising: a set of primer pairs of SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4. In some embodiments, the composition further comprises at least one probe selected from SEQ ID NOs: 5, 6, or 9. In some embodiments, the composition further comprises the primer pair of SEQ ID NOs: 7 and 8.

Additional embodiments provide a composition, comprising each the nucleic acids of SEQ ID NOs: 1-9. In some embodiments, the above compositions include or are substituted with one or more nucleic acid sequences selected from the SEQ ID NOs: 10-36.

Embodiments of the disclosure provide a kit, comprising: a) any of the aforementioned compositions, and b) at least one reagent for performing a nucleic acid amplification reaction (e.g., a nucleic acid polymerase; a plurality of dNTPS, a buffer, or an inactivation reagent). In some embodiments, the inactivation reagent comprises water, a detergent, an alcohol, and NaOH (e.g., isopropanol, sodium hydroxide, TWEEN-20, and water).

In other embodiments, the disclosure provides a reaction mixture, comprising: any of the aforementioned compositions or nucleic acids hybridized to a microbacterium tuberculosis (MTB) nucleic acid. In some embodiments, the MTB target nucleic acid is one or more (e.g., both) of insertion sequence (IS) 6110 and Protein Antigen B (PAB).

In further embodiments, the present disclosure provides a method of identifying an MTB nucleic acid in a biological sample, comprising: a) contacting a biological sample from a subject with any of the aforementioned nucleic acid primers or probes, and b) directly or indirectly detecting the binding of the nucleic acid primers or probes to the MTB nucleic acid. In some embodiments, the method further comprises the step of c) determining the presence of MTB in the sample when the binding is detected. In some embodiments, the detecting is via real time PCR detecting. In some embodiments, the method further comprises the step of inactivating MTB in the sample using the inactivation buffer. In some embodiments, the sample is sputum, bronchoalveolar lavage [BAL], or N-acetyl-L-cysteine

[NALC] sediments of sputum and BAL. In some embodiments, the method further comprises the step of extracting DNA from the sample following inactivation.

Yet other embodiments provide a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating the biological sample with an inactivating reagent to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting the DNA with one or more primer pairs and one or more nucleic acid probes; d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and e) identifying the presence of the targets in the sample.

Further embodiments provide a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating said biological sample with an inactivating reagent comprising isopropanol, sodium hydroxide, TWEEN-20, and water to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting the DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6; d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and e) identifying the presence of the targets in said sample.

Additional embodiments provide a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating said biological sample with an inactivating reagent to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting the DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6; d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and e) identifying the presence of the targets in the sample.

Other embodiments provide a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating the biological sample with an inactivating reagent to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting the DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6; d) performing a real time PCR assay to amplify one or more MTB nucleic acid targets; and e) identifying the presence of the targets in the sample.

Still other embodiments provide a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating the biological sample with an inactivating reagent to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting said DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and

SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6; d) performing an amplification assay to amplify one or more MTB nucleic acid targets selected from IS6110 and PAB; and e) identifying the presence of the targets in the sample.

In certain embodiments, the present disclosure provides a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating the biological sample with an inactivating reagent to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting the DNA with the primer pairs of SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and the nucleic acid probes of SEQ ID NOs: 5 and 6; d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and e) identifying the presence of the targets in the sample.

In some embodiments, the present disclosure provides a method of detecting an MTB nucleic acid in a biological sample, comprising: a) inactivating the biological sample with an inactivating reagent comprising isopropanol, sodium hydroxide, TWEEN-20, and water to generate an inactivated sample; b) extracting DNA from the inactivated sample; c) contacting the DNA with the primer pairs of SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and the nucleic acid probes of SEQ ID NOs: 5 and 6; d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and e) identifying the presence of the targets in the sample.

Additional embodiments are described herein.

DESCRIPTION OF FIGURES

Figure 1 shows an MTB assay work-flow diagram in some embodiments of the technology provided herein.

Figure 2 shows data from detection of 46 MTB phylogenetically and geographically diverse MTB isolates.

Figure 3 shows an MTB assay work-flow diagram for sample preparation.

Figure 4 shows mean cycle number values determined when MTB complex genomic DNAs were tested to determine assay inclusivity.

DETAILED DESCRIPTION

Provided herein are compositions and methods useful for the detection of MTB. In particular, provided herein are kits, reagents, reaction mixtures, and methods involving such for nucleic acid amplification and detection procedures, which specifically and sensitively detect MTB in samples.

In some embodiments, provided herein are polynucleotides that specifically hybridize with a nucleic acid sequence, or complement thereof, of MTB. These polynucleotides find use to amplify MTB, if present in a sample, and to specifically detect the presence of MTB. Exemplary polynucleotides are described, for example, by SEQ ID NOs: 1-9 or 10-36.

5 In some embodiments, assays described herein utilize multiple (e.g., two) different MTB-specific primer/probe sets. For example, in some embodiments, a first set is designed to detect the multi-copy insertion element, IS6110 (Thierry D, et al., *Nucleic Acids Res* 1990;18:188), and second set, the single copy gene, PAB (Anderson AB, Hansen EB, *Infect Immun* 1989;57:2481-2488). Because there have been reports of MTB strains that lack
10 IS6110 (Mathema B, et al., *Clinical Microbiology Reviews* 2006;19:658-685), or that have a deletion in the PAB gene (Gilpin CM, et al., *J Clin Microbiol* 2002;40:2305-2307), the use of both targets minimizes the risk of false negative results. Experiments described herein demonstrated that the dual target strategy results in the detection of MTB genomic DNA with high reliability.

15 The mycobacterial cell wall is resistant to conventional cell lysis techniques due to the complex structure of lipophilic molecules and polysaccharides. Thus, in some embodiments, MTB detection assays utilize a guanidinium thiocyanate-magnetic microparticle purification method with optimized incubation temperatures and mixing conditions for TB cell lyses and genomic DNA release. Experiments conducted during the development of embodiments of
20 the assays described herein showed that the sample DNA extraction method is comparable in efficiency to mechanical bead beating for TB cell lyses.

Further experiments demonstrated that all 66 MTB complex DNAs (including eight different MTB complex species) were detected by the assay. Reliability of the assay was assessed in four ways. First, specificity of the assay was assessed by testing 80 potentially
25 different cross-reactors. None of the potential cross-reactors were detected. Second, a carryover assessment was performed in which high positive MTB samples were processed alongside negative samples to determine if false positives, or carryover, were detected in the negative samples. No false positives were observed. Third, various potentially interfering substances were tested for their impact on assay performance. No interference was observed,
30 except with 8.3% and 5.0% bovine mucus, where interference was observed. This interference was removed when the mucus concentration was reduced to $\leq 2.5\%$. When clinical specimens were tested, the rate of specimens with invalid IC results was 0.3%, demonstrating that the sample preparation methods described herein removed PCR inhibitors in an effective manner. This provides evidence of the robustness of the protocol and indicates

that the impact of interference caused by bovine mucus is likely not significant to the assay. Finally, a reproducibility study was performed in which multiple users used multiple *m2000* instrument systems, or manual sample preparation, to test low positive (three times LOD) and negative panels. 100% reproducibility was observed. These data support the robust nature of the assay when used in analytical studies and clinical samples testing

The clinical utility of the MTB detection assays was assessed by testing sputum and NALC specimens collected from patients suspected of having TB in five countries using both archived samples and prospectively collected samples. Overall assay sensitivity was 93%, while it was 99% in smear positive culture positive samples and 81% in smear negative culture positive samples. Specificity was 97%. The results of the analytical specificity test and the sputum sample testing from non-TB suspect population from within the U.S. all showed 100% specificity. The clinical specificity was determined based on comparison of assay results with culture results.

Embodiments of the technology described herein provide high throughput, automated MTB detection with high sensitivity and specificity. Compared with conventional culture assays, the technology significantly improves the rapid diagnosis of TB by allowing the direct detection of mycobacteria in clinical specimens. The assays provide superior sensitivity and specificity compared to conventional acid-fast smears microscopic examination. A gap with current MTB diagnostic assays is the lack of sensitivity in culture positive and smear negative populations (with low TB concentration in samples). Embodiments of the technology provided herein fill that gap. Assays provided herein are robust with very low inhibition rate even if difficult to work with sputum samples. This reduces the time required for repeat testing of invalid samples. In some embodiments, a multi-copy MTB target is interrogated, providing greater target sensitivity and less chance of false negative assay results caused by mutations/deletions in the target region. Embodiments further provide unique and effective MTB inactivation methods.

The term "specifically hybridize" as used herein refers to the ability of a nucleic acid to bind detectably and specifically to a second nucleic acid. Polynucleotides specifically hybridize with target nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to non-specific nucleic acids. Stringent conditions that can be used to achieve specific hybridization are known in the art.

A "target sequence" or "target nucleic acid sequence" as used herein means a nucleic acid sequence of MTB or other sequence to be detected (e.g., HIV), or complement thereof, that is amplified, detected, or both amplified and detected using one or more of the

polynucleotides herein provided. Additionally, while the term target sequence sometimes refers to a double stranded nucleic acid sequence, those skilled in the art will recognize that the target sequence can also be single stranded. In cases where the target is double stranded, polynucleotide primer sequences preferably amplify both strands of the target sequence. A target sequence may be selected that is more or less specific for a particular organism. For example, the target sequence may be specific to an entire genus, to more than one genus, to a species or subspecies, serogroup, auxotype, serotype, strain, isolate or other subset of organisms.

The term "test sample" as used herein, means a sample taken from an organism, biological fluid, environmental sample, or other sample that is suspected of containing or potentially contains an MTB target sequence. The test sample can be taken from any biological source, such as for example, tissue, blood, saliva, sputa, N-acetyl-L-cysteine (NALC)-NaOH sediments of sputum, mucus, bronchial alveolar lavage (BAL), sweat, urine, urethral swabs, cervical swabs, urogenital or anal swabs, conjunctival swabs, ocular lens fluid, cerebral spinal fluid, milk, ascites fluid, synovial fluid, peritoneal fluid, amniotic fluid, fermentation broths, cell cultures, chemical reaction mixtures and the like. The test sample can be used (i) directly as obtained from the source or (ii) following a pre-treatment to modify the character of the sample. Thus, the test sample can be pre-treated prior to use by, for example, preparing plasma or serum from blood, disrupting cells or viral particles, preparing liquids from solid materials, diluting viscous fluids, filtering liquids, distilling liquids, concentrating liquids, inactivating interfering components, adding reagents, purifying nucleic acids, and the like.

The term "label" as used herein means a molecule or moiety having a property or characteristic which is capable of detection and, optionally, of quantitation. A label can be directly detectable, as with, for example (and without limitation), radioisotopes, fluorophores, chemiluminophores, enzymes, colloidal particles, fluorescent microparticles and the like; or a label may be indirectly detectable, as with, for example, specific binding members. It will be understood that directly detectable labels may require additional components such as, for example, substrates, triggering reagents, quenching moieties, light, and the like to enable detection and/or quantitation of the label. When indirectly detectable labels are used, they are typically used in combination with a "conjugate". A conjugate is typically a specific binding member that has been attached or coupled to a directly detectable label. Coupling chemistries for synthesizing a conjugate are well known in the art and can include, for example, any chemical means and/or physical means that does not destroy the specific binding property of

the specific binding member or the detectable property of the label. As used herein, "specific binding member" means a member of a binding pair, e.g., two different molecules where one of the molecules through, for example, chemical or physical means specifically binds to the other molecule. In addition to antigen and antibody specific binding pairs, other specific binding pairs include, but are not intended to be limited to, avidin and biotin; haptens and antibodies specific for haptens; complementary nucleotide sequences; enzyme cofactors or substrates and enzymes; and the like.

A polynucleotide is a nucleic acid polymer of ribonucleic acid (RNA), deoxyribonucleic acid (DNA), modified RNA or DNA, or RNA or DNA mimetics (such as, without limitation PNAs), and derivatives thereof, and homologues thereof. Thus, polynucleotides include polymers composed of naturally occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as polymers having non-naturally-occurring portions that function similarly. Such modified or substituted nucleic acid polymers are well known in the art and for the purposes of the present invention, are referred to as "analogues." For ease of preparation and familiarity to the skilled artisan, polynucleotides are preferably modified or unmodified polymers of deoxyribonucleic acid or ribonucleic acid.

Polynucleotide analogues that are useful include polymers with modified backbones or non-natural internucleoside linkages. Modified backbones include those retaining a phosphorus atom in the backbone, such as phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates, as well as those no longer having a phosphorus atom, such as backbones formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. An example of such a non-phosphorus containing backbone is a morpholino linkage (see, for example, U.S. Pat. Nos. 5,185,444, 5,034,506, and 5,142,047 all of which are herein incorporated by reference). Modified nucleic acid polymers (analogues) may contain one or more modified sugar moieties. For example, sugar moieties may be modified by substitution at the 2' position with a 2-methoxyethoxy (2-MOE) group (see, for example, Martin et al., (1995) *Helv. Chim. Acta*, 78:486-504).

Embodiments also contemplate analogues that are RNA or DNA mimetics, in which both the sugar and the internucleoside linkage of the nucleotide units are replaced with novel groups. In these mimetics the base units are maintained for hybridization with the target sequence. An example of such a mimetic, which has been shown to have excellent hybridization properties, is a peptide nucleic acid (PNA) (Nielsen et al., (1991) *Science*,

254:1497-1500; International Patent Application WO 92/20702, herein incorporated by reference). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, for example an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to the aza-nitrogen atoms of the amide
5 portion of the backbone.

Contemplated polynucleotides may further include derivatives wherein the nucleic acid molecule has been covalently modified by substitution, chemical, enzymatic, or other appropriate means with a moiety other than a naturally occurring nucleotide, for example with a moiety that functions as a label, as described herein.

10 The present invention further encompasses homologues of the polynucleotides having nucleic acid sequences set forth in SEQ ID NOs: 1-9 or 10-36. Homologues are nucleic acids having at least one alteration in the primary sequence set forth in any one of SEQ ID NOs: 1-9 or 10-36, that does not destroy the ability of the polynucleotide to specifically hybridize with a target sequence, as described above. Accordingly, a primary sequence can be altered,
15 for example, by the insertion, addition, deletion or substitution of one or more of the nucleotides of, for example, SEQ ID NOs: 1-9 or 10-36. Thus, homologues that are fragments of a sequence disclosed in SEQ ID NOs: 1-9 or 10-36 may have a consecutive sequence of at least about 7, 10, 13, 14, 15, 16, 17, 18, 19 20, 21, 22, 23 or more nucleotides of the nucleic acid sequences of SEQ ID NO: 1-9 or 10-36, and will retain the ability to specifically
20 hybridize with a target sequence, as described above. Ordinarily, the homologues will have a nucleic acid sequence having at least about 50%, 60%, 70%, 80%, 85%, 90% or 95% nucleic acid sequence identity with a nucleic acid sequence set forth in SEQ ID NOs: 1-9 or 10-36. Identity with respect to such sequences is defined herein as the percentage of nucleotides in the candidate sequence that are identical with the known polynucleotides after aligning the
25 sequences and introducing gaps, if necessary, to achieve the maximum percent identity. Terminal (5' or 3') or internal deletions, extensions or insertions into the nucleotide sequence shall not be construed as affecting identity.

In some embodiments, the polynucleotides comprise primers and probes that specifically hybridize to an MTB target sequence, for example the nucleic acid molecules
30 having any one of the nucleic acid sequences set forth in SEQ ID NOs: 1-9 or 10-36, including analogues and/or derivatives of said nucleic acid sequences, and homologues thereof, that can specifically hybridize with an MTB target sequence. As described below, polynucleotides find use as primers and/or probes to amplify or detect MTB.

The polynucleotides can be prepared by a variety of techniques. For example, the polynucleotides can be prepared using solid-phase synthesis using commercially available equipment, such as that available from Applied Biosystems USA Inc. (Foster City, Calif.), DuPont, (Wilmington, Del.), or Milligen (Bedford, Mass.). Modified polynucleotides, such as phosphorothioates and alkylated derivatives, can also be readily prepared (see, for example, 5 U.S. Pat. Nos. 5,464,746; 5,424,414; and 4,948,882).

The polynucleotides can be employed directly as probes for the detection, or quantitation, or both, of MTB nucleic acids in a test sample. The test sample is contacted with at least one of the polynucleotides under suitable hybridization conditions and the 10 hybridization between the target sequence and at least one of the polynucleotides is then detected. Detection can be direct or indirect. In some embodiments, a hybrid between the probe and target is detected directly. In some embodiments, the hybrid is detected indirectly, for example, by detecting reaction byproducts generated by an enzymatic reaction that occurs in the presence of a duplex between a probe and the MTB target.

15 The polynucleotides may incorporate one or more detectable labels. Detectable labels are molecules or moieties having a property or characteristic that can be detected directly or indirectly and are chosen such that the ability of the polynucleotide to hybridize with its target sequence is not adversely affected.

Detection labels have the same definition as "labels" previously defined and "capture 20 labels" are typically used to separate extension products, and probes associated with any such products, from other amplification reactants. Specific binding members (as previously defined) are well suited for this purpose. Also, probes used according to this method may be blocked at their 3' ends so that they are not extended under hybridization conditions. Methods for preventing extension of a probe are well known and are a matter of choice for one skilled 25 in the art.

In cases where labels are employed to detect primer-amplified products, primer sequences optionally can be labeled with either a capture label or a detection label. In some embodiments, primer comprise a 3' portion that hybridizing to an MTB target nucleic acid and a 5' portion that introduces a non-MTB sequence to extension products generated 30 therefrom. Such 5' portions may include a synthetic tag sequence for use, for example, in next-generation sequencing technologies.

In some embodiments, a probe is used to hybridize with the extension product or amplicon generated by the primer sequences, and typically hybridizes with a sequence that does not include the primer sequences. Similarly to the primer sequence, the probe sequence

can also be labeled with either a capture label or a detection label with the caveat that, in some embodiments, when the primer is labeled with a capture label, the probe is labeled with a detection label, and vice versa. Upon formation of the copy sequence/probe hybrids, the differential labels (i.e., capture and detection labels) on the copy sequence and probe sequence can be used to separate and detect such hybrids.

The polynucleotides are also suitable for use as capture probes in sandwich-type assays. Briefly, the polynucleotide capture probe is attached to a solid support and brought into contact with a test sample under suitable hybridization conditions such that a probe:target hybrid is formed between the capture probe and any target nucleic acid present in the test sample. After one or more appropriate washing steps, the probe:target hybrid is detected, usually by means of a second "disclosure" probe or by a specific antibody that recognizes the hybrid molecule.

Embodiments also contemplate the use of the polynucleotides in modified nucleic acid hybridization assays. For example, U.S. Pat. No. 5,627,030 discloses a method to amplify the detection signal in a nucleic acid hybridization assay. In the disclosed assay, a first polynucleotide probe sequence is hybridized under suitable conditions to a target sequence, the probe:target hybrid is subsequently immunocaptured and immobilized. A second polynucleotide probe which contains many repeating sequence units is then hybridized to the probe component of the probe:target hybrid. Detection is achieved by hybridization of many labeled nucleic acid sequence probes, one to each of the repeating sequence units present in the second probe. The attachment of multiple labeled probes to the second probe thus amplifies the detection signal and increases the sensitivity of the assay.

Amplification and Detection of MTB Nucleotide Sequences

The polynucleotides can be used as primers or probes to amplify and/or detect MTB in a test sample. The primer/probe sets provided herein comprise at least two primers and at least one probe. These primer/probe sets can be employed according to nucleic acid amplification techniques. Hence, the primers in any particular primer/probe set can be employed to amplify a target sequence. In most cases, the probe hybridizes to the copies of the target sequence generated by one or more of the primers and generally facilitates detecting any copies of the target sequence generated during the course of the amplification reaction. All of the primer/probe sets can be employed according to nucleic acid amplification procedures to specifically and sensitively detect MTB when the appropriate primers and probes are combined. It is contemplated that the individual primers and probes of

the primer/probe sets provided herein may alternatively be used in combination with primers and/or probes other than those described in the primer/probe sets provided herein. In some embodiments, two primer and probes sets are employed to detect two different MTB target sequences.

5 Amplification procedures include, but are not limited to, polymerase chain reaction (PCR), TMA, rolling circle amplification, nucleic acid sequence based amplification (NASBA), and strand displacement amplification (SDA). One skilled in the art will understand that for use in certain amplification techniques the primers may need to be modified, for example, for SDA the primer comprises additional nucleotides near its 5' end
10 that constitute a recognition site for a restriction endonuclease. Similarly, for NASBA the primer comprises additional nucleotides near the 5' end that constitute an RNA polymerase promoter.

In some embodiments, certain criteria are taken into consideration when selecting a primer for an amplification reaction. For example, when a primer pair is required for the
15 amplification reaction, the primers should be selected such that the likelihood of forming 3' duplexes is minimized, and such that the melting temperatures (T_M) are sufficiently similar to optimize annealing to the target sequence and minimize the amount of non-specific annealing.

In some embodiments, the amplification methods comprises (a) forming a reaction
20 mixture comprising nucleic acid amplification reagents, at least one primer/probe set, and a test sample suspected of containing at least one target sequence and (b) subjecting the mixture to amplification conditions to generate at least one copy of a nucleic acid sequence complementary to the target sequence. Step (b) of the above methods can be repeated any
25 suitable number of times (prior to step (c) in the detection method), e.g., by thermal cycling the reaction mixture between 10 and 100 times, typically between about 20 and about 60 times, more typically between about 25 and about 45 times.

Nucleic acid amplification reagents include but are not limited to, an enzyme having at least polymerase activity, enzyme cofactors such as magnesium or manganese; salts; nicotinamide adenine dinucleotide (NAD); and deoxynucleotide triphosphates (dNTPs) such
30 as for example deoxyadenine triphosphate, deoxyguanine triphosphate, deoxycytosine triphosphate and deoxythymine triphosphate.

Amplification conditions are conditions that generally promote annealing and extension of one or more nucleic acid sequences.

Specific amplicons produced by amplification of target nucleic acid sequences using the polynucleotides, as described above, can be detected by a variety of methods. For example, one or more of the primers used in the amplification reactions may be labeled such that an amplicon can be directly detected by conventional techniques subsequent to the amplification reaction. Alternatively, a probe consisting of a labeled version of one of the primers used in the amplification reaction, or a third polynucleotide distinct from the primer sequences that has been labeled and is complementary to a region of the amplified sequence, can be added after the amplification reaction is complete. The mixture is then submitted to appropriate hybridization and wash conditions and the label is detected by conventional methods.

The amplification product produced as above can be detected during or subsequently to the amplification of the target sequence. Methods for detecting the amplification of a target sequence during amplification (e.g., real-time PCR) are outlined above, and described, for example, in U.S. Pat. No. 5,210,015. Alternatively, amplification products are hybridized to probes, then separated from other reaction components and detected using microparticles and labeled probes.

It will be readily appreciated that a procedure that allows both amplification and detection of target nucleic acid sequences to take place concurrently in a single unopened reaction vessel would be advantageous. Such a procedure avoids the risk of "carry-over" contamination in the post-amplification processing steps, and also facilitates high-throughput screening or assays and the adaptation of the procedure to automation. Furthermore, this type of procedure allows "real-time" monitoring of the amplification reaction as well as "end-point" monitoring. Examples of probe molecules that are particularly well-suited to this type of procedure include molecular beacon probes and TAQMAN probes. TAQMAN probes are generally dual-labeled fluorogenic nucleic acid probes composed of a polynucleotide complementary to the target sequence that is labeled at the 5' terminus with a fluorophore and at the 3' terminus with a quencher. In the free probe, the close proximity of the fluorophore and the quencher ensures that the fluorophore is internally quenched. During the extension phase of the amplification reaction, the probe is cleaved by the 5' nuclease activity of the polymerase and the fluorophore is released. The released fluorophore can then fluoresce and thus produces a detectable signal.

In some embodiments, "molecular beacon" probes are employed. Molecular beacon probes are described, for example, in U.S. Pat. Nos. 6,150,097; 5,925,517 and 6,103,476 (herein incorporated by reference in their entireties). Basically, molecular beacons are

polynucleotide probes capable of forming a stem-loop (hairpin) structure. The loop is a single-stranded structure containing sequences complementary to the target sequence, whereas the stem typically is unrelated to the target sequence and self-hybridizes to form a double-stranded region. Nucleotides that are both complementary to the target sequence and that can self-hybridize may also form part of the stem region. Attached to one arm of the stem is a fluorophore moiety and to the other arm a quencher moiety. When the polynucleotide adopts a hairpin shape, the fluorophore and the quencher are in close proximity and the energy emitted by the fluorophore is thus absorbed by the quencher and given off as heat, resulting in internal quenching of the fluorophore. Upon binding of the polynucleotide to its target sequence, the fluorophore and the quencher become spatially separated and the fluorophore can fluoresce producing a detectable signal.

Examples of fluorophores that find use include, but are not limited to, fluorescein and fluorescein derivatives such as a dihalo-(C₁ to C₈)dialkoxycarboxyfluorescein, 5-(2'-aminoethyl)aminonaphthalene-1-sulphonic acid (EDANS), coumarin and coumarin derivatives, Lucifer yellow, Texas red, tetramethylrhodamine, tetrachloro-6-carboxyfluorescein, 5-carboxyrhodamine, cyanine dyes and the like. Quenchers include, but are not limited to, DABCYL, 4'-(4-dimethylaminophenylazo)benzoic acid (DABSYL), 4-dimethylaminophenylazophenyl-4'-maleimide (DABMI), tetramethylrhodamine, carboxytetramethylrhodamine (TAMRA), Black Hole Question (BHQ) dyes and the like.

In some embodiments, quantitative assays are employed. In some such embodiments, an internal standard is employed in the reaction. Such internal standards generally comprise a control target nucleic acid sequence and a control polynucleotide probe. The internal standard can optionally further include an additional pair of primers. The primary sequence of these control primers may be unrelated to the MTB polynucleotides and specific for the control target nucleic acid sequence. Alternatively, no additional primer need be used if the control target sequence is designed such that it binds the MTB primers. The amount of target nucleic acid in a test sample can be quantified using "end point" methods or "real time" methods.

In some embodiments, MTB detection assays are provided as high-throughput assays. For high-throughput assays, reaction components are usually housed in a multi-container carrier or platform, such as a multi-well microtiter plate, which allows a plurality of assay reactions containing different test samples to be monitored in the same assay. In some embodiments, highly automated high-throughput assays are employed to increase the efficiency of the screening or assay process. Many high-throughput screening or assay systems are now available commercially, as are automation capabilities for many procedures

such as sample and reagent pipetting, liquid dispensing, timed incubations, formatting samples into microarrays, microplate thermocycling and microplate readings in an appropriate detector, resulting in much faster throughput times. In some embodiments, reactions are performed in microfluidic devices (e.g., cards).

5 The polynucleotides, methods, and kits are useful in clinical or research settings for the detection and/or quantitation of MTB nucleic acids. Thus, in these settings the polynucleotides can be used in assays to diagnose MTB infection in a subject, or to monitor the quantity of an MTB target nucleic acid sequence in a subject infected with MTB. Monitoring the quantity of bacteria in a subject is particularly important in identifying or
10 monitoring response to anti-bacterial therapy.

 In some embodiments, a dual target assay is performed using real-time PCR, combined with sample inactivation. While a variety of sample may be used, highly clinically relevant sample include smear positive or smear negative specimens of sputum (induced or expectorated), bronchoalveolar lavage (BAL) samples, or N-Acetyl-LCysteine (NALC)-
15 treated sediments of sputum and BAL samples. Challenges presented with these samples include the molecular complexity of sputum, which contains numerous components that can interfere with molecular assays, cell lysis, and cell inactivation.

 In some embodiments, a sample inactivation step is performed to reduce the infection risk associated with clinical specimens that may contain MTB. Reduction of infection risk is
20 achieved, for example, by incubating clinical samples with inactivation reagent (see Example 3, below).

 In some embodiments, the assays are amenable for use with automated real-time PCR detection system, such as the Abbott *m2000sp* system. Thus, in some embodiments, prior to conducting an assay, the samples are prepared for use with such systems. For example, in
25 some embodiments, preparation of target DNA is performed using a magnetic microparticle-based technology (Abbott *mSample Preparation SystemDNA*). This can be performed using an Abbott *m2000sp* for automated sample preparation or using a manual sample preparation protocol. In some embodiments, an internal control (IC), positive control, and negative control are processed from the start of sample preparation to demonstrate that the process has
30 proceeded correctly.

 For amplification, in some embodiments, purified sample DNA and master mix are added to a 96-well PCR plate using an Abbott *m2000sp* instrument or manually. After addition, each plate is sealed and transferred to an Abbott *m2000rt* where PCR amplification is performed using DNA Polymerase.

In some embodiments, the presence of MTB amplification products is detected during the annealing/extension step by measuring the real-time fluorescence signal of the MTB probes. The presence of IC amplification products is detected by measuring the real-time fluorescence signal of the IC probe. In some embodiments, the MTB and IC probes are single-stranded DNA oligonucleotides consisting of the target-specific binding sequence, a fluorescent moiety covalently linked to the 5' end of the probe, and a quenching moiety covalently linked to the 3' end of the probe. In the absence of the MTB or IC target sequences, probe fluorescence is quenched. In the presence of MTB or IC target sequences, the MTB or IC probes specifically bind to their complementary sequences in the targets during the annealing/extension step, allowing fluorescent emission and detection. In some embodiments, the MTB probes are labeled with different fluorescent dyes (FAM™ for MTB target probes, Quasar® for IC), thus allowing the amplification products of MTB and IC to be simultaneously detected in the same reaction.

In some embodiments, steps are taken to avoid nucleic acid contamination. For example, in some embodiments, contamination is minimized because: PCR amplification and oligonucleotide hybridization occur in a sealed multi-well plates; detection is carried out automatically without the need to open the reaction vessels (e.g., plate wells); aerosol barrier pipette tips are used for all pipetting; the pipette tips are discarded after use; and separate dedicated areas are used to perform the MTB assay.

In some embodiments, the above reagents are provided in the form of a kit and/or system (e.g., systems comprising automated sample handling and assay instruments described herein). For example, in some embodiments, the kit comprises, consists essentially of, or consists of:

1. MTB Internal Control (4 vials, 0.4 mL per vial) < 0.01% noninfectious linearized DNA plasmid in a buffer solution with carrier DNA. Preservatives: Sodium azide and 0.15% ProClin® 950.
2. Amplification Reagent Pack (4 packs, 24 tests/pack). Each Reagent Pack contains: 1 bottle (0.078 mL) DNA Polymerase (5.4 to 5.9 units/μL) in buffered solution with stabilizers. 1 bottle (0.5314 mL) MTB Amplification Reagent. < 0.1% synthetic oligonucleotides (one or more target primer sets and probes; a primer set and probe for the internal control), and < 0.6% dNTPs in a buffered solution with a reference dye. Preservatives: Sodium azide and 0.15% ProClin® 950. 1 bottle (0.778 mL) Activation Reagent. 38

mM magnesium chloride in a buffered solution. Preservatives: Sodium azide and 0.15% ProClin® 950.

3. MTB Negative Control (8 vials, 1.6 mL per vial); Buffered solutions; Preservatives: Sodium azide and 0.15% ProClin® 950.
- 5 4. MTB Positive Control (8 vials, 1.6 mL per vial); < 0.01% noninfectious linearized DNA plasmid in a buffer solution with carrier DNA. Preservatives: Sodium azide and 0.15% ProClin® 950.

In some embodiments, all forms of MTB are detected (e.g., the primers and probes are selected to identify all MTB nucleic target sequences that might be present in a sample). In
10 some embodiments, specific MTB sequences are detected, such as antibiotic-resistant strains (e.g., rifampicin, isoniazid).

EXAMPLES

The following examples are for illustrative purposes only and should not be construed
15 to limit the scope of this invention in any way.

EXAMPLE 1

Exemplary assay workflow

20 This example describes a specific, efficient approach to conducting real-time PCR to detect MTB in a sample. In some embodiments, real-time PCR methods comprise or consist of the following steps:

- 25 1. Inactivation of MTB in samples (e.g., sputum, bronchoalveolar lavage [BAL], and N-acetyl-L-cysteine [NALC] sediments of sputum and BAL) using an inactivation reagent (IR). In some embodiments, the inactivation reagent comprises or consists isopropanol, sodium hydroxide, TWEEN-20, and water;
2. Sample preparation in which DNA is extracted from the inactivated samples using reagents; sample preparation is performed using the automated *m2000sp* instrument
30 (Abbott Molecular), or manually;
3. PCR assembly in which purified samples and assay PCR components are added together in a 96-well optical reaction plate or other multi-chamber reaction support; this is performed using the *m2000sp* or manually;

4. Manual sealing of the 96-well optical reaction plate and transfer of the plate to an *m2000rt* instrument.
5. Amplification and detection of PCR products using the automated *m2000rt* instrument; patient results are automatically reported on the *m2000rt* workstation.

5 A graphical summary of this workflow is shown in Figure 1.

EXAMPLE 2

Target selection and primer/probe design

10 In some embodiments, a dual target strategy is employed for detecting MTB complex. The two targets include: Insertion sequence (IS) 6110 and Protein Antigen B (PAB). See Table 1 below:

Table 1. Abbott RealTime MTB target selection:

<u>IS6110</u>	<p>- <u>Insertion sequence (IS) of the IS3 category</u></p> <ul style="list-style-type: none"> - Usually present in multiple copies per cell (e.g. Denmark, 50% 11-15 copies per cell) - Some TB strains have no or low copy numbers of IS6110
<u>PAB</u>	- <u>Single copy gene coding for protein antigen b</u>

15 Used of a dual target strategy prevents false negative results caused by target sequences mutation or deletion.

Probes and primers that find use in the detection of IS6110 and PAB target sequences include those in Table 2.

20 Table 2. RealTime MTB primer / probe sequences (FP = forward primer; RP = reverse primer; #pdU = 5' propynyl dU; *pdC = 5' propynyl dC; Fam = fluorescein dye; BHQ = Black Hole Quencher; IC = internal control):

Material	SEQ ID NO	Sequence
IS6110 (121) FP	SEQ ID NO:1	5' CCT GCG AGC GTA GGC GTC GGT GA 3'
IS6110 (121) RP	SEQ ID NO:2	5' CGT CCA GCG CCG CTT CGG ACC A 3'
PAB abt2 FPb	SEQ ID NO:3	5' GCA CCT CAA GCT GAA CGG AAA AGT CCT3'

PAB abt2 RPx	SEQ ID NO:4	5' CCG GGG TTG AGC GCA GCG ATC T 3'
IS6110 probe6	SEQ ID NO:5	5' 6-Fam-pdU#AG GpdUG AGG pdUpdC*pdU GpdCpdU ApdCpdC pdC-BHQ1 dT 3'
PAB probe 1	SEQ ID NO:6	5' 6-Fam-pdUApdC pdCAG GGpdC ApdCpdC ApdUpdC AAA-BHQ1 dT 3'
IC FP 196	SEQ ID NO:7	5' CTA CAG CAG AGT TGG CAG CTT CAC TTT C 3'
IC RP 310	SEQ ID NO:8	5' GTC TGG CCT TTC AGC AAG TTT C 3'
Internal Control Probe:	SEQ ID NO:9	5' Quasar-GApdC GAG pdUpdUpdC ApdUG AGG GpdCA-BHQ2 dT 3'

Table 3 provides alternative primers and probes for use in the detection MTB target sequences. In addition to IS6110 and PAB, additional targets include rPOB (single copy gene coding for β subunit of RNA polymerase, site of about 95% of rifampicin-resistance mutations), SenX3-RegXe (single copy gene coding for regulatory proteins), hsp65 (single copy gene coding for heat shock protein), and MPB64 (single copy gene coding for 23 KDA protein).

Table 3. Other primer / probe sequences:

Name	SEQ ID NO	Sequences
IS6110 (104) FP1	SEQ ID NO:10	5' GCCGCTTCGGACCACCAGCACCTAAC
IS6110 (104) RP1	SEQ ID NO:11	5' GTAGGCGTCGGTGACAAAGGCCACGTAG
IS6110 (104) probe	SEQ ID NO:12	5' TGCCCAGGTCGACACATA
IS6110 (80) FP1	SEQ ID NO:13	5' TACGACCACATCAACCGGGAGCCCA
IS6110 (80) RP1	SEQ ID NO:14	5' GCGTGGACGCGGCTGATGTGCTCCT
IS6110 (80) pr1	SEQ ID NO:15	5' CCGCGAGCTGCGCGATG
PAB abt1 FP	SEQ ID NO:16	5'-GCACGCTGCTCTACCCGCTGTTCAACCT
PAB abt1 RP	SEQ ID NO:17	5'-GTGCCCTGAGCGGTGATCGTGACGTT
PAB abt1 Probe:	SEQ ID NO:18	5' TCCGGCCTTTCACGAGA
nhsp65 FP1	SEQ ID NO:19	5' TCGGGGCTCGGGTAGAAGTT
nhsp65 RP1	SEQ ID NO:20	5' TCGTCAACTCGGGCAGCAAC

nhsp65 probe 1	SEQ ID NO:21	5' TACTCGGCTCACGCACG
vhsp65 FP1	SEQ ID NO:22	5' GGCTCGGGTAGAAGTTCGACTTGG
vhsp65 RP1	SEQ ID NO:23	5' GTCAACTCGGGCAGCAACGAC
vhsp65 probe 1	SEQ ID NO:24	5' CTCACGCACGGCGGTATTC
senX3 FP	SEQ ID NO:25	5' GGCAGCGGACTCGGGTT
senX3 RP	SEQ ID NO:26	5' ACCGCAGTTCGGGCTCTC
senX3 Pr	SEQ ID NO:27	5' TCACGACGACGAGCGAC
regX3 FP	SEQ ID NO:28	5' CGCTGATGACCAGTGTGTTGATT
regX3 RP	SEQ ID NO:29	5' GCAGCATCAGATCGAGCAGGAC
regX3 Probe	SEQ ID NO:30	5' ATGGTCCGGCAGCTCTC
MPB64 FP1	SEQ ID NO:31	5' CAACATCAGCCTGCCAGTTACTACC
MPB64 RP1	SEQ ID NO:32	5' CTTCGCGTGGAGTGGACGATG
MPB64 Probe1	SEQ ID NO:33	5' AAGTCGCTGGAAAATTACAT
rPOB FPa	SEQ ID NO:34	5' CGTGGAGGCGATCACACCGCAGACGTT
rPOB RPb	SEQ ID NO:35	5' CGTTGATCAACATCCGGCCGGTGGTC
rPOB probe5	SEQ ID NO:36	5' CGGTCTGTACGTGAGCGTGC

EXAMPLE 3

5 Sample Inactivation

This example describes exemplary reagents and methods for conducting a sample inactivation step.

Preparation of inactivation reagent (IR)

10 Materials Employed:

- Polypropylene or glass container
- 10M NaOH
- Isopropanol

- TWEEN-20
- Purified water

Preparation of IR:

Material Volume Required for 500 mL

- 5 10M NaOH 20 mL
 Purified water 179.1 mL
 Isopropanol 300 mL
 TWEEN-20 0.9 mL

1. Add 179.1 mL of water to an empty polypropylene or glass container (avoid use of a
 10 polystyrene container).
2. Add 0.9 mL of TWEEN-20 to the container.
3. Add 20 mL of 10M NaOH to the container.
4. Add 300 mL of isopropanol to the container.
5. Mix the components by inversion 20 times.
- 15 Use or store at ambient temperature for up to 1 month.

Inactivation procedure:

1. If frozen, thaw specimens at 15 to 30°C.
2. Estimate the volume of specimen to be inactivated.
- 20 3. Add IR at a ratio of 1:3 (e.g., 1 mL specimen + 3 mL IR) (the preferred specimen volume is 0.3 to 10 mL).
4. Invert the container to ensure contact between the IR and the specimen.
5. Vortex the mixture for 20 to 30 seconds.
6. Incubate the mixture at ambient temperature for at least 1 hour and preferably
 25 no more than 24 hours. Vortex the mixture one final time for 20 to 30 seconds at 20 to 30 minutes into the incubation period.

EXAMPLE 4

30 **Sample Preparation Method:**

The MTB assay of Example 1 uses an Abbott automated *m2000sp* instrument or manual method for processing sputum, BAL and NALC-NaOH sediment of sputum or BAL samples and uses an Abbott automated *m2000rt* instrument for amplification and detection. Both processes entail DNA extraction from samples, both DNA purifications are performed

using the DNA GPR (List 6K12-24) sample preparation reagents from the Abbott *mSample Preparation System*_{DNA}.

The sample preparation reagents and method (including lyses step, wash step, elution step, tip reuse arrangement etc.) were optimized to reduce the inhibitory effect on PCR reactions due to the inhibitory sputum or carryover of TB Inactivation reagent (IR): thus centrifugation to get rid of IR in the IR treated sample is not necessary. The procedure is also optimized to reduce carryover from high positives to nearby negative sample. The sample preparation is also optimized to ensure TB cell breakage for efficient DNA recovery and PCR.

Real-time PCR:

After PCR reaction assembly in a 96-well optical reaction plate (either manually or via the *m2000sp*), the 96-well plate is manually sealed and transferred to the *m2000rt* to perform the amplification and real-time fluorescence detection reaction. Patient results are automatically reported on the *m2000rt* workstation. The MTB assay detects an internal control nucleic acid sequence as sample validity control, sample extraction and amplification efficiency control. Table 4 provides exemplary PCR cycling conditions.

Table 4.

Stage	Cycle	Step	Temperature (°C)	Time
1	1	1	50	10 min
2	1	1	94	10 min
3	50	1	94	35 sec
		2	64	15 sec
Read		3	65	40 sec

For the data shown in the below examples, an assay cutoff of 42 was used. That is samples with Ct values < 42 are considered to be MTB Detected, while samples with assay Ct values > 42 are considered MTB Not Detected.

Assays run on the *m2000rt* are per the manufacturer’s recommend protocols. One such example includes the steps of:

1. 96 IR-treated samples are performed per run. One negative control and 1 positive control are included in each run, therefore allowing a maximum of 94 IR-treated samples to be processed per run.

2. Before use, vortex IR-treated samples for 3 to 5 seconds. Using a pipette, transfer the IR-treated samples to the reaction vessels. Minimize the transfer of visible particulates in the IR-treated samples during this step.

3. Thaw assay controls, IC, and amplification reagents at 2 to 8°C or 15 to 30°C. Once thawed, IC can be stored closed at 2 to 8°C for up to 14 days prior to use. Once thawed, controls can be stored at 2 to 8°C for up to 24 hours prior to use. If not using the optional amplification reagent extended use feature: Thaw new amplification reagents at 2 to 8°C or 15 to 30°C. Once thawed, the amplification reagents can be stored at 2 to 8°C for up to 24 hours, prior to use. If using the optional amplification reagent extended use feature: Select new and/or partial amplification reagent packs to be used in the run. Refer to Abbott *m2000sp* Operations Manual (List No. 9K20-06 or higher), Operating Instructions, for instructions pertaining to amplification reagent pack inventory management. Amplification reagent packs should have the same lot number.

4. Vortex each control 3 times for 2 to 3 seconds each time before use. Ensure that bubbles or foam are not created. If found, remove them with a new sterile pipette tip for each tube. Ensure that the contents of each vial are at the bottom after vortexing by tapping the vials on the bench to bring liquid to the bottom of the vial.

5. Gently invert the Abbott *mSample Preparation SystemDNA* bottles to ensure a homogeneous solution. If crystals are observed in any of the reagent bottles upon opening, allow the reagent to equilibrate at room temperature until the crystals disappear. Do not use the reagents until the crystals have dissolved. Ensure bubbles or foam are not generated; if present, remove with a sterile pipette tip, using a new tip for each bottle. NOTE: Before pouring the *mMicroparticlesDNA* into the 200 mL reagent vessel, vigorously mix or vortex until the *mMicroparticlesDNA* are fully resuspended.

6. Vortex the IC vial 3 times for 2 to 3 seconds each time before use. Ensure bubbles or foam are not generated; if present, remove with a sterile pipette tip.

7. Using a calibrated precision pipette dedicated for internal control use only, add 180 µL of IC to 1 bottle of *mLysisDNA* buffer. Mix by gently inverting the container 5 to 10 times to minimize foaming. Each bottle of *mLysisDNA* buffer supports up to 48 sample

preparations. Add 180 μ L of IC to a second bottle of *mLysisDNA* buffer for 49 to 96 samples. If using the optional amplification reagent extended use feature, partial vials of IC can be recapped and stored at 2 to 8°C for a second use.

5 8. Add 25 mL of USP grade 190 to 200 proof ethanol (95 to 100% ethanol) to the *mLysisDNA* buffer + IC reagent bottle. Do not use ethanol that contains denaturants. Gently invert the container to ensure homogeneous solution. For 49 to 96 samples, add 25 mL of ethanol to a second bottle of *mLysisDNA* buffer + IC. Gently invert to ensure a homogeneous solution.

10 9. Add 70 mL USP grade 190 to 200 proof ethanol (95 to 100% ethanol) to *mWash 2DNA* bottle. Do not use ethanol that contains denaturants. Each bottle of *mWash 2DNA* supports up to 48 reactions. Gently invert to ensure a homogeneous solution.

10. Place the negative and positive control and the patient specimens into the Abbott *m2000sp* sample rack.

15 11. Place the 5 mL Reaction Vessels into the Abbott *m2000sp* 1 mL subsystem carrier.

12. Load the carrier racks containing the Abbott *mSample Preparation SystemDNA* reagents and the Abbott 96-Deep-Well Plate on the Abbott *m2000sp* worktable as described in the Abbott *m2000sp* Operations Manual, Operating Instructions.

20 13. From the Run Sample Extraction screen, select and initiate the sample extraction protocol as described in the Abbott *m2000sp* Operations Manual, Operating Instruction.

NOTE: Change gloves before handling the amplification reagents.

25 14. Load the amplification reagent pack and master mix vial (if needed) on the Abbott *m2000sp* worktable after sample preparation is completed. Each amplification reagent pack supports up to 24 reactions. Thaw 1 set of reagents for 1 to 24 samples, 2 sets for 25 to 48 samples, 3 sets for 49 to 72 samples and 4 sets for 73 to 96 samples. Ensure the amplification reagents are thoroughly thawed before use. Ensure that the contents are at the bottom of the vials by tapping the vials in an upright position on the bench. Remove the amplification reagent vial caps. If using the optional amplification reagent extended use feature, a combination of new and partial reagent packs may be used. If not using the optional
30 amplification reagent extended use feature, only new reagent packs may be used. Ensure that the contents of new amplification reagent packs are at the bottom of the vials prior to opening the amplification reagents by tapping the vials in an upright position on the bench. Do not tap partial amplification reagent packs being used a second time. Tapping may result in loss of master mix volume in the cap. Remove caps. If a new amplification reagent pack is stored for

a second use, the vials are recapped for storage. If planning to reuse the original caps to recap the reagent vials, the original caps are saved and used. If planning to use fresh caps to recap the reagent vials, original caps are discarded. Partial amplification packs are loaded to the left of new amplification packs on the Abbott *m2000sp* worktable. Ensure that the amplification reagent packs are firmly seated on the instrument.

15. Select the appropriate deep-well plate from the Run Master Mix Addition screen that matches the corresponding sample preparation extraction. Initiate the Abbott *m2000sp* Master Mix Addition protocol. Follow the instructions as described in the Abbott *m2000sp* Operations Manual, Operating Instructions section. NOTE: The assembly of the amplification master mix and sample eluates into the Abbott 96-Well Optical Reaction Plate (step 15) should be initiated within 1 hour after completion of Sample Preparation. NOTE: The Abbott *m2000rt* protocol (step 20) should be started within 90 minutes of the initiation of the Master Mix Addition protocol. NOTE: If the run is aborted for any reason subsequent to step 15, the amplification reagents are to be discarded and a new 96-well PCR plate should be used if the Abbott *m2000sp* Master Mix Addition Protocol (step 15) will be repeated.

16. Switch on and initialize the Abbott *m2000rt* in the Amplification Area. NOTE: The Abbott *m2000rt* requires 15 minutes to warm up. NOTE: Change laboratory coats and gloves before returning to the sample preparation area.

17. Place the Abbott 96-Well Optical Reaction Plate into the Abbott Splash-Free Support Base after the Abbott *m2000sp* instrument has completed addition of samples and master mix.

18. Seal the Abbott 96-Well Optical Reaction Plate according to the Abbott *m2000sp* Operations Manual, Operating Instructions section. Export the completed PCR plate results to a CD (or directly to a mapped Abbott *m2000rt* via a network connection).

In some embodiments, a manual sample preparation method is employed. An example of such a method is as follows:

1. Thaw amplification reagents at 15 to 30°C or at 2 to 8°C. This step can be initiated before completion of the sample preparation procedure.

2. 12 samples are processed per set of magnetic racks. A negative control and a positive control are included in each run, therefore allowing a maximum of 10 specimens to be processed. Prepare the specimens for processing by following these steps: NOTE: Patient specimens should be inactivated prior to beginning sample extraction.

3. Thaw 1 tube of the MTB Negative Control, 1 tube of MTB Positive Control, and 1 vial of MTB Internal Control at 15 to 30°C or at 2 to 8°C. Once thawed, if IC is not being

processed immediately, store at 2 to 8°C for up to 14 days prior to use. Once thawed, if controls are not being processed immediately, store at 2 to 8°C for up to 24 hours prior to use. Vortex controls and IC 3 times for 2 to 3 seconds each time before use. Ensure that the contents of each vial are at the bottom after vortexing by tapping the vials on the bench to
5 bring liquid to the bottom of the vial. Ensure bubbles or foam are not generated; if present, remove with a sterile pipette tip, using a new tip for each vial.

4. Open the Abbott *mSample Preparation SystemDNA* reagent pack(s). If crystals are observed in any of the reagent bottles upon opening, allow the reagent to equilibrate at room temperature until the crystals disappear. Do not use the reagents until the crystals have
10 dissolved.

5. Prepare the *mWash 2DNA* by adding 70 mL of USP grade 190 to 200 proof ethanol (95 to 100% ethanol) to the *mWash 2DNA* bottle. Do not use ethanol that contains denaturants. Gently invert to ensure a homogeneous solution. NOTE: Mark the *mWash 2DNA* bottle to indicate that ethanol has already been added for extended use.

6. Prepare the *mLysisDNA* by adding 25 mL of USP grade 190 to 200 proof ethanol (95 to 100% ethanol) to the *mLysisDNA* bottle. Do not use ethanol that contains denaturants. Gently invert 5 to 10 times to mix and to minimize foaming. NOTE: Mark the *mLysisDNA* bottle to indicate that ethanol has already been added for extended use.

7. Calculate the volume of *mLysisDNA* solution required for the manual run: (1.85 mL of *mLysisDNA* × number of samples). Pipette the required volume of *mLysisDNA* solution into a polypropylene container large enough to hold the entire volume. Calculate the volume of IC required for the manual run: (3.51 µL of IC × number of samples). Use a precision pipette dedicated to internal control use only to add the required volume of IC into the polypropylene container containing the *mLysisDNA* solution required for the manual run.
20
25 Mix *mLysisDNA* solution and IC mixture by gentle inversion 10 to 15 times to minimize foaming. After initial use, partial IC vials maybe stored at 2 to 8°C for up to 14 days and used 1 additional time.

8. Gently invert all the reagent bottles, except the *mMicroparticlesDNA* bottle and the *mWash 1DNA* bottle, 5 to 10 times to ensure a homogenous solution prior to use. The
30 *mMicroparticlesDNA* bottle will be mixed in step 11.

9. Turn on the temperature-controlled dry heating blocks. Set the first block to 58°C. Set the second block to 80°C. NOTE: Check the temperature of the heating blocks. Do not proceed until the heating blocks are at the correct temperature.

10. Label all necessary tubes: One 5 mL reaction vessel per sample for the Lysis Incubation and *mWash 1DNA* steps. One 1.5 mL microfuge tube per sample for the first and second *mWash 2DNA* and Elution steps. One 1.5 mL microfuge tube per sample or a 96-well polypropylene plate for the eluate.

5 11. Place the labeled 5 mL reaction vessels for each sample in unheated stand. Resuspend *mMicroparticlesDNA* by vortexing or vigorously shaking until particles are in suspension and settled particles are no longer seen on the bottom of the bottle. After the particles are resuspended, use a precision pipettor and a sterile 200 μ L aerosol barrier pipette tip to add 50 μ L of *mMicroparticlesDNA* to each reaction vessel.

10 12. Using a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, add 1.75 mL (2 X 875 μ L) of *mLysisDNA* to the reaction vessels.

15 13. Add 0.8 mL of the controls, and specimens to the appropriate reaction vessels using a precision pipettor and a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample. Mix each sample/*mLysisDNA* mixture by aspirating and dispensing the 800 μ L volume 5 to 10 times until a uniform suspension is obtained. NOTE: Aspirate and dispense liquid slowly to avoid foaming.

14. Transfer the 5 mL reaction vessels into the 58°C heating block.

15. Start the timer and incubate for 15 minutes.

20 16. After incubation using a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, mix the mixture 5 times by aspirating and dispensing 800 μ L.

17. Start the timer and incubate for an additional 10 minutes in the 58°C heating block.

18. After incubation using a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, mix the mixture 5 times by aspirating and dispensing 800 μ L.

25 19. Start the timer and incubate for an additional 10 minutes in the 58°C heating block.

20. After the incubation is complete, place the reaction vessels in a magnetic capture stand for 2 minutes to allow the particles to be captured on the side of the reaction vessels.

30 21. With the reaction vessels in the magnetic capture stand, use a fresh, sterile 1000 μ L aerosol barrier pipette tip or disposable transfer pipette for each sample to carefully remove the *mLysisDNA* from each reaction vessel and discard the fluid into a liquid waste container. Remove the fluid as completely as possible. Do not disturb or aspirate the captured magnetic particles.

22. Remove the reaction vessels from the magnetic rack and transfer to a nonmagnetic rack. *mWash 1DNA* (Wash).

23. Using a precision pipettor and a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, add 800 μ L of *mWash 1DNA* to the samples and resuspend the magnetic
5 particles in the wash fluid by gently mixing 10 times by aspiration and dispense with a pipette tip. Wash the particles from the side of the reaction vessel, if necessary. NOTE: When adding *mWash 1DNA* wash, dispense liquid slowly to avoid splashing.

24. Transfer the wash fluid and particles to a labeled 1.5 mL microfuge tube.

25. Place the tubes in a magnetic capture stand for 1 minute to allow the particles to
10 be captured on the side of the tubes.

26. With the tubes in the magnetic capture stand, use a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample to carefully remove the *mWash 1DNA* from each tube and discard fluid into a liquid waste container. Remove the fluid as completely as possible. DO NOT disturb or aspirate the captured magnetic particles.

15 27. Remove the tubes from the magnetic rack and transfer to a nonmagnetic rack. *mWash 2DNA* (First Wash).

28. Using a precision pipettor and a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, add 800 μ L of *mWash 2DNA* to the samples and resuspend the magnetic
20 particles in the wash fluid by gently mixing 5 to 10 times by aspiration and dispense with a pipette tip. Wash the particles from the side of the tube, if necessary. NOTE: When adding *mWash 2DNA* wash, dispense liquid slowly to avoid splashing.

29. Place the tubes in a magnetic capture stand for 1 minute to allow the particles to be captured on the side of the tubes.

30 25 30. With the tubes in the magnetic capture stand, use a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample to carefully remove the *mWash 2DNA* from each tube and discard fluid into a liquid waste container. Remove the fluid as completely as possible. DO NOT disturb or aspirate the captured magnetic particles.

31. Remove the tubes from the magnetic rack and transfer to a nonmagnetic rack. *mWash 2DNA* (Second Wash).

30 32. Using a precision pipettor and a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, add 800 μ L of *mWash 2DNA* to the samples and resuspend the magnetic particles in the wash fluid by gently mixing 5 to 10 times by aspiration and dispense with a pipette tip. Wash the particles from the side of the tube, if necessary. NOTE: When adding *mWash 2DNA* wash, dispense liquid slowly to avoid splashing.

33. Place the tubes in a magnetic capture stand for 1 minute to allow the particles to be captured on the side of the tubes.

34. With the tubes in the magnetic capture stand, use a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample to carefully remove the *mWash 2DNA* from each tube and discard fluid into a liquid waste container. Remove the fluid as completely as possible. DO NOT disturb or aspirate the captured magnetic particles.

35. Remove the tubes from the magnetic rack and transfer to the 80°C heating block and incubate for 15 minutes with caps open to allow for the evaporation of the ethanol.

36. Using a precision pipettor and a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, add 250 μ L of *mElution BufferDNA* to the samples and resuspend the magnetic particles in the fluid by aspiration and dispense with the pipette tip. Wash the particles from the side of the tube, if necessary.

37. Place the tubes in the 80°C heating block, start the timer, and incubate for 4 minutes.

38. Remove the tubes from the 80°C heating block. Using a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample, mix the sample and *mElution BufferDNA* mixture 4 times by aspirating and dispensing 200 μ L.

39. Return the tubes to the 80°C heating block. Start the timer and incubate for 4 minutes.

40. Remove the tubes from the 80°C heating block and place in a magnetic capture stand for 1 minute to allow the particles to be captured on the side of the tubes.

41. With the tubes in the magnetic capture stand, use a fresh, sterile 1000 μ L aerosol barrier pipette tip for each sample to carefully remove the eluted sample from the tubes. Do not disturb or aspirate the captured microparticles. The eluted sample(s) can be placed into a fresh, labeled 1.5 mL microfuge tube or a 96-well polypropylene plate. NOTE: The assembly of the amplification master mix and sample eluates into the Abbott 96-Well Optical Reaction Plate (step 48) must be initiated within 1 hour after completion of Sample Preparation.

42. Switch on and initialize the Abbott *m2000rt* instrument. NOTE: The Abbott *m2000rt* requires 15 minutes to warm up.

43. Create the Abbott *m2000rt* test order. Refer to the Operating Instructions section of the Abbott *m2000rt* Operations Manual. From the Protocol screen, select the Abbott RealTime MTB assay application protocol. NOTE: Remove gloves before returning to the reagent preparation area.

44. Prepare the amplification master mix. NOTE: All reagent preparation should take place in the dedicated Reagent Preparation Area. Change gloves before handling the amplification reagents. Do not vortex or invert the amplification reagent pack. Each amplification reagent pack supports up to 24 reactions. Ensure the amplification reagents are thoroughly thawed before use. Prior to opening the amplification reagents, ensure that the contents of the amplification reagent pack are at the bottom by tapping the amplification reagent pack in an upright position on the bench to bring the liquid to the bottom of the vials. Identify the amplification reagents as follows: Activation Reagent (Reagent 1); MTB Amplification Reagent (Reagent 2); DNA Polymerase (Reagent 3); Remove and discard caps.
- 5
- 10 Using a calibrated precision pipette dedicated for reagent use only, add 298 μL of Activation Reagent (Reagent 1) and 418 μL of MTB Amplification Reagent (Reagent 2) to the DNA Polymerase bottle (Reagent 3) to make master mix. Mix by gently pipetting up and down 5 times. Avoid creating foam.
45. Pipette the contents of the master mix from the DNA Polymerase bottle into a 1.5 mL microfuge tube (List No. 4J71-50 or equivalent). Mix by gently pipetting up and down 5 times. Avoid creating foam.
- 15
46. Place an Abbott 96-Well Optical Reaction Plate in the Abbott Splash-Free Support Base to prevent contamination. Contamination of the bottom of the Abbott 96-Well Optical Reaction Plate with fluorescent materials could potentially interfere with the MTB assay. The Abbott 96-Well Optical Reaction Plate should be held and transported with the Abbott Splash-Free Support Base to minimize contamination.
- 20
47. Using a precision pipette dedicated for reagent use only, dispense 25 μL aliquots of the amplification master mix into each well of the Abbott 96-Well Optical Reaction Plate that will be used to run the samples and controls. A calibrated repeat pipettor may be used.
- 25 Add the master mix in an order starting with column 1 (from top to bottom), and moving to each consecutive column from left to right. Visually verify that 25 μL has been dispensed into each well. Transfer the Abbott 96-Well Optical Reaction Plate in the Abbott Splash-Free Support Base to the Sample Preparation Area.
48. Using a precision pipettor and a fresh, sterile 200 μL aerosol barrier pipette tip for each sample, transfer 25 μL of each eluted sample to the Abbott 96-Well Optical Reaction Plate. During the transfer of each sample, mix the final reaction by pipetting up and down 3 to 5 times. Visually verify that a total of 50 μL has been dispensed into each well.
- 30

49. Seal the Abbott 96-Well Optical Reaction Plate according to the instructions in the Abbott *m2000rt* Operations Manual, Operating Instructions section.

50. Centrifuge the Abbott 96-Well Optical Reaction Plate in the Abbott Splash-Free Support Base at 5000g for 5 minutes.

5 51. Transfer the Abbott 96-Well Optical Reaction Plate in the Abbott Splash-Free Support Base to the Amplification Area. NOTE: The Abbott *m2000rt* protocol (step 52) should be started within 90 minutes following the initiation of the master mix addition and PCR plate preparation (step 44).

10 52. Place the Abbott 96-Well Optical Reaction Plate in the Abbott *m2000rt* instrument, select the test order created (step 43), and initiate the Abbott RealTime MTB assay application protocol, as described in the Abbott *m2000rt* Operations Manual, Operating Instructions section. At the completion of the run, assay results are reported on the Abbott *m2000rt*.

15 **EXAMPLE 5**

Experimental data – Inactivation

The IR TB killing effectiveness was evaluated. In this experiment, MTB-containing samples (cultured MTB that was diluted to known MTB concentrations prior to inactivation, 20 as well as MTB-containing NALC-NaOH sediments) were subjected to the inactivation procedure of Example 3. Following inactivation, excess Inactivation Reagent was removed by centrifugation/washing and the surviving cells were placed into MGIT culture for up to 42 days or six weeks. This duration is the recommended longest time for MTB culture; most MTB positive specimens will result in detectable culture growth within 20 days of initiation 25 of culture). Table 5 below illustrates the results obtained when testing the cultured samples following inactivation. The Positive Control (PC), which consists of non-inactivated MTB, demonstrated growth within the expected 20-day timeframe, while the Negative Control (NC), showed no growth.

Table 5. Inactivation of cultured high concentration MTB cells.

Reduction of MTB Infection Risk Study Summary						
Study	Number of Isolate Samples	CFU/mL	Positive Samples Without IR Treatment		Positive Samples With IR Treatment	
			Number Samples and Replicates Tested	Culture Positive	Number Samples and Replicates Tested	Culture Positive
1	3 ^a	1 x 10 ⁸	3 x 1	3 of 3	3 x 3	0 of 9
		1 x 10 ⁷	3 x 1	3 of 3	3 x 3	0 of 9
		1 x 10 ⁶	3 x 1	3 of 3	3 x 3	0 of 9
2	20 ^b	N/A	20 x 1	20 of 20	20 x 1	0 of 20
3	31 ^b	N/A	31 x 1	31 of 31	31 x 1	0 of 31

^a Cultured Isolate Samples

^b Clinical Isolate Samples

5 These data demonstrate the effectiveness of the inactivation procedure for inactivation of MTB.

EXAMPLE 6

Analytical inclusivity

10 Eight subspecies and 20 samples of MTB complex (*M. tuberculosis*, *M. africanum*, *M. bovis*, *M. bovis BCG*, *M. canettii*, *M. microti*, *M. caprae*, *M. pinnipedii*.) were obtained from ATCC (*M. canettii* was received from the Public Health Research Institute) and were tested from 10 to 100 genomic DNA copies/reaction (See Table 6). All 8 subspecies were detected at both levels.

Table 6. MTB complex subspecies tested

Name	Name
Mycobacterium tuberculosis 25177D-5 (H37Ra)	M. pinnipedii BAA-688D
M. tuberculosis 25618D-5 (H37Rv)	M. bovis BCG 35747D
M. microti11152	M. caprae BAA-824D
M. microti 19422	M. tuberculosis BAA-2238D
M. africanum 25420	M. tuberculosis BAA-2237D
M. africanum 35711	M. tuberculosis 27294D
M. bovis 35735	M. tuberculosis BAA-2234D
M. bovis 19274	M. tuberculosis 35822D
M. bovis BCG 35746	M. tuberculosis 35838D
M. canettii	M. tuberculosis BAA-2235D

Forty six phylogenetically and geographically diverse MTB isolate DNAs (> 50% with MDR) obtained from the Public Health Research Institute were tested from 25 to 100 genomic DNA copies/reaction (Figure 2). All subspecies tested were detected.

EXAMPLE 7

Analytical specificity

Purified nucleic acid from different mycobacteria, viruses and other microorganisms (n=80) at targeted concentrations of 1e5 to 1e7 genomes/mL and cultured microorganisms at 1 x 10⁶ cfu/mL were added to MTB negative control to evaluate the effect of potential cross-reactants on MTB assay results for MTB negative specimens. Purified nucleic acid from different mycobacteria, viruses and other microorganisms at targeted concentrations of 1 x 10⁶ to 1 x 10⁷ genomes per milliliter and cultured microorganisms at 1e6 cfu/mL were added to MTB positive samples to evaluate the effect of potential cross-reactants on MTB assay results for MTB positive specimens. MTB positive samples were prepared by diluting heat inactivated MTB cell stock in negative control to a targeted concentration of 1000 copies/mL (quantitated using a genomic DNA curve). None of the MTB negative samples tested with the potential cross-reactants was detected. All 80 MTB positive samples tested with potential cross-reactants were detected.

Table 7
Microorganisms tested to determine analytical specificity

Species		
<i>Mycobacterium abscessus</i>	<i>Mycobacterium sphagni</i>	<i>Escherichia coli</i>
<i>Mycobacterium austroafricanum</i>	<i>Mycobacterium terrae</i>	Herpes simplex virus 1

<i>Mycobacterium avium</i>	<i>M. thermoresistibile</i>	<i>Klebsiella pneumoniae</i>
<i>Mycobacterium avium ssp. avium</i>	<i>Mycobacterium tokaiense</i>	<i>Lactobacillus delbrueckii</i>
<i>Mycobacterium avium ssp.</i>	<i>Mycobacterium ulcerans</i>	<i>Legionella pneumophila</i>
<i>Mycobacterium celatum</i>	<i>Mycobacterium vaccae</i>	<i>Neisseria gonorrhoeae</i>
<i>Mycobacterium chelonae</i>	<i>Mycobacterium xenopi</i>	<i>Neisseria meningitidis</i>
<i>Mycobacterium chitae</i>	<i>Acinetobacter baumannii</i>	<i>Porphyromonas gingivalis</i>
<i>Mycobacterium fallax</i>	<i>Aeromonas hydrophila</i>	<i>Proteus mirabilis</i>
<i>Mycobacterium flavescens</i>	<i>Bacillus cereus</i>	<i>Pseudomonas aeruginosa</i>
<i>Mycobacterium fortuitum</i>	<i>Bacillus subtilis</i>	<i>Salmonella choleraesuis</i>
<i>Mycobacterium gastri</i>	<i>Bordetella parapertussis</i>	<i>Serratia marcescens</i>
<i>Mycobacterium gordonae</i>	<i>Campylobacter jejuni</i>	<i>Staphylococcus aureus</i>
<i>Mycobacterium intracellulare</i>	<i>Candida albicans</i>	<i>Staphylococcus epidermidis</i>
<i>Mycobacterium kansasii</i>	<i>Chromobacterium violaceum</i>	<i>Staphylococcus haemolyticus</i>
<i>Mycobacterium lentiflavum</i>	<i>Chlamydia pneumoniae</i>	<i>Staphylococcus hominis</i>
<i>Mycobacterium marinum</i>	<i>Chlamydia trachomatis</i>	<i>Streptococcus agalactiae</i>
<i>Mycobacterium neoaurum</i>	<i>Citrobacter freundii</i>	<i>Streptococcus gordonae</i>
<i>Mycobacterium</i>	<i>Corynebacterium diphtheriae</i>	<i>Streptococcus mitis</i>
<i>Mycobacterium phlei</i>	<i>Corynebacterium xerosis</i>	<i>Streptococcus mutans</i>
<i>Mycobacterium pneumoniae</i>	<i>Cryptococcus neoformans</i>	<i>Streptococcus pneumoniae</i>
<i>Mycobacterium pulveris</i>	Cytomegalovirus	<i>Streptococcus pyogenes</i>
<i>Mycobacterium scrofulaceum</i>	<i>Enterobacter aerogenes</i>	<i>Streptomyces griseinus</i>
<i>Mycobacterium shimoidei</i>	<i>Enterobacter cloacae</i>	Varicella-zoster virus
<i>Mycobacterium simiae</i>	<i>Enterococcus faecalis</i>	<i>Veillonella parvula</i>
<i>Mycobacterium smegmatis</i>	<i>Enterococcus avium</i>	

EXAMPLE 8

Analytical sensitivity

5 A MTB panel, strain H37Rv at 40 cfu/mL was serially diluted in pooled MTB negative sputum to generate a sensitivity panel. Sixteen replicates of each dilution were tested. A detection rate of 100% was observed at all dilutions 160 fold and lower. Results are shown in Table 8.

10 Table 8. Analytical sensitivity determined by testing serial dilutions of a MTB panel.

Series dilution		AM RealTime MTB	
		Ct cut off 42	Ct cut off 39
<i>from 40 cfu/ml</i>	<i>Dil. Fold</i>	<u>Hit Rate</u>	<u>Hit Rate</u>
1	0	16/16	16/16
2	2	16/16	16/16
3	4	16/16	16/16
4	8	16/16	16/16
5	20	16/16	16/16
6	40	16/16	12/16
7	80	16/16	15/16
8	160	13/16	10/16
9	320	15/16	7/16
10	640	8/16	4/16
11	1276	4/16	0/16
12	2564	3/16	0/16
13	5128	1/16	0/16

EXAMPLE 9

Clinical specificity

5 Culture-negative NALC samples (n=155), sputum (n=23) and BAL (n=28) samples (NALC samples were from MTB suspect population. Sputum and BAL samples were from patients with no TB symptoms) were tested to determine clinical specificity (see data summarized in table 10 below). Specificity for sputum and BAL samples was 100%. Specificity for NALC samples was 98.7% with an overall specificity of 99%.

10

Table 9. Clinical specificity determined by testing TB negative samples.

<u>Specimen type</u>	<u>Specificity definition</u>	<u>Tested Numbers</u>	<u>Negative results</u>	<u>Positive results</u>	<u>Specificity</u>
NALC	TB suspect Culture negative	155	153	2	98.7%
Sputum	No TB symptom	23	23	0	100%
BAL	No TB symptom	28	28	0	100%

Total		206	204	2	99%
-------	--	-----	-----	---	-----

Table 10. TB Culture positive samples (including both smear positives and negatives) were tested by MTB assay (AM) vs. another comparator assay (Comparator).

TUBE #	SPECIMEN TYPE	CULTURE RESULTS	FAM Ct	MR	CY5 Ct	MR	AM	Comparator
39	SPUTUM	TB	15.91	0.218	34.50	0.166	Detected	high
38	SPUTUM	TB	16.76	0.225	34.55	0.176	Detected	high
35	SPUTUM	TB	19.88	0.238	33.34	0.175	Detected	high
23	SPUTUM	TB	20.77	0.233	33.21	0.169	Detected	med
30	BAL	TB	21.79	0.226	33.18	0.177	Detected	high
25	SPUTUM	TB	22.28	0.222	33.54	0.171	Detected	error
61	SPUTUM	TB	22.84	0.227	33.96	0.179	Detected	med
45	SPUTUM	TB	23.67	0.227	33.13	0.177	Detected	high
44	ASPIRATE	TB	25.06	0.243	33.25	0.181	Detected	high
34	SPUTUM	TB	25.50	0.230	33.08	0.184	Detected	med
63	SPUTUM	TB	27.55	0.245	32.99	0.141	Detected	low
21	SPUTUM	TB	27.61	0.229	33.54	0.187	Detected	med
33	ASPIRATE	TB	27.63	0.247	33.31	0.182	Detected	med
26	BAL	TB	28.05	0.248	33.70	0.187	Detected	med
58	SPUTUM	TB	28.26	0.215	33.88	0.177	Detected	med
32	SPUTUM	TB	29.26	0.225	36.49	0.177	Detected	low
28	SPUTUM	TB	29.43	0.241	33.15	0.197	Detected	low
42	SPUTUM	TB	29.82	0.228	35.19	0.184	Detected	med
60	SPUTUM	TB	30.14	0.222	34.24	0.179	Detected	low
59	SPUTUM	TB	30.21	0.229	34.70	0.185	Detected	low
29	SPUTUM	TB	30.68	0.229	34.19	0.188	Detected	low
57	SPUTUM	TB	31.31	0.231	34.40	0.192	Detected	low
24	SPUTUM	TB	31.70	0.239	35.37	0.191	Detected	low
31	SPUTUM	TB	32.73	0.189	-1	0.008	Detected	low
62	SPUTUM	TB	33.12	0.237	34.52	0.185	Detected	low
27	BRONCHIA WASH	TB	35.00	0.232	34.55	0.185	Detected	Not det
55	LUNG TISSUE	TB	36.67	0.212	34.76	0.176	Detected	not tested*
53	LUNG TISSUE	TB	37.71	0.277	35.94	0.177	Detected	Not det
51	SPUTUM	TB	38.19	0.142	-1	0.003	Detected	not tested*
46	SPUTUM	TB	38.58	0.161	34.58	0.185	Detected	Not det
22	BAL	TB	-1	0.003	34.05	0.174	Not det	Not det

36	SPUTUM	TB	-1	0.006	34.60	0.191	Not det	Not det
37	SPUTUM	TB	-1	0.004	35.90	0.178	Not det	Not det
40	SPUTUM	TB	-1	0.004	34.42	0.178	Not det	Not det
41	SPUTUM	TB	-1	0.002	34.79	0.182	Not det	Not det
43	SPUTUM	TB	-1	0.004	34.97	0.165	Not det	Not det
48	BAL	TB	-1	0.005	34.97	0.165	Not det	Not det
49	SPUTUM	TB	-1	0.001	34.58	0.168	Not det	Not det
50	SPUTUM	TB	-1	0.005	35.14	0.184	Not det	Not det
54	SPUTUM	TB	-1	0.006	34.64	0.173	Not det	Not det

* “not tested” sample was because of not enough volume

5 The RealTime MTB showed better sensitivity at low end samples comparing to the comparator’s assay.

EXAMPLE 10

Analytical and Clinical Performance of MTB Assay

10 This Example describes the analytical performance of the real time MTB detection assay.

Materials and methods

The work flow for the real time MTB assay is described in Figure 1.

15 *Sample inactivation*

500 mL of Inactivation reagent (IR) was prepared by combining the following components: 20 mL 10 M NaOH, 300 mL isopropanol, 0.9 mL Tween-20, and 179.1 mL purified water. Once prepared the IR was stable for up to one month at room temperature. If frozen, specimens (unprocessed specimens or processed NALCsediments) were thawed at 20 15° to 30°C. Approximately three volumes of IR were added to each volume of sample (the minimum allowable specimen volume is 0.3 mL). The same volume ratio of sample: IR was maintained notwithstanding the type of sample (unprocessed or NALC sediment). The mixture was vortexed twice for 20 to 30 seconds each during the first hour of room temperature incubation. The validated incubation time was one to 24 hours. The inactivation 25 process occurred under a biohood. Once completed, the inactivated samples were removed from under a biohood and then subjected to sample preparation outside of the biohood. The

inactivation process was demonstrated to effectively reduce MTB viability at three different laboratories using cultured MTB added to NALC sediments of sputum, MTB positive clinical NALC sediments, and MTB smear /culture positive sputum samples (Qi C., et al., Effectiveness of the sample inactivation procedure employed by the new Abbott RealTime assay for the detection of Mycobacterium tuberculosis, 24th European Congress of Clinical Microbiology and Infectious Diseases (ECCMID) 2014).

Sample preparation

IR-treated specimens and the assay controls were loaded onto an *m2000sp* instrument where DNA was isolated using guanidinium thiocyanate-magnetic microparticle technology to capture nucleic acids followed by washes to remove unbound components. An Internal Control (IC) was added at the start of sample preparation. The bound nucleic acids were eluted and transferred to a 96 deep-well plate. At the completion of sample preparation, the *m2000sp* was used to create an amplification master mix consisting of AmpliTaq Gold Polymerase, a magnesium chloride activation reagent, and oligonucleotide reagent containing primers, probes and dNTPs. The *m2000sp* was used to dispense 25 µl aliquots of the master mix followed by 25 µl aliquots of the extracted eluates to a 96-well optical reaction plate. The plate was sealed manually and transferred to the *m2000rt* for realtime PCR. As an alternative to the *m2000sp*, sample preparation, mastermix preparation, and PCR plate set-up can be performed manually.

Amplification and detection

The *m2000rt* instrument was used for amplification and realtime fluorescence detection. The detection of MTB complex members (Warren RM, et al., Int J Tuberc Lung Dis 2006;10:818–822) was achieved through the use of two sets of primers; one targeting the insertion element IS6110 (Thierry D, et al., Nucleic Acids Res 1990;18:188) and one the PAB gene (Anderson AB, Hansen EB Infect Immun 1989;57:2481-2488). Signal for MTB complex detection was generated with the use of fluorescent labelled probes. The MTB dual target probes are each labeled with the fluorophore FAM at the 5' end and the Black Hole Quencher (BHQ1) at the 3' end. Thus, MTB signals from both IS6110 and PAB are detected in the same FAM channel. The amplification cycle at which FAM fluorescent signal is detected is proportional to the log of the MTB DNA concentration present in the original sample. The probe for internal control (IC) is labelled with Quasar at the 5' and Black Hole

Quencher BHQ2 at the 3' end to allow IC and target signals to be distinguishable in a single PCR well.

Assay controls

5 A minimum of one replicate of the Negative Control and one replicate of the Positive Control were used to determine run validity. The Negative Control consisted of TE buffer and preservatives. The Positive Control consisted of plasmid DNA containing both the IS6110 and PAB target sequences diluted in TE buffer with 1.5 g/mL of poly dA:dT and preservatives. The IC consisted of plasmid DNA containing a pumpkin hydroxypyruvate reductase (HPR) sequence insert diluted in TE buffer with 1.5 g/mL of poly dA:dT and
10 preservatives. IC was added at the start of sample preparation, serving as a control for sample preparation recovery, sample inhibition, and amplification efficiency. The IC did not control for the inactivation procedure. The IC threshold cycle (Ct) value difference between each sample and the run controls was used to assess the validity of each sample result.

15

Panels and clinical specimens

MTB complex subspecies: Nineteen MTB complex subspecies DNA samples were obtained from the American Type Culture Collection (ATCC, Manassas, VA) and one (*M. canettii*) was provided kindly by Ibis Biosciences (Carlsbad, CA). A total of 20 MTB
20 complex strains was tested including *M. africanum* 25420, *M. africanum* 35711, *M. bovis* 35735, *M. bovis* 19274, *M. bovis* BCG 35746, *M. bovis* BCG 35747D, *M. canettii*, *M. caprae* BAA-824D, *M. microti* 11152, *M. microti* 19422, *M. pinnipedii* BAA-688D, MTB 25177D-5 (H37Ra), MTB 25618D-5 (H37Rv), MTB BAA-2236D, MTB BAA-2237D, MTB 27294D, MTB BAA-2234D, MTB 35822D, MTB 35838D, MTB BAA-2235D. Additionally, 46
25 strains of the MTB subspecies including the three principal genetic groups and nine genetic clusters were obtained from Dr. Barry Kreiswirth at the University of Medicine and Dentistry New Jersey (Newark, NJ) (Mathema B, et al., Current Insights, Clinical Microbiology Reviews 2006;19:658-685). The DNA of the 20 MTB complex subspecies obtained from ATCC and Ibis were directly tested using reported DNA concentrations as determined by the
30 PicoGreen® NanoDrop method. The other 46 DNA concentrations were determined using PicoGreen® NanoDrop measurements at Abbott Molecular with the exception of three samples where such measurements could not be obtained due to low volume and impurities. These three samples were diluted at a sample to water ratio of 1:600 and tested directly.

Limit of detection [LOD]: An MTB H37Rv panel targeted to 1×10^5 colony forming units (cfu)/mL was prepared by Zeptomatrix (Buffalo, NY). Three one mL aliquots of the Zeptomatrix panel were combined and centrifuged at $3,000 \times g$ for 15 minutes to remove free MTB DNA in the supernatant. The cell pellet was resuspended in three mL of TE buffer to
5 maintain the concentration of 1×10^5 cfu/mL. The cells were then added to a pool of sputum, which was homogenized using bead-beating, to make the following MTB-containing dilution panels: 80 cfu/mL, 50 cfu/mL, 25 cfu/mL, 10 cfu/mL, 5 cfu/mL, 1 cfu/mL, 0.50 cfu/mL, 0.10 cfu/mL, and 0.05 cfu/mL.

Analytical specificity: Analytical specificity panel members were collected as
10 follows: Cytomegalovirus, Herpes Simplex virus 1, and Varicella-zoster virus were obtained from Advanced Biotechnology Inc. (Columbia, MD), 69 mycobacterial and other microorganism species were obtained from ATCC, and eight bacterial isolates were cultured at Abbott Molecular.

Potentially interfering substances: The following materials were obtained for this
15 testing: blood, DNA from human cells, gastric acid, hypertonic saline, physiologic saline, culture media, NALC pellet material, five anti-TB medications (Isoniazid, Rifampicin, Streptomycin, Pyrazinamide, Ethambutol), and bovine mucus.

Carryover: Two samples were prepared: a high positive MTB sample containing 1×10^7 copies/mL of a plasmid containing the assay target sequences and a negative sample.

Reproducibility: Two samples were prepared: a positive sample containing an MTB
20 concentration of ~ three times the claimed assay LOD and a negative sample.

Clinical specimens: 198 sputum specimens were collected by Discovery Life
25 Sciences (Los Osos, CA) from TB suspect patients in Russia, South Africa, Uganda, and Vietnam. 150 sputum specimens from Vietnam were obtained from the specimen bank operated by the Foundation for Innovative New Diagnostics (FIND) (Geneva, Switzerland). 234 NALC specimens were obtained from Northwestern University Memorial Hospital (Chicago, IL). All patient specimens were collected under ethical guidelines. The HIV status of the patients was not determined. For all specimens smear (when available) and culture testing was performed near the collection site, while Abbott RealTime MTB assay testing was
30 performed at Abbott Molecular.

Results

MTB complex subspecies detection

This study was conducted to determine whether the specific primers and probes used in MTB assays would detect the following eight MTB complex sub-species: *M. africanum*, *M. bovis*, *M. bovis* BCG, *M. canettii*, *M. caprae*, *M. microti*, *M. pinnipedii*, and *M. tuberculosis*. Two sets of purified MTB complex DNAs were tested. The first set of 20 purified DNAs contained representatives of the previously mentioned MTB sub-species. Each purified DNA was tested at two concentrations (100 and 10 genomes/reaction) with four replicates being tested per concentration. At the 100 MTB genomes per reaction level, all four replicates of each of the 20 MTB strains were detected. At the 10 MTB genomes per reaction level, all four replicates of 17 of the strains were detected. For three strains (two *M. bovis* and one *M. bovis* BCG) two of the four replicates were detected (Figure 4). The second set of 46 MTB strains from the MTB sub-species were tested at two concentrations: 100 genomes/reaction and 25 genomes/reaction. Four replicates of each DNA were tested at each concentration. All the tested replicates were positive at both concentrations (Figure 2).

15 *Limit of Detection (LOD)*

A nine-level dilution series was made from MTB strain H37Rv cells diluted in a glass bead homogenized sputum pool. The panel members in the dilution series were targeted to the following concentrations: 80 cfu/mL, 50 cfu/mL, 25 cfu/mL, 10 cfu/mL, 5 cfu/mL, 1 cfu/mL, 0.50 cfu/mL, 0.10 cfu/mL, and 0.05 cfu/mL. Twenty replicates of each panel member were tested across four runs using the Abbott RealTime MTB assay. The study was conducted using one lot of MTB assay and control reagents. The significance level for this study was 0.05. The detection rate was calculated for each target concentration (Table 11). A Probit regression model was fitted, based on the target concentrations and the detection rate using PROC PROBIT in SAS, with the target concentration (X) as the independent variable and the detection rate P (Y=1) as the response variable. The Probit analysis of the data determined that the concentration of MTB detected with 95% probability was 2.45 cfu/mL (95% CI 1.44 – 6.10 cfu/mL). The claimed analytical sensitivity of the Abbott RealTime MTB assay is 17 cfu/mL in pooled homogenized sputum using the MTB H37Rv strain.

30 *Analytical specificity*

Each of the 80 potential cross-reactants was tested in both an MTB positive sample and an MTB negative sample. Nucleic acid from each potentially cross-reacting mycobacterium, virus, or other microorganism at a targeted concentration of 1×10^5 to 1×10^7 copies or genomes per mL was added to the MTB positive samples (containing 1,000

MTB genomes/mL) and the MTB negative samples. Cultured microorganisms at a target concentration of 1×10^6 cfu/mL were added to the MTB positive samples and the MTB negative samples. The assay results for all 80 negative samples were reported as “MTB Not Detected”. The assay results for all 80 MTB-containing samples were reported as “MTB
5 Detected” (Table 7).

Potentially interfering substances

The potential for interference in the test results was assessed with substances that may be present in the respiratory system. MTB negative and MTB positive (500 copies/mL)
10 samples were tested in the absence or the presence of each potentially interfering substance with elevated levels of bovine mucus, blood, DNA from human cells, gastric acid, hypertonic saline, physiological saline, culture media, NALC pellet material and five anti-TB medications (Isoniazid, Rifampicin, Streptomycin, Pyrazinamide, Ethambutol) (Table 12). The results showed no interference in the performance of the MTB assay in the presence of
15 high levels of blood, DNA from human cells, gastric acid, hypertonic saline, physiological saline, culture media, NALC pellet material and five anti-TB medications (Isoniazid, Rifampicin, Streptomycin, Pyrazinamide, Ethambutol). Interference of the Abbott RealTime MTB assay was observed in the presence of bovine mucus at 8.3% (all five replicates were false negative or inhibited) and 5.0% (one of five replicates was false negative). No
20 interference was found at bovine mucus concentrations of 2.5% or less.

Carryover

To evaluate the potential of carryover from high positive MTB samples to negative samples when using the Abbott RealTime MTB assay, five *m2000* system runs each
25 consisting of 96 samples (Positive Control, Negative Control, 46 high positive samples at 1×10^7 copies/mL and 46 negative samples) in which the high positive samples were interspersed among negative samples. The MTB concentration in the high positive sample of 1×10^7 copies/mL resulted in a Ct value that was earlier than 95% or more of the results obtained from specimens of the MTB positive population tested with the MTB assay. The assay did
30 not exhibit any carryover from high positive samples to the 230 negative samples in the five runs. A 96 sample run was completed in less than 8 hours.

Reproducibility

A reproducibility test was performed to evaluate the Abbott RealTime MTB assay repeatability in the *m2000* system and the compatibility between the Abbott *m2000sp* instrument and the manual sample preparation method. The study was performed with a positive panel at three times the claimed LOD level and a negative panel. The study was conducted by four operators using two lots of MTB amplification reagents: two operators performing runs using an Abbott *m2000sp* instrument and two operators performing runs using manual sample preparation. For each sample preparation method, the two operators each used one unique lot of Abbott RealTime MTB amplification reagents and tested each panel member in replicates of eight, once per day, for five days, for a total of 40 replicates per panel member (80 total replicates per panel member per method; 160 total tested with *m2000sp* instrument sample preparation and 160 total tested with manual sample preparation). The overall agreement with expected results was 100% (159/159, one sample was invalid because of an instrument error) with a lower 95% CI of 98.1% for samples prepared with the Abbott *m2000sp* instrument or with manual sample preparation. The MTB assay is compatible with both the Abbott *m2000sp* instrument and the Abbott manual sample preparation method.

Clinical sensitivity and specificity

One sputum or one NALC sediment was tested from each of 582 TB suspect patients. Samples were collected from Russia, South Africa, Uganda, the United States, and Vietnam. Each specimen was split to allow testing of MTB on one aliquot and smear and culture on the second aliquot. The testing samples were blinded and final result decoding was performed by AM statistical group. For MTB testing, two specimens generated an invalid IC result, and an additional four specimen results gave *m2000* error codes. The frequency of clinical specimens with invalid results measured by inhibition was 0.3% (2/582), while the invalid rate including both inhibition and instrument errors was 1.0% (6/582). Five culture negative specimens that were positive by both the MTB assay described herein and a commercially available MTB NAAT were excluded from the analysis. A total of 571 valid samples were included for data analysis. The overall MTB sensitivity versus culture was 93% (198/212). The assay sensitivity was 99% (147/149) in smear positive, culture positive specimens, and 81% (51/63) in smear negative, culture positive samples. The specificity was 97% (348/359) (Table 13). 76 of the MTB negative samples contained Non-Tuberculous Mycobacteria (NTM). Of these, 38 were MAC (*M. avium* complex), seven were *M. goodii*, five were *M. kansasii*, five were *M. chelonae/abscessus*, three were *M. xenopi*, and 18 contained other

mycobacterial species. With the MTB assay described herein, all of the NTM sample results were “MTB Not Detected” with the exception of two samples that produced “MTB Detected” results with late CN (>38) values as compared to the assay cutoff of 40. The specificity value of 97% resulted from testing the NTM population is similar to the specificity observed when testing the non-NTM population. Furthermore 500 non-TB suspect patient sputum samples collected from within the U.S. population showed 100% TB negative test results.

Table 11
Limit of Detection

10

Target Concentration (cfu/mL)	Number Tested	Number Detected	Detection Rate	Percent Detected
80	20	20	1.00	100
50	20	20	1.00	100
25	20	20	1.00	100
10	20	20	1.00	100
5	20	20	1.00	100
1	20	18	0.90	90
0.50	20	7	0.35	35
0.10	20	2	0.10	10
0.05	20	1	0.05	5

A probit analysis of the Abbott RealTime MTB data determined that the concentration of MTB detected with 95% probability was 2.45 cfu/mL at CN cutoff 40 (95% Confidence Interval of 1.44 – 6.10 cfu/mL).

15 Table 12
Potential interfering substances and their concentrations/percentages tested to determine susceptibility to interfering substances

Potentially Interfering		
Substance	Specimen Source	Concentration / Percentage
Mucus	Sputum	Mucin 5% (w/v)
Blood	Sputum or BAL	5% (v/v)
DNA from human cells	Sputum, BAL, NALC Sediments of Sputum/BAL	10 ⁶ cells/mL
Gastric acid	Sputum/BAL	pH 3 to 4 HCl in water, neutralized to pH 6 to 8 with sodium bicarbonate
Hypertonic saline used to induce sputum	Sputum	NaCl (5% w/v)
Physiologic saline used to collect BAL	BAL	NaCl (0.9% w/v)
Culture media	MTB culture	100%
Material used to resuspend NALC pellets	NALC pellets	0.067 M phosphate, pH 6.8
Isoniazid (Anti-TB medication)	Sputum or BAL	90 mg/mL
Rifampicin/Rifampin (Anti-TB medication)	Sputum or BAL	120 ug/mL
Streptomycin (Anti-TB medication)	Sputum or BAL	400 ug/mL
Pyrazinamide (Anti-TB medication)	Sputum or BAL	500 ug/mL
Ethambutol (Anti-TB medication)	Sputum or BAL	60 ug/mL

Table 13
Sensitivity and specificity obtained when testing clinical specimens

5

Culture / Smear results

10

	Sensitivity			Specificity
	C + / S +	C + / S -	C +	C -
RealTime MTB	99%	81%	93%	97%
	(147/149)	(51/63)	(198/212)	(348/359)

EXAMPLE 11

Inactivation Reagents

This example describes inactivation reagents for use in MTB detection assays. The assay is a NAAT for the detection of MTB complex DNA in respiratory specimens (sputum, bronchial alveolar lavage (BAL) and N-acetyl-L-cysteine (NALC) sediments of sputum and bronchial alveolar lavage (BAL). A sample inactivation reagent and procedure were developed to liquefy viscous samples and to reduce MTB viability to allow for safe testing of samples outside biosafety cabinets. The study was to assess the effectiveness of the sample inactivation procedure and to determine the stability of the Inactivation Reagent (IR).

For the viscosity reduction study, 150 sputum samples were mixed with IR (0.6% sodium hydroxide [w/v], 60% isopropanol [v/v], and 1.8% Tween-20 [v/v]) at a ratio of 1:2 or 1:3. The mixtures were vortexed vigorously and incubated at room temperature. The mixture was vortexed again after 20 to 30 minutes of incubation. Reduction of viscosity was assessed by visual examined after 30 minutes, 60 minutes, and 24 hours of incubation.

For the inactivation study, two MTB clinical isolates and MTB ATCC 27294 isolate were used to prepare mock MTB positive respiratory samples by mixing one mL of MTB cell suspension in the concentrations of 1×10^6 , 1×10^7 , or 1×10^8 cfu/mL with four mL of pooled MTB negative NALC treated respiratory sample. Each mock MTB NALC sample was then mixed with IR at ratios of 1:2 or 1:3. A mock sample treated with sterile PBS buffer at a sample to PBS ratio of 1:2 was used as the positive control. Negative controls were prepared by adding sterile PBS to the pooled MTB negative NALC sample at a ratio of PBS to NALC of 1:2. All samples/controls were vortexed vigorously and incubated for 60 minutes at room temperature. Vortexing was repeated 30 minutes into the incubation. At the end of the incubation, the IR treated samples were transferred into new 50 mL tubes, vortexed and centrifuged for 15 minutes at 3000 x g. The sediment was re-suspended in 10 mL sterile PBS and centrifuged for an additional 15 minutes at 3000 x g. Pellets were each re-suspended in 10 mL sterile PBS. One mL of the suspension was used to inoculate a Mycobacterial Growth Indicator Tube (MGIT). The final MTB added to each MGIT culture ranged from $1 - 2 \times 10^4$ to $1 - 2 \times 10^6$ cfu. In addition, a total of 51 MTB positive clinical NALC sediments of sputum, 20 from Northwestern Memorial hospital and 31 from Lancet Laboratories, were tested for growth after the IR treatment at a sample to IR ratio of 1:3 with the same procedure. Ten of the 20 samples from Northwestern Memorial Hospital were treated at a sample to IR ratio of 1:2. The remaining 41 samples were treated with a sample to IR ration of 1:3. Culture was performed with BACTEC MGIT 960 system (Becton Dickinson, Sparks, MD) for 42 days. Positive growth was identified with Gen-Probe Accuprobe® system (Gen-Probe Inc, San Diego, CA). Initial studies to demonstrate the inactivation efficiency of direct

respiratory samples (MTB smear and culture positive sputum samples) were also performed in combination with an IR stability study as described in the following paragraph.

To determine the optimal storage condition for IR, three aliquots of IR were stored for 39 days at storage conditions of 15-30°C and 33-37°C in glass or polypropylene bottles. Each
5 aliquot of IR at each storage condition was examined for changes in appearance and volume and tested for MTB inactivation efficacy after 39 days of storage with 12 MTB smear and culture positive sputum samples obtained from SAGE Bio Networks (Dhaka, Bangladesh) and Foundation for Innovative New Diagnostics (FIND) MTB specimen bank using a 1:3 sample to IR ratio. An MTB strain H37Rv cell panel obtained from Zeptomatrix Corporation
10 (Buffalo, New York) was used as the Positive Control.

The viscosity reduction study showed that 60 minutes of incubation was sufficient to reduce the viscosity of the samples. For the inactivation study, none of the mock MTB samples prepared with the three MTB isolates at 1×10^8 , 1×10^7 , and 1×10^6 cfu/mL showed
15 MTB growth after being treated with IR at a sample to IR ratio of 1:3. One IR-treated sample, prepared with 1×10^7 cfu/mL MTB and treated with IR at a sample to IR ratio of 1:2, showed growth after 27 days of incubation, although two repeats at the same bacterial concentration tested negative for growth. None of the 20 MTB positive NALC sputum sediments showed
20 MTB growth following the treatment with IR at a sample to IR ratio of 1:2 or 1:3. Further, 31 clinical NALC sputum sediments that had previously tested positive for MTB by culture were tested negative for MTB growth after receiving IR treatment at 1:3 sample to IR ratio.

Changes in appearance after storage were not observed after 39 days. A volume loss of 0 to 6% was observed after 39 days of storage. The highest volume loss of 6% was observed when IR was stored in polypropylene containers at 33-37°C. However, the efficiency of the IR solution to inactivate MTB was not affected after storage. The 12 MTB
25 positive sputum samples showed no growth after being treated with the IR stored under the various conditions described above.

It was noted that some MTB in clinical samples survived the recommended Cepheid GeneXpert MTB/RIF sample inactivation process (15 minute incubation period and a 1:2 sample to Sample Reagent ratio) (Banada, P.P., et al., 2010. J. Clin. Microbiol. 48:3551-
30 3557). The authors suggested complete MTB inactivation may require a longer incubation time. Experimental data generated by the study described herein demonstrated that sample inactivation performed for 60 minutes using the recommended vortexing steps was sufficient for complete MTB inactivation.

When a 1:2 sample to IR ratio was used, one replicate of a 1×10^7 MTB cfu/mL culture (2×10^5 cfu/mL in the MGIT culture) showed growth after 27 days of incubation of the MGIT culture. A previous study showed that MGIT cultures containing 10 cfu/mL of MTB became positive after 16 days of incubation, the result suggested that a very low number MTB

5 survived the inactivation process when a 1:2 sample to IR ratio was used (Tortoli, E., P. et al., J. Clin. Microbiol.37(11):3578-3582; Wallis, et al., 1999. Antimicrobial Agents and Chemotherapy, 43:2600-2606). To achieve the optimal inactivation efficiency, the sample to IR ratio of 1:3 was used for the rest of the inactivation experiments.

10 In conclusion, the IR evaluated in this study was able to liquefy sputum samples and to achieve effective inactivation of MTB in clinical specimens when treated with IR at a sample to IR ratio of 1:3 for 60 minutes. This inactivation procedure enables these samples to be safely handled outside of a biosafety cabinet after the proper inactivation procedure.

CLAIMS

We claim:

- 5 1. A composition, comprising: at least one primer pair selected from the group consisting of SEQ ID NOs: 1 and 2, SEQ ID NOs: 3 and 4, and SEQ ID NOs: 7 and 8.
2. The composition of claim 1, wherein said composition comprises SEQ ID NOs: 1-4 and 7-8.
- 10 3. The composition of claim 1, wherein said composition further comprises at least one probe selected from the group consisting of SEQ ID NOs: 5, 6, and 9.
4. A composition, comprising:
- 15 a set of primer pairs of SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4.
5. The composition of claim 1, wherein said composition further comprises at least one probe selected from the group consisting of SEQ ID NOs: 5, 6, and 9.
- 20 6. The composition of claim 4 or 5, wherein said composition further comprises the primer pair of SEQ ID NOs: 7 and 8.
7. A composition comprising the nucleic acids of SEQ ID NOs: 1-9.
- 25 8. The composition of any of claim 1-7, wherein one of more of the recited SEQ ID NOs comprises a label.
9. The composition of claim 8, wherein said label comprises a fluorophore.
- 30 10. The composition of claim 8, wherein said label comprises a fluorophore/quencher pair.

11. The composition of any one of claims 1 to 10, wherein said composition further comprises one or more nucleic acid sequences selected from the group consisting of SEQ ID NOs: 10-36.
- 5 12. The composition of any of claims 1-11, wherein said composition is a reaction mixture.
13. The composition of claim 12, wherein said reaction mixture comprises a sample.
- 10 14. The composition of claim 13, wherein said sample comprises a sample comprising a mycobacterium tuberculosis (MTB) target nucleic acid.
15. A kit, comprising:
- 15 a) the composition of any one of claims 1 to 11; and
- b) at least one reagent for performing a nucleic acid amplification reaction.
16. The kit of claim 15, wherein said reagent is selected from a nucleic acid polymerase; a plurality of dNTPS, a buffer, and an inactivation reagent.
- 20 17. The kit of claim 16, wherein said inactivation reagent comprises water, a detergent, an alcohol, and NaOH.
18. The kit of claim 16, wherein said inactivation reagent comprise isopropanol, sodium hydroxide, TWEEN-20, and water.
- 25 19. A reaction mixture comprising one or more primers or probes comprising a sequence of SEQ ID NOs:1-9 hybridized to a mycobacterium tuberculosis (MTB) nucleic acid.
20. The reaction mixture of claim 19, wherein one or more of said one or more primers or
- 30 probes comprises a label.
21. A reaction mixture comprising one or more primers or probes hybridized to each of two MTB target sequences: insertion sequence (IS) 6110 and Protein Antigen B (PAB).

22. A method of identifying an MTB nucleic acid in a biological sample, comprising:
a) contacting a biological sample from a subject with a nucleic acid primer or probe of SEQ ID NOs:1-9, and
b) detecting the binding of said nucleic acid primer or probe to said MTB nucleic acid.
23. The method of claim 22, further comprising the step of c) determining the presence of MTB in said sample when said binding is detected.
24. The method of claim 22, wherein said detecting comprises real time PCR.
25. The method of claim 22, wherein said method further comprises the step of inactivating said MTB in said sample using an inactivation reagent prior to said contacting.
26. The method of claim 25, wherein said inactivation reagent comprises water, a detergent, an alcohol, and NaOH.
27. The method of claim 25, wherein said inactivation reagent comprise isopropanol, sodium hydroxide, TWEEN-20, and water.
28. The method of claim 22, wherein said sample is sputum.
29. The method of claim 22, wherein said sample is bronchoalveolar lavage [BAL].
30. The method of claim 22, wherein said sample is N-acetyl-L-cysteine [NALC] sediments of sputum.
31. The method of claim 25, wherein said method further comprises the step of extracting DNA from said sample following said inactivation.
32. A method of detecting an MTB nucleic acid in a biological sample, comprising:
a) inactivating said biological sample with an inactivating reagent to generate an inactivated sample;
b) extracting DNA from said inactivated sample;

c) contacting said DNA with one or more primer pairs and one or more nucleic acid probes;

d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and

5 e) identifying the presence of said MTB in said sample.

33. The method of claim 32, wherein said amplification assay is real time PCR.

34. The method of claim 32, wherein said sample is sputum.

10

35. The method of claim 32, wherein said sample is bronchoalveolar lavage [BAL].

36. The method of claim 32, wherein said sample is N-acetyl-L-cysteine [NALC] sediments of sputum.

15

37. The method of claim 32, wherein two primer pairs and two probes are employed in said contacting step.

38. The method of claim 37, wherein said two primer pairs and two probes hybridize to
20 MTB target sequences insertion sequence (IS) 6110 and Protein Antigen B (PAB).

39. The method of claim 32, wherein said nucleic acid primers are SEQ ID NOs: 1-4.

40. The method of claim 32, wherein said nucleic acid probes comprise SEQ ID NOs: 5
25 and 6.

41. The method of claim 32, wherein said inactivation reagent comprises water, a detergent, an alcohol, and NaOH.

30 42. The method of claim 32, wherein said inactivation reagent comprises isopropanol, sodium hydroxide, TWEEN-20, and water.

43. A method of detecting an MTB nucleic acid in a biological sample, comprising:

- a) inactivating said biological sample with an inactivating reagent comprising isopropanol, sodium hydroxide, TWEEN-20, and water to generate an inactivated sample;
- b) extracting DNA from said inactivated sample;
- c) contacting said DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6;
- d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and
- e) identifying the presence of said targets in said sample.
- 10
44. A method of detecting an MTB nucleic acid in a biological sample, comprising:
- a) inactivating said biological sample with an inactivating reagent to generate an inactivated sample;
- b) extracting DNA from said inactivated sample;
- 15 c) contacting said DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6;
- d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and
- 20 e) identifying the presence of said targets in said sample.
45. A method of detecting an MTB nucleic acid in a biological sample, comprising:
- a) inactivating said biological sample with an inactivating reagent to generate an inactivated sample;
- 25 b) extracting DNA from said inactivated sample;
- c) contacting said DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6;
- d) performing a real time PCR assay to amplify one or more MTB nucleic acid
- 30 targets; and
- e) identifying the presence of said targets in said sample.
46. A method of detecting an MTB nucleic acid in a biological sample, comprising:

- 5 a) inactivating said biological sample with an inactivating reagent to generate an inactivated sample;
- b) extracting DNA from said inactivated sample;
- c) contacting said DNA with one or more primer pairs selected from SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and one or more nucleic acid probes selected from SEQ ID NOs: 5 and 6;
- d) performing an amplification assay to amplify one or more MTB nucleic acid targets selected from IS6110 and PAB; and
- e) identifying the presence of said targets in said sample.
- 10

47. A method of detecting an MTB nucleic acid in a biological sample, comprising:
- a) inactivating said biological sample with an inactivating reagent to generate an inactivated sample;
- 15 b) extracting DNA from said inactivated sample;
- c) contacting said DNA with the primer pairs of SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and the nucleic acid probes of SEQ ID NOs: 5 and 6;
- d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and
- 20 e) identifying the presence of said targets in said sample.

48. A method of detecting an MTB nucleic acid in a biological sample, comprising:
- a) inactivating said biological sample with an inactivating reagent comprising isopropanol, sodium hydroxide, TWEEN-20, and water to generate an inactivated sample;
- 25 b) extracting DNA from said inactivated sample;
- c) contacting said DNA with the primer pairs of SEQ ID NOs: 1 and 2 and SEQ ID NOs: 3 and 4; and the nucleic acid probes of SEQ ID NOs: 5 and 6;
- d) performing an amplification assay to amplify one or more MTB nucleic acid targets; and
- 30 e) identifying the presence of said targets in said sample.

FIG. 1


ASSAY WORK FLOW:

FIG. 2





FIG. 3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US15/39362

A. CLASSIFICATION OF SUBJECT MATTER IPC(8) - C12Q 1/68; G06F 19/22; C12N 15/09 (2015.01) CPC - C12Q 1/689; G06F 19/22 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC(8): C12Q 1/68; G06F 19/22; C12N 15/09 (2015.01) CPC: C12Q 1/689; G06F 19/22 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) PatSeer (US, EP, WO, JP, DE, GB, CN, FR, KR, ES, AU, IN, CA, INPADOC Data); Google/Google Scholar; NCBI/BLAST/PubMed; Dialog ProQuest; 'M tuberculosis,' primer, probe, inactivation, reagent, buffer, sputum, 'bronchoalveolar lavage,' 'NaOH,' detergent, 'Tween 20,' isopropanol, 'DNA extract'		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	WO 1995/031570 A1 (ABBOTT LABORATORIES); November 23, 1995; page 18, lines 10-20; page 20, lines 15-20	21 ----- 38
X --- Y	US 2013/0323224 A1 (OJHA, AK et al.); December 5, 2013; paragraphs [0008], [0038], [0053], [0056], [0127], [0168], [0216], [0283], [0292]	32-37 ----- 38, 41-42
Y	US 2009/0137406 A1 (KINOSHITA, K et al.); May 28, 2009; paragraphs [0073], [0187]-[0190]	41, 42
A	WO 2010/132054 A1 (IMMPORT THERAPEUTICS INC et al.); November 18, 2010; paragraphs [0008], [0024]; SEQ ID: NO:578	1-20, 22-31, 43-48
A	US 8143386 B2 (REED, SG et al.); March 27, 2012; column 2, lines 25-35; column 2, lines 45-55; SEQ ID NO: 5	1-20, 22-31, 43-48
A	EP 2035440 B1 (ABBOTT LABORATOIRES); March 18, 2009; paragraph [0023]; SEQ ID NOs: 40, 41	1-20, 22-31, 43-48
A	WO 2008/152636 A2 (QUARK PHARMACEUTICALS INC. et al.); December 18, 2008; Claim 16, SEQ ID NO:85	1-20, 22-31, 43-48
A	WO 2005/012560 A1 (UNIVERSITE LAVAL); February 10, 2005; SEQ ID NO: 7	1-20, 22-31, 43-48
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 13 October 2015 (13.10.2015)		Date of mailing of the international search report 30 NOV 2015
Name and mailing address of the ISA/ Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-8300		Authorized officer Shane Thomas PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774