

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



(10) International Publication Number

WO 2013/175009 A1

(43) International Publication Date

28 November 2013 (28.11.2013)

(51) International Patent Classification:

A61K 31/57 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/EP2013/060801

(22) International Filing Date:

24 May 2013 (24.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12305586.5 25 May 2012 (25.05.2012) EP

(71) Applicants: LABORATOIRE HRA-PHARMA [FR/FR]; 15 rue Béranger, F-75003 Paris (FR). INSERM (Institut National de la Santé et de la Recherche Médicale) [FR/FR]; 101 rue de Tolbiac, F-75013 Paris (FR).

(72) Inventors: RESCHE-RIGON, Michèle; 38 rue Richard Lenoir, F-75011 Paris (FR). LEVY, Delphine; 206 rue de Noisy le Sec, F-93170 Bagnolet (FR). GAINER, Erin; 18 rue de l'Atlas, F-75019 Paris (FR). GOMPEL, Anne; 32 rue de Bièvre, F-75005 Paris (FR). FORGEZ, Patricia; 3 rue de la Vigne de Lozère, F-91120 Palaiseau (FR).

(74) Agents: CHAJMOWICZ, Marion et al.; BECKER & ASSOCIES, 25 rue Louis Le Grand, F-75 002 Paris (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2013/175009 A1

(54) Title: ULIPIRISTAL ACETATE FOR PREVENTION AND TREATMENT OF BREAST TUMORS

(57) Abstract: The invention relates to the use of ulipristal acetate, or any of its metabolites, in preventing or treating a breast tumor in a patient, preferably a patient that carries a mutation in BRCA1 gene.


Ulipristal acetate for prevention and treatment of breast tumors

The present invention relates to prevention and therapy of breast tumor with ulipristal acetate.

5 **Background of the invention**

Ulipristal acetate (UPA) is a progesterone receptor modulator which efficiently binds and inhibits progesterone receptor in progesterone target tissues.

UPA, formerly known as CDB-2914, designates 17 α -acetoxy-11 β -[4-*N*, *N*-dimethylamino-phenyl]-19-norpregna-4, 9-diene-3, 20-dione, represented by formula I:



10 UPA, and methods for its preparation, are described e.g., in US Patent Nos. 4,954,490; 5,073,548; and 5,929,262, as well as in international patent applications WO2004/065405 and WO2004/078709.

15 UPA has been approved for emergency contraception (under tradename EllaOne \circledR), and for treatment of uterine fibroids (under tradename Esmyna \circledR). Various other potential clinical applications have been proposed in Chabbert-Buffet et al, Human Reproduction, 2005, 11(3): 293-307.

20 Other antiprogestins such as mifepristone and onapristone were developed for breast cancer treatment. Administration of mifepristone or onapristone in a second or third line of treatment, in 123 postmenopausal women with metastatic breast cancers induced objective response rate and disease stabilization in 11% and 43% of the patients, respectively (Romieu et al. 1987, Bull Cancer 74(4): 455-461; Klijn, et al. 1989, Cancer Res 49(11): 2851-2856). Unfortunately, clinical studies were not sustained because of the anti-glucocorticoid and liver toxic side effects of mifepristone and onapristone, respectively.

25 Long term UPA exposure may affect hormone-responsive tissues, and particularly breast tissue, as well. However, only gene reporter transactivation studies in T-47D breast cancer cell line were reported (Attardi et al. 2002, Mol Cell Endocrinol 188(1-2): 111-123; Attardi et al. 2004, J Steroid Biochem Mol Biol 88(3): 277-288).

30 There is still a need for a therapeutic agent for treating, and even preventing breast tumors, while causing no or reduced side-effects.

Summary of the invention

The inventors now propose a method for preventing or treating a breast tumor in a patient, which method comprises administering ulipristal acetate (UPA) or any of its metabolites.

5

It is therefore provided ulipristal acetate (UPA) or any of its metabolites for use in preventing or treating a breast tumor in a patient.

In a preferred embodiment, the patient carries a mutation in BRCA1 and/or BRCA2 gene.

10

In a particular preferred embodiment, ulipristal acetate (UPA) or any of its metabolites is used in preventing a breast tumor in a patient that carries a mutation in BRCA1 and/or BRCA2 gene.

15

In another preferred embodiment, ulipristal acetate (UPA) or any of its metabolites is used in treating a breast tumor in a patient, wherein the breast tumor preferably is a breast carcinoma.

Detailed description of the invention

20 The "patient" means any subject in need of the preventive or curative treatment of the invention, preferably a woman. However men are also encompassed, since they may be susceptible to breast tumors as well.

Preferably the patient has been diagnosed to carry a mutation in BRCA1 and/or BRCA2 gene.

25 BRCA1 and BRCA2 are tumor suppressor genes.

The mutational events of the BRCA1 or BRCA2 locus can involve deletions, insertions, and point mutations within the coding sequence and the non-coding sequence. Deletions may be the entire gene or only a portion of the gene. Point mutations may result in stop codons, frameshift mutations, or amino acid substitutions. Somatic mutations are those which occur 30 only in certain tissues, e.g., in the tumor tissue, and are not inherited in the germline. Germline mutations can be found in any of a body's tissues or cells and are inherited. If only a single allele is mutated, a predisposition to breast cancer is indicated.

In a particular embodiment, the patient may carry one copy or two copies of BRCA1 or BRCA2 predisposing alleles.

35 It is believed that BRCA1 or BRCA2 predisposing alleles are recessive to wild-type alleles; that is, cells that contain at least one wild-type BRCA1 or BRCA2 allele are not cancerous. However, cells that contain one wild-type BRCA1 or BRCA2 allele and one predisposing

allele may occasionally suffer the loss of the wild-type allele, either by random mutation or by chromosome loss during cell division. All the progeny of such a mutant cell lack the wild-type function of BRCA1 or BRCA2 and may develop into tumors. Thus, the predisposing alleles of BRCA1 or BRCA2 are susceptible to cancer, and the susceptibility is inherited in a dominant fashion.

The majority of mutant alleles are nonsense or frameshift and produce truncated proteins which are predicted to vary in length from 5% to 99% of the full-length protein. Many of these mutations reside in exon 11 of BRCA1 gene which comprises 61% of the BRCA1 coding region. The full-length BRCA1 cDNA sequence and the coding regions of BRCA1 gene have been described in U.S. Pat. No. 5,747,282.

10 The mutations in BRCA1 or BRCA2 may be detected by any method known in the art.

Myriad's current BRCA diagnostic test, BRACAnalysis®, uses a combination of two traditional technologies—Sanger sequencing and PCR—to identify mutations associated with a significant risk of breast cancer in the BRCA1 and BRCA2 genes.

15 Other methods for screening for BRCA1 and BRCA2 mutations include single stranded conformational polymorphism (SSCP) analysis and selected DNA sequencing of gene variants, or DHPLC and DNA sequencing of gene variants.

Still other methods have been disclosed, including protein-based systems for detecting mutations in the BRCA1 gene (U.S. Pat. No. 5,965,377; US6,514,713).

20 The term "tumor" refers to the presence of cells possessing characteristics such as atypical growth or morphology, including uncontrolled proliferation, immortality, metastatic potential, rapid growth and proliferation rate, and certain characteristic morphological features. "Tumor" includes both benign and malignant (i.e. cancerous) neoplasms. Adenomas and cysts are encompassed. Breast cancers include breast carcinomas. Breast cancer can begin in 25 different areas of the breast – the ducts, the lobules, or in some cases, the tissue in between. In the context of the present invention, any type of breast tumor or breast cancer is encompassed, including non-invasive, invasive, recurrent, and metastatic breast cancers.

In a particular embodiment, the patient has been diagnosed with a Hereditary Breast and Ovarian Cancer (HBOC).

30

It is herein described a preventive and a therapeutic method against breast tumor.

"Prevention" or "preventing" means that UPA or any of its metabolites is administered in a patient who is asymptomatic or does not show a breast tumor. More particularly the patient may be at risk of developing such breast tumor, e.g. in view of the family history or of the 35 BRCA1 and/or BRCA2 gene status. Such prevention aims at reducing the risk of developing a breast cancer.

“Therapeutic treatment” or “treating” means that UPA or any of its metabolites is administered in a patient who has been diagnosed with a breast tumor. The treatment alleviates the symptoms of the disease, slows down the progression of the disease, may lead to a remission or to a complete cure of the disease.


5 In particular, UPA was shown to antagonize the proliferation of tumor cells, in particular in lobules and galactophores (or ductal cells), in particular in BRCA1 mutant cells.

In a particular embodiment, the patient, who may be at risk of developing a breast tumor, but not necessarily, may be further in need of a contraception. This is particularly useful when the patient has no regular contraception. If so, UPA or any of its metabolites, may be


10 proposed in a form and dosage adapted to provide both regular contraception and breast tumor prevention. In another embodiment, the patient may have uterine fibroids, and be both in need of a treatment against uterine fibroids and in need of a therapeutic or preventive treatment against a breast tumor. In that case, UPA or any of its metabolites, may also be proposed in a form and dosage adapted to provide both treatment against uterine fibroids

15 and prevention or treatment of a breast tumor.


Ulipristal acetate (UPA) is preferably used. However metabolites of UPA could be used as well. Metabolites of ulipristal acetate, include those described in Attardi et al, Journal of Steroid Biochemistry & Molecular Biology, 2004, 88: 277-288, e.g. monodemethylated CDB-
20 2914 (CDB-3877) ;didemethylated CDB-2914 (CDB-3963) ; 17alpha-hydroxy CDB-2914 (CDB-3236) ; aromatic A-ring derivative of CDB-2914 (CDB-4183). Preferably the metabolite is monodemethylated CDB-2914 (CDB-3877).




monodemethylated CDB-2914



didemethylated CDB-2914



17α-hydroxy CDB-2914



aromatic A-ring CDB-2914

UPA or its metabolites may be administered by various routes, e.g., orally, intravenously, or transdermally. A preferred administration route is the oral route. An injection at the site of the tumor is also possible. Other routes of administration are encompassed, including the vaginal or intrauterine route. Devices, in particular vaginal or intrauterine devices, allowing sustained release of UPA or its metabolites may be particularly useful. Subcutaneous implants may be further contemplated.

Methods and compositions for making useful dosage units are well-known to those skilled in the art. For example, conventional techniques for making tablets and pills, containing active

ingredients, are described in the standard reference, Chase et al., Remington's Pharmaceutical Sciences, (16th ed., Mack Publishing Co., Easton. Pa., U.S.A., 1980) ("Remington's"), at pages 1553 through 1584. Conventional techniques for making powders, and their composition are described at pages 1535 through 1552 of the reference.

5 Conventional techniques for coating pharmaceutical dosage forms are described at pages 1585 to 1593 of Remington's.

Oral solid dosage forms are preferentially compressed tablets, that may be coated or uncoated, or capsules.

10 Capsules are solid dosage forms using preferentially either a hard or soft gelatine shell as a container for the mixture of the active ingredient and inert ingredients. Procedures for production and manufacture of hard gelatin and soft elastic capsules are well known in the art.

Compressed tablets may contain any excipients which are diluents to increase the bulk of the active ingredient so that production of a compressed tablet of practical size is possible.

15 Binders, which are agents which impart cohesive qualities to powdered materials are also necessary. Starch, gelatine, sugars such as lactose or dextrose, and natural and synthetic gums are used. Disintegrants are necessary in the tablets to facilitate break-up of the tablet. Disintegrants include starches, clays, celluloses, algins, gums and crosslinked polymers. Lastly small amounts of materials known as lubricants and glidants are included in the tablets

20 to prevent adhesion to the tablet material to surfaces in the manufacturing process and to improve the flow characteristics of the powder material during manufacture. Colloidal silicon dioxide is most commonly used as a glidant and compounds such as talc or stearic acids are most commonly used as lubricants. Procedures for the production and manufacture of compressed tablets are well known by those skilled in the art.

25 In a particular embodiment, ulipristal acetate or a metabolite thereof is used in form of an uncoated tablet wherein ulipristal acetate or a metabolite thereof is mixed with excipients that are lactose monohydrate, povidone (polyvinylpyrrolidone), croscarmellose sodium, and magnesium stearate (e.g. as described in international patent application WO2010/066749). Long term treatments are preferred, e.g. for a period of at least 3 months.

30 In the context of prevention, the treatment may be maintained several years.

In the context of a curative treatment, UPA may be administered from a few days, to several months or years, e.g. during at least 3 months to about 5 years.

35 Dosage may be adapted depending on the specific condition and severity of the tumor, and the gender and weight of the patient. A typical dosage would range from 0.1 mg to 150mg, preferably between 5 and 80mg, still preferably between 10 and 50mg. Daily administration(s) are preferred.

The examples and figures illustrate the invention without reducing its scope.

Legends to the Figures

5 Figure 1: UPA effects on MMTV reporter gene transactivation.

MMTV-Luc reporter gene was transfected in HBE cells: (A) alone, or in combination, (B) with hPR-A and hPR-B, (C) hPR-A, or (D) hPR-B isoform plasmids. (E) T-47D cells were transfected with MMTV-Luc. Cells were treated with P at 100 nM and/or UPA, or RU from 1 to 1 000 nM as indicated or with 100 nM when not specified. Results correspond to control 10 fold induction (mean \pm SEM, n = 3 for HBE and T-47D). *p < 0.05, **p < 0.001.

Figure 2: UPA effects on PR target gene expression.

11 In HBE and T-47D cells, mRNA expression was analyzed by Quantitative RT-PCR for (A) FASN, (B) cyclin A, (C) BCL2 and (D) ALPL. Cells were treated with P and/or UPA at 100 15 nM, and/or E2 at 10 nM. Results correspond to control fold induction (mean \pm SEM, n = 5 for HBE and n = 2 for T47D). *p < 0.05, **p < 0.01.

Figure 3: UPA effects on GRE reporter gene transactivation.

GRE-Luc reporter gene was transfected in (A) HBE cells and in (B) MCF-7 cells. Cells were 20 treated by DEX and/or UPA at 100 nM. Results correspond to control fold induction (mean \pm SEM, n = 13 for HBE and n = 3 for MCF-7). *p < 0.01, **p < 0.001.

Figure 4: UPA effects on GR target gene expression.

In HBE and MCF-7 cells, mRNA expression was analyzed by Quantitative RT-PCR for (A) 25 IEX-1, (B) G0S8, (C) cyclin A, (D) BCL2. Cells were treated with DEX and/or UPA at 100 nM. Results correspond to control fold induction (mean \pm SEM, n = 5 for HBE and n = 2 for MCF-7). *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 5: UPA effect on hormone mediated cell 1 proliferation.

30 Cell proliferation was measured by [3H] thymidine incorporation in (A) HBE cells, (B) T-47D cells and (C) MCF-7 cells. HBE cells were treated 96h and T-47D and MCF-7 cells were treated 48h at the concentration of 100 nM for P, DEX, UPA and 10 nM for E2. Results are expressed in percentage of control (mean \pm SEM, n = 7 for HBE, n = 4 for T47D and n = 3 for MCF-7). *p < 0.05, **p < 0.01, ***p < 0.001.

35

Figure 6: UPA effects on hormone mediated cell apoptosis.

Cell apoptosis was measured by flow cytometry quantification of sub-G1 phase in (A) HBE cells, (B) T-47D cells, and (C) MCF-7 cells. HBE cells were treated 96h and T-47D and MCF-7 cells were treated 48h at the concentration of 100 nM for P, DEX, UPA and 10 nM for E2. Results are expressed in percentage of control (mean \pm SEM, n = 9 for HBE, n = 3 for T-47D and n = 4 for MCF-7). *p < 0.05, **p < 0.001.

Figure 7: PR, ER and Ki67 expressions in breast tissue xenografts.

ER, PR and Ki67 expression were analyzed by immunohistochemistry. (A) Representative image for ER and PR labeling in original breast tissue before and in treated grafts in mice. (B) Ki67 expression in C, E2, E2+P, E2+P+UPA treatment groups of grafted tissues. Magnification x400. (C) Mitotic index (Ki67 positive cells percentage) control fold induction in lobules and ducts (mean \pm SEM, n = 4 in lobules and n = 6 in ducts). *p < 0.05.

Figure 8 shows induction of mitotic index (Ki67 positive cells percentage) in lobular cells of grafted tissues that carry BRCA1 mutations (BRCA1^{+/+}) or not (BRCA1⁺⁻), in response to various treatments. mean \pm SEM.

Figure 9 shows induction of mitotic index (Ki67 positive cells percentage) in ductal cells of grafted tissues that carry BRCA1 mutations (BRCA1^{+/+}) or not (BRCA1⁺⁻), in response to various treatments. mean \pm SEM.

Examples:

Example 1: Impact of UPA on proliferation and apoptosis of normal breast epithelial cells (HBE) and in breast cancer cell lines.

Materials and methods

Steroids

The progesterone receptor antagonist ulipristal acetate (UPA) and its mono-N4 demethylated metabolite CDB-3877 (CDB) were kindly provided by HRA-Pharma (Paris, France). 17 β estradiol (E2), progesterone (P), dexamethasone (DEX) and mifepristone/RU-486 (RU) were purchased from Sigma (St Quentin Fallavier, France).

Cell culture procedures

T-47D and MCF-7 cell lines were respectively maintained in RPMI 1640 and in DMEM medium supplemented with 10% fetal calf serum (PAA Laboratory, Les Mureaux, France). T-

47D cell line was derived from a human ductal breast carcinoma and constitutively expressed high amounts of PR (progesterone receptor) and ER (estradiol receptor). Normal human breast epithelial cell (HBE) primary cultures were obtained from 20 women (aged 17-50 years). The procedure used to culture HBE cells is described in detail by Gompel and 5 colleagues (Gompel, et al., 1986, *J Clin Endocrinol Metab* 63(5): 1174-1180). HBE cells were maintained in HAM F10 medium (PAA Laboratory, Les Mureaux, France) containing hydrocortisone (5 ng/ml), Triiodo-L-thyronine (6.5 ng/ml), cholera toxin (10 ng/ml), transferrin (5 mg/ml), insulin (0.016 U/ml), epidermal growth factor (10 ng/ml) (Sigma, St Quentin Fallavier, France) and 5% human serum (Etablissement Français du Sang). HBE primary 10 cultures express epithelial markers as well as low levels of estradiol receptor (ER) and estradiol induced PR (Malet, et al., 1991. *J Clin Endocrinol Metab* 73(1): 8-17; Courtin, et al., 2011, *Breast Cancer Res Treat.*).

Steroid treatments

15 After seeding, cells were cultured for 24h in serum and phenol red free medium. Then treatments were carried out in a phenol red free medium containing 5% dextran-charcoal stripped serum. Cells were treated with P or DEX (100 nM), alone or in combination with UPA, RU (1 nM to 1 μ M) or CDB-3877 (100 nM). Control cells were treated with ethanol, at 10 1:1000 final ethanol concentration alone as vehicle or E2 (10 nM).

20

Reporter enzyme assays

Cells were transfected with reporter gene plasmids containing glucocorticoid and progesterone responsive elements (GRE/PRE): 1) the MMTV-Luc is a Mouse Mammary Tumor Virus Long Terminal Repeat promoter containing one GRE/PRE palindrome and three 25 GRE/PRE hemi-palindromes upstream to firefly luciferase gene in pFC31 vector, 2) the GRELuc contains six copies of GRE/PRE palindrome upstream to firefly luciferase gene in pBL vector. When indicated, HBE cells were transfected with human PR isoforms hPR-A and hPR-B expression plasmids constructed in POP3 vectors. Rous Sarcoma Virus promoter upstream to beta galactosidase gene (pRSV- β -Gal) was transfected in each experiment as 30 control. Transfections were performed according to the manufacturer instructions using Lipofectamine or Lipofectamine LTX Reagents (Invitrogen, Cergy-Pontoise, France) for breast cancer cell lines or HBE cells, respectively. After 24h of transfection, breast cancer and HBE cells were treated with hormones for 24h or 48h, respectively. At the end of the experiment, cells were lysed and luciferase activity was determined using the Luciferase 35 Assay System (Promega, Charbonnières-les-bains, France). Beta galactosidase activity was assessed using the Galacto Star kit (Applied Biosystems, Courtaboeuf, France) to normalize luciferase activity data.

Real-time quantitative reverse transcription PCR (qRT-PCR)

Total RNA was extracted using TriZOL Reagent (Invitrogen, Cergy-Pontoise, France).

2 µg of total RNA was subjected to reverse transcription (RT) using random primers for 1h at

5 37°C. 2 µl of RT product was diluted (1:10) and subjected to quantitative PCR using sequence specific primers (300 nM) and Brilliant SYBR GREEN QPCR master mix (Fermentas, Saint- Rémy-lès-Chevreuse, France) on an Mx3000P apparatus (Agilent Technologies, Massy, France). Conditions were 1 cycle at 95°C for 10 min followed by 40 cycles at 95°C for 30 sec, 60°C for 1 min and 72°C for 30 sec. Gene expression values were

10 normalized to the housekeeping gene 36B4. Time of steroid treatment was chosen to get the optimal stimulation for a given gene. ALPL and G0S8 mRNA were analyzed after 6h of treatment. IEX-1, FASN, and BCL2 were analyzed after 24h of treatment. Cyclin A mRNA was analyzed after 24h for GR responses or 48h for PR responses.

15 **Tritiated Thymidine incorporation**

After 24h of hormonal treatment, cells were incubated with [methyl-1 3H] thymidine (Perkin Elmer, Courtaboeuf, France) for 48h or 20h for HBE or cancer cells, respectively at 37°C. After incubation cells were washed twice with PBS 1X and once with 5% trichloroacetic acid (TCA). Cells were incubated in 5% TCA for 15 min at 4°C and lysed in NaOH 0.1 N for 30 20 min at 37°C. The total cell lysate was added to 5 ml of Ecolite scintillation liquid (MP biomedical, Illkirch, France) and radioactivity was counted with a β-counter HIDEX 300SL(ScienceTec, Courtaboeuf, France).

Flow Cytometry analysis

25 After 96h or 48h of hormonal treatment, respectively, for HBE or MCF-7 and T-47D cells, cells were washed in PBS, matrix dissociated with accutase enzyme (PAA laboratory, Les Mureaux, France), and centrifuged 5 minutes at 1350 rpm. Cells were fixed and frozen at -20°C in 70% ethanol. Before analysis, cells were washed in PBS and stained with 10 µg/ml propidium iodide in PBS (containing 0.835 U/ml RNase A) (Sigma, St Quentin Fallavier, 30 France). For each sample at least 10 000 cells were counted in a BD LSR II flow cytometer (BD Biosciences, Le Pont de Claix, France). After gating out doublets and debris, cycle distribution was analyzed using the ModFit LT software (Verity Software House, USA).

Statistical analysis

35 Results were expressed in mean ± SEM. To determine the statistical significance of treatments, One-way ANOVA and Tukey-Kramer multiple comparisons tests were performed to compare the relative efficiency of each treatment with the Instat 3 software (GraphPad,

USA). When only two treatments were compared, an unpaired t test was performed. $p < 0.05$ was considered as significant.

Results

5

UPA effects on PR gene transactivation

In order to analyze UPA antagonist properties on PR-induced gene transactivation, normal human breast epithelial cells (HBE) and T-47D breast cancer cell line were transfected with the MMTV-Luc reporter gene (Fig. 1). In HBE cells which expressed low and variable 10 amounts of PR among patients (Malet, et al., 1991. *J Clin Endocrinol Metab* 73(1): 8-17), progesterone (P) induced a significant luciferase expression (1.38 ± 0.11 control fold induction, $p < 0.05$) (Fig. 1 A). To better study the dose effects of UPA on PR, HBE cells were co-transfected with hPR-A and hPR-B expression plasmids (Fig. 1 B). UPA and RU-486 (RU) inhibited P induced MMTV11 Luc transactivation in a dose dependent manner from 15 1 000 to 10 nM. HBE cells were then also transfected with only hPR-A or hPR-B, to discriminate UPA actions between PR isoforms. No difference with increasing the levels of each PR isoform was seen on UPA antagonist performance (Fig. 1 C, D). In the T-47D cell line, UPA and RU displayed similar and potent PR antagonist actions from 1 000 to 10 nM (Fig. 1 E). Partial antagonist responses were also detected at 1 nM (Fig. 1 E). In both HBE 20 and T-47D cells, UPA and RU did not exhibit progestational agonist properties on reporter gene transcription (Fig. 1 B-E). These results indicated that UPA acted as a potent P antagonist in normal and cancerous breast cells.

UPA activity on mRNA expression of PR target genes

25 A hundred nM was the lowest concentration of UPA necessary to exert a complete antagonist activity in HBE cells. This concentration was therefore, chosen to further study the effects of UPA on specific P target genes in HBE and T-47D cells. Estradiol (E2) was added to HBE cells, in order to increase PR expression as reported previously.

30 Fatty acid synthase (FASN) is implicated in normal breast cells differentiation as well as in mammary tumor progression. An up3 regulation of FASN mRNA by progestins through PR was previously demonstrated in normal and tumoral breast cells in vitro and in vivo. As shown in Figure 2 A, UPA was able to prevent P induction of FASN mRNA expression in HBE as in T-47D cells. We observed that P down regulated cyclin A mRNA expression in normal breast cells and in T-47-D cells (Fig. 2 B), as previously reported in MDA-MB-231 35 breast cancer cells transfected by PR (Lin, et al., 2003, *Endocrinology* 144(12): 5650-5657.). UPA reversed P-induced cyclin A mRNA down-regulation in HBE and T-47D (Fig. 2B). We previously reported a decrease of the anti-apoptotic B-cell CLL/lymphoma 2 (BCL2) protein

expression under progestin treatment in normal and tumoral breast cells (Kandouz, et al., 1996, *Int J Cancer* 68(1): 120-125; Gompel, et al., 2000, *Steroids* 65(10-11): 593-598). As shown in Figure 2 C, we also observed a decrease of BCL2 mRNA expression by P in HBE and T-47D cells.

5 Nevertheless, UPA did not reverse BCL2 mRNA down-regulation induced by P in HBE cells, whereas UPA antagonized the P inhibitory effect on this transcript in T-47D cells ($p < 0.01$) (Fig. 2 C). Tissue non-specific alkaline phosphatase (ALPL) is a P responsive gene implicated in metastasis of breast cancer. In T-47D, UPA totally inhibited the strong P mediated induction of ALPL transcript (Fig. 2 D). In HBE cells, ALPL mRNA expression was
10 not modified by P treatment (data not shown). UPA did not display any PR agonist activity on these genes.

UPA effects on GR gene transactivation

The study was performed in HBE and MCF-7 cells, which expressed high amounts of GR as
15 previously reported (Courtin, et al.; 2011, *Breast Cancer Res Treat.*). Cells were transfected with a GRE-Luc reporter gene and treated with dexamethasone (DEX), UPA, and its proximal mono-demethylated metabolite, CDB-3877 (CDB), known to exert less anti-glucocorticoid activity than UPA (Attardi, et al., 2004, *J Steroid Biochem Mol Biol* 88(3): 277-288.). We confirmed the reduced anti glucocorticoid activity of this metabolite, since UPA
20 inhibited by $41.3 \pm 5.8\%$ the DEX induced luciferase transactivation, whereas its metabolite failed to inhibit DEX activity in HBE cells ($18.1 \pm 11.8\%$) (Fig. 3 A). However, in MCF-7 cells, both UPA and CDB significantly antagonized by $59 \pm 3.4\%$ and 26.5 ± 9.1 , respectively, the DEX-induced luciferase transactivation (Fig. 3 B). These results suggest a stronger anti-glucocorticoid potency of UPA in MCF-7 breast cancer cells than in HBE.

25

UPA activity on mRNA expression of GR target genes

In order to better define UPA anti-glucocorticoid effects in HBE and MCF-7 cells, mRNA expressions of various glucocorticoid responsive genes were analyzed after DEX and UPA treatment. Immediate early response 3 (IEX-1) and regulator of G-protein signaling 2 (G0S8)
30 have been implicated in cell survival under stress conditions and in G-protein signaling, respectively. IEX-1 and G0S8 were characterized as glucocorticoid responsive genes. As shown in Figure 4 A, IEX-1 mRNA expression was down-regulated by DEX and UPA did not antagonize this response in HBE and MCF-7 cells. On the other hand, UPA exerted a slight and non-significant antagonist effect on DEX-induced up-regulation of G0S8 mRNA in HBE
35 cells, whereas its anti-glucocorticoid activity was strong in MCF-7 cells ($p < 0.001$) (Fig. 4 B). Cyclin A and BCL2 genes were previously shown to be regulated 1 by DEX in vitro in osteoblasts and neuroblastomas, respectively. DEX and UPA actions were therefore

analyzed on cyclin A and BCL2 mRNA expression in HBE and MCF-7 cells. Cyclin A was differentially regulated by DEX in the two types of cells as mRNA expression was induced in HBE cells and repressed in MCF-7 cells (Fig. 4 C) (Courtin, et al.; 2011, Breast Cancer Res Treat.). In both cellular models, UPA partially inhibited DEX effect on cyclin A mRNA expression. BCL2 mRNA was down-regulated by DEX in HBE and in MCF-7 cells (Fig. 4 D). However, UPA did not reverse the DEX down-regulation of BCL2 mRNA in HBE cells, whereas it partially antagonized this effect in MCF-7 cells. UPA did not display any glucocorticoid agonist activity on these genes (Fig 4).

10 UPA action on proliferation and apoptosis

P and glucocorticoid can induce different proliferative and survival effects on normal and tumoral breast cells. In order to evaluate the role of UPA on these cellular events, proliferation and apoptosis were measured respectively by tritiated thymidine incorporation (Fig. 5) and flow cytometry (Fig. 6) in the three cellular models. UPA effects on PR were studied in HBE and T-47D, but not in MCF-7 cells, because we previously demonstrated a lack of PR functionality in this cell line (Courtin, et al.; 2011, Breast Cancer Res Treat.). UPA actions on GR were only studied in HBE and MCF-7 cells as T-47D cells used herein do not express GR (Courtin, et al.; 2011, Breast Cancer Res Treat.). In HBE cells, UPA did not inhibit the anti-proliferative effect induced by P (Fig. 5 A). Conversely, UPA inhibited the strong proliferative activity induced by DEX (Fig. 5 A). In T-47D cells, UPA reversed P anti-proliferative activity (Fig. 5B). In MCF-7 cells, UPA tended to prevent the weak anti-proliferative effects of DEX (Fig. 5C). UPA treatment alone had no effect on proliferation in any type of cells (Fig. 5). Apoptosis was measured by quantification of sub-G1 percentage of cells by flow cytometry analysis. P exerted an intense pro-apoptotic effect in HBE cells, in contrast to DEX which displayed anti-apoptotic features (Fig. 6 A). When combined with UPA, both hormones activities were reversed. Similarly, P increased the percentage of sub-G1 cells and UPA tended to inhibit this effect in T-47D cells (Fig. 6 B). In MCF-7, DEX showed a pro-apoptotic ability which was totally inhibited by UPA (Fig. 6 C). UPA had no apoptotic or survival properties by itself, but counteracted most of the P and DEX effects in HBE, T-47D, and MCF-7 cells.

Example 2: Impact of UPA on an in vivo experimental model, with normal human breast tissues xenografted in the nude mice.

35 Materials and methods

Patients

Breast tissues samples were obtained from 6 women (aged 29-42 years) undergoing surgery for reduction mammoplasty, with their informed consent according to the French law on clinical experimentation. The patients had no history of breast disease and immunohistochemical studies including a hematoxylin-phloxine-saffron (HPS) stain only
5 showed normal breast tissue.

Human breast xenografts in mice and pellet treatments

Four weeks old ovariectomized female NMRI nu/nu 20 athymic mice were purchased from Janvier laboratory (Le Genest Saint Isle, France). Breast tissues samples were obtained
10 from 6 women (aged 29-42 years). Human mammary gland tissue was cut into 2x2x2 mm fragments and 4 fragments were then subcutaneously xenografted onto the back of 4 mice per group. Four treatment groups were performed: control, E2, E2+P and E2+P+UPA. Treatments were administered by grafting steroid pellets onto the neck of each mouse. The experimental conditions were initially determined by using a dose range of
15 hormones mixed with cholesterol into the pellets. Blood sample assays were performed after 2 weeks and 4 weeks of treatment to measure plasmatic hormonal concentrations. Finally, the dose of 0.3mg for E2 and 20 mg for P and UPA were used, as it provides the expected blood concentrations. For the control groups, pellets containing only cholesterol were used. To reproduce menstrual cycle conditions, mice were grafted on the first day of experiment
20 with cholesterol, E2 and/or UPA containing pellets and on the fourteen day with cholesterol or progesterone containing pellets in control and E2 groups, or in E+P and E2+P+UPA groups, respectively. Twenty eight days after the start of experiment, mice were sacrificed. Blood was collected for each mouse, and serum was frozen at -20°C until hormone concentrations

25 Analyses.

Breast tissue fragments were collected and immediately fixed in paraformaldehyde solution for immunohistochemical analysis. All study protocols and environmental conditions of the animal rooms were approved by the French Ethic committee for the care and use of laboratory animals Charles Darwin.

30

Hormone concentration analysis

Estradiol was measured by radioimmunoassay using Clinical Assays Estradiol-2 (Sorin Biomedica Diagnostics SpA, Saluggia, Italia). Progesterone levels were evaluated by UPLC-MSMS using Acquity UPLC and Quattro Premier XE (Waters, Milford MA, USA).

35 UPA concentrations were measured using LC-MS/MS technique by MPI Research (State College, Pennsylvania, USA).

Immunohistochemical analysis

Mitotic index was calculated using the Ki67 antibody, and was determined for each breast tissue grafted into mice. Immunohistochemical analyses were performed using the BOND-MAX workstation (Leica, Nanterre, France). Paraffin sections of breast tissue xenografts 5 were de-waxed and rehydrated before antigen retrieval using citrate retrieval solution (pH 6.0) for Ki67 antibody or EDTA retrieval solution (pH 9.0) for PR and ERalpha antibodies, for 30 min. Sections were then incubated with Ki67 at 1:100 (Novocastra, NCL5 L-Ki67-MM1), PR at 1:80 (Biogenex, MU-328-UC) or ER at 1:300 (Novocastra, NCL-L6 ER-6F11) monoclonal antibodies. For signal detection, the Kit Bond Polymer Refire Detection was 10 used. Reagents were purchased from Menarini-Diagnostic (Rungis, France). A negative control (omitting the first antibody) was included in each set. For each marker, determination of the ratio of positive cells was performed on a total of 1 000 lobular and 1 000 ductal luminal cells in the 4 breast tissue fragments grafted into each mouse. For each experiment, the final percentages for each treatment were the mean of percentages obtained in the four 15 mice per group.

Results

UPA action on breast tissue proliferation

20 The inventors developed an in vivo model to study UPA in long term administration on breast tissue. Human normal breast tissues samples were xenografted in athymic mice treated with E2, or E2+P, or E2+P+UPA, or cholesterol (control) (see experimental procedure). In order to reproduce the chronology of E2 and P secretions occurring in human female menstrual cycle, E2 pellets were grafted at the beginning of the experiment, whereas P pellets were 25 grafted at the fourteenth day. UPA pellets were grafted at the beginning of the experiment to mimic a chronic treatment. E2, P, and UPA concentrations in mice serum were measured to validate the treatment method. Average E2 concentrations in mice serum was 36.88 ± 4.25 pg/ml (mean \pm SEM) corresponding to the low range of the physiological E2 levels reported in follicular phase. The P level was 13.05 ± 1.14 ng/ml (mean \pm SEM) equivalently to the 30 average P plasma levels in women during the mid-luteal phase. UPA concentration was 63.49 ± 10.46 ng/ml (mean \pm SEM) in the same range that observed in clinical use. Hormones levels were undetectable in control mice (E2 < 0.8 pg/ml; P < 0.4 ng/ml; UPA < 0.5 ng/ml). As shown in Fig 7 A, estradiol receptor (ER) and PR expressions were maintained in the treated engrafted breast fragments at the end of the experiment compared to the original 35 breast tissue before grafts. We analyzed the mitotic Ki67 expression marker in order to determine the breast tissue proliferative activity in glandular lobules and ducts according to the treatment (Fig. 7 B, C). In the control group of grafted tissues, the rate of mitotic cells was

low and homogeneous, at $1.7 \pm 0.4\%$ in the lobules and $1.8 \pm 0.6\%$ in ducts. The proliferative activity was slightly but not significantly increased in lobules of E2 treated group when compared to the control group (Fig. 7 B, C). However, in ducts, the mitotic index was significantly elevated in E2 treated group (3.1 ± 0.7 fold induction compared to the control group, $p < 0.05$). No significant difference was observed in E2+P and E2+P+UPA groups compared to E2 treatment in lobular and ductal structures (Fig. 7 B, C). These results strongly suggest that proliferative activity in breast tissue is predominantly mediated by E2. We also showed that UPA does not significantly influence the proliferation rate of normal epithelial breast cells.

10

Example 3: Impact of UPA on BRCA1 human breast tissues xenografted on nude mice.Materials and methods

Mutated breast tissue samples were obtained from 5 women (36-57 years old) carrying mutations for BRCA1.

Xenografts were performed as described above (Example 2). Usually, 4 breast tissue fragments from a patient not carrying BRCA1 mutations and 4 breast fragments from a patient carrying BRCA1 mutations were grafted in the same mouse, on the left and right part of the spine respectively. The same groups were formed.

20

Results

The inventors used the same model as described in the above section to investigate the effects of hormones on the proliferation of human breast tissue collected in patient carrying BRCA1 mutations. A higher variability regarding ER α expression was observed in both lobular and ductal structures of patients with BRCA1 mutations before being grafted. The basal expression level of PR was lower in patients carrying BRCA1 mutation compared to patients with a wild type allele in the lobular structures. In contrast the BRCA1 status did not impact PR expression level in ductal structures. The proliferative status measure through the expression of Ki67 was higher in lobular cells from BRCA1 patients, but lower in the ductal cells, compared to wild type patients.

After the graft, but in the absence of treatment, no significant statistical modification of ER α expression was observed, whilst PR expression level in both lobular and ductal structures tended to be even lower than before the graft. In both structures the absence of hormonal stimulation resulted in decreased proliferation.

35 28 days treatment with E2, P, E2+P or E2+P+UPA did not result in significant variation of ER α expression level. Trends towards increased PR expression were observed in breast fragments grafted in mice receiving either E2, or E2+P or E2+P+UPA. This increase was

more particularly observed in the lobular structures of 3 patients carrying the BRCA1 mutation.

In each case where proliferation was stimulated by the combination of E2+P treatment, UPA exhibited antiproliferative activity (See Figures 8 and 9).

5

Example 4: Impact of UPA on human mammary breast tumor growth xenografted into nude mice

Material and methods

The xenograft model used in this study was HBCx-34. HBCx-34 is a mammary ductal carcinoma with wild type P53, no HER2 overexpression and PR and ER α overexpression. The tumor is highly responsive to adriamycin/cyclophosphamide and responsive to docetaxel and capecitabine. HBCx-34 has got no cachectic property.

Tumograft model induction: Pretreatment procedures

HBCx-34 tumors (P14.0.0/2) were transplanted subcutaneously onto 5-10 mice (donor mice, 10 passage (n-1), female athymic nude mice (Hsd:Athymic Nude-Fox1nu), 6- to 9-week-old, Harlan Laboratories (Gannat, France)). When these tumors reached 1000 to 2000 mm³ (between 60 to 78 days), donor mice were sacrificed by cervical dislocation, tumors were aseptically excised and dissected. After removing necrotic areas, tumors were cut into fragments measuring approximately 20 mm³ and transferred for 10 min maximum in sterile

20 DMEM/F12culture medium, without any additional components, before grafting.

Healthy mice aged 6 to 9 weeks and weighing at least 20 g were included in the study. Mice were anaesthetized with ketamine/xylazine, and then skin was aseptized with a chlorhexidine solution, incised at the level of the interscapular region, and a 20 mm³ tumor fragment was placed in the subcutaneous tissue. Skin was closed with clips.

25 Tumograft model induction: Treatment phase

Mice were allocated to different groups according to their tumor volume to give homogenous mean and median tumor volume in each treatment arms. Ten mice/group, with HBCx-34 tumors between 75 and 144 mm³, were randomized according to their tumor volume into experimental groups, and treatments (UPA 130 mg/kg, p.o. or control vehicle) were initiated 30 33 days post implantation of the tumor for a total duration of 42 days.

Analyses

Mice were observed for physical appearance, behavior and clinical changes. Animals were weighted biweekly during all the experimental period. Toxicity of the different treatments was determined as body weight loss.

35 Tumor volume was evaluated biweekly during all the experimental period. Tumors were collected at the end of treatment, weight and processed for analysis.

Results

UPA treatment was well tolerated and no significant body weight loss was recorded during the study. No treatment related clinical observation was reported during the experimental period.


Mean tumor volumes (TV) at initiation of treatment were 112.2 ± 7.1 and $102.4 \pm 7.7 \text{ mm}^3$ in 5 control and UPA-treated groups respectively.

At the end of treatment, a 4.63-fold increase in tumor volume was measured in the control group, with a mean tumor volume of $625.0 \pm 108.6 \text{ mm}^3$.

UPA exerted an antitumor activity; the mean TV was increased 2.17-fold in the UPA treated mice (mean TV at sacrifice : $332.0 \pm 61.0 \text{ mm}^3$). The ratio of TV in treated vs. control group 10 (T/C) was 53%.

CLAIMS

1. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor in a patient.
5
2. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to claim 1, wherein the patient carries a mutation in BRCA1 and/or BRCA2 gene.
- 10 3. Ulipristal acetate, or any of its metabolites, for use in preventing a breast tumor according to claim 1, wherein the patient carries a mutation in BRCA1 and/or BRCA2 gene.
- 15 4. Ulipristal acetate, or any of its metabolites, for use in treating a breast tumor according to any of claims 1 or 2, wherein the breast tumor is a breast carcinoma.
15
5. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to any of claims 1 to 4, wherein Ulipristal acetate, or any of its metabolites, is to be administered orally.
20
6. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to any of claims 1 to 4, wherein Ulipristal acetate, or any of its metabolites, is to be administered intravenously.
- 25 7. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to any of claims 1 to 4, wherein Ulipristal acetate, or any of its metabolites, is to be administered by the vaginal or intrauterine route.
8. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to any of claims 1 to 7, wherein the patient is a female patient.
30
9. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to any of claims 1 to 8, wherein the breast tumor is a carcinoma.
- 35 10. Ulipristal acetate, or any of its metabolites, for use in preventing or treating a breast tumor according to any of claims 1 to 8, wherein the breast tumor is a benign tumor.






FIGURE 1 cont



A FASN**B cyclin A**

Figure 2




C *BCL2*D *ALPL*

FIGURE 2 cont

A HBE



B MCF-7




Figure 3






A IEX-1

Figure 4

B GOS8**C cyclin A****D BCL2**

5/8








Figure 6






FIGURE 8




FIGURE 9

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/060801

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K31/57 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>WIEHLE ET AL.: "ANTI-PROGESTINS SUPPRESS THE GROWTH OF ESTABLISHED TUMORS INDUCED BY 7,12-DIMETHYLBENZ(A)ANTHRACENE: COMPARISON BETWEEN RU486 AND A NEW 21-SUBSTITUTED-19-NOR-PROGESTIN", ONCOLOGY REPORTS, vol. 18, no. 1, 30 July 2007 (2007-07-30), - 30 July 2007 (2007-07-30), pages 167-174, XP002683518, the whole document</p> <p style="text-align: center;">-----</p> <p style="text-align: center;">-/-</p>	1-10

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
14 June 2013	21/06/2013
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Trifilieff-Riolo, S

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/060801

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ATTARDI BARBARA J ET AL: "CDB-4124 and its putative monodemethylated metabolite, CDB-4453, are potent antiprogestins with reduced antiglucocorticoid activity: In vitro comparison to mifepristone and CDB-2914", MOLECULAR AND CELLULAR ENDOCRINOLOGY, ELSEVIER IRELAND LTD, IE, vol. 188, no. 1-2, 25 February 2002 (2002-02-25), pages 111-123, XP002496575, ISSN: 0303-7207, DOI: 10.1016/S0303-7207(01)00743-2 cited in the application page 123, last line - left-hand column -----	1-10
A	US 2007/213306 A1 (HAUSKNECHT RICHARD [US]) 13 September 2007 (2007-09-13) claims 1,6 -----	1-10
1		

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/EP2013/060801

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2007213306 A1	13-09-2007	US 2007213306 A1 US 2012115831 A1 WO 2007103510 A2	13-09-2007 10-05-2012 13-09-2007