

(19) **DANMARK**

(10) **DK/EP 2507372 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 9/34 (2006.01)** **C 12 P 7/06 (2006.01)** **C 12 P 19/14 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2015-05-11**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2015-02-25**
- (86) Europæisk ansøgning nr.: **10785298.0**
- (86) Europæisk indleveringsdag: **2010-11-30**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-10-10**
- (86) International ansøgning nr.: **US2010058438**
- (87) Internationalt publikationsnr.: **WO2011066576**
- (30) Prioritet: **2009-11-30 US 264977 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Danmark**
Novozymes North America, Inc., 77 Perry Chapel Church Road, Franklinton, North Carolina 27525, USA
- (72) Opfinder: **LANDVIK, Sara, Plantagekrogen 8, 2950 Vedbæk, Danmark**
MORANT, Marc Dominique, Oerestadsboulevard 61D 7th, 2300 København S, Danmark
AYABE, Keiichi, Konakadai-cho 323, Inage-ky, Chiba-shi, Chiba 263-0044, Japan
COWARD-KELLY, Guillermo, 3216 Coach Lantern Avenue, Wake Forest, North Carolina 27587, USA
- (74) Fuldmægtig i Danmark: **Novozymes A/S Patents, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (54) Benævnelse: **Polypeptider med glucoamylaseaktivitet og polynukleotider, der koder for dem**
- (56) Fremdragne publikationer:
WO-A2-2006/069289
WO-A2-2006/069290
WO-A2-2007/134207
WO-A2-2007/144424

DESCRIPTION

Reference to a Sequence Listing

[0001] This application contains a Sequence Listing in computer readable form.

Reference to a Deposit of Biological Material

[0002] This application contains a reference to a deposit of biological material. For complete information see last paragraph of the description.

Background of the Invention

Field of the Invention

[0003] The present invention relates to polypeptides having glucoamylase activity and polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods for producing and using the polypeptides, and to the use of glucoamylases of the invention for starch conversion to producing fermentation products, such as ethanol, and syrups, such as glucose. The invention also relates to a composition comprising a glucoamylase of the invention.

Description of the Related Art

[0004] Glucoamylase (1,4- α -D-glucan glucohydrolase, EC 3.2.1.3) is an enzyme, which catalyzes the release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules. Glucoamylases are produced by several filamentous fungi and yeast, with those from *Aspergillus* being commercially most important.

[0005] Commercially, glucoamylases are used to convert starchy material, which is already partially hydrolyzed by an α -amylase, to glucose. The glucose may then be converted directly or indirectly into a fermentation product using a fermenting organism. Examples of commercial fermentation products include alcohols (e.g., ethanol, methanol, butanol, 1,3-propanediol); organic acids (e.g., citric acid, acetic acid, itaconic acid, lactic acid, gluconic acid, gluconate, lactic acid, succinic acid, 2,5-diketo-D-gluconic acid); ketones (e.g., acetone); amino acids (e.g., glutamic acid); gases (e.g., H_2 and CO_2), and more complex compounds, including, for example, antibiotics (e.g., penicillin and tetracycline); enzymes; vitamins (e.g., riboflavin, B₁₂, beta-carotene); hormones, and other compounds which are difficult to produce synthetically. Fermentation processes are also commonly used in the consumable alcohol (e.g., beer and wine).

[0006] The end product may also be a syrup. For instance, the end product may be glucose, but may also be converted, e.g., by a glucose isomerase to fructose or a mixture composed almost equally of glucose and fructose. This mixture, or a mixture further enriched with fructose, is the most commonly used high fructose corn syrup (HFCS) commercialized throughout the world.

[0007] It is an object of the present invention to provide polypeptides having glucoamylase activity and polynucleotides encoding the polypeptides and which provide a high yield in fermentation product production processes, such as ethanol production processes, including one-step ethanol fermentation processes from un-gelatinized raw (or uncooked) starch.

[0008] Uniprot: B0CVJ1 discloses a polypeptide from *Laccaria bicolor* and WO 2006/069289 describes a glucoamylase from *Trametes cingulata*.

Summary of the Invention

[0009] Polypeptides produced by the fungus *Pycnoporus sanguineus* and having glucoamylase activity have been identified and characterized.

[0010] Accordingly, the present invention relates in a first aspect to an isolated polypeptide having glucoamylase activity, selected from the group consisting of: (a) a polypeptide comprising an amino acid sequence having preferably at least 90% more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity to amino acids 19 to 573 of SEQ ID NO: 2, of SEQ ID NO: 4 or of SEQ ID NO: 6; (b) a polypeptide comprising an amino acid sequence having preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity to the catalytic domain shown as amino acids 22 to 476 of SEQ ID NO: 2, of SEQ ID NO: 4 or of SEQ ID NO: 6; (c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity to the coding sequence of SEQ ID NO: 1, of SEQ ID NO: 3 or of SEQ ID NO: 5 encoding amino acids 19-573 of SEQ ID NO: 2, of SEQ ID NO: 4 or of SEQ ID NO: 6.

[0011] The present invention relates in a second aspect to an isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of the first aspect.

[0012] In further aspects the invention relates to a nucleic acid construct, a recombinant expression vector, a recombinant host cell, a transgenic plant, a plant part or plant cell comprising the polynucleotide of the second aspect.

[0013] In still further aspects the invention relates to a method of producing the polypeptide, uses of the polypeptide and a composition comprising an alpha-amylase and the polypeptide.

Definitions

[0014] **Glucoamylase:** The term glucoamylase (1,4-alpha-D-glucan glucohydrolase, EC 3.2.1.3) is defined as an enzyme, which catalyzes the release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules. For purposes of the present invention, glucoamylase activity is determined according to the procedure described in the 'Materials & Methods'-section below.

[0015] The polypeptides of the present invention have at least 20%, preferably at least 40%, preferably at least 45%, more preferably at least 50%, preferably at least 55%, more preferably at least 60%, preferably at least 65%, more preferably at least 70%, preferably at least 75%, more preferably at least 80%, preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 100% of the glucoamylase activity of the mature polypeptide of SEQ ID NO: 2 or a homologous sequence thereof, of SEQ ID NO: 4 or a homologous sequence thereof, or of SEQ ID NO: 6 or a homologous sequence thereof.

[0016] **Isolated polypeptide:** The term "isolated polypeptide" as used herein refers to a polypeptide that is isolated from a source. Preferably, the polypeptide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by SDS-PAGE.

[0017] **Substantially pure polypeptide:** The term "substantially pure polypeptide" denotes herein a polypeptide preparation that contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polypeptide material with which it is natively or recombinantly associated. It is, therefore, preferred that the substantially pure polypeptide is at least 92% pure, preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 96% pure, more preferably at least 97% pure, more preferably at least 98% pure, even more preferably at least 99%, most preferably at least 99.5% pure, and even most preferably 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides of the present invention are preferably in a substantially pure form, *i.e.*, that the polypeptide preparation is essentially free of other polypeptide material with which it is natively or recombinantly associated. This can be accomplished, for example, by preparing the polypeptide by well-known recombinant methods or by classical purification methods.

[0018] Mature polypeptide: The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. In one aspect, the mature polypeptide is amino acids 19 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 based on the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6) that predicts amino acids 1 to 18 of SEQ ID NO: 2, SEQ ID NO: 4, and SEQ ID NO: 6 are signal peptides. Preferably, the mature polypeptide is amino acids 19 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. The sequence defined by amino acids 22 to 476 of SEQ ID NO: 2, SEQ ID NO: 4, and SEQ ID NO: 6 is the catalytic domain. The sequence defined by amino acids 479 to 573 of SEQ ID NO: 2, SEQ ID NO: 4, and SEQ ID NO: 6 is a starch binding domain.

[0019] Mature polypeptide coding sequence: The term "mature polypeptide coding sequence" is defined herein as a nucleotide sequence that encodes a mature polypeptide having glucoamylase activity. Preferably, the mature polypeptide coding sequence is nucleotides defined by positions 55 to 159, 229 to 505, 573 to 877, 932 to 1207, 1269 to 1731, 1800 to 1895, 1962 to 2104 of SEQ ID NO: 1, or defined by positions 55 to 159, 229 to 504, 571 to 876, 942 to 1217, 1276 to 1738, 1806 to 1901, 1960 to 2102 of SEQ ID NO: 3 or of SEQ ID NO: 5.

[0020] Identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity".

[0021] For purposes of the present invention, the degree of identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends in Genetics 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0022] For purposes of the present invention, the degree of identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *supra*), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0023] Homologous sequence: The term "homologous sequence" is defined herein as a nucleotide sequence/polypeptide sequence having a degree of identity to the mature polypeptide encoding part of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 or to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, respectively, of at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, or even at least 99%.

[0024] Polypeptide fragment: The term "polypeptide fragment" is defined herein as a polypeptide having one or more (several) amino acids deleted from the amino and/or carboxyl terminus of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a homologous sequence thereof; wherein the fragment has glucoamylase activity. Preferably, a fragment contains at least 500 amino acid residues, more preferably at least 450 amino acid, and most preferably at least 400 amino acid residues, of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, or a homologous sequence thereof. A particular fragment is the sequence defined by amino acids 22 to 476 of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6 which comprise the catalytic domain of the polypeptide of the invention.

[0025] Subsequence: The term "subsequence" is defined herein as a nucleotide sequence having one or more (several) nucleotides deleted from the 5' and/or 3' end of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5; or a homologous sequence thereof; wherein the subsequence encodes a polypeptide fragment having glucoamylase activity. Preferably, a subsequence contains at least 1500 nucleotides, more preferably at least 1400 nucleotides, and most preferably at least 1200 nucleotides of the mature polypeptide coding sequence of SEQ ID NO: 1, of SEQ ID NO: 3 or of SEQ ID NO: 5; or a homologous sequence thereof.

[0026] Allelic variant: The term "allelic variant" denotes herein any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

[0027] Isolated polynucleotide: The term "isolated polynucleotide" as used herein refers to a polynucleotide that is isolated from a source. Preferably, the polynucleotide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by agarose electrophoresis.

[0028] Substantially pure polynucleotide: The term "substantially pure polynucleotide" as used herein refers to a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered protein production systems. Thus, a substantially pure polynucleotide contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polynucleotide material with which it is natively or recombinantly associated. A substantially pure polynucleotide may, however, include naturally occurring 5' and 3' untranslated regions, such as promoters and terminators. It is preferred that the substantially pure polynucleotide is at least 90% pure, preferably at least 92% pure, more preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, even more preferably at least 98% pure, most preferably at least 99%, and even most preferably at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form, *i.e.*, that the polynucleotide preparation is essentially free of other polynucleotide material with which it is natively or recombinantly associated. The polynucleotides may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

[0029] Coding sequence: When used herein the term "coding sequence" means a nucleotide sequence, which directly specifies the amino acid sequence of its protein product. The boundaries of the coding sequence are generally determined by an open reading frame, which usually begins with the ATG start codon or alternative start codons such as GTG and TTG and ends with a stop codon such as TAA, TAG, and TGA. The coding sequence may be a DNA, cDNA, synthetic, or recombinant nucleotide sequence.

[0030] cDNA: The term "cDNA" is defined herein as a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic cell. cDNA lacks intron sequences that are usually present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps before appearing as mature spliced mRNA. These steps include the removal of intron sequences by a process called splicing. cDNA derived from mRNA lacks, therefore, any intron sequences.

[0031] Nucleic acid construct: The term "nucleic acid construct" as used herein refers to a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or which is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic. The term nucleic acid construct is synonymous with the term "expression cassette" when the nucleic acid construct contains the control sequences required for expression of a coding sequence of the present invention.

[0032] Control sequences: The term "control sequences" is defined herein to include all components necessary for the expression of a polynucleotide encoding a polypeptide of the present invention. Each control sequence may be native or foreign to the nucleotide sequence encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleotide sequence encoding a polypeptide.

[0033] Operably linked: The term "operably linked" denotes herein a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of the polynucleotide sequence such that the control sequence directs the expression of the coding sequence of a polypeptide.

[0034] Expression: The term "expression" includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

[0035] Expression vector: The term "expression vector" is defined herein as a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide of the present invention and is operably linked to additional nucleotides that provide for its expression.

[0036] Host cell: The term "host cell", as used herein, includes any cell type that is susceptible to transformation, transfection, transduction, and the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention.

[0037] Modification: The term "modification" means herein any chemical modification of the polypeptide consisting of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a homologous sequence thereof; as well as genetic manipulation of the DNA encoding such a polypeptide. The modification can be a substitution, a deletion and/or an insertion of one or more (several) amino acids as well as replacements of one or more (several) amino acid side chains.

[0038] Variant: When used herein, the term "variant" means a polypeptide having glucoamylase activity comprising an alteration, *i.e.*, a substitution, insertion, and/or deletion of one or more (several) amino acid residues at one or more (several) positions. A substitution means a replacement of an amino acid occupying a position with a different amino acid; a deletion means removal of an amino acid occupying a position; and an insertion means adding 1-3 amino acids adjacent to an amino acid occupying a position.

Detailed Description of the Invention

Polypeptides Having Glucoamylase Activity

[0039] In a first aspect, the present invention relates to isolated polypeptides comprising an amino acid sequence having a degree of identity to the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6 of preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity, which have glucoamylase activity (hereinafter "homologous polypeptides"). Preferably, the homologous polypeptides have an amino acid sequence that differs by ten amino acids, preferably by five amino acids, more preferably by four amino acids, even more preferably by three amino acids, most preferably by two amino acids, and even most preferably by one amino acid from the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6.

[0040] A polypeptide of the present invention preferably comprises the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a fragment thereof having glucoamylase activity. In another preferred aspect, the polypeptide comprises the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a fragment thereof having glucoamylase activity. In another preferred aspect, the polypeptide consists of the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a fragment thereof having glucoamylase activity.

[0041] The present disclosure relates to isolated polypeptides having glucoamylase activity that are encoded by polynucleotides that hybridize under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5, (ii) the cDNA sequence contained in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5, (iii) a subsequence of (i) or (ii), or (iv) a full-length complementary strand of (i), (ii), or (iii) (J. Sambrook, E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning, A Laboratory Manual, 2d edition, Cold Spring Harbor, New York). A subsequence of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 contains at least 100 contiguous nucleotides or preferably at least 200 contiguous nucleotides. Moreover, the subsequence may encode a polypeptide fragment having glucoamylase activity. Preferably, the complementary strand is the full-length complementary strand of the mature polypeptide coding sequence of SEQ ID NO: 1, of SEQ ID NO: 3 or of SEQ ID NO: 5.

[0042] The nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5; or a subsequence thereof; as well as the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a fragment thereof; may be used to design nucleic acid probes to identify and clone DNA encoding polypeptides having glucoamylase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the

corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 14, preferably at least 25, more preferably at least 35, and most preferably at least 70 nucleotides in length. It is, however, preferred that the nucleic acid probe is at least 100 nucleotides in length. For example, the nucleic acid probe may be at least 200 nucleotides, preferably at least 300 nucleotides, more preferably at least 400 nucleotides, or most preferably at least 500 nucleotides in length. Even longer probes may be used, e.g., nucleic acid probes that are preferably at least 600 nucleotides, more preferably at least 700 nucleotides, even more preferably at least 800 nucleotides, or most preferably at least 900 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{33}P , ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

[0043] A genomic DNA or cDNA library prepared from such other strains may, therefore, be screened for DNA that hybridizes with the probes described above and encodes a polypeptide having glucoamylase activity. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that is homologous with SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5; or a subsequence thereof; the carrier material is preferably used in a Southern blot.

[0044] For purposes of the present disclosure, hybridization indicates that the nucleotide sequence hybridizes to a labeled nucleic acid probe corresponding to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5; the cDNA sequence contained in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5; its full-length complementary strand; or a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film.

[0045] Preferably, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5.

[0046] In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 2, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is the polynucleotide sequence contained in plasmid in *E. coli* strain DSM 23221, wherein the polynucleotide sequence thereof encodes a polypeptide having glucoamylase activity. In another preferred aspect, the nucleic acid probe is the mature polypeptide coding region contained in plasmid in *E. coli* strain DSM 23221.

[0047] In another preferred aspect, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 3. In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 4, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 3.

[0048] In another preferred aspect, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 5. In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 6, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 5.

[0049] For long probes of at least 100 nucleotides in length, very low to very high stringency conditions are defined as prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures for 12 to 24 hours optimally.

[0050] For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS preferably at 45°C (very low stringency), more preferably at 50°C (low stringency), more preferably at 55°C (medium stringency), more preferably at 60°C (medium-high stringency), even more preferably at 65°C (high stringency), and most preferably at 70°C (very high stringency).

[0051] For short probes that are about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at about 5°C to about 10°C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, Proceedings of the National Academy of Sciences USA 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures for 12 to 24 hours optimally.

[0052] For short probes that are about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in

6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated T_m .

[0053] In a third aspect, the present invention relates to isolated polypeptides having glucoamylase activity encoded by polynucleotides comprising or consisting of nucleotide sequences that have a degree of identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 of at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, most preferably at least 95%, and even most preferably 96%, 97%, 98%, or 99%, which encode an active polypeptide. See polynucleotide section herein.

[0054] The present invention also relates to artificial variants comprising a substitution, deletion, and/or insertion of one or more (or several) amino acids of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6; or a homologous sequence thereof. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

[0055] Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, In, *The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

[0056] In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline, and alpha-methyl serine) may be substituted for amino acid residues of a wild-type polypeptide. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, and preferably, are commercially available, and include pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

[0057] Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

[0058] Essential amino acids in the parent polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for glucoamylase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, *Science* 255: 306-312; Smith et al., 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver et al., 1992, *FEBS Lett.* 309: 59-64. The identities of essential amino acids can also be inferred from analysis of identities with polypeptides that are related to a polypeptide according to the invention.

[0059] Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, *Biochem.* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire et al., 1986, *Gene* 46: 145; Ner et al., 1988, *DNA* 7: 127).

[0060] Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard

methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

[0061] The total number of amino acid substitutions, deletions and/or insertions of the mature polypeptide, such as amino acids 19 to 573 of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, or the catalytic domain, such as amino acids 22 to 476 of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, is 10, preferably 9, more preferably 8, more preferably 7, more preferably at most 6, more preferably 5, more preferably 4, even more preferably 3, most preferably 2, and even most preferably 1.

Sources of Polypeptides Having glucoamylase Activity

[0062] A polypeptide of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a nucleotide sequence is produced by the source or by a strain in which the nucleotide sequence from the source has been inserted. Preferably, the polypeptide obtained from a given source is secreted extracellularly.

[0063] A polypeptide having glucoamylase activity of the present invention may also be bacterial polypeptide, or a yeast polypeptide, or more preferably a filamentous fungal polypeptide such as an *Acremonium*, *Agaricus*, *Alternaria*, *Artomyces*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Gloeophyllum*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotriconympha*, *Pycnoporus*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* polypeptide having glucoamylase activity.

[0064] In a more preferred aspect, the polypeptide is a *Pycnoporus* sp. polypeptide having glucoamylase activity. Particularly the *Pycnoporus* sp. is *Pycnoporus sanguineus*.

[0065] It will be understood that for the aforementioned species the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

[0066] Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

[0067] Furthermore, such polypeptides may be identified and obtained from other sources including microorganisms isolated from nature (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms from natural habitats are well known in the art. The polynucleotide may then be obtained by similarly screening a genomic or cDNA library of such a microorganism. Once a polynucleotide sequence encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are well known to those of ordinary skill in the art (see, e.g., Sambrook *et al.*, 1989, *supra*).

[0068] Polypeptides of the present invention also include fused polypeptides or cleavable fusion polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by fusing a nucleotide sequence (or a portion thereof) encoding another polypeptide to a nucleotide sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fused polypeptide is under control of the same promoter(s) and terminator.

[0069] A fusion polypeptide can further comprise a cleavage site. Upon secretion of the fusion protein, the site is cleaved releasing the polypeptide having glucoamylase activity from the fusion protein. Examples of cleavage sites include, but are not limited to, a Kex2 site that encodes the dipeptide Lys-Arg (Martin *et al.*, 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-76; Svetina *et al.*, 2000, *J. Biotechnol.* 76: 245-251; Rasmussen-Wilson *et al.*, 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward *et al.*, 1995, *Biotechnology* 13: 498-503; and Contreras *et al.*, 1991, *Biotechnology* 9: 378-381), an Ile-(Glu or Asp)-Gly-Arg site, which is cleaved by a Factor Xa protease after the arginine residue (Eaton *et al.*, 1986, *Biochem.* 25: 505-512); an Asp-Asp-Asp-Asp-Lys

site, which is cleaved by an enterokinase after the lysine (Collins-Racie et al., 1995, *Biotechnology* 13: 982-987); a His-Tyr-Glu site or His-Tyr-Asp site, which is cleaved by Genenase I (Carter et al., 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248); a Leu-Val-Pro-Arg-Gly-Ser site, which is cleaved by thrombin after the Arg (Stevens, 2003, *Drug Discovery World* 4: 35-48); a Glu-Asn-Leu-Tyr-Phe-Gln-Gly site, which is cleaved by TEV protease after the Gln (Stevens, 2003, *supra*); and a Leu-Glu-Val-Leu-Phe-Gln-Gly-Pro site, which is cleaved by a genetically engineered form of human rhinovirus 3C protease after the Gln (Stevens, 2003, *supra*).

Polynucleotides

[0070] The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that encode polypeptides having glucoamylase activity of the present invention.

[0071] Preferably, the nucleotide sequence comprises or consists of SEQ ID NO: 1. In another more preferred aspect, the nucleotide sequence comprises or consists of the sequence contained in plasmid which is contained in *E. coli* DSM 23221. In another preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence of SEQ ID NO: 1. In another more preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence contained in plasmid contained in *E. coli* DSM 23221.

[0072] The present invention also encompasses nucleotide sequences that encode polypeptides comprising or consisting of the amino acid sequence of SEQ ID NO: 2 or the mature polypeptide thereof, which differ from SEQ ID NO: 1 or the mature polypeptide coding sequence thereof by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 1 that encode fragments of SEQ ID NO: 2 that have glucoamylase activity.

[0073] The present invention also relates to mutant polynucleotides comprising or consisting of at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, in which the mutant nucleotide sequence encodes the mature polypeptide of SEQ ID NO: 2.

[0074] Preferably, the nucleotide sequence comprises or consists of SEQ ID NO: 3. In another preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence of SEQ ID NO: 3. The present invention also encompasses nucleotide sequences that encode polypeptides comprising or consisting of the amino acid sequence of SEQ ID NO: 4 or the mature polypeptide thereof, which differ from SEQ ID NO: 3 or the mature polypeptide coding sequence thereof by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 3 that encode fragments of SEQ ID NO: 4 that have glucoamylase activity.

[0075] The present invention also relates to mutant polynucleotides comprising or consisting of at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 3, in which the mutant nucleotide sequence encodes the mature polypeptide of SEQ ID NO: 4.

[0076] Preferably, the nucleotide sequence comprises or consists of SEQ ID NO: 5. In another preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence of SEQ ID NO: 5. The present invention also encompasses nucleotide sequences that encode polypeptides comprising or consisting of the amino acid sequence of SEQ ID NO: 6 or the mature polypeptide thereof, which differ from SEQ ID NO: 5 or the mature polypeptide coding sequence thereof by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 5 that encode fragments of SEQ ID NO: 6 that have glucoamylase activity.

[0077] The present invention also relates to mutant polynucleotides comprising or consisting of at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 5, in which the mutant nucleotide sequence encodes the mature polypeptide of SEQ ID NO: 6.

[0078] The techniques used to isolate or clone a polynucleotide encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the polynucleotides of the present invention from such genomic DNA can be effected, e.g., by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, e.g., Innis et al., 1990, *PCR: A Guide to Methods and Application*, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligated activated transcription (LAT) and nucleotide sequence-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Penicillium*, or another or related organism and thus, for example, may be an

allelic or species variant of the polypeptide encoding region of the nucleotide sequence.

[0079] The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that have a degree of identity to the mature polypeptide coding sequence of SEQ ID NO: 1, of SEQ ID NO: 3 or of SEQ ID NO: 5 of preferably at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% identity, which encode an active polypeptide.

[0080] Modification of a nucleotide sequence encoding a polypeptide of the present invention may be necessary for the synthesis of polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, e.g., artificial variants that differ in specific activity, thermostability, pH optimum, or the like. The variant sequence may be constructed on the basis of the nucleotide sequence presented as the mature polypeptide coding sequence of SEQ ID NO: 1, e.g., a subsequence thereof, and/or by introduction of nucleotide substitutions that do not give rise to another amino acid sequence of the polypeptide encoded by the nucleotide sequence, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford et al., 1991, *Protein Expression and Purification* 2: 95-107.

[0081] It will be apparent to those skilled in the art that such substitutions can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by an isolated polynucleotide of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, e.g., Cunningham and Wells, 1989, *supra*). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for glucoamylase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labeling (see, e.g., de Vos et al., 1992, *supra*; Smith et al., 1992, *supra*; Wlodaver et al., 1992, *supra*).

[0082] The present disclosure also relates to isolated polynucleotides encoding polypeptides of the present invention, which hybridize under very low stringency conditions, preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5, (ii) the cDNA sequence contained in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5, or (iii) a full-length complementary strand of (i) or (ii); or allelic variants and subsequences thereof (Sambrook et al., 1989, *supra*), as defined herein. Preferably, the complementary strand is the full-length complementary strand of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5.

[0083] The present disclosure also relates to isolated polynucleotides obtained by (a) hybridizing a population of DNA under very low, low, medium, medium-high, high, or very high stringency conditions with (i) the mature polypeptide coding sequence SEQ ID NO: 1, of SEQ ID NO: 3 or of SEQ ID NO: 5, (ii) the cDNA sequence contained in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having glucoamylase activity. Preferably, the complementary strand is the full-length complementary strand of the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5.

Hybrid enzymes

[0084] The present disclosure also relates to hybrid enzymes comprising a catalytic domain having enzyme activity (e.g., starch degrading enzyme activity, such as alpha-amylase, amylopullulanase, beta-amylase, CGTase, glucoamylase, isoamylase, maltogenic amylase, or pullulanase activity), and a carbohydrate-binding module (CBM). The hybrid enzyme may further comprise a linker.

[0085] The hybrid may be produced by fusing a first DNA sequence encoding a catalytic domain and a second DNA sequence encoding a carbohydrate-binding module, or the hybrid may be produced as a completely synthetic gene based on knowledge of the amino acid sequences of suitable CBMs, linkers and catalytic domains.

[0086] The term "hybrid enzyme" (also referred to as "fusion protein", "hybrid", hybrid polypeptide" or "hybrid protein") is used herein to characterize the hybrid polypeptides of the invention comprising a catalytic module having enzyme activity (e.g., starch degrading enzyme activity, such as alpha-amylase, amylopullulanase, beta-amylase, CGTase, glucoamylase, isoamylase, maltogenic amylase, or pullulanase activity) and a carbohydrate-binding module wherein the catalytic domain and the carbohydrate-binding module are derived from different sources. The term "source" includes, but is not limited to, a parent enzyme or a variant thereof, e.g., an amylase or glucoamylase, or other catalytic activity comprising a suitable catalytic module and/or a suitable CBM and/or a suitable linker. However the CBM may also be derived from a polypeptide having no catalytic activity. The catalytic domain and the carbohydrate binding module may be derived from the same microbial strain, from strains within the same species, from closely related species or less related organisms. Preferably the catalytic domain and the carbohydrate binding module of the hybrids are derived from different sources, e.g., from different enzymes from the same strain and/or species, or, e.g., from strains within different species.

[0087] In one aspect the hybrid enzyme comprises the CBM (also known as a carbohydrate binding domain or CBD) according to the invention and a catalytic domain. The catalytic domain is in a particular embodiment a glucoamylase catalytic domain.

Nucleic Acid Constructs

[0088] The present invention also relates to nucleic acid constructs comprising an isolated polynucleotide of the present invention operably linked to one or more (several) control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

[0089] An isolated polynucleotide encoding a polypeptide of the present invention may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide's sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotide sequences utilizing recombinant DNA methods are well known in the art.

[0090] The control sequence may be an appropriate promoter sequence, a nucleotide sequence that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter sequence contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any nucleotide sequence that shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0091] Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase); and mutant, truncated, and hybrid promoters thereof.

[0092] In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos et al., 1992, Yeast 8: 423-488.

[0093] The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleotide sequence encoding the polypeptide. Any terminator that is functional in the host cell of choice may be used in the present invention.

[0094] Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase,

Aspergillus niger glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.

[0095] Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*.

[0096] The control sequence may also be a suitable leader sequence, a nontranslated region of an mRNA that is important for translation by the host cell. The leader sequence is operably linked to the 5' terminus of the nucleotide sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used in the present invention.

[0097] Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

[0098] Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

[0099] The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3' terminus of the nucleotide sequence and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell of choice may be used in the present invention.

[0100] Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Fusarium oxysporum* trypsin-like protease, and *Aspergillus niger* alpha-glucosidase.

[0101] Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Molecular Cellular Biology* 15: 5983-5990.

[0102] The control sequence may also be a signal peptide coding sequence that codes for an amino acid sequence linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleotide sequence may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding sequence that is foreign to the coding sequence. The foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, the foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell of choice, *i.e.*, secreted into a culture medium, may be used in the present invention.

[0103] Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Rhizomucor miehei* aspartic proteinase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, and *Humicola lanuginosa* lipase.

[0104] It may also be desirable to add regulatory sequences that allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the *lac*, *tac*, *xyl* and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the nucleotide sequence encoding the polypeptide would be operably linked with the regulatory sequence.

Expression Vectors

[0105] The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleic acids and control sequences described herein may be joined together to produce a recombinant expression vector that may include one or more (several) convenient restriction sites to allow for insertion or substitution of the nucleotide sequence encoding the polypeptide at such sites. Alternatively, a polynucleotide sequence of the present invention may be expressed by inserting the nucleotide sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0106] The recombinant expression vector may be any vector (*e.g.*, a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about expression of the nucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids.

[0107] The vector may be an autonomously replicating vector, *i.e.*, a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

[0108] The vectors of the present invention preferably contain one or more (several) selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

[0109] Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*.

[0110] The vectors of the present invention preferably contain an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

[0111] For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleotide sequences for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which have a high degree of identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleotide sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

[0112] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" is defined herein as a nucleotide sequence that enables a plasmid or vector to replicate *in vivo*.

[0113] Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems et al., 1991, Gene 98: 61-67; Cullen et al., 1987, Nucleic Acids Research 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

[0114] More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of the gene product. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional

copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

[0115] The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, *e.g.*, Sambrook *et al.*, 1989, *supra*).

Host Cells

[0116] The present invention also relates to recombinant host cells, comprising an isolated polynucleotide of the present invention, which are advantageously used in the recombinant production of the polypeptides. A vector comprising a polynucleotide of the present invention is introduced into a host cell so that the vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

[0117] The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, *e.g.*, a prokaryote or a eukaryote.

[0118] The prokaryotic host cell may be any gram-positive bacterium or gram-negative bacterium. Gram-positive bacteria include, but are not limited to, *Bacillus*, *Enterococcus*, *Clostridium*, *Geobacillus*, *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, *Streptomyces*. Gram-negative bacteria include, but are not limited to, *E. coli*, *Campylobacter*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, and *Ureaplasma*.

[0119] The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

[0120] Preferably, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by Hawksworth *et al.*, In, Ainsworth and Bisby's Dictionary of The Fungi, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth *et al.*, 1995, *supra*, page 171) and all mitosporic fungi (Hawksworth *et al.*, 1995, *supra*).

[0121] In a more preferred aspect, the fungal host cell is a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in Biology and Activities of Yeast (Skinner, F.A., Passmore, S.M., and Davenport, R.R., eds, Soc. App. Bacteriol. Symposium Series No. 9, 1980).

[0122] In an even more preferred aspect, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

[0123] In a most preferred aspect, the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* cell. In another most preferred aspect, the yeast host cell is a *Kluyveromyces lactis* cell. In another most preferred aspect, the yeast host cell is a *Yarrowia lipolytica* cell. In another more preferred aspect, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

[0124] In an even more preferred aspect, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

[0125] In a most preferred aspect, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus*

fumigatus, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger* or *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermispora*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

[0126] Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 0238023 and Yelton et al., 1984, Proceedings of the National Academy of Sciences USA 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Malardier et al., 1989, Gene 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, 194: 182-187, Academic Press, Inc., New York ; Ito et al., 1983, Journal of Bacteriology 153: 163; and Hinnen et al., 1978, Proceedings of the National Academy of Sciences USA 75: 1920.

Methods of Production

[0127] The present disclosure also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. Preferably, the cell is of the genus *Pycnoporus*. In a more preferred aspect, the cell is of the species *Pycnoporus sanguineus*.

[0128] The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell, as described herein, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[0129] The present disclosure also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell under conditions conducive for production of the polypeptide, wherein the host cell comprises a mutant nucleotide sequence having at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, of SEQ ID NO: 3, or of SEQ ID NO: 5, wherein the mutant nucleotide sequence encodes a polypeptide that comprises or consists of the mature polypeptide of SEQ ID NO: 2, of SEQ ID NO: 4 or of SEQ ID NO: 6 and (b) recovering the polypeptide.

[0130] In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods well known in the art. For example, the cell may be cultivated by shake flask cultivation, and small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted into the medium, it can be recovered from cell lysates.

[0131] The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide as described herein.

[0132] The resulting polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction,

spray-drying, evaporation, or precipitation.

[0133] The polypeptides of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., Protein Purification, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure polypeptides.

Plants

[0134] The present invention also relates to plants, e.g., a transgenic plant, plant part, or plant cell, comprising an isolated polynucleotide encoding a polypeptide having glucoamylase activity of the present invention so as to express and produce the polypeptide in recoverable quantities. The polypeptide may be recovered from the plant or plant part. Alternatively, the plant or plant part containing the recombinant polypeptide may be used as such for improving the quality of a food or feed, e.g., improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

[0135] The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as *Festuca*, *Lolium*, temperate grass, such as *Agrostis*, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn).

[0136] Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

[0137] Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, e.g., epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilisation of the invention are also considered plant parts, e.g., embryos, endosperms, aleurone and seeds coats.

[0138] Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

[0139] The transgenic plant or plant cell expressing a polypeptide of the present invention may be constructed in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more (several) expression constructs encoding a polypeptide of the present invention into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

[0140] The expression construct is conveniently a nucleic acid construct that comprises a polynucleotide encoding a polypeptide of the present invention operably linked with appropriate regulatory sequences required for expression of the nucleotide sequence in the plant or plant part of choice. Furthermore, the expression construct may comprise a selectable marker useful for identifying host cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

[0141] The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences is determined, for example, on the basis of when, where, and how the polypeptide is desired to be expressed. For instance, the expression of the gene encoding a polypeptide of the present invention may be constitutive or inducible, or may be developmental, stage or tissue specific, and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are, for example, described by Tague et al., 1988, Plant Physiology 86: 506.

[0142] For constitutive expression, the 35S-CaMV, the maize ubiquitin 1, and the rice actin 1 promoter may be used (Franck et al., 1980, Cell 21: 285-294, Christensen et al., 1992, Plant Mol. Biol. 18: 675-689; Zhang et al., 1991, Plant Cell 3: 1155-1165). organ-specific promoters may be, for example, a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards & Coruzzi, 1990, Ann. Rev. Genet. 24: 275-303), or from metabolic sink tissues such as meristems (Ito et al., 1994, Plant Mol. Biol. 24: 863-878), a seed specific promoter such as the glutelin, prolamin, globulin, or albumin promoter from rice (Wu et al., 1998, Plant and Cell Physiology 39: 885-889), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein

gene from *Vicia faba* (Conrad et al., 1998, Journal of Plant Physiology 152: 708-711), a promoter from a seed oil body protein (Chen et al., 1998, Plant and Cell Physiology 39: 935-941), the storage protein *napA* promoter from *Brassica napus*, or any other seed specific promoter known in the art, e.g., as described in WO 91/14772. Furthermore, the promoter may be a leaf specific promoter such as the *rbcs* promoter from rice or tomato (Kyoizuka et al., 1993, Plant Physiology 102: 991-1000, the chlorella virus adenine methyltransferase gene promoter (Mitra and Higgins, 1994, Plant Molecular Biology 26: 85-93), or the *aldP* gene promoter from rice (Kagaya et al., 1995, Molecular and General Genetics 248: 668-674), or a wound inducible promoter such as the potato *pin2* promoter (Xu et al., 1993, Plant Molecular Biology 22: 573-588). Likewise, the promoter may be inducible by abiotic treatments such as temperature, drought, or alterations in salinity or induced by exogenously applied substances that activate the promoter, e.g., ethanol, oestrogens, plant hormones such as ethylene, abscisic acid, and gibberellic acid, and heavy metals.

[0143] A promoter enhancer element may also be used to achieve higher expression of a polypeptide of the present invention in the plant. For instance, the promoter enhancer element may be an intron that is placed between the promoter and the nucleotide sequence encoding a polypeptide of the present invention. For instance, Xu *et al.*, 1993, *supra*, disclose the use of the first intron of the rice actin 1 gene to enhance expression.

[0144] The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

[0145] The nucleic acid construct is incorporated into the plant genome according to conventional techniques known in the art, including *Agrobacterium*-mediated transformation, virus-mediated transformation, microinjection, particle bombardment, biolistic transformation, and electroporation (Gasser et al., 1990, Science 244: 1293; Potrykus, 1990, Bio/Technology 8: 535; Shimamoto et al., 1989, Nature 338: 274).

[0146] Presently, *Agrobacterium tumefaciens*-mediated gene transfer is the method of choice for generating transgenic dicots (for a review, see Hooykas and Schilperoort, 1992, Plant Molecular Biology 19: 15-38) and can also be used for transforming monocots, although other transformation methods are often used for these plants. Presently, the method of choice for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the transforming DNA) of embryonic calli or developing embryos (Christou, 1992, Plant Journal 2: 275-281; Shimamoto, 1994, Current Opinion Biotechnology 5: 158-162; Vasil et al., 1992, Bio/Technology 10: 667-674). An alternative method for transformation of monocots is based on protoplast transformation as described by Omirulleh et al., 1993, Plant Molecular Biology 21: 415-428.

[0147] Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well-known in the art. Often the transformation procedure is designed for the selective elimination of selection genes either during regeneration or in the following generations by using, for example, co-transformation with two separate T-DNA constructs or site specific excision of the selection gene by a specific recombinase.

[0148] The present invention also relates to methods of producing a polypeptide of the present invention comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide having glucoamylase activity of the present invention under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

Compositions

[0149] The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the glucoamylase activity of the composition has been increased, e.g., with an enrichment factor of at least 1.1.

[0150] The composition may comprise a polypeptide of the present invention as the major enzymatic component, e.g., a mono-component composition. Alternatively, the composition may comprise multiple enzymatic activities, such as an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, haloperoxidase, invertase, laccase, lipase, mannosidase, oxidase, pectinolytic enzyme, peptidoglutaminase, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase. The additional enzyme(s) may be produced, for example, by a microorganism belonging to the genus *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*; *Fusarium*, preferably *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*,

Fusarium roseum, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sulphureum*, *Fusarium toruloseum*, *Fusarium trichothecioides*, or *Fusarium venenatum*; *Humicola*, preferably *Humicola insolens* or *Humicola lanuginosa*; or *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride*.

[0151] The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art.

[0152] Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

Combination of glucoamylase and acid alpha-amylase

[0153] According to this aspect of the invention a glucoamylase of the invention may be combined with an alpha-amylase, preferably acid alpha-amylase in a ratio of between 0.3 and 5.0 AFAU/AGU. More preferably the ratio between acid alpha-amylase activity and glucoamylase activity is at least 0.35, at least 0.40, at least 0.50, at least 0.60, at least 0.7, at least 0.8, at least 0.9, at least 1.0, at least 1.1, at least 1.2, at least 1.3, at least 1.4, at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.85, or even at least 1.9 AFAU/AGU. However, the ratio between acid alpha-amylase activity and glucoamylase activity should preferably be less than 4.5, less than 4.0, less than 3.5, less than 3.0, less than 2.5, or even less than 2.25 AFAU/AGU. In AUU/AGI the activities of acid alpha-amylase and glucoamylase are preferably present in a ratio of between 0.4 and 6.5 AUU/AGI. More preferably the ratio between acid alpha-amylase activity and glucoamylase activity is at least 0.45, at least 0.50, at least 0.60, at least 0.7, at least 0.8, at least 0.9, at least 1.0, at least 1.1, at least 1.2, at least 1.3, at least 1.4, at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.9, at least 2.0, at least 2.1, at least 2.2, at least 2.3, at least 2.4, or even at least 2.5 AUU/AGI. However, the ratio between acid alpha-amylase activity and glucoamylase activity is preferably less than 6.0, less than 5.5, less than 4.5, less than 4.0, less than 3.5, or even less than 3.0 AUU/AGI.

[0154] The composition is suitable for use in a starch conversion process mentioned below for producing syrup and fermentation products, such as ethanol.

[0155] Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

Uses

[0156] The present invention is also directed to processes/methods for using the polypeptides having glucoamylase activity of the invention.

[0157] Uses according to the invention include starch conversion of starch to, e.g., syrup and fermentation products, including ethanol and beverages. Examples of processes where a glucoamylase of the invention may be used include the ones described in WO 92/20777, WO 03/066816, WO 03/066826, WO 2004/080923, and WO 2004/081193, which are hereby all incorporated by reference.

Production of fermentation products

Process for producing fermentation products from gelatinized starch containing material

[0158] In this aspect the present invention relates to a process for producing a fermentation product, especially ethanol, from starch-containing material, which process includes a liquefaction step and sequentially or simultaneously performed saccharification and fermentation steps.

[0159] The invention relates to a process for producing a fermentation product from starch-containing material comprising the steps of:

1. (a) liquefying starch-containing material; preferably using an alpha amylase
2. (b) saccharifying the liquefied material obtained in step (a) using a glucoamylase of the invention; and
3. (c) fermenting the saccharified material using a fermenting organism.

[0160] The fermentation product, such as especially ethanol, may optionally be recovered after fermentation, *e.g.*, by distillation. Suitable starch-containing starting materials are listed in the section "Starch-containing materials"-section below. Contemplated enzymes are listed in the "Enzymes"-section below. The liquefaction is preferably carried out in the presence of an alpha-amylase. The fermentation is preferably carried out in the presence of yeast, preferably a strain of *Saccharomyces*. Suitable fermenting organisms are listed in the "Fermenting Organisms"-section below. In preferred embodiments steps (b) and (c) are carried out sequentially or simultaneously (*i.e.*, as SSF process).

[0161] In a particular embodiment, the process of the invention further comprises, prior to step (a), the steps of:

- x) reducing the particle size of the starch-containing material, preferably by milling;
- y) forming a slurry comprising the starch-containing material and water.

[0162] The aqueous slurry may contain from 10-40 wt %, preferably 25-35 wt % starch-containing material. The slurry is heated to above the gelatinization temperature and alpha-amylase, preferably bacterial and/or acid fungal alpha-amylase, may be added to initiate liquefaction (thinning). The slurry may in an embodiment be jet-cooked to further gelatinize the slurry before being subjected to an alpha-amylase in step (a) of the invention.

[0163] More specifically liquefaction may be carried out as a three-step hot slurry process. The slurry is heated to between 60-95°C, preferably 80-85°C, and alpha-amylase is added to initiate liquefaction (thinning). Then the slurry may be jet-cooked at a temperature between 95-140°C, preferably 105-125°C, for 1-15 minutes, preferably for 3-10 minute, especially around 5 minutes. The slurry is cooled to 60-95°C and more alpha-amylase is added to finalize hydrolysis (secondary liquefaction). The liquefaction process is usually carried out at pH 4.5-6.5, in particular at a pH between 5 and 6. Milled and liquefied whole grains are known as mash.

[0164] The saccharification in step (b) may be carried out using conditions well known in the art. For instance, a full saccharification process may last up to from about 24 to about 72 hours, however, it is common only to do a pre-saccharification of typically 40-90 minutes at a temperature between 30-65°C, typically about 60°C, followed by complete saccharification during fermentation in a simultaneous saccharification and fermentation process (SSF process). Saccharification is typically carried out at temperatures from 30-65°C, typically around 60°C, and at a pH between 4 and 5, normally at about pH 4.5.

[0165] The most widely used process in fermentation product, especially ethanol, production is the simultaneous saccharification and fermentation (SSF) process, in which there is no holding stage for the saccharification, meaning that fermenting organism, such as yeast, and enzyme(s) may be added together. SSF may typically be carried out at a temperature between 25°C and 40°C, such as between 29°C and 35°C, such as between 30°C and 34°C, such as around 32°C. According to the invention the temperature may be adjusted up or down during fermentation.

[0166] In accordance with the present invention the fermentation step (c) includes, without limitation, fermentation processes used to produce alcohols (*e.g.*, ethanol, methanol, butanol); organic acids (*e.g.*, citric acid, acetic acid, itaconic acid, lactic acid, gluconic acid); ketones (*e.g.*, acetone); amino acids (*e.g.*, glutamic acid); gases (*e.g.*, H₂ and CO₂); antibiotics (*e.g.*, penicillin and tetracycline); enzymes; vitamins (*e.g.*, riboflavin, B12, beta-carotene); and hormones. Preferred fermentation processes include alcohol fermentation processes, as are well known in the art. Preferred fermentation processes are anaerobic fermentation processes, as are well known in the art.

Processes for producing fermentation products from un-gelatinized starch-containing

[0167] In this aspect the invention relates to processes for producing a fermentation product from starch-containing material without gelatinization of the starch-containing material (*i.e.*, uncooked starch-containing material). In one embodiment only a glucoamylase of the invention is used during saccharification and fermentation. According to the invention the desired fermentation product, such as ethanol, can be produced without liquefying the aqueous slurry containing the starch-containing material. In one embodiment a process of the invention includes saccharifying (milled) starch-containing material, *e.g.*, granular starch, below the gelatinization temperature in the presence of a glucoamylase of the invention to produce sugars that can be fermented into the desired fermentation product by a suitable fermenting organism.

[0168] Accordingly, in this aspect the invention relates to a process for producing a fermentation product from starch-containing material comprising:

1. (a) saccharifying starch-containing material with a mature glucoamylase comprised in SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, preferably the sequence shown as amino acids 19 to 573 in SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6 or a glucoamylase having at least 90%, more preferably at least 91%, more preferably at least 92%, even more preferably at least 93%, most preferably at least 94%, and even most preferably at least 95%, such as at least 96%, at least 97%, at least 98%, at least 99% or even 100% identity thereto, at a temperature below the initial gelatinization temperature of said starch-containing material,
2. (b) fermenting using a fermenting organism.

[0169] Steps (a) and (b) of the process of the invention may be carried out sequentially or simultaneously. In an embodiment a slurry comprising water and starch-containing material is prepared before step (a).

[0170] The fermentation process may be carried out for a period of 1 to 250 hours, preferably is from 25 to 190 hours, more preferably from 30 to 180 hours, more preferably from 40 to 170 hours, even more preferably from 50 to 160 hours, yet more preferably from 60 to 150 hours, even yet more preferably from 70 to 140 hours, and most preferably from 80 to 130 hours.

[0171] The term "initial gelatinization temperature" means the lowest temperature at which gelatinization of the starch commences. Starch heated in water begins to gelatinize between 50°C and 75°C; the exact temperature of gelatinization depends on the specific starch, and can readily be determined by the skilled artisan. Thus, the initial gelatinization temperature may vary according to the plant species, to the particular variety of the plant species as well as with the growth conditions. In the context of this invention the initial gelatinization temperature of a given starch-containing material is the temperature at which birefringence is lost in 5% of the starch granules using the method described by Gorinstein and Lii, 1992, *Starch/Stärke* 44(12): 461-466.

[0172] Before step (a) a slurry of starch-containing material, such as granular starch, having 10-55 wt. % dry solids, preferably 25-40 wt. % dry solids, more preferably 30-35 wt. % dry solids of starch-containing material may be prepared. The slurry may include water and/or process waters, such as stillage (backset), scrubber water, evaporator condensate or distillate, side stripper water from distillation, or other fermentation product plant process water. Because the process of the invention is carried out below the gelatinization temperature and thus no significant viscosity increase takes place, high levels of stillage may be used if desired. In an embodiment the aqueous slurry contains from about 1 to about 70 vol. % stillage, preferably 15-60 vol. % stillage, especially from about 30 to 50 vol. % stillage.

[0173] The starch-containing material may be prepared by reducing the particle size, preferably by dry or wet milling, to 0.05 to 3.0 mm, preferably 0.1-0.5 mm. After being subjected to a process of the invention at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or preferably at least 99% of the dry solids of the starch-containing material is converted into a soluble starch hydrolysate.

[0174] The process of the invention is conducted at a temperature below the initial gelatinization temperature. Preferably the temperature at which step (a) is carried out is between 30-75°C, preferably between 45-60°C.

[0175] In a preferred embodiment step (a) and step (b) are carried out as a sequential or simultaneous saccharification and fermentation process. In such preferred embodiment the process is typically carried at a temperature between 25°C and 40°C, such as between 29°C and 35°C, such as between 30°C and 34°C, such as around 32°C. According to the invention the temperature may be adjusted up or down during fermentation.

[0176] In an embodiment simultaneous saccharification and fermentation is carried out so that the sugar level, such as glucose

level, is kept at a low level such as below 6 wt. %, preferably below about 3 wt. %, preferably below about 2 wt. %, more preferred below about 1 wt. %, even more preferred below about 0.5 wt. %, or even more preferred 0.25 wt. %, such as below about 0.1 wt. %. Such low levels of sugar can be accomplished by simply employing adjusted quantities of enzyme and fermenting organism. A skilled person in the art can easily determine which quantities of enzyme and fermenting organism to use. The employed quantities of enzyme and fermenting organism may also be selected to maintain low concentrations of maltose in the fermentation broth. For instance, the maltose level may be kept below about 0.5 wt. % or below about 0.2 wt. %.

[0177] The process of the invention may be carried out at a pH in the range between 3 and 7, preferably from pH 3.5 to 6, or more preferably from pH 4 to 5.

Starch-containing materials

[0178] Any suitable starch-containing starting material, including granular starch, may be used according to the present invention. The starting material is generally selected based on the desired fermentation product. Examples of starch-containing starting materials, suitable for use in a process of present invention, include tubers, roots, stems, whole grains, corns, cobs, wheat, barley, rye, milo, sago, cassava, tapioca, sorghum, rice, peas, beans, or sweet potatoes, or mixtures thereof, or cereals, sugar-containing raw materials, such as molasses, fruit materials, sugar cane or sugar beet, potatoes, and cellulose-containing materials, such as wood or plant residues, or mixtures thereof. Contemplated are both waxy and non-waxy types of corn and barley.

[0179] The term "granular starch" means raw uncooked starch, *i.e.*, starch in its natural form found in cereal, tubers or grains. Starch is formed within plant cells as tiny granules insoluble in water. When put in cold water, the starch granules may absorb a small amount of the liquid and swell. At temperatures up to 50°C to 75°C the swelling may be reversible. However, with higher temperatures an irreversible swelling called "gelatinization" begins. Granular starch to be processed may be a highly refined starch quality, preferably at least 90%, at least 95%, at least 97% or at least 99.5% pure or it may be a more crude starch containing material comprising milled whole grain including non-starch fractions such as germ residues and fibers. The raw material, such as whole grain, is milled in order to open up the structure and allowing for further processing. Two milling processes are preferred according to the invention: wet and dry milling. In dry milling whole kernels are milled and used. Wet milling gives a good separation of germ and meal (starch granules and protein) and is often applied at locations where the starch hydrolysate is used in production of syrups. Both dry and wet milling is well known in the art of starch processing and is equally contemplated for the process of the invention.

[0180] The starch-containing material is reduced in particle size, preferably by dry or wet milling, in order to expose more surface area. In an embodiment the particle size is between 0.05 to 3.0 mm, preferably 0.1-0.5 mm, or so that at least 30%, preferably at least 50%, more preferably at least 70%, even more preferably at least 90% of the starch-containing material fit through a sieve with a 0.05 to 3.0 mm screen, preferably 0.1-0.5 mm screen.

Fermentation Products

[0181] The term "fermentation product" means a product produced by a process including a fermentation step using a fermenting organism. Fermentation products contemplated according to the invention include alcohols (*e.g.*, ethanol, methanol, butanol); organic acids (*e.g.*, citric acid, acetic acid, itaconic acid, lactic acid, gluconic acid); ketones (*e.g.*, acetone); amino acids (*e.g.*, glutamic acid); gases (*e.g.*, H₂ and CO₂); antibiotics (*e.g.*, penicillin and tetracycline); enzymes; vitamins (*e.g.*, riboflavin, B₁₂, beta-carotene); and hormones. In a preferred embodiment the fermentation product is ethanol, *e.g.*, fuel ethanol; drinking ethanol, *i.e.*, potable neutral spirits; or industrial ethanol or products used in the consumable alcohol industry (*e.g.*, beer and wine), dairy industry (*e.g.*, fermented dairy products), leather industry and tobacco industry. Preferred beer types comprise ales, stouts, porters, lagers, bitters, malt liquors, happoushu, high-alcohol beer, low-alcohol beer, low-calorie beer or light beer. Preferred fermentation processes used include alcohol fermentation processes, as are well known in the art. Preferred fermentation processes are anaerobic fermentation processes, as are well known in the art.

Fermenting Organisms

[0182] "Fermenting organism" refers to any organism, including bacterial and fungal organisms, suitable for use in a

fermentation process and capable of producing desired a fermentation product. Especially suitable fermenting organisms are able to ferment, *i.e.*, convert, sugars, such as glucose or maltose, directly or indirectly into the desired fermentation product. Examples of fermenting organisms include fungal organisms, such as yeast. Preferred yeast includes strains of *Saccharomyces* spp., in particular, *Saccharomyces cerevisiae*. Commercially available yeast include, *e.g.*, Red Star™/Lesaffre Ethanol Red (available from Red Star/Lesaffre, USA) FALI (available from Fleischmann's Yeast, a division of Burns Philp Food Inc., USA), SUPERSTART (available from Alltech), GERT STRAND (available from Gert Strand AB, Sweden) and FERMIOL (available from DSM Specialties).

ENZYMES

Glucoamylase

[0183] The glucoamylase is preferably a glucoamylase of the invention. However, as mentioned above a glucoamylase of the invention may also be combined with other glucoamylases. The term "glucoamylase" (1,4-alpha-D-glucan glucohydrolase, EC 3.2.1.3) is an enzyme, which catalyzes the release of D-glucose from the non-reducing ends of starch or related oligo- and polysaccharide molecules.

[0184] The glucoamylase may be added in an amount of 0.001 to 10 AGU/g DS, preferably from 0.01 to 5 AGU/g DS, such as around 0.1, 0.3, 0.5, 1 or 2 AGU/g DS, especially 0.1 to 0.5 AGU/g DS or 0.02-20 AGU/g DS, preferably 0.1-10 AGU/g DS.

Alpha-Amylase

[0185] The alpha-amylase may according to the invention be of any origin. Preferred are alpha-amylases of fungal or bacterial origin.

[0186] In a preferred embodiment the alpha-amylase is an acid alpha-amylase, *e.g.*, fungal acid alpha-amylase or bacterial acid alpha-amylase. The term "acid alpha-amylase" means an alpha-amylase (EC 3.2.1.1) which added in an effective amount has activity optimum at a pH in the range of 3 to 7, preferably from 3.5 to 6, or more preferably from 4-5.

Bacterial Alpha-Amylases

[0187] According to the invention a bacterial alpha-amylase may preferably be derived from the genus *Bacillus*.

[0188] In a preferred embodiment the *Bacillus* alpha-amylase is derived from a strain of *B. licheniformis*, *B. amyloliquefaciens*, *B. subtilis* or *B. stearothermophilus*, but may also be derived from other *Bacillus* sp. Specific examples of contemplated alpha-amylases include the *Bacillus licheniformis* alpha-amylase (BLA) shown in SEQ ID NO: 4 in WO 99/19467, the *Bacillus amyloliquefaciens* alpha-amylase (BAN) shown in SEQ ID NO: 5 in WO 99/19467, and the *Bacillus stearothermophilus* alpha-amylase (BSG) shown in SEQ ID NO: 3 in WO 99/19467. In an embodiment of the invention the alpha-amylase is an enzyme having a degree of identity of at least 60%, preferably at least 70%, more preferred at least 80%, even more preferred at least 90%, such as at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to any of the sequences shown as SEQ ID NOS: 1, 2, 3, 4, or 5, respectively, in WO 99/19467.

[0189] The *Bacillus* alpha-amylase may also be a variant and/or hybrid, especially one described in any of WO 96/23873, WO 96/23874, WO 97/41213, WO 99/19467, WO 00/60059, and WO 02/10355 (all documents hereby incorporated by reference). Specifically contemplated alpha-amylase variants are disclosed in US patent nos. 6,093,562, 6,297,038 or US patent no. 6,187,576 (hereby incorporated by reference) and include *Bacillus stearothermophilus* alpha-amylase (BSG alpha-amylase) variants having a deletion of one or two amino acid in position 179 to 182, preferably a double deletion disclosed in WO 96/23873 - see, *e.g.*, page 20, lines 1-10 (hereby incorporated by reference), preferably corresponding to delta(181-182) compared to the wild-type BSG alpha-amylase amino acid sequence set forth in SEQ ID NO: 3 disclosed in WO 99/19467 or deletion of amino acids 179 and 180 using SEQ ID NO: 3 in WO 99/19467 for numbering (which reference is hereby incorporated by reference). Even more preferred are *Bacillus* alpha-amylases, especially *Bacillus stearothermophilus* alpha-amylase, which have a double deletion corresponding to delta(181-182) and further comprise a N193F substitution (also denoted I181* + G182* + N193F) compared to the wild-type BSG alpha-amylase amino acid sequence set forth in SEQ ID NO: 3 disclosed in WO 99/19467.

[0190] The alpha-amylase may also be a maltogenic alpha-amylase. A "maltogenic alpha-amylase" (glucan 1,4-alpha-maltohydrolase, EC 3.2.1.133) is able to hydrolyze amylose and amylopectin to maltose in the alpha-configuration. A maltogenic alpha-amylase from *Bacillus stearotheophilus* strain NCIB 11837 is commercially available from Novozymes A/S, Denmark. The maltogenic alpha-amylase is described in U.S. Patent Nos. 4,598,048, 4,604,355 and 6,162,628, which are hereby incorporated by reference.

Bacterial Hybrid Alpha-Amylases

[0191] A hybrid alpha-amylase specifically contemplated comprises 445 C-terminal amino acid residues of the *Bacillus licheniformis* alpha-amylase (shown as SEQ ID NO: 3 in WO 99/19467) and the 37 N-terminal amino acid residues of the alpha-amylase derived from *Bacillus amyloliquefaciens* (shown as SEQ ID NO: 5 in WO 99/19467), with one or more, especially all, of the following substitution:

G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S (using the *Bacillus licheniformis* numbering). Also preferred are variants having one or more of the following mutations (or corresponding mutations in other *Bacillus* alpha-amylase backbones): H154Y, A181T, N190F, A209V and Q264S and/or deletion of two residues between positions 176 and 179, preferably deletion of E178 and G179 (using the SEQ ID NO: 5 numbering of WO 99/19467).

[0192] The bacterial alpha-amylase may be added in amounts as are well-known in the art. When measured in KNU units (described below in the "Materials & Methods"-section) the alpha-amylase activity is preferably present in an amount of 0.5-5,000 NU/g of DS, in an amount of 1-500 NU/g of DS, or more preferably in an amount of 5-1,000 NU/g of DS, such as 10-100 NU/g DS.

Fungal Alpha-Amylases

[0193] Fungal acid alpha-amylases include acid alpha-amylases derived from a strain of the genus *Aspergillus*, such as *Aspergillus oryzae*, *Aspergillus niger*, or *Aspergillus kawachii* alpha-amylases.

[0194] A preferred acid fungal alpha-amylase is a Fungamyl-like alpha-amylase which is preferably derived from a strain of *Aspergillus oryzae*. In the present disclosure, the term "Fungamyl-like alpha-amylase" indicates an alpha-amylase which exhibits a high identity, i.e., more than 70%, more than 75%, more than 80%, more than 85% more than 90%, more than 95%, more than 96%, more than 97%, more than 98%, more than 99% or even 100% identity to the mature part of the amino acid sequence shown in SEQ ID NO: 10 in WO 96/23874.

[0195] Another preferred acid alpha-amylase is derived from a strain *Aspergillus niger*. In a preferred embodiment the acid fungal alpha-amylase is the one from *A. niger* disclosed as "AMYA_ASPNG" in the Swiss-prot/TrEMBL database under the primary accession no. P56271 and described in more detail in WO 89/01969 (Example 3). The acid *Aspergillus niger* acid alpha-amylase is also shown as SEQ ID NO: 1 in WO 2004/080923 (Novozymes) which is hereby incorporated by reference. Also variants of said acid fungal amylase having at least 70% identity, such as at least 80% or even at least 90% identity, such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to SEQ ID NO: 1 in WO 2004/080923 are contemplated. A suitable commercially available acid fungal alpha-amylase derived from *Aspergillus niger* is SP288 (available from Novozymes A/S, Denmark).

[0196] In a preferred embodiment the alpha-amylase is derived from *Aspergillus kawachii* and disclosed by Kaneko et al., 1996, J. Ferment. Bioeng. 81: 292-298, "Molecular-cloning and determination of the nucleotide-sequence of a gene encoding an acid-stable alpha-amylase from *Aspergillus kawachii*"; and further as EMBL:#AB008370.

[0197] The fungal acid alpha-amylase may also be a wild-type enzyme comprising a carbohydrate-binding module (CBM) and an alpha-amylase catalytic domain (i.e., a none-hybrid), or a variant thereof. In an embodiment the wild-type acid alpha-amylase is derived from a strain of *Aspergillus kawachii*.

Fungal Hybrid Alpha-Amylases

[0198] In a preferred embodiment the fungal acid alpha-amylase is a hybrid alpha-amylase. Preferred examples of fungal hybrid alpha-amylases include the ones disclosed in WO 2005/003311 or U.S. Patent Publication no. 2005/0054071 (Novozymes) or US patent application no. 60/638,614 (Novozymes) which is hereby incorporated by reference. A hybrid alpha-amylase may comprise an alpha-amylase catalytic domain (CD) and a carbohydrate-binding domain/module (CBM) and optional a linker.

[0199] Specific examples of contemplated hybrid alpha-amylases include those disclosed in U.S. application no. 60/638,614 including Fungamyl variant with catalytic domain JA118 and *Athelia rolfsii* SBD (SEQ ID NO: 100 in U.S. application no. 60/638,614), *Rhizomucor pusillus* alpha-amylase with *Athelia rolfsii* AMG linker and SBD (SEQ ID NO: 101 in U.S. application no. 60/638,614) and *Meripilus giganteus* alpha-amylase with *Athelia rolfsii* glucoamylase linker and SBD (SEQ ID NO: 102 in U.S. application no. 60/638,614).

[0200] Other specific examples of contemplated hybrid alpha-amylases include those disclosed in U.S. Application Publication no. 2005/0054071, including those disclosed in Table 3 on page 15, such as *Aspergillus niger* alpha-amylase with *Aspergillus kawachii* linker and starch binding domain.

Commercial Alpha-Amylase Products

[0201] Preferred commercial compositions comprising alpha-amylase include MYCOLASE from DSM (Gist Brocades), BAN™, TERMAMYL™ SC, FUNGAMYL™, LIQUOZYME™ X and SAN™ SUPER, SAN™ EXTRA L (Novozymes A/S) and CLARASE™ L-40,000, DEX-LOT™, SPEZYME™ FRED, SPEZYME™ AA, SPEZYME™ Ethyl, GC358, GC980, SPEZYME™ RSL, and SPEZYME™ DELTA AA (Genencor Int.), and the acid fungal alpha-amylase sold under the trade name SP288 (available from Novozymes A/S, Denmark).

[0202] An acid alpha-amylases may according to the invention be added in an amount of 0.1 to 10 AFAU/g DS, preferably 0.10 to 5 AFAU/g DS, especially 0.3 to 2 AFAU/g DS.

Production of syrup

[0203] The present invention also provides a process of using a glucoamylase of the invention for producing syrup, such as glucose and the like, from a starch-containing material. Suitable starting materials are exemplified in the "Starch-containing materials"-section above. Generally, the process comprises the steps of partially hydrolyzing starch-containing material (liquefaction) in the presence of alpha-amylase and then further saccharifying the release of glucose from the non-reducing ends of the starch or related oligo- and polysaccharide molecules in the presence of glucoamylase of the invention.

[0204] Liquefaction and saccharification may be carried out as described above for fermentation product production.

[0205] The glucoamylase of the invention may also be used in immobilized form. This is suitable and often used for producing speciality syrups, such as maltose syrups, and further for the raffinate stream of oligosaccharides in connection with the production of fructose syrups, e.g., high fructose syrup (HFS).

[0206] Consequently, this aspect of the invention relates to a process of producing syrup from starch-containing material, comprising

1. (a) liquefying starch-containing material in the presence of an alpha-amylase, and
2. (b) saccharifying the material obtained in step (a) using a glucoamylase of the invention.

[0207] A syrup may be recovered from the saccharified material obtained in step (b).

[0208] Details on suitable conditions can be found above.

Brewing

[0209] A glucoamylase of the invention can also be used in a brewing process. The glucoamylases of the invention is added in effective amounts which can be easily determined by the skilled person in the art.

[0210] The present invention is further described by the following examples.

Materials & Methods

[0211] Yeast: RED STAR™ available from Red Star/Lesaffre, USA

Media and reagents:

[0212] Chemicals used as buffers and substrates were commercial products of at least reagent grade.

[0213] PDA: 39 g/L Potato Dextrose Agar, 20 g/L agar, 50 ml/L glycerol

Methods

[0214] Unless otherwise stated, DNA manipulations and transformations were performed using standard methods of molecular biology as described in Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology", John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990.

Glucoamylase activity

[0215] Glucoamylase activity may be measured in AGI units or in Glucoamylase Units (AGU).

Glucoamylase activity (AGI)

[0216] Glucoamylase (equivalent to amyloglucosidase) converts starch into glucose. The amount of glucose is determined here by the glucose oxidase method for the activity determination. The method described in the section 76-11 Starch-Glucoamylase Method with Subsequent Measurement of Glucose with Glucose Oxidase in "Approved methods of the American Association of Cereal Chemists". Vol. 1-2 AACC, from American Association of Cereal Chemists, (2000); ISBN: 1-891127-12-8.

[0217] One glucoamylase unit (AGI) is the quantity of enzyme which will form 1 micro mole of glucose per minute under the standard conditions of the method.

Standard conditions/reaction conditions:

[0218]

Substrate:	Soluble starch, concentration approx. 16 g dry matter/L.
Buffer:	Acetate, approx. 0.04 M, pH=4.3
pH:	4.3
Incubation temperature:	60°C
Reaction time:	15 minutes
Termination of the reaction:	NaOH to a concentration of approximately 0.2 g/L (pH=9)
Enzyme concentration:	0.15-0.55 AAU/mL.

[0219] The starch should be Lintner starch, which is a thin-boiling starch used in the laboratory as colorimetric indicator. Lintner starch is obtained by dilute hydrochloric acid treatment of native starch so that it retains the ability to color blue with iodine.

Glucoamylase activity (AGU)

[0220] The Novo Glucoamylase Unit (AGU) is defined as the amount of enzyme, which hydrolyzes 1 micromole maltose per minute under the standard conditions 37°C, pH 4.3, substrate: maltose 23.2 mM, buffer: acetate 0.1 M, reaction time 5 minutes.

[0221] An autoanalyzer system may be used. Mutarotase is added to the glucose dehydrogenase reagent so that any alpha-D-glucose present is turned into beta-D-glucose. Glucose dehydrogenase reacts specifically with beta-D-glucose in the reaction mentioned above, forming NADH which is determined using a photometer at 340 nm as a measure of the original glucose concentration.

AMG incubation:	
Substrate:	maltose 23.2 mM
Buffer:	acetate 0.1 M
pH:	4.30 ± 0.05
Incubation temperature:	37°C ± 1
Reaction time:	5 minutes
Enzyme working range:	0.5-4.0 AGU/mL
Color reaction:	
GlucDH:	430 U/L
Mutarotase:	9 U/L
NAD:	0.21 mM
Buffer:	phosphate 0.12 M; 0.15 M NaCl
pH:	7.60 ± 0.05
Incubation temperature:	37°C ± 1
Reaction time:	5 minutes
Wavelength:	340 nm

[0222] A folder (EB-SM-0131.02/01) describing this analytical method in more detail is available on request from Novozymes A/S, Denmark, which folder is hereby included by reference.

Alpha-amylase activity (KNU)

[0223] Alpha-amylase activity may be determined using potato starch as substrate. This method is based on the break-down of modified potato starch by the enzyme, and the reaction is followed by mixing samples of the starch/enzyme solution with an iodine solution. Initially, a blackish-blue color is formed, but during the break-down of the starch the blue color gets weaker and gradually turns into a reddish-brown, which is compared to a colored glass standard.

[0224] One Kilo Novo alpha amylase Unit (KNU) is defined as the amount of enzyme which, under standard conditions (*i.e.*, at 37°C ± 0.05; 0.0003 M Ca²⁺; and pH 5.6) dextrinizes 5260 mg starch dry substance Merck Amylum soluble.

[0225] A folder EB-SM-0009.02/01 describing this analytical method in more detail is available upon request to Novozymes A/S, Denmark, which folder is hereby included by reference.

Acid alpha-amylase activity

[0226] When used according to the present invention the activity of any acid alpha-amylase may be measured in AFAU (Acid Fungal Alpha-amylase Units). Alternatively activity of acid alpha-amylase may be measured in AAU (Acid Alpha-amylase Units).

Acid Alpha-amylase Units (AAU)

[0227] Acid alpha-amylase activity can be measured in AAU (Acid Alpha-amylase Units), which is an absolute method. One Acid Amylase Unit (AAU) is the quantity of enzyme converting 1 g of starch (100% of dry matter) per hour under standardized conditions into a product having a transmission at 620 nm after reaction with an iodine solution of known strength equal to the one of a color reference.

Standard conditions/reaction conditions:

[0228]

Substrate:	Soluble starch. Concentration approx. 20 g DS/L.
Buffer:	Citrate, approx. 0.13 M, pH=4.2
Iodine solution:	40.176 g potassium iodide + 0.088 g iodine/L
City water:	15°-20°dH (German degree hardness)
pH:	4.2
Incubation temperature:	30°C
Reaction time:	11 minutes
Wavelength:	620 nm
Enzyme concentration:	0.13-0.19 AAU/mL
Enzyme working range:	0.13-0.19 AAU/mL

[0229] The starch should be Lintner starch, which is a thin-boiling starch used in the laboratory as colorimetric indicator. Lintner starch is obtained by dilute hydrochloric acid treatment of native starch so that it retains the ability to color blue with iodine. Further details can be found in EP 0140410 B2, which disclosure is hereby included by reference.

Acid alpha-amylase activity (AFAU)

[0230] Acid alpha-amylase activity may be measured in AFAU (Acid Fungal Alpha-amylase Units), which are determined relative to an enzyme standard. 1 AFAU is defined as the amount of enzyme which degrades 5.260 mg starch dry matter per hour under the below mentioned standard conditions.

[0231] Acid alpha-amylase, an endo-alpha-amylase (1,4-alpha-D-glucan-glucanohydrolase, EC 3.2.1.1) hydrolyzes alpha-1,4-glucosidic bonds in the inner regions of the starch molecule to form dextrins and oligosaccharides with different chain lengths. The intensity of color formed with iodine is directly proportional to the concentration of starch. Amylase activity is determined using reverse colorimetry as a reduction in the concentration of starch under the specified analytical conditions.

ALPHA - AMYLASE



$\lambda = 590 \text{ nm}$

blue/violet t = 23 sec. decoloration

Standard conditions/reaction conditions:

[0232]

Substrate:	Soluble starch, approx. 0.17 g/L
Buffer:	Citrate, approx. 0.03 M
Iodine (I ₂):	0.03 g/L
CaCl ₂ :	1.85 mM
pH:	2.50 ± 0.05
Incubation temperature:	40°C
Reaction time:	23 seconds
Wavelength:	590 nm
Enzyme concentration:	0.025 AFAU/mL
Enzyme working range:	0.01-0.04 AFAU/mL

[0233] A folder EB-SM-0259.02/01 describing this analytical method in more detail is available upon request to Novozymes A/S, Denmark, which folder is hereby included by reference.

Example 1:

Simultaneous Saccharification and Fermentation (SSF) with *Pycnoporus sanguineus* AMG

[0234] The SSF performance of *Pycnoporus sanguineus* glucoamylases was tested at different enzyme doses. Fermentation was run under the following conditions:

Substrate: Ground corn was slurried with backset and adjusted its dry substance to approximately 32% (w/w). It was then liquefied at 85°C and pH 5.8. The liquefied mash had a DE of 13.4.

Temperature: 32°C

Initial pH: 5.0

Enzyme dose: *Pycnoporus sanguineus* AMG produced in *A. niger* at 30, 40, 55 and 70 micrograms enzyme protein/g DS. Enzymes were compared to a purified sample of the commercial *Talaromyces emersonii* AMG dosed at same dosages. The highest dose of *Talaromyces emersonii* AMG is equivalent to an industry relevant amount of 0.56 AGU/g DS. A control for maximum obtainable saccharification was prepared using excess amounts of commercial AMG and alpha-amylase.

Fermentation

[0235] To the substrate for SSF, 1000 ppm urea as nitrogen source and 3 ppm penicillin for bacterial control were added; the pH was adjusted to 5.0 with H₂SO₄. Aliquots of 5 g mash were transferred to 15 ml centrifuge tubes with a hole drilled at the top for CO₂ release. Enzymes and yeast were added and the tubes were placed in a water bath without stirring at 32°C for 54 hrs. Samples were analyzed in HPLC for determination of ethanol produced during fermentation. The results are shown in the tables below.

Table 1. Ethanol g/L produced during SSF with <i>Pycnoporus sanguineus</i> AMG at 30, 40, 55 and 70 micrograms enzyme protein/g DS as compared to <i>Talaromyces emersonii</i> AMG. Control resulted in 133.08 g/L ethanol				
	Enzyme dose (micrograms enzyme protein/g DS)			
	30	40	55	70
<i>Talaromyces emersonii</i> , SEQ ID NO: 10	110.3	119.8	124.9	126.9

Table 1. Ethanol g/L produced during SSF with *Pycnoporus sanguineus* AMG at 30, 40, 55 and 70 micrograms enzyme protein/g DS as compared to *Talaromyces emersonii* AMG. Control resulted in 133.08 g/L ethanol

	Enzyme dose (micrograms enzyme protein/g DS)			
	30	40	55	70
<i>Pycnoporus sanguineus</i> , SEQ ID NO: 2	112.8	119.5	126.0	130.7

Table 2. Ethanol g/L produced during SSF with *Pycnoporus sanguineus* AMG at 30, 40, 55 and 70 micrograms enzyme protein/g DS as compared to *Talaromyces emersonii* AMG. Control resulted in 131.9 g/L ethanol

	Enzyme dose (micrograms enzyme protein/g DS)			
	30	40	55	70
<i>Talaromyces emersonii</i> , SEQ ID NO: 10	110.4	118.9	123.4	125.6
<i>Pycnoporus sanguineus</i> , SEQ ID NO: 4	104.4	116.3	127.5	129.7
<i>Pycnoporus sanguineus</i> , SEQ ID NO: 6	109.4	115.7	126.8	129.5

Example 2:

Raw starch hydrolysis with *Pycnoporus* sp. AMG.

Materials

[0236] 3% raw starch suspension: Prepared to be 100 mM Na-acetate, 1 mM CaCl₂, 0.025% NaN₃, and 3% corn starch. Each component is prepared for a 160 ml preparation, and the volume adjusted to only 152 ml with milliQ water after pH adjustment, since then each component will have the right concentration when mixed with the enzyme as described below.

Glucose CII test kit (Wako)

[0237]

Purified *Pycnoporus* or *Talaromyces* AMG: The purified samples should not contain alpha-amylase.

Purified *T. cingulata* AMG (control): The purified samples should not contain alpha-amylase. The AGU activity should be known.

Purified JA126AN Alpha-Amylase A: Hybrid alpha-amylase consisting of *Rhizomucor pusillus* alpha-amylase with *Aspergillus niger* glucoamylase linker and SBD disclosed as V039 in Table 5 in WO 2006/069290 (Novozymes A/S).

Methods

[0238]

1) Dilute purified AMG and JA126AN with milliQ to the target conc. below.

AMG: A280 = 0.12

T. cingulata AMG: 0.34 AGU/ml (corresponding to A280=0.12)

JA126AN: A280 = 0.0024

Assay 1 (w/o JA126)

Sample: (20 microliters AMG + 20 microliters milliQ) x 4 wells

Control: (20 microliters *T. cingulata* AMG + 20 microliters milliQ) x 4 wells

Assay 2 (with JA126)

Sample: (20 μ microliters AMG + 20 microliters JA126) x 4 wells

Control: (20 μ microliters *T. cingulata* AMG + 20 microliters JA126) x 4 wells

2) Add 760 microliters of 3% raw starch suspension to the wells.

3) Plates were incubated at 32°C for 18h with shaking. The glucose conc. was measured using Glucose CII test kit (Wako) after proper dilution, before and 18 hours after incubation.

4) The amount of glucose produced in 18 hours was calculated. The RSH activity was expressed as a relative value to that of *T. cingulata* AMG.

Assay 1: RSH activity (% , w/o JA126) = (Glc. produced from AMG) / (Glc. produced from *T. cingulata* AMG)

Assay 2: RSH activity (% , with JA126) = (Glc produced from AMG+JA126) / (Glc produced from *T. cingulata* AMG+JA126)

Assay condition	AMG (A280=0.003)
	with or without JA126 (A280=0.00006)
	3% raw starch (corn)
	100 mM Na-acetate, pH 4.0
	1 mM CaCl ₂ , 0.025% NaN ₃
	32°C, 18 hours

[0239] The raw starch hydrolysis (RSH) activity of *Pycnoporus sanguineus* glucoamylases was tested in the following conditions:

Substrate: 3% raw starch (corn, Sigma catalog#S9679) suspended in 100 mM sodium acetate, 1 mM calcium chloride and 0.025% sodium azide, pH 4.0

Enzyme dose: Purified *Pycnoporus sanguineus* AMG adjusted to have a final A280 absorbance of 0.003. The hydrolysis tests were performed as comparative tests were the purified *Pycnoporus* or *Talaromyces emersonii* AMG was compared to a purified sample of *Trametes cingulata* AMG (A280=0.003) in the presence (with JA126) or absence (w/o JA126) of purified alpha-amylase JA126AN (A280=0.00006). The produced glucose during the reaction was determined.

Glucose measurement: Glucose CII test (Wako Chemical, catalog# 301-67002)

Temperature: 32°C

Incubation time: 18 hours

[0240] The results of the tests are shown in table 3 below.

Table 3. RSH tests	RSH (%)	
	w/o JA126	w JA126
<i>Pycnoporus sanguineus</i> AMG, P421 B - SEQ ID NO: 6	90%	110%
<i>Pycnoporus sanguineus</i> AMG, P2379 - SEQ ID NO: 4	98%	119%
<i>Talaromyces emersonii</i> T-AMG, P28N - SEQ ID NO: 10	75%	50%
<i>Pycnoporus sanguineus</i> AMG, P1TD - SEQ ID NO: 2	87%	106%

Table 3. RSH tests	RSH (%)	
	w/o JA126	w JA126
<i>Trametes cingulata</i> AMG G1, P13P - SEQ ID NO: 8	100%	100%

Deposit of Biological Material

[0241] The following biological material has been deposited under the terms of the Budapest Treaty with Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Mascheroder Weg 1 B, D-38124 Braunschweig, Germany, and given the following accession number:

Deposit: *E. coli* strain NN059222 with plasmid comprising sequence D4TU (SEQ ID NO: 1)

Accession Number: DSM 23221 Date of Deposit: January 13, 2010

[0242] The strain has been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by foreign patent laws to be entitled thereto. The deposit represents a substantially pure culture of the deposited strain. The deposit is available as required by foreign patent laws in countries wherein counterparts of the subject application or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

SEQUENCE LISTING

[0243]

<110> Landvik, Sara Morant , Marc Dominique Ayabe, Keiichi Coward-Kelly, Guillermo

<120> Polypeptides having glucoamylase activity and polynucleotides encoding same

<130> 11648-WO-PCT

<150> US 61/264,977

<151> 2009-11-30

<160> 10

<170> PatentIn version 3.5

<210> 1

<211> 2107

<212> DNA

<213> *Pycnoporus sanguineus*

<400> 1

```

atgcgcttca cactccttgc atctctggtc ggctctgctg tgggtgcgta tgcacagtgc      60
agcgcggccg acgcgtatgt cgcgtccgag tcccccatcg ccaagcaggg cgtcctgaac      120
aacatcggac cggacgggtc caagcccaac ggcgccaagg taggagatgc accgctctac      180
gtccaccttc ctctattgca agcgtgaca ttagactttg tatgacaggc cggcatcgtc      240
gtagcgagcc cgagcacctc gaacccaac tacctctata catggacgcg cgactcgtcc      300
ctcgtcttca agctgctcat cgaccagttc accagcggcg aggacacgag cctccgtgga      360
ctcatcgatg acttcacctc cgcagaggcc atcctccagc aggtcccgaa cccagcgga      420
accgtctcta ctggcggtt gggcgagccc aagttaaca ttgacgagac cgccttcacc      480
gaccttggg gccctcctca acgcggtacg agtcgcccgc ctctgacctg actttgtacc      540
gtatctcatt cccagtttgc tggatccac agatggcccgc gccctgcgcg cgacgtcgat      600
catcgcgtac cggaaactggc tgctcgacaa cgggaacacg acgtatgtgt cgaacacgct      660
ctggccggtc atccagctcg acctcgacta cgtcgcgaac aactggaacc agtccacatt      720
cgacctctgg gaggagatca actcgtcctc gttcttcacc accgcggtgc agcaccgccc      780
gctgcgcgag ggcgcgacgt tgcgtcccg catcggccag tgcctcgtcg tgagcggata      840
cacgactcag cgggacaacc tgctgtgctt ccttcagggtg acttgatctc ctctctcggt      900
tcgatgcgcg actgattgtg agcgtgcgta gtcgtactgg aaccctagtg gcggtacgt      960
gactgcaaac acgggcggcg gccgctccgg caaggacgca aacacggtgc tcacttccat      1020
ccataccttt gaccocgccg ccggtcgaga cgcgcgacg ttccagccat gctccgacaa      1080
ggcgctctcg aacctcaaag tctacgtcga cgcattccga tccatctata ccatcaacag      1140
cggtatcgcc tccaacgccg ccgtcgctac tggccgttac cccgaggact cgtaccaagg      1200

cggcaatgta agcgcgcgt ctctctgtat acggagacca tggttactca tacatagtcc      1260
cttgacagcc ctggtacctc accacgtccg cggtcgccga acagctctac gacgcgctct      1320
acgtctggga ccagctcggc gcgctcaacg tcacaagcac ctctcttgcc ttcttcagc      1380
agttctcgtc cgggatcagc acgggcacct acccgccctc ctctccacc tacgccacgc      1440
tcaccagcgc gatccgcacc ttgcggagc gcttctctgc ggtcaacgca aagtacacgc      1500
ccgcggacg cgggctcgcg gagcagtaca gcaggaaaga cggcacgccc ctgagcgag      1560
tcgacctcac gtggagctac gccgcgcgc tcacggcggtt tgcggcacgg gaggggaaga      1620
cgtatggcag ctggggcgcc ggggggttga cgtgcccgc gagctgctcg ggcagcgggg      1680
gcgctaccgt tgcggtcacg ttcaacgtcc aggcgacgac tgtctttggt ggtaagtggg      1740
atagagttcc ttacagagca tagagggagt gaaagggtt gctgaccgcg tacctgcaga      1800
gaacatctac atcacgggt ccgttgccgc cctgcagaac tggcccccg acaacgcgct      1860
catcctctcc gcccgaaact acccgacctg gagcagtaag taccctccct cgtccgcgtc      1920
cctcttcacc gcagcagcag gctgaccgcg cgcgcgtgca gtcacggtga acctccccgc      1980
gaacacggtc gtgcagtaca agtacatccg caagttcaac ggacaggtca cctgggagtc      2040
cgaccgaac aaccagatca cgacgccctc gggcgggtcg ttcacacaga acgacgtgtg      2100
gcggtga                                           2107

```

<210> 2

<211> 573

<212> PRT

<213> *Pycnopus sanguineus*

<400> 2

Met Arg Phe Thr Leu Leu Ala Ser Leu Val Gly Leu Val Val Gly Ala
 1 5 10 15
 Tyr Ala Gln Ser Ser Ala Ala Asp Ala Tyr Val Ala Ser Glu Ser Pro
 20 25 30
 Ile Ala Lys Gln Gly Val Leu Asn Asn Ile Gly Pro Asp Gly Ser Lys
 35 40 45
 Ala His Gly Ala Lys Ala Gly Ile Val Val Ala Ser Pro Ser Thr Ser
 50 55 60
 Asn Pro Asn Tyr Leu Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe
 65 70 75 80
 Lys Leu Leu Ile Asp Gln Phe Thr Ser Gly Glu Asp Thr Ser Leu Arg
 85 90 95
 Gly Leu Ile Asp Asp Phe Thr Ser Ala Glu Ala Ile Leu Gln Gln Val
 100 105 110
 Pro Asn Pro Ser Gly Thr Val Ser Thr Gly Gly Leu Gly Glu Pro Lys
 115 120 125
 Phe Asn Ile Asp Glu Thr Ala Phe Thr Asp Pro Trp Gly Arg Pro Gln
 130 135 140
 Arg Asp Gly Pro Ala Leu Arg Ala Thr Ser Ile Ile Arg Tyr Ala Asn
 145 150 155 160
 Trp Leu Leu Asp Asn Gly Asn Thr Thr Tyr Val Ser Asn Thr Leu Trp
 165 170 175
 Pro Val Ile Gln Leu Asp Leu Asp Tyr Val Ala Asn Asn Trp Asn Gln
 180 185 190
 Ser Thr Phe Asp Leu Trp Glu Glu Ile Asn Ser Ser Ser Phe Phe Thr
 195 200 205
 Thr Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Thr Phe Ala Ser
 210 215 220
 Arg Ile Gly Gln Ser Ser Val Val Ser Gly Tyr Thr Thr Gln Ala Asp
 225 230 235 240
 Asn Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr
 245 250 255
 Val Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr
 260 265 270
 Val Leu Thr Ser Ile His Thr Phe Asp Pro Ala Ala Gly Cys Asp Ala
 275 280 285
 Ala Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val
 290 295 300
 Tyr Val Asp Ala Phe Arg Ser Ile Tyr Thr Ile Asn Ser Gly Ile Ala
 305 310 315 320
 Ser Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Gln
 325 330 335
 Gly Gly Asn Pro Trp Tyr Leu Thr Thr Ser Ala Val Ala Glu Gln Leu
 340 345 350
 Tyr Asp Ala Leu Tyr Val Trp Asp Gln Leu Gly Ala Leu Asn Val Thr
 355 360 365

Ser Thr Ser Leu Ala Phe Phe Gln Gln Phe Ser Ser Gly Ile Ser Thr
 370 375 380
 Gly Thr Tyr Pro Ala Ser Ser Ser Thr Tyr Ala Thr Leu Thr Ser Ala
 385 390 395 400
 Ile Arg Thr Phe Ala Asp Gly Phe Leu Ala Val Asn Ala Lys Tyr Thr
 405 410 415
 Pro Ala Asp Gly Gly Leu Ala Glu Gln Tyr Ser Arg Asn Asp Gly Thr
 420 425 430
 Pro Leu Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ala Ala Leu Thr
 435 440 445
 Ala Phe Ala Ala Arg Glu Gly Lys Thr Tyr Gly Ser Trp Gly Ala Ala
 450 455 460
 Gly Leu Thr Val Pro Ala Ser Cys Ser Gly Ser Gly Gly Ala Thr Val
 465 470 475 480
 Ala Val Thr Phe Asn Val Gln Ala Thr Thr Val Phe Gly Glu Asn Ile
 485 490 495
 Tyr Ile Thr Gly Ser Val Ala Ala Leu Gln Asn Trp Ser Pro Asp Asn
 500 505 510
 Ala Leu Ile Leu Ser Ala Ala Asn Tyr Pro Thr Trp Ser Ile Thr Val
 515 520 525
 Asn Leu Pro Ala Asn Thr Val Val Gln Tyr Lys Tyr Ile Arg Lys Phe
 530 535 540
 Asn Gly Gln Val Thr Trp Glu Ser Asp Pro Asn Asn Gln Ile Thr Thr
 545 550 555 560
 Pro Ser Gly Gly Ser Phe Thr Gln Asn Asp Val Trp Arg
 565 570

<210> 3

<211> 2105

<212> DNA

<213> *Pycnopus sanguineus*

<400> 3

atgcgcttca cactccttgc ctctctcatc ggccctcgccg tcggtgcctt tgcacagtcg	60
agcgagtcg acgcgtacgt cgcgtccgag tcccccatcg ccaagcaggg cgtgctcaac	120
aacatcggac cgaacggatc caaggcacac ggccccaagg tcggcaatcc accgctctac	180

gtccaccttc	cgctcgtgca	ggcgctaaca	ccacactttg	tctgacaggg	cggcacgtc	240
gtagcaagcc	cgagcacaga	gaaccggac	tacctctaca	catggacgcg	cgactcctcc	300
ctcgtcttca	agctgctcat	cgaccagttc	accagcgggc	acgacacgtc	cctccgcggt	360
ctcatcgacg	acttcacctc	cgcggaggcc	atccttcagc	aggtctcgaa	ccctagcggg	420
accgtctcta	cgggtggctt	ggcggaacc	aagttcaaca	tgcacgagac	cgccttcacc	480
ggcgcatggg	gccgtcctca	acgcggtatg	aatgcgcccc	cttcaatatc	tctccgcgcc	540
gtgctgatct	ctggcttcgc	cacatatgca	gatggcccg	cctgcgcgc	gacgtccatc	600
atccgctacg	ggaactggct	gctcgacaac	ggaacacga	cgtaactgtc	gaacacgtc	660
tggcgggtca	tccagctcga	cctcgactac	gtcgccgaca	actggaacca	gtccacgttc	720
gacctatggg	aggaggtcga	ctcgtcgtcg	ttctttacca	ctgccgtgca	gcaccgcgcg	780
ctcgtgagg	gcgccacgtt	cgcgtcccg	atcgccaggt	cgtccgtcgt	gagcgggtac	840
accaccocag	cggacaacct	gctatgcttc	cttcagggtga	gccgatcccg	gcgctgtgtg	900
agtgagtga	gtgggcattg	actgactccg	cgtctgcaca	gtcgtactgg	aaccccagtg	960
gaggatacgt	gaccgcgaac	acgggcggcg	gacgtcccg	caaggactcc	aacaccgtgc	1020
tcacctccat	ccacaccttc	gaccccgctg	ccggttgcca	gcgcgcgacc	ttccagccat	1080
gctcggacaa	ggcgctctcg	aacctcaagg	tctacgtcga	cgttttcgc	tccatctaca	1140
ccatcaacaa	cgccatcgcc	tccaacgcgc	cgtcgtac	cggccgttac	cccaggagct	1200
cgtacatggg	agggaaacgta	agtaccattc	cctcacacag	catgccacgg	atactcacgt	1260
gcagtccttc	gacagccctg	gtacctcacc	acgtccgcgg	tcgcagagca	gctctacgac	1320
gcgctctacg	tctgggacca	gctcggcggg	ctcaacgtca	cgagcacctc	cctcgccttc	1380
ttccagcagt	tcgcgtctgg	gtcagcacg	ggcaccctact	cgcctcctc	gtccacgtac	1440
gccacgctca	cgagcgcgat	ccgcagcttc	gcggacggct	tctcgcgat	caacgcaaa	1500
tacacgcccg	cggacgggtg	gctcgcggag	cagtacagca	gaaacgaagg	cacgcgcgtg	1560
agcgcgggtc	acctcacgtg	gagctacgcc	gcggcgctca	cggcggttgc	ggcaaggagg	1620
gggaagacgt	acgggagctg	gggcgcggcg	gggctgacgg	tgcgcgcgag	ctgctcgggc	1680
ggcggaggcg	ctaccgttgc	ggtcacgttc	aacgtccagg	cgaactactgt	ctttggtggt	1740
aagtogattt	gtgttctttc	cagagcatgg	agggagtoga	cgggatgctg	acggcgtacc	1800
tgcagagAAC	atctacatca	cggctccgt	cgcgcctc	caaaactggt	ccccggacaa	1860
tgcgctcatt	ctctccgcg	cgaactaccc	gacctggagc	agtaactacc	ccgcctcatc	1920
tcttctcgcc	caccgcagca	ggtgacgag	cgcgcgcagt	cacggtgaac	ctgcgcgcga	1980
acacggctgt	gcagtacaag	tacatccgca	agttcaacgg	acaggtcacc	tgggagtcgg	2040
accogaacaa	ccagatcacg	acgccctcgg	gcgggtcgtt	cacacagaac	gacgtgtggc	2100
ggtaa						2105
<210>	4					
<211>	573					
<212>	PRT					
<213>	Pycnopus sanguineus					
<400>	4					

Met Arg Phe Thr Leu Leu Ala Ser Leu Ile Gly Leu Ala Val Gly Ala
 1 5 10 15
 Phe Ala Gln Ser Ser Ala Val Asp Ala Tyr Val Ala Ser Glu Ser Pro
 20 25 30
 Ile Ala Lys Gln Gly Val Leu Asn Asn Ile Gly Pro Asn Gly Ser Lys
 35 40 45
 Ala His Gly Ala Lys Ala Gly Ile Val Val Ala Ser Pro Ser Thr Glu
 50 55 60
 Asn Pro Asp Tyr Leu Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe
 65 70 75 80
 Lys Leu Leu Ile Asp Gln Phe Thr Ser Gly Asp Asp Thr Ser Leu Arg
 85 90 95
 Gly Leu Ile Asp Asp Phe Thr Ser Ala Glu Ala Ile Leu Gln Gln Val
 100 105 110
 Ser Asn Pro Ser Gly Thr Val Ser Thr Gly Gly Leu Gly Glu Pro Lys
 115 120 125
 Phe Asn Ile Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln
 130 135 140
 Arg Asp Gly Pro Ala Leu Arg Ala Thr Ser Ile Ile Arg Tyr Ala Asn
 145 150 155 160
 Trp Leu Leu Asp Asn Gly Asn Thr Thr Tyr Val Ser Asn Thr Leu Trp
 165 170 175
 Pro Val Ile Gln Leu Asp Leu Asp Tyr Val Ala Asp Asn Trp Asn Gln
 180 185 190
 Ser Thr Phe Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr
 195 200 205
 Thr Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Thr Phe Ala Ser
 210 215 220

Arg Ile Gly Gln Ser Ser Val Val Ser Gly Tyr Thr Thr Gln Ala Asp
 225 230 235 240
 Asn Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr
 245 250 255
 Val Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ser Asn Thr
 260 265 270
 Val Leu Thr Ser Ile His Thr Phe Asp Pro Ala Ala Gly Cys Asp Ala
 275 280 285
 Ala Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val
 290 295 300
 Tyr Val Asp Ala Phe Arg Ser Ile Tyr Thr Ile Asn Asn Gly Ile Ala
 305 310 315 320
 Ser Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Met
 325 330 335
 Gly Gly Asn Pro Trp Tyr Leu Thr Thr Ser Ala Val Ala Glu Gln Leu
 340 345 350
 Tyr Asp Ala Leu Tyr Val Trp Asp Gln Leu Gly Gly Leu Asn Val Thr
 355 360 365
 Ser Thr Ser Leu Ala Phe Phe Gln Gln Phe Ala Ser Gly Leu Ser Thr
 370 375 380
 Gly Thr Tyr Ser Ala Ser Ser Ser Thr Tyr Ala Thr Leu Thr Ser Ala
 385 390 395 400
 Ile Arg Ser Phe Ala Asp Gly Phe Leu Ala Ile Asn Ala Lys Tyr Thr
 405 410 415
 Pro Ala Asp Gly Gly Leu Ala Glu Gln Tyr Ser Arg Asn Asp Gly Thr
 420 425 430
 Pro Leu Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ala Ala Leu Thr
 435 440 445
 Ala Phe Ala Ala Arg Glu Gly Lys Thr Tyr Gly Ser Trp Gly Ala Ala
 450 455 460
 Gly Leu Thr Val Pro Ala Ser Cys Ser Gly Gly Gly Gly Ala Thr Val
 465 470 475 480
 Ala Val Thr Phe Asn Val Gln Ala Thr Thr Val Phe Gly Glu Asn Ile
 485 490 495
 Tyr Ile Thr Gly Ser Val Ala Ala Leu Gln Asn Trp Ser Pro Asp Asn
 500 505 510
 Ala Leu Ile Leu Ser Ala Ala Asn Tyr Pro Thr Trp Ser Ile Thr Val
 515 520 525
 Asn Leu Pro Ala Asn Thr Val Val Gln Tyr Lys Tyr Ile Arg Lys Phe
 530 535 540
 Asn Gly Gln Val Thr Trp Glu Ser Asp Pro Asn Asn Gln Ile Thr Thr
 545 550 555 560
 Pro Ser Gly Gly Ser Phe Thr Gln Asn Asp Val Trp Arg
 565 570

<210> 5

<211> 2105

<212> DNA

<213> Pycnopus sanguineus

<400> 5

```

atgcgcttca cactccttgc ctctctcacc ggctcgcgcg tcggtgcctt tgcacagtcg      60
agcgcagtcg acgcgtacgt cgcgtccgag tcccccatcg ccaagcaggc cgtgctcaac      120
aacatcggac cgaacggatc caaggcacac ggcgccaagg tcggcaatcc accgctctac      180
gtccaccttc cgtccgtgca ggcgctaaca ccacactttg tctgacaggc cggcatcgtc      240
gtagcaagcc cgagcacaga gaaccggac tacctctaca catggacgcg cgactcctcc      300
ctcgtcttca agctgctcat cgaccagttc accagcggcg acgacacgtc cctccgcggt      360
ctcatcgacg acttcacctc cgcggaggcc atccttcagc aggtctcgaa ccctagcggg      420
accgtctcta cgggtggctt gggcgaacc aagttcaaca tcgacgagac cgccttcacc      480
ggcgcagtcg gccgtcctca acgcgggatg aatgcgccc cttcaatatc tctccgcgcc      540
gtgctgatct ctggtctcgc cacatatgca gatggcccg ccctgcgcgc gacgtccatc      600
atccgctacg cgaactggct gctcgacaac ggaacacga cgtacgtgc gaacacgtc      660
tgcccggtca tccagctcga cctcgactac gtcgcgcaca actggaacca gtccacgttc      720
gacctatggg aggaggtcga ctcgtcgtcg ttctttacca ctgccgtgca gcaccgcgcg      780
ctccgtgagg gcgccacgtt cgcgtccgcg atcggccagt cgtccgtcgt gagcgggtac      840
accaccaggc cggacaacct gctgtgcttc ctccaggtga gccgatcccg gcgctgtgtg      900
agtgagtcaa gtgggcattg actgactcgc cgtcgcaca gtcgtactgg aaccccagtg      960
gaggatacgt gaccgcgaac acggcgggcg gacgctccgg caaggactcc aacaccgtgc      1020
tcacctctat ccacaccttc gaccccgctg ccggttgcca cgcgcgcacc ttccagccat      1080

gctcggacaa ggcgtctcgc aacctcaagg tctacgtcga cgttttcgcg tccatctaca      1140
ccatcaacaa cggcatcgcc tccaacgcgc cgtcgtctac cggcgtttac ccgaggact      1200
cgtacatggg aggaacacga agtaccattc cctcacacag catgccacgg atactcactg      1260
gcagtccttc gacagccctg gtacctcacc acgtccgcgc tcgcagagca gctctacgac      1320
gcgctctacg tctgggacca gctcggcggg ctcaacgtca cgagcacctc cctcgccttc      1380
ttccagcagt tcgctgtctg gctcagcacg ggcacctact ccgcctcctc gtccacgtac      1440
gccacgtcga cgagcgcgat ccgcagcttc gcggacggct tcctcgogat caacgcaaag      1500
tacacgcccg cggacggtgg gctcgcggag cagtacgca ggaacgacgg caccgcgctg      1560
agcgcggtcg acctcactg gagctacgcc gcggcgctca cggcgtttgt ggcaaggag      1620
gggaagacgt acgggagctg gggcgcgcg ggcgtgacgg tgcccgcgag ctgctcgggc      1680
ggcgggtggc ctaccgttgc ggtcacgttc aacgtccagg cgactactgt ctttgggtgt      1740
aagtcgattt gtgttcttcc ccgagcatgg agggagtcga cgggatgctg acagcgtacc      1800
tgcagagaac atctacatca ccggtccgt cgcgcgcctc caaaactggt ccccggaaca      1860
tgcgctcacc ctctccgcgc cgaactaccc gacctggagc agtacgtacc ccgcctcacc      1920
tcttctcgcc caccgcagca ggctgacgag cgcgcgcagt cacggtgaac ctgccggcga      1980
acacggtcgt gcagtacaag tacatccgca agttcaacgg acaggtcacc tgggagtcgg      2040
accgaacaa ccagatcacg acgcccctcg gcgggtcgtt cacacagaac gacgtgtggc      2100
ggtaa                                          2105

```

<210> 6

<211> 573

<212> PRT

<213> Pycnopus sanguineus

<400> 6

Met Arg Phe Thr Leu Leu Ala Ser Leu Ile Gly Leu Ala Val Gly Ala
 1 5 10 15
 Phe Ala Gln Ser Ser Ala Val Asp Ala Tyr Val Ala Ser Glu Ser Pro
 20 25 30
 Ile Ala Lys Gln Gly Val Leu Asn Asn Ile Gly Pro Asn Gly Ser Lys
 35 40 45
 Ala His Gly Ala Lys Ala Gly Ile Val Val Ala Ser Pro Ser Thr Glu
 50 55 60
 Asn Pro Asp Tyr Leu Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe
 65 70 75 80
 Lys Leu Leu Ile Asp Gln Phe Thr Ser Gly Asp Asp Thr Ser Leu Arg
 85 90 95
 Gly Leu Ile Asp Asp Phe Thr Ser Ala Glu Ala Ile Leu Gln Gln Val
 100 105 110
 Ser Asn Pro Ser Gly Thr Val Ser Thr Gly Gly Leu Gly Glu Pro Lys
 115 120 125
 Phe Asn Ile Asp Glu Thr Ala Phe Thr Gly Ala Trp Gly Arg Pro Gln
 130 135 140
 Arg Asp Gly Pro Ala Leu Arg Ala Thr Ser Ile Ile Arg Tyr Ala Asn
 145 150 155 160
 Trp Leu Leu Asp Asn Gly Asn Thr Thr Tyr Val Ser Asn Thr Leu Trp
 165 170 175
 Pro Val Ile Gln Leu Asp Leu Asp Tyr Val Ala Asp Asn Trp Asn Gln
 180 185 190
 Ser Thr Phe Asp Leu Trp Glu Glu Val Asp Ser Ser Ser Phe Phe Thr
 195 200 205
 Thr Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Thr Phe Ala Ser
 210 215 220
 Arg Ile Gly Gln Ser Ser Val Val Ser Gly Tyr Thr Thr Gln Ala Asp
 225 230 235 240
 Asn Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Ser Gly Gly Tyr
 245 250 255
 Val Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ser Asn Thr
 260 265 270
 Val Leu Thr Ser Ile His Thr Phe Asp Pro Ala Ala Gly Cys Asp Ala
 275 280 285
 Ala Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val
 290 295 300
 Tyr Val Asp Ala Phe Arg Ser Ile Tyr Thr Ile Asn Asn Gly Ile Ala
 305 310 315 320
 Ser Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Met
 325 330 335
 Gly Gly Asn Pro Trp Tyr Leu Thr Thr Ser Ala Val Ala Glu Gln Leu

340 345 350
 Tyr Asp Ala Leu Tyr Val Trp Asp Gln Leu Gly Gly Leu Asn Val Thr
 355 360 365
 Ser Thr Ser Leu Ala Phe Phe Gln Gln Phe Ala Ser Gly Leu Ser Thr
 370 375 380
 Gly Thr Tyr Ser Ala Ser Ser Ser Thr Tyr Ala Thr Leu Thr Ser Ala
 385 390 395 400
 Ile Arg Ser Phe Ala Asp Gly Phe Leu Ala Ile Asn Ala Lys Tyr Thr
 405 410 415
 Pro Ala Asp Gly Gly Leu Ala Glu Gln Tyr Ser Arg Asn Asp Gly Thr
 420 425 430
 Pro Leu Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ala Ala Leu Thr
 435 440 445
 Ala Phe Val Ala Arg Glu Gly Lys Thr Tyr Gly Ser Trp Gly Ala Ala
 450 455 460
 Gly Leu Thr Val Pro Ala Ser Cys Ser Gly Gly Gly Gly Ala Thr Val
 465 470 475 480
 Ala Val Thr Phe Asn Val Gln Ala Thr Thr Val Phe Gly Glu Asn Ile
 485 490 495
 Tyr Ile Thr Gly Ser Val Ala Ala Leu Gln Asn Trp Ser Pro Asp Asn
 500 505 510
 Ala Leu Ile Leu Ser Ala Ala Asn Tyr Pro Thr Trp Ser Ile Thr Val
 515 520 525
 Asn Leu Pro Ala Asn Thr Val Val Gln Tyr Lys Tyr Ile Arg Lys Phe
 530 535 540
 Asn Gly Gln Val Thr Trp Glu Ser Asp Pro Asn Asn Gln Ile Thr Thr
 545 550 555 560
 Pro Ser Gly Gly Ser Phe Thr Gln Asn Asp Val Trp Arg
 565 570

<210> 7

<211> 1725

<212> DNA

<213> Trametes cingulata

<400> 7

```

atgcgtttca cgtcctcac ctccctcctg ggctcgcgc toggcgcgtt cgcgcagtcg      60
agtgcggccg acgcgtacgt cgcgtccgaa tcgcccatcg ccaaggcggg tgtgctcgcc      120
aacatcgggc ccagcggctc caagtccaac ggagcaaagg caggcatcgt gattgcaagt      180
ccgagcacat ccaaccgaa ctacctgtac acatggacgc gcgactcgto cctcgtgttc      240
aaggcgctca tcgaccagtt caccactggc gaagatacct cgctccgaac tctgattgac      300
gagttcacct cggcggaggc catactccag caggtgccga acccgagcgg gacagtcagc      360
actggaggcc tcggcgagcc caagttcaac atcgacgaga ccgcgttcac ggatgcctgg      420
ggtcgtctc agcgcgatgg tcccgtctc cgggcgactg ccatcatcac ctacgccaac      480
tgggtcctcg acaacaagaa cagcacctac gtgaccaaca ctctctggcc tatcatcaag      540
ctcgacctcg actacgtcgc cagcaactgg aaccagtcca cgtttgatct ctgggaggag      600
attaactcct cgtcgttctt cactaccgcc gtccagcacc gtgctctcgc cgagggcgcg      660
actttcgcta atcgcatcg acaaacctcg gtggtcagcg ggtacaccac ccaagcaaac      720
aaccttctct gcttctcgca gtctactgg aacccaccg gcggctatat caccgcaaac      780
acgggcggcg gccgctctgg caaggacgcg aacaccgttc tcacgtcgat ccacaccttc      840
gacccggccg ctggatcgca cgtctgtacg ttccagccgt gctcggacaa ggcgctgtcg      900
aacttgaagg tgtactcga tgcgttcgc tgatctact ccatcaacag cgggatcgcc      960
tcgaatcgcg ccgttgctac cggcgcgtac ccgaggaca gctacatggg cggaaccaca      1020
tggtaacctc ccacctcgc cgtcgtcgtg cagctctacg atgcgctcat tgtgtggaac      1080
aaacttggcg cctgaacgt cagcgcacc tccctccct tcttcagca gttctcgtca      1140
ggcgtcaccg tcggcaccta tgcctcatcc tcgtccacct tcaagacgct cacttcgcc      1200
atcaagacct tcgcgcagg ctctctcgc gtcaacgcca agtacacgc ctcgaacggc      1260
ggccttgctg aacagtacg ccggagcaac ggctcgcgg tcagcgtgtg ggacctgacg      1320
tggagctatg ctgctgccct caoctgttt gctgcgcgt caggcaagac gtatgcgagc      1380
tggggcgcg cggtttgac tgtcccgacg acttgctcgg ggagtggcg tgcggggact      1440
gtggccgtca cttcaacgt gcaggcgacc accgtgttcg gcgagaacat ttacatcaca      1500
ggctcgttcc ccgtctcca gaactggtcg ccgacaacg cgctcatcct ctcagcggcc      1560
aactaccca cttggagcat caccgtgaac ctgccggcga gcacgacgat cgagtacaag      1620
tacattcgca agttcaacg cgcggtcacc tgggagtcgg accgaacaa ctcgatcacg      1680
acgcccgcga cgggcacgtt caccagaac gacacctggc ggtag      1725

```

<210> 8

<211> 574

<212> PRT

<213> Trametes cingulata

<400> 8

Met Arg Phe Thr Leu Leu Thr Ser Leu Leu Gly Leu Ala Leu Gly Ala
 1 5 10 15
 Phe Ala Gln Ser Ser Ala Ala Asp Ala Tyr Val Ala Ser Glu Ser Pro
 20 25 30
 Ile Ala Lys Ala Gly Val Leu Ala Asn Ile Gly Pro Ser Gly Ser Lys
 35 40 45
 Ser Asn Gly Ala Lys Ala Gly Ile Val Ile Ala Ser Pro Ser Thr Ser
 50 55 60
 Asn Pro Asn Tyr Leu Tyr Thr Trp Thr Arg Asp Ser Ser Leu Val Phe
 65 70 75 80
 Lys Ala Leu Ile Asp Gln Phe Thr Thr Gly Glu Asp Thr Ser Leu Arg
 85 90 95
 Thr Leu Ile Asp Glu Phe Thr Ser Ala Glu Ala Ile Leu Gln Gln Val
 100 105 110
 Pro Asn Pro Ser Gly Thr Val Ser Thr Gly Gly Leu Gly Glu Pro Lys
 115 120 125
 Phe Asn Ile Asp Glu Thr Ala Phe Thr Asp Ala Trp Gly Arg Pro Gln
 130 135 140
 Arg Asp Gly Pro Ala Leu Arg Ala Thr Ala Ile Ile Thr Tyr Ala Asn
 145 150 155 160
 Trp Leu Leu Asp Asn Lys Asn Thr Thr Tyr Val Thr Asn Thr Leu Trp
 165 170 175
 Pro Ile Ile Lys Leu Asp Leu Asp Tyr Val Ala Ser Asn Trp Asn Gln
 180 185 190
 Ser Thr Phe Asp Leu Trp Glu Glu Ile Asn Ser Ser Ser Phe Phe Thr
 195 200 205
 Thr Ala Val Gln His Arg Ala Leu Arg Glu Gly Ala Thr Phe Ala Asn
 210 215 220
 Arg Ile Gly Gln Thr Ser Val Val Ser Gly Tyr Thr Thr Gln Ala Asn
 225 230 235 240
 Asn Leu Leu Cys Phe Leu Gln Ser Tyr Trp Asn Pro Thr Gly Gly Tyr
 245 250 255

```

Ile Thr Ala Asn Thr Gly Gly Gly Arg Ser Gly Lys Asp Ala Asn Thr
260                               265                               270

Val Leu Thr Ser Ile His Thr Phe Asp Pro Ala Ala Gly Cys Asp Ala
275                               280                               285

Val Thr Phe Gln Pro Cys Ser Asp Lys Ala Leu Ser Asn Leu Lys Val
290                               295                               300

Tyr Val Asp Ala Phe Arg Ser Ile Tyr Ser Ile Asn Ser Gly Ile Ala
305                               310                               315                               320

Ser Asn Ala Ala Val Ala Thr Gly Arg Tyr Pro Glu Asp Ser Tyr Met
325                               330                               335

Gly Gly Asn Pro Trp Tyr Leu Thr Thr Ser Ala Val Ala Glu Gln Leu
340                               345                               350

Tyr Asp Ala Leu Ile Val Trp Asn Lys Leu Gly Ala Leu Asn Val Thr
355                               360                               365

Ser Thr Ser Leu Pro Phe Phe Gln Gln Phe Ser Ser Gly Val Thr Val
370                               375                               380

Gly Thr Tyr Ala Ser Ser Ser Ser Thr Phe Lys Thr Leu Thr Ser Ala
385                               390                               395                               400

Ile Lys Thr Phe Ala Asp Gly Phe Leu Ala Val Asn Ala Lys Tyr Thr
405                               410                               415

Pro Ser Asn Gly Gly Leu Ala Glu Gln Tyr Ser Arg Ser Asn Gly Ser
420                               425                               430

Pro Val Ser Ala Val Asp Leu Thr Trp Ser Tyr Ala Ala Ala Leu Thr
435                               440                               445

Ser Phe Ala Ala Arg Ser Gly Lys Thr Tyr Ala Ser Trp Gly Ala Ala
450                               455                               460

Gly Leu Thr Val Pro Thr Thr Cys Ser Gly Ser Gly Gly Ala Gly Thr
465                               470                               475                               480

Val Ala Val Thr Phe Asn Val Gln Ala Thr Thr Val Phe Gly Glu Asn
485                               490                               495

Ile Tyr Ile Thr Gly Ser Val Pro Ala Leu Gln Asn Trp Ser Pro Asp
500                               505                               510

Asn Ala Leu Ile Leu Ser Ala Ala Asn Tyr Pro Thr Trp Ser Ile Thr
515                               520                               525

Val Asn Leu Pro Ala Ser Thr Thr Ile Glu Tyr Lys Tyr Ile Arg Lys
530                               535                               540

Phe Asn Gly Ala Val Thr Trp Glu Ser Asp Pro Asn Asn Ser Ile Thr
545                               550                               555                               560

Thr Pro Ala Ser Gly Thr Phe Thr Gln Asn Asp Thr Trp Arg
565                               570

```

<210> 9

<211> 2076

<212> DNA

<213> Talaromyces emersonii

<400> 9

atggcgctccc	tcgttgctgg	cgctctctgc	atcctgggccc	tgacgcctgc	tgcatattgca	60
cgagcgcccc	ttgcagcgcg	agccaccggt	tccttggaact	cctttctcgc	aaccgaaact	120
ccaattgccc	tccaaggcgt	gctgaacaac	atcggggccc	atgggtgtga	tgtggcagga	180
gcaagcgccg	gcattgtggt	tgccagtcgg	agcagagcgg	accocaaattg	taggttcttt	240
cccaccagaa	attacttatt	taaatacagc	ctctgacagg	ttgaagattt	ctactcctgg	300
acacgtgacg	cagcgctcac	ggccaaatac	ctcgtogacg	ccttcacogc	gggcaacaag	360
gacctagagc	agaccatcca	gcagtacatc	agcgcgcagg	cgaaggtgca	aactatctcc	420
aatccgtccg	gagatttata	caccgggtggc	ttaggtgagc	ccaagttcaa	tgtgaatgag	480
acggctttta	cggggccctg	gggtcgtcca	cagaggggagc	gaccagcgtt	gagagcgacg	540
gccctcattg	cgtatgcgaa	ctatctcatc	gtaagcttct	gctcgcctgc	cttctctctg	600
ctcgtatgct	aagtagtcct	gtcaggacaa	cggcgaggct	tcgactgcgg	atgagatcat	660
ctggccgatt	gtccagaatg	atctgtccta	catcacccaa	tactggaact	catccacctt	720
cggtaggcaa	atgaatatct	cgcacacagc	gtgggtactaa	tttgattcag	acctctggga	780
agaagtagaa	ggatcctcat	tottcacaac	cgcgcgtgcaa	caccgcgcgc	tggtcgaagg	840
caatgcactg	gcaacaaggc	tgaaccacac	gtgctccaac	tgctgtctct	aggccctcca	900
ggtcctgtgt	ttcctgcagt	catactggac	cggatcgtat	gttctggcca	actttggtgg	960
cagcggctgt	tcgggcaagg	acgtgaattc	gattctgggc	agcatccaca	cctttgatcc	1020
cgcgcggaggc	tgtgacgact	cgaccttcca	gccgtgttcg	gccgcgtcct	tgggcaaatca	1080
caaggtggtc	accgactcgt	tcggagtat	ctatgcgac	aactcaggca	tcgcagaggg	1140
atctgccgtg	gcagtcggcc	gctaccctga	ggatgtctac	cagggcggga	accctcggtg	1200
cctggccaca	gcagcggctg	cagagcagct	ttacgacgcc	atctaccagt	ggaagaagat	1260
cggctcgata	agtatcacgg	acgttagtct	gccatttttc	caggatatct	acccttctgc	1320
cgcggtgggc	acctataact	ctggctccac	gactttcaac	gacatcatct	cggccgtcca	1380
gacgtatggt	gatggatata	tgagtattgt	cgtacgtttt	gccttagatt	ctcaggtgta	1440
aagaaaaaaa	tggaactaac	tcagttctag	gagaaatata	ctccctcaga	cggctctctt	1500
accgaacaat	tctcccgtag	agacggcact	cgcctttctg	cctctgccct	gacttggtcg	1560
tacgcttctc	tactaacccg	ttcggcccg	agacagtcgg	tcgtccctgc	ttcctggggc	1620
gaaagctccg	caagcagcgt	cctgcgcgtc	tgctctgcca	cctctgccac	gggcccatac	1680
agcacggcta	ccaacacccg	ctggccaagc	tctggctctg	gcagctcaac	aaccaccagt	1740
agcgcgccat	gcaccactcc	tacctctgtg	gctgtgacct	tcgacgaaat	cgtcagcacc	1800
agttacgggg	agacaatcta	cctggccggc	tcgatcccg	agctgggcaa	ctggtccacg	1860
gccagcgcca	tccccctccg	cgcggatgct	tacaccaaca	gcaaccgcgt	ctggtacgtg	1920
accgtcaatc	tgccccctgg	caccagcttc	gagtacaagt	tcttcaagaa	ccagacggac	1980
gggaccatcg	tctgggaaga	cgaaccgaac	cggctgtaca	cggctccago	gtactgtggg	2040
cagactaccg	ccattcttga	cgatagttgg	cagtga			2076

<210> 10

<211> 618

<212> PRT

<213> Talaromyces emersonii

<400> 10

Met Ala Ser Leu Val Ala Gly Ala Leu Cys Ile Leu Gly Leu Thr Pro
 1 5 10 15
 Ala Ala Phe Ala Arg Ala Pro Val Ala Ala Arg Ala Thr Gly Ser Leu
 20 25 30
 Asp Ser Phe Leu Ala Thr Glu Thr Pro Ile Ala Leu Gln Gly Val Leu
 35 40 45
 Asn Asn Ile Gly Pro Asn Gly Ala Asp Val Ala Gly Ala Ser Ala Gly
 50 55 60
 Ile Val Val Ala Ser Pro Ser Arg Ser Asp Pro Asn Tyr Phe Tyr Ser
 65 70 75 80
 Trp Thr Arg Asp Ala Ala Leu Thr Ala Lys Tyr Leu Val Asp Ala Phe
 85 90 95
 Ile Ala Gly Asn Lys Asp Leu Glu Gln Thr Ile Gln Gln Tyr Ile Ser
 100 105 110
 Ala Gln Ala Lys Val Gln Thr Ile Ser Asn Pro Ser Gly Asp Leu Ser
 115 120 125
 Thr Gly Gly Leu Gly Glu Pro Lys Phe Asn Val Asn Glu Thr Ala Phe
 130 135 140
 Thr Gly Pro Trp Gly Arg Pro Gln Arg Asp Gly Pro Ala Leu Arg Ala
 145 150 155 160
 Thr Ala Leu Ile Ala Tyr Ala Asn Tyr Leu Ile Asp Asn Gly Glu Ala
 165 170 175
 Ser Thr Ala Asp Glu Ile Ile Trp Pro Ile Val Gln Asn Asp Leu Ser
 180 185 190
 Tyr Ile Thr Gln Tyr Trp Asn Ser Ser Thr Phe Asp Leu Trp Glu Glu
 195 200 205
 Val Glu Gly Ser Ser Phe Phe Thr Thr Ala Val Gln His Arg Ala Leu
 210 215 220
 Val Glu Gly Asn Ala Leu Ala Thr Arg Leu Asn His Thr Cys Ser Asn
 225 230 235 240
 Cys Val Ser Gln Ala Pro Gln Val Leu Cys Phe Leu Gln Ser Tyr Trp
 245 250 255
 Thr Gly Ser Tyr Val Leu Ala Asn Phe Gly Gly Ser Gly Arg Ser Gly
 260 265 270
 Lys Asp Val Asn Ser Ile Leu Gly Ser Ile His Thr Phe Asp Pro Ala
 275 280 285
 Gly Gly Cys Asp Asp Ser Thr Phe Gln Pro Cys Ser Ala Arg Ala Leu
 290 295 300
 Ala Asn His Lys Val Val Thr Asp Ser Phe Arg Ser Ile Tyr Ala Ile
 305 310 315 320
 Asn Ser Gly Ile Ala Glu Gly Ser Ala Val Ala Val Gly Arg Tyr Pro
 325 330 335
 Glu Asp Val Tyr Gln Gly Gly Asn Pro Trp Tyr Leu Ala Thr Ala Ala
 340 345 350
 Ala Ala Glu Gln Leu Tyr Asp Ala Ile Tyr Gln Trp Lys Lys Ile Gly
 355 360 365
 Ser Ile Ser Ile Thr Asp Val Ser Leu Pro Phe Phe Gln Asp Ile Tyr
 370 375 380

```

Pro Ser Ala Ala Val Gly Thr Tyr Asn Ser Gly Ser Thr Thr Phe Asn
385                390                395                400

Asp Ile Ile Ser Ala Val Gln Thr Tyr Gly Asp Gly Tyr Leu Ser Ile
405                410                415

Val Glu Lys Tyr Thr Pro Ser Asp Gly Ser Leu Thr Glu Gln Phe Ser
420                425                430

Arg Thr Asp Gly Thr Pro Leu Ser Ala Ser Ala Leu Thr Trp Ser Tyr
435                440                445

Ala Ser Leu Leu Thr Ala Ser Ala Arg Arg Gln Ser Val Val Pro Ala
450                455                460

Ser Trp Gly Glu Ser Ser Ala Ser Ser Val Pro Ala Val Cys Ser Ala
465                470                475                480

Thr Ser Ala Thr Gly Pro Tyr Ser Thr Ala Thr Asn Thr Val Trp Pro
485                490                495

Ser Ser Gly Ser Gly Ser Ser Thr Thr Thr Ser Ser Ala Pro Cys Thr
500                505                510

Thr Pro Thr Ser Val Ala Val Thr Phe Asp Glu Ile Val Ser Thr Ser
515                520                525

Tyr Gly Glu Thr Ile Tyr Leu Ala Gly Ser Ile Pro Glu Leu Gly Asn
530                535                540

Trp Ser Thr Ala Ser Ala Ile Pro Leu Arg Ala Asp Ala Tyr Thr Asn
545                550                555                560

Ser Asn Pro Leu Trp Tyr Val Thr Val Asn Leu Pro Pro Gly Thr Ser
565                570                575

Phe Glu Tyr Lys Phe Phe Lys Asn Gln Thr Asp Gly Thr Ile Val Trp
580                585                590

Glu Asp Asp Pro Asn Arg Ser Tyr Thr Val Pro Ala Tyr Cys Gly Gln
595                600                605

Thr Thr Ala Ile Leu Asp Asp Ser Trp Gln
610                615

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2006069289A \[0008\]](#)
- [WO9517413A \[0059\]](#)
- [WO9522625A \[0059\]](#)
- [US5223409A \[0059\]](#)
- [WO9206204A \[0059\]](#)
- [WO0056900A \[0091\] \[0091\] \[0091\]](#)
- [WO9600787A \[0091\] \[0126\]](#)
- [WO0024883A \[0113\] \[0113\]](#)

- EP0238023A [0126]
- WO9114772A [0142]
- WO9220777A [0157]
- WO03066816A [0157]
- WO03066826A [0157]
- WO2004080923A [0157] [0195] [0195]
- WO2004081193A [0157]
- WO9919467A [0189] [0189] [0188] [0188] [0189] [0189] [0189] [0189] [0191] [0191] [0191]
- WO9623873A [0189] [0189]
- WO9623874A [0189] [0194]
- WO9741213A [0189]
- WO0060059A [0189]
- WO0210355A [0189]
- US6093562A [0189]
- US6297038B [0189]
- US6187576B [0189]
- US4598048A [0190]
- US4604355A [0190]
- US6162628A [0190]
- WO8901969A [0195]
- WO2005003311A [0198]
- US20050054071A [0198] [0200]
- US60638614B [0198]
- US638614P [0199] [0199] [0199] [0199]
- EP0140410B2 [0229]
- US61264977B [0243]

Non-patent literature cited in the description

- NIELSEN et al. Protein Engineering, 1997, vol. 10, 1-6 [0018]
- NEEDLEMANWUNSCHJ. Mol. Biol., 1970, vol. 48, 443-453 [0021]
- RICE et al. EMBOSS: The European Molecular Biology Open Software Suite Trends in Genetics, 2000, vol. 16, 276-277 [0021]
- RICE et al. EMBOSS: The European Molecular Biology Open Software Suite, 2000, [0022]
- J. SAMBROOKE.F. FRITSCHT. MANIATIS Molecular Cloning, A Laboratory Manual Cold Spring Harbor 19890000 [0041]
- BOLTON MCCARTHY Proceedings of the National Academy of Sciences USA, 1962, vol. 48, 1390- [0051]
- H. NEURATH R.L. HILL The Proteins, Academic Press 19790000 [0055]
- CUNNINGHAM WELLS Science, 1989, vol. 244, 1081-1085 [0058]
- HILTON et al. J. Biol. Chem., 1996, vol. 271, 4699-4708 [0058]
- DE VOS et al. Science, 1992, vol. 255, 306-312 [0058]
- SMITH et al. J. Mol. Biol., vol. 224, 899-904 [0058]
- WLODAVER et al. FEBS Lett., vol. 309, 59-64 [0058]
- REIDHAAR-OLSON SAUERS Science, 1988, vol. 241, 53-57 [0059]
- BOWIESAUER Proc. Natl. Acad. Sci. USA, 1989, vol. 86, 2152-2156 [0059]
- LOWMAN et al. Biochem., 1991, vol. 30, 10832-10837 [0059]
- DERBYSHIRE et al. Gene, 1986, vol. 46, 145- [0059]
- NER et al. DNA, 1988, vol. 7, 127- [0059]
- NESS et al. Nature Biotechnology, 1999, vol. 17, 893-896 [0060]
- MARTIN et al. J. Ind. Microbiol. Biotechnol., 2003, vol. 3, 568-76 [0069]
- SVETINA et al. J. Biotechnol., 2000, vol. 76, 245-251 [0069]
- RASMUSSEN-WILSON et al. Appl. Environ. Microbiol., 1997, vol. 63, 3488-3493 [0069]
- WARD et al. Biotechnology, vol. 13, 498-503 [0069]
- CONTRERAS et al. Biotechnology, 1991, vol. 9, 378-381 [0069]
- EATON et al. Biochem., 1986, vol. 25, 505-512 [0069]

- **COLLINS-RACIE et al.**Biotechnology, 1995, vol. 13, 982-987 [\[0069\]](#)
- **CARTER et al.**Proteins: Structure, Function, and Genetics, 1989, vol. 6, 240-248 [\[0069\]](#)
- **STEVENS**Drug Discovery World, 2003, vol. 4, 35-48 [\[0069\]](#)
- **INNIS et al.**PCR: A Guide to Methods and ApplicationAcademic Press19900000 [\[0078\]](#)
- **FORD et al.**Protein Expression and Purification, 1991, vol. 2, 95-107 [\[0090\]](#)
- **ROMANOS et al.**Yeast, 1992, vol. 8, 423-488 [\[0092\]](#)
- **GUOSHERMAN**Molecular Cellular Biology, 1995, vol. 15, 5983-5990 [\[0101\]](#)
- **GEMS et al.**Gene, 1991, vol. 98, 61-67 [\[0113\]](#)
- **CULLEN et al.**Nucleic Acids Research, 1987, vol. 15, 9163-9175 [\[0113\]](#)
- **HAWKSWORTH et al.**Ainsworth and Bisby's Dictionary of The FungiUniversity Press19950000 [\[0120\]](#)
- **Biology and Activities of Yeast**Soc. App. Bacteriol. Symposium Series No. 919800000 [\[0121\]](#)
- **YELTON et al.**Proceedings of the National Academy of Sciences USA, 1984, vol. 81, 1470-1474 [\[0126\]](#)
- **MALARDIER et al.**Gene, 1989, vol. 78, 147-156 [\[0126\]](#)
- **Guide to Yeast Genetics and Molecular Biology**BECKERGUARENTEMethods in EnzymologyAcademic Pressvol. 194, 182-187 [\[0126\]](#)
- **ITO et al.**Journal of Bacteriology, 1983, vol. 153, 163- [\[0126\]](#)
- **HINNEN et al.**Proceedings of the National Academy of Sciences USA, 1978, vol. 75, 1920- [\[0126\]](#)
- **Protein Purification**VCH Publishers19890000 [\[0133\]](#)
- **TAGUE et al.**Plant Physiology, 1988, vol. 86, 506- [\[0141\]](#)
- **FRANCK et al.**Cell, 1980, vol. 21, 285-294 [\[0142\]](#)
- **CHRISTENSEN et al.**Plant Mol. Biol., 1992, vol. 18, 675-689 [\[0142\]](#)
- **ZHANG et al.**Plant Cell, 1991, vol. 3, 1155-1165 [\[0142\]](#)
- **EDWARDSCORUZZI**Ann. Rev. Genet., 1990, vol. 24, 275-303 [\[0142\]](#)
- **ITO et al.**Plant Mol. Biol., 1994, vol. 24, 863-878 [\[0142\]](#)
- **WU et al.**Plant and Cell Physiology, 1998, vol. 39, 885-889 [\[0142\]](#)
- **CONRAD et al.**Journal of Plant Physiology, 1998, vol. 152, 708-711 [\[0142\]](#)
- **CHEN et al.**Plant and Cell Physiology, 1998, vol. 39, 935-941 [\[0142\]](#)
- **KYOZUKA et al.**Plant Physiology, 1993, vol. 102, 991-1000 [\[0142\]](#)
- **MITRAHIGGINS**Plant Molecular Biology, 1994, vol. 26, 85-93 [\[0142\]](#)
- **KAGAYA et al.**Molecular and General Genetics, 1995, vol. 248, 668-674 [\[0142\]](#)
- **XU et al.**Plant Molecular Biology, 1993, vol. 22, 573-588 [\[0142\]](#)
- **GASSER et al.**Science, 1990, vol. 244, 1293- [\[0145\]](#)
- **POTRYKUS**Bio/Technology, 1990, vol. 8, 535- [\[0145\]](#)
- **SHIMAMOTO et al.**Nature, 1989, vol. 338, 274- [\[0145\]](#)
- **HOYKASSCHILPEROORT**Plant Molecular Biology, 1992, vol. 19, 15-38 [\[0146\]](#)
- **CHRISTOU**Plant Journal, 1992, vol. 2, 275-281 [\[0146\]](#)
- **SHIMAMOTO**Current Opinion Biotechnology, 1994, vol. 5, 158-162 [\[0146\]](#)
- **VASIL et al.**Bio/Technology, 1992, vol. 10, 667-674 [\[0146\]](#)
- **OMIRULLEH et al.**Plant Molecular Biology, 1993, vol. 21, 415-428 [\[0146\]](#)
- **GORINSTEIN**LIStarch, 1992, vol. 44, 12461-466 [\[0171\]](#)
- **KANEKO et al.**J. Ferment. Bioeng., 1996, vol. 81, 292-298 [\[0196\]](#)
- **SAMBROOK et al.**Molecular cloning: A laboratory manualCold Spring Harbor lab.19890000 [\[0214\]](#)
- **Current protocols in Molecular Biology**John Wiley and Sons19950000 [\[0214\]](#)
- **Molecular Biological Methods for Bacillus**John Wiley and Sons19900000 [\[0214\]](#)
- **Starch-Glucoamylase Method with Subsequent Measurement of Glucose with Glucose Oxidase**Approved methods of the American Association of Cereal ChemistsAmerican Association of Cereal Chemists20000000vol. 1-2, [\[0216\]](#)

Patentkrav

1. Isoleret polypeptid med glucoamylaseaktivitet, der er udvalgt fra gruppen, som består af:

5 (a) et polypeptid, som omfatter en aminosyresekvens, der har mindst 90%, mere fortrinsvis mindst 91%, mere fortrinsvis mindst 92%, endnu mere fortrinsvis mindst 93%, mest fortrinsvis mindst 94% og allermost fortrinsvis mindst 95%, såsom mindst 96%, mindst 97%, mindst 98%, mindst 99% eller endda 100% identitet med aminosyrerne 19 til 573 ifølge SEQ ID NO: 2, SEQ ID NO: 4 eller SEQ ID NO: 6;

10 (b) et polypeptid, som omfatter en aminosyresekvens, der har mindst 90%, mere fortrinsvis mindst 91%, mere fortrinsvis mindst 92%, endnu mere fortrinsvis mindst 93%, mest fortrinsvis mindst 94% og allermost fortrinsvis mindst 95%, såsom mindst 96%, mindst 97%, mindst 98%, mindst 99% eller endda 100% identitet med det katalytiske domæne, som er vist som aminosyrerne 22 til 476 ifølge SEQ ID NO: 2, SEQ ID NO: 4 eller SEQ ID NO: 6;

15 (c) et polypeptid, der kodes for af et polynukleotid omfattende en nukleotidsekvens, som har mindst 90%, mere fortrinsvis mindst 91%, mere fortrinsvis mindst 92%, endnu mere fortrinsvis mindst 93%, mest fortrinsvis mindst 94% og allermost fortrinsvis mindst 95%, såsom endda mindst 96%, 97%, 98%, 99% eller 100% identitet med den kodende sekvens ifølge SEQ ID NO: 1, SEQ ID NO: 3 eller SEQ ID NO: 5, der koder for aminosyrerne 19
20 til 573 ifølge SEQ ID NO: 2, SEQ ID NO: 4 eller SEQ ID NO: 6.

2. Polypeptid ifølge krav 1, som omfatter eller består af aminosyrerne 19 til 573 ifølge SEQ ID NO: 2, SEQ ID NO: 4 eller SEQ ID NO: 6; eller et fragment deraf med glucoamylaseaktivitet.

25 3. Polypeptid ifølge krav 1 eller 2, der kodes for af et polynukleotid, som omfatter eller består af nukleotidsekvensen ifølge SEQ ID NO: 1, SEQ ID NO: 3 eller SEQ ID NO: 5; eller en delsekvens deraf, der koder for et fragment med glucoamylaseaktivitet.

30 4. Polypeptid ifølge et hvilket som helst af kravene 1-3, der kodes for af polynukleotidet, som indgår i plasmid indeholdt i *E. coli* DSM 23221.

5. Isoleret polynukleotid omfattende en nukleotidsekvens, der koder for polypeptidet ifølge et hvilket som helst af kravene 1-4.

6. Nukleinsyrekonstruktion, som omfatter polynukleotidet ifølge krav 9 operativt forbundet til en eller flere (adskillige) kontrolsekvenser, der styrer fremstillingen af polypeptidet i en ekspressionsvært.
- 5 7. Rekombinant ekspressionsvektor, som omfatter nukleinsyrekonstruktionen ifølge krav 6.
8. Rekombinant værtselle, som omfatter nukleinsyrekonstruktionen ifølge krav 6.
- 10 9. Fremgangsmåde til fremstilling af polypeptidet ifølge et hvilket som helst af kravene 1-4, hvilken fremgangsmåde omfatter: (a) dyrkning af en værtselle, som omfatter en nukleinsyrekonstruktion, hvilken nukleinsyrekonstruktion omfatter en nukleotidsekvens, der koder for polypeptidet, under betingelser, som er befordrende for fremstilling af polypeptidet; og (b) indvinding af polypeptidet.
- 15 10. Fremgangsmåde til fremstilling af polypeptidet ifølge et hvilket som helst af kravene 1-4, hvilken fremgangsmåde omfatter: (a) dyrkning af en transgen plante eller en plantecelle, som omfatter et polynukleotid, hvilket polynukleotid koder for polypeptidet, under betingelser, som er befordrende for fremstilling af polypeptidet; og (b) indvinding af polypeptidet.
- 20 11. Transgen plante, plantedel eller plantecelle omfattende et polynukleotid, hvilket polynukleotid koder for polypeptidet ifølge et hvilket som helst af kravene 1-4.
- 25 12. Anvendelse af et polypeptid ifølge et hvilket som helst af kravene 1-4 til fremstilling af sirup og/eller et fermenteringsprodukt.
13. Anvendelse ifølge krav 12, hvor udgangsmaterialet er gelatineret eller ugelatineret stivelsesholdigt materiale.
- 30 14. Anvendelse af et polypeptid ifølge et hvilket som helst af kravene 1-4 til brygning.
15. Sammensætning, som omfatter en alfa-amylase og et polypeptid ifølge et hvilket som helst af kravene 1-4.

16. Fremgangsmåde til fremstilling af et fermenteringsprodukt fra stivelsesholdigt materiale, hvilken fremgangsmåde omfatter følgende trin:

- (a) at gøre stivelsesholdigt materiale flydende;
- (b) at forsukre materialet, som er gjort flydende; og
- 5 (c) at fermentere med en fermenterende organisme;

hvor trin (b) udføres ved anvendelse af mindst en glucoamylase ifølge et hvilket som helst af kravene 1-4.

17. Fremgangsmåde til fremstilling af et fermenteringsprodukt fra stivelsesholdigt materiale, hvilken fremgangsmåde omfatter følgende trin:

- (a) at forsukre stivelsesholdigt materiale ved en temperatur, som er lavere end den laveste gelatineringstemperatur for det stivelsesholdige materiale; og
- (b) at fermentere med en fermenterende organisme,

hvor trin (a) udføres ved anvendelse af mindst en glucoamylase ifølge et hvilket som helst af kravene 1-4.