

(19)



(11) Publication number:

SG 181589 A1

(43) Publication date:

30.07.2012

(51) Int. Cl:

C07K 16/28, A61P 35/00, A61P 37/00;

(12)

Patent Application

(21) Application number: **2012042313**

(71) Applicant:

**F. HOFFMANN-LA ROCHE AG
GRENZACHER STRASSE 124, CH-4070
BASEL CH**

(22) Date of filing: **07.12.2010**

(30) Priority: **EP 09015310.7 10.12.2009**

(72) Inventor:

**DIMOUDIS, NIKOLAOS
LABERSTRASSE 9 82407 WIELENBACH
DE
FERTIG, GEORG ST.-KLARA-STRASSE
2 82377 PENZBERG DE
FIDLER, ALEXANDER
BARBARASTRASSE 10 82377
PENZBERG DE
KALUZA, KLAUS TANKENRAIN 3,
D-82362 WEILHEIM DE
PICKL, MARLENE
SAALANGERSTRASSE 5, D-82377
PENZBERG DE
RIES, CAROLA ST.-KLARA-STRASSE 9
82377 PENZBERG DE
SEEBER, STEFAN BREUNETSRIEDER
WEG 22 82377 PENZBERG DE
LANZENDOERFER, Martin -**

(54) Title:

**ANTIBODIES BINDING PREFERENTIALLY HUMAN CSF1R
EXTRACELLULAR DOMAIN 4 AND THEIR USE**

(57) Abstract:

The present invention relates to antibodies against human CSF-1R (anti-CSF-1R antibody), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 June 2011 (16.06.2011)

PCT

(10) International Publication Number
WO 2011/070024 A1

(51) International Patent Classification:
C07K 16/28 (2006.01) *A61P 37/00* (2006.01)
A61P 35/00 (2006.01)

(74) Agents: **BURGER, Alexander** et al.; Roche Diagnostics GmbH, Patent Department (LPP...6164), P.O. Box 11 52, 82372 Penzberg (DE).

(21) International Application Number:
PCT/EP2010/069090

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
7 December 2010 (07.12.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09015310.7 10 December 2009 (10.12.2009) EP
10173407.7 19 August 2010 (19.08.2010) EP

(71) Applicant (for all designated States except US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacher Strasse 124, CH-4070 Basel (CH).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventor: **LANZENDOERFER, Martin** (deceased).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DIMOUDIS, Nikolaos** [GR/DE]; Laberstrasse 9, 82407 Wielenbach (DE). **FERTIG, Georg** [DE/DE]; St.-Klara-Strasse 2, 82377 Penzberg (DE). **FIDLER, Alexander** [AT/DE]; Barbarastrasse 10, 82377 Penzberg (DE). **KALUZA, Klaus** [DE/DE]; Hochfeldanger 3, 83670 Bad Heilbrunn (DE). **PICKL, Marlene** [DE/DE]; Birkenstrasse 25, 82377 Penzberg (DE). **RIES, Carola** [DE/DE]; St.-Klara-Strasse 9, 82377 Penzberg (DE). **SEEBER, Stefan** [DE/DE]; Breunetsrieder Weg 22, 82377 Penzberg (DE).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))



WO 2011/070024 A1

(54) Title: ANTIBODIES BINDING PREFERENTIALLY HUMAN CSF1R EXTRACELLULAR DOMAIN 4 AND THEIR USE

(57) Abstract: The present invention relates to antibodies against human CSF-1R (anti-CSF-1R antibody), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

ANTIBODIES BINDING PREFERENTIALLY HUMAN CSF1R EXTRACELLULAR DOMAIN 4 AND THEIR USE

The present invention relates to antibodies against human CSF-1R (anti-CSF-1R antibodies), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

Background of the Invention

5 The human CSF-1 receptor (CSF-1R; colony stimulating factor 1 receptor; synonyms: M-CSF receptor; Macrophage colony-stimulating factor 1 receptor, Fms proto-oncogene, c-fms, SEQ ID NO: 62) is known since 1986 (Coussens, L., et al., Nature 320 (1986) 277-280). CSF-1R is a growth factor and encoded by the c-fms proto-oncogene (reviewed e.g. in Roth, P., and Stanley, E.R., Curr. Top. Microbiol.
10 Immunol. 181 (1992) 141-67).

CSF-1R is the receptor for CSF-1 (colony stimulating factor 1, also called M-CSF, macrophage colony-stimulating factor) and mediates the biological effects of this cytokine (Sherr, C.J., et al., Cell 41 (1985) 665-676). The cloning of the colony stimulating factor-1 receptor (CSF-1R) (also called c-fms) was described for the
15 first time in Roussel, M.F., et al., Nature 325 (1987) 549-552. In that publication, it was shown that CSF-1R had transforming potential dependent on changes in the C-terminal tail of the protein including the loss of the inhibitory tyrosine 969 phosphorylation which binds Cbl and thereby regulates receptor down regulation (Lee, P.S., et al., Embo J. 18 (1999) 3616-3628). Recently a second ligand for
20 CSF-1R termed interleukin-34 (IL-34) was identified (Lin, H., et al, Science 320 (2008) 807-811).

The cytokine CSF-1 (colony stimulating factor 1, also called M-CSF, macrophage) is found extracellularly as a disulfide-linked homodimer (Stanley, E.R. et al., Journal of Cellular Biochemistry 21 (1983) 151-159; Stanley, E.R. et al., Stem
25 Cells 12 Suppl. 1 (1995) 15-24).

The main biological effects of CSF-1R signaling are the differentiation, proliferation, migration, and survival of hematopoietic precursor cells to the macrophage lineage (including osteoclast). Activation of CSF-1R is mediated by its ligands, CSF-1 (M-CSF) and IL-34. Binding of CSF-1 (M-CSF) to CSF-1R
30 induces the formation of homodimers and activation of the kinase by tyrosine

phosphorylation (Li, W. et al, EMBO Journal.10 (1991) 277-288; Stanley, E.R., et al., Mol. Reprod. Dev. 46 (1997) 4-10).

5 The biologically active homodimer CSF-1 binds to the CSF-1R within the subdomains D1 to D3 of the extracellular domain of the CSF-1 receptor (CSF-1R-ECD). The CSF-1R-ECD comprises five immunoglobulin-like subdomains (designated D1 to D5). The subdomains D4 to D5 of the extracellular domain (CSF-1R-ECD) are not involved in the CSF-1 binding. (Wang, Z., et al Molecular and Cellular Biology 13 (1993) 5348-5359). The subdomain D4 is involved in dimerization (Yeung, Y-G., et al Molecular & Cellular Proteomics 2 (2003) 1143-1155; Pixley, F. J., et al., Trends Cell Biol 14 (2004) 628-638).

10 Further signaling is mediated by the p85 subunit of PI3K and Grb2 connecting to the PI3K/AKT and Ras/MAPK pathways, respectively. These two important signaling pathways can regulate proliferation, survival and apoptosis. Other signaling molecules that bind the phosphorylated intracellular domain of CSF-1R include STAT1, STAT3, PLC γ , and Cbl (Bourette, R.P. and Rohrschneider, L.R., Growth Factors 17 (2000) 155-166).

15 CSF-1R signaling has a physiological role in immune responses, in bone remodeling and in the reproductive system. The knockout animals for either CSF-1 (Pollard, J.W., Mol. Reprod. Dev. 46 (1997) 54-61) or CSF-1R (Dai, X.M., et al., Blood 99 (2002) 111-120) have been shown to have osteopetrotic, hematopoietic, tissue macrophage, and reproductive phenotypes consistent with a role for CSF-1R in the respective cell types.

20 Sherr, C.J., et al., Blood 73 (1989) 1786-1793 relates to some antibodies against CSF-1R that inhibit the CSF-1 activity (see Sherr, C.J. et al., Blood 73 (1989) 1786-1793). Ashmun, R.A., et al., Blood 73 (1989) 827-837 relates to CSF-1R antibodies. Lenda, D., et al., Journal of Immunology 170 (2003) 3254-3262 relates to reduced macrophage recruitment, proliferation, and activation in CSF-1-deficient mice results in decreased tubular apoptosis during renal inflammation. Kitaura, H., et al., Journal of Dental Research 87 (2008) 396-400 refers to an anti-CSF-1 antibody which inhibits orthodontic tooth movement. WO 2001/030381 mentions CSF-1 activity inhibitors including antisense nucleotides and antibodies while disclosing only CSF-1 antisense nucleotides. WO 2004/045532 relates to metastases and bone loss prevention and treatment of metastatic cancer by a CSF-1

antagonist disclosing as antagonist anti-CSF-1-antibodies only. WO 2005/046657 relates to the treatment of inflammatory bowel disease by anti-CSF-1-antibodies. US 2002/0141994 relates to inhibitors of colony stimulating factors. WO 2006/096489 relates to the treatment of rheumatoid arthritis by anti-CSF-1-antibodies. WO 2009/026303 and WO 2009/112245 relate to certain anti-CSF-1R antibodies binding to CSF-1R within the first three subdomains (D1 to D3) of the Extracellular Domain (CSF-1R-ECD).

Summary of the Invention

The invention comprises an antibody binding to human CSF-1R, characterized in that the antibody binds to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (SEQ ID NO: 64) with a ratio of 1:50 or lower.

The invention further comprises an antibody according to the invention characterized in that

- a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
- b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
- c) the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;
- d) the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;

or a humanized version thereof.

The invention further comprises an antibody according to the invention characterized in that

- a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
- b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;

or a humanized version thereof.

In one embodiment the antibody according to the invention is characterized in that

- a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
- 5 b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
- c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
- d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48, or
- 10 e) the heavy chain variable domain is SEQ ID NO:55 and the light chain variable domain is SEQ ID NO:56.

The invention further comprises an antibody according to the invention, characterized in that

- 15 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or
- 20 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or
- 25 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or
- 30 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of

SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

5 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

10 f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or

15 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54; or

20 h) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or

25 i) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.

Preferably the antibody according to the invention is of human IgG1 subclass or of human IgG4 subclass.

30 A further embodiment of the invention is a pharmaceutical composition comprising an antibody according to the invention.

The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of a CSF-1R mediated disease.

5 The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of cancer.

The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of bone loss.

The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of metastasis.

10 The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of inflammatory diseases.

The invention further comprises an antibody according to the invention for treatment of a CSF-1R mediated disease.

15 The invention further comprises an antibody according to the invention for treatment of cancer.

The invention further comprises an antibody according to the invention for treatment of bone loss.

20 The invention further comprises an antibody according to the invention for treatment of metastasis.

The invention further comprises an antibody according to the invention for treatment of inflammatory diseases.

A further embodiment of the invention is a nucleic acid encoding an antibody according to the invention characterized in that

25 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID

NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or,

5 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or

10 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

15 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

20 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

25 f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or

30 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of

SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54, or

5 h) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or

10 i) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.

A further embodiment of the invention is a nucleic acid encoding an antibody according to the invention characterized in that

15 a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
c) the heavy chain variable domain is SEQ ID NO:75 and the light chain
20 variable domain is SEQ ID NO:76;
d) the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;

or a humanized version thereof.

25 A further embodiment of the invention is a nucleic acid encoding an antibody according to the invention characterized in that

a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
30 c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or

d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48, or

e) the heavy chain variable domain is SEQ ID NO:55 and the light chain variable domain is SEQ ID NO:56.

5 The invention further provides expression vectors containing nucleic acid according to the invention capable of expressing said nucleic acid in a prokaryotic or eukaryotic host cell, and host cells containing such vectors for the recombinant production of an antibody according to the invention.

10 The invention further comprises a prokaryotic or eukaryotic host cell comprising a vector according to the invention.

The invention further comprises a method for the production of a recombinant human or humanized antibody according to the invention, characterized by expressing a nucleic acid according to the invention in a prokaryotic or eukaryotic host cell and recovering said antibody from said cell or the cell culture supernatant.

15 The invention further comprises the antibody obtained by such a recombinant method.

20 Antibodies according to the invention show benefits for patients in need of a CSF-1R targeting therapy. The antibodies according to the invention show efficient antiproliferative activity against ligand-independent and ligand-dependant proliferation and are therefore especially useful in the treatment of cancer and metastasis.

25 The invention further provides a method for treating a patient suffering from cancer, comprising administering to a patient diagnosed as having such a disease (and therefore being in need of such a therapy) an effective amount of an antibody according to the invention. The antibody is administered preferably in a pharmaceutical composition.

A further embodiment of the invention is a method for the treatment of a patient suffering from cancer characterized by administering to the patient an antibody according to the invention.

30 Surprisingly it has been found that, using a human CSF-1R fragment delD4 in which the D4 subdomain of human CSF-1R-ECD was deleted (SEQ ID NO:65),

the new anti-CSF-1R antibodies according to the invention could be selected. These antibodies show valuable properties like excellent ligand-dependant cell growth inhibition and at the same time ligand independent cell growth inhibition of NIH 3T3 cell, retrovirally infected with either an expression vector for full-length wildtype CSF-1R (SEQ ID NO:62) or mutant CSF-1R L301S Y969F (SEQ ID NO:63) whereby mutant CSF-1R recombinant cells are able to form spheroids independent of the CSF-1 ligand. Furthermore the antibodies according to the invention inhibit (both) human and cynomolgous macrophage differentiation, as they inhibit survival of human and cynomolgous monocytes.

10 **Detailed Description of the Invention**

The invention comprises an antibody binding to human CSF-1R, characterized in that the antibody binds to human CSF-1R fragment delD4 (comprising the extracellular subdomains D1 –D3 and D5) (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 –D5) (SEQ ID NO: 64) with a ratio of 1:50 or lower.

The invention further comprises an antibody according to the invention characterized in comprising as heavy chain variable domain CDR3 region a CDR3 region of SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO:23, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47 or SEQ ID NO:55.

20 The invention further comprises an antibody according to the invention characterized in that

- a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
 - b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
- 25 or a humanized version thereof.

The invention further comprises an antibody according to the invention characterized in that

- a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
- 30

- b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
- c) the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;
- 5 d) the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;

or a humanized version thereof.

The invention further comprises an antibody according to the invention characterized in that

- 10 the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8, or a humanized version thereof.

In one embodiment the antibody according to the invention is characterized in that

- a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
- 15 b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
- c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
- d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48, or
- 20 e) the heavy chain variable domain is SEQ ID NO:55 and the light chain variable domain is SEQ ID NO:56.

In one embodiment the antibody according to the invention is characterized in that

- a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
- 25 b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
- c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
- 30 d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48.

In one embodiment the antibody according to the invention is characterized in that

the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or

In one embodiment the antibody according to the invention is characterized in that

5 the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32.

In one embodiment the antibody according to the invention is characterized in that

the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40.

10 In one embodiment the antibody according to the invention is characterized in that

the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48.

The invention further comprises an antibody according to the invention characterized in that

15 the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16, or a humanized version thereof.

The invention further comprises an antibody according to the invention characterized in that

20 the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;

or a humanized version thereof.

The invention further comprises an antibody according to the invention characterized in that

25 the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;

or a humanized version thereof.

The invention further comprises an antibody according to the invention, characterized in that

5 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or,

10 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or

15 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

20 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

25 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

30 f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of

SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or

5 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54.

The invention further comprises an antibody according to the invention, characterized in that

10 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or,

15 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or

20 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

25 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

30 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID

NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

5 f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46,

10 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54;

15 h) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or

20 i) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.

In one embodiment the antibody according to the invention is characterized in that

25 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or

30 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID

NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.

In one embodiment the antibody according to the invention is characterized in that

- 5 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or
- 10 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or
- 15 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or
- 20 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or
- 25 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54.

30 In one embodiment the antibody according to the invention is characterized in that

5 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

10 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

15 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

20 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46.

In one embodiment the antibody according to the invention is characterized in that

25 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22.

In one embodiment the antibody according to the invention is characterized in that

30 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of

SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30.

In one embodiment the antibody according to the invention is characterized in that

5 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38.

In one embodiment the antibody according to the invention is characterized in that

10 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46.

15 In one embodiment the antibody binding to human CSF-1R, characterized in that the antibody binds to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R-ECD (SEQ ID NO: 64) with a ratio of 1:50 or lower, is further characterized in not binding to human CSF-1R fragment D1-D3 (SEQ ID NO: 66).

20 The term "antibody" encompasses the various forms of antibodies including but not being limited to whole antibodies, antibody fragments, human antibodies, humanized antibodies, chimeric antibodies, T cell epitope depleted antibodies, and further genetically engineered antibodies as long as the characteristic properties according to the invention are retained. "Antibody fragments" comprise a portion of a full length antibody, preferably the variable domain thereof, or at least the
25 antigen binding site thereof. Examples of antibody fragments include diabodies, single-chain antibody molecules, and multispecific antibodies formed from antibody fragments. scFv antibodies are, e.g., described in Houston, J.S., Methods in Enzymol. 203 (1991) 46-88). In addition, antibody fragments comprise single chain polypeptides having the characteristics of a V_H domain binding to CSF-1R,
30 namely being able to assemble together with a V_L domain, or of a V_L domain

binding to CSF-1R, namely being able to assemble together with a V_H domain to a functional antigen binding site and thereby providing the property.

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of a single amino acid composition.

5

The term "chimeric antibody" refers to a monoclonal antibody comprising a variable region, i.e., binding region, from mouse and at least a portion of a constant region derived from a different source or species, usually prepared by recombinant DNA techniques. Chimeric antibodies comprising a mouse variable region and a human constant region are especially preferred. Such rat/human chimeric antibodies are the product of expressed immunoglobulin genes comprising DNA segments encoding rat immunoglobulin variable regions and DNA segments encoding human immunoglobulin constant regions. Other forms of "chimeric antibodies" encompassed by the present invention are those in which the class or subclass has been modified or changed from that of the original antibody. Such "chimeric" antibodies are also referred to as "class-switched antibodies." Methods for producing chimeric antibodies involve conventional recombinant DNA and gene transfection techniques now well known in the art. See, e.g., Morrison, S.L., et al., Proc. Natl. Acad. Sci. USA 81 (1984) 6851-6855; US 5,202,238 and US 5,204,244.

10

15

20

The term "humanized antibody" refers to antibodies in which the framework or "complementarity determining regions" (CDR) have been modified to comprise the CDR of an immunoglobulin of different specificity as compared to that of the parent immunoglobulin. In a preferred embodiment, a murine CDR is grafted into the framework region of a human antibody to prepare the "humanized antibody." See e.g. Riechmann, L., et al., Nature 332 (1988) 323-327; and Neuberger, M.S., et al., Nature 314 (1985) 268-270. Optionally the framework region can be modified by further mutations. Also the CDRs can be modified by one or more mutations to generate antibodies according to the invention e.g. by mutagenesis based upon molecular modeling as described by Riechmann, L., et al., Nature 332 (1988) 323-327 and Queen, C., et al., Proc. Natl. Acad. Sci. USA 86 (1989) 10029-10033, or others. Particularly preferred CDRs correspond to those representing sequences recognizing the antigens noted above for chimeric antibodies. A "humanized

25

30

version of an antibody according to the invention” (which is e.g. of mouse origin) refers to an antibody, which is based on the mouse antibody sequences in which the V_H and V_L are humanized by standard techniques (including CDR grafting and optionally subsequent mutagenesis of certain amino acids in the framework region and the CDRs). Preferably such humanized version is chimerized with a human constant region (see e.g. Sequences SEQ ID NO:57-61).

Other forms of "humanized antibodies" encompassed by the present invention are those in which the constant region has been additionally modified or changed from that of the original antibody to generate the properties according to the invention, especially in regard to C1q binding and/or Fc receptor (FcR) binding.

In the following examples the terms “Mab” or “muMab” refer to murine monoclonal antibodies such as Mab 2F11 or Mab 2E10, whereas the term “hMab” refers to humanized monoclonal versions of such murine antibodies such as hMab 2F11-c11, hMab 2F11-d8, hMab 2F11-e7, hMab 2F11-f12, etc..

The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germ line immunoglobulin sequences. Human antibodies are well-known in the state of the art (van Dijk, M.A., and van de Winkel, J.G., *Curr. Opin. Chem. Biol.* 5 (2001) 368-374). Human antibodies can also be produced in transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire or a selection of human antibodies in the absence of endogenous immunoglobulin production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge (see, e.g., Jakobovits, A., et al., *Proc. Natl. Acad. Sci. USA* 90 (1993) 2551-2555; Jakobovits, A., et al., *Nature* 362 (1993) 255-258; Brueggemann, M., et al., *Year Immunol.* 7 (1993) 33-40). Human antibodies can also be produced in phage display libraries (Hoogenboom, H.R., and Winter, G.J. *Mol. Biol.* 227 (1992) 381-388; Marks, J.D., et al., *J. Mol. Biol.* 222 (1991) 581-597). The techniques of Cole, et al., and Boerner, et al., are also available for the preparation of human monoclonal antibodies (Cole, S.P.C., et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); and Boerner, P., et al., *J. Immunol.* 147 (1991) 86-95). As already mentioned for chimeric and humanized antibodies according to the invention the term “human antibody” as used herein

also comprises such antibodies which are modified in the constant region to generate the properties according to the invention, especially in regard to C1q binding and/or FcR binding, e.g. by "class switching" i.e. change or mutation of Fc parts (e.g. from IgG1 to IgG4 and/or IgG1/IgG4 mutation).

5 The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from a host cell such as a NS0 or CHO cell or from an animal (e.g. a mouse) that is transgenic for human immunoglobulin genes or antibodies expressed using a recombinant expression vector transfected into a
10 host cell. Such recombinant human antibodies have variable and constant regions in a rearranged form. The recombinant human antibodies according to the invention have been subjected to *in vivo* somatic hypermutation. Thus, the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germ line VH and VL sequences,
15 may not naturally exist within the human antibody germ line repertoire *in vivo*.

The antibodies according to the invention include, in addition, such antibodies having "conservative sequence modifications", nucleotide and amino acid sequence modifications which do not affect or alter the above-mentioned characteristics of the antibody according to the invention. Modifications can be introduced by
20 standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions include ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains
25 (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g. glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine,
30 phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a human anti-CSF-1R antibody can be preferably replaced with another amino acid residue from the same side chain family.

Amino acid substitutions can be performed by mutagenesis based upon molecular modeling as described by Riechmann, L., et al., *Nature* 332 (1988) 323-327 and Queen, C., et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 10029-10033.

5 The human CSF-1R (CSF-1 receptor; synonyms: M-CSF receptor; Macrophage colony-stimulating factor 1 receptor, *Fms* proto-oncogene, *c-fms*, SEQ ID NO: 22) is known since 1986 (Coussens, L., et al., *Nature* 320 (1986) 277-280). CSF-1R is a growth factor and encoded by the *c-fms* proto-oncogene (reviewed e.g. in Roth, P. and Stanley, E.R., *Curr. Top. Microbiol. Immunol.* 181 (1992) 141-67).

10 CSF-1R is the receptor for CSF-1 (macrophage colony stimulating factor, also called M-CSF) and IL-34 and mediates the biological effects of these cytokines (Sherr, C.J., et al., *Cell* 41 (1985) 665-676 (Lin, H., et al., *Science* 320 (2008) 807-811). The cloning of the colony stimulating factor-1 receptor (also called *c-fms*) was described for the first time in Roussel, M.F., et al., *Nature* 325 (1987) 549-552. In that publication, it was shown that CSF-1R had transforming potential
15 dependent on changes in the C-terminal tail of the protein including the loss of the inhibitory tyrosine 969 phosphorylation which binds *Cbl* and thereby regulates receptor down regulation (Lee, P.S., et al., *Embo J.* 18 (1999) 3616-3628).

20 CSF-1R is a single chain, transmembrane receptor tyrosine kinase (RTK) and a member of the family of immunoglobulin (Ig) motif containing RTKs characterized by 5 repeated Ig-like subdomains D1-D5 in the extracellular domain (ECD) of the receptor (Wang, Z., et al *Molecular and Cellular Biology* 13 (1993) 5348-5359). The human CSF-1R Extracellular Domain (CSF-1R-ECD) (SEQ ID NO: 64) comprises all five extracellular Ig-like subdomains D1 –D5. The human CSF-1R fragment delD4 (SEQ ID NO: 65) comprises the extracellular
25 Ig-like subdomains D1–D3 and D5, but is missing the D4 subdomain. The human CSF-1R fragment D1-D3 (SEQ ID NO: 66) comprises the respective subdomains D1-D3. The sequences are listed without the signal peptide MGS~~G~~PGVLLL LLVATAWHGQ G (SEQ ID NO: 67).

30 The intracellular protein tyrosine kinase domain is interrupted by a unique insert domain that is also present in the other related RTK class III family members that include the platelet derived growth factor receptors (PDGFR), stem cell growth factor receptor (*c-Kit*) and fins-like cytokine receptor (FLT3). In spite of the

structural homology among this family of growth factor receptors, they have distinct tissue-specific functions.

CSF-1R is mainly expressed on cells of the monocytic lineage and in the female reproductive tract and placenta. In addition expression of CSF-1R has been reported in Langerhans cells in skin, a subset of smooth muscle cells (Inaba, T., et al., J. Biol. Chem. 267 (1992) 5693-5699), B cells (Baker, A.H., et al., Oncogene 8 (1993) 371-378) and microglia (Sawada, M., et al., Brain Res. 509 (1990) 119-124). Cells with mutant human CSF-1R ((SEQ ID NO: 23) are known to proliferate independently of ligand stimulation.

As used herein, "binding to human CSF-1R" or "specifically binding to human CSF-1R" refers to an antibody specifically binding to the human CSF-1R antigen with a binding affinity of KD-value of 1.0×10^{-8} mol/l or lower at 35°C, in one embodiment of a KD-value of 1.0×10^{-9} mol/l or lower at 35°C. The binding affinity is determined with a standard binding assay at 35°C, such as surface plasmon resonance technique (BIAcore®, GE-Healthcare Uppsala, Sweden) A method for determining the KD-value of the binding affinity is described in Example 9. Thus an "antibody binding to human CSF-1R" as used herein refers to an antibody specifically binding to the human CSF-1R antigen with a binding affinity of KD 1.0×10^{-8} mol/l or lower (preferably 1.0×10^{-8} mol/l - 1.0×10^{-12} mol/l) at 35°C, preferably of a KD 1.0×10^{-9} mol/l or lower at 35°C (preferably 1.0×10^{-9} mol/l - 1.0×10^{-12} mol/l).

The "binding to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (SEQ ID NO: 64)" as used herein is measured by a Surface Plasmon Resonance assay (Biacore assay) as described in Example 4. The human CSF-1R fragment delD4 (SEQ ID NO: 65) or human CSF-1R Extracellular Domain (SEQ ID NO: 64), respectively, are captured to the surface (each to a separate surface) and the test antibodies were added (each in a separate measurement) and the respective binding signals (Response Units (RU)) were determined. Reference signals (blank surface) were subtracted. If signals of nonbinding test antibodies were slightly below 0 the values were set as 0. Then the ratio of the respective binding signals (binding signal (RU) to human CSF-1R fragment delD4 /binding signal (RU) to human CSF-1R Extracellular Domain (CSF-1R-ECD)) is determined. The antibodies according to the invention have a

ratio of the binding signals (RU(delD4) / RU(CSF-1R-ECD) of 1:50 or lower, preferably of 1:100 or lower (the lower included end is 0 (e.g. if the RU is 0, then the ratio is 0:50 or 0:100)).

5 This means that such anti-CSF-1R antibodies according to the invention do not bind to the human CSF-1R fragment delD4 (like the anti-CCR5 antibody m<CCR5>Pz03.1C5 (deposited as DSM ACC 2683 on 18.08.2004 at DSMZ) and have binding signals for binding to the human CSF-1R fragment delD4 in the range of the anti-CCR5 antibody m<CCR5>Pz03.1C5, which are below 20 RU (Response Units), preferably below 10 RU in a Surface Plasmon Resonance
10 (BIAcore) assay as shown in Example 4.

The term “binding to human CSF-1R fragment D1-D3” refers to a binding affinity determination by a Surface Plasmon Resonance assay (Biacore assay). The test antibody is captured to the surface and the human CSF-1R fragment D1-D3 (SEQ ID NO: 66) was added and the respective binding affinities were determined. The
15 term “not binding to human CSF-1R fragment D1-D3” denotes that in such an assay the detected signal was in the area of no more than 1.2 fold of background signal and therefore no significant binding could be detected and no binding affinity could be determined (see Example 10).

One embodiment of the invention is a screening method for selecting antibodies
20 according to the invention comprising the following steps:

- a) determining the binding signal (Response Units (RU)) of anti-CSF-1R antibodies to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (CSF-1R-ECD) (SEQ ID NO: 64) by a Surface Plasmon Resonance assay (Biacore assay).
- 25 b) selecting antibodies with ratio of the binding signals (human CSF-1R fragment delD4/ human CSF-1R Extracellular Domain (CSF-1R-ECD)) of 50:1 or lower.

In one embodiment the determination is performed at 25°C.

In one embodiment the screening method comprises as further steps the measuring
30 of the binding of anti-CSF-1R antibodies to human CSF-1R fragment D1-D3 (SEQ

ID NO: 66) (D1-D3) and the selecting of antibodies which show no binding to said fragment.

5 The term “epitope” denotes a protein determinant of human CSF-1R capable of specifically binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually epitopes have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents. Preferably an antibody according to the invention
10 binds specifically to native and to denatured CSF-1R.

The “variable domain” (variable domain of a light chain (V_L), variable domain of a heavy chain (V_H)) as used herein denotes each of the pair of light and heavy chain domains which are involved directly in binding the antibody to the antigen. The variable light and heavy chain domains have the same general structure and each
15 domain comprises four framework (FR) regions whose sequences are widely conserved, connected by three “hypervariable regions” (or complementary determining regions, CDRs). The framework regions adopt a β -sheet conformation and the CDRs may form loops connecting the β -sheet structure. The CDRs in each chain are held in their three-dimensional structure by the framework regions and
20 form together with the CDRs from the other chain the antigen binding site. The antibody’s heavy and light chain CDR3 regions play a particularly important role in the binding specificity/affinity of the antibodies according to the invention and therefore provide a further object of the invention.

The term “antigen-binding portion of an antibody” when used herein refer to the
25 amino acid residues of an antibody which are responsible for antigen-binding. The antigen-binding portion of an antibody comprises amino acid residues from the “complementary determining regions” or “CDRs”. “Framework” or “FR” regions are those variable domain regions other than the hypervariable region residues as herein defined. Therefore, the light and heavy chain variable domains of an
30 antibody comprise from N- to C-terminus the domains FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4. Especially, CDR3 of the heavy chain is the region which contributes most to antigen binding and defines the antibody’s properties. CDR and FR regions are determined according to the standard definition of Kabat et al.,

Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991) and/or those residues from a “hypervariable loop”.

5 The terms “nucleic acid” or “nucleic acid molecule”, as used herein, are intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

10 The term “amino acid” as used within this application denotes the group of naturally occurring carboxy α -amino acids comprising alanine (three letter code: ala, one letter code: A), arginine (arg, R), asparagine (asn, N), aspartic acid (asp, D), cysteine (cys, C), glutamine (gln, Q), glutamic acid (glu, E), glycine (gly, G), histidine (his, H), isoleucine (ile, I), leucine (leu, L), lysine (lys, K), methionine (met, M), phenylalanine (phe, F), proline (pro, P), serine (ser, S), threonine (thr, T), tryptophan (trp, W), tyrosine (tyr, Y), and valine (val, V).

15 In one embodiment the antibodies according to the invention inhibit CSF-1 binding to CSF-1R. In one embodiment with an IC₅₀ of 200 ng/ml or lower, in one embodiment with an IC₅₀ of 50 ng/ml or lower. The IC₅₀ of inhibition of CSF-1 binding to CSF-1R can be determined as shown in Example 2.

In one embodiment the antibodies according to the invention inhibit CSF-1-induced CSF-1R phosphorylation (in NIH3T3-CSF-1R recombinant cells).

20 In one embodiment with an IC₅₀ of 800 ng/ml or lower, in one embodiment with an IC₅₀ of 600 ng/ml or lower, in one embodiment with an IC₅₀ of 250 ng/ml or lower. The IC₅₀ of CSF-1-induced CSF-1R phosphorylation can be determined as shown in Example 3.

25 In one embodiment the antibodies according to the invention inhibit the growth of recombinant NIH3T3 cells expressing human CSF-1R (SEQ ID No: 62). In one embodiment with an IC₅₀ of 10 μ g/ml or lower, in one embodiment with an IC₅₀ of 5 μ g/ml or lower, in one embodiment with an IC₅₀ of 2 μ g/ml or lower. In one embodiment with an IC₃₀ of 10 μ g/ml or lower, in one embodiment with an IC₃₀ of 5 μ g/ml or lower, in one embodiment with an IC₃₀ of 2 μ g/ml or lower. The
30 IC₅₀ value, the IC₃₀ value or the % growth inhibition is determined as shown in Example 5.

In one embodiment the antibodies according to the invention inhibit the growth of recombinant NIH3T3 cells expressing human mutant CSF-1R L301S Y969F (SEQ ID No: 63). In one embodiment with an IC50 of 15 µg/ml or lower, in one embodiment with an IC50 of 10 µg/ml or lower. In one embodiment with an IC30 of 10 µg/ml or lower, in one embodiment with an IC50 of 5 µg/ml or lower; in one embodiment with an IC50 of 2 µg/ml or lower. The IC50 value, the IC30 value or the % growth inhibition is determined as shown in Example 5.

In one embodiment the antibodies according to the invention inhibit the growth of BeWo tumor cells (ATCC CCL-98) by 65 % or more (at an antibody concentration of 10 µg/ml ; and as compared to the absence of antibody). The % growth inhibition is determined as shown in Example 8. E.g. Mab 2F11 shows a growth inhibition of BeWo tumor cells of 70 %.

In one embodiment the antibodies according to the invention inhibit (both) human and cynomolgous macrophage differentiation (which is indicated by the inhibition of the survival of human and cynomolgous monocytes as shown in Examples 7 and 8). In one embodiment the antibodies according to the invention inhibit the survival of human monocytes with an IC50 of 0.15 µg/ml or lower, in one embodiment with an IC50 of 0.10 µg/ml or lower. The inhibition of the survival of human monocytes is determined as shown in Example 7. In one embodiment the antibodies according to the invention inhibit the survival of cynomolgous monocytes by 80 % or more, in one embodiment by 90 % or more (at an antibody concentration of 5 µg/ml ; and as compared to the absence of antibody). The inhibition of the survival of human monocytes is determined as shown in Example 8.

A further embodiment of the invention is a method for the production of an antibody against CSF-1R characterized in that the sequence of a nucleic acid encoding the heavy chain of a human IgG1 class antibody binding to human CSF-1R according to the invention said modified nucleic acid and the nucleic acid encoding the light chain of said antibody are inserted into an expression vector, said vector is inserted in a eukaryotic host cell, the encoded protein is expressed and recovered from the host cell or the supernatant.

The antibodies according to the invention are preferably produced by recombinant means. Therefore the antibody is preferably an isolated monoclonal antibody. Such recombinant methods are widely known in the state of the art and comprise protein

expression in prokaryotic and eukaryotic cells with subsequent isolation of the antibody polypeptide and usually purification to a pharmaceutically acceptable purity. For the protein expression, nucleic acids encoding light and heavy chains or fragments thereof are inserted into expression vectors by standard methods.
5 Expression is performed in appropriate prokaryotic or eukaryotic host cells like CHO cells, NS0 cells, SP2/0 cells, HEK293 cells, COS cells, yeast, or E.coli cells, and the antibody is recovered from the cells (supernatant or cells after lysis).

Recombinant production of antibodies is well-known in the state of the art and described, for example, in the review articles of Makrides, S.C., *Protein Expr. Purif.* 17 (1999) 183-202; Geisse, S., et al., *Protein Expr. Purif.* 8 (1996) 271-282;
10 Kaufman, R.J., *Mol. Biotechnol.* 16 (2000) 151-161; Werner, R.G., *Drug Res.* 48 (1998) 870-880.

The antibodies may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. Purification is performed in order to eliminate
15 other cellular components or other contaminants, e.g. other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis, and others well known in the art. See Ausubel, F., et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York (1987).

Expression in NS0 cells is described by, e.g., Barnes, L.M., et al., *Cytotechnology* 32 (2000) 109-123; and Barnes, L.M., et al., *Biotech. Bioeng.* 73 (2001) 261-270. Transient expression is described by, e.g., Durocher, Y., et al., *Nucl. Acids. Res.* 30 (2002) E9. Cloning of variable domains is described by Orlandi, R., et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 3833-3837; Carter, P., et al., *Proc. Natl. Acad. Sci. USA* 89 (1992) 4285-4289; and Norderhaug, L., et al., *J. Immunol. Methods* 204 (1997) 77-87. A preferred transient expression system (HEK 293) is described by
25 Schlaeger, E.-J., and Christensen, K., in *Cytotechnology* 30 (1999) 71-83 and by Schlaeger, E.-J., in *J. Immunol. Methods* 194 (1996) 191-199.

The control sequences that are suitable for prokaryotes, for example, include a
30 promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, enhancers and polyadenylation signals.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The monoclonal antibodies are suitably separated from the culture medium by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography. DNA and RNA encoding the monoclonal antibodies are readily isolated and sequenced using conventional procedures. The hybridoma cells can serve as a source of such DNA and RNA. Once isolated, the DNA may be inserted into expression vectors, which are then transfected into host cells such as HEK 293 cells, CHO cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of recombinant monoclonal antibodies in the host cells.

As used herein, the expressions "cell", "cell line", and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Variant progeny that have the same function or biological activity as screened for in the originally transformed cell are included.

The "Fc part" of an antibody is not involved directly in binding of an antibody to an antigen, but exhibit various effector functions. A "Fc part of an antibody" is a term well known to the skilled artisan and defined on the basis of papain cleavage of antibodies. Depending on the amino acid sequence of the constant region of their

heavy chains, antibodies or immunoglobulins are divided in the classes: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG1, IgG2, IgG3, and IgG4, IgA1, and IgA2. According to the heavy chain constant regions the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The Fc part of an antibody is directly involved in ADCC (antibody-dependent cell-mediated cytotoxicity) and CDC (complement-dependent cytotoxicity) based on complement activation, C1q binding and Fc receptor binding. Complement activation (CDC) is initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. While the influence of an antibody on the complement system is dependent on certain conditions, binding to C1q is caused by defined binding sites in the Fc part. Such binding sites are known in the state of the art and described e.g. by Boackle, R.J., et al., Nature 282 (1979) 742-743, Lukas, T.J., et al., J. Immunol. 127 (1981) 2555-2560, Brunhouse, R., and Cebra, J.J., Mol. Immunol. 16 (1979) 907-917, Burton, D.R., et al., Nature 288 (1980) 338-344, Thommesen, J.E., et al., Mol. Immunol. 37 (2000) 995-1004, Idusogie, E.E., et al., J. Immunol. 164 (2000) 4178-4184, Hezareh, M., et al., J. Virology 75 (2001) 12161-12168, Morgan, A., et al., Immunology 86 (1995) 319-324, EP 0307434. Such binding sites are e.g. L234, L235, D270, N297, E318, K320, K322, P331 and P329 (numbering according to EU index of Kabat, E.A., see below). Antibodies of subclass IgG1, IgG2 and IgG3 usually show complement activation and C1q and C3 binding, whereas IgG4 do not activate the complement system and do not bind C1q and C3.

In one embodiment the antibody according to the invention comprises a Fc part derived from human origin and preferably all other parts of the human constant regions. As used herein the term "Fc part derived from human origin" denotes a Fc part which is either a Fc part of a human antibody of the subclass IgG1, IgG2, IgG3 or IgG4, preferably a Fc part from human IgG1 subclass, a mutated Fc part from human IgG1 subclass (preferably with a mutation on L234A + L235A), a Fc part from human IgG4 subclass or a mutated Fc part from human IgG4 subclass (preferably with a mutation on S228P). Mostly preferred are the human heavy chain constant regions of SEQ ID NO: 58 (human IgG1 subclass), SEQ ID NO: 59 (human IgG1 subclass with mutations L234A and L235A), SEQ ID NO: 60 (human IgG4 subclass), or SEQ ID NO: 61 (human IgG4 subclass with mutation S228P).

Preferably the antibody according to the invention is of human IgG1 subclass or of human IgG4 subclass. In one embodiment the antibody according to the invention is of human IgG1 subclass. In one embodiment the antibody according to the invention is of human IgG4 subclass.

5 In one embodiment the antibody according to the invention is characterized in that the constant chains are of human origin. Such constant chains are well known in the state of the art and e.g. described by Kabat, E.A., (see e.g. Johnson, G. and Wu, T.T., Nucleic Acids Res. 28 (2000) 214-218). For example, a useful human heavy chain constant region comprises an amino acid sequence of SEQ ID NO: 58. For
10 example, a useful human light chain constant region comprises an amino acid sequence of a kappa-light chain constant region of SEQ ID NO: 57.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

15 a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
or a humanized version thereof.

20 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
25 c) the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;
d) the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;

or a humanized version thereof.

30 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8, or a humanized version thereof.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

- 5 a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
- b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
- 10 c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
- d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48, or
- e) the heavy chain variable domain is SEQ ID NO:55 and the light chain variable domain is SEQ ID NO:56.

15 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

- a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
- 20 b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
- c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
- d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48.

25 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or

30 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

5 the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

10 the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16, or a humanized version thereof.

15 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;

or a humanized version thereof.

20 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;

or a humanized version thereof.

25 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

- 5 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or,
- 10 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or
- 15 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or
- 20 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or
- 25 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or
- 30 f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46,

5 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54;

10 h) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or

15 i) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

20 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

25 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

30 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of

SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

5 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or

10 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

15 a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

20 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

25 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

30 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID

NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46.

5 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

10 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

15 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30.

Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

20 the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38.

25 Another aspect of the invention is an antibody binding to human CSF-1R, characterized in that

the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of

SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46.

5 The invention comprises a method for the treatment of a patient in need of therapy, characterized by administering to the patient a therapeutically effective amount of an antibody according to the invention.

The invention comprises the use of an antibody according to the invention for therapy.

10 One preferred embodiment of the invention are the CSF-1R antibodies of the present invention for use in the treatment of "CSF-1R mediated diseases" or the CSF-1R antibodies of the present invention for use for the manufacture of a medicament in the treatment of "CSF-1R mediated diseases", which can be described as follows:

15 There are 3 distinct mechanisms by which CSF-1R signaling is likely involved in tumor growth and metastasis. The first is that expression of CSF-ligand and receptor has been found in tumor cells originating in the female reproductive system (breast, ovarian, endometrium, cervical) (Scholl, S.M., et al., J. Natl. Cancer Inst. 86 (1994) 120-126; Kacinski, B.M., Mol. Reprod. Dev. 46 (1997) 71-74; Ngan, H.Y., et al., Eur. J. Cancer 35 (1999) 1546-1550; Kirma, N., et al., Cancer Res 67 (2007) 1918-1926) and the expression has been associated with
20 breast cancer xenograft growth as well as poor prognosis in breast cancer patients. Two point mutations were seen in CSF-1R in about 10-20% of acute myelocytic leukemia, chronic myelocytic leukemia and myelodysplasia patients tested in one study, and one of the mutations was found to disrupt receptor turnover (Ridge, S.A., et al., Proc. Natl. Acad. Sci USA 87 (1990) 1377-1380). However the
25 incidence of the mutations could not be confirmed in later studies (Abu-Duhier, F.M., et al., Br. J. Haematol. 120 (2003) 464-470). Mutations were also found in some cases of hepatocellular cancer (Yang, D.H., et al., Hepatobiliary Pancreat. Dis. Int. 3 (2004) 86-89) and idiopathic myelofibrosis (Abu-Duhier, F.M., et al., Br. J. Haematol. 120 (2003) 464-470). Recently, in the GDM-1 cell line derived
30 from a patient with myelomonoblastic leukemia the Y571D mutation in CSF-1R was identified (Chase, A., et al., Leukemia 23 (2009) 358-364).

Pigmented villonodular synovitis (PVNS) and Tenosynovial Giant cell tumors (TGCT) can occur as a result of a translocation that fuses the M-CSF gene to a collagen gene COL6A3 and results in overexpression of M-CSF (West, R.B., et al., Proc. Natl. Acad. Sci. USA 103 (2006) 690-695). A landscape effect is proposed to be responsible for the resulting tumor mass that consists of monocytic cells attracted by cells that express M-CSF. TGCTs are smaller tumors that can be relatively easily removed from fingers where they mostly occur. PVNS is more aggressive as it can recur in large joints and is not as easily controlled surgically.

The second mechanism is based on blocking signaling through M-CSF/CSF-1R at metastatic sites in bone which induces osteoclastogenesis, bone resorption and osteolytic bone lesions. Breast, multiple myeloma and lung cancers are examples of cancers that have been found to metastasize to the bone and cause osteolytic bone disease resulting in skeletal complications. M-CSF released by tumor cells and stroma induces the differentiation of hematopoietic myeloid monocyte progenitors to mature osteoclasts in collaboration with the receptor activator of nuclear factor kappa-B ligand-RANKL. During this process, M-CSF acts as a permissive factor by giving the survival signal to osteoclasts (Tanaka, S., et al., J. Clin. Invest. 91 (1993) 257-263). Inhibition of CSF-1R activity during osteoclast differentiation and maturation with a anti-CSF-1R antibody is likely to prevent unbalanced activity of osteoclasts that cause osteolytic disease and the associated skeletal related events in metastatic disease. Whereas breast, lung cancer and multiple myeloma typically result in osteolytic lesions, metastasis to the bone in prostate cancer initially has an osteoblastic appearance in which increased bone forming activity results in 'woven bone' which is different from typical lamellar structure of normal bone. During disease progression bone lesions display a significant osteolytic component as well as high serum levels of bone resorption and suggests that anti-resorptive therapy may be useful. Bisphosphonates have been shown to inhibit the formation of osteolytic lesions and reduced the number of skeletal-related events only in men with hormone-refractory metastatic prostate cancer but at this point their effect on osteoblastic lesions is controversial and bisphosphonates have not been beneficial in preventing bone metastasis or hormone responsive prostate cancer to date. The effect of anti-resorptive agents in mixed osteolytic/osteoblastic prostate cancer is still being studied in the clinic (Choueiri, M.B., et al., Cancer Metastasis Rev. 25 (2006) 601-609; Vessella, R.L. and Corey, E., Clin. Cancer Res. 12 (20 Pt 2) (2006) 6285s-6290s).

The third mechanism is based on the recent observation that tumor associated macrophages (TAM) found in solid tumors of the breast, prostate, ovarian and cervical cancers correlated with poor prognosis (Bingle, L., et al., *J. Pathol.* 196 (2002) 254-265; Pollard, J.W., *Nat. Rev. Cancer* 4 (2004) 71-78). Macrophages are recruited to the tumor by M-CSF and other chemokines. The macrophages can then contribute to tumor progression through the secretion of angiogenic factors, proteases and other growth factors and cytokines and may be blocked by inhibition of CSF-1R signaling. Recently it was shown by Zins et al (Zins, K., et al., *Cancer Res.* 67 (2007) 1038-1045) that expression of siRNA of Tumor necrosis factor alpha (TNF alpha), M-CSF or the combination of both would reduce tumor growth in a mouse xenograft model between 34% and 50% after intratumoral injection of the respective siRNA. SiRNA targeting the TNF alpha secreted by the human SW620 cells reduced mouse M-CSF levels and led to reduction of macrophages in the tumor. In addition treatment of MCF7 tumor xenografts with an antigen binding fragment directed against M-CSF did result in 40% tumor growth inhibition, reversed the resistance to chemotherapeutics and improved survival of the mice when given in combination with chemotherapeutics (Paulus, P., et al., *Cancer Res.* 66 (2006) 4349-4356).

TAMs are only one example of an emerging link between chronic inflammation and cancer. There is additional evidence for a link between inflammation and cancer as many chronic diseases are associated with an increased risk of cancer, cancers arise at sites of chronic inflammation, chemical mediators of inflammation are found in many cancers; deletion of the cellular or chemical mediators of inflammation inhibits development of experimental cancers and long-term use of anti-inflammatory agents reduce the risk of some cancers. A link to cancer exists for a number of inflammatory conditions among- those *H.pylori* induced gastritis for gastric cancer, Schistosomiasis for bladder cancer, HHVX for Kaposi's sarcoma, endometriosis for ovarian cancer and prostatitis for prostate cancer (Balkwill, F., et al., *Cancer Cell* 7 (2005) 211-217). Macrophages are key cells in chronic inflammation and respond differentially to their microenvironment. There are two types of macrophages that are considered extremes in a continuum of functional states: M1 macrophages are involved in Type 1 reactions. These reactions involve the activation by microbial products and consequent killing of pathogenic microorganisms that result in reactive oxygen intermediates. On the other end of the extreme are M2 macrophages involved in Type 2 reactions that

promote cell proliferation, tune inflammation and adaptive immunity and promote tissue remodeling, angiogenesis and repair (Mantovani, A., et al., Trends Immunol. 25 (2004) 677-686). Chronic inflammation resulting in established neoplasia is usually associated with M2 macrophages. A pivotal cytokine that mediates inflammatory reactions is TNF alpha that true to its name can stimulate anti-tumor immunity and hemorrhagic necrosis at high doses but has also recently been found to be expressed by tumor cells and acting as a tumor promoter (Zins, K., et al., Cancer Res. 67 (2007) 1038-1045; Balkwill, F., Cancer Metastasis Rev. 25 (2006) 409-416). The specific role of macrophages with respect to the tumor still needs to be better understood including the potential spatial and temporal dependence on their function and the relevance to specific tumor types.

Thus one embodiment of the invention are the CSF-1R antibodies of the present invention for use in the treatment of cancer. The term "cancer" as used herein may be, for example, lung cancer, non small cell lung (NSCL) cancer, bronchioloalviolar cell lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, hepatocellular cancer, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma, lymphoma, lymphocytic leukemia, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers. Preferably such cancer is a breast cancer, ovarian cancer, cervical cancer, lung cancer or prostate cancer. Preferably such cancers are further characterized by CSF-1 or CSF-1R expression or overexpression. One further embodiment the invention are the CSF-1R antibodies of the present invention for use in the simultaneous treatment of primary tumors and new metastases.

Thus another embodiment of the invention are the CSF-1R antibodies of the present invention for use in the treatment of periodontitis, histiocytosis X, osteoporosis, Paget's disease of bone (PDB), bone loss due to cancer therapy, periprosthetic osteolysis, glucocorticoid-induced osteoporosis, rheumatoid arthritis, psiratic arthritis, osteoarthritis, inflammatory arthritides, and inflammation.

Rabello, D., et al., *Biochem. Biophys. Res. Commun.* 347 (2006) 791-796 has demonstrated that SNPs in the CSF1 gene exhibited a positive association with aggressive periodontitis: an inflammatory disease of the periodontal tissues that causes tooth loss due to resorption of the alveolar bone.

Histiocytosis X (also called Langerhans cell histiocytosis, LCH) is a proliferative disease of Langerhans dendritic cells that appear to differentiate into osteoclasts in bone and extra osseous LCH lesions. Langerhans cells are derived from circulating monocytes. Increased levels of M-CSF that have been measured in sera and lesions where found to correlate with disease severity (da Costa, C.E., et al., *J. Exp. Med.* 201 (2005) 687-693). The disease occurs primarily in a pediatric patient population and has to be treated with chemotherapy when the disease becomes systemic or is recurrent.

The pathophysiology of osteoporosis is mediated by loss of bone forming osteoblasts and increased osteoclast dependent bone resorption. Supporting data has been described by Cenci et al showing that an anti-M-CSF antibody injection preserves bone density and inhibits bone resorption in ovariectomized mice (Cenci, S., et al., *J. Clin. Invest.* 105 (2000) 1279-1287). Recently a potential link between postmenopausal bone loss due to estrogen deficiency was identified and found that the presence of TNF alpha producing T-cell affected bone metabolism (Roggia, C., et al., *Minerva Med.* 95 (2004) 125-132). A possible mechanism could be the induction of M-CSF by TNF alpha in vivo. An important role for M-CSF in TNF-alpha-induced osteoclastogenesis was confirmed by the effect of an antibody directed against M-CSF that blocked the TNF alpha induced osteolysis in mice and thereby making inhibitors of CSF-1R signaling potential targets for inflammatory arthritis (Kitaura, H., et al., *J. Clin. Invest.* 115 (2005) 3418-3427).

Paget's disease of bone (PDB) is the second most common bone metabolism disorder after osteoporosis in which focal abnormalities of increased bone turnover lead to complications such as bone pain, deformity, pathological fractures and

deafness. Mutations in four genes have been identified that regulate normal osteoclast function and predispose individuals to PDB and related disorders: insertion mutations in TNFRSF11A, which encodes receptor activator of nuclear factor (NF) kappaB (RANK)-a critical regulator of osteoclast function, inactivating mutations of TNFRSF11B which encodes osteoprotegerin (a decoy receptor for RANK ligand), mutations of the sequestosome 1 gene (SQSTM1), which encodes an important scaffold protein in the NFkappaB pathway and mutations in the valosin-containing protein (VCP) gene. This gene encodes VCP, which has a role in targeting the inhibitor of NFkappaB for degradation by the proteasome (Doroszewska, A. and Ralston, S.H., *Nat. Clin. Pract. Rheumatol.* 2 (2006) 270-277). Targeted CSF-1R inhibitors provide an opportunity to block the deregulation of the RANKL signaling indirectly and add an additional treatment option to the currently used bisphosphonates.

Cancer therapy induced bone loss especially in breast and prostate cancer patients is an additional indication where a targeted CSF-1R inhibitor could prevent bone loss (Lester, J.E., et al., *Br. J. Cancer* 94 (2006) 30-35). With the improved prognosis for early breast cancer the long-term consequences of the adjuvant therapies become more important as some of the therapies including chemotherapy, irradiation, aromatase inhibitors and ovary ablation affect bone metabolism by decreasing the bone mineral density, resulting in increased risk for osteoporosis and associated fractures (Lester, J.E., et al., *Br. J. Cancer* 94 (2006) 30-35). The equivalent to adjuvant aromatase inhibitor therapy in breast cancer is androgen ablation therapy in prostate cancer which leads to loss of bone mineral density and significantly increases the risk of osteoporosis-related fractures (Stoch, S.A., et al., *J. Clin. Endocrinol. Metab.* 86 (2001) 2787-2791).

Targeted inhibition of CSF-1R signaling is likely to be beneficial in other indications as well when targeted cell types include osteoclasts and macrophages e.g. treatment of specific complications in response to joint replacement as a consequence of rheumatoid arthritis. Implant failure due to periprosthetic bone loss and consequent loosening of prostheses is a major complication of joint replacement and requires repeated surgery with high socioeconomic burdens for the individual patient and the health-care system. To date, there is no approved drug therapy to prevent or inhibit periprosthetic osteolysis (Drees, P., et al., *Nat. Clin. Pract. Rheumatol.* 3 (2007) 165-171).

Glucocorticoid-induced osteoporosis (GIOP) is another indication in which a CSF-1R inhibitor could prevent bone loss after longterm glucocorticosteroid use that is given as a result of various conditions among those chronic obstructive pulmonary disease, asthma and rheumatoid arthritis (Guzman-Clark, J.R., et al.,
5 Arthritis Rheum. 57 (2007) 140-146; Feldstein, A.C., et al., Osteoporos. Int. 16 (2005) 2168-2174).

Rheumatoid arthritis, psoriatic arthritis and inflammatory arthritides are in itself potential indications for CSF-1R signaling inhibitors in that they consist of a macrophage component and to a varying degree bone destruction (Ritchlin, C.T., et al.,
10 J. Clin. Invest. 111 (2003) 821-831). Osteoarthritis and rheumatoid arthritis are inflammatory autoimmune disease caused by the accumulation of macrophages in the connective tissue and infiltration of macrophages into the synovial fluid, which is at least partially mediated by M-CSF. Campbell, I., K., et al., J. Leukoc. Biol. 68 (2000) 144-150, demonstrated that M-CSF is produced by human-joint tissue cells
15 (chondrocytes, synovial fibroblasts) in vitro and is found in synovial fluid of patients with rheumatoid arthritis, suggesting that it contributes to the synovial tissue proliferation and macrophage infiltration which is associated with the pathogenesis of the disease. Inhibition of CSF-1R signaling is likely to control the number of macrophages in the joint and alleviate the pain from the associated bone
20 destruction. In order to minimize adverse affects and to further understand the impact of the CSF-1R signaling in these indications, one method is to specifically inhibit CSF-1R without targeting a myriad other kinases, such as Raf kinase.

Recent literature reports correlate increased circulating M-CSF with poor prognosis and atherosclerotic progression in chronic coronary artery disease (Saitoh, T., et al.,
25 J. Am. Coll. Cardiol. 35 (2000) 655-665; Ikonomidis, I., et al., Eur. Heart. J. 26 (2005) p. 1618-1624); M-CSF influences the atherosclerotic process by aiding the formation of foam cells (macrophages with ingested oxidized LDL) that express CSF-1R and represent the initial plaque (Murayama, T., et al., Circulation 99 (1999) 1740-1746).

30 Expression and signaling of M-CSF and CSF-1R is found in activated microglia. Microglia, which are resident macrophages of the central nervous system, can be activated by various insults, including infection and traumatic injury. M-CSF is considered a key regulator of inflammatory responses in the brain and M-CSF

levels increase in HIV-1, encephalitis, Alzheimer's disease (AD) and brain tumors. Microgliosis as a consequence of autocrine signaling by M-CSF/CSF-1R results in induction of inflammatory cytokines and nitric oxides being released as demonstrated by e.g. using an experimental neuronal damage model (Hao, A.J., et al., Neuroscience 112 (2002) 889-900; Murphy, G.M., Jr., et al., J. Biol. Chem. 273 (1998) 20967-20971). Microglia that have increased expression of CSF-1R are found to surround plaques in AD and in the amyloid precursor protein V717F transgenic mouse model of AD (Murphy, G.M., Jr., et al., Am. J. Pathol. 157 (2000) 895-904). On the other hand op/op mice with fewer microglia in the brain resulted in fibrillar deposition of A-beta and neuronal loss compared to normal control suggesting that microglia do have a neuroprotective function in the development of AD lacking in the op/op mice (Kaku, M., et al., Brain Res. Brain Res. Protoc. 12 (2003) 104-108).

Expression and signaling of M-CSF and CSF-1R is associated with inflammatory bowel disease (IBD) (WO 2005/046657). The term "inflammatory bowel disease" refers to serious, chronic disorders of the intestinal tract characterized by chronic inflammation at various sites in the gastrointestinal tract, and specifically includes ulcerative colitis (UC) and Crohn's disease.

The invention comprises an antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of cancer.

The invention comprises an antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of bone loss.

The invention comprises an antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the prevention or treatment of metastasis.

The invention comprises an antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or

alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for treatment of inflammatory diseases.

5 The invention comprises the use of an antibody characterized in comprising the antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of cancer or alternatively for the manufacture of a medicament for the treatment of cancer.

10 The invention comprises the use of an antibody characterized in comprising the antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of bone loss or alternatively for the manufacture of a medicament for the treatment of bone loss.

15 The invention comprises the use of an antibody characterized in comprising the antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the prevention or treatment of metastasis or alternatively for the manufacture of a medicament for the prevention or treatment of metastasis.

20

The invention comprises the use of an antibody characterized in comprising the antibody binding to human CSF-1R being characterized by the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for treatment of inflammatory diseases or alternatively for the manufacture of a medicament for the treatment of inflammatory diseases.

25

A further embodiment of the invention is a method for the production of an antibody against CSF-1R characterized in that the sequence of a nucleic acid encoding the heavy chain of a human IgG1 class antibody binding to human CSF-1R according to the invention said modified nucleic acid and the nucleic acid encoding the light chain of said antibody are inserted into an expression vector,

30

said vector is inserted in a eukaryotic host cell, the encoded protein is expressed and recovered from the host cell or the supernatant.

5 The antibodies according to the invention are preferably produced by recombinant means. Such methods are widely known in the state of the art and comprise protein expression in prokaryotic and eukaryotic cells with subsequent isolation of the antibody polypeptide and usually purification to a pharmaceutically acceptable purity. For the protein expression nucleic acids encoding light and heavy chains or fragments thereof are inserted into expression vectors by standard methods. Expression is performed in appropriate prokaryotic or eukaryotic host cells, such as
10 CHO cells, NS0 cells, SP2/0 cells, HEK293 cells, COS cells, yeast, or E. coli cells, and the antibody is recovered from the cells (from the supernatant or after cells lysis).

Recombinant production of antibodies is well-known in the state of the art and described, for example, in the review articles of Makrides, S.C., Protein Expr. Purif. 17 (1999) 183-202; Geisse, S., et al., Protein Expr. Purif. 8 (1996) 271-282;
15 Kaufman, R.J., Mol. Biotechnol. 16 (2000) 151-161; Werner, R.G., Drug Res. 48 (1998) 870-880.

The antibodies may be present in whole cells, in a cell lysate, or in a partially purified, or substantially pure form. Purification is performed in order to eliminate
20 other cellular components or other contaminants, e.g. other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis, and others well known in the art. See Ausubel, F., et al., ed. Current Protocols in Molecular Biology, Greene Publishing and Wiley Interscience, New York (1987).

25 Expression in NS0 cells is described by, e.g., Barnes, L.M., et al., Cytotechnology 32 (2000) 109-123; Barnes, L.M., et al., Biotech. Bioeng. 73 (2001) 261-270. Transient expression is described by, e.g., Durocher, Y., et al., Nucl. Acids. Res. 30 (2002) E9. Cloning of variable domains is described by Orlandi, R., et al., Proc. Natl. Acad. Sci. USA 86 (1989) 3833-3837; Carter, P., et al., Proc. Natl. Acad. Sci.
30 USA 89 (1992) 4285-4289; Norderhaug, L., et al., J. Immunol. Methods 204 (1997) 77-87. A preferred transient expression system (HEK 293) is described by Schlaeger, E.-J. and Christensen, K., in Cytotechnology 30 (1999) 71-83, and by Schlaeger, E.-J., in J. Immunol. Methods 194 (1996) 191-199.

Nucleic acid molecules encoding amino acid sequence variants of anti-CSF-1R antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by
5 oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of humanized anti-CSF-1R antibody.

The heavy and light chain variable domains according to the invention are combined with sequences of promoter, translation initiation, constant region, 3'
10 untranslated region, polyadenylation, and transcription termination to form expression vector constructs. The heavy and light chain expression constructs can be combined into a single vector, co-transfected, serially transfected, or separately transfected into host cells which are then fused to form a single host cell expressing both chains.

15 In another aspect, the present invention provides a composition, e.g. a pharmaceutical composition, containing one or a combination of monoclonal antibodies, or the antigen-binding portion thereof, of the present invention, formulated together with a pharmaceutically acceptable carrier.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents,
20 dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption/resorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for injection or infusion.

A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route
25 and/or mode of administration will vary depending upon the desired results.

Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. In addition to water, the carrier can be, for example,
30 an isotonic buffered saline solution.

Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient (effective amount). The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

The invention comprises the use of the antibodies according to the invention for the treatment of a patient suffering from cancer, especially from colon, lung or pancreas cancer.

The invention comprises also a method for the treatment of a patient suffering from such disease.

The invention further provides a method for the manufacture of a pharmaceutical composition comprising an effective amount of an antibody according to the invention together with a pharmaceutically acceptable carrier and the use of the antibody according to the invention for such a method.

The invention further provides the use of an antibody according to the invention in an effective amount for the manufacture of a pharmaceutical agent, preferably together with a pharmaceutically acceptable carrier, for the treatment of a patient suffering from cancer.

The invention also provides the use of an antibody according to the invention in an effective amount for the manufacture of a pharmaceutical agent, preferably together with a pharmaceutically acceptable carrier, for the treatment of a patient suffering from cancer.

- 5 The following examples, sequence listing and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the Sequences

10	SEQ ID NO: 1	heavy chain CDR3, Mab 2F11
	SEQ ID NO: 2	heavy chain CDR2, Mab 2F11
	SEQ ID NO: 3	heavy chain CDR1, Mab 2F11
	SEQ ID NO: 4	light chain CDR3, Mab 2F11
	SEQ ID NO: 5	light chain CDR2, Mab 2F11
15	SEQ ID NO: 6	light chain CDR1, Mab 2F11
	SEQ ID NO: 7	heavy chain variable domain, Mab 2F11
	SEQ ID NO: 8	light chain variable domain, Mab 2F11
	SEQ ID NO: 9	heavy chain CDR3, Mab 2E10
	SEQ ID NO: 10	heavy chain CDR2, Mab 2E10
20	SEQ ID NO: 11	heavy chain CDR1, Mab 2E10
	SEQ ID NO: 12	light chain CDR3, Mab 2E10
	SEQ ID NO: 13	light chain CDR2, Mab 2E10
	SEQ ID NO: 14	light chain CDR1, Mab 2E10
	SEQ ID NO: 15	heavy chain variable domain, Mab 2E10
25	SEQ ID NO: 16	light chain variable domain, Mab 2E10
	SEQ ID NO: 17	heavy chain CDR3, hMab 2F11-c11
	SEQ ID NO: 18	heavy chain CDR2, hMab 2F11-c11
	SEQ ID NO: 19	heavy chain CDR1, hMab 2F11-c11
	SEQ ID NO: 20	light chain CDR3, hMab 2F11-c11
30	SEQ ID NO: 21	light chain CDR2, hMab 2F11-c11
	SEQ ID NO: 22	light chain CDR1, hMab 2F11-c11
	SEQ ID NO: 23	heavy chain variable domain, hMab 2F11-c11
	SEQ ID NO: 24	light chain variable domain, hMab 2F11-c11

	SEQ ID NO: 25	heavy chain CDR3, hMab 2F11-d8
	SEQ ID NO: 26	heavy chain CDR2, hMab 2F11-d8
	SEQ ID NO: 27	heavy chain CDR1, hMab 2F11-d8
	SEQ ID NO: 28	light chain CDR3, hMab 2F11-d8
5	SEQ ID NO: 29	light chain CDR2, hMab 2F11-d8
	SEQ ID NO: 30	light chain CDR1, hMab 2F11-d8
	SEQ ID NO: 31	heavy chain variable domain, hMab 2F11-d8
	SEQ ID NO: 32	light chain variable domain, hMab 2F11-d8
	SEQ ID NO: 33	heavy chain CDR3, hMab 2F11-e7
10	SEQ ID NO: 34	heavy chain CDR2, hMab 2F11-e7
	SEQ ID NO: 35	heavy chain CDR1, hMab 2F11-e7
	SEQ ID NO: 36	light chain CDR3, hMab 2F11-e7
	SEQ ID NO: 37	light chain CDR2, hMab 2F11-e7
	SEQ ID NO: 38	light chain CDR1, hMab 2F11-e7
15	SEQ ID NO: 39	heavy chain variable domain, hMab 2F11-e7
	SEQ ID NO: 40	light chain variable domain, hMab 2F11-e7
	SEQ ID NO: 41	heavy chain CDR3, hMab 2F11-f12
	SEQ ID NO: 42	heavy chain CDR2, hMab 2F11-f12
	SEQ ID NO: 43	heavy chain CDR1, hMab 2F11-f12
20	SEQ ID NO: 44	light chain CDR3, hMab 2F11-f12
	SEQ ID NO: 45	light chain CDR2, hMab 2F11-f12
	SEQ ID NO: 46	light chain CDR1, hMab 2F11-f12
	SEQ ID NO: 47	heavy chain variable domain, hMab 2F11-f12
	SEQ ID NO: 48	light chain variable domain, hMab 2F11-f12
25	SEQ ID NO: 49	heavy chain CDR3, hMab 2F11-g1
	SEQ ID NO: 50	heavy chain CDR2, hMab 2F11-g1
	SEQ ID NO: 51	heavy chain CDR1, hMab 2F11-g1
	SEQ ID NO: 52	light chain CDR3, hMab 2F11-g1
	SEQ ID NO: 53	light chain CDR2, hMab 2F11-g1
30	SEQ ID NO: 54	light chain CDR1, hMab 2F11-g1
	SEQ ID NO: 55	heavy chain variable domain, hMab 2F11-g1
	SEQ ID NO: 56	light chain variable domain, hMab 2F11-g1
	SEQ ID NO: 57	human kappa light chain constant region
	SEQ ID NO: 58	human heavy chain constant region derived from IgG1

	SEQ ID NO: 59	human heavy chain constant region derived from IgG1 mutated on L234A and L235A
	SEQ ID NO: 60	human heavy chain constant region derived from IgG4
5	SEQ ID NO: 61	human heavy chain constant region derived from IgG4 mutated on S228P
	SEQ ID NO: 62	human wildtype CSF-1R (wt CSF-1R)
	SEQ ID NO: 63	human mutant CSF-1R L301S Y969F
	SEQ ID NO: 64	human CSF-1R Extracellular Domain
	SEQ ID NO: 65	human CSF-1R fragment delD4
10	SEQ ID NO: 66	human CSF-1R fragment D1-D3
	SEQ ID NO: 67	signal peptide
	SEQ ID NO: 68	Primer
	SEQ ID NO: 69	heavy chain CDR3, Mab 1G10
	SEQ ID NO: 70	heavy chain CDR2, Mab 1G10
15	SEQ ID NO: 71	heavy chain CDR1, Mab 1G10
	SEQ ID NO: 72	light chain CDR3, Mab 1G10
	SEQ ID NO: 73	light chain CDR2, Mab 1G10
	SEQ ID NO: 74	light chain CDR1, Mab 1G10
	SEQ ID NO: 75	heavy chain variable domain, Mab 1G10
20	SEQ ID NO: 76	light chain variable domain, Mab 1G10
	SEQ ID NO: 77	heavy chain CDR3, Mab 2H7
	SEQ ID NO: 78	heavy chain CDR2, Mab 2H7
	SEQ ID NO: 79	heavy chain CDR1, Mab 2H7
	SEQ ID NO: 80	light chain CDR3, Mab 2H7
25	SEQ ID NO: 81	light chain CDR2, Mab 2H7
	SEQ ID NO: 82	light chain CDR1, Mab 2H7
	SEQ ID NO: 83	heavy chain variable domain, Mab 2H7
	SEQ ID NO: 84	light chain variable domain, Mab 2H7

30 The following examples, sequence listing and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the Figures

- Figure 1** Growth inhibition of BeWo tumor cells in 3D culture under treatment with different anti-CSF-1R monoclonal antibodies at a concentration of 10 μ g/ml.
- 5 X axis: viability normalized mean relative light units (RLU) corresponding to the ATP-content of the cells (CellTiterGlo assay).
- Y axis: tested probes: Minimal Medium (0.5% FBS), mouse IgG1 (mIgG1, 10 μ g/ml), mouse IgG2a (mIgG2a 10 μ g/ml), CSF-1 only, Mab 2F11, Mab 2E10, Mab2H7, Mab1G10 and SC 2-4A5.
- 10 Highest inhibition of CSF-1 induced growth was observed with the anti-CSF-1R antibodies according to the invention.
- Figure 2a** Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R fragment delD4 (comprising the extracellular subdomains D1 –D3 and D5) (SEQ ID NO: 65) (y-axis: binding signal in Response Units (RU), baseline = 0 RU, x-axis: time in seconds (s)): While the antibodies Mab 3291 and sc 2-4A5 clearly show binding to this delD4 fragment, the antibodies according to the invention e.g. Mab 2F11, and Mab 2E10, did not bind to the CSF-1R fragment delD4. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did also not bind to the CSF-1R fragment delD4.
- 15
- 20
- Figure 2b** Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 – D5) (SEQ ID NO: 64) (y-axis: binding signal in Response Units (RU), baseline = 0 RU, x-axis: time in seconds (s)):
- 25 All anti-CSF-1R antibodies show binding to CSF-1R-ECD. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did not bind to the CSF-1R-ECD.
- 30
- Figure 2c** Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R fragment delD4 (comprising the extracellular subdomains D1 –D3 and D5) (SEQ ID NO: 65) (y-axis: binding signal in Response Units (RU), baseline = 0 RU, x-axis: time in seconds (s)): Mab 1G10, Mab
- 35

2H7 and humanized hMab 2F11-e7 did not bind to the CSF-1R fragment delD4. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did also not bind to the CSF-1R fragment delD4.

5 **Figure 2d**

Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 – D5) (SEQ ID NO: 64) (y-axis: binding signal in Response Units (RU), baseline = 0 RU, x-axis: time in seconds (s)): All anti-CSF-1R antibodies Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 showed binding to CSF-1R-ECD. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did not bind to the CSF-1R-ECD.

10

Figure 2e

Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R fragment delD4 (comprising the extracellular subdomains D1 –D3 and D5) (SEQ ID NO: 65) (y-axis: binding signal in Response Units (RU), baseline = 0 RU, x-axis: time in seconds (s)): All anti-CSF-1R antibodies 1.2.SM, CXIIG6, ab10676 and MAB3291 show binding to to the CSF-1R fragment delD4. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did also not bind to the CSF-1R fragment delD4.

15

20

Figure 2f

Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 – D5) (SEQ ID NO: 64) (y-axis: binding signal in Response Units (RU), baseline = 0 RU, x-axis: time in seconds (s)):

25

All anti-CSF-1R antibodies 1.2.SM, CXIIG6, ab10676 and MAB3291 show binding to CSF-1R-ECD. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did not bind to the CSF-1R-ECD.

30

Figure 3a-d

CSF-1 levels in Cynomolgus monkey after application of different dosages of anti-CSF-1R antibody according to the invention

Figure 4 In vivo efficacy – tumor growth inhibition of anti-CSF-1R antibodies according to the invention in breast cancer BT20 xenograft

5 **Example 1**

Generation of a hybridoma cell line producing anti-CSF-1R antibodies

Immunization procedure of NMRI mice

NMRI mice were immunized with an expression vector pDisplay™ (Invitrogen, USA) encoding the extracellular domain of huCSF-1R by utilizing electroporation.
10 Every mouse was 4 times immunized with 100µg DNA. When serum titers of anti-huCSF-1R were found to be sufficient, mice were additionally boosted once with 50µg of a 1:1 mixture huCSF-1R ECD/huCSF-1R ECDhuFc chimera in 200 µl PBS intravenously (i.v.) 4 and 3 days before fusion.

Antigen specific ELISA

15 Anti-CSF-1R titers in sera of immunized mice were determined by antigen specific ELISA.

0.3 µg/ml huCSF-1R-huFc chimera (soluble extracellular domain) was captured on a streptavidin plate (MaxiSorb; MicroCoat, DE, Cat.No. 11974998/MC1099) with 0.1 mg/ml biotinylated anti Fcγ (Jackson ImmunoResearch, Cat.No. 109-066-098)
20 and horse radish peroxidase (HRP)-conjugated F(ab')₂ anti mouse IgG (GE Healthcare, UK, Cat.No.NA9310V) diluted 1/800 in PBS/0.05% Tween20/0.5% BSA was added. Sera from all taps were diluted 1/40 in PBS/0.05% Tween20/0.5% BSA and serially diluted up to 1/1638400. Diluted sera were added to the wells. Pre-tap serum was used as negative control. A dilution series of mouse anti-human
25 CSF-1R Mab3291 (R&D Systems, UK) from 500 ng/ml to 0,25 ng/ml was used as positive control. All components were incubated together for 1,5 hours, Wells were washed 6 times with PBST (PBS/0.2% Tween20) and assays were developed with freshly prepared ABTS® solution (1 mg/ml) (ABTS: 2,2'-azino bis (3-ethylbenzthiazoline-6-sulfonic acid) for 10 minutes at RT. Absorbance was
30 measured at 405 nm.

Hybridoma generation

The mouse lymphocytes can be isolated and fused with a mouse myeloma cell line using PEG based standard protocols to generate hybridomas. The resulting hybridomas are then screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic derived lymphocytes from immunized mice are fused to Ag8 non-secreting mouse myeloma cells P3X63Ag8.653 (ATCC, CRL-1580) with 50% PEG. Cells are plated at approximately 10^4 in flat bottom 96 well micro titer plate, followed by about two weeks incubation in selective medium. Individual wells are then screened by ELISA for human anti-CSF-1R monoclonal IgM and IgG antibodies. Once extensive hybridoma growth occurs, the antibody secreting hybridomas are replated, screened again, and if still positive for human IgG, anti-CSF-1R monoclonal antibodies, can be subcloned by FACS. The stable subclones are then cultured in vitro to produce antibody in tissue culture medium for characterization. Antibodies according to the invention could be selected using the determination of the binding of anti-CSF-1R antibodies to human CSF-1R fragment delD4 and to human CSF-1R Extracellular Domain (CSF-1R-ECD) as described in Example 4, as well as the determination of growth inhibition of NIH3T3 cells transfected with wildtype CSF-1R (ligand dependent signalling) or mutant CSF-1R L301S Y969F (ligand independent signalling) under treatment with anti-CSF-1R monoclonal antibodies as described in Example 5.

Culture of hybridomas

Generated muMAb hybridomas were cultured in RPMI 1640 (PAN – Catalogue No. (Cat. No.) PO4-17500) supplemented with 2 mM L-glutamine (GIBCO - Cat. No.35050-038), 1 mM Na-Pyruvat (GIBCO - Cat. No.11360-039), 1x NEAA (GIBCO - Cat. No.11140-035), 10% FCS (PAA - Cat. No.A15-649), 1x Pen Strep (Roche - Cat. No.1074440), 1x Nutridoma CS (Roche - Cat. No.1363743), 50 μ M Mercaptoethanol (GIBCO - Cat. No.31350-010) and 50 U/ml IL 6 mouse (Roche - Cat. No.1 444 581) at 37°C and 5% CO₂. Some of the resulting mouse antibodies have been humanized (e.g. Mab 2F11) and been expressed recombinantly.

Example 2**Inhibition of CSF-1 binding to CSF-1R (ELISA)**

By setting-up this assay to first allow for anti-CSF-1R antibody binding to the CSF-1R -ECD followed by detection of ligand not bound to the receptor both-
5 ligand displacing antibodies and dimerization inhibitor anti-CSF-1R antibodies -
can be tested. The test was performed on 384 well microtiter plates (MicroCoat,
DE, Cat.No. 464718) at RT. After each incubation step plates were washed 3 times
with PBST.

At the beginning, plates were coated with 0.5 mg/ml goat F(ab')₂ biotinylated anti
10 Fc γ (Jackson ImmunoResearch., Cat.No.109-006-170) for 1 hour (h).

Thereafter the wells were blocked with PBS supplemented with 0.2% Tween[®]-20
and 2% BSA (Roche Diagnostics GmbH, DE) for 0.5 h. 75 ng/ml of huCSF-1R-
huFc chimera (which forms the dimeric soluble extracellular domain of huCSF-1R)
was immobilized to plate for 1 h. Then dilutions of purified antibodies in
15 PBS/0.05% Tween20/0.5% BSA were incubated for 1 h. After adding a mixture of
3 ng/ml CSF-1 (Biomol, DE, Cat.No.60530), 50ng/ml biotinylated anti CSF-1
clone BAF216 (R&D Systems,UK) and 1:5000 diluted streptavidin HRP (Roche
Diagnostics GmbH, DE, Cat.No.11089153001) for 1 h the plates were washed 6
times with PBST. Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US), which
20 inhibits the ligand- receptor interaction, was used as positive control. Plates were
developed with freshly prepared BM blue[®] POD substrate solution (BM blue[®]:
3,3'-5,5'-Tetramethylbenzidine, Roche Diagnostics GmbH, DE, Cat.No.
11484281001) for 30 minutes at RT. Absorbance was measured at 370 nm. A
decrease of absorbance is found, if the anti-CSF-1R antibody causes a release of
25 CSF-1 from the dimeric complex. All anti-CSF-1R antibodies showed significant
inhibition of the CSF-1 interaction with CSF-1R (see Table 1). Anti CSF-1R SC 2-
4A5 (Santa Cruz Biotechnology, US see also Sherr, C.J. et al., Blood 73 (1989)
1786-1793), which inhibits the ligand- receptor interaction, was used as reference
control.

Table 1:
Calculated IC₅₀ values for the inhibition of the CSF-1/CSF-1R interaction

CSF-1R Mab	IC₅₀ CSF-1 /CSF-1R Inhibition [ng/ml]
Mab 2F11	19.3
Mab 2E10	20.6
Mab 2H7	18.2
Mab 1G10	11.8
SC-2-4A5	35.2

Example 3

5 **Inhibition of CSF-1-induced CSF-1R phosphorylation in NIH3T3-CSF-1R recombinant cells**

4.5x10³ NIH 3T3 cells, retrovirally infected with an expression vector for full-length CSF-1R, were cultured in DMEM (PAA Cat. No.E15-011), 2mM L-glutamine (Sigma, Cat.No.G7513, 2mM Sodium pyruvate , 1x nonessential aminoacids, 10% FKS (PAA, Cat.No.A15-649) and 100µg/ml PenStrep (Sigma, Cat.No. P4333 [10mg/ml]) until they reached confluency. Thereafter cells were washed with serum-free DMEM media (PAA Cat.No.E15-011) supplemented with sodium selenite [5ng/ml] (Sigma, Cat.No. S9133), transferrin [10µg/ml] (Sigma, Cat.No. T8158), BSA [400µg/ml] (Roche Diagnostics GmbH, Cat.No. 10735078), 15 4mM L-glutamine (Sigma, Cat.No.G7513), 2mM sodium pyruvate (Gibco, Cat.No. 11360), 1x nonessential aminoacids (Gibco, Cat: 11140-035), 2-mercaptoethanol [0,05mM] (Merck, Cat.No. M7522), 100µg/ml and PenStrep (Sigma, Cat. No. P4333) and incubated in 30 µl of the same medium for 16 hours to allow for receptor up-regulation. 10 µl of diluted anti-CSF-1R antibodies were added to the cells for 1.5 h. Then cells were stimulated with 10 µl of 100 ng/ml huM-CSF-1 (Biomol Cat.No.60530) for 5 min. After the incubation, supernatant was removed, cells were washed twice with 80 µl of ice-cold PBS and 50 µl of freshly prepared ice-cold lysis buffer (150mM NaCl/ 20mM Tris pH 7.5 / 1mM EDTA/ 1mM EGTA/ 1% Triton X-100 / 1 protease inhibitor tablet (Roche Diagnostics GmbH Cat.No.1 836 170) per 10 ml buffer/10µl/ml phosphatase inhibitor cocktail 1 (Sigma Cat.No. P-2850, 100x Stock), 10µl/ml protease inhibitor 1 (Sigma Cat.No.P-5726, 100x Stock) /10µl/ml 1 M NaF) was added. After 30 minutes on 25

ice the plates were shaken vigourously on a plateshaker for 3 minutes and then centrifuged 10 minutes at 2200 rpm (Heraeus Megafuge 10).

The presence of phosphorylated and total CSF-1 receptor in the cell lysate was analyzed with Elisa. For detection of the phosphorylated receptor the kit from R&D Systems (Cat. No. DYC3268-2) was used according to the instructions of the supplier. For detection of total CSF-1R 10 μ l of the lysate was immobilized on plate by use of the capture antibody contained in the kit. Thereafter 1:750 diluted biotinylated anti CSF-1R antibody BAF329 (R&D Systems) and 1:1000 diluted streptavidin-HRP conjugate was added. After 60 minutes plates were developed with freshly prepared ABTS[®] solution and the absorbance was detected. Data were calculated as % of positive control without antibody and the ratio value phospho/total receptor expressed. The negative control was defined without addition of M-CSF-1. Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US, see also Sherr, C.J. et al., Blood 73 (1989) 1786-1793), which inhibits the ligand-receptor interaction, was used as reference control.

Table 2:

Calculated IC50 values for the inhibition of CSF-1 receptor phosphorylation.

CSF-1R Mab	IC50 CSF-1R Phosphorylation [ng/ml]
Mab 2F11	219.4
Mab 2E10	752.0
Mab 2H7	703.4
Mab 1G10	56.6
SC-2-4A5	1006.6

Example 4

Determination of the binding of anti-CSF-1R antibodies to human CSF-1R fragment delD4 and to human CSF-1R Extracellular Domain (CSF-1R-ECD)

Preparation of human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 –D5, hCSF-1R-ECD) of SEQ ID NO: 64:

pCMV-preS-Fc-hCSF-1R-ECD (7836bp) encodes the complete ECD of human CSF-1R (SEQ ID NO: 64) C-terminally fused to a PreScission protease cleavage

site, followed by aa100-330 of human IgG1 and a 6xHis-Tag, under the control of CMV promoter. The natural signal peptide has been varied by insertion of amino acids G and S after the first M, in order to create a BamHI restriction site.

Preparation of human CSF-1R fragment delD4 (comprising the extracellular subdomains D1 –D3 and D5, hCSF-1R-delD4) of SEQ ID NO: 65:

5

hCSF1R-delD4-V1-PreSc-hFc-His was cloned from pCMV-preS-Fc-hCSF-1R-ECD by means of the Stratagene QuikChange XL site-directed mutagenesis protocol, using delD4-for with sequence CACCTCCATGTTCTTCCGGTACCCCCAGAGGTAAG (SEQ ID NO: 68) as the forward primer and delD4-rev with the reverse complement sequence as the reverse primer. A protocol variation published in BioTechniques 26 (1999) 680 was used to extend both primers in separate reactions in three cycles preceding the regular Stratagene protocol:

10

15

Two separate 50 µl reaction mixtures were set up according to the manufacturer's manual, each containing 10 ng plasmid pCMV-preS-Fc-hCSF1R-ECD as the template and 10 pM of one of the primers delD4-for or delD4-rev, and 0,5 µl Pfu DNA polymerase as provided with the kit. Three PCR cycles 95 °C 30 sec / 55 °C 60 sec / 68 °C 8 min were run, then 25 µl each of both reaction mixtures were combined in a new tube and 0,5 µl fresh Pfu DNA polymerase were added. The regular PCR protocol with 18 temperature cycles as specified by Stratagene in the kit manual was carried out, followed by 2 hrs final digestion with the DpnI restriction enzyme provided with the kit. Clones bearing the deletion were detected by digestion with Cel II and Not I and verified by sequencing.

20

25

Protein was prepared by transient transfection in the Hek293 FreeStyle suspension cell system (Invitrogen) according to the manufacturer's specifications. After 1 week 500 ml supernatant was filtered and loaded onto a 1ml HiTrap MabSelect Xtra (GE healthcare) protein A column (0,2 ml /min). The column was washed first with PBS, then with 50 mM Tris/ 150 mM NaCl/ 1 mM EDTA/ pH 7,3. 75 µl PreScission Protease (GE #27-0843-01) diluted in 375 µl of the same buffer were loaded onto the column and the closed column was incubated over night at 4 °C with rolling. The column was mounted on top of a 1 ml GStrap FF column (GE healthcare) and the desired protein was eluted (0,2 ml/min, 0,2 ml fractions). Pooled

30

fractions were concentrated from 1,8 ml to 0,4 ml by centrifugal ultrafiltration via a 3k Nanosep and chromatographed over an S200 HR SEC in PBS (0,5 ml/min).

Human CSF-1R fragment delD4 was obtained in two fractions as a dimeric molecule (pool1, V=1,5 ml; c= 0,30 mg/ml; apparent mass on SDS page 83 kDa, reduced 62 kDa) and as the monomer (pool 2, V=1,4 ml; c = 0,25 mg/ml apparent mass on SDS page 62 kDa). The dimeric form was used for all experiments.

Determination of the binding of anti-CSF-1R antibodies to human CSF-1R fragment delD4 and to human CSF-1R Extracellular Domain (CSF-1R-ECD) (binding signals as Response Units (RU):

10 Instrument: Biacore T100 (GE Healthcare)
Software: T100 Control, Version 2.0.1
T100 Evaluation, Version 2.0.2
Assayformat Chip: CM5
Temperature: 25°C

15 CSF-1R fragments were immobilized via amine coupling. To compare the binding of different anti-CSF-1R antibodies according to the invention one concentration of the test antibody was injected. Anti CSF-1R Mab3291 (R&D-Systems) and SC 2-4A5 (Santa Cruz Biotechnology, US- see also Sherr, C.J. et al., Blood 73 (1989) 1786-1793), was used as reference control, anti-CCR5 m<CCR5>Pz03.1C5
20 (deposited as DSM ACC 2683 on 18.08.2004 at DSMZ) as negative control, all under the same conditions as the anti-CSF-1R antibodies according to the invention.

Amine coupling of CSF-1R fragments

Standard amine coupling according to the manufacturer's instructions: running
25 buffer: PBS-T (Roche: 11 666 789 + 0.05% Tween20: 11 332 465), activation by mixture of EDC/NHS, injection of human CSF-1R fragment delD4 (comprising the extracellular subdomains D1 –D3 and D5) (SEQ ID NO: 65) and human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 –D5) (SEQ ID NO: 64) for 600 seconds at flow rate 10µl/min; diluted in coupling
30 buffer NaAc, pH 5.0, c = 10 µg/mL; finally remaining activated carboxyl groups were blocked by injection of 1 M Ethanolamin.

Binding of <CSF-1R> Mab 2F11, Mab 2E10, Mab 3291 and sc2-4A5 and other anti-CSF-1R antibodies to human CSF-1R fragment delD4 and human CSF-1R Extracellular Domain (CSF-1R-ECD) at 25°C

Running buffer: PBS-T (Roche: 11 666 789 + 0.05% Tween20: 11 332 465)

5 Analyte sample:

Binding was measured at a flow rate of 30 μ L/min by one injection of the analyte with concentration $c = 10$ nM. (for Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 in second experiment) Each injection was 700 seconds long, followed by a dissociation phase of 180 seconds. Final regeneration was performed after each
10 cycle using 50 mM NaOH, contact time 60 seconds, flow rate 30 μ L/min.

Signals were measured by a report point 10 seconds after end of injection. Reference signals (signals from a blank reference flow cell (treated with EDC/NHS and ethanolamine, only) were subtracted to give the binding signals (as RU). If binding signals of nonbinding antibodies were slightly below 0 (Mab 2F11 = -3;
15 Mab 2E10 = -2; Mab 1G10 = - 6, Mab 2H7 = -9; and humanized hMab 2F11-e7 = - 7) the values were set as 0.

Table 3a:

Binding of <CSF-1R> MAbs to human CSF-1R fragment delD4 and CSF-1R-ECD and ratio at 25°C, measured by SPR

	<u>Binding to delD4 [RU]</u>	<u>Binding to CSF-1R-ECD [RU]</u>	<u>Ratio of binding of anti-CSF1R antibodies to CSF1R fragment delD4 / to CSF-1R-ECD</u>
Mab 3291	1015	627	1015/627= 1.61
sc2-4A5	374	249	374/249= 1.50
Mab 2F11	0	176	0/176 = 0
hMab 2F11-e7	0	237	0/237= 0
Mab 2E10	0	120	0/120 = 0
Mab 1G10	0	2708	0/2708 = 0
Mab 2H7	0	147	0/147 = 0
m<CCR5>Pz03.1C5	2	5	-

Mab 2F11 and Mab 2E10 showed binding to the human CSF-1R Extracellular Domain (CSF-1R-ECD) (see Fig. 2b); however no binding was detected to CSF-1R fragment delD4. (see Fig. 2a).

5 Sc2-4A5 and MAB3291 showed binding to CSF-1R-ECD and to del D4 (see Fig. 2b and 2a).

Thus the ratio of binding of anti-CSF1R antibodies Mab 2F11 and Mab 2E10 to CSF1R fragment delD4 / to CSF-1R-ECD was clearly below 1:50 (= 0.02), while the binding ratio of MAB3291 and Sc2-4A5 were 1.61 and 1.50, respectively and were highly above 1:50 (= 0.02). Negative control antibody m<CCR5>Pz03.1C5
10 did not show any binding (as expected).

Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 showed binding to the human CSF-1R Extracellular Domain (CSF-1R-ECD) (see Fig. 2d); however no binding was detected to CSF-1R fragment delD4. (see Fig. 2c). Thus the ratio of binding of anti-CSF1R antibodies Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 to
15 CSF1R fragment delD4 / to CSF-1R-ECD was clearly below 1:50 (= 0.02).

In a further experiment anti-CSF-1R antibodies 1.2.SM (ligand displacing CSF-1R antibody described in WO2009026303), CXIIG6 (ligand displacing CSF-1R antibody described in WO 2009/112245), the goat polyclonal anti-CSF-1R antibody ab10676 (abcam) were investigated. Anti-CSF-1R antibody Mab3291
20 (R&D-Systems) was used as reference control. Anti-CCR5 m<CCR5>Pz03.1C5 (deposited as DSM ACC 2683 on 18.08.2004 at DSMZ) was used as negative control.

Table 3b:
Binding of <CSF-1R> MAbs to human CSF-1R fragment delD4 and CSF-1R-ECD and ratio at 25°C, measured by SPR

	<u>Binding to delD4 [RU]</u>	<u>Binding to CSF-1R-ECD [RU]</u>	<u>Ratio of binding of anti-CSF1R antibodies to CSF1R fragment delD4 / to CSF-1R-ECD</u>
MAB3291	1790	1222	1790/1222= 1.47
1.2.SM	469	704	469/704 = 0.67
CXIIG6	1983	1356	1983/1356= 1.46
ab10676	787	547	787/547= 1.44
m<CCR5>Pz03.1C5	0	0	-

- 5 1.2.SM, CXIIG6, ab10676 and MAB3291 showed binding to CSF-1R-ECD and to del D4 (see Fig. 2f and 2e).

The binding ratio of 1.2.SM, CXIIG6, ab10676 and MAB3291 was highly above 1:50 (= 0.02). Negative control antibody m<CCR5>Pz03.1C5 did not show any binding (as expected).

10 **Example 5**

Growth inhibition of NIH3T3-CSF-1R recombinant cells in 3D culture under treatment with anti-CSF-1R monoclonal antibodies (CellTiterGlo-assay)

NIH 3T3 cells, retrovirally infected with either an expression vector for full-length wildtype CSF-1R (SEQ ID NO: 62) or mutant CSF-1R L301S Y969F (SEQ ID NO: 63), were cultured in DMEM high glucose media (PAA, Pasching, Austria) supplemented with 2mM L-glutamine, 2mM sodium pyruvate and non-essential amino acids and 10% fetal bovine serum (Sigma, Taufkirchen, Germany) on poly-HEMA (poly(2-hydroxyethylmethacrylate)) (Polysciences, Warrington, PA, USA)) coated dishes to prevent adherence to the plastic surface. Cells are seeded in medium replacing serum with 5ng/ml sodium selenite, 10mg/ml transferrin, 400µg/ml BSA and 0.05 mM 2-mercaptoethanol. When treated with 100ng/ml huCSF-1 (Biomol, Hamburg, Germany) wtCSF-1R (expressing cells form dense spheroids that grow three dimensionally, a property that is called anchorage independence. These spheroids resemble closely the three dimensional architecture

and organization of solid tumors in situ. Mutant CSF-1R recombinant cells are able to form spheroids independent of the CSF-1 ligand. Spheroid cultures were incubated for 3 days in the presence of different concentrations of antibody in order to determine an IC₅₀ (concentration with 50 percent inhibition of cell viability). The CellTiterGlo assay was used to detect cell viability by measuring the ATP-content of the cells.

Table 5a:

CSF-1R Mab	wtCSF-1R IC₅₀ [µg/ml]	Mutant CSF-1R IC₅₀ [µg/ml]
Mab 2F11	1.1	8.0
Mab 2E10	0.49	4.9
Mab 2H7	0.31	5.3
Mab 1G10	0.29	14.2
SC 2-4A5	10.0	10.0

Reference control Mab R&D-Systems 3291 did not show inhibition of mutant CSF-1R recombinant cell proliferation.

In a further experiment the anti-CSF-1R antibody according to the invention hMab 2F11-e7 and the anti-CSF-1R antibodies 1.2.SM (ligand displacing CSF-1R antibody described in WO2009026303), CXIIG6 (ligand displacing CSF-1R antibody described in WO 2009/112245), the goat polyclonal anti-CSF-1R antibody ab10676 (abcam), and SC 2-4A5 (Santa Cruz Biotechnology, US- see also Sherr, C.J. et al., Blood 73 (1989) 1786-1793) were investigated.

Spheroid cultures were incubated for 3 days in the presence of different concentrations of antibody in order to determine an IC₃₀ (concentration with 30 percent inhibition of cell viability). Maximum concentration was 20 µg/ml The CellTiterGlo assay was used to detect cell viability by measuring the ATP-content of the cells.

Table 5b:

CSF-1R Mab	wtCSF-1R IC₃₀ [µg/ml]	Mutant CSF-1R IC₃₀ [µg/ml]
hMab 2F11-e7	4.91	0.54
1.2.SM	1.19	> 20 µg/ml (-19% inhibition at 20 µg/ml = 19% stimulation)
CXIIG6	> 20 µg/ml (21% inhibition at 20 µg/ml)	> 20 µg/ml (-36% inhibition at 20 µg/ml = 36% stimulation)
ab10676	14.15	> 20 µg/ml (0% inhibition at 20 µg/ml)
SC 2-4A5	16.62	2.56

Example 6**Growth inhibition of BeWo tumor cells in 3D culture under treatment with anti-CSF-1R monoclonal antibodies (CellTiterGlo-assay)**

5

10

15

20

BeWo choriocarcinoma cells (ATCC CCL-98) were cultured in F12K media (Sigma, Steinheim, Germany) supplemented with 10% FBS (Sigma) and 2mM L-glutamine. 5×10^4 cells/well were seeded in 96-well poly-HEMA (poly(2-hydroxyethylmethacrylate)) coated plates containing F12K medium supplemented with 0.5 % FBS and 5% BSA. Concomitantly, 200 ng/ml huCSF-1 and 10µg/ml of different anti-CSF-1R monoclonal antibodies were added and incubated for 6 days. The CellTiterGlo assay was used to detect cell viability by measuring the ATP-content of the cells in relative light units (RLU). When BeWo spheroid cultures were treated with different anti-CSF-1R antibodies (10 µg/ml) inhibition of CSF-1 induced growth was observed. To calculate antibody-mediated inhibition the mean RLU value of unstimulated BeWo cells was subtracted from all samples. Mean RLU value of CSF-1 stimulated cells was set arbitrarily to 100%. Mean RLU values of cells stimulated with CSF-1 and treated with anti-CSF-1R antibodies were calculated in % of CSF-1 stimulated RLU. The Table 6 shows the calculated data of growth inhibition of BeWo tumor cells in 3D culture under treatment with anti-CSF-1R monoclonal antibodies; Fig.1a and b depicts normalized mean RLU values.

Table 6:

CSF-1R Mab	%inhibition 10µg/ml antibody concentration
CSF-1 only	0
Mab 2F11	70
Mab 2E10	102
Mab 2H7	103
Mab 1G10	99
SC 2-4A5	39

Example 7**Inhibition of human macrophage differentiation under treatment with anti-CSF-1R monoclonal antibodies (CellTiterGlo-assay)**

5

10

15

Human monocytes were isolated from peripheral blood using the RosetteSep™ Human Monocyte Enrichment Cocktail (StemCell Tech. - Cat. No.15028). Enriched monocyte populations were seeded into 96 well microtiterplates (2.5×10^4 cells/well) in 100 µl RPMI 1640 (Gibco - Cat. No.31870) supplemented with 10% FCS (GIBCO - Cat. No.011-090014M), 4 mM L-glutamine (GIBCO - Cat. No.25030) and 1x PenStrep (Roche Cat. No.1 074 440) at 37°C and 5% CO₂ in a humidified atmosphere. When 150 ng/ml huCSF-1 was added to the medium, a clear differentiation into adherent macrophages could be observed. This differentiation could be inhibited by addition of anti-CSF-1R antibodies. Furthermore, the monocyte survival is affected and could be analyzed by CellTiterGlo (CTG) analysis. From the concentration dependent inhibition of the survival of monocytes by antibody treatment, an IC₅₀ was calculated (see Table 7).

Table 7:

CSF-1R Mab	IC₅₀ [µg/ml]
Mab 2F11	0.08
Mab 2E10	0.06
Mab 2H7	0.03
Mab 1G10	0.06
SC 2-4A5	0.36

5 In a separate test series humanized versions of Mab 2 F11, e.g. hMab 2F11-c11, hMab 2F11-d8, hMab 2F11-e7, hMab 2F11-f12, showed IC₅₀ values of 0.07 µg/ml (hMab 2F11-c11), 0.07 µg/ml (hMab 2F11-d8), 0.04 µg/ml (hMab 2F11-e7) and 0.09 µg/ml (hMab 2F11-f12).

Example 8**Inhibition of cynomolgous macrophage differentiation under treatment with anti-CSF-1R monoclonal antibodies (CellTiterGlo-assay)**

10 Cynomolgous monocytes were isolated from peripheral blood using the CD14 MicroBeads non-human primate kit (Miltenyi Biotec - Cat.No. 130-091-097) according to the manufacturers description. Enriched monocyte populations were seeded into 96 well microtiterplates ($1-3 \times 10^4$ cells/well) in 100 µl RPMI 1640 (Gibco - Cat. No.31870) supplemented with 10% FCS (GIBCO - Cat. No.011-090014M), 4 mM L-glutamine (GIBCO - Cat. No.25030) and 1x PenStrep (Roche
15 Cat. No.1 074 440) at 37°C and 5% CO₂ in a humidified atmosphere. When 150 ng/ml huCSF-1 was added to the medium, a clear differentiation into adherent macrophages could be observed. This differentiation could be inhibited by addition of anti-CSF-1R antibodies. Furthermore, the monocyte survival is affected and
20 could be analyzed by CellTiterGlo (CTG) analysis. The viability was analyzed at a concentration of 5 µg/ml antibody treatment (see Table 8).

Table 8:

CSF-1R Mab	% survival	% inhibition (of survival) = (100% - %survival)
Mab 2F11	4 *	96
Mab 2E10	17 **	83
Mab 2H7	8	92
Mab 1G10	2	98
SC 2-4A5	31	69

* mean of four experiments (3 expts. using the murine, 1 expt. using the chimeric mAb)

** mean of two experiments using the murine mAb only

5 **Example 9**

Determination of the binding affinity of anti-CSF-1R antibodies to human CSF-1R

Instrument: BIACORE® A100

Chip: CM5 (Biacore BR-1006-68)

10 Coupling: amine coupling

Buffer: PBS (Biacore BR-1006-72), pH 7.4, 35°C

For affinity measurements 36 µg/ml anti mouse Fcγ antibodies (from goat, Jackson Immuno Reasearch JIR115-005-071) have been coupled to the chip surface for capturing the antibodies against CSF-1R. Human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1 –D5) (SEQ ID NO: 15 64) (R&D-Systems 329-MR or subcloned pCMV-presS-HisAvitag-hCSF-1R-ECD) was added in various concentrations in solution. Association was measured by an CSF-1R-injection of 1.5 minutes at 35 °C; dissociation was measured by washing the chip surface with buffer for 10 minutes at 35 °C. For calculation of 20 kinetic parameters the Langmuir 1:1 model was used.

Table 9:
Affinity data measured by SPR

CSF-1R Mab	K _D (nM)	k _a (1/Ms)	k _d (1/s)	t _{1/2} (min)
Mab 2F11	0.29	1.77E ⁺⁰⁵	5.18E ⁻⁰⁵	223
Mab 2E10	0.2	1.52E ⁺⁰⁵	2.97E ⁻⁰⁵	389
Mab 2H7	0.21	1.47E ⁺⁰⁵	3.12E ⁻⁰⁵	370
Mab 1G10	0.36	1.75E ⁺⁰⁵	6.28E ⁻⁰⁵	184

5 In a separate biacore binding assay using the CSF-1R ECD (data not shown) some competition of the antibodies Mab 2F11 and Mab 2E10 with the antibody Ab SC-2-4A5 was shown. However Mab 2F11/Mab 2E10 do not bind to the human CSF-1R fragment delD4, whereas Ab SC-2-4A5 binds to this delD4 fragment (see Example 4 and Fig 2a). Thus the binding region of Mab 2F11/Mab 2E10 is clearly distinct from the binding region of Ab SC-2-4A5, but probably located in a vicinity area. In such competition assay both antibodies Mab 2F11 and Mab 2E10 did not compete with Mab3291 from R&D-Systems (data not shown).

Example 10

Determination of the binding of anti-CSF-1R antibodies to human CSF-1R fragment D1-D3

15 Instrument: Biacore T100 (GE Healthcare)
Software: T100 Control, Version 1.1.11
B3000 Evaluation, Version 4.01
Scrubber, Version 2.0a

Assayformat Chip:CM5-Chip

20 Antibodies against CSF-1R were captured via amine coupled capture molecules. Using the single cycle kinetics five increasing concentrations of human CSF-1R fragment D1-D3 (SEQ ID NO: 66) were injected. Human CSF-1R fragment D1-D3 was subcloned into pCMV-presS-HisAvitag expression vector.

25 Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US; Sherr, C.J. et al., Blood 73 (1989) 1786-1793) which inhibits the ligand-receptor interaction, and Mab 3291 (R&D-Systems) were used as reference controls.

Capture molecules: Anti mouse Fc γ antibodies (from goat, Jackson Immuno Reasearch JIR115-005-071) for antibodies according to the invention and the R&D-Systems control Mab 3291 and Anti rat Fc γ antibodies (from goat, Jackson Immuno Reasearch JIR112-005-071) for the reference control anti CSF-1R SC 2-4A5.

5

Amine coupling of capture molecules

Standard amine coupling according to the manufacturer's instructions: running buffer: HBS-N buffer, activation by mixture of EDC/NHS, aim for ligand density of 2000 RU; the capture-Abs were diluted in coupling buffer NaAc, pH 4.5, c = 10 μ g/mL; finally remaining activated carboxyl groups were blocked by injection of 1 M Ethanolamin.

10

Kinetic characterization of human CSF-1R fragments D1-D3 binding to MAbs <CSF-1R> at 37°C

Running buffer: PBS (Biacore BR-1006-72)

15

Capturing of MAbs <CSF-1R> on flow cells 2 to 4: Flow 20 μ L/min, contact time 90 seconds, c(Abs<CSF-1R>) = 50 nM, diluted with running buffer + 1 mg/mL BSA;

Analyte sample:

Single Cycle Kinetics was measured at a flow rate of 30 μ L/min by five consecutive injections of the analyte with concentrations, c = 7.8 , 31.25, 125 500 and 2000 nM, without regeneration. Each injection was 30 seconds long and followed by a dissociation phase of 120 Seconds for the first four injections, and finally 1200 seconds for the highest concentration (=last injection).

20

Final regeneration was performed after each cycle using 10 mM Glycin pH 1.5 (Biacore BR-1003-54), contact time 60 seconds, flow rate 30 μ L/min.

25

Kinetic parameters were calculated by using the usual double referencing (control reference: binding of analyte to capture molecule; Flow Cell: subdomain CSF-1R concentration "0" as Blank) and calculation with model 'titration kinetics 1:1 binding with draft'.

Table 10:
Affinity data for binding of human CSF-1R fragment D1-D3 measured by SPR

CSF-1R Mab	Sub domain	K _D (nM)	k _a (1/Ms)	k _d (1/s)	t _{1/2} (min)
Mab 2F11	D1-D3	no binding			
Mab 2E10	D1-D3	no binding			
Mab 2H7	D1-D3	not determined			
Mab 1G10	D1-D3	no binding			
SC-2-4A5	D1-D3	no binding			
R&D-Systems 3291	D1-D3	5.4	2.2E ⁺⁵	1.2E ⁻³	9.6

5 The antibodies Mab 2F11, Mab 2E10 and Mab 1G10 showed no binding to human CSF-1R fragment D1-D3

Also reference control-Ab SC-2-4A5 did not bind to human CSF-1R fragment D1-D3.

10 The reference control Mab R&D-Systems 3291 showed binding to the human CSF-1R fragment D1-D3.

Example 11

CSF-1 level increase during CSF-1R inhibition in Cynomolgus monkey

15 Serum CSF-1 levels provide a pharmacodynamic marker of CSF-1R neutralizing activity of anti-human CSF-1R dimerization inhibitor hMab 2F11-e7. One male and one female cynomolgus monkey per dosage group (1 and 10 mg/kg) were intravenously administered anti-CSF1R antibody hMab 2F11-e7. Blood samples for analysis of CSF-1 levels were collected 1 week before treatment (pre-dose), 2, 24, 48, 72, 96, 168 hours post-dose and weekly for two additional weeks. CSF-1 levels were determined using a commercially available ELISA kit (Quantikine®
20 human M-CSF) according to the manufacturer's instructions (R&D Systems, UK). Monkey CSF-1 level were determined by comparison with CSF-1 standard curve samples provided in the kit.

Administration of hMab 2F11-e7 induced a dramatic increase in CSF-1 by ~ 1000-fold, which depending on the dose administered lasted for 48 hr (1mg/kg) or 15 days (10mg/kg). Hence, a dimerization inhibitor for CSF-1R offers the advantage to not directly compete with the dramatically upregulated ligand for binding to the receptor in contrast to a ligand displacing antibody.

Example 12

In vivo efficacy – tumor growth inhibition of anti-CSF-1R antibodies in breast cancer BT20 xenograft tumor cells in SCID beige mice

The human breast cancer cell line BT-20 expresses human CSF-1R but lacks CSF-1 expression (Sapi, E. et al Cancer Res 59 (1999) 5578-5585). Since the mouse derived CSF-1 fails to activate human CSF-1R on the tumor cells recombinant human CSF-1 (Biomol, Hamburg, Germany) was supplemented via osmotic minipumps (ALZET, Cupertino, CA) providing a continuous CSF-1 infusion rate of 2µg/day (Martin, T.A., Carcinogenesis 24 (2003) 1317-1323).

To directly compare the efficacy of an antibody interfering with dimerization of CSF-1R with a ligand displacing CSF-1R antibody we tested the chimeric anti-CSF-1R Mab 2F11 (antibody interfering with dimerization of CSF-1R) and 1.2.SM (ligand displacing CSF-1R antibody described in WO2009026303) in the BT-20 xenograft model.

SCID beige mice (Charles River, Sulzfeld, Germany) were subcutaneously coinjected with 1×10^7 cells BT-20 cells (ATCC HTB-19) and 100µl of Matrigel. Treatment of animals started at day of randomization at a mean tumor volume of 100 mm³. Mice are treated once weekly i.p. with the respective antibodies (see figure 4) in 20mM Histidine, 140 mM NaCl pH 6.0 buffer. The tumor dimensions are measured by caliper beginning on the staging day and subsequently 2 times per week during the whole treatment period. Tumor volume is calculated according to NCI protocol (Tumor weight = $1/2ab^2$, where “a” and “b” are the long and the short diameters of the tumor, respectively).

Tumor growth analysis is shown in Figure 4. Inhibition of human CSF-1R on tumor cells with the chimeric anti-CSF-1R Mab 2F11 was statistically more efficacious in mediating tumor growth inhibition than anti-CSF-1R antibody 1.2.SM (CSF-1R antibody described in WO2009026303).

Patent Claims

1. An antibody binding to human CSF-1R, characterized in that the antibody binds to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (SEQ ID NO: 64) with a ratio of 1:50 or lower.
5
2. The antibody according to claim 1, characterized in that
 - a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
 - b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
10
 - c) the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;
 - d) the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;
 - 15 or a humanized version thereof.
3. The antibody according to claim 1, characterized in that
 - a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
 - b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
20
 - c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
 - d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48, or
 - 25 e) the heavy chain variable domain is SEQ ID NO:55 and the light chain variable domain is SEQ ID NO:56.

4. The antibody according to claims 1, characterized in that
- a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or
- b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or
- c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or
- d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or
- e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or
- f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or

- 5 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54; or
- 10 h) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or
- 15 i) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.
5. The antibody according to any one of claims 1 to 4, characterized in that said antibody is of human IgG1 subclass or is of human IgG4 subclass.
6. A pharmaceutical composition characterized in comprising an antibody according to claims 1 to 5.
- 20 7. The antibody according to claims 1 to 5 for the treatment of cancer.
8. The antibody according to claims 1 to 5 for the treatment of bone loss.
9. The antibody according to claims 1 to 5 for the prevention or treatment of metastasis.
10. The antibody according to claims 1 to 5 for the treatment of inflammatory diseases.
- 25 11. A nucleic acid encoding a heavy chain of an antibody binding to CSF-1R, characterized in that said antibody comprises a variable domain according to claim 2 or 3.

12. An expression vector characterized in comprising a nucleic acid according to claim 11 for the expression of the antibody according to claims 1 to 5 in a prokaryotic or eukaryotic host cell.
- 5 13. A prokaryotic or eukaryotic host cell comprising a vector according to claim 12.
14. A method for the production of a recombinant antibody according to claims 1 to 5, characterized by expressing a nucleic acid according to claim 11 in a prokaryotic or eukaryotic host cell and recovering said antibody from said cell or the cell culture supernatant.
- 10 15. Use of the antibody according to claims 1 to 5 for the manufacture of a medicament for the treatment of cancer.
16. Use of the antibody according to claims 1 to 5 for the manufacture of a medicament for the treatment of bone loss.
- 15 17. Use of the antibody according to claims 1 to 5 for the manufacture of a medicament for the treatment of metastasis.
18. Use of the antibody according to claims 1 to 5 for the manufacture of a medicament for the treatment of inflammatory diseases.
19. A method for the treatment of a patient suffering from cancer characterized by administering to the patient an antibody according to claims 1 to 5.
- 20 20. A method for the treatment of a patient suffering from bone loss characterized by administering to the patient an antibody according to claims 1 to 5.
- 25 21. A method for the treatment of a patient suffering from metastasis characterized by administering to the patient an antibody according to claims 1 to 5.
22. A method for the treatment of a patient suffering from inflammatory diseases characterized by administering to the patient an antibody according to claims 1 to 5.

23. An antibody binding to human CSF-1R, characterized in that
- a) the heavy chain variable domain is SEQ ID NO:7 and the light chain variable domain is SEQ ID NO:8,
 - b) the heavy chain variable domain is SEQ ID NO:15 and the light chain variable domain is SEQ ID NO:16;
 - c) the heavy chain variable domain is SEQ ID NO:75 and the light chain variable domain is SEQ ID NO:76;
 - d) the heavy chain variable domain is SEQ ID NO:83 and the light chain variable domain is SEQ ID NO:84;
- or a humanized version thereof.
24. An antibody binding to human CSF-1R, characterized in that
- a) the heavy chain variable domain is SEQ ID NO:23 and the light chain variable domain is SEQ ID NO:24, or
 - b) the heavy chain variable domain is SEQ ID NO:31 and the light chain variable domain is SEQ ID NO:32, or
 - c) the heavy chain variable domain is SEQ ID NO:39 and the light chain variable domain is SEQ ID NO:40, or
 - d) the heavy chain variable domain is SEQ ID NO:47 and the light chain variable domain is SEQ ID NO:48, or
 - e) the heavy chain variable domain is SEQ ID NO:55 and the light chain variable domain is SEQ ID NO:56.
25. An antibody binding to human CSF-1R, characterized in that
- a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO:5, and a CDR1 region of SEQ ID NO:6, or

5 b) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:12, a CDR2 region of SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 14, or

10 c) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO:21, and a CDR1 region of SEQ ID NO:22, or

15 d) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 25, a CDR2 region of SEQ ID NO: 26, and a CDR1 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:28, a CDR2 region of SEQ ID NO: 29, and a CDR1 region of SEQ ID NO: 30, or

20 e) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38, or

25 f) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:41, a CDR2 region of SEQ ID NO: 42, and a CDR1 region of SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 44, a CDR2 region of SEQ ID NO:45, and a CDR1 region of SEQ ID NO:46, or

30 g) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:52, a CDR2 region of SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 54; or

5 h) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO:69, a CDR2 region of SEQ ID NO: 70, and a CDR1 region of SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region of SEQ ID NO: 72, a CDR2 region of SEQ ID NO:73, and a CDR1 region of SEQ ID NO:74, or

10 i) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 77, a CDR2 region of SEQ ID NO: 78, and a CDR1 region of SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region of SEQ ID NO:80, a CDR2 region of SEQ ID NO: 81, and a CDR1 region of SEQ ID NO: 82.

26. The antibody according to any one of claims 23 to 25, characterized in that said antibody is of human IgG1 subclass.

27. A pharmaceutical composition characterized in comprising an antibody according to claims 23 to 26.